



- (51) **International Patent Classification:** Not classified
- (21) **International Application Number:** PCT/IN20 12/0006 10
- (22) **International Filing Date:** 12 September 2012 (12.09.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
2570/MUM/201 1 13 September 201 1 (13.09.201 1) IN
2898/MUM/201 1 17 October 201 1 (17.10.201 1) IN
582/MUM/2012 5 March 2012 (05.03.2012) IN
- (71) **Applicant (for all designated States except US):** GLENMARK GENERICS LIMITED [IN/IN]; Glenmark House, HDO- Corporate Bldg, Wing-A, B. D. Sawant Marg, Chakala, Andheri (East), Mumbai 400099 (IN).
- (72) **Inventors; and**
- (75) **Inventors/ Applicants (for US only):** UPADHYE, Bhargav Krishnaji [IN/IN]; Plot No. 135, N-4, Sector-F, Cidco, Aurangabad 43 1003 (IN). JAGADALE, Shivaji Eknath [IN/IN]; Flat No. 003, Wing-A, Rajeshwari Park, Phase- 1, Opposite Chetna School, Malang Road, Kalyan (East) 421306 (IN). SONI, Mukesh [IN/IN]; House No.409, Lane No. 1, Devnagar, Taragad Road, Ajmer 305001 (IN).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17 :

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.1 7(H))
- of inventorship (Rule 4.17(iv))

Published:

- without international search report and to be republished upon receipt of that report (Rule 48.2(g))



WO 2013/072921 A2

(54) **Title:** PROCESS FOR PREPARATION OF SUBSTITUTED 3'-HYDRAZINO-BIPHENYL-3-CARBOXYLIC ACID COMPOUNDS

(57) **Abstract:** The present invention provides a process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds. The present invention further provides a process for the preparation of 3'-[N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino]-2'-hydroxybiphenyl-3-carboxylic acid, its intermediate compounds and pharmaceutically acceptable salts thereof.

PROCESS FOR PREPARATION OF SUBSTITUTED 3'-HYDRAZINO-BIPHENYL-3-CARBOXYLIC ACID COMPOUNDS

PRIORITY

[0001] This application claims the benefit to Indian Provisional Applications 2570/MUM/2011 filed on September 13, 2011, 2898/MUM/2011 filed on October 17, 2011 and 582/MUM/2012 filed on March 5, 2012 entitled "PROCESS FOR PREPARATION OF SUBSTITUTED 3'-HYDRAZINO-BIPHENYL-3-CARBOXYLIC ACID COMPOUNDS", which are incorporated herein by reference.


BACKGROUND OF THE INVENTION

10 Technical Field


[0002] The present invention relates to a process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds. The present invention relates to a process for the preparation of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid, its intermediate compounds and pharmaceutically acceptable salts thereof.

Description of the Related Art

[0003] Eltrombopag, also known as 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]-hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid is represented by the structure of formula Ia.



Ia




II

[0004] Eltrombopag olamine, compound of formula II, is a thrombopoietin receptor agonist indicated for the treatment of thrombocytopenia in patients with chronic immune (idiopathic) thrombocytopenic purpura who have had an insufficient response to corticosteroids, immunoglobulins, or splenectomy. Eltrombopag olamine is marketed under the brand name PROMACTA® in the United States (approved in


November 2008) and under the brand name 'REVOLADE[®]' in Europe (approved in March 2010).

[0005] Eltrombopag belongs to a class of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I,



5

(0006| United States Patent No. 7160870 (the '870 patent) discloses eltrombopag and its salts. United States Patent No. 7547719 discloses eltrombopag olamine, bisethanolamine salt of eltrombopag. The '870 patent discloses a process for the preparation of eltrombopag as schematically represented by Scheme I.




10

Scheme I


[0007] The object of the present invention is to provide a novel method, which is more convenient and more efficient than the previously known method for the synthesis of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds.

SUMMARY OF THE INVENTION


- 5 [0008] The present invention provides a process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I and salts thereof,



- wherein R represents hydrogen, linear or branched $C_{1,6}$ alkyl, $c_{3,8}$ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl; the process comprising:
- 10 a) reacting a compound of formula III,



- wherein R is as defined above, X is selected from the group consisting of Cl, Br, I, with a compound of formula IV.




15

- wherein R' represents boronic acid, boronic acid ester or halogen in the presence of a metal catalyst; and


(b) optionally, deprotecting the compound of formula I.

[0009] In another embodiment, the present invention provides a compound of formula III.




5 wherein X and R are as defined above.

[0010] In another embodiment, the present invention provides a compound of formula V;




wherein X, R and R'' are as defined above.

10 [0011] In another embodiment, the present invention provides a process for the preparation of eltrombopag, compound of formula Ia,




and pharmaceutically acceptable salts thereof, the process comprising:


(a) reacting 3-bromo-2-methoxyaniiline with ethylacetoacetate in the presence of alkali or alkaline earth metal nitrite and inorganic acid to give ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate, compound of formula Va;



5 (b) reacting the compound of formula Va with 3,4-dimethylphenylhydrazine or salt thereof to give 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone, compound of formula IIIa;




10 (c) reacting the compound of formula IIIa with 3-carboxyphenylboronic acid in the presence of a metal catalyst to give 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid, compound of formula Ib;




15 (d) deprotecting the compound of formula Ib using Lewis acid to give eltrombopag; and optionally, converting to its pharmaceutically acceptable salt.

[0012] In another embodiment, the present invention provides a process for the preparation of eltrombopag, a compound of formula Ia,



the process comprising subjecting the compound of formula I to a deprotection
5 reaction,



wherein R represents linear or branched C_{1-6} alkyl, C_{3-8} cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl.

10 [0013] In another embodiment, the present invention provides use of compound of formula Iia, Va, or Ib in the preparation of eltrombopag or salt thereof.

[0014] In another embodiment, the present invention provides an eltrombopag ammonium salt.

[0015] In another embodiment, the present invention provides a process for the
15 preparation of eltrombopag ammonium salt comprising reacting eltrombopag with a source of ammonia.

[0016] In another embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with excess of ethanolamine without using any additional reaction solvent.

[0017] In another embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising eltrombopag or a salt thereof with ethanolamine in an aqueous medium.

[0018] In another embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag ammonium salt with ethanolamine.

[0019] In another embodiment, the present invention provides use of eltrombopag ammonium salt in the preparation of eltrombopag olamine.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] Figure 1 is the proton NMR spectrum of ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate, compound of formula Va.


[0021] Figure 2 is the proton NMR spectrum of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone, compound of formula IIa.

[0022] Figure 3 is the proton NMR spectrum of 3'-{-2-[1-(ethoxycarbonyl)-2-oxopropylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid.

[0023] Figure 4 is the proton NMR spectrum of eltrombopag ammonium salt as obtained in Example 14c.


DETAILED DESCRIPTION OF THE INVENTION

[0024] The present invention provides a process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I and salts thereof,



wherein R represents hydrogen, linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl; the process comprising:

a) reacting a compound of formula III,



wherein R is as defined above, X is selected from the group consisting of Cl, Br, I, with a compound of formula IV, wherein R' represents boronic acid, boronic acid ester or halogen in the presence of a metal catalyst; and

10 (b) optionally, deprotecting the compound of formula I.

[0025] In the present application, the term "room temperature" means a temperature of about 25°C to about 30°C.

[0026] The term "linear or branched C₁₋₆ alkyl" includes groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl. The term "C₃₋₈ cycloalkyl" includes groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl. The term "optionally substituted benzyl" means benzyl which is optionally substituted with halo, alkyl, alkoxy or nitro group wherein halo includes Cl, Br, I; alkyl includes methyl, ethyl, propyl, butyl; alkoxy includes methoxy, ethoxy, propoxy. The term "linear or branched alkylalkoxy" includes groups such as methylethoxy, methylethoxy, ethylethoxy. The term "trialkylsilyl" includes groups such as trimethylsilyl, triethylsilyl, triisopropylsilyl, tert-butyldimethylsilyl. The term "acyl" includes groups such as acetyl, optionally substituted benzoyl, pivaloyl. The term "optionally substituted benzoyl" means benzoyl which is optionally substituted with halo or nitro group wherein halo includes Cl, Br, I.

25 [0027] In (a) of the above process, the compound of formula III, wherein X and R are as defined above is reacted with the compound of formula IV; wherein R' is as defined above, in the presence of a metal catalyst.

[0028] In one embodiment, the compound of formula III, wherein X and R are as defined above, is reacted with the compound of formula IV; wherein R' is boronic acid or boronic acid ester in the presence of a metal catalyst.

5 [0029] In one preferred embodiment, the compound of formula III, wherein R is C₁₋₆ alkyl, X is Br, is reacted with the compound of formula IV; wherein R' is boronic acid in the presence of a metal catalyst.

[0030] A suitable metal catalyst includes but is not limited to Pd(PPh₃)₄, PdCl₂(PPh₃)₂, PdCl₂(dppf), Pd(OAc)₂, NiCl₂(PPh₃)₂, PdCl₂(dppb). Preferably, the metal catalyst selected is PdCl₂(PPh₃)₂.

10 [0031] The reaction of compound of formula III with compound of formula IV may be carried out in the presence of base which includes organic base such as triethylamine, N-methylmorpholine, DBU; inorganic base such as lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate. Preferably, the base is selected from
15 inorganic base and more preferably the base is potassium hydroxide.

[0032] The reaction of compound of formula III with compound of formula IV may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to ethanol, methanol, 2-propanol, methyl acetate, ethyl acetate, acetone, ethylmethylketone, tetrahydrofuran, dioxane, toluene, dimethoxyethane, acetonitrile,
20 dimethylformamide, dimethyl sulfoxide; water or mixtures thereof. Preferably the solvent selected is ethanol-water mixture.

[0033] The reaction of compound of formula III with compound of formula IV may be carried out at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about 3 hours to
25 about 40 hours. Preferably the reaction is carried out at a temperature of about 70°C to about 85°C for a period of about 15 hour to about 30 hours.

[0034] In one embodiment, the compound of formula III, wherein X is Br and R is methyl, is reacted with the compound of formula IV; wherein R' is boronic acid, in the presence of PdCl₂(PPh₃)₂ catalyst and potassium hydroxide as base.

30 [0035] In one embodiment, the compound of formula III, wherein X and R are as defined above, is reacted with the compound of formula IV; wherein R' is halogen in the presence of a metal catalyst.

[0036] A suitable metal catalyst includes but is not limited to $\text{Pd}(\text{PPh}_3)_4$, $\text{PdCl}_2(\text{PPh}_3)_2$, $\text{NiCl}_2(\text{PPh}_3)_2$, $\text{Ni}(\text{PPh}_3)_2$. Preferably, the metal catalyst selected is $\text{PdCl}_2(\text{PPh}_3)_2$.

[0037] In one embodiment, the compound of formula III, wherein R is C_{1-6} alkyl, X is Br is reacted with the compound of formula IV; wherein R' is halogen selected from the group consisting of Cl, Br, I.

[0038] In one embodiment, the compound of formula III, wherein X is Br and R is methyl, is reacted with the compound of formula IV; wherein R' is Br, in the presence of $\text{PdCl}_2(\text{PPh}_3)_2$ catalyst.

[0039] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to ethanol, methanol, 1-propanol, 2-propanol, ethyl acetate, tetrahydrofuran, dioxane, acetone, toluene, dimethoxyethane, acetonitrile, dimethylformamide, dimethyl sulfoxide or mixtures thereof. Preferably the solvent selected is ethanol.

[0040] In (b) of the above process, the compound of formula I wherein $\text{R} \neq \text{H}$, is deprotected to give compound of formula Ia.

[0041] The deprotection reaction process includes any of the following:

(a) where R is C_{1-6} alkyl, C_{3-8} cycloalkyl, the deprotection of the compound of formula I is performed using protic acid such as hydroiodic acid, hydrobromic acid, hydrobromic acid/acetic acid, methanesulfonic acid, trifluoroacetic acid; Lewis acid selected from the group consisting of aluminium chloride, aluminium bromide, aluminium iodide, stannous chloride, stannous bromide, titanium chloride, boron trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium chloride, beryllium bromide, zinc chloride, zinc bromide, trimethylsilylchloride, trimethylsilylbromide, trimethylsilyliodide, lithium iodide, lithium iodide in refluxing 2,4,6-collidine, pyridine hydrochloride; sulphur compounds such as sodium ethylmercaptide, sodium trimethylsilanethiolate; alkali organomides such as sodium bis(trimethylsilyl)amide and lithium diisopropylamide; or

(b) where R is optionally substituted benzyl, methoxybenzyl, the deprotection of the compound of formula I is performed via hydrogenation reaction using hydrogen in the presence of a metal catalyst; or

(c) where R is linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, trityl, the deprotection of the compound of formula I is performed using an inorganic

acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; organic acid such as acetic acid; or

(d) where R is trialkylsilyl, the deprotection of the compound of formula I is performed using acids such as acetic acid or fluorides such as tetrabutylammonium fluoride; or

(e) where R is acyl, the deprotection of the compound of formula I is performed using an inorganic acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; inorganic base selected from the group consisting of sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate.

[0042] In one embodiment, the compound of formula I wherein R is methyl is deprotected using aqueous hydrobromic acid to give eltrombopag, compound of formula Ia.

[0043] The reaction may be carried out at a temperature in the range of about 25°C to about 125°C. The reaction is carried out for a period of about 2 hours to about 80 hours. Preferably the reaction is carried out at a temperature of about 110°C to about 115°C for a period of about 35 hours to about 65 hours.

[0044] In one embodiment, the compound of formula I wherein R is methyl is deprotected using Lewis acid such as aluminium chloride to give eltrombopag, compound of formula Ia.

[0045] In one embodiment, the compound of formula I wherein R is methyl is deprotected using Lewis acid such as aluminium chloride to give solid eltrombopag-aluminium complex which on acid treatment gives eltrombopag, compound of formula Ia.

[0046] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to chloroform, dichloromethane, dichloroethane, toluene, xylene, chlorobenzene, tetrahydrofuran, dioxane, tert-butylmethyl ether, dimethoxyethane. Preferably the solvent selected is toluene, tetrahydrofuran.

[0047] The reaction may be carried out at a temperature in the range of about 20°C to about 125°C. The reaction is carried out for a period of about 2 hours to about 80 hours. Preferably the reaction is carried out at a temperature of about 25°C to about 40°C for a period of about 35 hour to about 65 hours to give eltrombopag-aluminium complex.

[0048] The eltrombopag-aluminium complex on acid treatment gives eltrombopag, compound of formula Ia. The acid used includes hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, acetic acid. Preferably, the acid used is acetic acid. The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to tetrahydrofuran, methanol, ethanol, 2-propanol. The reaction may be carried out at a temperature in the range of about 25°C to about 125°C. The reaction is carried out for a period of about 2 hours to about 80 hours. Preferably the reaction is carried out at a temperature of about 40°C to about 120°C for a period of about 2 hour to about 24 hours.


10 [0049] In one embodiment, the present invention provides a process for the preparation of compound of formula I wherein R is H, the process comprising reacting compound of formula III wherein R is C₁₋₆ alkyl and X is Cl, Br, I, with compound of formula IV wherein R' is boronic acid in the presence of a metal catalyst to give compound of formula I wherein R is C₁₋₆ alkyl followed by deprotecting the compound of formula I wherein R is C₁₋₆ alkyl to give compound of formula I wherein R is H.

[0050] In one embodiment, the present invention provides a process for the preparation of compound of formula I wherein R is H, the process comprising reacting compound of formula III wherein R is methyl and X is Br. with compound of formula IV wherein R' is boronic acid in the presence of PdCl₂(PPh₃)₂ catalyst to give compound of formula I wherein R is methyl followed by deprotecting the compound of formula I wherein R is methyl using aqueous hydrobromic acid to give compound of formula I wherein R is H.

25 [0051] In one embodiment, the present invention provides a process for the preparation of compound of formula I wherein R is H, the process comprising reacting compound of formula III wherein R is methyl and X is Br, with compound of formula IV wherein R' is boronic acid in the presence of PdCl₂(PPh₃)₂ catalyst to give compound of formula I wherein R is methyl followed by deprotecting the compound of formula I wherein R is methyl using Lewis acid such as aluminium chloride to give compound of formula I wherein R is H.

[0052] In one embodiment, the present invention provides a process for isolating eltrombopag from an aqueous medium.


[0053] The present invention provides a compound of formula III,



wherein X and R are as defined above.

[0054] In one embodiment, the present invention provides a compound of formula III, wherein X is Br and R is methyl.

- 5 [0055] The present invention provides a process for the preparation of compound of formula III which comprises reacting a compound of formula V,



wherein X and R are as defined above and R'' represents hydrogen, Ci- & alkyl, with 3,4-dimethylphenylhydrazine or salt thereof to give the compound of formula III.

- 10 [0056j] The reaction may be carried out in the presence of alkali metal acetate such as sodium acetate, potassium acetate and the like; potassium carbonate, sodium carbonate. Preferably, alkali metal acetate is used; more preferably, sodium acetate is used.


- 15 [0057] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to acetic acid, methanol, ethanol, 2-propanol or mixtures thereof. Preferably the solvent selected is acetic acid.

- [0058] The reaction may be carried out at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about 1 hour to about 10 hours. Preferably the reaction is carried out at about the
20 reflux temperature of the solvent for a period of about 2 hours to about 5 hours.

[0059] In one embodiment, the present invention provides a process for the preparation of compound of formula III, the process comprising reacting a compound

of formula V with 3,4-dimethylphenylhydrazine or salt thereof to give the compound of formula III, wherein X is Br, R is methyl and R" is ethyl.


[0060] The present invention provides a compound of formula V,



5 wherein X, R and R" are as defined above.

[0061] In one embodiment, the present invention provides a compound of formula V, wherein X is Br, R is methyl and R" is ethyl.

[0062] The present invention provides a process for the preparation of compound of formula V which comprises reacting a compound of formula VI or its salt thereof,



10

wherein X and R are as defined above,

with alkyl acetoacetate or acetoacetic acid in the presence of alkali or alkaline earth metal nitrite and an inorganic acid in a solvent system to yield the compound of formula V.

15 [0063] The reaction may be carried out in the presence of alkali metal nitrite such as sodium nitrite, potassium nitrite and the like; alkaline earth metal nitrite such as calcium nitrite and the like. Preferably, sodium nitrite is used.


[0064] The inorganic acid is selected from the group consisting of hydrochloric acid, nitric acid, sulfuric acid, hydrobromic acid. Preferably, hydrochloric acid is used.

20 [0065] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to methanol, ethanol, 1-propanol, 2-propanol, water or mixtures thereof. Preferably the solvent selected is methanol, water, ethanol-water mixture.

[0066] The reaction may be carried out at a temperature in the range of about 0°C to about 10°C. The reaction is carried out for a period of about 2 hours to about 8 hours. Preferably the reaction is carried out at a temperature about 0°C to about 5°C for a period of about 2 hours to about 6 hours.


[0067] In one embodiment, the present invention provides a process for the preparation of compound of formula V, the process comprising reacting a compound of formula VI with alkyl acetoacetate in presence of sodium nitrite and hydrochloric acid to yield the compound of formula V, wherein X is Br, R is methyl and R" is ethyl.

[0068] The present invention provides a process for the preparation of eltrombopag, compound of formula Ia,




and pharmaceutically acceptable salts thereof, the process comprising:


- 10 (a) reacting 3-bromo-2-methoxyaniline with ethylacetoacetate in the presence of alkali or alkaline earth metal nitrite and inorganic acid to give ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate, compound of formula Va;



- 15 (b) reacting the compound of formula Va with 3,4-dimethylphenylhydrazine or salt thereof to give 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone, compound of formula Iia;



(c) reacting the compound of formula IIIa with 3-carboxyphenylboronic acid in the presence of a metal catalyst to give 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid, compound of formula Ib;



(d) deprotecting the compound of formula Ib using Lewis acid to give eltrombopag; and optionally, converting to its pharmaceutically acceptable salt.

[0069] In (a) of the above process, 3-bromo-2-methoxyaniline is reacted with ethylacetoacetate in presence of alkali or alkaline earth metal nitrite and inorganic acid to give ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate, compound of formula Va.

[0070] The reaction may be carried out in the presence of alkali metal nitrite such as sodium nitrite, potassium nitrite and the like; alkaline earth metal nitrite such as calcium nitrite and the like. Preferably, sodium nitrite is used. The inorganic acid is selected from the group consisting of hydrochloric acid, nitric acid, sulfuric acid, hydrobromic acid. Preferably, hydrochloric acid is used. The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not

limited to methanol, ethanol, 1-propanol, 2-propanol, water or mixtures thereof. Preferably the solvent selected is methanol, water, ethanol-water mixture.

[0071] In (b) of the above process, the compound of formula Va is reacted with 3,4-dimethylphenylhydrazine or salt thereof to give I-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone, compound of formula Iia.

[0072] The reaction may be carried out in the presence of alkali metal acetate such as sodium acetate, potassium acetate and the like; sodium carbonate, potassium carbonate. Preferably, the alkali metal acetate used is sodium acetate. The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to acetic acid, methanol, ethanol or mixtures thereof. Preferably the solvent selected is acetic acid. The reaction may be carried out at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about 1 hour to about 10 hours. Preferably the reaction is carried out at to about the reflux temperature of the solvent for a period of about 2h to about 5h.

[0073] In (c) of the above process, the compound of formula Iia with 3-carboxyphenylboronic acid in presence of metal catalyst to give 3'-{N'-[I-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid, compound of formula Ib.

[0074] A suitable metal catalyst includes but is not limited to Pd(PPh₃)₄, PdCl₂(PPh₃)₂, PdCl₂(dppf), Pd(OAc)₂, NiCl₂(PPh₃)₂, PdCl₂(dppb). Preferably, the metal catalyst selected is PdCl₂(PPh₃)₂. The reaction may carried out in the presence of base which includes organic base such as triethylamine, N-methylmorpholine, DBU; inorganic base such as lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate. Preferably, the base is selected from inorganic base and more preferably the base is potassium hydroxide. The reaction may carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to ethanol, methanol, 2-propanol, methyl acetate, ethyl acetate, acetone, ethylmethylketone, tetrahydrofuran, dioxane, toluene, dimethoxyethane, acetonitrile, dimethylformamide; water or mixtures thereof. Preferably the solvent selected is ethanol-water mixture. The reaction may be carried out at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about


3 hours to about 40 hours. Preferably the reaction is carried out at a temperature of about 70°C to about 85°C for a period of about 15 hours to about 30 hours.

[0075] In (d) of the above process, the compound of formula Ib is deprotected using Lewis acid to give eltrombopag.


5 [0076] In one embodiment, the compound of formula Ib is deprotected using Lewis acid selected from the group consisting of aluminium chloride, aluminium bromide, aluminium iodide, stannous chloride, stannous bromide, titanium chloride, boron trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium chloride, beryllium bromide, zinc chloride, zinc bromide, trimethylsilylchloride,
10 trimethylsilylbromide, trimethylsilyliodide.

[0077] In one embodiment, the compound of formula Ib is deprotected using Lewis acid such as aluminium chloride to give solid eltrombopag-aluminium complex which on acid treatment gives eltrombopag. The deprotection reaction process is as discussed supra.

15 [0078] The present invention provides a process for the preparation of eltrombopag, a compound of formula Ia,



the process comprising subjecting the compound of formula I to a deprotection reaction,



wherein R represents linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl.

5 [0079] In one embodiment, the present invention provides a process for the preparation of eltrombopag comprising subjecting the compound of formula I wherein R is C₁₋₆ alkyl, C₃₋₈ cycloalkyl, to a deprotection reaction using protic acid such as hydroiodic acid, hydrobromic acid, hydrobromic acid/acetic acid, methanesulfonic acid, trifluoroacetic acid; Lewis acid selected from the group consisting of aluminium chloride, aluminium bromide, aluminium iodide, stannous chloride, stannous bromide, titanium chloride, boron trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium chloride, beryllium bromide, zinc chloride, zinc bromide, trimethylsilylchloride, trimethylsilylbromide, trimethylsilyliodide, lithium iodide, lithium iodide in refluxing 2,4,6-collidine, pyridine hydrochloride; sulphur compounds such as sodium ethylmercaptide, sodium trimethylsilylanethiolate; alkali organometallics such as sodium bis(trimethylsilyl)amide and lithium diisopropylamide.

10

15

[0080] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein where R is optionally substituted benzyl, methoxybenzyl, to a deprotection reaction by hydrogenation using hydrogen in the presence of a metal catalyst.

20

[0081] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein R is linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, trityl, to a deprotection reaction using an inorganic acid selected

25

from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; organic acid such as acetic acid.

[0082] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein R is trialkylsilyl, to a deprotection reaction using acids such as acetic acid or fluorides such as tetrabutylammonium fluoride.

[0083] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein R is acyl, to a deprotection reaction using an inorganic acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; inorganic base selected from the group consisting of sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate.

[0084] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein R is methyl, to a deprotection reaction using hydrobromic acid.

[0085] In one embodiment, the present invention provides a process for the preparation of eltrombopag, the process comprising subjecting the compound of formula I wherein R is methyl, to a deprotection reaction using Lewis acid.

[0086] In one embodiment, the present invention provides a process for the preparation of eltrombopag, compound of formula Ia, the process comprising deprotecting the compound of formula I wherein R is methyl, using a Lewis acid such as aluminium chloride to give solid eltrombopag-aluminium complex, then on acid treatment gives eltrombopag. compound of formula Ia.

[0087] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to chloroform, dichloromethane, dichloroethane, toluene, xylene, chlorobenzene, tetrahydrofuran, dioxane, tert-butylmethyl ether, dimethoxyethane. Preferably the solvent selected is toluene, tetrahydrofuran.

[0088] The reaction may be carried out at a temperature in the range of about 20°C to about 125°C. The reaction is carried out for a period of about 2 hours to about 80 hours. Preferably the reaction is carried out at a temperature of about 25°C to about 40°C for a period of about 35 hours to about 65 hours to give eltrombopag-aluminium complex.

[0089] The eltrombopag-aluminium complex on acid treatment gives eltrombopag, compound of formula Ia. The acid used includes hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, acetic acid. Preferably, the acid used is acetic acid. The suitable solvent includes but is not limited to tetrahydrofuran, methanol, ethanol, 2-propanol. The reaction may be carried out at a temperature in the range of about 25°C to about 125°C. The reaction is carried out for a period of about 2 hours to about 80 hours. Preferably the reaction is carried out at a temperature of about 40°C to about 120°C for a period of about 2 hour to about 24 hours.

[0090] The present invention provides an eltrombopag-aluminium complex.


[0091] The present invention provides a solid eltrombopag-aluminium complex.

[0092] The present invention provides use of compound of formula III, or V in the preparation of eltrombopag or salt thereof.

[0093] The present invention provides use of compound of formula I, wherein R represents linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl in the preparation of eltrombopag or salt thereof.

[0094] In one embodiment, the present invention provides use of compound of formula IIIa, Va, or Ib in the preparation of eltrombopag and salt thereof.


[0095] The present invention provides a process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I and salts thereof,



I

wherein R represents hydrogen, linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl; the process comprising:

25 a) reacting a compound of formula VII,



VII

wherein R is as defined above and R'' represents hydrogen, C₁₋₆ alkyl,

with 3,4-dimethylphenylhydrazine or salt thereof to give the compound of formula I;

and

- 5 b) optionally, deprotecting the compound of formula I.


[0096] In (a) of the above process, the reaction may be carried out in the presence of
alkali metal acetate such as sodium acetate, potassium acetate and the like; potassium
carbonate. sodium carbonate. Preferably, alkali metal acetate is used; more preferably,
sodium acetate is used. The reaction may be carried out in the presence of a suitable
10 solvent. The suitable solvent includes but is not limited to acetic acid, methanol,
ethanol, 2-propanol or mixtures thereof. Preferably the solvent selected is acetic acid.
The reaction may be carried out at a temperature in the range of about 25°C to about
the reflux temperature of the solvent. The reaction is carried out for a period of about
1 hour to about 10 hours. Preferably the reaction is carried out at about the reflux
15 temperature of the solvent for a period of about 2h to about 5h.

[0097] In one embodiment, the compound of formula VII is reacted with 3,4-
dimethylphenylhydrazine or salt thereof to give the compound of formula I, wherein
R is methyl and R'' is ethyl.

[0098] In (b) of the above process, the compound of formula I wherein R≠H, is
20 deprotected to give eltrombopag, compound of formula Ia. The deprotection reaction
process is as discussed supra.

[0099] In one embodiment, the present invention provides a process for the
preparation of compound of formula I wherein R is H, the process comprising
reacting compound of formula VII wherein R is C₁₋₆ alkyl and R'' is hydrogen, C₁₋₆
25 alkyl with 3,4-dimethylphenylhydrazine or salt thereof to give the compound of
formula I wherein R is C₁₋₆ alkyl followed by deprotecting the compound of formula I
wherein R is C₁₋₆ alkyl using aqueous hydrobromic acid, Lewis acid such as
aluminium chloride to give compound of formula I wherein R is H.

[0100] The present invention provides a compound of formula VII,




VII

wherein R represents hydrogen, linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl; and R'' represents hydrogen, C₁₋₆ alkyl.

[0101] In one embodiment, the present invention provides a compound of formula V, wherein R is methyl and R'' is ethyl.

[0102] The present invention provides a process for the preparation of compound of formula VII, the process comprising:


a) reacting a compound of formula VI, wherein X and R are as defined above, with a compound of formula IV, wherein R' is as defined above,



VI

IV

15 in the presence of a metal catalyst to give a compound of formula VIII,



VIII

wherein R is as defined above; and

b) reacting the compound of formula VIII with alkyl acetoacetate or acetoacetic acid in the presence of alkali or alkaline earth metal nitrite and an inorganic acid to yield the compound of formula VII.

[0103] In (a) of the above process, the compound of formula VI, wherein X and R are as defined above, is reacted with the compound of formula IV; wherein R' is boronic

acid or boronic acid ester or halogen in the presence of a metal catalyst to give the compound of formula VIII, wherein R is as defined above.

[0104] A suitable metal catalyst includes but is not limited to Pd(PPh₃)₄, PdCl₂(PPh₃)₂, PdCl₂(dppf), Pd(OAc)₂, NiCl₂(PPh₃)₂, PdCl₂(dppb). Preferably, the metal catalyst selected is PdCl₂(PPh₃)₂. The reaction of compound of formula VI with compound of formula IV may be carried out in the presence of base which includes organic base such as triethylamine, N-methylmorpholine, DBU; inorganic base such as lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate. Preferably, the base is selected from inorganic base and more preferably the base is potassium hydroxide, potassium carbonate. The reaction of compound of formula VI with compound of formula IV may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to ethanol, methanol, 2-propanol, methyl acetate, ethyl acetate, acetone, ethylmethylketone, tetrahydrofuran, dioxane, toluene, dimethoxyethane, acetonitrile, dimethylformamide, dimethyl sulfoxide; water or mixtures thereof. Preferably the solvent selected is ethanol-water mixture. The reaction of compound of formula VI with compound of formula IV may be carried out at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about 3 hours to about 40 hours. Preferably the reaction is carried out at a temperature of about 70°C to about 85°C for a period of about 15 hour to about 30 hours.

[0105] In one embodiment, the compound of formula VI, wherein R is C₁₋₆ alkyl, X is Br, is reacted with the compound of formula IV; wherein R' is boronic acid in the presence of a metal catalyst.

[0106] In one embodiment, the compound of formula VI, wherein X is Br and R is methyl, is reacted with the compound of formula IV; wherein R' is boronic acid, in the presence of PdCl₂(PPh₃)₂ catalyst and potassium carbonate as base.


[0107] In one embodiment, the compound of formula VI, wherein R is C₁₋₆ alkyl, X is Br is reacted with the compound of formula IV; wherein R' is halogen selected from the group consisting of Cl, Br, I.

[0108] In (b) of the above process, the reaction may be carried out in the presence of alkali metal nitrite such as sodium nitrite, potassium nitrite and the like; alkaline earth metal nitrite such as calcium nitrite and the like. Preferably, sodium nitrite is used. The inorganic acid is selected from the group consisting of hydrochloric acid, nitric

acid, sulfuric acid, hydrobromic acid. Preferably, hydrochloric acid is used. The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to methanol, ethanol, 1-propanol, 2-propanol, water or mixtures thereof. Preferably the solvent selected is methanol-water mixture.

5 [0109] In one embodiment, the compound of formula VIII, wherein R is C₁₋₆ alkyl, is reacted with alkyl acetoacetate or acetoacetic acid in presence of sodium nitrite and hydrochloric acid to yield the compound of formula VII, wherein R is C₁₋₆ alkyl and R" is hydrogen. C₁₋₆ alkyl.

10 [0110] The present invention provides a process for the preparation of compound of formula VII which comprises reacting a compound of formula V,



15 wherein X, R and R" are as defined above, with a compound of formula IV, wherein R' is as defined above, in the presence of a metal catalyst to give the compound of formula VII.

[0111] In one embodiment, the compound of formula V, wherein X and R are as defined above, is reacted with the compound of formula IV; wherein R' is boronic acid or boronic acid ester or halogen in the presence of a metal catalyst.

20 [0112] A suitable metal catalyst includes but is not limited to Pd(PPh₃)₄, PdCl₂(PPh₃)₂, PdCl₂(dppf), Pd(OAc)₂, NiCl₂(PPh₃)₂, PdCl₂(dppb). Preferably, the metal catalyst selected is PdCl₂(PPh₃)₂. The reaction of compound of formula V with compound of formula IV may be carried out in the presence of base which includes organic base such as triethylamine, N-methylmorpholine, DBU; inorganic base such as lithium hydroxide, sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate, sodium bicarbonate, potassium bicarbonate. Preferably, the base is selected from inorganic base and more preferably the base is potassium hydroxide, potassium carbonate. The reaction of compound of formula V with compound of formula IV may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to ethanol, methanol, 2-propanol, methyl acetate,


ethyl acetate, acetone, ethylmethylketone, tetrahydrofuran, dioxane, toluene, dimethoxyethane, acetonitrile, dimethylformamide, dimethyl sulfoxide; water or mixtures thereof. Preferably the solvent selected is ethanol-water mixture. The reaction of compound of formula V with compound of formula IV may be carried out
 5 at a temperature in the range of about 25°C to about the reflux temperature of the solvent. The reaction is carried out for a period of about 3 hours to about 40 hours. Preferably the reaction is carried out at a temperature of about 70°C to about 85°C for a period of about 15 hour to about 30 hours.

[0113] In one embodiment, the compound of formula V, wherein R is C₁₋₆ alkyl, X is Br, is reacted with the compound of formula IV; wherein R' is boronic acid in the presence of a metal catalyst.
 10

[0114] In one embodiment, the compound of formula V, wherein X is Br and R is methyl, is reacted with the compound of formula IV; wherein R' is boronic acid, in the presence of PdCl₂(PPh₃)₂ catalyst and potassium carbonate as base.

[0115] In one embodiment, the compound of formula V, wherein R is C₁₋₆ alkyl, X is Br is reacted with the compound of formula IV; wherein R' is halogen selected from the group consisting of Cl, Br, I.
 15

[0116] The present invention provides a process for the deprotection of a compound of formula III,



20

wherein R represents linear or branched C₁₋₆ alkyl, C₃₋₈ cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl, and X is selected from the group consisting of Cl, Br, I.

[0117] The deprotection of compound of formula III may be carried out as described for compound of formula I wherein R≠H, as discussed supra.
 25

- [0118] In one embodiment, the present invention provides a process for the deprotection of the compound of formula III, the process comprising subjecting the compound of formula III wherein R is methyl, to a deprotection reaction using hydrobromic acid, Lewis acid such as aluminium chloride.
- 5 [0119] The present invention provides an eltrombopag ammonium salt.
- [0120] The present invention provides an eltrombopag ammonium salt characterized by a proton NMR spectrum having peaks at δ 14.81 (brs, 1H), 8.17 (s, 1H), 7.80-7.82(m, 3H), 7.65-7.68 (d, 1H), 7.34-7.42 (m, 2H), 7.01-7.07 (m, 2H), 6.87 (t, 1H), 2.37 (s, 3H), 2.22 (s, 3H), 2.18 (s, 3H).
- 10 [0121] The present invention provides an eltrombopag ammonium salt with ammonia content in the range of about 3.5% to about 8.5%.
- [0122] The present invention provides an eltrombopag ammonium salt which may be mono-ammonium salt or di-ammonium salt, preferably in crystal line form or amorphous form.
- 15 [0123] The present invention provides a process for the preparation of eltrombopag ammonium salt comprising reacting eltrombopag with a source of ammonia.
- [0124] A suitable source of ammonia includes but is not limited to ammonia water (aqueous ammonia), ammonium carbonate, ammonia gas, liquid ammonia. Preferably, the source of ammonia selected is ammonia water.
- 20 [0125] The reaction may be carried out in the presence of a suitable solvent. The suitable solvent includes but is not limited to methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, methyl acetate, ethyl acetate, n-propyl acetate, tert-butyl acetate, acetone, ethylmethylketone, methyl isobutyl ketone, chloroform, dichloromethane, dichloroethane, tetrahydrofuran, dioxane, tert-butylmethyl ether,
- 25 toluene, xylene, chlorobenzene, dimethoxyethane, acetonitrile, dimethylformamide; water or mixtures thereof. Preferably the solvent selected is tetrahydrofuran.
- [0126] The reaction may be carried out at a temperature in the range of about 10°C to about 40°C. The reaction is carried out for a period of about 1 hour to about 20 hours. Preferably the reaction is carried out at a temperature of about 20°C to about 35°C for
- 30 a period of about 1 hour to about 5 hours.
- [0127] In one embodiment, the present invention provides a process for isolating eltrombopag as an eltrombopag ammonium salt.

[0128] Eltrombopag may be converted to its pharmaceutically acceptable salts such as its monoethanolamine, bisethanolamine salts. Preferably, eltrombopag is converted to its bisethanolamine salt namely, eltrombopag olamine, compound of formula II.

5 [0129] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with excess of ethanolamine without using any additional reaction solvent.

[0130] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with ethanolamine in an aqueous medium.

10 [0131] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with ethanolamine in water.

[0132] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with ethanolamine in a solvent and isolating eltrombopag olamine by addition of anti-solvent.

[0133] The solvent that may be utilized for this step includes, but is not limited to alcohols such as methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, 1-pentanol, 1-octanol and the like; ethers such as tert-butylmethyl ether, tetrahydrofuran, dioxane and the like; esters such as methyl acetate, ethyl acetate, n-propyl acetate, and tert-butyl acetate and the like; ketones such as acetone, ethyl methyl ketone and methyl isobutyl ketone and the like; dimethyl sulfoxide; dimethyl formamide; dimethyl acetamide; water and mixtures thereof.

25 [0134] The anti-solvent that may be utilized for this step includes, but is not limited to ethers such as diethyl ether, dimethyl ether, diisopropyl ether, tetrahydrofuran, dioxane; hydrocarbons such as n-hexane, n-heptane, cyclohexane; water and mixtures thereof: preferably non-polar anti-solvent is used.

[0135] The present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag ammonium salt with ethanolamine.

30 [0136] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag ammonium salt with ethanolamine, preferably, without using any additional reaction solvent.

[0137] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine comprising reacting eltrombopag ammonium salt with ethanolamine in the presence of a solvent.

[0138] The solvent that may be utilized for this step includes, but is not limited to
5 methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-butanol, methyl acetate, ethyl acetate, n-propyl acetate, tert-butyl acetate, acetone, ethylmethylketone, methyl isobutyl ketone, tetrahydrofuran, dioxane, tert-butylmethyl ether, toluene, dimethoxyethane, acetonitrile, dimethylformamide; water or mixtures thereof.


[0139] The present invention provides use of eltrombopag ammonium salt in the
10 preparation of eltrombopag olamine.

[0140] In one embodiment, the present invention provides a process for the preparation of eltrombopag olamine, the process comprising: (a) treating a reaction mixture containing eltrombopag with a base to give eltrombopag salt, and (b) reacting the eltrombopag salt with ethanolamine to give eltrombopag olamine directly,
15 wherein eltrombopag is formed in-situ in the reaction. The eltrombopag salt obtained in this step includes, but is not limited to eltrombopag sodium salt, eltrombopag potassium salt, eltrombopag ammonium salt.


[0141] In one preferred embodiment, the present invention provides a process for the preparation of eltrombopag olamine, the process comprising: (a) treating a reaction
20 mixture containing eltrombopag with a source of ammonia to give eltrombopag ammonium salt, and (b) reacting the eltrombopag ammonium salt with ethanolamine to give eltrombopag olamine directly, wherein eltrombopag is formed in-situ in the reaction.

[0142] The present invention provides eltrombopag and a salt thereof, having a
25 compound of formula IIa in less than about 0.5%, preferably less than about 0.15%, more preferably less than about 0.05%.

[0143] The present invention provides eltrombopag and a salt thereof, having a compound of formula IIIb in less than about 0.5%, preferably less than about 0.15%, more preferably less than about 0.05%.




[0144] The present invention provides eltrombopag and a salt thereof, having a compound of formula IXa in less than about 0.5%, preferably less than about 0.15%, more preferably less than about 0.05%.



5

[0145] The present invention provides eltrombopag and a salt thereof, having a compound of formula IXb in less than about 0.5%, preferably less than about 0.15%, more preferably less than about 0.05%.



10 [0146] The present invention provides substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I, salts thereof and intermediate compounds, obtained by the above processes, as characterized and analyzed by following techniques:

A. Proton NMR spectra were recorded in CDCl_3 and DMSO-d_6 using NMR instrument- Varian 300 MHZ

B. IR spectra were recorded using IR instrument- Perkin Elmer Spectrum One FTIR and all the samples prepared in KBr.

C. Mass spectra were recorded using instrument- Thermofinnigan, LCQ DECA XP MAX

5 D. HPLC

E. Melting point

10 |0 147| The following examples are provided to enable one skilled in the art to practice the invention and are merely illustrative of the invention. The examples should not be read as limiting the scope of the invention as defined in the features and advantages.

15

20

25

30

EXAMPLES

[0148] EXAMPLE 1: Preparation of Ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate

To a solution of 3-bromo-2-methoxyaniline (20g) in 1N hydrochloric acid (400mL) was added a solution of sodium nitrite (7.2g in 720mL of water) at about 0°C to about 5°C under stirring. The reaction mixture was stirred for about 15 minutes at about 5°C. Then ethylacetoacetate (12.9g) was added to the reaction mixture and stirred for about 15 minutes at about 0°C to about 5°C. Sodium bicarbonate solution (27.5g in 300mL water) and ethanol (400mL) was then added to the reaction mixture. The reaction mixture was allowed to warm to about room temperature and stirred for about 2 hours. The mixture was filtered, washed with water (200mL) and dried to get yellowish solid. Yield: 33g; Melting point: 75.9-77.1°C; Purity (HPLC): 99.12%

IR: 3421, 1706, 1684, 1517, 1215, 1093, 980cm⁻¹; Mass: m/z 342.88 [M+] and 344.86 [M+2]

¹H NMR (300 MHz in CDCl₃): δ 12.86 (s,1H), 7.58-7.61 (d,1H), 7.30-7.33 (d,1H), 7.02-7.07 (m,1H), 4.31-4.43 (q,2H), 3.95 (s,3H), 2.5 (s,3H), 1.38-1.43 (t,3H)

[0149] EXAMPLE 2: Preparation of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone

A solution of ethyl-2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate (5g) prepared as in Example 1, and 3,4-dimethylphenylhydrazine hydrochloride (2.48g) and sodium acetate (1.4g) in glacial acetic acid (100mL) was stirred and heated to reflux for about 3 hours. The mixture was cooled to about room temperature and stirred for about 1 hour. It was filtered, washed with water (25mL) and dried in an oven at about 55°C to about 60°C for about 12 hours to obtain an orange solid product. Yield: 4.15g; Melting point: 197.4-198.2°C; Purity (HPLC): 99.33%

IR: 3444, 1559, 1337, 1256, 1114, 1019cm⁻¹; Mass: m/z 415 [M+] and 417 [M+2]

¹H NMR (300 MHz in CDCl₃): δ 13.7 (brs,1H), 7.64-7.73 (m,3H), 7.32-7.35 (d,1H), 7.16-7.19 (d,1H), 7.07-7.16 (t,1H), 4.01 (s,3H), 2.36 (s,3H), 2.31 (s,3H), 2.27 (s,3H)

[0150] EXAMPLE 3: Preparation of 3MN'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid

A mixture of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone (5g) prepared as in Example 2, and 3-carboxyphenylboronic acid (2.99g), bis(triphenylphosphine)palladium(II)chloride

[PdCl₂(PPh₃)₂] (0.42g), potassium hydroxide (1.74g) in ethanol (250mL) and water (30mL) was heated to about reflux temperature and stirred for about 24 hours. The hot mixture was filtered to remove the catalyst. The clear filtrate was concentrated under vacuum. To the residue, water (100mL) was added; acidified to pH 3-5 using hydrochloric acid and stirred for about 30 minutes at about room temperature. The slurry was filtered to give a yellowish orange solid product, which was washed with water (50mL) and dried in an oven at about 55°C to about 60°C for about 12 hours.

Yield: 4g; Melting point: 256.5-257.2°C; Mass: m/z 455.61 [M-1]

IR: 3413, 1687, 1546, 1501, 1258, 1003cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 13.7 (brs, 1H), 13 (brs, 1H), 8.17 (s, 1H), 7.98-8.01 (d, 1H), 7.87-7.96 (d, 1H), 7.78-7.84 (d, 1H), 7.58-7.75 (m, 3H), 7.36-7.52 (m, 1H), 7.29-7.34 (d, 1H), 7.18-7.26 (d, 1H), 3.44 (s, 3H), 2.32 (s, 3H), 2.25 (s, 3H), 2.21 (s, 3H)

[0151] EXAMPLE 4: Preparation of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid

A solution of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid (4g), prepared as in Example 3, in 48% aqueous hydrobromic acid (35mL) and glacial acetic acid (35mL) was stirred and heated under reflux for about 60 hours. The reaction mixture was concentrated under vacuum to a thick residue. Water (40mL) was added and basified to pH 7-8 using saturated solution of sodium bicarbonate (20mL). The mixture was filtered to give an orange solid, which was washed with water (25mL) and dried in an oven at 55°C about to about 60°C for about 12 hours.

Yield: 3g; Melting point: 231.6-232.9°C; Mass: m/z 443.16 [M+1]

IR: 3292, 1707, 1541, 1503, 1223, 1118, 1000cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 13.75 (brs, 1H), 13.09 (brs, 1H), 9.69 (s, 1H), 8.13 (s, 1H), 7.94-7.97 (d, 1H), 7.79-7.82 (d, 1H), 7.62-7.71 (m, 3H), 7.16-7.21 (m, 3H), 2.32 (s, 3H), 2.26 (s, 3H), 2.22 (s, 3H)

[0152] EXAMPLE 5: Preparation of eltrombopag olamine

To a solution of ethanolamine (0.9g) in ethanol (100mL) was added a solution of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid (3g), prepared as in Example 4, in tetrahydrofuran (40mL) at about room temperature over about 30 to 40 minutes and stirred for about 3 hours at about room temperature. The mixture was

filtered, washed with ethanol (25mL) and dried in an oven at about 55°C to about 60°C for about 12 hours. Yield: 3.7g

IR: 3421, 1637, 1508, 1377, 1347, 1293, 1273, 1255, 1227, 1193, 1117, 1064, 1015, 766, 747cm⁻¹

5 ¹H NMR (300 MHz in DMSO-d₆): δ 14.81 (brs,1H), 8.20(s,1H), 7.77-7.83 (m,4H), 7.64-7.67 (d,2H), 7.32-7.42 (m,4H), 7.01-7.07 (t, 3H), 6.87 (d,2H) 3.56(t, 4H), 2.82 (m, 4H), 2.37 (s,3H), 2.22 (s,3H), 2.18 (s,3H).

[0153] EXAMPLE 6: Preparation of S'-amino-Z'-methoxybiphenyl-S-carboxylic acid

10 A mixture of 3-bromo-2-methoxyaniline (3g), 3-carboxyphenylboronic acid (2.94g), bis(triphenylphosphine)palladium(II)chloride [PdCl₂(PPh₃)₂] (0.30g) and potassium carbonate (5.2g) in ethanol (100mL) and water (20mL) was heated to about reflux temperature and stirred for about 28 hours. The hot mixture was filtered to remove the catalyst. The clear filtrate was concentrated under vacuum. To the residue, water
15 (50mL) and methanol (50mL) was added, acidified to pH 3-5 using hydrochloric acid and stirred for about 30 minutes at about room temperature. The slurry was filtered, washed with n-hexane (50mL) and dried in an oven at about 55°C to about 60°C for about 12 hours. Yield: 1.8g

¹H NMR (300 MHz in DMSO-d₆): δ 8.08 (s,1H), 7.89-7.92 (d,1H), 7.73-7.75 (d,1H),
20 7.51-7.56 (t,1H), 6.87-6.92 (t,1H), 6.72-6.74 (d,1H), 6.50-6.53 (d,1H), 3.27 (s,3H)

[0154] EXAMPLE 7: Preparation of 3'-{-2-[1-(ethoxycarbonyl)-2-oxopropylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid

To a solution of 3'-amino-2'-methoxybiphenyl-3-carboxylic acid (5g) prepared as in Example 6, in hydrochloric acid (8.5mL) and methanol (90mL) was added a solution
25 of sodium nitrite (1.41g in 400mL of water) at about 0°C to about 5°C under stirring. The reaction mixture was stirred for about 15 minutes at about 5°C. Then ethyl acetoacetate (2.7g) was added and the reaction mixture was stirred for about 15 minutes at about 0°C to about 5°C. Solid sodium bicarbonate (15g) and ethanol (30mL) was added and the reaction mixture was allowed to warm to about room
30 temperature and stirred for about 2 hours. The mixture was filtered, washed with water (50mL) and dried under vacuum at about 35°C to about 40°C for about 6 hours. Yield: 4.2g; Mass: m/z 383.15 [M-1]

¹H NMR (300 MHz in DMSO-d₆): δ 14.74 (s, 1H), 12.56 (s, 1H), 8.14 (s, 1H), 7.97-8.00 (d, 1H), 7.83-7.85 (d, 1H), 7.60-7.71 (m, 2H), 7.26-7.34 (m, 1H), 7.19-7.23 (m, 1H), 4.10-4.34 (m, 2H), 3.40 (s, 3H), 2.52 (s, 3H), 1.32 (t, 3H)

[0155] EXAMPLE 8: Preparation of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid

A solution of 3'-{-2-[1-(ethoxycarbonyl)-2-oxopropylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid (3.5g) prepared as in Example 7, and 3,4-dimethylphenylhydrazine hydrochloride (1.72g) and sodium acetate (0.9g) in glacial acetic acid (120mL) was stirred and heated to about reflux temperature for about 4 hours. The mixture was cooled to about room temperature and stirred for about 1 hour. It was filtered, washed with water (25mL) and dried in an oven at about 55°C to about 60°C for about 12 hours to an orange solid.

Yield: 2.5g; Mass: m/z 455.61 [M-1]

¹H NMR (300 MHz in DMSO-d₆): δ 13.75 (brs, 1H), 8.17 (s, 1H), 7.99-8.01 (d, 1H), 7.84-7.87 (d, 1H), 7.64-7.79 (d, 1H), 7.62-7.67 (t, 1H), 7.34-7.40 (t, 1H), 7.27-7.29 (d, 1H), 7.19-7.22 (d, 1H), 3.34 (s, 3H), 2.32 (s, 3H), 2.26 (s, 3H), 2.22 (s, 3H)

[0156] EXAMPLE 9: Preparation of 3'-{-2-[1-(ethoxycarbonyl)-2-oxopropylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid

A mixture of ethyl-2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate (4g) prepared as in Example 1, 3-carboxyphenylboronic acid (2.30g), bis(triphenylphosphine)palladium(II)chloride [PdCl₂(PPh₃)₂] (0.30g), potassium carbonate (4.023 g) in ethanol (100mL) and water (30mL) was heated to about reflux temperature and stirred for about 20 hours. The hot mixture was filtered to remove the catalyst. The clear filtrate was concentrated under vacuum. To the residue, water (100mL) was added, acidified to pH 3-5 using hydrochloric acid and stirred for about 30 minutes at about room temperature. The slurry was filtered, washed with water (25ml) and dried in an oven at about 55°C to about 60°C for about 12 hours.

Yield: 1.9g; Mass: m/z 383.15 [M-1]

¹H NMR (300 MHz in CDCl₃): δ 13.7 (brs, 1H), 8.36 (s, 1H), 8.13-8.16 (d, 1H), 7.88-7.91 (d, 1H), 7.68-7.71 (d, 1H), 7.55-7.60 (m, 1H), 7.24-7.29 (m, 1H), 7.15-7.18 (d, 1H), 4.36-4.44 (m, 2H), 3.49 (s, 3H), 2.62 (s, 3H), 1.41 (t, 3H)

[0157] EXAMPLE 10: Preparation of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-hydroxyphenyl)hydrazono-5-pyrazolone

A solution of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone (5g) prepared as in Example 2, in 48% aqueous hydrobromic acid (200mL) and glacial acetic acid (50mL) was stirred and heated under reflux for about 80 hours. The reaction mixture was concentrated under vacuum to obtain a thick residue. Water (40mL) was added and basified to pH 7-8 using saturated solution of sodium bicarbonate (20mL). The mixture was filtered to give an orange solid, which was washed with water (25mL) and dried in an oven at about 55°C to about 60°C for about 12 hours. Yield: 3.5g

¹H NMR (300 MHz in DMSO -d₆): δ 13.45 (brs,1H), 10.98 (brs,1H), 7.69-7.71 (d,1H), 7.59-7.62 (d,2H), 7.12-7.21 (m,2H), 6.92-6.95 (d,1H), 2.31 (s,3H), 2.26 (s,3H), 2.22 (s,3H)

[0158] EXAMPLE 11: Preparation of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid

A mixture of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-hydroxyphenyl)hydrazono-5-pyrazolone (2g) prepared as in Example 10, 3-carboxyphenylboronic acid (1g), bis(triphenylphosphine)palladium(II)chloride [PdCl₂(PPh₃)₂] (0.2g), potassium hydroxide (1.50g) in ethanol (50mL) and water (20mL) was heated to about reflux temperature and stirred for about 20 hours. The hot mixture was filtered to remove the catalyst. The clear filtrate was concentrated under vacuum. To the residue, water (100mL) was added, acidified to pH 3-5 using hydrochloric acid and stirred for about 30 minutes at about room temperature. The slurry was filtered to give an orange solid, which was washed with water (20mL) and dried in an oven at about 55°C to about 60°C for about 12 hours.

Yield: 1.8g; Melting point: 231.6-232.9°C; Mass: m/z 443.16 [M+1]

TR: 3292, 1707, 1541, 1503, 1223, 1118, 1000cm⁻¹

¹H NMR (300 MHz in DMSO -d₆): δ 13.75 (brs,1H), 13.09 (brs,1H), 9.69 (s,1H), 8.13 (s,1H), 7.94-7.97 (d,1H), 7.79-7.82 (d,1H), 7.62-7.71 (m,3H), 7.16-7.21 (m,3H), 2.32 (s,3H), 2.26 (s,3H), 2.22 (s,3H)

[0159] EXAMPLE 12: Preparation of trombopag olamine

To a mixture of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid (2g) in water (40mL), ethanolamine (2.5g) was added and heated to about reflux temperature for about 2 hours. The mixture was cooled to about 0°C to about 10°C and stirred for about 1

hour. The reaction mixture was filtered, washed with water (10mL) and dried in an oven at about 55°C to about 60°C for about 12 hours to get purple solid.

Yield: 1.8g; Melting point: 237.3-237.6°C

IR: 3421, 1637, 1508, 1377, 1347, 1293, 1273, 1255, 1227, 1193, 1117, 1064, 1015, 766, 747cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 14.81 (brs, 1H), 8.20 (s, 1H), 7.77-7.83 (m, 4H), 7.64-7.67 (d, 2H), 7.32-7.42 (m, 4H), 7.01-7.07 (t, 3H), 6.87 (d, 2H), 3.56 (t, 4H), 2.82 (m, 4H), 2.37 (s, 3H), 2.22 (s, 3H), 2.18 (s, 3H)

[0160] EXAMPLE 13: Preparation of eltrombopag olamine

A mixture of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-hydroxybiphenyl-3-carboxylic acid (2g) and ethanolamine (10mL) was heated at about 55°C to about 60°C for 2 hours. The mixture was cooled to about room temperature and stirred for about 1 hour. It was filtered, washed with ethanol (10mL) and dried in an oven at about 55°C to about 60°C for about 12 hours to get purple solid. Yield: 1.7 g; Melting point: 237.3-237.6°C

IR: 3421, 1637, 1508, 1377, 1347, 1293, 1273, 1255, 1227, 1193, 1117, 1064, 1015, 766, 747cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 14.81 (brs, 1H), 8.20 (s, 1H), 7.77-7.83 (m, 4H), 7.64-7.67 (d, 2H), 7.32-7.42 (m, 4H), 7.01-7.07 (t, 3H), 6.87 (d, 2H), 3.56 (t, 4H), 2.82 (m, 4H), 2.37 (s, 3H), 2.22 (s, 3H), 2.18 (s, 3H)

[0161] EXAMPLE 14: Preparation of eltrombopag olamine

a) Preparation of eltrombopag-aluminium complex

A mixture of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid (10g) and aluminium chloride (20.4g) in toluene (50mL) was stirred for about 40 hours at about room temperature under nitrogen atmosphere. The reaction mass was poured into dilute hydrochloric acid (100mL) at about 10°C to about 15°C. The mixture was filtered to give a solid, which was washed with toluene (20mL) and dried under vacuum.

IR : 3418, 2924, 1588, 1490, 1407, 1349, 1288, 1172, 1123, 1072, 744, 603 cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 13 (brs, 1H), 8.57 (s, 1H), 8.0-8.02 (d, 1H), 7.88-7.91 (d, 1H), 7.54-7.63 (m, 4H), 7.41-7.44 (d, 1H), 7.23-7.26 (d, 1H), 6.80-6.85 (t, 1H), 2.33 (s, 3H), 2.28 (s, 3H), 2.25 (s, 3H)

Aluminum content: 3.52% by W/W

b) Preparation of eltrombopag

The above solid cake was added in tetrahydrofuran (100mL), acetic acid (50mL) and water (30mL). The reaction mixture was heated to about reflux temperature and stirred for about 6 hours. The reaction mixture was cooled to about room temperature and 20% sodium chloride solution (50mL) was added to it. The reaction mass was stirred and the two layers were separated. The organic layer contained the title compound.

c) Preparation of eltrombopag ammonium salt

25% aqueous ammonia (150mL) was added to the above organic layer and stirred for about 2 hours at about room temperature. The mixture was filtered to give a solid which was washed with water (20mL) and dried under vacuum.

Yield: 7g; HPLC Purity : 97.91%

IR: 3368, 3155, 1648, 1546, 1501, 1399, 1382, 1343, 1268, 1226, 1193, 1156, 1053, 1002, 790, 767. 741cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 14.81 (brs,1H), 8.17 (s,1H), 7.80-7.82(m,3H), 7.65-7.68 (d,1H), 7.34-7.42 (m,2H), 7.01-7.07 (m,2H), 6.87 (t,1H), 2.37 (s,3H), 2.22 (s,3H), 2.18 (s,3H)

Ammonia content: 4.7% by W/W

d) Preparation of eltrombopag olamine

To the above solid cake, ethanolamine (50mL) was added and stirred at about room temperature for about 2 hours. Ethanol (200mL) was then added to the reaction mixture. The reaction mixture was stirred and filtered to give eltrombopag olamine as a solid, which was washed with ethanol (20mL) and dried in vacuum oven at about 55°C to about 60°C for about 12 hours.- Yield: 8g

IR: 3421, 1637, 1508, 1377, 1347, 1293, 1273, 1255, 1227, 1193, 1117, 1064, 1015, 766, 747cm⁻¹

¹H NMR (300 MHz in DMSO-d₆): δ 14.81 (brs,1H), 8.20(s,1H), 7.77-7.83 (m,3H), 7.64-7.67 (d,1H), 7.32-7.42 (m,2H), 7.01-7.07 (t, 2H), 6.87 (m,1H) 3.56(t, 4H), 2.82 (m, 4H), 2.37 (s,3H), 2.22 (s,3H), 2.18 (s,3H)

[0162] EXAMPLE 15: Preparation of 3-bromo-2-methoxyaniline hydrochloride

To a stirred solution of 3-bromo-2-methoxyaniline (10g) in ethyl acetate (100mL) was added concentrated hydrochloric acid (7mL) at about room temperature. The reaction mixture was stirred for about 2 hours and filtered to give a solid, which was washed with ethyl acetate and dried at about 50°C to about 55°C. Yield: 10g

[0163] EXAMPLE 16: Preparation of Ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate

To a solution of 3-bromo-2-methoxyaniline hydrochloride (10g) prepared as in Example 15, in hydrochloric acid (13.5mL of concentrated hydrochloric acid in 136.5mL of water) was added a solution of sodium nitrite (3.47g in 10mL of water) at about -5°C to about 5°C under stirring. The reaction mixture was stirred for about 1 hour at about -5°C to about 5°C. Then ethylacetoacetate (6.55g) was added to the reaction mixture and stirred for about 3 hours at about -5°C to about 5°C. Sodium bicarbonate and methanol was then added to the reaction mixture. The reaction mixture was stirred at about -5°C to about 5°C for about 1 hour. The mixture was filtered to give a solid, which was washed with water and dried at about 50°C to about 55°C. Yield: 13.3g

[0164] EXAMPLE 17: Preparation of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone

A solution of ethyl-2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate (10g) prepared as in Example 15, and 3,4-dimethylphenylhydrazine hydrochloride (6.05g) and sodium acetate (2.63g) in acetic acid (150mL) was stirred and heated to about 80°C to about 90°C for about 3 hours. The reaction mixture was cooled to about room temperature and stirred for about 1 hour. The reaction mixture was filtered to give a solid, which was washed with acetic acid. The solid obtained was stirred in water for about 30 minutes at about room temperature, which was then filtered, washed with water and dried at about 55°C to about 60°C. Yield: 11g

[0165] EXAMPLE 18: Preparation of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid

A mixture of 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone (10g) prepared as in Example 16, 3-carboxyphenylboronic acid (5.2g), bis(triphenylphosphine)palladium(II)chloride [PdCl₂(PPh₃)₂] (0.84g), potassium hydroxide (6.72g) in ethanol (200mL) and water (15mL) was heated to about reflux temperature under nitrogen atmosphere and stirred for about 8 hours. The reaction mixture was cooled to about 30°C to about 35°C and stirred for about 4 hours at about the same temperature. The reaction mixture was treated with Norit™ charcoal and filtered over celite bed. The filtrate was treated with EDTA and Scavenger Type 2S. The mixture was stirred at about 30°C to about 40°C for about 20 hours and filtered over celite bed. To the filtrate, hydrochloric acid was added to adjust the pH to about 3 to about 5. The mixture was stirred for about 2

hours at about 80°C to about 85°C. The mixture was cooled to about 40°C to about 45°C and filtered. The wet cake was treated with potassium hydroxide in methanol and stirred to give a clear solution. The solution was treated with 50% tributylphosphene in ethyl acetate and stirred for about 20 hours at about room temperature. The pH of the solution was adjusted to about 2 to about 4 using hydrochloric acid and the solid obtained was filtered and purified using tetrahydrofuran. Yield: 9.3g

[0166] EXAMPLE 19: Preparation of eltrombopag olamine

a) Preparation of eltrombopag-aluminium complex

10 A mixture of 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid (10g) and aluminium chloride (29.3g) in toluene (50mL) was stirred for about 30 hours at about room temperature under nitrogen atmosphere. The reaction mass was poured into dilute hydrochloric acid (200mL) at about 25°C to about 45°C. The mixture was filtered to give a solid, which was washed with toluene (20mL) and dried under vacuum.

b) Preparation of eltrombopag

The above solid cake was added in tetrahydrofuran (200mL), acetic acid (50mL) and water (30mL). EDTA was added to the reaction mixture and the reaction mixture was heated to about 60°C to about 70°C and stirred for about 12 hours. The reaction mixture was cooled to about 30°C to about 40°C and 15% sodium chloride solution was added to it. The reaction mass was stirred and the two layers were separated. The organic layer was treated with Scavenger Type 2S and 50% tributylphosphene in ethyl acetate, stirred for about 15 hours at about 30°C to about 40°C and filtered over celite bed. The filtrate contained the title compound.

25 c) Preparation of eltrombopag ammonium salt


25% aqueous ammonia (10mL) was added to the above filtrate at about 30°C to about 40°C. The mixture was concentrated under vacuum at about 45°C to about 50°C. The residue obtained was degassed for about 1 hour.

d) Preparation of eltrombopag olamine


30 To the residue, ethanolamine (50mL) was added and stirred at about 20°C to about 30°C for about 2 hours. Methanol was then added to the reaction mixture. The reaction mixture was stirred and filtered to give eltrombopag olamine as solid which was washed with methanol and dried at about 55°C to about 60°C. Yield: 8g

CLAIMS:


1. A process for the preparation of substituted 3'-hydrazino-biphenyl-3-carboxylic acid compounds of formula I and salts thereof,



- 5 wherein R represents hydrogen, linear or branched C_{1-6} alkyl, C_{3-8} cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl; the process comprising:
- a) reacting a compound of formula III,




- 10 wherein R is as defined above, X is selected from the group consisting of Cl, Br, I, with a compound of formula IV,



wherein R' represents boronic acid, boronic acid ester or halogen in the presence of a metal catalyst; and


- 15 (b) optionally, deprotecting the compound of formula I.
2. The process of claim 1, wherein R is C_{1-6} alkyl, X is Br and R' is boronic acid.

3. The process of claim 1, wherein R is C₁₋₆ alkyl, X is Br and R' is halogen selected from the group consisting of Cl, Br, I.
4. The process of claim 1, wherein the metal catalyst is selected from the group consisting of Pd(PPh₃)₄, PdCl₂(PPh₃)₂, PdCl₂(dppf), Pd(OAc)₂, NiCl₂(PPh₃)₂, PdCl₂(dppb).
5. A compound of formula III,




wherein X and R are as defined above.

6. The compound of claim 5, wherein X is Br and R is methyl.
7. The process of claim 1, wherein the compound of formula III is obtained by reacting a compound of formula V,




wherein X and R are as defined above and R'' represents hydrogen, C₁₋₆ alkyl, with 3,4-dimethylphenylhydrazine or salt thereof to give the compound of formula III.

8. The process of claim 7, wherein X is Br, R is methyl and R'' is ethyl.
9. A compound of formula V,




wherein X, R and R'' are as defined above.


10. The compound of claim 9, wherein X is Br, R is methyl and R" is ethyl.
11. The process of claim 7, wherein the compound of formula V is obtained by reacting a compound of formula VI or its salt thereof,




- 5 wherein X and R are as defined above,
with alkyl acetoacetate or acetoacetic acid in the presence of alkali or alkaline earth metal nitrite and an inorganic acid in a solvent system to yield the compound of formula V.
12. The process of claim 11, wherein the inorganic acid is selected from the group
10 consisting of hydrochloric acid, nitric acid, sulfuric acid.
13. The process of claim 11, wherein the solvent system is selected from methanol, ethanol, water and mixtures thereof.
14. A process for the preparation of eltrombopag, compound of formula Ia,




15. and pharmaceutically acceptable salts thereof, the process comprising:
(a) reacting 3-bromo-2-methoxyaniline with ethylacetoacetate in the presence of alkali or alkaline earth metal nitrite and inorganic acid to give ethyl 2-[(3-bromo-2-methoxyphenyl)hydrazono]-3-oxobutanoate, compound of formula Va;



(b) reacting the compound of formula Va with 3,4-dimethylphenylhydrazine or salt thereof to give 1-(3,4-dimethylphenyl)-3-methyl-4-(3-bromo-2-methoxyphenyl)hydrazono-5-pyrazolone, compound of formula IIa;



5 (c) reacting the compound of formula IIa with 3-carboxyphenylboronic acid in the presence of a metal catalyst to give 3'-{N'-[1-(3,4-dimethylphenyl)-3-methyl-5-oxo-1,5-dihydro-pyrazol-4-ylidene]hydrazino}-2'-methoxybiphenyl-3-carboxylic acid, compound of formula Ib;




10 (d) deprotecting the compound of formula Ib using Lewis acid to give eltrombopag; and optionally, converting to its pharmaceutically acceptable salt.

15 15. The process of claim 14, wherein the Lewis acid used in (d) is selected from the group consisting of aluminium chloride, aluminium bromide, aluminium iodide, stannous chloride, stannous bromide, titanium chloride, boron trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium chloride, beryllium bromide, zinc chloride, zinc bromide, trimethylsilylchloride, trimethylsilylbromide, trimethylsilyliodide.


16. The process of claim 15, wherein the Lewis acid used is aluminium chloride.

17. A process for the preparation of eltrombopag, a compound of formula Ia,



Ia

the process comprising subjecting the compound of formula I to a deprotection reaction.



I

5 wherein R represents linear or branched C_{1-6} alkyl, C_{3-8} cycloalkyl, optionally substituted benzyl, linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, methoxybenzyl, trialkylsilyl, acyl, trityl.

18. The process of claim I and claim 17, wherein the deprotection reaction process includes any of the following:

- 10 (a) where R is C_{1-6} alkyl, C_{3-8} cycloalkyl, the deprotection of the compound of formula I is performed using protic acid such as hydroiodic acid, hydrobromic acid, hydrobromic acid/acetic acid, methanesulfonic acid, trifluoroacetic acid; Lewis acid selected from the group consisting of aluminium chloride, aluminium bromide, aluminium iodide, stannous chloride, stannous bromide, titanium chloride, boron trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium
- 15 trifluoride, boron tribromide, boron trifluoride-dimethylsulfide complex, beryllium chloride, beryllium bromide, zinc chloride, zinc bromide, trimethylsilylchloride, trimethylsilylbromide, trimethylsilyliodide, lithium iodide, lithium iodide in refluxing 2,4,6-collidine, pyridine hydrochloride; sulphur compounds such as sodium

ethylmercaptide, sodium trimethylsilylanethiolate; alkali organomides such as sodium bis(trimethylsilyl)amide and lithium diisopropylamide; or

(b) where R is optionally substituted benzyl, methoxybenzyl, the deprotection of the compound of formula I is performed via-hydrogenation reaction using hydrogen in the presence of a metal catalyst; or

(c) where R is linear or branched alkylalkoxy, tetrahydrofuranyl, tetrahydropyranyl, trityl, the deprotection of the compound of formula I is performed using an inorganic acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; organic acid such as acetic acid; or

(d) where R is trialkylsilyl, the deprotection of the compound of formula I is performed using acids such as acetic acid or fluorides such as tetrabutylammonium fluoride; or

(e) where R is acyl. the deprotection of the compound of formula I is performed using an inorganic acid selected from the group consisting of hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid; inorganic base selected from the group consisting of sodium hydroxide, potassium hydroxide, sodium carbonate, potassium carbonate:

19. The use of compound of formula IIa, Va, or Ib in the preparation of eltrombopag or salt thereof.

20. Eltrombopag ammonium salt.

21. A process for the preparation of eltrombopag ammonium salt comprising reacting eltrombopag with a source of ammonia.

22. The process of claim 21, wherein the source of ammonia is selected from the group consisting of ammonia water or ammonium carbonate or ammonia gas or liquid ammonia.

23. A process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with excess of ethanolamine without using any additional reaction solvent.

24. A process for the preparation of eltrombopag olamine comprising reacting eltrombopag or a salt thereof with ethanolamine in an aqueous medium.

25. A process for the preparation of eltrombopag olamine comprising reacting eltrombopag ammonium salt with ethanolamine.

26. The use of eltrombopag ammonium salt in the preparation of eltrombopag olamine.

Glenmark Generics Limited

Fig 1/4




Figure 1

Glenmark Generics Limited

Fig 2/4




Figure 2

Glenmark Generics Limited

Fig 3/4




Figure 3

Glenmark Generics Limited

Fig 4/4




Figure 4