



(86) Date de dépôt PCT/PCT Filing Date: 2004/06/26
(87) Date publication PCT/PCT Publication Date: 2004/12/29
(45) Date de délivrance/Issue Date: 2013/08/06
(85) Entrée phase nationale/National Entry: 2005/12/20
(86) N° demande PCT/PCT Application No.: AU 2004/000866
(87) N° publication PCT/PCT Publication No.: 2004/112687
(30) Priorités/Priorities: 2003/06/26 (AU2003903251);
2003/07/25 (AU2003903850);
2003/08/29 (AU2003904692);
2004/05/31 (AU2004902902)


(51) Cl.Int./Int.Cl. *C07C 279/22* (2006.01),
A61K 31/16 (2006.01), *A61K 31/165* (2006.01),
A61P 31/12 (2006.01), *C07C 279/18* (2006.01),
C07D 241/32 (2006.01), *C07D 307/54* (2006.01),
C07D 403/04 (2006.01)

(72) Inventeurs/Inventors:
GAGE, PETER WILLIAM, AU;
EWART, GARY DINNEEN, AU;
WILSON, LAUREN ELIZABETH, AU;
BEST, WAYNE, AU;
PREMKUMAR, ANITA, AU

(73) Propriétaire/Owner:
BIOTRON LIMITED, AU

(74) Agent: BLAKE, CASSELS & GRAYDON LLP

(54) Titre : COMPOSITIONS ET METHODES ANTIVIRALES
(54) Title: ANTIVIRAL COMPOUNDS AND METHODS



(57) Abrégé/Abstract:

The invention relates to compounds having antiviral activity and methods utilising the compounds to treat viral infections.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 December 2004 (29.12.2004)

PCT

(10) International Publication Number
WO 2004/112687 A2

(51) International Patent Classification⁷:

A61K

(74) Agent: SHELSTON IP; 60 Margaret Street, Sydney NSW 2000 (AU).

(21) International Application Number:

PCT/AU2004/000866

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 26 June 2004 (26.06.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

2003903251	26 June 2003 (26.06.2003)	AU
2003903850	25 July 2003 (25.07.2003)	AU
2003904692	29 August 2003 (29.08.2003)	AU
2004902902	31 May 2004 (31.05.2004)	AU

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): **BIOTRON LIMITED** [AU/AU]; LPO Box A315, Australian National University, Australian Capital Territory, 2601 (AU).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **GAGE, Peter, William** [AU/AU]; RMB 22, Powell Drive, Via Queanbeyan, NSW 2620 (AU). **EWART, Gary, Dinneen** [AU/AU]; 189 Phillip Avenue, Hackett, ACT 2602 (AU). **WILSON, Lauren, Elizabeth** [AU/AU]; 16 Ebden Street, Ainslie, ACT 2602 (AU). **BEST, Wayne** [AU/AU]; 4 Mimy Court, Gosnells, Western Australia 6110 (AU). **PREMKUMAR, Anita** [AU/AU]; 116 Kelleway Avenue, Nicholls, ACT 2915 (AU).

Declaration under Rule 4.17:

— of inventorship (*Rule 4.17(iv)*) for US only

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIVIRAL COMPOUNDS AND METHODS

(57) Abstract: The invention relates to compounds having antiviral activity and methods utilising the compounds to treat viral infections.



WO 2004/112687 A2

ANTIVIRAL COMPOUNDS AND METHODS

FIELD OF INVENTION

The present invention relates to methods for retarding, reducing or otherwise
5 inhibiting viral growth and/or functional activity. The invention also relates to
compounds and compositions suitable for use in the methods.

BACKGROUND OF THE INVENTION

Currently, there is a great need for the development of new treatments that are
effective against viral infections, particularly against viral infections which are
10 associated with high morbidity and mortality, and which impact on sizable
populations. Treatments currently available are inadequate or ineffective in large
proportions of infected patients.

For example, in ameliorating AIDS symptoms and prolonging life expectancy, a
measure of success has been achieved with drugs targeting the viral reverse
15 transcriptase and protease enzymes (Miller and Sarver, 1997; Mitsuya, 1992; Moore,
1997; and Thomas and Brady, 1997). However, no single treatment method is
completely effective against HIV infection. (Barry et al, 1998; Deeks, 1998; Miles,
1997; Miles, 1998; Moyle et al, 1998; Rachlis and Zarowny, 1998; Veil et al, 1997;
Volberding and Deeks, 1998; and Volberdin, 1998).

20 PCT Publication WO 2000/021538 describes the use of compounds 5-(N, N-
hexamethylene)-amiloride and 5-(N,N-dimethyl)-amiloride in the treatment of HIV
infection.

Another virus considered to be a significant human pathogen is the Hepatitis C
virus (HCV). This is a significant human pathogen in terms of both cost to human
25 health and associated economic costs. HCV causes chronic hepatitis and cirrhosis
and is the leading indicator for liver replacement surgery. In 2002 the Centre for
Disease Control and Prevention estimated that more than 4 million people were
infected in the USA alone and that approximately 8,000 to 10, 000 die as a result of
chronic HCV infection yearly. There is no known cure or vaccine. More effective
30 pharmacological agents are urgently required.

-2-

A further well-known family of pathogenic viruses are the Coronaviruses. Coronaviruses (Order *Nidovirales*, family *Coronaviridae*, Genus *Coronavirus*) are enveloped positive-stranded RNA viruses that bud from the endoplasmic reticulum-Golgi intermediate compartment or the *cis*-Golgi network (Fischer, Stegen et al. 1998; Maeda, Maeda et al. 1999; Corse and Machamer 2000; Maeda, Repass et al. 2001; Kuo and Masters 2003).

Coronaviruses infect humans and animals and it is thought that there could be a coronavirus that infects every animal. The two human coronaviruses, 229E and OC43, are known to be the major causes of the common cold and can occasionally cause pneumonia in older adults, neonates, or immunocompromised patients (Peiris, Lai et al. 2003). Animal coronaviruses can cause respiratory, gastrointestinal, neurological, or hepatic diseases in their host (Peiris, Lai et al. 2003). Several animal coronavirus are significant veterinary pathogens (Rota, Oberste et al. 2003).

Severe acute respiratory syndrome (SARS) is caused by a newly identified virus. SARS is a respiratory illness that has recently been reported in Asia, North America, and Europe (Peiris, Lai et al. 2003). The causative agent of SARS was identified as a coronavirus. (Drosten, Gunther et al. 2003; Ksiazek, Erdman et al. 2003; Peiris, Lai et al. 2003). The World Health Organization reports that the cumulative number of reported probable cases of SARS from 1 November 2002 to the 11th July 2003 is 8,437 with 813 deaths, nearly a 10% death rate. It is believed that SARS will not be eradicated, but will cause seasonal epidemics like the cold or influenza viruses (Vogel 2003).

To improve the prospect of treating and preventing viral infections, there is an on-going need to identify molecules capable of inhibiting various aspects of the viral life cycle.

It is an object of the present invention to overcome or ameliorate at least one of the disadvantages of the prior art, or to provide a useful alternative.

Any discussion of the prior art throughout the specification should in no way be considered as an admission that such prior art is widely known or forms part of common general knowledge in the field.

-3-

SUMMARY OF THE INVENTION

The inventors have surprisingly found that certain compounds that fall under the classification of substituted acylguanidines have antiviral activity against viruses from a range of different virus families. Without intending to be bound by any particular theory or mechanism of action, and despite current dogma, it appears possible that viral replication can be retarded by inhibiting or otherwise down-regulating the activity of ion channels expressed in the host cell. Thus, the negative impact of the compounds of the present invention on viral replication may be mediated by the inhibition or otherwise down-regulation of a membrane ion channel relied upon by the virus for replication. This membrane ion channel may be a viral membrane ion channel (exogenous to the host cell) or a host cell ion channel induced as a result of viral infection (endogenous to the host cell).


As an example, the compounds of the present invention may inhibit Vpu or p7 function and thereby inhibit the continuation of the respective HIV or HCV life cycle.

The SARS virus encodes an E protein which is shown for the first time, by the present inventors, to act as an ion channel. As similar E proteins are present in other coronaviruses, the compounds, compositions and methods of the present invention would have utility in the inhibition and/or treatment of infections by other coronaviruses.

The present invention is concerned with novel antiviral compounds that fall under the classification of substituted acylguanidines. It does not include in its scope the use of compounds 5-(N,N-hexamethylene)amiloride and 5-(N,N-dimethyl)-amiloride for retarding, reducing or otherwise inhibiting viral growth and/or functional activity of HIV.

Accordingly, a first aspect of the present invention provides an acylguanidine with antiviral activity.


According to a second aspect, the present invention provides an antiviral compound of Formula I




I

-4-

wherein R_1 - R_4 are independently aromatic groups, heteroaromatic groups, alkylaromatic groups, alkylheteroaromatic groups, alkenylaromatic groups, alkenylheteroaromatic groups, cycloalkylaromatic groups, cycloalkylheteroaromatic groups, aryloxyalkyl groups, heteroaryloxyalkyl groups, said groups are mono or polycyclic, and are optionally substituted with one or more substituents independently selected from hydrogen, hydroxy, nitro, halo, amino, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{3-6} cycloalkyl, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkyloxy, phenyl, C_{1-6} alkenyl, C_{3-6} cycloalkenyl, C_{1-6} alkeneoxy, benzo,







According to a third aspect, the present invention provides an antiviral compound of Formula I








or pharmaceutically acceptable salts thereof, wherein,




$R_1 =$


























 or  BOD^{IPY-FL}

and wherein


-6-

X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkenyl, C₃₋₆cycloalkenyl, C₁₋₆alkeneoxy, or benzo;

5 R_a, R_b, R_c, R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




10 R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;
R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



15

Preferably, the compounds of the invention include the following:


5-(N,N-hexamethylene)amiloride comprising the structure




20

-7-


5-(N,N-Dimethyl)amiloride hydrochloride comprising the structure



5-(N-methyl-N-isobutyl)amiloride comprising the structure




5-(N-ethyl-N-isopropyl)amiloride (herein referred to as EIPA), comprising the structure




-8-


N-(3,5-Diamino-6-chloro-pyrazine-2-carbonyl)-N'-phenyl-guanidine, comprising the structure




5 N-Benzyl -N'-(3,5-diamino-6-chloro-pyrazine-2-carbonyl)-guanidine, comprising the structure



3-methoxy amiloride comprising the structure comprising the structure




3-methoxy-5-(N,N-Hexamethylene)-amiloride comprising the structure




10

-9-


3-hydroxy-5-hexamethyleneimino-amiloride comprising the structure



Hexamethyleneimino-6-phenyl-2-pyrazinecarboxamide comprising the structure




N-amidino-3,5-diamino-6-phenyl-2-pyrazinecarboxamide comprising the structure




-10-


5-(N,N-hexamethylene)amiloride comprising the structure



N-amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide comprising the
5 structure




3'4 DichloroBenzamil comprising the structure




-11-


2'4 DichloroBenzamil HCl comprising the structure




5-(N-methyl-N-guanidinocarbonyl-methyl)amiloride comprising the structure



5 5-(N,N-Diethyl)amiloride hydrochloride comprising the structure




5-(N,N-Dimethyl)amiloride hydrochloride comprising the structure




-12-


5-tert-butylamino-amiloride comprising the structure




6-Iodoamiloride comprising the structure



Bodipy-FL Amiloride comprising the structure




5-(4-fluorophenyl)amiloride comprising the structure




-13-

1-naphthoylguanidine comprising the structure




2-naphthoylguanidine comprising the structure




5

N-(2-naphthoyl)-N'-phenylguanidine comprising the structure




N,N'-bis(2-naphthoyl)guanidine comprising the structure




10

-14-


N,N'-bis(1-naphthoyl)guanidine comprising the structure




N,N'-bis(2-naphthoyl)-N''-phenylguanidine comprising the structure



5 6-methoxy-2-naphthoylguanidine comprising the structure




N-Cinnamoyl-N',N'-dimethylguanidine comprising the structure




-15-

3-quinolinoylguanidine comprising the structure




cinnamoylguanidine comprising the structure




5

4-phenylbenzoylguanidine comprising the structure




N-(cinnamoyl)-N'phenylguanidine comprising the structure




10

-16-


(3-phenylpropanoyl)guanidine comprising the structure




N,N'-bis-(cinnamoyl)-N''-phenylguanidine comprising the structure



5 N-(3-phenylpropanoyl)-N'-phenylguanidine comprising the structure




N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine comprising the structure




-17-

trans-3-furanacryoylguanidine comprising the structure




N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine comprising the structure




5

(4-Phenoxybenzoyl)guanidine comprising the structure




10 N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide comprising the structure




15

-18-


6-bromo-2-naphthoylguanidine comprising the structure




1-bromo-2-naphthoylguanidine comprising the structure




2-(2-naphthyl)acetoyleguanidine comprising the structure



N''-Cinnamoyl-N,N'-diphenylguanidine comprising the structure




(Phenylacetyl)guanidine comprising the structure




-19-

N,N'-Bis(3-phenylpropanoyl)guanidine comprising the structure




Benzoylguanidine comprising the structure




5

(4-Chlorophenoxy-acetyl]guanidine comprising the structure




10

N-Benzoyl-N'-cinnamoylguanidine comprising the structure




[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine comprising the

15 structure




-20-

(4-Chlorocinnamoyl)guanidine comprising the structure




(4-Bromocinnamoyl)guanidine comprising the structure




5

(4-Methoxycinnamoyl)guanidine comprising the structure




(5-Phenyl-penta-2,4-dienoyl)guanidine comprising the structure




10

(3-Bromocinnamoyl)guanidine comprising the structure




15 (3-Methoxycinnamoyl)guanidine comprising the structure




-21-

(3-Chlorocinnamoyl)guanidine comprising the structure




(2-Chlorocinnamoyl)guanidine comprising the structure




5


(2-Bromocinnamoyl)guanidine comprising the structure




(2-Methoxycinnamoyl)guanidine comprising the structure



10 (trans-2-Phenylcyclopropanecarbonyl)guanidine comprising the structure




[3-(3-Pyridyl)acryloyl]guanidine comprising the structure




15

-22-

(4-Hydroxycinnamoyl)guanidine comprising the structure




(Quinoline-2-carbonyl)guanidine comprising the structure




5


(4-Nitrocinnamoyl)guanidine comprising the structure



10 (3-Nitrocinnamoyl)guanidine comprising the structure




(2-Nitrocinnamoyl)guanidine comprising the structure




15

-23-

(α -Methylcinnamoyl)guanidine comprising the structure




trans-3-(1-naphthyl)acryloylguanidine comprising the structure




5

4-phenylcinnamoylguanidine comprising the structure




3-(trifluoromethyl)cinnamoylguanidine comprising the structure




10

3-methylcinnamoylguanidine comprising the structure




-24-

4-(trifluoromethyl)cinnamoylguanidine comprising the structure




2-methylcinnamoylguanidine comprising the structure




5

2-(trifluoromethyl)cinnamoylguanidine comprising the structure




4-methylcinnamoylguanidine comprising the structure



10


4-isopropylcinnamoylguanidine comprising the structure




15

-25-


3-fluorocinnamoylguanidine comprising the structure




2-fluorocinnamoylguanidine comprising the structure




4-fluorocinnamoylguanidine comprising the structure



10 3,4-dichlorocinnamoylguanidine comprising the structure




2,4-dichlorocinnamoylguanidine comprising the structure




15

-26-


2,6-dichlorocinnamoylguanidine comprising the structure




4-ethoxycinnamoylguanidine comprising the structure




3,4-(methylenedioxy)cinnamoylguanidine comprising the structure



10 3-(2-naphthyl)acryloylguanidine comprising the structure




4-t-butylcinnamoylguanidine comprising the structure




-27-


3,4,5-trimethoxycinnamoylguanidine comprising the structure




2-(1-naphthyl)acetoyleguanidine comprising the structure




5 2,5-dimethylcinnamoylguanidine comprising the structure



2,3-difluorocinnamoylguanidine comprising the structure




3-phenylcinnamoylguanidine comprising the structure




-28-


3-(trans-hept-1-en-1-yl)cinnamoylguanidine comprising the structure




2-ethylcinnamoylguanidine comprising the structure



2-chloro-6-fluorocinnamoylguanidine comprising the structure




10 3-t-butylcinnamoylguanidine comprising the structure




-29-

3,4-difluorocinnamoylguanidine comprising the structure




5-bromo-2-fluorocinnamoylguanidine comprising the structure




5

3-(trifluoromethoxy)cinnamoylguanidine comprising the structure




2-ethoxycinnamoylguanidine comprising the structure



10


2-t-butylcinnamoylguanidine comprising the structure




15

-30-

3-(cyclohex-1-en-1-yl)cinnamoylguanidine comprising the structure




cinnamoylguanidine hydrochloride comprising the structure




5

2,3,5,6,-tetramethylcinnamoylguanidine comprising the structure (Bit 134)




2-cyclohexylcinnamoylguanidine comprising the structure




10

5-bromo-2-methoxycinnamoylguanidine comprising the structure




-31-

2,3-dimethylcinnamoylguanidine comprising the structure




3-ethoxycinnamoylguanidine comprising the structure




5

3-isopropylcinnamoylguanidine hydrochloride comprising the structure




.HCl

2-phenylcinnamoylguanidine comprising the structure




10

2-(cyclohex-1-en-1-yl)cinnamoylguanidine comprising the structure




-32-


2,4,6-trimethylcinnamoylguanidine comprising the structure




(5-Phenyl-penta-2,4-dienoyl)guanidine comprising the structure




5-(3'-bromophenyl)penta-2,4-dienoylguanidine comprising the structure



10 5-(2'-bromophenyl)penta-2,4-dienoylguanidine comprising the structure



Furanacryloyl comprising the structure



-33-

Preferably, the compounds of the invention are capable of reducing, retarding or otherwise inhibiting viral growth and/or replication.

Preferably, the antiviral activity of the compounds of the invention is against viruses such as those belonging to the Lentivirus family, and the Coronavirus family
5 family of viruses. For example, the compounds of the invention exhibit antiviral activity against viruses such as Human Immunodeficiency Virus (HIV), Severe Acute Respiratory Syndrome virus (SARS), Mouse Hepatitis virus (), and Hepatitis C virus (HCV).

According to a fourth aspect of the present invention, there is provided a
10 pharmaceutical composition comprising an antiviral compound according to any one of the first, second or third aspects, and optionally one or more pharmaceutical acceptable carriers or derivatives, wherein said compound is capable of reducing, retarding or otherwise inhibiting viral growth and/or replication.

Preferably, the antiviral activity of the compounds of the invention is against
15 viruses such as those belonging to the Lentivirus family, and the Coronavirus family of viruses. For example, the compounds of the invention exhibit antiviral activity against viruses such as Human Immunodeficiency Virus (HIV), Severe Acute Respiratory Syndrome virus (SARS), Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine
20 Respiratory Coronavirus (PRCV), Hepatitis C virus (HCV) and Equine Arteritis Virus (EAV).

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

The compositions of the invention may further comprise one or more known
25 antiviral compounds or molecules.

According to a fifth aspect, there is provided a method for reducing, retarding or otherwise inhibiting growth and/or replication of a virus comprising contacting a cell infected with said virus or exposed to said virus with a compound according to any one of the first, second or third aspects.

30 Preferably, the virus is from the Lentivirus family, or the Coronavirus family. More preferably, the virus is Human Immunodeficiency Virus (HIV), Severe Respiratory Syndrome virus (SARS), Human Coronavirus 229E, Human Coronavirus

-34-

OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV), Mouse Hepatitis virus (MHV), Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, the virus is HIV-1, HIV-2, the SARS virus, Coronaviruse 229E, Coronavirus OC43, PRCV, BCV, HCV, or
5 EAV.

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

According to a sixth aspect, there is provided a method for preventing the infection of a cell exposed to a virus comprising contacting said cell with a compound
10 according to any one of the first, second or third aspects.

Preferably, the virus is from the Lentivirus family, or the Coronavirus family. More preferably, the virus is Human Immunodeficiency Virus (HIV), Severe Respiratory Syndrome virus (SARS), Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine
15 Respiratory Coronavirus (PRCV), Mouse Hepatitis virus (MHV), Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, the virus is HIV-1, HIV-2, the SARS virus, Coronaviruse 229E, Coronavirus OC43, PRCV, BCV, HCV, EAV.

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

20 According to a seventh aspect of the invention, there is provided a method for the therapeutic or prophylactic treatment of a subject infected with or exposed to a virus, comprising the administration of a compound according to any one of the first, second or third aspects, to a subject in need of said treatment.

Preferably, infection with a virus or exposure to a virus occurs with viruses
25 belonging to the Lentivirus family, or the Coronavirus family. More preferably, infection or exposure occurs with HIV, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV), Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, infection or exposure occurs with HIV-1,
30 HIV-2, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV).

-35-

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

The subject of the viral inhibition is generally a mammal such as but not limited to human, primate, livestock animal (e.g. sheep, cow, horse, donkey, pig),
5 companion animal (e.g. dog, cat), laboratory test animal (e.g. mouse, rabbit, rat, guinea pig, hamster), captive wild animal (e.g. fox, deer). Preferably, the subject is a primate, or horse. Most preferably, the subject is a human.

According to a eighth aspect, there is provided a method of down regulating a membrane ion channel functional activity in a cell infected with a virus, comprising
10 contacting said cell with a compound according to any one of the first, second or third aspects.

The membrane ion channel may be endogenous to the cell or exogenous to the cell.

Preferably, the membrane ion channel of which functional activity is down
15 regulated is that which Lentiviruses, and Coronaviruses utilise for mediating viral replication and include, for example, the HIV membrane ion channel Vpu, the HCV membrane ion channel P7, the Coronavirus E protein membrane ion channel, and the SARS E protein membrane ion channel.

Preferably, infection with a virus or exposure to a virus occurs with viruses
20 belonging to the Lentivirus family, or the Coronavirus family. More preferably, infection or exposure occurs with HIV, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV), Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, infection or exposure occurs with HIV-1,
25 HIV-2, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV).

According to an ninth aspect of the present invention, there is provided a method of reducing, retarding or otherwise inhibiting growth and/or replication of a virus that has infected a cell, said method comprising contacting said infected cell with a
30 compound according to any one of the first, second or third aspects, wherein said compound down regulates functional activity of a membrane ion channel derived from said virus and expressed in said infected cell.

-36-

Preferably, infection occurs with a virus belonging to the Lentivirus family, or the Coronavirus family. More preferably, infection or exposure occurs with HIV, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV),
5 Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, infection or exposure occurs with HIV-1, HIV-2, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV).

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

10 Preferably, the membrane ion channel of which functional activity is down regulated is that which Lentiviruses, and Coronaviruses utilise for mediating viral replication and include, for example, the HIV membrane ion channel Vpu, the HCV membrane ion channel P7, and the Coronavirus E protein membrane ion channel.

According to an tenth aspect, the present invention provides a method of
15 reducing, retarding or otherwise inhibiting growth and/or replication of a virus that has infected a cell in a mammal, said method comprising administering to said mammal a compound according to any one of the first, second or third aspects, or a pharmaceutical composition according to the fourth aspect, wherein said compound or said composition down regulates functional activity of a membrane ion channel
20 expressed in said infected cell.

Preferably, infection occurs with a virus belonging to the Lentivirus family, or the Coronavirus family. More preferably, infection or exposure occurs with HIV, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV),
25 Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, infection or exposure occurs with HIV-1, HIV-2, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV).

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

30 Preferably, the membrane ion channel of which functional activity is down regulated is that which Lentiviruses, and Coronaviruses utilise for mediating viral

-37-

replication and include, for example, the HIV membrane ion channel Vpu, the HCV membrane ion channel P7, and the Coronavirus E protein membrane ion channel.

The subject of the viral inhibition is generally a mammal such as but not limited to human, primate, livestock animal (e.g. sheep, cow, horse, donkey, pig), companion
5 animal (e.g. dog, cat), laboratory test animal (e.g. mouse, rabbit, rat, guinea pig, hamster), captive wild animal (e.g. fox, deer). Preferably, the subject is a primate, or horse. Most preferably, the subject is a human.

According to a eleventh aspect, the present invention provides a method for the therapeutic or prophylactic treatment of a subject infected with or exposed to a virus
10 comprising administering to said subject a compound according to any one of the first, second or third aspects, or a pharmaceutical composition according to the fourth aspect, wherein said compound or said composition down-regulates functional activity of a membrane ion channel derived from said virus.

Preferably, infection occurs with a virus belonging to the Lentivirus family, or
15 the Coronavirus family of viruses. More preferably, infection or exposure occurs with HIV, SARS, Human Coronavirus 229E, Human Coronavirus OC43, Mouse Hepatitis virus (MHV), Bovine Coronavirus (BCV), Porcine Respiratory Coronavirus (PRCV), Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV). Most preferably, infection or exposure occurs with HIV-1, HIV-2, SARS, Human Coronavirus 229E, Human
20 Coronavirus OC43, Hepatitis C virus (HCV), or Equine Arteritis Virus (EAV).

Other Coronaviruses which can be inhibited or their infections treated by the compounds of the invention are those listed in Table 1.

Preferably, the membrane ion channel of which functional activity is down regulated is that which Lentiviruses, and Coronaviruses utilise for mediating viral
25 replication and include, for example, the HIV membrane ion channel Vpu, the HCV membrane ion channel P7, and the Coronavirus E protein membrane ion channel.

The subject of the viral inhibition is generally a mammal such as but not limited to human, primate, livestock animal (e.g. sheep, cow, horse, donkey, pig), companion
30 animal (e.g. dog, cat), laboratory test animal (e.g. mouse, rabbit, rat, guinea pig, hamster), captive wild animal (e.g. fox, deer). Preferably, the subject is a primate, or horse. Most preferably, the subject is a human.

-38-

According to a twelfth aspect, the invention provides an antiviral compound selected from the group consisting of:

- N-(3,5-Diamino-6-chloro-pyrazine-2-carbonyl)-N'-phenyl-guanidine,
- N-Benzyl-N'-(3,5-diamino-6-chloro-pyrazine-2-carbonyl)-guanidine,
- 5 3'4 DichloroBenzamil,
- 2'4 DichloroBenzamil,
- 5-(N-methyl-N-guanidinocarbonyl-methyl)amiloride,
- 5-(N-Methyl-N-isobutyl)amiloride,
- 5-(N-Ethyl-N-isopropyl)amiloride,
- 10 5-(N,N-Dimethyl)amiloride hydrochloride,
- 5-(N,N-hexamethylene)amiloride,
- 5-(N,N-Diethyl)amiloride hydrochloride,
- 6-Iodoamiloride,
- Bodipy-FL amiloride,
- 15 3-hydroxy-5-hexamethyleneimino-amiloride,
- 5-(4-fluorophenyl)amiloride,
- 5-tert-butylamino-amiloride,
- N-amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide,
- 3-methoxy-5-(N,N-Hexamethylene)-amiloride,
- 20 3-methoxy-amiloride,
- hexamethyleneimino-6-phenyl-2-pyrazinecarboximide,
- N-amidino-3,5-diamino-6-phenyl-2-pyrazinecarboxamide,
- 1-naphthoylguanidine,
- 2-naphthoylguanidine,
- 25 N-(2-naphthoyl)-N'-phenylguanidine,
- N,N'-bis(2-naphthoyl)guanidine,
- N,N'-bis(1-naphthoyl)guanidine,
- N,N'-bis(2-naphthoyl)-N''-phenylguanidine,
- 6-methoxy-2-naphthoylguanidine,
- 30 3-quinolinoylguanidine,
- cinnamoylguanidine,
- 4-phenylbenzoylguanidine,

-39-

- N-(cinnamoyl)-N'-phenylguanidine,
(3-phenylpropanoyl)guanidine,
N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
5 N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
trans-3-furanacryoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(4-Phenoxybenzoyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
10 N'-Cinnamoyl-N,N'-diphenylguanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(3-phenylpropanoyl)guanidine,
benzoylguanidine,
(4-Chlorophenoxy-acetyl]guanidine,
15 N-benzoyl-N'-cinnamoylguanidine,
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine,
(4-Chlorocinnamoyl)guanidine,
(4-Bromocinnamoyl)guanidine,
(4-Methoxycinnamoyl)guanidine,
20 (5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
25 (2-Bromocinnamoyl)guanidine,
(2-Methoxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
(4-Hydroxycinnamoyl)guanidine,
30 (Quinoline-2-carbonyl)guanidine,
or pharmaceutically acceptable salts thereof.

-40-

According to a thirteenth aspect, the present invention provides a pharmaceutical composition comprising a compound according to the twelfth aspect, and optionally one or more pharmaceutical acceptable carriers or derivatives.

Preferably, the pharmaceutical composition may further comprise one or more
5 known antiviral compounds or molecules.

Unless the context clearly requires otherwise, throughout the description and the claims, the words 'comprise', 'comprising', and the like are to be construed in an inclusive sense as opposed to an exclusive or exhaustive sense; that is to say, in the sense of "including, but not limited to".

10 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic representation of plasmids used for expression of Vpu in *E. coli*. A. The amino acid sequence ($< 400 > 1$) encoded by the *vpu* open reading frame (ORF) generated by PCR from an HIV-1 strain HXB2 cDNA clone. The *vpu* ORF was cloned in-frame at the 3' end of the GST gene in p2GEX to generate
15 p2GEXVpu (B). It was subsequently cloned into pPL451 to produce the plasmid pPL + Vpu (C).

Figure 2 is a photographic representation of the expression and purification of Vpu in *E. coli*. A. Western blotting after SDS-PAGE was used to detect expressed Vpu in *E. coli* extracts. Lanes 1-4 contain samples, at various stages of purity, of Vpu expressed
20 from p2GEXVpu: lane 1, GST-Vpu fusion protein isolated by glutathione-agarose affinity chromatography; lane 2, Vpu liberated from the fusion protein by treatment with thrombin; lane 3, Vpu purified by HPLC anion exchange chromatography; lane 4, Vpu after passage through the immunoaffinity column. Lanes 5 and 6, membrane vesicles prepared from 42°C induced cells containing pPL+Vpu or pPL451,
25 respectively. B. Silver stained SDS-PAGE gel: lane 1, Vpu purified by HPLC anion exchange chromatography; lane 2, Vpu after passage through the immunoaffinity column.

Figure 3 is a graphical representation of ion channel activity observed after exposure of lipid bilayers to aliquots containing purified Vpu. In A and B, the CIS chamber
30 contained 500mM NaCl and the TRANS chamber contained 50mM NaCl; both

-41-

solutions were buffered at pH 6.0 with 10 mM MES. **B** shows a current versus voltage curve generated from data similar to that shown in **A**.

Figure 4 is a photographic representation of bacterial cross-feeding assays. For all plates, the Met⁻, Pro⁻ auxotrophic strain was used to seed a soft agar overlay. Plates **A** and **B** contain minimal drop-out medium minus proline; in plate **C** the medium was minus methionine. To control for viability of the cells in the background lawn, the discs labelled **P** and **M** contained added proline or methionine, respectively. The discs labelled **C** and **V** were inoculated with Met⁺, Pro⁺ *E. coli* cells containing the plasmids pPL451 or pPL+Vpu, respectively. Plates were incubated at 37°C (**A** and **C**) or 30°C (**B**) for two days and photographed above a black background with peripheral illumination from a fluorescent light located below the plate. The images were recorded on a Novaline video gel documentation system. Light halos around the discs labelled **P** or **M** on all plates and around the disc labelled **V** on plate **A** indicate growth of the background lawn strain.

Figure 5 is a graphical representation of the screening of drugs for potential Vpu channel blockers. The photograph shows a section of a minimal medium-lacking adenine - agarose plate onto which a lawn of XL-I-blue *E. coli* cells containing the Vpu expression plasmid pPLVpu has been seeded. Numbers 6-11 are located at the sites of application of various drugs being tested, which were applied in 3µl drops and allowed to soak into the agarose. The plate was then incubated at 37°C for 48hr prior to being photographed. The background grey shade corresponds to areas of no bacterial growth. The bright circular area around "10" represents bacterial cell growth as a result of application of adenine at that location (positive control). The smaller halo of bacterial growth around "9" is due to the application of 5-(N,N-hexamethylene)-amiloride at that location.

Figure 6. SARS E protein ion channel activity observed in NaCl solutions after exposure of lipid bilayer to 3-10µg of E protein. **A.** The closed state is shown as solid line, openings are derivations from the line. Scale bar is 300ms and 5pA. The CIS chamber contained 50mM NaCl in 5mM HEPES buffer pH 7.2, the TRANS chamber contained 500mM NaCl in 5mM HEPES buffer pH 7.2. The CIS chamber was

-42-

earthed and the TRANS chamber was held at various potentials between -100 to $+100\text{mV}$. B. Largest single opening events of a single channel.

Figure 7. SARS E protein ion channel activity observed in NaCl solutions after exposure of lipid bilayer to $3\text{-}10\mu\text{g}$ of E protein. A. The closed state is shown as solid line, openings are derivations from the line. Scale bar is 300ms and 5pA . The CIS chamber contained 50mM NaCl in 5mM HEPES buffer pH 7.2, the TRANS chamber contained 500mM NaCl in 5mM HEPES buffer pH 7.2. The CIS chamber was earthed and the TRANS chamber was held at various potentials between -100 to $+100\text{mV}$. B. Largest single opening events of a single channel.

Figure 8. Cinnamoylguanidine (Bit036) inhibits SARS E protein ion channel activity in NaCl solution. A. Representative currents at holding potential of -40mV . Scale bar is 300mS and 5pA . E protein ion channel activity and E protein channel activity after the addition of $100\mu\text{M}$ Bit036. B. All points histogram at holding potential of -40mV . E protein ion channel activity before and after the addition of $100\mu\text{M}$ Bit036. C. Average current (pA), before formation of E protein ion channel, E protein ion channel activity and after addition of $100\mu\text{M}$ Bit036.

Figure 9 229E E protein Ion channel activity in lipid bilayers in KCl solutions.

Figure 10: Part A shows raw currents generated by the 229E-E protein ion channel in a planar lipid bilayer. The top trace shows current activity prior to drug addition and the lower trace shows the effect of addition of $100\mu\text{M}$ cinnamoylguanidine on channel activity. Part B is a graphical representation of the average current flowing across the bilayer (in arbitrary units), before and after addition of cinnamoylguanidine.

Figure 11: MHV E protein Ion channel activity in lipid bilayers NaCl solutions.

Figure 12: Part A shows raw currents generated by the MHV-E protein ion channel in a planar lipid bilayer. The top trace shows current activity prior to drug addition and the lower trace shows the effect of addition of $100\mu\text{M}$ cinnamoylguanidine on channel activity. Part B is a graphical representation of the average current flowing

-43-

across the bilayer (in arbitrary units), before and after addition of cinnamoylguanidine.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based, in part, on the surprising determination that
5 certain compounds that fall under the classification of substituted acylguanidines have antiviral activity against viruses from a range of different virus families. Without intending to be bound by any particular theory or mechanism of action, the negative impact of the compounds of the present invention on viral replication may be mediated by the inhibition or otherwise down-regulation of a membrane ion channel
10 relied upon by the virus for replication. This membrane ion channel may be a viral membrane ion channel (exogenous to the host cell) or a host cell ion channel induced as a result of viral infection (endogenous to the host cell).

As an example, the compounds of the present invention may inhibit Vpu or p7 function and thereby inhibit the continuation of the respective HIV or HCV life cycle.

15 The SARS virus encodes an E protein which is shown for the first time, by the present inventors, to act as an ion channel. As similar E proteins are present in other coronaviruses, the compounds, compositions and methods of the present invention would have utility in the inhibition and/or treatment of infections by other coronaviruses.

20 While the present invention is concerned with novel antiviral compounds falling under the classification of substituted acylguanidines, it does not include in its scope the use of compounds 5-(N,N-hexamethylene)amiloride and 5-(N,N-dimethyl)-amiloride for retarding, reducing or otherwise inhibiting viral growth and/or functional activity of HIV.

25 It will be understood by those skilled in the art that the compounds of the invention may be administered in the form of a composition or formulation comprising pharmaceutically acceptable carriers and excipients.

The pharmaceutical compositions of the invention may further comprise one or more known antiviral compounds or molecules. Preferably, the known antiviral
30 compounds are selected from the group consisting of Vidarabine, Acyclovir, Ganciclovir, Valganciclovir, Valacyclovir, Cidofovir, Fanciclovir, Ribavirin,

-44-

Amantadine, Rimantadine, Interferon, Oseltamivir, Palivizumab, Rimantadine, Zanamivir, nucleoside-analog reverse transcriptase inhibitors (NRTI) such as Zidovudine, Didanosine, Zalcitabine, Stavudine, Lamivudine and Abacavir, non-nucleoside reverse transcriptase inhibitors (NNRTI) such as Nevirapine, Delavirdine
 5 and Efavirenz, protease inhibitors such as Saquinavir, Ritonavir, Indinavir, Nelfinavir, Amprenavir, and other known antiviral compounds and preparations. Known antiviral compounds or molecules may in some cases act synergistically with the antiviral compounds of the invention.

10 Table 1 Known coronavirus isolates

Group 1 species
Canine coronavirus
Canine enteric coronavirus (strain INSAVC-1)
Canine enteric coronavirus (strain K378)
Feline coronavirus
Feline enteric coronavirus (strain 79-1683)
Feline infectious peritonitis virus (FIPV)
Human coronavirus 229E
Porcine epidemic diarrhea virus
Porcine epidemic diarrhea virus (strain Br1/87)
Porcine epidemic diarrhea virus (strain CV777)
Transmissible gastroenteritis virus
Porcine respiratory coronavirus
Porcine transmissible gastroenteritis coronavirus (STRAIN FS772/70)
Porcine transmissible gastroenteritis coronavirus (strain Miller)
Porcine transmissible gastroenteritis coronavirus (strain Neb72-RT)
Porcine transmissible gastroenteritis coronavirus (STRAIN PURDUE)
Group 2 species
Bovine coronavirus
Bovine coronavirus (STRAIN F15)
Bovine coronavirus (strain G95)
Bovine coronavirus (STRAIN L9)
Bovine coronavirus (strain LSU-94LSS-051)
Bovine coronavirus (STRAIN LY-138)
Bovine coronavirus (STRAIN MEBUS)
Bovine coronavirus (strain OK-0514-3)
Bovine coronavirus (strain Ontario)
Bovine coronavirus (STRAIN QUEBEC)
Bovine coronavirus (STRAIN VACCINE)
Bovine enteric coronavirus (strain 98TXSF-110-ENT)
Canine respiratory coronavirus

Chicken enteric coronavirus
Human coronavirus OC43
Murine hepatitis virus
Murine coronavirus (strain DVIM)
Murine hepatitis virus (strain A59)
Murine hepatitis virus (strain JHM)
Murine hepatitis virus (strain S)
Murine hepatitis virus strain 1
Murine hepatitis virus strain 2
Murine hepatitis virus strain 3
Murine hepatitis virus strain 4
Murine hepatitis virus strain ML-11
Porcine hemagglutinating encephalomyelitis virus
Porcine hemagglutinating encephalomyelitis virus (strain 67N)
Porcine hemagglutinating encephalomyelitis virus (strain IAF-404)
Puffinosis virus
Rat coronavirus
Rat coronavirus (strain 681)
Rat coronavirus (strain NJ)
Rat sialodacryoadenitis coronavirus
Group 3 species
Turkey coronavirus
Turkey coronavirus (strain Indiana)
Turkey coronavirus (strain Minnesota)
Turkey coronavirus (strain NC95)
Avian infectious bronchitis virus
Avian infectious bronchitis virus (STRAIN 6/82)
Avian infectious bronchitis virus (strain Arkansas 99)
Avian infectious bronchitis virus (strain Beaudette CK)
Avian infectious bronchitis virus (strain Beaudette M42)
Avian infectious bronchitis virus (strain Beaudette US)
Avian infectious bronchitis virus (strain Beaudette)
Avian infectious bronchitis virus (strain D1466)
Avian infectious bronchitis virus (strain D274)
Avian infectious bronchitis virus (strain D3896)
Avian infectious bronchitis virus (strain D41)
Avian infectious bronchitis virus (strain DE072)
Avian infectious bronchitis virus (strain GRAY)
Avian infectious bronchitis virus (strain H120)
Avian infectious bronchitis virus (strain H52)
Avian infectious bronchitis virus (strain KB8523)
Avian infectious bronchitis virus (strain M41)
Avian infectious bronchitis virus (strain PORTUGAL/322/82)
Avian infectious bronchitis virus (strain SAIB20)

-46-

Avian infectious bronchitis virus (strain UK/123/82)
Avian infectious bronchitis virus (strain UK/142/86)
Avian infectious bronchitis virus (strain UK/167/84)
Avian infectious bronchitis virus (strain UK/183/66)
Avian infectious bronchitis virus (strain UK/68/84)
Avian infectious bronchitis virus (strain V18/91)
Avian infectious bronchitis virus (strain Vic S)
Avian infectious laryngotracheitis virus
Preliminary Group 4 species
SARS coronavirus
SARS coronavirus Beijing ZY-2003
SARS coronavirus BJ01
SARS coronavirus BJ02
SARS coronavirus BJ03
SARS coronavirus BJ04
SARS coronavirus CUHK-Su10
SARS coronavirus CUHK-W1
SARS coronavirus Frankfurt 1
SARS coronavirus GZ01
SARS coronavirus HKU-39849
SARS coronavirus Hong Kong ZY-2003
SARS coronavirus Hong Kong/03/2003
SARS coronavirus HSR 1
SARS coronavirus Sin2500
SARS coronavirus Sin2677
SARS coronavirus Sin2679
SARS coronavirus Sin2748
SARS coronavirus Sin2774
SARS coronavirus Taiwan
SARS coronavirus Taiwan JC-2003
SARS coronavirus Taiwan TC1
SARS coronavirus Taiwan TC2
SARS coronavirus Tor2
SARS coronavirus TW1
SARS coronavirus TWC
SARS coronavirus Urbani
SARS coronavirus Vietnam
SARS coronavirus ZJ-HZ01
SARS coronavirus ZJ01
unclassified coronaviruses
Bovine respiratory coronavirus (strain 98TXSF-110-LUN)
Human enteric coronavirus 4408
Enteric coronavirus
Equine coronavirus

-47-

Equine coronavirus NC99

The present observations and findings now permit the use of agents such as certain substituted acylguanidines, as anti-viral agents for the therapy and prophylaxis of viral conditions caused by different viruses. The methods and compositions of the present invention may be particularly effective against viruses which rely on ion channel formation for their replication, however it will be understood that this is not the only mechanism relied on by viruses for replication and that the compounds and methods of the present invention are not limited to agents which exert their action by retarding or inhibiting the function of ion channels.

Reference to "membrane ion channel" should be understood as a reference to a structure which transports ions across a membrane. The present invention extends to ion channels which may function by means such as passive, osmotic, active or exchange transport. The ion channel may be formed by intracellular or extracellular means. For example, the ion channel may be an ion channel which is naturally formed by a cell to facilitate its normal functioning. Alternatively, the ion channel may be formed by extracellular means. Extracellular means would include, for example, the formation of ion channels due to introduced chemicals, drugs or other agents such as ionophores or due to the functional activity of viral proteins encoded by a virus which has entered a cell.

The ion channels which are the subject of certain embodiments of the present invention facilitate the transport of ions across membranes. Said membrane may be any membrane and is not limited to the outer cell wall plasma membrane. Accordingly, "membrane" as used herein encompasses the membrane surrounding any cellular organelle, such as the Golgi apparatus and endoplasmic reticulum, the outer cell membrane, the membrane surrounding any foreign antigen which is located within the cell (for example, a viral envelope) or the membrane of a foreign organism which is located extracellularly. The membrane is typically, but not necessarily, composed of a fluid lipid bilayer. The subject ion channel may be of any structure. For example, the Vpu ion channel is formed by Vpu which is an integral membrane protein encoded by HIV-1 which associates with, for example, the Golgi and endoplasmic reticulum membranes of infected cells. Reference hereinafter to "Vpu

-48-

ion channels" is a reference to all related ion channels for example P7 HCV and M2 of influenza and the like.

Reference to "HIV", "SARS", "Coronavirus" or "HCV" should be understood as a reference to any HIV, SARS, Coronavirus or HCV virus strain and including
5 homologues and mutants.

Reference to the "functional activity" of an ion channel should be understood as a reference to any one or more of the functions which an ion channel performs or is involved in. For example, the Vpu protein encoded ion channel, in addition to facilitating the transportation of Na^+ , K^+ , Cl^- and PO_4^{3-} , also plays a role in the
10 degradation of the CD4 molecule in the endoplasmic reticulum. Without wishing to be bound by a particular theory, the Vpu protein encoded ion channel is also thought to play a role in mediating the HIV life cycle. The present invention is not limited to treating HIV infection via the mechanism of inhibiting the HIV life cycle and, in particular, HIV replication. Rather, the present invention should be understood to
15 encompass any mechanism by which the compounds of the present invention exert their anti-viral activity and may include inhibition of HIV viability or functional activity. This also applies to HCV, Coronaviruses, and to other viruses.

Reference to the "functional activity" of a virus should be understood as a reference to any one or more of the functions which a virus performs or is involved
20 in.

Reference to the "viral replication" should be understood to include any one or more stages or aspects of the viral life cycle, such as inhibiting the assembly or release of virions. Ion channel mediation of viral replication may be by direct or indirect means. Said ion channel mediation is by direct means if the ion channel
25 interacts directly with the virion at any one or more of its life cycle stages. Said ion channel mediation is indirect if it interacts with a molecule other than those of the virion, which other molecule either directly or indirectly modulates any one or more aspects or stages of the viral life cycle. Accordingly, the method of the present invention encompasses the mediation of viral replication via the induction of a
30 cascade of steps which lead to the mediation of any one or more aspects or stages of the viral life cycle.

-49-

Reference to "down-regulating" ion channel functional activity, should be understood as a reference to the partial or complete inhibition of any one or more aspects of said activity by both direct and indirect mechanisms. For example, a suitable agent may interact directly with an ion channel to prevent replication of a virus or, alternatively, may act indirectly to prevent said replication by, for example, interacting with a molecule other than an ion channel. A further alternative is that said other molecule interacts with and inhibits the activity of the ion channel.

Screening for molecules that have antiviral activity can be achieved by the range of methodologies described herein.

Reference to a "cell" infected with a virus should be understood as a reference to any cell, prokaryotic or eukaryotic, which has been infected with a virus. This includes, for example, immortal or primary cell lines, bacterial cultures and cells *in situ*. In a suitable screening system for antiviral compounds, the preferred infected cells would be macrophages/monocytes or hepatocytes/lymphoid cells infected with either HIV or HCV respectively.

Without limiting the present invention to any one theory or mode of action, the compounds of the present invention are thought to inhibit viral replication or virion release from cells by causing ion channels, namely VPU of HIV, the E protein of SARS and other Coronaviruses, or P7 of HCV to become blocked. The present invention encompasses antiviral compounds that are substituted acylguanidines.

The present invention also includes the use of compounds 5-(N,N-hexamethylene)amiloride and 5-(N,N-dimethyl)-amiloride in the control of viral replication and/or growth other than HIV.

The subject of the viral inhibition is generally a mammal such as but not limited to human, primate, livestock animal (e.g. sheep, cow, horse, donkey, pig), companion animal (e.g. dog, cat), laboratory test animal (e.g. mouse, rabbit, rat, guinea pig, hamster), captive wild animal (e.g. fox, deer). Preferably, the subject is a human or primate. Most preferably, the subject is a human.

The method of the present invention is useful in the treatment and prophylaxis of viral infection such as, for example, but not limited to HIV infection, HCV infection and other viral infections. For example, the antiviral activity may be effected in subjects known to be infected with HIV in order to prevent replication of

-50-

HIV thereby preventing the onset of AIDS. Alternatively, the method of the present invention may be used to reduce serum viral load or to alleviate viral infection symptoms. Similarly, antiviral treatment may be effected in subjects known to be infected with, for example, HCV, in order to prevent replication of HCV, thereby
5 preventing the further hepatocyte involvement and the ultimate degeneration of liver tissue.

The method of the present invention may be particularly useful either in the early stages of viral infection to prevent the establishment of a viral reservoir in affected cells or as a prophylactic treatment to be applied immediately prior to or for
10 a period after exposure to a possible source of virus.

Reference herein to "therapeutic" and "prophylactic" is to be considered in their broadest contexts. The term "therapeutic" does not necessarily imply that a mammal is treated until total recovery. Similarly, "prophylactic" does not necessarily mean that the subject will not eventually contract a disease condition. Accordingly,
15 therapy and prophylaxis include amelioration of the symptoms of a particular condition or preventing or otherwise reducing the risk of developing a particular condition. The term "prophylaxis" may be considered as reducing the severity of onset of a particular condition. Therapy may also reduce the severity of an existing condition or the frequency of acute attacks.

20 In accordance with the methods of the present invention, more than one compound or composition may be co-administered with one or more other compounds, such as known anti-viral compounds or molecules. By "co-administered" is meant simultaneous administration in the same formulation or in two different formulations via the same or different routes or sequential administration by
25 the same or different routes. By "sequential" administration is meant a time difference of from seconds, minutes, hours or days between the administration of the two or more separate compounds. The subject antiviral compounds may be administered in any order.

Routes of administration include but are not limited to intravenously,
30 intraperitoneally, subcutaneously, intracranially, intradermally, intramuscularly, intraocularly, intrathecaly, intracerebrally, intranasally, transmucosally, by infusion,

-51-

orally, rectally, *via* iv drip, patch and implant. Intravenous routes are particularly preferred.

Compositions suitable for injectable use include sterile aqueous solutions (where water soluble) and sterile powders for the extemporaneous preparation of sterile injectable solutions. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol and liquid polyethylene glycol, and the like), suitable mixtures thereof and vegetable oils. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by, for example, filter sterilization or sterilization by other appropriate means. Dispersions are also contemplated and these may be prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, a preferred method of preparation includes vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution.

When the active ingredients are suitably protected, they may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsule, or it may be compressed into tablets. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 0.01 % by weight, more preferably 0.1% by weight, even more preferably 1% by weight of active compound.

-52-

The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 1 to about 99%, more preferably about 2 to about 90 %, even more preferably about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically useful compositions in such that a
5 suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 0.1 ng and 2000 mg of active compound.

The tablets, troches, pills, capsules and the like may also contain the components as listed hereafter: A binder such as gum, acacia, corn starch or gelatin;
10 excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a
15 liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Any material
20 used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

The present invention also extends to forms suitable for topical application such as creams, lotions and gels. In such forms, the anti-clotting peptides may need to
25 be modified to permit penetration of the surface barrier. Procedures for the preparation of dosage unit forms and topical preparations are readily available to those skilled in the art from texts such as *Pharmaceutical Handbook. A Martindale Companion Volume Ed. Ainley Wade Nineteenth Edition The Pharmaceutical Press London,*
30 *CRC Handbook of Chemistry and Physics Ed. Robert C. Weast Ph D. CRC Press Inc.; Goodman and Gilman's; The Pharmacological basis of Therapeutics. Ninth Ed.*

-53-

McGraw Hill; Remington; and The Science and Practice of Pharmacy. Nineteenth Ed. Ed. Alfonso R. Gennaro Mack Publishing Co. Easton Pennsylvania.

Pharmaceutically acceptable carriers and/or diluents include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and
5 absorption delaying agents and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, use thereof in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

10 It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with
15 the required pharmaceutical carrier. The specification for the novel dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active material and the particular therapeutic effect to be achieved and (b) the limitations inherent in the art of compounding.

Effective amounts contemplated by the present invention will vary depending
20 on the severity of the pain and the health and age of the recipient. In general terms, effective amounts may vary from 0.01 ng/kg body weight to about 100 mg/kg body weight.

Alternative amounts include for about 0.1 ng/kg body weight about 100 mg/kg body weight or from 1.0 ng/kg body weight to about 80 mg/kg body weight.

25 The subject of the viral inhibition is generally a mammal such as but not limited to human, primate, livestock animal (e.g. sheep, cow, horse, donkey, pig), companion animal (e.g. dog, cat), laboratory test animal (e.g. mouse, rabbit, rat, guinea pig, hamster), captive wild animal (e.g. fox, deer). Preferably, the subject is a human or primate. Most preferably, the subject is a human.

30 The methods of the present invention is useful in the treatment and prophylaxis of viral infection such as, for example, but not limited to HIV infection, HCV infection and other viral infections. For example, the antiviral activity may be

-54-


effected in subjects known to be infected with HIV in order to prevent replication of HIV thereby preventing the onset of AIDS. Alternatively, the methods of the present invention may be used to reduce serum viral load or to alleviate viral infection symptoms. Similarly, antiviral treatment may be effected in subjects known to be infected with, for example, HCV, in order to prevent replication of HCV, thereby preventing the further hepatocyte involvement and the ultimate degeneration of liver tissue.

The methods of the present invention may be particularly useful either in the early stages of viral infection to prevent the establishment of a viral reservoir in affected cells or as a prophylactic treatment to be applied immediately prior to or for a period after exposure to a possible source of virus.

The present invention will now be described in more detail with reference to specific but non-limiting examples describing studies of viral membrane ion channels and screening for antiviral activity. Some examples involve the use of the SARS virus. It will be clear from the description herein that other lentiviruses, and coronaviruses and other compounds may be used effectively in the context of the present invention. It is to be understood, however, that the detailed description is included solely for the purpose of exemplifying the present invention. It should not be understood in any way as a restriction on the broad description of the invention as set out above.

Example 1. Synthesis of the Compounds of the Invention.


The compounds of the present invention may be made from the corresponding acid chlorides or methyl esters as shown in Scheme 1. Both of these methods are well described in the literature.



-55-

The following examples show synthetic schemes for some compounds of the invention.

Example 2. Synthesis of Cinnamoylguanidine from Cinnamic acid Cinnamoyl chloride




To a solution of *trans*-cinnamic acid (1.50 g, 10.12 mmol) in dry benzene (30mL) containing a drop of *N,N*-dimethylformamide was added oxalyl chloride (5.14 g, 40.5 mmol) causing the solution to effervesce. After refluxing for 2 h, the solution was evaporated to dryness under reduced pressure. The resulting solid was dissolved in dry tetrahydrofuran (20mL) and added slowly to a solution of guanidine hydrochloride in 2M aqueous sodium hydroxide (25mL). The reaction was stirred at room temperature for 1h then extracted with ethyl acetate (3x50mL). The combined extracts were dried over magnesium sulfate and evaporated to give an orange oil. The crude product was purified by column chromatography. Elution with 10% to 20% methanol in dichloromethane gave *Cinnamoylguanidine* as a cream solid (0.829 g, 43%).

Example 3

Synthesis of N-amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide

Part 1




To a solution of methyl 3-amino-5,6-dichloro-2-pyrazinecarboxylate (0.444 g, 2.0 mmol) in tetrahydrofuran (5 mL) / water (10 mL) / toluene (20 mL) was added phenyl boronic acid (0.536 g, 4.4 mmol), sodium carbonate (0.699 g, 6.6 mmol) and tetrakis(triphenylphosphine)-palladium(0) (0.116 g, 0.10 mmol). The reaction was

-56-

evacuated and purged with nitrogen several times before being refluxed for 6 h. The organic layer was separated and the aqueous layer extracted with toluene (3 x 20 mL). The combined organic extracts were dried over magnesium sulfate, filtered and evaporated under reduced pressure to give *methyl 3-amino-6-chloro-5-*

5 *phenyl-2-pyrazinecarboxylate* as a yellow solid (0.43 g, 82%).

Part 2




To a solution of sodium (0.040 g, 1.74 mmol) dissolved in methanol (5 mL) was added guanidine hydrochloride (0.258 g, 2.70 mmol) and the mixture refluxed for 30 min after which it was filtered. To the filtrate was added methyl 3-amino-6-chloro-5-phenyl-2-pyrazinecarboxylate (0.264 g, 1.0 mmol) in *N,N*-dimethylformamide (5 mL) and the solution heated at 75°C for 12 h. The solvent was removed under reduced pressure and the residue chromatographed on silica gel eluting with 1% triethylamine / 5% methanol / dichloromethane. The resulting solid was suspended in chloroform, filtered and dried under high vacuum to give *N-Amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide* as a yellow solid (0.04 g, 14%).

Example 4.

20 Synthesis of hexamethyleneimino-6-phenyl-2-pyrazinecarboxamide

Part 1




To a solution of methyl 3-amino-5,6-dichloro-2-pyrazinecarboxylate (1.11 g, 5.0 mmol) in tetrahydrofuran (50 mL) was added hexamethylenimine (1.49 g, 15.0 mmol) and the reaction was refluxed for 1 h. The reaction was allowed to cool and the solid hexamethylenimine hydrochloride removed by filtration. The filtrate was

-57-


evaporated and the residue chromatographed over silica gel. Elution with dichloromethane gave *methyl 3-amino-6-chloro-5-hexamethyleneimino-2-pyrazinecarboxylate* as an off-white solid (1.20 g, 85%).

5 Part 2



To a solution of methyl 3-amino-6-chloro-5-hexamethyleneimino-2-pyrazinecarboxylate (0.350 g, 1.23 mmol) in dimethylsulfoxide (5 mL) was added phenyl boronic acid (0.166 g, 1.35 mmol), potassium carbonate (0.511 g, 3.70 mmol) and [1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II)-dichloromethane complex (0.041 g, 0.05 mmol). The reaction was heated at 90°C for 16 h before being poured into water (50 mL) and extracted with ethyl acetate (3 x 50 mL). The combined extracts were dried over magnesium sulfate, filtered and evaporated to give a brown oil which was purified by chromatography on silica gel. Elution with dichloromethane followed by 10% ethyl acetate/dichloromethane gave *methyl 3-amino-5-hexamethyleneimino-6-phenyl-2-pyrazinecarboxylate* as a yellow solid (0.309 g, 77%).

Part 3.



20

To a solution of sodium (0.090 g, 6.17 mmol) dissolved in methanol (8 mL) was added guanidine hydrochloride (0.598 g, 6.26 mmol) and the mixture was refluxed for 30 min after which it was filtered. To the filtrate was added methyl 3-amino-5-hexamethyleneimino-6-phenyl-2-pyrazinecarboxylate (0.310 g, 0.95 mmol) in tetrahydrofuran (10 mL) and the solution refluxed for 72 h. The solvent was removed under reduced pressure and the residue chromatographed on silica gel.

25

Elution with 5% methanol/dichloromethane gave *N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-2-pyrazinecarboxamide* as a yellow solid (0.116 g, 35%).

5 **Example 5. Viral Studies**

Construction of recombinant plasmids containing open reading frames encoding various virus proteins.

Complimentary DNA (cDNA) fragments for the various viral proteins listed in Table 2 were obtained either by PCR amplification from a parental virus genome clone, or by direct chemical synthesis of the polynucleotide sequence. For example, the open reading frame encoding Vpu (Fig 1a) was amplified by PCR from a cDNA clone of an Nde I fragment of the HIV-1 genome (isolate HXB2, McFarlane Burnet Centre, Melbourne, Australia) as follows: Native Pfu DNA polymerase (Stratagene; 0.035 U//II) was chosen to catalyse the PCR reaction to minimise possible PCR introduced errors by virtue of the enzyme's proofreading activity. The 5', sense, primer AGTAGGATCCATGCAACCTATACC (SEQ ID NO: 1) (< 400 > 2) introduces a BamH1 site (underlined) for cloning in-frame with the 3' end of the GST gene in p2GEX (41). This primer also repairs the start codon (bold T replaces a Q of the vpu gene which is a threonine codon in the HXB2 isolate. The 3', antisense, primer TCTGGAATTCTACAGATCAT CAAC (SEQ ID NO: 2) (<400> 3) introduces an EcoR1 site (underlined) to the other end of the PCR product to facilitate cloning. After 30 cycles of 94°C for 45 sec, 55°C for 1 min and 72°C for 1 min in 0.5 ml thin-walled eppendorf tubes in a Perkin-Elmer thermocycler, the 268bp fragment was purified, digested with BamH1 and EcoR1 and ligated to p2GEX prepared by digestion with the same two enzymes. The resultant recombinant plasmid is illustrated in Fig 1b. The entire Vpu open reading frame and the BamH1 and EcoR1 ligation sites were sequenced by cycle sequencing, using the Applied Biosystems dye-terminator kit, to confirm the DNA sequence. Other cDNAs were synthesised for us using state of the art methods by GenScript Corporation (New Jersey, USA). Codon sequences were optimised for expression in bacterial, insect or mammalian cells, as appropriate. Restriction endonuclease enzyme recognition sites were incorporated at the 5' and 3'

-59-

ends of the synthetic cDNAs to facilitate cloning into plasmid expression vectors, pcDNA3.1, pFastBac and pPL451 for expression of the encoded virus proteins in mammalian, insect or bacterial cells, respectively.

- 5 Standard techniques of molecular biology were used in cloning experiments. For example, to prepare the Vpu open reading frame for insertion into the pPL451 expression plasmid, p2GEXVpu was first digested with BamHI and the 5' base overhang was filled in the Klenow DNA polymerase in the presence of dNTPs. The Vpu-encoding fragment was then liberated by digestion with EcoRI, purified from an
10 agarose gel and ligated into pPL451 which had been digested with HpaI and EcoRI. Western blots subsequently confirmed that the pPLVpu construct (Fig 1c) expressed Vpu after induction of cultures at 42°C to inactivate the cI857 repressor of the PR and PL promoters.

Table 2 Source of viral cDNA or peptide sequences.

15

Target Protein	Source organism	Strain or Sequence Accession number
Vpu	HIV-1	strain HXB2
SARS-CoV E protein	SARS coronavirus	P59637
HCV p7	Hepatitis C virus H77 1a	NP_751922
MHV-E protein	Murine hepatitis virus	NP_068673
229E E protein	Human coronavirus 229E	NP_073554
Dengue M protein	Dengue virus type 1	Strain Singapore S275/90

Example 6. Purification of Recombinant Vpu from E. Coli.

- Cultures of *E. coli* strain XLI-blue cells containing p2GEXVpu were grown at 30°C with vigorous aeration in LB medium supplemented with glucose (6g/L) and
20 ampicillin (50mg/L) to a density of approximately 250 Klett units, at which time IPTG was added to a final concentration of 0.01mM and growth was continued for a further 4hr. The final culture density was approximately 280 Klett units. Since early experiments revealed that the majority of expressed GST-Vpu fusion protein was associated with both the cell debris and 30 membrane fractions, the method of
25 Varadhachary and Maloney (Varadhachary and Maloney, 1990) was adopted to isolate osmotically disrupted cell ghosts (combining both cell debris and membrane fractions) for the initial purification steps. Cells were harvested, washed, weighed and

-60-

resuspended to 10ml/g wet weight in MTPBS containing DTT (1mM) and $MgCl_2$ (10mM). Lysozyme (0.3 mg/ml; chicken egg white; Sigma) was added and incubated on ice for 30 min with gentle agitation followed by 5 min at 37°C. The osmotically sensitised cells were pelleted at 12,000g and resuspended to the original volume in water to burst the cells. The suspension was then made up to 1xMTPBS/DTT using a 10x buffer stock and the ghosts were isolated by centrifugation and resuspended in MTPBS/DTT to which was then sequentially added glycerol (to 20 % wt/vol) and CHAPS (to 2 % wt/vol) to give a final volume of one quarter the original volume. This mixture was stirred on ice for 1 hr and then centrifuged at 400,000g for 1hr to remove insoluble material. The GST-Vpu fusion protein was purified from the detergent extract by affinity chromatography on a glutathione agarose resin (Sigma). The resin was thoroughly washed in 50mM Tris pH 7.5 containing glycerol (5 %), DTT (1mM), and CHAPS (0.5 %) (Buffer A) and then the Vpu portion of the fusion protein was liberated and eluted from the resin-bound GST by treatment of a 50% (v/v) suspension of the beads with human thrombin (100U/ml; 37°C for 1hr). PMSF (0.5mM) was added to the eluant to eliminate any remaining thrombin activity. This Vpu fraction was further purified on a column of MA7Q anion exchange resin attached to a BioRad HPLC and eluted with a linear NaCl gradient (0-2M) in buffer A.

The Vpu was purified to homogeneity - as determined on silver stained gels - on an immunoaffinity column as follows: HPLC fractions containing Vpu were desalted on a NAP 25 column (Pharmacia) into buffer A and then mixed with the antibody-agarose beads for 1hr at room temperature. The beads were washed thoroughly and Vpu was eluted by increasing the salt concentration to 2M. Protein was quantitated using the BioRad dye binding assay.

Example 7. Expression and Purification of Vpu in E.Coli.

The plasmid p2GEXVpu (Fig. 1) was constructed to create an in-frame gene fusion between the GST and Vpu open-reading frames. This system enabled IPTG-inducible expression of the Vpu polypeptide fused to the C-terminus of GST and allowed purification of the fusion protein by affinity chromatography on glutathione agarose.

-61-

Optimal levels of GST-Vpu expression were obtained by growing the cultures at 30°C to a cell density of approximately 250-300 Klett units and inducing with low levels of IPTG (0.01mM). To purify the GST-Vpu, a combined cellular fraction containing the cell debris and plasma membrane was prepared by lysozyme treatment of the induced cells followed by a low-speed centrifugation. Approximately 50% of the GST-Vpu protein could be solubilised from this fraction using the zwitterionic detergent CHAPS. Affinity chromatography using glutathione-agarose beads was used to enrich the fusion protein and thrombin was used to cleave the fusion protein at the high affinity thrombin site between the fusion partners, liberating Vpu (Fig. 2A). In fractions eluted from the anion exchange column Vpu was the major protein visible on silver stained gels (Fig. 2B, lane 1). Finally, Vpu was purified to apparent homogeneity on an immunoaffinity column (Fig. 2B, lane 2). The N-terminal amino acid sequence of the protein band (excised from SDS-PAGE gels) corresponding to the immunodetected protein confirmed its identity as Vpu.

15

Example 8. Reconstitution of Vpu in Phospholipid Vesicles.

Proteoliposomes containing Vpu were prepared by the detergent dilution method (New, 1990). A mixture of lipids (PE:PC:PS; 5:3:2; 1mg total lipid) dissolved in chloroform was dried under a stream of nitrogen gas and resuspended in 0.1 ml of potassium phosphate buffer (50mM pH 7.4) containing DTT (1mM). A 25µl aliquot containing purified Vpu was added, followed by octylglucoside to a final concentration of 1.25 % (wt/vol). This mixture was subject to three rounds of freezing in liquid nitrogen, thawing and sonication in a bath type sonicator (20-30 sec) and was then rapidly diluted into 200 volumes of the potassium phosphate buffer. Proteoliposomes were collected by centrifugation at 400,000g for 1hr and resuspended in approximately 150µl of phosphate buffer.

20

25

Example 9. Assaying Vpu Ion Channel Activity

Purified Vpu was tested for its ability to induce channel activity in planar lipid bilayers using standard techniques as described elsewhere (Miller, 1986; and Piller et al, 1996). The solutions in the CIS and TRANS chambers were separated by a Delrin™ plastic wall containing a small circular hole of approximately 100µm

30

-62-

diameter across which a lipid bilayer was painted so as to form a high resistance electrical seal. Bilayers were painted from a mixture (8:2) of palmitoyl-oleoyl-phosphatidyl-ethanolamine and palmitoyl-oleoylphosphatidyl-choline (Avanti Polar Lipids, Alabaster, Alabama) in n-decane. The solutions in the two chambers
5 contained MES buffer (10mM, pH 6.0) to which various NaCl or KCl concentrations were added. Currents were recorded with an Axopatch™ 200 amplifier. The electrical potential between the two chambers could be manipulated between +/-200mV (TRANS relative to grounded CIS). Aliquots containing Vpu were added to the CIS chamber either as a detergent solution or after incorporation of the protein into
10 phospholipid vesicles. The chamber was stirred until currents were observed.

Example 10. Vpu Forms Ion Channels in Lipid Bilayers.

To assay for ion-channel formation by Vpu, reconstitution into planar lipid bilayers was performed. When samples (containing between 7 and 70ng of protein) of purified recombinant Vpu were added to the 1ml of buffer in the CIS chamber of the
15 bilayer apparatus, current fluctuations were detected after periods of stirring that varied from 2 to 30 min (Fig. 3). This time taken to observe channel activity approximately correlated with the amount of protein added to the chamber. No channels were detected when control buffer aliquots or control lipid vesicles were added to the CIS chamber. In those control experiments the chambers could be stirred
20 for more than an hour without appearance of channel activity.

Example 11. Properties of The Vpu Channels.

Channel activity was observed in over 40 individual experiments with Vpu samples prepared from five independent purifications. In different experiments, the amplitude of the currents varied over a large range and, again, seemed to
25 approximately correlate with the amount of protein added. The smallest and largest channels measured had conductances of 14 pS and 280 pS, respectively. The channels were consistently smaller when lipid vesicles containing Vpu were prepared and fused to the bilayer rather than when purified protein in detergent solution was added. This may be because the former method included treatment with high concentrations
30 of detergent and a dilution step that may have favoured the breakdown of large aggregates into monomers.

-63-

The relationship between current amplitude and voltage was linear and the reversal potential in solutions containing a ten-fold gradient of NaCl (500mM CIS; 50mM TRANS) was +30mV (Fig. 3B). A similar reversal potential was obtained when solutions contained KCl instead of NaCl. In 5 experiments with either NaCl or KCl in the solutions on either side of the membrane, the average reversal potential was 31.0 ± 1.2 mV (\pm -SEM). This is more negative than expected for a channel selectively permeable for the cations alone. Using ion activities in the Goldman-Hodgkin-Katz equation gives a P_{Na}/P_{Cl} ratio of about 5.5 indicating that the channels are also permeable to chloride ions. An attempt was made to reduce the anion current by substituting phosphate for chloride ions. When a Na-phosphate gradient (150mM Na^+ & 100mM phosphate CIS; 15mM Na^+ & 10mM phosphate TRANS, pH 6.8) was used instead of the NaCl gradient, the reversal potential was 37.1 ± 0.2 (\pm -SEM, $n=2$) again indicating a cation/anion permeability ratio of about 5. (For calculations involving the phosphate solutions, the summed activities of the mono and bivalent anions were used and it was assumed that the two species were equally permeable). The current-voltage curve now exhibited rectification that was not seen in the NaCl solutions. It can be concluded that the channels formed by Vpu are equally permeable to Na^+ and K^+ and are also permeable, though to a lesser extent, to chloride as well as phosphate ions.

Example 12. Bacterial Bio-Assay for Screening Potential Ion Channel-Blocking Drugs.

This bio-assay is based on the observation that expression of Vpu in *E. coli* results in an active Vpu channel located in the plasmalemma that dissipates the transmembrane sodium gradient. As a consequence of this Vpu channel activity, metabolites whose accumulation within the cells is mediated by a sodium dependent co-transporter (for example proline or adenine) leak out of the cell faster than they can be synthesised so that the metabolites' intracellular levels become limiting for growth of the cell. Thereby, an *E. coli* cell expressing Vpu is unable to grow in minimal drop-out media lacking adenine or proline. However, in the presence of a drug that blocks the Vpu channel, the cell is once again able to re-establish its transmembrane sodium gradient - due to the action of other ion pumps in the membrane - and the leakage of metabolites is prevented enabling growth.

-64-

Experiments to demonstrate that Vpu can form sodium channels in the plasma membrane of *E. coli* were performed as follows.

To express unfused Vpu in *E. coli*, the vpu open-reading frame was cloned into the plasmid pPL451 to create the recombinant plasmid pPL-Vpu (Fig. 1b). In this vector the strong P_L and P_R lambda promoters are used to drive expression of Vpu under control of the temperature sensitive c1857 repressor, such that when grown at 30°C expression is tightly repressed and can be induced by raising the temperature to between 37°C and 42°C. On agar plates, cells containing pPL-Vpu grew when incubated at 30°C and 37°C but not at 42°C, while control strains grew well at 42°C. Liquid cultures of cells containing pPL-Vpu were grown at 30°C to $OD_{600}=0.84$ then moved to grow at 42°C for two hours (the final cell density was $OD_{600}=0.75$). The plasma membrane fraction was prepared and western blotting, using an antibody that specifically binds to the C-terminus of Vpu, detected a single band at approximately 16kDa, indicating that Vpu was expressed and associated with the membranes (Fig. 2A, lane 5).

Example 13. Cross-Feeding Experiments Reveal That Proline Leaks Out of Cells Expressing Vpu.

Uptake of proline by *E. coli* is well characterised and active transport of the amino acid into the cells is known to use the sodium gradient as the energy source (Yamato *et al.*, 1994). To detect whether proline leakage occurs, the following cross-feeding assay was used: A lawn of an *E. coli* strain auxotrophic for proline and methionine ($Met^- Pro^-$), was seeded and poured as a soft agar overlay on minimal drop-out media plates lacking proline but containing methionine. Sterile porous filter discs were inoculated with a $Met^+ Pro^+$ strain (XL-1 blue) containing either the pPL451 control plasmid or pPL-Vpu and placed onto the soft agar. The plates were then incubated at 37°C or 30°C for two days. After than time a halo growth of the $Met^- Pro^-$ strain was clearly visible surrounding the disc inoculated with the cells containing pPL-Vpu incubated at 37°C (Fig. 4A). This growth can only be due to the leakage of proline from the Vpu-expressing cells on the disc. No such leakage was apparent from the control strain at 37°C nor around either strain on plates grown at 30°C (Fig. 4B).

-65-

In contrast to proline transport, the *E. coli* methionine permease is known to belong to the ABC transporter family (Rosen, 1987) and hence be energised by ATP. Identical crossfeeding experiments to those described above were set up except that the Met⁻ Pro⁻ strain was spread on minimal drop-out plates lacking methionine but
5 containing proline. No growth of this strain was evident around any of the discs (Fig. 4C), indicating that methionine was not leaking out of the XL-1 blue cells even when Vpu was being expressed.

Example 14. E.Coli Cells Expressing Vpu Require Adenine in the External Medium for Growth.

10 It was observed that, due to an uncharacterised mutation in the adenine synthesis pathway, growth of *E. coli* cells of the XL1-blue strain expressing Vpu at 37°C was dependant on the presence of adenine in the medium. This allowed the development of an even simpler bioassay for Vpu ion-channel activity than the proline cross-feeding assay described above: A lawn of XL1-blue cells containing the
15 pPL-Vpu plasmid is seeded onto an agarose plate lacking adenine in the medium, small aliquots of drugs to be tested for inhibition of the Vpu channel are spotted onto the agarose in discrete locations and the plates are incubated at 37°C for a suitable period of time (12-36 hours). Halos of growth around a particular drug application site indicate that the drug has inhibited expression of the Vpu ion channel activity that
20 prevents growth in the absence of the drug. (Figure 5).

Example 15

Assay of Compounds in Planar Lipid Bilayers for Vpu Channel Blocking Activity

25 Compounds were characterized for their ability to block Vpu ion channel activity reconstituted into planar lipid bilayers. Vpu N-terminal peptide (residues 1-32) dissolved in trifluoroethanol was added to the CIS chamber of the bilayer apparatus and the solutions was stirred until ion currents were observed, indicating incorporation of one or more Vpu ion channels into the bilayer. After recording the
30 channel activity for a few minutes, drugs were added to the solutions in the CIS and TRANS chambers – with stirring – to a final concentration of 100µM. Channel activity was then recorded for at least a further three minutes and the effect of drug addition on ion current was determined by comparing the channel activity before and

-66-

after drug addition. For each experiment, drug effect was classified into four categories: "Strong block", if current was inhibited approximately 90-100%; "weak block", approx. 50-90% inhibition; "partial block", <50%; and "no effect".

Experiments were disregarded if currents larger than ± 50 pA were generated after addition of Vpu N-peptide because in such cases it is possible that non-native peptide aggregates contribute to bilayer breakdown. Such aggregates, by virtue of their disorganized structure may not be specifically blocked by the drugs at the concentrations tested.

Table 3 summarises the results of the bilayer experiments. A novel outcome of these experiments was the strong blocking of Vpu channels observed with Phenamil. Phenamil has a phenyl group derivative at the guanidine group of amiloride. Amiloride itself is not a blocker of Vpu, whereas addition of the hexamethylene group at the 5- position of the pyrazine ring created a structure (HMA) that blocks the channel at concentrations as low as 25 μ M. These new results with Phenamil, however, now show that a bulky hydrophobic derivative at the opposite end of the molecule can also turn amiloride into an effective Vpu channel blocker. Interestingly, benzamil, with a very similar structure was much less effective at blocking the Vpu channel.

20

Table 3: Summary of Compounds Inhibiting the Vpu Ion Channel in Bilayers

Compound	No. of Expts.	Results
Phenamil	3	3x Strong block
MIA	2	1x Strong block; 1x weak
Benzamil	10	3x partial block; 7x no effect
EIPA	3	3x weak block;
HMA	1	1x Strong block;
(5-Phenyl-penta-2,4-dienoyl)guanidine	6	6x strong block
6-methoxy-2-naphthoylguanidine	5	5x strong block
(2-Chlorocinnamoyl)guanidine	6	4x strong; 2x partial blocks
3-(trifluoromethyl)cinnamoylguanidine	5	4x strong blocks; 1x no effect

-67-

N-{5-[3-(5-Guanidino-pentyloxymethyl)-benzyloxy]-pentyl}-guanidine	4	3x strong block; 1x no effect
4-phenylbenzoylguanidine	3	3x strong block
3-methylcinnamoylguanidine	4	2x strong block; 2x partial
(3-Chlorocinnamoyl)guanidine	4	2x strong block; 2x partial
N-(3-phenylpropanoyl)-N'-phenylguanidine	1	1x strong blocks
(3-Bromocinnamoyl)guanidine	3	3x partial-strong block
5-tert-butylamino-amiloride	3	3x partial block
N-amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide	3	3x partial block
3-methoxy-HMA	3	3x partial block
5-(N-Methyl-N-isobutyl)amiloride	1	1x partial block
5-(N-Ethyl-N-isopropyl)amiloride	1	1x partial block
2-naphthoylguanidine	7	7x weak block
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	7	7x weak block
cinnamoylguanidine	3	3x weak block
(5-Phenyl-penta-2,4-dienoyl)guanidine	6	6x strong block

Example 16 Compound Screening using the Bacterial Bio-Assay for the Vpu protein.

- 5 The halos of growth around the site of application of particular drugs – as described in example 14– were given a score between zero and six reflecting the size and density of the zone of bacterial cell growth. Scores greater than 3 represent strong inhibition of the Vpu protein; scores between 1.5 and 3 represent moderate inhibition and scores between 0.01 and 1.5 represent fair inhibition.

10

Table 4 lists the scores for inhibition of Vpu protein in the bacterial bio-assay.

Table 4

Compound	Vpu Inhibition (score / # of times tested)
(3-Chlorocinnamoyl)guanidine	4.38/4
(3-Bromocinnamoyl)guanidine	4.3/24
(2-Chlorocinnamoyl)guanidine	4.0/4
(2-Bromocinnamoyl)guanidine	3.7/2
3-(trifluoromethyl)cinnamoylguanidine	3.7/2

-68-

5-bromo-2-fluorocinnamoylguanidine	3.5/2
3-methylcinnamoylguanidine	3.4/2
2-methylcinnamoylguanidine	3.1/2
2,3-dimethylcinnamoylguanidine	3.1/2
cinnamoylguanidine	2.96/12
6-methoxy-2-naphthoylguanidine	2.9/4
trans-3-(1-naphthyl)acryloylguanidine	2.9/3
3,4-dichlorocinnamoylguanidine	2.9/3
2,6-dichlorocinnamoylguanidine	2.88/2
4-phenylbenzoylguanidine	2.75/5
2-ethylcinnamoylguanidine	2.75/2
(4-Chlorocinnamoyl)guanidine	2.7/5
2-naphthoylguanidine	2.7/11
2,5-dimethylcinnamoylguanidine	2.69/2
3-isopropylcinnamoylguanidine hydrochloride	2.6/2
(5-Phenyl-penta-2,4-dienoyl)guanidine	2.56/2
3-phenylcinnamoylguanidine	2.54/3
(4-Bromocinnamoyl)guanidine	2.5/4
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	2.5/2
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	2.5/2
3-(trifluoromethoxy)cinnamoylguanidine	2.44/2
2-(trifluoromethyl)cinnamoylguanidine	2.4/2
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	2.25/3
2-ethoxycinnamoylguanidine	2.25/2
N-(3-phenylpropanoyl)-N'-phenylguanidine	2.21/3
4-(trifluoromethyl)cinnamoylguanidine	2.2/2
(4-Methoxycinnamoyl)guanidine	2.13/3
2-t-butylcinnamoylguanidine	2.13/2
4-methylcinnamoylguanidine	2.1/2
2-fluorocinnamoylguanidine	2.1/2
2-phenylcinnamoylguanidine	2.1/2
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine	2.06/2
3-t-butylcinnamoylguanidine	2.06/2
3,4-difluorocinnamoylguanidine	2.06/2
5-(N,N-hexamethylene)amiloride	1.9/31
3-fluorocinnamoylguanidine	1.9/2
5-bromo-2-methoxycinnamoylguanidine	1.9/2
3-ethoxycinnamoylguanidine	1.9/2
3,4-(methylenedioxy)cinnamoylguanidine	1.88/2
(2-Methoxycinnamoyl)guanidine	1.7/4
2'4 DichloroBenzamil HCl	1.7/2
2,3,5,6,-tetramethylcinnamoylguanidine	1.6/2
3-(2-naphthyl)acryloylguanidine	1.56/2
2-(1-naphthyl)acetoyleguanidine	1.56/2
2,3-difluorocinnamoylguanidine	1.56/2
(3-Methoxycinnamoyl)guanidine	1.52/6
4-isopropylcinnamoylguanidine	1.4/2
2,4,6-trimethylcinnamoylguanidine	1.4/2

-69-

N-(cinnamoyl)-N'-phenylguanidine	1.25/3
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	1.2/2
2-(2-naphthyl)acetoylguanidine	1.19/2
(4-Hydroxycinnamoyl)guanidine	1.1/2
4-phenylcinnamoylguanidine	1.1/2
4-fluorocinnamoylguanidine	1.1/2
N,N'-bis-(cinnamoyl)-N''-phenylguanidine	0.94/2
(2-Furanacryloyl)guanidine	0.94/2
Phenamil methanesulfonate salt	0.9/5
Benzamil hydrochloride	0.9/3
(3-Nitrocinnamoyl)guanidine	0.9/1
Benzyoylguanidine	0.88/2
(4-Phenoxybenzoyl)guanidine	0.81/2
3-(trans-hept-1-en-1-yl)cinnamoylguanidine	0.81/2
5-(N-Methyl-N-isobutyl)amiloride	0.8/2
2-cyclohexylcinnamoylguanidine	0.8/2
4-ethoxycinnamoylguanidine	0.69/2
2,4-dichlorocinnamoylguanidine	0.63/2
5-(N-Ethyl-N-isopropyl)amiloride	0.6/3
N-amidino-3-amino-5-hexamethyleneimino-6-phenyl- 2-pyrazinecarboxamide	0.6/2
(a-Methylcinnamoyl)guanidine	0.6/2
cinnamoylguanidine hydrochloride	0.6/2
[(4-Chlorophenoxy-acetyl)guanidine	0.56/2
N-amidino-3-amino-5-phenyl-6-chloro-2- pyrazinecarboxamide	0.5/11
5-(4-fluorophenyl)amiloride	0.4/6
(trans-2-Phenylcyclopropanecarbonyl)guanidine	0.4/2
(2-Nitrocinnamoyl)guanidine	0.4/2
trans-3-Furanacryoylguanidine	0.38/2
1-naphthoylguanidine	0.3/2
5-tert-butylamino-amiloride	0.2/7
3-methoxy-HMA	0.2/4
(3-phenylpropanoyl)guanidine	0.2/4
4-t-butylcinnamoylguanidine	0.19/2
5-(N,N-Dimethyl)amiloride hydrochloride	0.1/2
N,N'-Bis(3-phenylpropanoyl)guanidine	0.1/2
N-Benzoyl-N'-cinnamoylguanidine	0.06/2
1-bromo-2-naphthoylguanidine	0.06/2

Example 17. Effect of Compounds on HIV Replication in Human Monocytes and Macrophages.

Human monocytes were isolated from peripheral blood and cultured either for
5 24hr (one day old monocytes) or for 7 days to allow differentiation into monocyte
derived macrophages (MDM). These cells were then exposed to cell-free preparations

-70-

of HIV isolates and allowed to absorb for 2hr before complete aspiration of the medium, washing once with virus-free medium and resuspension in fresh medium.

The cells were exposed to various concentration of compound either 24 hr prior to infection or after infection. Subsequent HIV replication, at various times after

- 5 infection, was compared in cells exposed to drugs and in cells not exposed to drugs (controls). The progression and extent of viral replication was assayed using either an HIV DNA PCR method (Fear et al, 1998) or an ELISA method to quantitate p24 in culture supernatants (Kelly et al, 1998).

10 Table 5 provides examples of results obtained using this assay and test antiviral compounds.

Table 5

Compound	Drug Conc. μ M	Percent of Positive Control
None – positive control		100%
4-phenylbenzoylguanidine	10	26
	5	4
	2.5	9
	1.	216
	0.625	10
None – positive control		100%
(3-Bromocinnamoyl)guanidine	10	3
	5	1
	2.5	9
	1.25	59
	0.625	116
None – positive control		100%
3-(trifluoro-methyl)cinnamoylguanidine	10	11
	5	8
	2.5	25
	1.25	27
	0.625	38
None – positive control		100%
5-(N,N-hexamethylene)amiloride	10	6
	5	21
	2.5	50
	1.25	19
	0.625	30

Example 18. SARS Coronavirus.**SARS E protein forms an ion channel****Peptide Synthesis**

A peptide corresponding to the full-length SARS-CoV (isolate Tor2 and
 5 Urbani) E protein (MYSFVSEETGTLIVNSVLLFLAFVVFLVTLAILTALRLCA
 YCCNIVNVSLVKPTVYVYSRVKNLNSSEGVPDLLV) (SEQ ID NO: 3) and a
 second peptide comprising the first 40 amino acids of the full length E protein which
 correspond to the transmembrane domain (MYSFVSEETGTLIVNSVLLFLAFVVF
 10 LLVTLAILTALRLC) (SEQ ID NO: 4) were synthesized manually using Fmoc
 chemistry and solid phase peptide synthesis. The synthesis was done at the
 Biomolecular Resource Facility (John Curtin School of Medical Research, ANU,
 Australia) using a Symphony^R Peptide Synthesiser from Protein Technologies
 Inc.(Tucson, AZ, USA) according to the manufacturers instructions.

Example 19. Peptide purification

15 Mass spectral analysis of the synthetic peptide revealed that the preparation
 contained significant amounts of material with lower m/z ratio than expected for the
 full-length product. The majority of these are presumably truncated peptides
 generated during the peptide synthesis process. To enrich the full-length E protein,
 the following procedure was used, which relies on differential solubility of the
 20 smaller molecules and full-length peptide. The crude preparation was suspended at
 12 mg/ml in 70% CH₃CN, 0.1 %TFA and vortexed for 10 minutes. This suspension
 was centrifuged at 10,000g for 10 minutes at 20°C. The supernatant was discarded
 and the insoluble fractions was extracted with 70% CH₃CN, 0.1% TFA, as above,
 two more times. The insoluble material containing the E protein was dried using
 25 Speedvac and the weight of the final product was used to calculate the yield. The
 purified peptide was analysed by Bruker Omnisflex MALDI-TOF mass spectrometry
 in HABA matrix at 2.5mg/ml in methanol at a 1:1 ratio and spectra were obtained in
 the positive linear mode. A clear peak at m/z ratio of 8,360.1 was seen as expected
 for the calculated molecular weight of full-length E protein and 4422.3 for the N-
 30 terminal E protein.

-72-

Example 20. Planar Lipid Bilayers

The SARS virus E protein was resuspended at 1mg/ml in 2,2,2-trifluoroethanol. The SARS virus E protein's ability to form ion channels was tested on a Warner (Warner instruments, Inc. 1125 Dixwell Avenue, Hamden, CT 06514) bilayer rig as follows; A lipid mix of 3:1:1, 1-Palmitoyl-2-oleoyl phosphatidyl Ethanolamine: 1-Palmitoyl-2-oleoyl phosphatidyl Serine: 1-Palmitoyl-2-oleoyl phosphatidyl choline in CHCl_3 was dried under N_2 gas and resuspended to 50mg/ml in n-decane. Bilayers were painted across a circular hole of approximately 100 μm diameter in a DelrinTM cup separating aqueous solution in the CIS and TRANS chambers. The CIS chamber contained a solution of 500mM NaCl or KCl, in a 5mM HEPES buffer pH 7.2, the TRANS chamber contained a solution of 50mM NaCl or KCl, in a 5mM HEPES buffer pH 7.2. Silver electrodes coated in chloride with 2% agarose bridges are placed in the CIS and TRANS chamber solutions. The SARS E protein full-length or N-terminal peptides (3-10ug) were added to the CIS chamber, which was stirred until channel activity was detected. The CIS chamber was earthed and the TRANS chamber was held at various holding potentials ranging between +100 to -100mV. Currents were recorded using a Warner model BD-525D amplifier, filtered at 1kHz, sampling at 5 kHz and digitally recorded on the hard disk of a PC using software developed in house.

Drugs to be tested for their ability to inhibit SARS E protein ion channel activity were made up at 50mM in a solution of 50% DMSO: 50% methanol. For experiments testing the ability of compounds to inhibit E protein ion channel activity, 100 μM to 400 μM of compound was added to the CIS chamber while stirring for 30 seconds. Bilayer currents were recorded before channel activity, during channel activity and after the addition of the drug.

Among the compounds tested was cinnamoylguanidine (Bit036), a compound which was shown in earlier experiments to be antiviral and to inhibit ion channel proteins from other viruses.

Example 20.1. Polyacrylamide gel electrophoresis

Purified E protein was dissolved to 1 mg/ml, 5 mg/ml and 10 mg/ml in, 6 M Urea, 10% Glycerol, 5% SDS, 500 mM DTT, 0.002% Bromophenol Blue, 62.5 mM Tris HCl (pH 8.3). Peptides in solutions were heated at 100°C for 20 minutes before

-73-

30 μ L samples were run on stacking gel 4-20% (Gradipore). SeeBlue® pre-stained standard (Invitrogen) was used for molecular weight markers.

Example 20.2 Results

To test if the SARS E protein forms ion channels the purified synthetic peptide was reconstituted into planar lipid bilayers (21). Typically, 3 μ g of SARS full-length E protein was added to the CIS chamber, while stirring. This CIS chamber contained 500 mM NaCl and the TRANS chamber contained 50 mM NaCl. In 60 experiments, ion currents due to SARS E protein ion channel activity were observed after about 5 –15 minutes of stirring. Activity was detected more rapidly and reliably with a holding potential of approximately –100mV across the bilayer. Currents recorded at –100mV, (A) and at –60mV (B) in one of these experiments are shown in Figure 6. In that experiment the reversal potential was about +48mV and the channel conductances were calculated to be 52pS and 26pS, respectively. This indicates that the current-voltage (IV) relationship is not linear. In ten other experiments, where no protein was added to the CIS chamber, no ion channel activity was detected, even after recording for over 1 hour.

Figure 7a shows typical current traces recorded over a range of potentials in NaCl solutions. In that experiment the direction of current flow reversed at +48mV (Fig 7b). The IV curve shows that at the lower voltages the average current flow across the bilayer is small but at higher potentials there is an increase in average current across the bilayer, resulting in a non-linear IV relationship. In seven independent experiments, the average reversal potential was $+48.3 \pm 2.3$ mV (mean \pm 1SEM), indicating that the channels were about 37 times more permeable to Na⁺ ion than to Cl[–] ions. The reversal potential is close to the Na⁺ equilibrium potential (+53mV), therefore the channel is selective for Na⁺ ions. For these 7 experiments the channel conductance varied between 95-164 pS; the average conductance was 130 ± 13 pS.

SARS E protein ion channel is slightly less selectivity for K⁺ ions than Na⁺ ions. Figure 8b shows recording of currents in KCl solutions at a range of potentials. In this experiment the currents reversed at +31 mV. In seven similar experiments E protein ion channel average reversal potential was $+34.5 \pm 2.5$ mV. Therefore the SARS E protein ion channel is about 7.2 times more permeable to K⁺ ions than Cl[–]

-74-

ions. In seven experiments, the channel conductance varied ranging between 24-166 pS, the average conductance was 83.4 ± 26 pS.

Similar results were obtained with a second synthetic peptide, which corresponded to the first forty N-terminal amino acids of the SARS E protein "N-terminal peptide"

- 5 (21). The average reversal potential in NaCl solution in four experiments was $+46.3 \pm 2.5$ mV, indicating that the ion channel formed by N-terminal peptide is about 25 times more permeable to Na^+ ion than to Cl^- ions. The SARS E protein N-terminal peptide was sufficient for the formation of ion channels with properties like those of the full length SARS E protein. Therefore, the selectivity filter for the SARS E
10 protein is most likely contained within the first forty amino acids of the N-terminal.

SARS E protein N-terminal peptide also formed ion channels in KCl solution that were similarly selective for K^+ ions compared to the full-length E protein. In five independent experiments the average channel reversal potential was $+39.5 \pm 3.6$ mV, therefore the channel is about 11 times more permeable to K^+ ions than Cl^- ions.

- 15 SDS-PAGE of the purified full-length E protein peptide showed bands corresponding to the full-length E protein (Data not shown). Larger bands of varying size up to about 20 kDa were detected, suggesting that SARS E protein may form homo-oligomers.

20 **Example 21. SARS E protein ion channel is blocked by cinnamoylguanidine and other compounds**

- E protein ion channel activity in NaCl solutions was significantly reduced ($p \geq 0.01$, $n=6$ experiments) by addition of 100 to 200 μM cinnamoylguanidine to the CIS chamber. The average current across the bilayer was reduced to baseline by
25 100 μM cinnamoylguanidine. In experiments when E protein ion channels had higher conductance, 100 to 200 μM cinnamoylguanidine reduced the average current across the bilayer about 4 fold. Similarly, in four other experiments, 100 to 200 μM cinnamoylguanidine blocked channels formed by full-length E protein in KCl solutions. In two additional experiments, the SARS E protein N-terminal peptide was
30 blocked by 100 to 200 μM cinnamoylguanidine, demonstrating that the cinnamoylguanidine drug-binding site is located within the first forty amino acids of

-75-

the E protein N-terminal domain. Other compounds tested in bilayers for their effect on the SARS E protein are shown in below in Table 6.

Table 6

Compound	% Reduction of average current by 100 μ M
5-(N,N-hexamethylene)amiloride	91 \pm 7
6-methoxy-2-naphthoylguanidine	92 \pm 16
2'4 DichloroBenzamil HCl	78 \pm 0
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	88 \pm 6
(3-Bromocinnamoyl)guanidine	87 \pm 11
(2-Bromocinnamoyl)guanidine	88 \pm 6
trans-3-(1-naphthyl)acryloylguanidine	66 \pm 2

Example 21.1 Results and Discussion.

We have shown that SARS E protein can form ion channels in lipid bilayer membranes. The ion currents reversed at positive potentials, which demonstrates that E protein ion channels are selective for monovalent cations over monovalent anions. E protein ion channels were about 37 times more selective for Na⁺ ions over Cl⁻ ions and about 7.2 times more selective for K⁺ ions over Cl⁻ ions. In over 60 experiments the Na⁺ conductance of the E protein ion channel varied from as low as 26 pS to as high as 164 pS. SDS-PAGE showed that the E protein forms homo-oligomers, and we surmised that the larger conductances were probably due to aggregation of the E protein peptide leading to larger ion channels or the synchronous opening of many ion channels. Single channel currents were observed in several experiments and from these the channel conductance was calculated to be voltage dependent.

The first 40 amino acids of the N-terminal which contains the hydrophobic domain of the SARS virus E protein is sufficient for the formation of ion channels on planar lipid bilayers. The N-terminal E protein ion channel has the same selectivity and conductance as the full-length E protein ion channel.

The SARS virus full length E protein ion channel activity and N-terminal domain E protein ion channel activity on planar lipid bilayers in NaCl and KCl solutions was inhibited by addition of between 100 μ M to 200 μ M cinnamoylguanidine to the CIS chamber. Inhibition or partial inhibition of the E

-76-

protein ion channel activity by cinnamoylguanidine has been observed in seven independent experiments in NaCl solution and four independent experiments in KCl solution.

All known coronaviruses encode an E protein with a hydrophobic N-terminus transmembrane domain therefore all coronaviruses E proteins could form ion channels on planar lipid bilayers. This indicates that the E protein could be a suitable target for antiviral drugs and potentially stop the spread of coronavirus from infected host cells. Drugs that block the E protein ion channel could be effective antiviral therapy for the treatment of several significant human and veterinary coronavirus diseases including SARS and the common cold.

Example 22. Bacterial Bio-Assay for Screening Potential SARS-CoV E protein Ion Channel-Blocking Drugs.

SARS-CoV E protein Ion Channel inhibits Bacterial Cell growth.

A bio-assay of SARS-CoV E protein function in bacterial cells was developed. A synthetic cDNA fragment encoding SARS-CoV E protein was cloned into the expression plasmid pPL451, creating a vector in which E protein expression is temperature inducible, as described in Example 4. Inhibition of the growth of E.coli cells expressing E protein at 37°C was observed as an indicator of p7 ion channel function dissipating the normal Na⁺ gradient maintained by the bacterial cells.

Example 23. Compound Screening using the Bacterial Bio-Assay for SARS coronavirus E protein.

The halos of growth around the site of application of particular drugs – as described in example 14 – were scored as described in example 15.

Table 7 lists the scores for inhibition of SARS-CoV E protein in the bacterial bio-assay.

Table 7

Compound	SARS E protein Inhibition (score / # of times tested)
----------	--

-77-

2,3-difluorocinnamoylguanidine	4.50 / 1
3,4-dichlorocinnamoylguanidine	4.15 / 2
4-t-butylcinnamoylguanidine	4.00 / 1
3-(2-naphthyl)acryloylguanidine	3.88 / 1
(3-Chlorocinnamoyl)guanidine	3.87 / 3
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	3.75 / 1
2,5-dimethylcinnamoylguanidine	3.63 / 1
trans-3-(1-naphthyl)acryloylguanidine	3.38 / 2
4-isopropylcinnamoylguanidine	3.16 / 2
(3-Bromocinnamoyl)guanidine	3.15 / 27
6-methoxy-2-naphthoylguanidine	3.13 / 3
5-(N-Methyl-N-isobutyl)amiloride	3.13 / 2
3-phenylcinnamoylguanidine	3.13 / 1
(2-Chlorocinnamoyl)guanidine	3.1 / 3
2'4 DichloroBenzamil HCl	3.00 / 2
4-phenylcinnamoylguanidine	2.75 / 2
4-(trifluoromethyl)cinnamoylguanidine	2.75 / 1
3-(trifluoromethoxy)cinnamoylguanidine	2.71 / 1
3-(trifluoromethyl)cinnamoylguanidine	2.67 / 1
2-ethoxycinnamoylguanidine	2.57 / 1
cinnamoylguanidine hydrochloride	2.50 / 1
4-ethoxycinnamoylguanidine	2.48 / 2
(2-Bromocinnamoyl)guanidine	2.47 / 3
2,6-dichlorocinnamoylguanidine	2.25 / 1
3,4,5-trimethoxycinnamoylguanidine	2.25 / 1
5-tert-butylamino-amiloride	2.01 / 2
3-t-butylcinnamoylguanidine	2.00 / 1
5-bromo-2-fluorocinnamoylguanidine	2.00 / 1
(4-Chlorocinnamoyl)guanidine	1.94 / 2
2-t-butylcinnamoylguanidine	1.86 / 1
2-cyclohexylcinnamoylguanidine	1.83 / 1
6-Iodoamiloride	1.75 / 2
3-(trans-hept-1-en-1-yl)cinnamoylguanidine	1.71 / 1
(4-Bromocinnamoyl)guanidine	1.69 / 2
(4-Hydroxycinnamoyl)guanidine	1.63 / 2
N-(3-phenylpropanoyl)-N'-phenylguanidine	1.57 / 2
(3-Nitrocinnamoyl)guanidine	1.51 / 2
3-fluorocinnamoylguanidine	1.50 / 1
2-(1-naphthyl)acetoyleguanidine	1.50 / 1
2-ethylcinnamoylguanidine	1.50 / 1
5-(N,N-Dimethyl)amiloride hydrochloride	1.38 / 2
2-naphthoylguanidine	1.38 / 2
5-(4-fluorophenyl)amiloride	1.38 / 1
2-(trifluoromethyl)cinnamoylguanidine	1.38 / 1
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine	1.35 / 3
(trans-2-Phenylcyclopropanecarbonyl)guanidine	1.34 / 3
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	1.33 / 3
1-naphthoylguanidine	1.32 / 3

-78-

Benzamil hydrochloride	1.32 / 2
3-methoxy-HMA	1.25 / 1
4-methylcinnamoylguanidine	1.25 / 1
4-fluorocinnamoylguanidine	1.25 / 1
3,4-(methylenedioxy)cinnamoylguanidine	1.25 / 1
5-(N,N-hexamethylene)amiloride	1.2 / 3
N-(cinnamoyl)-N'-phenylguanidine	1.19 / 2
5-(N-Ethyl-N-isopropyl)amiloride	1.07 / 2
3-methylcinnamoylguanidine	1.00 / 1
2-methylcinnamoylguanidine	1.00 / 1
2,3,5,6,-tetramethylcinnamoylguanidine	1.00 / 1
trans-3-Furanacryoylguanidine	0.88 / 2
(4-Methoxycinnamoyl)guanidine	0.88 / 2
(2-Furanacryloyl)guanidine	0.82 / 2
(3-phenylpropanoyl)guanidine	0.73 / 5
2-(2-naphthyl)acetoyleguanidine	0.71 / 1
cinnamoylguanidine	0.69 / 3
(2-Methoxycinnamoyl)guanidine	0.69 / 2
[3-(3-Pyridyl)acryloyl]guanidine	0.67 / 3
4-phenylbenzoylguanidine	0.63 / 2
2,4-dichlorocinnamoylguanidine	0.63 / 2
(3-Methoxycinnamoyl)guanidine	0.63 / 2
2-fluorocinnamoylguanidine	0.63 / 1
(4-Phenoxybenzoyl)guanidine	0.57 / 2
(a-Methylcinnamoyl)guanidine	0.50 / 1
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	0.5 / 1
(5-Phenyl-penta-2,4-dienoyl)guanidine	0.44 / 2
(Quinoline-2-carbonyl)guanidine	0.41 / 1
(Phenylacetyl)guanidine	0.32 / 3
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide	0.25 / 2
6-bromo-2-naphthoylguanidine	0.25 / 1
1-bromo-2-naphthoylguanidine	0.25 / 1
2-chloro-6-fluorocinnamoylguanidine	0.25 / 1
[(4-Chlorophenoxy-acetyl]guanidine	0.19 / 2
Phenamil methanesulfonate salt	0.13 / 2
N-Benzoyl-N'-cinnamoylguanidine	0.13 / 2
N-(2-naphthoyl)-N'-phenylguanidine	0.07 / 2

Example 24. SARS Antiviral Assay for testing compounds against replication of SARS coronavirus (SARS-CoV).

- 5 Compounds were tested against SARS-CoV (Hong Kong strain) using virus plaque purified three times in Vero cells. Stock virus was generated by infecting Vero cells at MOI = 1x TCID₅₀ per 100 cells.

Example 24.1 Screening for anti-viral activity using the virus microtitre assay

-79-

Monolayers of Vero cells grown in 25cm² flasks were infected at a multiplicity of 1:50 and treated immediately post infection with compounds at two concentrations, 10uM and 2uM. A control infected monolayer remained untreated. Samples of culture media were taken at 48 hours post infection. Two aliquots from each of the samples (titrations 1 and 2) were serially log diluted and 12 replicates of log dilutions -4 to -7 added to cells in microtitre plates. Four days later, wells in the microtitre plates were scored for cytopathic effect (CPE) and the titration values calculated based on the number of CPE positive wells at the 4 dilutions. Control titres were 4.8 and 5.9 TCID₅₀ x 10⁶ (average 5.35 x 10⁶)

Example 25: Effect of compounds in SARS CoV antiviral assay:

Three selected compounds were tested for activity against SARS-CoV according to the method described in example 21. For trans-3-(1-naphthyl)acryloylguanidine and cinnamoylguanidine a decrease in virus titre of approximately 80% was observed at a concentration of 10uM and a reduction of approximately 50% was seen to persist at 2uM trans-3-(1-naphthyl)acryloylguanidine.

Table 8 provides Virus titration data presented as % of a control (SARS CoV grown for 48 hours in the absence of compounds).

Table 8

Compound		Average TCID ₅₀ (x 10 ⁶)	Titre (% control)
Name	Concentration (uM)		
cinnamoylguanidine	10	1.3	24
	2	4.4	82
trans-3-(1-naphthyl)acryloylguanidine	10	1.15	22
	2	2.45	46
6-methoxy-2-naphthoylguanidine	10	5.95	111
	2	6.35	118
Control	0	5.35	100

Example 26, Human 229E Coronavirus**Synthesis and Purification of a Peptide Corresponding to the 229E-E Protein**

A peptide corresponding to the full-length 229E-E (sequence:

MFLKLVDDHALVVNVLLWCVVLIVILLVCITIIKLIKLCFTCHMFCNRTVYGPIK

5 NVYHIYQSYMHIDPFPPKRVIDF (SEQ ID NO: 5); accession number NP_073554)

was synthesized manually using Fmoc chemistry and solid phase peptide synthesis.

The synthesis was done at the Biomolecular Resource Facility (John Curtin School of Medical Research, ANU, Australia) using a Symphony^R Peptide Synthesiser from

Protein Technologies Inc. (Woburn, MS, USA) according to the manufacturers

10 instructions to give C-terminal amides, the coupling was done with HBTU and hydroxybenzotriazole in N-methylpyrrolidone. Each of the synthesis cycles used double coupling and a 4-fold excess of the amino acids. Temporary α -N Fmoc-protecting groups were removed using 20% piperidine in DMF.

The crude synthetic peptide was purified using the ProteoPlusTM kit (Qbiogene inc. CA),

15 following manufactures instructions. Briefly, the peptides were diluted in loading buffer (60mM Tris-HCl pH 8.3, 6M urea, 5% SDS, 10% glycerol, 0.2% Bromophenol blue, \pm 100 mM β -mercaptoethanol) and run on 4-20% gradient polyacrylamide gels

(Gradipore, NSW, Australia) in tris-glycine electrophoresis buffer (25 mM Tris, 250mM glycine, 0.1% SDS). The peptides were stained with gel code blue (Promega, NSW)

20 and the bands corresponding to the full-length peptide were excised out of the gel.

The gel slice was transferred to the ProteoPLUSTM tube and filled with tris-glycine electrophoresis buffer. The tubes were emerged in tris-glycine electrophoresis buffer and subjected to 100 volts for approximately 1 hour. The polarity of the electric current was reversed for 1 minute to increase the amount of protein recovered. The

25 peptides were harvested and centrifuged at 13,000 rpm for 1 minute. The purified peptides were dried in a Speedvac and the weight of the final product was used to calculate the yield.

-81-

Example 27. 229E-E protein forms ion channels in planar lipid bilayers.

Lipid bilayer studies were performed as described elsewhere (Sunstrom, 1996; Miller, 1986). A lipid mixture of palmitoyl-oleoyl-phosphatidylethanolamine, palmitoyl-oleoyl-phosphatidylserine and palmitoyl-oleoyl-phosphatidylcholine (5:3:2) (Avanti Polar Lipids, Alabaster, Alabama) was used. The lipid mixture was painted onto an aperture of 150-200 μm in the wall of a 1 ml delrin cup. The aperture separates two chambers, cis and trans, both containing salt solutions at different concentrations. The cis chamber was connected to ground and the trans chamber to the input of an Axopatch 200 amplifier. Normally the cis chamber contained either 500 mM NaCl or 500mM KCl and the trans 50 mM NaCl or 50mM KCl. The bilayer formation was monitored electrically by the amplitude of the current pulse generated by a current ramp. The potentials were measured in the trans chamber with respect to the cis. The synthetic peptide was added to the cis chamber and stirred until channel activity was seen. The currents were filtered at 1000 Hz, digitized at 5000 Hz and stored on magnetic disk.

The 229E E synthetic peptide was dissolved in 2,2,2-trifluoroethanol (TFE) at 0.05mg/ml to 1 mg/ml. 10 μl of this was added to the cis chamber (1ml aqueous volume) of the bilayer apparatus, which was stirred via a magnetic "flea". Ionic currents, indicating channel activity in the bilayer, were typically detected within 15-30 min. After channels were detected the holding potential across the bilayer was varied between -100mV and +100mV to characterise the size and polarity of current flow and enable the reversal potential to be determined.

In 15 experiments where the cis chamber contained 500mM NaCl solution and the trans chamber contained 50 mM NaCl solution, the average reversal potential of the channel activity was calculated to be 22 ± 7 (SEM) mV. In 13 experiments where the cis chamber contained 500mM KCl solution and the trans chamber contained 50 mM KCl solution, the average reversal potential of the channel activity was calculated to be 38 ± 4 (SEM) mV. These results indicate that the 229E E protein forms cation selective ion channels that are slightly more selective for K^+ than for Na^+ ions.

-82-

Figure 9 shows examples of raw current data for the 229E E ion channel at various holding potentials (cis relative to trans) in asymmetrical KCl solutions (500/50 mM). The graph is a representative plot of average bilayer current (pA; y-axis) versus holding potential (mV; x-axis).

5

Example 28. Chemical compounds inhibit the ion channel activity of the 229E E protein synthetic peptide.

To test compounds for their ability to block or otherwise inhibit the ion channel formed by 229E E protein, small aliquots of solutions containing the compounds were added to the aqueous solutions bathing planar lipids in which the peptide channel activity had been reconstituted and the effect of the compound addition on the ionic currents was recorded and measured.

Compound stock solutions were typically prepared at 500 mM in DMSO. This solution was further diluted to 50 mM, or lower concentration in 50% DMSO/50% methanol and 2 μ l of the appropriately diluted compound was added to the cis and/or trans chambers to yield the desired final concentration.

In the example shown in Figure 10, addition of 100 μ M cinnamoylguanidine to the cis chamber greatly reduced current flow through the 229E E ion channel.

20

Example. 29. Bacterial Bio-Assay for Screening Potential 229E-CoV E protein Ion Channel-Blocking Drugs.

229E-CoV E-protein Ion Channel inhibits Bacterial Cell growth.

A bio-assay of 229E-CoV E-protein function in bacterial cells was developed. A synthetic cDNA fragment encoding 229E-CoV E-protein was cloned into the expression plasmid pPL451, creating a vector in which E protein expression is temperature inducible, as described in Example 4. Inhibition of the growth of E.coli cells expressing E protein at 37°C was observed as an indicator of p7 ion channel function dissipating the normal Na⁺ gradient maintained by the bacterial cells.

30

35

Example 30 Compound Screening using the Bacterial Bio-Assay for 229E-CoV E-protein.

The halos of growth around the site of application of particular drugs – as described in example 14 – were scored as described in example 15.

5

Table 9 list the scores for inhibition of 229E-CoV E-protein in the bacterial bio-assay.

Table 9

Compound	229E E protein Inhibition (score)
4-isopropylcinnamoylguanidine	4.9
3,4-dichlorocinnamoylguanidine	4.4
3-(trifluoromethoxy)cinnamoylguanidine	4.1
4-t-butylcinnamoylguanidine	4.0
3-isopropylcinnamoylguanidine hydrochloride	4.0
3-t-butylcinnamoylguanidine	3.9
2-t-butylcinnamoylguanidine	3.9
trans-3-(1-naphthyl)acryloylguanidine	3.7
5-bromo-2-methoxycinnamoylguanidine	3.6
2,3-difluorocinnamoylguanidine	3.3
3-(2-naphthyl)acryloylguanidine	3.0
2-phenylcinnamoylguanidine	3.0
3-phenylcinnamoylguanidine	2.9
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	2.4
4-phenylbenzoylguanidine	2.3
3-(trifluoromethyl)cinnamoylguanidine	2.3
(4-Phenoxybenzoyl)guanidine	2.3
4-(trifluoromethyl)cinnamoylguanidine	2.3
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	2.3
(4-Bromocinnamoyl)guanidine	2.0
5-(N,N-hexamethylene)amiloride	1.9
1-naphthoylguanidine	1.9
5-(4-fluorophenyl)amiloride	1.8
(5-Phenyl-penta-2,4-dienoyl)guanidine	1.8
(3-Bromocinnamoyl)guanidine	1.7
2,5-dimethylcinnamoylguanidine	1.6
2-(trifluoromethyl)cinnamoylguanidine	1.5
6-methoxy-2-naphthoylguanidine	1.4
(4-Chlorocinnamoyl)guanidine	1.4
(3-Methoxycinnamoyl)guanidine	1.4
5-bromo-2-fluorocinnamoylguanidine	1.4
5-(N,N-Dimethyl)amiloride hydrochloride	1.3

-84-

cinnamoylguanidine	1.3
(2-Methoxycinnamoyl)guanidine	1.1
(<i>a</i> -Methylcinnamoyl)guanidine	1.0
4-phenylcinnamoylguanidine	1.0
2,6-dichlorocinnamoylguanidine	1.0
(2-Bromocinnamoyl)guanidine	0.9
2,4,6-trimethylcinnamoylguanidine	0.9
(trans-2-Phenylcyclopropanecarbonyl)guanidine	0.8
(3-Chlorocinnamoyl)guanidine	0.8
2-(1-naphthyl)acetoylguanidine	0.8
2-ethylcinnamoylguanidine	0.8
2-cyclohexylcinnamoylguanidine	0.8
(4-Hydroxycinnamoyl)guanidine	0.6
2-ethoxycinnamoylguanidine	0.6
3-methylcinnamoylguanidine	0.5
2-methylcinnamoylguanidine	0.5
3-fluorocinnamoylguanidine	0.5
cinnamoylguanidine hydrochloride	0.5
2,3-dimethylcinnamoylguanidine	0.5
2-fluorocinnamoylguanidine	0.4
4-fluorocinnamoylguanidine	0.4
3,4-difluorocinnamoylguanidine	0.4
5-tert-butylamino-amiloride	0.3
2-naphthoylguanidine	0.3
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide	0.3
N,N'-Bis(3-phenylpropanoyl)guanidine	0.3
4-methylcinnamoylguanidine	0.3
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	0.3
2,3,5,6,-tetramethylcinnamoylguanidine	0.3
3-ethoxycinnamoylguanidine	0.3
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	0.1
(4-Methoxycinnamoyl)guanidine	0.1
(2-Chlorocinnamoyl)guanidine	0.1
(3-Nitrocinnamoyl)guanidine	0.1
4-ethoxycinnamoylguanidine	0.1
3,4,5-trimethoxycinnamoylguanidine	0.1
2-(2-naphthyl)acetoylguanidine	0.1
N-(3-phenylpropanoyl)-N'-phenylguanidine	0.1

Example 31: Antiviral Assay for testing compounds against replication of human coronavirus 229E (229E).

5

To determine the antiviral activity of compounds against human coronavirus 229E replication (ATCC VR-740), an assay measuring reduction in the number of

-85-

plaques formed in monolayers of 229E infected MRC-5 cells (human lung fibroblasts ;ATCC CCL-171) was developed: First, a virus working stock was prepared by amplification in MRC-5 cells. This was then used to infect confluent monolayers of MRC-5 cells grown in 6-well tissue culture plates by exposure to the virus at an MOI of approx. 0.01 pfu/cell for 1 hour at 35°C in 5%CO₂. The infective inoculum was removed and replaced with fresh medium (DMEM supplemented with 10% fetal calf serum) containing various test concentrations of compounds or the appropriate level of solvent used for the compounds (control). Plates were subsequently incubated at 35°C (in 5% CO₂) for 3 –5 days post infection, after which time culture supernatant was removed and the cells were stained with 0.1% crystal violet solution in 20% ethanol for 10 minutes. Plaques were counted in all wells and the percentage reduction in plaque number compared to solvent control was calculated. Measurements were performed in duplicate to quadruplicate wells.

15 Table 10

Compound	Plaque Reduction (%control / # experiments)		
	5uM	2.5uM	1uM
2-t-butylcinnamoylguanidine	100 / 1	100 / 3	050 / 3
4-isopropylcinnamoylguanidine	100 / 1	100 / 2	057 / 2
3,4-dichlorocinnamoylguanidine	100 / 3	099 / 4	086 / 3
3-(trifluoromethoxy)cinnamoylguanidine	100 / 2	098 / 4	077 / 3
2,6-dichlorocinnamoylguanidine	100 / 1	097 / 3	066 / 2
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	100 / 1	097 / 3	021 / 1
2-cyclohexylcinnamoylguanidine	070 / 1	097 / 2	089 / 2
5-bromo-2-methoxycinnamoylguanidine	100 / 2	096 / 4	088 / 3
2-phenylcinnamoylguanidine	100 / 1	096 / 3	100 / 1
5-(2'-bromophenyl)penta-2,4-dienoylguanidine	100 / 2	095 / 3	079 / 2
4-t-butylcinnamoylguanidine	100 / 1	095 / 3	084 / 3
3-phenylcinnamoylguanidine		094 / 3	077 / 2
(3-Bromocinnamoyl)guanidine	100 / 2	093 / 3	072 / 2
(4-Bromocinnamoyl)guanidine	094 / 1	091 / 3	073 / 2
5-(N,N-hexamethylene)amiloride	089 /	091 / 2	033 / 1
trans-3-(1-naphthyl)acryloylguanidine	100 / 1	091 / 2	064 / 2

-86-

3-(2-naphthyl)acryloylguanidine	100 / 1	091 / 2	062 / 2
2,4-dichlorocinnamoylguanidine	100 / 2	090 / 4	064 / 3
(2-Nitrocinnamoyl)guanidine	085 / 2	090 / 2	046 / 2
3-(trifluoromethyl)cinnamoylguanidine	097 / 2	089 / 4	064 / 3
5-bromo-2-fluorocinnamoylguanidine	100 / 1	088 / 3	063 / 2
4-methylcinnamoylguanidine	091 / 2	087 / 4	063 / 2
(3-Chlorocinnamoyl)guanidine	100 / 1	086 / 3	009 / 1
(4-Methoxycinnamoyl)guanidine	100 / 1	085 / 4	057 / 3
(4-Chlorocinnamoyl)guanidine	100 / 2	084 / 2	051 / 2
3-fluorocinnamoylguanidine	095 / 1	083 / 3	051 / 2
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	100 / 1	082 / 3	063 / 2
(α -Methylcinnamoyl)guanidine	023 / 1	082 / 1	036 / 2
2,3,5,6-tetramethylcinnamoylguanidine	098 / 2	079 / 4	064 / 3
2-fluorocinnamoylguanidine	090 / 1	079 / 3	045 / 2
4-(trifluoromethyl)cinnamoylguanidine	100 / 1	079 / 1	052 / 1
(3-Nitrocinnamoyl)guanidine	100 / 1	079 / 1	045 / 1
2,5-dimethylcinnamoylguanidine	092 / 2	078 / 1	078 / 1
3- <i>t</i> -butylcinnamoylguanidine	100 / 1	077 / 4	030 / 3
(3-Methoxycinnamoyl)guanidine	089 / 1	075 / 2	030 / 1
3-methylcinnamoylguanidine	095 / 1	074 / 3	044 / 1
3-isopropylcinnamoylguanidine hydrochloride	089 / 1	074 / 3	014 / 1
(2-Bromocinnamoyl)guanidine	095 / 2	072 / 2	043 / 2
3-ethoxycinnamoylguanidine	100 / 1	072 / 3	057 / 1
(5-Phenyl-penta-2,4-dienoyl)guanidine	100 / 1	072 / 2	069 / 1
(2-Chlorocinnamoyl)guanidine	095 / 2	072 / 2	040 / 2
4-ethoxycinnamoylguanidine	073 / 1	069 / 2	057 / 1
4-fluorocinnamoylguanidine	100 / 1	067 / 3	034 / 2
3,4-difluorocinnamoylguanidine	085 / 1	065 / 3	042 / 2
N-(3-phenylpropanoyl)-N'-phenylguanidine	051 / 1	064 / 1	000 / 1
2,4,6-trimethylcinnamoylguanidine	075 / 2	063 / 3	062 / 2
2-methylcinnamoylguanidine	074 / 2	063 / 3	053 / 3
(trans-2-Phenylcyclopropanecarbonyl)-guanidine		063 / 2	022 / 1
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine		059 / 1	
N-Benzoyl-N'-cinnamoylguanidine		056 / 1	
4-phenylbenzoylguanidine	076 / 1	055 / 2	071 / 1
trans-3-Furanacryoylguanidine		055 / 2	018 / 1
(4-Phenoxybenzoyl)guanidine	069 / 1	054 / 3	040 / 2
(2-Methoxycinnamoyl)guanidine	051 / 1	053 / 2	024 / 1
N-amidino-3-amino-5-phenyl-6-	074 / 2	052 / 2	038 / 1

-87-

chloro-2-pyrazinecarboxamide			
N-(cinnamoyl)-N'phenylguanidine	084 / 1	048 / 2	035 / 1
cinnamoylguanidine	095 / 2	047 / 2	059 / 1
3,4-(methylenedioxy)cinnamoylguanidine	084 / 1	046 / 1	019 / 1
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide		045 / 1	
2,3-dimethylcinnamoylguanidine	073 / 1	044 / 2	024 / 1
5-(3'-bromophenyl)penta-2,4-dienoylguanidine		044 / 1	
N,N'-Bis(3-phenylpropanoyl)guanidine		041 / 1	
3-methoxy-amiloride	029 / 2	039 / 3	022 / 2
2,3-difluorocinnamoylguanidine		036 / 1	
1-naphthoylguanidine		036 / 1	
(3-phenylpropanoyl)guanidine		036 / 1	
6-methoxy-2-naphthoylguanidine	49 / 3	030 / 4	
5-(N,N-Dimethyl)amiloride hydrochloride		027 / 1	
2-ethoxycinnamoylguanidine		027 / 1	
2-naphthoylguanidine		027 / 1	
3,4,5-trimethoxycinnamoylguanidine		027 / 1	
3-methoxy-HMA		027 / 1	
Benzyoylguanidine		026 / 1	
2-(trifluoromethyl)cinnamoylguanidine		022 / 1	
N-amidino-3,5-diamino-6-phynyl-2-pyrazinecarboxamide		022 / 1	
cinnamoylguanidine hydrochloride		020 / 1	
(Quinoline-2-carbonyl)guanidine	015 / 3	019 / 3	006 / 2
(4-Hydroxycinnamoyl)guanidine		019 / 1	
5-(4-fluorophenyl)amiloride		018 / 1	
2-(1-naphthyl)acetoylguanidine		018 / 1	
(2-Furanacryloyl)guanidine		018 / 1	
[3-(3-Pyridyl)acryloyl]guanidine		018 / 1	
N-Cinnamoyl-N',N'-dimethylguanidine		015 / 1	
N-(2-naphthoyl)-N'-phenylguanidine		011 / 1	
2-(2-naphthyl)acetoylguanidine		009 / 1	
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine		009 / 1	
(Phenylacetyl)guanidine		009 / 1	

-88-

Example 32 Human OC43 Coronavirus**OC43 Antiviral Assay for testing compounds against replication of human coronavirus OC43.**

To determine the antiviral activity of compounds against human coronavirus OC43 replication (ATCC VR-759), an ELISA assay was developed measuring the release of the viral N-protein into culture supernatants from monolayers of OC43-infected MRC-5 cells (human lung fibroblasts ;ATCC CCL-171): First, a virus working stock was prepared by amplification in MRC-5 cells. This was then used to infect confluent monolayers of MRC-5 cells grown in 6-well tissue culture plates by exposure to the virus at an MOI of approx. 0.01 pfu/cell for 1 hour at 35°C in 5%CO₂. The infective inoculum was removed and replaced with fresh medium (DMEM supplemented with 10% fetal calf serum) containing various test concentrations of compounds or the appropriate level of solvent used for the compounds (control). Plates were subsequently incubated at 35°C (in 5% CO₂) for 5 days post infection, after which time culture supernatant was harvested and cellular debris removed by centrifugation at 5000 x g for 10 minutes. For N-antigen detection, 100µl samples of clarified culture supernatant were added to duplicate wells of a 96-well Maxi-Sorb plate; 100µl of RIPA buffer was added per well with mixing and the plate was covered and incubated at 4°C overnight to enable protein binding to the plastic wells. The next day, the coating solution was discarded, wells were washed thoroughly with PBST, and blocking of unoccupied protein binding sites was performed by incubation in 1% BSA in PBS for 1.5 hours. The antibody recognising OC43 N-protein was used at 1/800 dilution in PBS (1hr at 37°C) and the secondary antibody (goat-anti-mouse alkaline phosphatase) was used for the colour development reaction. Optical density of the wells was read at 405 nm and the effect of compounds determined by comparison of the level of signal in presence of compound to level of signal from the solvent control.

Example 33: Effect of compounds in OC43 antiviral assay

Compounds were screened for activity against OC43 replication according to the method described in example 22. Results are shown in Table 11.

Table 11

Compound	Virus inhibition at 2.5uM
3-methylcinnamoylguanidine	100
trans-3-(1-naphthyl)acryloylguanidine	100
(3-Bromocinnamoyl)guanidine	100
(2-Chlorocinnamoyl)guanidine	96
3,4-dichlorocinnamoylguanidine	90
3-(trifluoromethyl)cinnamoylguanidine	84
(trans-2-Phenylcyclopropanecarbonyl)guanidine	71
4-isopropylcinnamoylguanidine	68
cinnamoylguanidine	57
6-methoxy-2-naphthoylguanidine	47
2,4-dichlorocinnamoylguanidine	36
(4-Chlorocinnamoyl)guanidine	36
5-(N,N-hexamethylene)amiloride	30
(4-Bromocinnamoyl)guanidine	29
2,6-dichlorocinnamoylguanidine	27
5-bromo-2-methoxycinnamoylguanidine	24
(5-Phenyl-penta-2,4-dienoyl)guanidine	9
3-(trifluoromethoxy)cinnamoylguanidine	4
2-t-butylcinnamoylguanidine	4

5

Example 34. Mouse Hepatitis Virus (MHV).**Synthesis and Purification of a Peptide Corresponding to the MHV-A59 E protein.**

10 A peptide corresponding to the full-length MHV-A59 E protein (sequence: MFNLFLTDTVWYVGQIIFIFAVCLMVTIIVVAFLASIKLCIQLCGLCNTLVLSPIIYLYDRSKQLYKYYNEEMRLPLLEVDDI (SEQ ID NO:6); accession number NP_068673) was synthesized manually using Fmoc chemistry and solid phase peptide synthesis. The synthesis was done at the Bimolecular Resource Facility

15 (John Curtin School of Medical Research, ANU, Australia) using a Symphony^R Peptide Synthesiser from Protein Technologies Inc.(Woburn, MS, USA) according to the manufacturers instructions to give C-terminal amides, the coupling was done with HBTU and hydroxybenzotriazole in N-methylpyrrolidone. Each of the synthesis cycles used double coupling and a 4-fold excess of the amino acids. Temporary α -N-Fmoc- protecting groups were removed using 20% piperidine in DMF.

20

-90-

The crude synthetic peptide was purified using the ProteoPlus™ kit (Qbiogene inc. CA), following manufactures instructions. Briefly, the peptides were diluted in loading buffer (60mM Tris-HCl pH 8.3, 6M urea, 5% SDS, 10% glycerol, 0.2% Bromophenol blue, \pm 100 mM β -mercaptoethanol) and run on 4-20% gradient polyacrylamide gels (Gradipore, NSW, Australia) in tris-glycine electrophoresis buffer (25 mM Tris, 250 mM glycine, 0.1% SDS). The peptides were stained with gel code blue (Promega, NSW) and the bands corresponding to the full-length peptide were excised out of the gel.

The gel slice was transferred to the ProteoPLUS™ tube and filled with tris-glycine electrophoresis buffer. The tubes were emerged in tris-glycine electrophoresis buffer and subjected to 100 volts for approximately 1 hour. The polarity of the electric current was reversed for 1 minute to increase the amount of protein recovered. The peptides were harvested and centrifuged at 13, 000 rpm for 1 minute. The purified peptides were dried in a Speedvac and the weight of the final product was used to calculate the yield.

Example 35: MHV-E protein forms ion channels in planar lipid bilayers.

Lipid bilayer studies were performed as described elsewhere (Sunstrom, 1996; Miller, 1986). A lipid mixture of palmitoyl-oleoyl-phosphatidylethanolamine, palmitoyl-oleoyl-phosphatidylserine and palmitoyl-oleoyl-phosphatidylcholine (5:3:2) (Avanti Polar Lipids, Alabaster, Alabama) was used. The lipid mixture was painted onto an aperture of 150-200 μ m in the wall of a 1 ml delrin cup. The aperture separates two chambers, cis and trans, both containing salt solutions at different concentrations. The cis chamber was connected to ground and the trans chamber to the input of an Axopatch 200 amplifier. Normally the cis chamber contained either 500 mM NaCl or 500mM KCl and the trans 50 mM NaCl or 50mM KCl. The bilayer formation was monitored electrically by the amplitude of the current pulse generated by a current ramp. The potentials were measured in the trans chamber with respect to the cis. The synthetic peptide was added to the cis chamber and stirred until channel activity was seen. The currents were filtered at 1000 Hz, digitized at 5000 Hz and stored on magnetic disk.

-91-

The MHV E synthetic peptide was dissolved in 2,2,2-trifluoroethanol (TFE) at 0.05mg/ml to 1 mg/ml. 10 μ l of this was added to the cis chamber (1ml aqueous volume) of the bilayer apparatus, which was stirred via a magnetic "flea". Ionic currents, indicating channel activity in the bilayer, were typically detected within 15-30 min. After channels were detected the holding potential across the bilayer was varied between -100mV and +100mV to characterise the size and polarity of current flow and enable the reversal potential to be determined.

In 14 experiments where the cis chamber contained 500mM NaCl solution and the trans chamber contained 50 mM NaCl solution, the average reversal potential of the channel activity was calculated to be 49 ± 1 (SEM) mV. In 11 experiments where the cis chamber contained 500mM KCl solution and the trans chamber contained 50 mM KCl solution, the average reversal potential of the channel activity was calculated to be 13 ± 6 (SEM) mV. These results indicate that the MHV E protein forms cation selective ion channels that are more selective for Na^+ than for K^+ ions.

Figure 11 shows examples of raw current data for the MHV E ion channel at various holding potentials (cis relative to trans) in asymmetrical NaCl solutions (500/50 mM). The graph is a representative plot of average bilayer current (pA; y-axis) versus holding potential (mV; x-axis).

Example 36. Chemical compounds inhibit the ion channel activity of the MHV E protein synthetic peptide.

To test compounds for their ability to block or otherwise inhibit the ion channel formed by MHV E protein, small aliquots of solutions containing the compounds were added to the aqueous solutions bathing planar lipids in which the peptide channel activity had been reconstituted and the effect of the compound addition on the ionic currents was recorded and measured.

Compound stock solutions were typically prepared at 500 mM in DMSO. This solution was further diluted to 50 mM, or lower concentration in 50% DMSO/50% methanol and 2 μ l of the appropriately diluted compound was added to the cis and/or trans chambers to yield the desired final concentration.

-92-

In the example shown in Figure 12 below, addition of 100 μ M cinnamoylguanidine to the cis chamber greatly reduced current flow through the MHV E ion channel.

5

Example 37. Bacterial Bio-Assay for Screening Potential MHV E-protein Ion Channel-Blocking Drugs.

10 MHV E-protein Ion Channel inhibits Bacterial Cell growth.

A bio-assay of MHV E-protein function in bacterial cells was developed. A synthetic cDNA fragment encoding MHV E-protein was cloned into the expression plasmid pPL451, creating a vector in which E protein expression is temperature inducible, as described in Example 4. Inhibition of the growth of E.coli cells expressing E protein at 37°C was observed as an indicator of p7 ion channel function dissipating the normal Na⁺ gradient maintained by the bacterial cells.

15

Example 38. Compound Screening using the Bacterial Bio-Assay for MHV E protein.

20

The halos of growth around the site of application of particular drugs – as described in example 14 – were scored as described in example 15.

Table 12 lists the scores for inhibition of MHV E protein in the bacterial bio-assay.

25

Table 12

Compound	MHV E protein Inhibition (score)
4-isopropylcinnamoylguanidine	4.5
3-isopropylcinnamoylguanidine hydrochloride	4.2
4-t-butylcinnamoylguanidine	4.1
3-(trifluoromethoxy)cinnamoylguanidine	4.1
3-t-butylcinnamoylguanidine	4.0
3,4-dichlorocinnamoylguanidine	3.8
2,3-difluorocinnamoylguanidine	3.8
2-t-butylcinnamoylguanidine	3.8
3-phenylcinnamoylguanidine	3.7
2-phenylcinnamoylguanidine	3.4
5-bromo-2-methoxycinnamoylguanidine	3.3
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	3.3

-93-

3-(trifluoromethyl)cinnamoylguanidine	2.9
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	2.9
trans-3-(1-naphthyl)acryloylguanidine	2.8
4-(trifluoromethyl)cinnamoylguanidine	2.8
3-(2-naphthyl)acryloylguanidine	2.8
2-(trifluoromethyl)cinnamoylguanidine	2.7
(4-Phenoxybenzoyl)guanidine	2.4
(3-Bromocinnamoyl)guanidine	2.4
2,5-dimethylcinnamoylguanidine	2.3
5-bromo-2-fluorocinnamoylguanidine	2.1
6-methoxy-2-naphthoylguanidine	1.8
4-phenylbenzoylguanidine	1.8
(4-Bromocinnamoyl)guanidine	1.8
1-naphthoylguanidine	1.7
(5-Phenyl-penta-2,4-dienoyl)guanidine	1.4
(2-Bromocinnamoyl)guanidine	1.4
(4-Chlorocinnamoyl)guanidine	1.3
2-methylcinnamoylguanidine	1.2
2,6-dichlorocinnamoylguanidine	1.2
2,4,6-trimethylcinnamoylguanidine	1.2
5-(N,N-hexamethylene)amiloride	1.1
cinnamoylguanidine	1.1
cinnamoylguanidine hydrochloride	1.1
(α -Methylcinnamoyl)guanidine	1.0
2,3-dimethylcinnamoylguanidine	1.0
2-cyclohexylcinnamoylguanidine	0.9
N-(3-phenylpropanoyl)-N'-phenylguanidine	0.8
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	0.8
(3-Methoxycinnamoyl)guanidine	0.8
(2-Methoxycinnamoyl)guanidine	0.8
3-fluorocinnamoylguanidine	0.8
2-fluorocinnamoylguanidine	0.8
2,4-dichlorocinnamoylguanidine	0.8
2-ethylcinnamoylguanidine	0.8
(2-Chlorocinnamoyl)guanidine	0.7
(4-Hydroxycinnamoyl)guanidine	0.7
2-ethoxycinnamoylguanidine	0.7
2-naphthoylguanidine	0.6
(trans-2-Phenylcyclopropanecarbonyl)guanidine	0.6
5-(N,N-Dimethyl)amiloride hydrochloride	0.5
5-(4-fluorophenyl)amiloride	0.5
3-methylcinnamoylguanidine	0.5
(3-Chlorocinnamoyl)guanidine	0.4
4-methylcinnamoylguanidine	0.4
4-ethoxycinnamoylguanidine	0.4
2-(1-naphthyl)acetoyleguanidine	0.4

-94-

3,4-difluorocinnamoylguanidine	0.4
2-(2-naphthyl)acetoylguanidine	0.4
2,3,5,6,-tetramethylcinnamoylguanidine	0.4
(4-Methoxycinnamoyl)guanidine	0.3
3,4-(methylenedioxy)cinnamoylguanidine	0.3
3-ethoxycinnamoylguanidine	0.3
4-fluorocinnamoylguanidine	0.2
1-bromo-2-naphthoylguanidine	0.2
5-tert-butylamino-amiloride	0.1
(3-Nitrocinnamoyl)guanidine	0.1
3,4,5-trimethoxycinnamoylguanidine	0.1
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	0.1

5 **Example 39. MHV Antiviral Assay for testing compounds against replication of mouse hepatitis virus (MHV).**

To determine the antiviral activity of compounds against MHV replication (strain MHV-A59: ATCC VR-764), an assay measuring reduction in the number of plaques formed in monolayers of MHV infected L929 cells (ATCC CCL-a) was developed: First, a virus working stock was prepared by amplification in NCTC clone 1469 cells (ATCC CCL-9.1). This was then used to infect confluent monolayers of L929 cells grown in 6-well tissue culture plates by exposure to the virus at an MOI of 0.01 pfu/cell or 1 pfu/cell for 30 minutes at 37°C in 5%CO₂. The infective inoculum was removed and replaced with fresh medium (DMEM supplemented with 10% horse serum) containing various test concentrations of compounds or the appropriate level of solvent used for the compounds (control). Plates were subsequently incubated at 37°C (in 5% CO₂) for 16 – 24 hours post infection, after which time culture supernatant was removed and the cells were stained with 0.1% crystal violet solution in 20% ethanol for 10 minutes. Plaques were counted in all wells and the percentage reduction in plaque number compared to solvent control was calculated. Measurements were performed in duplicate to quadruplicate wells.

Example 40. Effect of compounds in MHV antiviral assay.

Table 13 provides the results obtained from this study.

Table 13

Compound	Percent reduction in Plaque number / # experiments		
	20uM	10uM	1uM
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	N/D	99 / 2	66 / 1
5-bromo-2-methoxycinnamoylguanidine	N/D	100 / 1	66 / 1
3-phenylcinnamoylguanidine	Toxic	86 / 2	64 / 3
2,3-difluorocinnamoylguanidine	Toxic	92 / 3	64 / 2
3-ethoxycinnamoylguanidine	100 / 1	89 / 2	58 / 1
5-(2'-bromophenyl)penta-2,4-dienoylguanidine	Toxic	100 / 1	57 / 1
cinnamoylguanidine hydrochloride	85 / 1	72 / 2	56 / 1
(2-Chlorocinnamoyl)guanidine	95 / 2	88 / 3	53 / 3
cinnamoylguanidine	97 / 8	88 / 8	52 / 7
(4-Bromocinnamoyl)guanidine	Toxic / 2	98 / 3	52 / 3
(2-Bromocinnamoyl)guanidine	91 / 2	89 / 3	52 / 3
(4-Methoxycinnamoyl)guanidine	98 / 4	96 / 4	51 / 3
(α -Methylcinnamoyl)guanidine	81 / 2	75 / 3	51 / 2
3,4-dichlorocinnamoylguanidine	91 / 2	96 / 1	50 / 2
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	N/D	97 / 1	50 / 1
3,4-difluorocinnamoylguanidine	Toxic	91 / 2	50 / 1
3-t-butylcinnamoylguanidine	Toxic	94 / 3	49 / 2
2-ethoxycinnamoylguanidine	93 / 2	85 / 3	48 / 2

-96-

trans-3-Furanacryoylguanidine	70 / 1	65 / 1	48 / 1
N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-2-pyrazinecarboxamide	84 / 1	52 / 2	48 / 1
(2-Nitrocinnamoyl)guanidine	97 / 1	77 / 2	47 / 1
4-(trifluoromethyl)cinnamoylguanidine	97 / 3	95 / 3	46 / 3
3,4-(methylenedioxy)cinnamoylguanidine	93 / 3	82 / 3	45 / 3
5-(N-Methyl-N-isobutyl)amiloride	92 / 1	85 / 1	44 / 2
(4-Chlorocinnamoyl)guanidine	97 / 2	88 / 2	43 / 3
2,4-dichlorocinnamolyguanidine	76 / 1	73 / 1	43 / 1
N-(3-phenylpropanoyl)-N'-phenylguanidine	80 / 1	65 / 1	43 / 1
(3-Nitrocinnamoyl)guanidine	95 / 2	77 / 3	42 / 3
2-phenylcinnamoylguanidine	N/D	100 / 1	42 / 1
4-isopropylcinnamoylguanidine	95 / 3	93 / 3	41 / 3
3-(trifluoromethoxy)cinnamoylguanidine	100 / 1	90 / 3	41 / 2
3-(trifluoromethyl)cinnamoylguanidine	98 / 1	83 / 1	40 / 1
(4-Nitrocinnamoyl)guanidine	97 / 1	75 / 3	40 / 3
3-(2-naphthyl)acryloylguanidine	93 / 1	93 / 1	40 / 1
4-ethoxycinnamoylguanidine	96 / 1	92 / 1	40 / 1
2,6-dichlorocinnamoylguanidine	91 / 1	70 / 1	40 / 1
2,5-dimethylcinnamoylguanidine	95 / 3	91 / 3	39 / 3
(3-Bromocinnamoyl)guanidine	95 / 2	90 / 3	39 / 3
(3-Chlorocinnamoyl)guanidine	94 / 1	86 / 2	39 / 2

-97-

3-methylcinnamoylguanidine	90 / 1	88 / 1	39 / 1
(3-Methoxycinnamoyl)guanidine	92 / 2	87 / 2	37 / 3
2-t-butylcinnamoylguanidine	N/D	98 / 2	37 / 1
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine	56 / 1	45 / 1	37 / 1
N,N'-bis(1-naphthoyl)guanidine	58 / 1	52 / 2	35 / 1
3-methoxy -HMA	15 / 1	31 / 1	35 / 1
5-tert-butylamino-amiloride	89 / 4	84 / 4	34 / 4
trans-3-(1-naphthyl)acryloylguanidine	95 / 2	86 / 3	34 / 3
6-methoxy-2-naphthoylguanidine	88 / 3	56 / 3	34 / 3
2-naphthoylguanidine	67 / 2	36 / 2	34 / 2
2-ethylcinnamoylguanidine	96 / 1	81 / 2	34 / 1
2,3-dimethylcinnamoylguanidine	95 / 1	79 / 2	34 / 1
N''-Cinnamoyl-N,N'-diphenylguanidine	97 / 1	72 / 2	34 / 1
3-isopropylcinnamoylguanidine hydrochloride	N/D	99 / 2	32 / 1
(4-Phenoxybenzoyl)guanidine	73 / 1	65 / 1	32 / 1
(trans-2-Phenylcyclopropanecarbonyl)guanidine	77 / 2	64 / 2	31 / 2
3-fluorocinnamoylguanidine	100 / 1	93 / 2	31 / 1
5-bromo-2-fluorocinnamoylguanidine	Toxic	81 / 2	31 / 1
N,N'-bis-(cinnamoyl)-N''-phenylguanidine	16 / 1	38 / 2	31 / 1
3-quinolinoylguanidine	27 / 1	36 / 2	30 / 1
2,4,6-trimethylcinnamoylguanidine	91 / 2	61 / 3	27 / 2

-98-

1-bromo-2-naphthoylguanidine	31 / 1	27 / 2	27 / 1
N-amidino-3,5-diamino-6-phenyl-2-pyrazinecarboxamide	53 / 1	39 / 2	25 / 1
N-Cinnamoyl-N',N'-dimethylguanidine	92 / 2	65 / 3	24 / 2
(2-Methoxycinnamoyl)guanidine	90 / 2	85 / 2	23 / 2
2-(2-naphthyl)acetylguanidine	52 / 1	20 / 2	23 / 1
4-phenylcinnamoylguanidine	53 / 1	36 / 1	21 / 3
[3-(3-Pyridyl)acryloyl]guanidine	81 / 2	73 / 2	21 / 2
3,4,5-trimethoxycinnamoylguanidine	84 / 1	84 / 1	21 / 1
4-methylcinnamoylguanidine	93 / 1	89 / 1	20 / 1
4-fluorocinnamoylguanidine	86 / 1	83 / 1	20 / 1
2-methylcinnamoylguanidine	91 / 1	82 / 1	20 / 1
6-bromo-2-naphthoylguanidine	65 / 1	37 / 2	19 / 1
5-(N,N-Dimethyl)amiloride hydrochloride	42 / 4	7 / 4	17 / 4
(5-Phenyl-penta-2,4-dienoyl)guanidine	27 / 1	24 / 1	17 / 1
2-cyclohexylcinnamoylguanidine	100 / 1	74 / 2	16 / 1
5-(4-fluorophenyl)amiloride	4 / 1	25 / 1	16 / 1
Benzoylguanidine	22 / 1	39 / 2	14 / 1
N-Benzoyl-N'-cinnamoylguanidine	0 / 1	0 / 1	14 / 1
5-(N,N-hexamethylene)amiloride	84 / 2	89 / 1	13 / 2
N-(cinnamoyl)-N'phenylguanidine	83 / 1	88 / 1	13 / 1
(4-Hydroxycinnamoyl)guanidine	19 / 1	15 / 1	13 / 1

-99-

2-(trifluoromethyl)cinnamoylguanidine	19 / 1	15 / 1	13 / 1
(Quinoline-2-carbonyl)guanidine	86 / 1	84 / 1	12 / 1
2-(1-naphthyl)acetoyleguanidine	19 / 1	02 / 1	11 / 1
2-chloro-6-fluorocinnamoylguanidine	100 / 1	84 / 2	9 / 1
N-amidino-3-amino-5-phenyl-6-chloro-2-pyrazinecarboxamide	20 / 1	20 / 2	9 / 1
4-phenylbenzoylguanidine	32 / 1	24 / 1	5 / 1
N,N'-bis(2-naphthoyl)guanidine	5 / 1	3 / 2	3 / 1
(Phenylacetyl)guanidine	35 / 1	22 / 1	3 / 1
1-naphthoylguanidine	71 / 3	62 / 3	2 / 3
N,N'-bis(3phenylpropanoyl)-N"-phenylguanidine	67 / 3	40 / 4	1 / 3
3-hydroxy-5-hexamethyleneimino-amiloride	16 / 1	22 / 2	1 / 1
2'4 DichloroBenzamil HCl	12 / 2	0 / 3	0 / 3
2,3,5,6,-tetramethylcinnamoylguanidine	N/D	68 / 2	0 / 1
Benzamil hydrochloride	0 / 1	26 / 1	0 / 1
6-Iodoamiloride	28 / 1	21 / 1	0 / 1
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide	19 / 1	16 / 1	0 / 1
[(4-Chlorophenoxy-acetyl)guanidine	19 / 1	16 / 1	0 / 1
(3-phenylpropanoyl)guanidine	51 / 1	03 / 1	0 / 1
2-fluorocinnamoylguanidine	76 / 1	73 / 1	
6-bromo-2-naphthoylguanidine		43 / 1	
(2-Furanacryloyl)guanidine	67 / 2	63 / 2	-3 / 2

-100-

N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine	43 / 1	39 / 1	-5 / 1
Amiloride.HCl	21 / 1	18 / 1	-5 / 1
3-(trans-hept-1-en-1-yl)cinnamoylguanidine	T / 1	23(T) / 1	-6 / 1
3-methoxy-amiloride	60 / 2	47 / 3	-7 / 2
N,N'-Bis(3-phenylpropanoyl)guanidine	41 / 3	30 / 4	-8 / 3
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	T / 1	2 / 2	-19 / 1

Example 41. Porcine Respiratory Coronavirus (PRCV)**Antiviral Assay for testing compounds against replication of porcine respiratory coronavirus (PRCV).**

To determine the antiviral activity of compounds against porcine respiratory coronavirus replication (ATCC VR-2384), an assay measuring reduction in the number of plaques formed in monolayers of PRCV infected ST cells (porcine fetal testis cell line, ATCC CRL-1746) was developed: Confluent ST cells in 6 well plates were infected with a quaternary passage of porcine respiratory virus (PRCV) strain AR310 at three dilutions 10^{-1} , 50^{-1} and 10^{-2} in PBS to provide a range of plaques numbers to count. 100ul of diluted virus was added per well in a volume of 1ml of media. Plates were incubated for one hour on a rocking platform at room temperature to allow virus to adsorb to cells. The viral supernatant was removed and 2ml/well of overlay containing 1% Seaplaque agarose in 1x MEM, 5% FCS was added to each well. Compounds to be tested were added to the overlay mixture by diluting the compounds from frozen stock to a concentration so that the same volume of compound/solvent would be added to the overlay for each concentration of compound. The volume of compound/solvent never exceeded 0.07% of the volume of the overlay. The solvent used to dissolve compounds was DMSO and methanol mixed in equal proportions. Compounds were tested for anti-plaque forming activity at four concentrations, 0.1uM, 1uM, 10uM and 20uM. Either duplicates or quadruplicates were performed at each concentration. Controls were performed

-101-

where the same volume of solvent was added to the overlay. The overlay was allowed to set at room temp for 20 mins. The plates were then incubated at 37°C for 2 days. The monolayers were then fixed and stained overnight at room temperature by adding 1ml/well of 0.5% methylene blue, 4% formaldehyde. Overlay agarose and stain was then rinsed off to visualize stained and fixed monolayer

Example 42: Effect of compounds in PRCV antiviral assay

Compounds were screened for activity against PRCV replication according to the method described in example 29. Table 14 provides EC50 values for some tested compounds.

Table 14

Compound	EC50 (uM)
5-(N,N-hexamethylene)amiloride	0.06
6-methoxy-2-naphthoylguanidine	0.04
cinnamoylguanidine	0.08
N-(3-phenylpropanoyl)-N'-phenylguanidine	19
3-methylcinnamoylguanidine	1.43
(3-Bromocinnamoyl)guanidine	11.2
(trans-2-Phenylcyclopropanecarbonyl)guanidine	17.2
trans-3-(1-naphthyl)acryloylguanidine	19.1
2-(2-naphthyl)acetoylguanidine	119.6

Example 43. Bovine Coronavirus.

Antiviral Assay for testing compounds against replication of bovine coronavirus (BCV).

To determine the antiviral activity of compounds against bovine coronavirus replication (ATCC VR-874), an assay measuring reduction in the number of plaques formed in monolayers of BCV infected MDBK cells (bovine kidney cell line ;ATCC CCL-22) was developed: Confluent MDBK cells in 6 well plates were infected with a secondary passage of BCV with serially diluted virus diluted to 10^{-3} , 5^{-5} and 10^{-4} in PBS to provide a range of plaques numbers to count. 100ul of diluted virus was

-102-

added per well in a volume of 1ml of media. Plates were incubated for one hr to allow virus to adsorb to cells. The viral supernatant was removed and 2ml/well of overlay containing 1% Seaplaque agarose in 1x MEM, 5% FCS was added to each well. Compounds to be tested were added to the overlay mixture by diluting the

5 compounds from a 0.5M frozen stock to a concentration so that the same volume of compound/solvent would be added to the overlay for each concentration of compound. The volume of compound/solvent never exceeded 0.07% of the volume of the overlay. The solvent used to dissolve compounds was DMSO and methanol mixed in equal proportions. Compounds were tested for anti-plaque forming activity

10 at four concentrations, 0.1uM, 1uM, 10uM and 20uM. Quadruplicates were performed at each concentration. Controls were performed where the same volume of solvent was added to the overlay. The overlay was allowed to set at room temp for 20 mins. The plates were then incubated at 37°C for 7 days. The monolayers were then fixed and stained by adding 1ml/well of 0.5% methylene blue, 4%

15 formaldehyde.

Example 44: Effect of compounds in BCV antiviral assay

Compounds were screened for activity against BCV replication according to the method described in example 31. Table 15 provides EC50 values for some tested

20 compounds.

Table 15

Compound	EC50 uM
(3-Bromocinnamoyl)guanidine	3
3-(trifluoromethyl)cinnamoylguanidine	3
6-methoxy-2-naphthoylguanidine	9
5-(N,N-hexamethylene)amiloride	9
trans-3-(1-naphthyl)acryloylguanidine	13
cinnamoylguanidine	42
(5-Phenyl-penta-2,4-dienoyl)guanidine	95
2-(2-naphthyl)acetoyleguanidine	99
(trans-2-Phenylcyclopropanecarbonyl)guanidine	109
N-(3-phenylpropanoyl)-N'-phenylguanidine	156
4-phenylbenzoylguanidine	190

Example 45 Hepatitis C Virus**Ion channel activity of Hepatitis C virus P7 Protein****Testing of a Synthetic P7 Peptide for channel activity in artificial lipid bilayers**

A peptide mimicking the protein P7 encoded by the hepatitis C virus (HCV)
 5 was synthesised having the following amino acid sequence (SEQ ID NO:7):

ALENLVILNAASLAGTHGLVSFLVFFCFAWYLGGRWVPGAVYAFYGMWPL
 LLLLLALPQRAYA

Lipid bilayer studies were performed as described elsewhere (Miller, 1986).
 10 A lipid mixture of palmitoyl-oleoyl-phosphatidylethanolamine, palmitoyl-oleoyl-phosphatidylserine and palmitoyl-oleoyl-phosphatidylcholine (5:3:2) (Avanti Polar Lipids, Alabaster, Alabama) was used. The lipid mixture was painted onto an aperture of 150-200 μm in the wall of a 1 ml delrin cup. The aperture separates two chambers, cis and trans, both containing salt solutions at different concentrations.
 15 The cis chamber was connected to ground and the trans chamber to the input of an Axopatch 200 amplifier. Normally the cis chamber contained 500 mM KCl and the trans 50 mM KCl. The bilayer formation was monitored electrically by the amplitude of the current pulse generated by a current ramp. The potentials were measured in the trans chamber with respect to the cis. The protein was added to the cis chamber and
 20 stirred until channel activity was seen. The currents were filtered at 1000 Hz, digitized at 2000 Hz and stored on magnetic disk. The P7 peptide was dissolved in 2,2,2-trifluoroethanol (TFE) at 10mg/ml. 10ul of this was added to the cis chamber of the bilayer which was stirred. Channel activity was seen within 15-20 min.

When the P7 peptide was added to the cis chamber and stirred, channel
 25 activity was recorded. The potential in the trans chamber was -80 mV and the currents are downwards. The currents reversed at +50 mV close to the potassium equilibrium potential in these solutions indicating that the channels were cation-selective. The amplitude of the open-channel peak is 1.7 pA corresponding to a channel conductance of about 14 pS. In most experiments, "single channels" had a
 30 much larger size, presumably because of aggregation of the P7 peptide. The currents reversed at about +40 mV in this experiment. In some experiments the solution in the

-104-

cis chamber was 150 mM KCl and 15 mM KCl in the trans chamber. The P7 peptide again produced currents that reversed.

Similar results were obtained when both chambers contained NaCl. Currents recorded in an experiment when the cis chamber contained 500 mM NaCl and the trans chamber 50 mM NaCl. Again the currents reversed between +40 and +60 mV, close to the Na^+ equilibrium potential indicating that channels were much more permeable to Na^+ than to K^+ .

10 The channels formed by the P7 peptide were blocked by 5-(N,N-hexamethylene) amiloride (HMA).

Addition of the P7 peptide produced channel activity: Addition of 2 μl of 50 μM HMA to the cis chamber followed by stirring resulted in disappearance of the channel activity. Block of channel activity produced by the P7 peptide with 100 μM HMA was recorded in 4 experiments. In 2 experiments, sodium channels (500/50) were blocked by 500 μM HMA

When 10 mM CaCl_2 was added to the cis chamber (K solutions) the reversal potential of the currents produced by P7 peptide shifted to more negative potentials indicating a decrease in the $P_{\text{K}}/P_{\text{Cl}}$ ratio.

When the cis chamber contained 500 mM CaCl_2 and the trans chamber 50 mM CaCl_2 , both positive and negative currents were seen at potentials around +20 mV and it was not possible to determine a reversal potential.

Example 46. Recombinant Expression of HCV p7 protein.

Two cDNA fragments, each encoding the same polypeptide corresponding to the amino acid sequence of the HCV-1a p7 protein, were synthesised commercially by GeneScript. The two cDNAs differed in nucleotide sequence such that in one cDNA ("cDp7.coli") the codons were optimised for expression of the p7 protein in E.coli while in the other cDNA ("cDp7.mam") codons were biased for expression in

-105-

mammalian cell lines. cDp7.coli was cloned into the plasmid pPL451 as a BamHI/EcoRI fragment for expression in E.coli. cDp7.mam was cloned into vectors (for example, pcDNA3.1 vaccinia virus, pfastBac-1) for expression of p7 in mammalian cell lines.

5

Example 47. Role of p7 in enhancement of Gag VLP Budding.

The budding of virus-like particles (VLP) from cultured HeLa cells results from the expression of retroviral Gag proteins in the cells and co-expression of small viral ion channels, such as M2, Vpu and 6K, with the Gag protein enhances budding. Interestingly, the viral ion channels can enhance budding of heterologous virus particles. Therefore, to assess budding enhancement by p7 it was co-expressed with the HIV-1 Gag protein in HeLa cells, and VLP release into the culture medium was measured by Gag ELISA. To achieve this, the plasmids pcDNAp7 (pc DNA3.1 = pcDp7.mam as described in Example 20, p7 expressed under control of the T7 promoter) and pcDNAGag (HIV-1 Gag protein expressed under control of the T7 promoter) were cotransfected into HeLa cells infected with the vaccinia virus strain vTF7.3 (expresses T7 RNA polymerase) and culture supernatants were collected for ELISA assay after 16 hours incubation.

Example 48. Assay of the ability of compounds to inhibit p7 ion channel functional activity.

The two methods of detecting p7 ion channel functional activity, described in Examples 33-35, were employed to assay the ability of compounds to inhibit the p7 channel. In the case of Example 33, compounds were tested for their ability to inhibit p7 channel activity in planar lipid bilayers. In the case of Example 35 compounds were tested for their ability to reduce the number of VLPs released from cells expressing both p7 and HIV-1 Gag.

30

-106-

Example 49.**Bacterial Bio-Assay for Screening Potential HCV p7 protein Ion Channel-Blocking Drugs.****HCV p7 Ion Channel inhibits Bacterial Cell growth.**

- 5 A bio-assay of p7 function in bacterial cells was developed. The p7-encoding synthetic cDNA fragment cDp7.coli was cloned into the expression plasmid pPL451, creating the vector pPLp7, in which p7 expression is temperature inducible, as described in Example 4. Inhibition of the growth of E.coli cells expressing p7 at 37°C was observed as an indicator of p7 ion channel function dissipating the normal
- 10 Na⁺ gradient maintained by the bacterial cells.

Example 50 Compound Screening using the Bacterial Bio-Assay for HCV p7 protein.

- The halos of growth around the site of application of particular drugs – as described
- 15 in example 14 – were scored as described in example 15.

Table 16 lists the scores for inhibition of HCV p7 protein in the bacterial bio-assay.

Table 16

Compound	HCV p7 protein Inhibition (score / # of times tested)
2,3-dimethylcinnamoylguanidine	3.88 /2
2,4,6-trimethylcinnamoylguanidine	3.75 / 1
5-bromo-2-fluorocinnamoylguanidine	3.73 /6
(4-Bromocinnamoyl)guanidine	3.47 /4
2,5-dimethylcinnamoylguanidine	3.43 /4
3-(trifluoromethyl)cinnamoylguanidine	3.34 /3
4-(trifluoromethyl)cinnamoylguanidine	3.33 /5
6-methoxy-2-naphthoylguanidine	3.33 /3
(2-Chlorocinnamoyl)guanidine	3.31 /6
(4-Chlorocinnamoyl)guanidine	3.16 /4
(2-Bromocinnamoyl)guanidine	3.00 /3
2,6-dichlorocinnamoylguanidine	3.00 /3
(3-Bromocinnamoyl)guanidine	2.92 /3
(3-Chlorocinnamoyl)guanidine	2.75 /3
2-(trifluoromethyl)cinnamoylguanidine	2.63 /3
(4-Phenoxybenzoyl)guanidine	2.63 /1

-107-

3,4-dichlorocinnamoylguanidine	2.59 /3
4-isopropylcinnamoylguanidine	2.51 /2
trans-3-(1-naphthyl)acryloylguanidine	2.44 /2
4-t-butylcinnamoylguanidine	2.42 /2
2-t-butylcinnamoylguanidine	2.36 /2
2-ethylcinnamoylguanidine	2.36 /2
4-methylcinnamoylguanidine	2.32 /2
5-bromo-2-methoxycinnamoylguanidine	2.32 /2
3-(trifluoromethoxy)cinnamoylguanidine	2.26 /2
2-cyclohexylcinnamoylguanidine	2.26 /2
1-naphthoylguanidine	2.25 /1
3-t-butylcinnamoylguanidine	2.23 /2
4-phenylbenzoylguanidine	2.19 /2
(5-Phenyl-penta-2,4-dienoyl)guanidine	2.13 /1
N-(cinnamoyl)-N'-phenylguanidine	2.13 /1
3-isopropylcinnamoylguanidine hydrochloride	2.00 /1
Benzamil hydrochloride	2.0 /1
N-(3-phenylpropanoyl)-N'-phenylguanidine	2.0 /1
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine	2.0 /1
3-(2-naphthyl)acryloylguanidine	1.93 /2
5-(N-Methyl-N-isobutyl)amiloride	1.88 /1
2'4 DichloroBenzamil HCl	1.88 /1
5-tert-butylamino-amiloride	1.88 /1
5-(N-Ethyl-N-isopropyl)amiloride	1.88 /1
(4-Methoxycinnamoyl)guanidine	1.88 /1
4-fluorocinnamoylguanidine	1.86 /1
(3-Nitrocinnamoyl)guanidine	1.71 /1
4-ethoxycinnamoylguanidine	1.63 /1
(4-Hydroxycinnamoyl)guanidine	1.63 /1
(trans-2-Phenylcyclopropanecarbonyl)guanidine	1.63 /1
3-ethoxycinnamoylguanidine	1.63 /1
2,3,5,6,-tetramethylcinnamoylguanidine	1.51 /2
4-phenylcinnamoylguanidine	1.5 /1
trans-3-Furanacryoylguanidine	1.38/1
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine	1.38 /1
(2-Furanacryloyl)guanidine	1.38 /1
3-(cyclohex-1-en-1-yl)cinnamoylguanidine	1.32 /2
cinnamoylguanidine hydrochloride	1.32 /2
5-(N,N-hexamethylene)amiloride	1.28 /4
2,3-difluorocinnamoylguanidine	1.24 /1
2-(1-naphthyl)acetoyleguanidine	1.14/ 1
(a-Methylcinnamoyl)guanidine	1.14 /1
(2-Nitrocinnamoyl)guanidine	1.14 /1
6-Iodoamiloride	1.13 /1
3,4-(methylenedioxy)cinnamoylguanidine	1.13 /1
2-ethoxycinnamoylguanidine	1.00 /1
cinnamoylguanidine	1.00 /1
2-phenylcinnamoylguanidine	1.00 /1

-108-

2-(cyclohex-1-en-1-yl)cinnamoylguanidine	1.00 / 1
2-naphthoylguanidine	1.0 / 3
3-phenylcinnamoylguanidine	1.0 / 1
5-(N,N-Dimethyl)amiloride hydrochloride	1.0 / 1
5-(4-fluorophenyl)amiloride	1.0 / 1
(3-Methoxycinnamoyl)guanidine	1.0 / 1
2-fluorocinnamoylguanidine	1.0 / 1
5-(3'-bromophenyl)penta-2,4-dienoylguanidine	1.0 / 1
[(4-Chlorophenoxy-acetyl)guanidine	1.0 / 1
(3-phenylpropanoyl)guanidine	1.0 / 1
2-chloro-6-fluorocinnamoylguanidine	0.88 / 1
3-fluorocinnamoylguanidine	0.86 / 1
2-methylcinnamoylguanidine	0.75 / 1
(2-Methoxycinnamoyl)guanidine	0.75 / 1
1-bromo-2-naphthoylguanidine	0.75 / 1
3,4,5-trimethoxycinnamoylguanidine	0.71 / 1
3-methylcinnamoylguanidine	0.63 / 1
3-(trans-hept-1-en-1-yl)cinnamoylguanidine	0.50 / 1
Amiloride.HCl	0.5 / 2
Phenamil methanesulfonate salt	0.5 / 1
2,4-dichlorocinnamolyguanidine	0.38 / 1
(4-Nitrocinnamoyl)guanidine	0.25 / 1
3,4-difluorocinnamoylguanidine	0.13 / 1
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine	0.03 / 4

Example 51: Equine Arteritis Virus (EAV)**Antiviral Assay for testing compounds against replication of equine arteritis virus (EAV).**

- 5 To determine the antiviral activity of compounds against EAV replication (strain Bucyrus; ATCC VR-796), an assay measuring reduction in the number of plaques formed in monolayers of EAV infected BHK-21 cells (ATCC CCL-10) was developed: A virus stock was amplified in RK-13 cells (ATCC CCL-37) and this was then used to infect confluent monolayers of BHK-21 cells grown in 6-well tissue
- 10 culture plates by exposure to the virus at an MOI of 5×10^{-3} pfu/cell for 1 hour at 37°C 5% CO₂. The infective inoculum was removed and the cells were overlaid with a 1% sea plaque overlay (Cambrex Bio Science) in MEM containing 10% FCS containing and 10, 5 or 1 μ M of compounds to be tested or the appropriate level of solvent used for the compounds (control). Plates were subsequently incubated at
- 15 37°C (in 5% CO₂) for 3 days post infection, after which time culture supernatant was removed and the cells were stained with 0.1% crystal violet solution in 20% ethanol

for 10 minutes. Plaques were counted in all wells and the percentage reduction in plaque number compared to solvent control was calculated. Measurements were performed in duplicate to quadruplicate wells.

5 **Example 52: Effect of compounds in EAV antiviral assay**

Compounds were screened for activity against EAV replication according to the method described in example 35. Results expressed as IC₅₀ values are shown in Table 17.

10 **Table 17**

Compound	IC ₅₀
5-(N,N-hexamethylene)amiloride	7.5 μ M
(3-Bromocinnamoyl)guanidine	10 μ M
trans-3-(1-naphthyl)acryloylguanidine	7.5 μ M
2-t-butylcinnamoylguanidine	1 μ M
2-(cyclohex-1-en-1-yl)cinnamoylguanidine	10 μ M

Example 53 Dengue Flavivirus

Peptide Synthesis of Dengue virus M Protein

15 The C-terminal 40 amino acids of the M protein of the Dengue virus type 1 strain Singapore S275/90 (Fu et al 1992)
 (ALRHPGFTVIALFLAHAIGTSITQKGIFILLMLVTPSMA) (SEQ ID NO: 8) was synthesised using the Fmoc method. The synthesis was done on a Symphony Peptide Synthesiser from Protein Technologies Inc (Tucson, Arizona) as used to give C-
 20 terminal amides, the coupling was done with HBTU and hydroxybenzotriazole in N-methylpyrrolidone. Each of the synthesis cycle used double coupling and a 4-fold excess of the amino acids. Temporary α -N Fmoc-protecting groups were removed using 20% piperidine in DMF.

Example 54. Incorporation of Dengue M virus protein into lipid bilayers.

25 Lipid bilayer studies were performed as described elsewhere (Sunstrom, 1996; Miller, 1986). A lipid mixture of palmitoyl-oleoyl-phosphatidylethanolamine,

-110-

palmitoyl-oleoyl-phosphatidylserine and palmitoyl-oleoyl-phosphatidylcholine (5:3:2) (Avanti Polar Lipids, Alabaster, Alabama) was used. The lipid mixture was painted onto an aperture of 150-200 μm in the wall of a 1 ml delrin cup. The aperture separates two chambers, cis and trans, both containing salt solutions at different concentrations. The cis chamber was connected to ground and the trans chamber to the input of an Axopatch 200 amplifier. Normally the cis chamber contained 500 mM KCl and the trans 50 mM KCl. The bilayer formation was monitored electrically by the amplitude of the current pulse generated by a current ramp. The potentials were measured in the trans chamber with respect to the cis. The protein was added to the cis chamber and stirred until channel activity was seen. The currents were filtered at 1000 Hz, digitized at 5000 Hz and stored on magnetic disk.

The dengue virus M protein C-terminal peptide (DMVC) was dissolved in 2,2,2-trifluoroethanol (TFE) at 0.05mg/ml to 1 mg/ml. 10 μl of this was added to the cis chamber of the bilayer which was stirred. Channel activity was seen within 15-30 min.

Example 55: Hexamethylene amiloride (HMA) to inhibits ion channel activity of the dengue virus M protein C-terminal peptide.

Solutions of 50 mM HMA were prepared by first making a 500 mM solution in DMSO. This solution was further diluted to 50 mM HMA using 0.1 M HCl. 2 μl of the 50 mM HMA was added to the cis chamber after channel activity was seen. The cis chamber contained 1 ml of solution making the final concentration of HMA 100 μM .

Example 56. Antiviral Assay for testing compounds against Effects of Dengue flavivirus against cytoproliferation.

Compounds were tested at 10, 5, 2.5, 1.25 and 0.625 μM for activity against Dengue 1 strain Hawaii using a cytoproliferation assay which measures the effect of dengue virus infection on proliferation of LLC-MK2, rhesus macaque monkey kidney cells. The LLC-MK2 cells were infected with a predetermined amount of virus, titrated such that cell proliferation in infected cultures would be significantly reduced

-111-

compared to uninfected controls. The infected cells were then plated at 1.5×10^3 cells per well in a 96 well plate. Negative controls (no virus, no experimental compound), positive controls (virus, no experimental compound), and cytotoxicity controls (experimental compound, no virus) were run with each drug assay. The cultures were allowed to grow for 7 days and then Alamar Blue, a fluorescent dye that measures the metabolism of the cultures (red/ox), was added to each culture and the fluorescence value for each culture was measured. The negative control without experimental compound or virus was fixed at 100%. The positive controls and the cultures with compound were scored by calculating their average fluorescence as a percentage of the negative control. At least six replicate wells were measured for each experimental condition.

Example 57 Effect of compounds in Dengue antiviral assay:

15 **Table 18**

Drug	Drug Conc. μ M	Antiviral Percent of Negative Control
cinnamoylguanidine		
Negative control	0	NA
Positive control	0	76.5%
	10	17.1%
	5	38.6%
	2.5	58.3%
	1.25	72.6%
	0.625	76.6%
(2-chlorocinnamoyl)guanidine		
Negative control	0	NA
Positive control	0	80.3%
	10	8.4%
	5	7.7%
	2.5	22.7%
	1.25	52.5%
	0.625	64.3%
Trans-3-(1-naphthyl)acryloylguanidine		
Negative control	0	NA
Positive control	0	80.4%
	10	6.8%

-112-

	5	12.4%
	2.5	38.7%
	1.25	73.7%
	0.625	77.7%

N.A. – not applicable

Example 58: Positive correlation between Bacterial Assay and Anti-viral Assays**5 Example 58.1 Positive correlation between Vpu Bacterial Assay and anti-HIV-1 Data.**

A correlative study was performed to measure correlation between the activity scores assigned to compounds tested in the Vpu bacterial assay and the ability of these compounds to inhibit HIV-1 in the anti-viral assay.

10 Example 58.2. Methodology

The p24-antigen data for twelve compounds representing various substituted acyl-guanidines was compared with the activity scores obtained for those compounds in the Vpu bacterial assay. The data from each assay was initially rank ordered for effectiveness. The rank order for the Vpu bacterial assay was determined from all
 15 activity scores, the highest score indicating the greatest effectiveness. The rank order for the anti-HIV-1 assay was determined based on the overall average value of p24 antigen measured in culture supernatants at all of the drug concentrations tested, with the lowest score indicating the greatest effectiveness. The two rank orders generated
 20 coefficient.

Example 58.3. Results and Conclusion

The Spearman's correlation coefficient was 0.785 which, by comparison with a statistical table of critical values (for n=12), indicates that the two rank orders are
 25 significantly positively correlated ($P < 0.01$) (Table 19a).

In addition, a different comparison of the Vpu Bacterial assay rank order with a yes/no score for whether the anti-viral data indicated a p24 reduction of at least one order of magnitude, aligned the 'yes' group of compounds with the top 6 compounds by the bacterial assay (Table 19b).

-113-

These results are indicative that a positive correlation exists between bacterial assays and the antiviral assays as performed according to the present invention. The bacterial assay may therefore be a useful tool in screening for compounds that exhibit anti-viral activity.

5

Table 19a. Comparison of Rank order of efficacy of 12 substituted acyl-guanidines in the Vpu bacterial assay and anti-HIV assay.

Compound	Bacterial assay rank order	p24 rank order	di ²
(3-bromocinnamoyl)guanidine	1	1	0
3-(trifluoromethyl)cinnamoylguanidine	2	2	0
3-methylcinnamoylguanidine	3	3	0
cinnamoylguanidine	4	4	0
trans -3-(1-naphthyl)acryloylguanidine	5.5	7	2.25
6-methoxy-2-naphthoylguanidine	5.5	5	0.25
4-phenylbenzoylguanidine	7	11	16
(5-phenyl-penta-2,4-dienoyl)guanidine	8	9	1
N-(3-phenylpropanoyl)-N'-phenylguanidine	9	12	9
Hexamethylene amiloride	10	6	16
2-(2-naphthyl)acetoylguanidine	11	10	1
(trans-2-phenylcyclopropanecarbonyl)guanidine	12	8	16
Sum di ²			61.5
Spearman correlation coefficient			0.785
P value			>0.01

Table 19b.

10

Compounds Ordered by p24 rank order	Vpu Bacterial score	Bacterial assay rank order	At least 1x log reduction seen in p24 assay?
(3-bromocinnamoyl)guanidine	4.3	1	yes
3-(trifluoromethyl)cinnamoylguanidine	3.7	2	yes
3-methylcinnamoylguanidine	3.4	3	yes
cinnamoylguanidine	3.0	4	yes
trans -3-(1-naphthyl)acryloylguanidine	2.9	5.5	yes
6-methoxy-2-naphthoylguanidine	2.9	5.5	yes
4-phenylbenzoylguanidine	2.8	7	no
(5-phenyl-penta-2,4-dienoyl)guanidine	2.6	8	no
N-(3-phenylpropanoyl)-N'-phenylguanidine	2.2	9	no
Hexamethylene amiloride	1.9	10	no
2-(2-naphthyl)acetoylguanidine	1.2	11	no

-114-

(trans-2-phenylcyclopropanecarbonyl)guanidine	0.4	12	no
---	-----	----	----

5 **Example 58.4. Correlation Between Percent inhibition of MHV plaque formation and MHV-E bacterial bio-assay score.**

A positive correlation was seen between the activity scores assigned to compounds when tested in the Mouse Hepatitis Virus E-protein bacterial bio-assay
 10 and the percent inhibition exhibited by these compounds in the Mouse Hepatitis Virus plaque assay.

Example 58.5. Method:

MHV plaque reduction activity data for 96 compounds screened were sorted from greatest to least percent plaque reduction and rank orders were assigned to the list of
 15 compounds. This was performed for the data generated by exposure to both 10 μ M and 1 μ M concentrations of the compounds, giving rise to two rank order lists. Similarly, a rank order list was generated for the MHVE bacterial bioassay scores for the same 96 compounds. Where one or more compounds had the same score, the rank values for that group were averaged.

20 Spearman's statistical test for [as described in "Mathematical Statistics with Applications" (2nd edn): Mendenhall, W., Scheaffer, RL., & Wackerly, DD. Duxbury Press, Boston Massachusetts – 1981] was used to compare rank orders. Briefly, this involved calculating the Sum of squares (SS) of the differences between rank values for each compound, and then generating the Spearman's Rank Correlation coefficient (Rs)
 25 according to the formula: $R_s = 1 - (6 \cdot SS / (n(n^2 - 1)))$, where n is the number of compounds ranked (96 in this case). Rs is then compared to a Table of critical values to determine statistical significance (P values).

Example 58.6. Summary of Results:

30 This table summarises the Rs and P values generated as a result of the indicated pairwise comparisons between rank orders.

-115-

Table 20

Comparison		Rs	P	+ive correlation
Plaque at 10 μ M	Plaque at 1 μ M	0.689	>99.5	Yes
Bacterial	Plaque at 10 μ M	0.444	>99	Yes
	Plaque at 1 μ M	0.406	>98.5	Yes
	<i>Randomised order</i>	-0.382	n/s	No

Example 58.7. Conclusions:

The rank order comparison of 96 compounds assayed in the bacterial bio-
 5 assay and the antiviral assay show that MHVE bacterial assay rank order for the
 compounds tested is significantly positively correlated with the rank orders generated
 by the MHV plaque reduction assay. The significant correlation between the assays
 is highly indicative that either assay may be utilised to identify compounds that may
 be useful. The bacterial assay may thereby be a useful tool in screening for
 10 compounds that exhibit anti-viral activity.

Example 58.8. Correlation Between Percent inhibition of 229E plaque formation and 229E-E bacterial bio-assay score.

A positive correlation was seen between the activity scores assigned to
 15 compounds when tested in the Human Coronavirus 229E E-protein bacterial bio-
 assay and the percent inhibition exhibited by these compounds in the Human
 Coronavirus 229E plaque assay.

Example 58.9. Method:

20 229E plaque reduction activity data for 97 compounds screened against 2.5 μ M
 compound concentration were sorted from greatest to least percent plaque reduction and
 rank orders were assigned to the list of compounds. Similarly, a rank order list was
 generated for the 229E E bacterial bioassay scores for the same 97 compounds. Where one
 or more compounds had the same score, the rank values for that group were averaged.

25

Spearman's statistical test for [as described in "Mathematical Statistics with Applications"
 (2nd edn): Mendenhall, W., Scheaffer, R.L., & Wackerly, D.D. Duxbury Press, Boston

-116-

Massachusetts – 1981] was used to compare rank orders. Briefly, this involves calculating the Sum of squares (SS) of the differences between rank values for each compound, and then generating the Spearman's Rank Correlation coefficient (Rs) according to the formula: $R_s = 1 - (6 \cdot SS / (n(n^2 - 1)))$, where n is the number of compounds ranked (97 in this case). Rs is then compared to a Table of critical values to determine statistical significance (P values).

Example 58.9.1. Summary of Results and Conclusions

10 This table summarises the Rs and P values generated as a result of the indicated pairwise comparisons between rank orders.

Table 21

Comparison		Rs	P	+ive correlation
Plaque at 2.5µM	Bacterial	0.584	>99.5	Yes
	<i>randomised</i>	-0.382	<i>n/s</i>	<i>No</i>

The results above indicate that the 229E bacterial assay rank order for the compounds tested is significantly positively correlated with the rank orders generated by the 229E plaque reduction assay. This result combined with that shown in Examples 49.1 and 49.4, provides strong evidence that either assay may be utilised to identify compounds that may be useful. The bacterial assay may thereby be a useful tool in screening for compounds that exhibit anti-viral activity.

20

Those skilled in the art will appreciate that the invention described herein is susceptible to variations and modifications other than those specifically described. It is to be understood that the invention includes all such variations and modifications. The invention also includes all of the steps, features, compositions and compounds referred to or indicated in this specification, individually or collectively, and any and all combinations of any two or more of said steps or features.

25

BIBLIOGRAPHY

- Barry, M., Mulcahy, F. and Back, D - J., *Br J Clin Pharmacol*, 45: 221 (1998)
- Deeks, S.G., *West J Med*, 168:133 (1998)
- Miles, S.A., *J Acquir Immune Defic Syndr Hum Retrovirol*, 16 Suppl 1: S36 (1997)
- 5 Miles, S.A., *J. Acquir Immune Defic Syndr Hum Retrovirol*, 16 Suppl 1: S 1 (1998)
- Moyle, G.J., Gazzard, B.G., Cooper, D.A. and Gatell, J., *Drugs*, 55:383 (1998)
- Rachlis, A.R. and Zarowny, D.P., *Cmaj*, 158:496 (1998)
- Vella, S., Fragola, V. and Palmisano, L., *J Biol Regul Homeost Agents*, 11:60 (1997)
- Volberding, P.A. and Deeks, S.G., *Jama*, 279:1343 (1998)
- 10 Volberding, P.A., *Hosp Pract (Off Ed)*, 33:81-4, 87-90, 95-6 passim. (1998)
- Miller, R.H. and Sarver, N., *Nat Med*, 3:389 (1997)
- Mitsuya, H., *Enzyme Inhibition*, 6:1 (1992)
- Moore, J.P., *Science*, 276:51 (1997)
- Thomas, M. and Brady, L., *Trends Biotechnol.*, 15:167 (1997)
- 15 Balliet, J.W., Kolson, D.L., Eiger, G., Kim, F.M., McGann, K.A., Srinivasan, A. and Collman, R., *Virology*, 200:623 (1994)
- Westervelt, P., Henkel, T., Trowbridge, D.B., Orenstein, J., Heuser, J., Gendelman, H.E. and Ratner, L., *J Virol*, 66:3925 (1992)
- Ewart, G.D., Sutherland, T., Gage, P.W. and Cox, G.B., *J Virol*, 70:7108 (1996)
- 20 Schubert, U., Henklein, P., Boldyreff, B., Wingender, E., Strebel, K. and Porstmann, T. *J Mol Biol*, 236:16 (1994)
- Friberg, J., Ladha, A., Gottlinger, H., Haseltine, W.A. and Cohen, E.A., *Journal of Acquired Immune Deficiency Syndromes & Human Retrovirology*, 8:10 (1995)
- Fu, J., Tan, B.H., Yap, E.H., Chan, Y.C. and Tan, Y.H. (1992) Full-length cDNA
- 25 sequence of dengue type 1 virus (Singapore strain S275/90). *Virology* 188 (2), 953-958
- Love, C.A., Lilley, P.E. and Dixon, N.E., *Escherichia coli*. Gene, in press. (1996)
- Yamato, I., Kotani, M., Oka, Y. and Anraku, Y., *Escherichia coli*. *Journal of Biological Chemistry*, 269:5729 (1994)
- 30 Rosen, B.R., ATP-coupled solute transport systems. *Escherichia coli and Salmonella typhimurium: Cellular and molecular biology* 1: (1987) Editor: Neidhardt, F.C.

-118-

American Society for Microbiology.

Piller, S.C., Ewart, G.D., Premkumar, A., Cox, G.B. and Gage, P.W., Proceedings of the National Academy of Sciences of the United States of America, 93:111 (1996)

Lu, Y.A., Clavijo, P., Galantino, M., Shen, Z.Y., Liu, W. and Tam, J.P., Molecular
5 Immunology, 28:623 (1991)

Harlow, E. and Lane, D., (1988) Antibodies: A laboratory manual. (ed). Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., 'Vol:'.

Varadhachary, A. and Maloney, P.-C., Molecular Microbiology, 4:1407 (1990) - 44

New, R.C.C., (1990). Liposomes: A practical approach. The Practical Approach
10 Series.

Kuhn RJ, Zhang W, Rossmann MG, Pletnev SV, Corver J, Lenches E, Jones CT, Mukhopadhyay S, Chipman PR, Strauss EG, Baker TS, Strauss JH. (2002). Structure of dengue virus: implications for flavivirus organization, maturation, and fusion. Cell. 2002 Mar 8;108(5):717-25.

15 Rickwood, D., Harries, B. D. (eds). IRL Press, Oxford.

Miller, C., (1986) Ion channel reconstitution. (ed). Plenum Press, New York and London.

Fear, W.R., Kesson, A.M., Naif, H., Lynch, G.W. and Cunningham, A.L., J. Virol, 72:1334 (1998)

20 Kelly, M.D., Naif, H., Adams, S.L., Cunningham, A.L. and Lloyd, A.R., J. Immunol, 160:3091 (1998)

New, R.C.C. (ed.), Liposomes: a practical approach. IRL Press, Oxford (1990) Grice,

A.L., Kerr, I.D. and Sansom, M.S., FEBS Lett, 405(3):299-304 (1997) Moore, P.B.,

Zhong, Q., Husslein, T. and Klein, M.L., FEBS Lett, 431(2):143-148 (1998)

25 Schubert, U., Bour, S., Ferrermonciel, A.V., Montal, M., Maldarelli, F. and Strebel, K., Journal of Virology, 70(2):809-819 (1996a)

Sunstrom NA, Premkumar LS, Premkumar A, Ewart G, Cox GB, Gage PW (1996),

Ion channels formed by NB, an influenza B virus protein. J Membr Biol. 1996

Mar;150(2):127-32


-119-

Willbold, D., Hoffmann, S. and Rosch, P., Eur J Biochem, 245(3):581-8 (1997)

Wray, V., Kinder, R., Federau, T., Henklein, P., Bechinger, B. and Schubert, U.,
Biochemistry, 38(16):5272-82 (1999)


THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A compound of Formula I




or pharmaceutically acceptable salts thereof,
wherein,

R₁ =




R₂, R₃ and R₄ are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkenyl, C₃₋₆cycloalkenyl, C₁₋₆alkeneoxy, or benzo;

- 10 R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,



R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;

- 15 R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein

when R_1 is $C_6H_5CH=CH$, R_2 is hydrogen and R_3 is phenyl, R_4 cannot be phenyl;

when R_1 is phenyl, R_2 is hydrogen, and R_3 is benzoyl, R_4 cannot be benzoyl;


when R_1 is phenyl, R_2 is substituted benzyl, R_3 is hydrogen and R_4 is hydrogen,


5 R_n , R_o and R_p cannot all be hydrogen;


when R_1 is phenyl, R_3 is hydrogen and R_4 is hydrogen, R_2 cannot be benzyl or phenyl;


when R_1 is phenyl, R_2 is hydrogen, R_3 cannot be phenyl together with R_4 as benzoyl;


10 when R_1 is phenyl, R_2 is hydrogen, R_3 and R_4 cannot both be benzyl;


when R_1 is , the compound is N-(3-phenylpropanoyl)-N'-phenylguanidine, N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine or N,N'-Bis(3-phenylpropanoyl)guanidine;


when R_1 is , the compound is cinnamoylguanidine;


15 when R_1 is  and R_i is hydrogen, the compound is trans-3-(1-naphthyl)acryloylguanidine;

20 when R_1 is , R_i is hydrogen and X is C_{1-6} alkyl, halo, halo-substituted C_{1-6} alkyl, hydrogen or nitro, the compound is 4-t-butylcinnamoylguanidine, 3-t-butylcinnamoylguanidine, 2-t-butylcinnamoylguanidine, 2-ethylcinnamoylguanidine, 4-isopropylcinnamoylguanidine, 3-isopropylcinnamoylguanidine hydrochloride, 4-methylcinnamoylguanidine, 3-fluorocinnamoylguanidine, 2-fluorocinnamoylguanidine, 2-(trifluoromethyl)cinnamoylguanidine or 3-(trifluoromethyl)cinnamoylguanidine;



when R_1 is , R_g is hydrogen and any one of R_d , R_e or R_f is halo or alkoxy, the compound is 5-bromo-2-methoxycinnamoylguanidine, 5-bromo-2-fluorocinnamoylguanidine or 3,4-difluorocinnamoylguanidine; and




when R_1 is  R_k hydrogen and R_j is hydrogen or C_{1-5} alkoxy, the compound is N-(2-naphthoyl)-N'-phenylguanidine, or N,N'-bis(2-naphthoyl)guanidine.

2. A pharmaceutical composition comprising an antiviral compound according to claim 1 and optionally one or more pharmaceutical acceptable carriers or derivatives.


3. The pharmaceutical composition according to claim 2, further comprising one or more known antiviral compounds or molecules.

4. Use of a compound according of Formula I




or pharmaceutically acceptable salts thereof,
wherein,

$R_1 =$




R_2 , R_3 and R_4 are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{3-6} cycloalkyl, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkyloxy, phenyl, C_{1-6} alkeneyl, C_{3-6} cycloalkeneyl, C_{1-6} alkeneoxy, or benzo;

- R_a , R_b , R_c , R_d , R_e , R_f , R_h , R_k , R_L , R_m , R_n , R_o , R_p independently =
hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl,
10 substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g , R_i independently = hydrogen, hydroxy, halo, or C_{1-5} alkyl;

- R_j = hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino,
15 aryl-substituted amino, PrS,



and wherein



when R₁ is R_c, R_a and R_c are amino, and R_b is halo, R₂, R₃ or R₄ cannot be hydrogen, benzyl or substituted benzyl or BODIPY-Fl;

when R₁ is C₆H₅CH=CH, R₂ is hydrogen and R₃ is phenyl, R₄ cannot be phenyl;

5 when R₁ is phenyl, R₂ is hydrogen, and R₃ is benzoyl, R₄ cannot be benzoyl;

when R₁ is phenyl, R₂ is substituted benzyl, R₃ is hydrogen and R₄ is hydrogen, R_n, R_o and R_p cannot all be hydrogen;

when R₁ is phenyl, R₃ is hydrogen and R₄ is hydrogen, R₂ cannot be benzyl or phenyl;

10 when R₁ is phenyl, R₂ is hydrogen, R₃ cannot be phenyl together with R₄ as benzoyl; and

when R₁ is phenyl, R₂ is hydrogen, R₃ and R₄ cannot both be benzyl,

for the manufacture of a medicament for reducing, retarding or otherwise inhibiting growth and/or replication of a virus.

15

5. The use according to claim 4, wherein said virus is a Lentivirus.

6. The use according to claim 5, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

20

7. The use according to claim 6, wherein said compound is selected from the group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,

trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoyleguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
4-isopropylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
N-(cinnamoyl)-N'phenylguanidine,
2-(cyclohex-1-en-1yl)cinnamoylguanidine,
2-(2-naphthyl)acetoyleguanidine,
(4-Hydroxycinnamoyl)guanidine,

4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamolyguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (a-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy-HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

8. The use according to claim 6, wherein said compound is selected from the group consisting of 4-phenylbenzoylguanidine, (3-bromocinnamoyl)guanidine, 3-(trifluoromethyl)cinnamoylguanidine, 5-(N,N-hexamethylene)amiloride, and
- 5 (5-Phenyl-penta-2,4-dienoyl)guanidine.

9. The use according to any one of claims 6 to 8, wherein said HIV is HIV-1.

10. The use according to claim 4, wherein said virus is a Coronavirus.

11. The use according to claim 10, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).

5

12. The use according to claim 11, wherein said compound is selected from the group consisting of

2,3-difluorocinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
(3-Chlorocinnamoyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2,5-dimethylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-isopropylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
6-methoxy-2-naphthoylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
3-phenylcinnamoylguanidine,
(2-Chlorocinnamoyl)guanidine,
2'4 DichloroBenazamil HCl
4-phenylcinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
cinnamoylguanidine hydrochloride,
4-ethoxycinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
5-tert-butylamino-amiloride,
3-t-butylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
6-Iodoamiloride,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,

(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,
1-bromo-2-naphthoylguanidine,

2-chloro-6-fluorocinnamoylguanidine,
 [(4-Chlorophenoxy-acetyl]guanidine,
 Phenamil methanesulfonate salt,
 N-Benzoyl-N'-cinnamoylguanidine and
 N-(2-naphthoyl)-N'-phenylguanidine.

13. The use according to claim 11, wherein said compound is selected from the group consisting of cinnamoylguanidine, trans-3-(1-naphthyl)acryloylguanidine, and 6-methoxy-2-naphthoylguanidine.

5

14. The use according to claim 10, wherein said Coronavirus is human Coronavirus 229E.

15. The use according to claim 14, wherein said compound is selected from the group
 10 consisting of

4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 4-t-butylcinnamoylguanidine,
 3-isopropylcinnamoylguanidine hydrochloride,
 3-t-butylcinnamoylguanidine,
 2-t-butylcinnamoylguanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 2,3-difluorocinnamoylguanidine,
 3-(2-naphthyl)acryloylguanidine,
 2-phenylcinnamoylguanidine,
 3-phenylcinnamoylguanidine,
 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 4-phenylbenzoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 1-naphthoylguanidine,
 5-(4-fluorophenyl)amiloride,
 (5-Phenyl-penta-2,4-dienoyl)guanidine,
 (3-Bromocinnamoyl)guanidine,

2,5-dimethylcinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
5-bromo-2-fluorocinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
(α -Methylcinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,4,6-trimethylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(1-naphthyl)acetylguanidine,
2-ethylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
2-ethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
3-fluorocinnamoylguanidine,
cinnamoylguanidine hydrochloride,
2,3-dimethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-tert-butylamino-amiloride,
2-naphthoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
N,N'-Bis(3-phenylpropanoyl)guanidine,
4-methylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
2,3,5,6-tetramethylcinnamoylguanidine,
3-ethoxycinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(2-naphthyl)acetylguanidine,

N-(3-phenylpropanoyl)-N'-phenylguanidine,
 5-(2'-bromophenyl)penta-2,4-
 dienoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,
 (4-Methoxycinnamoyl)guanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 [(E)-3-(4-Dimethylaminophenyl)-2-
 methylacryloyl]guanidine,
 N-Benzoyl-N'-cinnamoylguanidine,
 4-phenylbenzoylguanidine,
 trans-3-Furanacryoylguanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 Pyrazinecarboxamide,
 N-(cinnamoyl)-N'phenylguanidine,
 Cinnamoylguanidine,
 3-methoxy-amiloride,
 (3-phenylpropanoyl)guanidine,
 3-methoxy -HMA,
 Benzyoylguanidine,
 N-amidino-3,5-diamino-6-phynyl-2-
 Pyrazinecarboxamide,
 (Quinoline-2-carbonyl)guanidine,
 [3-(3-Pyridyl)acryloyl]guanidine,
 N-Cinnamoyl-N',N'-dimethylguanidine,
 N-(2-naphthoyl)-N'-phenylguanidine and
 (Phenylacetyl)guanidine.

16. The use according to claim 14, wherein said compound is selected from the group consisting of

2-t-butylcinnamoylguanidine,
 4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 2,6-dichlorocinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-cyclohexylcinnamoylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 2-phenylcinnamoylguanidine,
 4-t-butylcinnamoylguanidine,
 3-phenylcinnamoylguanidine,
 (3-Bromocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 trans-3-(1-naphthyl)acryloylguanidine,

3-(2-naphthyl)acryloylguanidine,
2,4-dichlorocinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
4-methylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(α -Methylcinnamoyl)guanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-t-butylcinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
3-methylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(2-Bromocinnamoyl)guanidine,
3-ethoxycinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-
Phenylguanidine,
2,4,6-trimethylcinnamoylguanidine,
2-methylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)-
guanidine,
(4-Phenoxybenzoyl)guanidine,
(2-Methoxycinnamoyl)guanidine,
Cinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-
Dicarboxamide,
2,3-dimethylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
N,N'-Bis(3-phenylpropanoyl)guanidine,
2,3-difluorocinnamoylguanidine,
1-naphthoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-ethoxycinnamoylguanidine,
2-naphthoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,

2-(trifluoromethyl)cinnamoylguanidine,
 cinnamoylguanidine hydrochloride,
 (4-Hydroxycinnamoyl)guanidine,
 5-(4-fluorophenyl)amiloride,
 2-(1-naphthyl)acetoylguanidine,
 (2-Furanacryloyl)guanidine,
 N-Cinnamoyl-N',N'-dimethylguanidine,
 2-(2-naphthyl)acetoylguanidine and
 N,N'-bis(3phenylpropanoyl)-N''-
 Phenylguanidine.

17. The use according to claim 10, wherein said Coronavirus is human Coronavirus OC43.

5 18. The use according to claim 17, wherein said compound is selected from the group consisting of

3-methylcinnamoylguanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 (3-Bromocinnamoyl)guanidine,
 (2-Chlorocinnamoyl)guanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 4-isopropylcinnamoylguanidine,
 Cinnamoylguanidine,
 6-methoxy-2-naphthoylguanidine,
 2,4-dichlorocinnamoylguanidine,
 (4-Chlorocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 (4-Bromocinnamoyl)guanidine,
 2,6-dichlorocinnamoylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 (5-Phenyl-penta-2,4-dienoyl)guanidine,
 3-(trifluoromethoxy)cinnamoylguanidine and
 2-t-butylcinnamoylguanidine.

19. The use according to claim 10, wherein said Coronavirus is porcine respiratory Coronavirus (PRCV).

20. The use according to claim 19, wherein said compound is selected from the group consisting of

5-(N,N-hexamethylene)amiloride,
6-methoxy-2-naphthoylguanidine,
Cinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
3-methylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
trans-3-(1-naphthyl)acryloylguanidine and
2-(2-naphthyl)acetoylguanidine.

21. The use according to claim 10, wherein said Coronavirus is bovine Coronavirus
5 (BCV).

22. The use according to claim 21, wherein said compound is selected from the group consisting of

(3-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-hexamethylene)amiloride,
trans-3-(1-naphthyl)acryloylguanidine,
Cinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine and
4-phenylbenzoylguanidine.

10 23. The use according to claim 10, wherein said Coronavirus is any one of the known
coronavirus isolates selected from the group consisting of canine enteric coronavirus
(strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus
(strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99,
porcine respiratory coronavirus, porcine transmissible gastroenteritis coronavirus
15 (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller),
porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine
transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus
(STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9),

bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus (STRAIN LY-138),
 bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine
 coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine
 coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-
 5 ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus
 (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus (strain JHM),
 murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus
 strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis
 virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N),
 10 porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat
 coronavirus (strain 681), rat coronavirus (strain NJ), rat sialodacryoadenitis coronavirus,
 turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey
 coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian
 infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain
 15 Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious
 bronchitis virus (strain Beaudette US), avian infectious bronchitis virus (strain
 Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis
 virus (strain D274), avian infectious bronchitis virus (strain D3896). avian infectious
 bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian
 20 infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120),
 avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain
 KB8523), avian infectious bronchitis virus (strain M41), avian infectious bronchitis virus
 (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian
 infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain
 25 UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious
 bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84),
 avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain
 Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003,
 SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS
 30 coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1,
 SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-

39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003,
 5 SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani, SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus
 10 (strain CV777).

24. The use according to claim 23, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 4-t-butylcinnamoylguanidine, and
 3-isopropylcinnamoylguanidine hydrochloride.

15 25. The use according to claim 4, wherein said virus is the Hepatitis C virus.

26. The use according to claim 25, wherein said compound is selected from the group consisting of

2,3-dimethylcinnamoylguanidine,
 2,4,6-trimethylcinnamoylguanidine,
 5-bromo-2-fluorocinnamoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 2,5-dimethylcinnamoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 6-methoxy-2-naphthoylguanidine,
 (2-Chlorocinnamoyl)guanidine,
 (4-Chlorocinnamoyl)guanidine,
 (2-Bromocinnamoyl)guanidine,
 2,6-dichlorocinnamoylguanidine,
 (3-Bromocinnamoyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,

2-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'-phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
3-ethoxycinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
4-phenylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(2-Furanacryloyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,
5-(N,N-hexamethylene)amiloride,
2,3-difluorocinnamoylguanidine,
2-(1-naphthyl)acetoyleguanidine,
(a-Methylcinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
6-Iodoamiloride,

3,4-(methylenedioxy)cinnamoylguanidine,
 2-ethoxycinnamoylguanidine,
 Cinnamoylguanidine,
 2-phenylcinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-naphthoylguanidine,
 3-phenylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 5-(4-fluorophenyl)amiloride,
 (3-Methoxycinnamoyl)guanidine,
 2-fluorocinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 (3-phenylpropanoyl)guanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 2-methylcinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 1-bromo-2-naphthoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 Phenamil methanesulfonate salt,
 2,4-dichlorocinnamoylguanidine,
 3,4-difluorocinnamoylguanidine and
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.


27. The use according to claim 4, wherein said virus is Equine Arteritis virus.

28. The use according to claim 27, wherein said compound is selected from the group
 5 consisting of


5-(N,N-hexamethylene)amiloride,
 (3-Bromocinnamoyl)guanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 2-t-butylcinnamoylguanidine and
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine.

29. The use according to any one of claims 4 to 28, wherein said medicament further comprises one or more known antiviral compounds or molecules.


10 30. Use of a compound according of Formula I



or pharmaceutically acceptable salts thereof,
wherein,




R_2 , R_3 and R_4 are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkeneyl, C₃₋₆cycloalkeneyl, C₁₋₆alkeneoxy, or benzo;

- 10 R_a, R_b, R_c, R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;

- 15 R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein



when R_1 is R_c , R_a and R_c are amino, and R_b is halo, R_2 , R_3 or R_4 cannot be hydrogen, benzyl or substituted benzyl or BODIPY-F1;

when R_1 is $C_6H_5CH=CH$, R_2 is hydrogen and R_3 is phenyl, R_4 cannot be phenyl;

5 when R_1 is phenyl, R_2 is hydrogen, and R_3 is benzoyl, R_4 cannot be benzoyl;

when R_1 is phenyl, R_2 is substituted benzyl, R_3 is hydrogen and R_4 is hydrogen, R_n , R_o and R_p cannot all be hydrogen;

when R_1 is phenyl, R_3 is hydrogen and R_4 is hydrogen, R_2 cannot be benzyl or phenyl;

10 when R_1 is phenyl, R_2 is hydrogen, R_3 cannot be phenyl together with R_4 as benzoyl; and

when R_1 is phenyl, R_2 is hydrogen, R_3 and R_4 cannot both be benzyl,

for the manufacture of a medicament for preventing the infection of a cell exposed to a virus.

15

31. The use according to claim 30, wherein said virus is a Lentivirus.

32. The use according to claim 31, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

20

33. The use according to claim 31, wherein said compound is selected from the group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,

trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoylguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
4-isopropylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
N-(cinnamoyl)-N'phenylguanidine,
2-(cyclohex-1-en-1yl)cinnamoylguanidine,
2-(2-naphthyl)acetoylguanidine,
(4-Hydroxycinnamoyl)guanidine,

4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamolyguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (α-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy-HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

34. The use according to claim 32, wherein said compound is selected from the group consisting of 4-phenylbenzoylguanidine, (3-bromocinnamoyl)guanidine, 3-(trifluoromethyl)cinnamoylguanidine, 5-(N,N-hexamethylene)amiloride, and
- 5 (5-Phenyl-penta-2,4-dienoyl)guanidine.

35. The use according to any one of claims 32 to 34, wherein said HIV is HIV-1.

36. The use according to claim 30, wherein said virus is a Coronavirus.

37. The use according to claim 36, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).

5

38. The use according to claim 37, wherein said compound is selected from the group consisting of

2,3-difluorocinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
(3-Chlorocinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2,5-dimethylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-isopropylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
6-methoxy-2-naphthoylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
3-phenylcinnamoylguanidine,
(2-Chlorocinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
cinnamoylguanidine hydrochloride,
4-ethoxycinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
5-tert-butylamino-amiloride,
3-t-butylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
6-Iodoamiloride,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,

(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,
1-bromo-2-naphthoylguanidine,
2-chloro-6-fluorocinnamoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,

Phenamil methanesulfonate salt,
N-Benzoyl-N'-cinnamoylguanidine and
N-(2-naphthoyl)-N'-phenylguanidine.

39. The use according to claim 37, wherein said compound is selected from the group consisting of cinnamoylguanidine, trans-3-(1-naphthyl)acryloylguanidine, and 6-methoxy-2-naphthoylguanidine.

5

40. The use according to claim 36, wherein said Coronavirus is human Coronavirus 229E.

41. The use according to claim 40, wherein said compound is selected from the group
10 consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
3-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2,3-difluorocinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-phenylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
4-phenylbenzoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
1-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,

6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
5-bromo-2-fluorocinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,4,6-trimethylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(1-naphthyl)acetoyleguanidine,
2-ethylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
2-ethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
3-fluorocinnamoylguanidine,
cinnamoylguanidine hydrochloride,
2,3-dimethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-tert-butylamino-amiloride,
2-naphthoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
N,N'-Bis(3-phenylpropanoyl)guanidine,
4-methylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-ethoxycinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(2-naphthyl)acetoyleguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
5-(2'-bromophenyl)penta-2,4-

dienoylguanidine,
(4-Bromocinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
(4-Methoxycinnamoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine,
N-Benzoyl-N'-cinnamoylguanidine,
4-phenylbenzoylguanidine,
trans-3-Furanacryoylguanidine,
N-amidino-3-amino-5-phenyl-6-chloro-2-Pyrazinecarboxamide,
N-(cinnamoyl)-N'phenylguanidine,
Cinnamoylguanidine,
3-methoxy-amiloride,
(3-phenylpropanoyl)guanidine,
3-methoxy -HMA,
Benzyoylguanidine,
N-amidino-3,5-diamino-6-phynyl-2-Pyrazinecarboxamide,
(Quinoline-2-carbonyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
N-Cinnamoyl-N',N'-dimethylguanidine,
N-(2-naphthoyl)-N'-phenylguanidine and
(Phenylacetyl)guanidine.

42. The use according to claim 40, wherein said compound is selected from the group consisting of

2-t-butylcinnamoylguanidine,
4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2-phenylcinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
trans-3-(1-naphthyl)acryloylguanidine,
3-(2-naphthyl)acryloylguanidine,
2,4-dichlorocinnamolyguanidine,

3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
4-methylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(α -Methylcinnamoyl)guanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-t-butylcinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
3-methylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(2-Bromocinnamoyl)guanidine,
3-ethoxycinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-
Phenylguanidine,
2,4,6-trimethylcinnamoylguanidine,
2-methylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)-
guanidine,
(4-Phenoxybenzoyl)guanidine,
(2-Methoxycinnamoyl)guanidine,
Cinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-
Dicarboxamide,
2,3-dimethylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
N,N'-Bis(3-phenylpropanoyl)guanidine,
2,3-difluorocinnamoylguanidine,
1-naphthoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-ethoxycinnamoylguanidine,
2-naphthoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,

(4-Hydroxycinnamoyl)guanidine,
 5-(4-fluorophenyl)amiloride,
 2-(1-naphthyl)acetoylguanidine,
 (2-Furanacryloyl)guanidine,
 N-Cinnamoyl-N',N'-dimethylguanidine,
 2-(2-naphthyl)acetoylguanidine and
 N,N'-bis(3phenylpropanoyl)-N''-
 Phenylguanidine.

43. The use according to claim 36, wherein said Coronavirus is human Coronavirus OC43.

5 44. The use according to claim 43, wherein said compound is selected from the group consisting of

3-methylcinnamoylguanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 (3-Bromocinnamoyl)guanidine,
 (2-Chlorocinnamoyl)guanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 4-isopropylcinnamoylguanidine,
 Cinnamoylguanidine,
 6-methoxy-2-naphthoylguanidine,
 2,4-dichlorocinnamoylguanidine,
 (4-Chlorocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 (4-Bromocinnamoyl)guanidine,
 2,6-dichlorocinnamoylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 (5-Phenyl-penta-2,4-dienoyl)guanidine,
 3-(trifluoromethoxy)cinnamoylguanidine and
 2-t-butylcinnamoylguanidine.

45. The use according to claim 36, wherein said Coronavirus is porcine respiratory Coronavirus (PRCV).

10

46. The use according to claim 45, wherein said compound is selected from the group consisting of

5-(N,N-hexamethylene)amiloride,
6-methoxy-2-naphthoylguanidine,
Cinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
3-methylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
trans-3-(1-naphthyl)acryloylguanidine and
2-(2-naphthyl)acetoyleguanidine.

47. The use according to claim 36, wherein said Coronavirus is bovine Coronavirus (BCV).

5 48. The use according to claim 47, wherein said compound is selected from the group consisting of

(3-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-hexamethylene)amiloride,
trans-3-(1-naphthyl)acryloylguanidine,
Cinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
2-(2-naphthyl)acetoyleguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine and
4-phenylbenzoylguanidine.

49. The use according to claim 36, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric coronavirus
10 (strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller), porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine
15 transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine

coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus (strain JHM),

5 murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N), porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat coronavirus (strain 681), rat coronavirus (strain NJ), rat sialodacryoadenitis coronavirus,

10 turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious bronchitis virus (strain Beaudette US), avian infectious bronchitis virus (strain

15 Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis virus (strain D274), avian infectious bronchitis virus (strain D3896). avian infectious bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120), avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain

20 KB8523), avian infectious bronchitis virus (strain M41), avian infectious bronchitis virus (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84),

25 avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003, SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1, SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-

30 39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS

coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003, SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani,
 5 SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus (strain CV777).

10 50. The use according to claim 49, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 4-t-butylcinnamoylguanidine, and
 3-isopropylcinnamoylguanidine hydrochloride.

51. The use according to claim 30, wherein said virus is the Hepatitis C virus.

15 52. The use according to claim 51, wherein said compound is selected from the group consisting of

2,3-dimethylcinnamoylguanidine,
 2,4,6-trimethylcinnamoylguanidine,
 5-bromo-2-fluorocinnamoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 2,5-dimethylcinnamoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 6-methoxy-2-naphthoylguanidine,
 (2-Chlorocinnamoyl)guanidine,
 (4-Chlorocinnamoyl)guanidine,
 (2-Bromocinnamoyl)guanidine,
 2,6-dichlorocinnamoylguanidine,
 (3-Bromocinnamoyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,
 2-(trifluoromethyl)cinnamoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3,4-dichlorocinnamoylguanidine,

4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'-phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
3-ethoxycinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
4-phenylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(2-Furanacryloyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,
5-(N,N-hexamethylene)amiloride,
2,3-difluorocinnamoylguanidine,
2-(1-naphthyl)acetoyleguanidine,
(a-Methylcinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
6-Iodoamiloride,
3,4-(methylenedioxy)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
Cinnamoylguanidine,

2-phenylcinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-naphthoylguanidine,
 3-phenylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 5-(4-fluorophenyl)amiloride,
 (3-Methoxycinnamoyl)guanidine,
 2-fluorocinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 (3-phenylpropanoyl)guanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 2-methylcinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 1-bromo-2-naphthoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 Phenamil methanesulfonate salt,
 2,4-dichlorocinnamoylguanidine,
 3,4-difluorocinnamoylguanidine and
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.


53. The use according to claim 30, wherein said virus is the Equine Arteritis virus.


54. The use according to claim 53, wherein said compound is selected from the group
 5 consisting of

5-(N,N-hexamethylene)amiloride,
 (3-Bromocinnamoyl)guanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 2-t-butylcinnamoylguanidine and
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine.

55. The use according to any one of claims 30 to 54, wherein said medicament further comprises one or more known antiviral compounds or molecules.

10 56. Use of a compound according of Formula I






and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkeneyl, C₃₋₆cycloalkeneyl, C₁₋₆alkeneoxy, or benzo;

- R_a, R_b, R_c, R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently =
hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl,
10 substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;

- R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted
15 aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein



when R₁ is R_c, R_a and R_c are amino, and R_b is halo, R₂, R₃ or R₄ cannot be hydrogen, benzyl or substituted benzyl or BODIPY-Fl;

when R₁ is C₆H₅CH=CH, R₂ is hydrogen and R₃ is phenyl, R₄ cannot be phenyl;

5 when R₁ is phenyl, R₂ is hydrogen, and R₃ is benzoyl, R₄ cannot be benzoyl;

when R₁ is phenyl, R₂ is substituted benzyl, R₃ is hydrogen and R₄ is hydrogen, R_n, R_o and R_p cannot all be hydrogen;

when R₁ is phenyl, R₃ is hydrogen and R₄ is hydrogen, R₂ cannot be benzyl or phenyl;

10 when R₁ is phenyl, R₂ is hydrogen, R₃ cannot be phenyl together with R₄ as benzoyl; and

when R₁ is phenyl, R₂ is hydrogen, R₃ and R₄ cannot both be benzyl,

for the manufacture of a medicament for the therapeutic or prophylactic treatment of a subject infected with or exposed to a virus.

15

57. The use according to claim 56, wherein said virus is a Lentivirus.

58. The use according to claim 57, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

20

59. The use according to claim 58, wherein said compound is selected from the group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,

trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoylguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
4-isopropylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
N-(cinnamoyl)-N'phenylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2-(2-naphthyl)acetoylguanidine,
(4-Hydroxycinnamoyl)guanidine,

4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamoylguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (α-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy -HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

60. The use according to claim 58, wherein said compound is selected from the group consisting of 4-phenylbenzoylguanidine, (3-bromocinnamoyl)guanidine, 3-(trifluoromethyl)cinnamoylguanidine, 5-(N,N-hexamethylene)amiloride, and
- 5 (5-Phenyl-penta-2,4-dienoyl)guanidine.

61. The use according to any one of claims 58 to 60, wherein said HIV is HIV-1.

62. The use according to claim 56, wherein said virus is a Coronavirus.

63. The use according to claim 62, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).

5

64. The use according to claim 63, wherein said compound is selected from the group consisting of

2,3-difluorocinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
(3-Chlorocinnamoyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2,5-dimethylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-isopropylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
6-methoxy-2-naphthoylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
3-phenylcinnamoylguanidine,
(2-Chlorocinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
4-phenylcinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
cinnamoylguanidine hydrochloride,
4-ethoxycinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
5-tert-butylamino-amiloride,
3-t-butylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
6-Iodoamiloride,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,

(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,
1-bromo-2-naphthoylguanidine,
2-chloro-6-fluorocinnamoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,

Phenamil methanesulfonate salt,
N-Benzoyl-N'-cinnamoylguanidine and
N-(2-naphthoyl)-N'-phenylguanidine.

65. The use according to claim 63, wherein said compound is selected from the group consisting of cinnamoylguanidine, trans-3-(1-naphthyl)acryloylguanidine, and 6-methoxy-2-naphthoylguanidine.

5

66. The use according to claim 62, wherein said Coronavirus is human Coronavirus 229E.

67. The use according to claim 66, wherein said compound is selected from the group
10 consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
3-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2,3-difluorocinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-phenylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
4-phenylbenzoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
1-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,

6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
5-bromo-2-fluorocinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,4,6-trimethylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(1-naphthyl)acetoyleguanidine,
2-ethylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
2-ethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
3-fluorocinnamoylguanidine,
cinnamoylguanidine hydrochloride,
2,3-dimethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-tert-butylamino-amiloride,
2-naphthoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
N,N'-Bis(3-phenylpropanoyl)guanidine,
4-methylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-ethoxycinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(2-naphthyl)acetoyleguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
5-(2'-bromophenyl)penta-2,4-

dienoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,
 (4-Methoxycinnamoyl)guanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine,
 N-Benzoyl-N'-cinnamoylguanidine,
 4-phenylbenzoylguanidine,
 trans-3-Furanacryoylguanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-Pyrazinecarboxamide,
 N-(cinnamoyl)-N'phenylguanidine,
 Cinnamoylguanidine,
 3-methoxy-amiloride,
 (3-phenylpropanoyl)guanidine,
 3-methoxy -HMA,
 Benzyoylguanidine,
 N-amidino-3,5-diamino-6-phynyl-2-Pyrazinecarboxamide,
 (Quinoline-2-carbonyl)guanidine,
 [3-(3-Pyridyl)acryloyl]guanidine,
 N-Cinnamoyl-N',N'-dimethylguanidine,
 N-(2-napthoyl)-N'-phenylguanidine and
 (Phenylacetyl)guanidine.

68. The use according to claim 66, wherein said compound is selected from the group consisting of

2-t-butylcinnamoylguanidine,
 4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 2,6-dichlorocinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-cyclohexylcinnamoylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 2-phenylcinnamoylguanidine,
 4-t-butylcinnamoylguanidine,
 3-phenylcinnamoylguanidine,
 (3-Bromocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 trans-3-(1-napthyl)acryloylguanidine,
 3-(2-napthyl)acryloylguanidine,
 2,4-dichlorocinnamolyguanidine,

3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
4-methylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(α -Methylcinnamoyl)guanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-t-butylcinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
3-methylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(2-Bromocinnamoyl)guanidine,
3-ethoxycinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-
Phenylguanidine,
2,4,6-trimethylcinnamoylguanidine,
2-methylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)-
guanidine,
(4-Phenoxybenzoyl)guanidine,
(2-Methoxycinnamoyl)guanidine,
Cinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-
Dicarboxamide,
2,3-dimethylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
N,N'-Bis(3-phenylpropanoyl)guanidine,
2,3-difluorocinnamoylguanidine,
1-naphthoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-ethoxycinnamoylguanidine,
2-naphthoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,

(4-Hydroxycinnamoyl)guanidine,
5-(4-fluorophenyl)amiloride,
2-(1-naphthyl)acetoylguanidine,
(2-Furanacryloyl)guanidine,
N-Cinnamoyl-N',N'-dimethylguanidine,
2-(2-naphthyl)acetoylguanidine and
N,N'-bis(3phenylpropanoyl)-N''-
Phenylguanidine.

69. The use according to claim 62, wherein said Coronavirus is human Coronavirus OC43.

5 70. The use according to claim 69, wherein said compound is selected from the group consisting of

3-methylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
4-isopropylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
2,4-dichlorocinnamolyguanidine,
(4-Chlorocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
(4-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-(trifluoromethoxy)cinnamoylguanidine and
2-t-butylcinnamoylguanidine.

71. The use according to claim 62, wherein said Coronavirus is porcine respiratory Coronavirus (PRCV).

10

72. The use according to claim 71, wherein said compound is selected from the group consisting of

5-(N,N-hexamethylene)amiloride,
6-methoxy-2-naphthoylguanidine,
Cinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
3-methylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
trans-3-(1-naphthyl)acryloylguanidine and
2-(2-naphthyl)acetoylguanidine.

73. The use according to claim 62, wherein said Coronavirus is bovine Coronavirus (BCV).

5 74. The use according to claim 73, wherein said compound is selected from the group consisting of

(3-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
5-(N,N-hexamethylene)amiloride,
trans-3-(1-naphthyl)acryloylguanidine,
Cinnamoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine and
4-phenylbenzoylguanidine.

75. The use according to claim 62, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric coronavirus
10 (strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller), porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine
15 transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine

coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine
 coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-
 ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus
 (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus (strain JHM),
 5 murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus
 strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis
 virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N),
 porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat
 coronavirus (strain 681), rat coronavirus (strain NJ), rat sialodacryoadenitis coronavirus,
 10 turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey
 coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian
 infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain
 Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious
 bronchitis virus (strain Beaudette US), avian infectious bronchitis virus (strain
 15 Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis
 virus (strain D274), avian infectious bronchitis virus (strain D3896). avian infectious
 bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian
 infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120),
 avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain
 20 KB8523), avian infectious bronchitis virus (strain M41), avian infectious bronchitis virus
 (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian
 infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain
 UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious
 bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84),
 25 avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain
 Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003,
 SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS
 coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1,
 SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-
 30 39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong
 Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS

- coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003, SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani,
- 5 SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus (strain CV777).
- 10 76. The use according to claim 75, wherein said compound is selected from the group consisting of
- 4-isopropylcinnamoylguanidine,
 - 3,4-dichlorocinnamoylguanidine,
 - 3-(trifluoromethoxy)cinnamoylguanidine,
 - 4-t-butylcinnamoylguanidine, and
 - 3-isopropylcinnamoylguanidine hydrochloride.
77. The use according to claim 56, wherein said virus is the Hepatitis C virus.
- 15 78. The use according to claim 77, wherein said compound is selected from the group consisting of
- 2,3-dimethylcinnamoylguanidine,
 - 2,4,6-trimethylcinnamoylguanidine,
 - 5-bromo-2-fluorocinnamoylguanidine,
 - (4-Bromocinnamoyl)guanidine,
 - 2,5-dimethylcinnamoylguanidine,
 - 3-(trifluoromethyl)cinnamoylguanidine,
 - 4-(trifluoromethyl)cinnamoylguanidine,
 - 6-methoxy-2-naphthoylguanidine,
 - (2-Chlorocinnamoyl)guanidine,
 - (4-Chlorocinnamoyl)guanidine,
 - (2-Bromocinnamoyl)guanidine,
 - 2,6-dichlorocinnamoylguanidine,
 - (3-Bromocinnamoyl)guanidine,
 - (3-Chlorocinnamoyl)guanidine,
 - 2-(trifluoromethyl)cinnamoylguanidine,
 - (4-Phenoxybenzoyl)guanidine,
 - 3,4-dichlorocinnamoylguanidine,

4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
3-ethoxycinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
4-phenylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
N-(6-Hydroxy-2-naphthyl)-N'-phenylguanidine,
(2-Furanacryloyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,
5-(N,N-hexamethylene)amiloride,
2,3-difluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
(a-Methylcinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
6-Iodoamiloride,
3,4-(methylenedioxy)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
Cinnamoylguanidine,

2-phenylcinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-naphthoylguanidine,
 3-phenylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 5-(4-fluorophenyl)amiloride,
 (3-Methoxycinnamoyl)guanidine,
 2-fluorocinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 (3-phenylpropanoyl)guanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 2-methylcinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 1-bromo-2-naphthoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 Phenamil methanesulfonate salt,
 2,4-dichlorocinnamoylguanidine,
 3,4-difluorocinnamoylguanidine and
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.


79. The use according to claim 56, wherein said virus is the Equine Arteritis virus.

80. The use according to claim 79, wherein said compound is selected from the group
5 consisting of


5-(N,N-hexamethylene)amiloride,
 (3-Bromocinnamoyl)guanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 2-t-butylcinnamoylguanidine and
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine.

81. The use according to any one of claims 56 to 80, wherein said medicament further comprises one or more known antiviral compounds or molecules.


10 82. Use of a compound according of Formula I



or pharmaceutically acceptable salts thereof,
wherein,




R_2 , R_3 and R_4 are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkeneyl, C₃₋₆cycloalkeneyl, C₁₋₆alkeneoxy, or benzo;

- R_a, R_b, R_c, R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently =
hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl,
10 substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;

- R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted
15 aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein



when R_1 is R_c , R_a and R_c are amino, and R_b is halo, R_2 , R_3 or R_4 cannot be hydrogen, benzyl or substituted benzyl or BODIPY-F1;

when R_1 is $C_6H_5CH=CH$, R_2 is hydrogen and R_3 is phenyl, R_4 cannot be phenyl;

5 when R_1 is phenyl, R_2 is hydrogen, and R_3 is benzoyl, R_4 cannot be benzoyl;

when R_1 is phenyl, R_2 is substituted benzyl, R_3 is hydrogen and R_4 is hydrogen, R_n , R_o and R_p cannot all be hydrogen;

when R_1 is phenyl, R_3 is hydrogen and R_4 is hydrogen, R_2 cannot be benzyl or phenyl;

10 when R_1 is phenyl, R_2 is hydrogen, R_3 cannot be phenyl together with R_4 as benzoyl; and

when R_1 is phenyl, R_2 is hydrogen, R_3 and R_4 cannot both be benzyl,

for the manufacture of a medicament for down regulating a membrane ion channel functional activity in a cell infected with a virus.

15

83. The use according to claim 82, wherein said virus is a Lentivirus.

84. The use according to claim 83, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

20

85. The use according to claim 84, wherein said membrane ion channel is the HIV Vpu membrane ion channel.

86. The use according to claim 85, wherein said compound is selected from the group
25 consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,

3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoyleguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,
4-isopropylcinnamoylguanidine,

2,4,6-trimethylcinnamoylguanidine,
 N-(cinnamoyl)-N'phenylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-(2-naphthyl)acetoylguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamolyguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (a-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy-HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

87. The use according to any one of claims 84 to 86, wherein said HIV is HIV-1.

88. The use according to claim 82, wherein said virus is a Coronavirus.

89. The use according to claim 88, wherein said membrane ion channel is the Coronavirus E protein.
- 5 90. The use according to claim 89, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).
91. The use according to claim 90, wherein said compound is selected from the group consisting of
- 2,3-difluorocinnamoylguanidine,
 - 3,4-dichlorocinnamoylguanidine,
 - 4-t-butylcinnamoylguanidine,
 - 3-(2-naphthyl)acryloylguanidine,
 - (3-Chlorocinnamoyl)guanidine,
 - 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 - 2,5-dimethylcinnamoylguanidine,
 - trans-3-(1-naphthyl)acryloylguanidine,
 - 4-isopropylcinnamoylguanidine,
 - (3-Bromocinnamoyl)guanidine,
 - 6-methoxy-2-naphthoylguanidine,
 - 5-(N-Methyl-N-isobutyl)amiloride,
 - 3-phenylcinnamoylguanidine,
 - (2-Chlorocinnamoyl)guanidine,
 - 2'4 DichloroBenazamil HCl,
 - 4-phenylcinnamoylguanidine,
 - 4-(trifluoromethyl)cinnamoylguanidine,
 - 3-(trifluoromethoxy)cinnamoylguanidine,
 - 3-(trifluoromethyl)cinnamoylguanidine,
 - 2-ethoxycinnamoylguanidine,
 - cinnamoylguanidine hydrochloride,
 - 4-ethoxycinnamoylguanidine,
 - (2-Bromocinnamoyl)guanidine,
 - 2,6-dichlorocinnamoylguanidine,
 - 3,4,5-trimethoxycinnamoylguanidine,
 - 5-tert-butylamino-amiloride,
 - 3-t-butylcinnamoylguanidine,
 - 5-bromo-2-fluorocinnamoylguanidine,
 - (4-Chlorocinnamoyl)guanidine,
 - 2-t-butylcinnamoylguanidine,
 - 2-cyclohexylcinnamoylguanidine,
 - 6-Iodoamiloride,
 - 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,

(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,

1-bromo-2-naphthoylguanidine,
2-chloro-6-fluorocinnamoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,
Phenamil methanesulfonate salt,
N-Benzoyl-N'-cinnamoylguanidine and
N-(2-naphthoyl)-N'-phenylguanidine.

92. The use according to claim 90, wherein said Coronavirus is human Coronavirus 229E.

5 93. The use according to claim 92, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
3-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2,3-difluorocinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-phenylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
4-phenylbenzoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
1-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,

5-bromo-2-fluorocinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 Cinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 (a-Methylcinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 2,6-dichlorocinnamoylguanidine,
 (2-Bromocinnamoyl)guanidine,
 2,4,6-trimethylcinnamoylguanidine,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,
 2-(1-naphthyl)acetoyleguanidine,
 2-ethylcinnamoylguanidine,
 2-cyclohexylcinnamoylguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 2-ethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 2-methylcinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 cinnamoylguanidine hydrochloride,
 2,3-dimethylcinnamoylguanidine,
 2-fluorocinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 3,4-difluorocinnamoylguanidine,
 5-tert-butylamino-amiloride,
 2-naphthoylguanidine,
 N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 4-methylcinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 2,3,5,6,-tetramethylcinnamoylguanidine,
 3-ethoxycinnamoylguanidine,
 N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
 (4-Methoxycinnamoyl)guanidine,
 (2-Chlorocinnamoyl)guanidine,
 (3-Nitrocinnamoyl)guanidine,
 4-ethoxycinnamoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 2-(2-naphthyl)acetoyleguanidine and
 N-(3-phenylpropanoyl)-N'-phenylguanidine.

94. The use according to claim 89, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric coronavirus

(strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller),

5 porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine

10 coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus (strain JHM), murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus

15 strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N), porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat coronavirus (strain 681), rat coronavirus (strain NJ), rat sialodacryoadenitis coronavirus, turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey

20 coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious bronchitis virus (strain Beaudette US), avian infectious bronchitis virus (strain Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis

25 virus (strain D274), avian infectious bronchitis virus (strain D3896). avian infectious bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120), avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain KB8523), avian infectious bronchitis virus (strain M41), avian infectious bronchitis virus

30 (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain

UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious
 bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84),
 avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain
 Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003,
 5 SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS
 coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1,
 SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-
 39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong
 Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS
 10 coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS
 coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003,
 SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus
 Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani,
 SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01,
 15 bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus
 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus
 (strain CV777).

95. The use according to claim 94, wherein said compound is selected from the group
 20 consisting of

4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 4-t-butylcinnamoylguanidine, and
 3-isopropylcinnamoylguanidine hydrochloride.

96. The use according to claim 82, wherein said virus is the Hepatitis C virus.

97. The use according to claim 96, wherein said membrane ion channel is the
 25 Hepatitis C virus p7 membrane ion channel.


98. The use according to claim 97, wherein said compound is selected from the group
 consisting of

2,3-dimethylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(2-Chlorocinnamoyl)guanidine,
(4-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,

3-ethoxycinnamoylguanidine,
 2,3,5,6,-tetramethylcinnamoylguanidine,
 4-phenylcinnamoylguanidine,
 trans-3-Furanacryoylguanidine,
 N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 cinnamoylguanidine hydrochloride,
 5-(N,N-hexamethylene)amiloride,
 2,3-difluorocinnamoylguanidine,
 2-(1-naphthyl)acetoyleguanidine,
 (a-Methylcinnamoyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 6-Iodoamiloride,
 3,4-(methylenedioxy)cinnamoylguanidine,
 2-ethoxycinnamoylguanidine,
 Cinnamoylguanidine,
 2-phenylcinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-naphthoylguanidine,
 3-phenylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 5-(4-fluorophenyl)amiloride,
 (3-Methoxycinnamoyl)guanidine,
 2-fluorocinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 (3-phenylpropanoyl)guanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 2-methylcinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 1-bromo-2-naphthoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 Phenamil methanesulfonate salt,
 2,4-dichlorocinnamolyguanidine,
 3,4-difluorocinnamoylguanidine and
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.

99. The use according to any one of claims 82 to 98, wherein said medicament further comprises one or more known antiviral compounds or molecules.


100. Use of a compound according of Formula I



or pharmaceutically acceptable salts thereof,
wherein,


5

R₁ =



10


R₂, R₃ and R₄ are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, C₃₋₆cycloalkyl, halo-substituted C₁₋₆alkyl, halo-substituted C₁₋₆alkyloxy, phenyl, C₁₋₆alkeneyl, C₃₋₆cycloalkeneyl, C₁₋₆alkeneoxy, or benzo;

- R_a, R_b, R_c, R_d, R_e, R_f, R_h, R_k, R_L, R_m, R_n, R_o, R_p independently =
hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted aryl,
10 substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g, R_i independently = hydrogen, hydroxy, halo, or C₁₋₅ alkyl;

- R_j = hydrogen, amino, halo, C₁₋₅alkyl, C₁₋₅alkyloxy, hydroxy, aryl, substituted
15 aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein



when R₁ is R_c, R_a and R_c are amino, and R_b is halo, R₂, R₃ or R₄ cannot be hydrogen, benzyl or substituted benzyl or BODIPY-Fl;

when R₁ is C₆H₅CH=CH, R₂ is hydrogen and R₃ is phenyl, R₄ cannot be phenyl;

5 when R₁ is phenyl, R₂ is hydrogen, and R₃ is benzoyl, R₄ cannot be benzoyl;

when R₁ is phenyl, R₂ is substituted benzyl, R₃ is hydrogen and R₄ is hydrogen, R_n, R_o and R_p cannot all be hydrogen;

when R₁ is phenyl, R₃ is hydrogen and R₄ is hydrogen, R₂ cannot be benzyl or phenyl;

10 when R₁ is phenyl, R₂ is hydrogen, R₃ cannot be phenyl together with R₄ as benzoyl; and

when R₁ is phenyl, R₂ is hydrogen, R₃ and R₄ cannot both be benzyl,

for the manufacture of a medicament for reducing, retarding or otherwise inhibiting growth and/or replication of a virus that has infected a cell, wherein said compound down
15 regulates functional activity of a membrane ion channel derived from said virus and expressed in said infected cell.

101. The use according to claim 100, wherein said virus is a Lentivirus.

20 102. The use according to claim 101, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

103. The use according to claim 102, wherein said membrane ion channel is the HIV Vpu membrane ion channel.

25

104. The use according to claim 103, wherein said compound is selected from the group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,

(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoyleguanidine,

2,3-difluorocinnamoylguanidine,
 (3-Methoxycinnamoyl)guanidine,
 4-isopropylcinnamoylguanidine,
 2,4,6-trimethylcinnamoylguanidine,
 N-(cinnamoyl)-N'phenylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-(2-naphthyl)acetoyleguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamolyguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (a-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy -HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

105. The use according to any one of claims 102 to 104, wherein said HIV is HIV-1.

106. The use according to claim 100, wherein said virus is a Coronavirus.
107. The use according to claim 106, wherein said membrane ion channel is the
5 Coronavirus E protein.
108. The use according to claim 107, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).
- 10 109. The use according to claim 108, wherein said compound is selected from the group consisting of
- 2,3-difluorocinnamoylguanidine,
 - 3,4-dichlorocinnamoylguanidine,
 - 4-t-butylcinnamoylguanidine,
 - 3-(2-naphthyl)acryloylguanidine,
 - (3-Chlorocinnamoyl)guanidine,
 - 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 - 2,5-dimethylcinnamoylguanidine,
 - trans-3-(1-naphthyl)acryloylguanidine,
 - 4-isopropylcinnamoylguanidine,
 - (3-Bromocinnamoyl)guanidine,
 - 6-methoxy-2-naphthoylguanidine,
 - 5-(N-Methyl-N-isobutyl)amiloride,
 - 3-phenylcinnamoylguanidine,
 - (2-Chlorocinnamoyl)guanidine,
 - 2'4 DichloroBenazamil HCl,
 - 4-phenylcinnamoylguanidine,
 - 4-(trifluoromethyl)cinnamoylguanidine,
 - 3-(trifluoromethoxy)cinnamoylguanidine,
 - 3-(trifluoromethyl)cinnamoylguanidine,
 - 2-ethoxycinnamoylguanidine,
 - cinnamoylguanidine hydrochloride,
 - 4-ethoxycinnamoylguanidine,
 - (2-Bromocinnamoyl)guanidine,
 - 2,6-dichlorocinnamoylguanidine,
 - 3,4,5-trimethoxycinnamoylguanidine,
 - 5-tert-butylamino-amiloride,
 - 3-t-butylcinnamoylguanidine,
 - 5-bromo-2-fluorocinnamoylguanidine,
 - (4-Chlorocinnamoyl)guanidine,
 - 2-t-butylcinnamoylguanidine,

2-cyclohexylcinnamoylguanidine,
6-Iodoamiloride,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoyleguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoyleguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,

(Phenylacetyl)guanidine,
 N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
 6-bromo-2-naphthoylguanidine,
 1-bromo-2-naphthoylguanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 Phenamil methanesulfonate salt,
 N-Benzoyl-N'-cinnamoylguanidine and
 N-(2-naphthoyl)-N'-phenylguanidine.

110. The use according to claim 107, wherein said Coronavirus is human Coronavirus 229E.

5 111. The use according to claim 110, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
 3,4-dichlorocinnamoylguanidine,
 3-(trifluoromethoxy)cinnamoylguanidine,
 4-t-butylcinnamoylguanidine,
 3-isopropylcinnamoylguanidine hydrochloride,
 3-t-butylcinnamoylguanidine,
 2-t-butylcinnamoylguanidine,
 trans-3-(1-naphthyl)acryloylguanidine,
 5-bromo-2-methoxycinnamoylguanidine,
 2,3-difluorocinnamoylguanidine,
 3-(2-naphthyl)acryloylguanidine,
 2-phenylcinnamoylguanidine,
 3-phenylcinnamoylguanidine,
 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 4-phenylbenzoylguanidine,
 3-(trifluoromethyl)cinnamoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 4-(trifluoromethyl)cinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 (4-Bromocinnamoyl)guanidine,
 5-(N,N-hexamethylene)amiloride,
 1-naphthoylguanidine,
 5-(4-fluorophenyl)amiloride,
 (5-Phenyl-penta-2,4-dienoyl)guanidine,
 (3-Bromocinnamoyl)guanidine,
 2,5-dimethylcinnamoylguanidine,
 2-(trifluoromethyl)cinnamoylguanidine,

6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
5-bromo-2-fluorocinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
(α -Methylcinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,4,6-trimethylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(1-naphthyl)acetoyleguanidine,
2-ethylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
2-ethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
3-fluorocinnamoylguanidine,
cinnamoylguanidine hydrochloride,
2,3-dimethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-tert-butylamino-amiloride,
2-naphthoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
N,N'-Bis(3-phenylpropanoyl)guanidine,
4-methylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
2,3,5,6-tetramethylcinnamoylguanidine,
3-ethoxycinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(2-naphthyl)acetoyleguanidine and
N-(3-phenylpropanoyl)-N'-phenylguanidine.

112. The use according to claim 107, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric coronavirus (strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller), porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus (strain JHM), murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N), porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat coronavirus (strain 681), rat coronavirus (strain NJ), rat sialodacryoadenitis coronavirus, turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious bronchitis virus (strain Beaudette US), avian infectious bronchitis virus (strain Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis virus (strain D274), avian infectious bronchitis virus (strain D3896), avian infectious bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120), avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain KB8523), avian infectious bronchitis virus (strain M41), avian infectious

bronchitis virus (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84), avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003, SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1, SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003, SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani, SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus (strain CV777).

20

113. The use according to claim 112, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine, and
3-isopropylcinnamoylguanidine hydrochloride.

114. The use according to claim 100, wherein said virus is the Hepatitis C virus.

25

115. The use according to claim 114, wherein said membrane ion channel is the Hepatitis C virus p7 membrane ion channel.


116. The use according to claim 115, wherein said compound is selected from the group consisting of

2,3-dimethylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(2-Chlorocinnamoyl)guanidine,
(4-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'-phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,

(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
3-ethoxycinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
4-phenylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(2-Furanacryloyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,
5-(N,N-hexamethylene)amiloride,
2,3-difluorocinnamoylguanidine,
2-(1-naphthyl)acetoyleguanidine,
(a-Methylcinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
6-Iodoamiloride,
3,4-(methylenedioxy)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
Cinnamoylguanidine,
2-phenylcinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2-naphthoylguanidine,
3-phenylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
5-(4-fluorophenyl)amiloride,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,
(3-phenylpropanoyl)guanidine,
2-chloro-6-fluorocinnamoylguanidine,
3-fluorocinnamoylguanidine,
2-methylcinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
1-bromo-2-naphthoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
Phenamil methanesulfonate salt,
2,4-dichlorocinnamolyguanidine,
3,4-difluorocinnamoylguanidine and
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.

117. The use according to any one of claims 100 to 116, wherein said medicament further comprises one or more known antiviral compounds or molecules.

118. Use of a compound according of Formula I




5

or pharmaceutically acceptable salts thereof,


wherein,

$\text{R}_1 =$



10


R_2 , R_3 and R_4 are independently hydrogen,



5 and wherein


X = hydrogen, hydroxy, nitro, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{3-6} cycloalkyl, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkyloxy, phenyl, C_{1-6} alkeneyl, C_{3-6} cycloalkeneyl, C_{1-6} alkeneoxy, or benzo;

10 R_a , R_b , R_c , R_d , R_e , R_f , R_h , R_k , R_L , R_m , R_n , R_o , R_p independently = hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g , R_i independently = hydrogen, hydroxy, halo, or C_{1-5} alkyl;

15 R_j = hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino, PrS,



and wherein



when R_1 is R_c , R_a and R_c are amino, and R_b is halo, R_2 , R_3 or R_4 cannot be hydrogen, benzyl or substituted benzyl or BODIPY-Fl;

5 when R_1 is $C_6H_5CH=CH$, R_2 is hydrogen and R_3 is phenyl, R_4 cannot be phenyl;

when R_1 is phenyl, R_2 is hydrogen, and R_3 is benzoyl, R_4 cannot be benzoyl;

when R_1 is phenyl, R_2 is substituted benzyl, R_3 is hydrogen and R_4 is hydrogen, R_n , R_o and R_p cannot all be hydrogen;

10 when R_1 is phenyl, R_3 is hydrogen and R_4 is hydrogen, R_2 cannot be benzyl or phenyl;

when R_1 is phenyl, R_2 is hydrogen, R_3 cannot be phenyl together with R_4 as benzoyl; and

when R_1 is phenyl, R_2 is hydrogen, R_3 and R_4 cannot both be benzyl,

for the manufacture of a medicament for reducing, retarding or otherwise inhibiting
15 growth and/or replication of a virus that has infected a cell in a mammal, wherein said compound down regulates functional activity of a membrane ion channel expressed in said infected cell.

119. The use according to claim 118, wherein said virus is a Lentivirus.

20

120. The use according to claim 119, wherein said Lentivirus is Human Immunodeficiency Virus (HIV).

121. The use according to claim 120, wherein said compound is selected from the
25 group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,

3-(trifluoromethyl)cinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoylguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,

4-isopropylcinnamoylguanidine,
 2,4,6-trimethylcinnamoylguanidine,
 N-(cinnamoyl)-N'phenylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-(2-naphthyl)acetoyleguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamolyguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (a-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy -HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

122. The use according to any one of claims 118 to 121, wherein said membrane ion channel is the HIV Vpu membrane ion channel.

123. The use according to any one of claims 120 to 122, wherein said HIV is HIV-1.
124. The use according to claim 118, wherein said virus is a Coronavirus.
- 5 125. The use according to claim 124, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).
126. The use according to claim 125, wherein said compound is selected from the group consisting of
- 2,3-difluorocinnamoylguanidine,
 - 3,4-dichlorocinnamoylguanidine,
 - 4-t-butylcinnamoylguanidine,
 - 3-(2-naphthyl)acryloylguanidine,
 - (3-Chlorocinnamoyl)guanidine,
 - 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 - 2,5-dimethylcinnamoylguanidine,
 - trans-3-(1-naphthyl)acryloylguanidine,
 - 4-isopropylcinnamoylguanidine,
 - (3-Bromocinnamoyl)guanidine,
 - 6-methoxy-2-naphthoylguanidine,
 - 5-(N-Methyl-N-isobutyl)amiloride,
 - 3-phenylcinnamoylguanidine,
 - (2-Chlorocinnamoyl)guanidine,
 - 2'4 DichloroBenazamil HCl,
 - 4-phenylcinnamoylguanidine,
 - 4-(trifluoromethyl)cinnamoylguanidine,
 - 3-(trifluoromethoxy)cinnamoylguanidine,
 - 3-(trifluoromethyl)cinnamoylguanidine,
 - 2-ethoxycinnamoylguanidine,
 - cinnamoylguanidine hydrochloride,
 - 4-ethoxycinnamoylguanidine,
 - (2-Bromocinnamoyl)guanidine,
 - 2,6-dichlorocinnamoylguanidine,
 - 3,4,5-trimethoxycinnamoylguanidine,
 - 5-tert-butylamino-amiloride,
 - 3-t-butylcinnamoylguanidine,
 - 5-bromo-2-fluorocinnamoylguanidine,
 - (4-Chlorocinnamoyl)guanidine,
 - 2-t-butylcinnamoylguanidine,
 - 2-cyclohexylcinnamoylguanidine,
 - 6-Iodoamiloride,
 - 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,

(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,

1-bromo-2-naphthoylguanidine,
2-chloro-6-fluorocinnamoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,
Phenamil methanesulfonate salt,
N-Benzoyl-N'-cinnamoylguanidine and
N-(2-naphthoyl)-N'-phenylguanidine.

127. The use according to any one of claim 124 to 126, wherein said membrane ion channel is the Coronavirus E protein.

5 128. The use according to claim 124, wherein said Coronavirus is human Coronavirus 229E.

129. The use according to claim 128, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
3-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2,3-difluorocinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-phenylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
4-phenylbenzoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
1-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,

2-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,
5-bromo-2-fluorocinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
(α -Methylcinnamoyl)guanidine,
4-phenylcinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,4,6-trimethylcinnamoylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(1-naphthyl)acetoyleguanidine,
2-ethylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
2-ethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
3-fluorocinnamoylguanidine,
cinnamoylguanidine hydrochloride,
2,3-dimethylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-tert-butylamino-amiloride,
2-naphthoylguanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
N,N'-Bis(3-phenylpropanoyl)guanidine,
4-methylcinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
2,3,5,6-tetramethylcinnamoylguanidine,
3-ethoxycinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
2-(2-naphthyl)acetoyleguanidine and
N-(3-phenylpropanoyl)-N'-phenylguanidine.

130. The use according to claim 128 or claim 129, wherein said membrane ion channel is the Coronavirus E protein.

5 131. The use according to claim 124, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric coronavirus (strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis
10 coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus (strain Miller), porcine transmissible gastroenteritis coronavirus (strain Neb72-RT), porcine transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus
15 (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain OK-0514-3), bovine coronavirus (strain Ontario), bovine coronavirus (STRAIN QUEBEC), bovine coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain 98TXSF-110-ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine coronavirus (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus
20 (strain JHM), murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine hepatitis virus strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4, murine hepatitis virus strain ML-11, porcine hemagglutinating encephalomyelitis virus (strain 67N), porcine hemagglutinating encephalomyelitis virus (strain IAF-404), puffinosis virus, rat coronavirus (strain 681), rat coronavirus (strain NJ), rat
25 sialodacryoadenitis coronavirus, turkey coronavirus (strain Indiana), turkey coronavirus (strain Minnesota), turkey coronavirus (strain NC95), avian infectious bronchitis virus (STRAIN 6/82), avian infectious bronchitis virus (strain Arkansas 99), avian infectious bronchitis virus (strain Beaudette CK), avian infectious bronchitis virus (strain Beaudette M42), avian infectious bronchitis virus (strain Beaudette US), avian infectious bronchitis
30 virus (strain Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious bronchitis virus (strain D274), avian infectious bronchitis virus (strain D3896). avian

infectious bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072), avian infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain H120), avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus (strain KB8523), avian infectious bronchitis virus (strain M41), avian infectious

5 bronchitis virus (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain SAIB20), avian infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis virus (strain UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84), avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis

10 virus (strain Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003, SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1, SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong

15 Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003, SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani,

20 SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus (strain CV777).

25 132. The use according to claim 131, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine, and
3-isopropylcinnamoylguanidine hydrochloride.

133. The use according to claim 131 or claim 132, wherein said membrane ion channel is the Coronavirus E protein.

134. The use according to claim 118, wherein said virus is the Hepatitis C virus.

5

135. The use according to claim 134, wherein said compound is selected from the group consisting of

2,3-dimethylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(2-Chlorocinnamoyl)guanidine,
(4-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'-phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,

5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
3-ethoxycinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
4-phenylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(2-Furanacryloyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
cinnamoylguanidine hydrochloride,
5-(N,N-hexamethylene)amiloride,
2,3-difluorocinnamoylguanidine,
2-(1-naphthyl)acetoyleguanidine,
(a-Methylcinnamoyl)guanidine,
(2-Nitrocinnamoyl)guanidine,
6-Iodoamiloride,
3,4-(methylenedioxy)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
Cinnamoylguanidine,
2-phenylcinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2-naphthoylguanidine,
3-phenylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
5-(4-fluorophenyl)amiloride,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
[(4-Chlorophenoxy-acetyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-chloro-6-fluorocinnamoylguanidine,
3-fluorocinnamoylguanidine,
2-methylcinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
1-bromo-2-naphthoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
3-methylcinnamoylguanidine,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,

Phenamil methanesulfonate salt,
2,4-dichlorocinnamolyguanidine,
3,4-difluorocinnamoylguanidine and
[(E)-3-(4-Dimethylaminophenyl)-2-
methylacryloyl]guanidine.

136. The use according to claim 135, wherein said membrane ion channel is the Hepatitis C virus p7 membrane ion channel.

5 137. The use according to any one of claims 118 to 130, wherein said mammal is a primate.

138. The use according to any one of claims 134 to 136, wherein said mammal is a primate.


10

139. The use according to claim 137 or claim 138, wherein said primate is human.

140. The use according to any one of claims 118 to 139, wherein said medicament further comprises one or more known antiviral compounds or molecules.

15


141. Use of a compound according of Formula I




or pharmaceutically acceptable salts thereof,

wherein,

20 $R_1 =$




R_2 , R_3 and R_4 are independently hydrogen,



and wherein


- 5 X = hydrogen, hydroxy, nitro, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, C_{3-6} cycloalkyl, halo-substituted C_{1-6} alkyl, halo-substituted C_{1-6} alkyloxy, phenyl, C_{1-6} alkenyl, C_{3-6} cycloalkenyl, C_{1-6} alkeneoxy, or benzo;

- R_a , R_b , R_c , R_d , R_e , R_f , R_h , R_k , R_L , R_m , R_n , R_o , R_p independently =
hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl,
10 substituted amino, mono or dialkyl-substituted amino, cycloalkyl-substituted amino, aryl-substituted amino,




R_g , R_i independently = hydrogen, hydroxy, halo, or C_{1-5} alkyl;

- R_j = hydrogen, amino, halo, C_{1-5} alkyl, C_{1-5} alkyloxy, hydroxy, aryl, substituted aryl, substituted amino, alkyl-substituted amino, cycloalkyl-substituted amino,
15 aryl-substituted amino, PrS,



and wherein



when R_1 is R_c , R_a and R_c are amino, and R_b is halo, R_2 , R_3 or R_4 cannot be hydrogen, benzyl or substituted benzyl or BODIPY-Fl;

when R_1 is $C_6H_5CH=CH$, R_2 is hydrogen and R_3 is phenyl, R_4 cannot be phenyl;

5 when R_1 is phenyl, R_2 is hydrogen, and R_3 is benzoyl, R_4 cannot be benzoyl;

when R_1 is phenyl, R_2 is substituted benzyl, R_3 is hydrogen and R_4 is hydrogen, R_n , R_o and R_p cannot all be hydrogen;

when R_1 is phenyl, R_3 is hydrogen and R_4 is hydrogen, R_2 cannot be benzyl or phenyl;

10 when R_1 is phenyl, R_2 is hydrogen, R_3 cannot be phenyl together with R_4 as benzoyl; and

when R_1 is phenyl, R_2 is hydrogen, R_3 and R_4 cannot both be benzyl,

for the manufacture of a medicament for the therapeutic or prophylactic treatment of a subject infected with or exposed to a virus, wherein said compound down-regulates

15 functional activity of a membrane ion channel derived from said virus.

142. The use according to claim 141, wherein said virus is a Lentivirus.

143. The use according to claim 142, wherein said Lentivirus is Human
20 Immunodeficiency Virus (HIV).

144. The use according to claim 143, wherein said membrane ion channel is the HIV Vpu membrane ion channel.

25 145. The use according to claim 144, wherein said compound is selected from the group consisting of

(3-Chlorocinnamoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
(2-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
3-(trifluoromethyl)cinnamoylguanidine,

5-bromo-2-fluorocinnamoylguanidine,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3-dimethylcinnamoylguanidine,
Cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
3,4-dichlorocinnamoylguanidine,
2,6-dichlorocinnamoylguanidine,
4-phenylbenzoylguanidine,
2-ethylcinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-naphthoylguanidine,
2,5-dimethylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
3-phenylcinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
2-ethoxycinnamoylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
(4-Methoxycinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
4-methylcinnamoylguanidine,
2-fluorocinnamoylguanidine,
2-phenylcinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
3-t-butylcinnamoylguanidine,
3,4-difluorocinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
3-fluorocinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-ethoxycinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
2,3,5,6,-tetramethylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-(1-naphthyl)acetoylguanidine,
2,3-difluorocinnamoylguanidine,
(3-Methoxycinnamoyl)guanidine,

4-isopropylcinnamoylguanidine,
 2,4,6-trimethylcinnamoylguanidine,
 N-(cinnamoyl)-N'-phenylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-(2-naphthyl)acetylguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 Phenamil methanesulfonate salt,
 Benzamil hydrochloride,
 (3-Nitrocinnamoyl)guanidine,
 Benzyoylguanidine,
 (4-Phenoxybenzoyl)guanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 5-(N-Methyl-N-isobutyl)amiloride,
 2-cyclohexylcinnamoylguanidine,
 4-ethoxycinnamoylguanidine,
 2,4-dichlorocinnamoylguanidine,
 5-(N-Ethyl-N-isopropyl)amiloride,
 N-amidino-3-amino-5-hexamethyleneimino-6-phenyl-
 2-pyrazinecarboxamide,
 (a-Methylcinnamoyl)guanidine,
 cinnamoylguanidine hydrochloride,
 [(4-Chlorophenoxy-acetyl]guanidine,
 N-amidino-3-amino-5-phenyl-6-chloro-2-
 pyrazinecarboxamide,
 5-(4-fluorophenyl)amiloride,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 trans-3-Furanacryoylguanidine,
 1-naphthoylguanidine,
 5-tert-butylamino-amiloride,
 3-methoxy -HMA,
 (3-phenylpropanoyl)guanidine,
 4-t-butylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 N-Benzoyl-N'-cinnamoylguanidine and
 1-bromo-2-naphthoylguanidine.

146. The use according to any one of claims 143 to 145, wherein said HIV is HIV-1.

147. The use according to claim 141, wherein said virus is a Coronavirus.

148. The use according to claim 147, wherein said membrane ion channel is the Coronavirus E protein.

5 149. The use according to claim 148, wherein said Coronavirus is the Severe Acute Respiratory Syndrome virus (SARS).

150. The use according to claim 149, wherein said compound is selected from the group consisting of

2,3-difluorocinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
(3-Chlorocinnamoyl)guanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
2,5-dimethylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-isopropylcinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
6-methoxy-2-naphthoylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
3-phenylcinnamoylguanidine,
(2-Chlorocinnamoyl)guanidine,
2'4 DichloroBenazamil HCl,
4-phenylcinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
2-ethoxycinnamoylguanidine,
cinnamoylguanidine hydrochloride,
4-ethoxycinnamoylguanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
3,4,5-trimethoxycinnamoylguanidine,
5-tert-butylamino-amiloride,
3-t-butylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Chlorocinnamoyl)guanidine,
2-t-butylcinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
6-Iodoamiloride,
3-(trans-hept-1-en-1-yl)cinnamoylguanidine,

(4-Bromocinnamoyl)guanidine,
(4-Hydroxycinnamoyl)guanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
(3-Nitrocinnamoyl)guanidine,
3-fluorocinnamoylguanidine,
2-(1-naphthyl)acetoylguanidine,
2-ethylcinnamoylguanidine,
5-(N,N-Dimethyl)amiloride hydrochloride,
2-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
2-(trifluoromethyl)cinnamoylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
1-naphthoylguanidine,
Benzamil hydrochloride,
3-methoxy -HMA,
4-methylcinnamoylguanidine,
4-fluorocinnamoylguanidine,
3,4-(methylenedioxy)cinnamoylguanidine,
5-(N,N-hexamethylene)amiloride,
N-(cinnamoyl)-N'phenylguanidine,
5-(N-Ethyl-N-isopropyl)amiloride,
3-methylcinnamoylguanidine,
2-methylcinnamoylguanidine,
2,3,5,6,-tetramethylcinnamoylguanidine,
trans-3-Furanacryoylguanidine,
(4-Methoxycinnamoyl)guanidine,
(2-Furanacryloyl)guanidine,
(3-phenylpropanoyl)guanidine,
2-(2-naphthyl)acetoylguanidine,
Cinnamoylguanidine,
(2-Methoxycinnamoyl)guanidine,
[3-(3-Pyridyl)acryloyl]guanidine,
4-phenylbenzoylguanidine,
2,4-dichlorocinnamolyguanidine,
(3-Methoxycinnamoyl)guanidine,
2-fluorocinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
(a-Methylcinnamoyl)guanidine,
5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(Quinoline-2-carbonyl)guanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
6-bromo-2-naphthoylguanidine,

1-bromo-2-naphthoylguanidine,
2-chloro-6-fluorocinnamoylguanidine,
[(4-Chlorophenoxy-acetyl]guanidine,
Phenamil methanesulfonate salt,
N-Benzoyl-N'-cinnamoylguanidine and
N-(2-naphthoyl)-N'-phenylguanidine.

151. The use according to claim 148, wherein said Coronavirus is human Coronavirus 229E.

5 152. The use according to claim 151, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
3-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
2,3-difluorocinnamoylguanidine,
3-(2-naphthyl)acryloylguanidine,
2-phenylcinnamoylguanidine,
3-phenylcinnamoylguanidine,
3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
4-phenylbenzoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
4-(trifluoromethyl)cinnamoylguanidine,
2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
5-(N,N-hexamethylene)amiloride,
1-naphthoylguanidine,
5-(4-fluorophenyl)amiloride,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
(3-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
2-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(4-Chlorocinnamoyl)guanidine,
(3-Methoxycinnamoyl)guanidine,

5-bromo-2-fluorocinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 Cinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 (a-Methylcinnamoyl)guanidine,
 4-phenylcinnamoylguanidine,
 2,6-dichlorocinnamoylguanidine,
 (2-Bromocinnamoyl)guanidine,
 2,4,6-trimethylcinnamoylguanidine,
 (trans-2-Phenylcyclopropanecarbonyl)guanidine,
 (3-Chlorocinnamoyl)guanidine,
 2-(1-naphthyl)acetoyleguanidine,
 2-ethylcinnamoylguanidine,
 2-cyclohexylcinnamoylguanidine,
 (4-Hydroxycinnamoyl)guanidine,
 2-ethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 2-methylcinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 cinnamoylguanidine hydrochloride,
 2,3-dimethylcinnamoylguanidine,
 2-fluorocinnamoylguanidine,
 4-fluorocinnamoylguanidine,
 3,4-difluorocinnamoylguanidine,
 5-tert-butylamino-amiloride,
 2-naphthoylguanidine,
 N,N'-Bis(amidino)naphthalene-2,6-dicarboxamide,
 N,N'-Bis(3-phenylpropanoyl)guanidine,
 4-methylcinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 2,3,5,6,-tetramethylcinnamoylguanidine,
 3-ethoxycinnamoylguanidine,
 N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
 (4-Methoxycinnamoyl)guanidine,
 (2-Chlorocinnamoyl)guanidine,
 (3-Nitrocinnamoyl)guanidine,
 4-ethoxycinnamoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 2-(2-naphthyl)acetoyleguanidine and
 N-(3-phenylpropanoyl)-N'-phenylguanidine.

153. The use according to claim 148, wherein said Coronavirus is any one of the known Coronavirus isolates selected from the group consisting of canine enteric

coronavirus (strain INSAVC-1), canine enteric coronavirus (strain K378), feline enteric
 coronavirus (strain 79-1683), feline infectious peritonitis virus (FIPV), Equine
 coronavirus NC99, porcine respiratory coronavirus, porcine transmissible gastroenteritis
 coronavirus (STRAIN FS772/70), porcine transmissible gastroenteritis coronavirus
 5 (strain Miller), porcine transmissible gastroenteritis coronavirus (strain Neb72-RT),
 porcine transmissible gastroenteritis coronavirus (STRAIN PURDUE), bovine
 coronavirus (STRAIN F15), bovine coronavirus (strain G95), bovine coronavirus
 (STRAIN L9), bovine coronavirus (strain LSU-94LSS-051), bovine coronavirus
 (STRAIN LY-138), bovine coronavirus (STRAIN MEBUS), bovine coronavirus (strain
 10 OK-0514-3), bovine coronavirus (strain Ontario), bovine coronavirus (STRAIN
 QUEBEC), bovine coronavirus (STRAIN VACCINE), bovine enteric coronavirus (strain
 98TXSF-110-ENT), canine respiratory coronavirus, chicken enteric coronavirus, murine
 coronavirus (strain DVIM), murine hepatitis virus (strain A59), murine hepatitis virus
 (strain JHM), murine hepatitis virus (strain S), murine hepatitis virus strain 1, murine
 15 hepatitis virus strain 2, murine hepatitis virus strain 3, murine hepatitis virus strain 4,
 murine hepatitis virus strain ML-11, porcine hemagglutinating encephalomyelitis virus
 (strain 67N), porcine hemagglutinating encephalomyelitis virus (strain IAF-404),
 puffinosis virus, rat coronavirus (strain 681), rat coronavirus (strain NJ), rat
 sialodacryoadenitis coronavirus, turkey coronavirus (strain Indiana), turkey coronavirus
 20 (strain Minnesota), turkey coronavirus (strain NC95), avian infectious bronchitis virus
 (STRAIN 6/82), avian infectious bronchitis virus (strain Arkansas 99), avian infectious
 bronchitis virus (strain Beaudette CK), avian infectious bronchitis virus (strain Beaudette
 M42), avian infectious bronchitis virus (strain Beaudette US), avian infectious bronchitis
 virus (strain Beaudette), avian infectious bronchitis virus (strain D1466), avian infectious
 25 bronchitis virus (strain D274), avian infectious bronchitis virus (strain D3896). avian
 infectious bronchitis virus (strain D41), avian infectious bronchitis virus (strain DE072),
 avian infectious bronchitis virus (strain GRAY), avian infectious bronchitis virus (strain
 H120), avian infectious bronchitis virus (strain H52), avian infectious bronchitis virus
 (strain KB8523), avian infectious bronchitis virus (strain M41), avian infectious
 30 bronchitis virus (strain PORTUGAL/322/82), avian infectious bronchitis virus (strain
 SAIB20), avian infectious bronchitis virus (strain UK/123/82), avian infectious bronchitis

virus (strain UK/142/86), avian infectious bronchitis virus (strain UK/167/84), avian infectious bronchitis virus (strain UK/183/66), avian infectious bronchitis virus (strain UK/68/84), avian infectious bronchitis virus (strain V18/91), avian infectious bronchitis virus (strain Vic S), avian infectious laryngotracheitis virus, SARS coronavirus Beijing ZY-2003, SARS coronavirus BJ01, SARS coronavirus BJ02, SARS coronavirus BJ03, SARS coronavirus BJ04, SARS coronavirus CUHK-Su10, SARS coronavirus CUHK-W1, SARS coronavirus Frankfurt 1, SARS coronavirus GZ01, SARS coronavirus HKU-39849, SARS coronavirus Hong Kong ZY-2003, SARS coronavirus Hong Kong/03/2003, SARS coronavirus HSR 1, SARS coronavirus Sin2500, SARS coronavirus Sin2677, SARS coronavirus Sin2679, SARS coronavirus Sin2748, SARS coronavirus Sin2774, SARS coronavirus Taiwan, SARS coronavirus Taiwan JC-2003, SARS coronavirus Taiwan TC1, SARS coronavirus Taiwan TC2, SARS coronavirus Tor2, SARS coronavirus TW1, SARS coronavirus TWC, SARS coronavirus Urbani, SARS coronavirus Vietnam, SARS coronavirus ZJ-HZ01, SARS coronavirus ZJ01, bovine respiratory coronavirus (strain 98TXSF-110-LUN), human enteric coronavirus 4408, porcine epidemic diarrhea virus (strain Br1/87) and porcine epidemic diarrhea virus (strain CV777).

154. The use according to claim 153, wherein said compound is selected from the group consisting of

4-isopropylcinnamoylguanidine,
3,4-dichlorocinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
4-t-butylcinnamoylguanidine, and
3-isopropylcinnamoylguanidine hydrochloride.

155. The use according to claim 141, wherein said virus is the Hepatitis C virus.

156. The use according to claim 155, wherein said membrane ion channel is the Hepatitis C virus p7 membrane ion channel.

157. The use according to claim 156, wherein said compound is selected from the group consisting of

2,3-dimethylcinnamoylguanidine,
2,4,6-trimethylcinnamoylguanidine,
5-bromo-2-fluorocinnamoylguanidine,
(4-Bromocinnamoyl)guanidine,
2,5-dimethylcinnamoylguanidine,
3-(trifluoromethyl)cinnamoylguanidine,
4-(trifluoromethyl)cinnamoylguanidine,
6-methoxy-2-naphthoylguanidine,
(2-Chlorocinnamoyl)guanidine,
(4-Chlorocinnamoyl)guanidine,
(2-Bromocinnamoyl)guanidine,
2,6-dichlorocinnamoylguanidine,
(3-Bromocinnamoyl)guanidine,
(3-Chlorocinnamoyl)guanidine,
2-(trifluoromethyl)cinnamoylguanidine,
(4-Phenoxybenzoyl)guanidine,
3,4-dichlorocinnamoylguanidine,
4-isopropylcinnamoylguanidine,
trans-3-(1-naphthyl)acryloylguanidine,
4-t-butylcinnamoylguanidine,
2-t-butylcinnamoylguanidine,
2-ethylcinnamoylguanidine,
4-methylcinnamoylguanidine,
5-bromo-2-methoxycinnamoylguanidine,
3-(trifluoromethoxy)cinnamoylguanidine,
2-cyclohexylcinnamoylguanidine,
1-naphthoylguanidine,
3-t-butylcinnamoylguanidine,
4-phenylbenzoylguanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
N-(cinnamoyl)-N'-phenylguanidine,
3-isopropylcinnamoylguanidine hydrochloride,
Benzamil hydrochloride,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
3-(2-naphthyl)acryloylguanidine,
5-(N-Methyl-N-isobutyl)amiloride,
2'4 DichloroBenazamil HCl,
5-tert-butylamino-amiloride,
5-(N-Ethyl-N-isopropyl)amiloride,
(4-Methoxycinnamoyl)guanidine,
4-fluorocinnamoylguanidine,
(3-Nitrocinnamoyl)guanidine,
4-ethoxycinnamoylguanidine,
(4-Hydroxycinnamoyl)guanidine,
(trans-2-Phenylcyclopropanecarbonyl)guanidine,

3-ethoxycinnamoylguanidine,
 2,3,5,6,-tetramethylcinnamoylguanidine,
 4-phenylcinnamoylguanidine,
 trans-3-Furanacryoylguanidine,
 N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
 (2-Furanacryloyl)guanidine,
 3-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 cinnamoylguanidine hydrochloride,
 5-(N,N-hexamethylene)amiloride,
 2,3-difluorocinnamoylguanidine,
 2-(1-naphthyl)acetoyleguanidine,
 (a-Methylcinnamoyl)guanidine,
 (2-Nitrocinnamoyl)guanidine,
 6-Iodoamiloride,
 3,4-(methylenedioxy)cinnamoylguanidine,
 2-ethoxycinnamoylguanidine,
 Cinnamoylguanidine,
 2-phenylcinnamoylguanidine,
 2-(cyclohex-1-en-1-yl)cinnamoylguanidine,
 2-naphthoylguanidine,
 3-phenylcinnamoylguanidine,
 5-(N,N-Dimethyl)amiloride hydrochloride,
 5-(4-fluorophenyl)amiloride,
 (3-Methoxycinnamoyl)guanidine,
 2-fluorocinnamoylguanidine,
 5-(3'-bromophenyl)penta-2,4-dienoylguanidine,
 [(4-Chlorophenoxy-acetyl)guanidine,
 (3-phenylpropanoyl)guanidine,
 2-chloro-6-fluorocinnamoylguanidine,
 3-fluorocinnamoylguanidine,
 2-methylcinnamoylguanidine,
 (2-Methoxycinnamoyl)guanidine,
 1-bromo-2-naphthoylguanidine,
 3,4,5-trimethoxycinnamoylguanidine,
 3-methylcinnamoylguanidine,
 3-(trans-hept-1-en-1-yl)cinnamoylguanidine,
 Phenamil methanesulfonate salt,
 2,4-dichlorocinnamolyguanidine,
 3,4-difluorocinnamoylguanidine and
 [(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine.

158. The use according to any one of claims 141 to 152, wherein said mammal is a primate.

159. The use according to any one of claims 155 to 157, wherein said mammal is a primate.

160. The use according to claim 158 or claim 159, wherein said primate is human.

5

161. A compound selected from the group consisting of

- 1-naphthoylguanidine,
N-(2-naphthoyl)-N'-phenylguanidine,
N,N'-bis(2-naphthoyl)guanidine,
10 N,N'-bis(1-naphthoyl)guanidine,
N,N'-bis(2-naphthoyl)-N''-phenylguanidine,
3-quinolinoylguanidine,
cinnamoylguanidine,
4-phenylbenzoylguanidine,
15 N-(cinnamoyl)-N'phenylguanidine,
N,N'-bis-(cinnamoyl)-N''-phenylguanidine,
N-(3-phenylpropanoyl)-N'-phenylguanidine,
N,N'-bis(3phenylpropanoyl)-N''-phenylguanidine,
N-(6-Hydroxy-2-naphthoyl)-N'-phenylguanidine,
20 (4-Phenoxybenzoyl)guanidine,
N,N'-Bis(amidino)napthalene-2,6-dicarboxamide,
N''-Cinnamoyl-N,N'-diphenylguanidine,
(Phenylacetyl)guanidine,
N,N'-Bis(3-phenylpropanoyl)guanidine,
25 benzyoylguanidine,
(4-Chlorophenoxy-acetyl)guanidine,
N-benzoyl-N'-cinnamoylguanidine,
[(E)-3-(4-Dimethylaminophenyl)-2-methylacryloyl]guanidine,
(5-Phenyl-penta-2,4-dienoyl)guanidine,
30 (4-Hydroxycinnamoyl)guanidine, and
(Quinoline-2-carbonyl)guanidine,

or pharmaceutically acceptable salts thereof.

162. A pharmaceutical composition comprising a compound according to claim 161 and one or more pharmaceutical acceptable carriers or derivatives.

5

163. The pharmaceutical composition according to claim 162, further comprising one or more known antiviral compounds.

10

10 * 20 * 30 * 40 * 50 * 60 * 70 * 80 *
 M Q P T P I V A I V A L V A L I I A I V V N S I V I I E Y R K I L R Q R K I D R L I D R L I E R A E D S C N E S E G E I S A L V E M G V E M G H H A P W D V D D L
 *

Figure 1a




Figure 1b




Figure 1c




Figure 2a




Figure 2b




Figure 3a




Figure 3b

Figure 4a


37°C, Pro⁻ plate

Figure 4b



30°C, Pro⁻ plate

Figure 4c

37°C, Met⁻ plate




Figure 5




Figure 6a




Figure 6b

9/14




Figure 7a




Figure 7b



Figure 8a




Figure 8b




Figure 8c

Figure 9:





Figure 10





Figure 11






Figure 12

A**B**