

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 April 2009 (09.04.2009)

PCT

(10) International Publication Number
WO 2009/046446 A2

(51) International Patent Classification:

A61K 9/14 (2006.01) C08G 83/00 (2006.01)
A61K 47/48 (2006.01) C08G 73/00 (2006.01)

(21) International Application Number:

PCT/US2008/078988

(22) International Filing Date: 6 October 2008 (06.10.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/997,987 5 October 2007 (05.10.2007) US
61/135,809 23 July 2008 (23.07.2008) US

(71) Applicant (for all designated States except US): WAYNE STATE UNIVERSITY [US/US]; 5057 Woodward, Suite 6306, Detroit, MI 48202 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): KANNAN, Rangaramanujam M. [US/US]; 44410 Midway Drive, Novi, MI 48375 (US). IEZZI, Raymond [US/US]; 3714 Balfour Avenue, Troy, MI 48084 (US). RAJAGURU, Bharath [IN/US]; 5200 Anthony Wayne Drive, 1216, Detroit, MI 48202 (US). KANNAN, Sujatha [US/US]; 44410 Midway Drive, Novi, MI 48375 (US).

(74) Agents: POTTER, Jane E., R. et al.; 1201 Third Avenue, Suite 2200, Seattle, WA 98101-3045 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

(54) Title: DENDRIMERS FOR SUSTAINED RELEASE OF COMPOUNDS



Figures 2A-2F

(57) Abstract: Dendrimer-based compositions and methods are provided, that are useful for administering pharmaceutical compositions to target cells and tissues for treatment of ocular diseases including macular degeneration, diabetic retinopathy, and retinitis pigmentosa.

WO 2009/046446 A2

DENDRIMERS FOR SUSTAINED RELEASE OF COMPOUNDS

FIELD OF THE INVENTION

The present invention relates to dendrimer-based compositions and methods
5 useful for administering pharmaceutical compositions to target cells and tissues for
suppressing neuroinflammation in disease states, including macular degeneration.

BACKGROUND

A common pathway in many human disease states is microglia-mediated
10 inflammation. Microglia are tissue-resident macrophages found in the retina and the
central nervous system. Microglial cells constitute about ten to twenty percent of the
cells in the adult brain. Under normal conditions, these cells are constitutively
suppressed by endogenous cortisol. The cells become activated in the form of
phagocytes and cytotoxic cells in the presence of a variety of stimuli. These stimuli
15 include trauma, infection, inflammation, ischemia, lipopolysaccharides, reactive oxygen
species, and damaged cell membranes. Once microglia are activated, they can
migrate and recruit other microglia to the original site of damage. Malfunctioning cells
can be killed by the release of tumor necrosis factor alpha (TNF- α), reactive oxygen
species (ROS), and proteases. The resulting cell debris is phagocytized by the
20 microglia cells.

Secondary cell damage occurs in a process referred to as bystander lysis:
nearby healthy cells are destroyed in the toxic extracellular milieu created by the
activated microglia. This amplifies the cell damage beyond the cells affected by the
underlying pathologic event, and turns the remedy – the activated microglial cells – into
25 a pathologic system in its own right.

The cascading pattern of primary pathology, response by microglial cells, and
subsequent secondary pathology has been observed in a broad range of human
diseases, including diseases of the eye. An essential element of sight is a functioning
retina. The retina can be likened to the “film” of the eye. It converts light rays into
30 electrical signals and sends them to the brain through the optic nerve. The sides of the
retina are responsible for peripheral vision. The center area, called the macula, is used
for fine central vision and color vision. The retina is where many problems leading to
vision loss occur. Three of the leading causes of blindness due to retina damage

associated with neuroinflammation are retinitis pigmentosa, macular degeneration and diabetic retinopathy. The leading cause of blindness in African Americans is glaucoma a degenerative process of the optic nerve and retina that involves neuroinflammation and microglial cell activation within the optic nerve that connects the retina to the brain.

- 5 Other important retinal diseases that are associated with neuroinflammation include uveitis, auto-immune photoreceptor degenerations and infection.

From a clinical perspective, retinitis pigmentosa, late-onset retinal degeneration, and age-related macular degeneration have significant impact on human health and quality of life. Nine million Americans suffer progressive vision loss due to retinal
10 neurodegenerative diseases. Retinitis pigmentosa affects one in four thousand individuals. It is the fourth leading cause of visual disability in the United States, after diabetic retinopathy, age-related macular degeneration, and glaucoma.

Age-related macular degeneration (AMD) is a neurodegenerative, neuroinflammatory disease of the macula, which is responsible for central vision loss.
15 AMD is the leading cause of vision loss in people over age 65. Eight million people are legally blind from macular degeneration worldwide, and as the population ages this number is expected to grow

The pathogenesis of age-related macular degeneration involves chronic neuroinflammation in the choroid (a blood vessel layer under the retina), the retinal
20 pigment epithelium (RPE), a cell layer under the neurosensory retina, Bruch's membrane and the neurosensory retina, itself. The disease first manifests as a dry form that involves the accumulation of drusen - cell debris and inflammatory material that form small masses within the RPE and Bruch's membrane. Drusen contain broken down cell membranes and other cell fragments and are highly antigenic, activating a
25 localized microglial and macrophage-mediated inflammatory response within the retina. Over time, the toxic mediators associated with this inflammation break down Bruch's membrane and the RPE and can lead directly to vision loss or may lead to the leakage of vascular endothelial-derived growth factor from the choroidal circulation into the subretinal space. This, in turn, can lead to the formation of abnormal blood vessels,
30 called choroidal neovascularization. Since these blood vessels are abnormal, they often leak serum, causing retina exudates and can sometimes bleed. Since fluid is involved, this is called, "wet" age-related macular degeneration. The "wet" and "dry" forms of age-related macular degeneration can co-exist, both involving neuroinflammation.

The microglial-mediated pathology is also common to a variety of central nervous system neurodegenerative diseases including Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, Huntington's disease, and acute spinal cord trauma. Brain injury is another cause of lifelong disability. For example,
5 brain injury in the perinatal period can lead to cerebral palsy, which also involves microglial cells in the peri-ventricular leukomalacia following the injury.

There is a strong and immediate need in the art for clinically effective treatments for all these diseases, and inhibiting the common pathway of microglial-mediated tissue destruction as provided by the present disclosure meets this need.

10

SUMMARY OF THE INVENTION

A composition is provided comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound. The biologically active component is selected from the group consisting of natural steroids
15 such as Cholesterol, Progestins Pregnenolone, 17-hydroxypregnenolone, Progesterone, 17-hydroxyprogesterone, Androgens, Androstenedione, 4-hydroxy-Androstenedione, 11 β -hydroxyandrostenedione, Androstenediol, Androsterone, Epiandrosterone, Adrenosterone, Dehydroepiandrosterone, Dehydroepiandrosterone Sulphate, Testosterone, Epitestosterone, 5 α -dihydrotestosterone, 5 β -
20 dihydrotestosterone, 11 β -hydroxytestosterone, 11-ketotestosterone, Estrogens, Estrone, Estradiol, Estriol, Corticosteroids, Corticosterone, Deoxycorticosterone, Cortisol, 11-Deoxycortisol, Cortisone, 18-hydroxycorticosterone, 1 α -hydroxycorticosterone, Aldosterone, synthetic steroids, anti-inflammatory agents, vitamins, peptides, growth factors, CNS stimulants, oligonucleotides, siRNAs, microRNAs, resolvins,
25 neurostimulants and protectants. The biologically active compound may be fluocinolone acetonide, ranibizumab, minocycline, rapamycin, methyl prednisone, dexamethasone, insulin, estradiol, CNTF, vitamin A, vitamin C, vitamin E, an antioxidant and an oligonucleotide, or a pharmaceutically acceptable salt thereof.

In the composition, the size of the nanoparticle is equal to or less than about
30 1000 nm, less than about 500 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 90 nm, less than about 80 nm, less than about 70 nm, less than about 60 nm, less than about 50 nm, less than about 40 nm, less than about 30 nm, less than about 20 nm, less than about 19 nm, less than about 18 nm, less than about 17 nm, less than about 16 nm, less than about 15 nm, less than about

14 nm, less than about 13 nm, less than about 12 nm, less than about 11 nm, less than about 10 nm, less than about 5 nm, less than about 4 nm, less than about 3 nm, less than about 2 nm, less than about 1 nm, or any value there-between or less.

The nanoparticle of the composition may be a soft nanoparticle, such as a dendrimer-branched or star-branched polymer, or dendrimer-polymer hybrid. The dendrimer-branched polymer may consist of polyamidoamine (PAMAM), priostar, polyester, polyether, polylysine, or polyethylene glycol (PEG), polypeptidedendrimers. The star-branched polymer may be a PEG star. The soft nanoparticle may have a diameter of 1.5 nanometers to 14.5 nanometers. Also provided is a composition comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound and the drug is incorporated into a hyperbranched formulation through encapsulation, complexation, or covalent linkage. The linkage may comprise a spacer consisting of a peptide, glutaric acid, or PEG to link the drug and the polymer.

In certain embodiments, the drug-nanoparticle is incorporated in a larger scale entity incorporating the drug-hyperbranched polymer, wherein the larger-scale entity consists of a polymer matrix, a microparticle, a nanoparticle, a liposome, a microcapsule, a nanocapsule, or a controlled-release implant.

In some embodiments, the dendrimer-drug nanoconjugate can be delivered alone or incorporated into a topical preparation an implanted device coating, and implanted drug-delivery system, an injectable or implantable hydrogel or may be incorporated into a contact lens. This may be injected into the systemic circulation, may be delivered to the surface of they eye in the form of a contact lens, applied as an eyedrop or delivered into the corneal stroma. It may applied to the subconjunctival space, the sub-tenons space, the episcleral space or intrasclerally. It may be delivered to the choroid, the suprachoroidal space, the sub-RPE space the sub-retinal space the epiretinal space, the intravitreal space or the anterior chamber.

In some embodiments, the nanoscale drug-hyperbranched polymer formulation is applied as a coating on an implantable device.

A composition is provided comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound, and wherein sustained release of the active compound occurs over a period of time. The release may occur over a period of minutes, hours, days, months, or years.

A composition is provided comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound, and wherein the composition provides sustained delivery of the compound to a targeted site in a patient. The targeted site may be the vitreous of the eye, and the sustained delivery may over a period of seconds, minutes, hours, days, weeks, or months.

A composition is provided comprising at least one anti-inflammatory compound conjugated to a dendrimer, wherein the composition is encapsulated in a biodegradable particle selected from the group consisting of a PLA nanoparticle or a PGLA microparticle. The dendrimer may be PAMAM-G4-OH and the anti-inflammatory compound may be selected from the group consisting of natural or synthetic steroids or steroid analogs such as fluocinolone acetonide, methyl prednisone or dexamethasone, an antioxidant, an antibiotic such as minocycline, an immunomodulator or an immunosuppressant.

A method of treatment of a neuroinflammation-related disorder is provided, comprising administering a composition comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound, and wherein said neuroinflammation-related disorder is a disease of the retina, optic nerve, central nervous system, the spinal cord or the peripheral nervous system. The disease may be selected from the group consisting of retinitis pigmentosa, age-related macular degeneration, cerebral palsy optic neuritis, blunt and penetrating injuries, infections, sarcoid, sickle cell disease, retinal detachment, temporal arteritis, retinal ischemia, arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, diabetic retinopathy, macular edema, stroke, uveitis, photoreceptor degeneration, autoimmune retinopathy, inherited photoreceptor degeneration, myopic retinal degeneration, retinal pigment epithelial degeneration, diabetic retinopathy, central serous retinopathy, acute zonal outer occult retinopathy, acute multifocal placoid pigment epitheliopathy, multiple evanescent white dot syndrome, cancer associated retinopathy, retinal vasculitis, Alzheimer's disease, Parkinson's disease, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism, aging, and neuronal loss.

Also provided is a method of treating progressive vision loss in a human in need thereof, the method comprising administering to the eye of said human a composition comprising a nanoscale drug-nanoparticle formulation, wherein the formulation

comprises at least one biologically active compound. The progressive vision loss may be associated with at least one condition selected from the group consisting of uveitis, age-related macular degeneration, diabetic retinopathy, and retinitis pigmentosa.

Further provided is a method of treating ocular neuroinflammation in a human in need thereof. The administering may occur once, or may occur two or more times over a period of days, weeks or months.

Further provided is a medical device comprising a nanoscale drug-nanoparticle formulation, wherein the formulation comprises at least one biologically active compound, and instructions for administering the composition.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1. Figure 1 shows microglial cell uptake of dendrimers by flow cytometry. Control cells are indicated and compared with cellular uptake of dendrimers having different functional groups (COOH, NH₂, and OH).

Figure 2. Figure 2A-F is a series of photographs showing retinal biodistribution of free-FITC and dendrimer-conjugated FITC (D-FITC) in healthy Sprague-Dawley (SD) and Royal College of Surgeons (RCS) retinal degeneration model rats with active neuroinflammation. Epi-fluorescence histology from retinal cryosections performed 24-hours and 10-days after intravitreal injection shows (A) D-FITC distribution in normal SD rats at 24-hours; (B) D-FITC distribution in the RCS retinal neurodegeneration model at 24-hours; (C) D-FITC retained within areas of neuroinflammation at 10-days; (D) Free-FITC distribution in SD at 24-hours; (E) Free-FITC uniformly distributed in the RCS rat retina at 24-hours; and (F) Free-FITC cleared from the retina in RCS rats 10-days after injection.

Figure 3. Figure 3A and B are bar graphs. (A) Mean ERG b-wave amplitudes in 9-week RCS rats, four-weeks after a single 1 μ l right-eye injection of 1 μ g or 3 μ g dendrimer-FA (D-FA) or unconjugated, free-FA. There is significant preservation of the ERG amplitude with dendrimer-FA treatment when compared to treatment with FA alone or the control untreated rats. (B) Outer Nuclear Layer cell densities in the same 9-week RCS rats with data from 0.2 μ g/day and 0.5 μ g/day IDIs.

Figure 4. Figure 4 shows the biodistribution of G4OH-⁶⁴Cu complex in mice. G4NH₂ and G4OH dendrimers were complexed with ⁶⁴Cu and injected into adult mice to determine biodistribution of the dendrimers with different surface charge. Regions of interest were drawn for various organs, and radioactivity normalized to dose injected

and weight of the animal was expressed as standard uptake value (SUV) for each of the organs. Adult mice were injected with 50uCi of dendrimer-⁶⁴Cu complex IV. The SUV is plotted over time for different organs.

Figure 5. Figure 5A-D shows a 24-hour post-injection histology of FITC. (A) Dendrimer-FITC (D-FITC) distribution in Sprague-Dawley (SD) retina; (B) D-FITC distribution in RCS rat retina; (C) unconjugated FITC distribution in SD; (D) PLGA-D-FITC microspheres in SD.

Figure 6. Figure 6 shows a schematic of the PAMAM-G4-OH dendrimer-fluocinolone acetonide conjugate synthesis. The glutaric acid spacer relieves the steric hindrance on the dendrimer surface, enabling intracellular drug release. The resultant fluocinolone conjugate is water soluble, even though the drug is not water soluble at comparable concentrations

Figure 7. Figure 7 is a bar graph showing microglia counts within the outer debris zone in the retina of nine week-old control rats and rats given different doses of fluocinolone acetonide.

Figure 8. Figure 8 shows a decrease in the [¹¹C] PK11195 uptake from the first ten minutes to the last ten minutes (similar to control pups) is seen in the postnatal day 5 pup exposed to endotoxin *in utero* that was treated with minocycline 15mg/kg for three days suggestive of a decrease in activated microglial cells. An increase in the PK11195 uptake is noted in the untreated endotoxin exposed pup in the last ten minutes when compared to the first ten minutes suggestive of continued presence of activated microglial cells in the untreated endotoxin pups. This indicates that minocycline treatment has resulted in inhibition of activated microglial cells in the endotoxin exposed pup.

Figure 9. Figure 9A and B are bar graphs showing (A) the mean ERG b-wave amplitudes in RCS rats treated with 0.2 µg/day and 0.5 µg/day sustained doses of FA over four-weeks in the right eye, compared to non-surgical and inactive drug-delivery implant groups, and (B) Mean outer nuclear cell density across the same four groups of RCS rats, according to retinal quadrant.

Figure 10. Figure 10(A-F) shows retinal biodistribution of free-FITC and dendrimer-conjugated FITC (D-FITC) in healthy Sprague-Dawley (SD) and Royal College of Surgeons retinal degeneration model (RCS) rats with active neuroinflammation. (A) D-FITC distribution in normal SD rats at 24-hours. (B) D-FITC distribution in the RCS retinal neurodegeneration model at 24-hours. (C), D-FITC

(RCS) at ten days; (D), Free-FITC (SD) at 24 hours; (E), Free-FITC (RCS) at 24 hours; and (F), Free-FITC (RCS) at ten days.

Figure 11. Figure 11(A-D) shows microglial uptake of D-FITC. (A) ED-1 immuno-histochemical labeling of inner-retinal microglial cells; (B) D-FITC uptake within inner retinal microglia (60x); (C) outer retinal ED-1 labeled activated microglia; and (D) D-FITC uptake within activated microglia.

Figure 12. Figure 12A shows glial-acidic fibrillary protein (GFAP) immunostaining of activated retinal astrocytes. Figure 12B shows dendrimer-FITC uptake by activated retinal astrocytes.

Figure 13. Figure 13A shows GFAP labeling of activated retinal Mueller cells in 5-week RCS rats (lateral view). Figure 13B shows D-FITC uptake by activated retinal Mueller cells (same field as in A). D-FITC uptake within the retinal capillary is shown by the arrow. Figure 13C shows GFAP labeling of retinal Mueller cells (axial view) at inner nuclear layer. Figure 13D shows D-FITC uptake by retinal Mueller cells (axial view).

Figure 14. Figure 14 is a photograph that shows uptake of D-FITC by retinal photoreceptors in 5-week RCS rats.

Figure 15. Figure 15(A and B) shows D-FITC uptake by retinal capillaries. (A) lateral view and (B) lateral view with view of vessel cross-section.

Figure 16. Figure 16(A and B) are photographs showing inner retinal nanoparticle biodistribution in S334-ter-4 rats, seventy two hours after intravitreal injection. Green: FITC-labeled PS nanoparticles, Red: Rhodamine GFAP. (A) 50nm FITC nanoparticles are seen within astrocyte somata. B) 200nm FITC nanoparticles remain confined to the pre-retinal vitreous, and do not appear to be taken into the cells that take up the dendrimer.

DETAILED DESCRIPTION OF THE INVENTION

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. One skilled in the art will recognize many methods and materials similar or equivalent to those described herein, which could be used in the practice of the present invention. Indeed, the present invention is in no way limited to the methods and materials described.

Introduction. Many effective pharmaceutical products and drugs fail to reach target tissue, or fail to remain in the target area, for long enough to achieve clinical

effectiveness. One example is the synthetic corticosteroid fluocinolone acetonide for severe uveitis. Uveitis refers to inflammation or swelling of the eye's structures responsible for its blood supply. These structures are collectively known as the uveal tract, and include the iris, ciliary body, and choroids. Uveitis is classified by the structures it affects, the underlying cause, and whether it is chronic (longer than six weeks), or acute in nature.

Implants for sustained intravitreal delivery of fluocinolone acetonide are FDA-approved for patients with uveitis. However, drawbacks of these implants include multiple incisional procedures for the surgical implantation, a fifty percent incidence of glaucoma, and the non-erodible character of the implants. Another exemplary drug is the anti-inflammatory agent minocycline, which has potential for treating neuroinflammation in the retina and brain.

One underlying goal of the present disclosure is to provide new materials and methods for reducing microglia and glial activation to a cytotoxic and/or phagocytic phenotype associated with neuroinflammation. Such activation is common to a variety of diseases, and by reducing it, the associated destruction of bystander normal tissue can be decreased or eliminated. By doing so, the pathological manifestations of the tissue damage can be significantly reduced, leading to reduced clinical severity of disease, and improved health and quality of life. Any disease in which activated microglia, glia or the infiltration of systemic macrophages play a role is amenable to treatment described herein. In addition, the present invention can address the role that blood vessels play in neuroinflammatory diseases. The examples of diseases described herein are not intended to be limiting, nor are the drugs and pharmaceutical compositions intended to be limiting.

One suitable disease is age-related macular degeneration (AMD). AMD is classified as either wet (neovascular) or dry (non-neovascular). About 10% of patients who suffer from macular degeneration have wet AMD. This type occurs when new vessels form to improve the blood supply to oxygen-deprived retinal tissue. However, the new vessels are very delicate and break easily, causing bleeding and damage to surrounding tissue. The dry type is much more common and is characterized by drusen and loss of pigment in the retina. Drusen are small, yellowish deposits that form within the layers of the retina. These contain antigenic material that can activate retinal neuroinflammation.

Another disease is diabetic retinopathy, which refers to the effect of diabetes on the eye. People with diabetes may develop eye problems such as cataracts and glaucoma, but the disease's effect on the retina is the main cause of vision loss. Over time, diabetes affects the circulatory system of the retina. The earliest phase of the disease is known as background diabetic retinopathy. In this phase, the arteries in the retina become weakened and leak, forming small, dot-like hemorrhages. These leaking vessels can lead to swelling or edema in the retina, causing decreased vision.

Proliferative diabetic retinopathy can follow, in which circulatory problems cause areas of the retina to become oxygen-deprived or ischemic. New, fragile, vessels develop as the circulatory system attempts to maintain adequate oxygen levels within the retina. This phase is called neovascularization and is characterized by delicate vessels that hemorrhage easily. Blood may leak into the retina and vitreous, causing spots or floaters, along with decreased vision. As the disease progresses, continued abnormal vessel growth and scar tissue may cause worsening problems such as retinal detachment and glaucoma. Thus, it is important to control and prevent the neovascularization and thereby eliminate the blood leakage. By virtue of their biodistribution, dendrimers can target and treat neuroinflammatory changes in blood vessels by delivering therapeutic molecules.

Retinitis pigmentosa (RP) refers to a rare, hereditary disease that causes the rod photoreceptors in the retina to undergo gradual degeneration. The disease may be X-linked (passed from a mother to her son), autosomal recessive (genes required from both parents) or autosomal dominant (gene required from one parent) trait. Since it is often a sex-linked disease, retinitis pigmentosa affects males more than females. The course of RP varies. For some people, the affect on vision may be mild. In others disease can progress to blindness.

To aid in describing the present methods of treatment, the following terms associated with eye structure and function are used. The "retina" is a multi-layered sensory tissue that lines the back of the eye. It contains millions of photoreceptors that capture light rays and convert them into electrical impulses. These impulses travel along the optic nerve to the brain where they are turned into images. There are two types of photoreceptors in the retina: rods and cones. The retina contains approximately six million cones. The cones are contained in the macula, the portion of the retina responsible for central vision. They are most densely packed within the

fovea, the very center portion of the macula. Cones function best in bright light and allow color recognition.

The "vitreous" is a thick, transparent substance that fills the center of the eye. It is composed mainly of water and comprises about two thirds of the eye's volume, giving it form and shape. The viscous properties of the vitreous allow the eye to return to its normal shape if compressed. In children, the vitreous has a consistency similar to an egg white. With age it gradually thins and becomes more liquid. The vitreous is firmly attached to certain areas of the retina. As the vitreous thins, it separates from the retina, often causing floaters.

The vitreous is within the posterior segment of the eye, and is one route for administering drugs that target abnormal blood vessel growth that characterizes several of the diseases discussed herein. For example, an antibody fragment (ranibizumab) directed to human vascular endothelial growth factor A (VEGF-A) is injected into the vitreous portion once a month. However, the injection procedure itself can cause serious adverse events such as inflammation of the interior of the eye, retinal detachment, retinal tear, and other problems. Thus, a treatment that requires fewer injections or administration procedures would be likely to decrease unwanted and traumatic adverse effects.

The "ciliary body" is located near the front of the eye, above and below the lens. It is targeted by drugs for treating glaucoma. It produces aqueous humor, so lowering aqueous humor production causes a decrease in the intraocular pressure.

Thus, depending on the location of the pathological event, such as overproduction of aqueous humor, growth of abnormal blood vessels, or unwanted activation of neuroinflammatory cells in the retina, the drug administration according to the present methods and compositions can be tailored to target the specific cells and tissues. Routes of administration to the eye have been studied and described, for example in Lee, T. W. *et al.*, *J. Ocular Pharm.* 20:43-53 and 55-64 (2004). Such routes of administration will be within the skill of the medical professionals by whom the present methods and compositions will be practiced.

Other related conditions and/or diseases that can be treated with particular embodiments disclosed herein include such conditions relating to neuroinflammation, and/or inflammation of the eye, including but not limited to: retinitis pigmentosa, macular degeneration, cerebral palsy, optic neuritis, blunt and penetrating injuries, infections, sarcoid, sickle cell disease, retinal detachment, temporal arteritis, retinal

ischemia, arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, diabetic retinopathy, macular edema, stroke, Alzheimer's disease, Parkinson's disease, brain or spinal cord trauma, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism, aging, and neuronal loss.

Microglia. Microglia are resident members of the dendritic immune system within the retina and central nervous system (CNS) and are activated by many stimuli, including bacterial cell wall lipopolysaccharides and gangliosides within damaged lipid membranes. (Jou, I. *et al.*, *The American Journal of Pathology* 2006; 168:1619-1630; Min, K.J. *et al.*, *Glia* 2004; 48:197-206; Pyo, H. *et al.*, *The Journal of Biological Chemistry* 1999; 274:34584-34589.)

In the case of eye disease, damaged photoreceptor cell membranes provide one antigenic stimulus for microglial activation. Upon activation, microglia assume a phagocytic phenotype, migrate towards the degenerating photoreceptors, and scavenge the debris. In addition, activated microglia orchestrate the recruitment and activation of other microglia via chemotactic cytokines such as CCL-5 (RANTES), macrophage inflammatory proteins (MIP-1 α and MIP-1 β), macrophage chemoattractant proteins MCP-1 and MCP-3.

After assuming the activated phenotype, microglia release cytotoxic free-radicals such as NO and superoxide anion as well as proinflammatory cytokines such as TNF- α , IL-1 and IL-6 incurring further photoreceptor cell damage (bystander lysis). Undampened, this induces a positive feedback cycle of microglia-mediated photoreceptor cell death which in turn exacerbates the bystander lysis of additional photoreceptors, thus coupling of photoreceptor apoptosis and necrosis, accelerating disease progression.

Microglial cells constitute about 10-20% of the cells in the adult brain and form part of the normal surveillance systems in the central nervous system. Microglial cells are known to be activated by stimuli such as trauma, infection, inflammation and ischemia. As a result of these stimuli, there is upregulation of a number of cell adhesion markers along with secretion of pro-inflammatory mediators, generation of reactive oxygen species and peroxynitrites that may lead to further neuronal damage. (Zeng, H.Y. *et al.*, *Invest. Ophthalmol. Vis Sci.* 46:2992-2999, 2005; Bell, M.J. *et al.*, *J. Neurosci. Res.* 70:570-579, 2002.)

One goal for treating neuroinflammatory diseases of the eye is to target drugs to the posterior segment of the eye. With aging, the retinal pigment epithelium (RPE) can sometimes lose its ability to process waste products produced by the photoreceptor cells. Deposits of this waste, called drusen, can distort and damage the retina leading to an eye condition called dry macular degeneration. Other potential sites for targeting drugs in the eye include blood vessels, neuroinflammatory cells, retinal pigment epithelium, optic nerve, cornea, iris, lens and the ciliary body.

The continued persistence of eye disease associated with microglial activation is evidence of the need for new treatments, and an important parameter for introducing a new treatment is its comparison to existing therapeutics. There have been attempts to introduce drugs using sustained release formulations, but none have achieved widespread success.

In the case of nanoparticles, the particle size affects intravitreal kinetics. Fluorescence-labeled-polystyrene micro and nanospheres (2 μm , 200 nm and 50 nm in diameter) were observed in the vitreous cavity of rabbits for over 1 month (Eiji Sakurai, H.O. *et al.*, *Ophthalmic Research* 33:31-36 (2001)). Histological studies using a fluorescence microscope revealed that 2 μm diameter particles were seen in the vitreous cavity and trabecular meshwork, while nanospheres with a diameter of smaller than 200 nm were also observed in the retina as well as the tissues (Eiji Sakurai, H.O. *et al.*, *Ophthalmic Research* 33:31-36 (2001)).

Bourges *et al.* reported studies of ocular drug delivery targeting the retina and retinal pigment epithelium using polylactide (PLA) nanoparticles (NP) (Jean-Louis Bourges, S. E. G. *et al.*, *Investigative Ophthalmology & Visual Science* 44:3562-3569 (2003)). The kinetics and localization of polylactide (PLA) nanoparticle within the intraocular tissues were studied. A single intravitreal injection (5 μl) of the NP suspension (2.2 mg/ml) encapsulating Rh-6G (fluorescent molecule) was performed on Lewis rats. The PLA NPs caused no adverse toxicity effects, and preferentially localized in the RPE cells. Encapsulated Rh-6G diffuses from the NPs and stains neuroretina and RPE cells. This suggested that specific targeting of these tissues is feasible. NPs were found within the RPE cells for four months after a single injection (Jean-Louis Bourges, S. E. G. *et al.*, *Investigative Ophthalmology & Visual Science* 44(8):3562-3569 (2003)). This result shows steady and continuous delivery of drugs can be achieved to RPE cells. If the NP size is small ($\approx < 200\text{nm}$), they may be taken in by the RPEs, and if the size is $\approx > 2 \mu\text{m}$, then they may stay in the vitreous chamber to a large extent.

At present, there are no effective treatments for RP and atrophic (dry) ARMD. Clinical studies have identified that the oral antioxidant, vitamin A palmitate 15,000 IU/d, slows the progression of ERG loss in patients with retinitis pigmentosa (Berson, E.L. *et al. Arch Ophthalmol* 111(11):1456-9 (1993)). The Age-Related Eye Disease Study

5 further identified that beta-carotene, vitamin E, vitamin C, zinc and copper taken orally, reduced the risk for patients with high-risk features of atrophic-ARMD of progressing to the neovascular form by approximately 25% (AREDS Study Group, 2001). In addition, a phase II clinical trial of ciliary-derived neurotrophic factor has been reported. This molecule has strong anti-apoptotic effects. In RP animal models that received the drug,

10 photoreceptor degeneration was slowed, but ERG amplitudes were not preserved (Liang, F.Q. *et al. Mol Ther* 4(5):461-72 (2001); Tao, W. *et al., Invest Ophthalmol Vis Sci* 43(10):3292-8 (2002); Sieving, P.A. *et al., Proc Natl Acad Sci U S A* 103(10):3896-901 (2006); Tao, W., *Expert Opin Biol Ther* 6(7):717-26 (2006); Zeiss, C.J. *et al., Exp Eye Res* 82(3):395-404 (2006)).

15 Steroids, including natural and synthetic glucocorticoids have been shown to exhibit neuroprotective properties through a number of mechanisms: 1) they intercalate within lipid membranes, mechanically stabilizing them (Ignarro, L.J., *J Pharmacol Exp Ther* 182(1):179-88 (1972); Horwitz, L.D. *et al., Free Radic Biol Med* 21(6):743-53 (1996); Wang, Y. *et al., J Pharmacol Exp Ther* 277(2):714-20 (1996)); 2) as

20 antioxidants, they inhibit lipid peroxidation (Eversole, R. R., *et al. Circ Shock* 40(2):125-31 (1993); Horwitz, L.D., *et al. Free Radic Biol Med* 21(6):743-53 (1996); Letteron, P., *et al. Am J Physiol* 272(5 Pt 1):G1141-50 (1997)) ; 3) they inhibit AP-1 (a pro-apoptotic signaling molecule) (Gonzalez, M. V., *et al. J Cell Biol* 150(5):1199-208 (2000); Wenzel, A., *et al. Invest Ophthalmol Vis Sci* 42(7):1653-9 (2001)); and 4)

25 through their potent anti-inflammatory effects, they suppress microglial activation and their ability to produce major histocompatibility complex (MHC) antigens, NO and TNF-alpha (Kiefer, R., *et al., J Neuroimmunol* 34(2-3):99-108 (1991); Hall 1993; Lehmann, C., *et al. Crit Care Med* 27(6):1164-7 (1999); Chang, J., *et al. Neurochem Res* 25(7):903-8 (2000); Drew, P. D. *et al., Brain Res Bull* 52(5):391-6 (2000); Dinkel, K. *et al., J Neurochem* 84(4):705-16 (2003); Lieb, K. *et al., Neurochem Int* 42(2):131-7 (2003); Glezer, I. *et al., Neuroscientist* 10(6):538-52 (2004)).

30

The neuroprotective, anti-apoptotic effects of antioxidants have been demonstrated in photoreceptors in a number of studies (Carmody, R. J. *et al., Exp Cell Res* 248(2):520-30 (1999); Sanvicens, N. *et al., J Biol Chem* 279(38) :39268-78 (2004);

Tanito, M. *et al.* *J Neurosci* 25(9) :2396-404 (2005)). Some glucocorticoid and non-glucocorticoid steroids exert anti-oxidant effects and are neuroprotective in acute and chronic retinal neurodegeneration models (Behl, C. *et al.*, *Mol Pharmacol* 51(4):535-41 (1997); Wenzel, A. *et al.*, *Invest Ophthalmol Vis Sci* 42(7):1653-9 (2001); Dykens, J. A. *et al.*, *Biochem Pharmacol* 68(10) :1971-84 (2004)). Estrogens have been shown to be neuroprotective against oxidative stressors *in vitro* and *in vivo* in transgenic RP animal models (Dykens, J. A. *et al.*, *Biochem Pharmacol* 68(10) :1971-84 (2004); Sanvicens, N. *et al.*, *J Biol Chem* 279(38) :39268-78 (2004)).

Glucocorticoids were found to inhibit Fas, which is a main apoptosis mediator in circulating immunologic cells, such as T lymphocytes and neutrophils, by specific downregulation of Fas gene expression (Cox, G. *J. Immunol.* 154(9):4719-25.1995; Yang, Y. *et al.*, *J. Exp. Med.* 181:1673-82, 1995; Chang, L.C. *et al.*, *J. Endocrinol.* 183:569-83.2004). The capacity of steroids to bind to free radicals was shown in trauma-associated spinal cord neural cell degeneration in humans (Hall 1993) and in oxidative stress-damaged uveal tissue in a Wistar rat model of chronic uveitis (Augustin, A.J. *et al.*, *Br J Ophthalmol* 80(5):451-7, 1996). Intravitreal injections of triamcinolone in albino rabbits (Dierks, Lei *et al.*, *Arch. Ophthalmol.* 123(11):1563-9 2005) enhanced ERG thresholds and were associated with improved retinal morphology.

Glucocorticoid receptors were also localized in apoptotic photoreceptor cells of mouse degenerating retina and can be activated in response to intraocular injection of dexamethasone. Once activated, they inhibit an activator protein (AP)-1 and reduce apoptotic reactions. In dexamethasone-treated eyes, a morphological preservation of retinal outer nuclear cell layer was observed (Wenzel, A. *et al.*, *Invest Ophthalmol Vis. Sci.* 42(7):1653-9 (2001)).

The reports discussed above confirm the need for more effective treatments of these diseases and conditions. Such treatments are provided by the compositions and methods disclosed herein.

Nanodevices and Nanosystems. As used herein, "nanodevices" and/or "nanosystems" may be used interchangeably, and may refer to microparticles or nanoparticles comprising dendrimers and at least one other therapeutic agent.

As used herein, the term "microparticle" or "microparticle system" generally refers to microparticles or microcapsules having a diameter of approximately less than 1 nm to approximately 2,000 nm with a diameter of preferably between 100-500 nm.

Further, "nanoparticles" typically have a diameter range of from less than 1 nm to 1000 nm. As provided herein, certain embodiments of the present invention relate to a series of biocompatible nanoparticle formulations in the form of nanodevices that can be used, for example, as drug delivery vehicles that have been designed to retain and/or delivery
5 drugs or other therapeutic agents over an extended period of time. These formulations permit modification to a desirable size, provide adequate mechanical strength and exhibit exceptional permeability and surface characteristics. Thus, the formulations may contain polymer matrices, liposomes, microcapsules, nanocapsules, controlled-release implants, and the like. A preferred nanoparticle is a soft nanoparticle.

10 In certain particular embodiments, the nanodevices described herein allow for a single dose of the therapeutic agent to the subject, on other embodiments, the nanodevices allow for multiple doses administered to a subject, preferably over an extended period of time. In certain embodiments, the nanodevices allow for a controlled-release of at least one therapeutic agent over a period of at least several
15 hours, at least several days, at least several weeks, or at least several months.

One advantage conferred by the present invention relates to improved control of the permeability of the particles and the release rate of drug conjugated or adsorbed to the nanoparticle periphery. Generally, the nanodevices comprise dendrimers, which may be formed from a variety of materials, including synthetic polymers and
20 biopolymers (e.g., proteins and polysaccharides) and can be used as carriers for other drugs and biotechnology products, such as growth factors and genes or may be used to carry imaging agents. These nanodevices may comprise a polymeric core shell, into which a pharmaceutical drug or other therapeutic agent may be incorporated by way of chemical and/or physical linking or attachment by way of adsorption or chemical
25 conjugation. Alternatively, the therapeutic agent may be conjugated to a polymer within the nanoparticle core. Non-charged small drugs may be attached to larger molecules, preferably charged polymers.

Prior to using the nanoparticles (comprising dendrimers), the particles may be cryoprotected or lyophilized to extend the therapeutic life of the nanoparticle.

30 Cryoprotecting the nanoparticles, with concomitant stabilization, is provided by means of lyophilization. The washed particles are then suspended in cryoprotective solution and lyophilization of the suspension is performed in a suitable lyophilization apparatus.

Dendrimers. According to the present disclosure, pharmaceutical drug compositions are administered to the eye in a nanodevice comprising polymeric

material to which the drug is associated or conjugated. In one embodiment, the polymeric material is in the form of dendrimers, which are manufactured to have a high degree of molecular uniformity, narrow molecular weight distribution, specific size and shape characteristics, and a highly-functionalized terminal surface. For example, ethylenediamine-core poly (amidoamine) (referred to as "PAMAM") dendrimers represent a class of macromolecular architecture called "dense star" polymers. In some embodiments, the dendrimer is a partially acetylated generation 4 or 5 (G4 or G5, respectively) polyamideamine (PAMAM) or polypropylamine (POPAM) dendrimer.

Unlike classical polymers, these dendrimers are manufactured in a series of repetitive steps starting with a central initiator core. Each subsequent growth step represents a new "generation" of polymer with a larger molecular diameter, twice the number of reactive surface sites, and approximately double the molecular weight of the preceding generation. The dendrimer-drug nanodevices can themselves be further packaged for enhanced sustained release. For example, they can be encapsulated into biodegradable poly(lactide-co-glycolide) (PLGA) microspheres as described in more detail herein and in the Examples.

Typically, dendrimer molecules have diameters ranging from 1.5 nanometers to 14.5 nanometers, such as 3 to 10 nanometers, which is comparable to the size of small proteins. They have a highly branched, three dimensional architecture, with high functionality and very low polydispersity (defined as the distribution of individual molecular weights in a batch of polymers). In view of these characteristics, they are capable of carrying many molecules such as pharmaceutical agents, and of possessing homogeneous size, making them suitable for specific modes of administration, such as intraocular.

The pharmaceutical agent of interest can be attached to the dendrimer through a permanent or cleavable bond, or physically encased inside the core of the dendrimer micelle. The dendrimer backbone can also have functional sites for incorporating targeting moieties to facilitate delivery to the desired biological site. The functional sites can also allow for modifying the dendrimer backbone, for example to increase the hydrophilicity and solubility in aqueous media, or to increase solubility in lipid regions.

Advantages of dendrimers include maintenance of drug levels at therapeutically desirable ranges; reducing or minimizing unwanted side effects; decreasing the amount of drug that has to be administered; decreased number of doses, and in the case of ocular administration, less invasive forms of dosing; and enhancing the administration

of drugs that have short half-lives. Some or all of these advantages come into play in the various compositions and methods disclosed herein.

Dendrimers are prepared according to known methods. For example, U.S. Patent No. 5,714,166, provides methods for preparing dendrimers having a variety of sizes and compositions. Yang, H. *et al.*, *J. Biomater. Sci. Polymer Edn.* 17:3-19 (2006) reviewed methods for associating therapeutic agents with dendrimers of the PAMAM compositions. The agent can be entrapped within the dendritic "box" which consists of a densely packed shell on the dendrimer surface. PEGylated dendrimers are useful for holding agents in a hydrophobic core and increasing the water solubility of hydrophobic agents. Dendrimer-drug conjugates can be prepared in which the drug is conjugated to the dendrimer surface, or to PEG which itself is attached to the dendrimer surface.

In the Examples described herein, PAMAM-G4-OH dendrimers were conjugated to fluocinolone acetonide (FA) for injection into the eye. The dendrimer administration of the drug offered advantages over the drug alone. A six-fold lower total FA dose gave greater functional and neuroprotective efficacy than sustained release FA alone, confirming that it is possible to administer less drug when conjugated to dendrimer, yet achieve the same or greater therapeutic effects. Once the dendrimer-FA was taken into the cells, it was unavailable for entering the systemic circulation in the short term, offering another advantage over free drug. Fellow-eye cross-over effects were lower in the dendrimer-FA treated eyes than free FA, which was also attributed to the enhanced cellular uptake of the dendrimer-FA conjugates.

Dendrimers suitable for the present methods and compositions are commercially available. For example, PAMAM dendrimers are available (Aldrich) having specific molecular weight, diameter, and surface groups, depending on the generation. In some Examples herein, PAMAM-G4-OH (fourth generation dendrimers with –OH terminal groups) were obtained from Aldrich. The choice of dendrimers will depend on several factors, including but not limited to: the route of administration, the target tissue, type of pharmaceutical drug utilized, the disease or condition to be treated, the overall health of the subject, the pharmaceutical drug formulation, and others.

"Associated with" means that the drug or pharmaceutical agent, or the imaging or targeting agent (collectively referred to as "agent") can be physically encapsulated or entrapped within the core of the dendrimer, dispersed partially or fully throughout the dendrimer, or attached or linked to the dendrimer or any combination thereof, whereby the attachment or linkage is by means of covalent bonding, hydrogen bonding,

adsorption, absorption, metallic bonding, van der Waals forces or ionic bonding, or any combination thereof.

The association of the agent and the dendrimer may optionally employ linkers, connectors and/or spacers to facilitate the preparation or use of the dendrimer conjugates. Suitable connecting groups are groups which link a targeting director to the dendrimer without significantly impairing the effectiveness of the director or the effectiveness of any other agent present in the dendrimer conjugate. These connecting groups may be cleavable or non-cleavable and are typically used in order to avoid steric hindrance between the target director and the dendrimer. Since the size, shape and functional group density of the dendrimers can be rigorously controlled, there are many ways in which the carried material can be associated with the dendrimer.

For example, (a) there can be covalent, coulombic, hydrophobic, or chelation type association between the agent(s) and entities, typically functional groups, located at or near the surface of the dendrimer; (b) there can be covalent, coulombic, hydrophobic, or chelation type association between the agent(s) and moieties located within the interior of the dendrimer; (c) the dendrimer can be prepared with an interior which is predominantly hollow, allowing for entrapment (*e.g.*, physically within or by association with the interior moieties of the dendrimer) of the agent(s) within the interior (void volume), (*e.g.*, magnetic or paramagnetic cores or domains created by the chelation and reduction of metal ions to the zero valence state within the dendrimer).

Dendrimers containing magnetic interiors can be used for harvesting various bioactive entities that can be complexed with various dendrimer surfaces by use of magnets and the like, wherein the release of the carried material can optionally be controlled by congesting the surface of the dendrimer with diffusion controlling moieties; or (d) various combinations of the aforementioned options can be employed.

Dendrimers useful in the present methods and compositions include the dense star polymers described in U.S. Pat. Nos. 4,507,466, 4,558,120, 4,568,737 or 4,587,329.

The pharmaceutical agents which are suitable for use in the dendrimer conjugates include any materials for *in vivo*, *ex vivo* or *in vitro* use for diagnostic or therapeutic treatment of mammals which can be associated with the dendrimers without appreciably disturbing the physical integrity of the dendrimer, for example, but not limited to fluocinolone acetonide, minocycline, a chemotherapeutic agent, an anti-oncogenic agent, an anti-angiogenic agent, a tumor suppressor agent, an anti-microbial agent, a nucleic acid (including RNA, DNA, cDNA, siRNA, microRNA, and chemical or

synthetic nucleic acid analogs), a protein, a polypeptide, a peptide, an amino acid (including naturally occurring and/or artificial amino acids or analogs), a vitamin, a mineral, a growth factor (such as epidermal growth factor, ciliary neurotrophic growth factor, TGF-beta, bone morphogenic protein, fibroblast growth factor, neurotrophins (NGF, BDNF, NT3, etc.), granulocyte-colony stimulating factor, granulocyte-macrophage colony stimulating factor, platelet-derived growth factor, erythropoietin, thrombopoietin, myostatin, growth differentiation factor 9, basic fibroblast growth factor, epidermal growth factor, hepatocyte growth factor, and others), an angiogenic factor (such as matrix metalloproteinases, vascular endothelial growth factor, angiopoietins, Notch family members, Delta-like ligands, etc.), integrins, an apoptotic factor, a cytokine, or an expression construct comprising a nucleic acid encoding a therapeutic protein, although the present invention is not limited by the nature of the therapeutic agent. It will be appreciated by the skilled artisan that proactive factors may be countered by anti-active factors (such as siRNA, microRNA, antisense, inhibiting antibodies, etc.) and these factors are also considered as part of the instant disclosure.

In further embodiments, the therapeutic agent is protected with a protecting group selected from photo-labile, radio-labile, and enzyme-labile protecting groups. In some embodiments, the chemotherapeutic agent is selected from a group consisting of, but not limited to, platinum complex, verapamil, podophyllotoxin, carboplatin, procarbazine, mechloroethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, bisulfan, nitrosurea, adriamycin, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, bleomycin, etoposide, tamoxifen, paclitaxel, taxol, transplatinum, 5-fluorouracil, vincristin, vinblastin, and methotrexate. In some embodiments, the anti-oncogenic agent comprises an antisense nucleic acid (e.g., RNA, molecule). In certain embodiments, the antisense nucleic acid comprises a sequence complementary to an RNA of an oncogene. In preferred embodiments, the oncogene includes, but is not limited to, abl, Bcl-2, Bcl-xL, erb, fms, gsp, hst, jun, myc, neu, raf; ras, ret, src, or trk. In some embodiments, the nucleic acid encoding a therapeutic protein encodes a factor including, but not limited to, a tumor suppressor, cytokine, receptor, inducer of apoptosis, or differentiating agent. In preferred embodiments, the tumor suppressor includes, but is not limited to, BRCA1, BRCA2, C-CAM, p16, p21, p53, p73, Rb, and p27. In preferred embodiments, the cytokine includes, but is not limited to, GMCSF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN-beta, IFN-gamma, and TNF. In certain

embodiments, the inducer of apoptosis includes, but is not limited to, AdE1B, Bad, Bak, Bax, Bid, Bik, Bim, Harakid, and ICE-CED3 protease. In some embodiments, the therapeutic agent comprises a short-half life radioisotope. }}

In some embodiments of the present invention, the nanodevice further
5 comprises an imaging agent comprising a radioactive label including, but not limited to ^{14}C , ^{36}Cl , ^{57}Co , ^{58}Co , ^{51}Cr , ^{125}I , ^{131}I , ^{111}Ln , ^{152}Eu , ^{59}Fe , ^{67}Ga , ^{32}P , ^{186}Re , ^{35}S , ^{75}Se , ^{175}Yb . In some embodiments, the imaging agent comprises a fluorescing entity. In a preferred embodiment, the imaging agent is fluorescein isothiocyanate.

For treatment of the wet form of macular degeneration, any of the currently
10 available therapeutic agents are amenable to administration using the dendrimer conjugates herein, including Macugen® (pegaptanib sodium injection); ranibizumab, bevacizumab, VEGF-trap, Retaane® (anecortave acetate); and Combretastatin A4 prodrug. Combination with other forms of ARMD treatment are also feasible, such as one of the preceding treatments in combination with light-activated therapy (for
15 example, photocoagulation, or photodynamic therapy), such as with Visudyne®, vitamins (particular vitamin A, vitamin C, vitamin E, or others), minerals (particularly zinc). Certain embodiments may also be combined with radiation therapy, thermal therapy, surgery, and the like.

The surface charge of the dendrimers can have a significant impact on the
20 intracellular transport and drug release from dendrimers (Kannan, S. *et al.*, *J of Biomaterials Science: Polymers Edition*. 2004; 15(3):311; Khandare, J. *et al.*, *Bioconjugate Chem*. 2005; 60:330-337; Perumal *et al.*, *Biomaterials*. 2008; 29:3469-3476). The use of such dendrimers to deliver and prolong the local presence of drugs is further supported by the recognition herein that dendrimers were taken up
25 preferentially by microglial cells both *in vitro* and *in vivo*, as described in detail in Example 1 herein. PAMAM-generation 4 dendrimers with -OH terminal groups (PAMAM-G4-OH) were effective for microglial cell uptake and intracellular drug release as measured in a rat model.

In vitro, microglial cells were treated with or without FITC-labeled PAMAM-G4
30 dendrimers having different functional groups, and the uptake was determined using a flow cytometer. Dendrimers with -OH functional groups showed a higher uptake than dendrimers with -NH₂ or -COOH functional groups. The -OH dendrimers were rapidly endocytosed by microglial cells.

Amino acid dendrimers have also been suggested. For example, Marano *et al.* investigated the ability of lipophilic amino acid dendrimers to deliver anti-vascular endothelial growth factor oligonucleotide (ODN-1) into rat eyes to inhibit laser-induced choroidal neovascularization (CNV). (Marano, R.J. *et al.*, *Nature Gene Therapy* 12:1544-1550 (2005)) The nanodevice showed no adverse effects and was tolerated long term. The dendrimer-ODN-1 conjugate was observed to penetrate the retinal cell layers to reach the retinal pigment epithelium. The conjugate showed appreciable efficacy compared to free nucleotide over a four to six month period. The conjugate showed significant efficacy and was present in RPEs at high levels after the first two months. For the delivery of ODN-1, the therapeutic efficacy was comparable to other delivery methods that required an injection every twenty eight days. However, these treatments have disadvantages as demonstrated by their failure to be embraced by the profession and widely used by patients.

The dendrimer-drug conjugates described herein are therefore preferable for treating ocular diseases in which uptake of drug into cells is desired. The drug is chosen on the basis of the disease to be treated, among other factors. The dendrimer size and composition is not limited to those tested herein. Dendrimer sizes suitable for ocular injection include less than about 500 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 90 nm, less than about 80 nm, less than about 70 nm, less than about 60 nm, less than about 50 nm, less than about 40 nm, less than about 30 nm, less than about 20 nm, less than about 19 nm, less than about 18 nm, less than about 17 nm, less than about 16 nm, less than about 15 nm, less than about 14 nm, less than about 13 nm, less than about 12 nm, less than about 11 nm, less than about 10 nm, less than about 5 nm, less than about 4 nm, less than about 3 nm, less than about 2 nm, less than about 1 nm, or any value there-between or less.

Dendrimers for ocular use may be composed of polymer micelles that incorporate a pharmaceutical drug. Such polymer micelles may be in the form of a nanoparticle comprising a hydrophilic polymer chain as a shell and a hydrophobic polymer chain as a core. The term "nanoscale" is used herein to denote the full range of sizes below the "microparticle" range, or below one micron (one micrometer) in diameter. It is not limited to a specific size cut-off, and the size ranges provided herein are exemplary. Particle sizes of the micelle include less than about 500 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 90 nm,

less than about 80 nm, less than about 70 nm, less than about 60 nm, less than about 50 nm, less than about 40 nm, less than about 30 nm, less than about 20 nm, less than about 10 nm, less than about 5 nm, less than about 4 nm, less than about 3 nm, less than about 2 nm, less than about 1 nm, or any value there-between or less.

5 Some non-limiting examples of hydrophilic polymers that may be used for such an embodiment include a polyalkylene oxide such as polyoxyethylene, polyethylene glycol, polymalic acid, polyaspartic acid and the like. Examples of the hydrophobic polymer which may be used include polylactone, hydrophobic polyamino acid, polystyrene, polymethacrylate ester and the like. A block copolymer may be formed
10 between the hydrophilic polymer chain and the hydrophobic polymer chain. An anionic or cationic charged polymer such as a polypeptide (such as polyaspartic acid), a polyamine or a polycarboxylic acid may be used to form the hydrophobic core. Also, a polyion complex of core-shell type having a core comprising the charged polymer chain and a polymer electrolyte such as a polypeptide, amino acid, or a polypseudo peptide
15 may also be used.

 Generally, non-biodegradable or biodegradable polymers may be used to form the microparticles. In the preferred embodiment, the microparticles are formed of a biodegradable polymer. Non-biodegradable polymers may be used for oral
administration. In general, synthetic polymers are preferred, although natural polymers
20 may be used and have equivalent or even better properties, especially some of the natural biopolymers which degrade by hydrolysis, such as some of the polyhydroxyalkanoates. Representative synthetic polymers are: poly(hydroxy acids) such as poly(lactic acid), poly(glycolic acid), and poly(lactic acid-co-glycolic acid), poly(lactide), poly(glycolide), poly(lactide-co-glycolide), polyanhydrides,
25 polyorthoesters, polyamides, polycarbonates, polyalkylenes such as polyethylene and polypropylene, polyalkylene glycols such as poly(ethylene glycol), polyalkylene oxides such as poly(ethylene oxide), polyalkylene terephthalates such as poly(ethylene terephthalate), polyvinyl alcohols, polyvinyl ethers, polyvinyl esters, polyvinyl halides such as poly(vinyl chloride), polyvinylpyrrolidone, polysiloxanes, poly(vinyl alcohols),
30 poly(vinyl acetate), polystyrene, polyurethanes and co-polymers thereof, derivativized celluloses such as alkyl cellulose, hydroxyalkyl celluloses, cellulose ethers, cellulose esters, nitro celluloses, methyl cellulose, ethyl cellulose, hydroxypropyl cellulose, hydroxy-propyl methyl cellulose, hydroxybutyl methyl cellulose, cellulose acetate, cellulose propionate, cellulose acetate butyrate, cellulose acetate phthalate,

carboxylethyl cellulose, cellulose triacetate, and cellulose sulfate sodium salt (jointly referred to herein as "synthetic celluloses"), polymers of acrylic acid, methacrylic acid or copolymers or derivatives thereof including esters, poly(methyl methacrylate), poly(ethyl methacrylate), poly(butylmethacrylate), poly(isobutyl methacrylate),

5 poly(hexylmethacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), and poly(octadecyl acrylate) (jointly referred to herein as "polyacrylic acids"), poly(butyric acid), poly(valeric acid), and poly(lactide-co-caprolactone), copolymers and blends thereof. As used herein, "derivatives" include polymers having substitutions,
10 additions of chemical groups and other modifications routinely made by those skilled in the art.

Examples of preferred natural polymers include proteins such as albumin, collagen, gelatin and prolamines, for example, zein, and polysaccharides such as alginate, cellulose derivatives and polyhydroxyalkanoates, for example,
15 polyhydroxybutyrate. The in vivo stability of the microparticles can be adjusted during the production by using polymers such as poly(lactide-co-glycolide) copolymerized with polyethylene glycol (PEG). If PEG is exposed on the external surface, it may increase the time these materials circulate due to the hydrophilicity of PEG.

The dendrimer delivery devices are designed to release molecules to be
20 encapsulated or attached over a period of days to weeks. Factors that affect the duration of release include pH of the surrounding medium (higher rate of release at pH 5 and below due to acid catalyzed hydrolysis of PLGA) and polymer composition. A liphasic polyesters differ in hydrophobicity and that in turn affects the degradation rate. Specifically the hydrophobic poly (lactic acid) (PLA), more hydrophilic poly (glycolic
25 acid) PGA and their copolymers, poly (lactide-co-glycolide) (PLGA) have various release rates. The degradation rate of these polymers, and often the corresponding drug release rate, can vary from days (PGA) to months (PLA) and is easily manipulated by varying the ratio of PLA to PGA.

Animal models. Diseases amenable to treatment herein include age-related
30 macular degeneration (ARMD) and retinitis pigmentosa (RP), which share common underlying pathology that can be studied in animals prior to clinical trials. Both diseases exhibit biochemical abnormalities that damage photoreceptors directly and/or indirectly as a consequence of impaired retinal pigment epithelial function. Photoreceptor damage ultimately occurs at the level of cell or mitochondrial

membranes. When mitochondrial outer membranes are damaged via lipid peroxidation (usually as a consequence of oxidative stress within the diseased retina), excessive influx of calcium and superoxide anions can occur into the mitochondrial matrix (Marchetti, P., *et al. J Exp Med* 184(3):1155-60 (1996); Green, D.R., *et al. Science* 305(5684):626-9 (2004); Spierings, D., *et al. Science* 310(5745):66-7 (2005)). In response to this superoxide and/or calcium overload (calpain-induced apoptosis), the mitochondria release cytochrome *c* into the cytosol where it interacts with APAF-1 and caspase-9 to form the apoptosome. This apoptotic process leads to the termination of DNA repair, replication, and transduction and cell death. Consequently, antioxidants and drugs that stabilize mitochondrial outer membranes are neuroprotective through their capacity to antagonize apoptotic cell death.

When photoreceptor-surface or other cell-surface membranes within the retina become damaged, microglial cells that are part of the retinal dendritic-cell-mediated immune system, and are normally quiescent under healthy conditions, can become activated to kill and phagocytize the damaged cells. Activated microglia, the mediators of retinal neuroinflammation, release cytotoxic proteins such as tumor necrosis factor- α (TNF- α), reactive oxygen species such as nitric oxide (NO), cytokines, chemokines, proteases and complement. These can all result in photoreceptor cell damage within the outer retina.

The Royal College of Surgeons (RCS) rat retinal neurodegenerative model is suitable for testing treatments for ARMD and RP disease mechanisms as discussed above. The model demonstrates significant apoptotic (Tso, M. *et al., Invest Ophthalmol Vis Sci* 35(6):2693-9 (1994); Katai, N. *et al., Invest Ophthalmol Vis Sci* 40(8):1802-7 (1999)) and necrotic cell death as a consequence of activated microglia (Thanos, S., *Brain Res* 588(1):21-8 (1992); Thanos, S. *et al., Int J Dev Neurosci* 11(5):671-80 (1993); de Kozak, Y. *et al., Ocul Immunol Inflamm* 5(2) :85-94 (1997); Akaishi, K. *et al. Jpn J Ophthalmol* 42(5) :357-62 (1998); Srinivasan, B. *et al., Science* 310(5745):66-7 (2004); Zeng, H.Y. *et al., Invest Ophthalmol Vis Sci* 46(8) :2992-9 (2005)). The neurosensory retina in the RCS rat is biochemically normal. These rats possess an MERTK tyrosine kinase mutation that resides within the RPE. This has been identified to cause RP in humans (Gal, A. *et al., Nat Genet* 26(3):270-1 (2000)).

The MERTK mutation impairs the RPEs capacity to phagocytize shed rod photoreceptor outer segments. The resulting accumulation of aged, peroxidized and damaged cell membranes within the subretinal space (called the outer debris zone) are

biochemically similar to drusen in patients with ARMD and act as a potent stimulus for retinal microglial activation. The microglial toll-like receptors bind to the damaged lipid membranes and microglial activation occurs. In the model, healthy photoreceptors die or undergo membrane damage as innocent bystanders when exposed to the cytotoxic proteins, NO, proteases and complement released by the activated microglia. These damaged or dying photoreceptors stimulate additional microglial-cell activation in a neuroinflammatory amplification process that leads to massive photoreceptor degeneration, vision loss and an extinguished electroretinogram (ERG). Phagocytic, rhodopsin-containing activated microglia have been immunohistochemically characterized within the postmortem retinas of patients with RP and ARMD (Gupta, N. *et al.*, *Exp Eye Res* 76(4):463-71 (2003)).

The RCS rat model described above is therefore a suitable animal model for the purposes of the present disclosure, and can be used to test dendrimer formulations and hybrid nanodevices and microdevices comprising dendrimer-drug conjugates and nanoparticles or microparticles encapsulating the conjugates. One of skill in the art will be familiar with use of appropriate controls, although guidance is also provided in the Examples herein.

To test the effect *in vivo*, healthy rats and rats with active neuroinflammation (retinal degeneration model) were treated with free-FITC and dendrimer-conjugated FITC. In the disease model, the dendrimer-FITC showed enhanced uptake into the outer retina, specially in and among the outer nuclear layer (ONL), activated microglia, and outer debris zone. This pattern of uptake was not observed in the healthy retinas. These data support one advantage of the present dendrimer-drug nanodevices: dendrimer-drug conjugates demonstrate enhanced uptake in retinal sublaminae undergoing active neuroinflammatory and other neurodegenerative processes.

The results from this experiment also showed that dendrimer conjugates were not cleared from the retina as rapidly as free-FITC. This result supports a second advantage of the nanodevices disclosed herein: dendrimer-drug conjugates were capable of prolonging the residence time of drugs in areas of active neuroinflammation such as glia, microglia, retinal pigment epithelium, and blood vessels, enhancing the pharmacodynamic efficacy, targeting specific retinal sublaminae, reducing the overall amount that must be delivered, and potentially reducing drug side-effects.

The FITC was a test molecule to determine dendrimer delivery parameters. Clinical drugs, such as FA and minocycline, are used to test delivery of a biologically

active molecule, and minocycline treatment led to suppression of microglial cell activation *in vivo*. Briefly, postnatal treatment with minocycline decreased the time course of microglial activation as determined by microPET imaging using the microglial specific ligand [^{11}C] PK11195. (Example 6.)

5 Exemplary drugs and therapeutic agents. The choice of drug or pharmaceutical composition will primarily depend on the disease to be treated or condition to be prevented. For example, the synthetic corticosteroid, fluocinolone acetonide (FA) is FDA-approved for chronic intravitreal delivery as part of a sustained-release system for patients with severe uveitis (Jaffe, G.J. *et al.*, *Ophthalmology* 2000;107:2024-2033;
10 Jaffe, G.J. *et al.*, *Ophthalmology* 2006;113:1020-1027; Jaffe, G.J. *et al.*, *Investigative Ophthalmology & Visual Science* 2000;41:3569-3575). Minocycline is known for its anti-inflammatory properties and is used for treating neuroinflammatory conditions.

Minocycline and doxycycline have been shown to reduce inflammation after brain injury. According to the dendrimer-based methods herein, delivering anti-
15 inflammatory agents such as minocycline specifically to microglial cells leads to a decrease in neuroinflammation and hence decreases the injury. Example 6 herein shows that minocycline decreases neuroinflammation by reducing activation of microglia cells..

The Examples were designed to show that the nanodevices can provide
20 enhanced drug delivery, because of their preferential cellular uptake by the microglial cells, and sustained intracellular drug release resulting in significant improvement in therapeutic effectiveness. Experiments were performed *in vivo* in the rat model of macular degeneration described above. The results are significant and have utility and industrial applicability because dendrimer-based nanodevices can now be developed
25 as a powerful platform in the treatment of neuroinflammation in various ocular pathologies.

An appropriate drug delivery system such as the approach disclosed herein can enhance and sustain the efficacy of these drugs as well as others developed for treating microglia-mediated and other inflammatory diseases of the eye. The results in
30 the Examples herein show that dendrimer-drug conjugates show better therapeutic efficacy in retinal neuroprotection compared to free drug alone or to sustained release drug implants. As one example of this improved route of treatment, the rat model of retinal degeneration was used to compare injected dendrimer-fluocinolone acetonide

(D-FA) nanodevices with free FA injections and sustained-release intravitreal drug delivery implants (IDDIs) that release free FA at a sustained daily rate.

FA is currently approved for ocular administration, for example as a sterile implant. The implant is designed to release FA locally to the posterior segment of the eye at a nominal initial rate of 0.6 $\mu\text{g/day}$, decreasing over the first month to a steady state between 0.3-0.4 $\mu\text{g/day}$ over approximately 30 months. The present compositions improve on the existing treatment, by eliminating the need for implantation and replacing it with a less invasive injection of a dendrimer-FA conjugate.

There was significant preservation of the ERG amplitude with dendrimer-FA treatment when compared to treatment with FA alone or the control untreated rats. Outer nuclear layer cell densities in 9-week RCS rats were compared with data from 0.2 $\mu\text{g/day}$ and 0.5 $\mu\text{g/day}$ IDDIs. Greater cell densities noted in rats treated with D-FA indicates significantly greater neuroprotection in D-FA eyes versus all forms of free-FA ($p < 0.001$ for 1 μg and 3 μg D-FA), including IDDIs. Fellow-eye crossover effects were lower in D-FA treated animals as compared to all animals receiving unconjugated FA, including IDDIs.

Without being bound by a mechanism, this seems to be due to the enhanced cell-uptake of D-FITC nanodevices in RCS rats. When D-FA is taken into cells, it is unavailable for re-distribution into the systemic circulation, unlike what is observed with unconjugated drugs. Consequently, the pharmacodynamic effects are enhanced at lower doses. The D-FA injections resulted in greater functional (ERG) and neuroprotective (ONL counts) efficacy at a six-fold lower total FA dose.

Another important issue in introducing a new treatment is the ability to image uptake and distribution of the targeted drug or device. In the present case of dendrimer nanodevices, dendrimer uptake by microglial cells can be imaged in real time using microPET imaging. To show this, biodistribution of $\text{G4OH-}^{64}\text{Cu}$ complex was examined in mice. (Figure 4.) G4NH_2 and G4OH dendrimers were complexed with ^{64}Cu and injected into adult mice to determine biodistribution of the dendrimers with different surface charge. Regions of interest were drawn for various organs, and radioactivity normalized to dose injected and weight of the animal was expressed as standard uptake value (SUV) for each of the organs. Adult mice were injected with 50 μCi of dendrimer- ^{64}Cu complex IV. G4-OH dendrimers showed better brain uptake than the G4-NH_2 dendrimers. This shows that imaging in real-time using micro-PET can be used

to quantify the uptake of dendrimer-minocycline ^{64}Cu complexes by microglial cells in the brain.

The dendrimer-conjugate compositions may optionally contain other active ingredients, additives and/or diluents. Injectable compositions may be either in suspension or solution form. In solution form, the complex is dissolved in a physiologically acceptable carrier. Such carriers comprise a suitable solvent, preservatives such as benzyl alcohol, if needed, and buffers. Useful solvents include, for example, water, aqueous alcohols, glycols, and phosphonate or carbonate esters. The dendrimer drug conjugate can be incorporated in vesicles or liposomes.

The conjugate can be encapsulated into a polymeric host system that could either be degradable (*i.e.*, lactic-glycolic acid copolymers or a polyanhydride polymer) or nondegradable (ethylene-vinylacetate copolymer). The conjugate can be incorporated into a hydrogel matrix comprising either poly(hydroxyethylmethacrylate) or poly(vinylalcohol).

Examples 7 and 11 herein provide and disclose microparticles for encapsulating dendrimer-drug conjugates. According to Example 7, PLGA (poly(lactic-co-glycolic acid) microspheres encapsulated dendrimer-FITC conjugates. Methods for manufacturing a PAMAM dendrimer with a protected core are known in the art. The present invention is not limited to any particular dendrimers of specific generations. For example, dendrimers of G1, G2, G3, G4, G5, G6, G7, G8, G9, G10 or greater may be used. Molecular weight and the number of terminal groups increase exponentially as a function of generation (the number of layers) of the polymer. Different types of dendrimers can be synthesized based on the core structure that initiates the polymerization process. In preferred embodiments, the dendrimers comprise G4 dendrimers.

The dendrimer core structures dictate several characteristics of the molecule such as the overall shape, density and surface functionality (Tomalia et al., *Chem. Int. Ed. Engl.* 29:5305 (1990)). Spherical dendrimers can have ammonia as a trivalent initiator core or ethylenediamine (EDA) as a tetravalent initiator core. Recently described rod-shaped dendrimers (Yin et al., *J. Am. Chem. Soc.* 120:2678 (1998)) use polyethyleneimine linear cores of varying lengths; the longer the core, the longer the rod. Dendritic macromolecules are available commercially in kilogram quantities and are produced under current good manufacturing processes (GMP) for biotechnology applications.

Dendrimers may be characterized by a number of techniques including, but not limited to, electrospray-ionization mass spectroscopy, ^{13}C nuclear magnetic resonance spectroscopy, ^1H nuclear magnetic resonance spectroscopy, high performance liquid chromatography, size exclusion chromatography with multi-angle laser light scattering, ultraviolet spectrophotometer, capillary electrophoresis and gel electrophoresis.

A variety of enteric coating systems can be employed to help the dendrimer drug conjugate pass through the stomach. The dendrimer drug conjugate can be formulated into a tablet using binders known to those skilled in the art. Such dosage forms are described in Remington's Pharmaceutical Sciences, 18th edition, 1990, Mack Publishing Company, Easton, PA. Suitable tablets include compressed tablets, sugar coated tablets, film-coated tablets, enteric-coated tablets, multiple compressed tablets, controlled-release tablets, and the like.

For treating abnormal or unwanted blood vessel growth, the therapeutic agent can be an anti-angiogenic agent. Examples include anecortave acetate, anti-VEGF aptamer, AMD-FAB, or protein kinase c inhibitor. Other anti-angiogenic agents known in the art may be used. Thus, other anti-angiogenic agents that may be used include, but are not limited to, steroids and angiostatic steroids, metalloproteinase inhibitors, and interferons.

In one non-limiting treatment regimen, PAMAM-G4-OH dendrimers are conjugated with FA and injected into the vitreous of a patient diagnosed with early stage ARMD. The specific concentration and/or formulation of the nanoparticles and/or dendrimers disclosed herein may vary according to any particular disease or condition to be treated. For example, formulations may be administered daily, weekly, monthly, or on an as needed basis, at a drug dose ranging from about at least 1 ng, at least about 1.5ng, at least 2.0 ng, and so on up to 2000 microgram, at least about 2.0 μm , at least about 2.5 μm , at least about 3.0 μm , at least about 5.0 μm , at least about 8.0 μm , at least about 10 μm , at least about 12 μm , at least about 15 μm , at least about 20 μm , at least about 25 μm , at least about 30 μm , at least about 35 μm , at least about 40 μm , at least about 45 μm , at least about 50 μm , at least about 60 μm , at least about 70 μm , at least about 80 μm , at least about 100 μm , at least about 250 μm , at least about 500 μm , at least about 750 μm , at least about 1000 μm , at least about 1500 μm , at least about 1800 μm , at least about 2000 μm , or any value therebetween.

Efficacy of the treatment can be evaluated according to conventional measurements based on industry standards, including but not limited to evaluation by

gross morphology, histology, microbiology, pathology, molecular biology (including analysis of DNA, RNA, or proteins obtained from the treatment subject), as well as input regarding symptoms provided by the subject, if possible.

In one particular embodiment, the dendrimer-FA conjugate is encapsulated in a PLGA nanoparticle and administered to the subject intravenously, orally, buccal, intraperitoneally, into a joint space, rectally, topically, transdermally (particularly for slow release preparations), subcutaneously, intramuscularly, intranasally, intravitreal, suprachoroidal, sub-retinal, episcleral, sub-tenons, intrascleral, epiretinal, injection, by aerosol, or other modes depending on the specific condition or disease to be treated, the organ to be treated, or other factors. Preferably, the nanoparticles comprising the dendrimers disclosed herein are delivered to the eye in the form of eye drops, by deposition of a pellet in or around the eye, by injection into any chamber within the eye, by direct infusion through the eye, and the like. In other embodiments, the inventive nanoparticles comprising the dendrimers disclosed herein are preferably delivered to the nervous system of a subject, preferably the central nervous system. In these particular embodiments, the target cells may include neurons, astrocytes, oligodendrocytes, glial cells or other cells or components associated with the nervous system of the subject.

Alternative nanoparticle formulations include: nanopowders, nanoclusters, nanocrystals, nanospheres, nanorods, nanocups, and other microscopic particles with at least one dimension less than 100 nm.

The Examples below are included for purposes of illustration only, and are not intended to limit the scope of the range of techniques and protocols in which the dendrimer compositions of the present invention may find utility, as will be appreciated by one of skill in the art and can be readily implemented.

EXAMPLES

EXAMPLE 1

DENDRIMER UPTAKE BY MICROGLIAL CELLS *IN VITRO* AND *IN VIVO*

The role of surface charge of dendrimers on the intracellular transport and drug release from dendrimers has been reported (Kannan, S. *et al.*, *J. Biomaterials Science: Polymers* Edition. 2004; 15:311; Khandare, J. *et al.*, *Bioconjugate Chem.* 2005; 60:330-

337), but not previously reported in relation to microglial cells, a target of the present therapeutic efforts.

To study the effect of dendrimer surface group on microglial cell uptake, microglial cells (5×10^5 cells/ml) were treated with or without (Ctrl) 10 μ g of FITC-labeled PAMAM-G4 dendrimers with different functional groups (-OH, -NH₂ and -COOH) at 37°C for 1 hour. The uptake of the dendrimers was estimated using a Beckton Dickinson flowcytometer. A significant increase in the fluorescence intensities with the control cells indicated that all three of the dendrimers entered the cells rapidly, with PANAN-G4-OH dendrimers showing a higher uptake than the other two. The results are shown in Figure 1, and indicate that the -OH dendrimers were rapidly endocytosed by microglial cells.

Retinal biodistribution of free-FITC and dendrimer-conjugated FITC (D-FITC) in healthy Sprague-Dawley (SD) and Royal College of Surgeons retinal degeneration model (RCS) rats with active neuroinflammation was studied. Epi-fluorescence histology from retinal cryosections performed 24-hours and ten days after intravitreal injection is shown in Figure 2A-F as follows. (A) D-FITC distribution in normal SD rats at 24-hours; (B) D-FITC distribution in the RCS retinal neurodegeneration model at 24-hours; the concentration of D-FITC within the outer nuclear layer is denoted by *, and a debris zone, where active neuroinflammation is present, is shown; (C) D-FITC is retained within areas of neuroinflammation at ten days; (D) Free-FITC distribution in SD at 24-hours; (E) Free-FITC is uniformly distributed in the RCS rat retina at 24-hours; (F) Free-FITC had cleared from the retina in RCS rats ten days after injection, as shown in Figure 2.

As evidenced by the RCS rat retina in Figure 2B, the D-FITC demonstrated enhanced uptake into the outer retina, precisely located in and among the ONL, activated microglia and outer debris zone. This pattern of uptake was not observed in any of the healthy SD retinas (Figures 2A, D). Red in the images is due to autofluorescence from the TRITC filter. The retinal pigment epithelium (RPE) demonstrates enhanced FITC fluorescence in retinas that received D-FITC-containing injections. No FITC fluorescence was observed in the RPE of eyes that received free FITC injections. This pattern was qualitatively similar at ten days post injection. The FITC signal present in Figure 2F is autofluorescence from the outer-debris zone. The RPE had separated from the retina in this image and was not visible.

These data suggest that cells that have high endocytotic rates, such as activated microglia and retinal pigment epithelium, demonstrate enhanced uptake of D-FITC conjugates. Consequently, the results support the conclusion that dendrimer-drug conjugates demonstrated enhanced uptake in retinal sublaminae undergoing active neuroinflammatory and other neurodegenerative processes. Furthermore, Figure 2C demonstrates that the dendrimer conjugates were not cleared from the retina as rapidly as free-FITC, Figure 2F.

This Example shows that dendrimer-drug conjugates are capable of prolonging the residence time of drugs in areas of active neuroinflammation, thereby enhancing the pharmacodynamic efficacy, targeting specific retinal sublaminae, and reducing the overall amount that must be delivered as well as potentially reducing drug side-effects.

EXAMPLE 2

PAMAM-G4-OH-FLUOCINOLONE ACETONIDE (D-FA) CONJUGATES (NANODEVICES) COMPARED TO SUSTAINED RELEASE IMPLANTS

Preparation and characterization of D-FA conjugates (nanodevice) were performed as follows. PAMAM-G4-OH dendrimers were chosen for nanodevice preparation. Jayanth Khandare, P.K. *et al.* (2005); P. Kolhe, J.K. *et al.*, *Biomaterials* 27:660-669 (2006). The dendrimers were obtained from Aldrich and were dialyzed to remove any small lower generation impurities that may be present. PAMAM-G4-OH (Mol. Wt = 14,217 Da, 64-OH end groups, size~ 5 nm) was conjugated using a 2-step dicyclohexyl carbodiimide (DCC) coupling reaction (Figure 6). The reaction mixture was stirred for three days and filtered to remove DCU.

The filtrate was dialysed against DMSO for three days by replacing DMSO after each day (dialysis membrane cutoff 1000 Da) to remove unreacted compounds. The dialyzed product was dried under vacuum to obtain conjugate. ¹H-NMR was used to characterize the conjugate. The conjugation ratio was 4.5 molecules of FA per molecule of dendrimer and they are soluble in water, as determined by the proton integration method. This suggests a nanodevice molecular weight of 16,700 Da.

These nanodevices were further characterized by MALDI-TOF mass spectrometry, which suggested a molecular weight of 17,000 Da, agreeing well with the NMR data. The stability of the conjugate in DMSO was tested by monitoring the drug release through a dialysis membrane. HPLC analysis of the samples passing to the

outside of the membrane collected over a period of 24 hours indicated that a negligible fraction of drug from the conjugate was released. This suggests that the conjugate is very stable, and that free (*i.e.*, unconjugated) FA in the conjugate was minimal after purification.

5 Neuroprotective effects of dendrimer-FA (D-FA) nanodevices were determined as follows. The neuroprotective effects of intravitreally-injected D-FA nanodevices to free-FA injections and IDDI's that release free-FA at a sustained daily rate were compared. Figure 3B shows ONL cell densities for 1 μ g and 3 μ g injections of D-FA and free-FA, in addition to the same data for IDDI's shown in Figure (9). Here the 1 μ g and
10 3 μ g refer to the FA content in D-FA. ONL cell densities were higher for D-FA than those measured for IDDI's and free FA injections. ONL cell densities were significantly higher in 1 μ g and 3 μ g D-FA injection groups than in free-FA 1 μ g ($p < 0.001$ for 1 μ g and 3 μ g D-FA). The fellow-eye crossover effects, described in Example 6 for IDDI's and that
15 seen with free-FA 1 μ g and 3 μ g injections in Figure 3B, were less pronounced for the D-FA injected eyes.

Without being bound by a specific mechanism, this is likely due to the enhanced cell-uptake of D-FITC nanodevices in RCS rats as demonstrated in Figure 5. When D-FA is taken into cells, it is unavailable for re-distribution into the systemic circulation as is observed with unconjugated drugs. Consequently, the pharmacodynamic effects are
20 enhanced at lower doses. The D-FA injections resulted in greater functional (ERG) and neuroprotective (ONL counts) efficacy at a six-fold lower total FA dose.

EXAMPLE 3

25 FLUOCINOLONE ACETONIDE WAS NEUROPROTECTIVE IN RCS RATS

According to this Example, the steroid fluocinolone acetonide (FA) was highly neuroprotective in slowing the retinal degeneration of RCS rats when delivered via sustained-release intravitreal drug delivery implants (IDDI's). Figure 9B shows outer
30 nuclear layer cell (photoreceptor cell body) densities among four-groups of nine-week RCS rats that received either FA 0.2 μ g/day, FA 0.5 μ g/day, inactive drug-delivery implants (in right eyes) and non-surgical controls during the four-week study period. Quantitative histological analysis showed that the ONL cell densities in FA 0.2 μ g/day-treated eyes were 2.38 ± 0.12 times greater than in the non-surgical controls ($p < 0.001$)
35 and 4.85 ± 0.24 times higher than in the inactive IDDI group ($p < 0.001$). In the FA

0.5µg/day-treated eyes, ONL counts were 1.78 ± 0.21 times greater than the non-surgical control eyes (p-value with Kruskal-Wallis test=0.02) and 3.56 ± 0.17 times higher than eyes that received inactive IDDIs (p<0.001). No significant differences were observed in the inner nuclear layer cell counts between groups.

5 In other experiments, fluocinolone acetonide preserved electroretinogram b-wave amplitudes in treated eyes. The ONL data were consistent with ERG findings in the same animals (Figure 9A). Comparison across groups showed that mean four-week postoperative ERG b-wave amplitudes recorded from 0.2 µg/day and 0.5 µg/day FA-treated eyes were significantly greater than those of the unoperated control and
10 placebo-treated eyes (p<0.001). ERGs supported the observation made in ONL cell count data that the lower daily dose group demonstrated greater neuroprotective and functional efficacy. Although mean b-wave amplitude values in the eyes treated with FA 0.5 µg/day were quantitatively lower than the values in the eyes treated with FA 0.2 µg/day, there was no statistically significant difference between them (p=0.113).

15 IDDI demonstrated a dose-related cross-over effect in RCS rats. Quantitatively, the left eye-ERG b-wave amplitudes of the FA 0.2µg/day-treated group at the four-week time point were greater than those of control right and left eyes and placebo group-right and left eyes. Mean ERG b-wave amplitudes of left eyes of FA 0.5 µg/day-group were significantly different from those of right and left eyes of unoperated control animals
20 (p<0.03, with power of analysis 0.050:0.810, Kruskal-Wallis p=0.006). No statistically significant differences in mean ERG b-wave amplitudes of left eyes of FA 0.2µg/day-treated group versus right or left eyes of placebo-treated group or right or left eyes of control animals was found. The fellow-eye effects were attributed to drug-crossover. This is commonly seen in small mammals in which the volume of distribution is small.
25 The observation of a dose-related fellow-eye treatment response indicates that free-drug is likely leaving the eye via the episcleral and/or vortex venous systems and becoming systemically distributed, after which it exhibits pharmacodynamic effects in the fellow-eye.

FA suppressed neuroinflammation in RCS rats. In examining the mechanism of
30 action of FA in preserving the ONL and ERG in the RCS rat, quantitative histological analysis of activated retinal microglial cells were performed using ED-1 immunolabelling. The photoreceptor layer and outer debris zone in RCS rats contain damaged lipid membranes that amplified photoreceptor damage by activating microglia to become phagocytic and release toxic substances such as tumor-necrosis factor

alpha (TNF- α). Activated microglial cell densities in the debris layer of FA 0.2 μ g/day-treated eyes showed a five-fold decrease compared to the non-surgical control eyes ($p < 0.001$) and nine-fold decrease compared to eyes that received inactive IDDs ($p < 0.001$). In the FA 0.5 μ g/day-treated eyes, the debris zone activated microglial density was four times lower than non-surgical control eyes ($p < 0.001$) and seven times lower than in eyes that received inactive IDDs ($p < 0.001$).

EXAMPLE 4

PAMAM-G4-OH DENDRIMERS SHOW ENHANCED UPTAKE IN THE OUTER NUCLEAR LAYER OF DEGENERATING RCS RATS, BUT NOT IN NORMAL SD RATS

Figure 5 shows epifluorescence histology from retinal cryosections in Sprague-Dawley (SD) and RCS rats. These were made twenty four hours after intravitreal injection with free, unconjugated FITC, dendrimer-conjugated FITC (D-FITC), and D-FITC encapsulated in PLGA microspheres (PLGA-(D-FITC)). In examining the RCS rat retina in Figure 5B, it is apparent that the D-FITC demonstrated enhanced uptake into the outer retina, precisely located in and among the ONL, activated microglia and outer debris zone. This pattern of uptake was not observed in any of the healthy SD retinas (Figures 5A, C, D). Red in the images is due to autofluorescence from the TRITC filter.

The retinal pigment epithelium (RPE) demonstrates enhanced FITC fluorescence in retinas that received D-FITC-containing injections. No FITC fluorescence was observed in the RPE of eyes that received free FITC injections. This pattern was qualitatively similar at 10-days post injection. These data suggest that cells that have high endocytotic rates such as activated microglia and retinal pigment epithelium demonstrate enhanced uptake of D-FITC conjugates. Consequently, Dendrimer-FA (D-FA) conjugates likely demonstrate enhanced uptake in retinal sublaminae undergoing active neuroinflammatory and other neurodegenerative processes.

EXAMPLE 5

PLGA MICROPARTICLES PREPARED WITH ENCAPSULATED DENDRIMER-FITC CONJUGATES

PLGA microparticles were prepared with encapsulated dendrimer-FITC conjugates as follows. The water-in-oil-in-water (w/o/w) method is suited to encapsulate water-soluble nanodevices like dendrimer-drug conjugates, unlike the o/w method which is ideal for water-insoluble drugs like such as FA. (Jayanth Panyam, M. M. D. *et*

al., *Journal of Controlled Release* 92:173-187 (2003); Jayanth Panyam, S. K. S. et al., *International Journal of Pharmaceutics* 262:1-11 (2003)). Appropriate amounts of PLGA (Mol. Wt. 90,000-125,000 Da.; 75% PLA/25% PGA) were dissolved in chloroform. Separately, a 2.5 % solution of PVA was prepared in cold distilled water, saturated with chloroform.

An aqueous solution of dendrimer-drug conjugates (10% W/V) was added, in two portions, to the PLGA solution with vortexing for one minute after each addition. It was then placed on an ice bath for five minutes and then emulsified using FS 20 Bath sonicator (44-48 KHz, Fisher scientific) for one minute to obtain water-in-oil emulsion.

The primary emulsion was then added in two portions to 8 ml of the PVA solution with intermittent vortexing to obtain the multiple w/o/w emulsion.

The emulsion was placed on an ice bath for five minutes and then sonicated for three minutes. The emulsion was stirred overnight on a magnetic stir plate to allow the evaporation of chloroform and the formation of the nanoparticles. The suspension was transferred into ultra-clear centrifuge tubes and centrifuged at 14,000 rpm for thirty minutes at 4° C in an ultracentrifuge. The pellet was resuspended in distilled water and sonicated for thirty seconds on an ice bath to disperse any aggregates. Centrifugation was repeated two more times at 14,000 rpm, thirty minutes to remove PVA and unencapsulated conjugates from the formulation. The scanning electron microscopy (SEM) and particle size analyzer were used to assess particles sizes. Even though the mean particle size was ~5-10 μm , there was some polydispersity. Fluorescence microscopy suggested that the D-FITC conjugate was "uniformly" encapsulated.

Drug release from PLGA microspheres was studied as follows. The drug release kinetics for free fluocinolone acetonide was analyzed for the same PLGA microspheres. For ~ 5 -10 μm microspheres (containing 5% by mass of FA), the release was ~0.5 $\mu\text{g/day}$ (average) for the first thirty days, with a cumulative release of ~30% in one month. There was 3% initial burst in the first three days. For dendrimer-FA conjugates, the initial burst may be capable of elimination due to their relatively larger size.

EXAMPLE 6

MINOCYCLINE TREATMENT LED TO SUPPRESSION OF MICROGLIAL CELL ACTIVATION IN VIVO

5 According to this Example, postnatal treatment with minocycline decreased the time course of microglial activation as determined by microPET imaging using the microglial specific ligand [^{11}C] PK11195. To demonstrate microglial inhibition with minocycline treatment, [^{11}C] PK11195 uptake in pups at postnatal day 5 with and without minocycline treatment was determined by PET imaging.

10 As shown in Figure 8, a decrease in the [^{11}C] PK11195 uptake from the first ten minutes to the last ten minutes (similar to control pups) was seen in the postnatal day five pup exposed to endotoxin in utero that was treated with minocycline 15mg/kg for three days suggestive of a decrease in activated microglial cells. An increase in the PK11195 uptake is noted in the untreated endotoxin exposed pup in the last ten
15 minutes when compared to the first ten minutes, suggestive of continued presence of activated microglial cells in the untreated endotoxin pups. This indicates that minocycline treatment resulted in inhibition of activated microglial cells in the endotoxin exposed pup.

Postnatal treatment with minocycline improved neurobehavioral outcome at
20 postnatal day eight, as evidenced by neurobehavioral testing of rabbit pups at postnatal day eight. The following animals were observed: (A) control pup, (B) pup exposed to endotoxin 20 $\mu\text{g/kg}$ *in utero* treated with PBS; and (C) endotoxin 20 $\mu\text{g/kg}$ pup treated with 15mg/kg of minocycline, at day eight of life. The endotoxin-exposed pups were more wobbly, with increased tone and decreased balance when compared to the
25 control pups. Hindlimbs were abducted with full range of motion in the control pup A and in the pup treated with 15mg/kg of minocycline (C) but not in the untreated pup (B). Hindlimb tone was greater in pup (B) leading to uncoordinated locomotion and decreased balance.

A dose response for free minocycline in this model showed that 15 mg/kg of
30 minocycline demonstrated greater improvement in motor deficits when compared to 5 mg/kg. A decrease in the white matter injury, which would result in the improvement in motor deficits due to suppression of microglial activation was confirmed by diffusion tensor imaging (DTI) and immunohistochemistry. In conclusion from this Example, minocycline has a specific and clinically relevant effect by reducing the endotoxin-

associated activation of microglial cells in rabbit pups *in utero*. This drug and others with similar effects are suitable for targeting microglial cells using dendrimer compositions disclosed herein.

5

EXAMPLE 7

RAT MODEL FOR MACULAR DEGENERATION

Dose-ranging of minocycline is studied for photoreceptor neuroprotection and suppression of retinal neuroinflammation. The goal of this Example is to determine the efficacy of dendrimer-minocycline nanodevices (D-Mino). Based on the efficacy
10 observed when conjugating fluocinolone acetonide to dendrimers, this Example is designed to evaluate the efficacy of D-Mino conjugates in suppressing retinal neuroinflammation associated with the RCS rat neurodegeneration model. Minocycline is used because it highly neuroprotective and suppresses microglia in the RCS rat retina. Chang, C. J. *et al.*, *Ophthalmic Res.* 2005;37:202-13; Hughes, E. H. *et al.*, *Exp*
15 *Eye Res.* 2004; 78(6): 1077-84; Shimazawa, M. *et al.*, *Brain Res.* 2005;1053:185-94; Zhang, C. *et al.*, *Invest Ophthalmol Vis Sci.* 2004;45:2753-9.

A dose-ranging study is conducted to identify the most efficacious dose of intravitreally-injected unconjugated minocycline (Free-Mino) for retinal neuroprotection in the RCS rat. D-Mino doses are determined based upon the optimal Free-Mino doses
20 and upon FA data. The same five-week homozygous recessive *rdy* albino RCS rat model used in Example 8 is used.

Four groups of five animals are established. These include intravitreal injection groups receiving 1µg Free-Mino, 10µg Free-Mino, a phosphate-buffered-saline (PBS)- (free-dendrimer), and a PBS injection control group. The PBS-(free-dendrimer) group
25 is used to rule out any neuroprotective effects from dendrimers alone. Animals undergo bilateral intravitreal injections, according to established methods. Animals are euthanized at one-month for quantitative histological analysis of photoreceptor and microglial cell counts within retinal sublaminae. Briefly, photoreceptors are counted from 6-µm H&E retinal sections. Retinal microglia are counted from IBA-1 and ED-1
30 labelled retinal slices and wholemounts.

The b-wave ERG amplitudes in RCS rats are recorded as a measure of photoreceptor function, once every two weeks, according to well-established methods. A second four-week dose-ranging study is conducted with Free-Mino, based upon the results of the first study to improve the optimal-dose measurement.

Subsequently, a four-group dose-ranging study for D-Mino is performed, using the same methods. Initial doses are based upon the relative potency of D-Mino to Free-Mino, according to the data for FA of Example 2. The D-Mino dose is further refined in a second study, based upon the degree of neuroprotection and upon the degree of microglial suppression observed in the minocycline studies, compared to the FA and Free-Mino results,

The FITC biodistribution data in Figure 2 demonstrate the improved localization (Figure 2B) and residence time (Figure 2F) that dendrimer conjugation confers to drugs. The optimal dosing interval for D-Mino is measured by performing a three-month study, similar to those outlined in this Example, however re-dosing D-Mino, based upon a 10% decline in ERG amplitude (measured once every two-weeks). In this way, a measure of retinal function is used to establish the first two-week time point when the prior D-Mino injection shows loss of efficacy. By re-dosing, based on retinal function, ERG stabilization should be restored. Over the course of the three-month study, the dose-interval that prevents progressive ERG amplitude loss will also be the optimal one. In addition, this will determine the duration of action for a single D-Mino injection.

EXAMPLE 8

BIODISTRIBUTION AND RESIDENCE TIME OF FREE-FITC AND D-FITC

The goal of this Example is to identify dendrimer-fluocinolone (D-FA) dosage requirements for sustained retinal neuroprotection using intravitreal injections of free-FITC and D-FITC. The experiments are performed in homozygous recessive rdy albino RCS rats of both genders, aged five-weeks, weighing 150-180 grams. Two-groups of animals are established. The free-FITC group consists of six animals that each receive bilateral injections of free-FITC, 1 μ g. Another group of ten animals receives intravitreal injections of D-FITC.

Prior to injection, the animals are anesthetized with ketamine (100 mg/kg) and xylazine (10 mg/kg) via IP injection. A 1 μ L volume of 1 μ g FITC dissolved in a small amount of DMSO is injected via a Hamilton syringe into the vitreous cavity of each eye. Animals injected with free-FITC are euthanized at 10-days, one-month and two-months. The eyes are enucleated, bisected, and the half globes are embedded in Tissue-Tek OCT Medium (Sakura Finetek, U.S.A., Torrance, Ca.) and then quick frozen in liquid nitrogen. Sections (14 μ) are cut on a cryostat (Leica Instruments, GmbH, Nussloch, Germany), mounted (still frozen) on slides with Vectashield (Vector Laboratories,

Burlingame, Ca.) and examined by epifluorescence digital imaging using separate FITC and TRITC filter cubes.

Pictures are taken using a MagnaFire digital camera (Olympus America, Melville, N.Y.). The images are evaluated for retinal sublaminae that are labelled for each time-point. In addition, the relative amount of fluorescence intensity is compared between slices, imaged at each time-point after the illuminator and the camera settings have been adjusted to pre-determined brightness and exposure settings. The epifluorescence illumination setting is adjusted according to the CCD image intensity value measured at pre-determined camera aperture and exposure settings, using a standardized low-concentration of FITC. This FITC-signal calibration process allows comparison of the relative intensity of FITC within each retinal sublamina as a relative measure of the tissue persistence of FITC. The same procedure is performed in animals that receive a 1 μ L volume of 1 μ g D-FITC. In this way, the relative differences in free-FITC and D-FITC biodistribution and tissue persistence are compared, since the CCD camera has a highly linear output.

The D-FITC group consists of ten animals that are euthanized and enucleated for cryosection epifluorescence imaging at 10-days, one-month, two-months, four-months and six-months. For both experimental groups, the fellow eye from each animal that has also undergone intravitreal injection of free-FITC or D-FITC is utilized for fluorophotometric assay of vitreous fluorescence and for FITC fluorescence of tissue homogenates. This allows determination of the relative amount of free-FITC or D-FITC within the vitreous and the tissue of each eye at each time-point. A third group of animals is included, concurrently to examine the biodistribution and tissue persistence of PLGA-(D-FITC). A fourth control group, consisting of one-animal per time-point, serves as bilateral vehicle-injected controls to establish baseline levels of tissue autofluorescence.

Estimation of D-FITC in the eye is performed as follows. For the biodistribution study, at monthly intervals, the eyeball is collected and the tissue is homogenized and dissolved in NaOH. Using a combination of solvent extraction with methanol (D-FITC is soluble in methanol), and further re-suspension in PBS, the D-FITC is extracted. The D-FITC content is quantitated, using a UV/Vis and fluorescence calibration curve previously prepared at various D-FITC concentrations.

Assessment of the duration of pharmacodynamic effect of single 0.2 μ g and 1 μ g D-FA injections, as well as the dose-interval require to maintain the retinal function of

RCS rats, *in vivo*, is performed as follows. One goal is to isolate the efficacy of D-FA nanodevices, as specifically as possible, with appropriate controls. This experiment is performed to examine the duration of action of single intravitreal injections of 0.2 μ g and 1 μ g D-FA in homozygous recessive rdy albino RCS rats of both genders, aged five-
5 weeks, weighing 150-180 grams. Here, 0.2 μ g and 1 μ g refer to the FA content in the D-FA injections.

Four groups of ten animals are established. These include an intravitreal injection group receiving 1 μ g D-FA, a group receiving 0.2 μ g D-FA, a group receiving PBS solution containing (2% DMSO with 1 μ g free-FA and free-dendrimer) in equivalent
10 amounts to the 1 μ g D-FA group, and a non-surgical control group. The PBS-(free-FA, free-dendrimer) group is used to isolate the efficacy of dendrimer-conjugated FA (D-FA). Animals undergo bilateral intravitreal injections, according to the methods described above. Five animals from each group are euthanized at one-month and at three-months for quantitative histological analysis of photoreceptor cell counts and
15 microglial cell counts within retinal sublaminae. These are performed using the methods described below.

The b-wave ERG amplitudes are recorded in RCS rats as a measure of photoreceptor function, once every two-weeks, according to the methods below. The time interval from the current injection to the prior injection is used as one datum in
20 determining the appropriate dosing interval and duration of action for the two D-FA doses, 0.2 μ g and 1 μ g when administered as single bolus intravitreal injections. In addition, the control injection group is re-injected at the same time points that the D-FA 1 μ g group is. This will assure that the control group receives that same number of injections that the D-FA 1 μ g group.

Electroretinography is performed as follows, using stimulation, recording and data analysis routines developed using Labview software (National Instruments, Austin Texas). Animals are dark-adapted for 12-hours overnight. Prior to testing, animals are anesthetized with an intraperitoneal injection of Ketamine 67mg/kg and Xylazine
25 10mg/kg. Pharmacological mydriasis is induced, bilaterally with 1% Tropicamide and 10mg/kg. Phenylephrine. Topical 0.9% saline is periodically applied to the corneas to
30 prevent dehydration. ERG responses are recorded using platinum wire loop corneal electrodes, from both eyes simultaneously. Reference platinum needle electrodes are placed in the ears. White clear LEDs CMD204 (UWC Series T-1), with a luminous intensity of 1000mcd, are employed for flash stimulation. Fifteen 1-millisecond flashes

are delivered to both eyes simultaneously with constant 10-second inter-stimulus intervals. The ERG responses are amplified using a gain of 5000, bandpassed between 10 and 100 Hz.

ERGs are averaged and mean b-wave amplitude is determined for each animal at each time point. If the ERG b-wave amplitude has declined by 10% from the average of previous two recording sessions, then a re-dose intravitreal injection is performed according to the methods outlined above.

For histology, the eyes are enucleated and placed in Karnovsky's fix overnight, at 4°C. After an 18-hour immersion period, eyes are rinsed with 0.01 M phosphate buffer solution. Eyes are prepared for sectioning, by making a transverse cut along the horizontal meridian. After grossing, eyes are dehydrated in serial dilutions of alcohol, cleared with Pro-Par (Xylene-substitute, Anatech Ltd., Battle Creek, MI) and embedded in paraffin containing DMSO (Fisher). Serial 6µm-thick whole-eye sections are obtained. Paraffin sections are mounted on Poly-L-lysine coated glass slides and stained using Harris' Haematoxylin and Eosin.

Microglia staining in whole-mounted retina is performed as follows. Rat eyes are enucleated and fixed briefly in buffered ten percent formalin. They are then bisected and the nasal section fixed for a further 4 hours at 4°C. The temporal section is embedded in OCT compound (Sakura Finetek, USA Inc., Torrance, CA) and flash-frozen in liquid nitrogen for cryo-sectioning. Orientation is preserved for each section.

After fixation, as detailed above, the retinas are carefully removed and placed in PBS containing 1% Triton X-100 for one hour at room temperature. Retinas are then incubated overnight at 4°C in a cocktail of Iba-1 antibody (Wako Chemicals USA Inc, Richmond, VA) at 1:200 dilution and ED-1 antibody (Serotec Ltd. Oxford, UK) at 1:100 dilution, in PBS containing 0.1% Triton X-100 and four percent normal goat serum. After several washes, retinas are incubated in the appropriate fluorescently labeled secondary antibody (Sigma-Aldrich, St. Louis, MO) solution for two hours at room temperature, mounted ILM side down on glass slides and cover-slipped with Vectashield (Vector Laboratories, Burlingame, CA).

Microglia staining in transverse retinal sections is performed as follows. Sections (10-micron) are cut on a cryostat (Leica Instruments GmbH., Nussloch, Germany), recovered on Vectabond (Vector Laboratories, Burlingame, CA) coated slides and fixed for one minute in acetone. Slides are allowed to air dry and are stored at 4°C until used. Sections are then re-hydrated in Tris buffered saline (pH 7.4) and

non-specific staining blocked with 5% normal goat serum. Sections are incubated in Iba-1 and ED-1 antibody solution, as detailed above, and developed for fluorescence microscopy using the appropriate fluorescently labeled secondary antibodies.

For each eye, ERG waveforms are averaged over fifteen traces for each recording session. The b-wave amplitude is defined as the peak-to-peak amplitude of the leading positive slope (from the lowest point of the a-wave to the highest point of the b-wave). This is measured from each of the averaged ERG waveforms. Mean and standard deviation b-wave amplitude are measured for each eye of each animal for each time point. Group means and standard deviations are also computed for right and left eyes at each time point. Parametric and non-parametric statistical analyses are performed on these data.

For histological cell count analysis, retinal photomicrographs are taken of both right and left eyes, using an Olympus B-MAX 50 microscope (Japan), with Olympus Magnafire PM30/PM20 digital camera, at 40× magnification. In all eyes, five photomicrographs are taken in each of the four retinal quadrants (i.e., twenty photomicrographs per eye): 1) superior temporal; 2) superior nasal; 3) inferior temporal; 4) inferior nasal. Care is taken to precisely correlate these regions between eyes.

Each photomicrograph is analyzed using a grid counting method. A masked observer counts the number of outer nuclear layer (ONL) and inner nuclear layer (INL) cells in each of the two grid regions defined per photomicrograph. The counts are averaged for the ONL and INL in each tested retinal region for right and left eyes in each group.

For microglial cell count analysis, positively stained microglial cells are counted using an Olympus B-MAX 50 microscope (Japan) with the appropriate filter sets. In retinal whole-mount preparations, counts are made in six standardized fields at three defined levels within the retina: a layer at the level of inner limiting membrane (ILM layer), a layer at the level of inner plexiform layer (MID layer), and a layer at the level of retinal photoreceptors (PHR layer). In retinal transverse sections, activated microglia are counted in the photoreceptor debris zone in two superior and two inferior retinal fields.

One way ANOVA with concomitant paired t-test, when valid, is performed for both the 4-week postoperative ERG data and the averaged ONL and INL cell count data across treated and untreated eyes in all groups. The measured microglial cell count values are averaged to calculate means and standard deviations for every

microglial cell layer among experimental groups. The analyzed data are graphed. Although power of analysis established that five animals per group are adequate to perform parametric statistical comparisons, a non-parametrical (Kruskal-Wallis or Mann-Whitney) analysis can be performed every time a one way ANOVA is applied.

5

EXAMPLE 9

ASSESSMENT OF SUSTAINED INTRAVITREAL D-FA EFFICACY *IN VIVO*

The present Example is designed to study sustained D-FA delivery via repeated intravitreal injections, at the appropriate dose and at the appropriate dosing interval for sustained retinal photoreceptor neuroprotection and preservation of the ERG b-wave amplitudes in RCS rats. Using either 0.2 μ g or 1 μ g D-FA injections (on a drug basis), RCS rats are divided into two groups of fifteen animals. Animals from one group undergo intravitreal injection of D-FA while the other group receives PBS containing (2% DMSO with free-FA and free-dendrimer). ERG recordings are performed once per month. D-FA and control re-injection are performed when ERG b-wave amplitudes in the D-FA group have declined by 10% as compared to the mean of the two-previous ERG recordings.

Five animals per group are euthanized at months three, seven, and nine for quantitative histological analysis of photoreceptor cell counts, total microglia and microglial localization, as well as number of activated microglia. By summing the total D-FA required to maintain ERG amplitudes, and dividing this number by the duration of neuroprotection, the required release rate of D-FA to maintain ERG amplitudes from sustained-release PLGA-(D-FA) is estimated.

From this Example, the dosage and the interval to maintain the function of the retina can be determined, since only a 10% reduction in the ERG is allowed. The control allows the effect of the nanodevice to be determined. The residence times of the dendrimers in the vitreous can be determined, as well as the cellular distribution in the retinal sublaminae. If the dendrimers are found in the outer retina beyond at least two to three weeks, the efficacy of the nanodevices could last for significantly longer than one month and provide long-term treatment. Therefore, this Example provides additional data for evaluating long-term clinical effectiveness of the devices disclosed herein.

EXAMPLE 10

INJECTABLE, SUSTAINED-RELEASE INTRAVITREAL DRUG-DELIVERY PLATFORM BASED UPON
HYBRID [PLGA-(D-FA) NANODEVICE] MICROSPHERES.

This Example is performed to determine the biodistribution and residence time of
5 PLGA-(D-FITC) microspheres. A group of ten animals is added to the biodistribution
study of Example 9, and these animals undergo bilateral intravitreal injection of PLGA-
(D-FITC) microparticles. These animals are euthanized at 10-days, one-month, two-
months, four-months and six-months.

PLGA-D-FA microspheres are prepared with the intent of achieving sustained
10 delivery of $\sim 0.03 \mu\text{g/day}$ of FA in D-FA for 180 days. One intravitreal injection of $1.0 \mu\text{g}$
D-FA is effective for one-month ($0.03 \mu\text{g/day}$). Thus, the dendrimer-FA conjugate is
estimated to clear the vitreous chamber in a few days. A significant portion of these
appear to be taken in by the ONL, where the drug is released over a period of time, in a
manner that suggests that the pharmacodynamic effect lasts for about a month.

15 Therefore, an appropriately designed PLGA microsphere encapsulation of the D-FA will
provide a controlled release of D-FA, which will then enter the retinal cells and release
the drug.

The size and molecular weight of the D-FA conjugates ($\sim 7 \text{ nm}$, $\sim 20 \text{ kDa}$) are
comparable to small proteins. In PBS ($\text{pH}=7.4$), the ester bond between the drug and
20 dendrimer is stable. The hydrolysis of this bond most likely occurs over a period of time
in the acidic lysosomal pH. Therefore, appreciable drug release from the dendrimer
inside the PLGA microsphere is not anticipated, especially with 75% PLA. (DD 1990; P.
Kolhe 2006).

It has been previously shown that PLA microspheres can last as long as long as
25 sixty days in PBS buffer of 7.4 pH. PLGA microspheres of size $\sim 10 \mu\text{m}$ and of
copolymer composition 90% PLA and 10% PGA (PLGA(90:10)) also resist degradation
for extended periods.(DD 1990). This size of the microsphere also significantly
improves the intravitreal residence time. PLGA (75:25) is used for sustained delivery of
the drug to the eye. Based on diffusion coefficients of larger molecules ($\sim 10^{-13}$
30 cm^2/sec) from PLGA microspheres, the D-FA diffusion is expected to take place over a
period of months, thereby providing sustained drug delivery over a six month period.
(Sandora 2001)

The PLGA microparticles (composition:75% PLA/25%PGA, molecular weight
 $\sim 90\text{-}125 \text{ kDa}$) containing dendrimer-FITC (for biodistribution studies) and dendrimer-FA

(for release studies and in vivo efficacy studies), are prepared using a water-in-oil-in water (w/o/w) method as described in the Examples. (Sanjeeb K Sahoo, J. P. *et al.*, *Journal of Controlled Release* 82:105-114 (2002); Jayanth Panyam 2003). The solution of particles is filtered to produce a particle size in the range of ~5 – 10 μm . The particle size is characterized by SEM and a Malvern particle sizer. The presence and distribution of nanodevice inside the microparticle is assessed with an encapsulated D-FITC nanodevice, using fluorescence and confocal microscopy.

For *in vitro* release studies, PLGA-dendrimer-FA microspheres are placed in a micro centrifuge tube containing 250 μl of 0.1M phosphate buffer (PBS pH 7.4) at 37°C in a shaker rotated at 100 rev/min. Initially, buffer solution is collected every day for drug analysis, and replaced with new buffer solution. After seven days, the sample is collected every alternate day, with the frequency decreasing to every five days at longer times. The collected samples are kept under the hood overnight to drive off the buffer solution.

The quantification of dendrimer-FA (D-FA) is performed using a C5 silica based HPLC column (250mm $\times$ 4.6 mm, 300 Å). (Mohammad T. Islam, X. S. *et al.*, *Anal. Chem.* 77:2063-2070 (2005)) The mobile phase for the elution of D-FA is a linear gradient beginning with 90:10 water/acetonitrile (ACN) at a flow rate of 1 mL/min, reaching 50:50 after 30 minutes. Trifluoroacetic acid (TFA) at 0.14 wt% concentration in water as well as in ACN is used as a counter-ion to make the dendrimer-conjugate surfaces hydrophobic. (Mohammad T. Islam, X. S. *et al.*, *Anal. Chem.* 77:2063-2070 (2005))

The conjugates are dissolved in the mobile phase (90:10 water/ACN). The detection of D-FA in eluted samples is performed at 210 and 238 nm, the λ_{max} of PAMAM-G4-OH and FA respectively. The calibration curve is prepared based on 238 nm for FA in the conjugated form. The elution time for D-FA in this system is expected to be at eight minutes. To quantify the possible presence of free FA, a C-18 reverse phase symmetry shield column from Waters is used (3.9 $\times$ 150 mm column, 5 μm ; W20881A). (Glenn J. Jaffe 2000; Glenn J. Jaffe 2000) A 1:1 mixture of acetonitrile and 0.02% sodium acetate (pH 4.0) is used as the mobile phase. The sample is injected at 1 ml/min and the drug FA elutes at around 3.0 minutes. The detection of eluted samples is performed at 238 nm (λ_{max} of FA), with an appropriately prepared calibration curve.

EXAMPLE 11

PHARMACODYNAMIC EFFICACY OF HYBRID [PLGA-(D-FA) NANODEVICE] MICROSPHERES IN
MAINTAINING ERG AMPLITUDES AND PHOTORECEPTORS IN THE RCS RAT

5 After collecting and analyzing six months of data from experiments as described
in Example 10, this Example is performed to examine the pharmacodynamic efficacy of
single PLGA-(D-FA) injections in maintaining ERG b-wave amplitudes and preserving
photoreceptor cell counts in aging RCS rats. For this study, PLGA-(D-FA)
microspheres are constructed that release D-FA at the release rate that is
10 experimentally validated to be efficacious according to Example 10.

 Using the same protocols outlined above for intravitreal injections, one group of
five-week RCS rats undergo intravitreal injection of PLGA-(D-FA) microspheres and
another group will receive intravitreal injection of PBS containing D-FA and free-PLGA
microparticles. Two experimental groups are established of fifteen animals per group.
15 ERG recordings are performed once per month. PLGA-(D-FA) and D-FA+free-PLGA
microsphere re-injection are performed when ERG b-wave amplitudes have declined by
10% as compared to the mean of the two-previous ERG recordings. Five animals per
group are euthanized at months three, six, and nine for quantitative histological analysis
of photoreceptor cell counts, total microglia and microglial localization, as well as
20 number of activated microglia.

 PLGA micro spheres (75%PLA/25%PGA) are designed that are intended to
release D-FA over a four to six month time, so that the efficacy can be maintained for at
least a six month period. A slow-release phase is expected for a sixty to seventy day
period, and a somewhat faster release at later times when the particle erodes
25 appreciably. A significantly slower sustained release profile is expected for the
nanodevice. Very little initial burst for the D-FA (17 kDa) is expected.
For biodistribution, compared to D-FITC, PLGA-D-FITC are expected to show
significantly higher intravitreal fluorescence, and significantly lower outer retinal
fluorescence at comparable times. D-FITC is expected to be present in the outer retina
30 at much longer times. If *in vitro* studies show that D-FA is released at a faster rate than
desired, then the composition of the PLGA can be changed to 90%PLA/ 10%PGA, or
pure PLA.

EXAMPLE 12

ELECTRORETINOGRAPHIC FINDINGS ASSOCIATED WITH INTRAVITREAL INJECTIONS OF
DENDRIMER-CONJUGATED AND UNCONJUGATED FLUOCINOLONE ACETONIDE IN RCS RATS

In this Example, the efficacy of D-FA nanodevices was compared to the free
5 drug and comparable drug doses. The efficacy was analyzed using electroretinography
(ERG) and histology. The results indicate that just one intravitreal injection of the
dendrimer nanodevice at the lowest administered dose, not only prevents ERG
reduction, it enhances the ERG levels substantially.

Thirty albino RCS rats aged five weeks were divided into six equal groups. The
10 experiments were terminated at nine weeks. Thus, the efficacy was evaluated over the
four-week peak period of retinal degeneration in the RCS rats. The groups were:

1. intravitreal injection of 3.0 μg dendrimer-conjugated fluocinolone acetonide
(FA) in 1.0 μL of Phosphate Buffered Solution (PBS);
- 15 2. intravitreal injection of 1.0 μg dendrimer-conjugated FA in 1.0 μL of PBS;
3. intravitreal injection of 3.0 μg FA in 1.0 μL of PBS;
4. intravitreal injection of 1.0 μg FA in 1.0 μL of PBS;
5. intravitreal injection of 1 μL of PBS (vehicle control);
6. light-exposed unoperated controls.

20 Comparison of initial (at five weeks of age) ERG a-wave amplitudes showed no
statistically significant difference between experimental groups ($p \geq 1.0$). Comparison of
end-point (at nine weeks of age) ERG a-wave amplitudes showed statistically
significant differences between right and left eyes of D-FA treated animals. Thus, D-FA
25 preserved photoreceptor response amplitudes in a dose-dependant manner. This was
not true for PBS-injected or free-FA injected eyes.

The dendrimer-conjugation enhanced FA's neuroprotective effects over the non-
conjugated drug. This phenomenon was also observed for ERG b-wave amplitudes.
All FA-treated animals demonstrated the greatest stabilization of ERG b-to-a-wave
30 amplitudes. ERG implicit times were significantly lower in D-FA treated animals as
compared to FA-treated, control or untreated animals. This is consistent with the
finding that dendrimer conjugation of FA improves its pharmacological efficacy.

The Table below shows ERG a-wave and b-wave amplitude change (Mean $\pm$ Standard Error) among six experimental groups at the end of the study, as compared to the ERG amplitudes at the beginning of the study (expressed as a percentage of loss (-)/gain (+) from the initial amplitude).

5

Experimental Group	ERG a-wave		ERG b-wave	
	OD	OS	OD	OS
Dendrimer-FA 3.0 μg	+ 17.8 $\pm$ 18.1 %	+ 30.9 $\pm$ 20.1 %	- 13.8 $\pm$ 9.8 %	- 34.7 $\pm$ 13.2 %
Dendrimer-FA 1.0 μg	+ 88.6 $\pm$ 34.3 %	+ 1.7 $\pm$ 11.7 %	- 15.8 $\pm$ 9.9 %	- 34.3 $\pm$ 9.2 %
Free FA 3.0 μg	- 12.5 $\pm$ 28.8 %	- 10.5 $\pm$ 30.1 %	- 42.6 $\pm$ 3.3 %	- 40.8 $\pm$ 3.5 %
Free FA 1.0 μg	- 41.9 $\pm$ 13.4 %	- 47.8 $\pm$ 7.7 %	- 64.2 $\pm$ 8.6 %	- 55.6 $\pm$ 3.6 %
PBS	- 73.9 $\pm$ 2.3 %	- 76.1 $\pm$ 2.4 %	- 62.9 $\pm$ 2.2 %	- 63.4 $\pm$ 2.9 %
Unoperated	- 73.9 $\pm$ 4.5 %	- 79.1 $\pm$ 3.9 %	- 60.8 $\pm$ 7.8 %	- 72.7 $\pm$ 4.8 %

The data disclosed in this Example provide additional evidence of the benefits of dendrimer conjugation, as shown in the Table above. For the first time, a-wave amplitudes increase over the period of time where a greater than 75% reduction in amplitudes was observed.

10

15

EXAMPLE 13

DENDRIMER BIODISTRIBUTION AND EFFICACY AND NANOPARTICLE BIODISTRIBUTION AND EFFICACY

20

According to this Example, PAMAM-G4-OH dendrimers showed enhanced uptake in the outer nuclear layer of degenerating RCS rats, but not in normal SD rats, suggesting that they can be used to target neuroinflammation. Retinal biodistribution of free-FITC and dendrimer-conjugated FITC (D-FITC) were studied in healthy Sprague-Dawley (SD) and Royal College of Surgeons retinal degeneration model (RCS) rats with active neuroinflammation.

25

The first set of results is shown in Figure 10, epi-fluorescence histology from retinal cryosections performed twenty four hours and ten days after intravitreal injection,

showing (A) D-FITC distribution in normal SD rats at twenty four hours, and (B) D-FITC distribution in the RCS retinal neurodegeneration model at twenty four hours. The concentration of D-FITC within the outer nuclear layer (denoted by *) and debris zone are significant findings.

5 In examining the RCS rat retina in Figure 10B, it is apparent that the D-FITC demonstrated enhanced uptake into the outer retina, precisely located in and among the ONL, activated microglia and outer debris zone. This pattern of uptake was not observed in any of the healthy SD retinas (Figures 10A, D). Red in the images is due to autofluorescence from the TRITC filter. The retinal pigment epithelium (RPE)
10 demonstrates enhanced FITC fluorescence in retinas that received D-FITC-containing injections. No FITC fluorescence was observed in the RPE of eyes that received free FITC injections. This pattern was qualitatively similar at ten days post injection. The FITC signal present in Figure 10F is autofluorescence from the outer-debris zone. The RPE has separated from the retina in this image and is not visible.

15 These data suggest that cells that have high endocytotic rates such as activated microglia and retinal pigment epithelium demonstrate enhanced uptake of D-FITC conjugates. Consequently, the results show that dendrimer-drug conjugates likely demonstrated enhanced uptake in retinal sublaminae undergoing active neuroinflammatory and other neurodegenerative processes. Furthermore, Figure 10C
20 demonstrates that the dendrimer conjugates are not cleared from the retina as rapidly as free-FITC, Figure 10F. Thus dendrimer-drug conjugates are capable of prolonging the residence time of drugs in areas of active neuroinflammation, enhancing the pharmacodynamic efficacy, targeting specific retinal sublaminae, reducing the overall amount that must be delivered and potentially reducing side-effects.

25 The D-FITC localization in the outer-retina was further analyzed by immunohistochemical labeling for activated microglial cells, astrocytes, and Mueller cells and the photoreceptors (as shown in Figures 11-15). The results suggest that dendrimers are selectively localizing in these cells associated with neuroinflammatory processes. There was minimal accumulation in the other cells in the inner and outer
30 retina. Figure 11 shows Microglial uptake of D-FITC. (A) ED-1 immuno-histochemical labeling of inner-retinal microglial cells; (B) D-FITC uptake within inner retinal microglia (60x); (C) outer retinal ED-1 labeled activated microglia; and (D) D-FITC uptake within activated microglia.

Figure 12 provides further results. (A) Glial-acidic fibrillary protein (GFAP) immunostaining of activated retinal astrocytes; and (B) Dendrimer-FITC uptake by activated retinal astrocytes.

Figure 13 shows that D-FITC is taken up by the activated retinal Mueller glial cells. Figure 13A shows GFAP labeling of activated retinal Mueller cells in five week RCS rats (lateral view); (B) shows D-FITC uptake by activated retinal Mueller cells (same field as in A), and D-FITC uptake within the retinal capillary (arrow); (C) shows GFAP labeling of retinal Mueller cells (axial view) at inner nuclear layer. The INL cell bodies are particularly notable (outlined by Mueller cells processes); and (D) shows D-FITC uptake by retinal Mueller cells (axial view).

GFAP staining in these cells was specific to the activated phenotype (Figure 12A). Imaging of the FITC epifluorescence (Figure 12B) in the same micrographic field shows significant D-FITC uptake. In addition, a small intra retinal capillary is also shown to uptake D-FITC (Figure 13B, marked by arrow). Co-localization of GFAP and D-FITC was also seen in Mueller cells as their processes pass the outer nuclear layer (Figure 13 C and D). In Figure 14, photoreceptors are labeled by D-FITC uptake.

The retinal vasculature demonstrated a high degree of D-FITC uptake (Figure 15), within the vessel walls. This was true for the inner retinal circulation as well as much deeper fine capillaries within the retinal parenchyma.

Nanoparticle biodistribution was also investigated. To study the role of the particles size and the hardness, monodispersed FITC-labeled PS nanoparticles, of size 50 nm and 200 nm were used. The results are shown in Figure 16, which shows inner retinal nanoparticle biodistribution in S334-ter-4 rats, seventy two hours after intravitreal injection. Green: FITC-labeled PS nanoparticles, Red: Rhodamine GFAP. (A) 50nm FITC nanoparticles are seen within astrocyte somata. (B) 200nm FITC nanoparticles remained confined to the pre-retinal vitreous, and did not appear to be taken into the cells that take up the dendrimer.

The results discussed in the examples above show that: (1) Upon intravitreal injection, the dendrimers selectively localize in the cells primarily associated with neuroinflammation, such as activated microglial cells, Mueller cells, astrocytes and macrophages; (2) the dendrimers appear to be present in these cells, even after weeks; (3) dendrimer-drug conjugate releases the conjugated drugs over a period of a month; (4) the *in vivo* efficacy of one injection of the dendrimer nanodevice in the vitreous

chamber is significantly better than both the free drug and a controlled release implant, over a one-month period. In fact, electroretinography (ERG) measurements suggest that the dendrimer nanodevice, not only prevents further degeneration, it may actually enhancing retinal health.

5

While the description above refers to particular embodiments of the present invention, it will be understood that many modifications may be made without departing from the spirit thereof. The presently disclosed embodiments are therefore to be
10 considered in all respects as illustrative and not restrictive. All patents and publications referenced are incorporated herein by reference.

CLAIMS

What is claimed is:

1. A composition comprising a nanoscale drug-nanoparticle formulation,
5 wherein said formulation comprises at least one biologically active compound.
2. The composition of claim 1, wherein the size of the nanoparticle is less than about 500 nm, less than about 200 nm, less than about 150 nm, less than about 100 nm, less than about 90 nm, less than about 80 nm, less than about 70 nm, less than about 60 nm, less than about 50 nm, less than about 40 nm, less than about 30
10 nm, less than about 20 nm, less than about 19 nm, less than about 18 nm, less than about 17 nm, less than about 16 nm, less than about 15 nm, less than about 14 nm, less than about 13 nm, less than about 12 nm, less than about 11 nm, less than about 10 nm, less than about 5 nm, less than about 4 nm, less than about 3 nm, less than about 2 nm, less than about 1 nm, or any value there-between or less.
- 15 3. The composition of claim 1, wherein said composition has the capacity to biodistribute in a preferential manner that targets multiple cell types involved in the process of neuroinflammation, including microglia, astrocytes, Mueller cells, macrophages, retinal pigment epithelial cells, and blood vessels.
4. The composition of claim 1, wherein said biologically active component is
20 selected from the group consisting of corticosteroids, anti-inflammatory agents, vitamins, peptides, growth factors, CNS stimulants, oligonucleotides, siRNAs, microRNAs, resolvins, neurostimulants and neuroprotectants.
5. The composition of claim 4, wherein said compound is selected from the group consisting of fluocinolone acetonide, ranibizumab, minocycline, rapamycin,
25 methyl prednisone, dexamethasone, insulin, estradiol, CNTF, vitamin A, vitamin C, vitamin E, and an oligonucleotide, or a pharmaceutically acceptable salt thereof.
6. The composition of claim 1, wherein said nanoparticle is a soft nanoparticle.
7. The composition of claim 6, wherein said nanoparticle is a dendrimer-
30 branched or star-branched polymer.
8. The composition of claim 7, wherein the dendrimer-branched polymers consists of polyamidoamine (PAMAM), priostar, polyester, polyether, polylysine, or polyethylene glycol (PEG) dendrimers, and the star-branched polymer is a PEG star.

9. The composition of claim 7, wherein said dendrimer has a diameter of 1.5 nanometers to 14.5 nanometers.

10. The composition of claim 7, wherein said dendrimer has a diameter of 2 to 10 nanometers

5 11. The composition of claim 1, wherein said drug is incorporated into the hyperbranched formulation through encapsulation, complexation, or covalent linkage.

12. The composition of claim 11, wherein said linkage comprises a spacer consisting of a peptide, glutaric acid, or PEG to link the drug and the polymer.

10 13. The composition of claim 9 wherein said drug-nanoparticle is incorporated in a larger scale entity incorporating the drug-hyperbranched polymer, wherein said larger-scale entity consists of a polymer matrix, a microparticle, a nanoparticle, a liposome, a microcapsule, a nanocapsule, injectible hydrogels, or controlled-release implants.

15 14. The composition of claim 1, wherein the nanoscale drug-hyperbranched polymer formulation is applied as a coating on an implantable device.

15. The composition of claim 1, wherein sustained release of the active compound occurs over a period of time.

16. The composition of claim 15, wherein said release occurs over a period of minutes, hours, days, months, or years.

20 17. The composition of claim 1, wherein the composition provides sustained delivery of said compound to a targeted site in a patient.

18. The composition of claim 17, wherein said targeted site is the vitreous of the eye.

25 19. The composition of claim 17, wherein said sustained delivery is over a period of hours.

20. The composition of claim 19, wherein said sustained delivery is over a period of days.

21. The composition of claim 19, wherein said sustained delivery is over a period of weeks.

30 22. The composition of claim 19, wherein said sustained delivery is over a period of months.

23. A composition comprising at least one anti-inflammatory compound conjugated to a dendrimer, wherein said composition is encapsulated in a

biodegradable particle selected from the group consisting of a PLA nanoparticle and a PGLA microparticle.

24. The composition of claim 23, wherein said dendrimer is –OH or –COOH terminated, belonging to PAMAM or Priostar family and said anti-inflammatory compound is selected from the group consisting of fluocinolone acetonide, minocycline, methyl prednisone and dexamethasone.

25. A method of treatment of a neuroinflammation-related disorder comprising administering the composition of claim 1, wherein said neuroinflammation-related disorder is a disease of the retina, central nervous system, the spinal cord, and/or the peripheral nervous system.

26. The method of claim 25, wherein for treating a disease of the retina, the composition is injected intravitreally, at an appropriate dose and interval, to treat age-related macular degeneration (ARMD)

27. The method of claim 25, wherein for treating a disease of the central nervous system, the composition is injected intrathecally, at an appropriate dose and interval, to treat cerebral palsy.

28. The method of claim 25, wherein said disease is selected from the group of eye diseases consisting of retinitis pigmentosa, age-related macular degeneration, optic neuritis, to the eye, infections, sarcoid, sickle cell disease, retinal detachment, temporal arteritis, retinal ischemia, arteriosclerotic retinopathy, hypertensive retinopathy, retinal artery blockage, retinal vein blockage, hypotension, diabetic retinopathy, and macular edema.

29. The method of claim 25, wherein said disease is selected from the group of central or peripheral nervous system diseases consisting of stroke, cerebral palsy, blunt and penetrating injuries to the brain, Alzheimer's disease, Parkinson's disease, brain or spinal cord trauma, sarcoid, sickle cell disease, AIDS dementia, age-related loss of cognitive function, memory loss, amyotrophic lateral sclerosis, seizure disorders, alcoholism, aging, and neuronal loss.

30. A method of treating progressive vision loss in a human in need thereof, said method comprising administering to the eye of said human a composition of claim 1.

31. The method of claim 30, wherein said progressive vision loss is associated with at least one condition selected from the group consisting of uveitis, age-related macular degeneration, diabetic retinopathy, and retinitis pigmentosa.


32. A method of treating ocular neuroinflammation in a human in need thereof, said method comprising administering to the eye of said human at least one composition of claim 23.

33. The method of claim 32, wherein said administering occurs once.


5 34. The method of claim 32, wherein said administering occurs two or more times over a period of days, weeks or months.

35. A medical device comprising the composition of claim 1 and instructions for administering said composition.


10 36. A medical device comprising the composition of claim 23 and instructions for administering said composition.



FIGURES 1A-1D



Figures 2A-2F



FIGURES 3A-3B




FIGURE 4




FIGURE 5

4/9




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9A





FIGURE 9B



FIGURES 10A-10F




FIGURE 11A-11D





FIGURE2 12A-12B



FIGURES 13A-13D






FIGURE 14



FIGURES 15A-15B



FIGURES 16A-16B