



US00RE47437E

(19) **United States**
 (12) **Reissued Patent**
Kawamoto et al.

(10) **Patent Number:** **US RE47,437 E**
 (45) **Date of Reissued Patent:** ***Jun. 18, 2019**

- (54) **PROLYL HYDROXYLASE INHIBITORS AND METHODS OF USE**
- (71) Applicant: **Akebia Therapeutics, Inc.**, Cambridge, MA (US)
- (72) Inventors: **Richard Masaru Kawamoto**, Divide, CO (US); **Shengde Wu**, West Chester, OH (US); **Artem G. Evdokimov**, Foristell, MO (US); **Kenneth D. Greis**, Fort Thomas, KY (US); **Angelique Sun Boyer**, West Chester, OH (US); **Namal C. Warshakoon**, Cary, NC (US)
- (73) Assignee: **Akebia Therapeutics, Inc.**, Cambridge, MA (US)
- (*) Notice: This patent is subject to a terminal disclaimer.
- (21) Appl. No.: **15/875,495**
- (22) Filed: **Jan. 19, 2018**

Related U.S. Patent Documents

Reissue of:

- (64) Patent No.: **9,598,370**
 Issued: **Mar. 21, 2017**
 Appl. No.: **14/854,080**
 Filed: **Sep. 15, 2015**

- (51) **Int. Cl.**
C07D 417/04 (2006.01)
C07D 405/10 (2006.01)
C07D 295/192 (2006.01)
C07D 213/65 (2006.01)
C07C 237/42 (2006.01)
C07C 255/57 (2006.01)
C07C 235/60 (2006.01)
C07D 213/81 (2006.01)
C07D 401/04 (2006.01)
C07D 405/04 (2006.01)
C07D 401/10 (2006.01)
A61K 31/4418 (2006.01)
A61K 31/4433 (2006.01)

- (52) **U.S. Cl.**
 CPC **A61K 31/4418** (2013.01); **A61K 31/4433** (2013.01); **C07C 235/60** (2013.01); **C07C 237/42** (2013.01); **C07C 255/57** (2013.01); **C07D 213/65** (2013.01); **C07D 213/81** (2013.01); **C07D 295/192** (2013.01); **C07D 401/04** (2013.01); **C07D 401/10** (2013.01); **C07D 405/04** (2013.01); **C07D 405/10** (2013.01); **C07D 417/04** (2013.01); **Y02A 50/401** (2018.01)

- (58) **Field of Classification Search**
 CPC **C07C 235/60**; **C07C 237/42**; **C07C 255/57**
 See application file for complete search history.

- (56) **References Cited**

U.S. PATENT DOCUMENTS

3,894,920 A 7/1975 Kondo et al.
 4,016,287 A 4/1977 Ebhardt et al.

5,397,799 A 3/1995 Kress et al.
 5,405,613 A 4/1995 Rowland et al.
 5,607,954 A 3/1997 Weidmann et al.
 5,610,172 A 3/1997 Weidmann et al.
 5,620,995 A 4/1997 Weidmann et al.
 5,620,996 A 4/1997 Weidmann et al.
 5,658,933 A 8/1997 Weidmann et al.
 5,719,164 A 2/1998 Weidmann et al.
 5,726,305 A 3/1998 Weidmann et al.
 6,020,350 A 2/2000 Weidmann et al.
 6,093,730 A 7/2000 Weidmann et al.
 6,420,427 B1 7/2002 Takahashi et al.
 6,566,088 B1 5/2003 McKnight et al.
 6,589,758 B1 7/2003 Zhu
 7,183,287 B2 2/2007 Durley
 7,588,824 B2 9/2009 Wudl et al.
 7,588,924 B2 9/2009 Evdokimov et al.
 7,811,595 B2 10/2010 Kawamoto et al.
 8,050,873 B2 11/2011 Evdokimov et al.
 8,124,582 B2 2/2012 Guenzler-Pukall et al.
 8,323,671 B2 12/2012 Wu et al.
 8,343,952 B2 1/2013 Kawamoto et al.
 8,530,404 B2 9/2013 Seeley et al.
 8,598,210 B2 * 12/2013 Kawamoto C07C 235/60
 514/338
 8,722,895 B2 * 5/2014 Kawamoto C07C 235/60
 546/298
 8,865,748 B2 10/2014 Shalwitz et al.
 8,940,773 B2 * 1/2015 Kawamoto C07C 235/60
 514/338
 9,145,366 B2 * 9/2015 Lanthier C07D 213/79
 9,598,370 B2 3/2017 Kawamoto et al.
 9,701,636 B2 7/2017 Copp et al.
 9,776,969 B2 10/2017 Lanthier et al.
 2002/0192737 A1 12/2002 Kaelin, Jr. et al.
 2003/0153503 A1 8/2003 Klaus et al.
 2003/0176317 A1 9/2003 Guenzler-Pukall et al.
 2004/0235082 A1 11/2004 Fournay et al.
 2004/0254215 A1 12/2004 Arend et al.
 2006/0276477 A1 12/2006 Klaus et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2098158 6/1993
 CA 2253282 11/1997
 EP 0650960 B1 3/1997
 EP 0650961 B1 3/1997
 EP 2044005 B1 10/2010

(Continued)

OTHER PUBLICATIONS

U.S. Appl. No. 60/816,522, filed Jun. 26, 2006, Kawamoto et al.

(Continued)

Primary Examiner — Bruce R Campell
 (74) *Attorney, Agent, or Firm* — Proskauer Rose LLP;
 Fangli Chen

- (57) **ABSTRACT**

The present disclosure relates to HIF-1 α prolyl hydroxylase inhibitors, compositions which comprise the HIF-1 α prolyl hydroxylase inhibitors described herein and to methods for controlling, inter alia, Peripheral Vascular Disease (PVD), Coronary Artery Disease (CAD), heart failure, ischemia, and anemia.

8 Claims, 2 Drawing Sheets**Specification includes a Sequence Listing.**

(56)

References Cited

U.S. PATENT DOCUMENTS

2007/0105899	A1	5/2007	Suzuki et al.	
2007/0154482	A1	7/2007	Sukhatme et al.	
2007/0213335	A1	9/2007	Fitch et al.	
2007/0299086	A1	12/2007	Kawamoto et al.	
2008/0124740	A1	5/2008	Evdokimov et al.	
2008/0213404	A1	9/2008	Johnson et al.	
2009/0082357	A1	3/2009	Fitch et al.	
2010/0331303	A1	12/2010	Kawamoto et al.	
2010/0331374	A1	12/2010	We et al.	
2012/0282627	A1	11/2012	Evdokimov et al.	
2012/0309977	A1	12/2012	Lanthier et al.	
2012/0316204	A1	12/2012	Shalwitz et al.	
2012/0329836	A1	12/2012	Marsh et al.	
2013/0203816	A1	8/2013	Kawamoto et al.	
2013/0245076	A1	9/2013	Kawamoto et al.	
2014/0045899	A1	2/2014	Kawamoto et al.	
2014/0057892	A1	2/2014	Kawamoto et al.	
2015/0119425	A1*	4/2015	Kawamoto	C07C 235/60 514/338
2015/0141467	A1*	5/2015	Copp	C07D 213/81 514/350
2015/0361043	A1*	12/2015	Lanthier	C07D 213/79 546/298
2016/0009648	A1	1/2016	Kawamoto et al.	
2016/0143891	A1	5/2016	Shalwitz et al.	
2017/0189387	A1	7/2017	Kawamoto et al.	
2017/0258773	A1	9/2017	Copp et al.	

FOREIGN PATENT DOCUMENTS

JP	H09221476	8/1997
JP	2001-48786	2/2001
WO	WO 97/41103	11/1997
WO	WO 97/44333	11/1997
WO	WO 99/48870	11/1999
WO	WO 02/074980	9/2002
WO	WO 02/074981	9/2002
WO	WO 02/083688	10/2002
WO	WO 03/028663	4/2003
WO	WO 03/032972	4/2003
WO	WO 03/049686	6/2003
WO	WO 03/053997	7/2003
WO	WO 2004/035812	4/2004
WO	WO 2004/048383	6/2004
WO	WO 2004/108121	12/2004
WO	WO 2005/007192	1/2005
WO	WO 2005/115984	12/2005
WO	WO 2005/118836	12/2005
WO	WO 2006/019831	2/2006
WO	WO 2006/030977	3/2006
WO	WO 2006/114213	11/2006
WO	WO 2007/038571	4/2007
WO	WO 2007/047194	4/2007
WO	WO 2007/070359	6/2007
WO	WO 2007/082899	7/2007
WO	WO 2007/084667	7/2007
WO	WO 2007/088571	8/2007
WO	WO 2007/103905	9/2007
WO	WO 2007/136990	11/2007
WO	WO 2007/150011	12/2007
WO	WO 2008/002576	1/2008
WO	WO 2008/089051	7/2008
WO	WO 2008/089052	7/2008
WO	WO 2008/130508	10/2008
WO	WO 2008/130527	10/2008
WO	WO 2008/137060	11/2008
WO	WO 2008/144266	11/2008
WO	WO 2009/019656	2/2009
WO	WO 2009/037570	3/2009
WO	WO 2009/039321	3/2009
WO	WO 2009/039323	3/2009
WO	WO 2007/038571	4/2009
WO	WO 2009/043093	4/2009
WO	WO 2009/049112	4/2009

WO	WO 2009/067790	4/2009
WO	WO 2009/070644	6/2009
WO	WO 2009/073497	6/2009
WO	WO 2009/073669	6/2009
WO	WO 2009/086044	7/2009
WO	WO 2009/086592	7/2009
WO	WO 2009/089547	7/2009
WO	WO 2016/153996	9/2016
WO	WO 2016/161094	10/2016

OTHER PUBLICATIONS

- U.S. Appl. No. 15/559,200, filed Sep. 18, 2017, Hanselmann et al.
 U.S. Appl. No. 15/563,169, filed Sep. 29, 2017, Smith et al.
 Alfred Burger, "Isosterism and bioisosterism in drug design," *Progress in Drug Research* (Ed. J.A. Salmon et al), pp. 287-371, 1991.
 Tim Cheeseright, "The identification of bioisosteres as drug development candidates," *Innovations in Pharmaceutical Technology*, 22-26, 2009.
 Stuart Enoch et al., "ABC of wound healing: non-surgical and drug treatments" *BMJ*: 332:900-903, 2006.
 Y.N. Gavhane et al., "Solid tumors: facts, challenges and solutions," *International Journal of Pharma Sciences and Research*, 2(1):1-12, 2011.
 Matthew J. Gaunt et al., "Rational design of benzyl-type protecting groups allows sequential deprotection of hydroxyl groups by catalytic hydrogenolysis," *J. Org. Chem.*, 63:4172-4173, 1998.
 Ian R. Hardcastle et al., "Discovery of potent chromen-4-one inhibitors of the DNA-dependent protein kinase (DNA-PK) using a small-molecule library approach," *J. Med. Chem.*, 48:7829-7846, 2005.
 Hippuric acid sodium salt. Science Lab.com: Chemicals & Laboratory Equipment. ("http://web.archive.org/web/20041107121553/http://www.sciencelab.com/page-e/SPVAR/10415/SLH2620", accessed Mar. 11, 2010).
 H. Hoeksema et al., "Structure of Rubradirin," *Journal of the American Chemical Society*, 104:5173-5181, 1982.
 Ingersoll et al. Organic Syntheses, CV 2, 328, "Hippuric acid". (<http://web.archive.org/web/20020724135719/http://www.orgsyn.org/orgsyn/o-rgsyn/prepContent.asp?prep=cv2p0328>, accessed Mar. 11, 2010).
 Qingdong Ke et al., "Hypoxia-inducible factor-1 (HIF-1)," *Molecular Pharmacology*, 70(5):1469-1480, 2006.
 So Yeon Kim et al., "Recent advances in developing inhibitors for hypoxia-inducible factor prolyl hydroxylases and their therapeutic implications," *Molecules*, 20(11), 20551-20568, 2015.
 Lida Moreira Lima et al., "Bioisosterism: a useful strategy for molecular modification and drug design," *Current Medicinal Chemistry*, 12:23-49, 2005.
 Pablo E. Pergola et al., "Vadadustat, a novel oral HIF stabilizer, provides effective anemia treatment in nondialysis-dependent chronic kidney disease," *Kidney International*, 90(5):1115-1122, 2016.
 Sirivipa Piyamongkol et al., "Amido-3-hydroxypyridin-4-ones as iron(III) ligands," *Chemistry, A European Journal*, 16:6374-6381, 2010.
 Michael Schelhaas et al., "Protecting group strategies in organic synthesis," *Angew. Chem. Int. Ed. Engl.*, 35:2056-2083, 1996.
 Gregg L. Semenza, "HIF-1 and human disease: one highly involved factor." *Genes & Dev.*, 14:1983-1991, 2000.
 Ambreena Siddiq et al., "Hypoxia-inducible factor prolyl 4-hydroxylase inhibition: a target for neuroprotection in the central nervous system," *Journal of Biological Chemistry*, 280:41732-41743, 2005.
 Silverman; *The Organic Chemistry of Drug Design and Drug Action*, 2nd Edition, 2004, Elsevier, Section 2.2, E.4, pp. 29-34.
 Carrie Hayes Sutter et al., "Hypoxia-inducible factor α protein expression is controlled by oxygen-regulated ubiquitination that is disrupted by deletions and missense mutations," *PNAS*, 97(9):4748-4753, 2000.
 C.W. Thornber, "Isosterism and molecular modification in drug design," *Chem. Soc. Rev.*, 8:563-580, 1979.
 European Patent No. 2044005, Opposition Filed, Jul. 20, 2011.

(56)

References Cited

OTHER PUBLICATIONS

- European Patent No. 2044005, Response to Opposition, Dec. 28, 2011.
- European Patent No. 2044005, Summons to Attend Oral Proceedings and Preliminary Opinion of the Opposition Division, Oct. 31, 2012.
- European Patent No. 2044005, Decision of the Opposition Division, Instruction, and Grounds for the Decision, May 3, 2013.
- European Patent No. 2044005, Patent Owner Statement of Grounds of Appeal, Sep. 13, 2013.
- European Patent No. 2044005, Opponent Statement of Grounds of Appeal, Sep. 13, 2013.
- European Patent No. 2044005, Opponent Reply to Appeal, Jan. 29, 2014.
- European Patent No. 2044005, Patent Owner Reply to Appeal, Jan. 29, 2014.
- European Patent No. 2044005, Patent Owner Response to the Opponent's Communication, Aug. 28, 2014.
- European Patent No. 2044005, Communication of the Board of Appeal, Aug. 29, 2017.
- European Patent No. 2044005, Response of Opponent to Communication of the Board of Appeal, Nov. 8, 2017.
- Withdrawal of an appeal by proprietor in the Opposition against EP2044005 (07835890.0) submitted to the European Patent Office on Feb. 27, 2018.
- Decision of the Board of Appeal of the European Patent Office in the Opposition against EP2044005 (07835890.0) issued on Mar. 6, 2018.
- Termination of Opposition Proceedings of Patent No. EP2044005 (07835890.0) issued on Apr. 18, 2018.
- "Notice of Opposition u/s 25(2)" filed on Sep. 25, 2018 by Dr. Reddy's Laboratories Limited against Indian Patent No. 287720.
- Hirsila et al., "Characterization of the Human Prolyl 4-Hydroxylases That Modify the Hypoxia-inducible Factor", *The Journal of Biological Chemistry*, 278(33): 30772-30780 (Aug. 15, 2003).
- Alesso et al., "Improving resins for solid phase synthesis: incorporation of 1-[2-(2-methoxyethoxy) ethoxy]4-vinyl-benzene" *Tetrahedron*: 59,7163-7169 (2003).
- Altschul et al., "Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search Programs," *Nucleic Acids Res.*, 25(27):3389-3402 (1997).
- Anderson et al., "Antileukemic Activity of Derivatives of 1,2-Dimethyl-3,4-bis(hydroxymethyl)-5-phenylpyrrole Bis(N-methylcarbamate)" *J. Med. Chem.*: vol. 22(8), 977-980 (1979).
- Annex et al., "Growth Factor-Induced Therapeutic Angiogenesis in the Heart: Protein Therapy," *Cardiovascular Research*, 65(3):649-655 (2005).
- Ardelt et al., "Estradiol Regulates Angiotensinogen-1 mRNA Expression Through Estrogen Receptor- α in a Rodent Experimental Stroke Model," *Stroke*, 36:337-341 (2005).
- Auerbach et al., "Angiogenesis Assays: A Critical Overview," *Clinical Chemistry*, 49:32-40 (2003).
- Barany et al., "Solid-phase Peptide Synthesis: A Silver Anniversary Report," *Int. J. Peptide Protein Res.*, 30(6):705-739 (1987).
- Bartlett et al., "Molecular Recognition in Chemical and Biological Problems," *Special Pub.*, Royal Chem. Soc., 78, 182-196 *Caveat: A Program to Facilitate the Structure-derived Design of Biologically Active Molecules* (Apr. 1989).
- Böhm, "The Computer Program LUDI: A New Method for the Novo Design of Enzyme Inhibitors," *J. Computer-Aided Molecular Design*, 6:61-78 (1992).
- Branden et al., "Introduction to Protein Structure Second Edition," Garland Publishing, Inc., New York, 1999, pp. 374-375.
- Bussolino, "Molecular Mechanisms of Blood Vessel Formation," *Trends Biochem. Sci.*, 22(7):251-256 (1997).
- Cunliffe et al., "Novel Inhibitors of Prolyl4-Hydroxylase 3 Inhibition by the Substrate Analogue N-Oxaloglycine and Its Derivatives," *J. Med. Chem.* 35:2652-2658 (1992).
- Elson et al., "Induction of Hypervascularity Without Leakage or Inflammation in Transgenic Mice Overexpressing Hypoxia-Inducible Factor- α ," *Genes & Dev.*, 15:2520-2532 (2001).
- European Patent Application No. 11 000 872.9; European Search Report, Apr. 28, 2011.
- European Patent Office, Interlocutory Decision in Opposition Proceedings, mailed May 3, 2013, 76 pages.
- European Patent Office, Minutes of the Oral Proceedings Before the Opposition Division, mailed May 3, 2013, 6 pages.
- Flower, "Modelling G-protein-coupled receptors for drug design," *Biochimica et Biophysica Acta*, 1422:207-234 (1999).
- Folkman et al., "Tumor Angiogenesis," *The Molecular Basis of Cancer*, Mendelsohn et al., eds., W. B. Saunders, Chapter 10, pp. 206-232 (1995).
- Franklin et al., "Approaches to the Design of Anti-Fibrotic Drugs," *Biochem. Soc. Trans.*, 19(4):812-5 (Nov. 1991).
- Goodford, "A Computational Procedure for Determining Energetically Favorable Binding Sites on Biologically Important Macromolecules," *J. Med. Chem.*, 28(7):849-857 (1985).
- Goodsell et al., "Automated Docking of Substrates to Proteins by Simulated Annealing," *Proteins: Structure, Function, and Genetics*, 8:195-202 (1990).
- International Union of Pure and Applied Chemistry; Glossary of Class Names of Organic Compounds and Reactive Intermediates Based on Structure; *Pure & D Appl. Chem.*, vol. 67, Nos. 8/9, pp. 1307-1375, (1995).
- Ivan et al., "HIF α Targeted for VHL-Mediated Destruction by Proline Hydroxylation: Implications for O₂ Sensing," *Science* 292, 464-468 (2001).
- Ivan et al., "Biochemical purification and pharmacological inhibition of a mammalian prolyl hydroxylase acting on hypoxia-inducible factor," *Proceedings of the National Academy of Science*; 99(21) 13459-13464 (2002).
- Jaakkola et al., "Targeting of HIF- α to the von Hippel-Lindau Ubiquitylation Complex by O₂ Regulated Prolyl Hydroxylation," *Science* 292, 468-472 (2001).
- Jones et al., "Molecular Recognition of Receptor Sites Using a Genetic Algorithm with a Description of Desolvation," *J. Mol. Biol.*, 245:43-53 (1995).
- Kaelin, "Proline Hydroxylation and Gene Expression," *Annu. Rev. Biochem.*, 74:115-125 (2005).
- Kawashima et al., "Suppressive effect of quinolinic acid and hippuric acid on bone marrow erythroid growth and lymphocyte blast formation in uremia," *Advances in Experimental Medicine and Biology*, (1987), vol. 223, p. 69-72.
- Krantz, "Erythropoietin," *Blood*, 77:419-434 (1991).
- Kuntz et al., "A Geometric Approach to Macromolecule-Ligand Interactions," *J. Mol. Biol.*, 161:269-288 (1982).
- Lee et al., "Structure of Human FIH-1 Reveals a Unique Active Site Pocket and Interaction Sites for HIF-1 and von Hippel Lindau," *JBC*, 278:7558-7563 (2003).
- Li et al., "PR39, A Peptide Regulator of Angiogenesis," *Nat Med.*, 6(1):49-55 (2000).
- Mancini et al., "Effect of Erythropoietin on Exercise Capacity in Patients with Moderate to Severe Chronic Heart Failure," *Circulation*, 107:294-299 (2003).
- McDonough et al., "Cellular Oxygen Sensing: Crystal Structure of Hypoxia-Inducible Factor Prolyl Hydroxylase (PHD2)," *PNAS*, 103(26):9814-9819 (2006).
- Miranker et al., "Functionality Maps of Binding Sites: A Multiple Copy Simultaneous Search Method," *Proteins: Structure, Function and Genetics*, 11:29-34 (1991).
- Nguyen et al., "Cellular Interactions in Vascular Growth and Differentiation," *Int. Review of Cytology*, 204:1-48 (2001).
- Nishibata et al., "Automatic Creation of Drug Candidate Structures Based on Receptor Structure. Starting Point for Artificial Lead Generation," *Tetrahedron*, 47(43):8985-8990 (1991).
- O'Reilly et al., "Angiostatin: A Novel Angiogenesis Inhibitor that Mediates the Suppression of Metastases by a Lewis Lung Carcinoma," *Cell*, 79:315-328 (1994).
- O'Reilly et al., "Endostatin: An Endogenous Inhibitor of Angiogenesis and Tumor Growth," *Cell*, 88:277-285 (1997).

(56)

References Cited

OTHER PUBLICATIONS

Peyssonaux et al., "HIF-1 α Expression Regulates the Bactericidal Capacity of Phagocytes," *J. Clinical Invest.*, 115(7):1806-1815 (2005).

Qian, Oral abstract FR-OR011, *J. Am. Soc. Nephrol.*, 24:38A (2013).

Schoneberg et al., "Structural Basis of G Protein-Coupled Receptor Function," *Molecular and Cellular Endocrinology*, 151:181-193 (1999).

Semenza et al., "Transcriptional Regulation of Genes Encoding Glycolytic Enzymes by Hypoxia-Inducible Factor 1," *J. Biol. Chem.*, 269:23757-23763 (1994).

Semenza, "Regulation of Erythropoietin Production: New Insights into Molecular Mechanisms of Oxygen Homeostasis," *Hematol. Oncol. Clin. North Am.*, 8:863-884 (1994).

Semenza, "Signal Transduction to Hypoxia-inducible Factor 1," *Biochem. Pharmacol.*, 64:993-998 (2002).

Sexton, "Recent advances in our understanding of peptide hormone receptors and RAMPS," *Current Opinion in Drug Discovery and Development*, 2(5):440-448 (1999).

Sheehan, "3-Hydroxypicolinic Acid and Some of its Derivatives," *J. Organic Chemistry* 31(3):636-638 (1996).

Teicher et al., "Potentiation of Cytotoxic Cancer Therapies by TNP-470 Alone and with Other Anti-Angiogenic Agents," *Int. J. Cancer*, 57:920-925 (1994).

Tzschucke et al., "Fluorous-Silica-Supported Perfluoro-Tagged Palladium Complexes Catalyze Suzuki Coupling in Water" *Helvetica Chimica Acta*; vol. 87 2882-2889 (2004).

Vickerstaffe et al., "Fully Automated Polymer-Assisted Synthesis of 1,5-Biaryl Pyrazoles," *J. Comb. Chem.*, 2004,6,332-33.

Vincent et al., "Angiogenesis is Induced in a Rabbit Model of Hindlimb Ischemia by Naked DNA Encoding an HIF-1 α /VP16 Hybrid Transcription Factor," *Circulation*, 102:2255-2261 (2000).

Warnecke et al., "Activation of the Hypoxia-Inducible Factor Pathway and Stimulation of Angiogenesis by Application of Prolyl Hydroxylase Inhibitors," *FASEB Journal*, 17: 1186-1188 (2003).

Warshakoon et al., "Design and synthesis of substituted pyridine derivatives as HIF-1 α prolyl hydroxylase inhibitors," *Bioorganic & Medicinal Chemistry Letters*, 16 (2006) 5616-5620.

Wax et al., "SM-20 is a Novel 20-kd Protein Whose Expression in the Arterial Wall is Restricted to Smooth Muscle," *Lab. Invest.*, 74(4):797-808 (1996).

Weidner et al., "Tumor Angiogenesis and Metastasis--Correlation in Invasive Breast Carcinoma," *New Eng. J. Med.*, 324(1):1-8 (1991).

Wright et al., "Activation of the Prolyl Hydroxylase Oxygen-Sensor Results in Induction of GLUT1, Heme Oxygenase-1, and Nitric-Oxide Synthase Proteins and Confers Protection from Metabolic Inhibition to Cardiomyocytes," *J. Bio. Chem.*, 278(22):20235-20239 (2003).

Office action mailed Oct. 7, 2013 in connection with U.S. Appl. No. 13/489,400.

Office action mailed Jan. 30, 2014 in connection with U.S. Appl. No. 13/489,400.

Office action mailed Dec. 12, 2013 in connection with U.S. Appl. No. 14/071,088.

"Standards of Medical Care in Diabetes-2006," *Diabetes Care*, 29:s4-s42 (2006).

Qian et al., "A Randomized, Double-Blind, Placebo Controlled Trial of FG-4592 for Correction of Anemia in Subjects with Chronic Kidney Disease in China," Oral Abstract FR-OR011, *J. Am. Soc. Nephrol.* 24:38A (2013).


Langsetmo, "Inhibition of HIF-Prolyl Hydroxylases with FG-4539 Is Neuroprotective in a Mouse Model of Permanent Focal Ischemia," International Stroke Conference, Kissimmee Florida, Presentation No. 427 (2006).

Rahtu-Korpela, "HIF Prolyl 4-Hydroxylase-2 Inhibition Improves Glucose and Lipid Metabolism and Protects Against Obesity and Metabolic Dysfunction," *Diabetes* 63:3324-3333 (2014).

Catrina et al., 2004, "Hyperglycemia Regulates Hypoxia-Inducible Factor-1 Protein Stability and Function," *Diabetes* 53:3226-3232.

* cited by examiner


Figure 1



Immunoblot analysis of nuclear extracts demonstrating stabilization of HIF-1 α in mouse liver by {{5-(3-chlorophenyl)-3-hydroxypyridine-2-carbonyl}amino}-acetic acid

Figure 2

Elevated erythropoietin levels in mouse serum following oral dosing of a compound, {[5-(3-chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid as compared with a vehicle.



**PROLYL HYDROXYLASE INHIBITORS AND
METHODS OF USE**

Matter enclosed in heavy brackets [] appears in the original patent but forms no part of this reissue specification; matter printed in italics indicates the additions made by reissue; a claim printed with strikethrough indicates that the claim was canceled, disclaimed, or held invalid by a prior post-patent action or proceeding.

PRIORITY

This application is a *Reissue of U.S. Pat. No. 9,598,370, which issued on Mar. 21, 2017, from U.S. application Ser. No. 14/854,080, filed Sep. 15, 2015, which is a Continuation of U.S. application Ser. No. 14/568,200, filed Dec. 12, 2014, which is a Continuation of U.S. application Ser. No. 14/062,011, filed Oct. 24, 2013, now U.S. Pat. No. 8,940,773, which is a Divisional application of U.S. application Ser. No. 13/681,876, filed on Nov. 20, 2012, now U.S. Pat. No. 8,598,210, which is a Continuation of U.S. application Ser. No. 12/860,073, filed on Aug. 20, 2010, now U.S. Pat. No. 8,343,952, which is a Continuation application of U.S. application Ser. No. 11/821,936, filed Jun. 26, 2007, now U.S. Pat. No. 7,811,595, which claims the benefit of Provisional Application Ser. No. 60/816,522 filed on Jun. 26, 2006, and the entire disclosures of Provisional Application Ser. No. 60/816,522 and U.S. application Ser. No. 11/821,936 are incorporated herein by reference in its entirety.*

FIELD OF THE DISCLOSURE

The present disclosure relates, in some aspects, to HIF-1 α prolyl hydroxylase inhibitor compounds and pharmaceutically acceptable salts thereof, compositions comprising the HIF-1 α prolyl hydroxylase inhibitor compounds, and to methods for treating or controlling, inter alia, Peripheral Vascular Disease (PVD), Coronary Artery Disease (CAD), heart failure, ischemia, and anemia.

BACKGROUND OF THE DISCLOSURE

HIF-1 α under normal healthy conditions wherein the cells have a sufficient supply of oxygen is readily converted to a degraded form by one of several prolyl hydroxylase enzymes, inter alia, EGLIN. When cells undergo hypoxia, this enzymatic transformation is slow or entirely stopped and HIF-1 α begins to build up in the cell. When this build up of HIF-1 α occurs, this protein combines with another factor, HIF-1 β which together form an active transcription factor complex. This transcription factor then activates several biological pathways which are present as a response to and a means for alleviating the body's state of hypoxia. These responses include, inter alia, angiogenic, erythropoietic (EPO), glucose metabolism, and matrix alteration responses.

In patients where there is a need for stimulating one or more of these responses, for example, in patients in need of increased tissue oxygen due to peripheral vascular disease (PVD), inhibiting the EGLIN enzyme will stimulate the body's own angiogenic response without the consequences of oxygen deficiency. In addition, in diseases of ischemia, inter alia, CAD and anemia, stimulation of angiogenic, erythropoietic, and metabolic adaptation would be expected to provide therapeutic benefits.

Therefore there continues to be a long felt need for compounds that inhibit prolyl hydroxylase enzymes and thereby regulate the concentration of HIF-1 α in cells so as to induce angiogenic or erythropoietic responses and therefore treat diseases related to hypoxia or anemia.

SUMMARY OF THE DISCLOSURE

The substituted aryl or heteroaryl amide compounds of the present disclosure are a new class of compounds that can inhibit HIF-1 α prolyl hydroxylase, thus resulting in improvement in blood flow, oxygen delivery and energy utilization in ischemic tissues, or upregulate the production of erythropoietin so as to treat anemia.

Disclosed herein are compounds and pharmaceutically acceptable salts thereof, and/or pharmaceutical compositions thereof comprising:

- a) an effective amount of one or more compounds according to the present disclosure; and
- b) an excipient.

The present disclosures also relate to methods for controlling, inter alia, Peripheral Vascular Disease (PVD), Coronary Artery Disease (CAD), heart failure, ischemia, and/or anemia.

The present disclosures also relate to methods for regulating blood flow, oxygen delivery and/or energy utilization in ischemic tissues, wherein the methods can comprise administering to a human an effective amount of one or more compounds or pharmaceutically acceptable salts disclosed herein.

These and other objects, features, and advantages will become apparent to those of ordinary skill in the art from a reading of the following detailed description and the appended claims. All documents cited herein are in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present disclosure.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. Immunoblot analysis of nuclear extracts demonstrating stabilization of HIF-1 α in mouse liver by {[5-(3-chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid

FIG. 2. An example of how the erythropoietin level is elevated versus vehicle in mouse serum after oral dosing of {[5-(3-chlorophenyl)-3-hydroxypyridine-2-carbonyl]-amino}-acetic acid.

DETAILED DESCRIPTION OF THE
DISCLOSURE

In this specification and in the claims that follow, reference will be made to a number of terms, which shall be defined to have the following meanings:

By "pharmaceutically acceptable" is meant a material that is not biologically or otherwise undesirable, i.e., the material can be administered to an individual along with the relevant active compound without causing clinically unacceptable biological effects or interacting in a deleterious manner with any of the other components of the pharmaceutical composition in which it is contained.

Throughout the description and claims of this specification the word "comprise" and other forms of the word, such as "comprising" and "comprises," means including but not limited to, and is not intended to exclude, for example, other additives, components, integers, or steps.

As used in the description and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a composition" includes mixtures of two or more such compositions.

"Optional" or "optionally" means that the subsequently described event or circumstance can or cannot occur, and that the description includes instances where the event or circumstance occurs and instances where it does not.

Ranges can be expressed herein as from "about" one particular value, and/or to "about" another particular value. When such a range is expressed, another aspect includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations, by use of the antecedent "about," it will be understood that the particular value forms another aspect. It will be further understood that the endpoints of each of the ranges are significant both in relation to the other endpoint, and independently of the other endpoint. It is also understood that there are a number of values disclosed herein, and that each value is also herein disclosed as "about" that particular value in addition to the value itself. For example, if the value "10" is disclosed, then "about 10" is also disclosed. It is also understood that when a value is disclosed, then "less than or equal to" the value, "greater than or equal to the value," and possible ranges between values are also disclosed, as appropriately understood by the skilled artisan. For example, if the value "10" is disclosed, then "less than or equal to 10" as well as "greater than or equal to 10" is also disclosed. It is also understood that throughout the application data are provided in a number of different formats and that this data represent endpoints and starting points and ranges for any combination of the data points. For example, if a particular data point "10" and a particular data point "15" are disclosed, it is understood that greater than, greater than or equal to, less than, less than or equal to, and equal to 10 and 15 are considered disclosed as well as between 10 and 15. It is also understood that each unit between two particular units are also disclosed. For example, if 10 and 15 are disclosed, then 11, 12, 13, and 14 are also disclosed.

The term "organic unit" as described herein refers to groups or moieties that comprise one or more carbon atoms and which form a portion of one of the compounds or pharmaceutically acceptable salts thereof. For example, many of the substituent units referred to elsewhere herein are organic units. In order to effectively function in the context of their presence in the compounds and/or salts disclosed herein, the organic units should often have variable ranges of restricted size and/or molecular weight, so as to provide desired binding to the target enzymes, solubility, bioabsorption characteristics. For example, organic unit can have, for example, 1-26 carbon atoms, 1-18 carbon atoms, 1-12 carbon atoms, 1-8 carbon atoms, or 1-4 carbon atoms. Organic units often have hydrogen bound to at least some of the carbon atoms of the organic units, and can optionally contain the common heteroatoms found in substituted organic compounds, such as oxygen, nitrogen, sulfur, and the like, or inorganic atoms such as halogens, phosphorus, and the like. One example, of an organic radical that comprises no inorganic atoms is a 5,6,7,8-tetrahydro-2-naphthyl radical. In some embodiments, an organic radical can contain 1-10 inorganic heteroatoms bound thereto or therein, including halogens, oxygen, sulfur, nitrogen, phosphorus, and the like. Examples of organic radicals include but are not limited to an alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, mono-substituted amino, di-substituted amino, acyloxy, cyano, carboxy, carboalkoxy, alkyl-

carboxamido, substituted alkylcarboxamido, dialkylcarboxamido, substituted dialkylcarboxamido, alkylsulfonfyl, alkylsulfanyl, thioalkyl, thiohaloalkyl, alkoxy, substituted alkoxy, haloalkyl, haloalkoxy, aryl, substituted aryl, heteroaryl, heterocyclic, or substituted heterocyclic radicals, wherein the terms are defined elsewhere herein. A few non-limiting examples of organic radicals that include heteroatoms include alkoxy radicals, trifluoromethoxy radicals, acetoxy radicals, dimethylamino radicals and the like.

Substituted and unsubstituted linear, branched, or cyclic alkyl units include the following non-limiting examples: methyl (C₁), ethyl (C₂), n-propyl (C₃), iso-propyl (C₃), cyclopropyl (C₃), n-butyl (C₄), sec-butyl (C₄), iso-butyl (C₄), tert-butyl (C₄), cyclobutyl (C₄), cyclopentyl (C₅), cyclohexyl (C₆), and the like; whereas substituted linear, branched, or cyclic alkyl, non-limiting examples of which includes, hydroxymethyl (C₁), chloromethyl (C₁), trifluoromethyl (C₁), aminomethyl (C₁), 1-chloroethyl (C₂), 2-hydroxyethyl (C₂), 1,2-difluoroethyl (C₂), 2,2,2-trifluoroethyl (C₃), 3-carboxypropyl (C₃), 2,3-dihydroxycyclobutyl (C₄), and the like.

Substituted and unsubstituted linear, branched, or cyclic alkenyl include, ethenyl (C₂), 3-propenyl (C₃), 1-propenyl (also 2-methylethenyl) (C₃), isopropenyl (also 2-methylethen-2-yl) (C₃), buten-4-yl (C₄), and the like; substituted linear or branched alkenyl, non-limiting examples of which include, 2-chloroethenyl (also 2-chlorovinyl) (C₂), 4-hydroxybuten-1-yl (C₄), 7-hydroxy-7-methyloct-4-en-2-yl (C₉), 7-hydroxy-7-methyloct-3,5-dien-2-yl (C₉), and the like.

Substituted and unsubstituted linear or branched alkynyl include, ethynyl (C₂), prop-2-ynyl (also propargyl) (C₃), propyn-1-yl (C₃), and 2-methyl-hex-4-yn-1-yl (C₇); substituted linear or branched alkynyl, non-limiting examples of which include, 5-hydroxy-5-methylhex-3-ynyl (C₇), 6-hydroxy-6-methylhept-3-yn-2-yl (C₈), 5-hydroxy-5-ethylhept-3-ynyl (C₉), and the like.

Substituted and unsubstituted "alkoxy" are used herein denotes a unit having the general formula —OR¹⁰⁰ wherein R¹⁰⁰ is an alkyl, alkylenyl, or alkynyl unit as defined herein above, for example, methoxy, methoxymethyl, methoxymethyl.

Substituted and unsubstituted "haloalkyl" are used herein denotes an alkyl unit having a hydrogen atom substituted by one or more halogen atoms, for example, trifluoromethyl, 1,2-dichloroethyl, and 3,3,3-trifluoropropyl.


The term "aryl" as used herein denotes cyclic organic units that comprise at least one benzene ring having a conjugated and aromatic six-membered ring, non-limiting examples of which include phenyl (C₆), naphthylen-1-yl (C₁₀), naphthylen-2-yl (C₁₀). Aryl rings can have one or more hydrogen atoms substituted by another organic or inorganic radical. Non-limiting examples of substituted aryl rings include: 4-fluorophenyl (C₆), 2-hydroxyphenyl (C₆), 3-methylphenyl (C₆), 2-amino-4-fluorophenyl (C₆), 2-(N,N-diethylamino)phenyl (C₆), 2-cyanophenyl (C₆), 2,6-di-tert-butylphenyl (C₆), 3-methoxyphenyl (C₆), 8-hydroxynaphthylen-2-yl (C₁₀), 4,5-dimethoxynaphthylen-1-yl (C₁₀), and 6-cyanonaphthylen-1-yl (C₁₀).

The term "heteroaryl" denotes an organic unit comprising a five or six membered conjugated and aromatic ring wherein at least one of the ring atoms is a heteroatom selected from nitrogen, oxygen, or sulfur. The heteroaryl rings can comprise a single ring, for example, a ring having 5 or 6 atoms wherein at least one ring atom is a heteroatom not limited to nitrogen, oxygen, or sulfur, such as a pyridine ring, a furan ring, or thiofuran ring. A "heteroaryl" can also


5

be a fused multicyclic and heteroaromatic ring system having wherein at least one of the rings is an aromatic ring and at least one atom of the aromatic ring is a heteroatom including nitrogen, oxygen, or sulfur

The following are non-limiting examples of heteroaryl rings according to the present disclosure:




The term "heterocyclic" denotes a ring system having from 3 to 10 atoms wherein at least one of the ring atoms is a heteroatom not limited to nitrogen, oxygen, or sulfur. The rings can be single rings, fused rings, or bicyclic rings. Non-limiting examples of heterocyclic rings include:



All of the aforementioned heteroaryl or heterocyclic rings can be optionally substituted with one or more substitutes for hydrogen as described herein further.

Throughout the description of the present disclosure the terms having the spelling "thiophene-2-yl and thiophene-3-yl" are used to describe the heteroaryl units having the respective formulae:



whereas in naming the compounds of the present disclosure, the chemical nomenclature for these moieties are typically spelled "thiophen-2-yl and thiophen-3-yl" respectively. Herein the terms "thiophene-2-yl and thiophene-3-yl" are used when describing these rings as units or moieties which make up the compounds of the present disclosure solely to make it unambiguous to the artisan of ordinary skill which rings are referred to herein.

The following are non-limiting examples of units which can substitute for hydrogen atoms on a hydrocarbyl or other unit:

6

- i) linear, branched, or cyclic alkyl, alkenyl, and alkynyl; for example, methyl (C₁), ethyl (C₂), n-propyl (C₃), iso-propyl (C₃), cyclopropyl (C₃), propylene-2-yl (C₃), propargyl (C₃), n-butyl (C₄), iso-butyl (C₄), sec-butyl (C₄), tert-butyl (C₄), cyclobutyl (C₄), n-pentyl (C₅), cyclopentyl (C₅), n-hexyl (C₆), and cyclohexyl (C₆);
- ii) substituted or unsubstituted aryl; for example, phenyl, 2-fluorophenyl, 3-chlorophenyl, 4-methylphenyl, 2-aminophenyl, 3-hydroxyphenyl, 4-trifluoromethylphenyl, and biphenyl-4-yl;
- iii) substituted or unsubstituted heterocyclic; examples of which are provided herein below;
- iv) substituted or unsubstituted heteroaryl; examples of which are provided herein below;
- v) $-(CR^{12a}R^{12b})_qOR^{11}$; for example, $-OH$, $-CH_2OH$, $-OCH_3$, $-CH_2OCH_3$, $-OCH_2CH_3$, $-CH_2OCH_2CH_3$, $-OCH_2CH_2CH_3$, and $-CH_2OCH_2CH_2CH_3$;
- vi) $-(CR^{12a}R^{12b})_qC(O)R^{11}$; for example, $-COCH_3$, $-CH_2COCH_3$, $-OCH_2CH_3$, $-CH_2COCH_2CH_3$, $-COCH_2CH_2CH_3$, and $-CH_2COCH_2CH_2CH_3$;
- vii) $-(CR^{12a}R^{12b})_qC(O)OR^{11}$; for example, $-CO_2CH_3$, $-CH_2CO_2CH_3$, $-CO_2CH_2CH_3$, $-CH_2CO_2CH_2CH_3$, $-CO_2CH_2CH_2CH_3$, and $-CH_2CO_2CH_2CH_2CH_3$;
- viii) $-(CR^{12a}R^{12b})_qC(O)N(R^{11})_2$; for example, $-CONH_2$, $-CH_2CONH_2$, $-CONHCH_3$, $-CH_2CONHCH_3$, $-CON(CH_3)_2$, and $-CH_2CON(CH_3)_2$;
- ix) $-(CR^{12a}R^{12b})_qOC(O)N(R^{11})_2$; for example, $-OC(O)NH_2$, $-CH_2OC(O)NH_2$, $-OC(O)NHCH_3$, $-CH_2OC(O)NHCH_3$, $-OC(O)N(CH_3)_2$, and $-CH_2OC(O)N(CH_3)_2$;
- x) $-(CR^{12a}R^{12b})_qN(R^{11})_2$; for example, $-NH_2$, $-CH_2NH_2$, $-NHCH_3$, $-N(CH_3)_2$, $-NH(CH_2CH_3)$, $-CH_2NHCH_3$, $-CH_2N(CH_3)_2$, and $-CH_2NH(CH_2CH_3)$;
- xi) halogen: $-F$, $-Cl$, $-Br$, and $-I$;
- xii) $-CH_mX_n$; wherein X is halogen, m is from 0 to 2, m+n=3; for example, $-CH_2F$, $-CHF_2$, $-CF_3$, $-CCl_3$, or $-CBr_3$;
- xiii) $-(CR^{12a}R^{12b})_qCN$; for example; $-CN$, $-CH_2CN$, and $-CH_2CH_2CN$;
- xiv) $-(CR^{12a}R^{12b})_qNO_2$; for example; $-NO_2$, $-CH_2NO_2$, and $-CH_2CH_2NO_2$;
- xv) $-(CR^{12a}R^{12b})_qSO_2R^{11}$; for example, $-SO_2H$, $-CH_2SO_2H$, $-SO_2CH_3$, $-CH_2SO_2CH_3$, $-SO_2C_6H_5$, and $-CH_2SO_2C_6H_5$; and
- xvi) $-(CR^{12a}R^{12b})_qSO_3R^{11}$; for example, $-SO_3H$, $-CH_2SO_3H$, $-SO_3CH_3$, $-CH_2SO_3CH_3$, $-SO_3C_6H_5$, and $-CH_2SO_3C_6H_5$;
- xvii) hydroxyl groups or thiol groups,
- xviii) amino groups, monosubstituted amino, or disubstituted amino,

wherein each R¹¹ is independently hydrogen, substituted or unsubstituted C₁-C₄ linear, branched, or cyclic alkyl; or two R¹¹ units can be taken together to form a ring comprising 3-7 atoms; R^{12a} and R^{12b} are each independently hydrogen or C₁-C₄ linear or branched alkyl; the index q is from 0 to 4.

The compounds and compositions recited herein can have a number of utilities, and address several unmet medical needs, inter alia;

- 1) Providing compositions effective as inhibitors of human protein prolyl hydroxylase, thereby stimulating an angiogenic response in human tissue, thereby providing a method for increasing blood flow, oxygen delivery and energy utilization in ischemic tissues;

7

- 2) Providing compositions effective as human protein HIF-1 α prolyl hydroxylase inhibitors, and thereby increasing the concentration of HIF-1 α leading to greater activation and sustaining the of various biological pathways that are the normal response to cellular hypoxia;
- 3) Providing compositions effective in stimulating an erythropoietic (EPO) response in cells and thereby enhancing the maintenance of red blood cells by controlling the proliferation and differentiation of erythroid progenitor cells into red blood cells;
- 4) Providing compositions effective in stimulating an angiogenic response and thereby increasing the number and density of blood vessels and thus alleviating the adverse consequences of hypertension and diabetes, inter alia, claudication, ischemic ulcers, accelerated hypertension, and renal failure;
- 5) Providing compositions that activate Vascular Endothelial Growth Factor (VEGF) gene transcription in hypoxic cells thus increasing stimulus of important biological responses, inter alia, vasodilation, vascular permeability, and endothelial cell migration and proliferation.


Therefore, these and other unmet medical needs are resolved by the HIF-1 α prolyl hydroxylase inhibitors of the present disclosure, which are capable of regulating blood flow, oxygen delivery and energy utilization in ischemic tissues that are caused by insufficient regulation of HIF-1 α prolyl hydroxylase. Those of skill in the art will also recognize that inhibition of HIF-1 α prolyl hydroxylase enzymes will have other positive medical effects on human tissue and the alleviation of symptoms and disease states other than those symptoms or diseases states that are specifically pointed out in the present disclosure. However, as greater details arise concerning disease states and conditions related to the angiogenic process, these yet undisclosed or yet unknown conditions will be positively affected by compositions which stimulate the body own response to hypoxia and other low blood oxygen conditions.

For the purposes of the present disclosure the terms "compound," "analog," and "composition of matter" stand equally well for the HIF-1 α prolyl hydroxylase enzyme inhibitors described herein, including all enantiomeric forms, diastereomeric forms, salts, and the like, and the terms "compound," "analog," and "composition of matter" are used interchangeably throughout the present specification.

The compounds disclosed herein include all salt forms, for example, salts of both basic groups, inter alia, amines, as well as salts of acidic groups, inter alia, carboxylic acids. The following are non-limiting examples of anions that can form pharmaceutically acceptable salts with basic groups: chloride, bromide, iodide, sulfate, bisulfate, carbonate, bicarbonate, phosphate, formate, acetate, propionate, butyrate, pyruvate, lactate, oxalate, malonate, maleate, succinate, tartrate, fumarate, citrate, and the like. The following are non-limiting examples of cations that can form pharmaceutically acceptable salts of the anionic form of acidic substituent groups on the compounds described herein: sodium, lithium, potassium, calcium, magnesium, zinc, bismuth, and the like.


The HIF-1 α prolyl hydroxylase inhibitor compounds described herein are substituted aryl or heteroaryl amides, having the core structure shown in Formula (I) below.

8




wherein X can be N or CH; L is an organic linking unit as further described, below, and Y, R, R¹ and R² can be any of the units further described below.

When X is a nitrogen atom the compounds of the present disclosure are 2-amidopyridines and when X is CH the compounds of the present disclosure are arylamides, as shown below:



arylamide



2-amidopyridines

R and R¹ are optional substituent groups that can be independently chosen from a wide variety of inorganic (hydrogen, hydroxyl, amino, halogen or the like) or organic substituent units, such as alkyls, cycloalkyls, heterocyclic, heteroaryl, and the like, wherein such substituent units can optionally have from 1 to 12 carbon atoms, or 1 to 10 carbon atoms, or 1 to six carbon atoms. In many aspects of the invention, R and R¹ can each be independently a chosen from:

- i) hydrogen;
- ii) substituted or unsubstituted phenyl; and
- iii) substituted or unsubstituted heteroaryl.

wherein the optional substituent units for the phenyl and heteroaryl rings can be chosen from a wide variety of inorganic and C₁-C₄ organic radicals, and there are typically zero, one, two, or three of such substituent groups. In many such aspects, one, two, or three substituents for the above-mentioned phenyl and heteroaryl rings can be independently selected from:

- i) C₁-C₄ linear, branched, or cyclic alkyl;
- ii) C₁-C₄ linear, branched, or cyclic alkoxy;
- iii) C₁-C₄ linear, branched, or cyclic haloalkyl;
- iv) halogen;
- v) —CN;
- vi) —NHC(O)R⁴
- vii) —C(O)NR^{5a}R^{5b};
- viii) heteroaryl; or

ix) two substituents can be taken together to form a fused ring having from 5 to 7 atoms;

wherein the above-mentioned R⁴ unit can be hydrogen or a C₁-C₄ linear, branched, or cyclic alkyl; and wherein the R^{5a} and R^{5b} units can be independently selected from:

- i) hydrogen;
- ii) C₁-C₄ linear, branched, or cyclic alkyl; or
- iii) R^{5a} and R^{5b} can be taken together to form a ring having from 3 to 7 atoms.

9

In some aspects of the compounds of Formula (I), the R units can be chosen from substituted or unsubstituted phenyl; or substituted or unsubstituted heteroaryls; and the R¹ units are hydrogen.

In other aspects of the compounds of Formula (I), R can be a substituted or unsubstituted phenyl, having one, two, or three optional inorganic or organic substituents, which in some embodiments are chosen from:


- i) C₁-C₄ linear, branched, or cyclic alkyl;
- ii) C₁-C₄ linear, branched, or cyclic alkoxy;
- iii) C₁-C₄ linear, branched, or cyclic haloalkyl;
- iv) halogen; or
- v) —CN.

Non-limiting examples of R units include 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-ethylphenyl, 3-ethylphenyl, 4-ethylphenyl, 2-iso-propylphenyl, 3-iso-propylphenyl, 4-iso-propylphenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 2-iso-propoxyphenyl, 3-iso-propoxyphenyl, 4-iso-propoxyphenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, and 4-trifluoromethylphenyl.

In additional aspects of the compounds of Formula (I) the R units can have the formula —NH(C(O)R⁴) where R⁴ is C₁-C₄ linear, branched, or cyclic alkyl. Non limiting examples of such R units include:

- i) —NH(C(O)CH₃);
- ii) —NH(C(O)CH₂CH₃);
- ii) —NH(C(O)CH₂CH₂CH₃);
- ii) —NH(C(O)CH(CH₃)₂);
- ii) —NH(C(O)(cyclopropyl)); and
- ii) —NH(C(O)CH₂CH₂CH₂CH₃).

In additional aspects of the compounds of Formula (I), the R units can have the formula:



wherein R¹⁰ has the formula —C(O)NR^{5a}R^{5b}; wherein R^{5a} and R^{5b} can be independently selected from hydrogen, C₁-C₄ linear or branched alkyls, or R^{5a} and R^{5b} are taken together to form a ring having 5 or 6 atoms. In some such aspects, the R¹⁰ units can have the formula: —C(O)NR^{5a}R^{5b} wherein R^{5a} and R^{5b} are each independently selected from hydrogen, methyl, ethyl, n-propyl, iso-propyl, and cyclopropyl. Non-limiting examples of such R¹⁰ units include:


- i) —C(O)NH₂;
- ii) —C(O)NHCH₃;
- iii) —C(O)N(CH₃)₂;
- iv) —C(O)NH(CH₂CH₃);
- v) —C(O)N(CH₂CH₃)₂;
- vi) —C(O)N(CH₃)(CH₂CH₃);
- vii) —C(O)NH(CH₂CH₂CH₃);
- viii) —C(O)N(CH₂CH₂CH₃)₂;
- ix) —C(O)NH[CH(CH₃)₂];
- x) —C(O)N[CH(CH₃)₂]₂;
- xi) —C(O)N(CH₂CH₂CH₃)[CH(CH₃)₂]; and
- xii) —C(O)NH(cyclopropyl).

10


In additional aspects of the compounds of Formula (I), R^{5a} and R^{5b} together to form a ring having 5 or 6 ring atoms, non-limiting examples of R¹⁰ units are heteroaryl units chosen from pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and morpholin-4-yl.

In additional aspects of the compounds of Formula (I), the R¹⁰ units can be heteroaryl units, non limiting examples of which are thiazol-2-yl, thiazol-4-yl, 1,2,3,4-tetrazol-5-yl, [1,2,4]triazol-5-yl, imidazol-2-yl, furan-2-yl, furan-3-yl, thiophene-2-yl, thiophene-3-yl, 1,2,3,4-tetrazol-5-yl, [1,2,4]triazol-5-yl, imidazol-2-yl, furan-2-yl, furan-3-yl, pyridin-2-yl, pyridin-3-yl, pyridin-4-yl, pyrimidin-2-yl, pyrimidin-4-yl, pyrimidin-5-yl, isoquinolin-1-yl, isoquinolin-3-yl, and isoquinolin-4-yl.

In additional aspects of the compounds of Formula (I), the R units can include substituted phenyl units wherein two substitutions can be taken together to form a fused ring having from 5 to 7 ring atoms, for example a 2,3-dihydrobenzo[1,4]dioxin-6-yl ring which would provide a compound having the formula:



Other examples of R include units wherein R is hydrogen and R¹ is hydrogen.



wherein R², X, Y, L, and R⁹ can be independently chosen in any manner as otherwise taught herein with respect to the compounds of Formula (I).

As described previously above, the R¹ substituents for compounds of Formula (I) can be chosen from a wide variety of inorganic and organic units. In some embodiments, R¹ is a phenyl ring, which can optionally be substituted with 1, 2, or 3 substituent units, independently selected from inorganic or C₁-C₄ organic units. In some are chosen from:

- i) C₁-C₄ linear, branched, or cyclic alkyl;
- ii) C₁-C₄ linear, branched, or cyclic alkoxy;
- iii) C₁-C₄ linear, branched, or cyclic haloalkyl;
- iv) halogen; or
- v) —CN.

Non-limiting examples of R¹ include 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 2-chlorophenyl, 3-chlorophenyl, 4-chlorophenyl, 2-methylphenyl, 3-methylphenyl, 4-methylphenyl, 2-ethylphenyl, 3-ethylphenyl, 4-ethylphenyl, 2-iso-propylphenyl, 3-iso-propylphenyl, 4-iso-propylphenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl,

11

2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 2-isopropoxyphenyl, 3-isopropoxyphenyl, 4-isopropoxyphenyl, 2-cyanophenyl, 3-cyanophenyl, 4-cyanophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, and 4-trifluoromethylphenyl.

Another example of R¹ units includes compounds wherein R is hydrogen and R¹ units are hydrogen.

In some aspects of the compounds of Formula (I), R is hydrogen and R¹ is a substituted or unsubstituted phenyl, wherein the substitutions are chosen from:

- i) C₁-C₄ linear, branched, or cyclic alkyl;
- ii) C₁-C₄ linear, branched, or cyclic alkoxy;
- iii) C₁-C₄ linear, branched, or cyclic haloalkyl;
- iv) halogen; and
- v) —CN.

In connection with the compounds of Formula (I), Y is a unit that can be chosen from a wide variety of inorganic units, such as H, —OH, —NH₂, or a halogen, and C₁-C₄ organic units. For example, Y can be chosen from:

- i) hydrogen;
- ii) —OR³, wherein R³ is hydrogen, or a lower alkyl, such as methyl, or ethyl.

In connection with the compounds of Formula (I), the R² units can be chosen from a wide variety of inorganic units, such as —OH, or —NH₂ units, or a variety of organic units.


In some aspects of the compounds of the invention, R² is chosen from:

- i) —OR⁶; or
- ii) —NR^{7a}R^{7b};

wherein R⁶ is hydrogen or C₁-C₄ linear, branched, or cyclic alkyl; and R^{7a} and R^{7b} are each independently chosen from:

- i) hydrogen;
- ii) C₁-C₄ linear, branched, or cyclic alkyl; or
- iii) R^{7a} and R^{7b} can be taken together to form a ring having from 3 to 7 atoms.

In certain favored aspects of the invention the R² units are hydroxyl (—OH) wherein the compounds are carboxylic acids, or may also be present in the form of a salt as the corresponding hydroxyl/carboxylate anion, i.e. R² can be a —O⁻ unit, so as to form a compound having a carboxylate group, as shown below.



wherein R, R¹, R², X, Y, L, and R⁹ can be independently chosen in any manner as otherwise taught herein with respect to the compounds of Formula (I).

Another example of R² includes compounds wherein R⁶ is C₁-C₄ linear, branched, or cyclic alkyl providing R² units which are alkoxy wherein the compounds formed are organic esters having C₁-C₄ linear, branched, or cyclic alkyl groups. Non-limiting examples of R² units are:

- i) —OCH₃;
- ii) —OCH₂CH₃; and
- iii) —OCH₂CH₂CH₃.

12

Yet further examples of R² units include compounds wherein R² has the formula —NR^{7a}R^{7b}; and R^{7a} and R^{7b} are each independently chosen from:

- i) hydrogen; and
 - ii) C₁-C₄ linear, branched, or cyclic alkyl.
- Non-limiting examples of R² units include:
- i) —NH₂;
 - ii) —NHCH₃;
 - iii) —N(CH₃)₂;
 - iv) —NH(CH₂CH₃);
 - v) —N(CH₂CH₃)₂;
 - vi) —N(CH₃)(CH₂CH₃);
 - vii) —NH(CH₂CH₂CH₃);
 - viii) —N(CH₂CH₂CH₃)₂;
 - ix) —NH[CH(CH₃)₂];
 - x) —N[CH(CH₃)₂]₂;
 - xi) —N(CH₂CH₂CH₃)[CH(CH₃)₂]; and
 - xii) —NH(cyclopropyl).

A yet further example of R² units includes compounds wherein R² has the formula —NR^{7a}R^{7b}; and R^{7a} and R^{7b} are taken together to form a ring having from 3 to 7 atoms, wherein the non-limiting examples of R² units include aziridin-1-yl, axetidin-1-yl, pyrrolidin-1-yl, piperidin-1-yl, piperazin-1-yl, and morpholin-4-yl.

In connection with the compounds of Formula (I), L is a unit that links the nitrogen atom of the carboxamide group to the neighboring carbonyl group. L is typically a C₁-C₆ or C₁-C₄ organic linking unit. In some embodiments, L comprises a one or more optionally substituted methylene units having the formula:




wherein R^{8a} and R^{8b} are each independently hydrogen, C₁-C₆ linear or branched alkyl, or phenyl; and the index y is from 1 to 4.

An example of L units includes units wherein R^{8a} and R^{8b} are both hydrogen and the index n is equal to 1, the L unit has the formula:




and are referred to herein as methylene linking units, so as to form compounds having the structure shown below:




wherein R, R¹, R², X, Y, and R⁹ can be independently chosen in any manner as otherwise taught herein with respect to the compounds of Formula (I).

Another example of L units includes units wherein R^{8a} and R^{8b} are each hydrogen or methyl and the index n is equal to 1, these units having the formula:



A further example of L units includes units wherein R^{8a} and R^{8b} units are hydrogen and the index n is equal to 2, these units having the formula:



and are referred to herein as ethylene linking units.


In connection with the compounds of Formula I, the R⁹ substituent for the amide nitrogen atom can be hydrogen or

13


a C₁-C₄ organic substituent, such as a C₁-C₄ alkyl group, such as methyl, or a C₁-C₄ haloalkyl, such as a trifluoromethyl group.

The compounds of Formula (I) can be organized into several categories for the strictly non-limiting purpose of describing alternatives for synthetic strategies for the preparation of subgenera of compounds within the scope of Formula (I) that are not expressly exemplified herein. This mental organization into categories does not imply anything with respect to increased or decreased biological efficacy with respect to any of the compounds or compositions of matter described herein.

One such subgenus of the compounds of Formula (I) relates to compounds having the formula:



which can be more specifically described by methyl ester compounds having the formula:



wherein R units can be substituted or unsubstituted phenyl, non-limiting examples of which are described in Table I herein below.

TABLE I


No.	R
1	2-fluorophenyl
2	3-fluorophenyl
3	4-fluorophenyl
4	2-chlorophenyl
5	3-chlorophenyl
6	4-chlorophenyl
7	2-cyanophenyl
8	3-cyanophenyl
9	4-cyanophenyl
10	2-methylphenyl
11	3-methylphenyl
12	4-methylphenyl
13	2-ethylphenyl
14	3-ethylphenyl
15	4-ethylphenyl
16	2-methoxyphenyl
17	3-methoxyphenyl
18	4-methoxyphenyl
19	2-ethoxyphenyl
20	3-ethoxyphenyl
21	4-ethoxyphenyl
22	2-iso-propoxyphenyl
23	3-iso-propoxyphenyl
24	4-iso-propoxyphenyl
25	2-carbamoylphenyl
26	3-carbamoylphenyl
27	4-carbamoylphenyl
28	2-(aziridine-1-carbonyl)phenyl
29	3-(aziridine-1-carbonyl)phenyl
30	4-(aziridine-1-carbonyl)phenyl

14


TABLE I-continued

No.	R
31	2-(azetidine-1-carbonyl)phenyl
32	3-(azetidine-1-carbonyl)phenyl
33	4-(azetidine-1-carbonyl)phenyl
34	2-(pyrrolidine-1-carbonyl)phenyl
35	3-(pyrrolidine-1-carbonyl)phenyl
36	4-(pyrrolidine-1-carbonyl)phenyl
37	2-(piperidine-1-carbonyl)phenyl
38	3-(piperidine-1-carbonyl)phenyl
39	4-(piperidine-1-carbonyl)phenyl
40	2-(acetylamino)phenyl
41	3-(acetylamino)phenyl
42	4-(acetylamino)phenyl
43	2-(ethanecarbonylamino)phenyl
44	3-(ethanecarbonylamino)phenyl
45	4-(ethanecarbonylamino)phenyl
46	2-(cyclopropanecarbonylamino)phenyl
47	3-(cyclopropanecarbonylamino)phenyl
48	4-(cyclopropanecarbonylamino)phenyl



Such compounds can be prepared by the procedure outlined in Scheme I and further described in Example 1 herein below.





15




Reagents and conditions: (c) GlyOMe.HCl, EDCl,
HOBt, DMF; 0° C. to rt, 3 days




Reagents and conditions: (d) H₂: Pd/C, MeOH, rt,
16 hr



Reagents and conditions: (e) (CF₃SO₂)₂NC₆H₅,
MeOH, rt, 16 hr



16



Reagents and conditions: (f) 3-chlorophenylboronic
acid, Pd(dppf)Cl₂, K₃PO₄, dioxane; 85° C., 16 hr

EXAMPLE 1

{[5-(3-Chloro-phenyl)-3-hydroxy-pyridine-2-carbo-
nyl]-amino}acetic acid methyl ester (6)

Preparation of 3,5-bis-benzyloxy-pyridine-2-carbonitrile
(1): To an 80 mL microwave pressure vessel is charged dry
THF (30 mL) and benzyl alcohol (6.32 mL, 61.1 mmol). The
solution is cooled to 0° C. and sodium hydride (2.44 g of a
60% dispersion in mineral oil, 61.1 mmol) is added in
portions. The reaction mixture is gradually allowed to warm
to room temperature with efficient stirring until the evolution
of hydrogen gas ceases. The solution is re-cooled to 0° C.
and 3,5-dichloro-2-cyanopyridine (5.00 g, 29.1 mmol) is
added, and the solution is transferred to an unfocused Mars
5 CEM microwave reactor to 190° C., 300 W and held for
5 hours. The reaction mixture is quenched with H₂O, con-
centrated under reduced pressure, diluted with EtOAc and
washed with 2M Na₂CO₃, H₂O and saturated aqueous NaCl.
The organic layer is dried (MgSO₄), filtered and concen-
trated under reduced pressure to give a brown solid. The
crude solid is purified over silica (EtOAc:heptane, gradient
1:1 to 1:0) to afford 8.6 g (94% yield) of the desired
compound as an orange solid. ¹H NMR (400 MHz, CDCl₃)
δ ppm 7.96 (1 H, d, J=2.2 Hz), 7.25-7.37 (10 H, m), 6.78 (1
H, d, J=2.2 Hz), 5.10 (2 H, s), 5.03 (2 H, s). HPLC-MS: m/z
317 [M+H]⁺.

Preparation of 3,5-bis-benzyloxy-pyridine-2-carboxylic
acid (2): To a solution of 3,5-bis-benzyloxy-pyridine-2-
carbonitrile, 1, (26.0 g, 82.3 mmol) in MeOH (217 mL) is
added 30% w/v sodium hydroxide (320 mL) and the reaction
mixture is refluxed for 16 hours. The solvent is removed
under reduced pressure and the resulting suspension is
acidified with conc. HCl until the pH is between 1 and 2. The
precipitate that results is collected by filtration, washed with
H₂O (10 mL) and dried overnight in a vacuum oven to afford
30 g (quantitative) of the desired product as the hydrochloride
salt. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 8.02 (1 H,
d, J=2.4 Hz), 7.29-7.53 (11 H, m), 5.96 (1 H, br s), 5.28 (4
H, s). HPLC-MS: m/z 336 [M+H]⁺.

Preparation of [(3,5-bis-benzyloxy-pyridine-2-carbonyl)-
amino]acetic acid methyl ester (3): To a solution of 3,5-
bis-benzyloxy-pyridine-2-carboxylic acid HCl, 2, (8.06 g,
21.7 mmol) in DMF (100 mL) at 0° C. under N₂ is added
diisopropylethylamine (11.35 mL, 65.1 mmol), 1-(3-dim-
ethylaminopropyl)-3-ethylcarbodiimide (EDCI) (6.23 g,
32.6 mmol) and 1-hydroxybenzotriazole (HOBt) (0.294 g,
2.2 mmol). The solution is stirred for 5 minutes and glycine
methyl ester hydrochloride (4.09 g, 32.6 mmol) is added.

17

The reaction is allowed to warm slowly to room temperature and stirred 3 days. The reaction volume is partially concentrated under reduced pressure then diluted with EtOAc and washed with saturated aqueous NaHCO₃ and saturated aqueous NaCl. The organic layer is dried (MgSO₄), filtered and concentrated under reduced pressure to afford a yellow oil that is purified over silica (EtOAc:heptane gradient 1:1 to 1:0) to afford 3.5 g (40% yield) of the desired product as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ ppm 8.12 (1 H, t, J=4.9 Hz), 7.95 (1 H, d, J=1.8 Hz), 7.38-7.44 (2 H, m), 7.22-7.35 (8 H, m), 6.85 (1 H, d, J=2.6 Hz), 5.14 (2 H, s), 5.03 (2 H, s), 4.18 (2 H, d, J=5.5 Hz), 3.69 (3 H, s). HPLC-MS: m/z 407 [M+H]⁺.

Preparation of [(3,5-dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid methyl ester (4): To a solution [(3,5-bis-benzyloxy-pyridine-2-carbonyl)-amino]-acetic acid methyl ester (3.50 g, 8.62 mmol) in MeOH (100 mL) is added 10% Pd/C (0.350 g, 0.862 mmol) and the reaction mixture stirred under an atmosphere of H₂ at room temperature for 16 hours. The suspension is filtered through Celite™ and the filtrate concentrated under reduced pressure. The crude material is purified over silica (MeOH:CH₂Cl₂ gradient 1% to 5%) to afford 1.95 g (quantitative yield) of the desired compound as an off-white solid. ¹H NMR (250 MHz, MeOD) δ ppm 7.62 (1 H, d, J=2.4 Hz), 6.53 (1 H, d, J=2.4 Hz), 4.04 (2 H, s), 3.64 (3 H, s). HPLC-MS: m/z 227 [M+H]⁺.

Preparation of [(3-hydroxy-5-trifluoromethanesulfonyloxy-pyridine-2-carbonyl)-amino]-acetic acid methyl ester (5): To a solution of [(3,5-dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid methyl ester, 4, (1.95 g, 8.62 mmol) in MeOH (60 mL) is added diisopropylethylamine (DIPEA) (1.62 mL, 9.3 mmol). The mixture is cooled to 0° C. and N-phenyl trifluoromethanesulfonimide (3.32 g, 9.3 mmol) is added. The resulting solution is slowly warmed to room temperature and stirred for an additional 16 hours. The solvent is removed under reduced pressure and the crude material is purified over silica (EtOAc:hexane 1:4) to afford 2.27 g (73% yield) of the desired product as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 12.17 (1 H, s), 8.27 (1 H, t, J=5.5 Hz), 8.08 (1 H, d, J=2.2 Hz), 7.28 (1 H, d, J=2.2 Hz), 4.24 (2 H, d, J=5.5 Hz), 3.82 (3 H, s). HPLC-MS: m/z 359 [M+H]⁺.


Preparation of {[5-(3-chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester (6): To a degassed solution of [(3-hydroxy-5-trifluoromethanesulfonyloxy-pyridine-2-carbonyl)-amino]-acetic acid methyl ester, 5, (0.30 g, 0.84 mmol) in 1,4-dioxane (10 mL) at room temperature under N₂ is added 3-chlorophenylboronic acid (0.196 g, 1.26 mmol), Pd(dppf)Cl₂ (0.068 g, 0.0084 mmol) and K₃PO₄ (0.195 g, 0.92 mmol). The resulting suspension is heated in a sealed tube at 85° C. for 16 hours. After this time, the mixture is cooled to room temperature and concentrated under reduced pressure. The residue is then treated with 1M HCl (1 mL) and diluted with EtOAc. The organic layer is separated, washed with H₂O, saturated aqueous NaCl and concentrated under reduced pressure. The crude material is purified over silica (EtOAc:heptane 3:7). The resulting solid can be crystallized from EtOAc/heptane to afford 0.143 g (53% yield) of the desired compound as a colorless solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 11.77 (1 H, s), 8.36 (1 H, t, J=5.7 Hz), 8.24 (1 H, d, J=1.8 Hz), 7.50-7.53 (1 H, m), 7.39-7.42 (2 H, m), 7.34-7.37 (2 H, m), 4.20 (2 H, d, J=5.9 Hz), 3.76 (3 H, s). HPLC-MS: m/z.

The procedure outlined in Scheme I can be modified by substituting in step (f) other reagents for 3-chlorophenylboronic acid. Non-limiting examples include 4-chlorophenylboronic acid, 2-chlorophenylboronic acid, 2-fluorophenylboronic acid, 3-fluorophenylboronic acid,


18

4-fluorophenylboronic acid, 2-methylphenylboronic acid, 3-methylphenylboronic acid, and 4-methylphenylboronic acid.


The following are further non-limiting examples of compounds encompassed within first aspect of Category I of the present disclosure.




{[5-(4-Chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ¹H NMR (400 MHz, CDCl₃) δ ppm 11.77 (1 H, s), 8.36 (1 H, t, J=5.5 Hz), 8.23 (1 H, d, J=1.8 Hz), 7.44-7.49 (2 H, m), 7.38-7.42 (3 H, m), 4.20 (2 H, d, J=5.9 Hz), 3.76 (3 H, s). HPLC-MS: m/z 321 [M+H]⁺.



{[5-(2-Chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ¹H NMR (400 MHz, MeOD) δ ppm 8.10 (1 H, d, J=1.8 Hz), 7.46 (1 H, dd, J=7.5, 2.4 Hz), 7.30-7.35 (4 H, m), 4.11 (2 H, s), 3.68 (3 H, s). HPLC-MS: m/z 321 [M+H]⁺.




{[5-(4-Fluorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ¹H NMR (250 MHz, CDCl₃) δ ppm 11.88 (1 H, s), 8.48 (1 H, t, J=5.6 Hz), 8.33 (1 H, d, J=2.1 Hz), 7.55-7.65 (2 H, m), 7.49 (1 H, d, J=2.1 Hz), 7.17-7.27 (2 H, m), 4.28-4.32 (2 H, m), 3.86 (3 H, s). HPLC-MS: m/z 305 [M+H]⁺.




19


{[3-Hydroxy-5-(4-methylphenyl)-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (400 MHz, CDCl_3) δ ppm 11.72 (1 H, s), 8.36 (1 H, t, $J=5.1$ Hz), 8.26 (1 H, d, $J=1.8$ Hz), 7.43 (2 H, d, $J=8.0$ Hz), 7.40 (1 H, d, $J=1.8$ Hz), 7.23 (2 H, d, $J=8.1$ Hz), 4.19 (2 H, d, $J=5.9$ Hz), 3.75 (3 H, s), 2.35 (3 H, s). HPLC-MS: m/z 301 $[\text{M}+\text{H}]^+$.




{[3-Hydroxy-5-(4-isopropylphenyl)-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.88 (1 H, s), 8.54 (1 H, t, $J=5.6$ Hz), 8.38 (1 H, d, $J=1.8$ Hz), 7.55-7.60 (2 H, m), 7.54 (1 H, d, $J=2.1$ Hz), 7.36-7.42 (2 H, m), 4.30 (2 H, d, $J=5.8$ Hz), 3.85 (3 H, s), 2.93-3.07 (1 H, m), 1.33 (6 H, d, $J=7.0$ Hz). HPLC-MS: m/z 329 $[\text{M}+\text{H}]^+$.



{[5-(4-Ethylphenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.86 (1 H, s), 8.51 (1 H, t, $J=5.8$ Hz), 8.37 (1 H, d, $J=1.8$ Hz), 7.56 (2 H, d, $J=8.2$ Hz), 7.53 (1 H, d, $J=1.8$ Hz), 7.36 (2 H, d, $J=8.5$ Hz), 4.30 (2 H, d, $J=5.8$ Hz), 3.85 (3 H, s), 2.75 (2 H, q, $J=7.6$ Hz), 1.31 (3 H, t, $J=7.6$ Hz). HPLC-MS: m/z 315 $[\text{M}+\text{H}]^+$.




{[3-Hydroxy-5-(3-trifluoromethyl-phenyl)-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.91 (1 H, s), 8.48 (1 H, t, $J=6.1$ Hz), 8.37 (1 H, d, $J=1.8$ Hz), 7.60-7.92 (4 H, m), 7.54 (1 H, d, $J=2.1$ Hz), 4.31 (2 H, d, $J=5.8$ Hz), 3.86 (3 H, s). HPLC-MS: m/z 355 $[\text{M}+\text{H}]^+$.




{[5-(4-Cyanophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (400 MHz,

20


CDCl_3) δ ppm 11.83 (1 H, s), 8.36 (1 H, t, $J=5.12$ Hz), 8.26 (1 H, d, $J=1.83$ Hz), 7.70-7.75 (2 H, m), 7.62-7.66 (2 H, m), 7.43 (1 H, d, $J=1.83$ Hz), 4.21 (2 H, d, $J=5.49$ Hz), 3.76 (3 H, s). HPLC-MS: m/z 312 $[\text{M}+\text{H}]^+$.



{[5-(3-Cyanophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, DMSO-d_6) δ ppm 12.30 (1 H, s), 9.51 (1 H, t, $J=5.8$ Hz), 8.55 (1 H, d, $J=1.8$ Hz), 8.32 (1 H, s), 8.14 (1 H, d, $J=8.5$ Hz), 7.89 (1 H, d, $J=7.8$ Hz), 7.81 (1 H, d, $J=1.9$ Hz), 7.68 (1 H, t, $J=7.8$ Hz), 4.06 (2 H, d, $J=6.1$ Hz), 3.63 (3 H, s). HPLC-MS: m/z 312 $[\text{M}+\text{H}]^+$.



{[5-(3-Carbamoylphenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, DMSO-d_6) δ ppm 12.30 (1 H, s), 9.54 (1 H, t, $J=6.0$ Hz), 8.61 (1 H, d, $J=1.8$ Hz), 8.29 (1 H, s), 8.18 (1 H, br s), 7.98 (2 H, t, $J=8.1$ Hz), 7.84 (1 H, d, $J=1.7$ Hz), 7.62 (1 H, t, $J=7.8$ Hz), 7.53 (1 H, br s), 4.12 (2 H, d, $J=6.0$ Hz), 3.69 (3 H, s). HPLC-MS: m/z 330 $[\text{M}+\text{H}]^+$.




{[3-Hydroxy-5-[3-(pyrrolidine-1-carbonyl)-phenyl]-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.80 (1 H, s), 8.44 (1 H, t, $J=5.3$ Hz), 8.35 (1 H, s), 7.77 (1 H, s), 7.47-7.70 (4 H, m), 4.28 (2 H, d, $J=5.7$ Hz), 3.83 (3 H, s), 3.64-3.76 (2 H, m), 3.42-3.55 (2 H, m), 1.84-2.07 (4 H, m). HPLC-MS: m/z 384 $[\text{M}+\text{H}]^+$.



21


{[5-[3-(Cyclopropanecarbonyl-amino)-phenyl]-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.80 (1 H, s), 8.45 (1 H, br s), 8.29 (1 H, s), 7.65-7.88 (2 H, m), 7.29-7.60 (4 H, m), 4.18-4.31 (3 H, m), 3.83 (3 H, s), 1.05-1.17 (2 H, m), 0.81-0.98 (2 H, m). HPLC-MS: m/z 370 $[\text{M}+\text{H}]^+$.

The following heteroaryl substituted phenyl compound can be prepared from {[5-(3-cyano-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}acetic acid methyl ester by treatment with trimethylsilyl azide and di-butyl tin oxide in DME and heating the mixture to 140°C ., 150 W, 200 psi in a microwave reactor.



{[3-Hydroxy-5-[3-(2 H-tetrazol-5-yl)-phenyl]-pyridine-2-carbonyl]-amino}-acetic acid methyl ester: ^1H NMR (400 MHz, DMSO-d_6) δ ppm 12.33 (1 H, s), 9.55 (1 H, t, $J=6.1$ Hz), 8.62 (1 H, d, $J=1.8$ Hz), 8.43 (1 H, s), 8.14 (1 H, d, $J=7.9$ Hz), 8.05 (1 H, d, $J=8.1$ Hz), 7.83 (1 H, d, $J=1.9$ Hz), 7.76 (1 H, t, $J=7.8$ Hz), 4.12 (2 H, d, $J=6.1$ Hz), 3.69 (3 H, s). HPLC-MS: m/z 355 $[\text{M}+\text{H}]^+$.

For the purposes of describing synthetic methods, another mental subgenus of the compounds of Formula (I) have the formula:



wherein R units which are substituted or unsubstituted heteroaryl, non-limiting examples of which are described in Table II herein below.

TABLE II


No.	R
49	pyridin-2-yl
50	pyridin-3-yl
51	pyridin-4-yl
52	pyrimidin-2-yl
53	pyrimidin-4-yl
54	pyrimidin-5-yl
55	isoquinolin-1-yl
56	isoquinolin-3-yl
57	isoquinolin-4-yl
58	thiazol-2-yl
59	thiazol-4-yl
60	1,2,3,4-tetrazol-5-yl
61	[1,2,4]triazol-5-yl
62	imidazol-2-yl
63	furan-2-yl
64	furan-3-yl
65	thiophene-2-yl
66	thiophene-3-yl

The compounds encompassed within the compounds described immediately above can be prepared by the procedure outlined in Scheme I and described in Example 1 herein above.


22

The procedure outlined in Scheme I can be modified by substituting in step (f) other reagents for 3-chlorophenylboronic acid. Non-limiting examples of substitutes include 3-(thiazol-2-yl)phenylboronic acid, 3-(thiazol-4-yl)phenylboronic acid, 4-(thiazol-2-yl)phenylboronic acid, 4-(thiazol-4-yl)phenylboronic acid, 3-(imidazol-2-yl)phenylboronic acid, 4-(imidazol-2-yl)phenylboronic acid, 3-(furan-2-yl)phenylboronic acid, 3-(furan-3-yl)phenylboronic acid, and 3-(thiophene-2-yl)phenylboronic acid.


The following are non-limiting examples of such compounds.



[(5-Hydroxy-[3,3']bipyridinyl-6-carbonyl)-amino]-acetic acid methyl ester ^1H NMR (250 MHz, CDCl_3) δ ppm 11.80 (1 H, s), 8.87 (1 H, d, $J=1.7$ Hz), 8.70 (1 H, dd, $J=4.8, 1.6$ Hz), 8.45 (1 H, t, $J=5.8$ Hz), 8.33 (1 H, d, $J=1.9$ Hz), 7.91 (1 H, ddd, $J=8.0, 2.3, 1.7$ Hz), 7.51 (1 H, d, $J=1.9$ Hz), 7.45 (1 H, ddd, $J=7.9, 4.8, 0.7$ Hz), 4.28 (2 H, d, $J=5.8$ Hz), 3.83 (3 H, s). HPLC-MS: m/z 288 $[\text{M}+\text{H}]^+$.




[(5'-Hydroxy-[2,3']bipyridinyl-6'-carbonyl)-amino]-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 8.70-8.79 (2 H, m), 8.49 (1 H, t, $J=5.8$ Hz), 7.72-7.92 (3 H, m), 7.31-7.39 (1 H, m), 4.28 (2 H, d, $J=5.8$ Hz), 3.83 (3 H, s). HPLC-MS: m/z 288 $[\text{M}+\text{H}]^+$.




{[3-(3-Hydroxy-5-pyrimidin-5-yl-pyridine-2-carbonyl)-amino]-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.90 (1 H, s), 9.32 (1 H, s), 9.00 (2 H, s), 8.45 (1 H, br s), 8.34 (1 H, d, $J=1.8$ Hz), 7.53 (1 H, d, $J=1.8$ Hz), 4.29 (2 H, d, $J=5.7$ Hz), 3.84 (3 H, s). HPLC-MS: m/z 289 $[\text{M}+\text{H}]^+$.

23




[(3-Hydroxy-5-isoquinolin-4-yl-pyridine-2-carbonyl)-amino]-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.80 (1 H, s), 9.38 (1 H, br s), 8.51 (2 H, t, $J=5.7$ Hz), 8.27 (1 H, s), 8.13 (1 H, d, $J=7.2$ Hz), 7.67-7.93 (3 H, m), 7.51 (1 H, s), 4.31 (2 H, d, $J=5.7$ Hz), 3.85 (3 H, s). HPLC-MS: m/z 338 $[\text{M}+\text{H}]^+$.



[(3-Hydroxy-5-thiazol-2-yl-pyridine-2-carbonyl)-amino]-acetic acid methyl ester: ^1H NMR (250 MHz, CDCl_3) δ ppm 11.80 (1 H, s), 8.67 (1 H, d, $J=1.8$ Hz), 8.39 (1 H, br s), 7.93 (1 H, d, $J=3.3$ Hz), 7.78 (1 H, d, $J=1.7$ Hz), 7.43 (1 H, d, $J=3.2$ Hz), 4.22 (2H, d, $J=5.8$ Hz), 3.78 (3 H, s). HPLC-MS: m/z 294 $[\text{M}+\text{H}]^+$.

Another mental subgenus of the compounds of Formula (I) encompasses compounds having the formula:



wherein R units which are substituted or unsubstituted phenyl, non-limiting examples of which are described in Table III herein below.

TABLE III

No.	R
67	2-fluorophenyl
68	3-fluorophenyl
69	4-fluorophenyl
70	2-chlorophenyl
71	3-chlorophenyl
72	4-chlorophenyl
73	2-cyanophenyl
74	3-cyanophenyl
75	4-cyanophenyl
76	2-methylphenyl
77	3-methylphenyl
78	4-methylphenyl
79	2-ethylphenyl
80	3-ethylphenyl
81	4-ethylphenyl
82	2-methoxyphenyl
83	3-methoxyphenyl
84	4-methoxyphenyl
85	2-ethoxyphenyl
86	3-ethoxyphenyl


24

TABLE III-continued

No.	R
87	4-ethoxyphenyl
88	2-iso-propoxyphenyl
89	3-iso-propoxyphenyl
90	4-iso-propoxyphenyl
91	2-carbamoylphenyl
92	3-carbamoylphenyl
93	4-carbamoylphenyl
94	2-(aziridine-1-carbonyl)phenyl
95	3-(aziridine-1-carbonyl)phenyl
96	4-(aziridine-1-carbonyl)phenyl
97	2-(azetidone-1-carbonyl)phenyl
98	3-(azetidone-1-carbonyl)phenyl
99	4-(azetidone-1-carbonyl)phenyl
100	2-(pyrrolidine-1-carbonyl)phenyl
101	3-(pyrrolidine-1-carbonyl)phenyl
102	4-(pyrrolidine-1-carbonyl)phenyl
103	2-(piperidine-1-carbonyl)phenyl
104	3-(piperidine-1-carbonyl)phenyl
105	4-(piperidine-1-carbonyl)phenyl
106	2-(acetylamino)phenyl
107	3-(acetylamino)phenyl
108	4-(acetylamino)phenyl
109	2-(ethanecarbonylamino)phenyl
110	3-(ethanecarbonylamino)phenyl
111	4-(ethanecarbonylamino)phenyl
112	2-(cyclopropanecarbonylamino)phenyl
113	3-(cyclopropanecarbonylamino)phenyl
114	4-(cyclopropanecarbonylamino)phenyl

The compounds described immediately above can be prepared by the procedure outlined in Scheme II and described in Example 2.

Scheme II



Reagents and conditions: (a) NaOH, $\text{H}_2\text{O}/\text{THF}$; rt, 1 hr

EXAMPLE 2


{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid (7)

Preparation of {[5-(3-chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid (7): To a solution of {[5-(3-


25

chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester, 6, (0.163 g, 0.509 mmol) in THF (5 mL) is added 1M NaOH (1.5 ml, 1.27 mmol) and the reaction mixture stirred at room temperature for 1 hour. The solution is acidified using 1M HCl (3 mL), the solvent removed under reduced pressure and the resulting solid suspended in CHCl₃:iso-propanol (1:1), filtered and the filtrate dried (MgSO₄), filtered and re-concentrated under reduced pressure. The crude material is triturated with a small amount of MeOH to afford 0.10 g (64% yield) of the desired product as a colorless solid. ¹H NMR (400 MHz, MeOD) δ ppm 8.31 (1H, d, J=1.8 Hz), 7.47 (2 H, d, J=1.8 Hz), 7.30-7.65 (4H, m), 4.07 (2H, s). HPLC-MS: m/z 307 [M+H]⁺.


The following are further non-limiting examples.



{[5-(4-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (400 MHz, MeOD) δ ppm 8.33 (1 H, d, J=1.5 Hz), 7.61 (2 H, d, J=8.4 Hz), 7.48 (1 H, d, J=1.8 Hz), 7.42 (2 H, d, J=8.4 Hz), 4.06 (2 H, s). HPLC-MS: m/z 307 [M+H]⁺.




{[5-(2-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (400 MHz, MeOD) δ ppm 8.10 (1 H, d, J=1.8 Hz), 7.40-7.56 (1 H, m), 7.09-7.40 (4 H, m), 4.07 (2H, s). HPLC-MS: m/z 307 [M+H]⁺.




{[5-(4-Fluorophenyl)-3-hydroxypyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.39 (1 H, br s), 9.38 (1 H, t, J=6.2 Hz), 8.53 (1 H, d, J=2.1 Hz), 7.91 (2 H, dd, J=8.8, 5.5 Hz), 7.74 (1 H, d, J=2.1 Hz), 7.38 (2 H, t, J=8.8 Hz), 4.02 (2 H, d, J=6.4 Hz). HPLC-MS: m/z 291 [M+H]⁺.


26




{[3-Hydroxy-5-(4-methylphenyl)pyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (400 MHz, MeOD) δ ppm 8.40 (1 H, s), 7.68 (1 H, s), 7.53 (2 H, d, J=8.42 Hz), 7.26 (2H, d, J=8.05 Hz), 4.10 (2 H, s), 2.31 (3 H, s). HPLC-MS: m/z 287 [M+H]⁺.



{[5-(4-Ethylphenyl)-3-hydroxypyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.40 (1 H, s), 9.35 (1 H, t, J=6.1 Hz), 8.52 (1 H, d, J=2.1 Hz), 7.76 (2 H, d, J=8.2 Hz), 7.71 (1 H, d, J=1.8 Hz), 7.38 (2 H, d, J=8.2 Hz), 4.02 (2 H, d, J=6.1 Hz), 2.68 (2 H, q, J=7.6 Hz), 1.22 (3 H, t, J=7.5 Hz). HPLC-MS: m/z 301 [M+H]⁺.




{[3-Hydroxy-5-(4-isopropylphenyl)pyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.40 (1 H, s), 9.36 (1 H, t, J=6.2 Hz), 8.52 (1 H, d, J=1.8 Hz), 7.70 (2 H, d, J=1.8 Hz), 7.76 (2 H, d, J=8.2 Hz), 7.41 (2 H, d, J=8.2 Hz), 4.02 (2 H, d, J=6.1 Hz), 2.97 (1 H, m, J=7.0 Hz), 1.25 (6 H, d, J=7.0 Hz). HPLC-MS: m/z [M+H]⁺ 315.




{[3-Hydroxy-5-(3-trifluoromethylphenyl)pyridine-2-carbonyl] amino}-acetic acid: ¹H NMR (250 MHz, DMSO-d₆)

27


δ ppm 12.42 (1 H, br s), 9.41 (1 H, t, $J=6.4$ Hz), 8.61 (1 H, d, $J=1.8$ Hz), 8.10-8.22 (2 H, m), 7.88 (1 H, d, $J=1.8$ Hz), 7.73-7.86 (2 H, m), 4.03 (2 H, d, $J=6.1$ Hz). HPLC-MS: m/z 341 $[M+H]^+$.




{[5-(4-Cyanophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid: 1H NMR (400 MHz, MeOD) δ ppm 8.38 (1 H, d, $J=1.8$ Hz), 7.75-7.83 (4 H, m), 7.56 (1 H, d, $J=1.8$ Hz), 4.06 (2 H, s). HPLC-MS: m/z 298 $[M+H]^+$.



{[5-(3-Cyanophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.40 (1 H, s), 9.40 (1 H, t, $J=6.17$ Hz), 8.59 (1 H, d, $J=1.71$ Hz), 8.37 (1 H, s), 8.19 (1 H, d, $J=7.77$ Hz), 7.93 (1 H, d, $J=7.88$ Hz), 7.86 (1 H, d, $J=1.94$ Hz), 7.73 (1 H, t, $J=7.77$ Hz), 4.00 (2 H, d, $J=6.17$ Hz). HPLC-MS: m/z 298 $[M+H]^+$.




{[5-(5-Chloro-2-methylphenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.35 (1 H, br s), 9.34 (1 H, t, $J=6.0$ Hz), 8.17 (1 H, d, $J=1.8$ Hz), 7.47 (1 H, d, $J=1.8$ Hz), 7.33-7.45 (3 H, m), 3.95 (2 H, d, $J=5.9$ Hz), 2.22 (3 H, s). HPLC-MS: m/z 321 $[M+H]^+$.




{[3-Hydroxy-5-(4-isopropoxyphenyl)pyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.36 (1 H, br s), 9.35 (1 H, t, $J=5.9$ Hz), 8.50 (1 H, d,

28


$J=1.8$ Hz), 7.71 (1 H, d, $J=1.8$ Hz), 7.27-7.46 (3 H, m), 7.02 (1 H, d, $J=8.5$ Hz), 4.70-4.85 (1 H, m), 4.01 (2 H, d, $J=6.1$ Hz), 1.29 (6 H, d, $J=5.9$ Hz). HPLC-MS: m/z 331 $[M+H]^+$.



{[5-[3-(Cyclopropanecarbonylamino)phenyl]-3-hydroxypyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.40 (1 H, br s), 10.39 (1 H, s), 9.37 (1 H, t, $J=6.1$ Hz), 8.43 (1 H, d, $J=1.9$ Hz), 7.97 (1 H, s), 7.64-7.71 (1 H, m), 7.60 (1 H, d, $J=1.8$ Hz), 7.41-7.49 (2 H, m), 3.99 (2 H, d, $J=6.1$ Hz), 1.73-1.86 (1 H, m), 0.74-0.84 (4 H, m). HPLC-MS: m/z 356 $[M+H]^+$.




{[3-Hydroxy-5-[3-(pyrrolidine-1-carbonyl)phenyl]-pyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.40 (1 H, s), 9.38 (1 H, t, $J=6.3$ Hz), 8.55 (1 H, d, $J=1.8$ Hz), 7.88-7.94 (2 H, m), 7.77 (1 H, d, $J=1.8$ Hz), 7.56-7.63 (2 H, m), 4.01 (2 H, d, $J=6.0$ Hz), 3.41-3.54 (4 H, m), 1.76-1.96 (4 H, m). HPLC-MS: m/z 370 $[M+H]^+$.



{[3-Hydroxy-5-[3-(2H-tetrazol-5-yl)phenyl]-pyridine-2-carbonyl]amino}-acetic acid: 1H NMR (250 MHz, DMSO- d_6) δ ppm 12.40 (1 H, s), 9.51 (1 H, br s), 8.50 (1 H, s), 8.20 (1 H, s), 7.94 (1 H, d, $J=1.5$ Hz), 7.88-7.99 (1 H, m), 7.72 (1 H, d, $J=1.9$ Hz), 7.34-7.53 (1 H, m), 7.48 (2 H, d, $J=2.2$ Hz), 7.00 (1 H, d, $J=1.8$ Hz), 6.38 (1 H, br s), 3.54 (2 H, br s). HPLC-MS: m/z 341 $[M+H]^+$.

The fourth aspect of Category I of the present disclosure encompasses compounds having the formula:




29

wherein R units are substituted or unsubstituted heteroaryl. Non-limiting examples of these R units are described in Table IV herein below.


TABLE IV

No.	R
115	pyridin-2-yl
116	pyridin-3-yl
117	pyridin-4-yl
118	pyrimidin-2-yl
119	pyrimidin-4-yl
120	pyrimidin-5-yl
121	isoquinolin-1-yl
122	isoquinolin-3-yl
123	isoquinolin-4-yl
124	thiazol-2-yl
125	thiazol-4-yl
126	1,2,3,4-tetrazol-5-yl
127	[1,2,4]triazol-5-yl
128	imidazol-2-yl
129	furan-2-yl
130	furan-3-yl
131	thiophene-2-yl
132	thiophene-3-yl

The compounds which encompass the fourth aspect of Category I of the present disclosure can be prepared by the procedure outlined in Scheme II and described in Example 2 herein beginning with compounds which are members of the first and second aspect of Category I. The following are non-limiting examples of compound encompassed by the fourth aspect of Category I.




[(5'-Hydroxy-[2,3']bipyridinyl-6'-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.45 (1 H, s), 9.45 (1 H, t, $J=6.1$ Hz), 8.91 (1 H, d, $J=1.8$ Hz), 8.73-8.79 (1 H, m), 8.20 (1 H, d, $J=8.0$ Hz), 8.07 (1 H, d, $J=1.8$ Hz), 8.01 (1 H, dt, $J=7.8, 1.8$ Hz), 7.52 (1 H, ddd, $J=7.5, 4.8, 0.9$ Hz), 4.01 (2 H, d, $J=6.1$ Hz). HPLC-MS: m/z 274 $[\text{M}+\text{H}]^+$.




[(5-Hydroxy-[3,3']bipyridinyl-6-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.44 (1 H, br s), 9.46 (1 H, t, $J=6.1$ Hz), 9.30 (1 H, s), 8.86 (1 H, d, $J=5.3$ Hz), 8.76 (1 H, d, $J=8.2$ Hz), 8.67 (1 H, d, $J=1.9$ Hz), 7.92-8.00 (2 H, m), 4.02 (2 H, d, $J=6.1$ Hz). HPLC-MS: m/z 274 $[\text{M}+\text{H}]^+$.


30




[(3-Hydroxy-5-pyrimidin-5-yl-pyridine-2-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.45 (1 H, br s), 9.45 (1 H, t, $J=5.9$ Hz), 9.27-9.33 (3 H, m), 8.67 (1 H, d, $J=1.8$ Hz), 7.97 (1 H, d, $J=1.9$ Hz), 4.02 (2 H, d, $J=6.2$ Hz). HPLC-MS: m/z 275 $[\text{M}+\text{H}]^+$.



[(3-Hydroxy-5-isoquinolin-4-yl-pyridine-2-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.53 (1 H, br s), 9.83 (1 H, s), 9.52 (1 H, t, $J=6.1$ Hz), 8.71 (1 H, s), 8.54 (1 H, d, $J=8.1$ Hz), 8.38 (1 H, d, $J=1.7$ Hz), 7.92-8.13 (3 H, m), 7.73 (1 H, d, $J=1.7$ Hz), 4.04 (2 H, d, $J=6.1$ Hz). HPLC-MS: m/z 324 $[\text{M}+\text{H}]^+$.



[(3-Hydroxy-5-thiazol-2-yl-pyridine-2-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.50 (1 H, s), 9.46 (1 H, t, $J=6.1$ Hz), 8.76 (1 H, d, $J=1.8$ Hz), 8.07 (1 H, d, $J=3.2$ Hz), 8.00 (1 H, d, $J=3.2$ Hz), 7.90 (1 H, d, $J=1.8$ Hz), 4.00 (2 H, d, $J=6.1$ Hz). HPLC-MS: m/z 280 $[\text{M}+\text{H}]^+$.




[[5-(2,3-Dihydro-benzo[1,4]dioxin-6-yl)-3-hydroxy-pyridine-2-carbonyl]-amino]-acetic acid: $^1\text{H NMR}$ (250 MHz, DMSO-d_6) δ ppm 12.83 (1 H, br s), 12.32 (1 H, s), 9.31 (1 H, t, $J=6.1$ Hz), 8.46 (1 H, d, $J=1.9$ Hz), 7.64 (1 H, d, $J=1.9$ Hz), 7.37 (1 H, d, $J=2.2$ Hz), 7.28-7.34 (1 H, m),


31

6.98 (1 H, d, J=8.5 Hz), 4.29 (4 H, s), 3.99 (2 H, d, J=6.1 Hz). HPLC-MS: m/z 331 [M+H]⁺.

Category II of the present disclosure relates to compounds having the formula:



wherein the first aspect encompasses compounds having the formula:




R units are substituted or unsubstituted phenyl. Non-limiting examples of these R units are described in Table V herein below.

TABLE V


No.	R
133	2-fluorophenyl
134	3-fluorophenyl
135	4-fluorophenyl
136	2-chlorophenyl
137	3-chlorophenyl
138	4-chlorophenyl
139	2-cyanophenyl
140	3-cyanophenyl
141	4-cyanophenyl
142	2-methylphenyl
143	3-methylphenyl
144	4-methylphenyl
145	2-ethylphenyl
146	3-ethylphenyl
147	4-ethylphenyl
148	2-methoxyphenyl
149	3-methoxyphenyl
150	4-methoxyphenyl
151	2-ethoxyphenyl
152	3-ethoxyphenyl
153	4-ethoxyphenyl
154	2-iso-propoxyphenyl
155	3-iso-propoxyphenyl
156	4-iso-propoxyphenyl
157	2-carbamoylphenyl
158	3-carbamoylphenyl
158	4-carbamoylphenyl
160	2-(aziridine-1-carbonyl)phenyl
161	3-(aziridine-1-carbonyl)phenyl
162	4-(aziridine-1-carbonyl)phenyl
163	2-(azetidine-1-carbonyl)phenyl
164	3-(azetidine-1-carbonyl)phenyl
165	4-(azetidine-1-carbonyl)phenyl
166	2-(pyrrolidine-1-carbonyl)phenyl
167	3-(pyrrolidine-1-carbonyl)phenyl
168	4-(pyrrolidine-1-carbonyl)phenyl
169	2-(piperidine-1-carbonyl)phenyl
170	3-(piperidine-1-carbonyl)phenyl
171	4-(piperidine-1-carbonyl)phenyl
172	2-(acetylamino)phenyl
173	3-(acetylamino)phenyl
174	4-(acetylamino)phenyl
175	2-(ethanecarbonylamino)phenyl
176	3-(ethanecarbonylamino)phenyl
177	4-(ethanecarbonylamino)phenyl
178	2-(cyclopropanecarbonylamino)phenyl
179	3-(cyclopropanecarbonylamino)phenyl
180	4-(cyclopropanecarbonylamino)phenyl

32


The compounds that encompass the first aspect of Category II of the present disclosure can be prepared by the procedure outlined in Scheme V and described in Example 3 herein below.




Reagents and conditions: (a) 4-chlorophenyl boronic acid, Pd(dppf)Cl₂, K₃PO₄, dioxane/MeOH; 80° C., 3 hr



Reagents and conditions: (b) 4-chlorophenyl boronic acid, Pd(dppf)Cl₂, K₃PO₄,





Reagents and conditions: (c) glycine ethyl ester
HCl, EDCI, HOBT, DIPEA, DMF, CH₂Cl₂; rt, 16
hr

EXAMPLE 3

[(4'-Chloro-3-methoxy-biphenyl-4-carbonyl)-
amino]-acetic acid ethyl ester (10)

Preparation of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid methyl ester (8): To a degassed solution of methyl 4-bromo-2-methoxybenzoate (0.70 g, 2.86 mmol) in 1,4-dioxane (10 mL) and MeOH (2.5 mL) at room temperature under nitrogen blanketing is added 4-chlorophenyl boronic acid (0.536 g, 3.43 mmol), Pd(dppf)Cl₂ (0.233 g, 0.286 mmol) and K₃PO₄ (0.728 g, 3.43 mmol). The resulting suspension is heated to 80° C. and stirred for 3 hours after which the reaction is cooled to room temperature and filtered through Celite™. The collected solids are washed with additional MeOH and the filtrate concentrated under reduced pressure. The crude material is purified over silica (hexanes: EtOAc gradient 6:1 to 4:1) to afford 0.614 g (78% yield) of the desired product as orange crystals. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.89 (1 H, d, J=8.0 Hz), 7.52-7.56 (2 H, m), 7.44 (2 H, d, J=8.7 Hz), 7.17 (1 H, d, J=8.0 Hz), 7.12 (1 H, d, J=1.6 Hz), 3.99 (3 H, s), 3.92 (3 H, s). HPLC-MS: m/z 277 [M+H]⁺.


Preparation of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid (9): To a solution of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid methyl ester, 8, (0.615 g, 2.22 mmol) in THF (20 mL) and H₂O (5 mL) at room temperature is added LiOH (0.932 g, 22.2 mmol). The resulting suspension is heated to reflux for 2 hours. The reaction is cooled and concentrated under reduced pressure. The crude product is acidified using conc. HCl and the resulting solid is collected by filtration washed with H₂O and dried to afford 0.532 g (91%) of the desired product as a grey solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 10.69 (1 H, br s), 8.26 (1 H, d, J=8.1 Hz), 7.53-7.58 (2 H, m), 7.44-7.50 (2 H, m), 7.33 (1 H, dd, J=8.1, 1.6 Hz), 7.20 (1 H, d, J=1.3 Hz), 4.17 (3 H, s). HPLC-MS: m/z 263 [M+H]⁺.

Preparation of [(4'-chloro-3-methoxy-biphenyl-4-carbonyl)-amino]-acetic acid ethyl ester (10): To a solution of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid, 9, (0.325 g, 1.24 mmol) in CH₂Cl₂ (5 mL) and DMF (1.5 mL) at room temperature under N₂ is added glycine ethyl ester hydrochloride (0.19 g, 1.36 mmol), 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide (EDCI) (0.261 g, 1.36 mmol), 1-hydroxybenzotriazole (HOBT) (0.033 g, 0.248 mmol) and diisopropylethylamine (DIPEA) (0.432 mL, 2.28 mmol). The resulting suspension is stirred for 16 hours after which the reaction mixture is diluted with EtOAc and washed with 1M HCl, 1M NaOH and saturated aqueous NaCl. The organic phase is separated, dried (MgSO₄), filtered and concentrated under reduced pressure. The crude material is purified over silica (hexanes:EtOAc 1:1) to afford 0.364 g (85% yield) of the desired product as a colorless solid. ¹H NMR (400 MHz,

34


CDCl₃) δ ppm 8.51 (1 H, br s), 8.28 (1 H, d, J=8.1 Hz), 7.53-7.57 (2 H, m), 7.42-7.46 (2 H, m), 7.27 (1 H, dd, J=8.1, 1.6 Hz), 7.14 (1 H, d, J=1.5 Hz), 4.29 (2 H, d, J=4.8 Hz), 4.28 (2 H, q, J=7.1 Hz), 4.09 (3 H, s), 1.34 (3 H, t, J=7.2 Hz).
HPLC-MS: m/z 348 [M+H]⁺.

A further non-limiting example includes the following.



[(3-Methoxy-4'-methyl-biphenyl-4-carbonyl)-amino]-acetic acid ethyl ester: ¹H NMR (400 MHz, CDCl₃) δ ppm 8.52 (br s, 1 H), 8.26 (d, J=8.14 Hz, 1 H), 7.53 (d, J=8.05 Hz, 2 H), 7.28-7.33 (m, 3 H), 7.17 (d, J=1.37 Hz, 1 H), 4.13-4.45 (m, 4 H), 4.08 (s, 3 H), 2.42 (s, 3 H), 1.33 (t, J=7.18 Hz, 3 H). HPLC-MS: m/z 328 [M+H]⁺.

The compounds of Formula (I) also encompasses compounds having the formula:



wherein R units are substituted or unsubstituted phenyl. Non-limiting examples of these R units are described in Table VI herein below.

TABLE VI


No.	R
181	2-fluorophenyl
182	3-fluorophenyl
183	4-fluorophenyl
184	2-chlorophenyl
185	3-chlorophenyl
186	4-chlorophenyl
187	2-cyanophenyl
188	3-cyanophenyl
189	4-cyanophenyl
190	2-methylphenyl
191	3-methylphenyl
192	4-methylphenyl
193	2-ethylphenyl
194	3-ethylphenyl
195	4-ethylphenyl
196	2-methoxyphenyl
197	3-methoxyphenyl
198	4-methoxyphenyl
199	2-ethoxyphenyl
200	3-ethoxyphenyl
201	4-ethoxyphenyl
202	2-iso-propoxyphenyl
203	3-iso-propoxyphenyl
204	4-iso-propoxyphenyl
205	2-carbamoylphenyl
206	3-carbamoylphenyl
207	4-carbamoylphenyl
208	2-(aziridine-1-carbonyl)phenyl
209	3-(aziridine-1-carbonyl)phenyl
210	4-(aziridine-1-carbonyl)phenyl
211	2-(azetidone-1-carbonyl)phenyl
212	3-(azetidone-1-carbonyl)phenyl
213	4-(azetidone-1-carbonyl)phenyl

TABLE VI-continued

No.	R
214	2-(pyrrolidine-1-carbonyl)phenyl
215	3-(pyrrolidine-1-carbonyl)phenyl
716	4-(pyrrolidine-1-carbonyl)phenyl
217	2-(piperidine-1-carbonyl)phenyl
218	3-(piperidine-1-carbonyl)phenyl
219	4-(piperidine-1-carbonyl)phenyl
220	2-(acetylamino)phenyl
221	3-(acetylamino)phenyl
222	4-(acetylamino)phenyl
223	2-(ethanecarbonylamino)phenyl
224	3-(ethanecarbonylamino)phenyl
225	4-(ethanecarbonylamino)phenyl
226	2-(cyclopropanecarbonylamino)phenyl
227	3-(cyclopropanecarbonylamino)phenyl
226	4-(cyclopropanecarbonylamino)phenyl

The compounds disclosed immediately above can be prepared by the procedure outlined in Scheme VI and described in Example 4 herein below.

Scheme VI




Reagents and conditions: (a) BBr_3 , CH_2Cl_2 ; rt, 3 days

EXAMPLE 4

[(4'-Chloro-3-hydroxy-biphenyl-4-carbonyl)-amino]-acetic acid (11)


Preparation of [(4'-chloro-3-hydroxy-biphenyl-4-carbonyl)-amino]-acetic acid (11): To a solution of [(4'-chloro-3-methoxy-biphenyl-4-carbonyl)-amino]-acetic acid ethyl ester (0.053 g, 0.152 mmol) in CH_2Cl_2 (2 mL) at room temperature under nitrogen is added BBr_3 (1.52 ml of a 1M solution in CH_2Cl_2 , 1.52 mmol) dropwise. The resulting mixture is stirred for 3 days after which time the reaction is quenched with H_2O (0.5 mL) then acidified to pH 1 with conc. HCl. The mixture is extracted with EtOAc ($\times 2$), the organic phase separated, dried (MgSO_4), filtered and concentrated under reduced pressure. The crude material is purified by preparative HPLC to afford 0.019 g (41% yield) of the desired product as a white solid. $^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ ppm 12.41 (1 H, s), 9.19 (1 H, s), 7.96 (1 H, d, $J=8.3$ Hz), 7.74 (2 H, d, $J=8.7$ Hz), 7.54 (2 H, d, $J=8.7$ Hz), 7.21 (1 H, d, $J=1.7$ Hz), 7.26 (1 H, dd, $J=8.3, 1.8$ Hz), 3.99 (2 H, d, $J=5.5$ Hz). HPLC-MS: m/z 306 $[\text{M}+\text{H}]^+$.

The following is a non-limiting example of the second aspect of Category II of the present disclosure.



[(3-Hydroxy-4'-methyl-biphenyl-4-carbonyl)-amino]-acetic acid: $^1\text{H NMR}$ (400 MHz, DMSO-d_6) δ ppm 12.37 (1 H, s), 9.17 (1 H, s), 7.94 (1 H, d, $J=8.32$ Hz), 7.60 (2 H, d, $J=8.14$ Hz), 7.29 (2 H, d, $J=7.96$ Hz), 7.22 (1 H, dd, $J=8.32, 1.83$ Hz), 7.18 (1 H, d, $J=1.74$ Hz), 4.00 (2 H, d, $J=5.67$ Hz), 2.35 (3 H, s). HPLC-MS: m/z 286 $[\text{M}+\text{H}]^+$.

Category III of the present disclosure relates to compounds having the formula:



wherein non-limiting examples of R, R^{6a} and R^{6b} are further described herein below in Table VII.


TABLE VII

No.	R	R^{6a}	R^{6b}
229	2-fluorophenyl	—H	—H
230	3-fluorophenyl	—H	—H
231	4-fluorophenyl	—H	—H
232	2-chlorophenyl	—H	—H
233	3-chlorophenyl	—H	—H
234	4-chlorophenyl	—H	—H
235	2-methylphenyl	—H	—H
236	3-methylphenyl	—H	—H
237	4-methylphenyl	—H	—H
238	2-fluorophenyl	— CH_3	—H
239	3-fluorophenyl	— CH_3	—H
240	4-fluorophenyl	— CH_3	—H
241	2-chlorophenyl	— CH_3	—H
242	3-chlorophenyl	— CH_3	—H
243	4-chlorophenyl	— CH_3	—H
244	2-methylphenyl	— CH_3	—H
245	3-methylphenyl	— CH_3	—H
246	4-methylphenyl	— CH_3	—H
247	2-fluorophenyl	— CH_3	— CH_3
248	3-fluorophenyl	— CH_3	— CH_3
249	4-fluorophenyl	— CH_3	— CH_3
250	2-chlorophenyl	— CH_3	— CH_3
251	3-chlorophenyl	— CH_3	— CH_3
252	4-chlorophenyl	— CH_3	— CH_3
253	2-methylphenyl	— CH_3	— CH_3
254	3-methylphenyl	— CH_3	— CH_3
255	4-methylphenyl	— CH_3	— CH_3
256	2-fluorophenyl	— CH_2CH_3	—H
257	3-fluorophenyl	— CH_2CH_3	—H
258	4-fluorophenyl	— CH_2CH_3	—H
259	2-chlorophenyl	— CH_2CH_3	—H
260	3-chlorophenyl	— CH_2CH_3	—H
261	4-chlorophenyl	— CH_2CH_3	—H
262	2-methylphenyl	— CH_2CH_3	—H
263	3-methylphenyl	— CH_2CH_3	—H
264	4-methylphenyl	— CH_2CH_3	—H

37

The compounds which are encompassed within Category III of the present disclosure can be prepared by the procedures outlined herein below in Schemes VII and VIII and described in Examples 5 and 6.

Scheme VII



Reagents and conditions: (a)


EXAMPLE 5

5-(3-Chlorophenyl)-N-(2-dimethylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide (12)

Preparation of 5-(3-chlorophenyl)-N-(2-dimethylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide (12): To a solution of {[5-(3-chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid, 7, (0.043 g, 0.139 mmol) in DMF (2 mL) at 0° C. under N₂ is added diisopropylethylamine (0.072 mL, 0.42 mmol), 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide (EDCI) (0.040 g, 0.21 mmol) and 1-hydroxybenzotriazole (HOBt) (0.002 g, 0.014 mmol). The resulting mixture is stirred for 5 minutes before dimethylamine (0.10 mL of a 2M solution in THF, 0.21 mmol) is added. The reaction is warmed slowly to room temperature and stirred for 3 days. The reaction is diluted with EtOAc, washed with H₂O, and saturated aqueous NaCl. The organic phase is dried (MgSO₄), filtered, concentrated under reduced pressure and the crude material is purified over silica (EtOAc) to afford 0.20 g (43% yield) of the desired product as a colorless solid. ¹H NMR (250 MHz, CDCl₃) δ ppm 11.90 (1 H, s), 8.83 (1 H, t, J=4.6 Hz), 8.25 (1 H, d, J=1.8 Hz), 7.51 (1 H, m), 7.31-7.44 (4 H, m), 4.19 (2 H, d, J=4.6 Hz), 2.99 (3 H, s), 2.98 (3 H, s). HPLC-MS: m/z 334 [M+H]⁺.


38

Scheme VIII




Reagents and conditions: (a) 2-aminoacetamide
HCl, EDCI, HOBt, DMF; 0° C. to rt, 3 days


Reagents and conditions: (b) H₂; Pd/C, MeOH, rt,
22 hr



Reagents and conditions: (c) (CF₃SO₂)₂NC₆H₅,
MeOH; rt, 24 hr



39



Reagents and conditions: (d) 3-chlorophenylboronic acid, Pd(dppf)Cl₂, K₃PO₄, dioxane; 85° C., 48 hr

EXAMPLE 6

5-(3-Chlorophenyl)-N-(2-amino-2-oxoethyl)-3-hydroxypyridin-2-yl amide (16)

Preparation of 3,5-bis-benzyloxy-N-(2-amino-2-oxoethyl)pyridin-2-yl amide (13): To a solution of 3,5-bis-benzyloxy-pyridine-2-carboxylic acid, 2, (1.00 g, 2.99 mmol) in DMF (20 mL) at room temperature under N₂ is added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDCI) (0.925 g, 5.97 mmol) and 1-hydroxybenzotriazole (HOBt) (0.806 g, 5.97 mmol). The resulting solution is stirred for 15 minutes then 2-aminoacetamide hydrochloride (0.66 g, 5.97 mmol) and diisopropylethylamine (1.56 mL, 8.96 mmol) are added. After 3 days the reaction mixture is concentrated under reduced pressure and H₂O added. The solid which forms is collected by filtration and is washed with water to afford 0.598 g (51% yield) of the desired product as a white solid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 8.39 (1 H, t, J=5.6 Hz), 8.01 (1 H, d, J=2.2 Hz), 7.28-7.56 (12 H, m), 7.11 (1 H, br s), 5.26 (2 H, s), 5.25 (2 H, s), 3.81 (2 H, d, J=5.6 Hz). HPLC-MS: m/z 392 [M+H]⁺.

Preparation of 3,5-dihydroxy-N-(2-amino-2-oxoethyl)pyridin-2-yl amide (14): A solution of 3,5-bis-benzyloxy-N-(2-amino-2-oxoethyl)pyridin-2-yl amide, 13, (0.598 g, 1.53 mmol) in EtOH (100 mL) containing 10% Pd/C (0.120 g) is stirred under an atmosphere of H₂ for 22 hours. The reaction solution is filtered through Celite™ and the collected solids are washed with hot MeOH. The combined filtrate and washings are concentrated under reduced pressure to afford 0.32 g (99% yield) of the desired product cream-colored solid. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 12.42 (1 H, s), 10.84 (1 H, br s), 8.75 (1 H, t, J=5.8 Hz), 7.75 (1 H, d, J=2.3 Hz), 7.47 (1 H, br s), 7.13 (1 H, br s), 6.67 (1 H, d, J=2.3 Hz), 3.85 (2 H, d, J=5.9 Hz). HPLC-MS: m/z 212 [M+H]⁺.


Preparation of 5-trifluoromethanesulfonyloxy-N-(2-amino-2-oxoethyl)-3-hydroxypyridin-2-yl amide (15): To a solution of 3,5-dihydroxy-N-(2-amino-2-oxoethyl)pyridin-2-yl amide, 14, (0.30 g, 1.42 mmol) in MeOH (10 mL) and DMF (5 mL) at 0° C. under N₂ is added diisopropylethylamine (0.247 mL, 1.42 mmol) followed by N-phenyltrifluoromethanesulfonamide (0.508 g, 1.42 mmol). The resulting mixture is warmed slowly to room temperature and stirring is continued for 24 hours. The solvent is then removed under reduced pressure and the crude material is purified over silica (2% MeOH:CH₂Cl₂) to afford 0.404 g (83% yield) of the desired product as a pale yellow solid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.85 (1 H, br s), 9.28 (1 H, t, J=5.9 Hz), 8.41 (1 H, d, J=2.3 Hz), 7.85 (1 H, d, J=2.4 Hz), 7.52

40


(1 H, br s), 7.18 (1 H, br s), 3.88 (2 H, d, J=6.1 Hz). HPLC-MS: m/z 344 [M+H]⁺.

Preparation of 5-(3-chlorophenyl)-N-(2-amino-2-oxoethyl)-3-hydroxypyridin-2-yl amide (16): To a degassed solution of 5-trifluoromethanesulfonyloxy-N-(2-amino-2-oxoethyl)-3-hydroxypyridin-2-yl amide, 15, (0.20 g, 0.58 mmol) in 1,4-dioxane (3.5 mL) at room temperature under N₂ is added 3-chlorophenyl boronic acid (0.109 g, 0.70 mmol), K₃PO₄ (0.148 g, 0.70 mmol) and Pd(dppf)Cl₂ (0.048 g, 0.06 mmol). The resulting suspension is heated to 90° C. in a sealed tube for 22 hours. The reaction is cooled to room temperature and additional 3-chlorophenyl boronic acid (0.055 g, 0.35 mmol) and Pd(dppf)Cl₂ (0.048 g, 0.06 mmol) is added and the reaction reheated to 90° C. for an additional 22 hours. After cooling, the reaction solution is filtered through Celite™ and the collected solids are washed with additional MeOH. The filtrate and washings are concentrated under reduced pressure and the residue dissolved in CH₂Cl₂ and washed with 10% citric acid. The organic layer is dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product is purified over silica (2% MeOH:CH₂Cl₂) to afford 0.033 g (18% yield) of the desired product as a colourless solid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.46 (1 H, s), 9.17 (1 H, t, J=5.9 Hz), 8.55 (1 H, d, J=2.0 Hz), 7.93 (1 H, d, J=0.9 Hz), 7.75-7.84 (2 H, m), 7.49-7.60 (3 H, m), 7.18 (1 H, s), 3.91 (2 H, d, J=5.9 Hz). HPLC-MS: m/z 306 [M+H]⁺.

The following are non-limiting examples of compounds encompassed within Category III of the present disclosure.




5-(3-Chlorophenyl)-N-(2-isopropylamino-2-oxoethyl)-3-hydroxy-pyridin-2-yl amide: ¹H NMR (250 MHz, CDCl₃) δ ppm 8.60 (1 H, t, J=6.4 Hz), 8.25 (1 H, d, J=2.0 Hz), 7.50-7.53 (1 H, m), 7.44 (1 H, d, J=2.0 Hz), 7.35-7.43 (3 H, m), 5.69 (1 H, br s), 4.04 (2 H, d, J=5.9 Hz), 3.41 (1 H, q, J=7.0 Hz), 1.13 (6 H, d, J=6.6 Hz). HPLC-MS: m/z 348 [M+H]⁺.




5-(4-Methylphenyl)-N-(2-methylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide: ¹H NMR (250 MHz, CDCl₃) δ ppm 11.90 (1 H, s), 8.47 (1 H, t, J=5.9 Hz), 8.25 (1 H, d, J=1.8 Hz), 7.33-7.56 (3 H, m), 7.12-7.31 (2 H, m), 6.07 (1 H, br s), 3.90-4.24 (2 H, m), 2.66-2.98 (3 H, m), 2.35 (3 H, s). HPLC-MS: m/z 300 [M+H]⁺.

41




5-(3-Chlorophenyl)-N-(2-methylamino-2-oxoethyl)-3-hydroxy-pyridin-2-yl amide: $^1\text{H NMR}$ (250 MHz, CDCl_3) δ ppm 11.79 (1 H, br s), 8.55 (1 H, br s), 8.31 (1 H, s), 7.60 (1 H, s), 7.41-7.53 (4 H, m), 6.01 (1 H, br s), 4.15 (2 H, d, $J=5.8$ Hz), 2.90 (3 H, d, $J=4.7$ Hz). HPLC-MS: m/z 320 $[\text{M}+\text{H}]^+$.

Category IV of the present disclosure relates to compounds having the formula:



wherein the first aspect relates to compounds having the formula:



non-limiting examples of R, R^5 , and L units are further described herein below in Table VIII.

TABLE VIII

No.	R^5	R	L
265	—H	2-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
266	—H	3-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
267	—H	4-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
268	—H	2-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
269	—H	3-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
270	—H	4-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
271	—H	2-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
272	—H	3-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
273	—H	4-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
274	— CH_3	2-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
275	— CH_3	3-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
276	— CH_3	4-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
277	— CH_3	2-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
278	— CH_3	3-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
279	— CH_3	4-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
280	— CH_3	2-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
281	— CH_3	3-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
282	— CH_3	4-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
283	— CH_2CH_3	2-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
284	— CH_2CH_3	3-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
255	— CH_2CH_3	4-fluorophenyl	— $\text{C}(\text{CH}_3)_2$ —
286	— CH_2CH_3	2-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
287	— CH_2CH_3	3-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
288	— CH_2CH_3	4-chlorophenyl	— $\text{C}(\text{CH}_3)_2$ —
289	— CH_2CH_3	2-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
290	— CH_2CH_3	3-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
291	— CH_2CH_3	4-methylphenyl	— $\text{C}(\text{CH}_3)_2$ —
292	—H	2-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
293	—H	3-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
294	—H	4-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —


42

TABLE VIII-continued


No.	R^5	R	L	
5	295	—H	2-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	296	—H	3-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	297	—H	4-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	298	—H	2-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	299	—H	3-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	300	—H	4-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
10	301	— CH_3	2-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
	302	— CH_3	3-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
	303	— CH_3	4-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
	304	— CH_3	2-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	305	— CH_3	3-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	306	— CH_3	4-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
15	307	— CH_3	2-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	308	— CH_3	3-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	309	— CH_3	4-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	310	— CH_2CH_3	2-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
	311	— CH_2CH_3	3-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
	312	— CH_2CH_3	4-fluorophenyl	— $\text{CH}(\text{CH}_3)$ —
20	313	— CH_2CH_3	2-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	314	— CH_2CH_3	3-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	315	— CH_2CH_3	4-chlorophenyl	— $\text{CH}(\text{CH}_3)$ —
	316	— CH_2CH_3	2-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	317	— CH_2CH_3	3-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
25	318	— CH_2CH_3	4-methylphenyl	— $\text{CH}(\text{CH}_3)$ —
	319	—H	2-fluorophenyl	— CH_2CH_2 —
	320	—H	3-fluorophenyl	— CH_2CH_2 —
	321	—H	4-fluorophenyl	— CH_2CH_2 —
	322	—H	2-chlorophenyl	— CH_2CH_2 —
	323	—H	3-chlorophenyl	— CH_2CH_2 —
	324	—H	4-chlorophenyl	— CH_2CH_2 —
30	325	—H	2-methylphenyl	— CH_2CH_2 —
	326	—H	3-methylphenyl	— CH_2CH_2 —
	327	—H	4-methylphenyl	— CH_2CH_2 —
	328	— CH_3	2-fluorophenyl	— CH_2CH_2 —
35	329	— CH_3	3-fluorophenyl	— CH_2CH_2 —
	330	— CH_3	4-fluorophenyl	— CH_2CH_2 —
	331	— CH_3	2-chlorophenyl	— CH_2CH_2 —
	332	— CH_3	3-chlorophenyl	— CH_2CH_2 —
	333	— CH_3	4-chlorophenyl	— CH_2CH_2 —
	334	— CH_3	2-methylphenyl	— CH_2CH_2 —
40	335	— CH_3	3-methylphenyl	— CH_2CH_2 —
	336	— CH_3	4-methylphenyl	— CH_2CH_2 —
	337	— CH_2CH_3	2-fluorophenyl	— CH_2CH_2 —
	338	— CH_2CH_3	3-fluorophenyl	— CH_2CH_2 —
	339	— CH_2CH_3	4-fluorophenyl	— CH_2CH_2 —
45	340	— CH_2CH_3	2-chlorophenyl	— CH_2CH_2 —
	341	— CH_2CH_3	3-chlorophenyl	— CH_2CH_2 —
	342	— CH_2CH_3	4-chlorophenyl	— CH_2CH_2 —
	343	— CH_2CH_3	2-methylphenyl	— CH_2CH_2 —
	344	— CH_2CH_3	3-methylphenyl	— CH_2CH_2 —
	345	— CH_2CH_3	4-methylphenyl	— CH_2CH_2 —

The compounds described immediately above wherein R^5 is C_1 - C_4 linear, branched, or cyclic alkyl can be prepared by the procedures outlined herein below in Scheme IX and described in Example 7.


Scheme IX




43




Reagents and conditions: (a)
2-amino-2-methylpropionic acid methyl ester,
EDCI, HOBt, DMF; 0° C. to rt, 3 days




Reagents and conditions: (b) H₂; Pd/C, MeOH, rt,
22 hr



Reagents and conditions: (c) (CF₃SO₂)₂NC₆H₅,
MeOH; rt, 24 hr



44



15 Reagents and conditions: (d) 3-chlorophenylboronic
acid, Pd(dppf)Cl₂, K₃PO₄, dioxane; 85° C., 48 hr

EXAMPLE 7

20 2-[[5-(3-Chloro-phenyl)-3-hydroxy-pyridine-2-car-
bonyl]-amino]-2-methyl-propionic acid methyl
ester (20)

25 Preparation of 2-[(3,5-bis-benzyloxy-pyridine-2-carbo-
nyl)-amino]-2-methyl-propionic acid methyl ester (17): To a
solution of 3,5-bis-benzyloxy-pyridine-2-carboxylic acid, 2,
(1.0 g, 2.99 mmol) in DMF (20 mL) at room temperature
under N₂ is added 1-(3-dimethylaminopropyl)-3-ethylcarbo-
diimide (EDCI) (0.925 g, 5.97 mmol) and 1-hydroxybenzo-
triazole (HOBt) (0.806 g, 5.97 mmol). The mixture is
stirred for 15 minutes after which α-aminoisobutyric acid
(1.56 mL, 5.97 mmol) and diisopropylethyl-amine (DIPEA)
(1.56 mL, 8.96 mmol) are added. The resulting solution is
stirred at room temperature for 16 hours then concentrated
under reduced pressure. The resulting brown oil is purified
over silica (EtOAc:heptane 1:1) to afford 0.58 g (45% yield)
of the desired compound as a colorless solid. ¹H NMR (250
MHz, CDCl₃) δ ppm 8.13 (1 H, s), 8.02 (1 H, d, J=2.3 Hz),
7.31-7.48 (10 H, m), 6.90 (1 H, d, J=2.3 Hz), 5.19 (2 H, s),
5.10 (2 H, s), 3.74 (3 H, s), 1.58 (6 H, s). HPLC-MS: m/z
435 [M+H]⁺.

Preparation of 2-[(3,5-dihydroxy-pyridine-2-carbonyl)-
amino]-2-methyl-propionic acid methyl ester (18): A solu-
tion of 2-[(3,5-bis-benzyloxy-pyridine-2-carbonyl)-amino]-
2-methylpropionic acid methyl ester, 17, (0.58 g, 1.34
mmol) in MeOH (100 mL) containing 10% Pd/C (0.116 g)
is stirred under an atmosphere of H₂ for 22 hours. The
reaction solution is filtered through Celite™ and the col-
lected solids are washed with hot MeOH. The combined
filtrate and washings are concentrated under reduced pres-
sure to afford 0.321 g (94% yield) of the desired compound
as a grey solid. ¹H NMR (250 MHz, MeOD) δ ppm 7.67
(1H, d, J=2.4 Hz), 6.58 (1H, d, J=2.4 Hz), 3.69 (3H, s), 1.56
(6H, s). HPLC-MS: m/z 255 [M+H]⁺.

Preparation of 2-[(3-hydroxy-5-trifluoromethanesulfo-
nyloxy-pyridine-2-carbonyl)-amino]-2-methyl-propionic acid
methyl ester (19): To a solution in 2-[(3,5-dihydroxy-pyri-
dine-2-carbonyl)-amino]-2-methyl-propionic acid methyl
ester, 18, (0.312 g, 1.23 mmol) in MeOH (10 mL) containing
diisopropylethylamine (0.214 mL, 1.23 mmol) at 0° C.
under N₂ is added N-phenyltrifluoromethanesulfonamide
(0.439 g, 1.23 mmol). The reaction is warmed slowly to
room temperature and stirred for 40 hours. The solvent is
removed under reduced pressure and the crude oil which
remains is purified over silica (EtOAc:heptane 1:9) to afford
0.170 g (36% yield) of the desired compound as a yellow oil.


45

¹H NMR (250 MHz, MeOD) δ ppm 8.85 (1 H, br s), 8.19 (1 H, d, J=2.4 Hz), 7.46 (1 H, d, J=2.3 Hz), 3.74 (3 H, s), 1.63 (6 H, s). HPLC-MS: m/z 387 [M+H]⁺.

Preparation of 2-[[5-(3-chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino]-2-methyl-propionic acid methyl ester (20): To a degassed solution of 2-[[3-hydroxy-5-trifluoromethanesulfonyloxy-pyridine-2-carbonyl]-amino]-2-methyl-propionic acid methyl ester (0.17 g, 0.44 mmol) in 1,4-dioxane (3 mL) at room temperature under N₂ is added 3-chlorophenyl boronic acid (0.082 g, 0.53 mmol), K₃PO₄ (0.112 g, 0.53 mmol) and Pd(dppf)Cl₂ (0.036 g, 0.04 mmol). The resulting suspension is heated to 85° C. in a sealed tube for 20 hours. After cooling, the reaction solution is filtered through Celite™ and the collected solids are washed with additional MeOH. The filtrate and washings are concentrated under reduced pressure and the residue dissolved in CH₂Cl₂ and washed with 10% citric acid. The organic layer is dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product is purified over silica (EtOAc: heptane 1:4) to afford 0.112 g (73% yield) of the desired compound as a colorless oil. ¹H NMR (250 MHz, CDCl₃) δ ppm 11.83 (1 H, br s), 8.29 (1 H, br s), 8.10 (1 H, d, J=2.0 Hz), 7.40 (1 H, m), 7.08-7.34 (4 H, m), 3.65 (3 H, s), 1.55 (6 H, s). HPLC-MS: m/z 349 [M+H]⁺.

The compounds of Formula (I) wherein R⁵ is hydrogen can be prepared by the procedures outlined herein below in Scheme X and described in Example 8

Scheme X



Reagents and conditions: (a) LiOH, THF/H₂O; rt, 3 days

EXAMPLE 8


2-[[5-(3-Chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino]-2-methyl-propionic acid (21)

Preparation of 2-[[5-(3-chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino]-2-methyl-propionic acid (21): To a solution of 2-[[5-(3-chlorophenyl)-3-hydroxy-pyridine-2-


46

carbonyl]-amino]-2-methyl-propionic acid methyl ester, 20, (0.082 g, 0.24 mmol) in THF (4 mL) is added LiOH (0.024 g, 0.98 mmol) and H₂O (1 mL) and the resulting solution stirred for 3 days at room temperature. The solvent is removed under reduced pressure and the pale yellow solid that remains is then acidified with 1M HCl to until the pH is approximately 1 and the solution extracted twice with EtOAc. The combined organic layers are combined, dried (Na₂SO₄), filtered and concentrated under reduced pressure to afford 0.064 g (81% yield) of the desired compound as a white solid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.99 (1 H, br s), 12.25 (1 H, s), 9.05 (1 H, s), 8.53 (1 H, d, J=2.0 Hz), 7.91 (1 H, s), 7.74-7.83 (2 H, m), 7.51-7.58 (2 H, m), 1.58 (6 H, s). HPLC-MS: m/z 335 [M+H]⁺.


Further non-limiting examples include the following.



3-[[3-Hydroxy-5-(4-methylphenyl)-pyridine-2-carbonyl]-amino]-propionic acid ethyl ester: ¹H NMR (250 MHz, CDCl₃) δ ppm 12.12 (1 H, s), 8.44 (1 H, t, J=5.9 Hz), 8.31 (1 H, d, J=1.8 Hz), 7.44-7.55 (3 H, m), 7.30 (2 H, d, J=7.9 Hz), 4.21 (2 H, q, J=7.2 Hz), 3.76 (2 H, q, J=6.4 Hz), 2.68 (2 H, t, J=6.2 Hz), 2.43 (3 H, s), 1.30 (3 H, t, J=7.2 Hz). HPLC-MS: m/z 329 [M+H]⁺.




3-[[3-Hydroxy-5-(3-chlorophenyl)-pyridine-2-carbonyl]-amino]-propionic acid ethyl ester: ¹H NMR (400 MHz, CDCl₃) δ ppm 12.16 (1 H, s), 8.45 (1 H, t, J=5.7 Hz), 8.25 (1 H, d, J=1.6 Hz), 7.55 (1 H, s), 7.37-7.47 (4 H, m), 4.20 (2 H, q, J=7.1 Hz), 3.75 (2 H, q, J=6.3 Hz), 2.68 (2 H, t, J=6.2 Hz), 1.28 (3 H, t, J=7.1 Hz). HPLC-MS: m/z 349 [M+H]⁺.




3-[[5-(3-Chlorophenyl)-3-hydroxy-pyridine-2-carbonyl]-amino]-propionic acid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.63 (1 H, br s), 12.37 (1 H, br s), 9.20 (1 H, t, J=5.6 Hz), 8.50 (1 H, d, J=1.8 Hz), 7.91 (1 H, s), 7.72-7.84 (2 H,

47

m), 7.54 (2 H, m), 3.54 (2 H, q, J=6.8 Hz), 2.58 (2 H, t, J=6.9 Hz). HPLC-MS: m/z 321 [M+H]⁺.




3-[(3-Hydroxy-5-(4-methylphenyl)-pyridine-2-carbonyl)-amino]-propionic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.59 (1 H, br s), 9.15 (1 H, t, J=5.9 Hz), 8.46 (1 H, d, J=1.8 Hz), 7.70 (2 H, d, J=8.2 Hz), 7.66 (1 H, d, J=2.0 Hz), 7.33 (2 H, d, J=8.1 Hz), 3.54 (2 H, q, J=6.7 Hz), 2.57 (2 H, t, J=7.0 Hz), 2.36 (3 H, s). HPLC-MS: m/z 301 [M+H]⁺.



2-[[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]-amino]-2-methyl-propionic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.99 (1 H, br s), 12.25 (1 H, s), 9.05 (1 H, s), 8.53 (1 H, d, J=2.0 Hz), 7.91 (1 H, s), 7.74-7.83 (2 H, m), 7.51-7.58 (2 H, m), 1.58 (6 H, s). HPLC-MS: m/z 335 [M+H]⁺.

The second aspect of Category IV relates to compounds having the formula:



wherein non-limiting examples of R, R^{6a}, R^{6b}, and L units are further described herein below in Table IX.

TABLE IX

No.	R ^{6a}	R ^{6b}	R	L
346	—H	—H	2-fluorophenyl	—C(CH ₃) ₂ —
347	—H	—H	3-fluorophenyl	—C(CH ₃) ₂ —
348	—H	—H	4-fluorophenyl	—C(CH ₃) ₂ —
349	—H	—H	2-chlorophenyl	—C(CH ₃) ₂ —
350	—H	—H	3-chlorophenyl	—C(CH ₃) ₂ —
351	—H	—H	4-chlorophenyl	—C(CH ₃) ₂ —
352	—H	—H	2-methylphenyl	—C(CH ₃) ₂ —
353	—H	—H	3-methylphenyl	—C(CH ₃) ₂ —
354	—H	—H	4-methylphenyl	—C(CH ₃) ₂ —
355	—CH ₃	—H	2-fluorophenyl	—C(CH ₃) ₂ —
356	—CH ₃	—H	3-fluorophenyl	—C(CH ₃) ₂ —
357	—CH ₃	—H	4-fluorophenyl	—C(CH ₃) ₂ —
358	—CH ₃	—H	2-chlorophenyl	—C(CH ₃) ₂ —
359	—CH ₃	—H	3-chlorophenyl	—C(CH ₃) ₂ —
360	—CH ₃	—H	4-chlorophenyl	—C(CH ₃) ₂ —
361	—CH ₃	—H	2-methylphenyl	—C(CH ₃) ₂ —
362	—CH ₃	—H	3-methylphenyl	—C(CH ₃) ₂ —


48

TABLE IX-continued

No.	R ^{6a}	R ^{6b}	R	L
363	—CH ₃	—H	4-methylphenyl	—C(CH ₃) ₂ —
364	—CH ₃	—CH ₃	2-fluorophenyl	—C(CH ₃) ₂ —
365	—CH ₃	—CH ₃	3-fluorophenyl	—C(CH ₃) ₂ —
366	—CH ₃	—CH ₃	4-fluorophenyl	—C(CH ₃) ₂ —
367	—CH ₃	—CH ₃	2-chlorophenyl	—C(CH ₃) ₂ —
368	—CH ₃	—CH ₃	3-chlorophenyl	—C(CH ₃) ₂ —
369	—CH ₃	—CH ₃	4-chlorophenyl	—C(CH ₃) ₂ —
370	—CH ₃	—CH ₃	2-methylphenyl	—C(CH ₃) ₂ —
371	—CH ₃	—CH ₃	3-methylphenyl	—C(CH ₃) ₂ —
372	—CH ₃	—CH ₃	4-methylphenyl	—C(CH ₃) ₂ —
373	—CH ₂ CH ₃	—H	2-fluorophenyl	—C(CH ₃) ₂ —
374	—CH ₂ CH ₃	—H	3-fluorophenyl	—C(CH ₃) ₂ —
375	—CH ₂ CH ₃	—H	4-fluorophenyl	—C(CH ₃) ₂ —
376	—CH ₂ CH ₃	—H	2-chlorophenyl	—C(CH ₃) ₂ —
377	—CH ₂ CH ₃	—H	3-chlorophenyl	—C(CH ₃) ₂ —
378	—CH ₂ CH ₃	—H	4-chlorophenyl	—C(CH ₃) ₂ —
379	—CH ₂ CH ₃	—H	2-methylphenyl	—C(CH ₃) ₂ —
380	—CH ₂ CH ₃	—H	3-methylphenyl	—C(CH ₃) ₂ —
381	—CH ₂ CH ₃	—H	4-methylphenyl	—C(CH ₃) ₂ —
382	—CH ₂ CH ₃	—CH ₂ CH ₃	2-fluorophenyl	—C(CH ₃) ₂ —
383	—CH ₂ CH ₃	—CH ₂ CH ₃	3-fluorophenyl	—C(CH ₃) ₂ —
384	—CH ₂ CH ₃	—CH ₂ CH ₃	4-fluorophenyl	—C(CH ₃) ₂ —
385	—CH ₂ CH ₃	—CH ₂ CH ₃	2-chlorophenyl	—C(CH ₃) ₂ —
386	—CH ₂ CH ₃	—CH ₂ CH ₃	3-chlorophenyl	—C(CH ₃) ₂ —
387	—CH ₂ CH ₃	—CH ₂ CH ₃	4-chlorophenyl	—C(CH ₃) ₂ —
388	—CH ₂ CH ₃	—CH ₂ CH ₃	2-methylphenyl	—C(CH ₃) ₂ —
389	—CH ₂ CH ₃	—CH ₂ CH ₃	3-methylphenyl	—C(CH ₃) ₂ —
390	—CH ₂ CH ₃	—CH ₂ CH ₃	4-methylphenyl	—C(CH ₃) ₂ —

Another subgenus of the Compounds of Formula (I) can be prepared by the procedures outlined herein below in Scheme XI and described in Example 9.

Scheme XI



Reagents and conditions: (a) CH₃NH₂HCl, EDCl, HOBt, DMF; 0° C. to rt, 2 days

EXAMPLE 9


5-(3-Chlorophenyl)-N-(2-methylamino-2-oxo-1,1-dimethylethyl)-3-hydroxy-pyridin-2-yl amide (22)

65 Preparation of 5-(3-chlorophenyl)-N-(2-methylamino-2-oxo-1,1-dimethylethyl)-3-hydroxy-pyridin-2-yl amide (22): To a solution of 2-[[5-(3-chlorophenyl)-3-hydroxy-pyri-


49

dine-2-carbonyl]-amino}-2-methyl-propionic acid, 21, (0.030 g, 0.09 mmol) in DMF (2 mL) at room temperature under N₂ is added 1-(3-dimethylamino-propyl)-3-ethyl-carbodiimide (EDCI) (0.021 g, 0.13 mmol), 1-hydroxybenzotriazole (HOBT) (0.012 g, 0.09 mmol) and diisopropylethylamine (DIPEA) (0.047 ml, 0.27 mmol). The reaction is stirred for 5 minutes then methylamine hydrochloride (0.09 g, 0.13 mmol) is added. After stirring for 2 days, the solvent is removed under reduced pressure and the residue partitioned between CH₂Cl₂ and H₂O. The organic layer is separated, washed with sat. NaCl, dried (Na₂SO₄), filtered and concentrated under reduced pressure. The crude product is purified over silica (MeOH:CH₂Cl₂ 1:99) to afford 0.025 g (80% yield) of the desired compound as a colorless oil. ¹H NMR (250 MHz, CDCl₃) δ ppm 11.93 (1 H, br s), 8.50 (1 H, s), 8.26 (1H, d, J=1.8 Hz), 7.56 (1 H, d, J=1.4 Hz), 7.38-7.50 (4 H, m), 6.50 (1 H, br s), 2.87 (3 H, d, J=4.7 Hz), 1.71 (6 H, s). HPLC-MS: m/z 348 [M+H]⁺.

Another subgenus of the compounds of Formula (I) relates to compounds having the formula:



which can be exemplified by compounds having the formula:



wherein the R¹ units are substituted or unsubstituted phenyl. Non-limiting examples of these units are described in Table X herein below.

TABLE X

No.	R ¹
391	2-fluorophenyl
392	3-fluorophenyl
393	4-fluorophenyl
394	2-chlorophenyl
395	3-chlorophenyl
396	4-chlorophenyl
397	2-cyanophenyl
398	3-cyanophenyl
399	4-cyanophenyl
400	2-methylphenyl
401	3-methylphenyl
402	4-methylphenyl
403	2-ethylphenyl
404	3-ethylphenyl
405	4-ethylphenyl
406	2-methoxyphenyl
407	3-methoxyphenyl
408	4-methoxyphenyl
409	2-ethoxyphenyl
410	3-ethoxyphenyl
411	4-ethoxyphenyl
412	2-iso-propoxyphenyl
413	3-iso-propoxyphenyl
414	4-iso-propoxyphenyl
415	2-carbamoylphenyl
416	3-carbamoylphenyl
417	4-carbamoylphenyl
418	2-(aziridine-1-carbonyl)phenyl
419	3-(aziridine-1-carbonyl)phenyl
420	4-(aziridine-1-carbonyl)phenyl


50

TABLE X-continued


No.	R ¹
421	2-(azetidine-1-carbonyl)phenyl
422	3-(azetidine-1-carbonyl)phenyl
423	4-(azetidine-1-carbonyl)phenyl
424	2-(pyrrolidine-1-carbonyl)phenyl
425	3-(pyrrolidine-1-carbonyl)phenyl
426	4-(pyrrolidine-1-carbonyl)phenyl
427	2-(piperidine-1-carbonyl)phenyl
428	3-(piperidine-1-carbonyl)phenyl
429	4-(piperidine-1-carbonyl)phenyl
430	2-(acetylamino)phenyl
431	3-(acetylamino)phenyl
432	4-(acetylamino)phenyl
433	2-(ethanecarbonylamino)phenyl
434	3-(ethanecarbonylamino)phenyl
435	4-(ethanecarbonylamino)phenyl
436	2-(cyclopropanecarbonylamino)phenyl
437	3-(cyclopropanecarbonylamino)phenyl
438	4-(cyclopropanecarbonylamino)phenyl

Another subgenus of the compounds of Formula (I) can be prepared by the procedure outlined in Scheme XII and described in Example 10 herein below.

Scheme XII



Reagents and conditions: (a) GlyOMe.HCl, EDCI, HOBT, CH₂Cl₂; rt, 16 hr



Reagents and conditions: (b) 4-methylphenyl boronic acid, Pd(dppf)Cl₂, K₃PO₄, dioxane/H₂O; 70° C., 16 hr

EXAMPLE 10

[(4-(4-Methylphenyl)pyridine-2-carbonyl)amino]-acetic acid methyl ester (24)


Preparation of [(4-iodo-pyridine-2-carbonyl)amino]-acetic acid methyl ester (23): To a solution of 4-iodo-picolinic

51


acid (1.41 g, 5.66 mmol) in CH_2Cl_2 (35 mL) at room temperature under N_2 is added 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide (EDCI) (1.62 g, 8.49 mmol) and 1-hydroxybenzotriazole (HOBT) (0.077 g, 0.57 mmol). The solution is stirred for 5 minutes and glycine methyl ester hydrochloride (1.07 g, 8.49 mmol) is added and the reaction is stirred 16 hours. The reaction volume is concentrated under reduced pressure and the crude material was partitioned between EtOAc and 1M K_2CO_3 . The aqueous phase is removed and the organic phase washed with H_2O , sat. NaCl, dried (MgSO_4), filtered and concentrated under reduced pressure to afford a brown oil which is purified over silica (EtOAc:heptane gradient 1:4) to afford 0.805 g (44% yield) of the desired product as a colorless solid. HPLC-MS: m/z 321 $[\text{M}+\text{H}]^+$.

Preparation of [[4-(4-methylphenyl)pyridine-2-carbonyl]amino]-acetic acid methyl ester (24): To a degassed solution of [[4-iodo-pyridine-2-carbonyl]-amino]-acetic acid methyl ester, 23, (0.150, 0.47 mmol) in 1,4-dioxane (4 mL) and MeOH (2 mL) is added K_3PO_4 (0.109 mg, 0.52 mmol), $\text{Pd}(\text{dppf})\text{Cl}_2$ (0.038 g, 0.047 mmol) and 4-methylphenyl boronic acid (0.064 g, 0.47 mmol). The reaction is heated to 70°C . in a sealed tube under N_2 for 16 hours. The solvents are then removed under reduced pressure and the solid which remains is partitioned between CH_2Cl_2 and 1M K_2CO_3 . The aqueous phase is removed and the organic phase washed with H_2O , sat. NaCl, dried (MgSO_4), filtered and concentrated under reduced pressure. The crude material is purified over silica (EtOAc:heptane gradient 1:4 to 3:7) to afford 0.113 g (85% yield) of the desired compound. ^1H NMR (400 MHz, CDCl_3) δ ppm 8.60 (1 H, dd, $J=5.1, 0.7$ Hz), 8.55 (1 H, t, $J=4.8$ Hz), 8.43-8.44 (1 H, m), 8.43 (1 H, s), 7.66 (1 H, dd, $J=5.1, 1.8$ Hz), 7.63 (2 H, d, $J=8.4$ Hz), 7.32 (2 H, d, $J=8.1$ Hz), 4.31 (1 H, d, $J=5.9$ Hz), 3.81 (2 H, s), 2.43 (3 H, s). HPLC-MS: m/z 285 $[\text{M}+\text{H}]^+$.

The following are non-limiting examples of compounds described immediately above.




[[4-(4-Cyanophenyl)pyridine-2-carbonyl]amino]-acetic acid methyl ester: ^1H NMR (400 MHz, CDCl_3) δ ppm 8.63 (1 H, d, $J=5.1$ Hz), 8.45 (1 H, t, $J=5.3$ Hz), 8.36 (1 H, d, $J=1.8$ Hz), 7.74 (4 H, s), 7.59 (1 H, dd, $J=5.1, 1.8$ Hz), 4.24 (2 H, d, $J=5.9$ Hz), 3.74 (3 H, s). HPLC-MS: m/z 296 $[\text{M}+\text{H}]^+$.




[[4-(4-Chlorophenyl)pyridine-2-carbonyl]amino]-acetic acid methyl ester: ^1H NMR (400 MHz, CDCl_3) δ ppm 8.56 (1 H, dd, $J=4.9, 0.9$ Hz), 8.47 (1 H, t, $J=5.1$ Hz), 8.33 (1 H,

52

dd, $J=2.0, 0.9$ Hz), 7.51-7.61 (3 H, m), 7.41 (2 H, m), 4.23 (2 H, d, $J=5.5$ Hz), 3.74 (3 H, s). HPLC-MS: m/z 305 $[\text{M}+\text{H}]^+$.




[[4-(3-Chlorophenyl)pyridine-2-carbonyl]amino]-acetic acid methyl ester: ^1H NMR (400 MHz, CDCl_3) δ ppm 8.57 (1 H, dd, $J=5.1, 0.7$ Hz), 8.47 (1 H, t, $J=5.1$ Hz), 8.33 (1 H, m), 7.62 (1 H, m), 7.57 (1 H, dd, $J=5.1, 1.8$ Hz), 7.47-7.54 (1 H, m), 7.31-7.42 (2 H, m), 4.24 (2 H, d, $J=5.9$ Hz), 3.74 (3 H, s). HPLC-MS: m/z 305 $[\text{M}+\text{H}]^+$.



[[4-(2-Chlorophenyl)pyridine-2-carbonyl]-amino]-acetic acid methyl ester: ^1H NMR (400 MHz, CDCl_3) δ ppm 8.57 (1 H, dd, $J=5.1, 0.7$ Hz), 8.46 (1 H, t, $J=4.6$ Hz), 8.20 (1 H, dd, $J=1.8, 0.7$ Hz), 7.50 (1 H, dd, $J=5.1, 1.8$ Hz), 7.42-7.46 (1 H, m), 7.30 (1 H, d, $J=1.8$ Hz), 7.28-7.32 (2 H, m), 4.23 (2 H, d, $J=5.5$ Hz), 3.74 (3 H, s). HPLC-MS: m/z 305 $[\text{M}+\text{H}]^+$.

The second aspect of Category V encompasses compounds having the formula:



wherein R^1 units are substituted or unsubstituted phenyl, non-limiting examples of which are described in Table XI herein below.

TABLE XI

No.	R^1
439	2-fluorophenyl
440	3-fluorophenyl
441	4-fluorophenyl
442	2-chlorophenyl
443	3-chlorophenyl
444	4-chlorophenyl
445	2-cyanophenyl
446	3-cyanophenyl
447	4-cyanophenyl
448	2-methylphenyl
449	3-methylphenyl
450	4-methylphenyl
451	2-ethylphenyl
452	3-ethylphenyl
453	4-ethylphenyl
454	2-methoxyphenyl


53

TABLE XI-continued

No.	R ¹
455	3-methoxyphenyl
456	4-methoxyphenyl
457	2-ethoxyphenyl
458	3-ethoxyphenyl
459	4-ethoxyphenyl
460	2-iso-propoxyphenyl
461	3-iso-propoxyphenyl
462	4-iso-propoxyphenyl
463	2-carbamoylphenyl
464	3-carbamoylphenyl
465	4-carbamoylphenyl
466	2-(aziridine-1-carbonyl)phenyl
467	3-(aziridine-1-carbonyl)phenyl
468	4-(aziridine-1-carbonyl)phenyl
469	2-(azetidine-1-carbonyl)phenyl
470	3-(azetidine-1-carbonyl)phenyl
471	4-(azetidine-1-carbonyl)phenyl
472	2-(pyrrolidine-1-carbonyl)phenyl
473	3-(pyrrolidine-1-carbonyl)phenyl
474	4-(pyrrolidine-1-carbonyl)phenyl
475	2-(piperidine-1-carbonyl)phenyl
476	3-(piperidine-1-carbonyl)phenyl
477	4-(piperidine-1-carbonyl)phenyl
478	2-(acetylamino)phenyl
479	3-(acetylamino)phenyl
480	4-(acetylamino)phenyl
481	2-(ethanecarbonylamino)phenyl
482	3-(ethanecarbonylamino)phenyl
483	4-(ethanecarbonylamino)phenyl
484	2-(cyclopropanecarbonylamino)phenyl
485	3-(cyclopropanecarbonylamino)phenyl
486	4-(cyclopropanecarbonylamino)phenyl

The compounds of the second aspect can be prepared by the procedure outlined in Scheme XIII and described in Example 11 herein below.

Scheme XIII



Reagents and conditions: (a) LiOH, THF, H₂O; rt, 16 hr

EXAMPLE 11

[(4-(4-Methyl-phenyl)pyridine-2-carbonyl)-amino]-acetic acid (25)


Preparation of [(4-(4-Methyl-phenyl)pyridine-2-carbonyl)-amino]-acetic acid (25): To a solution of [(4-(4-methyl-phenyl)pyridine-2-carbonyl)-amino]-acetic acid methyl ester, 24, (0.092 g, 0.32 mmol) in THF (2 mL) at room

54

temperature is added H₂O (1 mL) and LiOH.H₂O (0.027 g, 0.64 mmol). The reaction is stirred for 16 hours after which time the solution is acidified using 1M HCl. The solvents are removed under reduced pressure and the solid that remains is suspended in a mixture of THF: MeOH and filtered. The filtrate is concentrated under reduced pressure and the resulting solid is triturated with MeOH and collected by filtration to provide 0.012 g (12% yield) of the desired compound as a colorless solid. ¹H NMR (250 MHz, MeOD) δ ppm 8.69 (1 H, d, J=4.8 Hz), 8.38 (1 H, s), 7.86 (1 H, d, J=6.2 Hz), 7.72 (2 H, d, J=8.1 Hz), 7.38 (1 H, d, J=7.9 Hz), 4.21 (1 H, s), 2.44 (2 H, s). HPLC-MS: m/z 271 [M+H]⁺.


The following are non-limiting examples of compounds of Formula (I).

15




20
25
30
35
40
45
50
55
60

[[4-(4-Cyanophenyl)pyridine-2-carbonyl]amino]-acetic acid: ¹H NMR (250 MHz, MeOD) δ ppm 8.69 (1 H, d, J=6.0 Hz), 8.38 (1 H, br s), 7.71-8.01 (2 H, m), 4.10 (2 H, s). HPLC-MS: m/z 282 [M+H]⁺.




40
45
50
55
60

[[4-(4-Chlorophenyl)pyridine-2-carbonyl]-amino]-acetic acid: ¹H NMR (250 MHz, MeOD) δ ppm 8.75 (1 H, s), 8.44 (1 H, br s), 7.85 (3 H, m), 7.57 (2 H, d, J=7.8 Hz), 4.23 (2 H, s). HPLC-MS: m/z 291 [M+H]⁺.



55
60

[[4-(3-Chlorophenyl)pyridine-2-carbonyl]amino]-acetic acid: ¹H NMR (400 MHz, MeOD) δ ppm 8.63 (1 H, d, J=5.5 Hz), 8.27 (1 H, br s), 7.73 (1 H, s), 7.77 (1 H, d, J=4.0 Hz), 7.64 (1 H, d, J=7.0 Hz), 7.36-7.49 (3 H, m), 4.09 (2 H, s). HPLC-MS: m/z 291 [M+H]⁺.




60
65

[[4-(2-Chlorophenyl)pyridine-2-carbonyl]amino]-acetic acid: ¹H NMR (250 MHz, MeOD) δ ppm 8.63 (1 H, d, J=4.9

55

Hz), 8.05-8.10 (1 H, m), 7.55 (1 H, dd, J=4.9, 1.6 Hz), 7.46 (1 H, dt, J=3.9, 2.1 Hz), 7.35 (4 H, d, J=2.7 Hz), 4.08 (2 H, s). HPLC-MS: m/z 291 [M+H]⁺.

Another subgenus of the compounds of Formula (I) have the formula:




wherein non-limiting examples of R and R⁶ are further described herein below in Table XII.

TABLE XII


No.	R	R ⁶
487	—H	—H
488	—H	—CH ₃
489	—H	—CH ₂ CH ₃
490	—H	—CH ₂ CH ₂ CH ₃
491	—H	—C(CH ₃) ₃
492	—OH	—H
493	—OH	—CH ₃
494	—OH	—CH ₂ CH ₃
493	—OH	—CH ₂ CH ₂ CH ₃
496	—OH	—C(CH ₃) ₃
497	—Cl	—H
498	—Cl	—CH ₃
499	—Cl	—CH ₂ CH ₃
500	—Cl	—CH ₂ CH ₂ CH ₃
501	—Cl	—C(CH ₃) ₃
502	—OCH ₃	—H
503	—OCH ₃	—CH ₃
504	—OCH ₃	—CH ₂ CH ₃
505	—OCH ₃	—CH ₂ CH ₂ CH ₃
506	—OCH ₃	—C(CH ₃) ₃
507	—CN	—H
508	—CN	—CH ₃
509	—CN	—CH ₂ CH ₃
510	—CN	—CH ₂ CH ₂ CH ₃
511	—CN	—C(CH ₃) ₃
512	—F	—H
513	—F	—CH ₃
514	—F	—CH ₂ CH ₃
515	—F	—CH ₂ CH ₂ CH ₃
516	—F	—C(CH ₃) ₃

The compounds described above can be prepared by the procedures outlined in Schemes XIV-XVI and described in Examples 12-14 herein below.


Scheme XIV



56



Reagents and conditions: (a) glycine tert-butyl ester HCl, EDCI, HOBt, DIPEA, DMF; rt, 48 hr



Reagents and conditions: (b) H₂; Pd/C, EtOH, rt, 22 hr

EXAMPLE 12


[(3,5-Dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester (27)

Preparation of [(3,5-bis-benzyloxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester (26): To a solution of 3,5-bis-benzyloxy-pyridine-2-carboxylic acid, 2, (2.36 g, 6.36 mmol) in DMF (20 mL) at room temperature under N₂ is added 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide (EDCI) (1.83 g, 9.54 mmol) and 1-hydroxybenzo-triazole (HOBt) (0.086 g, 0.64 mmol). The mixture is stirred for 15 minutes after which time glycine tert-butyl ester hydrochloride (1.60 g, 9.54 mmol) and diisopropylethylamine (DIPEA) (3.32 ml, 19.08 mmol) are added. The resulting solution is stirred at room temperature for 48 hours then concentrated under reduced pressure. The resulting brown oil is purified over silica (EtOAc) to afford 3.04 g (99% yield) of the desired compound as a yellow solid. ¹H NMR (250 MHz, CDCl₃) δ ppm 8.19 (1 H, t, J=5.2 Hz), 8.01-8.08 (2 H, m), 7.27-7.54 (9 H, m), 6.97 (1 H, d, J=2.4 Hz), 5.24 (2 H, s), 5.13 (2 H, s), 4.17 (2 H, d, J=5.2 Hz), 1.51 (9 H, s). HPLC-MS: m/z 449 [M+H]⁺.

Preparation of [(3,5-dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester (27): A solution of [(3,5-bis-benzyloxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester, 26, (3.04 g, 6.79 mmol) in EtOH (100 mL) containing 10% Pd/C (0.300 g) is stirred under an atmosphere of H₂ for 22 hours. The suspension is then filtered through Celite™, concentrated under reduced pressure, and the crude product purified over silica (2.5% MeOH/CH₂Cl₂) to afford 1.20 g (66% yield) of the desired compound as a colorless oil. ¹H NMR (250 MHz, CDCl₃) δ ppm 11.90 (1 H, br s), 8.94 (1 H, br s), 8.20 (1 H, t, J=5.6 Hz), 7.76 (1 H, d, J=2.4 Hz), 6.77 (1 H, d, J=2.1 Hz), 4.13 (2 H, d, J=5.5 Hz), 1.53 (9 H, s). HPLC-MS: m/z 269 [M+H]⁺.

57

Scheme XV




Reagents and conditions: (a) TFA, CH₂Cl₂; rt, 16 hr

EXAMPLE 13


[(3,5-Dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid (29)

Preparation of [(3,5-dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid (29): To a solution of [(3,5-dihydroxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester, 28, (0.10 g, 0.37 mmol) in CH₂Cl₂ (4 mL) at room temperature is added trifluoroacetic acid (1 mL). The reaction is stirred for 16 hours at room temperature and then concentrated under reduced pressure. The solid that remains is collected by filtration, washed with Et₂O to afford 0.070 g (89% yield) of the desired compound as a colorless solid. ¹H NMR (250 MHz, DMSO-d₆) δ ppm 10.86 (1 H, br s), 9.00 (1 H, t, J=6.1 Hz), 7.77 (1 H, d, J=2.4 Hz), 6.69 (1 H, d, J=2.4 Hz), 3.95 (1 H, d, J=6.2 Hz). HPLC-MS: m/z 213 [M+H]⁺.

Scheme XVI




Reagents and conditions: (a) glycine tert-butyl ester HCl, EDCI, HOBt, DIPEA, DMF; rt, 3 days



58

-continued



Reagents and conditions: (b) TFA, CH₂Cl₂; rt, 5 hr


EXAMPLE 14

[(3-Hydroxy-pyridine-2-carbonyl)-amino]-acetic acid (30)

Preparation of [(3-hydroxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester (29): To a solution of 3-hydroxypicolinic acid (0.20 g, 1.44 mmol) in DMF (5 mL) at 0° C. under N₂ is added diisopropylethylamine (DIPEA) (0.75 ml, 4.3 mmol), 1-(3-dimethyl-aminopropyl)-3-ethylcarbodiimide (EDCI) (0.412 g, 2.9 mmol) and 1-hydroxybenzo-triazole (HOBt) (0.019 g, 0.14 mmol). The resulting mixture is stirred for 5 min before glycine tert-butyl ester HCl (0.36 g, 2.9 mmol) is introduced. The resulting solution is stirred at room temperature for 3 days then concentrated under reduced pressure. The reaction mixture is diluted with EtOAc then washed with 1M HCl, sat. NaCl, and the organic layer is dried (MgSO₄), filtered and concentrated under reduced pressure to a crude oil that is purified over silica (EtOAc/heptane 1:4) to afford 0.078 g (22% yield) of the desired compound as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 11.80 (1 H, s), 8.39 (1 H, br s), 8.02 (1 H, dd, J=4.4, 1.5 Hz), 7.27 (1 H, dd, J=8.8, 4.4 Hz), 7.25 (1 H, dd, J=8.4, 1.5 Hz), 4.06 (2 H, d, J=5.5 Hz), 1.44 (9 H, s). HPLC-MS: m/z 197 [M-tBu]⁺.

Preparation of [(3-Hydroxy-pyridine-2-carbonyl)-amino]-acetic acid (30): To a solution of [(3-hydroxy-pyridine-2-carbonyl)-amino]-acetic acid tert-butyl ester, 29, (0.070 g, 0.277 mmol) in CH₂Cl₂ (4 mL) is added TFA (1 mL). The resulting solution is stirred for 5 hours then concentrated under reduced pressure to afford 0.054 g (99% yield) of the desired compound as a colorless solid. ¹H NMR (400 MHz, MeOD) δ ppm 8.09 (1 H, d, J=3.3 Hz), 7.36-7.59 (2 H, m), 4.08 (2 H, s). HPLC-MS: m/z 197 [M+H]⁺.

Category VII of the present disclosure relates to compounds having the formula:




wherein non-limiting examples of R and R⁶ are further described herein below in Table XIII.

59

TABLE XIII


No.	R	R ⁶
517	3-fluorophenyl	—H
518	4-fluorophenyl	—H
519	3-chlorophenyl	—H
520	4-chlorophenyl	—H
521	3-cyanophenyl	—H
522	4-cyanophenyl	—H
523	3-methylphenyl	—H
524	4-methylphenyl	—H
525	3-ethylphenyl	—H
526	4-ethylphenyl	—H
527	3-methoxyphenyl	—H
528	4-methoxyphenyl	—H
529	3-ethoxyphenyl	—H
530	4-ethoxyphenyl	—H
531	3-fluorophenyl	—CH ₃
532	4-fluorophenyl	—CH ₃
533	3-chlorophenyl	—CH ₃
534	4-chlorophenyl	—CH ₃
535	3-cyanophenyl	—CH ₃
536	4-cyanophenyl	—CH ₃
537	3-methylphenyl	—CH ₃
538	4-methylphenyl	—CH ₃
539	3-ethylphenyl	—CH ₃
540	4-ethylphenyl	—CH ₃
541	3-methoxyphenyl	—CH ₃
542	4-methoxyphenyl	—CH ₃
543	3-ethoxyphenyl	—CH ₃
544	4-ethoxyphenyl	—CH ₃
545	3-fluorophenyl	—CH ₂ CH ₃
546	4-fluorophenyl	—CH ₂ CH ₃
547	3-chlorophenyl	—CH ₂ CH ₃
548	4-chlorophenyl	—CH ₂ CH ₃
549	3-cyanophenyl	—CH ₂ CH ₃
550	4-cyanophenyl	—CH ₂ CH ₃
551	3-methylphenyl	—CH ₂ CH ₃
552	4-methylphenyl	—CH ₂ CH ₃
553	3-ethylphenyl	—CH ₂ CH ₃
554	4-ethylphenyl	—CH ₂ CH ₃
555	3-methoxyphenyl	—CH ₂ CH ₃
556	4-methoxyphenyl	—CH ₂ CH ₃
557	3-ethoxyphenyl	—CH ₂ CH ₃
558	4-ethoxyphenyl	—CH ₂ CH ₃

The compounds which encompass Category VII of the present disclosure can be prepared by the procedures outlined in Schemes XVII and XVIII and described in Examples 15 and 16 herein below.




60


-continued



Reagents and conditions: (a) (4-ClC₆H₄)B(OH)₂, Pd(dppf)Cl₂, K₃PO₄, 1,4-dioxane, MeOH; 80° C., 3 hr



Reagents and conditions: (b) LiOH, THF, H₂O; reflux



Reagents and conditions: (c) GlyOEt.HCl, EDCl, HOBT, DIPEA, DMF, CH₂Cl₂; Rt


61

EXAMPLE 15

3-Methoxy-4'-methyl-biphenyl-4-carboxylic acid methyl ester (30)

Preparation of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid methyl ester (28): To a degassed solution of methyl 4-bromo-2-methoxybenzoate (0.70 g, 2.86 mmol) in 1,4-dioxane (10 mL) and MeOH (2.5 mL) is added 4-chlorophenyl boronic acid (0.536 g, 3.43 mmol), Pd(dppf)Cl₂ (0.233 g, 0.286 mmol) and K₃PO₄ (0.728 g, 3.43 mmol). The resulting suspension is heated to 80° C. and stirred for 3 hours. After this time, the reaction is cooled to room temperature and filtered through Celite™. The solids that form are collected and washed with additional MeOH before the filtrate is concentrated under reduced pressure. The crude material is purified over silica (hexanes:EtOAc; 6:1 to 4:1) to provide 0.615 g (78% yield) of the desired compound as orange crystals. ¹H NMR (400 MHz, CDCl₃) δ ppm 7.89 (1 H, d, J=8.0 Hz), 7.52-7.56 (2 H, m), 7.44 (2 H, d, J=8.7 Hz), 7.17 (1 H, d, J=8.0 Hz), 7.12 (1 H, d, J=1.6 Hz), 3.99 (3 H, s), 3.92 (3 H, s). HPLC-MS: m/z 277 [M+H]⁺.

Scheme XVIII



Reagents and conditions: (a) LiOH, THF, H₂O; reflux, 2 hr


EXAMPLE 16

4'-Chloro-3-methoxy-biphenyl-4-carboxylic acid


Preparation of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid (31): To a solution of 4'-chloro-3-methoxy-biphenyl-4-carboxylic acid methyl ester, 30, (0.615 g, 2.22 mmol) in THF (20 mL) and H₂O (5 mL) is added LiOH (0.932 g, 22.2 mmol). The resulting suspension is heated to reflux for 2 hours. The reaction is cooled and concentrated under reduced pressure. The crude product is acidified using conc. HCl and the resulting solid is collected by filtration, washed with H₂O to afford 0.532 g (91% yield) of the desired compound as a grey solid. ¹H NMR (400 MHz, CDCl₃) δ ppm 10.69 (1 H, br s), 8.26 (1 H, d, J=8.1 Hz), 7.53-7.58 (2 H, m), 7.44-7.50 (2 H, m), 7.33 (1 H, dd, J=8.1, 1.6 Hz), 7.20 (1 H, d, J=1.3 Hz), 4.17 (3 H, s). HPLC-MS: m/z 263 [M+H]⁺.

62

The following are non-limiting examples of additional compounds of Formula (I).




3-Methoxy-4'-methyl-biphenyl-4-carboxylic acid methyl ester: ¹H NMR (250 MHz, CDCl₃) δ ppm 7.89 (1 H, d, J=7.9 Hz), 7.52 (2 H, d, J=8.2 Hz), 7.25-7.32 (2 H, m), 7.15-7.24 (2 H, m), 3.99 (3 H, s), 3.92 (3 H, s), 2.42 (3 H, s). HPLC-MS: m/z 257 [M+H]⁺.



3-Methoxy-4'-methyl-biphenyl-4-carboxylic acid: ¹H NMR (400 MHz, CDCl₃) δ ppm 10.74 (1 H, br s), 8.24 (1 H, d, J=8.1 Hz), 7.52 (2 H, d, J=8.1 Hz), 7.36 (1 H, dd, J=8.1, 1.5 Hz), 7.30 (2 H, d, J=7.9 Hz), 7.23 (1 H, d, J=1.4 Hz), 4.16 (3 H, s), 2.43 (3 H, s). HPLC-MS: m/z 243 [M+H]⁺.

Additional compounds of Formula (I) the formula:




wherein non-limiting examples of R^{8a}, R^{8b}, R⁹ and R⁶ are further described herein below in Table.

TABLE XIV

No.	R	R ⁶	R ^{8a}	R ^{8b}	R ⁹
559	3-chlorophenyl	-H	-CH ₃	-H	-H
560	3-chlorophenyl	-CH ₃	-CH ₃	-H	-H
561	3-chlorophenyl	-H	-CH ₃	-CH ₃	-H
562	3-chlorophenyl	-CH ₃	-CH ₃	-CH ₃	-H
563	3-chlorophenyl	-H	-CH ₃	-CH ₃	-CH ₃
564	3-chlorophenyl	-CH ₃	-CH ₃	-CH ₃	-CH ₃
565	4-chlorophenyl	-H	-CH ₃	-H	-H
566	4-chlorophenyl	-CH ₃	-CH ₃	-H	-H
567	4-chlorophenyl	-H	-CH ₃	-CH ₃	-H
568	4-chlorophenyl	-CH ₃	-CH ₃	-CH ₃	-H
569	4-chlorophenyl	-H	-CH ₃	-CH ₃	-CH ₃
570	4-chlorophenyl	-CH ₃	-CH ₃	-CH ₃	-CH ₃
571	4-methylphenyl	-H	-CH ₃	-H	-H
572	4-methylphenyl	-CH ₃	-CH ₃	-H	-H
573	4-methylphenyl	-H	-CH ₃	-CH ₃	-H
574	4-methylphenyl	-CH ₃	-CH ₃	-CH ₃	-H
575	4-methylphenyl	-H	-CH ₃	-CH ₃	-CH ₃
576	4-methylphenyl	-CH ₃	-CH ₃	-CH ₃	-CH ₃

63

The compounds described above can be prepared by the procedures outlined herein below in Schemes XIX and XX and described in Examples 17 and 18 herein below.




Reagents and conditions: (a)
2-(N-methylamino)acetic acid ester HCl, EDCI,
HOBt, DIPEA, DMF; rt, 32 hr

EXAMPLE 17

[(3-Hydroxy-pyridine-2-carbonyl)-methyl-amino]-
acetic acid ethyl ester (32)

Preparation of [(3-hydroxy-pyridine-2-carbonyl)methyl-amino]acetic acid ethyl ester (31): To a solution of 3-hydroxy picolinic acid (0.40 g, 2.88 mmol) in DMF (5 mL) is added diisopropylethylamine (DIPEA) (1.50 ml, 8.63 mmol), 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide (EDCI) (0.825 g, 4.31 mmol) and 1-hydroxybenzotriazole (HOBt) (0.039 g, 0.29 mmol). The reaction mixture is stirred for 5 minutes then methylamino-acetic acid ester hydrochloride (0.663 g, 4.31 mmol) is added. The reaction is stirred at room temperature for 32 hours after which the solvent is removed under reduced pressure. The residue is partitioned between EtOAc and 1M HCl and the organic layer separated, dried (MgSO₄), filtered and concentrated under reduced pressure. The crude material is purified over silica (EtOAc:hexanes 1:1) to afford 0.10 g (15% yield) of the desired compound as a colourless solid. HPLC-MS: m/z 240 [M+H]⁺.



Reagents and conditions: (a) NaOH, THF, H₂O;
reflux, 3 hr

EXAMPLE 18


[(3-Hydroxy-pyridine-2-carbonyl)-methyl-amino]-
acetic acid (33)

Preparation of [(3-hydroxy-pyridine-2-carbonyl)-methyl-amino]-acetic acid (33): To a solution of [(3-hydroxy-pyri-


64

dine-2-carbonyl)methylamino]acetic acid ethyl ester, 32, (0.10 g, 0.42 mmol) in THF (4 mL) is added H₂O (1 mL) and NaOH (0.90 g, 2.25 mmol). The reaction is stirred for 3 hours then concentrated under reduced pressure. The remaining oil is acidified to pH ~1 with 1M HCl and the solution is concentrated under reduced pressure to give an off-white solid. The solid is suspended in CHCl₃:isopropanol (1:1) then collected by filtration. The solid is washed with additional CHCl₃:isopropanol (1:1) then transferred to a flask and triturated with Et₂O to afford 0.075 g (85% yield) of the desired compound as a pale yellow solid. ¹H NMR (250 MHz, MeOD) δ ppm (rotamers) 8.26 (1 H, br s), 7.63-7.74 (1 H, m), 7.56-7.63 (1 H, m), 4.38 (1 H, s), 4.32 (1 H, s), 3.20 (1.5 H, s), 3.12 (1.5 H, s). HPLC-MS: m/z 211 [M+H]⁺.


The following are non-limiting examples of the compounds described above.



2-(S)-[(3-Hydroxy-pyridine-2-carbonyl)-amino]-propionic acid methyl ester: ¹H NMR (250 MHz, CDCl₃) δ ppm 11.86 (1 H, s), 8.47 (1 H, br s), 8.10 (1 H, dd, J=4.1, 1.7 Hz), 7.28-7.43 (2 H, m), 4.63-4.84 (1 H, m), 3.81 (3 H, s), 1.57 (3 H, d, J=7.3 Hz). HPLC-MS: m/z 225 [M+H]⁺.



1-(3-Hydroxy-pyridine-2-carbonyl)-pyrrolidine-2-carboxylic acid methyl ester: ¹H NMR (250 MHz, CDCl₃) δ ppm (rotamers) 12.86 (0.67 H, br s), 12.44 (0.33 H, br s), 8.14 (0.33 H, t, J=2.9 Hz), 7.98 (0.67 H, dd, J=3.7, 2.1 Hz), 7.24-7.31 (2 H, m), 5.38 (0.67 H, dd, J=8.5, 3.4 Hz), 4.63-4.75 (0.33 H, m), 4.37 (0.67 H, t, J=6.7 Hz), 3.80-4.01 (1.33 H, m), 3.79 (1 H, s), 3.70 (2 H, s), 1.87-2.44 (4 H, m). HPLC-MS: m/z 251 [M+H]⁺.



2S-[(3-Hydroxy-pyridine-2-carbonyl)-amino]-propionic acid: ¹H NMR (250 MHz, DMSO-d₆) δ ppm 12.87 (1 H, br s), 12.28 (1 H, s), 9.19 (1 H, d, J=7.5 Hz), 8.19 (1 H, dd, J=4.3, 1.4 Hz), 7.50-7.65 (1 H, m), 7.37-7.49 (1 H, m), 3.95-4.95 (1 H, m), 1.45 (3 H, d, J=7.3 Hz). HPLC-MS: m/z 211 [M+H]⁺.

Administration of one or more of the compounds of Formula (I), alone in the form of pharmaceutical compositions, optionally in combination with other pharmaceutically active compounds or compositions, can be effective in treatment of the following disease states or conditions:

- i) as human protein HIF-1 α prolyl hydroxylase inhibitors; and thereby providing a means for regulating blood flow, oxygen delivery and energy utilization in ischemic tissues;
- ii) the compounds of the present disclosure are efficacious in regulating blood flow, oxygen delivery and energy utilization in ischemic tissues; and
- iii) the compounds of the present disclosure provide stabilized HIF-1 α by blocking a degradation pathway mediated by HIF prolyl hydroxylase.

Each of the disease states or conditions which the formulator desires to treat may require differing levels or amounts of the compounds described herein to obtain a therapeutic level. The formulator can determine this amount by any of the testing procedures known to the artisan or ordinary skill in the art.

The compounds of the present disclosure can be HIF-1 α prolyl hydroxylase inhibitors when administered in pharmaceutically effective amounts, and thereby provide increased angiogenic response or cellular responses which are activated by transcription factors that are directly or indirectly affected by an increase in cellular HIF-1 α concentration. Non-limiting examples of these diseases or disease states are listed herein below, inter alia, Peripheral Vascular Disease (PVD), Coronary Artery Disease (CAD), heart failure, ischemia, and anemia. The compounds disclosed herein are especially effective in the treatment of anemia.

Stimulation of EPO production: anemia

HIF-1 is a transcription factor that also regulates the hypoxia-inducible EPO gene. HIF-1 binding is required for EPO transcriptional activation in response to hypoxia (Semenza, G. L., "Regulation of erythropoietin production: New insights into molecular mechanisms of oxygen homeostasis", *Hematol. Oncol. Clin. North Am.*, Vol. 8, pp. 863-884 (1994)). In particular, HIF-1 α binds to the 3' hypoxia-response element of the EPO gene which results in the marked enhancement of EPO transcription (Semenza, G. L., et al. "Transcriptional regulation of genes encoding glycolytic enzymes by hypoxia-inducible factor 1", *J. Biol. Chem.*, Vol. 269, pp. 23757-63 (1994)). EPO, in turn, is essential for maintenance of red blood cells by controlling the proliferation and differentiation of erythroid progenitor cells into red blood cells (Krantz, S. B., "Erythropoietin," *Blood*, Vol. 77, pp 419-434 (1991)). During fetal development, the liver serves as the primary source of EPO. Shortly before birth, production of EPO in the liver decreases and the kidney becomes the primary source of EPO. However, in adults other organs such as the liver and brain produce small but significant amounts of EPO. A erythropoietin deficiency is associated with anemia. In humans, the most prevalent form of anemia is associated with kidney failure.

EPO has been described in the treatment of anemia: associated with chemotherapy; that occurs as a consequence of AIDS; and due to prematurity and autologous blood donation. EPO has even been suggested as a general use agent in pre-operative elective surgery.

Angiogenesis

Angiogenesis, the sprouting of new blood vessels from the pre-existing vasculature, plays a crucial role in a wide range of physiological and pathological processes (Nguyen, L. L. et al., *Int. Rev. Cytol.*, 204, 1-48, (2001)). Angiogenesis is a complex process, mediated by communication between the endothelial cells that line blood vessels and their surrounding environment. In the early stages of angiogenesis, tissue or tumor cells produce and secrete pro-angiogenic growth factors in response to environmental stimuli such as hypoxia. These factors diffuse to nearby endothelial cells

and stimulate receptors that lead to the production and secretion of proteases that degrade the surrounding extracellular matrix. The activated endothelial cells begin to migrate and proliferate into the surrounding tissue toward the source of these growth factors (Bussolino, F., *Trends Biochem. Sci.*, 22, 251-256, (1997)). Endothelial cells then stop proliferating and differentiate into tubular structures, which is the first step in the formation of stable, mature blood vessels. Subsequently, periendothelial cells, such as pericytes and smooth muscle cells, are recruited to the newly formed vessel in a further step toward vessel maturation.

Angiogenesis is regulated by a balance of naturally occurring pro- and anti-angiogenic factors. Vascular endothelial growth factor, fibroblast growth factor, and angiopoietin represent a few of the many potential pro-angiogenic growth factors. These ligands bind to their respective receptor tyrosine kinases on the endothelial cell surface and transduce signals that promote cell migration and proliferation. Whereas many regulatory factors have been identified, the molecular mechanisms of this process are still not fully understood.

There are many disease states driven by persistent unregulated or improperly regulated angiogenesis. In such disease states, unregulated or improperly regulated angiogenesis may either cause a particular disease or exacerbate an existing pathological condition. For example, ocular neovascularization has been implicated as the most common cause of blindness and underlies the pathology of approximately 20 eye diseases. In certain previously existing conditions such as arthritis, newly formed capillary blood vessels invade the joints and destroy cartilage. In diabetes, new capillaries formed in the retina invade the vitreous humor, causing bleeding and blindness.

Both the growth and metastasis of solid tumors are also angiogenesis-dependent (Folkman et al., "Tumor Angiogenesis," Chapter 10, 206-32, in *The Molecular Basis of Cancer*, Mendelsohn et al., eds., W. B. Saunders, (1995)). It has been shown that tumors which enlarge to greater than 2 mm in diameter must obtain their own blood supply and do so by inducing the growth of new capillary blood vessels. After these new blood vessels become embedded in the tumor, they provide nutrients and growth factors essential for tumor growth as well as a means for tumor cells to enter the circulation and metastasize to distant sites, such as liver, lung or bone (Weidner, *New Eng. J. Med.*, 324, 1, 1-8 (1991)). When used as drugs in tumor-bearing animals, natural inhibitors of angiogenesis may prevent the growth of small tumors (O'Reilly et al., *Cell*, 79, 315-28 (1994)). In some protocols, the application of such inhibitors leads to tumor regression and dormancy even after cessation of treatment (O'Reilly et al., *Cell*, 88, 277-85 (1997)). Moreover, supplying inhibitors of angiogenesis to certain tumors may potentiate their response to other therapeutic regimens (Teischer et al., *Int. J. Cancer*, 57, 920-25 (1994)).

Peripheral Vascular Disease

Peripheral vascular disease (PVD) is the term used to describe the clinical syndrome of vascular insufficiency outside of the coronary circulation and typically involving the circulation to the lower extremities. There are an estimated 8-12 million patients in the US with peripheral vascular disease; another 16.5 million undiagnosed. Atherosclerosis is by far the leading cause of peripheral vascular disease (PVD), although a number of discrete disease processes can contribute to its development and progression (i.e. diabetes, immune vasculitis and trauma). Atherosclerotic PVD can present in three ways:

- 1) Asymptomatic PVD diagnosed on the basis of noninvasive testing (usually physical exam);
- 2) Intermittent claudication with symptoms of leg pain with exercise; and
- 3) Critical limb ischemia with leg pain at rest and limb-threatening ischemic changes (usually non-healing or infected cutaneous ulcerations).

The age-adjusted (average age 66 years) prevalence of PVD in the US population is approximately 12%. Of those patients with claudication, 20-30% will have progressive symptoms and 10% will require amputation for critical limb ischemia. Although patients with symptomatic PVD suffer significant decreases in mobility, muscle mass, bone density and quality of life, there are currently no effective medical therapies available.

The present disclosure provides compounds which when administered in vivo inhibit HIF-1 α prolyl hydroxylase thereby leading to increased expression of HIF-regulated genes, inter alia, angiogenic factors, erythropoietin, and glycolytic enzymes thereby resulting in improvement in blood flow, oxygen delivery and energy utilization in ischemic tissues.

Although many disease states are driven by persistent unregulated or improperly regulated angiogenesis, some disease states could be treated by increased angiogenesis. Tissue growth and repair are biologic events wherein cellular proliferation and angiogenesis occur. Thus an important aspect of wound repair is the revascularization of damaged tissue by angiogenesis.

Wounds

Chronic, non-healing wounds are a major cause of prolonged morbidity in the aged human population. This is especially the case in bedridden or diabetic patients who develop severe, non-healing skin ulcers. In many of these cases, the delay in healing is a result of inadequate blood supply either as a result of continuous pressure or of vascular blockage. Poor capillary circulation due to small artery atherosclerosis or venous stasis contributes to the failure to repair damaged tissue. Such tissues are often infected with microorganisms that proliferate unchallenged by the innate defense systems of the body which require well vascularized tissue to effectively eliminate pathogenic organisms. As a result, most therapeutic intervention centers on restoring blood flow to ischemic tissues thereby allowing nutrients and immunological factors access to the site of the wound.

Atherosclerotic Lesions

Atherosclerotic lesions in large vessels may cause tissue ischemia that could be ameliorated by modulating blood vessel growth to the affected tissue. For example, atherosclerotic lesions in the coronary arteries may cause angina and myocardial infarction that could be prevented if one could restore blood flow by stimulating the growth of collateral arteries. Similarly, atherosclerotic lesions in the large arteries that supply the legs may cause ischemia in the skeletal muscle that limits mobility and in some cases necessitates amputation, which may also be prevented by improving blood flow with angiogenic therapy.

Diabetes/Hypertension

Diseases such as diabetes and hypertension are characterized by a decrease in the number and density of small blood vessels such as arterioles and capillaries. These small blood vessels are important for the delivery of oxygen and nutrients. A decrease in the number and density of these vessels contributes to the adverse consequences of hypertension and diabetes including claudication, ischemic ulcers, accelerated hypertension, and renal failure. These common disorders and many other less common ailments, such as

Burgers disease, could be ameliorated by increasing the number and density of small blood vessels using angiogenic therapy.

The present disclosure further relates to forms of the present compounds, which under normal human or higher mammalian physiological conditions, release the compounds described herein. One iteration of this aspect includes the pharmaceutically acceptable salts of the analogs described herein. The formulator, for the purposes of compatibility with delivery mode, excipients, and the like, can select one salt form of the present analogs over another since the compounds themselves are the active species which mitigate the disease processes described herein.

Formulations

The present disclosure also relates to compositions or formulations which comprise the human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure. In general, the compositions of the present disclosure comprise:

- a) an effective amount of one or more human protein HIF-1 α prolyl hydroxylase inhibitor according to the present disclosure which are effective for treating PVD, CAD, heart failure, ischemia, and anemia; and
- b) one or more excipients.

For the purposes of the present disclosure the term "excipient" and "carrier" are used interchangeably throughout the description of the present disclosure and said terms are defined herein as, "ingredients which are used in the practice of formulating a safe and effective pharmaceutical composition."

The formulator will understand that excipients are used primarily to serve in delivering a safe, stable, and functional pharmaceutical, serving not only as part of the overall vehicle for delivery but also as a means for achieving effective absorption by the recipient of the active ingredient. An excipient may fill a role as simple and direct as being an inert filler, or an excipient as used herein may be part of a pH stabilizing system or coating to insure delivery of the ingredients safely to the stomach. The formulator can also take advantage of the fact the compounds of the present disclosure have improved cellular potency, pharmacokinetic properties, as well as improved oral bioavailability.

Non-limiting examples of compositions according to the present disclosure include:

- a) from about 0.001 mg to about 1000 mg of one or more human protein HIF-1 α prolyl hydroxylase inhibitor according to the present disclosure; and
- b) one or more excipients.

Another example according to the present disclosure relates to the following compositions:

- a) from about 0.01 mg to about 100 mg of one or more human protein prolyl HIF-1 α prolyl hydroxylase inhibitor according to the present disclosure; and
- b) one or more excipients.

A further example according to the present disclosure relates to the following compositions:

- a) from about 0.1 mg to about 10 mg of one or more human protein HIF-1 α prolyl hydroxylase inhibitor according to the present disclosure; and
- b) one or more excipients.

The term "effective amount" as used herein means "an amount of one or more HIF-1 α prolyl hydroxylase inhibitors, effective at dosages and for periods of time necessary to achieve the desired or therapeutic result." An effective amount may vary according to factors known in the art, such

as the disease state, age, sex, and weight of the human or animal being treated. Although particular dosage regimes may be described in examples herein, a person skilled in the art would appreciate that the dosage regime may be altered to provide optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation. In addition, the compositions of the present disclosure can be administered as frequently as necessary to achieve a therapeutic amount.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating anemia.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating angiogenesis.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating peripheral vascular disease.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating wounds.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating atherosclerotic lesions.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating diabetes.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating hypertension.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating a disease affected by the level of VEGF, GAPDH and erythropoietin.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating a disorder chosen from Crohn's disease and ulcerative colitis, psoriasis, sarcoidosis, rheumatoid arthritis, hemangiomas, Osler-Weber-Rendu disease, or hereditary hemorrhagic telangiectasia, solid or blood borne tumors and acquired immune deficiency syndrome.

The present disclosure further relates to the use of one or more of the HIF-1 α prolyl hydroxylase inhibitors disclosed herein for making a medicament for treating a disorder chosen from diabetic retinopathy, macular degeneration, cancer, sickle cell anemia, sarcoid, syphilis, pseudoxanthoma elasticum, Paget's disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis/vitritis, mycobacterial infections, Lyme's disease, systemic lupus erythematosus, retinopathy of prematurity, Eales' disease, Behcet's disease, infections causing a retinitis or choroiditis, presumed ocular histoplasmosis, Best's disease, myopia, optic pits, Stargardt's disease, pars planitis, chronic retinal detachment, hyperviscosity syndrome, toxoplasmosis, trauma and post-laser complications, diseases associated with rubeosis, and proliferative vitreoretinopathy.

Method of Use

By increasing the transcription of these HIF-1 target genes, the HIF-1 α prolyl hydroxylase inhibitors of the present disclosure provide a method for increasing the vascularization of tissue in a subject. As used herein, "vas-

cularization of tissue" means a pro-angiogenic response whereby blood vessels or other vessels or ducts develop at or around the afflicted tissue. The afflicted tissue need not be hypoxic or ischemic per se, but rather the HIF-1 α prolyl hydroxylase inhibitors help to sustain or further stimulate the body's pro-angiogenic response to hypoxia. A non-limiting example of "vascularization" includes capillary proliferation in a non-healing wound or along the border of ischemic tissue. Thus, these compounds enhance the ability of the body to revascularize damaged tissues or increase vasculature (e.g. to prevent hypoxic damage). Non-limiting examples of "tissue" include: cardiac tissue, such as myocardium and cardiac ventricles; skeletal muscle; neurological tissue, such as from the cerebellum; internal organs, such as the stomach, intestine, pancreas, liver, spleen, and lung; and distal appendages such as fingers and toes.

Stimulated by a build up of the cellular concentration of HIF-1 α is the production of Vascular Endothelial Growth Factor (VEGF) which is known for its ability to induce vascular leakage. Oxygen tension has been shown to be a key regulator of VEGF gene expression, both in vitro and in vivo. During VEGF induction it is demonstrated that VEGF induces the formation of functional neo-vessels in the mouse cornea and enhanced blood flow in a dog model of coronary artery disease. The HIF-1 α prolyl hydroxylase inhibitors of the present disclosure provide enhancement in the expression of multiple hypoxia inducible genes including VEGF, GAPDH and erythropoietin (EPO). Additionally, the HIF-1 α prolyl hydroxylase inhibitors of the present disclosure provide enhanced the accumulation of HIF1- α in the cytoplasm and nucleus. Transgenic mice expressing a constitutively active HIF-1 α in the skin have increased dermal vascularity and had a 13-fold increase in VEGF levels

The present disclosure also relates to a method for controlling human protein HIF-1 α prolyl hydroxylase. The present method comprises the step of administering to a human or higher mammal an effective amount of a composition comprising one or more human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure.

The present disclosure also relates to the use of the human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure in the manufacture of a medicament for the treatment of atrial arrhythmias and related disorders.

The present disclosure also relates to hypoxia inducible factor HIF-1 α prolyl hydroxylase inhibition in myocardial remodeling and function, thereby providing means for inducing angiogenesis in a patient experiencing ischemia.

The present disclosure relates to a method for treating anemia comprising administering to a human or mammal in need of treatment an effective amount of one or more human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure.

The present disclosure relates to a method for regulating anemia comprising administering to a human or mammal in need of treatment an effective amount of one or more human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure.

The present disclosure relates to a method for preventing anemia comprising administering to a human or mammal in need of treatment an effective amount of one or more human protein HIF-1 α prolyl hydroxylase inhibitors according to the present disclosure.

EGLN-1 Activity Assay

The EGLN-1 (or EGLN-3) enzyme activity is determined using mass spectrometry (matrix-assisted laser desorption ionization, time-of-flight MS, MALDI-TOF MS- for assay details, see reference (Greis et al., 2006). Recombinant human EGLN-1-179/426 is prepared as described above and in the Supplemental Data. Full-length recombinant human EGLN-3 is prepared in a similar way, however it is necessary to use the His-MBP-TVMVEGLN-3 fusion for the assay due to the instability of the cleaved protein. For both enzymes, the HIF-1 α peptide corresponding to residues 556-574 (DLDLEALAPYIPADDDFQL) (SEQ ID NO. 1) is used as substrate. The reaction is conducted in a total volume of 50 μ L containing TrisCl (5 mM, pH 7.5), ascorbate (120 μ M), 2-oxoglutarate (3.2 μ M), HIF-1 α (8.6 μ M), and bovine serum albumin (0.01%). The enzyme, quantity predetermined to hydroxylate 20% of substrate in 20 minutes, is added to start the reaction. Where inhibitors are used, compounds are prepared in dimethyl sulfoxide at 10-fold final assay concentration. After 20 minutes at room temperature, the reaction is stopped by transferring 10 μ L of reaction mixture to 50 μ L of a mass spectrometry matrix solution (α -cyano-4-hydroxycinnamic acid, 5 mg/mL in 50% acetonitrile/0.1% TFA, 5 mM NH₄PO₄). Two microliters of the mixture is spotted onto a MALDI-TOF MS target plate for analysis with an Applied Biosystems (Foster City, Calif.) 4700 Proteomics Analyzer MALDI-TOF MS equipped with a Nd:YAG laser (355 nm, 3 ns pulse width, 200 Hz repetition rate). Hydroxylated peptide product is identified from substrate by the gain of 16 Da. Data defined as percent conversion of substrate to product is analyzed in GraphPad Prism 4 to calculate IC₅₀ values.

VEGF ELISA Assay

HEK293 cells are seeded in 96-well poly-lysine coated plates at 20,000 cells per well in DMEM (10% FBS, 1% NEAA, 0.1% glutamine). Following overnight incubation, the cells are washed with 100 μ L of Opti-MEM (Gibco, Carlsbad, Calif.) to remove serum. Compound in DMSO is serially diluted (beginning with 100 μ M) in Opti-MEM and added to the cells. The conditioned media is analyzed for VEGF with a Quantikine human VEGF immunoassay kit (R&D Systems, Minneapolis, Minn.). Optical density measurements at 450 nm are recorded using the Spectra Max 250 (Molecular Devices, Sunnyvale, Calif.). Data defined as % of DFO stimulation is used to calculate EC₅₀ values with GraphPad Prism 4 software (San Diego, Calif.).

Mouse Ischemic Hindlimb Study

All animal work is conducted in accordance with the Guide for the Care and Use of Laboratory Animals (National Academy of Sciences; Copyright© 1996) and the Institutional Animal Care and Use Committee guidelines at Procter and Gamble Pharmaceuticals. We studied 9-10 week old male C57Bl/6 mice from Charles River Laboratory (Portage, Mich.). The mice are orally dosed with vehicle (aqueous carbonate buffer, 50 mM; pH 9.0) or compound 1 in vehicle at 50 mg/kg or 100 mg/kg. The animals are dosed three times: day 1 at 8 am and 5 pm, day 2 at 8 am. One hour after the first dose, unilateral arterial ligation is performed under anesthesia using isoflurane. The femoral artery is ligated proximal to the origin of the popliteal artery. The contralateral limb underwent a sham surgical procedure. Ligation is performed in an alternating fashion between right and left hindlimbs. Two hours after 8 am dosing on day 2, we obtained blood by ventricular stick while the mice are anesthetized with isoflurane. Serum samples for EPO analysis are obtained using gel clot serum separation tubes. Heart, liver, and gastrocnemius muscles are harvested, snap-frozen in liquid nitrogen, and stored in -80° C. until use.


Mouse Serum EPO Assay

The mouse serum EPO is detected using Mouse Quantikine Erythropoietin ELISA kit from R&D Systems according to manufacturer's instructions.







Mouse Tissue HIF Western Blot Analysis

Tissues from mice stored at -80° C. are powdered with mortar and pestle chilled with liquid nitrogen. Nuclear extracts are prepared using an NE-PER kit (Pierce Biotechnology). For immunoprecipitation, nuclear extract is added to monoclonal antibody to HIF-1 α (Novus, Littleton, Colo.) at a tissue to antibody ratio of 200:1. The suspension is incubated in a conical micro centrifuge tube for 4 hours at 4° C. Protein A/G-coupled agarose beads (40 μ L of a 50% suspension) are then added to the tube. Following overnight tumbling at 4° C., the beads are washed 3 times with ice-cold phosphate buffered saline. The beads are then prepared for SDS-PAGE with 40 μ L of Laemmli sample buffer. Proteins separated on SDS-PAGE are transferred onto nitrocellulose sheets with XCell-II Blot Module system (Invitrogen, Carlsbad, Calif.). The blots are blocked with 5% BSA prior to incubation with a rabbit antibody to HIF-1 α at 1:100 dilution (Novus). The blots are then washed with Tris-buffered saline/Tween-20 buffer and incubated with horseradish peroxidase-conjugated goat anti-rabbit secondary antibody (Pierce, Rockford, Ill.). Blots are developed with the ECL reagent (Amersham, Piscataway, N.J.). Images of blots are captured with an Epson Expression 1600 scanner.






Table XV below provides non-limiting examples of the in vivo response for compounds according to the present disclosure.

Compound	EGLN1 IC ₅₀ μ M	EGLN3 IC ₅₀ μ M	VEGF EC ₅₀ μ M	EPO response
 {[5-(3-chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester	2.8	21.4	9.9 (39)	yes






-continued

Compound	EGLIN1 IC ₅₀ μM	EGLIN3 IC ₅₀ μM	VEGF EC ₅₀ μM	EPO response
 <p>{[5-(4-Chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	3.7	—	>100	ND*
 <p>{[5-(4-Cyano-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	5.8	—	>100	ND
 <p>[(3-Hydroxy-5-(4-methylphenyl)pyridine-2-carbonyl)amino]-acetic acid</p>	0.65	0.18	42.1	yes
 <p>[(3-Hydroxy-4'-methyl-biphenyl-4-carbonyl)-amino]-acetic acid</p>	20	—	>100	yes
 <p>[(3-Hydroxy-5-(4-methylphenyl)-pyridine-2-carbonyl)-amino]-acetic acid methyl ester</p>	4.1	52.3	8.2	ND
 <p>{[5-(2-Chloro-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	6.8	37.5	7.7	yes






-continued

Compound	EGLIN1 IC ₅₀ μM	EGLIN3 IC ₅₀ μM	VEGF EC ₅₀ μM	EPO response
 <p>{[5-(4-Cyanophenyl)-3-hydroxypyridine-2-carbonyl]amino)-acetic acid</p>	0099	0.56	4.3	yes
 <p>{[5-(4-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino)-acetic acid</p>	0.24	0.083	—	yes
 <p>{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino)-acetic acid trifluoroacetic acid salt</p>	0.41	—	7.6	ND
 <p>{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino)-acetic acid</p>	1.1	0.39	—	yes
 <p>{[5-(2-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino)-acetic acid</p>	1.1	0.39	—	yes




-continued

Compound	EGLIN1 IC ₅₀ μM	EGLIN3 IC ₅₀ μM	VEGF EC ₅₀ μM	EPO response
 <p>{[5-(4-Isopropylphenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid</p>	0.44	—	15	—
 <p>{[5-(4-Chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid</p>	0.3	—	>100	—
 <p>{[5-(4-Isopropoxyphenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid</p>	1.6	—	>100	—
 <p>{[5-(2-Methyl-5-chlorophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid</p>	2.5	—	3.3	—
 <p>5-(3-Chlorophenyl)-N-(2-methylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide</p>	1.2	1.3	1.4	yes

-continued

Compound	EGLIN1 IC ₅₀ μM	EGLIN3 IC ₅₀ μM	VEGF EC ₅₀ μM	EPO response
 <p>{[5-(3-Cyano-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	5.1	—	>100	ND
 <p>{[3-Hydroxy-5-[3-(2H-tetrazol-5-yl)-phenyl]-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	2.6	—	>100	ND
 <p>{[5-(3-Cyanophenyl)-3-hydroxypyridine-2-carbonyl]amino}-acetic acid</p>	0.19	—	>100	ND
 <p>{[5-(3-Carbamoyl-phenyl)-3-hydroxy-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	11.2	—	50.1	ND
 <p>{[3-Hydroxy-5-[3-(pyrrolidine-1-carbonyl)-phenyl]-pyridine-2-carbonyl]-amino}-acetic acid methyl ester</p>	2.5	—	>100	ND

-continued

Compound	EGLIN1 IC ₅₀ μM	EGLIN3 IC ₅₀ μM	VEGF EC ₅₀ μM	EPO response
 <p>{(3-Hydroxy-5-[3-(pyrrolidine-1-carbonyl)phenyl]-pyridine-2-carbonyl)amino)-acetic acid</p>	0.3	—	>100	ND
 <p>{(3-Hydroxy-5-[3-(2H-tetrazol-5-yl)phenyl]-pyridine-2-carbonyl)amino)-acetic acid</p>	0.4	—	>100	ND
 <p>{(5-[3-(Cyclopropanecarbonyl-amino)-phenyl]-3-hydroxy-pyridine-2-carbonyl)amino)-acetic acid methyl ester</p>	9.2	—	—	ND

*ND = not determined.

While particular embodiments of the present disclosure have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the disclosure. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this disclosure.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 1

<210> SEQ ID NO 1

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Unknown

<220> FEATURE:

<223> OTHER INFORMATION: HIF-1 alpha peptide corresponding to aa residues 556-574 for use as substrate to measure EGLN-1 (or EGLN-3) activity

<400> SEQUENCE: 1



Asp Leu Asp Leu Glu Ala Leu Ala Pro Tyr Ile Pro Ala Asp Asp Asp
1 5 10 15

Phe Gln Leu

83

What is claimed is:



1. A compound having the formula:



2. A pharmaceutical composition comprising:

- A) the compound according to claim 1; and
 B) one or more pharmaceutically acceptable excipients.


3. A compound having the formula:



4. A pharmaceutical composition comprising:




- A) the compound according to claim 3; and
 B) one or more pharmaceutically acceptable excipients.

5. A compound having the formula:



84

-continued



wherein:

R⁶ is hydrogen, methyl, or ethyl;R^{7a} and R^{7b} are each independently:

- i) hydrogen; or
 ii) methyl.

6. The compound according to claim 5, wherein the compound is:

{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}acetic acid methyl ester;

{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}acetic acid ethyl ester;

5-(3-Chlorophenyl)-N-(2-amino-2-oxoethyl)-3-hydroxypyridin-2-yl amide;

5-(3-Chlorophenyl)-N-(2-methylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide; and

5-(3-Chlorophenyl)-N-(2-dimethylamino-2-oxoethyl)-3-hydroxypyridin-2-yl amide.

7. A pharmaceutical composition comprising:

- A) one or more compounds according to claim 6; and
 B) one or more pharmaceutically acceptable excipients.

8. The composition according to claim 7, wherein the compound is:

{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}acetic acid methyl ester; or

{[5-(3-Chlorophenyl)-3-hydroxypyridine-2-carbonyl] amino}acetic acid ethyl ester.

* * * * *