



(86) Date de dépôt PCT/PCT Filing Date: 2012/11/06
(87) Date publication PCT/PCT Publication Date: 2013/05/10
(45) Date de délivrance/Issue Date: 2021/03/09
(85) Entrée phase nationale/National Entry: 2014/05/06
(86) N° demande PCT/PCT Application No.: US 2012/063670
(87) N° publication PCT/PCT Publication No.: 2013/067544
(30) Priorités/Priorities: 2011/11/06 (US61/556,247);
2011/11/06 (US61/556,241); 2011/11/06 (US61/556,264)

(51) Cl.Int./Int.Cl. *A01N 61/00* (2006.01),
A01N 25/30 (2006.01), *A01N 33/12* (2006.01),
A01N 37/00 (2006.01), *A01N 43/40* (2006.01),
A01N 59/00 (2006.01), *A01P 1/00* (2006.01)
(72) Inventeurs/Inventors:
SOOKRAM, BURT R., US;
VEENSTRA, JOHN W., US
(73) Propriétaire/Owner:
NBIP, LLC, US
(74) Agent: MARKS & CLERK

(54) Titre : COMPOSITIONS ANTIMICROBIENNES COMPRENANT UN BIOCIDÉ, UN TAMPON POUR PH ET UN
SURFACTIF
(54) Title: ANTIMICROBIAL COMPOSITIONS COMPRISING BIOCIDÉ, PH BUFFER AND SURFACTANT

(57) Abrégé/Abstract:

The claimed invention is directed to compositions and methods effective in preventing microbial contamination or reducing microbial count associated with a contaminated surface, comprising a biocidal system comprised of a primary biocide, a pH buffer agent, a surfactant, all in an aqueous based carrier, wherein the compositions may be considered environmentally friendly.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2013/067544 A1

(43) International Publication Date
10 May 2013 (10.05.2013)

- (51) **International Patent Classification:**
A01N 25/00 (2006.01) *A01N 25/30* (2006.01)
- (21) **International Application Number:**
PCT/US2012/063670
- (22) **International Filing Date:**
6 November 2012 (06.11.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/556,264 6 November 2011 (06.11.2011) US
61/556,241 6 November 2011 (06.11.2011) US
61/556,247 6 November 2011 (06.11.2011) US
- (71) **Applicant:** NBIP, LLC [US/US]; 5760 Legacy Drive,
Suite B3-411, Plano, TX 75024 (US).
- (72) **Inventors:** SOOKRAM, Burt, R.; 540 Sheridan Drive,
Palm Harbor, FL 36484 (US). VEENSTRA, John, W.;
5760 Legacy Drive, Suite B3-411, Plano, TX 75024 (US).
- (74) **Agents:** GOPALAKRISHNAN, Lekha et al.; Winstead
PC, P.O. Box 131851, Dallas, TX 75313 (US).

(81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** ANTI-MICROBIAL COMPOSITIONS AND RELATED METHODS

(57) **Abstract:** The claimed invention is directed to compositions and methods effective in preventing microbial contamination or reducing microbial count associated with a contaminated surface, comprising a biocidal system comprised of a primary biocide, a pH buffer agent, a surfactant, all in an aqueous based carrier, wherein the compositions may be considered environmentally friendly.



WO 2013/067544 A1

ANTIMICROBIAL COMPOSITIONS COMPRISING
BIOCIDE, PH BUFFER AND SURFACTANT

[0001]

5

FIELD OF THE INVENTION

[0002] The present disclosure relates to compositions and methods that are effective
10 in preventing microbial contamination or reducing the microbial count on or in a
contaminated surface.

BACKGROUND OF THE INVENTION

[0003] Mold, mildew, fungi, viral and bacterial (microbial) contamination is
undesirable on many types of surfaces. Control of such microbial contamination has largely
15 been based on the use of biocides that have limited efficacy and may have a harmful effect on
the environment. Therefore, the use of more effective biocides or derivatives thereof that
may be more effective in microbial control is desirable. An additional benefit may be that
compositions may be environmentally friendly which is also desirable in certain applications.

The anti-microbial properties of quaternary ammonium compounds are generally
20 known. Such compounds have been used extensively in cleaning compositions for domestic
and industrial applications. However, it would be preferable to use smaller amounts of these
compounds in cleaning compositions since they can be harmful to the environment and
minimally effective. Therefore, it would be desirable to create formulations comprising
compounds that exhibit anti-microbial properties and may at the same time be environmental
25 friendly.

SUMMARY OF THE INVENTION

[0004] The claimed invention relates to compositions and methods that are effective in preventing microbial contamination or reducing the microbial count on or in a contaminated surface.

5 [0005] The disclosed compositions comprise a biocidal system comprising a primary biocide, a pH buffer, and a surfactant in an aqueous based carrier.

[0006] An embodiment of the invention is directed to a composition the composition comprising:

- 10 a. from about 0.01% to about 20.0 % by weight of a biocidal system comprising:
- i. from about 0.01% to about 25% by weight of a primary biocide; and
- 15 ii. at least about 0.01% to about 25% by weight of a pH buffer, where the pH buffer is a biocidal, dermal, non-corrosive acid composition, having a proton count range of from 1.0×10^{20} to 9.9×10^{26} , an embodied conductivity range of from 250 mV to 1500mV and has a pH of less than 2.0 when the pH buffer is present at a concentration of 0.1% by weight.; and
- b. from about 0.01% to 4.0% by weight of a surfactant; and
- 20 c. the balance being an aqueous based carrier.

DETAILED DESCRIPTION OF EXEMPLARY EMBODIMENTS

[0007] In this specification and in the claims that follow, reference will be made to a number of terms that shall be defined to have the following meanings:

25 [0008] Throughout this specification, unless the context requires otherwise, the word “comprise,” or variations such as “comprises” or “comprising,” will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

[0009] It must be noted that, as used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a carrier” includes mixtures of two or more such carriers, and the like.

5 [00010] Ranges may be expressed herein as from “about” one particular value, and/or to “about” another particular value. When such a range is expressed, another aspect includes from the one particular value and/or to the other particular value. Similarly, when values are expressed as approximations by use of the antecedent “about,” it will be understood that the particular value forms another aspect. It will be further understood that
10 the endpoints of each of the ranges are significant both in relation to the other endpoint and independently of the other endpoint.

[00011] “Admixture” or “blend” as generally used herein means a physical combination of two or more different components

[00012] By “contacting” is meant an instance of applying a composition to a
15 contaminated surface.

[00013] “Contamination” is used herein to describe microbiological intrusions, such as the presence of toxins or pathogens in or on the surface of any material.

[00014] “Controlled release” as used herein means the use of a material to regulate the release of another substance.

20 [00015] “Effective amount” as used herein means an amount of a composition as disclosed herein, effective at dosages and for periods necessary to achieve the desired result.

[00016] “Environmentally friendly” as used herein refers to green, organic or natural compositions that are minimally harmful to the environment.

25 [00017] “Excipient” is used herein to include any other compound that may be contained in or on the microparticle that is not a therapeutically or biologically active compound. As such, an excipient should be pharmaceutically or biologically acceptable or relevant (for example, an excipient should generally be non-toxic to the subject). “Excipient” includes a single such compound and is also intended to include a plurality of excipients.

[00018] “Microbiological” as used herein refers to any inclusion or growth of harmful microorganisms such as mold, mildew, viral or bacterial contamination.

[00019] “Microbial Count” as used herein refers to the amount or number of microbiological contaminants present on any surface.

5 [00020] “Optional” or “optionally” means that the subsequently described event or circumstance can or cannot occur, and that the description includes instances where the event or circumstance occurs and instances where it does not.

[00021] “Primary biocide” is used herein to refer to compositions that are biologically active against microbial contaminants.

10 [00022] “Primary pathogen” is used herein to refer to mold, mildew, bacteria, viruses or other microorganisms that can cause contamination on a surface.

[00023] By “sufficient amount” and “sufficient time” means, an amount and time needed to achieve the desired result or results, e.g., control and/or prevention of microbial contamination.

15 [00024] “Surface” as used herein refers to the object that contains the microbiological contamination. The term surface can apply to the entire object, a portion or layer of the object, and down to the molecular structure of the object.

[00025] A “weight percent” of a component, unless specifically stated to the contrary, is based on the total weight of the formulation or composition in which the
20 component is included.

[00026] The present disclosure addresses solutions to several unmet needs as defined below:

1. Providing compositions effective in killing one or more bacterial pathogens. Non-limiting examples include Salmonella, Pseudomonas aeruginosa,
25 Staphylococcus aureus, Staphylococcus epidermidis, Streptococcus agalactiae, Pseudomonas species. Campylobacter jejuni, Escherichia coli, (E. coli) and Klebsiella pneumonia.

2. Providing compositions effective in killing one or more viral pathogens. Non-limiting examples which includes Cryptosporidium, Giardia lamblia, poliomyelitis, meningitis, and gastroenteritis, Rotaviruses, Norwalk, Norovirus, Hepatitis A, parasites.

5 3. Providing compositions effective in killing one or more mold and mildew spores pathogens. Non-limiting examples which includes Trichophyton mentagrophytes, Guignardia citricarpa, and Colletotrichum acutatum.

4. Providing compositions that are a replacement for chlorine and chlorine based products such as sodium hypochlorite based treatments.

10 **Treatment Compositions**

 [00027] The inventive composition generally includes a biocide system, which includes a primary biocide and a pH buffering component. The primary biocide is selected according to the nature of the microorganisms sought to be controlled, for example, a quaternary ammonium salt. The pH buffer is mechanically fused so as to aid in the delivery
15 of the primary biocide without causing damage to the surface being treated. By way of example, a suitable pH buffer is disclosed in U.S. Patent No. 7,824,524, or in a U.S. Patent Application No. 13/346,160, entitled "Reactive, non-corrosive and dermal-friendly composition and methods for manufacturing". In embodiments of the invention, the pH buffer is chosen for compatibility with the primary biocide.

20

 [00028] An embodiment of the invention relates to compositions that are effective in preventing or reducing microbial count, the composition comprising:

 a. from about 0.01% to about 20.0 % by weight of a biocidal system
25 comprising:

 i. from about 0.01% to about 25% by weight of a primary biocide; and

 ii. at least about 0.01% to about 25% by weight of a pH buffer, where the pH buffer is a biocidal, dermal, non-corrosive acid

composition, having a proton count range of from 1.0×10^{20} to 9.9×10^{26} , an embodied conductivity range of from 250 mV to 1500mV and has a pH of less than 2.0 when the pH buffer is present at a concentration of 0.1% by weight.; and

- 5 b. from about 0.01% to 4.0 % by weight of a surfactant; and
- c. the balance being an aqueous based carrier.

[00029] However, other non-limiting embodiments and combinations are possible as further disclosed herein.

Biocidal System

- 10 [00030] The disclosed compositions comprise a biocidal system. The biocidal system comprises a primary biocide and a pH buffer. In certain embodiments of the invention, the disclosed compositions comprise from about 0.01% to about 20.00% by weight of a biocidal system. In other embodiments, the biocidal system comprises at least about 25% by weight of a primary biocide and at least about 25% by weight of a pH buffer component.
- 15 The pH buffer is chosen for compatibility with the primary biocide.

Primary Biocide

- [00031] A first group of suitable biocides include quaternary ammonium compounds chosen from (C₁₂-C₁₄ alkyl)(C₁-C₂ dialkyl)benzyl ammonium salts, *N*-(C₁₂-C₁₈ alkyl)heteroaryl ammonium salts, and *N*-[(C₁₂-C₁₄ alkyl)(C₁-C₂ dialkyl)]heteroarylalkylene
- 20 ammonium salts. Non-limiting examples of the (C₁₂-C₁₄ alkyl)(C₁-C₂ dialkyl)benzyl ammonium salts include (C₁₂-C₁₄ alkyl)dimethyl-benzyl ammonium chloride, (C₁₂-C₁₄ alkyl)dimethylbenzyl ammonium bromide, and (C₁₂-C₁₄ alkyl)dimethylbenzyl ammonium hydrogen sulfate. Non-limiting examples of the *N*-(C₁₂-C₁₈ alkyl)heteroaryl ammonium salts include cetyl pyridinium chloride, cetyl pyridinium bromide, and cetyl pyridinium hydrogen
- 25 sulfide. For the *N*-(C₁₂-C₁₈ alkyl)heteroaryl ammonium salts other anions can be used.

[00032] Further examples of quaternary ammonium compounds suitable for use as the primary biocides include cetyltrimethylammonium chloride, stearyltrimethylammonium chloride, isostearyltrimethylammonium chloride, lauryltrimethylammonium chloride, behenyltrimethyl-ammonium chloride,

octadecyltrimethylammonium chloride, cocoyltriethylammonium chloride, cetyltrimethylammonium bromide, stearyltrimethylammonium bromide, lauryltrimethylammonium bromide, isostearyl lauryldimethylammonium chloride, dicetyldimethylammonium chloride, distearyldimethylammonium chloride, dicocoyldimethylammonium chloride, gluconamidopropyl dimethylhydroxyethylammonium chloride, di[polyoxyethylene(2)]oleylmethylammonium chloride, dodecyldimethylethylammonium chloride, octyldihydroxyethylmethylammonium chloride, tri[polyoxyethylene(5)]stearyl ammonium chloride, polyoxypropylenemethyldiethylammonium chloride, lauryldimethyl(ethylbenzyl)ammonium chloride, behenamidopropyl-N,N-dimethyl-N-(2,3-dihydroxypropyl) ammonium chloride, tallowdimethylammoniopropyltrimethylammonium dichloride, and benzalconium chloride.

[00033] A second group of suitable biocides include copper, zinc, silver, salts of chlorides, chlorites, perchlorates, hypochlorites, sulfates, sulfites, bisulfates, bisulfites, nitrates, nitrites and hydroxides. Also colloid metal such as silver, gold, copper and zinc have superior biocidal properties. Colloidal silver, gold, copper and zinc are extracted and created as ultrafine (0.010-0.001 micron) particles.

[00034] A third group of suitable biocides include organic acids which are safe under the FDA GRAS guidelines for food production yet still effective in controlling bacteria. The basic principle action of organic acids on bacteria is that non-dissociated organic acids can penetrate a bacterium cell wall and cause disruption due to the fact it cannot tolerate a wide internal and external pH gradient. This will cause the osmotic pressure inside the cell to increase which state is incompatible with bacterial survival.

[00035] A first group of suitable organic acids are Lactic acid, Acetic acid, Formic acid, Fumaric acid, Citric acid, Oxalic acid, Adipic acid and Uric acid.

[00036] A second group of suitable organic acids are the carboxylic acids, whose acidity is associated with their carboxyl group -COOH. Sulfonic acids, containing the group -SO₂OH, are relatively stronger acids. The relative stability of the conjugate base of the acid determines its acidity. In some biological systems more complex organic acids such as L-lactic, citric, and D-glucuronic acids are formed. These use the hydroxyl or carboxyl group.

[00037] A third group of suitable organic acids are Humic, Sebacic, Stearic, Gallic, Palmitic, Caffeic, Glyoxylic, Fulvic, Carnosic, Anthranilic, Ellagic, Lipoic, Chlorogenic, Rosmarinic, Phosphoric, Methacrylic, Oleanic, Nitrohumic, Florocinnamic, Hexaflorosilicic, Hydrofluoric, Hydroxycitric and Silicofluoric.

5 [00038] A fourth group of suitable organic acids is fruit acids. The acids in fruits are chiefly acetic, malic, citric, tartaric, oxalic, and in some instances boric. Malic acid is present in apples, pears, currants, blackberries, raspberries, quince, pineapple, cherries, and rhubarb. Citric acid is found in lemons, oranges, grapefruit, lemons, limes, quince, gooseberry, strawberry, raspberry, currant, and cranberry. Tartaric acid occurs in grapes.
10 Boric acid is found in many fresh fruits and vegetables. Mandelic acid is present in almonds.

[00039] A fifth group of suitable organic acids is beta hydroxy acids, which is a type of phenolic acid. Salicylic acid is a colorless crystalline organic acid whose main active ingredient obtained from this source is a monohydroxybenzoic acid.

[00040] A sixth group of suitable organic acids is a class of products that break
15 biofilm. Biofilms are the protective layer / barrier that surround bacteria. Some species are not able to attach to a surface on their own but are often able to anchor themselves to the matrix or the bacteria cells. It is during this colonization that the cells are able to communicate via its quorum sensing ability. Once colonization has begun, the biofilm grows through a combination of cell division and recruitment. The final stage of biofilm formation
20 is known as development and is the stage in which the biofilm is established and may only change in shape and size. The development of a biofilm may allow for an aggregate cell colony to be increasingly resistant. A biofilm's hard protective surface can be broken by *Lactobacillus sc Nisin* which is produced by fermentation using the bacterium *Lactococcus lactis*. This is obtained from the culturing of *Lactococcus lactis* on natural substrates, such as
25 milk or dextrose, and is not chemically synthesized. This is a peptide which is produced by the food grade dairy starter bacterium *Lactococcus lactis*.

[00041] A seventh group of suitable organic acids is natural enzymes. Enzymes are proteins that catalyze chemical reactions and range from just 62 amino acid residues. Typically these are protease, lipase, diastase and cellulase enzymes. Enzymes are usually
30 very specific as to which reactions they catalyze and the substrates that are involved in these reactions. The shape, charge and hydrophilic/hydrophobic nature characterize the enzymes.

pH Buffer

[00042] The pH buffer is a biocidal, dermal, non-corrosive acid composition, having a proton count range of from 1.0×10^{20} to 9.9×10^{26} , an embodied conductivity range of from 250 mV to 1500mV and a 0.1% solution of the composition having a pH of under 2.0.

Surfactant

[00043] The disclosed compositions comprise from about 0.05% to about 5.0 % by weight of a cationic surfactant having an HLB of from about five to about 30. One aspect of the disclosed compositions comprises a cationic or ionic surfactant having an HLB of from about 12 to about 18. A further aspect of the disclosed compositions comprises a cationic or ionic surfactant having an HLB of from about 13 to about 16. Another embodiment of the disclosed compositions comprise from about 0.1% to about 4.0 % by weight of a cationic or ionic surfactant.

In some embodiments, the surfactant is chosen from about 60% to about 99%, based on total polymer weight, of a quaternary diallyl dialkyl ammonium monomer, wherein alkyl groups are independently selected from alkyl groups of 1 to 18 carbon atoms, C_{1-4} alkyl, and wherein said quaternary diallyl dialkyl ammonium monomer's counterion is selected from the group consisting of conjugate bases of acids having an ionization constant greater than 10^{-13} , selected from the group consisting of fluoride, chloride, bromide, hydroxide, nitrate, acetate, hydrogen sulfate and primary phosphates.

In some embodiments, the surfactant is chosen from about 0.01% to about 4.0%, based on total polymer weight, of an anionic monomer selected from the group consisting of acrylic acid and methacrylic acid, wherein the average molecular weight of said polymer ranges from about 50,000 to about 10,000,000, as determined by gel permeation chromatography.

In another embodiment, the surfactant is chosen from a combination of one or more polymer bases, such as glycerol, glycerine, sortibal aloe, polyglycols, polyethylene glycol, polyoxyethylene or polyethylene oxide, in combination with ethoxylated partial glyceride fatty acid esters such as chain esters, ethoxylated partial glyceride fatty acid esters, protein derivatives, lanolin and lanolin derivatives, and fatty alcohol ethoxylates, emollient oils, fatty acids, fatty alcohols and their esters.

polyoxyethylene(branched) nonylphenyl ethers sold under the tradename TRITON™ N-101, polyoxyethylene(12) nonylphenyl ether sold under the tradename IGEPAL™ CO-720, polyoxyethylene(12) isooctylphenyl ether sold under the tradename IGEPAL™ CA-720, polyoxyethylene(40) nonylphenyl ether sold under the tradename IGEPAL™ CO-890, and
 5 polyoxyethylene(40) isooctylphenyl ether sold under the tradename TRITON™ X-405.

[00046] Another category of cationic or ionic surfactants for use in the disclosed compositions are polyoxyethylene sorbitan mono-, di-, and tri-(C₁₂-C₁₈)-alkanoates, non-limiting examples of which include polyoxyethylene(20) sorbitan trioleate sold under the tradename TWEEN™ 85, polyoxyethylene(20) sorbitan monooleate sold under the
 10 tradename TWEEN™ 80, polyoxy-ethylene(20) sorbitan monostearate sold under the tradename TWEEN™ 60, polyoxyethyl-ene(20) sorbitan monopalmitate sold under the tradename TWEEN™ 40, and polyoxyethyl-ene(20) sorbitan monolaurate sold under the tradename TWEEN™ 20.

[00047] A further category of cationic or ionic surfactants for use in the
 15 disclosed compositions are polyoxyethylene C₉-C₂₀ alkyl ethers, non-limiting examples of which include ethoxylate alcohols having the formula: RO(CH₂CH₂O)_mH, wherein R is a linear or branched alkyl group having from 6 to 20 carbon atoms and m is an integer of about 2 to about 20. On example of suitable ethoxylate alcohol surfactants are the NEODOL™ ethoxylated alcohols from Shell Chemicals. Non-limiting examples of suitable ethoxylated
 20 alcohols include NEODOL™ 91-5, NEODOL™ 91-6, NEODOL™ 91-8, NEODOL™ 91-9, NEODOL™ 23-6.5, NEODOL™ 25-5, NEODOL™ 25-7, NEODOL™ 25-9, NEODOL™ 25-12, NEODOL™ 45-7, and NEODOL™ 135-7.

Carriers

[00048] The balance of the disclosed compositions comprises a carrier. The
 25 carrier can be any suitable material that can dissolve the active ingredients and co-ingredients and deliver the biocidal system to the infected areas. Water is a convenient carrier for liquid embodiments of the disclosed composition.

Adjunct Ingredients

[00049] The disclosed compositions can further comprise one or more dyes at levels of from about 0.001% to 0.5%. Non-limiting examples of suitable dyes are Alizarine Light Blue B (C.I. 63010), Carta Blue VP (C.I. 24401), Acid Green 2G (C.I. 42085),
5 Astrogen Green D (C.I. 42040), Supranol Cyanine 7B (C.I. 42675, Maxilon Blue 3RL (C.I. Basic Blue 80), Drimarine Blue Z-RL (C.I. Reactive Blue 18), Alizarine Light Blue H-RL (C.I. Acid Blue 182), FD&C Blue No. 1 and FD&C Green No. 3. (See U.S. Patent No. 4,248,827 and U.S. Patent No. 4,200,606.

[00050] Other colors which can be Lakes that may be used are FD&C Blue No.
10 1 – Brilliant Blue FCF, (blue shade), FD&C Blue No. 2 – Indigotine, (dark blue shade), FD&C Green No. 3 – Fast Green FCF, (turquoise shade), FD&C Red No. 40 – Allura Red AC, (red shade), FD&C Red No. 3 – Erythrosine, (pink shade, commonly used in glacé cherries), FD&C Yellow No. 5 – Tartrazine, (yellow shade), FD&C Yellow No. 6 – Sunset Yellow FCF, E110 (orange shade)

15 [00051] Another adjunct ingredient suitable for use in the compositions disclosed herein includes fragrances.

Formulations

Table I

Ingredients	Amount (wt%)
Malic Acid	0.15-1.50
pH buffer	0.05 – 2.00
Thickener	0.50- 4.00
Carrier	balance
Adjunct Ingredients	trace

Table II

Ingredients	Amount (wt%)
Cetyl pyridinium chloride	0.10-1.0
pH buffer	0.05-2.0
PEG 6	0.35-5.0
Thickener	0.25 – 4.0
Carrier	balance
Adjunct Ingredients	trace

[00052] Table I represents an environmentally friendly composition formulated in accordance with embodiments of the invention comprising an organic acid in place of a quaternary ammonium salt.

5 Methods of Use

[00053] The disclosed compositions can be used in various applications. The application route and dosage regimen is dictated by the type and count of the microbial contaminant. As an example of possible applications of the invention, the compositions can be used on a variety of contaminated surfaces in order to reduce infection or contamination to humans and animals coming in contact with that surface. The compositions can be applied as a cleanser, scrub (cleanser with abrasive properties), spray, foam, fog, or gel.

[00054] An effective amount may vary according to factors known in the art, such as the type of microbial contaminant, the microbial count and the type of surface being treated. Although particular dosage regimes may be described in examples herein, a person skilled in the art would appreciate that the dosage regime may be altered to provide optimum therapeutic response. For example, several divided doses may be administered daily or the dose may be proportionally reduced as indicated by the exigencies of the contaminated surface. In addition, the compositions of the present disclosure can be administered as frequently as necessary to achieve desired results.

Working Examples

[00055] The following procedures were used to evaluate the disclosed compositions against various microbiologically contaminants. The composition tested in each of the below examples comprised 0.25 wt% malic acid, 0.10 wt% pH buffer, and 0.5 wt% thickener. The results below further indicate the effectiveness of the disclosed compositions.

[00056] Bacterial testing was done using *Staphylococcus aureus* ATCC #6538. The results are listed in Table A. The reduction in bacterial growth was a 99.9999 %.

Table A

Species	Control units	Results units
<i>Staphylococcus aureus</i>	7.9×10^7 cfu/ml	2.5 cfu/ml

10

[00057] Bacterial testing was completed at Biological Consulting Services of North Florida, Inc. on *E. coli* (ATCC 15597), *Salmonella enterica* (ATCC BAA-711), and Methicillin Resistant *Staphylococcus aureus* (MRSA; BAA-44). The results are listed in Table B.

15 Table B

Sample	Control units (cfu/ml)	Results units (cfu/ml)
<i>E. coli</i>	9.3×10^5	<0.5
<i>S. enterica</i>	1.1×10^6	<0.5
MRSA	1.0×10^6	<0.5

[00058] Viral tests were completed at Biological Consulting Services of North Florida, Inc. on the following on the following Orthomyxviridie virus ATCC type CCL-34, Influenza A/Equi 2(ATCC VR517) Poliovirus 1 (Chat; ATCC VR-1562), and Rhinovirus 39 (ATCC VR-340). The results of the viral tests are shown in Table C.

Table C

Sample	Control units	Results units
Orthomyxviridie virus	6.2×10^6	<200 cfu/ml
Influenza A (H1N1)	3.1×10^4	<200 cfu/ml
Poliovirus 1	1.6×10^5	2.1×10^1
Rhinovirus 39	6.7×10^5	< 200 cfu/ml

[00059] Mold and fungi tests were completed at Biological Consulting Services of North Florida, Inc. on the following on the following Trichophyton mentagrophytes ATCC 4807, Guignardia citricarpa ATCC 26254 , and Colletotrichum acutatum (ATCC 38689). The

5 results of the mold and fungi tests are shown in Table D.

Table D

Sample	Control units (cfu/ml)	Results units (cfu/ml)
Trichophyton mentagrophytes	1.1×10^4	< 1.0
Guignardia citricarpa	1.9×10^4	< 1.0
Colletotrichum acutatum	1.5×10^4	< 1.0

[00060] While particular embodiments of the present disclosure have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the disclosure. It is therefore intended to cover in the appended claims all such changes and

10 modifications that are within the scope of this disclosure.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A composition for preventing microbiological contamination or reducing the microbial count associated with a contaminated surface comprising:
 - a. from 0.01% to about 20.0% by weight of a biocidal system comprising:
 - i. from 0.01% to 25% by weight of a primary biocide where the primary biocide is cetyl pyridinium chloride or malic acid; and
 - ii. 0.01% to 25% by weight of a pH buffer, where the pH buffer is a biocidal, dermal, non-corrosive acid composition having a proton count range of from 1.0×10^{20} to 9.9×10^{26} , an embodied conductivity range of from 250 mV to 1500mV and has a pH of less than 2.0 when the pH buffer is present at a concentration of 0.1% by weight;
 - b. from 0.01% to 4.0% by weight of a surfactant where the surfactant is an ionic surfactant having an HLB of from 12 to 18 or cationic surfactant having an HLB of from about 5 to about 30; and
 - c. the balance being an aqueous based carrier.
2. The composition according to claim 1, where the surfactant is a combination of:
 - (i) a polymer base which is glycerol, glycerine, sorbitol aloe, a polyglycol, polyethylene glycol, polyoxyethylene or polyethylene oxide, or any combination thereof;
 - (ii) an ethoxylated partial glyceride fatty acid ester which is a chain ester, protein derivative, lanolin or a lanolin derivative, or any combination thereof; and
 - (iii) at least one fatty alcohol ethoxylate, emollient oil, fatty acid, fatty alcohol or their esters.