




**ΚΥΠΡΙΑΚΟ ΓΡΑΦΕΙΟ ΔΙΠΛΩΜΑΤΩΝ
ΕΥΡΕΣΙΤΕΧΝΙΑΣ
THE PATENT OFFICE OF CYPRUS**

**ΑΡΙΘΜΟΣ ΔΗΜΟΣΙΕΥΣΗΣ CY1616
PUBLICATION NUMBER**

ΑΡΙΘΜΟΣ ΔΗΜΟΣΙΕΥΣΗΣ
ΓΡΑΦΕΙΟΥ ΔΙΠΛΩΜΑΤΩΝ ΕΥΡΕΣΙΤΕΧΝΙΑΣ
ΗΝΩΜΕΝΟΥ ΒΑΣΙΛΕΙΟΥ
UK PATENT OFFICE
PUBLICATION NUMBER GB2181348

Το έγγραφο που παρουσιάζεται πιο κάτω καταχωρήθηκε στο «Γραφείο Διπλωμάτων Ευρεσιτεχνίας» στην Αγγλία σύμφωνα με το Νόμο Κεφ. 266 πριν την 1^η Απριλίου 1998. Δημοσίευση έγινε μετέπειτα από το Γραφείο Διπλωμάτων Ευρεσιτεχνίας του Ηνωμένου Βασιλείου μόνο στην Αγγλική γλώσσα.

**The document provided hereafter was filed at "The Patent Office"
in England under the law CAP.266 before the 1st of April 1998.
It was published afterwards by the UK patent office only in English.**

(12) UK Patent Application (19) GB (11) 2 181 348 (13) A

(43) Application published 23 Apr 1987

(21) Application No **8623048**

(22) Date of filing **25 Sep 1986**

(30) Priority data

(31) **8504721**

(32) **11 Oct 1985**

(33) **SE**

(51) INT CL⁴
A61K 9/14

(52) Domestic classification (Edition I)
A5B 180 422 423 42Y 431 43Y 586 58Y 800 828 829 832
835 G L
U1S 2415 A5B

(56) Documents cited
None

(58) Field of search
A5B
Selected US specifications from IPC sub-class A61K

(71) Applicant
Aktiebolaget Hassle,

(Incorporated in Sweden),

S—431 83 Molndal, Sweden

(72) Inventors
Ulf Erik Jonsson,
John Anders Sandberg,
John Albert Sjogren

(74) Agent and/or Address for Service
J. A. Kemp & Co., 14 South Square, Gray's Inn, London
WC1R 5EU

(54) **Metoprolol preparation**

(57) A controlled release preparation contains a number of beads comprising a salt of metoprolol as the main soluble component, the beads being coated with a polymeric membrane containing derivatives of cellulose without protolysable groups so that at least 75% of the dose of metoprolol is released within 20 hours virtually independent of the pH in the range 1—8. The beads may be filled into hard gelatin capsules or compressed into tablets.

Formulae in the printed specification were reproduced from drawings submitted after the date of filing, in accordance with Rule 20(14) of the Patents Rules 1982.

FIGURE 1 . IN VITRO RELEASE OF METOPROLOL
RATE PLOT (%/h) VERSUS TIME




FIGURE 2 . MEAN ($n=10$) PLASMA CONCENTRATIONS OF METOPROLOL AFTER SINGLE-DOSE ADMINISTRATION. DOSE: METOPROLOL CORRESP. 200 MG TARTRATE SALT





FIGURE 3 . MEAN $\pm$ SEM (n=12) REDUCTION OF EXERCISE HEART RATE ON DAY 5 AFTER ONCE DAILY ADMINISTRATION DOSE: METOPROLOL CORRESP. 100 MG TARTRATE SALT



SPECIFICATION


New pharmaceutical preparation

FIELD OF THE INVENTION

The present invention is related to a new pharmaceutical preparation with controlled release of metoprolol, to a method for the manufacture of such a preparation and to a method for treatment of cardiovascular disorders using the new pharmaceutical preparation.

BACKGROUND OF THE INVENTION

Metoprolol, which has the structural formula



is known from e.g. DE—2 106 209. The drug, which is a β -adrenoceptor antagonist has preferably been used as a salt, e.g. the tartrate.

Metoprolol blocks the adrenergic stimulation of the heart and thus reduces the oxygen demand of the cardiac tissue. Apparently, this explains its beneficial effects in angina pectoris and cardioprotective action in myocardial infarction. In addition metoprolol normalizes blood pressure in a large proportion of patients with arterial hypertension which probably is due to an additional action on the control of peripheral resistance to blood-flow.

For patients suffering from cardiovascular disorders it is advantageous to have a constant concentration of the administered drug in the blood. Thus, a controlled release of the drug over a long period of time is desirable. According to the most common treatment, the patients are ordered one fast dissolving tablet twice a day. This gives a varying concentration with high peak and trough values of the drug during the day.

For dosage once a day metoprolol has been incorporated in controlled release tablets of the insoluble matrix type, e.g. Durules®. However the drug release from the matrix tablets is not satisfying as about 50 per cent of the dose is released within a few hours after administration. It has thus been a demand to find a way to obtain a drug preparation having a more constant controlled release of the active component for approximately 20—24 hours, whereby smoother blood concentration and effect profiles will be obtained over the entire dosage interval.

A drug delivery system named Oros® may be used to obtain a controlled release of e.g. metoprolol for once daily dosage. The system is described in U.S. Patent 4 036 227 and in a supplement to British Journal of Clinical Pharmacology (1985), 19, 695—765 by Theeuwes F et al. Oros® is a single-unit system consisting of an osmotically active core composed mainly of the drug substance surrounded by a semipermeable membrane through which a single small opening is drilled. The release of the drug from the system remains constant as long as a steady osmotic pressure is maintained across the membrane. 50—60% of the total content of the drug is released at a constant rate.

In SE—A—8 400 085 it has been proposed to prepare an enteric coated product, containing e.g. metoprolol, and with slow release of the active compound close to the colon. Such a preparation does not give the constant and slow pH-independent release of metoprolol, which the preparation according to this invention gives.

Depot preparations comprising a large number of smaller units are also known e.g. from EP 13263. This patent describes pharmaceutically indifferent cores covered by the active compound. The cores are made of soluble material e.g. lactose.

There are advantages of a depot preparation comprising a large number of smaller units, each of which releases the drug at a controlled rate over a depot preparation consisting of one single unit, e.g. a matrix tablet or a tablet surrounded by a coating, which controls the release. It is for example possible to obtain a reproducible emptying of units from the stomach when the particles used are smaller than 1—2 mm. Cf. Bogentoft C et al.: Influence of food on the absorption of acetyl-salicylic acid from enteric-coated dosage forms. Europ J Clin Pharmacol 1978, 14, 351—355. Dispersion over a great area in the gastrointestinal canal gives a more reproducible total time of the passage which is of advantage for the absorption. Cf. Edgar B, Bogentoft C and Lagerström P O: Comparison of two enteric-coated acetylsalicylic acid preparations by monitoring steady-state levels of salicylic acid and its metabolites in plasma and urine. Biopharmaceutics & Drug Disposition 1984, 5, 251—260. In addition a multiple unit preparation is preferable to one single drug

unit as the dose is spread out in the intestine. The risk of local irritation and accumulation of several doses due to constriction in the alimentary canal is also considerably lower.

A further advantage with a multiple unit preparation is that it may be divided into smaller portions all having the same absorption properties. This makes it possible to obtain greater flexibility in selecting the size of the dose.

OUTLINE OF THE INVENTION

The present invention is related to a preparation containing metoprolol as active ingredient and having a controlled rate of drug release during at least 15 hours. By making a preparation containing a large number of small compact particles all comprising a salt of metoprolol as the main soluble component and coated with a polymeric membrane containing derivatives of cellulose without protolysable groups it has been possible to prepare a suitable dosage form having a controlled rate of release of metoprolol, virtually independent of pH, during 16—24 hours.

The small particles, beads, containing metoprolol have a size of 0.25—2 mm, preferably 0.35—1.0 mm.

The beads may contain metoprolol alone or may consist of insoluble cores coated with metoprolol. The beads have a very high content of metoprolol, preferably 95—100 w/w% of the soluble part of the beads. The insoluble cores have a size of 0.1—1.0 mm, preferably 0.15—0.3 mm. Examples of insoluble cores according to the invention are silicon dioxide and small particles of glass.

The beads according to the invention are compact, which means that their porosity is less than 15 per cent.

As can be seen from figure 1 the new preparation is characterized in that at least 75% of the metoprolol is released within 20 hours and at least 50% of the dose of metoprolol is released at the rate 3—7 w/w %/hour.

Metoprolol used in the preparation may be in the form of the racemate, or one of the enantiomers, namely the S-isomer. (Preferably, the concentration of the S-isomer should be at least 95% in relation to the R-form.)

Suitable soluble salts of metoprolol have a solubility less than 600 mg/ml in water at 25°C, preferably 30—600 mg/ml in water at 25°C. Examples of suitable salts are salts formed of organic carboxylic acids, preferably of low molecular weight. Especially preferred are the succinate, fumarate or benzoate of racemic metoprolol and the benzoate or sorbate of the S-enantiomer of metoprolol.

Very soluble salts, e.g. tartrate, hydrochloride are less suitable according to the present invention. Examples of suitable polymeric materials are ethyl cellulose or a mixture of ethyl cellulose with hydroxypropylmethyl cellulose, hydroxypropyl cellulose, Eudragit RL or Eudragit RS.

Ethyl cellulose is available in variants having different grades of viscosity. According to the invention it is suitable to use ethyl cellulose having a viscosity between 10—50 cps, but also other types of ethyl cellulose may be used.

Eudragit® is the trade name for a number of filmcoating substances on an acrylic resin basis produced by Röhm Pharma. Eudragit RL and RS are copolymers synthesized from acrylic and methacrylic acid esters with a low content of quaternary ammonium groups. The molar ratio of these ammonium groups to the remaining neutral (meth)acrylic acid esters is 1:20 for Eudragit RL and 1:40 for Eudragit RS, resulting in different permeability characteristics.

Plasticizers and/or pigments may be added to the polymeric layer in order to improve the technical properties of the layer or change the permeability of the coating. Examples of suitable plasticizers are citrate esters, acetylated monoglycerides and glycerinetriacetate, especially preferred is acetyltributylcitrate.

The polymeric membrane is made of one or more polymers and gives a membrane with virtually pH-independent permeability characteristics within the pH range 1.0—8.0.

Each coated bead of metoprolol according to this invention forms an individual controlled release unit, releasing the drug at a predetermined rate. Therefore, the coated beads according to this invention make it possible to formulate and administer the preparation in different dosage forms. They can be filled into e.g. hard gelatin capsules or sachets or compressed to tablets and still give the desired plasma concentration profile and duration of the effect.

When the small coated particles of metoprolol are tableted they are mixed with additives e.g. microcrystalline cellulose such as Avicel®, which improves the tableting properties and facilitates the disintegration of the tablet, whereby the individual beads are liberated.

The invention makes it possible to formulate pharmaceutical dosage forms which can be given once daily and still produce almost constant concentrations of the drug in the blood during the entire dosage interval until the next dose is administered.

PREPARATION

A process for the manufacture of the controlled release preparation represents a further aspect of the invention. After the initial forming of the beads containing metoprolol, the beads obtained are coated with the polymeric layer described in the examples. The polymeric mixture is dissolved in an organic solvent such as ethanol, isopropyl alcohol and/or methylene chloride. The spraying can be carried out in a coating pan, but is preferably carried out in a fluidized bed. Ethyl cellulose can also be applied from an aqueous dispersion (latex).

The preparation according to the invention is particularly advantageous in the treatment of cardiovascular disorders, and a method for the treatment of such conditions represents a further aspect of the invention.

The invention is described in detail in the following examples:

5	EXAMPLE 1	EXAMPLES	5
	Metoprolol fumarate	1440 g	
	Methylene chloride	9618 g	
	Ethanol 95%	3888 g	
10	SiO ₂ (0.15—0.25 mm)	375 g	10
	<u>Polymeric layer</u>		
	Ethyl cellulose 10 cps	265.6 g	
	Hydroxypropylmethyl cellulose	58.4 g	
	Acetyltributylcitrate	36.0 g	
15	Methylene chloride	6141 g	15
	Isopropyl alcohol	1544 g	
20	In a fluidized bed granulator metoprolol fumarate was sprayed onto the cores of silicon dioxide from a solution of ethanol 95%. 400 g of the beads (granules) so formed (fraction 0.4—0.63 mm) were covered with the polymeric layer containing ethyl cellulose 10 cps, hydroxypropylmethyl cellulose and acetyltributylcitrate by spraying a solution of the mentioned substances in methylene chloride and isopropyl alcohol. The coated beads were then filled into hard gelatine capsules.		
	20		20
	EXAMPLE 2		
	Metoprolol succinate	1440 g	
	Methylene chloride	9618 g	
25	Ethanol 95%	3888 g	25
	SiO ₂ (0.15—0.25 mm)	375 g	
	<u>Polymeric layer</u>		
	Ethylcellulose 50 cps	168.1 g	
	Hydroxypropylmethyl cellulose	36.9 g	
30	Acetyltributylcitrate	22.8 g	30
	Methylene chloride	4167 g	
	Isopropyl alcohol	815 g	
	<u>Tablet additives</u>		
	Microcrystalline cellulose	470.3 g	
35	Maize starch	117.6 g	35
	Potato starch	10.6 g	
	Water purified	342.2 g	
	Magnesium stearate	1.2 g	

Metoprolol succinate was sprayed onto the cores of silicon dioxide according to the process described in Example 1. 400 g of the granules so formed were coated with a polymeric film containing ethyl cellulose 50 cps, hydroxypropylmethyl cellulose and acetyltributylcitrate. An additional tablet mass was made by wet granulation of the dry mixture of microcrystalline cellulose and maize starch with the potato starch-water solution in a planetary mixer. Equal amounts (600 g) of the active and additional granules were finally mixed with Mg-stearate 0.1% and compressed to tablets.

EXAMPLE 3

Metoprolol succinate 100% in the form of compact spherical granules and having an average particle size of 0.42 mm.

400 g of the metoprolol succinate granules above with particles less than 0.63 mm were coated with

	Ethylcellulose 10 cps	177.1 g	
	Hydroxypropylmethyl cellulose	38.9 g	
	Acetyltributylcitrate	24.0 g	
	Methylene chloride	4094 g	
15	Isopropyl alcohol	1029 g	15

The beads obtained were formed into pharmaceutical preparations as described above.

EXAMPLE 4

	Metoprolol succinate	1440 g	
	Methylene chloride	9618 g	
20	Ethanol 95%	3888 g	20
	SiO ₂ (0.15—0.25 mm)	375 g	

Polymeric layer

	Ethylcellulose N—10	166.2 g	
	Hydroxypropylmethyl cellulose	39.0 g	
25	Acetyltributylcitrate	22.8 g	25
	Methylene chloride	3889 g	
	Isopropyl alcohol	978 g	

Tablet additives

	Microcrystalline cellulose	429.3 g	
30	Maize starch	67.1 g	30
	Lactose powder	40.3 g	
	Polyvidone	55.5 g	
	Water purified	314.7 g	
	Magnesium stearate	1.2 g	

Tablet coating (12.500 tablets)

	Hydroxypropylmethyl cellulose	159.6 g	
	Polyethylene glycol 6000	39.9 g	
	Colour Titanium Dioxide	39.9 g	
5	Water purified	1356 g	5
	Paraffin special	1.6 g	

Metoprolol succinate was sprayed onto the cores of silicon dioxide according to the process described in Examples 1 and 2 above. 400 g of the so obtained beads (fraction 0.4—0.63 mm) were coated with the polymeric mixture also described above. The coated beads of metoprolol succinate obtained were mixed with the additives in equal portions and after addition of Mg-stearate 0.1%, the dry mixture was compressed to tablets. Finally, the tablets were coated in a coating pan with the tablet coating described above.

EXAMPLE 5

S-enantiomeric metoprolol sorbate in the form of compact spherical granules in the fraction 0.4—0.63 mm.

40 g of the metoprolol sorbate granules above with particles less than 0.63 mm together with 360 g of non-pareil granules with particles between 0.75—1.0 mm were coated with

	Ethylcellulose 10 cps	57.7 g	
	Hydroxypropylmethyl cellulose	11.3 g	
20	Acetyltributylcitrate	7.0 g	20
	Methylene chloride	1194 g	
	Isopropyl alcohol	300 g	

The beads obtained were formed into pharmaceutical preparations as described above.

BIOPHARMACEUTICAL STUDIES

The plasma concentrations of metoprolol after a single dose of a controlled release preparation containing metoprolol succinate 190 mg according to example 4 of the description and the plasma concentrations after a single dose of Durules® containing 200 mg of metoprolol tartrate are shown in the attached Figure 2. 190 mg of the succinate salt is equivalent to 200 mg of metoprolol tartrate. The comparison has been carried out in 10 subjects. Each point represents the mean data from the 10 subjects. As can be seen the preparation according to the invention gives an almost constant concentration of metoprolol during more than 20 hours, whereas the insoluble matrix preparation gives unwanted high plasma concentration during the first hours after administration.

REDUCTION OF EXERCISE HEART RATE

12 subjects were given an ordinary tablet containing 100 mg of metoprolol tartrate once a day and the reduction of the exercise heart rate on day 5 of the treatment was measured and compared with the reduction of the exercise heart rate on day 5 in subjects given a controlled release preparation according to Example 4 of the invention containing 95 mg metoprolol succinate (equivalent to 100 mg metoprolol tartrate). The reduction of the heart rate is illustrated in Fig. 3. As can be seen the preparation according to the invention gives an even pharmacodynamic effect for 24 hours.

The best mode of carrying out the invention is at present considered to be Example 4.

Table 1 illustrates the in vitro release of metoprolol from the compositions according to Examples 1—4. As can be seen from the table at least 50% of the dose of metoprolol is released at a rate varying between 3—7 w/w %/hour.

TABLE 1
Cumulative in vitro dissolution of metoprolol in a phosphate buffer pH 6.8.
Method: USP apparatus No. II, rotating paddle at 100 rpm.

Example No.	Preparation	Per cent released over time (h)											
		1	2	3	4	6	8	10	12	14	16	18	20
1	capsule	1	2	5	11	25	39	52	62	69	74	78	81
2	tablet	7	11	16	19	29	40	50	59	68	76	82	86
3	capsule	3	7	12	17	27	37	44	52	60	67	74	80
4	tablet	7	13	18	23	33	43	52	61	69	76	82	86
5	capsule	4	—	—	21	—	47	—	67	—	80	—	88

CLAIMS

1. Controlled release preparation containing a number of beads comprising a salt of metoprolol as the main soluble component, the beads being coated with a polymeric membrane containing derivatives of cellulose without protolysable groups so that at least 75% of the dose of metoprolol is released within 20 hours virtually independent of the pH in the range 1—8. 5
2. Preparation according to claim 1 wherein the diameter of the beads is 0.25—2.0 mm.
3. Preparation according to claim 2 wherein the diameter of the beads is 0.35—1.0 mm.
4. Preparation according to any one of the preceding claims wherein at least 50% of the dose of metoprolol is released at the rate 3—7 w/w %/hour. 10
5. Preparation according to any one of the preceding claims wherein the salt of metoprolol has a solubility less than 600 mg/ml in water at 25°C, preferably a solubility of 30—600 mg/ml.
6. Preparation according to any one of the preceding claims wherein the salt is a salt of an organic carboxylic acid.
7. Preparation according to claim 6 wherein the salt of metoprolol is the succinate or fumarate of racemic metoprolol. 15
8. Preparation according to claim 6 wherein the salt of metoprolol is the benzoate or sorbate of the S-enantiomer of metoprolol.
9. Preparation according to any one of the preceding claims wherein the uncoated beads comprise at least 95 w/w% of metoprolol. 20
10. Preparation according to any one of the preceding claims wherein the uncoated beads comprise a salt of metoprolol applied on an insoluble core.
11. Preparation according to claim 10 wherein the insoluble core has a diameter of 0.1—1.0 mm, preferably 0.15—0.3 mm.
12. Preparation according to claim 10 wherein the insoluble core has a diameter of 0.15—0.3 mm and is covered by a salt of metoprolol to give a bead having a diameter of 0.35—1.0 mm, such beads being coated with the polymeric membrane. 25
13. Preparation according to any one of claims 1 to 9 wherein the bead has a diameter of 0.35—1.0 mm, the bead comprises a salt of metoprolol and is coated with the polymeric membrane.
14. Preparation according to any one of the preceding claims wherein the polymeric membrane contains ethyl cellulose. 30
15. Preparation according to any one of the preceding claims wherein the polymeric membrane contains ethyl cellulose together with hydroxypropylmethyl cellulose.
16. Preparation according to claim 1 substantially as hereinbefore described with reference to any one of the Examples. 35
17. Method of producing a preparation as defined in any one of claims 1 to 16 wherein beads comprising the salt of metoprolol as the main soluble component are spray coated with a membrane-forming solution containing derivatives of cellulose.
18. A method according to claim 17 substantially as hereinbefore described with reference to any one of the Examples. 40
19. A preparation obtained by a process according to claim 17 or 18.
20. A pharmaceutical preparation comprising the controlled release preparation according to any one of claims 1 to 16 or 19 filled into hard gelatine capsules.
21. A pharmaceutical preparation comprising the controlled release preparation according to any one of claims 1 to 16 or 19 wherein the coated beads together with pharmaceutical additives are compressed to 45

tablets which disintegrate into the primary formed coated beads when the tablets are brought into contact with gastro-intestinal fluids.

22. A pharmaceutical preparation according to claim 20 or 21 substantially as hereinbefore described with reference to any one of the Examples.

5 23. A preparation according to any one of Claims 1 to 16 or 19 or 20 to 22 for use in a method of treatment of the human or animal body by surgery or therapy or of diagnosis practised on the human or animal body. 5

Printed for Her Majesty's Stationery Office by Courier Press, Leamington Spa, 4/1987. Demand No. 8991685.
Published by the Patent Office, 25 Southampton Buildings, London, WC2A 1AY, from which copies may be obtained.