

(43) International Publication Date
31 July 2014 (31.07.2014)(10) International Publication Number
WO 2014/116880 A1

(51) International Patent Classification:

C07K 19/00 (2006.01) A61P 37/00 (2006.01)
A61K 39/395 (2006.01) A61P 27/02 (2006.01)

(21) International Application Number:

PCT/US2014/012831

(22) International Filing Date:

23 January 2014 (23.01.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/755,968	23 January 2013 (23.01.2013)	US
61/755,960	23 January 2013 (23.01.2013)	US
61/771,565	1 March 2013 (01.03.2013)	US
61/771,560	1 March 2013 (01.03.2013)	US

(72) Inventors; and

(71) Applicants : **ROHRER, Baerbel** [US/US]; 1000 Cummings Circle, Mount Pleasant, SC 29464 (US). **HOLERS, Michael** [US/US]; 242 S. Hudson Street, Denver, CO 80246 (US). **TOMLINSON, Stephen** [US/US]; 1287 Abercorn Trace, Mount Pleasant, SC 29466 (US). **THURMAN, Joshua, M.** [US/US]; 2 Red Fox Lane, Greenwood Village, CO 80111 (US). **ATKINSON, Carl** [GB/US]; 3249 Beaconsfield Road, Park West, Mount Pleasant, SC29466 (US). **KULIK, Liudmila** [BY/US]; 14934 E. Hampden Circle, D5, Denver, CO 80014-5227 (US).(74) Agents: **XIAO, Jian** et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: TARGETING CONSTRUCTS BASED ON NATURAL ANTIBODIES AND USES THEREOF

Figure 3

A



(57) Abstract: The present invention provides targeted delivery methods and constructs for treating inflammatory diseases and/or detecting in vivo tissue injuries in an individual. The targeted delivery approach utilizes an antibody that recognizes an epitope found to be present at sites of inflammation. The invention also provides methods of inhibiting complement-driven inflammation in the eye in an individual, comprising administering to the individual an antibody or a fragment thereof or compositions thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid. Also provided are related methods of treating a complement-associated ocular disease or an ocular disease involving oxidative damage. Additionally, the invention provides methods of detecting complement-mediated injury in an eye tissue of an individual, comprising administering to the individual a construct or compositions thereof, wherein the construct comprises (a) an antibody or fragment thereof that specifically binds to Annexin IV or phospholipid; and (b) a detectable moiety.

Published:

— *with international search report (Art. 21(3))*

TARGETING CONSTRUCTS BASED ON NATURAL ANTIBODIES AND USES THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This patent application claims priority benefit of United States Provisional Patent Application No. 61/755,960 filed January 23, 2013, United States Provisional Patent Application No. 61/755,968, filed January 23, 2013, United States Provisional Patent Application No. 61/771,560, filed March 1, 2013, and of United States Provisional Patent Application No. 61/771,565 filed March 1, 2013, the entire content of each of which is incorporated herein by reference.

TECHNICAL FIELD

[0002] This application pertains to targeting constructs based on natural antibodies and uses thereof. This application also pertains to compositions and methods of treating ocular disease, specifically ocular diseases associated with oxidative stress.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0003] This invention was made with Government support under Grant (Contract) Nos.: U.S. Army Medical Research and Materiel Command (MRMC) Awards W81XWH-06-1-0520 and W81XWH-07-1-0286, and National Institutes of Health R01 AI31105, C06 RR015455, and R01EY019320 and Department of Veterans Affairs I01 RX000444; Beckman Institute for Macular Research.

BACKGROUND

[0004] Natural antibodies exist in an immune competent individual and can be found in the serum or plasma of an individual not known to have been stimulated by a specific antigen to which the antibody binds. Previous studies by the present inventors and colleagues have shown that certain types of natural antibodies recognize epitopes on ischemic tissue and catalyze the initiation and subsequent development of ischemia-reperfusion injury (Fleming et al., 2002, J. Immunol. 169:2126-2133; Rehrig et al., 2001, J. Immunol. 167:5921-5927). Ischemia-reperfusion injury, as well as hypovolemic shock and subsequent tissue damage, is known to be caused by complement and Fc receptor activation and the recruitment and activation of

neutrophils and other inflammatory cells (Rehrig et al., 2001, *supra*). It had also been shown that single monoclonal antibodies that react broadly with phospholipids and other extracellular or intracellular antigens such as DNA can cause ischemia-reperfusion injury in mice that lack other antibodies (i.e., B cell-deficient mice).

[0005] Ischemia-reperfusion (IR) injury refers to damage to a tissue caused when the blood supply returns to the tissue after a period of ischemia (restriction in blood supply). The absence of oxygen and nutrients from the blood creates a condition in which the restoration of circulation results in inflammation and oxidative damage, rather than restoration of normal function. Ischemia-reperfusion injury can be associated with traumatic injury, including hemorrhagic shock, as well as many other medical conditions such as stroke or large vessel occlusion, and is a major medical problem. More particularly, ischemia-reperfusion injury is important in heart attacks, stroke, kidney failure following vascular surgery, post-transplantation injury and chronic rejection, as well as in various types of traumatic injury, where hemorrhage will lead to organ hypoperfusion, and then subsequent reperfusion injury during fluid resuscitation. Ischemia-reperfusion injury, or an injury due to reperfusion and ischemic events, is also observed in a variety of autoimmune and inflammatory diseases. Independently of other factors, ischemia-reperfusion injury leads to increased mortality.

[0006] There is also increasing evidence of reperfusion injury that can be found in autoimmune and inflammatory diseases that are not traditionally thought of as reperfusion injury-related. For example, the synovium in rheumatoid arthritis patients is a site that is subjected to constant reperfusion stress (e.g., low pH, lots of tissue pressure and poor perfusion). The higher quantity of synovial fluid found in hypermobile patients having this disease causes an increase in the intra-articular pressure, which is then exacerbated by joint motion. This may aggravate local inflammation through a hypoxic/reperfusion mechanism, which in turn causes oxidative injury due to intermittent ischemia (e.g., see Punzi et al., *Rheumatology* 2001; 40: 202-204; Pianon et al., *Reumatismo* 1996; 48(Suppl. 1):93; and Jawed et al., *Ann Rheum Dis* 1997; 56:686-9). A variety of inflammatory and autoimmune diseases can also be associated with similar changes in cell stress responses of local cells that are similar to, or mimic, some changes in reperfusion injury.

[0007] Kulik et al. showed that pathogenic natural antibodies recognizing Annexin IV are required to develop intestinal ischemia-reperfusion injury. *J. Immunol.* 2009;

182:5363-5373. U.S. Patent Application Publication No. 2011/0014270 discloses lipids, annexins, and lipid-annexin complexes for use in the prevention and/or treatment of ischemia-reperfusion injury and reperfusion injury associated with a variety of diseases and conditions.

[0008] Natural antibodies are also involved in the pathology of ocular diseases. Age-related macular degeneration (AMD), which is characterized by progressive loss of central vision resulting from damage to the photoreceptor cells in the central area of the retina, the macula, is the leading cause of vision loss in the elderly of industrialized nations (Council, N. (1999) Vision research - a national plan: 1999-2003, executive summary. (National Eye Institute, N. i. o. h. ed., Washington, DC). Although AMD occurs in two forms, neovascular (wet) and atrophic (dry), both are associated with pathological lesions at the retinal pigmented epithelium (RPE)/choroid interface in the macular region (Nowak, J. Z. (2006) *Pharmacol Rep* **58**, 353-363). Early AMD is characterized by a thickening of Bruch's membrane, which includes basal linear deposits and drusen (Hageman, G. S., Luthert, P. J., Victor Chong, N. H., Johnson, L. V., Anderson, D. H., and Mullins, R. F. (2001) *Prog Retin Eye Res* **20**, 705-732). Additionally, changes in RPE morphology, pigmentation and deterioration of its function as a blood-retina barrier have been reported (McLeod, D. S., Taomoto, M., Otsuji, T., Green, W. R., Sunness, J. S., and Lutty, G. A. (2002) *Invest Ophthalmol Vis Sci* **43**, 1986-1993). Advanced AMD is characterized by additional subtype-specific morphological features exacerbating the early pathological damage (Bhutto, I., and Lutty, G. (2012) *Mol Aspects Med* **33**, 295-317). Dry AMD, or geographic atrophy, results from the loss of RPE followed by the loss of photoreceptors; whereas wet AMD is associated with choroidal neovascularization and leakage of these new vessels.

[0009] AMD is a complex disease with both genetic and environmental risk factors. The main environmental risk factor is persistent oxidative stress (Snodderly, D. M. (1995) *Am J Clin Nutr* **62**, 1448S-1461S), whether that might be caused by smoking, nutritional deficits or even light exposure. A main genetic risk factor for the disease is polymorphisms in genes for complement proteins, including complement factor H (CFH), complement factor B (CFB), complement component 2 (C2) and complement component 3 (C3) (reviewed in (Charbel Issa, P., Chong, N. V., and Scholl, H. P. (2011) *Graefes Arch Clin Exp Ophthalmol* **249**, 163-174)). Discovering complement genes as risk factors was consistent with prior clinical studies, which demonstrated

that the complement system activation products were found locally in the eye in all stages of AMD (Hageman, G. S., Anderson, D. H., Johnson, L. V., Hancox, L. S., Taiber, A. J., Hardisty, L. I., Hageman, J. L., Stockman, H. A., Borchardt, J. D., Gehrs, K. M., Smith, R. J., Silvestri, G., Russell, S. R., Klaver, C. C., Barbazetto, I., Chang, S., Yannuzzi, L. A., Barile, G. R., Merriam, J. C., Smith, R. T., Olsh, A. K., Bergeron, J., Zernant, J., Merriam, J. E., Gold, B., Dean, M., and Allikmets, R. (2005) *Proc Natl Acad Sci U S A* **102**, 7227-7232). Follow-up experiments in animal models, in particular of wet AMD, further support the hypothesis that inadequate control of complement-driven inflammation may be a major factor in the disease pathogenesis of AMD (*e.g.*, (Nozaki, M., Raisler, B. J., Sakurai, E., Sarma, J. V., Barnum, S. R., Lambris, J. D., Chen, Y., Zhang, K., Ambati, B. K., Baffi, J. Z., and Ambati, J. (2006) *Proc Natl Acad Sci USA* **103**, 2328-2333; Bora, P. S., Sohn, J. H., Cruz, J. M., Jha, P., Nishihori, H., Wang, Y., Kaliappan, S., Kaplan, H. J., and Bora, N. S. (2005) *J Immunol* **174**, 491-497; Rohrer, B., Coughlin, B., Kunchithapautham, K., Long, Q., Tomlinson, S., Takahashi, K., and Holers, V. M. (2011) *Mol Immunol* **48**, e1-8)). Although the current understanding of AMD is that chronic oxidative damage over time leads to alterations in photoreceptors, RPE/Bruch's membrane and the choriocapillaris complex, in particular in the macula, resulting in chronic inflammation and complement activation (Zarbin, M. A., and Rosenfeld, P. J. (2010) *Retina* **30**, 1350-1367), it is unclear which components of the complement cascade are involved in causing damage, and what ligands or age-related changes in these tissues enable complement activation. The complement cascade, an evolutionarily ancient and highly conserved system, is part of the innate and adaptive immune system, consisting of >40 soluble and membrane-bound components (Muller-Eberhard, H. J. (1988) *Annu Rev Biochem* **57**, 321-347). Its normal role is to complement the ability of antibodies and phagocytic cells to eliminate pathogens. To spot these microorganisms, pattern recognition molecules complexed to inactive serum proteases circulate in the blood. Upon ligand interaction, the protease becomes activated to initiate the complement cascade. This results in the production of anaphylatoxins to recruit phagocytic cells, opsonins to tag material for removal, and the generation of the membrane attack complex (MAC) to rupture membranes of cells and leading to pro-inflammatory signaling in the target cell. Self cells are protected by either membrane-bound or soluble complement inhibitors. However, under

pathological conditions, complement inhibition might be compromised, resulting in complement activation on self surfaces.

[0010] The complement system can be activated by one of three pathways, the classical, lectin and alternative pathway, each with its unique pattern recognition molecules. The classical pathway (CP) is activated when C1q binds to its ligands that include C-reactive protein (CRP); serum amyloid protein or IgG and IgM molecules present as immune complexes; the lectin pathway (LP) when mannan-binding lectin (MBL) or ficolin (H-ficolin, L-ficolin or M-ficolin) binds to specific carbohydrates or acetylated molecules on foreign cells or IgM molecules bound to antigens; and finally, the alternative pathway (AP) is spontaneously and continuously activated at a low level in a process called tickover, as well as when C3b is generated on cell surfaces by the CP or LP and becomes a substrate for the AP. All three pathways lead to the generation of a pathway-specific C3 convertase that then triggers the common terminal pathway with its above-described biological effects.

[0011] The disclosures of all publications, patents, patent applications and published patent applications referred to herein are hereby incorporated herein by reference in their entirety.

BRIEF SUMMARY OF THE INVENTION

[0012] In one aspect, the present disclosure provides a method of inhibiting complement-mediated inflammation in a tissue having non-ischemic injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor.

[0013] In another aspect, the present disclosure provides a method of treating an inflammatory disease in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor.

[0014] In some embodiments of the methods above, the complement inhibitor is selected from the group consisting of an anti-C5 antibody, anti-MASP antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7

antibody, an anti-factor B antibody, an anti-factor D antibody, and an anti-properdin antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding. In some embodiments, the complement inhibitor is a specific inhibitor of the alternative pathway. In some embodiments, the complement inhibitor is a specific inhibitor of the lectin pathway.

[0015] In another aspect, the present disclosure provides a method of detecting complement-mediated injury in a tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated tissue injury. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the detectable moiety is selected from the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.

[0016] In some embodiments of the methods above, the tissue injury results from any of from inflammatory disorders, transplant rejection, pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders. In some embodiments, the tissue is any one of eye, joint, kidney, brain, heart, spinal cord, and liver. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the disease is any one of an ocular disease, arthritis, or renal injury.

[0017] In some embodiments of the methods above, the antibody or fragment thereof specifically binds to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as monoclonal antibody B4 (such as B4/14/12, ATCC Deposit No. PTA-13522). In some embodiments, the Annexin IV is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing non-ischemic injury.

[0018] In some embodiments of the methods, the antibody or fragment thereof specifically binds to a phospholipid. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid.

In some embodiments, the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody C2 (such as C2/19/8, ATCC Deposit No. PTA-13523).

In some embodiments, the phospholipid is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is MDA.

[0019] In some embodiments of the methods above, construct is a fusion protein.

In some embodiments, the antibody or fragment thereof and the complement inhibitor or detectable moiety are linked via a peptide linker. In some embodiments, the antibody or fragment thereof is a scFv. In some embodiments, the antibody or fragment thereof is Fab, Fab', or F(ab')₂.

[0020] In another aspect, the present disclosure provides a construct comprising: (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1 or 7, a sequence of SEQ ID NO:2 or 8, or a sequence of SEQ ID NO:3 or 9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4 or 10, a sequence of SEQ ID NO:5 or 11, or a sequence of SEQ ID NO:6 or 12. In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1 or 7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2 or 8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3 or 9. In some embodiments, the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4 or 10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5 or 11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6 or 12. In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:13 or 14. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:15 or 16. In some embodiments, the antibody or fragment is an scFv having the sequence of SEQ ID NO:17 or 18. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as

monoclonal antibody B4. In some embodiments, the Annexin IV is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing or is at risk of undergoing tissue injury.

[0021] In another aspect, the present disclosure provides a construct comprising: (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25 or 31, a sequence of SEQ ID NO:26 or 32, or a sequence of SEQ ID NO:27 or 33; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25 or 31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26 or 32; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27 or 33. In some embodiments, the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:34 or 35. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:36. In some embodiments, the antibody or fragment is an scFv having the sequence of SEQ ID NO:37 or 38. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid. In some embodiments, the antibody or fragment thereof binds to the same epitope as monoclonal antibody C2.

[0022] In some embodiments, the phospholipid is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing or is at risk of undergoing tissue injury. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is MDA.

[0023] In some embodiments, the construct comprises a complement modulator. In some embodiments, the complement modulator is a complement inhibitor. In some embodiments, the complement inhibitor is selected from the group consisting of an anti-C5 antibody, anti-MASP antibody, an Eculizumab, an pexelizumab, an anti-C3b

antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-factor D antibody, and an anti-properdin antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding. In some embodiments, the complement inhibitor is a specific inhibitor of the alternative pathway. In some embodiments, the complement inhibitor is a specific inhibitor of the lectin pathway.

[0024] In some embodiments, the construct comprises a detectable moiety. In some embodiments, the detectable moiety is selected from the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.

[0025] In some embodiments, the construct is a fusion protein. In some embodiments, the antibody or fragment thereof and the complement modulator or detectable moiety are linked by a peptide linker.

[0026] In another aspect, the present disclosure provides a pharmaceutical composition comprising a construct described above. Also provided is a method of inhibiting complement-mediated inflammation in an individual, comprising administering to the individual an effective amount of a pharmaceutical composition described herein. Additionally, the present disclosure provides a method of treating an inflammatory disease in an individual, comprising administering to the individual an effective amount of a pharmaceutical composition herein. Also provided herein is a composition comprising a construct comprising a detectable moiety, *e.g.*, a radioisotope, a fluorescent dye, an electron-dense reagent, an enzyme, a biotin, a paramagnetic agent, a magnetic agent, and a nanoparticle. Also provided is a method of detecting complement-mediated injury in a tissue of an individual, comprising administering to the individual an effective amount of a composition of a construct comprising a detectable moiety.

[0027] In one aspect, the present disclosure provides a method of inhibiting complement-driven inflammation in the eye in an individual, comprising administering to the individual an effective amount of (a) an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid; or (b) a composition comprising a construct, wherein the

construct comprises: (i) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent.

[0028] In another aspect, the present disclosure provides a method of treating a complement-associated ocular disease or an ocular disease involving oxidative damage, comprising administering to the individual an effective amount of (a) an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid; or (b) a composition comprising a construct, wherein the construct comprises: (i) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent.

[0029] In some embodiments, the methods comprise administration of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid.

[0030] In some embodiments, the methods comprise administration of a composition comprising a construct, wherein the construct comprises: (i) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent. In some embodiments, the therapeutic agent is a complement inhibitor. In some embodiments, the complement inhibitor is selected from, but not limited to, the group consisting of an anti-C5 antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-MASP antibody, an anti-factor D antibody, and an anti-properdin antibody, an anti-MBL antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding. In some embodiments, the complement inhibitor is a human complement inhibitor (e.g., a human MCP, a human DAF, a human CD59, a human CR1, a human Factor H, or another complement inhibitor derived from humans). In some embodiments, the complement inhibitor is a human complement inhibitor (e.g., a mouse DAF, a mouse CD59 (also known as isoform A), a mouse CD59 isoform B, a mouse Crry, a mouse Factor H, or another complement inhibitor derived from

mouse). Complement inhibitors from other species and variant complement inhibitors are also contemplated.

[0031] In some embodiments of the methods, the ocular disease is selected from, but not limited to, the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0032] In another aspect, the present disclosure provides a method of detecting complement-mediated injury in an eye tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated eye tissue injury. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the detectable moiety is selected from, but not limited to, the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.

[0033] In some embodiments of the methods above, the antibody or fragment thereof specifically binds to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody B4 (such as B4/14/12, ATCC Deposit No. PAT-13522). In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane, or in a pathological structure in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage.

[0034] In some embodiments of the methods above, the antibody or fragment thereof specifically binds to a phospholipid. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid. In some embodiments, the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody C2 (such as C2/19/8, ATCC Deposit No. PAT-13523). In some embodiments, the phospholipid is present on the surface of a

cell, a basement membrane, or in a pathological structure in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments the phospholipid is MDA.

[0035] In some embodiments of any of the methods above, the antibody or fragment thereof is a scFv. In some embodiments, the antibody or fragment thereof is Fab, Fab', or F(ab')₂. In some embodiments, the construct is a fusion protein. In some embodiments, the antibody or a fragment thereof and the therapeutic agent or detectable moiety are joined by a linker. In some embodiments, the antibody or a fragment thereof and the therapeutic agent or detectable moiety are joined directly.

[0036] In some embodiments of any of the methods above, the administration is by injection into the eye. In some embodiments, the individual is human.

[0037] Also provided are unit dosage forms, kits, and articles of manufacture that are useful for methods described herein.

[0038] It is to be understood that one, some, or all of the properties of the various embodiments described herein may be combined to form other embodiments of the present invention.

BRIEF DESCRIPTION OF THE FIGURES

[0039] **Figure 1** is a series of graphs showing characterization of B4scFv and B4scFv-Crry proteins. **A)** B4scFv, but not control C2scFv, bound directly to recombinant annexin IV *in vitro*. **B)** B4scFv competitively inhibited binding of B4 mAb to annexin IV. **C)** B4scFv-Crry inhibited complement activation *in vitro* similarly to positive control CR2-Crry.

[0040] **Figure 2** is a series of graphs showing that administration of B4 mAb and C2 mAb overcame protection from spinal cord injury (SCI) due to ischemic injury in Rag1^{-/-} mice and cerebral ischemia reperfusion injury. **A)** Locomotor activity after experimentally induced SCI in Rag1^{-/-} mice administered different mAbs. Rag1^{-/-} mice (filled squares) were protected from SCI 21 days post-injury while wild-type mice (filled circles) showed a reduction of locomotor activity 21 days post-injury. Injury in Rag1^{-/-} mice was reconstituted to levels comparable to wild-type mice when administered B4 mAb (empty circles) or C2 mAb (filled triangles). In comparison, Rag1^{-/-} mice administered control F632 mAb (empty diamond) remained protected from SCI 21 days post-injury. P<0.05, n=6-9. **B)** Post-ischemic infarct volume (TTC

staining) in a mouse model of ischemic stroke (cerebral ischemia-reperfusion injury) after administering a targeting construct. Rag1^{-/-} mice were protected from cerebral infarct compared to C57Bl/6 wild-type mouse controls (#p<0.001). Reconstitution with increasing amounts of C2 or B4 mAbs restored injury to Rag1^{-/-} mice (@p=0.01). Reconstitution with 100 µg control antibody did not restore injury in Rag1^{-/-} mice. n=8-12.

[0041] Figure 3 is a series of graphs showing that B4scFv-Crry targeting construct protected mice from SCI. **A)** Locomotor activity after experimentally induced SCI in wild-type mice administered a targeting construct. Targeted complement inhibitor B4scFv-Crry (0.2 mg), shown as B4-Crry, protected wild-type mice against SCI as compared to wild-type mice administered PBS. All mice had a score of 0 immediately after impact. p<0.05 from day 3. n=6. **B)** Morphometric analysis of tissue sparing 3 days after traumatic injury. Cross sectional area of 120 µm increments (H&E). B4-Crry indicates wild-type treated mice, WT indicates wild-type mice treated with PBS, all others were Rag1^{-/-} mice reconstituted with the indicated IgM mAbs, C2 mAb, B4 mAb, and F632 mAb or administered PBS (shown as Rag1^{-/-}). p < 0.01 up to 1.2 mm each side of injury site. Mean +/- SD, n = 6 per group.

[0042] Figure 4 is immunofluorescence confocal analysis of IgM and C3 deposition. **A-C)** Spinal cord sections from untreated wild-type mice stained for **A)** IgM or **B)** C3 at 24 hours after injury. **C)** Merged image of IgM and C3 staining. **D and E)** Sections from wild-type mice treated with B4scFv-Crry and stained for **D)** IgM or **E)** C3 at 24 hours after injury.

[0043] Figure 5 is a graph showing animal survival in a cecal ligation and puncture model of acute septic peritonitis after administration of B4scFv-Crry. The absence of C3 in C3 deficient mice (C3^{-/-}) resulted in death within 48 hours. There was no significant difference in survival seen between B4scFv-Crry (treated with 0.2mg dose immediately post procedure) and wild-type untreated controls. n=5-6.

[0044] Figure 6 shows localization of B4 mAb in transplanted hearts after administration of B4 mAb to Rag1^{-/-} recipients. **A)** B4 mAb bound to endothelial cells of the transplanted heart but not the native heart. **B)** Co-localization of C3d (left panel) and endogenous IgM (middle panel) in a graft of wild-type mouse. Overlay image (right panel) shows co-localization of C3d and endogenous IgM.

[0045] Figure 7 shows **A-D)** a protective effect of B4scFv and B4scFv-Crry against cardiac ischemia reperfusion injury in transplanted heart. **A)** Graph showing

decreased serum levels of cardiac troponin I with a single dose of B4scFv or B4scFv-Crry administered to recipient after transplant. ns= not significant. **B)** Graph summarizing histological scoring of cardiac damage showed a significant reduction in injury histology score with a single dose of B4scFv or B4scFv-Crry. ns= not significant. **C)** Immunofluorescence imaging of B4scFv-HisTag *in vivo* showing binding to endothelial cells. **D)** Immunofluorescent staining for C3d in allograft heart transplants after treatment of recipient with either 0.2mg B4scFv-Crry (lower panels) or PBS control (upper panels). Only weak C3d vascular staining was seen in B4scFv-Crry treated animals 48 hours post treatment. Images representative of n=3. **E and F)** Shows the effect of B4scFv and B4scFvCrry treatment on IgM and C3d deposition in transplanted allografts. Recipient mice were treated with PBS, B4scFv or B4scFvCrry and allografts were isolated at 6 hours or 48 hours post-transplantation for analysis. Representative images of IgM and C3d deposition in grafts are shown. Semi-quantitative histological scoring of **E)** IgM and **F)** C3d deposition at 6 and 48 hours post-transplantation. * $P < 0.01$, ** $P < 0.001$, Mean $\pm$ SEM, n=6–8.

[0046] **Figure 8** is a series of graphs showing reduction of specific cytokines in allograft recipients after treatment with B4scFv or B4scFv-Crry. Cytokine levels for **A)** MCP-1, **B)** IL-6, **C)** KC, **D)** CXCL9, and **E)** IFN- γ . n = 4-10, # $p < 0.01$.

[0047] **Figure 9** is graph showing biodistribution of ^{125} I-labeled B4scFv-Crry in recipient mice administered immediately after heart transplantation and analyzed 6 hours later.

[0048] **Figure 10** shows anti-IgM immunofluorescence images of B4 mAb binding to **A-B)** a mouse brain endothelial cell line (bEnd.3) and **C-D)** human umbilical vein endothelial cells (HUVEC), both subjected to 3 hours hypoxia (left panels) and 1 hour re-oxygenation (right panels). B4 mAb did not bind to either cell type not exposed to hypoxic conditions.

[0049] **Figure 11** is a series of graphs showing that administration of B4 mAb and C2 mAb overcame protection from hepatic ischemic reperfusion injury in Rag1^{-/-} mice. **A)** Serum ALT levels 6 hours post I/R in sham (Rag1^{-/-}), wild-type, Rag1^{-/-} mice, and Rag1^{-/-} mice injected with 25 μ g C2 mAb or B4 mAb. Mean $\pm$ SD, n=4-9. **B)** Necrotic index 6 hours post IR in sham (Rag1^{-/-}), wild-type, Rag1^{-/-} mice and Rag1^{-/-} mice injected with 25 μ g C2 mAb or B4 mAb. Mean $\pm$ SD, n = 7. ##, $P < 0.001$ vs. wild-type; **, $P < 0.001$ vs. Rag1^{-/-}; @, $P < 0.01$ vs. Rag1^{-/-}; *, $P < 0.05$ vs. Rag1^{-/-}.

[0050] **Figure 12** is a series of graphs showing that administration of B4 mAb and C2 mAb stimulated liver regeneration following 70% partial hepatectomy in Rag1^{-/-} mice. **A)** Serum ALT levels 48 hours after PHx in Rag1^{-/-} mice and Rag1^{-/-} mice treated with 10 µg C2 mAb or B4 mAb. **, @@ P<0.01 vs. rag IR injury. Mean ± SD, n= 6. **B)** Necrotic index score (H&E staining) 48 hours post PHx in Rag1^{-/-} mice and Rag1^{-/-} mice treated with 10 µg C2 mAb or B4 mAb. ##, P<0.01 vs. wild-type; * and @, P<0.05 vs. Rag1^{-/-}. Mean ± SD, n=4. **C)** Mitotic index 48 hours post PHx in Rag1^{-/-} mice and Rag1^{-/-} mice treated with 10 µg C2 mAb or B4 mAb. ##, P<0.01 vs. wild-type; **, P<0.01 vs. Rag1^{-/-}; @@, P<0.01 vs. Rag1^{-/-}. Mean ± SD, n=4.

[0051] **Figure 13** is a series of graphs showing that administration of B4 mAb and C2 mAb stimulated liver regeneration following 70% partial hepatectomy in Rag1^{-/-} mice. **A)** Liver weight restitution 48 hours after PHx in Rag1^{-/-} mice and Rag1^{-/-} mice treated with 10 µg C2 mAb or B4 mAb. **B)** Brdu-positive cells 48 hours after PHx in Rag1^{-/-} mice and Rag1^{-/-} mice treated with 10 µg C2 mAb or B4 mAb. ***, P<0.001; **, P<0.01; *, P<0.05.

[0052] **Figure 14** is **A)** a series of immunohistochemistry pictures of IgM staining in liver sections 6 hours post IR in wild-type mice (wt), Rag1^{-/-} mice, and Rag1^{-/-} mice treated with C2 mAb or B4 mAb; **B)** a series of immunofluorescence images showing localization of endogenous IgM (left panel) and C3d (middle panel) in Rag1^{-/-} mice reconstituted with B4 mAb following hepatic IR. Overlay image (right panel) shows co-localization of C3d and endogenous IgM; **C)** immunohistochemical staining of IgM and C3d showed IgM and C3d staining in a sinusoidal pattern in WT and Rag1^{-/-} mice reconstituted with B4 mAb or C2 mAb 48 hours after 70% PHx. No staining was seen in Rag1^{-/-} treated with PBS. Representative images from 3 animals per group, magnification x400; and **D)** immunofluorescence images showing localization of IgM (left panel) and C3d (middle panel) in Rag1^{-/-} mice reconstituted with B4 mAb following 70% PHx. Overlay image (right panel) shows co-localization of C3d and endogenous IgM. Representative image from 3 animals, magnification x520.

[0053] **Figure 15** is a series of graphs showing *in vivo* kinetics of B4scFv-Crry, biodistribution of IgM antibodies following IR or 70% Phx in Rag1^{-/-} mice, and *in vivo* binding of B4 scFv construct. **A)** B4scFc-Crry had an initial rapid phase of elimination from the circulation with a half-life ($t_{1/2}$) of 27 minutes; **B)** a second prolonged phase with a half-life of 6.5 hours; **C and D)** Rag1^{-/-} animals were reconstituted with I¹²⁵ radiolabeled B4, C2, or isotype control (F632) IgMs following

IR, PHx or sham surgeries. Tissues were harvested and radioactivity was measured at 6 h post surgery. **C)** Biodistribution of IgMs following hepatic IR showed increased levels of B4 and C2 in the liver compared to sham controls. Isotype control antibody F632 did not accumulate in any tissue. **D)** Biodistribution of IgMs in Rag1-/- following 70% PHx. B4 and C2 radiolabeled mAbs accumulated in the liver specifically. There was no tissue accumulation of antibody in sham or F632 isotype control treated animals following 70% PHx. Data was representative of 2 independent experiments, n=2 for each group; and **E and F)** Biodistribution of I¹²⁵ radiolabeled B4 scFv in Rag1-/- following IR, PHx, or sham operation. Rag1-/- mice were given radiolabeled B4 scFv intraperitoneally immediately following surgeries, tissues were harvested and measured for radioactivity at 6 h post surgery. **E)** Biodistribution of B4 scFv in Rag1-/- mice following IR or sham operation showed accumulation of B4 scFv mainly in the liver of mice that underwent IR with no appreciable B4 scFv accumulation in sham animals. n=3 for each group, mean $\pm$ SD. **F)** Biodistribution of radiolabeled B4 scFv in Rag1-/- following 70% PHx or sham operation, showed accumulation of B4 scFv in the liver of mice given 70% PHx with no accumulation in sham operated mice n=2 for each group. Biodistribution studies were representative of 2 separate experiments.

[0054] **Figure 16** shows the protective effect of B4scFv and B4scFv-Crry against hepatic ischemic reperfusion injury. **A)** Serum ALT levels 24 hours post IR in sham, wild-type and wild-type mice injected with 25 μ g B4scFv or B4scFv-Crry. n=3. **B-E)** H&E staining in liver 24 hours post reperfusion in **B)** sham, **C)** wild-type mouse, **D)** B4scFv treated mouse, and **E)** B4scFv-Crry treated mouse. Images taken at 40 x zoom.

[0055] **Figure 17** is imaging analysis of B4 IgM deposition in human liver. **A)** Human liver section stained for IgM post-reperfusion of transplanted liver. **B)** Immunofluorescence confocal images of CD31 (endothelial marker, left panel) or B4 IgM (middle panel) stained ischemic non-reperfused human liver section. Right panel shows merged image of CD31 and B4 IgM staining.

[0056] **Figure 18** is a series of graphs showing human serum antibody levels after liver transplantation. **A)** total IgM antibodies, **B)** anti-albumin 2 antibodies, and **C)** total IgG antibodies.

[0057] **Figure 19** is a series of graphs showing human serum antibody levels after liver transplantation. **A)** anti-PE antibodies, **B)** anti-Annexin IV antibodies, **C)** anti-PC-BSA antibodies, and **D)** anti-cardiolipin antibodies.

[0058] **Figure 20** shows B cell depletion with an anti-CD20 antibody reduced glomerular IgM deposition in mice with adriamycin nephropathy. Mice were injected with anti-CD20 monoclonal antibody to deplete their B cells prior to the induction of adriamycin nephropathy. **A)** Four weeks after induction of adriamycin nephropathy immunofluorescence microscopy was performed on kidneys to assess the abundance of glomerular IgM. Kidneys from three to five mice per group were examined. Thirty glomeruli per section were visualized, and the average for each mouse was determined. Mice injected with adriamycin demonstrated a substantial increase in glomerular IgM deposition compared to control animals. Mice treated with anti-CD20 demonstrated less glomerular IgM. Mice with adriamycin nephropathy that had been injected with the anti-CD20 but were also reconstituted with purified IgM eluted from the kidneys of mice with adriamycin nephropathy (Adria/anti-CD20/IgM) demonstrated an apparent increase in glomerular IgM. Glomeruli are indicated with arrowheads. Original magnification X200. Scale bar = 100 μ M. After converting the images to grayscale, the brightness and contrast of the shown images were adjusted. All images were adjusted equally. **B)** Quantitative analysis of the glomerular IgM in the different treatment groups confirmed that glomerular IgM deposition was increased after injection with adriamycin, but that this increase was attenuated in mice injected with anti-CD20 therapy.

[0059] **Figure 21** shows treatment with anti-CD20 reduced glomerular complement activation in mice with adriamycin nephropathy. Mice were injected with anti-CD20 monoclonal antibody to deplete B cells prior to the induction of adriamycin nephropathy. Four weeks after injection with adriamycin, complement activation in the glomeruli was examined by immunofluorescence microscopy. **A)** Staining for C4 demonstrated that glomerular C4 deposition increased after injection with adriamycin, but that treatment of the mice with anti-CD20 prevented this increase. **B)** Staining for C3 demonstrated that glomerular C3 deposition increased after injection with adriamycin. Treatment of the mice with anti-CD20 prevented this increase in C3 deposition. Glomeruli are indicated with arrowheads. Kidneys from three to five mice per group were examined. Thirty glomeruli per section were visualized, and the average for each mouse was determined. Original magnification X200. Scale bar =

100 μ M. After converting the images to grayscale, the brightness and contrast of the shown images were adjusted. All images were adjusted equally.

[0060] **Figure 22** shows that treatment with anti-CD20 reduced albuminuria in mice with adriamycin nephropathy. Mice were injected with anti-CD20 mAb to deplete B cells prior to the induction of adriamycin nephropathy. **A)** Urine albumin/creatinine levels were measured. Treatment of mice with adriamycin caused high-level albuminuria, but this was significantly attenuated by treatment with anti-CD20. Mice treated with anti-CD20 that were re-injected with IgM purified from diseased kidneys had levels of albuminuria similar to those seen in adriamycin treated mice. **B)** Glomerulosclerosis in the different treatment groups was assessed. Kidneys from three to five mice per group were examined. Forty glomeruli per section were visualized. Glomeruli are indicated with arrowheads. **C)** There was less glomerulosclerosis in mice treated with adriamycin and anti-CD20 compared to adriamycin treated mice, but this reduction was not statistically significant. **D)** Staining of kidneys for collagen IV demonstrated that injection of mice with adriamycin caused an increase in glomerular collagen IV deposition, and this was not significantly affected by treatment with anti-CD20. Kidneys from three mice per group were examined. Thirty glomeruli per section were visualized, and the average for each mouse was determined. Representative glomeruli from mice in each group are shown. Original magnification X400. Scale bar = 100 μ M.

[0061] **Figure 23** shows depletion of peritoneal B cells reduced the glomerular deposition of IgM, C3, and C4 in mice with adriamycin nephropathy. Peritoneal cells were depleted with hypotonic shock, starting two weeks prior to inducing adriamycin nephropathy. **A)** Immunofluorescence microscopy demonstrated that depletion of the peritoneal cells attenuated the glomerular deposition of IgM. **B)** Immunofluorescence microscopy for C4 demonstrated that depletion of the peritoneal cells also reduced C4 deposition within the glomeruli, although this did not reach statistical significance. **C)** Immunofluorescence microscopy for C3 demonstrated that depletion of the peritoneal cells prevented complement C3 activation within the glomerulus. Kidneys from three to four mice per group were examined. Thirty glomeruli per section were visualized, and the average for each mouse was determined. Glomeruli are indicated with arrowheads. Original magnification X200. Scale bar = 100 μ M.

[0062] **Figure 24** shows depletion of peritoneal B cells reduced albuminuria in mice with adriamycin nephropathy. Peritoneal cells were depleted with hypotonic

shock, starting two weeks prior to the induction of adriamycin nephropathy. Urine albumin/creatinine levels were measured. Peritoneal depletion of B cells significantly attenuated the level of albuminuria one week (**A**) and four weeks (**B**) after injection of the adriamycin. **C**) Peritoneal cell depletion significantly reduced the degree of glomerulosclerosis compared to control mice that also had adriamycin nephropathy. Kidneys from three to seven mice per group were examined. Forty glomeruli per section were visualized, and the average for each mouse was determined. Representative glomeruli from mice in each group are shown. Glomeruli are indicated with arrowheads. Original magnification x400. Scale bar = 100 μ M. **D**) Staining of kidneys for collagen IV demonstrated that injection of mice with adriamycin caused an increase in glomerular collagen IV deposition, and this was not significantly affected by depletion of peritoneal B cells. Kidneys from four mice per group were examined. Thirty glomeruli per section were visualized, and the average for each mouse was determined. Representative glomeruli from mice in each group are shown. Original magnification x400.

[0063] **Figure 25** shows IgM and C3d co-localized in glomeruli of patients with FSGS. **A**) Available tissue from 19 patients with FSGS was dual-stained for IgM and C3d. In biopsies that contained both IgM and C3d, the two immune factors co-localized within the glomeruli. **B**) Tissue from 19 patients with FSGS was dual-stained for IgM and C4. In biopsies that contained both IgM and C4, the two immune factors appeared to co-localize within the glomeruli. Original magnification X400. Scale bar = 100 μ M. The brightness and contrast of the shown images were adjusted to improve localization of the factors in the overlay.

[0064] **Figure 26** shows IgM deposited in mouse glomeruli after renal ischemia/reperfusion (I/R). Immunofluorescence microscopy revealed that IgM was present in the mesangium of mice 24 hours after sham treatment (**A**, **C**) or renal IR (**B**, **D**). **E**) Quantitative analysis confirmed that mesangial IgM deposition was increased after ischemia. **F**) Western blot analysis under reducing conditions of lysates made from kidneys subjected to sham treatment or IR also demonstrated IgM increase after ischemia. Arrowheads indicate glomeruli. **A and B**, Original magnification 3400; **C and D**, original magnification 3100.

[0065] **Figure 27** shows anti-IgM immunofluorescence images of **A**) B4 mAb and **B**) C2 mAb localization to glomeruli in a renal IR injury mouse model.

[0066] **Figure 28** shows immunofluorescence images of C3 deposition in glomerulus of **A)** wild-type (wt) mice and **B)** factor H deficient (fH^{-/-}) mice.

[0067] **Figure 29** shows immunofluorescence images of IgM deposition in glomerulus of **A)** wild-type (wt) mice at 3 months (left panel), 6 months (middle panel), and 9 months (right panel) of age; and **B)** factor H deficient (fH^{-/-}) mice at 3 months (left panel), 6 months (middle panel), and 9 months (right panel) of age.

[0068] **Figure 30** shows immunofluorescence images of C3 and IgM deposition in glomerulus of factor H deficient (fH^{-/-}) mice at **A)** 3 months and **B)** 9 months of age. **C)** shows immunofluorescence images of C3 and IgM deposition in glomerulus of factor H deficient (fH/ μ MT) mice at 9 months of age.

[0069] **Figure 31** shows immunofluorescence images of **A)** IgM deposition and co-localization with synaptopodin in glomerulus of factor H deficient (fH^{-/-}) mice and **B)** IgM deposition and co-localization with BM marker in glomerulus of factor H deficient (fH^{-/-}) mice.

[0070] **Figure 32** shows immunofluorescence images of C3 and C4 deposition in glomerulus of factor H deficient (fH^{-/-}) mice at **A)** 3 months and **B)** 9 months of age.

[0071] **Figure 33** is a series of graphs showing **A)** serum urea nitrogen (SUN) levels and **B)** urine albumin to creatinine (Cr) ratio in wild-type, fH^{-/-}, and fH/ μ MT mice at 9 months of age.

[0072] **Figure 34** shows **A-D)** a series of flow cytometry histograms of *in vitro* IgM and complement protein binding experiments with mesangial cells. **A)** IgM bound to mesangial cells, **B)** IgG did not bind to mesangial cells, **C)** C3 bound to mesangial cells, and **D)** C4 bound to mesangial cells. **E-J)** Monoclonal IgM clones demonstrated selective binding to glomerular cells *in vivo* and *in vitro*. B cell deficient mice were given an intravenous injection of monoclonal IgM. Kidney sections were assessed for presence of IgM by immunofluorescence. **G)** IgM deposition occurred following injection of the monoclonal IgM clone C2 into a B cell deficient mouse. Representative glomeruli are shown and are marked with arrows. **H)** The corresponding hematoxylin stained section highlighting the location of glomeruli with arrows. **I)** Mice injected with the monoclonal IgM clone D5 did not demonstrate evidence of IgM deposition. **J)** Corresponding hematoxylin stained section. Original magnification x 200. Cultured murine mesangial cells were incubated with polyclonal IgM or seven different monoclonal IgM clones. Following incubation, cells were analyzed by flow cytometry to determine the degree of IgM binding. **E)** IgM

antibodies that exhibited positive binding to mesangial cells are shown. F) The remaining five monoclonal IgM clones all of which did not demonstrate binding to mesangial cells are shown. Isotype control is represented by the shaded histogram.

[0073] **Figure 35** is a graph showing that administration of B4 mAb significantly worsened arthritic symptoms in a model of rheumatoid arthritis.

[0074] **Figure 36** provides a complement pathway analysis in oxidatively-stressed ARPE-19 cell monolayers. **A)** Oxidative stress was induced by treating cells with 500 μ M H_2O_2 , which sensitizes monolayers to complement attack when treating cells with 10% normal human serum (NHS) (Thurman, J. M., Renner, B., Kunchithapautham, K., Ferreira, V. P., Pangburn, M. K., Ablonczy, Z., Tomlinson, S., Holers, V. M., and Rohrer, B. (2009) *J Biol Chem* **284**, 16939-16947). Pathway analysis was performed using serum depleted of specific complement components. Results shown are percentage of starting value in the presence of factor B-, C1q-, MBL-, and C1q/MBL-depleted sera, revealing that complement activation on H_2O_2 -treated cells is triggered by lectin and amplified by the alternative pathway. **B)** Lectin pathway activation was probed in the absence of complement factors C2 or C4 to examine a potential bypass of these components. Elimination of either C2 or C4 from NHS eliminated the effect of H_2O_2 + serum on TER, indicating that both components were for activity. Specificity of the depleted sera was confirmed by reconstituting with purified C2 and C4 protein, respectively.

[0075] **Figure 37** provides an analysis of the pattern recognition receptors involved in lectin pathway activation. **A)** Serum passed over the mannan column was analyzed for components of the lectin pathway. Western blot analysis showed depletion of MBL, MASP-2, and M-, L-, and H-ficolin in two samples (lanes 2 and 3), whereas C3 levels were unaffected. Normal human serum (lane 1) was used as control. **B)** Ficolin binding to ARPE monolayer was examined using NHS as the source in oxidatively-stressed cells, followed by specific antibody binding. Specific, saturable binding could be documented for M-ficolin and H-ficolin, whereas saturable binding was not seen for L-ficolin. Concentration of ficolins was calculated based on known concentrations in human serum. **C)** Reconstitution assays were performed to examine whether ligands on ARPE-19 cells triggering complement activation were recognized by ficolin or MBL, using TER as the readout. TER is reduced by H_2O_2 + serum, but eliminated when serum is passed over a mannan binding column (MBL depl).

Reduction in TER is reconstituted by adding MASP-2 together with one of the pattern recognition receptors; adding all three was not found to be additive.

[0076] Figure 38 shows the results of experiments performed to determine whether natural antibodies activate the lectin pathway. **A)** TER measurements were performed using serum from which either all antibodies (NHS depleted of all Igs) or IgM and IgD (serum from *rag1*^{-/-} mice) was used, indicating that antibodies are required for complement activation in this assay. **B)** IgM binding to ARPE monolayer was examined on ARPE-19 cells cultured as monolayers in 96-well plates, using serum as a source of IgM, followed by colorimetric detection of IgM binding with anti-IgM antibodies. No difference in overall binding of IgM to ARPE-19 cells could be detected when comparing control and oxidatively-stressed cells. **C)** When using an IgM antibody specific for oxidative stress epitopes (IgM-C2), specific binding to cells was observed, which was augmented in the presence of oxidative stress. **D)** Reconstitution assays were performed to determine whether an IgM antibody known to recognize neoepitopes generated by oxidative stress, can activate the complement cascade in Ig-depleted serum. Reduction in TER is obliterated in Ig-depleted serum. Ig-depleted human serum used in the presence of IgM natural antibody, C2, was found to activate the complement cascade in this assay, whereas the control antibody, F1102, was ineffective.

[0077] Figure 39 provides an epitope analysis of IgM-C2 natural antibody. **A)** ELISA analysis was performed, coating plates with BSA coupled to phosphatidylcholine (PC). Specific binding could be observed for IgM-C2 to this ligand as reported previously (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* **188**, 1460-1468), whereas the control IgM specific for annexin IV (IgM-B4) showed no binding. **B)** Since malondialdehyde (MDA) is generated on lipids by oxidative stress, it was examined whether specific binding of IgM-C2 to MDA-BSA could be documented. ELISA assays revealed binding of IgM-C2 to MDA-BSA, albeit possibly at a lower apparent affinity when compared to PC-BSA, or the MDA-BSA wells might have less capacity. No binding could be documented for the control IgM (IgMB4). **C and D)** To test whether reconstitution of Ig-depleted serum using IgM-C2 antibody is mediated by MDA-binding **C)** or unmodified lipid binding **D)**, IgM-C2 was preabsorbed with either BSA-MDA or PC-BSA. BSA-MDA or PC-BSA was added

to Ig-depleted serum as control. Reduction in TER can be mediated through binding of the IgM-C2 antibody to either one of the two lipid ligands.

[0078] **Figure 40** shows the results of experiments performed to determine whether Malondialdehyde (MDA)-neoepitopes are present on oxidatively-stressed ARPE-19 cells. **A)** Immunofluorescence staining of ARPE cells using α -MDA in the presence and absence of H_2O_2 . Specific staining was revealed in oxidatively-stressed cells when compared to control cells. Incubation without primary antibody was performed as a negative control. **B)** Both the anti-MDA (red; α -MDA) and the IgM-C2 antibody (green; α -C2) recognized epitopes present in a punctate fashion on ARPE cells.

[0079] **Figure 41** shows the results of experiments performed to determine whether MDA-neoepitopes are present on oxidatively-stressed primary fetal human cells. Primary fetal human RPE cells were grown in monolayers and TER-assessed in response to 500 μM H_2O_2 and 10% normal human serum (NHS). **A)** Elimination of immunoglobulin (Ig-depleted serum) significantly reduced the drop in TER. The Ig-depleted serum could be reconstituted using the IgM-C2 and the IgM-B4 antibody, and not with a control (IgM-F1102) antibody. **B)** Reconstitution of Ig-depleted serum using IgM-C2 antibody is in part mediated by MDA-binding, as preabsorbing with BSA-MDA eliminated the effect. Oxidative stress-mediated generation of phospholipid neoepitopes is a more general phenomenon for RPE cells.

[0080] **Figure 42** shows the results of experiments performed to determine whether neoepitopes generated by oxidative stress serve as ligands for complement factor H (CFH) on oxidatively-stressed ARPE-19 cells. Malondialdehyde (MDA) has been postulated to serve as a ligand on cell surfaces to recruit CFH and prevent complement-mediated damage (Weismann, D., Hartvigsen, K., Lauer, N., Bennett, K. L., Scholl, H. P., Charbel Issa, P., Cano, M., Brandstatter, H., Tsimikas, S., Skerka, C., Superti-Furga, G., Handa, J. T., Zipfel, P. F., Witztum, J. L., and Binder, C. J. (2011) *Nature* **478**, 76-81). Transepithelial resistance (TER) measurements were performed upon addition of 500 μM H_2O_2 , + 25% normal human serum (NHS); H_2O_2 , + 25% NHS supplemented with 375 μg of purified CFH; cells treated with H_2O_2 , which was removed prior to the addition of HNH + exogenous CFH; or H_2O_2 , + 25% NHS supplemented with 50 μg of CR2-FH (a targeted inhibitory protein of the alternative pathway that targets the inhibitory domain of CFH to sites of C3d deposition). Only CR2-FH was able to block TER reduction induced by H_2O_2 , +

NHS, suggesting that neoepitopes generated by oxidative stress do not recruit CFH to the cell surface for protection.

[0081] **Figure 43** shows the results of experiments performed to determine whether C2-IgM neoepitopes are generated in choroidal neovascularization (CNV) lesions and whether such serve to augment CNV growth in antibody-deficient mice reconstituted with C2-IgM. **A)** Immunofluorescence staining of CNV lesions using the IgM-C2 antibody. Specific staining was revealed when compared to controls in which the primary antibody was omitted. **B)** Antibody-deficient *rag1*^{-/-} mice were reconstituted with three injections of C2-IgM, B4-IgM, or control IgM (F1102 and F632) during the course of CNV development. Both C2-IgM and B4-IgM injections resulted in a significant increase in lesion size when compared to the control antibody. CNV lesion size in wild type mice was unaffected.

DETAILED DESCRIPTION OF THE INVENTION

[0082] The present invention provides targeted delivery methods and constructs for treating inflammatory diseases and/or detecting in vivo tissue injuries in an individual. The targeted delivery approach utilizes an antibody that recognizes an epitope found to be present at sites of inflammation. Specifically, it was found that monoclonal antibodies B4 and C2, initially identified as pathogenic IgM natural antibodies, recognize epitopes widely distributed on organs undergoing ischemia-reperfusion injury as well as sites of inflammation that are undergoing non-ischemic injury. These observations demonstrate the involvement of IgM natural antibodies in inflammatory disorders that go beyond ischemia-reperfusion injury, and suggest a widespread role of these natural antibodies in innate immune recognition of such disorders. The present application thus provides targeted delivery methods and constructs for treating inflammatory diseases and/or detecting in vivo tissue injury based on the binding properties of such natural antibodies.

[0083] Thus, the present application in one aspect provides a method of treating an inflammatory disease in an individual comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement modulator (such as a complement inhibitor).

[0084] In another aspect, there is provided a method of detecting injury or inflammation in a tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety.

[0085] In another aspect, there is provided a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement modulator or a detectable moiety.

[0086] Also provided are methods of delivering a complement modulator (such as a complement inhibitor) or a detectable moiety to a site of tissue injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement modulator (such as a complement inhibitor) or a detectable moiety.

[0087] The present invention also provides methods and compositions for treating ocular diseases. Using a combination of *in vitro* and *in vivo* techniques, it was shown that certain neoepitopes in the eye, namely, Annexin IV and phospholipid-based neoepitopes, are involved during the development of ocular diseases such as age-related macular degeneration (“AMD”). The neoepitopes are recognized by natural antibodies, such as antibodies recognizing the same epitopes as IgM monoclonal antibodies B4 and C2. The binding of the natural antibodies to their respective epitopes in turn lead to the activation of lectin complement pathway and the alternative complement pathway. The present application thus for the first time defines mechanisms of complement activation in oxidatively stressed eye tissue (such as retinal pigmented epithelial monolayers (“RPE”), which provides a basis for the development of therapeutics and diagnostic agents for ocular diseases.

[0088] Thus, the present application in one aspect provides a method of inhibiting inflammation in the eye or treating an ocular disease in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof: (i) specifically binds to

Annexin IV (e.g., an epitope of Annexin IV); or (ii) specifically binds to a phospholipid (e.g., an epitope on a phospholipid).

[0089] In another aspect, there is provided a method of inhibiting inflammation in the eye or treating an ocular disease in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof: (i) specifically binds to Annexin IV (e.g., an epitope of Annexin IV); or (ii) specifically binds to a phospholipid (e.g., an epitope on a phospholipid); and (b) a therapeutic agent (such as a complement inhibitor).

[0090] In another aspect, there is provided a method of detecting injury or inflammation in the eye in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof: (i) specifically binds to Annexin IV (e.g., an epitope of Annexin IV); or (ii) specifically binds to a phospholipid (e.g., an epitope on a phospholipid); and (b) detectable moiety, wherein the presence of the detectable moiety in the eye is indicative of injury or inflammation in the eye.

[0091] Also provided are methods of delivering a complement modulator (such as a complement inhibitor) or a detectable moiety to a site of tissue injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV and competitively inhibits the binding of monoclonal antibody B4 to Annexin IV; and (b) a complement modulator (such as a complement inhibitor) or a detectable moiety.

[0092] Also provided are unit dosage forms, kits, and articles of manufacture that are useful for methods described herein.

Definitions

[0093] The term “individual” refers to a mammal, including humans. An individual includes, but is not limited to, human, bovine, horse, feline, canine, rodent, or primate. In some embodiments, the individual is human. In some embodiments, the individual is a human.

[0094] As used herein, “treatment” or “treating” is an approach for obtaining beneficial or desired results including clinical results. For purposes of this invention, beneficial or desired clinical results include, but are not limited to, one or more of the following: alleviating one or more symptoms resulting from the disease, diminishing the extent of the disease, stabilizing the disease (*e.g.*, preventing or delaying the worsening of the disease), preventing or delaying the spread of the disease, preventing or delaying the recurrence of the disease, delay or slowing the progression of the disease, ameliorating the disease state, providing a remission (partial or total) of the disease, decreasing the dose of one or more other medications required to treat the disease, delaying the progression of the disease, increasing or improving the quality of life, increasing weight gain, and/or prolonging survival. Also encompassed by “treatment” is a reduction of pathological consequence of the disease. The methods of the invention contemplate any one or more of these aspects of treatment.

[0095] The term “effective amount” used herein refers to an amount of a compound or composition sufficient to treat a specified disorder, condition or disease such as ameliorate, palliate, lessen, and/or delay one or more of its symptoms.

[0096] As used herein, by “combination therapy” is meant that a first agent be administered in conjunction with another agent. “In conjunction with” refers to administration of one treatment modality in addition to another treatment modality, such as administration of a nanoparticle composition described herein in addition to administration of the other agent to the same individual. As such, “in conjunction with” refers to administration of one treatment modality before, during, or after delivery of the other treatment modality to the individual.

[0097] As used herein, by “pharmaceutically acceptable” or “pharmacologically compatible” is meant a material that is not biologically or otherwise undesirable, *e.g.*, the material may be incorporated into a pharmaceutical composition administered to an individual or patient without causing any significant undesirable biological effects or interacting in a deleterious manner with any of the other components of the composition in which it is contained. Pharmaceutically acceptable carriers or excipients have preferably met the required standards of toxicological and manufacturing testing and/or are included on the Inactive Ingredient Guide prepared by the U.S. Food and Drug administration.

[0098] As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly indicates otherwise.

[0099] Reference to “about” a value or parameter herein includes (and describes) embodiments that are directed to that value or parameter per se. For example, description referring to “about X” includes description of “X.”

[0100] It is understood that aspects, variations, and embodiments of the invention described herein include “consisting” and/or “consisting essentially of” aspects, variations, and embodiments.

Methods of treating diseases

[0101] The present application in some embodiments provides a method of inhibiting complement activation, inhibiting inflammation, or treating an inflammatory disease in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor. In some embodiments, the composition is administered by injection, such as parenteral, intravenous, subcutaneous, intraocular, intra-articular, or intramuscular injections. In some embodiments, there is provided a method of delivering a complement modulator (such as a complement inhibitor) to a site of tissue injury (such as non-ischemic tissue injury) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor.

[0102] In some embodiments, there is provided a method of inhibiting complement activation (or inhibiting inflammation for example complement-mediated inflammation) in a tissue having a non-ischemic injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual

that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders. In some embodiments, at least about 10% (including for example at least about any of 2%, 3%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0103] In some embodiments, there is provided a method of inhibiting complement activation (or inhibiting inflammation for example complement-mediated inflammation) in a tissue having a non-ischemic injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some

embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders. In some embodiments, at least about 10% (including for example at least about any of 2-%, 3-%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0104] In some embodiments, there is provided a method of inhibiting complement activation (or inhibiting inflammation, for example complement-mediated inflammation) in a tissue having an oxidative damage in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral

nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders. In some embodiments, at least about 10% (including for example at least about any of 2-%, 3-%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0105] In some embodiments, there is provided a method of inhibiting complement activation (or inhibiting inflammation, for example complement-mediated inflammation) in a tissue having an oxidative damage in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets

and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders. In some embodiments, at least about 10% (including for example at least about any of 2-%, 3-%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0106] In some embodiments, there is provided a method of treating an inflammatory disease (or a disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease is any of inflammatory disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft injection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0107] In some embodiments, there is provided a method of treating an inflammatory disease (or a disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and

(b) a complement inhibitor. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease is any of inflammatory disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft rejection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0108] In some embodiments, the disease to be treated is an ocular disease. In some embodiments, the disease is an ocular disease associated with complement activation. In some embodiments, the disease is age-related macular degeneration (“AMD”), including wet AMD and dry AMD. Other ocular diseases that can be treated by methods described herein include, but are not limited to, CMV retinitis, macular edema, uveitis, glaucoma, diabetic retinopathy, retinitis pigmentosa, retinal detachment, proliferative vitreoretinopathy and ocular melanoma.

[0109] In some embodiments, the disease to be treated is inflammatory arthritis.

[0110] In some embodiments, the disease to be treated is a kidney disease, including but is not limited to, acute kidney injury, glomerulonephritis, chronic kidney disease, and focal segmental glomerulosclerosis.

[0111] In some embodiments, the disease to be treated is an inflammatory disorder, which include, but is not limited to, burns, endotoxemia, septic shock, adult respiratory distress syndrome, cardiopulmonary bypass, hemodialysis, anaphylactic

shock, asthma, angioedema, Crohn's disease, sickle cell anemia, poststreptococcal glomerulonephritis, membranous nephritis, and pancreatitis.

[0112] In some embodiments, the disease to be treated is a pregnancy-related disease, which includes, but is not limited to, HELLP (Hemolytic anemia, elevated liver enzymes, and low platelet count), recurrent fetal loss, and pre-eclampsia.

[0113] In some embodiments, the disease to be treated is an autoimmune or immune complex disorder, which include, but is not limited to, myasthenia gravis, Alzheimer's disease, multiple sclerosis, neuromyelitis optica, rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, lupus nephritis, IgG4 associated diseases, insulin-dependent diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, antiphospholipid antibody syndrome, thrombotic thrombocytopenic purpura, autoimmune hepatitis, Crohn's disease, Goodpasture's syndromes, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, idiopathic thrombocytopenic purpura, pemphigus, Sjogren's syndrome, Takayasu's arteritis, autoimmune glomerulonephritis, membranoproliferative glomerulonephritis type II, membranous disease, paroxysmal nocturnal hemoglobinuria, age-related macular degeneration, diabetic maculopathy, uveitis, retinal degeneration disorders, diabetic nephropathy, focal segmental glomerulosclerosis, ANCA associated vasculitis, hemolytic uremic syndrome, Shiga-toxin-associated hemolytic uremic syndrome, and atypical hemolytic uremic syndrome. In some embodiments, the disease to be treated is an autoimmune glomerulonephritis, which includes, but is not limited to, immunoglobulin A nephropathy or membranoproliferative glomerular nephritis type I.

Methods of treating ocular diseases

[0114] The present application in some embodiments provides a method of inhibiting complement activation in the eye, inhibiting inflammation in the eye, or treating an ocular disease (for example an ocular disease involving oxidative damage or a complement-associated ocular disease) in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof: (i) specifically binds to Annexin IV; or (ii) specifically binds to a phospholipid.

[0115] In some embodiments, there is provided a method of inhibiting complement activation or inhibiting inflammation (such as complement-driven inflammation) in

the eye in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0116] In some embodiments, there is provided a method of inhibiting oxidative damage to eye (including for example oxidative damage to the photoreceptors, RPE/Bruch's membrane, macula, and/or choriocapillary complex) in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, at least about 10% (including for

example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) oxidative damage is inhibited.

[0117] In some embodiments, there is provided a method of treating an ocular disease (such as a complement-associated ocular disease or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0118] In some embodiments, there is provided a method of inhibiting complement activation or inhibiting inflammation (such as complement-driven inflammation) in the eye in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to a phospholipid. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is

present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0119] In some embodiments, there is provided a method of inhibiting oxidative damage to eye (including for example oxidative damage to the photoreceptors, RPE/Bruch's membrane, macula, and/or choriocapillary complex) in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to a phospholipid. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) oxidative damage is inhibited.

[0120] In some embodiments, there is provided a method of treating an ocular disease (such as a complement-associated ocular disease or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to a phospholipid. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0121] In another aspect, there is provided a method of inhibiting inflammation in the eye or treating an ocular disease in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof: (i) specifically binds to Annexin IV (*e.g.*, an epitope of Annexin IV); or (ii) specifically binds to a phospholipid (*e.g.*, an epitope on a phospholipid); and (b) a therapeutic agent (such as a complement inhibitor).

[0122] In some embodiments, there is provided a method of inhibiting complement activation or inhibiting inflammation (such as complement-driven inflammation) in the eye in an individual, comprising administering to the individual an effective

amount of a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0123] In some embodiments, there is provided a method of inhibiting oxidative damage to eye (including for example oxidative damage to the photoreceptors, RPE/Bruch's membrane, macula, and/or choriocapillary complex) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, at least about 10% (including for example at least about any of 20%,

30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) oxidative damage is inhibited.

[0124] In some embodiments, there is provided a method of treating an ocular disease (such as a complement-associated ocular disease or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0125] In some embodiments, there is provided a method of inhibiting complement activation or inhibiting inflammation (such as complement-driven inflammation) in the eye in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to a phospholipid, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some

embodiments, the phospholipid is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) complement activation or inflammation is inhibited.

[0126] In some embodiments, there is provided a method of inhibiting oxidative damage to eye (including for example oxidative damage to the photoreceptors, RPE/Bruch's membrane, macula, and/or choriocapillary complex) in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to a phospholipid, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, at least about 10% (including for example at least about any of 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, or 100%) oxidative damage is inhibited.

[0127] In some embodiments, there is provided a method of treating an ocular disease (such as a complement-associated ocular disease or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to a phospholipid, and (b) a therapeutic agent (such as a complement inhibitor). In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (*e.g.*, Bruch's membrane), or in a pathological structure (*e.g.*, drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0128] The methods described herein are also useful for any one of more of the following: (1) inhibiting, preventing, or delaying the formation of drusen in the eye; (2) inhibiting, preventing, or delaying loss of photoreceptor cells; (3) inhibiting, preventing, or delaying neovascularization associated with an ocular disease (such as AMD); (3) inhibiting, preventing, or delaying retinal detachment; (4) inhibiting, preventing, or delaying oxidative stress-mediated injury; and (5) improving visual acuity or visual field in the eye of an individual.

Methods of detecting tissue injury

[0129] The present application in some embodiments provides a method of detecting a complement-mediated tissue injury or inflammation or diagnosing an inflammatory disease in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety. In some embodiments, there is provided a method of delivering a detectable moiety to a site of complement-mediated tissue injury or inflammation in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the composition is administered by injection, such as parenteral, intravenous, subcutaneous, intraocular, intra-articular, or intramuscular injections.

[0130] In some embodiments, there is provided a method of detecting a complement-mediated tissue injury or inflammation or diagnosing an inflammatory disease in an individual, comprising contacting a tissue of the individual with a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety. In some embodiments, the method further comprises detecting the detectable moiety.

[0131] In some embodiments, there is provided a method of detecting complement-mediated injury (or detecting inflammation for example complement-mediated inflammation) in a tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated tissue injury (or complement-mediated inflammation). In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments,

the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0132] In some embodiments, there is provided a method of detecting complement-mediated injury (or detecting inflammation for example complement-mediated inflammation) in a tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated tissue injury (or complement-mediated inflammation). In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of

undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0133] In some embodiments, there is provided a method of detecting oxidative damage (or an inflammatory disease involving oxidative damage) in a tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a oxidative damage in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein.

In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the oxidative damage is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0134] In some embodiments, there is provided a method of detecting oxidative damage (or an inflammatory disease involving oxidative damage) in a tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a oxidative damage in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver

or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the oxidative damage is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0135] In some embodiments, there is provided a method of detecting non-ischemic tissue injury (or an inflammatory disease involving non-ischemic tissue injury) in a tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of non-ischemic tissue injury. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the non-ischemic tissue injury is

associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0136] In some embodiments, there is provided a method of detecting non-ischemic tissue injury (or an inflammatory disease involving non-ischemic tissue injury) in a tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of non-ischemic tissue injury. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (*e.g.*, drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the tissue is any one of liver or portal tract, heart, muscle, brain, central or peripheral nervous system, gastrointestinal tract, lung, limb, arterial or venous vascular system, skin, bone marrow cells including red blood cells, platelets and nucleated cells, pancreas, eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the tissue is any one of eye, joint, and kidney. In some embodiments, the non-ischemic tissue injury is associated with tissue damage resulting from inflammatory disorders, transplant rejection (cellular or antibody

mediated), pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.

[0137] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an tissue-specific inflammatory disease (or a disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of the inflammatory disease in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease to be diagnosed is any of inflammatory disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft injection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0138] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an tissue-specific inflammatory disease (or a disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of the inflammatory disease in the tissue. In some embodiments, the method further comprises detecting

the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (e.g., drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease to be diagnosed is any of inflammatory disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft rejection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0139] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an tissue-specific inflammatory disease (or a disease involving oxidative damage) in an individual, comprising contacting a tissue sample from an individual with an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of the inflammatory disease in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual

that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease to be diagnosed is any of inflammatory disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft rejection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0140] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an tissue-specific inflammatory disease (or a disease involving oxidative damage) in an individual, comprising contacting a tissue sample from an individual with an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of the inflammatory disease in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell (or in a pathological structure (e.g., drusen)) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the inflammatory disease to be diagnosed is any of inflammatory

disorders, transplant rejection (cellular or antibody mediated, such as hyperacute xenograft rejection), pregnancy-related diseases, adverse drug reactions (such as drug allergy and IL-2 induced vascular leakage syndrome), autoimmune or immune complex disorders.

[0141] In some embodiments, the disease to be diagnosed is an ocular disease. In some embodiments, the disease is an ocular disease associated with complement activation. In some embodiments, the disease is age-related macular degeneration (“AMD”), including wet AMD and dry AMD. Other diseases that can be diagnosed by methods described herein include, but are not limited to, CMV retinitis, macular edema, uveitis, glaucoma, diabetic retinopathy, retinitis pigmentosa, retinal detachment, proliferative vitreoretinopathy and ocular melanoma.

[0142] In some embodiments, the disease to be diagnosed is inflammatory arthritis.

[0143] In some embodiments, the disease to be diagnosed is a kidney disease, including but is not limited to, acute kidney injury, glomerulonephritis, chronic kidney disease, and focal segmental glomerulosclerosis.

[0144] In some embodiments, the disease to be diagnosed is an inflammatory disorder, which include, but is not limited to, burns, endotoxemia, septic shock, adult respiratory distress syndrome, cardiopulmonary bypass, hemodialysis, anaphylactic shock, asthma, angioedema, Crohn's disease, sickle cell anemia, poststreptococcal glomerulonephritis, membranous nephritis, and pancreatitis.

[0145] In some embodiments, the disease to be diagnosed is a pregnancy-related disease, which includes, but is not limited to, HELLP (Hemolytic anemia, elevated liver enzymes, and low platelet count), recurrent fetal loss, and pre-eclampsia.

[0146] In some embodiments, the disease to be diagnosed is an autoimmune or immune complex disorder, which include, but is not limited to, myasthenia gravis, Alzheimer's disease, multiple sclerosis, neuromyelitis optica, rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, lupus nephritis, IgG4 associated diseases, insulin-dependent diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, antiphospholipid antibody syndrome, thrombotic thrombocytopenic purpura, autoimmune hepatitis, Crohn's disease, Goodpasture's syndromes, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, idiopathic thrombocytopenic purpura, pemphigus, Sjogren's syndrome, Takayasu's arteritis, autoimmune glomerulonephritis, membranoproliferative glomerulonephritis type II, membranous disease, paroxysmal nocturnal hemoglobinuria, age-related macular

degeneration, diabetic maculopathy, uveitis, retinal degeneration disorders, diabetic nephropathy, focal segmental glomerulosclerosis, ANCA associated vasculitis, hemolytic uremic syndrome, Shiga-toxin-associated hemolytic uremic syndrome, and atypical hemolytic uremic syndrome. In some embodiments, the disease to be diagnosed is an autoimmune glomerulonephritis, which includes, but is not limited to, immunoglobulin A nephropathy or membranoproliferative glomerular nephritis type I.

[0147] Methods of detecting tissue injury in the eye

[0148] Also provided herein are methods of detecting injury or inflammation in the eye in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises: (a) an antibody or a fragment thereof, wherein the antibody or fragment thereof: (i) specifically binds to Annexin IV or (ii) specifically binds to phospholipid; and (b) detectable moiety, wherein the presence of the detectable moiety in the eye is indicative of injury or inflammation in the eye. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the composition is administered by injection, such as intraocular injections.

[0149] In some embodiments, the method is useful for detecting deterioration of the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea. In some embodiments, the method is useful for detecting inflammation in the of the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea. In some embodiments, the method is useful for detecting oxidative damage of the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea. In some embodiments, the method is useful for detecting pathological lesions at the RPE/choroid interface in the macula.

[0150] In some embodiments, there is provided a method of detecting complement-mediated injury (or detecting inflammation for example complement-mediated inflammation) in an eye tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated eye tissue injury (or complement-mediated inflammation). In

some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the detectable moiety are linked via a linker (such as a peptide linker). In some embodiments, the eye tissue is the photoreceptor cell, retinal ganglion cell, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea. In some embodiments, there is provided a method of detecting oxidative damage in an eye tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a oxidative damage in the eye tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the detectable moiety are linked via a linker (such as a peptide

linker). In some embodiments, the eye tissue is the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea.

[0151] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an inflammatory ocular disease (or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a detectable moiety, wherein the presence of the detectable moiety at the eye tissue is indicative of the inflammatory ocular disease (or an ocular disease involving oxidative damage) in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the detectable moiety are linked via a linker (such as a peptide linker). In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0152] In some embodiments, there is provided a method of detecting complement-mediated injury (or detecting inflammation for example complement-mediated inflammation) in an eye tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the

construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated eye tissue injury (or complement-mediated inflammation). In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the eye tissue is the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea.

[0153] In some embodiments, there is provided a method of detecting oxidative damage in an eye tissue in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a oxidative damage in the eye tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue

undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the eye tissue is the photoreceptor cells, retinal ganglion cells, RPE, Bruch's membrane, choriocapillary complex, macula, or cornea.

[0154] In some embodiments, there is provided a method of diagnosing (or assisting in diagnosing) an inflammatory ocular disease (or an ocular disease involving oxidative damage) in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the eye tissue is indicative of the inflammatory ocular disease (or an ocular disease involving oxidative damage) in the tissue. In some embodiments, the method further comprises detecting the detectable moiety. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal

wound healing, corneal transplants, and ocular melanoma. In some embodiments, the ocular disease is AMD.

[0155] Targeting constructs

[0156] The present application in some embodiments provides targeted constructs which can be useful, but are not limited, any one or more of the methods described herein. It is to be understood that any of the constructs described in the section herein can be used for any of the methods described in the sections above. The present application further provides methods of delivering any of the complement modulator or detectable moiety disclosed herein to a site of complement activation, a site of tissue injury (such as non-ischemic tissue injury), or a site of complement-associated disease in an individual by administering to the individual any one of the target constructs described herein.

[0157] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a therapeutic agent or a detectable moiety. In some embodiments, the construct comprises a therapeutic agent (such as a complement modulator, for example a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the antibody or fragment thereof (hereinafter also referred to as the “targeting moiety” and the detectable moiety (hereinafter also referred to as “the active moiety” are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0158] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1, a sequence of SEQ ID NO:2, or a sequence of SEQ ID NO:3; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4, a sequence of SEQ ID NO:5, or a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a

composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7, a sequence of SEQ ID NO:8, or a sequence of SEQ ID NO:9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:10, a sequence of SEQ ID NO:11, or a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0159] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the

antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0160] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof

competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0161] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (vi) heavy chain variable domain

comprising a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0162] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:1; (ii) a light chain CDR2 of SEQ ID NO:2; (iii) a light chain CDR3 of SEQ ID NO:3; (iv) heavy chain CDR1 of SEQ ID NO:4; (v) heavy chain CDR2 of SEQ ID NO:5; and (vi) heavy chain CDR3 of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:7; (ii) a light chain CDR2 of SEQ ID NO:8; (iii) a light chain CDR3 of SEQ ID NO:9; (iv) heavy chain CDR1 of SEQ ID NO:10; (v) heavy chain CDR2 of SEQ ID NO:11; and (vi) heavy chain CDR3 of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a

pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0163] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:13. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:15. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:14. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:16. In some embodiments, the antibody or

fragment thereof competitively inhibits the binding of a pathogenic monoclonal antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0164] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:13; and (ii) heavy chain variable domain of SEQ ID NO:15. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:14; and (ii) heavy chain variable domain of SEQ ID NO:16. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface

of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0165] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:17. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:18. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting

moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0166] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25, a sequence of SEQ ID NO:26, or a sequence of SEQ ID NO:27; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31, a sequence of SEQ ID NO:32, or a sequence of SEQ ID NO:33; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor).

In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0167] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments,

the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0168] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0169] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some

embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0170] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:25; (ii) a light chain CDR2 of SEQ ID NO:26; (iii) a light chain CDR3 of SEQ ID NO:27; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:31; (ii) a light chain CDR2 of SEQ ID NO:32; (iii) a light chain CDR3 of SEQ ID NO:33; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and

phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0171] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:34. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety a detectable moiety), wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:36. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:35. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an

individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0172] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:34; and (ii) heavy chain variable domain of SEQ ID NO:36.

In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:35; and (ii) heavy chain variable domain of SEQ ID NO:36.

In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid.

In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid.

In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some

embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0173] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:37. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or detectable moiety), wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:38. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is

positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0174] In some embodiments, the targeting moiety and the active moiety are directly bonded, covalently bonded, or, reversibly bonded.

[0175] A “construct” or “targeting construct” used herein refers to a non-naturally occurring molecule comprising a “targeting moiety” and an “active moiety”. The targeting moiety is capable of specifically binding to Annexin IV. The targeting moiety of the targeting construct is responsible for targeted delivery of the molecule to the sites of, e.g., complement activation. The active moiety is responsible for therapeutic activity, e.g., specifically inhibiting complement activation, or detection, e.g., permitting the detection and or localization of the targeting moiety. The targeting moiety and the active moiety of a targeting construct molecule can be linked together by any methods known in the art, as long as the desired functionalities of the two portions are maintained.

[0176] The targeting construct described herein thus generally has the dual functions of binding to an epitope recognized by an antibody described herein and exerting therapeutic activity or allowing detection. A “epitope of monoclonal antibody B4 antibody” refers to any molecule that binds to a naturally occurring B4 or C2 antibody, which include, epitopes that bind to a B4 or C2 antibody with a binding affinity that is about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of the epitope that naturally binds a B4 antibody. Binding affinity can be determined by any method known in the art, including for example, surface plasmon resonance, calorimetry titration, ELISA, and flow cytometry.

[0177] In some embodiments, a targeting construct described herein is generally capable of inhibiting complement activation (for example inhibiting activation of the alternative pathway and/or lectin pathway). The targeting construct may be a more potent complement inhibitor than the naturally occurring antibody as described herein. For example, in some embodiments, the targeting construct has a complement inhibitory activity that is about any of 1.5, 2, 2.5, 3, 3.5, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 40, or more fold of that of a B4 or C2 antibody. In some embodiments,

the targeting construct has an EC₅₀ of less than about any of 100 nM, 90 nM, 80 nM, 70 nM, 60 nM, 50 nM, 40 nM, 30 nM, 20 nM, or 10 nM, inclusive, including any values in between these numbers. In some embodiments, the targeting construct molecule has an EC₅₀ of about 5 to 60 nM, including for example any of 8 to 50 nM, 8 to 20 nM, 10 to 40 nM, and 20 to 30 nM. In some embodiments, the targeting construct molecule has complement inhibitory activity that is about any of 50%, 60%, 70%, 80%, 90%, or 100% of that of a B4 or C2 antibody.

[0178] Complement inhibition can be evaluated based on any methods known in the art, including for example, in vitro zymosan assays, assays for lysis of erythrocytes, antibody or immune complex activation assays, alternative pathway activation assays, and mannan activation assays.

[0179] In some embodiments, the targeting construct is a fusion protein. “Fusion protein” used herein refers to two or more peptides, polypeptides, or proteins operably linked to each other. In some embodiments, the targeting moiety and the active moiety are directly fused to each other. In some embodiments, the targeting moiety and the active moiety are linked by an amino acid linker sequence. Examples of linker sequences are known in the art, and include, for example, (Gly4Ser), (Gly4Ser)₂, (Gly4Ser)₃, (Gly3Ser)₄, (SerGly4), (SerGly4)₂, (SerGly4)₃, and (SerGly4)₄. Linking sequences can also comprise “natural” linking sequences found between different domains of complement factors. The order of targeting moiety and active moiety in the fusion protein can vary. For example, in some embodiments, the C-terminus of the targeting moiety is fused (directly or indirectly) to the N-terminus of the active moiety of the targeting construct. In some embodiments, the N-terminus of the targeting moiety is fused (directly or indirectly) to the C-terminus of the active moiety of the targeting construct.

[0180] In some embodiments, the targeting moiety of a targeting construct is encoded by a polynucleotide comprising a nucleic acid sequence of any of SEQ ID NOs: 19-24 and 57. In some embodiments, the targeting construct molecule is encoded by a polynucleotide comprising a nucleic acid sequence that is at least about 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to that of any of SEQ ID NOs: 19-24 and 57.

[0181] In some embodiments, the targeting moiety of a targeting construct is encoded by a polynucleotide comprising a nucleic acid sequence of any of SEQ ID NOs: 37 or 38. In some embodiments, the targeting construct molecule is encoded by

a polynucleotide comprising a nucleic acid sequence that is at least about 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to that of any of SEQ ID NOs: 37 or 38.

[0182] In some embodiments, the targeting construct comprises a targeting moiety and an active moiety linked via a chemical cross-linker. Linking of the two portions can occur on reactive groups located on the two moieties. Reactive groups that can be targeted using a crosslinker include primary amines, sulfhydryls, carbonyls, carbohydrates, and carboxylic acids, or active groups that can be added to proteins. Examples of chemical linkers are well known in the art and include, but are not limited to, bismaleimido hexane, maleimidobenzoyl-N-hydroxysuccinimide ester, NHS-Esters-Maleimide Crosslinkers such as SPDP, carbodiimide, glutaraldehyde, MBS, Sulfo-MBS, SMPB, sulfo-SMPB, GMBS, Sulfo-GMBS, EMCS, Sulfo-EMCS, imidoester crosslinkers such as DMA, DMP, DMS, DTBP, EDC and DTME.

[0183] In some embodiments, the targeting moiety and the active moiety are non-covalently linked. For example, the two portions may be brought together by two interacting bridging proteins (such as biotin and streptavidin), each linked to a targeting moiety or an active moiety.

[0184] In some embodiments, the targeting construct comprises two or more (same or different) targeting moieties described herein. In some embodiments, the targeting construct comprises two or more (same or different) active moieties described herein. These two or more targeting (or active) moieties may be tandemly linked (such as fused) to each other. In some embodiments, the targeting construct comprises a targeting moiety and two or more (such as three, four, five, or more) active moieties. In some embodiments, the targeting construct comprises an active moiety and two or more (such as three, four, five, or more) targeting moieties. In some embodiments, the targeting construct comprises two or more targeting moieties and two or more active moieties.

[0185] In some embodiments, there is provided an isolated targeting construct. In some embodiments, the targeting constructs form dimers or multimers.

[0186] The active moiety and the targeting moiety in the targeting construct can be from the same species (such as human or mouse), or from different species.

[0187] Also provided herein are targeting constructs and compositions (such as pharmaceutical compositions) comprising a targeting construct molecule. The present application further provides methods of delivering any of the complement modulator

or detectable moiety disclosed herein to a site of complement activation, a site of tissue injury (such as non-ischemic tissue injury), or a site of complement-associated disease in an individual by administering to the individual any one of the target constructs described herein.

[0188] A “targeting construct” used herein refers to a non-naturally occurring molecule comprising a “targeting moiety” and an “active moiety”. In certain embodiments, the targeting moiety is capable of binding to Annexin IV. In certain embodiments, the targeting moiety is capable of binding a phospholipid, such as PC, PE, and/or CL. The targeting moiety of the targeting construct is thus responsible for targeted delivery of the molecule to the sites of, e.g., complement activation. The active moiety is responsible for therapeutic activity, e.g., specifically inhibiting complement activation, or detection, e.g., permitting the detection and or localization of the targeting moiety. The targeting moiety and the active moiety of a targeting construct molecule can be linked together by any methods known in the art, as long as the desired functionalities of the two portions are maintained.

[0189] The targeting construct described herein thus generally has the dual functions of binding to an epitope recognized by an antibody described herein and exerting therapeutic activity or allowing detection. A “epitope of a B4 or C2 antibody” refers to any molecule that binds to a naturally occurring B4 or C2 antibody, which include, epitopes that bind to a B4 or C2 antibody with a binding affinity that is about any of 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% of the epitope that naturally binds a B4 or C2 antibody. Binding affinity can be determined by any method known in the art, including for example, surface plasmon resonance, calorimetry titration, ELISA, and flow cytometry.

[0190] In some embodiments, a targeting construct described herein is generally capable of inhibiting complement activation (for example inhibiting activation of the alternative pathway and/or lectin pathway). The targeting construct may be a more potent complement inhibitor than the naturally occurring antibody as described herein. For example, in some embodiments, the targeting construct has a complement inhibitory activity that is about any of 1.5, 2, 2.5, 3, 3.5, 4, 5, 6, 7, 8, 9, 10, 12, 14, 16, 18, 20, 25, 30, 40, or more fold of that of a B4 or C2 antibody. In some embodiments, the targeting construct has an EC₅₀ of less than about any of 100 nM, 90 nM, 80 nM, 70 nM, 60 nM, 50 nM, 40 nM, 30 nM, 20 nM, or 10 nM. In some embodiments, the targeting construct molecule has an EC₅₀ of about 5-60 nM, including for example

any of 8-50 nM, 8-20 nM, 10-40 nM, and 20-30 nM. In some embodiments, the targeting construct molecule has complement inhibitory activity that is about any of 50%, 60%, 70%, 80%, 90%, or 100% of that of a B4 or C2 antibody.

[0191] Complement inhibition can be evaluated based on any methods known in the art, including for example, in vitro zymosan assays, assays for lysis of erythrocytes, immune complex activation assays, and mannan activation assays.

[0192] In some embodiments, the targeting construct is a fusion protein. “Fusion protein” used herein refers to two or more peptides, polypeptides, or proteins operably linked to each other. In some embodiments, the targeting moiety and the active moiety are directly fused to each other. In some embodiments, the targeting moiety and the active moiety are linked by an amino acid linker sequence. Examples of linker sequences are known in the art, and include, for example, (Gly₄Ser), (Gly₄Ser)₂, (Gly₄Ser)₃, (Gly₃Ser)₄, (SerGly₄), (SerGly₄)₂, (SerGly₄)₃, and (SerGly₄)₄. Linking sequences can also comprise “natural” linking sequences found between different domains of complement factors. The order of targeting moiety and active moiety in the fusion protein can vary. For example, in some embodiments, the C-terminus of the targeting moiety is fused (directly or indirectly) to the N-terminus of the active moiety of the targeting construct. In some embodiments, the N-terminus of the targeting moiety is fused (directly or indirectly) to the C-terminus of the active moiety of the targeting construct.

[0193] In some embodiments, the targeting moiety of a targeting construct is encoded by a polynucleotide comprising a nucleic acid sequence of any of SEQ ID NOs: 19-24, 39-43, 57 and 58. In some embodiments, the targeting construct molecule is encoded by a polynucleotide comprising a nucleic acid sequence that is at least about 50%, 60%, 70%, 80%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to that of any of SEQ ID NOs: 19-24 and 39-43.

[0194] In some embodiments, the targeting construct comprises a targeting moiety and an active moiety linked via a chemical cross-linker. Linking of the two portions can occur on reactive groups located on the two moieties. Reactive groups that can be targeted using a crosslinker include primary amines, sulfhydryls, carbonyls, carbohydrates, and carboxylic acids, or active groups that can be added to proteins. Examples of chemical linkers are well known in the art and include, but are not limited to, bismaleimido hexane, maleimidobenzoyl-N-hydroxysuccinimide ester, NHS-Esters-Maleimide Crosslinkers such as SPDP, carbodiimide, glutaraldehyde,

MBS, Sulfo-MBS, SMPB, sulfo-SMPB, GMBS, Sulfo-GMBS, EMCS, Sulfo-EMCS, imidoester crosslinkers such as DMA, DMP, DMS, DTBP, EDC and DTME.

[0195] In some embodiments, the targeting moiety and the active moiety are non-covalently linked. For example, the two portions may be brought together by two interacting bridging proteins (such as biotin and streptavidin), each linked to a targeting moiety or an active moiety.

[0196] In some embodiments, the targeting construct comprises two or more (same or different) targeting moieties described herein. In some embodiments, the targeting construct comprises two or more (same or different) active moieties described herein. These two or more targeting (or active) moieties may be tandemly linked (such as fused) to each other. In some embodiments, the targeting construct comprises a targeting moiety and two or more (such as three, four, five, or more) active moieties. In some embodiments, the targeting construct comprises an active moiety and two or more (such as three, four, five, or more) targeting moieties. In some embodiments, the targeting construct comprises two or more targeting moieties and two or more active moieties.

[0197] In some embodiments, there is provided an isolated targeting construct. In some embodiments, the targeting constructs form dimers or multimers.

[0198] The active moiety and the targeting moiety in the targeting construct can be from the same species (such as human or mouse), or from different species.

[0199] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a therapeutic agent or a detectable moiety. In some embodiments, the construct comprises a therapeutic agent (such as a complement modulator, for example a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the antibody or fragment thereof (hereinafter also referred to as the “targeting moiety” and the detectable moiety (hereinafter also referred to as “the active moiety” are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0200] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the

construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1, a sequence of SEQ ID NO:2, or a sequence of SEQ ID NO:3; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4, a sequence of SEQ ID NO:5, or a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7, a sequence of SEQ ID NO:8, or a sequence of SEQ ID NO:9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:10, a sequence of SEQ ID NO:11, or a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0201] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a

light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0202] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a

pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0203] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or

a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0204] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:1; (ii) a light chain CDR2 of SEQ ID NO:2; (iii) a light chain CDR3 of SEQ ID NO:3; (iv) heavy chain CDR1 of SEQ ID NO:4; (v) heavy chain CDR2 of SEQ ID NO:5; and (vi) heavy chain CDR3 of SEQ ID NO:6. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable

moiety, wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:7; (ii) a light chain CDR2 of SEQ ID NO:8; (iii) a light chain CDR3 of SEQ ID NO:9; (iv) heavy chain CDR1 of SEQ ID NO:10; (v) heavy chain CDR2 of SEQ ID NO:11; and (vi) heavy chain CDR3 of SEQ ID NO:12. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0205] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:13. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:15. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable

moiety, wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:14. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:16. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic monoclonal antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0206] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:13; and (ii) heavy chain variable domain of SEQ ID NO:15. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator

or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:14; and (ii) heavy chain variable domain of SEQ ID NO:16. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0207] In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:17. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:18. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody B4) to Annexin IV. In some embodiments, the antibody or fragment thereof binds to the same epitope as a pathogenic antibody (such as a monoclonal antibody B4) to Annexin IV. In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury

(such as non-ischemic injury) and/or oxidative damage. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein. In some embodiments, the construct comprises a complement modulator (such as a complement inhibitor). In some embodiments, the construct comprises a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker). In some embodiments, the targeting moiety and the active moiety are directly linked.

[0208] The present application in some embodiments provides various targeting constructs for targeted delivery. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25, a sequence of SEQ ID NO:26, or a sequence of SEQ ID NO:27; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, there is provided a construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31, a sequence of SEQ ID NO:32, or a sequence of SEQ ID NO:33; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such

as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0209] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in

a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0210] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments,

the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0211] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement

membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0212] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:25; (ii) a light chain CDR2 of SEQ ID NO:26; (iii) a light chain CDR3 of SEQ ID NO:27; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:31; (ii) a light chain CDR2 of SEQ ID NO:32; (iii) a light chain CDR3 of SEQ ID NO:33; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as

monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0213] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:34. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:36. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:35. In

some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0214] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:34; and (ii) heavy chain variable domain of SEQ ID NO:36.

In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:35; and (ii) heavy chain variable domain of SEQ ID NO:36.

In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid.

In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0215] In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or a detectable moiety), wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:37. In some embodiments, there is provided a targeting construct (or a composition comprising the construct such as a pharmaceutical composition), wherein the targeting construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid (such as PE, CL, MDA, and/or PC); and (b) an active moiety (e.g., a therapeutic moiety or detectable moiety), wherein the antibody or fragment is a scFv having the sequence of SEQ ID NO:38. In some embodiments, the antibody or fragment thereof competitively inhibits the binding of a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the antibody or antibody fragment thereof binds to the same epitope as a pathogenic antibody (such as monoclonal antibody C2) to phospholipid. In some embodiments, the phospholipid is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure

(e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury (such as non-ischemic injury) and/or oxidative damage. In some embodiments, the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC). In some embodiments, the phospholipid is malondialdehyde (MDA). In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid is oxidized. In some embodiments, the construct comprises an active moiety that comprises a therapeutic moiety (such as a complement inhibitor). In some embodiments, the construct comprises an active moiety that is a detectable moiety. In some embodiments, the construct is a fusion protein. In some embodiments, the targeting moiety and the active moiety are linked via a linker (such as a peptide linker).

[0216] In some embodiments, the targeting moiety and the active moiety are directly bonded, covalently bonded, or, reversibly bonded.

[0217] Targeting moieties (e.g., antibodies recognizing an injury-associated neoepitope)

[0218] The antibody or fragment thereof described herein (also referred to as the targeting moiety when provided in the context of a targeting construct) specifically bind to Annexin IV or a phospholipid.

[0219] The antibody or fragment thereof described herein (also referred to as the targeting moiety when provided in the context of a targeting construct) in some embodiments specifically bind to Annexin IV.

[0220] Annexin IV belongs to a family of proteins that are Ca^{2+} and phospholipid proteins. The structure of annexins consists of a conserved Ca^{2+} and membrane binding core of four annexin repeats (eight for annexin IV) and variable N-terminal regions. Annexins are soluble cytosolic proteins, but despite the lack of obvious signal sequences and the apparent inability to enter the classical secretory pathway, annexins have been identified in extracellular fluids or associated with the external cell surface through poorly understood binding sites. Annexin IV is predominantly produced by epithelial cells and is also found at high levels in lung, intestine, pancreas, liver, photoreceptors, and kidney. Rescher et al., J. Cell Sci., (2004), 117:2631-2639, Kulik et al., (2009) J Immunol. 182(9):5363-73, and Zernii et al., Biochemistry (Mosc.), (2003), 68(1):129-60. It is also present in drusen, the

hallmarks of age-related macular degeneration (AMD) (Rayborn, M. E., Sakaguchi, S., Shadrach, K., Crabb, J. W., and Hollyfield, J. G. (2006) *Ret Degen Dis Adv Exp Med Bio* 572, 75-78).

[0221] In some embodiments, the Annexin IV is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury. In some embodiments, the Annexin IV is present on the surface of a cell of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) non-ischemic injury. In some embodiments, the Annexin IV is present on the surface of a cell of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage. In some embodiments, the Annexin IV is present on the surface of a cell of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ischemia-reperfusion injury. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein.

[0222] In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) non-ischemic injury. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage. In some embodiments, the Annexin IV is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ischemia-reperfusion injury. In some embodiments, the Annexin IV is produced by a nucleated cell (such as a mammalian cell). In some embodiments, the Annexin IV is recombinant protein.

[0223] In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of

undergoing) tissue injury but not on the surface of a cell that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) tissue injury. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) non-ischemic injury but not on the surface of a cell that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) non-ischemic injury. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage but not on the surface of a cell that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) oxidative damage. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell (and/or in a pathological structure) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ischemia-reperfusion injury but is not present on the surface of a cell that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) ischemia reperfusion injury.

[0224] In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) tissue injury. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) non-ischemic injury but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) non-ischemic injury. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological

structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) oxidative damage. In some embodiments, the epitope on Annexin IV for the antibody or fragment thereof is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ischemia-reperfusion injury but is not present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) ischemia reperfusion injury.

[0225] In some embodiments, the antibody or fragment thereof described herein (also referred to as the targeting moiety when provided in the context of a targeting construct) specifically binds to a phospholipid, which include, but is not limited to, phosphatidylethanolamine (PE), cardiolipin (CL), phosphatidylcholine (PC), and malondialdehyde (MDA). PE, CL, and PC are classes of phospholipids found in biological membranes. Phosphatidylcholine is more commonly found in the exoplasmic or outer leaflet of a cell membrane. It is thought to be transported between membranes within the cell by phosphatidylcholine transfer protein (PCTP). The phospholipid is composed of a choline head group and glycerophosphoric acid with a variety of fatty acids, one being a saturated fatty acid and one being an unsaturated fatty acid. PE consists of a combination of glycerol esterified with two fatty acids and phosphoric acid. Whereas the phosphate group is combined with choline in phosphatidylcholine, it is combined with the ethanolamine in PE. The two fatty acids may be the same, or different, and are usually in the 1,2 positions (though they can be in the 1,3 positions). Cardiolipin (IUPAC name "1,3-bis(sn-3'-phosphatidyl)-sn-glycerol") is an important component of the inner mitochondrial membrane, where it constitutes about 20% of the total lipid composition. Cardiolipin (CL) is a kind of diphosphatidylglycerol lipid, in which two phosphatidylglycerols connect with a glycerol backbone in the center to form a dimeric structure. In most animal tissues, cardiolipin contains 18-carbon fatty alkyl chains with 2 unsaturated bonds on each of them. It has been proposed that the (18:2)₄ acyl chain configuration is an important structural requirement for the high affinity of CL to inner membrane proteins in mammalian mitochondria. Phospholipid accumulation has been shown in eyes with age-related macular degeneration (Lommatzsch, et al. (2008) Graefes Arch Clin Exp Ophthalmol. 246(6):803-10).

[0226] Malondialdehyde (MDA) is generated from reactive oxygen species (ROS), and as such is often assayed in vivo as a bio-marker of oxidative stress. Reactive oxygen species degrade polyunsaturated lipids, forming malondialdehyde. This compound is a reactive aldehyde and is one of the many reactive electrophile species that cause toxic stress in cells and form covalent protein adducts referred to as advanced lipoxidation end-products (ALE). The production of this aldehyde is also used as a biomarker to measure the level of oxidative stress in an organism. MDA modifications have been shown in eyes with age-related macular degeneration and in the mouse laser-induced CNV model of wet AMD (Weissman et al. (2011) *Nature*. 478(7367):76-81).

[0227] In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell (or in a pathological structure, e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell (or in a pathological structure, e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ocular disease. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell (or in a pathological structure, e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage. In some embodiments, the phospholipid is neutral. In some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is oxidized.

[0228] In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to an ocular tissue undergoing (or is at risk of undergoing) tissue injury. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) of an individual that is in or adjacent to an ocular tissue undergoing (or is at risk of undergoing) ocular disease. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) of an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage. In some embodiments, the phospholipid is neutral. In

some embodiments, the phospholipid is positively charged. In some embodiments, the phospholipid (such as PE, CL, MDA, and/or PC) is oxidized.

[0229] In some embodiments, the epitope of phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) tissue injury but not on the surface of a cell or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) tissue injury. In some embodiments, the epitope of phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ocular disease but not on the surface of a cell or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) non-ocular disease. In some embodiments, the epitope on phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage but not on the surface of a cell or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) oxidative damage.

[0230] In some embodiments, the epitope of phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a ocular tissue undergoing (or is at risk of undergoing) tissue injury but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a ocular tissue not undergoing (or is not at risk of undergoing) tissue injury. In some embodiments, the epitope of phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) in an individual that is in or adjacent to a tissue undergoing (or is at risk of undergoing) ocular disease but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of

undergoing) non-ocular disease. In some embodiments, the epitope on phospholipid (such as PE, CL, MDA, and/or PC) to which the antibody or fragment thereof binds is present on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue undergoing (or is at risk of undergoing) oxidative damage but not on the surface of a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a tissue not undergoing (or is not at risk of undergoing) oxidative damage.

[0231] As described herein, a cell (and/or a pathological structure) that is in or adjacent to a particular tissue as described herein includes a cell (and/or a pathological structure, e.g., drusen) that is part of a tissue or organ, or adjacent to (near, directly next to, in the microenvironment of, bordering, flanking, adjoining) a tissue or organ, in which a certain event (such as non-ischemic injury or oxidative damage) is going to occur, is likely to occur, or is beginning to occur. As described herein, a cell, a basement membrane (e.g., Bruch's membrane), or in a pathological structure (e.g., drusen) that is in or adjacent to a particular tissue as described herein includes a cell that is part of a tissue or organ, or adjacent to (near, directly next to, in the microenvironment of, bordering, flanking, adjoining) a tissue or organ, in which a certain event (such as non-ischemic injury or oxidative damage) is going to occur, is likely to occur, or is beginning to occur. In the case of an adjacent cell, the cell is sufficiently within the microenvironment of the specific tissue or organ such that conditions of oxidative damage and/or inflammation affect the adjacent cell, as well as the specific tissue or organ. Such a cell may display signs of stress, including, but not limited to, the display of "stress proteins" (e.g., heat shock proteins and other proteins associated with a cellular stress response, including annexins) or other molecules on the cell surface (phospholipids, carbohydrate moieties), including the display of abnormal levels of proteins, modified proteins, or other molecules on the cell surface. Such a cell may be undergoing apoptosis or showing signs of apoptosis, such signs including morphological changes in the cell, chromatin condensation, changes in cellular signal transduction protein interactions, changes in intracellular calcium levels, externalization of phospholipids, cell detachment, loss of cell surface structures, etc.

[0232] As used herein, the term "selectively binds to" refers to the specific binding of one protein to another protein, to a lipid, or to a carbohydrate moiety (e.g., the

binding of an antibody, a fragment thereof, or binding partner to an antigen), wherein the level of binding, as measured by any standard assay (e.g., an immunoassay), is statistically significantly higher than the background control for the assay. For example, when performing an immunoassay, controls typically include a reaction well/tube that contain antibody or antigen binding fragment alone (i.e., in the absence of antigen), wherein an amount of reactivity (e.g., non-specific binding to the well) by the antibody or antigen binding fragment thereof in the absence of the antigen is considered to be background. Binding can be measured using a variety of methods standard in the art, including, but not limited to: Western blot, immunoblot, enzyme-linked immunosorbant assay (ELISA), radioimmunoassay (RIA), immunoprecipitation, surface plasmon resonance, chemiluminescence, fluorescent polarization, phosphorescence, immunohistochemical analysis, matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry, microcytometry, microarray, microscopy, fluorescence activated cell sorting (FACS), and flow cytometry.

[0233] According to the present invention, an "epitope" of a given protein or peptide or other molecule is generally defined, with regard to antibodies, as a part of or site on a larger molecule to which an antibody or antigen-binding fragment thereof will bind, and against which an antibody will be produced. The term epitope can be used interchangeably with the term "antigenic determinant", "antibody binding site", or "conserved binding surface" of a given protein or antigen. More specifically, an epitope can be defined by both the amino acid residues involved in antibody binding and also by their conformation in three-dimensional space (e.g., a conformational epitope or the conserved binding surface). An epitope can be included in peptides as small as about 4-6 amino acid residues, or can be included in larger segments of a protein, and need not be comprised of contiguous amino acid residues when referring to a three dimensional structure of an epitope, particularly with regard to an antibody-binding epitope. Antibody-binding epitopes are frequently conformational epitopes rather than a sequential epitope (i.e., linear epitope), or in other words, an epitope defined by amino acid residues arrayed in three dimensions on the surface of a protein or polypeptide to which an antibody binds. As mentioned above, the conformational epitope is not comprised of a contiguous sequence of amino acid residues, but instead, the residues are perhaps widely separated in the primary protein sequence, and are

brought together to form a binding surface by the way the protein folds in its native conformation in three dimensions.

[0234] Competition assays can be performed using standard techniques in the art (e.g., competitive ELISA or other binding assays). For example, competitive inhibitors can be detected and quantitated by their ability to inhibit the binding of an antigen to a known, labeled antibody (e.g., the mAb B4) or to sera or another composition that is known to contain antibodies against the particular antigen (e.g., sera known to contain natural antibodies against the antigen).

[0235] According to the present invention, antibodies are characterized in that they comprise immunoglobulin domains and as such, they are members of the immunoglobulin superfamily of proteins. Generally speaking, an antibody molecule comprises two types of chains. One type of chain is referred to as the heavy or H chain and the other is referred to as the light or L chain. The two chains are present in an equimolar ratio, with each antibody molecule typically having two H chains and two L chains. The two H chains are linked together by disulfide bonds and each H chain is linked to an L chain by a disulfide bond. There are only two types of L chains referred to as lambda (λ) and kappa (κ) chains. In contrast, there are five major H chain classes referred to as isotypes. The five classes include immunoglobulin M (IgM or μ), immunoglobulin D (IgD or δ), immunoglobulin G (IgG or γ), immunoglobulin A (IgA or α), and immunoglobulin E (IgE or ϵ). The distinctive characteristics between such isotypes are defined by the constant domain of the immunoglobulin and are discussed in detail below. Human immunoglobulin molecules comprise nine isotypes, IgM, IgD, IgE, four subclasses of IgG including IgG1 (γ 1), IgG2 (γ 2), IgG3 (γ 3) and IgG4 (γ 4), and two subclasses of IgA including IgA1 (α 1) and IgA2 (α 2). In humans, IgG subclass 3 and IgM are the most potent complement activators (classical complement system), while IgG subclass 1 and to an even lesser extent, 2, are moderate to low activators of the classical complement system. IgG4 subclass does not activate the complement system (classical or alternative). The only human immunoglobulin isotype known to activate the alternative complement system is IgA. In mice, the IgG subclasses are IgG1, IgG2a, IgG2b and IgG3. Murine IgG1 does not activate complement, while IgG2a, IgG2b and IgG3 are complement activators.

[0236] Each H or L chain of an immunoglobulin molecule comprises two regions referred to as L chain variable domains (VL domains) and L chain constant domains

(CL domains), and H chain variable domains (VH domains) and H chain constant domains (CH domains). A complete CH domain comprises three sub-domains (CH1, CH2, CH3) and a hinge region. Together, one H chain and one L chain can form an arm of an immunoglobulin molecule having an immunoglobulin variable region. A complete immunoglobulin molecule comprises two associated (e.g., di-sulfide linked) arms. Thus, each arm of a whole immunoglobulin comprises a VH+L region, and a CH+L region. As used herein, the term "variable region" or "V region" refers to a VH+L region (also known as an Fv fragment), a VL region or a VH region. Also as used herein, the term "constant region" or "C region" refers to a CH+L region, a CL region or a CH region.

[0237] The antigen specificity of an immunoglobulin molecule is conferred by the amino acid sequence of a variable, or V, region. As such, V regions of different immunoglobulin molecules can vary significantly depending upon their antigen specificity. Certain portions of a V region are more conserved than others and are referred to as framework regions (FR regions). In contrast, certain portions of a V region are highly variable and are designated hypervariable regions. When the VL and VH domains pair in an immunoglobulin molecule, the hypervariable regions from each domain associate and create hypervariable loops that form the antigen binding sites (antigen combining sites). Thus, the hypervariable loops determine the specificity of an immunoglobulin and are termed complementarity-determining regions (CDRs) because their surfaces are complementary to antigens.

[0238] Both an L chain and H chain V gene segment contain three regions of substantial amino acid sequence variability. Such regions are referred to as L chain CDR1, CDR2 and CDR3, and H chain CDR1, CDR2 and CDR3, respectively. The length of an L chain CDR1 can vary substantially between different VL regions. For example, the length of CDR1 can vary from about 7 amino acids to about 17 amino acids. In contrast, the lengths of L chain CDR2 and CDR3 typically do not vary between different VL regions. The length of an H chain CDR3 can vary substantially between different VH regions. For example, the length of CDR3 can vary from about 1 amino acid to about 20 amino acids. Each H and L chain CDR region is flanked by FR regions.

[0239] Limited digestion of an immunoglobulin with a protease may produce two fragments. An antigen binding fragment is referred to as an Fab, an Fab', or an F(ab')₂ fragment. A fragment lacking the ability to bind to antigen is referred to as an Fc

fragment. A Fab fragment comprises one arm of an immunoglobulin molecule containing a L chain (VL+CL domains) paired with the VH region and a portion of the CH region (CH1 domain). An Fab' fragment corresponds to an Fab fragment with part of the hinge region attached to the CH1 domain. An F(ab')₂ fragment corresponds to two Fab' fragments that are normally covalently linked to each other through a disulfide bond, typically in the hinge regions.

[0240] Isolated antibodies of the present invention can include serum containing such antibodies, or antibodies that have been purified to varying degrees. Whole antibodies of the present invention can be polyclonal or monoclonal. Alternatively, functional equivalents of whole antibodies, such as antigen binding fragments in which one or more antibody domains are truncated or absent (e.g., Fv, Fab, Fab', or F(ab')₂ fragments), as well as genetically-engineered antibodies or antigen binding fragments thereof, including single chain antibodies (e.g., scFv), humanized antibodies, antibodies that can bind to more than one epitope (e.g., bi-specific antibodies), or antibodies that can bind to one or more different antigens (e.g., bi- or multi-specific antibodies), may also be employed in the invention.

[0241] In some embodiments, the targeting moiety of the targeting constructs provided herein comprises an antibody. In some embodiments, the targeting moiety is a scFv. In some embodiments, the targeting moiety is a scFv comprising a (i) a light chain variable domain of SEQ ID NO:13; and/or (ii) heavy chain variable domain of SEQ ID NO:15. In some embodiments, the targeting moiety is a scFv comprising (i) a light chain variable domain of SEQ ID NO:14; and/or (ii) heavy chain variable domain of SEQ ID NO:16. In some embodiments, the targeting moiety is a scFv having the sequence of SEQ ID NO:17. In some embodiments, the targeting moiety is a scFv having the sequence of SEQ ID NO:18.

[0242] In some embodiments, the targeting moiety is a scFv comprising a (i) a light chain variable domain of SEQ ID NO:34; and/or (ii) heavy chain variable domain of SEQ ID NO:36. In some embodiments, the targeting moiety is a scFv comprising (i) a light chain variable domain of SEQ ID NO:35; and/or (ii) heavy chain variable domain of SEQ ID NO:36. In some embodiments, the targeting moiety is a scFv having the sequence of SEQ ID NO:37. In some embodiments, the targeting moiety is a scFv having the sequence of SEQ ID NO:38.

[0243] In one embodiment, targeting constructs of the present invention include humanized antibodies or a fragment thereof (such as a humanized scFv). A

humanized antibody or fragment thereof are molecules having an antigen binding site derived from an immunoglobulin from a non-human species, the remaining immunoglobulin-derived parts of the molecule being derived from a human immunoglobulin. The antigen binding site may comprise either complete variable regions fused onto human constant domains or only the complementarity determining regions (CDRs) grafted onto appropriate human framework regions in the variable domains. A humanized antibody or fragment thereof can be produced, for example, by modeling the antibody variable domains, and producing the antibodies using genetic engineering techniques, such as CDR grafting. A description various techniques for the production of humanized antibodies is found, for example, in Morrison et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-55; Whittle et al. (1987) Prot. Eng. 1:499-505; Co et al. (1990) J. Immunol. 148:1149-1154; Co et al. (1992) Proc. Natl. Acad. Sci. USA 88:2869-2873; Carter et al. (1992) Proc. Natl. Acad. Sci. 89:4285-4289; Routledge et al. (1991) Eur. J. Immunol. 21:2717-2725 and PCT Patent Publication Nos. WO 91/09967; WO 91/09968 and WO 92/113831.

[0244] In some embodiments, the antibody or fragment thereof does not activate complement activation. Methods of modifying antibodies or fragments thereof by reducing or eliminating their complement activation activities are known in the art (Tan et al. (1990) Proc Natl Acad Sci USA 87, 162-166).

[0245] In some embodiments, the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1, a sequence of SEQ ID NO:2, or a sequence of SEQ ID NO:3; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4, a sequence of SEQ ID NO:5, or a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7, a sequence of SEQ ID NO:8, or a sequence of SEQ ID NO:9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:10, a sequence of SEQ ID NO:11, or a sequence of SEQ ID NO:12.

[0246] In some embodiments, the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3. In some embodiments, the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence

of SEQ ID NO:8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9.

[0247] In some embodiments, the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:12.

[0248] In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:12.

[0249] In some embodiments, the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:1; (ii) a light chain CDR2 of SEQ ID NO:2; (iii) a light chain CDR3 of SEQ ID NO:3; (iv) heavy chain CDR1 of SEQ ID NO:4; (v) heavy chain CDR2 of SEQ ID NO:5; and (vi) heavy chain CDR3 of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof comprises: (i) a light chain CDR1 of SEQ ID NO:7; (ii) a light chain CDR2 of SEQ ID NO:8; (iii) a light chain CDR3 of SEQ ID NO:9; (iv) heavy chain CDR1 of SEQ ID NO:10; (v) heavy chain CDR2 of SEQ ID NO:11; and (vi) heavy chain CDR3 of SEQ ID NO:12.

[0250] In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:13. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:15. In

some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:14. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:16.

[0251] In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:13; and (ii) heavy chain variable domain of SEQ ID NO:15. In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:14; and (ii) heavy chain variable domain of SEQ ID NO:16.

[0252] In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:17. In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:18.

[0253] In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25, a sequence of SEQ ID NO:26, or a sequence of SEQ ID NO:27; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31, a sequence of SEQ ID NO:32, or a sequence of SEQ ID NO:33; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30.

[0254] In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27. In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33.

[0255] In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (ii) heavy chain variable domain comprising a sequence of SEQ

ID NO:29; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:30.

[0256] In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:32; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:33; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:30.

[0257] In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain CDR1 of SEQ ID NO:25; (ii) a light chain CDR2 of SEQ ID NO:26; (iii) a light chain CDR3 of SEQ ID NO:27; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30. In some embodiments, the antibody or fragment thereof specifically binds to a phospholipid and comprises: (i) a light chain CDR1 of SEQ ID NO:31; (ii) a light chain CDR2 of SEQ ID NO:32; (iii) a light chain CDR3 of SEQ ID NO:33; (iv) heavy chain CDR1 of SEQ ID NO:28; (v) heavy chain CDR2 of SEQ ID NO:29; and (vi) heavy chain CDR3 of SEQ ID NO:30.

[0258] In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:34. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:36. In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:35.

[0259] In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:34; and (ii) heavy chain variable domain of SEQ ID NO:36. In some embodiments, the antibody or fragment thereof

comprises: (i) a light chain variable domain of SEQ ID NO:35; and (ii) heavy chain variable domain of SEQ ID NO:36.

[0260] In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:37. In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:38.

[0261] In some embodiments, the antibody or fragment thereof does not activate complement activation. Methods of modifying antibodies or fragments thereof by reducing or eliminating their complement activation activities are known in the art (Tan et al. (1990) Proc Natl Acad Sci USA 87, 162-166).

[0262] In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1, a sequence of SEQ ID NO:2, or a sequence of SEQ ID NO:3; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4, a sequence of SEQ ID NO:5, or a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7, a sequence of SEQ ID NO:8, or a sequence of SEQ ID NO:9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:10, a sequence of SEQ ID NO:11, or a sequence of SEQ ID NO:12.

[0263] In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3. In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9.

[0264] In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID

NO:10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:12.

[0265] In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:4; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:5; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:8; (iii) a light chain variable domain comprising a sequence of SEQ ID NO:9; (iv) heavy chain variable domain comprising a sequence of SEQ ID NO:10; (v) heavy chain variable domain comprising a sequence of SEQ ID NO:11; and (vi) heavy chain variable domain comprising a sequence of SEQ ID NO:12.

[0266] In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain CDR1 of SEQ ID NO:1; (ii) a light chain CDR2 of SEQ ID NO:2; (iii) a light chain CDR3 of SEQ ID NO:3; (iv) heavy chain CDR1 of SEQ ID NO:4; (v) heavy chain CDR2 of SEQ ID NO:5; and (vi) heavy chain CDR3 of SEQ ID NO:6. In some embodiments, the antibody or fragment thereof specifically binds to Annexin IV and comprises: (i) a light chain CDR1 of SEQ ID NO:7; (ii) a light chain CDR2 of SEQ ID NO:8; (iii) a light chain CDR3 of SEQ ID NO:9; (iv) heavy chain CDR1 of SEQ ID NO:10; (v) heavy chain CDR2 of SEQ ID NO:11; and (vi) heavy chain CDR3 of SEQ ID NO:12.

[0267] In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:13. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:15. In some embodiments, the antibody or fragment thereof comprises a light chain variable domain of SEQ ID NO:14. In some embodiments, the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:16.

[0268] In some embodiments, the antibody or fragment thereof comprises: (i) a light chain variable domain of SEQ ID NO:13; and (ii) heavy chain variable domain of SEQ ID NO:15. In some embodiments, the antibody or fragment thereof

comprises: (i) a light chain variable domain of SEQ ID NO:14; and (ii) heavy chain variable domain of SEQ ID NO:16.

[0269] In some embodiments, the antibody or fragment thereof is an scFv. In some embodiments, the targeting moiety is an scFv comprising a (i) a light chain variable domain of SEQ ID NO: 34; and/or (ii) heavy chain variable domain of SEQ ID NO:36. In some embodiments, scFv comprises (i) a light chain variable domain of SEQ ID NO:35; and/or (ii) heavy chain variable domain of SEQ ID NO:36. In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:37. In some embodiments, the antibody or fragment is a scFv having the sequence of SEQ ID NO:38.

[0270] Therapeutic moieties and Complement modulators

[0271] In certain embodiments, the targeting construct comprises an active moiety that is a therapeutic moiety. In some embodiments, the therapeutic moiety is targeted or presented to the near space of the binding partner, e.g., Annexin IV, PE, PC, MDA, and/or CL, recognized by the targeting moiety.

[0272] In some embodiments the therapeutic moiety comprises an anti-VEGF drug, such as for treating macular edema, including, but not limited to, e.g., the monoclonal antibody bevacizumab (Avastin), derivatives of bevacizumab such as ranibizumab (Lucentis), orally-available small molecules that inhibit the tyrosine kinases stimulated by VEGF (e.g., lapatinib (Tykerb), sunitinib (Sutent), sorafenib (Nexavar), axitinib, and pazopanib), and aflibercept (EYLEA), a recombinant fusion protein consisting of portions of human VEGF receptors 1 and 2 extracellular domains fused to the Fc portion of human IgG1.

[0273] In certain embodiments, the therapeutic moiety in the targeting construct is an anti-viral agent, such as for the treatment of CMV retinitis, including, but not limited to, e.g., Ganciclovir, Foscarnet, Fomivirsen, Valganciclovir, and cidofovir. In certain embodiments, the therapeutic moiety in the targeting construct is a corticosteroid or an anti-TNFalpha agent, such as for the treatment of uveitis, including, but not limited to, e.g., prednisone, prednisolone, infliximab (Remicade), adalimumab (Humira), certolizumab pegol (Cimzia), and golimumab (Simponi), etanercept (Enbrel), xanthine derivatives (e.g., pentoxifylline) and Bupropion. In some embodiments, the therapeutic agent is an agent used for the treatment of glaucoma, including, but not limited to, e.g., a prostaglandins, a prostaglandin analog

(such as misoprostol, lantanprost, bimatoprost, or travoprost), a beta blocker (such as timolol, levobumamol, or betaxolol), a carbonic anhydrase inhibitor (such as dorzolamide (Trusopt), brinzolamide (Azopt), or acetazolamide (Diamox)), a miotic agent (such as pilocarpine), a cholinergic agent, and a neurotrophic factor to prevent retinal ganglion cell degeneration.

[0274] In certain embodiments, the therapeutic agent is an agent that is used in the treatment of diabetic retinopathy, including, but not limited to, e.g., an anti-VEGF drug or a Protein Kinase C inhibitor. In certain embodiments, the therapeutic agent is an agent that is used in the treatment of retinitis pigmentosa, e.g., ciliary neurotrophic factor (CNTF), the carbonic anhydrase inhibitor Acetazolamide, calcium channel blockers (such as Diltiazem), and immunosuppressive agents (if anti-retinal antibodies are present).

[0275] In certain embodiments, the therapeutic agent is an agent that is used in the treatment of proliferative vitreoretinopathy, including, but not limited to, e.g., minocyclin and Daunorubicin.

[0276] Methods of conjugating a targeting moiety, such as a B4 or C2 antibody or an antibody binding fragment thereof, to a drug, such a drug described herein, are known in the art and described in, e.g., McDonagh et al. (2006) *Prot Engineer Design Select* 19(7): 299-307; Hurwitz et al. (1975) *Cancer Res* 35:1175-1181; Garnett et al. (1983) *Int J Cancer* 31(5):661-670; Kovtun, et al. (2007) *Cancer Letters* 255(2): 232-40; Teicher et al. (2012) *N Engl J Med* 367(19):1847-8, and others.

[0277] In some embodiments, a therapeutic moiety is a complement modulator. An example of a complement modulator is a complement inhibitor. In some embodiments, such therapeutic moiety is a complement inhibitor. Accordingly, as used herein, the term “a therapeutic moiety” can encompass both a complement modulator and a complement inhibitor.

[0278] The constructs described herein in some embodiments comprise a complement modulator, such as a complement inhibitor.

[0279] As used herein, the term “complement inhibitor” refers to any compound, composition, or protein that reduces or eliminates complement activity. The reduction in complement activity may be incremental (e.g., a 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% reduction in activity) or complete. For example, in some embodiments, a complement inhibitor can inhibit complement activity by at least 10 (e.g., at least 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, or

95 or greater) % in a standard in vitro red blood cell hemolysis assay or an in vitro CH50eq assay. See, e.g., Kabat and Mayer (eds), "Experimental Immunochemistry, 2nd Edition," 135-240, Springfield, IL, CC Thomas (1961), pages 135-139, or a conventional variation of that assay such as the chicken erythrocyte hemolysis method as described in, e.g., Hillmen et al. (2004) N Engl J Med 350(6):552.

[0280] The CH50eq assay is a method for measuring the total classical complement activity in serum. This test is a lytic assay, which uses antibody-sensitized erythrocytes as the activator of the classical complement pathway and various dilutions of the test serum to determine the amount required to give 50% lysis (CH50). The percent hemolysis can be determined, for example, using a spectrophotometer. The CH50eq assay provides an indirect measure of terminal complement complex (TCC) formation, since the TCC themselves are directly responsible for the hemolysis that is measured.

[0281] The assay is well known and commonly practiced by those of skill in the art. Briefly, to activate the classical complement pathway, undiluted serum samples (e.g., human serum samples) are added to microassay wells containing the antibody-sensitized erythrocytes to thereby generate TCC. Next, the activated sera are diluted in microassay wells, which are coated with a capture reagent (e.g., an antibody that binds to one or more components of the TCC). The TCC present in the activated samples bind to the monoclonal antibodies coating the surface of the microassay wells. The wells are washed and, to each well, is added a detection reagent that is detectably labeled and recognizes the bound TCC. The detectable label can be, e.g., a fluorescent label or an enzymatic label. The assay results are expressed in CH50 unit equivalents per milliliter (CH50 U Eq/mL).

[0282] Additional methods for detecting and/or measuring complement activity in vitro are set forth and exemplified in the working examples.

[0283] The complement inhibitor described herein in some embodiments is a specific inhibitor of the lectin pathway. In some embodiments, the complement inhibitor is a specific inhibitor of the alternative pathway. In some embodiments, the complement inhibitor is a specific inhibitor of the classical pathway.

[0284] In some embodiments, the complement inhibitor is a soluble or membrane-bound protein such as, for example, membrane cofactor protein (MCP), decay accelerating factor (DAF/CD55), CD59, mouse complement receptor 1-related gene/protein y (Crry), human complement receptor 1 (CRI) or factor H, or Factor I, or

an antibody specific for a component of a complement pathway such as, for example, eculizumab (an anti-CS antibody marketed under the trade name Soliris®), pexelizumab (the antigen-binding fragment of eculizumab), an anti-factor B antibody (such as the monoclonal antibody 1379 produced by ATCC Deposit No. PTA-6230), an anti-properdin antibody, an anti-factor D antibody, an anti-MASP antibody, an anti-MBL-antibody, and the like (see below). Alternatively, a complement inhibitor may be a small molecule or a linear or cyclic peptide such as, for example, compstatin, N-acetylaspartylglutamic acid (NAAGA), and the like. In some embodiments, the complement inhibitor is selected from the group consisting of: an anti-C5 antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-factor D antibody, and an anti-properdin antibody, a human membrane cofactor protein (MCP), a human decay accelerating factor (DAF), a mouse decay accelerating factor (DAF), a human CD59, a mouse CD59, a mouse CD59 isoform B, a mouse Crry, a human CR1, a Factor I, a human factor H, a mouse factor H, and a biologically active fragment of any the preceding.

[0285] As used herein, the term “membrane cofactor protein,” “MCP,” or “CD46” refers to a widely distributed C3b/C4b-binding cell surface glycoprotein which inhibits complement activation on host cells and serves as a cofactor for the factor I-mediated cleavage of C3b and C4b, including homologs thereof. T.J. Oglesby et al., J. Exp. Med. (1992) 175:1547-1551. MCP belongs to a family known as the regulators of complement activation (“RCA”). Family members share certain structural features, comprising varying numbers of short consensus repeat (SCR) domains, which are typically between 60 and 70 amino acids in length. Beginning at its amino-terminus, MCP comprises four SCRs, a serine/threonine/proline-enriched region, an area of undefined function, a transmembrane hydrophobic domain, a cytoplasmic anchor and a cytoplasmic tail. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that human MCP or biologically active fragments thereof encompass all species and strain variations.

[0286] SEQ ID NO:44 represents the full-length human MCP amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P15529). Amino acids 1-34 correspond to the signal peptide, amino acids 35-343 correspond to the extracellular domain, amino acids 344-366 correspond to the transmembrane domain, and amino acids 367-392 correspond to the cytoplasmic domain. In the extracellular domain,

amino acids 35-96 correspond to SCR 1, amino acids 97-159 correspond to SCR 2, amino acids 160-225 correspond to SCR 3, amino acids 226-285 correspond to SCR 4, and amino acids 302-326 correspond to the serine/threonine-rich domain. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that MCP or biologically active fragments thereof encompass all species and strain variations. As used herein, the term “biologically active” fragment of MCP refers to any soluble fragment lacking both the cytoplasmic domain and the transmembrane domain, including fragments comprising, consisting essentially of or consisting of 1, 2, 3, or 4 SCR domains, with or without the serine/threonine-rich domain, having some or all of the complement inhibitory activity of the full-length human MCP protein. In some embodiments, the complement inhibitor portion comprises full-length human MCP (amino acids 35-392 of SEQ ID NO:44), the extracellular domain of human MCP (amino acids 35-343 of SEQ ID NO:44), or SCRs 1-4 of human MCP (amino acids 35-285 of SEQ ID NO:44).

[0287] Decay accelerating factor, also referred to as CD55 (DAF/CD55) (SEQ ID NO:45 and SEQ ID NO:46), is an ~70 kiloDalton (kDa) membrane-bound glycoprotein which inhibits complement activation on host cells. Like several other complement regulatory proteins, DAF comprises several approximately 60 amino acid repeating motifs termed short consensus repeats (SCR).

[0288] As used herein, the term “decay accelerating factor,” “DAF,” or “CD55” refers to a seventy kilodalton (“kDa”) membrane glycoprotein comprising four short consensus repeat (SCR) domains followed by a heavily O-glycosylated serine/threonine-rich domain at the C-terminus that elevates the molecule from the membrane surface, followed by a glycosylphosphatidylinositol (“GPI”) anchor. DAF protects the cell surface from complement activation by dissociating membrane-bound C3 convertases that are required to cleave complement protein C3 and to amplify the complement cascade. DAF prevents assembly or accelerates decay of both the C3- and C5-convertases of the alternative and classical complement pathways.

[0289] SEQ ID NO:45 represents the full-length human DAF amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P08173); SEQ ID NO:46 represents the full-length mouse DAF amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. Q61475). In the human DAF sequence, amino acids 1-34 correspond

to the signal peptide, amino acids 35-353 appear in the mature protein, and amino acids 354-381 are removed from the polypeptide after translation. Within the mature protein, amino acids 35-96 correspond to SCR 1, amino acids 96- 160 correspond to SCR 2, amino acids 161-222 correspond to SCR 3, amino acids 223-285 correspond to SCR 4, and amino acids 287-353 correspond to the O-glycosylated serine/threonine-rich domain. The GPI anchor is attached to human DAF at a serine at position 353. In the mouse DAF sequence, amino acids 1-34 correspond to the signal peptide, amino acids 35-362 appear in the mature protein, and amino acids 363-390 are removed from the polypeptide after translation. Within the mature protein, amino acids 35-96 correspond to SCR 1, amino acids 97-160 correspond to SCR 2, amino acids 161-222 correspond to SCR 3, amino acids 223-286 correspond to SCR 4, and amino acids 288-362 correspond to the O-glycosylated serine/threonine-rich domain. The GPI anchor is attached to mouse DAF at a serine at position 362. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that DAF or biologically active fragments thereof encompass all species and strain variations. As used herein, the term “biologically active” fragment of DAF refers to any fragment of DAF lacking a GPI anchor and/or the amino acid to which it is attached (i.e., Ser-353), including any fragments of the full-length DAF protein comprising, consisting essentially of or consisting of 1, 2, 3, or 4 SCR domains, with or without the O-glycosylated serine/threonine-rich domain, having some or all the complement inhibitory activity of the full-length DAF protein.

[0290] As used herein, the term “CD59” refers to a membrane-bound 128 amino acid glycoprotein that potently inhibits the membrane attack complex (MAC) of complement. CD59 acts by binding to the C8 and/or C9 components of the MAC during assembly, ultimately preventing incorporation of the multiple copies of C9 required for complete formation of the osmolytic pore at the heart of the MAC. CD59 is both N- and O-glycosylated. The N-glycosylation comprises primarily bi- or tri-antennary structures with and without lactosamine and outer arm fucose residues, with variable sialylation present at some sites. Like DAF, CD59 is anchored in the cell membrane by a glycosylphosphatidylinositol (“GPI”) anchor, which is attached to an asparagine at amino acid 102. Soluble forms of CD59 (sCD59) have been produced, but they generally have low functional activity in vitro, particularly in the presence of serum, suggesting that unmodified sCD59 has little or no therapeutic efficacy. See, e.g., S. Meri et al., “Structural composition and functional characterization of soluble

CD59: heterogeneity of the oligosaccharide and glycosphosphoinositol (GPI) anchor revealed by laser-desorption mass spectrometric analysis,” *Biochem. J.* 316:923-935 (1996).

[0291] SEQ ID NO:47 represents the full-length human CD59 amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P13987); SEQ ID NO:48 represents the full-length mouse CD59 sequence, isoform A (see, e.g., UniProtKB/Swiss-Prot. Accession No. 055186); SEQ ID NO:49 represents the full-length mouse CD59 sequence, isoform B (see, e.g., UniProtKB/SwissProt. Accession No. P58019). In the human CD59 sequence, amino acids 1-25 of SEQ ID NO:47 correspond to the leader peptide, amino acids 26-102 of SEQ ID NO:47 correspond to the mature protein, and amino acids 103-128 of SEQ ID NO:47 are removed after translation. The GPI anchor is attached to CD59 at an asparagine at position 102 of SEQ ID NO:47. In isoform A of the mouse CD59 sequence, amino acids 1-23 of SEQ ID NO:48 correspond to the leader peptide, amino acids 24-96 of SEQ ID NO: 48 correspond to the mature protein, and amino acids 97-123 of SEQ ID NO:48 are removed after translation. The GPI anchor is attached to CD59 at a serine at position 96 of SEQ ID NO: 48. In isoform B of the mouse CD59 sequence, amino acids 1-23 of SEQ ID NO: 49 correspond to the leader peptide, amino acids 24-104 of SEQ ID NO: 49 correspond to the mature protein, and amino acids 105-129 of SEQ ID NO:49 are removed after translation. The GPI anchor is attached to CD59 at an asparagine at position 104 of SEQ ID NO:49. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that CD59 or biologically active fragments thereof encompass all species and strain variations.

[0292] As used herein, the term “biologically active” fragment of human CD59 refers to any fragment of human CD59 lacking a GPI anchor and/or the amino acid to which it is attached (i.e., Asn-102), including any fragments of the full-length human CD59 protein having some or all the complement inhibitory activity of the full-length CD59 protein; and the term “biologically active” fragment of mouse CD59 refers to any fragment of mouse CD59 isoform A or isoform B lacking a GPI anchor and/or the amino acid to which it is attached (i.e., Ser-96 of isoform A, or Asp-104 of isoform B), including any fragments of either full-length mouse CD59 protein isoform having some or all the complement inhibitory activity of the full-length CD59 protein.

[0293] As used herein, the term “mouse complement receptor 1-related gene/protein y” or “Crry” refers to a membrane-bound mouse glycoprotein that regulates

complement activation, including homologs thereof. Crry regulates complement activation by serving as a cofactor for complement factor I, a serine protease which cleaves C3b and C4b deposited on host tissue. Crry also acts as a decay-accelerating factor, preventing the formation of C4b2a and C3bBb, the amplification convertases of the complement cascade.

[0294] SEQ ID NO:50 represents the full-length mouse Crry protein amino acid sequence. Amino acids 1-40 correspond to the leader peptide, amino acids 41-483 of SEQ ID NO:50 correspond to the mature protein, comprising amino acids 41-405 of SEQ ID NO:50, corresponding to the extracellular domain, amino acids 406-426 of SEQ ID NO:50, corresponding to the transmembrane domain, and amino acids 427-483 of SEQ ID NO:50, corresponding to the cytoplasmic domain. In the extracellular domain, amino acids 83-143 of SEQ ID NO:50 correspond to SCR 1, amino acids 144-205 of SEQ ID NO:50 correspond to SCR 2, amino acids 206-276 of SEQ ID NO:50 correspond to SCR 3, amino acids 277-338 of SEQ ID NO:50 correspond to SCR 4, and amino acids 339-400 of SEQ ID NO:50 correspond to SCR 5. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that mouse Crry protein or biologically active fragments thereof encompasses all species and strain variations. As used herein, the term “biologically active” fragment of mouse Crry protein refers to any soluble fragment of mouse Crry lacking the transmembrane domain and the cytoplasmic domain, including fragments comprising, consisting essentially of or consisting of 1, 2, 3, 4, or 5 SCR domains, including any fragments of the full-length mouse Crry protein having some or all the complement inhibitory activity of the full-length Crry protein.

[0295] As used herein, the term “complement receptor 1,” “CR1,” or “CD35” refers to a human gene encoding a protein of 2039 amino acids, with a predicted molecular weight of 220 kilodaltons (“kDa”), including homologs thereof. The gene is expressed principally on erythrocytes, monocytes, neutrophils, and B cells, but is also present on some T lymphocytes, mast cells, and glomerular podocytes. CR1 protein is typically expressed at between 100 and 1000 copies per cell. CR1 is the main system for processing and clearance of complement-opsonized immune complexes. CR1 negatively regulates the complement cascade, mediates immune adherence and phagocytosis, and inhibits both the classic and alternative complement pathways. The full-length CR1 protein comprises a 42 amino acid signal peptide, an

extracellular domain of 1930 amino acids, a 25 amino acid transmembrane domain, and a 43 amino acid C-terminal cytoplasmic domain. The extracellular domain of CR1 has 25 potential N-glycosylation signal sequences, and comprises 30 short consensus (“SCR”) domains, also known as complement control protein (CCP) repeats, or sushi domains, each 60 to 70 amino acids long. The sequence homology between SCRs ranges between 60-99 percent. The 30 SCR domains are further grouped into four longer regions termed long homologous repeats (“LHRs”), each encoding approximately 45 kDa segments of the CR1 protein, designated LHR-A, -B, -C, and -D. The first three comprise seven SCR domains each, while LHR-D comprises 9 SCR domains. The active sites on the extracellular domain of CR1 protein include a C4b-binding site with lower affinity for C3b in SCRs 1-4 comprising amino acids 42-295, a C3b-binding site with lower affinity for C4b in SCRs 8-11 comprising amino acids 490-745, a C3b-binding site with lower affinity for C4b in SCRs 15-18 comprising amino acids 940-1196, and a C1q-binding site in SCRs 22-28 comprising amino acids 1394-1842.

[0296] SEQ ID NO:51 represents the full-length human CR1 amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P17927). Amino acids 1-41 correspond to the signal peptide, amino acids 42-2039 correspond to the mature protein, comprising amino acids 42-1971, corresponding to the extracellular domain, amino acids 1972-1996, corresponding to the transmembrane domain, and amino acids 1997-2039, corresponding to the cytoplasmic domain. In the extracellular domain, amino acids 42-101 correspond to SCR 1, 102-163 correspond to SCR2, amino acids 164-234 correspond to SCR3, amino acids 236-295 correspond to SCR4, amino acids 295-355 correspond to SCR5, amino acids 356-418 correspond to SCR6, amino acids 419-489 correspond to SCR7, amino acids 491-551 correspond to SCR8, amino acids 552- 613 correspond to SCR9, amino acids 614-684 correspond to SCR10, amino acids 686-745 correspond to SCR11, amino acids 745-805 correspond to SCR12, amino acids 806-868 correspond to SCR13, amino acids 869-939 correspond to SCR14, amino acids 941-1001 correspond to SCR15, amino acids 1002-1063 correspond to SCR16, amino acids 1064-1134 correspond to SCR17, amino acids 1136-1195 correspond to SCR18, amino acids 1195-1255 correspond to SCR 19, amino acids 1256-1318 correspond to SCR 20, amino acids 1319-1389 correspond to SCR 21, amino acids 1394-1454 correspond to SCR 22, amino acids 1455-1516 correspond to SCR 23, amino acids 1517-1587 correspond to SCR 24,

amino acids 1589-1648 correspond to SCR 25, amino acids 1648-1708 correspond to SCR 26, amino acids 1709-1771 correspond to SCR 27, amino acids 1772-1842 correspond to SCR 28, amino acids 1846-1906 correspond to SCR 29, amino acids 1907-1967 correspond to SCR 30. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that CR1 protein or biologically active fragments thereof encompass all species and strain variations. As used herein, the term “biologically active” fragment of CR1 protein refers to any soluble fragment of CR1 lacking the transmembrane domain and the cytoplasmic domain, including fragments comprising, consisting essentially of or consisting of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 SCR domains, including any fragments of the full-length CR1 protein having some or all the complement inhibitory activity of the full-length CR1 protein.

[0297] As used herein, the term “complement factor H,” “factor H,” or “FH” refers to complement factor H, a single polypeptide chain plasma glycoprotein, including homologs thereof. The protein is composed of 20 conserved short consensus repeat (SCR) domains of approximately 60 amino acids, arranged in a continuous fashion like a string of beads, separated by short linker sequences of 2-6 amino acids each. Factor H binds to C3b, accelerates the decay of the alternative pathway C3-convertase (C3bBb), and acts as a cofactor for the proteolytic inactivation of C3b. In the presence of factor H, proteolysis by factor I results in the cleavage and inactivation of C3b. Factor H has at least three distinct binding domains for C3b, which are located within SCRs 1-4, SCRs 5-8, and SCRs 19-20. Each domain binds to a distinct region within the C3b protein: the N-terminal sites bind to native C3b; the second site, located in the middle region of factor H, binds to the C3c fragment and the site located within SCR19 and 20 binds to the C3d region. In addition, factor H also contains binding sites for heparin, which are located within SCR 7, SCRs 5-12, and SCR 20 of factor H and overlap with those of the C3b binding sites. Structural and functional analyses have shown that the domains for the complement inhibitory activity of factor H are located within the first four N-terminal SCR domains.

[0298] SEQ ID NO:52 represents the full-length human factor H amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P08603); SEQ ID NO:53 represents the full-length mouse factor H amino acid sequence (see, e.g., UniProtKB/Swiss-Prot. Accession No. P06909). In the human factor H sequence,

amino acids 1-18 of SEQ ID NO:52 correspond to the signal peptide, and amino acids 19-1231 of SEQ ID NO:52 correspond to the mature protein. Within that protein, amino acids 21-80 of SEQ ID NO:52 correspond to SCR 1, amino acids 85-141 of SEQ ID NO:52 correspond to SCR 2, amino acids 146-205 of SEQ ID NO:52 correspond to SCR 3, amino acids 210-262 of SEQ ID NO:52 correspond to SCR 4, and amino acids 267-320 of SEQ ID NO:52 correspond to SCR 5. In the mouse factor H sequence, amino acids 1-18 of SEQ ID NO:53 correspond to the signal peptide, and amino acids 19-1234 of SEQ ID NO:53 correspond to the mature protein. Within that protein, amino acids 19-82 of SEQ ID NO:53 correspond to SCR 1, amino acids 83-143 of SEQ ID NO:53 correspond to SCR 2, amino acids 144-207 of SEQ ID NO:53 correspond to SCR 3, amino acids 208-264 of SEQ ID NO:53 correspond to SCR 4, and amino acids 265-322 of SEQ ID NO:53 correspond to SCR 5. It is understood that species and strain variations exist for the disclosed peptides, polypeptides, and proteins, and that factor H or biologically active fragments thereof encompass all species and strain variations.

[0299] As used herein, the term “biologically active” fragment of factor H refers to any portion of a factor H protein having some or all the complement inhibitory activity of the full-length factor H protein, and includes, but is not limited to, factor H fragments comprising SCRs 1-4, SCRs 1-5, SCRs 1-8, SCRs 1-18, SCRs 19-20, or any homolog of a naturally-occurring factor H or fragment thereof, as described in detail below. In some embodiments, the biologically active fragment of factor H has one or more of the following properties: (1) binding to C-reactive protein (CRP), (2) binding to C3b, (3) binding to heparin, (4) binding to sialic acid, (5) binding to endothelial cell surfaces, (6) binding to cellular integrin receptor, (7) binding to pathogens, (8) C3b co-factor activity, (9) C3b decay-acceleration activity, and (10) inhibiting the alternative complement pathway.

[0300] Thus, in some embodiments, the therapeutic moiety of a targeting construct described herein comprises a complement inhibitor or biologically active fragment thereof. In some embodiments, the complement inhibitor is selected from the group consisting of human MCP, human DAF, mouse DAF, human CD59, mouse CD59 isoform A, mouse CD59 isoform B, mouse Crry protein, human CR1, human factor H, or mouse factor H, a Factor I, or a biologically active fragment thereof.

[0301] In some embodiments, the complement inhibitor portion of the targeting construct comprises full-length human MCP (SEQ ID NO:44). In some

embodiments, the complement inhibitor portion of the targeting construct comprises a biologically active fragment of human MCP (SEQ ID NO:44). In some embodiments, the biologically active fragment of human MCP is selected from the group consisting of SCRs 1-4 (amino acids 35-285 of SEQ ID NO:44), SCRs 1-4 plus the serine/threonine-rich domain (amino acids 35-326 of SEQ ID NO:44), and the extracellular domain of MCP (amino acids 35-343 of SEQ ID NO:44).'

[0302] In some embodiments, the complement inhibitor portion of the targeting construct comprises full-length human DAF. In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of human DAF (SEQ ID NO:45). In some embodiments, the biologically active fragment of human DAF is selected from the group consisting of SCRs 1-4 (amino acids 25-285 of SEQ ID NO:45) and SCRs 1-4 plus the O- glycosylated serine/threonine-rich domain (amino acids 25-353 of SEQ ID NO:45). In some embodiments, the complement inhibitor portion of the construct comprises full-length mouse DAF (SEQ ID NO:46). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of mouse DAF. In some embodiments, the biologically active fragment of mouse DAF is selected from the group consisting of SCRs 1-4 (amino acids 35-286 of SEQ ID NO:46) and SCRs 1-4 plus the O- glycosylated serine/threonine-rich domain (amino acids 35-362 of SEQ ID NO:46).

[0303] In some embodiments, the complement inhibitor portion of the construct comprises full-length human CD59 (SEQ ID NO:47). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of human CD59 (SEQ ID NO:47). In some embodiments, the biologically active fragment of human CD59 comprises the extracellular domain of human CD59 lacking its GPI anchor (amino acids 26-101 of SEQ ID NO:47). In some embodiments, the complement inhibitor portion of the construct comprises full-length mouse CD59, isoform A (SEQ ID NO:48). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of mouse CD59, isoform A (SEQ ID NO:48). In some embodiments, the biologically active fragment of mouse CD59, isoform A comprises the extracellular domain of mouse CD59, isoform A lacking its GPI anchor (amino acids 24-95 of SEQ ID NO:48). In some embodiments, the complement inhibitor portion of the construct comprises full-length mouse CD59, isoform B (SEQ ID NO:49). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active

fragment of mouse CD59, isoform B (SEQ ID NO:49). In some embodiments, the biologically active fragment of mouse CD59, isoform B comprises the extracellular domain of mouse CD59, isoform B lacking its GPI anchor (amino acids 24-103 of SEQ ID NO:49).

[0304] In some embodiments, the complement inhibitor portion of the construct comprises full-length mouse Crry protein (SEQ ID NO:50). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of mouse Crry protein (SEQ ID NO:50). In some embodiments, the biologically active fragment of mouse Crry protein is selected from the group consisting of SCRs 1-5 (amino acids 41-400 of SEQ ID NO:50) and the extracellular domain of mouse Crry protein (amino acids 41-405 of SEQ ID NO:50).

[0305] In some embodiments, the complement inhibitor portion of the construct comprises full-length human CR1 protein (SEQ ID NO:51). In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of human CR1 protein (SEQ ID NO:51). In some embodiments, the biologically active fragment of human CR1 protein is selected from the group consisting of SCRs 1-4 (amino acids 42-295 of SEQ ID NO:51), SCRs 1-10 (amino acids 42-684 of SEQ ID NO:51), SCRs 8-11 (amino acids 490-745 of SEQ ID NO:51), SCRs 15-18 (amino acids 940-1196 of SEQ ID NO:51), and SCRs 22-28 (amino acids 1394-1842 of SEQ ID NO:51).

[0306] In some embodiments, the complement inhibitor portion of the construct comprises full-length human (SEQ ID NO:52) or mouse (SEQ ID NO:53) factor H. In some embodiments, the complement inhibitor portion of the construct comprises a biologically active fragment of human (SEQ ID NO:52) or mouse (SEQ ID NO:53) factor H. In some embodiments, the biologically active fragment of human factor H (SEQ ID NO:52) is selected from the group consisting of SCRs 1-4 (amino acids 21-262 of SEQ ID NO:52), SCRs 1-5 of factor H (amino acids 21-320 of SEQ ID NO:52), SCRs 1-8 of factor H (amino acids 21-507 of SEQ ID NO:52), and SCRs 1-18 of factor H (amino acids 21-1104 of SEQ ID NO:52). In some embodiments, the biologically active fragment of mouse factor H (SEQ ID NO:53) is selected from the group consisting of SCRs 1-4 (amino acids 19-264 of SEQ ID NO:53), SCRs 1-5 of factor H (amino acids 19-322 of SEQ ID NO:53), SCRs 1-8 of factor H (amino acids 19-507 of SEQ ID NO:53), and SCRs 1-18 of factor H (amino acids 19-1109 of SEQ ID NO:53). In some embodiments, the biologically active fragment of human (SEQ

ID NO:52) or mouse (SEQ ID NO:53) factor H comprises (and in some embodiments consists of or consists essentially of) at least the first four N-terminal SCR domains of factor H, including for example, at least any of the first 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, or more N-terminal SCR domains of factor H.

[0307] In some embodiments, the complement inhibitor portion of the targeting construct is a homolog of any of the complement inhibitors described herein or a biologically active fragment thereof. Homologs of the complement inhibitors (or biologically active fragments thereof) include proteins which differ from a naturally occurring complement inhibitor (or biologically-active fragment thereof) in that at least one or a few, but not limited to one or a few, amino acids have been deleted (e.g., a truncated version of the protein, such as a peptide or fragment), inserted, inverted, substituted and/or derivatized (e.g., by glycosylation, phosphorylation, acetylation, myristoylation, prenylation, palmitation, amidation and/or addition glycosylphosphatidyl inositol). For example, homologue of a complement inhibitor may have an amino acid sequence that is at least about 70% identical to the amino acid sequence of a naturally complement inhibitor (e.g., SEQ ID NOs:44-53), for example at least about any of 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identical to the amino acid sequence of a naturally occurring complement inhibitor (e.g., SEQ ID NOs:44-53). Amino acid sequence identity can be determined in various ways, for example, using publicly available computer software such as BLAST, BLAST-2, ALIGN or MEGALIGN™ (DNAST AR) software. One skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared.

[0308] In certain embodiments, a homologue of complement inhibitor (or a biologically active fragment thereof) retains all the alternative complement pathway inhibitory activity of the complement inhibitor (or a biologically active fragment thereof) from which it is derived. In certain embodiments, the homologue of a complement inhibitor (or a biologically-active fragment thereof) retains at least about 50%, for example, at least about any of 60%, 70%, 80%, 90%, or 95% of the complement inhibition activity the complement inhibitor (or a biologically-active fragment thereof) from which is derived.

[0309] In some embodiments, the complement inhibitor is an antibody (or an antigen binding fragment thereof) that binds to a complement component, e.g., a complement component selected from the group consisting of C1, C1q, C1s, C2, C2a, C3, C3a, C3b, C4, C4b, C5, C5a, C5b, C6, C7, C8, and C9. The complement polypeptides to which the antibodies or antigen binding fragments thereof bind can be, in some embodiments, human polypeptides, e.g., human C1, C1q, C1s, C2, C2a, C3, C3a, C3b, C4, C4b, C5, C5a, C5b, C6, C7, C8, C9, factor B, factor D, or properdin polypeptides. The amino acid sequences for the foregoing complement proteins are well-known in the art as are methods for preparing the proteins or fragments thereof for use in preparing an antibody (or antigen-binding fragment thereof) specific for one or more of the complement proteins. Suitable methods are also described and exemplified herein.

[0310] Exemplary anti-complement protein antibodies, which are suitable for incorporation into the targeting constructs described herein and for subsequent use in any of the methods described herein, are also well known in the art. For example, antibodies that bind to complement component C5 and inhibit the cleavage of C5 into fragments C5a and C5b include, e.g., eculizumab (Soliris®; Alexion Pharmaceuticals, Inc., Cheshire, CT) and pexelizumab (Alexion Pharmaceuticals, Inc., Cheshire, CT). See, e.g., Kaplan (2002) *Curr Opin Investig Drugs* 3(7):1017-23; Hill (2005) *Clin Adv Hematol Oncol* 3(11):849-50; Rother et al. (2007) *Nature Biotechnol* 25(11):1256-1488; Whiss (2002) *Curr Opin Investig Drugs* 3(6):870-7; Patel et al. (2005) *Drugs Today (Barc)* 41(3):165-70; and Thomas et al. (1996) *Mol Immunol* 33(17- 18):1389-401.

[0311] In some embodiments, the anti-C5 antibody can bind to an epitope in the alpha chain of the human complement component C5 protein. Antibodies that bind to the alpha chain of C5 are described in, for example, PCT application publication no. WO 2010/136311 and U.S. patent no. 6,355,245.

[0312] In some embodiments, the anti-C5 antibody can bind to an epitope in the beta chain of the human complement component C5 protein. Antibodies that bind to the C5 beta chain are described in, e.g., Moongkarndi et al. (1982) *Immunobiol* 162:397; Moongkarndi et al. (1983) *Immunobiol* 165:323; and Mollnes et al. (1988) *Scand J Immunol* 28:307-312.

[0313] Additional anti-C5 antibodies, and antigen-binding fragments thereof, suitable for use in the targeting constructs described herein are described in, e.g., PCT

application publication no. WO 2010/015608, the disclosure of which is incorporated herein by reference in its entirety.

[0314] Antibodies that bind to C3b and, for example, inhibit the C3b convertase are also well known in the art. For example, PCT application publication nos. WO 2010/136311, WO2009/056631, and WO 2008/154251, the disclosures of each of which are incorporated herein by reference in their entirety. Antagonistic anti-C6 antibodies and anti-C7 antibodies have been described in, e.g., Brauer et al. (1996) *Transplantation* 61(4):S88-S94 and U.S. patent no. 5,679,345.

[0315] In some embodiments, the complement inhibitor is an anti-factor B antibody (such as the monoclonal antibody 1379 produced by ATCC Deposit No. PTA-6230). Anti-factor B antibodies are also described in, e.g., Ueda et al. (1987) *J Immunol* 138(4):1143-9; Tanhehco et al. (1999) *Transplant Proc* 31(5):2168-71; U.S. patent application publication nos. 20050260198 and 2008029911; and PCT publication no. WO 09/029669.

[0316] In some embodiments, the complement inhibitor is an anti-factor D antibody, e.g., an antibody described in Pascual et al. (1990) *J Immunol Methods* 127:263-269; Sahu et al. (1993) *Mol Immunol* 30(7):679-684; Pascual et al. (1993) *Eur J Immunol* 23:1389-1392; Niemann et al. (1984) *J Immunol* 132(2):809-815; U.S. patent no. 7,439,331; or U.S. patent application publication no. 20080118506.

[0317] In some embodiments, the complement inhibitor is an anti-properdin antibody. Suitable anti-properdin antibodies are also well-known in the art and include, e.g., U.S. patent application publication nos. 20110014614 and PCT application publication no. W02009110918.

[0318] In some embodiments, the complement inhibitor portion is an anti-MBL antibody. Mannose-binding mannan-binding lectin (MBL), a plasma protein, forms a complex with proteins known as MBL-associated serine proteases (MASPs). MBL binds to several monosaccharides that are uncharacteristic of mammalian proteins, e.g., mannose, N-acetylglucosamine, N-acetylmannoseamine, L-fucose and glucose, whereas sialic acid and galactose are not bound. When the MBL-MASP complex binds to microorganisms, the proenzymic forms of the serine proteases are activated and mediate the activation of complement components C4 and C2, thereby generating the C3 convertase C4b2b and leading to opsonization by the deposition of C4b and C3b fragments. MASP-2 has been shown to cleave C4 and C2, while MASP-1 may be responsible for direct cleavage of C3. The functions of MASP-3 and MASP-19 are

less well understood. Studies have shown a clear link between low levels of MBL and opsonic deficiency, as well as clinical manifestations such as severe diarrhea, chronic hepatitis and HIV infection, and autoimmune disease. See, Petersen et al., *J. Immunological Methods*, 257:107-16 (2001); Petersen et al., *Molecular Immunology*, 38:133-49 (2001). Anti-mannan-binding lectin antibodies are known in the art (see, e.g., Pradhan et al. (2012) *Rheumatol. Int.* epubliished September, 2012) and commercially available (AbCam).

[0319] In some embodiments, the complement inhibitor portion is an anti-MASP antibody. The mannan-binding lectin-associated serine proteases (MASPs) are a family of at least three proteins (mannan-binding lectin-associated serine protease-1, -2 and -3 (MASP-1, MASP-2 and MASP-3, respectively)), which have been taught to play a significant role in modulation of the lectin pathway of complement activation. Petersen et al., *Molecular Immunology* 38:133-149 (2001).

[0320] MASP-1 has a histidine loop structure of the type found in trypsin and trypsin-like serine proteases. MASP-1 has been found to be involved in complement activation by MBL. A cDNA clone encoding MASP-1 has been reported that encodes a putative leader peptide of 19 amino acids followed by 680 amino acid residues predicted to form the mature peptide. MASP-2 (MBL-associated serine protease 2) is a serine protease also similar in structure to C1 r and C1 s of the complement pathway. Like these, and contrary to MASP-1, it has no histidine loop structure of the type found in trypsin and trypsin-like serine proteases. It has been theorized that MASP-1 can cleave C3, generating C3b, which may be deposited on an activated cell or tissue surface

[0321] It has been shown that MASP-2, cleaves C4 and C2, giving rise to the C3 convertase, C4b2b (Thiel et al., *Nature*, 386:506-10 (1997)). The MASP-2 protein comprises of a number of domains namely the CUB1, EGF, CUB2, CCP1, CCP2 and serine protease domains. It is believed that the domain responsible for association with MBL is situated in the N-terminus, whereas the serine protease domain is responsible for the serine protease activity of MASP-2. sMAP, also known as MAP19, is a 19 kd is derived from the same gene as MASP-2, which lacks the serine protease domain and a major part of the A chain. Skjoedt et al., *Immunobiology*, 215:921-31 (2010). Recently, a third member of the family, MASP-3 was identified, which shares a high degree of homology with MASP-1, such that it appears that

MASP-1 and MASP-3 are generated as a result of alternative splicing of primary mRNA transcripts.

[0322] Antibodies against MBL, MASP-1, MASP-2, MASP-3 and the MBL/MASP complex, and their use for inhibiting the adverse effects of complement activation, such as ischemia-reperfusion injury, have been disclosed, for example, in WO04/075837; US 2009/0017031.

[0323] Other antibodies to MASP-2 have been described previously, as well. See, e.g., WO 02/06460, US2007/0009528, Peterson et al., *Mol. Immunol.* 37:803-11 (2000), Moller-Kristensen et al., *J. of Immunol. Methods* 282:159-67 (2003), Petersen et al., *Mol. Immunol.* 35:409, and WO 04/106384.

[0324] An additional related protein, MBL/Ficolin Associated Protein (MAP-1), which is present in low serum levels compared to MASP-1 and MASP-3, has been reported to function as a local lectin pathway specific complement inhibitor. Skjodt et al., *Molecular Immunology*, 47:2229-30 (2010). Accordingly MAP-1 itself, or fragments of MAP-1, may be useful in the present invention as an inhibitor of MASP, and accordingly, as a lectin-pathway-specific inhibitor of complement activation. Finally, the ficolin family of proteins are characterized by carbohydrate binding and opsonic activities, sharing a structure similar to MBL. Like MBL, the ficolins have been shown to associate with MASPs in serum and may mediate complement activation in response to pathogenic, necrotic, or apoptotic cell-specific carbohydrate markers. Accordingly, inhibitors of the ficolin family or functional fragments thereof may be useful in the present invention as an inhibitor of MASPs and as a lectin-pathway specific inhibitor of complement activation. US Patent 6,333,034 and US Patent 7,423,128; see also, WO 2008/154018 and WO 2009/110918.

[0325] In some embodiments, the complement inhibitor portion is an antibody (or antigen binding fragment thereof) that specifically binds to a human complement component protein (e.g., human C5, C6, C7, C8, or C9). The terms “specific binding” or “specifically binds” refer to two molecules forming a complex (e.g., a complex between an antibody and a complement component protein) that is relatively stable under physiologic conditions. Typically, binding is considered specific when the association constant (K_a) is higher than 10^6 M^{-1} . Thus, an antibody can specifically bind to a C5 protein with a K_a of at least (or greater than) 10^6 (e.g., at least or greater than 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , or 10^{15} or higher) M^{-1} . Examples of antibodies that specifically bind to a human complement component C5 protein are

described in, e.g., U.S. patent no. 6,355,245 and PCT application publication no. WO 2010/015608.

[0326] Methods for determining whether an antibody binds to a protein antigen and/or the affinity for an antibody to a protein antigen are known in the art and described herein. For example, the binding of an antibody to a protein antigen can be detected and/or quantified using a variety of techniques such as, but not limited to, Western blot, dot blot, surface plasmon resonance method (e.g., BIAcore system; Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, N.J.), or enzyme-linked immunosorbent assay (ELISA) assays. See, e.g., Harlow and Lane (1988) “Antibodies: A Laboratory Manual” Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; Benny K. C. Lo (2004) “Antibody Engineering: Methods and Protocols,” Humana Press (ISBN: 1588290921); Borrebaek (1992) “Antibody Engineering, A Practical Guide,” W.H. Freeman and Co., NY; Borrebaek (1995) “Antibody Engineering,” 2nd Edition, Oxford University Press, NY, Oxford; John et al. (1993) *J Immunol Meth.* 160:191- 198; Jonsson et al. (1993) *Ann Biol Clin* 51:19-26; and Jonsson et al. (1991) *Biotechniques* 11:620-627. See also, U.S. Patent No. 6,355,245.

[0327] In any of the embodiments described herein, the targeting construct also includes an amino acid linker sequence linking the targeting moiety and the therapeutic moiety (or a biologically active fragment thereof).

[0328] In some embodiments, the targeting moiety of the targeting construct is joined (e.g., directly or by way of a linker) to the amino-terminus of the therapeutic moiety. In some embodiments, the targeting moiety of the targeting construct is joined (e.g., directly or by way of a linker) to the carboxy-terminus of the therapeutic moiety (e.g., a complement inhibitor or drug described herein).

[0329] In some embodiments, a targeting construct described herein comprises more than one (e.g., two, three, four, five, six, or seven or more) therapeutic moiety, e.g., more than one complement inhibitor polypeptide or drug described herein. The two or more therapeutic moieties can be the same or different. For example, a targeting construct described herein can comprise, in some embodiments, two or more soluble CD59 portions (e.g., soluble human CD59 portions) or two or more beta blockers. In another example, a targeting construct described herein can contain two or more complement inhibitor polypeptide portions, wherein one is a soluble human CD59 and another is soluble human MCP. In another example, a targeting construct

described herein can contain a complement inhibitor and a drug, e.g., one soluble CD59 portion and one corticosteroid. Thus, e.g., a targeting construct described herein can comprise: (a) a targeting moiety (e.g., a C2 antibody, a B4 antibody, or an antigen-binding fragment of either of the foregoing); (b) a first therapeutic moiety (e.g., a soluble form of CD59, e.g., human CD59); and (c) a second therapeutic moiety (e.g., a soluble form of DAF, e.g., a soluble form of human DAF, or a corticosteroid such as prednisone). The therapeutic moiety can be, e.g., any of those described herein including variants and biologically active fragments of the complement inhibitors described herein.

[0330] In some embodiments, the light chain of the targeting moiety of the targeting construct comprises at least one therapeutic moiety and the heavy chain comprises at least therapeutic moiety. The two or more complement inhibitor polypeptides can be the same or different. For example, in some embodiments, the targeting construct comprises the Fab fragment of a targeting moiety described herein, wherein: (i) the light chain of the Fab fragment comprises (at its C-terminal end) a complement inhibitor polypeptide such as DAF, CD59, or any of the complement inhibitor polypeptides described herein and (ii) the heavy chain of the Fab fragment comprises (at its C-terminal end) the same or a different therapeutic moiety as in (i), e.g., a complement inhibitor or a drug described herein. Appropriate pairing of the two chains can be expected to occur as an inherent property of the Fab. The complement inhibitor portion and the light chain or heavy chain of the Fab can be joined together directly or by way of a linker sequence (such as any of those described herein).

[0331] Detectable moieties

[0332] In some embodiments, the targeting construct comprises a targeting moiety fused to an active moiety that is detectable moiety. In some embodiments, the detectable moiety can be a paramagnetic molecule, a paramagnetic nanoparticle, an ultrasmall superparamagnetic iron oxide (“USPIO”) nanoparticle, or a USPIO nanoparticle aggregate. In certain embodiments, the USPIO nanoparticle aggregate is between about 10 nm and about 150 nm in diameter, between about 65 nm and about 85 nm in diameter, or about 75 nm in diameter. In certain embodiments, the USPIO nanoparticle aggregate is about 150 nm in diameter. In certain embodiments, the USPIO nanoparticle aggregate is coated with dextran or an amphiphilic polymer, or the USPIO nanoparticle aggregate is encapsulated with phospholipid. In certain

embodiments, the phospholipid is PEGylated. In certain embodiments, the PEGylated phospholipid is amine-functionalized or carboxylic acid-functionalized. In certain embodiments the PEGylated, amine-functionalized phospholipid is 1,2-distearoyl-sn-glycero-3-phosphoethanolamine-PEG2000.

[0333] In certain embodiments, the a paramagnetic nanoparticle is a superparamagnetic iron oxide (“SPIO”) nanoparticle, an SPIO nanoparticle aggregate, standard superparamagnetic iron oxide (“SSPIO”), an SSPIO nanoparticle aggregate, polydisperse superparamagnetic iron oxide (“PSPIO”), a PSPIO nanoparticle aggregate, monocrystalline SPIO, a monocrystalline SPIO aggregate, a monocrystalline iron oxide nanoparticle, monocrystalline iron oxide, or any other nanoparticle contrast agent known to one of skill in the art. Methods of conjugating such particles to a targeting moiety and detecting paramagnetic particle-conjugated targeting moieties are known in the art and described in, e.g., Richardson et al. (2001) *Biosens and Bioelect* 16:989-993; Boyaci et al. (2005) *Anal Bioanal Chem* 382(5):1234-41; Toma et al. (2005) *Br J Cancer* 93(1):131-6; and others.

[0334] In some embodiments, the detectable moiety is a liposome or another delivery vehicle containing Gadolinium chelate (“Gd-chelate”) molecules. In some embodiments, the detectable moiety is an electron-dense reagent, such as Gadolinium, an iodinated contrast agent, barium sulfate, thorium dioxide, gold, a gold nanoparticle, a gold nanoparticle aggregate. In some embodiments, the detectable moiety is a biocolloid, or a microbubble. In some embodiments, the detectable moiety is a radioisotope or a radionuclide, including, but not limited to, e.g., ³²P, carbon-11, nitrogen-13, oxygen-15, fluorine-18, rubidium-82, fluorodeoxyglucose, a gamma ray emitting radionuclide, radiolabeled glucose, radiolabeled water, or radiolabeled ammonia. In certain embodiments, the detectable moiety is a positron-emitting radionuclide. In certain embodiments, the detectable moiety is a hapten, a protein (such as biotin), an enzyme, a digoxigenin, or a fluorophore, a two-photon fluorophore, a fluorescent dye, or a fluorescent moiety (e.g., a fluorescein, fluorescein isothiocyanate, or a fluorescein derivative). Methods of conjugating such detectable moieties to a targeting moiety are known in the art and are described elsewhere herein.

[0335] Targeting construct comprising a detectable moiety can be used in noninvasive methods of detecting complement-mediated inflammation or complement activation in the eye of an individual in need thereof. As noted elsewhere herein, the

methods include administering to the individual a composition comprising an effective amount of a targeting construct comprising a detectable moiety and measuring the presence of the detectable moiety using an instrument and/or method (e.g. MRI, CT, SPECT, radiography, spectroscopy, microscopy, PET, ultrasound, or any other detection method described herein) capable of detecting the presence of the detectable moiety.

[0336] MRI can be used to non-invasively acquire tissue images with high resolution. Paramagnetic agents or USPIO nanoparticles or aggregates thereof enhance signal attenuation on T2-weighted magnetic resonance images, and conjugation of such nanoparticles to, e.g., an antibody described herein (or a fragment thereof) or a construct described herein, permits the detection of specific molecules at the cellular level. For example, MRI with nanoparticle detection agents can image cell migration (J. W. Bulte et al., 2001, Nat. Biotechnol. 19:1141-1147), apoptosis (M. Zhao et al., 2001, Nat. Med. 7:1241-1244), and can detect small foci of cancer. See e.g., Y. W. Jun et al., 2005, J. Am. Chem. Soc. 127:5732-5733; Y. M. Huh et al., 2005, J. Am. Chem. Soc. 127:12387-12391. Contrast-enhanced MRI is well-suited for the dynamic non-invasive imaging of macromolecules or of molecular events, but it requires ligands that specifically bind to the molecule of interest. J. W. Bulte et al., 2004, NMR Biomed. 17:484-499. Fluorescent dyes and fluorophores e.g. fluorescein, fluorescein isothiocyanate, and fluorescein derivatives) can be used to non-invasively acquire tissue images with high resolution, with for example spectrophotometry, two-photon fluorescence, two-photon laser microscopy, or fluorescence microscopy (e.g. of tissue biopsies). MRI can be used to non-invasively acquire tissue images with high resolution, with for example paramagnetic molecules, paramagnetic nanoparticles, ultrasmall superparamagnetic iron oxide ("USPIO") nanoparticles, USPIO nanoparticle aggregates, superparamagnetic iron oxide ("SPIO") nanoparticles, SPIO nanoparticle aggregates, monocrystalline iron oxide nanoparticles, monocrystalline iron oxide, other nanoparticle contrast agents. MRI can be used to non-invasively acquire tissue images with high resolution, with for example Gadolinium, including liposomes or other delivery vehicles containing Gadolinium chelate ("Gd-chelate") molecules. Positron emission tomography (PET), PET/computed tomography (CT), single photon emission computed tomography (SPECT), and SPECT/CT can be used to non-invasively acquire tissue images with high resolution, with for example radionuclides (e.g. carbon-11, nitrogen-13, oxygen-

15, fluorine-18, rubidium-82), fluorodeoxyglucose (e.g. fluorine-18 labeled), any gamma ray emitting radionuclides, positron-emitting radionuclide, radiolabeled glucose, radiolabeled water, radiolabeled ammonia. Ultrasound (ultrasonography) and contrast enhanced ultrasound (contrast enhanced ultrasonography) can be used to non-invasively acquire tissue images with high resolution, with for example biocolloids or microbubbles (e.g. including microbubble shells including albumin, galactose, lipid, and/or polymers; microbubble gas core including air, heavy gas(es), perfluorocarbon, nitrogen, octafluoropropane, perflubron lipid microsphere, perflubron, etc.). X-ray imaging (radiography) or CT can be used to non-invasively acquire tissue images with high resolution, with for example iodinated contrast agents (e.g. iohexol, iodixanol, ioversol, iopamidol, ioxilan, iopromide, diatrizoate, metrizoate, ioxaglate), barium sulfate, thorium dioxide, gold, gold nanoparticles, or gold nanoparticle aggregates. These detection methods and instruments and detectable moieties capable of measured or detected by the corresponding method are non-limiting examples.

[0337] As used herein, the term "ultrasmall superparamagnetic iron oxide nanoparticle" or "USPIO nanoparticle" refers to superparamagnetic iron oxide particles ranging from 1 to 50 nm in diameter, more typically between 5 and 40 nm in diameter (excluding any coating applied after synthesis). USPIO nanoparticles are commonly made of maghemite (Fe_2O_3) or magnetite (Fe_3O_4) having crystal-containing regions of unpaired spins. Those magnetic domains are disordered in the absence of a magnetic field, but when a field is applied (i.e., while taking an MRI), the magnetic domains align to create a magnetic moment much greater than the sum of the individual unpaired electrons without resulting in residual magnetization of the particles. When injected into the blood stream, USPIO nanoparticles are taken up by macrophages and accumulate in inflamed tissues. Their iron moiety negatively enhances signal attenuation on T2-weighted images, and their relative concentrations can be assessed by decreased T2-signal intensity or, more precisely, by decreased spin-spin T2-relaxation time. The decreased T2-relaxation time (the transverse relaxation time) can thus be used to detect inflammation. The shortened T2 relaxation time results in a darkening of the magnetic resonance image where the particles are located, thereby generating "negative contrast." This approach has been successfully utilized to detect renal inflammation in several models. In some cases, USPIO nanoparticles may be aggregated after synthesis to produce aggregates thereof (referred to herein as "ultrasmall superparamagnetic iron oxide" ("USPIO"))

nanoparticle aggregates" or "USPIO nanoparticle aggregates") of 25 nm, 50 nm, 75 nm, 100 nm, or 150 nm, in diameter, or even larger.

[0338] The USPIO nanoparticles or aggregates thereof may be coated with a wide variety of materials, including natural or synthetic polymers, surfactants, phospholipids, or inorganic materials, any of which may be modified or derivatized to permit attachment of targeting groups, either directly or via different types of linkers, including peptides, polypeptides, proteins, or other chemical groups, or uncoated. Possible coatings include synthetic polymers, such as those based on poly(ethylene-co-vinyl acetate), polyvinylpyrrolidone ("PVP"), poly(lactic-co-glycolic acid) ("PLGA"), polyethylene glycol ("PEG"), polyvinyl alcohol ("PVA"), polyacrylic acid, and the like; natural polymers, such as gelatin, dextran, chitosan, pullulan, and the like; surfactants, such as sodium oleate, dodecylamine, sodium carboxymethylcellulose, and the like; inorganic materials, such as gold or silica; and biological materials, such as phospholipids.

[0339] Thus, a complement-mediated inflammation (such as in the eye) can be detected in an individual in a non-invasive manner by administering an the antibody-targeted USPIO nanoparticle or nanoparticle aggregate compositions and/or USPIO nanoparticle- or USPIO nanoparticle aggregate-conjugated targeting constructs provided herein, and taking a magnetic resonance taking a magnetic resonance image of the individual, or of the individual's eye. In some of the embodiments described herein, the composition administered to the individual is a pharmaceutical composition comprising any of the antibody (or antigen-binding fragment thereof) and/or construct described herein. In some of the embodiments described herein, the composition administered to the individual is a pharmaceutical composition comprising any of the antibody-targeted USPIO nanoparticle aggregate compositions described herein.

[0340] As used herein, the term "magnetic resonance imaging "or "MRI" refers to a non-invasive medical imaging technique commonly used to visualize the structure and function of the body that provides detailed images of the body in any plane. MRI provides much greater contrast between the different soft tissues of the body than other non-invasive imaging methods, such as computed tomography (CT), making it especially useful in neurological, musculoskeletal, cardiovascular, and oncological (cancer) imaging. Unlike CT, it does not require ionizing radiation, instead using a powerful magnetic field to align the nuclear magnetization of hydrogen atoms in

water in the body. Radiofrequency fields are used to systematically alter the alignment of this magnetization, causing the hydrogen nuclei to produce a rotating magnetic field detectable by the scanner. This signal can be manipulated by additional magnetic fields to build up enough information to reconstruct an image of the body or a portion thereof, e.g., the eye.

[0341] When an individual lies in a scanner, the hydrogen nuclei (i.e., protons) found in abundance in water molecules throughout the individual's body, align with the strong main magnetic field. A second electromagnetic field, which oscillates at radiofrequencies and is perpendicular to the main field, is then pulsed to push a proportion of the protons out of alignment with the main field. These protons then drift back into alignment with the main field, emitting a detectable radiofrequency signal as they do so. Since protons in different body tissues (e.g., fat vs. muscle) realign at different speeds, different body structures can be imaged. Contrast agents may be injected intravenously to enhance the appearance of blood vessels, organs (e.g., the eye), tumors or sites of inflammation.

[0342] As used herein, an "effective amount" or "diagnostically effective amount" of an antibody-targeted ultrasmall superparamagnetic iron oxide ("USPIO") nanoparticle or nanoparticle aggregate composition (including any of the pharmaceutical compositions described herein) is an amount sufficient to produce a clinically useful magnetic resonance image of complement-mediated inflammation such as in the eye. A clinically useful magnetic resonance image is one containing sufficient detail to enable an experienced clinician to assess the degree and/or extent of inflammation for purposes of diagnosis, monitoring the efficacy of a therapeutic intervention, and the like. As used herein, an "effective amount" or "diagnostically effective amount" of an antibody-targeted detectable moiety or a targeting construct comprising a detectable moiety (including any of the pharmaceutical compositions described herein) is an amount sufficient to produce a clinically useful characterization or measurement of complement-mediated inflammation or complement activation (e.g. in an individual, patient, human, mammal, clinical sample, tissue, biopsy) when coupled with a detection method capable of detecting the antibody (or fragment thereof) and/or the targeting moiety. A clinically useful characterization or measurement of complement-mediated inflammation or complement activation is one containing sufficient detail to enable an experienced clinician to assess the degree and/or extent of inflammation or complement activation

for purposes of diagnosis, monitoring the efficacy of a therapeutic intervention, and the like.

[0343] Delivery of ultrasmall superparamagnetic iron oxide ("USPIO") nanoparticles, USPIO nanoparticle aggregates, superparamagnetic iron oxide ("SPIO") nanoparticles, SPIO nanoparticle aggregates or other nanoparticle contrast agents (examples of detectable moieties) to the sites of active inflammation via delivery of antibodies and/or targeting construct to sites of complement activation permits non-invasive magnetic resonance imaging of such inflammation, enabling the specific detection of complement activation throughout the body, and distinguishing complement-mediated inflammation from other types of inflammation.

[0344] Accordingly, in one aspect, the invention provides compositions comprising nanoparticle contrast agents conjugated to an antibody (or antigen binding fragment thereof) or a construct described herein for non-invasive medical or diagnostic imaging applications. In certain embodiments, the nanoparticle contrast agent-conjugated antibody (or antigen binding fragment thereof) or a construct comprises USPIO nanoparticles or aggregates thereof. In certain embodiments, the nanoparticle contrast agent-conjugated antibody (or antigen binding fragment thereof) or a construct comprise liposomes or other vehicles containing Gadolinium chelate ("Gdchelate") molecules. Ultrasmall super paramagnetic iron oxide ("USPIO") nanoparticles or aggregates are examples of detectable moieties that can be conjugated to a targeting construct described herein.

[0345] At least two physicochemical characteristics of ultrasmall super paramagnetic iron oxide ("USPIO") nanoparticles or aggregates thereof vary with the size of the individual nanoparticles or nanoparticle aggregates. First, the ability of USPIO nanoparticle preparations to enhance contrast in MRI imaging and the degree of contrast enhancement both vary with nanoparticle diameter, because the magnetic moment of individual USPIO nanoparticles also varies with particle diameter. Iron oxide nanoparticles with diameters up to approximately 15 nm (preferably less than 10 nm) remain super paramagnetic, but larger iron oxide nanoparticles lose their superparamagnetic properties. Thus, there is an upper limit to the diameter of USPIO nanoparticles suitable for use as MRI contrast reagents. This limitation can be overcome by use of multiparticle aggregates of smaller individual USPIO nanoparticles. Such USPIO nanoparticle aggregates effectively enhance MRI contrast because the magnetic moments of the individual nanoparticles within each

nanoparticle aggregate are additive. Unlike individual iron oxide nanoparticles, aggregates of ultra small super paramagnetic iron oxide nanoparticles do not lose their paramagnetic properties with increased size.

[0346] Second, the in vivo half-life (e.g., circulating plasma or blood half-life and tissue half-life) and biodistribution of USPIO nanoparticles or aggregates thereof varies with nanoparticle or aggregate size. For example, USPIO nanoparticles ~10 nm or less in diameter (monocrystalline iron oxide nanoparticles) have a circulating blood half-life of ~81 minutes (R. Weissleder et al., 1990, *Radial.* 175(2):489-493), USPIO nanoparticles ~50 nm in diameter have a circulating half-life of ~30 minutes (D. Pouliquen et al., 1991, *Magnet. Resonance Imag.* 9(3):275-283), USPIO nanoparticles ~150 nm in diameter are thought to have a circulating half-life of less than ~30 minutes, and USPIO nanoparticles ~80 nm in diameter have a tissue half-life on the order of one to several days (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, or more days) and a whole body half-life of ~45 days (R. Weissleder et al., 1989, *Am. J. Roentgenol.* 152(1):167-173). Effective targeted MRI contrast-enhancing reagents must circulate in the vasculature long enough to recognize and bind the desired target (e.g., annexin IV or a phospholipid such as PE, PC, CL, or MDA) while still being cleared quickly enough to minimize any potential toxicity. Optimal USPIO nanoparticle or nanoparticle aggregate sizes for generating clinically useful magnetic resonance images vary depending on the organ (e.g., the kidney, eye, retina), tissue, and/or physiological phenomenon (e.g., complement-mediated inflammation) to be imaged.

[0347] The circulating half-life of USPIO nanoparticles or nanoparticle aggregates can also be altered (i.e., reduced or extended) by coating them with different materials. For instance, USPIO nanoparticles or nanoparticle aggregates can be coated with natural or synthetic polymers, surfactants, or phospholipids, among other materials, any of which may be modified or derivatized to permit attachment of an antibody (or antigen binding fragment thereof) and/or a construct described herein, either directly or indirectly via different types of linkers, including peptides, polypeptides, proteins, or other chemical groups. In some cases, the coatings may be further modified to incorporate synthetic polymers, natural polymers, amphiphilic polymers, or other molecules (e.g., polyvinylpyrrolidone ("PVP"), poly (lactic-coglycolic acid) ("PLGA"), polyethylene glycol ("PEG"), polyvinyl alcohol ("PYA"), polyacrylic acid, and the like) suitable for stabilizing the aggregates or minimizing their susceptibility to extravasation, opsonization, phagocytosis, endocytosis or other

modes of physiological clearance. In some cases, USPIO nanoparticles or nanoparticle aggregates conjugated to an antibody (or antigen-binding fragment thereof) or construct can be phospholipid-encapsulated. As with USPIO nanoparticle or nanoparticle aggregate size, the particular coating, modification or derivatization suitable for targeting the nanoparticles or nanoparticle aggregates to a desired organ (e.g., the kidney, eye, retina), tissue, and/or physiological phenomenon (e.g., complement-mediated inflammation) may be determined empirically.

[0348] Variants of targeting constructs

[0349] Also encompassed are variants of the targeting constructs. A variant of the targeting construct described herein may be: (i) one in which one or more of the amino acid residues of the targeting moiety and/or the active moiety (i.e., wherein the active moiety comprises a protein) are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code; or (ii) one in which one or more of the amino acid residues in the targeting and/or active moiety includes a substituent group, or (iii) one in which the targeting construct is fused with another compound, such as a compound to increase the half-life of the targeting construct (for example, polyethylene glycol), or (iv) one in which additional amino acids are fused to the targeting construct (such as the targeting moiety or the active moiety, wherein the active moiety comprises a protein), such as a leader or secretory sequence or a sequence which is employed for purification of the targeting construct, or (v) one in which the targeting construct is fused with a larger polypeptide, i.e., human albumin, an antibody or Fc, for increased duration of effect. Such variants are deemed to be within the scope of those skilled in the art from the teachings herein.

[0350] In some embodiments, the variant of the targeting construct contains conservative amino acid substitutions (defined further below) made at one or more predicted, preferably nonessential amino acid residues. A “nonessential” amino acid residue is a residue that is altered from the wild-type sequence of a protein without altering the biological activity, whereas an “essential” amino acid residue is required for biological activity. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain.

[0351] Twenty amino acids are commonly found in proteins. Those amino acids can be grouped into nine classes or groups based on the chemical properties of their

side chains. Substitution of one amino acid residue for another within the same class or group is referred to herein as a “conservative” substitution. Conservative amino acid substitutions can frequently be made in a protein without significantly altering the conformation or function of the protein. Substitution of one amino acid residue for another from a different class or group is referred to herein as a “non-conservative” substitution. In contrast, non-conservative amino acid substitutions tend to disrupt conformation and function of a protein. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). (See Table 1 below.)

Table 1: Example of amino acid classification

Small/Aliphatic residues:	Gly, Ala, Val, Leu, Ile
Cyclic Imino Acid:	Pro
Hydroxyl-containing Residues:	Ser, Thr
Acidic Residues:	Asp, Glu
Amide Residues:	Asn, Gln
Basic Residues:	Lys, Arg
Imidazole Residue:	His
Aromatic Residues:	Phe, Tyr, Trp
Sulfur-containing Residues:	Met, Cys

[0352] In some embodiments, the conservative amino acid substitution comprises substituting any of glycine (G), alanine (A), isoleucine (I), valine (V), and leucine (L) for any other of these aliphatic amino acids; serine (S) for threonine (T) and vice versa; aspartic acid (D) for glutamic acid (E) and vice versa; glutamine (Q) for asparagine (N) and vice versa; lysine (K) for arginine (R) and vice versa; phenylalanine (F), tyrosine (Y) and tryptophan (W) for any other of these aromatic amino acids; and methionine (M) for cysteine (C) and vice versa. Other substitutions can also be considered conservative, depending on the environment of the particular amino acid and its role in the three-dimensional structure of the protein. For example, glycine (G) and alanine (A) can frequently be interchangeable, as can alanine (A) and valine (V). Methionine (M), which is relatively hydrophobic, can frequently be interchanged with leucine and isoleucine, and sometimes with valine. Lysine (K) and

arginine (R) are frequently interchangeable in locations in which the significant feature of the amino acid residue is its charge and the differing pKs of these two amino acid residues are not significant. Still other changes can be considered “conservative” in particular environments (see, e.g., Biochemistry at pp. 13-15, 2nd ed. Lubert Stryer ed. (Stanford University); Henikoff et al., Proc. Nat’l Acad. Sci. USA (1992) 89:10915-10919; Lei et al., J. Biol. Chem. (1995) 270(20):11882-11886).

[0353] Amino acid substitutions in the targeting moiety and/or the active moiety of the targeting construct is introduced to improve the functionality of the targeting construct. For example, amino acid substitutions can be introduced into the targeting moiety of targeting construct to increase binding affinity of the targeting moiety to its ligand(s), increase binding specificity of the targeting construct to its ligand(s), improve targeting of the targeting construct to desired sites, increase dimerization or multimerization of the targeting construct, and improve pharmacokinetics of the targeting construct. Similarly, amino acid substitutions can be introduced into the active moiety of the targeting construct to increase the functionality of the targeting construct molecule and improve pharmacokinetics of the targeting construct.

[0354] In some embodiments, the targeting construct is fused with another compound, such as a compound to increase the half-life of the targeting construct and/or to reduce potential immunogenicity of the targeting construct (for example, polyethylene glycol, “PEG”). The PEG can be used to impart increased stability, water solubility, size, slow rate of kidney clearance, and reduced immunogenicity to the targeting construct. See e.g., U.S. Pat. No. 6,214,966; Lee et al. (1999) Bioconjug Chem 10(6): 973-8; Kinstler et al. (2002) Advanced Drug Deliveries Reviews 54:477-485; and Roberts et al. (2002) Advanced Drug Delivery Reviews 54:459-476. The stabilization moiety can improve the stability, or retention of, the polypeptide by at least 1.5 (e.g., at least 2, 5, 10, 15, 20, 25, 30, 40, or 50 or more) fold. In the case of PEGylations, the fusion of a targeting construct described herein to PEG can be accomplished by any means known to one skilled in the art. For example, PEGylation can be accomplished by first introducing a cysteine mutation into the targeting moiety or the active moiety (i.e., wherein the active moiety comprises a protein), followed by site-specific derivatization with PEG-maleimide. The cysteine can be added to the C-terminus of the targeting construct. See, e.g., Tsutsumi et al. (2000) Proc. Natl. Acad. Sci. USA 97(15):8548-8553. Another modification which can be made to the targeting construct involves biotinylation. In certain instances, it may be useful to

have the targeting construct biotinylated so that it can readily react with streptavidin. Methods for biotinylation of proteins are well known in the art. Additionally, chondroitin sulfate can be linked with the targeting construct.

[0355] In some embodiments, the targeting construct is fused to another moiety which further increases the targeting efficiency of the targeting construct. For example, a targeting construct comprising a B4 antibody can be fused to, e.g., a C2 antibody or another antibody that has the capability to bind or otherwise attach to an endothelial cell of a blood vessel (referred to as “vascular endothelial targeting amino acid ligand”). Exemplary vascular endothelial targeting ligands include, but are not limited to, VEGF, FGF, integrin, fibronectin, I-CAM, PDGF, or an antibody to a molecule expressed on the surface of a vascular endothelial cell.

[0356] In some embodiments, the targeting construct is conjugated (such as fused) to a ligand for intercellular adhesion molecules. For example, the target construct molecule can be conjugated to one or more carbohydrate moieties that bind to an intercellular adhesion molecule. The carbohydrate moiety facilitates localization of the target construct molecule to the site of injury. The carbohydrate moiety can be attached to the target construct molecule by means of an extracellular event such as a chemical or enzymatic attachment, or can be the result of an intracellular processing event achieved by the expression of appropriate enzymes. In some embodiments, the carbohydrate moiety binds to a particular class of adhesion molecules such as integrins or selectins, including E-selectin, L-selectin or P-selectin. In some embodiments, the carbohydrate moiety comprises an N-linked carbohydrate, for example the complex type, including fucosylated and sialylated carbohydrates. In some embodiments, the carbohydrate moiety is related to the Lewis X antigen, for example the sialylated Lewis X antigen.

[0357] For treatment of eye diseases such as AMD, the targeting construct can be conjugated (such as fused) to an antibody that recognizes an epitope of the drusen. Other targeting molecules such as small targeting peptide can also be used. Other modifications of the targeting construct include, for example, glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, and the like.

[0358] The targeting construct may include the addition of an immunologically active domain, such as an antibody epitope or other tag, to facilitate targeting or purification of the polypeptide. The use of 6xHis and GST (glutathione S transferase)

as tags is well known. Inclusion of a cleavage site at or near the fusion junction will facilitate removal of the extraneous polypeptide from the targeting construct after purification. Other amino acid sequences that may be included in the targeting construct include functional domains, such as active sites from enzymes such as a hydrolase, glycosylation domains, and cellular targeting signals.

[0359] Variants of the targeting construct include polypeptides having an amino acid sequence sufficiently similar to the amino acid sequence of a targeting construct described herein. The term “sufficiently similar” means a first amino acid sequence that contains a sufficient or minimum number of identical or equivalent amino acid residues relative to a second amino acid sequence such that the first and second amino acid sequences have a common structural domain and/or common functional activity. For example, amino acid sequences that contain a common structural domain that is at least about 45%, preferably about 75% through 98%, identical are defined herein as sufficiently similar. Variants include variants of targeting constructs encoded by a polynucleotide that hybridizes to a polynucleotide of this invention or a complement thereof under stringent conditions. Such variants generally retain the functional activity of the targeting constructs of this invention. Libraries of fragments of the polynucleotides can be used to generate a variegated population of fragments for screening and subsequent selection. For example, a library of fragments can be generated by treating a double-stranded PCR fragment of a polynucleotide with a nuclease under conditions wherein nicking occurs only about once per molecule, denaturing the double-stranded DNA, renaturing the DNA to form double-stranded DNA which can include sense/antisense pairs from different nicked products, removing single-stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, one can derive an expression library that encodes N-terminal and internal fragments of various sizes of the targeting constructs of this invention.

[0360] Variants include targeting constructs that differ in amino acid sequence due to mutagenesis. In addition, bioequivalent analogs of the targeting constructs may also be constructed by making various substitutions on residues or sequences in the targeting moiety and/or the active moiety.

[0361] In some embodiments, targeting construct is fused at its N-terminus to a signal peptide. Such signal peptides are useful for the secretion of the targeting construct. Suitable signal peptides include, for example, the signal peptide of the

CD5 protein (such as signal peptide of the human CD5 protein MPMGSLQPLATLYLLGMLVAS, SEQ ID NO:54). In some embodiments, the signal peptide of the CR2 protein is used. For example, in some embodiments, the signal peptide of the human CR2 protein (MGAAGLLGVFLALVAPG, SEQ ID NO:55 or MGAAGLLGVFLALVAPGVLG, SEQ ID NO:56) is used.

[0362] Targeting construct production methods

[0363] The targeting construct described herein can be produced using a variety of techniques known in the art of molecular biology and protein chemistry. For example, a nucleic acid encoding a targeting construct described herein can be inserted into an expression vector that contains transcriptional and translational regulatory sequences, which include, e.g., promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, transcription terminator signals, polyadenylation signals, and enhancer or activator sequences. The regulatory sequences include a promoter and transcriptional start and stop sequences. In addition, the expression vector can include more than one replication system such that it can be maintained in two different organisms, for example in mammalian or insect cells for expression and in a prokaryotic host for cloning and amplification.

[0364] Several possible vector systems are available for the expression of targeting constructs from nucleic acids in mammalian cells. One class of vectors relies upon the integration of the desired gene sequences into the host cell genome. Cells which have stably integrated DNA can be selected by simultaneously introducing drug resistance genes such as *E. coli* gpt (Mulligan and Berg (1981) *Proc Natl Acad Sci USA* 78:2072) or Tn5 neo (Southern and Berg (1982) *Mol Appl Genet* 1:327). The selectable marker gene can be either linked to the DNA gene sequences to be expressed, or introduced into the same cell by co-transfection (Wigler et al. (1979) *Cell* 16:77). A second class of vectors utilizes DNA elements which confer autonomously replicating capabilities to an extrachromosomal plasmid. These vectors can be derived from animal viruses, such as bovine papillomavirus (Sarver et al. (1982) *Proc Natl Acad Sci USA*, 79:7147), polyoma virus (Deans et al. (1984) *Proc Natl Acad Sci USA* 81: 1292), or SV 40 virus (Lusky and Botchan (1981) *Nature* 293:79).

[0365] The expression vectors can be introduced into cells in a manner suitable for subsequent expression of the nucleic acid. The method of introduction is largely

dictated by the targeted cell type, discussed below. Exemplary methods include CaP04 precipitation, liposome fusion, lipofectin, electroporation, viral infection, dextran-mediated transfection, polybrene-mediated transfection, protoplast fusion, and direct microinjection.

[0366] Appropriate host cells for the expression of the targeting constructs include yeast, bacteria, insect, plant, and, as described above, mammalian cells. Of interest are bacteria such as *E. coli*, fungi such as *Saccharomyces cerevisiae* and *Pichia pastoris*, insect cells such as SF9, mammalian cell lines (e.g., human cell lines), as well as primary cell lines (e.g., primary mammalian cells). In some embodiments, the targeting constructs can be expressed in Chinese hamster ovary (CHO) cells or in a suitable myeloma cell line such as (NSO). Suitable cell lines also include, for example, BHK-21 (baby hamster kidney) cells; 293 (human embryonic kidney) cells; HMEpC (Human Mammary Epithelial cells); 3T3 (mouse embryonic fibroblast) cells.

[0367] The targeting moiety and the one or more active moieties may optionally be directly joined to each other, or may optionally be joined via a linker. Where the targeting and active moieties are directly joined, the hybrid vector is made where the DNA encoding the targeting and active moieties are themselves directly ligated to each other using known scientific methods. Where a linker is used, the hybrid vector is made where the DNA encoding the targeting moiety is ligated to DNA encoding one end of the linker; and the DNA encoding the active moiety is ligated to the other end of the linker. Methods are known for performing such ligations in proper orientation. Such ligation may be performed either in series, or as a three way ligation. Examples of sequences which may serve as the linker sequence in the present invention include short peptides of about 2 to about 16 amino acids in length. Among the peptide sequences useful as linkers in the present invention are (Gly-Ser) $_n$, where $n = 1$ to 8; (GlyGlyGlySer) $_n$, where $n = 1$ to 4; (GlySerSerGly) $_n$, where $n = 1$ to 4. Other examples of sequences useful as the linker sequence in the present invention include one or more short conserved region (SCR) domains from one or more of the following complement-related proteins: Factor H; complement receptor 1; complement receptor 2; Factor B; DAF; and others.

[0368] As will be recognized by the skilled artisan, many active moieties which may be used in the present invention occur in nature as secreted proteins in conjunction with a signal or leader peptide and/or as a pro-peptide which undergoes further intra- or extra-cellular processing. In such cases, the hybrid vectors of the

present invention may include one or more DNA sequences encoding such signal or leader peptides and/or one or more DNA sequences encoding such propeptide sequence, depending upon whether such secretion and/or processing is desired.

Alternatively, the hybrid vectors of the present disclosure may include DNA sequences encoding a different signal or leader peptide and/or pro-peptide sequence chosen to optimize the expression and localization of the targeting construct. In most cases, the signal peptide may be omitted, as the targeting moiety will supply sufficient information for targeting of the active moiety to the desired tissue and cells within the subject's body.

[0369] In some embodiments, a targeting construct described herein can be expressed in, and purified from, transgenic animals (e.g., transgenic mammals). For example, a targeting construct described herein can be produced in transgenic non-human mammals (e.g., rodents, sheep or goats) and isolated from milk as described in, e.g., Houdebine (2002) *Curr Opin Biotechnol* 13(6):625-629; van Kuik-Romeijn et al. (2000) *Transgenic Res* 9(2): 155-159; and Pollock et al. (1999) *J Immunol Methods* 231(1-2):147-157. Additional methods for producing proteins in mammalian milk products are described in, e.g., U.S. patent application publication nos. 200600105347 and 20040006776 and U.S. patent no. 7,045,676.

[0370] The targeting constructs described herein can be produced from cells by culturing a host cell transformed with the expression vector containing nucleic acid encoding the antibodies, under conditions, and for an amount of time, sufficient to allow expression of the proteins. Such conditions for protein expression will vary with the choice of the expression vector and the host cell, and will be easily ascertained by one skilled in the art through routine experimentation. For example, polypeptides expressed in *E. coli* can be refolded from inclusion bodies (see, e.g., Hou et al. (1998) *Cytokine* 10:319-30). Bacterial expression systems and methods for their use are well known in the art (see *Current Protocols in Molecular Biology*, Wiley & Sons, and *Molecular Cloning--A Laboratory Manual--3rd Ed.*, Cold Spring Harbor Laboratory Press, New York (2001)). The choice of codons, suitable expression vectors and suitable host cells will vary depending on a number of factors, and may be easily optimized as needed. A targeting construct described herein can be expressed in mammalian cells or in other expression systems including but not limited to yeast, baculovirus, and in vitro expression systems (see, e.g., Kaszubska et al. (2000) *Protein Expression and Purification* 18:213-220).

[0371] Following expression, the targeting construct can be isolated. The term “purified” or “isolated” as applied to any of the proteins described herein (e.g., a targeting construct, a targeting moiety, and/or an active moiety) refers to a polypeptide that has been separated or purified from components (e.g., proteins or other naturally-occurring biological or organic molecules) which naturally accompany it, e.g., other proteins, lipids, and nucleic acid in a prokaryote expressing the proteins. Typically, a polypeptide is purified when it constitutes at least 60 (e.g., at least 65, 70, 75, 80, 85, 90, 92, 95, 97, or 99) %, by weight, of the total protein in a sample.

[0372] A targeting construct described herein can be isolated or purified in a variety of ways known to those skilled in the art depending on what other components are present in the sample. Standard purification methods include electrophoretic, molecular, immunological, and chromatographic techniques, including ion exchange, hydrophobic, affinity, and reverse-phase HPLC chromatography. For example, a targeting construct can be purified using a standard anti-targeting construct antibody affinity column. Ultrafiltration and diafiltration techniques, in conjunction with protein concentration, are also useful. See, e.g., Scopes (1994) “Protein Purification, 3rd edition,” Springer-Verlag, New York City, New York. The degree of purification necessary will vary depending on the desired use. In some instances, no purification of the expressed polypeptide thereof will be necessary.

[0373] Methods for determining the yield or purity of a purified polypeptide are known in the art and include, e.g., Bradford assay, UV spectroscopy, Biuret protein assay, Lowry protein assay, amido black protein assay, high pressure liquid chromatography (HPLC), mass spectrometry (MS), and gel electrophoretic methods (e.g., using a protein stain such as Coomassie Blue or colloidal silver stain).

[0374] In some embodiments, a targeting construct described herein can be synthesized de novo in whole or in part, using chemical methods well known in the art. For example, the component amino acid sequences can be synthesized by solid phase techniques, cleaved from the resin, and purified by preparative high performance liquid chromatography followed by chemical linkage to form a desired polypeptide. The composition of the synthetic peptides may be confirmed by amino acid analysis or sequencing.

[0375] Once expressed and/or purified, a targeting construct described herein can be assayed for any one of a number of desired properties using in vitro or in vivo assays such as any of those described herein. For example, a targeting construct

described herein can be assayed for its ability to inhibit complement activity as described in.

[0376] In some embodiments, endotoxin can be removed from the targeting construct preparations. Methods for removing endotoxin from a protein sample are known in the art. For example, endotoxin can be removed from a protein sample using a variety of commercially available reagents including, without limitation, the ProteoSpin™ Endotoxin Removal Kits (Norgen Biotek Corporation), Detoxi-Gel Endotoxin Removal Gel (Thermo Scientific; Pierce Protein Research Products), MiraCLEAN® Endotoxin Removal Kit (Mirus), or Acrodisc™- Mustang® E membrane (Pall Corporation).

[0377] Methods for detecting and/or measuring the amount of endotoxin present in a sample (both before and after purification) are known in the art and commercial kits are available. For example, the concentration of endotoxin in a protein sample can be determined using the QCL-1000 Chromogenic kit (BioWhittaker), the limulus amoebocyte lysate (LAL)-based kits such as the Pyrotell®, Pyrotell®-T, Pyrochrome®, Chromo-LAL, and CSE kits available from the Associates of Cape Cod Incorporated.

[0378] Following expression and purification, the targeting constructs described herein can be modified. The modifications can be covalent or non-covalent modifications. Such modifications can be introduced into the targeting constructs by, e.g., reacting targeted amino acid residues in the targeting moiety and/or the active moiety with an organic derivatizing agent that is capable of reacting with selected side chains or terminal residues. Suitable sites for modification can be chosen using any of a variety of criteria including, e.g., structural analysis or amino acid sequence analysis of the targeting constructs described herein.

[0379] In some embodiments, a targeting construct described herein can be conjugated to a heterologous moiety. In embodiments where the heterologous moiety is a polypeptide, a targeting construct and a corresponding heterologous moiety described herein can be joined by way of fusion protein. The heterologous moiety can be, e.g., a heterologous polypeptide, a therapeutic agent (e.g., a toxin or a drug), or a detectable label such as, but not limited to, a radioactive label, an enzymatic label, a fluorescent label, or a luminescent label. Suitable heterologous polypeptides include, e.g., an antigenic tag (e.g., FLAG, polyhistidine, hemagglutinin (HA), glutathione-S-transferase (GST), or maltose-binding protein (MBP)) for use in

purifying the targeting constructs. Heterologous polypeptides also include polypeptides that are useful as diagnostic or detectable markers, for example, luciferase, green fluorescent protein (GFP), or chloramphenicol acetyl transferase (CAT). Where the heterologous moiety is a polypeptide, the moiety can be incorporated into a fusion protein described herein, resulting in a fusion protein.

[0380] In some embodiments, a targeting construct described herein can be conjugated to a detectable moiety. In embodiments where the detectable moiety is a polypeptide (e.g., GFP), a targeting moiety and a corresponding detectable moiety described herein can be joined by way of fusion protein. The detectable moiety can be, e.g., a heterologous polypeptide, a therapeutic agent (e.g., a toxin or a drug), or a detectable label such as, but not limited to, a radioactive label, an enzymatic label, a fluorescent label, or a luminescent label. Suitable heterologous polypeptides include, e.g., an antigenic tag (e.g., FLAG, polyhistidine, hemagglutinin (HA), glutathione-S-transferase (GST), or maltose-binding protein (MBP)) for use in purifying the targeting constructs. Heterologous polypeptides also include polypeptides that are useful as diagnostic or detectable markers, for example, luciferase, green fluorescent protein (GFP), or chloramphenicol acetyl transferase (CAT).

[0381] Conjugates

[0382] In some embodiments, the fusion molecules described herein are created by linkage of two independently produced polypeptide fragments, e.g., an antibody (e.g., a Fab fragment of a B4 or C2 antibody) and a complement modulator polypeptide (e.g., a soluble form of CD59). In certain embodiments, the targeting moiety is conjugated to the active moiety through a lysine, cysteine, glutamate, aspartate, or arginine amino acid. A targeting moiety can be conjugated to an active moiety through, e.g., a reaction comprising a thiolated targeting moiety, and a maleoyl-activated amine of the active moiety; an EDC/NHS-activated targeting moiety, and an amine of the active moiety; or an EDC/NHS-activated carboxylic acid of the active moiety and an amine of the targeting moiety. Two proteins (e.g., a targeting construct described herein and a heterologous moiety or the two constituent parts of a targeting construct) can, in some embodiments, be chemically cross-linked using any of a number of known chemical cross linkers. Examples of such cross linkers are those which link two amino acid residues via a linkage that includes a “hindered” disulfide bond. In these linkages, a disulfide bond within the cross-linking unit is protected (by

hindering groups on either side of the disulfide bond) from reduction by the action, for example, of reduced glutathione or the enzyme disulfide reductase. One suitable reagent, 4-succinimidylloxycarbonyl-methyl-2-pyridyldithio toluene (SMPT), forms such a linkage between two proteins utilizing a terminal lysine on one of the proteins and a terminal cysteine on the other. Heterobifunctional reagents that cross-link by a different coupling moiety on each protein can also be used. Other useful cross-linkers include, without limitation, reagents which link two amino groups (e.g., N-5-azido-2-nitrobenzoyloxysuccinimide), two sulfhydryl groups (e.g., 1,4-bis-maleimidobutane), an amino group and a sulfhydryl group (e.g., m-maleimidobenzoyl-N-hydroxysuccinimide ester), an amino group and a carboxyl group (e.g., 4-[pazidosalicylamido]butylamine), and an amino group and a guanidinium group that is present in the side chain of arginine (e.g., p-azidophenyl glyoxal monohydrate).

[0383] In some embodiments, a fusion protein described herein can contain a heterologous moiety which is chemically linked to the fusion protein. For example, in some embodiments, a drug described herein, a fluorescent label, a paramagnetic label, a radioactive label, etc., can be directly conjugated to the amino acid backbone of the targeting construct and/or targeting moiety (e.g., for use of the labeled targeting construct for in vivo imaging studies).

[0384] In some embodiments, the targeting constructs can be modified, e.g., with a moiety that improves the stabilization and/or retention of the targeting constructs in circulation, e.g., in blood, serum, or other tissues. For example, a targeting construct described herein can be PEGylated as described in, e.g., Lee et al. (1999) *Bioconjug Chem* 10(6): 973-8; Kinstler et al. (2002) *Advanced Drug Deliveries Reviews* 54:477-485; and Roberts et al. (2002) *Advanced Drug Delivery Reviews* 54:459-476. The stabilization moiety can improve the stability, or retention of, targeting construct by at least 1.5 (e.g., at least 2, 5, 10, 15, 20, 25, 30, 40, or 50 or more) fold.

[0385] In some embodiments, the targeting constructs described herein can be glycosylated. In some embodiments, a targeting construct described herein can be subjected to enzymatic or chemical treatment, or produced from a cell, such that the targeting construct, targeting moiety, and/or active moiety has reduced or absent glycosylation. Methods for producing polypeptides with reduced glycosylation are known in the art and described in, e.g., U.S. patent no. 6,933,368; Wright et al. (1991) *EMBO J* 10(10):2717-2723; and Co et al. (1993) *Mol Immunol* 30:1361-1367.

[0386] Pharmaceutical compositions

[0387] Also provided herein are pharmaceutical compositions comprising an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) described herein and a pharmaceutically acceptable carrier. The pharmaceutical compositions may be suitable for a variety of modes of administration described herein, including for example systemic or localized administration. The pharmaceutical compositions can be in the form of eye drops, injectable solutions, or in a form suitable for inhalation (either through the mouth or the nose) or oral administration. The pharmaceutical compositions described herein can be packaged in single unit dosages or in multidosage forms.

[0388] In some embodiments, the pharmaceutical compositions comprise an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) described herein and a pharmaceutically acceptable carrier suitable for administration to human. In some embodiments, the pharmaceutical compositions comprise a targeting construct and a pharmaceutically acceptable carrier suitable for intraocular injection. In some embodiments, the pharmaceutical compositions comprise a targeting construct and a pharmaceutically acceptable carrier suitable for topical application to the eye. In some embodiments, the pharmaceutical compositions comprise a targeting construct and a pharmaceutically acceptable carrier suitable for intravenous injection. In some embodiments, the pharmaceutical compositions comprise a targeting construct and a pharmaceutically acceptable carrier suitable for injection into the arteries (such as renal arteries).

[0389] The compositions are generally formulated as sterile, substantially isotonic, and in full compliance with all Good Manufacturing Practice (GMP) regulations of the U.S. Food and Drug Administration. In some embodiments, the composition is free of pathogen. For injection, the pharmaceutical composition can be in the form of liquid solutions, for example in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the targeting construct pharmaceutical composition can be in a solid form and redissolved or suspended immediately prior to use. Lyophilized compositions are also included.

[0390] For oral administration, the pharmaceutical compositions can take the form of, for example, tablets or capsules prepared by conventional means with pharmaceutically acceptable excipients such as binding agents (e.g., pregelatinized

maize starch, polyvinylpyrrolidone or hydroxypropyl methylcellulose); fillers (e.g., lactose, microcrystalline cellulose or calcium hydrogen phosphate); lubricants (e.g., magnesium stearate, talc or silica); disintegrants (e.g., potato starch or sodium starch glycolate); or wetting agents (e.g., sodium lauryl sulfate). Liquid preparations for oral administration can take the form of, for example, solutions, syrups or suspensions, or they can be presented as a dry product for constitution with water or other suitable vehicle before use. Such liquid preparations can be prepared by conventional means with pharmaceutically acceptable additives such as suspending agents (e.g., sorbitol syrup, cellulose derivatives or hydrogenated edible fats); emulsifying agents (e.g., lecithin or acacia); non-aqueous vehicles (e.g., oil, oily esters, ethyl alcohol or fractionated vegetable oils); and preservatives (e.g., methyl or propyl-p-hydroxybenzoates or sorbic acid). The preparations can also contain buffer salts, flavoring, coloring and sweetening agents as appropriate.

[0391] The present invention in some embodiments provides compositions comprising an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) and a pharmaceutically acceptable carrier suitable for administration to the eye. Such pharmaceutical carriers can be sterile liquids, such as water and oil, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, and the like. Saline solutions and aqueous dextrose, polyethylene glycol (PEG) and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, sodium state, glycerol monostearate, glycerol, propylene, water, and the like. The pharmaceutical composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. The targeting construct and other components of the composition may be encased in polymers or fibrin glues to provide controlled release of the targeting construct. These compositions can take the form of solutions, suspensions, emulsions, ointment, gel, or other solid or semisolid compositions, and the like. The compositions typically have a pH in the range of 4.5 to 8.0. The compositions must also be formulated to have osmotic values that are compatible with the aqueous humor of the eye and ophthalmic tissues. Such osmotic values will generally be in the range of from about 200 to about 400 milliosmoles per kilogram of water ("mOsm/kg"), but will preferably be about 300 mOsm/kg. The retina is considered to have an osmotic value of ~283 mOsm/kg.

[0392] In some embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for injection intravenously, intraperitoneally, or intravitreally. Typically, compositions for injection are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

[0393] The compositions may further comprise additional ingredients, for example preservatives, buffers, tonicity agents, antioxidants and stabilizers, nonionic wetting or clarifying agents, viscosity-increasing agents, and the like.

[0394] Suitable preservatives for use in a solution include polyquaternium-1, benzalkonium chloride, thimerosal, chlorobutanol, methyl paraben, propyl paraben, phenylethyl alcohol, edetate disodium, sorbic acid, benzethonium chloride, and the like. Typically (but not necessarily) such preservatives are employed at a level of from 0.001% to 1.0% by weight.

[0395] Suitable buffers include boric acid, sodium and potassium bicarbonate, sodium and potassium borates, sodium and potassium carbonate, sodium acetate, sodium biphosphate and the like, in amounts sufficient to maintain the pH at between about pH 6 and pH 8, and preferably, between about pH 7 and pH 7.5.

[0396] Suitable tonicity agents are dextran 40, dextran 70, dextrose, glycerin, potassium chloride, propylene glycol, sodium chloride, and the like, such that the sodium chloride equivalent of the ophthalmic solution is in the range 0.9 plus or minus 0.2%.

[0397] Suitable antioxidants and stabilizers include sodium bisulfite, sodium metabisulfite, sodium thiosulfite, thiourea and the like. Suitable wetting and clarifying agents include polysorbate 80, polysorbate 20, poloxamer 282 and tyloxapol. Suitable viscosity-increasing agents include dextran 40, dextran 70, gelatin, glycerin, hydroxyethylcellulose, hydroxymethylpropylcellulose, lanolin,

methylcellulose, petrolatum, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose and the like.

[0398] The use of viscosity enhancing agents to provide topical compositions with viscosities greater than the viscosity of simple aqueous solutions may be desirable to increase ocular absorption of the active compounds by the target tissues or increase the retention time in the eye. Such viscosity building agents include, for example, polyvinyl alcohol, polyvinyl pyrrolidone, methyl cellulose, hydroxy propyl methylcellulose, hydroxyethyl cellulose, carboxymethyl cellulose, hydroxy propyl cellulose or other agents known to those skilled in the art. Such agents are typically employed at a level of from 0.01% to 2% by weight.

[0399] In some embodiments, there is provided a pharmaceutical composition for delivery of a nucleotide encoding a targeting construct. The pharmaceutical composition for gene therapy can be in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle or compound is imbedded. Alternatively, where the complete gene delivery system can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical composition can comprise one or more cells which produce the gene delivery system.

[0400] In clinical settings, a gene delivery system for a gene therapeutic can be introduced into a subject by any of a number of methods. For instance, a pharmaceutical composition of the gene delivery system can be introduced systemically, e.g., by intravenous injection, and specific transduction of the protein in the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the receptor gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter, See U.S. Pat. 5,328,470, or by stereotactic injection, Chen et al. (1994), Proc. Natl. Acad. Sci., USA 91: 3054-3057. A polynucleotide encoding a targeting construct can be delivered in a gene therapy construct by electroporation using techniques described, Dev et al. (1994), Cancer Treat. Rev. 20:105-115.

[0401] In some embodiments, there is provided a pharmaceutical composition for gene delivery to the eye. Ophthalmic solutions useful for storing and/or delivering expression vectors have been disclosed, for example, in WO03077796A2.

[0402] Diseases to be treated

[0403] The treatment and diagnosis methods described herein can be used for treating or diagnosing a variety of diseases, including, but not limited to, inflammatory diseases, transplant rejections, pregnancy-related diseases, adverse drug reactions, tissue damage resulting from ischemia-reperfusion injury, ocular diseases, kidney diseases, joint diseases, and autoimmune or immune complex disorders. In some embodiments, the disease to be treated or diagnosed include, but not limited to, systemic lupus erythematosus and glomerulonephritis, rheumatoid arthritis, cardiopulmonary bypass and hemodialysis, hyperacute rejection in organ transplantation, myocardial infarction, ischemia/reperfusion injury, antibody-mediated allograft rejection, for example, in the kidneys, and adult respiratory distress syndrome. Moreover, other inflammatory conditions and autoimmune/immune complex diseases are also closely associated with complement activation, including, but not limited to, thermal injury, severe asthma, anaphylactic shock, bowel inflammation, urticaria, angioedema, vasculitis, multiple sclerosis, myasthenia gravis, myocarditis, membranoproliferative glomerulonephritis, atypical hemolytic uremic syndrome, Sjogren's syndrome, renal and pulmonary ischemia/reperfusion, and other organ-specific inflammatory disorders. Accordingly, in some embodiments, the methods described herein are particularly useful for treating or diagnosing a complement-mediated disease including, but not limited to, inflammatory disease, a transplant rejection, pregnancy-related disease, adverse drug reaction, tissue damage resulting from ischemia-reperfusion injury, ocular disease, kidney disease, joint disease, or an autoimmune or immune complex disorder. In some embodiments, also provided herein are methods of treating or diagnosing a complement-mediated disease in an individual, comprising administering to the individual an effective amount of any of the compositions (such as composition comprising a targeting construct) described herein.

[0404] The methods described herein are particularly useful for treating or diagnosing inflammatory diseases including, but not limited to, burns, endotoxemia, septic shock, adult respiratory distress syndrome, cardiopulmonary bypass, hemodialysis, anaphylactic shock, asthma, angioedema, Crohn's disease, sickle cell anemia, poststreptococcal glomerulonephritis, membranous nephritis, pancreatitis, rheumatoid arthritis, inflammatory arthritis, inflammatory bowel disease, acute lung injury, and disseminated intravascular coagulation (DIC). In some embodiments, the

inflammation (such as complement mediated inflammation) is associated with tissue damage resulting from inflammatory or autoinflammatory disorders, transplant rejection (cellular or antibody mediated), pregnancy-related diseases, adverse drug reactions, degenerative, neovascular, hemolytic, thrombotic, vasculitic, arthritic, regenerative, traumatic, autoimmune or immune complex disorders.

[0405] The compositions described herein are also useful for treating or diagnosing a transplant rejection including, but not limited to, a hyperacute transplant rejection, antibody-mediated transplant rejection, cellular-mediated transplant rejection, acute transplant rejection, and chronic transplant rejection. In some embodiments, the transplant is a xenograft, an allograft, or an isograft. In some embodiments, the transplant is a fluid, a cell, a tissue or an organ. In some embodiments, the transplant is selected from the group consisting of: a heart, liver, kidney, lung, pancreas, intestine, stomach, testis, hand, arm, leg, uterus, ovary, and thymus. In some embodiments, the transplant is selected from the group consisting of: a bone, tendons, cornea, skin, heart valve, islets of Langerhans, bone marrow, hematopoietic stem cell, blood transfusion, or vein. In some embodiments, the transplant is a heart, liver or kidney. Transplant rejections can result in several complications such as graft-versus-host disease. In some embodiments, a complement-mediated disease is graft-versus-host disease.

[0406] The methods described herein are also particularly useful for treating or diagnosing a pregnancy-related disease including, but not limited to, HELLP (Hemolytic anemia, elevated liver enzymes, and low platelet count), recurrent fetal loss, atypical hemolytic uremic syndrome, fetal hypoxia syndrome, hypertensive disease, and pre-eclampsia.

[0407] In addition, the methods described herein are useful for treating or diagnosing an adverse drug reaction including, but not limited to, a drug allergy, a radiographic contrast media allergy, and IL-2 induced vascular leakage.

[0408] The methods described herein are also useful for treating or diagnosing tissue damage resulting from ischemia-reperfusion injury following, but not limited to, acute myocardial infarction, aneurysm, aneurysm repair, deep hypothermic circulatory arrest, tourniquet use, solid organ transplant, stroke including perinatal stroke, hemorrhagic shock, crush injury, multiple organ failure, hemodialysis, hypovolemic shock, spinal cord injury, traumatic brain injury, intestinal ischemia, retinal ischemia, cardiopulmonary bypass, emergency coronary surgery for failed

percutaneous transluminal coronary angioplasty (PCTA), and any vascular surgery with blood vessel cross clamping, pancreatitis after manipulation of pancreatic or bile duct. In some embodiments, tissue damage can be treated before, during, or after the ischemic event (such as intestinal ischemia) that triggers ischemia-reperfusion injury.

[0409] In some embodiments, tissue damage is treated or diagnosed with any of the methods disclosed herein by administering a targeting construct (or a composition comprising the targeting construct) disclosed herein before reperfusion. In some embodiments, tissue damage is treated or diagnosed with any of the methods disclosed herein by administering a targeting construct (or a composition comprising the targeting construct) disclosed herein after reperfusion. In some embodiments, the ischemia-reperfusion injury is selected from the group consisting of: myocardial ischemia-reperfusion, renal ischemia-reperfusion injury, gastrointestinal ischemia-reperfusion injury, hepatic ischemia-reperfusion injury, skeletal muscle ischemia-reperfusion injury, cerebral ischemia-reperfusion injury, pulmonary ischemia-reperfusion injury, intestine ischemia-reperfusion injury, retinal ischemia-reperfusion injury, and joint ischemia-reperfusion injury. In some embodiments, tissue damage is caused by oxidative damage.

[0410] There are instances when a therapy or surgery induces a reperfusion but not an ischemia (referred herein as non-ischemia reperfusion injury). Such therapy or surgery includes, but is not limited to, pharmacological thrombolysis, including intravenous and endovascular therapies for stroke, acute coronary syndromes, peripheral arterial occlusion, pulmonary embolus, renal artery occlusion, mechanical thrombolysis, e.g. percutaneous coronary intervention, peripheral arterial angioplasty, visceral arterial angioplasty, coronary artery bypass grafting, carotid endarterectomy, mesenteric ischemia, shock including hemorrhagic, cardiogenic, neurogenic, anaphylactic, flap-failure, e.g. plastic surgery, re-implantation of digits and limbs, and strangulated bowel. Accordingly, in some embodiments, tissue damage resulting from non-ischemia reperfusion injury is treated or diagnosed with any of the methods disclosed herein by administering a targeting construct (or a composition comprising the targeting construct) disclosed herein.

[0411] The methods described herein are also particularly useful for treating or diagnosing a kidney disease including, but not limited to, acute kidney injury, hemolytic uremic syndrome, glomerulonephritis, membranous glomerulonephritis, mesangioproliferative glomerulonephritis, acute postinfectious glomerulonephritis

(such as poststreptococcal glomerulonephritis), cryoglobulinemic glomerulonephritis, lupus nephritis, membranoproliferative glomerulonephritis (such as mesangiocapillary glomerulonephritis), dense deposit disease, minimal change disease, diabetic nephropathy, Henoch-Schonlein purpura nephritis, IgA nephropathy, chronic kidney disease, delayed graft function of a kidney transplant, acute and chronic renal transplant rejection, proteinuric renal disease and nephrotic syndrome, hypertensive kidney disease, and focal segmental glomerulosclerosis. In some embodiments, the kidney disease is a glomerular disease. For example, the methods are useful for treating or diagnosing glomerular disease that leads to binding of natural IgM to damaged glomerulus. In some embodiments, damaged glomerulus can be a result of mechanical, metabolic, chemical, oxidative or immunologic stress. In some embodiments, damaged glomerulus can be a result of ischemia, diabetes, hypertension, and secondary focal segmental glomerulosclerosis. Symptoms of damaged glomerulus include an inflammatory response such as cytokine release and fibrosis such as collagen mesangial matrix deposition, tubular cell damage, and tubulointerstitial fibrosis. The methods are also useful for treating or diagnosing kidney disease such a glomerulonephritis which is inflammation of the glomerulus. Glomerulonephritis is commonly associated with deposition of electron dense material in the glomerulus which contains complement components, including C3. The methods are also useful for treating or diagnosing acute kidney injury associated with renal ischemia. Ischemia is the leading cause of acute kidney injury. Ischemia and subsequent reperfusion elicit acute kidney injury through endothelial dysfunction, leukocyte-mediated inflammation and decreased microvascular blood flow that can lead to rarefaction of the peritubular capillaries, shifting the fragile balance of oxygen supply and demand to the corticomedullary junction toward a negative oxygen balance. The shift in balance causes a hypoxic environment and can lead to accumulation of fibrosis and subsequent development of chronic kidney disease. In some embodiments, the kidney disease is due to a factor H deficiency.

[0412] The methods described herein are also useful for treating or diagnosing a joint disease including, but not limited to, arthritis (such as rheumatoid arthritis) and joint inflammation associated with infection (such as hepatitis B infection), inflammatory disease (such as inflammatory bowel disease) or autoimmune disease (such as systemic lupus erythematosus). In some embodiments, methods provided herein are useful for treating or diagnosing a joint disease including, but not limited

to, arthritis, amyloid arthropathy, amyloidosis, ankylosing spondylitis, carpal tunnel syndrome, temporal arteritis, polymyalgia rheumatica, polyarthralgia, tendinitis, Whipple's disease, bursitis, trigeminal neuralgia, fibromyoma, fibrositis, autoimmune arthritis, rheumatoid arthritis, juvenile arthritis, psoriatic arthritis, lupus arthritis, polyarthritis, inflammatory arthritis not resulting from an autoimmune disease or disorder, such as an infectious arthritis, i.e., joint pain, soreness, stiffness and swelling caused by an infectious agent such as bacteria (including mycoplasma), viruses, fungi, septic arthritis, or osteoarthritis. Joint disease can be associated with symptoms such as joint stiffness, pain, weakness, joint fatigue, tenderness and swelling. Accordingly, in some embodiments, symptoms of joint disease can be treated or diagnosed with any of the methods disclosed herein by administering a targeting construct (or a composition comprising the targeting construct) disclosed herein. For example, the compositions are useful for treating or diagnosing arthritis or symptoms of arthritis. In some embodiments, the arthritis is selected from the group consisting of: rheumatoid arthritis, juvenile onset rheumatoid arthritis, psoriatic arthritis, and lupus arthritis. In some embodiments the arthritis is osteoarthritis. In some embodiments, the arthritis is infectious arthritis caused by a bacterial pathogen, such as *Haemophilus influenzae*, *Gonococcus* spp., *Mycoplasma* spp. *Meingococcus* spp., *Pneumococcus* spp., *Streptococcus* spp., *Staphylococcus* spp., *Salmonella* spp., *Brucella* spp., *Neisseria* spp., *Streptobacillus moniliformis* (Haverhill fever), *Mycobacterium tuberculosis*, *Treponema pallidum* (syphilis), *Treponema pertenue* (yaws), or *Rickettsia* spp. In some embodiments, the arthritis is infectious arthritis caused by a viral pathogen, such as a rubella virus, a mumps virus, a varicella-zoster virus, an adenovirus, an echovirus, a herpes simplex virus, a cytomegalovirus, a parvovirus, a retrovirus, and alphavirus, or a hepatitis virus. In some embodiments, the arthritis is infectious arthritis caused by a fungus, such as *Coccidioides* spp., *Histoplasma* spp., *Blastomyces* spp., *Cryptococcus* spp., *Candida* spp., or *Sporothrix* spp. As another example, the compositions are useful for treating or diagnosing a joint disease or symptoms of a joint disease. In some embodiments, the joint disease is arthritis, amyloid arthropathy, amyloidosis, ankylosing spondylitis, carpal tunnel syndrome, temporal arteritis, polymyalgia rheumatica, polyarthralgia, tendinitis, Whipple's disease, bursitis, trigeminal neuralgia, fibromyoma, and fibrositis. In some embodiments, the joint disease is associated with arthritis. In some embodiments, the

joint disease precedes the development of arthritis. In some embodiments, the joint disease develops due to the onset of arthritis.

[0413] Rheumatoid arthritis affects approximately 1% of the population, with women affected three times more commonly than men. Rheumatoid arthritis and juvenile onset rheumatoid arthritis are systemic diseases with numerous pathologic manifestations in addition to their joint inflammatory aspects. In rheumatoid arthritis, these manifestations include vasculitis (inflammation of the blood vessels), which can affect nearly any organ system and can cause numerous pathologic sequelae including polyneuropathy, cutaneous ulceration, and visceral infarction. Pleuropulmonary manifestations include pleuritis, interstitial fibrosis, pleuropulmonary nodules, pneumonitis, and arteritis. Other manifestations include the development of inflammatory rheumatoid nodules on a variety of periarticular structures such as extensor surfaces, as well as on pleura and meninges. Weakness and atrophy of skeletal muscle are common. Many patients with systemic lupus erythematosus also develop joint inflammation referred to as lupus arthritis. Systemic lupus erythematosus is an autoimmune disease of unknown cause in which numerous different cells, tissues, and organs are damaged by pathogenic autoantibodies and immune complexes. Clinical manifestations of systemic lupus erythematosus are numerous and include a variety of maculopapular rashes, nephritis, cerebritis, vasculitis, hematologic abnormalities including cytopenias and coagulopathies, pericarditis, myocarditis, pleurisy, gastrointestinal symptoms, and the aforementioned joint inflammation. Osteoarthritis represents the most common chronic joint disease. It is manifested by pain, stiffness, and swelling of the involved joints. Articular cartilage, responsible for the most critical mechanical functions of the joint, is the major target tissue of osteoarthritis and the breakdown of articular cartilage in osteoarthritis is mediated by various enzymes such as metalloproteinases, plasmin, and cathepsin, which are in turn stimulated by various factors that can also act as inflammatory mediators. These factors include cytokines such as interleukin-1, which is known to activate the pathogenic cartilage and synovial proteases. Synovial inflammation becomes more frequent as the disease progresses. Psoriatic arthritis is a chronic inflammatory joint disorder that affects 5 to 8% of people with psoriasis. A significant percentage of these individuals (one- fourth) develop progressive destructive disease. Twenty five percent of psoriasis patients with joint inflammation develop symmetric

joint inflammation resembling the joint inflammation manifestations of rheumatoid arthritis, and over half of these go on to develop varying degrees of joint destruction.

[0414] The methods described herein are useful for treating or diagnosing an autoimmune or immune complex including, but not limited to, but is not limited to, myasthenia gravis, Alzheimer's disease, multiple sclerosis, emphysema, obesity, neuromyelitis optica, rheumatoid arthritis, osteoarthritis, systemic lupus erythematosus, lupus nephritis, IgG4 associated diseases, insulin-dependent diabetes mellitus, acute disseminated encephalomyelitis, Addison's disease, antiphospholipid antibody syndrome, thrombotic thrombocytopenic purpura, autoimmune hepatitis, Crohn's disease, Goodpasture's syndromes, Graves' disease, Guillain-Barre syndrome, Hashimoto's disease, idiopathic thrombocytopenic purpura, pemphigus, Sjogren's syndrome, Takayasu's arteritis, autoimmune glomerulonephritis, dense deposit disease (also known as membranoproliferative glomerulonephritis type II), membranous disease, paroxysmal nocturnal hemoglobinuria, age-related macular degeneration, diabetic maculopathy, uveitis, retinal degeneration disorders, diabetic nephropathy, focal segmental glomerulosclerosis, ANCA associated vasculitis, hemolytic uremic syndrome, Shiga-toxin-associated hemolytic uremic syndrome, atypical hemolytic uremic syndrome, and inflammation associated cardiopulmonary bypass and hemodialysis. In some embodiments, the disease to be treated or diagnosed is an autoimmune glomerulonephritis, which includes, but is not limited to, immunoglobulin A nephropathy or membranoproliferative glomerulonephritis type I. In some embodiments, an autoimmune or immune complex disorder is an inflammatory disease.

[0415] The methods described herein are particularly useful for treating or diagnosing ocular diseases including, but not limited to, age-related macular degeneration ("AMD"), including wet AMD and dry AMD, CMV retinitis, macular edema, uveitis, glaucoma, diabetic retinopathy, retinitis pigmentosa, retinal detachment, proliferative vitreoretinopathy and ocular melanoma. For example, the methods are useful for treating or diagnosing age-related macular degeneration (AMD). AMD is clinically characterized by progressive loss of central vision which occurs as a result of damage to the photoreceptor cells in an area of the retina called the macula. AMD has been broadly classified into two clinical states: a wet form and a dry form, with the dry form making up to 80-90% of total cases. The dry form is characterized clinically by the presence of macular drusen, which are localized

deposits between the retinal pigment epithelium (RPE) and the Bruch's membrane, and by geographic atrophy characterized by RPE cell death with overlying photoreceptor atrophy. Wet AMD, which accounts for approximately 90% of serious vision loss, is associated with neovascularization in the area of the macular and leakage of these new vessels. The accumulation of blood and fluid can cause retinal detachment followed by rapid photoreceptor degeneration and loss of vision. It is generally accepted that the wet form of AMD is preceded by and arises from the dry form.

[0416] Analysis of the contents of drusen in AMD patients has shown a large number of inflammatory proteins including amyloid proteins, coagulation factors, and a large number of proteins of the complement pathway. A genetic variation in the complement factor H substantially raises the risk of age-related macular degeneration (AMD), suggesting that uncontrolled complement activation underlies the pathogenesis of AMD. Edward et al., Science 2005, 308:421; Haines et al., Science 2005, 308:419; Klein et al., Science 308:385-389; Hageman et al., Proc. Natl. Acad. Sci. USA 2005, 102:7227.

[0417] In some embodiments, the methods described herein can be used to treat or diagnose cytomegalovirus (CMV) retinitis. CMV retinitis is an infection that causes inflammation of the photoreceptor cells in the retina. CMV is typically rare in immunocompetent individuals. However, individuals who are immunocompromised, e.g., by diseases, transplants, or chemotherapy, are particularly susceptible to CMV retinitis. Retinitis usually begins in one eye, but often progresses to the other eye. Without treatment, progressive damage to the retina can lead to blindness in 4-6 months or less.

[0418] In some embodiments, the methods described herein can be used to treat or diagnose macular edema. Macular edema occurs when fluid and protein deposits collect on or under the macula of the eye, causing it to thicken and swell. The swelling may distort an individual's central vision, as the macula holds tightly packed cones that provide sharp, clear central vision to enable a person to see detail, form, and color that is directly in the direction of gaze. Macular edema can be classified into two types. Cystoid macular edema (CME) involves fluid accumulation in the outer plexiform layer secondary to abnormal perifoveal retinal capillary permeability. Diabetic macular edema (DME) is similarly caused by leaking macular capillaries.

DME is the most common cause of visual loss in both proliferative, and non-proliferative diabetic retinopathy.

[0419] In certain embodiments, the methods described herein can be used to treat or diagnose uveitis, i.e., inflammation of the uvea (the iris, ciliary body, and choroid of the eye beneath the sclera). Uveitis is typically associated with eye infections, eye injuries, and/or autoimmune disorders. However, in many cases, the cause is unknown. The most common form of uveitis is anterior uveitis, which involves inflammation in iris. Posterior uveitis affects the choroid, a layer of blood vessels and connective tissue in the middle part of the eye. Another form of uveitis is pars planitis. This inflammation affects the narrowed area (pars plana) between the iris and the choroid.

[0420] In certain embodiments, the methods described herein can be used to treat or diagnose glaucoma, a group of eye conditions that lead to damage to the optic nerve, and loss of vision. The nerve damage involves loss of retinal ganglion cells in a characteristic pattern. The many different subtypes of glaucoma can all be considered to be a type of optic neuropathy. Raised intraocular pressure (above 21 mmHg or 2.8 kPa) is the most important and only modifiable risk factor for glaucoma. Intraocular pressure is a function of production of liquid aqueous humor by the ciliary processes of the eye, and its drainage through the trabecular meshwork. Aqueous humor flows from the ciliary processes into the posterior chamber, bounded posteriorly by the lens and the zonules of Zinn, and anteriorly by the iris. It then flows through the pupil of the iris into the anterior chamber, bounded posteriorly by the iris and anteriorly by the cornea. From here, the trabecular meshwork drains aqueous humor via Schlemm's canal into scleral plexuses and general blood circulation.

[0421] In open/wide-angle glaucoma, flow is reduced through the trabecular meshwork, due to the degeneration and obstruction of the trabecular meshwork, whose original function is to absorb the aqueous humor. Loss of aqueous humor absorption leads to increased resistance and thus a chronic, painless buildup of pressure in the eye. In close/narrow-angle, the iridocorneal angle is completely closed because of forward displacement of the final roll and root of the iris against the cornea, resulting in the inability of the aqueous fluid to flow from the posterior to the anterior chamber and then out of the trabecular network. This accumulation of aqueous humor causes an acute increase of pressure and pain.

[0422] In some embodiments, the methods described herein can be used to treat or diagnose diabetic retinopathy, a complication of diabetes that causes damage that results from microvascular retinal changes. Small blood vessels, such as those in the eye, are especially vulnerable to poor blood sugar control. An over accumulation of glucose and/or fructose damages the tiny blood vessels in the retina. Hyperglycemia-induced pericyte death and thickening of the basement membrane lead to increased permeability of the vascular walls, which changes the formation of the blood-retinal barrier. In some individuals, diabetic retinopathy is accompanied by macular edema. As diabetic retinopathy progresses, the lack of oxygen in the retina causes fragile, new, blood vessels to grow along the retina and in the vitreous humour. Without timely treatment, these new blood vessels can bleed, cloud vision, and destroy the retina and/or cause tractional retinal detachment.

[0423] In certain embodiments, the methods described herein can be used to treat or diagnose retinitis pigmentosa (RP), a group of inherited, degenerative eye diseases that cause severe vision impairment and blindness. Mutations in more than 60 genes are known to cause retinitis pigmentosa. Approximately 20% of RP is autosomal dominant (ADRP), 20% is autosomal recessive (ARRP), and 10% is X linked (XLRP), while the remaining 50% is found in patients without any known affected relatives. The genes associated with retinitis pigmentosa play essential roles in the structure and function of photoreceptors in the retina, and the progressive degeneration of these cells causes vision loss.

[0424] In certain embodiments, the methods described herein can be used to treat or diagnose proliferative vitreoretinopathy, i.e., the formation of scar tissue within the eye that is often a complication of rhegmatogenous retinal detachment. During rhegmatogenous retinal detachment, fluid from the vitreous humor enters a retinal hole. The accumulation of fluid in the subretinal space and the tractional force of the vitreous on the retina result in rhegmatogenous retinal detachment. During this process the retinal cell layers come in contact with vitreous cytokines, which trigger the proliferation and migration of retinal pigmented epithelium (RPE). The RPE cells undergo epithelial-mesenchymal transition (EMT) and develop the ability to migrate out into the vitreous. During this process the RPE cell layer-neural retinal adhesion and RPE-ECM (extracellular matrix) adhesions are lost. The RPE cells lay down fibrotic membranes while they migrate and these membranes contract and pull

at the retina, and this can lead to secondary retinal detachment after primary retinal detachment surgery.

[0425] In certain embodiments, the treatment methods described herein can be used in conjunction with, e.g., surgery for the repair of a retinal tear, hole or detachment, or with, e.g., radiation therapy for the treatment of ocular melanoma.

[0426] In certain embodiments, the compositions and methods described herein can be used to treat and/or improve the outcome of corneal wound healing and/or corneal transplantation. The corneal wound healing response is a complex cascade involving cytokine mediated interactions between the epithelial cells, stromal keratocytes, corneal nerves, lacrimal glands, tear film and cells of the immune system. The response of the tissue changes depends on the inciting injury. For example, incisional, lamellar and surface scrape injuries, like the ones used in keratorefractive surgery procedures, are followed by typical wound healing responses that are similar in some respects, but different in others. For example, elsewhere in the body, wound healing culminates in scar formation and vascularisation whereas one of the most crucial aspects of corneal wound healing is how the healing processes aim to minimize these end results, which would otherwise have serious visual consequences. Causes of corneal scarring include almost any disruption to normal corneal structure and function, whether from infection, laser refractive surgery, corneal transplantation, ocular trauma (chemical or physical) or corneal dystrophies.

[0427] Corneal transplantation, also known as corneal grafting, is a surgical procedure where a damaged or diseased cornea is replaced by donated corneal tissue (the graft) in its entirety (penetrating keratoplasty) or in part (lamellar keratoplasty). The graft is taken from a recently deceased individual with no known diseases or other factors that may affect the viability of the donated tissue or the health of the recipient. Since the cornea has no blood vessels (it takes its nutrients from the aqueous humor) it heals much more slowly than a cut on the skin. The risks are similar to other intraocular procedures, but additionally include graft rejection (lifelong), detachment or displacement of lamellar transplants and primary graft failure. There is also a risk of infection.

[0428] The present invention provides methods of treating or diagnosing an ocular disease described herein by administering an effective amount of a composition comprising a targeting construct. In some embodiments, the invention provides methods of treating or diagnosing one or more aspects or symptoms of the ocular

diseases described herein, including, but not limited to, formation of ocular drusen, inflammation in the eye or eye tissue, loss of photoreceptor cells, loss of vision (including for example visual acuity and visual field), neovascularization (such as choroidal neovascularization or CNV), and retinal detachment. Other related aspects, such as photoreceptor degeneration, RPE degeneration, retinal degeneration, chorioretinal degeneration, cone degeneration, retinal dysfunction, retinal damage in response to light exposure (such as constant light exposure), damage of the Bruch's membrane, loss of RPE function, gain or RPE function, loss of integrity of the histoarchitecture of the cells and/or extracellular matrix of the normal macular, loss of function of the cells in the macula, photoreceptor dystrophy, mucopolysaccharidoses, rod-cone dystrophies, cone-rod dystrophies, anterior and posterior uveitis, and diabetic neuropathy, are also included.

[0429] In some embodiments, there are provided methods of treating or diagnosing a drusen-associated disease. The term "drusen-associated disease" refers to any disease in which formation of drusen or drusen-like extracellular disease plaque takes place, and for which drusen or drusen-like extracellular disease plaque causes or contributes to thereto or represents a sign thereof. For example, AMD, characterized by the formation of macular drusen, is considered as a drusen-associated disease. Non-ocular drusen-related disease include, but are not limited to, amyloidosis, elastosis, dense deposit disease, and/or atherosclerosis.

Ocular diseases

[0430] The compositions and methods described herein are particularly useful for treating ocular diseases. For example, the compositions and methods can be useful in detecting and/or treating ischemia/reperfusion (I/R) injury to the eye. As used herein, the term "ischemia/reperfusion injury" refers to inflammatory injury to the endothelium and underlying parenchymal tissues following reperfusion of hypoxic tissues. It is a general syndrome that is responsible for both acute and chronic injury to various tissues including, for example, myocardium, central nervous system, hind limb, intestine, and eye. Ischemia reperfusion injury can result in necrosis and irreversible cell injury. The complement pathway (including the alternative complement pathway) is a major mediator of I/R injury. The noninvasive methods provided herein are thus useful for detection of complement-mediated inflammation associated with ischemia reperfusion that occurs in eye.

[0431] The compositions and methods provided herein may also be used to detect and/or treat complement-mediated inflammation in drusen-associated diseases. For example, age-related macular degeneration (AMD), characterized by the formation of macular drusen, is considered a drusen-associated disease. AMD is clinically characterized by progressive loss of central vision which occurs as a result of damage to the photoreceptor cells in an area of the retina called the macula. AMD has been broadly classified into two clinical states: a wet form and a dry form, with the dry form making up to 80-90% of total cases. The dry form is characterized clinically by the presence of macular drusen, which are localized deposits between the retinal pigment epithelium (RPE) and the Bruch's membrane, and by geographic atrophy characterized by RPE cell death with overlying photoreceptor atrophy. Wet AMD, which accounts for approximately 90% of serious vision loss, is associated with neovascularization in the area of the macular and leakage of these new vessels. The accumulation of blood and fluid can cause retinal detachment followed by rapid photoreceptor degeneration and loss of vision. It is generally accepted that the wet form of AMD is preceded by and arises from the dry form.

[0432] Analysis of the contents of drusen in AMD patients has shown a large number of inflammatory proteins including amyloid proteins, coagulation factors, and a large number of proteins of the complement pathway. A genetic variation in the complement factor H substantially raises the risk of age-related macular degeneration (AMD), suggesting that uncontrolled complement activation underlies the pathogenesis of AMD. Edward et al., *Science* 2005, 308:421; Haines et al., *Science* 2005, 308:419; Klein et al., *Science* 308:385-389; Hageman et al., *Proc. Natl. Acad. Sci. USA* 2005, 102:7227. In addition, lipid accumulation and modifications, as well as the presence of Annexin II, has been reported in pathological structures associated with AMD (*see, e.g.*, references cited above). Animal models of AMD respond favorably to complement therapeutics such as those described here (Rohrer, et al. (2009) *Invest Ophthalmol Vis Sci.* 50(7):3056-64; Rohrer, et al. (2012) *J Ocul Pharmacol Ther.* 28(4):402-9).

[0433] In some embodiments, the compositions and methods described herein can be used to detect and/or treat cytomegalovirus (CMV) retinitis. CMV retinitis is an infection that causes inflammation of the photoreceptor cells in the retina. CMV is typically rare in immunocompetent individuals. However, individuals who are immunocompromised, *e.g.*, by diseases, transplants, or chemotherapy, are particularly

susceptible to CMV retinitis. Retinitis usually begins in one eye, but often progresses to the other eye. Without treatment, progressive damage to the retina can lead to blindness in 4-6 months or less.

[0434] In some embodiments, the compositions and methods described herein can be used to detect and/or treat macular edema. Macular edema occurs when fluid and protein deposits collect on or under the macula of the eye, causing it to thicken and swell. The swelling may distort an individual's central vision, as the macula holds tightly packed cones that provide sharp, clear central vision to enable a person to see detail, form, and color that is directly in the direction of gaze. Macular edema can be classified into two types. Cystoid macular edema (CME) involves fluid accumulation in the outer plexiform layer secondary to abnormal perifoveal retinal capillary permeability. Diabetic macular edema (DME) is similarly caused by leaking macular capillaries. DME is the most common cause of visual loss in both proliferative, and non-proliferative diabetic retinopathy.

[0435] In certain embodiments, the compositions and methods described herein can be used to detect and/or treat uveitis, *i.e.*, inflammation of the uvea (the iris, ciliary body, and choroid of the eye beneath the sclera). Uveitis is typically associated with eye infections, eye injuries, and/or autoimmune disorders. However, in many cases, the cause is unknown. The most common form of uveitis is anterior uveitis, which involves inflammation in iris. Posterior uveitis affects the choroid, a layer of blood vessels and connective tissue in the middle part of the eye. Another form of uveitis is pars planitis. This inflammation affects the narrowed area (pars plana) between the iris and the choroid.

[0436] In certain embodiments, the compositions and methods described herein can be used to detect and/or treat glaucoma, a group of eye conditions that lead to damage to the optic nerve, and loss of vision. The nerve damage involves loss of retinal ganglion cells in a characteristic pattern. The many different subtypes of glaucoma can all be considered to be a type of optic neuropathy. Raised intraocular pressure (above 21 mmHg or 2.8 kPa) is the most important and only modifiable risk factor for glaucoma. Intraocular pressure is a function of production of liquid aqueous humor by the ciliary processes of the eye, and its drainage through the trabecular meshwork. Aqueous humor flows from the ciliary processes into the posterior chamber, bounded posteriorly by the lens and the zonules of Zinn, and anteriorly by the iris. It then flows through the pupil of the iris into the anterior chamber, bounded posteriorly by

the iris and anteriorly by the cornea. From here, the trabecular meshwork drains aqueous humor via Schlemm's canal into scleral plexuses and general blood circulation.

[0437] In open/wide-angle glaucoma, flow is reduced through the trabecular meshwork, due to the degeneration and obstruction of the trabecular meshwork, whose original function is to absorb the aqueous humor. Loss of aqueous humor absorption leads to increased resistance and thus a chronic, painless buildup of pressure in the eye. In close/narrow-angle, the iridocorneal angle is completely closed because of forward displacement of the final roll and root of the iris against the cornea, resulting in the inability of the aqueous fluid to flow from the posterior to the anterior chamber and then out of the trabecular network. This accumulation of aqueous humor causes an acute increase of pressure and pain.

[0438] In some embodiments, the compositions and methods described herein can be used to detect and/or treat diabetic retinopathy, a complication of diabetes that causes damage that results from microvascular retinal changes. Small blood vessels, such as those in the eye, are especially vulnerable to poor blood sugar control. An overaccumulation of glucose and/or fructose damages the tiny blood vessels in the retina. Hyperglycemia-induced pericyte death and thickening of the basement membrane lead to increased permeability of the vascular walls, which changes the formation of the blood-retinal barrier. In some individuals, diabetic retinopathy is accompanied by macular edema. As diabetic retinopathy progresses, the lack of oxygen in the retina causes fragile, new, blood vessels to grow along the retina and in the vitreous humour. Without timely treatment, these new blood vessels can bleed, cloud vision, and destroy the retina and/or cause tractional retinal detachment.

[0439] In certain embodiments, the compositions and methods described herein can be used to detect and/or treat retinitis pigmentosa (RP), a group of inherited, degenerative eye diseases that cause severe vision impairment and blindness. Mutations in more than 60 genes are known to cause retinitis pigmentosa. Approximately 20% of RP is autosomal dominant (ADRP), 20% is autosomal recessive (ARRP), and 10% is X linked (XLRP), while the remaining 50% is found in patients without any known affected relatives. The genes associated with retinitis pigmentosa play essential roles in the structure and function of photoreceptors in the retina, and the progressive degeneration of these cells causes vision loss.

[0440] In certain embodiments, the compositions and methods described herein can be used to detect and/or treat proliferative vitreoretinopathy, *i.e.*, the formation of scar tissue within the eye that is often a complication of rhegmatogenous retinal detachment. During rhegmatogenous retinal detachment, fluid from the vitreous humor enters a retinal hole. The accumulation of fluid in the subretinal space and the tractional force of the vitreous on the retina result in rhegmatogenous retinal detachment. During this process the retinal cell layers come in contact with vitreous cytokines, which trigger the proliferation and migration of retinal pigmented epithelium (RPE). The RPE cells undergo epithelial-mesenchymal transition (EMT) and develop the ability to migrate out into the vitreous. During this process the RPE cell layer-neural retinal adhesion and RPE-ECM (extracellular matrix) adhesions are lost. The RPE cells lay down fibrotic membranes while they migrate and these membranes contract and pull at the retina, and this can lead to secondary retinal detachment after primary retinal detachment surgery.

[0441] In certain embodiments, the compositions described herein can be used in conjunction with, *e.g.*, surgery for the repair of a retinal tear, hole or detachment, or with, *e.g.*, radiation therapy for the treatment of ocular melanoma.

[0442] In certain embodiments, the compositions and methods described herein can be used to treat and/or improve the outcome of corneal wound healing and/or corneal transplantation. The corneal wound healing response is a complex cascade involving cytokine mediated interactions between the epithelial cells, stromal keratocytes, corneal nerves, lacrimal glands, tear film and cells of the immune system. The response of the tissue changes depends on the inciting injury. For example, incisional, lamellar and surface scrape injuries, like the ones used in keratorefractive surgery procedures, are followed by typical wound healing responses that are similar in some respects, but different in others. For example, elsewhere in the body, wound healing culminates in scar formation and vascularisation whereas one of the most crucial aspects of corneal wound healing is how the healing processes aim to minimize these end results, which would otherwise have serious visual consequences. Causes of corneal scarring include almost any disruption to normal corneal structure and function, whether from infection, laser refractive surgery, corneal transplantation, ocular trauma (chemical or physical) or corneal dystrophies.

[0443] Corneal transplantation, also known as corneal grafting, is a surgical procedure where a damaged or diseased cornea is replaced by donated corneal tissue

(the graft) in its entirety (penetrating keratoplasty) or in part (lamellar keratoplasty). The graft is taken from a recently deceased individual with no known diseases or other factors that may affect the viability of the donated tissue or the health of the recipient. Since the cornea has no blood vessels (it takes its nutrients from the aqueous humor) it heals much more slowly than a cut on the skin. The risks are similar to other intraocular procedures, but additionally include graft rejection (lifelong), detachment or displacement of lamellar transplants and primary graft failure. There is also a risk of infection.

[0444] The present invention provides methods of detecting and/or treating an ocular disease described herein by administering an effective amount of a composition comprising a targeting construct. In some embodiments, the invention provides methods of treating or preventing one or more aspects or symptoms of the ocular diseases described herein, including, but not limited to, formation of ocular drusen, inflammation in the eye or eye tissue, loss of photoreceptor cells, loss of vision (including for example visual acuity and visual field), neovascularization (such as choroidal neovascularization or CNV), and retinal detachment. Other related aspects, such as photoreceptor degeneration, RPE degeneration, retinal degeneration, chorioretinal degeneration, cone degeneration, retinal dysfunction, retinal damage in response to light exposure (such as constant light exposure), damage of the Bruch's membrane, loss of RPE function, gain or RPE function, loss of integrity of the histoarchitecture of the cells and/or extracellular matrix of the normal macular, loss of function of the cells in the macula, photoreceptor dystrophy, mucopolysaccharidoses, rod-cone dystrophies, cone-rod dystrophies, anterior and posterior uveitis, and diabetic neuropathy, are also included. In some embodiments, the invention provides methods of improving corneal wound healing and/or improving the outcome of corneal transplantations.

[0445] In some embodiments, there are provided methods of treating an ocular disease in an individual, *e.g.*, an ocular disease described herein, comprising administering to the individual an effective amount of a composition comprising a targeting construct: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety comprising a therapeutic moiety or a fragment thereof. In certain embodiments, the ocular disease is AMD. In certain embodiments, the AMD is wet AMD. In certain embodiments, the AMD is dry AMD. In addition to macular degeneration, other eye diseases that can be treated by

methods of the present invention include, for example, retinitis pigmentosa, diabetic retinopathy, and other eye diseases that involve a local inflammatory process. In some embodiments, the eye disease is uveitis (anterior and posterior). In some embodiments, the eye disease is retinitis pigmentosa. In some embodiments, the eye disease involves the cornea. In some embodiments, the eye disease is proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, or ocular melanoma.

[0446] In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) formation of drusen and other extracellular deposits in the eye of an individual, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *e.g.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) inflammation in the eye of an individual, comprising administering to the individual an effective amount of a composition comprising targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *e.g.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) loss of photoreceptors cells in an individual, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *e.g.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) loss of photoreceptors cells in an individual, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *e.g.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) neovascularization associated with AMD, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an

active moiety, *i.e.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of treating (such as reducing, delaying, eliminating, or preventing) retinal detachment, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *e.g.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of improving (including for example decreasing, delaying, or blocking loss of) visual acuity or visual field in the eye of an individual, comprising administering to the individual an effective amount of a composition comprising a targeting construct comprising: a) a targeting moiety comprising a B4 or C2 antibody or a fragment thereof, and b) an active moiety, *i.e.*, one or more therapeutic moieties described herein. In some embodiments, there are provided methods of improving corneal wound healing or the outcome of corneal transplantations in the eye of an individual, comprising administering to the individual an effective amount of a targeting construct described herein.

[0447] In some embodiments, there are provided methods of treating a drusen-associated disease. The term “drusen-associated disease” refers to any disease in which formation of drusen or drusen-like extracellular disease plaque takes place, and for which drusen or drusen-like extracellular disease plaque causes or contributes to thereto or represents a sign thereof. For example, AMD, characterized by the formation of macular drusen, is considered as a drusen-associated disease. Non-ocular drusen-related disease include, but are not limited to, amyloidosis, elastosis, dense deposit disease, and/or atherosclerosis.

Modes of administration

[0448] The compositions described herein can be administered to an individual via any route, including, but not limited to, intravenous (*e.g.*, by infusion pumps), intraperitoneal, intraocular, intra-arterial, intrapulmonary, oral, inhalation, intravesicular, intramuscular, intra-tracheal, subcutaneous, intraocular, intrathecal, transdermal, transpleural, intraarterial, topical, inhalational (*e.g.*, as mists or sprays), mucosal (such as via nasal mucosa), subcutaneous, transdermal, gastrointestinal, intraarticular, intracisternal, intraventricular, rectal (*i.e.*, via suppository), vaginal (*i.e.*, via pessary), intracranial, intraurethral, intrahepatic, and intratumoral. In some

embodiments, the compositions are administered systemically (for example by intravenous injection). In some embodiments, the compositions are administered locally (for example by intraarterial or intraocular injection).

[0449] In some embodiments, the compositions are administered directly to the eye or the eye tissue. As used herein, the term “eye” refers to any and all anatomical tissues and structures associated with an eye. The eye has a wall composed of three distinct layers: the outer sclera, the middle choroid layer, and the inner retina. The chamber behind the lens is filled with a gelatinous fluid referred to as the vitreous humor. At the back of the eye is the retina, which detects light. The cornea is an optically transparent tissue, which conveys images to the back of the eye. The cornea includes one pathway for the permeation of drugs into the eye. Other anatomical tissue structures associated with the eye include the lacrimal drainage system, which includes a secretory system, a distributive system and an excretory system. The secretory system comprises secretors that are stimulated by blinking and temperature change due to tear evaporation and reflex secretors that have an efferent parasympathetic nerve supply and secrete tears in response to physical or emotional stimulation. The distributive system includes the eyelids and the tear meniscus around the lid edges of an open eye, which spread tears over the ocular surface by blinking, thus reducing dry areas from developing.

[0450] In some embodiments, the compositions are administered directly to the eye or the eye tissue. In some embodiments, the compositions are administered topically to the eye, for example, in eye drops. In some embodiments, the compositions are administered by injection to the eye (intraocular injection) or to the tissues associated with the eye. The compositions can be administered, for example, by intraocular injection, periocular injection, subretinal injection, intravitreal injection, trans-septal injection, subcleral injection, intrachoroidal injection, intracameral injection, subconjunctival injection, subconjunctival injection, sub-Tenon's injection, retrobulbar injection, peribulbar injection, or posterior juxtасleral delivery. These methods are known in the art. For example, for a description of exemplary periocular routes for retinal drug delivery, see Periocular routes for retinal drug delivery, Raghava et al. (2004), Expert Opin. Drug Deliv. 1(1):99-114. The compositions may be administered, for example, to the vitreous, aqueous humor, sclera, conjunctiva, the area between the sclera and conjunctiva, the choroid tissues, the retina choroids tissues, macula, or other area in or proximate to the eye of an individual. The

compositions can also be administered to the individual as an implant. Preferred implants are biocompatible and/or biodegradable sustained release formulations which gradually release the compounds over a period of time. Ocular implants for drug delivery are well-known in the art. See, e.g., U.S. Pat. No. 5,501,856, 5,476,511, and 6,331,313. The compositions can also be administered to the individual using iontophoresis, including, but are not limited to, the ionophoretic methods described in U.S. Pat. No. 4,454,151 and U.S. Pat. App. Pub. No. 2003/0181531 and 2004/0058313.

[0451] In some embodiments, the compositions are administered intravascularly, such as intravenously (IV) or intraarterially. In some embodiments (for example for the treatment of renal diseases), the compositions are administered directly into arteries (such as renal arteries).

[0452] In some embodiments, the compositions are administered directly into the joint tissue. In some embodiments, the compositions are administered to the synovium.

[0453] The optimal effective amount of the compositions can be determined empirically and will depend on the type and severity of the disease, route of administration, disease progression and health, mass and body area of the individual. Such determinations are within the skill of one in the art. The effective amount can also be determined based on in vitro complement activation assays. Examples of dosages of antibodies (or antigen-binding fragments thereof) and/or constructs (e.g., targeting constructs) which can be used for methods described herein include, but are not limited to, an effective amount within the dosage range of any of about 0.01 $\mu\text{g/kg}$ to about 300 mg/kg , or within about 0.1 $\mu\text{g/kg}$ to about 40 mg/kg , or with about 1 $\mu\text{g/kg}$ to about 20 mg/kg , or within about 1 $\mu\text{g/kg}$ to about 10 mg/kg . For example, when administered intraocularly, the composition may be administered at low microgram ranges, including for example about 0.1 $\mu\text{g/kg}$ or less, about 0.05 $\mu\text{g/kg}$ or less, or 0.01 $\mu\text{g/kg}$ or less. In some embodiments, the amount of an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) administered to an individual is about 10 μg to about 500 mg per dose, including for example any of about 10 μg to about 50 μg , about 50 μg to about 100 μg , about 100 μg to about 200 μg , about 200 μg to about 300 μg , about 300 μg to about 500 μg , about 500 μg to about 1 mg , about 1 mg to about 10 mg , about 10 mg to about 50 mg ,

about 50 mg to about 100 mg, about 100 mg to about 200 mg, about 200 mg to about 300 mg, about 300 mg to about 400 mg, or about 400 mg to about 500 mg per dose.

[0454] The antibody (or antigen-binding fragment thereof) and/or construct (e.g., targeting construct) compositions may be administered in a single daily dose, or the total daily dose may be administered in divided dosages of two, three, or four times daily. The compositions can also be administered less frequently than daily, for example, six times a week, five times a week, four times a week, three times a week, twice a week, once a week, once every two weeks, once every three weeks, once a month, once every two months, once every three months, or once every six months. The compositions may also be administered in a sustained release formulation, such as in an implant which gradually releases the composition for use over a period of time, and which allows for the composition to be administered less frequently, such as once a month, once every 2-6 months, once every year, or even a single administration. The sustained release devices (such as pellets, nanoparticles, microparticles, nanospheres, microspheres, and the like) may be administered by injection or surgically implanted in various locations in the eye or tissue associated with the eye, such as intraocular, intravitreal, subretinal, periocular, subconjunctival, or sub-tenons.

[0455] The antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) compositions (e.g., pharmaceutical compositions) can be administered alone or in combination with other molecules known to have a beneficial effect on retinal attachment or damaged retinal tissue, including molecules capable of tissue repair and regeneration and/or inhibiting inflammation. Examples of useful cofactors include anti-VEGF agents (such as an antibody against VEGF), basic fibroblast growth factor (bFGF), ciliary neurotrophic factor (CNTF), axokine (a mutein of CNTF), leukemia inhibitory factor (LIF), neurotrophin 3 (NT-3), neurotrophin-4 (NT-4), nerve growth factor (NGF), insulin-like growth factor II, prostaglandin E₂, 30 kD survival factor, taurine, and vitamin A. Other useful cofactors include symptom-alleviating cofactors, including antiseptics, antibiotics, antiviral and antifungal agents and analgesics and anesthetics.

[0456]

[0457] Gene Therapy

[0458] The targeting constructs can also be delivered by expression of the targeting construct fusion protein in vivo, which is often referred to as “gene therapy”. For

example, cells may be engineered with a polynucleotide (DNA or RNA) encoding for the fusion protein ex vivo, the engineered cells are then provided to an individual to be treated with the fusion protein. Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding for the fusion protein of the present invention.

[0459] Local delivery of the targeting construct of the present invention using gene therapy may provide the therapeutic agent to the target area, for example to the eye or the eye tissue.

[0460] The an antibodies (or antigen-binding fragments thereof) and/or constructs (e.g., a targeting constructs) can also be delivered by expression of the antibody and/or targeting construct in vivo, which is often referred to as “gene therapy”. For example, cells may be engineered with a polynucleotide (DNA or RNA) encoding for the antibody and/or targeting construct ex vivo, the engineered cells are then provided to an individual to be treated with the antibody and/or targeting construct. Such methods are well-known in the art. For example, cells may be engineered by procedures known in the art by use of a retroviral particle containing RNA encoding for the antibody and/or targeting construct of the present invention.

[0461] Local delivery of the antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) of the present invention using gene therapy may provide the therapeutic agent to the target area, for example to the eye or the eye tissue.

[0462] Methods of gene delivery are known in the art. These methods include, but are not limited to, direct DNA transfer, see, e.g., Wolff et al. (1990) Science 247: 1465-1468; 2) Liposome-mediated DNA transfer, see, e.g., Caplen et al. (1995) Nature Med. 3:39-46; Crystal (1995) Nature Med. 1:15-17; Gao and Huang (1991) Biochem. Biophys. Res. Comm. 179:280-285; 3) Retrovirus-mediated DNA transfer, see, e.g., Kay et al. (1993) Science 262:117-119; Anderson (1992) Science 256:808-813; 4) DNA Virus-mediated DNA transfer. Such DNA viruses include adenoviruses (preferably Ad2 or Ad5 based vectors), herpes viruses (preferably herpes simplex virus based vectors), and parvoviruses (preferably “defective” or non-autonomous parvovirus based vectors, more preferably adeno-associated virus based vectors, most preferably AAV-2 based vectors). See, e.g., Ali et al. (1994) Gene Therapy 1:367-

384; U.S. Pat. No. 4,797,368, incorporated herein by reference, and U.S. Pat. No. 5,139,941.

[0463] Retroviruses from which the retroviral plasmid vectors hereinabove mentioned may be derived include, but are not limited to, Moloney Mouse Leukemia Virus, spleen necrosis virus, retroviruses such as Rous Sarcoma Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, adenovirus, Myeloproliferative Sarcoma Virus, and mammary tumor virus. In one embodiment, the retroviral plasmid vector is derived from Moloney Mouse Leukemia Virus.

[0464] Adenoviruses have the advantage that they have a broad host range, can infect quiescent or terminally differentiated cells, such as neurons or hepatocytes, and appear essentially non-oncogenic. See, e.g., Ali et al. (1994), *supra*, p. 367. Adenoviruses do not appear to integrate into the host genome. Because they exist extrachromosomally, the risk of insertional mutagenesis is greatly reduced. Ali et al. (1994), *supra*, p. 373.

[0465] Adeno-associated viruses exhibit similar advantages as adenoviral-based vectors. However, AAVs exhibit site-specific integration on human chromosome 19 (Ali et al. (1994), *supra*, p. 377).

[0466] The gene therapy vectors include one or more promoters. In some embodiments, the vector has a promoter that drives expression in multiple cell types. In some embodiments, the vector has a promoter that drives expression in specific cell types (such as cells of retina or cells in the kidney). Suitable promoters which may be employed include, but are not limited to, the retroviral LTR; the SV40 promoter; and the human cytomegalovirus (CVM) promoter described in Miller et al. (1989) *Biotechniques* 7(9):980-990, or any other promoter (e.g., cellular promoters such as eukaryotic cellular promoters including, but not limited to, the histone, pol III, and .beta.-actin promoters). Other viral promoters which may be employed include, but are not limited to, adenovirus promoters, thymidine kinase (TK) promoters, and B19 parvovirus promoters. The selection of a suitable promoter will be apparent to those skilled in the art from the teachings contained herein.

[0467] The nucleic acid sequence encoding an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) is under the control of a suitable promoter. Suitable promoters which may be employed include, but are not limited to, adenoviral promoters, such as the adenoviral major late promoter; or heterologous

promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs (including the modified retroviral LTRs hereinabove described); the β -actin promoter; and human growth hormone promoter.

[0468] Retroviral plasmid vectors can be employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which maybe transfected are described in Miller (1990) Human Gene Therapy 1:5-14. The vectors may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO₄ precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a lipid, and then administered to a host. The producer cell line generates infectious retroviral vector particles which include the nucleic acid sequence(s) encoding the polypeptides. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either in vitro or in vivo. The transduced eukaryotic cells will express the nucleic acid sequence(s) encoding the polypeptide. Eukaryotic cells which may be transduced include, but are not limited to, embryonic stem cells, embryonic carcinoma cells, as well as hematopoietic stem cells, hepatocytes, fibroblasts, myoblasts, keratinocytes, endothelial cells, and bronchial epithelial cells.

[0469] In some embodiments, gene delivery vectors which direct expression of an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) in the eye are used. Vectors for gene delivery to the eye are known in the art, and have been disclosed, for example, in U.S. Patent No. 6,943,153, and U.S. Patent Application Publication Nos. US20020194630, US20030129164, US200600627165.

[0470] In some embodiments, the complement activation is inhibited by contacting a body fluid with a composition comprising an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) ex vivo under conditions that permit the antibody (or antigen-binding fragment thereof) and/or construct (e.g., targeting construct) to function to inhibit complement activation. Suitable body fluids include those that can be returned to the individual, such as blood, plasma, or lymph. Affinity adsorption apheresis is described generally in Nilsson et al. (1988) Blood

58(1):38-44; Christie et al. (1993) Transfusion 33:234-242; Richter et al. (1997) ASAIO J. 43(1):53-59; Suzuki et al. (1994) Autoimmunity 19: 105-112; U.S. Pat. No. 5,733,254; Richter et al. (1993) Metabol. Clin. Exp. 42:888-894; and Wallukat et al. (1996) Int'l J. Card. 54:1910195.

[0471] Accordingly, the invention include methods of treating one or more diseases described herein in an individual comprising treating the individual's blood extracorporeally (i.e., outside the body or ex vivo) with a composition comprising a targeting construct under conditions that permit the molecule to function to inhibit complement activation, and returning the blood to the individual.

[0472] Unit dosages, articles of manufacture, and kits

[0473] Also provided are unit dosage forms of an antibody (or antigen-binding fragment thereof) and/or construct (e.g., targeting construct) compositions, each dosage containing from about 0.01 mg to about 50 mg, including for example any of about 0.1 mg to about 50 mg, about 1 mg to about 50 mg, about 5 mg to about 40 mg, about 10 mg to about 20 mg, or about 15 mg of the targeting construct. In some embodiments, the unit dosage forms of an antibody (or antigen-binding fragment thereof) and/or construct (e.g., a targeting construct) composition comprises about any of 0.01 mg-0.1mg, 0.1 mg-0.2 mg, 0.2 mg-0.25 mg, 0.25 mg-0.3 mg, 0.3 mg-0.35 mg, 0.35 mg-0.4 mg, 0.4 mg-0.5 mg, 0.5 mg-1.0 mg, 10 mg-20 mg, 20mg -50 mg, 50 mg-80 mg, 80 mg-100 mg, 100 mg-150 mg, 150 mg-200 mg, 200 mg-250 mg, 250 mg-300 mg, 300 mg-400 mg, or 400 mg-500 mg targeting construct. In some embodiments, the unit dosage form comprises about 0.25 mg targeting construct. The term "unit dosage form" refers to a physically discrete unit suitable as unitary dosages for an individual, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect, in association with a suitable pharmaceutical carrier, diluent, or excipient. These unit dosage forms can be stored in a suitable packaging in single or multiple unit dosages and may also be further sterilized and sealed.

[0474] Also provided are articles of manufacture comprising the compositions described herein in suitable packaging. Suitable packaging for compositions (such as ophthalmic compositions) described herein are known in the art, and include, for example, vials (such as sealed vials), vessels, ampules, bottles, jars, flexible

packaging (e.g., sealed Mylar or plastic bags), and the like. These articles of manufacture may further be sterilized and/or sealed.

[0475] The present invention also provides kits comprising compositions (or unit dosages forms and/or articles of manufacture) described herein and may further comprise instruction(s) on methods of using the composition, such as uses described herein. The kits described herein may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, syringes, and package inserts with instructions for performing any methods described herein.

EXAMPLES

[0476] Example 1: Generation of targeting constructs.

[0477] To identify self-reactive monoclonal antibodies (mAbs) that recognize neo-epitopes on ischemic tissues, fresh isolated intestinal epithelial cells (IECs) were used to screen hybridomas obtained by the fusion of peritoneal, lymph node, and spleen cells from wild-type C57BL/6 mice with the Sp2/0-Ag14 myeloma cell line as described in Kulik et al., *J Immunol.*, (2009), 182:5363-5373. Briefly, hybridoma fusions were screened based on reactivity by Western blot analysis on IEC lysates and positive surface staining of IECs as detected by flow cytometric analysis. For flow cytometry analysis, isolated IECs were washed in staining buffer (2% FCS/0.01% NaN₃/PBS) then resuspended in staining buffer containing hybridoma supernatant, and incubated for 30 min at room temperature. After incubation, cells were washed in staining buffer three times and then incubated with secondary goat anti-mouse IgM (μ -chain specific) antibodies (Jackson ImmunoResearch Laboratories) for 30 minutes at room temperature. Following incubation, cells were washed as described above and then resuspended in the staining buffer. Flow cytometry was performed using a BD Biosciences FACSCalibur. For Western blot analysis, IECs were lysed on ice for 20 min in a buffer containing 0.5% Triton X-100, 0.5% Chaps, 20 mM Tris-HCl (pH 7.5), 1 mM EDTA, 10 μ g/ml leupeptin, and protease inhibitor mixture (Roche Molecular Biochemicals). Lysates were cleared by centrifugation at 8000 x g for 5 minutes. After separation by 8% Tris-glycine SDS-PAGE, the proteins were transferred to a polyvinylidene difluoride membrane. The membrane was blocked overnight with 5% nonfat milk dissolved in PBS. The membrane was washed in PBS and then probed with an antibody from hybridoma supernatant for 1–2 hours in 2% milk/PBS, washed, and then incubated with HRP-conjugated secondary antibodies. A

positive signal was visualized using the ECL system (PerkinElmer). Candidate hybridomas were subsequently serially recloned to obtain monoclonal cell lines stably producing a single mAb. To purify candidate mAbs, antibodies from the exhausted supernatants of cultured hybridomas were affinity purified on a column of agarose beads with goat anti-human IgM (Sigma-Aldrich). Bound antibody was eluted with a buffer containing 0.1 M glycine (pH 2.3) and collected into a buffer containing 1.5 M Tris (pH 8.8). Eluted mAb was dialyzed against PBS (pH 7.4) for 48 hours and concentrated using centrifugal filtration on Centricon Plus-20 (Millipore). Antibody concentration was determined by measuring the A280 nm of the sample, and purity was confirmed by analysis on a 10% SDS-PAGE gel.

[0478] A hybridoma of interest produced an IgM κ isotype antibody designated as mAb B4. Flow cytometric analysis demonstrated that mAb B4 bound to a surface epitope on IEC but not to freshly isolated splenocytes or thymocytes. By Western blot analysis, mAb B4 recognized a protein with a molecular weight of 37 kDa in IEC lysates but not lysates from freshly isolated splenocytes or thymocytes. When other tissue lysates were probed by Western blot with mAb B4, the epitope was found to be widely distributed, including in the lung, liver, and kidney. To identify the protein recognized by mAb B4 on IEC, 2D gel separation of proteins according to their molecular weight and charge was conducted based on the method described in Vossenaar et al., *Arthritis Res. Ther.*, (2004), 6:R142-R150 and Kulik et al., *J Immunol.*, (2009), 182:5363-5373. After 2D separation, proteins reactive with mAb B4 were characterized by electrospray liquid chromatography MS analysis as described in Kulik et al., *J Immunol.*, (2009), 182:5363-5373, and it was determined that mouse annexin IV was a protein recognized by mAb B4.

[0479] Another hybridoma of interest produced an IgM antibody designated as mAb C2 which did not react with Western blots of IECs. To determine reactivity of mAb C2 to phospholipids, which have been suggested to be exposed in apoptotic and ischemic cells, ELISA analysis was performed using phospholipid as a binding partner as described in Elvington et al., *J Immunol.*, (2012), 188:1460-1468. Briefly, microtiter plates (Immulon 1B; Dynatech Laboratories, Chatilly, VA) coated with 100 μ l/well of 50 μ g/ml phospholipid in methanol were dried under blowing air to allow the organic solvent to evaporate, and the wells were then washed with PBS and blocked with 1% BSA. Supernatant from the mAb C2 hybridoma was added to wells and bound antibody was detected by alkaline phosphatase-conjugated goat anti-mouse

IgM (Jackson ImmunoResearch Laboratories, West Grove, PA). Phospholipids assayed included phosphatidylserine (PS)-1,2-distearoyl-sn-glycerol-3-[phospho-L-serine] (Avanti Polar-lipids, Alabaster, AL) (referred to as PS), cardiolipin from bovine heart (referred to as CL), phosphatidylethanolamine (PE)-1,2-diacyl-sn-glycerol-3-phosphoethanolamine (referred to as PE), phosphatidylglycerol (PG)-1,2-diacyl-sn-glycerol-3-phospho-(1-rac-glycerol) from yolk lecithin (referred to as PG), and phosphorylcholine (PC)(10)-BSA (Biosearch Technologies, Novato, CA) (referred to as PC-BSA). mAb C2 was shown to recognize a subset of phospholipids that included PC-BSA, PE, and CL, but not PG or PS.

[0480] Since B4 and C2 mAbs were identified as antibodies that recognize neo-epitopes of ischemic tissue that can contribute to complement pathway mediated injury, targeting constructs comprising a targeting moiety, in this case scFv isolated from the IgM-B4 antibody or IgM-C2 antibody, and an active moiety, in this case a complement modulator, were generated and tested for their ability to protect ischemic tissue from injury. For generation of targeting constructs a scFv from both IgM-B4 antibody and IgM-C2 antibody was prepared by amplifying the isolated VH and VL genes from cDNA and by linking using overlap extension PCR and expressed as a scFv with an N-terminal His-tag. Purified scFv had the expected molecular weight by SDS-PAGE. The purified scFv retained parent IgM binding specificity as demonstrated by its ability to directly bind to its binding partner, which in the case of B4scFv, was the ability to directly bind to recombinant annexin IV in vitro as well as to competitively inhibit binding of B4 mAb to annexin IV (Fig. 1A and B). The targeting moiety B4scFv or C2scFv encoding sequences were linked with the artificial linker (G4S1)₂ to an active moiety encoding sequence selected from the complement modulators: complement receptor 1-like protein (Crry), complement factor H (fH), or CD59 molecule complement regulatory protein (CD59). For Crry and CD59, the sequence encoding the extracellular domain of the proteins was used. For fH, the sequence encoding the N-terminal 5 short consensus repeat domains (the active region) was used. The targeting construct encoding sequences were inserted into an expression vector and expression plasmids were transfected into Chinese hamster ovary (CHO) cells. Positive high expressing clones were selected by a limiting dilution assay and targeting construct proteins, specifically B4scFv-Crry, B4scFv-fH, B4scFv-CD59, C2scFv-Crry, C2scFv-fH, and C2scFv-CD59 proteins were produced, recovered from culture supernatant, and purified by affinity chromatography using

either antibodies to the active moiety or to the His tag. The purified targeting construct had the expected molecular weight by SDS-PAGE and the activity of the complement modulator was retained. In the case of B4scFv-Crry, in vitro complement modulatory activity was tested in a standard assay that measured C3 deposition on zymosan particles and compared to positive control CR2-Crry (Fig. 1C).

[0481] Mouse monoclonal antibody B4 was deposited at the ATCC with ATCC Deposit No. PTA-13522 (B4/14/12). Mouse monoclonal antibody C2 was deposited at the ATCC with ATCC Deposit No. PTA-13523 (C2/19/8).

[0482] Example 2: Reduction of spinal cord injury by treatment with a targeting construct.

[0483] The role of self-reactive C2 or B4 mAbs in spinal cord injury (SCI) and cerebral ischemia reperfusion injury (ischemic stroke) was investigated. Rag1^{-/-} mice, which produce no mature T-cells or B-cells and are therefore antibody-deficient, have been previously reported to be protected from ischemia reperfusion (IR) injury. See Williams et al., J. Appl. Physiol. (1999), 86:938-942. For these studies, mouse models of SCI and middle cerebral artery occlusion (MCAo) (with 60 minutes ischemia and 24 hours reperfusion) were used as described in Qiao et al, Am. J. Pathol, (2006), 169:1039-1047 and Atkinson et al, J Immunol., (2006), 177:7266-7274. Antibodies were administered intravenously 60 minutes after SCI or at time of reperfusion for MCAo model. Locomotor function was assessed after SCI using the Basso, Beattie and Bresnahan (BBB) rating scale developed for rats, but adapted for mice (Noble et al., J. Neuroscience, (2002), 22:7526-7535). To measure infarct volume after MCAo, coronal section from isolated brains were stained with 2% triphenyltetrazolium (TTC) and infarcted area (excluding dye) was determined using NIH image analysis software. It was determined that IgM antibody deficient Rag1^{-/-} mice were protected from SCI as measured by locomotor activity, but injury was reconstituted in Rag1^{-/-} mice to wild-type mouse levels by intravenous administration of C2 or B4 mAbs, but not with control F632 mAb (Fig. 2A). Furthermore, B4 mAb and C2 mAb, but not control antibody, reconstituted cerebral ischemia and reperfusion injury in Rag1^{-/-} mice (Fig. 2B).

[0484] C57Bl/6 wild-type mice were administered targeting construct B4scFV-Crry, to test a possible protective effect of this targeting construct against self-reactive

antibodies in an ischemic injury model of SCI. In this SCI model, 0.2 mg B4-Crry or phosphate buffered saline (PBS) was injected intravenously 60 minutes post-injury. Assessment of locomotor activity showed that B4-Crry treated mice had significantly improved recovery and protected mice from SCI as compared to PBS treated mice (Fig. 3A). Tissue sparing was also analyzed at 3 days post SCI, and there was significantly increased tissue sparing in B4scFv-Crry treated C57Bl/6 wild-type mice (Fig. 3B, filled triangles) as compared to control C57Bl/6 wild-type mice administered PBS (Fig. 3B, filled circles). Tissue sparing of B4scFv-Crry treated C57Bl/6 wild-type mice was comparable to tissue sparing in Rag1^{-/-} mice treated with PBS (Fig. 3B, filled circles, upper line) or control F632 mAb (Fig. 3B, empty squares) but was significantly increased as compared to Rag1^{-/-} mice reconstituted with C2 mAb (Fig. 3B, filled squares) or B4 mAb (Fig. 3B, open circles). Furthermore, IgM binding to the spinal cord following SCI in C57Bl/6 wild type mice was confirmed, and IgM and C3 was shown to colocalize in spinal cords from wild-type mice 24 hours after injury (Fig. 4A-C). Administration of B4scFv-Crry reduced IgM and C3 deposition 24 hours after injury (Fig. 4D and E).

[0485] Example 3: Administration of a targeting construct did not increase host susceptibility to infection.

[0486] A major potential advantage of targeted versus systemic complement inhibition is that a targeted approach is less likely to impact physiologically important roles of complement. One such important role, and one that has some relevance to a transplant patient, is host defense against infection. The effect of B4scFv-Crry on susceptibility to infection using a well characterized model of polymicrobial sepsis was investigated with a 0.2 mg dose that was therapeutic in animal models of SCI as well as animal models of hepatic and cardiac ischemia reperfusion injury. Cecal ligation and puncture was performed in C3 deficient mice or in wild-type mice treated with either B4scFv-Crry or PBS, and survival was monitored. B4scFv-Crry had no effect on host susceptibility to infection in a model of acute septic peritonitis, and mice treated with B4scFv-Crry survived significantly longer following cecal ligation puncture, and survival was not significantly different compared to PBS treated wild-type mice (Fig. 5). In contrast, all mice deficient in C3 died within 48 hours, a similar result that was previously obtained for both C3 deficient mice and mice treated with a therapeutic dose of Crry-Ig, a systemic counterpart of B4scFv-Crry.

[0487] Example 4: Reduction of cardiac ischemia reperfusion injury by treatment with a targeting construct.

[0488] The role of self-reactive natural IgM in post-transplant cardiac ischemia reperfusion injury was investigated in isografts after heart transplantation from C57BL/6 wild-type donors to either antibody deficient Rag1^{-/-} recipients or Rag1^{-/-} recipients reconstituted with C2 mAb or B4 mAb. mAbs were administered intravenously to recipient immediately following reperfusion of transplanted heart. Immunofluorescence for endothelial markers, IgM and C3 binding was performed on 8 micron cryosections stained with appropriate fluorescently labeled antibodies, and imaged with a confocal microscope. For histopathology, sections were stained with H&E and scored by an observer blinded to experimental groups. Sections were scored on a scale of 0-3 as described in Atkinson et al., J. Immunol., (2010), 185:7007-7013. Compared to grafts in wild-type recipients (Balb/C to C57BL/6), hearts transplanted into antibody-deficient Rag1^{-/-} recipient mice by heterotopic abdominal heart transplantation were protected from post-transplant cardiac IR injury as evidenced by reduced histologic injury and inflammation and decreased serum levels of cardiac troponin I, an index of cardiac cell damage. However, reconstitution with either B4 mAb or C2 mAb, but not control F632 IgM, in Rag1^{-/-} recipients restored cardiac IR injury in the transplanted hearts to that seen in wild-type recipients. The graft specificity and binding characteristics of B4 mAb and C2 mAb was investigated by immunofluorescence analysis of graft sections. In B4 mAb (Fig. 6A) and C2 mAb treated heart recipient Rag1^{-/-} mice, both mAbs bound to endothelial cells of arterioles, capillaries, and microvessels within the myocardium of the transplanted heart, but not the native heart. Additional mAb binding in grafts was seen on myocytes, with a preferential localization to injured myocytes in epicardium. No IgM immunostaining could be detected in the graft or native heart in animals reconstituted with F623 control mAb (Fig. 6A). Analysis of IgM binding and complement activation showed that B4 IgM co-localized with C3d (complement activation product) in hearts transplanted in Rag1^{-/-} recipients and that endogenous IgM co-localized with C3d in grafts of wild-type recipients (Fig. 6B).

[0489] In a therapeutic protocol, a 0.2mg dose of B4scFV-Crry or B4scFv was administered immediately post-transplantation in a wild-type allograft transplant model (Balb/C to C57BL/6). Allografts were harvested at 6 hours (after intravenous

B4scFv treatment) or 48 hours (after intravenous B4scFV-Crry treatment) after reperfusion and analyzed. When administered immediately after reperfusion, administration of B4scFv or B4scFv-Crry resulted in a significant reduction in myocardial IRI as evidenced by decreased cardiac troponin I, an index of cardiac cell damage levels (Fig. 7A). Grafts were also assessed for histological evidence of injury and inflammatory cell infiltration. In accordance with cardiac troponin I levels, grafts from B4scFv-Crry treated recipients had a significantly lower injury and inflammation histology score than grafts from vehicle treated recipients (Fig. 7B). Reduced graft injury and inflammation in B4scFv treated recipients was also observed as compared to recipients receiving PBS vehicle (Fig. 7B). In His-tagged B4scFv treated allografts, graft sections stained positive with anti-His antibodies (Fig. 7C), and B4scFv co-localized with a pan-endothelial marker (Endo-1) with B4scFv binding seen in post-ischemic vessels of all sizes. In addition, B4scFv-Crry reduced C3 deposition as shown by a reduced C3 immunostaining (Fig. 7D). Quantitative analysis of the images demonstrated that IgM binding and C3d deposition occurred in grafts from recipients treated with PBS, with staining predominantly in the microvasculature at 6 hours post-transplantation, and in the microvasculature and on myocytes at 48 hours post-transplantation (Fig. 7E and F). Treatment of recipients with either B4scFv or B4scFv-Crry significantly reduced or eliminated both IgM and C3d deposition in grafts at 6 hours post-transplantation. At 48 hours post-transplantation, however, significant reductions in IgM binding and C3d deposition was seen only in grafts from B4scFv-Crry treated recipients (Fig. 7E and F). To further investigate the inflammatory environment modulated by B4scFv-Crry, the effect of this molecule on graft levels of specific cytokines and chemokines was assessed. B4scFv-Crry treatment of recipients significantly reduced graft levels of the cardiotoxic cytokine IL-6 and MCP-1 as compared to PBS vehicle treatment. Graft levels of KC at 48 hours post-transplantation were not altered by treatment with either B4scFv-Crry (Fig. 8). These molecules also reduced inflammatory cell infiltration as compared to PBS control treated mice.

[0490] To determine the circulatory half-life ($t_{1/2}$) of B4scFV-Crry, 100 μ g B4scFv-Crry was injected intravenously into C57BL/6 recipient mice, and plasma concentration of the protein was determined at different time points by ELISA using an anti-Crry antibody. There was a two-phase elimination profile; an initial rapid phase with a $t_{1/2}$ of 9.7 minutes, and a second prolonged phase with a $t_{1/2}$ of 5.5

hours. A similar two-phase elimination profile has been shown for other biologics, including untargeted Crry (Crry-Ig), although the second phase $t_{1/2}$ for Crry-Ig was considerably longer (40 hrs). Significant therapeutic efficacy was seen with a 0.2 mg dose of B4scFv-Crry, and since preferential accumulation at the target site would be expected to translate to a lower dose requirement for therapeutic benefit, a biodistribution study was performed. ^{125}I -labeled B4scFv-Crry was injected into recipient mice immediately after heart transplantation, and tissue distribution of radiolabel determined 6 hours later. In addition to being present in the circulation, ^{125}I -B4 scFv-Crry localized primarily to the transplanted graft (Fig.9).

[0491] Example 5: Species cross-reactivity of self-reactive IgM.

[0492] Reconstitution of Rag1^{-/-} mice with human natural IgM reconstituted ischemia reperfusion injury (IRI) via IgM mediated complement activation, demonstrating species cross reactivity. Furthermore, there were high levels of anti-annexin IV Abs (B4 mAb specificity) present in human serum. Cross species reactivity of mouse B4 mAb and C2 mAb was further investigated by exposing human umbilical vein endothelial cells and a mouse brain endothelial cell line to 3 hours of hypoxia followed by 12 hours normoxia in the presence of either B4 mAb or C2 mAb in vitro. Flow cytometry and immunofluorescence microscopy demonstrated that both mAbs bound to mouse (Fig. 10A) and human cells (Fig. 10C) exposed to hypoxia, but not to control normoxic cells (Fig. 10B and D), indicating that B4 and C2 show cross specificity for human neoepitopes.

[0493] Example 6: Reduction of hepatic ischemia reperfusion injury by treatment with a targeting construct.

[0494] Complement activation plays an important role in both hepatic ischemia/reperfusion injury (IRI) and in the priming phase of liver regeneration. Utilizing separate models of hepatic IRI and 70% partial hepatectomy (Phx), the role of IgM in both hepatic IRI and liver regeneration was investigated. For IRI experiments, saline or a 25 μg dose of C2 mAb or B4 mAb was injected into antibody deficient (Rag1^{-/-}) mice following 30 minutes total warm ischemia and just prior to reperfusion. Samples were taken after 6 hours of reperfusion. Serum ALT levels were measured by ELISA and histological necrosis was quantified in H&E stained sections (scale 0-3). For 70% partial hepatectomy experiments, saline or a 10 μg dose

of C2 mAb or B4 mAb was injected into antibody deficient (Rag1^{-/-}) mice immediately following 70% Phx. Samples were taken 48 hours post Phx. Mitotic index was calculated as a measure of hepatic regeneration (Mitotic figures/total cells in 10 hpf). Serum ALT levels were measured and histological necrosis was quantified (scale 0-3). The results showed that antibody deficient Rag1^{-/-} mice were protected from hepatic IRI, and reconstitution of Rag1^{-/-} mice with either B4 mAb or C2 mAb restored IRI to a level close to that seen in wild-type mice as determined by serum ALT levels (Fig. 11A) and histological score (Fig. 11B). Following 70% PHx in Rag1^{-/-} mice, there was increased injury in the remnant liver compared to wild-type mice, and reconstitution of Rag1^{-/-} mice with either B4 mAb or C2 mAb was protective as determined by serum ALT levels (Fig. 12A) and histological score (Fig. 12B). Rag1^{-/-} mice had an impaired regenerative response after 70% PHx compared to wild-type mice, but reconstitution with B4 mAb or C2 mAb restored regeneration as assessed by mitotic index (Fig. 12C), liver weight restitution (Fig. 13A), and increase of Brdu-positive cells (Fig. 13B). Analysis of liver sections showed that IgM was deposited in the liver of wild-type mice after ischemia and reperfusion, and both B4 mAb and C2 mAb were deposited in livers of Rag1^{-/-} mice after IR (Fig. 14A) and after PHx (Fig. 14C). Furthermore, analysis of liver sections after either IR injury (Fig. 14B) or PHx (Fig. 14D) showed IgM deposition co-localized with C3 deposition and there was no detectable IgM or C3 in sections from untreated Rag1^{-/-} mice.

[0495] To determine the circulatory half-life ($t_{1/2}$) of B4scFV-Crry, 100 μ g B4scFv-Crry was injected intravenously into mice, and plasma concentration of the protein was determined at different time points by ELISA using an anti-Crry antibody. There was a two-phase elimination profile; an initial rapid phase with a $t_{1/2}$ of 27 minutes, and a second prolonged phase with a $t_{1/2}$ of 6.5 hours (Fig. 15A and B). To further determine the specificity of the C2 and B4 IgMs for injured tissues, biodistribution studies were performed in Rag1^{-/-} mice. Rag1^{-/-} mice were injected intravenously with ¹²⁵Iodine radiolabeled C2, B4, or isotype control antibody F632 after one of the following procedures: sham, 70% PHx, or IR. Organs and blood were harvested 6 hours after surgical procedures. Blood was removed by cardiac puncture and the animals were perfused with PBS before the heart, brain, liver, intestine, lung, kidney, and spleen were removed. Tissues were rinsed with PBS, shredded, weighed and then radioactivity was measured with a Hewlett-Packard 5780 γ counter. Results were

recorded in $\mu\text{Ci/g}$ of tissue. Both C2 and B4 antibodies targeted mainly to the liver following either IR or PHx (Fig. 15C and D). Additionally, some binding of both C2 and B4 was seen in the kidneys following IR or PHx and in the intestines following IR. Isotype control antibody F632 was not found in higher quantities in any organ as compared to sham animals. These findings suggested that similar epitopes were expressed following both IR and PHx in the liver, and that these epitopes were only found on injured tissues. Given the specificity of these epitopes for injured tissues they may serve as a ligand to specifically target a therapy to liver following either PHx or IRI. To confirm the role of IgM, B4 scFv was administered to the animals. B4 scFv had the same biodistribution profile following IRI or Phx as the B4 IgM mAb used in the above studies (Fig. 15E and F).

[0496] The *in vivo* activity of the targeting moiety B4scFv and the targeting construct B4scFv-Crry was assessed in the mouse model of hepatic ischemia reperfusion injury. Administration of B4scFv or B4scFv-Crry in mice subjected to 35 minutes of ischemia followed with 24 hours of reperfusion protected mice against hepatic ischemia reperfusion injury as demonstrated by serum ALT levels (Fig. 16A) and immunohistochemistry (Fig. 16B-E).

[0497] To determine if B4 mAb bound to human ischemic liver tissues, frozen human tissue samples were harvested just prior to transplantation. These livers were therefore ischemic but not yet reperfused. Sectioned liver was also stained for IgM post-reperfusion (Fig. 17A). It was observed that B4 mAb bound to human ischemic tissue and colocalized with the vascular endothelial marker CD31 demonstrating that B4 mAb bound to ischemic tissue in both humans and mice (Fig. 17B).

[0498] To determine if serum levels of natural IgMs were decreased following liver transplantation in human, serum samples were taken from patients undergoing liver transplantation at three time points: just prior to transplantation, 1 hours post-transplantation and 24 hours post-transplantation. Using ELISA to measure natural IgMs, it was observed that patient serum levels of IgMs specific for phospholipids and annexin IV were decreased at 1 and 24 hrs after transplantation as compared to the levels seen just prior to transplantation (Fig. 19). Total IgM levels were not significantly different between the time points (Fig. 18). The antigen specificities of the antibodies depleted in human serum were the same as the specificity of the mouse mAbs B4 and C2. To test if the depletion of these antibodies from the serum was a result of them binding to neoepitopes exposed in the post ischemic liver, levels of

bound IgM in the liver were measured prior to ischemia and then following reperfusion. There was little IgM detected in normally perfused livers. However, livers that were ischemic and then reperfused had high levels of IgM bound in a characteristic sinusoidal pattern. These data indicate that similar events occur in murine and human hepatic IRI, in that specific circulating natural IgMs recognized and bound to neoepitopes expressed following ischemia.

[0499] Example 7: In vivo study of IgM effects in a chemical and ischemic reperfusion mouse model of renal injury.

[0500] The role of natural IgM antibodies in a chemically induced mouse model of renal injury was investigated. Adriamycin nephropathy was induced in Balb/c mice in 8 to 10 week old male mice with a single intravenous injection of 11 mg/kg adriamycin and the abundance of glomerular IgM was examined (Fig.20). Injection of the mice with adriamycin caused a significant increase in the abundance of glomerular IgM. Pre-treatment of the mice with anti-CD20 prevented an increase in glomerular IgM after injection with adriamycin, and levels of glomerular IgM in mice that received both anti-CD20 and adriamycin were similar to those seen in healthy controls. To determine whether IgM eluted specifically from kidneys with adriamycin nephropathy would bind to glomerular epitopes when re-injected into mice, IgM was purified from the kidneys of adriamycin treated mice and then injected into mice with adriamycin nephropathy that had previously undergone B cell depletion with the anti-CD20. This was repeated every week during the course of the study. A trend towards greater glomerular IgM was seen in the reconstituted mice (Fig. 20A and B). In mice with adriamycin nephropathy, low levels of glomerular C4d deposition were detected in control mice but significantly increased in mice treated with adriamycin (Fig. 21A). Treatment of mice with anti-CD20 reduced glomerular C4d, confirming that the glomerular C4d deposits were caused by immunoglobulin-mediated complement activation. Immunostaining of kidneys for C3 deposits showed a similar pattern. C3 deposition increased in mice after treatment with adriamycin, and treatment of the mice with anti-CD20 prevented this increase (Fig. 21B). Urine albumin/creatinine ratios were measured as a marker of glomerular injury. Mice with adriamycin nephropathy were grossly albuminuric (Fig. 22A), and treatment of the mice with anti-CD20 reduced the degree of albuminuria in mice with adriamycin nephropathy. When anti-CD20 treated mice were treated with adriamycin

and then re-injected with purified glomerular IgM, however, the degree of albuminuria was comparable to that seen in mice that received adriamycin alone. This demonstrated that IgM is an important mediator of injury in this model. There was less glomerulosclerosis in mice treated with the anti-CD20 antibody (Figure 22B-C). Mice with adriamycin demonstrated greater glomerular collagen IV deposition than control mice, but treatment with anti-CD20 did not significantly reduce glomerular collagen IV deposition (Figure 22D). Natural antibody IgM is predominantly produced by peritoneal B-1a cells. Treatment of mice with anti-CD20 depletes peritoneal B cells, although less effectively than it depletes splenic B cells. In order to effectively reduce the peritoneal B cells without affecting splenic B cell function, peritoneal cells were lysed by hypotonic shock. This treatment caused lysis of all peritoneal cells, including macrophages, but it did not affect splenic B cells. Reduction in circulating B cells was observed using this strategy, suggesting that the peritoneal compartment is the source of some circulating B cells. Although all peritoneal cells were reduced by this method, the treatment did not reduce the number of B-1a cells as a percentage of total peritoneal cells. Consequently, the effects of this treatment may not have been specific to depletion of B-1a cells. The level of total IgM in the serum was unaffected by this treatment. In order to deplete the peritoneal B cells in mice with adriamycin nephropathy, the peritoneal cells were lysed by hypotonic shock every three days for two weeks prior to injection with adriamycin in order to give sufficient time for pre-formed IgM to turn over. Depletion of the cells was maintained by intra-peritoneal injections of distilled water every three days during the course of the study. Although this treatment is very effective at immediately reducing the peritoneal B cell numbers, the cells re-accumulated and the depletion at day 14 was not complete. As controls, another group of mice received peritoneal injections with PBS according to the same schedule. When reported as a percentage of total peritoneal cells the B-1a cells were not reduced by this treatment because the treatment reduced all peritoneal cells equally. In mice that underwent peritoneal B cell depletion, the accumulation of glomerular IgM was significantly attenuated after injection of the mice with adriamycin (Fig. 23A). Depletion of the peritoneal cells showed a trend towards reduction of glomerular C4 deposition, and glomerular C3 deposition was reduced by this treatment (Fig. 23B-C). Tubulointerstitial C3 deposition was unaffected by this treatment. These results demonstrated that peritoneal B cells generated a significant proportion of the IgM

deposited in the glomeruli in this model. Levels of circulating IgM were not affected by the peritoneal cell depletion even though glomerular deposition was reduced, suggesting that the glomerular IgM deposits were comprised of IgM that bound to specific glomerular antigens. As with treatment of mice with anti-CD20, depletion of the peritoneal B cells reduced glomerular complement activation. Urine albumin/creatinine ratios were measured in samples collected 1 or 4 weeks after injection with adriamycin (Fig. 24A). The level of albuminuria was significantly lower in mice that had undergone peritoneal cell depletion at both the early and the later time-point. The reduction in albuminuria at the 1-week time-point indicated that glomerular IgM contributed to injury at an early phase of disease in this model. Some glomeruli in the mice that underwent peritoneal cell depletion appeared normal (Fig. 24B). As was seen in mice treated with anti-CD20, however, depletion of the peritoneal B cells did not significantly attenuate the overall degree of collagen IV accumulation in adriamycin treated mice, however there was less glomerulosclerosis (Fig. 24B-D). Previous series have reported that glomerular IgM and/or C3 may be detected in up to 90% of patients with FSGS. The biopsy reports of 174 cases of focal segmental glomerulosclerosis (FSGS) evaluated over the past eight years was reviewed. Of those biopsies, approximately 23% demonstrated glomerular IgM without C3, approximately 2% demonstrated glomerular C3 without IgM, and 7% of the biopsies demonstrated glomerular IgM and C3. Dual staining for IgM and for C3d on tissue in which both factors were detectable was performed (Fig. 25A). C3d and IgM showed a similar pattern of distribution throughout the glomerulus. Dual staining for IgM and C4 was also performed (Fig. 25B). This staining demonstrated similar co-localization of these factors within the glomerulus.

[0501] Sham-treated mice and mice subjected to renal IR injury were examined for tissue deposition of IgG and IgM. See Renner et al., *J. Immunol.*, (2010), 185:4393-440 for a description for the renal IR injury mouse model. Briefly, mice were anesthetized with 300 μ l 2,2,2-Tribromoethanol (Sigma-Aldrich) by intraperitoneal injection. Laparotomies were performed, and the renal pedicles were located and isolated by blunt dissection. The pedicles were clamped with surgical clips (Miltex Instrument, Bethpage, NY), and occlusion of blood flow was confirmed by visual inspection of the kidneys. The clamps were left in place for 24 min and then released. The kidneys were observed for 1 min to ensure blood reflow, and then fascia and skin were sutured with 4-0 silk (U.S. Surgical, Norwalk, CT). Sham surgery was

performed in an identical manner, except that the renal pedicles were not clamped. The mice were volume resuscitated with 0.5 ml normal saline subcutaneous injection. After 8, 24, 48, or 72 h of reperfusion, the mice were anesthetized, and blood was obtained by cardiac puncture. Laparotomy was performed, and the kidneys were harvested. IgG was not seen in any of the kidneys, but IgM was seen in the mesangium of sham-treated kidneys (Fig. 26A and C) and kidney that underwent IR (Fig. 26B and D). The level of mesangial IgM increased after renal I/R (Fig. 26E-F), suggesting that circulating IgM bound to the mesangium during reperfusion. IgM was not seen along the tubular basement membrane. Peritoneal B cells in wild-type mice were depleted for 2 weeks by injecting distilled water into the peritoneum, and control mice were injected with an equal volume of PBS. Depletion of the B-1 population was confirmed by flow-cytometry analysis of B220, CD5, and CD19. Lysis of the peritoneal B-1 cells did not alter the overall levels of circulating IgM, but it did reduce levels of mesangial IgM after sham treatment and after renal I/R compared with control mice. Depletion of peritoneal B-1 cells was also associated with a significantly attenuated increase in serum urea nitrogen (SUN) after 24 h of reperfusion. SUN levels were not significantly different from control mice by 48 hours of reperfusion. Although the decrease in mesangial IgM was associated with protection of renal function, mice that underwent peritoneal B cell depletion still demonstrated tubular necrosis comparable to that seen in wild-type mice.

[0502] The role of self-reactive C2 or B4 mAbs in a renal IR injury model was investigated. See Renner et al., *J. Immunol.*, (2010), 185:4393-440 for a description for the renal IR injury mouse model. Briefly, mice were anesthetized with 300 μ l 2,2,2-Tribromoethanol (Sigma-Aldrich) by intraperitoneal injection. Laparotomies were performed, and the renal pedicles were located and isolated by blunt dissection. The pedicles were clamped with surgical clips (Miltex Instrument, Bethpage, NY), and occlusion of blood flow was confirmed by visual inspection of the kidneys. The clamps were left in place for 24 min and then released. The kidneys were observed for 1 min to ensure blood reflow, and then fascia and skin were sutured with 4-0 silk (U.S. Surgical, Norwalk, CT). Sham surgery was performed in an identical manner, except that the renal pedicles were not clamped. The mice were volume resuscitated with 0.5 ml normal saline subcutaneous injection. After 8, 24, 48, or 72 h of reperfusion, the mice were anesthetized, and blood was obtained by cardiac puncture. A 100 mcg dose of B4 mAb was injected into a B cell deficient (Mu) mouse subjected

to unilateral renal ischemia reperfusion and kidneys were harvested for immunohistochemistry analysis. In this model, B4 mAb and C2 mAb localized in glomeruli consistent with the ability of B4 mAb to recognize neoepitopes due to ischemia (Fig. 27A and B).

[0503] Example 8: In vivo study of IgM effects in a systemic factor H deficiency mouse model of renal disease.

[0504] Patients with systemic factor H deficiency primarily develop renal disease with complement deposition within the glomerulus. Factor H knockout mice develop similar disease with glomerular deposition of C3 and albuminuria. See Pickering et al., *Nature Genetics.*, (2002)., 31:424-28. The role of natural IgM deposition and complement activation in the fH knockout model (fH^{-/-} mice), which is not an immune complex model, was investigated. Double knockout mice that lacked factor H and B (fH/ μ MT mice) were also generated and studied. The kidneys were harvested from wild-type, fH^{-/-}, and fH/ μ MT mice for immunohistochemistry analysis. Immunofluorescence staining demonstrated increased glomerular C3 deposition in fH/ μ MT and fH^{-/-} mice as compared to wild-type mice (Fig. 28A and B). Immunofluorescence staining also demonstrated increased glomerular IgM deposition in fH^{-/-} mice as compared to wild-type mice which increased with age (Fig. 29A and B) which followed C3 deposition (Fig. 30A and B).

Immunofluorescence staining of glomerulus from fH/ μ MT showed C3 deposition but no IgM deposition (Fig. 30C). Ultrastructural analysis of immunofluorescence images of glomerulus from fH^{-/-} mice showed co-localization with a marker for the glomerular basement membrane (Fig. 31). Additional immunofluorescence staining of glomerulus from fH^{-/-} mice showed that both C3 and C4 were deposited (Fig. 32). Histopathology section of the kidneys from fH^{-/-} mice showed basement membrane thickening and foot process effacement. Serum urea nitrogen (SUN) and creatinine (Cr) were similar among the wild-type, fH^{-/-}, and fH/ μ MT mice (Fig. 33A). However, fH^{-/-} mice developed albuminuria while fH/ μ MT mice showed protection from albuminuria (Fig. 33B).

[0505] To identify the deposition sites of IgM, an in vitro experiment was conducted by incubating a murine mesangial cell line with wild-type mouse serum. The cells were then analyzed by flow cytometry for the presence of C3, C4, IgM, and IgG. Flow cytometric analysis demonstrated that IgM but not IgG bound to mesangial cells

(Fig. 34A and B). Furthermore, both C3 and C4 were bound by mesangial cells (Fig. 34C and D).

[0506] In order to characterize the glomerular epitopes, monoclonal natural IgM antibodies were screened for their ability to bind to mesangial cells. Two of the monoclonal antibodies tested, C2 and F632, bound to mesangial cells in vitro (Figure 34E). The five other clones that were tested did not bind to the mesangial cells (Figure 34F). The binding of these mAbs to mesangial cells from C57BL/6 mice that were grown in primary culture and to conditionally immortalized mesangial cells that were developed from H-2Kb-tsA58 transgenic mice was also tested. C2 and F632 also bound to these mesangial cells lines, whereas the other IgM clones did not. Next, the ability of these antibodies to bind in vivo was tested. Three μ MT mice were each injected with 100 μ g of mAb C2, F632, or D5 (D5 did not bind to mesangial cells in vitro). Deposits of IgM were seen in the kidneys of all of the mice injected with C2 (Figure 34, panels G and H) and F632, but no deposits were seen in the mice injected with D5 (Figure 34, panels I and J). The identical pattern of mAb binding seen with all three mesangial cell types and with the in vivo experiment supported the specificity of IgM binding in these assays. It also suggested that the target antigen may be the same in all of these assays.

[0507] Example 9: In vivo study of IgM effects in an ischemic reperfusion mouse model of arthritis.

[0508] The role of natural IgM antibodies in an ischemic reperfusion mouse model of arthritis was investigated. In these experiments, passive arthritis was induced by intravenous transfer of a submaximal dose of a cocktail of monoclonal antibodies to CII (Arthrogen-CIA®, Chemicon) and/or the monoclonal antibody B4 mAb as well as purified total IgM from wild-type mice. An IgM monoclonal antibody to trinitrophenol-KLH (anti-TNP; BD PharMingen) was administered as a negative control. Arthrogen was titrated to determine the dose that would yield sub-maximal disease in animals for use in combination with test and control antibodies. An intraperitoneal injection of 50 micrograms/mouse of LPS followed three days after administration of each antibody. From days 1 through 14 after the initial transfer, mice were scored daily by an individual blinded to their treatments for signs of arthritis in the paws based on the following scale: 0=no redness or swelling, 1=one digit swollen, 2=two digits swollen, 3=three digits affected, and 4=entire paw swollen

with ankylosis. The scores for each of four paws of a mouse were totaled to give a final score with a maximal severity of 16. Analysis of arthritic signs demonstrated that administration of B4 mAb resulted in a significantly greater arthritic score than poly IgM or a submaximal dose of the cocktail of arthritis-inducing monoclonal antibodies, Arthrogen (Fig. 35).

[0509] Example 10: In vitro and in vivo study of IgM effects in mouse model of age-related macular degeneration.

[0510] Methods and Materials

[0511] Reagents

[0512] Pooled normal human serum (NHS) from Quidel Corporation (Santa Clara, CA) was used as a source of complement. To probe the involvement of the classical pathway, complement C2- and C4-depleted serum as well as complement proteins C2 and C4 were purchased from Complement Technology, Inc. (CompTech; Tyler, TX), and C1q-depleted serum was contributed by Deborah Fraser, (University of California Irvine, Irvine, CA) (Fraser, D. A., Laust, A. K., Nelson, E. L., and Tenner, A. J. (2009) *J Immunol* 183, 6175-6185). To probe the involvement of the lectin pathway, recombinant M-ficolin, H-ficolin, and mannan-binding lectin (MBL)-associated serine protease 2 (MASP-2) (Frederiksen, P. D., Thiel, S., Larsen, C. B., and Jensenius, J. C. (2005) *Scand J Immunol* 62, 462-473; Thiel, S., Kolev, M., Degn, S., Steffensen, R., Hansen, A. G., Ruseva, M., and Jensenius, J. C. (2009) *J Immunol* 182, 2939-2947; Zacho, R. M., Jensen, L., Terp, R., Jensenius, J. C., and Thiel, S. (2012) *J Biol Chem* 287, 8071-8081), purified L-ficolin (Lacroix, M., Dumestre-Perard, C., Schoehn, G., Houen, G., Cesbron, J. Y., Arlaud, G. J., and Thielens, N. M. (2009) *J Immunol* 182, 456-465) and recombinant MBL (Jensenius, J. C., Jensen, P. H., McGuire, K., Larsen, J. L., and Thiel, S. (2003) *Biochem Soc Trans* 31, 763-767) were used. Finally, to determine the requirement for the alternative pathway, factor B-depleted serum was purchased from CompTech. To analyze the involvement of immunoglobulins (Ig) in triggering the lectin pathway, Ig-depleted serum was obtained from Sunnyslab (Sittingbourne, UK) or collected from rag1^{-/-} mice; and reconstitution experiments were performed by adding back antigen-specific (IgM-C2) (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* 188, 1460-1468) or control IgMs (F1102; raised against 4-Hydroxy-3-nitrophenylacetyl hapten conjugated to KLH and purified in the

same way as IgM-C2). Primary antibodies included a mouse monoclonal anti-MBL from Millipore (Billerica, MA), a rabbit polyclonal anti-MBL from Abcam (Cambridge, MA), a rabbit polyclonal MASP-2, a goat antibody against human C3 (CompTech), and monoclonal antibodies to human H-Ficolin, L-Ficolin and M-Ficolin from Santa Cruz Biotechnology (Santa Cruz, CA). Species-specific secondary antibodies were from Zymed Laboratories (Invitrogen; Carlsbad, CA).

[0513] Mice and collection of serum

[0514] C57BL/6 (B6) and B6 rag1^{-/-} mice were generated from breeding pairs (Jackson Laboratory; Bar Harbor, ME). For collection of serum, mice were deeply anesthetized (ketamine/xylazine, 80/10 mg/kg). Blood was collected in BD vacutainer tubes by cardiac puncture and serum was collected after clot formation (2 hours on ice) and centrifugation (1000-1400 rcf, 4°C for 10 min).

[0515] Cell culture

[0516] ARPE-19 cells were expanded in Dulbecco's modified Eagle's medium F12 (Invitrogen) with 10% fetal bovine serum (FBS) and antibiotics as described before (Thurman, J. M., Renner, B., Kunchithapautham, K., Ferreira, V. P., Pangburn, M. K., Ablonczy, Z., Tomlinson, S., Holers, V. M., and Rohrer, B. (2009) *J Biol Chem* 284, 16939-16947). Human fetal retinal pigment epithelium (RPE) cells were prepared and expanded in Minimum Essential Medium (MEM; Sigma-Aldrich, St. Louis, MO) with 15% FBS, following our published protocol (Bandyopadhyay, M., and Rohrer, B. (2012) *Invest Ophthalmol Vis Sci* 53, 1953-1961). Globes were supplied by Advanced Bioscience Recourses (Alameda, CA), and experiments adhered to the Declaration of Helsinki on the ethical principles for medical research involving human materials.

[0517] Transepithelial resistance (TER) measurements

[0518] ARPE-19 cells or human fetal RPE were grown as mature monolayers on 6-well Transwell inserts (Corning, 0.4 µm PET, 24 mm insert) in the presence of 5% FBS for 2-3 weeks (Ablonczy, Z., and Crosson, C. E. (2007) *Exp Eye Res* 85, 762-771). For the final 2-3 days prior to the experiments, cells were changed to serum-free media. Complement activation was induced as reported previously (Thurman, J. M., Renner, B., Kunchithapautham, K., Ferreira, V. P., Pangburn, M. K., Ablonczy, Z., Tomlinson, S., Holers, V. M., and Rohrer, B. (2009) *J Biol Chem* 284, 16939-16947), exposing cells to 0.5 mM H₂O₂ in the presence of 10% normal human serum (NHS). It was previously shown that sublytic complement activation results in VEGF

release, which in turn reduces barrier function (Thurman, J. M., Renner, B., Kunchithapautham, K., Ferreira, V. P., Pangburn, M. K., Ablonczy, Z., Tomlinson, S., Holers, V. M., and Rohrer, B. (2009) *J Biol Chem* 284, 16939-16947), TER measurements are a convenient readout for the level of activity in the complement cascade. TER was determined by measuring the resistance across the monolayer with an EVOM volt-ohmmeter (World Precision Instruments, Sarasota, FL). The value for cell monolayers was determined by subtracting the TER for filters without cells and the percentage calculated using the starting value as reference.

[0519] Binding assays

[0520] For testing the binding of ficolin, non-specific IgM or antigen-specific IgM (IgMC2) binding, ARPE-19 cells were grown as monolayers in 96-well plates. Cells were changed to serum-free media 2 days before the experiments. To identify ficolin binding, normal human serum was used as the source of ficolins. Cells were incubated with serial dilutions of serum for 1 hour at 37°C, washed and fixed in PBS containing 4% paraformaldehyde, and non-specific binding sites were blocked with 1% BSA in PBS. Bound ficolins were detected with corresponding antibodies followed by alkaline phosphatase conjugated secondary antibody and color development using the pNPP phosphatase substrate system (KPL; Gaithersburg, MD). To characterize non-specific IgM or antigen-specific IgM (IgM-C2) binding to control and oxidatively-stressed ARPE-19 cells, neo-epitopes were generated by exposing cells to 0.5 mM H₂O₂ for 10 minutes prior to incubation with serial dilutions of serum (for detection of IgM binding) or C2 antibody (in PBS). Bound IgMs were detected with alkaline phosphatase conjugated secondary antibody and color development using the pNPP phosphatase substrate system.

[0521] Depletion of MBL

[0522] MBL-depleted human serum was prepared using mannan-agarose (Sigma-Aldrich, St. Louis, MO) as depleting beads according to published protocols (Rajagopalan, R., Salvi, V. P., Jensenius, J. C., and Rawal, N. (2009) *Immunol Lett* 123, 114-124). A 2.0 mL column of mannan-agarose was prepared and equilibrated with Veronal buffer (Lonza; Allendale, NJ) containing calcium chloride (3 mM) and magnesium chloride (10 mM). Normal human serum was passed through the column

and the flow through was collected. The depletion of MBL was confirmed by ELISA and Western blot.

[0523] MBL ELISA

[0524] Microtiter (Immulon2; Dynatech Laboratories, Chatilly, VA) plates were first coated with 10 µg/mL polyclonal rabbit anti-MBL capture antibody overnight at 4°C. The plates were then washed three-times with PBS and blocked with 3% milk in PBS for 1 hr at room temperature, followed by exposure to the antigen (normal human serum or MBL-depleted human serum) for 2 hrs at 37°C. The plates were again washed and incubated with monoclonal antibody to MBL followed by peroxidase conjugated secondary antibody and color development using Turbo-TMB ELISA (Pierce; Thermo Scientific, Rockford, IL).

[0525] Characterization of IgM-C2 epitopes

[0526] ELISAs to determine reactivity of the IgM-C2 to phospholipids were performed as described (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* 188, 1460-1468). In short, microtiter plates (Immulon 1B) were coated with 100 µL/well 50 µg/mL phospholipid in methanol. After the plates were air-dried, the wells were washed with PBS and blocked with 1% BSA. IgM was added to wells and bound-Ab detected by alkaline-phosphatase-conjugated goat anti-mouse IgM (Jackson ImmunoResearch Laboratories, West Grove, PA). Relative units of Ab were calculated by comparing OD at 405 nm for individual titrated serum with a standard curve of OD measurements established previously (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* 188, 1460-1468). Binding of IgM-C2 was compared against a control IgM known to cross-react with annexin IV (IgM-B4) (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* 188, 1460-1468). Two phospholipids were assayed; phosphorylcholine (PC)(10)-BSA (Biosearch Technologies, Novato, CA) and malondialdehyde (MDA)-BSA (Cell Biolabs, San Diego, CA).

[0527] SDS-polyacrylamide gel electrophoresis and Western blotting

[0528] Protein concentration was determined by BCATM protein assay according to the manufacturer's instructions (Pierce, Rockford, IL). For each sample, 40 µg of

protein was denatured, subjected to SDS-polyacrylamide gel electrophoresis and analyzed by immunoblotting with appropriate antibodies.

[0529] Immunofluorescence staining

[0530] Surface exposure of phospholipid-specific epitopes was examined by immunofluorescence microscopy. ARPE-19 cells were grown on 35-mm lysine-coated glass-bottom culture dishes (MatTek Corporation; Ashland, MA), treated with H₂O₂ for 10 minutes, fixed in PBS containing 4% paraformaldehyde and nonspecific binding sites were blocked with 1% normal goat serum and 3% BSA in PBS (preabsorption buffer) for 2 hours. The cells were incubated overnight at 4°C with either rabbit anti-MDA polyclonal antibody (1:200 in PBS) followed by incubation for 1 hr at room temperature with FITC-conjugated goat anti-rabbit IgG (1:200; Zymed Laboratories, Invitrogen); or with IgM natural antibody (IgM-C2) followed by goat anti-mouse IgM (1:200; Zymed Laboratories, Invitrogen). As a negative control, primary antibodies were omitted. Staining was also performed ex vivo on eyes with CNV lesions (see below). Eyecups were fixed in 4% paraformaldehyde, washed, preabsorbed and incubated over night at 4°C with C2-IgM (1:200), followed by anti-mouse IgM (1:200) as described above. Omission of primary antibody staining served as the negative control. Staining of cells and flatmounts was examined by confocal microscopy (Olympus FluoView).

[0531] In vivo CNV induction and assessment

[0532] B6 rag1^{-/-} and C57BL/6 mice were housed in the Medical University of South Carolina animal care facility under a 12:12 hour light:dark cycle with access to food and water ad libitum. All experiments were performed in accordance with the Association for Research in Vision and Ophthalmology and were approved by the Institutional Animal Care and Use Committee. CNV lesions (four spots in each eye surrounding the optic nerve) were generated as described previously using argon laser photocoagulation (532 nm; 100 µm spot size; 0.1 s duration; 100 mW) (Rohrer, B., Long, Q., Coughlin, B., Wilson, R. B., Huang, Y., Qiao, F., Tang, P. H., Kunchithapautham, K., Gilkeson, G. S., and Tomlinson, S. (2009) *Invest Ophthalmol Vis Sci* 50, 3056-3064). Animals (n=6-8 per treatment group) were treated on days 0, 2 and 4 with C2-IgM or control F1102-IgM (100 µg diluted in 400 µL PBS) using intraperitoneal (IP) injections. IP injections have been shown to be effective for antibody delivery to CNV lesions (Campa, C., Kasman, I., Ye, W., Lee, W. P., Fuh, G., and Ferrara, N. (2008) *Invest Ophthalmol Vis Sci* 49, 1178-1183). Relative CNV

size was determined in flatmount preparations of RPE-choroid stained with ICAM2 (Campa, C., Kasman, I., Ye, W., Lee, W. P., Fuh, G., and Ferrara, N. (2008) *Invest Ophthalmol Vis Sci* 49, 1178-1183). Staining, flatmounting, imaging and analysis of fluorescence measurements by confocal microscopy were performed as reported previously (Rohrer, B., Long, Q., Coughlin, B., Wilson, R. B., Huang, Y., Qiao, F., Tang, P. H., Kunchithapautham, K., Gilkeson, G. S., and Tomlinson, S. (2009) *Invest Ophthalmol Vis Sci* 50, 3056-3064). Data are expressed as mean $\pm$ SEM per eye.

[0533] Statistics

[0534] For data consisting of multiple groups, one-way ANOVA followed by Fisher's post hoc test ($P < 0.05$) was used; single comparisons were analyzed by Student t test analysis ($P < 0.05$).

[0535] Sublytic complement activation as a function of complement status.

[0536] Using a combination of serum-depletion strategies, complement activation pathways involved in TER reduction were analyzed (Fig 36A). ARPE-19 cells grown as monolayers on Transwell filters develop TER levels of 40-45 Ω /cm², a value that is not affected over the course of a 4-hour exposure to 0.5 mM of H₂O₂ or 10% of normal human serum (NHS). However, the co-treatment with H₂O₂ + NHS reduced TER by >40% (i.e., resulting in <60% baseline TER values; $P < 0.001$). TER reduction in control serum did not differ significantly from that elicited in the presence of C1q-depleted serum, whereas both MBL- or factor B-depleted serum were found to be ineffective in reducing TER (i.e., resulting in ~95% baseline TER values after 4 hours of exposure; n.s.). Taken together, these results allow the conclusion that the lectin pathway is responsible for triggering the complement attack on oxidatively-stressed RPE cells, followed by the amplification by the alternative pathway.

[0537] The typical activity of the lectin pathway serine protease MASP-2 is to split complement C2 and C4 into their respective a- (C2a and C4a) and b-components (C2b and C4b), resulting in the formation of the C3 convertase (C4b2a complex) (Takahashi, M., Mori, S., Shigeta, S., and Fujita, T. (2007) *Adv Exp Med Biol* 598, 93-104). However, recently, bypass mechanisms have been observed; MASP-2 has been shown to activate the alternative pathway via a C2-bypass pathway (Tateishi, K., and Matsushita, M. (2011) *Microbiol Immunol* 55, 817-821), and Schwaeble and colleagues have described a lectin pathway-dependent C4-bypass (Schwaeble, W. J., Lynch, N. J., Clark, J. E., Marber, M., Samani, N. J., Ali, Y. M., Dudler, T., Parent,

B., Lhotta, K., Wallis, R., Farrar, C. A., Sacks, S., Lee, H., Zhang, M., Iwaki, D., Takahashi, M., Fujita, T., Tedford, C. E., and Stover, C. M. (2011) *Proc Natl Acad Sci U S A* 108, 7523-7528). Removing C2 or C4 from normal human serum attenuated the effect of H₂O₂ + serum on TER (~80% baseline TER values) (Fig 36B), although not to the level of MBL-depleted serum (~95% baseline TER values). Addition of physiological levels of C2 (10 µg/mL) and C4 (400 µg/mL) to their respective depleted serum reconstituted the effect of serum on TER in both C2- and C4- depleted sera (Fig 36B). Based on the partial effect on TER reduction of the C2- and C4-depleted sera, the results suggest that while activity in the lectin pathway involves the generation of the regular C4b2a complex, a contribution by MASP-2 mediated activation of the alternative pathway cannot be excluded as has been described for MASP-1 and MASP-3 (Banda, N. K., Takahashi, M., Takahashi, K., Stahl, G. L., Hyatt, S., Glogowska, M., Wiles, T. A., Endo, Y., Fujita, T., Holers, V. M., and Arend, W. P. (2011) *Mol Immunol* 49, 281-289).

[0538] Pattern recognition molecules in the Lectin Pathway

[0539] Complement activation requires the binding of a ligand by a pattern recognition molecule. For the lectin pathway, those entry molecules are mannan-binding lectin (MBL) and the ficolins (H-ficolin, L-ficolin and M-ficolin), which then activate the MBL-associated serine protease, MASP-2. Here, a combination of binding assays and reconstitution experiments was employed to determine which pattern recognition molecule could be employed to recognize ligands on oxidatively-stressed RPE cells to activate the lectin pathway.

[0540] In the blood, MBL or ficolin are both complexed with inactive MASP; thus, if MBL is removed from serum using a mannan-agarose column, MASP levels might also be affected. In addition L-ficolin has been shown to bind directly to cyanogen-activated Sepharose beads (Tan, S. M., Chung, M. C., Kon, O. L., Thiel, S., Lee, S. H., and Lu, J. (1996) *Biochem J* 319 (Pt 2), 329-332), a problem that might also apply to H-ficolin and Mficolin. Western blot analysis confirmed that serum passed over a mannan-agarose column is depleted of MBL, MASP-2, M-ficolin and L-ficolin, or levels are below detection level, whereas H ficolin levels were drastically reduced (Fig 37A). As a positive control, complement C3 levels were unaffected by the depletion process (Fig 37A). To narrow down which ficolins recognize binding sites on ARPE-19 cells, monolayers were exposed to serial dilutions of serum in the presence 0.5 mM H₂O₂ (1 hr at 37°C) and bound ficolins were detected using

subtype-specific antibodies (Fig 37B). In oxidatively-stressed cells, M- and H-ficolin binding was found to be saturable at physiological concentrations (mean serum concentrations of ficolins are M-ficolin 1.1, L-ficolin 3.3, and H-ficolin 18.4 $\mu\text{g/mL}$; (Zacho, R. M., Jensen, L., Terp, R., Jensenius, J. C., and Thiel, S. (2012) *J Biol Chem* 287, 8071-8081)), whereas L-ficolin binding was not saturable, but did appear to bind. However, since L-ficolin has been shown to bind non-specifically to BSA in solid-phase binding assays, a possible contribution of L-ficolin cannot be ruled out completely (Faro, J., Chen, Y., Jhaveri, P., Oza, P., Spear, G. T., Lint, T. F., and Gewurz, H. (2008) *Clin Exp Immunol* 151, 275-283).

[0541] Thus, the flow-through from the mannan-agarose column was utilized for reconstitution experiments, comparing MBL with M- and H-ficolin for their ability to reconstitute activity in the TER assay (Fig 37C). Exposure of monolayers to H_2O_2 + MBL-depleted serum resulted in ~95% baseline TER values after 4 hours of exposure, whereas H_2O_2 + NHS reduced TER to <60% of baseline values. Addition of MASP-2 (50 ng/mL) in combination with either one of the three pattern recognition molecules at physiological levels increased activity in the TER assay ($P < 0.05$) to levels that were not significantly different from those of complete normal human serum. Adding MBL to either M- or H-ficolin or both did not further increase the activity. Thus, the lectin pathway can be activated in oxidatively-stressed cells by either MBL/MASP or ficolin/MASP.

[0542] Activation of the lectin pathway by natural IgM

[0543] The lectin pathway has historically been recognized as a pathway that is activated by ficolin/MASP or MBL/MASP recognizing specific carbohydrates or acetylated molecules on pathogen surfaces; however, more recently, it has been shown that IgM molecules recognizing epitopes generated during ischemia reperfusion injury can activate the lectin pathway (Zhang, M., Takahashi, K., Alicot, E. M., Vorup-Jensen, T., Kessler, B., Thiel, S., Jensenius, J. C., Ezekowitz, R. A., Moore, F. D., and Carroll, M. C. (2006) *J Immunol* 177, 4727-4734). This experiment demonstrates that ficolin/MASP or MBL/MASP requires immunoglobulins, and specifically natural IgM to initiate complement activation on the cell surface of RPE cells.

[0544] TER reduction was tested in complement-sufficient serum (mouse or human) and compared to serum from which Igs had been eliminated either genetically (rag1-/- mice) or by depletion (human Ig-depleted serum) (Fig 38A). Both human and mouse

complement-sufficient sera reduced TER by 40-50%, whereas Ig-depleted serum was ineffective. To characterize non-specific IgM binding to control and oxidatively-stressed ARPE-19 cells, cells were exposed to serial dilutions of serum followed by detection of bound IgMs using an IgM-specific secondary antibody coupled to alkaline phosphatase. However, just like for the ficolins and MASP, binding under both conditions was indistinguishable (Fig 38B). When epitope-specific IgM-C2 binding to control and oxidatively-stressed ARPE-19 cells was compared, exposing cells to serial dilutions of IgM-C2 antibody followed by colorimetric detection of bound IgMs, ~2.5-fold higher binding of IgM-C2 could be documented when comparing control and oxidative-stress conditions (Fig 38C). Addition of the IgM-C2, but not a control antibody (IgM F1102, raised against dinitrophenol) was able to reconstitute activity to levels indistinguishable from normal human serum (Fig 38D).

[0545] Identification of ligands for natural IgMs on RPE Cells

[0546] Cell binding assays have shown that IgM-C2 binding and injury is augmented under oxidative-stress conditions (Fig 38C). Oxidative damage of membrane phospholipids results in the formation of malondialdehyde (MDA), an end-product of lipid peroxidation. The natural ligand of IgM-C2 on oxidatively-stressed RPE cells was further characterized.

[0547] ELISAs to determine the reactivity of IgM-C2 to ligands were performed using microtiter plates coated with phosphorylcholine (PC)-BSA or malondialdehyde (MDA)-BSA in methanol. IgMC2 recognized both PC and MDA (Fig 39A, B). To determine which ligand is relevant for IgM-C2 binding to oxidatively-stressed RPE cells, IgM-C2 was preabsorbed with either MDA-BSA or PCBSA. IgM-C2 preabsorbed with either MDA-BSA or PC-BSA completely abolished its ability to reconstitute Ig-depleted serum (Fig 39C,D). MDA and C2-IgM neoepitopes could be identified in a punctate fashion on H₂O₂-treated cells when compared to control cells. Both the anti-MDA antibody and the IgM-C2 antibody recognize epitopes present in puncta across the apical surface of the ARPE-19 cells (Fig 40).

[0548] To confirm that the same neoepitopes are generated in primary human RPE cells, primary fetal RPE cells were grown into monolayers with a stable TER level of 250-300 cm². These monolayers are susceptible to complement attack (Fig 41), albeit not to the same degree as ARPE-19 cells. The combined treatment of H₂O₂ + 10% NHS significantly decreased TER (P<0.001), which was attenuated by the elimination of all immunoglobulins (Ig-depleted serum; P<0.01). IgM-C2 (P<0.01),

but not IgM-F1102 was able to reconstitute Ig-depleted serum, an effect that could be eliminated by preabsorption with MDA-BSA.

[0549] MDA has recently been shown to bind CFH. In a human acute monocytic leukemia cell line, malondialdehyde-acetaldehyde-BSA elicited a proinflammatory response as determined by IL-8 secretion, which was inhibited by physiological concentrations of CFH. If MDA represents a ligand for CFH in ARPE-19 cells, physiological concentrations of CFH might inhibit complement activation to prevent TER reduction. Since all the experiments thus far have been executed with 10% NHS, the experiments were repeated with higher NHS concentrations as well as in the presence of exogenous CFH. Average CFH concentration in serum is ~500 µg/mL. Hence, TER experiments were performed in the presence of 25% NHS (Fig 42) or 25% NHS supplemented with 375 µg of CFH. Both combinations were unable to prevent TER reduction by H₂O₂ + NHS. To ensure that potential modification of CFH by H₂O₂ does not impair its binding to its ligand on ARPE-19 cells, the supernatant containing H₂O₂ was removed after 5 minutes of stimulation and NHS + exogenous CFH was added, which also did not prevent TER deterioration. However, CR2-fH (at 10 µg/mL), a CFH mimetic that consists of the complement receptor-2 binding domain for C3bi and C3d coupled to the inhibitory domain of CFH was able to inhibit TER deterioration induced by H₂O₂ + NHS as previously reported (Thurman, J. M., Renner, B., Kunchithapautham, K., Ferreira, V. P., Pangburn, M. K., Ablonczy, Z., Tomlinson, S., Holers, V. M., and Rohrer, B. (2009) J Biol Chem 284, 16939-16947). Thus, on ARPE-19 cells, using H₂O₂ as the oxidant stimulus, MDA does not appear to serve as a ligand for CFH.

Identification of roles for IgM-C2 in vivo

[0550] Mouse CNV laser lesions were examined by immunohistochemistry for labeling with the C2-IgM antibody. Specific labeling could be identified in CNV lesions as opposed to the area surrounding the lesion, when compared to the secondary antibody-only control (Fig 43A). The MDA-specific antibody labeling was indistinguishable from that shown previously (Weismann, D., Hartvigsen, K., Lauer, N., Bennett, K. L., Scholl, H. P., Charbel Issa, P., Cano, M., Brandstatter, H., Tsimikas, S., Skerka, C., Superti-Furga, G., Handa, J. T., Zipfel, P. F., Witztum, J. L., and Binder, C. J. (2011) Nature 478, 76-81).

[0551] To address the physiological relevance of the C2-IgM autoantibodies in CNV lesions, it was examined whether C2-IgM reconstitution experiments would alter

injury in *rag1*^{-/-} mice. CNV lesions were examined in *rag1*^{-/-} mice after 3 administrations every 48 hours of either PBS, the F1102-IgM, the F632-IgM, the B4-IgM, or the C2-IgM (100 µg/mouse in 100 µL of PBS). While the control IgM antibodies (F1102-IgM and F632-IgM) had no effect on the size of the CNV lesion when compared to PBS-injected animals, CNV lesions were approximately twice the size after C2-IgM injections ($P < 0.01$) or B4-IgM. Neither C2-IgM or B4-IgM injections had any effect in wildtype mice (Fig 43B).

[0552] The principal findings obtained here, studying complement activation in oxidatively stressed RPE monolayers, can be summarized as follows: (1) oxidative stress generates neoepitopes on RPE cell surfaces that contain phospholipids, including MDA; (2) specific autoantibodies present in normal serum recognize these surface epitopes, and consequently trigger the activation of the complement cascade using the lectin pathway; (3) both MBL and ficolin can serve as the pattern recognition receptors for the lectin pathway; (4) this basal activity generated by the lectin pathway is subsequently amplified by the alternative pathway to generate the maximal effect; and finally, (5) the C2-IgM antibody recognizes neoepitopes in mouse CNV lesions and augments CNV development in antibody-deficient, *rag1*^{-/-} mice.

[0553] In the present example, TER was used as a convenient, rapid and sensitive readout of complement activation. The availability of complement- and Ig-depleted serum as well as purified protein and individual IgMs allowed us to dissect the activation pathway for the terminal complement cascade on these oxidatively-stressed RPE cells. Since the 3 complement pathways require unique entry or activator molecules, pathway-specific serum can be generated (C1q-depleted serum, no CP; MASP-2-depleted serum, no LP; factor B-depleted serum, no AP; and C1q-MASP-2-double-depleted serum, AP-only) (Fig 36). Both the LP and the AP are necessary for generating the loss of TER. AP plays a role in amplifying the complement cascade on oxidatively stressed RPE cells. Since the serum depleted for the LP pathway components was found to be depleted for MASP-2, MBL and all three ficolins, (Fig 37A), reconstitution studies could be performed with the individual pattern recognition molecules. Roles for MBL, M-ficolin and H-ficolin could be demonstrated utilizing the TER assay. Since specific binding of M- and H-ficolin to RPE cells could be demonstrated in the context of complete serum, LP can be activated by the presence of a specific antibody/antigen complex. While Ig-

dependence was demonstrated utilizing Ig-depleted serum (Fig 38A), H₂O₂-dependent changes in total IgM (Fig 38B) or total IgG (data not shown) could not be demonstrated. However, H₂O₂-dependent changes in binding of a phospholipid antigen-specific IgM (IgM-C2) was revealed (Fig 38C), and in reconstitution assays, IgM-C2 antibodies, but not a control antibody, F1102, resulted in the restoration of the effect on TER in Ig-depleted serum (Fig 38D). Control antibody, F1102, when tested in binding assays, also showed no saturable binding to RPE cells (data not shown). Finally, antigen-specificity for C2-IgM was further refined by ELISA (Fig 39A) and TER experiments (Fig 39B, C). Since the original characterization of the ligands (Elvington, A., Atkinson, C., Kulik, L., Zhu, H., Yu, J., Kindy, M. S., Holers, V. M., and Tomlinson, S. (2012) *J Immunol* 188, 1460-1468) was performed in the absence of H₂O₂, but phospholipids can undergo peroxidation, binding was compared between phosphatidylcholine (non-oxidized) and malondialdehyde (MDA; oxidized) coupled to bovine serum albumin (BSA). Specific binding could be documented by ELISA, and preabsorption of the antibody with either MDA-BSA or PC-BSA interfered with activity in the TER assay. H₂O₂-dependent binding could be documented for both a MDA-specific antibody, as well as the neoepitope-specific IgM on RPE cell surfaces (Fig 40). Finally, the neoepitopes recognizable by IgM-C2 and IgM-B4 are also present on oxidatively-stressed primary fetal human RPE cells (Fig 41).

[0554] The results of the Examples show that alterations in barrier function produced by oxidative stress-mediated complement activation requires a phospholipid as a ligand, LP initiation molecules, and alternative pathway amplification, followed by activation of the terminal pathway, including transient membrane attack complex activation. This is the first report that identifies a potential ligand, the pattern recognition receptor, and the pathway required for activation in a model relevant for AMD.

[0555] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is apparent to those skilled in the art that certain minor changes and modifications will be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention.

SEQUENCE CHART

Table 2. Summary of sequences

SEQ ID NO:	Function	Type of Sequence
1.	B4 CDR-L1	Amino acid
2.	B4 CDR-L2	Amino acid
3.	B4 CDR-L3	Amino acid
4.	B4 CDR-H1	Amino acid
5.	B4 CDR-H2	Amino acid
6.	B4 CDR-H3	Amino acid
7.	B4 CDR-L1	Amino acid
8.	B4 CDR-L2	Amino acid
9.	B4 CDR-L3	Amino acid
10.	B4 CDR-H1	Amino acid
11.	B4 CDR-H2	Amino acid
12.	B4 CDR-H3	Amino acid
13.	B4 light chain variable domain	Amino acid
14.	B4 light chain variable domain	Amino acid
15.	B4 heavy chain variable domain	Amino acid
16.	B4 heavy chain variable domain	Amino acid
17.	B4 scFV	Amino acid
18.	B4 scFV	Amino acid
19.	B4 light chain variable domain	Nucleic acid
20.	B4 light chain variable domain	Nucleic acid
21.	B4 heavy chain variable domain	Nucleic acid
22.	B4 heavy chain variable domain	Nucleic acid
23.	B4 scFV	Nucleic acid
24.	B4 scFV, optimized sequence for CHO expression	Nucleic acid
25.	C2 CDR-L1	Amino acid
26.	C2 CDR-L2	Amino acid
27.	C2 CDR-L3	Amino acid
28.	C2 CDR-H1	Amino acid
29.	C2 CDR-H2	Amino acid
30.	C2 CDR-H3	Amino acid
31.	C2 CDR-L1	Amino acid
32.	C2 CDR-L2	Amino acid
33.	C2 CDR-L3	Amino acid
34.	C2 light chain variable domain	Amino acid
35.	C2 light chain variable domain	Amino acid
36.	C2 heavy chain variable domain	Amino acid
37.	C2 scFV	Amino acid
38.	C2 scFV	Amino acid
39.	C2 light chain variable domain	Nucleic acid
40.	C2 light chain variable domain	Nucleic acid
41.	C2 heavy chain variable domain	Nucleic acid
42.	C2 scFV	Nucleic acid
43.	C2 scFV, optimized sequence for CHO expression	Nucleic acid
44.	Full-length human membrane cofactor protein (MCP)	Amino acid
45.	Full-length human decay accelerating factor (DAF)	Amino acid
46.	Full-length mouse decay accelerating factor (DAF)	Amino acid
47.	Full-length human CD59	Amino acid
48.	Full-length mouse CD59	Amino acid
49.	Full-length mouse CD59, isoform B	Amino acid
50.	Full-length mouse Cry	Amino acid
51.	Full-length human CR1	Amino acid
52.	Full-length human factor H	Amino acid

53.	Full-length mouse factor H	Amino acid
54.	Signal peptide of the human CD5 protein	Amino acid
55.	Signal peptide of the human CR2 protein	Amino acid
56.	Signal peptide of the human CR2 protein	Amino acid
57.	B4 scFV	Nucleic acid
58.	C2 scFV	Nucleic acid

SEQUENCES

B4 CDR-L1 amino acid sequence

SSISSNY (SEQ ID NO:1)

B4 CDR-L2 amino acid sequence

RTS (SEQ ID NO:2)

B4 CDR-L3 amino acid sequence

QQGSSIPRTRSEGAPSWK (SEQ ID NO:3)

B4 CDR-H1 amino acid sequence

GYTFTSYW (SEQ ID NO:4)

B4 CDR-H2 amino acid sequence

IGPNSSGT (SEQ ID NO:5)

B4 CDR-H3 amino acid sequence

ARRMVKGCYGLLGPRDHGHRLL (SEQ ID NO:6)

B4 CDR-L1 amino acid sequence

QSIVHSNGNTY (SEQ ID NO:7)

B4 CDR-L2 amino acid sequence

KVS (SEQ ID NO:8)

B4 CDR-L3 amino acid sequence

FQGSHPYPT (SEQ ID NO:9)

B4 CDR-H1 amino acid sequence

GYTFTDYY (SEQ ID NO:10)

B4 CDR-H2 amino acid sequence

INPNNGGT (SEQ ID NO:11)

B4 CDR-H3 amino acid sequence

ARYDYAWYFDV (SEQ ID NO:12)

B4 VL amino acid sequence

DIELTQSPTTMAASPGEKITITCSASSISSNYLHWYQQKPGFSPKLLIYRTSNL
ASGVPARFSGSGSGTSYSLTIGTMEAEDVATYYCQQGSSIPRTRSEGAPSWK
(SEQ ID NO:13)

B4 VL amino acid sequence

DVLMTQTPLSLPVS LGDQASISCRSSQSIVHSNGNTYLEWYLQKPGQSPKLLIY
KVS NRFS GVPDRFS GSGSGTDFTLKISRVEAEDLGVYYCFQGSHVPYTFGGGT
KLEIK (SEQ ID NO:14)

B4 VH amino acid sequence

VKLQESGAELVKPGASVKLSCKASGYTFTSYWMHWVKQRPGRGLEWIGRIG
PNSGGTKYNEKFKSKATLTVDKPSSTAYMQLSSLTSEDSAVYYCARRMVKG
CYGLLGPRDHGHRLL (SEQ ID NO:15)

B4 VH amino acid sequence

VKLQESGP ELVKPGASVKISCKASGYTFTDYYMNWVKQSHGKSLEWIGDINP
NNGGTSYNQKFKGKATLTVDKSSSTAYMELRSLTSEDSAVYYCARYDYAWY
FDVWGQGTTVTVSS (SEQ ID NO:16)

B4 scFV amino acid sequence

HHHHHHV KLQESGAELVKPGASVKLSCKASGYTFTSYWMHWVKQRPGRGL
EWIGRIGPNSGGTKYNEKFKSKATLTVDKPSSTAYMQLSSLTSEDSAVYYCA
RRMVKG CYGLLGPRDHGHRLLKGRIPAHWRPLLVDPSVPSLASGGGGGSG
GGGSWISAEFALDIELTQSPTTMAASPGEKITITCSASSISSNYLHWYQQKPGF
SPKLLIYRTSNLASGVPARFS GSGSGTSYSLTIGTMEAEDVATYYCQQGSSIPR
TRSEGAPSWK (SEQ ID NO:17)

B4 scFV amino acid sequence

MSVPTQVLG LLLLWLTDARCVKLQESGP ELVKPGASVKISCKASGYTFTDYY
MNWVKQSHGKSLEWIGDINPNNGGTSYNQKFKGKATLTVDKSSSTAYMELR
SLTSEDSAVYYCARYDYAWYFDVWGQGTTVTVSSGGGGSGGGGSGGGGDV
LMTQTPLSLPVS LGDQASISCRSSQSIVHSNGNTYLEWYLQKPGQSPKLLIYKV
SNRFS GVPDRFS GSGSGTDFTLKISRVEAEDLGVYYCFQGSHVPYTFGGGT KL
EIKRIEGRHHHHHH (SEQ ID NO:18)

B4 VL nucleic acid sequence

GACATTGAGCTCACCCAGTCTCCAACCACCATGGCTGCATCTCCCGGGGA
GAAGATCACTATCACCTGCAGTGCCAGCTCAAGTATAAGTTCCAATTACTT
GCATTGGTATCAGCAGAAGCCAGGATTCTCCCCTAAACTCTTGATTTATAG
GACATCCAATCTGGCTTCTGGAGTCCCAGCTCGCTTCAGTGGCAGTGGGTC
TGGGACCTCTTACTCTCTCACAATTGGCACCATGGAGGCTGAAGATGTTGC
CACTTACTACTGCCAGCAGGGTAGTAGTATACCACGTACACGTTCCGAGG
GGGCACCAAGCTGGAAA (SEQ ID NO:19)

B4 VL nucleic acid sequence

GATGTTTTGATGACCCAAACTCCACTCTCCCTGCCTGTCAGTCTTGAGAT
CAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCATTGTACATAGTAATGGA

AACACCTATTTAGAATGGTACCTGCAGAAACCAGGCCAGTCTCCAAAGCT
CCTGATCTACAAAGTTTCCAACCGATTTTCTGGGGTCCCAGACAGGTTTCAG
TGGCAGTGGATCAGGGACAGATTTCACTCAAGATCAGCAGAGTGGAGG
CTGAGGATCTGGGAGTTTATTACTGCTTTCAAGGTTACATGTTCCGTACA
CGTTCGGAGGGGGGACCAAGCTGGAAATAAAACG (SEQ ID NO:20)

B4 VH nucleic acid sequence

GTGAACTGCAGGAGTCAGGGGCTGAGCTTGTGAAGCCTGGGGCTTCAGT
GAAGCTGTCCTGCAAGGCTTCTGGCTACACCTTCACCAGCTACTGGATGCA
CTGGGTGAAGCAGAGGCCTGGACGAGGCCTTGAGTGGATTGGAAGGATTG
GTCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAAGAGCAAGGCC
ACACTGACTGTAGACAAACCCTCCAGCACAGCCTACATGCAGCTCAGCAG
CCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGAAGAATGGTAA
AGGGGTGCTATGGACTACTGGGGCCAAGGGACCACGGTCACCGTCTCCTC
A (SEQ ID NO:21)

B4 VH nucleic acid sequence

GTGAAGCTGCAGGAGTCTGGACCTGAGCTGGTGAAGCCTGGGGCTTCAGT
GAAGATATCCTGTAAGGCTTCTGGATACACGTTCACTGACTACTACATGA
ACTGGGTGAAGCAGAGCCATGGAAAGAGCCTTGAGTGGATTGGAGATATT
AATCCTAACAATGGTGGTACTAGCTACAACCAGAAGTTCAAGGGCAAGGC
CACATTGACTGTAGACAAGTCCTCCAGCACAGCCTACATGGAGCTCCGCA
GCCTGACATCTGAGGACTCTGCAGTCTATTACTGTGCAAGATATGATTACG
CTTGGTACTTCGATGTCTGGGGCCAAGGGACCACGGTCACCGTCTCCTCA
(SEQ ID NO:22)

B4 scFV nucleic acid sequence

GCCGCCACCATGAGTGTGCCCACTCAGGTCCTGGGGTTGCTGCTGCTGTGG
CTTACAGATGCCAGATGTGTGAAGCTGCAGGAGTCTGGACCTGAGCTGGT
GAAGCCTGGGGCTTCAGTGAAGATATCCTGTAAGGCTTCTGGATACACGT
TCACTGACTACTACATGAACTGGGTGAAGCAGAGCCATGGAAAGAGCCTT
GAGTGGATTGGAGATATTAATCCTAACAATGGTGGTACTAGCTACAACCA
GAAGTTCAAGGGCAAGGCCACATTGACTGTAGACAAGTCCTCCAGCACAG
CCTACATGGAGCTCCGCAGCCTGACATCTGAGGACTCTGCAGTCTATTACT
GTGCAAGATATGATTACGCTTGGTACTTCGATGTCTGGGGCCAAGGGACC
ACGGTCACCGTCTCCTCAGGCGGAGGTGGGTGGGTGGCGGCGGATCTGG
CGGAGGTGGGGATGTTTTGATGACCCAACTCCACTCTCCCTGCCTGTCAG
TCTTGGAGATCAAGCCTCCATCTCTTGCAGATCTAGTCAGAGCATTGTACA
TAGTAATGGAAACACCTATTTAGAATGGTACCTGCAGAAACCAGGCCAGT
CTCCAAAGCTCCTGATCTACAAAGTTTCCAACCGATTTTCTGGGGTCCCAG
ACAGGTTCAGTGGCAGTGGATCAGGGACAGATTTCACTCAAGATCAGC
AGAGTGGAGGCTGAGGATCTGGGAGTTTATTACTGCTTTCAAGGTTACA
TGTTCCGTACACGTTTCGGAGGGGGGACCAAGCTGGAAATAAAACGGATCG
AAGGCCGGCATCACCATCATCACCCTGATAG (SEQ ID NO:23)

CHO optimized B4 scFV nucleic acid sequence

ATGTCCGTGCCTACCCAGGTGCTCGGACTCCTGCTGCTGTGGCTCACCGAC
GCCAGGTGTGTGAAGCTGCAGGAGAGCGGACCCGAGCTGGTGAAGCCTG
GAGCCTCCGTGAAGATCAGCTGCAAGGCTTCCGGATACACCTTCACCGAC
TACTATATGAACTGGGTGAAGCAGAGCCACGGCAAGAGCCTGGAGTGGAT
CGGCGACATCAACCCTAACAACGGCGGCACCTCCTACAACCAGAAGTTCA
AGGGCAAGGCTACACTGACCGTGGACAAGTCCTCCAGCACCGCCTACATG
GAGCTCAGGAGCCTGACCTCCGAGGATTCCGCCGTCTATTACTGTGCCCCG
GTACGACTACGCCTGGTATTTCGACGTGTGGGGCCAGGGCACAACCGTCA
CAGTCTCCAGCGGAGGAGGAGGAAGCGGCGGCGGAGGATCCGGAGGCGG
AGGCGATGTCCTGATGACACAGACACCTCTGAGCCTCCCCGTGAGCCTGG
GAGACCAAGCCTCCATCTCCTGCAGGTCCTCCCAGTCCATCGTGACAGC
AATGGCAACACCTACCTGGAGTGGTATCTGCAGAAGCCTGGCCAGTCCCC
CAAGCTGCTGATCTACAAGGTGTCCAACCGGTTTCAGCGGCGTCCCTGACA
GGTTCTCCGGATCCGGAAGCGGCACAGATTTCACCCTGAAGATCAGCAGG
GTCGAGGCCGAGGACCTGGGAGTGTACTACTGCTTCCAGGGCTCCCATGT
CCCTTACACCTTCGGCGGCGGCACCAAACTGGAGATCAAGCGGATCGAGG
GCAGGCATCACCACCATCACCCTGA (SEQ ID NO:24)

C2 CDR-L1 amino acid sequence

KSVSTSGYSY (SEQ ID NO:25)

C2 CDR-L2 amino acid sequence

LVS (SEQ ID NO:26)

C2 CDR-L3 amino acid sequence

QHIRELTRSEGGPSWK (SEQ ID NO:27)

C2 CDR-H1 amino acid sequence

GYTFTSYW (SEQ ID NO:28)

C2 CDR-H2 amino acid sequence

INPSNGGT (SEQ ID NO:29)

C2 CDR-H3 amino acid sequence

ARRGIRLRHFDY (SEQ ID NO:30)

C2 CDR-L1 amino acid sequence

QDVGTA (SEQ ID NO:31)

C2 CDR-L2 amino acid sequence

WAS (SEQ ID NO:32)

C2 CDR-L3 amino acid sequence

QQYSSYPLT (SEQ ID NO:33)

C2 VL amino acid sequenceDIVMTQSPASLAVSLGQRATISYRASKSVSTSGYSYMHWNQQKPGQPPRLLIY
LVSNLESGVPARFSGSGSGTDFTLNIHPVEEEDAATYYCQHIRELTRSEGGPS
WK (SEQ ID NO:34)C2 VL amino acid sequenceDIQMTQSPKFMSTSVGDRVSITCKASQDVGTAVAWYQQKPGQSPKLLIYWAS
TRHTGVPDRFTGSGSGTDFTLTISNVQSEDLADYFCQQYSSYPLTFGAGTKLE
LK (SEQ ID NO:35)C2 VH amino acid sequenceVKLQESGTELVKPGASVKLSCKASGYTFTSYWMHWVKQRPGQGLEWIGNIN
PSNGGTNYNEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARRGIRLR
HFDYWGQGTTVTVS (SEQ ID NO:36)C2 scFV amino acid sequenceVKLQESGTELVKPGASVKLSCKASGYTFTSYWMHWVKQRPGQGLEWIGNIN
PSNGGTNYNEKFKSKATLTVDKSSSTAYMQLSSLTSEDSAVYYCARRGIRLR
HFDYWGQGTTVTVSSRANSADIIHTGGRSSMHLEGPIRPIVSRISGGGGGSGG
GGSWISAEFALDIVMTQSPASLAVSLGQRATISYRASKSVSTSGYSYMHWNQ
QKPGQPPRLLIYLVSNLESGVPARFSGSGSGTDFTLNIHPVEEEDAATYYCQHI
RELTRSEGGPSWK (SEQ ID NO:37)C2 scFV amino acid sequenceMSVPTQVLGLLLLWLTDARCVKLQESGTELVKPGASVKLSCKASGYTFTSY
WMHWVKQRPGQGLEWIGNINPSNGGTNYNEKFKSKATLTVDKSSSTAYMQ
LSSLTSEDSAVYYCARRGIRLRHFDYWGQGTTVTVSSGGGGGSGGGGSGGGGS
DIQMTQSPKFMSTSVGDRVSITCKASQDVGTAVAWYQQKPGQSPKLLIYWAS
TRHTGVPDRFTGSGSGTDFTLTISNVQSEDLADYFCQQYSSYPLTFGAGTKLE
LKRIEGRHHHHHH (SEQ ID NO:38)C2 VL nucleic acid sequenceGACATTGTGATGACACAGTCTCCTGCTTCCTTAGCTGTATCTCTGGGGCAG
AGGGCCACCATCTCATACAGGGCCAGCAAAAGTGTCAGTACATCTGGCTA
TAGTTATATGCACTGGAACCAACAGAAACCAGGACAGCCACCCAGACTCC
TCATCTATCTTGTATCCAACCTAGAATCTGGGGTCCCTGCCAGGTTTCAGTG
GCAGTGGGTCTGGGACAGACTTCACCCTCAACATCCATCCTGTGGAGGAG
GAGGATGCTGCAACCTATTACTGTCAGCACATTAGGGAGCTTACACGTTT
GGAGGGGGGACCAAGCTGGAAA (SEQ ID NO:39)

C2 VL nucleic acid sequence

GACATCCAGATGACCCAGTCTCCCAAATTCATGTCCACATCAGTAGGAGA
CAGGGTCAGCATCACCTGCAAGGCCAGTCAGGATGTGGGTACTGCTGTAG
CCTGGTATCAACAGAAACCAGGGCAATCTCCTAAACTACTGATTTACTGG
GCATCCACCCGGCACACTGGAGTCCCTGATCGCTTCACAGGCAGTGGATC
TGGGACAGATTTCACTCTCACCATTAGCAATGTGCAGTCTGAAGACTTGGC
AGATTATTTCTGTCAGCAATATAGCAGCTATCCTCTCACGTTCCGGTGCTGG
GACCAAGCTGGAGCTGAAAC (SEQ ID NO:40)

C2 VH nucleic acid sequence

GTGAAACTGCAGGAGTCTGGGACTGAACTGGTGAAGCCTGGGGCTTCAGT
GAAGCTGTCCTGCAAGGCTTCTGGCTACACCTTCACCAGCTACTGGATGCA
CTGGGTGAAGCAGAGGCCTGGACAAGGCCTTGAGTGGATTGGAATATTA
ATCCTAGCAATGGTGGTACTAACTACAATGAGAAGTTCAAGAGCAAGGCC
ACACTGACTGTAGACAAATCCTCCAGCACAGCCTACATGCAGCTCAGCAG
CCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGAAGAGGCATAC
GGTTACGACACTTTGACTACTGGGGCCAAGGGACCACGGTCACCGTCTCC
(SEQ ID NO:41)

C2 scFV nucleic acid sequence

GCCGCCACCATGAGTGTGCCCACTCAGGTCCTGGGGTTGCTGCTGCTGTGG
CTTACAGATGCCAGATGTGTGAAACTGCAGGAGTCTGGGACTGAACTGGT
GAAGCCTGGGGCTTCAGTGAAGCTGTCCTGCAAGGCTTCTGGCTACACCTT
CACCAGCTACTGGATGCACTGGGTGAAGCAGAGGCCTGGACAAGGCCTTG
AGTGGATTGGAATATTAATCCTAGCAATGGTGGTACTAACTACAATGAG
AAGTTCAAGAGCAAGGCCACACTGACTGTAGACAAATCCTCCAGCACAGC
CTACATGCAGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTATTG
TGCAAGAAGAGGCATACGGTTACGACACTTTGACTACTGGGGCCAAGGGA
CCACGGTCACCGTCTCCTCTGGCGGAGGTGGGTCTGGGTGGCGGCGGATCT
GGCGGAGGTGGGTCTGGACATCCAGATGACCCAGTCTCCCAAATTCATGTC
CACATCAGTAGGAGACAGGGTCAGCATCACCTGCAAGGCCAGTCAGGATG
TGGGTACTGCTGTAGCCTGGTATCAACAGAAACCAGGGCAATCTCCTAAA
CTACTGATTTACTGGGCATCCACCCGGCACACTGGAGTCCCTGATCGCTTC
ACAGGCAGTGGATCTGGGACAGATTTCACTCTCACCATTAGCAATGTGCA
GTCTGAAGACTTGGCAGATTATTTCTGTCAGCAATATAGCAGCTATCCTCT
CACGTTCCGGTGCTGGGACCAAGCTGGAGCTGAAACGGATCGAAGGCCGGC
ATCACCATCATCACCCTGATAG (SEQ ID NO:42)

CHO optimized C2 scFV nucleic acid sequence

ATGAGCGTGCCTACACAGGTGCTCGGCCTGCTGCTCCTCTGGCTGACAGA
CGCCCGGTGTGTGAAGCTGCAGGAGTCCGGAACCGAGCTGGTGAAACCTG
GCGCCAGCGTGAAACTGAGCTGCAAAGCCAGCGGATACACCTTCACCTCC
TACTGGATGCACTGGGTGAAACAGAGGCCTGGCCAGGGCCTGGAATGGAT
TGGCAACATCAACCCCAGCAACGGCGGCACCAACTACAATGAGAAGTTCA

AGAGCAAGGCCACCCTGACCGTGGATAAGTCCTCCTCCACCGCCTACATG
CAGCTGTCCTCCCTCACCTCCGAGGACAGCGCCGTCTATTACTGTGCCAGG
CGGGGCATCAGGCTGAGGCACTTCGACTACTGGGGCCAAGGCACAACCGT
CACCGTGAGCTCCGGAGGAGGAGGCAGCGGAGGCGGAGGCTCCGGCGGA
GGCGGAAGCGACATTCAGATGACCCAGAGCCCCAAGTTCATGTCCACCTC
CGTCGGCGACAGGGTGAGCATCACCTGTAAGGCCAGCCAGGATGTCCGGA
CAGCTGTGGCCTGGTACCAGCAGAAGCCCGGCCAGTCCCCCAAGCTGCTG
ATCTACTGGGCTTCCACAAGGCATACCGGCGTCCCCGATAGGTTACAGG
CTCCGGCTCCGGCACCGACTTCACACTCACCATCAGCAACGTCCAGTCCG
AGGACCTGGCCGACTACTTCTGCCAGCAGTACTCCAGCTACCCCCTCACCT
TCGGCGCTGGCACCAAGCTGGAAGCTCAAGCGGATCGAGGGCAGGCATCAC
CACCATCACCCTGATAG (SEQ ID NO: 43)

Full-length human membrane cofactor protein (MCP)

MEPPGRRECPFPSWRFPGLLLAAMVLLLYSFSDACEEPPTFEAMELIGKPKPY
YEIGERVDYKCKKGYFYIPPLATHTICDRNHTWLPVSDDACYRETCPIYRDPL
NGQAVPANGTYEFGYQMHFICNEGYYLIGEEILYCEKGSVAIWSGKPPICEK
VLCTPPPPIKNGKHTFSEVEVFEYLDVAVTYSDDPAPGPDPSLIGESTIYCGDN
SVWSRAAPECKVVKCRFPVVENGGKQISGFGKKFYKATVMFECDKGFYLDG
SDTIVCDNSTWDPPVPKCLKVLPPSSSTKPPALSH
SVSTSSTTKSPASSASGPRPTYKPPVSNYPGYPKPEEGILDSLDVWVIAVIVIAI
VVGVAVICVVPYRYLQRRKKKGTYLTDETHREVKFTSL (SEQ ID NO: 44)

Full-length human decay accelerating factor (DAF)

MTVARPSVPAALPLLGLPRLLLVLLCLPAVWGDCLPPDVPNAQPALEGR
TSFPEDTVITYKCEESFVKIPGEKDSVICLKGSQWSDIEEFCNRSCEVPTRLNSA
SLKQPYITQNYFPVGTVEYECRPGYRREPSLSPKLTCLQNLKWSTAVEFCKK
KSCPNGEIRNGQIDVPGGILFGATISFSCNTGYKLFGSTSSFCLISGSSVQWSD
PLPECREIYCPAPPQIDNGHIIQGERDHYGYRQSVTYACNKGFTMIGEHSIYCTV
NNDEGEWSGPPPECRGKSLTSKVPPTVQKPTTVNVPTTEVSPTSQKTTTCTTT
PNAQATRSTPVSRTTKHFHETTPNKGSGTTSCTTRLLSGHTCFTLTGLLGTL
VTMGLLT (SEQ ID NO: 45)

Full-length mouse decay accelerating factor (DAF)

MIRGRAPRTRPSPPPPLLPLLSLLLLSPTVRGDCGPPPDIPNARPILGRHSKFA
EQSKVAYSCNNGFKQVPDKSNIVVCLENGQWSSHETFCCKSCVAPERLSFAS
LKKEYLNMNFFPVGTIVEYECRPGFRKQPPLPGKATCLEDLVWSPVAQFCKK
KSCPNNPKDLNNGHINIPTGILFGSEINFSCNPGYRLVGVSSSTFCSTGNTVDWD
DEFPVCTEIHCPPEPKINNGIMRGESDSYYSQVVITYSCDKGFILVGNASIYCT
VSKSDVGQWSSPPPRCIEKSKVPTTKPTINVPSTGTPSTPQKPTTESVNPBGDQ
PTPQKPSTVKVSATQHVPVTKTTVRHPIRTSTDKGEPNTGGDRYIYGHTCLITL
TVLHVMLSLIGYLT (SEQ ID NO: 46)

Full-length human CD59

MGIQGGSVLFGLLLVLAVFCHSGHSLQCYNCPNPTADCKTAVNCSSDFDACL
ITKAGLQVYNKCWKFEHCNFDVTTTRLRENELTYYCCKKDLNCFNEQLENG
GTSLSEKTVLLLVTPLAAA WSLHP (SEQ ID NO: 47)

Full-length mouse CD59

MRAQRGLILLLLLLAVFCSTAVSLTCYHCFQPVVSSCNMNSTCSPDQDSCLYA
VAGMQVYQRCWKQSDCHGEIIMDQLEETKLKFRCCQFNLCNKSDGSLGKTP
LLGTSVLVAILNLCFLSHL (SEQ ID NO: 48)

Full-length mouse CD59, isoform B

MRAQRGLILLLLLLAVFCSTAVSLKCYNCFQFVSSCKINTTCSPNLDSCLYAV
AGRQVYQQCWKLSDCNSNYIMSRDLVAGIQSKCCQWGLCNKNLDGLEEPN
NAETSSLRKTALLGTSVLVAILKFCF (SEQ ID NO: 49)

Full-length mouse complement receptor 1-related gene/protein y (Crry)

MEVSSRSSEPLDPVWLLVAFGRGGVKLEVLLLFLLPFTLGELRGGLGKHGHT
VHREPAVNRLCADSKRWSGLPVSAQRPFPMGHCPAPSQLPSAKPINLTDESM
FPIGTYYLLECLPGYIKRQFSITCKQDSTWTS AEDKCIRKQCKTPSPENGLVH
VHTGIQFGSRINYTCNQGYRLIGSSSAVCVITDQSV DWDTEAPICEWIPCEIPPG
IPNGDFFSSTREDFHYGMVVITYRCNTDARGKALFNLVGEPSLYCTSNDEIGV
WSGPPPQCIELNKCTPPPYVENAVMLSENRSLSLRDIVEFRCHPGFIMKGASS
VHCQSLNKWEPELPSCFKGVICRLPQEMSGFQKGLGMKKEYYYGENVTLEC
EDGYTLEGSSQSQCQSDGSWNPLAKCVSRISGLIVGIFIGIIVFILVIIVFIWMI
LKYKKRNTTDEKYKEVGIHLNYKEDSCVRLQSLTTSQENSSTTSPARNSLTQE
V S (SEQ ID NO:50)

Full-length human complement receptor 1(CR1)

MGASSPRSPPEVGPAPGLPFCCGGSLLAVVLLALPV AWGQCNAPEWL PFA
RPTNL TDEFEP IGT YL NYE CRPGYSGRPFSIICL KNSVWTGAKDR CR RKSCRN
PPDPVNGMVHVIKGIQFGSQIKYSCTKGYRLIGSSSATCIISGDTV IWDNETPIC
DRIPCGLPPTITNGDFISTNREN FHYGSVVITYRCNPGSGGRKVFELVGEPSIYC
TSNDDQVGIWSGPAPQCIIPNKCTPPNVENGILVSDNRSLSLNEVVEFRCPG
FVMKGPRRVKQCQALNKWEPELPSCSRVCQPPPDVLHAERTQRDKDNFSPGQE
VFYSCEPGYDLRGAASMRCTPQGDWSPAAPTCEVKSCDDFMGQLLN GRVL F
PVNLQLGAKVDFVCDEGFQLKGSSASYCVLAGMESLWNSSVPVCEQIFCPS
PVIPNGRHTGKPLEVFPFGKAVNYTCDPHPDRGTSFDLIGESTIRCTSDPQGNG
VWSSPAPRCGILGHCQAPDHFLFAKLKTQTNASDFPIGTSLKYECRPEYYGRP
FSITCLDNLVWSSPKDVCKRKSCKTPPDPVNGMVHVITDIQVGS RINYSCTTG
HRLIGHSSAECILSGNAAHWSTKPPICQRI PCGLPPTIANGDFISTNREN FHYGS
VVITYRCNPGSGGRKVFELVGEPSIYCTSNDDQVGIWSGPAPQCIIPNKCTPPN
VENGILVSDNRSLSLNEVVEFRCPG FVMKGPRRVKQCQALNKWEPELPSCS
RVCQPPPDVLHAERTQRDKDNFSPGQEVFYSCEPGYDLRGAASMRCTPQGD
WSPAAPTCEVKSCDDFMGQLLN GRVLFPVNLQLGAKVDFVCDEGFQLKGSS

ASYCVLAGMESLWNSSVPVCEQIFCPSPPVIPNGRHTGKPLEVFPFGKAVNYT
CDPHPDRGTSFDLIGESTIRCTSDPQNGVWSSPAPRCGILGHCQAPDHFLFAK
LKTQTNASDFPIGTSCLKYECRPEYYGRPFSITCLDNLVWSSPKDVCKRKSKT
PPDPVNGMVHVITDIQVGSRINYSC TTGHR LIGHSSAECILSGNTAHWSTKPPI
CQRIPCGLPPTIANGDFISTNREN FHYG SVVTYRCNLGSRGRKVFELVGEPSIY
CTSNDQVGIWSGPAPQCIIPNKCTPPNVENGILVSDNRSLFSLNEVVEFRCQP
GFVMKGPRRVKCQALNKWEPELPSCSRVCQPPPEILHGEHTPSHQDNFSPGQE
VFYSCEPGYDLRGAASLHCTPQGDWSPEAPRC AVKSCDDFLGQLPHGRVLP
LNLQLGAKVSFVCDEGFRLKGSSVSHCVLVGM RSLWNNSVPVCEHIFCPNPP
AILNGRHTGTPSGDIPYGKEISYTC DPHPDRGMTFNLIGESTIRCTSDPHGNGV
WSSPAPRCELSVRAGHCKTPEQFPFASPTIPINDFEFPVGTSLNYECRPGYFGK
MFSISCLENLVWSSVEDNCRRKSCGPPPEPFNGMVHINTDTQFGSTVNYSCNE
GFRLIGSPSTTCLVSGNNVTW DKKAPICEIISCEPPPTISNGDFYSNNRTSFHNG
TVVTYQCHTGP DGEQLFELVGERSIYCTSKDDQVG V WSSPPPRCISTNKCTAP
EVENAIRVPGNRSFFSLTEIIRFCQPGFVMVGSHTVQCQTNGRWGPKLPHCS
RVCQPPPEILHGEHTLSHQDNFSPGQEVFYSCEPSYDLRGAASLHCTPQGDWS
PEAPRCTVKSCDDFLGQLPHGRVLLPLNLQLGAKVSFVCDEGFRLKGRSASH
CVLAGMKALWNSSVPVCEQIFCPNPPAILNGRHTGTPFGDIPYGKEISYACDT
HPDRGMTFNLIGESSIRCTSDPQNGVWSSPAPRCELSVPAACPHPPKIQNG
HYIGGHVSLYLPGMTISYTCDPGYLLVGKGFIFCTDQGIWSQLDHYCKE VNC S
FPLFMNGISKELEMKKVYHYGDYVTLKCEDGYTLEGSPWSQCQADDRWDPP
LAKCTSR AHDALIVGTLSGTIFFILLIIFLSWILKHRKGNNAHENPKEVAIHLHS
QGGSSVHPRTLQTNEENSRLP (SEQ ID NO: 51)

Full-length human factor H

MRL LAKIICLMLWAICVAEDCNELPPRRNTEILTGSWSDQTYPEGTQAIYKCR
PGYRSLGNVIMVCRKGEWVALNPLRK CQKRPCGHPGDTPFGTFTLTGGNVFE
YGVKAVYTCNEG YQLLGEINYRECDTDGWTNDIPICEVVKCLPVTAPENGKI
VSSAMEPDREYHFGQAVRFVCNSGYKIEGDEEMHCSDDGFWSKEKPKC V EIS
CKSPDVINGSPISQKIIYKENERFQYKCNMGY EY SERGDAVCTESGWRPLPSC
EEKSCDNPYIPNGDYSPLRIKHRTGDEITYQCRNGFY PATRGNTAKCTSTGWI
PAPRCTLKPCDYPDIKHGGLYHENMRRPYFPVAVGKYYSY CDEHFETPSGS
YWDHIHCTQDGWSPA VPCLRKCYFPYLENGYNQNYGRKFVQGKSIDVACHP
GYALPKAQTTVT CMENGWSPTPRCIRVKTCSSIDIENGFISESQYTYALKEK
AKYQCKLGYVTADGETSGSITCGKDGWSAQPTCIKSCDIPVFMNARTKNDFT
WFKLNDTLDYECHDGYESNTGSTTGSIVCGYNGWSDLPICYERECELPKIDV
HLVPDRKKDQYKVGEVLKFSCKPGFfiVGPNSVQCYHFGLSPDLPICKEQVQS
CGPPPELLNGNVKEKTKEEYGHSEVVEYYCNPRFLMKGP NKIQCV DGEWTTL
PVCIVEESTCGDIPELEHGW AQLSSPPY YYGDSVEFNCSESFTMIGHRSITCIHG
VWTQLPQCVAIDKLKKCKSSNLIIEHLKNKKEFDHNSNIRYRCRGKEGWIH
TVCINGRWDPEVNCSMAQIQLC P P P P QIPNSHNMTTTLNYRDGEKVS V L C Q E
NYLIQEGEEITCKDGRWQSIPLC V E K I P C S Q P P Q I E H G T I N S S R S S Q E S Y A H G T K
LSYTCEGGFRISEENETTCYMGKWSSPPQCEGLPCKSPPEISHGVVAHMSDSY
QYGEEV TYKCFEGFGIDGPAIAKCLGEKWSHPPSCIKTDCLSLPSFENAIPMGE

KKDVYKAGEQVITYTCATYYKMDGASNVTCINSRWTGRPTCRDTSCVNPPTV
 QNAYIVSRQMSKYPSEGERVRYQCRSPYEMFGDEEVMCLNGNWTEPPQCKDS
 TGKCGPPPIDNGDITSFPLSVYAPASSVEYQCQNLYQLEGNKRITCRNGQWS
 EPPKCLHPCVISREIMENYNIALRWTAQKQLYSRTGESVEFVCKRGYRLSSRS
 HTLR TTCWDGKLEYPTCAKR (SEQIDNO:52)

Full-length mouse factor H

MRLSARIWLILWTVCAAEDCKGPPPRENSEILSGSWSEQLYPEGTQATYKCR
 PGYRTLGTIVKVCKNGKWVASNPSRICRKKPCGHPGDTFPGSFRLAVGSQFEF
 GAKVVYTCDDGYQLLGEIDYRECGADGWINDIPLCEVVKCLPVTELENGRIV
 SGAAETDQEYYFGQVVRFECSNGFKIEGHKEIHCSENGLWSNEKPRCVEILCT
 PPRVENGDGINVKPVYKENERYHYKCKHGYVPKERGDAVCTGSGWSSQPF
 EEKRCSPPIYILNGIYTPHRIIHRSDDEIRYECNYGFYPVTGSTVSKCTPTGWIPV
 PRCTLKPCEFPQFKYGRLYYEESLRPNFPV SIGNKYSYKCDNGFSPPSGYSWD
 YLRCTAQWEPEVPCVRKCVFHYVENGD SAYWEKVYVQGGQSLKVQCYNGY
 SLQNGQDTMTCTENGWSPPPKCIKTCASDIHIDNGFLSESSSIYALNRETSY
 RCKQGYVTNTGEISGSITCLQNGWSPQPSCIKSCDMPVFENSITKNTRTWFKL
 NDKLDYECLVGFENEYKHTKGSITCTYYGWS DTPSCYEREC SVPTLDRKLVV
 SPRKEYR VGD LLEFSCHSGHRVGPDSVQCYHFGWSPGFPTCKGQVASCAPP
 LEILNGEINGAKKVEYSHGEVVKYDCKPRFLLKGP NKIQCV DGNWTTLPVCIE
 EERTCGDIPELEHGS AKCSVPPYHHGDSVEFICEENfMIGHGSVSCISGKWTQ
 LPKCVATDQLEKCRVLKSTGIEAIKPKLTEffHNSTMDYKCRDKQEYERSICIN
 GKWDPEPNCTSKTSCPPPPQIPNTQVIETTVKYLDGEKLSVLCQDNYLTQDSE
 EMVCKDGRWQSLPRCIEKIPCSQPPTIEHGSINLPRSSEERRDSIESSSHEHGTTF
 SYVCD DGFRIPEENRITCYMGKWSTPPRCVGLPCGPPPSIPLGTVSLELESYQH
 GEEV TYHCSTGFGIDGPAFIICEGGKWS DPPKCIKTDCDVLPTVKNAIIRGKSK
 KSYRTGEQVTFRCQSPYQMNGSDTVTCVNSRWIGQPVCKDNSCVDPPHVPN
 ATIVTRTKNKYLHGDRVRYECNKPLELFGQVEVMCENGIWTEKPKCRDSTG
 KCGPPPIDNGDITSLSLPVYEPLSSVEYQCQKYLLKGKKTITCTNGKWSEPP
 TCLHACVIPENIMESHNIILKWRHTEKIYSHSGEDIEFGCKYGYKARDSPPFR
 TKCINGTINYPTCV (SEQIDNO:53)

Signal peptide of the human CD5 protein

MPMGS LQPLATLYLLGMLVAS (SEQ ID NO: 54)

Signal peptide of the human CR2 protein

MGAAGLLGVFLALVAPG (SEQ ID NO:55)

Signal peptide of the human CR2 protein

MGAAGLLGVFLALVAPGVLG (SEQ ID NO:56)

B4 scFV nucleic acid sequence

GACACGTGATCAGCCGCCACCATGCCCATGGGGTCTCTGCAACCGCTGGC
 CACCTTG TACCTGCTGGGGATGCTGGTCGCTTCCGTGCTAGCGCATCATCA

TCATCATCATGTGAAACTGCAGGAGTCAGGGGCTGAGCTTGTGAAGCCTG
GGGCTTCAGTGAAGCTGTCTGCAAGGCTTCTGGCTACACCTTCACCAGCT
ACTGGATGCACTGGGTGAAGCAGAGGCCTGGACGAGGCCTTGAGTGGATT
GGAAGGATTGGTCCTAATAGTGGTGGTACTAAGTACAATGAGAAGTTCAA
GAGCAAGGCCACACTGACTGTAGACAAACCCTCCAGCACAGCCTACATGC
AGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGA
AGAATGGTAAAGGGGTGCTATGGACTACTGGGGCCAAGGGACCACGGTC
ACCGTCTCCTCAAAGGGCGAATTCCAGCACACTGGCGGCCGTTACTAGTG
GATCCGAGCTCGGTACCAAGCTTGGCGTCAGGAGGCGGTGGCGGCTCGGG
TGGCGGCGGCTCTTGGATATCTGCAGAATTCGCCCTTGACATTGAGCTCAC
CCAGTCTCCAACCACCATGGCTGCATCTCCCGGGGAGAAGATCACTATCA
CCTGCAGTGCCAGCTCAAGTATAAGTTCCAATTACTTGCATTGGTATCAGC
AGAAGCCAGGATTCTCCCCTAAACTCTTGATTTATAGGACATCCAATCTGG
CTTCTGGAGTCCCAGCTCGCTTCAGTGGCAGTGGGTCTGGGACCTCTTACT
CTCTCACAATTGGCACCATGGAGGCTGAAGATGTTGCCACTTACTACTGCC
AGCAGGGTAGTAGTATACCACGTACACGTTTCGGAGGGGGCACCAAGCTGG
AAATAATAGACTAGTCGTGCG (SEQ ID NO:57)

C2 scFV nucleic acid sequence

GACACGAAGCTTGCCGCCACCATGCCCATGGGGTCTCTGCAACCGCTGGC
CACCTTGTACCTGCTGGGGATGCTGGTCGCTTCCGTGCTAGCGCATCATCA
TCATCATCATGTCAAGCTGCAGGAGTCTGGGACTGAACTGGTGAAGCCTG
GGGCTTCAGTGAAGCTGTCTGCAAGGCTTCTGGCTACACCTTCACCAGCT
ACTGGATGCACTGGGTGAAGCAGAGGCCTGGACAAGGCCTTGAGTGGATT
GGAATATTAATCCTAGCAATGGTGGTACTAACTACAATGAGAAGTTCAA
GAGCAAGGCCACACTGACTGTAGACAAATCCTCCAGCACAGCCTACATGC
AGCTCAGCAGCCTGACATCTGAGGACTCTGCGGTCTATTATTGTGCAAGA
AGAGGCATACGGTTACGACACTTTGACTACTGGGGCCAAGGGACCACGGT
CACCGTCTCCTCAAGGGCGAATTCTGCAGATATCCATCACACTGGCGGCC
GCTCGAGCATGCATCTAGAGGGCCCAATTTCGCCCTATAGTGAGTCGTATA
TCAGGAGGCGGTGGCGGCTCGGGTGGCGGCGGCTCTTGGATATCTGCAGA
ATTCGCCCTTGACATTGTGATGACACAGTCTCCTGCTTCCTTAGCTGTATCT
CTGGGGCAGAGGGCCACCATCTCATAAGGGCCAGCAAAAGTGTCAGTAC
ATCTGGCTATAGTTATATGCACTGGAACCAACAGAAACCAGGACAGCCAC
CCAGACTCCTCATCTATCTTGTATCCAACCTAGAATCTGGGGTCCCTGCCA
GGTTCAGTGGCAGTGGGTCTGGGACAGACTTCACCCTCAACATCCATCCT
GTGGAGGAGGAGGATGCTGCAACCTATTACTGTCAGCACATTAGGGAGCT
TACACGTTTCGGAGGGGGGACCAAGCTGGAAATAATAGCCCGGGCGTGCG
(SEQ ID NO:58)

CLAIMS

What is claimed is:

1. A method of inhibiting complement-mediated inflammation in a tissue having non-ischemic injury in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor.
2. A method of treating an inflammatory disease in an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or a phospholipid; and (b) a complement inhibitor.
3. The method of claim 1 or claim 2, wherein the complement inhibitor is selected from the group consisting of an anti-C5 antibody, anti-MASP antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-factor D antibody, and an anti-properdin antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding.
4. The method of claim 1 or 2, wherein the complement inhibitor is a specific inhibitor of the alternative pathway.
5. The method of claim 1 or 2, wherein the complement inhibitor is a specific inhibitor of the lectin pathway.
6. A method of detecting complement-mediated injury in a tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV or phospholipid; and (b) a detectable moiety,

wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated tissue injury.

7. The method of claim 6, wherein the method further comprises detecting the detectable moiety.
8. The method of claim 6 or claim 7, wherein the detectable moiety is selected from the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.
9. The method of any one of claims 1 and 3-8, wherein the tissue injury results from any of from inflammatory disorders, transplant rejection, pregnancy-related diseases, adverse drug reactions, autoimmune or immune complex disorders.
10. The method of any one of claims 1 and 3-8, wherein the tissue is any one of eye, joint, kidney, brain, heart, spinal cord, and liver.
11. The method of claim 10, wherein the tissue is any one of eye, joint, and kidney.
12. The method of any one of claims 2-5 and 7-8, wherein the disease is any one of an ocular disease, arthritis, or renal injury.
13. The method of any one of claims 1-12, wherein the antibody or fragment thereof specifically binds to Annexin IV.
14. The method of claim 13, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV.
15. The method of claim 13 or 14, wherein the antibody or antibody fragment thereof binds to the same epitope as monoclonal antibody B4.
16. The method of any one of claims 13-15, wherein the Annexin IV is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing non-ischemic injury.
17. The method of any one of claims 1-12, wherein the antibody or fragment thereof specifically binds to a phospholipid.
18. The method of claim 17, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid.
19. The method of claim 17 or 18, wherein the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody C2.

20. The method of any one of claims 17-19, wherein the phospholipid is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage.
21. The method of any one of claims 17-20, wherein the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC).
22. The method of any one of claims 17-20, wherein the phospholipid is MDA.
23. The method of any one of claims 1-22, wherein the construct is a fusion protein.
24. The method of claim 23, wherein the antibody or fragment thereof and the complement inhibitor or detectable moiety are linked via a peptide linker.
25. The method of any one of claims 1-24, wherein the antibody or fragment thereof is a scFv.
26. The method of any one of claims 1-24, wherein the antibody or fragment thereof is Fab, Fab', or F(ab')₂.
27. A construct comprising: (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to Annexin IV; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1 or 7, a sequence of SEQ ID NO:2 or 8, or a sequence of SEQ ID NO:3 or 9; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:4 or 10, a sequence of SEQ ID NO:5 or 11, or a sequence of SEQ ID NO:6 or 12.
28. The construct of claim 27, wherein the antibody or fragment thereof comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:1 or 7; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:2 or 8; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:3 or 9.
29. The construct of claims 27 or 28, wherein the antibody or fragment thereof comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:4 or 10; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:5 or 11; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:6 or 12.

30. The construct of any one of claims 27-29, wherein the antibody or fragment there of comprises a light chain variable domain of SEQ ID NO:13 or 14.
31. The construct of any one of claims 27-30, wherein the antibody or fragment there of comprises a heavy chain variable domain of SEQ ID NO:15 or 16.
32. The construct of any one of claims 27-31, wherein the antibody or fragment is an scFv having the sequence of SEQ ID NO:17 or 18.
33. The construct of any one of claims 27-32, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV.
34. The construct of any one of claims 27-33, wherein the antibody or fragment thereof binds to the same epitope as monoclonal antibody B4.
35. The construct of any one of claims 27-34, the Annexin IV is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing or is at risk of undergoing tissue injury.
36. A construct comprising: (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof specifically binds to a phospholipid; and (b) a complement modulator or a detectable moiety, wherein the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25 or 31, a sequence of SEQ ID NO:26 or 32, or a sequence of SEQ ID NO:27 or 33; and/or (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:28, a sequence of SEQ ID NO:29, or a sequence of SEQ ID NO:30.
37. The construct of claim 36, wherein the antibody or fragment there of comprises: (i) a light chain variable domain comprising a sequence of SEQ ID NO:25 or 31; (ii) a light chain variable domain comprising a sequence of SEQ ID NO:26 or 32; and (iii) a light chain variable domain comprising a sequence of SEQ ID NO:27 or 33.
38. The construct of claims 36 or 37, wherein the antibody or fragment there of comprises: (i) heavy chain variable domain comprising a sequence of SEQ ID NO:28; (ii) heavy chain variable domain comprising a sequence of SEQ ID NO:29; and (iii) heavy chain variable domain comprising a sequence of SEQ ID NO:30.
39. The construct of any one of claims 36-38, wherein the antibody or fragment there of comprises a light chain variable domain of SEQ ID NO:34 or 35.

40. The construct of any one of claims 36-39, wherein the antibody or fragment thereof comprises a heavy chain variable domain of SEQ ID NO:36.
41. The construct of any one of claims 36-40, wherein the antibody or fragment is an scFv having the sequence of SEQ ID NO:37 or 38.
42. The construct of any one of claims 36-41, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid.
43. The construct of any one of claims 36-42, wherein the antibody or fragment thereof binds to the same epitope as monoclonal antibody C2.
44. The construct of any one of claims 36-43, wherein the phospholipid is present on the surface of a cell in an individual that is in or adjacent to a tissue undergoing or is at risk of undergoing tissue injury.
45. The method of any one of claims 36-44, wherein the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC).
46. The method of any one of claims 36-44, wherein the phospholipid is MDA.
47. The construct of any one of claims 27-46, wherein the construct comprises a complement modulator.
48. The construct of claim 47, wherein the complement modulator is a complement inhibitor.
49. The construct of claim 48, wherein the complement inhibitor is selected from the group consisting of an anti-C5 antibody, anti-MASP antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-factor D antibody, and an anti-properdin antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding.
50. The construct of claim 48, wherein the complement inhibitor is a specific inhibitor of the alternative pathway.
51. The construct of claim 48, wherein the complement inhibitor is a specific inhibitor of the lectin pathway.
52. The construct of any one of claims 27-46, wherein the construct comprises a detectable moiety.

53. The method of claim 52, wherein the detectable moiety is selected from the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.
54. The construct of any one of claims 27-53, wherein the construct is a fusion protein.
55. The construct of claim 54, wherein the antibody or fragment thereof and the complement modulator or detectable moiety are linked by a peptide linker.
56. A pharmaceutical composition comprising a construct of any one of claims 27-51 and 54-55.
57. A method of inhibiting complement-mediated inflammation in an individual, comprising administering to the individual an effective amount of a composition of claim 56.
58. A method of treating an inflammatory disease in an individual, comprising administering to the individual an effective amount of a composition of claim 56.
59. A composition comprising a construct of any one of claims 52 or 53.
60. A method of detecting complement-mediated injury in a tissue of an individual, comprising administering to the individual an effective amount of a composition of claim 59.
61. A method of inhibiting complement-driven inflammation in the eye in an individual, comprising administering to the individual an effective amount of (a) an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid; or (b) a composition comprising a construct, wherein the construct comprises: (i) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent.
62. A method of treating a complement-associated ocular disease or an ocular disease involving oxidative damage, comprising administering to the individual an effective amount of (a) an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid; or (b) a composition comprising a construct, wherein the construct comprises: (i) an antibody or a fragment thereof, wherein the

- antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent.
63. The method of claim 61 or claim 62, wherein the method comprises administration of an antibody or fragment thereof, wherein the antibody or fragment thereof does not activate complement activation, and wherein the antibody or fragment thereof specifically binds to Annexin IV or a phospholipid.
64. The method of claim 61 or claim 62, wherein the method comprises administration of a composition comprising a construct, wherein the construct comprises: (i) an antibody or a fragment thereof, wherein the antibody or fragment thereof specifically binds to Annexin IV or phospholipid, and (ii) a therapeutic agent.
65. The method of claim 64, wherein the therapeutic agent is a complement inhibitor.
66. The method of claim 65, wherein the complement inhibitor is selected from the group consisting of an anti-C5 antibody, an Eculizumab, an pexelizumab, an anti-C3b antibody, an anti-C6 antibody, an anti-C7 antibody, an anti-factor B antibody, an anti-MASP antibody, an anti-factor D antibody, and an anti-properdin antibody, a membrane cofactor protein (MCP), a decay accelerating factor (DAF), a CD59, a Crry, a CR1, a factor H, a Factor I, a linear peptide, a cyclic peptide, a compstatin, an N-acetylaspartylglutamic acid (NAAGA), and a biologically active fragment of any the preceding.
67. The method of any one of claims 62-66, wherein the ocular disease is selected from the group consisting of age-related macular degeneration (AMD) (including wet AMD and dry AMD), cytomegalovirus (CMV) retinitis, macular edema, uveitis (anterior and posterior), glaucoma, open/wide-angle glaucoma, close/narrow-angle glaucoma, retinitis pigmentosa (RP), proliferative vitreoretinopathy, retinal detachment, corneal wound healing, corneal transplants, and ocular melanoma.
68. The method of claim 67, wherein the ocular disease is AMD.
69. A method of detecting complement-mediated injury in an eye tissue of an individual, comprising administering to the individual an effective amount of a composition comprising a construct, wherein the construct comprises (a) an antibody or a fragment thereof, wherein the antibody or a fragment thereof

specifically binds to Annexin IV or a phospholipid; and (b) a detectable moiety, wherein the presence of the detectable moiety at the tissue is indicative of a complement-mediated eye tissue injury.

70. The method of claim 69, wherein the method further comprises detecting the detectable moiety.
71. The method of claim 69 or 70, wherein the detectable moiety is selected from the group consisting of radioisotopes, fluorescent dyes, electron-dense reagents, enzymes, biotins, paramagnetic agents, magnetic agents, and nanoparticles.
72. The method of any one of claims 61-71, wherein the antibody or fragment thereof specifically binds to Annexin IV.
73. The method of claim 72, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody B4 to Annexin IV.
74. The method of claim 72 or claim 73, wherein the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody B4.
75. The method of any one of claims 72-74, wherein the Annexin IV is present on the surface of a cell, a basement membrane, or in a pathological structure in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage.
76. The method of any one of claims 61-71, wherein the antibody or fragment thereof specifically binds to a phospholipid.
77. The method of claim 76, wherein the antibody or fragment thereof competitively inhibits the binding of monoclonal antibody C2 to phospholipid.
78. The method of claim 76 or 77, wherein the antibody or fragment thereof binds to the same epitope as that of monoclonal antibody C2.
79. The method of any one of claims 76-78, wherein the phospholipid is present on the surface of a cell, a basement membrane, or in a pathological structure in an individual that is in or adjacent to a tissue undergoing tissue injury and/or oxidative damage.
80. The method of any one of claims 76-79, wherein the phospholipid is selected from the group consisting of phosphatidylethanolamine (PE), cardiolipin (CL), and phosphatidylcholine (PC).
81. The method of any one of claims 76-79, wherein the phospholipid is MDA.

82. The method of any one of claims 61-81, wherein the antibody or fragment thereof is a scFv.
83. The method of any one of claims 61-81, wherein the antibody or fragment thereof is Fab, Fab', or F(ab')₂.
84. The method of any one of claims 64-83, wherein the construct is a fusion protein.
85. The method of any one of claims 64-83, wherein the antibody or a fragment thereof and the therapeutic agent or detectable moiety are joined by a linker.
86. The method of any one of claims 64-83, wherein the antibody or a fragment thereof and the therapeutic agent or detectable moiety are joined directly.
87. The method of any one of claims 61-86, wherein the administration is by injection into the eye.
88. The method of any one of claims 61-87, wherein the individual is human.

Figure 1




Figure 2





Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 7





Figure 8






Figure 9

Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 14




Figure 15




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 22




Figure 23




Figure 24




Figure 24




Figure 25





Figure 26







Figure 27

Figure 28




Figure 29




Figure 30




Figure 31




Figure 32




Figure 33





Figure 34






Figure 34

Figure 35





Figure 36







Figure 37

Figure 38






Figure 40

Figure 41




Figure 42





Figure 43



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2014/012831

A. CLASSIFICATION OF SUBJECT MATTER

C07K 19/00 (2006.01) A61K 39/395 (2006.01) A61P 37/00 (2006.01) A61P 27/02 (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Applicant/Inventor Search in PubMed and Espacenet; Keyword search in WPI, EPODOC, and MEDLINE: antibody, Annexin-IV, phospholipid, cardiolipin, MDA and complement, as well as similar/related keywords; Sequence search of SEQ ID NO. 1-12 and 25-33 depicted on pp. 211 and 214-215 with keywords Annexin or phospholipid or similar as appropriate.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Documents are listed in the continuation of Box C	

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 22 April 2014	Date of mailing of the international search report 22 April 2014
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA Email address: pct@ipaustalia.gov.au Facsimile No.: +61 2 6283 7999	Authorised officer Christina van Broekhoven AUSTRALIAN PATENT OFFICE (ISO 9001 Quality Certified Service) Telephone No. 0262833196

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/US2014/012831
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Kulik, L. et al. "Pathogenic Natural Antibodies Recognizing Annexin IV Are Required to Develop Intestinal Ischemia-Reperfusion Injury", J. Immunol. 2009; vol. 182, pp. 5363-5373 see Figure 1 and Discussion	27-35, 52-56 and 59
X	WO 2007/112403 A2 (The Regents of the University of Colorado et al.) 04 October 2007 see pp. 10, pp. 13, pp. 37-39, pp. 44, pp. 50, Example 1 and claims	27-46, 52-56 and 59
A	Elvington, A. et al. "Pathogenic Natural Antibodies Propagate Cerebral Injury Following Ischemic Stroke in Mice". J. Immunol. 2012; vol. 188, pp. 1460-1468 see abstract	1-88
A	CN 101235089 A (Zhang Xiuru) 06 August 2008 see abstract and claims	1-88

Form PCT/ISA/210 (fifth sheet) (July 2009)

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
the subject matter listed in Rule 39 on which, under Article 17(2)(a)(i), an international search is not required to be carried out, including
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a)

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See Supplemental Box for Details

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

Supplemental Box**Continuation of: Box III**

This International Application does not comply with the requirements of unity of invention because it does not relate to one invention or to a group of inventions so linked as to form a single general inventive concept.

This Authority has found that there are different inventions based on the following features that separate the claims into distinct groups:

- Claims 1-5, 9-51, 54-58, 61-68, 72-88 are directed to constructs comprising an antibody to Annexin-IV or phospholipids and a complement modulator, compositions comprising the same and methods of using the same. The feature of constructs comprising an antibody to Annexin-IV or phospholipids and a complement modulator is specific to this group of claims.
- Claims 6-35, 52-60, 69-75 and 82-88 are directed to constructs comprising an antibody to Annexin-IV and a detectable moiety, compositions comprising the same and methods of using the same. The feature of constructs comprising an antibody to Annexin-IV and a detectable moiety is specific to this group of claims.
- Claims 6-26, 36-46, 52-60, 69-71, 76-88 are directed to constructs comprising an antibody to phospholipids and a detectable moiety, compositions comprising the same and methods of using the same. The feature of constructs comprising an antibody to phospholipids and a detectable moiety is specific to this group of claims.
- Claims 61-88 are directed to methods relating to complement associated ocular inflammation/disease/injury comprising administering an antibody that does not activate complement and specifically binds to Annexin-IV or a phospholipid; or (b) a construct comprising: (i) an antibody that specifically binds to Annexin-IV or phospholipid, and (ii) a therapeutic agent or a detectable moiety. The feature of methods relating to complement associated ocular inflammation/disease/injury comprising administering an antibody that specifically binds to Annexin-IV or a phospholipid, or a construct based thereon is specific to this group of claims.

PCT Rule 13.2, first sentence, states that unity of invention is only fulfilled when there is a technical relationship among the claimed inventions involving one or more of the same or corresponding special technical features. PCT Rule 13.2, second sentence, defines a special technical feature as a feature which makes a contribution over the prior art.

When there is no special technical feature common to all the claimed inventions there is no unity of invention.

In the above groups of claims, the identified features may have the potential to make a contribution over the prior art but are not common to all the claimed inventions and therefore cannot provide the required technical relationship. The only feature common to all of the claimed inventions and which provides a technical relationship among them is an antibody to Annexin-IV or phospholipids.

However this feature does not make a contribution over the prior art because it is disclosed in:

D2 discloses detectably labelled antibodies to Annexin-IV or phospholipids, but fails to disclose other constructs or the use of the same in methods relating to inflammation or the complement cascade (see Box V for further details).

Therefore in the light of this document this common feature cannot be a special technical feature. Therefore there is no special technical feature common to all the claimed inventions and the requirements for unity of invention are consequently not satisfied *a posteriori*.

INTERNATIONAL SEARCH REPORT Information on patent family members		International application No. PCT/US2014/012831	
This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.			
Patent Document/s Cited in Search Report		Patent Family Member/s	
Publication Number	Publication Date	Publication Number	Publication Date
WO 2007/112403 A2	04 Oct 2007	US 2011014270 A1 US 2013171236 A1	20 Jan 2011 04 Jul 2013
CN 101235089 A	06 Aug 2008	None	
End of Annex			
Due to data integration issues this family listing may not include 10 digit Australian applications filed since May 2001. Form PCT/ISA/210 (Family Annex)(July 2009)			

SEQUENCE LISTING

<110> MUSC Foundation for Research Development
The Regents of the University of Colorado, a body corporate
Department of VETERANS Affairs (US)
Rohrer, Baerbel M.
Holers, Michael
Tomlinson, Stephen
Thurman, Joshua M.
Atkinson, Carl
Kulik, Liudmila

<120> TARGETING CONSTRUCTS BASED ON NATURAL ANTIBODIES AND USES THEREOF

<130> 48498-559001WO

<140> WO PCT/US2014/012831
<141> 2014-01-23

<150> US 61/755,968
<151> 2013-01-23

<150> US 61/755,960
<151> 2013-01-23

<150> US 61/771,565
<151> 2013-03-01

<150> US 61/771,560
<151> 2013-03-01

<160> 58

<170> PatentIn version 3.5

<210> 1
<211> 7
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic polypeptide

<400> 1

Ser Ser Ile Ser Ser Asn Tyr
1 5

<210> 2
<211> 3
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic polypeptide

<400> 2

Arg Thr Ser
1

<210> 3
 <211> 18
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 3

Gln Gln Gly Ser Ser Ile Pro Arg Thr Arg Ser Glu Gly Ala Pro Ser
 1 5 10 15

Trp Lys

<210> 4
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 4

Gly Tyr Thr Phe Thr Ser Tyr Trp
 1 5

<210> 5
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 5

Ile Gly Pro Asn Ser Gly Gly Thr
 1 5

<210> 6
 <211> 22
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 6

Ala Arg Arg Met Val Lys Gly Cys Tyr Gly Leu Leu Gly Pro Arg Asp
 1 5 10 15

His Gly His Arg Leu Leu
 20

<210> 7
 <211> 11
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 7

Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr
 1 5 10

<210> 8
 <211> 3
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 8

Lys Val Ser
 1

<210> 9
 <211> 9
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 9

Phe Gln Gly Ser His Val Pro Tyr Thr
 1 5

<210> 10
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 10

Gly Tyr Thr Phe Thr Asp Tyr Tyr
 1 5

<210> 11
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 11

Ile Asn Pro Asn Asn Gly Gly Thr
1 5

<210> 12

<211> 11

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 12

Ala Arg Tyr Asp Tyr Ala Trp Tyr Phe Asp Val
1 5 10

<210> 13

<211> 107

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 13

Asp Ile Glu Leu Thr Gln Ser Pro Thr Thr Met Ala Ala Ser Pro Gly
1 5 10 15

Glu Lys Ile Thr Ile Thr Cys Ser Ala Ser Ser Ser Ile Ser Ser Asn
20 25 30

Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Phe Ser Pro Lys Leu Leu
35 40 45

Ile Tyr Arg Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser
50 55 60

Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Gly Thr Met Glu
65 70 75 80

Ala Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Gly Ser Ser Ile Pro
85 90 95

Arg Thr Arg Ser Glu Gly Ala Pro Ser Trp Lys
100 105

<210> 14

<211> 112

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 14

Asp	Val	Leu	Met	Thr	Gln	Thr	Pro	Leu	Ser	Leu	Pro	Val	Ser	Leu	Gly
1				5					10					15	

Asp	Gln	Ala	Ser	Ile	Ser	Cys	Arg	Ser	Ser	Gln	Ser	Ile	Val	His	Ser
			20					25					30		

Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser
		35					40					45			

Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro
	50					55					60				

Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
65					70					75					80

Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Gly
				85					90					95	

Ser	His	Val	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys
			100					105					110		

<210> 15

<211> 117

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 15

Val	Lys	Leu	Gln	Glu	Ser	Gly	Ala	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser
1				5					10					15	

Val	Lys	Leu	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Trp
			20					25					30		

Met	His	Trp	Val	Lys	Gln	Arg	Pro	Gly	Arg	Gly	Leu	Glu	Trp	Ile	Gly
		35					40					45			

Arg	Ile	Gly	Pro	Asn	Ser	Gly	Gly	Thr	Lys	Tyr	Asn	Glu	Lys	Phe	Lys
	50					55					60				

Ser	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Pro	Ser	Ser	Thr	Ala	Tyr	Met
65					70					75					80

Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala
85 90 95

Arg Arg Met Val Lys Gly Cys Tyr Gly Leu Leu Gly Pro Arg Asp His
100 105 110

Gly His Arg Leu Leu
115

<210> 16
<211> 117
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic polypeptide

<400> 16

Val Lys Leu Gln Glu Ser Gly Pro Glu Leu Val Lys Pro Gly Ala Ser
1 5 10 15

Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr Tyr
20 25 30

Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile Gly
35 40 45

Asp Ile Asn Pro Asn Asn Gly Gly Thr Ser Tyr Asn Gln Lys Phe Lys
50 55 60

Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr Met
65 70 75 80

Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala
85 90 95

Arg Tyr Asp Tyr Ala Trp Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr
100 105 110

Val Thr Val Ser Ser
115

<210> 17
<211> 272
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic polypeptide

<400> 17

His His His His His His Val Lys Leu Gln Glu Ser Gly Ala Glu Leu
1 5 10 15

Val Lys Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr
20 25 30

Thr Phe Thr Ser Tyr Trp Met His Trp Val Lys Gln Arg Pro Gly Arg
35 40 45

Gly Leu Glu Trp Ile Gly Arg Ile Gly Pro Asn Ser Gly Gly Thr Lys
50 55 60

Tyr Asn Glu Lys Phe Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Pro
65 70 75 80

Ser Ser Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser
85 90 95

Ala Val Tyr Tyr Cys Ala Arg Arg Met Val Lys Gly Cys Tyr Gly Leu
100 105 110

Leu Gly Pro Arg Asp His Gly His Arg Leu Leu Lys Gly Arg Ile Pro
115 120 125

Ala His Trp Arg Pro Leu Leu Val Asp Pro Ser Ser Val Pro Ser Leu
130 135 140

Ala Ser Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Trp Ile Ser
145 150 155 160

Ala Glu Phe Ala Leu Asp Ile Glu Leu Thr Gln Ser Pro Thr Thr Met
165 170 175

Ala Ala Ser Pro Gly Glu Lys Ile Thr Ile Thr Cys Ser Ala Ser Ser
180 185 190

Ser Ile Ser Ser Asn Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Phe
195 200 205

Ser Pro Lys Leu Leu Ile Tyr Arg Thr Ser Asn Leu Ala Ser Gly Val
210 215 220

Pro Ala Arg Phe Ser Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr
225 230 235 240

Ile Gly Thr Met Glu Ala Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln
 245 250 255

Gly Ser Ser Ile Pro Arg Thr Arg Ser Glu Gly Ala Pro Ser Trp Lys
 260 265 270

<210> 18
 <211> 274
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 18

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Leu Trp Leu Thr
 1 5 10 15

Asp Ala Arg Cys Val Lys Leu Gln Glu Ser Gly Pro Glu Leu Val Lys
 20 25 30

Pro Gly Ala Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe
 35 40 45

Thr Asp Tyr Tyr Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu
 50 55 60

Glu Trp Ile Gly Asp Ile Asn Pro Asn Asn Gly Gly Thr Ser Tyr Asn
 65 70 75 80

Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser
 85 90 95

Thr Ala Tyr Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val
 100 105 110

Tyr Tyr Cys Ala Arg Tyr Asp Tyr Ala Trp Tyr Phe Asp Val Trp Gly
 115 120 125

Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly Gly
 130 135 140

Gly Gly Ser Gly Gly Gly Gly Asp Val Leu Met Thr Gln Thr Pro Leu
 145 150 155 160

Ser Leu Pro Val Ser Leu Gly Asp Gln Ala Ser Ile Ser Cys Arg Ser
 165 170 175

Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu Trp Tyr

180

185

190

Leu Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Lys Val Ser
 195 200 205

Asn Arg Phe Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly
 210 215 220

Thr Asp Phe Thr Leu Lys Ile Ser Arg Val Glu Ala Glu Asp Leu Gly
 225 230 235 240

Val Tyr Tyr Cys Phe Gln Gly Ser His Val Pro Tyr Thr Phe Gly Gly
 245 250 255

Gly Thr Lys Leu Glu Ile Lys Arg Ile Glu Gly Arg His His His His
 260 265 270

His His

<210> 19
 <211> 321
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 19
 gacattgagc tcacccagtc tccaaccacc atggctgcat ctcccgggga gaagatcact 60
 atcacctgca gtgccagctc aagtataagt tccaattact tgcattggta tcagcagaag 120
 ccaggattct cccctaaact cttgatttat aggacatcca atctggcttc tggagtccca 180
 gctcgcttca gtggcagtggt gtctgggacc tottactctc tcacaattgg caccatggag 240
 gctgaagatg ttgccactta ctactgccag cagggtagta gtataccacg tacacgttcg 300
 gagggggcac caagctggaa a 321

<210> 20
 <211> 338
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 20
 gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120
 tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180

tctgggggtcc cagacaggtt cagtggcagt ggatcagggg cagatttcac actcaagatc 240
 agcagagtgg aggctgagga tctgggagtt tattactgct ttcaagggtc acatgttccg 300
 tacacgttcg gaggggggac caagctggaa ataaaacg 338

<210> 21
 <211> 352
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 21
 gtgaaactgc aggagtcagg ggctgagctt gtgaagcctg gggcttcagt gaagctgtcc 60
 tgcaaggctt ctggctacac cttcaccagc tactggatgc actgggtgaa gcagaggcct 120
 ggacgaggcc ttgagtggat tgggaaggatt ggtcctaata gtggtggtac taagtacaat 180
 gagaagttca agagcaaggc cacactgact gtagacaaac cctccagcac agcctacatg 240
 cagctcagca gcctgacatc tgaggactct gcggtctatt attgtgcaag aagaatggta 300
 aaggggtgct atggactact ggggccaaagg gaccacggtc accgtctcct ca 352

<210> 22
 <211> 351
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 22
 gtgaagctgc aggagtctgg acctgagctg gtgaagcctg gggcttcagt gaagatatcc 60
 tgtaaggctt ctggatacac gttcactgac tactacatga actgggtgaa gcagagccat 120
 ggaaagagcc ttgagtggat tggagatatt aatcctaaca atggtggtac tagctacaac 180
 cagaagttca agggcaaggc cacattgact gtagacaagt cctccagcac agcctacatg 240
 gagctccgca gcctgacatc tgaggactct gcagtctatt actgtgcaag atatgattac 300
 gcttgggtact tcgatgtctg gggccaaggg accacggtca ccgctctctc a 351

<210> 23
 <211> 837
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 23
 gccgccacca tgagtgtgcc cactcaggtc ctgggggtgc tgctgctgtg gcttacagat 60

```

gccagatgtg tgaagctgca ggagtctgga cctgagctgg tgaagcctgg ggcttcagtg      120
aagatatcct gtaaggcttc tggatacacg ttactgact actacatgaa ctgggtgaag      180
cagagccatg gaaagagcct tgagtggatt ggagatatta atcctaacaa tgggtggtact      240
agctacaacc agaagttcaa gggcaaggcc acattgactg tagacaagtc ctccagcaca      300
gcctacatgg agctccgcag cctgacatct gaggactctg cagtctatta ctgtgcaaga      360
tatgattacg cttggtactt cgatgtctgg ggccaaggga ccacgggtcac cgtctcctca      420
ggcggaggtg ggtcgggtgg cggcgatct ggcgaggtg gggatgttt gatgacccaa      480
actccactct ccctgcctgt cagtcttggga gatcaagcct ccatctcttg cagatctagt      540
cagagcattg tacatagtaa tggaaacacc tatttagaat ggtacctgca gaaaccaggc      600
cagtctcaa agctcctgat ctacaaagtt tccaaccgat tttctgggt cccagacagg      660
ttcagtggca gtggatcagg gacagatttc aactcaaga tcagcagagt ggaggctgag      720
gatctgggag tttattactg ctttcaaggt tcacatgttc cgtacacgtt cggagggggg      780
accaagctgg aaataaaacg gatcgaaggc cggcatcacc atcatcacca ctgatag      837

```

```

<210> 24
<211> 825
<212> DNA
<213> Artificial sequence

```

```

<220>
<223> Synthetic polynucleotide

```

```

<400> 24
atgtccgtgc ctaccaggt gctcggactc ctgctgctgt ggctcaccga cgccaggtgt      60
gtgaagctgc aggagagcgg acccgagctg gtgaagcctg gaggctccgt gaagatcagc      120
tgcaaggctt ccggatacac cttcacgcac tactatatga actgggtgaa gcagagccac      180
ggcaagagcc tggagtggat cggcgacatc aaccctaaca acggcggcac ctctacaac      240
cagaagttca agggcaaggc tacactgacc gtggacaagt cctccagcac cgcctacatg      300
gagctcagga gcctgacctc cgaggattcc gccgtctatt actgtgcccg gtacgactac      360
gcctggtatt tcgacgtgtg gggccagggc acaaccgtca cagtctccag cggaggagga      420
ggaagcggcg gcggaggatc cggaggcgga ggcgatgtcc tgatgacaca gacacctctg      480
agcctccccg tgagcctggg agaccaagcc tccatctcct gcaggtcctc ccagtcctc      540
gtgcacagca atggcaacac ctacctggag tggatatctgc agaagcctgg ccagtcctcc      600
aagctgctga tctacaaggt gtccaaccgg ttcagcggcg tccctgacag gttctccgga      660
tccggaagcg gcacagattt caccctgaag atcagcaggg tcgaggccga ggacctggga      720
gtgtactact gcttccaggc ctcccatgtc ccttacacct tcggcggcgg caccaaactg      780
gagatcaagc ggatcgaggg caggcatcac caccatcacc actga      825

```

<210> 25
 <211> 10
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 25

Lys	Ser	Val	Ser	Thr	Ser	Gly	Tyr	Ser	Tyr
1				5					10

<210> 26
 <211> 3
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 26

Leu	Val	Ser
1		

<210> 27
 <211> 16
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 27

Gln	His	Ile	Arg	Glu	Leu	Thr	Arg	Ser	Glu	Gly	Gly	Pro	Ser	Trp	Lys
1				5					10					15	

<210> 28
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 28

Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Trp
1				5			

<210> 29
 <211> 8
 <212> PRT
 <213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 29

Ile Asn Pro Ser Asn Gly Gly Thr
1 5

<210> 30

<211> 12

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 30

Ala Arg Arg Gly Ile Arg Leu Arg His Phe Asp Tyr
1 5 10

<210> 31

<211> 6

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 31

Gln Asp Val Gly Thr Ala
1 5

<210> 32

<211> 3

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 32

Trp Ala Ser
1

<210> 33

<211> 9

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 33

Gln Gln Tyr Ser Ser Tyr Pro Leu Thr
1 5

<210> 34
 <211> 108
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 34

Asp Ile Val Met Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

Gln Arg Ala Thr Ile Ser Tyr Arg Ala Ser Lys Ser Val Ser Thr Ser
 20 25 30

Gly Tyr Ser Tyr Met His Trp Asn Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

Arg Leu Leu Ile Tyr Leu Val Ser Asn Leu Glu Ser Gly Val Pro Ala
 50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
 65 70 75 80

Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys Gln His Ile Arg
 85 90 95

Glu Leu Thr Arg Ser Glu Gly Gly Pro Ser Trp Lys
 100 105

<210> 35
 <211> 107
 <212> PRT
 <213> Artificial sequence

<220>
 <223> Synthetic polypeptide

<400> 35

Asp Ile Gln Met Thr Gln Ser Pro Lys Phe Met Ser Thr Ser Val Gly
 1 5 10 15

Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Thr Ala
 20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
 35 40 45

Tyr Trp Ala Ser Thr Arg His Thr Gly Val Pro Asp Arg Phe Thr Gly

50

55

60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Asn Val Gln Ser
65 70 75 80

Glu Asp Leu Ala Asp Tyr Phe Cys Gln Gln Tyr Ser Ser Tyr Pro Leu
85 90 95

Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100 105

<210> 36

<211> 117

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 36

Val Lys Leu Gln Glu Ser Gly Thr Glu Leu Val Lys Pro Gly Ala Ser
1 5 10 15

Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Pro Thr Ser Tyr Trp
20 25 30

Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile Gly
35 40 45

Asn Ile Asn Pro Ser Asn Gly Gly Thr Asn Tyr Asn Glu Lys Phe Lys
50 55 60

Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr Met
65 70 75 80

Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys Ala
85 90 95

Arg Arg Gly Ile Arg Leu Arg His Phe Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Thr Val Thr Val Ser
115

<210> 37

<211> 275

<212> PRT

<213> Artificial sequence

<220>

<223> Synthetic polypeptide

<400> 37

Val	Lys	Leu	Gln	Glu	Ser	Gly	Thr	Glu	Leu	Val	Lys	Pro	Gly	Ala	Ser
1				5					10					15	

Val	Lys	Leu	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr	Trp
			20					25					30		

Met	His	Trp	Val	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	Gly
		35					40					45			

Asn	Ile	Asn	Pro	Ser	Asn	Gly	Gly	Thr	Asn	Tyr	Asn	Glu	Lys	Phe	Lys
	50					55					60				

Ser	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr	Met
65					70					75					80

Gln	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	Ala
				85					90					95	

Arg	Arg	Gly	Ile	Arg	Leu	Arg	His	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr
			100					105					110		

Thr	Val	Thr	Val	Ser	Ser	Arg	Ala	Asn	Ser	Ala	Asp	Ile	His	His	Thr
		115					120					125			

Gly	Gly	Arg	Ser	Ser	Met	His	Leu	Glu	Gly	Pro	Ile	Arg	Pro	Ile	Val
	130					135					140				

Ser	Arg	Ile	Ser	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Trp
145					150					155					160

Ile	Ser	Ala	Glu	Phe	Ala	Leu	Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Ala
				165					170					175	

Ser	Leu	Ala	Val	Ser	Leu	Gly	Gln	Arg	Ala	Thr	Ile	Ser	Tyr	Arg	Ala
			180					185					190		

Ser	Lys	Ser	Val	Ser	Thr	Ser	Gly	Tyr	Ser	Tyr	Met	His	Trp	Asn	Gln
		195					200					205			

Gln	Lys	Pro	Gly	Gln	Pro	Pro	Arg	Leu	Leu	Ile	Tyr	Leu	Val	Ser	Asn
	210					215					220				

Leu	Glu	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr
225					230					235					240

Asp Phe Thr Leu Asn Ile His Pro Val Glu Glu Glu Asp Ala Ala Thr
245 250 255

Tyr Tyr Cys Gln His Ile Arg Glu Leu Thr Arg Ser Glu Gly Gly Pro
260 265 270

Ser Trp Lys
275

<210> 38
<211> 271
<212> PRT
<213> Artificial sequence

<220>
<223> Synthetic polypeptide

<400> 38

Met Ser Val Pro Thr Gln Val Leu Gly Leu Leu Leu Trp Leu Thr
1 5 10 15

Asp Ala Arg Cys Val Lys Leu Gln Glu Ser Gly Thr Glu Leu Val Lys
20 25 30

Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
35 40 45

Thr Ser Tyr Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
50 55 60

Glu Trp Ile Gly Asn Ile Asn Pro Ser Asn Gly Gly Thr Asn Tyr Asn
65 70 75 80

Glu Lys Phe Lys Ser Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser
85 90 95

Thr Ala Tyr Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val
100 105 110

Tyr Tyr Cys Ala Arg Arg Gly Ile Arg Leu Arg His Phe Asp Tyr Trp
115 120 125

Gly Gln Gly Thr Thr Val Thr Val Ser Ser Gly Gly Gly Gly Ser Gly
130 135 140

Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Ile Gln Met Thr Gln Ser
145 150 155 160

Pro Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Ile Thr Cys
 165 170 175

Lys Ala Ser Gln Asp Val Gly Thr Ala Val Ala Trp Tyr Gln Gln Lys
 180 185 190

Pro Gly Gln Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg His
 195 200 205

Thr Gly Val Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe
 210 215 220

Thr Leu Thr Ile Ser Asn Val Gln Ser Glu Asp Leu Ala Asp Tyr Phe
 225 230 235 240

Cys Gln Gln Tyr Ser Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys
 245 250 255

Leu Glu Leu Lys Arg Ile Glu Gly Arg His His His His His His
 260 265 270

<210> 39
 <211> 324
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 39
 gacattgtga tgacacagtc tcttgcttcc ttagctgtat ctctggggca gagggccacc 60
 atctcataca gggccagcaa aagtgtcagt acatctggct atagttatat gcactggaac 120
 caacagaaac caggacagcc acccagactc ctcatctatc ttgtatccaa cctagaatct 180
 ggggtccctg ccaggttcag tggcagtggt tctgggacag acttcaccct caacatccat 240
 cctgtggagg aggaggatgc tgcaacctat tactgtcagc acattaggga gcttacacgt 300
 tcggaggggg gaccaagctg gaaa 324

<210> 40
 <211> 322
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 40
 gacatccaga tgaccagtc tcccaaattc atgtccacat cagtaggaga cagggtcagc 60
 atcacctgca aggcagtc ggaatgtgggt actgctgtag cctgggtatca acagaaacca 120

gggcaatctc ctaaactact gatttactgg gcatccaccc ggcacactgg agtccctgat 180
 cgcttcacag gcagtggatc tgggacagat ttactctca ccattagcaa tgtgcagtct 240
 gaagacttgg cagattatct ctgtcagcaa tatagcagct atcctctcac gttcgggtgct 300
 gggaccaagc tggagctgaa ac 322

<210> 41
 <211> 351
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 41
 gtgaaactgc aggagtctgg gactgaactg gtgaagcctg gggcttcagt gaagctgtcc 60
 tgcaaggctt ctggctacac cttcaccagc tactggatgc actgggtgaa gcagaggcct 120
 ggacaaggcc ttgagtggat tggaaatatt aatcctagca atggtggtac taactacaat 180
 gagaagttca agagcaaggc cactgact gtagacaaat cctccagcac agcctacatg 240
 cagctcagca gcctgacatc tgaggactct gcggtctatt attgtgcaag aagaggcata 300
 cggttacgac actttgacta ctggggccaa gggaccacgg tcaccgtctc c 351

<210> 42
 <211> 828
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 42
 gccgccacca tgagtgtgcc cactcaggtc ctgggggttc tgctgctgtg gcttacagat 60
 gccagatgtg tgaaactgca ggagtctggg actgaactgg tgaagcctgg ggcttcagtg 120
 aagctgtcct gcaaggcttc tggctacacc ttaccagct actggatgca ctgggtgaag 180
 cagaggcctg gacaaggcct tgagtggatt ggaaatatta atcctagcaa tgggtgtact 240
 aactacaatg agaagttcaa gagcaaggcc aactgactg tagacaaatc ctccagcaca 300
 gcctacatgc agctcagcag cctgacatct gaggactctg cggctctatta ttgtgcaaga 360
 agaggcatac ggttacgaca ctttgactac tggggccaag ggaccacggt caccgtctcc 420
 tctggcggag gtgggtcggg tggcggcgga tctggcggag gtgggtcggg catccagatg 480
 acccagtctc ccaaattcat gtccacatca gtaggagaca gggtcagcat cacctgcaag 540
 gccagtcagg atgtgggtac tgctgtagcc tggatatcaac agaaaccagg gcaatctcct 600
 aaactactga ttactgggc atccaccgg cactgagag tccctgatcg cttcacaggc 660
 agtggatctg ggacagattt cactctcacc attagcaatg tgcagtctga agacttgga 720

gattatttct gtcagcaata tagcagctat cctctcacgt tcggtgctgg gaccaagctg 780
gagctgaaac ggatcgaagg ccggcatcac catcatcacc actgatag 828

<210> 43
<211> 819
<212> DNA
<213> Artificial sequence

<220>
<223> Synthetic polynucleotide

<400> 43
atgagcgtgc ctacacaggt gctcggcctg ctgctcctct ggctgacaga cgcccgggtgt 60
gtgaagctgc aggagtccgg aaccgagctg gtgaaacctg gcgccagcgt gaaactgagc 120
tgcaaagcca gcggatacac cttcacctcc tactggatgc actgggtgaa acagaggcct 180
ggccagggcc tggaatggat tggcaacatc aaccccagca acggcggcac caactacaat 240
gagaagttca agagcaaggc caccctgacc gtggataagt cctcctccac cgcctacatg 300
cagctgtcct ccctcacctc cgaggacagc gccgtctatt actgtgccag gcggggcatc 360
aggctgaggg acttcgacta ctggggccaa ggcacaaccg tcaccgtgag ctccggagga 420
ggaggcagcg gaggcggagg ctccggcgga ggcggaagcg acattcagat gaccagagc 480
cccaagttca tgtccacctc cgtcggcgac agggtgagca tcacctgtaa ggccagccag 540
gatgtcggca cagctgtggc ctggtaccag cagaagcccg gccagtcccc caagctgctg 600
atctactggg cttccacaag gcataccggc gtccccgata ggttcacagg ctccggctcc 660
ggcaccgact tcacactcac catcagcaac gtccagtccg aggacctggc cgactacttc 720
tgccagcagt actccagcta cccctcacc ttcggcgctg gcaccaagct ggaactcaag 780
cggatcgagg gcaggcatca ccaccatcac cactgatag 819

<210> 44
<211> 392
<212> PRT
<213> Homo sapiens

<400> 44

Met Glu Pro Pro Gly Arg Arg Glu Cys Pro Phe Pro Ser Trp Arg Phe
1 5 10 15

Pro Gly Leu Leu Leu Ala Ala Met Val Leu Leu Leu Tyr Ser Phe Ser
20 25 30

Asp Ala Cys Glu Glu Pro Pro Thr Phe Glu Ala Met Glu Leu Ile Gly
35 40 45

Lys	Pro	Lys	Pro	Tyr	Tyr	Glu	Ile	Gly	Glu	Arg	Val	Asp	Tyr	Lys	Cys		
50						55					60						
Lys	Lys	Gly	Tyr	Phe	Tyr	Ile	Pro	Pro	Leu	Ala	Thr	His	Thr	Ile	Cys		
65					70					75					80		
Asp	Arg	Asn	His	Thr	Trp	Leu	Pro	Val	Ser	Asp	Asp	Ala	Cys	Tyr	Arg		
				85					90					95			
Glu	Thr	Cys	Pro	Tyr	Ile	Arg	Asp	Pro	Leu	Asn	Gly	Gln	Ala	Val	Pro		
			100					105					110				
Ala	Asn	Gly	Thr	Tyr	Glu	Phe	Gly	Tyr	Gln	Met	His	Phe	Ile	Cys	Asn		
		115					120					125					
Glu	Gly	Tyr	Tyr	Leu	Ile	Gly	Glu	Glu	Ile	Leu	Tyr	Cys	Glu	Leu	Lys		
	130					135					140						
Gly	Ser	Val	Ala	Ile	Trp	Ser	Gly	Lys	Pro	Pro	Ile	Cys	Glu	Lys	Val		
145					150					155					160		
Leu	Cys	Thr	Pro	Pro	Pro	Lys	Ile	Lys	Asn	Gly	Lys	His	Thr	Phe	Ser		
				165					170					175			
Glu	Val	Glu	Val	Phe	Glu	Tyr	Leu	Asp	Ala	Val	Thr	Tyr	Ser	Cys	Asp		
			180					185					190				
Pro	Ala	Pro	Gly	Pro	Asp	Pro	Phe	Ser	Leu	Ile	Gly	Glu	Ser	Thr	Ile		
		195					200					205					
Tyr	Cys	Gly	Asp	Asn	Ser	Val	Trp	Ser	Arg	Ala	Ala	Pro	Glu	Cys	Lys		
	210					215					220						
Val	Val	Lys	Cys	Arg	Phe	Pro	Val	Val	Glu	Asn	Gly	Lys	Gln	Ile	Ser		
225					230					235					240		
Gly	Phe	Gly	Lys	Lys	Phe	Tyr	Tyr	Lys	Ala	Thr	Val	Met	Phe	Glu	Cys		
				245					250					255			
Asp	Lys	Gly	Phe	Tyr	Leu	Asp	Gly	Ser	Asp	Thr	Ile	Val	Cys	Asp	Ser		
			260					265					270				
Asn	Ser	Thr	Trp	Asp	Pro	Pro	Val	Pro	Lys	Cys	Leu	Lys	Val	Leu	Pro		
		275					280					285					
Pro	Ser	Ser	Thr	Lys	Pro	Pro	Ala	Leu	Ser	His	Ser	Val	Ser	Thr	Ser		
	290					295					300						

Ser Thr Thr Lys Ser Pro Ala Ser Ser Ala Ser Gly Pro Arg Pro Thr
305 310 315 320

Tyr Lys Pro Pro Val Ser Asn Tyr Pro Gly Tyr Pro Lys Pro Glu Glu
325 330 335

Gly Ile Leu Asp Ser Leu Asp Val Trp Val Ile Ala Val Ile Val Ile
340 345 350

Ala Ile Val Val Gly Val Ala Val Ile Cys Val Val Pro Tyr Arg Tyr
355 360 365

Leu Gln Arg Arg Lys Lys Lys Gly Thr Tyr Leu Thr Asp Glu Thr His
370 375 380

Arg Glu Val Lys Phe Thr Ser Leu
385 390

<210> 45
<211> 381
<212> PRT
<213> Homo sapiens

<400> 45

Met Thr Val Ala Arg Pro Ser Val Pro Ala Ala Leu Pro Leu Leu Gly
1 5 10 15

Glu Leu Pro Arg Leu Leu Leu Leu Val Leu Leu Cys Leu Pro Ala Val
20 25 30

Trp Gly Asp Cys Gly Leu Pro Pro Asp Val Pro Asn Ala Gln Pro Ala
35 40 45

Leu Glu Gly Arg Thr Ser Phe Pro Glu Asp Thr Val Ile Thr Tyr Lys
50 55 60

Cys Glu Glu Ser Phe Val Lys Ile Pro Gly Glu Lys Asp Ser Val Ile
65 70 75 80

Cys Leu Lys Gly Ser Gln Trp Ser Asp Ile Glu Glu Phe Cys Asn Arg
85 90 95

Ser Cys Glu Val Pro Thr Arg Leu Asn Ser Ala Ser Leu Lys Gln Pro
100 105 110

Tyr Ile Thr Gln Asn Tyr Phe Pro Val Gly Thr Val Val Glu Tyr Glu
115 120 125

Cys Arg Pro Gly Tyr Arg Arg Glu Pro Ser Leu Ser Pro Lys Leu Thr
 130 135 140

Cys Leu Gln Asn Leu Lys Trp Ser Thr Ala Val Glu Phe Cys Lys Lys
 145 150 155 160

Lys Ser Cys Pro Asn Pro Gly Glu Ile Arg Asn Gly Gln Ile Asp Val
 165 170 175

Pro Gly Gly Ile Leu Phe Gly Ala Thr Ile Ser Phe Ser Cys Asn Thr
 180 185 190

Gly Tyr Lys Leu Phe Gly Ser Thr Ser Ser Phe Cys Leu Ile Ser Gly
 195 200 205

Ser Ser Val Gln Trp Ser Asp Pro Leu Pro Glu Cys Arg Glu Ile Tyr
 210 215 220

Cys Pro Ala Pro Pro Gln Ile Asp Asn Gly Ile Ile Gln Gly Glu Arg
 225 230 235 240

Asp His Tyr Gly Tyr Arg Gln Ser Val Thr Tyr Ala Cys Asn Lys Gly
 245 250 255

Phe Thr Met Ile Gly Glu His Ser Ile Tyr Cys Thr Val Asn Asn Asp
 260 265 270

Glu Gly Glu Trp Ser Gly Pro Pro Pro Glu Cys Arg Gly Lys Ser Leu
 275 280 285

Thr Ser Lys Val Pro Pro Thr Val Gln Lys Pro Thr Thr Val Asn Val
 290 295 300

Pro Thr Thr Glu Val Ser Pro Thr Ser Gln Lys Thr Thr Thr Lys Thr
 305 310 315 320

Thr Thr Pro Asn Ala Gln Ala Thr Arg Ser Thr Pro Val Ser Arg Thr
 325 330 335

Thr Lys His Phe His Glu Thr Thr Pro Asn Lys Gly Ser Gly Thr Thr
 340 345 350

Ser Gly Thr Thr Arg Leu Leu Ser Gly His Thr Cys Phe Thr Leu Thr
 355 360 365

Gly Leu Leu Gly Thr Leu Val Thr Met Gly Leu Leu Thr
 370 375 380

<210> 46
 <211> 390
 <212> PRT
 <213> Mus musculus

<400> 46

Met Ile Arg Gly Arg Ala Pro Arg Thr Arg Pro Ser Pro Pro Pro Pro
 1 5 10 15

Leu Leu Pro Leu Leu Ser Leu Ser Leu Leu Leu Ser Pro Thr Val
 20 25 30

Arg Gly Asp Cys Gly Pro Pro Pro Asp Ile Pro Asn Ala Arg Pro Ile
 35 40 45

Leu Gly Arg His Ser Lys Phe Ala Glu Gln Ser Lys Val Ala Tyr Ser
 50 55 60

Cys Asn Asn Gly Phe Lys Gln Val Pro Asp Lys Ser Asn Ile Val Val
 65 70 75 80

Cys Leu Glu Asn Gly Gln Trp Ser Ser His Glu Thr Phe Cys Glu Lys
 85 90 95

Ser Cys Val Ala Pro Glu Arg Leu Ser Phe Ala Ser Leu Lys Lys Glu
 100 105 110

Tyr Leu Asn Met Asn Phe Phe Pro Val Gly Thr Ile Val Glu Tyr Glu
 115 120 125

Cys Arg Pro Gly Phe Arg Lys Gln Pro Pro Leu Pro Gly Lys Ala Thr
 130 135 140

Cys Leu Glu Asp Leu Val Trp Ser Pro Val Ala Gln Phe Cys Lys Lys
 145 150 155 160

Lys Ser Cys Pro Asn Pro Lys Asp Leu Asp Asn Gly His Ile Asn Ile
 165 170 175

Pro Thr Gly Ile Leu Phe Gly Ser Glu Ile Asn Phe Ser Cys Asn Pro
 180 185 190

Gly Tyr Arg Leu Val Gly Val Ser Ser Thr Phe Cys Ser Val Thr Gly
 195 200 205

Asn Thr Val Asp Trp Asp Asp Glu Phe Pro Val Cys Thr Glu Ile His
 210 215 220

Cys Pro Glu Pro Pro Lys Ile Asn Asn Gly Ile Met Arg Gly Glu Ser
225 230 235 240

Asp Ser Tyr Thr Tyr Ser Gln Val Val Thr Tyr Ser Cys Asp Lys Gly
245 250 255

Phe Ile Leu Val Gly Asn Ala Ser Ile Tyr Cys Thr Val Ser Lys Ser
260 265 270

Asp Val Gly Gln Trp Ser Ser Pro Pro Pro Arg Cys Ile Glu Lys Ser
275 280 285

Lys Val Pro Thr Lys Lys Pro Thr Ile Asn Val Pro Ser Thr Gly Thr
290 295 300

Pro Ser Thr Pro Gln Lys Pro Thr Thr Glu Ser Val Pro Asn Pro Gly
305 310 315 320

Asp Gln Pro Thr Pro Gln Lys Pro Ser Thr Val Lys Val Ser Ala Thr
325 330 335

Gln His Val Pro Val Thr Lys Thr Thr Val Arg His Pro Ile Arg Thr
340 345 350

Ser Thr Asp Lys Gly Glu Pro Asn Thr Gly Gly Asp Arg Tyr Ile Tyr
355 360 365

Gly His Thr Cys Leu Ile Thr Leu Thr Val Leu His Val Met Leu Ser
370 375 380

Leu Ile Gly Tyr Leu Thr
385 390

<210> 47
<211> 128
<212> PRT
<213> Homo sapiens

<400> 47

Met Gly Ile Gln Gly Gly Ser Val Leu Phe Gly Leu Leu Leu Val Leu
1 5 10 15

Ala Val Phe Cys His Ser Gly His Ser Leu Gln Cys Tyr Asn Cys Pro
20 25 30

Asn Pro Thr Ala Asp Cys Lys Thr Ala Val Asn Cys Ser Ser Asp Phe
35 40 45

Asp Ala Cys Leu Ile Thr Lys Ala Gly Leu Gln Val Tyr Asn Lys Cys
50 55 60

Trp Lys Phe Glu His Cys Asn Phe Asn Asp Val Thr Thr Arg Leu Arg
65 70 75 80

Glu Asn Glu Leu Thr Tyr Tyr Cys Cys Lys Lys Asp Leu Cys Asn Phe
85 90 95

Asn Glu Gln Leu Glu Asn Gly Gly Thr Ser Leu Ser Glu Lys Thr Val
100 105 110

Leu Leu Leu Val Thr Pro Phe Leu Ala Ala Ala Trp Ser Leu His Pro
115 120 125

<210> 48
<211> 123
<212> PRT
<213> Mus musculus

<400> 48

Met Arg Ala Gln Arg Gly Leu Ile Leu Leu Leu Leu Leu Ala Val
1 5 10 15

Phe Cys Ser Thr Ala Val Ser Leu Thr Cys Tyr His Cys Phe Gln Pro
20 25 30

Val Val Ser Ser Cys Asn Met Asn Ser Thr Cys Ser Pro Asp Gln Asp
35 40 45

Ser Cys Leu Tyr Ala Val Ala Gly Met Gln Val Tyr Gln Arg Cys Trp
50 55 60

Lys Gln Ser Asp Cys His Gly Glu Ile Ile Met Asp Gln Leu Glu Glu
65 70 75 80

Thr Lys Leu Lys Phe Arg Cys Cys Gln Phe Asn Leu Cys Asn Lys Ser
85 90 95

Asp Gly Ser Leu Gly Lys Thr Pro Leu Leu Gly Thr Ser Val Leu Val
100 105 110

Ala Ile Leu Asn Leu Cys Phe Leu Ser His Leu
115 120

<210> 49
<211> 129

<212> PRT

<213> Mus musculus

<400> 49

Met Arg Ala Gln Arg Gly Leu Ile Leu Leu Leu Leu Leu Ala Val
 1 5 10 15

Phe Cys Ser Thr Ala Val Ser Leu Lys Cys Tyr Asn Cys Phe Gln Phe
 20 25 30

Val Ser Ser Cys Lys Ile Asn Thr Thr Cys Ser Pro Asn Leu Asp Ser
 35 40 45

Cys Leu Tyr Ala Val Ala Gly Arg Gln Val Tyr Gln Gln Cys Trp Lys
 50 55 60

Leu Ser Asp Cys Asn Ser Asn Tyr Ile Met Ser Arg Leu Asp Val Ala
 65 70 75 80

Gly Ile Gln Ser Lys Cys Cys Gln Trp Gly Leu Cys Asn Lys Asn Leu
 85 90 95

Asp Gly Leu Glu Glu Pro Asn Asn Ala Glu Thr Ser Ser Leu Arg Lys
 100 105 110

Thr Ala Leu Leu Gly Thr Ser Val Leu Val Ala Ile Leu Lys Phe Cys
 115 120 125

Phe

<210> 50

<211> 483

<212> PRT

<213> Mus musculus

<400> 50

Met Glu Val Ser Ser Arg Ser Ser Glu Pro Leu Asp Pro Val Trp Leu
 1 5 10 15

Leu Val Ala Phe Gly Arg Gly Gly Val Lys Leu Glu Val Leu Leu Leu
 20 25 30

Phe Leu Leu Pro Phe Thr Leu Gly Glu Leu Arg Gly Gly Leu Gly Lys
 35 40 45

His Gly His Thr Val His Arg Glu Pro Ala Val Asn Arg Leu Cys Ala
 50 55 60

Asp Ser Lys Arg Trp Ser Gly Leu Pro Val Ser Ala Gln Arg Pro Phe
 65 70 75 80

 Pro Met Gly His Cys Pro Ala Pro Ser Gln Leu Pro Ser Ala Lys Pro
 85 90 95

 Ile Asn Leu Thr Asp Glu Ser Met Phe Pro Ile Gly Thr Tyr Leu Leu
 100 105 110

 Tyr Glu Cys Leu Pro Gly Tyr Ile Lys Arg Gln Phe Ser Ile Thr Cys
 115 120 125

 Lys Gln Asp Ser Thr Trp Thr Ser Ala Glu Asp Lys Cys Ile Arg Lys
 130 135 140

 Gln Cys Lys Thr Pro Ser Asp Pro Glu Asn Gly Leu Val His Val His
 145 150 155 160

 Thr Gly Ile Gln Phe Gly Ser Arg Ile Asn Tyr Thr Cys Asn Gln Gly
 165 170 175

 Tyr Arg Leu Ile Gly Ser Ser Ser Ala Val Cys Val Ile Thr Asp Gln
 180 185 190

 Ser Val Asp Trp Asp Thr Glu Ala Pro Ile Cys Glu Trp Ile Pro Cys
 195 200 205

 Glu Ile Pro Pro Gly Ile Pro Asn Gly Asp Phe Phe Ser Ser Thr Arg
 210 215 220

 Glu Asp Phe His Tyr Gly Met Val Val Thr Tyr Arg Cys Asn Thr Asp
 225 230 235 240

 Ala Arg Gly Lys Ala Leu Phe Asn Leu Val Gly Glu Pro Ser Leu Tyr
 245 250 255

 Cys Thr Ser Asn Asp Gly Glu Ile Gly Val Trp Ser Gly Pro Pro Pro
 260 265 270

 Gln Cys Ile Glu Leu Asn Lys Cys Thr Pro Pro Pro Tyr Val Glu Asn
 275 280 285

 Ala Val Met Leu Ser Glu Asn Arg Ser Leu Phe Ser Leu Arg Asp Ile
 290 295 300

 Val Glu Phe Arg Cys His Pro Gly Phe Ile Met Lys Gly Ala Ser Ser
 305 310 315 320

Val His Cys Gln Ser Leu Asn Lys Trp Glu Pro Glu Leu Pro Ser Cys
325 330 335

Phe Lys Gly Val Ile Cys Arg Leu Pro Gln Glu Met Ser Gly Phe Gln
340 345 350

Lys Gly Leu Gly Met Lys Lys Glu Tyr Tyr Tyr Gly Glu Asn Val Thr
355 360 365

Leu Glu Cys Glu Asp Gly Tyr Thr Leu Glu Gly Ser Ser Gln Ser Gln
370 375 380

Cys Gln Ser Asp Gly Ser Trp Asn Pro Leu Leu Ala Lys Cys Val Ser
385 390 395 400

Arg Ser Ile Ser Gly Leu Ile Val Gly Ile Phe Ile Gly Ile Ile Val
405 410 415

Phe Ile Leu Val Ile Ile Val Phe Ile Trp Met Ile Leu Lys Tyr Lys
420 425 430

Lys Arg Asn Thr Thr Asp Glu Lys Tyr Lys Glu Val Gly Ile His Leu
435 440 445

Asn Tyr Lys Glu Asp Ser Cys Val Arg Leu Gln Ser Leu Leu Thr Ser
450 455 460

Gln Glu Asn Ser Ser Thr Thr Ser Pro Ala Arg Asn Ser Leu Thr Gln
465 470 475 480

Glu Val Ser

<210> 51
<211> 2039
<212> PRT
<213> Homo sapiens

<400> 51

Met Gly Ala Ser Ser Pro Arg Ser Pro Glu Pro Val Gly Pro Pro Ala
1 5 10 15

Pro Gly Leu Pro Phe Cys Cys Gly Gly Ser Leu Leu Ala Val Val Val
20 25 30

Leu Leu Ala Leu Pro Val Ala Trp Gly Gln Cys Asn Ala Pro Glu Trp
35 40 45

Leu Pro Phe Ala Arg Pro Thr Asn Leu Thr Asp Glu Phe Glu Phe Pro
50 55 60

Ile Gly Thr Tyr Leu Asn Tyr Glu Cys Arg Pro Gly Tyr Ser Gly Arg
65 70 75 80

Pro Phe Ser Ile Ile Cys Leu Lys Asn Ser Val Trp Thr Gly Ala Lys
85 90 95

Asp Arg Cys Arg Arg Lys Ser Cys Arg Asn Pro Pro Asp Pro Val Asn
100 105 110

Gly Met Val His Val Ile Lys Gly Ile Gln Phe Gly Ser Gln Ile Lys
115 120 125

Tyr Ser Cys Thr Lys Gly Tyr Arg Leu Ile Gly Ser Ser Ser Ala Thr
130 135 140

Cys Ile Ile Ser Gly Asp Thr Val Ile Trp Asp Asn Glu Thr Pro Ile
145 150 155 160

Cys Asp Arg Ile Pro Cys Gly Leu Pro Pro Thr Ile Thr Asn Gly Asp
165 170 175

Phe Ile Ser Thr Asn Arg Glu Asn Phe His Tyr Gly Ser Val Val Thr
180 185 190

Tyr Arg Cys Asn Pro Gly Ser Gly Gly Arg Lys Val Phe Glu Leu Val
195 200 205

Gly Glu Pro Ser Ile Tyr Cys Thr Ser Asn Asp Asp Gln Val Gly Ile
210 215 220

Trp Ser Gly Pro Ala Pro Gln Cys Ile Ile Pro Asn Lys Cys Thr Pro
225 230 235 240

Pro Asn Val Glu Asn Gly Ile Leu Val Ser Asp Asn Arg Ser Leu Phe
245 250 255

Ser Leu Asn Glu Val Val Glu Phe Arg Cys Gln Pro Gly Phe Val Met
260 265 270

Lys Gly Pro Arg Arg Val Lys Cys Gln Ala Leu Asn Lys Trp Glu Pro
275 280 285

Glu Leu Pro Ser Cys Ser Arg Val Cys Gln Pro Pro Pro Asp Val Leu

290		295		300											
His	Ala	Glu	Arg	Thr	Gln	Arg	Asp	Lys	Asp	Asn	Phe	Ser	Pro	Gly	Gln
305					310					315					320
Glu	Val	Phe	Tyr	Ser	Cys	Glu	Pro	Gly	Tyr	Asp	Leu	Arg	Gly	Ala	Ala
				325					330					335	
Ser	Met	Arg	Cys	Thr	Pro	Gln	Gly	Asp	Trp	Ser	Pro	Ala	Ala	Pro	Thr
			340					345					350		
Cys	Glu	Val	Lys	Ser	Cys	Asp	Asp	Phe	Met	Gly	Gln	Leu	Leu	Asn	Gly
		355					360					365			
Arg	Val	Leu	Phe	Pro	Val	Asn	Leu	Gln	Leu	Gly	Ala	Lys	Val	Asp	Phe
	370					375					380				
Val	Cys	Asp	Glu	Gly	Phe	Gln	Leu	Lys	Gly	Ser	Ser	Ala	Ser	Tyr	Cys
385					390					395					400
Val	Leu	Ala	Gly	Met	Glu	Ser	Leu	Trp	Asn	Ser	Ser	Val	Pro	Val	Cys
				405					410						415
Glu	Gln	Ile	Phe	Cys	Pro	Ser	Pro	Pro	Val	Ile	Pro	Asn	Gly	Arg	His
			420					425					430		
Thr	Gly	Lys	Pro	Leu	Glu	Val	Phe	Pro	Phe	Gly	Lys	Ala	Val	Asn	Tyr
		435					440					445			
Thr	Cys	Asp	Pro	His	Pro	Asp	Arg	Gly	Thr	Ser	Phe	Asp	Leu	Ile	Gly
	450					455					460				
Glu	Ser	Thr	Ile	Arg	Cys	Thr	Ser	Asp	Pro	Gln	Gly	Asn	Gly	Val	Trp
465					470					475					480
Ser	Ser	Pro	Ala	Pro	Arg	Cys	Gly	Ile	Leu	Gly	His	Cys	Gln	Ala	Pro
				485					490					495	
Asp	His	Phe	Leu	Phe	Ala	Lys	Leu	Lys	Thr	Gln	Thr	Asn	Ala	Ser	Asp
			500					505					510		
Phe	Pro	Ile	Gly	Thr	Ser	Leu	Lys	Tyr	Glu	Cys	Arg	Pro	Glu	Tyr	Tyr
		515					520					525			
Gly	Arg	Pro	Phe	Ser	Ile	Thr	Cys	Leu	Asp	Asn	Leu	Val	Trp	Ser	Ser
	530					535					540				

Pro Lys Asp Val Cys Lys Arg Lys Ser Cys Lys Thr Pro Pro Asp Pro
 545 550 555 560

Val Asn Gly Met Val His Val Ile Thr Asp Ile Gln Val Gly Ser Arg
 565 570 575

Ile Asn Tyr Ser Cys Thr Thr Gly His Arg Leu Ile Gly His Ser Ser
 580 585 590

Ala Glu Cys Ile Leu Ser Gly Asn Ala Ala His Trp Ser Thr Lys Pro
 595 600 605

Pro Ile Cys Gln Arg Ile Pro Cys Gly Leu Pro Pro Thr Ile Ala Asn
 610 615 620

Gly Asp Phe Ile Ser Thr Asn Arg Glu Asn Phe His Tyr Gly Ser Val
 625 630 635 640

Val Thr Tyr Arg Cys Asn Pro Gly Ser Gly Gly Arg Lys Val Phe Glu
 645 650 655

Leu Val Gly Glu Pro Ser Ile Tyr Cys Thr Ser Asn Asp Asp Gln Val
 660 665 670

Gly Ile Trp Ser Gly Pro Ala Pro Gln Cys Ile Ile Pro Asn Lys Cys
 675 680 685

Thr Pro Pro Asn Val Glu Asn Gly Ile Leu Val Ser Asp Asn Arg Ser
 690 695 700

Leu Phe Ser Leu Asn Glu Val Val Glu Phe Arg Cys Gln Pro Gly Phe
 705 710 715 720

Val Met Lys Gly Pro Arg Arg Val Lys Cys Gln Ala Leu Asn Lys Trp
 725 730 735

Glu Pro Glu Leu Pro Ser Cys Ser Arg Val Cys Gln Pro Pro Pro Asp
 740 745 750

Val Leu His Ala Glu Arg Thr Gln Arg Asp Lys Asp Asn Phe Ser Pro
 755 760 765

Gly Gln Glu Val Phe Tyr Ser Cys Glu Pro Gly Tyr Asp Leu Arg Gly
 770 775 780

Ala Ala Ser Met Arg Cys Thr Pro Gln Gly Asp Trp Ser Pro Ala Ala
 785 790 795 800

Pro Thr Cys Glu Val Lys Ser Cys Asp Asp Phe Met Gly Gln Leu Leu
805 810 815

Asn Gly Arg Val Leu Phe Pro Val Asn Leu Gln Leu Gly Ala Lys Val
820 825 830

Asp Phe Val Cys Asp Glu Gly Phe Gln Leu Lys Gly Ser Ser Ala Ser
835 840 845

Tyr Cys Val Leu Ala Gly Met Glu Ser Leu Trp Asn Ser Ser Val Pro
850 855 860

Val Cys Glu Gln Ile Phe Cys Pro Ser Pro Pro Val Ile Pro Asn Gly
865 870 875 880

Arg His Thr Gly Lys Pro Leu Glu Val Phe Pro Phe Gly Lys Ala Val
885 890 895

Asn Tyr Thr Cys Asp Pro His Pro Asp Arg Gly Thr Ser Phe Asp Leu
900 905 910

Ile Gly Glu Ser Thr Ile Arg Cys Thr Ser Asp Pro Gln Gly Asn Gly
915 920 925

Val Trp Ser Ser Pro Ala Pro Arg Cys Gly Ile Leu Gly His Cys Gln
930 935 940

Ala Pro Asp His Phe Leu Phe Ala Lys Leu Lys Thr Gln Thr Asn Ala
945 950 955 960

Ser Asp Phe Pro Ile Gly Thr Ser Leu Lys Tyr Glu Cys Arg Pro Glu
965 970 975

Tyr Tyr Gly Arg Pro Phe Ser Ile Thr Cys Leu Asp Asn Leu Val Trp
980 985 990

Ser Ser Pro Lys Asp Val Cys Lys Arg Lys Ser Cys Lys Thr Pro Pro
995 1000 1005

Asp Pro Val Asn Gly Met Val His Val Ile Thr Asp Ile Gln Val
1010 1015 1020

Gly Ser Arg Ile Asn Tyr Ser Cys Thr Thr Gly His Arg Leu Ile
1025 1030 1035

Gly His Ser Ser Ala Glu Cys Ile Leu Ser Gly Asn Thr Ala His
1040 1045 1050

Trp	Ser	Thr	Lys	Pro	Pro	Ile	Cys	Gln	Arg	Ile	Pro	Cys	Gly	Leu
1055						1060					1065			
Pro	Pro	Thr	Ile	Ala	Asn	Gly	Asp	Phe	Ile	Ser	Thr	Asn	Arg	Glu
1070						1075					1080			
Asn	Phe	His	Tyr	Gly	Ser	Val	Val	Thr	Tyr	Arg	Cys	Asn	Leu	Gly
1085						1090					1095			
Ser	Arg	Gly	Arg	Lys	Val	Phe	Glu	Leu	Val	Gly	Glu	Pro	Ser	Ile
1100						1105					1110			
Tyr	Cys	Thr	Ser	Asn	Asp	Asp	Gln	Val	Gly	Ile	Trp	Ser	Gly	Pro
1115						1120					1125			
Ala	Pro	Gln	Cys	Ile	Ile	Pro	Asn	Lys	Cys	Thr	Pro	Pro	Asn	Val
1130						1135					1140			
Glu	Asn	Gly	Ile	Leu	Val	Ser	Asp	Asn	Arg	Ser	Leu	Phe	Ser	Leu
1145						1150					1155			
Asn	Glu	Val	Val	Glu	Phe	Arg	Cys	Gln	Pro	Gly	Phe	Val	Met	Lys
1160						1165					1170			
Gly	Pro	Arg	Arg	Val	Lys	Cys	Gln	Ala	Leu	Asn	Lys	Trp	Glu	Pro
1175						1180					1185			
Glu	Leu	Pro	Ser	Cys	Ser	Arg	Val	Cys	Gln	Pro	Pro	Pro	Glu	Ile
1190						1195					1200			
Leu	His	Gly	Glu	His	Thr	Pro	Ser	His	Gln	Asp	Asn	Phe	Ser	Pro
1205						1210					1215			
Gly	Gln	Glu	Val	Phe	Tyr	Ser	Cys	Glu	Pro	Gly	Tyr	Asp	Leu	Arg
1220						1225					1230			
Gly	Ala	Ala	Ser	Leu	His	Cys	Thr	Pro	Gln	Gly	Asp	Trp	Ser	Pro
1235						1240					1245			
Glu	Ala	Pro	Arg	Cys	Ala	Val	Lys	Ser	Cys	Asp	Asp	Phe	Leu	Gly
1250						1255					1260			
Gln	Leu	Pro	His	Gly	Arg	Val	Leu	Phe	Pro	Leu	Asn	Leu	Gln	Leu
1265						1270					1275			
Gly	Ala	Lys	Val	Ser	Phe	Val	Cys	Asp	Glu	Gly	Phe	Arg	Leu	Lys

1280		1285		1290
Gly Ser Ser Val Ser His Cys Val Leu Val Gly Met Arg Ser Leu	1295	1300		1305
Trp Asn Asn Ser Val Pro Val Cys Glu His Ile Phe Cys Pro Asn	1310	1315		1320
Pro Pro Ala Ile Leu Asn Gly Arg His Thr Gly Thr Pro Ser Gly	1325	1330		1335
Asp Ile Pro Tyr Gly Lys Glu Ile Ser Tyr Thr Cys Asp Pro His	1340	1345		1350
Pro Asp Arg Gly Met Thr Phe Asn Leu Ile Gly Glu Ser Thr Ile	1355	1360		1365
Arg Cys Thr Ser Asp Pro His Gly Asn Gly Val Trp Ser Ser Pro	1370	1375		1380
Ala Pro Arg Cys Glu Leu Ser Val Arg Ala Gly His Cys Lys Thr	1385	1390		1395
Pro Glu Gln Phe Pro Phe Ala Ser Pro Thr Ile Pro Ile Asn Asp	1400	1405		1410
Phe Glu Phe Pro Val Gly Thr Ser Leu Asn Tyr Glu Cys Arg Pro	1415	1420		1425
Gly Tyr Phe Gly Lys Met Phe Ser Ile Ser Cys Leu Glu Asn Leu	1430	1435		1440
Val Trp Ser Ser Val Glu Asp Asn Cys Arg Arg Lys Ser Cys Gly	1445	1450		1455
Pro Pro Pro Glu Pro Phe Asn Gly Met Val His Ile Asn Thr Asp	1460	1465		1470
Thr Gln Phe Gly Ser Thr Val Asn Tyr Ser Cys Asn Glu Gly Phe	1475	1480		1485
Arg Leu Ile Gly Ser Pro Ser Thr Thr Cys Leu Val Ser Gly Asn	1490	1495		1500
Asn Val Thr Trp Asp Lys Lys Ala Pro Ile Cys Glu Ile Ile Ser	1505	1510		1515

Cys	Glu	Pro	Pro	Pro	Thr	Ile	Ser	Asn	Gly	Asp	Phe	Tyr	Ser	Asn
1520						1525					1530			
Asn	Arg	Thr	Ser	Phe	His	Asn	Gly	Thr	Val	Val	Thr	Tyr	Gln	Cys
1535						1540					1545			
His	Thr	Gly	Pro	Asp	Gly	Glu	Gln	Leu	Phe	Glu	Leu	Val	Gly	Glu
1550						1555					1560			
Arg	Ser	Ile	Tyr	Cys	Thr	Ser	Lys	Asp	Asp	Gln	Val	Gly	Val	Trp
1565						1570					1575			
Ser	Ser	Pro	Pro	Pro	Arg	Cys	Ile	Ser	Thr	Asn	Lys	Cys	Thr	Ala
1580						1585					1590			
Pro	Glu	Val	Glu	Asn	Ala	Ile	Arg	Val	Pro	Gly	Asn	Arg	Ser	Phe
1595						1600					1605			
Phe	Ser	Leu	Thr	Glu	Ile	Ile	Arg	Phe	Arg	Cys	Gln	Pro	Gly	Phe
1610						1615					1620			
Val	Met	Val	Gly	Ser	His	Thr	Val	Gln	Cys	Gln	Thr	Asn	Gly	Arg
1625						1630					1635			
Trp	Gly	Pro	Lys	Leu	Pro	His	Cys	Ser	Arg	Val	Cys	Gln	Pro	Pro
1640						1645					1650			
Pro	Glu	Ile	Leu	His	Gly	Glu	His	Thr	Leu	Ser	His	Gln	Asp	Asn
1655						1660					1665			
Phe	Ser	Pro	Gly	Gln	Glu	Val	Phe	Tyr	Ser	Cys	Glu	Pro	Ser	Tyr
1670						1675					1680			
Asp	Leu	Arg	Gly	Ala	Ala	Ser	Leu	His	Cys	Thr	Pro	Gln	Gly	Asp
1685						1690					1695			
Trp	Ser	Pro	Glu	Ala	Pro	Arg	Cys	Thr	Val	Lys	Ser	Cys	Asp	Asp
1700						1705					1710			
Phe	Leu	Gly	Gln	Leu	Pro	His	Gly	Arg	Val	Leu	Leu	Pro	Leu	Asn
1715						1720					1725			
Leu	Gln	Leu	Gly	Ala	Lys	Val	Ser	Phe	Val	Cys	Asp	Glu	Gly	Phe
1730						1735					1740			
Arg	Leu	Lys	Gly	Arg	Ser	Ala	Ser	His	Cys	Val	Leu	Ala	Gly	Met
1745						1750					1755			

Lys	Ala	Leu	Trp	Asn	Ser	Ser	Val	Pro	Val	Cys	Glu	Gln	Ile	Phe
1760						1765					1770			
Cys	Pro	Asn	Pro	Pro	Ala	Ile	Leu	Asn	Gly	Arg	His	Thr	Gly	Thr
1775						1780					1785			
Pro	Phe	Gly	Asp	Ile	Pro	Tyr	Gly	Lys	Glu	Ile	Ser	Tyr	Ala	Cys
1790						1795					1800			
Asp	Thr	His	Pro	Asp	Arg	Gly	Met	Thr	Phe	Asn	Leu	Ile	Gly	Glu
1805						1810					1815			
Ser	Ser	Ile	Arg	Cys	Thr	Ser	Asp	Pro	Gln	Gly	Asn	Gly	Val	Trp
1820						1825					1830			
Ser	Ser	Pro	Ala	Pro	Arg	Cys	Glu	Leu	Ser	Val	Pro	Ala	Ala	Cys
1835						1840					1845			
Pro	His	Pro	Pro	Lys	Ile	Gln	Asn	Gly	His	Tyr	Ile	Gly	Gly	His
1850						1855					1860			
Val	Ser	Leu	Tyr	Leu	Pro	Gly	Met	Thr	Ile	Ser	Tyr	Thr	Cys	Asp
1865						1870					1875			
Pro	Gly	Tyr	Leu	Leu	Val	Gly	Lys	Gly	Phe	Ile	Phe	Cys	Thr	Asp
1880						1885					1890			
Gln	Gly	Ile	Trp	Ser	Gln	Leu	Asp	His	Tyr	Cys	Lys	Glu	Val	Asn
1895						1900					1905			
Cys	Ser	Phe	Pro	Leu	Phe	Met	Asn	Gly	Ile	Ser	Lys	Glu	Leu	Glu
1910						1915					1920			
Met	Lys	Lys	Val	Tyr	His	Tyr	Gly	Asp	Tyr	Val	Thr	Leu	Lys	Cys
1925						1930					1935			
Glu	Asp	Gly	Tyr	Thr	Leu	Glu	Gly	Ser	Pro	Trp	Ser	Gln	Cys	Gln
1940						1945					1950			
Ala	Asp	Asp	Arg	Trp	Asp	Pro	Pro	Leu	Ala	Lys	Cys	Thr	Ser	Arg
1955						1960					1965			
Ala	His	Asp	Ala	Leu	Ile	Val	Gly	Thr	Leu	Ser	Gly	Thr	Ile	Phe
1970						1975					1980			
Phe	Ile	Leu	Leu	Ile	Ile	Phe	Leu	Ser	Trp	Ile	Ile	Leu	Lys	His
1985						1990					1995			

Arg Lys Gly Asn Asn Ala His Glu Asn Pro Lys Glu Val Ala Ile
2000 2005 2010

His Leu His Ser Gln Gly Gly Ser Ser Val His Pro Arg Thr Leu
2015 2020 2025

Gln Thr Asn Glu Glu Asn Ser Arg Val Leu Pro
2030 2035

<210> 52
<211> 1231
<212> PRT
<213> Homo sapiens

<400> 52

Met Arg Leu Leu Ala Lys Ile Ile Cys Leu Met Leu Trp Ala Ile Cys
1 5 10 15

Val Ala Glu Asp Cys Asn Glu Leu Pro Pro Arg Arg Asn Thr Glu Ile
20 25 30

Leu Thr Gly Ser Trp Ser Asp Gln Thr Tyr Pro Glu Gly Thr Gln Ala
35 40 45

Ile Tyr Lys Cys Arg Pro Gly Tyr Arg Ser Leu Gly Asn Val Ile Met
50 55 60

Val Cys Arg Lys Gly Glu Trp Val Ala Leu Asn Pro Leu Arg Lys Cys
65 70 75 80

Gln Lys Arg Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Thr Phe
85 90 95

Thr Leu Thr Gly Gly Asn Val Phe Glu Tyr Gly Val Lys Ala Val Tyr
100 105 110

Thr Cys Asn Glu Gly Tyr Gln Leu Leu Gly Glu Ile Asn Tyr Arg Glu
115 120 125

Cys Asp Thr Asp Gly Trp Thr Asn Asp Ile Pro Ile Cys Glu Val Val
130 135 140

Lys Cys Leu Pro Val Thr Ala Pro Glu Asn Gly Lys Ile Val Ser Ser
145 150 155 160

Ala Met Glu Pro Asp Arg Glu Tyr His Phe Gly Gln Ala Val Arg Phe
165 170 175

Val	Cys	Asn	Ser	Gly	Tyr	Lys	Ile	Glu	Gly	Asp	Glu	Glu	Met	His	Cys		
			180					185					190				
Ser	Asp	Asp	Gly	Phe	Trp	Ser	Lys	Glu	Lys	Pro	Lys	Cys	Val	Glu	Ile		
		195					200					205					
Ser	Cys	Lys	Ser	Pro	Asp	Val	Ile	Asn	Gly	Ser	Pro	Ile	Ser	Gln	Lys		
	210					215					220						
Ile	Ile	Tyr	Lys	Glu	Asn	Glu	Arg	Phe	Gln	Tyr	Lys	Cys	Asn	Met	Gly		
225					230					235					240		
Tyr	Glu	Tyr	Ser	Glu	Arg	Gly	Asp	Ala	Val	Cys	Thr	Glu	Ser	Gly	Trp		
				245					250					255			
Arg	Pro	Leu	Pro	Ser	Cys	Glu	Glu	Lys	Ser	Cys	Asp	Asn	Pro	Tyr	Ile		
			260					265					270				
Pro	Asn	Gly	Asp	Tyr	Ser	Pro	Leu	Arg	Ile	Lys	His	Arg	Thr	Gly	Asp		
		275					280					285					
Glu	Ile	Thr	Tyr	Gln	Cys	Arg	Asn	Gly	Phe	Tyr	Pro	Ala	Thr	Arg	Gly		
	290					295					300						
Asn	Thr	Ala	Lys	Cys	Thr	Ser	Thr	Gly	Trp	Ile	Pro	Ala	Pro	Arg	Cys		
305					310					315					320		
Thr	Leu	Lys	Pro	Cys	Asp	Tyr	Pro	Asp	Ile	Lys	His	Gly	Gly	Leu	Tyr		
				325					330					335			
His	Glu	Asn	Met	Arg	Arg	Pro	Tyr	Phe	Pro	Val	Ala	Val	Gly	Lys	Tyr		
			340					345					350				
Tyr	Ser	Tyr	Tyr	Cys	Asp	Glu	His	Phe	Glu	Thr	Pro	Ser	Gly	Ser	Tyr		
		355					360					365					
Trp	Asp	His	Ile	His	Cys	Thr	Gln	Asp	Gly	Trp	Ser	Pro	Ala	Val	Pro		
	370					375					380						
Cys	Leu	Arg	Lys	Cys	Tyr	Phe	Pro	Tyr	Leu	Glu	Asn	Gly	Tyr	Asn	Gln		
385					390					395					400		
Asn	Tyr	Gly	Arg	Lys	Phe	Val	Gln	Gly	Lys	Ser	Ile	Asp	Val	Ala	Cys		
				405					410					415			
His	Pro	Gly	Tyr	Ala	Leu	Pro	Lys	Ala	Gln	Thr	Thr	Val	Thr	Cys	Met		

420

425

430

Glu Asn Gly Trp Ser Pro Thr Pro Arg Cys Ile Arg Val Lys Thr Cys
 435 440 445

Ser Lys Ser Ser Ile Asp Ile Glu Asn Gly Phe Ile Ser Glu Ser Gln
 450 455 460

Tyr Thr Tyr Ala Leu Lys Glu Lys Ala Lys Tyr Gln Cys Lys Leu Gly
 465 470 475 480

Tyr Val Thr Ala Asp Gly Glu Thr Ser Gly Ser Ile Thr Cys Gly Lys
 485 490 495

Asp Gly Trp Ser Ala Gln Pro Thr Cys Ile Lys Ser Cys Asp Ile Pro
 500 505 510

Val Phe Met Asn Ala Arg Thr Lys Asn Asp Phe Thr Trp Phe Lys Leu
 515 520 525

Asn Asp Thr Leu Asp Tyr Glu Cys His Asp Gly Tyr Glu Ser Asn Thr
 530 535 540

Gly Ser Thr Thr Gly Ser Ile Val Cys Gly Tyr Asn Gly Trp Ser Asp
 545 550 555 560

Leu Pro Ile Cys Tyr Glu Arg Glu Cys Glu Leu Pro Lys Ile Asp Val
 565 570 575

His Leu Val Pro Asp Arg Lys Lys Asp Gln Tyr Lys Val Gly Glu Val
 580 585 590

Leu Lys Phe Ser Cys Lys Pro Gly Phe Phe Ile Val Gly Pro Asn Ser
 595 600 605

Val Gln Cys Tyr His Phe Gly Leu Ser Pro Asp Leu Pro Ile Cys Lys
 610 615 620

Glu Gln Val Gln Ser Cys Gly Pro Pro Pro Glu Leu Leu Asn Gly Asn
 625 630 635 640

Val Lys Glu Lys Thr Lys Glu Glu Tyr Gly His Ser Glu Val Val Glu
 645 650 655

Tyr Tyr Cys Asn Pro Arg Phe Leu Met Lys Gly Pro Asn Lys Ile Gln
 660 665 670

Cys Val Asp Gly Glu Trp Thr Thr Leu Pro Val Cys Ile Val Glu Glu
 675 680 685

Ser Thr Cys Gly Asp Ile Pro Glu Leu Glu His Gly Trp Ala Gln Leu
 690 695 700

Ser Ser Pro Pro Tyr Tyr Tyr Gly Asp Ser Val Glu Phe Asn Cys Ser
 705 710 715 720

Glu Ser Phe Thr Met Ile Gly His Arg Ser Ile Thr Cys Ile His Gly
 725 730 735

Val Trp Thr Gln Leu Pro Gln Cys Val Ala Ile Asp Lys Leu Lys Lys
 740 745 750

Cys Lys Ser Ser Asn Leu Ile Ile Leu Glu Glu His Leu Lys Asn Lys
 755 760 765

Lys Glu Phe Asp His Asn Ser Asn Ile Arg Tyr Arg Cys Arg Gly Lys
 770 775 780

Glu Gly Trp Ile His Thr Val Cys Ile Asn Gly Arg Trp Asp Pro Glu
 785 790 795 800

Val Asn Cys Ser Met Ala Gln Ile Gln Leu Cys Pro Pro Pro Pro Gln
 805 810 815

Ile Pro Asn Ser His Asn Met Thr Thr Thr Leu Asn Tyr Arg Asp Gly
 820 825 830

Glu Lys Val Ser Val Leu Cys Gln Glu Asn Tyr Leu Ile Gln Glu Gly
 835 840 845

Glu Glu Ile Thr Cys Lys Asp Gly Arg Trp Gln Ser Ile Pro Leu Cys
 850 855 860

Val Glu Lys Ile Pro Cys Ser Gln Pro Pro Gln Ile Glu His Gly Thr
 865 870 875 880

Ile Asn Ser Ser Arg Ser Ser Gln Glu Ser Tyr Ala His Gly Thr Lys
 885 890 895

Leu Ser Tyr Thr Cys Glu Gly Gly Phe Arg Ile Ser Glu Glu Asn Glu
 900 905 910

Thr Thr Cys Tyr Met Gly Lys Trp Ser Ser Pro Pro Gln Cys Glu Gly
 915 920 925

Leu Pro Cys Lys Ser Pro Pro Glu Ile Ser His Gly Val Val Ala His
 930 935 940

Met Ser Asp Ser Tyr Gln Tyr Gly Glu Glu Val Thr Tyr Lys Cys Phe
 945 950 955 960

Glu Gly Phe Gly Ile Asp Gly Pro Ala Ile Ala Lys Cys Leu Gly Glu
 965 970 975

Lys Trp Ser His Pro Pro Ser Cys Ile Lys Thr Asp Cys Leu Ser Leu
 980 985 990

Pro Ser Phe Glu Asn Ala Ile Pro Met Gly Glu Lys Lys Asp Val Tyr
 995 1000 1005

Lys Ala Gly Glu Gln Val Thr Tyr Thr Cys Ala Thr Tyr Tyr Lys
 1010 1015 1020

Met Asp Gly Ala Ser Asn Val Thr Cys Ile Asn Ser Arg Trp Thr
 1025 1030 1035

Gly Arg Pro Thr Cys Arg Asp Thr Ser Cys Val Asn Pro Pro Thr
 1040 1045 1050

Val Gln Asn Ala Tyr Ile Val Ser Arg Gln Met Ser Lys Tyr Pro
 1055 1060 1065

Ser Gly Glu Arg Val Arg Tyr Gln Cys Arg Ser Pro Tyr Glu Met
 1070 1075 1080

Phe Gly Asp Glu Glu Val Met Cys Leu Asn Gly Asn Trp Thr Glu
 1085 1090 1095

Pro Pro Gln Cys Lys Asp Ser Thr Gly Lys Cys Gly Pro Pro Pro
 1100 1105 1110

Pro Ile Asp Asn Gly Asp Ile Thr Ser Phe Pro Leu Ser Val Tyr
 1115 1120 1125

Ala Pro Ala Ser Ser Val Glu Tyr Gln Cys Gln Asn Leu Tyr Gln
 1130 1135 1140

Leu Glu Gly Asn Lys Arg Ile Thr Cys Arg Asn Gly Gln Trp Ser
 1145 1150 1155

Glu Pro Pro Lys Cys Leu His Pro Cys Val Ile Ser Arg Glu Ile
 1160 1165 1170

Met Glu Asn Tyr Asn Ile Ala Leu Arg Trp Thr Ala Lys Gln Lys
 1175 1180 1185

Leu Tyr Ser Arg Thr Gly Glu Ser Val Glu Phe Val Cys Lys Arg
 1190 1195 1200

Gly Tyr Arg Leu Ser Ser Arg Ser His Thr Leu Arg Thr Thr Cys
 1205 1210 1215

Trp Asp Gly Lys Leu Glu Tyr Pro Thr Cys Ala Lys Arg
 1220 1225 1230

<210> 53
 <211> 1234
 <212> PRT
 <213> Mus musculus

<400> 53

Met Arg Leu Ser Ala Arg Ile Ile Trp Leu Ile Leu Trp Thr Val Cys
 1 5 10 15

Ala Ala Glu Asp Cys Lys Gly Pro Pro Pro Arg Glu Asn Ser Glu Ile
 20 25 30

Leu Ser Gly Ser Trp Ser Glu Gln Leu Tyr Pro Glu Gly Thr Gln Ala
 35 40 45

Thr Tyr Lys Cys Arg Pro Gly Tyr Arg Thr Leu Gly Thr Ile Val Lys
 50 55 60

Val Cys Lys Asn Gly Lys Trp Val Ala Ser Asn Pro Ser Arg Ile Cys
 65 70 75 80

Arg Lys Lys Pro Cys Gly His Pro Gly Asp Thr Pro Phe Gly Ser Phe
 85 90 95

Arg Leu Ala Val Gly Ser Gln Phe Glu Phe Gly Ala Lys Val Val Tyr
 100 105 110

Thr Cys Asp Asp Gly Tyr Gln Leu Leu Gly Glu Ile Asp Tyr Arg Glu
 115 120 125

Cys Gly Ala Asp Gly Trp Ile Asn Asp Ile Pro Leu Cys Glu Val Val
 130 135 140

Lys Cys Leu Pro Val Thr Glu Leu Glu Asn Gly Arg Ile Val Ser Gly
 145 150 155 160

Ala Ala Glu Thr Asp Gln Glu Tyr Tyr Phe Gly Gln Val Val Arg Phe
 165 170 175

Glu Cys Asn Ser Gly Phe Lys Ile Glu Gly His Lys Glu Ile His Cys
 180 185 190

Ser Glu Asn Gly Leu Trp Ser Asn Glu Lys Pro Arg Cys Val Glu Ile
 195 200 205

Leu Cys Thr Pro Pro Arg Val Glu Asn Gly Asp Gly Ile Asn Val Lys
 210 215 220

Pro Val Tyr Lys Glu Asn Glu Arg Tyr His Tyr Lys Cys Lys His Gly
 225 230 235 240

Tyr Val Pro Lys Glu Arg Gly Asp Ala Val Cys Thr Gly Ser Gly Trp
 245 250 255

Ser Ser Gln Pro Phe Cys Glu Glu Lys Arg Cys Ser Pro Pro Tyr Ile
 260 265 270

Leu Asn Gly Ile Tyr Thr Pro His Arg Ile Ile His Arg Ser Asp Asp
 275 280 285

Glu Ile Arg Tyr Glu Cys Asn Tyr Gly Phe Tyr Pro Val Thr Gly Ser
 290 295 300

Thr Val Ser Lys Cys Thr Pro Thr Gly Trp Ile Pro Val Pro Arg Cys
 305 310 315 320

Thr Leu Lys Pro Cys Glu Phe Pro Gln Phe Lys Tyr Gly Arg Leu Tyr
 325 330 335

Tyr Glu Glu Ser Leu Arg Pro Asn Phe Pro Val Ser Ile Gly Asn Lys
 340 345 350

Tyr Ser Tyr Lys Cys Asp Asn Gly Phe Ser Pro Pro Ser Gly Tyr Ser
 355 360 365

Trp Asp Tyr Leu Arg Cys Thr Ala Gln Gly Trp Glu Pro Glu Val Pro
 370 375 380

Cys Val Arg Lys Cys Val Phe His Tyr Val Glu Asn Gly Asp Ser Ala
 385 390 395 400

Tyr Trp Glu Lys Val Tyr Val Gln Gly Gln Ser Leu Lys Val Gln Cys

Asn Gly Ala Lys Lys Val Glu Tyr Ser His Gly Glu Val Val Lys Tyr
645 650 655

Asp	Cys	Lys	Pro	Arg	Phe	Leu	Leu	Lys	Gly	Pro	Asn	Lys	Ile	Gln	Cys		
			660					665					670				
Val	Asp	Gly	Asn	Trp	Thr	Thr	Leu	Pro	Val	Cys	Ile	Glu	Glu	Glu	Arg		
		675					680					685					
Thr	Cys	Gly	Asp	Ile	Pro	Glu	Leu	Glu	His	Gly	Ser	Ala	Lys	Cys	Ser		
	690					695					700						
Val	Pro	Pro	Tyr	His	His	Gly	Asp	Ser	Val	Glu	Phe	Ile	Cys	Glu	Glu		
705					710					715					720		
Asn	Phe	Phe	Met	Ile	Gly	His	Gly	Ser	Val	Ser	Cys	Ile	Ser	Gly	Lys		
			725						730					735			
Trp	Thr	Gln	Leu	Pro	Lys	Cys	Val	Ala	Thr	Asp	Gln	Leu	Glu	Lys	Cys		
			740					745					750				
Arg	Val	Leu	Lys	Ser	Thr	Gly	Ile	Glu	Ala	Ile	Lys	Pro	Lys	Leu	Thr		
		755					760					765					
Glu	Phe	Phe	His	Asn	Ser	Thr	Met	Asp	Tyr	Lys	Cys	Arg	Asp	Lys	Gln		
	770					775					780						
Glu	Tyr	Glu	Arg	Ser	Ile	Cys	Ile	Asn	Gly	Lys	Trp	Asp	Pro	Glu	Pro		
785					790					795					800		
Asn	Cys	Thr	Ser	Lys	Thr	Ser	Cys	Pro	Pro	Pro	Pro	Gln	Ile	Pro	Asn		
				805					810					815			
Thr	Gln	Val	Ile	Glu	Thr	Thr	Val	Lys	Tyr	Leu	Asp	Gly	Glu	Lys	Leu		
			820					825					830				
Ser	Val	Leu	Cys	Gln	Asp	Asn	Tyr	Leu	Thr	Gln	Asp	Ser	Glu	Glu	Met		
		835					840					845					
Val	Cys	Lys	Asp	Gly	Arg	Trp	Gln	Ser	Leu	Pro	Arg	Cys	Ile	Glu	Lys		
	850					855					860						
Ile	Pro	Cys	Ser	Gln	Pro	Pro	Thr	Ile	Glu	His	Gly	Ser	Ile	Asn	Leu		
865					870					875					880		
Pro	Arg	Ser	Ser	Glu	Glu	Arg	Arg	Asp	Ser	Ile	Glu	Ser	Ser	Ser	His		
				885					890					895			
Glu	His	Gly	Thr	Thr	Phe	Ser	Tyr	Val	Cys	Asp	Asp	Gly	Phe	Arg	Ile		
			900					905					910				

Pro Glu Glu Asn Arg Ile Thr Cys Tyr Met Gly Lys Trp Ser Thr Pro
 915 920 925

Pro Arg Cys Val Gly Leu Pro Cys Gly Pro Pro Pro Ser Ile Pro Leu
 930 935 940

Gly Thr Val Ser Leu Glu Leu Glu Ser Tyr Gln His Gly Glu Glu Val
 945 950 955 960

Thr Tyr His Cys Ser Thr Gly Phe Gly Ile Asp Gly Pro Ala Phe Ile
 965 970 975

Ile Cys Glu Gly Gly Lys Trp Ser Asp Pro Pro Lys Cys Ile Lys Thr
 980 985 990

Asp Cys Asp Val Leu Pro Thr Val Lys Asn Ala Ile Ile Arg Gly Lys
 995 1000 1005

Ser Lys Lys Ser Tyr Arg Thr Gly Glu Gln Val Thr Phe Arg Cys
 1010 1015 1020

Gln Ser Pro Tyr Gln Met Asn Gly Ser Asp Thr Val Thr Cys Val
 1025 1030 1035

Asn Ser Arg Trp Ile Gly Gln Pro Val Cys Lys Asp Asn Ser Cys
 1040 1045 1050

Val Asp Pro Pro His Val Pro Asn Ala Thr Ile Val Thr Arg Thr
 1055 1060 1065

Lys Asn Lys Tyr Leu His Gly Asp Arg Val Arg Tyr Glu Cys Asn
 1070 1075 1080

Lys Pro Leu Glu Leu Phe Gly Gln Val Glu Val Met Cys Glu Asn
 1085 1090 1095

Gly Ile Trp Thr Glu Lys Pro Lys Cys Arg Asp Ser Thr Gly Lys
 1100 1105 1110

Cys Gly Pro Pro Pro Pro Ile Asp Asn Gly Asp Ile Thr Ser Leu
 1115 1120 1125

Ser Leu Pro Val Tyr Glu Pro Leu Ser Ser Val Glu Tyr Gln Cys
 1130 1135 1140

Gln Lys Tyr Tyr Leu Leu Lys Gly Lys Lys Thr Ile Thr Cys Thr
 1145 1150 1155

Asn Gly Lys Trp Ser Glu Pro Pro Thr Cys Leu His Ala Cys Val
1160 1165 1170

Ile Pro Glu Asn Ile Met Glu Ser His Asn Ile Ile Leu Lys Trp
1175 1180 1185

Arg His Thr Glu Lys Ile Tyr Ser His Ser Gly Glu Asp Ile Glu
1190 1195 1200

Phe Gly Cys Lys Tyr Gly Tyr Tyr Lys Ala Arg Asp Ser Pro Pro
1205 1210 1215

Phe Arg Thr Lys Cys Ile Asn Gly Thr Ile Asn Tyr Pro Thr Cys
1220 1225 1230

Val

<210> 54
<211> 21
<212> PRT
<213> Homo sapiens

<400> 54

Met Pro Met Gly Ser Leu Gln Pro Leu Ala Thr Leu Tyr Leu Leu Gly
1 5 10 15

Met Leu Val Ala Ser
20

<210> 55
<211> 17
<212> PRT
<213> Homo sapiens

<400> 55

Met Gly Ala Ala Gly Leu Leu Gly Val Phe Leu Ala Leu Val Ala Pro
1 5 10 15

Gly

<210> 56
<211> 20
<212> PRT
<213> Homo sapiens

<400> 56

Met Gly Ala Ala Gly Leu Leu Gly Val Phe Leu Ala Leu Val Ala Pro
 1 5 10 15

Gly Val Leu Gly
 20

<210> 57
 <211> 927
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 57
 gacacgtgat cagccgccac catgcccatg ggggtctctgc aaccgctggc caccttgtag 60
 ctgctgggga tgctggtcgc ttccgtgcta ggcacatcatc atcatcatca tgtgaaactg 120
 caggagtcag gggctgagct tgtgaagcct ggggcttcag tgaagctgtc ctgcaaggct 180
 tctggctaca ccttcaccag ctactggatg cactgggtga agcagaggcc tggacgaggc 240
 cttgagtgga ttggaaggat tggctcctaag agtgggtgga ctaagtacaa tgagaagttc 300
 aagagcaagg ccacactgac tgtagacaaa ccctccagca cagcctacat gcagctcagc 360
 agcctgacat ctgaggactc tgcggtctat tattgtgcaa gaagaatggg aaaggggtgc 420
 tatggactac tggggccaag ggaccacggt caccgtctcc tcaaagggcg aattccagca 480
 cactggcggc cggtactagt ggatccgagc tcggtaccaa gcttggcgtc aggaggcggg 540
 ggcggctcgg gtggcggcgg ctcttgata tctgcagaat tcgcccttga cattgagctc 600
 acccagtctc caaccaccat ggctgcatct cccggggaga agatcactat cacctgcagt 660
 gccagctcaa gtataagttc caattacttg cattgggtatc agcagaagcc aggattctcc 720
 cctaaactct tgatttatag gacatccaat ctggcttctg gaggcccagc tcgcttcagt 780
 ggcagtgggt ctgggacctc ttactctctc acaattggca ccatggaggc tgaagatgtt 840
 gccacttact actgccagca gggtagtagt ataccacgta cacgttcgga gggggcacca 900
 agctggaaat aatagactag tcgtgcg 927

<210> 58
 <211> 954
 <212> DNA
 <213> Artificial sequence

<220>
 <223> Synthetic polynucleotide

<400> 58
 gacacgaagc ttgccgccac catgcccatg ggggtctctgc aaccgctggc caccttgtag 60
 ctgctgggga tgctggtcgc ttccgtgcta ggcacatcatc atcatcatca tgtcaagctg 120

caggagtctg ggactgaact ggtgaagcct ggggcttcag tgaagctgtc ctgcaaggct	180
tctggctaca ccttcaccag ctactggatg cactgggtga agcagaggcc tggacaaggc	240
cttgagtgga ttggaaatat taatcctagc aatggtggta ctaactacaa tgagaagttc	300
aagagcaagg ccacactgac tgtagacaaa tcctccagca cagcctacat gcagctcagc	360
agcctgacat ctgaggactc tgcggtctat tattgtgcaa gaagaggcat acggttacga	420
cactttgact actggggcca agggaccacg gtcaccgtct cctcaagggc gaattctgca	480
gatatccatc aactggcgg ccgctcgagc atgcatctag agggcccaat tcgccctata	540
gtgagtcgta tatcaggagg cgggtggcggc tcgggtggcg gcggctcttg gatattctgca	600
gaattcgccc ttgacattgt gatgacacag tctcctgctt ccttagctgt atctctgggg	660
cagagggcca ccatctcata cagggccagc aaaagtgtca gtacatctgg ctatagttat	720
atgcactgga accaacagaa accaggacag ccaccagac tcctcatcta tcttgtatcc	780
aacctagaat ctgggggtccc tgccaggttc agtggcagtg ggtctgggac agacttcacc	840
ctcaacatcc atcctgtgga ggaggaggat gctgcaacct attactgtca gcacattagg	900
gagcttacac gttcggaggg gggaccaagc tggaaataat agcccgggcg tgcg	954

摘要

本發明提供了用於治療個體體內的炎性疾病和/或檢測體內組織損傷的靶向遞送方法和構建物。靶向遞送方法利用識別發現存在炎症部位的表位的抗體。本發明還提供了在個體的眼睛中抑制補體驅動炎症的方法，包括給個體施用抗體或其片段或它們的組合物，其中該抗體或其片段特異性結合到膜聯蛋白 IV 或磷脂。還提供了治療補體相關眼疾病或涉及氧化性損傷的眼疾病的相關方法。此外，本發明提供了在個體的眼組織中檢測補體介導損傷的方法，包括向個體施用構建體或其組合物，其中該構建體包括（a）抗體或其片段特異性結合到膜聯蛋白 IV 或磷脂；（b）可檢測的部分。