

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2013237330 C1**

(54) Title
Chemical process for preparing spiroindolones and intermediates thereof

(51) International Patent Classification(s)
C07D 405/06 (2006.01) **C07D 209/34** (2006.01)
C07D 209/12 (2006.01) **C07D 471/20** (2006.01)
C07D 209/14 (2006.01) **C12P 17/10** (2006.01)

(21) Application No: **2013237330** (22) Date of Filing: **2013.03.22**

(87) WIPO No: **WO13/139987**

(30) Priority Data

(31) Number	(32) Date	(33) Country
PCT/CN2012/000359	2012.03.23	CN

(43) Publication Date: **2013.09.26**

(44) Accepted Journal Date: **2015.12.03**

(44) Amended Journal Date: **2016.08.25**

(71) Applicant(s)
Novartis AG

(72) Inventor(s)
Crowe, Michael;Foulkes, Michael;Francese, Giancarlo;Grimler, Dominique;Kuesters, Ernst;Laumen, Kurt;Li, Yunzhong;Lin, Changxue;Nazor, Jovana;Ruch, Thomas;Smith, Derek;Song, Shiwei;Teng, Shangjun

(74) Agent / Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000

(56) Related Art
WO 2009/132921 A1
WO 2012/007548 A1
EP 0 120 208 A2
ASANO Y ET AL:"HIGH YIELD SYNTHESIS OF L-AMINO ACIDS
BY PHENYLALANINE DEHYDROGENASE FROM SPOROSARCINA
UREAE" ,AGRICULTURAL AND BIOLOGICAL CHEMISTRY, JAPAN SOC. FOR
BIOSCIENCE, BIOTECHNOLOGY AND AGROCHEM, vol . 51, no. 7, 1 Jan 1987
WO 2010/081053 A2
AT 507 050 A1
US 4 073 795 A
EP 0 036 741 A2
EP 0 135 846 A2
EP 0 135 846 A2
GB 974 895 A
US 3 282 959 A
TABUSHI I ET AL: "A-(modified B6)-B-[.omega.-amino(ethylamino)]-.beta.-
cyclodextrin as an artificial B6 enzyme for chiral aminotransfer reaction"
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol . 107, no. 19, 1
September 1985, pages 5545-5546
NICHOLS D E ET AL JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN
CHEMICAL SOCIETY, US, vol . 31, no. 7, 1 January 1988, pages 1406-1412



(43) International Publication Date
26 September 2013 (26.09.2013)

WIPO | PCT

(10) International Publication Number
WO 2013/139987 A1

(51) International Patent Classification:

C07D 405/06 (2006.01) **C07D 209/14** (2006.01)
C07D 471/20 (2006.01) **C07D 209/34** (2006.01)
C07D 209/12 (2006.01) **C12P 17/10** (2006.01)

(21) International Application Number:

PCT/EP2013/056170

(22) International Filing Date:

22 March 2013 (22.03.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PCT/CN2012/000359 23 March 2012 (23.03.2012) CN

(71) Applicant (for all designated States except US): **NOVARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(71) Applicants (for US only): **CROWE, Michael** [GB/SG]; 20 Upper Circular Road, #13-03, The Riverwalk, Singapore 058416 (SG). **FOULKES, Michael** [GB/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **FRANCESE, Giancarlo** [IT/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **GRIMLER, Dominique** [FR/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **KUESTERS, Ernst** [DE/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **LAUMEN, Kurt** [DE/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **LI, Yunzhong** [CN/CN]; Suzhou Novartis Pharma Technology Co., Ltd., #18 Tonglian Road, Riverside Industrial Park, Changshu Economic Development Zone, Changshu, Jiangsu 215537 (CN). **LIN, Changxue** [CN/CN]; Suzhou Novartis Pharma Technology Co., Ltd., #18 Tonglian Road, Riverside Industrial Park, Changshu Economic Development Zone, Changshu, Jiangsu 215537 (CN). **NAZOR, Jovana** [HR/US]; Codexis, Inc., 200 Penobscot Drive, Redwood City, California 94063 (US).

RUCH, Thomas [DE/CH]; Novartis Pharma AG, Postfach, CH-4002 Basel (CH). **SMITH, Derek** [GB/SG]; 12 Holland Avenue, #10-31, Singapore 272012 (SG). **SONG, Shiwei** [CN/SG]; Blk 60 Teban Gardens, #11-462, Singapore 600060 (SG). **TENG, Shangjun** [CN/CN]; Suzhou Novartis Pharma Technology Co., Ltd., #18 Tonglian Road, Riverside Industrial Park, Changshu Economic Development Zone, Changshu, Jiangsu 215537 (CN).

(74) Agent: **STUTTLE, James**; Novartis Pharma AG, Patent Department, CH-4002 Basel (CH).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: CHEMICAL PROCESS FOR PREPARING SPIROINDOLONES AND INTERMEDIATES THEREOF

(57) Abstract: The present invention relates to processes and intermediates useful for the manufacture of spiroindolone compounds such as (1'R,3'S)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3, 4-b]indol]-2-one and salts and hydrates and solvates thereof.



WO 2013/139987 A1


CHEMICAL PROCESS FOR PREPARING SPIROINDOLONES AND INTERMEDIATES THEREOF

REFERENCE TO SEQUENCE LISTING, TABLE OR COMPUTER PROGRAM

The official copy of the Sequence Listing is submitted concurrently with the specification as an ASCII formatted text file via EFS-Web, with a file name of "PAT055051_seq|2.txt", a creation date of March 22, 2013, and a size of 447 kilobytes. The Sequence Listing filed via EFS-Web is part of the specification and is incorporated in its entirety by reference herein.

Prior art:

(1'*R*, 3'*S*)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-*b*]indol]-2-one (eg. a compound of formula (IV), which comprises a spiroindolone moiety) and a 6-steps synthetic method for preparing, including known chiral amine intermediate compound (IIA) are known (WO 2009/132921):




Invention:


The present invention is directed to an improved method of synthesizing spiroindolone compounds, in particular, (1'*R*, 3'*S*)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-*b*]indol]-2-one, and intermediates used in the improved method.

2013237330 27 Oct 2015

In one aspect of the present invention, there is provided a process for preparing a compound of formula (II), or a salt or solvate or hydrate thereof,



comprising enzymatically converting a compound of formula (I) to compound of formula (II), or a salt, solvate or hydrate thereof,




wherein:

the dashed line is absent;

A is C=O

2013237330 27 Oct 2015



R^4 and R^7 are each, independently, H or $-Cl$;


R^5 is H, $-OH$, $-CH_3$, $-OCH_3$, $-F$, $-Cl$, $-CF_3$ or $-CN$;

R^6 is H, $-OH$, $-OCH_3$, $-F$ or $-Cl$;

R^8 is H, $-CH_3$, $-CH_2CH_3$, $-CH_2OH$, $-CO_2H$, $-CO_2CH_3$, $-CO_2CH_2CH_3$ or $-CF_3$; and

n is 1 or 2.


In another aspect of the present invention, there is provided a process for preparing a compound of formula (IV), or a salt or hydrate or solvate thereof,



comprising,


reacting a compound of formula (III), or a salt or hydrate or solvate thereof,

2013237330 27 Oct 2015




(III)

with a compound of formula (II), or a salt or hydrate or solvate thereof,



(II)

wherein the compound of formula (II), or a salt or hydrate or solvate thereof, is prepared by enzymatically converting a compound of formula (I), or a salt or hydrate or solvate thereof,



(I)

wherein:

the dashed line is absent;





A is C=O;

R^{1'} is hydrogen or (C₁-C₆)alkyl;


2013237330 27 Oct 2015

$R^{4'}$, $R^{5'}$, $R^{6'}$ and $R^{7'}$, are each, independently hydrogen, halo, hydroxy, amino, alkylamino, dialkylamino, (C_1-C_6) alkyl, and (C_1-C_6) alkyloxy;

R^1 is H, $-CH_3$,



or




;

R^4 and R^7 are each, independently, H or $-Cl$; R^5 is H, $-OH$, $-CH_3$, $-OCH_3$, $-F$, $-Cl$, $-CF_3$ or $-CN$; R^6 is H, $-OH$, $-OCH_3$, $-F$ or $-Cl$;

R^8 is H, $-CH_3$, $-CH_2CH_3$, $-CH_2OH$, $-CO_2H$, $-CO_2CH_3$, $-CO_2CH_2CH_3$ or $-CF_3$;

and n is 1 or 2.

In yet another aspect of the present invention, there is provided a process for preparing a compound of formula (IVA), or a salt or solvate or hydrate thereof,




(IVA)

comprising


reacting a compound of formula (IIIA) or a salt or solvate or hydrate thereof,

2013237330 27 Oct 2015



(IIIA)


with a compound of formula (IIA) or a salt or solvate or hydrate thereof,



(IIA)


to provide the compound of formula (IVA), or a salt or solvate or hydrate thereof,

wherein the compound of formula (IIA) is prepared by enzymatically converting a compound of formula (IA), or a salt or hydrate or solvate thereof,



(IA).


In a first embodiment, the invention is a process for preparing a compound of formula (II), or a salt or solvate or hydrate thereof,



(II)


2013237330 27 Oct 2015

comprising converting a compound of formula (I) to compound of formula (II), or a salt, solvate or hydrate thereof,




(I)

wherein: the dashed line is a bond or absent; A is selected from C=O and C=NH; or when the dashed line is a double bond A-R⁸ is:




R^4 and R^7 are each, independently, H or -Cl; R^5 is H, -OH, -CH₃, -OCH₃, -F, -Cl, -CF₃ or -CN; R^6 is H, -OH, -OCH₃, -F or -Cl; R^8 is H, -CH₃, -CH₂CH₃, -CH₂OH, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃ or -CF₃; and n is 1 or 2.

In a second embodiment, the invention is a process for preparing a compound of formula (IIA) or a salt or hydrate or solvate thereof,



(IIA)


comprising enzymatically transaminating a compound of formula (IA) or a salt or solvate or hydrate thereof,



(IA).


to provide the compound of formula (IIA).

In a third embodiment, the invention is a process for preparing a compound of formula (IV), or a salt or hydrate or solvate thereof,




(IV)

comprising, reacting a compound of formula (III), or a salt or hydrate or solvate thereof,




(III)

with a compound of formula (II), or a salt or hydrate or solvate thereof,




(II)

wherein the compound of formula (II), or a salt or hydrate or solvate thereof, is prepared from a compound of formula (I), or a salt or hydrate or solvate thereof,




(I)

wherein: the dashed line is a bond or absent; A is selected from C=O and C=NH; or when the dashed line is a double bond A-R⁸ is:




; R^a is C₁₋₆ alkyl; R¹ is (C₁₋₆) alkyl, optionally substituted with an amino, (C₁₋₆) alkyl amino, (C₁₋₆) alkyl di-alkyl amino or (C₁₋₆) alkyl C(O)NH (C₁₋₆) alkyl; R⁴ and R⁷ are each, independently, H or halo; R⁵ and R⁶ are each, independently hydrogen, halo, hydroxyl, (C₁₋₆) alkyl, trihalo (C₁) alkyl, cyano or (C₁₋₆) alkoxy; R⁸ is (C₁₋₆) alkyl, optionally substituted with a hydroxyl; n is 1 or 2; R^{1'} is hydrogen or (C₁₋₆)alkyl; and R^{4'}, R^{5'}, R^{6'} and R^{7'}, are each, independently hydrogen, halo, hydroxy, amino, alkylamino, dialkylamino, (C₁₋₆)alkyl, and (C₁₋₆)alkyloxy.

In a fourth embodiment, the invention is a process for preparing a compound of formula (IVA), or a salt or solvate or hydrate thereof,




(IVA)

comprising: reacting a compound of formula (IIIA) or a salt or solvate or hydrate thereof,



(IIIA)


with a compound of formula (II) or a salt or solvate or hydrate thereof,




(IIA)

to provide the compound of formula (IVA), or a salt or solvate or hydrate thereof,

wherein the compound of formula (IIA) is prepared from the compound of formula (IA), or a salt or hydrate or solvate thereof,



In a fifth embodiment, the invention is a compound of formula (IC):





(IC)

wherein: R^1 is (C_1-C_6) alkyl, optionally substituted with an amino, (C_1-C_6) alkyl amino, (C_1-C_6) alkyl di-alkyl amino or (C_1-C_6) alkyl $C(O)NH$ (C_1-C_6) alkyl; R^5 and R^6 are each, independently hydrogen, halo, hydroxyl, (C_1-C_6) alkyl, trihalo (C_1) alkyl, cyano or (C_1-C_6)

alkoxy; R^8 is (C_1-C_6) alkyl, optionally substituted with a hydroxyl; n is 1 or 2; or a pharmaceutically acceptable salt or hydrate or solvate thereof.

In a sixth embodiment the invention is a compound selected from:



hydrate thereof.

Field of the invention


The invention relates to novel processes a novel process step and a novel intermediate useful for the preparation of spiroindolone compounds useful for the treatment of parasitic diseases comprising e.g. a spiroindolone moiety, such as (1'*R*,3'*S*)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-*b*]indol]-2-one.

Background of the invention

The present invention relates to processes for the preparation of spiroindolone compounds, such as (1'*R*,3'*S*)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-*b*]indol]-2-one.


(1'*R*,3'*S*)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-*b*]indol]-2-one is useful in the treatment and/or prevention of infections such as those caused by *Plasmodium falciparum*, *Plasmodium vivax*, *Plasmodium malariae*,

Plasmodium ovale, *Trypanosoma cruzi* and parasites of the *Leishmania* genus such as, for example, *Leishmania donovani*., and it has the following structure:




(1'R,3'S)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-b]indol]-2-one and a synthesis thereof are described in WO 2009/132921 A1 in particular in Example 49 therein.

There is a need to provide new process for the preparation of (1'R,3'S)-5,7'-dichloro-6'-fluoro-3'-methyl-2',3',4',9'-tetrahydrospiro[indoline-3,1'-pyrido[3,4-b]indol]-2-one in order to improve the overall efficiency of the synthesis to make it suitable for manufacturing. In particular, there is a need to increase the efficiency of synthesizing the chiral amine intermediate (IIA):



Detailed Description of the Invention


The process(es), according to the present invention, for producing spiroindolone compounds, such as compounds according to formula (IV), or salt or hydrate or solvate thereof, and intermediates, as defined herein, are summarized in Scheme 1.




Scheme 1

Namely, a compound of formula (I), or salt or hydrate or solvate thereof, is converted into a compound of formula (II), or salt or hydrate or solvate thereof, according to methods 1, 2, 3, 4, 5 or 6 wherein

- method 1 comprises
 - a) Enzymatic transamination to convert a compound of formula (I), or salt or hydrate or solvate thereof, into a compound of formula (II), or salt or hydrate or solvate thereof




- method 2 comprises;
 - a) Chemical asymmetric catalysis to convert a compound of formula (I), or salt or hydrate or solvate thereof, into a compound of formula (II), or salt thereof;




- method 3 comprises

a) Chemical asymmetric reduction to convert a compound of formula (I), or salt or hydrate or solvate thereof, into a compound of formula (II), or salt or hydrate or solvate thereof;




- method 4 comprises

a) Reduction followed by chiral resolution to convert a compound of formula (I), or salt or hydrate or solvate thereof, into a compound of formula (II), or salt or hydrate or solvate thereof;



- method 5 comprises

- a) Lipase resolution to convert a racemate of a compound of formula (II), or salt or hydrate or solvate thereof, into a single enantiomer compound of formula (II), or salt or hydrate or solvate thereof;



Novozym 435: *Candida antarctica* Lipase B immobilized on an acrylic resin

- method 6 comprises

- a) A combination of two or more of methods 1-5 to convert a compound of formula (I), or salt or hydrate or solvate thereof, into a compound of formula (II), or salt or hydrate or solvate thereof.

A compound of formula (II), or salt thereof, may be converted into a compound formula (IV), or salt thereof, for example, as described in WO 2009/132921 in particular as described in the relevant claims and examples, which are incorporated by reference herein.

The invention specially relates to the processes described in each section. The invention likewise relates, independently, to every single step described in a process sequence within the corresponding section. Therefore, each and every single step of any process, consisting of a sequence of steps, described herein is itself a preferred embodiment of the present invention. Thus, the invention also relates to those embodiments of the process, according to which a compound obtainable as an intermediate in any step of the process is used as a starting material.


The invention likewise relates to novel starting materials which have been specifically developed for the preparation of the compounds according to the invention, to their use and to processes for their preparation.

The invention also relates to intermediates which have been specifically developed for the preparation of the compounds according to the invention, to their use and to processes for their preparation.


It is noted that in the present application usually explanations made in one section are also applicable for other sections, unless otherwise stated. For example, the explanations for the residue R^1 in formula (I) given in section A also apply if formula (I) occurs in other sections, such as Section B, unless otherwise stated.

Section A: Preparation of a compound of formula (I)

A compound of formula (I), or salt thereof, or hydrate or solvate thereof, may be prepared as described below and/or according to Examples 1-3 contained herein.




or




Section B: Conversion of a compound of formula (I) into a compound of formula (II).


In a first embodiment, the invention is a process for preparing a compound of formula (II), or a salt or solvate or hydrate thereof,



comprising converting a compound of formula (I) to compound of formula (II), or a salt, solvate or hydrate thereof,



wherein: the dashed line is a bond or absent; A is selected from C=O and C=NH; or when the dashed line is a double bond A-R⁸ is:



R⁴ and R⁷ are each, independently, H or -Cl; R⁵ is H, -OH, -CH₃, -OCH₃, -F, -Cl, -CF₃ or -CN; R⁶ is H, -OH, -OCH₃, -F or -Cl; R⁸ is H, -CH₃, -CH₂CH₃, -CH₂OH, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃ or -CF₃; and n is 1 or 2.


In a first alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, R⁵ and R⁶ are fluoro when: R⁸ is -CH₃, and n is 1.

In a second alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, R^5 and R^6 are fluoro and chloro, when: R^8 is $-CH_3$, and n is 1.

In a third alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, R⁵ and R⁶ are hydrogen when: R⁸ is -CH₃ and n is 1.

In a fourth alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, R⁵ is fluoro when: n is 1, and R⁶ is hydrogen.

In a sixth alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, the compound of formula (II) is of formula (IIA), or a salt or solvate or hydrate thereof,




(IIA).

In a seventh alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, the compound of formula (II) is converted from the compound of formula (I) under a condition selected from enzymatic transamination, chemical asymmetric catalysis, asymmetric reduction and chiral resolution, or a combination of two or more conditions.

In an eighth alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, A is C=O.


In a ninth alternative embodiment, the invention is a process for converting a compound of formula (I) into a compound of formula (II), wherein, the compound of formula (I) is a compound of formula (IA) or a salt or hydrate or solvate thereof,



(IA).


In an exemplary embodiment, the enzyme is SEQ ID NO: 134.

In a second embodiment, the invention is a process for preparing a compound of formula (IIA) or a salt or hydrate or solvate thereof,



(IIA)

comprising enzymatically transaminating a compound of formula (IA) or a salt or solvate or hydrate thereof,



(IA).


to provide the compound of formula (IIA).

Typically, the ketone, compound (I) is dissolved in an organic solvent, *e.g.* glycol, and added to an aqueous mixture of isopropyl amine HCL and pyridoxalphosphate, followed by the addition of TEA buffer. The pH is then adjusted to neutral with an appropriate base, *e.g.* NaOH followed by warming and addition of the transaminase. The reaction is allowed to stir at temperature for approximately 24 hrs. The solid chiral amine product (compound of formula (II))) is isolated by means known to one of skill in the art, and/or according to examples 10-12.

In an exemplary embodiment, the enzyme is SEQ ID NO: 134.


Section C: Conversion of a compound of formula (II) into a compound of formula (IV)

In a third embodiment, the invention is a process for preparing a compound of formula (IV), or a salt or hydrate or solvate thereof,




(IV)

comprising, reacting a compound of formula (III), or a salt or hydrate or solvate thereof,




(III)

with a compound of formula (II), or a salt or hydrate or solvate thereof,




(II)

wherein the compound of formula (II), or a salt or hydrate or solvate thereof, is prepared from a compound of formula (I), or a salt or hydrate or solvate thereof,



(I)

wherein: the dashed line is a bond or absent; A is selected from C=O and C=NH; or when the dashed line is a double bond A-R⁸ is:




; R^a is C₁₋₆ alkyl; R¹ is (C₁-C₆) alkyl, optionally substituted with an amino, (C₁-C₆) alkyl amino, (C₁-C₆) alkyl di-alkyl amino or (C₁-C₆) alkyl C(O)NH (C₁-C₆) alkyl; R⁴ and R⁷ are each, independently, H or halo; R⁵ and R⁶ are each, independently hydrogen, halo, hydroxyl, (C₁-C₆) alkyl, trihalo (C₁) alkyl, cyano or (C₁-C₆) alkoxy; R⁸ is (C₁-C₆) alkyl, optionally substituted with a hydroxyl; n is 1 or 2, R^{1'} is hydrogen or (C₁-C₆)alkyl; and R^{4'}, R^{5'}, R^{6'} and R^{7'}, are each, independently hydrogen, halo, hydroxy, amino, alkylamino, dialkylamino, (C₁-C₆)alkyl, and (C₁-C₆)alkyloxy.

In a tenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, R⁵ and R⁶ are fluoro when: R⁸ is -CH₃, and n is 1.

In an eleventh alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, R⁵ and R⁶ are fluoro and chloro when: R⁸ is -CH₃, and n is 1.


In a twelve alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, R⁵ and R⁶ are hydrogen when: R⁸ is -CH₃ and n is 1.

In a fourteenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, R⁵ is fluoro when: n is 1, and R⁶ is hydrogen.



R^4 and R^7 are each, independently, H or -Cl; R^5 is H, -OH, -CH₃, -OCH₃, -F, -Cl, -CF₃ or -CN; R^6 is H, -OH, -OCH₃, -F or -Cl; R^8 is H, -CH₃, -CH₂CH₃, -CH₂OH, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃ or -CF₃; and n is 1 or 2.

In a fifteenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, the compound of formula (II) is of formula (IIA), or a salt or solvate or hydrate thereof,




(IIA).

In a sixteenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, wherein $R^{5'}$ is a halo.

In a seventeenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, $R^{5'}$ is a chloro, and $R^{1'}$, $R^{4'}$, $R^{6'}$ and $R^{7'}$ are each hydrogen.


In an eighteenth alternative embodiment, the invention is a process for preparing a compound of formula (IV) wherein, the compound of formula (III) is of formula (IIIA), or a salt or hydrate or solvate thereof,



(IIIA).


In an exemplary embodiment, the enzyme is SEQ ID NO: 134.

In a fourth embodiment, the invention is a process for preparing a compound of formula (IVA), or a salt or solvate or hydrate thereof,




(IVA)

comprising: reacting a compound of formula (IIIA) or a salt or solvate or hydrate thereof,



(IIIA)


with a compound of formula (II) or a salt or solvate or hydrate thereof,




(IIA)

to provide the compound of formula (IVA), or a salt or solvate or hydrate thereof,

wherein the compound of formula (IIA) is prepared from the compound of formula (IA), or a salt or hydrate or solvate thereof,



In a nineteenth alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the compound of formula (IIA) is converted to a salt of formula (IIB):




prior to reaction with the compound of formula (IIIA).

In an exemplary embodiment, the enzyme is SEQ ID NO: 134.

In a twentieth alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the reaction is carried out under basic conditions.

In a twenty-first alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the reaction is carried out in the presence of triethyl amine.

In a twenty-second alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the compound of formula (IVA) is isolated as a salt of formula (IVB):



In a twenty-third alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the salt of formula (IVB) is converted to the compound of formula (IVA) in free base.

In a twenty-fourth alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the salt of formula (IVB) is converted to the compound of formula (IVA) in free base with sodium carbonate.


In a twenty-fifth alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the compound of formula (IVA) is a hydrate.

In a twenty-sixth alternative embodiment, the invention is a process for preparing a compound of formula (IVA) wherein, the compound of formula (IVA) is a $\frac{1}{2}$ hydrate.

In a twenty-seventh alternative embodiment, the invention is a process for preparing a compound of formula (IVA) $\frac{1}{2}$ hydrate wherein, the compound of formula (IVA) $\frac{1}{2}$ hydrate is milled after isolation.

Section D: Use of the Novel and Inventive Compounds of Formula (IC).

In a fifth embodiment, the invention is a compound of formula (IC):



wherein: R^1 is (C₁-C₆) alkyl, optionally substituted with an amino, (C₁-C₆) alkyl amino, (C₁-C₆) alkyl di-alkyl amino or (C₁-C₆) alkyl C(O)NH (C₁-6) alkyl; R^5 and R^6 are each, independently hydrogen, halo, hydroxyl, (C₁-C₆) alkyl, trihalo (C₁) alkyl, cyano or (C₁-C₆) alkoxy; R^8 is (C₁-C₆) alkyl, optionally substituted with a hydroxyl; n is 1 or 2; or a pharmaceutically acceptable salt or hydrate or solvate thereof.


In a twenty-eighth alternative embodiment, the invention is a compound of formula (IC) wherein, R^5 and R^6 are fluoro when: R^8 is -CH₃, and n is 1.

In a twenty-ninth alternative embodiment, the invention is a compound of formula (IC) wherein, R^5 and R^6 are fluoro and chloro, when: R^8 is -CH₃, and n is 1.

In a thirtieth alternative embodiment, the invention is a compound of formula (IC) wherein, R^5 and R^6 are hydrogen when: R^8 is -CH₃ and n is 1.


In a thirty-first alternative embodiment, the invention is a compound of formula (IC) wherein, R^5 is fluoro when: n is 1, and R^6 is hydrogen.

In a thirty-second alternative embodiment, the invention is a compound of formula (IC)





R⁵ is H, -OH, -CH₃, -OCH₃, -F, -Cl, -CF₃ or -CN; R⁶ is H, -OH, -OCH₃, -F or -Cl; R⁸ is H, -CH₃, -CH₂CH₃, -CH₂OH, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃ or -CF₃; and n is 1 or 2.

In a thirty-third alternative embodiment, the invention is a compound of formula (I) wherein, the compound is of formula (IA):



or a pharmaceutically acceptable salt or hydrate or solvate thereof.

In a sixth embodiment the invention is a compound selected from:



, or a pharmaceutically acceptable salt, solvate or

hydrate thereof.

Section I: General terms

Listed below are definitions of various terms used to describe the novel intermediates and synthetic steps of the present invention. These definitions, either by replacing one, more than one or all general expressions or symbols used in the present disclosure and thus yielding embodiments of the invention, in particular apply to the terms as they are used throughout the specification unless they are otherwise defined in specific instances either individually or as part of a larger group. Thus, the general definitions used above and below, unless defined differently, have the following meanings:

The term “C₁-C₂₀” defines a moiety with up to and including maximally 20, especially up to and including maximally 7 carbon atoms, said moiety being branched (one or more times) or straight-chained and bound via a terminal or a non-terminal carbon.

Alkyl being a radical or part of a radical is a straight or branched (one or, if desired and possible, more times) carbon chain, and is especially C₁-C₇-alkyl, such as C₁-C₄-alkyl, in particular branched C₁-C₄-alkyl, such as isopropyl. The term “lower” or “C₁-C₇” defines a moiety with up to and including maximally 7, especially up to and including maximally 4, carbon atoms, said moiety being branched (one or more times) or straight-chained and

bound via a terminal or a non-terminal carbon. Lower or C₁-C₇-alkyl, for example, is n-pentyl, n-hexyl or n-heptyl or preferably C₁-C₄-alkyl, especially as methyl, ethyl, n-propyl, iso-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl, in particular methyl, ethyl, n-propyl, iso-propyl, n-butyl, isobutyl, sec-butyl, tert-butyl; preferably methyl.

Alkylamino and dialkylamino refer to alkyl-NH- and (alkyl)₂N-, respectively, wherein alkyl may be linear or branched. The alkyl group for example comprises 1 to 7 and in particular 1 to 4 C atoms. Some examples are methylamino, dimethylamino, ethylamino, and diethylamino; preferably methylamino.

Halo or halogen is preferably fluoro, chloro, bromo or iodo, preferably fluoro or chloro; where halo is mentioned as a substituent, where possible, one or more (e.g. up to three or one) halogen atoms may be present, e.g. in halo-C₁-C₇-alkyl, such as trifluoromethyl, 2,2-difluoroethyl or 2,2,2-trifluoroethyl.

Halo-C₁-C₇-alkyl may be linear or branched and in particular comprises 1 to 4 C atoms, for example 1 or 2 C atoms. Examples are fluoromethyl, difluoromethyl, trifluoromethyl, chloromethyl, dichloromethyl, trichloromethyl, 2-chloroethyl and 2,2,2-trifluoroethyl; preferably trifluoromethyl.

Alkoxy, being a radical or part of a radical, refers to alkyl-O-, wherein the term alkyl is as defined herein, and includes, for example, C₁-C₂₀-alkoxy (-O-C₁-C₂₀alkyl), preferably C₁-C₇-alkoxy (-O-C₁-C₇alkyl). In particular, alkoxy includes, for example, methoxy, ethoxy, n-propyloxy, isopropyloxy, n-butyloxy, isobutyloxy, sec-butyloxy, tert-butyloxy, pentyloxy, hexyloxy and heptyloxy radicals; preferably methoxy.

The term "optically active base" describes, for example, chiral amines, preferably chiral tertiary amines, more preferably cinchona alkaloids, such as quinidine and quinine, most preferably modified cinchona alkaloids. Examples of such modified cinchona alkaloids are detailed, for example, in Tian, S.-K.; Chen, Y.; Hang, J.; Tang, L.; McDiad, P.; Deng, L. *Acc. Chem. Res.* **2004**, 37, 621-631 and references cited therein.

The term "phase transfer catalyst" as used herein refers to a catalytic amount of a chemical agent that enhances the rate of a reaction between chemical species located in different phases (eg. immiscible liquids or solid and liquid) by extracting one of the reactants, most commonly an anion, across the interface into the other phase. These catalysts include quaternary ammonium or phosphonium salts (e.g. tetraalkylammonium

salts, wherein alkyl can be same or different), or agents that complex inorganic cations (e.g. crown ethers or other cryptands). The catalyst cation is not consumed in the reaction although an anion exchange does occur. In particular, suitable phase transfer catalysts to be used according to the present invention are quaternary ammonium salts, for example of the formula $R_mR_nR_lR_kNX$, wherein $R_mR_nR_lR_k$ are, either the same or different, alkyl, and X is halo (eg. chloride, bromide, iodide) or hydroxide, for example, tetra-n-butylammonium hydroxide.

A "heterogeneous" catalyst as used herein refers to a catalyst supported on a carrier, typically although not necessarily a substrate comprised of an inorganic material, for example, a porous material such as carbon, silicon and/or aluminum oxide.

A "homogeneous" catalyst as used herein refers to a catalyst that is not supported on a carrier.

The term "chiral" refers to molecules which have the property of non-superimposability on their mirror image partner, while the term "achiral" refers to molecules which are superimposable on their mirror image partner.

The term "catalyst" means any substance that affects the rate of a chemical reaction by lowering the activation energy for the chemical reaction.

The term "powder catalyst" means a catalyst with a water contain of from 0 to 30 mass%.

The term "substrate to catalyst ratio" (S/C) refers to the molar ratio of starting compounds, or salts thereof, to "transition metal catalyst".

The term "work-up" means the work of isolation and/or purification which is carried out once the reaction is finished.

As used herein, unless specified otherwise, the term "room temperature" or "ambient temperature" means a temperature of from 15 to 30°C, such as of from 20 to 30°C, such as of from 20 to 25°C.

The term "inert" as used throughout this application, means unreactive with any of the reactants, solvents, or other components of the reaction mixture. Such inert conditions are generally accomplished by using inert gas such as carbon dioxide, helium, nitrogen, argon, among other gases.

Bonds with the asterisk (*) denote point of binding to the rest of the molecule.

The compounds of the present invention can possess one or more asymmetric centers. The preferred absolute configurations are as indicated herein specifically. However, any possible pure enantiomer, pure diastereoisomer, or mixtures thereof, e.g., mixtures of enantiomers, such as racemates, are encompassed by the present invention.

In the formulae of the present application the term "", "" or "—" on a C-sp³ represents a covalent bond wherein the stereochemistry of the bond is not defined. This means that the term "" or "—" on a C-sp³ comprises an (S) configuration as well as an (R) configuration of the respective chiral centre. Furthermore, mixtures are also encompassed, e.g., mixtures of enantiomers, such as racemates, are encompassed by the present invention.

In the formulae of the present application the term "" or "" on a C-sp² represents a covalent bond, wherein the stereochemistry or the geometry of the bond is not defined. This means that the term "" on a C-sp² comprises a *cis* (Z) configuration as well as a *trans* (E) configuration of the respective double bond. Furthermore, mixtures are also encompassed, e.g., mixtures of double bond isomers are encompassed by the present invention.

In the formulae of the present application, the term "=====" indicates a Csp³-Csp³ bond or a Csp²-Csp² bond.

The compounds of the present invention can possess one or more asymmetric centers. The preferred absolute configurations are as indicated herein specifically.

In the formulae of the present application the term "" on a C-sp³ indicates the absolute stereochemistry, either (R) or (S).

In the formulae of the present application the term "" on a C-sp³ indicates the absolute stereochemistry, either (R) or (S).

The term "stereomeric purity" at a given percentage means that the designated stereoisomer predominates at that given percentage in a mixture of stereoisomers.

The term "stereoisomer" means one of the absolute configurations of a single organic molecule having at least one asymmetric carbon. Included within the definition of a stereoisomer are enantiomers and diastereomers.

The term "resolution" refers to the separation or concentration or depletion of one of the stereoisomers of a molecule.

Salts are especially pharmaceutically acceptable salts or generally salts of any of the intermediates mentioned herein, where salts are not excluded for chemical reasons the skilled person will readily understand. They can be formed where salt forming groups, such as basic or acidic groups, are present that can exist in dissociated form at least partially, e.g. in a pH range from 4 to 10 in aqueous solutions, or can be isolated especially in solid, especially crystalline, form.

Such salts are formed, for example, as acid addition salts, preferably with organic or inorganic acids, from compounds or any of the intermediates mentioned herein with a basic nitrogen atom (e.g. imino or amino), especially the pharmaceutically acceptable salts. Suitable inorganic acids are, for example, halogen acids, such as hydrochloric acid, sulfuric acid, or phosphoric acid. Suitable organic acids are, for example, carboxylic, phosphonic, sulfonic or sulfamic acids, for example acetic acid, propionic acid, lactic acid, fumaric acid, succinic acid, citric acid, amino acids, such as glutamic acid or aspartic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, benzoic acid, methane- or ethane-sulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 2-naphthalenesulfonic acid, 1,5-naphthalene-disulfonic acid, N-cyclohexylsulfamic acid, N-methyl-, N-ethyl- or N-propyl-sulfamic acid, or other organic protonic acids, such as ascorbic acid.

In the presence of negatively charged radicals, such as carboxy or sulfo, salts may also be formed with bases, e.g. metal or ammonium salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium, magnesium or calcium salts, or ammonium salts with ammonia or suitable organic amines for example triethylamine or tri(2-hydroxyethyl)amine, N-ethyl-piperidine, N,N'-dimethylpiperazine, t-butylamine, n-butylamine, phenylethylamine, dicyclohexylamine or cyclohexylamine.

When a basic group and an acid group are present in the same molecule, any of the intermediates mentioned herein may also form internal salts.

For isolation or purification purposes of any of the intermediates mentioned herein it is also possible to use pharmaceutically unacceptable salts, for example picrates or perchlorates.

Preferred salts forms include, for example, acid addition salts. The compounds having at least one acid group (e.g., COOH or 5-tetrazolyl) can also form salts with bases. Suitable salts with bases are, e.g., metal salts, such as alkali metal or alkaline earth metal salts, e.g., sodium, potassium, calcium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, e.g., ethyl-, tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethylpropylamine, or a mono-, di- or trihydroxy lower alkylamine, e.g., mono-, di- or tri-ethanolamine. Corresponding internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, e.g., for the isolation or purification of free compounds I or their pharmaceutically acceptable salts, are also included. Most preferably the salt of formula (IV) is the camphorsulfonic acid salt.

In particular, the term "salt of a compound of formula (IV)" refers, for example, to an amine salt thereof, an alkali salt thereof or an earth alkali metal salt thereof (eg. sodium salt, potassium salt, calcium salt, magnesium salt, etc). In particular, the term "amine" in the expression "amine salt thereof", for example when referring to an amine salt of the compound of formula (IV), means tertiary amine of formula NR₉R₁₀R₁₁, secondary amine of formula NHR₉R₁₀R or primary amine of formula NH₂R₉, wherein R₉, R₁₀ and R₁₁ are, independently from one another, alkyl, aryl, cycloalkyl or heterocyclyl, as defined herein, preferably alkyl or cycloalkyl. The term "amine" is, for example, diphenylamine, diisopropylamine, dimethylamine, triethylamine, diisopropylethylamine, dicyclohexylamine, t-butylamine, n-butylamine or cyclohexylamine, in particular, t-butylamine, n-butylamine or cyclohexylamine, more preferably n-butylamine or cyclohexylamine.

As used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to "a polypeptide" includes more than one polypeptide.

Similarly, "comprise," "comprises," "comprising" "include," "includes," and "including" are interchangeable and not intended to be limiting.

It is to be further understood that where descriptions of various embodiments use the term "comprising," those skilled in the art would understand that in some specific instances, an embodiment can be alternatively described using language "consisting essentially of" or "consisting of."

It is to be understood that both the foregoing general description, including the drawings, and the following detailed description are exemplary and explanatory only and are not restrictive of this disclosure.

The section headings used herein are for organizational purposes only and not to be construed as limiting the subject matter described.

The abbreviations used for the genetically encoded amino acids are conventional and are as follows:

Amino Acid	Three-Letter	One-Letter Abbreviation
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartate	Asp	D
Cysteine	Cys	C
Glutamate	Glu	E
Glutamine	Gln	Q
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V

When the three-letter abbreviations are used, unless specifically preceded by an “L” or a “D” or clear from the context in which the abbreviation is used, the amino acid may be in either the L- or D-configuration about α -carbon (C_α). For example, whereas “Ala” designates alanine without specifying the configuration about the α -carbon, “D-Ala” and “L-Ala” designate D-alanine and L-alanine, respectively. When the one-letter

abbreviations are used, upper case letters designate amino acids in the L-configuration about the α -carbon and lower case letters designate amino acids in the D-configuration about the α -carbon. For example, "A" designates L-alanine and "a" designates D-alanine. When polypeptide sequences are presented as a string of one-letter or three-letter abbreviations (or mixtures thereof), the sequences are presented in the amino (N) to carboxy (C) direction in accordance with common convention.

The abbreviations used for the genetically encoding nucleosides are conventional and are as follows: adenosine (A); guanosine (G); cytidine (C); thymidine (T); and uridine (U). Unless specifically delineated, the abbreviated nucleotides may be either ribonucleosides or 2'-deoxyribonucleosides. The nucleosides may be specified as being either ribonucleosides or 2'-deoxyribonucleosides on an individual basis or on an aggregate basis. When nucleic acid sequences are presented as a string of one-letter abbreviations, the sequences are presented in the 5' to 3' direction in accordance with common convention, and the phosphates are not indicated.

In reference to the present disclosure, the technical and scientific terms used in the descriptions herein will have the meanings commonly understood by one of ordinary skill in the art, unless specifically defined otherwise. Accordingly, the following terms are intended to have the following meanings:


"Protein", "polypeptide," and "peptide" are used interchangeably herein to denote a polymer of at least two amino acids covalently linked by an amide bond, regardless of length or post-translational modification (e.g., glycosylation, phosphorylation, lipidation, myristilation, ubiquitination, etc.). Included within this definition are D- and L-amino acids, and mixtures of D- and L-amino acids.

"Polynucleotide" or "nucleic acid" refers to two or more nucleosides that are covalently linked together. The polynucleotide may be wholly comprised ribonucleosides (*i.e.*, an RNA), wholly comprised of 2' deoxyribonucleotides (*i.e.*, a DNA) or mixtures of ribo- and 2' deoxyribonucleosides. While the nucleosides will typically be linked together via standard phosphodiester linkages, the polynucleotides may include one or more non-standard linkages. The polynucleotide may be single-stranded or double-stranded, or may include both single-stranded regions and double-stranded regions. Moreover, while a polynucleotide will typically be composed of the naturally occurring encoding nucleobases (*i.e.*, adenine, guanine, uracil, thymine and cytosine), it may include one or

more modified and/or synthetic nucleobases, such as, for example, inosine, xanthine, hypoxanthine, *etc.* Preferably, such modified or synthetic nucleobases will be encoding nucleobases.

“Aminotransferase” and “transaminase” are used interchangeably herein to refer to a polypeptide having an enzymatic capability of transferring an amino group (NH₂) from a primary amine to a carbonyl group (C=O) of an acceptor molecule. Transaminases as used herein include naturally occurring (wild type) transaminase as well as non-naturally occurring engineered polypeptides generated by human manipulation.

“Amino acceptor” and “amine acceptor,” “keto substrate,” “keto,” and “ketone” are used interchangeably herein to refer to a carbonyl (keto, or ketone) compound which accepts an amino group from a donor amine. In some embodiments, amino acceptors are molecules of the following general formula,




amino acceptor

in which each of R^α and R^β, when taken independently, is an alkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, which can be unsubstituted or substituted with one or more enzymatically acceptable groups. R^α may be the same or different from R^β in structure or chirality. In some embodiments, R^α and R^β, taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. Amino acceptors include keto carboxylic acids and alkanones (ketones). Typical keto carboxylic acids are α-keto carboxylic acids such as glyoxalic acid, pyruvic acid, oxaloacetic acid, and the like, as well as salts of these acids. Amino acceptors also include substances which are converted to an amino acceptor by other enzymes or whole cell processes, such as fumaric acid (which can be converted to oxaloacetic acid), glucose (which can be converted to pyruvate), lactate, maleic acid, and others. Amino acceptors that can be used include, by way of example and not limitation, 3,4-dihydronaphthalen-1(2H)-one, 1-phenylbutan-2-one, 3,3-dimethylbutan-2-one, octan-2-one, ethyl 3-oxobutanoate, 4-phenylbutan-2-one, 1-(4-bromophenyl)ethanone, 2-methyl-cyclohexanone, 7-methoxy-2-tetralone, 1-hydroxybutan-2-one, pyruvic acid, acetophenone, 3'-

hydroxyacetophenone, 2-methoxy-5-fluoroacetophenone, levulinic acid, 1-phenylpropan-1-one, 1-(4-bromophenyl)propan-1-one, 1-(4-nitrophenyl)propan-1-one, 1-phenylpropan-2-one, 2-oxo-3-methylbutanoic acid, 1-(3-trifluoromethylphenyl)propan-1-one, hydroxypropanone, methoxyoxypropanone, 1-phenylbutan-1-one, 1-(2,5-dimethoxy-4-methylphenyl)butan-2-one, 1-(4-hydroxyphenyl)butan-3-one, 2-acetylnaphthalene, phenylpyruvic acid, 2-ketoglutaric acid, and 2-ketosuccinic acid, including both (*R*) and (*S*) single isomers where possible.

“Amino donor” or “amine donor” refers to an amino compound which donates an amino group to the amino acceptor, thereby becoming a carbonyl species. In some embodiments, amino donors are molecules of the following general formula,



amino donor

in which each of R^ϵ and R^δ , when taken independently, is an alkyl, cycloalkyl, heterocycloalkyl, aryl, or heteroaryl, which is unsubstituted or substituted with one or more enzymatically non-inhibiting groups. R^ϵ can be the same or different from R^δ in structure or chirality. In some embodiments, R^ϵ and R^δ , taken together, may form a ring that is unsubstituted, substituted, or fused to other rings. Typical amino donors that can be used include chiral and achiral amino acids, and chiral and achiral amines. Amino donors that can be used include, by way of example and not limitation, isopropylamine (also referred to as 2-aminopropane), α -phenethylamine (also termed 1-phenylethanamine), and its enantiomers (*S*)-1-phenylethanamine and (*R*)-1-phenylethanamine, 2-amino-4-phenylbutane, glycine, L-glutamic acid, L-glutamate, monosodium glutamate, L-alanine, D-alanine, D,L-alanine, L-aspartic acid, L-lysine, D,L-ornithine, β -alanine, taurine, n-octylamine, cyclohexylamine, 1,4-butanediamine (also referred to as putrescine), 1,6-hexanediamine, 6-aminohexanoic acid, 4-aminobutyric acid, tyramine, and benzyl amine, 2-aminobutane, 2-amino-1-butanol, 1-amino-1-phenylethane, 1-amino-1-(2-methoxy-5-fluorophenyl)ethane, 1-amino-1-phenylpropane, 1-amino-1-(4-hydroxyphenyl)propane, 1-amino-1-(4-bromophenyl)propane, 1-amino-1-(4-nitrophenyl)propane, 1-phenyl-2-aminopropane, 1-(3-trifluoromethylphenyl)-2-aminopropane, 2-aminopropanol, 1-amino-1-phenylbutane, 1-phenyl-2-aminobutane, 1-

(2,5-dimethoxy-4-methylphenyl)-2-aminobutane, 1-phenyl-3-aminobutane, 1-(4-hydroxyphenyl)-3-aminobutane, 1-amino-2-methylcyclopentane, 1-amino-3-methylcyclopentane, 1-amino-2-methylcyclohexane, 1-amino-1-(2-naphthyl)ethane, 3-methylcyclopentylamine, 2-methylcyclopentylamine, 2-ethylcyclopentylamine, 2-methylcyclohexylamine, 3-methylcyclohexylamine, 1-aminotetralin, 2-aminotetralin, 2-amino-5-methoxytetralin, and 1-aminoindan, including both (*R*) and (*S*) single isomers where possible and including all possible salts of the amines.

“Chiral amine” refers to amines of general formula $R^1\text{-CH(NH}_2\text{)-R}^1$ and is employed herein in its broadest sense, including a wide variety of aliphatic and alicyclic compounds of different, and mixed, functional types, characterized by the presence of a primary amino group bound to a secondary carbon atom which, in addition to a hydrogen atom, carries either (i) a divalent group forming a chiral cyclic structure, or (ii) two substituents (other than hydrogen) differing from each other in structure or chirality. Divalent groups forming a chiral cyclic structure include, for example, 2-methylbutane-1,4-diyl, pentane-1,4-diyl, hexane-1,4-diyl, hexane-1,5-diyl, 2-methylpentane-1,5-diyl. The two different substituents on the secondary carbon atom (R^1 and R^2 above) also can vary widely and include alkyl, aralkyl, aryl, halo, hydroxy, lower alkyl, lower alkoxy, lower alkylthio, cycloalkyl, carboxy, carbalkoxy, carbamoyl, mono- and di-(lower alkyl) substituted carbamoyl, trifluoromethyl, phenyl, nitro, amino, mono- and di-(lower alkyl) substituted amino, alkylsulfonyl, arylsulfonyl, alkylcarboxamido, arylcarboxamido, etc., as well as alkyl, aralkyl, or aryl substituted by the foregoing.

“Pyridoxal-phosphate,” “PLP,” “pyridoxal-5'-phosphate,” “PYP,” and “P5P” are used interchangeably herein to refer to the compound that acts as a coenzyme in transaminase reactions. In some embodiments, pyridoxal phosphate is defined by the structure 1-(4'-formyl-3'-hydroxy-2'-methyl-5'-pyridyl)methoxyphosphonic acid, CAS number [54-47-7], Pyridoxal-5'-phosphate can be produced *in vivo* by phosphorylation and oxidation of pyridoxol (also known as Vitamin B₆). In transamination reactions using transaminase enzymes, the amine group of the amino donor is transferred to the coenzyme to produce a keto byproduct, while pyridoxal-5'-phosphate is converted to pyridoxamine phosphate. Pyridoxal-5'-phosphate is regenerated by reaction with a different keto compound (the amino acceptor). The transfer of the amine group from pyridoxamine phosphate to the amino acceptor produces a chiral amine and regenerates the coenzyme. In some embodiments, the pyridoxal-5'-phosphate can be replaced by

other members of the vitamin B₆ family, including pyridoxine (PN), pyridoxal (PL), pyridoxamine (PM), and their phosphorylated counterparts; pyridoxine phosphate (PNP), and pyridoxamine phosphate (PMP).

“Coding sequence” refers to that portion of a nucleic acid (*e.g.*, a gene) that encodes an amino acid sequence of a protein.

“Naturally-occurring” or “wild-type” refers to the form found in nature. For example, a naturally occurring or wild-type polypeptide or polynucleotide sequence is a sequence present in an organism that can be isolated from a source in nature and which has not been intentionally modified by human manipulation.

“Recombinant” or “engineered” or “non-naturally occurring” when used with reference to, *e.g.*, a cell, nucleic acid, or polypeptide, refers to a material, or a material corresponding to the natural or native form of the material, that has been modified in a manner that would not otherwise exist in nature, or is identical thereto but produced or derived from synthetic materials and/or by manipulation using recombinant techniques. Non-limiting examples include, among others, recombinant cells expressing genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise expressed at a different level.

“Percentage of sequence identity” and “percentage homology” are used interchangeably herein to refer to comparisons among polynucleotides and polypeptides, and are determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference sequence for optimal alignment of the two sequences. The percentage may be calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Alternatively, the percentage may be calculated by determining the number of positions at which either the identical nucleic acid base or amino acid residue occurs in both sequences or a nucleic acid base or amino acid residue is aligned with a gap to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity. Those of skill in the art appreciate

that there are many established algorithms available to align two sequences. Optimal alignment of sequences for comparison can be conducted, *e.g.*, by the local homology algorithm of Smith and Waterman, 1981, *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the GCG Wisconsin Software Package), or by visual inspection (see generally, *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)). Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., 1990, *J. Mol. Biol.* 215: 403-410 and Altschul et al., 1977, *Nucleic Acids Res.* 3389-3402, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length *W* in the query sequence, which either match or satisfy some positive-valued threshold score *T* when aligned with a word of the same length in a database sequence. *T* is referred to as, the neighborhood word score threshold (Altschul et al, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters *M* (reward score for a pair of matching residues; always >0) and *N* (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity *X* from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters *W*, *T*, and *X* determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (*W*) of 11, an expectation (*E*) of 10, *M*=5, *N*=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (*W*) of 3, an expectation (*E*) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff, 1989, *Proc Natl Acad Sci USA* 89:10915).

Exemplary determination of sequence alignment and % sequence identity can employ the BESTFIT or GAP programs in the GCG Wisconsin Software package (Accelrys, Madison WI), using default parameters provided.

“Reference sequence” refers to a defined sequence used as a basis for a sequence comparison. A reference sequence may be a subset of a larger sequence, for example, a segment of a full-length gene or polypeptide sequence. Generally, a reference sequence is at least 20 nucleotide or amino acid residues in length, at least 25 residues in length, at least 50 residues in length, or the full length of the nucleic acid or polypeptide. Since two polynucleotides or polypeptides may each (1) comprise a sequence (*i.e.*, a portion of the complete sequence) that is similar between the two sequences, and (2) may further comprise a sequence that is divergent between the two sequences, sequence comparisons between two (or more) polynucleotides or polypeptide are typically performed by comparing sequences of the two polynucleotides or polypeptides over a “comparison window” to identify and compare local regions of sequence similarity. In some embodiments, a “reference sequence” can be based on a primary amino acid sequence, where the reference sequence is a sequence that can have one or more changes in the primary sequence. For instance, a “reference sequence based on SEQ ID NO:4 having at the residue corresponding to X14 a valine” or X14V refers to a reference sequence in which the corresponding residue at X14 in SEQ ID NO:4, which is a tyrosine, has been changed to valine.

“Comparison window” refers to a conceptual segment of at least about 20 contiguous nucleotide positions or amino acids residues wherein a sequence may be compared to a reference sequence of at least 20 contiguous nucleotides or amino acids and wherein the portion of the sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The comparison window can be longer than 20 contiguous residues, and includes, optionally 30, 40, 50, 100, or longer windows.

“Substantial identity” refers to a polynucleotide or polypeptide sequence that has at least 80 percent sequence identity, at least 85 percent identity and 89 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 residue positions, frequently over a window of at least 30-50 residues, wherein the percentage of sequence identity is

calculated by comparing the reference sequence to a sequence that includes deletions or additions which total 20 percent or less of the reference sequence over the window of comparison. In specific embodiments applied to polypeptides, the term "substantial identity" means that two polypeptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least 80 percent sequence identity, preferably at least 89 percent sequence identity, at least 95 percent sequence identity or more (*e.g.*, 99 percent sequence identity). Preferably, residue positions which are not identical differ by conservative amino acid substitutions.

"Corresponding to", "reference to" or "relative to" when used in the context of the numbering of a given amino acid or polynucleotide sequence refers to the numbering of the residues of a specified reference sequence when the given amino acid or polynucleotide sequence is compared to the reference sequence. In other words, the residue number or residue position of a given polymer is designated with respect to the reference sequence rather than by the actual numerical position of the residue within the given amino acid or polynucleotide sequence. For example, a given amino acid sequence, such as that of an engineered transaminase, can be aligned to a reference sequence by introducing gaps to optimize residue matches between the two sequences. In these cases, although the gaps are present, the numbering of the residue in the given amino acid or polynucleotide sequence is made with respect to the reference sequence to which it has been aligned.

"Amino acid difference" or "residue difference" refers to a change in the amino acid residue at a position of a polypeptide sequence relative to the amino acid residue at a corresponding position in a reference sequence. The positions of amino acid differences generally are referred to herein as "X_n," where n refers to the corresponding position in the reference sequence upon which the residue difference is based. For example, a "residue difference at position X14 as compared to SEQ ID NO: 4" refers to a change of the amino acid residue at the polypeptide position corresponding to position 14 of SEQ ID NO:4. Thus, if the reference polypeptide of SEQ ID NO: 4 has a tyrosine at position 14, then a "residue difference at position X14 as compared to SEQ ID NO:4" an amino acid substitution of any residue other than tyrosine at the position of the polypeptide corresponding to position 14 of SEQ ID NO: 4. In most instances herein, the specific amino acid residue difference at a position is indicated as "X_nY" where "X_n" specified the corresponding position as described above, and "Y" is the single letter identifier of the

amino acid found in the engineered polypeptide (*i.e.*, the different residue than in the reference polypeptide). In some embodiments, there more than one amino acid can appear in a specified residue position, the alternative amino acids can be listed in the form XnY/Z, where Y and Z represent alternate amino acid residues. In some instances (*e.g.*, in Table **2A and 2B**), the present disclosure also provides specific amino acid differences denoted by the conventional notation “AnB”, where A is the single letter identifier of the residue in the reference sequence, “n” is the number of the residue position in the reference sequence, and B is the single letter identifier of the residue substitution in the sequence of the engineered polypeptide. Furthermore, in some instances, a polypeptide of the present disclosure can include one or more amino acid residue differences relative to a reference sequence, which is indicated by a list of the specified positions where changes are made relative to the reference sequence. The present disclosure includes engineered polypeptide sequences comprising one or more amino acid differences that include either/or both conservative and non-conservative amino acid substitutions.

“Conservative amino acid substitution” refers to a substitution of a residue with a different residue having a similar side chain, and thus typically involves substitution of the amino acid in the polypeptide with amino acids within the same or similar defined class of amino acids. By way of example and not limitation, an amino acid with an aliphatic side chain may be substituted with another aliphatic amino acid, *e.g.*, alanine, valine, leucine, and isoleucine; an amino acid with hydroxyl side chain is substituted with another amino acid with a hydroxyl side chain, *e.g.*, serine and threonine; an amino acid having aromatic side chains is substituted with another amino acid having an aromatic side chain, *e.g.*, phenylalanine, tyrosine, tryptophan, and histidine; an amino acid with a basic side chain is substituted with another amino acid with a basic side chain, *e.g.*, lysine and arginine; an amino acid with an acidic side chain is substituted with another amino acid with an acidic side chain, *e.g.*, aspartic acid or glutamic acid; and a hydrophobic or hydrophilic amino acid is replaced with another hydrophobic or hydrophilic amino acid, respectively. Exemplary conservative substitutions are provided in Table 1 below.

Table 1

Residue	Possible Conservative Substitutions
A, L, V, I	Other aliphatic (A, L, V, I) Other non-polar (A, L, V, I, G, M)

G, M	Other non-polar (A, L, V, I, G, M)
D, E	Other acidic (D, E)
K, R	Other basic (K, R)
N, Q, S, T	Other polar
H, Y, W, F	Other aromatic (H, Y, W, F)
C, P	None

“Non-conservative substitution” refers to substitution of an amino acid in the polypeptide with an amino acid with significantly differing side chain properties. Non-conservative substitutions may use amino acids between, rather than within, the defined groups and affects (a) the structure of the peptide backbone in the area of the substitution (*e.g.*, proline for glycine), (b) the charge or hydrophobicity, or (c) the bulk of the side chain. By way of example and not limitation, an exemplary non-conservative substitution can be an acidic amino acid substituted with a basic or aliphatic amino acid; an aromatic amino acid substituted with a small amino acid; and a hydrophilic amino acid substituted with a hydrophobic amino acid.

“Deletion” refers to modification to the polypeptide by removal of one or more amino acids from the reference polypeptide. Deletions can comprise removal of 1 or more amino acids, 2 or more amino acids, 5 or more amino acids, 10 or more amino acids, 15 or more amino acids, or 20 or more amino acids, up to 10% of the total number of amino acids, or up to 20% of the total number of amino acids making up the reference enzyme while retaining enzymatic activity and/or retaining the improved properties of an engineered transaminase enzyme. Deletions can be directed to the internal portions and/or terminal portions of the polypeptide. In various embodiments, the deletion can comprise a continuous segment or can be discontinuous.

“Insertion” refers to modification to the polypeptide by addition of one or more amino acids from the reference polypeptide. In some embodiments, the improved engineered transaminase enzymes comprise insertions of one or more amino acids to the naturally occurring transaminase polypeptide as well as insertions of one or more amino acids to other improved transaminase polypeptides. Insertions can be in the internal portions of the polypeptide, or to the carboxy or amino terminus. Insertions as used herein include fusion proteins as is known in the art. The insertion can be a contiguous segment of amino acids or separated by one or more of the amino acids in the naturally occurring polypeptide.

"Fragment" as used herein refers to a polypeptide that has an amino-terminal and/or carboxy-terminal deletion, but where the remaining amino acid sequence is identical to the corresponding positions in the sequence. Fragments can be at least 14 amino acids long, at least 20 amino acids long, at least 50 amino acids long or longer, and up to 70%, 80%, 90%, 95%, 98%, and 99% of the full-length transaminase polypeptide, for example the polypeptide of SEQ ID NO:2 or engineered transaminase of SEQ ID NO:34.

"Isolated polypeptide" refers to a polypeptide which is substantially separated from other contaminants that naturally accompany it, *e.g.*, protein, lipids, and polynucleotides. The term embraces polypeptides which have been removed or purified from their naturally-occurring environment or expression system (*e.g.*, host cell or *in vitro* synthesis). The improved transaminase enzymes may be present within a cell, present in the cellular medium, or prepared in various forms, such as lysates or isolated preparations. As such, in some embodiments, the improved transaminase enzyme can be an isolated polypeptide.

"Substantially pure polypeptide" refers to a composition in which the polypeptide species is the predominant species present (*i.e.*, on a molar or weight basis it is more abundant than any other individual macromolecular species in the composition), and is generally a substantially purified composition when the object species comprises at least about 50 percent of the macromolecular species present by mole or % weight. Generally, a substantially pure transaminase composition will comprise about 60 % or more, about 70% or more, about 80% or more, about 90% or more, about 95% or more, and about 98% or more of all macromolecular species by mole or % weight present in the composition. In some embodiments, the object species is purified to essential homogeneity (*i.e.*, contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500 Daltons), and elemental ion species are not considered macromolecular species. In some embodiments, the isolated improved transaminases polypeptide is a substantially pure polypeptide composition.

"Stereoselectivity" refers to the preferential formation in a chemical or enzymatic reaction of one stereoisomer over another. Stereoselectivity can be partial, where the formation of one stereoisomer is favored over the other, or it may be complete where only one stereoisomer is formed. When the stereoisomers are enantiomers, the

stereoselectivity is referred to as enantioselectivity, the fraction (typically reported as a percentage) of one enantiomer in the sum of both. It is commonly alternatively reported in the art (typically as a percentage) as the enantiomeric excess (e.e.) calculated therefrom according to the formula $[\text{major enantiomer} - \text{minor enantiomer}] / [\text{major enantiomer} + \text{minor enantiomer}]$. Where the stereoisomers are diastereoisomers, the stereoselectivity is referred to as diastereoselectivity, the fraction (typically reported as a percentage) of one diastereomer in a mixture of two diastereomers, commonly alternatively reported as the diastereomeric excess (d.e.). Enantiomeric excess and diastereomeric excess are types of stereomeric excess.

“Highly stereoselective” refers to a chemical or enzymatic reaction that is capable of converting a substrate, *e.g.*, Compound (IA), to its corresponding chiral amine product, *e.g.*, Compound (IIA), with at least about 85% stereomeric excess.

“Improved enzyme property” refers to a transaminase polypeptide that exhibits an improvement in any enzyme property as compared to a reference transaminase. For the engineered transaminase polypeptides described herein, the comparison is generally made to the wild-type transaminase enzyme, although in some embodiments, the reference transaminase can be another improved engineered transaminase. Enzyme properties for which improvement is desirable include, but are not limited to, enzymatic activity (which can be expressed in terms of percent conversion of the substrate), thermo stability, solvent stability, pH activity profile, cofactor requirements, refractoriness to inhibitors (*e.g.*, substrate or product inhibition), stereospecificity, and stereoselectivity (including enantioselectivity).

“Increased enzymatic activity” refers to an improved property of the engineered transaminase polypeptides, which can be represented by an increase in specific activity (*e.g.*, product produced/time/weight protein) or an increase in percent conversion of the substrate to the product (*e.g.*, percent conversion of starting amount of substrate to product in a specified time period using a specified amount of transaminase) as compared to the reference transaminase enzyme. Exemplary methods to determine enzyme activity are provided in the Examples. Any property relating to enzyme activity may be affected, including the classical enzyme properties of K_m , V_{max} or k_{cat} , changes of which can lead to increased enzymatic activity. Improvements in enzyme activity can be from about 1.2 times the enzymatic activity of the corresponding wild-type transaminase enzyme, to as much as 2 times, 5 times, 10 times, 20 times, 25 times, 50 times, 75

times, 100 times, or more enzymatic activity than the naturally occurring transaminase or another engineered transaminase from which the transaminase polypeptides were derived. Transaminase activity can be measured by any one of standard assays, such as by monitoring changes in spectrophotometric properties of reactants or products. In some embodiments, the amount of products produced can be measured by High-Performance Liquid Chromatography (HPLC) separation combined with UV absorbance or fluorescent detection following derivatization, such as with o-phthaldialdehyde (OPA). Comparisons of enzyme activities are made using a defined preparation of enzyme, a defined assay under a set condition, and one or more defined substrates, as further described in detail herein. Generally, when lysates are compared, the numbers of cells and the amount of protein assayed are determined as well as use of identical expression systems and identical host cells to minimize variations in amount of enzyme produced by the host cells and present in the lysates.

“Conversion” refers to the enzymatic conversion of the substrate(s) to the corresponding product(s). “Percent conversion” refers to the percent of the substrate that is converted to the product within a period of time under specified conditions. Thus, the “enzymatic activity” or “activity” of a transaminase polypeptide can be expressed as “percent conversion” of the substrate to the product.

“Thermostable” refers to a transaminase polypeptide that maintains similar activity (more than 60% to 80% for example) after exposure to elevated temperatures (*e.g.*, 40-80°C) for a period of time (*e.g.*, 0.5-24 hrs) compared to the wild-type enzyme.

“Solvent stable” refers to a transaminase polypeptide that maintains similar activity (more than *e.g.*, 60% to 80%) after exposure to varying concentrations (*e.g.*, 5-99%) of solvent (ethanol, isopropyl alcohol, dimethylsulfoxide (DMSO), tetrahydrofuran, 2-methyltetrahydrofuran, acetone, toluene, butyl acetate, methyl tert-butyl ether, etc.) for a period of time (*e.g.*, 0.5-24 hrs) compared to the wild-type enzyme.

“Thermo- and solvent stable” refers to a transaminase polypeptide that is both thermostable and solvent stable.

“Stringent hybridization” is used herein to refer to conditions under which nucleic acid hybrids are stable. As known to those of skill in the art, the stability of hybrids is reflected in the melting temperature (T_m) of the hybrids. In general, the stability of a hybrid is a function of ion strength, temperature, G/C content, and the presence of chaotropic

agents. The T_m values for polynucleotides can be calculated using known methods for predicting melting temperatures (see, *e.g.*, Baldino et al., *Methods Enzymology* 168:761-777; Bolton et al., 1962, *Proc. Natl. Acad. Sci. USA* 48:1390; Bresslauer et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:8893-8897; Freier et al., 1986, *Proc. Natl. Acad. Sci. USA* 83:9373-9377; Kierzek et al., *Biochemistry* 25:7840-7846; Rychlik et al., 1990, *Nucleic Acids Res* 18:6409-6412 (erratum, 1991, *Nucleic Acids Res* 19:698); Sambrook et al., *supra*); Suggs et al., 1981, In *Developmental Biology Using Purified Genes* (Brown et al., eds.), pp. 683-693, Academic Press; and Wetmur, 1991, *Crit Rev Biochem Mol Biol* 26:227-259. All publications incorporate herein by reference). In some embodiments, the polynucleotide encodes the polypeptide disclosed herein and hybridizes under defined conditions, such as moderately stringent or highly stringent conditions, to the complement of a sequence encoding an engineered transaminase enzyme of the present disclosure.

"Hybridization stringency" relates to hybridization conditions, such as washing conditions, in the hybridization of nucleic acids. Generally, hybridization reactions are performed under conditions of lower stringency, followed by washes of varying but higher stringency. The term "moderately stringent hybridization" refers to conditions that permit target-DNA to bind a complementary nucleic acid that has about 60% identity, preferably about 75% identity, about 85% identity to the target DNA, with greater than about 90% identity to target-polynucleotide. Exemplary moderately stringent conditions are conditions equivalent to hybridization in 50% formamide, 5× Denhart's solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.2×SSPE, 0.2% SDS, at 42°C. "High stringency hybridization" refers generally to conditions that are about 10°C or less from the thermal melting temperature T_m as determined under the solution condition for a defined polynucleotide sequence. In some embodiments, a high stringency condition refers to conditions that permit hybridization of only those nucleic acid sequences that form stable hybrids in 0.018M NaCl at 65°C (*i.e.*, if a hybrid is not stable in 0.018M NaCl at 65°C, it will not be stable under high stringency conditions, as contemplated herein). High stringency conditions can be provided, for example, by hybridization in conditions equivalent to 50% formamide, 5× Denhart's solution, 5×SSPE, 0.2% SDS at 42°C, followed by washing in 0.1×SSPE, and 0.1% SDS at 65°C. Another high stringency condition is hybridizing in conditions equivalent to hybridizing in 5X SSC containing 0.1% (w:v) SDS at 65°C and washing in 0.1x SSC containing 0.1% SDS at 65°C. Other high

stringency hybridization conditions, as well as moderately stringent conditions, are described in the references cited above.

"Heterologous" polynucleotide refers to any polynucleotide that is introduced into a host cell by laboratory techniques, and includes polynucleotides that are removed from a host cell, subjected to laboratory manipulation, and then reintroduced into a host cell.

"Codon optimized" refers to changes in the codons of the polynucleotide encoding a protein to those preferentially used in a particular organism such that the encoded protein is efficiently expressed in the organism of interest. Although the genetic code is degenerate in that most amino acids are represented by several codons, called "synonyms" or "synonymous" codons, it is well known that codon usage by particular organisms is nonrandom and biased towards particular codon triplets. This codon usage bias may be higher in reference to a given gene, genes of common function or ancestral origin, highly expressed proteins versus low copy number proteins, and the aggregate protein coding regions of an organism's genome. In some embodiments, the polynucleotides encoding the transaminase enzymes may be codon optimized for optimal production from the host organism selected for expression.

"Preferred, optimal, high codon usage bias codons" refers interchangeably to codons that are used at higher frequency in the protein coding regions than other codons that code for the same amino acid. The preferred codons may be determined in relation to codon usage in a single gene, a set of genes of common function or origin, highly expressed genes, the codon frequency in the aggregate protein coding regions of the whole organism, codon frequency in the aggregate protein coding regions of related organisms, or combinations thereof. Codons whose frequency increases with the level of gene expression are typically optimal codons for expression. A variety of methods are known for determining the codon frequency (*e.g.*, codon usage, relative synonymous codon usage) and codon preference in specific organisms, including multivariate analysis, for example, using cluster analysis or correspondence analysis, and the effective number of codons used in a gene (see GCG CodonPreference, Genetics Computer Group Wisconsin Package; CodonW, John Peden, University of Nottingham; McInerney, J. O, 1998, *Bioinformatics* 14:372-73; Stenico et al., 1994, *Nucleic Acids Res.* 22:2437-46; Wright, F., 1990, *Gene* 87:23-29). Codon usage tables are available for a growing list of organisms (see for example, Wada et al., 1992, *Nucleic Acids Res.* 20:2111-2118; Nakamura et al., 2000, *Nucl. Acids Res.* 28:292; Duret, et al., *supra*;

Henaut and Danchin, "Escherichia coli and Salmonella," 1996, Neidhardt, et al. Eds., ASM Press, Washington D.C., p. 2047-2066. The data source for obtaining codon usage may rely on any available nucleotide sequence capable of coding for a protein. These data sets include nucleic acid sequences actually known to encode expressed proteins (*e.g.*, complete protein coding sequences-CDS), expressed sequence tags (ESTS), or predicted coding regions of genomic sequences (see for example, Mount, D., Bioinformatics: Sequence and Genome Analysis, Chapter 8, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 2001; Uberbacher, E. C., 1996, Methods Enzymol. 266:259-281; Tiwari et al., 1997, Comput. Appl. Biosci. 13:263-270).

"Control sequence" is defined herein to include all components, which are necessary or advantageous for the expression of a polynucleotide and/or polypeptide of the present disclosure. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide.

"Operably linked" is defined herein as a configuration in which a control sequence is appropriately placed (*i.e.*, in a functional relationship) at a position relative to a polynucleotide of interest such that the control sequence directs or regulates the expression of the polynucleotide and/or polypeptide of interest.

"Promoter sequence" refers to a nucleic acid sequence that is recognized by a host cell for expression of a polynucleotide of interest, such as a coding sequence. The promoter sequence contains transcriptional control sequences, which mediate the expression of a polynucleotide of interest. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

"Suitable reaction conditions" refer to those conditions in the biocatalytic reaction solution (*e.g.*, ranges of enzyme loading, substrate loading, cofactor loading, temperature, pH, buffers, co-solvents, etc.) under which a transaminase polypeptide of

the present disclosure is capable of converting a substrate compound to a product compound (e.g., conversion of Compound (IA) to Compound (IIA)). Exemplary "suitable reaction conditions" are provided in the present disclosure and illustrated by the Examples.

"Loading", such as in "compound loading" or "enzyme loading" or "cofactor loading" refers to the concentration or amount of a component in a reaction mixture at the start of the reaction.

"Substrate" in the context of a biocatalyst mediated process refers to the compound or molecule acted on by the biocatalyst. For example, an exemplary substrate for the transaminase biocatalyst in the process disclosed herein is compound (IA).

"Product" in the context of a biocatalyst mediated process refers to the compound or molecule resulting from the action of the biocatalyst. For example, an exemplary product for the transaminase biocatalyst in the process disclosed herein is compound (IIA).

The present disclosure provides methods of using polypeptides having transaminase activity for the synthesis of (S)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine in enantiomeric excess of (R)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine. Where the description relates to polypeptides, it is to be understood that it also describes the polynucleotides encoding the polypeptides.

Aminotransferases, also known as transaminases, catalyze the transfer of an amino group from a primary amine of an amino donor substrate to the carbonyl group (e.g., a keto or aldehyde group) of an amino acceptor molecule. Aminotransferases have been identified from various organisms, such as *Alcaligenes denitrificans*, *Bordetella bronchiseptica*, *Bordetella parapertussis*, *Brucella melitensis*, *Burkholderia malle*, *Burkholderia pseudomallei*, *Chromobacterium violaceum*, *Oceanicola granulosus* HTCC2516, *Oceanobacter* sp. RED65, *Oceanospirillum* sp. MED92, *Pseudomonas putida*, *Ralstonia solanacearum*, *Rhizobium meliloti*, *Rhizobium* sp. (strain NGR234), *Vibrio fluvialis*, *Bacillus thuringensis*, and *Klebsiella pneumoniae* (Shin et al., 2001, Biosci. Biotechnol. Biochem. 65:1782-1788).

Transaminases are useful for the chiral resolution of racemic amines by exploiting the ability of the transaminases to carry out the reaction in a stereospecific manner, i.e., preferential conversion of one enantiomer to the corresponding ketone, thereby resulting in a mixture enriched in the other enantiomer (see, e.g., Koselewski et al., 2009, Org

Lett. 11(21):4810-2). The stereoselectivity of transaminases in the conversion of a ketone to the corresponding amine also make these enzymes useful in the asymmetric synthesis of optically pure amines from the corresponding keto compounds (see, e.g., Höhne et al., "Biocatalytic Routes to Optically Active Amines," Chem Cat Chem 1(1):42 – 51; Zua and Hua, 2009, Biotechnol J. 4(10):1420-31).

The ω -transaminase from *Vibrio fluvialis* (ω -VfT) displays high enantioselectivity for (S)-enantiomer of chiral amines and has distinctive substrate specificity for chiral aromatic amines (Shin and Kim, 2001, J. Org. Chem. 67:2848-2853). The high enantioselectivity of ω -VfT has been applied to chiral resolution of amines (H. Yun, B.-K. Cho, B.-G. Kim, Biotechnol. Bioeng. 2004, 87, 772–778; J.-S. Shin, B.-G. Kim, Biotechnol. Bioeng. 1997, 55, 348–358; M. Hçhne, K. Robins, U. T. Bornscheuer, Adv. Synth. Catal. 2008, 350,802–807). The enzyme has also been used in the asymmetric synthesis of optically pure amines using a prochiral ketone substrate. However, limitation in asymmetric synthesis is the unfavorable equilibrium of the reverse reaction (Shin and Kim, 1999, Biotechnol. Bioeng. 65, 206–211), inhibition of ω -VfT enzyme by the amine product (Shin et al., 2001, Biotechnol Bioeng 73:179–187; Yun and Kim, 2008, Biosci. Biotechnol. Biochem. 72(11):3030-3033); low activity on amino acceptors having bulky side chains, such as aromatic groups (Shin and Kim, 2002, J. Org. Chem. 67:2848-2853); and low enzyme stability (Yun and Kim, supra).


Engineered transaminases derived from the transaminase of *Vibrio fluvialis* having increased resistance to aliphatic ketones are described in Yun et al., 2005, Appl Environ Microbiol., 71(8):4220–4224) while ω -VfTs with broadened amino donor substrate specificity are described in Cho et al., 2008, Biotechnol Bioeng. 99(2):275-84. Patent publications WO2010081053 and US20100209981, incorporated by reference herein, describe engineered ω -VfTs having increased stability to temperature and/or organic solvent and enzymatic activity towards structurally different amino acceptor molecules.

The present disclosure relates to methods of using engineered transaminase polypeptides derived from *V. fluvialis* that efficiently mediate conversion of 1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-one, to product compound (S)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine in enantiomeric excess. Significantly, the disclosure identifies amino acid residue positions and corresponding mutations in the transaminase polypeptide that increase the enzymatic activity, enantioselectivity, stability and refractoriness to the product inhibition in the conversion of 1-(1H-5-fluoro-6-chloro-indol-

3-yl)propan-2-one, to product compound (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine.

Accordingly, in one aspect, the present disclosure relates to methods of using polypeptides that are capable of converting the substrate converting substrate compound (IA), 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-one, to product compound (IIA), (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine,

Scheme 1



in the presence of an amino donor, where the (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine is produced in enantiomeric excess of compound (IIC), (R)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine.

In some embodiments, the polypeptides used in the methods are non-naturally occurring transaminases engineered for improved properties as compared to the wild-type *V. fluvialis* polypeptide of SEQ ID NO:2, or another engineered polypeptide, for example SEQ ID NO:4. These engineered transaminase polypeptides adapted for efficient conversion of conversion of 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-one to (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine have one or more residue differences as compared to the amino acid sequence of SEQ ID NO:2 or a reference engineered transaminase polypeptide, such as the reference polypeptide of SEQ ID NO:4. The residue differences are associated with enhancements in enzyme properties, including enzymatic activity, enzyme stability, and resistance to inhibition by the product amine.

In some embodiments, the engineered transaminase polypeptides used in the instant disclosure show increased activity in the conversion of the substrate 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-one to product (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine product in enantiomeric excess in a defined time with the same amount of

enzyme as compared to the wild type or a reference engineered enzyme. In some embodiments, the engineered transaminase polypeptide has at least about 1.2 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 10 fold, 20 fold, 30 fold, 40 fold, or 50 fold or more the activity as compared to the polypeptide represented by SEQ ID NO:4 under suitable reaction conditions.

In some embodiments, the engineered transaminase polypeptides used in the instant disclosure have increased stability to temperature and/or solvents used in the conversion reaction as compared to the wild type or a reference engineered enzyme. In some embodiments, the engineered transaminase polypeptide has at least 1.2 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, 10 fold or more the stability as compared to the polypeptide of SEQ ID NO:4 under suitable reaction conditions.

In some embodiments, the engineered transaminase polypeptides used in the instant disclosure have increased refractoriness or resistance to product amine (S)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine as compared to the wild type or a reference engineered enzyme. In some embodiments, the engineered transaminase polypeptide has at least 1.2 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, or more resistant to inhibition by 1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine, in particular (S)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine, as compared to the polypeptide represented by SEQ ID NO:4 under suitable reaction conditions, as further described below.

In some embodiments, the engineered transaminase polypeptides used in the instant disclosure are capable of converting the substrate 1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-one to product (S)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2-amine in enantiomeric excess of greater than 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5 or greater over (R)-1-(1H-5-fluoro-6-chloro-indol-3-yl)propan-2- under suitable reaction conditions.

In some embodiments, the engineered transaminase polypeptides used in the instant disclosure are capable of converting compound (IA) to compound (IIA) with increased tolerance for the presence of substrate relative to the reference polypeptide of SEQ ID NO: 4 under suitable reaction conditions. Thus, in some embodiments the engineered transaminase polypeptides are capable of converting the substrate compound (IIA) to product compound (IA) in the presence of a substrate loading concentration of at least about 1 g/L, about 5 g/L, about 10 g/L, about 20 g/L, about 30 g/L, about 40 g/L, about 50 g/L, about 70 g/L, about 100 g/L, about 125 g/L, about 150 g/L, about 175 g/L or

about 200 g/L or more with a percent conversion of at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 98%, or at least about 99%, in a reaction time of about 72 h or less, about 48 h or less, about 36 h or less, or about 24 h or less, under suitable reaction conditions.

The suitable reaction conditions under which the above-described improved properties of the engineered polypeptides carry out the conversion can be determined with respect to concentrations or amounts of polypeptide, substrate, cofactor, buffer, co-solvent, pH, and/or conditions including temperature and reaction time, as further described below and in the Examples.

The exemplary engineered polypeptides used in the instant disclosure have associated with their improved properties for conversion of compound **(IA)** to compound **(IIA)** and which include one or more residue differences as compared to SEQ ID NO:2 at the following residue positions: X9; X14; X18; X21; X26; X31; X33; X41; X45; X47; X57; X70; X86; X88; X107; X113; X132; X133; X146; X147; X148; X153; X163; X168; X173; X177; X203; X211; X233T; X235; X244; X250; X284; X294; X314; X315; X318; X323; X324; X324; X346; X383; X391; X395; X398; X400; X417; X419; X420; X423; X424; X427; X448; and X451. The specific amino acid differences at each of these positions that are associated with the improved properties of the exemplary polypeptides of **Table 2A** and **2B** include: X9T; X14V; X18A; X21H; X26R; X31M; X31S; X33T; X41L; X45H; X47N; X57F; X57Y; X70A; X86D; X86Y; X88A; X88L; X107P; X113L; X113V; X132F; X133R; X146L; X147K; X148Q; X148R; X153S; X163F; X163I; X163L; X163R; X163V; X168K; X168S; X173A; X177L; X203S; X211K; X233T; X235P; X244T; X250A; X284A; X294V; X314N; X315G; X318D; X323T; X324G; X324H; X346L; X383V; X391A; X395P; X398L; X398V; X398W; X400G; X417M; X419S; X420N; X423I; X424V; X424A; X427Y; X448E; and X451D.

The residue differences as compared to the engineered transaminase represented by SEQ ID NO:4 includes those at residue positions: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X113; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X420; X423; X424; X448; and X451. The specific amino acid differences at these positions include: X14V; X26R; X31S; X33T; X41L; X47N; X57F; X57Y; X70A; X86D; X88A; X88L; X107P; X113L; X113V; X132F; X148Q; X148R; X163I; X163L; X163R; X163V; X168K; X168S; X173A; X203S;

X250A; X284A; X314N; X315G; X324H; X346L; X395P; X398L; X398V; X398W; X400G; X417M; X419S; X420N; X423I; X424V; X448E; and X451D. Although residue differences compared to SEQ ID NO:4 also occur at residue positions X153 and X383, these differences represent reversions to the amino acid residue present on the wild type sequence of SEQ ID NO:2, indicating that interconversions between amino acids S and V at residue position X153 and between amino acids A and V at residue position X383 have no significant deleterious effects on the engineered enzyme properties.

The structure and function information for exemplary non-naturally occurring (or engineered) transaminase polypeptides used in the method of the present disclosure are shown below in **Table 2A** and **2B**. The odd numbered sequence identifiers (*i.e.*, SEQ ID NOs) refer to the nucleotide sequence encoding the amino acid sequence provided by the even numbered SEQ ID NOs, and the sequences are provided in the electronic sequence listing file accompanying this disclosure, which is hereby incorporated by reference herein. The amino acid residue differences are based on comparison to the reference sequence of SEQ ID NO: 4, which is an engineered transaminases derived from the wild type ω -VfT polypeptide having the following 24 amino acid residue differences A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y relative to SEQ ID NO:2. The activity of each engineered polypeptide relative to the reference polypeptide of SEQ ID NO: 4 was determined as conversion of the ketone substrate 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-one, to product amine compound (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine over a set time period and temperature in a high-throughput (HTP) assay, which was used as the primary screen. The HTP assay values in **Table 2A** were determined using *E. coli* clear cell lysates in 96 well-plate format of ~200 μ L volume per well following assay reaction conditions as noted in the table and the Examples. In some instances, a shake-flask powder (SFP) and/or downstream processed (DSP) powder assay were used as a secondary screen to assess the properties of the engineered transaminases, the results of which are provided in **Table 2B**. The SFP and DSP forms provide a more purified powder preparation of the engineered polypeptides. For example, the engineered transaminase in the SFP preparations is approximately 30% of the total protein while the DSP preparations can contain the engineered transaminases that are approximately 80% of total protein.

The levels of activity (*i.e.*, “+” “++”, etc.) are defined as follows: “+” indicates 1.2 fold or greater activity as compared to that of SEQ ID NO: 4 for engineered transaminase polypeptide SEQ ID NO: 6 to 14, and 1.2 or greater activity as compared to that of SEQ ID NO: 8 for engineered transaminase polypeptide SEQ ID NO: 16 to 154; and a “++” indicates 5 fold or greater activity as compared to that of SEQ ID NO:4 for engineered transaminase polypeptides SEQ ID NO: 6-14, and 5 fold or greater activity as compared that of SEQ ID NO:8 for engineered transaminase polypeptides SEQ ID NO: 16 to 154. The stability data is obtained by including the following amounts of product (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine in the assay and comparing the activity to that of a reference enzyme under the same conditions: 14 g/L for analysis of the engineered transaminase polypeptides of SEQ ID NO. 6 to 14, and 16 g/L for analysis of the engineered transaminases polypeptides SEQ ID NO. 16 to 154. Assessment of stability was made by comparing activities at two different temperatures, 55°C and 50°C.

Table 2A: Engineered Polypeptides and Relative Enzyme Improvements Using HTP Preparations

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^{a, b}	Stability ^b	Product Toleranc ^{c, d} _e
1/2	n/a		Nd	nd	nd
3/4	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;		"Control"	"Control"	"Control"
5/6	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	F163L;	+	+	+
7/8	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	F163I;	+	+	nd
9/10	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163R; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	F163R;	+	nd	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a _b	Stability _b	Product Toleranc ^{c, d} _e
11/12	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y86D;	+	+	+
13/14	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163V; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	F163V;	+	nd	nd
15/16	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; R88L; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; R88L; Y113L; F163L;	+	+	++
17/18	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; F163L;	Nd	+	+
19/20	A9T; Y14V; G18A; D21H; V31M; V33T; N45H; W57F; F86Y; R88L; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y; V448E;	Y14V; V33T; W57F; R88L; Y113L; F163L; V448E;	Nd	+	nd
21/22	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; R88A; Y113L; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R203S; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; R88A; Y113L; N148Q; L163I; R203S;	+	+	+
23/24	A9T; G18A; D21H; V31M; N45H; W57F; F86Y; R88L; Y113L; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y; V448E;	W57F; R88L; Y113L; N148Q; L163I; V448E;	+	+	+
25/26	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; F163L;	Nd	nd	+

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a _b	Stability _b	Product Toleranc ^{c, d} _e
27/28	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; R88A; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; R88A; N148Q; L163I;	+	+	+
29/30	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; R88L; Y113V; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; R88L; Y113V; F163L;	+	+	+
31/32	A9T; Y14V; G18A; D21H; V31M; N45H; W57F; F86Y; Y113V; A133R; R146L; W147K; N148Q; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; W57F; Y113V; N148Q; F163L;	+	+	nd
33/34	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; N148Q; L163I;	Nd	+	+
35/36	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; F163L;	Nd	+	+
37/38	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; Y113L; F163L;	Nd	+	+
39/40	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; L163I;	Nd	+	+
41/42	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; R88L; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; R88L; L163I;	Nd	+	+

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a _b	Stability _b	Product Toleranc ^{c, d} _e
43/44	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; Y113V; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; Y113V; L163I;	Nd	+	nd
45/46	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; R88L; Y113L; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; R88L; Y113L; N148Q; L163I;	Nd	+	+
47/48	A9T; Y14V; G18A; D21H; V31M; N45H; W57F; F86Y; R88L; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; W57F; R88L; N148Q; L163I;	+	+	+
49/50	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; R88L; Y113V; A133R; R146L; W147K; N148R; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; R88L; Y113V; N148R; F163L;	Nd	+	nd
51/52	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; R88A; Y113V; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; R88A; Y113V; L163I;	+	+	+
53/54	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; D250A; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; F163L; D250A;	Nd	nd	+
55/56	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; R88L; Y113L; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; R88L; Y113L; L163I;	+	+	+

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
57/58	A9T; Y14V; G18A; D21H; V31M; N45H; W57F; F86Y; R88A; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; W57F; R88A; Y113L; F163L;	Nd	nd	+
59/60	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; Y113L; A133R; R146L; W147K; N148R; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; Y113L; N148R; F163L;	Nd	+	+
61/62	A9T; Y14V; G18A; D21H; V31M; N45H; W57F; F86Y; Y113L; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; W57F; Y113L; L163I;	+	+	+
63/64	A9T; G18A; D21H; V31M; N45H; F86Y; R88L; A133R; R146L; W147K; V153S; K163I; L173A; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; A383V; T391A; S400G; L417M; C424V; F427Y;	D86Y; R88L; F163I; L173A; S400G; A424V;	+	+	nd
65/66	A9T; G18A; D21H; V31M; V33T; N45H; F86Y; R88L; A133R; R146L; W147K; V153S; K163F; L173A; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; T391A; S400G; C424A; F427Y;	V33T; D86Y; R88L; L173A; V383A; S400G;	+	+	nd
67/68	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324H; A383V; T391A; S400G; C424A; F427Y;	Y86D; G324H; S400G;	+	+	nd
69/70	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; V168K; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324H; T391A; S400G; C424A; F427Y;	Y86D; V168K; G324H; V383A; S400G;	Nd	+	nd
71/72	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; L173A; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324H; A383V; T391A;	Y86D; L173A; G324H;	Nd	+	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
	C424A; F427Y;				
73/74	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; T391A; S400G; C424A; F427Y;	Y86D; V383A; S400G;	Nd	+	+
75/76	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; L173A; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; S400G; C424A; F427Y; V448E;	Y86D; L173A; S400G; V448E;	Nd	+	+
77/78	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324H; A383V; T391A; S400G; C424A; F427Y;	Y86D; G324H; S400G;	Nd	+	nd
79/80	A9T; G18A; D21H; V31M; N45H; F86D; A133R; R146L; W147K; V153S; K163F; L173A; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324H; A383V; T391A; S400G; C424A; F427Y;	Y86D; L173A; G324H; S400G;	Nd	+	nd
81/82	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; D70A; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A383V; T391A; S398V; C424A; F427Y; E451D;	I41L; W57Y; D70A; D107P; H132F; F163L; I314N; S398V; E451D;	Nd	+	nd
83/84	A9T; G18A; D21H; V31M; I41L; N45H; R47N; W57Y; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A383V; T391A; S398V; L423I; C424A; F427Y;	I41L; R47N; W57Y; F163L; E315G; S398V; L423I;	Nd	+	nd
85/86	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A383V; T391A; S398V; C424A; F427Y;	I41L; W57Y; F163L; E315G; S398V;	+	+	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
87/88	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A383V; T391A; G395P; S398W; C424A; F427Y;	I41L; W57Y; D107P; H132F; F163L; E315G; G395P; S398W;	+	+	nd
89/90	A9T; G18A; D21H; V31M; I41L; N45H; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; S398V; C424A; F427Y; E451D;	I41L; D107P; H132F; F163L; S398V; E451D;	+	+	nd
91/92	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; E315G; P318D; A323T; S324G; A383V; T391A; S398V; L423I; C424A; F427Y; E451D;	I41L; W57Y; D107P; H132F; F163L; I314N; E315G; S398V; L423I; E451D;	+	+	nd
93/94	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A383V; T391A; S398V; L423I; C424A; F427Y;	I41L; W57Y; D107P; H132F; F163L; E315G; S398V; L423I;	Nd	+	nd
95/96	A9T; G18A; D21H; V31M; I41L; N45H; W57Y; D70A; F86Y; D107P; H132F; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A383V; T391A; S398W; Q419S; C424A; F427Y;	I41L; W57Y; D70A; D107P; H132F; F163L; I314N; S398W; Q419S;	Nd	+	nd
97/98	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A346L; A383V; T391A; G395P; S398W; C424A; F427Y; E451D;	M31S; W57F; L163I; I314N; A346L; G395P; S398W; E451D;	+	+	+
99/100	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398L; Q419S; C424A; F427Y;	M31S; W57F; L163I; E315G; A346L; S398L; Q419S;	+	+	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
101/102	A9T; G18A; D21H; V31S; I41L; N45H; W57F; F86Y; R88A; A133R; R146L; W147K; V153S; K163I; V168S; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y; E451D;	M31S; I41L; W57F; R88A; L163I; V168S; A346L; S398V; E451D;	Nd	+	nd
103/104	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V168S; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; Q419S; C424A; F427Y;	M31S; W57F; F163L; V168S; E315G; A346L; S398V; Q419S;	+	+	nd
105/106	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A383V; T391A; S398W; C424A; F427Y;	M31S; W57F; F163L; I314N; S398W;	+	+	nd
107/108	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	M31S; W57F; F163L;	+	+	nd
109/110	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V168K; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398W; C424A; F427Y; E451D;	M31S; W57F; L163I; V168K; E315G; A346L; S398W; E451D;	+	+	nd
111/112	A9T; Y14V; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y;	Y14V; M31S; W57F; F163L; I314N; A346L; S398V;	+	+	nd
113/114	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398L; C424A; F427Y;	M31S; W57F; L163I; E315G; A346L; S398L;	+	+	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
115/116	A9T; G18A; D21H; V31S; N45H; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y; E451D;	M31S; L163I; E315G; A346L; S398V; E451D;	+	+	nd
117/118	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398W; C424A; F427Y; E451D;	M31S; W57F; L163I; E315G; A346L; S398W; E451D;	+	+	nd
119/120	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y;	M31S; W57F; L163I; E315G; A346L; S398V;	188, 190, 193, 194, 196,	+	nd
121/122	A9T; G18A; D21H; V31M; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A346L; A383V; T391A; S398L; C424A; F427Y;	W57F; F163L; A346L; S398L;	Nd	+	+
123/124	A9T; G18A; D21H; V31M; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V168K; V177L; R211K; P233T; A235P; P244T; M294V; I314N; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y;	W57F; F163L; V168K; I314N; E315G; A346L; S398V;	+	+	++
125/126	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; L423I; C424A; F427Y;	M31S; W57F; L163I; E315G; A346L; S398V; L423I;	+	+	nd
127/128	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398V; Q419S; C424A; F427Y;	M31S; W57F; F163L; I314N; E315G; A346L; S398V; Q419S;	+	+	nd

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
129/130	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; A383V; T391A; S400G; L417M; S420N; C424A; F427Y;	Y14V; H26R; L163L; S284A; S400G; L417M; S420N;	Nd	nd	+
131/132	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; L173A; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; S400G; S420N; C424A; F427Y;	Y14V; F163L; L173A; S400G; S420N;	+	+	++
133/134	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; A383V; T391A; C424V; F427Y;	Y14V; Y113L; F163L; S284A; A424V;	+	+	++
135/136	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; R88L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324H; A383V; T391A; S400G; L417M; C424A; F427Y;	Y14V; R88L; F163L; G324H; S400G; L417M;	+	+	nd
137/138	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; H26R; W57F; Y113L; F163L; S284A;	Nd	nd	+
139/140	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; T391A; S400G; C424A; F427Y;	Y14V; H26R; F163L; S284A; V383A; S400G;	Nd	nd	++
141/142	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; T391A; C424V; F427Y;	Y14V; H26R; Y113L; F163L; V383A; A424V;	+	+	nd
143/144	A9T; G18A; D21H; V31M; N45H; F86Y; R88L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A;	R88L; F163L;	+	+	#VALUE!

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a _b	Stability _b	Product Toleranc _{c, d} ^e
	F427Y;				
145/146	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; T391A; S400G; S420N; C424A; F427Y;	Y14V; H26R; N148Q; L163I; S284A; V383A; S400G; S420N;	Nd	+	++
147/148	A9T; Y14V; G18A; D21H; V31M; V33T; N45H; W57F; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	Y14V; V33T; W57F; Y113L; F163L;	+	+	nd
149/150	A9T; Y14V; G18A; D21H; V31M; N45H; W57F; F86Y; R88L; Y113L; A133R; R146L; W147K; N148Q; V153S; K163I; L173A; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; T391A; S400G; C424A; F427Y;	Y14V; W57F; R88L; Y113L; N148Q; L163I; L173A; V383A; S400G;	Nd	+	nd
151/152	A9T; G18A; D21H; V31M; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	W57F; F163L;	Nd	+	+
153/154	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324H; A383V; T391A; S400G; C424A; F427Y;	Y14V; H26R; F163L; S284A; G324H; S400G;	+	+	nd

^a HTP Assay Condition 1: Cells grown in 96 well plates were lysed with 200 μ L of Lysis Buffer (1 mg/mL lysozyme, 0.5 mg/mL polymyxin B sulfate, 1 mM PLP, 0.1 M triethanolamine (TEA), pH 7.0). The reaction conditions comprised: 10 g/L (44.4 mM) Compound (IA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 100 mM triethanolamine (TEA), pH 7.0; 5% PEG 200 v/v; 10 μ L Lysate; and 50°C for 24 h.

^b HTP Assay Condition 2: Cells grown in 96 well plates were lysed with 400 μ L of Lysis Buffer (1 mg/mL lysozyme, 0.5 mg/mL polymyxin B sulfate, 1 mM PLP, 0.1 M triethanolamine (TEA), pH 7.0). The reaction conditions comprised: 10 g/L (44.4 mM) Compound (IA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 100 mM triethanolamine (TEA), pH 7.0; 5% PEG 200 v/v; 10 μ L Lysate; and

SEQ ID NO:	Active Mutations (relative to SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Activity ^a , b	Stability ^b	Product Toleranc ^{c, d} e
50°C for 24 h.					
^c Stability Assay conditions: 10 g/L (44.4 mM) Compound (IA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 100 mM triethanolamine (TEA), pH 7.0; 5% PEG 200 v/v; 10 μ L Lysate (prepared according to HTP Assay Condition 1 or 2); and 55°C for 24 h. Lysates were prepared according to HTP Assay Condition 1 (for SEQ ID NO: 4-90) or HTP Assay Condition 2 (for SEQ ID NO:92-230).					
^d Product Inhibition Assay Condition 1: 10 g/L (44.4 mM) Compound (IA); 14 g/L Compound (IIA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 100 mM triethanolamine (TEA), pH 7.0; 5% PEG 200 v/v; 10 μ L Lysate; and 50°C. Lysates were prepared according to HTP Assay Condition 1.					
^e Product Inhibition Assay Condition 2: 10 g/L (44.4 mM) Compound (IA); 16 g/L Compound (IIA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 100 mM triethanolamine (TEA), pH 7.0; 5% PEG 200 v/v; 10 μ L Lysate; and 50°C. Lysates were prepared according to HTP Assay Condition 2.					
nd: not determined					

Table 2B: Engineered Polypeptides and Relative Enzyme Improvements Using Shake Flask and DSP Preparations

SEQ ID NO:	Active Mutations (relative to Wild-type of SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Shake Flask Activity ^a		DSP Activity ^b	
			% ee		% ee	
1/2	n/a		nd	Nd	nd	nd
3/4	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y		"Control"	"Control"	"Control"	"Control"
5/6	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y;	F163L	99.7	+	nd	nd
7/8	A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T;	F163I	99.5	+	nd	nd

SEQ ID NO:	Active Mutations (relative to Wild-type of SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Shake Flask		DSP	
			% ee	Activity ^a	% ee	Activity ^b
	A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y					
45/46	A9T; Y14V; G18A; D21H; H26R; V31M; N45H; W57F; F86Y; R88L; Y113L; A133R; R146L; W147K; N148Q; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y	Y14V; H26R; W57F; R88L; Y113L; N148Q; L163I	99.6	+	99.4	+
99/100	A9T; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163I; V177L; R211K; P233T; A235P; P244T; M294V; E315G; P318D; A323T; S324G; A346L; A383V; T391A; S398L; Q419S; C424A; F427Y	M31S; W57F; L163I; E315G; A346L; S398L; Q419S	99.8	+	nd	nd
111/112	A9T; Y14V; G18A; D21H; V31S; N45H; W57F; F86Y; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; I314N; P318D; A323T; S324G; A346L; A383V; T391A; S398V; C424A; F427Y	Y14V; M31S; W57F; F163L; I314N; A346L; S398V	99.7	+	nd	nd
133/134	A9T; Y14V; G18A; D21H; V31M; N45H; F86Y; Y113L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; S284A; M294V; P318D; A323T; S324G; A383V; T391A; C424V; F427Y	Y14V; Y113L; F163L; S284A; A424V	nd	Nd	99.8	+
143/144	A9T; G18A; D21H; V31M; N45H; F86Y; R88L; A133R; R146L; W147K; V153S; K163L; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y	R88L; F163L	99	Nd	nd	nd

SEQ ID NO:	Active Mutations (relative to Wild-type of SEQ ID NO: 2)	Active Mutations (relative to SEQ ID NO: 4)	Shake Flask		DSP	
			Activity		Activity	
			% ee	^a	% ee	^b
^a Shake Flask Assay Conditions: 25 g/L (or 50 or 100 g/L) Compound (IA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 5% v/v PEG200; 100 mM triethanolamine (TEA), pH 7.0; 2 g/L protein of transaminase-containing shake flask preparation; and 50°C for a reaction time of 24 h.						
^b DSP Assay Conditions: 25 g/L (or 50 or 100 g/L) Compound (IA); 1 mM pyridoxal-5-phosphate (PLP); 2 M isopropylamine (IPM), pH 7; 5% v/v PEG200; 100 mM triethanolamine (TEA), pH 7; 2 g/L protein from transaminase-containing DSP preparation; and 50°C for a reaction time of 24 h.						

From an inspection of the exemplary polypeptides useful in the methods of the present invention, improvements in enzyme properties are associated with residue differences as compared to SEQ ID NO:4 at residue positions X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X113; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X420; X423; X424; X448; and X451. The specific residue differences at each of these positions that are associated with the improved properties include: X14V; X26R; X31S; X33T; X41L; X47N; X57F; X57Y; X70A; X86D; X88A; X88L; X107P; X113L; X113V; X132F; X148Q; X148R; X163I; X163L; X163R; X163V; X168K; X168S; X173A; X203S; X250A; X284A; X314N; X315G; X324H; X346L; X395P; X398L; X398V; X398W; X400G; X417M; X419S; X420N; X423I; X424V; X448E; and X451D.

Accordingly, in some embodiments, the engineered transaminase polypeptide useful in the methods of the instant disclosure are capable of converting substrate Compound (IA), 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-one, to the product Compound (IIA), (S)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine with improved properties as compared to SEQ ID NO:4, comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to reference sequence SEQ ID NO:2 and one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, where the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M.

In some embodiments, the engineered transaminase polypeptide useful in the instantly disclosed methods is capable of converting substrate Compound (IA) to the product Compound (IIA) with improved properties as compared to SEQ ID NO:4, comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to reference sequence SEQ ID NO:4 and one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, wherein the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M. In some embodiments, the engineered transaminase polypeptides are capable of carrying out the conversion with the enantioselectivities described above, *e.g.*, $\geq 90\%$ ee.

In some embodiments, the transaminase polypeptide useful in the instantly disclosed methods is capable of converting substrate Compound (IA) to product Compound (IIA) with improved properties as compared to SEQ ID NO:4, comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity to a reference sequence selected from SEQ ID NO: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154, and one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, where the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M. In some embodiments, the reference sequence is selected from SEQ ID NO: 4, 8, 14, 16, 132, 134, and 146. In some embodiments, the reference sequence is SEQ ID NO:4. In some embodiments, the reference sequence is SEQ ID NO:8. In some embodiments, the reference sequence is SEQ ID NO:134. In some embodiments, the reference sequence is SEQ ID NO:146.

In some embodiments, the transaminase polypeptide capable of converting substrate Compound (IA) to product Compound (IIA) with improved properties as compared to SEQ ID NO:4, comprises an amino acid sequence selected from SEQ ID NO: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154, and having one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, where the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M. In some embodiments, the amino acid sequence is selected from SEQ ID NO: 4, 8, 14, 16, 132, 134, and 146. In some embodiments, the amino acid sequence is SEQ ID NO:4. In some embodiments, the amino acid sequence is SEQ ID NO:8. In some embodiments, the amino acid sequence is SEQ ID NO:134. In some embodiments, the amino acid sequence is SEQ ID NO:146.

In some embodiments, the engineered transaminase polypeptide is capable of converting the substrate Compound (IA) to the product Compound (IIA) with at least 1.2 fold the activity relative to SEQ ID NO:4 comprises an amino acid sequence selected from: SEQ ID NO: 6, 8, 10, 12, 14, 16, 22, 24, 28, 30, 32, 48, 52, 56, 62, 64, 66, 68, 86, 88, 90, 92, 98, 100, 104, 106, 108, 110, 112, 114, 116, 118, 124, 126, 128, 132, 134, 136, 142, 144, 148, and 154.

In some embodiments, the engineered transaminase polypeptide is capable of converting the substrate Compound (IA) to the product Compound (IIA) with at least 5 fold the activity relative to SEQ ID NO:4 and comprises an amino acid sequence having one or more residue differences selected from: X14V, X26R; X33T; X88A/L; X163I/L; and X284A.

In some embodiments, the engineered transaminase polypeptide useful in the instantly disclosed methods is capable of converting the substrate Compound (IA) to the product Compound (IIA) with at least 5 fold the activity relative to SEQ ID NO:4 comprises an amino acid sequence selected from: SEQ ID NO: 6, 8, 10, 14, 16, 22, 24, 28, 30, 32, 48,

52, 56, 62, 64, 86, 88, 90, 92, 98, 100, 104, 106, 108, 110, 112, 114, 116, 118, 124, 126, 128, 132, 134, 136, 142, 144, 148, and 154.

In some embodiments, the engineered transaminase polypeptide has at least 1.2 fold, 1.5 fold, 2 fold, 3 fold, 4 fold, 5 fold, or more refractoriness to inhibition by 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine, in particular (*S*)-1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine, as compared to the polypeptide represented by SEQ ID NO:4 in the conversion of Compound (IA) to Compound (IIA). Generally, the increase in refractoriness or resistance to inhibition by the product compound can be measured under HTP assay conditions in presence of 14 g/L or 16 g/L of Compound (IIA), as described in **Table 2A** and **2B** and the Examples. In some embodiments, the engineered transaminase polypeptide having at least 1.2 fold or greater refractoriness to inhibition by 1-(1*H*-5-fluoro-6-chloro-indol-3-yl)propan-2-amine comprises an amino acid sequence having one or more residue differences as compared to SEQ ID NO:4 selected from: X26R; X70A; X86D; X88A/L; X132F; X163L; X315G; X395P; X398L; and X419S.

In some embodiments, the engineered transaminase polypeptide with improved properties in the conversion of Compound (IA) to Compound (IIA) has an amino acid sequence comprising a sequence selected from SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154.

In some embodiments, the engineered transaminase capable of converting Compound (IA) to Compound (IIA) under suitable reaction conditions, comprises an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% identity to one of SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154, and the amino acid residue differences as compared to SEQ ID NO:4 present in any one of SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112,

114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154, as provided in **Tables 2A** and **2B**.

In some embodiments, the engineered polypeptides can be in various forms, for example, such as an isolated preparation, as a substantially purified enzyme, whole cells transformed with gene(s) encoding the enzyme, and/or as cell extracts and/or lysates of such cells. The enzymes can be lyophilized, spray-dried, precipitated or be in the form of a crude paste, as further discussed below.

In some embodiments, the transaminase polypeptides can be bound on a physical substrate. The transaminase polypeptide can be bound non-covalently or covalently on a physical substrate such that they retain their improved activity, stereoselectivity, product tolerance, stability, and/or other improved properties relative to the reference polypeptide of SEQ ID NO: 4. In such embodiments, the immobilized polypeptides can facilitate the biocatalytic conversion of the suitable ketone substrates, e.g., Compound (IA) or structural analogs thereof, to the corresponding amine product, e.g., Compound (IIA) or corresponding structural analogs, and after the reaction is complete are easily retained (e.g., by retaining beads on which polypeptide is immobilized) and then reused or recycled in subsequent reactions. Such immobilized enzyme processes allow for further efficiency and cost reduction. Methods of enzyme immobilization are well-known in the art. Various methods for conjugation to substrates, e.g., membranes, beads, glass, etc. are described in, among others, Hermanson, G.T., *Bioconjugate Techniques*, Second Edition, Academic Press; (2008), and *Bioconjugation Protocols: Strategies and Methods*, In *Methods in Molecular Biology*, C.M. Niemeyer ed., Humana Press (2004); the disclosures of which are incorporated herein by reference.

Polynucleotides Encoding Engineered Transaminases, Expression Vectors and Host Cells

In another aspect, the present disclosure provides polynucleotides encoding the engineered transaminase polypeptides described herein. The polynucleotides may be operatively linked to one or more heterologous regulatory sequences that control gene expression to create a recombinant polynucleotide capable of expressing the polypeptide. Expression constructs containing a heterologous polynucleotide encoding the engineered transaminase can be introduced into appropriate host cells to express the corresponding transaminase polypeptide.

As will be apparent to the skilled artisan, availability of a protein sequence and the knowledge of the codons corresponding to the various amino acids provide a description of all the polynucleotides capable of encoding the subject polypeptides. The degeneracy of the genetic code, where the same amino acids are encoded by alternative or synonymous codons, allows an extremely large number of nucleic acids to be made, all of which encode the improved transaminase enzymes. Thus, having knowledge of a particular amino acid sequence, those skilled in the art could make any number of different nucleic acids by simply modifying the sequence of one or more codons in a way which does not change the amino acid sequence of the protein. In this regard, the present disclosure specifically contemplates each and every possible variation of polynucleotides that could be made encoding the polypeptides described herein by selecting combinations based on the possible codon choices, and all such variations are to be considered specifically disclosed for any polypeptide described herein, including the amino acid sequences presented in **Tables 2A** and **2B**, and disclosed in the sequence listing incorporated by reference herein as SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154.

In various embodiments, the codons are preferably selected to fit the host cell in which the protein is being produced. For example, preferred codons used in bacteria are used to express the gene in bacteria; preferred codons used in yeast are used for expression in yeast; and preferred codons used in mammals are used for expression in mammalian cells. In some embodiments, all codons need not be replaced to optimize the codon usage of the transaminases since the natural sequence will comprise preferred codons and because use of preferred codons may not be required for all amino acid residues. Consequently, codon optimized polynucleotides encoding the transaminase enzymes may contain preferred codons at about 40%, 50%, 60%, 70%, 80%, or greater than 90% of codon positions of the full length coding region.

An isolated polynucleotide encoding an improved transaminase polypeptide may be manipulated in a variety of ways to provide for expression of the polypeptide. In some embodiments, the polynucleotides encoding the polypeptides can be provided as expression vectors where one or more control sequences is present to regulate the

expression of the polynucleotides and/or polypeptides. Manipulation of the isolated polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides and nucleic acid sequences utilizing recombinant DNA methods are well known in the art. Guidance is provided in Sambrook et al., 2001, *Molecular Cloning: A Laboratory Manual*, 3rd Ed., Cold Spring Harbor Laboratory Press; and *Current Protocols in Molecular Biology*, Ausubel, F. ed., Greene Pub. Associates, 1998, updates to 2006.

In the embodiments herein, the improved polypeptides and corresponding polynucleotides can be obtained using methods used by those skilled in the art. The parental polynucleotide sequence encoding the wild-type polypeptide of *Vibrio fluvialis* is described in Shin *et al.*, 2003, *Appl. Microbiol. Biotechnol.* 61(5-6):463-471, and methods of generating engineered transaminase polypeptides with improved stability and substrate recognition properties are disclosed in patent application publications WO2010081053 and US20100209981, incorporated herein by reference.

Where the sequence of the engineered polypeptide is known, the polynucleotides encoding the enzyme can be prepared by standard solid-phase methods, according to known synthetic methods. In some embodiments, fragments of up to about 100 bases can be individually synthesized, then joined (*e.g.*, by enzymatic or chemical ligation methods, or polymerase mediated methods) to form any desired continuous sequence. For example, polynucleotides and oligonucleotides of the invention can be prepared by chemical synthesis using, *e.g.*, the classical phosphoramidite method described by Beaucage et al., 1981, *Tet Lett* 22:1859-69, or the method described by Matthes et al., 1984, *EMBO J.* 3:801-05, *e.g.*, as it is typically practiced in automated synthetic methods. According to the phosphoramidite method, oligonucleotides are synthesized, *e.g.*, in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors. In addition, essentially any nucleic acid can be obtained from any of a variety of commercial sources.

Accordingly, in some embodiments, a method for preparing the engineered transaminases polypeptide can comprise: (a) synthesizing a polynucleotide encoding a polypeptide comprising an amino acid sequence selected from SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136,

138, 140, 142, 144, 146, 148, 150, 152, and 154 and having one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168; X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, wherein the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M.; and (b) expressing the transaminase polypeptide encoded by the polynucleotide.

The expressed engineered transaminase can be measured for the desired improved property, *e.g.*, activity, enantioselectivity, stability, and product tolerance, in the conversion of compound (IA) to compound (IIA) by any of the assay conditions described herein.

In some embodiments, any of the engineered transaminase enzymes expressed in a host cell can be recovered from the cells and or the culture medium using any one or more of the well known techniques for protein purification, including, among others, lysozyme treatment, sonication, filtration, salting-out, ultra-centrifugation, and chromatography. Suitable solutions for lysing and the high efficiency extraction of proteins from bacteria, such as *E. coli*, are commercially available under the trade name CellLytic B™ from Sigma-Aldrich of St. Louis MO.

Chromatographic techniques for isolation of the transaminase polypeptide include, among others, reverse phase chromatography high performance liquid chromatography, ion exchange chromatography, gel electrophoresis, and affinity chromatography. Conditions for purifying a particular enzyme will depend, in part, on factors such as net charge, hydrophobicity, hydrophilicity, molecular weight, molecular shape, etc., and will be apparent to those having skill in the art.

In some embodiments, affinity techniques may be used to isolate the improved transaminase enzymes. For affinity chromatography purification, any antibody which specifically binds the transaminase polypeptide may be used. For the production of antibodies, various host animals, including but not limited to rabbits, mice, rats, etc., may be immunized by injection with a transaminase polypeptide, or a fragment thereof. The transaminase polypeptide or fragment may be attached to a suitable carrier, such as BSA, by means of a side chain functional group or linkers attached to a side chain functional group. Various adjuvants may be used to increase the immunological

response, depending on the host species, including but not limited to Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, dinitrophenol, and potentially useful human adjuvants such as BCG (bacilli Calmette Guerin) and *Corynebacterium parvum*.

Methods of Using the Engineered Transaminase Enzymes

In another aspect, the transaminases described herein can be used in a process for carrying out transaminase reactions in which an amino group from an amino donor is transferred to an amino acceptor, e.g., ketone substrate, to produce an amine. Use of a prochiral ketone acceptor can result in the production of a chiral amine in enantiomeric excess. Generally, the process for performing the transamination reaction comprises contacting or incubating an amino donor and an amino acceptor with an engineered transaminase polypeptide of the disclosure under reaction conditions suitable for converting the amino acceptor to an amine.

In some embodiments, the transaminases can be used in the conversion of substrate compound of formula (I) to product compound of formula (II). Accordingly, in some embodiments, a process for preparing the compound of formula (II) in enantiomeric excess comprises contacting the compound of formula (I) in the presence of an amino donor under suitable reaction conditions with an engineered transaminase polypeptide described herein. In some embodiments of the process, the compound of formula (II) can be formed in at least 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater enantiomeric excess.

For the foregoing processes, any of the engineered transaminases described herein can be used. By way of example and without limitation, in some embodiments, the process can use an engineered transaminases polypeptide comprising an amino acid sequence having at least 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity to a reference sequence selected from SEQ ID NO: 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154, and one or more residue differences as compared to SEQ ID NO:4 at residue positions selected from: X14; X26; X31; X33; X41; X47; X57; X70; X86; X88; X107; X132; X148; X163; X168;

X173; X203; X250; X284; X314; X315; X324; X346; X395; X398; X400; X417; X419; X423; X448; and X451, wherein the residue differences at residue positions X31; X57; X86; X163; X168; X314; X324; X398; and X417 are selected from: X31S; X57Y; X86D; X163I; X163L; X163R; X163V; X168S; X314N; X324H; X398L; X398V; X398W; and X417M. In some embodiments, the reference sequence is selected from SEQ ID NO: 4, 8, 14, 16, 132, 134, and 146. In some embodiments, the reference sequence is SEQ ID NO:4. In some embodiments, the reference sequence is SEQ ID NO:8. In some embodiments, the reference sequence is SEQ ID NO:134. In some embodiments, the reference sequence is SEQ ID NO:146.

In some embodiments, exemplary transaminases capable of carrying out the processes herein can be a polypeptide comprising an amino acid sequence selected from SEQ ID NO: 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, 44, 46, 48, 50, 52, 54, 56, 58, 60, 62, 64, 66, 68, 70, 72, 74, 76, 78, 80, 82, 84, 86, 88, 90, 92, 94, 96, 98, 100, 102, 104, 106, 108, 110, 112, 114, 116, 118, 120, 122, 124, 126, 128, 130, 132, 134, 136, 138, 140, 142, 144, 146, 148, 150, 152, and 154. Guidance on the choice and use of the engineered transaminases is provided in the descriptions herein, for example **Table 2** and the Examples.

In the embodiments herein and illustrated in the Examples, various ranges of suitable reaction conditions that can be used in the processes, including but not limited, to ranges of amino donor, pH, temperature, buffer, solvent system, substrate loading, polypeptide loading, cofactor loading, pressure, and reaction time. Further suitable reaction conditions for carrying out the process for biocatalytic conversion of substrate compounds to product compounds using an engineered transaminase polypeptide described herein can be readily optimized in view of the guidance provided herein by routine experimentation that includes, but is not limited to, contacting the engineered transaminase polypeptide and substrate compound under experimental reaction conditions of concentration, pH, temperature, solvent conditions, and detecting the product compound.

In the embodiments described herein, the transaminase polypeptide uses an amino donor to form the product compounds. In some embodiments, the amino donor in the reaction condition can be selected from isopropylamine (also referred to herein as "IPM"), putrescine, L-lysine, α -phenethylamine, D-alanine, L-alanine, or D,L-alanine, or D,L-ornithine. In some embodiments, the amino donor is selected from the group

consisting of IPM, putrescine, L-lysine, D- or L-alanine. In some embodiments, the amino donor is IPM. In some embodiments, the suitable reaction conditions comprise the amino donor, in particular IPM, present at a concentration of at least about 0.1 to about 3.0 M, 0.2 to about 2.5 M, about 0.5 to about 2 M or about 1 to about 2 M. In some embodiments, the amino donor is present at a concentration of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 1, 1.5, 2, 2.5 or 3 M. Higher concentrations of amino donor, *e.g.*, IPM, can be used to shift the equilibrium towards amine product formation.

Suitable reaction conditions using the engineered transaminase polypeptides also typically comprise a cofactor. Cofactors useful in the processes using the engineered transaminase enzymes include, but are not limited to, pyridoxal-5'-phosphate (also known as pyridoxal-phosphate, PLP, P5P), pyridoxine (PN), pyridoxal (PL), pyridoxamine (PM), and their phosphorylated counterparts; pyridoxine phosphate (PNP), and pyridoxamine phosphate (PMP). In some embodiments, the cofactor PLP is present naturally in the cell extract and does not need to be supplemented. In some embodiments of the methods, the suitable reaction conditions comprise cofactor added to the enzyme reaction mixture, for example, when using partially purified, or purified transaminase enzyme. In some embodiments, the suitable reaction conditions can comprise the presence of a cofactor selected from PLP, PN, PL, PM, PNP, and PMP, at a concentration of about 0.1 g/L to about 10 g/L, about 0.2 g/L to about 5 g/L, about 0.5 g/L to about 2.5 g/L. In some embodiments, the reaction conditions comprise a PLP concentration of about 0.1 g/L or less, 0.2 g/L or less, 0.5 g/L or less, 1 g/L or less, 2.5 g/L or less, 5 g/L or less, or 10 g/L or less. In some embodiments, the cofactor can be added either at the beginning of the reaction and/or additional cofactor is added during the reaction.

Substrate compound in the reaction mixtures can be varied, taking into consideration, for example, the desired amount of product compound, the effect of substrate concentration on enzyme activity, stability of enzyme under reaction conditions, and the percent conversion of substrate to product. In some embodiments, the suitable reaction conditions comprise a substrate compound loading of at least about 0.5 to about 200 g/L, 1 to about 200 g/L, 5 to about 150 g/L, about 10 to about 100 g/L, 20 to about 100 g/L or about 50 to about 100 g/L. In some embodiments, the suitable reaction conditions comprise a substrate compound loading of at least about 0.5 g/L, at least about 1 g/L, at least about 5 g/L, at least about 10 g/L, at least about 15 g/L, at least about 20 g/L, at

least about 30 g/L, at least about 50 g/L, at least about 75 g/L, at least about 100 g/L, at least about 150 g/L or at least about 200 g/L, or even greater.

The improved activity and/or stereoselectivity of the engineered transaminase polypeptides disclosed herein provides for processes wherein higher percentage conversion can be achieved with lower concentrations of the engineered polypeptide. In some embodiments of the process, the suitable reaction conditions comprise an engineered polypeptide concentration of about 0.01 to about 50 g/L; about 0.05 to about 50 g/L; about 0.1 to about 40 g/L; about 1 to about 40 g/L; about 2 to about 40 g/L; about 5 to about 40 g/L; about 5 to about 30 g/L; about 0.1 to about 10 g/L; about 0.5 to about 10 g/L; about 1 to about 10 g/L; about 0.1 to about 5 g/L; about 0.5 to about 5 g/L; or about 0.1 to about 2 g/L. In some embodiments, the transaminase polypeptide is concentration at about 0.01, 0.05, 0.1, 0.2, 0.5, 1, 2, 5, 10, 15, 20, 25, 30, 35, 40, or 50 g/L.

During the course of the transamination reactions, the pH of the reaction mixture may change. The pH of the reaction mixture may be maintained at a desired pH or within a desired pH range. This may be done by the addition of an acid or a base, before and/or during the course of the reaction. Alternatively, the pH may be controlled by using a buffer. Accordingly, in some embodiments, the reaction condition comprises a buffer. Suitable buffers to maintain desired pH ranges are known in the art and include, by way of example and not limitation, borate, carbonate, phosphate, triethanolamine buffer, and the like. In some embodiments, the buffer is borate. In some embodiments of the process, the suitable reaction conditions comprise a buffer solution of triethanolamine, where the triethanolamine concentration is from about 0.01 to about 0.4 M, 0.05 to about 0.4 M, 0.1 to about 0.3 M, or about 0.1 to about 0.2 M. In some embodiments, the reaction condition comprises a triethanolamine concentration of about 0.01, 0.02, 0.03, 0.04, 0.05, 0.07, 0.1, 0.12, 0.14, 0.16, 0.18, 0.2, 0.3, or 0.4 M. In some embodiments, the reaction conditions comprise water as a suitable solvent with no buffer present.

In the embodiments of the process, the reaction conditions can comprise a suitable pH. The desired pH or desired pH range can be maintained by use of an acid or base, an appropriate buffer, or a combination of buffering and acid or base addition. The pH of the reaction mixture can be controlled before and/or during the course of the reaction. In some embodiments, the suitable reaction conditions comprise a solution pH comprise a pH from about 6 to about 12, pH from about 6 to about 10, pH from about 6 to about 8,

pH from about 7 to about 10, pH from about 7 to about 9, or pH from about 7 to about 8. In some embodiments, the reaction conditions comprise a solution pH of about 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 11.5 or 12.

In the embodiments of the processes herein, a suitable temperature can be used for the reaction conditions, for example, taking into consideration the increase in reaction rate at higher temperatures, and the activity of the enzyme during the reaction time period. For example, the engineered polypeptides of the present disclosure have increased stability relative to naturally occurring transaminase polypeptide *e.g.*, the wild type polypeptide of SEQ ID NO: 2, which allow the engineered polypeptides to be used at higher temperatures for increased conversion rates and improved substrate solubility characteristics for the reaction. Accordingly, in some embodiments, the suitable reaction conditions comprise a temperature of about 10°C to about 70°C, about 10°C to about 65°C, about 15°C to about 60°C, about 20°C to about 60°C, about 20°C to about 55°C, about 30°C to about 55°C, or about 40°C to about 50°C. In some embodiments, the suitable reaction conditions comprise a temperature of about 10°C, 15°C, 20°C, 25°C, 30°C, 35°C, 40°C, 45°C, 50°C, 55°C, 60°C, 65°C, or 70°C. In some embodiments, the temperature during the enzymatic reaction can be maintained at a temperature throughout the course of the reaction. In some embodiments, the temperature during the enzymatic reaction can be adjusted over a temperature profile during the course of the reaction.

The processes of the disclosure are generally carried out in a solvent. Suitable solvents include water, aqueous buffer solutions, organic solvents, polymeric solvents, and/or co-solvent systems, which generally comprise aqueous solvents, organic solvents and/or polymeric solvents. The aqueous solvent (water or aqueous co-solvent system) may be pH-buffered or unbuffered. In some embodiments, the processes using the engineered transaminase polypeptides are generally carried out in an aqueous co-solvent system comprising an organic solvent (*e.g.*, ethanol, isopropanol (IPA), dimethyl sulfoxide (DMSO), ethyl acetate, butyl acetate, 1-octanol, heptane, octane, methyl t butyl ether (MTBE), toluene, and the like), ionic or polar solvents (*e.g.*, 1 ethyl 4 methylimidazolium tetrafluoroborate, 1 butyl 3 methylimidazolium tetrafluoroborate, 1 butyl 3 methylimidazolium hexafluorophosphate, glycerol, polyethylene glycol, and the like). In some embodiments, the co-solvent can be a polar solvent, such as a polyol, dimethylsulfoxide, DMSO, or lower alcohol. The non-aqueous co-solvent component of

an aqueous co-solvent system may be miscible with the aqueous component, providing a single liquid phase, or may be partly miscible or immiscible with the aqueous component, providing two liquid phases. Exemplary aqueous co-solvent systems can comprise water and one or more co-solvents selected from an organic solvent, polar solvent, and polyol solvent. In general, the co-solvent component of an aqueous co-solvent system is chosen such that it does not adversely inactivate the transaminase enzyme under the reaction conditions. Appropriate co-solvent systems can be readily identified by measuring the enzymatic activity of the specified engineered transaminase enzyme with a defined substrate of interest in the candidate solvent system, utilizing an enzyme activity assay, such as those described herein.

In some embodiments of the process, the suitable reaction conditions comprise an aqueous co-solvent, where the co-solvent comprises a polyol solvent, particularly glycols. Examples of suitable polyol solvents include, by way of example and not limitation, polyethylene glycol, polyethylene glycol methyl ether, diethylene glycol dimethyl ether, triethylene glycol dimethyl ether, and polypropylene glycol. In some embodiments, the aqueous co-solvent comprises polyethylene glycol, which is available in different molecular weights. Particularly useful are lower molecular weight glycols, such as PEG200 to PEG600. Accordingly, in some embodiments, the aqueous co-solvent comprises PEG200 of about 1% to about 40% v/v; about 1% to about 40% v/v; about 2% to about 40% v/v; about 5% to about 40% v/v; 2% to about 30% v/v; 5% to about 30% v/v; 1 to about 20% v/v; about 2% to about 20% v/v; about 5% to about 20% v/v; about 1% to about 10% v/v; about 2% to about 10% v/v. In some embodiments, the suitable reaction conditions comprises an aqueous co-solvent comprising PEG200 at about 1%, 2%, 5%, 10%, 15%, 20%; 25%; 30%; 35%; 35% or about 40% v/v.

In some embodiments of the process, the suitable reaction conditions comprise an aqueous co-solvent, where the co-solvent comprises DMSO at about 1% to about 80% (v/v), about 1 to about 70% (v/v), about 2% to about 60% (v/v), about 5% to about 40% (v/v), 10% to about 40% (v/v), 10% to about 30% (v/v), or about 10% to about 20% (v/v). In some embodiments of the process, the suitable reaction conditions comprise an aqueous co-solvent comprising DMSO at least about 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, or 80% (v/v).

The quantities of reactants used in the transamination reaction will generally vary depending on the quantities of product desired, and concomitantly the amount of

transaminase substrate employed. Those having ordinary skill in the art will readily understand how to vary these quantities to tailor them to the desired level of productivity and scale of production.

In some embodiments, the order of addition of reactants is not critical. The reactants may be added together at the same time to a solvent (*e.g.*, monophasic solvent, biphasic aqueous co-solvent system, and the like), or alternatively, some of the reactants may be added separately, and some together at different time points. For example, the cofactor, transaminase, and transaminase substrate may be added first to the solvent.

The solid reactants (*e.g.*, enzyme, salts, etc.) may be provided to the reaction in a variety of different forms, including powder (*e.g.*, lyophilized, spray dried, and the like), solution, emulsion, suspension, and the like. The reactants can be readily lyophilized or spray dried using methods and equipment that are known to those having ordinary skill in the art. For example, the protein solution can be frozen at -80°C in small aliquots, then added to a pre-chilled lyophilization chamber, followed by the application of a vacuum.

For improved mixing efficiency when an aqueous co-solvent system is used, the transaminase, and cofactor may be added and mixed into the aqueous phase first. The organic phase may then be added and mixed in, followed by addition of the transaminase substrate. Alternatively, the transaminase substrate may be premixed in the organic phase, prior to addition to the aqueous phase.

The transamination reaction is generally allowed to proceed until further conversion of ketone substrate to amine product does not change significantly with reaction time, *e.g.*, less than 10% of substrate being converted, or less than 5% of substrate being converted). In some embodiments, the reaction is allowed to proceed until there is complete or near complete conversion of substrate ketone to product amine.

Transformation of substrate to product can be monitored using known methods by detecting substrate and/or product. Suitable methods include gas chromatography, HPLC, and the like. Conversion yields of the chiral amine product generated in the reaction mixture are generally greater than about 50%, may also be greater than about 60%, may also be greater than about 70%, may also be greater than about 80%, may also be greater than 90%, and are often greater than about 97%. In some embodiments, the methods for preparing compounds of formula (II) using an engineered transaminase polypeptide under suitable reaction conditions results in at least about 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater conversion of ketone substrate, *e.g.*,

compound of formula (I), to the amine product compound, e.g., compound of formula (II) in about 48h or less, in about 36 h or less, in about 24 h or less, or even less time.

In some embodiments of the process, the suitable reaction conditions comprise a substrate loading of substrate compound of at least about 20 g/L, 30 g/L, 40 g/L, 50 g/L, 60 g/L, 70 g/L, 100 g/L, or more, and wherein the method results in at least about 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater conversion of substrate compound to product compound in about 48h or less, in about 36 h or less, or in about 24 h or less.

The engineered transaminase polypeptides of the present disclosure when used in the process under suitable reaction conditions result in an enantiomeric excess of the chiral amine in at least 97%, 98, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or 99.9%. e.e.

In a further embodiment of the methods for converting substrate compound to amine product compound using the engineered transaminase polypeptides, the suitable reaction conditions comprise an initial substrate loading to the reaction solution which is then contacted by the polypeptide. This reaction solution is then further supplemented with additional substrate of compound as a continuous addition over time at a rate of at least about 1 g/L/h, at least about 2 g/L/h, at least about 4 g/L/h, at least about 6 g/L/h, or higher. Thus, according to these suitable reaction conditions polypeptide is added to a solution having an initial substrate loading of at least about 20 g/L, 30 g/L, or 40 g/L. This addition of polypeptide is then followed by continuous addition of further substrate to the solution at a rate of about 2 g/L/h, 4 g/L/h, or 6 g/L/h until a much higher final substrate loading of at least about 30 g/L, 40 g/L, 50 g/L, 60 g/L, 70 g/L, 100 g/L, 150 g/L, 200 g/L or more, is reached. Accordingly, in some embodiments of the method, the suitable reaction conditions comprise addition of the polypeptide to a solution having an initial substrate loading of at least about 20 g/L, 30 g/L, or 40 g/L followed by addition of further substrate to the solution at a rate of about 2 g/L/h, 4 g/L/h, or 6 g/L/h until a final substrate loading of at least about 30 g/L, 40 g/L, 50 g/L, 60 g/L, 70 g/L, 100 g/L or more, is reached. This substrate supplementation reaction condition allows for higher substrate loadings to be achieved while maintaining high rates of conversion of ketone substrate to amine product of at least about 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or greater. In some embodiments of this method, the further substrate added is in a solution comprising isopropylamine or isopropylamine acetate at a concentration

of at least about 0.5 M, at least about 1.0 M, at least about 2.5 M, at least about 5.0 M, at least about 7.5 M, at least about 10.0 M.

In some embodiments of the processes, the transamination reaction using an engineered transaminase polypeptide can comprise the following suitable reaction conditions: (a) substrate loading at about 5 g/L to 200 g/L; (b) about 0.1 to 50 g/L of transaminases polypeptide; (c) about 0.1 to 4 M of isopropylamine (IPM); (d) about 0.1 to 10 g/L of pyridoxal phosphate (PLP) cofactor; (e) pH of about 6 to 9; and (f) temperature of about 30 to 60°C.

In some embodiments of the processes, the transamination reaction using an engineered transaminase polypeptide can comprise the following suitable reaction conditions: (a) substrate loading at about 5 to about 20 g/L; (b) about 0.05 to 2 g/L of transaminases polypeptide; (c) about 1 to 10% v/v of PEG200; (d) about 1 to 2 M of isopropylamine (IPM); (e) about 0.1 to 1 g/L of pyridoxal phosphate (PLP) cofactor; (f) about 0.1 to about 0.5 M of triethanolamine (TEA); (g) pH of about 6 to 8; and (h) temperature of about 45 to 55°C.

In some embodiments of the processes, the transamination reaction using an engineered transaminase polypeptide can comprise the following suitable reaction conditions: (a) substrate loading of about 25 to about 100 g/L; (b) about 0.5 to 10 g/L of transaminases polypeptide; (c) about 1 to 10% v/v of PEG200; (d) about 1 to 2 M of isopropylamine (IPM); (e) about 0.1 to 1 g/L of pyridoxal phosphate (PLP) cofactor; (f) about 0.1 to about 0.5 M of triethanolamine; (g) pH of about 6 to 8; and (h) temperature of about 45 to 55°C.

In some embodiments, additional reaction components or additional techniques carried out to supplement the reaction conditions. These can include taking measures to stabilize or prevent inactivation of the enzyme, reduce product inhibition, shift reaction equilibrium to product amine. formation.

Accordingly, in some embodiments of the process for preparing an amine, such as a chiral amine, additional quantities of the amino acceptor can be added (up to saturation) and/or the amino acceptor (ketone) formed can be continuously removed from the reaction mixture. For example, a solvent bridge or a two phase co-solvent system can be used to move the amine product to an extraction solution, and thereby reduce inhibition

by amine product and also shift the equilibrium towards product formation (see, e.g., Yun and Kim, 2008, Biosci. Biotechnol. Biochem. 72(11):3030-3033).

In some embodiments of the processes, the suitable reaction conditions comprise the presence of the reduced cofactor, nicotinamide adenine dinucleotide (NADH), which can act to limit the inactivation of the transaminase enzyme (see e.g., van Ophem et al., 1998, Biochemistry 37(9):2879-88). In such embodiments where NADH is present, a cofactor regeneration system, such as glucose dehydrogenase (GDH) and glucose or formate dehydrogenase and formate can be used to regenerate the NADH in the reaction medium.

In some embodiments, the process can further comprise removal of the carbonyl by-product formed from the amino group donor when the amino group is transferred to the amino group acceptor. Such removal *in situ* can reduce the rate of the reverse reaction such that the forward reaction dominates and more substrate is then converted to product. Removal of the carbonyl by-product can be carried in a number of ways. Where the amino group donor is an amino acid, such as alanine, the carbonyl by product, a keto acid, can be removed by reaction with a peroxide (see, e.g., US 2008/0213845, incorporated herein by reference). Peroxides which can be used include, among others, hydrogen peroxide; peroxyacids (peracids) such as peracetic acid ($\text{CH}_3\text{CO}_3\text{H}$), trifluoroperacetic acid and metachloroperoxybenzoic acid; organic peroxides such as t-butyl peroxide ($(\text{CH}_3)_3\text{COOH}$), or other selective oxidants such as tetrapropylammonium perruthenate, MnO_2 , KMnO_4 , ruthenium tetroxide and related compounds. Alternatively, pyruvate removal can be achieved via its reduction to lactate by employing lactate dehydrogenase to shift equilibrium to the product amine (see, e.g., Koszelewski et al., 2008, Adv. Syn. Catal. 350: 2761-2766). Pyruvate removal can also be achieved via its decarboxylation by employing pyruvate decarboxylase (see, e.g., Höhne et al., 2008, Chem BioChem 9: 363-365) or acetolactate synthase (see, Yun and Kim, supra).

Alternatively, in embodiments where an amino acid is used as amino group donor, the keto acid carbonyl by-product can be recycled back to the amino acid by reaction with ammonia and NADH using an appropriate dehydrogenase enzyme, e.g., amino acid dehydrogenase, in presence of an amine donor, such as ammonia, thereby replenishing the amino group donor.

In some embodiments, where the choice of the amino donor results in a carbonyl by-product that has a vapor pressure higher than water (e.g., a low boiling co-product such

as a volatile organic carbonyl compound), the carbonyl by-product can be removed by sparging the reaction solution with a non-reactive gas or by applying a vacuum to lower the reaction pressure and removing the carbonyl by-product present in the gas phase. A non-reactive gas is any gas that does not react with the reaction components. Various non-reactive gases include nitrogen and noble gases (*e.g.*, inert gases). In some embodiments, the non-reactive gas is nitrogen gas. In some embodiments, the amino donor used in the process is isopropylamine (IPM), which forms the carbonyl by-product acetone upon transfer of the amino group to the amino group acceptor. The acetone can be removed by sparging with nitrogen gas or applying a vacuum to the reaction solution and removing the acetone from the gas phase by an acetone trap, such as a condenser or other cold trap. Alternatively, the acetone can be removed by reduction to isopropanol using a transaminase.

In some embodiments of the processes above where the carbonyl by-product is removed, the corresponding amino group donor can be added during the transamination reaction to replenish the amino group donor and/or maintain the pH of the reaction. Replenishing the amino group donor also shifts the equilibrium towards product formation, thereby increasing the conversion of substrate to product. Thus, in some embodiments wherein the amino group donor is isopropylamine and the acetone product is removed *in situ*, isopropylamine can be added to the solution to replenish the amino group donor lost during the acetone removal and to maintain the pH of the reaction (*e.g.*, at about 8.5).

In further embodiments, any of the above described process for the conversion of substrate compound to product compound can further comprise one or more steps selected from: extraction; isolation; purification; and crystallization of product compound. Methods, techniques, and protocols for extracting, isolating, purifying, and/or crystallizing the product amine from biocatalytic reaction mixtures produced by the above disclosed methods are known to the ordinary artisan and/or accessed through routine experimentation. Additionally, illustrative methods are provided in the Examples below.

Section J: Examples

The following Examples serve to illustrate the invention without limiting the scope thereof, while they on the other hand represent preferred embodiments of the reaction steps, intermediates and/or the process of the present invention.

Abbreviations:

δ	chemical shift
μl	microlitre
μm	micrometre
aq	aqueous
(+)CSA	(+) Camphorsulphonic Acid
ESTP	Ethyl Acetate
IPA	Isopropyl Acetate
TRMA	Treithyl Amine
Ac	acetyl
AcOH	acetic acid
br	broad
brm	broad multiplet
br. m.	broad multiplet
br. mult.	broad multiplet
br. s.	broad signal
br. sign.	broad signal
cat.	catalytic amount
compl. m	complex multiplet
compl. mult.	complex multiplet
copl. m.	complex multiplet
cpl. m.	complex multiplet
CDCl_3	deuterated chloroform
CH_2Cl_2	dichloromethane
CH_2O	formaldehyde
CO_2	carbon dioxide
d	doublet
dd	doublet of doublet
de	diastereomeric excess
dr	diastereomeric ratio
DCM	dichloromethane
DEA	diethyl amine
DMSO	dimethylsulfoxide
$\text{DMSO-}d_6$	dimethylsulfoxide deuterated


ee	enantiomeric excess
equiv	equivalent
ES	electrospray
ES+	positive electrospray ionisation
ESI	electrospray ionisation
Et	ethyl
Et ₂ O	diethyl ether
EtOAc	ethyl acetate
EtOH	ethanol
FTIR	fourier transform infrared
g	gram(s)
GC	gass chromatography
h	hour(s)
HCl	hydrochloric acid
HCl _(aq)	hydrogen chloride aqueous solution
HNMR	proton nuclear magnetic resonance
¹ HNMR	proton nuclear magnetic resonance
HPLC	high performance liquid chromatography
H ₃ PO ₄	phosphoric acid
HRMS	high resolution mass spectroscopy
H ₂ SO ₄	sulfuric acid
Hz	hertz
iPr	isopropyl
iPr ₂ NEt	N-ethyldiisopropylamine
iPrOAc	isopropyl acetate
iPrOH	isopropanol
IR	infrared
<i>J</i>	coupling constant
K ₂ CO ₃	potassium carbonate
L	litre
LC-MS	liquid chromatography-mass spectrometry
LCMS	liquid chromatography-mass spectrometry
LRMS	low resolution mass spectroscopy
m	multiplet

m/e	mass-to-charge ratio
mg	milligram
min	minute(s)
ml	millilitre
mL	millilitre
mmol(s)	millimole(s)
mol(s)	mole(s)
monosub.	monosubstituted
mp	melting point
m.p.	melting point
mult. d	multiplet doublets
m/z	mass-to-charge ratio
M	molarity/molar
Me	methyl
2-MeTHF	2-methyltetrahydrofuran
MeOH	methanol
MgSO ₄	magnesium sulfate
MS	mass spectrometry
[MS] ⁺	mass of proton adduct
MTBE	tertbutylmethylether
nm	nanometre
N	nitrogen atom
N ₂	nitrogen
NaCl	sodium chloride
Na ₂ CO ₃	sodium carbonate
NaHCO ₃	sodium bicarbonate
NaHMDS	sodium bis(trimethylsilyl)amide
NaOH	sodium hydroxide
Na ₂ SO ₄	sodium sulfate
NH ₄ Cl	ammonium chloride
NMR	nuclear magnetic resonance
oct.	octet
ppm	parts per million
psi	pounds per square inch

Pd/C	palladium on carbon
Pd(PPh ₃) ₄	tetrakis(triphenylphosphine)palladium(0)
Pt/C	platinum on carbon
q	quartet
Rh/C	rhodium on carbon
RT = rt	room temperature
s	singlet
sev. brm	several broad multiplets
sev. dd	several doublets of doublets
t	triplet
temp.	temperature
T	temperature
t _R	retention time
tBu	tertiary-butyl
TEA	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
Tol	toluene
UV	ultraviolet light
vol.	volume
wt.	weight

Example 1: Synthesis of Ketone Intermediate

6-Chloro-5-fluoro-3-hydroxy-3-(2-oxo-propyl)-1,3-dihydro-indol-2-one




To the suspension of 6-Chloro-5-fluoro-isatin(25 g, 125 mmol) in acetone (620 ml), Et₂NH (0.9 g, 12.5 mmol) and K₂CO₃ (1.7g, 12.5 mmol) were added. The reaction mixture was heated to reflux for 2 hours. After cooled down to the r.t. , the reaction mixture was filtrated and the solvent was removed under reduced pressure. The crude

product was re-dissolved in MeOH (300 ml) at 50 °C and water (300 ml) was added. With addition of water, the solid was precipitate. MeOH (250 mL) was removed under reduce pressure at 50 °C; The mixture was cooled down to 0-5 °C and stirred for 30mins; The solid was filtered and dried to produce 6-Chloro-5-fluoro-3-hydroxy-3-(2-oxo-propyl)-1,3-dihydro-indol-2-one (27g). ¹H NMR (DMSO-d₆): 2.00(3H, s, CH₃), 3.09(1H, d, CH, *J* =16Hz), 3.39(1H, d, CH, *J* =16Hz), 6.16(1H, s, OH); 6.88(1H, d, CH, *J* =8Hz), 7.36(1H, d, CH, *J* =12Hz), 10.37(1H, s, NH). MS (ESI) *m/z* 258.0 (M+H)⁺

HPLC method : Column: Agilent Extend-C18; 150×3.0mm; 3.5 μm. Mobile Phase A (0.1 % H₃PO₄) in water; Mobile Phase B (Acetonitrile). Gradient: 0 min (95 % A); 0.5 min (95% A), 14 min (5% A), 19.5 min (5% A),. Flow rate: 0.5 ml min⁻¹ . Wavelength: 210 nm. Temperature: 40 °C. Retention time: 7.52 min.


6-Chloro-5-fluoro-3-hydroxy-3-(2-methyl-[1,3]dioxolan-2-ylmethyl)-1,3-dihydro-indol-2-one



To the suspension of 6-chloro-5-fluoro-3-hydroxy-3-(2-oxopropyl)indolin-2-one (25g, 97 mmol) in ethylene glycol (250 mL), triethyl orthoformate (43g, 291mmol) and toluene-4-sulfonic acid monohydrate (0.9g, 4.8mmol) were added. The reaction mixture was heated at 40-50 °C for 2 hours. MeTHF (500 mL) and 10% NaCl aqueous solution (500 mL) were successively added. The organic layer was separated and washed with water twice. After the solvent was evaporated under vacuum, the white solid was washed with TBME (400 mL) and dried to produce 6-Chloro-5-fluoro-3-hydroxy-3-(2-methyl-[1,3]dioxolan-2-ylmethyl)-1,3-dihydro-indol-2-one (27.8 g). ¹H NMR (DMSO-d₆): 1.00(3H, s, CH₃), 2.37(2H, d, CH₂), 3.13(1H, q, CHH), 3.58(1H, q, CHH), 3.59(1H, q, CHH), 3.68(1H, q, CHH), 6.00(1H, s, OH); 6.85(1H, d, CH, *J* =8Hz), 7.31(1H, d, CH, *J* =12Hz), 10.24(1H, s, NH). MS (ESI) *m/z* 302.0 (M+H)⁺ HPLC method :Column: Agilent Extend-C18; 150×3.0mm; 3.5 μm. Mobile Phase A (0.1 % H₃PO₄) in water; Mobile Phase B (Acetonitrile). Gradient: 0 min (95 % A); 0.5 min (95% A), 14 min (5% A), 19.5

min (5% A),. Flow rate: 0.5 ml min⁻¹ . Wavelength: 210 nm. Temperature: 40 °C.
Retention time: 7.94 min.

1-(6-Chloro-5-fluoro-1H-indol-3-yl)-propan-2-one



To the solution of 6-chloro-5-fluoro-3-hydroxy-3-((2-methyl-1,3-dioxolan-2-yl)methyl)indolin-2-one (27.8g, 92mmol) in THF (250 mL), Red-Al (79.8g, 276mmol) in THF (100 mL) was added slowly at 60 °C under N₂ atmosphere. After the addition, the mixture was stirring at r.t. for 3 hours. 20 % HCl (400 mL) was added and the mixture was stirred for 20mins. Ethyl acetate (500 mL) was added and stirred for 10mins. The organic layer was separated and washed with water (100ml x 2); After treated with Na₂SO₄ (50 g) and active carbon (5 g) at r.t., the mixture was filtered and the filtration was concentrated. The crude product was recrystallized with MeOH (80 mL) and water (60 mL) to produce 1-(6-Chloro-5-fluoro-1H-indol-3-yl)-propan-2-one (15.5 g).


¹H NMR (DMSO-d₆): 2.21(3H, s, CH₃), 3.78(2H, s, CH₂), 7.13(1H, d, CH, J = 4Hz); 7.23(1H, d, CH, J = 8Hz), 7.33(1H, d, CH, J = 8Hz), 8.23(1H, s, NH).

MS (ESI) m/z 226.0 (M+H)⁺

HPLC method :Column: Agilent Extend-C18; 150×3.0mm; 3.5 μm. Mobile Phase A (0.1 % H₃PO₄) in water; Mobile Phase B (Acetonitrile). Gradient: 0 min (95 % A); 0.5 min (95% A), 14 min (5% A), 19.5 min (5% A),. Flow rate: 0.5 ml min⁻¹. Wavelength: 210 nm. Temperature: 40 C Retention time: 10.75 min.

Example 2: Second Alternative Synthesis of Ketone Starting Material


1-(6-Chloro-5-fluoro-1H-indol-3-yl)-propan-2-one



Under N₂, Pt/C (10 g, 5% Pt on Carbon) was charged to the suspension of 6-Chloro-5-fluoro-3-((Z)-2-nitro-propenyl)-1H-indole (100 g, 0.39 mol) in ethyl acetate (500 mL). The reaction mixture vacuumed and purged with H₂ three times and stirred at r.t. for overnight. The catalyst was filtered off and the organic solvent was removed under reduced pressure to give brown oil. The oil was re-dissolved in EtOH (500 mL) and sodium bisulfite (81.7 g, 0.79 mol) in H₂O () was added. The suspension was heated up to 80 °C for overnight. The precipitate was filtered off and ethanol was removed under pressure. The residue was diluted with water (500 mL) and extracted with ethyl acetate (300mL X2). The combined organic layer was dried over Na₂SO₄ and condensed on the rota vapor to give brown oil (53 g, 60 % yield).

Example 3: Second Alternative Synthesis of Ketone Intermediate

1-(6-Chloro-5-fluoro-1H-indol-3-yl)-propan-2-one



Raney/Ni (0.75g) and acetic acid (1.1 mL, 19.6 mmol) were charged to the suspension of 6-Chloro-5-fluoro-3-((Z)-2-nitro-propenyl)-1H-indole (5g, 19.6 mmol) in the mixture of MeOH and H₂O (2:1, 50 mL) at r.t. The reaction mixture vacuumed and purged with H₂ three times and heated at 50 °C for overnight. MeOH (50 mL) was added to the reaction mixture and the catalyst was filtered off over celite. The solvent was concentrated to 50 mL and H₂O (300 ml) was added. With the addition of H₂O, off-white solid was produced. The solid was filtered and washed by water. After dried in the oven, 2.53 g off-white solid was isolated.

Example 4: Synthesis, Optimization, and Screening Engineered Transaminase Polypeptides

Gene synthesis and optimization: The polynucleotide sequence encoding the reported wild-type omega transaminase polypeptide from *Vibrio fluvialis* of SEQ ID NO: 2 was codon optimized and synthesized as the gene of SEQ ID NO: 1. The synthetic gene of SEQ ID NO: 1 was cloned into a pCK110900 vector system (see e.g., US Patent Application Publication 20060195947, which is hereby incorporated by reference herein) and subsequently expressed in *E. coli* W3110 *fhuA*. The *E. coli* W3110 expresses the transaminase polypeptides as an intracellular protein under the control of the *lac* promoter. The polynucleotide (SEQ ID NO:3) encoding the engineered transaminase polypeptide of SEQ ID NO: 4 was obtained by directed evolution of the codon-optimized gene SEQ ID NO:1. The polypeptide of SEQ ID NO:4 has 24 amino acid residue differences (A9T; G18A; D21H; V31M; N45H; F86Y; A133R; R146L; W147K; V153S; K163F; V177L; R211K; P233T; A235P; P244T; M294V; P318D; A323T; S324G; A383V; T391A; C424A; F427Y) relative to SEQ ID NO:2. This synthetic gene SEQ ID NO: 3 (encoding the polypeptide of SEQ ID NO: 4) was used as the starting backbone for further optimization using standard methods of directed evolution via iterative variant library generation by gene synthesis followed by screening and sequencing of the hits to generate genes encoding engineered transaminases capable of converting compound (IA) to compound (IIA) with improved enzyme properties relative to the polypeptides SEQ ID NOs: 2 and 4. The resulting engineered transaminase polypeptide sequences and specific mutations and relative activities are listed in **Table 2A**.

Example 5: Production of Engineered Transaminases

The engineered transaminase polypeptides were produced in *E. coli* W3110 as an intracellular protein expressed under the control of the *lac* promoter. The polypeptide accumulates primarily as a soluble cytosolic active enzyme. A shake-flask procedure is used to generate engineered polypeptide powders that can be used in activity assays or biocatalytic process disclosed herein.

High-throughput growth & expression. Cells were picked and grown overnight in LB media containing 1% glucose and 30 µg/mL chloramphenicol (CAM), 30°C, 200 rpm, 85% humidity. 20 µL of overnight growth were transferred to a deep well plate containing 380 µL 2xYT growth media containing 30 µg/mL CAM, 1 mM IPTG, 1 mM MgSO₄, and incubated for ~18 h at 30°C, 200 rpm, 85% humidity. Subculture TB media was made up

of TB media (380 μ L/well), 30 μ g/mL CAM, 1mM MgSO_4 , and 1mM IPTG. Cell cultures were centrifuged at 4000 rpm, 4°C for 10 min., and the media discarded. Cell pellets were resuspended in 200 or 400 μ L Lysis Buffer (0.1 M triethanolamine (TEA) buffer, pH 9.0, containing 1 mM MgSO_4 , 400 μ g/mL PMBS and 500 μ g/mL Lysozyme), as described below.

Production of shake flask powders (SFP): A shake-flask procedure was used to generate engineered transaminase polypeptide powders used in secondary screening assays or in the biocatalytic process disclosed herein. Shake flask powder (SFP) includes approximately 30% total protein and accordingly provide a more purified preparation of an engineered enzyme as compared to the cell lysate used in HTP assays. A single microbial colony of *E. coli* containing a plasmid encoding an engineered transaminase of interest is inoculated into 50 mL Luria Bertani broth containing 30 μ g/ml chloramphenicol and 1% glucose. Cells are grown overnight (at least 16 hours) in an incubator at 30°C with shaking at 250 rpm. The culture is diluted into 250 mL Terrific Broth (12 g/L bacto-tryptone, 24 g/L yeast extract, 4 mL/L glycerol, 65 mM potassium phosphate, pH 7.0, 1 mM MgSO_4) containing 30 μ g/ml chloramphenicol, in a 1 liter flask to an optical density at 600 nm (OD600) of 0.2 and allowed to grow at 30°C. Expression of the transaminase gene is induced by addition of isopropyl- β -D-thiogalactoside ("IPTG") to a final concentration of 1 mM when the OD600 of the culture is 0.6 to 0.8 and incubation is then continued overnight (at least 16 hours). Cells are harvested by centrifugation (5000 rpm, 15 min, 4°C) and the supernatant discarded. The cell pellet is resuspended with an equal volume of cold (4°C) 100 mM triethanolamine (chloride) buffer, pH 7.0 (optionally including 2 mM MgSO_4), and harvested by centrifugation as above. The washed cells are resuspended in two volumes of the cold triethanolamine (chloride) buffer and passed through a French Press twice at 12,000 psi while maintained at 4°C. Cell debris is removed by centrifugation (9000 rpm, 45 minutes, 4°C). The clear lysate supernatant was collected and stored at -20°C. Lyophilization of frozen clear lysate provides a dry shake-flask powder of crude transaminase polypeptide. Alternatively, the cell pellet (before or after washing) can be stored at 4°C or -80°C.

Production of downstream process (DSP) powders: DSP powders contains approximately 80% total protein and accordingly provide a more purified preparation of the engineered transaminase enzyme as compared to the cell lysate used in the

high throughput assay. Larger-scale (~100 – 120 g) fermentation of the engineered transaminase for production of DSP powders can be carried out as a short batch followed by a fed batch process according to standard bioprocess methods. Briefly, transaminase expression is induced by addition of IPTG to a final concentration of 1 mM. Following fermentation, the cells are harvested and resuspended in 100 mM Triethanolamine-H₂SO₄ buffer, then mechanically disrupted by homogenization. The cell debris and nucleic acid are flocculated with polyethylenimine (PEI) and the suspension clarified by centrifugation. The resulting clear supernatant is concentrated using a tangential cross-flow ultrafiltration membrane to remove salts and water. The concentrated and partially purified enzyme concentrate can then be dried in a lyophilizer and packaged (e.g., in polyethylene containers).

Example 6: Analytical Procedures

HPLC Analysis of HTP Reactions: An aliquot of the quenched reaction was subject to HPLC analysis under the following conditions.

Column	Mightysil RP-18 GP Aqua 150x4.6 mm, 5 μ m
Temperature	30°C
Mobile Phase	Isocratic, 60% 20 mM NH ₄ Ac (pH 6.6)/40% acetonitrile
Flow Rate	2.5 mL/min
Detection	254 nm
Retention Times	Amine Product compound (IIA): 0.96 min; Substrate Impurity: 1.8 min; Ketone Substrate compound (IA): 2.7 min

Conversion of compound (**IA**) to compound (**IIA**) was determined from the resulting chromatograms as follows:

$$\text{Conversion (\%)} = \text{Product Area} / (\text{Product Area} + \text{Substrate Area} \times 0.73) \times 100\%$$

HPLC Analysis of 5 mL and 100 mL scale reactions: Aliquots of the quenched reaction was subject to HPLC analysis under the following conditions.

Column	Mightysil RP-18 GP Aqua 250x4.6 mm, 5 μ m
Temperature	30°C
Mobile Phase	Isocratic: 60% 20 mM NH ₄ Ac (pH 6.6)/40% acetonitrile
Flow Rate	2.0 mL/min
Detection	254 nm

Retention Times	Amine Product compound (IIA): 2.1 min Ketone Substrate compound (IA): 7.6 min

Conversion of compound (**IA**) to compound (**IIA**) was determined from the chromatograms as follows:

$$\text{Conversion (\%)} = \text{Product Area} / (\text{Product Area} + \text{Substrate Area} \times 0.73) \times 100\%$$

Determination of product chiral purity (%ee): The chiral purity or enantiomeric excess of compound (**IIA**) was assessed by HPLC using the following conditions.

Column	Astec Chirobiotic TAG column
Temperature	15°C
Mobile Phase	Methanol/Acetic acid/Triethylamine (100/0.2/0.1)
Flow Rate	1.0 mL /min
Detection Wavelength	225 nm
Retention Times	Ketone substrate: 3.6 min <i>R</i> -product: 17.9 min; <i>S</i> -product: 18.9 min

Determination of product purity: The purity of product was determined by HPLC using the following conditions.

Column	Mightysil Rp-18 GP aqua 250 x 4.6 mm, 5 µm	
Temperature	30 °C	
Mobile Phase	Gradient. A: 20 mM NH ₄ Ac (pH 6.6); B: acetonitrile	
	Time	Composition
	0 min	5% B
	1.5 min	5% B
	35 min	70% B
	40 min	70% B
	45 min	5% B
	60 min	5% B
Flow Rate	1.5 mL/min	
Detection Wavelength	254 nm	
Retention Times	Major impurity: 22.2 min; amine product: 17.5 min; ketone substrate: 25.4 min	

Example 7: High Throughput (HTP) Screening of Transaminases for Conversion of Compound (IA) to Compound (IIA)

HTP Screening Assays: High-throughput screening used to guide primary selection of variants was carried out in 96-well plates using cell lysates under assay conditions of 10 g/L compound (IA); 1 mM pyridoxal phosphate (PLP); 2 M isopropylamine (IPM), pH 7.0; 0.1 M triethanolamine (TEA), pH 7; 5% v/v PEG200; 10 uL lysate; and 50 or 55°C.

Cells were grown in 96-well plates as described above and lysates prepared by dispensing 200 uL (for Round 1) or 400 uL (for Round 2) of Lysis Buffer (1 mg/mL lysozyme, 0.5 mg/mL polymyxin B sulfate, 1 mM PLP, 0.1 M triethanolamine (TEA), pH 7.0) into each well. Plates were sealed, shaken for 2 h, and then centrifuged for 20 min at 4,000 rpm at 4 °C to pellet the cell debris.

A 10 uL of stock substrate solution (200 g/L compound (IIA) dissolved in PEG200) was added to each well of a 96-well plate followed by 180 uL of a stock solution of isopropylamine (IPM)/pyridoxal phosphate (PLP) (2.2 M IPM and 1.06 mM PLP in 100 mM TEA, pH 7.0. For assessing refractoriness to product compound (IIA) inhibition, compound (IIA) was added to the reaction mixture to a final 14 g/L for Round 1 and 16 g/L Round 2 assays. Reactions were initiated by adding 10 uL of cell lysate/well. Plates were sealed and incubated with shaking at 50 or 55°C for 24 h. Reactions were quenched with 600 uL of acetonitrile and samples examined by HPLC as described in Example 5.

Example 8: Process for Conversion of Compound (IA) to Compound (IIA) in 5 mL Scale

A 5 ml scale reactions were carried out as follows. To a 20 mL glass vial with equipped a cross-shaped magnetic stirring bar was added 0.75 mL (or 0.5 mL in case of 10 % v/v PEG 200 concentration) of 100 mM TEA buffer (pH 7.0). 2 mL of 5 M IPM•HCl stock solution was added to the vial followed by 1 mL of 5 mM PLP stock solution. The mixture was stirred at 500 rpm (magnetic stirring). The pH of the mixture was then adjusted to 7.0 using 1 M NaOH solution. 125 mg (or 250 mg for 50 g/L or 500 mg for 100 g/L concentration) substrate (solid) was then added to the vial. 0.25 mL (or 0.5 mL for 10 % v/v) PEG 200 was then added to the mixture. Final concentrations of components were: 25 g/L (or 50 or 100 g/L) of compound (IA); 1 mM PLP; 2 M IPM; 5 % v/v (or 10 %) PEG 200; 2 g/L TA enzyme; 100 mM TEA, pH 7.0). The mixture was then stirred and heated to 50°C.

Reactions were initiated by adding 1 mL of the enzyme stock solution (10 g/L). 20 μ L samples were taken at different time points and diluted with 750 μ L methanol and analyzed by HPLC. After 24 h, the reaction mixtures were quenched with 5 mL acetonitrile and samples analyzed by HPLC to get the final %conversion.


Example 9: Process for Conversion of Compound (IA) to Compound (IIA) in 100 mL Scale

A 100 mL scale reaction was carried out as follows. To a 250 mL round bottom flask with an anchor-shaped stirring blade was added 15 mL of 100 mM TEA buffer (pH 7). 40 mL of the 5 M IPM•HCl stock solution was added to the round bottom flask followed by 20 mL of the 5 mM PLP stock solution. The mixture was stirred at 100 rpm (overhead stirring) and the pH adjusted to 7.0 using 10 M NaOH. Solid substrate Compound (IA) was then added to the round bottom flask over ~5 min with stirring. PEG 200 (5 mL) was then added and the mixture heated to 50°C. 20 mL of the enzyme stock solution (10 g/L) was then added to start the reaction. The reaction was stirred at 200 rpm (overhead stirring). 20 μ L samples were taken at different time points and diluted with 750 μ L methanol and analyzed by HPLC. In some cases, where work up was not pursued, after 24 h, the reaction mixtures were quenched with 100 mL acetonitrile and samples analyzed by HPLC to get the final %conversion (Figure 9). Concentration of different components in the reaction: ketone: 25 g/L (or 50 or 100 g/L); PLP: 1 mM; IPM: 2 M; PEG200: 5 % v/v; TA enzyme: 2 g/L; buffer: 100 mM TEA, pH 7.0. Conversion of substrate to product was analyzed by HPLC as described in Example 3.

Product Workup: After 24 h reaction under the conditions described above (up to point 11) the reaction mixture was cooled to room temperature. The mixture was filtered through a standard G4 sintered glass funnel with a piece of filter paper (Whatman 1, pore size 11 μ m). The round bottom flask was rinsed with 20 mL deionized water which was then filtered through the same funnel. The filter cake was washed twice with 20 mL deionized water. The pH of the filtrate was adjusted from 6.8 to 3 using a 5 M HCl solution. The filtrate was then transferred into a separatory funnel and extracted with 100 mL MTBE. The biphasic mixture was allowed to separate. The MTBE layer containing unreacted substrate and impurities was discarded. The aqueous layer was transferred into a beaker and 100 mL MTBE was added. The pH of the aqueous layer was then adjusted from 3 to 10 using 10 M NaOH solution. The mixture was transferred into a separatory funnel and the phases allowed to separate.

The aqueous layer was then extracted with 100 mL MTBE three times (nb: repeated until product was not present in aqueous layer). The MTBE phases from the three extractions were combined and evaporated to dryness using a rotary evaporator. The crude product was further dried under vacuum for 48 h.

Example 10: Process for Conversion of Compound (IA) to Compound (IIA) in 30g Scale



152.48g *iso*-propylamine hydrochloride and 0.204g pyridoxalphosphate monohydrate were dissolved in 495ml water while stirring. To this yellow clear solution a solution of 30.0g ketone in 85ml poly ethylene glycol (average mol weight 200) within 15 minutes. Upon addition the ketone precipitates as fine particles which are evenly distributed in the reaction media. To the suspension 180ml triethanolamine buffer (0.1 mol/l, pH 7) were added and the pH was adjusted to 7 by addition of aqueous sodium hydroxide solution (1 mol/l). The reaction mixture is heated to 50°C and a solution of 1.62g transaminase SEQ ID NO: 134 dissolved in 162ml triethanolamine buffer (0.1 mol/l, pH 7) is added. The reaction mixture is continuously kept at pH 7 by addition of 1 mol/l aqueous sodium hydroxide solution. The reaction mixture is stirred 24h at 50°C and a stream of Nitrogen is blown over the surface of the reaction mixture to strip off formed acetone. The reaction mixture is then cooled to 25°C and filtered over a bed of cellulose flock. The pH of the filtrate is adjusted to ≈ 1 by addition of concentrated sulfuric acid. The acidified filtrate is extracted with 250 ml *iso*-Propyl acetate. The layers are separated and the pH of the aqueous phase is adjusted to ≈ 10 by addition of concentrated aqueous sodium hydroxide

solution. The basified aqueous phase is extracted with *iso*-propyl acetate. The layers are separated and the organic phase is washed with 100 ml water. The organic phase is concentrated by distillation to 2/3 of its origin volume. In a second reactor 33.98g (+)-camphor sulfonic acid is dissolved in 225 ml *iso*-propyl acetate upon refluxing and the concentrated organic phase is added within 10 minutes. After complete addition the formed thin suspension is cooled to 0°C within 2 hours and kept at 0°C for 15 hours. The precipitated amine-(+)-camphor sulfonate salt is filtered, washed with 70 ml *iso*-propyl acetate and dried at 40°C in vacuum yielding 51.57g of colourless crystals (84.5% yield t.q.)

Analytical Data

IR:

$\tilde{\nu}$ (cm⁻¹)=3296, 3061, 2962, 2635, 2531, 2078, 1741, 1625, 1577, 1518, 1461, 1415, 1392, 1375, 1324, 1302, 1280, 1256, 1226, 1170, 1126, 1096, 1041, 988, 966, 937, 868, 834, 814, 790, 766, 746, 719, 669, 615.

LC-MS (ESI +):

Ammonium ion: m/z =227 ([M+H]), 268 ([M+H+CH₃CN]), 453 ([2M+H]).

Camphorsulfonate ion: m/z =250 ([M+NH₄]), 482 ([2M+NH₄]).

LC-MS (ESI -):

Camphorsulfonate ion: m/z=231 ([M-H]), 463 ([2M-H]).

¹H-NMR (DMSO-d₆, 400 MHz):


11.22 (br. s., 1H), 7.75 (br. s., 3H), 7.59 (d, *J* = 10.3 Hz, 1H), 7.54 (d, *J* = 6.5 Hz, 1H), 7.36 (d, *J* = 2.3 Hz, 1H), 3.37 - 3.50 (m, 1H), 2.98 (dd, *J* = 14.3, 5.8 Hz, 1H), 2.91 (d, *J* = 14.8 Hz, 1H), 2.81 (dd, *J* = 14.3, 8.0 Hz, 1H), 2.63 - 2.74 (m, 1H), 2.41 (d, *J* = 14.6 Hz, 1H), 2.24 (dt, *J* = 18.3, 3.8 Hz, 1H), 1.94 (t, *J* = 4.4 Hz, 1H), 1.86 (dt, *J* = 7.4, 3.6 Hz, 1H), 1.80 (d, *J* = 18.1 Hz, 1H), 1.23 - 1.35 (m, 2H), 1.15 (d, *J* = 6.3 Hz, 3H), 1.05 (s, 3H), 0.74 (s, 3H)

Free Amine (obtained by evaporating the *iso*-Propylacetate layer after extraction of the basified aqueous layer):

¹H NMR (400MHz, DMSO-d₆):

11.04 (br. s., 1H), 7.50 (d, J = 10.5 Hz, 1H), 7.48 (d, J = 6.5 Hz, 1H), 7.25 (s, 1H), 3.03 (sxt, J = 6.3 Hz, 1H), 2.61 (dd, J = 14.3, 6.5 Hz, 1H), 2.57 (dd, J = 14.1, 6.5 Hz, 1H), 1.36 (br. s., 2H), 0.96 (d, J = 6.3 Hz, 3H)

Example 11: Process for Conversion of Compound (IIA) to Compound (IVB)



13.62 g 5-chloroisatin is suspended in 35 ml *iso*-propanol and 2.3 g triethyl amine is added. The suspension is heated to reflux and a solution of 34.42g amine-(+)-camphor sulfonate salt dissolved in 300 ml *iso*-propanol is added within 50 minutes. The reaction mixture is stirred at reflux for 17 hours. The reaction mixture is cooled to 75°C and 17.4g (+)-camphorsulfonic acid are added to the reaction mixture. Approximately 300 ml *iso*-propanol are removed by vacuum distillation. Distilled off *iso*-propanol is replaced by *iso*-propyl acetate and vacuum distillation is continued. This distillation is repeated a second time. To the distillation residue 19 ml ethanol and 265 ml ethyl acetate is added and the mixture is heated to reflux. The mixture is cooled in ramps to 0°C and kept at 0°C for 24 hours. The beige to off white crystals are filtered off, washed with 3 portions (each 25 ml) precooled (0°C) ethylacetate and dried in vacuum yielding 40.3 g beige to off white crystals. (86.3% yield t.q.)

IR:

$\tilde{\nu}$ (cm^{-1}) = 3229, 3115, 3078, 3052, 2971, 2890, 2841, 2772, 2722, 2675, 2605, 2434, 1741, 1718, 1621, 1606, 1483, 1460, 1408, 1391, 1372, 1336, 1307, 1277, 1267, 1238, 1202, 1184, 1162, 1149, 1128, 1067, 1036, 987, 973, 939, 919, 896, 871, 857, 843, 785, 771, 756, 717, 690, 678, 613.

LC-MS (ESI +):

Ammonium ion: m/z = 390 ($[\text{M}+\text{H}]$), 431 ($[\text{M}+\text{H}+\text{CH}_3\text{CN}]$)

Camphorsulfonate ion: $m/z = 250$ ($[M+NH_4]$), 482 ($[2M+NH_4]$)


LC-MS (ESI -):

Camphorsulfonate ion: $m/z = 231$ ($[M-H]$), 463 ($[2M-H]$)

^1H NMR (DMSO- d_6 , 600 MHz):

11.49 (s, 1H), 11.23 (s, 1H), 10.29 - 10.83 (m, 1H), 9.78 - 10.31 (m, 1H), 7.55 - 7.60 (m, 2H), 7.52 (s, 1H), 7.40 (d, $J = 6.2$ Hz, 1H), 7.16 (d, $J = 8.8$ Hz, 1H), 4.52 - 4.63 (m, 1H), 3.20 (dd, $J = 16.3, 4.2$ Hz, 1H), 2.96 (dd, $J = 16.1, 11.3$ Hz, 1H), 2.90 (d, $J = 15.0$ Hz, 1H), 2.56 - 2.63 (m, 1H), 2.39 (d, $J = 14.6$ Hz, 1H), 2.21 (dt, $J = 18.0, 3.8$ Hz, 1H), 1.89 - 1.93 (m, 1H), 1.81 (ddd, $J = 15.3, 7.8, 3.7$ Hz, 1H), 1.76 (d, $J = 18.3$ Hz, 1H), 1.53 (d, $J = 6.6$ Hz, 3H), 1.20 - 1.33 (m, 2H), 0.98 (s, 3H), 0.70 (s, 3H)

Example 12: Process for Preparing a Compound of formula (IVA) $\frac{1}{2}$ Hydrate



In a 750ml reactor with impeller stirrer 50g of compound (IVB) salt were dissolved in 300ml Ethanol (ALABD) and 100 ml deionised Water (WEM). The clear, yellowish solution was heated to 58°C internal temperature. To the solution 85 g of a 10% aqueous sodium carbonate solution was added within 10 minutes. The clear solution was particle filtered into a second reaction vessel. Vessel and particle filter were each rinsed with 25 ml of a mixture of ethanol:water (3:1 v/v) in the second reaction vessel. The combined particle filtered solution is heated to 58°C internal temperature and 200ml water (WEM) were added dropwise within 15 minutes. Towards the end of the addition the solution gets turbid. The mixture is stirred for 10 minutes at 58°C internal temperature and is then cooled slowly to room temperature within 4hours 30 minutes forming a thick, well stirable white suspension. To the suspension 200 ml water are added and the mixture is stirred for additional 15hours 20 minutes at room temperature. The suspension is filtered and the filter cake is washed twice with 25 ml portions of a mixture of ethanol:water 9:1 (v/v). The colourless crystals are dried at 60°C in vacuum yielding 26.23g (=91.2% yield).

2013237330 27 Oct 2015

¹H NMR (400 MHz, DMSO-d₆)


10.70 (s, 1H), 10.52 (s, 1H), 7.44 (d, J = 10.0 Hz, 1H), 7.33 (dd, J = 8.4, 2.1 Hz, 1H), 7.26 (d, J = 6.5 Hz, 1H), 7.05 (d, J = 2.3 Hz, 1H), 6.93 (d, J = 8.3 Hz, 1H), 3.83 - 4.00 (m, 1H), 3.13 (d, J = 6.0 Hz, 1H), 2.77 (dd, J = 15.1, 3.8 Hz, 1H), 2.38 (dd, J = 15.1, 10.5 Hz, 1H), 1.17 (d, J = 6.3 Hz, 3H).

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.


20132237330 23 Mar 2016

The Claims Defining the Invention are as Follows:

1. A process for preparing a compound of formula (II), or a salt or solvate or hydrate thereof,



comprising enzymatically converting a compound of formula (I) to compound of formula (II), or a salt, solvate or hydrate thereof,




wherein:

the dashed line is absent;

A is C=O;

20132237330 23 Mar 2016



R^4 and R^7 are each, independently, H or -Cl;

R^5 is H, -OH, -CH₃, -OCH₃, -F, -Cl, -CF₃ or -CN;


R^6 is H, -OH, -OCH₃, -F or -Cl;

R^8 is H, -CH₃, -CH₂CH₃, -CH₂OH, -CO₂H, -CO₂CH₃, -CO₂CH₂CH₃ or -CF₃; and
n is 1 or 2;

wherein the enzyme is SEQ ID NO: 134.


2. The process of claim 1, wherein R^5 and R^6 are fluoro when: R^8 is -CH₃, and n is 1.
3. The process of claim 1, wherein R^5 and R^6 are fluoro and chloro, when: R^8 is -CH₃, and n is 1.
4. The process of claim 1, wherein R^5 and R^6 are hydrogen when: R^8 is -CH₃ and n is 1.
5. The process of claim 1, wherein R^5 is fluoro when: n is 1, and R^6 is hydrogen.

6. The process of claim 1, wherein the compound of formula (II) is of formula (IIA), or a salt or solvate or hydrate thereof,




(IIA).

7. The process according to claim 1 or 6, wherein the compound of formula (I) is a compound of formula (IA) or a salt or hydrate or solvate thereof,



(IA).

8. A process for preparing a compound of formula (IV), or a salt or hydrate or solvate thereof,




(IV)

comprising,


reacting a compound of formula (III), or a salt or hydrate or solvate thereof,

2013237330 23 Mar 2016




(III)

with a compound of formula (II), or a salt or hydrate or solvate thereof,



(II)

wherein the compound of formula (II), or a salt or hydrate or solvate thereof, is prepared by enzymatically converting a compound of formula (I), or a salt or hydrate or solvate thereof,



(I)

wherein:

the dashed line is absent;





A is C=O;

R^{1'} is hydrogen or (C₁-C₆)alkyl;


2013237330 23 Mar 2016

R^4 , R^5 , R^6 and R^7 , are each, independently hydrogen, halo, hydroxy, amino, alkylamino, dialkylamino, (C_1-C_6) alkyl, and (C_1-C_6) alkyloxy;

R^1 is H, $-CH_3$,



or




;

R^4 and R^7 are each, independently, H or $-Cl$; R^5 is H, $-OH$, $-CH_3$, $-OCH_3$, $-F$, $-Cl$, $-CF_3$ or $-CN$; R^6 is H, $-OH$, $-OCH_3$, $-F$ or $-Cl$;

R^8 is H, $-CH_3$, $-CH_2CH_3$, $-CH_2OH$, $-CO_2H$, $-CO_2CH_3$, $-CO_2CH_2CH_3$ or $-CF_3$;

and n is 1 or 2.

9. A process for preparing a compound of formula (IVA), or a salt or solvate or hydrate thereof,




(IVA)

comprising


reacting a compound of formula (IIIA) or a salt or solvate or hydrate thereof,

20132237330 23 Mar 2016



(IIIA)


with a compound of formula (IIA) or a salt or solvate or hydrate thereof,



(IIA)

to provide the compound of formula (IVA), or a salt or solvate or hydrate thereof,


wherein the compound of formula (IIA) is prepared by enzymatically converting a compound of formula (IA), or a salt or hydrate or solvate thereof,



(IA).

10. The process according to claim 8 or claim 9, wherein the enzyme is SEQ ID NO: 134.

11. The compound of formula (IA):




(IA);


or a pharmaceutically acceptable salt or hydrate or solvate thereof.

2013237330 23 Mar 2016


12. A compound selected from:



,



or



, or a pharmaceutically acceptable salt, solvate or

hydrate thereof.

eolf-seql.txt
SEQUENCE LISTING

<110> Novartis AG
CROWE, Michael
FOULKES, Michael
FRANCESE, Giancarlo
GRIMLER, Dominique
KUESTERS, Ernst
LAUMEN, Kurt
LI, Yunzhong
LIN, Changxue
NAZOR, Jovana
RUCH, Thomas
SMITH, Derek
SONG, Shiwei
TENG, Shangjun

<120> CHEMICAL PROCESS FOR PREPARING
SPIROINDOLONES AND INTERMEDIATES THEREOF

<130> PAT055051-WO-PCT
<140>
<141>

<150> PCT/CN2012/000359
<151> 2012-03-23

<160> 154

<170> PatentIn version 3.5

<210> 1
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Codon optimized transaminase gene from *Vibrio fluvialis*

<400> 1
atgaacaaac cgcagagctg ggaagcgcgt gcggaaacct atagcctgta tggctttacc 60
gatatgccga gcctgcatca gcgtggcacc gtggtggtga cccatggcga aggcccgtat 120
atcgtgggatg tgaacggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttttg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatgcgg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacccgttg gaacgcgtat catggcgtga ccgcggtgag cgcgagcatg 480
accggcaaac cgtataacag cgtgtttggc ctgccgctgc cgggctttgt gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtggcgcgt 600
ctggcccgtg aactggaaga aaccattcag cgtgaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgccggcggt gttattccgc cggcgaaagg ctattttcag 720
gcgattctgc cgatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggcc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgcg 840

eolf-seql.txt

```
attattagca gcaaaaacct gaccgcgggt ttttttccga tgggcgcggt gattctgggt      900
ccggaactga gcaaactgtct ggaaaccgcg attgaagcga tcgaagaatt tccgcatggt      960
tttaccgcga gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gcattggctt tatgtgggcg     1140
ctggaagcgg tgaaagataa agcgagcaaa accccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggcctgattt gccgtccgct gggccagagc     1260
gtggttctgt gcccgccgtt tattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362
```

<210> 2
 <211> 453
 <212> PRT
 <213> *Vibrio fluvialis*

<400> 2

Met Asn Lys Pro Gln Ser Trp Glu Ala Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Gly Phe Thr Asp Met Pro Ser Leu His Gln Arg Gly Thr Val Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val Asn Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Phe Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Ala Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Arg Trp Asn Ala Tyr His Gly Val Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Lys Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

eolf-seql.txt

Val His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Arg Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Pro Pro Ala Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Pro Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Met Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Pro His Gly
305 310 315 320

Phe Thr Ala Ser Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Thr Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Cys Pro Pro Phe Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

eolf-seql.txt

Phe Ala Glu Val Ala
450

<210> 3
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 3
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggctttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 4
<211> 453
<212> PRT
<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 4

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1      5      10      15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
      20      25      30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
      35      40      45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
      50      55      60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65      70      75      80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
      85      90      95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
      100     105     110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
      115     120     125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
      130     135     140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145     150     155     160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
      165     170     175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
      180     185     190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
      195     200     205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
      210     215     220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225     230     235     240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
      245     250     255

```


eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 5
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 5
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattgccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240

eolf-seql.txt

```

ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcggggcgt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctgctg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgctg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatattgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 6
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 6

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

eolf-seql.txt

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

eolf-seql.txt

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 7
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 7
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcattc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900

eolf-seql.txt

```

ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg      1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg      1080
aaacatatatg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

```

<210> 8
<211> 453
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Engineered variant of V. fluvialis transaminase

```

```

<400> 8

```

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1          5          10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
          20          25          30

```

```

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

```

```

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
          50          55          60

```

```

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

```

```

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
          85          90          95

```

```

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
          100          105          110

```

```

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
          115          120          125

```

```

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
          130          135          140

```

```

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145          150          155          160

```

eof-seql.txt

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 9
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 9
 atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
 cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
 atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccgtc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
 tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
 gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
 ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 10
 <211> 453
 <212> PRT
 <213> Artificial Sequence

eolf-seql.txt

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 10

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Arg Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 11
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 11
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatgggtg 180

eolf-seql.txt

```

gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggt gttattacc cgcctaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 12
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 12

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

eofl-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 13
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 13
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atgggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcgtgc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840

eolf-seql.txt

```
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggcct tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttcttg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362
```

<210> 14
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 14

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

eolf-seql.txt

Thr Gly Val Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 15
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 15
 atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
 cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
 atcgtggatg tgcattggccg tcgttatctg gatgcgaaca ggggcctgtg gaacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
 ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
 aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
 tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatggg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
 gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 16
 <211> 453

eolf-seql.txt

<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 16

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

eolf-seql.txt

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 17
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 17
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60

cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120

eolf-seql.txt

```

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg      180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg      240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgccggcggt gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 18
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 18

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

eolf-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 19
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 19
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtgacca cccatggcga aggcccgat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780

eolf-seql.txt

```

ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agagtttgcg gaagtggcgt aa                          1362

```

<210> 20
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 20

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1          5          10          15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
          20          25          30

Thr Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
          50          55          60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
          85          90          95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
          100         105         110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
          115         120         125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
          130         135         140

```

eofl-seql.txt

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

eolf-seql.txt

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Glu
435 440 445

Phe Ala Glu Val Ala
450

<210> 21
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 21
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtaa cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cgccatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggccagtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
aaacatattg cggaactgcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

eolf-seql.txt

<210> 22
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 22

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Ala Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Ser Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

eolf-seql.txt

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 23
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 23
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc

eolf-seql.txt

```

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgtgc cgggctttct gcattctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgccggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccattcctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
aaacatattg cggaactcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agagtttgcg gaagtggcgt aa 1362

```

<210> 24
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 24

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

eofl-seql.txt

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320

eolf-seql.txt

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Glu
435 440 445

Phe Ala Glu Val Ala
450

<210> 25

<211> 1362

<212> DNA

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 25

atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatgggtg 180

gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240

ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300

gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360

aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420

cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480

accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540

tgcccgcatt attggcggtta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600

ctggcccgtg aactggaaga aaccattcag aaggaaggcg cgataccat tgcgggcttt 660

tttgcggaac cgggtgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720

eolf-seql.txt

```

gcgattctga ccacctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaactct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 26
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 26

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

eolf-seql.txt

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

eolf-seql.txt

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 27
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 27
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cgccatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
aaacatattg cggaactcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

eolf-seql.txt

<210> 28
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 28

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Ala Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

eolf-seql.txt

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
 385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 29

<211> 1362

<212> DNA

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

eolf-seql.txt

```

<400> 29
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc      60
cacatgccga gcctacgtca gcgtggcacc atgggtggtga cccatggcga aggcccgtat      120
atcgtgggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttgtga ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

```

<210> 30
<211> 453
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Engineered variant of V. fluvialis transaminase

```

```

<400> 30

```

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1           5           10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
          20          25          30

```

```

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

```

eolf-seql.txt

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Val Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320

eolf-seql.txt

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 31
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 31
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcatggccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttgtga ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cgataccat tgcgggcttt 660

eolf-seql.txt

```

tttgcggaac cggtgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggt gattttgggg      900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 32
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 32

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Val Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

eolf-seql.txt

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

eolf-seql.txt

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 33
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 33
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgaggg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320

ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa

1362

<210> 34

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 34

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

eolf-seql.txt

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 35
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>

<223> Engineered variant of *V. fluvialis* transaminase

<400> 35

```

atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc      60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat      120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgttttacgg ccgatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cgggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggt gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 36

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of *V. fluvialis* transaminase

<400> 36

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1           5           10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
          20          25          30

```

co1f-seql.txt

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

eolf-seql.txt

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 37
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 37
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600

eolf-seql.txt

```

ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctatttttcag      720
gcgattctga ccacacctgcg caaatatgat attccgggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 38
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 38

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

eolf-seql.txt

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335
 Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350
 Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365
 Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380
 Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
 385 390 395 400

eolf-seql.txt

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 39
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 39
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccggt aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgccggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260

eolf-seql.txt

gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 40
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase
<400> 40

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

eofl-seql.txt

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 41
<211> 1362
<212> DNA
<213> Artificial Sequence

eolf-seql.txt

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 41

```
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc      60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgatat      120
atcgtgggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg      240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgtgc cgggctttct gcatttgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggt gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362
```

<210> 42

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 42

```
Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1           5           10           15
```

```
Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
          20           25           30
```

eolf-seql.txt

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

eolf-seql.txt

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 43
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 43
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttgtga ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540

eolf-seql.txt

```

tgcccgcat attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 44
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 44

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

eo1f-seql.txt

Val Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

eofl-seql.txt

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 45
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 45
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200

eolf-seql.txt

gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 46
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 46

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

eolf-seql.txt

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 47
<211> 1362

eolf-seql.txt

<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 47
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atgggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgtttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatatttggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
ctggaagtgg tgaaagataa agcgagcaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 48
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 48

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

eofl-seql.txt

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30
 Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45
 Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

eolf-seql.txt

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 49
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 49
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtaggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttgtga ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcgtagcgtat catggctcca ccgcggtgag cgcgagcatg 480

eolf-seql.txt

```

accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgctg      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcggtt ttttttcag taggagcggg gattttgggg      900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgttttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 50
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 50

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

eolf-seql.txt

Val Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Arg Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

eolf-seql.txt

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 51
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 51
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgttttacgg cgccatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttgtga ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgaggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccattcctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140

eolf-seql.txt

ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttcttg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 52
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 52

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Ala Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Val Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

eofl-seql.txt

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

eolf-seql.txt

<210> 53
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 53
 atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
 cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
 atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
 gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgtttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
 tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatgg tgcgggcggg gttattaccg cgccaaaagg ctattttcag 720
 gcgattctga ccatcctgcg caaatatgct attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 54
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 54

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

eolf-seql.txt

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30
 Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45
 Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Ala Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

eolf-seql.txt

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 55
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 55
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattgccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420

eolf-seql.txt

```

cgtaaaattc tgaccctgaa gaacgcgtat catggctcca cgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgtgc cgggctttct gcatctgacc      540
tgcccgcatc attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcbc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcbt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcb     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 56
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 56

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95

eo1f-seq1.txt

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

eolf-seql.txt

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 57
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 57
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacg gccatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcat attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgcgggcggt gttattacc cgcgaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080

eolf-seql.txt

```

aaacatatg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

```

<210> 58
<211> 453
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Engineered variant of V. fluvialis transaminase

```

```

<400> 58

```

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1          5          10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
          20          25          30

```

```

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

```

```

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
          50          55          60

```

```

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

```

```

Gly Tyr His Ala Phe Tyr Gly Ala Met Ser Asp Gln Thr Val Met Leu
          85          90          95

```

```

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
          100          105          110

```

```

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
          115          120          125

```

```

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
          130          135          140

```

```

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
          145          150          155          160

```

```

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
          165          170          175

```

```

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
          180          185          190

```


eolf-seql.txt

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

eolf-seql.txt

<210> 59
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 59
 atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
 cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
 atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
 aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gcgtgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgtgc cgggctttct gcattctgacc 540
 tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatgg tgccggcggt gttattacc cgcctaaagg ctattttcag 720
 gcgattctga ccattcctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 60
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 60

eolf-seql.txt

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Arg Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

eolf-seql.txt

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
 385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 61
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 61
 atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
 cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
 atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360

eolf-seql.txt

```

aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 62
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 62

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

eolf-seql.txt

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

eolf-seql.txt

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 63
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 63
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcatctgacc 540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cgataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagcg ccaaaaacct aaccgcgggt tttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020

eolf-seql.txt

```

atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg      1080
aacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gtcgaccgat gggccagagc      1260
gtggttctgg tgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 64
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 64

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
165 170 175

eolf-seql.txt

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335
 Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350
 Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365
 Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380
 Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
 385 390 395 400
 Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415
 Met Gly Gln Ser Val Val Leu Val Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430
 Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
450

<210> 65
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

```
<400> 65
atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc atggtgacca cccatggcga aggcccgat      120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg      240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcattctgacc      540
tgcccgcatc attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt      600
ctggcccggt aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgccggcggt gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgag      840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag     1140
ctggaagccg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362
```

<210> 66
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

eolf-seql.txt

<400> 66

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Thr Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270

eolf-seql.txt

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 67
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 67
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300

eolf-seql.txt

```

gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatattgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 68
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 68

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

eofl-seql.txt

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

eolf-seql.txt

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 69
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 69
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttgatgg ccgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggctttc cgtataacag caagtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttgggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960

eolf-seql.txt

```

tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg      1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg      1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagccg tgaaagataa agcgagcaaa gcccgttttg atggcaacct gagcgtgggt      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 70
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 70

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Lys Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

eolf-seql.txt

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335
 Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350
 Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365
 Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
 370 375 380
 Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
 385 390 395 400
 Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415
 Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430
 Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
450

<210> 71
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

```

<400> 71
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat      120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcatctgacc      540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccggt aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggt gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgttttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 72
<211> 453
<212> PRT
<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 72

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1      5      10      15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
      20      25      30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
      35      40      45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
      50      55      60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65      70      75      80

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
      85      90      95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
      100      105      110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
      115      120      125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
      130      135      140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145      150      155      160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
      165      170      175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
      180      185      190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
      195      200      205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
      210      215      220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225      230      235      240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
      245      250      255

```

eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 73
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 73
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattgccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240

eolf-seql.txt

```

ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcggggcgt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatattgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagccg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 74
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 74

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

eolf-seql.txt

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

eolf-seql.txt

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 75
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 75
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggctttc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900

eolf-seql.txt

```

ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg      1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg      1080
aaacatatatg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agagtttgcg gaagtggcgt aa                          1362

```

<210> 76
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 76

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

eof-seql.txt

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Glu
435 440 445

Phe Ala Glu Val Ala
450

<210> 77
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 77
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggctttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 78
<211> 453
<212> PRT
<213> Artificial Sequence

eolf-seql.txt

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 78

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 79

<211> 1362

<212> DNA

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 79

atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatgggtg 180

eolf-seql.txt

```

gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttgatgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggctttc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggg gttattacc cgcctaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagcg ccaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgct gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 80
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 80

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

eofl-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Asp Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Phe Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 81
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 81
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
ctggtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgta taacatggtg 180
gcgggctttg atcataaagg cctgattgcg gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840

eolf-seql.txt

```
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga atgaggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggcct tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttcttg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa                        1362
```

<210> 82
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 82

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Ala Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

eolf-seql.txt

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Asp Val Ala
 450

<210> 83
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 83
 atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
 cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
 ctggtggatg tgcattggca cgttatctg gatgcgaaca gcggcctgta taacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
 tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatgg tgcgggcggg gttattacc cgcctaaagg ctattttcag 720
 gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaactctt ggaaaccgcg attgaagcga ttggggaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
 aaacatattg cggaacttcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct ggttgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttatcg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 84
 <211> 453

eolf-seql.txt

<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 84

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Asn Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

eolf-seql.txt

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Ile Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 85
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 85
atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
Page 113

eolf-seql.txt

```

ctggtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgta taacatgggtg      180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg      240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 86
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 86

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
 50 55 60

eolf-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 87
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 87
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
ctggtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgta taacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattacc ccgcaaaagg ctattttcag 720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780

eolf-seql.txt

```

ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atccaaacct gtgggtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacg gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 88
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 88

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1          5          10          15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
          20          25          30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
          35          40          45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
          50          55          60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
          85          90          95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
          100         105         110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
          115         120         125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
          130         135         140

```


eofl-seql.txt

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Pro Asn Leu Trp Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

eolf-seql.txt

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 89
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 89
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
ctggtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcat attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccggt aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga ttgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
aaacatattg cggaactgcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct ggttgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa 1362

eolf-seql.txt

<210> 90
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 90

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

eolf-seql.txt

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 91
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 91
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc

eolf-seql.txt

```

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
ctggtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgta taacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgccggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactgtc ggaaaccgcg attgaagcga atggggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gggtgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttatcg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa 1362

```

<210> 92
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 92

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

eolf-seql.txt

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Gly Glu Phe Asp His Gly
305 310 315 320

eolf-seql.txt

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Ile Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 93

<211> 1362

<212> DNA

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 93

atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120

ctggtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgta taacatgggtg 180

gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240

ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300

gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360

aacgatacca tgggtgaaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420

cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480

accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540

tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600

ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660

tttgcggaac cgggtgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720

eolf-seql.txt

```

gcgattctga ccacctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttatcg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 94
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 94

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

eolf-seql.txt

Thr	Leu	Lys	Asn	Ala	Tyr	His	Gly	Ser	Thr	Ala	Val	Ser	Ala	Ser	Met
145					150					155					160
Thr	Gly	Leu	Pro	Tyr	Asn	Ser	Val	Phe	Gly	Leu	Pro	Leu	Pro	Gly	Phe
				165					170					175	
Leu	His	Leu	Thr	Cys	Pro	His	Tyr	Trp	Arg	Tyr	Gly	Glu	Glu	Gly	Glu
			180					185					190		
Thr	Glu	Glu	Gln	Phe	Val	Ala	Arg	Leu	Ala	Arg	Glu	Leu	Glu	Glu	Thr
		195					200					205			
Ile	Gln	Lys	Glu	Gly	Ala	Asp	Thr	Ile	Ala	Gly	Phe	Phe	Ala	Glu	Pro
	210					215					220				
Val	Met	Gly	Ala	Gly	Gly	Val	Ile	Thr	Pro	Pro	Lys	Gly	Tyr	Phe	Gln
225					230					235					240
Ala	Ile	Leu	Thr	Ile	Leu	Arg	Lys	Tyr	Asp	Ile	Pro	Val	Ile	Ser	Asp
				245					250					255	
Glu	Val	Ile	Cys	Gly	Phe	Gly	Arg	Thr	Gly	Asn	Thr	Trp	Gly	Cys	Val
			260					265					270		
Thr	Tyr	Asp	Phe	Thr	Pro	Asp	Ala	Ile	Ile	Ser	Ser	Lys	Asn	Leu	Thr
		275					280					285			
Ala	Gly	Phe	Phe	Pro	Val	Gly	Ala	Val	Ile	Leu	Gly	Pro	Glu	Leu	Ser
	290					295					300				
Lys	Arg	Leu	Glu	Thr	Ala	Ile	Glu	Ala	Ile	Gly	Glu	Phe	Asp	His	Gly
305					310					315					320
Phe	Thr	Thr	Gly	Gly	His	Pro	Val	Gly	Cys	Ala	Ile	Ala	Leu	Lys	Ala
				325					330					335	
Ile	Asp	Val	Val	Met	Asn	Glu	Gly	Leu	Ala	Glu	Asn	Val	Arg	Arg	Leu
			340					345					350		
Ala	Pro	Arg	Phe	Glu	Glu	Arg	Leu	Lys	His	Ile	Ala	Glu	Arg	Pro	Asn
		355					360					365			
Ile	Gly	Glu	Tyr	Arg	Gly	Ile	Gly	Phe	Met	Trp	Ala	Leu	Glu	Val	Val
	370					375					380				
Lys	Asp	Lys	Ala	Ser	Lys	Ala	Pro	Phe	Asp	Gly	Asn	Leu	Val	Val	Ser
385					390					395					400
Glu	Arg	Ile	Ala	Asn	Thr	Cys	Thr	Asp	Leu	Gly	Leu	Ile	Cys	Arg	Pro
				405					410					415	

eolf-seql.txt

Leu Gly Gln Ser Val Val Ile Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 95
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 95
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
ctggtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgta taacatggtg 180
gcgggctttg atcataaagg cctgattgcg gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttcc aagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgtttcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga atgaggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
aaacatattg cggaactgcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gtgggtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggcagcagc 1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

eolf-seql.txt

<210> 96
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 96

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Tyr Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Pro Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu Phe Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

eolf-seql.txt

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Glu Glu Phe Asp His Gly
 305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Trp Val Ser
 385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415

Leu Gly Ser Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 97

<211> 1362

<212> DNA

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

eolf-seql.txt

```

<400> 97
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat      120
atcgtgggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatgggtg      180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtgggtt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggaagaaga ggcgaaccg aagaacagtt tgcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga atgaggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaactctg      1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atccaaacct gtgggtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtggttgcg gatgtggcgt aa                        1362

```

```

<210> 98
<211> 453
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Engineered variant of V. fluvialis transaminase

```

<400> 98

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1           5           10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
          20          25          30

```

```

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

```

eolf-seql.txt

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Glu Glu Phe Asp His Gly
 305 310 315 320

eolf-seql.txt

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Pro Asn Leu Trp Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 99
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 99
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cgataccat tgcgggcttt 660

eolf-seql.txt

```

tttgcggaac cggtgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggt gattttgggg      900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct gttggtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggcagcagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 100
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 100

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

eolf-seql.txt

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Leu Val Ser
385 390 395 400

eolf-seql.txt

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Ser Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 101
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 101
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgat 120
ctggtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cgccatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cagctttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag 720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgctttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gggtgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320

ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa

1362

<210> 102

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 102

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Leu Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Ala Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Ser Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

eolf-seql.txt

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 103
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 103

```

atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgat      120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgttttacgg ccgatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cagctttggc ctgccgctgc cgggctttct gcatctgacc      540
tgccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggg gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggcagcagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 104

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 104

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1          5          10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
          20          25          30

```

co1f-seql.txt

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Ser Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

eolf-seql.txt

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Ser Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 105
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 105
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcatgtgtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600

eolf-seql.txt

```

ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctatttttcag      720
gcgattctga ccacctctgcg caaatatgat attccgggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga atgaggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gtgggtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 106
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 106

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

eolf-seql.txt

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Trp Val Ser
385 390 395 400

eolf-seql.txt

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 107
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 107
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260

eolf-seql.txt

gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 108
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase
<400> 108

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

eofl-seql.txt

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 109
<211> 1362
<212> DNA
<213> Artificial Sequence

eolf-seql.txt

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 109

```

atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat      120
atcgtgggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg      240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag caagtttggc ctgccgtgc cgggctttct gcattctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgccggcggt gttattacc cgcctaaagg ctattttcag      720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaactct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaactctg     1080
aaacatattg cggaactcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gtgggtgagc     1200
gaacgtattg cgaacacct caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa                        1362

```

<210> 110

<211> 453

<212> PRT

<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 110

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1          5          10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
          20          25          30

```

eolf-seql.txt

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Lys Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

eolf-seql.txt

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Trp Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 111
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 111
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540

eolf-seql.txt

```

tgcccgcat attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga atgaggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 112
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 112

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

eolf-seql.txt

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

eolf-seql.txt

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 113
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 113
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga ttggggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtgggtg 1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gttggtgagc 1200

eolf-seql.txt

gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 114
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 114

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

eolf-seql.txt

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Leu Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 115
<211> 1362

eolf-seql.txt

<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

```
<400> 115
atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat      120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaattg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gggtgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtggttgcg gatgtggcgt aa                        1362
```

<210> 116
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 116

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

eofl-seql.txt

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30
 Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45
 Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

eolf-seql.txt

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 117
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 117
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gctggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcaggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480

eolf-seql.txt

```

accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcggt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcggtt tttttccag taggagcggt gattttgggg      900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga ttggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gtgggtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gatgtggcgt aa                        1362

```

<210> 118
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 118

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

eolf-seql.txt

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335
 Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
 340 345 350
 Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365
 Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380

eolf-seql.txt

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Trp Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Asp Val Ala
450

<210> 119
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 119
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgaggg tgcgggcggt gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccattcctgc caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140

eolf-seql.txt

ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttcttg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 120
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 120

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

eofl-seql.txt

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

eolf-seql.txt

<210> 121
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 121
 atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
 cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
 atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
 tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatgg tgccggcggt gttattaccg cgccaaaagg ctattttcag 720
 gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttgaggaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gttggtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 122
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 122

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

eolf-seql.txt

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30
 Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45
 Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60
 His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285

eolf-seql.txt

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Leu Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 123
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 123
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattgccg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420

eolf-seql.txt

```

cgtaaaattc tgaccctgaa gaacgcgtat catggctcca cgcggtgag cgcgagcatg      480
accggccttc cgtataacag caagtttggc ctgccgtgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga atggggaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct ggttgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 124
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 124

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

eofl-seql.txt

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Lys Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

eolf-seql.txt

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 125
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 125
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgttttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcat attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga ttggggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080

eolf-seql.txt

```

aaacatatg cggaacgtcc gaacattggc gaatatcgtg gtattggcctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggtaacct ggttgtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttatcg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

```

<210> 126
<211> 453
<212> PRT
<213> Artificial Sequence

```

```

<220>
<223> Engineered variant of V. fluvialis transaminase
<400> 126

```

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1          5          10          15

```

```

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
          20          25          30

```

```

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

```

```

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50          55          60

```

```

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

```

```

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
          85          90          95

```

```

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
          100          105          110

```

```

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
          115          120          125

```

```

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
          130          135          140

```

```

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
          145          150          155          160

```

```

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
          165          170          175

```

```

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
          180          185          190

```

eolf-seql.txt

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Ile Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

eolf-seql.txt

<210> 127
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 127
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc tcagtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga atggggaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
atgaacgaag gcctgctgga aaacgtgcgt cgtctggccc cgcgttttga agaactcttg 1080
aaacatattg cggaactgcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggtaacct gggtgtgagc 1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggcagcagc 1260
gtggttctgg cgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 128
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 128

eo1f-seq1.txt

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Ser Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

eolf-seql.txt

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Asn Gly Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Leu Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Val Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Ser Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 129
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 129
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360

eolf-seql.txt

```

aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggccccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggtgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagcg ccaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gtcgaccgat gggccagAAC     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 130
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 130

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

eolf-seql.txt

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

eolf-seql.txt

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Met Gly Gln Asn Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 131
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 131
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cgataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gatatttgggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020

eolf-seql.txt

```

atgaacgaag gcctggccga aaacgtgctg cgtctggccc cgcgttttga agaacgtctg      1080
aacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gtcgaccgct gggccagAAC      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 132
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 132

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
165 170 175

eolf-seql.txt

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Asn Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 133
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

```

<400> 133
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc      60
cacatgccga gcctacatca gcgtggcacc atggtggttaa cccatggcga agggccgtat      120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg      240
ggctatcatg cgttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatc attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggag     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg tgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 134
<211> 453
<212> PRT
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

eolf-seql.txt

<400> 134

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270

eolf-seql.txt

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Val Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 135
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 135
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggttaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300

eolf-seql.txt

```

gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggtgatggg tgcgggcggt gttattaccg cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gtcgaccgat gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 136
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 136

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

eof-seql.txt

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

eolf-seql.txt

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Met Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 137
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 137
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctggt caacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg ccgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgacctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcacgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatgg tgcgggcggt gttattacc cgcctaaagg ctattttcag 720
gcgattctga ccattctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960

eolf-seql.txt

```

tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg      1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg      1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 138
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 138

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175

eolf-seql.txt

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335
 Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
 340 345 350
 Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
 355 360 365
 Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
 370 375 380
 Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
 385 390 395 400
 Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
 405 410 415
 Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
 420 425 430
 Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
450

<210> 139
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of *V. fluvialis* transaminase

```

<400> 139
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc      60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgatat      120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg      180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg      240
ggctatcatg cgtttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcgcta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggccccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctatttttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggcttttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagcg ccaaaaacct aaccgcgggt tttttttccag taggagcggg gatatttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgtttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagccg tgaaagataa agcgagcaaa gcccgttttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttcttg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 140
<211> 453
<212> PRT
<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 140

```

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1          5          10          15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
          20          25          30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
          35          40          45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50          55          60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65          70          75          80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
          85          90          95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100          105          110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115          120          125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130          135          140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145          150          155          160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165          170          175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180          185          190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195          200          205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210          215          220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225          230          235          240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
          245          250          255

```

eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 141
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 141
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
atcgtggatg tgcattgccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240

eolf-seql.txt

```

ggctatcatg cgttttacgg ccgcatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcggggcgt gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg      840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatattgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagccg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg tgccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 142
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 142

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

eolf-seql.txt

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

eolf-seql.txt

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Val Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 143
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 143
atgaacaaac cgcagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcggggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900

eolf-seql.txt

```

ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg      1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg      1080
aaacatatatg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg      1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc      1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc      1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa      1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 144
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 144

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

eof-seq1.txt

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 145
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 145
 atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
 cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
 atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tgggtgaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggcatcc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
 tgcccgcatt attggcgtaa tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatggg tgcgggcggt gttattaccc cgccaaaagg ctattttcag 720
 gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagcg ccaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttgggg 900
 ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg 1080
 aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg 1140
 ctggaagccg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gtcgaccgct gggccagaac 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 146
 <211> 453
 <212> PRT
 <213> Artificial Sequence

eolf-seql.txt

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 146

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

eolf-seql.txt

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Asn Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 147
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 147
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60

cacatgccga gcctacatca gcgtggcacc atggtgacca cccatggcga aggcccgtat 120

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180

eolf-seql.txt

```

gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg      240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg      360
aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catgggtcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc      540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgcgggcggg gttattaccc cgccaaaagg ctattttcag      720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagca gtaaaaacct aaccgcgggt tttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgagc     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                          1362

```

<210> 148
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 148

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Thr Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

eofl-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80
 Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
 85 90 95
 Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110
 Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125
 Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140
 Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160
 Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
 165 170 175
 Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
 180 185 190
 Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
 195 200 205
 Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
 210 215 220
 Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
 225 230 235 240
 Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
 245 250 255
 Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
 260 265 270
 Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
 275 280 285
 Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
 290 295 300
 Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
 305 310 315 320
 Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
 325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 149
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 149
atgaacaaac cgcagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60
cacatgccga gcctacatca gcgtggcacc atggtggtaa cccatggcga aggcccgtat 120
atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtt caacatgggtg 180
gcgggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgttttccg 240
ggctatcatg cgtttttacgg cctgatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
gtggaagtga gcccgtttga tagcggccgt gtgtttctga ccaacagcgg cagcgaagcg 360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
cgtaaaattc tgaccctgaa gcaggcgtat catggctcca ccgcggtgag cgcgagcatg 480
accggcatcc cgtataacag cgtgtttggc ctgccggctc cgggctttct gcatctgacc 540
tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgtcgcgcgt 600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
tttgcggaac cggatgatggg tgcgggcggg gttattaccc cgccaaaagg ctattttcag 720
gcgattctga ccatcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg 840

eolf-seql.txt

```
attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gattttgggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggcct tatgtgggcg     1140
ctggaagccg tgaaagataa agcgagcaaa gccccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttcttg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362
```

<210> 150
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 150

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
 50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
 65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Leu Met Ser Asp Gln Thr Val Met Leu
 85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
 100 105 110

Leu Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
 115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
 130 135 140

Thr Leu Lys Gln Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
 145 150 155 160

eolf-seql.txt

Thr Gly Ile Pro Tyr Asn Ser Val Phe Gly Leu Pro Ala Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Ala Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

eolf-seql.txt

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
 435 440 445

Phe Ala Glu Val Ala
 450

<210> 151
 <211> 1362
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase

<400> 151
 atgaacaaac cgagagctg ggaaacgcgt gcggaaacct atagcctgta tgcctttacc 60
 cacatgccga gcctacatca gcgtggcacc atggtggtga cccatggcga aggcccgat 120
 atcgtggatg tgcattggcg tcgttatctg gatgcgaaca gcggcctgtt caacatggtg 180
 gcgggctttg atcataaagg cctgattgat gcggcgaaag cgagtatga acgttttccg 240
 ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg 300
 gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg 360
 aacgatacca tggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa 420
 cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg 480
 accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcattctgacc 540
 tgcccgcatt attggcgta tggcgaagaa ggcgaaaccg aagaacagtt tgcgcgcgt 600
 ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt 660
 tttgcggaac cggatgatggg tgcgggcggg gttattaccg cgccaaaagg ctattttcag 720
 gcgattctga ccattcctgcg caaatatgat attccggtga tcagcgatga agtgatttgc 780
 ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccgatgacg 840
 attattagca gtaaaaacct aaccgcgggt ttttttccag taggagcggg gatatttggg 900
 ccggaactga gcaaactctt ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt 960
 tttaccacag gcggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg 1020
 atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaactctg 1080
 aaacatattg cggaactcc gaacattggc gaatatcgtg gtattggctt tatgtgggag 1140
 ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgagc 1200
 gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc 1260
 gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa 1320
 ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa 1362

<210> 152
 <211> 453

eolf-seql.txt

<212> PRT
<213> Artificial Sequence

<220>

<223> Engineered variant of V. fluvialis transaminase

<400> 152

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Tyr Ser Leu
1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu His Gln Arg Gly Thr Met Val
20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Phe Asn Met Val Ala Gly Phe Asp
50 55 60

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

eolf-seql.txt

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ser Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr Gly Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Ser
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450

<210> 153
<211> 1362
<212> DNA
<213> Artificial Sequence

<220>
<223> Engineered variant of V. fluvialis transaminase

<400> 153
atgaacaaac cgagagctg ggaaacgcgt gcggaaaccg tgagcctgta tgcctttacc 60

cacatgccga gcctacgtca gcgtggcacc atggtggtga cccatggcga aggcccgtat 120
Page 204

eolf-seql.txt

```

atcgtggatg tgcattggccg tcgttatctg gatgcgaaca gcggcctgtg gaacatgggtg      180
gcggggctttg atcataaagg cctgattgat gcggcgaaag cgcagtatga acgtttttccg      240
ggctatcatg cgttttacgg ccgtatgagc gatcagaccg tgatgctgtc tgaaaaactg      300
gtggaagtga gcccgtttga tagcggccgt gtgttttata ccaacagcgg cagcgaagcg      360
aacgatacca tgggtgaaaat gctgtggttt ctgcatcgtg cggaaggcaa accgcagaaa      420
cgtaaaattc tgaccctgaa gaacgcgtat catggctcca ccgcggtgag cgcgagcatg      480
accggccttc cgtataacag cgtgtttggc ctgccgctgc cgggctttct gcatctgacc      540
tgcccgcatc attggcggtt tggcgaagaa ggcgaaccg aagaacagtt tgtcgcgcgt      600
ctggcccgtg aactggaaga aaccattcag aaggaaggcg cggataccat tgcgggcttt      660
tttgcggaac cggatgatgg tgccggcggt gttattacc ccgcaaaagg ctattttcag      720
gcgattctga ccacctctgc caaatatgat attccggtga tcagcgatga agtgatttgc      780
ggctttggtc gtaccggcaa cacctggggc tgcgtgacct atgattttac cccggatgcg      840
attattagcg ccaaaaacct aaccgcgggt tttttccag taggagcggg gatatttggg      900
ccggaactga gcaaacgtct ggaaaccgcg attgaagcga tcgaagaatt tgatcatggt      960
tttaccacac acggccatcc ggtgggttgt gcgattgcgc tgaaagcgat tgatgtggtg     1020
atgaacgaag gcctggccga aaacgtgcgt cgtctggccc cgcgttttga agaacgtctg     1080
aaacatattg cggaacgtcc gaacattggc gaatatcgtg gtattggctt tatgtgggcg     1140
ctggaagtgg tgaaagataa agcgagcaaa gcccgtttg atggcaacct gagcgtgggt     1200
gaacgtattg cgaacacctg caccgatctg ggactgattt gccgaccgtt gggccagagc     1260
gtggttctgg caccgccgta cattctgacc gaagcgcaga tggatgaaat gttcgataaa     1320
ctggaaaaag cgctggataa agtgtttgcg gaagtggcgt aa                        1362

```

<210> 154
 <211> 453
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Engineered variant of V. fluvialis transaminase
 <400> 154

Met Asn Lys Pro Gln Ser Trp Glu Thr Arg Ala Glu Thr Val Ser Leu
 1 5 10 15

Tyr Ala Phe Thr His Met Pro Ser Leu Arg Gln Arg Gly Thr Met Val
 20 25 30

Val Thr His Gly Glu Gly Pro Tyr Ile Val Asp Val His Gly Arg Arg
 35 40 45

Tyr Leu Asp Ala Asn Ser Gly Leu Trp Asn Met Val Ala Gly Phe Asp
 50 55 60

eolf-seql.txt

His Lys Gly Leu Ile Asp Ala Ala Lys Ala Gln Tyr Glu Arg Phe Pro
65 70 75 80

Gly Tyr His Ala Phe Tyr Gly Arg Met Ser Asp Gln Thr Val Met Leu
85 90 95

Ser Glu Lys Leu Val Glu Val Ser Pro Phe Asp Ser Gly Arg Val Phe
100 105 110

Tyr Thr Asn Ser Gly Ser Glu Ala Asn Asp Thr Met Val Lys Met Leu
115 120 125

Trp Phe Leu His Arg Ala Glu Gly Lys Pro Gln Lys Arg Lys Ile Leu
130 135 140

Thr Leu Lys Asn Ala Tyr His Gly Ser Thr Ala Val Ser Ala Ser Met
145 150 155 160

Thr Gly Leu Pro Tyr Asn Ser Val Phe Gly Leu Pro Leu Pro Gly Phe
165 170 175

Leu His Leu Thr Cys Pro His Tyr Trp Arg Tyr Gly Glu Glu Gly Glu
180 185 190

Thr Glu Glu Gln Phe Val Ala Arg Leu Ala Arg Glu Leu Glu Glu Thr
195 200 205

Ile Gln Lys Glu Gly Ala Asp Thr Ile Ala Gly Phe Phe Ala Glu Pro
210 215 220

Val Met Gly Ala Gly Gly Val Ile Thr Pro Pro Lys Gly Tyr Phe Gln
225 230 235 240

Ala Ile Leu Thr Ile Leu Arg Lys Tyr Asp Ile Pro Val Ile Ser Asp
245 250 255

Glu Val Ile Cys Gly Phe Gly Arg Thr Gly Asn Thr Trp Gly Cys Val
260 265 270

Thr Tyr Asp Phe Thr Pro Asp Ala Ile Ile Ser Ala Lys Asn Leu Thr
275 280 285

Ala Gly Phe Phe Pro Val Gly Ala Val Ile Leu Gly Pro Glu Leu Ser
290 295 300

Lys Arg Leu Glu Thr Ala Ile Glu Ala Ile Glu Glu Phe Asp His Gly
305 310 315 320

Phe Thr Thr His Gly His Pro Val Gly Cys Ala Ile Ala Leu Lys Ala
325 330 335

eolf-seql.txt

Ile Asp Val Val Met Asn Glu Gly Leu Ala Glu Asn Val Arg Arg Leu
340 345 350

Ala Pro Arg Phe Glu Glu Arg Leu Lys His Ile Ala Glu Arg Pro Asn
355 360 365

Ile Gly Glu Tyr Arg Gly Ile Gly Phe Met Trp Ala Leu Glu Val Val
370 375 380

Lys Asp Lys Ala Ser Lys Ala Pro Phe Asp Gly Asn Leu Ser Val Gly
385 390 395 400

Glu Arg Ile Ala Asn Thr Cys Thr Asp Leu Gly Leu Ile Cys Arg Pro
405 410 415

Leu Gly Gln Ser Val Val Leu Ala Pro Pro Tyr Ile Leu Thr Glu Ala
420 425 430

Gln Met Asp Glu Met Phe Asp Lys Leu Glu Lys Ala Leu Asp Lys Val
435 440 445

Phe Ala Glu Val Ala
450