



(51) International Patent Classification:

A01N 43/38 (2006.01) C07D 209/58 (2006.01)
A61K 31/40 (2006.01)

(21) International Application Number:

PCT/US20 13/042 148

(22) International Filing Date:

22 May 2013 (22.05.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/650,039 22 May 2012 (22.05.2012) US
61/757,876 29 January 2013 (29.01.2013) US

(71) Applicant: TRUSTEES OF DARTMOUTH COLLEGE

[US/US]; Technology Transfer Office, 11 Rope Ferry
Road, Hanover, NH 03755-1404 (US).

(72) Inventor: WU, Jimmy; 17 Carriage Lane, Hanover, NH
03755 (US).

(74) Agents: LICATA, Jane, Massey et al; Licata & Tyrrell
P.C., 66 E. Main Street, Marlton, NJ 08053 (US).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every

kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: METHOD FOR SYNTHESIZING CYCLOALKANYL(5) INDOLES, CYCLOALKANYL(5) BENZOFURANS, CYCLOALKANYL(5)BENZOTHIOPHENES, COMPOUNDS AND METHODS OF USE

(57) Abstract: A method of synthesizing cycloalkanyl [b] indoles, as well as related cycloalkanyl [b] benzofurans and cycloalkanyl [b] benzothiophenes is provided. The method is a single, multicomponent reaction that combines (1) an indole, benzofuran, or benzothiophene, (2) an aldehyde, ketone, or ketal, and (3) a diene in the presence of an acid, in particular a Ga(III) or In (III) salt. Compositions and methods of using these compounds to stimulate secretion and/or production of glucagon-like peptide-1 or inhibit the activity of Calcitonin Gene-Related Peptide receptor are also provided.



WO 2013/177241 A1

**Method for Synthesizing Cycloalkanyl [\Rightarrow] indoles ,
Cycloalkanyl [b]benzofurans , Cycloalkanyl [b]benzothiophenes ,
Compounds and Methods of Use**

Introduction


[0001] This invention was made with government support under contract numbers CHE-1052824 awarded by the National Science Foundation. The government has certain rights in the invention.

Background of the Invention



[0002] Cyclohepta [b] indoles (1) exhibit a broad spectrum of biological activity. For instance, compound 2 is a potent inhibitor ($IC_{50} = 63$ nM) of SIRT1, a member of the class III histone deacetylases (HDAC; Napper, et al. (2005) *J. Med. Chem.* 48:8045). SIRT1 can effectively deacetylate p53 and has also been implicated in the regulation of apoptosis. With an IC_{50} of 100 nM, compound 3 inhibits the production of leukotriene B_4 (LTB_4), which is involved in various inflammatory responses (Kuehm-Caubere, et al. (1999) *Eur. J. Med. Chem.* 34:51). A third molecule, compound 4, inhibits adipocyte fatty-acid binding protein (A-FABP) with an IC_{50} of 450 nM (Barf, et al. (2009) *Bioorg. Med. Chem. Lett.* 19:1745).



1, cyclohepta[b]indole



2, SIRT1 Inhibitor




3, A-FABP Inhibitor**4, LTB₄ Production Inhibitor**

[0003] The biology of cyclohepta [*b*]indoles, as well as cyclopenta- and cyclohexa [*b*]indoles, has attracted considerable interest from the pharmaceutical industry as potential therapeutics. In this respect, various compounds with this structural motif have been described. See, e.g., WO 2010/036998, US 2011/0152306, US 2009/0170923, US 2009/0156621, WO 2004/063156, WO 2010/054382, WO 2006/047017, WO 2006/034090, WO 2004/069831, EP 1184373, WO 2009/120720, WO 2005/023245, WO 2004/110999, WO 2005/094833, WO 03/091257, US 2008/0027090, WO 2011/044134, US 2011/0003737, WO 2010/111483, US 2007/0037791, EP 1505061, WO 2006/055760 and WO 2008/021364.

[0004] The preparation of cycloalka [*b*]indoles often includes the Fisher indole synthesis which, while quite useful, possesses certain limitations (Ambekar (1983) *Curr. Sci.* 52:578; Robinson (1969) *Chem. Rev.* 69:227; Inman & Moody (2011) *Chem. Commun.* 47:788). These include the need to make the requisite hydrazine and ketone starting materials. Regioselectivity with unsymmetrical ketones can also be problematic. Finally, electron-withdrawing groups on the aromatic hydrazine can substantially attenuate reactivity. Other methods for preparing cycloalka [*b*]indoles are known but have not been extensively explored (Willis, et al. (2005) *Angew. Chem. Int. Ed.* 44:403; Barluenga, et al. (2007) *Angew. Chem. Int. Ed.* 46:1529; Sun, et al. (2011) *Angew. Chem. Int. Ed.* 50:1702; Liu & Widenhoefer (2004) *J. Am. Chem. Soc.* 126:10250; Ragains & Winkler (2006) *Org. Lett.* 8:4437; Silvanus, et al. (2009) *Org. Lett.* 11:1175; Ishikura & Kato (2002) *Tetrahedron* 58: 9827).

[0005] (4+3) Cycloaddition reactions have been explored in some detail (Harmata (2010) *Chem. Comm.* 46:8886; Harmata (2001) *Acc. Chem. Res.* 34:595; Harmata (2006) *Adv. Synth.*

Catal. 348:2297; Battiste, et al. (2006) *Chem. Eur. J.* 12:3438; Hartung & Hoffman (2004) *Angew. Chem., Int. Ed.* 43:1934; Harmata (1997) *Tetrahedron* 53:6235; Chung, et al. (2009) *J. Am. Chem. Soc.* 131:4556; Harmata, et al. (1996) *J. Am. Chem. Soc.* 118:2860; Nilson & Funk (2011) *J. Am. Chem. Soc.* 133:12451; Yu, et al. (2010) *Org. Lett.* 12:5135; Liu & Chiu (2011) *Chem. Commun.* 47:3416; Lee, et al. (1998) *J. Org. Chem.* 63:2804; Lee & Cha (2000) *Tetrahedron* 56:10175; Davies & Dai (2004) *J. Am. Chem. Soc.* 126:2692). A stabilizing group is usually present at C2 of the 2n component (Scheme 1).




SCHEME 1

[0006] Heteroatom substitution with sulfur, oxygen, and halide at the terminal ends of the allylic cation are known (Xiong, et al. (2003) *J. Am. Chem. Soc.* 125:12694; Lohse & Hsung (2011) *Chem. Eur. J.* 17:3812; Lohse, et al. (2011) *J. Org. Chem.* 76:3246; Harmata (2010) *Chem. Commun.* 46:8904; Jeffrey, et al. (2011) *J. Am. Chem. Soc.* 133:7688; Lee & Cha (2001) *J. Am. Chem. Soc.* 123:3243; Aungst, Jr. & Funk (2001) *Org. Lett.* 3:3553; Harmata & Wacharasindhu (2005) *Org. Lett.* 7:2563; Lee & Cha (1999) *Org. Lett.* 1:523; Blackburn, et al. (1983) *Can. J. Chem.* 61:1981; Harmata, et al. (2004) *Heterocycles* 62:583; Harmata & Gamlath (1988) *J. Org. Chem.* 53:6154; Hardinger, et al. (1995) *J. Org. Chem.* 60:1104; Sasaki, et al. (1982) *Tetrahedron Lett.* 23:1693). However, cycloaddition reactions utilizing substrates with nitrogen-based substituents have are rare (Xiong, et al. (2003) *J. Am.*


Chem. Soc. 125:12694; Lohse & Hsung (2011) *Chem. Eur.* 17:3812; Lohse, et al. (2011) *J. Org. Chem.* 76:3246).

Summary of the Invention


[0007] This invention is a method of synthesizing a cycloalkanyl ^[a]indole, cycloalkanyl ^[b]benzofuran or cycloalkanyl ^[b]benzothiophene by combining, in a single reaction, (a) an indole, benzofuran or benzothiophene; (b) a ketone, aldehyde or ketal; and (c) a coupling partner and adding an acid catalyst thereby producing a cycloalkanyl compounds of Formula Va, Vb, Vc or Vd.




Formula Va



Formula Vb




Formula Vc




Formula Vd

In certain embodiments, the acid catalyst is a Lewis acid, e.g., a metal halide or triflate salt of Ga(III) or In(III); a chiral Lewis acid complex; a Brønsted acid; a chiral phosphoric acid; or other chiral Brønsted acid.


[0008] A compound of Formula Ve, Vf or Vg is also provided,



Formula Ve



Formula Vf




Formula Vg

as are methods of using such a compound to stimulate the secretion and/or production of glucagon-like peptide-1 (GLP-1)

and/or inhibit the activity of the Calcitonin Gene-Related Peptide (CGRP) receptor, and treat disease.

Detailed Description of the Invention


[0009] It has now been found that certain metal salts and Brønsted acids effectively mediate regio- and diastereoselective three-component (4+3) cycloaddition reactions to furnish cyclohepta [b]indoles in high yields (Scheme 2). These reactions occur in a single step at room temperature without the need for Schlenk techniques, glove boxes, or inert atmosphere. Because each of the three coupling components (*i.e.*, indole/furan/thiophene, aldehyde/ketone, coupling partner) can be independently varied, this methodology provides rapid access to a library of diverse cycloalkanyl [b]indole/benzofuran/benzothiophene derivatives.




SCHEME 2

[0010] Given the data presented herein, this invention is a method for synthesizing cycloalkanyl [b]indoles, as well as related cycloalkanyl [b]benzofurans and cycloalkanyl [b]benzothiophenes, in acid-catalyzed three-component (4+3), (3+3) and (3+2) cycloaddition reactions. The method of the invention involves combining, in a single,


multi component reaction, (1) an indole/furan/thiophene; (2) a ketone, or aldehyde or ketal; and (3) an appropriate coupling partner in the presence of an acid catalyst to produce a compound of Formula Va, Vb, Vc or Vd.




Formula Va



Formula Vb




Formula Vc




Formula Vd


[0011] More specifically, the method of the invention includes the steps of combining in a single reaction: (1) a compound of Formula I; (2) a ketone of Formula II, or an aldehyde or ketal of Formula III; and (3) a coupling partner in the presence of an acid catalyst to produce a compound of Formula Va, Vb, Vc or Vd.




[0012] Coupling partners of use in the present invention include, but are not limited to, those represented by Formula IVa-IVf .




Formula IVa




Formula IVb




Formula IVc



Formula IVd



Formula IVe



Formula IVf

[0013] In reference to Formulae I-V, each X is the same or different and is independently $c-(R^1)_n$, OR^1 , S or NR^1 , wherein each R^1 is independently hydrogen, alkyl, alkenyl, alkynyl or aryl;

each Y is the same or different and is independently N or C;

each Z is the same or different and is independently N or C and wherein Z can optionally be connected to any other Z by a chain of one or more atoms independently selected from substituted or unsubstituted c, N, O, S, or a combination thereof, thereby forming an additional ring(s).

dashed bonds are independently present or absent such that this invention includes cyclopenta-, cyclohexa- and cyclohepta- compounds;

each R^2 is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7$ -cycloalkyl, aryl, $-NHR^7$, Het, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(O)R^8$, $-C(O)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(O)NR^8R^9$, $-C(O)aryl$, $-C(O)NR^8aryl$, $-C(O)Het$, $-C(O)NHR^7Het$, $-R^7C(O)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(O)_2NR^8R^9$, $-S(O)_2NR^8aryl$, $-R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

R^3 and R^4 are independently hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl,

cycloalkenyl, Het, aryl, $-NHR^7$, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(0)R^8$, $-C(0)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(0)NR^8R^9$, $-C(0)aryl$, $-C(0)NR^8aryl$, $-C(0)Het$, $-C(0)NHR^7Het$, $-R^7C(0)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(0)_2NR^8R^9$, $-S(0)_2N$ R^8aryl , $-R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(0)_mR^8$, cyano, nitro, or azido group, or

R^3 and R^4 together form a $C_{4,6}$ cycloalkyl;

R^5 is hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het or aryl group;

each R^6 is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, $-NHR^7$, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(0)R^8$, $-C(0)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(0)NR^8R^9$, $-C(0)aryl$, $-C(0)NR^8aryl$, $-C(0)Het$, $-C(0)NHR^7Het$, $-R^7C(0)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(0)_2NR^8R^9$, $-S(0)_2N$ R^8aryl , $-R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(0)_mR^8$, cyano, nitro, or azido group;

each R^7 is the same or different and is independently selected from an alkylene, cycloalkylene, alkenylene, cycloalkenylene or alkynylene group;

each of R^8 and R^9 are the same or different and are independently selected from the group of hydrogen, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7cycloalkyl$, $-R^7OH$, $-R^7(OR^7)_w$, or $-R^7NR^{10}R^{11}$ group;

each of R^{10} and R^{11} are the same or different and are independently selected from the group of an alkyl, cycloalkyl, alkenyl, cycloalkenyl, or alkynyl group;

each R^{12} is independently is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl,

cycloalkenyl, Het, aryl, $-NHR^7$, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(0)R^8$, $-C(0)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(0)NR^8R^9$, $-C(0)aryl$, $-C(0)NR^8aryl$, $-C(0)Het$, $-C(0)NHR^7Het$, $-R^7C(0)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(0)_2NR^8R^9$, $-S(0)_2N$ R^8aryl , $-R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(0)_mR^8$, cyano, nitro, or azido group;

each R^{13} is independently $C-(R^1)_n$ or OR^1 ;

p is selected from 0, 1, 2, 3, or 4;

q is selected from 0, 1, 2, 3, or 4;

each n independently is 0, 1 or 2;

each m independently is 0, 1 or 2;

w is 1-10; and

LG is leaving group.

[0014] In some embodiments, at least two of the three dashed bonds in Formula Va are present. In other embodiments, all three dashed bonds in Formula Va are present.

[0015] Het represents an optionally substituted 5- or 6-membered heterocyclyl or heteroaryl group.

[0016] As used herein, the term "alkyl" refers to a straight or branched chain hydrocarbon, preferably having from one to twelve carbon atoms. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl, n-butyl, tert-butyl, isopentyl, n-pentyl, and the like.

[0017] The term "alkenyl" refers to a straight or branched chain aliphatic hydrocarbon containing one or more carbon-to-carbon double bonds. Examples include, but are not limited to, vinyl, allyl, and the like.

[0018] As used herein, the term "alkynyl" refers to a straight or branched chain aliphatic hydrocarbon containing one or more carbon-to-carbon triple bonds. Examples include, but are not limited to, ethynyl and the like.

[0019] The term "alkylene" refers to a straight or branched chain divalent hydrocarbon radical, preferably having from one to ten carbon atoms. Alkylene groups as defined herein may optionally be substituted. Examples of "alkylene" as used herein include, but are not limited to, methylene, ethylene, n-propylene, n-butylene.

[0020] As used herein, the term "alkenylene" refers to a straight or branched chain divalent hydrocarbon radical, preferably having from one to ten carbon atoms, containing one or more carbon-to-carbon double bonds that may be optionally substituted. Examples include, but are not limited to, vinylene, allylene or 2-propenylene, and the like.

[0021] The term "alkynylene" refers to a straight or branched chain divalent hydrocarbon radical, preferably having from one to ten carbon atoms, containing one or more carbon-to-carbon triple bonds that may be optionally substituted. Examples include, but are not limited to, ethynylene and the like.

[0022] As used herein, the term "cycloalkyl" refers to an optionally substituted non-aromatic cyclic hydrocarbon ring, which optionally includes an alkylene linker through which the cycloalkyl may be attached. Exemplary "cycloalkyl" groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and substituted versions thereof. As used herein, the term "cycloalkyl" includes an optionally substituted fused polycyclic hydrocarbon saturated ring and aromatic ring system, namely polycyclic hydrocarbons with less than maximum number of non-cumulative double bonds, for example where a saturated hydrocarbon ring (such as a cyclopentyl ring) is fused with an aromatic ring (herein "aryl," such as a benzene ring) to form, for example, groups such as indane.

[0023] As used herein, the term "cycloalkenyl" refers to an optionally substituted non-aromatic cyclic hydrocarbon ring

containing one or more carbon-to-carbon double bonds which optionally includes an alkylene linker through which the cycloalkenyl may be attached. Exemplary "cycloalkenyl" groups include, but are not limited to, cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, and substituted versions thereof.

[0024] The term "cycloalkylene" refers to a divalent, optionally substituted non-aromatic cyclic hydrocarbon ring. Exemplary "cycloalkylene" groups include, but are not limited to, cyclopropylene, cyclobutylene, cyclopentylene, cyclohexylene, cycloheptylene, and the like.

[0025] As used herein, the term "cycloalkenylene" refers to a divalent optionally substituted non-aromatic cyclic hydrocarbon ring containing one or more carbon-to-carbon double bonds. Exemplary "cycloalkenylene" groups include, but are not limited to, cyclopropenylene, cyclobutenylene, cyclopentenylene, cyclohexenylene, cycloheptenylene, and the like.

[0026] As used herein, the term "heterocycle" or "heterocyclyl" refers to an optionally substituted mono- or polycyclic ring system containing one or more degrees of unsaturation and also containing one or more heteroatoms. Preferred heteroatoms include N, O, and/or S, including N-oxides, sulfur oxides, and dioxides. Preferably the ring is three to twelve-membered and is either fully saturated or has one or more degrees of unsaturation. Such rings may be optionally fused to one or more of another "heterocyclic" ring(s) or cycloalkyl ring(s). Examples of "heterocyclic" groups include, but are not limited to, tetrahydrofuran, pyran, 1,4-dioxane, 1,3-dioxane, piperidine, pyrrolidine, morpholine, tetrahydrothiopyran, and tetrahydrothiophene.

[0027] As used herein, the term "aryl" refers to an optionally substituted benzene ring or to an optionally substituted fused

benzene ring system, for example anthracene, phenanthrene, or naphthalene ring systems. Examples of "aryl" groups include, but are not limited to, phenyl, 2-naphthyl, 1-naphthyl, and the like.

[0028] As used herein, the term "heteroaryl" refers to an optionally substituted monocyclic five to seven membered aromatic ring, or to an optionally substituted fused bicyclic aromatic ring system comprising two of such aromatic rings. These heteroaryl rings contain one or more nitrogen, sulfur, and/or oxygen atoms, where N-oxides, sulfur oxides, and dioxides are permissible heteroatom substitutions. Examples of "heteroaryl" groups used herein include, but should not be limited to, furan, thiophene, pyrrole, imidazole, pyrazole, triazole, tetrazole, thiazole, oxazole, isoxazole, oxadiazole, thiadiazole, isothiazole, pyridine, pyridazine, pyrazine, pyrimidine, quinoline, isoquinoline, benzofuran, benzothiophene, indole, indazole, benzimidazolyl, imidazopyridinyl, pyrazolopyridinyl, pyrazolopyrimidinyl, and the like.

[0029] As used herein, the term "halogen" refers to fluorine, chlorine, bromine, or iodine. In this respect, the term "haloalkyl" refers to an alkyl group, as defined herein, that is substituted with at least one halogen. Examples of branched or straight chained "haloalkyl" groups useful in the present invention include, but are not limited to, methyl, ethyl, propyl, isopropyl, n-butyl, and t-butyl substituted independently with one or more halogens, e.g., fluoro, chloro, bromo, and iodo. The term "haloalkyl" should be interpreted to include such substituents as perfluoroalkyl groups and the like.

[0030] As used herein, the term "alkoxy" refers to the group -O-alkyl. The term "nitro" refers to the group -NO₂. The term "cyano" refers to the group -CN. The term "azido" refers to

the group $-N_3$. The term "acyl" refers to the group $R^aC(=O)-$, where R^a is alkyl, aryl, heteroaryl, or heterocyclyl, as each is defined herein.

[0031] As used herein throughout the specification, the phrase "optionally substituted" or variations thereof denote an optional substitution, including multiple degrees of substitution, with one or more substituent group. The phrase should not be interpreted so as to be imprecise or duplicative of substitution patterns herein described or depicted specifically. Rather, those of ordinary skill in the art will appreciate that the phrase is included to provide for obvious modifications, which are encompassed within the scope of the appended claims.

[0032] Exemplary optional substituent groups include acyl; alkyl; alkenyl; alkynyl; alkylsulfonyl; alkoxy; alkoxy-carbonyl; cyano; halogen; haloalkyl; hydroxy; nitro; aryl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroaryl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; arylsulfonyl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroarylsulfonyl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; aryloxy, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroaryloxy, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; or aryloxy-carbonyl, which may be further


substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro.

[0033] According to the instant method, the multicomponent reaction is carried out in the presence of an acid catalyst. In some embodiments, the acid is a Lewis acid, e.g., a chiral Lewis acid complex. In some embodiments, the reaction is carried out at room temperature. In other embodiments, the reaction is carried out at 0 to 100°C. In still other embodiments, the acid is a **Brønsted** acid. To provide enantioselective variants of compounds of Formula V, other embodiments of the invention include the use of a chiral phosphoric acid or other chiral **Brønsted** acid.


[0034] Lewis acids include, but are not limited to, metal cations of aluminum (III), gallium(III), ferric(III), indium(III), antimony(V), tin(IV), titanium(IV), zinc(II), boron (III), and copper (II); and electron-deficient molecules such as boron trifluoride and aluminum trichloride. In some embodiments, the Lewis acid is a metal halide. Examples of metal halides include, but are not limited to, AlBr_3 , **AlCl_3** , GaBr_3 , GaCl_3 , FeCl_3 , InI_3 , InBr_3 , InCl_3 , SbCl_5 , SnCl_4 , TiCl_4 , **ZnCl_2** , BF_3 , BCl_3 and lanthanide halides. In other embodiments, the Lewis acid is a metal salt. Examples of metal salts include, but are not limited to, triflates such as $\text{Ga}(\text{OTf})_3$, $\text{In}(\text{OTf})_3$, $\text{Sc}(\text{OTf})_3$, $\text{Ln}(\text{OTf})_3$ or $\text{Cu}(\text{OTf})_2$. In particular embodiments the Lewis acid is a metal halide or triflate salt of Ga (III) or In (III).

[0035] **Brønsted** acids of use in the method of this invention include, but are not limited to, TFA, TfOH, AcOH, TsOH, Tf_2NH , MsOH and the like.

[0036] Chiral phosphoric acids of this invention have the structure of Formula Via or VIb, or a racemic mixture thereof,



Formula Via



Formula VIb

wherein,


R^{12} is selected from groups including, but not limited to, a hydrogen, p-biphenyl, anthryl, $SiPh_3$, 3,5-bis (trifluoromethyl) phenyl, 2,4,6-triisopropylphenyl and $NO_2C_6H_5$. Such chiral phosphoric acid catalysts are known and available from commercial sources such as Sigma-Aldrich (St. Louis, MO) and include, but are not limited to (*R*)-(-)-1,1'-Binaphthyl-2,2'-diyl hydrogenphosphate, (*S*)-(+)-1,1'-Binaphthyl-2,2'-diyl hydrogenphosphate, (*R*)-(-)-3,3'-Bis (triphenylsilyl) -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, (*S*)-(+)-3,3'-Bis (triphenylsilyl) -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, (*R*)-3,3'-Bis [3,5-bis (trifluoromethyl) phenyl] -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, (*S*)-3,3'-Bis [3,5-bis (trifluoromethyl) phenyl] -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, (*R*)-3,3'-Bis (2,4,6-triisopropylphenyl) -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, (*S*)-3,3'-Bis (2,4,6-triisopropylphenyl) -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate, and (*R*)-3,3'-Bis (9-anthracenyl) -1,1'-binaphthyl-2,2'-diyl hydrogenphosphate. See also, Cheon & Yamamoto (2008) *J. Am. Chem. Soc.* 130:9246-7.

[0037] Compounds of the invention may be provided as mixtures of stereoisomers as well as purified enantiomers or enantiomerically/diastereomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds represented by Formula Va, Vb, Vc and Vd as well as any wholly or partially equilibrated mixtures

thereof. Moreover, the instant compounds may be provided as pharmaceutically acceptable salts or solvates thereof.

[0038] Compounds produced by the method of this invention have been shown to stimulate the secretion and/or production of GLP-1 and/or inhibit the activity of the CGRP receptor. Therefore, the compounds of this invention can be provided in pharmaceutical compositions and used in methods for stimulating the secretion and/or production of GLP-1 and/or inhibiting the activity of the CGRP receptor.

[0039] In particular embodiments, a compound of use in the compositions and method of the invention has the structure of Formula Ve, Vf or Vg:



wherein X is C-(R¹)_n, OR¹, S or NR¹, wherein each R¹ is independently hydrogen, alkyl, alkenyl, alkynyl or aryl;

each Y is the same or different and is independently N or C;

each Z is the same or different and is independently N or C and wherein Z can optionally be connected to any other Z by a chain of one or more atoms independently selected from substituted or unsubstituted C, N, O, S, or a combination thereof, thereby forming an additional ring(s).

each R² is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, -R⁷-cycloalkyl, aryl, -NHR⁷, Het, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -CO₂R⁸, -R⁷CO₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -

$R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(O)_2NR^8R^9$, $-S(O)_2NR^8R^9$, $-R^7S(O)_2NHR^8$, $-R^7S(O)_2NR^8R^9$, $-R^7S(O)_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^6 is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, $-NHR^7$, Het, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(O)R^8$, $-C(O)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(O)NR^8R^9$, $-C(O)aryl$, $-C(O)NR^8aryl$, $-C(O)Het$, $-C(O)NHR^7Het$, $-R^7C(O)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(O)_2NR^8R^9$, $-S(O)_2NR^8R^9$, $-R^7S(O)_2NHR^8$, $-R^7S(O)_2NR^8R^9$, $-R^7S(O)_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^7 is the same or different and is independently selected from an alkylene, cycloalkylene, alkenylene, cycloalkenylene or alkynylene group;

each of R^8 and R^9 are the same or different and are independently selected from the group of hydrogen, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7cycloalkyl$, $-R^7OH$, $-R^7(OR^7)_w$, or $-R^7NR^{10}R^{11}$ group;

each of R^{10} and R^{11} are the same or different and are independently selected from the group of an alkyl, cycloalkyl, alkenyl, cycloalkenyl, or alkynyl group;

p is selected from 0, 1, 2, 3, or 4;

q is selected from 0, 1, 2, 3, or 4;

each n independently is 0, 1 or 2;

each m independently is 0, 1 or 2; and

w is 1-10.

[0040] Any one of the compounds of this invention can be prepared as a salt, solvate or physiologically functional derivative. Typically, but not absolutely, the salts of the compounds of this invention are pharmaceutically acceptable salts. Salts encompassed within the term "pharmaceutically

acceptable salts" refer to non-toxic salts of the compounds of this invention. Salts of the compounds of the invention may include acid addition salts. Representative salts include acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, calcium edetate, camsylate, carbonate, clavulanate, citrate, dihydrochloride, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycollylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isethionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylsulfate, monopotassium maleate, mucate, napsylate, nitrate, N-methylglucamine, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, potassium, salicylate, sodium, stearate, subacetate, succinate, sulfate, tannate, tartrate, teoclate, tosylate, triethiodide, trimethylammonium, and valerate salts. Other salts, which are not pharmaceutically acceptable, may be useful in the preparation of compounds of this invention and these should be considered to form a further aspect of the invention.

[0041] As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of Formula Va-Vg, or a salt or physiologically functional derivative thereof) and a solvent. Such solvents, for the purpose of the invention, should not interfere with the biological activity of the solute. Non-limiting examples of suitable solvents include, but are not limited to water, methanol, ethanol, and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Non-limiting examples of suitable pharmaceutically acceptable solvents include water, ethanol, and acetic acid. Most preferably the solvent used is water.

[0042] As used herein, the term "physiologically functional derivative" refers to any pharmaceutically acceptable derivative of a compound of this invention that, upon administration to a mammal, is capable of providing (directly or indirectly) a compound of the invention or an active metabolite thereof. Such derivatives, for example, esters and amides, will be clear to those skilled in the art, without undue experimentation. Reference may be made to the teaching of Burger's Medicinal Chemistry and Drug Discovery, 5th Edition, Vol 1: Principles and Practice.

[0043] Pharmaceutical compositions can be prepared by methods and contain carriers which are well-known in the art. A generally recognized compendium of such methods and ingredients is Remington: The Science and Practice of Pharmacy, Alfonso R. Gennaro, editor, 20th ed. Lippincott Williams & Wilkins: Philadelphia, PA, 2000. A pharmaceutically acceptable carrier or vehicle, such as a liquid or solid filler, diluent, excipient, or solvent encapsulating material, is involved in carrying or transporting the subject agent from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be acceptable in the sense of being compatible with the other ingredients of the formulation and not injurious to the patient.

[0044] Examples of materials which can serve as pharmaceutically acceptable carriers include sugars, such as lactose, glucose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and

polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; pH buffered solutions; polyesters, polycarbonates and/or polyanhydrides; and other non-toxic compatible substances employed in pharmaceutical formulations. Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions .

[0045] When used *in vivo*, i.e., administered directly to a subject, the compounds of this invention can be administered by a variety of methods, e.g., orally, by injection (e.g. subcutaneous, intravenous, intraperitoneal, etc.), topically including via inhalation, transdermally, intranasally, intravaginally, or rectally according to standard medical practices .

[0046] Depending on the route of administration, the active compounds may be coated in a material to protect the compound from the action of acids and other natural conditions which may inactivate the compound. They may also be administered by continuous perfusion/infusion into a subject or provided as a prodrug.

[0047] To administer the therapeutic compound by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation . For example, the therapeutic compound may be administered to a patient in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Liposomes include water-in-oil-in-

water CGF emulsions as well as conventional liposomes (Strejan, et al. (1984) *J. Neuroimmunol.* 7:27).

[0048] The therapeutic compound may also be administered parenterally, intraperitoneally, intraspinally, or intracerebrally. Dispersions can be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

[0049] Sterile injectable solutions can be prepared by incorporating a compound of the invention in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating a compound of the invention into a sterile pharmaceutically acceptable carrier which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0050] A compound of the invention can be orally administered, for example, with an inert diluent or an assimilable edible carrier. The active agent and other ingredients may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the patient's diet. For oral therapeutic administration, the active agent may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. The percentage of the active ingredient in the compositions and

preparations may, of course, be varied. The amount of the active ingredient in such therapeutically useful compositions is such that a suitable dosage will be obtained.

[0051] It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the patients to be treated; each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding an active ingredient for the treatment of a selected condition in a patient.

[0052] Active compounds are administered at a therapeutically effective dosage sufficient to treat a condition associated with a given patient. For example, the efficacy of a compound of this invention can be evaluated in an animal model system that may be predictive of efficacy in treating the disease in humans .

[0053] The actual dosage amount of a compound of the present disclosure or composition comprising a compound of the present disclosure administered to a patient may be determined by physical and physiological factors such as age, sex, body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the patient and on the route of administration. These factors may be determined by a skilled artisan. The practitioner responsible for administration will typically determine the concentration of active ingredient (s) in a

composition and appropriate dose(s) for the individual patient. The dosage may be adjusted by the individual physician in the event of any complication.

[0054] An effective amount typically will vary from about 0.001 mg/kg to about 1000 mg/kg, from about 0.01 mg/kg to about 750 mg/kg, from about 100 mg/kg to about 500 mg/kg, from about 1.0 mg/kg to about 250 mg/kg, from about 10.0 mg/kg to about 150 mg/kg in one or more dose administrations daily, for one or several days (depending of course of the mode of administration and the factors discussed above). Other suitable dose ranges include 1 mg to 10000 mg per day, 100 mg to 10000 mg per day, 500 mg to 10000 mg per day, and 500 mg to 1000 mg per day. In some particular embodiments, the amount is less than 10,000 mg per day with a range of 750 mg to 9000 mg per day.

[0055] The effective amount may be less than 1 mg/kg/day, less than 500 mg/kg/day, less than 250 mg/kg/day, less than 100 mg/kg/day, less than 50 mg/kg/day, less than 25 mg/kg/day or less than 10 mg/kg/day. It may alternatively be in the range of 1 mg/kg/day to 200 mg/kg/day. In other non-limiting examples, a dose may also comprise from about 1 microgram/kg/body weight, about 5 microgram/kg/body weight, about 10 microgram/kg/body weight, about 50 microgram/kg/body weight, about 100 microgram/kg/body weight, about 200 microgram/kg/body weight, about 350 microgram/kg/body weight, about 500 microgram/kg/body weight, about 1 milligram/kg/body weight, about 5 milligram/kg/body weight, about 10 milligram/kg/body weight, about 50 milligram/kg/body weight, about 100 milligram/kg/body weight, about 200 milligram/kg/body weight, about 350 milligram/kg/body weight, about 500 milligram/kg/body weight, to about 1000 mg/kg/body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from

the numbers listed herein, a range of about 5 mg/kg/body weight to about 100 mg/kg/body weight, about 5 microgram/kg/body weight to about 500 milligram/kg/body weight, etc., can be administered, based on the numbers described above.

[0056] Single or multiple doses of a compound described herein are contemplated. Desired time intervals for delivery of multiple doses can be determined by one of ordinary skill in the art employing no more than routine experimentation. As an example, patients may be administered two doses daily at approximately 12 hour intervals. In some embodiments, the compound is administered once a day.

[0057] The compound may be administered on a routine schedule. As used herein, a routine schedule refers to a predetermined designated period of time. The routine schedule may encompass periods of time which are identical or which differ in length, as long as the schedule is predetermined. For instance, the routine schedule may involve administration twice a day, every day, every two days, every three days, every four days, every five days, every six days, a weekly basis, a monthly basis or any set number of days or weeks there-between. Alternatively, the predetermined routine schedule may involve administration on a twice daily basis for the first week, followed by a daily basis for several months, etc. In other embodiments, the invention provides that the compound may be taken orally and that the timing of which is or is not dependent upon food intake.

[0058] Glucagon-like peptide 1 (GLP-1) is secreted by the intestinal L-cells and is a potent antihyperglycemic hormone. GLP-1 functions by inducing glucose-dependent stimulation of insulin secretion and at the same time suppressing glucagon secretion. It has been reported that GLP-1 can also restore the glucose sensitivity of pancreatic β -cells. As a result, GLP-1 secretagogues are potential therapeutic agents for type

2 diabetes. GLP-1 is also known to inhibit both gastric secretion and gastric emptying, thereby contributing to the feeling of satiety. Having demonstrated the stimulation of GLP-1 secretion and/or production, the compounds of this invention can also be used in the preparation of a dietary supplement, a food or feed product, or a beverage product for helping to sustain energy, helping control appetite, helping control blood sugar levels, reducing the risks associated with metabolic syndrome, reducing the risk associated with obesity and diabetes, reducing the risk associated with diabetes, helping to maintain healthy glucose and fat metabolism, or for helping to normalize production and release of GLP1 necessary for healthy glucose and fat metabolism, in a subject during and/or between meals or feedings comprising said dietary supplement, a food or feed product, or a beverage product.

[0059] In this respect, the invention also provides a method for stimulating GLP-1 secretion and/or production for helping to sustain energy, helping control appetite, helping control blood sugar levels, reducing the risks associated with metabolic syndrome, reducing the risk associated with obesity and diabetes, reducing the risk associated with diabetes, helping to maintain healthy glucose and fat metabolism, or for helping to normalize production and release of GLP-1 necessary for healthy glucose and fat metabolism, said method comprising administering to a subject an effective amount of a compound of the invention during and/or between meals or feedings. In some embodiments, a compound of the invention may provide a general improvement to human health and can protect against the harmful health effects associated with metabolic syndrome. It is also contemplated that a compound of the invention can protect against the harmful health effects associated with type 2 diabetes, and against the harmful health effects associated with obesity. The effect on GLP-1 secretion and/or

production may be confirmed in an *in vivo* experiment such as described in Examples 5 and 6.

[0060] Furthermore, having demonstrated CGRP receptor inhibitory activity, the invention also provides methods for inhibiting CGRP receptor activity and in the treatment of pathophysiologic conditions where excessive CGRP receptor activation has occurred. Some of these include neurogenic vasodilation, neurogenic inflammation, migraine, cluster headache and other headaches, thermal injury, circulatory shock, menopausal flushing, and asthma. CGRP receptor activation has been implicated in the pathogenesis of migraine headache (Edvinsson (2001) *CNS Drugs* 15(10):745-53; Williamson (2001) *Microsc. Res. Tech.* 53:167-178; Grant (2002) *Brit. J. Pharmacol.* 135:356-362). Serum levels of CGRP are elevated during migraine (Goadsby, et al. (1990) *Ann. Neurol.* 28:183-7) and treatment with antimigraine drugs returns CGRP levels to normal coincident with alleviation of headache (Gallai, et al. (1995) *Cephalalgia* 15:384-90). Migraineurs exhibit elevated basal CGRP levels compared to controls (Ashina, et al. (2000) *Pain* 86(1-2):133-8). Intravenous CGRP infusion produces lasting headache in migraineurs (Lassen, et al. (2002) *Cephalalgia* 22(1):54-61). Preclinical studies in dog and rat report that systemic CGRP blockade with the peptide antagonist CGRP (8-37) does not alter resting systemic hemodynamics nor regional blood flow (Shen, et al. (2001) *J. Pharmacol. Exp. Ther.* 298:551-8). Thus, CGRP-receptor antagonists may present a novel treatment for migraine that avoids the cardiovascular liabilities of active vasoconstriction associated with non-selective 5-HT_{1B/1D} agonists.

[0061] The invention is described in greater detail by the following non-limiting examples.

Example 1: Lewis Acid and Brønsted Acid Catalyzed Synthesis of Cyclohepta [1,2-b]Indoles

[0062] It has been shown that treatment of 2-furfuryl alcohols with stoichiometric TiCl_4 promotes the formation of furfuryl cations for (4+3) cycloaddition reactions (Winne, et al. (2011) *Angew. Chem. Int. Ed.* 50:11900; Pattenden & Winne (2009) *Tetrahedron Lett.* 50:7310). Therefore, it was speculated that nucleophilic addition of indole at C3 to either an aldehyde or ketone would furnish alcohol **12** (Scheme 2). In the presence of an appropriate Lewis acid, water would be ejected to generate the requisite allylic cation **13** for (4+3) cycloaddition reactions. Ga(III)-catalyzed processes have been described (Lauer, et al. (2011) *J. Am. Chem. Soc.* 133:9119; Han & Wu (2010) *Org. Lett.* 12:5780; Robertson & Wu (2010) *Org. Lett.* 12:2668) and therefore these Lewis acids were used in the title reaction. Ga(III) salts, and in particular $\text{Ga}(\text{OTf})_3$ (Olah, et al. (1988) *J. Am. Chem. Soc.* 11:2560; Prakash, et al. (2007) *Proc. Natl. Acad. Sci. USA* 104:3703; Prakash, et al. (2007) *Org. Lett.* 9:179; Prakash, et al. (2003) *Catal. Lett.* 85:1; Yan, et al. (2005) *Catal. Lett.* 103:165) are stable to air and moisture and therefore ideal in the present case because an equivalent of water is formed in the reaction.

TABLE 1

Entry ^a	Catalyst	Comments ^d
1	$\text{Ga}(\text{OTf})_3$	Full conversion, impurities hard to separate
2	GaBr_3	Full conversion, clean reaction (85% yield)
3 ^b	GaBr_3	Slow reaction
4 ^c	GaBr_3	Only 12 observed

5	GaCl ₃	Mostly 12 observed
6	In(OTf) ₃	Substantial decomposition
7	InI ₃	Full conversion, clean reaction
8	InBr ₃	Slow reaction
9	InCl ₃	Mostly 12 observed
10	Sc(OTf) ₃	Slow reaction
11	Cu(OTf) ₂	Only 12 observed
12	TFA	Only 12 observed
13	TfOH	Full conversion, impurities hard to separate

^a10 mol% catalyst, 2 equiv. **9a**, 5 equiv **10a**, rt, CH₂Cl₂.

^bToluene for EtOAc as solvent.

^cTHF or Et₂O as solvent.


















^dDiastereoselectivities were 3:1 to 5:1 for all reactions in which product was observed (as determined by ¹H NMR spectroscopy).


[0063] As is evident from Table 1, both Ga(OTf)₃ and GaBr₃ were effective in promoting the desired reaction. Ga(OTf)₃ was qualitatively the more reactive catalyst. But because it also generated a by-product (not formed with the use of GaBr₃) that was difficult to separate from **11a**, the scope of the transformation was analyzed using GaBr₃. InI₃ was also effective but several other Lewis acids (entries 6-11) resulted in either slow reaction rates or varying amounts of alcohol **12**.

[0064] The scope of the three-component (4+3) cycloaddition reaction was quite broad (Table 2). A variety of combinations involving indoles **8a-e**, aldehydes **9a-e** and **16**, acetal **15**, ketones **17a-c**, and dienes **10a-e** were surveyed. The reaction was tolerant to N-H indoles as well as alkylation at nitrogen. Aldehydes, ketones, and acetals were appropriate electrophilic components. Electron-withdrawing or -donating groups on either the indole or carbonyl components were compatible. Electron-rich substrates such as **8d**, **9b**, and **9d-e** appeared to accelerate reaction rate. With less reactive substrates (**Hi**, **11k-11n**) the use of Ga(OTf)₃ was beneficial for achieving higher yields and shorter reaction times. The presence of halides in **11s-u** offers a convenient handle for additional

elaboration of the indole core through transitional metal-catalyzed cross-coupling reactions.

TABLE 2

 <p>8a; R = Me 8b; R = Bn 8c; R = H</p>		 <p>8d; R = MeO 8e; R = MeO₂C 8f-h; R = Cl, Br, I</p>		
 <p>9a; R = H 9b; R = MeO 9c; R = CF₃</p>	 <p>9d; X = O 9e; X = S</p>	 <p>15</p>  <p>16</p>	 <p>17a; n = 1 17b; n = 2 17c; n = 3</p>	
 <p>10a</p>	 <p>10b</p>	 <p>10c</p>	 <p>10d</p>	 <p>10e</p>
 <p>11a (94%, 4:1 dr)^a</p>	 <p>11b (75%, 10:1 dr)^a</p>	 <p>11c (85%, 2:1 dr)^a</p>	 <p>11d (83%, >10:1 dr)^a</p>	
 <p>11e (83%, 10:1 dr)^a</p>	 <p>11f (82%, 1:1 dr)^a</p>	 <p>11g (94%)^a</p>	 <p>11h (84%)^a</p>	



^aindole (1 equiv), electrophile (2 equiv), diene (5 equiv), GaBr₃ (10 mol%), RT.

^bindole (1 equiv), electrophile (2 equiv), diene (5 equiv), Ga(OTf)₃ (10 mole%), RT.

^cindole (1 equiv), electrophile (1.1 equiv), diene (5 equiv), Ga(OTf)₃ (20 mole%), RT.

^dYield with 20% TfOH.

^eStructure confirmed by single crystal X-ray analysis.


^f2 mmol scale, 5 mol% GaBr₃, all else same.

[0065] For products **11a-f**, diastereoselectivities were observed in the range of >10:1 to 1:1. Electron-rich aldehydes appeared to couple with the highest selectivities while electron-poor and linear alkyl substrates furnished products with lower selectivities. When the use of the unsymmetrical diene, isoprene (**10c**), was examined, the desired product **11m** was isolated as a single regioisomer. The structures of **11j**, **11n** and **11p** were confirmed by single-crystal X-ray crystallography.

[0066] To gain insight into the mechanism of this reaction, density-functional-theory (DFT) calculations were performed using the M06 functional (Zhao & Truhlar (2008) *Acc. Chem. Res.* 41:157; Zhao & Truhlar (2003) *Theor. Chem. Acc.* 120:215) and the 6-311G**++ basis set (Krishnan, et al. (1980) *J. Chem. Phys.* 72:650; McLean & Chandler (1980) *J. Chem. Phys.* 72:5639; Clark, et al. (1983) *J. Comput. Chem.* 4:294; Frisch, et al. (1984) *J. Chem. Phys.* 80:3265) as implemented in the Jaguar program. For the reaction of cyclopentadiene (10a) with the indolyl cation derived from indole (8c) and ketal 15, the lowest energy pathway was identified as a two-step process. Rather than a concerted pericyclic reaction, an intermediate was formed by initial C-C bond formation between the terminal carbon of the indolyl cation and a terminal diene carbon, followed by closure of the 7-membered ring. Transition states for both steps were located, and an overall reaction free energy profile was generated. A stepwise mechanism was observed as proposed for related (4+3) reactions (Winne, et al. (2011) *supra*).

Example 2: Chiral Phosphoric Acid Catalyzed Synthesis of Cyclohepta [↔] Indoles

[0067] An investigation to develop enantioselective variants of this transformation was also initiated. This analysis indicated that chiral phosphoric acid 18 catalyzed the reaction between indole (8c), anisaldehyde (9b), and (10) to furnish the desired product 11v in 64% ee (eq 1) (Scheme 4).




SCHEME 4


[0068] Several chiral Ga(III) and In(III) complexes are known (Li, et al. (2005) *Adv. Synth. Catal.* 347:1247; Li, et al. (2002) *Chem Commun.* 2994; Lv, et al. (2011) *Angew. Chem. Int. Ed.* 50:6612; Teo, et al. (2005) *Org. Lett.* 7:2743; Gutierrez, et al. (2011) *Org. Lett.* 13:5754) and have been utilized to promote transformations with excellent *ee*'s. Therefore, these catalysts may also be of use in the synthesis of the title reaction.

Example 3: Lewis Acid and Brønsted Acid Catalyzed Synthesis of Cyclopenta [b]Indoles

[0069] Cyclopenta [b]indoles were also synthesized using the method of this invention. Schemes 5 and 6 illustrate the synthesis of two exemplary cyclopenta [b]indoles prepared with TfOH.



SCHEME 5







SCHEME 6

[0070] Table 3 provides examples of additional compounds and their respective yields using TfOH.

TABLE 3


Compound	Product	Yield with TfOH	Reaction Time
19c		80%	40 min.
19d		62%	1 hour
19e		64%	3 hours
19f		64%	3 hours







19g		83%	15 min.
19h		52%	1 hour
19i		58%	40 min.
19j		50%	2 hours

Reaction conditions: 1 equiv indole, 4 equiv ketone, 5 equiv alkene, 0.5 equiv TfOH, room temperature in dichloroethane as solvent .

[0071] In similar reactions using $\text{Bi}(\text{OTf})_3$, additional cyclopenta [b] indoles were synthesized (Table 4) .


TABLE 4

Compound	Product	Yield with $\text{Bi}(\text{OTf})_3$
19k ^a		51%

19l ^d		46%
19m ^c		51%
19n ^b		13%
19o ^c		41%
19p ^c		25%
19q ^c		32%

^a10% catalyst used; ^b15% catalyst used; ^c20% catalyst used, 40°C.

[0072] Scheme 7 illustrates the synthesis of another exemplary cyclopenta [b] indole prepared with Ga(OTf)₃.



SCHEME 7

[0073] The use of In(OTf)_3 in the preparation of cyclopenta [b] indoles was also demonstrated (Table 5).

TABLE 5



Compound	Product	Yield with In(OTf)_3
19s		64%
19t^a		65%

^a10% Catalyst, 40°C.

Example 4: *In Vitro* Activity

[0074] Selected compounds of this invention were screened for K-Ras Wnt synthetic lethal, anti-angiogenic activity, Wnt pathway modulatory activity, insulin and GLP-1 secretion modulatory activity, GPR119 receptor agonist activity, mGlu2R antagonist activity, CGRP receptor antagonist activity, Apelin receptor agonist activity and hexokinase 2 inhibitory activity using the PD² and TD² Screening Panel (Eli Lilly & Company). See Lee, et al. (2011) *J. Biomol. Screen.* 16:588-602. The results of this analysis are presented in Table 6.

TABLE 6

Activity	Assay	Compound	
		 11s	 11t
K-Ras Wnt	HCT116 KrasSL % Inhibit @0.2 μM @2 μM @20 μM	2.3% -38% 96.4%	5.3% -11.5% 103.9%
	RKO KrasSL % Inhibit @0.2 μM @2 μM @20 μM	-2.8%* 14.4%* 101.7%*	-5.8%* 17.6%* 102.2%*
	Colo320 KrasSL % Inhibit @0.2 μM @2 μM @20 μM	33.3% 15.8 150.3%	-6.6% 23.1% 151.5%
	SNU-C1 KrasSL % Inhibit @0.2 μM @2 μM @20 μM	15.7% 5.5% 106.4%	-1% -19.1% 107.2%
Anti-Angiogenic	Angio Tube Area % Inhibit @2 μM @10 μM	21.7% 21.8%	8% 38%
Wnt Pathway	Osteo bCat % Stimulation @2 μM @10 μM	1.3% 9.3%	1.8% 5.5%
Insulin Secretion	Secretion Hi Gluc % Stimulation @2 μM @10 μM	-1.2% 15.8%	0.1% 19.1%
GLP-1 Secretion	hNCI GLP-1 Secretion % Stimulation @2 μM @20 μM EC ₅₀ (μM)	-1.7% 108% 34.828	0% 35.2% 29.052
GPR 119 Receptor Agonist	hGPR119 agonism % Stimulation	-5.1%	-2.3%

	@10 μ M		
mGlu2R Antagonist	hMGLUR2 % Inhibition @50 μ M	-20.4%	-19.7%
CGRP Receptor Antagonist	hCGRP1 Antagonism % Inhibition @30 μ M IC ₅₀ (nM)	77.1% 0	67.7% 0
Apelin Receptor Agonist	hApelin agonism % Stimulation @30 μ M	-46.2%	-52.8%
Hexokinase 2 Inhibitor	hHK2 ADP-FP % Inhibition @20 μ M	5.5%	3.2%





^Average of three independent experiments . ^Average of three independent experiments.




[0075] This analysis indicated that compounds **11s** and **11t** exhibited both stimulatory activity for the secretion and/or production of GLP-1, and inhibitory activity for the CGRP receptor with EC₅₀ values of 35 and 29 μ M, respectively.

[0076] In light of the above results, additional phenotypic assays were carried out using hNCI-H716 and mSTC-1 cell lines to monitor GLP-1 secretion (single point (SP) and concentration response curves (CRC)) as well as a secondary assay with pituitary cells. The results of these assays are presented in Table 1.

TABLE 7

Compound	Primary SP hNCI-H716 (% Stim.)		Primary CRC		Pituitary Cells (Growth Hormone Secretion, % Stimulation)	
			hNCI- H716	mSTC- 1		
	@ 2 μ M	@ 20 μ M	EC ₅₀ (μ M)	EC ₅₀ (μ M)	@ 3.3 μ M	@ 11 μ M
11a	-2.8	-0.5				
11b	-0.5	-2.4				
11c	1	0.7				
11d	-2.5	-1				
11e	-0.5	-2.4				
11f	-1.1	-2.4				

11g	-0.8	-2.7				
11h	-2.2	-3.3				
11i	0	-6				
11j	-0.3	-3.3				
11k	3.1	-1.9				
11l	-2	-3.3				
11m	-2.2	-1				
11n	10.3	-3.5				
11o	-0.3	-2.1				
11p	-0.5	23.3				
11q	1.3	5.9				
11r	-1.1	13.4				
11u	0	57	>40.0 14.7 24.5	2.0 3.2	9.3 14.5	6.5 -15.2
11t	0	35.2	29.1 17.6 16.4	2.4 4.1 2.2	-12.6 -7.1	2.4 -8.1
11s	-1.7	108	34.8 33.5 23.2	1.9 3.6 5.0	-10.0 -17.7	-22.6 -37.6
 19u	-0.8	-3.8				
 19v	-2	-2.4				
 19w	-0.8	-1.9				
 19x	-3.1	-3.3				

 <p>11w</p>	1.4	41.3				
 <p>11x</p>	3.4	-0.8				
 <p>11y</p>	-0.4	55.9				

Example 5: In Vivo Activity in Animal Model

[0077] Mice are fasted overnight (16 hours) by being placed in a clean cage without food. The next day, all mice are weighed and assigned to groups of similar body weights. Compounds are dosed in a time-dependent manner and cardiac sticks are done on CO₂ euthanized mice at 0.25, 0.50, 1.50, and 3.00 hours post-dose (option 1) or 0.5, 1.5, 3, and 6 hours post-dose (option 2). Blood (600+ μ l) from the cardiac stick is placed in EDTA-plasma tubes with inhibitors (DPP4 inhibitor and aprotinin), and stored on ice. Plasma is separated by centrifugation and aliquoted for compound exposure and up to three analytes, including total GIP, GLP-1 (7-36) amide, and/or total PYY. Plates for compound exposure are analyzed for absorption, distribution, metabolism, and excretion, while plates for GLP-1, GIP, and/or PYY are analyzed using conventional kits. Resulting raw data are used to generate area under the curve analyses (AUC).

Example 6 : In Vivo Activity in Human Subjects

[0078] Healthy volunteers that have fasted for 10 hours are given enteral feeding by a duodenal tube. Each volunteer is given two different liquid meals (bolus, 55-65 ml) on four different days (A-D) ,

Day A : Test Compound in 50 ml glycerol + 5 ml ethanol

Day B : oleic acid (1.54 g) in 50 ml glycerol + 5 ml ethanol (= control Day A)

Day c : Test Compound (2 g) + glucose (10-20g in 10 ml water) in 50 ml glycerol + 5 ml ethanol

Day D : oleic acid (1.54 g) + glucose (10-20g in 10 ml water) in 50 ml glycerol + 5 ml ethanol (=control Day c)


[0079] Blood samples are collected at -15, -10, 0, 10, 20, 30, 40, 50, 60, 75, 90, 120, 150, 180, and 240 minutes. In addition, a 1 mL sample is collected from the duodenal lumen at 15 and 30 minutes. Insulin and C-peptide levels are measured in the serum; glucose, GLP-1, glucose-dependent insulinotropic polypeptide (GIP), glucagon, peptide-YY, and cholecystokinin are measured in plasma; and bilirubin levels in duodenal samples are measured. The methods for these assays are known in the art. See, e.g., Hojberg, et al. (2008) *Diabet. Med.* 25:1268-1275. GLP-1 secretion and/or production is expected to be stimulated in subjects receiving a compound described herein.

What is claimed is:

1. A method for synthesizing a cycloalkanyl [b]indole, cycloalkanyl [b]benzofuran or cycloalkanyl [b]benzothiophene comprising


(a) combining, in a single reaction,

(i) a compound of Formula I,




Formula I

(ii) a compound of Formula II or III, and




Formula II




Formula III

(iii) a coupling partner; and


(b) adding an acid catalyst thereby producing a compound of Formula Va, Vb, Vc or Vd




Formula Va



Formula Vb



Formula Vc



Formula Vd

wherein,

each X is the same or different and is independently C-(R¹)_n, OR¹, S or NR¹, wherein each R¹ is independently hydrogen, alkyl, alkenyl, alkynyl or aryl;

each Y is the same or different and is independently N or C;

dashed bonds are independently present or absent;

each R² is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, -R⁷-cycloalkyl, aryl, -NHR⁷, Het, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -CO₂R⁸, -R⁷C(O)₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -R⁷C(S)NR⁸R⁹, -R⁷(NH)NR⁸R⁹, -C(NH)NR⁸R⁹, -R⁷C(NH)NR⁸R⁹, -S(O)₂NR⁸R⁹, -S(O)₂N R⁸aryl, -R⁷S(O)₂NHCOR⁸, -R⁷S(O)₂NR⁸R⁹, -R⁷S(O)₂R⁸, -S(O)_mR⁸, cyano, nitro, or azido group;

R³ and R⁴ are independently hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, -NHR⁷, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -CO₂R⁸, -R⁷C(O)₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -R⁷C(S)NR⁸R⁹, -R⁷(NH)NR⁸R⁹, -C(NH)NR⁸R⁹, -R⁷C(NH)NR⁸R⁹, -S(O)₂NR⁸R⁹, -S(O)₂N R⁸aryl, -R⁷S(O)₂NHCOR⁸, -R⁷S(O)₂NR⁸R⁹, -R⁷S(O)₂R⁸, -S(O)_mR⁸, cyano, nitro, or azido group, or

R³ and R⁴ together form a C₄₋₆ cycoalkyl;

R⁵ is hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het or aryl group;

each R⁶ is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het aryl, -NHR⁷, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -C(O)₂R⁸, -R⁷C(O)₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -R⁷C(S)NR⁸R⁹, -R⁷(NH)NR⁸R⁹, -C(NH)NR⁸R⁹, -R⁷C(NH)NR⁸R⁹, -S(O)₂NR⁸R⁹, -S(O)₂N R⁸aryl, -

$R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^7 is the same or different and is independently selected from an alkylene, cycloalkylene, alkenylene, cycloalkenylene or alkynylene group;

each of R^8 and R^9 are the same or different and are independently selected from the group of hydrogen, an alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7cycloalkyl$, $-R^7OH$, $-R^7(OR^7)_w$, or $-R^7NR^{10}R^{11}$ group;

each of R^{10} and R^{11} are the same or different and are independently selected from the group of an alkyl, cycloalkyl, alkenyl, cycloalkenyl, or alkynyl group;

p is selected from 0, 1, 2, 3, or 4;

q is selected from 0, 1, 2, 3, or 4;

each n independently is 0, 1 or 2;




each m independently is 0, 1 or 2; and


w is 1-10.

2. The method of claim 1, wherein the acid catalyst is a Lewis acid, chiral Lewis acid complex, Brønsted acid, chiral phosphoric acid, or other chiral Brønsted acid.


3. The method of claim 2, wherein the Lewis acid comprises a metal halide or triflate salt of Ga(III) or In(III).

4. The method of claim 1, wherein the coupling partner comprises a compound having the structure of Formula IVa-IVf:






Formula IVd



Formula IVe



Formula IVf

wherein X is C-(R¹)_n, OR¹, S or NR¹, wherein each R¹ is independently hydrogen, alkyl, alkenyl, alkynyl or aryl;

each R⁶ is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, -NHR⁷, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -CO₂R⁸, -R⁷CO₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -R⁷C(S)NR⁸R⁹, -R⁷(NH)NR⁸R⁹, -C(NH)NR⁸R⁹, -R⁷C(NH)NR⁸R⁹, -S(O)₂NR⁸R⁹, -S(O)₂NR⁸aryl, -R⁷SO₂NHCO⁸, -R⁷SO₂NR⁸R⁹, -R⁷SO₂R⁸, -S(O)_mR⁸, cyano, nitro, or azido group;

each R⁷ is the same or different and is independently selected from an alkylene, cycloalkylene, alkenylene, cycloalkenylene or alkynylene group;

each of R⁸ and R⁹ are the same or different and are independently selected from the group of hydrogen, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, -R⁷cycloalkyl, -R⁷OH, -R⁷(OR⁷)_w, or -R⁷NR¹⁰R¹¹ group;

each of R¹⁰ and R¹¹ are the same or different and are independently selected from the group of an alkyl, cycloalkyl, alkenyl, cycloalkenyl, or alkynyl group;

each R¹² is independently the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, -NHR⁷, -NHHet, -NHR⁷Het, -OR⁸, -O-aryl, -OHet, -R⁷OR⁸, -NR⁸R⁹, -NR⁸-aryl, -R⁷NR⁸R⁹, -R⁷NR⁸-aryl, -R⁷C(O)R⁸, -C(O)R⁸, -CO₂R⁸, -R⁷CO₂R⁸, -C(O)NR⁸R⁹, -C(O)aryl, -C(O)NR⁸aryl, -C(O)Het, -C(O)NHR⁷Het, -R⁷C(O)NR⁸R⁹, -C(S)NR⁸R⁹, -R⁷C(S)NR⁸R⁹, -R⁷(NH)NR⁸R⁹, -C(NH)NR⁸R⁹, -R⁷C(NH)NR⁸R⁹, -S(O)₂NR⁸R⁹,

$-S(O)_2N R^8$ aryl, $-R^7SO_2NHCOR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^{13} is independently $C-(R^1)_n$ or OR^1 ;

q is selected from 0, 1, 2, 3, or 4;


each n independently is 0, 1 or 2;

each m independently is 0, 1 or 2;


w is 1-10; and

LG is leaving group.


5. A compound having the structure of Formula Ve, Vf or Vg, or a pharmaceutically acceptable salt or solvate thereof:



Formula Ve



Formula Vf



Formula Vg

wherein

X is $C-(R^1)_n$, OR^1 , S or NR^1 , wherein each R^1 is independently hydrogen, alkyl, alkenyl, alkynyl or aryl;

each Y is the same or different and is independently N or C ;

each Z is the same or different and is independently N or C and wherein Z can optionally be connected to any other Z by a chain of one or more atoms independently selected from substituted or unsubstituted C , N , O , S , or a combination thereof, thereby forming an additional ring(s).

each R^2 is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7$ -cycloalkyl, aryl, $-NHR^7$, Het, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(O)R^8$, $-C(O)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(O)NR^8R^9$, $-C(O)aryl$, $-C(O)NR^8$ aryl, $-C(O)Het$, $-C(O)NHR^7Het$, $-R^7C(O)NR^8R^9$, $-C(S)NR^8R^9$, $-$

$R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(O)_2NR^8R^9$, $-S(O)_2NR^8R^9$, $-R^7SO_2NHR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^6 is the same or different and is independently selected from the group of hydrogen, a halogen, haloalkyl, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, Het, aryl, $-NHR^7$, Het, $-NHHet$, $-NHR^7Het$, $-OR^8$, $-O$ -aryl, $-OHet$, $-R^7OR^8$, $-NR^8R^9$, $-NR^8$ -aryl, $-R^7NR^8R^9$, $-R^7NR^8$ -aryl, $-R^7C(O)R^8$, $-C(O)R^8$, $-CO_2R^8$, $-R^7CO_2R^8$, $-C(O)NR^8R^9$, $-C(O)aryl$, $-C(O)NR^8aryl$, $-C(O)Het$, $-C(O)NHR^7Het$, $-R^7C(O)NR^8R^9$, $-C(S)NR^8R^9$, $-R^7C(S)NR^8R^9$, $-R^7(NH)NR^8R^9$, $-C(NH)NR^8R^9$, $-R^7C(NH)NR^8R^9$, $-S(O)_2NR^8R^9$, $-S(O)_2NR^8R^9$, $-R^7SO_2NHR^8$, $-R^7SO_2NR^8R^9$, $-R^7SO_2R^8$, $-S(O)_mR^8$, cyano, nitro, or azido group;

each R^7 is the same or different and is independently selected from an alkylene, cycloalkylene, alkenylene, cycloalkenylene or alkynylene group;

each of R^8 and R^9 are the same or different and are independently selected from the group of hydrogen, alkyl, alkoxy, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, $-R^7cycloalkyl$, $-R^7OH$, $-R^7(OR^7)_w$, or $-R^7NR^{10}R^{11}$ group;

each of R^{10} and R^{11} are the same or different and are independently selected from the group of an alkyl, cycloalkyl, alkenyl, cycloalkenyl, or alkynyl group;

p is selected from 0, 1, 2, 3, or 4;

q is selected from 0, 1, 2, 3, or 4;

each n independently is 0, 1 or 2;

each m independently is 0, 1 or 2; and

w is 1-10.

6. A pharmaceutical composition comprising a compound of claim 5 in admixture with a pharmaceutically acceptable carrier.

7. A method for stimulating the secretion and/or production of glucagon-like peptide-1 (GLP-1) or inhibiting the activity of Calcitonin Gene-Related Peptide (CGRP) receptor comprising contacting a cell with a compound of claim 5 so that the secretion and/or production of GLP-1 is stimulated or the activity of the CGRP receptor is inhibited.

INTERNATIONAL SEARCH REPORT

013/042148 18-10-2013

International application No.

PCT/US 13/42148

A. CLASSIFICATION OF SUBJECT MATTER
 IPC(8) - A01N 43/38; A61K 31/40; C07D 209/58 (2013.01)
 USPC - 514/411; 548/427

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8)- A01N 43/38; A61K 31/40; C07D 209/58 (2013.01)

USPC- 514/411; 548/427

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 514/415, 443, 468

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Patbase, Freepatentsonline (us pat, ppub, epo, jpo, wipo, npl), Google Scholar (pi, npl); Search Terms; 4+3 cycloaddition, Indium, gallium, Lewis acid, triflate, cyclopentabindole, benzofuran, thiophene, pyrrole, allyl cation, diene, dienophile, Calcitonin Gene-Related Peptide glucagon-like peptide-1

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2004/069831 A1 (BALASUBRAMANIAN et al.) 19 August 2004 (19.08.2004) pg 8-10, Formula I	5-7 -----
Y		1-4
Y	IOVEL et al. 'HYDROXYMETHYLATION OF FURANI AND ITS DERIVATIVES IN THE PRESENCE OF CATION-EXCHANGE RESINS', journal of Molecular Catalysis, 1989, Vol.57, pp 91 - 103. [Downloaded from www.sciencedirect.com on 10 October 2013] Abstract; pg 92, para 4 - pg 94, para 1; Table 1-2; pg 95, scheme	1-4
Y	WINNIE et al., Scope and Mechanism of the (4+3) Cycloaddition Reaction of Furfuryl Cations, Angew. Chem. Int. Ed. 2011, Vol.50, pp 11990-11993. [Downloaded from http://onlinelibrary.wiley.com on 10 October 2013] pg 12196, col 2, para 3 to pg 12197, col 1, para 2, scheme 1; Table 1	1-4
X,P	HAN et al. 'Gallium(III)-Catalyzed Three-Component (4+3) Cycloaddition Reactions', Angew. Chem. Int. Ed. 2012, Vol.51, pp 10390 -10393. Published online: September 11, 2012 [Downloaded from http://onlinelibrary.wiley.com on 10 October 2013] Entire Document	1-4
A	Harmata, Exploration of Fundamental and Synthetic Aspects of the Intramolecular 4 + 3 Cycloaddition Reaction, Acc. Chem. Res. 2001, Vol.34, pp 595-605. pg 598, col 2, para 3 and Scheme 5.	1-4

Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

8 October 2013 (08.10.2013)

Date of mailing of the international search report

18 OCT 2013

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, Virginia 22313-1450
 Facsimile No. 571-273-3201

Authorized officer: .

Lee W. Young

PCT Helpdesk: 571-272-4300
 PCT OSP: 571-272-7774