



(51) International Patent Classification:

C12Q 1/04 (2006.01) *C12N 5/00* (2006.01)
C12Q 1/68 (2006.01) *C12N 5/073* (2010.01)

(21) International Application Number:

PCT/US2014/042880

(22) International Filing Date:

18 June 2014 (18.06.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

13/922,430 20 June 2013 (20.06.2013) US
13/922,445 20 June 2013 (20.06.2013) US
14/026,040 13 September 2013 (13.09.2013) US

(71) Applicant: **ELWHA LLC** [US/US]; 11235 SE 6th Street, Suite 200, Bellevue, Washington 98004-6481 (US).

(72) Inventors: **BANGERA, Mahalaxmi Gita**; 1810 Bremer-ton Ave NE, Renton, Washington 98059 (US). **ECK-HOFF, Philip A.**; 4230 105th Ave NE, Kirkland, Wash-ington 98033 (US). **HYDE, Roderick A.**; 9915 161st Ave NE, Redmond, Washington 98052 (US). **JUNG, Edward K.Y.**; 10091 Park Run Drive #200, Las Vegas, Nevada 89145 (US). **KINDSVOGEL, Wayne R.**; 6014 24th Ave NE, Seattle, Washington 98115 (US). **WOOD, Lowell L., Jr.**; 989 112th Ave NE #2310, Bellevue, Washington 98004 (US).

(74) Agents: **MALASKA, Stephen L.** et al.; 11235 SE 6th Street, Suite 200, Bellevue, Washington 98004-6481 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: RAPID BREEDING OF ANIMALS



(57) Abstract: Various embodiments disclosed herein include systems, methods, compositions, and products by process for in vitro cycling of gametes for rapid breeding of animals. In certain embodiments, the systems and/or methods are at least partially auto-mated.

Rapid Breeding of Animals

All subject matter of the Priority Application(s) is incorporated herein by reference to the extent such subject matter is not inconsistent herewith.

SUMMARY

5 Described herein for various embodiments include systems, methods, compositions, and products by process for rapid breeding of plants. In certain embodiments, the systems and/or methods are at least partially automated.

The foregoing summary is illustrative only and is not intended to be in any way limiting. In addition to the illustrative aspects, embodiments, and features described
10 above, further aspects, embodiments, and features will become apparent by reference to the drawings and the following detailed description.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 illustrates a partial view of an embodiment disclosed herein.

FIG. 2 illustrates a partial view of a seed.

15 **FIG. 3** illustrates a partial view of an embodiment disclosed herein.

FIG. 4 illustrates a partial view of an embodiment disclosed herein.

FIG. 5 illustrates a partial view of an embodiment disclosed herein.

FIG. 6 illustrates a partial view of a sexual reproduction cycle in plants.

FIG. 7 illustrates a partial view of an embodiment disclosed herein.

20 **FIG. 8** illustrates a partial view of an embodiment disclosed herein.

FIG. 9 illustrates a partial view of an embodiment disclosed herein.

FIG. 10 illustrates a partial view of an embodiment disclosed herein.

FIG. 11 illustrates a partial view of an embodiment disclosed herein.

FIG. 12 illustrates a partial view of spermatogenesis.

25 **FIG. 13** illustrates a partial view of oogenesis.

DETAILED DESCRIPTION

In the following detailed description, reference is made to the accompanying drawings, which form a part hereof. In the drawings, similar symbols typically identify similar components, unless context dictates otherwise. The illustrative embodiments

described in the detailed description, drawings, and claims are not meant to be limiting. Other embodiments may be utilized, and other changes may be made, without departing from the spirit or scope of the subject matter presented here.

It is highly beneficial to evolve plant breeding to include characteristics that are
5 desirable without the time-scales usually required for growing plants to maturity for such plants that reproduce sexually. Described herein include various embodiments related to a focused means for selecting and breeding desired cultivars with pre-determined or naturally resulting desired characteristics. Various embodiments disclosed herein provide a higher rate of improvement of desired characteristics, such as increased yield, or drought
10 resistance, among others, which assists in sustainable agriculture practices. In this way, certain embodiments disclosed herein realize multi-generation sequences of gametes by the classic sexual reproductive route, without intervening phenotypes with their attendant complexities and latencies. Furthermore, various embodiments disclosed herein may be utilized to derive new cultivars particularly amenable to certain environmental conditions,
15 or needs for obtaining plant products. Additionally, the disclosed embodiments herein allow for a reduction of labor and field resources for plant breeding, increased capacity to rapidly breed a large number of plants per field unit, as well as an increased capacity to analyze a larger number of plants prior to breeding for desirable characteristics.

Various embodiments disclosed herein relate to genotype-to-genotype generational
20 stepping without an intervening phenotype but instead result from generating male and female gametes directly from plant stem cells. For example, first generation stem cells are induced to derive gametes, which are bred. The resulting fertilized seed includes an embryo from which second generation stem cells may be obtained, if additional breeding is desired. This multi-generational cycling of gametes *in vitro* (CoGiV) allows for allele
25 valuation and selection for rapid and efficient directed evolution of the particular plant species toward one or more desired characteristics. For example, for specific agricultural crops it is desirable to increase the number of progeny and decrease the amount of time needed to screen and select the particular progeny. In another example, it is desirable to increase the precision of breeding-generated phenotypes or the breadth of accessible
30 genetic variation. As described herein, multiple cycles of breeding and/or selecting are conducted depending on genetic complexity, and desired characteristics. In an embodiment, the first round of fertilization is referred to as “first generation fertilization” and the gamete therefrom is referred to as “first generation gamete” etc. while the second

round of fertilization is referred to as “second generation fertilization” and the gamete therefrom is referred to as a “second generation gamete,” etc.

As disclosed herein, in an embodiment described are an increase in efficiency of Rapid-breeding, and do not require genetic or other modification of the plant genome.

5 However, in an embodiment, modification of plant cells (genetic, epigenetic, transcriptional or translational, etc.) is utilized with the Rapid-breeding process.

At any number of steps, standard breeding techniques may be implemented in conjunction with the steps of the Rapid-breeding methods and systems disclosed herein. For example, in conjunction with the steps of Rapid-breeding of plants, conventional plant
10 crossing may be conducted for selecting starting material (e.g., plant stem cells, gametophytes, or sporophytes, etc.), or once a first, second, third, etc. generation of progeny have been developed. Standard plant crossing includes, for example, obtaining seeds of the first parent plant (may be modified, such as transgenic, or not modified) and a second parent plant and growing the seeds into second generation mature parent plants.
15 Next, pollinating a flower from the first parent with pollen from the second parent and harvesting seeds produced on the parent plant bearing the fertilized flower is conducted. As described herein, introgressing target genes by backcrossing may be accomplished, for example, by first crossing a superior inbred plant (recurrent parent A) to a donor inbred plant (non-recurrent parent B), which carries the suitable gene(s) for the desired
20 characteristic, and the progeny of this cross are selected for the desired characteristic for transference from parent B and selected progeny are mated back to parent A. Following five or more backcross generations with selection for the desired characteristic, the progeny are hemizygous for loci controlling the characteristics being transferred but are similar to the recurrent parent A for almost all other genes. The final backcross generation
25 is self-pollinated to generate progeny which are purebred for the desired gene. However, as described herein, this conventional backcross breeding technique is laborious and time consuming, while the Rapid-breeding methods and systems disclosed herein are able to accomplish the same task in a much more efficient manner. Still, in certain instances one or more backcrosses may be implemented, for example, in conjunction with the Rapid-
30 breeding steps of certain disclosed embodiments.

In an embodiment, a method for plant breeding includes selecting one or more plant stem cells; inducing at least one of the plant stem cells to differentiate into at least one first gamete; combining the at least one first gamete with at least one second gamete

of opposite gender for fertilization; and molecularly or microscopically analyzing the progeny resulting from the fertilization.

In an embodiment, the plant stem cell includes a cell isolated from the meristem of a plant (e.g., the apical meristem or lateral meristem). In an embodiment, the plant stem
5 cells are isolated from meristematic tissues such as the root apical meristem, shoot apical meristem, or vascular system ((pro) cambium or vascular meristem, for example). In an embodiment, plant stem cells are isolated from cambium.

As described herein with regard to various embodiments, plant stem cells, similarly to animal stem cells, are totipotent cells that facilitate plant growth and production of plant
10 tissues. Plant stem cells have the ability to self-renew, as well as differentiate into the various different cell types of the plant.

In an embodiment, callus (dedifferentiated cells) is utilized to give rise to totipotent embryogenic cells for utilization with one or more processes described herein. Callus can be initiated from various plant tissues, including but not limited to, immature embryos or
15 parts of embryos, seedling apical meristems, microspores, etc. and result in recipients of genetic information and give rise to fertile plants.

In an embodiment, the cell wall of the plant cell is disrupted or removed (e.g., enzymatically) prior to modification or other use with Rapid-breeding. In an embodiment, meristematic tissue including one or more plant stem cells is utilized for modification as
20 described in the various embodiments disclosed herein.

In an embodiment, a method includes selecting one or more plant stem cells; inducing at least one of the plant stem cells to differentiate into at least one first gamete; combining the at least one first gamete with at least one second gamete of opposite gender for fertilization; and molecularly or microscopically analyzing the embryo resulting from
25 the fertilization.

In an embodiment, microfluidics devices or systems may be utilized for selecting plant stem cells (e.g., by markers) or selecting progeny generated by various embodiments disclosed herein.

In an embodiment, a method includes selecting one or more plant stem cells;
30 inducing at least one of the plant stem cells to differentiate into at least one first gamete of first genetic composition; combining the nucleus of the at least one first gamete with the nucleus of the at least one second gamete of second genetic composition for fertilization; and molecularly or microscopically analyzing the seed resulting from the fertilization. In

this particular embodiment, the gametes may be from the same or opposite gender. In an embodiment, the gamete may be from different genetic composition if diversity is desired, or the same genetic composition if the “ideal” or goal genetic composition has already been determined or obtained, or there are particular characteristics desired to be preserved and passed on to the progeny. In an embodiment, the progeny is/are generated by combining the nuclei containing the chromosomes that yield a cell having the ability to sustain itself long enough to be utilized in an additional round of breeding, and/or to go on to develop into a plant. For example, nuclei may be extracted from pollen by purification from the pollen wall by enzymatic, chemical, mechanical, or osmotic means. *See*, for example, Dewitte, et al., Ch.4, “Use of 2n Gametes in Plant Breeding,” (2012) available online at intechopen.com, the subject matter of which is incorporated herein by reference. For example, the outer exine layer on the pollen surface is a biopolymer highly resistant to enzymatic or hydrolytic breakdown, as it has evolved to withstand environmental stresses. However, several physical techniques, such as bead beating or chopping of pollen grains, has allowed for release of nuclei from pollen grains sufficient to study the nuclei by flow cytometry. *Id.*

In an embodiment, inducing gametes includes inducing plant stem cells first to differentiate to sporophyte stage. In an embodiment, gametes are derived from sporophyte stages without first being induced from stem cells. In an embodiment, inducing gametes includes inducing plant stem cells first to differentiate to gametophyte stage. In an embodiment, gametes are derived from gametophyte stage without first being induced from stem cells.

For example, culturing microspores gives rise to pollen, doubled haploid embryos, and apoptotic cells, depending on the culture conditions, including hormones, and sugars. *See* Wang et al., Plant Physiol. (2000), 124: 523-530, which is incorporated herein by reference. Furthermore, since induction of differentiation is considered a stress condition, specific stress hormones, such as abscisic acid (ABA) and its signal transduction pathway are important factors, as well as lipid transfer proteins, heat shock proteins, and peroxiredoxin anti-oxidant. *Id.*

In an embodiment, the gametes and/or stem cells are genetically analyzed. Genetic analysis for embodiments described herein includes genetic sequence analysis, ploidy analysis, or marker selection, for example. In an embodiment, mRNA sequence information is obtained at one or more steps in the plant breeding process, for one or more

cell types (e.g., stem cells, fertilized seed, gametes, etc.). In an embodiment, marker assisted selection is utilized to screen genetic markers in the very early progeny (zygote, blastocyst, embryo, etc. stage).

In an embodiment, the plant stem cells are treated with one or more transcription factors in order to derive gametes therefrom. For example, MADS-box genes, encoding MADS-domain family of transcription factors that bind DNA, are involved in controlling various cell development and proliferation processes. *See* Gramzow and Theissen, *Genome Biol.*, 2010, 11:214, which is hereby incorporated by reference. For example, two main types of MADS-domain transcription factors have been characterized, including Type I and Type II, whereas Type I MADS-box genes have usually one or two exons, Type II genes have an average of seven. *Id.* Likewise, MIKC-type MADS-box genes have been identified as being involved in plant reproductive organ formation, particularly MIKC* and MIKC^c. *Id.* The MADS-box genes are regulated in various ways, including transcriptional regulation by transcription factors (including feedback and feed-forward loops), epigenetic control, and microRNA regulation. *Id.*

In an embodiment, selecting one or more plant stem cells includes choosing at least one plant stem cell tissue (e.g., meristem tissue) containing plant stem cells based at least one criteria. In an embodiment, selecting one or more plant stem cells includes choosing from at least one plant stem cell tissue, one or more single cells isolated from the remaining tissue. In an embodiment, one or more single plant cells may be isolated from the plant stem cell tissue, for example, by physical or chemical means (e.g., dissection, flow cytometry, enzymatic extraction, lysis of tissue, etc.).

In an embodiment, inducing at least one of the plant stem cells to differentiate into at least one first gamete includes inducing to differentiate to at least one of a plant ovule, megagametophyte, or egg cell in the case of a female gamete; or at least one pollen grain, microgametophyte, or plant sperm cell in the case of a male gamete.

For example, light intensity, light/dark period, and temperature influence sexual reproduction in plants. *See* for example, Hohe, et al. (2002), *Plant Biol.* 4: 595-602, which is incorporated herein by reference. As another example, MADS-box genes are involved in development of sexual reproduction plant components as well as seed development, and root, flower, and fruit development.

In an embodiment, a plant somatic cell is inverse-differentiated into a germ cell precursor and forward-specialized into pre-gametes of either sex and matured in

appropriate cellular milieu into fertilization-ready gametes. In an embodiment, plant stem cells are utilized in the process, rather than inverse-differentiated somatic cells. Post-fertilization, in an embodiment the zygote resulting from fertilization is analyzed directly, or cultured to at least a blastula stage, at which stage multiple cells are provided for next-generation pre-gametes. Following maturation into gametes of both sexes, each of these may be employed (in potentially quite wide) crossings aimed at realization of specific allelic traits and/or the exclusion of others, as determined in near-real time by endosperm and/or polar body analysis. As described herein, such Rapid-breeding and analyses may be done at each generation step, and in parallel and/or with highly-automated means that allows for rapid trait selection at each generational step. In an embodiment, the germination viability of the analyzed plant or plant tissue (such as a seed portion or in its entirety) is maintained.

For any given round of breeding, the method includes selecting naturally occurring characteristics that emerge in the crossing of the parent cells. Multiple rounds of such a selection process without the necessity of culturing the resulting offspring yields a rapid-breeding approach that reduces the time needed for natural selection of desirable characteristics to a fraction of that required for traditional plant breeding that allows the offspring to grow up to plantlets or sexually mature adult plants.

For any given round of breeding, one or more genes may be inserted, deleted, or mutated in order to produce a resulting seed bearing the gene or stacks of genes of interest. In an embodiment, for example, the plant stem cells are modified prior to induction of gametes. In another embodiment, the gametes are modified prior to fertilization. In an embodiment, the plant stem cells are modified transgenically by one or more gene insertions, deletions, or mutations. In an embodiment, the plant stem cells are modified epigenetically by altering gene expression through changes in methylation, for example. See for example, Downen et al., PNAS, pp. 1-9, 2012 available online at pnas.org. For example, in *Arabidopsis*, DNA methylation is deposited at CC, CHG, and CHH sequences (where H is A, C, or T) through three genetically separable pathways to regulate transposon silencing, genomic imprinting, and stable gene silencing. *Id.* Furthermore, simultaneous disruption of the chromatin remodeling enzymes KYP, SUVH5, and SUVH6 in *Arabidopsis* results in concomitant decrease in cytosine methylation and H3K9me2 levels, and consequently transcriptional reactivation of heterochromatic transposons. Cytosine methylation is established in all sequence contexts by de novo

methyltransferases (DRM1/2) through a small RNA-directed DNA methylation (RdDM) pathway. *Id.* DICER-dependent 21 to 24 nucleotide siRNAs guide Argonaute proteins (AGO4/AGO6) to complementary sequences within the genome, likely through a siRNA:nascent RNA base pairing mechanism, to direct cytosine methylation. *Id.*

- 5 Methylation of CGs and CHGs are maintained through DNA replication by MET1, a homologue of the mammalian DNA methyltransferase DNMT1, and the plant-specific CMT3 methyltransferase, respectively. *Id.* Conversely, active demethylation of methylcytosines is catalyzed by DEMETER (DME) family of DNA glycosylases. *Id.* Moreover, methylation alterations also assist in immune response to pathogens. *Id.*

- 10 In an embodiment, the one or more modifications of plant cells described herein are reversible. In an embodiment, the one or more modifications occur in DNA of the plant cell nucleus. In an embodiment, the one or more modifications occur in DNA of the mitochondria of the plant cells. In an embodiment, the plant stem cells or gametes are modified by the addition, deletion, or mutation of one or more mRNA molecules or
 15 proteins utilized in cell transcription or translation, respectively. In an embodiment, the genetic or epigenetic modification of the plant cells yields transformed progeny plants with a genome that has been altered by the stable integration of a recombinant DNA.

20 Plants

- Various embodiments described herein are applicable to a number of plants, including but not limited to grass, fruit, vegetable, flowering trees and plants (e.g., ornamental plants, fruit plants, such as apple and cherry, etc.), grain crops (e.g., corn, soybean, alfalfa, wheat, rye, oats, barley, etc.), other food or fiber crops (e.g., canola,
 25 cotton, rice, peanut, coffee, bananas, sugar cane, melon, cucumber, sugar beet, quinoa, cassava, potato, onion, tomato, strawberry, cannabis, tobacco, etc.), or other plants (including but not limited to banana, bean, broccoli, castorbean, citrus, clover, coconut, Douglas fir, Eucalyptus, Loblolly pine, linseed, olive, palm, pea, pepper, poplar, truf, Arabidopsis thaliana, Radiata pine, rapeseed, sorghum, or Southern pine. Most of the
 30 calories consumed by humans come from members of the grass family (e.g., wheat, corn [maize], rice, oats, barley, sorghum, millet, rye, etc.), and grasses make up at least a quarter of all vegetation on Earth, rendering these important food crops worldwide. Various embodiments described herein are applicable to plant cells, seeds, pollen, fruit,

zygotes, etc., as disclosed. In addition, in an embodiment, a hybrid of two different plants is generated by the fertilization of gametes from each plant. In an embodiment, the one or more gametes utilized in such a fertilization event are modified in order to increase their compatibility with each other thus resulting in a successful fertilization.

5 Various embodiments disclosed herein are utilized to improve one or more characteristics of a plant by way of rapid selection of desirable naturally occurring characteristics. Various embodiments disclosed herein are utilized to improve one or more characteristics of a modified plant (e.g., genetic, or epigenetic modification) and include rapid selection of desirable characteristics introduced into the plant by modification as
10 described. For example, such characteristics include but are not limited to increased telomere(s), improved agronomic characteristic (e.g., improved yield, nutritional content, physiology, growth or development, stress resistance or tolerance (e.g., disease, pest, or chemical resistance or tolerance), increased seed oil or protein content. In an embodiment, for example, such characteristics include one or more of resistance or tolerance to drought,
15 fungal, bacterial, or viral disease, temperature fluctuation, light fluctuation, exposure to extreme temperature (e.g., heat exposure, or cold exposure), exposure to extreme light conditions (e.g., abundance of light, shade or too little light), insect or worm infestation, low nutrient environment (e.g., low nitrogen or phosphorous availability), osmotic stress, or high plant density. For example, the plant yield includes such properties as plant
20 height, plant structure, assimilation of nutrients (e.g., carbon), pod number or position on the plant, pod shatter, grain size, number of internodes, efficiency of nitrogen fixation, resistance to abiotic or biotic stress, seedling vigor, overall percentage of seed germination, resistance to lodging, growth rate, seed number or size, or seed composition.

 In an embodiment, analyzing the resulting progeny (e.g., fertilized seed or resulting
25 plant(let) therefrom) includes analyzing for an improvement in a particular characteristic, whether such characteristic is attributed to a naturally occurring trait or is the result of a modification (e.g., physical, chemical, genetic, epigenetic intervention, etc.). In an embodiment an improvement in a particular characteristic includes an increased level of a particular characteristic in the progeny, relative to a control plant (in certain instances, the
30 control plant may be a parent plant). For example, an increased level of a particular characteristic may be measured as described herein, including increased seed size or increased number of seeds per unit measured.

In an embodiment, analyzing the resulting progeny includes evaluating one or more characteristics including a genetic marker, a single nucleotide polymorphism, a simple sequence repeat, a restriction fragment length polymorphism, a haplotype, a tag single nucleotide polymorphism, a gene, an allele of a genetic marker, a DNA-derived sequence, an RNA-derived sequence, promoter, 5'-untranslated region of a gene, 3'-untranslated region of a gene, microRNA, siRNA, a QTL, satellite marker, transgene, mRNA, high-resolution chromosomal structure and content analyses with microarrays, double stranded mRNA, transcriptional profile, or methylation pattern.

In an embodiment, an improvement in a particular characteristic includes a decreased level of a particular characteristic in the progeny, relative to a control plant (again, in certain instances, the control plant is the parent plant). For example, a decreased level of a particular characteristic may be measured as described herein, including decreased use of water or nutrients from the soil.

The various characteristics described may be analyzed for improvement in several ways. For example, chemistry, cell biology, or molecular biology techniques may be utilized for measuring telomere changes, ploidy, or gene, protein, or mRNA changes, seed biomass, seed content (e.g., oil, carbohydrate, protein, nitrogen, phosphorous, etc. or ratios of one or more of these seed components), epigenetic changes, etc. As another example, ploidy may be evaluated by flow cytometry or microscopic analysis. In an embodiment, the desired ploidy plants or plant tissues (including seeds) are selected and those that are not of the desired ploidy are removed from the desired ploidy plants/plant tissues. In an embodiment, the plant tissue is analyzed for one or more alleles, or ploidy level of at least one locus.

As another example, an immunoreactive antibody may be utilized to detect the presence or absence of a protein that is expressed or suppressed in the progeny seed or plant tissue.

Other changes, including improvements, in the various characteristics described may be measured by physical means, for example, plant weight, seed weight, seed number per plant, seed number or seed weight per acre, bushels per acre, tons per acre, kilo per hectare, etc. As another example, a small portion of tissue is removed from the fertilized seed without disturbing the viability of the seeds, and such tissue is analyzed by means described herein.

In an embodiment, when autopolyploid plants (such as alfalfa) are utilized in the described methods, gametes are optionally selected depending on their ploidy, for example, gametes that are bivalents or quadrivalents resulting from meiosis may be selected (univalent or trivalents resulting from triploids are usually sterile). If polyploidy
5 plants are desired for use with various embodiments described herein, inhibition of mitosis artificially induces polyploids. For example, high temperatures or use of chemicals to inhibit mitosis (such as colchicine, EMS, GA₃, temperature, trifluralin, N₂O, or dinitroanilines) may be used to induce polyploids.

In an embodiment, a polyploidy plant may be desired due to the increased cell size
10 or cell volume (including, for example, enlarged plant organs) that is a result of additional chromosomes. For example, tetraploid grapes yield greater juice content than their diploid counterparts, and ornamental crops with increased numbers of chromosomes have higher quality and size of blossoms. See, for example, Olmo (1952) American. Soc. Hort. Sci. 59: 285-290, which is incorporated herein by reference.

Likewise, in an embodiment doubled haploid cells are desired and may be
15 generated in over 250 plant species. See, for example, Malunzyski, et al., 2003 Doubled haploid production in crop plants. A manual. Kluwer Acad. Pub. Dordrecht, Boston, London, which is incorporated herein by reference. Doubled haploid plants may be produced *in vivo* or *in vitro*, for example. Haploid embryos are produced *in vivo*, for
20 example, by parthenogenesis, pseudogamy, or chromosome elimination following wide crossing.

Doubled haploid plants save time in generating homozygous lines essentially without the need for conventional breeding. In certain embodiments disclosed herein, doubled haploids are utilized for Rapid-breeding. For example, typically the first step for
25 doubled haploid production includes haploidization of the plant with production of haploid seed. Non-homozygous lines are crossed with an inducer parent, resulting in the production of haploid seed, and selection of seed that contains a haploid embryo independent of the ploidy of the endosperm (the process of which normally results in triploid endosperm). Subsequently, selected haploid seeds (that is haploid embryo in the
30 seed) undergo chromosome doubling to generate doubled haploid seeds. Spontaneous chromosome doubling leads to normal gamete production or production of unreduced gametes from haploid cell lineages, while addition of colchicine or other mitotic disruptor can also be used to increase the rate of diploidization. The chimeric plants are self-

pollinated to produce diploid (doubled haploid) seed, which is cultivated and/or evaluated and used in hybrid testcross production.

In an embodiment, plant tissue from the doubled haploid is analyzed, and utilized further in methods described herein. Furthermore, by utilizing endosperm tissue derived from a diploid plant, the ploidy level of the particular genetic marker is determined, with a
5 diploid level indicating maternal inheritance and a haploid level indicating paternal inheritance. This is due to the endosperm tissue being triploid, with two copies of the gene derived from the female gamete, which allows for analysis of linkage phase of the parental line by evaluating heterozygous progeny genotypes that indicates different allele
10 frequencies in DNA samples.

For example, maize embryo zygoty includes triploid endosperm and diploid embryo tissue, thus endosperm copy number indicates the zygoty of the embryo, that is, a homozygous endosperm accompanies a homozygous embryo, while a heterozygous endosperm indicates a heterozygous embryo. For example, endosperm that is
15 homozygous for an internal control gene will contain three copies of the gene, whereas the gene of interest varies from 0 to 3, depending on the heritability and zygoty of the resulting progeny.

It is useful to determine zygoty of one or more loci during plant breeding in order to evaluate the degree of fixation, or inbreeding, segregation distortion (i.e., in transgenic
20 germplasm, maternal inheritance testing of for loci that affect gametes).

For example, determination of allelic frequencies assists in determining parental linkages for a specific marker, and comparing allele frequency data between two or more germplasm pools provides guidance with selection of targeted desired characteristics in such a way that as alleles increasing in frequency in conjunction with a shift in distribution
25 of one or more characteristics, such is presumed to be linked to the trait of interest. Furthermore, determining relative allele frequency data assists in construction of genetic linkage maps. For example, several species of plants have had full or partial genomic sequence analysis conducted, including rice varieties, eucalyptus varieties, strawberry varieties, apple varieties, grape varieties, tomato varieties, potato varieties, wheat varieties,
30 maize varieties, oil plants, cereal crops, biodiesel crops, various algae, and others. Thus, genetic sequences for particular plants, or the homologs from other species, are readily available or attainable for use with various embodiments described herein.

In certain embodiments, selected progeny are bulked, depending on the analysis of the resulting genotype. Conversely, if multiple characteristics are analyzed by way of QLT or genomic selection, for example, with varying effects being selected for a particular population, the progeny may keep individual identity preserved, and cultivate
5 the progeny for determination of various combinations of the desired QTL.

In an embodiment, methods disclosed herein relate to introgressing a characteristic into a plant, for example, by removing a sample of cells or nucleic acids from plant tissue (e.g., seeds), and analyzing the nucleic acids extracted from each seed for the presence or absence of at least one genetic marker, selecting seeds from the population based on the
10 results of the nucleic acid analysis, and cultivating a plant for use with a second generation of breeding.

In an embodiment, cycling of gametes *in vitro*, as described herein, accelerates the introgression of a specific gene or genes into a plant and allow for greater precision by more cycles of recombination.

Likewise, genetic recombination can involve “gene conversion” or the non-reciprocal transfer of DNA from one genotype to another. Utilizing various embodiments disclosed herein, vast populations for rare zygotes with gene conversion at a target locus may be screened to identify a true isogenic derivative rather than a nearly-isogenic backcross derivative in which flanking alleles from the donor may confer undesirable
15 characteristics. Gene conversion may be especially useful for manipulating exotic germplasm in which a desirable gene is often surrounded by undesirable ones and thus methods disclosed herein assist in avoiding intermediate phenotypes.

Sexual plant breeding results in the addition and transfer of chromosomes, which allows for increased diversity in the progeny, and allows for selection of particular
25 desirable characteristics. Chromosome doubling *in vitro* induces polyploidy, by utilizing mitotic inhibitors such as dinitroaniles and colchicine. As described herein, ploidy status may be determined in a number of cell or molecular biology techniques, including, for example, chloroplast count in guard cells, examining morphological features such as leaf, flower, or pollen size (termed the gigas effect) and flow cytometry.

Further as described herein, various embodiments disclosed herein utilize
30 polyploidy induction for Rapid-breeding techniques. For example, allopolyploids may be generated from hybridization of two or more genomes followed by chromosome doubling or by fusion with unreduced gametes between or within species. For example, meiosis or

mitosis takes a diploid cell and reduces it to 4 haploid cells (one copy of each gene and chromosome). However, an unreduced gamete has not undergone the reduction (e.g., by inhibiting cell division) and remains diploid. *See* Figures disclosed herein. In an embodiment, a method disclosed herein includes inducing at least one plant stem cell to
5 differentiate into at least one first gamete. In an embodiment, the at least one first gamete is unreduced (i.e. has not yet undergone meiosis) such that ploidy is induced upon fertilization with a second gamete. In an embodiment, the nuclei are combined instead of the entire gamete(s).

For example, a dominant mutation coding for AGO104, a member of the
10 ARGONAUTE family of proteins, results in formation of functional unreduced gametes. *See* Singh, et al., Plant Cell Vol.23: 443-458 (2011), which is incorporated herein by reference. The mutant shows defects in chromatin condensation during meiosis and subsequent failure to segregate chromosomes when screened in *Zea mays*. *Id.* AGO104 is needed for non-CG methylation of centromeric and knob-repeat DNA. Thus, in an
15 embodiment, a mutant AGO104 is utilized for differentiating unreduced gametes.

In certain instances, the hybridized genomes may differ in their degree of homology such that some genomes are able to pair during mitosis and/or meiosis, while others are not. For example, in certain instances only segments of the chromosomes of the combining genomes differ (termed segmental allopolyploidy), and in order to achieve a “pure
20 cross” a bridge cross, or intervening cross, may be utilized to achieve the desired hybrid. For example, in an embodiment a cow may be generated that includes the desired characteristics of both dairy cattle and beef cattle such that the single animal generated by the method(s) described includes characteristics of both and thus both meat and milk products can be obtained from the same animal.

25 Likewise, the multiallelic nature of loci in polyploids allows for masking deleterious alleles that may arise during modification or polyploidy induction. Further, polyploids are able to tolerate deleterious allele modifications post-mutation and they have increased mutation frequency due to their large genomes. Polyploids typically have larger cells and/or plant organs than their traditional counterparts, however the larger cell volume
30 is usually due to higher water retention, rather than an increase in biomass.

In an embodiment, the progeny are analyzed for example, by statistical sample testing of plant tissue(s) (including seeds) for further use. In an embodiment, a portion of the plant tissue(s) are utilized while maintaining the germination viability of the progeny.

For example, a portion of the endosperm, zygote, embryo, blastocyst, or other seed portion may be obtained by mechanical or chemical means and utilized for analysis. For example, an automated seed sampler may be used with various embodiments disclosed herein. *See* for example, U.S. Patent No. 7,941,969, which is incorporated herein by reference.

5 In an embodiment, the one or more desired characteristics are quantified based on the analyzed plant tissue. In an embodiment, the one or more quantified desired characteristics are compared with control plant tissue samples or known germplasm pools in order to identify frequency shifts. Likewise, DNA and RNA may be extracted from plant tissue(s) sufficient to yield DNA or RNA adequate for PCR, RT-PCR, TaqMan
10 assays, sequencing, arrays or blots, or other amplification.

 Several criteria may be established in order to determine whether additional breeding is desired. For example, a small flake from the fertilized seed (optionally retaining the embryo intact for planting, if desired) is removed for genetic analysis of a desired characteristic, whether the corresponding gene(s) are naturally occurring in the
15 seed or artificially inserted, deleted, or mutated.

 As described herein, in certain aspects a method disclosed includes molecularly modifying a plant cell. In an embodiment, a DNA construct is utilized to modify a plant cell. Assembly of DNA constructs may be done using standard methods, typically with a promoter operably coupled to the DNA, and optionally other regulatory elements such as
20 5' leaders, introns, 3' untranslated regions (such as polyadenylation signals), signal peptides, or repressor or enhancer sequences.

 For example, promoter elements include non-constitutive promoters (such as spatially specific promoters, temporally specific promoters, inducible promoters, etc.), or constitutive promoters. For example, spatially specific promoters include organelle, cell,
25 tissue, or organ-specific promoters functional in a plant (such as plastid-specific, root-specific, pollen-specific, or seed-specific promoter for suppressing expression of the target RNA in plastids, roots, pollen, or seeds, respectively). In an embodiment, a seed-specific, embryo-specific, gamete-specific, stem cell-specific, aleurone-specific, or endosperm-specific promoter is utilized with an embodiment. For example, temporally specific
30 promoters can include promoters that favor promoter expression during certain developmental stages in a plant's growth or reproductive cycle, or during different times of day or night, or different seasons of the year. Inducible promoters include promoters induced by chemicals such as exogenous or synthetic chemicals as well as endogenous

pheromones and other signaling molecules) or by environmental conditions such as but not limited to biotic or abiotic stress (e.g., water deficit or drought, temperature fluctuations or extremes, nutrient fluctuations or extremes, salt levels, light levels, pest or pathogen infection, or physical damage, etc.). An expression-specific promoter also
5 includes generally constitutively expressed promoters that vary on the degree of expression.

In addition to promoter elements, in certain instances terminator elements may be utilized in the embodiments disclosed herein, including for example, polyadenylation signals as described herein. Additionally, spacer DNA segments (found, for example,
10 between parts of a gene suppression element or between different gene suppression elements) may include translatable DNA encoding a target gene, translatable DNA encoding a marker or reporter gene, transcribable DNA derived from an intron, transcribable DNA encoding RNA that forms a structure such as a loop, stem, or aptamer capable of binding to a specific ligand, spliceable DNA including introns and self-splicing
15 ribozymes, transcribable DNA encoding a target gene detectable by molecular biology techniques.

Promoter and enhancer elements that are active in plants include but are not limited to nopaline synthase (NOS) promoter, octopine synthase (OCS), CaMV35S promoter (cauliflower mosaic virus), rice actin promoter, maize chloroplast aldolase promoter,
20 maize aldolase promoter, napin, maize L3 oleosin, zein Z27, globulin I, glutelin 1, peroxiredoxin antioxidant (Per 1), maize nicotianamine synthase promoter, etc. Numerous other promoters that function in plant cells (whether plant cell or another cell was the original source) may be utilized in assembling constructs for use in modifying plant cells as described herein.

25 Likewise, various enhancer sequences, such as 5' introns of rice actin 1 and rice actin 2 genes, maize alcohol dehydrogenase gene intron, maize heat shock protein 70 gene intron, maize shrunken 1 gene, etc. may be used to assist in increasing gene expression.

In various aspects, one or more genes or other modifications may be implemented with various embodiments disclosed herein, by for example, making multiple
30 modifications to a single cell or by making single mutations in a cell and crossing those cells to result in a transformed plant. Such modifications may be stacked into the resulting plant with various modifications. For example, characteristics for increasing drought resistance may be stacked with characteristics for increasing yield, or tolerating herbicides.

For example, transgenic plants have been produced that have demonstrated tolerance to glyphosate, dicamba, glufosinate, sulfonyleurea, bromoxynil, and norflurazon, and such plants may be further modified utilizing various embodiments described herein.

Some non-limiting examples of 3' polyadenylation signals include nos 3', tml 3',
5 tmr 3', tms 3', ocs 3', tr7 3', and others from *Agrobacterium tumefaciens* genes, heat shock protein 17 3' from wheat (*triticum aestivum*), wheat ubiquitin gene, wheat fructose-1, 6-biphosphatase gene, rice glutelin gene, rice lactate dehydrogenase gene, rice beta-tubulin gene, pea ribulose biphosphate carboxylase gene (rbs 3'), etc.

With regard to gene suppression, various modes may be used, for example,
10 including anti-sense, co-suppression, and RNA interference (miRNA, dsRNA, siRNA, and other forms of double stranded RNA that block translation). Several microRNA genes have been characterized in plants, including promoters for the same. *See* for example, U.S. Patent No. 8,395,023, which is incorporated herein by reference. Transposable elements can also be utilized to suppress gene function. In certain instances, mutations by
15 insertion of one or more transposable elements disrupts protein translation from a particular gene.

Modification of plant cells by transformation with molecular constructs may be used with various embodiments described herein. For example, agrobacterium-mediated transformation and microprojectile bombardment have been described for use with
20 soybean, corn, wheat, rice, and sugar beet. *See* U.S. Patent No. 8,410,336, which is incorporated herein by reference. Culturing conditions for transforming plant cells includes, for example, nutrient media, kept at conditions which facilitate transformation of the cells. Plant cell targets for transformation include at least one of meristem cells, calli, hypocotyls, gametes (microspores, pollen, sperm, egg, etc.), or immature embryos. In
25 addition, callus may be differentiated from tissues such as immature embryos, hypocotyls, microspores, seedling apical meristems, etc. and can be grown into a plantlet or mature plant. Since typically not all cells are transformed when modified in this manner, one or more markers (e.g., marker genes) are generally utilized to provide an efficient system for identification of the cells that are stably transformed with DNA constructs. Further, if
30 conferring resistance to an antibiotic or herbicide, for example, transformation may be screened by application of the antibiotic or herbicide to the plant (or plant cells).

In an embodiment, a target sequence includes at least one of amino acid catabolic genes (such as maize LKR/SDH), maize zein genes, fatty acid synthesis genes (such as

plant microsomal fatty acid desaturases or plant acyl-ACP thioesterases), genes involved in multi-step biosynthesis pathways (such as genes encoding enzymes for polyhydroxyalkanoate biosynthesis), genes encoding cell-cycle control proteins (such as proteins with cyclin-dependent kinase (CDK) inhibitor-like activity, genes from
 5 undesirable proteins (such as allergens or toxins), or enzymes for the biosynthesis of undesirable compounds (such as undesirable flavor or odor components). For example, a modified plant disclosed herein includes a plant with decreased allergenicity (such as peanut), regulated fruit ripening, a pest or pathogen resistant plant, or an herbicide tolerant plant.

10 For example, several non-limiting examples of pathogens include but are not limited to fungal, bacterial, or viral pathogens. For example, *Phakospora pachirhizi* (Asian soy rust), *Puccinia sorghi* (corn common rust), *Puccinia polysora* (corn Southern rust), *Fusarium oxysporum* and other *Fusarium* spp., *Alternaria* spp., *Penicillium* spp., *Pythium aphanidermatum* and other *Pythium* spp., *Rhizoctonia solani*, *Exserohilum*
 15 *turcicum* (Northern corn leaf blight), *Bipolaris maydis* (Southern corn leaf blight), *Ustilago maydis* (corn smut), *Fusarium graminearum* (*Gibberella zeae*), *Fusarium verticillioides* (*Gibberella moniliformis*), *F. proliferatum* (*G. fujikuroi* var. *intermedia*), *F. subglutinans* (*G. subglutinans*), *Diplodia maydis*, *Sporisorium holci-sorghii*, *Colletotrichum graminicola*, *Setosphaeria turcica*, *Aureobasidium zeae*, *Phytophthora*
 20 *infestans*, *Phytophthora sojae*, *Sclerotinia sclerotiorum*, *Pseudomonas avenae*, *Pseudomonas andropogonis*, *Erwinia stewartii*, *Pseudomonas syringae* pv. *syringae*, maize dwarf mosaic virus (MDMV), sugarcane mosaic virus (SCMV, formerly MDMV strain B), wheat streak mosaic virus (WSMV), maize chlorotic dwarf virus (MCDV), barley yellow dwarf virus (BYDV), banana bunchy top virus (BBTV), etc. See for example, U.S.
 25 Patent No. 8,395,023, which is incorporated herein by reference.

For example, several non-limiting examples of pests capable of destroying plants include but are not limited to northern corn rootworm (*Diabrotica barberi*), southern corn rootworm (*Diabrotica undecimpunctata*), Western corn rootworm (*Diabrotica virgifera*), corn root aphid (*Anuraphis maidiradicis*), black cutworm (*Agrotis ipsilon*), glassy
 30 cutworm (*Crymodes devastator*), dingy cutworm (*Feltia ducens*), claybacked cutworm (*Agrotis gladiaria*), wireworm (*Melanotus* spp., *Aeolus mellillus*), wheat wireworm (*Aeolus mancus*), sand wireworm (*Horistonotus uhlerii*), maize billbug (*Sphenophorus maidis*), timothy billbug (*Sphenophorus zeae*), bluegrass billbug (*Sphenophorus parvulus*),

southern corn billbug (*Sphenophorus callosus*), white grubs (*Phyllophaga* spp.), seedcorn maggot (*Delia platura*), grape colaspis (*Colaspis brunnea*), seedcorn beetle (*Stenolophus lecontei*), and slender seedcorn beetle (*Clivinia impressifrons*), as well as parasitic nematodes. *Id.*

5 For example, several non-limiting examples of target genes related to pests include but are not limited to major sperm protein, alpha tubulin, beta tubulin, vacuolar ATPase, glyceraldehyde-3-phosphate dehydrogenase, RNA polymerase II, chitin synthase, cytochromes, miRNAs, miRNA precursor molecules, miRNA promoters, etc. *Id.*

10 In an embodiment, modification of plant cells is conducted with one or more aptamers, which include DNA or RNA sequences that recognize and specifically bind to a particular ligand or molecule, often with high affinity. Aptamers have been utilized similarly to antibodies for binding specific antigens or receptors as well as for protein antagonists, as molecular escorts for delivery of an agent to a particular cell or tissue, as well as part of a riboswitch. Riboswitches include complex folded RNA sequences that
15 contain an aptamer domain for a specific ligand and can operate as “cis” or “trans” elements. Cis riboswitches may control gene expression by harnessing allosteric structural changes caused by ligand binding, whereas trans riboswitches control expression not operably linked to the riboswitch itself. For example, a riboswitch defaults to “on” position during gene expression, and the riboswitch is turned to the “off” position when an
20 increased concentration of the ligand is bound by the aptamer domain of the riboswitch and the riboswitch terminates transcription or translation of the gene under its control. Thus, in certain embodiments described herein, plant cells are modified and may result in stable transgenic plants as determined by the ligand being bound to an aptamer and the resulting expression or suppression of a target sequence.

25 Another class of RNA aptamers useful with an embodiment disclosed herein include “thermoswitches” which do not bind a ligand but that determine the aptamer’s conformation by temperature, so as to be thermally responsive.

 In an embodiment, modifying plant cells to reduce damage caused by disease or pest includes a transgene that transcribes to an RNA aptamer capable of binding to a
30 ligand that is at least part of a molecule endogenous to the pest or pathogen source of the disease or destruction. In this way, binding to the ligand reduces damage to the plant relative to a plant without the transgene. In an embodiment, the transgene includes at least partial sequence of a gene found in a pathogen or in the gut of a pest.

In an embodiment, a target sequence includes a gene native to the particular plant utilized in various embodiments disclosed herein, a transgene in a transgenic plant, a gene native to a pest or pathogen, etc. For example, plant cells modified as described herein may be modified to provide resistance to a particular disease or pest. In order to convey
5 such resistance, a gene native to the particular causal agent may be incorporated into the plant when modified as disclosed in various embodiments herein. Alternatively, in another example, multiple copies of a native gene to the plant may be incorporated, for example, to increase yield or nutritional content. A target sequence includes a sequence that expresses a specific gene or a sequence that suppresses a specific gene. Moreover, the
10 target sequence includes a translatable sequence (coding sequence) or a non-coding sequence (non-coding regulatory sequence), or both. The target sequence includes sequences from the plant itself, its species, or another species including any eukaryote or prokaryote, depending on the desired characteristic.

In an embodiment, regulatory RNA is utilized to modify plant cells as described
15 herein, and include ribozymes (self-cleaving ribozymes, hammerhead ribozymes, hairpin ribozymes, etc.) and may include sense or anti-sense segments capable of hybridizing to form an intramolecular double-stranded RNA.

In an embodiment, clustered regularly interspersed short palindromic repeats (CRISPR) is utilized for the one or more modifications of plant cells for Rapid-breeding of
20 plants as described. *See* van der Oost, Science, (2013) pp. 768-770, Vol.339; and Mali et al., Science (2013), pp. 823-826, Vol.339; and Cong et al., Science (2013), pp. 819-823, Vol.339; each of which is hereby incorporated by reference. For example, CRISPR utilizes RNA-guided DNA nuclease for highly specific gene targeting (e.g. for modification of genes). *Id.* For example, precise genome engineering may be conducted
25 based on the RNA-guided Cas9 nuclease, and can be utilized for multiple gene recognition sites. *Id.*

For example, in an embodiment CRISPR is utilized for targeting based on Watson-Crick complementarity, and in an embodiment CRISPR machinery can be reprogrammed to target a different DNA sequence through the use of a different crRNA. In an
30 embodiment, CRISPR is utilized for targeting a complementary 24-48 RNA sequence, with little to no mismatches.

As described herein, plant stem cells are utilized in various embodiments disclosed. However, various other plant cells may be utilized from the resulting plant

generated by methods disclosed herein, or in the process of generating the resulting plant. In this regard, plants may be crossed by modifying one gamete or nucleus and not the second gamete or nucleus utilized in sexual fertilization of the new progeny. In certain aspects, markers associated with one gamete, nucleus, or modification (e.g., gene
5 modification) are utilized with various embodiments disclosed herein. For example, a selectable marker may be linked to a DNA construct, a marker for herbicide tolerance may be tested by application of the herbicide to the progeny plant, or antibiotic resistance may serve as a marker for stable integration of a construct in the plant cell.

Transformation of DNA constructs into plant cells is likely to result in a percentage
10 of target plant cells that do not receive the construct or achieve expression of the construct. Common selective marker genes include those that confer resistance to antibiotics such as kanamycin and paromomycin, hygromycin B, spectinomycin, gentamycin, for example, or resistance to herbicides such as glufosinate, dicamba, or glyphosate. As with animal cells, markers such as green fluorescent protein or luciferase may be employed to indicate that
15 gene expression from the construct is occurring, or for example, a gene expressing beta-glucuronidase or uidA gene may be utilized, in conjunction with chromogenic substrates to visually screen modified plant cells.

As described herein, if plant cells are modified as part of various embodiments disclosed, subsequent to fertilization occurs and the resulting progeny plant seed is
20 obtained, the seed may be tested or optionally cultured to a plantlet stage and then tested. In the case where the fertilized seed is tested, in an embodiment a portion of the endosperm is removed for analysis without disrupting the embryo of the seed. In an embodiment, a portion of the embryo is analyzed without disrupting the viability of the embryo. In an embodiment, a small portion of the fertilized seed is analyzed without
25 disrupting the viability of the embryo.

As described herein, an optional step in various embodiments disclosed include culturing the progeny seed under conditions that allow for the seed to grow to multiple cells, to a plantlet stage, or to a mature plant. Plant seeds that have been modified, in certain embodiments, grow to plants that have an improved characteristic as compared
30 with a control plant. As described herein, modified seeds or plants with improved characteristics are selected by evaluating the seeds or plants at a microscopic, molecular, genotype, or phenotype level.

In other embodiments, the seed and or plant generated by various embodiments disclosed herein harbor naturally occurring improvements over control plants, and such improved seeds or plants are derived by the Rapid-breeding processes disclosed.

Improvements include, for example, higher seed quality, better water efficiency, better
5 nutrient efficiency, greater temperature tolerance, higher yield, increased seed protein, or increased seed oil, level of fermentable starch, increased metabolite, level of fatty acids, level of amino acids or proteins, as well as other characteristics described herein.

As described herein, the progeny of various embodiments disclosed herein may be analyzed by various means, including, for example, microscopic or molecular analysis. In
10 addition to the molecular or microscopic analysis of the progeny seeds or plants, in certain aspects where the seeds are grown to plantlet or mature plant stage, several screening assays may be utilized, depending on the characteristic desired to be analyzed. For example, plantlets or mature plants may be screened in greenhouses or field trial or research centers, and may include detecting changes in morphology of the plant,
15 physiological characteristics, biomass, or chemical composition. For example, chemical composition of the progeny seeds or plants may be analyzed by evaluating seed composition for protein, free amino acids, oil, free fatty acids, starch, tocopherols, and other components. For example, biomass may be measured by evaluating plant height, stem diameter, root and shoot weights, leaf or shoot lengths or diameters, etc., whereas
20 plant morphology may be evaluated by screening for days to pollen shed, days to silking, leaf extension rate, chlorophyll content, leaf temperature, stand, seedling vigor, internode length, plant height, leaf number, leaf area, tillering, brace roots, stay green, stalk lodging, root lodging, plant health, barrenness or prolificacy, green snap, pest resistance, kernels per row, number of rows of kernels on an ear, kernel abortion, kernel weight, kernel size,
25 kernel density, physical grain quality, bushiness, height, thicker or narrower leaves, striped leaves, knotted trait, chlorosis, albino, anthocyanin production, altered tassels, altered ears, altered roots, or similar characteristics. As described herein, in addition to molecular or microscopic/cellular analysis, seeds or plants can be analyzed by testing against stress conditions, such as water use efficiency, enhanced cold tolerance, shade tolerance,
30 herbicide resistance, etc., by growing plants/plantlets under the stress condition to be analyzed and evaluating the response against control plants/plantlets.

For example, marker-assisted selection (MAS) relies on quantitative trait loci (QTL) studies derived from the analysis of a few segregating populations and may be

employed for the analysis as described herein with regard to several claimed embodiments. Likewise, genomic selection relies on phenotyping and high-density genotyping of a sufficiently large and representative sample of the population, with the majority of loci regulating a quantitative trait in linkage disequilibrium with one or more molecular markers. *See*, for example, Resende, et al., *New Phytologist* (2012) 193: 617-624, which is incorporated herein by reference. In certain embodiments, the MAS or genomic selection is utilized on a very early plant (zygote, blastocyst, embryo, etc.), while in certain other embodiments the MAS or genomic selection is utilized at a plantlet or mature plant stage. For example, age and genotype x environment interaction affect expression of certain complex genetic characteristics, and need a more mature plant in order to accurately measure such characteristics. Likewise, for certain characteristics that are expressed in the early plant, simple molecular testing of particular markers at the early plant stage are sufficient for establishing presence or absence of the trait. As described herein, in certain plants the time-frame between the beginning of breeding and the production of improved seeds can span multiple decades. *Id.* Thus, in an embodiment, the techniques of MAS and genomic selection may be employed with certain embodiments disclosed herein for Rapid-breeding methods.

In an embodiment, one or more QTLs are linked with at least one of herbicide tolerance, disease resistance, insect or pest resistance, altered fatty acid, protein or carbohydrate metabolism, increased grain yield, increased oil, increased nutritional content, increased growth rates, enhanced stress tolerance, preferred maturity, enhanced organoleptic properties, altered morphological characteristics, increased consumer appeal, industrial use characteristics, or other characteristics as described herein. In an embodiment, one or more progeny are selected based on at least one characteristic indicative of a recurrent parent that assists in selection for marker-assisted backcrossing.

In an embodiment, additional transcription factors for use in various embodiments disclosed, or additional genes of interest associated with desired characteristics may be derived by techniques, such as differential screening using, for example, subtractive hybridization of a cDNA library, PCR, RT-PCR, Northern blot, Reverse Northern blot, Southern blot, Western blot, DNA fingerprinting, dot-blots, new generation sequencing, gene cloning or Random Primed RT-PCR, etc. readily assists in identification of genes or proteins of interest in eukaryotic cells, including plants.

For example, the genomes of many different plant varieties have been sequenced in full or in large part, thus enabling ready identification of particular genes or proteins as determined by differential screening related to various cell types, for example. In an embodiment, a differential screening for a particular plant is conducted with at least two or
5 more of a gametophyte, gamete, spore, or sporophyte. In an embodiment, a differential screening is conducted between a male gamete and a female gamete of the same species. In an embodiment, the differential screening is conducted between related but different species, or related but different cultivars, in order to more readily identify the genes or proteins more intricately involved in differentiation of different cell types. In an
10 embodiment, identification of a homolog is sufficient for utilization in differentiation of a cell type or development of a sufficient factor therefor.

For example, genomic sequence information for rice, maize, cucumber, oilseed, legumes, cocoa, eucalyptus, wild strawberry, apple, grape, tomato, potato, wheat, barley, banana, cannabis, grasses, and other crops and trees have been obtained in whole or in
15 large part. With new generation sequencing, the relevant sequence information of other plants is readily identifiable.

Animals

Various embodiments described herein are applicable to a number of animals,
20 including but not limited to domesticated or wild agricultural animals, companion animals, rodents or vermin, or other domesticated or wild animals including but not limited to cow, horse, goat, sheep, goat, llama, alpaca, pig, hog, boar, bison, yak, buffalo, worm, chicken, turkey, goose, duck, fish, crab, lobster, oyster, shrimp, mussels, other shell fish, donkey, camel, mule, oxen, dog, cat, mouse, rat, hamster, rabbit, chinchilla, guinea pig, gerbil,
25 ferret, elephant, bear, tiger, lion, dolphin, alligator, crocodile, whale, frog, toad, lizard, gecko, chameleon, raccoon, cougar, mountain lion, monkey, chimpanzee, gorilla, orangutan, ape, baboon, or other primate, giraffe, pigeon, pheasant, grouse, zebra, ostrich, bullock, water buffalo, carabao, snake, reindeer, carabiau, elk, insect, spider, antelope, deer, moose, pony, chiliquene, cormorant, parrot, parakeet, etc. or any hybrid thereof. In
30 an embodiment, one or more gametes are modified such that hybrids, including cross-species hybrids, are generated from the fertilization. In an embodiment, the animal includes one or more reptile, amphibian, mammal, fish, or bird.

Various embodiments disclosed herein are utilized to improve one or more characteristics of an animal by way of rapid selection of desirable naturally occurring characteristics through cycling of gametes. Various embodiments disclosed herein are utilized to improve one or more characteristics of a modified animal (e.g., genetic, or
5 epigenetic modification) and include rapid selection of desirable characteristics introduced into the animal by modification as described. For example, such characteristics include but are not limited to increased telomere(s) (e.g., length, quality, etc.), improved agricultural characteristic. For example, in an embodiment, the desired characteristic includes one or more of increased size, faster maturation of progeny; increased muscle
10 mass; increased lifespan; increased fertility (for example, increase in number and/or frequency of germ cell production or maturation including sperm or eggs); decreased gestation time; increased number of offspring over the animal's lifetime (for example, increase in the number of progeny any particular animal parents during its lifetime); decreased susceptibility to disease or death; increased quality of fur, hair, or feathers;
15 increased docility; increased quality of skin, hide, or feathers; increased ability to obtain energy or absorb nutrients from feed; alteration in microbiome of one or more of gut, skin, or other body part (e.g., for reduced contamination of food supplies with dangerous bacteria, prions, viruses, fungi, worms or other parasites or microorganisms); increased tolerance to antibiotics or other anti-microbial agents; increased milk production;
20 increased fatty acid production; increased protein production; alteration in type of fatty acid production; increased immunological compatibility of tissues or organs with the human counterpart tissue or organ (e.g., for use in cross-species organ transplants particularly into humans); or increased adaptability to environment. For example, a published genomic listing is utilized that identifies all presently identified cell surface
25 antigens on human cells. *See*, for example, Cunha, et al. PNAS, 106: 16752-16757 (2009), which is incorporated herein by reference. In an embodiment, utilizing this genetic sequence listing of cell surface antigens, one or more can be manipulated such that organs or tissues can be generated that are tolerogenic when transplanted into humans.

In an embodiment, analyzing the resulting progeny includes analyzing for an
30 improvement in a particular characteristic, whether such characteristic is attributed to a naturally occurring trait or is the result of an additional assisted modification (e.g., physical, chemical, genetic, epigenetic intervention, etc.). In an embodiment an improvement in a particular characteristic includes an increased level of a particular

characteristic in the progeny, relative to a control animal (in certain instances, the control animal may be a parent animal). For example, an increased level of a particular characteristic may be measured as described herein, including increased muscle mass or milk production.

5 In an embodiment, analyzing the resulting progeny includes evaluating one or more characteristics including a genetic marker, a single nucleotide polymorphism, a simple sequence repeat, a restriction fragment length polymorphism, a haplotype, a tag single nucleotide polymorphism, a gene, an allele of a genetic marker, a DNA-derived sequence, an RNA-derived sequence, promoter, 5'-untranslated region of a gene, 3'-
10 untranslated region of a gene, microRNA, siRNA, a QTL, satellite marker, telomere length, characterizing DNA from the cell nucleus, characterizing DNA from the cell mitochondria, transgene, mRNA, high-resolution chromosomal structure and content analyses with microarrays, double stranded mRNA, transcriptional profile, or methylation pattern.

15 In an embodiment, an improvement in a particular characteristic includes a decreased level of a particular characteristic in the progeny, relative to a control animal (again, in certain instances, the control animal is the parent animal). For example, a decreased level of a particular characteristic or an increased level of a particular characteristic may be measured as described herein, including for example, increased
20 muscle mass.

The various characteristics described may be analyzed for improvement in several ways. For example, chemistry, cell biology, or molecular biology techniques may be utilized for measuring telomere changes, ploidy, or gene, protein, or mRNA changes, epigenetic changes, etc. As another example, cell surface antigens may be evaluated by
25 flow cytometry, blotting, other cell staining or microscopic analysis. In an embodiment, the desired tissues are selected and those that are not of the desired phenotype or genotype are removed from the desired animals or animal tissues. In an embodiment, the animal tissue is analyzed for one or more alleles of at least one locus.

As another example, an immunoreactive antibody may be utilized to detect the
30 presence or absence of a protein that is expressed or suppressed in the progeny animal or animal tissue.

Other changes, including improvements, in the various characteristics described may be measured by physical means, for example, animal weight, animal height, volume

of an animal product (e.g. milk, fur, hair, feathers, meat, hide, skin, hoof, etc.). As another example, a small portion of tissue is removed from the fertilized embryo without disturbing the viability of the embryo(s), and such tissue is analyzed by means described herein.

5 In an example, a larger portion of tissue is removed or swapped from one fertilized embryo to another fertilized embryo, or from an adult animal to a fertilized embryo. For example, imaginal disc tissue, blood island tissue, neural crest tissue, or other selected animal tissue can be transplanted from one animal to another or even one species to another.

10 It is useful to determine zygosity of one or more loci during breeding in order to evaluate the degree of fixation, or inbreeding, segregation distortion (i.e., in transgenic germplasm, maternal inheritance testing of for loci that affect gametes).

 For example, determination of allelic frequencies assists in determining parental linkages for a specific marker, and comparing allele frequency data between two or more
15 germplasm pools provides guidance with selection of targeted desired characteristics in such a way that as alleles increasing in frequency in conjunction with a shift in distribution of one or more characteristics, such is presumed to be linked to the trait of interest. Furthermore, determining relative allele frequency data assists in construction of genetic linkage maps. For example, several species of animals have had full or partial genomic
20 sequence analysis conducted, as described herein. Thus, genetic sequences for particular animals, or the homologs from other species, are readily available or attainable for use with various embodiments described herein.

 In certain embodiments, selected progeny are bulked, depending on the analysis of the resulting genotype. Conversely, if multiple characteristics are analyzed by way of
25 QTL or genomic selection, for example, with varying effects being selected for a particular population, the progeny may keep individual identity preserved, and culture the progeny for determination of various combinations of the desired QTL.

 In an embodiment, cycling of gametes *in vitro*, as described herein, accelerates the introgression of a specific gene or genes into an animal and allows for greater precision by
30 more cycles of recombination.

 Likewise, genetic recombination can involve “gene conversion” or the non-reciprocal transfer of DNA from one genotype to another. Utilizing various embodiments disclosed herein, vast populations for rare zygotes with gene conversion at a target locus

may be screened to identify a true isogenic derivative rather than a nearly-isogenic backcross derivative in which flanking alleles from the donor may confer undesirable characteristics. Gene conversion may be especially useful for manipulating exotic germplasm in which a desirable gene is often surrounded by undesirable ones and thus
5 methods disclosed herein assist in avoiding intermediate phenotypes.

Sexual breeding results in the addition and transfer of chromosomes, which allows for increased diversity in the progeny, and allows for selection of particular desirable characteristics. In an embodiment, a method described herein is utilized to preserve an endangered species, or possibly revive an extinct species through use of DNA obtained
10 from preserved fragments or portions of an animal or animal tissue. In an embodiment, the DNA is deposited in a cell that is used as a surrogate host for replication of the cellular DNA. In an embodiment, a hybrid of the endangered or extinct species may be utilized in order to generate progeny animal or animal cells that are further utilized to generate the desired phenotypic progeny. For example, DNA from mastodon and woolly mammoth
15 has been extracted and sequenced from tissue samples. (*See for example, Rohland, et al., PLOS Biology, Dec. 2010, Vol. 8, No. 12, which is incorporated herein by reference.*) In an embodiment, the DNA of a modern day elephant is modified for a desired characteristic, for example of a desired characteristic of a woolly mammoth or mastodon, based on the genetic sequence, and progeny are selected based on the desired
20 characteristic.

In an embodiment, the progeny are analyzed for example, by statistical sample testing of animal tissue(s) (including embryos) for further use. In an embodiment, a portion of the animal tissue(s) are utilized while maintaining the viability of the progeny. For example, a portion of the zygote, embryo, blastocyst, or other tissue portion may be
25 obtained by mechanical or chemical means and utilized for analysis.

In an embodiment, the one or more desired characteristics are quantified based on the analyzed animal tissue. In an embodiment, the one or more quantified desired characteristics are compared with control animal tissue samples or known germplasm pools in order to identify frequency shifts. Likewise, DNA and RNA may be extracted
30 from animal tissue(s) sufficient to yield DNA or RNA adequate for PCR, RT-PCR, TaqMan assays, sequencing, arrays or blots, or other amplification.

Several criteria may be established in order to determine whether additional breeding is desired. For example, one or more cells from an embryo (e.g., blastocyst) is

removed for genetic analysis of a desired characteristic, whether the corresponding gene(s) are naturally occurring in the embryo or artificially inserted, deleted, or mutated.

As described herein, in certain aspects a method disclosed includes molecularly modifying an animal cell. In an embodiment, a DNA construct is utilized to modify an animal cell. Assembly of DNA constructs may be done using standard methods, typically with a promoter operably coupled to the DNA, and optionally other regulatory elements such as 5' leaders, introns, 3' untranslated regions (such as polyadenylation signals), signal peptides, or repressor or enhancer sequences. In an embodiment, at least one epigenetic modification (e.g., DNA methylation) is conducted. In an embodiment, at least one telomere modification or addition to the cell is conducted.

For example, promoter elements include non-constitutive promoters (such as spatially specific promoters, temporally specific promoters, inducible promoters, etc.), or constitutive promoters. For example, spatially specific promoters include organelle, cell, tissue, or organ-specific promoters functional in an animal (such as involucrin or K-14 for skin/epidermis, SM-22 for arterial tissue, mucin-2 for intestinal tissue, B29 promoter for B cells, CD14 promoter for monocytes, CD43 promoter for leukocytes and platelets, CD45 promoter for hematopoietic cells, CD68 promoter for macrophages, desmin promoter for muscle, elastase-1 promoter for pancreatic acinar cells, endoglin promoter for endothelial cells, fibronectin promoter for differentiating tissues, flt-1 promoter for endothelial cells, GFAP promoter for astrocytes, GP11 promoter for megakaryocytes, ICAM-2 promoter for endothelial cells, INF-beta promoter for hematopoietic cells, Mb promoter for muscle, Nphs1 promoter for podocytes, OG-2 promoter for osteoblasts, SP-B promoter for lung, SYN1 promoter for neurons, WASP promoter for hematopoietic cells, etc. *See* for example, Zhai et al., J. Animal Sci. and Biotech. 2012, 3:32; Lin et al., Abstract Br. J. Dermatol., 144(1):34-9, 2001; each of which is incorporated herein by reference. In an embodiment, an embryo-specific promoter is utilized with an embodiment. For example, temporally specific promoters can include promoters that favor promoter expression during certain developmental stages in an embryo's growth or reproductive cycle, or during different stages of the animal's life.

Inducible promoters include promoters induced by chemicals such as exogenous or synthetic chemicals as well as endogenous transcription factors and other signaling molecules, and several standard inducible promoters can be adapted for use in various embodiments described herein. An expression-specific promoter also includes generally

constitutively expressed promoters that vary on the degree of expression. For example, cumate switches, thymidine kinase, antibiotics, and other promoters inducible by a compound have been developed and can be adapted for use in the various embodiments described herein.

5 In addition to promoter elements, in certain instances terminator elements may be utilized in the embodiments disclosed herein, including for example, polyadenylation signals as described herein. Additionally, spacer DNA segments (found, for example, between parts of a gene suppression element or between different gene suppression elements) may include translatable DNA encoding a target gene, translatable DNA
10 encoding a marker or reporter gene, transcribable DNA derived from an intron, transcribable DNA encoding RNA that forms a structure such as a loop, stem, or aptamer capable of binding to a specific ligand, spliceable DNA including introns and self-splicing ribozymes, transcribable DNA encoding a target gene detectable by molecular biology techniques.

15 Numerous other promoters and enhancers that function in animal cells (whether plant cell or another cell was the original source) may be utilized in assembling constructs for use in modifying animal cells as described herein.

 In various aspects, one or more genes or other modifications may be implemented with various embodiments disclosed herein, by for example, making multiple
20 modifications to a single cell or by making single mutations in a cell and crossing those cells to result in a transformed animal. Such modifications may be stacked into the resulting progeny animal with various modifications. For example, characteristics for increasing muscle mass may be stacked with characteristics for increasing milk production, or tolerating antibiotics.

25 With regard to gene suppression, various modes may be used, for example, including anti-sense, co-suppression, and RNA interference (miRNA, dsRNA, siRNA, and other forms of double stranded RNA that block translation). Several microRNA genes have been characterized in animals, including promoters for the same, and can be adapted for use with animals. Transposable elements can also be utilized to suppress gene function.
30 In certain instances, mutations by insertion of one or more transposable elements disrupts protein translation from a particular gene.

 Modification of animal cells by transformation with molecular constructs may be used with various embodiments described herein. Culturing conditions for transforming

animal cells includes, for example, nutrient media, kept at conditions which facilitate transformation of the cells. Animal cell targets for transformation include at least one of gametes (sperm, egg, etc.), or immature embryos. Since typically not all cells are transformed when modified in this manner, one or more markers (e.g., marker genes) are generally utilized to provide an efficient system for identification of the cells that are stably transformed with DNA constructs. Further, if conferring resistance to an antibiotic, for example, transformation may be screened by application of the antibiotic to the animal (or animal cells).

In an embodiment, one or more gametes are induced from pluripotent stem cells, or primordial germ cells. For example, cell differentiation of gametes in vitro can be adapted for use with various embodiments disclosed herein. See for example, Zhu et al., Abstract Asian J. Androl. 2012 14(4):574-9; Imamura et al., Abstract Mol. Reprod. Dev. 2010 Sep:77(9):802-11; Equizabal, et al., Abstract Stem Cells 2011 Aug 29(8):1186-95, each of which is incorporated herein by reference.

For example, several non-limiting examples of pathogens include but are not limited to fungal, bacterial, or viral pathogens. For example, pathogens include but are not limited to Streptococcus, Escherichia coli, Salmonella, Vibrio, Streptococcus, Spirillum, Shigella, Mycoplasma, yeast, or other pathogens.

In an embodiment, modification of the microbiome includes modifying one or more regions of the animal such that the region(s) encourage growth or sustenance of one or more non-pathogenic microorganisms including but not limited to lactobacillus, bacillus, bifidobacterium, or other non-pathogenic microorganisms.

In an embodiment, modification of animal cells is conducted with one or more aptamers, which include DNA or RNA sequences that recognize and specifically bind to a particular ligand or molecule, often with high affinity. Aptamers have been utilized similarly to antibodies for binding specific antigens or receptors as well as for protein antagonists, as molecular escorts for delivery of an agent to a particular cell or tissue, as well as part of a riboswitch. Riboswitches include complex folded RNA sequences that contain an aptamer domain for a specific ligand and can operate as “cis” or “trans” elements. Cis riboswitches may control gene expression by harnessing allosteric structural changes caused by ligand binding, whereas trans riboswitches control expression not operably linked to the riboswitch itself. For example, a riboswitch defaults to “on” position during gene expression, and the riboswitch is turned to the “off” position when an

increased concentration of the ligand is bound by the aptamer domain of the riboswitch and the riboswitch terminates transcription or translation of the gene under its control. Thus, in certain embodiments described herein, animal cells are modified and may result in stable transgenic animals or animal cells as determined by the ligand being bound to an aptamer and the resulting expression or suppression of a target sequence.

Another class of RNA aptamers useful with an embodiment disclosed herein include “thermoswitches” which do not bind a ligand but that determine the aptamer’s conformation by temperature, so as to be thermally responsive.

In an embodiment, regulatory RNA is utilized to modify animal cells as described herein, and include ribozymes (self-cleaving ribozymes, hammerhead ribozymes, hairpin ribozymes, etc.) and may include sense or anti-sense segments capable of hybridizing to form an intramolecular double-stranded RNA.

In an embodiment, clustered regularly interspersed short palindromic repeats (CRISPR) is utilized for the one or more modifications of animal cells for Rapid-breeding of animals as described. *See* van der Oost, Science, (2013) pp. 768-770, Vol.339; and Mali et al., Science (2013), pp. 823-826, Vol.339; and Cong et al., Science (2013), pp. 819-823, Vol.339; each of which is hereby incorporated by reference. For example, CRISPR utilizes RNA-guided DNA nuclease for highly specific gene targeting (e.g. for modification of genes). *Id.* For example, precise genome engineering may be conducted based on the RNA-guided Cas9 nuclease, and can be utilized for multiple gene recognition sites. *Id.*

For example, in an embodiment CRISPR is utilized for targeting based on Watson-Crick complementarity, and in an embodiment CRISPR machinery can be reprogrammed to target a different DNA sequence through the use of a different crRNA. In an embodiment, CRISPR is utilized for targeting a complementary 24-48 RNA sequence, with little to no mismatches.

As described herein, animal stem cells are utilized in various embodiments disclosed. However, various other animal cells may be utilized from the resulting animal generated by methods disclosed herein, or in the process of generating the resulting animal. In this regard, animals may be crossed by modifying one gamete or nucleus and not the second gamete or nucleus utilized in sexual fertilization of the new progeny. In certain aspects, markers associated with one gamete, nucleus, or modification (e.g., gene modification) are utilized with various embodiments disclosed herein. For example, a

selectable marker may be linked to a DNA construct, a marker for antibiotic resistance may be tested by application of the antibiotic to the progeny animal.

Transformation of DNA constructs into animal cells is likely to result in a percentage of target animal cells that do not receive the construct or achieve expression of the construct. Common selective marker genes include those that confer resistance to antibiotics such as kanamycin and paromomycin, hygromycin B, spectinomycin, gentamycin, for example. In addition, colorimetric markers such as green fluorescent protein or luciferase may be employed to indicate that gene expression from the construct is occurring.

As described herein, if animal cells are modified as part of various embodiments disclosed, subsequent to fertilization occurs and the resulting progeny animal embryo is obtained, the embryo may be tested or optionally cultured to a later stage and then tested. In the case where the fertilized embryo is tested, in an embodiment a portion of the embryo is removed for analysis without disrupting the viability of the embryo itself.

As described herein, an optional step in various embodiments disclosed include culturing the progeny embryo under conditions that allow for the embryo to grow to multiple cells, to a later developmental stage, or (subsequent to implantation) to a newborn animal progeny. Animal cells that have been modified, in certain embodiments, grow to animals that have an improved characteristic as compared with a control animal. As described herein, modified animals with improved characteristics are selected by evaluating the embryos or animals at a microscopic, molecular, genotype, or phenotype level.

In other embodiments, the embryo and animal generated by various embodiments disclosed herein harbor naturally occurring improvements over control animals, and such improved animals are derived by the Rapid-breeding processes disclosed. Improvements include, for example, higher tolerogenicity for human organs, better efficiency of nutrient or energy use, increased level of fatty acids, increased level of amino acids or proteins, as well as other characteristics described herein.

As described herein, the progeny of various embodiments disclosed herein may be analyzed by various means, including, for example, microscopic or molecular analysis. In addition to the molecular or microscopic analysis of the progeny embryos or animal(s) or animal cell(s), in certain aspects where the embryos are grown to a later developmental stage, several screening assays may be utilized, depending on the characteristic desired to

be analyzed. For example, marker-assisted selection (MAS) relies on quantitative trait loci (QTL) studies derived from the analysis of a few segregating populations and may be employed for the analysis as described herein with regard to several claimed embodiments.

5 Likewise, genomic selection relies on phenotyping and high-density genotyping of a sufficiently large and representative sample of the population, with the majority of loci regulating a quantitative trait in linkage disequilibrium with one or more molecular markers. In certain embodiments, the MAS or genomic selection is utilized on a very early animal (zygote, blastocyst, embryo, etc.), while in certain other embodiments the
10 MAS or genomic selection is utilized with a mature animal or newly born animal stage. In addition, maternal imprinting may be considered, measured, and manipulated by, for example, modifying methylation of one or more cells, as described herein. Thus, in an embodiment, the techniques of MAS and genomic selection may be employed with certain embodiments disclosed herein for Rapid-breeding methods.

15 In an embodiment, one or more QTLs are linked with at least one of antibiotic resistance, altered fatty acid, protein or carbohydrate metabolism, increased size (e.g., height, weight, fat composition, muscle mass, etc.), increased tolerogenicity between human and animal organs or tissues, or other characteristics as described herein. In an embodiment, one or more progeny are selected based on at least one characteristic
20 indicative of a recurrent parent that assists in selection for marker-assisted backcrossing.

 In an embodiment, additional transcription factors for use in various embodiments disclosed, or additional genes of interest associated with desired characteristics may be derived by techniques, such as differential screening using, for example, subtractive hybridization of a cDNA library, PCR, RT-PCR, Northern blot, Reverse Northern blot,
25 Southern blot, Western blot, DNA fingerprinting, dot-blot, new generation sequencing, gene cloning or Random Primed RT-PCR, etc. readily assists in identification of genes or proteins of interest in eukaryotic cells, including animal cells.

 For example, the genomes of many different animals or animal breeds have been sequenced in full or in large part, thus enabling ready identification of particular genes or
30 proteins as determined by differential screening related to various cell types, for example. In an embodiment, a differential screening for a particular animal is conducted with at least two or more of a gamete, or stem cell. In an embodiment, a differential screening is conducted between a male gamete and a female gamete of the same species. In an

embodiment, the differential screening is conducted between related but different species, or related but different animals or animal breeds, in order to more readily identify the genes or proteins more intricately involved in differentiation of different cell types. In an embodiment, identification of a homolog is sufficient for utilization in differentiation of a cell type or development of a sufficient factor therefor.

For example, genomic sequence information for many animals has been determined, including porifera (including sponges), ctenophore, placozoa, cnidarian, deuterostomia (including echinoderms, hemichordates, urochordates, cephalochordates, cyclostomes, cartilaginous fish, bony fish, amphibians, reptiles, birds, and mammals), protostomia (including insects, coleopteran, diptera, hemiptera, hymenoptera, Lepidoptera, phthiraptera), crustaceans, Chelicerates, molluscs, Platyhelminthes, nematodes, and annelids, etc. in whole or in large part. With new generation sequencing, the relevant sequence information of other animals is readily identifiable.

It is highly beneficial to evolve animal breeding to include characteristics that are desirable without the time-scales usually required for growing animals to maturity for sexual reproduction. Described herein include various embodiments related to a focused means for selecting and breeding desired animals or animal breeds with pre-determined or naturally resulting desired characteristics. Various embodiments disclosed herein provide a higher rate of improvement of desired characteristics, as described herein.

In this way, certain embodiments disclosed herein realize multi-generation sequences of gametes by the classic sexual reproductive route, without intervening phenotypes with their attendant complexities and latencies. Furthermore, various embodiments disclosed herein may be utilized to derive new animals or animal breeds particularly amenable to certain environmental conditions, or needs for obtaining animal products. Additionally, the disclosed embodiments herein allow for a reduction of labor and field resources for animal breeding, increased capacity to rapidly breed a large number of animals in a short time period, as well as an increased capacity to analyze a larger number of animals prior to breeding for desirable characteristics.

Various embodiments disclosed herein relate to genotype-to-genotype generational stepping without an intervening phenotype but instead result from generating male and female gametes directly from animal stem cells. For example, first generation stem cells are induced to derive gametes, which are bred. The resulting fertilized embryo includes second generation stem cells may be obtained, if additional breeding is desired. This

multi-generational cycling of gametes *in vitro* (CoGiV) allows for allele valuation and selection for rapid and efficient directed evolution of the particular animal species toward one or more desired characteristics. For example, for specific agricultural animals it is desirable to increase the number of progeny and decrease the amount of time needed to
5 screen and select the particular progeny. In another example, it is desirable to increase the precision of breeding-generated phenotypes or the breadth of accessible genetic variation. As described herein, multiple cycles of breeding and/or selecting are conducted depending on genetic complexity, and desired characteristics.

As disclosed herein, in an embodiment described are an increase in efficiency of
10 Rapid-breeding, and do not require genetic or other modification of the animal genome. However, in an embodiment, modification of animal cells (genetic, epigenetic, transcriptional or translational, etc.) is utilized with the Rapid-breeding process.

At any number of steps, standard breeding techniques may be implemented in conjunction with the steps of the Rapid-breeding methods and systems disclosed herein.
15 For example, in conjunction with the steps of Rapid-breeding of animals, conventional animal crossing may be conducted for selecting starting material (e.g., animal stem cells, primordial germ cells, etc.), or once a first, second, third, etc. generation of progeny have been developed. Standard animal crossing includes, for example, obtaining sperm (or conversely eggs and in an embodiment each are removed from the parents) of one parent
20 (may be modified, such as transgenic, or not modified) and optionally the other parent and growing resulting offspring into second generation mature parent animals. One advantage of various embodiments described herein relates to the ability to cycle through gametes from somatic cells, stem cells, or primordial germ cells and produce progeny that are screened and either utilized for further breeding while still *in vitro* or implanted into a host
25 (mother) animal for further growth and development.

In an embodiment, a method for animal breeding includes selecting one or more animal stem cells; inducing at least one of the animal stem cells to differentiate into at least one first gamete; combining the at least one first gamete with at least one second gamete (optionally of opposite gender) for fertilization; and molecularly or
30 microscopically analyzing the progeny resulting from the fertilization prior to implantation.

In an embodiment, the animal stem cell includes a cell isolated from the early stage embryo (e.g., blastocyst), or the cell is induced from a somatic cell of the animal. Such

induction has been conducted and can be adapted for use with the various embodiments herein. As described herein with regard to various embodiments, animal stem cells are totipotent cells that facilitate growth and production of tissues. Animal stem cells have the ability to self-renew, as well as differentiate into the various different cell types of the animal.

In an embodiment, a method includes selecting one or more animal stem cells; inducing at least one of the animal stem cells to differentiate into at least one first gamete; combining the at least one first gamete with at least one second gamete (optionally of opposite gender) for fertilization; and molecularly or microscopically analyzing the embryo resulting from the fertilization.

In an embodiment, microfluidics devices or systems may be utilized for selecting animal stem cells (e.g., by markers) or selecting progeny generated by various embodiments disclosed herein.

In an embodiment, a method includes selecting one or more animal stem cells; inducing at least one of the stem cells to differentiate into at least one first gamete of first genetic composition; combining the nucleus of the at least one first gamete with the nucleus of the at least one second gamete of second genetic composition for fertilization; and molecularly or microscopically analyzing the embryo resulting from the fertilization. In this particular embodiment, the gametes may be from the same or opposite gender. In an embodiment, the gamete may be from different genetic composition if diversity is desired, or the same genetic composition if the “ideal” or goal genetic composition has already been determined or obtained, or there are particular characteristics desired to be preserved and passed on to the progeny. In an embodiment, the progeny is/are generated by combining the containing chromosomes that yield a cell having the ability to sustain itself long enough to be utilized in an additional round of breeding, and/or to go on to develop into a live birthed progeny or even to a sexually mature adult animal.

In an embodiment, inducing gametes includes inducing animal stem cells first to differentiate to a primordial germ cell stage. In an embodiment, gametes are derived from primordial germ cells without first being induced from stem cells. In an embodiment, inducing gametes includes inducing animal stem cells first to differentiate to any stage along the germ cell to gamete maturation pathway.

In an embodiment, the gametes and/or stem cells are genetically analyzed. Genetic analysis for embodiments described herein includes genetic sequence analysis, ploidy

analysis, or marker selection, for example. In an embodiment, mRNA sequence information is obtained at one or more steps in the rapid-breeding process, for one or more cell types (e.g., gametes, etc.). In an embodiment, marker assisted selection is utilized to screen genetic markers in the very early progeny (zygote, blastocyst, embryo, etc. stage).

5 In mammals, the primordial germ cells arise in the embryo from the proximal epiblast, a region near the first blood lineages in the yolk sac. Several transcription factors are known to function in the differentiation of germ cells and gametes and can be adapted for use in various embodiments disclosed herein. For example, TFAP2C, Blimp1/Prdm1, Prdm14, LIF, BMP4, FGF2, KL, and others. *See* for example, De Felici et al., Abstract
10 Curr. Stem Cell Res. Ther. 2009 May (2):87-97, which is incorporated herein by reference.

In an embodiment, selecting one or more animal stem cells includes choosing at least one animal stem cell tissue containing animal stem cells based at least one criteria. In an embodiment, selecting one or more animal stem cells includes choosing from at least
15 one animal stem cell tissue, one or more single cells isolated from the remaining tissue. In an embodiment, one or more single animal cells may be isolated from the animal stem cell tissue, for example, by physical or chemical means (e.g., dissection, flow cytometry, enzymatic extraction, lysis of tissue, etc.).

In an embodiment, inducing at least one of the animal stem cells to differentiate
20 into at least one first gamete includes inducing to differentiate to at least one of an ovum (egg), oocyte, ootid, oogonium, spermatozoa, spermatid, or spermatocyte.

In an embodiment, a plant somatic cell is inverse-differentiated into a germ cell precursor and forward-specialized into pre-gametes of either sex and matured in appropriate cellular milieu into fertilization-ready gametes. In an embodiment, stem cells
25 are utilized in the process, rather than inverse-differentiated somatic cells. Post-fertilization, in an embodiment the zygote resulting from fertilization is analyzed directly, or cultured to at least a blastula stage, at which stage multiple cells are provided for next-generation pre-gametes. Following maturation into gametes of both sexes, each of these may be employed (in potentially quite wide) crossings aimed at realization of specific
30 allelic traits and/or the exclusion of others, as determined in near-real time by endosperm and/or polar body analysis. As described herein, such Rapid-breeding and analyses may be done at each generation step, and in parallel and/or with highly-automated means that allows for rapid trait selection at each generational step. In an embodiment, the

germination viability of the analyzed animal or animal tissue (such as a portion of an embryo or in its entirety) is maintained.

For any given round of breeding, the method includes selecting naturally occurring characteristics that emerge in the crossing of the parent cells. Multiple rounds of such a selection process without the necessity of culturing the resulting offspring yields a rapid-breeding approach that reduces the time needed for natural selection of desirable characteristics to a fraction of that required for traditional animal breeding that allows the offspring to grow into viable progeny or sexually mature adult animals.

For any given round of breeding, one or more genes may be inserted, deleted, or mutated in order to produce a resulting progeny bearing the gene or stacks of genes of interest. In an embodiment, for example, the animal stem cells are modified prior to induction of gametes. In another embodiment, the gametes are modified prior to fertilization. In an embodiment, the animal stem cells are modified transgenically by one or more gene insertions, deletions, or mutations. In an embodiment, the animal stem cells are modified epigenetically by altering gene expression through changes in methylation, for example. For example, CpG islands are associated with at least half of all cellular genes and are normally without methylation. *See* Baylin and Bestor, Abstract Cancer Cell 2002, 1(4) pp. 299-305, which is incorporated herein by reference. Methylation of the CpG islands is associated with tumorigenesis. *Id.* Dense methylation of cytosine residues within islands causes strong and heritable transcriptional silencing. *Id.* The silencing usually occurs with genes subject to genomic imprinting or X chromosome inactivation. *Id.*

In an embodiment, the one or more modifications of animal cells described herein are reversible. In an embodiment, the one or more modifications occur in DNA of the animal cell nucleus. In an embodiment, the one or more modifications occur in DNA of the mitochondria of the animal cells. In an embodiment, the animal stem cells or gametes are modified by the addition, deletion, or mutation of one or more mRNA molecules or proteins utilized in cell transcription or translation, respectively. In an embodiment, the genetic or epigenetic modification of the animal cells yields transformed progeny animals with a genome that has been altered by the stable integration of a recombinant DNA.

Figures

As illustrated in Figure 1, an embodiment includes a method 100 for Rapid-breeding of plants. As illustrated, cells (stem cells, gametophytes, sporophytes, somatic cells, etc.) are extracted 105 from Parent 1 and Parent 2 and induced to differentiate *in vitro* tissue culture 110, deriving gametes or transferable nuclei 115 that when combined 120, give rise to one or more zygote progeny 185. Next, the progeny are analyzed 130 as described herein utilizing cell, molecular, or microscopic methods, for example, or optionally cultured 125 to give rise to more than one cell. As illustrated, if the progeny are optionally cultured 125, they may be subsequently analyzed 130 for example, by removing a small portion of plant tissue or clonal cells to evaluate the genotype as described herein.

Following analysis of the progeny 130, the progeny are selected 135 based on the analysis. Optionally, the selected progeny may be cultured 125, or cultivated 140 to plantlet stage or optionally to mature plant stage 150. Alternatively, following selection 135 of the progeny based on the analysis, the progeny cells (which include multipotent, pluripotent, and/or totipotent cells) are utilized to repeat 195 the Rapid-breeding process. Optionally, cultured plant cells 125, cultivated plantlets 140 or cultivated mature plants 150 may be optionally validated through a second analysis 145 following the growth in culture or through cultivation. Cells may be obtained from progeny at any step following analysis for one or more desired characteristics, as illustrated in Figure 1. For example, in an embodiment, culturing plant cells or plant progeny includes growing them in a tissue culture plate. In an embodiment, cultivating a plant includes growing the progeny or other plants in a greenhouse, field, test plot, or small plant containers located indoors or outdoors.

As illustrated in Figure 2, in an embodiment, a portion of the plant tissue (such as a seed) is removed for analysis as described herein. In an embodiment, a portion of the endosperm is removed for analysis, leaving the embryo intact and maintaining the germination viability of the seed. In an embodiment, the endosperm and embryo are derived from sperm that share the same genotype, thus eliminating possible variation in inheritance from the sperm fertilizing the polar body that gives rise to the endosperm and the sperm fertilizing the egg that gives rise to the embryo.

As illustrated in Figure 3, in an embodiment, the progeny generated from the rapid-breeding disclosed herein are analyzed genetically (including genotypically, for example) or epigenetically, as described herein, and further analyzed phenotypically in tissue culture plates, or as plantlets/plants in a greenhouse or field. As illustrated in Figure 3, one or

more cameras 300 are utilized to survey a plot, a row, or a specific plant/dish, while a location marker (e.g., global positioning system indicator, etc.) 370 is placed at various positions around or near the growing plants/plant cells. As described herein, the cameras include one or more of infrared cameras, visual cameras, x-ray cameras, or similar.

5 Information detected from these or other sensors are transmitted (optionally wirelessly) to a central database, as indicated by “1,” “2,” or “3,” and the data collected 397 is evaluated by a user that is a person 395, or computer 380, for example. The user evaluates the data and optionally enters additional observed data via an input/output device 390. Based on the data evaluation, a Desired Characteristic Value from one or more samples is assigned,
10 and any experimental progeny (310, 330, 350, etc.) that include desired characteristics may be assessed further. Likewise, any experimental progeny that lack the desired characteristics, or that bear a phenotype no better than the control progeny (320, 340, 360, etc.) may be culled from further research. In an embodiment, a locator indicator 398 is utilized to locate a particular progeny for further assessment or for culling based at least in
15 part on the location marker 370 in the field or greenhouse. In an embodiment, for example, the location marker 370 includes a transmitter, and the location indicator 398 includes a receiver. In an embodiment, for example, the location marker 370 includes a receiver, and the location indicator 398 includes a transmitter. In an embodiment, one or more components described in Figure 7 are implemented in Figure 3. In an embodiment,
20 an integrated sensor-to-database collection may be utilized, such as a wireless sensor network, for example that integrates collected information, optionally encrypts information shared wirelessly and optionally through an internet interface. *See Bencini et al., intechopen, pp. 1-18, which is incorporated herein by reference.* In an embodiment, the information contributes to and/or compares the information with a database. In an
25 embodiment, the database includes genomic sequence information for the particular organism.

As illustrated in Figure 4, included in the analysis of the progeny of the rapid-breeding disclosed herein, the phenotype may be assessed by utilizing one or more sensors, for example, that contact the plant or plant cells, or are in the vicinity of the plant
30 or plant cells. In an embodiment, the one or more sensors include at least one of a receiver or transmitter. In an example, the sensors detect particular environmental conditions or plant/plant cell conditions and transmit the information to a user (as shown in Figure 3) where the information is collected and/or evaluated. In an embodiment, as described in

Figure 3, one or more progeny are culled or further assessed depending on the information collected or evaluated from the one or more sensors.

For example, a collar 400 includes a flexible band with a force sensor or strain sensor incorporated that senses stalk diameter may be utilized for sensing plant growth. In an embodiment, the collar 400, is in a fixed state and erupts or breaks apart as the plant reaches a threshold size, wherein the breaking of the collar sends a signal. In an embodiment, a sensor in the form of a patch 410 is attached to a leaf and is capable of measuring leaf morphology or growth. In an embodiment, a capture strip 420A or B is utilized for detecting emitted plant chemicals, such as carbon dioxide, oxygen, pheromones, stress hormones, reactive oxygen species, and others. In an embodiment, the capture strip 420A is located on the plant or plant cells. In an embodiment, the capture strip is located in the vicinity of the plant or plant cells, and may include, for example, the cover 420B to a tissue culture dish or 420C the floor of the tissue culture plate. For example, a colorimetric oxygen sensor, or other gas sensor may be employed with the tissue culture dishes. In another example, a pH indicator or nutrient indicator is employed that includes a transmitter or receiver in order to share information relating to the tissue culture plates. In another embodiment (not shown), remote spectral sensing is employed that includes sharing of information or evaluating the information against a data base, as described herein. For example, remote spectral sensing of crops has been utilized through imagery of a field or plot, where the incident electromagnetic radiation is generally sunlight. *See* Li et al., *Electrochem. Soc. Winter 2010*, pp. 41-46, which is incorporated herein by reference. For example, measuring sunlight of the crop and soil that is reflected, absorbed, and/or transmitted depending on the wavelength and the surface it strikes, indicates differences in physical or chemical properties of the crops, e.g., leaf color, texture, shape, etc. In an embodiment, the spectral reflectance measurements are utilized for analysis, including the ratio of reflected energy to incident energy as a function of wavelength, and can be evaluated based on spatial or temporal resolution. In an embodiment, one or more of a spectrometer, radiometer, or digital camera may be mounted on a variety of platforms either on the ground, aerial, or space to gather data. *Id.*

In an embodiment, one or more sensors include soil and/or air sensors 470 that detect environmental conditions. In an embodiment, one or more root sensors 480 are banded on the plant roots with a strain sensor or threshold sensor as described for the plant stalk, or set at a goal distance for root growth to approach and provide a signal. In an

embodiment, one or more cameras 300 are placed on or near the vicinity of the plants or plant cells in order to gather visual information about the plants or plant cells. As described in Figure 3, the cameras include one or more of infrared cameras, visual cameras, x-ray cameras, or similar.

5 In an embodiment, one or more sensors including Raman and Fourier Transform Infrared spectroscopy, capacitance probes, reflectometers, ultrasonic ranging sensors, pH soil-based sensors, Eddy covariance sensors, fluorescence-based optical sensor, optical or microwave sensors, or other sensors may be employed in certain embodiments. *See* Pajares, Sensors (2011) 11:8930-8932, which is incorporated herein by reference.

10 In an embodiment, one or more “smart transducers” are utilized with certain embodiments disclosed. Specifically, “smart transducers” include sensors or actuators equipped with microcontrollers to provide local “intelligence” and network capability and can combine sensing, computing, and communication. *See* Wang et al., Comp. and Elect. In Ag. (2006)50: 1-14, which is incorporated herein by reference.

15 In an embodiment, a system disclosed herein includes a network of research stations or plant testing stations that are configured to share information regarding the one or more measured characteristics of the plantlet or mature plant developed according to methods described herein. In an embodiment, the network of stations includes means to transmit and/or receive information regarding the one or more measured characteristics of
20 the plantlet or mature plant developed according to methods described herein.

As described in Figure 5, a system or method 500 includes culturing or cultivating the progeny 501 of the rapid-breeding process, and analyzing the progeny 515. In an embodiment, genotype information is obtained from the progeny and input 510 into the system. In an embodiment, sensed information as described herein, relating to phenotype
25 505 is input into the system. Utilizing the input information, next a comparator for the genotype and/or phenotype input is employed with a Desired Characteristic Value dataset 520. In an embodiment, the comparator is operably coupled to a database related to one or more of at least one parent plant of the progeny tested, or other progeny of a breeding cycle with at least one parent in common as the progeny tested. In an embodiment, the
30 database is related to progeny of an earlier breeding cycle as the progeny tested. In an embodiment, the database is related to progeny of a later breeding cycle as the progeny tested (for example, when the progeny tested was allowed to grow for a length of time that surpassed the amount of time needed for at least one additional breeding cycle). In an

embodiment, a generator for the Desired Characteristic Value 525 is employed based on the comparison. From the generated Desired Characteristic Value, in an embodiment, the Desired Characteristic Value exceeds a (optionally predetermined) threshold value 530 and an output is generated indicating that the progeny should be continued to be used in the process 535 (e.g., allowed to grow further or be subjected to further analysis). In an embodiment, the Desired Characteristic Value does not exceed a (optionally predetermined) threshold 540 such that an output is generated indicating that the progeny should be discontinued from the process (e.g., culled from the group). *See* for example, Figure 1. In an embodiment, one round of fertilization are performed, and the progeny are from that first round of fertilization. In an embodiment, multiple rounds of fertilization are performed, and the progeny from one or more of the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, or more round.

In an embodiment described herein, where an assessment threshold is met, then one or more stem cells or somatic cells may be selected from the organism at that stage for additional modification or utilization in cycled fertilization as described herein.

For example, a sensed response for a particular progeny being tested generates a Desired Characteristic Value dataset (or a starting dataset is derived from “ideal” Desired Characteristic Values based on quantification of genotype and phenotype characteristics). Based on the queries of the database from further sensors reporting additional information regarding the progeny, the Desired Characteristic Value may increase or decrease as time goes on. Further as described, such database generation or inquiry may occur in real-time, and may provide feedback in the form of an output or other information. For example, as described, once a progeny exceeds a particular threshold, an alert is generated in the form of an output (e.g., visual, audio, tactile, etc.) indicating that the threshold has been exceeded and the progeny should continue to be used in the process. Additionally, any change itself in the Desired Characteristic Value may be alerted as to whether the change has exceeded a threshold (optionally predetermined) for a particularly exuberant progeny. As indicated, in an embodiment, the Desired Characteristic Value (DCV) dataset may be derived from various sources, and may evolve with additional queries or inputs. *See* for example, Figures 5 and 6. In this way, the progeny may be evaluated phenotypically, as well as genotypically or epigenetically, over time. As described, the first analysis of progeny is conducted very early following fertilization, and optionally progeny are selected for culturing in the lab or cultivating in a greenhouse or field. Thus, in an

embodiment the DCV is dynamic, and the first input related to the DCV may include the first genotyping or other analysis of the progeny very early following fertilization, and this information may be maintained over time either as stored information or for use with comparison for other progeny being tested. As indicated herein, any number of steps in the various disclosed rapid-breeding embodiments may be automated. For example, shortly after fertilization when the first analysis is conducted on the progeny, several progeny show positive genotypes for a particular desired characteristic and may start out with a high Desired Characteristic Value. Next, the positive genotype progeny are selected for further culturing and over time additional positive or negative phenotypic or epigenetic changes occur that increase or decrease the progeny's Desired Characteristic Value depending on the "ideal" desired characteristic. Thus, the various characteristics of the progeny may be quantified and compared in order to determine the Desired Characteristic Value over time.

As illustrated in Figure 6, sexual reproduction in plants involves meiosis and mitosis, for example, when gametophytes give rise to sperm (haploid) or egg (haploid) cells. When fertilization occurs, and the male sperm and female egg cells are combined, the resulting zygote (diploid) undergoes mitosis, producing a sporophyte, which in turn undergoes meiosis to produce spores (haploid) that undergo mitosis to produce the gametophyte. As described herein, certain rapid-breeding embodiments bypasses various portions of this process and as a result, increases the efficiency and effectiveness of plant breeding.

As described herein, Figure 7 illustrates system components that may be utilized with any embodiment disclosed herein and includes an input/output device 700 operably coupled with a computing device 720 that includes a processing unit 721, a system memory 722, and a system bus 723 that couples various system components including the system memory 722 to the processing unit 721. The system bus 723 may be any of several types of bus structures including a memory bus or memory controller, a peripheral bus, and a local bus using any of a variety of bus architectures. The system bus 723 may be any of several types of bus structures including a memory bus or memory controller, a peripheral bus, and a local bus using any of a variety of bus architectures. By way of example, and not limitation, such architectures include Industry Standard Architecture (ISA) bus, Micro Channel Architecture (MCA) bus, Enhanced ISA (EISA) bus, Video

Electronics Standards Association (VESA) local bus, and Peripheral Component Interconnect (PCI) bus, also known as Mezzanine bus.

The system memory includes read-only memory (ROM) 724 and random access memory (RAM) 725. A basic input/output system (BIOS) 726, containing the basic
5 routines that help to transfer information between sub-components within the thin computing device 720, such as during start-up, is stored in the ROM 724. A number of program modules may be stored in the ROM 724 or RAM 725, including an operating system 728, one or more application programs 729, other program modules 730 and program data 731.

10 A user may enter commands and information into the computing device 720 through input devices, such as a number of switches and buttons, illustrated as hardware buttons 744, connected to the system via a suitable interface 745. Input devices may further include a touch-sensitive display with suitable input detection circuitry, illustrated as a display 732 and screen input detector 733. The output circuitry of the touch-sensitive
15 display 732 is connected to the system bus 723 via a video driver 737. Other input devices may include a microphone 734 connected through a suitable audio interface 735, and a physical hardware keyboard (not shown). Output devices may include at least one the display 732, or a projector display 736.

In addition to the display 732, the computing device 720 may include other
20 peripheral output devices, such as at least one speaker 738. Other external input or output devices 739, such as a joystick, game pad, satellite dish, scanner or the like may be connected to the processing unit 721 through a USB port 740 and USB port interface 741, to the system bus 723. Alternatively, the other external input and output devices 739 may be connected by other interfaces, such as a parallel port, game port or other port. The
25 computing device 720 may further include or be capable of connecting to a flash card memory (not shown) through an appropriate connection port (not shown). The computing device 720 may further include or be capable of connecting with a network through a network port 742 and network interface 743, and through wireless port 746 and corresponding wireless interface 747 may be provided to facilitate communication with
30 other peripheral devices, including other computers, printers, and so on (not shown). It will be appreciated that the various components and connections shown are examples and other components and means of establishing communication links may be used.

The computing device 720 may be designed to include a user interface. The user interface may include a character, a key-based, or another user data input via the touch sensitive display 732. The user interface may include using a stylus (not shown).

Moreover, the user interface is not limited to an actual touch-sensitive panel arranged for directly receiving input, but may alternatively or in addition respond to another input device such as the microphone 734. For example, spoken words may be received at the microphone 734 and recognized. Alternatively, the computing device 720 may be designed to include a user interface having a physical keyboard (not shown).

In certain instances, one or more components of the computing device 720 may be deemed not necessary and omitted. In other instances, one or more other components may be deemed necessary and added to the computing device.

In certain instances, the computing system typically includes a variety of computer-readable media products. Computer-readable media may include any media that can be accessed by the computing device 720 and include both volatile and nonvolatile media, removable and non-removable media. By way of example, and not of limitation, computer-readable media may include computer storage media. By way of further example, and not of limitation, computer-readable media may include a communication media.

Computer storage media includes volatile and nonvolatile, removable and non-removable media implemented in any method or technology for storage of information such as computer-readable instructions, data structures, program modules, or other data. Computer storage media includes, but is not limited to, random-access memory (RAM), read-only memory (ROM), electrically erasable programmable read-only memory (EEPROM), flash memory, or other memory technology, CD-ROM, digital versatile disks (DVD), or other optical disk storage, magnetic cassettes, magnetic tape, magnetic disk storage, or other magnetic storage devices, or any other medium which can be used to store the desired information and which can be accessed by the computing device 720. In a further embodiment, a computer storage media may include a group of computer storage media devices. In another embodiment, a computer storage media may include an information store. In another embodiment, an information store may include a quantum memory, a photonic quantum memory, or atomic quantum memory. Combinations of any of the above may also be included within the scope of computer-readable media.

Communication media may typically embody computer-readable instructions, data structures, program modules, or other data in a modulated data signal such as a carrier wave or other transport mechanism and include any information delivery media. The term "modulated data signal" means a signal that has one or more of its characteristics set or
5 changed in such a manner as to encode information in the signal. By way of example, and not limitation, communication media include wired media, such as a wired network and a direct-wired connection, and wireless media such as acoustic, RF, optical, and infrared media.

The computing device 720 may also include other removable/non-removable,
10 volatile/nonvolatile computer storage media products. For example, such media includes a non-removable non-volatile memory interface (hard disk interface) 745 reads from and writes for example to non-removable, non-volatile magnetic media, or a removable non-volatile memory interface 750 that, for example, is coupled to a magnetic disk drive 751 that reads from and writes to a removable, non-volatile magnetic disk 752, or is coupled to
15 an optical disk drive 755 that reads from and writes to a removable, non-volatile optical disk 756, such as a CD ROM. Other removable/nonremovable, volatile/non-volatile computer storage media that can be used in the example operating environment include, but are not limited to, magnetic tape cassettes, memory cards, flash memory cards, DVDs, digital video tape, solid state RAM, and solid state ROM. The hard disk drive 757 is
20 typically connected to the system bus 723 through a non-removable memory interface, such as the interface 745, and magnetic disk drive 751 and optical disk drive 755 are typically connected to the system bus 723 by a removable non-volatile memory interface, such as interface 750.

The drives and their associated computer storage media discussed above provide
25 storage of computer-readable instructions, data structures, program modules, and other data for the computing device 720.

A user may enter commands and information into the computing device 720 through input devices such as a microphone, keyboard, or pointing device, commonly referred to as a mouse, trackball, or touch pad. Other input devices (not shown) may
30 include at least one of a touch sensitive display, joystick, game pad, satellite dish, and scanner. These and other input devices are often connected to the processing unit through a user input interface that is coupled to the system bus, but may be connected by other

interface and bus structures, such as a parallel port, game port, or a universal serial bus (USB).

The computing system may operate in a networked environment using logical connections to one or more remote computers, such as a remote computer 780. The
5 remote computer 780 may be a personal computer, a server, a router, a network PC, a peer device, or other common network node, and typically includes many or all of the elements described above relative to the computing device 720, although only a memory storage device. The network logical connections include a local area network (LAN) and a wide
10 area network (WAN), and may also include other networks such as a personal area network (PAN) (not shown). Such networking environments are commonplace in offices, enterprise-wide computer networks, intranets, and the Internet. Included herein is a network for field research station(s).

When used in a networking environment, the computing system is connected to the network 771 through a network interface, such as the network interface 770, the modem
15 772, or the wireless interface 793. The network may include a LAN network environment, or a WAN network environment, such as the Internet. In a networked environment, program modules depicted relative to the computing device 720, or portions thereof, may be stored in a remote memory storage device. By way of example, and not limitation, remote application programs 785 as residing on computer medium 781. It will be
20 appreciated that the network connections shown are examples and other means of establishing communication link between the computers may be used.

In certain instances, one or more elements of the computing device 720 may be deemed not necessary and omitted. In other instances, one or more other components may be deemed necessary and added to the computing device 720.

25 The signal generator 790 includes a signal generator configured to generate a signal indicative of the desired characteristic of the plant or plant tissue. In one embodiment, the signal may include a raw data signal, i.e., a capacitance measurement, a change in one or more measurements of a phenotypic characteristic, or an indicator that a desired characteristic has exceeded a (predetermined) threshold. In one embodiment, the
30 signal generator may include a processor circuit 792, a threshold circuit 794, an output circuit 796, or a communications circuit 798. In one embodiment, the communications circuit may be operable to communicate using an electrical conductor or using a wireless

transmission. In one embodiment, the signal generator may include an instance of the thin computing device 720 and the processor circuit may be the processing unit 721.

In one embodiment, the system actively monitors (e.g., detects, tracks, etc.) a plant or plant tissue located by using at least one of computerized axial tomography, fiber optic
5 thermometry, infrared thermography, magnetic resonance imaging, magnetic resonance spectroscopy, microwave thermography, microwave dielectric spectroscopy, positron emission tomography, ultrasound reflectometry, spectroscopic imaging, visual imaging, infrared imaging, single photon emission computed tomography, global positioning system, satellite imaging, or the like.

10 As illustrated in Figure 8, an embodiment includes a method 800 for Rapid-breeding of animals. As illustrated, cells (stem cells, somatic cells, progenitor cells, etc.) are extracted 805 from Parent 1 and Parent 2 and induced to differentiate *in vitro* tissue culture 810, deriving gametes or transferable nuclei 815 that when combined, and give rise to one or more zygote progeny 885. Next, the progeny are analyzed 830 as described
15 herein utilizing cell, molecular, or microscopic methods, for example, or optionally cultured 825 to give rise to more than one cell. As illustrated, if the progeny are optionally cultured 825, they may be subsequently analyzed 830 for example, by removing a small portion of tissue or clonal cells to evaluate the genotype as described herein. Following analysis of the progeny 830, the progeny are selected 835 based on the analysis.
20 Optionally, the selected progeny may be briefly cultured for expansion 825, or cultured 840 to embryo stage or a later stage. Alternatively, following selection 835 of the progeny based on the analysis, the progeny cells (which include multipotent, pluripotent, and/or totipotent cells) are utilized to repeat 895 the Rapid-breeding process. Optionally, cultured animal cells 825, or cultured embryos 840 may be optionally validated through a second
25 analysis 845 following the growth in culture. Cells may be obtained from progeny at any step following analysis for one or more desired characteristics, as illustrated in Figure 8. For example, in an embodiment, culturing animal cells, tissues, or embryos includes growing them *in vitro*. In an embodiment, cells, tissues, or embryos are grown on feeder cell layers or optionally *in vivo* prior to repeating the Rapid-breeding process.

30 As illustrated in Figure 9, in an embodiment, the progeny generated from the rapid-breeding disclosed herein are analyzed genetically (including genotypically, for example) or epigenetically, as described herein, and further analyzed phenotypically in tissue culture plates, or as tissues/embryos. As illustrated in Figure 9, one or more cameras 901 are

utilized to survey a row or shelf 900, 999, a specific experimental or control group 910, 920, etc. or a specific tissue culture dish, while a location marker (e.g., global positioning system indicator, etc.) (indicated as “ ”) is placed at various positions around or near the growing animal cells. As described herein, the cameras include one or more of infrared
5 cameras, visual cameras, x-ray cameras, or similar. Information detected from these or other sensors are transmitted (optionally wirelessly) to a central database, as indicated by “1,” “2,” or “3,” and the data collected 975 is evaluated by a user that is a person 905, or computer 903, for example. The user evaluates the data and optionally enters additional observed data via an input/output device 904. Based on the data evaluation, a Desired
10 Characteristic Value from one or more samples is assigned, and any experimental progeny that include desired characteristics may be assessed further. Likewise, any experimental progeny that lack the desired characteristics, or that bear a phenotype no better than the control progeny may be culled from further research. In an embodiment, a locator indicator 902 is utilized to locate a particular progeny for further assessment or for culling
15 based at least in part on the location marker in the tissue culture room, shelf, building, or incubator. In an embodiment, for example, the location marker includes a transmitter, and the location indicator 902 includes a receiver. In an embodiment, for example, the location marker includes a receiver, and the location indicator 902 includes a transmitter. In an embodiment, one or more components described in Figure 7 are implemented in
20 Figure 9. In an embodiment, an integrated sensor-to-database collection may be utilized, such as a wireless sensor network, for example that integrates collected information, optionally encrypts information shared wirelessly and optionally through an internet interface. *See Bencini et al., intechopen, pp. 1-18, which is incorporated herein by reference.* In an embodiment, the information contributes to and/or compares the
25 information with a database. In an embodiment, the database includes genomic sequence information for the particular

As illustrated in Figure 10, included in the analysis of the progeny of the rapid-breeding disclosed herein, the phenotype may be assessed by utilizing one or more sensors, for example, that contact the animal cells, or are in the vicinity of the animal cells.
30 In an embodiment, the one or more sensors include at least one of a receiver or transmitter. In an example, the sensors detect particular environmental conditions or animal cell conditions and transmit the information to a user (as shown in Figure 9) where the information is collected and/or evaluated. In an embodiment, as described in Figure 9, one

or more progeny are culled or further assessed depending on the information collected or evaluated from the one or more sensors.

In an embodiment, a capture strip 1020 is utilized for detecting emitted chemicals, such as carbon dioxide, glucose, urea, ammonia, oxygen, pheromones, stress hormones, reactive oxygen species, and others. In an embodiment, the capture strip 1020 is located on the cells. In an embodiment, the capture strip is located in the vicinity of the animal cells, and may include, for example, the cover to a tissue culture dish or the floor of the tissue culture plate. For example, a colorimetric oxygen sensor, or other gas sensor may be employed with the tissue culture dishes. In another example, a pH indicator or nutrient indicator is employed that includes a transmitter or receiver in order to share information relating to the tissue culture plates. In another embodiment (not shown), remote spectral sensing is employed that includes sharing of information or evaluating the information against a data base, as described herein.

In an embodiment, one or more sensors including Raman and Fourier Transform Infrared spectroscopy, capacitance probes, reflectometers, ultrasonic ranging sensors, pH based sensors, Eddy covariance sensors, fluorescence-based optical sensor, optical or microwave sensors, or other sensors may be employed in certain embodiments. *See* Pajares, Sensors (2011) 11:8930-8932, which is incorporated herein by reference.

In an embodiment, one or more “smart transducers” are utilized with certain embodiments disclosed. Specifically, “smart transducers” include sensors or actuators equipped with microcontrollers to provide local “intelligence” and network capability and can combine sensing, computing, and communication. *See* Wang et al., Comp. and Elect. In Ag. (2006)50: 1-14, which is incorporated herein by reference.

In an embodiment, “real-time” comparisons can be made between the control 1000 and experimental 1010 tissue culture plates, for example by utilizing sensors as described herein or cameras 901.

As described in Figure 11, a system or method 1100 includes culturing the progeny 1101 of the rapid-breeding process, and analyzing the progeny by way of genotyping 1110 or phenotyping 1105. In an embodiment, genotype information is obtained from the progeny and input 1110 into the system. In an embodiment, sensed information as described herein, relating to phenotype 1105 is input into the system. Utilizing the input information, next a comparator for the genotype and/or phenotype input is employed with a Desired Characteristic Value dataset 1120. In an embodiment, the comparator is

operably coupled to a database related to one or more of at least one parent animal of the progeny tested, or other progeny of a breeding cycle with at least one parent in common as the progeny tested. In an embodiment, the database is related to progeny of an earlier breeding cycle as the progeny tested. In an embodiment, the database is related to progeny of a later breeding cycle as the progeny tested (for example, when the progeny tested was allowed to grow for a length of time that surpassed the amount of time needed for at least one additional breeding cycle). In an embodiment, a generator for the Desired Characteristic Value 1125 is employed based on the comparison. From the generated Desired Characteristic Value, in an embodiment, the Desired Characteristic Value exceeds a (optionally predetermined) threshold value 1130 and an output is generated indicating that the progeny should be continued to be used in the process 1135 (e.g., allowed to grow further or be subjected to further analysis). In an embodiment, the Desired Characteristic Value does not exceed a (optionally predetermined) threshold 1140 such that an output 1145 is generated indicating that the progeny should be discontinued from the process (e.g., culled from the group), as described herein at other figures. In an embodiment, one round of fertilization is performed, and the progeny are from that first round of fertilization. In an embodiment, multiple rounds of fertilization are performed, and the progeny from one or more of the second, third, fourth, fifth, sixth, seventh, eighth, ninth, tenth, or more round.

For example, a sensed response for a particular progeny being tested generates a Desired Characteristic Value dataset (or a starting dataset is derived from “ideal” Desired Characteristic Values based on quantification of genotype and phenotype characteristics). Based on the queries of the database from further sensors reporting additional information regarding the progeny, the Desired Characteristic Value may increase or decrease as time goes on. Further as described, such database generation or inquiry may occur in real-time, and may provide feedback in the form of an output or other information. For example, as described, once a progeny exceeds a particular threshold, an alert is generated in the form of an output (e.g., visual, audio, tactile, etc.) indicating that the threshold has been exceeded and the progeny should continue to be used in the process. Additionally, any change itself in the Desired Characteristic Value may be alerted as to whether the change has exceeded a threshold (optionally predetermined) for a particularly exuberant progeny. As indicated, in an embodiment, the Desired Characteristic Value (DCV) dataset may be derived from various sources, and may evolve with additional queries or inputs, as

described herein. In this way, the progeny may be evaluated phenotypically, as well as genotypically or epigenetically, over time. As described, the first analysis of progeny is conducted very early following fertilization, and optionally progeny are selected for culturing in the lab or cultured further. Thus, in an embodiment the DCV is dynamic, and the first input related to the DCV may include the first genotyping or other analysis of the progeny very early following fertilization, and this information may be maintained over time either as stored information or for use with comparison for other progeny being tested. As indicated herein, any number of steps in the various disclosed rapid-breeding embodiments may be automated. For example, shortly after fertilization when the first analysis is conducted on the progeny, several progeny show positive genotypes for a particular desired characteristic and may start out with a high Desired Characteristic Value. Next, the positive genotype progeny are selected for further culturing and over time additional positive or negative phenotypic or epigenetic changes occur that increase or decrease the progeny's Desired Characteristic Value depending on the "ideal" desired characteristic. Thus, the various characteristics of the progeny may be quantified and compared in order to determine the Desired Characteristic Value over time.

As illustrated in Figure 12, spermatogenesis involves both mitosis and meiosis, during which the chromosome number in the maturing cell is reduced from $2n$ to n . Likewise, as illustrated in Figure 13, oogenesis involves both mitosis and meiosis as well, during which the chromosome number in the maturing cell is also reduced from $2n$ to n .

The various embodiments disclosed herein allow for Rapid-breeding of plants and animals by utilizing advanced technologies to circumvent the as-of-now requirements for growth of organisms into mature adults (reproductively competent). Several prophetic examples are illustrated below.

25

Prophetic Examples

Prophetic Example 1: Cycling of Gametes in Vitro (CoGiV)

1. Producing pluripotent plant stem cells in vitro.

Plant cell explants are cultured in vitro (dedifferentiated) to generate callus cells which contain pluripotent plant stem cells. Plant pluripotent stem cells may be obtained from in vitro plant tissue cell cultures which are initiated from somatic or embryonic tissues. For example, a leaf tissue explant, of approximately 1 cm^2 , is cultured on solid media (containing agar) which contains essential salts, vitamins and hormones. The media

30

contain plant hormones, auxins (e.g., 2,4 dichlorophenoxyacetic acid (2,4-D)) and cytokinins known to induce growth of a callus which is comprised of pluripotent stem cells (see e.g., He et al., PLoS Genetics 8: e1002911. doi:10.1371/journal.pgen.1002911 and Plant Cell and Tissue Culture, p. 3-15, Indra K. Vasil, Trevor A. Thorpe (eds.), 1994
5 Kluwer Acad. Publ., Dordrecht, The Netherlands, each of which is incorporated herein by reference). Callus cells which arise in culture are propagated in suspension culture and used to generate gametes for in vitro fertilization.

2. Inducing differentiation of plant stem cells to gametes in vitro.

Pluripotent callus cells are differentiated in vitro to undergo meiosis and generate
10 haploid gametes. The stem cells are cocultured in vitro with plant tissues and media which promote gametogenesis. For example, male gametes may be derived by coculture with anther-derived tissues and culture media which promote meiosis and gamete differentiation. For example, anther tissues from *Arabidopsis thaliana* including the endothecium, middle cell layer and tapetum, are cocultured with the pluripotent stem cells
15 to initiate and support gamete formation (see e.g., Wilson et al., *Reproduction* **128**: 483-492, 2004 which is incorporated herein by reference). Media and methods to promote meiosis and microspore culture may be optimized from similar described (see e.g., International Publication No. WO 03/017753 by Dirks et al., published on March 6, 2003 and Wang et al., *Plant Physiol.* **124**: 523-530, 2000 each of which is incorporated herein
20 by reference). To induce female gamete differentiation in pluripotent cells tissues from the ovule, including the nucellus and integument, the tissues are cocultured with the stem cells. Methods to induce meiosis and megaspore differentiation may be optimized from similar described (see e.g., Wilson et al., *Ibid.* and Wang et al., 2000, *Ibid.*). Gametes obtained by in vitro culture and differentiation of pluripotent stem cells are “bred” by
25 electrofusion in vitro.

3. Breeding of haploid gametes to generate hundreds of diploid plant zygotes which are sorted and cultured in vitro.

Gametes derived from pluripotent stem cells are “bred” by fusing pairs of gametes and culturing the diploid hybrids (i.e., zygotes) in vitro to generate cell lines and/or plant
30 embryos. Gamete pairs are fused by applying a brief electric charge (50 msec, 1 kV/cm) to the gametes which are aligned by dielectrophoresis. Electrofusion and culture of gamete pairs in vitro may yield hundreds of plant embryos. For example, electrofusion of egg cells and sperm from maize may be done on microscope coverslips with

approximately 2000 gamete pairs and yield fusion products, i.e., zygotes, which are capable of growing to become plant embryos (see *e.g.*, Kranz et al., *The Plant Cell* **5**: 739-746, 1993 which is incorporated herein by reference). Zygotes obtained by electrofusion are propagated in vitro and sorted using a fluorescence activated cell sorter (FACS). For example, multicellular structures obtained approximately 5 days after electrofusion of maize gametes are stained with fluorescent DNA stains and fluorescently labeled antibodies, and sorted into individual wells of a microtiter plate. Zygotes may be stained with propidium iodide to determine DNA content, (*e.g.*, diploid, haploid, tetraploid, or aneuploid) and antibodies which detect cell wall components or plasma membrane components. Methods and instrumentation for flow cytometry sorting of plant cells and plant cell protoplasts are described (see *e.g.*, Gaurav, Vishal, "Flow Cytometry of Cultured Plant Cells for Characterization of Culture Heterogeneity and Cell Sorting Applications" (2011), Open Access Dissertations, Paper 370 available online at: scholarworks.umass.edu/open_access_dissertations/370/ which is incorporated herein by reference). Zygotes with the appropriate number of chromosomes, *e.g.*, 2n, and expressing desired cell wall or cell membrane components are cultured in vitro to generate cells for genomic selection.

4. Genomic selection of zygotes

Zygotes selected by cell sorting are propagated in vitro and then selected by genomic selection. Genomic selection of zygotes is done by determining the DNA sequence of single nucleotide polymorphisms (SNPs) which mark specific chromosomal loci, haplotypes and alleles. For example, SNPs may be determined by whole genome sequencing (see *e.g.*, Resende Jr. et al., *New Phytologist* **193**: 617-624, 2012 which is incorporated herein by reference). SNPs may provide markers for specific haplotypes, quantitative trait loci and individual alleles that are present in the zygotes. Next generation sequencing technology combined with other technologies is used to determine SNPs present in the individual zygote DNAs. (See *e.g.*, Siva P. Kumpatla et al., (2012) *Genomics-Assisted Plant Breeding in the 21st Century: Technological Advances and Progress*, Plant Breeding, pp. 131-184, Abdurakhmonov (Ed.) InTech Publ., Rijeka, Croatia, which is included herein by reference.) For example Illumina's BeadArray platform (available from Illumina, San Diego, CA) uses beads with oligonucleotides attached to determine 3072 SNPs in a single reaction. Zygotes with combinations of

preferred chromosomes are selected for another round of breeding and/or embryogenesis and optionally, plantlet development.

5. Cell culture of selected zygotes to obtain stem cells and plantlets

Selected zygotes are cultured in vitro to obtain stem cells and/or plantlets. The
5 zygotes resulting from electrofusion are pluripotent and give rise to flowering plants if
cultured in vitro. Media and methods to propagate the zygotes in vitro may be optimized
from similar described (e.g., see Kranz et al., *Ibid*). Alternatively to start another round of
breeding the selected zygotes are differentiated in vitro to gametes (see Step 2 above) and
enter another cycle of breeding, and genomic selection followed by in vitro culture to
10 generate zygotes with improved traits. Repeated rounds of gamete cycling combined with
genomic selection will reduce the time required to create highly selected plants with
optimal traits.

Prophetic Example 2: Rapid Breeding of Apple Trees Using Gamete Cycling

15 **In Vitro**

A rapid breeding method is used to create apple trees that produce apples with
desired traits in a short time frame. The rapid breeding method is used in combination
with genetic marker selection to rapidly produce and select apple trees bearing multiple
genetic markers that determine desired apple traits. Stem cells (apical meristem cells) are
20 cultured in vitro and differentiated to generate gametes which are mated to produce
embryos which serve as a source of stem cells for another round of breeding. Multiple
embryos produced at each round of breeding are genotyped using next generation DNA
sequencing technology to identify DNA markers associated with the desired apple traits.

The rapid breeding method may be initiated by in vitro fertilization using gametes
25 from elite strains of apple trees with desired apple fruit traits. For example male gametes
from a semi-sweet apple, e.g., Granny Smith, may be electrofused with egg cells from a
sweet apple, e.g., Golden Delicious. Electrofusion and culture of gamete pairs in vitro
may yield hundreds of plant embryos (see e.g., Kranz et al., *The Plant Cell* 5: 739-746,
1993 which is incorporated herein by reference). The resulting “Granny Smith” X
30 “Golden Delicious” embryos are genotyped to identify embryos with a desired
combination of genes from Granny Smith and Golden Delicious apple trees. For example,
if a Granny Smith apple with the sweetness of a Golden Delicious apple is desired, then an
embryo with a set of “sweetness genes” derived from Golden Delicious apples is selected.

DNA sequencing of the apple genome has identified a set of genes associated with carbohydrate metabolism in Golden Delicious. For example, multiple copies are found of the gene encoding sorbitol-dehydrogenase which converts sorbitol to fructose in the fruit. Moreover there are 71 sorbitol metabolism and transport genes in the apple genome which may be candidates for “sweetness genes”. See *e.g.*, Velasco et al., *Nature Genetics*, **42**: 833-839, 2010 which is incorporated by reference herein. Genotyping of the embryos may be done by using next generation sequencing technology to sequence specific loci, *e.g.*, the sorbitol metabolism genes, or to survey DNA markers (*e.g.*, single nucleotide polymorphisms (SNPs)) genome-wide. Genetic markers may be associated with desired traits in plants by whole genome sequencing or exome sequencing (see *e.g.*, Resende Jr. et al., *New Phytologist* **193**: 617-624, 2012 which is incorporated herein by reference). Genomic selection of apple tree embryos may be done approximately 10-14 days after electrofusion of gametes when a meristem part of the embryo enlarges and cells for DNA isolation and sequencing may be obtained as well as cells for induction of gametes.

Preferred embryos are selected by genomic selection and the corresponding stem cells are isolated for induction of gametes and a second round of breeding. Apple stem cells are isolated from apical meristem cells by dissection of apple embryos thus eliminating the time required for sexual maturation of trees (apple trees require approximately 10 years to mature sexually). The isolation and propagation of meristem cells are described (see *e.g.*, U.S. Patent Appl. No. 2004/0016015 by Nguyen et al., published on Jan. 22, 2004 which is incorporated herein by reference). For example, embryo meristem cells may be obtained 8-10 days following electrofusion of gametes (see Kranz et al., *Ibid.*). To induce gametogenesis in plant stem cells, in vitro cultures are established in conjunction with gene expression vectors encoding transcription factors essential to gametogenesis. Methods to induce gamete production from embryonic stem cells in mice are described (see *e.g.*, Hayashi et al., *Science* **338**: 971-975, 2012 and Nayernia et al., *Developmental Cell* **11**: 125-132, 2006 which are incorporated herein by reference). To generate apple male gametes, an in vitro culture system for apple apical meristem cells is established with plant hormones and nutrients. Methods to culture meristem cells in vitro are known (see *e.g.*, U.S. Patent Appl. No. 2004/0016015, *Ibid.*). To promote differentiation of the stem cells to gametes, vectors encoding transcription factors essential for male gamete production are transfected into the stem cells. For example, MADS-box genes encoding essential transcription factors include:

SPOROCYTELESS (SPL)/NOZZLE (NZZ), AGAMOUS-LIKE 66 (AGL66) and AGL104 (see *e.g.*, Gramzow et al., *Genome Biology* **11**: 214, 2010 available online at: <http://genomebiology.com/2010/11/6/214> which is included herein by reference). Plasmid expression vectors containing MADS-box genes may be derived from *Agrobacterium tumefaciens* Ti plasmids (see *e.g.*, U.S. Patent No. 7,612,258 issued to He et al. on Nov. 3, 2009 which is incorporated herein by reference). Expression vectors with promoter sequences, MADS-box genes, selectable markers (*e.g.* Hygromycin resistance gene) and regulatory sequences may be transferred into stem cells by microprojectile bombardment or electroporation (see *e.g.*, U.S. Patent Appl. 2004/0016015, *Ibid.*) Similarly a vector encoding AGL23, may be used to promote female gamete (*i.e.*, egg cell) differentiation in apple stem cells. MADS-box genes essential to gametogenesis are described (see *e.g.*, Gramzow et al., *Ibid.*)

Gametes derived from the meristem cells of genetically selected apple tree embryos are bred again to improve the apple tree genotype. For example, zygotes bearing “sweetness genes”, *e.g.*, the Golden Delicious alleles for sorbitol metabolism and transport may be crossed with Golden Delicious gametes to capture both alleles at each “sweetness gene” locus. Domesticated apple trees such as Golden Delicious are highly heterozygous, and in addition multiple genes may be required to control a single trait such as sweetness (see *e.g.*, Velasco et al., *Ibid.*). A second round of electrofusion generates hundreds of zygotes which are cultured in vitro for 10-14 days and then genetically selected by DNA sequencing of sorbitol metabolism and transport genes or associated SNP’s (as described above). Progeny are selected that contain Golden Delicious alleles at the sorbitol gene loci and Granny Smith alleles at other loci. Meristem cells may be isolated and induced to generate male and female gametes for another round of breeding.

Repeated breeding cycles coupled with genetic selection using next generation sequencing technology allows rapid development of apple trees with traits such as sweetness which are controlled by multiple heterozygous genes. For example, apple tree embryos with multiple sorbitol metabolism genes derived from Golden Delicious on a background of Granny Smith genes may be selected in multiple crosses which require approximately 11-15 days each. In vitro culture and plant outgrowth of multiple preferred apple tree embryos may be done by optimizing similarly described methods (see *e.g.*, U.S. Patent No. 7,612,258, *Ibid.*) and optimized apple trees bearing fruit may be obtained in less than one year using grafting techniques.

Prophetic Example 3: Rapid Breeding of Cotton by In Vitro Gamete Cycling

To develop improved cotton cultivars a rapid breeding method is employed that induces stem cells to differentiate directly to gametes which are mated and propagated in vitro. An elite cotton cultivar is crossed with a second elite cultivar to select for a desired trait, for example, drought resistance. Rapid breeding is combined with genomic selection to identify preferred progeny emanating from an initial cross and multiple backcrosses.

The rapid breeding method may be initiated by in vitro fertilization using gametes from elite cotton cultivars, *e.g.*, *Gossypium hirsutum*, cv. Siv'on (GH) and *Gossypium barbadense*, cv. F-177 (GB) which carry quantitative trait loci (QTL) conferring drought resistance (see *e.g.*, Levi et al. *Mol. Breeding* **23**: 179-195, 2009 which is incorporated herein by reference). For example male gametes from GH may be electrofused with gametes from GB. Electrofusion and culture of gamete pairs in vitro may yield hundreds of plant embryos (see *e.g.*, Kranz et al., *The Plant Cell* **5**: 739-746, 1993 which is incorporated herein by reference). The resulting "GH X GB" embryos are genotyped to identify embryos bearing chromosomal loci conferring draught resistance: For example QTL on chromosome 6 and chromosome 2 of GB are associated with drought resistance, and QTL on chromosome 25 and chromosome 22 of GH were also linked to drought resistance (see Levi et al., *Ibid.*). Genotyping of the embryos may be done by using next generation sequencing technology to sequence specific loci, *e.g.*, QTL, or to survey DNA markers (*e.g.*, single nucleotide polymorphisms (SNPs)) genome-wide. For example SNPs identifying GH chromosomes versus GB chromosomes may be used for genomic selection (see *e.g.*, Yu et al., *G3, Genes; Genomes; Genetics* **2**: 43-58, 2012 and Resende Jr. et al., *New Phytologist* **193**: 617-624, 2012 which are incorporated herein by reference).

Genotyping of cotton embryos is done using next generation sequencing technology. For example the Illumina Infinium assay (available from Illumina, San Diego, CA) is used to determine cotton SNPs using software also available from Illumina. Genomic selection of cotton embryos may be done approximately 10-14 days after electrofusion of gametes when a meristem part of the embryo enlarges and cells for DNA isolation and sequencing may be obtained as well as cells for induction of gametes (see Kranz et al., *Ibid.*).

Alternatively polymerase chain reaction (PCR) and agarose gels may be used to identify embryos containing a specific gene (see *e.g.*, Wang et al., *Plant Breeding* **130**: 569-573, 2011 which is incorporated herein by reference).

Cotton embryo hybrids bearing GH chromosomes conferring drought resistance are backcrossed to the elite cotton cultivar, GB, to restore the GB genetic background. Gamete cycling and genomic selection are used to expedite the breeding process and eliminate the time needed for sexual maturation of the cotton embryos. For example the
5 generation time for cotton plants in the field is approximately 130-140 days (*e.g.*, see Wang et al., 2011, *Ibid.*) versus a generation time of approximately 11-15 days using gamete cycling in vitro.

Hybrid embryos (GH X GB) bearing the drought-resistance loci are dissected to obtain pluripotent meristem cells which are cultured in vitro to induce gamete formation.
10 Meristem cells that are pluripotent stem cells are isolated from the hybrid embryos, where they appear 8-10 days after electrofusion of gametes (see *e.g.*, Kranz et al., *Ibid.*). The stem cells are cocultured in vitro with plant tissues and media which promote gametogenesis. For example, male gametes may be derived by coculture with anther-derived tissues and culture media which promote meiosis and gamete differentiation. For
15 example, anther tissues from GH including the endothecium, middle cell layer and tapetum are cocultured with the hybrid stem cells to support gamete formation (see *e.g.*, Wilson et al., *Reproduction* **128**: 483-492, 2004 which is incorporated herein by reference). Media and methods to promote meiosis and microspore culture have been
20 described (see *e.g.*, International Publication No. WO 03/017753 by Dirks et al., published on March 6, 2003 and Wang et al., *Plant Physiol.* **124**: 523-530, 2000 which are incorporated herein by reference). For example media for cotton anther and ovule culture in vitro are described (see *e.g.*, Memon et al., *World Applied Sciences Journal* **8**: 76-79, 2010 which is incorporated herein by reference). Murashige and Skoog media with 2,4
25 dichlorophenoxyacetic acid (2,4-D), indole-3-acetic acid (IAA), indole-3-butyric acid (IBA), and kinetin (6-furfuryl-aminopurine) (all available from Sigma-Aldrich Corp., St. Louis, MO) may be used to culture ovule and anther tissues. The plant hormone abscisic acid may be added to promote androgenesis (see *e.g.*, Wang et al., 2000, *Ibid.*). Gametes derived from the stem cells of genetically selected embryos are bred again to improve the cotton cultivar. For example the preferred hybrid embryos (GH X GB) may be
30 backcrossed with GB to create an improved drought-resistant strain with all the positive traits of GB.

Gametes derived from genomically selected GH X GB embryo's meristem cells are bred with GB gametes to obtain cotton with predominantly GB genes and multiple GH

genes conferring drought-resistance. Multiple chromosomal loci may be required to confer a single trait such as drought resistance (see *e.g.*, Levi et al., *Ibid.*). A second round of electrofusion generates hundreds of zygotes which are cultured in vitro for 10-14 days and then genetically selected by DNA sequencing of specific loci or the corresponding SNP's (as described above). Progeny are selected that retain GH drought-resistance loci and carry predominately GB chromosomes otherwise. The selected embryos may be used as a source of stem cells and gametes for another round of breeding. For example, a breeding program may include four backcross generations to select an optimal cultivar with an improved trait. The rapid cycling of gametes method requires approximately 60 days to complete four backcrosses versus approximately 360 days for four generations using alternative methods (see *e.g.*, Wang et al., 2011, *Ibid.*).

Prophetic Example 4: Rapid Breeding of Cattle to Optimize Milk and Beef Production in a Single Breed

Cattle are bred and selected using a rapid breeding method and DNA sequencing to select progeny with chromosomal loci associated with milk and beef production. Rapid breeding of breeds with superior production of milk (*e.g.*, Holstein) and beef (*e.g.*, Angus) is accomplished by cycling of gametes in vitro (CoGiV) and selecting from multiple progeny using next generation DNA sequencing (NGS) technology. Using CoGiV and NGS multiple rounds of breeding and selection are completed in a relatively short timeframe, and superior breeds combining multiple desired chromosomal loci are created.

Bovine embryonic stem cells (ESC) are differentiated to gametes which are mated in vitro to obtain bovine embryos for selection. Embryonic stem cells are initially obtained from bovine embryos using established methods. For example, oocytes are obtained from an Angus cow at a slaughterhouse and fertilized in vitro with sperm obtained from a Holstein bull. Zygotes are cultured for approximately 8-10 days until blastocysts hatch. The blastocysts are cultured over feeder cells (*e.g.*, mouse embryo fibroblasts) with selected growth factors (*e.g.*, leukemia inhibitory factor (LIF), basic fibroblast growth factor (bFGF), and stem cell factor (SCF) (all are available from Life Technologies, Grand Island, NY). Blastocysts attach to feeder cell layers and develop inner cell masses which are isolated mechanically and cultured in the same media over feeder cells. Approximately 120 blastocysts may be obtained by in vitro fertilization of oocytes harvested from bovine ovaries and approximately 20 ESC-like colonies may be

obtained (see *e.g.*, Kim et al., *Cellular Reprogramming* **14** (6) 2012, DOI: 10.1089/cell.2012.0038 which is incorporated herein by reference). Methods to establish bovine ESC lines are described (see *e.g.*, Jin et al., *Cytotechnology* **64**: 379-389, 2012 and Renard et al., *Theriogenology* **68S**: S196-S205, 2007 which are incorporated herein by
5 reference). Next ESC lines, derived from in vitro fertilization of Angus oocytes with Holstein sperm, are screened to identify clones with the optimal combination of preferred chromosomal loci.

DNA sequencing is used to identify ESC clones with preferred chromosomes from Holstein and Angus. Preferred chromosomal loci in cattle have been mapped as
10 quantitative trait loci (QTL) which correspond to traits such as: milk yield, milk fat content, carcass yield, intramuscular fat content, and weight gain. For example loci on multiple chromosomes have been associated with a single trait, *e.g.*, milk yield (see *e.g.*, Polineni et al., *BMC Bioinformatics* **7**: 283, 2006; doi:10.1186/1471-2105-7-283 which is incorporated herein by reference). Moreover the bovine QTL have been tabulated in a
15 database and linked to DNA sequence data for the bovine genome see Polineni et al., *Ibid.*). Furthermore DNA sequence data derived from different breeds of cattle has allowed construction of a map of single nucleotide polymorphisms (SNPs) which discriminate chromosome pairs in different breeds of cattle (see *e.g.*, The Bovine HapMap Consortium, *Science* **324**: 528-532, 2009 which is incorporated herein by reference). Thus
20 specific haplotypes for a desired trait can be identified by their unique SNPs. DNA sequencing of the ESC clones may be done by using next generation sequencing technology to sequence specific loci, *e.g.*, QTL, or to survey DNA markers (*e.g.*, single nucleotide polymorphisms (SNPs)) genome-wide. The Illumina Infinium assay (available from Illumina, San Diego, CA) may be used to determine SNPs using software also
25 available from Illumina. For example analysis of Angus and Holstein chromosomes revealed 1 SNP every 714 base pairs for pairs of Angus or Holstein chromosomes, *i.e.* heterozygous pairs of chromosomes. Identifying SNPs associated with QTL in preferred breeds allows one to select chromosomes based on DNA sequence of the SNP. For example, one may select specific chromosomes associated with high milk production from
30 Holstein which display very high milk production (see *e.g.*, Publication L-5206 Texas Agricultural Extension Service, Texas A&M University, College Station, TX which is incorporated herein by reference). To summarize, ESC clones containing one or more

SNPs (i.e., SNP DNA sequences) which are associated with high milk yield in Holstein are selected for differentiation to gametes and in vitro fertilization.

Rapid breeding of selected ESC clones is accomplished by differentiation of the ESCs to gametes and then mating them in vitro. For example an ESC clone carrying preferred chromosomes from Holstein and Angus associated with high milk production and high quality meat production respectively is differentiated in vitro: first to a primordial germ-like cell and second to a mature oocyte competent for fertilization in vitro. Methods to differentiate mammalian ESC to oocytes are described (see e.g., Hayashi et al., *Science* **338**: 971-975, 2012 and Hubner et al., *Science* **300**: 1251-1256, 2003 which are incorporated herein by reference). Briefly, ESCs are cultured in plates with medium containing activin A, bFGF and KSR (serum replacement) (all available from Life Technologies, Grand Island, NY) for approximately 2 days to derive epiblast like cells which are then cultured in nonadherent plates in serum-free media containing: bone morphogenetic protein 4 (BMP4), LIF, SCF, BMP8a and epidermal growth factor (EGF) (all are available from R&D Systems, Minneapolis, MN). Primordial germ-like cells are obtained after approximately 2 days of culture and they are combined with embryonic gonadal somatic cells (obtained from the slaughterhouse) whereupon they undergo meiosis and complete oogenesis. The oocytes obtained are competent for fertilization, embryonic development and generation of offspring. See e.g., Hayashi et al., *Ibid*.

The ESC-derived oocytes may be fertilized with native sperm from Angus or Holstein or with sperm derived from an ESC clone. Methods to differentiate ESCs in vitro to become sperm cells are known. See for example, Nayernia et al., *Developmental Cell* **11**: 125-132, 2006 and Geijsen et al., *Nature* **427**: 148-154, 2004 which are incorporated herein by reference. For example the sperm may derive from the same ESC clone differentiated to an oocyte (see above). ESCs are cultured with mouse embryo fibroblasts and media which promote formation of embryoid bodies. Embryoid bodies are dispersed and grown in the presence of retinoic acid (available from Sigma Aldrich, St. Louis, MO) which results in proliferation of primordial germ cells (PGC) See Geijsen et al., *Ibid*. Culture of PGC in approximately 1 μ M retinoic acid results in meiosis and differentiation of sperm suitable for fertilization.

In vitro fertilization with ESC-derived sperm cells yields embryos which can be used for another round of breeding (i.e., CoGiV), or transferred to a foster mother for development to term. For example, microinjection of oocytes with ESC-derived sperm

- cells leads to blastocyst development in approximately 20 % of the injected oocytes (see e.g., Geijsen et al., *Ibid.*) and preimplantation embryos which can be transferred to a foster mother and brought to term (Nayernia et al., *Ibid.*). Multiple rounds of breeding and selection can be accomplished rapidly to allow development of superior breeds with
- 5 genotypes enriched for preferred alleles from multiple breeds and from a number of QTL corresponding to several preferred traits. For example, CoGiV may require approximately 2 months to complete a cycle of ESC culture, NGS selection, differentiation of gametes, in vitro fertilization and embryo development(see e.g., Hayashi and Saito, *Nature Protocols* 8: 1513-1524, 2013 and Kim et al., *Ibid.* which are incorporated herein by reference).
- 10 While the minimum generation time for cattle is approximately 25 months (10 months gestation and 15 months to reach sexual maturity). Multiple rounds of breeding and comprehensive selection by DNA sequence of hundreds of progeny will allow rapid development of superior breeds homozygous at multiple loci with preferred alleles and superior traits.
- 15 In an embodiment, one or more cells (somatic, stem cells, primordial germ cells, etc.) are utilized from a progeny for an additional fertilization cycle deriving an additional generation. In an embodiment, the selection of one or more cells from a progeny (e.g., embryo or zygote) occurs prior to implanting into a host mother, or prior to birth. In an embodiment, the selection of one or more cells from a progeny occurs prior to discarding
- 20 the progeny.
- While various aspects and embodiments have been disclosed herein, other aspects and embodiments will be apparent to those skilled in the art. The various aspects and embodiments disclosed herein are for purposes of illustration and are not intended to be limiting, with the true scope and spirit being indicated by the following claims.

25

CLAIMS

What is claimed is:

1. A method for animal breeding comprising:
selecting one or more animal stem cells;
5 inducing at least one of the animal stem cells to differentiate into at least one first gamete;
combining the at least one first gamete with at least one second gamete for fertilization; and
molecularly analyzing the genotype of the progeny resulting from the fertilization
10 prior to implantation into a host mother.
2. The method of claim 1, wherein the second gamete results from the induced differentiation of a second selected stem cell.
3. The method of claim 1, wherein molecularly analyzing the genotype of the progeny includes one or more of genetic sequencing, mRNA sequencing, or
15 protein sequencing.
4. The method of claim 1, further including utilizing at least a portion of the progeny to induce the resident stem cells of the progeny to a second round of differentiation into at least one gamete.
5. The method of claim 4, further including allowing the progeny to develop to birth,
20 wherein the second round of differentiation occurs prior to birth of the progeny.
6. The method of claim 4, further including implanting the progeny into a host animal, wherein the second round of differentiation occurs prior to implantation of the progeny into the host animal.
7. The method of claim 1, further including identifying at least one desired genetic,
25 transcription, or translation characteristic from the sequence of the progeny.
8. The method of claim 7, further including utilizing at least a portion of the progeny having the identified desired characteristic to induce the resident stem cells of the progeny to a second round of differentiation into at least one gamete.

9. The method of claim 8, further including combining the second round gamete with at least one gamete for a second round of fertilization, resulting in second round progeny.
10. The method of claim 9, further including molecularly analyzing the genotype of at least a portion of the second progeny resulting from the second round of fertilization.
11. The method of claim 9, wherein at least the second round of fertilization is performed prior to implantation of the progeny into a host mother.
12. The method of claim 1, wherein the one or more animal stem cells are obtained from at least one of a zygote, an embryo, or a fetus.
13. The method of claim 1, wherein the one or more animal stem cells are induced stem cells derived from a selected animal somatic cell.
14. The method of claim 13, further including modifying the selected animal somatic cell prior to inducing the animal somatic cell to differentiate into the animal stem cell.
15. The method of claim 1, further including at least one additional round of fertilization wherein the one or more gametes are obtained from primordial germ cells of the progeny from the fertilization of the first round.
16. The method of claim 1, wherein selecting the one or more animal stem cells includes selecting one or more cells based on at least one of genetic sequence analysis, mRNA sequence analysis, or gene or protein marker selection.
17. The method of claim 1, further including modifying the at least one selected animal stem cell by at least one of gene insertion; gene deletion; gene mutation; alteration in methylation; disrupting or removing the outer cell membrane or cell nucleus membrane; protein insertion; or manipulation of cell cytoplasm, at least one organelle, telomere, or cell nucleus.
18. The method of claim 1, further including molecularly analyzing at least one polar body resulting from the fertilization.

19. The method of claim 1, wherein at least one step of the method is automated.
20. The method of claim 1, further including selecting the progeny with at least one desired characteristic.
21. The method of claim 1, further including inducing a resident stem cell from the
5 selected progeny to a second round of differentiation into at least one third gamete.
22. The method of claim 1, further including culturing the progeny.
23. The method of claim 1, wherein inducing at least one of the animal stem cells to differentiate into at least one first gamete includes inducing to differentiate to at least one egg cell or at least one sperm cell.
- 10 24. A system, comprising:
- one or more input/output devices having a non-transitory signal bearing medium operable to
- select one or more animal stem cells;
- provide a first protocol or instruct a computing device to execute the first protocol
15 for induction of at least one of the animal stem cells to differentiate into at least one first gamete;
- provide a second protocol or instruct a computing device to execute the second protocol for combining the at least one first gamete and at least one second gamete for fertilization; and
- 20 accept a first input information related to the genotype of the first generation progeny animal or animal cells resulting from the fertilization;
- receive at least one second input information relating to one or more desired characteristics of the progeny animal or animal cells;
- compare the at least one second input information relating to the one or more
25 desired characteristics of the progeny animal or animal cells with a Desired Characteristic Value dataset;


generate a Desired Characteristic Value based on the comparison;

reinitiate receiving at least one second input information relating to one or more desired characteristics of progeny animal or animal cells resulting from a second round of fertilization of gametes differentiated from animal cells, comparing the at least one second input with the Desired Characteristic Value dataset, generating a Desired Characteristic Value based on the comparison until an assessment threshold is satisfied; and

communicate an output indicative of the Desired Characteristic Value of the progeny animal or animal cells.

- 10 25. The system of claim 24, wherein where the assessment threshold is met, further including selecting one or more stem cells from the progeny for an additional round of differentiation of cells.
- 15 26. The system of claim 24, further including receiving at least one third input information relating to one or more desired characteristics of a second generation of progeny animal or animal cells resulting from a second round of fertilization of at least one first generation progeny animal cell.
27. The system of claim 24, wherein the progeny animal or animal cells include multiple cells from the same round of fertilization.
- 20 28. The system of claim 24, wherein the assessment threshold includes one or more of a period of time, characteristic value, progeny location, or disease or death of the progeny.
29. The system of claim 24, wherein the Desired Characteristic Value includes a quantitative evaluation of at least one desired phenotypic characteristic, epigenetic characteristic, or genetic characteristic.
- 25 30. The system of claim 29, wherein the at least one desired phenotypic characteristic includes at least one of increased size, faster maturation of progeny; increased muscle mass; increased lifespan; increased fertility; decreased gestation time; decreased susceptibility to disease or death; increased quality of fur, hair, or

- 5 feathers; increased docility; increased quality of skin, hide, or feathers; increased ability to obtain energy or absorb nutrients from feed; alteration in microbiome of one or more of gut, skin, or other body part; increased tolerance to antibiotics or other anti-microbial agents; increased number of offspring over the animal's lifetime; increased milk production; increased fatty acid production; increased protein production; alteration in type of fatty acid production; increased immunological compatibility of tissues or organs with the human counterpart tissue or organ; or increased adaptability to environment.
- 10 31. The system of claim 30, wherein the increased quality of fur, hair, or feathers includes increased density of fur, hair, or feathers on the animal; increased tensile strength of fur, hair or feathers; increased durability of fur, hair, or feathers; increased size of fur, hair, or feather shaft; increased keratin level in the fur, hair, or feathers; increased length or diameter of the fur, hair, or feathers; increased occurrence of a desired color of fur, hair, or feathers; increased softness of fur, hair, or feathers; increased luster of fur, hair, or feathers; decreased shrinkage of the fur, hair, or feathers; increased grade of fur, hair, or feathers; or increased yield of fur, hair, or feathers.
- 15 32. The system of claim 24, further including inserting one or more markers into the progeny to track it in the environment.
- 20 33. The system of claim 32, wherein the marker includes one or more of a colorimetric marker, electronic marker, protein marker, or genetic marker.
34. The system of claim 24, wherein the input information includes at least one of information obtained directly from a sensor, or information observed by a worker.
- 25 35. The system of claim 34, wherein the worker includes at least one of a farm implement, an unmanned aerial vehicle, or a person.
36. The system of claim 24, wherein the one or more input/output devices is a computer operable to convert information into electronic signals that include digitized or weighted protocols.






FIG.2




FIG.3




FIG.4




FIG. 5







FIG.6

FIG. 7







௧௮௮௮




FIG. 10




FIG.11

12/13




FIG. 12




FIG. 13

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2014/042880**A. CLASSIFICATION OF SUBJECT MATTER****C12Q 1/04(2006.01)i, C12Q 1/68(2006.01)i, C12N 5/00(2006.01)i, C12N 5/073(2010.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12Q 1/04; C12N 5/06; C12N 5/10; C12N 15/89; C12Q 1/68; C12N 5/00; C12N 5/073

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: breeding, stem cell, gamete, fertilization, progeny, cycling

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	HAYASHI et al., `Offspring from oocytes derived from in vitro primordial germ cell-like cells in mice` Science, Vol.338, No.6109, pp.971-975 (2012) See abstract; pages 972-975; and figures 1-3.	1-36
A	HAYASHI et al., `Reconstitution of the mouse germ cell specification pathway in culture by pluripotent stem cells` Cell, Vol.146, No.4, pp.519-532 (2011) See abstract; pages 519-520, 526-527, 529, 531; and figures 1-6.	1-36
A	WO 2006-108284 A1 (UNIVERSITY OF GUELPH) 19 October 2006 See abstract and claims 1-35.	1-36
A	US 2012-0167242 A1 (WILES et al.) 28 June 2012 See abstract and claims 1-56.	1-36
A	WO 00-29602 A1 (CEDARS-SINAI MEDICAL CENTER et al.) 25 May 2000 See abstract and claims 1-132.	1-36



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

08 October 2014 (08.10.2014)

Date of mailing of the international search report

08 October 2014 (08.10.2014)

Name and mailing address of the ISA/KR

International Application Division
Korean Intellectual Property Office
189 Cheongsa-ro, Seo-gu, Daejeon Metropolitan City, 302-701,
Republic of Korea

Facsimile No. +82-42-472-7140

Authorized officer

KIM, Seung Beom

Telephone No. +82-42-481-3371



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2014/042880

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
PX	MURRAY et al., 'A proposal to use gamete cycling in vitro to improve crops and livestock' Nature Biotechnology, Vol.31, No.10, pp.877-880 (Epub. October 2013) See pages 878-879 and figure 1.	1-36

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2014/042880

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2006-108284 A1	19/10/2006	None	
US 2012-0167242 A1	28/06/2012	US 2013-0298269 A1 US 8658155 B2 WO 2012-092284 A1	07/11/2013 25/02/2014 05/07/2012
WO 00-29602 A1	25/05/2000	CA 2309904 A1 CA 2350829 A1 CA 2368620 A1 EP 1030929 A1 EP 1030929 B1 EP 1047792 A1 EP 1129205 A1 EP 1190086 A2 JP 2001-523468 A JP 2002-543814 A JP 2003-525581 A JP 2003-526326 A US 2002-0053092 A1 US 2002-0056148 A1 US 2002-0083479 A1 US 2002-0129398 A1 US 2002-0133835 A1 US 2002-0138865 A1 US 2005-0034177 A1 US 2008-0178311 A1 US 6316692 B1 US 6734338 B1 US 7294755 B1 WO 00-29601 A1 WO 00-69257 A2 WO 00-69257 A3 WO 99-25863 A1	27/05/1999 25/05/2000 23/11/2000 30/08/2000 18/03/2009 02/11/2000 05/09/2001 27/03/2002 27/11/2001 24/12/2002 02/09/2003 09/09/2003 02/05/2002 09/05/2002 27/06/2002 12/09/2002 19/09/2002 26/09/2002 10/02/2005 24/07/2008 13/11/2001 11/05/2004 13/11/2007 25/05/2000 23/11/2000 09/08/2001 27/05/1999