



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

Publication number:

**0 135 376  
B1**

12

## EUROPEAN PATENT SPECIFICATION

45 Date of publication of patent specification: 07.11.90

51 Int. Cl.<sup>5</sup>: A 61 K 39/39, A 61 K 47/00

71 Application number: 84305687.0

72 Date of filing: 21.08.84

54 Polyoxypropylene-polyoxyethylene block polymer based adjuvants and vaccines using them.

30 Priority: 22.08.83 US 525190

43 Date of publication of application:  
27.03.85 Bulletin 85/13

45 Publication of the grant of the patent:  
07.11.90 Bulletin 90/45

84 Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

56 References cited:  
DE-A-2 603 321  
US-A-4 370 265

JOURNAL OF IMMUNOLOGY, vol. 127, no. 3,  
September 1981, pages 1244-1250, American  
Association of Immunologists, US; R. HUNTER  
et al.: "The adjuvant activity of nonionic block  
polymer surfactants. I. The role of hydrophile-  
lipophile balance"

L. CHEDID et al., editors: "Immunostimulation",  
1980, pages 111-128, Springer Verlag, Berlin,  
DE; M. PARANT: "Biologic properties of a new  
synthetic adjuvant, muramyl dipeptide (MDP)"

73 Proprietor: SYNTEX (U.S.A.) INC.  
3401 Hillview Avenue  
Palo Alto California 94304 (US)

72 Inventor: Allison, Anthony Clifford  
2513 Hastings Drive  
Belmont California 94002 (US)  
Inventor: Byars, Noelene Elva  
1091 Syracuse Drive  
Sunnyvale California 94087 (US)

74 Representative: Armitage, Ian Michael et al  
MEWBURN ELLIS & CO. 2/3 Cursitor Street  
London EC4A 1BQ (GB)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Courier Press, Learnington Spa, England.

EP 0 135 376 B1

## Description

This invention relates to adjuvants and vaccines. More particularly, this invention relates to adjuvants and vaccines based on polyoxypropylene-polyoxyethylene block polymers.

Freund's discovery that the immunogenicity of antigens could be potentiated by emulsifying an aqueous antigen solution with mineral oil alone or with mineral oil and *M. tuberculosis*, formed the basis of the concept of using a secondary material to increase a subject's humoral and cell-mediated immune responses to an antigen. An essential component of Freund's complete and incomplete adjuvant is mineral oil. This component plays a central role in effecting an increased humoral response to the antigen. However, mineral oil is only acceptable for use in research circumstances. The mycobacteria in complete Freund's adjuvant are essential for significantly enhanced cellular immunity.

Though little attention was initially paid to the role the surfactant may play in Freund's incomplete or complete adjuvant, subsequent research has indicated that in several instances a surfactant may demonstrate adjuvant properties in and of itself. A number of naturally occurring surface active agents such as the lipid A portion of endotoxin of gram negative bacteria and trehalose dimycolate of mycobacteria are among the most potent adjuvants of these naturally occurring surfactants. A constituent of mammalian cells, the phospholipid lysolecithin also has been shown to have adjuvant activity. (B. Arnold et al, *Eur. J. Immunol.*, 9:363—366 (1979).)

In addition, several synthetic surfactants, for example, dimethyldioctadecyl ammonium bromide (DDA) and certain polyoxypropylene-polyoxyethylene block polymers have been reported as having adjuvant activity. (See H. Snippe et al, *Int. Archs. Allergy Appl. Immun.*, 65:390—398 (1981). In addition, R. Hunter et al, have reported in the *Journal of Immunology*, 127:1244—1250 that polyoxypropylene-polyoxyethylene block polymers, when used as the surfactant component of an oil-in-water based adjuvant formulation, increase antibody formation to BSA in mice.

While these natural and synthetic surfactants demonstrate a certain degree of adjuvanticity, results so far published demonstrate that, except for one specific test methodology for DDA, none of the surfactants when used alone matches the immunopotentiating activity found when using complete or incomplete Freund's adjuvant. However, it is not possible to use either Freund's incomplete or complete adjuvant for general vaccination purposes because both mineral oil and mycobacteria have deleterious side effects when injected subcutaneously as a result of which it has not been authorized for domestic animal or human use by governmental regulatory agencies. Mineral oil is limited to use in experimental animals.

However, there is a substantial need for some means of potentiating the immunogenicity of antigens. This is particularly true because virus subunit and other protein antigens are now being prepared by recombinant DNA technology. Moreover, naturally occurring or synthetic peptide fragments from larger proteins known to be antigenic are being administered rather than whole protein or a mixture of materials containing the whole protein.

To elicit useful immune responses, antigenic proteins and haptens must be administered with some type of adjuvant. Neither mineral oil nor mycobacteria can be used, as noted above. Glycopeptides should be able to provide the needed immunopotentiation, but these materials are most effective when presented to the subject as an emulsion. Since mineral oil may not be used due to its toxicity, an alternative emulsion-forming material is needed for administering antigens.

It has now been found that when immunopotentiating glycopeptides and an antigen are confected with a non-toxic polyoxypropylene-polyoxyethylene block polymer and a multiphase stabilizing amount of a glycol ether-based non-toxic surfactant, the immunogenicity of the antigen is increased in the same manner and to the same degree as when mineral oil is used. It has been found that the block polymer is critical to achieving an immune response but that a maximal response is most effectively achieved only when the multiphase system is stabilized by some detergent such as a non-ionic glycol ether-based surfactant. The presence of a metabolizable oil may enhance the effectiveness of these formulations as well. Because the polyoxypropylene-polyoxyethylene block polymers and glycol ether surfactants are non-toxic, this adjuvant formulation may be safely used as a vehicle for enhancing the immunogenicity of antigens administered to birds and mammals.

This invention is drawn to an adjuvant vehicle for vaccines comprising an immunopotentiating amount of glycopeptide, a multiphase-forming amount of a non-toxic polyoxypropylene-polyoxyethylene block copolymer, a multiphase-stabilizing amount of a glycol ether-based surfactant, optionally a metabolizable oil of 6 to 30 carbon atoms, and buffered, isoosmotic aqueous solution in a quantity sufficient to make volume.

In a second aspect, this invention relates to a vaccine composition comprising an effective amount of an antigen, an immunopotentiating amount of a glycopeptide, a multiphase-forming amount of a non-toxic polyoxypropylene-polyoxyethylene block copolymer, a multiphase-stabilizing amount of a glycol ether-based surfactant, optionally a metabolizable oil of 6 to 30 carbon atoms, and buffered, isoosmotic aqueous solution in a quantity sufficient to make volume.

In yet another aspect, this invention relates to a method for enhancing the immunogenicity of an antigen which method comprises emulsifying an antigen with an adjuvant vehicle comprising an immunopotentiating amount of glycopeptide, a multi-phase forming amount of a non-toxic polyoxypropylene-polyoxyethylene block copolymer, a multiphase-stabilizing amount of a glycol ether-based surfactant,

## EP 0 135 376 B1

optionally a metabolizable oil of 6 to 30 carbon atoms, and buffered, isoosmotic aqueous solution in a quantity sufficient to make volume and administering said emulsion to a bird or mammal.

As pointed out hereinabove the uniqueness of this invention lies in the use of polyoxypropylene-polyoxyethylene block copolymers in combination with a second surfactant such as a glycol ether-based surfactant as an adjuvant vehicle which, when formed into an emulsion with an immunopotentiating glycopeptide and an antigen, potentiates the immunogenicity of an antigen. This formulation is further unique in that it represents a formulation which can be safely administered to birds and mammals. Thus it is possible to prepare injectable vaccines wherein the antigen will elicit humoral and cell mediated immune responses comparable with those that can be obtained if the antigen was administered in Freund's incomplete or complete adjuvant. A vaccine with such properties will serve to reduce the number of times an antigen must be administered in order to develop a protective response in the subject. In addition, the amount of antigen needed to be administered to elicit a protective response can be reduced.

The polyoxypropylene-polyoxyethylene (POP-POE) block copolymers of this invention are a widely available material commercially known by the trademark Pluronic® polyols. These compounds are made by the sequential addition of propylene oxide and then ethylene oxide to a low molecular weight, reactive hydrogen compound, usually propylene glycol. The characteristics of these polyols are determined by the molecular weight of the polyoxyethylene glycol nucleus and of the ratio of polyoxypropylene to polyoxyethylene in the product. Polyoxypropylene glycols with a molecular weight of approximately 900 or more are water insoluble and impart hydrophobic characteristics to the polyol. The polyethylene glycol, which usually constitutes from approximately 10% to 90% of the total weight, imparts water-soluble properties to the polymer, thus resulting in a compound having surface active properties.

POP-POE block polymers may be represented empirically by the formula:



wherein a and c are statistically equal. These compounds can be prepared by the methods set out in U.S. Patent 2,674,619 issued to Lunsted. The most common means of making the compound is to react a bifunctional initiator, propylene glycol, with propylene oxide in the presence of a catalytic amount of anhydrous sodium hydroxide and the reaction allowed to proceed until the desired molecular weight is obtained. A set amount of ethylene oxide is then added to the same reaction pot to make the block copolymer.

These block polymer polyols range in molecular weight from approximately 1,000 up to 16,000. The polymers of particular interest to this invention are those having an average molecular weight between about 5,000 and 15,500. Such materials are commercially available from several sources, for example, BASF-Wyandotte Corp., Parsippany NJ. 07054 and E. I. Du pont.

POP-POE polyols available from BASF-Wyandotte under the name Pluronic® are identified by a letter prefix followed by a two or a three digit number. The letter prefixes are L, P and F and refer to the physical form of each polymer, L-liquid, P-paste, or F-flakeable solid. The two and three digit numbers are used to designate the average molecular weight of the polyoxypropylene hydrophobic base in comparison with the percent polyoxyethylene in the total molecule. Take, for example, Pluronic® L-101. The first two digits refer to the fact the polyoxypropylene base has a typical molecular weight of 3,250 while the third digit indicates 10% polyoxyethylene is present in the polymer. A second example would be Pluronic® F-108. Here, again, the "10" designates a polyoxypropylene base which has a typical molecular of 3,250 while the "8" designates that polyoxyethylene comprises 80% of the total molecule.

The block polymers of greatest interest to this invention are those which are liquid over a temperature range between about 15°-40°C. In addition, polymer mixtures of liquid and paste, liquid, paste and flakeable solid or liquid and flakeable solid mixtures which are liquid within the specified temperature range may have utility in this invention.

Preferred block polymers are those having a POP base ranging in molecular weight between about 2250 and 4300 and POE in an amount between about 1 and 30%. More preferred are those polymers wherein POP has a molecular weight falling between 3250 and 4000 and the POE component comprises 10-20%. The Pluronic® polyols L-101, L-121 and L-122 fall within this definition. Most preferred are the polymers wherein POP has a molecular weight of 4000 and POE in an amount of 10% or POP has a molecular weight of 3250 and POE in an amount of 10 eg. Pluronic® polyols L-121 and L-101 respectively.

A multi-phase forming amount of polymer is that quantity which will form micelles, or more broadly, a system characterized as an emulsion or suspension. For the purposes of this invention that is some amount between 0.2% and 49% by volume. A more preferred amount would be between 0.2% and 20% (V/V) though up to 5% is even more preferred and 2.5% is most preferred.

The efficacy of these compositions can best be realized by employing a non-toxic, non-ionic detergent to stabilize the formed multi-phase composition. If the block polymer alone is used, the antigen will show some increased immunogenicity but the effect will be smaller in comparison with the effect observed when the non-ionic detergent is present. This non-ionic detergent could also be characterized as an emulsifying or suspending agent in that it is present for achieving those two purposes and is present in a minor amount

## EP 0 135 376 B1

in comparison to the block polymer, which could be characterized as the "oil" in the compositions of this invention.

There are a substantial number of emulsifying and suspending agents generally used in the pharmaceutical sciences. There are naturally derived materials such as gums from trees, vegetable protein, sugar-based polymers such as alginates and cellulose, and the like. Certain oxypolymers or polymers having a hydroxide or other water soluble substituent substituted on the carbon backbone have surfactant activity, for example, povidone, polyvinyl alcohol, and glycol ether-based mono- and poly-functional compounds. Long chain fatty acid-derived compounds form a third substantial group of emulsifying and suspending agents which could be used in this invention. Though any of the foregoing surfactants could be used so long as they are non-toxic, the glycol ether derived compounds are the emulsifying agents of choice in this invention.

The simplest member of the family of compounds based on the glycol ethers is ethylene oxide. The internal or cyclic ethers of the simplest glycol, ethylene glycol, mono- and diethers of ethylene glycol are also well-known. One important group of glycol ethers is the polyethylene glycols. The compounds of most interest herein are the PEG 200, 300, 400, 600 and 900.

A particularly useful group of surfactants are the sorbitan-based non-ionic surfactants. These surfactants are prepared by dehydration of sorbitol to give 1,4-sorbitan which is then reacted with one or more equivalents of a fatty acid or the fatty acid substituted moiety is further reacted with ethylene oxide to give a second group of surfactants.

The fatty acid substituted sorbitan surfactants are made by reacting 1,4-sorbitan with a fatty acid such as lauric acid, palmitic acid, stearic acid, oleic acid, or a similar long chain fatty acid to give the 1,4-sorbitan mono-ester, 1,4-sorbitan sesquiester or 1,4-sorbitan triester. The common name for these surfactants are for example sorbitan monolaurate, sorbitan monopalmitate, sorbitan monostearate, sorbitan monooleate, sorbitan sesquioleate, sorbitan trioleate. These surfactants are commercially available under the name SPAN or ARLACEL, usually with a letter or number designation which distinguishes between the various mono-, di- and triester substituted sorbitans.

SPAN and ARLACEL surfactants are hydrophilic and are generally soluble or dispersible in oil and tend to form water-in-oil emulsions. They are also soluble in most organic solvents. In water they are generally insoluble but dispersible. Generally these surfactants will have a hydrophilic-lipophilic balance (HLB) number between about 1.8 to 8.6. Such surfactants can be readily made by means known in the art or are commercially available from, for example, ICI, America's Inc., Wilmington, Delaware under the registered mark Atlas®.

A related group of surfactants are the polyoxyethylene sorbitan monoesters and polyoxyethylene sorbitan triesters. These materials are prepared by addition of ethylene oxide to a 1,4-sorbitan monoester or triester. The addition of polyoxyethylene converts the lipophilic sorbitan mono- or triester surfactant to a hydrophilic surfactant generally soluble or dispersible in water and soluble in varying degrees in organic liquids.

These materials, commercially available under the mark TWEEN®, are useful for preparing oil-in-water emulsions, dispersions or for the solubilization of oils and making anhydrous ointments water-soluble or washable. The TWEEN surfactants may be combined with a related sorbitan monoester or triester surfactants to promote emulsion stability. TWEEN surfactants generally have a HLB value falling between 9.6 to 16.7. TWEEN surfactants are commercially available from a number of manufacturers for example ICI, America's Inc., Wilmington, Delaware under the registered mark Atlas® surfactants.

A third group of non-ionic surfactants which could be used alone or in conjunction with SPAN, ARLACEL and TWEEN surfactants are the polyoxyethylene fatty acids made by the reaction of ethylene oxide with a long-chain fatty acid. The most commonly available surfactant of this type is sold under the name MYRJ® and is a polyoxyethylene derivative of stearic acid. MYRJ® surfactants are hydrophilic and soluble or dispersible in water like TWEEN® surfactants. The MYRJ® surfactants may be blended with TWEEN® surfactants or with TWEEN®/SPAN® or ARLACEL® mixtures for use in forming emulsions. MYRJ® surfactants can be made by methods known in the art or are available commercially from ICI America's Inc.

A fourth group of polyoxyethylene based non-ionic surfactants are the polyoxyethylene fatty acid ethers derived from lauryl, acetyl, stearyl and oleyl alcohols. These materials are prepared as before by addition of ethylene oxide to a fatty alcohol. The commercial name for these surfactants is BRIJ®. BRIJ® surfactants may be hydrophilic or lipophilic depending on the size of the polyoxyethylene moiety in the surfactant. While the preparation of these compounds is available from the art, they are also readily available from such commercial sources as ICI America's Inc.

Other non-ionic surfactants which potentially could be used in the practice of this invention are for example: polyoxyethylene, polyol fatty acid esters, polyoxyethylene ether, polyoxypropylene fatty ethers, bee's wax derivatives containing polyoxyethylene, polyoxyethylene lanolin derivatives, polyoxyethylene fatty glycerides, glycerol fatty acid esters or other polyoxyethylene acid alcohol or ether derivatives of long-chain fatty acids of 12—21 carbon atoms.

As the adjuvant and the vaccine formulations of this invention are intended to be multi-phase systems, it is preferable to choose an emulsion-forming non-ionic surfactant which has an HLB value in the range of about 7 to 16. This value may be obtained through the use of a single non-ionic surfactant such as a TWEEN® surfactant or may be achieved by the use of a blend of surfactants such as with a sorbitan mono-

EP 0 135 376 B1


di- or triester with a compatible polyoxyethylene sorbitan mono- or triester based surfactant; a sorbitan ester-polyoxyethylene fatty acid; a sorbitan ester in combination with a polyoxyethylene lanolin derived surfactant; a sorbitan ester surfactant in combination with a high HLB polyoxyethylene fatty ether surfactant; or a polyethylene fatty ether surfactant or polyoxyethylene sorbitan fatty acid.

It is more preferred to use a single non-ionic surfactant, most particularly a TWEEN® surfactant, as the emulsion stabilizing non-ionic surfactant in the practice of this invention. Most preferably the surfactant named TWEEN® 80 otherwise known as polysorbate 80 or polyoxyethylene 20 sorbitan monooleate is the most preferred of the foregoing surfactants.

Multiphase stabilization usually can be effected by having the surfactant present in an amount of 0.05 to 2.5% by weight (w/v) or volume (v/v), depending on whether the material is a solid or liquid at room temperature. An amount of 0.2% to 1% is preferred.

The immune response stimulating glycopeptides of this invention are a group of compounds related to and derived from the N-acetylmuramyl-L-alanyl-D-isoglutamine, which was determined by Ellouz et al, *Biochem. & Biophys. Res. Comm.*, Vol 59, 4, 1317 (1974) to be the smallest effective unit possessing immunological adjuvant activity in *M. tuberculosis*, the mycobacterial component of Freund's complete adjuvant. A number of dipeptide- and polypeptide-substituted muramic acid derivatives were subsequently developed and found to have immunostimulating activity.

Though these glycopeptides are a diverse group of compounds, they can be generally represented by the formula




wherein the pyran ring oxygens are substituted by hydrogen, alkyl, or acyl or the like, or may be replaced by nitrogen-based substituent, particularly the 6-position oxygen; the 2-amino group is an acyl group or some other amide, the lactyl side chain is modified, i.e. is ethyl or another two-position alkyl moiety; and the peptide function is a dipeptide or polypeptide. Furanosyl analogs of the pyranosyl compounds also have immunopotentiating activity and are useful in this invention.

Among the glycopeptides of this invention are those disaccharides and tetrasaccharides linked by meso- $\alpha$ - $\epsilon$ -diaminopimelic acid such as described in U.S. patents 4,235,771 and 4,186,194.

Immune response stimulating glycopeptides which may be used in the practice of this invention are disclosed in U.S. Patents 4,094,971; 4,101,536; 4,153,684; 4,235,771; 4,323,559; 4,327,085; 4,185,089; 4,082,736; 4,369,178; 4,314,998 and 4,082,735; and 4,186,194. The compounds of Japanese patent applications J5 4079-227, J5 4079-228, J5 41206-696 and JA-028012 would also be useful in the practice of this invention.

Methods for preparing these compounds are disclosed and well-known in the art. Preparative process exemplification can be found in U.S. Patent 4,082,736 and 4,082,735. Additional, similar preparative processes may be found in the U.S. patents referenced in the preceding paragraph.

Preferred glycopeptides are those having the Formula I



wherein

## EP 0 135 376 B1

R and R<sub>1</sub> are the same or different and are hydrogen or an acyl radical containing from 1 to 22 carbon atoms;

R<sub>2</sub> is an unsubstituted or substituted alkyl radical containing from 1 to 22 carbon atoms, or an unsubstituted or substituted aryl radical containing from 6 to 10 carbon atoms;

R<sub>3</sub> is hydrogen, alkyl, or aryl of 7 to 10 carbon atoms;

R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms;

X is an aminoacyl moiety selected from the group consisting of alanyl, valyl, leucyl, isoleucyl, α-amino-butyryl, threonyl, methionyl, cysteinyl, glutamyl, glutaminyl, aspartyl, phenylalanyl, tyrosyl, tryptophanyl, lysyl, ornithinyl, arginyl, histidyl, asparaginyl, prolyl, hydroxyprolyl, seryl, or glycy;

R<sub>5</sub> denotes an optionally esterified or amidated carboxyl group; and

R<sub>6</sub> denotes an optionally esterified or amidated carboxyl group;

Alkyl is a straight or branched radical comprised of 1 to 7 carbon atoms exemplified by methyl, ethyl, propyl, butyl, pentyl, hexyl or heptyl or and isomer. Lower alkyl is a radical of 1 to 4 carbon atoms.

An optionally substituted benzyl radical is that benzyl radical which is optionally mono-substituted, di-substituted, or poly-substituted in the aromatic nucleus, for example, by lower alkyl, free, etherified or esterified hydroxyl or mercapto groups, for example, lower alkoxy or lower alkylene dioxy groups, as well as lower alkyl mercapto or trifluoromethyl groups and/or halogen atoms.

An optionally esterified or amidated carboxyl group is, the carboxyl group itself or a carboxyl group esterified with a lower alkanol, such as methanol, ethanol, propanol, butanol, or the carbamoyl group, which, on the nitrogen atom, is unsubstituted or mono-substituted or di-substituted by alkyl, especially lower alkyl, aryl, particularly phenyl, or aralkyl, particularly benzyl. The carbamoyl group may also be substituted with an alkylidene radical such as butylidene or pentylidene radical. In addition, the carbamoyl group R<sub>5</sub> may also be substituted with a carbamoylmethyl group on the nitrogen atom.

Particularly preferred compounds are those of Formula 1 wherein R and R<sub>1</sub> are the same or different and are selected from the group consisting of hydrogen or an acyl radical containing from 1 to 22 carbon atoms; R<sub>2</sub> is methyl; R<sub>3</sub> is hydrogen; X is an aminoacyl moiety selected from the group consisting of L-seryl, L-alanyl, L-valyl, L-leucyl, L-isoleucyl, L-α-aminobutyryl, L-seryl, L-threonyl, L-methionyl, L-cysteinyl, L-phenylalanyl, L-tyrosyl, L-tryptophenyl, L-lysyl, L-ornithyl, L-arginyl, L-histidyl, L-glutamyl, L-glutaminyl, L-aspartyl, L-asparaginyl, L-prolyl, or L-hydroxyprolyl.

A more preferred group of glycopeptides are the compounds of Formula 1 wherein R, R<sub>1</sub> are hydrogen or acyl of 1 to 22 carbon atoms, R<sub>3</sub> is hydrogen, R<sub>3</sub> and R<sub>4</sub> are methyl, and X is L-valyl, L-seryl, L-alanyl, L-threonyl or L-α-aminobutyryl.

Most particularly preferred are the following compounds:

N-acetylmuramyl-L-α-aminobutyryl-D-isoglutamine;

6-O-stearoyl-N-acetylmuramyl-L-α-aminobutyryl-D-isoglutamine;

N-acetylmuramyl-L-threonyl-D-isoglutamine;

N-acetylmuramyl-L-valyl-D-isoglutamine;

N-acetylmuramyl-L-alanyl-D-isoglutamine;

N-acetyl-desmethylmuramyl-L-alanyl-D-isoglutamine;

n-butyl N-acetylmuramyl-L-alanyl-D-glutamate; and

N-acetylmuramyl-L-seryl-D-isoglutamine.

An effective amount of immunostimulating glycopeptide will be that amount which effects an increase in titer level when administered in conjunction with an antigen over that titer level observed when the glycopeptide has not been included in the adjuvanted composition. As can be appreciated, each glycopeptide may have an effective dose range that may differ from the other glycopeptides. Therefore, a single dose range cannot be prescribed which will have a precise fit for each possible glycopeptide within the scope of this invention. However, as a general rule, the glycopeptide will preferably be present in an amount of between 0.01 and 2% by weight/volume (w/v). A more preferred amount is 0.01 to 1% (w/v). The glycopeptide may be N-acetylmuramyl-L-threonyl-D-isoglutamine present in an amount of 0.5% (w/v) with copolymer present in an amount of 2.5%.

Another component of these formulations, which may be present at the option of the formulator, is a metabolizable, non-toxic oil, preferably one of 6 to 30 carbon atoms including, but not limited to, alkanes, alkenes, alkynes, and their corresponding acids and alcohols, the ethers and esters thereof, and mixtures thereof. The oil may be any vegetable oil, fish oil, animal oil or synthetically prepared oil which can be metabolized by the body of the animal or bird to which the adjuvant is been administered and which is not toxic in some manner to the organism. Mineral oil and similar petroleum distillate oils are expressly excluded from this invention.

The oil component of this invention may be any long chain alkane, alkene or alkyne, or an acid or alcohol derivative thereof either as the free acid, its salt or an ester such as a mono-, di- or triester, such as the triglycerides and esters of 1,2-propanediol or similar multi-hydroxy alcohol. Alcohols may be acylated employing a mono or poly functional acid, for example acetic acid, propanoic acid, citric acid and the like. Ethers derived from long chain alcohols which are oils and meet the other criteria set forth herein may also be used.

The individual alkane, alkene or alkyne moiety and its acid or alcohol derivatives will have 6—30 carbon atoms. The moiety may have a straight or branched chain structure. It may be fully saturated or

## EP 0 135 376 B1

have one or more double bonds or acetylenic bonds. Where mono or poly ester- or ether-based oils are employed, the limitation of 6—30 carbons applies to the individual fatty acid or fatty alcohol moiety, not the total carbon count.

Any metabolizable oil, particularly from an animal, fish or vegetable source, may be used herein. It is essential that the oil be metabolized by the animal or bird to which it is administered, otherwise the oil component may cause abscesses, granulomas or even carcinomas, or may make the meat of vaccinated birds and animals unacceptable for human consumption because of the deleterious effect the unmetabolized oil may have on the consumer.

Nuts, seeds and grains are the sources for the so called vegetable oils. Peanut oil, soybean oil, coconut oil, and olive oil, the most commonly available, exemplary the nut oils. Seed oils include safflower oil, cottonseed oil, sunflower seed oil, sesame seed oil and the like. In the grain group, corn oil is readily available but while the oil of other cereal grains such as wheat, oats, rye, rice, triticale and the like may be used, such oils, except for wheat germ oil, are not readily available and therefore of less interest, but only for the reason cereal grain oils may be more difficult to obtain commercially.

The technology for obtaining the vegetable oils is well developed and well known. The compositions of these and other similar oils may be found in, for example, the Merck Index, and source materials on foods, nutrition and food technology.

The 6—10 carbon fatty acid esters of glycerol and 1,2-propanediol, while not occurring naturally in seed oils, may be prepared by hydrolysis, separation and esterification of the appropriate materials starting from the nut and seed oils. These products are commercially available under the name "Neobee" from PVO International, Inc., Chemical Specialities Division, 416 Division Street, Boonton, New Jersey and others. Reference is made to U.S. Patent application Serial No. 341,403, filed January 21, 1982 for methods for making these latter materials.

Oils from any animal source, including birds, may be employed in the adjuvants and vaccines of this invention. Animal oils and fats usually are solids at physiological temperatures due to the fact they exist as triglycerides and have a higher degree of saturation than oils from fish or vegetables. However, fatty acids are obtainable from animal fats by partial or complete triglyceride saponification which gives the free fatty acids. Fats and oils from mammalian milk are metabolizable and therefore could be used in the practice of this invention. The procedures for separation, purification, saponification and other means necessary for obtaining pure oils from animal sources are well known in the art.

Another source of oils are the fish oils. Most fish contain metabolizable oils which may be readily recovered. For example, cod liver oil, shark liver oils, and whale oil such as spermaceti exemplary several of the fish oils which may be used herein. Shark liver oil contains a branched, unsaturated oil known as squalene, 2,6,10,15,19,23-hexamethyl-2,6,10,14,18,22-tetracosahexane which is particularly preferred herein. Squalane, the saturated analog of squalene is also a particularly preferred oil herein. Fish oils, including squalene and squalane, are readily available from commercial sources or may be obtained by methods known in the art.

The oil component of these adjuvants and vaccine formulations will be present in an amount between 1 to 30% by volume/volume but preferably in an amount of 1 to 10% v/v. It is most preferred to use a 5% v/v concentration of oil.

The aqueous portion of these adjuvant compositions is buffered, isoosmotic saline. Because these compositions are intended for parenteral administration, it is preferable to make up these solutions so that the tonicity, i.e., osmolality, is essentially the same as normal physiological fluids in order to prevent post-administration swelling or rapid absorption of the composition because of differential ion concentrations between composition and physiological fluids. It is also preferable to buffer the saline in order to maintain a pH compatible with normal physiological conditions. Also, in certain instances, it may be necessary to maintain the pH at a particular level in order to insure the stability of certain composition components, such as the glycopeptides.

Any physiologically acceptable buffer may be used herein but it has been found that it is most convenient to use a phosphate buffer. Any other acceptable buffer such as acetate, tris, bicarbonate, carbonate, or the like could be used as a substitute for a phosphate buffer.

The pH of the aqueous component will preferably be between 6.0—8.0 though it is preferable to adjust the pH of the system to 6.8 where that pH is compatible with maintaining the stability of other composition components or for any other physiologically suitable reason.

The quantity of buffered saline employed in these compositions will be that amount necessary to bring the value of the composition to unity. That is, a quantity of buffered saline sufficient to make 100% will be mixed in with the other components listed hereinabove in order to bring the solution to volume.

The word antigen refers to any substance, usually a protein or protein-polysaccharide, protein-lipopolsaccharide, polysaccharide or lipopolysaccharide which, when foreign to the blood stream of a bird or animal, on gaining access to the tissue of such an animal, stimulates the formation of specific antibodies and reacts specifically in vivo or in vitro with an homologous antibody. Moreover, it stimulates the proliferation of T-lymphocytes with receptors for the antigen, and can react with the lymphocytes to initiate the series of responses designated cell-mediated immunity.

A hapten is within the scope of this definition. A hapten is that portion of an antigenic molecule or antigenic complex that determines its immunological specificity. Commonly, a hapten is a peptide or poly-

## EP 0 135 376 B1

saccharide in naturally occurring antigens. In artificial antigens it may be a low molecular weight substance such as, for example, an arsanilic acid derivative. A hapten will react specifically in vivo and in vitro with homologous antibodies or T-lymphocytes. Alternative descriptors are antigenic determinant, antigenic structural grouping and haptenic grouping.

5 The formulation of a vaccine using the adjuvant compositions described herein will employ an effective amount of antigenic material. That is, there will be included an amount of antigen which, in combination with the adjuvants, will cause the subject to produce a specific and sufficient immunological response so as to impart protection to the subject to the subsequent exposure to the material or organism against which the vaccine is intended to be effective. Alternatively, the antibody will combine with a

10 hormone or naturally occurring material in such a way as to alter biological processes such as growth. No single dose designation can be assigned which will provide specific guidance for each and every antigen which may be employed in this invention. The effective amount of antigen will obviously be a factor of its inherent activity and molecular weight and will be a function of the degree to which the specific antigen can be purified from its source. It is contemplated that the adjuvant compositions of this invention

15 may be used in conjunction with whole cell or virus vaccines as well as purified antigens or subunit or peptide vaccines prepared by recombinant DNA techniques or synthesis.

Adjuvant preparations are readily made by well known art methods. For example, one can make a 2-fold concentrated solution of the antigen and glycopeptide in the buffered saline. A two-fold concentration of the block polymer, oil, and multiphase stabilizing surfactant is mixed with buffered saline and sonicated;

20 then the first and second solution are mixed and sonicated.

A further understanding of the invention may be had from the following non-limiting examples.

### Example 1

The activity of the adjuvant compositions was examined by means of a delayed hypersensitivity test and by an egg albumin (EA) antibody production test in guinea pigs. These two assays measure the ability of the adjuvant compositions to stimulate delayed hypersensitivity (DH, cell mediated response) and antibody synthesis (Ab, humoral immune response) in response to specific antigens in guinea pigs. The two tests were as follows: groups of 8 female guinea pigs were injected subcutaneously with EA emulsified in the adjuvant composition. Appropriate controls were included. The animals were subsequently skin

30 tested with EA to measure delayed hypersensitivity, and were bled to obtain serum for antibody titrations.

Standard bioassay procedures were set up as follows: EA (200 micrograms per animal) was dissolved in saline and then emulsified with each of the six following adjuvant compositions.

TABLE 1  
First Test Composition

<u>Components</u>	<u>Quantity</u>
N-acetylmuramyl-L-threonyl-D-isoglutamine	250 µg
Tween 80	2 µl
Pluronic® L-121	25 µl
Squalene	50 µl
Phosphate buffered saline QS	1,000 µl

TABLE 2  
Second Test Composition

<u>Components</u>	<u>Quantity</u>
N-acetylmuramyl-L-threonyl-D-isoglutamine	250 µg
Tween 80	2 µl
Pluronic® L-121	25 µl
Phosphate buffered saline QS	1,000 µl

EP 0 135 376 B1

TABLE 3  
Third Test Composition

<u>Components</u>	<u>Quantity</u>
N-acetylmuramyl-L-threonyl-D-Isoglutamine	250 µg
Tween 80	2 µl
Squalene	50 µl
Phosphate buffered saline QS	1,000 µl

TABLE 4  
Fourth Test Composition

<u>Components</u>	<u>Quantity</u>
N-acetylmuramyl-L-threonyl-D-Isoglutamine	250 µg
Pluronic® L-121	25 µl
Squalene	50 µl
Phosphate buffered saline QS	1,000 µl

TABLE 5  
Fifth Test Composition

<u>Components</u>	<u>Quantity</u>
Pluronic® L-121	250 µl
Squalene	50 µl
Phosphate buffered saline QS	1,000 µl

TABLE 6  
Sixth Test Composition

<u>Components</u>	<u>Quantity</u>
N-acetylmuramyl-L-threonyl-D-Isoglutamine	250 µg
Pluronic® L-121	25 µl
Phosphate buffered saline QS	1,000 µl

0.2 ml of this emulsion were administered per guinea pig. A booster injection of egg albumin in saline was given at 4 weeks. Blood samples were drawn at three and six weeks to determine EA antibody titers. The egg albumin skin test was carried out at six weeks. Results of these studies are as follows:

# EP 0 135 376 B1

TABLE 7  
Delayed Hypersensitivity and Antibody Production

TEST COMPOSITION	Antibody Titers		DH	
	Ab <sub>3</sub>	Ab <sub>6</sub>	Diam.	Inf.
First	3.83	7.50	15.16	2.33
Second	3.14	6.14	14.14	2.07
Third	1.85	5.42	10.64	1.29
Fourth	3.42	7.85	13.35	1.64
Fifth	2.83	5.33	10.75	1.33
Sixth	2.33	5.00	11.00	1.33

Ab<sub>3</sub> — Antibody titer level at 3 weeks.  
 Ab<sub>6</sub> — Antibody titer level at 6 weeks.  
 DH — Delayed hypersensitivity.  
 Diam. — Diameter of reaction in mm.  
 Inf. — Infiltration, scored on a scale of 1—3.

## Example 2

The effect of squalene on antibody activity was determined in the experiment set forth below. The procedure described in Example 1 was used in this study.

TABLE 8

Components	Quantity
N-acetylmuramyl-L-Thr-D-Iglu	250 µg
Tween 80	2 µl
Pluronic L-121	25 µl
Squalene	50 µl
PBS	to 1,000 µl

TABLE 9

Components	Quantity
N-acetylmuramyl-L-Thr-D-Iglu	250 µg
Tween 80	2 µl
Pluronic L-121	25 µl
PBS	to 1,000 µl

EP 0 135 376 B1

TABLE 10

Components	Quantity
Tween 80	2 $\mu$ l
Pluronic L-121	25 $\mu$ l
PBS	to 1,000 $\mu$ l

Egg albumin was mixed with the above formulations and 0.2 ml injected (S.C.) per guinea pig. A booster injection of egg albumin was given at 4 weeks. Animals were bled at 4 and 6 weeks to determine antibody titres, and skin tested with egg albumin at 6 weeks to measure delayed hypersensitivity. The results of these studies are as follows:

TABLE 11  
Delayed Hypersensitivity and Antibody Production

Test Composition	Antibody Titers		Diam.	DH	
	Ab <sub>4</sub>	Ab <sub>6</sub>			Inf.
First	5.00 $\pm$ 0.37	8.25 $\pm$ 0.31	17.70 $\pm$ 1.12		1.94
Second	2.87 $\pm$ 0.22	6.62 $\pm$ 0.26	12.37 $\pm$ 0.88		1.38
Third	2.12 $\pm$ 0.22	6.00 $\pm$ 0.33	10.25 $\pm$ 0.84		1.00

Example 3

Groups of 6 or 7 female guinea pigs, 400—500 g, were immunized with 200  $\mu$ g of egg albumin (EA) in one of 4 vehicles. The basic vehicle mixture consisted of 0.2 ml of phosphate buffered saline with 0.2% TWEEN<sup>®</sup> 80 containing 10  $\mu$ l of squalene (SQE) or squalane (SQA) and 5  $\mu$ l of Pluronic<sup>®</sup> L-121. Two test groups were also given 50  $\mu$ g/animal of the glycodipeptide N-acetylmuramyl-L- $\alpha$ -aminobutyryl-D-isoglutamine. The animals were boosted with 50  $\mu$ g EA in saline on Day 28. On days 21, 35, and 50 the animals were bled by heart puncture and the sera assayed using the passive hemagglutination technique. On days 35 and 50 the animals were also skin tested with 10  $\mu$ g EA injected ID. The skin tests were measured at 24 hours. The data obtained in this experiment are summarized in Table 12.

From the observations in these three Examples, it is clear that to elicit a powerful cell mediated and humoral response, the combination of glycopeptide and Pluronic<sup>®</sup> polyol is essential. While these two components are essential, cell mediated and humoral response can be maximized by employing a metabolizable oil.

TABLE 12

Delayed Hypersensitivity and Antibody Production


	Antibody Titers			DH			
	Ab <sub>3</sub>	Ab <sub>5</sub>	Ab <sub>7</sub>	35 Days		49 Days	
				Diam.	Inf.	Diam.	Inf.
PBS Control	2.18 $\pm$ .17	2.83 $\pm$ .31	2.83 $\pm$ .31	6.17 $\pm$ 2.31	1.17	3.67 $\pm$ 1.66	1.00
L-121 Control	2.00 $\pm$ .00	4.86 $\pm$ .74	5.57 $\pm$ .30	9.00 $\pm$ 1.61	1.14	8.00 $\pm$ 1.39	1.14
SQE/Glycodipeptide	2.43 $\pm$ .37	7.43 $\pm$ .48	7.83 $\pm$ .31	17.77 $\pm$ 1.38	2.14	12.50 $\pm$ 1.29	1.64
SQA/Glycodipeptide	2.29 $\pm$ .18	6.67 $\pm$ .42	7.83 $\pm$ .31	18.83 $\pm$ 0.71	2.33	13.83 $\pm$ 0.80	1.92

#Infiltration Score: Infiltration is scored from 1 (none) to 3 (strongly infiltrated).

# EP 0 135 376 B1

## Claims

1. An adjuvant composition comprising:
  - an immunopotentiating amount of an immunostimulating glycopeptide;
  - a multi-phase-forming amount of a non-toxic polyoxypropylene-polyoxyethylene block polymer in an amount of 0.2 to 49% by volume (v/v);
  - a multi-phase-stabilizing amount of a glycol ether-based surfactant in an amount between 0.05 to 2.5% by volume (v/v); and
  - buffered isoosmotic saline in a quantity sufficient to make volume.
2. The composition of Claim 1 wherein said glycopeptide is present in an amount between 0.01 to 2% by weight/volume.
3. A composition according to Claim 1 or Claim 2 wherein said glycopeptide is a compound of the formula




wherein

- R and R<sub>1</sub> are the same or different and are hydrogen or an acyl radical containing from 1 to 22 carbon atoms;
  - R<sub>2</sub> is an unsubstituted or substituted alkyl radical containing from 1 to 22 carbon atoms, or an unsubstituted or substituted aryl radical containing from 6 to 10 carbon atoms;
  - R<sub>3</sub> is hydrogen, alkyl, or aryl of 7 to 10 carbon atoms;
  - R<sub>4</sub> is hydrogen or alkyl of 1 to 6 carbon atoms;
  - X is an aminoacyl moiety selected from the group consisting of alanyl, valyl, leucyl, isoleucyl, α-amino-butyryl, threonyl, methionyl, cysteinyl, glutamyl, glutaminyl, aspartyl, phenylalanyl, tyrosyl, tryptophanyl, lysyl, ornithinyl, arginyl, histidyl, asparaginyl, prolyl, hydroxyprolyl, seryl, or glycylyl;
  - R<sub>5</sub> denotes an optionally esterified or amidated carboxyl group; and
  - R<sub>6</sub> denotes an optionally esterified or amidated carboxyl group;
- said block polymer has a polyoxypropylene base of molecular weight between 3,250 and 4,000 and the percent polyoxyethylene in the total molecule comprises 0.2 to 20% (Volume/Volume); and said saline is phosphate buffered saline.
4. The composition of Claim 3 wherein the glycopeptide of Formula I is substituted as follows: R and R<sub>1</sub> are the same or different and are selected from the group consisting of hydrogen or an acyl radical containing from 1 to 22 carbon atoms; R<sub>2</sub> is methyl; R<sub>3</sub> is hydrogen; X is an aminoacyl moiety selected from the group consisting of L-seryl, L-alanyl, L-valyl, L-leucyl, L-isoleucyl, L-α-aminobutyryl, L-seryl, L-threonyl, L-methionyl, L-cysteinyl, L-phenylalanyl, L-tyrosyl, L-tryptophenyl, L-lysyl, L-ornithyl, L-arginyl, L-histidyl, L-glutamyl, L-glutaminyl, L-aspartyl, L-asparaginyl, L-prolyl, or L-hydroxyprolyl, being present in an amount between 0.01 to 1% (w/v);
    - said block copolymer has a polyoxypropylene base of molecular weight between 2250 and 4300 and wherein the percentage of polyoxyethylene in the total molecule is 5—20%, said copolymer being present in an amount between 0.2 to 20% (v/v); and
    - said glycol ether surfactant is a polysorbate surfactant present in an amount of 0.2 to 1% (v/v).
  5. The composition of Claim 4 wherein said glycopeptide is a compound of Formula 1 wherein R, R<sub>1</sub> and R<sub>3</sub> are hydrogen, R<sub>2</sub> is methyl, R<sub>4</sub> is methyl or hydrogen, X is L-valyl, L-alanyl, L-seryl, L-threonyl or L-α-aminobutyryl, R<sub>5</sub> is carboxyl, carbamoyl or n-butyl carboxylate and R<sub>6</sub> is carboxyl or carbamoyl.
  6. A composition according to Claim 5 wherein said glycopeptide is N-acetylmuramyl-L-threonyl-D-isoglutamine, N-acetyl-L-α-aminobutyryl-D-isoglutamine, 6-O-stearoyl-N-acetylmuramyl-L-α-aminobutyryl-D-isoglutamine, or N-acetyl-desmethylmuramyl-L-alanyl-D-isoglutamine present in an amount between 0.01 to 1% (w/v);
    - said block polymer is comprised of a polyoxypropylene base of molecular weight 4000 and has a percentage polyoxyethylene in the total molecule of 10% and is present in an amount between 0.2 to 5% (v/v); and
    - said polysorbate is TWEEN 80®.

7. A composition according to Claim 5 wherein said glycopeptide is n-butyl-N-acetylmuramyl-L-alanyl-D-glutamate, N-acetylmuramyl-L-alanyl-D-isoglutamine, or N-acetylmuramyl-L-seryl-D-isoglutamine; and said copolymer is comprised of a polyoxypropylene base of molecular weight 4000 and which contains 10% polyoxyethylene as a percentage of the total molecule.
8. A composition according to Claim 6 wherein said glycopeptide is N-acetylmuramyl-L-threonyl-D-isoglutamine present in an amount of 0.5% (w/v), and said copolymer is present in an amount of 2.5%.
9. A composition according to any one of the preceding claims which further comprises a metabolizable, non-toxic oil in an amount between 1 to 30% by volume (v/v).
10. The composition according to any one of claims 4 to 8 which further comprises squalene or squalane present in an amount between 1 to 10% (v/v).
11. A vaccine composition comprising:  
 an effective amount of an antigen;  
 an immunopotentiating amount of an immunostimulating glycopeptide;  
 a non-toxic polyoxypropylene-polyoxyethylene block copolymer in an amount of 0.2 to 49% by volume (v/v);  
 a glycol ether based surfactant in an amount of 0.05 to 2.5% by volume (v/v); and  
 buffered isoosmotic saline in a quantity sufficient to make volume.
12. A vaccine composition of claim 11, further comprising a metabolizable, non-toxic oil in an amount between 1 to 30% by volume (v/v).
13. A vaccine composition comprising an effective amount of an antigen and an adjuvant composition according to any one of claims 1 to 10.
14. An adjuvant composition according to any one of claims 1 to 10 for use in enhancing the immunogenicity of an antigen mixed therewith as a multi-phase system for administering to a bird or mammal.
15. An adjuvant composition according to claim 1 wherein the glycopolymer comprises 0.01—2% (w/v) N-acetyl-muramyl-L-threonyl-D-isoglutamine.
16. An adjuvant composition according to claim 1 or claim 15 which comprises 2.5% (w/v) polyoxypropylene-polyoxyethylene block copolymer which has a polyoxypropylene base of molecular weight between 3250 and 4000 and wherein polyoxyethylene comprises 10% of the total molecule.
17. An adjuvant composition according to claim 1, claim 15 or claim 16 wherein the glycol ether-based surfactant comprises 0.2 to 1% (w/v) polyoxyethylene 20 sorbitan monooleate surfactant.

Patentansprüche

1. Adjuvans-Zusammensetzung, umfassend:  
 eine immunpotenzierende Menge eines immunstimulierenden Glycopeptids;  
 eine Multi-Phasen-bildende Menge eines nicht-toxischen Polyoxypropylen-Polyoxyethylen-Blockpolymeren in einer Menge von 0,2 bis 49 Volumen-% (Vol/Vol);  
 eine Multi-Phasen-stabilisierende Menge eines Tensids auf Glykoether-Basis in einer Menge von 0,05 bis 2,5 Volumen-% (Vol/Vol);  
 gepufferte isoosmotische Kochsalzlösung in einer Menge, die genügt, Volumen zu erzeugen.
2. Zusammensetzung nach Anspruch 1, worin das Glycopeptid in einer Menge zwischen 0,01 bis 2 Gewichts/Volumen-% anwesend ist.
3. Zusammensetzung nach Anspruch 1 oder Anspruch 2, worin das Glycopeptid eine Verbindung der Formel



- ist, worin
- R und R<sub>1</sub> gleich oder verschieden sind und Wasserstoff oder ein Acylrest mit 1 bis 22 Kohlenstoffatomen sind;
- R<sub>2</sub> ein unsubstituierter oder substituierter Alkylrest mit 1 bis 22 Kohlenstoffatomen oder ein unsubstituierter oder substituierter Arylrest mit 6 bis 10 Kohlenstoffatomen ist;
- R<sub>3</sub> Wasserstoff, Alkyl oder Aryl mit 7 bis 10 Kohlenstoffatomen ist;
- R<sub>4</sub> Wasserstoff oder Alkyl mit 1 bis 6 Kohlenstoffatomen ist;

X eine Aminoacyl-Einheit, ausgewählt aus der Gruppe bestehend aus Alanyl, Valyl, Leucyl, Isoleucyl,  $\alpha$ -Aminobutyryl, Threonyl, Methionyl, Cysteinyl, Glutamyl, Glutaminyl, Aspartyl, Phenylalanyl, Tyrosyl, Tryptophanyl, Lysyl, Ornithyl, Arginyl, Histidyl, Asparaginy, Prolyl, Hydroxyprolyl, Seryl oder Glycyl, ist;

<sup>5</sup>  $R_5$  eine gegebenenfalls veresterte oder amidierte Carboxylgruppe darstellt; und

$R_6$  eine gegebenenfalls veresterte oder amidierte Carboxylgruppe darstellt;

das Blockpolymer eine Polyoxypropylen-Basis mit einem Molekulargewicht zwischen 3250 und 4000 aufweist und der Prozentsatz an Polyoxyethylen im gesamten Molekül 0,2 bis 20% (Volumen/Volumen) umfaßt; und

die Kochsalzlösung Phosphat-gepufferte Kochsalzlösung ist.

<sup>10</sup> 4. Zusammensetzung nach Anspruch 3, worin das Glycopeptid der Formel (I) wie folgt substituiert ist:  $R$  und  $R_1$  sind gleich oder verschieden und sind ausgewählt aus der Gruppe bestehend aus Wasserstoff oder einem Acylrest mit 1 bis 22 Kohlenstoffatomen;  $R_2$  ist Methyl;  $R_3$  ist Wasserstoff;  $X$  ist eine Aminoacyl-Einheit, ausgewählt aus der Gruppe bestehend aus L-Seryl, L-Alanyl, L-Valyl, L-Leucyl, L-Isoleucyl, L- $\alpha$ -Aminobutyryl, L-Seryl, L-Threonyl, L-Methionyl, L-Cysteinyl, L-Phenylalanyl, L-Tyrosyl, L-Tryptophanyl, L-

<sup>15</sup> Lysyl, L-Ornithyl, L-Arginyl, L-Histidyl, L-Glutamyl, L-Glutaminyl, L-Aspartyl, L-Asparaginy, L-Prolyl oder L-Hydroxyprolyl; wobei es in einer Menge zwischen 0,01 bis 1% (Gew/Vol) anwesend ist;

das Blockpolymer eine Polyoxypropylen-Basis mit einem Molekulargewicht zwischen 2250 und 4300 aufweist und worin der Prozentsatz an Polyoxyethylen im gesamten Molekül 5—20% ist, wobei dieses Copolymer in einer Menge zwischen 0,2 bis 20% (Vol/Vol) anwesend ist; und

<sup>20</sup> das Glykolether-Tensid ein Polysorbat-Tensid ist, das in einer Menge von 0,2 bis 1% (Vol/Vol) anwesend ist.

5. Zusammensetzung nach Anspruch 4, worin das Glycopeptid eine Verbindung der Formel 1 ist, worin

$R$ ,  $R_1$  und  $R_3$  Wasserstoff sind,  $R_2$  Methyl ist,  $R_4$  Methyl oder Wasserstoff ist,  $X$  L-Valyl, L-Alanyl, L-Seryl, L-Threonyl oder L- $\alpha$ -Aminobutyryl ist,  $R_5$  Carboxyl, Carbamoyl oder n-Butylcarboxylat ist und  $R_6$  Carboxyl oder Carbamoyl ist.

<sup>25</sup> 6. Zusammensetzung nach Anspruch 5, worin das Glycopeptid N-Acetylmuramyl-L-threonyl-D-isoglutamin, N-Acetyl-L- $\alpha$ -aminobutyryl-D-isoglutamin, 6-O-Stearoyl-N-acetylmuramyl-L- $\alpha$ -aminobutyryl-D-isoglutamin oder N-Acetyl-desmethylmuramyl-L-alanyl-D-isoglutamin ist, anwesend in einer Menge zwischen 0,01 bis 1% (Gew/Vol);

<sup>30</sup> das Blockpolymer eine Polyoxypropylen-Basis des Molekulargewichts 4000 umfaßt und einen Prozentsatz an Polyoxyethylen im gesamten Molekül von 10% aufweist und in einer Menge zwischen 0,2 bis 5% (Vol/Vol) anwesend ist; und

das Polysorbat TWEEN 80<sup>®</sup> ist.

<sup>35</sup> 7. Zusammensetzung nach Anspruch 5, worin das Glycopeptid n-Butyl-N-acetylmuramyl-L-alanyl-D-gutaminat, N-Acetylmuramyl-L-alanyl-D-isoglutamin oder N-Acetylmuramyl-L-seryl-D-isoglutamin ist; und das Copolymer eine Polyoxypropylen-Basis des Molekulargewichts 4000 umfaßt, und das 10% Polyoxyethylen als Prozentsatz des gesamten Moleküls enthält.

8. Zusammensetzung nach Anspruch 6, worin das Glycopeptid N-Acetylmuramyl-L-threonyl-D-isoglutamin, anwesend in einer Menge von 0,5% (Gew/Vol), ist, und das Copolymer in einer Menge von <sup>40</sup> 2,5% anwesend ist.

9. Zusammensetzung nach irgendeinem der vorangehenden Ansprüche, die weiter ein metabolisierbares, nichttoxisches Öl in einer Menge zwischen 1 bis 30 Volumen-% (Vol/Vol) umfaßt.

10. Zusammensetzung nach irgendeinem der Ansprüche 4 bis 8, die weiter Squalen oder Squalan, anwesend in einer Menge zwischen 1 bis 10% (Vol/Vol), umfaßt.

<sup>45</sup> 11. Impfzusammensetzung, umfassend:

eine wirksame Menge eines Antigens;

eine immunpotenzierende Menge eines immunstimulierenden Glycopeptids;

ein nicht-toxisches Polyoxypropylen-Polyoxyethylen-Blockcopolymer in einer Menge von 0,2 bis 49 <sup>50</sup> Volumen-% (Vol/Vol);

ein Tensid auf Glykolether-Basis in einer Menge von 0,05 bis 2,5 Volumen-% (Vol/Vol); und

gepufferte isoosmotische Kochsalzlösung in einer Menge, die ausreicht, Volumen zu erzeugen.

12. Impfzusammensetzung nach Anspruch 11, weiter ein metabolisierbares, nicht-toxisches Öl in einer <sup>55</sup> Menge zwischen 1 bis 30 Volumen-% (Vol/Vol) umfassend.

13. Impfzusammensetzung, eine effektive Menge eines Antigens und einer Adjuvans-

<sup>55</sup> Zusammensetzung nach irgendeinem der Ansprüche 1 bis 10 umfassend.

14. Adjuvans-Zusammensetzung nach irgendeinem der Ansprüche 1 bis 10 zur Verwendung bei der Verstärkung der Immunisierungsfähigkeit eines Antigens, das damit in Form eines Multiphasen-System zur <sup>60</sup> Verabreichung an einen Vogel oder ein Säugetier gemischt wird.


15. Adjuvans-Zusammensetzung nach Anspruch 1, worin das Glycopolypeptid 0,01 bis 2% (Gew/Vol) <sup>60</sup> N-Acetyl-muramyl-L-threonyl-D-isoglutamin umfaßt.

16. Adjuvans-Zusammensetzung nach Anspruch 1 oder Anspruch 15, umfassend 2,5% (Gew/Vol) <sup>65</sup> Polyoxypropylen-Polyoxyethylen-Blockcopolymer, das eine Polyoxypropylen-Basis eines Molekulargewichts zwischen 3250 und 4000 aufweist und worin Polyoxyethylen 10% des gesamten Moleküls umfaßt.

17. Adjuvans-Zusammensetzung nach Anspruch 1, Anspruch 15 oder Anspruch 16, worin das Tensid <sup>65</sup> auf Glykolether-Basis 0,2 bis 1% (Gew/Vol) Polyoxyethylen-20-Sorbitanmonooleat-Tensid umfaßt.

Revendications

1. Composition d'adjuvant comprenant:  
 une quantité immunopotentialisatrice d'un glycopeptide immunostimulant;  
 une quantité, formant des phases multiples, d'un polymère séquencé polyoxypropylène-polyoxyéthylène non toxique, comprise dans l'intervalle de 0,2 à 49% en volume (v/v);  
 une quantité, stabilisant les phases multiples, d'un surfactant, à base d'éther de glycol, comprise dans l'intervalle de 0,05 à 2,5% en volume (v/v); et  
 du sérum physiologique iso-osmotique tamponné, en une quantité suffisante pour compléter au volume.
2. Composition suivant la revendication 1, dans laquelle le glycopeptide est présent en une quantité de 0,01 à 2% en poids/volume.
3. Composition suivant la revendication 1 ou la revendication 2, dans laquelle le glycopeptide est un composé de formule



I

4. Composition suivant la revendication 3, dans laquelle le glycopeptide de formule I est substitué de la manière suivante: R et R<sub>1</sub> sont identiques ou différents et représentent l'hydrogène ou un radical acyle contenant 1 à 22 atomes de carbone;  
 R<sub>2</sub> représente un radical alkyle non substitué ou substitué, contenant 1 à 22 atomes de carbone, ou bien un radical aryle non substitué ou substitué, contenant 6 à 10 atomes de carbone;  
 R<sub>3</sub> représente l'hydrogène, un groupe alkyle ou aryle ayant 7 à 10 atomes de carbone;  
 R<sub>4</sub> représente l'hydrogène ou un groupe alkyle ayant 1 à 6 atomes de carbone;  
 X représente un groupe amino-acyle choisi entre les groupes alanyl, valyl, leucyl, isoleucyl, α-aminobutyryl, thréonyle, méthionyle, cystéinyle, glutamyle, glutaminyle, aspartyle, phénylalanyle, tyrosyle, tryptophanyle, lysyle, ornithyle, arginyle, histidyle, asparaginyle, prolyle, hydroxyprolyle, séryle ou glycyle;  
 R<sub>5</sub> représente un groupe carboxyle facultativement estérifié ou amidé; et  
 R<sub>6</sub> représente un groupe carboxyle facultativement estérifié ou amidé;  
 le polymère séquencé possédant une base de polyoxypropylène ayant un poids moléculaire de 3250 à 4000 et la pourcentage de polyoxyéthylène dans la molécule totale étant compris dans l'intervalle de 0,2 à 20% (volume/volume); et  
 le sérum physiologique étant du sérum physiologique tamponné avec un phosphate.
5. Composition suivant la revendication 3, dans laquelle le glycopeptide de formule I est substitué de la manière suivante: R et R<sub>1</sub> sont identiques ou différents et sont choisis dans le groupe comprenant l'hydrogène et un radical acyle contenant 1 à 22 atomes de carbone; R<sub>2</sub> représente un groupe méthyle, R<sub>3</sub> représente l'hydrogène; X représente un groupe amino-acyle choisi entre les groupes L-séryle, L-alanyl, L-valyl, L-leucyl, L-isoleucyl, L-α-aminobutyryl, L-séryle, L-thréonyle, L-méthionyle, L-cystéinyle, L-phénylalanyle, L-tyrosyle, L-tryptophanyle, L-lysyle, L-ornithyle, L-arginyle, L-histidyle, L-glutamyle, L-glutaminyle, L-aspartyle, L-asparaginyle, L-prolyle et L-hydroxyprolyle, présent en une quantité de 0,01 à 1% (en poids/volume);  
 le copolymère séquencé possédant une base de polyoxypropylène ayant un poids moléculaire de 2250 à 4300, et le pourcentage de polyoxyéthylène dans la molécule totale étant compris dans l'intervalle de 5 à 20%, ledit copolymère étant présent en une quantité de 0,2 à 20% (en volume/volume); et  
 l'éther de glycol servant de surfactant étant un polysorbate présent en une quantité de 0,2 à 1% (en volume/volume).
6. Composition suivant la revendication 4, dans laquelle le glycopeptide est un composé de formule I dans laquelle R, R<sub>1</sub> et R<sub>3</sub> représentent l'hydrogène, R<sub>2</sub> représente un groupe méthyle, R<sub>4</sub> représente un groupe méthyle ou l'hydrogène; X représente un groupe L-valyl, L-alanyl, L-séryle, L-thréonyle ou L-α-aminobutyryl, R<sub>5</sub> représente un groupe carboxyle, carboxyle ou n-butylcarboxylate et R<sub>6</sub> représente un groupe carboxyl ou carbamoyl.
7. Composition suivant la revendication 5, dans laquelle le glycopeptide est la N-acétylmuramyl-L-

## EP 0 135 376 B1

thréonyl-D-isoglutamine, la N-acétyl-L- $\alpha$ -aminobutyryl-D-isoglutamine, la 6-O-stéaroyl-N-acétylmuramyl-L- $\alpha$ -aminobutyryl-D-isoglutamine ou la N-acétyldesméthylmuramyl-L-alanyl-D-isoglutamine, présente en une quantité de 0,01 à 1% (en poids/volume);

5 le copolymère séquencé est constitué d'une base de polyoxypropylène ayant un poids moléculaire de 4000, possède un pourcentage de polyoxyéthylène dans la molécule totale égal à 10% et est présent en une quantité de 0,2 à 5% (en volume/volume); et

le polysorbate est le TWEEN 80®.

7. Composition suivant la revendication 5, dans laquelle le glycopeptide est le n-butyl-N-acétylmuramyl-L-alanyl-D-glutamate, la N-acétylmuramyl-L-alanyl-D-isoglutamine, ou la N-acétylmuramyl-L-séryl-D-isoglutamine; et le copolymère est constitué d'une base de polyoxypropylène ayant un poids moléculaire de 4000 et qui contient 10% de polyoxyéthylène, en pourcentage de la molécule totale.

8. Composition suivant la revendication 6, dans laquelle le glycopeptide est la N-acétylmuramyl-L-thréonyl-D-isoglutamine, présente en une quantité de 0,5% (en poids/volume), et le copolymère est présent en une quantité de 2,5%.

9. Composition suivant l'une quelconque des revendications précédentes, qui comprend en outre une huile non toxique, métabolisable, en une quantité de 1 à 30% en volume (v/v).

10. Composition suivant l'une quelconque des revendications 4 à 8, qui comprend en outre du squalène ou du squalane présent en une quantité de 1 à 10% (v/v).

11. Composition de vaccin comprenant:

20 une quantité efficace d'antigène;

une quantité immunopotentialisatrice d'un glycopeptide immunostimulant;

un copolymère séquencé polyoxypropylène-polyoxyéthylène non toxique, en une quantité de 0,2 à 49% en volume (v/v);

un surfactant à base d'éther de glycol, en une quantité de 0,05 à 2,5% en volume (v/v); et

25 du sérum physiologique iso-osmotique tamponné, en une quantité suffisante pour compléter au volume.

12. Composition de vaccin suivant la revendication 11, comprenant en outre une huile non toxique métabolisable, en une quantité de 1 à 30% en volume (v/v).

13. Composition de vaccin comprenant une quantité efficace d'un antigène et une composition d'adjuvant suivant l'une quelconque des revendications 1 à 10.

30 14. Composition d'adjuvant suivant l'une quelconque des revendications 1 à 10, destinée à être utilisée pour accroître le pouvoir immunogène d'un antigène mélangé à cette composition sous forme d'une composition polyphasique destinée à être administrée à un oiseau ou un mammifère.

15. Composition d'adjuvant suivant la revendication 1, dans laquelle le glycopolypeptide comprend 35 0,01 à 2% (en poids/volume) de N-acétyl-muramyl-L-thréonyl-D-isoglutamine.

16. Composition d'adjuvant suivant la revendication 1 ou la revendication 15, qui comprend 2,5% (en poids/volume) d'un copolymère séquencé polyoxypropylène-polyoxyéthylène qui possède une base de polyoxypropylène ayant un poids moléculaire de 3250 à 4000 et dans lequel le polyoxyéthylène représente 10% de la molécule totale.

40 17. Composition d'adjuvant suivant la revendication 1, la revendication 15 ou la revendication 16, dans laquelle le surfactant à base d'éther de glycol comprend 0,2 à 1% (en poids/volume) d'un surfactant polyoxyéthylène 20-monooléate de sorbitanne.

45

50

55

60

65

REGISTER ENTRY FOR EP0135376 ✓

European Application No EP84305687.0 filing date 21.08.1984 ✓

Priority claimed:

22.08.1983 in United States of America - doc: 525190

Designated States BE CH DE FR GB IT LI LU NL SE AT

Title POLYOXYPROPYLENE-POLYOXYETHYLENE BLOCK POLYMER BASED ADJUVANTS AND  
VACCINES USING THEM

Applicant/Proprietor ✓

SYNTEX (U.S.A.) INC., 3401 Hillview Avenue, Palo Alto, California 94304,  
United States of America [ADP No. 50562982002]

Inventors

ANTHONY CLIFFORD ALLISON, 2513 Hastings Drive, Belmont California 94002,  
United States of America [ADP No. 53227351001]

NOELENE ELVA BYARS, 1091 Syracuse Drive, Sunnyvale California 94087,  
United States of America [ADP No. 53227369001]

Classified to

A5B  
A61K

Address for Service

MEWBURN ELLIS, 2 Cursitor Street, LONDON, EC4A 1BQ, United Kingdom  
[ADP No. 00000109001]

EPO Representative

IAN MICHAEL ARMITAGE, MEWBURN ELLIS & CO. 2/3 Cursitor Street, London EC4A  
1BQ, United Kingdom [ADP No. 50329846001]

Publication No EP0135376 dated 27.03.1985

Publication in English

Examination requested 21.08.1984

Patent Granted with effect from 07.11.1990 (Section 25(1)) with title  
POLYOXYPROPYLENE-POLYOXYETHYLENE BLOCK POLYMER BASED ADJUVANTS AND  
VACCINES USING THEM.

---

24.12.1986 EPO: Search report published on 30.12.1986

Entry Type 25.11 Staff ID.

Auth ID. EPT

01.10.1990 Notification from EPO of change of Applicant/Proprietor details  
from

SYNTEX (U.S.A.) INC., 3401 Hillview Avenue, Palo Alto, California  
94304, United States of America [ADP No. 50562982002]

to

SYNTEX (U.S.A.) INC., 3401 Hillview Avenue, Palo Alto California  
94304, United States of America [ADP No. 50562982003]

Entry Type 25.14 Staff ID. RD06 Auth ID. EPT

TIMED: 02/07/92 12:00:44

REGISTER ENTRY FOR EP0135376 (Cont.)

PAGE: 2

08.10.1990 MEWBURN ELLIS, 2 Cursitor Street, LONDON, EC4A 1BQ, United Kingdom  
[ADP No. 00000109001]

registered as address for service

Entry Type 8.11 Staff ID. IW1 Auth ID. AA

\*\*\*\* END OF REGISTER ENTRY \*\*\*\*

OA80-01  
EP

OPTICS - PATENTS

02/07/92 12:01:55  
PAGE: 1

RENEWAL DETAILS

PUBLICATION NUMBER

EP0135376

PROPRIETOR(S)

SYNTEX (U.S.A.) INC., 3401 Hillview Avenue, Palo Alto California  
94304, United States of America

DATE FILED

21.08.1984

DATE GRANTED

07.11.1990

DATE NEXT RENEWAL DUE

21.08.1992

DATE NOT IN FORCE

DATE OF LAST RENEWAL

08.08.1991

YEAR OF LAST RENEWAL

08

STATUS

PATENT IN FORCE