





US 20140113005A1

(19) **United States**(12) **Patent Application Publication**  
**Charrier et al.**(10) **Pub. No.: US 2014/0113005 A1**(43) **Pub. Date: Apr. 24, 2014**(54) **COMPOUNDS USEFUL AS INHIBITORS OF  
ATR KINASE**(71) Applicant: **Vertex Pharmaceuticals Incorporated,**  
(US)(72) Inventors: **Jean-Damien Charrier**, Grove (GB);  
**Steven John Durrant**, Abingdon (GB);  
**David Kay**, Purton (GB); **Michael**  
**O'Donnell**, Abingdon (GB); **Ronald**  
**Marcellus Alphonsus Knegt**,  
Abingdon (GB); **Somhairle**  
**MacCormick**, Abingdon (GB); **Joanne**  
**Pinder**, Didcot (GB); **Stephen Clinton**  
**Young**, Oxford (GB); **Haley Marie**  
**Binch**, Encinitas, CA (US); **Thomas**  
**Cleveland**, San Diego, CA (US); **Lev**  
**Tyler Dewey Fanning**, San Marcos, CA  
(US); **Dennis James Hurley**, San  
Marcos, CA (US); **Pramod Joshi**, San  
Diego, CA (US); **Urvi Jagdishbhai**  
**Sheth**, San Diego, CA (US); **Alina**  
**Silina**, San Diego, CA (US); **Philip**  
**Michael Reaper**, Shillingford (GB);  
**Anisa Nizarali Virani**, Thatcham (GB)(73) Assignee: **Vertex Pharmaceuticals Incorporated,**  
Cambridge, MA (US)(21) Appl. No.: **13/742,948**(22) Filed: **Jan. 16, 2013****Related U.S. Application Data**(60) Division of application No. 13/104,291, filed on Oct.  
27, 2011, now Pat. No. 8,410,112, which is a continu-  
ation of application No. PCT/US2009/063922, filed  
on Nov. 10, 2009.(60) Provisional application No. 61/163,655, filed on Mar.  
26, 2009, provisional application No. 61/146,463,  
filed on Jan. 22, 2009, provisional application No.  
61/114,204, filed on Nov. 13, 2008, provisional appli-  
cation No. 61/112,906, filed on Nov. 10, 2008.**Publication Classification**(51) **Int. Cl.**  
**C07D 403/14** (2006.01)  
**A61K 31/497** (2006.01)  
**C07D 403/04** (2006.01)  
**A61K 31/541** (2006.01)**C07D 417/14** (2006.01)  
**A61K 45/06** (2006.01)  
**C07D 413/04** (2006.01)  
**C07D 241/28** (2006.01)  
**C07D 405/14** (2006.01)  
**C07D 471/04** (2006.01)  
**C07D 491/113** (2006.01)  
**C07D 409/04** (2006.01)  
**A61K 31/551** (2006.01)  
**C07D 417/04** (2006.01)(52) **U.S. Cl.**CPC ..... **C07D 403/14** (2013.01); **A61K 31/551**  
(2013.01); **A61K 31/497** (2013.01); **C07D**  
**403/04** (2013.01); **A61K 31/541** (2013.01);  
**C07D 417/14** (2013.01); **C07D 417/04**  
(2013.01); **C07D 413/04** (2013.01); **C07D**  
**241/28** (2013.01); **C07D 405/14** (2013.01);  
**C07D 471/04** (2013.01); **C07D 491/113**  
(2013.01); **C07D 409/04** (2013.01); **A61K**  
**45/06** (2013.01)USPC ... **424/649**; 514/218; 514/255.05; 514/233.8;  
514/252.11; 514/228.2; 514/236.2; 514/188;  
514/34; 514/49; 514/47; 514/2.9(57) **ABSTRACT**The present disclosure relates to pyrazine compounds of for-  
mula I:

wherein L, n, R<sup>1</sup>, and R<sup>2</sup> are as described in the specification. These compounds are useful as inhibitors of ATR protein kinase. The disclosure also relates to pharmaceutically acceptable compositions comprising the compounds of the disclosure; methods of treating of various diseases, disorders, and conditions using the compounds of the disclosure; processes for preparing the compounds of the disclosure; intermediates for the preparation of the compounds of the disclosure; and methods of using the compounds in in vitro applications, such as the study of kinases in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such kinases; and the comparative evaluation of new kinase inhibitors.

FIGURE 1



## COMPOUNDS USEFUL AS INHIBITORS OF ATR KINASE

### BACKGROUND OF THE INVENTION

[0001] ATR (“ATM and Rad3 related”) kinase is a protein kinase involved in cellular responses to DNA damage. ATR kinase acts with ATM (“ataxia telangiectasia mutated”) kinase and many other proteins to regulate a cell’s response to DNA damage, commonly referred to as the DNA Damage Response (“DDR”). The DDR stimulates DNA repair, promotes survival and stalls cell cycle progression by activating cell cycle checkpoints, which provide time for repair. Without the DDR, cells are much more sensitive to DNA damage and readily die from DNA lesions induced by endogenous cellular processes such as DNA replication or exogenous DNA damaging agents commonly used in cancer therapy.

[0002] Healthy cells can rely on a host of different proteins for DNA repair including the DDR kinase ATR. In some cases these proteins can compensate for one another by activating functionally redundant DNA repair processes. On the contrary, many cancer cells harbour defects in some of their DNA repair processes, such as ATM signaling, and therefore display a greater reliance on their remaining intact DNA repair proteins which include ATR. In addition, many cancer cells express activated oncogenes or lack key tumour suppressors, and this can make these cancer cells prone to dysregulated phases of DNA replication which in turn cause DNA damage.

[0003] ATR has been implicated as a critical component of the DDR in response to disrupted DNA replication. As a result, these cancer cells are more dependent on ATR activity for survival than healthy cells. Accordingly, ATR inhibitors may be useful for cancer treatment, either used alone or in combination with DNA damaging agents, because they shut down a DNA repair mechanism that is more important for cellular survival in many cancer cells than in healthy normal cells. In fact, ATR inhibition has been shown to be effective in cancer cells as single agents and as potent sensitizers to radiotherapy and genotoxic chemotherapy.

[0004] Additionally, ATR is also implicated in other diseases, such as HIV1: (see Ward et al, 2009 PLOS Pathogens 5:e1000613, Lai et al, 2005 J Virol 79:1544, and Daniel et al 2003 PNAS 100:4778); Hepatitis B (see e.g., Zhao et al, 2008 World J Gastroenterol 14:6163, Wang et al 2008 JBC 283: 25455); Adenovirus (see e.g., Nichols et al, 2009 J Virol 83: 5987), and Psoriasis (see e.g., Derheimer et al., 2009 Mol Pharmacol 75:599).

[0005] Accordingly, there is a need for the development of potent and selective ATR inhibitors to treat these diseases. More specifically, there is a need for the development of potent and selective ATR inhibitors for the treatment of cancer, either as single agents or as combination therapies with radiotherapy or genotoxic chemotherapy.


### DESCRIPTION OF THE FIGURES

[0006] FIG. 1: A pictorial representation illustrating the position of the hydrogen bond acceptor of J. The hydrogen bond acceptor is located in a sphere wherein the center of the sphere is located 6 Å from C<sup>5</sup> on the line defined by the C<sup>2</sup>-C<sup>5</sup> axis in the direction away from C<sup>2</sup> and the radius of the sphere is 4 Å;

### SUMMARY OF THE INVENTION

[0007] The present invention relates to pyrazine compounds useful as inhibitors of ATR protein kinase. The invention also relates to pharmaceutically acceptable compositions comprising the compounds of this invention; methods of treating various diseases, disorders, and conditions using the compounds of this invention; processes for preparing the compounds of this invention; intermediates for the preparation of the compounds of this invention; and methods of using the compounds in in vitro applications, such as the study of kinases in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such kinases; and the comparative evaluation of new kinase inhibitors.


[0008] The compounds of this invention have formula I:



wherein the variables are as defined herein.

### DETAILED DESCRIPTION OF THE INVENTION

[0009] One aspect of this invention provides a compound of formula I:



or a pharmaceutically acceptable salt thereof wherein:

[0010] R<sup>1</sup> is a 5-6 membered monocyclic aryl or heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein said monocyclic aryl or heteroaryl ring is optionally fused to another ring to form a 8-10 membered bicyclic aryl or heteroaryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R<sup>1</sup> is optionally substituted with 1-5 J<sup>1</sup> groups;

[0011] C<sup>2</sup>, C<sup>3</sup> and C<sup>5</sup> are carbon;

[0012] R<sup>2</sup> is -Q or -Q-Q<sup>1</sup>;

[0013] Q is a 3-7 membered monocyclic saturated or partially unsaturated non-aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; Q can be optionally fused to Q<sup>1</sup>; each Q is independently and optionally substituted with 1-4 J<sup>2</sup> groups; Q is substituted with one to two occurrences of J or contains J as a ring member;

[0014] Q<sup>1</sup> is a 5-8 membered monocyclic saturated or unsaturated ring having 0-4 heteroatoms independently

selected from nitrogen, oxygen, or sulfur; each  $Q^1$  is independently and optionally substituted with 1-4  $J^{Q1}$  groups;

[0015] J is fluoro or a moiety containing a hydrogen bond acceptor:

[0016] wherein the hydrogen bond acceptor is located in a sphere wherein

[0017] a) the center of the sphere is located 6 Å from  $C^5$  on the line defined by the  $C^2$ - $C^5$  axis in the direction away from  $C^2$  and

[0018] b) the radius of the sphere is 4 Å;

[0019] L is  $-C(O)NH-$ ;

[0020] n is 0 or 1;

[0021] Each  $J^1$ ,  $J^2$ , and  $J^{Q1}$  is independently halo,  $-CN$ ,  $-NO_2$ ,  $V-R$ , or  $-(V)_m-Q^2$ ;

[0022] V is a  $C_{1-10}$  aliphatic chain wherein 0-3 methylene units are optionally replaced with oxygen, nitrogen, sulfur, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of  $J^V$ ;

[0023] m is 0 or 1;

[0024]  $Q^2$  is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $Q^2$  is optionally substituted with 1-5  $J^{Q2}$ ;

[0025] each  $J^V$  is independently  $NH_2$ ,  $NH(C_{1-4} \text{aliphatic})$ ,  $N(C_{1-4} \text{aliphatic})_2$ , halogen,  $C_{1-4} \text{aliphatic}$ , OH,  $O(C_{1-4} \text{aliphatic})$ ,  $NO_2$ , CN,  $CO_2H$ ,  $CO_2(C_{1-4} \text{aliphatic})$ ,  $O(haloC_{1-4} \text{aliphatic})$ , or  $haloC_{1-4} \text{aliphatic}$ ;

[0026] each  $J^{Q2}$  is independently halogen,  $NO_2$ , CN, or  $C_{1-6} \text{aliphatic}$  wherein up to 1 methylene unit is optionally replaced with  $N(R)_2$ , OR, SR, COR,  $CO_2R$ ,  $CON(R)_2$ , SOR,  $SO_2R$ ,  $SO_2N(R)_2$ , OCOR, NRCOR, NRCOOR, NRSOR,  $NRSO_2R$ ,  $NRSO_2N(R)_2$ ,  $OCON(R)_2$ , or  $NRCON(R)_2$ ; wherein said  $C_{1-6} \text{aliphatic}$  is optionally substituted with 1-4 substituents selected from  $NH_2$ ,  $NH(C_{1-4} \text{aliphatic})$ ,  $N(C_{1-4} \text{aliphatic})_2$ , halogen,  $C_{1-4} \text{aliphatic}$ , OH,  $O(C_{1-4} \text{aliphatic})$ ,  $NO_2$ , CN,  $CO_2H$ ,  $CO_2(C_{1-4} \text{aliphatic})$ ,  $O(haloC_{1-4} \text{aliphatic})$ , or  $haloC_{1-4} \text{aliphatic}$ ;

[0027] R is H or  $C_{1-4} \text{alkyl}$  wherein said  $C_{1-6} \text{alkyl}$  is optionally substituted with 1-4 halo.

[0028] In one aspect of the invention, Q is substituted with one occurrence of J. In some embodiments, J is oxo or  $(V^1)-R''$ ;  $V^1$  is  $C_{1-6} \text{aliphatic}$  chain wherein up to three methylene unit of the aliphatic chain may be optionally replaced with  $-NR'$ ,  $-O-$ ,  $-S-$ ,  $-C(O)-$ ,  $-S(O)-$ , or  $-S(O)_2-$  wherein the first or second methylene group away from the point of attachment is replaced with CO, SO,  $SO_2$ , S, or O;  $R'$  is H or  $C_{1-4} \text{alkyl}$ ; and  $R''$  is H or a 5-6 membered monocyclic ring containing 0-2 heteroatoms selected from O, N, or S; wherein said  $R''$  is optionally substituted with 1-3 occurrences of halo,  $C_{1-3} \text{alkyl}$ , CN, OH,  $O(C_{1-3} \text{alkyl})$ ,  $NH_2$ ,  $NH(C_{1-3} \text{alkyl})$ ,  $N(C_{1-3} \text{alkyl})_2$ , or acetyl. In some embodiments, up to 2 methylene units of the aliphatic chain are optionally replaced with heteroatom selected from O, N, or S.

[0029] In another aspect of the invention, Q contains J as a ring member. In some embodiment, J is selected from sulfur, nitrogen, or oxygen. Examples of such Q rings include, but are not limited to, morpholinyl and thiomorpholinyl.

[0030] In some embodiments, n is 1; in other embodiments, n is 0.


[0031] In some embodiments,  $R^1$  is a monocyclic 6-membered ring, such as phenyl or pyridyl. In other embodiments,

$R^1$  is a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered bicyclic heteroaryl ring having 1-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $R^1$  is optionally substituted with 1-5  $J^1$  groups.

[0032] In some embodiments, n is 1 and  $R^1$  is a monocyclic 6-membered ring. In some embodiments, said 6-membered ring is phenyl or pyridyl.


[0033] In other embodiments, n is 0 and  $R^1$  is a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered bicyclic heteroaryl ring having 1-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $R^1$  is optionally substituted with 1-5  $J^1$  groups. In some embodiments,  $R^1$  is a monocyclic 5-membered heteroaryl ring containing 1-3 heteroatoms selected from oxygen, nitrogen, or sulfur; or  $R^1$  is a 5-membered heteroaryl ring containing 1-2 heteroatoms selected from oxygen, nitrogen, or sulfur fused to a 6-membered aryl or heteroaryl ring containing 0-4 nitrogen atoms.


[0034] In some embodiments,  $R^1$  is bonded to  $C^3$  via a carbon atom and wherein  $R^1$  contains at least one ring heteroatom. In some embodiments, said ring heteroatom (shown as G) is located next to the carbon atom bonded to  $C^3$  as shown below in formula II:



II


[0035] In some embodiments,  $R^1$  is a 5-membered ring optionally fused to another ring. In some embodiments,  $R^1$  is a monocyclic 5-membered ring containing 1-3 heteroatoms selected from oxygen, nitrogen, or sulfur. In some embodiments,  $R^1$  is a 5-membered ring and is selected from the following:






wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ .


[0036] In some embodiments,  $R^1$  is



[0037] In other embodiments,  $R^1$  is a 5-membered ring fused to another ring. In some embodiments,  $R^1$  is selected from the following:




wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ ; A is carbon or nitrogen, provided that at least two occurrences of A are carbon; and  $X_1$  is O, N, or S. In some embodiments,  $R^1$  is




wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ ; A is carbon; and  $X_1$  is selected from O, N, or S.

[0038] In another aspect of this invention,  $R^2$  can be -Q or -Q- $Q^1$ . In one embodiment,  $R^2$  is Q (i.e. monocyclic ring). In some embodiments, Q is a 6 membered ring and can contain


0, 1, or 2 double bonds. In some embodiments, Q is  $C_{3-7}$  cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl, homopiperidinyl, morpholinyl, or thiomorpholinyl. In some embodiments, Q is selected from the following:




[0039] In some embodiments, Q is




In other embodiments, Q is



In some embodiments, when J is oxo,  $R^2$  is




[0040] In another aspect of this invention, J is bonded in the position of Q as shown below:



[0041] In some embodiments, J is a  $C_{1-6}$  aliphatic chain wherein up to 2 methylene units of  $C_{1-6}$  aliphatic are optionally replaced with a heteroatom selected from O, N, or S. In other embodiments, J is selected from oxo, —OH, —O( $C_{1-6}$  aliphatic), CN,  $CH_2CN$ , —C(O)( $C_{1-6}$  aliphatic), —C(O)(phenyl), —C(O)(benzyl), —C(O)CH<sub>2</sub>O(benzyl), —C(O)(pyridyl), —C(O)(pyrrolidinyl), —C(O)(piperidinyl), —C(O)(piperazinyl), —C(O)(morpholinyl), —C(O)(tetrahydropyranyl), —C(O)O( $C_{1-6}$  aliphatic), —C(O)O(phenyl), —C(O)O(benzyl), —C(O)NH<sub>2</sub>, —C(O)NH( $C_{1-6}$  aliphatic), —C(O)NH( $C_{1-6}$  aliphatic)<sub>2</sub>, —C(O)NH(phenyl), —C(O)NH(benzyl), —S(O)<sub>2</sub>( $C_{1-6}$  aliphatic), —S(O)<sub>2</sub>(phenyl), —S(O)<sub>2</sub>(benzyl), —S(O)<sub>2</sub>(pyridyl), —S(O)<sub>2</sub>(furanyl), —S(O)<sub>2</sub>(imidazole), —S(O)<sub>2</sub>(thienyl), —S(O)<sub>2</sub>NH( $C_{1-6}$  aliphatic), —S(O)<sub>2</sub>N( $C_{1-6}$  aliphatic)<sub>2</sub>, —S(O)<sub>2</sub>NH(phenyl), —S(O)<sub>2</sub>NH(benzyl), —NHC(O)( $C_{1-6}$  aliphatic); wherein said phenyl, benzyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, or  $C_{1-6}$  aliphatic is optionally substituted with halo,  $C_{1-3}$  alkyl, CN, OH, O( $C_{1-3}$  alkyl), NH<sub>2</sub>, NH( $C_{1-3}$  alkyl), N( $C_{1-3}$  alkyl)<sub>2</sub>, acetyl. In yet other embodiments, J is selected from oxo, —OH, —O( $C_{1-4}$  alkyl), —C(O)( $C_{1-4}$  alkyl), —C(O)NH<sub>2</sub>, —C(O)NH( $C_{1-4}$  alkyl), —C(O)NH( $C_{1-4}$  alkyl)<sub>2</sub>, —S(O)<sub>2</sub>( $C_{1-4}$  alkyl), —S(O)<sub>2</sub>NH( $C_{1-4}$  alkyl), or —S(O)<sub>2</sub>N( $C_{1-4}$  alkyl)<sub>2</sub>.

[0042] Another embodiment provides a compound of formula I:



or a pharmaceutically acceptable salt thereof wherein:

[0043] R<sup>1</sup> is a 5-6 membered monocyclic aryl or heteroaryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein said monocyclic aryl or heteroaryl ring is optionally fused to another ring to form a 8-10 membered bicyclic aryl or heteroaryl ring having

0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R<sup>1</sup> is optionally substituted with 1-5 J<sup>1</sup> groups;

[0044] C<sup>2</sup>, C<sup>3</sup> and C<sup>5</sup> are carbon;

[0045] R<sup>2</sup> is —Q or —Q-Q<sup>1</sup>;

[0046] Q is a 3-7 membered monocyclic saturated or unsaturated non-aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q is independently and optionally substituted with 1-4 J<sup>Q</sup> groups; Q is substituted with zero to two occurrences of J or contains J as a ring member;

[0047] Q is optionally fused to Q<sup>1</sup> to form a fused bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup> are optionally joined together at a carbon atom to form a spirocyclic bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup>, taken together, form a bridged bicyclic ring Q-Q<sup>1</sup> wherein said bridge is 1-3 atoms long; Q-Q<sup>1</sup> can be optionally substituted with 1-2 J groups; wherein J can be bonded to either Q or Q<sup>1</sup>;

[0048] Q<sup>1</sup> is a 3-8 membered monocyclic saturated or unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>1</sup> is independently and optionally substituted with 1-4 J<sup>Q1</sup> groups;

[0049] J is fluoro, oxo, or a moiety containing a hydrogen bond acceptor:

[0050] wherein the hydrogen bond acceptor is located in a sphere wherein

[0051] a) the center of the sphere is located 6 Å from C<sup>5</sup> on the line defined by the C<sup>2</sup>-C<sup>5</sup> axis in the direction away from C<sup>2</sup> and

[0052] b) the radius of the sphere is 4 Å;

[0053] L is —C(O)NH— or —C(O)N( $C_{1-6}$  alkyl)-;

[0054] n is 0 or 1;

[0055] each J<sup>Q</sup> and J<sup>Q1</sup> is independently halo, oxo, —CN, —NO<sub>2</sub>, V—R, or —(V)<sub>m</sub>-Q<sup>2</sup>;

[0056] J<sup>1</sup> is halo, —CN, —NO<sub>2</sub>, V—R, or —(V<sup>2</sup>)<sub>m</sub>-Q<sup>3</sup>;

[0057] V is a  $C_{1-10}$  aliphatic chain wherein 0-3 methylene units are optionally and independently replaced with oxygen, nitrogen, sulfur, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of J<sup>V</sup>;

[0058] V<sup>2</sup> is a  $C_{1-10}$  aliphatic chain wherein 0-3 methylene units are optionally and independently replaced with oxygen, nitrogen, sulfur, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of J<sup>V2</sup>;

[0059] m is 0 or 1;


[0060] Q<sup>2</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>2</sup> is optionally substituted with 1-5 J<sup>Q2</sup>;

[0061] Q<sup>3</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>3</sup> is optionally substituted with 1-5 J<sup>Q3</sup>;


[0062] each J<sup>V</sup> and J<sup>V2</sup> is independently halogen, CN, NH<sub>2</sub>, NO<sub>2</sub>,  $C_{1-4}$  aliphatic, NH( $C_{1-4}$  aliphatic), N( $C_{1-4}$  aliphatic)<sub>2</sub>, OH, O( $C_{1-4}$  aliphatic), CO<sub>2</sub>H, CO<sub>2</sub>( $C_{1-4}$  aliphatic), C(O)NH<sub>2</sub>, C(O)NH( $C_{1-4}$  aliphatic), C(O)N( $C_{1-4}$  aliphatic)<sub>2</sub>, NHCO( $C_{1-4}$  aliphatic), N( $C_{1-4}$  aliphatic)CO( $C_{1-4}$  aliphatic), SO<sub>2</sub>( $C_{1-4}$  aliphatic), NHSO<sub>2</sub>( $C_{1-4}$  aliphatic), or

- $N(C_{1-4}\text{aliphatic})SO_2(C_{1-4}\text{aliphatic})$ , wherein said  $C_{1-4}\text{aliphatic}$  is optionally substituted with halo;
- [0063] each  $J^{Q2}$  and  $J^{Q3}$  is independently halo, oxo, CN,  $NO_2X-R$ , or  $-(X)_p-Q^4$ ;
- [0064]  $p$  is 0 or 1;
- [0065]  $X$  is  $C_{1-10}\text{aliphatic}$ ; wherein 1-3 methylene units of said  $C_{1-6}\text{aliphatic}$  are optionally replaced with  $-NR$ ,  $-O-$ ,  $-S-$ ,  $C(O)$ ,  $S(O)_2$ , or  $S(O)$ ; wherein  $X$  is optionally and independently substituted with 1-4 occurrences of  $NH_2$ ,  $NH(C_{1-4}\text{aliphatic})$ ,  $N(C_{1-4}\text{aliphatic})_2$ , halogen,  $C_{1-4}\text{aliphatic}$ ,  $OH$ ,  $O(C_{1-4}\text{aliphatic})$ ,  $NO_2$ ,  $CN$ ,  $CO_2H$ ,  $CO_2(C_{1-4}\text{aliphatic})$ ,  $C(O)NH_2$ ,  $C(O)NH(C_{1-4}\text{aliphatic})$ ,  $C(O)N(C_{1-4}\text{aliphatic})_2$ ,  $SO(C_{1-4}\text{aliphatic})$ ,  $SO_2(C_{1-4}\text{aliphatic})$ ,  $SO_2NH(C_{1-4}\text{aliphatic})$ ,  $SO_2NH(C_{1-4}\text{aliphatic})_2$ ,  $NHC(O)(C_{1-4}\text{aliphatic})$ ,  $N(C_{1-4}\text{aliphatic})C(O)(C_{1-4}\text{aliphatic})$ , wherein said  $C_{1-4}\text{aliphatic}$  is optionally substituted with 1-3 occurrences of halo;
- [0066]  $Q^4$  is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $Q^4$  is optionally substituted with 1-5  $J^{Q4}$ ;  $J^{Q4}$  is halo, CN, or  $C_{1-4}\text{alkyl}$  wherein up to 2 methylene units are optionally replaced with  $O$ ,  $N$ ,  $S$ ,  $C(O)$ ,  $S(O)$ , or  $S(O)_2$ ;
- [0067]  $R$  is  $H$  or  $C_{1-4}\text{alkyl}$  wherein said  $C_{1-4}\text{alkyl}$  is optionally substituted with 1-4 halo.
- [0068] In some embodiments,
- [0069]  $R^1$  is a 5-6 membered monocyclic aryl or heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein said monocyclic aryl or heteroaryl ring is optionally fused to another ring to form a 8-10 membered bicyclic aryl or heteroaryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;  $R^1$  is optionally substituted with 1-5  $J^1$  groups;
- [0070]  $Q$  is a 3-7 membered monocyclic saturated or partially unsaturated non-aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;  $Q$  can be optionally fused to  $Q^1$ ; each  $Q$  is independently and optionally substituted with 1-4  $J^Q$  groups;  $Q$  is substituted with one to two occurrences of  $J$  or contains  $J$  as a ring member;
- [0071]  $Q^1$  is a 5-8 membered monocyclic saturated or unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $Q^1$  is independently and optionally substituted with 1-4  $J^{Q1}$  groups;
- [0072]  $J$  is fluoro or a moiety containing a hydrogen bond acceptor;
- [0073]  $L$  is  $-C(O)NH-$ ;
- [0074] each  $J^Q$  and  $J^{Q1}$  is independently halo,  $-CN$ ,  $-NO_2$ ,  $V-R$ , or  $-(V)_m-Q^2$ ;
- [0075] each  $J^{Q2}$  and  $J^{Q3}$  is independently halogen,  $NO_2$ ,  $CN$ , or  $C_{1-6}\text{aliphatic}$  wherein up to 1 methylene unit is optionally replaced with  $N(R)_2$ ,  $OR$ ,  $SR$ ,  $COR$ ,  $CO_2R$ ,  $CON(R)_2$ ,  $SOR$ ,  $SO_2R$ ,  $SO_2N(R)_2$ ,  $OCOR$ ,  $NRCOR$ ,  $NRCOOR$ ,  $NRSOR$ ,  $NRSO_2R$ ,  $NRSO_2N(R)_2$ ,  $OCON(R)_2$ , or  $NRCON(R)_2$ ; wherein said  $C_{1-6}\text{aliphatic}$  is optionally substituted with 1-4 substituents selected from  $NH_2$ ,  $NH(C_{1-4}\text{aliphatic})$ ,  $N(C_{1-4}\text{aliphatic})_2$ , halogen,  $C_{1-4}\text{aliphatic}$ ,  $OH$ ,  $O(C_{1-4}\text{aliphatic})$ ,  $NO_2$ ,  $CN$ ,  $CO_2H$ ,  $CO_2(C_{1-4}\text{aliphatic})$ ,  $O(\text{halo}C_{1-4}\text{aliphatic})$ , or halo  $C_{1-4}\text{aliphatic}$ ;


- [0076] each  $J^V$  is independently  $NH_2$ ,  $NH(C_{1-4}\text{aliphatic})$ ,  $N(C_{1-4}\text{aliphatic})_2$ , halogen,  $C_{1-4}\text{aliphatic}$ ,  $OH$ ,  $O(C_{1-4}\text{aliphatic})$ ,  $NO_2$ ,  $CN$ ,  $CO_2H$ ,  $CO_2(C_{1-4}\text{aliphatic})$ ,  $O(\text{halo}C_{1-4}\text{aliphatic})$ , or halo  $C_{1-4}\text{aliphatic}$ .
- [0077] In other embodiments,  $J$  is halo, oxo,  $-CN$ ,  $-NO_2$ ,  $V^1-R''$ , or  $-(V)_m-R^*$ . In other embodiments, the moiety containing a hydrogen-bond acceptor is selected from oxo or  $(V^1)-R''$ .
- [0078] In some embodiments,
- [0079]  $J$  is oxo or  $(V^1)-R''$ ;
- [0080]  $V^1$  is  $C_{1-6}\text{aliphatic}$  chain wherein up to three methylene unit of the aliphatic chain may be optionally replaced with  $-NR'$ ,  $-O-$ ,  $-S-$ ,  $-C(O)-$ ,  $-S(O)-$ , or  $-S(O)_2-$ ; wherein the first or second methylene group away from the point of attachment is replaced with  $CO$ ,  $SO$ ,  $SO_2$ ,  $S$ , or  $O$ ;
- [0081]  $R'$  is  $H$  or  $C_{1-4}\text{alkyl}$ ;
- [0082]  $R''$  is  $H$  or a 5-6 membered monocyclic ring containing 0-2 heteroatoms selected from  $O$ ,  $N$ , or  $S$ ; wherein said  $R''$  is optionally substituted with 1-3 occurrences of halo,  $C_{1-3}\text{alkyl}$ ,  $CN$ ,  $OH$ ,  $O(C_{1-3}\text{alkyl})$ ,  $NH_2$ ,  $NH(C_{1-3}\text{alkyl})$ ,  $N(C_{1-3}\text{alkyl})_2$ , or acetyl.
- [0083] In other embodiments,  $R^1$  is



- [0084] In some embodiments,  $Q$  is  $C_{3-7}\text{cycloalkyl}$ , pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl, homopiperidinyl, morpholinyl, or thiomorpholinyl. In some embodiments,  $Q$  is



- [0085] In other embodiments,  $J$  is  $C_{1-6}\text{aliphatic}$  chain wherein up to 2 methylene units of  $C_{1-6}\text{aliphatic}$  optionally replaced with heteroatom selected from  $O$ ,  $N$ , or  $S$ .
- [0086] Another embodiment provides a compound of formula Ia:



Ia

or a pharmaceutically acceptable salt thereof wherein:

- [0087]  $R^1$  is a 5-6 membered monocyclic aryl or heteroaryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein said monocyclic aryl or heteroaryl ring is optionally fused to another ring to form

- a 8-10 membered bicyclic aryl or heteroaryl ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R<sup>1</sup> is optionally substituted with 1-5 J<sup>1</sup> groups;
- [0088] R<sup>2</sup> is -Q or -Q-Q<sup>1</sup>;
- [0089] Q is a 3-7 membered monocyclic saturated or unsaturated non-aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; Q is optionally substituted with 1-4 J<sup>Q</sup> groups; Q is optionally substituted with 1-2 J groups;
- [0090] Q<sup>1</sup> is a 3-8 membered monocyclic saturated or unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>1</sup> is independently and optionally substituted with 1-4 J<sup>Q1</sup> groups;
- [0091] Q is optionally fused to Q<sup>1</sup> to form a fused bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup> are optionally joined together at a carbon atom to form a spirocyclic bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup>, taken together, form a bridged bicyclic ring Q-Q<sup>1</sup> wherein said bridge is 1-3 atoms long; Q-Q<sup>1</sup> can be optionally substituted with 1-2 J groups; wherein J can be bonded to either Q or Q<sup>1</sup>;
- [0092] each J<sup>Q</sup> and J<sup>Q1</sup> is independently halo, oxo, -CN, -NO<sub>2</sub>, V-R, or -(V)<sub>m</sub>-Q<sup>2</sup>;
- [0093] J is halo, oxo, -CN, -NO<sub>2</sub>, V<sup>1</sup>-R'', or -(V)<sub>m</sub>-R\*;
- [0094] L is -C(O)NH- or -C(O)N(C<sub>1-6</sub>alkyl)-;
- [0095] n is 0 or 1;
- [0096] J<sup>1</sup> is halo, -CN, -NO<sub>2</sub>, V-R, or -(V<sup>2</sup>)<sub>m</sub>-Q<sup>3</sup>;
- [0097] V is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally replaced with -NR-, -O-, -S-, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of J<sup>V</sup>;
- [0098] V<sup>1</sup> is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally replaced with -NR'-, -O-, -S-, C(O), S(O), or S(O)<sub>2</sub>; V<sup>1</sup> is optionally substituted with 1-6 occurrences of J<sup>V1</sup>;
- [0099] V<sup>2</sup> is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally and independently replaced with -NR-, -O-, -S-, C(O), S(O), or S(O)<sub>2</sub>; V<sup>2</sup> is optionally substituted with 1-6 occurrences of J<sup>V2</sup>;
- [0100] m is 0 or 1;
- [0101] Q<sup>2</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>2</sup> is optionally substituted with 1-5 J<sup>Q2</sup>;
- [0102] Q<sup>3</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>3</sup> is optionally substituted with 1-5 J<sup>Q3</sup>;
- [0103] R'' is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each R'' is optionally substituted with 1-5 J<sup>1</sup>;
- [0104] each J<sup>V</sup>, J<sup>V1</sup>, and J<sup>V2</sup> is independently halogen, CN, NH<sub>2</sub>, NO<sub>2</sub>, C<sub>1-4</sub>aliphatic, NH(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, OH, O(C<sub>1-4</sub>aliphatic), CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-4</sub>aliphatic), C(O)NH<sub>2</sub>, C(O)NH(C<sub>1-4</sub>aliphatic), C(O)N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, NHCO(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)CO(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), NHSO<sub>2</sub>(C<sub>1-4</sub>aliphatic), or N(C<sub>1-4</sub>aliphatic)SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), wherein said C<sub>1-4</sub>aliphatic is optionally substituted with halo;
- [0105] each J<sup>Q2</sup>, J<sup>Q3</sup>, and J<sup>1</sup> is independently halo, oxo, CN, NO<sub>2</sub>, X-R, or -(X)<sub>p</sub>-Q<sup>4</sup>;
- [0106] p is 0 or 1;
- [0107] X is a C<sub>1-10</sub>aliphatic; wherein 1-3 methylene units of said C<sub>1-6</sub>aliphatic are optionally replaced with -NR-, -O-, -S-, C(O), S(O)<sub>2</sub>, or S(O); wherein X is optionally and independently substituted with 1-4 occurrences of NH<sub>2</sub>, NH(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, halogen, C<sub>1-4</sub>aliphatic, OH, O(C<sub>1-4</sub>aliphatic), NO<sub>2</sub>, CN, CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-4</sub>aliphatic), C(O)NH<sub>2</sub>, C(O)NH(C<sub>1-4</sub>aliphatic), C(O)N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, SO(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>NH(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, NHC(O)(C<sub>1-4</sub>aliphatic), or N(C<sub>1-4</sub>aliphatic)C(O)(C<sub>1-4</sub>aliphatic), wherein said C<sub>1-4</sub>aliphatic is optionally substituted with 1-3 occurrences of halo;
- [0108] Q<sup>4</sup> is a 3-6 membered monocyclic ring containing 0-2 heteroatoms selected from O, N, or S;
- [0109] each Q<sup>4</sup> is optionally substituted with halo, CN, or C<sub>1-4</sub>alkyl wherein up to 2 methylene units are optionally replaced with O, N, S, C(O), S(O), or S(O)<sub>2</sub>;
- [0110] each R\*, R', and R is independently H or C<sub>1-4</sub>alkyl wherein said C<sub>1-4</sub>alkyl is optionally substituted with 1-4 halo.
- [0111] In one embodiment, Q is an unsubstituted ring and Q<sup>1</sup> does not exist. In some embodiments, Q is pyrrolidinyl or piperidinyl. In some embodiments, each J<sup>Q</sup> and J<sup>Q1</sup> is independently halo, oxo, -CN, -NO<sub>2</sub>, or V-R.
- [0112] In some embodiments, Q or Q<sup>1</sup> is substituted with 1-2 occurrences of J. In other embodiments, Q or Q<sup>1</sup> is substituted with 1 occurrence of J. In other embodiments, Q or Q<sup>1</sup> is substituted with zero occurrences of JQ and J<sup>Q1</sup> and 1 occurrence of J. In some embodiments, Q-Q<sup>1</sup> is substituted with 1 occurrence of J.
- [0113] In one embodiment,
- [0114] each J is independently oxo, (V<sup>1</sup>)-R'' or -(V)<sub>m</sub>-R\*;
- [0115] each V and V<sup>1</sup> is independently a C<sub>1-6</sub>aliphatic chain wherein up to three methylene unit of the aliphatic chain may be optionally replaced with -NR'-, -O-, -S-, -C(O)-, -S(O)-, or -S(O)<sub>2</sub>-; wherein the first or second methylene group away from the point of attachment is replaced with -C(O)-, -S(O)-, -S(O)<sub>2</sub>-, -S-, or -O-;
- [0116] each J<sup>V</sup> and J<sup>V1</sup> is independently halo or C<sub>1-4</sub>alkyl;
- [0117] R' is H or C<sub>1-4</sub>alkyl;
- [0118] R\* is H or C<sub>1-4</sub>alkyl;
- [0119] R'' is a 3-7 membered monocyclic ring containing 0-2 heteroatoms selected from O, N, or S; wherein said R'' is optionally substituted with 1-3 occurrences of J<sup>1</sup>;
- [0120] J<sup>1</sup> is oxo, halo, C<sub>1-3</sub>alkyl, haloC<sub>1-3</sub>alkyl, CN, OH, O(C<sub>1-3</sub>alkyl), NH<sub>2</sub>, NH(C<sub>1-3</sub>alkyl), N(C<sub>1-3</sub>alkyl)<sub>2</sub>, or acetyl.
- [0121] In some embodiments, Q contains sulfur, nitrogen, or oxygen as a ring member. In some embodiments, Q contains nitrogen as a ring member. In some embodiments, said nitrogen of Q is substituted with one occurrence of J<sup>Q</sup>; wherein J<sup>Q</sup> is phenyl, pyridyl, pyrimidyl, thiazolyl, C<sub>1-6</sub>ali-

phatic, or benzyl; and wherein  $J^Q$  is optionally substituted with 1-3 occurrences of  $J^{Q2}$ ; and  $J^{Q2}$  is halo, halo $C_{1-6}$ aliphatic, CN,  $NO_2$ , or  $C_{1-6}$ aliphatic, wherein 1-3 methylene units of said  $C_{1-6}$ aliphatic are optionally replaced with  $-NH-$ ,  $-O-$ ,  $-S-$ ,  $C(O)$ ,  $S(O)_2$ , or  $S(O)$ .

[0122] One aspect provides compounds wherein n is 1.


[0123] In some embodiments,  $R^1$  is a bicyclic ring, according to some embodiments, said bicyclic ring is a 8-9 membered ring. In other embodiments, said ring is a 8-9 membered heteroaryl containing 1-4 heteroatoms selected from O, N, or S. In yet other embodiments, said ring is selected from benzimidazolyl, benzoxazolyl, indazolyl, benzothiazolyl, indolyl, benzotriazolyl, pyrrolopyridyl, imidazopyridyl, or triazolopyridyl. In other embodiments, said ring is a 8-9 membered heteroaryl containing 1-4 nitrogen atoms. In some embodiments, said ring is selected from benzimidazolyl or indazolyl.

[0124] In other embodiments,  $R^1$  is a monocyclic 5-6-membered ring. In some embodiments,  $R^1$  is pyrazolyl, phenyl, pyridyl, or pyrazinyl. In other embodiments,  $R^1$  is 2-pyridyl, 3-pyridyl, or 4-pyridyl.

[0125] Another aspect provides compounds wherein n is 0. In some embodiments,  $R^1$  is a 5-6 membered monocyclic heteroaryl ring having 1-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered bicyclic heteroaryl ring having 1-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each  $R^1$  is optionally substituted with 1-5  $J^1$  groups.

[0126] In some embodiments,  $R^1$  is a 6-membered monocyclic ring selected from pyrimidyl or pyridyl. In other embodiments,  $R^1$  is a monocyclic 5-membered heteroaryl ring containing 1-3 heteroatoms selected from oxygen, nitrogen, or sulfur; or  $R^1$  is a 5-membered heteroaryl ring containing 1-2 heteroatoms selected from oxygen, nitrogen, or sulfur fused to a 6-membered aryl or heteroaryl ring containing 0-4 nitrogen atoms. In some embodiments,  $R^1$  is bonded to  $C^3$  via a carbon atom and wherein  $R^1$  contains at least one ring heteroatom.


[0127] According to one embodiment, said ring heteroatom (shown as G) is located next to the carbon atom bonded to  $C^3$  as shown below in formula II:



[0128] In some embodiments,  $R^1$  is a 5-membered ring optionally fused to another ring.


[0129] In other embodiments,  $R^1$  is a monocyclic 5-membered ring containing 1-3 heteroatoms selected from oxygen, nitrogen, or sulfur.

[0130] In some embodiments,  $R^1$  is a 5-membered ring and is selected from the following:




wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ .


[0131] In other embodiments,  $R^1$  is selected from the following:




[0132] In some embodiments,  $R^1$  is



[0133] In other embodiments,  $R^1$  is



[0134] In yet other embodiments,  $R^1$  is




[0135] In some embodiments,  $J^1$  is  $C_{1-4}$ alkyl, cyclohexyl, phenyl, thienyl, furanyl, or NH-phenyl. In certain embodiments,  $J^1$  is phenyl.

[0136] In some embodiments, the phenyl of  $J^1$  is optionally substituted with 1-3 occurrences of  $J^{Q3}$ ; wherein  $J^{Q3}$  is selected from halo, CN,  $NO_2$ ,  $X-R$ , or  $-(X)_p-Q^4$ ;  $p$  is 0-1; and  $X$  is a  $C_{1-10}$ aliphatic; wherein 1-3 methylene units of said  $C_{1-6}$ aliphatic are optionally replaced with  $-NR-$ ,  $-O-$ ,  $-S-$ ,  $C(O)$ ,  $S(O)_2$ , or  $S(O)$ ;  $R$  is H; and  $Q^4$  is a 3-6 membered monocyclic ring containing 0-2 heteroatoms selected from O or N, wherein  $X$  is optionally substituted with 1-3 occurrences of halo or CN.

[0137] In some embodiments,  $J^{Q3}$  is a  $C_{1-10}$ aliphatic chain wherein 1-2 methylene units of  $X$  are replaced with  $-O-$  or  $-NR-$ . In some embodiments,  $Q^4$  is a 3-6 membered cycloaliphatic ring. In some embodiments,  $Q^4$  is cyclopropyl.


[0138] In another embodiment,  $J^{Q3}$  is halo, OH,  $NH_2$ ,  $CH_2NH_2$ ,  $CH(CH_3)NH_2$ ,  $CH(CH_3)NHCH_3$ ,  $C(CH_3)_2NH_2$ ,  $CH_2CH_2NH_2$ ,  $CH_2NHCH_3$ ,  $CH_2N(CH_3)_2$ ,  $CH(CH_3)NH_2$ ,  $CH_2NHC(CH_3)_2$ ,  $CH_2NHCH_2CHF_2$ ,  $CH_2NHCH_2CH(CH_3)OH$ ,  $CH_2NHCH_2CH_2N(CH_3)_2$ ,  $CH_2NHCH(CH_2CH_3)_3$ ,  $CH_2NHCH_3$ ,  $CH_2NHCH_2CH_3$ ,  $CH_2NHCH_2CH_2CH_3$ ,  $CH_2NH-cyclopropyl$ ,  $CH_2NHCH_2CH_2OH$ ,  $CH_2NHCH_2CH_2OCH_3$ ,  $CH_2NHCH_2CH_2-OCH_2CH_2OH$ , azetidyl, or pyrrolidinyl.

[0139] In another embodiment,  $R^1$  is



[0140] In yet another embodiment,  $R^1$  is selected from benzimidazolyl, imidazopyridyl, triazolopyridyl, benzofuranyl, or benzothiazolyl.

[0141] In some embodiments,  $R^1$  is selected from:




wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ ;

[0142]  $A$  is C or N, provided that at least two occurrence of  $A$  are carbon; and

[0143]  $X_1$  is O, N, or S.

[0144] In other embodiments,  $R^1$  is




wherein each  $R^1$  is optionally substituted with 1-2 occurrences of  $J^1$ ;  $A$  is carbon or nitrogen; and  $X_1$  is selected from O, N, or S.

[0145] In certain embodiments,  $R^1$  is benzimidazolyl.


[0146] Another aspect provides compounds wherein  $R^2$  is  $Q$  (i.e. monocyclic ring). Each  $Q$  is optionally substituted with 1-4  $J^Q$  groups and 1-2  $J$  groups. It should be understood that  $Q$ ,  $Q^1$ , or  $Q-Q^1$  structures depicted herein showing no substituents or showing only  $J$ -substituents (see e.g., paragraphs [0070], and [0073] to 0075]), are also intended to be optionally substituted with 1-4 occurrences of  $J^Q$  or  $J^{Q1}$ .

[0147] In some embodiments, Q is selected from the following:




wherein each Q is optionally substituted with 1-4  $J^Q$  groups and 1-2 J groups.


[0148] In other embodiments, Q is selected from the following:



wherein each Q is optionally substituted with 1-4  $J^Q$  groups and 1-2 J groups. In yet another embodiment, Q is selected from the following:



In yet another embodiment, Q is




[0149] In yet other embodiments, Q is  $C_{3-7}$  cycloalkyl, pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl, homopiperidinyl, morpholinyl, or thiomorpholinyl.

[0150] According to one embodiment, Q is bonded through a carbon atom. In some embodiments, Q,  $Q^1$ , or  $Q-Q^1$  is optionally substituted with 1-2 occurrences of  $J^Q$  or  $J^{Q^1}$  respectively and optionally with 1 occurrence of J.


[0151] In some embodiments, Q or  $Q-Q^1$  is selected from the following:

[0152]  $C_{3-8}$  cycloaliphatic ring,



wherein said  $C_{1-4}$ aliphatic bonded to the pyridone is optionally substituted with halo, halo $C_{1-3}$ alkyl, CN, OH, O( $C_{1-3}$ alkyl),  $NH_2$ ,  $NH(C_{1-3}$ alkyl),  $N(C_{1-3}$ alkyl) $_2$ , or acetyl.


[0153] In some embodiments, Q is




[0154] According to another embodiment, Q is bonded through a nitrogen atom.

[0155] In some embodiments, Q is azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, homopiperazinyl, homopiperidinyl, morpholinyl, or thiomorpholinyl, wherein said Q is optionally fused to  $Q^1$ ; or Q and  $Q^1$  optionally joined together at a carbon atom to form a spirocyclic bicyclic ring, or Q and  $Q^1$ , taken together, form a bridged bicyclic ring  $Q-Q^1$  wherein said bridge is 1-3 atoms long; wherein  $Q-Q^1$  is optionally substituted with 1-2 J groups; wherein J can be bonded to either Q or  $Q^1$ .

[0156] In other embodiments, Q is selected from the following:





[0157] In yet other embodiments, Q and  $Q^1$  together combine to form a bicyclic ring selected from the group consisting of




wherein each Q or Q<sup>1</sup> is optionally substituted with 1-2 occurrences of J<sup>Q</sup> or J<sup>Q1</sup> respectively and optionally with 1 occurrence of J.

**[0158]** In some embodiments, J is bonded in the position of Q as shown below:



**[0159]** In other embodiments, J is bonded in the position of Q or Q<sup>1</sup> as shown below:



**[0160]** In some embodiments, J is C<sub>1-6</sub> aliphatic chain wherein up to 2 methylene units of C<sub>1-6</sub> aliphatic optionally replaced with heteroatom selected from O, N, or S.

**[0161]** In yet another embodiment, J is oxo, V<sup>1</sup>—R<sup>\*</sup>, or —(V)<sub>m</sub>—R<sup>\*</sup>; wherein

**[0162]** each V and V<sup>1</sup> is independently a C<sub>1-6</sub> alkylidene chain wherein 0-3 methylene units are replaced with O, N, S, CO, SO, or SO<sub>2</sub>;

**[0163]** R<sup>\*</sup> is H or C<sub>1-6</sub> alkyl;

**[0164]** m is 0-1, and

**[0165]** R" is a 3-7 membered saturated or unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

**[0166]** In some embodiments, R<sup>2</sup> is piperazine or tetrahydropyridinyl; m is 0; and J is oxo, V<sup>1</sup>—R", or —(V)<sub>m</sub>—R\* wherein R" is thiazolyl, pyridyl, pyrimidyl, phenyl; and R\* is optionally substituted with halo, O(C<sub>1-4</sub>alkyl), haloC<sub>1-4</sub>alkyl, or CN.

**[0167]** In yet another embodiment, V is O, O(C<sub>1-6</sub>alkyl), (C<sub>1-4</sub>alkyl)O, C(O)O, C(O)O(C<sub>1-6</sub>alkyl), C(O)O(C<sub>1-6</sub>alkyl)O(C<sub>1-6</sub>alkyl), C(O)(C<sub>1-6</sub>alkyl), C(O)(C<sub>1-6</sub>alkyl)N, C(O)(C<sub>1-6</sub>alkyl)NH(C<sub>1-6</sub>alkyl), C(O)(C<sub>1-6</sub>alkyl)C(O)O, C(O)(C<sub>1-6</sub>alkyl)O, C(O)(C<sub>1-6</sub>alkyl)O(C<sub>1-6</sub>alkyl), C(O)(C<sub>1-6</sub>alkyl)NH(C<sub>1-6</sub>alkyl), C(O)(C<sub>1-6</sub>alkyl)O(C<sub>1-4</sub>alkyl)O(C<sub>1-6</sub>alkyl), C(O)NH, C(O)NH(C<sub>1-6</sub>alkyl), C(O)N(C<sub>1-6</sub>alkyl), NH, N(C<sub>1-6</sub>alkyl), N(C<sub>1-6</sub>alkyl)O, NHC(O), NHC(O)C<sub>1-6</sub>alkyl, NHC(O)(C<sub>1-6</sub>alkyl)O, NHC(O)(C<sub>1-6</sub>alkyl)O(C<sub>1-6</sub>alkyl), NHC(O)(C<sub>1-6</sub>alkyl)O(C<sub>1-6</sub>alkyl)O(C<sub>1-6</sub>alkyl), NHC(O)(C<sub>1-6</sub>alkyl)NH(C<sub>1-6</sub>alkyl), C(O)N(C<sub>1-6</sub>alkyl)-, C(O)N(C<sub>1-6</sub>alkyl)<sub>2</sub>, SO<sub>2</sub>, S(O)<sub>2</sub>(C<sub>1-6</sub>alkyl)-, S(O)<sub>2</sub>(C<sub>1-6</sub>alkyl)NH, S(O)<sub>2</sub>NH(C<sub>1-6</sub>alkyl)-, S(O)<sub>2</sub>N(C<sub>1-6</sub>alkyl)<sub>2</sub>, NHSO<sub>2</sub>, or NHSO<sub>2</sub>N(C<sub>1-6</sub>alkyl)<sub>2</sub>;

**[0168]** R\* is H or C<sub>1-6</sub>alkyl; and

**[0169]** R" is C<sub>3</sub>–C<sub>8</sub> cycloaliphatic, imidazolyl, thienyl, thiazolyl, furanyl, pyrazolyl, triazolyl, pyrrolyl, phenyl, pyridyl, pyrimidyl, pyrazinyl, pyrrolidinyl, piperidinyl, piperazinyl, homopiperidinyl, homopiperazinyl, morpholinyl, thiomorpholinyl, or tetrahydropyranyl.

**[0170]** In yet another embodiment V is O, OCH<sub>2</sub>, CH<sub>2</sub>O, C(O), C(O)CH<sub>2</sub>, C(O)CH<sub>2</sub>CH<sub>2</sub>NH, C(O)CH<sub>2</sub>CH<sub>2</sub>NHCH<sub>2</sub>, C(O)CH<sub>2</sub>CH<sub>2</sub>C(O)O, C(O)CH<sub>2</sub>O, C(O)CH<sub>2</sub>OCH<sub>2</sub>, C(O)CH<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>, C(O)CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>, C(O)CH(CH<sub>3</sub>)CH<sub>2</sub>, C(O)CH(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>, C(O)O, C(O)OCH<sub>2</sub>, C(O)OCH<sub>2</sub>CH<sub>2</sub>, C(O)OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>, C(O)OCH<sub>2</sub>CEC-CH<sub>2</sub>, C(O)NH, C(O)NHCH<sub>2</sub>, C(O)N(CH<sub>3</sub>)—, C(O)N(CH<sub>3</sub>)CH<sub>2</sub>—, NH, N(C<sub>1-6</sub>aliphatic), N(CH<sub>3</sub>)CH<sub>2</sub>, NHC(O), NHC(O)CH<sub>2</sub>, NHC(O)CH<sub>2</sub>O, NHC(O)CH<sub>2</sub>OCH<sub>2</sub>, NHC(O)CH<sub>2</sub>OCH<sub>2</sub>CH<sub>2</sub>OCH<sub>2</sub>, NHC(O)CH<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>, NHC(O)C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>, NHC(O)CH(CH<sub>2</sub>CH<sub>3</sub>)CH<sub>2</sub>CH<sub>2</sub>, SO<sub>2</sub>, S(O)<sub>2</sub>CH<sub>2</sub>, S(O)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, S(O)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH, S(O)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH, S(O)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, S(O)<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, S(O)<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>, NHSO<sub>2</sub>, or NHSO<sub>2</sub>N(CH<sub>3</sub>)CH<sub>2</sub>;

**[0171]** R is H or C<sub>1-6</sub>alkyl; and

**[0172]** Q<sup>2</sup> is cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, imidazolyl, thienyl, thiazolyl, furanyl, pyrazolyl, triazolyl, tetrazolyl, pyrrolyl, phenyl, pyridyl, pyrimidyl, pyrazinyl, pyrrolidinyl, piperidinyl, piperazinyl, homopiperidinyl, homopiperazinyl, morpholinyl, thiomorpholinyl, or tetrahydropyranyl; wherein said Q<sup>2</sup> is optionally substituted with C<sub>1-6</sub>alkyl, CN, halo, haloC<sub>1-4</sub>alkyl, NH(C<sub>1-4</sub>alkyl), N(C<sub>1-4</sub>alkyl)<sub>2</sub>, or O(C<sub>1-6</sub>alkyl).

**[0173]** In some embodiments, J is selected from C<sub>1-6</sub>aliphatic, oxo, —OH, —O(C<sub>1-6</sub>aliphatic), CN, —C(O)(C<sub>1-6</sub>aliphatic), —C(O)(phenyl), —C(O)(benzyl), —C(O)CH<sub>2</sub>O(phenyl), —C(O)CH<sub>2</sub>O(C<sub>1-6</sub>aliphatic), —C(O)CH<sub>2</sub>O(benzyl), —C(O)(pyridyl), —C(O)(pyrrolidinyl), —C(O)(piperidinyl), —C(O)(piperazinyl), —C(O)(homopiperazinyl), —C(O)(morpholinyl), —C(O)(tetrahydropyranyl), —C(O)O(C<sub>1-6</sub>aliphatic), —C(O)O(phenyl), —C(O)O(benzyl), —C(O)NH<sub>2</sub>, —C(O)NH(C<sub>1-6</sub>aliphatic), —C(O)N(C<sub>1-6</sub>aliphatic)<sub>2</sub>, —C(O)NH(phenyl), —C(O)NH(benzyl), —S(O)<sub>2</sub>(C<sub>1-6</sub>aliphatic), —S(O)<sub>2</sub>(phenyl), —S(O)<sub>2</sub>(benzyl), —S(O)<sub>2</sub>(pyridyl), —S(O)<sub>2</sub>(furanyl), —S(O)<sub>2</sub>(imidazolyl), —S(O)<sub>2</sub>(thienyl), —S(O)<sub>2</sub>NH(C<sub>1-6</sub>

aliphatic), —S(O)<sub>2</sub>N(C<sub>1-6</sub>aliphatic)<sub>2</sub>, —S(O)<sub>2</sub>NH(phenyl), —S(O)<sub>2</sub>NH(benzyl), —NHC(O)(C<sub>1-6</sub>aliphatic), —NHC(O)(pyridyl), NHS(O)<sub>2</sub>N(C<sub>1-6</sub>aliphatic)<sub>2</sub>, NH<sub>2</sub>, NH(C<sub>1-6</sub>aliphatic), N(C<sub>1-6</sub>aliphatic)<sub>2</sub>; wherein said phenyl, benzyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, tetrahydropyranyl, or C<sub>1-6</sub>aliphatic is optionally substituted with halo, C<sub>1-3</sub>alkyl, CN, OH, O(C<sub>1-3</sub>alkyl), NH<sub>2</sub>, NH(C<sub>1-3</sub>alkyl), N(C<sub>1-3</sub>alkyl)<sub>2</sub>, acetyl, SO<sub>2</sub>(C<sub>1-3</sub>alkyl).

**[0174]** In other embodiments J is selected from, oxo, —OH, —NHC(O)(C<sub>1-4</sub>alkyl), —C(O)(C<sub>1-4</sub>alkyl), —C(O)NH<sub>2</sub>, —C(O)NH(C<sub>1-4</sub>alkyl), —C(O)NH(C<sub>1-4</sub>alkyl)<sub>2</sub>, —S(O)<sub>2</sub>(C<sub>1-4</sub>alkyl), —S(O)<sub>2</sub>NH(C<sub>1-4</sub>alkyl), or —S(O)<sub>2</sub>N(C<sub>1-4</sub>alkyl)<sub>2</sub>.

**[0175]** In yet other embodiments J is selected from —C(O)CH<sub>2</sub>(phenyl), NH<sub>2</sub>, NH(C<sub>1-3</sub>alkyl), N(C<sub>1-3</sub>alkyl)<sub>2</sub>, —NHS(O)<sub>2</sub>(phenyl), —NHC(O)(phenyl), —NHC(O)(pyridyl), —C(O)CH<sub>2</sub>O(phenyl), —C(O)CH<sub>2</sub>O(C<sub>1-3</sub>alkyl), —NHC(O)CH<sub>2</sub>O(C<sub>1-3</sub>alkyl), —NHC(O)CH<sub>2</sub>O(phenyl), —C(O)CH<sub>2</sub>N(C<sub>1-3</sub>alkyl)<sub>2</sub>, —NHC(O)CH<sub>2</sub>N(C<sub>1-3</sub>alkyl)<sub>2</sub>, —C(O)CH<sub>2</sub>O(C<sub>1-3</sub>alkyl)O(C<sub>1-3</sub>alkyl), —NHC(O)CH<sub>2</sub>O(C<sub>1-3</sub>alkyl)O(C<sub>1-3</sub>alkyl), —C(O)O(C<sub>1-3</sub>alkyl)O(C<sub>1-3</sub>alkyl), phenyl, pyridyl, thiazolyl, CF<sub>3</sub>, pyrimidyl, wherein said phenyl, pyridyl, or pyrimidyl group is optionally substituted with halo, O(C<sub>1-6</sub>aliphatic), CN, CF<sub>3</sub>, C(O)(furanyl).

**[0176]** In one embodiment, two J groups, attached to the same carbon atom, taken together form 1,3 dioxalane.

**[0177]** In some embodiments, the variables are as depicted in the compounds of the following Tables. Another embodiment provides a compound selected from the following tables:

TABLE 1






TABLE 1-continued




I-3


TABLE 1-continued




I-7




I-4




I-8




I-5



I-9



I-6



I-10

TABLE 1-continued




TABLE 1-continued




TABLE 1-continued




TABLE 1-continued




TABLE 2




TABLE 2-continued





TABLE 3




TABLE 3-continued







TABLE 3-continued







TABLE 3-continued







TABLE 3-continued







TABLE 3-continued







TABLE 3-continued




TABLE 3-continued




TABLE 3-continued




TABLE 3-continued







TABLE 3-continued




I-65



I-66




I-67




I-68


TABLE 3-continued



I-69



I-70



I-71

TABLE 3-continued







TABLE 3-continued










TABLE 3-continued




I-80



I-81




I-82




I-83


TABLE 3-continued



I-84




I-85




I-86

TABLE 3-continued




I-87


TABLE 3-continued




I-90




I-88



I-91



I-89



I-92

TABLE 3-continued






TABLE 3-continued








TABLE 3-continued




I-99


TABLE 3-continued




I-102




I-100



I-103



I-101



I-104

TABLE 3-continued




TABLE 3-continued










TABLE 3-continued




I-111


TABLE 3-continued




I-114




I-112



I-115




I-113




I-116

TABLE 3-continued




I-117


TABLE 3-continued




I-120




I-118



I-121



I-119



I-122

TABLE 3-continued






TABLE 3-continued








TABLE 3-continued




I-129


TABLE 3-continued




I-132




I-130



I-133



I-131



I-134

TABLE 3-continued




TABLE 3-continued











TABLE 3-continued




I-142


TABLE 3-continued




I-145




I-143



I-146



I-144



I-147

TABLE 3-continued






TABLE 3-continued






TABLE 3-continued




TABLE 3-continued




TABLE 3-continued






TABLE 3-continued










TABLE 3-continued



I-168




I-169




I-170


TABLE 3-continued



I-171




I-172



I-173

TABLE 4



I-174

TABLE 4-continued






TABLE 4-continued






TABLE 4-continued






TABLE 4-continued






TABLE 4-continued






TABLE 4-continued








TABLE 4-continued




I-193


TABLE 4-continued




I-196




I-194



I-197



I-195



I-198

TABLE 4-continued






TABLE 4-continued






TABLE 4-continued






TABLE 4-continued







TABLE 4-continued






TABLE 4-continued








TABLE 4-continued




I-218


TABLE 4-continued




I-221




I-219



I-222



I-220



I-223

TABLE 4-continued






TABLE 4-continued








TABLE 4-continued




I-230


TABLE 4-continued




I-233




I-231



I-234




I-232




I-235

TABLE 4-continued




I-236


TABLE 4-continued




I-239




I-237



I-240



I-238



I-241

TABLE 4-continued




TABLE 4-continued




TABLE 4-continued






TABLE 4-continued








TABLE 4-continued



I-256



I-257






TABLE 4-continued



I-259



I-260






TABLE 4-continued



I-262



I-263




TABLE 4-continued







TABLE 4-continued







TABLE 4-continued







TABLE 4-continued







TABLE 4-continued





TABLE 5




TABLE 5-continued






TABLE 5-continued








TABLE 5-continued




I-289


TABLE 5-continued




I-292




I-290



I-293



I-291



I-294

TABLE 5-continued




TABLE 5-continued








TABLE 5-continued






TABLE 5-continued










TABLE 5-continued




I-308



I-309




I-310




I-311


TABLE 5-continued



I-312



I-313



I-314

TABLE 5-continued





TABLE 5-continued







TABLE 5-continued






TABLE 5-continued








TABLE 5-continued




I-327


TABLE 5-continued



I-329



I-328



I-330

TABLE 5-continued




TABLE 5-continued




TABLE 5-continued






TABLE 5-continued






TABLE 5-continued







TABLE 5-continued



I-345



I-347



Ⓢ indicates text missing or illegible when filed

TABLE 6






TABLE 6-continued




TABLE 6-continued






TABLE 6-continued




TABLE 6-continued






TABLE 6-continued






TABLE 6-continued





TABLE 6-continued






TABLE 6-continued





TABLE 6-continued






TABLE 6-continued

I-375



I-376





TABLE 6-continued

I-377



Ⓢ indicates text missing or illegible when filed


**[0178]** Compounds of this invention include those described generally herein, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75<sup>th</sup> Ed. Additionally, general principles of organic chemistry are described in “Organic Chemistry”, Thomas Sorrell, University Science Books, Sausalito: 1999, and “March’s Advanced Organic Chemistry”, 5<sup>th</sup> Ed., Ed.: Smith, M. B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

**[0179]** As described herein, a specified number range of atoms includes any integer therein. For example, a group having from 1-4 atoms could have 1, 2, 3, or 4 atoms.

**[0180]** As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally herein, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase “optionally substituted” is used interchangeably with the phrase “substituted or unsubstituted.” In general, the term “substituted”, whether preceded by the term “optionally” or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are

preferably those that result in the formation of stable or chemically feasible compounds.

**[0181]** Unless otherwise indicated, a substituent connected by a bond drawn from the center of a ring means that the substituent can be bonded to any position in the ring. In example i below, for instance, J<sup>1</sup> can be bonded to any position on the pyridyl ring. For bicyclic rings, a bond drawn through both rings indicates that the substituent can be bonded from any position of the bicyclic ring. In example ii below, for instance, J<sup>1</sup> can be bonded to the 5-membered ring (on the nitrogen atom, for instance), and to the 6-membered ring.



**[0182]** The term “stable”, as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40° C. or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

**[0183]** The term “aliphatic” or “aliphatic group”, as used herein, means a straight-chain (i.e., unbranched), branched, or cyclic, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation that has a single point of attachment to the rest of the molecule.

**[0184]** Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. Aliphatic groups may be linear or branched, substituted or unsubstituted alkyl, alkenyl, or alkynyl groups. Specific examples include, but are not limited to, methyl, ethyl, isopropyl, n-propyl, sec-butyl, vinyl, n-butenyl, ethynyl, and tert-butyl.

**[0185]** The term “cycloaliphatic” (or “carbocycle” or “carbocyclyl”) refers to a monocyclic  $C_3$ - $C_8$  hydrocarbon or bicyclic  $C_8$ - $C_{12}$  hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Examples of cycloaliphatic groups include, but are not limited to, cycloalkyl and cycloalkenyl groups. Specific examples include, but are not limited to, cyclohexyl, cyclopropenyl, and cyclobutyl.

**[0186]** The term “heterocycle”, “heterocyclyl”, or “heterocyclic” as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members are an independently selected heteroatom. In some embodiments, the “heterocycle”, “heterocyclyl”, or “heterocyclic” group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

**[0187]** Examples of heterocycles include, but are not limited to, 3-1H-benzimidazol-2-one, 3-(1-alkyl)-benzimidazol-2-one, 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholino, 3-morpholino, 4-morpholino, 2-thiomorpholino, 3-thiomorpholino, 4-thiomorpholino, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-tetrahydropiperazinyl, 2-tetrahydropiperazinyl, 3-tetrahydropiperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 1-pyrazolinyl, 3-pyrazolinyl, 4-pyrazolinyl, 5-pyrazolinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 2-thiazolidinyl, 3-thiazolidinyl, 4-thiazolidinyl, 1-imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 5-imidazolidinyl, indolyl, tetrahydroquinolyl, tetrahydroisoquinolyl, benzothiolane, benzodithiane, and 1,3-dihydro-imidazol-2-one.

**[0188]** Cyclic groups, (e.g. cycloaliphatic and heterocycles), can be linearly fused, bridged, or spirocyclic.

**[0189]** The term “heteroatom” means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or  $NR^+$  (as in N-substituted pyrrolidinyl)).

**[0190]** The term “unsaturated”, as used herein, means that a moiety has one or more units of unsaturation. As would be known by one of skill in the art, unsaturated groups can be partially unsaturated or fully unsaturated. Examples of par-

tially unsaturated groups include, but are not limited to, butene, cyclohexene, and tetrahydropyridine. Fully unsaturated groups can be aromatic, anti-aromatic, or non-aromatic. Examples of fully unsaturated groups include, but are not limited to, phenyl, cyclooctatetraene, pyridyl, thienyl, and 1-methylpyridin-2(1H)-one. For the avoidance of doubt, it should be understood that rings such as phenyl, pyridyl, and thienyl are aromatic, while rings such as cyclooctatetraene and 1-methylpyridin-2(1H)-one are not aromatic.

**[0191]** The term “aromatic” as used herein, is used to describe a moiety that satisfies Hückel’s rule, having  $4n+2$  electrons in a delocalized cloud.

**[0192]** The term “alkoxy”, or “thioalkyl”, as used herein, refers to an alkyl group, as previously defined, attached through an oxygen (“alkoxy”) or sulfur (“thioalkyl”) atom.

**[0193]** The terms “haloalkyl”, “haloalkenyl”, “haloaliphatic”, and “haloalkoxy” mean alkyl, alkenyl or alkoxy, as the case may be, substituted with one or more halogen atoms. This term includes perfluorinated alkyl groups, such as  $-CF_3$  and  $-CF_2CF_3$ .

**[0194]** The terms “halogen”, “halo”, and “hal” mean F, Cl, Br, or I.

**[0195]** The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term “aryl” may be used interchangeably with the term “aryl ring”.

**[0196]** The term “heteroaryl”, used alone or as part of a larger moiety as in “heteroaralkyl” or “heteroarylalkoxy”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term “heteroaryl” may be used interchangeably with the term “heteroaryl ring” or the term “heteroaromatic”. Examples of heteroaryl rings include, but are not limited to, 2-furanyl, 3-furanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, benzimidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, N-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, pyridazinyl (e.g., 3-pyridazinyl), 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, tetrazolyl (e.g., 5-tetrazolyl), triazolyl (e.g., 2-triazolyl and 5-triazolyl), 2-thienyl, 3-thienyl, benzofuryl, benzothiophenyl, indolyl (e.g., 2-indolyl), pyrazolyl (e.g., 2-pyrazolyl), isothiazolyl, 1,2,3-oxadiazolyl, 1,2,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, purinyl, pyrazinyl, 1,3,5-triazinyl, quinolinyl (e.g., 2-quinolinyl, 3-quinolinyl, 4-quinolinyl), and isoquinolinyl (e.g., 1-isoquinolinyl, 3-isoquinolinyl, or 4-isoquinolinyl).

**[0197]** The term “hydrogen bond acceptor” refers to a relatively electronegative atom that can form a hydrogen bond with a hydrogen atom. Examples of hydrogen bond acceptors include, but are not limited to, fluorine, oxygen, nitrogen, and sulfur.

**[0198]** The term “protecting group” and “protective group” as used herein, are interchangeable and refer to an agent used to temporarily block one or more desired functional groups in a compound with multiple reactive sites. In certain embodiments, a protecting group has one or more, or preferably all, of the following characteristics: a) is added selectively to a


functional group in good yield to give a protected substrate that is b) stable to reactions occurring at one or more of the other reactive sites; and c) is selectively removable in good yield by reagents that do not attack the regenerated, deprotected functional group. As would be understood by one skilled in the art, in some cases, the reagents do not attack other reactive groups in the compound. In other cases, the reagents may also react with other reactive groups in the compound. Examples of protecting groups are detailed in Greene, T. W., Wuts, P. G in "Protective Groups in Organic Synthesis", Third Edition, John Wiley & Sons, New York: 1999 (and other editions of the book), the entire contents of which are hereby incorporated by reference. The term "nitrogen protecting group", as used herein, refers to an agent used to temporarily block one or more desired nitrogen reactive sites in a multifunctional compound. Preferred nitrogen protecting groups also possess the characteristics exemplified for a protecting group above, and certain exemplary nitrogen protecting groups are also detailed in Chapter 7 in Greene, T. W., Wuts, P. G in "Protective Groups in Organic Synthesis", Third Edition, John Wiley & Sons, New York: 1999, the entire contents of which are hereby incorporated by reference.

**[0199]** In some embodiments, a methylene unit of an alkyl or aliphatic chain is optionally replaced with another atom or group. Examples of such atoms or groups include, but are not limited to, nitrogen, oxygen, sulfur,  $-\text{C}(\text{O})-$ ,  $-\text{C}(=\text{N}-\text{CN})-$ ,  $-\text{C}(=\text{NR})-$ ,  $-\text{C}(=\text{NOR})-$ ,  $-\text{SO}-$ , and  $-\text{SO}_2-$ . These atoms or groups can be combined to form larger groups. Examples of such larger groups include, but are not limited to,  $-\text{OC}(\text{O})-$ ,  $-\text{C}(\text{O})\text{CO}-$ ,  $-\text{CO}_2-$ ,  $-\text{C}(\text{O})\text{NR}-$ ,  $-\text{C}(=\text{N}-\text{CN})-$ ,  $-\text{NRCO}-$ ,  $-\text{NRC}(\text{O})\text{O}-$ ,  $-\text{SO}_2\text{NR}-$ ,  $-\text{NRSO}_2-$ ,  $-\text{NRC}(\text{O})\text{NR}-$ ,  $-\text{OC}(\text{O})\text{NR}-$ , and  $-\text{NRSO}_2\text{NR}-$ , wherein R is, for example, H or  $\text{C}_{1-6}$  aliphatic. It should be understood that these groups can be bonded to the methylene units of the aliphatic chain via single, double, or triple bonds. An example of an optional replacement (nitrogen atom in this case) that is bonded to the aliphatic chain via a double bond would be  $-\text{CH}_2\text{CH}=\text{N}-\text{CH}_3$ . In some cases, especially on the terminal end, an optional replacement can be bonded to the aliphatic group via a triple bond. One example of this would be  $\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{N}$ . It should be understood that in this situation, the terminal nitrogen is not bonded to another atom.


**[0200]** Unless otherwise indicated, the optional replacements form a chemically stable compound. Optional replacements can occur both within the chain and/or at either end of the chain; i.e. both at the point of attachment and/or also at the terminal end. Two optional replacements can also be adjacent to each other within a chain so long as it results in a chemically stable compound. For example, a  $\text{C}_3$  aliphatic can be optionally replaced by 2 nitrogen atoms to form  $-\text{C}\text{f}\text{N}=\text{N}$ . The optional replacements can also completely replace all of the carbon atoms in a chain. For example, a  $\text{C}_3$  aliphatic can be optionally replaced by  $-\text{NR}-$ ,  $-\text{C}(\text{O})-$ , and  $-\text{NR}-$  to form  $-\text{NRC}(\text{O})\text{NR}-$  (a urea).

**[0201]** Unless otherwise indicated, if the replacement occurs at the terminal end, the replacement atom is bound to a hydrogen atom on the terminal end. For example, if a methylene unit of  $-\text{CH}_2\text{CH}_2\text{CH}_3$  were optionally replaced with  $-\text{O}-$ , the resulting compound could be  $-\text{OCH}_2\text{CH}_3$ ,  $-\text{CH}_2\text{OCH}_3$ , or  $-\text{CH}_2\text{CH}_2\text{OH}$ . It should be understood that if the terminal atom does not contain any free valence electrons, then a hydrogen atom is not required at the terminal end (e.g.,  $-\text{CH}_2\text{CH}_2\text{CH}=\text{O}$  or  $-\text{CH}_2\text{CH}_2\text{C}\equiv\text{N}$ ).

**[0202]** Unless otherwise indicated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, geometric, conformational, and rotational) forms of the structure. For example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers are included in this invention. As would be understood to one skilled in the art, a substituent can freely rotate around any rotatable bonds. For example, a substituent drawn as



also represents



**[0203]** Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, geometric, conformational, and rotational mixtures of the present compounds are within the scope of the invention.

**[0204]** Unless otherwise indicated, all tautomeric forms of the compounds of the invention are within the scope of the invention.

**[0205]** Additionally, unless otherwise indicated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a  $^{13}\text{C}$ - or  $^{14}\text{C}$ -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

#### Pharmaceutically Acceptable Salts

**[0206]** The compounds of this invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable salt.

**[0207]** A "pharmaceutically acceptable salt" means any non-toxic salt of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. As used herein, the term "inhibitorily active metabolite or residue thereof" means that a metabolite or residue thereof is also an inhibitor of the ATR protein kinase.

**[0208]** Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. These salts can be prepared in situ

during the final isolation and purification of the compounds. Acid addition salts can be prepared by 1) reacting the purified compound in its free-based form with a suitable organic or inorganic acid and 2) isolating the salt thus formed.

[0209] Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, glycolate, gluconate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

[0210] Base addition salts can be prepared by 1) reacting the purified compound in its acid form with a suitable organic or inorganic base and 2) isolating the salt thus formed. Salts derived from appropriate bases include alkali metal (e.g., sodium, lithium, and potassium), alkaline earth metal (e.g., magnesium and calcium), ammonium and  $N^+(C_{1-4}alkyl)_4$  salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

[0211] Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate. Other acids and bases, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid or base addition salts.

#### ABBREVIATIONS

- [0212] The following abbreviations are used:
- [0213] DMSO dimethyl sulfoxide
- [0214] ATP adenosine triphosphate
- [0215] DTT dithiothreitol
- [0216]  $^1H$ NMR proton nuclear magnetic resonance
- [0217] HPLC high performance liquid chromatography
- [0218] LCMS liquid chromatography-mass spectrometry
- [0219] TLC thin layer chromatography
- [0220] Rt retention time

#### Compound Uses

[0221] One aspect of this invention provides compounds that are inhibitors of ATR kinase, and thus are useful for treating or lessening the severity of a disease, condition, or disorder where ATR is implicated in the disease, condition, or disorder.

[0222] Another aspect of this invention provides compounds that are useful for the treatment of diseases, disorders, and conditions characterized by excessive or abnormal cell proliferation. Such diseases include, a proliferative or hyperproliferative disease. Examples of proliferative and hyperproliferative diseases include, without limitation, cancer and myeloproliferative disorders.

[0223] The term "cancer" includes, but is not limited to the following cancers. Oral: buccal cavity, lip, tongue, mouth, pharynx; Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell or epidermoid, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, larynx, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel or small intestines (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel or large intestines (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma), colon, colon-rectum, colorectal; rectum, Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma, biliary passages; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondroblastoma), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma), breast; Hematologic: blood (myeloid leukemia [acute and chronic], acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma [malignant lymphoma] hairy cell; lymphoid disorders; Skin: malignant melanoma, basal cell

carcinoma, squamous cell carcinoma, Kaposi's sarcoma, keratoacanthoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis, Thyroid gland: papillary thyroid carcinoma, follicular thyroid carcinoma, undifferentiated thyroid cancer, medullary thyroid carcinoma, multiple endocrine neoplasia type 2A, multiple endocrine neoplasia type 2B, familial medullary thyroid cancer, pheochromocytoma, paraganglioma; and Adrenal glands: neuroblastoma.

**[0224]** Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above-identified conditions. In some embodiments, the cancer is selected from colorectal, thyroid, or breast cancer.

**[0225]** The term "myeloproliferative disorders", includes disorders such as polycythemia vera, thrombocythemia, myeloid metaplasia with myelofibrosis, hypereosinophilic syndrome, juvenile myelomonocytic leukemia, systemic mast cell disease, and hematopoietic disorders, in particular, acute-myelogenous leukemia (AML), chronic-myelogenous leukemia (CML), acute-promyelocytic leukemia (APL), and acute lymphocytic leukemia (ALL).

#### Pharmaceutically Acceptable Derivatives or Prodrugs

**[0226]** In addition to the compounds of this invention, pharmaceutically acceptable derivatives or prodrugs of the compounds of this invention may also be employed in compositions to treat or prevent the herein identified disorders.

**[0227]** The compounds of this invention can also exist as pharmaceutically acceptable derivatives.

**[0228]** A "pharmaceutically acceptable derivative" is an adduct or derivative which, upon administration to a patient in need, is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof. Examples of pharmaceutically acceptable derivatives include, but are not limited to, esters and salts of such esters.

**[0229]** A "pharmaceutically acceptable derivative or pro-drug" means any pharmaceutically acceptable ester, salt of an ester or other derivative or salt thereof of a compound, of this invention which, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof. Particularly favoured derivatives or prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a patient (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological compartment (e.g., the brain or lymphatic system) relative to the parent species.

**[0230]** Pharmaceutically acceptable prodrugs of the compounds of this invention include, without limitation, esters, amino acid esters, phosphate esters, metal salts and sulfonate esters.

#### Pharmaceutical Compositions

**[0231]** The present invention also provides compounds and compositions that are useful as inhibitors of ATR kinase.

**[0232]** One aspect of this invention provides pharmaceutically acceptable compositions that comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle.

**[0233]** The pharmaceutically acceptable carrier, adjuvant, or vehicle, as used herein, includes any and all solvents,

diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component (s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention.

**[0234]** Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such as propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

#### Combination Therapies

**[0235]** Another aspect of this invention is directed towards a method of treating cancer in a subject in need thereof, comprising administration of a compound of this invention or a pharmaceutically acceptable salt thereof, and an additional therapeutic agent. In some embodiments, said method comprises the sequential or co-administration of the compound or a pharmaceutically acceptable salt thereof, and the additional therapeutic agent.

**[0236]** In some embodiments, said additional therapeutic agent is an anti-cancer agent. In other embodiments, said additional therapeutic agent is a DNA-damaging agent. In yet other embodiments, said additional therapeutic agent is selected from radiation therapy, chemotherapy, or other agents typically used in combination with radiation therapy or chemotherapy, such as radiosensitizers and chemosensitizers.

**[0237]** As would be known by one of skill in the art, radiosensitizers are agents that can be used in combination with radiation therapy. Radiosensitizers work in various different ways, including, but not limited to, making cancer cells more

sensitive to radiation therapy, working in synergy with radiation therapy to provide an improved synergistic effect, acting additively with radiation therapy, or protecting surrounding healthy cells from damage caused by radiation therapy. Likewise chemosensitizers are agents that can be used in combination with chemotherapy. Similarly, chemosensitizers work in various different ways, including, but not limited to, making cancer cells more sensitive to chemotherapy, working in synergy with chemotherapy to provide an improved synergistic effect, acting additively to chemotherapy, or protecting surrounding healthy cells from damage caused by chemotherapy.

**[0238]** Examples of DNA-damaging agents that may be used in combination with compounds of this invention include, but are not limited to radiation, radiomimetic neocarzinostatin, Platinating agents, such as Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, Lobaplatin, Triplatin Tetranitrate, Picoplatin, Satraplatin, ProLindac, Aroplatin and other derivatives; Topo I inhibitors, such as Camptothecin, Topotecan, Irinotecan/SN38, Rubitecan, Belotecan, and other derivatives; Topo II inhibitors, such as Etoposide, Daunorubicin, Doxorubicin, Aclarubicin, Epirubicin, Idarubicin, Amrubicin, Pirarubicin, Valrubicin, Zorubicin and Teniposide; Antimetabolites, such as Folic family (Methotrexate, Pemetrexed and relatives); Purine antagonists and Pyrimidine antagonists (Thioguanine, Fludarabine, Cladribine, Cytarabine, Gemcitabine, 6-Mercaptopurine, 5-Fluorouracil (5FU) and relatives); Some other examples of antimetabolites: (Aminopterin, Raltitrexed, Pentostatin, Cladribine, Clofarabine, Fludarabine, Mercaptopurine, Fluorouracil, Capecitabine, Tegafur, Carmofur, Floxuridine, Azacitidine and Hydroxyurea); Alkylating agents, such as Nitrogen mustards (Cyclophosphamide, Melphalan, Chlorambucil, mechlorethamine, Ifosfamide and relatives); nitrosoureas (eg Carmustine); Triazines (Dacarbazine, temozolomide); Alkyl sulphonates (eg Busulfan); Procarbazine and Aziridines; Some other examples of alkylating agents: (Mechlorethamine, Cyclophosphamide, Ifosfamide, Trofosfamide, Chlorambucil, Melphalan, Prednimustine, Bendamustine, Uramustine, Estramustine, Lomustine, Semustine, Fotemustine, Nimustine, Ranimustine, Streptozocin, Busulfan, Mannosulfan, Treosulfan, Carboquone, ThioTEPA, Triaziquone, Triethyl-enemelamine, Procarbazine, Altretamine, Mitobronitol, Actinomycin, Bleomycin, Mitomycin and Plicamycin). Antibiotics, such as Hydroxyurea, Anthracyclines (doxorubicin, daunorubicin, epirubicin and other derivatives); Anthracenediones (Mitoxantrone and relatives); *Streptomyces* family (Bleomycin, Mitomycin C, actinomycin); and Ultraviolet light.

**[0239]** Other therapies or anticancer agents that may be used in combination with the inventive agents of the present invention include surgery, radiotherapy (in but a few examples, gamma-radiation, neutron beam radiotherapy, electron beam radiotherapy, proton therapy, brachytherapy, and systemic radioactive isotopes, to name a few), endocrine therapy, biologic response modifiers (interferons, interleukins, and tumor necrosis factor (TNF) to name a few), hyperthermia and cryotherapy, agents to attenuate any adverse effects (e.g., antiemetics), and other approved chemotherapeutic drugs, including, but not limited to, the DNA damaging agents listed herein, spindle poisons (Vinblastine, Vincristine, Vinorelbine, Paclitaxel), podophyllotoxins (Etoposide, Irinotecan, Topotecan), nitrosoureas (Carmustine, Lomustine), inorganic ions (Cisplatin, Carboplatin), enzymes (As-

paraginase), and hormones (Tamoxifen, Leuprolide, Flutamide, and Megestrol), Gleevec™, adriamycin, dexamethasone, and cyclophosphamide.

**[0240]** A compound of the instant invention may also be useful for treating cancer in combination with any of the following therapeutic agents: abarelix (Plenaxis Depot®); aldesleukin (Prokin®); Aldesleukin (Proleukin®); Alemtuzumab (Campath®); alitretinoin (Panretin®); allopurinol (Zyloprim®); altretamine (Hexalen®); amifostine (Ethyol®); anastrozole (Arimidex®); arsenic trioxide (Trisenox®); asparaginase (Elspar®); azacitidine (Vidaza®); bevacuzimab (Avastin®); bexarotene capsules (Targretin®); bexarotene gel (Targretin®); bleomycin (Blenoxane®); bortezomib (Velcade®); busulfan intravenous (Busulfex®); busulfan oral (Myleran®); calusterone (Methosarb®); capecitabine (Xeloda®); carboplatin (Paraplatin®); carmustine (BCNU®, BiCNU®); carmustine (Gliadel®); carmustine with Polifeprosan 20 Implant (Gliadel Wafer®); celecoxib (Celebrex®); cetuximab (Erbix®); chlorambucil (Leukeran®); cisplatin (Platinol®); cladribine (Leustatin®, 2-CdA®); clofarabine (Clolar®); cyclophosphamide (Cytosan®, Neosar®); cyclophosphamide (Cytosan Injection®); cyclophosphamide (Cytosan Tablet®); cytarabine (Cytosar-U); cytarabine liposomal (DepoCyt®); dacarbazine (DTIC-Dome®); dactinomycin, actinomycin D (Cosmegen®); Darbepoetin alfa (Aranesp®); daunorubicin liposomal (DanuoXome®); daunorubicin, daunomycin (Daunorubicin®); daunorubicin, daunomycin (Cerubidine®); Denileukin difitox (Ontak®); dexrazoxane (Zincard®); docetaxel (Taxotere®); doxorubicin (Adriamycin PFS®); doxorubicin (Adriamycin®, Rubex®); doxorubicin (Adriamycin PFS Injection®); doxorubicin liposomal (Doxil®); dromostanolone propionate (Dromostanolone®); dromostanolone propionate (masterone Injection®); Elliott's B Solution (Elliott's B Solution®); epirubicin (Ellence®); Epoetin alfa (Epogen®); erlotinib (Tarceva®); estramustine (Emcyt®); etoposide phosphate (Etopophos®); etoposide, VP-16 (Vepesid®); exemestane (Aromasin®); Filgrastim (Neupogen®); floxuridine (intraarterial) (FUDR®); fludarabine (Fludara®); fluorouracil, 5-FU (Adrucil®); fulvestrant (Faslodex®); gefitinib (Iressa®); gemcitabine (Gemzar®); gemtuzumab ozogamicin (Mylotarg®); goserelin acetate (Zoladex Implant®); goserelin acetate (Zoladex®); histrelin acetate (Histrelin Implant®); hydroxyurea (Hydrea®); Ibriumomab Tiuxetan (Zevalin®); idarubicin (Idamycin®); ifosfamide (IFEX®); imatinib mesylate (Gleevec®); interferon alfa 2a (Roferon A®); Interferon alfa-2b (Intron A®); irinotecan (Camptosar®); lenalidomide (Revlimid®); letrozole (Femara®); leucovorin (Wellcovorin®, Leucovorin®); Leuprolide Acetate (Eligard®); levamisole (Ergamisol®); lomustine, CCNU (CeeBU); mechlorethamine, nitrogen mustard (Mustargen®); megestrol acetate (Megace®); melphalan, L-PAM (Alkeran®); mercaptopurine, 6-MP (Purinethol®); mesna (Mesnex®); mesna (Mesnex Tabs®); methotrexate (Methotrexate®); methoxsalen (Uvadex®); mitomycin C (Mutamycin®); mitotane (Lysodren®); mitoxantrone (Novantrone®); nandrolone phenpropionate (Duralbolin-50®); nelarabine (Arranon®); Nofetumomab (Verluma®); Oprelvekin (Neumega®); oxaliplatin (Eloxatin®); paclitaxel (Paxene®); paclitaxel (Taxol®); paclitaxel protein-bound particles (Abraxane®); palifermin (Kepivance®); pamidronate (Aredia®); pegademase (Adagen (Pegademase Bovine®)); pegaspargase (Oncaspar®); Pegfilgrastim (Neulasta®); pemetrexed disodium (Alimta®); pentostatin

(Nipent®); pipobroman (Vercyte®); plicamycin, mithramycin (Mithracin®); porfimer sodium (Photofrin®); procabazine (Matulane®); quinacrine (Atabrine®); Rasburicase (Elitek®); Rituximab (Rituxan®); sargramostim (Leukine®); Sargramostim (Prokine®); sorafenib (Nexavar®); streptozocin (Zanosar®); sunitinib maleate (Sutent®); talc (Sclerosol®); tamoxifen (Nolvadex®); temozolomide (Temodar®); teniposide, VM-26 (Vumon®); testolactone (Teslac®); thioguanine, 6-TG (Thioguanine®); thiotepa (Thioplex®); topotecan (Hycamtin®); toremifene (Fareston®); Tositumomab (Bexxar®); Tositumomab/I-131 tositumomab (Bexxar®); Trastuzumab (Herceptin®); tretinoin, ATRA (Vesanoid®); Uracil Mustard (Uracil Mustard Capsules®); valrubicin (Valstar®); vinblastine (Velban®); vincristine (Oncovin®); vinorelbine (Navelbine®); zoledronate (Zometa®) and vorinostat (Zolinza®).

**[0241]** For a comprehensive discussion of updated cancer therapies see, <http://www.nci.nih.gov/>, a list of the FDA approved oncology drugs at <http://www.fda.gov/cder/cancer/druglistframe.htm>, and The Merck Manual, Seventeenth Ed. 1999, the entire contents of which are hereby incorporated by reference.

Compositions for Administration into a Subject

**[0242]** The ATR kinase inhibitors or pharmaceutical salts thereof may be formulated into pharmaceutical compositions for administration to animals or humans. These pharmaceutical compositions, which comprise an amount of the ATR inhibitor effective to treat or prevent the diseases or conditions described herein and a pharmaceutically acceptable carrier, are another embodiment of the present invention.

**[0243]** The exact amount of compound required for treatment will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression “dosage unit form” as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term “patient”, as used herein, means an animal, preferably a mammal, and most preferably a human.

**[0244]** In some embodiments, these compositions optionally further comprise one or more additional therapeutic agents. For example, chemotherapeutic agents or other antiproliferative agents may be combined with the compounds of this invention to treat proliferative diseases and cancer. Examples of known agents with which these compositions can be combined are listed above under the “Combination Therapies” section and also throughout the specification. Some embodiments provide a simultaneous, separate or sequential use of a combined preparation.

Modes of Administration and Dosage Forms

**[0245]** The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

**[0246]** Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

**[0247]** Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

**[0248]** The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

**[0249]** In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot

injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

**[0250]** Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

**[0251]** Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar—agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

**[0252]** Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

**[0253]** The active compounds can also be in microencapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed

manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

**[0254]** Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

**[0255]** The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term “parenteral” as used herein includes, but is not limited to, subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

**[0256]** Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents which are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

**[0257]** The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include, but are not limited to, lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use,

the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

**[0258]** Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

**[0259]** The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

**[0260]** Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

**[0261]** For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

**[0262]** For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

**[0263]** The pharmaceutical compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

**[0264]** The amount of protein kinase inhibitor that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01-100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

**[0265]** It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the

severity of the particular disease being treated. The amount of inhibitor will also depend upon the particular compound in the composition.

**Administering with Another Agent**

**[0266]** Depending upon the particular protein kinase-mediated conditions to be treated or prevented, additional drugs, which are normally administered to treat or prevent that condition, may be administered together with the compounds of this invention.

**[0267]** Those additional agents may be administered separately, as part of a multiple dosage regimen, from the protein kinase inhibitor-containing compound or composition. Alternatively, those agents may be part of a single dosage form, mixed together with the protein kinase inhibitor in a single composition.

**[0268]** Another aspect of this invention is directed towards a method of treating cancer in a subject in need thereof, comprising the sequential or co-administration of a compound of this invention or a pharmaceutically acceptable salt thereof, and an anti-cancer agent. In some embodiments, said anti-cancer agent is selected from Platinating agents, such as Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, or Satraplatin and other derivatives; Topo I inhibitors, such as Camptothecin, Topotecan, irinotecan/SN38, rubitecan and other derivatives; Antimetabolites, such as Folic family (Methotrexate, Pemetrexed and relatives); Purine family (Thioguanine, Fludarabine, Cladribine, 6-Mercaptopurine and relatives); Pyrimidine family (Cytarabine, Gemcitabine, 5-Fluorouracil and relatives); Alkylating agents, such as Nitrogen mustards (Cyclophosphamide, Melphalan, Chlorambucil, mechlorethamine, Ifosfamide, and relatives); nitrosoureas (e.g. Carmustine); Triazines (Dacarbazine, temozolomide); Alkyl sulphonates (e.g. Busulfan); Procarbazine and Aziridines; Antibiotics, such as Hydroxyurea; Anthracyclines (doxorubicin, daunorubicin, epirubicin and other derivatives); Anthracenediones (Mitoxantrone and relatives); *Streptomyces* family (Bleomycin, Mitomycin C, actinomycin) and Ultraviolet light.

#### Biological Samples

**[0269]** As inhibitors of ATR kinase, the compounds and compositions of this invention are also useful in biological samples. One aspect of the invention relates to inhibiting ATR kinase activity in a biological sample, which method comprises contacting said biological sample with a compound of formula I or a composition comprising said compound. The term "biological sample", as used herein, means an *in vitro* or an *ex vivo* sample, including, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

**[0270]** Inhibition of ATR kinase activity in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, blood transfusion, organ-transplantation, and biological specimen storage.

#### Study of Protein Kinases

**[0271]** Another aspect of this invention relates to the study of protein kinases in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such protein kinases; and the comparative evaluation of new protein kinase inhibitors. Examples of such uses

include, but are not limited to, biological assays such as enzyme assays and cell-based assays.

**[0272]** The activity of the compounds as protein kinase inhibitors may be assayed in vitro, in vivo or in a cell line. In vitro assays include assays that determine inhibition of either the kinase activity or ATPase activity of the activated kinase. Alternate in vitro assays quantitate the ability of the inhibitor to bind to the protein kinase and may be measured either by radiolabelling the inhibitor prior to binding, isolating the inhibitor/kinase complex and determining the amount of radiolabel bound, or by running a competition experiment where new inhibitors are incubated with the kinase bound to known radioligands. Detailed conditions for assaying a compound utilized in this invention as an inhibitor of ATR is set forth in the Examples below.

**[0273]** Another aspect of the invention provides a method for modulating enzyme activity by contacting a compound of formula I with ATR kinase.

#### Methods of Treatment

**[0274]** In one aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where ATR kinase is implicated in the disease state. In another aspect, the present invention provides a method for treating or lessening the severity of an ATR kinase disease, condition, or disorder where inhibition of enzymatic activity is implicated in the treatment of the disease. In another aspect, this invention provides a method for treating or lessening the severity of a disease, condition, or disorder with compounds that inhibit enzymatic activity by binding to the ATR kinase. Another aspect provides a method for treating or lessening the severity of a kinase disease, condition, or disorder by inhibiting enzymatic activity of ATR kinase with an ATR kinase inhibitor.

**[0275]** One aspect of the invention relates to a method of inhibiting ATR kinase activity in a patient, which method comprises administering to the patient a compound of formula I, or a composition comprising said compound. In some embodiments, said method is used to treat or prevent a condition selected from proliferative and hyperproliferative diseases, such as cancer. In another embodiment, said method is used to treat or prevent HIV, hepatitis, adenovirus, or psoriasis. In some embodiments, said method is used to treat or prevent HIV 1, hepatitis B, adenovirus, or psoriasis.

**[0276]** Another aspect of this invention provides a method for treating, preventing, or lessening the severity of proliferative or hyperproliferative diseases comprising administering an effective amount of a compound, or a pharmaceutically acceptable composition comprising a compound, to a subject in need thereof. In some embodiments, said subject is a patient. The term "patient", as used herein, means an animal, preferably a human.

**[0277]** In some embodiments, said method is used to treat or prevent cancer. In some embodiments, said method is used to treat or prevent a type of cancer with solid tumors. In yet another embodiment, said cancer is selected from the following cancers: Oral: buccal cavity, lip, tongue, mouth, pharynx; Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell or epidermoid, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal:

esophagus (squamous cell carcinoma, larynx, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel or small intestines (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel or large intestines (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma), colon, colon-rectum, colorectal; rectum, Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma, biliary passages; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondroblastoma), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma), breast; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, keratoacanthoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis, Thyroid gland: papillary thyroid carcinoma, follicular thyroid carcinoma; medullary thyroid carcinoma, multiple endocrine neoplasia type 2A, multiple endocrine neoplasia type 2B, familial medullary thyroid cancer, pheochromocytoma, paraganglioma; and Adrenal glands: neuroblastoma. In some embodiments, the cancer is selected from the cancers described herein. In some embodiments, said cancer is lung cancer, head and neck cancer, pancreatic cancer, gastric cancer, or brain cancer.

**[0278]** In certain embodiments, an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective in order to treat said disease. The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of said disease.

**[0279]** One aspect provides a method for inhibiting ATR in a patient comprising administering a compound of formula I as described herein. Another embodiment provides a method

of treating cancer comprising administering to a patient a compound of formula I, wherein the variables are as defined herein.

**[0280]** Some embodiments comprising administering to said patient an additional therapeutic agent selected from a DNA-damaging agent; wherein said additional therapeutic agent is appropriate for the disease being treated; and said additional therapeutic agent is administered together with said compound as a single dosage form or separately from said compound as part of a multiple dosage form.

**[0281]** In some embodiments, said DNA-damaging agent is selected from ionizing radiation, radiomimetic neocarzinostatin, a platinating agent, a Topo I inhibitor, a Topo II inhibitor, an antimetabolite, an alkylating agent, an alkyl sulphonates, an antimetabolite, or an antibiotic. In other embodiments, said DNA-damaging agent is selected from ionizing radiation, a platinating agent, a Topo I inhibitor, a Topo II inhibitor, or an antibiotic. In yet other embodiments, said DNA-damaging agent is a platinating agent or ionizing radiation.

**[0282]** Examples of Platinating agents include Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, Satraplatin and other derivatives. Other platinating agents include Lobaplatin, and Triplatin. Other platinating agents include Tetranitrate, Pico-platin, Satraplatin, ProLindac and Aroplatin.

**[0283]** Examples of Topo I inhibitor include Camptothecin, Topotecan, irinotecan/SN38, rubitecan and other derivatives. Other Topo I inhibitors include Belotecan.

**[0284]** Examples of Topo II inhibitors include Etoposide, Daunorubicin, Doxorubicin, Aclarubicin, Epirubicin, Idarubicin, Amrubicin, Pirarubicin, Valrubicin, Zorubicin and Teniposide.

**[0285]** Examples of Antimetabolites include members of the Folic family, Purine family (purine antagonists), or Pyrimidine family (pyrimidine antagonists). Examples of the Folic family include methotrexate, pemetrexed and relatives; examples of the Purine family include Thioguanine, Fludarabine, Cladribine, 6-Mercaptopurine, and relatives; examples of the Pyrimidine family include Cytarabine, gemcitabine, 5-Fluorouracil (5FU) and relatives. Some other specific examples of antimetabolites include Aminopterin, Methotrexate, Pemetrexed, Raltitrexed, Pentostatin, Cladribine, Clofarabine, Fludarabine, Thioguanine, Mercaptopurine, Fluorouracil, Capecitabine, Tegafur, Carmofur, Floxuridine, Cytarabine, Gemcitabine, Azacitidine and Hydroxyurea.

**[0286]** Examples of alkylating agents include Nitrogen mustards, Triazenes, alkyl sulphonates, Procarbazine and Aziridines. Examples of Nitrogen mustards include Cyclophosphamide, Melphalan, Chlorambucil and relatives; examples of nitrosoureas include Carmustine; examples of triazenes include Dacarbazine and temozolomide; examples of alkyl sulphonates include Busulfan. Other specific examples of alkylating agents include Mechlorethamine, Cyclophosphamide, Ifosfamide, Trofosfamide, Chlorambucil, Melphalan, Prednimustine, Bendamustine, Uramustine, Estramustine, Carmustine, Lomustine, Semustine, Fotemustine, Nimustine, Ranimustine, Streptozocin, Busulfan, Mannosulfan, Treosulfan, Carboquone, ThioTEPA, Triaziquone, Triethylenemelamine, Procarbazine, Dacarbazine,

Temozolomide, Altretamine, Mitobronitol, Actinomycin, Bleomycin, Mitomycin and Plicamycin.

**[0287]** Examples of antibiotics include Mitomycin, Hydroxyurea; Anthracyclines, Anthracenediones, *Streptomyces* family. Examples of Anthracyclines include doxorubicin, daunorubicin, epirubicin and other derivatives; examples of Anthracenediones include Mitoxantrone and relatives; examples of *Streptomyces* family include Bleomycin, Mitomycin C, and actinomycin.

**[0288]** In certain embodiments, said platinating agent is Cisplatin or Oxaliplatin; said Topo I inhibitor is Camptothecin; said Topo II inhibitor is Etoposide; and said antibiotic is Mitomycin. In other embodiments, said platinating agent is selected from Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, or Satraplatin; said Topo I inhibitor is selected from Camptothecin, Topotecan, irinotecan/SN38, rubitecan; said Topo II inhibitor is selected from Etoposide; said antimetabolite is selected from a member of the Folic Family, the Purine Family, or the Pyrimidine Family; said alkylating agent is selected from nitrogen mustards, nitrosoureas, triazenes, alkyl sulphonates, Procarbazine, or aziridines; and said antibiotic is selected from Hydroxyurea, Anthracyclines, Anthracenediones, or *Streptomyces* family.

**[0289]** Another embodiment provides a method of promoting cell death in cancer cells comprising administering to a patient a compound of formula I, or a composition comprising said compound.

**[0290]** Yet another embodiment provides a method of preventing cell repair of DNA damage in cancer cells comprising administering to a patient a compound of formula I, or a composition comprising said compound. Yet another embodiment provides a method of preventing cell repair caused by of DNA damage in cancer cells comprising administering to a patient a compound of formula I, or composition comprising said compound.

**[0291]** Another embodiment provides a method of sensitizing cells to DNA damaging agents comprising administering to a patient a compound of formula I, or a composition comprising said compound.


**[0292]** In some embodiments, said cell is a cancer cell having defects in the ATM signaling cascade. In some embodiments, said defect is altered expression or activity of one or more of the following: ATM, p53, CHK2, MRE11, RAD50, NBS1, 53BP1, MDC1 or H2AX. In other embodiments, said cell is a cancer cell expressing DNA damaging oncogenes. In some embodiments, said cancer cell has altered expression or activity of one or more of the following: K-Ras, N-Ras, H-Ras, Raf, Myc, Mos, E2F, Cdc25A, CDC4, CDK2, Cyclin E, Cyclin A and Rb.

**[0293]** Yet another embodiment provides use of a compound of formula I as a radio-sensitizer or a chemo-sensitizer. Yet other embodiment provides use of a compound of formula I as a single agent (monotherapy) for treating cancer. In some embodiments, the compounds of formula I are used for treating patients having cancer with a DNA-damage response (DDR) defect. In other embodiments, said defect is a mutation or loss of ATM, p53, CHK2, MRE11, RAD50, NBS1, 53BP1, MDC1, or H2AX.

## Synthetic Methodology

[0294]

GENERAL SCHEME A



[0295] The compounds of this invention may be made according to the general schemes shown above. Unless otherwise indicated, all variables in the schemes are as defined herein.

[0296] Intermediate **ii**, rapidly obtained by bromination of aminoester **i**, can be used as a building block to generate

various advanced intermediates **iii.1-iii.4** useful for the preparation of the compounds of this invention. Compound **ii** can be used to generate advanced intermediate **iii.1** to make compounds wherein  $R^2$  is linked to the aminopyrimidine moiety through a nitrogen atom. This can be done by (a) using the bromine atom as a handle for nucleophilic displacement with a cyclic amine followed by (b) an ester hydrolysis to generate


acid iii.1. The carboxylic acid moiety can be manipulated in multiple ways, using methodology known in the art to generate compounds iv of this invention where  $n$  is 0 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring. For compounds iv where an H-bond acceptor is not yet present, ring Q can be further manipulated and derivatised into compounds v of this invention that bear a H-bond acceptor feature. Additionally, the carboxylic acid group in iii.1 can be transformed into amides vi of this invention where  $n$  is 1 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring. This can be done using methods known in the art for amide bond formation. For compounds vi where an H-bond acceptor is not yet present, ring Q can be further manipulated and derivatised into compounds vii of this invention that bear a H-bond acceptor feature.


[0297] In a similar step sequence compound ii can be used to generate advanced intermediate iii.4 to make compounds where  $R^2$  is linked to the aminopyrazine moiety through a carbon atom. This can be done by (a) using the bromine atom as a handle for metal-catalysed cross-coupling (eg palladium catalysed coupling with a boronic acid) followed by (b) an ester hydrolysis to generate acid iii.4. The carboxylic acid moiety can be manipulated in multiple ways, using methodology known in the art to generate compounds viii of this invention where  $n$  is 0 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring. For compounds viii where an H-bond acceptor is not yet present, ring Q can be further manipulated and derivatised into compounds ix of this invention that bear a H-bond acceptor feature. Additionally, the carboxylic acid group in iii.4 can be transformed into amides x of this invention where  $n$  is 1 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring; This has been performed using methods known in the art for amide bond formation. For compounds x where an H-bond acceptor is not yet present, ring Q can be further manipulated and derivatised into compounds xi of this invention that bear a H-bond acceptor feature.

[0298] Alternatively, the synthetic sequence can be modified and compound ii can be used to prepare advanced intermediate iii.2 and iii.3;

[0299] Compound iii.2 is prepared by hydrolysis of the ester in ii, followed by a variety of methodologies known in the art to transform the carboxylic acid moiety into advanced intermediate iii.2 where  $n$  is 0 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring. The bromine atom in iii.2 can then be used as a handle for nucleophilic displacement with a cyclic amine to generate compounds iv of this invention where  $R^2$  is linked to the aminopyrazine moiety through a nitrogen atom. The bromine atom in iii.2 can also be used as a handle for metal-catalysed cross-coupling (e.g., palladium-catalyzed coupling with a boronic acid) to generate compounds viii of this invention where  $R^2$  is linked to the aminopyrazine moiety through a carbon atom

[0300] Hydrolysis of the ester in ii, followed by amide bond formation using methods known in the art afforded advanced intermediate iii.3 where  $n$  is 1 and  $R^1$  is a monocyclic or bicyclic (hetero)aromatic ring. The bromine atom in iii.3 can then be used as a handle for nucleophilic displacement with a cyclic amine to generate compounds vi of this invention where  $R^2$  is linked to the aminopyrazine moiety through a nitrogen atom. The bromine atom in iii.3 can also be used as a handle for metal-catalyzed cross-coupling (e.g., palladium catalyzed coupling with a boronic acid) to generate compounds x of this invention where  $R^2$  is linked to the aminopyrazine moiety through a carbon atom.





[0301] Scheme B depicts a general method for making compounds of formula III wherein one of the  $J^1$  groups is  $\text{CH}_2\text{N}(\text{R}^B)_2$ , wherein  $\text{R}^B$  is defined such that  $\text{CH}_2\text{N}(\text{R}^B)_2$  is within the scope of  $J^{Q3}$  as defined herein. The hydrazide of formula B-ii is formed from an ester of formula B-i in the presence of  $\text{NH}_2\text{NH}_2$  and a suitable solvent. The result carbohydrazide of formula B-ii can then be combined with a variety of different carboxylic acids to form oxadiazoles of formula B-iii, which can undergo a displacement reaction to form amines of formula B-iv. The amines of formula B-iv can then be protected with suitable amine protecting groups ("PG") to form compounds of formula B-v, which can either undergo an  $\text{S}_{\text{N}}\text{Ar}$  reaction or a metal-mediated coupling reaction to form compounds of formula B-vi. Deprotection of the nitrogen protecting group is then done under suitable deprotecting conditions to form compounds of formula III.

[0302] Suitable hydrazide formation solvents include, but are not limited to alcohol solvents, such as EtOH and MeOH. Suitable condensation reaction conditions include, but are not limited to the following:  $\text{PPh}_3\text{Br}_2$  in the presence of  $\text{CH}_3\text{CN}$  and DIEA.

[0303] The  $\text{S}_{\text{N}}\text{Ar}$  reaction is used to prepare compounds wherein  $\text{R}^2$  is bonded to the pyrazine via a nitrogen atom. The bromo-aminopyrazine of B-v is combined with suitable  $\text{R}^2$  cyclic amines under suitable reduction  $\text{S}_{\text{N}}\text{Ar}$  conditions to form compounds of formula B-v. Suitable reduction  $\text{S}_{\text{N}}\text{Ar}$  conditions include, but are not limited to a suitable solvent (e.g., DMSO or DMF), a suitable base (e.g., TEA), and heat.

[0304] Metal-mediated coupling conditions are used to prepare compounds wherein  $\text{R}^2$  is bonded to the pyrazine via a carbon atom. The bromo-aminopyrazine of B-v is combined with suitable  $\text{R}^2$  boronic acids or esters under suitable metal-mediated coupling conditions to form compounds of formula B-v. Suitable metal-mediated coupling conditions include, but are not limited to, a catalyst (such as  $\text{Pd}(\text{PPh}_3)_4$ ), a salt (such as  $\text{Na}_2\text{CO}_3$ ), and suitable solvents (such as  $\text{CH}_3\text{CN}$ , DMF, DME, or water).

[0305] Suitable deprotection conditions, include, but are not limited to TFA/DCM or dioxane/HCl for a BOC protecting group.

#### EXAMPLES

[0306] The compounds of this invention may be prepared in light of the specification using steps generally known to those of ordinary skill in the art. Those compounds may be analyzed by known methods, including but not limited to LCMS (liquid chromatography mass spectrometry) and NMR (nuclear magnetic resonance). Compounds of this invention may be also tested according to these examples. It should be understood that the specific conditions shown below are only examples, and are not meant to limit the scope of the conditions that can be used for making, analyzing, or testing the

compounds of this invention. Instead, this invention also includes conditions known to those skilled in that art for making, analyzing, and testing the compounds of this invention.

#### HPLC Methods

[0307] As used herein, the term "Rt(min)" refers to the HPLC retention time, in minutes, associated with the compound. Unless otherwise indicated, the HPLC method utilized to obtain the reported retention time is as follows:

[0308] Column: ACE C8 column, 4.6x150 mm

[0309] Gradient: 0-100% acetonitrile+methanol 60:40 (20 mM Tris phosphate)

[0310] Flow rate: 1.5 mL/minute

[0311] Detection: 225 nm.

#### HNMR Methods

[0312]  $^1\text{H}$ -NMR spectra were recorded at 400 MHz using a Bruker DPX 400 instrument.

#### Mass Spectrometry Methods

[0313] Mass spec. samples were analyzed on a MicroMass Quattro Micro mass spectrometer operated in single MS mode with electrospray ionization. Samples were introduced into the mass spectrometer using chromatography. Mobile phase for all mass spec. analyses consisted of 10 mM pH 7 ammonium acetate and a 1:1 acetonitrile-methanol mixture, column gradient conditions are 5%-100% acetonitrile-methanol over 3.5 mins gradient time and 5 mins run time on an ACE C8 3.0x75 mm column. Flow rate is 1.2 ml/min.


[0314] The following compounds of formula I were prepared and analyzed as follows.


#### Example 1

1-(4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-1,4-diazepan-1-yl)ethanone (Compound I-1)

[0315]

#### SCHEME I





## Method A

Step 1: Methyl  
3-amino-6-bromopyrazine-2-carboxylate

**[0316]** A mixture of methyl 3-aminopyrazine-2-carboxylate (8.35 g, 54.53 mmol) and N-bromo-succinimide (9.705 g, 54.53 mmol) was stirred in MeCN (100 mL) at room temp overnight. The resultant precipitate was filtered, washed with MeCN and dried to give the desired product as a yellow solid (11.68 g, 92% Yield) <sup>1</sup>H NMR (400.0 MHz, DMSO) 3.85 (s, 3H), 7.55 (br s, 2H) and 8.42 (s, 1H) ppm; MS (ES<sup>+</sup>) 233

Step 2: 3-amino-6-bromopyrazine-2-carboxylic acid

**[0317]** A mixture of methyl 3-amino-6-bromo-pyrazine-2-carboxylate (3 g, 12.93 mmol) and lithium hydroxide (1.548 g, 64.65 mmol) in MeOH (11.74 mL) and H<sub>2</sub>O (11.74 mL) was heated to 90° C. for 2 hours. The reaction mixture was allowed to cool, neutralised with HCl and diluted with water, and the resultant precipitate collected by filtration (2.2 g, 78% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) 7.57 (br s, 2H) and 8.39 (s, 1H), 13.41 (br s, 1H) ppm.

Step 3: 5-bromo-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine

**[0318]** A mixture of 3-amino-6-bromo-pyrazine-2-carboxylic acid (2.5 g, 11.47 mmol), 4-methylbenzene-1,2-diamine (1.401 g, 11.47 mmol), diethoxyphosphoryl-formonitrile (2.058 g, 1.871 mL, 12.62 mmol) and triethylamine (2.321 g, 3.197 mL, 22.94 mmol) was heated in DME (75.00 mL) at 170° C. in the microwave for 1 hour. The reaction mixture was diluted with EtOAc, filtered and washed with water and brine, dried over MgSO<sub>4</sub> and concentrated to a brown solid. The mixture was slurried in DCM and treated with diethyl ether/petroleum ether. The resultant solid was washed with ether and dried to afford the product as an orange/brown solid (1.6 g, 46% Yield). MS (ES<sup>+</sup>) 305.

Step 4: 1-(4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-1,4-diazepan-1-yl)ethanone  
(Compound I-1)

**[0319]** A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-amine (100 mg, 0.3288 mmol) and 1-(1,4-diazepan-1-yl)ethanone (701 mg, 4.932 mmol) in NMP (200.0 μL) was heated at 190° C. in the microwave for 2 hours. The reaction mixture was partitioned between DCM and water and the organics separated and concentrated to dryness. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μm, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound (67.7 mg, 54% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.80-1.94 (2H, m), 1.78 (3H, s), 2.43-2.46 (3H, s), 3.40 (2H, m), 3.63-3.83 (5H, m), 3.92 (1H, m), 7.04-7.11 (3H, m), 7.38-7.59 (2H, m), 7.90 (1H, m), 12.37 (1H, m) ppm; MS (ES<sup>+</sup>) 366

**[0320]** The following compounds were all prepared using Method A.

**[0321]** Compound I-2 (1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidin-4-yl)methanol

**[0322]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.22 (2H, m), 1.59 (1H, m), 1.79 (2H, m), 2.45 (3H, s), 2.72 (2H, m), 3.31 (1H, m), 4.30 (2H, d), 4.52 (1H, s), 7.04 (1H, m), 7.18 (2H, s), 7.37-7.57 (3H, m), 7.98 (1H, s), 12.44 (1H, s) ppm; MS (ES<sup>+</sup>) 339

**[0323]** Compound I-3 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-morpholinopyrazin-2-amine

**[0324]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.45 (3H, s), 3.45 (4H, m), 3.79 (4H, m), 7.08 (1H, s), 7.27-7.58 (4H, m), 7.98 (1H, s), 12.50 (1H, s) ppm; MS (ES<sup>+</sup>) 311

**[0325]** Compound I-4 (S)-N-(1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)ethanamide

**[0326]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.22 (2H, m), 1.59 (1H, m), 1.79 (2H, m), 2.45 (3H, s), 2.72 (2H, m), 3.31 (1H, m), 4.30 (2H, d), 4.52 (1H, s), 7.04 (1H, m), 7.18 (2H, s), 7.37-7.57 (3H, m), 7.98 (1H, s), 12.44 (1H, s) ppm; MS (ES<sup>+</sup>) 352

**[0327]** Compound I-5 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidin-4-ol

**[0328]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.48 (2H, m), 1.86 (2H, m), 2.43, 2.45 (3H, s), 3.02 (2H, m), 3.69 (1H, m), 4.02 (2H, m), 4.72 (1H, s), 7.09 (1H, d), 7.18 (2H, s), 7.36-7.57 (2H, m), 7.99 (1H, s), 12.45 (1H, s) ppm; MS (ES<sup>+</sup>) 325

**[0329]** Compound I-6 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.31 (3H, t), 2.50 (3H, s), 3.20 (2H, m), 3.35 (4H, m), 3.66 (4H, m), 7.12 (1H, m), 7.35-7.66 (4H, m), 8.09 (1H, s), 12.61 (1H, m) ppm; MS (ES<sup>+</sup>) 402

**[0330]** Compound I-7 4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-2-one; MS (ES<sup>+</sup>) 324

**[0331]** Compound I-8 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-thiomorpholinopyrazin-2-amine; MS (ES<sup>+</sup>) 327

**[0332]** Compound I-9 (R)-N-(1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)ethanamide; MS (ES<sup>+</sup>) 352


**[0333]** Compound I-10 1-(4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)ethanone; MS (ES<sup>+</sup>) 352

- [0334] Compound I-11 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidin-4-one; MS (ES+) 323
- [0335] Compound I-12 5-(4-methoxypiperidin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES+) 339
- [0336] Compound I-13 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N-methylpiperidine-4-carboxamide; MS (ES+) 366
- [0337] Compound I-14 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidine-4-carboxamide; MS (ES+) 352
- [0338] Compound I-30 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-phenylpiperazin-1-yl)pyrazin-2-amine 1H NMR (400.0 MHz, DMSO)  $\delta$  2.49, 2.50 (3H, 2xs), 3.31 (4H, m), 3.64 (4H, m), 6.91 (1H, m), 7.05, 7.07 (2H, 2xs), 7.17, 7.25 (4H, m), 7.35-7.60 (2H, m), 8.05 (1H, 2xs), 12.55, 12.58 ppm; MS (ES+) 386.06
- [0339] Compound I-31 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-(trifluoromethyl) piperidin-1-yl)pyrazin-2-amine; MS (ES+) 377.05
- [0340] Compound I-32 ethyl 4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate; MS (ES+) 382.05
- [0341] Compound I-33 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(1,4-dioxo-8-azaspiro[4.5]decan-8-yl)pyrazin-2-amine; MS (ES+) 367.14
- [0342] Compound I-34 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidine-4-carbonitrile
- [0343] 1H NMR (400.0 MHz, DMSO)  $\delta$  1.90 (2H, m), 2.07 (2H, m), 2.49, 2.52 (3H, 2xs), 3.16 (1H, m), 3.37 (2H, m), 3.86 (2H, m), 7.10, 7.16 (1H, 2xd), 7.31 (2H, br s), 7.42, 7.56 (1H, 2xs), 7.52, 7.64 (1H, 2xd), 8.07 (1H, s), 12.54 (1H, 2xs) ppm; MS (ES+) 334.09
- [0344] Compound I-35 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-(pyrimidin-2-yl)piperazin-1-yl)pyrazin-2-amine; MS (ES+) 388.17
- [0345] Compound I-43 N-(1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidin-4-yl)ethanamide; MS (ES+) 366.08
- [0346] Compound I-44 1-(1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-4-phenylpiperidin-4-yl)ethanone; MS (ES+) 427.2
- [0347] Compound I-45 5-(4-(2-fluorophenyl)piperazin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES+) 404.16
- [0348] Compound I-46 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-(thiazol-2-yl)piperazin-1-yl)pyrazin-2-amine; MS (ES+) 393.1
- [0349] Compound I-47 5-(4-(3-fluorophenyl)piperazin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES+) 404.16
- [0350] Compound I-48 5-(4-(3-methoxyphenyl)piperazin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; 1H NMR (400.0 MHz, DMSO)  $\delta$  2.44+2.45 (3H, 2xs), 3.32 (4H, m), 3.63 (4H, m), 3.75 (3H, s), 6.41 (1H, dd), 6.56 (1H, s), 6.65 (1H, d), 7.06+7.11 (1H, 2xd), 7.16 (1H, t), 7.28 (2H, br s), 7.37+7.51 (1H, 2xs), 7.47+7.59 (1H, 2xd), 8.05, 8.06 (1H, 2xs), 12.12 (1H, 2xs); MS (ES+) 416.17
- [0351] Compound I-49 5-(4-(dimethylamino)piperidin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES+) 352.11
- [0352] Compound I-50 2-(4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)benzonitrile; MS (ES+) 411.13
- [0353] Compound I-51 5-(3-(dimethylamino)piperidin-1-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES+) 352.11
- [0354] Compound I-52 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-(3-(trifluoromethyl) pyridin-2-yl)piperazin-1-yl)pyrazin-2-amine; MS (ES+) 455.04
- [0355] Compound I-53 4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl(furan-2-yl)methanone; MS (ES+) 404.13
- [0356] Compound I-56 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-amine; 1H NMR (400.0 MHz, DMSO)  $\delta$  1.45 (3H, t), 3.17 (2H, q), 3.37-3.4 (4H, m), 3.65-3.7 (4H, m), 7.3-7.35 (1H, m), 8.12 (1H, d), 8.15 (1H, s), 8.42-8.45 (1H, m) ppm; MS (ES+) 389
- [0357] Compound I-63 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperidin-4-yl(phenyl)methanone; 1H NMR (400.0 MHz, DMSO)  $\delta$  1.70 (2H, m), 1.92 (2H, m), 2.43+2.45 (3H, 2xs), 2.99 (2H, m), 2.72 (1H, m), 4.34 (2H, m), 7.07 (1H, dd), 7.23 (2H, br s), 7.35-7.67 (5H, m), 8.03 (3H, m), 12.50 (1H, d) ppm; MS (ES+) 413.1
- [0358] Compound I-64 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(4-(pyridin-2-yl)piperazin-1-yl)pyrazin-2-amine; 1H NMR (400.0 MHz, DMSO)  $\delta$  2.43+2.46 (3H, 2xs), 3.60 (4H, m), 3.68 (4H, m), 6.68 (1H, m), 6.95 (1H, d), 7.09 (1H, dd), 7.28 (2H, br s), 7.51-7.61 (3H, m), 8.06 (1H, d), 8.17 (1H, d), 12.50 (1H, d) ppm; MS (ES+) 387.06
- [0359] Compound I-65 1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-1,4-diazepan-5-one; 1H NMR (400.0 MHz, DMSO)  $\delta$  2.43+2.46 (3H, 2xs), 2.55 (2H, m), 3.24 (2H, m), 3.80 (4H, m), 7.06 (1H, dd), 7.12 (2H, br s), 7.36-7.65 (3H, m), 7.99 (1H, s), 12.40 (1H, s) ppm; MS (ES+) 338.03
- [0360] Compound I-70 (S)-1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N-methylpyrrolidine-3-carboxamide; 1H NMR (400.0 MHz, DMSO)  $\delta$  2.15 (2H, m), 2.43+2.45 (3H, 2xs), 2.63+2.65 (3H, 2xs), 3.06 (1H, m), 3.46 (2H, m), 3.63 (1H, m), 3.77 (1H, m), 7.01 (3H, m), 7.35-7.64 (2H, m), 7.69 (1H, s), 8.05 (1H, m), 12.35 (1H, m) ppm; MS (ES+) 352.03
- [0361] Compound I-75 1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-2-one 1H NMR (400.0 MHz, DMSO)  $\delta$  2.05-2.15 (2H, m), 2.62 (2H, t), 4.18 (2H, t), 7.25-7.3 (2H, m), 7.7-7.75 (2H, m), 9.03 (1H, s), 13.15 (1H, vbrs) ppm; MS (ES+) 295
- [0362] Compound I-77 (R)-1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N-methylpyrrolidine-3-carboxamide; MS (ES+) 352.08
- [0363] Compound I-158 5-(5-(ethylsulfonyl)hexahydro-pyrrolo[3,4-c]pyrrol-2(1H)-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine 1H NMR (400.0 MHz, DMSO)  $\delta$  1.23 (3H, m), 2.43+2.46 (3H, 2xs), 3.11-3.20 (6H, m), 3.34-3.47 (2H, m), 3.61 (4H, m), 7.00-7.10 (3H, m), 7.35-7.59 (2H, m), 7.73 (1H, br s), 12.37+12.40 (1H, 2xs) ppm; MS (ES+) 428.01 For compound I-281 below, please see Tables 7 and 8 for analytical data.
- [0364] Compound I-281 5-(5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)-3-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine

## Example 2

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(1-methyl-1H-benzo[d]imidazol-2-yl)pyrazine-2-amine (Compound I-15)

[0365]



## Method B

## Step 1:

3-amino-6-bromo-N-methylpyrazine-2-carboxamide

[0366] Methyl 3-amino-6-bromo-pyrazine-2-carboxylate (7.5 g, 32.32 mmol) was suspended in 40% aqueous methylaniline and the resulting mixture stirred vigorously at RT for 4 hours. The resultant precipitate was collected, washed with water and dried to give the desired product (6.73 g, 90% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.76 (s, 3H), 7.75 (br s, 2H), 8.34 (s, 1H), 8.58 (b rs, 1H) ppm; MS (ES<sup>+</sup>) 232.

## Step 2: 6-bromo-3-methylpteridin-4(3H)-one

[0367] A mixture of triethyl orthoformate (32.23 g, 36.17 mL, 217.5 mmol) and acetic anhydride (38.49 g, 35.57 mL,

377.0 mmol) was treated with 3-amino-6-bromo-N-methylpyrazine-2-carboxamide (6.7 g, 29.00 mmol) and the resulting solution was heated at reflux for 2 hours. After cooling to RT, the resultant precipitate was collected, washed with EtOAc and dried to give the product as a beige solid (5.52 g, 79% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 3.54 (s, 3H), 8.72 (s, 1H), 9.17 (s, 1H) ppm; MS (ES<sup>+</sup>) 242.

## Step 3: 6-(4-(ethylsulfonyl)piperazin-1-yl)-3-methylpteridin-4(3H)-one

[0368] A solution of 6-bromo-3-methylpteridin-4-one (5.516 g, 22.88 mmol) in 2-methoxyethanol (50.81 mL) was treated with 1-ethylsulfonylpiperazine (6.118 g, 34.32 mmol) and the resulting suspension heated for 2 hours at 100° C. The reaction mixture was cooled in an ice-water bath and resultant precipitate collected and washed with cold methanol to give the product as a yellow solid (5.02 g, 65% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.25 (t, 3H), 3.10 (q, 2H), 3.34 (m, 4H), 3.49 (s, 3H), 3.85 (m, 4H), 8.37 (s, 1H), 8.84 (s, 1H) ppm; MS (ES<sup>+</sup>) 339

## Step 4: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid

[0369] A solution of 6-(4-(ethylsulfonyl)piperazin-1-yl)-3-methylpteridin-4-one (5 g, 14.78 mmol) in methanol (51.00 mL) was treated with sodium hydroxide (45.92 mL of 10% w/v, 114.8 mmol) and the reaction mixture stirred at RT overnight. The reaction mixture was concentrated in vacuo and the residue diluted with water and neutralised with formic acid. The resulting precipitate was filtered off and the filtrate acidified to pH4 by the addition of formic acid and extracted with dichloromethane. The combined organics were washed with brine, dried over MgSO<sub>4</sub> and concentrated in vacuo to give the product as a bright yellow solid (4.66 g, 52% Yield). <sup>1</sup>H NMR (400.0 MHz, CDCl<sub>3</sub>) δ 1.45 (3H, t), 3.05 (2H, q), 3.54 (8H, m), 6.05 (2H, br s), 8.21 (1H, s) ppm; MS (ES<sup>+</sup>) 316

## Step 5: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(1-methyl-1H-benzo[d]imidazol-2-yl)pyrazine-2-amine (Compound I-15)

[0370] A mixture of 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid (100 mg, 0.3171 mmol), N-methylbenzene-1,2-diamine (42.61 mg, 39.64 μL, 0.3488 mmol), diethoxyphosphorylformonitrile (56.89 mg, 51.72 μL, 0.3488 mmol) and triethylamine (64.17 mg, 88.39 μL, 0.6342 mmol) in DME (3.000 mL) was heated in the microwave at 150° C. for 20 mins. The reaction mixture was allowed to cool and concentrated in vacuo. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound as a yellow solid (79 mg, 60% Yield).

[0371] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.24 (3H, t), 3.13 (2H, q), 3.31 (masked signal), 3.50 (4H, m), 4.23 (3H, s), 7.30 (1H, m), 7.36 (1H, m), 7.44 (2H, br s), 7.67 (1H, m), 7.73 (1H, m), 8.10 (1H, s) ppm; MS (ES<sup>+</sup>) 402

[0372] The following compounds were all prepared using the method described for Compound I-15 above.

[0373] Compound I-16 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-amine was also prepared using Method B, Steps 1-5. <sup>1</sup>H NMR (400.0 MHz,

DMSO) d 1.25 (3H, t), 3.12 (2H, q), 3.34 (4H, m), 3.60 (4H, m), 7.25 (2H, m), 7.66 (2H, m), 8.06 (1H, s) ppm; MS (ES<sup>+</sup>) 388


**[0374]** Compound I-308 5-(4-ethylsulfonylpiperazin-1-yl)-3-(1-phenylbenzimidazol-2-yl)pyrazin-2-amine 1H NMR (400.0 MHz, CDCl<sub>3</sub>) d 1.37 (t, 3H), 2.76 (t, 4H), 2.93 (q, 2H), 3.11-3.14 (m, 4H), 7.04 (d, 1H), 7.24-7.28 (m, 3H), 7.32-7.35 (m, 3H), 7.43-7.54 (m, 3H), 7.64 (s, 1H) and 7.83 (d, 1H); MS (ES<sup>+</sup>) 464.0

### Example 3

3-(benzo[d]thiazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine (Compound I-17)

**[0375]**

#### SCHEME III



Compound I-17

**[0376]** Compound I-17 was prepared using Method B, Steps 1-4 followed by Method C, Steps 1 and 2.

#### Method C


Step 1: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid

**[0377]** Thionyl chloride (1.887 g, 1.157 mL, 15.86 mmol) was added to a solution of 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid (1.25 g, 3.964 mmol) in chloroform (50 mL) and DMF (1 mL) and the resulting dark solution heated under reflux for 1 hour. The reaction mixture was allowed to cool to RT, concentrated in vacuo and taken through crude to the next stage.

Step 2: 3-(benzo[d]thiazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine (Compound I-17)

**[0378]** A mixture of 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid (100 mg, 0.2996 mmol) and 2-aminobenzenethiol (112.5 mg, 0.9698 mmol) in acetonitrile was heated at 70° C. overnight. The reaction mixture was allowed to cool to RT, filtered and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were combined but still contained an impurity. The mixture was therefore purified on silica gel by flash column chromatography (50% EtOAc/Hexanes) to afford the title compound as a bright orange solid (23.2 mg, 20% Yield). 1H NMR (400.0 MHz, DMSO) d 1.24 (3H, t), 3.12 (2H, q), 3.36 masked signal, 3.54 (4H, m), 7.39 (2H, br s), 7.44-7.48 (1H, d), 7.52-7.57 (1H, d), 8.06 (1H, m), 8.11 (1H, m), 8.24 (1H, s) ppm; MS (ES<sup>+</sup>) 405

-continued




## Example 4

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(3-phenyl-1,2,4-oxadiazol-5-yl)pyrazin-2-amine (Compound I-19)

[0379]

## SCHEME IV

Compound I-19



[0380] Compound I-19 was prepared using Method B, Steps 1-4 followed by Method D, Step 1.

## METHOD D

Step 1: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazine-2-carboxylic acid (Compound I-19)

[0381] CDI (102.8 mg, 0.6342 mmol) was added to a solution of 3-amino-6-(4-ethylsulfonylpiperazin-1-yl)pyrazine-2-carboxylic acid (100 mg, 0.3171 mmol) and N'-hydroxybenzamide (86.35 mg, 0.6342 mmol) in DMF (3.000 mL) and the resulting solution stirred at RT for 3 hours and then heated at 100° C. overnight. The mixture was allowed to cool, poured in water and extracted with EtOAc. The organic extract was washed with water (2×), brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The resultant

residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound as a bright yellow solid (53.6 mg, 30% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.24 (3H, t), 3.11 (2H, q), 3.33 (4H, m), 3.55 (4H, m), 7.18 (2H, br s), 7.59-7.66 (3H, m), 8.19 (2H, m), 8.42 (1H, s) ppm; MS (ES<sup>+</sup>) 416


## Example 5


5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-18)

[0382]

## SCHEME V

Compound I-18





**[0383]** Compound I-18 was prepared using Method B, Steps 1-4 followed by Method E, Steps 1-2.

#### Method E

Step 1: 3-amino-N'-benzoyl-6-(4-ethylsulfonylpiperazin-1-yl)pyrazine-2-carbohydrazide

**[0384]** TBTU (763.9 mg, 2.379 mmol) and triethylamine (160.5 mg, 221.1  $\mu$ L, 1.586 mmol) were added to a solution of 3-amino-6-(4-ethylsulfonylpiperazin-1-yl)pyrazine-2-carboxylic acid (500 mg, 1.586 mmol) and benzohydrazide (259.1 mg, 1.903 mmol) in DMF and the resulting solution stirred overnight at RT. The mixture was diluted with EtOAc and water and the layers separated. Aqueous layer extracted further with ethyl acetate (2 $\times$ 20 mL) and combined organics washed with water (3 $\times$ 20 mL), dried (MgSO<sub>4</sub>) and concentrated in vacuo. Purified by column chromatography on silica eluting with ethyl acetate. Product fractions combined and concentrated in vacuo to leave the product as a yellow solid (522 mg, 76% Yield).

**[0385]** <sup>1</sup>H NMR (400.0 MHz, DMSO)  $\delta$  1.36 (3H, t), 3.24 (2H, q), 3.41 (4H, m), 3.65 (4H, m), 6.95 (2H, br s), 7.58 (2H, m), 7.65 (1H, m), 8.05 (2H, m), 8.45 (1H, m), 10.41 (1H, s), 10.55 (1H, s) ppm; MS (ES<sup>+</sup>) 434

Step 2: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-18)

**[0386]** A mixture of 3-amino-N'-benzoyl-6-(4-ethylsulfonylpiperazin-1-yl)pyrazine-2-carbohydrazide (50 mg, 0.1153 mmol) in phosphorus oxychloride (884.0 mg, 537.4  $\mu$ L, 5.765 mmol) was heated at 100° C. for 3 hours. The mixture was allowed to cool to RT and added carefully to ice water. The mixture was extracted with EtOAc (3 $\times$ 10 mL) and the combined organics washed with 1N NaOH and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (48.8 mg, 80% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO)  $\delta$  1.24 (3H, t), 3.11 (2H, q), 3.33-3.36 (4H, m), 3.56 (4H, masked signal), 6.95 (2H, br s), 7.63-7.69 (3H, m), 8.11-8.13 (2H, m), 8.28 (1H, s) ppm; MS (ES<sup>+</sup>) 416.01

**[0387]** The following compounds were all prepared using the same method or a similar method to the one described for Compound I-18 above.

**[0388]** Compound I-69 5-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)-N-phenyl-1,3,4-oxadiazol-2-amine <sup>1</sup>H NMR (400.0 MHz, DMSO)  $\delta$  1.24 (t, 3H), 3.17 (d, J=5.2 Hz, 2H), 3.35 (br s, 4H), 3.48-3.50 (m, 4H), 6.87 (s, 2H), 7.03 (t, 1H), 7.35-7.39 (m, 2H), 7.65-7.67 (m, 2H), 8.14 (s, 1H) and 10.89 (s, 1H) ppm; MS (ES<sup>+</sup>) 430.99

**[0389]** Compound I-283 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(3-fluorophenyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 3.11 (q, 2H), 3.33 (m, 4H), 3.56 (m, 4H), 6.97 (br s, 2H), 7.53-7.58 (m, 1H), 7.69-7.75 (m, 1H), 7.86-7.90 (m, 1H), 7.96-7.98 (m, 1H) and 8.29 (1H, s) ppm; MS (ES<sup>+</sup>) 434.16

**[0390]** Compound I-287 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(o-tolyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 2.70 (s, 3H), 3.10 (q, 2H), 3.30 (m, 4H), 3.54 (m, 4H), 6.96 (br s, 2H), 7.45-7.50 (m, 2H), 7.57 (m, 1H), 8.03 (m, 1H) and 8.28 (s, 1H) ppm; MS (ES<sup>+</sup>) 430.20

**[0391]** Compound I-289 3-[5-(2,6-difluorophenyl)-1,3,4-oxadiazol-2-yl]-5-(4-ethylsulfonylpiperazin-1-yl)pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.23 (t, 3H), 3.10 (q, 2H), 3.32 (m, 4H), 3.52 (m, 4H), 6.98 (br s, 2H), 7.45 (m, 1H), 7.77-7.85 (m, 1H) and 8.31 (s, 1H) ppm; MS (ES<sup>+</sup>) 452.12

**[0392]** Compound I-292 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(4-fluorophenyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (m, 3H), 3.11 (t, 2H), 3.31 masked signal, 3.56 (m, 4H), 6.96 (br s, 2H), 7.50 (m, 2H), 8.18 (m, 2H) and 8.28 (1H, s) ppm; MS (ES<sup>+</sup>) 434.12

**[0393]** Compound I-294 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(3-methyl-2-thienyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.23 (t, 3H), 2.63 (s, 3H), 3.10 (q, 2H), 3.30 (m, 4H), 3.54 (m, 4H), 6.92 (br s, 2H), 7.19 (d, 1H), 7.87 (d, 1H) and 8.27 (s, 1H) ppm; MS (ES<sup>+</sup>) 436.11

**[0394]** Compound I-297 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(2-furyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 3.11 (q, 2H), 3.30 (m, 4H), 3.52 (m, 4H), 6.85 (m, 1H), 6.95 (br s, 2H), 7.47 (m, 1H), 8.12 (m, 1H) and 8.27 (1H, s) ppm; MS (ES<sup>+</sup>) 406.15

**[0395]** Compound I-299 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(2-fluorophenyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 3.11 (t, 2H), 3.30 (m, 4H), 3.53 (m, 4H), 6.97 (br s, 2H), 7.47-7.56 (m, 2H), 7.75 (m, 1H), 8.11 (m, 1H) and 8.29 (1H, s) ppm; MS (ES<sup>+</sup>) 434.15


**[0396]** Compound I-312 5-(4-ethylsulfonylpiperazin-1-yl)-3-[5-(2-thienyl)-1,3,4-oxadiazol-2-yl]pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 3.11 (q, 2H), 3.34 (m, 4H), 3.55 (m, 4H), 6.93 (br s, 2H), 7.35 (m, 1H), 7.94 (dd, 1H), 8.01 (dd, 1H) and 8.27 (1H, s) ppm; MS (ES<sup>+</sup>) 422.10

## Example 6

3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-phenylpyrazine-2-carboxamide (Compound I-20)

[0397]

## SCHEME VI



[0398] Compound I-20 was prepared using Method A, Steps 1-2 followed by Method F, Steps 1-2.

## Method F

## Step 1:

3-amino-6-bromo-N-phenylpyrazine-2-carboxamide

[0399] A mixture of 3-amino-6-bromo-pyrazine-2-carboxylic acid (3.5 g, 16.05 mmol), 1,1'-carbonyldiimidazole (5.205 g, 32.10 mmol), DIPEA (2.282 g, 3.075 mL, 17.66 mmol) and DMAP (98.04 mg, 0.8025 mmol) were combined in DMSO (131.2 mL) and stirred for 30 min. Aniline (1.495 g, 1.463 mL, 16.05 mmol) was then added and the resulting solution stirred at RT for 18 hours. After this time water was added and the product collected by filtration to give a brown powder (3.5 g, 74% Yield).

[0400]  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  7.04 (1H, m), 7.29 (2H, m), 7.72 (4H, m), 8.36 (1H, s), 10.22 (2H) ppm; MS ( $\text{ES}^+$ ) 295.

Step 2: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-phenylpyrazine-2-carboxamide

[0401] 3-amino-6-bromo-N-phenylpyrazine-2-carboxamide (50 mg, 0.1706 mmol) was heated in neat 1-ethylsulfonylpiperazine (excess) at  $190^\circ\text{C}$ . for 115 mins in the microwave. The mixture was partitioned between water and EtOAc and the aqueous layer separated. The organic phase was dried ( $\text{MgSO}_4$ ) and concentrated in vacuo. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{M}$ , 100  $\text{\AA}$  column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (41.2 mg, 58% Yield).

[0402]  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.24 (t,  $J=7.4$  Hz, 3H), 3.11 (t,  $J=7.3$  Hz, 2H), 3.34-3.31 (m, 4H), 3.58-3.56 (m, 4H), 6.94 (s, 2H), 7.13 (s, 1H), 7.40-7.36 (m, 2H), 7.77-7.75 (m, 2H), 8.29 (s, 1H) and 9.99 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 391


## Example 7

5-(1,1-dioxo-1,4-thiazinan-4-yl)-3-(6-methyl-1H-benzimidazol-2-yl)pyrazine-2-amine (Compound I-21)


[0403]

## SCHEME VII

Compound I-21



-continued



[0404] Compound I-21 was prepared using Method A, Steps 1-3 followed by Method G, Steps 1-2.

## Method G

Step 1: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-bromopyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate

[0405] A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-amine (800 mg, 2.630 mmol), di-tert-butyl dicarbonate (2.584 g, 2.720 mL, 11.84 mmol) and DMAP (32.13 mg, 0.2630 mmol) in a mixture of acetonitrile (15 mL) and THF (10 mL) was allowed to stir overnight at RT. The mixture was concentrated in vacuo and purified on silica gel by flash column chromatography (30% EtOAc/Hexanes) to afford the title compound as a light yellow solid (1.10 g, 69% Yield). MS (ES<sup>+</sup>) 606

Step 2: 5-(1,1-dioxo-1,4-thiazinan-4-yl)-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-amine

[0406] A solution of tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]-6-methyl-benzimidazole-1-carboxylate (70 mg, 0.1158 mmol) in DMF (2 mL) was treated with thiomorpholine (59.74 mg, 0.5790 mmol) and heated to 60° C. for 3 hours. The mixture was then diluted with EtOAc, washed with water, brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was dissolved in DCM (10 mL), treated with mCPBA (59.95 mg, 0.3474 mmol) and stirred at RT for 1 hour. After washing with a 1:1 Na<sub>2</sub>CO<sub>3</sub>/Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution, TFA (264.1 mg, 178.4 μL, 2.316 mmol) was added and the mixture stirred at RT for 1 hour. This was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μm, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (13.1 mg, 31% Yield).

[0407] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.45 (3H, s), 3.15-3.2 (4H, m), 4.15-4.2 (4H, m), 7.1-7.15 (1H, m), 7.4-7.45 (1H, m), 7.5-7.55 (1H, m), 8.15 (1H, s) ppm; MS (ES<sup>+</sup>) 359


## Example 8

3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)pyrazin-2-amine (Compound I-22)

[0408]

## SCHEME VIII

Compound I-22



[0409] Compound I-22 was prepared using Method A, Steps 1-3 followed by Method H, Step 1.

## Method H

Step 1: 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)pyrazin-2-amine

[0410] A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-amine (150 mg, 0.4932 mmol), 1,4-dioxaspiro[4.5]dec-7-en-8-ylboronic acid (181.5 mg, 0.9864 mmol), dichloropalladium; triethylphosphane (10.20 mg, 0.02466 mmol) and Na<sub>2</sub>CO<sub>3</sub> (740.0 μL of 2 M, 1.480 mmol) were heated in DMF (1.500 mL) at 130° C. in a microwave for 30 min. EtOAc was added and the organic phase washed with saturated aqueous Na<sub>2</sub>CO<sub>3</sub>, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The resulting oil was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μm, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound as a yellow solid (113 mg, 63% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.86 (t, J=6.4 Hz, 2H), 2.45 (m, 5H), 2.76 (s, 2H), 3.95 (s, 4H), 6.61 (t, J=3.9 Hz, 1H), 7.07 (dd, J=1.3, 8.3 Hz, 1H), 7.38 (s, 1H), 7.48 (d, J=8.2 Hz, 1H), 7.54


(s, 1H), 7.62 (d, J=8.3 Hz, 1H), 8.31 (d, J=4.0 Hz, 1H) and 8.37 (s, 1H) ppm; MS (ES<sup>+</sup>) 364

### Example 9


4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide (Compound I-23)

[0411]

#### SCHEME IX



-continued



[0412] Compound I-23 was prepared using Method A, Steps 1-3 followed by Method I, Steps 1-3.

Method I: Step 1: Tert-butyl 4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate

[0413] A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-amine (500 mg, 1.644 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (660.8 mg, 2.137 mmol), dichloropalladium; triethylphosphane (34.00 mg, 0.08220 mmol) and Na<sub>2</sub>CO<sub>3</sub> (2.466 mL of 2 M, 4.932 mmol) in DMF (5.000 mL) were heated to 130° C. for 60 min in a microwave. EtOAc was added and the organic phase washed with saturated aqueous Na<sub>2</sub>CO<sub>3</sub>, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The mixture was taken through crude to the next stage. MS (ES<sup>+</sup>) 407

Step 2: 3-(6-methyl-1H-benzo[d]imidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0414] Tert-butyl 4-[5-amino-6-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate (620 mg, 1.525 mmol) was stirred in DCM (20 mL) with TFA (4 mL) for 2 hours. The solvent was removed in vacuo and the residue taken up in EtOAc and washed with saturated aqueous Na<sub>2</sub>CO<sub>3</sub>. DCM was added and the resultant solid was collected by filtration and taken through to the next stage. MS (ES<sup>+</sup>) 307

Step 3: 4-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide (Compound I-23)

[0415] A solution of 3-(6-methyl-1H-benzimidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (58 mg, 0.1893 mmol) and triethylamine (38.31 mg, 52.77 μL, 0.3786 mmol) in DMSO (1.450 mL) was treated with N,N-dimethylsulfamoyl chloride (35.34 mg, 26.43 μL, 0.2461 mmol). The solution was stirred at RT for 16 hours and then partitioned between EtOAc and water. The aqueous layer was separated, organics dried over MgSO<sub>4</sub> and concentrated in vacuo to give a yellow solid which was washed with MeOH and filtered to give the desired product (47 mg, 57% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.46 (s, 3H), 2.77 (s, 2H),


2.80 (s, 6H), 3.48 (s, 2H), 3.96 (d, J=3.0 Hz, 2H), 6.76 (s, 1H), 7.10 (s, 1H), 7.61-7.40 (m, 3H) and 8.34 (s, 1H) ppm; MS (ES<sup>+</sup>) 414

### Example 10

1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyridin-4(1H)-one (Compound I-24)

[0416]

#### SCHEME X



[0417] Compound I-24 was prepared using Method A, Steps 1-3 followed by Method J, Step 1.

#### Method J

1-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyridin-4(1H)-one

[0418] A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-ylpyridin-4(1H)-one (80 mg, 0.263 mmol), pyridin-4(1H)-one (287 mg, 2.63 mmol), cesium carbonate (257 mg, 0.8 mmol) and copper (5 mg, 0.08 mmol) were heated in NMP (2 mL) at 150° C. for 1 hour. The reaction mixture was diluted with EtOAc, washed with 1N NaOH, and the organic layer filtered through a pad of celite and concentrated in vacuo. The resulting solid was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (24.7 mg, 30% Yield).

[0419] <sup>1</sup>H NMR (400.0 MHz, MeOD) δ 2.5 (3H, s), 6.65 (2H, d), 7.1-7.2 (1H, m), 7.4-7.5 (1H, m), 7.5-7.6 (1H, m), 8.45 (1H, s), 8.6 (2H, d) ppm; MS (ES<sup>+</sup>) 319

[0420] The following compounds were all prepared using a similar method to the one described for Compound I-24 in Example 10 above.

[0421] Compound I-310 1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyridin-2(1H)-one <sup>1</sup>H NMR (400 MHz, DMSO) δ 6.45 (t, 1H), 6.55 (d, 1H), 7.3-7.35 (m, 2H), 7.5-7.55 (m, 1H), 7.6-7.7 (br s, 2H), 8.05-8.1 (m, 1H), 8.55 (s, 1H) and 13.15 (br s, 1H) ppm; MS (ES<sup>+</sup>) 305.1

[0422] Compound I-311 1-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]pyridin-4-one


[0423] <sup>1</sup>H NMR (400 MHz, DMSO) δ 6.34 (d, 2H), 7.26-7.37 (m, 2H), 7.62 (d, 1H), 7.80 (d, 1H), 8.56 (d, 2H), 8.64 (s, 1H) and 13.17 (s, 1H) ppm; MS (ES<sup>+</sup>) 305.15

### Example 11

5-(5-amino-6-(1-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-1-methylpyridin-2(1H)-one (Compound I-25)

[0424]

#### SCHEME XI



[0425] Compound I-25 was prepared using Method A, Steps 1-3 followed by Method K, Step 1.

#### Method K

Step 1: 5-(5-amino-6-(6-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-1-methylpyridin-2(1H)-one

[0426] A mixture of 5-bromo-3-(6-methyl-1H-benzimidazol-2-yl)pyrazin-2-ylpyridin-4(1H)-one (100 mg, 0.3288 mmol), 1-methyl-


5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2(1H)-one (154.6 mg, 0.6578 mmol), potassium carbonate (136.4 mg, 0.9867 mmol) and [1,1-bis(diphenylphosphino)ferrocene]dichloropalladium (26.86 mg, 0.03289 mmol) in DMF (3 mL) were heated to 110° C. for 2 hours. The mixture was diluted with EtOAc, washed with water, and the organic layer concentrated in vacuo. The resulting residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (47.4 mg, 40% Yield). MS (ES<sup>+</sup>) 333

## Example 12




3-(1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine  
(Compound I-71)

[0427]

## SCHEME XII



Compound I-71



[0428] Compound I-71 was prepared using Method A, Steps 1-3 followed by Method G, Step 1 followed by Method L, Steps 1-3.

## Method L

Step 1: Tert-butyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0429] A mixture of tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]benzimidazole-1-carboxylate (5 g, 8.468 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (3.142 g, 10.16 mmol), 4-ditert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (59.96 mg, 0.08468 mmol) and K<sub>2</sub>CO<sub>3</sub> (2.341 g, 16.94 mmol) in toluene (45.00 mL) and water (5.000 mL) was heated to 100° C. overnight. The reaction mixture was allowed to cool to RT and diluted with DCM and water. The DCM layer was separated, washed with 1M NaOH solution, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to dryness to give a brown oil. This was purified on silica gel by flash column chromatography (0-30% EtOAc/hexanes) to afford the title compound (4.818 g, 82% Yield).

[0430] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.16 (18H, s), 1.38 (9H, s), 1.43 (9H, s), 2.63 (2H, m), 3.61 (2H, m), 4.02 (2H, m), 7.01 (1H, m), 7.49 (1H, m), 7.52 (1H, m), 7.70 (1H, m), 8.05 (1H, m), 9.02 (1H, m) ppm; MS (ES<sup>+</sup>) 693.38

## Method L

Step 2: 3-(1H-benzo[d]imidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0431] To a solution of tert-butyl 2-[3-(bis(tert-butoxycarbonyl)amino)-6-(1-tert-butoxycarbonyl)-3,6-dihydro-2H-py-

ridin-4-yl)pyrazin-2-yl]benzimidazole-1-carboxylate (5.808 g, 8.383 mmol) in DCM (58.08 mL) was added TFA (7.646 g, 5.166 mL, 67.06 mmol) and the mixture allowed to stir at RT for 30 minutes. Analysis showed there was SM remaining therefore a further 10 mL of TFA was added and reaction mixture left to stir for 24 hours. The reaction mixture was evaporated in vacuo to obtain the product as a yellow solid (4.362 g, 84% Yield).

**[0432]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.90 (s, 2H), 3.39 (s, 2H), 3.87 (s, 2H), 6.80 (s, 1H), 7.30 (s, 2H), 7.70 (s, 2H), 8.44 (s, 1H), 8.88 (s, 2H) and 12.90 (br s, 1H) ppm; MS (ES<sup>+</sup>) 293.0

#### Method L

##### Step 3: 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

**[0433]** Ethanesulfonyl chloride (16.30 mg, 12.01 μL, 0.1268 mmol) was added to a solution of 3-(1H-benzimidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (60 mg, 0.1153 mmol) and DIPEA (44.71 mg, 60.26 μL, 0.3459 mmol) in N,N-dimethylformamide (1 mL) and the resulting solution stirred at ambient temperature. Analysis after 30 minutes showed SM still remaining therefore the reaction was treated with further ethanesulfonyl chloride (4 μL, 0.04222 mmol). The mixture was allowed to stir for a further 10 minutes and then partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc (2×), DCM (2×), dried (MgSO<sub>4</sub>), filtered and concentrated. The mixture was triturated with DCM and the resultant yellow solid dried under vacuum at 50° C. (44.33 mg, 57% Yield).

**[0434]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.25 (t, J=7.3 Hz, 3H), 2.79 (br s, 2H), 3.16 (q, 2H), 3.51 (t, J=5.6 Hz, 2H), 4.01 (br d, 2H), 6.79 (s, 1H), 7.26 (t, 1H), 7.32 (t, 1H), 7.62 (d, J=7.8 Hz, 1H), 7.76 (d, J=7.9 Hz, 1H), 8.37 (s, 1H) and 12.85 (s, 1H) ppm; MS (ES<sup>+</sup>) 385.0

**[0435]** The following compounds were all prepared using same method or a method similar to the one described for Compound I-71 in Example 12 above.

**[0436]** Compound I-26 4-(5-amino-6-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide

**[0437]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 8.48-8.44 (m, 2H), 8.13 (d, J=7.9 Hz, 1H), 7.38 (dd, J=4.9, 7.6 Hz, 1H), 6.84 (s, 1H), 3.96 (d, J=2.8 Hz, 2H), 3.47 (s, 2H), 3.47 (s, 6H) ppm; MS (ES<sup>+</sup>) 401

**[0438]** Compound I-27 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-amine

**[0439]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 8.48-8.44 (m, 2H), 8.13 (d, J=7.9 Hz, 1H), 7.38 (dd, J=4.9, 7.6 Hz, 1H), 6.84 (s, 1H), 4.00 (d, J=2.8 Hz, 2H), 3.50 (t, J=5.7 Hz, 2H), 3.16 (q, J=7.4 Hz, 2H), 2.81 (d, J=1.5 Hz, 2H) and 1.25 (t, J=7.3 Hz, 3H) ppm; MS (ES<sup>+</sup>) 386

**[0440]** Compound I-58 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 386

**[0441]** Compound I-60 4-(5-amino-6-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide MS (ES<sup>+</sup>) 401

**[0442]** Compound I-66 3-(1H-benzo[d]imidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.90 (s, 2H), 3.39 (s, 2H), 3.87 (s, 2H),

6.80 (s, 1H), 7.30 (s, 2H), 7.70 (s, 2H), 8.44 (s, 1H), 8.88 (s, 2H) and 12.90 (br s, 1H) ppm; MS (ES<sup>+</sup>) 293

**[0443]** Compound I-67 1-(4-(5-amino-6-(3H-imidazo[4,5-b]pyridin-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-methylpropan-1-one <sup>1</sup>H NMR (400.0 MHz, MeOH) δ 8.51 (d, J=5.6 Hz, 1H), 8.42-8.38 (m, 1H), 8.34 (s, 1H), 7.66-7.63 (m, 1H), 6.61 (s, 1H), 4.35 (s, 1H), 4.24 (d, J=1.2 Hz, 1H), 3.83 (dd, J=5.8, 13.7 Hz, 2H), 3.05 (dt, J=23.4, 7.5 Hz, 1H), 2.74 (s, 1H), 2.68 (s, 1H) and 1.17 (dd, J=6.8, 12.1 Hz, 6H) ppm; MS (ES<sup>+</sup>) 364

**[0444]** Compound I-76 2-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)ethanenitrile <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.78-2.82 (2H, m), 2.95-3.0 (2H, m), 3.42-3.45 (2H, m), 4.05 (2H, s), 6.8-6.83 (1H, m), 7.3-7.33 (2H, m), 7.65-7.7 (2H, m), 8.19 (1H, s) ppm; MS (ES<sup>+</sup>) 332.1

**[0445]** Compound I-78 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl-(1-methylpiperidin-4-yl)methanone <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.65 (4H, m), 2.06 (2H, m), 2.23 (3H, s), 2.67-2.85 (4H, m), 3.73 (2H, m), 4.19, 4.29 (2H, 2×m), 6.75 (1H, d), 7.27 (2H, m), 7.61 (1H, d), 7.75 (1H, d), 8.24 (1H, s), 8.36 (1H, d), 12.85 (1H, br s) ppm; MS (ES<sup>+</sup>) 418.05

**[0446]** Compound I-79 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(thiophen-2-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 439.1

**[0447]** Compound I-82 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)(phenyl)methanone; MS (ES<sup>+</sup>) 396.2

**[0448]** Compound I-89 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide; MS (ES<sup>+</sup>) 399.2

**[0449]** Compound I-93 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(cyclopropylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 396.1

**[0450]** Compound I-94 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(1-methyl-1H-imidazol-4-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 436.1

**[0451]** Compound I-105 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(thiophen-3-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 438.1

**[0452]** Compound I-136 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)ethanone; MS (ES<sup>+</sup>) 334.2

**[0453]** Compound I-137 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-phenoxyethanone; MS (ES<sup>+</sup>) 426.2

**[0454]** Compound I-138 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-methoxyethanone; MS (ES<sup>+</sup>) 364.2

**[0455]** Compound I-139 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)(pyridin-2-yl)methanone; MS (ES<sup>+</sup>) 397.2

**[0456]** Compound I-140 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-(2-methoxyethoxy)ethanone; MS (ES<sup>+</sup>) 408.2

**[0457]** Compound I-161 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-(4-methylpiperazin-1-yl)methanone <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.19 (3H, s), 2.32 (4H, m), 2.73 (2H, m), 3.19 (4H, m), 3.42 (2H, m), 3.96 (2H, m), 6.74 (1H, s), 7.30 (2H, m), 7.61 (1H, d), 7.75 (1H, d), 8.36 (1H, s), 12.82 (1H, s) ppm; MS (ES<sup>+</sup>) 419.06

**[0458]** Compound I-173 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(4-(dimethylamino)piperidin-1-yl)methanone <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.35 (2H, m), 1.76 (2H, d), 2.19 (6H, s), 2.26 (1H, m), 2.74 (4H, m), 3.41 (2H, m), 3.64 (2H, d), 3.94 (2H, s), 6.74 (1H, s), 7.29 (2H, dt), 7.61 (1H, d), 7.75 (1H, d), 8.36 (1H, s), 12.82 (1H, s) ppm; MS (ES<sup>+</sup>) 447.09

For compounds I-186 to I-280 below, please see Tables 7 and 8 for analytical data.

**[0459]** Compound I-186 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-(dimethylamino)ethanone

**[0460]** Compound I-191 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(piperidin-3-yl)methanone

**[0461]** Compound I-200 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(cyclobutyl)methanone

**[0462]** Compound I-206 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(cyclopentyl)methanone

**[0463]** Compound I-210 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(phenylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

**[0464]** Compound I-213 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-phenylethanone

**[0465]** Compound I-214 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2,2-dimethylpropan-1-one

**[0466]** Compound I-215 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(2,6-difluorophenyl)methanone

**[0467]** Compound I-227 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(morpholino)methanone

**[0468]** Compound I-234 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-carboxamide

**[0469]** Compound I-235 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(tetrahydro-2H-pyran-4-yl)methanone

**[0470]** Compound I-241 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(pyridin-3-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

**[0471]** Compound I-243 2-methoxyethyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0472]** Compound I-244 methyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0473]** Compound I-249 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(piperidin-1-yl)methanone

**[0474]** Compound I-250 ethyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0475]** Compound I-254 but-2-ynyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0476]** Compound I-259 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(cyclohexyl)methanone

**[0477]** Compound I-261 benzyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0478]** Compound I-263 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(2,6-dimethylphenyl)methanone

**[0479]** Compound I-278 (R)-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)(3-(dimethylamino)pyrrolidin-1-yl)methanone

**[0480]** Compound I-279 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(1,4-diazepan-1-yl)methanone

**[0481]** Compound I-280 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(piperazin-1-yl)methanone

**[0482]** Compound I-284 [4-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]-cyclopropyl-methanone <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.75-0.78 (m, 4H), 2.00-2.05 (m, 1H), 2.67 (s, 1H), 2.80 (s, 1H), 3.75 (s, 1H), 3.94 (s, 1H), 4.21 (s, 1H), 4.47 (s, 1H), 6.78 (s, 1H), 7.24 (s, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.37 (d, 1H) and 12.85 (s, 1H) ppm; MS (ES<sup>+</sup>) 361.21

**[0483]** Compound I-291 3-(1H-benzimidazol-2-yl)-5-(1-propylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazin-2-amine <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.99-1.03 (m, 3H), 1.69-1.79 (m, 2H), 2.80 (br s, 2H), 3.10-3.20 (m, 2H), 3.49-3.50 (m, 2H), 3.95 (s, 2H), 6.79 (s, 1H), 7.24-7.34 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.37 (s, 1H) and 12.85 (s, 1H) ppm; MS (ES<sup>+</sup>) 399.14

**[0484]** Compound I-293 1-[4-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]-2-methylpropan-1-one <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.04-1.07 (m, 6H), 2.67 (br s, 1H), 2.73 (br s, 1H), 2.89-3.04 (m, 1H), 3.75 (s, 2H), 4.20 (s, 1H), 4.30 (s, 1H), 6.76 (br s, 1H), 7.24-7.33 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.36 (s, 1H) and 12.84 (s, 1H) ppm; MS (ES<sup>+</sup>) 363.22

**[0485]** Compound I-309 1-[4-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.01-1.06 (m, 3H), 2.33-2.47 (m, 2H), 2.67 (s, 1H), 2.76 (s, 1H), 3.67-3.75 (m, 2H), 4.19-4.23 (m, 2H), 6.76 (d, 1H), 7.24-7.33 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.36 (d, 1H) and 12.84 (s, 1H) ppm; MS (ES<sup>+</sup>) 349.19


## Example 13

3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide (Compound I-55)

[0486]

## SCHEME XIII

Compound I-55



[0487] Compound I-55 was prepared using Method A, Step 1 followed by Method M, Steps 1-5.

## Method M

Step 1: Tert-butyl 4-(3-methyl-4-oxo-3,4-dihydropteridin-6-yl)piperazine-1-carboxylate

[0488] To a solution of 6-bromo-3-methyl-pteridin-4-one (17.75 g, 73.64 mmol) in 2-methoxyethanol (163.5 mL) was

added tert-butyl piperazine-1-carboxylate (24.70 g, 132.6 mmol) and the resulting suspension heated for 2 hours at 100° C. The reaction mixture was allowed to cool in an ice-water bath and the precipitate that formed collected and washed with cold methanol to give the desired product as a yellow solid (25.08 g, 98% Yield).

[0489] <sup>1</sup>H NMR (400.0 MHz, CDCl<sub>3</sub>) δ 1.50 (9H, m), 3.49 (3H, s), 3.62 (4H, br m), 3.83 (4H, br m), 8.09 (1H, s), 8.58 (1H, s) ppm; MS (ES<sup>+</sup>) 347.08

## Method M

## Step 2: 3-amino-6-(4-(tert-butoxycarbonyl)piperazin-1-yl)pyrazine-2-carboxylic acid

[0490] Tert-butyl 4-(3-methyl-4-oxo-pteridin-6-yl)piperazine-1-carboxylate (25 g, 72.17 mmol) was stirred in NaOH (250 mL of 10% w/w.) and MeOH (250 mL) overnight at RT. The resulting suspension was concentrated and then acidified with formic acid followed by extraction of the aqueous with DCM. The mixture was treated with water, taken to pH 10 and filtered. The aqueous layer was reacidified and the resulting precipitated product was extracted into DCM and concentrated to give the product as a yellow solid (8 g, 24.74 mmol, 34% Yield). MS (ES<sup>+</sup>) 324.02

## Method M

## Step 3: Tert-butyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate

[0491] To a mixture of 3-amino-6-(4-tert-butoxycarbonylpiperazin-1-yl)pyrazine-2-carboxylic acid (8.08 g, 24.99 mmol), di(imidazol-1-yl)methanone (4.863 g, 29.99 mmol), DMAP (152.7 mg, 1.250 mmol) and N-ethyl-N-isopropylpropan-2-amine (8.074 g, 10.88 mL, 62.47 mmol) in DMSO (80.80 mL) was added pyridin-2-amine (4.704 g, 49.98 mmol) and the resulting solution stirred at 110° C. for 20 hours. Water (500 mL) was added and the mixture extracted into EtOAc (3×300 mL), dried and concentrated to give a residue which was triturated with EtOAc and filtered to give the required product as a yellow solid (5.48 g, 55% Yield).

[0492] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 10.08 (s, 1H), 8.38 (dd, J=0.9, 5.0 Hz, 1H), 8.36 (s, 1H), 8.21 (d, J=8.3 Hz, 1H), 7.88 (s, 1H), 7.20-7.17 (m, 1H), 7.01 (s, 2H), 3.50 (s, 4H), 3.41 (d, J=5.2 Hz, 4H) and 1.43 (s, 9H) ppm; MS (ES<sup>+</sup>) 400.08

## Method M

## Step 4: 3-amino-6-(piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0493] Tert-butyl 4-[5-amino-6-(2-pyridylcarbamoyl)pyrazin-2-yl]piperazine-1-carboxylate (5.43 g, 13.59 mmol) was suspended in DCM (50 mL) and stirred at 0° C. during addition of TFA (10 mL). The reaction mixture was warmed to RT over 2 hours but analysis showed little deprotection. The reaction was treated with a further 30 mL of TFA and stirred at RT for 15 minutes. The mixture was concentrated under reduced pressure to give a dark red oil which was dissolved in MeOH/DCM and azeotroped with toluene to give an intense orange solid which was filtered, washed with ether and dried under high vacuum (5.41 g, 73% Yield).

[0494] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 3.29 (4H, m), 3.65 (4H, m), 7.21 (1H, m), 7.89 (1H, m), 8.21 (1H, d), 8.38 (2H, m), 8.88 (2H, br s), 10.10 (1H, s) ppm; MS (ES<sup>+</sup>) 300.0

## Method M

## Step 5: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0495] 3-amino-6-piperazin-1-yl-N-(2-pyridyl)pyrazine-2-carboxamide (100 mg, 0.1896 mmol) was dissolved in DCM (5 mL) and treated with Et<sub>3</sub>N (57.56 mg, 79.28 μL, 0.5688 mmol) followed by ethanesulfonyl chloride (26.82

mg, 19.76 μL, 0.2086 mmol). The reaction mixture was allowed to stir for 2 hours and then diluted with dichloromethane and washed with water. The aqueous layer was extracted further with dichloromethane and the organics washed with saturated aqueous NaHCO<sub>3</sub>, dried and concentrated in vacuo. The residue was purified on silica gel by flash column chromatography (0-5% MeOH/DCM) to afford the title compound (74.2 mg, 37% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.24 (t, J=7.3 Hz, 3H), 3.13 (q, J=7.4 Hz, 2H), 3.34-3.38 (m, 4H), 3.52-3.54 (m, 4H), 7.04 (s, 2H), 7.18-7.21 (m, 1H), 7.88 (s, 1H), 8.21 (d, J=8.4 Hz, 1H), 8.38 (d, J=3.2 Hz, 2H) and 10.08 (s, 1H) ppm; MS (ES<sup>+</sup>) 392.0

[0496] The following compounds were all prepared using the same method or a method similar to the one described for Compound I-55 in Example 13 above.

[0497] Compound I-42 3-amino-6-(piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0498] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 3.29 (4H, m), 3.65 (4H, m), 7.21 (1H, m), 7.89 (1H, m), 8.21 (1H, d), 8.38 (2H, m), 8.88 (2H, br s), 10.10 (1H, s); MS (ES<sup>+</sup>) 300

[0499] Compound I-68 3-amino-6-(4-(2-methylpropanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.03 (d, J=6.7 Hz, 6H), 2.96 (m, 1H), 3.40-3.48 (m, 4H), 3.62-3.71 (m, 4H), 7.02 (s, 2H), 7.18-7.21 (m, 1H), 7.86-7.90 (m, 1H), 8.22 (d, 1H), 8.38 (s, 2H) and 10.09 (s, 1H) ppm; MS (ES<sup>+</sup>) 370.1

[0500] Compound I-81 3-amino-6-(4-(N,N-dimethylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 406.2

[0501] Compound I-109 3-amino-6-(4-(phenylcarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 403.2

[0502] Compound I-113 ethyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate; MS (ES<sup>+</sup>) 371.2

[0503] Compound I-114 3-amino-6-(4-(morpholine-4-carbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 412.2

[0504] Compound I-116 3-amino-N-(pyridin-2-yl)-6-(4-(thiophen-2-ylsulfonyl)piperazin-1-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 445.1

[0505] Compound I-117 3-amino-N-(pyridin-2-yl)-6-(4-(thiophen-3-ylsulfonyl)piperazin-1-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 445.1

[0506] Compound I-118 3-amino-6-(4-(cyclopropylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 403.1

[0507] Compound I-119 3-amino-6-(4-(furan-3-ylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide MS (ES<sup>+</sup>) 429.1

[0508] Compound I-141 3-amino-6-(4-(2-phenylethanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 417.2

[0509] Compound I-142 3-amino-6-(4-ethanoylpiperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 341.2

[0510] Compound I-143 3-amino-6-(4-(2-phenoxyethanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 433.2

[0511] Compound I-144 3-amino-6-(4-(cyclopentanecarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 395.2

[0512] Compound I-145 3-amino-6-(4-(cyclobutanecarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 381.2

[0513] Compound I-146 3-amino-N-(pyridin-2-yl)-6-(4-(pyridine-3-carbonyl)piperazin-1-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 404.2

[0514] Compound I-147 3-amino-6-(4-(2-methoxyethanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 371.2

[0515] Compound I-148 3-amino-N-(pyridin-2-yl)-6-(4-(pyridine-2-carbonyl)piperazin-1-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 404.2

[0516] Compound I-149 3-amino-6-(4-(2,6-difluorophenylcarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 439.2

[0517] Compound I-150 3-amino-6-(4-(2-(dimethylamino)ethanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 384.2

[0518] Compound I-151 3-amino-6-(4-(2-(2-methoxyethoxy)ethanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide; MS (ES<sup>+</sup>) 415.2

For compounds I-207 to I-262 below, please see Tables 7 and 8 for analytical data.

[0519] Compound I-207 3-amino-6-(4-(phenylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0520] Compound I-209 3-amino-6-(4-(2,2-dimethylpropanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0521] Compound I-212 3-amino-6-(4-(2-ethylbutanoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0522] Compound I-219 3-amino-6-(4-(methylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0523] Compound I-229 3-amino-6-(4-(dimethylcarbamoyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0524] Compound I-230 3-amino-N-(pyridin-2-yl)-6-(4-(tetrahydro-2H-pyran-4-carbonyl)piperazin-1-yl)pyrazine-2-carboxamide

[0525] Compound I-233 3-amino-N-(pyridin-2-yl)-6-(4-(pyridin-3-ylsulfonyl)piperazin-1-yl)pyrazine-2-carboxamide

[0526] Compound I-237 2-methoxyethyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate

[0527] Compound I-239 3-amino-N-(pyridin-2-yl)-6-(4-(pyrrolidine-1-carbonyl)piperazin-1-yl)pyrazine-2-carboxamide

[0528] Compound I-240 3-amino-6-(4-(propylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0529] Compound I-242 methyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate

[0530] Compound I-247 3-amino-6-(4-(piperidine-1-carbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0531] Compound I-248 3-amino-6-(4-(cyclopentylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0532] Compound I-252 but-2-ynyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate

[0533] Compound I-253 3-amino-6-(4-(benzylsulfonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0534] Compound I-256 3-amino-6-(4-(cyclohexanecarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide

[0535] Compound I-260 benzyl 4-(5-amino-6-(pyridin-2-ylcarbamoyl)pyrazin-2-yl)piperazine-1-carboxylate


[0536] Compound I-262 3-amino-6-(4-(2,6-dimethylphenylcarbonyl)piperazin-1-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide


#### Example 14

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-18)

[0537]

SCHEME XIV





**[0538]** Compound I-18 was also prepared using Method A, Steps 1-2 followed by Method N, Steps 1-6.

#### Method N

Step 1: 3-amino-6-bromo-N'-(phenylcarbonyl)pyrazine-2-carbohydrazide

**[0539]** TBTU (22.09 g, 68.80 mmol) and Et<sub>3</sub>N (4.642 g, 6.394 mL, 45.87 mmol) were added to a solution of 3-amino-6-bromo-pyrazine-2-carboxylic acid (10 g, 45.87 mmol) and benzohydrazide (7.494 g, 55.04 mmol) in DMF (100.0 mL) and the resulting solution stirred at ambient temperature overnight. The reaction mixture was poured into water and the resultant precipitate was filtered and dried under vacuum. This was triturated with hot EtOAc and the resulting precipitate filtered and dried to give the desired product (13.11 g, 85% Yield).

**[0540]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 7.53 (t, 2H), 7.60 (d, 1H), 7.69 (br s, 2H), 7.91 (d, 2H), 8.44 (s, 1H), 10.48 (s, 1H) and 10.55 (s, 1H) ppm; MS (ES<sup>+</sup>) 338.92

#### Method N

Step 2: 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0541]** Polyphosphonic acid (25 mL of 84% w/v, 256.1 mmol) was heated to 100° C. and treated with 3-amino-N'-benzoyl-6-bromo-pyrazine-2-carbohydrazide (6.82 g, 20.29 mmol) in portions. Analysis after 1 hour showed SM remaining therefore further polyphosphonic acid was added (25 mL of 84% w/v, 256.1 mmol). The reaction was heated for a total of 3 hours and then allowed to cool. Water was added and the mixture allowed to stir to obtain a precipitate which was filtered and dried under vacuum (5.02 g, 78% Yield). MS (ES<sup>+</sup>) 317.86

#### Method N

Step 3: Tert-butyl 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate

**[0542]** 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (5.02 g, 15.78 mmol) was suspended in DCM (75 mL) and DMAP (192.8 mg, 1.578 mmol) was added. BOC anhydride (10.33 g, 47.34 mmol) was added in portions and the reaction allowed to stir at ambient

temperature. Further BOC anhydride (3.444 g, 3.625 mL, 15.78 mmol) was added followed by triethylamine (2.395 g, 3.299 mL, 23.67 mmol) and the mixture heated to 60° C. The reaction was allowed to cool to ambient temperature and the solvent removed in vacuo. The mixture was redissolved in DCM, washed with 1M HCl, NaHCO<sub>3</sub>, brine, dried over MgSO<sub>4</sub>, filtered and concentrated to dryness. The resultant residue was purified on silica gel by flash column chromatography (0-40% EtOAc/Petrol) to afford the title compound as a yellow solid (5.12 g, 78% Yield); <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.29 (s, 9H), 7.57-7.75 (m, 4H), 8.12-8.14 (m, 2H) and 9.16 (s, 1H) ppm; MS (ES<sup>+</sup>) 420.01

#### Method N

Step 4: Tert-butyl 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-ylcarbamate

**[0543]** Tert-butyl 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (2.56 g, 6.121 mmol) was dissolved in DMF (20 mL) and piperazine (5.272 g, 61.21 mmol) was added. The mixture was heated to 100° C. for 30 minutes, allowed to cool to ambient temperature and the solvent removed in vacuo. The mixture was partitioned between DCM/Water and the organic layer washed with brine, dried over MgSO<sub>4</sub>, filtered and concentrated to give a yellow oil. The resultant residue was purified on silica gel by flash column chromatography (0-10% MeOH/DCM) to afford the title compound (2.29 g, 89% Yield).

**[0544]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.30 (s, 9H), 2.60 (t, J=5.0 Hz, 2H), 2.83 (d, J=5.0 Hz, 2H), 3.21 (t, J=4.7 Hz, 2H), 3.55 (d, J=5.1 Hz, 2H), 7.63-7.69 (m, 3H), 8.06-8.07 (m, 2H), 8.35 (s, 1H) and 9.55 (s, 1H) ppm; MS (ES<sup>+</sup>) 424.17

#### Method N

Step 5: 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-amine

**[0545]** Tert-butyl 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-ylcarbamate (2.29 g, 5.408 mmol) was dissolved in DCM (20 mL) and treated with TFA (5 mL, 64.90 mmol) was added. The reaction was allowed to stir for 1 hour at ambient temperature and then the solvent removed in vacuo. The mixture was treated with NaHCO<sub>3</sub>/brine and EtOAc and the 2 layers separated. The aqueous layer was extracted with EtOAc, followed by DCM and CHCl<sub>3</sub>.

The mixture was dried over  $\text{MgSO}_4$ , filtered and concentrated to give the product as an orange solid (1.44 g, 82% Yield).  $^1\text{H}$ NMR (400.0 MHz, DMSO)  $\delta$  3.00 (4H, br m), 3.43 (4H, br m), 6.05 (2H, br s), 7.20 (1H, m), 7.34 (2H, m), 7.90 (1H, m), 8.13 (2H, m) ppm; ( $\text{ES}^+$ ) 324.03

#### Method N

Step 6: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0546]** Ethanesulfonyl chloride (37.51 mg, 27.64  $\mu\text{L}$ , 0.2917 mmol) was added to a solution of 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-piperazin-1-yl-pyrazin-2-amine (116 mg, 0.2652 mmol) and triethylamine (53.67 mg, 73.93  $\mu\text{L}$ , 0.5304 mmol) in dichloromethane (5 mL) at  $0^\circ\text{C}$ . and the resulting solution stirred at room temperature for 1.5 hours. The mixture was diluted with dichloromethane, washed with water and the aqueous layer extracted further with dichloromethane. The organics were washed with saturated aqueous  $\text{NaHCO}_3$ , dried over  $\text{MgSO}_4$  and concentrated in vacuo. The resultant residue was purified on silica gel by flash column chromatography (3% MeOH/DCM) to afford the title compound still impure. This was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{M}$ , 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (25.2 mg, 18% Yield).

**[0547]**  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.24 (3H, t), 3.11 (2H, q), 3.33-3.36 (4H, m), 3.56 (4H, m), 6.97 (2H, br s), 7.63-7.68 (3H, m), 8.11-8.13 (2H, m), 8.28 (1H, s) ppm; MS ( $\text{ES}^+$ ) 416.01

**[0548]** The following compounds were all prepared using the same method or a method similar to the one described for Compound I-18 in Example 14 above.

**[0549]** Compound I-37 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-amine  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  3.27 (s, 4H), 3.67-3.69 (m, 4H), 7.03 (s, 1H), 7.64-7.70 (m, 3H), 8.10-8.13 (m, 2H), 8.31 (s, 1H) and 8.87 (br s, 2H) ppm; MS ( $\text{ES}^+$ ) 323.99

**[0550]** Compound I-72 3-(5-cyclohexyl-1,3,4-oxadiazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine;  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.24 (3H, t), 1.38-1.47 (3H, m), 1.60 (2H, m), 1.66 (2H, m), 1.79 (2H, m), 2.06 (2H, m), 3.10 (2H, q), 6.89 (2H, br s), 8.22 (1H, s) ppm; MS ( $\text{ES}^+$ ) 422.03

**[0551]** Compound I-83 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-N,N-dimethylpiperazine-1-sulfonamide; MS ( $\text{ES}^+$ ) 430.2

**[0552]** Compound I-87 5-(4-(1-methyl-1H-imidazol-4-ylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 467.2

**[0553]** Compound I-90 5-(4-(furan-3-ylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 453.1

**[0554]** Compound I-92 5-(4-(cyclopentylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 455.2

**[0555]** Compound I-96 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(4-(thiophen-2-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 469.1

**[0556]** Compound I-98 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(4-(thiophen-3-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 469.1

**[0557]** Compound I-99 5-(4-(benzylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS ( $\text{ES}^+$ ) 477.2

**[0558]** Compound I-101 6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(phenyl)methanone; MS ( $\text{ES}^+$ ) 427.2

For compounds I-178 to I-258 below, please see Tables 7 and 8 for analytical data.

**[0559]** Compound I-178 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-(2-methoxyethoxy)ethanone

**[0560]** Compound I-179 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-methoxyethanone

**[0561]** Compound I-180 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)ethanone

**[0562]** Compound I-183 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(pyridin-3-yl)methanone

**[0563]** Compound I-187 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(pyridin-2-yl)methanone

**[0564]** Compound I-189 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(cyclobutyl)methanone

**[0565]** Compound I-190 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-methylpropan-1-one

**[0566]** Compound I-194 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(cyclopentyl)methanone

**[0567]** Compound I-195 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-phenoxyethanone

**[0568]** Compound I-196 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(4-(phenylsulfonyl)piperazin-1-yl)pyrazin-2-amine

**[0569]** Compound I-197 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-phenylethanone

**[0570]** Compound I-198 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2,2-dimethylpropan-1-one

**[0571]** Compound I-202 5-(4-(2,6-difluorophenylsulfonyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0572]** Compound I-203 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(2,6-difluorophenyl)methanone

**[0573]** Compound I-216 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(morpholino)methanone

**[0574]** Compound I-218 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-N,N-dimethylpiperazine-1-carboxamide

**[0575]** Compound I-221 2-methoxyethyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

**[0576]** Compound I-223 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(pyrrolidin-1-yl)methanone

**[0577]** Compound I-225 methyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0578] Compound I-231 (4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(piperidin-1-yl) methanone

[0579] Compound I-232 ethyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0580] Compound I-246 but-2-ynyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0581] Compound I-251 (4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(cyclohexyl) methanone

[0582] Compound I-255 benzyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0583] Compound I-258 (4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(2,6-dimethylphenyl)methanone

### Example 15

5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine  
(Compound I-28)

[0584]

[0585] Compound I-28 was prepared using Method A, Steps 1-2 followed by Method N, Steps 1-2 followed by Method O, Steps 1-4.


### Method O

Step 1: Tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

[0586] 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (3.8348 g, 12.05 mmol) was suspended in DCM (57.29 mL) and THF (57.29 mL) and DMAP (147.2 mg, 1.205 mmol) was added followed by BOC anhydride (7.890 g, 8.305 mL, 36.15 mmol) in portions. The reaction was allowed to stir at RT. After 1 hour the reaction was treated with triethylamine (2.52 mL, 18.08 mmol) and the mixture allowed to stir. Analysis showed SM still remaining therefore a further 0.1 eq of DMAP was added followed by 5 g of BOC anhydride (1.9 eq) and the reaction left to stir at RT for 2 hours. The solvent was removed in vacuo and the mixture redissolved in DCM, washed with NaHCO<sub>3</sub> and brine, dried over MgSO<sub>4</sub>, filtered and concentrated to give a brown oil which was purified on silica gel by flash column chromatog-

SCHEME XV

Compound I-28



raphy (0-25% EtOAc/Petrol) to afford the title compound as a pale yellow solid (3.96 g, 63% Yield). MS (ES<sup>+</sup>) 519.95

#### Method O

Step 2: tert-butyl 4-(5-(bis(tert-butoxycarbonyl)amino)-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

**[0587]** Tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (2.4633 g, 4.752 mmol) was dissolved in DMF (19.24 mL) and tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (2.204 g, 7.128 mmol) and Pd(PPh<sub>3</sub>)<sub>2</sub>Cl<sub>2</sub> (196.6 mg, 0.4752 mmol) were added followed by Na<sub>2</sub>CO<sub>3</sub> (7.130 mL of 2 M, 14.26 mmol) and the reaction heated at 100° C. under nitrogen for 15 minutes. The reaction mixture was allowed to cool to RT and the solvent removed in vacuo. The mixture was diluted with EtOAc/water and the organic layer washed with water (×4) followed by brine (×3). After drying over MgSO<sub>4</sub>, the reaction was filtered and evaporated to dryness to give a dark brown oil which was purified on silica gel by flash column chromatography (0-40% EtOAc/Petrol) to afford the title compound (2.40 g, 81% Yield).

**[0588]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.02 (27H, m), 2.67 (2H, m), 3.60 (2H, m), 4.13 (2H, m), 6.88 (1H, m), 7.70 (2H, m), 8.15 (2H, m), 8.87 (1H, m) ppm

#### Method O

Step 3: 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

**[0589]** Tert-butyl 4-(5-(bis(tert-butoxycarbonyl)amino)-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate (2.4 g, 3.867 mmol) was dissolved in DCM (40 mL) and treated with TFA (8 mL, 103.8 mmol). The reaction was allowed to stir overnight and then the solvent was removed in vacuo. The mixture was azeotroped with DCM (×3) and dried under vacuum to give the desired product as an orange solid (1.89 g, 100% Yield).

**[0590]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.83 (br d, 2H), 3.39 (br d, 2H), 3.91 (br d, 2H), 6.66 (s, 1), 7.66-7.73 (m, 3H), 8.11-8.14 (m, 2H), 8.63 (s, 1H) and 8.86 (s, 1H) ppm; MS (ES<sup>+</sup>) 321.0

#### Method O

Step 4: 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0591]** 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (50 mg, 0.1151 mmol) was dissolved in DCM (3 mL) and Et<sub>3</sub>N (34.94 mg, 48.13 μL, 0.3453 mmol) followed by ethanesulfonyl chloride (16.28 mg, 12.00 μL, 0.1266 mmol) were added. The reaction was allowed to stir for 30 mins and then diluted with dichloromethane and washed with water. The aqueous layer was extracted further with dichloromethane and the organics washed with saturated aqueous NaHCO<sub>3</sub>, dried over MgSO<sub>4</sub> and concentrated in vacuo. The resultant residue was purified on silica gel by flash column chromatography (0-5% MeOH/DCM) to afford the title compound (47.47 mg, 60% Yield).

**[0592]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.24 (t, J=7.4 Hz, 3H), 2.74 (br d, 2H), 3.15 (q, 2H), 3.50 (t, 2H), 4.00 (br d, 2H), 6.68 (br s, 1H), 7.67-7.69 (m, 5H), 8.14 (d, 2H) and 8.58 (s, 1H) ppm; MS (ES<sup>+</sup>) 413.0

**[0593]** The following compounds were all prepared using the same method or a similar method to the one described for Compound I-28 in Example 15 above.

**[0594]** Compound I-41 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.83 (br d, 2H), 3.39 (br d, 2H), 3.91 (br d, 2H), 6.66 (s, 1H), 7.66-7.73 (m, 3H), 8.11-8.14 (m, 2H), 8.63 (s, 1H) and 8.86 (s, 1H) ppm; MS (ES<sup>+</sup>) 321

**[0595]** Compound I-54 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 413

**[0596]** Compound I-84 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-sulfonamide; MS (ES<sup>+</sup>) 427.1

**[0597]** Compound I-100 5-(1-(benzylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 474.2

**[0598]** Compound I-102 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1-(phenylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 460.1

**[0599]** Compound I-103 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-yl (phenyl)methanone; MS (ES<sup>+</sup>) 424.2

**[0600]** Compound I-115 but-2-ynyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate; MS (ES<sup>+</sup>) 416.2

**[0601]** Compound I-120 5-(1-(1-methyl-1H-imidazol-4-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 464.1

**[0602]** Compound I-121 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1-(thiophen-3-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 466.1

**[0603]** Compound I-122 5-(1-(cyclopropylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 424.1

**[0604]** Compound I-123 5-(1-(furan-3-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 450.1

**[0605]** Compound I-124 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1-(propylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 426.2

**[0606]** Compound I-125 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)ethanone; MS (ES<sup>+</sup>) 362.2

**[0607]** Compound I-126 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-phenoxyethanone; MS (ES<sup>+</sup>) 454.2

**[0608]** Compound I-127 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-yl (cyclopentyl)methanone; MS (ES<sup>+</sup>) 416.2

**[0609]** Compound I-128 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-yl (cyclobutyl)methanone; MS (ES<sup>+</sup>) 402.2

**[0610]** Compound I-129 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-yl (pyridin-3-yl)methanone; MS (ES<sup>+</sup>) 425.2

**[0611]** Compound I-130 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-methoxyethanone; MS (ES<sup>+</sup>) 392.2

[0612] Compound I-131 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-ethylbutan-1-one; MS (ES<sup>+</sup>) 418.2

[0613] Compound I-132 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl (pyridin-2-yl)methanone; MS (ES<sup>+</sup>) 425.2

[0614] Compound I-133 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl (2,6-difluorophenyl)methanone; MS (ES<sup>+</sup>) 460.2

[0615] Compound I-134 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-(dimethylamino)ethanone; MS (ES<sup>+</sup>) 405.2

[0616] Compound I-135 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-(2-methoxyethoxy)ethanone; MS (ES<sup>+</sup>) 436.2

[0617] Compound I-163 Tert-butyl 4-(5-amino-6-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate; <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.44 (9H, s), 2.60 (2H, m), 3.57 (2H, m), 4.06 (2H, m), 6.70 (1H, s), 7.30 (2H, br s), 7.69 (2H, m), 7.77 (1H, m), 8.25 (2H, m), 8.51 (1H, s) ppm; MS (ES<sup>+</sup>) 421.02

For compounds I-192 to I-257 below, please see Tables 7 and 8 for analytical data.

[0618] Compound I-192 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-methylpropan-1-one

[0619] Compound I-201 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-phenylethanone

[0620] Compound I-217 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl (morpholino)methanone

[0621] Compound I-220 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-N,N-dimethyl-5,6-dihydropyridine-1(2H)-carboxamide

[0622] Compound I-222 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1-(pyridin-3-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0623] Compound I-224 2-methoxyethyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0624] Compound I-226 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl (pyrrolidin-1-yl)methanone

[0625] Compound I-228 methyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0626] Compound I-236 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl (piperidin-1-yl)methanone

[0627] Compound I-238 ethyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0628] Compound I-245 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(1-(thiophen-2-ylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0629] Compound I-257 benzyl 4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0630] Compound I-285 [4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]-cyclopropyl-methanone <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.74-0.78 (m, 4H), 1.99 (br s, 0.5H), 2.09 (br s, 0.5H), 2.61 (br s, 1H), 2.68 (br s, 1H), 3.73 (br s, 1H), 3.93 (br s, 1H), 4.19 (br

s, 1H), 4.47 (br s, 1H), 6.67 (br s, 1H), 7.63-7.70 (m, 5H), 8.13 (d, 2H) and 8.59 (s, 1H) ppm; MS (ES<sup>+</sup>) 389.16

[0631] Compound I-301 1-[4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.04 (t, 3H), 2.36-2.45 (m, 2H), 2.60 (br s, 1H), 2.70 (br s, 1H), 3.65-3.72 (m, 2H), 4.17 (s, 1H), 4.21 (s, 1H), 6.65 (s, 1H), 7.62-7.69 (m, 5H), 8.11-8.13 (d, 2H) and 8.56 (s, 1H) ppm; MS (ES<sup>+</sup>) 377.17


### Example 16

3-(1H-benzo[d]imidazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine (Compound I-16)


[0632]

### SCHEME XVI

Compound I-16



-continued



**[0633]** Compound I-16 was prepared using Method A, Steps 1-3 followed by Method P, Steps 1-3.

## Method P

Step 1: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-bromopyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate

**[0634]** 3-(1H-benzimidazol-2-yl)-5-bromo-pyrazin-2-amine (5 g, 17.23 mmol) was dissolved in acetonitrile (70 mL) and THF (120 mL) and treated with a few crystals of DMAP followed by addition of BOC anhydride (16.92 g, 17.81 mL, 77.54 mmol) in 4 portions over 15 minutes. The reaction was allowed to stir at RT overnight and concentrated to an oil, then redissolved and concentrated from DCM twice. The resultant residue was purified on silica gel by flash column chromatography (5-30% EtOAc/Petroleum ether) to afford the title compound (9.94 g, 98% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.10 (27H, m), 7.45 (1H, m), 7.52 (1H, m), 7.75 (1H, m), 8.09 (1H, m) and 9.10 (1H, s) ppm; MS (ES<sup>+</sup>) 591.90

## Method P

Step 2: 3-(1H-benzo[d]imidazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-amine

**[0635]** Tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]benzimidazole-1-carboxylate (12 g, 20.32 mmol) was dissolved in NMP (30 mL) followed by addition of DIPEA (5.252 g, 7.078 mL, 40.64 mmol) and piperazine (12.25 g, 142.2 mmol). The reaction was heated to 100° C. for 4 hours and diluted with EtOAc, washed with water, followed by brine and concentrated. The resultant residue was purified on silica gel by flash column chromatography (0-50% MeOH/EtOAc followed by 3% NH<sub>4</sub>OH in MeOH) to afford the BOC-protected product. This was dissolved in DCM/methanol and HCl (4M in dioxane) (50.80 mL of 4 M, 203.2 mmol) was added and the mixture stirred overnight at RT. The reaction was concentrated to a solid and redissolved in EtOAc, washed with sat. aqueous NaHCO<sub>3</sub> and brine. The organic layer was dried over MgSO<sub>4</sub> and concentrated to a give a solid which was slurried in DCM and titrated by slow addition of ether to give the desired product as an orange solid (7.07 g, 85% Yield). <sup>1</sup>H NMR (400.0 MHz,

DMSO) δ 2.95 (4H, m), 3.52 (4H, m), 7.10 (2H, m), 7.55 (1H, m), 7.65 (1H, m), 8.15 (1H, m) and 12.61 (1H, br s) ppm; MS (ES<sup>+</sup>) 296.02

## Method P

Step 3: 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine

**[0636]** 3-(1H-benzo[d]imidazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-amine (50 mg, 0.169 mmol) was dissolved in DCM (3 mL) and Et<sub>3</sub>N (70.0 μL, 0.507 mmol) followed by ethanesulfonyl chloride (17.6 μL, 0.186 mmol) were added. The reaction was allowed to stir for 30 mins and then diluted with dichloromethane and washed with water. The aqueous layer was extracted further with dichloromethane and the organics washed with saturated aqueous NaHCO<sub>3</sub>, dried over MgSO<sub>4</sub> and concentrated in vacuo. The resultant residue was purified on silica gel by flash column chromatography (0-5% MeOH/DCM) to afford the title compound (42.6 mg, 65% Yield). MS (ES<sup>+</sup>) 388.0

**[0637]** The following compounds were all prepared using the same method or a similar method to the one described for Compound I-16 in Example 16 above.

**[0638]** Compound I-39 (S)-5-(3-aminopyrrolidin-1-yl)-3-(1H-benzo[d]imidazol-2-yl)pyrazin-2-amine

**[0639]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.7-1.75 (1H, m), 2.05-2.1 (1H, m), 3.35-3.4 (1H, m), 3.4-3.5 (1H, m), 3.55-3.65 (3H, m), 7.02 (2H, s), 7.3-7.4 (2H, m), 7.55 (1H, d), 7.65-7.7 (2H, m), 12.5 (1H, vbrs) ppm; MS (ES<sup>+</sup>) 296

**[0640]** Compound I-40 3-(1H-benzo[d]imidazol-2-yl)-5-(piperazin-1-yl)pyrazin-2-amine

**[0641]** <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.9-2.95 (4H, m), 3.4-3.5 (4H, m), 7.3-7.4 (4H, m), 7.6 (1H, d), 7.7 (1H, d), 7.95 (1H, s), 12.7 (1H, vbrs) ppm; MS (ES<sup>+</sup>) 296

**[0642]** Compound I-80 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(thiophen-3-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 441.1

**[0643]** Compound I-85 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)propane-2-N, N-dimethylsulfamide; MS (ES<sup>+</sup>) 402.2

**[0644]** Compound I-86 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(methylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 373.1

**[0645]** Compound I-88 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-N,N-dimethylpiperazine-1-sulfonamide; MS (ES<sup>+</sup>) 402.2

**[0646]** Compound I-91 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(1-methyl-1H-imidazol-4-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 439.2

**[0647]** Compound I-95 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(propylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 401.2

**[0648]** Compound I-97 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(isopropylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 401.2

**[0649]** Compound I-104 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(thiophen-2-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 441.1

**[0650]** Compound I-106 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(benzylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 449.2

**[0651]** Compound I-107 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)benzenesulfonamide; MS (ES<sup>+</sup>) 435.2

[0652] Compound I-108 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)benzamide; MS (ES<sup>+</sup>) 399.2

[0653] Compound I-110 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(phenylsulfonyl)piperazin-1-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 435.2

[0654] Compound I-111 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2,2-dimethylpropanamide; MS (ES<sup>+</sup>) 379.2

[0655] Compound I-112 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(cyclohexyl)methanone; MS (ES<sup>+</sup>) 405.2

For compounds I-174 to I-275, please see Tables 7 and 8 for analytical data.

[0656] Compound I-174 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(cyclobutyl)methanone

[0657] Compound I-175 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-ethylbutan-1-one

[0658] Compound I-176 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(pyridin-2-yl)methanone

[0659] Compound I-177 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(2,6-difluorophenyl)methanone

[0660] Compound I-181 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2-(2-methoxyethoxy)ethanamide

[0661] Compound I-182 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2-methoxyethanamide

[0662] Compound I-184 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2-(dimethylamino)ethanamide

[0663] Compound I-185 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)ethanamide

[0664] Compound I-188 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)pyridine-3-carboxamide

[0665] Compound I-193 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)pyridine-2-carboxamide

[0666] Compound I-199 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-methylpropan-1-one

[0667] Compound I-204 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)cyclopentanecarboxamide

[0668] Compound I-205 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2-phenoxyethanamide

[0669] Compound I-208 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(phenyl)methanone

[0670] Compound I-211 (S)—N-(1-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)pyrrolidin-3-yl)-2-ethylbutanamide

[0671] Compound I-264 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-(2-methoxyethoxy)ethanone

[0672] Compound I-265 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-methoxyethanone

[0673] Compound I-266 2-methoxyethyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0674] Compound I-267 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(pyrrolidin-1-yl)methanone

[0675] Compound I-268 methyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0676] Compound I-269 3-(1H-benzo[d]imidazol-2-yl)-5-(4-(furan-3-ylsulfonyl)piperazin-1-yl)pyrazin-2-amine

[0677] Compound I-270 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(piperidin-1-yl)methanone

[0678] Compound I-271 ethyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0679] Compound I-272 but-2-ynyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0680] Compound I-273 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)-2-phenoxyethanone

[0681] Compound I-274 benzyl 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazine-1-carboxylate

[0682] Compound I-275 (4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)piperazin-1-yl)(2,6-dimethylphenyl)methanone

[0683] Compound I-295 3-(1H-benzimidazol-2-yl)-5-(2-ethylsulfonyl-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-5-yl)pyrazin-2-amine MS (ES<sup>+</sup>) 414.2


#### Example 17


3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(2-methylpyridin-4-yl)pyrazine-2-carboxamide (Compound I-166)

[0684]

#### SCHEME XVII

Compound I-166





**[0685]** Compound I-166 was prepared using Method A, Step 1 followed by Method Q, Steps 1-4.

#### Method Q

Step 1: Methyl 3-amino-6-(1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxylate

**[0686]** A mixture of methyl 3-amino-6-bromo-pyrazine-2-carboxylate (6 g, 25.86 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (7.996 g, 25.86 mmol),  $\text{Na}_2\text{CO}_3$  (38.79 mL of 2 M, 77.58 mmol) and dichloropalladium; triphenylphosphane (544.5 mg, 0.7758 mmol) was stirred in DME (120.0 mL) at 100° C. for 5 hours. After this time water and EtOAc were added and the organics separated, dried and concentrated in vacuo. The residue was redissolved in DCM (20 mL) and TFA (20 mL) and the resulting solution stirred at ambient temperature for 2 hours. After this time saturated  $\text{NaHCO}_3$  was added and the organics separated, dried and concentrated. The solid was then triturated with MeOH, filtered and washed with MeOH to give the desired product as a yellow solid (4.15 g, 69%).  $^1\text{H}$  NMR (400.0 MHz, DMSO) d 2.43 (3H, s), 3.41

(2H, m), 3.80-4.02 (5H, m), 6.52 (1H, s), 7.33 (2H, s), 8.52 (1H, s) ppm; MS ( $\text{ES}^+$ ) 234.97

#### Method Q

Step 2: Methyl 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxylate

**[0687]** A solution of methyl 3-amino-6-(1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxylate (4.15 g, 17.72 mmol) and N-ethyl-N-isopropyl-propan-2-amine (4.580 g, 6.173 mL, 35.44 mmol) in DMF (68.89 mL) was treated with ethanesulfonyl chloride (2.506 g, 1.847 mL, 19.49 mmol) and allowed to stir at ambient temperature for 16 hours. The reaction mixture was treated with water and the resulting precipitate collected and washed with water (5.5 g, 95% Yield). MS ( $\text{ES}^+$ ) 327.01

#### Method Q

Step 3: 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxylic acid

**[0688]** A solution of methyl 3-amino-6-(1-ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylate (5.5 g, 16.85 mmol) in MeOH (30 mL) was treated with a solution of NaOH (8.425 mL of 2 M, 16.85 mmol) in  $\text{H}_2\text{O}$  (30 mL). The resulting mixture was heated to 60° C. for 2 hours. After this time the solution was allowed to cool, neutralised with HCl and the resultant precipitate filtered (3.6 g, 65% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO) d 1.20 (3H, t), 2.65 (2H, br s), 3.11 (2H, q), 3.44 (2H, m), 3.94 (2H, m), 6.60 (1H, m), 7.51 (2H, br s), 7.95 (1H, s), 8.52 (1H, s) ppm; MS ( $\text{ES}^+$ ) 312.97

#### Method Q

Step 4: 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(2-methylpyridin-4-yl)pyrazine-2-carboxamide

**[0689]** A solution of 3-amino-6-(1-ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylic acid (79.87 mg, 0.255 mmol), CDI (82.92 mg, 0.511 mmoles), triethylamine (107  $\mu\text{L}$ , 0.767 mmoles), and DMAP (1.56 mg, 0.012 mmol) was treated with 2-methylpyridin-4-amine (55.30 mg, 0.511 mmol). The reaction was allowed to stir at 100° C. for 16 hours, filtered and purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{M}$ , 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the title compound (30.4 mg, 21% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO) d 11.10 (s, 1H), 8.69-8.67 (m, 2H), 8.25-8.22 (m, 2H), 7.72 (s, 2H), 6.77 (d, J=3.3 Hz, 1H), 3.99 (d, J=2.9 Hz, 2H), 3.48 (t, J=5.7 Hz, 2H), 3.16 (q, J=7.4 Hz, 2H), 2.79 (s, 2H), 2.68 (s, 3H) and 1.25 (t, J=7.3 Hz, 3H) ppm; MS ( $\text{ES}^+$ ) 403.0

**[0690]** The following compounds were all prepared using the same method or a method similar to the one described for Compound I-166 in Example 17 above.

**[0691]** Compound I-165 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(2-methoxypyridin-3-yl)pyrazine-2-carboxamide; MS ( $\text{ES}^+$ ) 419

**[0692]** Compound I-167 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(4-methylpyridin-2-yl)pyrazine-2-carboxamide MS ( $\text{ES}^+$ ) 403

**[0693]** Compound I-168 3-amino-N-(1,3-dimethyl-1H-pyrazol-5-yl)-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxamide MS (ES+) 406

**[0694]** Compound I-169 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(1H-indazol-3-yl)pyrazine-2-carboxamide MS (ES+) 428

**[0695]** Compound I-170 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(pyrazin-2-yl)pyrazine-2-carboxamide MS (ES+) 390


**[0696]** Compound I-171 3-amino-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-N-(pyridin-2-yl)pyrazine-2-carboxamide MS (ES+) 389

**[0697]** Compound I-172 3-amino-N-(1H-benzo[d]imidazol-2-yl)-6-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazine-2-carboxamide MS (ES+) 428


### Example 18

3-(5-(3-aminophenyl)-4H-1,2,4-triazol-3-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine  
(Compound I-57)

**[0698]**



-continued



Compound I-57

**[0699]** Compound I-57 was prepared using Method A, Step 1 followed by Method R, Steps 1-3.

### Method R


Step 1: 3-amino-6-bromopyrazine-2-carbohydrazide

**[0700]** A mixture of methyl 3-amino-6-bromo-pyrazine-2-carboxylate (5 g, 21.55 mmol) and hydrazine hydrate (5.397 g, 5.245 mL, 107.8 mmol) were heated to 100° C. for 1.5 hours. Water was added and the product collected by filtration, washed with methanol and dried to give 3-amino-6-bromo-pyrazine-2-carbohydrazide as a yellow solid (5.02 g, 100.4% Yield). MS (ES+) 233.78

### Method R

Step 2: 3-(5-(3-aminophenyl)-4H-1,2,4-triazol-3-yl)-5-bromopyrazin-2-amine

**[0701]**




**[0702]** A mixture of 3-amino-6-bromo-pyrazine-2-carbohydrazide (5 g, 21.55 mmol), 3-aminobenzamidine dihydrochloride (4.484 g, 21.55 mmol) and NaOEt (4.399 g, 64.65 mmol) were combined in DMF (50.00 mL) and heated to 100° C. in the microwave for 20 min. Water was added and the precipitate collected. Used without further purification. Assumed 100% Yield. MS (ES+) 334.02

## Method R

Step 3: 3-(5-(3-aminophenyl)-4H-1,2,4-triazol-3-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine

[0703]





[0704] 3-[5-(3-aminophenyl)-4H-1,2,4-triazol-3-yl]-5-bromo-pyrazin-2-amine (220 mg, 0.6623 mmol) was heated in neat 1-ethanesulfonyl piperazine (590.4 mg, 3.312 mmol) in the microwave at 190° C. for 2 hours. The reaction was treated with EtOAc and H<sub>2</sub>O, the two layers separated and the organics dried and concentrated to give a yellow oil which was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound as a yellow solid (23.9 mg, 8% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 8.38 (s, 2H), 8.04 (s, 1H), 7.33 (s, 2H), 7.24 (s, 1H), 7.09 (s, 1H), 6.96 (s, 1H), 6.60 (s, 1H), 3.55 (s, 4H), 3.28 (s, 4H), 3.08 (d, J=5.7 Hz, 2H) and 1.20 (s, 3H) ppm; MS (ES<sup>+</sup>) 430.0

## Example 19

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(imidazo[2,1-b]thiazol-6-yl)pyrazin-2-amine (Compound I-62)

[0705]



Compound I-62

[0706] Compound I-62 was prepared using Method A, Step 1 followed by Method B, Steps 1-4 followed by Method S, Steps 1-4.

#### Method S

Step 1: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-methoxy-N-methylpyrazine-2-carboxamide

[0707] A mixture of 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazine-2-carboxylic acid (1 g, 3.171 mmol), N-methoxymethanamine hydrochloride (371.2 mg, 3.805 mmol) and triethylamine (705.9 mg, 972.3  $\mu$ L, 6.976 mmol) were dissolved in DMF (10 mL) and treated with TBTU (1.527 g, 4.756 mmol). The mixture was allowed to stir at ambient temperature for 1 hour. The reaction was treated with water, extracted with EtOAc (4 $\times$ ), and the organics separated, dried ( $\text{MgSO}_4$ ) and concentrated. The resultant residue was purified on silica gel by flash column chromatography (0-100% EtOAc/Petrol) to afford the title compound as a yellow solid (864 mg, 76% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.21 (t, 3H), 3.10 (q, 2H), 3.27-3.30 (m, 4H), 3.34 (s, 3H), 3.38-3.40 (m, 4H), 3.68 (s, 3H), 5.87 (br s, 2H) and 7.99 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 359.0

#### Method S

Step 2: 1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)ethanone

[0708] 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-methoxy-N-methylpyrazine-2-carboxamide (864 mg, 2.411 mmol) was dissolved in THF (10 mL) and cooled to  $-20^\circ\text{C}$ . under nitrogen.  $\text{MeMgBr}$  (2.411 mL of 3 M, 7.233 mmol) was added dropwise and the reaction was left to stir at this temperature for 15 minutes after which time the reaction was allowed to slowly warm to ambient temperature. A further  $\text{MeMgBr}$  (803.7  $\mu$ L of 3 M, 2.411 mmol) was added and analysis showed after 10 minutes showed no SM remaining. The mixture was quenched with 10 mL 1M HCl, stirred for 5 mins and then extracted with EtOAc (3 $\times$ ), dried ( $\text{MgSO}_4$ ), filtered and the solvent removed in vacuo to give the product as an orange solid (510 mg, 68% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.23 (t, 3H), 2.54 (s, 3H), 3.11 (q, 2H), 3.30-3.32 (m, 4H), 3.47-3.49 (m, 4H), 7.42 (br s, 2H) and 8.34 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 314.0

#### Method S

Step 3: 1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)-2-bromoethanone

[0709] A solution of 1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)ethanone (510 mg, 1.627 mmol) in AcOH (2 mL) and HBr in AcOH (0.5 mL of 33% w/v, 2.039 mmol) was treated with pyridinium tribromide (544.5 mg, 1.708 mmol) and the mixture allowed to stir at ambient temperature for 2 hours. The reaction was poured into ether (50 mL), partitioned between EtOAc/water and extract with EtOAc (3 $\times$ ). The mixture was dried ( $\text{MgSO}_4$ ), filtered, evapo-

rated down and the resultant residue was purified on silica gel by flash column chromatography (0-100% EtOAc/Petrol) to afford the title compound as a orange oil (68 mg, 11% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.23 (t, 3H), 3.12 (q, 2H), 3.31 (m, 4H), 3.52 (m, 4H), 4.86 (s, 2H), 7.21 (s, 1H), 8.34 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 393.91

#### Method S


Step 4: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(imidazo[2,1-b]thiazol-6-yl)pyrazin-2-amine

[0710] 1-[3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl]-2-bromo-ethanone (35 mg, 0.08922 mmol) was suspended in EtOH (2 mL) and 2 aminothiazole (8.935 mg, 0.08922 mmol) was added. The mixture was heated at reflux for 4 hours and then allowed to cool to ambient temperature. The solution was filtered through a  $0.45\ \mu\text{m}$  PTFE filter and the solvent removed under a stream of nitrogen. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{m}$ , 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (55 mg, 10% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.14 (t,  $J=7.3$  Hz, 3H), 3.01 (q,  $J=7.4$  Hz, 2H), 3.21-3.23 (m, 4H), 3.34-3.36 (m, 4H), 7.28 (d,  $J=4.4$  Hz, 1H), 7.67 (s, 1H), 7.89 (d,  $J=4.4$  Hz, 1H) and 8.23 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 394.0


#### Example 20

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(6-phenylpyrimidin-4-yl)pyrazin-2-amine (Compound I-74)

[0711]



-continued



**[0712]** Compound I-74 was prepared using Method A, Step 1 followed by Method B, Steps 1-4 followed by Method S, Steps 1-2 followed by Method T, Steps 1-2.

## Method T

Step 1: (E)-1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)-3-phenylprop-2-en-1-one

**[0713]** A solution of NaOH in methanol (191.5  $\mu$ L of 10% w/v, 0.4787 mmol) was added to a solution of 1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)ethanone (150 mg, 0.4787 mmol) and benzaldehyde (50.80 mg, 48.66  $\mu$ L, 0.4787 mmol) in methanol (4 mL) (1 mL of dichloromethane was added to aid solubility). The reaction mixture was allowed to stir overnight at RT and then filtered and washed with a small amount of methanol. The bright orange coloured solid was taken through crude to the next stage. (104.6 mg, 54% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.21 (t, 3H), 3.13 (q, 2H), 3.35-3.37 (m, 4H), 3.54-3.57 (m, 4H), 7.47-7.51 (m, 3H), 7.55 (br s, 2H), 7.72 (d, 1H), 7.79-7.81 (m, 2H), 8.28 (d, 1H), 8.41 (s, 1H) ppm; MS (ES<sup>+</sup>) 402.01

## Method T

Step 2: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(6-phenylpyrimidin-4-yl)pyrazin-2-amine

**[0714]** A mixture of 1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)-3-phenylprop-2-en-1-one (45 mg, 0.1121 mmol), formamidine hydrochloride (9.026 mg, 0.1121 mmol) and NaOH (4.932 mg, 0.1233 mmol) were heated at reflux for 1 hour. The reaction mixture was allowed to cool to ambient temperature and the solvent removed in vacuo. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (73 mg, 4% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.25 (t, 3H), 3.12 (q, 2H), 3.36-3.38 (m, 4H), 3.56-3.59 (m, 4H), 7.15 (br s, 2H), 7.60-7.61 (m, 3H), 8.25-8.26 (m, 3H), 8.76 (s, 1H) and 9.31 (s, 1H) ppm; MS (ES<sup>+</sup>) 426.0

**[0715]** Compound I-73 was also prepared using Method A, Step 1 followed by Method B, Steps 1-4 followed by Method S, Steps 1-2 followed by Method T, Step 1.


**[0716]** Compound I-73 (E)-1-(3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-yl)-3-phenylprop-2-en-1-one;  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  1.21 (t, 3H), 3.13 (q, 2H), 3.35-3.37 (m, 4H), 3.54-3.57 (m, 4H), 7.47-7.51 (m, 3H), 7.55 (br s, 2H), 7.72 (d, 1H), 7.79-7.81 (m, 2H), 8.28 (d, 1H), 8.41 (s, 1H) ppm; MS (ES<sup>+</sup>) 402.01


## Example 21

3-(1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine  
(Compound I-160)

**[0717]**

SCHEME XXI






[0718] Compound I-160 was prepared using Method U, Steps 1-7.

#### Method U

##### Step 1:

5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-amine

[0719]



[0720] (Trimethylsilyl)acetylene (1.845 g, 2.655 mL, 18.78 mmol) was added dropwise to a solution of 3,5-dibromopyrazin-2-amine (5 g, 19.77 mmol), triethylamine (10.00 g, 13.77 mL, 98.85 mmol), Copper(I) iodide (451.7 mg, 2.372 mmol) and Pd(PPh<sub>3</sub>)<sub>4</sub> (1.142 g, 0.9885 mmol) in DMF (25.00 mL) and the resulting solution stirred at ambient temperature for 30 minutes. The reaction was diluted with EtOAc and water and the layers separated. The aqueous layer was extracted further with EtOAc and the combined organics washed with water, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The mixture was purified on silica gel by flash column chromatography (0-15% EtOAc/Petrol) to afford the product as a yellow solid (3.99 g, 75% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.30 (s, 9H), 8.06 (s, 1H) ppm; MS (ES<sup>+</sup>) 271.82

#### Method U

Step 2: Tert-butyl N-tert-butoxycarbonyl-N-[5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-yl]carbamate

[0721] 5-bromo-3-(2-trimethylsilyl)ethynylpyrazin-2-amine (480 mg, 1.776 mmol) was dissolved in DCM (15 mL) and treated with BOC anhydride (1.163 g, 1.224 mL, 5.328 mmol) followed by DMAP (21.70 mg, 0.1776 mmol). The mixture was allowed to stir at ambient temperature overnight. The reaction was washed with NaHCO<sub>3</sub>, extracted with DCM (3×), dried (MgSO<sub>4</sub>), filtered and concentrated. The resultant brown oil was purified (ISCO Companion™, 40 g column, 0-10% EtOAc/Petrol) to afford the product as a colourless oil (641 mg, 77% Yield).

[0722] <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.00 (s, 9H), 1.11 (s, 18H) and 8.63 (s, 1H) ppm

#### Method U

Step 3: Tert-butyl N-tert-butoxycarbonyl-5-(piperazin-1-yl)-3-((trimethylsilyl)ethynyl)pyrazin-2-ylcarbamate

[0723] Tert-butyl N-tert-butoxycarbonyl-N-[5-bromo-3-((trimethylsilyl)ethynyl)pyrazin-2-yl]carbamate (640 mg, 1.360 mmol) was dissolved in DMF (5 mL) and treated with piperazine (351.4 mg, 4.080 mmol). The reaction was allowed to heat at 100° C. for 15 minutes and then allowed to cool to ambient temperature and diluted with EtOAc/water. The aqueous layer was extracted with EtOAc and the com-

bined organics washed with water (2×) and brine (2×). This was dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by column chromatography (ISCO Companion™, 12 g column, 0-100% EtOAc/Petrol to 10% MeOH/DCM/1% NH<sub>4</sub>OH) to afford the title compound as a yellow oil (494 mg, 76% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 0.03 (s, 9H), 1.18 (s, 18H), 2.99 (m, 4H), 3.29 (m, 4H) and 7.88 (s, 1H) ppm; MS (ES<sup>+</sup>) 476.08

#### Method U

Step 4: Tert-butyl N-tert-butoxycarbonyl-5-(4-(ethylsulfonyl)piperazin-1-yl)-3-((trimethylsilyl)ethynyl)pyrazin-2-ylcarbamate

[0724] Ethanesulfonyl chloride (147.0 mg, 108.3 μL, 1.143 mmol) was added slowly to a stirred solution of tert-butyl N-tert-butoxycarbonyl-5-(piperazin-1-yl)-3-((trimethylsilyl)ethynyl)pyrazin-2-ylcarbamate (494 mg, 1.039 mmol) and DIPEA (161.2 mg, 217.3 μL, 1.247 mmol) in DMF (5 mL) at ambient temperature. The reaction was allowed to stir for 5 minutes and then partitioned between EtOAc/water. The aqueous layer was extracted with EtOAc (3×) and the combined organics washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated to give the product as a terracotta oil (645 mg, quantitative yield). MS (ES<sup>+</sup>) 590.05

#### Method U

Step 5: Tert-butyl N-tert-butoxycarbonyl-N-[5-(4-ethylsulfonylpiperazin-1-yl)-3-ethynylpyrazin-2-yl]carbamate

[0725] Tert-butyl N-tert-butoxycarbonyl-5-(4-(ethylsulfonyl)piperazin-1-yl)-3-((trimethylsilyl)ethynyl)pyrazin-2-ylcarbamate (589 mg, 1.037 mmol) was dissolved in anhydrous MeOH (10 mL) and treated with K<sub>2</sub>CO<sub>3</sub> (14.33 mg, 0.1037 mmol). The reaction was allowed to stir at ambient temperature for 45 minutes. The solvent was removed in vacuo and the residue redissolved in DCM and washed with water followed by brine. The solution was dried (MgSO<sub>4</sub>), filtered and concentrated and the residue was purified by column chromatography (ISCO Companion™, 24 g column, 0-100% EtOAc/Petrol) to afford the title compound as an orange solid (287 mg, 60% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.23 (t, J=7.4 Hz, 3H), 1.38 (s, 18H), 3.11 (q, 2H), 3.26-3.34 (m, 4H), 3.69-3.71 (m, 4H), 4.65 (s, 1H) and 8.27 (s, 1H) ppm; MS (ES<sup>+</sup>) 396.0

#### Method U

Step 6: Tert-butyl N-tert-butoxycarbonyl-3-(1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-ylcarbamate

[0726] A mixture of tert-butyl N-tert-butoxycarbonyl-N-[5-(4-ethylsulfonylpiperazin-1-yl)-3-ethynylpyrazin-2-yl]carbamate (50 mg, 0.1009 mmol) and 4-Azidoaniline hydrochloride (17.32 mg, 0.1009 mmol) were suspended in t-BuOH (1 mL)/Water (1.000 mL). The reaction was treated with (+)-Sodium L-ascorbate (1.999 mg, 0.01009 mmol) followed by CuSO<sub>4</sub>·0.5H<sub>2</sub>O (0.2519 mg, 0.1103 μL, 0.001009 mmol) and the mixture allowed to stir at ambient temperature for 16 hours. The reaction was treated with DIPEA (13.04 mg, 17.57 μL, 0.1009 mmol) and allowed to heat to 40° C. for 5 hours and then cooled to ambient temperature. The mixture

was partitioned between EtOAc/NaHCO<sub>3</sub> and the aqueous layer extracted with EtOAc (2×), dried (MgSO<sub>4</sub>), filtered and concentrated. The material was used crude in the next step (quantitative yield assumed). MS (ES<sup>+</sup>) 630.00

#### Method U

Step 7: 3-(1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine

[0727] Tert-butyl N-tert-butoxycarbonyl-3-(1-(4-aminophenyl)-1H-1,2,3-triazol-4-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-ylcarbamate (64 mg, 0.1016 mmol) was dissolved in DCM (5 mL) and treated with TFA (500  $\mu$ L, 6.490 mmol). The reaction was allowed to stir at ambient temperature overnight and then the solvent removed in vacuo.


The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ m, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (24.3 mg, 30% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO)  $\delta$  1.24 (t, J=7.4 Hz, 3H), 3.11 (q, J=7.4 Hz, 2H), 3.30-3.33 (m, 4H), 3.50-3.52 (m, 4H), 6.77 (d, J=8.6 Hz, 2H), 7.67 (d, J=8.8 Hz, 2H), 7.88 (s, 1H) and 9.09 (s, 1H) ppm; MS (ES<sup>+</sup>) 430.0

#### Example 22

3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (Compound I-276)

[0728]

#### SCHEME XXII



[0729] Compound I-276 was prepared using Method A, Step 1 followed by Method V, Steps 1-5.

#### Method V

Step 1: 3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-bromopyrazin-2-amine

[0730] A mixture of pyridin-1-ium-1,2-diamine (185 mg, 0.7805 mmol), methyl 3-amino-6-bromo-pyrazine-2-carboxylate (217.3 mg, 0.9366 mmol) and NaOH (49.96 mg, 1.249 mmol) in EtOH (7 mL) were heated at 80° C. After 1 hour further methyl 3-amino-6-bromo-pyrazine-2-carboxylate (50 mg, 0.2155 mmol) was added and the reaction allowed to heat for a further 1.5 hours. The reaction was allowed to cool to ambient temperature and the solvent removed in vacuo. The mixture was partitioned between DCM/Water and the layers separated. The organics were dried (MgSO<sub>4</sub>), filtered and concentrated to give the product as an off-white solid (134 mg, 59% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 7.34 (td, J=6.9, 2.7 Hz, 1H), 7.77-7.81 (m, 1H), 7.87 (br s, 2H), 7.97 (d, 1H), 8.31 (s, 1H) and 9.09 (d, 1H) ppm; MS (ES<sup>+</sup>) 292.8

#### Method V

Step 2: Tert-butyl N-tert-butoxycarbonyl-3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-bromopyrazin-2-ylcarbamate

[0731] A solution of 3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-bromopyrazin-2-amine (168 mg, 0.5771 mmol) in DCM (15 mL) was treated with BOC anhydride (377.8 mg, 397.7 μL, 1.731 mmol) followed by DMAP (7.050 mg, 0.05771 mmol). The mixture was allowed to stir at ambient temperature for 16 hours. Analysis showed SM remaining therefore further BOC anhydride (377.8 mg, 397.7 μL, 1.731 mmol) and DMAP (7.050 mg, 0.05771 mmol) were added and the mixture allowed to stir for a further 16 hours. The reaction was washed with NaHCO<sub>3</sub>, extracted with DCM (3×), dried (MgSO<sub>4</sub>), filtered and concentrated. The residue was purified by column chromatography (ISCO Companion™, 24 g column, 0-100% EtOAc/Petrol) to afford the title compound as a white solid (167 mg, 59% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.28 (s, 18H), 7.38-7.42 (m, 1H), 7.82-7.86 (m, 1H), 7.99-8.02 (m, 1H), 9.06 (s, 1H) and 9.05-9.10 (m, 1H) ppm; MS (ES<sup>+</sup>) 493.0

#### Method V

Step 3: Tert-butyl 4-(6-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-(bis(tert-butoxycarbonyl)amino)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0732] Tert-butyl N-tert-butoxycarbonyl-3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-bromopyrazin-2-ylcarbamate (92 mg, 0.1872 mmol) was dissolved in DMF (1.5 mL) and treated with tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (86.83 mg, 0.2808 mmol), PdCl<sub>2</sub>(PPh<sub>3</sub>)<sub>2</sub> (13.14 mg, 0.01872 mmol) and K<sub>2</sub>CO<sub>3</sub> (280.8 μL of 2 M, 0.5616 mmol). The mixture was allowed to heat at 100° C. for 1 hour and then allowed to cool to ambient temperature. The reaction was treated with EtOAc/water, the organics separated, washed with water (2×) followed by brine (2×), dried (MgSO<sub>4</sub>), filtered and concentrated to a brown oil. The residue was purified by column

chromatography (ISCO Companion™, 12 g column, 0-100% EtOAc/Petrol) to afford the title compound as a white solid (77 mg, 69% Yield). MS (ES<sup>+</sup>) 594.0

#### Method V

Step 4: 3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0733] Tert-butyl 4-[5-(bis(tert-butoxycarbonyl)amino)-6-([1,2,4]triazolo[1,5-a]pyridin-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate (77 mg, 0.1297 mmol) was dissolved in DCM (1.6 mL) and treated with TFA (400 μL, 5.192 mmol). The mixture was allowed to stir at ambient temperature for 2 hours and the solvent removed in vacuo. The mixture was azeotroped with DCM (3×) and Ether (3×) and dried under vacuum to furnish the product as a yellow solid (104 mg, quantitative yield). MS (ES<sup>+</sup>) 294.0

#### Method V

Step 5: 3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine


[0734] 5-(1,2,3,6-tetrahydropyridin-4-yl)-3-([1,2,4]triazolo[1,5-a]pyridin-2-yl)pyrazin-2-amine (82 mg, 0.1291 mmol) was dissolved in DCM (4 mL) and treated with triethylamine (78.38 mg, 108.0 μL, 0.7746 mmol) followed by ethanesulfonyl chloride (18.26 mg, 13.46 μL, 0.1420 mmol). The reaction mixture was allowed to stir at ambient temperature for 1 hour when analysis showed SM remaining. The mixture was treated with further ethanesulfonyl chloride (16.60 mg, 12.23 μL, 0.1291 mmol) and stirred for a further hour. The reaction was diluted with DCM, washed with water and the aqueous layer extracted further with dichloromethane (2×). The combined organics were washed with saturated aqueous NaHCO<sub>3</sub>, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The resultant residue was purified by column chromatography (ISCO Companion™, 12 g column, 0-2.5% MeOH/DCM) to afford the title compound which was freeze-dried to give a yellow solid (9 mg, 17% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.24 (t, J=7.4 Hz, 3H), 2.71 (br s, 2H), 3.14 (q, J=7.3 Hz, 2H), 3.48 (t, J=5.7 Hz, 2H), 3.98 (d, J=2.9 Hz, 2H), 6.62 (br s, 1H), 7.30-7.33 (m, 1H), 7.72 (br s, 2H), 7.75-7.79 (m, 1H), 7.98 (d, J=8.8 Hz, 1H), 8.40 (s, 1H) and 9.11 (d, J=6.7 Hz, 1H) ppm; MS (ES<sup>+</sup>) 386.0


#### Example 23

3-(benzofuran-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine (Compound I-29)

[0735]

#### SCHEME XXIII





[0736] Compound I-29 was prepared using Method W, Steps 1-3.

#### Method W

##### Step 1:

##### 3-(benzofuran-2-yl)-5-bromopyrazin-2-amine

[0737] A solution of 3,5-dibromopyrazin-2-amine (200 mg, 0.7908 mmol), benzofuran-2-ylboronic acid (140.9 mg, 0.8699 mmol),  $\text{NaHCO}_3$  (199.3 mg, 2.372 mmol) and palladium; triphenylphosphane (91.38 mg, 0.07908 mmol) in dimethoxyethane (2.000 mL) and water (1.000 mL) was heated at 120° C. in the microwave for 10 minutes. The reaction was diluted with EtOAc/Water and the layers separated. The aqueous layer was extracted further with EtOAc (2×), dried ( $\text{MgSO}_4$ ) and concentrated in vacuo. The mixture was purified on silica gel by flash column chromatography (0-20% EtOAc/Petrol) to afford the product as a bright yellow solid (191 mg, 83% Yield). MS ( $\text{ES}^+$ ) 291.89

#### Method W

##### Step 2: Tert-butyl

##### 3-(benzofuran-2-yl)-5-bromopyrazin-2-ylcarbamate

[0738] A solution of 3-(benzofuran-2-yl)-5-bromopyrazin-2-amine (179 mg, 0.6170 mmol) was treated with

di-tert-butylidicarbonate (404.0 mg, 425.3  $\mu\text{L}$ , 1.851 mmol) and 4-(dimethylamino)pyridine (7.538 mg, 0.06170 mmol) in dichloromethane (5 mL) and the reaction mixture allowed to stir overnight at ambient temperature. The reaction was diluted with DCM (10 mL) and washed with 1N HCl (1×), sat aqueous  $\text{NaHCO}_3$  (1×) and brine (1×), dried ( $\text{MgSO}_4$ ) and concentrated in vacuo. The material was used crude in the next step (235 mg, 98% Yield).

#### Method W

##### Step 3: 3-(benzofuran-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine


[0739] A solution of tert-butyl N-[3-(benzofuran-2-yl)-5-bromo-pyrazin-2-yl]carbamate, 1-ethylsulfonylpiperazine (164.5 mg, 0.9228 mmol) and 2-methoxyethanol (3 mL) were heated at 100° C. for 3 hours. The reaction mixture was allowed to cool to RT, treated with HCl in dioxane (384.5  $\mu\text{L}$  of 4 M, 1.538 mmol) and allowed to stir at RT for 3 hours. The mixture was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{M}$ , 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound as an orange solid (15.5 mg, 12% Yield).  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$  1.24 (3H, t), 3.11 (2H, q), 3.45 masked signal, 3.51 (4H, m), 6.12 (2H, br s), 7.30 (1H, m), 7.34 (1H, m), 7.47 (1H, s), 7.71 (2H, m), 7.98 (1H, s) ppm; MS ( $\text{ES}^+$ ) 388.01


#### Example 24

##### 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyloxazol-2-yl)pyrazin-2-amine (Compound I-36)

[0740]

#### SCHEME XXIV





**[0741]** Compound I-36 was prepared using Method A, Step 1 followed by Method B, Steps 1-4 followed by Method X, Steps 1-2.

#### Method X

Step 1: 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-(2-oxo-2-phenylethyl)pyrazine-2-carboxamide

**[0742]** 2-amino-1-phenyl-ethanone hydrochloride (136.1 mg, 0.7928 mmol) was added to a solution of 3-amino-6-(4-

ethylsulfonylpiperazin-1-yl)pyrazine-2-carboxylic acid (250 mg, 0.7928 mmol) and DIPEA (512.3 mg, 690.4  $\mu$ L, 3.964 mmol) in  $\text{CH}_2\text{Cl}_2$  (3.332 mL) and DMF (833.2  $\mu$ L) and the resulting solution stirred at room temperature for 1 hour. The reaction was diluted with EtOAc/water and the layers separated. The aqueous layer was extracted further with EtOAc (2 $\times$ ) and the combined organics washed with water (3 $\times$ ), dried ( $\text{MgSO}_4$ ) and concentrated in vacuo. The mixture was purified on silica gel by flash column chromatography (100% EtOAc) to afford the product (229 mg, 67% Yield).  $^1\text{H}$  NMR ( $\text{CDCl}_3$ )  $\delta$  1.28 (3H, t), 3.06 (2H, q), 3.51-3.53 (4H, m), 3.56-3.59 (4H, m), 4.95 (2H, d), 6.2 (2H, br s), 7.56 (2H, m), 7.68 (1H, m), 8.05 (3H, m), 8.65 (1H, m) ppm; MS ( $\text{ES}^+$ ) 433.0

#### Method X

Step 2: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-phenyloxazol-2-yl)pyrazine-2-amine


**[0743]** 3-amino-6-(4-(ethylsulfonyl)piperazin-1-yl)-N-phenacylpyrazine-2-carboxamide (80 mg, 0.1850 mmol) in  $\text{H}_2\text{SO}_4$  (1.5 mL) was stirred at ambient temperature for 16 hours. The reaction was neutralised by addition of  $\text{NaHCO}_3$  solution and then extracted with EtOAc (3 $\times$ ), combined organics dried ( $\text{MgSO}_4$ ) and concentrated in vacuo. The resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound as an orange solid (4.5 mg, 6% Yield).  $^1\text{H}$  NMR ( $\text{DMSO}$ )  $\delta$  1.25 (3H, t), 3.11 (2H, q), 3.35 masked signal, 3.54 (4H, m), 7.00 (2H, br s), 7.42 (1H, m), 7.53 (2H, m), 7.84 (2H, m), 7.99 (1H, s), 8.14 (1H, s) ppm; MS ( $\text{ES}^+$ ) 415.0


#### Example 25

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-methyl-1,2,4-oxadiazol-3-yl)pyrazine-2-amine (Compound I-38)

**[0744]**

#### SCHEME XXV





**[0745]** Compound I-38 was prepared using Method Y, Steps 1-7.

#### Method Y

##### Step 1: 3-amino-6-bromopyrazine-2-carbonitrile

**[0746]** 3-amino-6-bromo-pyrazine-2-carboxamide (500 mg, 2.304 mmol) was dissolved in pyridine (5 mL) and treated with  $\text{POCl}_3$  (666  $\mu\text{L}$ , 7.145 mmol) dropwise at  $0^\circ\text{C}$ . After the addition was complete the reaction was allowed to warm to ambient temperature and stirred for 1.5 hours. The solvent was removed in vacuo and the reaction carefully treated with  $\text{NaHCO}_3$  to basify. The mixture was diluted with EtOAc and the layers separated. The aqueous layer was further extracted with EtOAc (1 $\times$ ) and the combined organics washed with brine (1 $\times$ ), 0.1 M HCl (1 $\times$ ) and water (1 $\times$ ), dried ( $\text{MgSO}_4$ ), filtered and concentrated to furnish the product as a brown solid (197 mg, 43% Yield).  $^1\text{H}$  NMR (DMSO)  $\delta$  7.62 (2H, br s), 8.45 (1H, s) ppm

#### Method Y

##### Step 2: (E)-3-amino-6-bromo-N-hydroxypyrazine-2-carboximidamide

**[0747]** 3-amino-6-bromo-pyrazine-2-carbonitrile (4.5958 g, 23.09 mmol) was dissolved in MeOH (91.92 mL), cooled to  $0^\circ\text{C}$  and treated with hydroxylamine hydrochloride (1.605 g, 23.09 mmol) and triethylamine (2.336 g, 3.218 mL, 23.09 mmol). The reaction was allowed to warm to ambient temperature, stirred for 48 hours and then evaporated to dryness. Methanol (a minimum amount) was added, the mixture sonicated, filtered and washed with MeOH to give the product as a brown solid 5.27 g, 98% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  5.88 (s, 2H), 7.64 (br s, 2H), 8.14 (s, 1H) and 10.38 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 233.75

#### Method Y

##### Step 3: (E)-3-amino-6-bromo-N-(ethanoyloxy)pyrazine-2-carboximidamide

**[0748]** A suspension of 3-amino-6-bromo-N'-hydroxypyrazine-2-carboximidamide (5.2697 g, 22.71 mmol) in DCM (68.46 mL) and triethylamine (2.528 g, 3.482 mL, 24.98 mmol) was treated with acetyl chloride (1.961 g, 1.776 mL, 24.98 mmol) and allowed to stir at ambient temperature for 15 minutes. The solvent and volatile reagents were removed in vacuo and the residue triturated from MeOH/

ether. The resultant precipitate was isolated by filtration to furnish the product as a brown solid (8.40 g, quantitative yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  2.18 (3H, s), 6.90 (2H, br s), 7.78 (2H, br s), 8.29 (1H, s) ppm; MS ( $\text{ES}^+$ ) 275.78

#### Method Y

##### Step 4: 5-bromo-3-(5-methyl-1,2,4-oxadiazol-3-yl)pyrazin-2-amine

**[0749]** A mixture of (E)-3-amino-6-bromo-N'-(ethanoyloxy)pyrazine-2-carboximidamide (8.405 g, 22.71 mmol) and AcOH (30 mL) was heated to  $70^\circ\text{C}$  for 2 hours. The reaction mixture was allowed to cool, diluted with water and then neutralised with solid  $\text{NaHCO}_3$ . The resultant dark coloured precipitate was filtered off (3.83 g, 66% Yield).  $^1\text{H}$  NMR (400.0 MHz, DMSO)  $\delta$  2.68 (s, 3H), 7.31 (br s, 2H) and 8.34 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 258.80

#### Method Y

##### Step 5: Tert-butyl N-[5-bromo-3-(5-methyl-1,2,4-oxadiazol-3-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

**[0750]** BOC anhydride (1.355 g, 1.426 mL, 6.210 mmol) and DMAP (25.29 mg, 0.2070 mmol) were added to a solution of 5-bromo-3-(5-methyl-1,2,4-oxadiazol-3-yl)pyrazin-2-amine (530 mg, 2.070 mmol) in DCM (7.918 mL) and the reaction mixture allowed to stir overnight at ambient temperature. The mixture was diluted with DCM and washed with water (1 $\times$ ), 1N HCl (1 $\times$ ), saturated  $\text{NaHCO}_3$  (1 $\times$ ), dried over  $\text{MgSO}_4$  and concentrated in vacuo. The resultant residue was purified on silica gel by flash column chromatography (0-30% EtOAc/Petrol) to afford the product as a beige solid (686 mg, 73% Yield).  $^1\text{H}$  NMR (400.0 MHz,  $\text{CDCl}_3$ )  $\delta$  1.37 (18H, s), 2.73 (3H, s), 8.74 (1H, s) ppm

#### Method Y

##### Step 6: Tert-butyl N-[3-(5-methyl-1,2,4-oxadiazol-3-yl)-5-piperazin-1-yl-pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

**[0751]** A mixture of tert-butyl N-[5-bromo-3-(5-methyl-1,2,4-oxadiazol-3-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (200 mg, 0.4383 mmol) in DMF (1.562 mL) was treated with piperazine (377.5 mg, 4.383 mmol) and the reaction mixture heated at  $100^\circ\text{C}$  for 30 minutes. The reaction was allowed to cool to RT and diluted with EtOAc/Water and

the layers separated. The organic layer was washed with water (3×), dried (MgSO<sub>4</sub>), filtered and concentrated in vacuo. The material was used crude in the next step (150 mg, 67% Yield). <sup>1</sup>H NMR (400.0 MHz, CDCl<sub>3</sub>) δ 1.37 (18H, s), 2.73 (3H, s), 3.14 (4H, m) ppm; MS (ES<sup>+</sup>) 462.14

#### Method Y



##### Step 7: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(5-methyl-1,2,4-oxadiazol-3-yl)piperazin-2-amine

**[0752]** Ethanesulfonyl chloride (41.38 mg, 30.49 μL, 0.3218 mmol) was added to a solution of tert-butyl N-tert-butoxycarbonyl-N-[3-(5-methyl-1,2,4-oxadiazol-3-yl)-5-piperazin-1-yl-pyrazin-2-yl]carbamate (150 mg, 0.2925 mmol) and triethylamine (59.20 mg, 81.54 μL, 0.5850 mmol) in dichloromethane (5.818 mL) at 0° C. and the resulting solution stirred at ambient temperature for 1.5 hours. The reaction was treated with HCl in dioxane (365.8 μL of 4 M, 1.463 mmol) and allowed to stir at ambient temperature overnight. The reaction was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound (19.8 mg, 12% Yield). <sup>1</sup>H NMR (DMSO) δ 1.23 (3H, t), 2.70 (3H, s), 3.08-3.13 (2H, q), 3.31 (4H, m), 3.48 (4H, m), 6.50 (2H, br s), 8.17 (1H, s) ppm; MS (ES<sup>+</sup>) 353.99

#### Example 26

##### 3-(benzo[b]thiophen-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine (Compound I-59)

**[0753]**



**[0754]** Compound I-59 was prepared using Method Z, Steps 1-3.

#### Method Z

##### Step 1: Tert-butyl N-[3-(benzothiophen-2-yl)-5-bromo-pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

**[0755]** A mixture of 3,5-dibromopyrazin-2-amine (200 mg, 0.7908 mmol), benzo[b]thiophen-2-ylboronic acid (140.8 mg, 0.7908 mmol), NaHCO<sub>3</sub> (199.3 mg, 2.372 mmol) and palladium; triphenylphosphane (45.69 mg, 0.03954 mmol) in dimethoxyethane (2.000 mL) and water (1.000 mL) was heated at 120° C. in the microwave for 10 minutes. The reaction was diluted with EtOAc/water and the layers separated. The aqueous layer was extracted further with EtOAc (2×), dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was taken up in DCM (5.0 mL), treated with BOC anhydride (517.7 mg, 544.9 μL, 2.372 mmol) and DMAP (9.661 mg, 0.07908 mmol) and allowed to stir overnight at ambient temperature. The reaction was concentrated in vacuo and the resultant residue was purified on silica gel by flash column chromatography (0-10% EtOAc/Petrol) to afford the product as a bright yellow solid (302 mg, 75% Yield). <sup>1</sup>H NMR (DMSO) δ 1.33 (18H, s), 7.42 (2H, m), 7.89 (2H, m), 8.06 (1H, s), 8.49 (1H, s) ppm; MS (ES<sup>+</sup>) 507.79

## Method Z

Step 2: Tert-butyl N-[3-(benzothiophen-2-yl)-5-piperazin-1-yl-pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

[0756] Tert-butyl N-[3-(benzothiophen-2-yl)-5-bromopyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (100 mg, 0.1975 mmol) was dissolved in DMF (781.2  $\mu$ L) and treated with piperazine (170.1 mg, 1.975 mmol) and the reaction mixture heated at 100° C. for 30 minutes. The reaction was allowed to cool to ambient temperature and diluted with EtOAc/water and the layers separated. The organic layer was washed with water (3 $\times$ ), dried (MgSO<sub>4</sub>) and concentrated in vacuo. The material was used crude in the next step (92 mg, 91% Yield). <sup>1</sup>H NMR (DMSO)  $\delta$  1.23 (18H, s), 2.98 (4H, m), 3.59 (2H, m), 3.63 (2H, m), 7.29 (2H, m), 7.74 (3H, m), 7.89 (1H, s) ppm; MS (ES<sup>+</sup>) 511.98

## Method Z



Step 3: 3-(benzo[b]thiophen-2-yl)-5-(4-(ethylsulfonyl)piperazin-1-yl)pyrazin-2-amine

[0757] Ethanesulfonyl chloride (25.43 mg, 18.74  $\mu$ L, 0.1978 mmol) was added to a solution of tert-butyl N-[3-(benzothiophen-2-yl)-5-piperazin-1-yl-pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (92 mg, 0.1798 mmol) and triethylamine (36.39 mg, 50.12  $\mu$ L, 0.3596 mmol) in dichloromethane (3.965 mL) at 0° C. and the resulting solution stirred at room temperature for 1.5 hours. The reaction was then treated with HCl in dioxane (224.8  $\mu$ L of 4 M, 0.8990 mmol) and allowed to stir at ambient temperature overnight. The mixture was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound as a yellow solid (14.4 mg, 20% Yield). <sup>1</sup>H NMR (DMSO)  $\delta$  1.24 (3H, t), 2.51 (2H, q), 3.30 masked signal, 3.52 (4H, m), 5.91 (2H, br s), 7.36-7.40 (2H, m), 7.84-7.86 (1H, m), 7.92 (1H, s), 7.95 (1H, m), 8.06 (1H, s) ppm; MS (ES<sup>+</sup>) 404.02

## Example 27

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(1H-indol-2-yl)pyrazin-2-amine (Compound I-61)

[0758]



[0759] Compound I-61 was prepared using Method AA, Steps 1-3.

## Method AA

Step 1: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-bromopyrazin-2-yl)-1H-indole-1-carboxylate

[0760] A mixture of 3,5-dibromopyrazin-2-amine (200 mg, 0.7908 mmol), tert-butyl 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indole-1-carboxylate (271.4 mg, 0.7908 mmol), NaHCO<sub>3</sub> (199.3 mg, 2.372 mmol) and palladium; triphenylphosphane (45.69 mg, 0.03954 mmol) in

dimethoxyethane (2.000 mL) and water (1.000 mL) was heated at 120° C. in the microwave for 10 minutes. The reaction was diluted with EtOAc/water and the layers separated. The aqueous layer was extracted further with EtOAc (2×), dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was taken up in DCM (5 mL), treated with BOC anhydride (517.7 mg, 544.9 μL, 2.372 mmol) and DMAP (9.661 mg, 0.07908 mmol) and allowed to stir overnight at ambient temperature. The reaction was concentrated in vacuo and the resultant residue was purified on silica gel by flash column chromatography (0-10% EtOAc/Petrol) to afford the product as a bright yellow solid (254 mg, 55% Yield). <sup>1</sup>H NMR (DMSO) δ 1.33 (18H, s), 1.53 (9H, s), 6.90 (1H, s), 7.26 (1H, m), 7.38 (1H, m), 7.58 (1H, m), 8.21 (1H, m) 8.56 (1H, s); MS (ES<sup>+</sup>) 590.89

#### Method AA

Step 2: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-(piperazin-1-yl)pyrazin-2-yl)-1H-indole-1-carboxylate

**[0761]** A mixture of tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]indole-1-carboxylate (100 mg, 0.1696 mmol) and piperazine (146.1 mg, 1.696 mmol) in DMF (2 mL) was heated at 100° C. for 30 minutes and then allowed to cool to ambient temperature. The reaction was diluted with EtOAc/water and the organic layer washed further with water (2×), dried (MgSO<sub>4</sub>) and concentrated in vacuo. The material was used crude in the next step (quantitative yield assumed). MS (ES<sup>+</sup>) 595.09

#### Method AA

Step 3: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(1H-indol-2-yl)pyrazin-2-amine


**[0762]** Ethanesulfonyl chloride (43.25 mg, 31.87 μL, 0.3364 mmol) was added to a solution of tert-butyl 2-[3-(bis(tert-butoxycarbonyl)amino)-6-piperazin-1-yl-pyrazin-2-yl]indole-1-carboxylate (100 mg, 0.1682 mmol) and triethylamine (18.72 mg, 25.79 μL, 0.1850 mmol) in DCM (2 mL) at 0° C. and the reaction mixture allowed to stir at ambient temperature for 30 minutes. The reaction was then treated with HCl in dioxane (210.2 μL of 4 M, 0.8410 mmol) and allowed to stir at ambient temperature overnight. The mixture was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound as a yellow solid (5.5 mg, 8% Yield). <sup>1</sup>H NMR (DMSO) δ 1.25 (3H, t), 3.12 (2H, q), 3.30 masked signal, 3.59 (4H, m), 5.66 (2H, br s), 7.00-7.04 (1H, m), 7.14-7.18 (2H, m), 7.51 (1H, m), 7.58 (1H, m), 7.81 (1H, s), 11.25 (1H, br s) ppm; MS (ES<sup>+</sup>) 386.87

#### Example 28

5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(3-phenylisoxazol-5-yl)pyrazin-2-amine (Compound I-162)

**[0763]**

SCHEME XXVIII



Compound I-162

**[0764]** Compound I-162 was prepared using Method U, Steps 1-5 followed by Method AB, Step 1.

#### Method AB

Step 1: 5-(4-(ethylsulfonyl)piperazin-1-yl)-3-(3-phenylisoxazol-5-yl)pyrazin-2-amine


**[0765]** Triethylamine (10.21 mg, 14.06 μL, 0.1009 mmol) was added to a solution of tert-butyl N-tert-butoxycarbonyl-N-[5-(4-ethylsulfonylpiperazin-1-yl)-3-ethynyl-pyrazin-2-yl]carbamate (50 mg, 0.1009 mmol) and N-hydroxybenzimi-

doyle chloride (15.70 mg, 0.1009 mmol) in THF (4 mL) and the reaction mixture allowed to stir at ambient temperature for 1 hour. After this time the reaction mixture was heated under reflux for 3 hours and allowed to cool to RT. The reaction was treated with HCl in dioxane (126.1  $\mu$ L of 4 M, 0.5045 mmol), allowed to stir overnight at ambient temperature, concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected and freeze-dried to give the title compound as an orange solid (9.9 mg, 18% Yield). <sup>1</sup>H NMR (DMSO)  $\delta$  1.03 (3H, t), 2.90 (2H, q), 3.11 (4H, m), 3.33 (4H, m), 5.88 (2H, br s), 7.34-7.36 (4H, m), 7.77 (2H, m), 7.92 (1H, s) ppm; MS (ES<sup>+</sup>) 414.98


## Example 29

3-(1H-benzo[d]imidazol-2-yl)-5-(1-phenyl-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (Compound I-159)

[0766]



-continued



[0767] Compound I-159 was prepared using Method A, Steps 1-3 followed by Method AC, Steps 1-3.

## Method AC

Step 1: tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-bromopyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate

[0768] 3-(1H-benzimidazol-2-yl)-5-bromo-pyrazin-2-amine (8.0443 g, 27.73 mmol) was dissolved in acetonitrile (100.6 mL) and THF (100.6 mL) and treated with DMAP (338.8 mg, 2.773 mmol) followed by portionwise addition of BOC anhydride (27.24 g, 28.67 mL, 124.8 mmol). The reaction was allowed to stir at ambient temperature overnight. The mixture was concentrated in vacuo, diluted with DCM and washed with NaHCO<sub>3</sub> (3 $\times$ ). The aqueous layer was further extracted with DCM (2 $\times$ ) and the combined organics washed with brine, dried (MgSO<sub>4</sub>), filtered and concentrated in vacuo. The resultant residue was purified by column chromatography (ISCO Companion<sup>TM</sup>, 330 g column, 0-30%

EtOAc/Petrol) to afford the title compound which was freeze-dried to give a yellow solid (13.05 g, 80% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 1.26 (s, 18H), 1.45 (s, 9H), 7.45-7.56 (m, 2H), 7.79-7.81 (m, 1H), 8.02-8.04 (m, 1H) and 9.08 (s, 1H) ppm; MS (ES<sup>+</sup>) 592.09

#### Method AC

Step 2: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-(1-phenyl-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate

[0769] A mixture of tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]benzimidazole-1-carboxylate (100 mg, 0.1694 mmol), 1-phenyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine (58 mg, 0.2034 mmol), 4-di-tert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (2 mg, 0.002825 mmol) and K<sub>2</sub>CO<sub>3</sub> (28 mg, 0.2026 mmol) in toluene (1000 μL) and water (120 μL) were heated to 100° C. overnight. The reaction was diluted with DCM, washed with water followed by 1M NaOH, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The material was used crude in the next step (quantitative yield assumed). MS (ES<sup>+</sup>) 669.02

#### Method AC


Step 3: 3-(1H-benzo[d]imidazol-2-yl)-5-(1-phenyl-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0770] A solution of tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-(1-phenyl-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate (100 mg, 0.1495 mmol) in DCM (1 mL) was treated with TFA (1 mL, 12.98 mmol) and allowed to stir at ambient temperature for 40 minutes. The reaction was concentrated in vacuo and the resultant residue was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 μM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound as a pale brown solid (8.0 mg, 14% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 2.84 (2H, m), 3.53 (2H, m), 3.94 (2H, m), 6.76-6.90 (3H, m), 7.20-7.69 (6H, m), 7.70-9.39 (4H, m), 12.88 (1H, m) ppm; MS (ES<sup>+</sup>) 369.02

#### Example 30

3-(1H-benzo[d]imidazol-2-yl)-5-(1-(thiazol-2-yl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (Compound I-164)


[0771]




#### SCHEME XXX

Method A  
Steps 1-3  
Method AC  
Step 1


-continued



Method AD  
Step 1




Method AD  
Step 2



Method AD  
Step 3

-continued



Compound I-164

[0772] Compound I-164 was prepared using Method A, Steps 1-3 followed by Method AC, Step 1 followed by Method AD, Steps 1-3.

## Method AD

Step 1: Tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-(1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate

[0773] A mixture of tert-butyl 2-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]benzimidazole-1-carboxylate (1.7068 g, 1.734 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (643.5 mg, 2.081 mmol), 4-ditert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (12.28 mg, 0.01734 mmol) and  $K_2CO_3$  (479.3 mg, 3.468 mmol) in toluene (9.216 mL) and water (1.024 mL) was heated to 100° C. for 16 hours. The reaction mixture was allowed to cool to RT and partitioned between DCM and water. The organic layer was washed with 1M NaOH solution, dried ( $Na_2SO_4$ ), filtered and concentrated in vacuo to give a brown oil. This was purified by column chromatography (ISCO Companion™, 40 g column, 0-20% EtOAc/Petrol) to afford the title compound as a yellow oil (1.028 g, 60% Yield). MS ( $ES^+$ ) 693.38

## Method AD

Step 2: 3-(1H-benzo[d]imidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

[0774] A solution of tert-butyl 2-(3-(bis(tert-butoxycarbonyl)amino)-6-(1-(tert-butoxycarbonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-yl)-1H-benzo[d]imidazole-1-carboxylate (1.7248 g, 2.490 mmol) in DCM (17.25 mL) was treated with TFA (5 mL, 64.90 mmol) and left to stir at ambient temperature for 4 hours. The reaction mixture was concentrated in vacuo and triturated from MeOH/Toluene and the resultant orange solid was filtered off (622 mg, 48% Yield). 1H NMR (400.0 MHz, DMSO)  $\delta$  2.90 (2H, m), 3.41

(2H, m), 3.96 (2H, m), 6.80 (1H, m), 7.09-7.30 (4H, m), 7.82 (2H, br s), 8.52 (1H, s) ppm; MS ( $ES^+$ ) 293.03

## Method AD

Step 3: 3-(1H-benzo[d]imidazol-2-yl)-5-(1-(thiazol-2-yl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine


[0775] A solution of 3-(1H-benzo[d]imidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (100 mg, 0.1922 mmol) in DMSO (2.000 mL) was treated with triethylamine (58.35 mg, 80.37  $\mu$ L, 0.5766 mmol) and 2-chlorothiazole (27.57 mg, 0.2306 mmol) and heated in the microwave for 40 minutes at 160° C. The reaction mixture was filtered and purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $CH_3CN$ ) over 16 minutes at 25 mL/min]. The fractions were collected, passed through a sodium bicarbonate cartridge and freeze-dried to give the title compound (20.6 mg, 15% Yield). 1H NMR (400.0 MHz, DMSO)  $\delta$  2.89 (s, 2H), 3.80 (t,  $J=5.7$  Hz, 2H), 4.21 (d,  $J=2.5$  Hz, 2H), 6.86 (s, 1H), 6.94-6.96 (m, 1H), 7.28-7.31 (m, 3H), 7.69 (s, 2H) and 8.41 (s, 1H) ppm; MS ( $ES^+$ ) 376.0

## Example 31


5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine (Compound I-157)

[0776]

## SCHEME XXXI



-continued



Compound I-157

[0777] Compound I-157 was prepared using Method A, Step 1 followed by Method Q, Steps 1-3 followed by Method AE, Step 1.

## Method AE

Step 1: 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(5-methyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine

[0778] A mixture of 3-amino-6-(1-ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylic acid (100 mg, 0.320 mmol), 4-methylbenzene-1,2-diamine (39.12 mg, 0.320 mmol), diethoxyphosphorylformonitrile (47.83 uL, 0.320 mmol) and triethylamine (44.63 uL, 0.320 mmol) in DME (1 mL) were heated in a microwave at 170° C. for 90 min. After this time the reaction mixture was filtered and purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the title compound (46.8 mg, 21% Yield). <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 8.34 (s, 1H), 7.57 (d, J=8.1 Hz, 1H), 7.46 (s, 1H), 7.11 (d, J=8.0 Hz, 1H), 6.78 (s, 1H), 4.00 (d, J=2.8 Hz, 2H), 3.50 (t, J=5.7 Hz, 2H), 3.16 (q, J=7.4 Hz, 2H), 2.78 (s, 2H), 2.46 (s, 3H) and 1.25 (t, J=7.4 Hz, 3H) ppm; MS (ES<sup>+</sup>) 399

[0779] The following compounds were all prepared using the same method or a method similar to the one described for Compound I-157 in Example 31 above.

[0780] Compound I-152 3-(6,7-dimethyl-1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine <sup>1</sup>H NMR (400.0 MHz, DMSO) δ 12.68 (s, 1H), 8.33 (s, 1H), 7.32 (d, J=7.1 Hz, 1H), 7.11 (d, J=8.2 Hz, 1H), 6.79 (s, 1H), 4.00 (d, J=2.7 Hz, 2H), 3.50 (t, J=5.7 Hz, 2H), 3.16 (q, J=7.4 Hz, 2H), 2.79 (s, 2H), 2.54 (s, 3H), 2.36 (s, 3H) and 1.25 (t, J=7.3 Hz, 3H) ppm; MS (ES<sup>+</sup>) 413

[0781] Compound I-153 3-(6,7-difluoro-1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 421

[0782] Compound I-154 3-(5,6-dimethyl-1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 413

[0783] Compound I-155 5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(4-fluoro-1H-benzo[d]imidazol-2-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 403

[0784] Compound I-156 3-(5-tert-butyl-1H-benzo[d]imidazol-2-yl)-5-(1-(ethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine; MS (ES<sup>+</sup>) 441

For compounds I-191 and I-192 below, please see Tables 7 and 8 for analytical data.

[0785] Compound I-191 4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl(pyridin-3-yl)methanone


[0786] Compound I-192 1-(4-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-2-methylpropan-1-one


## Example 32

1-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridine-4-carboxylic acid (Compound I-288)

[0787]

## SCHEME XXXII





**[0788]** Compound I-288 was prepared using Method A, Steps 1-2, followed by Method N Steps 1-2 followed by Method O Step 1 and Method AF Step 1.

#### Method AF

Step 1: 1-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridine-4-carboxylic acid

**[0789]** A mixture of 1,2,3,6-tetrahydropyridine-4-carboxylic acid (36.18 mg, 0.2846 mmol) and tert-butyl N-tert-butoxycarbonyl-5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (80 mg, 0.1423 mmol) and Et<sub>3</sub>N (43.

20 mg, 59.50  $\mu$ L, 0.4269 mmol) in DMF (800.3  $\mu$ L) were heated at 100° C. in the microwave for 20 mins. The reaction mixture was diluted with ethylacetate, washed with water and concentrated to an oil. The oil was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and TFA (162.3 mg, 109.7  $\mu$ L, 1.423 mmol) added and the resulting solution stirred at room temperature for 1 hr. The reaction mixture was concentrated in vacuo to leave an oil which was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the product (15 mg, 20% yield) <sup>1</sup>H NMR (400 MHz, MeOD)  $\delta$  1.70-1.75 (m, 2H), 2.85 (t, 2H), 3.35-3.4 (m, 2H), 6.25 (br s, 1H), 6.65 (d, 1H), 6.7-6.8 (m, 4H) and 7.25-7.4 (m, 4H); MS (ES<sup>+</sup>) 365.1

**[0790]** The following compounds were all prepared using the method described for Compound I-288 in Example 32 above.


**[0791]** Compound I-290 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(4-thiazol-2-yl-3,6-dihydro-2H-pyridin-1-yl)pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, MeOD)  $\delta$  1.95-2.0 (m, 2H), 3.0 (t, 2H), 3.35-3.4 (m, 2H), 5.95-6.0 (m, 1H), 6.65 (d, 1H), 6.7-6.8 (m, 4H) and 7.25-7.35 (m, 4H) ppm; MS (ES<sup>+</sup>) 404.18


#### Example 33

N-(1-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)azetidin-3-yl)ethanamide (Compound I-282)

**[0792]**

#### SCHEME XXXIII





**[0793]** Compound I-282 was prepared using Method A, Steps 1-2, followed by Method N Steps 1-2 followed by Method O Step 1 and Method AH Steps 1-2.

#### Method AG


Step 1: 5-(3-aminoazetidin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0794]** A mixture of tert-butyl N-tert butoxycarbonyl-5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (200 mg, 0.3858 mmol) and tert-butyl N-(azetidin-3-yl)carbamate (66.44 mg, 0.3858 mmol) and DIPEA (99.72 mg, 134.4  $\mu$ L, 0.7716 mmol) in DMF (5 mL) was heated at 90° C. After this time, the reaction mixture was cooled to room temperature and diluted with ethylacetate (5 mL), washed with water (5 mL) and concentrated in vacuo to leave a solid. The solid was dissolved in dichloromethane (10 mL) and trifluoroacetic acid (1.319 g, 891.2  $\mu$ L, 11.57 mmol) was added and the resulting solution stirred at room temperature for 1 hr. The reaction mixture was concentrated in vacuo to leave an oil. The oil was redissolved in dichloromethane and concentrated in vacuo to leave the product, which was used crude in the next step.

#### Method AG

Step 2: N-[1-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]azetidin-3-yl]acetamide

**[0795]** Acetyl chloride (12.69 mg, 11.49  $\mu$ L, 0.1616 mmol) was added to a solution of 5-(3-aminoazetidin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (50 mg, 0.1616 mmol) and DIPEA (27.15 mg, 36.59  $\mu$ L, 0.2101 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and the resulting solution stirred at room temperature for 20 mins. After this time, the reaction mixture was washed with saturated aqueous  $\text{NaHCO}_3$  solution (1  $\times$  5 mL), dried over  $\text{MgSO}_4$  and concentrated in vacuo to leave a solid. The product was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the product (12.5 mg, 21% yield);  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  1.98 (s, 3H), 3.4 (q, 1H), 3.7-3.76 (m, 2H), 4.4-4.45 (m, 2H), 4.75-4.85 (m, 1H), 6.2 (s, 2H), 7.5-7.6 (m, 3H), 7.7 (s, 1H) and 8.2-8.3 (m, 2H) ppm; MS ( $\text{ES}^+$ ) 352.1



**[0796]** The following compounds were all prepared using a similar method to the one described for Compound I-282 in Example 34 above


**[0797]** Compound I-302 N-[1-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]azetidin-3-yl]ethanesulfonamide  $^1\text{H}$  NMR (400 MHz, DMSO)  $\delta$  1.3 (t, 3H), 3.1 (q, 2H), 3.8-3.85 (m, 2H), 4.35-4.42 (m, 3H), 7.0 (m, 2H), 7.7-7.8 (m, 3H), 7.8-7.85 (m, 2H) and 8.05-8.1 (m, 2H); MS ( $\text{ES}^+$ ) 402.2


#### Example 34

5-(2-ethylsulfonyl-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-5-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-305)

**[0798]**

#### SCHEME XXXIV





**[0799]** Compound I-305 was prepared using Method A, Steps 1-2, followed by Method N Steps 1-2 followed by Method O Step 1 and followed by Method AI Step 1.

#### Method AH

Step 1: 5-(2-ethylsulfonyl-1,3,3a,4,6,6a-hexahydro-pyrrolo[3,4-c]pyrrol-5-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0800]** 2-ethylsulfonyl-3,3a,4,5,6,6a-hexahydro-1H-pyrrolo[3,4-c]pyrrole (36.34 mg, 0.1779 mmol) and Et<sub>3</sub>N (54.01


mg, 74.39  $\mu$ L, 0.5337 mmol) were added to a solution of tert-butyl N-tert-butoxycarbonyl-5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (100 mg, 0.1779 mmol) in DMF (2 mL) and the resulting solution heated at 100° C. for 2 h. The reaction mixture was cooled to room temperature and diluted with ethyl acetate (5 mL) and washed successively with water (2 $\times$ 5 mL) followed by brine (1 $\times$ 5 mL). The organic layer was concentrated in vacuo to leave a solid. The solid was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and trifluoroacetic acid (405.7 mg, 274.1  $\mu$ L, 3.558 mmol) was added. The reaction mixture was stirred at room temperature for 1 h and the reaction mixture then concentrated in vacuo to an oil. The oil was redissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and concentrated in vacuo to leave a solid. The solid was purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ M, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the product (34.4 mg, 40% yield); <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.25 (t, 3H), 3.1-3.2 (m, 4H), 3.25-3.3 (m, 2H), 3.4-3.45 (m, 2H), 3.5-3.7 (m, 4H), 6.75 (s, 2H), 7.6-7.65 (m, 3H) and 8.05-8.1 (m, 2H) ppm; MS (ES<sup>+</sup>) 442.1


#### Example 35

5-[4-(2-methylsulfonyl-ethyl)piperazin-1-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-296)

**[0801]**

#### SCHEME XXXV





**[0802]** Compound I-296 was prepared using Method A, Steps 1-2, followed by Method N Steps 1-2 followed by Method O Step 1 and followed by Method AJ Steps 1-2.

#### Method AI

Step 1: tert-butyl N-tert-butoxycarbonyl-5-(4-(2-(methylsulfonyl)ethyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate

**[0803]** tert-butyl N-tert-butoxycarbonyl-5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (75 mg, 0.13 mmol) was dissolved in DMF (1 mL) and 1-(2-(methylsulfonyl)ethyl)piperazine (77 mg, 0.40 mmol) was added. The reaction mixture was heated at 100° C. in a sealed tube overnight. The reaction mixture was cooled to room temperature and partitioned between CH<sub>2</sub>Cl<sub>2</sub> and water and the layers separated using a phase separation cartridge. The organic layer was concentrated in vacuo to give the product (84.01 mg, 0.13 mmol)

#### Method AI

Step 2: 5-[4-(2-methylsulfonyl)ethyl]piperazin-1-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0804]** tert-butyl N-tert-butoxycarbonyl-5-(4-(2-(methylsulfonyl)ethyl)piperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-ylcarbamate (84.01 mg, 0.13 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (10 mL) and trifluoroacetic acid (2 mL) added and the reaction mixture stirred at room temperature overnight. The reaction mixture was concentrated in vacuo and purified by reverse phase preparative HPLC [Waters Sunfire C18, 10 uM, 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The product was passed through a bicarbonate cartridge and the cartridge washed with methanol and the filtrate concentrated in vacuo to leave a solid. The solid was taken up in acetonitrile and water and freeze dried to give the product (20.6 mg, 34% yield); <sup>1</sup>H NMR (400 MHz, DMSO) δ 2.60 (m, 5H), 2.77 (m, 2H), 3.06 (s, 3H), 3.31 (m, 2H), 3.36 (masked signal, 3.56 (m, 4H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H) and 8.25 (s, 1H) ppm; MS (ES<sup>+</sup>) 430.2

**[0805]** The following compounds were all prepared using the method described for Compound I-296 in Example 36 above

**[0806]** Compound I-298 3-[4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]piperazin-1-yl]propanenitrile <sup>1</sup>H NMR (400 MHz, DMSO) δ 2.60-2.66 (m, 6H), 2.75 (m, 2H), 3.46 (m, 4H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H) and 8.24 (s, 1H) ppm; MS (ES<sup>+</sup>) 377.2

**[0807]** Compound I-300 5-(4-benzylpiperazin-1-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine <sup>1</sup>H NMR (400 MHz, DMSO) δ 3.31 (masked signal, 3.46 (m, 4H) 3.55 (m, 2H), 6.88 (br s, 2H), 7.27 (m, 1H), 7.36 (m, 4H), 7.64-7.68 (m, 3H), 8.11 (m, 2H) and 8.23 (s, 1H) ppm; MS (ES<sup>+</sup>) 414.2

**[0808]** Compound I-303 2-[4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]piperazin-1-yl]-1-morpholino-ethanone <sup>1</sup>H NMR (400 MHz, DMSO) δ 2.59 (m, 4H), 3.24 (s, 2H), 3.46 (m, 6H) 3.56 (m, 6H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H) and 8.23 (s, 1H) ppm; MS (ES<sup>+</sup>) 451.3


**[0809]** Compound I-306 2-[4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]piperazin-1-yl]ethanol <sup>1</sup>H NMR (400 MHz, DMSO) δ 2.46 (m, 2H), 2.59 (m, 4H), 3.46 (m, 4H), 3.56 (m, 2H), 4.47 (t, 1H), 6.90 (br s, 2H), 7.64-7.69 (m, 3H), 8.11 (m, 2H) and 8.24 (s, 1H) ppm


#### Example 36

1-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]pyridin-4-one (Compound I-286)

**[0810]**

#### SCHEME XXXVI





**[0811]** Compound I-286 was prepared using Method A, Steps 1-2, Method N Steps 1-2 followed by Method AK step 1.

#### Method AJ

Step 1: 1-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]pyridin-4-one



**[0812]** To a solution of 5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (150.0 mg, 0.4715 mmol) in DMF (3.000 mL) was added copper (5.992 mg, 0.09430 mmol) and cesium carbonate (307.2 mg, 0.9430 mmol) followed by pyridin-4-ol (448.4 mg, 4.715 mmol) and the reaction mixture heated at 160° C. in the microwave for 1 h. The reaction mixture was filtered and washed with DMF (2 mL). Ethyl acetate and water were added to the filtrate and the layers separated. The aqueous layer was extracted further with ethyl acetate (3×5 mL) and the combined organic extracts were washed with water (3×5 mL). The organic extracts were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated in vacuo. The material was triturated with a mixture of methanol and ether to leave the product. (42.6 mg, 25% yield); <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 6.33 (d, 2H), 7.69 (s, 2H), 8.15-8.18 (m, 2H), 8.35 (d, 2H) and 8.78 (s, 1H) ppm; MS (ES<sup>+</sup>) 333.11

#### Example 37

2-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]isoindolin-1-one (Compound I-304)

**[0813]**

#### SCHEME XXXVII



**[0814]** Compound I-304 was prepared using Method A, Steps 1-3, followed by Method AL Steps 1-2.

#### Method AK

Step 1: tert-butyl N-tert-butoxycarbonyl-5-bromo-3-(1-tosyl-1H-benzo[d]imidazol-2-yl)pyrazin-2-ylcarbamate

**[0815]** Sodium hydride (496.4 mg, 551.6 μL, 12.41 mmol) was added to a solution of 3-(1H-benzimidazol-2-yl)-5-bromo-pyrazin-2-amine (3 g, 10.34 mmol) THF (20 mL) at 0° C. The resulting suspension was stirred for 5 mins and 4-methylbenzenesulfonylchloride (2.168 g, 11.37 mmol) was added. The reaction was stirred for a further 30 mins and then quenched with saturated NH<sub>4</sub>Cl aqueous solution and extracted with ethylacetate (3×50 mL). The organic layer was washed with water (1×20 mL) and concentrated in vacuo to a solid. The solid was dissolved in THF (15 mL) and di-tert-butyl dicarbonate (5.190 g, 5.463 mL, 23.78 mmol) and DMAP (126.3 mg, 1.034 mmol) were added. The mixture was stirred overnight at room temperature and then concentrated in vacuo to leave an oil. The oil was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and the product precipitated by slow addition of ether and isolated by filtration (3.1 g, 47% yield);

[0816]  $^1\text{H}$  NMR (400 MHz, DMSO)  $\delta$  1.75 (m, 18H), 2.15 (s, 3H), 7.70-7.80 (m, 4H), 7.90 (m, 1H), 8.00 (m, 1H), 8.20 (m, 2H) and 8.60 (s, 1H); MS ( $\text{ES}^+$ ) 646.20

#### Method AK

Step 2: 2-[5-amino-6-(1H-benzimidazol-2-yl)pyrazin-2-yl]isoindolin-1-one

[0817]  $\text{C}_{28}\text{H}_{30}\text{BrN}_5\text{O}_6\text{S}$  (100 mg, 0.1552 mmol), isoindolin-1-one (31.00 mg, 0.2328 mmol), (5-diphenylphosphanyl-9,9-dimethyl-xanthen-4-yl)-diphenyl-phosphane (13.47 mg, 0.02328 mmol), 1,5-diphenylpenta-1,4-dien-3-one palladium (7.106 mg, 0.007760 mmol) and cesium carbonate (101.1 mg, 0.3104 mmol) were heated in dioxane (3 mL) at 70° C. for 1 hr. The reaction mixture was cooled to room temperature and diluted with ethyl acetate (10 mL) and washed with water (1×10 mL) and brine (1×10 mL), dried over  $\text{MgSO}_4$  and concentrated in vacuo. The residue was


dissolved in  $\text{CH}_2\text{Cl}_2$  (10 mL) and trifluoroacetic acid (353.9 mg, 239.1  $\mu\text{L}$ , 3.104 mmol) added and the reaction mixture stirred at room temperature for 1 h. The reaction mixture was concentrated in vacuo and the residue purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu\text{M}$ , 100 Å column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B:  $\text{CH}_3\text{CN}$ ) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried to give the product (7.5 mg, 13.4% yield);  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  5.25 (s, 2H), 7.3-7.35 (m, 2H), 7.55-7.6 (m, 1H), 7.63-7.7 (m, 4H), 7.9 (d, 1H) and 9.22 (s, 1H) ppm; MS ( $\text{ES}^+$ ) 343.1

#### Example 38

5-(1-ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)-3-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-amine  
(Compound I-307)

[0818]

#### SCHEME XXXVIII



[0819] Compound I-307 was prepared using Method Y, Steps 1-4 followed by Method AM steps 1-3.

#### Method AL

Step 1: tert-butyl 4-(5-amino-6-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-yl)pyrazin-2-yl)-5,6-dihydropyridine-1(2H)-carboxylate

[0820] 5-bromo-3-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-amine (150 mg, 0.4715 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (189.5 mg, 0.6130 mmol) and sodium carbonate (707.0  $\mu$ L of 2 M, 1.414 mmol) in DMF (1.500 mL) were heated to 120° C. for 45 min in the microwave. After this time the reaction mixture was diluted with water (5 mL) and ethyl acetate (5 mL) and the layers separated. Organic layer washed with water (3 $\times$ 5 mL), dried (MgSO<sub>4</sub>) and concentrated in vacuo to leave the title compound as a yellow/orange solid (175.6 mg, 88% yield). <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.44 (s, 9H), 2.60 (m, 2H), 3.57 (m, 2H), 4.06 (m, 2H), 6.70 (s, 1H), 7.30 (br s, 2H), 7.69 (m, 2H), 7.77 (m, 1H), 8.25 (m, 2H) and 8.51 (1H, s) ppm; MS (ES<sup>+</sup>) 421.02

#### Method AL

Step 2: 3-(5-phenyl-1,2,4-oxadiazol-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine


[0821] A solution of HCl in dioxane (505.5  $\mu$ L of 4 M, 2.022 mmol) was added to a solution of tert-butyl 4-[5-amino-6-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate (170 mg, 0.4043 mmol) in methanol (5 mL) at rt and the reaction mixture stirred at room temperature for 48 h. The reaction mixture was concentrated in vacuo to give the title compound (129 mg, 99% yield); MS (ES<sup>+</sup>) 321.03

#### Method AL

Step 3: 5-(1-ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)-3-(5-phenyl-1,2,4-oxadiazol-3-yl)pyrazin-2-amine


[0822] Ethanesulfonyl chloride (62.13 mg, 45.78  $\mu$ L, 0.4832 mmol) and triethyl amine (122.2 mg, 168.3  $\mu$ L, 1.208 mmol) were added to a solution of 3-(5-phenyl-1,2,4-oxadiazol-3-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (129 mg, 0.4027 mmol) in dichloromethane (5 mL) at 0° C. and the resulting solution stirred at room temperature for 2 h. The reaction mixture was concentrated in vacuo and purified by reverse phase preparative HPLC [Waters Sunfire C18, 10  $\mu$ m, 100 A column, gradient 10%-95% B (solvent A: 0.05% TFA in water, solvent B: CH<sub>3</sub>CN) over 16 minutes at 25 mL/min]. The fractions were collected, and freeze-dried. The solid was taken up in dichloromethane (5 mL) and washed with saturated aqueous NaHCO<sub>3</sub> solution (2 $\times$ 5 mL), dried over MgSO<sub>4</sub> and concentrated in vacuo to leave the product (65 mg, 40% yield) <sup>1</sup>H NMR (400 MHz, DMSO)  $\delta$  1.24 (t, 3H), 2.68 (m, 2H), 3.13 (q, 2H), 3.48 (m, 2H), 3.99 (m, 2H), 6.63 (m, 1H), 7.29 (br s, 2H), 7.68 (m, 2H), 7.77 (m, 1H), 8.26 (m, 2H) and 8.52 (1H, s) ppm; MS (ES<sup>+</sup>) 412.97

III



[0823] Illustrated below are exemplary methods for the preparation of compounds of formula III.

[0824] Procedure for the Preparation of Bromo Intermediate:



#### Method BF

Step 1: 3-Amino-6-bromopyrazine-2-carbohydrazide

[0825] To a suspension of methyl 3-amino-6-bromo-pyrazine-2-carboxylate (2.5 g, 10.8 mmol) in EtOH (50 mL) was added hydrazine hydrate (3.2 g, 3.1 mL, 64.6 mmol) and the reaction mixture was heated at 70° C. for 1.5 h forming a thick yellow solid. The solid was isolated by filtration, washed with water (20 mL) and ethanol (40 mL). The solid was dried under

high vacuum to yield 3-amino-6-bromo-pyrazine-2-carbohydrazide (2.35 g, 94%) as a light yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.78 (s, 1H), 8.31 (s, 1H), 7.62 (s, 2H), 4.53 (d,  $J=3.5$  Hz, 2H). LC/MS  $m/z$  233.1  $[\text{M}+\text{H}]^+$ .

Step 2: 5-Bromo-3-(5-(4-(bromomethyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0826]** To a suspension of 3-amino-6-bromo-pyrazine-2-carbohydrazide (200 mg, 0.86 mmol) and 4-(bromomethyl) benzoic acid (185 mg, 0.86 mmol) in MeCN (4 mL) at room temperature was added dibromo(triphenyl)phosphorane (1.7 g, 4.14 mmol) and the resulting reaction mixture was stirred for 1 h. The reaction mixture was diluted with acetonitrile (2 mL) and DIPEA (668 mg, 900  $\mu\text{L}$ , 5.2 mmol) was added dropwise and the suspension was allowed to stir overnight. The suspension was filtered, washed with  $\text{CH}_3\text{CN}$  and hexane, and air dried to provide 5-bromo-3-(5-(4-(bromomethyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (240 mg, 68%) as a yellow solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  8.45 (s, 1H), 8.11 (d,  $J=8.1$  Hz, 2H), 7.72 (d,  $J=8.2$  Hz, 2H), 4.82 (s, 2H). LC/MS  $m/z$  412.1  $[\text{M}+\text{H}]^+$ .

Step 3: tert-Butyl N-[5-bromo-3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

**[0827]** To a solution of 5-bromo-3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine (240 mg, 0.58 mmol) in THF (10 mL) was added DIPEA (302 mg, 407  $\mu\text{L}$ , 2.3 mmol) followed by the addition of di-tert-butyl dicarbonate (637 mg, 671  $\mu\text{L}$ , 2.9 mmol) and 4-dimethylaminopyridine (7 mg, 0.06 mmol). The reaction mixture was stirred at  $45^\circ\text{C}$ . overnight. The reaction mixture was concentrated in vacuo and purified by silica gel column chromatography using 10-70% ethyl acetate in hexane to yield tert-butyl N-[5-bromo-3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (120 mg, 34%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.16 (s, 1H), 8.12 (d,  $J=8.2$  Hz, 2H), 7.74 (d,  $J=8.2$  Hz, 2H), 4.83 (s, 2H), 1.29 (s, 18H).


Procedures for the Preparation of Amine Intermediates.

**[0828]** We have provided several methods for preparing amine intermediates. The methods provided herein involve the use of nitrogen-protecting groups and deprotecting agents that are used under a variety of conditions. One of skill in the art would understand that the methods provided herein (e.g., Methods BG, BG-1, BG-2, BH, BH-1 BI,) can be applied to a variety of amines and are not limited to the specific examples shown herein. In general, suitable amine protecting groups that could be used include, but are not limited to, [F-moc, Boc, Cbz, Bn, Ts]. Suitable solvents for these reactions include, but are not limited to [dioxane, DMF,  $\text{H}_2\text{O}$ , EtOAc,  $\text{CH}_2\text{Cl}_2$ ]. Suitable deprotecting groups include, but are not limited to, [Piperidine, TFA, HCl,  $\text{H}_2/\text{Pd}-\text{C}$ ,  $\text{BBr}_3$ ,  $\text{HBr}-\text{AcOH}$ ].

Amine Intermediate: Example 1

1-(2,7-Diazaspiro[4.4]nonan-2-yl)propan-1-one

**[0829]**



Method BG

Step 1: tert-Butyl 7-propionyl-2,7-diazaspiro[4.4]nonane-2-carboxylate


**[0830]** To a solution of tert-butyl 3,7-diazaspiro[4.4]nonane-3-carboxylate (127 mg, 0.56 mmol) in anhydrous DCM (1 mL), cooled in an ice water bath was added  $\text{Et}_3\text{N}$  (57 mg, 78  $\mu\text{L}$ , 0.56 mmol), followed by the addition of propanoyl chloride (57 mg, 54  $\mu\text{L}$ , 0.62 mmol) as a solution in DCM (0.2 mL). The reaction mixture was stirred for 20 minutes, diluted with DCM and was washed with 1M HCl ( $\times 2$ ) and brine ( $\times 1$ ). The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under reduced pressure to yield tert-butyl 3-propanoyl-3,7-diazaspiro[4.4]nonane-7-carboxylate (154 mg, 97%) as an orange oil.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  3.73-3.11 (m, 8H), 2.34-2.17 (m, 2H), 1.97-1.70 (m, 4H), 1.46 (s, 9H), 1.16 (t,  $J=7.4$  Hz, 3H). LC/MS  $m/z$  283.5  $[\text{M}+\text{H}]^+$ .

Step 2:


1-(2,7-Diazaspiro[4.4]nonan-2-yl)propan-1-one

**[0831]** To a solution of tert-butyl 3-propanoyl-3,7-diazaspiro[4.4]nonane-7-carboxylate (153 mg, 0.54 mmol) in DCM (0.5 mL) was added a solution of hydrogen chloride (1.4 mL of 4 M, 5.4 mmol) in dioxane. The reaction mixture was stirred at room temperature for 1 h. The solvents were removed under reduced pressure and the crude product was dried on the high vacuum for 1 h to yield 1-(3,7-diazaspiro[4.4]nonan-3-yl)propan-1-one as an off white gum. LC/MS  $m/z$  183.3  $[\text{M}+\text{H}]^+$ .


[0832] The following amines were prepared using the procedure described above.




[0833] 2-Ethylsulfonyl-3,3a,4,5,6,6a-hexahydro-1H-pyrrolo[3,4-c]pyrrole. LC/MS  $m/z$  205.3  $[M+H]^+$ .




[0834] 2-(Ethylsulfonyl)-2,7-diazaspiro[4.4]nonane.




[0835] 1-(2,2-Dimethylpiperazin-1-yl)propan-1-one. LC/MS  $m/z$  171.5  $[M+H]^+$ .




[0836] 1-(Ethylsulfonyl)-2,2-dimethylpiperazine. LC/MS  $m/z$  207.1  $[M+H]^+$ .




[0837] 1-(Piperazin-1-yl)propan-1-one. LC/MS  $m/z$  143.3  $[M+H]^+$ .




[0838] tert-Butyl (3R)-3-isopropyl-4-propanoyl-piperazine-1-carboxylate. LC/MS  $m/z$  185.3  $[M+H]^+$ .




[0839] tert-Butyl (3S)-3-isopropyl-4-propanoyl-piperazine-1-carboxylate. LC/MS  $m/z$  185.3  $[M+H]^+$ .




[0840] 1-[(1S,4R)-2,5-Diazabicyclo[2.2.1]heptan-5-yl]propan-1-one.









[0841] 1-[(2R)-2-Methylpiperazin-1-yl]propan-1-one.




[0842] 1-[(2S)-2-Methylpiperazin-1-yl]propan-1-one.



[0843] (2S)-1-Ethylsulfonyl-2-methyl-piperazine.

[0850] N-(Piperidin-4-yl)propionamide. LC/MS m/z 157.3 [M+H]<sup>+</sup>.[0844] N-[(3R)-Pyrrolidin-3-yl]methyl]acetamide. LC/MS m/z 143.1 [M+H]<sup>+</sup>.[0851] 1-(4,7-Diazaspiro[2.5]octan-4-yl)propan-1-one. LC/MS m/z 169.1 [M+H]<sup>+</sup>.[0845] N-[(3S)-Pyrrolidin-3-yl]methyl]acetamide. LC/MS m/z 143.3 [M+H]<sup>+</sup>.[0846] N-((1R,6S)-3-Azabicyclo[3.1.0]hexan-6-yl)propionamide. LC/MS m/z 155.3 [M+H]<sup>+</sup>.


[0847] N-((1R,6S)-3-Azabicyclo[3.1.0]hexan-6-yl)propane-2-sulfonamide. LC/MS m/z 205.3

[0848] [M+H]<sup>+</sup>.[0849] N-Methyl-N-(piperidin-4-yl)propionamide. LC/MS m/z 171.3 [M+H]<sup>+</sup>.

Amine Intermediate: Example 2

(R)—N-Ethylpyrrolidine-2-carboxamide

[0852]



Method BG-1

Step 1: (R)-tert-Butyl  
2-(ethylcarbamoyl)pyrrolidine-1-carboxylate

[0853] To a solution of (2R)-1-tert-butoxycarbonylpyrrolidine-2-carboxylic acid (810 mg, 3.8 mmol) and HATU (1.40 g, 3.8 mmol) in DMF (7.5 mL) was added ethanamine (3.8 mL of 2 M, 7.5 mmol) followed by DIEA (1.0 g, 1.4 mL, 8.0 mmol) at room temperature. The reaction mixture was stirred at room temperature for 16 h. The reaction mixture was diluted with EtOAc and washed sequentially with 1M HCl, saturated NaHCO<sub>3</sub> and brine solution. The organic layer was

dried ( $\text{Na}_2\text{SO}_4$ ) filtered and concentrated under reduced pressure to obtain tert-butyl (2R)-2-(ethylcarbamoyl)pyrrolidine-1-carboxylate (624 mg, 68%). LC/MS  $m/z$  243.4  $[\text{M}+\text{H}]^+$ .


Step 2: (R)—N-Ethylpyrrolidine-2-carboxamide

**[0854]** To tert-butyl (2R)-2-(ethylcarbamoyl)pyrrolidine-1-carboxylate (561 mg, 2.3 mmol) was added hydrogen chloride (580  $\mu\text{L}$  of 4 M, 2.3 mmol) in dioxane at room temperature under an atmosphere of nitrogen. The reaction mixture was stirred for 3 h. The solvent was removed under reduced pressure to obtain (2R)—N-ethylpyrrolidine-2-carboxamide (225 mg, 54%). LC/MS  $m/z$  143.2  $[\text{M}+\text{H}]^+$ .

**[0855]** The following amines were prepared using the procedure described above.

(S)—N-Ethylpyrrolidine-2-carboxamide

**[0856]**




**[0857]** (S)—N-Ethylpyrrolidine-2-carboxamide. LC/MS  $m/z$  143.2  $[\text{M}+\text{H}]^+$ .

Amine Intermediate: Example 3

(S)-Pyrrolidin-3-yl ethylcarbamate

**[0858]**



Method BG-2

Step 1: (S)-1-(tert-Butoxycarbonyl)pyrrolidin-3-yl 1H-imidazole-1-carboxylate

**[0859]** To a solution of tert-butyl (3S)-3-hydroxypyrrolidine-1-carboxylate (360 mg, 1.9 mmol) in THF (8 mL) was added triethylamine (253 mg, 349  $\mu\text{L}$ , 2.5 mmol) followed by the addition of CDI (374 mg, 2.3 mmol) portionwise. The reaction mixture was heated at 50° C. for 3 h, cooled to room temperature and diluted with EtOAc. The organic layer was washed with 1 M HCl solution, dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude material was purified via silica gel column chromatography using 20 to 100% EtOAc in hexanes to obtain (S)-1-(tert-butoxycarbonyl)pyrrolidin-3-yl 1H-imidazole-1-carboxylate (345 mg, 64%).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.14 (s, 1H), 7.42 (d,  $J=1.2$  Hz, 1H), 7.09 (s, 1H), 4.46 (s, 1H), 3.66 (dd,  $J=33.8$ , 10.7 Hz, 1H), 3.56-3.29 (m, 2H), 2.25 (s, 1H), 2.04-1.87 (m, 2H), 1.48 (t,  $J=4.0$  Hz, 9H). LC/MS  $m/z$  282.4  $[\text{M}+\text{H}]^+$ .

Step 2: tert-Butyl (3S)-3-(Ethylcarbamoyloxy)pyrrolidine-1-carboxylate

**[0860]** To a solution of [(3S)-1-tert-butoxycarbonylpyrrolidin-3-yl]imidazole-1-carboxylate (173 mg, 0.61 mmol) in THF (2.5 mL) at room temperature under an atmosphere of nitrogen was added a solution of ethanamine (369 mg, 461  $\mu\text{L}$  of 2 M, 0.9 mmol) in THF followed by the addition of DIEA (119 mg, 161  $\mu\text{L}$ , 0.9 mmol). The reaction mixture was heated at 60° C. for 1.5 h, cooled to room temperature and diluted with EtOAc. The organic layer was washed with 5 mL of 1M HCl solution, dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure to obtain tert-butyl (3S)-3-(ethylcarbamoyloxy)pyrrolidine-1-carboxylate. LC/MS  $m/z$  259.2  $[\text{M}+\text{H}]^+$ .


Step 3: (S)-Pyrrolidin-3-yl ethylcarbamate


**[0861]** Crude tert-butyl (S)-3-(ethylcarbamoyloxy)pyrrolidine-1-carboxylate from step 2 was treated with hydrogen chloride (461  $\mu\text{L}$  of 4 M, 1.8 mmol) in dioxane. The reaction mixture was stirred for 1.5 h at room temperature. The solvent was removed under reduced pressure to obtain (S)-pyrrolidin-3-yl ethylcarbamate (88 mg, 73%).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.56-8.98 (m, 2H), 4.38 (s, 1H), 3.29-3.05 (m, 4H), 3.03-2.94 (m, 2H), 1.95-1.75 (m, 3H), 1.29 (dd,  $J=13.8$ , 6.6 Hz, 1H), 1.01 (t,  $J=7.2$  Hz, 1H). LC/MS  $m/z$  159.4  $[\text{M}+\text{H}]^+$ .

Amine Intermediate: Example 4

(3aS,6aR)-1,2,3,3a,4,5,6,6a-Octahydrocyclopenta[c]pyrrol-4-ol

**[0862]**





## Method BH

## Step 1: (3aS,6aR)-2-Benzylhexahydrocyclopenta[c]pyrrol-4(5H)-one

**[0863]** A round bottom flask purged with argon was charged with N-benzyl-1-methoxy-N-(trimethylsilylmethyl) methanamine (31.2 g, 131.4 mmol) and cyclopent-2-ene-1-one (9.0 g, 9.2 mL, 109.5 mmol). The reaction mixture was cooled to 0° C. and a solution of trifluoroacetic acid (1.25 g, 847  $\mu$ L, 11.0 mmol) in DCM (11 mL) was added to it. The reaction mixture was stirred at 0° C. for 4 h, warmed to room temperature and stirred for 18 h. The reaction was quenched by the addition of saturated sodium bicarbonate and the layers were separated. The aqueous layer was extracted with DCM and the combined organic layers was washed with brine, dried

over sodium sulfate and concentrated under reduced pressure. The crude material was purified via silica gel column chromatography using 40 to 80% EtOAc in hexanes to obtain (3aR,6aS)-2-benzyl-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-6-one (19.8 g, 84%) as a light yellow oil. LC/MS  $m/z$  216.5  $[M+H]^+$ .

## Step 2: (3aS,6aR)-Benzyl 4-oxohexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

**[0864]** A solution of (3aR,6aS)-2-benzyl-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-6-one (19.8 g, 92 mmol) in dichloromethane (770 mL) was treated with benzyl chloroformate (26.7 g, 22.3 mL, 156 mmol) at room temperature. The reaction mixture was stirred overnight at room temperature. The solvent was removed under reduced pressure and the crude material obtained was purified using silica gel column chromatography using 40 to 99% EtOAc in hexanes to obtain benzyl (3aR,6aS)-6-oxo-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrole-2-carboxylate (11.2 g, 43.19 mmol, 47%) as a light yellow oil.  $^1H$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.41-7.23 (m, 5H), 5.11 (d,  $J=4.2$  Hz, 2H), 3.82-3.52 (m, 3H), 3.29-3.14 (m,  $J=5.6$  Hz, 1H), 3.13-2.93 (m,  $J=11.1$ , 6.8 Hz, 1H), 2.81-2.68 (m,  $J=8.6$ , 3.6 Hz, 1H), 2.37 (t,  $J=7.8$  Hz, 2H), 2.26-2.10 (m, 1H), 1.86 (s, 1H). LC/MS  $m/z$  260.1  $[M+H]^+$ .


## Step 3: (3aS,6aR)-Benzyl 4-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate

**[0865]** A solution of benzyl (3aR,6aS)-6-oxo-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrole-2-carboxylate (850 mg, 3.3 mmol) in THF (10 mL) was added dropwise to a suspension of lithium borohydride (86 mg, 4 mmol) in THF (5 mL) at -78° C. The reaction mixture was allowed to warm to room temperature and was stirred overnight. The reaction mixture was then cooled to 0° C. and quenched with  $H_2O_2$  (30% in  $H_2O$ ) and then poured onto  $H_2O$ . The aqueous layer was extracted with EtOAc (2x) and the combined organic layer was washed with brine, dried over  $Na_2SO_4$ , filtered and concentrated under reduced pressure to obtain benzyl (3aR,6aS)-6-hydroxy-3,3a,4,5,6a-hexahydro-1H-cyclopenta[c]pyrrole-2-carboxylate (850 mg, 99%) as a colorless oil. LC/MS  $m/z$  262.1  $[M+H]^+$ .

Step 4:  
(3aS,6aR)-Octahydrocyclopenta[c]pyrrol-4-ol

**[0866]** A mixture of benzyl (3aR,6aS)-6-hydroxy-3,3a,4,5,6a-hexahydro-1H-cyclopenta[c]pyrrole-2-carboxylate (850 mg, 3.3 mmol) and Pd/C (200 mg) in THF (5.000 mL) was hydrogenated at 25-30 psi in a Parr shaker. After 12 h, the reaction mixture was diluted with MeOH and filtered through a plug of celite. The solvents were removed under reduced pressure to obtain (3aS,6aR)-1,2,3,3a,4,5,6a-octahydrocyclopenta[c]pyrrol-4-ol in quantitative yield. LC/MS  $m/z$  128.1  $[M+H]^+$ .

## Amine Intermediate: Example 5

2-(Isopropylsulfonyl)-2,6-diazaspiro[3.4]octane  
[0867]

## Method BH-1

Step 1: Benzyl  
2,6-diazaspiro[3.4]octane-6-carboxylate

[0868] A solution of 6-benzyl 2-tert-butyl 2,6-diazaspiro[3.4]octane-2,6-dicarboxylate (705 mg, 2.0 mmol) in DCM (1 mL) was treated with hydrogen chloride (5 mL of 4 M, 20 mmol) in dioxane. The reaction mixture was stirred at room temperature for 1 h. The solvent was removed under reduced pressure to obtain benzyl 2,7-diazaspiro[3.4]octane-7-carboxylate. LC/MS  $m/z$  247.3  $[M+H]^+$ .

Step 2: Benzyl 2-(isopropylsulfonyl)-2,6-diazaspiro  
[3.4]octane-6-carboxylate


[0869] A solution of benzyl 2,7-diazaspiro[3.4]octane-7-carboxylate (192 mg, 0.68 mmol) in DCM (1 mL) was treated with triethylamine (137 mg, 189  $\mu$ L, 1.36 mmol) and isopropylsulfonyl chloride (107 mg, 84  $\mu$ L, 0.75 mmol) at 0° C. The reaction mixture was stirred for 20 minutes. The reaction mixture was diluted with DCM and washed with 1N HCl and brine. The organic layer was dried over  $Na_2SO_4$  filtered and concentrated under reduced pressure to obtain benzyl 2-isopropylsulfonyl-2,6-diazaspiro[3.4]octane-6-carboxylate (230 mg, 96%) as a colorless oil. LC/MS  $m/z$  353.3  $[M+H]^+$ .


Step 3:  
2-(Isopropylsulfonyl)-2,6-diazaspiro[3.4]octane

[0870] A solution of benzyl 2-isopropylsulfonyl-2,6-diazaspiro[3.4]octane-6-carboxylate (230 mg, 0.65 mmol) in AcOH (2 mL) was treated with Pd (7.0 mg, 0.06 mmol) (10% on activated carbon) and stirred under an atmosphere of hydrogen for 1.5 h. The reaction mixture was diluted with DCM, filtered and washed with 1N HCl and brine. The organic layer was dried over  $Na_2SO_4$ , filtered and concen-

trated to obtain 2-isopropylsulfonyl-2,6-diazaspiro[3.4]octane in quantitative yield. LC/MS  $m/z$  219.3  $[M+H]^+$ .

## Amine Intermediate: Example 5

8-(Ethylsulfonyl)-3,8-diazabicyclo[3.2.1]octane  
[0871]



## Method BI

Step 1: 3-(9H-Fluoren-9-yl)methyl 8-tert-butyl 3,8-diazabicyclo[3.2.1]octane-3,8-dicarboxylate

**[0872]** A solution of tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylate (105 mg, 0.5 mmol) in dichloromethane (3 mL) and triethylamine (100 mg, 138  $\mu$ L, 1.0 mmol) was treated with 9H-fluoren-9-ylmethyl carbonochloridate (128.0 mg, 0.5 mmol). The reaction mixture was stirred for 1 h. The reaction mixture was washed with 1 N HCl, water, 50% saturated bicarbonate and brine. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to obtain a solution of 3-(9H-fluoren-9-yl)methyl 8-tert-butyl 3,8-diazabicyclo[3.2.1]octane-3,8-dicarboxylate in DCM (2 mL), which was used directly for the next step. LC/MS  $m/z$  435.5  $[\text{M}+\text{H}]^+$ .

Step 2: (9H-Fluoren-9-yl)methyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylate

**[0873]** A solution of 3-(9H-fluoren-9-yl)methyl 8-tert-butyl 3,8-diazabicyclo[3.2.1]octane-3,8-dicarboxylate in DCM (2 mL) from the step 1 was treated dropwise with TFA (2.2 g, 1.5 mL, 19.5 mmol) at room temperature and the reaction mixture was stirred for 1 h. The solvents were removed under reduced pressure and the crude material was dissolved in 1 N HCl (2 mL) and washed with ether. The solution was basified with saturated bicarbonate solution and the aqueous layer was extracted with DCM. The organic layer was dried over

$\text{Na}_2\text{SO}_4$ , filtered and concentrated to obtain a solution of (9H-fluoren-9-yl)methyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylate in DCM (2 mL), which was used directly for the next step. LC/MS  $m/z$  335.5  $[\text{M}+\text{H}]^+$ .

Step 3: (9H-Fluoren-9-yl)methyl 8-(ethylsulfonyl)-3,8-diazabicyclo[3.2.1]octane-3-carboxylate

**[0874]** To the solution of (9H-fluoren-9-yl)methyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylate in DCM (2 mL) from step 2 was added triethylamine (100 mg, 138  $\mu$ L, 1.0 mmol) and ethanesulfonyl chloride (190 mg, 140  $\mu$ L, 1.4 mmol). The reaction mixture was stirred for 16 h. The reaction mixture was washed with 50% saturated bicarbonate solution and brine. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo to obtain (9H-fluoren-9-yl)methyl 8-(ethylsulfonyl)-3,8-diazabicyclo[3.2.1]octane-3-carboxylate, which was used directly for the next step. LC/MS  $m/z$  427.5  $[\text{M}+\text{H}]^+$ .

Step 4:


8-(Ethylsulfonyl)-3,8-diazabicyclo[3.2.1]octane

**[0875]** Crude 9H-fluoren-9-ylmethyl 8-ethylsulfonyl-3,8-diazabicyclo[3.2.1]octane-3-carboxylate obtained from step 3 was dissolved in DMF (500  $\mu$ L) and treated with piperidine (4 mg, 5  $\mu$ L, 0.05 mmol). The reaction mixture was stirred for 2 h at room temperature. The reaction mixture was diluted with methanol/DCM and treated with carbonate resin. The reaction mixture was filtered and concentrated under reduced pressure to yield 8-ethylsulfonyl-3,8-diazabicyclo[3.2.1]octane which was used directly in the next step without further purification. LC/MS  $m/z$  205.3  $[\text{M}+\text{H}]^+$ .


## Example 32

5-(1-(2-Aminoethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(1H-benzo[d]imidazol-2-yl)pyrazin-2-amine (Compound I-313)

**[0876]**



-continued



**[0877]** Compound I-313 was prepared using Method A, Steps 1-3 followed by Method G, Step 1 followed by Method L, Steps 1-2, followed by Method BJ, Steps 1-2.

#### Method BJ

Step 1: 2-(2-(4-(5-Amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)sulfonyl)ethyl)isoindoline-1,3-dione

**[0878]** To a solution of 3-(1H-benzimidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (73 mg, 0.25 mmol) and triethylamine (51 mg, 70  $\mu$ L, 0.50 mmol) in DMSO (1 mL) at 0° C. was added 2-(1,3-dioxoisindolin-2-yl)ethanesulfonyl chloride (68 mg, 0.25 mmol). The reaction mixture was warmed to room temperature, stirred for 3 h and then partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc, dried ( $\text{MgSO}_4$ ), filtered and concentrated. The crude material was dissolved in DMSO and purified by prep HPLC to obtain: 2-(2-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)sulfonyl)ethyl)isoindoline-1,3-dione (23 mg, 17%). LC/MS  $m/z$  530.0  $[\text{M}+\text{H}]^+$ .

#### Method BJ

Step 2: 5-(1-(2-Aminoethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(1H-benzo[d]imidazol-2-yl)pyrazin-2-amine

**[0879]** To a suspension of 2-(2-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-

ylsulfonyl)ethyl)isoindoline-1,3-dione (24 mg, 0.045 mmol) in ethanol (240  $\mu$ L) was added hydrazine (1.5 mg, 1.4  $\mu$ L, 0.045 mmol), and the reaction mixture was heated at 80° C. for 2 h. The reaction mixture was cooled to room temperature and was dissolved in 0.5 ml of DMSO and purified via reverse phase HPLC using 1 to 99% MeOH in  $\text{H}_2\text{O}$  (0.05% HCl as modifier) to obtain 5-(1-(2-aminoethylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(1H-benzo[d]imidazol-2-yl)pyrazin-2-amine (4 mg, 18%).  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  8.59 (s, 1H), 8.12-8.03 (m, 2H), 7.78 (dd,  $J=6.1, 3.2$  Hz, 2H), 7.16 (s, 1H), 4.43 (d,  $J=2.1$  Hz, 2H), 3.95 (t,  $J=5.6$  Hz, 2H), 3.81-3.67 (m, 4H), 3.12 (s, 2H). LC/MS  $m/z$  400.0  $[\text{M}+\text{H}]^+$ .


**[0880]** Compound I-331 was prepared using the method described for Compound I-313 in Example 32 above.

**[0881]** Compound I-331: 5-(1-(3-aminopropylsulfonyl)-1,2,3,6-tetrahydropyridin-4-yl)-3-(1H-benzo[d]imidazol-2-yl)pyrazin-2-amine.  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  8.31 (s, 1H), 7.81 (dd,  $J=6.1, 3.1$  Hz, 2H), 7.52 (dd,  $J=6.2, 3.2$  Hz, 2H), 6.90 (s, 1H), 4.12 (s, 2H), 3.63 (t,  $J=5.6$  Hz, 2H), 3.26 (t,  $J=7.3$  Hz, 2H), 3.18-3.10 (m, 2H), 2.83 (d,  $J=1.5$  Hz, 2H), 2.32-2.05 (m, 2H). LC/MS  $m/z$  414.2  $[\text{M}+\text{H}]^+$ .

#### Example 33


4-(4-(5-Amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-4-oxobutanoic acid. (Compound I-324)

**[0882]**



Method BK  
Step 1

-continued



**[0883]** Compound I-324 was prepared using Method A, Steps 1-3 followed by Method G, Step 1 followed by Method L, Steps 1-2, followed by Method BK, Step 1.



## Method BK

Step 1: 4-(4-(5-Amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-4-oxobutanoic acid

**[0884]** To a solution of 3-(1H-benzimidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (50 mg, 0.12 mmol) in DCM (2 mL) at 0° C. was added methyl 4-chloro-4-oxobutanoate (19 mg, 15  $\mu$ L, 0.12 mmol) followed by the addition of triethylamine (50 mg, 69  $\mu$ L, 0.5 mmol). The reaction mixture was stirred at room temperature for 2 h and then partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc, dried ( $\text{MgSO}_4$ ), filtered and concentrated. The crude material was dissolved in DMSO and purified by reverse phase HPLC using 1-99% MeOH in  $\text{H}_2\text{O}$  to obtain methyl 4-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-4-oxobutanoate, which was re-dissolved in THF (1.0 mL), treated with NaOH (996  $\mu$ L of 2.5 M, 2.5 mmol) and stirred for 18 h at room temperature. The reaction mixture was then acidified with conc. HCl till pH7 and extracted with DCM. The organic layer was dried ( $\text{MgSO}_4$ ), filtered and concentrated. The crude material was purified via reverse phase HPLC using 1-99% MeOH in  $\text{H}_2\text{O}$  (0.05% HCl modifier) to obtain 4-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-4-oxobutanoic acid (5 mg, 11%).  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  8.24 (s, 1H), 7.66 (d,  $J=4.1$  Hz, 2H), 7.27 (dd,  $J=6.0, 3.1$  Hz, 2H), 6.68 (d,  $J=7.9$  Hz, 1H), 4.34 (s, 1H), 4.27 (s, 1H), 3.84 (t,  $J=5.6$  Hz, 2H), 2.85 (s, 1H), 2.80-2.65 (m, 3H), 2.56-2.47 (m, 2H). LC/MS  $m/z$  393.0  $[\text{M}+\text{H}]^+$ .

## Example 34

1-(4-(5-Amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-3-(methylamino)propan-1-one (Compound I-346)

**[0885]**

**[0886]** Compound I-346 was prepared using Method A, Steps 1-3 followed by Method G, Step 1 followed by Method L, Steps 1-2, followed by Method BL, Step 1.

#### Method BL

Step 1: 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-3-(methylamino)propan-1-one


**[0887]** To a solution of 3[tert-butoxycarbonyl(methyl)amino]propanoic acid (14 mg, 0.07 mmol) in DMSO (445  $\mu$ L) was added HATU (26 mg, 0.07 mmol) followed by the addition of triethylamine (28 mg, 38  $\mu$ L, 0.27 mmol) and

**[0889]** Compound I-338 3-amino-1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)propan-1-one.  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  8.26 (d,  $J=16.6$  Hz, 1H), 7.68 (dd,  $J=5.9, 3.2$  Hz, 2H), 7.31 (dd,  $J=6.0, 3.1$  Hz, 2H), 6.73 (d,  $J=36.2$  Hz, 1H), 4.30 (dd,  $J=22.0, 2.6$  Hz, 2H), 3.84 (dt,  $J=44.5, 5.7$  Hz, 2H), 3.25-3.20 (m, 2H), 2.93-2.72 (m, 4H). LC/MS  $m/z$  364.5  $[\text{M}+\text{H}]^+$ .

#### Example 35

5-(4-(Ethylamino)cyclohex-1-en-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (Compound I-314)

**[0890]**



3-(1H-benzimidazol-2-yl)-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine (20 mg, 0.07 mmol) at room temperature. The reaction mixture was stirred at room temperature for 18 h and then partitioned between EtOAc and water. The aqueous layer was extracted with EtOAc, dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated. To the crude material was added 1:1 DCM: TFA (3 ml) and it was stirred for 1 h. The solvents were evaporated under reduced pressure and the crude material obtained was dissolved in DMSO and purified by reverse phase chromatography using 1-99% MeOH in  $\text{H}_2\text{O}$  (0.05% HCl modifier) to obtain 1-(4-(5-amino-6-(1H-benzo[d]imidazol-2-yl)pyrazin-2-yl)-5,6-dihydropyridin-1(2H)-yl)-3-(methylamino)propan-1-one (2 mg, 7%).  $^1\text{H}$  NMR (400 MHz, MeOD)  $\delta$  8.31 (d,  $J=15.9$  Hz, 1H), 7.81 (dd,  $J=6.1, 3.1$  Hz, 2H), 7.52 (dd,  $J=6.0, 3.1$  Hz, 2H), 6.88 (d,  $J=20.4$  Hz, 1H), 4.33 (dd,  $J=13.5, 2.6$  Hz, 2H), 3.85 (dt,  $J=39.3, 5.6$  Hz, 2H), 2.95 (dt,  $J=24.1, 6.0$  Hz, 2H), 2.82 (s, 1H), 2.75 (s, 4H). LC/MS  $m/z$  378.0  $[\text{M}+\text{H}]^+$ .

**[0888]** Compound I-338 was prepared using the method described for Compound I-346 in Example 34 above.

**[0891]** Compound I-314 was prepared using Method A, Steps 1-2 followed by Method N, Step 1-2 followed by Method O, Step 1, followed by Method BM, Steps 1-3.

#### Method BM

Step 1: tert-Butyl N-tert-butoxycarbonyl-N-[5-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]carbamate

**[0892]** A solution of tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonylcarbamate (250 mg, 0.48 mmol), 2-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (128 mg, 0.48 mmol), 4-ditert-butylphosphanyl-N,N-dimethylaniline; dichloropalladium (3.5 mg, 0.005 mmol) and  $\text{K}_2\text{CO}_3$  (133 mg, 0.96 mmol) in toluene (2 mL) and water (250.0  $\mu$ L) was heated to  $80^\circ\text{C}$  for 18 h. The reaction mixture was cooled to room temperature and diluted with DCM (30 mL) and water (30 mL). The layers were separated and the organic layer was washed with 1M NaOH solution, dried ( $\text{Na}_2\text{SO}_4$ ),

filtered and concentrated. The crude material was purified via silica gel column chromatography using 0-30% EtOAc/Hexanes to obtain tert-butyl N-tert-butoxycarbonyl-N-[5-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]carbamate (180 mg, 65%) as a yellow solid. LC/MS  $m/z$  478.26  $[M+H]^+$ .

#### Method BM

##### Step 2: 4-[5-Amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]cyclohex-3-en-1-one

**[0893]** A solution of tert-butyl N-tert-butoxycarbonyl-N-[5-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]carbamate in (40 mg, 0.069 mmol) AcOH (832 mg, 788  $\mu$ L, 13.85 mmol) and H<sub>2</sub>O (50 mg, 50  $\mu$ L, 2.8 mmol) was heated at 80° C. for 2 h. The reaction mixture was diluted with 10 ml of EtOAc and pH was adjusted to 8 with 1N NaOH. The reaction mixture was washed with 20 ml of NaHCO<sub>3</sub>. The layers were separated and the organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to obtain 4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]cyclohex-3-en-1-one (19 mg, 82%), which was used as crude for the next step. LC/MS  $m/z$  334.0  $[M+H]^+$ .


##### Method BM Step 3: 5-[4-(Ethylamino)cyclohexen-1-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0894]** To a solution of 4-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]cyclohex-3-en-1-one (15 mg, 0.045 mmol) in DCE (500  $\mu$ L) was added a solution of ethaneamine (34  $\mu$ L of 2 M, 0.07 mmol) in THF and the reaction mixture was stirred for 5 min. Thereafter Na(OAc)<sub>3</sub>BH (15 mg, 0.07 mmol) was added followed by addition of AcOH (3 mg, 3  $\mu$ L, 0.05 mmol) and the reaction mixture was stirred for 2 h at room temperature. The crude material was dissolved in DMSO and purified via reverse phase HPLC using 1-99 MeOH/H<sub>2</sub>O (0.05% HCl modifier) to obtain 5-[4-(ethylamino)cyclohexen-1-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (2 mg, 9%) as yellow solid. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.84 (s, 1H), 8.56 (s, 1H), 8.12 (d, J=7.9 Hz, 2H), 7.68 (d, J=7.0 Hz, 3H), 7.60 (s, 1H), 6.58 (s, 1H), 3.05 (dd, J=13.2, 6.6 Hz, 2H), 2.76 (m, 3H), 2.38 (m, 1H), 2.27 (d, J=10.9 Hz, 1H), 1.78 (m, 1H), 1.26 (t, J=7.2 Hz, 3H). LC/MS  $m/z$  363.05  $[M+H]^+$ .

#### Example 36

##### Ethyl 2-(5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)cyclopropanecarboxylate (Compound I-337)

**[0895]**



[0896] Compound I-337 was prepared using Method A, Steps 1-2 followed by Method N, Step 1-2 followed by Method O, Step 1, followed by Method BN, Steps 1-3.

#### Method BN

Step 1: tert-Butyl N-tert-butoxycarbonyl-N-[3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-yl]carbamate

[0897] To a solution of tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonylcarbamate (500 mg, 0.96 mmol) in toluene (2 mL) and ethanol (2 mL) was added pyridine; 2,4,6-trivinyl-1,3,5,2,4,6-trioxatriborinane (290 mg, 1.2 mmol),  $K_2CO_3$  (533 mg, 4 mmol) and  $Pd(PPh_3)_4$  (112 mg, 0.1 mmol). The reaction mixture was heated at 60° C. under an atmosphere of nitrogen for 1 h. The reaction mixture was filtered, concentrated and the crude material obtained was purified via silica gel column chromatography using EtOAc/hex gradient to obtain tert-butyl N-tert-butoxycarbonyl-N-[3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-yl]carbamate (449 mg, 100%).

#### Method BN

Step 2: 3-(5-Phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-amine

[0898] To a solution of tert-butyl N-tert-butoxycarbonyl-N-[3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-yl]carbamate (138 mg, 0.3 mmol) in DCM (0.7 mL) at room temperature under an atmosphere of  $N_2$  was added TFA (338 mg, 228  $\mu$ L, 3 mmol) and reaction mixture was stirred for 20

minutes. The reaction mixture was diluted with DCM and slowly quenched with satd.  $NaHCO_3$ . The layers were separated and the organic layer was dried over  $Na_2SO_4$ , filtered and concentrated to yield 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-amine (67 mg, 85%) as a yellow solid. LC/MS  $m/z$  266.4 [M+H]

#### Method BN


Step 3: Ethyl 2-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]cyclopropanecarboxylate

[0899] To a vial charged with 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-vinyl-pyrazin-2-amine (26 mg, 0.01 mmol) under an atmosphere of nitrogen at room temperature was slowly added a solution of ethyl 2-diazoacetate (34 mg, 31  $\mu$ L, 0.1 mmol) in toluene (350  $\mu$ L) over a period of 10 min. The reaction mixture was heated at 110° C. for 1 h. The reaction mixture was cooled to room temperature and the solvent was removed under reduced pressure. The crude material was dissolved in DMSO (1 mL), filtered and purified via reverse phase column chromatography using 10 to 99% MeOH— $H_2O$  (TFA modifier) to obtain ethyl 2-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]cyclopropanecarboxylate.  $^1H$  NMR (400.0 MHz,  $CDCl_3$ )  $\delta$  8.23-8.15 (m, 3H), 7.59-7.50 (m, 3H), 4.19 (q,  $J=7.1$  Hz, 2H), 2.64 (dd,  $J=6.2$ , 12.7 Hz, 1H), 2.25 (dd,  $J=5.5$ , 12.4 Hz, 1H), 1.66-1.61 (m, 2H) and 1.39-1.24 (m, 3H) ppm; LC/MS  $m/z$  352.2 [M+H]

#### Example 37

3-(5-Phenyl-1,3,4-oxadiazol-2-yl)-5-(2,7-diazaspiro[4.4]nonan-2-yl)pyrazin-2-amine (Compound I-340)

[0900]



[0901] Compound I-340 was prepared using Method A, Steps 1-2 followed by Method N, Steps 1-2 followed by Method O, Step 1, followed by Method BO Steps 1-2.

#### Method BO

Step 1: tert-Butyl 7-[5-[bis(tert-butoxycarbonyl)amino]-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,7-diazaspiro[4.4]nonane-3-carboxylate

[0902] A mixture of tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonylcarbamate (70 mg, 0.14 mmol), tert-butyl 3,7-diazaspiro[4.4]nonane-3-carboxylate (31 mg, 0.14 mmol) and DIEA (35 mg, 47  $\mu$ L, 0.27 mmol) in DMF (1 mL) was heated at 90° C. for 2 h. The reaction mixture was allowed to cool to room temperature and diluted with EtOAc. The organic layer was washed with brine (2 $\times$ ), dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure. The crude material was purified by flash chromatography using 0-35% EtOAc in hexanes to yield tert-butyl 7-[5-[bis(tert-butoxycarbonyl)amino]-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,7-diazaspiro[4.4]nonane-3-carboxylate (56 mg, 62%) as a yellow solid. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.15-8.04 (m, 3H), 7.71-7.60 (m, J=7.5 Hz, 3H), 3.74-3.62 (m, 2H), 3.59-3.53 (m, 2H), 3.39 (s, 2H), 2.89 (s, 1H), 2.73 (s, 1H), 2.10-1.99 (m, 2H), 1.98-1.84 (m, 2H), 1.41 (d, J=7.3 Hz, 9H), 1.29 (s, 18H). LC/MS m/z 664.7 [M+H]<sup>+</sup>.

#### Method BO

Step 2: 3-(5-Phenyl-1,3,4-oxadiazol-2-yl)-5-(2,7-diazaspiro[4.4]nonan-2-yl)pyrazin-2-amine

[0903] tert-Butyl 7-[5-[bis(tert-butoxycarbonyl)amino]-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,7-diazaspiro

[4.4]nonane-3-carboxylate (50.9 mg, 0.07668 mmol) was treated with a solution of hydrogen chloride (192  $\mu$ L of 4 M, 0.77 mmol) in dioxane. The reaction mixture was stirred for 2 h at room temperature. The solvent was removed under reduced pressure and the residue was dissolved in DMF and purified by reverse phase HPLC using (1-99% ACN/H<sub>2</sub>O (5 mM HCl)) to yield 3-(5-phenyl-1,3,4-oxadiazol-2-yl)-5-(2,7-diazaspiro[4.4]nonan-2-yl)pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.34 (s, 2H), 8.25-7.99 (m, 2H), 7.89 (s, 1H), 7.66 (d, J=6.6 Hz, 3H), 3.60-3.47 (m, 4H), 3.36-3.25 (m, 2H), 3.18 (tt, J=12.1, 6.0 Hz, 2H), 2.19-1.83 (m, J=17.8, 12.7, 5.9 Hz, 4H); LC/MS m/z 364.1 [M+H]<sup>+</sup>.


[0904] The following compounds were all prepared using the method described for Compound I-340 in Example 37 above.

[0905] Compound I-318: (3aR,6aS)-2-[5-Amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-ol. LC/MS m/z 365.1 [M+H]<sup>+</sup>.

#### Example 38

(3aS,6aR)-2-(5-Amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)hexahydrocyclopenta[c]pyrrol-4(5H)-one (Compound I-319)

[0906]



[0907] Compound I-319 was prepared using Method A, Steps 1-2 followed by Method N, Steps 1-2 followed by Method O, Step 1, followed by Method BP Steps 1-3.

#### Method BP

Step 1: tert-Butyl N-[5-[(3aR,6aS)-6-hydroxy-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

[0908] A solution of tert-butyl N-[5-bromo-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (115 mg, 0.22 mmol), (3aS,6aR)-1,2,3,3a,4,5,6,6a-octahydrocyclopenta[c]pyrrol-4-ol (73 mg, 0.57 mmol) and triethyl amine (34 mg, 46  $\mu$ L, 0.33 mmol) in DMF was heated to 90° C. for 90 minutes. The solvent was removed under reduced pressure and the crude material was purified by silica gel column chromatography using 5-60% EtOAc in DCM to yield tert-butyl N-[5-[(3aR,6aS)-6-hydroxy-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (104 mg, 83%). LC/MS  $m/z$  465.5 [M+H-Boc]<sup>+</sup>.

#### Method BP

Step 2: tert-Butyl N-[5-[(3aR,6aS)-6-oxo-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate

[0909] A solution of tert-butyl N-[5-[(3aR,6aS)-6-hydroxy-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (104 mg, 0.18 mmol) in DCM (1

mL) at 0° C. was treated with a solution of Dess-Martin periodinane (921  $\mu$ L 0.3 M, 0.28 mmol) in DCM. The reaction mixture was warmed to room temperature and stirred overnight. The solvent was evaporated under reduced pressure and the crude material was purified by silica gel column chromatography using 5-60% EtOAc in DCM to yield tert-butyl N-[5-[(3aR,6aS)-6-oxo-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (57 mg, 55%) as a yellow solid. LC/MS  $m/z$  463.5 [M+H-Boc]<sup>+</sup>.

#### Method BP


Step 3: (3aR,6aS)-2-[5-Amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-6-one

[0910] A solution of tert-butyl N-[5-[(3aR,6aS)-6-oxo-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-2-yl]-3-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (57 mg, 0.10 mmol) in DCM (1 mL) was treated with TFA (1.5 g, 1 mL, 13 mmol). The reaction mixture was stirred at room temperature for 1 h. The solvents were evaporated and the residue was dissolved in DMF and purified by reverse phase column chromatography to obtain (3aR,6aS)-2-[5-amino-6-(5-phenyl-1,3,4-oxadiazol-2-yl)pyrazin-2-yl]-1,3,3a,4,5,6a-hexahydrocyclopenta[c]pyrrol-6-one. LC/MS  $m/z$  363.1 [M+H]<sup>+</sup>


#### Example 39

1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one (Compound I-342)

[0911]



-continued



**[0912]** Compound I-342 was prepared using Method BF, Steps 1-2 followed by Method BQ, Steps 1-4

#### Method BQ

Step 1: 5-Bromo-3-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine

**[0913]** A suspension of 5-bromo-3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine (2 g, 4.9 mmol) and  $\text{Na}_2\text{CO}_3$  (1.55 g, 14.60 mmol) in tetrahydrofuran (30 mL) was treated with methanamine (3.6 mL of 2 M, 7.3 mmol) in THF. The reaction mixture was heated at 60° C. for 1 h. The reaction was cooled, diluted with water and the organic layer was extracted with dichloromethane. The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo to yield 5-bromo-3-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (1.7 g, 97%) as a yellow solid, which was used without purification in the next step. LC/MS  $m/z$  363.5  $[\text{M}+\text{H}]^+$ .

#### Method BQ

Step 2: tert-Butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

**[0914]** To a solution of 5-bromo-3-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine (2.1

g, 5.8 mmol) and DMAP (71 mg, 0.58 mmol) in tetrahydrofuran (62 mL) was added a solution of tert-butoxycarbonyl tert-butyl carbonate (6.3 g, 6.7 mL, 29 mmol) in tetrahydrofuran (12 mL). The reaction mixture was heated at 45° C. for 16 h, cooled to room temperature and concentrated in vacuo. The crude material was purified using silica gel column chromatography (10-40% ethyl acetate/hexane) to obtain tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (2.6 g, 67% yield) as a white foam.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.15 (s, 1H), 8.12 (d,  $J=8.1$  Hz, 2H), 7.50 (d,  $J=8.1$  Hz, 2H), 4.50 (s, 2H), 2.84 (s, 3H), 1.28 (s, 27H). LC/MS  $m/z$  663.5  $[\text{M}+\text{H}]^+$ .

#### Method BQ

Step 3: tert-Butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(3-propanoyl-3,7-diazaspiro[4.4]nonan-7-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

**[0915]** To a solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (153 mg, 0.23 mmol) and 1-(3,7-diazaspiro[4.4]nonan-3-yl)propan-1-one

(101 mg, 0.46 mmol) in DMF (2 mL) was added Et<sub>3</sub>N (70 mg, 97  $\mu$ L, 0.70 mmol). The reaction vessel was sealed and heated at 90° C. for 40 minutes. The reaction mixture was cooled to room temperature, diluted with EtOAc and washed with brine (2 $\times$ ). The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography using 0-60% EtOAc in DCM to yield tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(3-propanoyl-3,7-diazaspiro[4.4]nonan-7-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (133 mg, 75%). LC/MS m/z 763.9 [M+H]<sup>+</sup>.

#### Method BQ

Step 4: 1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one

**[0916]** A solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(3-propanoyl-3,7-diazaspiro[4.4]nonan-7-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (129 mg, 0.17 mmol) in dioxane (0.3 mL) was treated with a solution of hydrogen chloride (423  $\mu$ L of 4 M, 1.7 mmol) in dioxane and the reaction mixture was allowed to stir for 40 minutes. The solvents were removed under reduced pressure and the residue was dissolved in DMF (1 mL) and purified by reverse phase HPLC using (10-99% ACN/H<sub>2</sub>O (5 mM HCl modifier)) to yield 1-(7-(5-amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.44 (s, 2H), 8.13 (dd, J=8.3, 2.4 Hz, 2H), 7.91 (t, J=4.5 Hz, 1H), 7.81 (d, J=7.5 Hz, 2H), 4.26-4.20 (m, 2H), 3.58 (dt, J=13.6, 6.8 Hz, 3H), 3.51-3.29 (m, 5H), 2.57 (t, J=5.3 Hz, 3H), 2.24 (dt, J=15.1, 7.5 Hz, 2H), 1.94 (ddt, J=32.7, 13.2, 6.6 Hz, 4H), 0.99 (dd, J=16.5, 7.5 Hz, 3H). LC/MS m/z 463.5 [M+H]<sup>+</sup>.

**[0917]** The following compounds were all prepared using the method described for Compound I-342 in Example 39 above.

**[0918]** Compound I-332: 5-(2-Ethylsulfonyl-1,3,3a,4,6,6a-hexahydropyrrolo[3,4-c]pyrrol-5-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.57 (d, J=4.9 Hz, 2H), 8.13 (d, J=8.2 Hz, 2H), 7.95 (s, 1H), 7.83 (d, J=8.2 Hz, 2H), 4.23 (t, J=5.8 Hz, 2H), 3.65 (dd, J=10.2, 6.9 Hz, 2H), 3.58 (dd, J=9.8, 7.2 Hz, 2H), 3.43 (dd, J=10.5, 3.0 Hz, 2H), 3.21 (dd, J=10.0, 3.7 Hz, 2H), 3.13 (dt, J=11.0, 5.5 Hz, 4H), 2.57 (t, J=5.3 Hz, 3H), 1.22 (t, J=7.4 Hz, 3H). LC/MS m/z 485.5 [M+H]<sup>+</sup>.

**[0919]** Compound I-330: 5-(3-Ethylsulfonyl-3,7-diazaspiro[4.4]nonan-7-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.45 (s, 2H), 8.13 (d, J=8.1 Hz, 2H), 7.91 (s, 1H), 7.81 (d, J=8.0 Hz, 2H), 4.26-4.21 (m, 2H), 3.58 (t, J=6.9 Hz, 2H), 3.54-3.46 (m, 2H), 3.44 (t, J=6.8 Hz, 2H), 3.30 (q, J=9.6 Hz, 2H), 3.16 (q, J=7.3 Hz, 2H), 2.57 (t, J=5.2 Hz, 3H), 2.09-1.90 (m, 4H), 1.23 (dd, J=13.1, 5.7 Hz, 3H). LC/MS m/z 499.2 [M+H]<sup>+</sup>.

**[0920]** Compound I-329: 1-[4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2,2-dimethyl-piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.34 (s, 2H), 8.14 (d, J=7.8 Hz, 3H), 7.80 (d, J=8.2 Hz, 2H), 4.24 (t, J=5.8 Hz, 2H), 3.81-3.76 (m, 2H),

3.71 (s, 2H), 3.62 (t, J=5.3 Hz, 2H), 2.58 (t, J=5.3 Hz, 3H), 2.34 (q, J=7.4 Hz, 2H), 1.43 (s, 6H), 0.98 (t, J=7.3 Hz, 3H). LC/MS m/z 451.5 [M+H]<sup>+</sup>.

**[0921]** Compound I-328: 5-(4-Ethylsulfonyl-3,3-dimethyl-piperazin-1-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.15 (s, 2H), 8.27 (s, 1H), 8.17 (d, J=8.1 Hz, 2H), 7.77 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.7 Hz, 2H), 3.60-3.54 (m, 4H, partially obscured by H<sub>2</sub>O peak), 3.44 (s, 2H, partially obscured by H<sub>2</sub>O peak), 3.14 (q, J=7.3 Hz, 2H), 2.60 (t, J=5.2 Hz, 3H), 1.48 (s, 6H), 1.24 (t, J=7.3 Hz, 3H). LC/MS m/z 487.3 [M+H]<sup>+</sup>.

**[0922]** Compound I-344: 8-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2-methyl-2,8-diazaspiro[4.5]decan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.29 (s, 2H), 8.28 (s, 1H), 8.14 (d, J=8.2 Hz, 2H), 7.78 (d, J=8.2 Hz, 2H), 4.24 (s, 2H), 4.10 (d, J=12.8 Hz, 2H), 3.32 (t, J=6.8 Hz, 2H), 3.00 (t, J=12.4 Hz, 2H), 2.75 (s, 3H), 2.58 (t, J=4.8 Hz, 3H), 2.00 (t, J=6.8 Hz, 2H), 1.77 (t, J=11.5 Hz, 2H), 1.47 (d, J=13.2 Hz, 2H). LC/MS m/z 449.5 [M+H]<sup>+</sup>.

**[0923]** Compound I-341: 5-(8-Ethylsulfonyl-3,8-diazabicyclo[3.2.1]octan-3-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.25 (s, 2H), 8.16 (d, J=8.3 Hz, 3H), 7.78 (d, J=8.2 Hz, 2H), 4.32 (s, 2H), 4.25 (t, J=5.7 Hz, 2H), 3.99 (d, J=10.5 Hz, 2H), 3.19 (q, J=7.1 Hz, 2H), 3.02 (d, J=11.3 Hz, 2H), 2.59 (t, J=5.2 Hz, 3H), 1.98-1.87 (m, 2H), 1.87-1.78 (m, 2H), 1.25 (t, J=7.2 Hz, 3H). LC/MS m/z 485.5 [M+H]<sup>+</sup>.

**[0924]** Compound I-334: 1-[(1S,4R)-2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2,5-diazabicyclo[2.2.1]heptan-5-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, MeOD)  $\delta$  8.27 (d, J=8.1 Hz, 2H), 7.86 (s, 1H), 7.77 (d, J=8.2 Hz, 2H), 5.09-4.94 (m, 2H), 4.33 (s, 2H), 3.82-3.64 (m, 2H), 3.63-3.44 (m, 2H), 2.79 (s, 3H), 2.49 (ddd, J=45.5, 15.6, 7.7 Hz, 1H), 2.32-1.96 (m, 3H), 1.10 (dt, J=36.9, 7.5 Hz, 3H). LC/MS m/z 435.54 [M+H]<sup>+</sup>.

**[0925]** Compound I-315: 1-[(2R)-4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2-methyl-piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, MeOD)  $\delta$  8.28 (d, J=8.2 Hz, 2H), 8.08 (s, 1H), 7.76 (d, J=8.2 Hz, 2H), 4.53-4.36 (m, 1H), 4.33 (s, 2H), 4.27-4.04 (m, 2H), 3.99-3.89 (m, 1H), 3.69-3.55 (m, 1H), 3.19 (d, J=11.7 Hz, 1H), 3.12-2.86 (m, 1H), 2.79 (s, 3H), 2.59-2.39 (m, 2H), 1.34 (d, J=41.1 Hz, 3H), 1.16 (t, J=7.1 Hz, 3H). LC/MS m/z 435.54 [M+H]<sup>+</sup>.

**[0926]** Compound I-321: 5-Isoindolin-2-yl-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.44 (s, 2H), 8.17 (d, J=8.1 Hz, 2H), 8.05 (s, 1H), 7.83 (d, J=8.2 Hz, 2H), 7.45 (dd, J=5.3, 3.2 Hz, 2H), 7.33 (dd, J=5.5, 3.1 Hz, 2H), 4.84 (s, 4H), 4.25 (t, J=5.7 Hz, 2H), 2.59 (t, J=5.2 Hz, 3H). LC/MS m/z 400.5 [M+H]<sup>+</sup>.

**[0927]** Compound I-345: 5-[(3S)-4-Ethylsulfonyl-3-methyl-piperazin-1-yl]-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.25 (d, J=5.0 Hz, 2H), 8.28 (s, 1H), 8.17 (d, J=8.2 Hz, 2H), 7.78 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.7 Hz, 2H), 4.17 (d, J=12.1 Hz, 1H), 4.09 (dd, J=8.4, 4.2 Hz, 1H), 4.01 (d, J=12.6 Hz, 1H), 3.40-3.30 (m, 2H), 3.21-3.09 (m, 2H), 3.09-3.00 (m, 1H), 2.86 (td, J=12.0, 3.2 Hz, 1H), 2.59 (t, J=5.3 Hz, 3H), 1.31 (d, J=6.7 Hz, 3H), 1.23 (t, J=7.3 Hz, 3H). LC/MS m/z 473.3 [M+H]<sup>+</sup>.

**[0928]** Compound I-347: 1-[(2R)-4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2-methyl-piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.32 (d, J=4.2 Hz, 2H), 8.28 (s, 1H), 8.16 (d, J=8.1 Hz, 2H), 7.79 (d, J=8.1 Hz, 2H), 4.44-4.29 (m, 1H), 4.25 (t, J=5.8 Hz, 2H), 4.21-4.11 (m, 1H), 4.02 (d, J=11.4 Hz, 1H), 3.90-3.76 (m, 2H), 2.99 (s, 2H), 2.90-2.64 (m, 1H), 2.59 (t, J=5.3 Hz, 3H), 2.43-2.22 (m, 2H), 1.27 (s, 1H), 1.22-1.09 (m, 1H), 1.02 (t, J=7.2 Hz, 3H). LC/MS m/z 438.34 [M+H]<sup>+</sup>.

**[0929]** Compound I-323: N-[(3S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-3-yl]methylacetamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.17 (s, 2H), 8.15 (d, J=8.0 Hz, 2H), 8.04 (t, J=5.5 Hz, 1H), 7.90 (s, 1H), 7.77 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.9 Hz, 2H), 3.59 (dd, J=20.4, 10.2 Hz, 2H), 3.44 (dd, J=17.2, 7.3 Hz, 1H), 3.22-3.01 (m, 3H), 2.63-2.53 (m, 3H), 2.46-2.39 (m, 1H), 2.06 (dd, J=12.3, 5.4 Hz, 1H), 1.88 (d, J=26.8 Hz, 3H), 1.73 (dd, J=12.6, 7.1 Hz, 1H). LC/MS m/z 423.5 [M+H]<sup>+</sup>.

**[0930]** Compound I-317: N-[(3R)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-3-yl]methylacetamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.06 (s, 2H), 8.15 (d, J=8.2 Hz, 2H), 8.04 (s, 1H), 7.90 (s, 1H), 7.76 (d, J=8.2 Hz, 2H), 4.25 (s, 2H), 3.58 (dd, J=20.3, 10.4 Hz, 2H), 3.44 (dd, J=17.0, 6.9 Hz, 1H), 3.14 (s, 3H), 2.60 (d, J=5.3 Hz, 3H), 2.33 (s, 1H), 2.06 (d, J=6.8 Hz, 1H), 1.84 (s, 3H), 1.73 (dd, J=11.9, 7.3 Hz, 1H). LC/MS m/z 423.5 [M+H]<sup>+</sup>.

**[0931]** Compound I-316: 3-[5-[4-(Methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-(3-phenylpyrrolidin-1-yl)pyrazin-2-amine. LC/MS m/z 428.2 [M+H]<sup>+</sup>.

**[0932]** Compound I-322: 5-(4-Ethylpiperazin-1-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 10.78 (s, 1H), 9.38 (s, 2H), 8.34 (s, 1H), 8.17 (d, J=8.2 Hz, 2H), 7.80 (d, J=8.2 Hz, 2H), 4.37-4.15 (m, 3H), 3.59 (d, J=12.5 Hz, 3H), 3.36-3.24 (m, 3H), 3.20-3.04 (m, 3H), 2.58 (t, J=5.3 Hz, 3H), 1.30 (t, J=7.2 Hz, 3H). LC/MS m/z 395.0 [M+H]<sup>+</sup>.

**[0933]** Compound I-327: 1-[4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.33 (s, 2H), 8.27 (s, 1H), 8.16 (d, J=8.1 Hz, 2H), 7.80 (d, J=8.1 Hz, 2H), 4.24 (t, J=5.4 Hz, 2H), 3.62 (s, 4H), 3.46 (d, J=21.5 Hz, 4H), 2.58 (t, J=5.3 Hz, 3H), 2.39 (q, J=7.3 Hz, 2H), 1.02 (t, J=7.4 Hz, 3H). LC/MS m/z 423.5 [M+H]<sup>+</sup>.

**[0934]** Compound I-325: 5-(4-Ethylsulfonylpiperazin-1-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.24 (s, 2H), 8.28 (s, 1H), 8.17 (d, J=8.0 Hz, 2H), 7.78 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.8 Hz, 2H), 3.59-3.51 (m, 4H), 3.42-3.30 (m, 4H), 3.12 (q, J=7.4 Hz, 2H), 2.59 (t, J=5.2 Hz, 3H), 1.24 (t, J=7.3 Hz, 3H). LC/MS m/z 459.5 [M+H]<sup>+</sup>.

**[0935]** Compound I-333: 1-[(2R)-4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2-isopropyl-piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.30 (s, 2H), 8.25 (s, 1H), 8.13 (d, J=8.1 Hz, 2H), 7.79 (d, J=8.2 Hz, 2H), 4.50-4.38 (m, 2H), 4.31-4.21 (m, 3H), 4.12 (dd, J=8.6, 6.4 Hz, 1H), 3.85 (d,

J=13.4 Hz, 0.6H), 3.65 (d, J=10.9 Hz, 0.4H), 3.26 (t, J=11.4 Hz, 0.6H), 2.87 (ddd, J=15.2, 9.2, 5.6 Hz, 1H), 2.81-2.65 (m, 1.5H), 2.58 (t, J=5.2 Hz, 3H), 2.48-2.26 (m, 3H), 2.25-2.00 (m, 1H), 1.14-0.99 (m, 6H), 0.80 (d, J=6.6 Hz, 1.2H), 0.74 (d, J=6.7 Hz, 1.8H). LC/MS m/z 465.7 [M+H]<sup>+</sup>.

**[0936]** Compound I-326: 1-[(2S)-4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-2-isopropyl-piperazin-1-yl]propan-1-one. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.30 (s, 2H), 8.25 (s, 1H), 8.13 (d, J=8.1 Hz, 2H), 7.79 (d, J=8.2 Hz, 2H), 4.50-4.38 (m, 2H), 4.31-4.21 (m, 3H), 4.12 (dd, J=8.6, 6.4 Hz, 1H), 3.85 (d, J=13.4 Hz, 0.6H), 3.65 (d, J=10.9 Hz, 0.4H), 3.26 (t, J=11.4 Hz, 0.6H), 2.87 (ddd, J=15.2, 9.2, 5.6 Hz, 1H), 2.81-2.65 (m, 1.5H), 2.58 (t, J=5.2 Hz, 3H), 2.48-2.26 (m, 3H), 2.25-2.00 (m, 1H), 1.14-0.99 (m, 6H), 0.80 (d, J=6.6 Hz, 1.2H), 0.74 (d, J=6.7 Hz, 1.8H). LC/MS m/z 465.5 [M+H]<sup>+</sup>.

**[0937]** Compound I-343: 3-[5-[4-(Aethylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-pyrrolidin-1-yl-pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.00 (s, 2H), 8.15 (d, J=8.2 Hz, 2H), 7.91 (s, 1H), 7.74 (d, J=8.2 Hz, 2H), 4.26 (s, 2H), 3.47 (t, J=6.3 Hz, 4H), 2.63 (s, 3H), 1.97 (t, J=6.4 Hz, 4H). LC/MS m/z 428.2 [M+H]<sup>+</sup>.

**[0938]** Compound I-348: (2S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-ethyl-pyrrolidine-2-carboxamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.85 (s, 2H), 8.19 (d, J=8.1 Hz, 2H), 8.00 (t, J=5.7 Hz, 1H), 7.87 (s, 1H), 7.73 (d, J=8.2 Hz, 2H), 4.36-4.22 (m, 3H), 3.52-3.39 (m, 2H), 3.05 (dtd, J=19.9, 13.4, 6.3 Hz, 2H), 2.64 (dd, J=13.5, 8.3 Hz, 3H), 2.25-2.14 (m, 1H), 2.04-1.92 (m, 3H), 0.91 (t, J=7.1 Hz, 3H).

**[0939]** Compound I-352: [(2S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-2-yl]methanol. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.87 (s, 2H), 8.16 (d, J=8.2 Hz, 2H), 7.99 (s, 1H), 7.73 (d, J=8.2 Hz, 2H), 6.69 (s, 2H), 4.26 (t, J=5.7 Hz, 2H), 4.05-3.99 (m, 1H), 3.64-3.55 (m, 3H), 3.44-3.32 (m, 2H), 2.62 (t, J=5.1 Hz, 3H), 1.98-1.92 (m, 3H); LC/MS m/z 382.4 [M+H]<sup>+</sup>.

**[0940]** Compound I-356: [(2R)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-2-yl]methanol. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.34 (s, 2H), 8.14 (d, J=8.1 Hz, 1H), 7.98 (t, J=5.8 Hz, 2H), 7.80 (d, J=8.2 Hz, 1H), 7.67 (d, J=8.2 Hz, 1H), 4.26-4.17 (m, 3H), 4.09-3.97 (m, 1H), 3.66-3.49 (m, 2H), 3.45-3.32 (m, 1H), 3.31-3.22 (m, 1H), 2.57 (dd, J=11.3, 5.6 Hz, 3H), 1.99-1.90 (m, 3H); LC/MS m/z 382.2 [M+H]<sup>+</sup>.

**[0941]** Compound I-357: (2R)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-ethyl-pyrrolidine-2-carboxamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.03 (s, 2H), 8.18 (d, J=8.1 Hz, 2H), 8.02 (t, J=4.9 Hz, 1H), 7.87 (s, 1H), 7.76 (d, J=8.2 Hz, 2H), 4.37-4.21 (m, 3H), 3.13-2.95 (m, 2H), 2.61 (t, J=5.3 Hz, 3H), 2.27-2.12 (m, 2H), 2.07-1.90 (m, 4H), 0.91 (t, J=7.2 Hz, 3H); LC/MS m/z 423.2 [M+H]<sup>+</sup>.

**[0942]** Compound I-361: 3-[5-[4-(Methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-[(2S)-2-methylpyrrolidin-1-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.85 (s, 2H), 8.14 (d, J=8.1 Hz, 2H), 7.92 (s, 1H), 7.73 (d,

J=8.2 Hz, 2H), 4.26 (s, 2H), 4.14 (s, 1H), 2.67 (s, 1H), 2.62 (t, J=5.3 Hz, 3H), 2.45 (s, 1H), 2.33 (s, 1H), 2.06 (s, 3H), 1.70 (s, 1H), 1.24 (d, J=6.1 Hz, 3H). LC/MS m/z 366.4 [M+H]<sup>+</sup>.

**[0943]** Compound I-362: 3-[5-[4-(Methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-[(2R)-2-methylpyrrolidin-1-yl]pyrazin-2-amine. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.85 (s, 2H), 8.14 (d, J=8.2 Hz, 2H), 7.92 (s, 1H), 7.73 (d, J=8.1 Hz, 2H), 4.26 (s, 2H), 4.15 (s, 1H), 2.67 (s, 1H), 2.62 (t, J=5.3 Hz, 3H), 2.33 (s, 1H), 2.06 (s, 3H), 1.70 (s, 1H), 1.24 (d, J=6.2 Hz, 3H). LC/MS m/z 366.2 [M+H]<sup>+</sup>.

**[0944]** Compound I-359: (R)-1-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-3-methylpiperazin-1-yl)propan-1-one. <sup>1</sup>H NMR (400 MHz, MeOD) δ 8.27 (d, J=8.3 Hz, 2H), 8.15 (d, J=3.7 Hz, 1H), 7.76 (d, J=8.3 Hz, 2H), 4.63-4.39 (m, 1H), 4.33 (s, 2H), 4.07 (m, 2H), 3.82 (dd, J=42.9, 13.5 Hz, 1H), 3.71-3.36 (m, 3H), 3.09 (m, 1H), 2.79 (s, 3H), 2.57-2.44 (m, 2H), 1.36-1.07 (m, 8H). LC/MS m/z 437.5 [M+H]<sup>+</sup>.

**[0945]** Compound I-360: 845-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,8-diazaspiro[4.5]decan-1-one; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.40 (d, J=4.5 Hz, 2H), 8.28 (s, 1H), 8.14 (d, J=8.1 Hz, 2H), 7.80 (d, J=8.2 Hz, 2H), 7.63 (s, 1H), 4.23 (t, J=5.8 Hz, 2H), 4.10 (d, J=13.1 Hz, 2H), 3.21 (t, J=6.7 Hz, 2H), 3.00 (t, J=11.5 Hz, 2H), 2.57 (t, J=5.3 Hz, 3H), 2.04 (t, J=6.7 Hz, 2H), 1.76 (td, J=12.8, 4.1 Hz, 2H), 1.48 (d, J=13.2 Hz, 2H); LC/MS m/z 435.5 [M+H]<sup>+</sup>.

**[0946]** Compound I-363: 1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-4,7-diazaspiro[2.5]octan-4-yl)propan-1-one; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.39 (s, 2H), 8.20 (s, 1H), 8.16 (d, J=8.1 Hz, 2H), 7.80 (d, J=8.1 Hz, 2H), 4.29-4.17 (m, J=5.8 Hz, 4H), 3.82-3.45 (m, 4H), 3.34 (s, 3H), 2.58 (t, J=5.3 Hz, 3H), 1.23 (s, 1H), 1.11-0.97 (m, 7H). LC/MS m/z 449.3 [M+H]<sup>+</sup>.

**[0947]** Compound I-364: 1-(1-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-4-yl)pyrrolidin-2-one; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.40 (d, J=4.6 Hz, 2H), 8.28 (s, 1H), 8.15 (d, J=8.2 Hz, 2H), 7.80 (d, J=8.2 Hz, 2H), 4.30 (d, J=12.9 Hz, 2H), 4.24 (t, J=5.8 Hz, 2H), 4.04-3.92 (m, 1H), 3.31 (t, J=6.9 Hz, 2H), 2.88 (t, J=11.5 Hz, 2H), 2.57 (t, J=5.3 Hz, 3H), 2.24 (t, J=8.0 Hz, 2H), 1.96-1.84 (m, 2H), 1.77-1.65 (m, 4H); LC/MS m/z 449.3 [M+H]<sup>+</sup>.

**[0948]** Compound I-365: N-(1-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-4-yl)-N-methylpropionamide; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.41 (s, 2H), 8.28 (d, J=5.1 Hz, 1H), 8.15 (d, J=8.0 Hz, 2H), 7.80 (d, J=7.7 Hz, 2H), 4.31 (d, J=12.6 Hz, 3H), 4.24 (t, J=5.7 Hz, 2H), 2.88 (dd, J=24.8, 12.7 Hz, 2H), 2.80 (s, 2H), 2.69 (s, 1H), 2.57 (t, J=5.3 Hz, 3H), 2.44-2.38 (m, J=7.4 Hz, 1H), 2.31 (dd, J=14.6, 7.2 Hz, 1H), 1.89-1.67 (m, J=14.0, 10.5 Hz, 3H), 1.57 (d, J=10.5 Hz, 1H), 1.04-0.95 (m, J=11.4, 7.4 Hz, 3H). LC/MS m/z 451.3 [M+H]<sup>+</sup>.

**[0949]** Compound I-366: N-(1-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-4-yl)propionamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.24 (s, 1H), 8.03 (d, J=8.1 Hz, 2H), 7.74 (d, J=7.7 Hz, 1H), 7.58 (d, J=8.2 Hz, 2H), 6.85 (s, 2H), 4.10 (d,

J=13.0 Hz, 2H), 3.87-3.65 (m, 3H), 2.95 (t, J=11.1 Hz, 2H), 2.29 (s, 3H), 2.10-2.02 (m, 2H), 1.84 (d, J=10.2 Hz, 2H), 1.46 (dd, J=20.2, 10.9 Hz, 2H), 0.99 (t, J=7.6 Hz, 3H). LC/MS m/z 437.5 [M+H]<sup>+</sup>.

**[0950]** Compound I-368: 5-(2-(Isopropylsulfonyl)-2,6-diazaspiro[3.4]octan-6-yl)-3-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-amine; <sup>1</sup>H NMR (400 MHz, DMSO) δ 9.14 (s, 2H), 8.16 (d, J=8.1 Hz, 2H), 7.93 (s, 1H), 7.77 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.8 Hz, 2H), 3.94 (d, J=7.8 Hz, 2H), 3.86 (d, J=7.8 Hz, 2H), 3.68 (s, 2H), 3.52 (t, J=6.8 Hz, 2H), 3.31-3.20 (m, 1H), 2.60 (t, J=5.2 Hz, 3H), 2.22 (t, J=6.8 Hz, 2H), 1.25 (d, J=6.8 Hz, 6H); LC/MS m/z 499.1 [M+H]<sup>+</sup>.

**[0951]** Compound I-358: N-[(1S,5R)-3-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexan-6-yl]propanamide; <sup>1</sup>H NMR (400 MHz, DMSO) δ 9.11 (s, 2H), 8.17 (d, J=8.1 Hz, 2H), 7.99 (d, J=3.6 Hz, 1H), 7.91 (s, 1H), 7.77 (d, J=8.2 Hz, 2H), 4.25 (t, J=5.8 Hz, 3H), 3.83 (d, J=10.0 Hz, 2H), 3.41 (d, J=9.1 Hz, 2H), 2.60 (t, J=5.2 Hz, 3H), 2.05 (d, J=7.6 Hz, 2H), 1.79 (s, 2H), 0.99 (t, J=7.5 Hz, 3H); LC/MS m/z 435.5 [M+H]<sup>+</sup>.

**[0952]** Compound I-369: N-[(1S,5R)-3-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexan-6-yl]ethanesulfonamide; <sup>1</sup>H NMR (400 MHz, DMSO) δ 9.03 (s, 2H), 8.17 (d, J=8.3 Hz, 2H), 7.91 (s, 1H), 7.76 (d, J=8.3 Hz, 2H), 7.56 (s, 1H), 6.76 (s, 2H), 4.24-4.22 (m, 2H), 3.82 (d, J=10.0 Hz, 2H), 3.43 (s, 2H), 3.11 (t, J=7.3 Hz, 2H), 2.61 (t, J=5.1 Hz, 3H), 2.33 (s, 1H), 1.96 (s, 2H), 1.21 (t, J=7.3 Hz, 3H); LC/MS m/z 471.3 [M+H]<sup>+</sup>.

**[0953]** Compound I-370: N-[(1S,5R)-3-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexan-6-yl]propane-2-sulfonamide; <sup>1</sup>H NMR (400 MHz, DMSO) δ 9.05 (s, 2H), 8.16 (d, J=9.4 Hz, 2H), 7.91 (s, 1H), 7.76 (d, J=8.2 Hz, 2H), 7.55 (s, 1H), 4.25 (t, J=5.8 Hz, 2H), 3.81 (d, J=10.2 Hz, 2H), 3.42 (d, J=9.8 Hz, 2H), 3.38-3.31 (m, 1H), 2.61 (t, J=5.3 Hz, 3H), 2.33 (s, 1H), 1.96 (s, 2H), 1.24 (d, J=5.2 Hz, 6H); LC/MS m/z 485.5 [M+H]<sup>+</sup>.

**[0954]** Compound I-371: (3S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-3-ol; LC/MS m/z 368.1 [M+H]<sup>+</sup>.


**[0955]** Compound I-372: [(3S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-3-yl]N-ethylcarbamate; <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 8.21 (s, 1H), 8.04 (d, J=8.1 Hz, 2H), 7.92 (d, J=15.9 Hz, 1H), 7.62 (t, J=7.1 Hz, 2H), 7.19 (t, J=5.5 Hz, 0.5H), 6.72 (s, 1.5H), 5.27 (s, 1H), 3.82 (s, 2H), 3.75-3.46 (m, 3H), 3.00 (dt, J=13.3, 6.8 Hz, 2H), 2.34 (s, 3H), 2.28-2.17 (m, 1.5H), 2.16-2.04 (m, 1.5H), 1.13 (t, J=7.0 Hz, 0.5H), 0.99 (t, J=7.2 Hz, 3H), 0.82 (d, J=6.9 Hz, 0.5H). LC/MS m/z 439.14 [M+H]<sup>+</sup>.

**[0956]** Compound I-373: [(3S)-1-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]pyrrolidin-3-yl]N-(2-methoxyethyl)carbamate. LC/MS m/z 469.2 [M+H]<sup>+</sup>.


## Example 40

[(3aR,6aS)-2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]methanesulfonate (Compound I-320)

[0957]



-continued



**[0958]** Compound I-320 was prepared using Method BF, Steps 1-2, followed by Method BQ Steps 1-3, followed by Method BR Steps 1-2.

#### Method BR

Step 1: [(3aR,6aS)-2-[5-[bis(tert-Butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]methanesulfonate

**[0959]** To a solution of tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-hydroxy-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (70 mg, 0.10 mmol) and DMAP (1.2 mg, 0.01 mmol) in DCM at 0° C. was added triethylamine (11 mg, 15  $\mu$ L, 0.11 mmol) followed by the addition of methane sulfonyl chloride (12 mg, 8.  $\mu$ L, 0.10 mmol). The reaction mixture was warmed to room temperature over a period of 1.5 h and used directly in the next step.

#### Method BR


Step 2: [(3aR,6aS)-2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]methanesulfonate

**[0960]** To the crude reaction mixture from step 1 was added TFA (1 mL). The reaction mixture was stirred for 30 minutes at room temperature. The solvent was evaporated and the crude material was redissolved in DMSO and purified via reverse phase HPLC to obtain [(3aR,6aS)-2-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]methanesulfonate (8 mg, 23%). LC/MS  $m/z$  486.2 [M+H]<sup>+</sup>.

#### Example 41

N-(3aS,6aR)-2-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)octahydrocyclopenta[c]pyrrol-4-yl)ethanesulfonamide (Compound I-354)

**[0961]**



[0962] Compound I-354 was prepared using Method BF, Steps 1-2, followed by Method BQ Steps 1-3, followed by Method BR Step 1, followed by Method BS Steps 1-4.

#### Method BS

Step 1: tert-Butyl N-[[4-[5-[6-[(3aR,6aS)-6-azido-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0963] A solution of [(3aR,6aS)-2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]methanesulfonate (460 mg, 0.58 mmol) and sodium azide (190 mg, 573  $\mu$ L, 2.9 mmol) in DMF (4 mL) was heated at 70° C. for 48 h. The solvent was removed under reduced pressure and the residue was dissolved in EtOAc. The organic layer was washed with H<sub>2</sub>O, followed by washing with brine. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure to obtain tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-azido-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (401 mg, 93%). LC/MS m/z 733.7 [M+H]<sup>+</sup>.

#### Method BS

Step 2: tert-Butyl N-[[4-[5-[6-[(3aR,6aS)-6-amino-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0964] A flask charged with tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-azido-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (400 mg, 0.55 mmol) and Pd/C (108 mg, 1.0 mmol) was flushed under N<sub>2</sub> followed by evacuating under vacuum. EtOH (3 mL) was added under inert atmosphere followed by evacuating under vacuum. The reaction mixture was stirred for 4.5 h under an atmosphere of H<sub>2</sub>. The reaction mixture was filtered and the solvent was evaporated under reduced pressure to obtain tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-amino-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (300 mg, 78%). LC/MS m/z 707.5 [M+H]<sup>+</sup>.

#### Method BS

Step 3: tert-Butyl N-[[4-[5-[6-[(3aR,6aS)-6-(ethylsulfonylamino)-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0965] To a solution of tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-amino-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (150 mg, 0.22 mmol) and triethylamine (24 mg, 33  $\mu$ L, 0.23 mmol) in DCM (1 mL) at 0° C. was added ethanesulfonyl chloride (29 mg, 214, 0.22 mmol). The reaction mixture was stirred for 30 min. The solvent was evaporated and the crude material was purified via silica gel column chromatography using 10 to 70% EtOAc in DCM to obtain tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-(ethylsulfonylamino)-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (110 mg, 650%). LC/MS m/z 799.5 [M+H]<sup>+</sup>.

#### Method BS

Step 4: N-[(3aR,6aS)-2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]ethanesulfonamide

[0966] A solution of tert-butyl N-[[4-[5-[6-[(3aR,6aS)-6-(ethylsulfonylamino)-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-2-yl]-3-[bis(tert-butoxycarbonyl)amino]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (110 mg, 0.14 mmol) in DCM (1 mL) was treated with TFA (1.0 g, 676  $\mu$ L, 8.8 mmol). The reaction mixture was stirred for 1 hour. The solvent was evaporated and the residue was purified via reverse phase HPLC using 10 to 99% ACN in water (5 mM HCl modifier) to obtain N-[(3aR,6aS)-2-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,3a,4,5,6,6a-hexahydro-1H-cyclopenta[c]pyrrol-6-yl]ethanesulfonamide. LC/MS m/z 499.2 [M+H]<sup>+</sup>.


[0967] Compound I-355: was prepared using the method described for Compound I-354 in Example 41 above.

[0968] Compound I-355: N-((3aS,6aR)-2-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)octahydrocyclopenta[c]pyrrol-4-yl)propanamide. LC/MS m/z 463.2 [M+H]<sup>+</sup>.


#### Example 42

1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one (Compound I-349)

[0969]



-continued



[0970] Compound I-349 was prepared using Method BF, Steps 1-2 followed by Method BQ, Steps 1-3, followed by the Method BT, Steps 1-2.

#### Method BT

Step 1: (1R,5S)-3-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid

[0971] To a solution of ethyl (1R,5S)-3-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylate (270 mg, 0.37 mmol) in of 2:1 THF: MeOH (1 mL) was added NaOH (550  $\mu$ L of 1 M, 0.55 mmol). The reaction mixture was stirred for 30 minutes at room temperature. The solvent was evaporated to obtain (1R,5S)-3-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid, which was used for the next step without further purification.

#### Method BT


Step 2: (1R,5S)-3-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid

[0972] To a solution of (1R,5S)-3-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid (30 mg, 0.04 mmol) in DCM (0.3 mL) was added TFA (500  $\mu$ L, 6.50 mmol) and reaction was stirred at room temperature for 2 h. The solvent was removed under reduced pressure and the residue was dissolved in DMSO, filtered and purified via reverse phase HPLC using a gradient of 10 to 99% MeOH—H<sub>2</sub>O (TFA modifier) to obtain (1R,5S)-3-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid. LC/MS  $m/z$  408.4 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  12.23 (s, 1H), 8.85 (s, 2H), 8.17 (d, J=8.2 Hz, 2H), 7.91 (s, 1H), 7.73 (d, J=8.2 Hz, 2H), 6.77 (s, 2H), 4.26 (t, J=5.7 Hz, 2H), 3.87 (d, J=10.4 Hz, 2H), 3.47-3.46 (m, 2H), 2.63 (t, J=5.2 Hz, 3H), 2.19 (s, 2H), 1.49 (t, J=2.9 Hz, 1H).


#### Example 43

1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one (Compound I-351)

[0973]



-continued



[0974] Compound I-351 was prepared using Method BF, Steps 1-2 followed by Method BQ, Step 1-3, followed by the Method BT, Step 1, followed by Method BU, Steps 1-2.

#### Method BU

Step 1: tert-Butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[(1R,5S)-6-(dimethylcarbamoyl)-3-azabicyclo[3.1.0]hexan-3-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0975] To a solution of (1R,5S)-3-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3-azabicyclo[3.1.0]hexane-6-carboxylic acid (127 mg, 0.18 mmol) and HATU (68 mg, 0.18 mmol) in DMF (720  $\mu$ L) at room temperature was added DIEA (49 mg, 66  $\mu$ L, 0.38 mmol) followed by the addition of a solution of N-methylmethanamine (253 mg, 270  $\mu$ L of 2 M, 0.54 mmol) in THF. The reaction mixture was stirred at room temperature for 1.5 h. The reaction mixture was diluted with ethyl acetate and washed with brine solution. The organics were separated, dried over  $\text{MgSO}_4$ , filtered and concentrated under reduced pressure. The crude material was purified via silica gel column chromatography using 2 to 15% MeOH in DCM to obtain tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[(1R,5S)-6-(dimethylcarbamoyl)-3-azabicyclo[3.1.0]hexan-3-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (81 mg, 61%).

#### Method BU

Step 2: (1R,5S)-3-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N,N-dimethyl-3-azabicyclo[3.1.0]hexane-6-carboxamide

[0976] A solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[(1R,5S)-6-(dimethylcarbamoyl)-3-

azabicyclo[3.1.0]hexan-3-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (75 mg, 0.1 mmol) in 0.8 mL of 50% TFA-DCM mixture was stirred under an atmosphere of nitrogen at room temperature for 1 h. The solvents were evaporated and the crude material was purified via reverse phase HPLC using 10 to 99% methanol in water (TFA modifier) to obtain (1R,5S)-3-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N,N-dimethyl-3-azabicyclo[3.1.0]hexane-6-carboxamide.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.87 (s, 2H), 8.16 (d,  $J=8.2$  Hz, 2H), 7.91 (s, 1H), 7.73 (d,  $J=8.2$  Hz, 2H), 6.76 (s, 2H), 4.26 (t,  $J=5.7$  Hz, 2H), 3.86 (d,  $J=10.2$  Hz, 2H), 3.49-3.46 (m, 2H), 3.10 (s, 3H), 2.84 (s, 3H), 2.63 (t,  $J=5.2$  Hz, 3H), 2.09 (s, 2H), 1.90 (t,  $J=3.1$  Hz, 1H). LC/MS  $m/z$  435.2  $[\text{M}+\text{H}]^+$ .

[0977] Compound I-350 was prepared using the method described for Compound I-351 in Example 43 above.


[0978] Compound I-350: (1R,5S)-3-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-ethyl-3-azabicyclo[3.1.0]hexane-6-carboxamide.

$^1\text{H}$  NMR (400 MHz,  $\text{DMSO-d}_6$ )  $\delta$  8.86 (s, 2H), 8.17 (d,  $J=8.1$  Hz, 2H), 8.00 (t,  $J=5.4$  Hz, 1H), 7.95-7.88 (m, 1H), 7.73 (d,  $J=8.2$  Hz, 2H), 6.77 (s, 2H), 4.26 (t,  $J=5.6$  Hz, 2H), 3.85 (t,  $J=11.5$  Hz, 2H), 3.09 (dt,  $J=14.0$ , 7.0 Hz, 2H), 2.64 (dd,  $J=14.4$ , 9.4 Hz, 4H), 2.33 (s, 1H), 2.19 (s, 1H), 1.52 (dd,  $J=9.1$ , 6.2 Hz, 1H), 1.01 (t,  $J=7.2$  Hz, 2H). LC/MS  $m/z$  435.4  $[\text{M}+\text{H}]^+$ .


#### Example 44

1-(7-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)-2,7-diazaspiro[4.4]nonan-2-yl)propan-1-one (Compound I-335)

[0979]



-continued



**[0980]** Compound I-335 was prepared using Method BF, Steps 1-3 followed by Method BV, Steps 1-4.

## Method BV

Step 1: tert-Butyl 4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate

**[0981]** To a solution of tert-butyl N-[5-bromo-3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-tert-butoxycarbonyl-carbamate (500 mg, 0.82 mmol) in toluene (4.4 mL)/water (484  $\mu$ L) was added tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (304 mg, 1.0 mmol) followed by the addition of 4-ditert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (58 mg, 0.082 mmol) and  $K_2CO_3$  (226 mg, 1.6 mmol). The reaction mixture was heated at 60° C. for 3 h. The reaction mixture was partitioned between DCM and saturated aqueous  $NaHCO_3$ , the layers were separated and the aqueous layer was extracted with DCM. The combined organics were dried over  $Na_2SO_4$ , filtered and concentrated to brown oil. The crude material was purified by silica gel column chromatography using 0-20% EtOAc in hexanes to yield tert-butyl 4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate (350 mg, 60%).  $^1H$

NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.74 (s, 1H), 8.22 (d,  $J=33.8$  Hz, 2H), 7.59 (d,  $J=8.2$  Hz, 2H), 6.92 (s, 1H), 4.54 (s, 2H), 4.24 (s, 2H), 3.73 (t,  $J=5.5$  Hz, 2H), 2.79 (d,  $J=0.8$  Hz, 2H), 1.37 (s, 18H), 1.24 (s, 9H).

## Method BV

Step 2: 3-[5-[4-(Bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine

**[0982]** tert-Butyl 4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridine-1-carboxylate (54 mg, 0.07 mmol) was treated with a solution of 4 M hydrogen chloride (376  $\mu$ L of 4 M, 1.5 mmol) in dioxane and stirred at room temperature for 1 h. The solvent was removed under reduced pressure to yield 3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-(1,2,3,6-tetrahydropyridin-4-yl)pyrazin-2-amine as an off white solid, which was used in the next step without further purification. LC/MS  $m/z$  415.5  $[M+H]^+$ .

## Method BV

Step 3: 1-[4-[5-Amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one

**[0983]** To a solution of 3-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]-5-(1,2,3,6-tetrahydropyridin-4-yl)

pyrazin-2-amine (34 mg, 0.07 mmol) in anhydrous DCM (500  $\mu$ L) cooled in an ice water bath under an atmosphere of nitrogen was added DIEA (49 mg, 66  $\mu$ L, 0.38 mmol) followed by the dropwise addition of a solution of propanoyl chloride (8 mg, 7  $\mu$ L, 0.08 mmol) in DCM (0.2 mL). The reaction mixture was stirred for 15 minutes. The reaction mixture was diluted with DCM and washed three times with 1M aqueous HCl solution, twice with saturated aqueous NaHCO<sub>3</sub> solution and then twice with a solution of brine. The layers were separated and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness under reduced pressure to yield 1-[4-[5-amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one (20 mg, 57%). LC/MS m/z 471.3 [M+H]<sup>+</sup>.

#### Method BV

Step 4: 1-[4-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one

**[0984]** A solution of 1-[4-[5-amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one (20 mg, 0.04 mmol) and a solution of methanamine (107  $\mu$ L of 2 M, 0.21 mmol) in THF

was heated at 65° C. for 6 h. The reaction mixture was allowed to cool to room temperature and the solvents were evaporated under reduced pressure. The crude material obtained was dissolved in DMF (1 mL), filtered and purified by mass directed reverse phase LC/MS (1-99% ACN/H<sub>2</sub>O (5 mM HCl)) to yield 1-[4-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-3,6-dihydro-2H-pyridin-1-yl]propan-1-one (9 mg, 45%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.28 (s, 2H), 8.58 (s, 1H), 8.18 (d, J=7.7 Hz, 2H), 7.80 (d, J=8.0 Hz, 2H), 7.63 (s, 2H), 6.66 (s, 1H), 4.30-4.13 (m, 4H), 3.70 (dt, J=16.2, 5.4 Hz, 2H), 2.71 (s, 1H), 2.59 (t, J=5.2 Hz, 4H), 2.41 (dt, J=21.8, 7.3 Hz, 2H), 1.03 (q, J=7.0 Hz, 3H). LC/MS m/z 420.5 [M+H]<sup>+</sup>.


**[0985]** Compound I-339 was prepared using the method described for Compound I-335 in Example 44 above.

**[0986]** Compound I-339: 5-(1-Ethylsulfonyl-3,6-dihydro-2H-pyridin-4-yl)-3-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-amine. LC/MS m/z 456.1 [M+H]<sup>+</sup>.

#### Example 45

3-(5-(4-((Methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)-5-(4-methylcyclohex-1-enyl)pyrazin-2-amine. (Compound I-336)

**[0987]**



[0988] Compound I-336 was prepared using Method BF, Steps 1-2 followed by Method BR, Steps 1-2, followed by Method BW, Step 1.

Method BW


Step 1: 3-(5-(4-((Methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)-5-(4-methylcyclohex-1-enyl)pyrazin-2-amine


[0989] A suspension of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (70 mg, 0.11 mmol), 4-ditert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (0.75 mg, 0.001 mmol) and  $K_2CO_3$  (29 mg, 0.2 mmol) in toluene (630  $\mu$ L) and water (70  $\mu$ L) was heated to 80° C. for 18 h. The reaction mixture was cooled to room temperature and diluted with DCM (30 ml) and water (30 ml). The layers were separated and the organic layer was washed with 1M NaOH solution, dried over ( $Na_2SO_4$ ), filtered and concentrated to dryness. The crude material obtained was stirred in 1:1 DCM:TFA mixture (3 mL) for 1 h. The solvents were evaporated under reduced pressure and the crude material was purified by reverse phase chromatography using 1 to 99% MeOH in water (0.05% HCl) to obtain 3-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)-5-(4-methylcyclohex-1-enyl)pyrazin-2-amine.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  8.33-8.21 (m, 3H), 7.79-7.68 (m, 2H), 6.65 (s, 1H), 4.32 (d, J=5.3 Hz, 2H), 3.78-3.61 (m, 2H), 3.34 (d, J=5.4 Hz, 1H), 2.78 (d, J=5.3 Hz, 3H), 2.71 (d, J=14.9 Hz, 1H), 2.42 (ddd, J=22.1, 18.2, 5.7 Hz, 2H), 2.00-1.83 (m, 2H), 1.82-1.69 (m, 1H), 1.40 (dt, J=16.6, 4.6 Hz, 1H), 1.05 (t, J=5.9 Hz, 3H). LC/MS m/z 377.0[M+H] $^+$ .

Example 46

N-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)cyclohex-3-enyl)propionamide. (Compound I-367)

[0990]





[0991] Compound I-367 was prepared using Method BF, Steps 1-2 followed by Method BR, Steps 1-2, followed by Method BV, Step 1, followed by Method BX, Steps 1-5.

#### Method BX

Step 1: tert-Butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-oxocyclohexen-1-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0992] To a solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(1,4-dioxaspiro[4.5]dec-7-en-8-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (680 mg, 0.94 mmol) in THF (2.0 mL) was added 1M aqueous HCl (1.9 mL of 1 M, 1.9 mmol) and the reaction mixture was heated at 60° C. for 6 h. The reaction was cooled to room temperature and diluted with EtOAc, the layers were separated and the organic layer was washed once with aqueous NaHCO<sub>3</sub>, then once with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to yield tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-oxocyclohexen-1-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (487 mg, 76%) and was used without further purification. LC/MS m/z 677.5 [M+H]<sup>+</sup>.

#### Method BX

Step 2: tert-Butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-hydroxycyclohexen-1-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0993] A solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-oxocyclohexen-1-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (431 mg, 0.64 mmol) in MeOH (2 mL) was cooled in an ice water bath. Sodium boranuide (27 mg, 0.70 mmol) was added and the reaction mixture was stirred for 40 minutes at 0° C. upon which the ice bath was removed. After stirring at room temperature for 8 h, the reaction mixture was quenched with water and aq. NaHCO<sub>3</sub> and diluted with EtOAc. The layers were separated and the aqueous layer was extracted once more with EtOAc. The combined organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated. The crude material was purified by silica gel column chromatography using 0-40% EtOAc in DCM to yield tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-hydroxycyclohexen-1-yl)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (126 mg, 41%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>) δ 9.04 (s, 1H), 8.11 (d, J=8.0 Hz, 2H), 7.49 (d, J=8.1 Hz, 2H), 7.01 (s, 1H), 4.81 (d, J=3.8 Hz, 1H), 4.49 (s, 2H), 3.93-3.86 (m, 1H), 2.83 (s, 5H), 2.64-2.53 (m, J=19.3 Hz, 2H), 1.97-1.90 (m, J=12.4 Hz, 1H), 1.71 (dd, J=14.0, 8.7 Hz, 1H), 1.45 (s, 3H), 1.36 (s, 6H), 1.26 (s, 18H); LC/MS m/z 579.9 [M+H]<sup>+</sup>.

#### Method BX

Step 3: [4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclohex-3-en-1-yl]methanesulfonate

[0994] To a solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-(4-hydroxycyclohexen-1-yl)

pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (124 mg, 0.18 mmol) in anhydrous DCM (2 mL) was added DMAP (2 mg, 0.02 mmol), followed by the addition of Et<sub>3</sub>N (20 mg, 28 μL, 0.20 mmol) under an atmosphere of nitrogen. The reaction mixture was cooled in an ice water bath and then treated with a solution of methanesulfonyl chloride (22 mg, 15 μL, 0.19 mmol) in DCM (0.1 mL) dropwise. The reaction mixture was stirred for 45 minutes and then partitioned between DCM/1M aqueous HCl. The layers were separated and the organic layer was washed twice with aqueous 1M HCl, once with aqueous NaHCO<sub>3</sub>, then brine solution. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness under reduced pressure to yield [4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclohex-3-en-1-yl]methanesulfonate (138 mg, 100%) and was used without any further purification in the next step. LC/MS m/z 757.5 [M+H]<sup>+</sup>.

#### Method BX

Step 4: tert-Butyl N-[[4-[5-[6-(4-azidocyclohexen-1-yl)-3-(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0995] To a solution of [4-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclohex-3-en-1-yl]methanesulfonate (138 mg, 0.18 mmol) in anhydrous DMF (1 mL) was added sodium azide (59 mg, 0.91 mmol) and the reaction mixture was heated at 70° C. under an atmosphere of nitrogen for 15 h. The reaction mixture was cooled to room temperature, quenched with water and diluted with EtOAc. An aqueous saturated brine solution was added, the layers were separated and the organic layer was washed twice more with an aqueous saturated brine solution. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated under reduced pressure to a brown oil to provide tert-butyl N-[[4-[5-[6-(4-azidocyclohexen-1-yl)-3-(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (103 mg, 94%), which was used without any further purification in the next step. LC/MS m/z 604.7 [M+H]<sup>+</sup>.

#### Method BX

Step 5: tert-Butyl N-[[4-[5-[6-(4-aminocyclohexen-1-yl)-3-(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

[0996] To a solution of tert-butyl N-[[4-[5-[6-(4-azidocyclohexen-1-yl)-3-(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (103 mg, 0.17 mmol) in anhydrous THF (1.5 mL) was added triphenylphosphine (67 mg, 0.26 mmol) and the reaction mixture was stirred at room temperature for 1 h. Water (61 μL, 3.41 mmol) was added after 1 h, and the reaction mixture was heated at 65° C. for 3 h. The reaction mixture was allowed to cool to room temperature and was partitioned between EtOAc/aqueous brine. The layers were separated and the aqueous layer was extracted once more with EtOAc. The combined organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness. The crude material was purified by silica gel column chromatography using 20% MeOH in EtOAc to yield tert-butyl N-[[4-[5-[6-(4-aminocyclohexen-1-yl)-3-

(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (21 mg, 21%). LC/MS  $m/z$  578.3  $[M+H]^+$ .

#### Method BX

Step 6: tert-Butyl N-[[4-[5-[3-(tert-butoxycarbonylamino)-6-[4-(propanoylamino)cyclohexen-1-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate

**[0997]** To a solution of tert-butyl N-[[4-[5-[6-(4-aminocyclohexen-1-yl)-3-(tert-butoxycarbonylamino)pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (20 mg, 0.03 mmol) in anhydrous DCM (200  $\mu$ L) was added  $Et_3N$  (5 mg, 7  $\mu$ L, 0.05 mmol) and the reaction mixture was cooled in an ice water bath. A solution of propanoyl chloride (3.843 mg, 3.625  $\mu$ L, 0.04154 mmol) in DCM (0.1 mL) was added dropwise to the above reaction mixture. The reaction mixture was diluted with DCM and washed once with water. The organic layer was dried over  $Na_2SO_4$ , filtered and concentrated under reduced pressure. The crude material was purified by silica gel column chromatography using 10% MeOH in EtOAc to yield tert-butyl N-[[4-[5-[3-(tert-butoxycarbonylamino)-6-[4-(propanoylamino)cyclohexen-1-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (21 mg, 96%). LC/MS  $m/z$  634.3  $[M+H]^+$ .

#### Method BX


Step 7: N-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)cyclohex-3-enyl)propionamide


**[0998]** To a solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonylamino)-6-[4-(propanoylamino)cyclohexen-1-yl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (21 mg, 0.03 mmol) in DCM (220  $\mu$ L) was added TFA (326 mg, 220  $\mu$ L, 2.9 mmol) and the reaction mixture was stirred for 40 minutes at room temperature. The reaction mixture was concentrated under reduced pressure. The residue was dissolved in DMF (1 mL) and purified by Gilson reverse phase HPLC using (5-99% ACN/ $H_2O$  (5 mM HCl)) to yield N-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)cyclohex-3-enyl)propionamide.  $^1H$  NMR (400 MHz, MeOD)  $\delta$  8.41 (s, 1H), 8.31 (d,  $J=8.4$  Hz, 2H), 7.77 (d,  $J=8.3$  Hz, 2H), 6.64 (s, 1H), 4.34 (s, 2H), 4.11-3.98 (m, 1H), 2.81 (s, 4H), 2.63 (d,  $J=17.8$  Hz, 2H), 2.26 (q,  $J=7.7$  Hz, 3H), 2.15-2.04 (m, 1H), 1.78 (dd,  $J=11.3, 5.0$  Hz, 1H), 1.18 (t,  $J=7.6$  Hz, 3H). LC/MS  $m/z$  434.2  $[M+H]^+$ .

#### Example 42

1-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-1-yl)propan-1-one (Compound I-353)

**[0999]**





**[1000]** Compound I-353 was prepared using Method BY Steps 1-8.

#### Method BY

Step 1: Methyl 3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazine-2-carboxylate

**[1001]** Methyl 3-amino-6-bromo-pyrazine-2-carboxylate (3 g, 12.93 mmol) and DMAP (158.0 mg, 1.3 mmol) were suspended in dichloromethane (45 mL) and treated dropwise with a solution of di-tert-butyl dicarbonate (11.3 g, 11.9 mmol, 51.72 mmol) in dichloromethane (10 mL). The reaction mixture was stirred for 30 minutes, then washed with 0.2 N HCl, 50% saturated sodium bicarbonate solution, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated in vacuo. The crude product was purified by silica gel column chromatography using 10-30% ethyl acetate/hexane to obtain methyl 3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazine-2-carboxylate (5.40 g, 97%) as a white solid.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.11 (s, 1H), 3.90 (s, 3H), 1.35 (s, 18H). LC/MS  $m/z$  434.5  $[\text{M}+\text{H}]^+$ .

#### Method BY

Step 2: Methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylate

**[1002]** A suspension of tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (858 mg, 2.8 mmol), 4-di-tert-butylphosphanyl-N,N-dimethyl-aniline; dichloropalladium (164 mg, 0.23 mmol),

methyl 3-[bis(tert-butoxycarbonyl)amino]-6-bromo-pyrazine-2-carboxylate (1.0 g, 2.3 mmol), and  $\text{Na}_2\text{CO}_3$  (490 mg, 4.6 mmol) under an atmosphere of nitrogen in toluene (15 mL) and water (2 mL) was heated to  $95^\circ\text{C}$ . for 2 h. The reaction mixture was diluted with ethyl acetate, washed with 1N HCl, 50% saturated sodium bicarbonate solution, and brine. The organic layer was dried over anhydrous  $\text{Na}_2\text{SO}_4$ , filtered, and concentrated. The crude product was purified via silica gel column chromatography using 10-50% ethyl acetate/hexane to provide methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylate (1.2 g, 97%) as a amber oil.  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.03 (s, 1H), 6.97 (s, 1H), 4.12 (s, 2H), 3.89 (s, 3H), 3.57 (t,  $J=5.2$  Hz, 2H), 2.61 (s, 2H), 1.43 (s, 9H), 1.34 (s, 18H). LC/MS  $m/z$  535.5  $[\text{M}+\text{H}]^+$ .

#### Method BY

Step 3: Methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-4-piperidyl)pyrazine-2-carboxylate

**[1003]** A solution of methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-3,6-dihydro-2H-pyridin-4-yl)pyrazine-2-carboxylate (700 mg, 1.3 mmol) in methanol (28 mL) was stirred with 10% Pd/C (wet) (557 mg, 0.26 mmol) at  $35^\circ\text{C}$ . under an atmosphere of hydrogen for 16 h. The reaction mixture was filtered, concentrated in vacuo, and purified via silica gel column chromatography using 10-70% ethyl acetate/hexane to provide methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-4-piperidyl)pyrazine-2-carboxylate (285 mg) as an light orange sticky solid.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  8.52 (s, 1H), 4.28 (bs, 2H),

3.97 (s, 3H), 3.06 (t, J=12.0 Hz, 1H), 2.86 (t, J=12.0 Hz, 2H), 1.98 (d, J=12.5 Hz, 2H), 1.79 (qd, J=12.6, 3.9 Hz, 2H), 1.48 (s, 9H), 1.41 (s, 18H). LC/MS m/z 537.7 [M+H]<sup>+</sup>.

#### Method BY

##### Step 4: Methyl

##### 3-amino-6-(4-piperidyl)pyrazine-2-carboxylate

**[1004]** To a solution of methyl 3-[bis(tert-butoxycarbonyl)amino]-6-(1-tert-butoxycarbonyl-4-piperidyl)pyrazine-2-carboxylate (285 mg) in DCM (7 mL) was added TFA (3 mL) dropwise. The reaction mixture was stirred at room temperature for 1 h, then diluted with acetonitrile and concentrated in vacuo. The reaction mixture was diluted with ethyl acetate, washed with 50% saturated sodium bicarbonate solution (2×20 mL), water, and brine. The product was not able to be extracted from aqueous layer with ethyl acetate, and was taken forward directly to the subsequent acylation step using Schotten-Baumen conditions LC/MS m/z 237.1 [M+H]<sup>+</sup>.

#### Method BY

##### Step 5: Methyl 3-amino-6-(1-propionylpiperidin-4-yl)pyrazine-2-carboxylate

**[1005]** To a rapidly stirring biphasic solution of methyl 3-amino-6-(4-piperidyl)pyrazine-2-carboxylate (135 mg, 0.57 mmol) in 50% saturated sodium bicarbonate and dichloromethane (20 mL) at 25° C. was added propanoyl chloride (79 mg, 75  $\mu$ L, 0.86 mmol) dropwise. The reaction was stirred at room temperature for 30 minutes. The organic layer was separated, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated in vacuo to provide methyl 3-amino-6-(1-propionylpiperidin-4-yl)pyrazine-2-carboxylate (111 mg, 66%). <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  8.25 (s, 1H), 7.19 (s, 2H), 4.51 (d, J=12.9 Hz, 1H), 3.95 (d, J=13.6 Hz, 1H), 3.84 (s, 3H), 3.08 (t, J=12.2 Hz, 1H), 2.88 (ddd, J=15.5, 7.9, 3.7 Hz, 1H), 2.61 (t, J=11.7 Hz, 1H), 2.34 (q, J=7.5 Hz, 2H), 1.80 (t, J=13.7 Hz, 2H), 1.69-1.39 (m, 2H), 1.00 (t, J=7.4 Hz, 3H). LC/MS m/z 293.5 [M+H]<sup>+</sup>.

#### Method BY

##### Step 6: 3-amino-6-(1-propionylpiperidin-4-yl)pyrazine-2-carbohydrazide

**[1006]** Methyl 3-amino-6-(1-propanoyl-4-piperidyl)pyrazine-2-carboxylate (110 mg, 0.38 mmol) was suspended in EtOH (2 mL) and hydrazine hydrate (94 mg, 92  $\mu$ L, 1.9 mmol) was added and the reaction mixture was heated at 70° C. for 1.5 h, followed by 16 h heating at 35° C. forming a thick yellow solid. The solid was isolated by filtration, washed with water (20 mL) and ethanol (40 mL). The solid was dried under high vacuum to yield 3-amino-6-(1-propionylpiperidin-4-yl)pyrazine-2-carbohydrazide (105 mg, 95%) as a light yellow solid. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.70 (s, 1H), 8.12 (s, 1H), 7.27 (bs, 2H), 4.57-4.43 (m, 3H), 3.94 (d, J=13.6 Hz, 1H), 3.33 (bs, 2H), 3.13-2.97 (m, 1H), 2.90-2.76 (m, 1H),

2.58 (t, J=11.7 Hz, 1H), 2.34 (q, J=7.4 Hz, 2H), 1.83-1.70 (m, 3H), 1.69-1.55 (m, 1H), 1.01 (t, J=7.4 Hz, 3H). LC/MS m/z 293.3 [M+H]<sup>+</sup>.

#### Method BY

##### Step 7: 1-[4-[5-Amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-1-piperidyl]propan-1-one

**[1007]** Dibromo(triphenyl)phosphorane (682 mg, 1.6 mmol) was added to a suspension of 3-amino-6-(1-propanoyl-4-piperidyl)pyrazine-2-carbohydrazide (105 mg, 0.36 mmol) and 4-(bromomethyl)benzoic acid (77 mg, 0.36 mmol) in MeCN (2 mL) at room temperature and the resulting solution was stirred for 1 h under an atmosphere of nitrogen. DIPEA (278.5 mg, 375.3  $\mu$ mol, 2.155 mmol) was added dropwise and the solution was stirred for 2 h. The reaction mixture was diluted with ethyl acetate, washed with 50% saturated sodium bicarbonate solution (2×20 mL), water, and brine. The solution was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, concentrated in vacuo to obtain 1-[4-[5-amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-1-piperidyl]propan-1-one, which was taken to the next step without further purification. LC/MS m/z 472.5 [M+H]<sup>+</sup>.

#### Method BY

##### Step 8: 1-(4-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-1-yl)propan-1-one

**[1008]** A solution of 1-[4-[5-amino-6-[5-[4-(bromomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-1-piperidyl]propan-1-one and Na<sub>2</sub>CO<sub>3</sub> (114 mg, 1.1 mmol) in tetrahydrofuran was treated with a solution of methylamine (1.3 mL of 2 M, 2.5 mmol) in THF and heated at 65° C. for 30 minutes. The reaction mixture was diluted with ethyl acetate, washed with 50% saturated sodium bicarbonate solution (2×20 mL), water, and brine. The solution was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated in vacuo. The crude material was purified via reverse phase HPLC (1-50% CH<sub>3</sub>CN/5 mM HCl) to provide 1-(4-(5-amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)piperidin-1-yl)propan-1-one.


**[1009]** <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.16 (s, 2H), 8.29 (s, 1H), 8.17 (d, J=8.1 Hz, 2H), 7.78 (d, J=8.1 Hz, 2H), 4.56 (d, J=12.5 Hz, 1H), 4.25 (t, J=5.7 Hz, 2H), 4.00 (d, J=13.3 Hz, 1H), 3.14 (t, J=12.5 Hz, 1H), 2.98 (t, J=11.9 Hz, 1H), 2.67 (t, J=11.9 Hz, 1H), 2.60 (t, J=5.2 Hz, 3H), 2.37 (q, J=7.5 Hz, 2H), 1.91 (t, J=12.1 Hz, 2H), 1.64 (ddd, J=33.0, 20.8, 11.9 Hz, 2H), 1.02 (t, J=7.4 Hz, 3H). LC/MS m/z 422.5 [M+H]<sup>+</sup>.

#### Example 43

##### (2-(5-Amino-6-(5-(4-((methylamino)methyl)phenyl)-1,3,4-oxadiazol-2-yl)pyrazin-2-yl)cyclopropyl)((R)-3-aminopyrrolidin-1-yl)methanone (Compound I-374)

**[1010]**

175



[1011] Compound I-374 was prepared using Method BF, Steps 1-2, followed by Method BQ Steps 1-2, followed by Method BN Step 1, followed by Method BZ Steps 1-4

#### Method BZ

Step 1: Ethyl 2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylate

[1012] To a solution of tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-vinyl-pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate (850 mg, 1.4 mmol) in toluene (0.2 mL) at 110° C. was added a solution of ethyl 2-diazoacetate (637 mg, 587  $\mu$ L, 5.6 mol) in toluene (0.7 mL) slowly over a period of 30 minutes. The reaction mixture showed incomplete conversion to product, when stirred at room temperature, so another 4 equivalents of ethyl 2-diazoacetate (637 mg, 587  $\mu$ L, 5.6 mmol) was added and the reaction mixture was heated at 110° C. for 20 minutes. The reaction mixture was cooled to room temperature and the solvent was evaporated under reduced pressure. The residue obtained was purified via silica gel column chromatography using 5 to 100% EtOAc in hexanes to yield ethyl 2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylate (921 mg, 1.326 mmol, 94.96%). LC/MS m/z 595.6 [M+H-Boc]<sup>+</sup>.

#### Method BZ

Step 2: 2-[5-[bis(tert-Butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylic acid

[1013] To a solution of ethyl 2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylate (825 mg, 1.2 mmol) in THF:MeOH solvent mixture (3.2 mL:1.6 mL) at room temperature was added aqueous NaOH (594  $\mu$ L of 4 M, 2.37 mmol) and reaction mixture was heated at 50° C. for 20 minutes. The reaction mixture was cooled to room temperature and diluted with EtOAc and washed with 1 M HCl solution. The layers were separated and the organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to obtain 2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylic acid. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  10.19 (s, 1H), 8.73 (s, 1H), 8.07 (d, J=8.2 Hz, 2H), 7.47 (t, J=6.7 Hz, 2H), 4.49 (s, 2H), 2.83 (s, 5H), 1.91 (s, 2H), 1.49-1.33 (m, 27H). LC/MS m/z 567.6 [M+H]<sup>+</sup>. (less 1 Boc)

#### Method BZ

Step 3: [2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropyl]-[(3R)-3-aminopyrrolidin-1-yl]methanone

[1014] To a solution of 2-[5-[bis(tert-butoxycarbonyl)amino]-6-[5-[4-[[tert-butoxycarbonyl(methyl)amino]methyl]phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropanecarboxylic acid (100 mg, 0.15 mmol) and tert-butyl

N-[(3R)-pyrrolidin-3-yl]carbamate (31 mg, 0.17 mmol) in DMF (0.6 mL) at room temperature was added HATU (57 mg, 0.15 mmol) followed by the addition of DIEA (48 mg, 65  $\mu$ L, 0.37 mmol) and reaction mixture was stirred at room temperature overnight. The reaction mixture was diluted with EtOAc and washed with 1 M HCl solution, and then with brine. The layers were separated and the organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to obtain tert-butyl N-[[4-[5-[3-[bis(tert-butoxycarbonyl)amino]-6-[2-[(3R)-3-(tert-butoxycarbonylamino)pyrrolidine-1-carbonyl]cyclopropyl]pyrazin-2-yl]-1,3,4-oxadiazol-2-yl]phenyl]methyl]-N-methyl-carbamate, which was dissolved in 50% TFA-DCM (1 mL) mixture and stirred for 25 minutes at room temperature. The reaction mixture was concentrated under reduced pressure. The crude material was dissolved in DMSO (1 mL), filtered and purified using reverse phase HPLC (10 to 99% MeOH in H<sub>2</sub>O with HCl modifier) to obtain [2-[5-amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]cyclopropyl]-[(3R)-3-aminopyrrolidin-1-yl]methanone. LC/MS m/z 435.4 [M+H]<sup>+</sup>.

[1015] The following compounds were all prepared using the method described for Compound I-374 in Example 43 above.

[1016] Compound I-375: 2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-ethyl-cyclopropanecarboxamide. LC/MS m/z 394.05 [M+H]<sup>+</sup>.

[1017] Compound I-376: 2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-(2,3-dihydroxypropyl)cyclopropanecarboxamide. LC/MS m/z 440.2 [M+H]<sup>+</sup>.

[1018] Compound I-377: 2-[5-Amino-6-[5-[4-(methylaminomethyl)phenyl]-1,3,4-oxadiazol-2-yl]pyrazin-2-yl]-N-(2-methoxyethyl)-N-methyl-cyclopropanecarboxamide. <sup>1</sup>H NMR (400 MHz, DMSO-d<sub>6</sub>)  $\delta$  9.23 (s, 2H), 8.26 (s, 0.3H, minor), 8.18 (d, J=8.1 Hz, 2H), 8.12 (d, J=5.1 Hz, 1H, minor), 7.79 (d, J=8.1 Hz, 2H), 7.47 (bs, 1H), 4.25 (t, J=5.8 Hz, 2H), 3.54-3.46 (m, 1H), 3.43 (s, 1H), 3.25 (s, 1H), 3.16 (d, J=8.6 Hz, 1H), 3.08-2.93 (m, 1H), 2.88 (s, 1H), 2.71 (dt, J=15.5, 7.6 Hz, 1H), 2.60 (dd, J=13.9, 8.7 Hz, 4H), 2.41 (dd, J=14.6, 8.1 Hz, 1H), 2.33 (s, 0.5H, minor), 2.21 (dd, J=14.3, 8.3 Hz, 1H), 1.74 (dd, J=12.0, 5.6 Hz, 1H), 1.52-1.42 (m, 1H), 1.27 (dd, J=9.0, 4.1 Hz, 1H). LC/MS m/z 438.22 [M+H]<sup>+</sup>.

[1019] Tables 7 and 8 below include some analytical data for some compounds of the instant disclosure.

TABLE 7

| Cmpd No. | LCMS ES+ | LCMS Rt (min) |
|----------|----------|---------------|
| I-174    | 378.2    | 1.09          |
| I-175    | 394.23   | 1.17          |
| I-176    | 401.18   | 0.98          |
| I-177    | 436.16   | 1.14          |
| I-178    | 440.2    | 0.97          |
| I-179    | 396.17   | 0.96          |
| I-180    | 366.16   | 0.95          |
| I-181    | 412.2    | 0.97          |
| I-182    | 368.18   | 0.95          |
| I-183    | 429.17   | 0.97          |
| I-184    | 381.21   | 0.89          |
| I-185    | 338.17   | 0.91          |
| I-186    | 378.2    | 0.88          |
| I-187    | 429.17   | 1.01          |
| I-188    | 401.18   | 0.95          |

TABLE 7-continued

| Cmpd No. | LCMS ES+ | LCMS Rt (min) |
|----------|----------|---------------|
| I-189    | 406.19   | 1.15          |
| I-190    | 394.19   | 1.10          |
| I-191    | 398.17   | 1.00          |
| I-192    | 391.18   | 1.15          |
| I-193    | 401.18   | 1.08          |
| I-194    | 420.21   | 1.21          |
| I-195    | 458.19   | 1.18          |
| I-196    | 464.14   | 1.25          |
| I-197    | 442.19   | 1.17          |
| I-198    | 408.21   | 1.2           |
| I-199    | 366.2    | 1.05          |
| I-200    | 375.19   | 1.15          |
| I-201    | 439.18   | 1.20          |
| I-202    | 500.12   | 1.25          |
| I-203    | 464.16   | 1.18          |
| I-204    | 392.21   | 1.11          |
| I-205    | 430.19   | 1.14          |
| I-206    | 389.2    | 1.21          |
| I-207    | 440.14   | 1.14          |
| I-208    | 400.18   | 1.11          |
| I-209    | 384.21   | 1.07          |
| I-210    | 433.14   | 1.25          |
| I-211    | 394.23   | 1.12          |
| I-212    | 398.22   | 1.10          |
| I-213    | 411.19   | 1.17          |
| I-214    | 377.2    | 1.19          |
| I-215    | 433.15   | 1.17          |
| I-216    | 437.2    | 1.01          |
| I-217    | 434.19   | 1.05          |
| I-218    | 395.19   | 1.04          |
| I-219    | 378.13   | 0.92          |
| I-220    | 392.18   | 1.09          |
| I-221    | 426.18   | 1.10          |
| I-222    | 462.13   | 1.14          |
| I-223    | 421.2    | 1.12          |
| I-224    | 423.17   | 1.14          |
| I-225    | 382.15   | 1.11          |
| I-226    | 418.19   | 1.16          |
| I-227    | 406.19   | 1.03          |
| I-228    | 379.14   | 1.16          |
| I-229    | 371.19   | 0.93          |
| I-230    | 412.2    | 0.89          |
| I-231    | 435.22   | 1.21          |
| I-232    | 396.17   | 1.19          |
| I-233    | 441.14   | 0.99          |

TABLE 7-continued

| Cmpd No. | LCMS ES+ | LCMS Rt (min) |
|----------|----------|---------------|
| I-234    | 364.18   | 1.06          |
| I-235    | 405.2    | 1.02          |
| I-236    | 432.21   | 1.26          |
| I-237    | 402.18   | 0.97          |
| I-238    | 393.16   | 1.24          |
| I-239    | 397.2    | 1.0           |
| I-240    | 406.16   | 1.05          |
| I-241    | 434.13   | 1.12          |
| I-242    | 358.15   | 0.98          |
| I-243    | 395.18   | 1.12          |
| I-244    | 351.15   | 1.12          |
| I-245    | 467.09   | 1.27          |
| I-246    | 420.17   | 1.22          |
| I-247    | 411.22   | 1.09          |
| I-248    | 432.17   | 1.12          |
| I-249    | 404.21   | 1.21          |
| I-250    | 365.16   | 1.19          |
| I-251    | 434.22   | 1.26          |
| I-252    | 396.17   | 1.10          |
| I-253    | 454.16   | 1.13          |
| I-254    | 389.16   | 1.23          |
| I-255    | 458.19   | 1.33          |
| I-256    | 410.22   | 1.15          |
| I-257    | 455.18   | 1.37          |
| I-258    | 456.21   | 1.24          |
| I-259    | 403.22   | 1.25          |
| I-260    | 434.19   | 1.22          |
| I-261    | 427.18   | 1.32          |
| I-262    | 432.21   | 1.13          |
| I-263    | 425.2    | 1.23          |
| I-264    | 412.2    | 0.95          |
| I-265    | 368.18   | 0.94          |
| I-266    | 398.19   | 1.05          |
| I-267    | 393.21   | 1.06          |
| I-268    | 354.16   | 1.05          |
| I-269    | 426.13   | 1.12          |
| I-270    | 407.22   | 1.14          |
| I-271    | 368.18   | 1.12          |
| I-272    | 392.18   | 1.17          |
| I-273    | 430.19   | 1.14          |
| I-274    | 430.19   | 1.27          |
| I-275    | 428.21   | 1.18          |
| I-295    | 414.2    | 3.5           |

TABLE 8

| Cmpd No. | LCMS ES+ | LCMS Rt (min) | HNMR                                                                                                                                                                                                                                                                                               |
|----------|----------|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| I-276    | 386      | 3             | H NMR (400.0 MHz, DMSO) d 1.24 (t, J = 7.4 Hz, 3H), 2.71 (br s, 2H), 3.14 (q, J = 7.3 Hz, 2H), 3.48 (t, J = 5.7 Hz, 2H), 3.98 (d, J = 2.9 Hz, 2H), 6.62 (br s, 1H), 7.30-7.33 (m, 1H), 7.72 (br s, 2H), 7.75-7.79 (m, 1H), 7.98 (d, J = 8.8 Hz, 1H), 8.40 (s, 1H) and 9.11 (d, J = 6.7 Hz, 1H) ppm |
| I-277    | 379      | 3.70          | 3.57-3.59 (m, 4H), 3/65-3.67 (m, 4H), 6.91-6.94 (m, 1H), 7.23-7.26 (m, 3H), 7.64 (br s, 2H), 8.07 (2, 1H)                                                                                                                                                                                          |
| I-278    | 433.05   | 3.38          | dmso d6 1.64 (1H, m), 2.00 (1H, m), 2.17 (6H, s), 2.60 (1H, m), 2.72 (2H, m), 3.25 (2H, m), 3.41 (3H, m), 3.61 (1H, m), 3.95 (2H, m), 6.74 (1H, s), 7.29 (2H, m), 7.62 (1H, d), 7.75 (1H, d), 8.36 (1H, s)                                                                                         |
| I-279    | 419.03   | 3.05          | dmso d6 1.85 (2H, s), 2.74 (2H, s), 2.88 (2H, s), 2.99 (2H, s), 3.40 (6H, m), 3.90 (2H, s), 6.76 (1H, s), 7.28 (2H, m), 7.62 (1H, d), 7.76 (1H, d), 8.36 (1H, s)                                                                                                                                   |
| I-280    | 405.03   | 3.05          | DMSO D6 2.73 (6H, m), 3.17 (4H, m), 3.42 (2H, m), 3.96 (2H, s), 6.75 (1H, s), 7.29 (2H, m), 7.62 (1H, d), 7.75 (1H, d), 8.36 (1H, s)                                                                                                                                                               |
| I-281    | 348      | 3.29          | MeOH 2.5 (3H, s), 4.15-4.2 (2H, m), 4.4-4.45 (2H, m), 5.05 (2H, s), 7.25 (1H, d), 7.52 (1H, s), 7.6 (1H, d), 8.2 (1H, s), 8.8 (1H, s)                                                                                                                                                              |

TABLE 8-continued

| Cmpd No. | LCMS ES+ | LCMS Rt (min) | HNMR                                                                                                                                                                                                                                                           |
|----------|----------|---------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| I-282    | 352.1    | 2.93          | CDCl <sub>3</sub> 1.98 (3H, s), 3.4 (1H, q), 3.7-3.76 (2H, m), 4.-4.45 (2H, m), 4.75-4.85 (1H, m), 6.2 (2H, s), 7.5-7.6 (3H, m), 7.7 (1H, s), 8.2-8.3 (2H, m)                                                                                                  |
| I-283    | 434.16   | 3.62          | (DMSO) 1.24 (3H, t), 3.11 (2H, q), 3.33 (4H, masked signal), 3.56 (4H, m), 6.97 (2H, br s), 7.53-7.58 (1H, m), 7.69-7.75 (1H, m), 7.86-7.90 (1H, m), 7.96-7.98 *1H, m), 8.29 (1H, s)                                                                           |
| I-284    | 361.21   | 3.62          | H NMR (400.0 MHz, DMSO) d 0.75-0.78 (m, 4H), 2.00-2.05 (m, 1H), 2.67 (s, 1H), 2.80 (s, 1H), 3.75 (s, 1H), 3.94 (s, 1H), 4.21 (s, 1H), 4.47 (s, 1H), 6.78 (s, 1H), 7.24 (s, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.37 (d, 1H) and 12.85 (s, 1H) ppm                 |
| I-285    | 389.16   | 3.67          | H NMR (400.0 MHz, DMSO) d 0.74-0.78 (m, 4H), 1.99 (br s, 0.5H), 2.09 (br s, 0.5H), 2.61 (br s, 1H), 2.68 (br s, 1H), 3.73 (br s, 1H), 3.93 (br s, 1H), 4.19 (br s, 1H), 4.47 (br s, 1H), 6.67 (br s, 1H), 7.63-7.70 (m, 5H), 8.13 (d, 2H) and 8.59 (s, 1H) ppm |
| I-286    | 333.11   | 2.97          | H NMR (400.0 MHz, DMSO) d 6.33 (d, J = 7.8 Hz, 2H), 7.69 (s, 2H), 8.15-8.18 (m, 2H), 8.35 (d, J = 7.9 Hz, 2H) and 8.78 (s, 1H) ppm                                                                                                                             |
| I-287    | 430.2    | 3.72          | (DMSO) 1.24 (3H, t), 2.70 (3H, s), 3.10 (2H, q), 3.30 (4H-hidden by water signal) 3.54 (4H, m), 6.96 (2H, br s), 7.45-7.50 (2H, m), 7.57 (1H, m), 8.03 (1H, m), 8.28 (1H, s)                                                                                   |
| I-288    | 365.1    | 3             | MeOH 1.7-1.75 (2H, m), 2.85 (2H, t), 3.35-3.4 (2H, m), 6.25 (1H, brs), 6.65 (1H, d), 6.7-6.8 (4H, m), 7.25-7.4 (4H, m),                                                                                                                                        |
| I-289    | 452.12   | 3.45          | (DMSO) 1.23 (3H, t), 3.10 (2H, q), 3.32 (4H, m), 3.52 (4H, m), 6.98 (2H, br s), 7.45 (1H, m), 7.77-7.85 (1H, m), 8.31 (1H, s)                                                                                                                                  |
| I-290    | 404.18   | 3.64          | MeOH 1.95-2.0 (2H, m), 3.0 (2H, t), 3.35-3.4 (2H, m), 5.95-6.0 (1H, m), 6.65 (1H, d), 6.7-6.8 (4H, m), 7.25-7.35 (4H, m),                                                                                                                                      |
| I-291    | 399.14   | 3.62          | H NMR (400.0 MHz, DMSO) d 0.99-1.03 (m, 3H), 1.69-1.79 (m, 2H), 2.80 (br s, 2H), 3.10-3.20 (m, 2H), 3.49-3.50 (m, 2H), 3.95 (s, 2H), 6.79 (s, 1H), 7.24-7.34 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.37 (s, 1H) and 12.85 (s, 1H) ppm                           |
| I-292    | 434.12   | 3.59          | (DMSO) 1.24 (3H, m), 3.11 (2H, t), 3.56 (4H, m), 6.96 (2H, br s), 7.50 (2H, m), 8.18 (2H, m), 8.28 (1H, s) 4H of piperazine hidden by water peak                                                                                                               |
| I-293    | 363.22   | 3.7           | H NMR (400.0 MHz, DMSO) d 1.04-1.07 (m, 6H), 2.67 (br s, 1H), 2.73 (br s, 1H), 2.89-3.04 (m, 1H), 3.75 (s, 2H), 4.20 (s, 1H), 4.30 (s, 1H), 6.76 (br s, 1H), 7.24-7.33 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.36 (s, 1H) and 12.84 (s, 1H) ppm                 |
| I-294    | 436.11   | 3.6           | (DMSO) 1.23 (3H, t), 2.63 (3H, s), 3.10 (2H, q), 3.30 (4H, masked signal with water peak), 3.54 (4H, m), 6.92 (2H, br s), 7.19 (1H, d), 7.87 (1H, d), 8.27 (1H, s)                                                                                             |
| I-296    |          |               | H NMR (400.0 MHz, DMSO) d 2.60 (m, 5H), 2.77 (m, 2H), 3.06 (s, 3H), 3.31 (m, 2H), 3.36 masked signal, 3.56 (m, 4H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H), 8.25 (s, 1H)                                                                             |
| I-297    | 406.15   | 3.27          | (DMSO) 1.24 (3H, t), 3.11 (2H, q), 3.30 (4H, masked signal), 3.52 (4H, m), 6.85 (1H, m), 6.95 (2H, br s), 7.47 (1H, m), 8.12 (1H, m), 8.27 (1H, s)                                                                                                             |
| I-298    |          |               | H NMR (400.0 MHz, DMSO) d 2.60-2.66 (m, 6H), 2.75 (m, 2H), 3.46 (m, 4H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H), 8.24 (s, 1H)                                                                                                                        |
| I-299    | 434.15   | 3.48          | (DMSO) 1.24 (3H, t), 3.11 (2H, t), 3.30 (4H, masked signal), 3.53 (4H, m), 6.97 (2H, br s), 7.47-7.56 (2H, m), 7.75 (1H, m), 8.11 (1H, m), 8.29 (1H, s)                                                                                                        |
| I-300    |          |               | H NMR (400.0 MHz, DMSO) d 3.31 masked signal, 3.46 (m, 4H) 3.55 (m, 2H), 6.88 (br s, 2H), 7.27 (m, 1H), 7.36 (m, 4H), 7.64-7.68 (m, 3H), 8.11 (m, 2H), 8.23 (s, 1H)                                                                                            |
| I-301    | 377.17   | 3.59          | H NMR (400.0 MHz, DMSO) d 1.04 (t, 3H), 2.36-2.45 (m, 2H), 2.60 (br s, 1H), 2.70 (br s, 1H), 3.65-3.72 (m, 2H), 4.17 (s, 1H), 4.21 (s, 1H), 6.65 (s, 1H), 7.62-7.69 (m, 5H), 8.11-8.13 (d, 2H) and 8.56 (s, 1H) ppm                                            |
| I-302    | 402.2    | 3.17          | DMSO 1.3 (3H, t), 3.1 (2H, q), 3.8-3.85 (2H, m), 4.35-4.42 (3H, m), 7.0 (2H, m), 7.7-7.8 (3H, m), 7.8-7.85 (2H, m), 8.05-8.1 (2H, m)                                                                                                                           |
| I-303    |          |               | H NMR (400.0 MHz, DMSO) d 2.59 (m, 4H), 3.24 (s, 2H), 3.46 (m, 6H) 3.56 (m, 6H), 6.90 (br s, 2H), 7.64-7.68 (m, 3H), 8.11 (m, 2H), 8.23 (s, 1H)                                                                                                                |
| I-304    | 343.1    | 3.6           | MeOH 5.25 (2H, s), 7.3-7.35 (2H, m), 7.55-7.6 (1H, m), 7.63-7.7 (4H, m), 7.9 (1H, d), 9.22 (1H, s),                                                                                                                                                            |

TABLE 8-continued

| Cmpd No. | LCMS ES+ | LCMS Rt (min) | HNMR                                                                                                                                                                                                                                                                      |
|----------|----------|---------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| I-305    | 442.1    | 3.42          | DMSO 1.25 (3H, t), 3.1-3.2 (4H, m), 3.25-3.3 (2H, m), 3.4-3.45 (2H, m), 3.5-3.7 (4H, m), 6.75 (2H, s), 7.6-7.65 (3H, m), 8.05-8.1 (2H, m)                                                                                                                                 |
| I-306    |          |               | H NMR (400.0 MHz, DMSO) d 2.46 (m, 2H), 2.59 (m, 4H), 3.46 (m, 4H), 3.56 (m, 2H), 4.47 (t, 1H), 6.90 (br s, 2H), 7.64-7.69 (m, 3H), 8.11 (m, 2H), 8.24 (s, 1H)                                                                                                            |
| I-307    | 412.97   | 3.55          | (DMSO) 1.24 (3H, t), 2.68 (2H, m), 3.13 (2H, q), 3.48 (2H, m), 3.99 (2H, m), 6.63 (1H, m), 7.29 (2H, br s), 7.68 (2H, m), 7.77 (1H, m), 8.26 (2H, m), 8.52 (1H, s)                                                                                                        |
| I-308    | 464      | 3.54          | H NMR (400.0 MHz, CDCl <sub>3</sub> ) d 7.83 (d, J = 8.0 Hz, 1H), 7.64 (s, 1H), 7.54-7.43 (m, 3H), 7.35-7.32 (m, 3H), 7.28-7.24 (m, 3H), 7.04 (d, J = 8.0 Hz, 1H), 3.14-3.11 (m, 4H), 2.93 (q, J = 7.4 Hz, 2H), 2.76 (t, J = 5.0 Hz, 4H) and 1.37 (t, J = 7.4 Hz, 3H) ppm |
| I-309    | 349.19   | 3.57          | H NMR (400.0 MHz, DMSO) d 1.01-1.06 (m, 3H), 2.33-2.47 (m, 2H), 2.67 (s, 1H), 2.76 (s, 1H), 3.67-3.75 (m, 2H), 4.19-4.23 (m, 2H), 6.76 (d, 1H), 7.24-7.33 (m, 2H), 7.62 (d, 1H), 7.76 (d, 1H), 8.36 (d, 1H) and 12.84 (s, 1H) ppm                                         |
| I-310    | 305.1    | 3.09          | DMSO 6.45 (1H, t), 6.55 (1H, d), 7.3-7.35 (2H, m), 7.5-7.55 (1H, m), 7.6-7.7 (2H, brs), 8.05-8.1 (1H, m), 8.55 (1H, s), 13.15 (1H, vbrs)                                                                                                                                  |
| I-311    | 305.15   | 2.97          | H NMR (400.0 MHz, DMSO) d 6.34 (d, J = 7.8 Hz, 2H), 7.26-7.37 (m, 2H), 7.62 (d, 1H), 7.80 (d, 1H), 8.56 (d, J = 7.8 Hz, 2H), 8.64 (s, 1H) and 13.17 (s, 1H) ppm                                                                                                           |
| I-312    | 422.1    | 3.45          | (DMSO) 1.24 (3H, t), 3.11 (2H, q), 3.34 (4H, m), 3.55 (4H, m), 6.93 (2H, br s), 7.35 (1H, m), 7.94 (1H, dd), 8.01 (1H, dd), 8.27 (1H, s)                                                                                                                                  |

## Examples 32

## Cellular ATR Inhibition Assay

**[1020]** Compounds were screened for their ability to inhibit intracellular ATR using an immunofluorescence microscopy assay to detect phosphorylation of the ATR substrate histone H2AX in hydroxyurea treated cells. HT29 cells were plated at 14,000 cells per well in 96-well black imaging plates (BD 353219) in McCoy's 5A media (Sigma M8403) supplemented with 10% foetal bovine serum (JRH Biosciences 12003), Penicillin/Streptomycin solution diluted 1:100 (Sigma P7539), and 2 mM L-glutamine (Sigma G7513), and allowed to adhere overnight at 37° C. in 5% CO<sub>2</sub>. Compounds were then added to the cell media from a final concentration of 25  $\mu$ M in 3-fold serial dilutions and the cells were incubated at 37° C. in 5% CO<sub>2</sub>. After 15 min, hydroxyurea (Sigma H8627) was added to a final concentration of 2 mM.

**[1021]** After 45 min of treatment with hydroxyurea, the cells were washed in PBS, fixed for 10 min in 4% formaldehyde diluted in PBS (Polysciences Inc 18814), washed in 0.2% Tween-20 in PBS (wash buffer), and permeabilised for 10 min in 0.5% Triton X-100 in PBS, all at room temperature. The cells were then washed once in wash buffer and blocked for 30 min at room temperature in 10% goat serum (Sigma G9023) diluted in wash buffer (block buffer). To detect H2AX phosphorylation levels, the cells were then incubated for 1 h at room temperature in primary antibody (mouse monoclonal anti-phosphorylated histone H2AX Ser139 antibody; Upstate 05-636) diluted 1:250 in block buffer. The cells were then washed five times in wash buffer before incubation for 1 h at room temperature in the dark in a mixture of secondary antibody (goat anti-mouse Alexa Fluor 488 conjugated antibody; Invitrogen A11029) and Hoechst stain (Invitrogen H3570); diluted 1:500 and 1:5000, respectively, in

wash buffer. The cells were then washed five times in wash buffer and finally 100  $\mu$ l PBS was added to each well before imaging.

**[1022]** Cells were imaged for Alexa Fluor 488 and Hoechst intensity using the BD Pathway 855 Bioimager and Attovision software (BD Biosciences, Version 1.6/855) to quantify phosphorylated H2AX Ser139 and DNA staining, respectively. The percentage of phosphorylated H2AX-positive nuclei in a montage of 9 images at 20 $\times$  magnification was then calculated for each well using BD Image Data Explorer software (BD Biosciences Version 2.2.15). Phosphorylated H2AX-positive nuclei were defined as Hoechst-positive regions of interest containing Alexa Fluor 488 intensity at 1.75-fold the average Alexa Fluor 488 intensity in cells not treated with hydroxyurea. The percentage of H2AX positive nuclei was finally plotted against concentration for each compound and IC50s for intracellular ATR inhibition were determined using Prism software (GraphPad Prism version 3.0cx for Macintosh, GraphPad Software, San Diego Calif., USA). Compound I-27 was found to have an IC50 value of <1  $\mu$ M. Compounds I-24 and I-26 were found to have an IC50 value of  $\geq$ 1  $\mu$ M but <5  $\mu$ M. Compounds I-25 and I-28 were found to have an IC50 value of  $\geq$ 5  $\mu$ M but <10  $\mu$ M. Compound I-16 was found to have an IC50 value of  $\geq$ 10  $\mu$ M but <30  $\mu$ M.

**[1023]** The compounds described herein can also be tested according to other methods known in the art (see Sarkaria et al, "Inhibition of ATM and ATR Kinase Activities by the Radiosensitizing Agent, Caffeine: *Cancer Research* 59: 4375-5382 (1999); Hickson et al, "Identification and Characterization of a Novel and Specific Inhibitor of the Ataxia-Telangiectasia Mutated Kinase ATM" *Cancer Research* 64: 9152-9159 (2004); Kim et al, "Substrate Specificities and Identification of Putative Substrates of ATM Kinase Family Members" *The Journal of Biological Chemistry*, 274(53): 37538-37543 (1999); and Chiang et al, "Determination of the

catalytic activities of mTOR and other members of the phosphoinositide-3-kinase-related kinase family” *Methods Mol. Biol.* 281:125-41 (2004)).

### Example 33

#### ATR Inhibition Assay

**[1024]** Test compounds were screened for their ability to inhibit ATR kinase using a radioactive-phosphate incorporation assay. Assays were carried out in a mixture of 50 mM Tris/HCl (pH 7.5), 10 mM MgCl<sub>2</sub> and 1 mM DTT. Final substrate concentrations were 10  $\mu$ M [ $\gamma$ -33P]ATP (3 mCi 33P ATP/mmol ATP, Perkin Elmer) and 800  $\mu$ M target peptide (ASELPASQPQPFSAKKK). Assays were carried out at 25° C. in the presence of 5 nM full-length ATR. An assay stock buffer solution was prepared containing all of the reagents listed above, with the exception of ATP and the test compound of interest. 13.5  $\mu$ L of the stock solution was placed in a 96 well plate followed by addition of 2  $\mu$ L of DMSO stock containing serial dilutions of the test compound (typically starting from a final concentration of 15  $\mu$ M with 3-fold serial dilutions) in duplicate (final DMSO concentration 7%). The plate was pre-incubated for 10 minutes at 25° C. and the reaction initiated by addition of 15  $\mu$ L [ $\gamma$ -33P]ATP (final concentration 10  $\mu$ M).

**[1025]** The reaction was stopped after 24 hours by the addition of 30  $\mu$ L 0.1M phosphoric acid containing 2 mM ATP. A multiscreen phosphocellulose filter 96-well plate (Millipore, Cat no. MAPHN0B50) was pretreated with 100  $\mu$ L 0.2M phosphoric acid prior to the addition of 454 of the stopped assay mixture. The plate was washed with 5 $\times$ 2004, 0.2M phosphoric acid. After drying, 100  $\mu$ L Optiphase ‘SuperMix’ liquid scintillation cocktail (Perkin Elmer) was added to the well prior to scintillation counting (1450 Microbeta Liquid Scintillation Counter, Wallac).

**[1026]** After removing mean background values for all of the data points, Ki(app) data were calculated from non-linear regression analysis of the initial rate data using the Prism software package (GraphPad Prism version 3.0cx for Macintosh, GraphPad Software, San Diego Calif., USA).

**[1027]** Below is a chart showing the Ki value of compounds of the disclosure. Compounds with a Ki value of  $\leq$ 100 nM are marked with “+++.” Compounds with a Ki value  $>$ 100 nM but  $\leq$ 1  $\mu$ M are marked with “++.” Compounds with a Ki value  $>$ 1  $\mu$ M but  $<$ 10  $\mu$ M are marked with “+.”

| Cmpd No. | ATR Ki |
|----------|--------|
| I-1      | +      |
| I-2      | +      |
| I-3      | ++     |
| I-4      | ++     |
| I-5      | +      |
| I-6      | ++     |
| I-7      | ++     |
| I-8      | ++     |
| I-9      | +      |
| I-10     | ++     |
| I-11     | +      |
| I-12     | +      |
| I-13     | +      |
| I-14     | +      |
| I-15     | ++     |
| I-16     | ++     |

-continued

| Cmpd No. | ATR Ki |
|----------|--------|
| I-17     | ++     |
| I-18     | +++    |
| I-19     | ++     |
| I-20     | ++     |
| I-21     | +      |
| I-22     | ++     |
| I-23     | +++    |
| I-24     | +++    |
| I-25     | +++    |
| I-26     | +++    |
| I-27     | +++    |
| I-28     | +++    |
| I-29     | ++     |
| I-30     | ++     |
| I-31     | +      |
| I-32     | +      |
| I-33     | +      |
| I-34     | +      |
| I-35     | +      |
| I-36     | ++     |
| I-37     | +      |
| I-38     | +      |
| I-39     | +      |
| I-40     | +      |
| I-41     | ++     |
| I-42     | +      |
| I-43     | ++     |
| I-44     | +      |
| I-45     | +      |
| I-46     | ++     |
| I-47     | +      |
| I-48     | ++     |
| I-49     | +      |
| I-50     | +      |
| I-51     | +      |
| I-52     | +      |
| I-53     | ++     |
| I-54     | +++    |
| I-55     | ++     |
| I-56     | ++     |
| I-57     | ++     |
| I-58     | +++    |
| I-59     | ++     |
| I-60     | +++    |
| I-61     | +      |
| I-62     | +      |
| I-63     | ++     |
| I-64     | +      |
| I-65     | +      |
| I-66     | ++     |
| I-67     | +++    |
| I-68     | ++     |
| I-69     | ++     |
| I-70     | +      |
| I-71     | +++    |
| I-72     | +      |
| I-73     | +      |
| I-74     | ++     |
| I-75     | +      |
| I-76     | ++     |
| I-77     | +      |
| I-78     | ++     |
| I-79     | +++    |
| I-80     | ++     |
| I-81     | +      |
| I-82     | +++    |
| I-83     | ++     |
| I-84     | +++    |
| I-85     | +      |
| I-86     | ++     |
| I-87     | ++     |
| I-88     | ++     |
| I-89     | +++    |
| I-90     | ++     |

| -continued  |           | -continued  |           |
|-------------|-----------|-------------|-----------|
| Cmpd<br>No. | ATR<br>Ki | Cmpd<br>No. | ATR<br>Ki |
| I-91        | ++        | I-165       | +++       |
| I-92        | ++        | I-166       | ++        |
| I-93        | +++       | I-167       | +++       |
| I-94        | +++       | I-168       | ++        |
| I-95        | ++        | I-169       | +++       |
| I-96        | ++        | I-170       | ++        |
| I-97        | ++        | I-171       | ++        |
| I-98        | ++        | I-172       | +++       |
| I-99        | ++        | I-173       | ++        |
| I-100       | +++       | I-174       | +         |
| I-101       | ++        | I-175       | +         |
| I-102       | +++       | I-176       | ++        |
| I-103       | +++       | I-177       | ++        |
| I-104       | ++        | I-178       | ++        |
| I-105       | +++       | I-179       | ++        |
| I-106       | ++        | I-180       | ++        |
| I-107       | ++        | I-181       | +         |
| I-108       | ++        | I-182       | +         |
| I-109       | +         | I-183       | ++        |
| I-110       | ++        | I-184       | +         |
| I-111       | ++        | I-185       | ++        |
| I-112       | ++        | I-186       | ++        |
| I-113       | +         | I-187       | ++        |
| I-114       | +         | I-188       | ++        |
| I-115       | +++       | I-189       | ++        |
| I-116       | ++        | I-190       | ++        |
| I-117       | ++        | I-191       | ++        |
| I-118       | ++        | I-192       | +++       |
| I-119       | ++        | I-193       | ++        |
| I-120       | +++       | I-194       | ++        |
| I-121       | +++       | I-195       | +         |
| I-122       | +++       | I-196       | ++        |
| I-123       | +++       | I-197       | ++        |
| I-124       | +++       | I-198       | ++        |
| I-125       | +++       | I-199       | +         |
| I-126       | +++       | I-200       | +++       |
| I-127       | +++       | I-201       | +++       |
| I-128       | +++       | I-202       | ++        |
| I-129       | +++       | I-203       | ++        |
| I-130       | +++       | I-204       | ++        |
| I-131       | +++       | I-205       | ++        |
| I-132       | +++       | I-206       | ++        |
| I-133       | +++       | I-207       | ++        |
| I-134       | ++        | I-208       | ++        |
| I-135       | +++       | I-209       | +         |
| I-136       | +++       | I-210       | +++       |
| I-137       | ++        | I-211       | +         |
| I-138       | +++       | I-212       | +         |
| I-139       | +++       | I-213       | ++        |
| I-140       | ++        | I-214       | ++        |
| I-141       | +         | I-215       | +++       |
| I-142       | +         | I-216       | ++        |
| I-143       | +         | I-217       | +++       |
| I-144       | ++        | I-218       | ++        |
| I-145       | ++        | I-219       | ++        |
| I-146       | +         | I-220       | +++       |
| I-147       | +         | I-221       | ++        |
| I-148       | +         | I-222       | +++       |
| I-149       | ++        | I-223       | ++        |
| I-150       | +         | I-224       | +++       |
| I-151       | +         | I-225       | ++        |
| I-152       | ++        | I-226       | +++       |
| I-153       | +++       | I-227       | ++        |
| I-154       | ++        | I-228       | +++       |
| I-155       | +++       | I-229       | ++        |
| I-156       | +         | I-230       | +         |
| I-157       | +++       | I-231       | ++        |
| I-158       | ++        | I-232       | ++        |
| I-159       | ++        | I-233       | ++        |
| I-160       | ++        | I-234       | ++        |
| I-161       | ++        | I-235       | ++        |
| I-162       | ++        | I-236       | +++       |
| I-163       | ++        | I-237       | +         |
| I-164       | +++       | I-238       | +++       |


| -continued  |           | -continued  |           |
|-------------|-----------|-------------|-----------|
| Cmpd<br>No. | ATR<br>Ki | Cmpd<br>No. | ATR<br>Ki |
| I-239       | +         | I-313       | ++        |
| I-240       | ++        | I-314       | ++        |
| I-241       | +++       | I-315       | ++        |
| I-242       | +         | I-316       | +++       |
| I-243       | ++        | I-317       | +++       |
| I-244       | ++        | I-318       | +++       |
| I-245       | +++       | I-319       | ++        |
| I-246       | ++        | I-320       | +++       |
| I-247       | +         | I-321       | +         |
| I-248       | ++        | I-322       | ++        |
| I-249       | ++        | I-323       | +++       |
| I-250       | ++        | I-324       | ++        |
| I-251       | ++        | I-325       | +++       |
| I-252       | +         | I-326       | +++       |
| I-253       | ++        | I-327       | +++       |
| I-254       | ++        | I-328       | ++        |
| I-255       | +         | I-329       | +         |
| I-256       | ++        | I-330       | +         |
| I-257       | ++        | I-331       | +++       |
| I-258       | ++        | I-332       | +         |
| I-259       | +++       | I-333       | ++        |
| I-260       | +         | I-334       | ++        |
| I-261       | ++        | I-335       | ++        |
| I-262       | ++        | I-336       | ++        |
| I-263       | +++       | I-337       | ++        |
| I-264       | +         | I-338       | ++        |
| I-265       | +         | I-339       | ++        |
| I-266       | +         | I-340       | +         |
| I-267       | +         | I-341       | ++        |
| I-268       | +         | I-342       | ++        |
| I-269       | ++        | I-343       | +++       |
| I-270       | +         | I-344       | +++       |
| I-271       | +         | I-345       | +++       |
| I-272       | +         | I-346       | ++        |
| I-273       | +         | I-347       | ++        |
| I-274       | +         | I-348       | +         |
| I-275       | +         | I-349       | ++        |
| I-276       | +         | I-350       | ++        |
| I-277       | ++        | I-351       | ++        |
| I-278       | ++        | I-352       | ++        |
| I-279       | ++        | I-353       | ++        |
| I-280       | ++        | I-354       | ++        |
| I-281       | +         | I-355       | ++        |
| I-282       | +         | I-356       | ++        |
| I-283       | ++        | I-357       | ++        |
| I-284       | +++       | I-358       | ++        |
| I-285       | +++       | I-359       | ++        |
| I-286       | +++       | I-360       | ++        |
| I-287       | ++        | I-361       | ++        |
| I-288       | ++        | I-362       | ++        |
| I-289       | ++        | I-363       | ++        |
| I-290       | ++        | I-364       | ++        |
| I-291       | +++       | I-365       | ++        |
| I-292       | ++        | I-366       | ++        |
| I-293       | ++        | I-367       | +++       |
| I-294       | +++       | I-368       | ++        |
| I-295       | +         | I-369       | +++       |
| I-296       | ++        | I-370       | ++        |
| I-297       | ++        | I-371       | ++        |
| I-298       | +         | I-372       | ++        |
| I-299       | ++        | I-373       | ++        |
| I-300       | +         | I-374       | +         |
| I-301       | +++       | I-375       | ++        |
| I-302       | ++        | I-376       | ++        |
| I-303       | ++        | I-377       | +         |
| I-304       | ++        |             |           |
| I-305       | ++        |             |           |
| I-306       | +         |             |           |
| I-307       | ++        |             |           |
| I-308       | ++        |             |           |
| I-309       | +++       |             |           |
| I-310       | +++       |             |           |
| I-311       | +++       |             |           |
| I-312       | ++        |             |           |

[1028] While we have described a number of embodiments of this invention, it is apparent that our basic examples may be altered to provide other embodiments that utilize the compounds, methods, and processes of this invention. Therefore, it will be appreciated that the scope of this invention is to be

defined by the appended claims rather than by the specific embodiments that have been represented by way of example herein.

1-71. (canceled)

72. A method of treating a disease selected from cancer, HIV, hepatitis, adenovirus, or psoriasis comprising administering to a patient a compound of formula I,



or a pharmaceutically acceptable salt thereof; wherein:

R<sup>1</sup> is a 5-6 membered monocyclic aryl or heteroaryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein said monocyclic aryl or heteroaryl ring is optionally fused to another ring to form a 8-10 membered bicyclic aryl or heteroaryl ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; R<sup>1</sup> is optionally substituted with 1-5 J<sup>1</sup> groups;

C<sup>2</sup>, C<sup>3</sup> and C<sup>5</sup> are carbon;

R<sup>2</sup> is -Q or -Q-Q<sup>1</sup>;

Q is a 3-7 membered monocyclic saturated or unsaturated non-aromatic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q is independently and optionally substituted with 1-4 J<sup>Q</sup> groups; Q is substituted with zero to two occurrences of J;

Q is optionally fused to Q<sup>1</sup> to form a fused bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup> are optionally joined together at a carbon atom to form a spirocyclic bicyclic ring Q-Q<sup>1</sup>; or Q and Q<sup>1</sup>, taken together, form a bridged bicyclic ring Q-Q<sup>1</sup> wherein said bridge is 1-3 atoms long; Q-Q<sup>1</sup> can be optionally substituted with 1-2 J groups; wherein J can be bonded to either Q or Q<sup>1</sup>;

Q<sup>1</sup> is a 3-8 membered monocyclic saturated or unsaturated ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>1</sup> is independently and optionally substituted with 1-4 J<sup>Q1</sup> groups;

J is halo, oxo, -CN, -NO<sub>2</sub>, V<sup>1</sup>-R<sup>1</sup>, or -(V)<sub>m</sub>-R<sup>\*</sup>;

L is -C(O)NH- or -C(O)N(C<sub>1-6</sub>alkyl)-;

n is 0 or 1;

each J<sup>Q</sup> and J<sup>Q1</sup> is independently halo, oxo, -CN, -NO<sub>2</sub>, V-R, or -(V)<sub>m</sub>-Q<sup>2</sup>;

J<sup>1</sup> is halo, -CN, -NO<sub>2</sub>, V-R, or -(V<sup>2</sup>)<sub>m</sub>-Q<sup>3</sup>;

V is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally and independently replaced with oxygen, nitrogen, sulfur, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of J<sup>V</sup>;

V<sup>1</sup> is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally replaced with -NR<sup>1</sup>-, -O-, -S-, C(O), S(O), or S(O)<sub>2</sub>; V<sup>1</sup> is optionally substituted with 1-6 occurrences of J<sup>V1</sup>;

m is 0 or 1;

V<sup>2</sup> is a C<sub>1-10</sub>aliphatic chain wherein 0-3 methylene units are optionally and independently replaced with oxygen,

nitrogen, sulfur, C(O), S(O), or S(O)<sub>2</sub>; V is optionally substituted with 1-6 occurrences of J<sup>V2</sup>;

m is 0 or 1;

Q<sup>2</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>2</sup> is optionally substituted with 1-5 J<sup>Q2</sup>;

Q<sup>3</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>3</sup> is optionally substituted with 1-5 J<sup>Q3</sup>;

each J<sup>V</sup> and J<sup>V2</sup> is independently halogen, CN, NH<sub>2</sub>, NO<sub>2</sub>, C<sub>1-4</sub>aliphatic, NH(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)<sub>7</sub>, OH, O(C<sub>1-4</sub>aliphatic), CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-4</sub>aliphatic), C(O)NH<sub>2</sub>, C(O)NH(C<sub>1-4</sub>aliphatic), C(O)N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, NHCO(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)CO(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), NHSO<sub>2</sub>(C<sub>1-4</sub>aliphatic), or N(C<sub>1-4</sub>aliphatic)SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), wherein said C<sub>1-4</sub>aliphatic is optionally substituted with halo;

each J<sup>1</sup>, J<sup>Q2</sup> and J<sup>Q3</sup> is independently halo, oxo, CH, NO<sub>2</sub>, X-R, or -(X)<sub>n</sub>-Q<sup>4</sup>,

p is 0 or 1;

X is C<sub>1-10</sub>aliphatic; wherein 1-3 methylene units of said C<sub>1-6</sub>aliphatic are optionally replaced with -NR, -O-, -S-, C(O), S(O)<sub>2</sub>, or S(O); wherein X is optionally and independently substituted with 1-4 occurrences of NH<sub>2</sub>, NH(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, halogen, C<sub>1-4</sub>aliphatic, OH, O(C<sub>1-4</sub>aliphatic), NO<sub>2</sub>, CN, CO<sub>2</sub>H, CO<sub>2</sub>(C<sub>1-4</sub>aliphatic), C(O)NH<sub>2</sub>, C(O)NH(C<sub>1-4</sub>aliphatic), C(O)N(C<sub>1-4</sub>aliphatic)<sub>2</sub>, SO(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>NH(C<sub>1-4</sub>aliphatic), SO<sub>2</sub>NH(C<sub>1-4</sub>aliphatic), NHC(O)(C<sub>1-4</sub>aliphatic), N(C<sub>1-4</sub>aliphatic)C(O)(C<sub>1-4</sub>aliphatic), wherein said C<sub>1-4</sub>aliphatic is optionally substituted with 1-3 occurrences of halo;

Q<sup>4</sup> is a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each Q<sup>4</sup> is optionally substituted with 1-5 J<sup>Q4</sup>;

J<sup>Q4</sup> is halo, CN, or C<sub>1-4</sub>alkyl wherein up to 2 methylene units are optionally replaced with O, N, S, C(O), S(O), or S(O)<sub>2</sub>;

each R<sup>\*</sup>, R<sup>1</sup>, and R is independently H or C<sub>1-4</sub>alkyl wherein said C<sub>1-4</sub>alkyl is optionally substituted with 1-4 halo;

R<sup>1</sup> is a H or a 3-8 membered saturated or unsaturated monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or a 8-10 membered saturated or unsaturated bicyclic ring having 0-6 heteroatoms independently selected from nitrogen, oxygen, or sulfur; each R<sup>1</sup> is optionally substituted with 1-5 J<sup>1</sup>.

73. The method of claim 72, wherein said disease is cancer.

74. The method of claim 73, comprising administering to said patient an additional therapeutic agent selected from a DNA-damaging agent; wherein said additional therapeutic

agent is appropriate for the disease being treated; and said additional therapeutic agent is administered together with said compound as a single dosage form or separately from said compound as part of a multiple dosage form.

**75.** The method of claim **74**, wherein said DNA-damaging agent is selected from ionizing radiation, radiomimetic neocarzinostatin, a platinating agent, a Topo I inhibitor, a Topo II inhibitor, an antimetabolite, an alkylating agent, an alkyl sulphonates, an antimetabolite, or an antibiotic.

**76.** The method of claim **75**, wherein said DNA-damaging agent is selected from ionizing radiation, a platinating agent, a Topo I inhibitor, a Topo II inhibitor, or an antimetabolite.

77. The method of claim 75, wherein said platinating agent is selected from Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, Lobaplatin, Triplatin Tetranitrate, Picoplatin, Satraplatin, ProLindac and Aroplatin; said Topo I inhibitor is selected from Camptothecin, Topotecan, Irinotecan/SN38, Rubitecan and Belotecan; said Topo II inhibitor is selected from Etoposide, Daunorubicin, Doxorubicin, Aclarubicin, Epirubicin, Idarubicin, Amrubicin, Pirarubicin, Valrubicin, Zorubicin and Teniposide; said antimetabolite is selected from Aminopterin, Methotrexate, Pemetrexed, Raltitrexed, Pentostatin, Cladribine, Clofarabine, Fludarabine, Thioguanine, Mercaptopurine, Fluorouracil, Capecitabine, Tegafur, Carmofur, Floxuridine, Cytarabine, Gemcitabine, Azacitidine and Hydroxyurea; said alkylating agent is selected from Mechlorethamine, Cyclophosphamide, Ifosfamide, Trofosfamide, Chlorambucil, Melphalan, Prednimustine, Bendamustine, Uramustine, Estramustine, Carmustine, Lomustine, Semustine, Fotemustine, Nimustine, Ranimustine, Streptozocin, Busulfan, Mannosulfan, Treosulfan, Carboquone, ThioTEPA, Triaziquone, Triethylenemelamine, Procarbazine, Dacarbazine, Temozolomide, Altretamine, Mitobranitol, Actinomycin, Bleomycin, Mitomycin and Plicamycin.

78. The method of claim 75, wherein said platinating agent is selected from Cisplatin, Oxaliplatin, Carboplatin, Nedaplatin, or Satraplatin; said Topo I inhibitor is selected from Camptothecin, Topotecan, irinotecan/SN38, rubitecan; said Topo II inhibitor is selected from Etoposide; said antimetabolite is selected from methotrexate, pemetrexed, Thioguanine, Fludarabine, Cladribine, Cytarabine, gemcitabine, 6-Mercaptopurine, or 5-Fluorouracil; said alkylating agent is selected from nitrogen mustards, nitrosoureas, triazenes, alkyl sulfonates, Procarbazine, or aziridines; and said antibiotic is selected from Hydroxyurea, Anthracyclines, Anthracenediones, or *Streptomyces* family.


**79.** The method of claim 76, wherein said DNA-damaging agent is a platinating agent or ionizing radiation.

**80.** The method of claim **73**, wherein said cancer is a solid tumor selected from the following cancers: Oral: buccal cavity, lip, tongue, mouth, pharynx; Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell or epidermoid, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, larynx, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel or small intestines (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma,

lipoma, neurofibroma, fibroma), large bowel or large intestines (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma), colon, colon-rectum, colorectal; rectum, Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor [nephroblastoma], lymphoma), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma, biliary passages; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocytoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteocartilaginous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma [pinealoma], glioblastoma multiforme, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma [serous cystadenocarcinoma, mucinous cystadenocarcinoma, unclassified carcinoma], granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma), breast; Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, keratoacanthoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis, Thyroid gland: papillary thyroid carcinoma, follicular thyroid carcinoma; medullary thyroid carcinoma, multiple endocrine neoplasia type 2A, multiple endocrine neoplasia type 2B, familial medullary thyroid cancer, pheochromocytoma, paraganglioma; and Adrenal glands: neuroblastoma.


**81.** The method of claim **80** wherein said cancer is selected from the group consisting of lung cancer, head and neck cancer, pancreatic cancer, gastric cancer, and brain cancer.

**82.** A method of promoting cell death in cancer cells comprising administering to a patient a compound of formula I,



wherein the variables are as defined according to claim 72.


83. A method of preventing repair of DNA damage in cancer cells comprising administering to a patient a compound of formula I,



I

wherein the variables are as defined according to claim 72.

**84.** A method of inhibiting ATR in a biological sample comprising the step of contacting a compound of formula I




I

wherein the variables are as defined according to claim 72 with said biological sample.

**85.** The method of claim 0, wherein said biological sample is a cell.

**86.** A method of sensitizing cells to DNA damaging agents comprising administering to a patient a compound of formula I,



I

wherein the variables are as defined according to claim 72.

**87.** The method of claim 83, wherein said cell is a cancer cell having defects in the ATM signaling cascade.

**88.** The method of claim 0, wherein said defect is altered expression or activity of one or more of the following: ATM, p53, CHK2, MRE11, RAD50, NBS1, 53BP1, MDC1 or H2AX.

**89.** The method of claim 83, wherein said cell is a cancer cell expressing DNA damaging oncogenes.

**90.** The method of claim 0, wherein said cancer cell has altered expression or activity of one or more of the following: K-Ras, N-Ras, H-Ras, Raf, Myc, Mos, E2F, Cdc25A, CDC4, CDK2, Cyclin E, Cyclin A and Rb.

**91-94.** (canceled)

**95.** The method of claim 80, wherein the cancer is cancer of the pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, or vipoma).

**96.** The method of claim 80, wherein the cancer is lung cancer (bronchogenic carcinoma (squamous cell or epidermoid, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, or mesothelioma).

\* \* \* \* \*