



US00PP27485P2

(12) **United States Plant Patent**
Rijk

(10) **Patent No.:** **US PP27,485 P2**

(45) **Date of Patent:** **Dec. 20, 2016**

(54) **BEGONIA PLANT NAMED ‘FILUCKSTR’**

(50) Latin Name: *Begonia hybrida*
Varietal Denomination: **Filuckstr**

(71) Applicant: **Nadine Rijk**, De Lier (NL)

(72) Inventor: **Nadine Rijk**, De Lier (NL)

(73) Assignee: **Fides B.V.**, De Lier (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/545,700**

(22) Filed: **Jun. 6, 2015**

(51) **Int. Cl.**
A01H 5/02 (2006.01)

(52) **U.S. Cl.**
USPC **Plt./343**

(58) **Field of Classification Search**
USPC Plt./343
CPC A01H 5/0238
See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Begonia* plant named ‘Filuckstr’, characterized by its upright to spreading and mounded plant habit; relatively compact habit; freely basal branching habit; dark green-colored leaves; freely and continuously flowering habit; and semi-double flowers that are yellow in color with apricot-colored margins.

1 Drawing Sheet

1

Botanical designation: *Begonia hybrida*.
Cultivar denomination: ‘FILUCKSTR’.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct cultivar of *Begonia* plant, botanically known as *Begonia hybrida*, and hereinafter referred to by the name ‘Filuckstr’.

The new *Begonia* plant is a product of a planned breeding program conducted by the Inventor in Aalsmeer, The Netherlands. The objective of the breeding program was to develop new freely branching and flowering *Begonia* plants with unique and attractive flower colors.

The new *Begonia* plant originated from a cross-pollination made by the Inventor in 2011 of a proprietary selection of *Begonia hybrida* identified as code number 208207-003, not patented, as the female, or seed, parent and a proprietary selection of *Begonia hybrida* identified as code number 210005, not patented, as the male, or pollen, parent. The new *Begonia* plant was discovered and selected by the Inventor as a single flowering plant from within the progeny of the stated cross-pollination in a controlled greenhouse environment in Aalsmeer, The Netherlands in 2011.

Asexual reproduction of the new *Begonia* plant by vegetative tip cuttings in a controlled greenhouse environment in Aalsmeer, The Netherlands since 2011 has shown that the unique features of this new *Begonia* plant are stable and reproduced true to type in successive generations.

SUMMARY OF THE INVENTION

Plants of the new *Begonia* have not been observed under all possible combinations of environmental conditions and cultural practices. The phenotype may vary somewhat with variations in environment such as temperature and light intensity, without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘Filuckstr’. These characteristics in combination distinguish ‘Filuckstr’ as a new and distinct *Begonia* plant:

2

1. Upright to spreading and mounded plant habit; relatively compact.
2. Freely basal branching habit.
3. Dark green-colored leaves.
4. Freely and continuously flowering habit.
5. Semi-double flowers that are yellow in color with apricot-colored margins.

Plants of the new *Begonia* can be compared to plants of the proprietary parent selections. Plants of the new *Begonia* differ from plants of the parent selections in the following characteristics:

1. Plants of the new *Begonia* are more uniform than plants of the parent selections.
2. Plants of the new *Begonia* are more freely flowering than plants of the parent selections.

Plants of the new *Begonia* can be compared to plants of *Begonia hybrida* ‘Fimissmo’, disclosed in a U.S. Plant Patent application filed concurrently, 14/545,695. Plants of the new *Begonia* differ primarily from plants of ‘Fimissmo’ in the following characteristics:

1. Plants of the new *Begonia* are more freely branching than plants of ‘Fimissmo’.
2. Plants of the new *Begonia* have smaller flowers than plants of ‘Fimissmo’.
3. Plants of the new *Begonia* have semi-double flowers whereas plants of ‘Fimissmo’ have double flowers.
4. Plants of the new *Begonia* have yellow-colored flowers with apricot-colored margins whereas plants of ‘Fimissmo’ have cream-colored flowers with light pink-colored margins.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new *Begonia* plant showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may

differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Begonia* plant.

The photograph is a side perspective view of a typical flowering plant of 'Filuckstr' grown in a container.

DETAILED BOTANICAL DESCRIPTION

Plants used for the aforementioned photograph and following observations and measurements were grown in 12-cm containers during the summer in a glass-covered greenhouse in Rheinberg, Germany. During the production of the plants, day temperatures ranged from 17° C. to 30° C. and night temperatures ranged from 10° C. to 20° C. Plants were eight weeks old when the photograph and the description were taken. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 2007 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Begonia hybrida* 'Filuckstr'.

Parentage:

Female, or seed, parent.—Proprietary selection of *Begonia hybrida* identified as code number 208207-003, not patented.

Male, or pollen, parent.—Proprietary selection of *Begonia hybrida* identified as code number 210005, not patented.

Propagation:

Type.—By vegetative tip cuttings.

Time to initiate roots, summer.—About 18 days at temperatures about 22° C. to 30° C.

Time to initiate roots, winter.—About 21 days at temperatures about 22° C. to 30° C.

Time to produce a rooted young plant, summer.—About 25 days at temperatures about 22° C. to 30° C.

Time to produce a rooted young plant, winter.—About 28 days at temperatures about 20° C. to 25° C.

Root description.—Medium in thickness, fibrous; whitish grey in color; plants of the new *Begonia* have not been observed to form tubers.

Rooting habit.—Moderately freely branching; medium density.

Plant description:

Plant form and growth habit.—Upright to spreading and mounded plant habit; relatively compact; pendulous with development; freely basal branching with about 15 primary branches; moderately vigorous to vigorous growth habit.

Plant height.—About 24 cm.

Plant width.—About 36 cm.

Lateral branch description.—Length: About 15 cm to 16 cm. Diameter: About 1 cm. Internode length: About 1 cm to 3 cm. Texture: Smooth, glabrous. Color: Close to 144B.

Leaf description.—Arrangement: Alternate, simple. Length: About 8 cm to 11 cm. Width: About 3.5 cm to 5 cm. Shape: Ovate to lanceolate. Apex: Acute. Base: Cordate. Margin: Serrate. Texture, upper and lower surfaces: Smooth, glabrous. Venation pattern: Palmate; reticulate. Color: Developing leaves, upper surface: Close to 183A. Developing leaves, lower surface: Close to 183B. Fully expanded leaves, upper surface: Close to N137A; venation, close to 139B. Fully expanded leaves, lower surface: Close to 144A and N137A; venation, close to 144A and N137A.

Petioles: Length: About 2.5 cm to 5.5 cm. Diameter: About 3 mm to 5 mm. Texture, upper and lower surfaces: Smooth, glabrous. Color, upper and lower surfaces: Close to 144B.

5 Flower description:

Flowering habit.—Semi-double flowers arranged in axillary cymes; freely flowering habit with numerous flowers developing per plant; flowers pendulous and face outwardly to downwardly.

10 *Fragrance.*—None detected.

Natural flowering season.—Plants in full flower about eight weeks after planting; long flowering period, in the garden plants flower freely and continuously throughout the summer in Northern Europe and can be flowered year-round in greenhouses.

Flower longevity.—Individual flowers last about four weeks on the plant; flowers persistent.

15 *Inflorescence height (including peduncle).*—About 8 cm to 9 cm.

Inflorescence diameter.—About 5 cm.

Inflorescence peduncles.—Length: About 7 cm. Diameter: About 3 mm to 5 mm. Angle: About 30° to 45° from vertical. Strength: Moderately strong. Texture: Smooth, glabrous. Color: Close to 144B.

20 *Female flower buds.*—Length: About 1.5 cm to 2 cm. Diameter: About 1 cm. Shape: Ovoid. Texture: Smooth, glabrous. Color: Close to 29C.

Female flowers.—Diameter: About 4 cm to 4.5 cm. Depth (height): About 3 cm to 4 cm.

30 *Female flower tepals.*—Quantity per flower and arrangement: Typically four per flower arranged in a single whorl. Length: About 3 cm to 4 cm. Width: About 1.5 cm to 2 cm. Shape: Ovate. Apex: Acute. Base: Cordate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper surface: Close to 12C; towards the margins, close to 29B; color does not fade with development. When opening and fully opened, lower surface: Close to 12C; towards the margins, close to 29B; color does not fade with development.

Female flower tepaloids.—None observed.

Female flower pedicels.—Length: About 3 cm to 4 cm. Diameter: About 2 mm to 3 mm. Aspect: About 90° from peduncle axis. Strength: Moderately strong. Texture: Smooth, glabrous. Color: Close to 144B.

Female flowers reproductive organs.—Quantity of pistils per flower: Three. Pistil length: About 5 mm. Style length: About 5 mm. Style color: Close to 145A. Stigma shape: Curled. Stigma color: Close to 13A. Ovary color: Close to 145A. Fruits and seeds: Fruit and seed development have not been observed on plants of the new *Begonia*.

55 *Male flower buds.*—Length: About 3 cm. Diameter: About 2 cm to 3 cm. Shape: Ovoid. Texture: Smooth, glabrous. Color: Close to 29C.

Male flowers.—Diameter: About 6 cm to 6.5 cm. Depth (height): About 3 cm.

60 *Male flower tepals.*—Quantity per flower and arrangement: Typically about four or five per flower arranged in a single whorl. Length: About 1 cm to 3 cm. Width: About 0.5 cm to 3 cm. Shape: Ovate. Apex: Acute. Base: Cordate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper sur-

face: Close to 12C; towards the margins, close to 29B; color does not fade with development. When opening and fully opened, lower surface: Close to 12C; towards the margins, close to 29B; color does not fade with development.

Male flower tepaloids.—Quantity per flower and arrangement: Typically about 25 per flower arranged in multiple whorls. Length: About 1 cm to 2 cm. Width: About 4 mm to 6 mm. Shape: Roughly oval. Apex: Mostly obtuse. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Smooth, glabrous. Color: When opening and fully opened, upper surface: Close to 29A and 15A; color does not fade with development. When opening and fully opened, lower surface: Close to 29A and 15A; color does not fade with development.

Male flower pedicels.—Length: About 2.5 cm to 3 cm. Diameter: About 2 mm to 4 mm. Aspect: About 90° from peduncle axis. Strength: Moderately strong. Texture: Smooth, glabrous. Color: Close to 144C.

Male flowers reproductive organs.—None observed; all transformed into tepaloids.

Disease & pest resistance: Resistance to pathogens and pests common to *Begonia* plants has not been observed on plants of the new *Begonia*.

10 Temperature tolerance: Plants of the new *Begonia* have been observed to tolerate temperatures ranging from about 4° C. to about 35° C.

It is claimed:

15 1. A new and distinct *Begonia* plant named 'Filuckstr' as illustrated and described.

* * * * *

