




(72) RICE, DOUGLAS A., US


(71) PIONEER HI-BRED INTERNATIONAL, INC., US

(51) Int.Cl.⁶ C12N 15/29, C12N 15/82, C12N 15/53, A01H 5/00, C12N 5/10,
A01H 5/10

(30) 1998/02/26 (60/076,075) US


(54) **PROMOTEURS CONSTITUTIFS DU MAIS**

(54) **CONSTITUTIVE MAIZE PROMOTERS**



(57) La présente invention concerne des compositions et des méthodes permettant de réguler l'expression de séquences nucléotidiques hétérologues chez une plante. Les compositions de cette invention sont de nouvelles séquences nucléotidiques de promoteurs constitutifs isolés à partir de gènes de maïs codant pour la histone H2B, la métallothionéine, l'alpha-tubuline 3, le facteur d'allongement efla, la protéine ribosomique rps8, la

(57) The present invention provides compositions and methods for regulating expression of heterologous nucleotide sequences in a plant. Compositions are novel nucleotide sequences for constitutive promoters isolated from maize genes encoding histone H2B, metallothionein, alpha-tubulin 3, elongation factor efla, ribosomal protein rps8, chlorophyll a/b binding protein, and glyceraldehyde-3-phosphate dehydrogenase. A



(21) (A1) **2,315,546**
(86) 1999/02/25
(87) 1999/09/02


protéine de liaison chlorophylle a/b, et la glycéraldéhyde-3-phosphate déhydrogénase. L'une des méthodes de cette invention permet d'exprimer de manière constitutive une séquence nucléotidique hétérologue chez une plante à l'aide des séquences promoteurs susmentionnées, cette méthode consistant tout d'abord à transformer une cellule végétale pour que celle-ci renferme une séquence nucléotidique hétérologue liée de manière opérationnelle à l'un des promoteurs constitutifs de la présente invention, et enfin à régénérer une plante transformée de manière stable à partir de ladite cellule végétale transformée.

method for constitutively expressing a heterologous nucleotide sequence in a plant using the promoter sequences disclosed herein is provided. The method comprises transforming a plant cell to comprise a heterologous nucleotide sequence operably linked to one of the constitutive promoters of the present invention and regenerating a stably transformed plant from the transformed plant cell.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00	A2	(11) International Publication Number: WO 99/43797 (43) International Publication Date: 2 September 1999 (02.09.99)
(21) International Application Number: PCT/US99/04203 (22) International Filing Date: 25 February 1999 (25.02.99) (30) Priority Data: 60/076,075 26 February 1998 (26.02.98) US (71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50319 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): RICE, Douglas, A. [US/US]; 4334 Woodland Avenue, Des Moines, IA 50312 (US). (74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report. With an indication in relation to deposited biological material furnished under Rule 13bis separately from the description.</i>

(54) Title: CONSTITUTIVE MAIZE PROMOTERS**(57) Abstract**

The present invention provides compositions and methods for regulating expression of heterologous nucleotide sequences in a plant. Compositions are novel nucleotide sequences for constitutive promoters isolated from maize genes encoding histone H2B, metallothionein, alpha-tubulin 3, elongation factor e1a, ribosomal protein rps8, chlorophyll a/b binding protein, and glyceraldehyde-3-phosphate dehydrogenase. A method for constitutively expressing a heterologous nucleotide sequence in a plant using the promoter sequences disclosed herein is provided. The method comprises transforming a plant cell to comprise a heterologous nucleotide sequence operably linked to one of the constitutive promoters of the present invention and regenerating a stably transformed plant from the transformed plant cell.

CONSTITUTIVE MAIZE PROMOTERS

CROSS-REFERENCE TO RELATED APPLICATION

This application claims the benefit of U.S. Provisional Application Serial No. 60/076,075, filed February 26, 1998, herein incorporated by reference.

5 FIELD OF THE INVENTION

The present invention relates to the field of plant molecular biology, more particularly to regulation of gene expression in plants.

BACKGROUND OF THE INVENTION

10 Expression of heterologous DNA sequences in a plant host is dependent upon the presence of an operably linked promoter that is functional within the plant host. Choice of the promoter sequence will determine when and where within the organism the heterologous DNA sequence is expressed. Thus, where continuous expression is desired throughout the cells of a plant, constitutive promoters are utilized. In contrast,
15 where gene expression in response to a stimulus is desired, inducible promoters are the regulatory element of choice. In either case, additional regulatory sequences upstream and/or downstream from the core promoter sequence may be included in expression constructs of transformation vectors to bring about varying levels of constitutive or inducible expression of heterologous nucleotide sequences in a
20 transgenic plant.

Frequently it is desirable to have constitutive expression of a DNA sequence throughout the cells of an organism. For example, increased resistance of a plant to infection by soil- and air-borne pathogens might be accomplished by genetic manipulation of the plant's genome to comprise a constitutive promoter operably
25 linked to a heterologous pathogen-resistance gene such that pathogen-resistance proteins are continuously expressed throughout the plant's tissues.

Alternatively, it might be desirable to inhibit expression of a native DNA sequence within a plant's tissues to achieve a desired phenotype. In this case, such inhibition might be accomplished with transformation of the plant to comprise a

constitutive promoter operably linked to an antisense nucleotide sequence, such that constitutive expression of the antisense sequence produces an RNA transcript that interferes with translation of the mRNA of the native DNA sequence.

Thus, isolation and characterization of constitutive promoters that can serve as
5 regulatory regions for constitutive expression of heterologous nucleotide sequences of interest are needed for genetic manipulation of plants to exhibit specific phenotypic traits.

SUMMARY OF THE INVENTION

10 Compositions and methods for regulating expression of heterologous nucleotide sequences in a plant are provided. Compositions are novel nucleotide sequences for constitutive plant promoters, more particularly promoters isolated from maize genes encoding histone H2B, metallothionein, alpha-tubulin 3, elongation factor efla, ribosomal protein rps8, chlorophyll a/b binding protein, and
15 glyceraldehyde-3-phosphate dehydrogenase. A method for constitutively expressing a heterologous nucleotide sequence in a plant using the promoter sequences disclosed herein is provided. The method comprises transforming a plant cell with a transformation vector that comprises a heterologous nucleotide sequence operably linked to one of the plant promoters of the present invention and regenerating a stably
20 transformed plant from the transformed plant cell.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 shows the plasmid vector PHP3953 comprising the GUS gene operably linked to the ubiquitin promoter. Promoter fragments of the present
25 invention were recloned into this plasmid in place of the ubiquitin promoter, and the resulting plasmid DNA was available for use in transformation studies to test for promoter activity.

DETAILED DESCRIPTION OF THE INVENTION

30 Compositions of the present invention are nucleic acid molecules comprising novel nucleotide sequences for plant promoters, more particularly constitutive promoters for the maize genes encoding histone H2B; metallothionein-1; alpha-tubulin 3-18; elongation factor efla-11, efla-15, and efla-16; ribosomal protein rps8; chlorophyll a/b binding protein cab-10 and cab-20; and glyceraldehyde-3-phosphate

dehydrogenase gpc4. The nucleotide sequences for these promoters are set forth in SEQ ID NOs: 1-10, respectively. In particular, the present invention provides for isolated nucleic acid molecules comprising nucleotide sequences encoding the DNA sequences deposited in a bacterial host as ATCC Accession Nos. _____, and
5 variants and fragments thereof. The promoters for these maize genes were isolated from the 5' untranslated region flanking their respective transcription initiation sites. Methods for isolation of promoter regions are well known in the art. The specific method used to obtain the promoters of the present invention is described in Example 1 below.

10 Plasmids containing the promoter nucleotide sequences of the invention were deposited with American Type Culture Collection (ATCC), Manassas, Virginia, and assigned Accession Nos. _____. These deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. These deposits were made
15 merely as a convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

The invention encompasses isolated or substantially purified nucleic acid compositions. An "isolated" or "purified" nucleic acid molecule, or biologically active portion thereof, is substantially free of other cellular material, or culture
20 medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (*i.e.*, sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is
25 derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived.

By "promoter" is intended a regulatory region of DNA usually comprising a
30 TATA box capable of directing RNA polymerase II to initiate RNA synthesis at the appropriate transcription initiation site for a particular coding sequence. A promoter may additionally comprise other recognition sequences generally positioned upstream or 5' to the TATA box, referred to as upstream promoter elements, which influence the transcription initiation rate. It is recognized that having identified the nucleotide

sequences for the promoter regions disclosed herein, it is within the state of the art to isolate and identify further regulatory elements in the 5' untranslated region upstream from the particular promoter regions identified herein. Thus the promoter regions disclosed herein may further comprise upstream regulatory elements that confer
5 tissue-specific expression of any heterologous nucleotide sequence operably linked to one of the disclosed promoter sequences. See particularly Australian Patent No. AU-A-77751/94 and U.S. Patent Nos. 5,466,785 and 5,635,618.

The maize promoter sequences of the present invention, when assembled within a DNA construct such that the promoter is operably linked to a heterologous
10 nucleotide sequence of interest, enable constitutive expression of the heterologous nucleotide sequence in the cells of a plant stably transformed with this DNA construct. By "heterologous nucleotide sequence" is intended a sequence that is not naturally occurring with the promoter sequence. While this nucleotide sequence is heterologous to the promoter sequence, it may be homologous or native or
15 heterologous or foreign to the plant host. By "constitutive" is intended expression in the cells throughout a plant at most times and in most tissues.

The isolated promoter sequences of the present invention can be classified as providing for a range of constitutive expression. Thus, some are weak constitutive promoters, and others are strong constitutive promoters. Generally, by "weak
20 promoter" is intended a promoter that drives expression of a coding sequence at a low level. By "low level" is intended at levels of about 1/10,000 transcripts to about 1/100,000 transcripts to about 1/500,000 transcripts. Conversely, a strong promoter drives expression of a coding sequence at a high level, or at about 1/10 transcripts to about 1/100 transcripts to about 1/1,000 transcripts.

25 The nucleotide sequences for the constitutive promoters of the present invention may be the naturally occurring sequences or any sequence having substantial homology. By "substantial homology" is intended a sequence exhibiting substantial functional and structural equivalence with the native or naturally occurring sequence. Any functional or structural differences between substantially homologous
30 sequences do not effect the ability of the sequence to function as a promoter as disclosed in the present invention. Thus, any sequence having substantial sequence homology with the sequence of a particular constitutive promoter of the present invention will direct constitutive expression of an operably linked heterologous nucleotide sequence. Two promoter nucleotide sequences are considered

substantially homologous when they have at least about 50%, 60%, to 70%, generally about 80%, preferably about 85%, 90%, up to 98% sequence homology.

Substantially homologous sequences of the present invention include variants of the disclosed sequences, such as those that result from site-directed mutagenesis, as well as synthetically derived sequences. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods Enzymol.* 154:367-382; U.S. Patent No. 4,873,192; Walker *et al.*, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Thus, the promoter nucleotide sequences of the invention include both the naturally occurring sequences as well as mutant and synthetically derived forms. Generally, nucleotide sequence variants of the invention will have at least about 50%, 60%, to 70%, generally about 80%, preferably about 85%, 90%, up to 98% sequence identity to its respective native nucleotide sequence.

Fragments of the promoter nucleotide sequences disclosed herein are also encompassed by the present invention. By "fragment" is intended a portion of the promoter nucleotide sequence. Fragments of a promoter nucleotide sequence may retain their biological activity. Thus, for example, less than the entire promoter sequences disclosed herein may be utilized to drive expression of an operably linked nucleotide sequence of interest, such as a nucleotide sequence encoding a heterologous protein. It is within skill in the art to determine whether such fragments decrease expression levels or alter the nature of expression, *i.e.*, constitutive expression. Alternatively, fragments of a promoter nucleotide sequence that are useful as hybridization probes generally do not retain this biological activity.

Nucleic acid molecules that are fragments of a promoter nucleotide sequence comprise at least 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 325, 350, 375, 400, 425, 450, 500, 550, 600, 650, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, 1,400, 1,500, 1,600, or 1,700 nucleotides, or up to the number of nucleotides present in a full-length promoter nucleotide sequence disclosed herein (*i.e.*, 1392, 1300, 166, 1023, 1022, 1769, 461, 467, 467, and 565 for SEQ ID NOs: 1-10, respectively). Generally, fragments of a promoter sequence that retain their biological activity comprise at least 30, 35, 40 contiguous nucleotides, preferably at least 50 contiguous nucleotides, more preferably at least 75 contiguous nucleotides, still more preferably at least 100 contiguous nucleotides of the particular promoter nucleotide sequence disclosed herein. Preferred

fragment lengths depend upon the objective and will also vary depending upon the particular promoter sequence.

The nucleotides of such fragments will usually comprise the TATA recognition sequence of the particular promoter sequence. Such fragments may be obtained by use of restriction enzymes to cleave the naturally occurring promoter nucleotide sequence disclosed herein; by synthesizing a nucleotide sequence from the naturally occurring sequence of the promoter DNA sequence; or may be obtained through the use of PCR technology. See particularly, Mullis *et al.* (1987) *Methods Enzymol.* 155:335-350, and Erlich, ed. (1989) *PCR Technology* (Stockton Press, New York). Variants of these promoter fragments, such as those resulting from site-directed mutagenesis, are encompassed by the compositions of the present invention.

The disclosed promoter nucleotide sequences can be used to isolate homologous promoter sequences in other plant species. Methods are readily available in the art for the hybridization of nucleic acid sequences. Promoter sequences from other plants may be isolated according to well-known techniques based on their sequence homology to the promoter sequences set forth herein. In these techniques all or part of the known nucleotide sequence is used as a probe which selectively hybridizes to other promoter nucleotide sequences present in a population of cloned genomic DNA fragments or cDNA fragments (*i.e.*, genomic or cDNA libraries) from a chosen organism.

To obtain other homologous promoter sequences, the entire promoter nucleotide sequence or portions thereof may be used as probes capable of specifically hybridizing to corresponding promoter sequences. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique and are preferably at least about 10 nucleotides in length, and most preferably at least about 20 nucleotides in length. Such probes may be used to amplify the promoter sequences of interest from a chosen organism by the well-known process of polymerase chain reaction (PCR). This technique may be used to isolate additional promoter sequences from a desired organism or as a diagnostic assay to determine the presence of promoter sequences in an organism.

Such techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, *e.g.*, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York) and amplification by PCR using oligonucleotide primers (see, *e.g.*, Innis *et al.*,

eds. (1990) *PCR Protocols, a Guide to Methods and Applications*, Academic Press, New York).

For example, hybridization of such sequences may be carried out under conditions of reduced stringency, medium stringency, or even stringent conditions (e.g., conditions represented by a wash stringency of 35-40% Formamide with 5x Denhardt's solution, 0.5% SDS and 1x SSPE at 37°C; conditions represented by a wash stringency of 40-45% Formamide with 5x Denhardt's solution, 0.5% SDS, and 1x SSPE at 42°C; and conditions represented by a wash stringency of 50% Formamide with 5x Denhardt's solution, 0.5% SDS and 1x SSPE at 42°C, respectively) to DNA for the promoter sequences disclosed herein in a standard hybridization assay. See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2d ed., Cold Spring Harbor Laboratory Press, Plainview, New York). In general, homologous sequences that are promoter sequences and hybridize to the promoter sequences disclosed herein will be at least 40% to 50% homologous, 60% to 70% homologous, and even 80%, 85%, 90%, 95% homologous or more with the disclosed sequence. That is, the sequence similarity of sequences may range, sharing at least about 40%, 50%, about 60%, 70%, and even about 80%, 85%, 90%, 95%, 98% sequence similarity.

The following terms are used to describe the sequence relationships between two or more nucleic acids: (a) "reference sequence", (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity".

(a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence.

(b) As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (*i.e.*, gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a

reference sequence due to inclusion of gaps in the polynucleotide sequence, a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison may be conducted by the local
5 homology algorithm of Smith *et al.* (1981) *Adv. Appl. Math.* 2:482; by the homology
alignment algorithm of Needleman *et al.* (1970) *J. Mol. Biol.* 48:443; by the search
for similarity method of Pearson *et al.* (1988) *Proc. Natl. Acad. Sci.* 85:2444; by
computerized implementations of these algorithms, including, but not limited to:
CLUSTAL in the PC/Gene program by Intelligenetics, Mountain View, California;
10 GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software
Package, Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin;
the CLUSTAL program is well described by Higgins *et al.* (1988) *Gene* 73:237-244
(1988); Higgins *et al.* (1989) *CABIOS* 5:151-153; Corpet *et al.* (1988) *Nuc. Acids*
Res. 16:10881-90; Huang *et al.* (1992) *Computer Applications in the Biosciences*
15 8:155-65, and Person *et al.* (1994) *Meth. Mol. Biol.* 24:307-331; preferred computer
alignment methods also include the BLASTP, BLASTN, and BLASTX algorithms
(see Altschul *et al.* (1990) *J. Mol. Biol.* 215:403-410). Alignment is also often
performed by inspection and manual alignment.

(c) As used herein, "sequence identity" or "identity" in the context of two
20 nucleic acid sequences makes reference to the residues in the two sequences that are
the same when aligned for maximum correspondence over a specified comparison
window.

(d) As used herein, "percentage of sequence identity" means the value
determined by comparing two optimally aligned sequences over a comparison
25 window, wherein the portion of the polynucleotide sequence in the comparison
window may comprise additions or deletions (*i.e.*, gaps) as compared to the reference
sequence (which does not comprise additions or deletions) for optimal alignment of
the two sequences. The percentage is calculated by determining the number of
positions at which the identical nucleic acid base occurs in both sequences to yield the
30 number of matched positions, dividing the number of matched positions by the total
number of positions in the window of comparison, and multiplying the result by 100
to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that
a polynucleotide comprises a sequence that has at least 70% sequence identity,

preferably at least 80%, more preferably at least 90% and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters.

Another indication that nucleotide sequences are substantially identical is if
5 two molecules hybridize to each other under stringent conditions. Generally, stringent conditions are selected to be about 5°C to about 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent wash
10 conditions are those in which the salt concentration is about 0.02 molar at pH 7 and the temperature is at least about 50, 55, or 60°C.

The nucleotide sequences for the constitutive promoters disclosed in the present invention, as well as variants and fragments thereof, are useful in the genetic manipulation of any plant when assembled within a DNA construct such that the
15 promoter sequence is operably linked with a heterologous nucleotide sequence whose constitutive expression is to be controlled to achieve a desired phenotypic response. By "operably linked" is intended the transcription or translation of the heterologous nucleotide sequence is under the influence of the promoter sequence. In this manner, the nucleotide sequences for the promoters of the invention are provided in expression
20 cassettes along with heterologous nucleotide sequences for expression in the plant of interest. It is recognized that the promoter sequences of the invention may also be used with their native coding sequences to increase or decrease expression of the native coding sequence, thereby resulting in a change in phenotype in the transformed plant.

25 Such expression cassettes will comprise a transcriptional initiation region comprising one of the promoter nucleotide sequences of the present invention, or variants or fragments thereof, operably linked to the heterologous nucleotide sequence whose expression is to be controlled by the constitutive promoters disclosed herein. Such an expression cassette is provided with a plurality of restriction sites for
30 insertion of the nucleotide sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

In order to increase transcription levels, enhancers may be utilized in combination with the promoter regions of the invention. Enhancers are nucleotide

sequences that act to increase the expression of a promoter region. Enhancers are known in the art and include the SV40 enhancer region, the 35S enhancer element, and the like.

The transcriptional cassette will include in the 5'-to-3' direction of
5 transcription, a transcriptional and translational initiation region, a heterologous nucleotide sequence of interest, and a transcriptional and translational termination region functional in plants. The termination region may be native with the transcriptional initiation region comprising one of the promoter nucleotide sequences of the present invention, may be native with the DNA sequence of interest, or may be
10 derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158;
15 Ballas *et al.* 1989) *Nuc. Acids Res.* 17:7891-7903; Joshi *et al.* (1987) *Nuc. Acid Res.* 15:9627-9639.

The expression cassette comprising the promoter sequence of the present invention operably linked to a heterologous nucleotide sequence may also contain at least one additional nucleotide sequence for a gene to be cotransformed into the
20 organism. Alternatively, the additional sequence(s) can be provided on another expression cassette.

Where appropriate, the heterologous nucleotide sequence whose expression is to be under the control of the promoter sequence of the present invention and any additional nucleotide sequence(s) may be optimized for increased expression in the
25 transformed plant. That is, these nucleotide sequences can be synthesized using plant-preferred codons for improved expression. Methods are available in the art for synthesizing plant-preferred nucleotide sequences. See, for example, U.S. Patent Nos. 5,380,831 and 5,436,391, and Murray *et al.* (1989) *Nuc. Acids Res.* 17:477-498, herein incorporated by reference.

30 Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene expression. The G-C content of the heterologous nucleotide sequence may be adjusted to levels

average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein *et al.* (1989) *Proc. Nat. Acad. Sci. USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986)); MDMV leader (Maize Dwarf Mosaic Virus) (*Virology* 154:9-20); human immunoglobulin heavy-chain binding protein (BiP) (Macejak and Sarnow (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling and Gehrke (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) *Molecular Biology of RNA*, pages 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also Della-Cioppa *et al.* (1987) *Plant Physiology* 84:965-968. Other methods known to enhance translation and/or mRNA stability can also be utilized, for example, introns, and the like.

In those instances where it is desirable to have the constitutively expressed product of the heterologous nucleotide sequence directed to a particular organelle, such as the chloroplast or mitochondrion, or secreted at the cell's surface or extracellularly, the expression cassette may further comprise a coding sequence for a transit peptide. Such transit peptides are well known in the art and include, but are not limited to, the transit peptide for the acyl carrier protein, the small subunit of RUBISCO, plant EPSP synthase, and the like.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, for example, transitions and transversions, may be involved.

The expression cassette comprising the particular promoter sequence of the present invention operably linked to a heterologous nucleotide sequence of interest can be used to transform any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. Transformation protocols may vary depending on the type of plant or plant cell, *i.e.*, monocot or dicot, targeted for transformation. Suitable methods of transforming plant cells include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606), *Agrobacterium*-mediated transformation (Hinchey *et al.* (1988) *Biotechnology* 6:915-921), direct gene transfer (Paszowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.* U.S. Patent 4,945,050; Tomes *et al.* (1995) in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Annual Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839; Hooydaas-Van Slogteren and Hooykaas (1984) *Nature (London)* 311:763-764; Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. G. P. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418; and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or

more generations may be grown to ensure that constitutive expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure constitutive expression of the desired phenotypic characteristic has been achieved.

5 The promoter nucleotide sequences and methods disclosed herein are useful in regulating constitutive expression of any heterologous nucleotide sequence in a host plant in order to vary the phenotype of a plant. Various changes in phenotype are of interest including modifying the fatty acid composition in a plant, altering the amino acid content of a plant, altering a plant's pathogen defense mechanism, and the like.
10 These results can be achieved by providing expression of heterologous products or increased expression of endogenous products in plants. Alternatively, the results can be achieved by providing for a reduction of expression of one or more endogenous products, particularly enzymes or cofactors in the plant. These changes result in a change in phenotype of the transformed plant.

15 Genes of interest are reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest change, and as developing nations open up world markets, new crops and technologies will emerge also. In addition, as our understanding of agronomic traits and characteristics such as yield and heterosis increase, the choice of genes for transformation will
20 change accordingly. General categories of genes of interest include for example, those genes involved in information, such as zinc fingers, those involved in communication, such as kinases, and those involved in housekeeping, such as heat shock proteins. More specific categories of transgenes, for example, include genes encoding important traits for agronomics, insect resistance, disease resistance,
25 herbicide resistance, sterility, grain characteristics, and commercial products. Genes of interest include, generally, those involved in oil, starch, carbohydrate, or nutrient metabolism as well as those affecting kernel size, sucrose loading, and the like.

 Agronomically important traits such as oil, starch, and protein content can be genetically altered in addition to using traditional breeding methods. Modifications
30 include increasing content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur, providing essential amino acids, and also modification of starch. Hordothionin protein modifications are described in U.S. Patent Application Serial Nos. 08/838,763, filed April 10, 1997; 08/824,379, filed March 26, 1997; 08/824,382, filed March 26, 1997; and U.S. Patent No. 5,703,409; herein incorporated

by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in U.S. Serial No. 08/618,911, filed March 20, 1996, and the chymotrypsin inhibitor from barley, Williamson *et al.* (1987) *Eur. J. Biochem.* 165:99-106, the disclosures of which are herein incorporated by reference.

5 Derivatives of the coding sequences can be made by site directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL) is derived from barley chymotrypsin inhibitor, U.S. Serial No. 08/740,682 filed November 1, 1996, and PCT/US97/20441, filed October 31, 1997, the disclosures of each are
10 incorporated herein by reference. Other proteins include methionine-rich plant proteins such as from sunflower seed (Lilley *et al.* (1989) *Proceedings of the World Congress on Vegetable Protein Utilization in Human Foods and Animal Feedstuffs*, ed. Applewhite (American Oil Chemists Society, Champaign, Illinois), pp. 497-502; herein incorporated by reference)); corn (Pedersen *et al.* (1986) *J. Biol. Chem.*
15 261:6279; Kirihaara *et al.* (1988) *Gene* 71:359; both of which are herein incorporated by reference); and rice (Musumura *et al.* (1989) *Plant Mol. Biol.* 12:123, herein incorporated by reference). Other agronomically important genes encode latex, Flourey 2, growth factors, seed storage factors, and transcription factors.

 Insect resistance genes may encode resistance to pests that have great yield
20 drag such as rootworm, cutworm, European Corn Borer, and the like. Such genes include, for example *Bacillus thuringiensis* toxic protein genes (U.S. Patent Nos. 5,366,892; 5,747,450; 5,737,514; 5,723,756; 5,593,881; Geiser *et al.* (1986) *Gene* 48:109); lectins (Van Damme *et al.* (1994) *Plant Mol. Biol.* 24:825); and the like.

 Genes encoding disease resistance traits include detoxification genes, such as
25 against fumonosin (U.S. Patent Application Serial No. 08/484,815, filed June 7, 1995); avirulence (avr) and disease resistance (R) genes (Jones *et al.* (1994) *Science* 266:789; Martin *et al.* (1993) *Science* 262:1432; Mindrinos *et al.* (1994) *Cell* 78:1089); and the like.

 Herbicide resistance traits may include genes coding for resistance to
30 herbicides that act to inhibit the action of acetolactate synthase (ALS), in particular the sulfonylurea-type herbicides (*e.g.*, the acetolactate synthase (ALS) gene containing mutations leading to such resistance, in particular the S4 and/or Hra mutations), genes coding for resistance to herbicides that act to inhibit action of glutamine synthase, such as phosphinothricin or basta (*e.g.*, the bar gene), or other

such genes known in the art. The *bar* gene encodes resistance to the herbicide basta, the *nptII* gene encodes resistance to the antibiotics kanamycin and geneticin, and the ALS gene encodes resistance to the herbicide chlorsulfuron.

5 Sterility genes can also be encoded in an expression cassette and provide an alternative to physical detasseling. Examples of genes used in such ways include male tissue-preferred genes and genes with male sterility phenotypes such as QM, described in U.S. Patent No. 5,583,210. Other genes include kinases and those encoding compounds toxic to either male or female gametophytic development.

10 The quality of grain is reflected in traits such as levels and types of oils, saturated and unsaturated, quality and quantity of essential amino acids, and levels of cellulose. In corn, modified hordothionin proteins, described in U.S. Patent Application Serial Nos. 08/838,763, filed April 10, 1997; 08/824,379, filed March 26, 1997; 08/824,382, filed March 26, 1997; and U.S. Patent No. 5,703,409 issued December 30, 1997, provide descriptions of modifications of proteins for desired
15 purposes.

Commercial traits can also be encoded on a gene or genes that could increase for example, starch for ethanol production, or provide expression of proteins. Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Patent No. 5,602,321 issued
20 February 11, 1997. Genes such as B-Ketothiolase, PHBase (polyhydroxybutyrate synthase) and acetoacetyl-CoA reductase (see Schubert *et al.* (1988) *J. Bacteriol.* 170:5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs).

Exogenous products include plant enzymes and products as well as those from other sources including procaryotes and other eukaryotes. Such products include
25 enzymes, cofactors, hormones, and the like. The level of proteins, particularly modified proteins having improved amino acid distribution to improve the nutrient value of the plant, can be increased. This is achieved by the expression of such proteins having enhanced amino acid content.

Thus, the heterologous nucleotide sequence operably linked to one of the
30 constitutive promoters disclosed herein may be a structural gene encoding a protein of interest. Examples of such heterologous genes include, but are not limited to, genes encoding proteins conferring resistance to abiotic stress, such as drought, temperature, salinity, and toxins such as pesticides and herbicides, or to biotic stress, such as attacks by fungi, viruses, bacteria, insects, and nematodes, and development of

diseases associated with these organisms. More particularly, the constitutive promoters disclosed herein and identified as weak constitutive promoters are useful in transforming plants to constitutively express an avirulence gene as disclosed in the copending applications both entitled "*Methods for Enhancing Disease Resistance in*
5 *Plants*," U.S. Application Serial No. 60/075,151, filed February 26, 1998, and U.S. Application Serial No. 60/092,464, filed July 11, 1998, both of which are herein incorporated by reference. Such weak promoters may cause activation of the plant defense system short of hypersensitive cell death. Thus, there is an activation of the plant defense system at levels sufficient to protect from pathogen invasion. In this
10 state, there is at least a partial activation of the plant defense system wherein the plant produces increased levels of antipathogenic factors such as PR proteins, *i.e.*, PR-1, chitinases, α -glucanases, etc.; secondary metabolites; phytoalexins; reactive oxygen species; and the like.

Alternatively, the heterologous nucleotide sequence operably linked to one of
15 the constitutive promoters disclosed herein may be an antisense sequence for a targeted gene. By "antisense DNA nucleotide sequence" is intended a sequence that is in inverse orientation to the 5' to 3' normal orientation of that nucleotide sequence. When delivered into a plant cell, expression of the antisense DNA sequence prevents normal expression of the DNA nucleotide sequence for the targeted gene. The
20 antisense nucleotide sequence encodes an RNA transcript that is complementary to and capable of hybridizing to the endogenous messenger RNA (mRNA) produced by transcription of the DNA nucleotide sequence for the targeted gene. In this case, production of the native protein encoded by the targeted gene is inhibited to achieve a desired phenotypic response. Thus the promoter sequences disclosed herein may be
25 operably linked to antisense DNA sequences to reduce or inhibit expression of a native protein in the plant.

The following examples are offered by way of illustration and not by way of limitation.

30

EXPERIMENTAL

Promoter regions for maize genes encoding histone H2B, metallothionein, α -tubulin 3-18, elongation factor e1a (e1a-11, e1a-15, and e1a-16), ribosomal protein rps8, chlorophyll a/b binding protein (cab-10 and cab-20), and glyceraldehyde-3-phosphate gpc4 were isolated from maize plants and cloned. These

genes were selected as sources of constitutive promoters based on the developmental and spatial expression of their gene products. The method for their isolation is described below.

Histones are classified on the basis of their role in chromatin structure. The histone H2B is one of four histones contributing to the nucleosome core. Of the core histones H2A, H2B, H3, and H4, H2B is the least conserved from one species to another. In maize, it exists in the form of multiple variants encoded by a large multigenic family. The maize H2B histones are constitutively expressed at a high level in meristematic tissues throughout the plant. The novel promoter sequence for a maize gene encoding one of these H2B histones is set forth in SEQ ID NO: 1.

Metallothioneins make up a class of proteins that are constitutively expressed in low copy number in cells throughout a plant. Expression of these proteins in mammalian cells is inducible in response to stress and to elevated concentrations of trace elements and hormones. The novel promoter sequence for a maize gene encoding one of these metallothioneins, met-1, is set forth in SEQ ID NO: 2.

Alpha-tubulins play a key role in the cytoskeleton of a plant, contributing to microtubule formation. In maize, these proteins are encoded by a family of genes. Generally, these genes are constitutively expressed in all meristematic tissues, though some are preferentially expressed in certain tissues (see Montoliu *et al.* (1989) *Plant Mol. Biol.* 14:1-15; Montoliu *et al.* (1990) *Gene* 94:201-207). The novel promoter sequence for a maize gene encoding alpha-tubulin 3-18 is set forth in SEQ ID NO: 3.

Elongation factor efla is one of four subunits of the translation elongation factor 1 protein. This subunit plays a role in binding aminoacyl tRNA to the ribosome's acceptor site during protein synthesis. However, it has been found to play a number of other diverse roles in the cell, and hence is a multifunctional protein (see Durso and Cyr (1994) *Protoplasma* 180:99-105 for a review). In maize, efla is encoded by a family of genes comprising at least 6 members (see Berberich *et al.* (1995) *Plant Mol. Biol.* 29:611-615). Novel promoter sequences for three maize genes encoding efla proteins, efla-11, efla-15, and efla-16, are set forth in SEQ ID NOs: 4, 5, and 6, respectively.

Ribosomal proteins contribute to the structure of ribosomes. A novel promoter sequence for a gene encoding a maize ribosomal protein rps8 is set forth in SEQ ID NO: 7.

Chlorophyll a/b binding protein is associated with photosystem II in the thylakoid membranes. Expression of this protein in leaf tissues is elevated in the light. Novel promoter sequences for two maize genes encoding maize chlorophyll a/b binding proteins, cab-10 and cab-20, are set forth in SEQ ID NOs: 8 and 9, respectively. The cab-10 promoter sequence shares homology (97% in an overlap from nt 1-271 of SEQ ID NO: 8) with a maize cab-m7 gene sequence (emb Accession No. X53398/ZMCABM7). The C-terminal end of the cab-10 promoter sequence disclosed herein is unique from nt 272-467. The cab-20 promoter sequence shares homology (98% in an overlap from nt 1-85 of SEQ ID NO:9) with this same cab-m7 gene sequence. The C-terminal end of the cab-20 promoter sequence disclosed herein is unique from nt 86-467. Fragments comprising contiguous nucleotides of the cab-10 and cab-20 promoters disclosed herein are preferably obtained from the unique C-terminal end of these promoter sequences, but fragments may overlap the N-terminal regions of homology, and thus include contiguous nucleotides within this region of overlap.

Glyceraldehyde-3-phosphate dehydrogenase is a glycolytic enzyme. A novel promoter sequence for a maize gene encoding glyceraldehyde-3-phosphate dehydrogenase gpc4 is set forth in SEQ ID NO: 10.

Example 1: Isolation of Promoter Sequences

The procedure for promoter isolation is described in the User Manual for the GenomeWalker kit sold by Clontech Laboratories, Inc., Palo Alto, California. Genomic DNA from maize line A63 was extracted by grinding 10-day old seedling leaves in liquid nitrogen, and the DNA prepared as described by Chen and Dellaporta (1994) in *The Maize Handbook*, ed. Freeling and Walbot (Springer-Verlag, Berlin). RNase A was added to 10 µg/ml and then incubated at 37°C for 1 hr. The DNA was then extracted once with phenol-chloroform, then chloroform, then ethanol precipitated and resuspended in TE (10 mM Tris pH 8.0, 1 mM EDTA). The DNA was then used exactly as described in the GenomeWalker User Manual (Clontech PT3042-1 version PR68687). Briefly, the DNA was digested separately with restriction enzymes DraI, EcoRV, PvuII, ScaI, and StuI, all blunt end cutters. The GenomeWalker adapters were then ligated onto the ends of the restricted DNA. The resulting DNA is referred to as DL1-DL5, respectively.

For isolation of specific promoter regions, two nonoverlapping gene-specific primers (usually 27 bp in length) were designed from the sequences of abundant ESTs in the Pioneer/HGS database. The primers were designed to amplify the region upstream of the coding sequence, *i.e.*, the 5' untranslated region and promoter region of the chosen gene. The sequences of the primers are given below for each promoter described. The first round of PCR was performed on each DNA sample (DL1-5) with Clontech primer AP1 (sequence 5'-gtaatacgactcactatagggc-3'; SEQ ID NO: 11) and the gene-specific primer (gsp)1 with the following sequences:

- 10 histone H2B gsp1 sequence: 5'-cgcgggctcctcctccgccggcttctt-3' (SEQ ID NO: 12)
- metallothionein gsp1 sequence: 5'-gtacttcttgccgcacttgacagcttga-3' (SEQ ID NO: 13)
- alpha-tubulin 3 gsp1 sequence: 5'-ggctctgtcaccgggcatctgaccatc-3' (SEQ ID NO: 14)
- efla gsp1 sequence: 5'-gtcctgtggtggtcgacttgccagagt-3' (SEQ ID NO: 15)
- rps8 gsp1 sequence: 5'-ttcttctccaggccttctgctttcca-3' (SEQ ID NO: 16)
- 15 cab gsp1 sequence: 5'-aggagacgacgacggcacgttcacggc-3' (SEQ ID NO: 17)
- gpc4 sequence: 5'-taatcggtgctgatgaaggggtcgttc-3' (SEQ ID NO: 18)

PCR was performed in a model PTC-100 thermal cycler with HotBonnet from MJ Research (Watertown, Massachusetts) using reagents supplied with the GenomeWalker kit. The following cycle parameters were used: 7 cycles of 94°C for 20 2 sec, then 72°C for 3 min, followed by 32 cycles of 94°C for 2 sec, and 67°C for 3 min. Finally, the samples were held at 67°C for 4 min, then at 4°C until further analysis.

As described in the User Manual, the DNA from the first round of PCR was then diluted and used as a template in a second round of PCR using the Clontech AP2 25 primer (sequence 5'-actatagggcacgcgttgtt-3'; SEQ ID NO: 19) and gene-specific primer (gsp)2 with the following sequences:

- histone H2B gsp2 sequence: 5'-gggcttcttctcggccttgggcgccat-3' (SEQ ID NO: 20)
- 30 metallothionein gsp2 sequence: 5'-agccgcagcttgatccgcagttgcaag-3' (SEQ ID NO: 21)
- alpha-tubulin 3 gsp2 sequence: 5'-cccagcacgcgtttccgacctggatac-3' (SEQ ID NO: 22)
- efla gsp2 sequence: 5'-tggccaataaccacaatgttgatgtg-3' (SEQ ID NO: 23)
- rps8 gsp2 sequence: 5'-cgcttgatgcacgagtcacgcgagata-3' (SEQ ID NO: 24)

cab gsp2 sequence: 5'-gaaggctgtggaggagagggccatggt-3' (SEQ ID NO: 25)

gpc4 gsp2 sequence: 5'-acccattgatcccgatcttgatct-3' (SEQ ID NO: 26)

The cycle parameters for the second round were: 5 cycles of 94°C for 2 sec,
5 then 72°C for 3 min, followed by 20 cycles of 94°C for 2 sec, and 67°C for 3 min.
Finally, the samples were held at 67°C for 4 min, and then held at 4°C.

Approximately 10 µl of each reaction were run on a 0.8% agarose gel, and bands
(usually 500bp or larger) were excised, purified with the Sephaglas BandPrep kit
(Pharmacia, Piscataway, New Jersey), and cloned into the TA vector pCR2.1
10 (Invitrogen, San Diego, California). Clones were sequenced for verification, and then
the mini-prep DNA was diluted 1:30 in water and 1 µl was amplified with gene-
specific primer (gsp)3 (sequences below) and the Clontech AP2 primer with the
following cycle parameters: five cycles of 94°C for 2 sec, 46°C for 30 sec, 72°C for
min, followed by 20 cycles of 94°C for 2 sec, 67°C for 3 min, then 67°C for 4 min,
15 and held at 4°C.

histone H2B gsp3 sequence: 5'-gaccatggtgtcgtgtggatccgatgcggctgct-3' (SEQ ID
NO: 27)

metallothionein gsp3 sequence: 5'-gaccatggtgtcgtgtggatcccttggtgct-3' (SEQ ID
20 NO: 28)

alpha-tubulin gsp3 sequence: 5'-gaccatggtgtcgtgtggatccggtgtgtgaacg-3' (SEQ ID
NO: 29)

efla-11 and efla-15 gsp3 sequence: 5'-gaccatggtgtcgtgtggatccgtgagattgaac-3' (SEQ
ID NO: 30)

efla-16 gsp3 sequence: 5'-gaccatggtgtcgtgtggatccgtgaagcttaa-3' (SEQ ID NO:
25 31)

rps8 gsp3 sequence: 5'-gaccatggtgtcgtgtggatccgcctgctcctgtc-3' (SEQ ID NO: 32)

cab gsp3 sequence: 5'-gaccatggtgtcgtgtggatccctgcactgctac-3' (SEQ ID NO: 33)

gpc4 gsp3 sequence: 5'-gaccatggtgtcgtgtggatccacaaacacaagc-3' (SEQ ID NO: 34)

30

Ten µl of the resulting amplified DNA was run out on a 0.8% agarose gel,
purified with Sephaglas, and cloned into the PCR2.1 TA vector. Final sequences
were determined on the resulting plasmids.

Example 2: Transient Gene Expression Data Using Promoter Sequences

A transient gene expression assay was used to test the cloned DNAs for promoter activity. The promoters were recloned into plasmid PHP3953 (Figure 1) digested with BglII and SphI or with BglII and PvuII to remove the ubiquitin promoter. The new promoter fragments were inserted in place of the ubiquitin promoter such that the ubiquitin 5' untranslated region and intron were now between the test promoter and the GUS reporter gene.

Experiments were performed with immature embryos, essentially the scutellar surface. Immature GS3 maize embryos were isolated from ears 9-11 days after pollination using a scalpel. Prior to embryo isolation, pollinated ears were surface-sterilized with a microdetergent and 25% commercial bleach mixture, then washed with 3 exchanges of sterile H₂O. Isolated immature embryos (1.4-1.7 mm) were placed on bombardment medium and aligned in a target grid in preparation for bombardment.

Immature embryos were then transformed by the tungsten particle biolistic method (Tomes *et al.* (1995) in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); Koziel *et al.* (1993) *Bio/Technology* 11:194-200) using a high pressure particle delivery system (Biolistic Particle Delivery System Model PDS-100 by DuPont). Plasmid DNA comprising a promoter sequence of the invention operably linked to the GUS gene was bombarded into the maize immature embryos. Following culture for 40 hours, bombarded embryos were stained with X-Gluc staining solution (McCabe *et al.* (1988) *Bio/Technology* 6(87):923-926) for 12 h at 37°C in the dark. GUS activity was then measured, using the ubiquitin promoter as a control, by counting blue spots after staining for GUS activity as described elsewhere (see Jefferson (1987) *Plant Mol. Biol. Rep.* 5:387-405). Transient expression data are shown in Table 1.

Table 1: Promoter activity as measured by transient expression of GUS (see Christensen and Quail (1989) *Plant Mol. Biol.* 12:619-632 for details of ubiquitin promoter activity).

<u>Gene</u>	<u>#Blue spots</u>	<u>Average</u>
Ubiquitin (control)	395, 155, 307, 416	318
histone H2B	2127, 2826, 2830	2594
tub3-18 (alpha-tubulin 3)	2542, 1832, 2727	2367
met-1 (methallothionein)	163, 111, 27	100
efla-16 (elongation factor 1a)	136, 2, 0	46
efla-15 (another family member)	81, 38, 25	48

5

Promoter activity of the promoter sequences for the alpha-tubulin and histone H2B genes was significantly stronger than that of the ubiquitin promoter, indicating these as strong constitutive promoters. Promoter activity of the promoter sequence for the metallothionein gene was considerably lower than that of the ubiquitin promoter, indicating this is a weak constitutive promoter.

10

Example 3: Transformation and Regeneration of Transgenic Plants

Immature GS3 maize embryos were bombarded with a plasmid containing the promoter sequence operably linked to the GUS gene plus a plasmid containing the selectable marker gene PAT (Wohlleben *et al.* (1988) *Gene* 70:25-37) that confers resistance to the herbicide Bialaphos. Transformation was performed as described in Example 1. See Appendix for examples of bombardment (560Y), selection (560R), hormone-containing regeneration (288J), and hormone-free (272V) media.

15

One day after bombardment, the embryos were transferred from bombardment medium to selection medium containing 3 mg/liter Bialaphos and subcultured on selection medium every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones were transferred to hormone-containing medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos were transferred to medium for germination and

20

transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets were transferred to hormone-free medium in tubes for 7-10 days until plantlets were well established. Plants were then transferred to inserts in flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic 600 pots (1.6 gallon) and grown to maturity.

Promoter activity in transformed plant tissues was assessed by measuring expression of GUS. Plant tissue samples were placed into the wells of 12-well cell culture clusters and stained with X-Gluc staining solution for 12 h at 37°C in the dark. Samples were then scored for GUS staining. GUS staining score ranged from 0 (negative) – 6 (highest). These staining scores served as a measure of level of GUS expression. Results are shown in Table 2.

Six transformation events have been evaluated for the H2B promoter. Overall, five out of six events scored strong GUS expression in maize plants, with one event scoring median expression. GUS expression was particularly strong in leaf, root, and tassel, with these tissues having GUS staining scores of 5-6 in all events. GUS expression in husk and silk was slightly lower, with these two tissues mostly having staining scores of 4-5. GUS staining with kernels was conducted with event TC7394 and it scored strong. In general, H2B is a strong promoter in maize embryos, callus, and different tissues.

Four transformation events have been evaluated for the met-1 promoter. Overall, three events scored strong GUS expression and one event scored median expression. GUS expression was found in all plant tissues, including leaf, root, husk, silk, and tassel, with these tissues mostly having staining scores of 4-5. The expression pattern seems similar to that observed with the tub3-18 promoter.

Five transformation events have been evaluated for the tub3-18 promoter. Overall, three events scored strong GUS expression and two scored median expression. GUS expression was found in all plant tissues, including leaf, root, husk, silk, and tassel, with these tissues mostly having staining scores of 3-5. GUS staining with kernels was conducted with event TC7406 and it scored very strong.

Five transformation events have been evaluated the efla-15 promoter. Overall, three events scored strong GUS expression, one event scored median expression, and one event scored weak expression. GUS expression was found in all plant tissues,

including leaf, root, husk, silk, and tassel, with these tissues mostly having staining scores of 3-5.

The ubiquitin promoter was used as a control. Five transformation events have been evaluated. Overall, two events scored strong GUS expression and three scored weak expression. GUS expression was found in all plant tissues, including leaf, root, husk, silk, and tassel. With those transformation events exhibiting strong expression, GUS was expressed in all different tissues, with tissues mostly having staining scores of 4-5. With those transformation events exhibiting weak expression, GUS was expressed in leaf, root, and tassel, with these tissues mostly having staining scores of 1-3. There was very little GUS expression in husk and silk tissues in those transformation events that exhibited weak expression.

Table 2. Gus expression as driven by various promoters of the invention.

Promoters	Event	GUS expression	GUS PCR
H2B	TC4703	strong	na
	TC4701	strong	na
	TC4694	strong	na
	TC7392	median	positive
	TC7394	strong	positive
	TC7396	strong	positive
	TC7413	strong	positive
	TC7414	weak	positive
	TC7415	strong	positive
	TC7416	strong	positive
tub3-18	TC7406	strong	positive
	TC7407	strong	positive
	TC7408	strong	positive
	TC7409	median	positive
	TC7410	median	positive
efla-15	TC7418	median	positive
	TC7419	strong	positive
	TC7420	strong	positive
	TC7421	weak	positive
	TC7422	strong	positive
ubi	TC7390	strong	positive
	TC7386	strong	positive
	TC7387	weak	positive
	TC7388	weak	positive
	TC7389	weak	positive

APPENDIX**272 V**

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts (GIBCO 11117-074)	4.300	g
Myo-Inositol	0.100	g
MS Vitamins Stock Solution ##	5.000	ml
Sucrose	40.000	g
Bacto-Agar @	6.000	g

Directions:

- 5 @ = Add after bringing up to volume

Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occurs, then make fresh stock.

- 15 Total Volume (L) = 1.00

WO 99/43797

27

PCT/US99/04203

288 J

Ingredient	Amount	Unit
D-I H ₂ O	950.000	ml
MS Salts	4.300	g
Myo-Inositol	0.100	g
MS Vitamins Stock Solution ##	5.000	ml
Zeatin .5mg/ml	1.000	ml
Sucrose	60.000	g
Gelrite @	3.000	g
Indoleacetic Acid 0.5 mg/ml #	2.000	ml
.1mM Absciscic Acid	1.000	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

- 5 Dissolve ingredients in polished D-I H₂O in sequence

Adjust to pH 5.6

Bring up to volume with polished D-I H₂O after adjusting pH

Sterilize and cool to 60°C.

Add 3.5g/L of Gelrite for cell biology.

- 10 ## = Dissolve 0.100 g of Nicotinic Acid; 0.020 g of Thiamine.HCL; 0.100 g of Pyridoxine.HCL; and 0.400 g of Glycine in 875.00 ml of polished D-I H₂O in sequence. Bring up to volume with polished D-I H₂O. Make in 400 ml portions. Thiamine.HCL & Pyridoxine.HCL are in Dark Desiccator. Store for one month, unless contamination or precipitation occurs, then make fresh stock.

- 15 Total Volume (L) = 1.00

WO 99/43797

28

PCT/US99/04203

560 R

Ingredient	Amount	Unit
D-I Water. Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	30.000	g
2, 4-D 0.5mg/ml	4.000	ml
Gelrite @	3.000	g
Silver Nitrate 2mg/ml #	0.425	ml
Bialaphos 1mg/ml #	3.000	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 Total Volume (L) = 1.00

WO 99/43797

29

PCT/US99/04203

560 Y

Ingredient	Amount	Unit
D-I Water. Filtered	950.000	ml
CHU (N6) Basal Salts (SIGMA C-1416)	4.000	g
Eriksson's Vitamin Mix (1000X SIGMA-1511)	1.000	ml
Thiamine.HCL 0.4mg/ml	1.250	ml
Sucrose	120.000	g
2,4-D 0.5mg/ml	2.000	ml
L-Proline	2.880	g
Gelrite @	2.000	g
Silver Nitrate 2mg/ml #	4.250	ml

Directions:

@ = Add after bringing up to volume

5 # = Add after sterilizing and cooling to temp.

Dissolve ingredients in D-I H₂O in sequence

Adjust to pH 5.8 with KOH

Bring up to volume with D-I H₂O

Sterilize and cool to room temp.

10 ** Autoclave less time because of increased sucrose**

Total Volume (L) = 1.00

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains.

All publications and patent applications are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically
5 and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

WO 99/43797

PCT/US99/04203

1

SEQUENCE LISTING

<110> Rice, Douglas A.

<120> Constitutive Maize Promoters

<130> 5718-33, 035718/175218

<140>

<141>

<150> 60/076,075

<151> 1998-02-26

<160> 34

<170> PatentIn Ver. 2.0

<210> 1

<211> 1392

<212> DNA

<213> Zea mays

<400> 1

```
cgacggcccg ggctggtatc gataaatggt tccacataga ttttgcatac cataatgatg 60
tttgctcgttc cgtatctatg tttcatacaa aattttttacg catatcgcaa cacatgggca 120
catacctagt gactgtataa ctctgcatgt atgagtgtat gactatatga ttagtagtaact 180
aataagaagg gtagacattt gagtgattct tttattcctg gacttgtaag acttgacatt 240
tctgccttga gtgcgataca tcatatggac aggggttatg catacactgc ttgtttgttg 300
tttatgttct aagagcatct ccaacaacgt gacatatgaa aatgccctac aattttaaaaa 360
tggttatatt ttataaaatt tagggcataa ataaaacatc ccgctccaac attaaagcct 420
taaattctatt ataggggaagc ccactatgat atagtatatt tgaggcactt tagaggggtgc 480
cctataatth tttgaccatt tttttatgaa atgagacact attggagtat tttttttccg 540
tagagcacca tatttcaatt tgagacacca atttaaggca ttgttggaga tgttctaaat 600
gttggtttat tttgtctgta tcgttgtggt tttgatagtg gtgcctttgc aatgtacatc 660
ttacattgac aataataata ggtaaaactc taaaaatttt ttatctaata gactcttgta 720
tgaaacattg tacttgacac catctgatgt aaacactgca tacttttaac agtgacaaga 780
ttctgtttca ttttagggct agtttgggaa ccaaatttta ttagggtttt tatttttctaa 840
gaaaaagtaa tttattttac cttgagaaaa tataaattac ttgagaaaat agagttccaa 900
actagctctt atctttgtcg aatcctcctc tattcaaagt tgacatttct ggcacgtgac 960
aactggtgat gttgtagact gtgttaagta atacgtgtca ttattactaa atgccatttt 1020
agtaaagtgt gagtatgtac tctactacag taagtattat tgggtgtattt acactagaca 1080
gttggcggcc tggcgggtaa agttatcctg tagaaagttg ggccaggcca aaaccaaccg 1140
ccaaaggaaa ggccttccgg ccgcgccacc tttgcgcgcc gaaggtcagt tccttcagtc 1200
tcctcccgtc tcagactctg accacgtcga caatccgggc cgaaacacat ctgcaccgtc 1260
cacttgcgac agattgaaca caccatttct atccacgtca gcgatccgtg gcactagccc 1320
ttccaccaat cagcccaagt tgccccttct ctttaaattc gccgcacca ttgctcttct 1380
cacggccata ga 1392
```

WO 99/43797

2

PCT/US99/04203

<210> 2

<211> 1300

<212> DNA

<213> Zea mays

<400> 2

```

gaatataggc aacaccccac actggtaggt aaagctccgg gattgtgact gagatgggga 60
tgatgacgtt gtagagaata caggcagccg tcggcgctac aactctcctc acaaagaaga 120
agctatagag catatagaac ttcttgacga gtgatataat cttagcggaac aacaagtggg 180
gctaattttt agattttccc ctgcgaaagg catgatttgc cagaatggga acctgggcag 240
atatatgctc accttggcaa acaaaacatc ctttgccatc ttacggaata agtttgcacc 300
acctagacca ccggaattgc tgccgacagt aggccttgta agtggacggc agttcactct 360
tgacctgtaa atataacatg tttgcatagt caaatgatca catcaaatat atttaatttt 420
ggcaacactc ttatatgtca aaatctactc actctaactg ctccaacata tacgaatttc 480
cagcccttta gtgttgctcg aaccgccaag tccatgcctt caactgtagt tcggtccttc 540
caacctcctg catctcttat ggctcctgta cgccacactc cagcagttcc tagaatagac 600
ataccatcag gaagaattta tggctctgaa attgacattc tgggggtggtt cccctctttt 660
ggggaatttg tttgtactct ttagagggtt caaaagatgt gtaccgttga aactgaagaa 720
ggcaaaggtc gctgatcctg cttcttggtc aactttgaag tggtagtcgt aaaacatctt 780
ttgtaccctt gtcagcaggc ttgtcgtgtc attcactgca aaattctgtc aattaagaat 840
tggaccgatc agagtgaata aaaatactat atcgctcgag aagcttatgc actagcagtt 900
ctctttttca tgttatatca tctttgaaca gttctatgcc atctttctgg catcatcatt 960
aaagaatgtg aaaatatatt ttattgtcat cattatgaaa cgtgatgcct attttataag 1020
cccaaaacag tgaggttgtc tgttccaagt tccttgtgct atttgtaaga actatctgtg 1080
cacttctcac aagaaaaaca tctgaaaatg gagaaagtat tccattgtgg tggataactg 1140
aatgtggga gagcatatct gacaaatttg acatgagggt taagaaatat ttcccatagt 1200
atttttctat tccataaatt aatcatgtgc atttgttata gacgggtagc tggaaaagtg 1260
tgctcaagaa ctagcaggga atggaccagc caggccgtcg 1300

```

<210> 3

<211> 166

<212> DNA

<213> Zea mays

<400> 3

```

cgacggcccg ggctgggtcca acctccaagc ctacaaccag atcagccggt ctggtactgt 60
gcttctgtgg cacgaaaaaa ccgaccgttg cacggacgag aaaccogaac cgtcgaaaca 120
atcgtaatcc ccagggggtc aaacgcaaaa caccgtccgc tcccct 166

```

<210> 4

<211> 1023

<212> DNA

<213> Zea mays

<400> 4

```

tgtaacaag acgaatcaat gattagaaac actagtaaata taatactacc agtaacgaca 60
caatttatcg gatgggatca gaaactagat ctaacgttgt cgccggagtt atgagcttga 120

```

WO 99/43797

3

PCT/US99/04203

```

aaaaagcttg atacacataa gcagtgtaaa aataaagaag caaatattaa acagagcaat 180
caatgtaata taaaaaataa agcaggcata gaaaataaag ggcaaactga taagcgtgcg 240
cacatgattt tgagccagta agacgaattc aaaattaggc acaacaactc tataaagcag 300
atcaggaatt tggtcattat atgaggttac aactcttact gcgcatgaac tacaaagaca 360
agtatcggtc aaatagaaac tacagcagct ctacacctca aaaattcgat ctgactttat 420
taagaccacg taacacgacc gcaagcgcaa ttaagaagtc gtccaactac ggtggcacgg 480
tgcgtatgta atttatcaaa tccagaagca actatccgta agaatcaggt catccaaaca 540
cggaagcaaa caaaaccaca aaaactaaga gaatggagac gcagtagcta caattaatcg 600
caaacagaag gccgaatcac ttcaggcacg gggaggcaac agagcacagc tttgaaccac 660
tgcctatctc atcgagcacc tcaaggacaa agatcgagcg acagaggatt agctaccttg 720
caagggacag aggccgaaat ggaggcgcac aaggcagacg gcgagttggg gaggaagggtg 780
ctggcctcta gggttcggga ctttatataa ggggtgtcga gtggccggac cagcccaata 840
cttgacttcc gaatggcggc ccagggccca taaacggagc gagacatagt atatgggatg 900
actgaatgac aatgttatth gctcttaata ttgtatggca gcacttttag gtatatgcc 960
caatttctct aatactctac tctttcatat tctaattgca tctcatgtta tgttggccta 1020
aat

```

1023

<210> 5

<211> 1022

<212> DNA

<213> Zea mays

<400> 5

```

gttaacaaga cgaatcaacg attagaaaca ctagtaaata ttaatactac cagtaacgac 60
acaattttatc agatgggatc agaaactaga tctaacgttg tcaccggagt tatgagcttg 120
aaaaaagctt gatacacata agcagtgtaa aaataaagaa gcaaatatta aacagagcaa 180
tcaatgtaat ataaaaaaat aaagcaggca tagaaaataa agagcaaact aataagtgtg 240
cgcacataat tttgagccag taagacgaat tcaaaattag gcacaacaac tctataaagc 300
agatcaggaa tttggtcata ttatgagggtt acaactctta ctgcgcatga actacaaaga 360
caagtatcgg tcaaatagaa actacagcag ctctacacct caaaaattcg atctgacttt 420
attaagacca cgtagcacga ccgcaagcgc aattaagaag tcgtccaact acggtggcac 480
ggtgcgtatg taattttatca aatccagaag caactatccg taagaatcag gtcattccaa 540
cacggaagca aacaaaacca caaaaaacta agagaatgga gacgcagtag ctacaattaa 600
tcgcaaacag aagaccgaat cacttcaggc acggggaggc aacagagcac agctttgaac 660
cactgcctat ctcatcgagc acctcaagga caaagatcga gcgacagagg attagctacc 720
ttgcaaggga cagaggccga aatggaggcg cagaaggcag acggcgagtt ggggaggaag 780
gtgcttgctt ctagggttcg ggactttata taagggtgtc ggtgtggccg gaccagccca 840
atgtgacttc cgattttcga atggcgggcc agggccataa acggagcgag acaggggtgtg 900
tttggttcgg tttttttctg accagcttat atgaaaagct ggttgtgggg aaaagctggc 960
tgttgggaaa aactggtggc taaaatttag gtgtttgggt cgccagacca gcccgggccg 1020
tc

```

1022

<210> 6

<211> 1769

<212> DNA

<213> Zea mays

<400> 6

WO 99/43797

4

PCT/US99/04203

```

cgacggcccg ggctgggtctg acttcagtgt aattcaaagc cggaacagat aaaagtgtgt 60
ctttgccgat gatgttaaat agagtaatac ctgattattg ctgtcaagtg gttgtgttag 120
taattggaat actgttggag tatgtttccc tttatctttc cattttatca tctgtcagtg 180
ctcgtgactt tatggctctt aaattttctta gatagttcat gtcgtacat gaaatgtgtg 240
tctgtaacat tgggtcaacca tattgttcat gtcttcaact ctatttcgaa gtcacaaact 300
gatcacttgg agtttcggac gtgaagcaac gacaggcatc aatgcctact tcatatataa 360
aactgcttgc ttgttgcatt ctgttacagc acctaaagacc taagtacata cttgcttcct 420
tttcttattt gtatgctggg tcccatggac cccgttttgt tttgggtttc cctacacatt 480
gtccccggac ttacagatga caaatatgat gtgttctatg agtcacgtcg aaagaaagaa 540
atgtacttta gcatttcact gtacataagt ccatcgattt cacattatac tcttctacct 600
aaciaagccc aaaattgagc tacacatctt tttgatatta gtacgttaat gttagtcaat 660
tggtatcagg cagcgatgtc tgattataga tctcaaagta agactgtcat tcccccttag 720
ttcagttagt atagttgctt aaattcagca agtctgaata cttcaacaag gcacaataag 780
ggaggttcag cacgtaagtt agtataggac catgtgtaat ctatttaaca gagaaaatag 840
gcataaaaac gactagcctg cagtcgtgcg gatatttgaa tataagtgtt cagatgggcg 900
agcgcccttg tgctttctct attcgctaaa attcttgaaa agcatgtaag cttttcgcaa 960
ggttcgtag cagtgtactc ctgtttgtag tacatgaaca ttagtatttt acttccctca 1020
gttcaccttt cttgtgcttg caatacaata cattattttt tacgggtgaat ttataaacca 1080
tatttggtaa gttatattta tactagccta atgcactcga ctcttctatt tagattaagt 1140
ttggtgaata atttatatgt ttgtgtagcg aaagactcat atttatttag tttttttctt 1200
gtcaaggata ggacataatt tgggtgctttg cttgtgtaat tatgtctacg cacttaaaaa 1260
atggcacttt ccttacgcct tgggtacagt cttcaataat gagtggataa atgtttttatg 1320
aagttgatct tatttttttg agaattcagc attaataatc atgcgccaca ttatccaaag 1380
cttatggtat tctttatatg atttttagtg taataatgtt gtgccactac agggcggtgc 1440
cggctctgat ctggtttata aagtagcaaa ttttgaattt gttttgttgg ctctaaattc 1500
tgtgtgaaca cttattagtc tgctctgttt attcttttagt gcttcttata ttatattgtt 1560
tgctctattg aatgtttcta ggtctgcttt atcccttgct aacatgtacc atgctttttc 1620
taagctgatt acagtagttg caactccaaa tttagtttct aattccatct aataattttt 1680
tttgtagtg gcttttagtat tgtgtgattt ttacttgctg tcctaataat ttattcgttt 1740
gattaacagt ttaagcttca cggatccac 1769

```

<210> 7

<211> 461

<212> DNA

<213> Zea mays

<400> 7

```

ggtacagaag tgagagaagg aaaccctagg tatcgagtgt gtatatagga ggaggggaagg 60
tggggcgcct gggccgcata aaatgtggaa aggctctgaa ccgacaaatt tggatgttta 120
acagacctgt aaatgggtct gtactctaaa atatatagta tagagaatac atgcatcttt 180
ttatctccaa caaaagtctg taaagaatca tctaaagttt agagcacggc atgcatccgc 240
taaaaatgaa gggcgtagcg ttctactcta tacactgtgc ttatttattg tgtttgctca 300
ttcacatgca catgaccggt aaaaatcaat atagaatgtg aaatatggta cactattgat 360
atagaggatg aaatttagag gacattgctg gagatggaga aaatatagag aacaaaattt 420
tttagagagt gctgtaaagg actgagaata ttcttttang g 461

```

<210> 8

<211> 467

WO 99/43797

5

PCT/US99/04203

<212> DNA

<213> Zea mays

<400> 8

```

gtagcgagac ttgcagatcg atggtggcctt gaggtgctgt ggacttgtgg aggcgacgaa 60
ggtattttaa gcgacggagg gaaacaccgc cgcgagcaag ttctatcttg aatgttgcca 120
gaaaataaaa aataaatctg atgcttggat gccattgggt cgtccgggat gctggaacac 180
ttggcgcgac gggagcctca gggggacgat gaggcagatg tgggctgtgg ctgaggctag 240
gaatctccgt ggagaagggt gattgggcgc cgtaaatact actatatacc tttgttccaa 300
gatgtctgtt ttaacttctc taactttaac tgagttcata taaaaaatcc aaacatatta 360
atcaacttan ttataaattt gcctaaattt aaataagctc tacttacgac aaaaccatat 420
taaacgtttt aaaacgtcca caacccccaa ttttgcaatg caaaagg 467

```

<210> 9

<211> 467

<212> DNA

<213> Zea mays

<400> 9

```

gtagcgagac ttgcagatcg atggtggcctt gagatgctgt ggacttgtgg aggcgacgaa 60
ggtattttaa gcgacggagg gaaacctaca atacggagta tgagctaate taggcggcat 120
ctcccagtc ccttgatggt gccaatgaat aattatttag ttcgttttat tggtatcatt 180
tatttttgta agactcttcc actatgtaat aatgattatg acatttatct ctatacactt 240
tgtcattata tgtgatgttc tcctttggcg cacatatgag acgcacccgt ctttatccct 300
taaatttggg tgtgacactc cgaaagccct atacaatcac tataaaatct gccaaagact 360
gggatttgat tgttgctcct agctcgaaag tgacattgta cccagggaat tcggctgccc 420
atttgctat tctggcagat gccttgagaa ttctaaatan gtctccc 467

```

<210> 10

<211> 565

<212> DNA

<213> Zea mays

<400> 10

```

taacagtacc aacttgcaca tcagaagccc cgcgagcaca gaagtgtata gaacatatgg 60
atctgctagc aagtattcct taagaagtat ttctgcttta tacttatagg ccttagctgc 120
catgattggg tcaacaggaa aatccataat ctgatccatc aagtcaacag tcagattggt 180
gaacagagtg gttccattcc ccacaatcca taagcaatgc ctacaacaaa ttgttaaaaa 240
ttaaatagtt ttattatcat gcaccagata aaagattatg aagtaattga agtgaaattt 300
aaatgcttgg cgcagaactt acttggcccc tgtaacaat atatagtata gatatgtgct 360
cgtgcgttgc gatggaagta aaaaatttgt atgaaacatt gatacggaac gacaaaaatc 420
actataatat gcaaaatata cgtggaaaat attatcaata tcacacaaat taattctaaa 480
aagttaattt agaattttta attacagaca aatatgtcga caactaatca gctaatecct 540
tttagcgata ccagcccggg ccgtc 565

```

<210> 11

<211> 22

<212> DNA

WO 99/43797

6

PCT/US99/04203

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR Primer

<400> 11

gtaatacgcac tcactatagg gc

22

<210> 12

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 12

cgcgggctcc tcctccgccg gcttctt

27

<210> 13

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 13

gtacttcttg ccgcacttgc agcttga

27

<210> 14

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 14

ggctcttgatc ccgggcatct gaccatc

27

<210> 15

<211> 27

<212> DNA

<213> Artificial Sequence

WO 99/43797

7

PCT/US99/04203

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 15

gtcctgtggt ggtcgacttg ccagagt

27

<210> 16

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 16

ttcttcctcc aggccttctg ctttcca

27

<210> 17

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 17

aggagacgac gacggcacgt tcacggc

27

<210> 18

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 18

taatcggtgc tgatgaaggg gtcgttc

27

<210> 19

<211> 19

<212> DNA

<213> Artificial Sequence

WO 99/43797

8

PCT/US99/04203

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 19

actatagggc acgcgtggt

19

<210> 20

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: PCR primer

<400> 20

gggcttcttc tcggccttgg gcgccat

27

<210> 21

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 21

agccgcagct tgatccgcag ttgcaag

27

<210> 22

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 22

cccagcacgc gtttccgacc tggatac

27

<210> 23

<211> 26

<212> DNA

<213> Artificial Sequence

WO 99/43797

PCT/US99/04203

9

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 23

tggccaataa ccacaatgtt gatgtg

26

<210> 24

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 24

cgcttgtgca tcgagtcacg cgagata

27

<210> 25

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 25

gaaggctgtg gaggagaggg ccatggt

27

<210> 26

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 26

accattgat ccgatcttg atct

24

<210> 27

<211> 34

<212> DNA

<213> Artificial Sequence

WO 99/43797

PCT/US99/04203

10

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 27

gaccatggtg tcgtgtggat ccgatgcggc tgct

34

<210> 28

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 28

gaccatggtg tcgtgtggat ccccttgtgg tgc

33

<210> 29

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 29

gaccatggtg tcgtgtggat ccggtgttgt tgaacg

36

<210> 30

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 30

gaccatggtg tcgtgtggat ccgtgagatt gaac

34

<210> 31

<211> 36

<212> DNA

<213> Artificial Sequence

WO 99/43797

PCT/US99/04203

11

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 31

gaccatggtg tcgtcgtgtg gatccgtgaa gcttaa

36

<210> 32

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 32

gaccatggtg tcgtgtggat ccgcctgctc cttgtc

36

<210> 33

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR Primer

<400> 33

gaccatggtg tcgtgtggat cctgcactgc tac

33

<210> 34

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Maize
gene-specific PCR primer

<400> 34

gaccatggtg tcgtgtggat ccacaaacac aagc

34

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule having a nucleotide sequence for a promoter that is capable of initiating constitutive transcription in a plant cell,
5 wherein said nucleotide sequence is selected from the group consisting of:
 - a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10;
 - b) a nucleotide sequence selected from the group consisting of sequences deposited as ATCC Accession Nos. _____;
 - 10 c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10; and
 - d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).
- 15 2. A DNA construct comprising a nucleotide sequence of claim 1 operably linked to a heterologous nucleotide sequence of interest.
3. A vector comprising the DNA construct of claim 2.
- 20 4. A host cell having stably incorporated in its genome the DNA construct of claim 2.
5. An isolated nucleic acid molecule having a nucleotide sequence for a promoter that is capable of initiating constitutive transcription in a plant cell,
25 wherein said nucleotide sequence is selected from the group consisting of:
 - a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 8;
 - b) the nucleotide sequence deposited as ATCC Accession No. _____;
 - 30 c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 8, wherein said C-terminal end comprises nucleotides 272-467 of SEQ ID NO: 8; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

6. A DNA construct comprising a nucleotide sequence of claim
5 5 operably linked to a heterologous nucleotide sequence of interest.

7. A vector comprising the DNA construct of claim 6.

8. A host cell having stably incorporated in its genome the DNA
10 construct of claim 6.

9. An isolated nucleic acid molecule having a nucleotide sequence for a promoter that is capable of initiating constitutive transcription in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:

15 a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 9;

b) the nucleotide sequence deposited as ATCC Accession No. _____;

20 c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 9, wherein said C-terminal end comprises nucleotides 86-467 of SEQ ID NO: 9; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

25 10. A DNA construct comprising a nucleotide sequence of claim 9 operably linked to a heterologous nucleotide sequence of interest.

11. A vector comprising the DNA construct of claim 10.

30 12. A host cell having stably incorporated in its genome the DNA construct of claim 10.

13. A method for constitutively expressing a heterologous nucleotide sequence in a plant, said method comprising transforming a plant cell with a DNA construct comprising said heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in a plant cell and
5 regenerating a stably transformed plant from said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

- a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10;
- b) a nucleotide sequence selected from the group consisting of
10 sequences deposited as ATCC Accession Nos. _____;
- c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10; and
- d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b) or c).

15

14. The method of claim 13, wherein said plant is a monocot.

15. The method of claim 14, wherein said monocot is maize.

20 16. The method of claim 13, wherein said plant is a dicot.

17. A method for constitutively expressing a heterologous nucleotide sequence in a plant, said method comprising transforming a plant cell with a DNA construct comprising said heterologous nucleotide sequence operably linked to a
25 promoter that is capable of initiating constitutive transcription in a plant cell and regenerating a stably transformed plant from said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

- a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 8;
- b) the nucleotide sequence deposited as ATCC Accession No.
30 _____;

c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 8, wherein said C-terminal end comprises nucleotides 272-467 of SEQ ID NO: 8; and

d) a nucleotide sequence that hybridizes under stringent
5 conditions to a sequence of a), b), or c).

18. A method for constitutively expressing a heterologous nucleotide sequence in a plant, said method comprising transforming a plant cell with a DNA construct comprising said heterologous nucleotide sequence operably linked to a
10 promoter that is capable of initiating constitutive transcription in a plant cell and regenerating a stably transformed plant from said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 9;

15 b) the nucleotide sequence deposited as ATCC Accession No. _____;

c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 9, wherein said C-terminal end comprises nucleotides 86-467 of SEQ ID NO: 9; and

20 d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

19. A plant cell stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of
25 initiating constitutive transcription in said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10;

b) a nucleotide sequence selected from the group consisting of
30 sequences deposited as ATCC Accession Nos. _____;

c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

20. The plant cell of claim 19, wherein said plant cell is from a
5 monocot.

21. The plant cell of claim 20, wherein said monocot is maize.

22. The plant cell of claim 19, wherein said plant cell is from a dicot.
10

23. A plant cell stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

15 a) a nucleotide sequence comprising the sequence set forth in
SEQ ID NO: 8;

b) the nucleotide sequence deposited as ATCC Accession No.
_____;

c) a nucleotide sequence comprising at least 40 contiguous
20 nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 8,
wherein said C-terminal end comprises nucleotides 272-467 of SEQ ID NO: 8; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

25 24. A plant cell stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in said plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of:

a) a nucleotide sequence comprising the sequence set forth in
30 SEQ ID NO: 9;

b) the nucleotide sequence deposited as ATCC Accession No.
_____;

c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 9, wherein said C-terminal end comprises nucleotides 86-467 of SEQ ID NO: 9; and

d) a nucleotide sequence that hybridizes under stringent
5 conditions to a sequence of a), b), or c).

25. A plant stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in a plant cell, wherein said promoter comprises
10 a nucleotide sequence selected from the group consisting of :

a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10;

b) a nucleotide sequence selected from the group consisting of sequences deposited as ATCC Accession Nos. _____;

15 c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the sequence set forth in SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, or 10; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

20 26. The plant of claim 25, wherein said plant is a monocot.

27. The plant of claim 26, wherein said monocot is maize.

28. The plant of claim 25, wherein said plant is a dicot.

25

29. A plant stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in a plant cell, wherein said promoter comprises a nucleotide sequence selected from the group consisting of :

30 a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 8;

b) the nucleotide sequence deposited as ATCC Accession No. _____;

c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 8, wherein said C-terminal end comprises nucleotides 272-467 of SEQ ID NO: 8; and

5 d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

30. A plant stably transformed with a DNA construct comprising a heterologous nucleotide sequence operably linked to a promoter that is capable of initiating constitutive transcription in a plant cell, wherein said promoter comprises
10 a nucleotide sequence selected from the group consisting of:

a) a nucleotide sequence comprising the sequence set forth in SEQ ID NO: 9;

b) the nucleotide sequence deposited as ATCC Accession No. _____;


15 c) a nucleotide sequence comprising at least 40 contiguous nucleotides of the C-terminal end of the sequence set forth in SEQ ID NO: 9, wherein said C-terminal end comprises nucleotides 86-467 of SEQ ID NO: 9; and

d) a nucleotide sequence that hybridizes under stringent conditions to a sequence of a), b), or c).

20

31. Seed of the plant of any one of claims 25-30.

1/1

**FIG. 1.**

