



(51) International Patent Classification:

C07D 403/12 (2006.01) C07D 409/12 (2006.01)  
C07D 405/12 (2006.01) A01N 43/10 (2006.01)

(21) International Application Number:

PCT/EP2013/058070

(22) International Filing Date:

18 April 2013 (18.04.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12356010.4	20 April 2012 (20.04.2012)	EP
12356015.3	5 July 2012 (05.07.2012)	EP
61/675,001	24 July 2012 (24.07.2012)	US

(71) Applicant: BAYER CROPSCIENCE AG [DE/DE]; Alfred-Nobel-Strasse 50, 40789 Monheim (DE).

(72) Inventors: BRAUN, Christoph; Benderstrasse 18, 40625 Dusseldorf (DE). CRISTAU, Pierre; 4 rue Marietton, F-69009 Lyon (FR). DAHMEN, Peter; AltebruckerstraBe 63, 41470 Neuss (DE). DESBORDES, Philippe; 14 rue du Docteur Mouisset, F-69006 Lyon (FR). ES-SAYED, Mazen; Ricarda-Huch-StraBe 36, 40764 Langenfeld (DE). LACHAISE, Helene; 11 chemin de Montpellas, F-69009 Lyon (FR). RINOLFI, Philippe; Residence le Coteau, Lieu-dit la Colletiere, F-69380 Chatillon-d'Azergues (FR). SCHMIDT, Jan-Peter; 32 chemin de la Begonniere, F-69230 Saint-Genis-Laval (FR). TSUCHYIA, Tomoh; 2 rue Rivet, F-69001 Lyon (FR). VORS, Jean-Pierre; 6 allée Adanson 691 10, F-691 10 Sainte-Foy-Les-Lyon (FR). WACHENDORFF-NEUMANN, Ulrike; Oberer Markeweg 85, 56566 Neuwied (DE).

(74) Agent: GUITTON, Carole; Bayer SAS, Patents & Licensing, 14 impasse Pierre Baizet, F-69009 Lyon (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.1 7(H))

Published:

— with international search report (Art. 21(3))

(54) Title: N-CYCLOALKYL-N-[(HETEROCYCLYLPHENYL)METHYLENE]-(THIO)CARBOXAMIDE DERIVATIVES

(57) Abstract: The present invention relates to fungicidal N-cycloalkyl-N-[(heterocyclylphenyl)methylene] carboxamide derivatives and their thiocarbonyl derivatives, their process of preparation and intermediate compounds for their preparation, their use as fungicides, particularly in the form of fungicidal compositions and methods for the control of phytopathogenic fungi of plants using these compounds or their compositions.




**N-CYCLOALKYL-N-[(HETEROCYCLYLPHENYL)METHYLENE]-(THIO)CARBOXAMIDE  
DERIVATIVES**

**DESCRIPTION**

5 The present invention relates to fungicidal N-cycloalkyl-N-[(heterocyclylphenyl)methylene] carboxamide derivatives and their thiocarbonyl derivatives, their process of preparation and intermediate compounds for their preparation, their use as fungicides, particularly in the form of fungicidal compositions and methods for the control of phytopathogenic fungi of plants using these compounds or their compositions.


10

In international patent application WO-2007/087906 certain N-cycloalkyl-N-benzyl-carboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:




15 wherein A represents a carbo-linked, partially saturated or unsaturated, 5-membered heterocyclyl group, Z<sup>1</sup> represents a substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group, n is equal to 1 to 5 and X can represent various substituents among which a substituted or non-substituted pyridinyl or pyridinyloxy group. However, there is no explicit disclosure or suggestion to select in this document of any such derivative wherein X can represent a substituted or non-substituted heterocyclyl group.

20 In international patent application WO-2009/01 6220 certain N-cycloalkyl-N-benzyl-thiocarboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:




25 wherein A represents a carbo-linked, partially saturated or unsaturated, 5-membered heterocyclyl group, T can represent S, Z<sup>1</sup> represents a substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group, n is equal to 1 to 5 and X can represent various substituents among which a substituted or non-substituted pyridinyl or pyridinyloxy group. However, there is no explicit disclosure or suggestion to select in this document of any such derivative wherein X can represent a substituted or non-substituted heterocyclyl group.

30 In international patent application WO-2010/1 30767 certain N-cycloalkyl-N-benzyl-carboxamides or thiocarboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:



wherein X<sup>1</sup> and X<sup>2</sup> represent a fluorine or a chlorine atom, T can represent O or S, Z<sup>1</sup> represents a substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group, Y can represent CR<sup>5</sup> and R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> or R<sup>5</sup> can, independently, represent various substituents among which a substituted or non-substituted pyridinyl or pyridinyloxy group. However, there is no explicit disclosure or suggestion to select in this document of any such derivative wherein R<sup>1</sup> or R<sup>2</sup> or R<sup>3</sup> or R<sup>4</sup> or R<sup>5</sup> can represent a substituted or non-substituted heterocycyl group.


In international patent application WO-2009/024342 certain N-(substituted phenyl)methylene-carboxamides are generically embraced in a broad disclosure of numerous compounds of the following formula:



wherein R<sub>1</sub> represents a C<sub>1</sub>-C<sub>4</sub>-alkyl group, R<sub>2</sub> represents a C<sub>i</sub>-C<sub>4</sub>-haloalkyl group, B can represent a phenyl group that can be substituted by various substituents among which a substituted or non substituted phenyl group or a substituted or non substituted phenoxy group, and R<sub>15</sub> can represent hydrogen or a C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group with the proviso that B is different from phenyl if R<sub>15</sub> represents a C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group. Therefore, there is no disclosure in this document of any such derivative wherein R<sub>15</sub> can represent a C<sub>3</sub>-C<sub>7</sub>-cycloalkyl group and B be simultaneously a substituted phenyl group.

20

Accordingly, the present invention provides a N-cycloalkyl-N-[(heterocyclylphenyl)methylene](thio)carboxamide of formula (I)



(I)

25 wherein

- A represents a carbo-linked, unsaturated or partially saturated, 5-membered heterocyclyl group that can be substituted by up to four groups R that can be the same or different ;
- T represents O or S ;
- n represents 0, 1, 2, 3 or 4 ;
- 5 • L represents a direct bond, CZ<sup>4</sup>Z<sup>5</sup>, O, S, SO, SO<sub>2</sub> or NZ<sup>6</sup> ; B represents a saturated, partially saturated or unsaturated, monocyclic or fused bicyclic 3-, 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered ring comprising from 1 up to 4 heteroatoms selected in the list consisting of N, O, S, that can be substituted by up to 6 groups Y which can be the same or different ; with the proviso that B does not represent a pyridyl ring when L represents a direct bond or an oxygen atom
- 10 • Z<sup>1</sup> represents a non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl or a C<sub>3</sub>-C<sub>7</sub>-cycloalkyl substituted by up to 10 atoms or groups that can be the same or different and that can be selected in the list consisting of halogen atoms, cyano, C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, C<sub>1</sub>-C<sub>8</sub>-alkoxy, C<sub>1</sub>-C<sub>8</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>8</sub>-halogenoalkoxycarbonyl comprising up to 9 halogen atoms that can be the same or different, C<sub>1</sub>-C<sub>8</sub>-alkylaminocarbonyl or di-Ci-C<sub>8</sub>-alkylaminocarbonyl ;
- 15 • Z<sup>2</sup> and Z<sup>3</sup>, which can be the same or different, represent a hydrogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyl ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyl ; cyano ; isonitrile ; nitro ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxy ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyloxy ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfanyl ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfinyl ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylamino ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbamoyl ; substituted or non-substituted di-CrC<sub>8</sub>-alkylcarbamoyl ; or substituted or non-substituted N-C<sub>1</sub>-C<sub>8</sub>-alkyl-Ci-C<sub>8</sub>-alkoxy-carbamoyl ; or
- 20 • Z<sup>2</sup> and Z<sup>3</sup> together with the carbon atom to which they are linked can form a substituted or non-substituted C<sub>3</sub>-C<sub>7</sub> cycloalkyl ; or
- 30 • Z<sup>3</sup> and the substituent X vicinal to the point of attachment of the phenyl ring, together with the consecutive carbon atoms to which they are linked, can form a substituted or non-substituted 5-, 6- or 7-membered, partly saturated, carbo- or heterocycle comprising up to 3 heteroatoms and Z<sup>2</sup> is as herein-described ; Z<sup>4</sup> and Z<sup>5</sup> independently represent a hydrogen atom ; a halogen atom ; cyano ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxy ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfanyl ; or substituted or non-substituted Ci-C<sub>8</sub>-alkoxycarbonyl ; or
- 35

- Z<sup>4</sup> and Z<sup>5</sup> together with the carbon atom to which they are linked can form a C(=O) carbonyl group ;
- Z<sup>6</sup> represents a hydrogen atom ; a substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkyl ; a C<sub>1-C<sub>8</sub></sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>2-C<sub>8</sub></sub>-alkenyl ; C<sub>2-C<sub>8</sub></sub>-halogenoalkenyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>3-C<sub>8</sub></sub>-alkynyl ; C<sub>3-C<sub>8</sub></sub>-halogenoalkynyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>3-C<sub>7</sub></sub>-cycloalkyl ; C<sub>3-C<sub>7</sub></sub>-halogeno-cycloalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>3-C<sub>7</sub></sub>-cycloalkyl-CrC<sub>8</sub>-alkyl ; formyl ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-halogenoalkylcarbonyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkoxy carbonyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkoxy carbonyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkylsulfonyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkylsulfonyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted benzyl ; or substituted or non-substituted phenylsulfonyl ; X independently represents a halogen atom ; nitro ; cyano ; isonitrile ; hydroxy ; amino ; sulfanyl ; pentafluoro-A<sup>6</sup>-sulfanyl ; formyl ; formyloxy ; formylamino ; substituted or non-substituted (hydroxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (Ci-C<sub>8</sub>-alkoxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (C<sub>2-C<sub>8</sub></sub>-alkenyloxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (C<sub>2-C<sub>8</sub></sub>-alkynyloxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (benzyloxyimino)-Ci-C<sub>8</sub>-alkyl ; carboxy ; carbamoyl ; N-hydroxycarbamoyl ; carbamate ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2-C<sub>8</sub></sub>-alkenyl ; C<sub>2-C<sub>8</sub></sub>-halogenoalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2-C<sub>8</sub></sub>-alkynyl ; C<sub>2-C<sub>8</sub></sub>-halogenoalkynyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkoxy ; C<sub>1-C<sub>8</sub></sub>-halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkylsulfanyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkylsulfanyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfonyl ; Ci-C<sub>8</sub>-halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkylamino ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>2-C<sub>8</sub></sub>-alkenyloxy ; C<sub>2-C<sub>8</sub></sub>-halogenoalkenyloxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>3-C<sub>8</sub></sub>-alkynyloxy ; C<sub>3-C<sub>8</sub></sub>-halogenoalkynyloxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>3-C<sub>7</sub></sub>-cycloalkyl ; C<sub>3-C<sub>7</sub></sub>-halogenocycloalkyl having 1 to 9 halogen atoms ; substituted or non-substituted (C<sub>3-C<sub>7</sub></sub>-cycloalkyl)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>4-C<sub>7</sub></sub>-cycloalkenyl ; C<sub>4-C<sub>7</sub></sub>-halogenocycloalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted (C<sub>3-C<sub>7</sub></sub>-cycloalkyl)-C<sub>2-C<sub>8</sub></sub>-alkenyl ; substituted or non-substituted (C<sub>3-C<sub>7</sub></sub>-cycloalkyl)-C<sub>2-C<sub>8</sub></sub>-alkynyl ; substituted or non-substituted tri(CrC<sub>8</sub>)alkylsilyl ; substituted or non-substituted tri(Ci-C<sub>8</sub>)alkylsilyl-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>1-C<sub>8</sub></sub>-alkylcarbonyl ; C<sub>1-C<sub>8</sub></sub>-halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted

- or non-substituted  $\text{CrC}_8$ -alkylcarbonyloxy ;  $\text{C}_1$ - $\text{C}_8$ -halogenoalkylcarbonyloxy having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkylcarbonylamino ;  $\text{CrC}_8$ -halogenoalkylcarbonylamino having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkoxycarbonyl ;  $\text{CrC}_8$ -halogenoalkoxycarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{Ci-C}_8$ -alkyloxycarbonyloxy ;  $\text{Ci-C}_8$ -halogenoalkoxycarbonyloxy having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylcarbamoyle ; substituted or non-substituted di- $\text{CrC}_8$ -alkylcarbamoyle ; substituted or non-substituted  $\text{CrC}_8$ -alkylaminocarbonyloxy ; substituted or non-substituted di- $\text{Ci-C}_8$ -alkylaminocarbonyloxy ; substituted or non-substituted N-( $\text{CrC}_8$ -alkyl)hydroxy carbamoyle ; substituted or non-substituted  $\text{CrC}_8$ -alkoxycarbamoyle ; substituted or non-substituted N-( $\text{Ci-C}_8$ -alkyl)- $\text{CrC}_8$ -alkoxycarbamoyle ; aryl that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{CrC}_8$ -alkyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{C}_2$ - $\text{C}_8$ -alkenyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{C}_2$ - $\text{C}_8$ -alkynyl that can be substituted by up to 6 groups Q which can be the same or different ; aryloxy that can be substituted by up to 6 groups Q which can be the same or different ; arylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; arylamino that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{CrC}_8$ -alkyloxy that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{Ci-C}_8$ -alkylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl- $\text{CrC}_8$ -alkylamino that can be substituted by up to 6 groups Q which can be the same or different ; pyridinyl which can be substituted by up to 4 groups Q ; or pyridinyloxy which can be substituted by up to 4 groups Q ;
- Y independently represents a halogen atom ; cyano ; hydroxy ; amino ; sulfanyl ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkyl ;  $\text{CrC}_8$ -halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{Ci-C}_8$ -alkoxy ;  $\text{Ci-C}_8$ -halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylsulfanyl ;  $\text{C}_1$ - $\text{C}_8$ -halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylsulfanyl ;  $\text{C}_1$ - $\text{C}_8$ -halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylsulfonyl ;  $\text{C}_1$ - $\text{C}_8$ -halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylamino ; substituted or non-substituted di- $\text{Ci-C}_8$ -alkylamino ; substituted or non-substituted  $\text{CrC}_8$ -alkylcarbonyl ;  $\text{CrC}_8$ -halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkoxycarbonyl ;  $\text{CrC}_8$ -halogenoalkoxycarbonyl having 1 to 9 halogen atoms ; or aryl that can be substituted by up to 6 groups Q which can be the same or different ;
  - Q independently represents a halogen atom, cyano, nitro, substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkyl,  $\text{C}_1$ - $\text{C}_8$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, substituted or non-substituted  $\text{Ci-C}_8$ -alkoxy,  $\text{C}_1$ - $\text{C}_8$ -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, substituted or non-substituted  $\text{CrC}_8$ -alkylsulfanyl,  $\text{C}_1$ - $\text{C}_8$ -halogenoalkylsulfanyl comprising up to 9 halogen atoms that can be the same or different, substituted or non-substituted tri( $\text{C}_1$ - $\text{C}_8$ )alkylsilyl, substituted or non-

substituted tri(Ci-C<sub>8</sub>)alkylsilyl-Ci-C<sub>8</sub>-alkyl, substituted or non-substituted (Ci-C<sub>8</sub>-alkoxyimino)-C<sub>1</sub>-C<sub>8</sub>-alkyl, or substituted or non-substituted (benzyloxyimino)-Ci-C<sub>8</sub>-alkyl ;

- R independently represents hydrogen atom ; halogen atom ; nitro ; cyano ; hydroxy ; amino ; sulfanyl ; pentafluoro-A<sup>6</sup>-sulfanyl ; substituted or non-substituted (Ci-C<sub>8</sub>-alkoxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (benzyloxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted Ci-C<sub>8</sub>-alkyl ; Ci-C<sub>8</sub>-halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyl ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyl ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkynyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkoxy ; CrC<sub>8</sub>-halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkylsulfanyl ; CrC<sub>8</sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkylsulfanyl ; CrC<sub>8</sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfonyl ; CrC<sub>8</sub>-halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkylamino ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; substituted or non-substituted C<sub>3</sub>-C<sub>8</sub>-alkynyloxy ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; C<sub>3</sub>-C<sub>7</sub>-halogenocycloalkyl having 1 to 9 halogen atoms ; substituted or non-substituted tri(Ci-C<sub>8</sub>)alkylsilyl ; substituted or non-substituted CrC<sub>8</sub>-alkylcarbonyl ; CrC<sub>8</sub>-halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkoxycarbonyl ; CrC<sub>8</sub>-halogenoalkoxycarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbamoyl ; substituted or non-substituted di-CrC<sub>8</sub>-alkylcarbamoyl ; phenoxy ; phenylsulfanyl ; phenylamino ; benzyloxy ; benzylsulfanyl ; or benzylamino ;

as well as its salts, N-oxides, metal complexes, metalloid complexes and optically active isomers or geometric isomers thereof.

Unless indicated otherwise, a group or a substituent that is substituted according to the invention can be substituted by one or more of the following groups or atoms: a halogen atom ; nitro ; hydroxyl ; cyano ; isonitrile ; amino ; sulfanyl ; a pentafluoro-λ<sup>6</sup>-sulfanyl group ; formyl ; formyloxy ; formylamino ; carbamoyl ; N-hydroxycarbamoyl ; carbamate ; (hydroxyimino)-Ci-C<sub>6</sub>-alkyl ; Ci-C<sub>8</sub>-alkyl ; a tri(CrC<sub>8</sub>-alkyl)silyl ; C<sub>3</sub>-C<sub>8</sub>-cycloalkyl ; CrC<sub>8</sub>-halogenoalkyl having 1 to 5 halogen atoms ; a C<sub>3</sub>-C<sub>8</sub>-halogenocycloalkyl having 1 to 5 halogen atoms ; C<sub>2</sub>-C<sub>8</sub>-alkenyl ; C<sub>2</sub>-C<sub>8</sub>-alkynyl ; C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; C<sub>2</sub>-C<sub>8</sub>-alkynyloxy ; CrC<sub>8</sub>-alkylamino ; di-CrC<sub>8</sub>-alkylamino ; CrC<sub>8</sub>-alkoxy ; CrC<sub>8</sub>-halogenoalkoxy having 1 to 5 halogen atoms ; CrC<sub>8</sub>-alkylsulfanyl ; CrC<sub>8</sub>-halogenoalkylsulfanyl having 1 to 5 halogen atoms ; C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkenyloxy having 1 to 5 halogen atoms ; C<sub>3</sub>-C<sub>8</sub>-alkynyloxy ; C<sub>3</sub>-C<sub>8</sub>-halogenoalkynyloxy having 1 to 5 halogen atoms ; CrC<sub>8</sub>-alkylcarbonyl ; CrC<sub>8</sub>-halogenoalkylcarbonyl having 1 to 5 halogen atoms ; CrC<sub>8</sub>-alkylcarbamoyl ; di-CrC<sub>8</sub>-alkylcarbamoyl ; N-CrC<sub>8</sub>-alkyloxycarbamoyl ; CrC<sub>8</sub>-alkoxycarbamoyl ; N-CrC<sub>8</sub>-alkyl-Ci-C<sub>8</sub>-alkoxycarbamoyl ; CrC<sub>8</sub>-alkoxycarbonyl ; CrC<sub>8</sub>-halogenoalkoxycarbonyl having 1 to 5 halogen atoms ; CrC<sub>8</sub>-alkylcarbonyloxy ; CrC<sub>8</sub>-halogenoalkylcarbonyloxy having 1 to 5 halogen atoms ; Ci-C<sub>8</sub>-alkylcarbonylamino ; Ci-C<sub>8</sub>-halogenoalkylcarbonylamino having 1 to 5 halogen atoms ; C<sub>1</sub>-

$C_{5-}$ -alkylaminocarbonyloxy ; di-Ci- $C_{8-}$ -alkylaminocarbonyloxy ;  $C_{1-C_{8-}}$ -alkyloxycarbonyloxy ;  $C_{rC_{8-}}$ -alkylsulfanyl ;  $C_{1-C_{8-}}$ -halogenoalkylsulfanyl having 1 to 5 halogen atoms ;  $C_{rC_{8-}}$ -alkylsulfanyl ;  $C_{rC_{8-}}$ -halogenoalkylsulfanyl having 1 to 5 halogen atoms ; Ci- $C_{8-}$ -alkylsulfonyl ; Ci- $C_{8-}$ -halogenoalkylsulfonyl having 1 to 5 halogen atoms ;  $C_{1-C_{8-}}$ -alkylaminosulfamoyl ; di-Ci- $C_{8-}$ -alkylaminosulfamoyl ;  
 5 (C<sub>1-C<sub>6</sub></sub>-alkoxyimino)-Ci- $C_{6-}$ -alkyl ; (C<sub>1-C<sub>6</sub></sub>-alkenyloxyimino)-Ci- $C_{6-}$ -alkyl ; (C<sub>1-C<sub>6</sub></sub>-alkynyloxyimino)-Ci- $C_{6-}$ -alkyl ; 2-oxopyrrolidin-1-yl ; (benzyloxyimino)-Ci- $C_{6-}$ -alkyl ;  $C_{1-C_{8-}}$ -alkoxyalkyl ;  $C_{rC_{8-}}$ -halogenoalkoxyalkyl having 1 to 5 halogen atoms ; benzyloxy ; benzylsulfanyl ; benzylamino ; aryloxy ; arylsulfanyl or arylamino.

10 According to the invention, the following generic terms are generally used with the following meanings:

- halogen means fluorine, chlorine, bromine or iodine ,  
 carboxy means -C(=O)OH ;  
 carbonyl means -C(=O)- ;  
 carbamoyl means -C(=O)NH<sub>2</sub> ;  
 15 N-hydroxycarbamoyl means -C(=O)NHOH ;  
 SO represents a sulfoxide group ;  
 SO<sub>2</sub> represents a sulfone group ;  
 heteroatom means sulfur, nitrogen or oxygen ;  
 methylene means the diradical -CH<sub>2</sub>- ;
- 20 • an alkyl group, an alkenyl group and an alkynyl group as well as moieties containing these terms, can be linear or branched ;
- halogenated groups, notably haloalkyl, haloalkoxy and cycloalkyl groups, can comprise up to nine identical or different halogen atoms ;
- the term "aryl" means phenyl or naphthyl ;
- 25 • In the case of an amino group or the amino moiety of any other amino-containing group, substituted by two substituents that can be the same or different, the two substituents together with the nitrogen atom to which they are linked can form a heterocyclyl group, preferably a 5- to 7-membered heterocyclyl group, that can be substituted or that can include other heteroatoms, for example a morpholino group or piperidinyl group.
- 30 • Where a compound of the invention can be present in tautomeric form, such a compound is understood hereinabove and hereinbelow also to include, where applicable, corresponding tautomeric forms, even when these are not specifically mentioned in each case.

Any of the compounds of the present invention can exist in one or more optical or chiral isomer forms  
 35 depending on the number of asymmetric centres in the compound. The invention thus relates equally to all the optical isomers and to their racemic or scalemic mixtures (the term "scalemic" denotes a mixture of enantiomers in different proportions) and to the mixtures of all the possible stereoisomers, in all proportions. The diastereoisomers and/or the optical isomers can be separated according to the methods which are known *per se* by the man ordinary skilled in the art.

40 Any of the compounds of the present invention can also exist in one or more geometric isomer forms


depending on the number of double bonds in the compound. The invention thus relates equally to all geometric isomers and to all possible mixtures, in all proportions. The geometric isomers can be separated according to general methods, which are known *per se* by the man ordinary skilled in the art.

5 Any of the compounds of the present invention can also exist in one or more geometric isomer forms depending on the relative position (syn/anti or cis/trans) of the substituents of the chain or ring. The invention thus relates equally to all syn/anti (or cis/trans) isomers and to all possible syn/anti (or cis/trans) mixtures, in all proportions. The syn/anti (or cis/trans) isomers can be separated according to general methods, which are known *per se* by the man ordinary skilled in the art.

10

Preferred compounds according to the invention are compounds of formula (I) wherein A is selected in the list consisting of:

- a heterocycle of formula (A<sup>1</sup>)



(A<sup>1</sup>)


15

wherein :

R<sup>1</sup> to R<sup>3</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; CrC<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted CrC<sub>5</sub>-alkoxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

20

- a heterocycle of formula (A<sup>2</sup>)



(A<sup>2</sup>)


25

wherein :

R<sup>4</sup> to R<sup>6</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; CrC<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted Ci-Cs-alkoxy or Ci-Cs-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

30

- a heterocycle of formula (A<sup>3</sup>)




(A<sup>3</sup>)

wherein :

R<sup>7</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkoxy or C<sub>i</sub>-C<sub>s</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>8</sup> represents a hydrogen atom or a substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;


- a heterocycle of formula (A<sup>4</sup>)

(A<sup>4</sup>)

wherein :

R<sup>9</sup> to R<sup>11</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkyl ; amino ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkoxy ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>i</sub>-C<sub>s</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>5</sup>)

(A<sup>5</sup>)


wherein :

R<sup>12</sup> and R<sup>13</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; amino ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>14</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; amino ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>i</sub>-C<sub>s</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>6</sup>)

10

(A<sup>6</sup>)


wherein :

R<sup>15</sup> represents a hydrogen atom ; a halogen atom ; a cyano ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>16</sup> and R<sup>18</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxycarbonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>17</sup> represent a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;

- a heterocycle of formula (A7)


(A<sup>7</sup>)

wherein :

R<sup>19</sup> represents a hydrogen atom or a C<sub>1</sub>-C<sub>5</sub>-alkyl

R<sup>20</sup> to R<sup>22</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A8)


(A<sup>8</sup>)

wherein :

R<sup>23</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>24</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different;

- a heterocycle of formula (A<sup>9</sup>)




wherein :

5 R<sup>25</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>26</sup> represents a hydrogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

10 - a heterocycle of formula (A<sup>10</sup>)




wherein :

15 R<sup>27</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>28</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(C<sub>1</sub>-C<sub>5</sub>-alkyl)amino ;

20

- a heterocycle of formula (A<sup>11</sup>)



wherein :

25 R<sup>29</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

30 R<sup>30</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(C<sub>1</sub>-C<sub>5</sub>-alkyl)amino ;


- a heterocycle of formula (A<sup>12</sup>)

(A<sup>12</sup>)

wherein :

- 5 R<sup>31</sup> represents a hydrogen atom or a substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl  
 R<sup>32</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  
 R<sup>33</sup> represents a hydrogen atom ; a halogen atom ; a nitro ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms  
 10 that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>13</sup>)


(A<sup>13</sup>)

wherein :

- R<sup>34</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>3</sub>-C<sub>8</sub>-cycloalkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; substituted or  
 20 non-substituted C<sub>2</sub>-C<sub>5</sub>-alkynyloxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;  
 R<sup>35</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; a cyano ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or  
 25 different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(C<sub>1</sub>-C<sub>5</sub>-alkyl)amino;  
 R<sup>36</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;

30 -a heterocycle of formula (A<sup>14</sup>)

13


(A<sup>14</sup>)

wherein :

R<sup>37</sup> and R<sup>38</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ;  
 5 substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that  
 can be the same or different ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy or a substituted or non-  
 substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ;

R<sup>39</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;


10 - a heterocycle of formula (A<sup>15</sup>)

(A<sup>15</sup>)

wherein :

R<sup>40</sup> and R<sup>41</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ;  
 15 substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms  
 that can be the same or different ;

- a heterocycle of formula (A<sup>16</sup>)

(A<sup>16</sup>)


20

wherein :

R<sup>42</sup> and R<sup>43</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ;  
 substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that  
 can be the same or different or amino ;

25


-a heterocycle of formula (A<sup>17</sup>)

(A<sup>17</sup>)

wherein :

R<sup>44</sup> and R<sup>45</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

5 - a heterocycle of formula (A<sup>18</sup>)




wherein :

10 R<sup>47</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>46</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ;

15 - a heterocycle of formula (A<sup>19</sup>)



wherein :

20 R<sup>49</sup> and R<sup>48</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>20</sup>)


25



wherein :

30 R<sup>50</sup> and R<sup>51</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;


-a heterocycle of formula (A<sup>21</sup>)

(A<sup>21</sup>)

wherein :

- 5 R<sup>52</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.


-a heterocycle of formula (A<sup>22</sup>)

(A<sup>22</sup>)

wherein :

R<sup>53</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.

15 -a heterocycle of formula (A<sup>23</sup>)


(A<sup>23</sup>)

wherein :

- 20 R<sup>54</sup> and R<sup>56</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  
R<sup>55</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;

-a heterocycle of formula (A<sup>24</sup>)

25

(A<sup>24</sup>)


wherein :

$R^{57}$  and  $R^{59}$  that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted  $C_{1-5}$ -alkyl or  $C_{1-5}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

$R^{58}$  represents a hydrogen atom or substituted or non-substituted  $C_{1-5}$ -alkyl ;

5

-a heterocycle of formula (A<sup>25</sup>)


(A<sup>25</sup>)

wherein :

10  $R^{60}$  and  $R^{61}$  that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted  $C_{1-5}$ -alkyl or  $C_{1-5}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

$R^{62}$  represents a hydrogen atom or substituted or non-substituted  $C_{1-5}$ -alkyl ;

15 -a heterocycle of formula (A<sup>26</sup>)

(A<sup>26</sup>)

wherein :

20  $R^{65}$  represents a hydrogen atom ; a halogen atom ; substituted or non-substituted  $C_{1-5}$ -alkyl ; substituted or non-substituted  $C_{3-5}$ -cycloalkyl ;  $C_{1-5}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted d-C<sub>5</sub>-alkoxy ; substituted or non-substituted  $C_{2-5}$ -alkynyloxy or  $C_{1-5}$ -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

25  $R^{63}$  represents a hydrogen atom ; a halogen atom ; substituted or non-substituted  $C_{1-5}$ -alkyl ; a cyano ; substituted or non-substituted  $C_{1-5}$ -alkoxy ; substituted or non-substituted  $C_{1-5}$ -alkylsulfanyl ;  $C_{1-5}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  $C_{1-5}$ -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted  $C_{1-5}$ -alkylamino or di( $C_{1-5}$ -alkyl)amino;

$R^{64}$  represents a hydrogen atom or substituted or non-substituted  $C_{1-5}$ -alkyl.

30

More preferred compounds according to the invention are compounds of formula (I) wherein A is selected in the list consisting of A<sup>2</sup> ; A<sup>5</sup> ; A<sup>6</sup> ; A<sup>10</sup> and A<sup>13</sup> as herein-defined.

Even more preferred compounds according to the invention are compounds of formula (I) wherein A

represents  $A^{13}$  wherein  $R^{34}$  represents a substituted or non-substituted  $C_{1-5}$ -alkyl,  $C_{1-5}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted  $C_{1-5}$ -alkoxy ;  $R^{35}$  represents a hydrogen atom or a halogen atom and  $R^{36}$  represents a substituted or non-substituted  $C_{1-5}$ -alkyl.

5

Even more preferred compounds according to the invention are compounds of formula (I) wherein A represents  $A^{13}$  wherein  $R^{34}$  represents  $C_{1-5}$ -alkyl,  $C_{1-5}$ -halogenoalkyl comprising up to 3 halogen atoms that can be the same or different ;  $R^{35}$  represents a hydrogen atom ; a chlorine atom ; or a fluorine atom ; and  $R^{36}$  represents a methyl.

10

Other preferred compounds according to the invention are compounds of formula (I) wherein T represents O.

Other preferred compounds according to the invention are compounds of formula (I) wherein  $Z^1$  represents a substituted or non-substituted cyclopropyl.

15

Other more preferred compounds according to the invention are compounds of formula (I) wherein  $Z^1$  represents a non-substituted cyclopropyl or a 2- $C_{1-5}$ -alkylcyclopropyl.

Other even more preferred compounds according to the invention are compounds of formula (I) wherein  $Z^1$  represents a non-substituted cyclopropyl.

20

Other even more preferred compounds according to the invention are compounds of formula (I) wherein  $Z^1$  represents a 2-methylcyclopropyl.

25

Other preferred compounds according to the invention are compounds of formula (I) wherein  $Z^2$  and  $Z^3$  independently represent a hydrogen atom or a methyl.

More preferred compounds according to the invention are compounds of formula (I) wherein  $Z^2$  represents a hydrogen atom and  $Z^3$  represents a hydrogen atom or a methyl.

30

Other preferred compounds according to the invention are compounds of formula (I) wherein n represents 0, 1 or 2.

Other preferred compounds according to the invention are compounds of formula (I) wherein L represents a direct bond.

35

Other preferred compounds according to the invention are compounds of formula (I) wherein L represents an oxygen atom.

40

Other preferred compounds according to the invention are compounds of formula (I) wherein L represents a methylene or a carbonyl group.

5 Other preferred compounds according to the invention are compounds of formula (I) wherein B represents a substituted or non-substituted oxiranyl ring; a substituted or non-substituted pyrrolyl ring; a substituted or non-substituted pyrazolyl ring; a substituted or non-substituted thienyl ring; a substituted or non-substituted benzothienyl ring; a substituted or non-substituted quinoliny ring; a substituted or non-substituted isoquinoliny ring ; a substituted or non-substituted furan ring; or a substituted or non-substituted benzofuran ring.

10

Other more preferred compounds according to the invention are compounds of formula (I) wherein B represents a substituted or non-substituted thienyl ring. Other more preferred compounds according to the invention are compounds of formula (I) wherein B represents a substituted or non-substituted benzothienyl ring.

15

Other preferred compounds according to the invention are compounds of formula (I) wherein X independently, represents a halogen atom ; substituted or non-substituted  $C_{1-C_8}$ -alkyl ;  $C_{rC_8}$ -halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted **C3-C7**-cycloalkyl ; tri(Ci-Ce-alkyl)silyl; substituted or non-substituted  $C_{i-C_8}$ -alkoxy; or  
20 substituted or non-substituted  $C_{rC_8}$ -alkylsulfanyl.

20

Other preferred compounds according to the invention are compounds of formula (I) wherein Y independently, represents a halogen atom ; substituted or non-substituted  $C_{1-C_8}$ -alkyl ;  $C_{rC_8}$ -halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or  
25 non-substituted  $C_{rC_8}$ -alkoxy ;  $C_{rC_8}$ -halogenoalkoxy comprising up to 9 halogen atoms which can be the same or different.

25

The above mentioned preferences with regard to the substituents of the compounds according to the invention can be combined in various manners. These combinations of preferred features thus provide  
30 sub-classes of compounds according to the invention. Examples of such sub-classes of preferred compounds according to the invention are:

30

- preferred features of A with preferred features of T, Z<sup>1</sup> to Z<sup>3</sup>, n, X, L, B and Y;
- preferred features of T with preferred features of A, Z<sup>1</sup> to Z<sup>3</sup>, n, X, L, B and Y;
- preferred features of Z<sup>1</sup> with preferred features of A, T, Z<sup>2</sup>, Z<sup>3</sup>, n, X, L, B and Y;
- 35 - preferred features of Z<sup>2</sup> with preferred features of A, T, Z<sup>1</sup>, Z<sup>3</sup>, n, X, L, B and Y;
- preferred features of Z<sup>3</sup> with preferred features of A, T, Z<sup>1</sup>, Z<sup>2</sup>, n, X, L, B and Y;
- preferred features of n with preferred features of A, T, Z<sup>1</sup> to Z<sup>3</sup>, X, L, B and Y;
- preferred features of X with preferred features of A, T, Z<sup>1</sup> to Z<sup>3</sup>, n, L, B and Y;
- preferred features of L with preferred features of A, T, Z<sup>1</sup> to Z<sup>3</sup>, n, X, B and Y;
- 40 - preferred features of B with preferred features of A, T, Z<sup>1</sup> to Z<sup>3</sup>, n, X, L and Y;


40

- preferred features of Y with preferred features of A, T, Z<sup>1</sup> to Z<sup>3</sup>, n, X, L and B.


In these combinations of preferred features of the substituents of the compounds according to the invention, the said preferred features can also be selected among the more preferred features of each of A, T, Z<sup>1</sup> to Z<sup>3</sup>, n, X, L, B and Y so as to form most preferred subclasses of compounds according to the invention.

The present invention also relates to a process for the preparation of the compound of formula (I).

Thus, according to a further aspect of the present invention there is provided a process P1 for the preparation of a compound of formula (I) as herein-defined and wherein T represents O and that comprises reaction of an amine of formula (II) or one of its salts:



wherein Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X, L and B are as herein-defined; with a carboxylic acid derivative of formula (III):



wherein A is as herein-defined and U<sup>1</sup> represents a leaving group selected in the list consisting of a halogen atom, a hydroxyl group, -OR<sup>a</sup>, -OC(=O)R<sup>a</sup>, R<sup>a</sup> being a substituted or non-substituted CrC<sub>6</sub>-alkyl, a substituted or non-substituted CrC<sub>6</sub>-haloalkyl, a benzyl, 4-methoxybenzyl or pentafluorophenyl group, or a group of formula O-C(=O)A; in the presence when needed or useful of a catalyst and in the presence of a condensing agent in case U<sup>1</sup> represents a hydroxyl group, and in the presence of an acid binder in case U<sup>1</sup> represents a halogen atom.

N-substituted amine derivatives of formula (II) are known or can be prepared by known processes such as reductive amination of aldehydes or ketones (Bioorganics and Medicinal Chemistry Letters (2006), 16, 2014), or reduction of imines (Tetrahedron (2005), 61, 11689), or nucleophilic substitution of a halogen, mesylate or tosylate (Journal of Medicinal Chemistry (2002), 45, 3887).

Carboxylic acid derivatives of formula (III) are known or can be prepared by known processes.

In case U<sup>1</sup> represents a hydroxy group, process P1 according to the present invention is conducted in the presence of condensing agent. Suitable condensing agent may be selected in the non limited list consisting of acid halide former, such as phosgene, phosphorous tribromide, phosphorous trichloride, phosphorous pentachloride, phosphorous trichloride oxide or thionyl chloride; anhydride former, such as ethyl chloroformate, methyl chloroformate, isopropyl chloroformate, isobutyl chloroformate or methanesulfonyl chloride; carbodiimides, such as N,N'-dicyclohexylcarbodiimide (DCC) or other customary condensing agents, such as phosphorous pentoxide, polyphosphoric acid, N,N'-carbonyl-

diimidazole, 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline (EEDQ), triphenylphosphine/tetrachloromethane, 4-(4,6-dimethoxy[1,3,5]-triazin-2-yl)-4-methylmorpholinium chloride hydrate, bromotripyrrolidinophosphoniumhexafluorophosphate or propanephosphonic anhydride (T3P).

Process P<sub>1</sub> according to the present invention may be conducted in the presence of a catalyst.

5 Suitable catalyst may be selected in the list consisting of N,N-dimethylpyridin-4-amine, 1-hydroxybenzotriazole or N,N-dimethylformamide.

In case U<sup>1</sup> represents a halogen atom, process P<sub>1</sub> according to the present invention is conducted in the presence of an acid binder. Suitable acid binders for carrying out process P<sub>1</sub> according to the

10 invention are in each case all inorganic and organic bases that are customary for such reactions.

Preference is given to using alkaline earth metal, alkali metal hydride, alkali metal hydroxides or alkali metal alkoxides, such as sodium hydroxide, sodium hydride, calcium hydroxide, potassium hydroxide, potassium tert-butoxide or other ammonium hydroxide, alkali metal carbonates, such as caesium carbonate, sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate,

15 alkali metal or alkaline earth metal acetates, such as sodium acetate, potassium acetate, calcium acetate and also tertiary amines, such as trimethylamine, triethylamine, diisopropylethylamine, tributylamine, N,N-dimethylaniline, pyridine, N-methylpiperidine, N,N-dimethylpyridin-4-amine, diazabicyclooctane (DABCO), diazabicyclo-nonene (DBN) or diazabicycloundecene (DBU).

20 It is also possible to work in the absence of an additional condensing agent or to employ an excess of the amine component, so that it simultaneously acts as acid binder agent.

Suitable solvents for carrying out process P<sub>1</sub> according to the invention can be customary inert organic solvents. Preference is given to using optionally halogenated aliphatic, alicyclic or aromatic

25 hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile; amides, such as

30 N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone, or hexamethylphosphoric triamide; alcohols such as methanol, ethanol, propanol, iso-propanol; esters, such as methyl acetate or ethyl acetate, sulfoxides, such as dimethyl sulfoxide, or sulfones, such as sulfolane.


35 When carrying out process P<sub>1</sub> according to the invention, the amine derivative of formula (II) can be employed as its salt, such as chlorhydrate or any other convenient salt.

When carrying out process P<sub>1</sub> according to the invention, 1 mole or an excess of the amine derivative of formula (II) and from 1 to 3 moles of the acid binder can be employed per mole of the reagent of

40 formula (III).

It is also possible to employ the reaction components in other ratios. Work-up is carried out by known methods.

- 5 According to a further aspect of the present invention, there is provided a second process P2 for the preparation of a compound of formula (I) wherein L represents a direct bond or a methylene group, as illustrated by the following reaction scheme:




10

wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X and B are as herein-defined, L represents a direct bond or a methylene group, U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group and W<sup>1</sup> represents a boron derivative such as a boronic acid, a boronic ester or a potassium trifluoroborate derivative.


15

The present invention therefore refers to a process P2 for preparing a compound of formula (I)




20

wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X and B are as defined in any one of claims 1 to 14 and L represents a direct bond or a methylene group, characterized in that a compound of formula (IV)



wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X are as defined above and U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group, is reacted with a compound of formula (V)

25



(V)

wherein L and B are as defined above and  $W^1$  represents a boron derivative such as a boronic acid, a boronic ester or a potassium trifluoroborate derivative.

5 Process P2 can be performed in the presence of a transition metal catalyst such as palladium and if appropriate in the presence of a phosphine ligand or a N-heterocyclic carbene ligand and in the presence of a base and if appropriate in the presence of a solvent.

10 Compounds of formula (IV) can be prepared by known processes (WO-2007/087906 and WO-2010/130767) and the preparation of compounds of formula (V) is well known.

Process P2 according to the invention can be carried out in the presence of a catalyst, such as a metal salt or complex. Suitable metal derivatives for this purpose are transition metal catalysts such as palladium. Suitable metal salts or complexes for this purpose are for example, palladium chloride, palladium acetate, tetrakis(triphenylphosphine)palladium, bis(dibenzylideneacetone)palladium, bis(triphenyl phosphine)palladium dichloride or 1,1'-bis(diphenylphosphino) ferrocenepalladium(I) chloride.

20 It is also possible to generate a palladium complex in the reaction mixture by separate addition to the reaction of a palladium salt and a ligand or salt, such as triethylphosphine, tri-tert-butylphosphine, tri-tert-butylphosphonium tetrafluoroborate, tricyclohexylphosphine, 2-(dicyclohexylphosphine)biphenyl, 2-(di-tert-butylphosphine)biphenyl, 2-(dicyclohexylphosphine)-2'-(N,N-dimethylamino)biphenyl, triphenyl-phosphine, tris-(o-tolyl)phosphine, sodium 3-(diphenylphosphino)benzenesulfonate, tris-2-(methoxy-phenyl)phosphine, 2,2'-bis(diphenylphosphine)-1,1'-binaphthyl, 1,4-bis(diphenylphosphine)butane, 1,2-bis(diphenylphosphine) ethane, 1,4-bis(dicyclohexylphosphine)butane, 1,2-bis(dicyclohexylphosphine)-ethane, 2-(dicyclohexyl phosphine)-2'-(N,N-dimethylamino)-biphenyl, 1,1'-bis(diphenylphosphino)-ferrocene, (R)-(-)-1-[(S)-2-diphenylphosphino]ferrocenyl]ethyl dicyclohexylphosphine, tris-(2,4-tert-butyl-phenyl)phosphite or 1,3-bis(2,4,6-trimethylphenyl)imidazolium chloride.

30 It is also advantageous to choose the appropriate catalyst and/or ligand from commercial catalogues such as "Metal Catalysts for Organic Synthesis" by Strem Chemicals or "Phosphorous Ligands and Compounds" by Strem Chemicals.

35 Suitable bases for carrying out process P2 according to the invention can be inorganic and organic bases which are customary for such reactions. Preference is given to using alkaline earth metal or alkali metal hydroxides, such as sodium hydroxide, calcium hydroxide, potassium hydroxide or other ammonium hydroxide derivatives ; alkaline earth metal, alkali metal or ammonium fluorides such as

potassium fluoride, caesium fluoride or tetrabutylammonium fluoride ; alkaline earth metal or alkali metal carbonates, such as sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate or caesium carbonate ; alkali metal or alkaline earth metal acetates, such as sodium acetate, potassium acetate or calcium acetate ; alkali metal alcoholates, such as potassium ter-  
5 butoxide or sodium ter-butoxide ; alkali metal phosphates, such as tri-potassium phosphate ; tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dicyclohexylmethylamine, N-methylpiperidine, N,N-dimethylaminopyridine, diazabicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU) ; and also aromatic bases, such as pyridine, picolines, lutidines or collidines.

10

Suitable solvents for carrying out process P2 according to the invention can be customary inert organic solvents. Preference is given to using optionally halogen atomated aliphatic, alicyclic or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin ; chlorobenzene, dichlorobenzene, dichloromethane, chloroform,  
15 carbon tetrachloride, dichlorethane or trichlorethane ; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole ; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile ; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide ; esters, such as methyl acetate or ethyl  
20 acetate, sulfoxides, such as dimethyl sulfoxide, or sulfones, such as sulfolane.

It can also be advantageous to carry out process P2 according to the invention, with a co-solvent such as water or an alcohol such as methanol, ethanol, propanol, i-propanol or t-butanol.


25

When carrying out process P2 according to the invention, 1 mole or an excess of compound of formula (V) and from 1 to 5 moles of base and from 0.01 to 20 mole percent of a palladium complex can be employed per mole of compound of formula (IV).

It is also possible to employ the reaction components in other ratios. Work-up is carried out by known methods.

30

According to a further aspect of the present invention, there is provided a third process P3 for the preparation of a compound of formula (I) wherein L represents a heteroatom, as illustrated by the following reaction scheme:



wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X and B are as herein-defined, U<sup>3</sup> represents a halogen atom and L  
 5 represents an oxygen atom, a sulfur atom or a mono-substituted or non-substituted nitrogen atom,

Process P3 according to the invention is performed in the presence of a base.

Compounds of formula (VI) can be prepared by known processes (WO-2007/087906 and WO-  
 10 201 0/1 30767) and the preparation of compounds of formula (VI I) is well known.

Suitable bases for carrying out process P3 according to the invention can be inorganic and organic  
 bases which are customary for such reactions. Preference is given to using alkaline earth metal or  
 alkali metal hydroxides, such as sodium hydroxide, calcium hydroxide, potassium hydroxide or other  
 15 ammonium hydroxide derivatives ; alkaline earth metal, alkali metal or ammonium fluorides such as  
 potassium fluoride, caesium fluoride or tetrabutylammonium fluoride ; alkaline earth metal or alkali  
 metal carbonates, such as sodium carbonate, potassium carbonate, potassium bicarbonate, sodium  
 bicarbonate or caesium carbonate ; alkali metal or alkaline earth metal acetates, such as sodium  
 acetate, potassium acetate or calcium acetate ; alkali metal alcoholates, such as potassium ter-  
 20 butoxide or sodium ter-butoxide ; alkali metal phosphates, such as tri-potassium phosphate ; tertiary  
 amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-  
 dicyclohexylmethylamine, N-methylpiperidine, N,N-dimethylaminopyridine, diazabicyclooctane  
 (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU) ; and also aromatic bases, such  
 as pyridine, picolines, lutidines or collidines.

25

Other suitable bases for carrying out process P3 according to the invention can be amides or  
 organometallic derivatives. Preference is given to alkali metal amides, such as sodium amide or  
 potassium amide ; organic amides, such as lithium diisopropylamine (LDA), lithium  
 hexamethyldisilazane (LiHMDS), potassium hexamethyldisilazane (KHMDS) or sodium  
 30 hexamethyldisilazane (NaHMDS) ; organolithium derivatives, such as methyllithium, phenyllithium,  
 butyllithium, sec-butyllithium, iso-butyllithium or tert-butyllithium.


Suitable solvents for carrying out process P3 according to the invention can be customary inert  
 organic solvents. Preference is given to using optionally halogen atomated aliphatic, alicyclic or  
 35 aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane,  
 benzene, toluene, xylene or decalin ; chlorobenzene, dichlorobenzene, dichloromethane, chloroform,

carbon tetrachloride, dichlorethane or trichlorethane ; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole ; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile ; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide ; esters, such as methyl acetate or ethyl acetate, sulfoxides, such as dimethyl sulfoxide, or sulfones, such as sulfolane.

When carrying out process P3 according to the invention, 1 mole or an excess of compound of formula (VII) and from 1 to 5 moles of base can be employed per mole of compound of formula (VI).

It is also possible to employ the reaction components in other ratios. Work-up is carried out by known methods.

According to a further aspect according to the invention, there is provided a fourth process P4 for the preparation of a compound of formula (I), according to the following reaction scheme :



wherein A, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X, L and B are as herein-defined and W<sup>2</sup> represents a boron derivative such as a boronic acid, a boronic ester or a potassium trifluoroborate derivative when p is 1, or W<sup>2</sup> represents Bi when p is 3.

Compounds of formula (IX) can be prepared by known processes.

Process P4 according to the invention is performed in the presence of copper(I) acetate.

Suitable bases for carrying out process P4 according to the invention can be inorganic and organic bases which are customary for such reactions. Preference is given to using alkaline earth metal or alkali metal carbonates, such as sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate or caesium carbonate ; alkali metal phosphates, such as tri-potassium phosphate ; tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dicyclohexylmethylamine, N-methylpiperidine, N,N-dimethylaminopyridine ; and also aromatic bases, such as pyridine, picolines, lutidines or collidines.

Other suitable bases for carrying out process P4 according to the invention can be organic bases which are customary for such reactions. Preference is given to using organic amides such as lithium diisopropylamine (LDA), lithium hexamethyldisilazane (LiHMDS), potassium hexamethyldisilazane (KHMDS) or sodium hexamethyldisilazane (NaHMDS).

5

Suitable ligands for carrying out process P4 according to the invention can be organic bases which are customary for such reactions. Preference is given to using aromatic bases such as pyridine, picolines, lutidines, collidines, 2,2'-bipyridine or 1,10-phenanthroline ; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dicyclohexylmethylamine, N-methylpiperidine, N,N-dimethylaminopyridine.

10

Suitable solvents for carrying out process P4 according to the invention can be customary inert organic solvents. Preference is given to using optionally halogen atomated aliphatic, alicyclic or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin ; chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane ; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-dimethoxyethane or anisole ; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile ; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylpyrrolidone or hexamethylphosphoric triamide ; esters, such as methyl acetate or ethyl acetate, sulfoxides, such as dimethyl sulfoxide, or sulfones, such as sulfolane.

15

20

When carrying out process P4 according to the invention, 1 mole or an excess of compound of formula (IX) can be employed per mole of compound of formula (VII).

25

When carrying out process P4 according to the invention, 1 to 3 mole or an excess of base can be employed per mole of compound of formula (VII). If suitable, a combination of bases in various ratio can also be employed.

30

When carrying out process P4 according to the invention, no ligand or 0.1 to 1 mole of ligand can be employed per mole of compound of formula (VII).


When carrying out process P4 according to the invention, 0.1 to 2.5 mole or an excess of copper(I) acetate can be employed per mole of compound of formula (VII).

35

It is also possible to employ the reaction components in other ratios. Work-up is carried out by known methods.

Secondary amides of formula (VI) can be, for example, prepared according to the following reaction scheme :


40



wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X and B are as herein-defined, and

- L represents a direct bond or a methylene group when U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group, and W<sup>1</sup> represents a boron derivative such as a boronic acid, a boronic ester or a potassium trifluoroborate derivative (step a); or
- L represents an oxygen atom, a sulfur atom or a mono-substituted or non-substituted nitrogen atom when U<sup>3</sup> represents a halogen atom (step b).

- 10 Tertiary amides of formula (IV) can be, for example, prepared according to the following reaction scheme :




- Wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n and X are as herein-defined, U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group, and
- W<sup>2</sup> represents a boron derivative such as a boronic acid, a boronic ester or a potassium trifluoroborate derivative when p is 1, or
  - W<sup>2</sup> represents Bi when p is 3.

- 20 Derivatives of formula (X) or formula (XI) are known or can be prepared by known processes (WO-2011/2052490).

Step a can be performed according to process P2, step b can be performed according to process P3 and step c can be performed according to process P4.

According to a further aspect according to the invention, there is provided a fourth process P5 for the preparation of a compound of formula (I) wherein T represents S, starting from a compound of formula (I) wherein T represents O and illustrated according to the following reaction scheme :

5



Process P5

wherein A, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X, L and B are as herein-defined.

10

Process P5 according to the invention is performed in the presence of a thionating agent.

Starting amide derivatives of formula (I) wherein T represents O can be prepared according to processes P1, P2, P3 and P4.

15

Suitable thionating agents for carrying out process P5 according to the invention can be sulfur (S), sulfhydic acid (H<sub>2</sub>S), sodium sulfide (Na<sub>2</sub>S), sodium hydrosulfide (NaHS), boron trisulfide (B<sub>2</sub>S<sub>3</sub>), bis(diethylaluminium) sulfide ((AlEt<sub>2</sub>)<sub>2</sub>S), ammonium sulfide ((NH<sub>4</sub>)<sub>2</sub>S), phosphorous pentasulfide (P<sub>2</sub>S<sub>5</sub>), Lawesson's reagent (2,4-bis(4-methoxyphenyl)-1,2,3,4-dithiadiphosphetane 2,4-disulfide) or a polymer-supported thionating reagent such as described in Journal of the Chemical Society, Perkin 1 (2001), 358, in the optionally presence of a catalytic or stoichiometric or excess amount, quantity of a base such as an inorganic and organic base. Preference is given to using alkali metal carbonates, such as sodium carbonate, potassium carbonate, potassium bicarbonate, sodium bicarbonate ; heterocyclic aromatic bases, such as pyridine, picoline, lutidine, collidine ; and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylpyridin-4-amine or N-methyl-piperidine.

25

Suitable solvents for carrying out process P5 according to the invention can be customary inert organic solvents. Preference is given to using optionally halogenated aliphatic, alicyclic or aromatic hydrocarbons, such as petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichlorethane or trichlorethane, ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane or 1,2-diethoxyethane, nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile, sulfurous solvents, such as sulfolane or carbon disulfide.

35

When carrying out process P5 according to the invention, 1 mole or an excess of the sulfur equivalent of the thionating agent and from 1 to 3 moles of the base can be employed per mole of the amide reactant (I).

5 It is also possible to employ the reaction components in other ratios. Work-up is carried out by known methods.

Processes P1, P2, P3, P4 and P5 according to the invention are generally carried out under atmospheric pressure. It is also possible to operate under elevated or reduced pressure.


10 When carrying out processes P1, P2, P3, P4 and P5 according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, these processes are carried out at temperatures from -75 °C to 200 °C, preferably from 10 °C to 150 °C. A way to control the temperature for the processes according to the invention is to use microwave technology.

15 In general, the reaction mixture is concentrated under reduced pressure. The residue that remains can be freed by known methods, such as chromatography or crystallization, from any impurities that can still be present.

20 Work-up is carried out by customary methods. Generally, the reaction mixture is treated with water and the organic phase is separated off and, after drying, concentrated under reduced pressure. If appropriate, the remaining residue can, be freed by customary methods, such as chromatography, crystallization or distillation, from any impurities that may still be present.

25 The compound according to the present invention can be prepared according to the general processes of preparation described above. It will nevertheless be understood that, on the basis of his general knowledge and of available publications, the skilled worker will be able to adapt this method according to the specifics of each of the compounds, which it is desired to synthesize.

30 Still in a further aspect, the present invention relates to compounds of formula (II) useful as intermediate compounds or materials for the process of preparation according to the invention. The present invention thus provides compounds of formula (Ma.) as well as their acceptable salts:



wherein  $Z^2$ ,  $Z^3$ ,  $n$  and  $X$  are as herein-defined, and  $L$  is a direct bond and  $B^1$  is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothieryl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring,

5 providing that (IIa) does not represent :

- N-[2-(2-furyl)benzyl]cyclopropanamine
- N-[2-(2-furyl)-4,5-dimethoxybenzyl]cyclopropanamine
- N-[2-(3-bromo-2-furyl)-4,5-dimethoxybenzyl]cyclopropanamine
- N-[2-(3-bromo-2-furyl)benzyl]cyclopropanamine


10 - N-[2-(3-furyl)benzyl]cyclopropanamine

- N-[2-(2-thienyl)benzyl]cyclopropanamine

- N-[4,5-dimethoxy-2-(2-thienyl)benzyl]cyclopropanamine.

The present invention thus provides compounds of formula (IIb) as well as their acceptable salts:

15



wherein  $Z^2$ ,  $Z^3$ ,  $n$  and  $X$  are as herein-defined, and  $L$  is a direct bond and  $B^1$  is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothieryl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring,

20

providing that (IIb) does not represent :

- N-[3-(2-furyl)benzyl]cyclopropanamine
- N-[5-(2-furyl)-2-methoxybenzyl]cyclopropanamine
- 25 - N-[4-fluoro-3-(2-furyl)benzyl]cyclopropanamine
- N-[2-fluoro-5-(2-furyl)benzyl]cyclopropanamine
- N-[2-fluoro-5-(3-furyl)benzyl]cyclopropanamine
- N-[3-(3-bromo-2-furyl)-4-fluorobenzyl]cyclopropanamine
- N-[3-(3-bromo-2-furyl)benzyl]cyclopropanamine

30

- N-[5-(3-bromo-2-furyl)-2-fluorobenzyl]cyclopropanamine
- N-[5-(3-bromo-2-furyl)-2-methoxybenzyl]cyclopropanamine
- N-[3-(3-furyl)benzyl]cyclopropanamine

- N-[3-(2-thienyl)benzyl]cyclopropanamine

- N-[4-fluoro-3-(2-thienyl)benzyl]cyclopropanamine

35

- N-[2-fluoro-5-(2-thienyl)benzyl]cyclopropanamine

- N-[2-methoxy-5-(2-thienyl)benzyl]cyclopropanamine

- 5-{5-[(cyclopropylamino)methyl]-2-fluorophenyl}thiophene-2-carbonitrile.

N-cyclopropyl derivatives of formula (IIa) or formula (IIb) can be prepared according to known methods.

5

Preferred compounds of formula (IIa) according to the invention are :

- N-[5-chloro-2-(2-furyl)benzyl]cyclopropanamine
- N-[5-chloro-2-(2-thienyl)benzyl]cyclopropanamine
- N-[2-(3-thienyl)benzyl]cyclopropanamine

10

- N-[1-[2-(3-thienyl)phenyl]ethyl]cyclopropanamine
- N-[5-chloro-2-(3-thienyl)benzyl]cyclopropanamine
- N-[2-(5-chloro-2-thienyl)benzyl]cyclopropanamine
- N-[5-chloro-2-(5-chloro-2-thienyl)benzyl]cyclopropanamine.

15


Preferred compounds of formula (IIb) according to the invention are :

- N-[2-chloro-5-(2-furyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(2-thienyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(3-thienyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(5-chloro-2-thienyl)benzyl]cyclopropanamine.

20

Still in a further aspect, the present invention relates to compounds of formula (X) useful as intermediate compounds or materials for the process of preparation according to the invention.

The present invention thus provides compounds of formula (Xa) :



(Xa)

25

wherein Z<sup>2</sup> and Z<sup>3</sup> are as herein-defined, A represents A<sup>13</sup>, X<sup>a1</sup> represents a halogen atom ; substituted or non-substituted Ci-Cs-alkyl ; Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; tri(Ci-C<sub>8</sub>-alkyl)silyl; substituted or non-substituted CrC<sub>8</sub>-alkoxy; or substituted or non-substituted CrC<sub>8</sub>-alkylsulfanyl, X<sup>a2</sup> represents hydrogen or X<sup>a1</sup>, and U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group, providing that (Xa) does not represent :

30

- N-(2,5-dichlorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
- N-[1-(2,5-dichlorophenyl)ethyl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
- N-[2-(2,5-dichlorophenyl)propan-2-yl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide

35

- N-[1 -(2,5-dichlorophenyl)cyclopropyl]-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide
- 5-chloro-N-[1 -(2,5-dichlorophenyl)ethyl]-3-(difluoromethyl)-1 -methyl-1 H-pyrazole-4-carboxamide
- 5-chloro-N-[2-(2,5-dichlorophenyl)propan-2-yl]-3-(difluoromethyl)-1 -methyl-1 H-pyrazole-4-carboxamide
- 5 - 5-chloro-N-[1 -(2,5-dichlorophenyl)cyclopropyl]-3-(difluoromethyl)-1 -methyl-1 H-pyrazole-4-carboxamide
- 5-chloro-N-(2,5-dichlorobenzyl)-3-(difluoromethyl)-1 -methyl-1 H-pyrazole-4-carboxamide
- N-(5-bromo-2-fluorobenzyl)-1 ,3,5-trimethyl-1 H-pyrazole-4-carboxamide
- 10 - N-(5-bromo-2-fluorobenzyl)-3,5-dimethyl-1 H-pyrazole-4-carboxamide
- 1-methyl-N-[1 -(2,4,5-trichlorophenyl)ethyl]-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide
- 3-(difluoromethyl)-1 -methyl-N-[1 -(2,4,5-trichlorophenyl)ethyl]-1 H-pyrazole-4-carboxamide
- N-(5-chloro-2-fluorobenzyl)-1 -methyl-5-propoxy-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide
- N-(2,5-dichlorobenzyl)-1 -methyl-5-propoxy-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide.

15

Preferred compounds of formula (Xa) according to the invention are :

- N-(5-chloro-2-ethylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide
- N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide
- N-[5-chloro-2-(trifluoromethyl)benzyl]-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide
- 20 - N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide.

Most preferred compounds of formula (Xa) according to the invention are :

- 25 - N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide.

In a further aspect, the present invention also relates to a fungicide composition comprising an effective and non-phytotoxic amount of an active compound of formula (I).

The expression "effective and non-phytotoxic amount" means an amount of composition according to the invention that is sufficient to control or destroy the fungi present or liable to appear on the crops and that does not entail any appreciable symptom of phytotoxicity for the said crops. Such an amount can vary within a wide range depending on the fungus to be controlled, the type of crop, the climatic conditions and the compounds included in the fungicide composition according to the invention. This amount can be determined by systematic field trials that are within the capabilities of a person skilled in the art.

35

Thus, according to the invention, there is provided a fungicide composition comprising, as an active ingredient, an effective amount of a compound of formula (I) as herein defined and an agriculturally acceptable support, carrier or filler.

40

According to the invention, the term "support" denotes a natural or synthetic, organic or inorganic compound with that the active compound of formula (I) is combined or associated to make it easier to apply, notably to the parts of the plant. This support is thus generally inert and should be agriculturally acceptable. The support can be a solid or a liquid. Examples of suitable supports include clays, natural  
5 or synthetic silicates, silica, resins, waxes, solid fertilisers, water, alcohols, in particular butanol, organic solvents, mineral and plant oils and derivatives thereof. Mixtures of such supports can also be used.

The composition according to the invention can also comprise additional components. In particular, the  
10 composition can further comprise a surfactant. The surfactant can be an emulsifier, a dispersing agent or a wetting agent of ionic or non-ionic type or a mixture of such surfactants. Mention can be made, for example, of polyacrylic acid salts, lignosulfonic acid salts, phenolsulfonic or naphthalenesulfonic acid salts, polycondensates of ethylene oxide with fatty alcohols or with fatty acids or with fatty amines, substituted phenols (in particular alkylphenols or arylphenols), salts of sulfosuccinic acid esters,  
15 taurine derivatives (in particular alkyl taurates), phosphoric esters of polyoxyethylated alcohols or phenols, fatty acid esters of polyols and derivatives of the above compounds containing sulfate, sulfonate and phosphate functions. The presence of at least one surfactant is generally essential when the active compound and/or the inert support are water-insoluble and when the vector agent for the application is water. Preferably, surfactant content can be comprised from 5% to 40% by weight of  
20 the composition.

Optionally, additional components can also be included, e.g. protective colloids, adhesives, thickeners, thixotropic agents, penetration agents, stabilisers, sequestering agents. More generally, the active  
25 compounds can be combined with any solid or liquid additive, that complies with the usual formulation techniques.

In general, the composition according to the invention can contain from 0.05 to 99% by weight of active compound, preferably 10 to 70% by weight.

30 Compositions according to the invention can be used in various forms and formulations such as aerosol dispenser, capsule suspension, cold fogging concentrate, dustable powder, emulsifiable concentrate, emulsion oil in water, emulsion water in oil, encapsulated granule, fine granule, flowable concentrate for seed treatment, gas (under pressure), gas generating product, granule, hot fogging concentrate, macrogranule, microgranule, oil dispersible powder, oil miscible flowable concentrate, oil  
35 miscible liquid, paste, plant rodlet, powder for dry seed treatment, seed coated with a pesticide, soluble concentrate, soluble powder, solution for seed treatment, suspension concentrate (flowable concentrate), ultra low volume (ULV) liquid, ultra low volume (ULV) suspension, water dispersible granules or tablets, water dispersible powder for slurry treatment, water soluble granules or tablets, water soluble powder for seed treatment and wettable powder. These compositions include not only  
40 compositions that are ready to be applied to the plant or seed to be treated by means of a suitable

device, such as a spraying or dusting device, but also concentrated commercial compositions that must be diluted before application to the crop.

The formulations can be prepared in a manner known per se, for example by mixing the active ingredients  
5 with at least one customary extender, solvent or diluent, adjuvant, emulsifier, dispersant, and/or binder or fixative, wetting agent, water repellent, if appropriate desiccants and UV stabilizers and, if appropriate, dyes and pigments, antifoams, preservatives, inorganic and organic thickeners, adhesives, gibberellins and also further processing auxiliaries and also water. Depending on the formulation type to be prepared further processing steps are necessary, e.g. wet grinding, dry grinding and granulation.

10 The inventive active ingredients may be present as such or in their (commercial) formulations and in the use forms prepared from these formulations as a mixture with other (known) active ingredients, such as insecticides, attractants, sterilants, bactericides, acaricides, nematocides, fungicides, growth regulators, herbicides, fertilizers, safeners and/or semiochemicals.

15 The compounds of formula (I) and the fungicide composition according to the invention can be used to curatively or preventively control the phytopathogenic fungi of plants or crops.

Thus, according to a further aspect of the invention, there is provided a method for curatively or preventively controlling the phytopathogenic fungi of plants or crops characterised in that a compound  
20 of formula (I) or a fungicide composition according to the invention is applied to the seed, the plant or to the fruit of the plant or to the soil wherein the plant is growing or wherein it is desired to grow.

The method of treatment according to the invention can also be useful to treat propagation material such as tubers or rhizomes, but also seeds, seedlings or seedlings pricking out and plants or plants pricking out. This method of treatment can also be useful to treat roots. The method of treatment  
25 according to the invention can also be useful to treat the overground parts of the plant such as trunks, stems or stalks, leaves, flowers and fruit of the concerned plant.

According to the invention all plants and plant parts can be treated. By plants is meant all plants and plant populations such as desirable and undesirable wild plants, cultivars and plant varieties (whether  
30 or not protectable by plant variety or plant breeder's rights). Cultivars and plant varieties can be plants obtained by conventional propagation and breeding methods which can be assisted or supplemented by one or more biotechnological methods such as by use of double haploids, protoplast fusion, random and directed mutagenesis, molecular or genetic markers or by bioengineering and genetic engineering methods. By plant parts is meant all above ground and below  
35 ground parts and organs of plants such as shoot, leaf, blossom and root, whereby for example leaves, needles, stems, branches, blossoms, fruiting bodies, fruits and seed as well as roots, corms and rhizomes are listed. Crops and vegetative and generative propagating material, for example cuttings, corms, rhizomes, runners and seeds also belong to plant parts.

Among the plants that can be protected by the method according to the invention, mention may be made of major field crops like corn, soybean, cotton, *Brassica* oilseeds such as *Brassica napus* (e.g. canola),  
40 *Brassica rapa*, *B. juncea* (e.g. mustard) and *Brassica carinata*, rice, wheat, sugarbeet, sugarcane, oats,

rye, barley, millet, triticale, flax, vine and various fruits and vegetables of various botanical taxa such as *Rosaceae* sp. (for instance pip fruit such as apples and pears, but also stone fruit such as apricots, cherries, almonds and peaches, berry fruits such as strawberries), *Ribesioideae* sp., *Juglandaceae* sp., *Betulaceae* sp., *Anacardiaceae* sp., *Fagaceae* sp., *Moraceae* sp., *Oleaceae* sp., *Actinidaceae* sp.,  
5 *Lauraceae* sp., *Musaceae* sp. (for instance banana trees and plantings), *Rubiaceae* sp. (for instance coffee), *Theaceae* sp., *Sterculiaceae* sp., *Rutaceae* sp. (for instance lemons, oranges and grapefruit) ;  
*Solanaceae* sp. (for instance tomatoes, potatoes, peppers, eggplant), *Liliaceae* sp., *Compositae* sp. (for instance lettuce, artichoke and chicory - including root chicory, endive or common chicory), *Umbelliferae*  
10 sp. (for instance carrot, parsley, celery and celeriac), *Cucurbitaceae* sp. (for instance cucumber - including pickling cucumber, squash, watermelon, gourds and melons), *Alliaceae* sp. (for instance onions and leek),  
*Cruciferae* sp. (for instance white cabbage, red cabbage, broccoli, cauliflower, brussel sprouts, pak choi, kohlrabi, radish, horseradish, cress, Chinese cabbage), *Leguminosae* sp. (for instance peanuts, peas and beans - such as climbing beans and broad beans), *Chenopodiaceae* sp. (for instance mangold, spinach beet, spinach, beetroots), *Malvaceae* (for instance okra), *Asparagaceae* (for instance asparagus);  
15 horticultural and forest crops; ornamental plants; as well as genetically modified homologues of these crops.

The method of treatment according to the invention can be used in the treatment of genetically modified organisms (GMOs), e.g. plants or seeds. Genetically modified plants (or transgenic plants) are plants of  
20 which a heterologous gene has been stably integrated into genome. The expression "heterologous gene" essentially means a gene which is provided or assembled outside the plant and when introduced in the nuclear, chloroplastic or mitochondrial genome gives the transformed plant new or improved agronomic or other properties by expressing a protein or polypeptide of interest or by downregulating or silencing other gene(s) which are present in the plant (using for example, antisense technology, cosuppression technology  
25 or RNA interference - RNAi - technology). A heterologous gene that is located in the genome is also called a transgene. A transgene that is defined by its particular location in the plant genome is called a transformation or transgenic event.

Depending on the plant species or plant cultivars, their location and growth conditions (soils, climate, vegetation period, diet), the treatment according to the invention may also result in superadditive  
30 ("synergistic") effects. Thus, for example, reduced application rates and/or a widening of the activity spectrum and/or an increase in the activity of the active compounds and compositions which can be used according to the invention, better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salt content, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, bigger fruits, larger plant height, greener leaf  
35 color, earlier flowering, higher quality and/or a higher nutritional value of the harvested products, higher sugar concentration within the fruits, better storage stability and/or processability of the harvested products are possible, which exceed the effects which were actually to be expected.

At certain application rates, the active compound combinations according to the invention may also have a strengthening effect in plants. Accordingly, they are also suitable for mobilizing the defense system of the plant against attack by unwanted microorganisms. This may, if appropriate, be one of the reasons of the enhanced activity of the combinations according to the invention, for example against fungi. Plant-strengthening (resistance-inducing) substances are to be understood as meaning, in the present context, those substances or combinations of substances which are capable of stimulating the defense system of plants in such a way that, when subsequently inoculated with unwanted microorganisms, the treated plants display a substantial degree of resistance to these microorganisms. In the present case, unwanted microorganisms are to be understood as meaning phytopathogenic fungi, bacteria and viruses. Thus, the substances according to the invention can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period of time within which protection is effected generally extends from 1 to 10 days, preferably 1 to 7 days, after the treatment of the plants with the active compounds.

Plants and plant cultivars which are preferably to be treated according to the invention include all plants which have genetic material which impart particularly advantageous, useful traits to these plants (whether obtained by breeding and/or biotechnological means).

Plants and plant cultivars which are also preferably to be treated according to the invention are resistant against one or more biotic stresses, i.e. said plants show a better defense against animal and microbial pests, such as against nematodes, insects, mites, phytopathogenic fungi, bacteria, viruses and/or viroids.

Examples of nematode resistant plants are described in e.g. US Patent Application Nos 11/765,491, 11/765,494, 10/926,819, 10/782,020, 12/032,479, 10/783,417, 10/782,096, 11/657,964, 12/192,904, 11/396,808, 12/166,253, 12/166,239, 12/166,124, 12/166,209, 11/762,886, 12/364,335, 11/763,947, 12/252,453, 12/209,354, 12/491,396 or 12/497,221.

Plants and plant cultivars which may also be treated according to the invention are those plants which are resistant to one or more abiotic stresses. Abiotic stress conditions may include, for example, drought, cold temperature exposure, heat exposure, osmotic stress, flooding, increased soil salinity, increased mineral exposure, ozone exposure, high light exposure, limited availability of nitrogen nutrients, limited availability of phosphorus nutrients, shade avoidance.

Plants and plant cultivars which may also be treated according to the invention, are those plants characterized by enhanced yield characteristics. Increased yield in said plants can be the result of, for example, improved plant physiology, growth and development, such as water use efficiency, water retention efficiency, improved nitrogen use, enhanced carbon assimilation, improved photosynthesis, increased germination efficiency and accelerated maturation. Yield can furthermore be affected by improved plant architecture (under stress and non-stress conditions), including but not limited to, early flowering, flowering control for hybrid seed production, seedling vigor, plant size, internode number and distance, root growth, seed size, fruit size, pod size, pod or ear number, seed number per pod or ear, seed mass, enhanced seed filling, reduced seed dispersal, reduced pod dehiscence and lodging resistance. Further yield traits include seed composition, such as carbohydrate content, protein

content, oil content and composition, nutritional value, reduction in anti-nutritional compounds, improved processability and better storage stability.

Examples of plants with the above-mentioned traits are non-exhaustively listed in Table A.

5 Plants that may be treated according to the invention are hybrid plants that already express the characteristic of heterosis or hybrid vigor which results in generally higher yield, vigor, health and resistance towards biotic and abiotic stresses). Such plants are typically made by crossing an inbred male-sterile parent line (the female parent) with another inbred male-fertile parent line (the male parent). Hybrid seed is typically harvested from the male sterile plants and sold to growers. Male  
10 sterile plants can sometimes (e.g. in corn) be produced by detasseling, i.e. the mechanical removal of the male reproductive organs (or males flowers) but, more typically, male sterility is the result of genetic determinants in the plant genome. In that case, and especially when seed is the desired product to be harvested from the hybrid plants it is typically useful to ensure that male fertility in the hybrid plants is fully restored. This can be accomplished by ensuring that the male parents have  
15 appropriate fertility restorer genes which are capable of restoring the male fertility in hybrid plants that contain the genetic determinants responsible for male-sterility. Genetic determinants for male sterility may be located in the cytoplasm. Examples of cytoplasmic male sterility (CMS) were for instance described in Brassica species (WO 92/05251, WO 95/09910, WO 98/27806, WO 05/002324, WO 06/021972 and US 6,229,072). However, genetic determinants for male sterility can also be located in  
20 the nuclear genome. Male sterile plants can also be obtained by plant biotechnology methods such as genetic engineering. A particularly useful means of obtaining male-sterile plants is described in WO 89/10396 in which, for example, a ribonuclease such as barnase is selectively expressed in the tapetum cells in the stamens. Fertility can then be restored by expression in the tapetum cells of a ribonuclease inhibitor such as barstar (e.g. WO 91/02069).

25

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may be treated according to the invention are herbicide-tolerant plants, i.e. plants made tolerant to one or more given herbicides. Such plants can be obtained either by genetic transformation, or by selection of plants containing a mutation imparting such herbicide tolerance.

30 Herbicide-resistant plants are for example glyphosate-tolerant plants, i.e. plants made tolerant to the herbicide glyphosate or salts thereof. Plants can be made tolerant to glyphosate through different means. For example, glyphosate-tolerant plants can be obtained by transforming the plant with a gene encoding the enzyme 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS). Examples of such EPSPS genes are the AroA gene (mutant CT7) of the bacterium *Salmonella typhimurium* (Comai et al., 1983, Science 221, 370-371), the CP4 gene of the bacterium *Agrobacterium sp.* (Barry et al., 1992, Curr. Topics Plant Physiol. 7, 139-145), the genes encoding a Petunia EPSPS (Shah et al., 1986, Science 233, 478-481), a Tomato EPSPS (Gasser et al., 1988, J. Biol. Chem. 263, 4280-4289), or an Eleusine EPSPS (WO 01/66704). It can also be a mutated EPSPS as described in for example EP 0837944, WO 00/66746, WO 00/66747 or WO02/26995. Glyphosate-tolerant plants can also be  
40 obtained by expressing a gene that encodes a glyphosate oxido-reductase enzyme as described in

U.S. Patent Nos. 5,776,760 and 5,463,175. Glyphosate-tolerant plants can also be obtained by expressing a gene that encodes a glyphosate acetyl transferase enzyme as described in for example WO 02/36782, WO 03/092360, WO 05/012515 and WO 07/024782. Glyphosate-tolerant plants can also be obtained by selecting plants containing naturally-occurring mutations of the above-mentioned genes, as described in for example WO 01/024615 or WO 03/013226. Plants expressing EPSPS genes that confer glyphosate tolerance are described in e.g. US Patent Application Nos 11/517,991, 10/739,610, 12/139,408, 12/352,532, 11/312,866, 11/315,678, 12/421,292, 11/400,598, 11/651,752, 11/681,285, 11/605,824, 12/468,205, 11/760,570, 11/762,526, 11/769,327, 11/769,255, 11/943801 or 12/362,774. Plants comprising other genes that confer glyphosate tolerance, such as decarboxylase genes, are described in e.g. US patent applications 11/588,811, 11/185,342, 12/364,724, 11/185,560 or 12/423,926.

Other herbicide resistant plants are for example plants that are made tolerant to herbicides inhibiting the enzyme glutamine synthase, such as bialaphos, phosphinothricin or glufosinate. Such plants can be obtained by expressing an enzyme detoxifying the herbicide or a mutant glutamine synthase enzyme that is resistant to inhibition, e.g. described in US Patent Application No 11/760,602. One such efficient detoxifying enzyme is an enzyme encoding a phosphinothricin acetyltransferase (such as the bar or pat protein from *Streptomyces* species). Plants expressing an exogenous phosphinothricin acetyltransferase are for example described in U.S. Patent Nos. 5,561,236; 5,648,477; 5,646,024; 5,273,894; 5,637,489; 5,276,268; 5,739,082; 5,908,810 and 7,112,665. Further herbicide-tolerant plants are also plants that are made tolerant to the herbicides inhibiting the enzyme hydroxyphenylpyruvatedioxygenase (HPPD). Hydroxyphenylpyruvatedioxygenases are enzymes that catalyze the reaction in which para-hydroxyphenylpyruvate (HPP) is transformed into homogentisate. Plants tolerant to HPPD-inhibitors can be transformed with a gene encoding a naturally-occurring resistant HPPD enzyme, or a gene encoding a mutated or chimeric HPPD enzyme as described in WO 96/38567, WO 99/24585, WO 99/24586, WO 2009/144079, WO 2002/046387, or US 6,768,044. Tolerance to HPPD-inhibitors can also be obtained by transforming plants with genes encoding certain enzymes enabling the formation of homogentisate despite the inhibition of the native HPPD enzyme by the HPPD-inhibitor. Such plants and genes are described in WO 99/34008 and WO 02/36787. Tolerance of plants to HPPD inhibitors can also be improved by transforming plants with a gene encoding an enzyme having prephenate deshydrogenase (PDH) activity in addition to a gene encoding an HPPD-tolerant enzyme, as described in WO 2004/024928. Further, plants can be made more tolerant to HPPD-inhibitor herbicides by adding into their genome a gene encoding an enzyme capable of metabolizing or degrading HPPD inhibitors, such as the CYP450 enzymes shown in WO 2007/103567 and WO 2008/150473.

Still further herbicide resistant plants are plants that are made tolerant to acetolactate synthase (ALS) inhibitors. Known ALS-inhibitors include, for example, sulfonylurea, imidazolinone, triazolopyrimidines, pyrimidinyoxy(thio)benzoates, and/or sulfonaminocarbonyltriazolinone herbicides. Different mutations in the ALS enzyme (also known as acetohydroxyacid synthase, AHAS) are known to confer tolerance to different herbicides and groups of herbicides, as described for example in Tranel and

Wright (2002, Weed Science 50:700-712), but also, in U.S. Patent No. 5,605,011, 5,378,824, 5,141,870, and 5,013,659. The production of sulfonylurea-tolerant plants and imidazolinone-tolerant plants is described in U.S. Patent Nos. 5,605,011; 5,013,659; 5,141,870; 5,767,361; 5,731,180; 5,304,732; 4,761,373; 5,331,107; 5,928,937; and 5,378,824; and international publication WO 96/33270. Other imidazolinone-tolerant plants are also described in for example WO 2004/040012, WO 2004/106529, WO 2005/020673, WO 2005/093093, WO 2006/007373, WO 2006/015376, WO 2006/024351, and WO 2006/060634. Further sulfonylurea- and imidazolinone-tolerant plants are also described in for example WO 07/024782 and US Patent Application No. 61/288958.

Other plants tolerant to imidazolinone and/or sulfonylurea can be obtained by induced mutagenesis, selection in cell cultures in the presence of the herbicide or mutation breeding as described for example for soybeans in U.S. Patent 5,084,082, for rice in WO 97/41218, for sugar beet in U.S. Patent 5,773,702 and WO 99/057965, for lettuce in U.S. Patent 5,198,599, or for sunflower in WO 01/065922.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are insect-resistant transgenic plants, i.e. plants made resistant to attack by certain target insects. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such insect resistance.

An "insect-resistant transgenic plant", as used herein, includes any plant containing at least one transgene comprising a coding sequence encoding:

- 1) an insecticidal crystal protein from *Bacillus thuringiensis* or an insecticidal portion thereof, such as the insecticidal crystal proteins listed by Crickmore et al. (1998, Microbiology and Molecular Biology Reviews, 62: 807-813), updated by Crickmore et al. (2005) at the *Bacillus thuringiensis* toxin nomenclature, online at: [http://www.lifesci.sussex.ac.uk/Home/Neil\\_Crickmore/Bt/](http://www.lifesci.sussex.ac.uk/Home/Neil_Crickmore/Bt/), or insecticidal portions thereof, e.g., proteins of the Cry protein classes Cry1 Ab, Cry1 Ac, Cry1 B, Cry1 C, Cry1 D, Cry1 F, Cry2Ab, Cry3Aa, or Cry3Bb or insecticidal portions thereof (e.g. EP 1999141 and WO 2007/107302), or such proteins encoded by synthetic genes as e.g. described in and US Patent Application No. 12/249,016; or
- 2) a crystal protein from *Bacillus thuringiensis* or a portion thereof which is insecticidal in the presence of a second other crystal protein from *Bacillus thuringiensis* or a portion thereof, such as the binary toxin made up of the Cry34 and Cry35 crystal proteins (Moellenbeck et al. 2001, Nat. Biotechnol. 19: 668-72; Schnepf et al. 2006, Applied Environm. Microbiol. 71, 1765-1774) or the binary toxin made up of the Cry1 A or Cry1 F proteins and the Cry2Aa or Cry2Ab or Cry2Ae proteins (US Patent Appl. No. 12/214,022 and EP 08010791.5); or
- 3) a hybrid insecticidal protein comprising parts of different insecticidal crystal proteins from *Bacillus thuringiensis*, such as a hybrid of the proteins of 1) above or a hybrid of the proteins of 2) above, e.g., the Cry1 A.105 protein produced by corn event MON89034 (WO 2007/027777); or
- 4) a protein of any one of 1) to 3) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect

species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation, such as the Cry3Bb1 protein in corn events MON863 or MON8801 7, or the Cry3A protein in corn event MIR604; or

5 5) an insecticidal secreted protein from *Bacillus thuringiensis* or *Bacillus cereus*, or an insecticidal portion thereof, such as the vegetative insecticidal (VIP) proteins listed at: [http://www.lifesci.sussex.ac.uk/home/Neil\\_Crickmore/Bt/vip.html](http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/vip.html), e.g., proteins from the VIP3Aa protein class; or

10 6) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a second secreted protein from *Bacillus thuringiensis* or *B. cereus*, such as the binary toxin made up of the VIP1A and VIP2A proteins (WO 94/21 795); or

7) a hybrid insecticidal protein comprising parts from different secreted proteins from *Bacillus thuringiensis* or *Bacillus cereus*, such as a hybrid of the proteins in 1) above or a hybrid of the proteins in 2) above; or

15 8) a protein of any one of 5) to 7) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding an insecticidal protein), such as the VIP3Aa protein in cotton event COT1 02; or

20 9) a secreted protein from *Bacillus thuringiensis* or *Bacillus cereus* which is insecticidal in the presence of a crystal protein from *Bacillus thuringiensis*, such as the binary toxin made up of VIP3 and Cry1 A or Cry1 F (US Patent Appl. No. 61/126083 and 61/19501 9), or the binary toxin made up of the VIP3 protein and the Cry2Aa or Cry2Ab or Cry2Ae proteins (US Patent Appl. No. 12/21 4,022 and EP 0801 0791 .5).

25 10) a protein of 9) above wherein some, particularly 1 to 10, amino acids have been replaced by another amino acid to obtain a higher insecticidal activity to a target insect species, and/or to expand the range of target insect species affected, and/or because of changes introduced into the encoding DNA during cloning or transformation (while still encoding an insecticidal protein)

30 Of course, an insect-resistant transgenic plant, as used herein, also includes any plant comprising a combination of genes encoding the proteins of any one of the above classes 1 to 10. In one embodiment, an insect-resistant plant contains more than one transgene encoding a protein of any one of the above classes 1 to 10, to expand the range of target insect species affected when using  
35 different proteins directed at different target insect species, or to delay insect resistance development to the plants by using different proteins insecticidal to the same target insect species but having a different mode of action, such as binding to different receptor binding sites in the insect.

An "insect-resistant transgenic plant", as used herein, further includes any plant containing at least one  
40 transgene comprising a sequence producing upon expression a double-stranded RNA which upon

ingestion by a plant insect pest inhibits the growth of this insect pest, as described e.g. in WO 2007/0801 26, WO 2006/1 29204, WO 2007/074405, WO 2007/0801 27 and WO 2007/035650.

Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which  
5 may also be treated according to the invention are tolerant to abiotic stresses. Such plants can be obtained by genetic transformation, or by selection of plants containing a mutation imparting such stress resistance. Particularly useful stress tolerance plants include:

1) plants which contain a transgene capable of reducing the expression and/or the activity of poly(ADP-ribose) polymerase (PARP) gene in the plant cells or plants as described in WO  
10 00/041 73, WO/2006/045633, EP 04077984.5, or EP 06009836.5.

2) plants which contain a stress tolerance enhancing transgene capable of reducing the expression and/or the activity of the PARG encoding genes of the plants or plants cells, as described e.g. in WO 2004/0901 40.

3) plants which contain a stress tolerance enhancing transgene coding for a plant-functional  
15 enzyme of the nicotineamide adenine dinucleotide salvage synthesis pathway including nicotinamidase, nicotinate phosphoribosyltransferase, nicotinic acid mononucleotide adenyl transferase, nicotinamide adenine dinucleotide synthetase or nicotine amide phosphorybosyltransferase as described e.g. in EP 04077624.7, WO 2006/1 33827, PCT/EP07/002433, EP 1999263, or WO 2007/1 07326.

20 Plants or plant cultivars (obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention show altered quantity, quality and/or storage-stability of the harvested product and/or altered properties of specific ingredients of the harvested product such as :

25 1) transgenic plants which synthesize a modified starch, which in its physical-chemical characteristics, in particular the amylose content or the amylose/amylopectin ratio, the degree of branching, the average chain length, the side chain distribution, the viscosity behaviour, the gelling strength, the starch grain size and/or the starch grain morphology, is changed in comparison with the synthesised starch in wild type plant cells or plants, so that this is better  
30 suited for special applications. Said transgenic plants synthesizing a modified starch are disclosed, for example, in EP 0571 427, WO 95/04826, EP 071 9338, WO 96/1 5248, WO 96/1 9581 , WO 96/27674, WO 97/1 1188, WO 97/26362, WO 97/32985, WO 97/42328, WO 97/44472, WO 97/45545, WO 98/2721 2, WO 98/40503, W099/58688, WO 99/58690, WO 99/58654, WO 00/081 84, WO 00/081 85, WO 00/081 75, WO 00/28052, WO 00/77229, WO  
35 01/12782, WO 01/12826, WO 02/1 01059, WO 03/071 860, WO 2004/056999, WO 2005/030942, WO 2005/030941 , WO 2005/095632, WO 2005/09561 7, WO 2005/09561 9, WO 2005/09561 8, WO 2005/1 23927, WO 2006/01 831 9, WO 2006/1 031 07, WO 2006/1 08702, WO 2007/009823, WO 00/221 40, WO 2006/063862, WO 2006/072603, WO 02/034923, EP 060901 34.5, EP 06090228.5, EP 06090227.7, EP 07090007.1 , EP 07090009.7, WO  
40 01/14569, WO 02/7941 0, WO 03/33540, WO 2004/078983, WO 01/19975, WO 95/26407, WO 96/34968, WO 98/201 45, WO 99/1 2950, WO 99/66050, WO 99/53072, US 6,734,341 , WO

00/1 1192, WO 98/22604, WO 98/32326, WO 01/98509, WO 01/98509, WO 2005/002359, US 5,824,790, US 6,013,861, WO 94/04693, WO 94/09144, WO 94/11520, WO 95/35026, WO 97/20936

2) transgenic plants which synthesize non starch carbohydrate polymers or which synthesize non starch carbohydrate polymers with altered properties in comparison to wild type plants without genetic modification. Examples are plants producing polyfructose, especially of the inulin and levan-type, as disclosed in EP 0663956, WO 96/01904, WO 96/21023, WO 98/39460, and WO 99/24593, plants producing alpha-1,4-glucans as disclosed in WO 95/31553, US 2002031826, US 6,284,479, US 5,712,107, WO 97/47806, WO 97/47807, WO 97/47808 and WO 00/14249, plants producing alpha-1,6 branched alpha-1,4-glucans, as disclosed in WO 00/73422, plants producing alternan, as disclosed in e.g. WO 00/47727, WO 00/73422, EP 06077301.7, US 5,908,975 and EP 0728213,

3) transgenic plants which produce hyaluronan, as for example disclosed in WO 2006/032538, WO 2007/039314, WO 2007/039315, WO 2007/039316, JP 2006304779, and WO 2005/012529.

4) transgenic plants or hybrid plants, such as onions with characteristics such as 'high soluble solids content', 'low pungency' (LP) and/or 'long storage' (LS), as described in US Patent Appl. No. 12/020,360 and 61/054,026.

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as cotton plants, with altered fiber characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered fiber characteristics and include:

- a) Plants, such as cotton plants, containing an altered form of cellulose synthase genes as described in WO 98/00549
- b) Plants, such as cotton plants, containing an altered form of rsw2 or rsw3 homologous nucleic acids as described in WO 2004/053219
- c) Plants, such as cotton plants, with increased expression of sucrose phosphate synthase as described in WO 01/17333
- d) Plants, such as cotton plants, with increased expression of sucrose synthase as described in WO 02/45485
- e) Plants, such as cotton plants, wherein the timing of the plasmodesmatal gating at the basis of the fiber cell is altered, e.g. through downregulation of fiber-selective  $\beta$ -1,3-glucanase as described in WO 2005/017157, or as described in EP 08075514.3 or US Patent Appl. No. 61/128,938
- f) Plants, such as cotton plants, having fibers with altered reactivity, e.g. through the expression of N-acetylglucosaminetransferase gene including nodC and chitin synthase genes as described in WO 2006/136351

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered oil profile characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered oil profile

5 characteristics and include:

- a) Plants, such as oilseed rape plants, producing oil having a high oleic acid content as described e.g. in US 5,969,169, US 5,840,946 or US 6,323,392 or US 6,063,947
- b) Plants such as oilseed rape plants, producing oil having a low linolenic acid content as described in US 6,270,828, US 6,169,190, or US 5,965,755
- 10 c) Plant such as oilseed rape plants, producing oil having a low level of saturated fatty acids as described e.g. in US Patent No. 5,434,283 or US Patent Application No 12/668303

Plants or plant cultivars (that can be obtained by plant biotechnology methods such as genetic engineering) which may also be treated according to the invention are plants, such as oilseed rape or related Brassica plants, with altered seed shattering characteristics. Such plants can be obtained by genetic transformation, or by selection of plants contain a mutation imparting such altered seed shattering characteristics and include plants such as oilseed rape plants with delayed or reduced seed shattering as described in US Patent Appl. No. 61/135,230 WO09/068313 and W010/006732.

Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or combination of transformation events, that are the subject of petitions for non-regulated status, in the United States of America, to the Animal and Plant Health Inspection Service (APHIS) of the United States Department of Agriculture (USDA) whether such petitions are granted or are still pending. At any time this information is readily available from APHIS (4700 River Road Riverdale, MD 20737, USA), for instance on its internet site (URL

25 [http://www.aphis.usda.gov/brs/not\\_reg.html](http://www.aphis.usda.gov/brs/not_reg.html)). On the filing date of this application the petitions for nonregulated status that were pending with APHIS or granted by APHIS were those listed in table B which contains the following information:

- Petition : the identification number of the petition. Technical descriptions of the transformation events can be found in the individual petition documents which are obtainable from APHIS, for example on the APHIS website, by reference to this petition number. These descriptions are herein incorporated by reference.
- Extension of Petition : reference to a previous petition for which an extension is requested.
- Institution : the name of the entity submitting the petition.
- 35 - Regulated article : the plant species concerned.
- Transgenic phenotype : the trait conferred to the plants by the transformation event.
- Transformation event or line : the name of the event or events (sometimes also designated as lines or lines) for which nonregulated status is requested.
- APHIS documents : various documents published by APHIS in relation to the Petition and which can be requested with APHIS.

40

Additional particularly useful plants containing single transformation events or combinations of transformation events are listed for example in the databases from various national or regional regulatory agencies (see for example [http://gmoinfo.jrc.it/gmp\\_browse.aspx](http://gmoinfo.jrc.it/gmp_browse.aspx) and <http://www.agbios.com/dbase.php>).

5 Particularly useful transgenic plants which may be treated according to the invention are plants containing transformation events, or a combination of transformation events, and that are listed for example in the databases for various national or regional regulatory agencies including Event 1143-14A (cotton, insect control, not deposited, described in WO 2006/1 28569); Event 1143-51 B (cotton, insect control, not deposited, described in WO 2006/1 28570); Event 1445 (cotton, herbicide tolerance, 10 not deposited, described in US-A 2002-1 20964 or WO 02/034946); Event 17053 (rice, herbicide tolerance, deposited as PTA-9843, described in WO 201 0/1 17737); Event 1731 4 (rice, herbicide tolerance, deposited as PTA-9844, described in WO 201 0/1 17735); Event 281 -24-236 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, described in WO 2005/1 03266 or US-A 2005-2 16969); Event 3006-21 0-23 (cotton, insect control - herbicide tolerance, deposited as PTA-6233, 15 described in US-A 2007-1 43876 or WO 2005/1 03266); Event 3272 (corn, quality trait, deposited as PTA-9972, described in WO 2006/098952 or US-A 2006-230473); Event 4041 6 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-1 1508, described in WO 201 1/075593); Event 43A47 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-1 1509, described in WO 201 1/075595) ; Event 5307 (corn, insect control, deposited as ATCC PTA-9561 , described in WO 20 201 0/07 781 6); Event ASR-368 (bent grass, herbicide tolerance, deposited as ATCC PTA-481 6, described in US-A 2006-1 62007 or WO 2004/053062); Event B 16 (corn, herbicide tolerance, not deposited, described in US-A 2003-1 26634); Event BPS-CV1 27-9 (soybean, herbicide tolerance, deposited as NCIMB No. 4 1603, described in WO 201 0/080829); Event CE43-67B (cotton, insect control, deposited as DSM ACC2724, described in US-A 2009-21 7423 or WO2006/1 28573); Event 25 CE44-69D (cotton, insect control, not deposited, described in US-A 201 0-0024077); Event CE44-69D (cotton, insect control, not deposited, described in WO 2006/1 28571 ); Event CE46-02A (cotton, insect control, not deposited, described in WO 2006/1 28572); Event COT1 02 (cotton, insect control, not deposited, described in US-A 2006-1 301 75 or WO 2004/039986); Event COT202 (cotton, insect control, not deposited, described in US-A 2007-067868 or WO 2005/054479); Event COT203 (cotton, 30 insect control, not deposited, described in WO 2005/054480); Event DAS40278 (corn, herbicide tolerance, deposited as ATCC PTA-1 0244, described in WO 201 1/022469); Event DAS-591 22-7 (corn, insect control - herbicide tolerance, deposited as ATCC PTA 11384 , described in US-A 2006-0701 39); Event DAS-591 32 (corn, insect control - herbicide tolerance, not deposited, described in WO 2009/1 001 88); Event DAS6841 6 (soybean, herbicide tolerance, deposited as ATCC PTA-1 0442, 35 described in WO 201 1/066384 or WO 201 1/066360); Event DP-0981 40-6 (corn, herbicide tolerance, deposited as ATCC PTA-8296, described in US-A 2009-1 37395 or WO 2008/1 1201 9); Event DP-305423-1 (soybean, quality trait, not deposited, described in US-A 2008-31 2082 or WO 2008/054747); Event DP-321 38-1 (corn, hybridization system, deposited as ATCC PTA-91 58, described in US-A 2009-021 0970 or WO 2009/1 03049); Event DP-356043-5 (soybean, herbicide tolerance, deposited as 40 ATCC PTA-8287, described in US-A 201 0-01 84079 or WO 2008/002872); Event EE-1 (brinjal, insect

control, not deposited, described in WO 2007/091 277); Event FI117 (corn, herbicide tolerance, deposited as ATCC 209031, described in US-A 2006-059581 or WO 98/0441 40); Event GA21 (corn, herbicide tolerance, deposited as ATCC 209033, described in US-A 2005-08671 9 or WO 98/0441 40); Event GG25 (corn, herbicide tolerance, deposited as ATCC 209032, described in US-A 2005-1 88434 or WO 98/0441 40); Event GHB1 19 (cotton, insect control - herbicide tolerance, deposited as ATCC PTA-8398, described in WO 2008/1 51780); Event GHB61 4 (cotton, herbicide tolerance, deposited as ATCC PTA-6878, described in US-A 201 0-050282 or WO 2007/01 7186); Event GJ11 (corn, herbicide tolerance, deposited as ATCC 209030, described in US-A 2005-1 88434 or WO 98/0441 40); Event GM RZ1 3 (sugar beet, virus resistance, deposited as NCIMB-41 601, described in WO 201 0/07621 2); Event H7-1 (sugar beet, herbicide tolerance, deposited as NCIMB 41158 or NCIMB 41159, described in US-A 2004-1 72669 or WO 2004/074492); Event JOPLIN 1 (wheat, disease tolerance, not deposited, described in US-A 2008-064032); Event LL27 (soybean, herbicide tolerance, deposited as NCIMB41 658, described in WO 2006/1 08674 or US-A 2008-32061 6); Event LL55 (soybean, herbicide tolerance, deposited as NCIMB 41660, described in WO 2006/1 08675 or US-A 2008-1 961 27); Event LLCotton25 (cotton, herbicide tolerance, deposited as ATCC PTA-3343, described in WO 03/01 3224 or US-A 2003-097687); Event LLRICE06 (rice, herbicide tolerance, deposited as ATCC-23352, described in US 6,468,747 or WO 00/026345); Event LLRICE601 (rice, herbicide tolerance, deposited as ATCC PTA-2600, described in US-A 2008-2289060 or WO 00/026356); Event LY038 (corn, quality trait, deposited as ATCC PTA-5623, described in US-A 2007-028322 or WO 2005/061 720); Event MIR1 62 (corn, insect control, deposited as PTA-81 66, described in US-A 2009-300784 or WO 2007/1 42840); Event MIR604 (corn, insect control, not deposited, described in US-A 2008-1 67456 or WO 2005/1 03301); Event MON1 5985 (cotton, insect control, deposited as ATCC PTA-251 6, described in US-A 2004-25031 7 or WO 02/1 001 63); Event MON81 0 (corn, insect control, not deposited, described in US-A 2002-1 02582); Event MON863 (corn, insect control, deposited as ATCC PTA-2605, described in WO 2004/01 1601 or US-A 2006-095986); Event MON87427 (corn, pollination control, deposited as ATCC PTA-7899, described in WO 201 1/062904); Event MON87460 (corn, stress tolerance, deposited as ATCC PTA-891 0, described in WO 2009/1 11263 or US-A 201 1-01 38504); Event MON87701 (soybean, insect control, deposited as ATCC PTA-81 94, described in US-A 2009-130071 or WO 2009/064652); Event MON87705 (soybean, quality trait - herbicide tolerance, deposited as ATCC PTA-9241, described in US-A 201 0-0080887 or WO 201 0/03701 6); Event MON87708 (soybean, herbicide tolerance, deposited as ATCC PTA9670, described in WO 201 1/034704); Event MON87754 (soybean, quality trait, deposited as ATCC PTA-9385, described in WO 201 0/024976); Event MON87769 (soybean, quality trait, deposited as ATCC PTA-891 1, described in US-A 201 1-00671 41 or WO 2009/1 02873); Event MON8801 7 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-5582, described in US-A 2008-028482 or WO 2005/0591 03); Event MON8891 3 (cotton, herbicide tolerance, deposited as ATCC PTA-4854, described in WO 2004/072235 or US-A 2006-059590); Event MON89034 (corn, insect control, deposited as ATCC PTA-7455, described in WO 2007/1 40256 or US-A 2008-260932); Event MON89788 (soybean, herbicide tolerance, deposited as ATCC PTA-6708, described in US-A 2006-28291 5 or WO 2006/1 30436); Event MS1 1 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC

PTA-850 or PTA-2485, described in WO 01/031 042); Event MS8 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in WO 01/041 558 or US-A 2003-188347); Event NK603 (corn, herbicide tolerance, deposited as ATCC PTA-2478, described in US-A 2007-292854) ; Event PE-7 (rice, insect control, not deposited, described in WO 2008/1 14282); Event  
 5 RF3 (oilseed rape, pollination control - herbicide tolerance, deposited as ATCC PTA-730, described in WO 01/041 558 or US-A 2003-1 88347); Event RT73 (oilseed rape, herbicide tolerance, not deposited, described in WO 02/036831 or US-A 2008-070260); Event T227-1 (sugar beet, herbicide tolerance, not deposited, described in WO 02/44407 or US-A 2009-26581 7); Event T25 (corn, herbicide tolerance, not deposited, described in US-A 2001-02901 4 or WO 01/051 654); Event T304-40 (cotton,  
 10 insect control - herbicide tolerance, deposited as ATCC PTA-8171, described in US-A 2010-077501 or WO 2008/1 22406); Event T342-1 42 (cotton, insect control, not deposited, described in WO 2006/1 28568) ; Event TC1 507 (corn, insect control - herbicide tolerance, not deposited, described in US-A 2005-039226 or WO 2004/099447); Event VIP1034 (corn, insect control - herbicide tolerance, deposited as ATCC PTA-3925., described in WO 03/052073), Event 3231 6 (corn, insect control-  
 15 herbicide tolerance, deposited as PTA-1 1507, described in WO 2011/084632), Event 4114 (corn, insect control-herbicide tolerance, deposited as PTA-1 1506, described in WO 2011/084621 ).

Among the diseases of plants or crops that can be controlled by the method according to the  
 20 invention, mention can be made of :

Powdery mildew diseases such as :

Blumeria diseases, caused for example by *Blumeria graminis* ;

Podosphaera diseases, caused for example by *Podosphaera leucotricha* ;

Sphaerotheca diseases, caused for example by *Sphaerotheca fuliginea* ;

25 Uncinula diseases, caused for example by *Uncinula necator* ;

Rust diseases such as :

Gymnosporangium diseases, caused for example by *Gymnosporangium sabinae* ;

Hemileia diseases, caused for example by *Hemileia vastatrix* ;

Phakopsora diseases, caused for example by *Phakopsora pachyrhizi* or *Phakopsora meibomiae* ;

30 Puccinia diseases, caused for example by *Puccinia recondite*, *Puccinia graminis* or *Puccinia striiformis*;

Uromyces diseases, caused for example by *Uromyces appendiculatus* ;

Oomycete diseases such as :

Albugo diseases caused for example by *Albugo Candida*;

35 Bremia diseases, caused for example by *Bremia lactucae* ;

Peronospora diseases, caused for example by *Peronospora pisi* or *P. brassicae* ;

Phytophthora diseases, caused for example by *Phytophthora infestans* ;

Plasmopara diseases, caused for example by *Plasmopara viticola* ;

Pseudoperonospora diseases, caused for example by *Pseudoperonospora humuli* or

40 *Pseudoperonospora cubensis* ;

Pythium diseases, caused for example by *Pythium ultimum* ;

Leafspot, leaf blotch and leaf blight diseases such as :

Alternaria diseases, caused for example by *Alternaria solani* ;

Cercospora diseases, caused for example by *Cercospora beticola* ;

Cladosporium diseases, caused for example by *Cladosporium cucumerinum* ;

5 Cochliobolus diseases, caused for example by *Cochliobolus sativus* (Conidiaform: Drechslera, Syn: Helminthosporium) or *Cochliobolus miyabeanus* ;

Colletotrichum diseases, caused for example by *Colletotrichum lindemuthanium* ;

Cycloconium diseases, caused for example by *Cycloconium oleaginum* ;

Diaporthe diseases, caused for example by *Diaporthe citri* ;

10 Elsinoe diseases, caused for example by *Elsinoe fawcettii* ;

Gloeosporium diseases, caused for example by *Gloeosporium laeticolor* ;

Glomerella diseases, caused for example by *Glomerella cingulata* ;

Guignardia diseases, caused for example by *Guignardia bidwelli* ;

Leptosphaeria diseases, caused for example by *Leptosphaena maculans* ; *Leptosphaena nodorum* ;

15 Magnaporthe diseases, caused for example by *Magnaporthe grisea* ;

Mycosphaerella diseases, caused for example by *Mycosphaerella graminicola* ; *Mycosphaerella arachidicola* ; *Mycosphaerella fijiensis* ;

Phaeosphaeria diseases, caused for example by *Phaeosphaeria nodorum* ;

Pyrenophora diseases, caused for example by *Pyrenophora teres*, or *Pyrenophora tritici repentis* ;

20 Ramularia diseases, caused for example by *Ramularia collocygni* , or *Ramularia areola* ;

Rhynchosporium diseases, caused for example by *Rhynchosporium secalis* ;

Septoria diseases, caused for example by *Septoria apii* or *Septoria lycopersici* ;

Typhula diseases, caused for example by *Typhula incarnata* ;

Venturia diseases, caused for example by *Venturia inaequalis* ;

25 Root, Sheath and stem diseases such as :

Corticium diseases, caused for example by *Corticium graminearum* ;

Fusarium diseases, caused for example by *Fusarium oxysporum* ;

Gaeumannomyces diseases, caused for example by *Gaeumannomyces graminis* ;

Rhizoctonia diseases, caused for example by *Rhizoctonia solani* ;

30 Sarocladium diseases caused for example by *Sarocladium oryzae*;

Sclerotium diseases caused for example by *Sclerotium oryzae*;

Tapesia diseases, caused for example by *Tapesia acuformis* ;

Thielaviopsis diseases, caused for example by *Thielaviopsis basicola* ;

Ear and panicle diseases such as :

35 Alternaria diseases, caused for example by *Alternaria spp.* ;

Aspergillus diseases, caused for example by *Aspergillus flavus* ;

Cladosporium diseases, caused for example by *Cladosporium spp.* ;

Claviceps diseases, caused for example by *Claviceps purpurea* ;

Fusarium diseases, caused for example by *Fusarium culmorum* ;

40 Gibberella diseases, caused for example by *Gibberella zeae* ;

Monographella diseases, caused for example by *Monographella nivalis* ;

Smut and bunt diseases such as :

Sphacelotheca diseases, caused for example by *Sphacelotheca reiliana* ;

Tilletia diseases, caused for example by *Tilletia caries* ;

5 Urocystis diseases, caused for example by *Urocystis occulta* ;

Ustilago diseases, caused for example by *Ustilago nuda* ;

Fruit rot and mould diseases such as :

Aspergillus diseases, caused for example by *Aspergillus flavus* ;

Botrytis diseases, caused for example by *Botrytis cinerea* ;

10 Penicillium diseases, caused for example by *Penicillium expansum* ;

Rhizopus diseases caused by example by *Rhizopus stolonifer*

Sclerotinia diseases, caused for example by *Sclerotinia sclerotiorum* ;

Verticillium diseases, caused for example by *Verticillium alboarum* ;

Seed and soilborne decay, mould, wilt, rot and damping-off diseases :

15 Alternaria diseases, caused for example by *Alternaria brassicicola* ;

Aphanomyces diseases, caused for example by *Aphanomyces euteiches* ;

Ascochyta diseases, caused for example by *Ascochyta lentis* ;

Aspergillus diseases, caused for example by *Aspergillus flavus* ;

Cladosporium diseases, caused for example by *Cladosporium herbarum* ;

20 Cochliobolus diseases, caused for example by *Cochliobolus sativus* ;

(Conidiaform: *Drechslera*, *Bipolaris* Syn: *Helminthosporium*) ;

Colletotrichum diseases, caused for example by *Colletotrichum coccodes* ;

Fusarium diseases, caused for example by *Fusarium culmorum* ;

Gibberella diseases, caused for example by *Gibberella zeae* ;

25 Macrophomina diseases, caused for example by *Macrophomina phaseolina* ;

Monographella diseases, caused for example by *Monographella nivalis* ;

Penicillium diseases, caused for example by *Penicillium expansum* ;

Phoma diseases, caused for example by *Phoma lingam* ;

Phomopsis diseases, caused for example by *Phomopsis sojae* ;

30 Phytophthora diseases, caused for example by *Phytophthora cactorum* ;

Pyrenophora diseases, caused for example by *Pyrenophora graminea* ;

Pyricularia diseases, caused for example by *Pyricularia oryzae* ;

Pythium diseases, caused for example by *Pythium ultimum* ;

Rhizoctonia diseases, caused for example by *Rhizoctonia solani* ;

35 Rhizopus diseases, caused for example by *Rhizopus oryzae* ;

Sclerotium diseases, caused for example by *Sclerotium rolfsii* ;

Septoria diseases, caused for example by *Septoria nodorum* ;

Typhula diseases, caused for example by *Typhula incarnata* ;

Verticillium diseases, caused for example by *Verticillium dahliae* ;

40 Canker, broom and dieback diseases such as :

Nectria diseases, caused for example by *Nectria galligena* ;

Blight diseases such as :

Monilinia diseases, caused for example by *Monilinia laxa* ;

Leaf blister or leaf curl diseases such as :

5 Exobasidium diseases caused for example by *Exobasidium vexans* ;

Taphrina diseases, caused for example by *Taphrina deformans* ;

Decline diseases of wooden plants such as :

Esca diseases, caused for example by *Phaemoniella clamydospora* ;

Eutypa dyeback, caused for example by *Eutypa lata* ;

10 Ganoderma diseases caused for example by *Ganoderma boninense* ;

Rigidoporus diseases caused for example by *Rigidoporus lignosus* ;

Diseases of Flowers and Seeds such as :

Botrytis diseases caused for example by *Botrytis cinerea* ;

Diseases of Tubers such as :

15 Rhizoctonia diseases caused for example by *Rhizoctonia solani* ;

Helminthosporium diseases caused for example by *Helminthosporium solani* ;

Club root diseases such as :

Plasmodiophora diseases, cause for example by *Plasmodiophora brassicae* ;

Diseases caused by Bacterial Organisms such as :

20 Xanthomonas species for example *Xanthomonas campestris* pv. *oryzae* ;

Pseudomonas species for example *Pseudomonas syringae* pv. *lachrymans* ;

Erwinia species for example *Erwinia amylovora*.

The composition according to the invention may also be used against fungal diseases liable to grow  
25 on or inside timber. The term "timber" means all types of species of wood, and all types of working of  
this wood intended for construction, for example solid wood, high-density wood, laminated wood, and  
plywood. The method for treating timber according to the invention mainly consists in contacting one  
or more compounds according to the invention or a composition according to the invention; this  
includes for example direct application, spraying, dipping, injection or any other suitable means.

30

The dose of active compound usually applied in the method of treatment according to the invention is  
generally and advantageously from 10 to 800 g/ha, preferably from 50 to 300 g/ha for applications in  
foliar treatment. The dose of active substance applied is generally and advantageously from 2 to 200 g  
per 100 kg of seed, preferably from 3 to 150 g per 100 kg of seed in the case of seed treatment.

35

It is clearly understood that the doses indicated herein are given as illustrative examples of the method  
according to the invention. A person skilled in the art will know how to adapt the application doses,  
notably according to the nature of the plant or crop to be treated.

40

The compounds or mixtures according to the invention can also be used for the preparation of  
composition useful to curatively or preventively treat human or animal fungal diseases such as, for

example, mycoses, dermatoses, trichophyton diseases and candidiasis or diseases caused by *Aspergillus spp.*, for example *Aspergillus fumigatus*.

The present invention further relates to the use of compounds of the formula (I) as herein defined for the control of phytopathogenic fungi.

The present invention further relates to the use of compounds of the formula (I) as herein defined for the treatment of transgenic plants.


The present invention further relates to the use of compounds of the formula (I) as herein defined for the treatment of seed and of seed of transgenic plants.

The present invention further relates to a process for producing compositions for controlling phytopathogenic harmful fungi, characterized in that derivatives of the formula (I) as herein defined are mixed with extenders and/or surfactants.

The various aspects of the invention will now be illustrated with reference to the following table of compound examples and the following preparation or efficacy examples.

15


Table 1 illustrates in a non-limiting manner examples of compounds of formula (I) according to the invention :



20

(I)

wherein A can be selected in the list consisting of the following groups : A-G1 , A-G2, A-G3, A-G4, A-G5, A-G6 and A-G7 :



wherein \* denotes the point of attachment the group A to the (thio)carbonyl moiety.

- 5 In table 1, unless otherwise specified, M+H (ApCl<sup>+</sup>) means the molecular ion peak plus 1 a.m.u. (atomic mass unit) as observed in mass spectroscopy via positive atmospheric pressure chemical ionisation.

In table 1, the logP values were determined in accordance with EEC Directive 79/831 Annex V.A8 by HPLC (High Performance Liquid Chromatography) on a reversed-phase column (C 18), using the  
10 method described below :

Temperature: 40 °C ; Mobile phases : 0.1% aqueous formic acid and acetonitrile ; linear gradient from 10% acetonitrile to 90% acetonitrile.


Calibration was carried out using unbranched alkan-2-ones (comprising 3 to 16 carbon atoms) with known logP values (determination of the logP values by the retention times using linear interpolation  
15 between two successive alkanones). lambda-max-values were determined using UV-spectra from 200 nm to 400 nm and the peak values of the chromatographic signals.

In table 1, "position" denotes the point of attachment of the -L-B residue to the phenyl ring.


Tab le 1.:


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.1	A-G1	O	cyclopropyl	H	H	-	2-	oxetan-3-yl	380	1.51	table 5
I.2	A-G2	O	cyclopropyl	H	H	-	2-	oxetan-3-yl	396	1.56	table 5
I.3	A-G1	O	cyclopropyl	H	H	-	2-	2-methyloxiran-2-yl	380	2.68	table 5
I.4	A-G1	O	cyclopropyl	H	H	5-Cl	2-	2-methyloxiran-2-yl	414	3.19	table 5
I.5	A-G1	O	cyclopropyl	H	H	5-Me	2-	2-methyloxiran-2-yl	394	3.04	table 5
I.6	A-G3	O	cyclopropyl	H	H	5-F	2-	2-methyloxiran-2-yl	380	2.53	table 5
I.7	A-G1	O	cyclopropyl	H	H	5-F	2-	2-methyloxiran-2-yl	398	2.86	table 5
I.8	A-G2	O	cyclopropyl	H	H	5-F	2-	2-methyloxiran-2-yl	414	3.00	table 5
I.9	A-G1	O	cyclopropyl	H	H	5,6-diCl	2-	2-methyloxiran-2-yl	448	3.55	table 5
I.10	A-G2	O	cyclopropyl	H	H	5-Cl-6-F	2-	2-methyloxiran-2-yl	448	3.37	table 5


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.11	A-G2	O	cyclopropyl	H	H	5,6-diCl	2-	2-methyloxiran-2-yl	464	3.67	table 5
I.12	A-G1	O	cyclopropyl	H	H	6-F-5-Me	2-	2-methyloxiran-2-yl	412	3.17	table 5
I.13	A-G1	O	cyclopropyl	H	H	5-Cl-6-F	2-	2-methyloxiran-2-yl	432	3.29	table 5
I.14	A-G1	O	cyclopropyl	H	H	5-Me	2-	2-methyloxiran-2-yl	428	3.41	table 5
I.15	A-G1	O	cyclopropyl	H	H	-	2-	3-methyloxetan-3-yl	394	1.94	table 5
I.16	A-G2	O	cyclopropyl	H	H	-	2-	3-methyloxetan-3-yl	410	2.08	table 5
I.17	A-G1	O	cyclopropyl	H	H	2-Cl	5-	6-oxabicyclo[3.1.0]hex-1-yl	440	3.59	table 5
I.18	A-G1	O	cyclopropyl	H	H	5-Cl	2-	6-oxabicyclo[3.1.0]hex-1-yl	440	3.60	
I.19	A-G1	O	cyclopropyl	H	H	-	2-	6-oxabicyclo[3.1.0]hex-1-yl	406	3.12	
I.20	A-G1	O	cyclopropyl	H	H	-	2-	2-methyl-1,3-dioxolan-2-yl	410	2.94	table 5
I.21	A-G1	O	cyclopropyl	H	H	4-Me	2-	2-methyl-1,3-dioxolan-2-yl	424	3.25	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
l.22	A-G1	O	cyclopropyl	H	H	2-Cl	5-	7-oxabicyclo[4.1.0]hept-1-yl	454	3.99	table 5
l.23	A-G1	O	cyclopropyl	H	H	5-Cl	2-	7-oxabicyclo[4.1.0]hept-1-yl	454	3.99	table 5
l.24	A-G1	O	cyclopropyl	H	H	-	2-	7-oxabicyclo[4.1.0]hept-1-yl	420	3.48	table 5
l.25	A-G1	O	cyclopropyl	H	H	5-Cl	2-	8-oxabicyclo[5.1.0]oct-1-yl	468	4.44	
l.26	A-G1	O	cyclopropyl	H	H	-	2-	8-oxabicyclo[5.1.0]oct-1-yl	434	3.94	
l.27	A-G1	O	cyclopropyl	H	H	5-F	2-	3-furyl	408	3.33	table 5
l.28	A-G1	O	cyclopropyl	H	H	5-Cl	2-	3-furyl	424	3.64	table 5
l.29	A-G1	O	cyclopropyl	H	H	-	2-	3-furyl	390	3.23	table 5
l.30	A-G2	O	cyclopropyl	H	H	-	2-	2-furyl	406	3.46	table 5
l.31	A-G3	O	cyclopropyl	H	H	-	2-	2-furyl	372	2.96	table 5
l.32	A-G1	O	cyclopropyl	H	H	-	2-	2-furyl	390	3.31	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
1.33	A-G4	O	cyclopropyl	H	H	-	2-	2-furyl	450	4.01	table 5
1.34	A-G1	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	424	3.81	table 5
1.35	A-G2	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	440	4.01	table 5
1.36	A-G3	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	406	3.44	table 5
1.37	A-G2	O	cyclopropyl	H	H	5-Me	2-	3-furyl	420	3.71	table 5
1.38	A-G3	O	cyclopropyl	H	H	5-Me	2-	3-furyl	386	3.19	table 5
1.39	A-G1	O	cyclopropyl	H	H	5-Me	2-	3-furyl	404	3.55	table 5
1.40	A-G5	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	385	3.90	table 5
1.41	A-G4	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	484	4.61	table 5
1.42	A-G5	O	cyclopropyl	H	H	-	2-	2-furyl	371	3.44	table 5
1.43	A-G3	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	3-furyl	400	3.50	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.44	A-G1	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	3-furyl	418	3.80	table 5
I.45	A-G2	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	3-furyl	434	3.92	table 5
I.46	A-G6	O	cyclopropyl	H	H	2-Cl	5-	2-furyl	423	3.99	table 5
I.47	A-G1	O	cyclopropyl	H	H	5-F	2-	1-methyl-1H-pyrrol-3-yl	421	3.37	table 5
I.48	A-G1	O	cyclopropyl	H	H	5-Cl	2-	1-methyl-1H-pyrrol-3-yl	437	3.71	table 5
I.49	A-G1	O	cyclopropyl	H	H	-	2-	1-methyl-1H-pyrrol-3-yl	403	3.25	table 5
I.50	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1-methyl-1H-pyrrol-3-yl	445	3.99	table 5
I.51	A-G1	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrrol-3-yl	417	3.55	table 5
I.52	A-G3	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrrol-3-yl	399	3.19	table 5
I.53	A-G2	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrrol-3-yl	433	3.69	table 5
I.54	A-G1	O	cyclopropyl	H	H	-	2-	1-methyl-1H-pyrazol-4-yl	404	2.34	table 5


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
1.55	A-G1	O	cyclopropyl	H	H	5-Cl	2-	1-methyl-1H-pyrazol-4-yl	438	2.80	table 5
1.56	A-G1	O	cyclopropyl	H	H	5-F	2-	1-methyl-1H-pyrazol-4-yl	422	2.50	table 5
1.57	A-G2	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	434	2.71	table 5
1.58	A-G1	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	418	2.63	table 5
1.59	A-G3	O	cyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	400	2.37	table 5
1.60	A-G1	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	432	2.96	table 5
1.61	A-G2	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	448	3.06	table 5
1.62	A-G3	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	414	2.66	table 5
1.63	A-G1	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	432	2.82	table 5
1.64	A-G2	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	1-methyl-1H-pyrazol-4-yl	448	2.86	table 5
1.65	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1-methyl-1H-pyrazol-4-yl	446	3.13	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
1.66	A-G1	O	cyclopropyl	H	H	5-Cl	2-	3-thienyl	440	3.99	table 5
1.67	A-G1	O	cyclopropyl	H	H	-	2-	3-thienyl	406	3.55	table 5
1.68	A-G1	O	cyclopropyl	H	H	5-Cl	2-	2-thienyl	440	4.11	table 5
1.69	A-G1	O	cyclopropyl	H	H	-	2-	2-thienyl	406	3.64	table 5
1.70	A-G3	O	cyclopropyl	H	H	-	2-	3-thienyl	388	3.17	table 5
1.71	A-G3	O	cyclopropyl	H	H	5-Cl	2-	2-thienyl	422	3.69	table 5
1.72	A-G3	O	cyclopropyl	H	H	-	2-	2-thienyl	388	3.25	table 5
1.73	A-G1	S	cyclopropyl	H	H	-	2-	3-thienyl	422	4.21	table 5
1.74	A-G1	S	cyclopropyl	H	H	5-Cl	2-	2-thienyl	456	4.69	table 5
1.75	A-G1	S	cyclopropyl	H	H	-	2-	2-thienyl	422	4.29	table 5
1.76	A-G2	O	cyclopropyl	H	H	-	2-	3-thienyl	422	3.71	table 5


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
1.77	A-G2	O	cyclopropyl	H	H	5-Cl	2-	2-thienyl	456	4.29	table 5
1.78	A-G2	O	cyclopropyl	H	H	-	2-	2-thienyl	422	3.78	table 5
1.79	A-G2	S	cyclopropyl	H	H	5-Cl	2-	2-thienyl	472	4.92	table 5
1.80	A-G2	S	cyclopropyl	H	H	-	2-	3-thienyl	438	4.39	table 5
1.81	A-G2	S	cyclopropyl	H	H	-	2-	2-thienyl	438	4.49	table 5
1.82	A-G1	O	cyclopropyl	Me	H	-	2-	3-thienyl	420	3.78	table 5
1.83	A-G1	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	440	4.06	table 5
1.84	A-G3	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	422	3.68	table 5
1.85	A-G2	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	456	4.29	table 5
1.86	A-G1	O	2-methyl cyclopropyl	H	H	-	2-	3-thienyl	420	3.83	table 5
1.87	A-G3	O	2-methyl cyclopropyl	H	H	-	2-	3-thienyl	402	3.48	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
1.88	A-G2	O	2-methyl cyclopropyl	H	H	-	2-	3-thienyl	436	3.92	table 5
1.89	A-G4	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	500	4.78	table 5
1.90	A-G5	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	421	4.16	table 5
1.91	A-G4	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	500	4.91	table 5
1.92	A-G5	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	421	4.06	table 5
1.93	A-G3	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	422	3.59	table 5
1.94	A-G2	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	456	4.16	table 5
1.95	A-G1	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	440	3.96	table 5
1.96	A-G6	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	439	4.14	table 5
1.97	A-G6	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	439	4.26	table 5
1.98	A-G7	O	cyclopropyl	H	H	2-Cl	5-	3-thienyl	386	4.59	table 5


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.99	A-G7	O	cyclopropyl	H	H	2-Cl	5-	2-thienyl	386	4.72	table 5
I.100	A-G1	O	cyclopropyl	H	H	-	2-	4-methyl-2-thienyl	420	3.99	table 5
I.101	A-G1	O	cyclopropyl	H	H	5-F	2-	4-methyl-2-thienyl	438	4.09	table 5
I.102	A-G1	O	cyclopropyl	H	H	5-F	2-	4-methyl-3-thienyl	438	3.94	table 5
I.103	A-G1	O	cyclopropyl	H	H	5-Cl	2-	4-methyl-2-thienyl	454	4.51	table 5
I.104	A-G1	O	cyclopropyl	H	H	5-Cl	2-	4-methyl-3-thienyl	454	4.34	table 5
I.105	A-G1	O	cyclopropyl	H	H	-	2-	4-methyl-3-thienyl	420	3.90	table 5
I.106	A-G1	O	cyclopropyl	H	H	2- <i>i</i> Pr	5-	4-methyl-2-thienyl	462	4.93	table 5
I.107	A-G2	O	cyclopropyl	H	H	5-Me	2-	4-methyl-3-thienyl	450	4.41	table 5
I.108	A-G1	O	cyclopropyl	H	H	5-Me	2-	4-methyl-3-thienyl	434	4.25	table 5
I.109	A-G3	O	cyclopropyl	H	H	5-Me	2-	4-methyl-3-thienyl	416	3.83	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.110	A-G1	O	cyclopropyl	H	H	2-iPr	5-	4-methyl-3-thienyl	462	4.72	table 5
I.111	A-G1	O	cyclopropyl	H	H	5-Me	2-	3,5-dimethyl-4,5-dihydro-1,2-oxazol-5-yl	435	2.98	table 5
I.112	A-G1	O	cyclopropyl	H	H	5-F	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	450	2.66	table 5
I.113	A-G1	O	cyclopropyl	H	H	5-Cl	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	466	2.98	table 5
I.114	A-G1	O	cyclopropyl	H	H	-	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	432	2.52	table 5
I.115	A-G1	O	cyclopropyl	H	H	5-Me	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	446	2.71	table 5
I.116	A-G3	O	cyclopropyl	H	H	5-Me	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	428	2.45	table 5
I.117	A-G2	O	cyclopropyl	H	H	5-Me	2-	1,3,5-trimethyl-1H-pyrazol-4-yl	462	2.80	table 5
I.118	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1,3,5-trimethyl-1H-pyrazol-4-yl	474	3.23	table 5
I.119	A-G1	O	cyclopropyl	H	H	5-F	2-	3-methoxy-2-thienyl	454	3.63	table 5
I.120	A-G1	O	cyclopropyl	H	H	-	2-	3-methoxy-2-thienyl	436	3.52	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.121	A-G1	O	cyclopropyl	H	H	5-Cl	2-	3-methoxy-2-thienyl	470	3.99	table 5
I.122	A-G1	O	cyclopropyl	H	H	4-F	2-	3-methoxy-2-thienyl	454	3.70	table 5
I.123	A-G1	O	cyclopropyl	H	H	4-Cl	2-	3-methoxy-2-thienyl	470	4.14	table 5
I.124	A-G3	O	cyclopropyl	H	H	5-Me	2-	3-methoxy-2-thienyl	432	3.46	table 5
I.125	A-G1	O	cyclopropyl	H	H	5-Me	2-	3-methoxy-2-thienyl	450	3.83	table 5
I.126	A-G2	O	cyclopropyl	H	H	5-Me	2-	3-methoxy-2-thienyl	466	3.99	table 5
I.127	A-G1	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	3-methoxy-2-thienyl	464	4.21	table 5
I.128	A-G2	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	3-methoxy-2-thienyl	480	4.41	table 5
I.129	A-G1	O	cyclopropyl	H	H	5-F	2-	1-benzofuran-3-yl	458	4.06	table 5
I.130	A-G1	O	cyclopropyl	H	H	5-Cl	2-	1-benzofuran-3-yl	474	4.44	table 5
I.131	A-G1	O	cyclopropyl	H	H	-	2-	1-benzofuran-3-yl	440	3.96	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.132	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1-benzofuran-3-yl	482	4.90	table 5
I.133	A-G2	O	cyclopropyl	H	H	5-Me	2-	1-benzofuran-3-yl	470	4.49	table 5
I.134	A-G1	O	cyclopropyl	H	H	5-Me	2-	1-benzofuran-3-yl	454	4.32	table 5
I.135	A-G3	O	cyclopropyl	H	H	5-Me	2-	1-benzofuran-3-yl	436	3.89	table 5
I.136	A-G1	O	cyclopropyl	H	H	5-F	2-	2-chloro-3-thienyl	458	4.04	table 5
I.137	A-G1	O	cyclopropyl	H	H	5-Cl	2-	2-chloro-3-thienyl	474	4.41	table 5
I.138	A-G1	O	cyclopropyl	H	H	-	2-	2-chloro-3-thienyl	440	3.92	table 5
I.139	A-G1	O	cyclopropyl	H	H	-	2-	5-chloro-2-thienyl	440	4.32	table 5
I.140	A-G3	O	cyclopropyl	H	H	-	2-	5-chloro-2-thienyl	422	3.87	table 5
I.141	A-G2	O	cyclopropyl	H	H	-	2-	5-chloro-2-thienyl	456	4.51	table 5
I.142	A-G2	S	cyclopropyl	H	H	-	2-	5-chloro-2-thienyl	472	5.19	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.143	A-G1	O	cyclopropyl	H	H	2-Cl	5-	5-chloro-2-thienyl	474	4.83	table 5
I.144	A-G3	O	cyclopropyl	H	H	2-Cl	5-	5-chloro-2-thienyl	456	4.41	table 5
I.145	A-G2	O	cyclopropyl	H	H	2-Cl	5-	5-chloro-2-thienyl	490	5.06	table 5
I.146	A-G5	O	cyclopropyl	H	H	5-Cl	2-	5-chloro-2-thienyl	455	4.85	table 5
I.147	A-G4	O	cyclopropyl	H	H	5-Cl	2-	5-chloro-2-thienyl	534	5.68	table 5
I.148	A-G2	O	cyclopropyl	H	H	5-Me	2-	2-chloro-3-thienyl	470	4.44	table 5
I.149	A-G3	O	cyclopropyl	H	H	5-Me	2-	2-chloro-3-thienyl	436	3.85	table 5
I.150	A-G1	O	cyclopropyl	H	H	5-Me	2-	2-chloro-3-thienyl	454	4.26	table 5
I.151	A-G6	O	cyclopropyl	H	H	2-Cl	5-	5-chloro-2-thienyl	473	5.00	table 5
I.152	A-G7	O	cyclopropyl	H	H	2-Cl	5-	5-chloro-2-thienyl	420	5.54	table 5
I.153	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1-methyl-1H-indol-3-yl	495	4.74	table 5


Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.154	A-G1	O	cyclopropyl	H	H	5-F	2-	1-benzothiophen-3-yl	474	4.39	table 5
I.155	A-G1	O	cyclopropyl	H	H	5-Cl	2-	1-benzothiophen-3-yl	490	4.74	table 5
I.156	A-G1	O	cyclopropyl	H	H	-	2-	1-benzothiophen-3-yl	456	4.27	table 5
I.157	A-G1	O	cyclopropyl	H	H	2-iPr	5-	1-benzothiophen-3-yl	498	5.17	table 5
I.158	A-G1	O	cyclopropyl	H	H	4-F	2-	1-benzothiophen-3-yl	474	4.44	table 5
I.159	A-G1	O	cyclopropyl	H	H	4-Cl	2-	1-benzothiophen-3-yl	490	4.92	table 5
I.160	A-G1	O	cyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	470	4.64	table 5
I.161	A-G3	O	cyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	452	4.15	table 5
I.162	A-G2	O	cyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	486	4.83	table 5
I.163	A-G1	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	484	4.98	table 5
I.164	A-G2	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	500	5.19	table 5

Example	A	T	Z <sup>1</sup>	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP	NMR
I.165	A-G3	O	rel-(1R,2R)-2-methylcyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	466	4.46	table 5
I.166	A-G1	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	484	4.92	table 5
I.167	A-G2	O	rel-(1R,2S)-2-methylcyclopropyl	H	H	5-Me	2-	1-benzothiophen-3-yl	500	5.05	table 5
I.168	A-G1	O	cyclopropyl	H	H	-	2-	(4,6-dichloro pyrimidin-2-yl)oxy	486	3.50	table 5
I.169	A-G1	O	cyclopropyl	H	H	-	2-	(tetrahydro-2H-pyran-4-yl)oxy	424	2.80	
I.170	A-G1	O	cyclopropyl	H	H	-	2-	[3-chloro-5-(trifluoromethyl)pyridin-2-yl]sulfanyl			
I.171	A-G1	O	cyclopropyl	Me	H	-	2-	(5-chloropyridin-2-yl)methyl	463	3.68	

Note: Me : methyl ; iPr : isopropyl

Table 2 illustrates in a non limiting manner examples of compounds of formula (Ha) according to the invention,

wherein L is a direct bond and B<sup>1</sup> is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothieryl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring,



In table 2, M+H (Apcl<sup>+</sup>) and logP are defined as for table 1.

10

In table 2, "position" denotes the point of attachment of the -L-B<sup>1</sup> residue to the phenyl ring.


Table 2:

Example	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP
Ila.1	H	H	5-Cl	2-	2-furyl		0.99
Ila.2	H	H	5-Cl	2-	2-thienyl		1.10
Ila.3	H	H	-	2-	3-thienyl	230	1.34
Ila.4	Me	H	-	2-	3-thienyl	244	1.42
Ila.5	H	H	5-Cl	2-	3-thienyl		1.12
Ila.6	H	H	-	2-	5-chloro-2-thienyl		1.28
Ila.7	H	H	5-Cl	2-	5-chloro-2-thienyl		1.51

15

Table 3 illustrates in a non limiting manner examples of compounds of formula (Iib) according to the invention,

wherein L is a direct bond and B<sup>1</sup> is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothieryl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring




5

In table 3, M+H (Apcl+) and logP are defined as for table 1.

In table 3, "position" denotes the point of attachment of the -L-B<sup>1</sup> residue to the phenyl ring.

10

Table 3:


Example	Z <sup>2</sup>	Z <sup>3</sup>	(X) <sub>n</sub>	position		M+H	logP
IIb.1	H	H	2-Cl	5-	2-furyl		1.08
IIb.2	H	H	2-Cl	5-	2-thienyl		1.29
IIb.3	H	H	2-Cl	5-	3-thienyl		1.28
IIb.4	H	H	2-Cl	5-	5-chloro-2-thienyl		1.58

15 Table 4 illustrates in a non limiting manner examples of compounds of formula (Xa) according to the invention,

wherein Z<sup>2</sup> and Z<sup>3</sup> are as herein-defined, A represents A<sup>13</sup>, X<sup>a1</sup> represents a halogen atom ; substituted or non-substituted Ci-Cs-alkyl ; Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; tri(CrC<sub>8</sub>-alkyl)silyl; substituted or non-substituted C<sub>1</sub>-Cs-alkoxy; or substituted or non-substituted Ci-Cs-alkylsulfanyl, X<sup>a2</sup> represents hydrogen or X<sup>a1</sup>, and U<sup>2</sup> represents a halogen atom such as chlorine, bromine or iodine, or a triflate group,

20


70



(Xa)

In table 4, M+H (Apcl+) and logP are defined as for table 1.

- 5 wherein A can be selected in the list consisting of the following groups : A-G1 .



A-G1

Table 4 :

Example	A	Z <sup>2</sup>	Z <sup>3</sup>	X <sup>a1</sup>	X <sup>a2</sup>	U <sup>2</sup>	M+H	logP
Xa.1	A-G1	H	H	2-iPr	-	5-Cl	360	3.46
Xa.2	A-G1	H	H	2-Et	-	5-Cl		
Xa.3	A-G1	H	H	2-CF <sub>3</sub>	-	5-Cl		
Xa.4	A-G1	H	H	2-CF <sub>3</sub>	6-F	5-Cl		

10

Note: Et : ethyl ; iPr : isopropyl

Table 5 provides the NMR data (<sup>1</sup>H) of a selected number of compounds from table 1.

- 15 The <sup>1</sup>H-NMR data of selected examples are stated in the form of <sup>1</sup>H-NMR peak lists. For each signal peak, the δ value in ppm and the signal intensity in brackets are listed.

Intensity of sharp signals correlates with the height of the signals in a printed example of a NMR spectrum in cm and shows the real relations of signal intensities. From broad signals several peaks or the middle of the signal and their relative intensity in comparison to the most intensive signal in the spectrum can be shown.

20

The  $^1\text{H-NMR}$  peak lists are similar to classical  $^1\text{H-NMR}$  prints and contain therefore usually all peaks, which are listed at classical NMR-interpretation. Additionally they can show like classical  $^1\text{H-NMR}$  prints signals of solvents, stereoisomers of the target compounds, which are also object of the invention, and/or peaks of impurities. To show compound signals in the delta-range of solvents and/or water the usual peaks of solvents, for example peaks of DMSO in DMSO-d6 and the peak of water are shown in our  $^1\text{H-NMR}$  peak lists and have usually on average a high intensity.

The peaks of stereoisomers of the target compounds and/or peaks of impurities have usually on average a lower intensity than the peaks of target compounds (for example with a purity >90%). Such stereoisomers and/or impurities can be typical for the specific preparation process. Therefore their peaks can help to recognize the reproduction of our preparation process via "side-products-fingerprints".

An expert, who calculates the peaks of the target compounds with known methods (MestreC, ACD-simulation, but also with empirically evaluated expectation values), can isolate the peaks of the target compounds as needed optionally using additional intensity filters. This isolation would be similar to relevant peak picking at classical  $^1\text{H-NMR}$  interpretation.

Further details of NMR-data description with peak lists can be found in the publication "Citation of NMR Peaklist Data within Patent Applications" of the Research Disclosure Database Number 564025.

Table 5 : NMR peak lists

Example	Solvent	Frequency (Mhz)	$^1\text{H-NMR}$
I.1	$\text{CDCl}_3$	250	7.2603 (23.03); 7.2507 (2.23); 7.2334 (3.16); 7.2186 (0.64); 7.2082 (0.72); 7.1897 (4.43); 7.1794 (2.18); 7.1696 (5.21); 7.1533 (4.18); 7.1406 (3.36); 7.1055 (0.49); 7.0893 (2.54); 7.0729 (1.94); 7.0536 (1.25); 6.9259 (6.30); 6.7111 (3.10); 4.5321 (1.39); 4.5140 (1.24); 4.4909 (2.41); 4.4735 (2.79); 4.4005 (2.54); 4.3636 (2.89); 4.3575 (2.25); 4.3215 (1.82); 3.9602 (2.38); 3.8994 (3.57); 3.8404 (16.00); 3.6650 (3.70); 3.6042 (2.47); 3.1925 (3.38); 3.1771 (1.34); 3.1526 (3.62); 2.8133 (1.27); 2.7931 (2.16); 2.7619 (1.83); 2.7437 (1.02); 1.8466 (0.54); 1.8242 (1.27); 1.8058 (2.32); 1.7863 (1.31); 1.7665 (0.64); 1.5642 (11.51); 1.2665 (2.30); 0.9074 (0.63); 0.8819 (2.03); 0.8543 (0.71); 0.5486 (1.18); 0.5277 (3.47); 0.5191 (3.96); 0.5130 (2.87); 0.5013 (4.63); 0.4931 (4.09); 0.4793 (3.26); 0.4697 (2.71); 0.4595 (1.91); 0.4406 (2.13); 0.4263 (1.70); 0.4078 (1.36); 0.3986 (0.60); 0.3904 (0.62); 0.3759 (0.40); 0.0700 (0.45); 0.0128 (0.57); -0.0002 (14.42); -0.0134 (0.43)
I.2	DMSO-d6	400	7.2994 (1.59); 7.2877 (2.13); 7.2801 (2.19); 7.2706 (1.78); 7.1918 (3.25); 7.1837 (4.40); 7.1740 (3.65); 7.1369 (4.80); 7.0032 (1.33); 5.7708 (4.82); 4.4065 (0.92); 4.3975 (1.05); 4.3808 (2.00); 4.3713 (1.96); 4.3422 (1.78); 4.3181 (2.34); 4.2931 (0.93); 3.9225 (16.00); 3.8950 (2.07); 3.8570 (2.52); 3.6026 (2.51); 3.5648 (1.98); 3.3475 (21.85); 3.3244 (0.75); 3.1860 (1.38); 3.1740 (1.37); 3.1483 (1.87); 3.1175 (1.75); 2.7515 (1.45); 2.7442 (1.51); 2.7234 (1.36); 2.7162 (1.29); 2.5111 (42.60); 1.8171 (1.78); 0.4935 (1.81); 0.4793 (2.06); 0.4555 (1.12); 0.4295 (0.93); 0.4207 (1.05); 0.3936 (1.50); 0.3066 (1.40)
I.3	DMSO-d6	500	9.6384 (2.36); 7.3995 (3.46); 7.3969 (3.55); 7.3852 (4.55); 7.3822 (4.71); 7.3516 (0.60); 7.3447 (1.05); 7.3312 (5.78); 7.3185 (4.29); 7.3154 (4.02); 7.3038 (3.94); 7.3003 (3.36); 7.2949 (3.57); 7.2920 (3.97); 7.2800 (3.97); 7.2776 (4.02); 7.2652 (1.46); 7.1566 (4.60); 7.1424 (3.84); 7.1170 (1.25); 7.1106 (3.55); 7.1048 (1.20); 7.0975 (0.99); 7.0029 (7.07); 6.9894 (1.88); 6.8951 (3.57); 6.8817 (0.94); 4.8943 (1.19); 4.8619 (1.94); 4.7952 (0.63); 4.7737 (4.53); 4.7649 (2.19); 4.7417 (4.09); 4.7103 (0.53); 4.0645 (0.44); 4.0565 (0.46); 4.0504 (0.95); 4.0363 (0.88); 4.0253 (0.37); 4.0220 (0.35); 3.8243 (13.53); 3.8029 (6.99); 3.7804 (0.42); 3.3129 (105.38); 3.2891 (16.75); 3.0316 (4.75); 3.0213 (5.25); 2.8411 (3.93); 2.8311

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(3.86); 2.6484 (0.68); 2.6443 (0.57); 2.6149 (0.56); 2.5991 (0.36); 2.5848 (0.33); 2.5525 (0.80); 2.5202 (32.84); 2.5168 (67.60); 2.5133 (92.23); 2.5097 (67.03); 2.3744 (0.56); 2.0008 (2.67); 1.7142 (0.35); 1.5686 (16.00); 1.3573 (0.40); 1.3436 (4.63); 1.3296 (4.59); 1.3122 (0.38); 1.2723 (0.87); 1.2545 (2.63); 1.2026 (1.03); 1.1884 (1.72); 1.1741 (0.87); 0.8860 (0.65); 0.8723 (1.67); 0.8670 (0.89); 0.8581 (0.87); 0.8530 (0.55); 0.6879 (5.11); 0.6746 (5.51); 0.6396 (0.48); 0.6059 (1.70); 0.5987 (2.00); 0.5913 (1.63); 0.5750 (3.56); 0.5686 (3.59)
I.4	CDCl <sub>3</sub>	500	7.3528 (3.69); 7.3364 (4.80); 7.2646 (51.88); 7.2511 (0.43); 7.2398 (3.53); 7.2356 (3.82); 7.2234 (2.62); 7.2192 (2.98); 7.1634 (5.24); 7.1597 (4.72); 7.0528 (0.32); 6.9732 (1.06); 6.8639 (2.16); 6.7545 (1.10); 5.3051 (9.73); 4.9418 (0.73); 4.9107 (1.94); 4.8778 (2.00); 4.8465 (0.77); 3.8466 (12.44); 3.0045 (4.18); 2.9939 (4.78); 2.8686 (2.64); 2.8590 (2.17); 2.5836 (0.33); 1.6250 (16.00); 1.5777 (30.04); 1.2522 (0.77); 0.7083 (2.57); 0.6951 (3.12); 0.6711 (2.51); 0.6643 (2.32); 0.6580 (2.28); 0.6485 (2.04); 0.6378 (1.18); 0.0768 (0.39); 0.0697 (8.80); 0.0624 (0.35); 0.0063 (1.42); -0.0002 (34.47); -0.0068 (1.29)
I.5	CDCl <sub>3</sub>	300	16.8730 (0.33); 7.4203 (0.33); 7.2992 (10.37); 7.1056 (3.68); 7.0220 (4.93); 6.9124 (2.08); 6.7333 (1.10); 4.9992 (0.93); 4.9558 (2.43); 4.9207 (2.53); 4.8599 (0.89); 4.8372 (0.53); 4.7848 (0.34); 4.5739 (0.50); 4.4074 (0.35); 4.1430 (0.43); 4.1193 (0.33); 4.0885 (0.33); 3.8704 (11.32); 3.0135 (3.70); 2.9011 (4.13); 2.5090 (0.48); 2.4435 (1.09); 2.3489 (16.00); 2.2212 (0.39); 2.0838 (1.37); 2.0530 (0.38); 1.9422 (0.32); 1.7524 (0.58); 1.6536 (13.23); 1.5954 (56.89); 1.5478 (0.59); 1.4996 (0.42); 1.4890 (0.36); 1.4457 (0.48); 1.2938 (2.50); 1.1232 (0.32); 0.9206 (0.89); 0.8148 (0.48); 0.7579 (0.96); 0.6899 (8.46); -3.4594 (0.32)
I.6	CDCl <sub>3</sub>	300	7.7871 (0.94); 7.3860 (2.03); 7.3668 (2.21); 7.3578 (2.39); 7.3386 (2.29); 7.2899 (1.07); 7.2809 (1.91); 7.1080 (2.06); 6.9511 (1.04); 6.9425 (1.21); 6.9240 (2.71); 6.9150 (2.24); 6.8957 (0.96); 6.8871 (1.06); 6.8471 (1.66); 6.8391 (1.43); 6.8141 (1.68); 6.8060 (1.41); 4.9821 (0.99); 4.9285 (3.58); 4.8915 (2.66); 4.8378 (0.74); 3.9788 (12.07); 3.9289 (0.38); 2.9906 (3.45); 2.9730 (4.55); 2.8994 (0.51); 2.8858 (1.12); 2.8760 (1.55); 2.8634 (3.95); 2.8491 (2.86); 2.8277 (0.71); 2.0428 (1.15); 1.6147 (16.00); 1.2814 (0.33); 1.2576 (0.64); 0.8184 (0.38); 0.7888 (2.90); 0.7665 (3.80); 0.7479 (1.82); 0.7414 (1.42); 0.7165 (2.92); 0.7087 (2.88); 0.7023 (2.91); 0.6890 (1.45); 0.6746 (0.81); 0.6637 (0.72); 0.0771 (0.54); -0.0002 (0.97)
I.7	CDCl <sub>3</sub>	400	7.3903 (2.12); 7.3759 (2.37); 7.3692 (2.40); 7.3548 (2.28); 7.2875 (1.51); 6.9981 (1.29); 6.9550 (1.25); 6.9484 (1.49); 6.9343 (2.22); 6.9277 (2.66); 6.9135 (1.15); 6.9069 (1.42); 6.8928 (2.20); 6.8865 (1.78); 6.8675 (2.74); 6.8616 (4.28); 6.7250 (1.34); 4.9635 (0.68); 4.9237 (2.29); 4.8945 (2.26); 4.8547 (0.66); 3.8379 (12.10); 2.9982 (3.95); 2.9850 (4.76); 2.9000 (1.02); 2.8726 (2.43); 2.8597 (1.91); 2.0435 (0.48); 1.6268 (16.00); 1.2755 (0.37); 1.2638 (0.64); 1.2581 (0.69); 0.8805 (0.91); 0.8629 (0.39); 0.7103 (2.39); 0.6939 (3.04); 0.6769 (1.89); 0.6537 (2.72); 0.6284 (0.85); 0.6197 (0.58); 0.0811 (0.50); -0.0002 (1.02)
I.8	CDCl <sub>3</sub>	400	7.3810 (1.96); 7.3668 (2.46); 7.3610 (2.68); 7.3467 (2.17); 7.2808 (4.24); 7.0151 (3.46); 7.0088 (3.99); 6.9899 (3.52); 6.9836 (3.98); 6.9515 (2.84); 6.9449 (2.40); 6.9308 (5.19); 6.9242 (4.43); 6.9102 (3.56); 6.9035 (2.44); 6.7768 (2.95); 6.6398 (1.48); 4.9653 (1.57); 4.9283 (1.58); 4.1465 (0.34); 4.1286 (1.05); 4.1108 (1.06); 4.0929 (0.39); 3.9243 (16.00); 3.8783 (0.59); 3.8580 (0.50); 3.0128 (3.08); 2.9999 (3.81); 2.9300 (2.13); 2.9178 (1.80); 2.8919 (3.36); 2.8795 (2.87); 2.0445 (4.83); 1.6396 (15.65); 1.5210 (0.35); 1.2761 (1.75); 1.2583 (3.64); 1.2404 (1.43); 0.8975 (0.75); 0.8807 (2.18); 0.8630 (1.03); 0.6010 (8.12); 0.5843 (5.26); 0.0777 (1.49); -0.0002 (2.81)
I.9	CDCl <sub>3</sub>	300	7.4530 (2.69); 7.4252 (5.26); 7.3694 (4.85); 7.3416 (2.73); 7.2996 (0.82); 7.1060 (1.53); 6.9242 (3.25); 6.7423 (1.65); 5.1236 (0.44); 5.0745 (4.65); 5.0657 (5.09); 5.0167 (0.52); 3.8173 (10.75); 3.0374 (2.57); 3.0201 (4.05); 2.9672 (2.33); 2.9501 (1.54); 2.6532 (0.50); 2.0094 (2.03); 1.6222 (16.00); 0.7063 (0.48); 0.6760 (0.81); 0.6526 (0.49); 0.6430 (0.60); 0.6020 (1.95); 0.5852 (1.84); 0.5791 (2.01); 0.5603 (1.13); 0.5520 (1.00); 0.5374 (0.60); 0.5220 (0.55); 0.0842 (0.47); -0.0002 (0.34)
I.10	CDCl <sub>3</sub>	500	7.3506 (1.27); 7.3343 (1.99); 7.3195 (1.56); 7.2606 (10.07); 7.2216 (1.32); 7.2055

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.95); 7.1913 (0.35); 6.8684 (1.34); 6.7591 (2.82); 6.6498 (1.45); 3.8961 (2.39); 3.8865 (16.00); 3.0364 (1.37); 3.0269 (1.45); 2.9285 (0.84); 2.8399 (0.40); 1.9997 (5.81); 1.8279 (1.14); 1.6494 (6.86); 1.2583 (0.33); 0.6238 (1.17); 0.5778 (0.81); 0.5732 (0.82); 0.5647 (0.91); 0.5611 (0.86); 0.5521 (0.82); 0.5398 (0.57); -0.0002 (6.43)
I.11	CDCl <sub>3</sub>	500	7.4438 (2.03); 7.4271 (2.86); 7.3603 (0.44); 7.3513 (1.14); 7.3438 (0.65); 7.3346 (0.86); 7.3262 (2.18); 7.2599 (19.96); 6.9129 (0.46); 6.8881 (1.01); 6.7786 (2.13); 6.6692 (1.11); 5.2961 (0.42); 4.0689 (0.35); 3.8924 (16.00); 3.3576 (0.34); 3.3384 (0.35); 3.0547 (1.30); 3.0446 (1.44); 2.9250 (0.93); 2.9169 (0.85); 2.7708 (0.35); 1.9999 (13.81); 1.8374 (1.99); 1.7973 (1.10); 1.6527 (0.36); 1.6269 (8.07); 1.5519 (3.66); 1.2574 (0.34); 0.4860 (0.41); 0.4473 (1.25); 0.3271 (0.33); 0.3197 (0.38); 0.3152 (0.39); 0.3116 (0.39); 0.3075 (0.41); 0.0063 (0.55); -0.0002 (12.18); -0.0069 (0.42)
I.12	CDCl <sub>3</sub>	300	9.6302 (0.46); 7.2736 (2.91); 7.1685 (0.67); 7.1444 (7.62); 7.1257 (2.33); 7.1042 (1.48); 7.0880 (0.46); 7.0399 (0.36); 7.0202 (0.38); 6.9225 (2.33); 6.9026 (0.35); 6.8580 (0.44); 6.7407 (1.14); 4.9522 (5.55); 4.7434 (0.33); 3.8056 (13.24); 2.9871 (2.72); 2.9696 (3.86); 2.8898 (2.14); 2.8725 (1.62); 2.6298 (0.56); 2.2978 (0.83); 2.2845 (1.34); 2.2666 (12.09); 2.2592 (10.18); 2.0070 (1.77); 1.7954 (1.01); 1.6145 (16.00); 1.5315 (0.37); 1.3877 (0.82); 1.3644 (0.81); 1.2583 (0.53); 0.7031 (0.99); 0.6802 (2.26); 0.6599 (3.88); 0.6393 (4.28); 0.6036 (0.70); 0.0753 (0.99); -0.0002 (1.47)
I.13	CDCl <sub>3</sub>	300	8.0202 (1.64); 7.3723 (1.44); 7.3481 (1.82); 7.3446 (2.33); 7.3206 (2.22); 7.2639 (15.65); 7.2397 (2.56); 7.2355 (2.38); 7.2118 (1.59); 7.2076 (1.55); 7.0775 (1.43); 6.8957 (3.02); 6.7140 (1.50); 5.0291 (0.84); 4.9798 (2.71); 4.9388 (2.73); 4.8893 (0.85); 3.8264 (0.35); 3.8097 (11.04); 3.0089 (2.80); 2.9917 (3.85); 2.9589 (14.30); 2.9008 (2.33); 2.8861 (13.85); 2.8846 (13.40); 2.7271 (0.43); 2.6996 (0.58); 2.2361 (0.50); 1.6266 (16.00); 1.5965 (1.05); 1.2556 (0.78); 0.7193 (1.03); 0.7041 (3.13); 0.6836 (3.98); 0.6577 (1.58); 0.6456 (1.10); 0.6157 (0.54); 0.0697 (7.15); 0.0574 (0.33); 0.0106 (0.41); -0.0002 (14.06); -0.0111 (0.65)
I.14	CDCl <sub>3</sub>	300	9.6562 (0.33); 7.3044 (1.85); 7.2782 (3.38); 7.2656 (8.28); 7.2198 (3.02); 7.1936 (1.67); 7.1316 (1.26); 6.9496 (2.45); 6.7677 (1.24); 5.0765 (6.69); 5.0073 (0.53); 3.8092 (10.73); 3.0142 (2.35); 2.9967 (3.77); 2.9425 (2.06); 2.9251 (1.38); 2.6075 (0.33); 2.5720 (0.42); 2.4180 (1.02); 2.3970 (16.00); 2.3641 (1.36); 1.7807 (0.78); 1.6056 (15.61); 1.5581 (0.35); 1.3900 (0.65); 1.3667 (0.56); 1.2816 (0.44); 1.2561 (0.94); 0.6669 (0.67); 0.6571 (0.68); 0.6233 (0.56); 0.6045 (0.41); 0.5861 (0.33); 0.5408 (2.38); 0.5236 (2.44); 0.0712 (2.75); -0.0002 (3.97)
I.15	CDCl <sub>3</sub>	250	7.3280 (1.07); 7.3124 (1.30); 7.2915 (1.76); 7.2598 (7.72); 7.2102 (0.50); 7.1882 (2.43); 7.1713 (2.15); 7.1608 (2.09); 7.1519 (2.60); 7.1324 (0.54); 7.0785 (1.70); 7.0545 (2.42); 7.0433 (0.89); 6.8398 (3.18); 6.6252 (1.54); 4.3804 (0.68); 4.3378 (4.89); 4.3255 (4.56); 4.2830 (0.63); 3.9177 (1.30); 3.8577 (2.42); 3.8328 (10.27); 3.7082 (2.49); 3.6485 (1.45); 3.1092 (1.67); 3.0629 (1.93); 2.5253 (2.38); 2.4791 (2.05); 1.8085 (0.34); 1.7935 (0.70); 1.7828 (0.94); 1.7678 (1.37); 1.7537 (0.82); 1.7437 (0.78); 1.7275 (0.44); 1.5681 (4.39); 1.5472 (0.43); 1.3196 (16.00); 1.2657 (4.18); 1.1608 (0.91); 0.9066 (1.11); 0.8817 (3.12); 0.8540 (1.17); 0.5343 (0.35); 0.4988 (2.87); 0.4726 (3.06); 0.4576 (1.01); 0.4499 (1.34); 0.4268 (1.65); 0.4073 (1.95); 0.3912 (1.46); 0.3708 (0.70); 0.3566 (0.42); -0.0002 (4.85)
I.16	DMSO-d <sub>6</sub>	400	7.4708 (1.02); 7.4547 (1.24); 7.2664 (0.36); 7.2473 (1.76); 7.2382 (1.72); 7.2289 (2.06); 7.2101 (0.48); 7.1775 (2.13); 7.1638 (0.91); 7.0432 (1.60); 6.9097 (0.80); 4.4148 (1.22); 4.3886 (1.91); 4.3050 (1.85); 4.2787 (1.19); 3.9666 (9.65); 3.8887 (1.02); 3.8513 (1.55); 3.7216 (1.59); 3.6841 (1.06); 3.4025 (16.00); 3.3790 (0.49); 3.1161 (1.25); 3.0877 (1.38); 2.5663 (23.90); 2.5295 (1.44); 2.1408 (0.41); 1.8456 (0.80); 1.8380 (1.00); 1.8304 (0.82); 1.3586 (8.77); 0.5177 (1.24); 0.4187 (0.47); 0.3865 (0.94); 0.3485 (0.91); -0.0002 (0.86)
I.17	CDCl <sub>3</sub>	500	7.3443 (4.52); 7.3278 (6.10); 7.2738 (4.40); 7.2697 (5.22); 7.2587 (17.14); 7.2377 (3.82); 7.2334 (3.16); 7.2212 (2.85); 7.2168 (2.45); 6.9553 (1.71); 6.8460 (3.56); 6.7367 (1.82); 5.2943 (6.11); 4.7998 (4.63); 3.8134 (16.00); 3.5024 (6.35); 2.7635

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.74); 2.1940 (0.44); 2.1738 (0.49); 2.1666 (2.08); 2.1600 (0.47); 2.1519 (4.08); 2.1466 (4.76); 2.1340 (3.87); 2.1304 (3.26); 2.1240 (1.05); 2.1180 (2.12); 2.1072 (0.79); 2.1016 (2.28); 2.0901 (2.36); 2.0744 (2.49); 1.8134 (0.67); 1.8112 (0.67); 1.7966 (1.05); 1.7939 (1.30); 1.7903 (0.95); 1.7857 (0.67); 1.7833 (0.63); 1.7757 (1.27); 1.7734 (1.32); 1.7689 (1.09); 1.7663 (1.21); 1.7629 (0.74); 1.7486 (1.30); 1.7463 (1.25); 1.7383 (0.89); 1.7338 (0.91); 1.7248 (0.76); 1.7205 (0.98); 1.7135 (1.06); 1.7091 (1.23); 1.6968 (1.31); 1.6825 (0.67); 1.6785 (0.65); 1.6182 (0.51); 1.6016 (0.95); 1.5981 (0.79); 1.5941 (0.53); 1.5858 (0.78); 1.5821 (1.41); 1.5779 (1.17); 1.5739 (0.83); 1.5656 (0.89); 1.5616 (1.16); 1.5573 (1.20); 1.5533 (0.59); 1.5430 (8.41); 1.5367 (0.85); 1.5200 (0.37); 0.6541 (5.34); 0.6487 (5.42); 0.0063 (0.57); -0.0002 (15.35); -0.0068 (0.75)
I.20	DMSO-d <sub>6</sub>	400	7.5221 (5.61); 7.5185 (5.67); 7.5035 (6.73); 7.4995 (6.87); 7.4593 (0.74); 7.4402 (0.76); 7.3192 (2.23); 7.3048 (5.61); 7.3008 (5.60); 7.2865 (5.68); 7.2821 (5.28); 7.2766 (5.69); 7.2723 (6.35); 7.2577 (6.20); 7.2540 (6.75); 7.2393 (2.56); 7.2357 (2.36); 7.2054 (0.38); 7.1905 (0.38); 7.1380 (6.16); 7.1222 (9.73); 6.9877 (11.17); 6.8531 (5.54); 4.8923 (10.87); 4.0560 (1.12); 4.0382 (2.86); 4.0283 (3.67); 4.0201 (5.82); 4.0099 (10.47); 3.9943 (5.14); 3.9791 (1.56); 3.9750 (1.45); 3.9698 (1.18); 3.9601 (0.93); 3.9310 (0.67); 3.9145 (0.65); 3.8169 (16.00); 3.7060 (0.72); 3.6630 (8.67); 3.6480 (4.10); 3.6283 (1.23); 3.3064 (72.61); 3.2566 (0.45); 2.8540 (1.67); 2.6779 (0.51); 2.6737 (0.96); 2.6690 (1.46); 2.6644 (1.11); 2.5486 (0.51); 2.5225 (3.89); 2.5177 (6.53); 2.5091 (80.68); 2.5046 (169.46); 2.5000 (235.65); 2.4955 (173.85); 2.4910 (88.61); 2.4505 (3.63); 2.3783 (0.44); 2.3315 (1.16); 2.3269 (1.58); 2.3223 (1.22); 2.0273 (0.63); 2.0168 (0.52); 1.9876 (9.91); 1.6361 (13.66); 1.6128 (6.56); 1.2478 (0.72); 1.1923 (2.79); 1.1745 (5.68); 1.1567 (2.86); 0.8584 (1.05); 0.8406 (0.41); 0.6652 (6.77); 0.6516 (7.11); 0.5992 (0.59); 0.5151 (6.80); 0.3743 (0.39); 0.3693 (0.62); 0.3598 (0.48); 0.3531 (0.49); 0.2698 (0.65); 0.2613 (0.63); 0.2559 (0.46); -0.0002 (1.32)
I.21	DMSO-d <sub>6</sub>	400	7.3913 (1.90); 7.3716 (2.13); 7.3190 (3.44); 7.1251 (1.31); 7.1158 (2.35); 7.0920 (2.13); 7.0671 (1.33); 7.0473 (1.18); 7.0217 (3.19); 7.0019 (2.18); 6.9905 (2.68); 6.9816 (3.67); 6.9276 (2.27); 6.8558 (1.30); 6.8470 (1.80); 4.8438 (3.52); 4.0560 (0.64); 4.0383 (1.99); 4.0204 (2.40); 4.0117 (1.72); 3.9960 (5.42); 3.9803 (2.01); 3.9602 (0.40); 3.9555 (0.40); 3.8148 (6.80); 3.6618 (3.30); 3.6474 (2.73); 3.3062 (41.44); 2.8233 (0.77); 2.6736 (0.59); 2.6691 (0.78); 2.6645 (0.59); 2.5504 (0.36); 2.5225 (1.86); 2.5090 (47.29); 2.5046 (96.98); 2.5000 (131.41); 2.4956 (92.78); 2.4911 (43.51); 2.4549 (0.65); 2.4504 (0.71); 2.3314 (0.64); 2.3269 (0.89); 2.3224 (0.66); 2.2861 (16.00); 2.2557 (11.43); 2.0274 (0.50); 2.0168 (0.43); 1.9876 (7.78); 1.6151 (6.36); 1.5897 (1.75); 1.2793 (0.57); 1.2476 (2.78); 1.1923 (2.18); 1.1745 (4.31); 1.1567 (2.08); 0.8752 (1.22); 0.8585 (4.18); 0.8408 (1.63); 0.6518 (3.26); 0.6382 (3.36); 0.5077 (3.25); -0.0002 (0.67)
I.22	CDCl <sub>3</sub>	500	7.3358 (4.20); 7.3194 (5.93); 7.2616 (10.84); 7.2596 (6.13); 7.2419 (4.04); 7.2375 (3.00); 7.2255 (2.82); 7.2211 (2.33); 6.9601 (1.81); 6.8508 (3.79); 6.7415 (1.94); 5.2943 (10.15); 4.8338 (0.39); 4.7936 (3.01); 4.7628 (0.60); 3.8137 (16.00); 3.0179 (3.20); 2.7554 (0.75); 2.2411 (0.90); 2.2303 (1.16); 2.2242 (1.10); 2.2124 (1.89); 2.2007 (1.71); 2.1945 (1.56); 2.1835 (1.62); 2.0928 (1.00); 2.0818 (1.87); 2.0712 (1.26); 2.0633 (0.79); 2.0519 (1.34); 2.0416 (0.84); 2.0125 (0.32); 1.9972 (1.43); 1.9941 (1.51); 1.9911 (1.46); 1.9848 (2.61); 1.9811 (3.58); 1.9772 (3.08); 1.9692 (2.67); 1.9657 (2.65); 1.9602 (1.52); 1.9462 (0.37); 1.9433 (0.38); 1.9402 (0.34); 1.6163 (0.33); 1.6062 (0.43); 1.6042 (0.40); 1.5992 (0.57); 1.5952 (0.90); 1.5905 (1.04); 1.5853 (1.09); 1.5780 (4.70); 1.5732 (1.67); 1.5686 (1.86); 1.5635 (1.63); 1.5569 (1.90); 1.5517 (1.28); 1.5462 (1.59); 1.5415 (1.49); 1.5311 (1.08); 1.5198 (0.60); 1.5153 (0.52); 1.4897 (0.56); 1.4784 (1.02); 1.4747 (1.12); 1.4703 (0.76); 1.4640 (1.21); 1.4587 (1.15); 1.4529 (1.02); 1.4471 (1.11); 1.4391 (0.69); 1.4322 (0.55); 1.3492 (0.37); 1.3419 (0.43); 1.3348 (0.47); 1.3323 (0.49); 1.3275 (0.87); 1.3213 (0.85); 1.3153 (0.73); 1.3117 (1.00); 1.3074 (1.06); 1.3048 (0.95); 1.3017 (0.79); 1.2989 (0.86); 1.2960 (0.79); 1.2908 (0.82); 1.2852 (0.76); 1.2817 (0.62); 1.2786 (0.48); 1.2704 (0.32); 1.2684 (0.32); 1.2645 (0.44); 0.6493 (5.86); -0.0002 (6.87); -0.0068 (0.36)
I.23	CDCl <sub>3</sub>	300	7.3328 (4.05); 7.3055 (6.65); 7.2705 (4.68); 7.2279 (3.65); 7.2209 (4.11); 7.2006 (2.16); 7.1936 (2.68); 7.1375 (5.34); 7.1310 (4.67); 7.0394 (1.65); 6.8574 (3.40);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			6.6755 (1.71); 5.2999 (15.52); 4.9331 (1.17); 4.8801 (2.32); 4.7826 (3.34); 4.7298 (1.68); 3.8432 (16.00); 3.1759 (2.94); 2.8685 (1.33); 2.1698 (0.52); 2.1477 (1.20); 2.1283 (1.44); 2.0975 (2.47); 2.0781 (2.99); 2.0574 (1.98); 2.0466 (1.24); 2.0374 (1.80); 2.0267 (1.81); 2.0161 (1.10); 2.0056 (1.16); 1.9959 (0.55); 1.9862 (0.68); 1.9753 (0.73); 1.9591 (0.99); 1.9401 (1.47); 1.9188 (1.02); 1.8886 (0.98); 1.8684 (0.63); 1.6631 (2.13); 1.6505 (0.35); 1.6258 (0.72); 1.6057 (1.25); 1.5945 (1.43); 1.5853 (1.67); 1.5761 (1.88); 1.5581 (1.92); 1.5515 (1.81); 1.5396 (1.27); 1.5284 (1.63); 1.5072 (0.76); 1.4831 (0.73); 1.4617 (1.41); 1.4522 (1.39); 1.4285 (1.94); 1.4164 (1.69); 1.4070 (1.87); 1.3962 (1.41); 1.3833 (1.22); 1.3621 (0.79); 1.2573 (0.65); 0.7037 (3.05); 0.6803 (3.91); 0.6661 (4.05); 0.6487 (4.51); 0.6119 (0.81); -0.0002 (3.87)
I.24	CDCl <sub>3</sub>	300	7.3980 (2.51); 7.3919 (2.11); 7.3855 (2.08); 7.3770 (2.92); 7.3682 (3.86); 7.2968 (0.51); 7.2896 (1.05); 7.2730 (7.16); 7.2647 (6.80); 7.2531 (11.62); 7.2417 (7.55); 7.2342 (4.68); 7.2169 (1.32); 7.2098 (0.68); 7.1976 (0.44); 7.1879 (0.69); 7.1733 (4.14); 7.1641 (3.24); 7.1555 (1.93); 7.1489 (2.13); 7.1436 (2.62); 7.0610 (1.14); 6.8791 (2.25); 6.6970 (1.13); 5.2936 (13.04); 4.9947 (0.42); 4.9751 (0.99); 4.9233 (1.78); 4.8128 (3.08); 4.7610 (1.68); 3.8270 (16.00); 3.2064 (3.18); 2.8425 (1.50); 2.1895 (0.64); 2.1672 (1.18); 2.1346 (1.85); 2.1155 (2.49); 2.1014 (2.56); 2.0825 (3.19); 2.0590 (2.85); 2.0489 (2.73); 2.0383 (2.65); 2.0277 (1.68); 2.0171 (1.73); 2.0069 (0.99); 1.9969 (1.43); 1.9870 (1.68); 1.9748 (1.93); 1.9538 (1.29); 1.9222 (1.08); 1.9035 (0.74); 1.7509 (2.11); 1.6550 (0.41); 1.6316 (1.00); 1.6113 (1.76); 1.6021 (1.94); 1.5840 (2.83); 1.5634 (2.96); 1.5448 (1.92); 1.5368 (2.30); 1.5154 (1.12); 1.4920 (1.07); 1.4708 (1.93); 1.4615 (1.96); 1.4449 (2.55); 1.4378 (2.73); 1.4257 (2.37); 1.4161 (2.59); 1.4049 (2.01); 1.3926 (1.71); 1.3714 (1.08); 1.3483 (0.44); 1.2585 (0.43); 0.6493 (7.91); -0.0002 (2.89)
I.27	CDCl <sub>3</sub>	300	7.4932 (5.48); 7.4702 (4.81); 7.2697 (4.15); 7.0199 (5.00); 6.9876 (5.53); 6.9582 (1.33); 6.8409 (2.45); 6.6587 (1.23); 6.5122 (4.44); 5.2991 (0.77); 4.7636 (8.31); 3.8162 (16.00); 2.7208 (2.08); 1.6830 (1.71); 0.5678 (4.29); 0.5163 (6.67); -0.0002 (0.97)
I.28	CDCl <sub>3</sub>	300	7.4924 (7.69); 7.2825 (6.15); 7.2599 (20.54); 7.2297 (0.76); 7.0192 (1.23); 6.8372 (2.44); 6.6551 (1.25); 6.5200 (4.30); 5.3013 (4.72); 4.7627 (8.05); 3.8192 (16.00); 2.7002 (1.80); 1.5630 (6.77); 0.5669 (3.74); 0.5196 (6.03); -0.0002 (8.79)
I.29	CDCl <sub>3</sub>	300	7.4902 (9.88); 7.3108 (20.30); 7.2638 (2.84); 7.0368 (1.04); 6.8546 (2.04); 6.6723 (1.03); 6.5454 (3.69); 5.2966 (1.15); 4.8067 (6.27); 3.8020 (16.00); 2.6472 (1.77); 1.6586 (2.22); 0.5085 (7.78); -0.0002 (1.66)
I.30	DMSO-d <sub>6</sub>	400	7.9529 (1.55); 7.8218 (6.92); 7.7824 (0.34); 7.6573 (3.07); 7.3763 (6.74); 7.3676 (8.07); 7.3617 (8.99); 7.3535 (10.23); 7.3439 (4.22); 7.3333 (7.42); 7.3221 (4.35); 7.3102 (2.48); 7.1603 (1.35); 7.0256 (2.70); 6.9878 (0.33); 6.8909 (1.41); 6.8063 (3.94); 6.7691 (0.38); 6.7611 (0.33); 6.6430 (6.57); 4.8521 (4.29); 4.7661 (0.35); 4.6539 (0.34); 4.6408 (0.60); 3.8982 (16.00); 3.8612 (1.29); 3.8051 (0.49); 3.3368 (322.79); 3.3019 (0.89); 3.2842 (0.48); 3.2706 (0.32); 3.1759 (0.33); 2.8906 (9.89); 2.7598 (2.64); 2.7316 (9.67); 2.6715 (0.87); 2.5068 (89.44); 2.5025 (114.38); 2.4984 (86.83); 2.3293 (0.76); 1.2345 (0.74); 1.1953 (0.35); 1.1653 (0.35); 0.6977 (0.34); 0.6798 (0.39); 0.4496 (8.06); -0.0002 (3.32)
I.31	DMSO-d <sub>6</sub>	400	8.4144 (1.47); 7.9527 (0.91); 7.8039 (4.41); 7.8004 (4.21); 7.7890 (0.33); 7.6636 (2.16); 7.6572 (2.00); 7.6491 (1.45); 7.6452 (1.83); 7.6409 (2.46); 7.3671 (0.88); 7.3534 (2.69); 7.3481 (3.88); 7.3392 (4.35); 7.3302 (3.95); 7.3247 (2.59); 7.3112 (0.85); 7.3063 (0.58); 7.2834 (1.73); 7.2258 (2.25); 7.2100 (1.60); 7.2037 (1.51); 7.1475 (3.65); 7.0117 (1.78); 6.7758 (3.66); 6.7676 (4.05); 6.6399 (3.27); 6.6353 (3.48); 6.6315 (3.22); 6.6269 (2.86); 4.8275 (9.82); 3.9908 (0.66); 3.9355 (16.00); 3.8995 (0.45); 3.3731 (0.38); 3.3356 (146.74); 3.3020 (0.44); 2.9384 (1.13); 2.8902 (6.13); 2.7311 (5.18); 2.5067 (39.59); 2.5023 (49.63); 2.4979 (36.55); 2.3290 (0.32); 0.7218 (0.83); 0.7043 (3.39); 0.6911 (3.47); 0.6744 (1.08); 0.5520 (1.31); 0.5408 (3.63); 0.5345 (4.00); 0.5261 (3.21); 0.5130 (0.92); -0.0002 (1.64)
I.32	DMSO-d <sub>6</sub>	400	7.9528 (0.93); 7.7991 (7.19); 7.7959 (7.15); 7.6637 (2.43); 7.6569 (2.13); 7.6524 (2.20); 7.6414 (2.68); 7.3912 (0.71); 7.3857 (1.27); 7.3727 (4.93); 7.3694 (5.80);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			7.3599 (7.36); 7.3506 (6.13); 7.3469 (4.97); 7.3342 (1.29); 7.3284 (0.65); 7.2593 (3.77); 7.2456 (2.44); 7.2367 (2.41); 7.1026 (2.59); 6.9678 (5.75); 6.8332 (2.93); 6.7759 (3.19); 6.7690 (3.33); 6.6363 (6.21); 6.6317 (6.27); 6.6279 (5.58); 6.6233 (5.23); 4.8294 (11.80); 3.7992 (16.00); 3.3323 (210.84); 3.3100 (0.69); 2.8905 (7.12); 2.7316 (7.03); 2.6805 (0.40); 2.6758 (0.54); 2.6712 (0.65); 2.6668 (0.48); 2.5245 (1.81); 2.5112 (32.49); 2.5068 (64.04); 2.5022 (83.37); 2.4976 (59.61); 2.4932 (28.16); 2.3335 (0.38); 2.3290 (0.53); 2.3244 (0.39); 1.7212 (0.42); 1.6074 (0.41); 1.4365 (0.79); 1.2343 (0.48); 0.6130 (1.12); 0.5950 (4.67); 0.5816 (4.58); 0.5653 (1.69); 0.5442 (0.74); 0.5333 (0.60); 0.5272 (0.57); 0.5045 (2.09); 0.4939 (5.72); 0.4859 (5.70); 0.4652 (1.15); -0.0002 (6.91)
I.33	DMSO-d6	400	7.9526 (2.33); 7.8260 (6.60); 7.8005 (0.94); 7.7896 (0.79); 7.7687 (4.03); 7.7553 (4.33); 7.7231 (0.52); 7.6550 (3.22); 7.6475 (3.15); 7.5748 (0.46); 7.5673 (0.51); 7.5622 (0.47); 7.5544 (0.43); 7.4578 (3.07); 7.3794 (8.65); 7.3719 (8.58); 7.3654 (8.27); 7.3576 (6.81); 7.3191 (0.47); 7.3104 (0.36); 7.1089 (0.34); 7.0783 (3.16); 7.0671 (3.13); 6.9325 (0.91); 6.9190 (0.86); 6.8050 (3.94); 6.7706 (0.66); 6.7546 (0.33); 6.6491 (5.76); 4.8477 (8.80); 4.6387 (0.41); 3.3943 (0.43); 3.3366 (306.92); 2.9838 (3.05); 2.8899 (16.00); 2.8234 (3.06); 2.7305 (15.29); 2.7211 (2.84); 2.7104 (3.97); 2.7017 (2.73); 2.6829 (1.20); 2.6761 (0.89); 2.6714 (0.97); 2.5622 (0.39); 2.5480 (0.65); 2.5066 (96.99); 2.5022 (122.40); 2.4978 (89.06); 2.3289 (0.77); 1.2343 (0.79); 0.6874 (0.40); 0.6738 (0.45); 0.6438 (0.36); 0.6179 (0.43); 0.6045 (0.44); 0.5782 (0.35); 0.5557 (0.47); 0.5392 (0.50); 0.4013 (6.96); 0.3734 (5.43); -0.0002 (5.56)
I.34	DMSO-d6	400	7.9536 (1.65); 7.7823 (5.86); 7.7793 (5.92); 7.6779 (2.43); 7.6727 (2.97); 7.6571 (3.03); 7.6520 (4.31); 7.6352 (5.30); 7.5467 (5.58); 7.5260 (4.14); 7.1302 (2.29); 6.9955 (5.14); 6.9563 (4.78); 6.9481 (5.06); 6.8608 (2.58); 6.6302 (4.26); 6.6257 (4.62); 6.6217 (4.51); 6.6172 (4.07); 4.7318 (10.21); 3.8098 (16.00); 3.3340 (141.14); 2.8913 (11.15); 2.7322 (9.77); 2.7174 (0.72); 2.6946 (0.73); 2.6769 (0.81); 2.6724 (0.82); 2.6679 (0.70); 2.5077 (46.17); 2.5033 (58.77); 2.4989 (44.07); 2.3301 (0.39); 1.7221 (0.33); 1.4372 (0.70); 1.2332 (0.37); 0.6884 (0.85); 0.6699 (3.87); 0.6537 (4.27); 0.6416 (2.57); 0.6107 (4.92); 0.5823 (0.93); -0.0002 (4.97)
I.35	DMSO-d6	400	7.9535 (2.32); 7.7803 (6.76); 7.7779 (6.66); 7.6708 (4.51); 7.6541 (13.06); 7.5678 (0.43); 7.5475 (3.29); 7.5254 (2.48); 7.1778 (1.47); 7.0430 (2.92); 6.9413 (4.59); 6.9334 (4.69); 6.9087 (1.55); 6.6293 (5.81); 6.6248 (6.13); 6.6208 (5.84); 6.6163 (5.46); 4.7588 (1.35); 4.6483 (0.33); 3.9066 (12.32); 3.3320 (190.91); 2.8910 (16.00); 2.8004 (1.35); 2.7320 (13.63); 2.6762 (0.45); 2.6717 (0.61); 2.5073 (67.43); 2.5028 (86.60); 2.4984 (63.77); 2.3341 (0.46); 2.3298 (0.59); 2.3252 (0.45); 1.2333 (0.56); 0.5777 (6.98); -0.0002 (5.66)
I.36	DMSO-d6	400	8.4276 (1.77); 7.9532 (1.38); 7.7645 (3.96); 7.7611 (3.84); 7.6615 (1.74); 7.6560 (1.89); 7.6482 (0.34); 7.6407 (2.36); 7.6352 (2.66); 7.5720 (3.67); 7.5668 (3.31); 7.5399 (0.43); 7.5335 (5.22); 7.5190 (0.36); 7.5126 (3.88); 7.3043 (1.44); 7.1684 (3.40); 7.0325 (1.62); 6.9587 (3.42); 6.9509 (3.54); 6.6157 (3.18); 6.6112 (3.22); 6.6072 (3.10); 6.6027 (2.96); 4.7373 (8.51); 4.0013 (0.61); 3.9344 (16.00); 3.9006 (0.38); 3.3738 (0.33); 3.3368 (143.64); 2.8909 (11.55); 2.8728 (0.75); 2.7316 (9.04); 2.5253 (0.77); 2.5121 (16.90); 2.5076 (33.75); 2.5030 (44.35); 2.4985 (31.73); 2.4940 (15.01); 0.7702 (0.63); 0.7522 (2.62); 0.7392 (2.67); 0.7348 (2.31); 0.7222 (0.91); 0.6846 (0.43); 0.6456 (1.20); 0.6351 (2.97); 0.6278 (2.99); 0.6063 (0.67); -0.0002 (2.24)
I.37	DMSO-d6	400	7.9592 (0.78); 7.8815 (2.08); 7.7731 (3.07); 7.2909 (1.20); 7.2719 (1.54); 7.2447 (0.32); 7.1707 (0.76); 7.1416 (4.44); 7.1218 (1.88); 7.0367 (1.25); 6.9011 (0.64); 6.7676 (2.13); 4.7149 (2.08); 3.9076 (7.06); 3.3326 (47.29); 2.8969 (4.71); 2.7375 (5.21); 2.6773 (0.37); 2.5085 (50.34); 2.3777 (1.03); 2.3351 (1.00); 2.3153 (16.00); 0.4651 (1.98); 0.4508 (2.22); 0.4053 (2.64)
I.38	DMSO-d6	400	8.3504 (0.94); 7.9597 (0.79); 7.8428 (3.83); 7.7535 (3.03); 7.7492 (5.11); 7.7452 (2.96); 7.2828 (1.37); 7.2733 (3.24); 7.2539 (4.06); 7.1468 (2.92); 7.1229 (2.07); 7.1034 (1.61); 7.0120 (4.08); 6.7395 (3.33); 6.7374 (3.33); 4.6908 (7.80); 3.9348 (13.25); 3.3316 (22.87); 2.8970 (6.05); 2.8540 (0.79); 2.7377 (5.02); 2.5307 (0.91);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			2.5174 (15.50); 2.5131 (30.80); 2.5086 (40.58); 2.5040 (29.77); 2.4996 (14.82); 2.3353 (0.33); 2.3031 (16.00); 0.6744 (0.67); 0.6564 (2.86); 0.6433 (2.95); 0.6390 (2.48); 0.6262 (0.88); 0.4949 (0.98); 0.4838 (2.86); 0.4773 (3.06); 0.4687 (2.54); 0.4557 (0.72)
I.39	DMSO-d <sub>6</sub>	600	7.9522 (0.59); 7.8404 (0.87); 7.7522 (2.62); 7.7495 (4.43); 7.7467 (2.51); 7.2734 (0.89); 7.2608 (1.05); 7.1318 (1.73); 7.1189 (1.42); 7.0517 (0.72); 7.0436 (3.13); 6.9623 (1.29); 6.8727 (0.66); 6.7305 (0.99); 4.6857 (2.42); 3.8003 (3.97); 3.3149 (19.29); 2.8900 (4.98); 2.7311 (4.13); 2.7302 (4.07); 2.6920 (0.39); 2.6154 (0.33); 2.6124 (0.44); 2.5217 (0.82); 2.5186 (1.09); 2.5154 (1.24); 2.5067 (23.05); 2.5037 (49.31); 2.5006 (67.72); 2.4976 (47.55); 2.4945 (21.45); 2.3848 (0.40); 2.3039 (16.00); 0.5619 (1.59); 0.5527 (1.59); 0.4442 (0.78); 0.4367 (2.36); 0.4325 (2.51); 0.4269 (2.11); 0.4184 (0.56)
I.40	DMSO-d <sub>6</sub>	400	7.9534 (1.10); 7.7587 (3.19); 7.7577 (3.20); 7.7545 (3.30); 7.6453 (1.46); 7.6399 (1.60); 7.6245 (1.98); 7.6191 (2.30); 7.5448 (3.15); 7.5397 (2.87); 7.5203 (4.52); 7.4994 (5.22); 7.2597 (1.22); 7.1221 (2.68); 7.1171 (5.07); 6.9734 (1.32); 6.9322 (2.98); 6.9238 (3.15); 6.6092 (2.66); 6.6047 (2.67); 6.6007 (2.57); 6.5962 (2.49); 4.7241 (7.31); 3.7270 (0.35); 3.6925 (16.00); 3.3248 (12.54); 2.8905 (9.30); 2.7317 (7.15); 2.5243 (0.85); 2.5111 (14.03); 2.5066 (27.83); 2.5020 (36.72); 2.4975 (26.80); 2.4930 (13.33); 0.7771 (0.56); 0.7590 (2.35); 0.7461 (2.42); 0.7417 (2.09); 0.7290 (0.80); 0.6332 (0.91); 0.6225 (2.49); 0.6154 (2.66); 0.6074 (2.04); 0.5938 (0.63)
I.41	DMSO-d <sub>6</sub>	400	7.9592 (2.60); 7.7994 (11.48); 7.7963 (11.38); 7.7784 (4.19); 7.6995 (6.26); 7.6797 (4.98); 7.6589 (5.62); 7.5487 (3.37); 7.5293 (2.80); 7.1065 (2.57); 6.9955 (5.74); 6.6199 (6.70); 6.6155 (7.43); 6.6117 (6.91); 6.6073 (5.50); 4.7622 (5.83); 3.3318 (20.28); 2.8959 (16.00); 2.7374 (15.55); 2.6815 (0.52); 2.6770 (0.58); 2.5121 (55.09); 2.5078 (64.25); 2.5035 (48.09); 2.3349 (0.43); 2.3304 (0.34); 1.2372 (0.48); 0.7800 (0.47); 0.7663 (0.47); 0.7217 (0.37); 0.5020 (5.95)
I.42	DMSO-d <sub>6</sub>	400	9.7884 (0.51); 7.9529 (0.84); 7.8116 (3.53); 7.8105 (3.57); 7.8072 (3.42); 7.6646 (1.74); 7.6581 (1.32); 7.6553 (1.35); 7.6467 (1.26); 7.6418 (2.10); 7.5799 (0.32); 7.5748 (0.33); 7.4970 (1.94); 7.3539 (0.73); 7.3406 (2.30); 7.3372 (2.92); 7.3277 (3.55); 7.3189 (2.90); 7.3145 (2.35); 7.3017 (0.61); 7.2398 (1.33); 7.2143 (1.70); 7.2004 (1.22); 7.1917 (1.15); 7.1165 (2.55); 7.1123 (2.49); 7.0969 (2.91); 6.9537 (1.40); 6.7750 (3.05); 6.7666 (3.54); 6.6433 (2.98); 6.6387 (3.06); 6.6348 (2.70); 6.6303 (2.50); 4.8167 (7.71); 3.7163 (1.38); 3.6911 (16.00); 3.3253 (17.96); 2.9709 (0.61); 2.9632 (0.81); 2.9540 (1.09); 2.9450 (0.82); 2.9372 (0.61); 2.8898 (6.42); 2.7313 (5.25); 2.7305 (5.13); 2.5237 (0.82); 2.5104 (13.53); 2.5060 (26.66); 2.5014 (34.92); 2.4969 (25.36); 2.4924 (12.35); 0.7307 (0.63); 0.7127 (2.55); 0.6998 (2.63); 0.6954 (2.21); 0.6827 (0.78); 0.5516 (0.95); 0.5407 (2.62); 0.5340 (2.89); 0.5256 (2.41); 0.5125 (0.84); 0.4819 (0.35)
I.43	CDCl <sub>3</sub>	300	7.4607 (4.94); 7.4549 (5.19); 7.4495 (2.92); 7.4417 (2.23); 7.2625 (9.72); 7.2259 (2.46); 7.2142 (0.69); 7.2060 (1.61); 7.1985 (4.18); 7.1139 (3.99); 7.0956 (1.68); 7.0236 (2.47); 6.8412 (1.25); 6.5191 (2.34); 5.2991 (9.18); 4.7552 (4.52); 3.9478 (11.42); 2.3625 (16.00); 2.1716 (0.82); 2.1603 (0.93); 2.1497 (0.95); 2.0451 (0.34); 1.6018 (7.96); 0.8443 (0.59); 0.8244 (0.70); 0.8133 (0.61); 0.7939 (0.44); 0.7008 (2.36); 0.6675 (1.54); 0.6536 (1.26); 0.6488 (1.26); 0.6355 (1.36); 0.6224 (0.82); 0.6175 (0.81); 0.6041 (0.64); 0.3109 (0.80); 0.2902 (0.78); -0.0002 (5.69)
I.44	CDCl <sub>3</sub>	300	7.4858 (2.46); 7.4803 (4.56); 7.4748 (3.16); 7.4482 (0.75); 7.2611 (12.44); 7.2077 (1.08); 7.1814 (2.45); 7.1117 (2.00); 7.0860 (1.27); 6.8891 (0.33); 6.5069 (0.66); 5.2994 (7.60); 4.8632 (0.34); 4.7786 (0.80); 4.7277 (0.53); 3.7973 (5.56); 2.6829 (0.53); 2.3540 (16.00); 1.5692 (6.99); 0.8327 (1.76); 0.7472 (0.33); 0.6752 (0.55); 0.1044 (0.74); 0.0665 (0.34); 0.0107 (0.38); -0.0002 (11.41); -0.0110 (0.52)
I.45	CDCl <sub>3</sub>	300	7.4905 (2.73); 7.3640 (1.25); 7.3300 (0.99); 7.2632 (2.82); 7.2263 (0.66); 7.2005 (1.09); 7.1492 (1.39); 7.1143 (1.09); 7.0934 (1.33); 6.9376 (0.38); 6.7554 (0.75); 6.6993 (0.54); 6.5730 (0.40); 6.5314 (1.31); 6.3849 (0.89); 5.2938 (1.84); 5.2547 (0.46); 4.6980 (0.35); 4.6451 (0.65); 4.5258 (0.97); 4.4862 (0.52); 4.4713 (0.48); 3.8879 (4.48); 3.8459 (3.30); 3.8138 (0.34); 2.6880 (0.46); 2.6656 (1.04); 2.6484

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(1.06); 2.6268 (0.52); 2.3536 (16.00); 1.6832 (2.16); 1.1595 (0.33); 1.1393 (0.33); 1.0710 (2.02); 1.0526 (1.50); 0.8556 (2.02); 0.8364 (2.56); 0.7800 (0.52); 0.7573 (0.61); 0.7328 (0.55); 0.7095 (0.49); 0.6880 (0.39); 0.5930 (0.54); 0.5824 (0.46); 0.5713 (0.52); 0.1711 (0.46); 0.1540 (0.47); 0.0337 (0.62); 0.0164 (0.60); -0.0002 (2.58)
I.46	CDCl <sub>3</sub>	300	7.5753 (1.96); 7.5695 (3.07); 7.5623 (2.27); 7.5552 (0.87); 7.5403 (0.96); 7.5348 (2.26); 7.5277 (1.64); 7.4819 (2.68); 7.4797 (2.86); 7.4760 (2.81); 7.4738 (2.61); 7.4112 (3.19); 7.3837 (2.25); 7.3612 (1.38); 7.2720 (0.37); 7.2615 (27.91); 7.2524 (0.55); 7.2509 (0.45); 7.2493 (0.37); 7.2479 (0.33); 7.1819 (0.67); 7.1536 (1.10); 6.9721 (2.28); 6.7908 (1.14); 6.6740 (2.39); 6.6718 (2.46); 6.6627 (2.77); 6.6605 (2.72); 6.4918 (2.57); 6.4858 (2.64); 6.4805 (2.44); 6.4745 (2.31); 4.8455 (4.35); 2.7974 (5.96); 2.7749 (16.00); 2.7124 (0.45); 2.6942 (0.95); 2.6761 (1.36); 2.6687 (0.65); 2.6584 (0.89); 2.6395 (0.47); 2.0097 (4.09); 0.7368 (4.62); 0.7195 (8.28); 0.0106 (0.54); -0.0002 (23.95); -0.0093 (0.92); -0.0110 (1.18)
I.47	CDCl <sub>3</sub>	300	7.2620 (7.09); 7.0187 (0.93); 6.9619 (3.68); 6.9327 (4.11); 6.8382 (1.86); 6.6242 (7.90); 6.1853 (3.06); 5.3001 (7.65); 4.8504 (5.78); 3.8168 (13.12); 3.6935 (16.00); 2.7254 (1.87); 1.5884 (5.02); 1.2406 (1.95); 0.5017 (6.90); -0.0002 (3.57)
I.48	CDCl <sub>3</sub>	300	7.2806 (1.73); 7.2611 (10.26); 7.2200 (7.92); 7.0141 (0.88); 6.8328 (1.79); 6.6366 (6.18); 6.2008 (2.75); 5.3003 (3.49); 4.8554 (5.28); 3.8196 (12.74); 3.6922 (16.00); 2.7058 (1.62); 1.9266 (1.95); 1.5805 (4.09); 1.2407 (12.75); 0.5003 (5.97); -0.0002 (4.97)
I.49	CDCl <sub>3</sub>	300	7.3337 (2.15); 7.2573 (6.93); 7.2414 (8.58); 7.0260 (0.61); 6.8434 (1.23); 6.6582 (2.86); 6.6308 (3.96); 6.2308 (1.92); 5.2972 (5.61); 4.9009 (3.45); 3.8046 (10.69); 3.6919 (16.00); 2.6623 (1.35); 1.6042 (2.29); 1.2384 (2.09); 0.4866 (6.23); -0.0002 (2.72)
I.50	CDCl <sub>3</sub>	300	7.4766 (0.93); 7.4705 (1.08); 7.4498 (1.34); 7.4438 (1.63); 7.4010 (1.79); 7.3371 (2.35); 7.3100 (1.79); 7.3057 (2.31); 7.1544 (0.55); 6.9724 (1.13); 6.9348 (1.51); 6.9283 (2.65); 6.9218 (1.64); 6.7904 (0.57); 6.6665 (1.44); 6.6583 (2.45); 6.6503 (1.53); 6.4592 (1.80); 6.4530 (2.12); 6.4504 (2.06); 6.4442 (1.67); 4.8320 (2.11); 3.8437 (6.85); 3.7227 (16.00); 3.2078 (0.33); 3.1870 (0.43); 3.1646 (0.36); 2.7044 (0.49); 2.0338 (5.88); 1.2943 (10.92); 1.2717 (10.82); 0.7198 (3.38); 0.7064 (2.90); 0.0506 (1.07)
I.51	DMSO-d <sub>6</sub>	400	7.9598 (0.68); 7.2267 (0.93); 7.2081 (1.13); 7.1053 (0.73); 7.0547 (1.52); 7.0354 (1.23); 6.9706 (1.55); 6.9489 (2.70); 6.8736 (0.99); 6.8357 (0.84); 6.7668 (1.70); 6.7611 (2.60); 6.7553 (1.52); 6.1728 (1.05); 4.7472 (2.68); 3.8167 (4.28); 3.6595 (16.00); 3.6403 (0.43); 3.3309 (23.39); 2.8970 (4.75); 2.7381 (4.46); 2.5167 (17.18); 2.5126 (31.70); 2.5081 (39.45); 2.5035 (28.34); 2.4992 (13.88); 2.2755 (11.85); 0.5608 (1.60); 0.5468 (1.64); 0.4286 (0.73); 0.4178 (2.18); 0.4114 (2.33); 0.3906 (0.53)
I.52	DMSO-d <sub>6</sub>	400	8.3346 (0.47); 7.9596 (0.90); 7.2902 (0.90); 7.2132 (2.20); 7.1938 (2.75); 7.1541 (2.05); 7.0385 (1.44); 7.0184 (2.10); 6.9134 (2.42); 6.8573 (2.34); 6.7560 (1.74); 6.7500 (2.76); 6.7442 (1.60); 6.1709 (1.53); 6.1657 (2.06); 6.1604 (1.42); 4.7456 (4.91); 3.9378 (7.99); 3.6419 (16.00); 3.3324 (32.83); 2.8969 (6.57); 2.8633 (0.62); 2.7381 (5.28); 2.7374 (5.37); 2.5172 (14.76); 2.5129 (28.15); 2.5084 (35.99); 2.5038 (26.12); 2.4994 (12.87); 2.2682 (10.96); 0.6548 (0.49); 0.6370 (2.08); 0.6237 (2.16); 0.6196 (1.82); 0.6065 (0.62); 0.4649 (0.70); 0.4538 (2.08); 0.4474 (2.23); 0.4389 (1.84); 0.4258 (0.52)
I.53	DMSO-d <sub>6</sub>	400	7.9595 (1.68); 7.7725 (0.43); 7.2291 (1.56); 7.2105 (1.80); 7.1624 (0.93); 7.1417 (0.78); 7.1217 (0.46); 7.0563 (4.82); 7.0314 (2.53); 6.8945 (2.94); 6.7735 (2.79); 6.1892 (2.11); 4.7671 (2.84); 4.7173 (0.41); 3.9130 (8.80); 3.8318 (0.41); 3.8127 (0.37); 3.6704 (13.18); 3.6402 (0.48); 3.3325 (46.22); 2.8967 (11.92); 2.7544 (1.01); 2.7377 (11.11); 2.6817 (0.33); 2.6771 (0.41); 2.6728 (0.32); 2.5168 (24.68); 2.5126 (47.42); 2.5082 (60.91); 2.5037 (43.99); 2.3776 (0.50); 2.3348 (0.66); 2.3301 (0.52); 2.3155 (2.55); 2.2810 (16.00); 0.4445 (2.49); 0.4277 (2.49); 0.3809

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(2.87)
I.54	CDCl <sub>3</sub>	300	7.6745 (1.76); 7.6429 (1.33); 7.5584 (4.92); 7.4581 (5.63); 7.2966 (18.31); 7.2675 (10.51); 7.0390 (1.12); 6.8556 (2.06); 6.6727 (1.05); 4.8130 (6.98); 4.1334 (0.60); 4.1127 (0.64); 3.9414 (16.00); 3.8061 (14.90); 2.6403 (2.25); 2.0501 (2.35); 1.6214 (4.80); 1.2601 (1.50); 0.5022 (9.16); -0.0002 (6.25)
I.55	CDCl <sub>3</sub>	300	7.7048 (0.91); 7.6752 (1.24); 7.6417 (1.13); 7.5366 (4.62); 7.4898 (1.44); 7.4493 (4.61); 7.2590 (19.14); 7.2194 (0.72); 7.0180 (1.05); 6.8371 (2.17); 6.6551 (1.09); 4.7678 (7.25); 4.1339 (0.64); 4.1107 (0.67); 3.9425 (16.00); 3.8193 (13.91); 2.6972 (1.65); 2.0479 (2.61); 1.6036 (8.38); 1.2851 (0.74); 1.2571 (1.73); 1.2360 (0.83); 0.5630 (3.29); 0.5130 (5.17); -0.0002 (7.14)
I.56	CDCl <sub>3</sub>	300	8.0938 (0.89); 7.5302 (4.33); 7.4325 (4.41); 7.2686 (5.06); 7.2406 (2.19); 7.0067 (4.47); 6.9751 (4.53); 6.8368 (2.36); 6.6547 (1.19); 5.3020 (0.40); 4.7615 (8.13); 3.9491 (16.00); 3.8180 (15.06); 2.7138 (2.22); 2.0099 (0.50); 0.5673 (4.40); 0.5114 (6.58); -0.0002 (2.02)
I.57	DMSO-d <sub>6</sub>	400	7.9592 (1.54); 7.9397 (2.72); 7.6259 (2.70); 7.2784 (1.29); 7.2581 (1.58); 7.1739 (0.74); 7.1066 (5.06); 7.0903 (2.14); 7.0391 (1.46); 6.9047 (0.74); 4.7208 (2.45); 3.9114 (8.19); 3.8914 (15.60); 3.3368 (19.88); 2.8966 (8.83); 2.7514 (1.17); 2.7377 (8.26); 2.6772 (0.39); 2.5123 (44.29); 2.5081 (55.48); 2.5037 (40.81); 2.3350 (0.44); 2.3302 (0.38); 2.3007 (16.00); 0.4586 (2.12); 0.4445 (2.35); 0.4025 (2.87)
I.58	DMSO-d <sub>6</sub>	400	7.9596 (0.70); 7.9083 (1.19); 7.5981 (1.39); 7.2713 (0.93); 7.2522 (1.15); 7.1092 (2.22); 7.0906 (1.23); 7.0043 (2.60); 6.9737 (1.61); 6.8390 (0.80); 4.7029 (2.98); 3.8771 (16.00); 3.8133 (4.91); 3.3331 (44.12); 2.8971 (4.76); 2.7378 (4.50); 2.7225 (0.51); 2.6823 (0.32); 2.6777 (0.34); 2.5128 (26.97); 2.5085 (34.60); 2.5041 (25.58); 2.2964 (11.57); 0.5685 (1.64); 0.5543 (1.67); 0.4509 (0.69); 0.4399 (2.11); 0.4332 (2.26); 0.4120 (0.48)
I.59	DMSO-d <sub>6</sub>	600	7.9524 (0.83); 7.8815 (2.48); 7.6352 (0.54); 7.6330 (0.36); 7.6296 (0.52); 7.6266 (0.73); 7.6238 (0.78); 7.6218 (0.82); 7.6178 (0.74); 7.6155 (0.78); 7.6041 (0.84); 7.6017 (0.55); 7.5817 (2.83); 7.5676 (0.62); 7.5626 (0.63); 7.5599 (0.38); 7.5552 (0.64); 7.5505 (0.70); 7.2434 (1.95); 7.2306 (2.55); 7.1424 (1.27); 7.0817 (1.13); 7.0689 (0.98); 7.0520 (0.62); 6.9673 (1.95); 4.6923 (4.10); 3.9296 (6.05); 3.8480 (16.00); 3.3159 (15.89); 2.8898 (7.02); 2.8506 (0.37); 2.7307 (5.46); 2.5217 (0.63); 2.5186 (0.86); 2.5154 (1.02); 2.5067 (16.87); 2.5037 (35.76); 2.5007 (48.66); 2.4976 (34.17); 2.4946 (15.40); 2.3849 (0.33); 2.2825 (9.85); 0.6517 (0.41); 0.6422 (1.45); 0.6397 (1.78); 0.6311 (1.77); 0.6281 (1.54); 0.6195 (0.49); 0.4764 (0.59); 0.4687 (1.64); 0.4649 (1.75); 0.4620 (1.61); 0.4588 (1.57); 0.4502 (0.44)
I.60	CDCl <sub>3</sub>	300	7.5337 (1.57); 7.4215 (1.57); 7.2722 (2.24); 7.2170 (1.17); 7.2031 (0.50); 7.1896 (1.92); 7.1076 (2.05); 7.0999 (2.48); 7.0128 (0.67); 6.8307 (1.35); 6.6488 (0.68); 5.2957 (1.72); 4.8672 (0.37); 4.8174 (0.62); 4.6927 (1.16); 4.6419 (0.68); 3.9256 (16.00); 3.8060 (6.71); 2.3600 (11.69); 2.2974 (0.62); 2.2865 (0.82); 2.2753 (0.84); 2.2640 (0.66); 1.8647 (1.15); 1.2588 (4.11); 1.2530 (1.40); 1.2356 (1.14); 0.8002 (0.50); 0.6674 (1.71); 0.6233 (0.97); 0.6069 (0.86); 0.5919 (0.60); 0.5759 (0.42); 0.2660 (0.49); 0.2481 (0.47); -0.0002 (1.35)
I.61	CDCl <sub>3</sub>	300	7.5576 (1.59); 7.4358 (1.64); 7.2659 (4.42); 7.2179 (1.91); 7.1936 (1.37); 7.1044 (1.18); 7.0784 (0.76); 6.9265 (0.40); 6.7441 (0.79); 6.5618 (0.40); 5.2983 (4.79); 4.6681 (0.35); 3.9578 (16.00); 3.9011 (5.13); 2.3585 (10.72); 2.3287 (0.63); 1.7048 (1.76); 1.2650 (5.39); 1.2531 (1.80); 1.2387 (1.91); 0.7833 (0.35); 0.7631 (0.40); 0.7524 (0.35); 0.5932 (0.40); 0.5756 (0.64); 0.5497 (2.94); 0.5299 (2.46); 0.1758 (0.58); 0.1544 (0.55); -0.0002 (2.61)
I.62	CDCl <sub>3</sub>	300	7.5407 (3.81); 7.4385 (3.22); 7.3336 (3.58); 7.2732 (3.14); 7.2307 (2.30); 7.2137 (1.65); 7.2055 (3.76); 7.1243 (2.19); 7.0961 (4.67); 7.0317 (2.68); 6.8495 (1.32); 5.2975 (3.55); 4.8326 (0.46); 4.7815 (1.71); 4.7496 (2.46); 4.6988 (0.64); 3.9383 (12.23); 3.9089 (0.72); 3.8259 (13.05); 2.3647 (16.00); 2.0556 (0.76); 2.0438

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.73); 1.8653 (2.78); 1.2582 (1.03); 0.8473 (0.51); 0.8366 (0.65); 0.8261 (0.67); 0.8166 (0.76); 0.8057 (0.64); 0.7870 (0.45); 0.6684 (2.51); 0.6546 (2.85); 0.6424 (2.26); 0.6231 (1.65); 0.6099 (0.99); 0.5918 (0.70); 0.2912 (0.85); 0.2717 (0.81); -0.0002 (1.92)
I.63	CDCl <sub>3</sub>	300	7.5125 (0.75); 7.4094 (0.77); 7.2634 (8.70); 7.1947 (0.85); 7.1552 (2.21); 7.1048 (2.07); 7.0794 (1.33); 6.8911 (0.37); 5.2994 (12.70); 4.8164 (1.32); 3.9354 (12.74); 3.7981 (5.86); 2.6943 (0.64); 2.3519 (16.00); 1.6317 (5.20); 1.2671 (0.86); 1.2411 (0.65); 0.8280 (1.98); 0.6640 (0.62); 0.0968 (0.87); -0.0002 (8.08); -0.0111 (0.39)
I.64	CDCl <sub>3</sub>	400	7.5469 (0.79); 7.4345 (0.85); 7.3921 (0.60); 7.3401 (0.74); 7.2636 (4.23); 7.2113 (0.42); 7.1922 (0.59); 7.1514 (0.67); 7.1191 (0.49); 7.1033 (0.62); 7.0819 (0.75); 6.7513 (0.42); 5.2966 (3.90); 4.6849 (0.35); 4.5078 (0.39); 3.9578 (16.00); 3.8891 (2.58); 3.8437 (1.87); 2.6800 (0.69); 2.3497 (9.85); 2.0425 (0.35); 1.6592 (5.01); 1.2651 (1.72); 1.2579 (0.57); 1.2394 (0.87); 1.0615 (1.14); 1.0480 (0.98); 0.8446 (1.28); 0.8308 (1.47); 0.7428 (0.37); 0.7251 (0.35); 0.5880 (0.33); 0.0256 (0.39); 0.0085 (0.45); -0.0002 (3.79)
I.65	CDCl <sub>3</sub>	300	7.7260 (3.58); 7.7237 (3.55); 7.5794 (3.45); 7.4016 (0.74); 7.3958 (0.80); 7.3748 (1.45); 7.3689 (1.67); 7.3242 (2.70); 7.3093 (1.87); 7.3040 (1.78); 7.2980 (1.73); 7.2643 (4.18); 7.0832 (0.77); 6.9012 (1.62); 6.7191 (0.81); 4.8002 (2.31); 3.9360 (16.00); 3.8117 (6.34); 3.1835 (0.34); 3.1612 (0.45); 3.1387 (0.36); 2.7090 (0.41); 2.0033 (0.61); 1.2607 (10.84); 1.2381 (10.80); 0.6769 (3.55); 0.6602 (3.32); -0.0002 (3.42)
I.66	CDCl <sub>3</sub>	300	7.4033 (3.57); 7.3954 (3.80); 7.3097 (5.59); 7.2623 (16.39); 7.2217 (5.17); 7.1115 (3.53); 7.0120 (1.24); 6.8290 (2.45); 6.6482 (1.22); 5.3011 (4.07); 4.7248 (8.52); 3.8115 (16.00); 2.6399 (2.25); 1.5785 (5.06); 1.2585 (0.51); 0.5233 (4.27); 0.4602 (6.72); -0.0002 (6.14)
I.67	CDCl <sub>3</sub>	300	7.3900 (3.64); 7.3806 (3.93); 7.3228 (12.34); 7.2639 (3.10); 7.2220 (4.01); 7.1392 (3.20); 7.0292 (1.12); 6.8466 (2.12); 6.6634 (1.07); 5.2970 (5.52); 4.7695 (6.91); 3.7949 (16.00); 2.5871 (2.26); 1.6282 (2.40); 0.4467 (7.29); -0.0002 (1.88)
I.68	CDCl <sub>3</sub>	300	7.3743 (3.85); 7.3175 (9.17); 7.2850 (4.09); 7.2618 (4.10); 7.0994 (3.52); 7.0890 (3.50); 7.0201 (4.90); 6.8324 (2.40); 6.6502 (1.20); 5.2989 (5.34); 4.8069 (8.06); 3.8126 (16.00); 2.6760 (2.18); 1.6250 (2.12); 0.5455 (4.01); 0.4782 (6.22); -0.0002 (1.74)
I.69	CDCl <sub>3</sub>	300	7.3903 (3.75); 7.3456 (12.06); 7.3034 (2.52); 7.2621 (4.41); 7.0965 (3.78); 7.0867 (3.84); 7.0300 (4.61); 6.8481 (2.07); 6.6664 (1.03); 5.2987 (1.92); 4.8494 (6.62); 3.7979 (16.00); 2.6254 (2.26); 1.6041 (3.51); 0.4654 (7.08); -0.0002 (2.79)
I.70	CDCl <sub>3</sub>	300	7.4954 (3.62); 7.3731 (3.29); 7.3657 (3.59); 7.3169 (20.53); 7.2625 (8.06); 7.2236 (4.29); 7.1413 (3.12); 7.0731 (2.33); 6.8910 (1.14); 5.3005 (5.99); 4.7858 (8.43); 3.9388 (16.00); 2.5001 (2.00); 1.5998 (8.32); 0.5433 (4.36); 0.4959 (6.42); -0.0002 (4.30)
I.71	CDCl <sub>3</sub>	300	7.5855 (3.26); 7.3485 (4.53); 7.3217 (4.04); 7.2893 (9.38); 7.2629 (13.29); 7.0874 (3.11); 7.0705 (4.21); 7.0214 (3.95); 6.8844 (1.19); 4.8145 (9.12); 3.9614 (16.00); 2.5857 (1.97); 1.5815 (15.13); 0.6250 (3.55); 0.6040 (3.76); 0.5274 (5.12); 0.0696 (0.36); -0.0002 (7.00)
I.72	CDCl <sub>3</sub>	300	7.5359 (2.63); 7.4123 (2.09); 7.3907 (3.20); 7.3237 (14.14); 7.2659 (3.18); 7.0792 (5.49); 7.0283 (4.17); 6.9019 (1.24); 5.2937 (0.85); 4.8514 (8.86); 3.9300 (16.00); 2.5461 (2.02); 1.7337 (2.36); 0.5719 (4.31); 0.5162 (6.14); -0.0002 (0.91)
I.73	CDCl <sub>3</sub>	400	7.4187 (3.07); 7.4112 (3.75); 7.4064 (3.74); 7.3990 (3.89); 7.3844 (0.43); 7.3595 (1.18); 7.3562 (1.26); 7.3550 (1.25); 7.3435 (5.73); 7.3375 (18.38); 7.3203 (1.36); 7.3136 (1.03); 7.3004 (1.01); 7.2958 (1.01); 7.2595 (5.82); 7.2560 (3.76); 7.2528 (4.00); 7.2486 (3.50); 7.2454 (3.36); 7.2221 (0.41); 7.2036 (0.39); 7.1699 (3.89); 7.1667 (3.84); 7.1619 (2.95); 7.1577 (3.77); 7.1545 (3.34); 7.0936 (0.47); 7.0892

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.44); 7.0247 (4.96); 6.9869 (0.42); 6.9751 (0.41); 6.8875 (2.52); 6.6408 (0.44); 5.2918 (7.26); 4.8063 (0.39); 3.7821 (16.00); 3.7052 (2.12); 2.8187 (0.36); 2.8120 (0.64); 2.8068 (1.03); 2.8011 (1.01); 2.7955 (1.42); 2.7905 (1.30); 2.7837 (0.99); 2.7787 (1.05); 2.7738 (0.65); 2.7668 (0.36); 1.5988 (1.62); 0.8753 (0.44); 0.8590 (0.46); 0.6462 (0.54); 0.6365 (0.52); 0.5750 (0.40); 0.5529 (3.04); 0.5446 (4.68); 0.5338 (4.01); 0.5284 (3.48); 0.5221 (2.93); 0.5162 (2.44); 0.5101 (2.44); 0.5042 (2.56); 0.4827 (0.41); -0.0002 (3.38)
1.74	CDCl <sub>3</sub>	400	7.4043 (3.55); 7.4017 (3.82); 7.3915 (3.91); 7.3888 (4.04); 7.3738 (3.24); 7.3535 (5.81); 7.3223 (3.54); 7.3183 (4.98); 7.3080 (4.33); 7.3027 (3.03); 7.2878 (2.03); 7.2824 (1.54); 7.2622 (7.55); 7.1913 (0.33); 7.1459 (2.22); 7.1246 (2.71); 7.1158 (3.68); 7.1119 (2.87); 7.1030 (3.41); 7.0567 (3.78); 7.0541 (3.91); 7.0480 (3.01); 7.0454 (2.83); 7.0087 (4.57); 6.8715 (2.43); 5.4500 (0.33); 5.3475 (0.33); 5.2980 (9.30); 4.8413 (0.33); 3.8029 (16.00); 3.7188 (1.05); 2.8966 (0.36); 2.8828 (1.06); 2.8690 (1.42); 2.8583 (1.05); 2.8551 (1.03); 2.8447 (0.36); 1.5776 (3.50); 0.6868 (0.33); 0.6796 (0.32); 0.5987 (15.22); 0.5844 (10.35); -0.0002 (5.05)
1.75	CDCl <sub>3</sub>	400	7.4437 (2.38); 7.4403 (1.54); 7.4265 (3.45); 7.4233 (2.29); 7.4113 (0.38); 7.4069 (0.35); 7.3897 (4.45); 7.3869 (4.71); 7.3768 (4.78); 7.3740 (5.34); 7.3665 (1.87); 7.3620 (3.59); 7.3523 (9.20); 7.3418 (3.70); 7.3322 (2.03); 7.3240 (1.87); 7.3185 (1.15); 7.3140 (1.67); 7.3016 (0.70); 7.2602 (9.29); 7.2458 (0.44); 7.1640 (2.37); 7.1233 (2.84); 7.1145 (3.99); 7.1105 (2.97); 7.1017 (3.74); 7.0862 (0.36); 7.0660 (4.02); 7.0632 (4.21); 7.0573 (3.06); 7.0545 (2.88); 7.0268 (4.87); 6.8896 (2.50); 6.8713 (0.49); 6.8637 (0.45); 6.6552 (0.47); 5.4889 (0.36); 5.2963 (1.08); 4.8847 (0.56); 3.7883 (16.00); 3.7105 (2.21); 2.8446 (0.35); 2.8381 (0.63); 2.8328 (1.01); 2.8272 (1.01); 2.8214 (1.42); 2.8164 (1.30); 2.8096 (0.99); 2.8046 (1.06); 2.7928 (0.36); 1.5734 (5.63); 0.9027 (0.46); 0.8859 (0.48); 0.6866 (0.58); 0.6769 (0.56); 0.5979 (0.34); 0.5667 (4.70); 0.5561 (4.35); 0.5518 (3.97); 0.5452 (3.00); 0.5389 (2.48); 0.5343 (2.60); 0.5273 (2.71); 0.5060 (0.44); -0.0002 (6.11)
1.76	CDCl <sub>3</sub>	300	7.4544 (0.78); 7.4298 (1.24); 7.4166 (2.21); 7.4066 (2.40); 7.4003 (2.37); 7.3903 (2.39); 7.3758 (1.09); 7.3640 (1.11); 7.3577 (1.38); 7.3462 (2.34); 7.3215 (2.95); 7.3129 (5.33); 7.2960 (1.86); 7.2620 (15.68); 7.2335 (1.37); 7.1606 (1.09); 7.1481 (1.08); 6.9107 (0.52); 6.7273 (0.86); 6.5449 (0.44); 5.3000 (16.00); 4.8196 (1.73); 3.8859 (7.43); 2.6546 (0.75); 2.6368 (1.05); 2.6196 (0.87); 1.6035 (2.99); 0.3898 (3.92); -0.0002 (9.43); -0.0111 (0.41)
1.77	CDCl <sub>3</sub>	300	7.4566 (2.69); 7.4003 (4.12); 7.3966 (4.16); 7.3832 (4.70); 7.3795 (4.61); 7.3510 (1.98); 7.3237 (4.26); 7.2863 (4.57); 7.2794 (4.27); 7.2635 (13.10); 7.2521 (2.07); 7.1257 (2.50); 7.1139 (3.56); 7.1089 (2.80); 7.0970 (2.98); 7.0366 (2.11); 6.9239 (0.87); 6.7414 (1.66); 6.5588 (0.82); 5.3001 (16.00); 4.8548 (2.26); 3.9024 (12.53); 2.7151 (1.46); 2.6992 (1.22); 1.6097 (12.04); 0.4364 (6.96); -0.0002 (7.61); -0.0112 (0.36)
1.78	CDCl <sub>3</sub>	300	7.4624 (1.67); 7.4377 (2.96); 7.4157 (3.04); 7.3912 (6.50); 7.3844 (9.83); 7.3805 (8.16); 7.3672 (12.81); 7.3632 (11.23); 7.3427 (3.62); 7.3369 (2.60); 7.3223 (4.78); 7.2981 (4.23); 7.2936 (3.79); 7.2741 (1.48); 7.2623 (14.13); 7.1224 (3.25); 7.1107 (4.82); 7.1059 (3.83); 7.0939 (4.12); 7.0479 (2.76); 6.9183 (1.16); 6.7353 (2.04); 6.5529 (1.02); 5.2971 (16.00); 4.9052 (4.35); 3.8875 (15.85); 2.6807 (1.55); 2.6630 (2.23); 2.6461 (1.86); 1.6409 (11.46); 1.2576 (0.80); 0.5635 (0.38); 0.5459 (0.39); 0.5197 (0.34); 0.4130 (8.93); -0.0002 (8.24); -0.0111 (0.40)
1.79	CDCl <sub>3</sub>	300	7.4779 (3.20); 7.4713 (3.41); 7.4184 (2.88); 7.4146 (3.00); 7.4013 (3.62); 7.3974 (3.60); 7.3783 (3.24); 7.3507 (5.98); 7.3206 (0.36); 7.3105 (3.08); 7.3034 (2.87); 7.2958 (1.35); 7.2933 (1.43); 7.2835 (1.97); 7.2758 (1.51); 7.2620 (21.82); 7.2438 (0.81); 7.1424 (2.74); 7.1307 (3.67); 7.1253 (2.55); 7.1136 (3.35); 7.1060 (0.67); 7.0943 (0.63); 7.0888 (0.51); 7.0771 (0.55); 7.0607 (3.66); 7.0568 (3.70); 7.0490 (2.93); 7.0451 (2.80); 7.0385 (2.57); 6.8557 (3.88); 6.8491 (0.83); 6.8451 (0.66); 6.8370 (0.55); 6.8334 (0.51); 6.8196 (0.40); 6.6726 (2.47); 6.6382 (0.76); 6.4568 (0.38); 5.7191 (2.47); 5.6674 (3.05); 5.3010 (0.64); 5.2568 (3.30); 5.2052 (2.69); 4.9090 (0.35); 4.8552 (0.57); 4.7080 (0.63); 4.6539 (0.37); 3.8923 (16.00); 3.8199 (3.03); 3.7058 (0.32); 2.9310 (0.42); 2.9159 (0.91); 2.9080 (0.92); 2.8934 (1.76); 2.8825 (0.78); 2.8779 (1.04); 2.8692 (0.98); 2.8541 (0.50); 1.5651 (17.03); 0.6562

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.33); 0.6456 (0.44); 0.6347 (0.51); 0.6269 (0.53); 0.6217 (0.60); 0.6149 (0.71); 0.6092 (0.86); 0.6035 (1.51); 0.5947 (0.96); 0.5894 (1.39); 0.5783 (1.44); 0.5724 (1.44); 0.5666 (1.62); 0.5627 (1.72); 0.5574 (2.17); 0.5534 (2.03); 0.5415 (1.61); 0.5310 (3.32); 0.5259 (4.15); 0.5165 (1.77); 0.5029 (3.64); 0.4939 (0.93); 0.4881 (0.64); 0.4790 (0.39); 0.0107 (0.40); -0.0002 (12.43); -0.0111 (0.50)
I.80	DMSO-d <sub>6</sub>	400	7.6976 (2.06); 7.6903 (2.50); 7.6854 (2.26); 7.6781 (2.56); 7.6526 (0.65); 7.6452 (0.70); 7.6403 (0.70); 7.6329 (0.72); 7.6073 (2.60); 7.6040 (2.83); 7.6000 (2.41); 7.5967 (2.21); 7.4263 (0.74); 7.4231 (0.81); 7.4190 (0.75); 7.4158 (0.73); 7.4077 (0.33); 7.3935 (0.52); 7.3898 (0.60); 7.3751 (0.59); 7.3662 (1.37); 7.3537 (16.00); 7.3427 (1.34); 7.3232 (0.59); 7.3203 (0.66); 7.3044 (0.55); 7.2902 (0.86); 7.2865 (0.80); 7.2752 (2.60); 7.2719 (2.80); 7.2630 (2.46); 7.2597 (2.37); 7.2248 (0.57); 7.2052 (0.44); 7.1654 (1.40); 7.0872 (0.71); 7.0840 (0.70); 7.0749 (0.69); 7.0717 (0.67); 7.0306 (2.50); 6.9279 (0.36); 6.8965 (1.59); 6.7941 (0.83); 6.6603 (0.42); 5.7591 (0.79); 5.4815 (1.96); 5.4429 (2.41); 5.1709 (2.44); 5.1323 (1.98); 4.7637 (0.86); 4.7471 (0.87); 3.8789 (12.46); 3.7715 (3.79); 3.3326 (27.96); 3.0486 (0.37); 2.8784 (0.80); 2.8706 (0.54); 2.8640 (1.28); 2.8587 (0.59); 2.8506 (0.85); 2.8353 (0.33); 2.5108 (1.74); 2.5064 (3.64); 2.5019 (4.95); 2.4974 (3.53); 2.4930 (1.64); 0.8648 (0.54); 0.8599 (0.62); 0.8467 (0.54); 0.8418 (0.66); 0.5922 (0.41); 0.5813 (0.43); 0.5659 (0.42); 0.5551 (0.41); 0.5157 (0.38); 0.5073 (0.82); 0.4916 (3.08); 0.4857 (2.34); 0.4805 (2.60); 0.4783 (2.55); 0.4741 (2.49); 0.4655 (2.62); 0.4522 (1.31); 0.4415 (0.59); -0.0002 (2.06)
I.81	CDCl <sub>3</sub>	300	7.4885 (1.33); 7.4829 (1.62); 7.4594 (2.36); 7.4478 (1.76); 7.4423 (1.53); 7.4246 (2.89); 7.4178 (2.94); 7.4040 (3.08); 7.4002 (3.13); 7.3927 (1.90); 7.3868 (5.57); 7.3832 (4.56); 7.3687 (4.35); 7.3627 (3.87); 7.3458 (4.06); 7.3393 (3.15); 7.3227 (2.12); 7.3172 (1.73); 7.3088 (0.83); 7.2974 (1.48); 7.2838 (0.44); 7.2781 (0.41); 7.2612 (12.17); 7.1413 (2.66); 7.1296 (3.63); 7.1242 (2.40); 7.1126 (3.40); 7.1042 (0.91); 7.0925 (0.94); 7.0870 (0.82); 7.0751 (1.14); 7.0702 (3.71); 7.0663 (3.72); 7.0585 (2.82); 7.0547 (2.57); 7.0374 (2.41); 6.8544 (4.35); 6.8422 (1.38); 6.8391 (0.95); 6.6715 (2.44); 6.6605 (1.15); 6.4790 (0.58); 5.6689 (2.35); 5.6179 (3.40); 5.4015 (3.56); 5.3505 (2.49); 5.2989 (13.10); 4.9459 (0.50); 4.8927 (0.86); 4.7558 (0.90); 4.7023 (0.52); 3.9180 (0.42); 3.8810 (16.00); 3.8149 (4.61); 3.0148 (0.46); 2.8861 (0.43); 2.8715 (0.92); 2.8620 (0.99); 2.8479 (1.66); 2.8331 (1.08); 2.8234 (1.03); 2.8091 (0.52); 2.0461 (0.75); 1.5780 (8.35); 1.2595 (0.74); 0.8811 (0.45); 0.8629 (0.32); 0.7150 (0.40); 0.6982 (0.47); 0.6908 (0.32); 0.6767 (0.42); 0.6573 (0.39); 0.6099 (0.36); 0.6050 (0.43); 0.5995 (0.51); 0.5918 (0.51); 0.5844 (0.59); 0.5769 (1.23); 0.5715 (1.42); 0.5625 (1.35); 0.5573 (1.66); 0.5541 (1.56); 0.5501 (1.83); 0.5463 (1.65); 0.5396 (1.78); 0.5360 (1.77); 0.5238 (0.83); 0.5173 (0.82); 0.5006 (1.91); 0.4740 (3.85); 0.4661 (2.00); 0.4494 (3.63); 0.4426 (1.74); 0.4309 (0.65); 0.4152 (0.33); -0.0002 (6.73); -0.0111 (0.33)
I.82	DMSO-d <sub>6</sub>	300	7.6565 (2.31); 7.6519 (2.45); 7.6312 (2.73); 7.6271 (2.80); 7.5811 (3.33); 7.5713 (3.94); 7.5648 (3.67); 7.5550 (3.98); 7.4281 (4.04); 7.4240 (5.10); 7.4185 (4.83); 7.4145 (3.64); 7.3994 (2.89); 7.3938 (2.83); 7.3748 (2.60); 7.3690 (3.85); 7.3639 (2.46); 7.3448 (3.21); 7.3399 (3.06); 7.3203 (1.56); 7.3156 (1.29); 7.2558 (3.74); 7.2506 (3.17); 7.2316 (2.71); 7.2259 (2.18); 7.1510 (3.67); 7.1470 (3.48); 7.1346 (3.47); 7.1306 (3.23); 7.0567 (2.32); 6.8768 (4.84); 6.6970 (2.57); 5.7583 (1.32); 5.5704 (0.67); 5.5470 (2.50); 5.5236 (2.53); 5.5001 (0.72); 3.7983 (0.63); 3.7817 (0.86); 3.7513 (16.00); 3.3323 (79.42); 2.5141 (2.40); 2.5081 (5.08); 2.5020 (6.92); 2.4960 (5.01); 2.4901 (2.39); 2.2452 (1.05); 2.2318 (1.45); 2.2224 (1.46); 2.2093 (1.13); 2.1999 (0.70); 1.5503 (0.37); 1.5256 (9.71); 1.5021 (9.65); 1.2567 (0.54); 1.2339 (0.58); 0.8514 (0.45); 0.6700 (1.08); 0.6558 (1.16); 0.6352 (1.05); 0.6215 (1.13); 0.5212 (0.37); 0.4900 (1.01); 0.4672 (3.24); 0.4476 (3.26); 0.4246 (0.98); 0.3927 (0.33); 0.2002 (0.50); 0.1862 (1.17); 0.1683 (1.15); 0.1525 (1.23); 0.1475 (1.21); 0.1384 (1.03); -0.0002 (1.60)
I.83	DMSO-d <sub>6</sub>	400	7.6543 (2.40); 7.6487 (2.56); 7.6336 (3.32); 7.6279 (3.68); 7.6090 (4.06); 7.6066 (4.25); 7.5964 (4.33); 7.5940 (4.35); 7.5523 (4.27); 7.5482 (4.11); 7.5385 (6.41); 7.5175 (4.98); 7.5135 (4.87); 7.5110 (4.54); 7.5043 (4.55); 7.5019 (4.22); 7.1724 (4.05); 7.1632 (4.15); 7.1597 (4.10); 7.1506 (3.60); 7.1288 (2.32); 6.9940 (5.14); 6.8593 (2.58); 4.7338 (10.66); 3.8105 (16.00); 3.3354 (136.50); 2.8912 (1.43); 2.7323 (1.81); 2.6767 (0.46); 2.6724 (0.52); 2.5116 (22.74); 2.5076 (42.76);

$\delta$	$\delta$	$\delta$	$^1\text{H-NMR}$
	S <sub>e</sub>		2.5032 (54.35); 2.4987 (39.28); 2.3298 (0.36); 0.6982 (0.80); 0.6793 (3.80); 0.661 1 (3.84); 0.6505 (2.03) ; 0.6328 (1.12) ; 0.61 4 7 (4.26); -0.0002 (3.50)
1.84	DMSO -d <sub>6</sub>	400	8.4383 (2.00) ; 7.9531 (1.27); 7.6323 (1.63); 7.6266 (1.77); 7.61 15 (2.38); 7.6058 (2.57); 7.5882 (2.71) ; 7.5857 (2.85) ; 7.5756 (2.90); 7.5731 (2.91) ; 7.5324 (0.48); 7.5246 (5.48) ; 7.51 2 6 (0.55); 7.5037 (3.87); 7.4989 (3.1 8); 7.4963 (3.04); 7.4871 (6.14); 7.481 6 (3.38); 7.3031 (1.43) ; 7.1 671 (3.42); 7.1 568 (2.93); 7.1 477 (2.86); 7.1 442 (2.83) ; 7.1 351 (2.55); 7.031 2 (1.60); 4.7353 (8.60); 4.0028 (0.73); 3.9365 (16.00) ; 3.3358 (133.71) ; 2.8909 (10.35); 2.731 7 (8.1 6); 2.5250 (0.88); 2.51 18 (16.39) ; 2.5074 (32.26); 2.5029 (42.03); 2.4984 (30.1 6); 2.4939 (14.35); 0.781 8 (0.61) ; 0.7639 (2.64); 0.7507 (2.70) ; 0.7465 (2.33); 0.7336 (0.85); 0.6453 (1.1 6); 0.6345 (2.93); 0.6275 (3.02) ; 0.61 9 7 (2.22) ; 0.6059 (0.64); -0.0002 (1.48)
1.85	DMSO -d <sub>6</sub>	400	7.9520 (2.1 7) ; 7.6571 (3.04); 7.651 7 (3.68); 7.6363 (4.83); 7.6308 (5.52); 7.6098 (6.1 2); 7.6074 (6.78); 7.5970 (6.92) ; 7.5947 (7.56); 7.5806 (1.61) ; 7.5553 (8.43); 7.5499 (8.91) ; 7.5408 (6.34); 7.5206 (4.24); 7.4993 (6.21) ; 7.491 7 (6.69); 7.4449 (0.48); 7.4246 (0.33); 7.1 7 1 8 (6.87) ; 7.1 626 (6.81) ; 7.1 592 (7.34); 7.1 501 (6.69); 7.041 4 (3.76) ; 6.9067 (1.88); 4.8291 (0.50); 4.7580 (1.91) ; 4.6485 (0.48); 4.6352 (0.44); 4.6023 (0.36); 3.8998 (14.91) ; 3.8399 (0.78); 3.3922 (0.36) ; 3.3824 (0.37); 3.3705 (0.66); 3.3296 (446.02); 3.31 4 9 (58.1 2); 3.2498 (0.42) ; 2.8904 (16.00); 2.8742 (1.83); 2.8000 (1.78); 2.7470 (0.46); 2.7307 (13.68); 2.71 4 9 (1.87); 2.6755 (0.96); 2.6708 (1.30); 2.6665 (1.1 0); 2.5240 (3.1 9) ; 2.51 0 8 (67.26); 2.5065 (139.91) ; 2.5020 (193.78); 2.4975 (157.95); 2.4931 (102.24); 2.3333 (0.87); 2.3288 (1.23) ; 2.3239 (1.01) ; 1.2329 (0.59); 0.5809 (9.05) ; 0.3909 (0.33); 0.3798 (0.32) ; 0.3741 (0.34) ; 0.291 0 (0.33); 0.0082 (0.43) ; -0.0002 (12.46); -0.0085 (1.32); -0.01 7 4 (1.15)
1.86	DMSO -d <sub>6</sub>	300	7.661 9 (0.94) ; 7.6552 (3.05); 7.6455 (4.27); 7.6389 (3.35); 7.6291 (3.83); 7.5601 (2.83); 7.3903 (0.63); 7.3753 (1.44) ; 7.3599 (2.36); 7.3539 (2.44); 7.3370 (8.90); 7.3303 (10.89); 7.3200 (5.1 4); 7.3002 (2.96); 7.2733 (1.71) ; 7.2444 (2.35); 7.2307 (2.29); 7.1 1 6 2 (1.60); 6.9369 (3.38) ; 6.7576 (1.66) ; 5.7585 (16.00) ; 4.7825 (1.1 4); 4.7308 (1.87) ; 4.5964 (3.74); 4.5451 (2.30); 4.0407 (0.45); 4.01 6 8 (0.45); 3.7932 (11.40); 3.7691 (2.40); 3.3246 (65.38); 2.51 3 4 (7.55); 2.5074 (16.59); 2.501 4 (23.21) ; 2.4953 (17.28); 2.4894 (8.57); 2.2340 (1.03) ; 1.9889 (1.92) ; 1.2454 (0.74); 1.1980 (0.56) ; 1.1743 (1.04); 1.1505 (0.53); 0.8583 (0.81) ; 0.8354 (0.40); 0.691 2 (1.76); 0.6787 (1.71) ; 0.61 1 2 (3.86) ; 0.5388 (1.1 1) ; 0.5221 (1.65); 0.5085 (1.84); 0.4935 (1.20) ; 0.4778 (0.92); 0.31 7 9 (0.69); 0.2990 (1.35); 0.2793 (1.36); 0.2591 (0.60); 0.01 0 7 (0.61) ; -0.0002 (19.88); -0.01 1 2 (1.32); -0.0300 (0.53)
1.87	DMSO -d <sub>6</sub>	300	8.2069 (2.28) ; 7.6450 (0.60); 7.6354 (2.67); 7.6258 (2.96); 7.61 9 1 (2.98); 7.6095 (2.95); 7.5589 (2.73); 7.5546 (3.1 9) ; 7.5492 (2.90); 7.5451 (2.46); 7.3785 (0.45); 7.3638 (1.02) ; 7.3503 (1.1 2); 7.3424 (1.95); 7.3203 (9.82); 7.31 0 2 (6.62); 7.281 9 (1.14); 7.2558 (1.86); 7.2466 (2.71) ; 7.2424 (2.76); 7.2303 (2.54); 7.2262 (2.55); 7.1 8 1 1 (0.38); 7.0746 (3.43); 6.8938 (1.69); 5.7588 (10.28); 4.9653 (0.37); 4.91 4 0 (0.40); 4.661 0 (7.72); 4.5009 (0.61) ; 4.4485 (0.44) ; 3.91 0 8 (16.00); 3.3252 (54.81) ; 2.51 3 1 (8.04); 2.5072 (17.47); 2.501 2 (24.33); 2.4952 (18.05); 2.4893 (9.03); 2.3828 (1.07) ; 1.9889 (0.73); 1.2460 (0.45); 1.1742 (0.38); 1.11 6 2 (0.32); 0.8579 (0.46); 0.7279 (0.80); 0.71 7 0 (0.82) ; 0.7074 (1.08); 0.6972 (0.94); 0.651 4 (7.60); 0.631 9 (4.26) ; 0.5295 (0.71) ; 0.51 2 6 (1.19); 0.4995 (1.38) ; 0.4831 (0.89); 0.4700 (0.71) ; 0.3797 (0.79); 0.361 6 (1.36) ; 0.3390 (1.44); 0.31 9 5 (0.58); 0.0667 (0.34); 0.01 0 7 (0.60); -0.0002 (20.43) ; -0.01 1 0 (1.06)
1.88	DMSO -d <sub>6</sub>	300	7.6782 (1.85) ; 7.6685 (2.60); 7.6625 (2.51) ; 7.6526 (2.56); 7.5882 (2.71) ; 7.4648 (0.34); 7.401 0 (0.81) ; 7.3907 (0.89) ; 7.3527 (7.1 6); 7.3351 (7.55); 7.2708 (2.64); 7.2568 (2.37); 7.1 855 (1.02); 7.0056 (1.90); 6.8260 (1.01) ; 5.7590 (16.00); 4.7793 (0.43); 4.661 4 (0.34); 4.61 2 1 (0.34) ; 4.5572 (0.48); 4.5270 (0.47); 4.4856 (0.50); 4.4481 (0.36); 3.8948 (10.42); 3.8731 (2.22); 3.8076 (1.07); 3.3281 (26.01) ; 2.51 3 6 (3.24) ; 2.5078 (6.87); 2.501 8 (9.46); 2.4958 (7.04); 2.4900 (3.51) ; 2.3032 (1.12); 1.9891 (0.97); 1.2345 (0.33) ; 1.1744 (0.50); 0.9057 (0.62); 0.8842 (0.61) ; 0.8583 (0.44) ; 0.6684 (1.80); 0.4787 (6.36); 0.4597 (5.71) ; 0.2306 (0.65); 0.2096 (1.27); 0.1 892 (1.22); 0.1 691 (0.55); 0.01 0 6 (0.35) ; -0.0002 (7.26)
1.89	DMSO	400	7.9535 (2.1 3) ; 7.8799 (2.96); 7.7769 (2.93); 7.7635 (3.05); 7.6908 (2.54); 7.6854

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
	-d6		(3.73); 7.6795 (5.12); 7.6721 (6.35); 7.6660 (16.00); 7.6599 (8.98); 7.5405 (4.30); 7.5300 (4.86); 7.5120 (1.73); 7.1156 (1.35); 4.7634 (3.00); 3.3266 (34.64); 2.8906 (15.72); 2.7591 (1.05); 2.7498 (1.04); 2.7321 (13.81); 2.7310 (12.75); 2.6714 (0.38); 2.5246 (1.35); 2.5113 (19.93); 2.5069 (38.48); 2.5023 (49.73); 2.4978 (36.03); 2.4933 (17.79); 2.3291 (0.32); 1.2328 (0.43); 0.4968 (3.04)
I.90	DMSO-d6	400	7.9592 (1.10); 7.6231 (1.40); 7.6177 (1.53); 7.6022 (2.07); 7.5969 (2.30); 7.5859 (2.54); 7.5732 (2.62); 7.5180 (6.73); 7.4972 (2.85); 7.4822 (2.69); 7.4733 (2.95); 7.4594 (3.30); 7.4545 (3.22); 7.2633 (1.23); 7.1559 (2.13); 7.1466 (2.47); 7.1433 (2.47); 7.1340 (3.10); 7.1209 (4.33); 6.9770 (1.31); 4.7256 (8.01); 3.7009 (16.00); 3.3383 (77.93); 2.9336 (1.08); 2.8967 (6.96); 2.7379 (6.17); 2.5126 (31.37); 2.5085 (39.58); 2.5044 (30.67); 0.7949 (0.61); 0.7774 (2.79); 0.7637 (2.80); 0.7475 (0.81); 0.6375 (0.93); 0.6261 (2.82); 0.6199 (3.18); 0.5983 (0.66)
I.91	DMSO-d6	400	7.9538 (2.05); 7.7909 (3.14); 7.7777 (3.25); 7.6529 (2.55); 7.6322 (3.49); 7.5910 (11.01); 7.5871 (11.30); 7.5754 (5.37); 7.5432 (6.60); 7.5364 (7.64); 7.5182 (2.11); 7.1674 (5.91); 7.1582 (6.06); 7.1548 (6.00); 7.1456 (5.42); 7.1330 (0.46); 7.1050 (1.72); 4.7589 (4.07); 3.3295 (98.26); 2.8910 (16.00); 2.7715 (1.32); 2.7321 (13.67); 2.6765 (0.36); 2.6719 (0.47); 2.6676 (0.35); 2.5250 (1.44); 2.5119 (24.40); 2.5075 (48.20); 2.5029 (63.23); 2.4984 (46.18); 2.4940 (22.61); 2.3296 (0.41); 1.2326 (0.51); 0.5041 (4.27)
I.92	DMSO-d6	400	7.9597 (1.26); 7.8454 (2.23); 7.8424 (2.46); 7.8383 (2.56); 7.8353 (2.41); 7.6759 (2.16); 7.6682 (3.31); 7.6631 (3.69); 7.6561 (2.32); 7.6469 (1.93); 7.6413 (2.06); 7.5325 (3.85); 7.5274 (5.00); 7.5118 (4.38); 7.4911 (3.28); 7.4794 (2.48); 7.4764 (2.47); 7.4668 (2.20); 7.4638 (2.16); 7.2705 (1.22); 7.1269 (3.62); 7.1215 (2.70); 7.1170 (2.48); 6.9840 (1.30); 4.7338 (7.52); 3.6956 (16.00); 3.3346 (51.15); 2.9300 (0.93); 2.8969 (9.24); 2.7379 (7.77); 2.5173 (14.52); 2.5131 (28.12); 2.5086 (36.52); 2.5042 (26.80); 2.5001 (13.38); 0.7795 (0.55); 0.7616 (2.41); 0.7483 (2.49); 0.7315 (0.78); 0.6405 (0.89); 0.6296 (2.56); 0.6227 (2.73); 0.6011 (0.58)
I.93	DMSO-d6	400	8.4388 (2.24); 7.9530 (1.61); 7.8652 (2.49); 7.8621 (2.87); 7.8581 (3.01); 7.8550 (2.80); 7.6800 (1.88); 7.6748 (4.02); 7.6675 (2.49); 7.6620 (3.59); 7.6548 (4.49); 7.5667 (3.50); 7.5618 (3.29); 7.5176 (4.38); 7.4970 (5.96); 7.4853 (2.52); 7.4822 (2.48); 7.3073 (1.38); 7.1714 (3.09); 7.0355 (1.51); 4.7403 (8.70); 3.9310 (16.00); 3.3248 (17.80); 2.8902 (11.57); 2.7310 (9.38); 2.5062 (33.21); 2.5018 (42.78); 2.4975 (32.22); 0.7647 (0.65); 0.7471 (2.82); 0.7337 (2.80); 0.7172 (0.95); 0.6793 (0.37); 0.6503 (1.14); 0.6396 (3.12); 0.6323 (3.25); 0.6111 (0.64)
I.94	DMSO-d6	400	7.9530 (2.26); 7.8319 (4.22); 7.6915 (6.95); 7.6843 (9.05); 7.6792 (6.95); 7.6717 (7.06); 7.6693 (5.53); 7.6635 (5.64); 7.6251 (7.36); 7.6201 (6.57); 7.5334 (3.25); 7.5130 (2.74); 7.4907 (4.22); 7.4792 (3.85); 7.1809 (1.46); 7.0460 (3.08); 6.9114 (1.55); 4.7651 (1.67); 4.6924 (0.43); 3.8939 (13.42); 3.3240 (35.59); 2.8903 (16.00); 2.7993 (1.31); 2.7481 (0.35); 2.7317 (13.32); 2.7307 (12.96); 2.6755 (0.47); 2.6709 (0.60); 2.6665 (0.46); 2.5240 (1.85); 2.5106 (34.74); 2.5063 (67.19); 2.5019 (87.11); 2.4973 (66.33); 2.4930 (35.18); 2.3332 (0.43); 2.3286 (0.58); 2.3242 (0.44); 1.2338 (0.36); 0.5785 (6.98)
I.95	DMSO-d6	400	7.8724 (3.92); 7.8691 (4.24); 7.8652 (4.48); 7.8619 (4.06); 7.7070 (2.74); 7.7014 (3.28); 7.6959 (4.41); 7.6884 (5.36); 7.6832 (6.61); 7.6810 (4.92); 7.6760 (4.66); 7.6389 (3.99); 7.5370 (5.94); 7.5192 (5.71); 7.5162 (8.88); 7.5068 (4.06); 7.5035 (3.80); 7.1417 (2.50); 7.0070 (5.71); 6.8723 (2.86); 4.7423 (9.72); 3.8089 (16.00); 3.3396 (135.49); 2.8965 (0.47); 2.7378 (0.74); 2.7082 (0.58); 2.6869 (0.52); 2.6822 (0.60); 2.6776 (0.64); 2.6729 (0.52); 2.5308 (1.09); 2.5175 (19.44); 2.5130 (38.21); 2.5085 (49.62); 2.5039 (35.56); 2.4994 (16.86); 2.3353 (0.33); 0.6902 (0.78); 0.6712 (3.37); 0.6583 (3.81); 0.6438 (2.43); 0.6282 (2.19); 0.6170 (4.68); 0.5883 (0.65)
I.96	CDCl <sub>3</sub>	300	7.5207 (2.06); 7.5145 (2.67); 7.4954 (1.37); 7.4883 (0.75); 7.4679 (2.55); 7.4595 (2.93); 7.4541 (2.12); 7.4489 (2.71); 7.4443 (2.59); 7.4243 (4.23); 7.4184 (2.13); 7.4085 (1.38); 7.4016 (3.21); 7.3969 (2.24); 7.3919 (2.25); 7.3675 (0.33); 7.3558 (2.84); 7.3513 (2.55); 7.3391 (1.71); 7.3346 (1.51); 7.2615 (9.91); 7.1397 (1.17);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			6.9580 (2.41); 6.7762 (1.20); 4.8623 (4.86); 2.7801 (1.30); 2.7666 (16.00); 2.7120 (0.40); 2.6939 (0.89); 2.6759 (1.29); 2.6680 (0.60); 2.6584 (0.81); 2.6393 (0.39); 2.0079 (11.92); 0.7344 (4.35); 0.7175 (8.10); 0.0105 (0.35); -0.0002 (8.97); -0.0110 (0.35)
I.97	CDCl <sub>3</sub>	300	7.5025 (5.18); 7.4782 (2.47); 7.4707 (1.56); 7.4080 (2.68); 7.4033 (1.42); 7.3833 (0.96); 7.3787 (1.48); 7.3193 (1.68); 7.3156 (2.70); 7.3079 (3.05); 7.3037 (2.76); 7.2990 (4.11); 7.2955 (4.06); 7.2917 (1.46); 7.2608 (24.60); 7.1435 (1.15); 7.1030 (2.44); 7.0904 (2.30); 7.0865 (2.23); 7.0739 (1.93); 6.9620 (2.39); 6.7804 (1.18); 4.8492 (4.66); 2.7721 (16.00); 2.7193 (0.37); 2.7015 (0.81); 2.6893 (0.64); 2.6832 (1.24); 2.6751 (0.56); 2.6662 (0.75); 2.6465 (0.38); 2.0084 (2.62); 1.7796 (0.35); 0.7405 (3.59); 0.7245 (7.49); 0.0106 (0.60); -0.0002 (21.57); -0.0111 (0.98)
I.98	DMSO-d <sub>6</sub>	400	7.8743 (2.77); 7.8709 (2.94); 7.8670 (3.08); 7.8636 (2.84); 7.6876 (2.76); 7.6802 (4.11); 7.6750 (4.56); 7.6678 (2.89); 7.6591 (2.12); 7.6534 (2.42); 7.5976 (3.55); 7.5922 (3.01); 7.5178 (4.72); 7.5031 (3.23); 7.4996 (3.77); 7.4972 (4.24); 7.4906 (2.75); 7.4872 (2.60); 6.3721 (1.50); 5.7602 (14.39); 4.6978 (8.14); 3.3151 (49.04); 2.7528 (0.61); 2.6811 (0.41); 2.6764 (0.54); 2.6720 (0.40); 2.5581 (0.40); 2.5298 (1.51); 2.5165 (30.75); 2.5120 (62.21); 2.5075 (83.77); 2.5029 (58.25); 2.4985 (26.57); 2.4616 (1.48); 2.3389 (0.53); 2.3342 (0.84); 2.3194 (16.00); 2.2336 (13.16); 2.2176 (1.03); 1.9948 (0.53); 1.2547 (0.39); 0.8653 (0.56); 0.6528 (2.11); 0.6468 (1.88); 0.6357 (2.80); 0.6308 (3.13); 0.6200 (3.53); 0.5902 (0.34)
I.99	DMSO-d <sub>6</sub>	400	7.6490 (1.63); 7.6433 (1.71); 7.6282 (2.22); 7.6225 (2.35); 7.5973 (2.55); 7.5947 (2.77); 7.5846 (2.70); 7.5820 (2.77); 7.5255 (4.82); 7.5167 (2.99); 7.5140 (3.00); 7.5075 (3.75); 7.5048 (6.17); 7.4914 (3.39); 7.4858 (3.10); 7.1698 (2.75); 7.1607 (2.73); 7.1572 (2.73); 7.1481 (2.44); 6.3657 (1.60); 6.2137 (0.38); 5.7605 (9.99); 4.6910 (8.22); 3.3149 (17.81); 2.7776 (0.60); 2.6810 (0.33); 2.6766 (0.46); 2.6720 (0.34); 2.5299 (1.31); 2.5165 (26.66); 2.5120 (54.08); 2.5075 (72.89); 2.5031 (51.73); 2.4987 (24.30); 2.4615 (2.44); 2.3338 (16.00); 2.2410 (12.92); 2.2166 (1.59); 0.6872 (0.47); 0.6825 (0.40); 0.6682 (2.23); 0.6556 (2.13); 0.6508 (2.41); 0.6394 (1.38); 0.6234 (0.67); 0.6167 (1.20); 0.6067 (3.05); 0.6006 (2.35); 0.5773 (0.55)
I.100	CDCl <sub>3</sub>	300	7.5598 (0.33); 7.4026 (1.57); 7.3791 (2.40); 7.3674 (0.62); 7.3469 (1.14); 7.3413 (1.73); 7.3268 (5.40); 7.3215 (6.84); 7.3101 (3.13); 7.3008 (1.77); 7.2870 (1.66); 7.2772 (1.62); 7.2616 (41.92); 7.0396 (0.56); 6.9313 (2.84); 6.9275 (4.05); 6.9236 (3.01); 6.9106 (0.38); 6.8822 (0.47); 6.8573 (1.24); 6.8339 (2.08); 6.6758 (0.54); 5.3014 (3.67); 4.8646 (3.03); 4.7952 (0.58); 3.8063 (10.47); 2.6539 (0.83); 2.6458 (0.89); 2.2918 (15.81); 2.2889 (16.00); 1.5597 (19.64); 0.6581 (0.64); 0.4789 (2.90); 0.0106 (0.81); -0.0002 (26.70); -0.0111 (1.10)
I.101	CDCl <sub>3</sub>	300	7.3715 (1.32); 7.3472 (2.46); 7.3262 (1.58); 7.2615 (15.00); 7.0253 (3.52); 6.9880 (3.87); 6.9503 (1.42); 6.9271 (4.79); 6.8397 (2.22); 6.8043 (4.06); 6.6597 (1.05); 4.8134 (7.09); 4.7607 (0.54); 3.8169 (14.92); 2.7091 (1.89); 2.2885 (16.00); 1.5596 (8.55); 1.2605 (1.67); 0.8797 (0.83); 0.8591 (0.45); 0.6600 (0.41); 0.5524 (3.49); 0.4914 (5.35); -0.0002 (9.45)
I.102	CDCl <sub>3</sub>	300	7.2624 (6.14); 7.1745 (1.31); 7.1503 (2.75); 7.1294 (1.87); 7.0819 (3.94); 7.0464 (7.19); 7.0127 (4.74); 6.9660 (1.20); 6.8272 (2.46); 6.6449 (1.24); 4.5122 (7.66); 3.8127 (16.00); 2.6894 (1.85); 2.0376 (15.34); 1.5828 (3.72); 1.2619 (4.30); 0.8819 (2.24); 0.8592 (1.11); 0.5384 (3.51); 0.4652 (5.60); -0.0002 (3.73)
I.103	CDCl <sub>3</sub>	300	7.3355 (1.66); 7.3081 (4.02); 7.2917 (4.52); 7.2630 (8.88); 7.2455 (1.38); 7.0234 (0.92); 6.9413 (4.10); 6.8401 (2.24); 6.8256 (3.45); 6.6591 (0.91); 4.8230 (5.47); 3.8201 (12.86); 2.6909 (1.23); 2.2878 (16.00); 1.5802 (5.23); 1.3044 (0.53); 1.2633 (3.61); 0.9007 (1.06); 0.8817 (2.40); 0.8585 (1.11); 0.5699 (1.88); 0.5529 (2.37); 0.4952 (3.78); -0.0002 (4.50)
I.104	CDCl <sub>3</sub>	300	7.3169 (5.01); 7.2950 (2.48); 7.2625 (18.19); 7.1359 (3.87); 7.1090 (3.03); 7.0849 (3.46); 7.0543 (4.12); 7.0089 (1.27); 6.8263 (2.50); 6.6441 (1.24); 4.5144 (6.99); 3.8168 (16.00); 2.6692 (1.42); 2.0418 (15.49); 1.5658 (10.62); 1.2640 (6.31);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			0.9003 (1.90); 0.8819 (3.91); 0.8595 (1.83); 0.5584 (2.63); 0.5393 (2.78); 0.4661 (4.51); -0.0002 (9.52)
I.105	CDCl <sub>3</sub>	300	7.3831 (0.44); 7.3389 (8.98); 7.2967 (2.55); 7.2619 (8.78); 7.1981 (3.49); 7.1748 (2.36); 7.0887 (3.36); 7.0389 (4.84); 6.8432 (2.04); 6.6615 (1.02); 4.5570 (5.60); 3.8020 (16.00); 2.6317 (1.63); 2.0489 (14.61); 1.5768 (3.83); 1.2609 (3.80); 0.8982 (0.99); 0.8818 (1.79); 0.8592 (0.86); 0.5043 (3.23); 0.4527 (5.69); -0.0002 (5.74)
I.106	CDCl <sub>3</sub>	300	7.5153 (1.30); 7.5089 (1.49); 7.4884 (1.72); 7.4820 (2.06); 7.4395 (3.06); 7.4336 (2.55); 7.3328 (3.26); 7.3058 (2.49); 7.2604 (6.38); 7.0912 (4.65); 7.0882 (4.75); 6.9076 (2.40); 6.8274 (3.35); 6.7256 (1.21); 4.7972 (4.16); 3.8110 (10.32); 3.1818 (0.50); 3.1599 (0.67); 3.1378 (0.55); 2.6784 (0.70); 2.2822 (13.42); 2.2799 (13.20); 1.6394 (0.40); 1.2553 (16.00); 1.2326 (15.85); 0.6925 (5.89); 0.6751 (6.50); -0.0002 (3.82)
I.107	DMSO-d <sub>6</sub>	400	7.9596 (0.98); 7.3502 (1.46); 7.3057 (1.80); 7.1624 (2.04); 7.1526 (1.95); 7.1312 (2.03); 7.0761 (1.66); 7.0571 (1.19); 7.0222 (1.11); 6.8876 (0.51); 4.4113 (1.50); 3.8981 (5.87); 3.3335 (55.00); 2.8972 (7.15); 2.7376 (6.12); 2.6775 (0.92); 2.6732 (0.90); 2.6684 (0.85); 2.5173 (19.48); 2.5131 (38.11); 2.5087 (49.97); 2.5042 (36.96); 2.4999 (18.73); 2.3357 (16.00); 2.0823 (0.83); 2.0052 (5.45); 0.4339 (1.53); 0.4197 (1.54); 0.2997 (2.09)
I.108	DMSO-d <sub>6</sub>	400	7.9594 (0.89); 7.3347 (1.62); 7.3275 (2.04); 7.2913 (2.72); 7.2889 (2.75); 7.2835 (2.10); 7.2810 (1.88); 7.1501 (1.40); 7.1309 (2.22); 7.0874 (3.28); 7.0706 (3.96); 7.0514 (2.45); 6.9550 (2.86); 6.8203 (1.44); 4.4051 (4.73); 3.8768 (0.42); 3.7975 (7.62); 3.3326 (49.39); 2.8972 (6.87); 2.7382 (5.94); 2.7370 (5.39); 2.6773 (0.32); 2.6087 (0.54); 2.5305 (0.96); 2.5173 (18.02); 2.5128 (35.60); 2.5083 (46.25); 2.5037 (32.95); 2.4992 (15.63); 2.3308 (16.00); 1.9912 (6.89); 0.5426 (1.81); 0.5280 (1.81); 0.3573 (0.82); 0.3461 (2.57); 0.3399 (2.76); 0.3321 (2.32); 0.3188 (0.61)
I.109	DMSO-d <sub>6</sub>	400	8.3015 (1.25); 7.9594 (1.22); 7.3304 (2.67); 7.3227 (3.42); 7.2765 (2.93); 7.2740 (3.10); 7.2686 (2.97); 7.2661 (3.10); 7.1296 (3.44); 7.1128 (2.51); 7.0635 (3.99); 7.0541 (4.77); 7.0349 (2.71); 6.9931 (1.18); 4.4111 (7.96); 3.9227 (14.49); 3.3348 (115.25); 2.8971 (8.53); 2.7378 (7.98); 2.6819 (0.38); 2.6775 (0.46); 2.6730 (0.35); 2.5129 (52.43); 2.5085 (66.66); 2.5040 (50.12); 2.3246 (16.00); 1.9905 (12.44); 1.2422 (2.19); 0.8604 (0.36); 0.6343 (0.69); 0.6166 (3.11); 0.6032 (3.15); 0.5863 (0.88); 0.3938 (0.99); 0.3820 (3.02); 0.3765 (3.34); 0.3680 (2.89); 0.3549 (0.79)
I.110	CDCl <sub>3</sub>	300	7.3756 (1.26); 7.3489 (4.44); 7.3319 (2.76); 7.3261 (3.00); 7.3051 (0.74); 7.2994 (1.05); 7.2668 (3.17); 7.2589 (5.45); 7.1606 (3.07); 7.1497 (3.59); 7.0686 (1.11); 7.0156 (2.30); 7.0122 (2.44); 7.0084 (1.73); 7.0047 (2.31); 7.0013 (2.19); 6.8867 (2.27); 6.7048 (1.15); 4.8120 (4.07); 3.7987 (10.32); 3.2218 (0.46); 3.1999 (0.61); 3.1783 (0.50); 2.6886 (0.66); 2.2719 (13.90); 2.2694 (13.95); 2.1144 (1.12); 2.1119 (1.13); 1.9994 (1.91); 1.6300 (0.33); 1.2809 (16.00); 1.2582 (15.79); 1.2276 (0.55); 1.2048 (0.38); 0.6836 (4.34); 0.6677 (6.38); -0.0002 (3.28)
I.111	CDCl <sub>3</sub>	300	7.3456 (0.97); 7.3201 (1.19); 7.2644 (6.54); 7.0856 (0.36); 7.0556 (2.11); 7.0291 (2.75); 7.0101 (3.07); 6.8475 (1.77); 6.6653 (0.89); 5.2992 (0.41); 4.8579 (4.08); 3.8348 (6.18); 3.3616 (0.64); 3.3053 (0.90); 3.0424 (2.60); 2.9864 (2.02); 2.9441 (0.67); 2.3022 (16.00); 2.0545 (0.32); 2.0279 (15.56); 1.9779 (0.42); 1.6889 (7.51); 1.2571 (0.54); 0.6785 (4.58); 0.6658 (4.53); 0.0710 (0.38); -0.0002 (4.90)
I.112	CDCl <sub>3</sub>	300	7.2678 (4.22); 7.1154 (0.68); 7.0922 (1.59); 7.0713 (1.15); 7.0323 (1.87); 7.0041 (2.87); 6.9665 (0.65); 6.8310 (1.39); 6.6482 (0.69); 4.6718 (0.63); 4.6210 (0.90); 4.3924 (1.87); 4.3398 (1.31); 3.8202 (9.35); 3.7865 (10.51); 2.7141 (1.03); 2.0745 (16.00); 1.7033 (0.44); 0.5201 (2.05); 0.4613 (3.03); -0.0002 (2.80)
I.113	CDCl <sub>3</sub>	300	7.3075 (3.16); 7.2849 (1.65); 7.2652 (5.72); 7.0754 (1.89); 7.0489 (1.48); 7.0079 (0.68); 6.8252 (1.35); 6.6446 (0.68); 5.3021 (1.13); 4.6670 (0.62); 4.6160 (0.84); 4.3973 (1.82); 4.3459 (1.23); 3.8188 (9.25); 3.7810 (10.18); 2.6837 (0.90); 2.0744

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(16.00); 0.5156 (1.88); 0.4530 (2.71); -0.0002 (2.98)
I.114	CDCl <sub>3</sub>	300	8.1178 (1.51); 7.4683 (0.33); 7.3267 (7.47); 7.3017 (1.93); 7.2936 (1.98); 7.2636 (14.03); 7.1355 (2.24); 7.1130 (1.74); 7.0259 (0.74); 6.8433 (1.44); 6.6630 (0.70); 5.3018 (0.91); 4.6987 (0.76); 4.6479 (0.98); 4.4543 (2.07); 4.4042 (1.42); 3.8071 (12.83); 3.7949 (16.00); 2.6466 (1.03); 2.0884 (13.90); 2.0107 (1.37); 1.3277 (0.34); 0.4458 (3.69); -0.0002 (9.19)
I.115	DMSO-d <sub>6</sub>	400	7.9594 (1.42); 7.6339 (0.36); 7.1278 (1.68); 7.1085 (2.27); 7.0972 (1.15); 7.0745 (3.63); 7.0014 (2.98); 6.9823 (2.38); 6.9623 (2.34); 6.8276 (1.17); 4.4328 (0.40); 4.3937 (0.89); 4.3465 (1.80); 4.3081 (0.77); 3.8008 (8.50); 3.6939 (13.64); 3.3755 (0.67); 3.3357 (62.91); 2.8971 (10.91); 2.7376 (9.15); 2.6777 (0.38); 2.6020 (0.49); 2.5827 (0.66); 2.5712 (0.78); 2.5573 (1.28); 2.5434 (0.77); 2.5309 (1.09); 2.5174 (20.71); 2.5131 (41.31); 2.5086 (54.41); 2.5040 (39.66); 2.4996 (19.50); 2.3480 (0.45); 2.3349 (0.91); 2.3202 (16.00); 2.0206 (4.76); 1.9755 (0.34); 1.9326 (5.27); 1.2422 (0.62); 1.1652 (2.45); 1.0757 (0.39); 0.5028 (1.66); 0.4873 (1.61); 0.3243 (2.27)
I.116	DMSO-d <sub>6</sub>	600	8.3021 (0.40); 7.9525 (1.84); 7.2309 (0.33); 7.1408 (0.65); 7.0994 (1.66); 7.0868 (1.97); 7.0448 (3.41); 6.9764 (3.66); 6.9637 (3.06); 4.4137 (0.48); 4.3877 (1.21); 4.3601 (2.73); 4.3345 (1.07); 3.9173 (11.50); 3.6683 (15.05); 3.3157 (36.34); 2.8902 (15.37); 2.7312 (12.38); 2.7304 (12.59); 2.7072 (0.36); 2.6156 (0.42); 2.6126 (0.56); 2.6096 (0.39); 2.5219 (0.87); 2.5188 (1.14); 2.5157 (1.26); 2.5069 (29.43); 2.5039 (63.77); 2.5009 (87.61); 2.4978 (61.31); 2.4948 (27.81); 2.3881 (0.37); 2.3850 (0.51); 2.3820 (0.40); 2.3070 (16.00); 2.0052 (4.73); 1.9250 (6.45); 0.5836 (2.52); 0.5808 (2.90); 0.5722 (2.48); 0.5692 (3.04); 0.5559 (0.33); 0.3832 (0.39); 0.3784 (0.36); 0.3685 (1.30); 0.3625 (1.52); 0.3532 (1.73); 0.3469 (1.54); 0.3395 (0.36); 0.3361 (0.38); 0.3327 (0.40); -0.0002 (0.34)
I.117	DMSO-d <sub>6</sub>	600	7.9524 (1.60); 7.1537 (1.69); 7.1167 (1.41); 7.1051 (1.87); 7.0186 (1.05); 6.9964 (1.35); 6.9838 (1.18); 6.9291 (0.54); 4.4248 (0.44); 4.3290 (0.44); 4.3040 (0.34); 3.9118 (0.36); 3.8923 (6.00); 3.7054 (8.25); 3.3153 (30.96); 2.8902 (13.20); 2.7305 (10.84); 2.6275 (0.74); 2.6188 (0.69); 2.6157 (0.83); 2.6127 (0.83); 2.6096 (0.61); 2.5219 (0.89); 2.5188 (1.13); 2.5157 (1.18); 2.5069 (28.74); 2.5039 (62.25); 2.5009 (85.23); 2.4978 (60.12); 2.4948 (27.18); 2.3881 (0.37); 2.3850 (0.51); 2.3820 (0.37); 2.3185 (16.00); 2.0244 (6.09); 1.9382 (6.06); 0.3733 (1.48); 0.3618 (1.51); 0.2715 (1.53); -0.0002 (0.39)
I.118	CDCl <sub>3</sub>	300	7.3643 (1.73); 7.3377 (2.22); 7.2631 (9.67); 7.1747 (1.23); 7.1688 (1.39); 7.1482 (0.92); 7.1422 (1.11); 7.0994 (2.20); 7.0938 (1.89); 7.0674 (0.69); 6.8854 (1.40); 6.7036 (0.71); 4.8063 (2.49); 3.8027 (6.99); 3.7789 (14.27); 3.1896 (0.40); 3.1673 (0.33); 2.7057 (0.43); 2.5779 (0.60); 2.2381 (14.57); 2.2263 (16.00); 1.2794 (10.10); 1.2567 (10.08); 1.2261 (0.37); 0.6811 (2.70); 0.6671 (4.04); -0.0002 (8.38); -0.0111 (0.38)
I.119	CDCl <sub>3</sub>	300	7.4937 (1.61); 7.3654 (0.73); 7.3426 (1.72); 7.3207 (2.50); 7.2971 (2.02); 7.2645 (6.65); 7.2480 (5.27); 7.0573 (1.71); 7.0217 (2.37); 6.9824 (1.81); 6.9552 (0.97); 6.9103 (2.07); 6.8930 (1.96); 6.8402 (0.97); 6.7769 (0.32); 6.6590 (0.48); 4.7619 (1.13); 4.7155 (3.13); 3.8043 (13.63); 2.6533 (1.22); 1.9765 (2.45); 1.6424 (1.85); 1.2592 (2.23); 1.2370 (16.00); 0.8942 (0.56); 0.8813 (1.02); 0.8584 (0.52); 0.5244 (1.98); 0.4729 (3.40); 0.3044 (0.59); 0.2426 (0.46); -0.0002 (1.68)
I.120	CDCl <sub>3</sub>	300	7.4691 (0.48); 7.3340 (7.25); 7.2621 (11.83); 7.2438 (3.15); 7.0411 (0.39); 6.9112 (2.39); 6.8936 (2.22); 6.8605 (0.77); 6.6780 (0.43); 5.3011 (1.33); 4.7626 (2.86); 3.7936 (12.36); 2.5852 (1.27); 1.9265 (1.43); 1.5809 (4.68); 1.2415 (16.00); 0.8999 (0.41); 0.8819 (0.75); 0.8591 (0.39); 0.4581 (4.03); -0.0002 (5.84)
I.121	CDCl <sub>3</sub>	300	7.5065 (0.93); 7.3756 (0.45); 7.3521 (1.18); 7.3256 (5.64); 7.2600 (15.39); 7.0208 (0.64); 6.9101 (2.87); 6.8922 (2.66); 6.8393 (1.30); 6.6572 (0.64); 5.2952 (2.13); 4.7194 (4.07); 3.8033 (16.00); 2.6295 (1.61); 2.0451 (0.34); 2.0056 (0.55); 1.6784 (1.93); 1.2579 (1.02); 1.2342 (3.75); 0.5226 (2.58); 0.4697 (4.47); 0.2968 (0.35); -0.0002 (1.16)

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
I.122	CDCl <sub>3</sub>	300	7.7214 (0.33); 7.7074 (0.37); 7.7027 (0.54); 7.6969 (0.34); 7.6801 (0.39); 7.6752 (0.35); 7.6400 (0.32); 7.5781 (0.33); 7.5733 (0.35); 7.5540 (0.66); 7.5486 (0.73); 7.5297 (0.95); 7.5245 (0.95); 7.4932 (0.79); 7.4890 (0.68); 7.4860 (0.69); 7.4835 (0.69); 7.4733 (0.57); 7.4684 (0.63); 7.4632 (0.57); 7.4586 (0.58); 7.4452 (0.41); 7.4402 (0.41); 7.4354 (0.41); 7.4161 (0.40); 7.3971 (0.94); 7.3939 (0.82); 7.3740 (1.01); 7.3516 (2.36); 7.3266 (2.96); 7.3074 (1.71); 7.2856 (7.06); 7.2669 (10.04); 7.2649 (9.45); 7.2508 (2.15); 7.2262 (0.95); 7.0627 (5.07); 7.0533 (0.91); 7.0343 (7.02); 7.0080 (2.05); 6.9989 (1.15); 6.9113 (5.05); 6.8929 (4.65); 6.8353 (1.16); 6.6538 (0.58); 4.7678 (0.83); 4.7111 (3.18); 3.8292 (9.48); 3.7868 (16.00); 2.5563 (0.97); 2.5427 (1.24); 2.5350 (1.35); 2.5214 (1.01); 1.9993 (3.38); 1.6818 (11.65); 0.4810 (1.95); 0.4334 (3.44); 0.3107 (0.40); -0.0002 (4.05)
I.123	CDCl <sub>3</sub>	300	7.3317 (5.22); 7.3278 (5.51); 7.3004 (7.98); 7.2944 (10.43); 7.2894 (10.23); 7.2707 (8.20); 7.2654 (7.02); 7.0154 (0.59); 6.9103 (5.45); 6.8918 (5.05); 6.8338 (1.18); 6.6517 (0.59); 4.7070 (3.36); 3.8274 (9.73); 3.7913 (16.00); 2.5767 (1.02); 2.5631 (1.29); 2.5550 (1.42); 2.5413 (1.07); 2.1703 (0.83); 2.0033 (0.83); 1.6703 (1.60); 0.4860 (2.02); 0.4373 (3.54); -0.0002 (3.36)
I.124	DMSO-d <sub>6</sub>	600	8.2839 (0.39); 7.9524 (1.38); 7.5143 (6.12); 7.5051 (6.27); 7.3474 (0.33); 7.2185 (0.41); 7.1477 (2.53); 7.1349 (4.66); 7.1287 (0.95); 7.1034 (2.14); 7.1018 (2.24); 7.0889 (1.31); 7.0740 (3.21); 7.0648 (3.11); 7.0514 (0.43); 7.0421 (0.58); 7.0379 (0.45); 7.0218 (3.31); 4.5605 (6.04); 3.9174 (12.77); 3.8831 (2.24); 3.7621 (5.84); 3.3154 (27.90); 2.8899 (11.98); 2.7468 (0.34); 2.7310 (10.02); 2.7303 (9.92); 2.6154 (0.34); 2.6124 (0.47); 2.6094 (0.33); 2.5217 (0.75); 2.5186 (0.96); 2.5155 (0.96); 2.5067 (24.01); 2.5037 (52.34); 2.5007 (71.84); 2.4976 (50.77); 2.4946 (23.12); 2.3848 (0.44); 2.3046 (16.00); 0.6131 (0.67); 0.6037 (2.37); 0.6011 (2.92); 0.5924 (2.90); 0.5895 (2.51); 0.5809 (0.78); 0.4094 (0.82); 0.4017 (2.48); 0.3980 (2.68); 0.3951 (2.52); 0.3918 (2.42); 0.3833 (0.67)
I.125	DMSO-d <sub>6</sub>	600	7.9524 (1.12); 7.5347 (5.02); 7.5255 (5.16); 7.1595 (0.96); 7.1468 (1.73); 7.1193 (2.18); 7.1063 (1.20); 7.0910 (1.22); 7.0822 (1.18); 7.0491 (2.99); 6.9518 (0.68); 6.8615 (0.34); 4.5520 (1.52); 3.7933 (6.63); 3.7769 (2.68); 3.3151 (18.46); 2.8901 (9.24); 2.7309 (7.53); 2.6154 (0.69); 2.6125 (0.83); 2.6094 (0.70); 2.5217 (0.93); 2.5186 (1.24); 2.5154 (1.57); 2.5067 (26.08); 2.5038 (54.00); 2.5007 (72.59); 2.4977 (51.21); 2.4947 (23.27); 2.3879 (0.32); 2.3849 (0.44); 2.3107 (16.00); 0.5223 (1.18); 0.3627 (1.94)
I.126	DMSO-d <sub>6</sub>	600	7.9523 (1.51); 7.5533 (1.75); 7.5442 (1.91); 7.1668 (0.81); 7.1540 (1.44); 7.1388 (1.76); 7.1159 (2.34); 7.1067 (2.62); 7.0216 (0.98); 6.9320 (0.48); 4.5534 (1.17); 3.8938 (5.62); 3.8440 (0.35); 3.7884 (5.94); 3.3151 (27.68); 2.8900 (12.48); 2.7306 (9.89); 2.6545 (0.74); 2.6154 (0.43); 2.6125 (0.57); 2.6094 (0.39); 2.5217 (0.91); 2.5187 (1.14); 2.5156 (1.18); 2.5068 (29.16); 2.5038 (63.51); 2.5007 (87.52); 2.4977 (61.42); 2.4946 (27.96); 2.3880 (0.37); 2.3849 (0.52); 2.3818 (0.36); 2.3155 (16.00); 0.4134 (1.53); 0.4041 (1.61); 0.3183 (1.82)
I.127	CDCl <sub>3</sub>	300	7.2613 (8.47); 7.2327 (3.85); 7.2141 (5.50); 7.1438 (2.06); 7.1121 (2.14); 7.0861 (1.40); 7.0187 (0.41); 6.9009 (2.84); 6.8824 (2.58); 6.8365 (0.81); 6.6544 (0.39); 5.2977 (10.68); 4.8496 (0.49); 4.7986 (0.72); 4.5979 (1.26); 4.5469 (0.95); 3.8024 (11.69); 2.3605 (16.00); 2.2686 (0.83); 2.2585 (0.86); 1.9415 (0.86); 1.6005 (6.80); 1.2391 (7.80); 0.7717 (0.61); 0.6657 (2.49); 0.5884 (0.80); 0.5760 (0.76); 0.2372 (0.61); 0.2201 (0.59); -0.0002 (5.30)
I.128	CDCl <sub>3</sub>	300	7.2760 (2.12); 7.2610 (21.38); 7.2478 (4.50); 7.2386 (1.78); 7.2295 (4.89); 7.2127 (2.05); 7.1947 (0.41); 7.1069 (1.68); 7.0806 (1.16); 6.9286 (0.76); 6.9175 (2.14); 6.8991 (1.95); 6.7471 (1.22); 6.5648 (0.63); 5.2995 (10.81); 4.8790 (0.39); 4.5651 (0.35); 4.5239 (0.32); 3.8953 (8.05); 3.8383 (0.82); 3.8070 (9.81); 3.7048 (0.54); 2.3600 (16.00); 2.3113 (0.85); 2.2991 (0.96); 2.2876 (0.96); 1.9188 (0.70); 1.5686 (17.00); 1.2692 (0.40); 1.2558 (0.39); 1.2409 (5.62); 0.9681 (0.45); 0.7559 (0.56); 0.7351 (0.63); 0.5474 (4.29); 0.5278 (3.89); 0.1806 (0.41); 0.1599 (0.91); 0.1381 (0.88); 0.1169 (0.39); 0.0106 (0.45); -0.0002 (13.04); -0.0111 (0.54)
I.129	CDCl <sub>3</sub>	300	7.6351 (6.68); 7.5756 (3.53); 7.5489 (4.51); 7.4334 (2.70); 7.4124 (3.53); 7.3874

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(4.29); 7.3617 (7.80); 7.3390 (4.69); 7.2956 (3.27); 7.2639 (6.49); 7.1235 (3.61); 7.0902 (5.63); 7.0619 (3.84); 7.0349 (1.83); 6.9947 (1.50); 6.8128 (2.95); 6.6311 (1.46); 5.2969 (2.91); 4.7080 (11.02); 3.7811 (16.00); 2.6746 (2.78); 1.6259 (3.93); 0.4876 (5.12); 0.4326 (8.57); -0.0002 (2.23)
I.130	CDCl <sub>3</sub>	300	7.6454 (6.14); 7.5748 (3.19); 7.5479 (4.16); 7.4382 (2.33); 7.4119 (3.26); 7.3917 (8.59); 7.3633 (4.24); 7.3402 (16.00); 7.2992 (3.05); 7.2677 (4.20); 6.9910 (1.31); 6.8092 (2.64); 6.6271 (1.33); 5.2955 (6.54); 4.7100 (9.75); 3.7792 (14.39); 2.6567 (2.26); 1.6329 (2.03); 1.2575 (0.33); 0.4846 (4.24); 0.4298 (7.06); -0.0002 (1.93)
I.131	CDCl <sub>3</sub>	300	7.6488 (6.20); 7.5730 (3.79); 7.5456 (4.90); 7.4080 (16.00); 7.3561 (5.80); 7.3304 (3.02); 7.2890 (3.34); 7.2613 (12.36); 7.2411 (1.78); 7.0072 (1.15); 6.8257 (2.27); 6.6429 (1.16); 5.2985 (11.19); 4.7539 (8.35); 3.7643 (13.72); 2.6060 (2.41); 1.5884 (5.73); 1.2606 (0.43); 0.4186 (9.48); -0.0002 (5.97)
I.132	CDCl <sub>3</sub>	300	7.8324 (1.79); 7.8304 (1.74); 7.8079 (2.11); 7.8041 (1.90); 7.7701 (7.57); 7.6025 (1.38); 7.5964 (1.50); 7.5757 (1.99); 7.5696 (2.33); 7.5617 (2.00); 7.5596 (2.05); 7.5373 (1.85); 7.5338 (2.54); 7.4991 (3.25); 7.4936 (2.93); 7.4534 (3.54); 7.4266 (2.52); 7.3769 (0.87); 7.3724 (0.96); 7.3527 (1.92); 7.3485 (1.81); 7.3264 (1.60); 7.3212 (1.41); 7.3110 (1.80); 7.3067 (1.81); 7.2855 (2.07); 7.2819 (2.08); 7.2582 (6.90); 7.0821 (1.23); 6.9002 (2.57); 6.7183 (1.29); 4.8599 (4.24); 3.7937 (10.83); 3.2470 (0.52); 3.2245 (0.69); 3.2031 (0.56); 2.7440 (0.68); 1.6244 (0.34); 1.3057 (16.00); 1.2830 (15.90); 0.7058 (6.26); 0.6884 (6.42); -0.0002 (3.97)
I.133	DMSO-d <sub>6</sub>	400	8.1518 (1.96); 7.9592 (1.72); 7.6958 (1.97); 7.6753 (2.27); 7.4884 (0.93); 7.4700 (1.16); 7.4155 (1.13); 7.3955 (2.17); 7.3770 (1.36); 7.3505 (1.16); 7.3330 (2.48); 7.3155 (1.77); 7.2965 (0.93); 7.2615 (2.12); 7.2445 (2.35); 7.2248 (1.53); 7.1416 (0.50); 7.0063 (0.99); 6.8709 (0.50); 4.6566 (1.52); 3.8834 (5.46); 3.3338 (71.58); 2.8967 (11.50); 2.7376 (10.46); 2.6772 (0.40); 2.6731 (0.32); 2.6318 (0.91); 2.5125 (44.84); 2.5083 (56.92); 2.5040 (41.41); 2.3777 (16.00); 2.3351 (0.59); 0.3667 (1.68); 0.3532 (1.81); 0.3062 (2.20)
I.134	DMSO-d <sub>6</sub>	400	8.1078 (2.14); 7.9593 (1.34); 7.6782 (2.69); 7.6578 (3.10); 7.4627 (0.91); 7.4436 (1.14); 7.4073 (1.10); 7.4044 (1.19); 7.3867 (2.28); 7.3689 (1.59); 7.3658 (1.45); 7.3416 (2.02); 7.3229 (4.44); 7.3057 (2.42); 7.2872 (0.93); 7.2439 (2.22); 7.2244 (1.53); 7.1778 (3.48); 7.0694 (0.48); 6.9342 (0.94); 6.7992 (0.48); 4.6353 (4.61); 3.7660 (3.99); 3.3318 (45.16); 2.8970 (10.57); 2.7377 (8.88); 2.6776 (0.40); 2.5999 (0.58); 2.5306 (1.37); 2.5175 (21.93); 2.5131 (43.82); 2.5085 (57.92); 2.5040 (42.14); 2.4996 (20.52); 2.3737 (16.00); 2.3395 (0.36); 2.3354 (0.46); 2.3309 (0.35); 0.4849 (1.69); 0.4703 (1.70); 0.3469 (2.48); 0.3400 (2.66); 0.3203 (0.59)
I.135	DMSO-d <sub>6</sub>	400	8.0895 (5.93); 7.9589 (4.86); 7.6336 (2.71); 7.6134 (3.12); 7.4568 (1.70); 7.4379 (2.21); 7.3960 (1.20); 7.3781 (2.36); 7.3576 (1.55); 7.3236 (3.91); 7.3042 (5.27); 7.2889 (1.06); 7.2325 (3.43); 7.2130 (1.72); 7.1730 (3.92); 7.0964 (1.98); 6.9607 (0.99); 4.6390 (7.98); 3.8549 (14.29); 3.3331 (25.97); 2.8970 (11.94); 2.7377 (10.82); 2.6737 (1.17); 2.6675 (1.17); 2.5129 (37.76); 2.5089 (50.17); 2.5049 (39.97); 2.3716 (16.00); 2.3355 (0.41); 0.5672 (0.68); 0.5503 (3.07); 0.5365 (2.99); 0.5202 (0.88); 0.3980 (1.00); 0.3867 (3.03); 0.3803 (3.33); 0.3588 (0.71)
I.136	CDCl <sub>3</sub>	300	7.2629 (7.33); 7.2385 (1.71); 7.2223 (2.89); 7.1981 (4.42); 7.1883 (3.67); 7.1790 (3.30); 7.0990 (2.54); 7.0641 (3.72); 7.0275 (2.90); 7.0172 (2.57); 6.8981 (2.94); 6.8329 (2.52); 6.6506 (1.25); 5.3013 (6.33); 4.6068 (7.30); 3.8084 (16.00); 2.6672 (2.04); 2.0470 (0.50); 1.5786 (5.17); 1.2611 (0.79); 0.5466 (3.64); 0.4663 (5.92); -0.0002 (4.76)
I.137	CDCl <sub>3</sub>	300	7.3695 (4.96); 7.3296 (2.26); 7.3006 (3.36); 7.2627 (9.84); 7.2023 (6.07); 7.1803 (5.22); 7.0112 (1.29); 6.8939 (3.12); 6.8300 (2.53); 6.6483 (1.27); 5.3016 (7.17); 4.6054 (7.60); 3.8095 (16.00); 2.6454 (2.07); 1.5697 (6.96); 1.2584 (0.57); 0.5450 (3.77); 0.4648 (6.06); -0.0002 (6.55)
I.138	CDCl <sub>3</sub>	300	7.3820 (7.51); 7.3405 (2.20); 7.3311 (2.06); 7.2622 (13.85); 7.1877 (3.03); 7.1689

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(3.45); 7.0314 (1.08); 6.9276 (2.54); 6.9097 (2.39); 6.8496 (2.22); 6.6672 (1.08); 5.3009 (8.56); 4.6497 (5.34); 3.7978 (16.00); 2.5998 (1.64); 1.5690 (6.76); 0.5114 (3.06); 0.4536 (5.27); -0.0002 (8.07)
I.139	CDCl <sub>3</sub>	300	7.3400 (11.41); 7.2634 (3.97); 7.0324 (1.19); 6.9026 (4.94); 6.8494 (2.50); 6.7952 (3.79); 6.6680 (1.18); 5.2991 (4.39); 4.8370 (7.39); 3.8068 (16.00); 2.6631 (2.18); 1.6114 (2.98); 0.5420 (4.37); 0.4920 (6.92); -0.0002 (2.34)
I.140	CDCl <sub>3</sub>	300	7.5543 (3.30); 7.3321 (10.41); 7.2641 (7.26); 7.0833 (2.45); 6.9015 (1.37); 6.8738 (4.24); 6.7961 (3.89); 5.3011 (8.17); 4.8415 (9.19); 3.9545 (16.00); 3.8650 (0.34); 2.5510 (2.09); 1.6077 (7.32); 0.5980 (4.42); 0.5486 (6.24); -0.0002 (3.60)
I.141	CDCl <sub>3</sub>	300	7.4485 (1.16); 7.4253 (2.53); 7.4017 (2.19); 7.3960 (2.29); 7.3794 (2.99); 7.3736 (4.05); 7.3616 (2.00); 7.3482 (3.37); 7.3407 (3.80); 7.3191 (3.92); 7.2936 (2.65); 7.2627 (26.89); 6.9523 (0.33); 6.9292 (5.11); 6.9169 (6.09); 6.8608 (0.32); 6.8480 (0.35); 6.8122 (1.86); 6.7415 (1.68); 6.5585 (0.79); 5.3009 (16.00); 4.8900 (3.30); 3.8956 (12.53); 2.6963 (1.36); 2.6810 (1.19); 1.5897 (27.95); 1.2551 (0.38); 0.4454 (6.42); 0.4306 (5.71); 0.0107 (0.50); -0.0002 (15.96); -0.0111 (0.67)
I.142	CDCl <sub>3</sub>	300	7.4639 (1.38); 7.4396 (2.15); 7.4334 (2.09); 7.4105 (0.37); 7.4012 (1.53); 7.3943 (2.09); 7.3880 (1.89); 7.3734 (4.12); 7.3687 (4.80); 7.3632 (2.89); 7.3519 (2.47); 7.3431 (2.77); 7.3371 (2.56); 7.3213 (2.45); 7.3133 (2.06); 7.3044 (1.20); 7.2900 (0.57); 7.2763 (0.60); 7.2621 (11.81); 7.0401 (2.15); 6.9466 (4.58); 6.9341 (5.74); 6.9122 (1.25); 6.8997 (1.22); 6.8571 (4.00); 6.8354 (5.66); 6.8229 (4.58); 6.6739 (2.94); 6.6132 (1.12); 6.6007 (1.04); 6.4914 (0.47); 5.6253 (2.20); 5.5741 (3.41); 5.4036 (3.54); 5.3525 (2.29); 5.2999 (15.80); 4.9305 (0.44); 4.8773 (0.76); 4.7509 (0.79); 4.6980 (0.44); 3.8869 (16.00); 3.8212 (3.85); 3.0096 (0.39); 2.9206 (0.42); 2.9062 (0.88); 2.8966 (0.97); 2.8826 (1.63); 2.8679 (1.04); 2.8581 (0.98); 2.8436 (0.48); 1.5780 (4.45); 1.2572 (0.41); 0.7470 (0.37); 0.7306 (0.40); 0.7091 (0.35); 0.6375 (0.33); 0.6318 (0.42); 0.6243 (0.55); 0.6192 (0.58); 0.6107 (0.99); 0.6044 (1.64); 0.5902 (2.81); 0.5853 (1.86); 0.5760 (1.83); 0.5704 (1.96); 0.5626 (0.93); 0.5559 (0.66); 0.5362 (1.17); 0.5233 (0.91); 0.5154 (2.17); 0.5095 (3.63); 0.4855 (3.01); 0.4799 (1.78); 0.4692 (0.66); 0.4554 (0.35); -0.0002 (6.41)
I.143	CDCl <sub>3</sub>	300	7.3977 (3.80); 7.3931 (5.78); 7.3900 (5.16); 7.3881 (4.97); 7.3691 (8.85); 7.3595 (6.87); 7.3535 (4.33); 7.3309 (1.04); 7.3253 (1.04); 7.2632 (17.00); 7.0512 (9.12); 7.0381 (8.69); 6.8925 (8.85); 6.8795 (7.87); 6.8700 (4.57); 6.6879 (2.17); 5.3007 (3.95); 4.8247 (6.36); 3.8303 (16.00); 2.8258 (0.84); 1.5955 (14.53); 0.7069 (2.96); 0.6822 (6.01); 0.6734 (5.51); 0.0106 (0.35); -0.0002 (10.19); -0.0110 (0.42)
I.144	CDCl <sub>3</sub>	300	7.6998 (2.07); 7.3921 (1.26); 7.3791 (3.22); 7.3732 (3.90); 7.3644 (7.08); 7.3542 (5.03); 7.3480 (3.29); 7.3263 (0.88); 7.3200 (0.75); 7.2816 (1.58); 7.2632 (17.08); 7.0992 (3.16); 7.0400 (5.22); 7.0270 (6.18); 6.9167 (1.59); 6.8858 (6.13); 6.8728 (5.23); 5.3013 (10.35); 4.8286 (8.77); 3.9775 (16.00); 2.7972 (0.39); 2.7832 (0.88); 2.7747 (0.98); 2.7611 (1.68); 2.7546 (0.87); 2.7475 (1.03); 2.7389 (0.92); 2.7253 (0.48); 1.5904 (19.55); 0.8060 (0.39); 0.7791 (2.23); 0.7628 (2.32); 0.7567 (2.40); 0.7529 (2.11); 0.7489 (2.08); 0.7416 (1.91); 0.7327 (2.33); 0.7184 (4.15); 0.7087 (2.29); 0.7048 (2.49); 0.6922 (0.66); 0.6777 (0.42); 0.0107 (0.34); -0.0002 (10.09); -0.0112 (0.39)
I.145	CDCl <sub>3</sub>	300	7.4842 (2.91); 7.3946 (1.36); 7.3669 (7.32); 7.3588 (6.54); 7.3523 (5.27); 7.3313 (1.09); 7.3243 (1.06); 7.2630 (34.03); 7.0673 (6.12); 7.0543 (7.05); 6.9496 (1.55); 6.9110 (0.41); 6.8917 (6.86); 6.8788 (5.97); 6.7671 (3.16); 6.5848 (1.62); 5.3019 (16.00); 4.8850 (1.70); 3.9091 (14.33); 2.8478 (0.92); 1.5746 (48.59); 1.2597 (0.55); 0.8820 (0.55); 0.7237 (0.39); 0.6482 (3.92); 0.0101 (0.63); -0.0002 (18.11); -0.0101 (0.95)
I.146	CDCl <sub>3</sub>	300	7.3753 (1.55); 7.3668 (2.60); 7.3618 (2.85); 7.3603 (2.95); 7.3476 (6.47); 7.3322 (3.42); 7.3255 (2.43); 7.3044 (0.85); 7.2977 (0.75); 7.2610 (10.63); 7.1579 (2.64); 7.1568 (2.67); 7.0314 (4.39); 7.0184 (5.19); 7.0092 (0.40); 6.9917 (1.92); 6.9659 (1.46); 6.8913 (2.16); 6.8854 (2.07); 6.8786 (1.17); 6.8721 (5.29); 6.8591 (4.42); 5.2989 (10.35); 4.8298 (8.74); 3.6900 (16.00); 2.8278 (0.34); 2.8131 (0.69);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			2.8059 (0.75); 2.8012 (0.64); 2.7917 (1.46); 2.7837 (0.67); 2.7774 (0.82); 2.7699 (0.75); 2.7558 (0.41); 1.5834 (8.28); 0.7700 (1.83); 0.7663 (1.63); 0.7617 (2.15); 0.7557 (1.75); 0.7476 (2.00); 0.7396 (4.03); 0.7346 (3.84); 0.7297 (2.40); 0.7253 (3.61); 0.7206 (3.24); 0.7124 (2.00); 0.6980 (0.45); -0.0002 (6.53)
I.147	CDCl <sub>3</sub>	300	7.6075 (2.25); 7.4473 (5.12); 7.4291 (5.55); 7.3966 (1.51); 7.3635 (10.97); 7.3570 (8.95); 7.3357 (1.66); 7.3294 (1.67); 7.3165 (0.48); 7.3104 (0.41); 7.2999 (0.35); 7.2615 (19.43); 7.0855 (13.99); 7.0725 (16.00); 7.0238 (0.32); 6.9623 (2.53); 6.9458 (2.62); 6.9104 (0.58); 6.9042 (0.62); 6.8851 (9.25); 6.8721 (8.36); 5.2981 (2.92); 4.8960 (4.20); 4.8580 (1.18); 2.7181 (2.05); 2.7038 (1.71); 2.0453 (0.56); 1.5980 (12.24); 1.2826 (0.42); 1.2590 (1.26); 0.9027 (0.62); 0.8811 (1.51); 0.8579 (0.80); 0.7398 (0.33); 0.7164 (0.36); 0.6887 (0.41); 0.5703 (6.60); 0.0107 (0.35); -0.0002 (11.88); -0.0111 (0.59)
I.148	DMSO-d <sub>6</sub>	400	7.9608 (1.20); 7.5970 (2.02); 7.5830 (2.16); 7.2152 (1.49); 7.1967 (1.18); 7.1771 (2.30); 7.1516 (2.11); 7.1320 (1.00); 7.0903 (1.24); 7.0769 (1.14); 7.0537 (0.34); 7.0193 (0.93); 6.8840 (0.46); 4.4982 (1.19); 3.8940 (5.52); 3.8095 (0.34); 3.3326 (43.61); 2.8985 (9.01); 2.7387 (7.54); 2.6790 (0.36); 2.6202 (0.73); 2.5322 (0.99); 2.5188 (20.27); 2.5144 (40.74); 2.5099 (53.71); 2.5053 (39.29); 2.5009 (19.29); 2.3750 (0.56); 2.3456 (16.00); 2.3184 (0.34); 2.3125 (0.52); 0.4211 (1.41); 0.4087 (1.43); 0.3703 (0.40); 0.3123 (1.83)
I.149	DMSO-d <sub>6</sub>	400	8.2764 (1.58); 7.9594 (1.23); 7.5403 (4.34); 7.5262 (4.61); 7.2666 (1.09); 7.1789 (1.23); 7.1594 (2.62); 7.1275 (7.50); 7.1074 (2.26); 7.0607 (3.35); 7.0465 (3.18); 6.9948 (1.19); 4.4995 (7.50); 3.9185 (14.83); 3.3315 (47.59); 2.8972 (8.52); 2.7379 (7.48); 2.6819 (1.10); 2.6774 (1.11); 2.5127 (44.39); 2.5083 (57.28); 2.5039 (41.97); 2.3369 (16.00); 0.6244 (0.61); 0.6067 (2.82); 0.5934 (2.86); 0.5764 (0.78); 0.4174 (0.91); 0.4060 (2.74); 0.3998 (3.01); 0.3913 (2.51); 0.3779 (0.68)
I.150	DMSO-d <sub>6</sub>	600	7.9524 (1.13); 7.5548 (6.19); 7.5455 (6.72); 7.1857 (1.30); 7.1729 (2.11); 7.1424 (1.80); 7.1341 (3.70); 7.1213 (2.15); 7.0472 (1.27); 7.0353 (1.74); 6.9453 (2.29); 6.8558 (1.17); 4.4845 (2.43); 3.7854 (7.18); 3.3149 (24.39); 2.8902 (9.64); 2.7312 (7.88); 2.7303 (7.94); 2.6155 (0.36); 2.6125 (0.51); 2.6095 (0.37); 2.5433 (0.46); 2.5218 (0.99); 2.5187 (1.24); 2.5156 (1.26); 2.5069 (26.88); 2.5038 (58.49); 2.5008 (81.00); 2.4977 (56.89); 2.4947 (25.71); 2.3880 (0.35); 2.3850 (0.49); 2.3819 (0.35); 2.3351 (16.00); 0.5206 (1.34); 0.5116 (1.35); 0.3681 (0.72); 0.3605 (2.28); 0.3566 (2.44); 0.3509 (2.13); 0.3424 (0.56)
I.151	CDCl <sub>3</sub>	300	7.4143 (0.95); 7.4059 (2.59); 7.3860 (5.06); 7.3825 (5.32); 7.3759 (2.81); 7.3531 (0.43); 7.3477 (0.42); 7.2612 (20.50); 7.1329 (1.19); 7.0643 (3.59); 7.0513 (4.20); 6.9512 (2.45); 6.9043 (4.16); 6.8913 (3.55); 6.7694 (1.23); 4.8383 (4.95); 2.7845 (2.78); 2.7754 (16.00); 2.7248 (0.41); 2.7090 (0.65); 2.7032 (0.65); 2.6979 (0.60); 2.6887 (1.34); 2.6801 (0.61); 2.6734 (0.74); 2.6672 (0.66); 2.6524 (0.40); 2.0091 (2.69); 1.2548 (0.55); 0.7494 (1.99); 0.7247 (4.17); 0.0108 (0.47); -0.0002 (17.87); -0.0111 (0.82)
I.152	DMSO-d <sub>6</sub>	400	12.3603 (0.35); 7.6093 (2.02); 7.5883 (3.48); 7.5427 (3.67); 7.5384 (4.07); 7.5219 (2.08); 7.5175 (2.32); 7.4331 (4.97); 7.4139 (3.10); 7.4088 (3.76); 7.4041 (3.46); 7.3990 (3.74); 7.2003 (3.19); 7.1951 (3.93); 7.1905 (3.15); 7.1853 (3.39); 6.3690 (2.46); 6.2144 (1.21); 5.7633 (5.60); 5.7596 (5.29); 5.7580 (7.04); 4.6801 (9.22); 3.3172 (13.48); 3.3124 (16.92); 2.7667 (1.06); 2.6786 (0.71); 2.6745 (0.73); 2.5099 (110.05); 2.5055 (112.26); 2.4602 (6.40); 2.4139 (0.45); 2.3976 (0.33); 2.3265 (16.00); 2.2924 (0.65); 2.2385 (15.49); 2.2167 (4.74); 1.9925 (0.34); 1.2518 (0.56); 0.8628 (0.39); 0.6777 (0.66); 0.6619 (3.27); 0.6467 (3.84); 0.6033 (4.79)
I.153	CDCl <sub>3</sub>	300	7.9261 (1.41); 7.8996 (1.51); 7.6126 (0.90); 7.6066 (0.99); 7.5859 (1.16); 7.5798 (1.30); 7.5042 (1.63); 7.4999 (1.53); 7.4065 (2.09); 7.3794 (2.42); 7.3501 (1.98); 7.2975 (0.85); 7.2939 (0.89); 7.2743 (1.27); 7.2707 (1.46); 7.2546 (5.21); 7.2470 (1.07); 7.2437 (0.90); 7.2134 (4.73); 7.1835 (0.99); 7.1798 (0.98); 7.1568 (1.46); 7.1534 (1.06); 7.1337 (0.67); 7.1301 (0.64); 7.1112 (0.59); 6.9293 (1.14); 6.7473 (0.57); 4.8419 (2.12); 3.8511 (0.39); 3.8369 (16.00); 3.8092 (0.40); 3.7844 (6.71);

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			3.1986 (0.38); 2.7257 (0.43); 1.9960 (5.11); 1.2938 (9.79); 1.2712 (9.73); 1.2374 (3.28); 1.2259 (0.50); 1.2032 (0.35); 0.7016 (3.50); 0.6858 (2.91); -0.0002 (4.20)
I.154	CDCl <sub>3</sub>	400	7.9321 (3.21); 7.9306 (3.49); 7.9277 (4.31); 7.9138 (2.91); 7.9104 (4.29); 7.9079 (3.46); 7.4470 (1.49); 7.4285 (2.12); 7.4105 (1.48); 7.4067 (1.81); 7.3928 (4.01); 7.3889 (3.16); 7.3728 (5.59); 7.3685 (6.17); 7.3533 (3.20); 7.3500 (3.47); 7.3327 (7.77); 7.3227 (3.84); 7.3082 (3.64); 7.3017 (3.99); 7.2872 (3.80); 7.2591 (10.07); 7.1406 (2.74); 7.1340 (3.30); 7.1160 (2.80); 7.1094 (3.29); 7.0878 (2.22); 7.0810 (1.77); 7.0671 (3.90); 7.0604 (3.07); 7.0464 (1.85); 7.0396 (1.47); 6.9365 (1.10); 6.7999 (2.23); 6.6633 (1.12); 5.2933 (16.00); 4.5669 (1.92); 4.1288 (0.50); 4.1109 (0.51); 3.7747 (12.55); 2.6434 (0.36); 2.6333 (0.79); 2.6271 (1.31); 2.6163 (1.68); 2.6100 (1.75); 2.6005 (1.31); 2.5941 (0.83); 2.5839 (0.39); 2.0426 (2.33); 1.6176 (5.76); 1.2754 (0.74); 1.2576 (1.52); 1.2397 (0.65); 0.8811 (0.47); 0.4311 (2.13); 0.3680 (3.90); 0.0741 (0.44); -0.0002 (6.29)
I.155	CDCl <sub>3</sub>	300	7.9329 (3.16); 7.9088 (3.36); 7.4510 (2.03); 7.4043 (8.79); 7.3742 (7.48); 7.3410 (10.73); 7.2901 (4.99); 7.2610 (12.31); 6.9804 (1.32); 6.7982 (2.64); 6.6158 (1.33); 5.2997 (11.30); 4.5684 (4.05); 3.7772 (16.00); 2.5911 (2.32); 1.5691 (4.90); 1.2608 (0.45); 0.4239 (3.77); 0.3606 (6.17); -0.0002 (6.94)
I.156	CDCl <sub>3</sub>	300	7.9248 (3.74); 7.9010 (4.01); 7.4247 (11.02); 7.3525 (14.27); 7.3426 (14.55); 7.2566 (2.64); 6.9995 (1.19); 6.8172 (2.40); 6.6349 (1.20); 5.2885 (6.47); 4.6079 (2.39); 3.7545 (16.00); 2.5474 (2.66); 1.6704 (2.32); 1.2580 (0.40); 0.3472 (7.80); -0.0002 (1.73)
I.157	CDCl <sub>3</sub>	300	7.9785 (3.29); 7.9681 (2.41); 7.9585 (2.50); 7.9484 (3.93); 7.9371 (0.69); 7.5942 (1.40); 7.5904 (1.41); 7.5633 (2.83); 7.5102 (7.42); 7.4843 (2.31); 7.4499 (0.74); 7.4234 (9.67); 7.4144 (3.98); 7.4015 (3.35); 7.3959 (2.23); 7.3782 (0.64); 7.3726 (0.46); 7.3063 (2.16); 7.2520 (0.36); 7.1274 (1.30); 6.9455 (2.66); 6.9327 (0.39); 6.7637 (1.33); 4.9117 (5.08); 4.7923 (0.47); 3.8542 (0.98); 3.8231 (12.97); 3.3144 (0.59); 3.2928 (0.79); 3.2711 (0.64); 2.7957 (0.81); 2.0340 (6.21); 1.3683 (16.00); 1.3456 (15.90); 1.2784 (1.53); 1.2558 (1.51); 1.2366 (0.34); 0.7522 (5.02); 0.7377 (7.48); 0.0545 (1.14)
I.158	CDCl <sub>3</sub>	300	7.9375 (1.52); 7.9356 (1.60); 7.9301 (2.11); 7.9157 (1.24); 7.9096 (1.84); 7.9073 (1.93); 7.9052 (1.80); 7.4868 (0.82); 7.4600 (1.28); 7.4436 (1.59); 7.4243 (1.82); 7.4150 (1.89); 7.4032 (2.38); 7.3965 (3.25); 7.3882 (2.66); 7.3805 (4.54); 7.3713 (2.64); 7.3645 (5.80); 7.3584 (2.55); 7.3397 (0.73); 7.3347 (0.52); 7.2608 (5.49); 7.1596 (0.99); 7.1504 (1.28); 7.1317 (1.65); 7.1224 (2.21); 7.1034 (0.84); 7.0941 (1.35); 7.0884 (2.17); 7.0794 (1.58); 7.0578 (2.12); 7.0488 (1.56); 6.9794 (0.78); 6.7973 (1.56); 6.6152 (0.79); 5.2959 (16.00); 4.5649 (1.00); 3.7557 (8.16); 2.5162 (0.40); 2.5080 (0.72); 2.4940 (0.91); 2.4852 (0.95); 2.4721 (0.77); 2.4636 (0.48); 1.6217 (5.54); 0.3641 (1.41); 0.3195 (2.67); -0.0002 (3.62)
I.159	CDCl <sub>3</sub>	300	7.9377 (3.05); 7.9320 (3.74); 7.9177 (2.32); 7.9114 (3.47); 7.9094 (3.49); 7.4755 (1.53); 7.4496 (2.42); 7.4252 (2.97); 7.4191 (2.37); 7.4064 (4.68); 7.3982 (8.73); 7.3914 (13.19); 7.3833 (11.89); 7.3797 (11.56); 7.3649 (11.08); 7.3552 (9.55); 7.3484 (6.43); 7.2609 (20.97); 6.9768 (1.44); 6.7947 (2.99); 6.6127 (1.51); 5.2988 (0.47); 4.5533 (1.31); 3.7638 (16.00); 2.5562 (0.33); 2.5341 (1.39); 2.5200 (1.75); 2.5112 (1.82); 2.4984 (1.47); 2.4899 (0.93); 2.4759 (0.45); 1.5833 (21.55); 1.2571 (0.40); 0.3807 (2.71); 0.3310 (5.05); 0.0106 (0.43); -0.0002 (13.92); -0.0111 (0.69)
I.160	DMSO-d <sub>6</sub>	400	8.0787 (2.13); 8.0607 (2.28); 7.9593 (1.58); 7.7145 (3.36); 7.4447 (0.55); 7.4340 (1.38); 7.4235 (1.70); 7.4151 (1.51); 7.4039 (3.47); 7.3969 (4.93); 7.2299 (9.18); 7.2117 (0.36); 7.1808 (3.57); 7.0563 (0.72); 6.9218 (1.50); 6.7869 (0.75); 4.4654 (4.37); 3.7617 (5.90); 3.3346 (74.64); 2.8969 (10.41); 2.7377 (9.49); 2.5127 (38.43); 2.5085 (48.85); 2.5045 (37.23); 2.3838 (16.00); 2.3353 (0.32); 0.4321 (1.76); 0.4182 (1.81); 0.2546 (2.81)
I.161	DMSO-d <sub>6</sub>	400	8.0410 (1.34); 8.0353 (1.74); 8.0258 (1.01); 8.0197 (2.43); 8.0165 (1.68); 7.9597 (1.84); 7.8345 (3.54); 7.7045 (7.06); 7.4431 (0.54); 7.4311 (0.98); 7.4217 (1.92); 7.4035 (7.29); 7.3912 (1.89); 7.2380 (0.49); 7.2166 (7.53); 7.1902 (3.56); 7.0769

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(1.99); 6.9409 (1.01); 4.4927 (1.12); 3.8407 (14.62); 3.3316 (52.49); 2.8972 (14.45); 2.7382 (11.79); 2.6776 (0.42); 2.5802 (0.85); 2.5724 (1.11); 2.5635 (0.95); 2.5552 (0.75); 2.5449 (0.57); 2.5308 (1.44); 2.5176 (25.47); 2.5131 (51.33); 2.5085 (67.98); 2.5040 (49.58); 2.4995 (24.68); 2.3841 (16.00); 2.3398 (0.34); 2.3353 (0.47); 2.3308 (0.36); 0.4996 (2.24); 0.4827 (2.19); 0.3059 (2.16)
I.162	DMSO-d <sub>6</sub>	600	7.7433 (0.36); 7.2007 (0.48); 7.0188 (0.70); 3.6621 (0.76); 3.1064 (5.09); 2.6809 (3.13); 2.5220 (2.43); 2.5212 (2.47); 2.2978 (5.26); 2.2948 (11.57); 2.2917 (16.00); 2.2886 (11.24); 2.2856 (5.06); 2.1722 (3.18)
I.163	CDCl <sub>3</sub>	300	7.9256 (1.48); 7.9236 (1.53); 7.9185 (1.97); 7.9029 (1.19); 7.8974 (1.79); 7.8955 (1.83); 7.8932 (1.69); 7.4839 (0.68); 7.4627 (0.89); 7.4027 (0.58); 7.3970 (0.76); 7.3791 (1.82); 7.3733 (1.69); 7.3613 (1.90); 7.3554 (3.03); 7.3480 (1.78); 7.3365 (1.47); 7.3315 (1.42); 7.3021 (2.70); 7.2589 (9.55); 7.2451 (1.70); 7.2193 (6.72); 7.1870 (2.59); 7.1814 (1.63); 7.1620 (0.99); 6.9770 (0.65); 6.7950 (1.26); 6.6129 (0.64); 4.6770 (0.36); 4.4140 (0.34); 3.7723 (5.26); 2.4196 (16.00); 2.2406 (0.67); 2.2299 (0.92); 2.2189 (0.94); 2.2076 (0.83); 2.0033 (0.62); 1.6073 (0.94); 0.7033 (0.64); 0.6018 (2.14); 0.4653 (0.70); 0.4518 (0.73); 0.4388 (0.58); 0.1057 (0.53); -0.0002 (6.04)
I.164	CDCl <sub>3</sub>	300	7.9394 (1.26); 7.9332 (1.49); 7.9160 (0.97); 7.9101 (1.72); 7.4988 (0.72); 7.4733 (1.01); 7.4104 (0.51); 7.4056 (0.55); 7.3872 (1.33); 7.3820 (1.45); 7.3636 (2.30); 7.3559 (2.29); 7.3441 (2.68); 7.3092 (2.99); 7.2605 (14.68); 7.2467 (1.32); 7.2207 (2.32); 7.1844 (1.95); 7.1592 (0.90); 6.9002 (0.57); 6.7184 (1.15); 6.5361 (0.60); 4.7742 (0.37); 3.8785 (7.09); 2.4205 (12.55); 2.2987 (0.48); 2.2867 (0.85); 2.2753 (0.94); 2.2632 (0.89); 2.2514 (0.53); 2.0068 (16.00); 1.5731 (3.02); 0.6916 (0.50); 0.6835 (0.51); 0.6724 (0.56); 0.4912 (3.66); 0.4712 (3.12); 0.4210 (0.61); 0.4065 (0.60); 0.0106 (0.79); -0.0002 (9.62)
I.165	CDCl <sub>3</sub>	300	7.8717 (1.32); 7.8657 (1.32); 7.8522 (1.02); 7.8454 (1.47); 7.8429 (1.54); 7.5421 (0.76); 7.5180 (0.95); 7.4167 (0.45); 7.4111 (0.69); 7.3934 (1.89); 7.3871 (1.84); 7.3808 (2.08); 7.3718 (3.94); 7.3620 (2.02); 7.3556 (2.48); 7.3494 (4.90); 7.3325 (0.77); 7.3274 (0.51); 7.2612 (6.54); 7.2395 (5.78); 7.2241 (3.13); 7.1980 (0.78); 7.1550 (0.86); 6.9724 (1.69); 6.7899 (0.85); 6.4613 (0.68); 3.7537 (14.83); 2.4376 (16.00); 2.3393 (0.39); 2.0017 (2.75); 1.8652 (0.61); 0.7217 (0.56); 0.6162 (2.43); 0.4951 (0.52); 0.1533 (0.60); -0.0002 (3.65)
I.166	CDCl <sub>3</sub>	400	7.9225 (1.94); 7.9027 (2.19); 7.4548 (0.45); 7.3963 (0.84); 7.3929 (0.81); 7.3759 (1.91); 7.3592 (1.51); 7.3557 (1.71); 7.3451 (1.40); 7.3268 (1.76); 7.3088 (1.30); 7.2586 (12.00); 7.2371 (0.88); 7.2183 (1.39); 7.1835 (2.10); 7.1648 (1.04); 6.8399 (0.33); 5.2959 (12.80); 3.7531 (2.34); 2.5544 (0.82); 2.5428 (0.83); 2.4096 (16.00); 1.5651 (13.89); 0.5580 (1.39); 0.0080 (0.45); -0.0002 (11.36); -0.0085 (0.43); -0.0942 (0.54)
I.167	CDCl <sub>3</sub>	300	7.9363 (2.42); 7.9335 (2.89); 7.9115 (2.20); 7.9073 (3.67); 7.4867 (2.62); 7.4676 (1.38); 7.4074 (1.05); 7.4002 (1.07); 7.3799 (2.52); 7.3532 (3.03); 7.3256 (1.70); 7.3080 (3.67); 7.2589 (8.12); 7.2387 (2.33); 7.2198 (1.64); 7.1971 (2.24); 7.1724 (2.97); 6.8843 (0.80); 6.7020 (1.64); 6.5198 (0.84); 5.2941 (13.42); 5.2270 (0.49); 5.1776 (0.45); 4.5250 (0.37); 4.4721 (0.57); 4.2835 (0.37); 4.2384 (0.37); 3.8593 (7.24); 3.7857 (1.68); 2.6111 (0.47); 2.5936 (0.48); 2.5188 (0.75); 2.5027 (0.74); 2.4778 (0.46); 2.4088 (16.00); 1.6160 (5.38); 1.2579 (0.55); 0.8568 (1.81); 0.8403 (1.57); 0.4602 (3.71); 0.0106 (0.40); -0.0002 (6.53); -0.0111 (0.47); -0.1467 (0.94); -0.1614 (0.92)
I.168	CDCl <sub>3</sub>	400	8.0183 (1.43); 7.4242 (2.02); 7.4177 (1.49); 7.4057 (3.24); 7.3984 (3.48); 7.3941 (2.08); 7.3890 (1.23); 7.3790 (2.61); 7.3750 (1.64); 7.3651 (0.49); 7.3401 (2.44); 7.3376 (2.43); 7.3211 (2.75); 7.3189 (2.97); 7.3106 (0.51); 7.3078 (0.53); 7.3028 (1.06); 7.2919 (0.46); 7.2894 (0.47); 7.2673 (16.08); 7.1456 (0.50); 7.1296 (2.98); 7.1237 (2.13); 7.1110 (2.53); 6.9791 (1.59); 6.8426 (3.50); 6.8329 (12.58); 6.7062 (1.66); 5.3028 (10.12); 4.6495 (5.18); 3.7978 (16.00); 2.9604 (11.16); 2.8861 (9.89); 2.6992 (1.12); 1.6348 (3.62); 1.2541 (0.48); 0.6615 (2.18); 0.6445 (2.61); 0.6089 (1.79); 0.5983 (4.10); 0.5881 (3.45); 0.0078 (0.38); -0.0002 (9.35); -0.0084

Example	Solvent	Frequency (Mhz)	<sup>1</sup> H-NMR
			(0.38)

The following examples illustrate in a non-limiting manner the preparation and efficacy of the compounds of formula (I) according to the invention.

- 5 Preparation example 1 : preparation of N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[2-(3-methyloxetan-3-yl)benzyl]-1 H-pyrazole-4-carboxamide (compound 1.15)

Step 1 : preparation of N-[2-(3-methyloxetan-3-yl)benzyl]cyclopropanamine

To a cooled solution of 1.94 g (34 mmol) of cyclopropylamine and 2.44 mL (42 mmol) of acetic acid, together with 9 g of 4A molecular sieves, in 50 mL of methanol, are added 3 g (17 mmol) of 2-(3-methyloxetan-3-yl)benzaldehyde. The reaction mixture is stirred for 3 hrs at reflux. The reaction mixture is then cooled to 0 °C and 1.6 g (25 mmol) of sodium cyanoborohydride are slowly added and the reaction mixture is further stirred for 1 hr at reflux. The cooled reaction mixture is then filtered over a cake of diatomaceous earth. The cake is washed twice by 80 mL of methanol and the combined methanolic extracts are concentrated under vacuum. 100 mL of water are then added to the residue and the pH is adjusted to 12 with a 1 N aqueous solution of sodium hydroxide. The watery layer is extracted with 2 x 50 mL of ethyl acetate. The organic layer is washed twice by a concentrated aqueous solution of NaCl and dried over magnesium sulfate to yield after concentration 4.37 g of a yellow oil. Column chromatography on silica gel (gradient heptane/ethyl acetate) yields 2.44 g (62% yield) of N-[2-(3-methyloxetan-3-yl)benzyl]cyclopropanamine as a colorless oil. (M+H) = 218.

Step 2 : preparation of N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[2-(3-methyloxetan-3-yl)benzyl]-1 H-pyrazole-4-carboxamide

At ambient temperature, a solution of 195 mg (0.92 mmol) of 3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carbonyl chloride in 2 mL of dry tetrahydrofuran is added to a mixture of 200 mg (0.92 mmol) of N-[2-(3-methyloxetan-3-yl)benzyl]cyclopropanamine and 0.141 mL (1.01 mmol) of triethylamine in 3 mL of dry tetrahydrofuran. The reaction mixture is stirred at ambient temperature for four days. The reaction mixture is then quenched by 0.1 mL of a 1 N solution of HCl in diethylether and solids are filtered off. The organic phase is concentrated under vacuum to yield 400 mg of a crude residue. Column chromatography on silica gel (gradient n-heptane/ethyl acetate) yields 314 mg (82% yield) of N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[2-(3-methyloxetan-3-yl)benzyl]-1 H-pyrazole-4-carboxamide as an oil. LogP = 1.94. (M+H) = 394.

- 35 Preparation example 2 : preparation of N-[5-chloro-2-(2-methyloxiran-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide (compound 1.4)

Step 1 : preparation of N-[5-chloro-2-(prop-1-en-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide

To a suspension of 1.55 g (3.55 mmol) of N-(2-bromo-5-chlorobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide in a 20/1 mixture of toluene / water degassed with argon, are successively added 753 mg (4.25 mmol) of 4,4,5,5-tetramethyl-2-(prop-1-en-2-yl)-1,3,2-dioxaborolane, 205 mg (0.71 mmol) of tricyclohexylphosphine, 167 mg (0.355 mmol) of palladium (II) acetate and 4.65 g (21.3 mmol) of tri-potassium phosphate. The mixture is heated at 100 °C for 3 hrs. The reaction mixture is cooled to ambient temperature. 60 mL of water are added and the reaction mixture is extracted 3 times by ethyl acetate. The combined organic extracts are washed by a saturated aqueous solution of NaCl and dried over a phase-separator filter. The organic phase is concentrated under vacuum and the crude residue is purified by column chromatography on silica gel (gradient n-heptane/ethyl acetate) to yield 1.01 g (72% yield) of a 68/32 mixture of N-[5-chloro-2-(prop-1-en-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide and N-cyclopropyl-3-(difluoromethyl)-N-[2,5-di(prop-1-en-2-yl)benzyl]-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide used as such in the next step. LogP = 3.96. (M+H) = 398.

Step 2 : preparation of N-[5-chloro-2-(2-methyloxiran-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide

To a cooled solution of 0.5 g (1.26 mmol - 68% purity) of N-[5-chloro-2-(prop-1-en-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide in 20 mL of dichloromethane, are slowly added 1.65 g (6.69 mmol) of 70% wt. m-chloroperbenzoic acid. The reaction mixture is stirred at ambient temperature for 2 hrs. The excess of oxidant is neutralized by addition of an aqueous solution of 2.47 g (12.6 mmol) of sodium hydrosulfite and the mixture is then extracted 3 times by dichloromethane. The combined organic extracts are washed by an aqueous solution of sodium hydrogenocarbonate and dried over a phase-separator filter. The organic phase is concentrated under vacuum and the crude residue is purified by column chromatography on silica gel (gradient n-heptane/ethyl acetate) to yield 183 mg (33% yield) of N-[5-chloro-2-(2-methyloxiran-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide. LogP = 3.19. (M+H) = 414.

Preparation example 3 : preparation of N-[5-(1-benzothiophen-3-yl)-2-isopropylbenzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1H-pyrazole-4-carboxamide (compound 1.157)

Step 1 : preparation of 1-(5-chloro-2-isopropylphenyl)-N-hydroxymethanimine

To a solution of 1 g (5.47 mmol) of 5-chloro-2-isopropylbenzaldehyde in 12 mL of methanol are successively added 0.5 g (6 mmol) of sodium acetate and 419 mg (6 mmol) of hydroxylamine hydrochloride. The reaction mixture is stirred at ambient temperature for 6 hrs. The solvent is removed under vacuum and the residue is extracted by dichloromethane. The organic phase is filtered over a ChemElut™ cartridge and concentrated under vacuum to yield 1.08 g (89% yield) of 1-(5-chloro-2-isopropylphenyl)-N-hydroxymethanimine as a 8/92 mixture of two isomers. LogP = 2.90 (minor isomer) and 3.17 (major isomer). (M+H) = 198.

Step 2 : preparation of 1-(5-chloro-2-isopropylphenyl)methanamine

To a cooled solution of 1.08 g (5.46 mmol) of 1-(5-chloro-2-isopropylphenyl)-N-hydroxymethanimine in 18 mL of tetrahydrofuran are slowly added 21 mL (21 mmol) of a 1 M solution of the

borane:tetrahydrofuran complex in tetrahydrofuran. The reaction mixture is heated at reflux for 1 day. The excess of hydride is destroyed by addition of ethanol. A few mL of a 1 N aqueous solution of sodium hydroxide are added and the reaction mixture is further heated at 70 °C for 30 min. The reaction mixture is diluted by water, extracted by tertbutylmethylether. The organic phase is filtered over a ChemElut™  
5 cartridge and concentrated under vacuum to yield 1.30 g of 1-(5-chloro-2-isopropylphenyl)methanamine used as such in the next step. LogP = 1.20. (M+H) = 184.

Step 3 : preparation of N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide (compound Xa.1)

10 At ambient temperature, 824 mg (3.87 mmol) of 3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carbonyl chloride are added to a mixture of 647 mg (3.52 mmol) of 1-(5-chloro-2-isopropylphenyl)methanamine and 0.590 mL (4.23 mmol) of triethylamine in 7 mL of dry tetrahydrofuran. The reaction mixture is heated at reflux for 2 hrs. A few mL of water are added and the reaction mixture is filtered over a ChemElut™ cartridge and concentrated to give 1.5 g of crude material as a yellow oil. The  
15 residue is purified by preparative HPLC (gradient acetonitrile/water + 0.1% HCO<sub>2</sub>H) to yield 360 mg (27% yield) of N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide as a white solid. LogP = 3.46. (M+H) = 360.

Step 4 : preparation of N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1 -methyl-  
20 1H-pyrazole-4-carboxamide

In a dried Radleys™ vial, are added 63 mg (0.19 mmol) of N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide, 35 mg (0.19 mmol) of copper(II) acetate and 31 mg (0.096 mmol) of caesium carbonate. The vial is flushed with dry air and vigorously stirred for a few minutes. 0.5 mL of toluene is added together with 46 mg (0.58 mmol) of pyridine and 65 mg (0.38 mmol)  
25 of 2-cyclopropyl-4,4,5,5-tetramethyl-1,3,2-dioxaborolane. The vial is flushed again with dry air, sealed and heated at 100 °C for 1 day. HPLC-mass analysis of the reaction shows the formation of 4% (7% cv) of N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide. LogP = 4.06. (M+H) = 400.

30 Step 5 : preparation of N-[5-(1-benzothiophen-3-yl)-2-isopropylbenzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide

In a 5 mL microwave vial, 200 mg (0.5 mmol) of N-(5-chloro-2-isopropylbenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide are dissolved in 1 mL of acetonitrile. The vial is degassed with argon and 104 mg (0.75 mmol) of potassium carbonate dissolved in 1 mL of water are  
35 further added, followed by 106 mg (0.6 mmol) of 1-benzothiophen-3-ylboronic acid and 6.5 mg (0.01 mmol) of 1,1'-Bis(di-ter-butylphosphino)(ferrocene)dichloropalladium(II). The microwave apparatus [Biotage Initiator™] is then heated at 130 °C for 20 minutes. The reaction mixture is cooled to ambient temperature, filtered, and extracted by ethyl acetate. The organic extract is dried over a ChemElut™ cartridge and concentrated under vacuum. The residue is purified by preparative HPLC (gradient acetonitrile/water + 0.1% HCO<sub>2</sub>H) to yield 75 mg (28% yield) of N-[5-(1-benzothiophen-3-yl)-2-isopropylbenzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1 -methyl-1 H-pyrazole-4-carboxamide as a brown oil.  
40 LogP = 5.17. (M+H) = 498.

Preparation example 4 : preparation of N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[2-(4-methyl-2-thienyl)benzyl]-1 H-pyrazole-4-carboxamide (compound 1.100)

In a Radleys™ vial, 200 mg (0.49 mmol) of N-(2-bromobenzyl)-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide are dissolved in 5 mL of dioxan together with 137 mg (0.99 mmol) of potassium carbonate dissolved in 0.5 mL of water. The vial is degassed with argon for 10 min. Are then added, 106 mg (0.74 mmol) of (4-methyl-2-thienyl)boronic acid and 115 mg (0.1 mmol) of tetrakis-(triphenylphosphine)palladium(O). The reaction mixture is heated at 100 °C for 8 hrs. The cooled mixture is filtered over a cake of diatomaceous earth. The filtrate is washed by an aqueous solution of potassium carbonate followed by a saturated aqueous solution of NaCl and dried over a ChemElut™ cartridge. The organic layer is concentrated to leave a residue that is purified by column chromatography on silica gel (gradient n-heptane/ethyl acetate) to yield 165 mg (71 % yield) of N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-N-[2-(4-methyl-2-thienyl)benzyl]-1 H-pyrazole-4-carboxamide as a colorless oil. LogP = 3.99. (M+H) = 420.

General preparation example 5 : thionation of amide of formula (I) on Chemspeed™ apparatus

In a 13 mL Chemspeed™ vial is weighted 0.27 mmol of phosphorous pentasulfide (P<sub>2</sub>S<sub>5</sub>). 3 mL of a 0.18 M solution of the amide (I) (0.54 mmol) in dioxane is added and the mixture is heated at reflux for two hours. The temperature is then cooled to 80 °C and 2.5 mL of water are added. The mixture is heated at 80 °C for one more hour. 2 mL of water are then added and the reaction mixture is extracted twice by 4 mL of dichloromethane. The organic phase is deposited on a basic alumina cartridge (2 g) and eluted twice by 8 mL of dichloromethane. The solvents are removed and the crude thioamide derivative is analyzed by LCMS and NMR. Insufficiently pure compounds are further purified by preparative LC.

Example A : *in vivo* preventive test on *Blumeria praminis* (barley)

Solvent: 49 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl/polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has been dried, the plants are dusted with spores of *Blumeria graminis f.sp. hordei*.

The plants are placed in the greenhouse at a temperature of approximately 18 °C and a relative atmospheric humidity of approximately 80% to promote the development of mildew pustules.

The test is evaluated 7 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, excellent (at least 90%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table A :

Table A:

Example	Efficacy
I.3	100
I.4	94
I.100	100
I.105	100
I.157	100

Example B : *in vivo* preventive test on *Alternaria brassicae* (leaf spot on radish)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

Radish plants ("Pernod Clair" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 17 °C, are treated at the cotyledon stage by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the cotyledons with an aqueous suspension of *Alternaria brassicae* spores (50 000 spores per mL). The spores are collected from a 15-day-old culture. The contaminated radish plants are incubated at 20 °C and at 100% relative humidity.

Grading (% of efficacy) is carried out 6 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table B :

Table B:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.28	96	I.101	83	I.132	97
I.29	82	I.102	83	I.136	70
I.50	96	I.104	83	I.137	70
I.54	83	I.106	100	I.139	73
I.68	71	I.129	97	I.140	90
I.72	70	I.130	96	I.154	75
I.74	83	I.131	91	I.157	100

Example C : *in vivo* preventive test on *Botrytis cinerea* (grey mould)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO and then diluted with water to obtain the desired active material concentration.

Gherkin plants ("Vert petit de Paris" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 24 °C, are treated at the Z11 cotyledon stage by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the cotyledons with an aqueous suspension of cryopreserved *Botrytis cinerea* spores (50 000 spores per mL). The spores are suspended in a nutrient solution composed of 10 g/L of PDB, 50 g/L of D-Fructose, 2 g/L of NH<sub>4</sub>NO<sub>3</sub> and 1 g/L of KH<sub>2</sub>PO<sub>4</sub>. The contaminated gherkin plants are incubated at 17 °C and at 90% relative humidity.

Grading (% of efficacy) is carried out 4 to 5 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table C :

5

Table C:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.29	100	I.101	99	I.138	100
I.49	99	I.102	94	I.139	100
I.50	99	I.103	98	I.141	90
I.54	97	I.104	93	I.143	98
I.66	91	I.106	80	I.144	98
I.73	100	I.122	100	I.146	100
I.74	100	I.123	100	I.156	97
I.75	100	I.129	92	I.157	100
I.78	78	I.131	100	I.158	100
I.100	80	I.132	99	I.159	100

Under the same conditions, good (at least 70%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table C2 :

10

Table C2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.31	81	I.59	81	I.125	78
I.32	94	I.82	97	I.145	100
I.34	99	I.84	100	I.150	90
I.35	94	I.108	100		

Example D : *in vivo* preventive test on *Leptosphaena nodorum* test (wheat)

Solvent: 49 parts by weight of N,N-dimethylacetamide

15 Emulsifier: 1 part by weight of alkylaryl/polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound or active compound combination is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

20 To test for preventive activity, young plants are sprayed with the preparation of active compound or active compound combination at the stated rate of application.

After the spray coating has been dried, the plants are sprayed with a spore suspension of *Leptosphaena nodorum*. The plants remain for 48 hours in an incubation cabinet at approximately 20 °C and a relative atmospheric humidity of approximately 100%.

25 The plants are placed in the greenhouse at a temperature of approximately 22 °C and a relative atmospheric humidity of approximately 80%.

The test is evaluated 8 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.

Under these conditions, high (at least 80%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table D:

Table D:

Example	Efficacy
I.3	92
I.4	80
I.100	83
I.157	100

5

Example E : in vivo preventive test on *Pyrenophora teres* (barley)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

Barley plants ("Plaisant" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 22 °C, are treated at the 1 leaf stage (10 cm height) by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the leaves with an aqueous suspension of *Pyrenophora teres* spores (12 000 spores per mL). The spores are collected from a 12-day-old culture. The contaminated barley plants are incubated for 48 hours at 20 °C and at 100% relative humidity, and then for 12 days at 20 °C at 70-80% relative humidity.

Grading (% of efficacy) is carried out 14 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table E :

Table E:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.3	88	I.100	95	I.130	79
I.4	96	I.101	95	I.131	93
I.5	93	I.103	95	I.132	88
I.27	93	I.104	70	I.136	86
I.28	79	I.105	97	I.137	79
I.29	93	I.106	93	I.138	79
I.47	71	I.114	79	I.139	100
I.50	100	I.119	71	I.143	79
I.67	79	I.120	79	I.154	71
I.73	71	I.121	86	I.157	93
I.86	79	I.129	86		

Under the same conditions, good (at least 70%) to excellent (at least 95%) protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table E2 :

25

Table E2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.7	97	I.83	86	I.134	71
I.34	81	I.84	81	I.144	79
I.36	71	I.93	88	I.159	75
I.57	79	I.95	88		

Under the same conditions, good (at least 70%) protection to high (at least 80%) protection is observed at a dose of 100 ppm of active ingredient with compounds of example I.84 and I.144, whereas no protection is observed with the pyridine analogue CMP1 disclosed in patent application WO-2007/087906 as in table E3:

Table E3:

Example	dose (ppm)	Efficacy
I.84 from this patent	100	81
I.144 from this patent	100	79
CMP1 from WO-2007/087906	100	0

CMP1 disclosed in international patent WO-2007/087906 corresponds to N-[2-chloro-4-(pyridin-3-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide. These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds disclosed in WO-2007/087906.

Example F : In vivo preventive test on *Pyricularia oryzae* (rice blast)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

Rice plants ("Koshihikari" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 26 °C, are treated at the 2 leaf stage (10 cm height) by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the leaves with an aqueous suspension of *Pyricularia oryzae* spores (40 000 spores per mL). The spores are collected from a 15-day-old culture and are suspended in water containing 2.5 g/L of gelatin. The contaminated rice plants are incubated at 25 °C and at 80% relative humidity.

Grading (% of efficacy) is carried out 6 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to high (at least 80%) protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table F :

Table F:

Example	Efficacy
1.106	80
1.119	70
1.132	80

Example G : *in vivo* preventive test on *Puccinia recondita* (brown rust on wheat)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

Wheat plants ("Scipion" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 22 °C, are treated at the 1 leaf stage (10 cm height) by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the leaves with an aqueous suspension of *Puccinia recondita* spores (100 000 spores per mL). The spores are collected from an infected plant and are suspended in water containing 2.5 mL/L of Tween 80 at 10%. The contaminated wheat plants are incubated for 24 hours at 20 °C and at 100% relative humidity, and then for 10 days at 20 °C and at 70-80% relative humidity.

Grading (% of efficacy) is carried out 12 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table G :

20 Table G:

Example	Efficacy	Example	Efficacy	Example	Efficacy
1.1	75	1.75	94	1.122	98
1.3	94	1.76	93	1.123	94
1.4	98	1.77	93	1.130	81
1.5	98	1.78	93	1.131	75
1.27	100	1.79	75	1.132	83
1.28	94	1.86	98	1.136	98
1.29	81	1.87	75	1.137	88
1.48	75	1.88	94	1.138	94
1.50	100	1.100	100	1.139	94
1.54	80	1.101	100	1.141	86
1.55	95	1.102	100	1.143	88
1.56	78	1.103	95	1.144	75
1.66	94	1.104	95	1.145	88
1.67	100	1.105	100	1.154	75
1.68	94	1.112	83	1.155	75
1.69	94	1.114	72	1.156	81
1.70	78	1.119	100	1.157	98
1.73	100	1.120	100	1.158	81
1.74	94	1.121	94	1.168	81

Under the same conditions, good (at least 70%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table G2 :

Table G2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.6	81	I.37	81	I.111	88
I.7	100	I.39	88	I.125	81
I.8	94	I.82	81	I.150	81
I.32	88	I.83	75		

5

Under the same conditions, average (at least 60%) to excellent (at least 90%) protection is observed at a dose of 100 ppm and 500 ppm of active ingredient with compounds of example 1.77, 1.78 or 1.79, whereas poor (less than 40%) to no protection is observed with the pyridine analogue CMP2 disclosed in patent application WO-201 0/1 30767 as in table G3:

10

Table G3:

Example	dose (ppm)	Efficacy
I.77 from this patent	500	93
	100	93
I.78 from this patent	500	93
	100	43
I.79 from this patent	500	75
	100	63
CMP2 from WO-2010/130767	500	38
	100	0

CMP2 disclosed in international patent WO-201 0/1 30767 corresponds to 5-chloro-N-[2-chloro-4-(pyridin-3-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide.

15 These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds disclosed in WO-201 0/1 30767.

Example H : *in vivo* preventive test on *Septoria tritici* (wheat)

20 The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

Wheat plants ("Scipion" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 22 °C, are treated at the 1 leaf stage (10 cm height) by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

25 After 24 hours, the plants are contaminated by spraying the leaves with an aqueous suspension of cryopreserved *Septoria tritici* spores (500 000 spores per mL). The contaminated wheat plants are incubated for 72 hours at 18 °C and at 100% relative humidity, and then for 21 days at 90% relative humidity.

Grading (% of efficacy) is carried out 24 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table H :

5

Table H:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.3	100	I.76	75	I.129	90
I.4	100	I.78	83	I.130	93
I.5	97	I.79	75	I.131	96
I.16	90	I.86	79	I.132	100
I.27	80	I.88	79	I.136	93
I.28	97	I.100	100	I.137	97
I.29	96	I.101	100	I.138	100
I.48	86	I.102	95	I.139	96
I.49	96	I.103	95	I.140	96
I.50	97	I.104	88	I.141	83
I.54	100	I.105	80	I.143	100
I.55	95	I.106	80	I.144	100
I.56	100	I.112	100	I.145	94
I.67	97	I.113	70	I.154	96
I.68	83	I.114	100	I.155	86
I.69	97	I.119	90	I.156	96
I.70	86	I.120	80	I.157	90
I.71	90	I.121	80	I.158	98
I.72	80	I.122	100	I.159	88
I.75	86	I.123	94	I.168	80

Under the same conditions, good (at least 70%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table H2 :

10

Table H2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.6	92	I.82	97	I.125	96
I.7	100	I.84	71	I.134	79
I.39	80	I.93	86	I.150	80
I.58	97	I.108	86	I.160	71

Example I : *in vivo* preventive test on *Sphaerotheca fulipinea* (powdery mildew on cucurbits)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO and then diluted with water to obtain the desired active material concentration.

15

Gherkin plants ("Vert petit de Paris" variety), sown in starter cups on a 50/50 peat soil-pozzolana substrate and grown at 24 °C, are treated at the Z11 cotyledon stage by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

After 24 hours, the plants are contaminated by spraying the cotyledons with an aqueous suspension of *Sphaerotheca fuliginea* spores (100 000 spores per mL). The spores are collected from infected plants. The contaminated gherkin plants are incubated at about 20 °C and at 70-80% relative humidity.

Grading (% of efficacy) is carried out 12 days after the contamination, in comparison with the control plants.

Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table I :

Table I:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.1	93	I.74	100	I.130	100
I.3	100	I.75	90	I.131	98
I.4	100	I.86	98	I.132	100
I.5	100	I.100	100	I.136	100
I.27	100	I.101	100	I.137	100
I.47	94	I.102	100	I.138	100
I.49	78	I.103	100	I.139	100
I.50	100	I.104	100	I.140	90
I.54	95	I.105	100	I.141	95
I.55	100	I.106	100	I.143	100
I.56	98	I.112	100	I.144	100
I.66	95	I.113	98	I.145	94
I.67	100	I.114	94	I.154	98
I.68	94	I.119	100	I.156	95
I.69	100	I.120	100	I.157	100
I.70	85	I.121	100	I.158	100
I.71	98	I.122	100	I.159	100
I.72	90	I.123	100	I.168	83
I.73	100	I.129	100		

10

Under the same conditions, good (at least 70%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table I2 :

Table I2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.7	94	I.108	75	I.150	100
I.82	98	I.111	100	I.160	98
I.83	98	I.125	95		
I.95	98	I.134	90		

15

Under the same conditions, total protection is observed at a dose of 100 ppm of active ingredient with compound of example I.144, whereas no protection is observed with the pyridine analogue CMP1 disclosed in patent application WO-2007/087906 as in table I3:

Table I3:

Example	dose (ppm)	Efficacy
I.144 from this patent	100	100
CMP1 from WO-2007/087906	100	0

CMP1 disclosed in international patent WO-2007/087906 corresponds to N-[2-chloro-4-(pyridin-3-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide.

- 5 These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds disclosed in WO-2007/087906.

Under the same conditions, excellent (at least 90%) to total protection is observed at a dose of 500 ppm of active ingredient with compound of example I.139 (respectively I.140, I.141, I.144 and I.145) whereas  
 10 poor (less than 40%) protection or plant damages is observed with the pyridine analogue CMP3 (respectively CMP4, CMP5, CMP6 and CMP7) claimed in patent applications WO-2007/087906 or WO-2010/130767 as in table I4:

Table I4:

Example	dose (ppm)	Efficacy
I.139 from this invention	500	100
CMP3 from WO-2010/130767	500	33
I.140 from this invention	500	90
CMP4 from WO-2007/087906	500	33
I.141 from this invention	500	95
CMP5 from WO-2010/130767	500	22
I.144 from this invention	500	100
CMP6 from WO-2007/087906	500	0
I.145 from this invention	500	94
CMP7 from WO-2010/130767	500	0

15

CMP3 disclosed in international patent WO-2010/130767 corresponds to N-[2-(6-chloropyridin-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide,

CMP4 disclosed in international patent WO-2007/087906 corresponds to N-[2-(6-chloropyridin-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide,

- 20 CMP5 disclosed in international patent WO-2010/130767 corresponds to 5-chloro-N-[2-(6-chloropyridin-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide,

CMP6 disclosed in international patent WO-2007/087906 corresponds to N-[5-chloro-2-(6-chloropyridin-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide, and

CMP7 disclosed in international patent WO-2010/130767 corresponds to 5-chloro-N-[5-chloro-2-(6-chloropyridin-2-yl)benzyl]-N-cyclopropyl-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide.

These results show that the compounds according to the invention have a much better biological activity than the structurally closest compounds disclosed in WO-2007/087906.

5

Example J : *in vivo* preventive test on *Uromyces appendiculatus* (bean rust)

The active ingredients tested are prepared by homogenization in a mixture of acetone/tween/DMSO, and then diluted with water to obtain the desired active material concentration.

10 Bean plants ("Saxa" variety), sown in starter cups on a 50/50 peat soil pozzolana substrate and grown at 24 °C, are treated at the 2 leaf stage (9 cm height) by spraying with the active ingredient prepared as described above. Plants, used as controls, are treated with the mixture of acetone/tween/DMSO/water not containing the active material.

15 After 24 hours, the plants are contaminated by spraying the leaves with an aqueous suspension of *Uromyces appendiculatus* spores (150 000 spores per mL). The spores are collected from infected plants and are suspended in water containing 2.5 mL/L of Tween 80 at 10%. The contaminated bean plants are incubated for 24 hours at 20 °C and at 100% relative humidity, and then for 10 days at 20 °C and at 70-80% relative humidity.

Grading (% of efficacy) is carried out 11 days after the contamination, in comparison with the control plants.

20 Under these conditions, good (at least 70%) to total protection is observed at a dose of 500 ppm of active ingredient with the following compounds from table J :

Table J:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.1	83	I.72	83	I.122	100
I.3	100	I.73	100	I.123	99
I.4	100	I.74	100	I.129	96
I.27	100	I.75	100	I.130	85
I.28	99	I.76	96	I.132	77
I.29	100	I.77	84	I.136	100
I.48	96	I.78	85	I.138	100
I.49	100	I.79	100	I.139	100
I.50	99	I.86	97	I.140	73
I.54	97	I.87	84	I.144	94
I.55	99	I.100	100	I.145	100
I.56	85	I.101	100	I.155	99
I.66	100	I.102	100	I.157	100
I.67	100	I.103	94	I.158	91
I.68	100	I.104	100	I.159	84
I.69	100	I.105	100	I.168	99
I.70	96	I.119	100		
I.71	81	I.120	100		

25 Under the same conditions, good (at least 70%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table J2 :

Table J2:

Example	Efficacy	Example	Efficacy	Example	Efficacy
I.6	73	I.37	83	I.143	93
I.7	100	I.83	80	I.160	77
I.8	95	I.111	100		
I.32	74	I.134	77		

Example K : *in vivo* preventive test on *Venturis inaequalis* (apples)

- 5 Solvent: 24.5 parts by weight of acetone  
 24.5 parts by weight of N,N-dimethylacetamide  
 Emulsifier: 1 part by weight of alkylarylpolyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for preventive activity, young plants are sprayed with the preparation of active compound at the stated rate of application. After the spray coating has dried on, the plants are inoculated with an aqueous conidia suspension of the causal agent of apple scab (*Venturis inaequalis*) and then remain for 1 day in  
 15 an incubation cabinet at approximately 20 °C and a relative atmospheric humidity of 100%.

The plants are then placed in a greenhouse at approximately 21 °C and a relative atmospheric humidity of approximately 90%.

The test is evaluated 10 days after the inoculation. 0% means an efficacy which corresponds to that of the untreated control, while an efficacy of 100% means that no disease is observed.


20 Under these conditions, excellent (at least 90%) to total protection is observed at a dose of 100 ppm of active ingredient with the following compounds from table K :

Table K:

Example	Efficacy
I.3	96
I.4	100
I. 139	100
I. 157	100

## CLAIMS

1. A compound of formula (I)



5

(I)

wherein

- A represents a carbo-linked, unsaturated or partially saturated, 5-membered heterocyclyl group that can be substituted by up to four groups R that can be the same or different ;
- T represents O or S ;
- n represents 0, 1, 2, 3 or 4 ;
- L represents a direct bond, CZ<sup>4</sup>Z<sup>5</sup>, O, S, SO, SO<sub>2</sub> or NZ<sup>6</sup> ; B represents a saturated, partially saturated or unsaturated, monocyclic or fused bicyclic 3-, 4-, 5-, 6-, 7-, 8-, 9-, or 10-membered ring comprising from 1 up to 4 heteroatoms selected in the list consisting of N, O, S, that can be substituted by up to 6 groups Y which can be the same or different ; with the proviso that B does not represent a pyridyl ring when L represents a direct bond or an oxygen atom
- Z<sup>1</sup> represents a non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl or a C<sub>3</sub>-C<sub>7</sub>-cycloalkyl substituted by up to 10 atoms or groups that can be the same or different and that can be selected in the list consisting of halogen atoms, cyano, CrC<sub>8</sub>-alkyl, CrC<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different, CrC<sub>8</sub>-alkoxy, CrC<sub>8</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, CrC<sub>8</sub>-alkoxycarbonyl, CrC<sub>8</sub>-halogenoalkoxycarbonyl comprising up to 9 halogen atoms that can be the same or different, Ci-C<sub>8</sub>-alkylaminocarbonyl or di-Ci-C<sub>8</sub>-alkylaminocarbonyl ;
- Z<sup>2</sup> and Z<sup>3</sup>, which can be the same or different, represent a hydrogen atom ; substituted or non-substituted Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyl ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyl ; cyano ; isonitrile ; nitro ; a halogen atom ; substituted or non-substituted CrC<sub>8</sub>-alkoxy ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyloxy ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfanyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfinyl ; amino ; substituted or non-substituted CrC<sub>8</sub>-alkylamino ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxycarbonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbamoyl ; substituted or non-substituted di-CrC<sub>8</sub>-alkylcarbamoyl ; or substituted or non-substituted N-CrC<sub>8</sub>-alkyl-Ci-C<sub>8</sub>-alkoxy-carbamoyl ; or

30

- Z<sup>2</sup> and Z<sup>3</sup> together with the carbon atom to which they are linked can form a substituted or non-substituted C<sub>3</sub>-C<sub>7</sub> cycloalkyl ; or
- Z<sup>3</sup> and the substituent X vicinal to the point of attachment of the phenyl ring, together with the consecutive carbon atoms to which they are linked, can form a substituted or non-substituted 5-,  
5 6- or 7-membered, partly saturated, carbo- or heterocycle comprising up to 3 heteroatoms and Z<sup>2</sup> is as herein-described ; Z<sup>4</sup> and Z<sup>5</sup> independently represent a hydrogen atom ; a halogen atom ; cyano ; substituted or non-substituted CrC<sub>8</sub>-alkyl ; CrC<sub>8</sub>-halogenoalkyl having 1 to 5 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkoxy; substituted or non-substituted CrC<sub>8</sub>-alkylsulfanyl ; or substituted or non-substituted CrC<sub>8</sub>-alkoxycarbonyl ; or
- 10 • Z<sup>4</sup> and Z<sup>5</sup> together with the carbon atom to which they are linked can form a C(=O) carbonyl group ;
- Z<sup>6</sup> represents a hydrogen atom ; a substituted or non-substituted CrC<sub>8</sub>-alkyl ; a CrC<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted **C2**-C<sub>8</sub>-alkenyl ; **C2**-C<sub>8</sub>-halogenoalkenyl comprising up to 9 halogen atoms that can  
15 be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>8</sub>-alkynyl ; C<sub>3</sub>-C<sub>8</sub>-halogenoalkynyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; C<sub>3</sub>-C<sub>7</sub>-halogeno-cycloalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl-Ci-C<sub>8</sub>-alkyl ; formyl ; substituted or non-substituted CrC<sub>8</sub>-alkylcarbonyl ; CrC<sub>8</sub>-halogenoalkylcarbonyl comprising up  
20 to 9 halogen atoms that can be the same or different ; substituted or non-substituted Ci-C<sub>8</sub>-alkoxycarbonyl ; CrC<sub>8</sub>-halogenoalkoxycarbonyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfonyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylsulfonyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted benzyl ; or substituted or non-substituted phenylsulfonyl ; X  
25 independently represents a halogen atom ; nitro ; cyano ; isonitrile ; hydroxy ; amino ; sulfanyl ; pentafluoro-A<sup>6</sup>-sulfanyl ; formyl ; formyloxy ; formylamino ; substituted or non-substituted (hydroxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (C<sub>1</sub>-**C8**-alkoxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (**C2-C8**-alkenyloxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (C<sub>2</sub>-C<sub>8</sub>-alkynyloxyimino)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted (benzyloxyimino)-Ci-C<sub>8</sub>-alkyl ;  
30 carboxy ; carbamoyl ; N-hydroxycarbamoyl ; carbamate ; substituted or non-substituted CrC<sub>8</sub>-alkyl ; CrC<sub>8</sub>-halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyl ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkynyl ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkynyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkoxy ; CrC<sub>8</sub>-halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted  
35 C<sub>1</sub>-C<sub>8</sub>-alkylsulfanyl ; CrC<sub>8</sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfinyl ; Ci-C<sub>8</sub>-halogenoalkylsulfinyl having 1 to 9 halogen atoms ; substituted or non-substituted CrC<sub>8</sub>-alkylsulfonyl ; CrC<sub>8</sub>-halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted di-CrC<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>2</sub>-C<sub>8</sub>-alkenyloxy ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkenyloxy  
40 having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>3</sub>-C<sub>8</sub>-alkynyloxy ; C<sub>2</sub>-C<sub>8</sub>-halogenoalkynyloxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-


- cycloalkyl ; C<sub>3</sub>-C<sub>7</sub>-halogenocycloalkyl having 1 to 9 halogen atoms ; substituted or non-substituted (C<sub>3</sub>-C<sub>7</sub>-cycloalkyl)-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>4</sub>-C<sub>7</sub>-cycloalkenyl ; C<sub>4</sub>-C<sub>7</sub>-halogenocycloalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted (C<sub>3</sub>-C<sub>7</sub>-cycloalkyl)-C<sub>2</sub>-C<sub>8</sub>-alkenyl ; substituted or non-substituted (C<sub>3</sub>-C<sub>7</sub>-cycloalkyl)-C<sub>2</sub>-C<sub>8</sub>-alkynyl ; substituted or non-substituted tri(Ci-C<sub>8</sub>)alkylsilyl ; substituted or non-substituted tri(Ci-C<sub>8</sub>)alkylsilyl-Ci-C<sub>8</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyloxy ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylcarbonyloxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbonylamino ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylcarbonylamino having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxy carbonyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkoxy carbonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkyloxycarbonyloxy ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkyloxycarbonyloxy having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylcarbonyl ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylcarbonyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylaminocarbonyloxy ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylaminocarbonyloxy ; substituted or non-substituted N-(Ci-C<sub>8</sub>-alkyl)hydroxy carbonyl ; substituted or non-substituted Ci-C<sub>8</sub>-alkoxy carbonyl ; substituted or non-substituted N-(Ci-C<sub>8</sub>-alkyl)-Ci-C<sub>8</sub>-alkoxy carbonyl ; aryl that can be substituted by up to 6 groups Q which can be the same or different ; aryl-Ci-C<sub>8</sub>-alkyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl-C<sub>2</sub>-C<sub>8</sub>-alkenyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl-C<sub>2</sub>-C<sub>8</sub>-alkynyl that can be substituted by up to 6 groups Q which can be the same or different ; aryloxy that can be substituted by up to 6 groups Q which can be the same or different ; arylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; arylamino that can be substituted by up to 6 groups Q which can be the same or different ; aryl-Ci-C<sub>8</sub>-alkyloxy that can be substituted by up to 6 groups Q which can be the same or different ; aryl-CrC<sub>8</sub>-alkylsulfanyl that can be substituted by up to 6 groups Q which can be the same or different ; aryl-Ci-C<sub>8</sub>-alkylamino that can be substituted by up to 6 groups Q which can be the same or different ; pyridinyl which can be substituted by up to 4 groups Q ; or pyridinyloxy which can be substituted by up to 4 groups Q ;
- Y independently represents a halogen atom ; cyano ; hydroxy ; amino ; sulfanyl ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxy ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfanyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted Ci-C<sub>8</sub>-alkylsulfanyl ; Ci-C<sub>8</sub>-halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylsulfonyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylamino ; substituted or non-substituted di-Ci-C<sub>8</sub>-alkylamino ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkylcarbonyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkoxy carbonyl ; C<sub>1</sub>-C<sub>8</sub>-halogenoalkoxy carbonyl having 1 to 9 halogen atoms ; or aryl that can be substituted by up to 6 groups Q which can be the same or different ;
  - Q independently represents a halogen atom, cyano, nitro, substituted or non-substituted C<sub>1</sub>-C<sub>8</sub>-alkyl, C<sub>1</sub>-C<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different,

substituted or non-substituted  $\text{CrC}_8$ -alkoxy,  $\text{C}_1$ - $\text{C}_8$ -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different, substituted or non-substituted  $\text{Ci-C}_8$ -alkylsulfanyl,  $\text{Ci-C}_8$ -halogenoalkylsulfanyl comprising up to 9 halogen atoms that can be the same or different, substituted or non-substituted  $\text{tri}(\text{CrC}_8)$ alkylsilyl, substituted or non-substituted  $\text{tri}(\text{CrC}_8)$ alkylsilyl- $\text{C}_1$ - $\text{C}_8$ -alkyl, substituted or non-substituted  $(\text{Ci-C}_8\text{-alkoxyimino})\text{-Ci-C}_8$ -alkyl, or substituted or non-substituted  $(\text{benzyloxyimino})\text{-Ci-C}_8$ -alkyl ;

- R independently represents hydrogen atom ; halogen atom ; nitro ; cyano ; hydroxy ; amino ; sulfanyl ; pentafluoro- $\text{A}^6$ -sulfanyl ; substituted or non-substituted  $(\text{Ci-C}_8\text{-alkoxyimino})\text{-Ci-C}_8$ -alkyl ; substituted or non-substituted  $(\text{benzyloxyimino})\text{-Ci-C}_8$ -alkyl ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkyl ;  $\text{CrC}_8$ -halogenoalkyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{C}_2$ - $\text{C}_8$ -alkenyl ;  $\text{C}_2$ - $\text{C}_8$ -halogenoalkenyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{C}_2$ - $\text{C}_8$ -alkynyl ;  $\text{C}_2$ - $\text{C}_8$ -halogenoalkynyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkoxy ;  $\text{CrC}_8$ -halogenoalkoxy having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{Ci-C}_8$ -alkylsulfanyl ;  $\text{Ci-C}_8$ -halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylsulfanyl ;  $\text{CrC}_8$ -halogenoalkylsulfanyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylsulfonyl ;  $\text{CrC}_8$ -halogenoalkylsulfonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkylamino ; substituted or non-substituted  $\text{di-C}_1$ - $\text{C}_8$ -alkylamino ; substituted or non-substituted  $\text{C}_2$ - $\text{C}_8$ -alkenyloxy ; substituted or non-substituted  $\text{C}_3$ - $\text{C}_8$ -alkynyloxy ; substituted or non-substituted  $\text{C}_3$ - $\text{C}_7$ -cycloalkyl ;  $\text{C}_3$ - $\text{C}_7$ -halogenocycloalkyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{tri}(\text{Ci-C}_8)$ alkylsilyl ; substituted or non-substituted  $\text{C}_1$ - $\text{C}_8$ -alkylcarbonyl ;  $\text{CrC}_8$ -halogenoalkylcarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkoxycarbonyl ;  $\text{CrC}_8$ -halogenoalkoxycarbonyl having 1 to 9 halogen atoms ; substituted or non-substituted  $\text{CrC}_8$ -alkylcarbamoyl ; substituted or non-substituted  $\text{di-CrC}_8$ -alkylcarbamoyl ; phenoxy ; phenylsulfanyl ; phenylamino ; benzyloxy ; benzylsulfanyl ; or benzylamino ;

as well as its salts, N-oxides, metal complexes, metalloid complexes and optically active isomers or geometric isomers thereof.


2. A compound according to claim 1 wherein A is selected in the list consisting of:  
- a heterocycle of formula (A<sup>1</sup>)



wherein :

- $\text{R}^1$  to  $\text{R}^3$  that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted  $\text{Ci-Cs}$ -alkyl ;  $\text{Ci-Cs}$ -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted  $\text{CrC}_5$ -alkoxy or  $\text{C}_1$ - $\text{C}_5$ -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;


- a heterocycle of formula (A<sup>2</sup>)



wherein :

- 5 R<sup>4</sup> to R<sup>6</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

10 - a heterocycle of formula (A<sup>3</sup>)




wherein :

- 15 R<sup>7</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>8</sup> represents a hydrogen atom or a substituted or non-substituted d-Cs-alkyl ;


20 - a heterocycle of formula (A<sup>4</sup>)



wherein :

- 25 R<sup>9</sup> to R<sup>11</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted d-Cs-alkyl ; amino ; substituted or non-substituted d-Cs-alkoxy ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

30 - a heterocycle of formula (A<sup>5</sup>)




(A<sup>5</sup>)

wherein :

R<sup>12</sup> and R<sup>13</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; substituted or non-substituted CrC<sub>5</sub>-alkoxy ; amino ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>14</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; amino ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>6</sup>)

(A<sup>6</sup>)


wherein :

R<sup>15</sup> represents a hydrogen atom ; a halogen atom ; a cyano ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; substituted or non-substituted Ci-Cs-alkoxy ; C<sub>1</sub>-Cs-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>16</sup> and R<sup>18</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkoxycarbonyl ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>17</sup> represent a hydrogen atom or substituted or non-substituted CrC<sub>5</sub>-alkyl ;

- a heterocycle of formula (A<sup>7</sup>)

(A<sup>7</sup>)


wherein :

R<sup>19</sup> represents a hydrogen atom or a CrC<sub>5</sub>-alkyl

R<sup>20</sup> to R<sup>22</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted Ci-Cs-alkyl or Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

115


- a heterocycle of formula (A<sup>8</sup>)



wherein :

- 5 R<sup>23</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  
 R<sup>24</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different;


10 - a heterocycle of formula (A<sup>9</sup>)



wherein :

- 15 R<sup>25</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  
 R<sup>26</sup> represents a hydrogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>10</sup>)




20

wherein :

- R<sup>27</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;  
 25 R<sup>28</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(C<sub>1</sub>-C<sub>5</sub>-alkyl)amino ;

30 - a heterocycle of formula (A<sup>11</sup>)




wherein :

R<sup>29</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub> -alkyl ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>30</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(Ci-C<sub>5</sub>-alkyl)amino ;

- a heterocycle of formula (A<sup>12</sup>)



(A<sup>12</sup>)


wherein :

R<sup>31</sup> represents a hydrogen atom or a substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl

R<sup>32</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub> -alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>33</sup> represents a hydrogen atom ; a halogen atom ; a nitro ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub> -alkyl ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

- a heterocycle of formula (A<sup>13</sup>)



(A<sup>13</sup>)


wherein :

R<sup>34</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>s</sub>-C<sub>s</sub> -cycloalkyl ; C<sub>i</sub>-C<sub>s</sub> -halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkoxy ; substituted or non-substituted C<sub>2</sub>-C<sub>5</sub>-alkynyloxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>35</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkyl ; a cyano ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkoxy ; substituted or non-substituted C<sub>1</sub>C<sub>5</sub>-alkylsulfanyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>i</sub>-C<sub>s</sub> -halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylamino or substituted or non-substituted di(Ci-C<sub>5</sub>-alkyl)amino ;

R<sup>36</sup> represents a hydrogen atom or substituted or non-substituted CrC<sub>5</sub>-alkyl ;

-a heterocycle of formula (A<sup>14</sup>)

(A<sup>14</sup>)

5


wherein :

R<sup>37</sup> and R<sup>38</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted CrC<sub>5</sub>-alkoxy or a substituted or non-substituted CrC<sub>5</sub>-alkylsulfanyl ;

10

R<sup>39</sup> represents a hydrogen atom or substituted or non-substituted CrC<sub>5</sub>-alkyl ;

- a heterocycle of formula (A<sup>15</sup>)

(A<sup>15</sup>)


15

wherein :

R<sup>40</sup> and R<sup>41</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted Ci-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

20

- a heterocycle of formula (A<sup>16</sup>)


(A<sup>16</sup>)

wherein :

R<sup>42</sup> and R<sup>43</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted CrC<sub>5</sub>-alkyl ; Ci-Cs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or amino ;

25

-a heterocycle of formula (A<sup>17</sup>)

(A<sup>17</sup>)


30

wherein :

R<sup>44</sup> and R<sup>45</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

5

- a heterocycle of formula (A<sup>18</sup>)




wherein :

10 R<sup>47</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>46</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkylsulfanyl ;

15

- a heterocycle of formula (A<sup>19</sup>)




wherein :

20 R<sup>49</sup> and R<sup>48</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

25

- a heterocycle of formula (A<sup>20</sup>)




wherein :

30 R<sup>50</sup> and R<sup>51</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkoxy ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

119


-a heterocycle of formula (A<sup>21</sup>)



wherein :

- 5 R<sup>52</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.

-a heterocycle of formula (A<sup>22</sup>)




10

wherein :

R<sup>53</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different.

15 -a heterocycle of formula (A<sup>23</sup>)




wherein :

- 20 R<sup>54</sup> and R<sup>56</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl or C<sub>1</sub>-C<sub>5</sub>-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>55</sup> represents a hydrogen atom or substituted or non-substituted C<sub>1</sub>-C<sub>5</sub>-alkyl ;

-a heterocycle of formula (A<sup>24</sup>)

25




wherein :

R<sup>57</sup> and R<sup>59</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkyl or CrCs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>58</sup> represents a hydrogen atom or substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl ;

5

-a heterocycle of formula (A<sup>25</sup>)


(A<sup>25</sup>)

wherein :

10 R<sup>60</sup> and R<sup>61</sup> that can be the same or different represent a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>i</sub>-C<sub>s</sub>-alkyl or CrCs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ;

R<sup>62</sup> represents a hydrogen atom or substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl ;

15 -a heterocycle of formula (A<sup>26</sup>)

(A<sup>26</sup>)

wherein :

20 R<sup>65</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl ; substituted or non-substituted C<sub>3</sub>-C<sub>5</sub>-cycloalkyl ; CrCs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; substituted or non-substituted d-Cs-alkoxy ; substituted or non-substituted **C2-C5**-alkynyloxy or C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ;

R<sup>63</sup> represents a hydrogen atom ; a halogen atom ; substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl ; a cyano ; substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkoxy ; substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkylsulfanyl ; **C1-C5**-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; C<sub>1</sub>-C<sub>5</sub>-halogenoalkoxy comprising up to 9 halogen atoms that can be the same or different ; amino ; substituted or non-substituted C<sub>1</sub>-C<sub>s</sub>-alkylamino or di(Ci-**C5**-alkyl)amino;


25 R<sup>64</sup> represents a hydrogen atom or substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl.

30 3. A compound according to claim 2 wherein A is selected in the list consisting of A<sup>2</sup> ; A<sup>5</sup> ; A<sup>6</sup> ; A<sup>10</sup> and A<sup>13</sup>.

4. A compound according to claim 3 wherein A represents A<sup>13</sup> wherein R<sup>34</sup> represents a substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl, CrCs-halogenoalkyl comprising up to 9 halogen atoms that can be the same or different ; or substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkoxy ; R<sup>35</sup> represents a hydrogen atom or a halogen atom and R<sup>36</sup> represents a substituted or non-substituted C<sub>r</sub>C<sub>5</sub>-alkyl.

35


5. A compound according to anyone of claims 1 to 4 wherein T represents O.
6. A compound according to anyone of claims 1 to 5 wherein  $Z^1$  represents a substituted or non-substituted cyclopropyl.
7. A compound according to claim 6 wherein  $Z^1$  represents a non-substituted cyclopropyl or a 2-CrC<sub>5</sub>-alkylcyclopropyl.
8. A compound according to anyone of claims 1 to 7 wherein  $Z^2$  and  $Z^3$  independently represent a hydrogen atom or a methyl.
9. A compound according to anyone of claims 1 to 8 wherein n represents 0, 1 or 2.
10. A compound according to anyone of claims 1 to 9 wherein L represents a direct bond, an oxygen atom, a methylene or a carbonyl group.
11. A compound according to anyone of claims 1 to 10 wherein B represents a substituted or non-substituted oxiranyl ring; a substituted or non-substituted pyrrolyl ring; a substituted or non-substituted pyrazolyl ring; a substituted or non-substituted thienyl ring; a substituted or non-substituted benzothienyl ring; a substituted or non-substituted quinoliny ring; a substituted or non-substituted isoquinoliny ring ; a substituted or non-substituted furan ring; or a substituted or non-substituted benzofuran ring.
12. A compound according to claim 11 wherein B represents a substituted or non-substituted thienyl ring or a substituted or non-substituted benzothienyl ring.
13. A compound according to anyone of claims 1 to 12 wherein X independently, represents a halogen atom ; substituted or non-substituted CrC<sub>8</sub>-alkyl ; CrC<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted C<sub>3</sub>-C<sub>7</sub>-cycloalkyl ; tri(Ci-Ce-alkyl)silyl; substituted or non-substituted CrC<sub>8</sub>-alkoxy; or substituted or non-substituted CrC<sub>8</sub>-alkylsulfanyl.
14. A compound according to anyone of claims 1 to 13 wherein Y independently, represents a halogen atom ; substituted or non-substituted CrC<sub>8</sub>-alkyl ; CrC<sub>8</sub>-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted Ci-C<sub>8</sub>-alkoxy ; Ci-C<sub>8</sub>-halogenoalkoxy comprising up to 9 halogen atoms which can be the same or different.
15. Process for preparing a compound of formula (I)



122


(I)

wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X and B are as defined in any one of claims 1 to 14 and L represents a direct bond or a methylene group, characterized in that a compound of formula (IV)



(IV)

- 5 wherein A, T, Z<sup>1</sup>, Z<sup>2</sup>, Z<sup>3</sup>, n, X are as defined above and U<sup>2</sup> represents a halogen atom or a triflate group, is reacted with a compound of formula (V)




(V)

wherein L and B are as defined above and W<sup>1</sup> represents a boron derivative.

10

16. Compound of formula (IIa.)



(IIa)

- wherein Z<sup>2</sup>, Z<sup>3</sup>, n and X are as defined in anyone of claims 1 to 14, and L is a direct bond and B<sup>1</sup> is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothienyl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring,

providing that (Ma.) does not represent :


- N-[2-(2-furyl)benzyl]cyclopropanamine
  - 20 - N-[2-(2-furyl)-4,5-dimethoxybenzyl]cyclopropanamine
  - N-[2-(3-bromo-2-furyl)-4,5-dimethoxybenzyl]cyclopropanamine
  - N-[2-(3-bromo-2-furyl)benzyl]cyclopropanamine
  - N-[2-(3-furyl)benzyl]cyclopropanamine
  - N-[2-(2-thienyl)benzyl]cyclopropanamine
  - 25 - N-[4,5-dimethoxy-2-(2-thienyl)benzyl]cyclopropanamine,
- as well as salts thereof.

17. A compound of formula (IIa) according to claim 16 selected in the list consisting of:

- N-[5-chloro-2-(2-furyl)benzyl]cyclopropanamine

- N-[5-chloro-2-(2-thienyl)benzyl]cyclopropanamine
- N-[2-(3-thienyl)benzyl]cyclopropanamine
- N-{1-[2-(3-thienyl)phenyl]ethyl}cyclopropanamine
- N-[5-chloro-2-(3-thienyl)benzyl]cyclopropanamine
- 5 - N-[2-(5-chloro-2-thienyl)benzyl]cyclopropanamine
- N-[5-chloro-2-(5-chloro-2-thienyl)benzyl]cyclopropanamine.

18. Compound of formula (Iib)



10

wherein  $Z^2$ ,  $Z^3$ ,  $n$  and  $X$  are as defined in any one of claims 1 to 14, and  $L$  is a direct bond and  $B^1$  is a substituted or non-substituted oxiranyl ring, a substituted or non-substituted thienyl ring, a substituted or non-substituted benzothienyl ring, a substituted or non-substituted furan ring, or a substituted or non-substituted benzofuran ring,

15

providing that (Iib) does not represent :

20

25

30


- N-[3-(2-furyl)benzyl]cyclopropanamine
  - N-[5-(2-furyl)-2-methoxybenzyl]cyclopropanamine
  - N-[4-fluoro-3-(2-furyl)benzyl]cyclopropanamine
  - N-[2-fluoro-5-(2-furyl)benzyl]cyclopropanamine
  - N-[2-fluoro-5-(3-furyl)benzyl]cyclopropanamine
  - N-[3-(3-bromo-2-furyl)-4-fluorobenzyl]cyclopropanamine
  - N-[3-(3-bromo-2-furyl)benzyl]cyclopropanamine
  - N-[5-(3-bromo-2-furyl)-2-fluorobenzyl]cyclopropanamine
  - N-[5-(3-bromo-2-furyl)-2-methoxybenzyl]cyclopropanamine
  - N-[3-(3-furyl)benzyl]cyclopropanamine
  - N-[3-(2-thienyl)benzyl]cyclopropanamine
  - N-[4-fluoro-3-(2-thienyl)benzyl]cyclopropanamine
  - N-[2-fluoro-5-(2-thienyl)benzyl]cyclopropanamine
  - N-[2-methoxy-5-(2-thienyl)benzyl]cyclopropanamine
  - 5-{5-[(cyclopropylamino)methyl]-2-fluorophenyl}thiophene-2-carbonitrile.
- as well as salts thereof.

19. A compound of formula (Iib) according to claim 18 selected in the list consisting of:

35

- N-[2-chloro-5-(2-furyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(2-thienyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(3-thienyl)benzyl]cyclopropanamine
- N-[2-chloro-5-(5-chloro-2-thienyl)benzyl]cyclopropanamine.

20. Compound of formula (Xa)



(Xa)

5

wherein  $Z^2$  and  $Z^3$  are as defined in any one of claims 1 to 14,

A represents  $A^{13}$ ,

$X^{a1}$  represents a halogen atom ; substituted or non-substituted **Ci-C<sub>8</sub>**-alkyl ; **Ci-C<sub>8</sub>**-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted **C<sub>3</sub>-C<sub>7</sub>**-cycloalkyl ; tri(**CrC<sub>8</sub>**-alkyl)silyl; substituted or non-substituted **CrC<sub>8</sub>**-alkoxy; or substituted or non-substituted **C<sub>1</sub>-C<sub>8</sub>**-alkylsulfanyl,

10

$X^{a2}$  represents hydrogen, a halogen atom ; substituted or non-substituted **CrC<sub>8</sub>**-alkyl ; **C<sub>1</sub>-C<sub>8</sub>**-halogenoalkyl comprising up to 9 halogen atoms which can be the same or different ; substituted or non-substituted **C<sub>3</sub>-C<sub>7</sub>**-cycloalkyl ; tri(**CrC<sub>8</sub>**-alkyl)silyl; substituted or non-substituted **CrC<sub>8</sub>**-alkoxy; or substituted or non-substituted **CrC<sub>8</sub>**-alkylsulfanyl, and

15

$U^2$  represents a halogen atom or a triflate group,

providing that (Xa) does not represent :

- N-(2,5-dichlorobenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
  - N-[1-(2,5-dichlorophenyl)ethyl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
  - 20 - N-[2-(2,5-dichlorophenyl)propan-2-yl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
  - N-[1-(2,5-dichlorophenyl)cyclopropyl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
  - 5-chloro-N-[1-(2,5-dichlorophenyl)ethyl]-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide
  - 5-chloro-N-[2-(2,5-dichlorophenyl)propan-2-yl]-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide
  - 5-chloro-N-[1-(2,5-dichlorophenyl)cyclopropyl]-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide
  - 25 - 5-chloro-N-(2,5-dichlorobenzyl)-3-(difluoromethyl)-1-methyl-1 H-pyrazole-4-carboxamide
  - N-(5-bromo-2-fluorobenzyl)-1,3,5-trimethyl-1 H-pyrazole-4-carboxamide
  - N-(5-bromo-2-fluorobenzyl)-3,5-dimethyl-1 H-pyrazole-4-carboxamide
  - 1-methyl-N-[1-(2,4,5-trichlorophenyl)ethyl]-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide
  - 3-(difluoromethyl)-1-methyl-N-[1-(2,4,5-trichlorophenyl)ethyl]-1 H-pyrazole-4-carboxamide
  - 30 - N-(5-chloro-2-fluorobenzyl)-1-methyl-5-propoxy-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide
  - N-(2,5-dichlorobenzyl)-1-methyl-5-propoxy-3-(trifluoromethyl)-1 H-pyrazole-4-carboxamide,
- as well as salts thereof.

21. A compound according to claim 20 selected in the list consisting of

35

- N-(5-chloro-2-ethylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
- N-(5-chloro-2-isopropylbenzyl)-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide
- N-[5-chloro-2-(trifluoromethyl)benzyl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide

- N-[3-chloro-2-fluoro-6-(trifluoromethyl)benzyl]-3-(difluoromethyl)-5-fluoro-1-methyl-1 H-pyrazole-4-carboxamide.

5 **22.** A fungicide composition comprising, as an active ingredient, an effective amount of a compound of formula (I) according to claims 1 to 14 and an agriculturally acceptable support, carrier or filler.

**23.** A method for controlling phytopathogenic fungi of crops, characterized in that an agronomically effective and substantially non-phytotoxic quantity of a compound according to any one of claims 1 to 14 or a composition according to claim 21 is applied to the soil where plants grow or are capable of growing,  
10 to the leaves and/or the fruit of plants or to the seeds of plants.

**24.** Use of compounds of the formula (I) according to any one of claims 1 to 14 for the control of phytopathogenic harmful fungi.

15 **25.** Process for producing compositions for controlling phytopathogenic harmful fungi, characterized in that derivatives of the formula (I) according to any one of claims 1 to 14 are mixed with extenders and/or surfactants.

**26.** Use of compounds of the formula (I) according to any one of claims 1 to 14 for treatment of plants, seeds, transgenic plants or transgenic seeds..

INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2013/058070

A. CLASSIFICATION OF SUBJECT MATTER  
INV. C07D403/12 C07D405/12 C07D409/12 A01N43/10  
ADD.  
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
Minimum documentation searched (classification system followed by classification symbols)  
C07D A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EPO-Internal , WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2009/024342 A2 (SYNGENTA PARTICIPATIONS AG [CH]; STIERLI DANIEL [CH]; WALTER HARALD [C]) 26 February 2009 (2009-02-26) Table .1, p. 20ff ;; claims 1, 3 -----	1-26
X	wo 2007/087906 A1 (BAYER CROPSCIENCE SA [FR]; MANSFIELD DARREN [DE]; COQUERON PIERRE-YVES) 9 August 2007 (2007-08-09) cited in the application claim 1 -----	1-26
X	wo 2009/016220 A1 (BAYER CROPSCIENCE SA [FR]; DESBORDES PHILIPPE [FR]; GARY STEPHANIE [FR]) 5 February 2009 (2009-02-05) cited in the application claim 1 -----	1-26
	-/-	

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search <b>27 May 2013</b>	Date of mailing of the international search report <b>07/06/2013</b>
---	---

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  <b>Wolf, Claudia</b>
--	--

## INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2013/058070

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	wo 2010/130767 A2 (BAYER CROPSCIENCE AG [DE] ; BARTELS GUENTER [DE] ; BECKER ANGELA [DE] ; B) 18 November 2010 (2010-11-18) cited in the application claim 1 -----	1-26
X,P	wo 2012/052490 AI (BAYER CROPSCIENCE AG [DE] ; BENTING JUERGEN [DE] ; CRISTAU PIERRE [FR] ; ) 26 April 2012 (2012-04-26) p. 51, compounds 7 and 8; -----	20
X	wo 2005/070889 AI (DU PONT [US] ; HANAGAN MARY ANN [US] ; SELBY THOMAS PAUL [US] ; SHARPE PA) 4 August 2005 (2005-08-04) p. 109, compound 85; -----	20
X	us 2010/249094 AI (YEUNG KAP-SUN [US] ET AL) 30 September 2010 (2010-09-30) p. 52, 4th compound; -----	18

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/058070

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
wo 2009024342	A2	26-02-2009	NONE
-----			
wo 2007087906	AI	09-08-2007	AR 056882 AI 31-10-2007
		AU 2006337483	AI 09-08-2007
		BR PI0620513	A2 16-11-2011
		CA 2632870	AI 09-08-2007
		CN 101360730	A 04-02-2009
		EP 1981866	AI 22-10-2008
		EP 2351747	AI 03-08-2011
		JP 5052531	B2 17-10-2012
		JP 2009525290	A 09-07-2009
		KR 20080090438	A 08-10-2008
		US 2010286221	AI 11-11-2010
		wo 2007087906	AI 09-08-2007
		ZA 200805608	A 27-01-2010
-----			
wo 2009016220	AI	05-02-2009	AT 499835 T 15-03 -2011
		AU 2008281772	AI 05-02 -2009
		CA 2692767	AI 05-02 -2009
		CN 101820759	A 01-09 -2010
		CO 6160263	A2 20-05 -2010
		EP 2182800	AI 12-05 -2010
		ES 2359877	T3 27-05 -2011
		JP 2010534715	A 11-11 -2010
		PT 2182800	E 16-05 -2011
		RU 2010107171	A 10-09 -2011
		US 2010144817	AI 10-06 -2010
		US 2010189584	AI 29-07 -2010
		wo 2009016220	AI 05-02 -2009
		ZA 200908563	A 25-08 -2010
-----			
wo 2010130767	A2	18-11-2010	AR 076839 AI 13-07 -2011
		AU 2010247385	AI 22-12 -2011
		CA 2761269	AI 18-11 -2010
		CN 102421757	A 18-04 -2012
		CO 6460782	A2 15-06 -2012
		EA 201171409	AI 30-05 -2012
		EP 2430000	A2 21-03 -2012
		JP 2012526768	A 01-11 -2012
		KR 20120023095	A 12-03 -2012
		TW 201043138	A 16-12 -2010
		US 2012065164	AI 15-03 -2012
		wo 2010130767	A2 18-11 -2010
-----			
wo 2012052490	AI	26-04-2012	NONE
-----			
wo 2005070889	AI	04--08 -2005	GT 200500013 A 10-08-2005
		wo 2005070889	AI 04-08-2005
-----			
us 2010249094	AI	30--09 -2010	US 2010249094 AI 30-09-2010
		US 2011256099	AI 20-10-2011
		US 2012201783	AI 09-08-2012
		US 2012316126	AI 13-12-2012
-----			