



(86) **Date de dépôt PCT/PCT Filing Date:** 2005/06/24
(87) **Date publication PCT/PCT Publication Date:** 2006/01/05
(45) **Date de délivrance/Issue Date:** 2015/05/12
(85) **Entrée phase nationale/National Entry:** 2006/12/21
(86) **N° demande PCT/PCT Application No.:** US 2005/022768
(87) **N° publication PCT/PCT Publication No.:** 2006/002421
(30) **Priorités/Priorities:** 2004/06/24 (US60/582,676);
2004/11/22 (US60/630,127); 2004/12/13 (US60/635,674);
2005/03/03 (US60/658,219); 2005/03/11 (US60/661,311)


(51) **Cl.Int./Int.Cl.** **C07D 401/12** (2006.01),
C07D 209/08 (2006.01), **C07D 215/56** (2006.01),
C07D 405/12 (2006.01), **C07D 409/12** (2006.01),
C07D 413/12 (2006.01), **C07D 417/12** (2006.01)

(72) **Inventeurs/Inventors:**
HADIDA-RUAH, SARA, US;
HAZLEWOOD, ANNA R., US;
GROOTENHUIS, PETER D.J., US;
VAN GOOR, FREDRICK F., US;
SINGH, ASHVANI K., US;
ZHOU, JINGLAN, US;
MCCARTNEY, JASON, US

(73) **Propriétaire/Owner:**

(54) **Titre : MODULATEURS DE TRANSPORTEURS DE CASSETTE DE LIAISON A L'ATP**

(54) **Title: MODULATORS OF ATP-BINDING CASSETTE TRANSPORTERS**



(57) **Abrégé/Abstract:**

The present invention relates to processes for making (see formula I and II), which are intermediates that may be used in the preparation of modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including Cystic Fibrosis Transmembrane Conductance Regulator. G₄ and G₅ are as defined in the specification. The invention also relates to (see formula III).




(73) **Propriétaires(suite)/Owners(continued):** VERTEX PHARMACEUTICALS INCORPORATED, US

(74) **Agent:** SMART & BIGGAR


53568-20

Abstract

The present invention relates to processes for making



are intermediates that may be used in the preparation of modulators of ATP-Binding Cassette
 5 (“ABC”) transporters or fragments thereof, including Cystic Fibrosis Transmembrane
 Conductance Regulator. G₄ and G₅ are as defined in the specification. The invention also



DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**


THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.

53568-20

MODULATORS OF ATP-BINDING CASSETTE TRANSPORTERS

[001] The present patent relates to processes for making



are intermediates that may be used in the preparation of modulators of ATP-Binding Cassette
5 (“ABC”) transporters or fragments thereof, including Cystic Fibrosis Transmembrane
Conductance Regulator. Other inventions are also disclosed. Thus, any reference to “the
present invention” and the like may refer to the invention of this patent or one of the other
inventions disclosed.

TECHNICAL FIELD OF THE INVENTION

10 [002] The present invention relates to modulators of ATP-Binding Cassette
 (“ABC”) transporters or fragments thereof, including cystic fibrosis transmembrane
conductance regulator (“CFTR”), compositions thereof, and methods therewith. The present
invention also relates to methods of treating ABC transporter mediated diseases using such
modulators.

BACKGROUND OF THE INVENTION

15 [003] ABC transporters are a family of membrane transporter proteins that
regulate the transport of a wide variety of pharmacological agents, potentially toxic drugs, and
xenobiotics, as well as anions. ABC transporters are homologous membrane proteins that bind
and use cellular adenosine triphosphate (ATP) for their specific activities. Some of these
20 transporters were discovered as multidrug resistance proteins (like the MDR1-P glycoprotein,
or the multidrug resistance protein, MRP1), defending malignant cancer cells against
chemotherapeutic agents. To date, 48 ABC Transporters have been identified and grouped
into 7 families based on their sequence identity and function.

[004] ABC transporters regulate a variety of important physiological roles within the body and provide defense against harmful environmental compounds. Because of this, they represent important potential drug targets for the treatment of diseases associated with defects in the transporter, prevention of drug transport out of the target cell, and intervention in other diseases in which modulation of ABC transporter activity may be beneficial.

[005] One member of the ABC transporter family commonly associated with disease is the cAMP/ATP-mediated anion channel, CFTR. CFTR is expressed in a variety of cells types, including absorptive and secretory epithelia cells, where it regulates anion flux across the membrane, as well as the activity of other ion channels and proteins. In epithelia cells, normal functioning of CFTR is critical for the maintenance of electrolyte transport throughout the body, including respiratory and digestive tissue. CFTR is composed of approximately 1480 amino acids that encode a protein made up of a tandem repeat of transmembrane domains, each containing six transmembrane helices and a nucleotide binding domain. The two transmembrane domains are linked by a large, polar, regulatory (R)-domain with multiple phosphorylation sites that regulate channel activity and cellular trafficking.

[006] The gene encoding CFTR has been identified and sequenced (See Gregory, R. J. et al. (1990) Nature 347:382-386; Rich, D. P. et al. (1990) Nature 347:358-362), (Riordan, J. R. et al. (1989) Science 245:1066-1073). A defect in this gene causes mutations in CFTR resulting in cystic fibrosis ("CF"), the most common fatal genetic disease in humans. Cystic fibrosis affects approximately one in every 2,500 infants in the United States. Within the general United States population, up to 10 million people carry a single copy of the defective gene without apparent ill effects. In contrast, individuals with two copies of the CF associated gene suffer from the debilitating and fatal effects of CF, including chronic lung disease.

[007] In patients with cystic fibrosis, mutations in CFTR endogenously expressed in respiratory epithelia leads to reduced apical anion secretion causing an imbalance in ion and fluid transport. The resulting decrease in anion transport contributes to enhanced mucus accumulation in the lung and the accompanying microbial infections that ultimately cause death in CF patients. In addition to respiratory disease, CF patients typically suffer from gastrointestinal problems and pancreatic insufficiency that, if left untreated, results in death. In addition, the majority of males with cystic fibrosis are infertile and fertility is decreased among females with cystic fibrosis. In contrast to the severe effects of two copies of the CF associated

gene, individuals with a single copy of the CF associated gene exhibit increased resistance to cholera and to dehydration resulting from diarrhea – perhaps explaining the relatively high frequency of the CF gene within the population.

[008] Sequence analysis of the CFTR gene of CF chromosomes has revealed a variety of disease causing mutations (Cutting, G. R. et al. (1990) *Nature* 346:366-369; Dean, M. et al. (1990) *Cell* 61:863:870; and Kerem, B-S. et al. (1989) *Science* 245:1073-1080; Kerem, B-S et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:8447-8451). To date, > 1000 disease causing mutations in the CF gene have been identified (<http://www.genet.sickkids.on.ca/cftr/>). The most prevalent mutation is a deletion of phenylalanine at position 508 of the CFTR amino acid sequence, and is commonly referred to as $\Delta F508$ -CFTR. This mutation occurs in approximately 70% of the cases of cystic fibrosis and is associated with a severe disease.

[009] The deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly. This results in the inability of the mutant protein to exit the ER, and traffic to the plasma membrane. As a result, the number of channels present in the membrane is far less than observed in cells expressing wild-type CFTR. In addition to impaired trafficking, the mutation results in defective channel gating. Together, the reduced number of channels in the membrane and the defective gating lead to reduced anion transport across epithelia leading to defective ion and fluid transport. (Quinton, P. M. (1990), *FASEB J.* 4: 2709-2727). Studies have shown, however, that the reduced numbers of $\Delta F508$ -CFTR in the membrane are functional, albeit less than wild-type CFTR. (Dalemans et al. (1991), *Nature Lond.* 354: 526-528; Denning et al., *supra*; Pasyk and Foskett (1995), *J. Cell. Biochem.* 270: 12347-50). In addition to $\Delta F508$ -CFTR, other disease causing mutations in CFTR that result in defective trafficking, synthesis, and/or channel gating could be up- or down-regulated to alter anion secretion and modify disease progression and/or severity.

[010] Although CFTR transports a variety of molecules in addition to anions, it is clear that this role (the transport of anions) represents one element in an important mechanism of transporting ions and water across the epithelium. The other elements include the epithelial Na^+ channel, ENaC, $\text{Na}^+/\text{2Cl}^-/\text{K}^+$ co-transporter, Na^+/K^+ -ATPase pump and the basolateral membrane K^+ channels, that are responsible for the uptake of chloride into the cell.

[011] These elements work together to achieve directional transport across the epithelium via their selective expression and localization within the cell. Chloride absorption

takes place by the coordinated activity of ENaC and CFTR present on the apical membrane and the Na^+/K^+ -ATPase pump and Cl^- channels expressed on the basolateral surface of the cell. Secondary active transport of chloride from the luminal side leads to the accumulation of intracellular chloride, which can then passively leave the cell via Cl^- channels, resulting in a vectorial transport. Arrangement of $\text{Na}^+/\text{2Cl}^-/\text{K}^+$ co-transporter, Na^+/K^+ -ATPase pump and the basolateral membrane K^+ channels on the basolateral surface and CFTR on the luminal side coordinate the secretion of chloride via CFTR on the luminal side. Because water is probably never actively transported itself, its flow across epithelia depends on tiny transepithelial osmotic gradients generated by the bulk flow of sodium and chloride.

[012] In addition to cystic fibrosis, modulation of CFTR activity may be beneficial for other diseases not directly caused by mutations in CFTR, such as secretory diseases and other protein folding diseases mediated by CFTR. These include, but are not limited to, chronic obstructive pulmonary disease (COPD), dry eye disease, and Sjögren's Syndrome. COPD is characterized by airflow limitation that is progressive and not fully reversible. The airflow limitation is due to mucus hypersecretion, emphysema, and bronchiolitis. Activators of mutant or wild-type CFTR offer a potential treatment of mucus hypersecretion and impaired mucociliary clearance that is common in COPD. Specifically, increasing anion secretion across CFTR may facilitate fluid transport into the airway surface liquid to hydrate the mucus and optimized periciliary fluid viscosity. This would lead to enhanced mucociliary clearance and a reduction in the symptoms associated with COPD. Dry eye disease is characterized by a decrease in tear aqueous production and abnormal tear film lipid, protein and mucin profiles. There are many causes of dry eye, some of which include age, Lasik eye surgery, arthritis, medications, chemical/thermal burns, allergies, and diseases, such as cystic fibrosis and Sjögren's syndrome. Increasing anion secretion via CFTR would enhance fluid transport from the corneal endothelial cells and secretory glands surrounding the eye to increase corneal hydration. This would help to alleviate the symptoms associated with dry eye disease. Sjögren's syndrome is an autoimmune disease in which the immune system attacks moisture-producing glands throughout the body, including the eye, mouth, skin, respiratory tissue, liver, vagina, and gut. Symptoms, include, dry eye, mouth, and vagina, as well as lung disease. The disease is also associated with rheumatoid arthritis, systemic lupus, systemic sclerosis, and polymyositis/dermatomyositis. Defective protein trafficking is believed to cause the disease,

for which treatment options are limited. Modulators of CFTR activity may hydrate the various organs afflicted by the disease and help to elevate the associated symptoms.

[013] As discussed above, it is believed that the deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly, resulting in the inability of this mutant protein to exit the ER, and traffic to the plasma membrane. As a result, insufficient amounts of the mature protein are present at the plasma membrane and chloride transport within epithelial tissues is significantly reduced. Infact, this cellular phenomenon of defective ER processing of ABC transporters by the ER machinery, has been shown to be the underlying basis not only for CF disease, but for a wide range of other isolated and inherited diseases. The two ways that the ER machinery can malfunction is either by loss of coupling to ER export of the proteins leading to degradation, or by the ER accumulation of these defective/misfolded proteins [Aridor M, *et al.*, *Nature Med.*, 5(7), pp 745- 751 (1999); Shastri, B.S., *et al.*, *Neurochem. International*, 43, pp 1-7 (2003); Rutishauser, J., *et al.*, *Swiss Med Wkly*, 132, pp 211-222 (2002); Morello, JP *et al.*, *TIPS*, 21, pp. 466- 469 (2000); Bross P., *et al.*, *Human Mut.*, 14, pp. 186-198 (1999)]. The diseases associated with the first class of ER malfunction are cystic fibrosis (due to misfolded $\Delta F508$ -CFTR as discussed above), hereditary emphysema (due to $\alpha 1$ -antitrypsin; non Piz variants), hereditary hemochromatosis, hoagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, Mucopolysaccharidoses (due to lysosomal processing enzymes), Sandhof/Tay-Sachs (due to β -hexosaminidase), Crigler-Najjar type II (due to UDP-glucuronyl-sialyc-transferase), polyendocrinopathy/hyperinsulemia, Diabetes mellitus (due to insulin receptor), Laron dwarfism (due to growth hormone receptor), myleoperoxidase deficiency, primary hypoparathyroidism (due to preproparathyroid hormone), melanoma (due to tyrosinase). The diseases associated with the latter class of ER malfunction are Glycanosis CDG type 1, hereditary emphysema (due to $\alpha 1$ -Antitrypsin (PiZ variant), congenital hyperthyroidism, osteogenesis imperfecta (due to Type I, II, IV procollagen), hereditary hypofibrinogenemia (due to fibrinogen), ACT deficiency (due to $\alpha 1$ -antichymotrypsin), Diabetes insipidus (DI), neurophyseal DI (due to vasopvessin hormone/V2-receptor), neprogenic DI (due to aquaporin II), Charcot-Marie Tooth syndrome (due to peripheral myelin protein 22), Perlizaeus-

Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease (due to β APP and presenilins), Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders such as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluyian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease (due to lysosomal α -galactosidase A) and Straussler-Scheinker syndrome (due to Prp processing defect).

[014] In addition to up-regulation of CFTR activity, reducing anion secretion by CFTR modulators may be beneficial for the treatment of secretory diarrheas, in which epithelial water transport is dramatically increased as a result of secretagogue activated chloride transport. The mechanism involves elevation of cAMP and stimulation of CFTR.

[015] Although there are numerous causes of diarrhea, the major consequences of diarrheal diseases, resulting from excessive chloride transport are common to all, and include dehydration, acidosis, impaired growth and death.

[016] Acute and chronic diarrheas represent a major medical problem in many areas of the world. Diarrhea is both a significant factor in malnutrition and the leading cause of death (5,000,000 deaths/year) in children less than five years old.

[017] Secretory diarrheas are also a dangerous condition in patients of acquired immunodeficiency syndrome (AIDS) and chronic inflammatory bowel disease (IBD). 16 million travelers to developing countries from industrialized nations every year develop diarrhea, with the severity and number of cases of diarrhea varying depending on the country and area of travel.

[018] Diarrhea in barn animals and pets such as cows, pigs and horses, sheep, goats, cats and dogs, also known as scours, is a major cause of death in these animals. Diarrhea can result from any major transition, such as weaning or physical movement, as well as in response to a variety of bacterial or viral infections and generally occurs within the first few hours of the animal's life.

[019] The most common diarrheal causing bacteria is enterotoxogenic E.coli (ETEC) having the K99 pilus antigen. Common viral causes of diarrhea include rotavirus and

coronavirus. Other infectious agents include cryptosporidium, giardia lamblia, and salmonella, among others.

[020] Symptoms of rotaviral infection include excretion of watery feces, dehydration and weakness. Coronavirus causes a more severe illness in the newborn animals, and has a higher mortality rate than rotaviral infection. Often, however, a young animal may be infected with more than one virus or with a combination of viral and bacterial microorganisms at one time. This dramatically increases the severity of the disease.

[021] Accordingly, there is a need for modulators of an ABC transporter activity, and compositions thereof, that can be used to modulate the activity of the ABC transporter in the cell membrane of a mammal.

[022] There is a need for methods of treating ABC transporter mediated diseases using such modulators of ABC transporter activity.

[023] There is a need for methods of modulating an ABC transporter activity in an *ex vivo* cell membrane of a mammal.


[024] There is a need for modulators of CFTR activity that can be used to modulate the activity of CFTR in the cell membrane of a mammal.

[025] There is a need for methods of treating CFTR-mediated diseases using such modulators of CFTR activity.

[026] There is a need for methods of modulating CFTR activity in an *ex vivo* cell membrane of a mammal.

SUMMARY OF THE INVENTION


[027] It has now been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are useful as modulators of ABC transporter activity. These compounds have the general formula **I**:




or a pharmaceutically acceptable salt thereof, wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , and Ar^1 are described generally and in classes and subclasses below.

53568-20


[027a] In an embodiment, the invention relates to a compound which is B-24

**B-24**

[027b] In another embodiment, the invention relates to a process for the preparation of
5 a compound having the formula LL

**LL**

comprising contacting the compound having the formula KK



10


KK

with Raney Ni in the presence of H₂ gas, wherein


G₄ is CF₃, t-butyl, 2-methoxyphenyl, 2-ethoxyphenyl, (4-C(O)NH(CH₂)₂-NMe₂)-phenyl, 2-methoxy-4-chloro-phenyl, pyridine-3-yl, 4-isopropylphenyl, 2,6-dimethoxyphenyl, sec-butylaminocarbonyl, ethyl, or piperidin-1-ylcarbonyl.

15 **[027c]** In a further embodiment, the invention relates to a process for the preparation of Compound B-24

53568-20

**B-24**


comprising contacting 4-*tert*-butylaniline with N-bromosuccinimide to produce 2-bromo-4-*tert*-butylaniline



5

2-bromo-4-*tert*-butylaniline


contacting 2-bromo-4-*tert*-butylaniline with a mixture of KNO₃ and sulfuric acid to produce 2-bromo-4-*tert*-butyl-5-nitroaniline



10

2-bromo-4-*tert*-butyl-5-nitroaniline

contacting 2-bromo-4-*tert*-butyl-5-nitroaniline with trimethylsilylacetylene, Pd(PPh₃)Cl₂, CuI and triethylamine to produce 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline




15

4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline


53568-20

contacting 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline with CuI to produce 5-*tert*-butyl-6-nitro-1H-indole




5-*tert*-butyl-6-nitro-1H-indole

5 contacting 5-*tert*-butyl-6-nitro-1H-indole with Raney Ni in the presence of hydrogen gas to produce 6-amino-5-*tert*-butyl-1H-indole




6-amino-5-*tert*-butyl-1H-indole (B-24)

[027d] In another invention embodiment, the invention relates to a process for the
10 preparation of a compound having the formula V-B-5a




V-B-5a

comprising contacting the compound having the formula A-II



7c


53568-20

A-IIwith Compound **A-1****A-1**


- 5 in the presence of a coupling reagent, base and solvent, wherein G_4 is hydrogen, halo, CN, CF_3 , CHF_2 , CH_2F , optionally substituted C_1 - C_6 aliphatic, aryl- C_1 - C_6 alkyl, or a phenyl, wherein G_4 is optionally substituted with up to 4 WR^W substituents; wherein up to two methylene units of said C_1 - C_6 aliphatic or C_1 - C_6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-; G_5 is hydrogen or an optionally substituted C_1 - C_6 aliphatic; and wherein said indole ring system is further optionally substituted with up to 3 substituents independently selected from WR^W , wherein W is a bond or is an optionally substituted C_1 - C_6 alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -S-, -SO-, -SO₂-, -NR'-, -SO₂NR'-, NR'SO₂-, -NR'SO₂NR'-; R^W is independently R' , halo, NO₂, CN, CF_3 , or OCF₃; and R' is independently selected from hydrogen or an optionally substituted group selected from a C_1 - C_8 aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

53568-20


[027e] In a further invention embodiment, the invention relates to a process for the preparation of Compound **381**

**381**

5 comprising contacting Compound **B-24**

**B-24**

with Compound **A-1**

**A-1**


10

in the presence of a coupling reagent, base and solvent.

53568-20

DETAILED DESCRIPTION OF THE INVENTION[029] *I. General Description of Compounds of the Invention:*

[030] The present invention relates to compounds of formula I useful as modulators of ABC transporter activity:



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is a 5-6 membered aromatic monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is optionally fused to a 5-12 membered monocyclic or bicyclic, aromatic, partially unsaturated, or saturated ring, wherein

each ring contains 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Ar^1 has m substituents, each independently selected from $-\text{WR}^{\text{W}}$;

W is a bond or is an optionally substituted $\text{C}_1\text{-C}_6$ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by $-\text{CO}-$, $-\text{CS}-$, $-\text{COCO}-$, $-\text{CONR}'-$, $-\text{CONR}'\text{NR}'-$, $-\text{CO}_2-$, $-\text{OCO}-$, $-\text{NR}'\text{CO}_2-$, $-\text{O}-$, $-\text{NR}'\text{CONR}'-$, $-\text{OCONR}'-$, $-\text{NR}'\text{NR}'$, $-\text{NR}'\text{NR}'\text{CO}-$, $-\text{NR}'\text{CO}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}'-$, $-\text{SO}_2\text{NR}'-$, $\text{NR}'\text{SO}_2-$, or $-\text{NR}'\text{SO}_2\text{NR}'-$;

R^{W} is independently R' , halo, NO_2 , CN , CF_3 , or OCF_3 ;

m is 0-5;

each of R^1 , R^2 , R^3 , R^4 , and R^5 is independently $-\text{X}-\text{R}^{\text{X}}$;

X is a bond or is an optionally substituted $\text{C}_1\text{-C}_6$ alkylidene chain wherein up to two methylene units of X are optionally and independently replaced by $-\text{CO}-$, $-\text{CS}-$, $-\text{COCO}-$, $-\text{CONR}'-$, $-\text{CONR}'\text{NR}'-$, $-\text{CO}_2-$, $-\text{OCO}-$, $-\text{NR}'\text{CO}_2-$, $-\text{O}-$, $-\text{NR}'\text{CONR}'-$, $-\text{OCONR}'-$, $-\text{NR}'\text{NR}'$, $-\text{NR}'\text{NR}'\text{CO}-$, $-\text{NR}'\text{CO}-$, $-\text{S}-$, $-\text{SO}-$, $-\text{SO}_2-$, $-\text{NR}'-$, $-\text{SO}_2\text{NR}'-$, $\text{NR}'\text{SO}_2-$, or $-\text{NR}'\text{SO}_2\text{NR}'-$;


R^{X} is independently R' , halo, NO_2 , CN , CF_3 , or OCF_3 ;

R^6 is hydrogen, CF_3 , $-\text{OR}'$, $-\text{SR}'$, or an optionally substituted C_{1-6} aliphatic group;

R^7 is hydrogen or a C_{1-6} aliphatic group optionally substituted with $-\text{X}-\text{R}^{\text{X}}$;

R' is independently selected from hydrogen or an optionally substituted group selected from a $\text{C}_1\text{-C}_8$ aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[031] In certain other embodiments, compounds of formula **I** are provided:



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is a 5-6 membered aromatic monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein said ring is optionally fused to a 5-12 membered monocyclic or bicyclic, aromatic, partially unsaturated, or saturated ring, wherein each ring contains 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein Ar¹ has m substituents each independently selected from -WR^W;

W is a bond or is an optionally substituted C₁-C₆ alkylidene chain wherein up to two methylene units of W are optionally and independently replaced by -CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -S-, -SO-, -SO₂-, -NR'-, -SO₂NR'-, NR'SO₂-, -NR'SO₂NR'-;

R^W is independently R', halo, NO₂, CN, CF₃, or OCF₃;

m is 0-5;

each of R¹, R², R³, R⁴, and R⁵ is independently -X-R^X;

X is a bond or is an optionally substituted C₁-C₆ alkylidene chain wherein up to two methylene units of X are optionally and independently replaced by -CO-, -CS-, -COCO-, -CONR'-, -CONR'NR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'NR', -NR'NR'CO-, -NR'CO-, -S-, -SO-, -SO₂-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-;

R^X is independently R', halo, NO₂, CN, CF₃, or OCF₃;

R⁶ is hydrogen, CF₃, -OR', -SR', or an optionally substituted C₁-C₈ aliphatic group;

R⁷ is hydrogen or a C₁-C₆ aliphatic group optionally substituted with -X-R^X;

R' is independently selected from hydrogen or an optionally substituted group selected from a C₁-C₈ aliphatic group, a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

provided that:

i) when $R^1, R^2, R^3, R^4, R^5, R^6$ and R^7 are hydrogen, then Ar^1 is not phenyl, 2-methoxyphenyl, 4-methoxyphenyl, 2-methylphenyl, 2,6-dichlorophenyl, 2,4-dichlorophenyl, 2-bromophenyl, 4-bromophenyl, 4-hydroxyphenyl, 2,4-dinitrophenyl, 3,5-dicarboxylic acid phenyl, 2,4-dimethylphenyl, 2,6-dimethylphenyl, 2-ethylphenyl, 3-nitro-4-methylphenyl, 3-carboxylic-acid phenyl, 2-fluorophenyl, 3-fluorophenyl, 3-trifluoromethylphenyl, 3-ethoxyphenyl, 4-chlorophenyl, 3-methoxyphenyl, 4-dimethylaminophenyl, 3,4-dimethylphenyl, 2-ethylphenyl, or 4-ethoxycarbonylphenyl;

ii) when R^1, R^2, R^3, R^5, R^6 and R^7 are hydrogen, and R^4 is methoxy, then Ar^1 is not 2-fluorophenyl or 3-fluorophenyl;

iii) when R^1, R^3, R^4, R^5, R^6 and R^7 are hydrogen, R^2 is 1,2,3,4-tetrahydroisoquinolin-1-yl-sulfonyl, then Ar^1 is not 3-trifluoromethylphenyl;

iv) when R^1, R^2, R^3, R^4, R^5 and R^7 are hydrogen, R^6 is methyl, then Ar^1 is not phenyl;

v) when R^1, R^4, R^5, R^6 and R^7 are hydrogen, R^2 and R^3 , taken together, are methylenedioxy, then Ar^1 is not 4-chlorophenyl, 4-bromophenyl, 4-nitrophenyl, 4-carboethoxyphenyl, 6-ethoxy-benzothiazol-2-yl, 6-carboethoxy-benzothiazol-2-yl, 6-halo-benzothiazol-2-yl, 6-nitro-benzothiazol-2-yl, or 6-thiocyano-benzothiazol-2-yl.

vi) when R^1, R^4, R^5, R^6 and R^7 are hydrogen, R^2 and R^3 , taken together, are methylenedioxy, then Ar^1 is not 4-substituted phenyl wherein said substituent is $-SO_2NHR^{xx}$, wherein R^{xx} is 2-pyridinyl, 4-methyl-2-pyrimidinyl, 3,4-dimethyl-5-isoxazolyl;

vii) when $R^1, R^2, R^3, R^4, R^5, R^6$, and R^7 are hydrogen, then Ar^1 is not thiazol-2-yl, 1*H*-1,2,4-triazol-3-yl, or 1*H*-1,3,4-triazol-2-yl;

viii) when R^1, R^2, R^3, R^5, R^6 , and R^7 are hydrogen, and R^4 is CF_3 , OMe, chloro, SCF_3 , or OCF_3 , then Ar^1 is not 5-methyl-1,2-oxazol-3-yl, thiazol-2-yl, 4-fluorophenyl, pyrimidin-2-yl, 1-methyl-1,2-(1*H*)-pyrazol-5-yl, pyridine-2-yl, phenyl, N-methyl-imidazol-2-yl, imidazol-2-yl, 5-methyl-imidazol-2-yl, 1,3-oxazol-2-yl, or 1,3,5-(1*H*)-triazol-2-yl;

ix) when $R^1, R^2, R^3, R^4, R^5, R^6$, and R^7 each is hydrogen, then Ar^1 is not pyrimidin-2-yl, 4,6-dimethyl-pyrimidin-2-yl, 4-methoxy-6-methyl-1,3,5-triazin-2-yl; 5-bromo-pyridin-2-yl, pyridin-2-yl, or 3,5-dichloro-pyridin-2-yl;

x) when R^1, R^2, R^3, R^4, R^5 and R^7 each is hydrogen, R^6 is hydroxy, then Ar^1 is not 2,6-dichloro-4-aminosulfonyl-phenyl;

xi) when R^2 or R^3 is an optionally substituted N-piperazyl, N-piperidyl, or N-morpholinyl, then Ar^1 is not an optionally substituted ring selected from thiazol-2-yl, pyridyl, phenyl, thiadiazolyl, benzothiazol-2-yl, or indazolyl;

xii) when R^2 is optionally substituted cyclohexylamino, then Ar^1 is not optionally substituted phenyl, pyridyl, or thiadiazolyl;

xiii) Ar^1 is not optionally substituted tetrazolyl;

xiv) when R^2 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, and R^1 and R^3 both are simultaneously CF_3 , chloro, methyl, or methoxy, then Ar^1 is not 4,5-dihydro-1,3-thiazol-2-yl, thiazol-2-yl, or [3,5-bis(trifluoromethyl)-1*H*-pyrazol-1-yl]phenyl;

xv) when R^1 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, and Ar^1 is thiazol-2-yl, then neither R^2 nor R^3 is isopropyl, chloro, or CF_3 ;

xvi) when Ar^1 is 4-methoxyphenyl, 4-trifluoromethylphenyl, 2-fluorophenyl, phenyl, or 3-chlorophenyl, then:

a) when R^1 , R^2 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, then R^3 is not methoxy; or

b) when R^1 , R^3 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, then R^2 is not chloro; or

c) when R^1 , R^2 , R^3 , R^5 , R^6 , and R^7 each is hydrogen, then R^4 is not methoxy; or



d) when when R^1 , R^3 , R^4 , R^6 , and R^7 each is hydrogen, and R^5 is ethyl, then R^2 is not chloro;

e) when R^1 , R^2 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, then R^3 is not chloro;

xvi) when R^1 , R^3 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, and R^2 is CF_3 or OCF_3 , then Ar^1 is not [3,5-bis(trifluoromethyl)-1*H*-pyrazol-1-yl]phenyl;

xvii) when R^1 , R^2 , R^4 , R^5 , R^6 , and R^7 each is hydrogen, and R^3 is hydrogen or CF_3 , then Ar^1 is not a phenyl substituted with $-OCH_2CH_2Ph$, $-OCH_2CH_2(2\text{-trifluoromethyl-phenyl})$, $-OCH_2CH_2-(6,7\text{-dimethoxy-1,2,3,4-tetrahydroisoquinolin-2-yl})$, or substituted 1*H*-pyrazol-3-yl; and

xviii) the following two compounds are excluded:



and

[032] 2. *Compounds and Definitions:*

[033] Compounds of this invention include those described generally above, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated.

[034] The term "ABC-transporter" as used herein means an ABC-transporter protein or a fragment thereof comprising at least one binding domain, wherein said protein or fragment thereof is present *in vivo* or *in vitro*. The term "binding domain" as used herein means a domain on the ABC-transporter that can bind to a modulator. See, e.g., Hwang, T. C. *et al.*, J. Gen. Physiol. (1998): 111(3), 477-90.

[035] The term "CFTR" as used herein means cystic fibrosis transmembrane conductance regulator or a mutation thereof capable of regulator activity, including, but not limited to, $\Delta F508$ CFTR and G551D CFTR (see, e.g., <http://www.genet.sickkids.on.ca/cftr/>, for CFTR mutations).

[036] The term "modulating" as used herein means increasing or decreasing by a measurable amount.

[037] For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and

53568-20

"March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M.B. and March, J., John Wiley & Sons, New York: 2001.

[038] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted", whether preceded by the term "optionally" or not, refers to the replacement of hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term "stable", as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and preferably their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[039] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "carbocycle" "cycloaliphatic" or "cycloalkyl"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-20 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-10 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-8 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-6 aliphatic carbon atoms, and in yet other embodiments aliphatic groups contain 1-4 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" (or "carbocycle" or "cycloalkyl") refers to a monocyclic C₃-C₈ hydrocarbon or bicyclic or tricyclic

C₈-C₁₄ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl. Suitable cycloaliphatic groups include cycloalkyl, bicyclic cycloalkyl (e.g., decalin), bridged bicycloalkyl such as norbornyl or [2.2.2]bicyclo-octyl, or bridged tricyclic such as adamantyl.

[040] The term "heteroaliphatic", as used herein, means aliphatic groups wherein one or two carbon atoms are independently replaced by one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon. Heteroaliphatic groups may be substituted or unsubstituted, branched or unbranched, cyclic or acyclic, and include "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" groups.

[041] The term "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic, or tricyclic ring systems in which one or more ring members is an independently selected heteroatom. In some embodiments, the "heterocycle", "heterocyclyl", "heterocycloaliphatic", or "heterocyclic" group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 7 ring members.

[042] The term "heteroatom" means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl)).

[043] The term "unsaturated", as used herein, means that a moiety has one or more units of unsaturation.

[044] The term "alkoxy", or "thioalkyl", as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen ("alkoxy") or sulfur ("thioalkyl") atom.

[045] The terms “haloaliphatic” and “haloalkoxy” means aliphatic or alkoxy, as the case may be, substituted with one or more halo atoms. The term “halogen” or “halo” means F, Cl, Br, or I. Examples of haloaliphatic include $-\text{CHF}_2$, $-\text{CH}_2\text{F}$, $-\text{CF}_3$, $-\text{CF}_2-$, or perhaloalkyl, such as, $-\text{CF}_2\text{CF}_3$.

[046] The term “aryl” used alone or as part of a larger moiety as in “aralkyl”, “aralkoxy”, or “aryloxyalkyl”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term “aryl” may be used interchangeably with the term “aryl ring”. The term “aryl” also refers to heteroaryl ring systems as defined hereinbelow.

[047] The term “heteroaryl”, used alone or as part of a larger moiety as in “heteroaralkyl” or “heteroarylalkoxy”, refers to monocyclic, bicyclic, and tricyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms, and wherein each ring in the system contains 3 to 7 ring members. The term “heteroaryl” may be used interchangeably with the term “heteroaryl ring” or the term “heteroaromatic”.

[048] An aryl (including aralkyl, aralkoxy, aryloxyalkyl and the like) or heteroaryl (including heteroaralkyl and heteroarylalkoxy and the like) group may contain one or more substituents. Suitable substituents on the unsaturated carbon atom of an aryl or heteroaryl group are selected from halo; $-\text{R}^\circ$; $-\text{OR}^\circ$; $-\text{SR}^\circ$; 1,2-methylene-dioxy; 1,2-ethylenedioxy; phenyl (Ph) optionally substituted with R° ; $-\text{O}(\text{Ph})$ optionally substituted with R° ; $-(\text{CH}_2)_{1-2}(\text{Ph})$, optionally substituted with R° ; $-\text{CH}=\text{CH}(\text{Ph})$, optionally substituted with R° ; $-\text{NO}_2$; $-\text{CN}$; $-\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\circ$; $-\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{OC}(\text{O})\text{N}(\text{R}^\circ)_2$; $-\text{S}(\text{O})_2\text{R}^\circ$; $-\text{SO}_2\text{N}(\text{R}^\circ)_2$; $-\text{S}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{SO}_2\text{N}(\text{R}^\circ)_2$; $-\text{NR}^\circ\text{SO}_2\text{R}^\circ$; $-\text{C}(=\text{S})\text{N}(\text{R}^\circ)_2$; $-\text{C}(=\text{NH})-\text{N}(\text{R}^\circ)_2$; or $-(\text{CH}_2)_{0-2}\text{NHC}(\text{O})\text{R}^\circ$ wherein each independent occurrence of R° is selected from hydrogen, optionally substituted C_{1-6} aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, $-\text{O}(\text{Ph})$, or $-\text{CH}_2(\text{Ph})$, or, notwithstanding the definition above, two independent occurrences of R° , on the same substituent or different substituents, taken together with the atom(s) to which each R° group is bound, form a 3-8-membered cycloalkyl, heterocyclyl, aryl, or heteroaryl ring having 0-3 heteroatoms

independently selected from nitrogen, oxygen, or sulfur. Optional substituents on the aliphatic group of R° are selected from NH_2 , $NH(C_{1-4} \text{aliphatic})$, $N(C_{1-4} \text{aliphatic})_2$, halo, $C_{1-4} \text{aliphatic}$, OH , $O(C_{1-4} \text{aliphatic})$, NO_2 , CN , CO_2H , $CO_2(C_{1-4} \text{aliphatic})$, $O(\text{halo } C_{1-4} \text{aliphatic})$, or $\text{halo } C_{1-4} \text{aliphatic}$, wherein each of the foregoing $C_{1-4} \text{aliphatic}$ groups of R° is unsubstituted.


[049] An aliphatic or heteroaliphatic group, or a non-aromatic heterocyclic ring may contain one or more substituents. Suitable substituents on the saturated carbon of an aliphatic or heteroaliphatic group, or of a non-aromatic heterocyclic ring are selected from those listed above for the unsaturated carbon of an aryl or heteroaryl group and additionally include the following: $=O$, $=S$, $=NNHR^*$, $=NN(R^*)_2$, $=NNHC(O)R^*$, $=NNHCO_2(\text{alkyl})$, $=NNHSO_2(\text{alkyl})$, or $=NR^*$, where each R^* is independently selected from hydrogen or an optionally substituted C_{1-6} aliphatic. Optional substituents on the aliphatic group of R^* are selected from NH_2 , $NH(C_{1-4} \text{aliphatic})$, $N(C_{1-4} \text{aliphatic})_2$, halo, $C_{1-4} \text{aliphatic}$, OH , $O(C_{1-4} \text{aliphatic})$, NO_2 , CN , CO_2H , $CO_2(C_{1-4} \text{aliphatic})$, $O(\text{halo } C_{1-4} \text{aliphatic})$, or $\text{halo}(C_{1-4} \text{aliphatic})$, wherein each of the foregoing $C_{1-4} \text{aliphatic}$ groups of R^* is unsubstituted.


[050] Optional substituents on the nitrogen of a non-aromatic heterocyclic ring are selected from $-R^+$, $-N(R^+)_2$, $-C(O)R^+$, $-CO_2R^+$, $-C(O)C(O)R^+$, $-C(O)CH_2C(O)R^+$, $-SO_2R^+$, $-SO_2N(R^+)_2$, $-C(=S)N(R^+)_2$, $-C(=NH)-N(R^+)_2$, or $-NR^+SO_2R^+$; wherein R^+ is hydrogen, an optionally substituted C_{1-6} aliphatic, optionally substituted phenyl, optionally substituted $-O(\text{Ph})$, optionally substituted $-CH_2(\text{Ph})$, optionally substituted $-(CH_2)_{1-2}(\text{Ph})$; optionally substituted $-CH=CH(\text{Ph})$; or an unsubstituted 5-6 membered heteroaryl or heterocyclic ring having one to four heteroatoms independently selected from oxygen, nitrogen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R^+ , on the same substituent or different substituents, taken together with the atom(s) to which each R^+ group is bound, form a 3-8-membered cycloalkyl, heterocyclyl, aryl, or heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Optional substituents on the aliphatic group or the phenyl ring of R^+ are selected from NH_2 , $NH(C_{1-4} \text{aliphatic})$, $N(C_{1-4} \text{aliphatic})_2$, halo, $C_{1-4} \text{aliphatic}$, OH , $O(C_{1-4} \text{aliphatic})$, NO_2 , CN , CO_2H , $CO_2(C_{1-4} \text{aliphatic})$, $O(\text{halo } C_{1-4} \text{aliphatic})$, or $\text{halo}(C_{1-4} \text{aliphatic})$, wherein each of the foregoing $C_{1-4} \text{aliphatic}$ groups of R^+ is unsubstituted.

[051] The term "alkylidene chain" refers to a straight or branched carbon chain that may be fully saturated or have one or more units of unsaturation and has two points of

attachment to the rest of the molecule. The term “spirocycloalkylidene” refers to a carbocyclic ring that may be fully saturated or have one or more units of unsaturation and has two points of attachment from the same ring carbon atom to the rest of the molecule.


[052] As detailed above, in some embodiments, two independent occurrences of R^0 (or R^+ , or any other variable similarly defined herein), are taken together together with the atom(s) to which each variable is bound to form a 3-8-membered cycloalkyl, heterocyclyl, aryl, or heteroaryl ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Exemplary rings that are formed when two independent occurrences of R^0 (or R^+ , or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound include, but are not limited to the following: a) two independent occurrences of R^0 (or R^+ , or any other variable similarly defined herein) that are bound to the same atom and are taken together with that atom to form a ring, for example, $N(R^0)_2$, where both occurrences of R^0 are taken together with the nitrogen atom to form a piperidin-1-yl, piperazin-1-yl, or morpholin-4-yl group; and b) two independent occurrences of R^0 (or R^+ , or any other variable similarly defined herein) that are bound to different atoms and are taken together with both of those atoms to form a ring, for example where a phenyl group is substituted with two

occurrences of OR^0 , these two occurrences of R^0 are taken together with the oxygen atoms to which they are bound to form a fused 6-membered oxygen containing ring:




. It will be appreciated that a variety of other rings can be formed when two independent occurrences of R^0 (or R^+ , or any other variable similarly defined herein) are taken together with the atom(s) to which each variable is bound and that the examples detailed above are not intended to be limiting.

[053] A substituent bond in, e.g., a bicyclic ring system, as shown below, means that the substituent can be attached to any substitutable ring atom on either ring of the bicyclic ring system:




[054] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. E.g., when R⁵ in compounds of formula I is hydrogen, compounds of formula I may exist as tautomers:



Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ¹³C- or ¹⁴C-enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

[055] 3. *Description of Exemplary Compounds:*

[056] In some embodiments of the present invention, Ar¹ is selected from:



wherein ring A₁ 5-6 membered aromatic monocyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or

A₁ and A₂, together, is an 8-14 aromatic, bicyclic or tricyclic aryl ring, wherein each ring contains 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[057] In some embodiments, A_1 is an optionally substituted 6 membered aromatic ring having 0-4 heteroatoms, wherein said heteroatom is nitrogen. In some embodiments, A_1 is an optionally substituted phenyl. Or, A_1 is an optionally substituted pyridyl, pyrimidinyl, pyrazinyl or triazinyl. Or, A_1 is an optionally substituted pyrazinyl or triazinyl. Or, A_1 is an optionally substituted pyridyl.

[058] In some embodiments, A_1 is an optionally substituted 5-membered aromatic ring having 0-3 heteroatoms, wherein said heteroatom is nitrogen, oxygen, or sulfur. In some embodiments, A_1 is an optionally substituted 5-membered aromatic ring having 1-2 nitrogen atoms. In one embodiment, A_1 is an optionally substituted 5-membered aromatic ring other than thiazolyl.


[059] In some embodiments, A_2 is an optionally substituted 6 membered aromatic ring having 0-4 heteroatoms, wherein said heteroatom is nitrogen. In some embodiments, A_2 is an optionally substituted phenyl. Or, A_2 is an optionally substituted pyridyl, pyrimidinyl, pyrazinyl, or triazinyl.
























[060] In some embodiments, A_2 is an optionally substituted 5-membered aromatic ring having 0-3 heteroatoms, wherein said heteroatom is nitrogen, oxygen, or sulfur. In some embodiments, A_2 is an optionally substituted 5-membered aromatic ring having 1-2 nitrogen atoms. In certain embodiments, A_2 is an optionally substituted pyrrolyl.


[061] In some embodiments, A_2 is an optionally substituted 5-7 membered saturated or unsaturated heterocyclic ring having 1-3 heteroatoms independently selected from nitrogen, sulfur, or oxygen. Exemplary such rings include piperidyl, piperazyl, morpholinyl, thiomorpholinyl, pyrrolidinyl, tetrahydrofuranyl, etc.

[062] In some embodiments, A_2 is an optionally substituted 5-10 membered saturated or unsaturated carbocyclic ring. In one embodiment, A_2 is an optionally substituted 5-10 membered saturated carbocyclic ring. Exemplary such rings include cyclohexyl, cyclopentyl, etc.

[063] In some embodiments, ring A_2 is selected from:



**v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi****xvii****xviii****xix****xx****xxi****xxii****xxiii****xxiv****xxv****xxvi****xxviii**



wherein ring A₂ is fused to ring A₁ through two adjacent ring atoms.

[064] In other embodiments, W is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR', S, SO, SO₂, or COO, CO, SO₂NR', NR'SO₂, C(O)NR', NR'C(O), OC(O), OC(O)NR', and R^W is R' or halo. In still other embodiments, each occurrence of WR^W is independently -C1-C3 alkyl, C1-C3 perhaloalkyl, -O(C1-C3alkyl), -CF₃, -OCF₃, -SCF₃, -F, -Cl, -Br, or -COOR', -COR', -O(CH₂)₂N(R')(R'), -O(CH₂)N(R')(R'), -CON(R')(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted monocyclic or bicyclic aromatic ring, optionally substituted arylsulfone, optionally substituted 5-membered heteroaryl ring, -N(R')(R'), -(CH₂)₂N(R')(R'), or -(CH₂)N(R')(R').

[065] In some embodiments, m is 0. Or, m is 1. Or, m is 2. In some embodiments, m is 3. In yet other embodiments, m is 4.

[066] In one embodiment, R⁵ is X-R^X. In some embodiments R⁵ is hydrogen. Or, R⁵ is an optionally substituted C₁₋₈ aliphatic group. In some embodiments, R⁵ is optionally substituted C₁₋₄ aliphatic. Or, R⁵ is benzyl.

[067] In some embodiments R⁶ is hydrogen. Or, R⁶ is an optionally substituted C₁₋₈ aliphatic group. In some embodiments, R⁶ is optionally substituted C₁₋₄ aliphatic. In certain other embodiments, R⁶ is -(O-C₁₋₄ aliphatic) or -(S-C₁₋₄ aliphatic). Preferably, R⁶ is -OMe or -SMe. In certain other embodiments, R₆ is CF₃.

[068] In one embodiment of the present invention, R¹, R², R³, and R⁴ are simultaneously hydrogen. In another embodiment, R⁶ and R⁷ are both simultaneously hydrogen.

[069] In another embodiment of the present invention, R¹, R², R³, R⁴, and R⁵ are simultaneously hydrogen. In another embodiment of the present invention, R¹, R², R³, R⁴, R⁵ and R⁶ are simultaneously hydrogen.

[070] In another embodiment of the present invention, R^2 is $X-R^X$, wherein X is $-SO_2NR'$, and R^X is R' ; i.e., R^2 is $-SO_2N(R')_2$. In one embodiment, the two R' therein taken together form an optionally substituted 5-7 membered ring with 0-3 additional heteroatoms selected from nitrogen, oxygen, or sulfur. Or, R^1 , R^3 , R^4 , R^5 and R^6 are simultaneously hydrogen, and R^2 is $SO_2N(R')_2$.

[071] In some embodiments, X is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two non-adjacent methylene units are optionally and independently replaced by O, NR' , S, SO_2 , or COO, CO, and R^X is R' or halo. In still other embodiments, each occurrence of XR^X is independently $-C_{1-3}alkyl$, $-O(C_{1-3}alkyl)$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, $-Br$, OH , $-COOR'$, $-COR'$, $-O(CH_2)_2N(R')(R')$, $-O(CH_2)N(R')(R')$, $-CON(R')(R')$, $-(CH_2)_2OR'$, $-(CH_2)OR'$, optionally substituted phenyl, $-N(R')(R')$, $-(CH_2)_2N(R')(R')$, or $-(CH_2)N(R')(R')$.

[072] In some embodiments, R^7 is hydrogen. In certain other embodiment, R^7 is C_{1-4} straight or branched aliphatic.

[073] In some embodiments, R^W is selected from halo, cyano, CF_3 , CHF_2 , $OCHF_2$, Me, Et, $CH(Me)_2$, $CHMeEt$, n-propyl, t-butyl, OMe, OEt, OPh, O-fluorophenyl, O-difluorophenyl, O-methoxyphenyl, O-tolyl, O-benzyl, SMe, SCF_3 , $SCHF_2$, SEt, CH_2CN , NH_2 , $NHMe$, $N(Me)_2$, $NHEt$, $N(Et)_2$, $C(O)CH_3$, $C(O)Ph$, $C(O)NH_2$, SPh, SO_2 -(amino-pyridyl), SO_2NH_2 , SO_2Ph , SO_2NHPh , SO_2 -N-morpholino, SO_2 -N-pyrrolidyl, N-pyrrolyl, N-morpholino, 1-piperidyl, phenyl, benzyl, (cyclohexyl-methylamino)methyl, 4-Methyl-2,4-dihydro-pyrazol-3-one-2-yl, benzimidazol-2-yl, furan-2-yl, 4-methyl-4H-[1,2,4]triazol-3-yl, 3-(4'-chlorophenyl)-[1,2,4]oxadiazol-5-yl, $NHC(O)Me$, $NHC(O)Et$, $NHC(O)Ph$, $NHSO_2Me$, 2-indolyl, 5-indolyl, $-CH_2CH_2OH$, $-OCF_3$, O-(2,3-dimethylphenyl), 5-methylfuryl, $-SO_2$ -N-piperidyl, 2-tolyl, 3-tolyl, 4-tolyl, O-butyl, $NHCO_2C(Me)_3$, $CO_2C(Me)_3$, isopropenyl, n-butyl, O-(2,4-dichlorophenyl), $NHSO_2PhMe$, O-(3-chloro-5-trifluoromethyl-2-pyridyl), phenylhydroxymethyl, 2,5-dimethylpyrrolyl, $NHCOCH_2C(Me)_3$, O-(2-tert-butyl)phenyl, 2,3-dimethylphenyl, 3,4-dimethylphenyl, 4-hydroxymethyl phenyl, 4-dimethylaminophenyl, 2-trifluoromethylphenyl, 3-trifluoromethylphenyl, 4-trifluoromethylphenyl, 4-cyanomethylphenyl, 4-isobutylphenyl, 3-pyridyl, 4-pyridyl, 4-isopropylphenyl, 3-isopropylphenyl, 2-methoxyphenyl, 3-methoxyphenyl, 4-methoxyphenyl, 3,4-methylenedioxyphenyl, 2-ethoxyphenyl, 3-ethoxyphenyl, 4-ethoxyphenyl, 2-methylthiophenyl, 4-methylthiophenyl, 2,4-dimethoxyphenyl, 2,5-dimethoxyphenyl, 2,6-

dimethoxyphenyl, 3,4-dimethoxyphenyl, 5-chloro-2-methoxyphenyl, 2-OCF₃-phenyl, 3-trifluoromethoxy-phenyl, 4-trifluoromethoxyphenyl, 2-phenoxyphenyl, 4-phenoxyphenyl, 2-fluoro-3-methoxy-phenyl, 2,4-dimethoxy-5-pyrimidyl, 5-isopropyl-2-methoxyphenyl, 2-fluorophenyl, 3-fluorophenyl, 4-fluorophenyl, 3-cyanophenyl, 3-chlorophenyl, 4-chlorophenyl, 2,3-difluorophenyl, 2,4-difluorophenyl, 2,5-difluorophenyl, 3,4-difluorophenyl, 3,5-difluorophenyl, 3-chloro-4-fluoro-phenyl, 3,5-dichlorophenyl, 2,5-dichlorophenyl, 2,3-dichlorophenyl, 3,4-dichlorophenyl, 2,4-dichlorophenyl, 3-methoxycarbonylphenyl, 4-methoxycarbonyl phenyl, 3-isopropoxyloxycarbonylphenyl, 3-acetamidophenyl, 4-fluoro-3-methylphenyl, 4-methanesulfinyl-phenyl, 4-methanesulfonyl-phenyl, 4-N-(2-N,N-dimethylaminoethyl)carbamoylphenyl, 5-acetyl-2-thienyl, 2-benzothienyl, 3-benzothienyl, furan-3-yl, 4-methyl-2-thienyl, 5-cyano-2-thienyl, N'-phenylcarbonyl-N-piperazinyl, -NHCO₂Et, -NHCO₂Me, N-pyrrolidinyl, -NHSO₂(CH₂)₂ N-piperidine, -NHSO₂(CH₂)₂ N-morpholine, -NHSO₂(CH₂)₂N(Me)₂, COCH₂N(Me)COCH₂NHMe, -CO₂Et, O-propyl, -CH₂CH₂NHCO₂C(Me)₃, hydroxy, aminomethyl, pentyl, adamantyl, cyclopentyl, ethoxyethyl, C(Me)₂CH₂OH, C(Me)₂CO₂Et, -CHOHMe, CH₂CO₂Et, -C(Me)₂CH₂NHCO₂C(Me)₃, O(CH₂)₂OEt, O(CH₂)₂OH, CO₂Me, hydroxymethyl, , 1-methyl-1-cyclohexyl, 1-methyl-1-cyclooctyl, 1-methyl-1-cycloheptyl, C(Et)₂C(Me)₃, C(Et)₃, CONHCH₂CH(Me)₂, 2-aminomethyl-phenyl, ethenyl, 1-piperidinylcarbonyl, ethynyl, cyclohexyl, 4-methylpiperidinyl, -OCO₂Me, -C(Me)₂CH₂NHCO₂CH₂CH(Me)₂, -C(Me)₂CH₂NHCO₂CH₂CH₂CH₃, -C(Me)₂CH₂NHCO₂Et, -C(Me)₂CH₂NHCO₂Me, -C(Me)₂CH₂NHCO₂CH₂C(Me)₃, -CH₂NHCOCF₃, -CH₂NHCO₂C(Me)₃, -C(Me)₂CH₂NHCO₂(CH₂)₃CH₃, C(Me)₂CH₂NHCO₂(CH₂)₂OMe, C(OH) (CF₃)₂, -C(Me)₂CH₂NHCO₂CH₂-tetrahydrofurane-3-yl, C(Me)₂CH₂O(CH₂)₂OMe, or 3-ethyl-2,6-dioxopiperidin-3-yl.

[074] In one embodiment, R' is hydrogen.



[075] In one embodiment, R' is a C1-C8 aliphatic group, optionally substituted with up to 3 substituents selected from halo, CN, CF₃, CHF₂, OCF₃, or OCHF₂, wherein up to two methylene units of said C1-C8 aliphatic is optionally replaced with -CO-, -CONH(C1-C4 alkyl)-, -CO₂-, -OCO-, -N(C1-C4 alkyl)CO₂-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, -SO₂N(C1-C4 alkyl)-, N(C1-C4 alkyl)SO₂-, or -N(C1-C4 alkyl)SO₂N(C1-C4 alkyl)-.

[076] In one embodiment, R' is a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF₃, CHF₂, OCF₃, OCHF₂, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with -CO-, -CONH(C1-C4 alkyl)-, -CO₂-, -OCO-, -N(C1-C4 alkyl)CO₂-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, -SO₂N(C1-C4 alkyl)-, N(C1-C4 alkyl)SO₂-, or -N(C1-C4 alkyl)SO₂N(C1-C4 alkyl)-.






[077] In one embodiment, R' is an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF₃, CHF₂, OCF₃, OCHF₂, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with -CO-, -CONH(C1-C4 alkyl)-, -CO₂-, -OCO-, -N(C1-C4 alkyl)CO₂-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, -SO₂N(C1-C4 alkyl)-, N(C1-C4 alkyl)SO₂-, or -N(C1-C4 alkyl)SO₂N(C1-C4 alkyl)-.

[078] In one embodiment, two occurrences of R' are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, wherein R' is optionally substituted with up to 3 substituents selected from halo, CN, CF₃, CHF₂, OCF₃, OCHF₂, or C1-C6 alkyl, wherein up to two methylene units of said C1-C6 alkyl is optionally replaced with -CO-, -CONH(C1-C4 alkyl)-, -CO₂-, -OCO-, -N(C1-C4 alkyl)CO₂-, -O-, -N(C1-C4 alkyl)CON(C1-C4 alkyl)-, -OCON(C1-C4 alkyl)-, -N(C1-C4 alkyl)CO-, -S-, -N(C1-C4 alkyl)-, -SO₂N(C1-C4 alkyl)-, N(C1-C4 alkyl)SO₂-, or -N(C1-C4 alkyl)SO₂N(C1-C4 alkyl)-.

[079] According to one embodiment, the present invention provides compounds of formula **IIA** or formula **IIB**:

**IIA****IIB**

[080] According to another embodiment, the present invention provides compounds of formula **IIIA**, formula **IIIB**, formula **IIIC**, formula **IIID**, or formula **IIIE**:


**IIIA****IIIB****IIIC****IIID****IIIE**

wherein each of X_1 , X_2 , X_3 , X_4 , and X_5 is independently selected from CH or N; and X_6 is O, S, or NR' .


[081] In one embodiment, compounds of formula **IIIA**, formula **IIIB**, formula **IIIC**, formula **IIID**, or formula **IIIE** have y occurrences of substituent $X-R^X$, wherein y is 0-4. Or, y is 1. Or, y is 2.


[082] In some embodiments of formula **IIIA**, X_1 , X_2 , X_3 , X_4 , and X_5 taken together with WR^W and m is optionally substituted phenyl.





















[083] In some embodiments of formula **IIIA**, X_1 , X_2 , X_3 , X_4 , and X_5 taken together is an optionally substituted ring selected from:















































[084] In some embodiments of formula **IIIB**, formula **IIIC**, formula **IIID**, or formula **IIIE**, X_1 , X_2 , X_3 , X_4 , X_5 , or X_6 , taken together with ring A_2 is an optionally substituted ring selected from:






**b-xxxv****b-xxxvi****b-xxxvii****b-xxxviii****bxxxix****b-xL****b-xLi****b-xLii****b-xLiii****b-xLiv****b-xLv****b-xLvi****b-xLviii****b-xLviii****b-xLix****b-L****b-Li****b-Lii****b-Liii****b-Liv**

**b-Lv****b-Lvi****b-Lvii****b-Lviii****b-Lix****b-Lx****b-Lxi****b-Lxii****b-Lxiii****b-Lxiv****b-Lxv****b-Lxvi****b-Lxvii****b-Lxviii****b-Lxix****b-Lxx****b-Lxxi****b-Lxxii****b-Lxxiii****b-Lxxiv**




**b-Lxxv****b-Lxxvi****b-Lxxvii****b-Lxxviii****b-Lxxix****b-Lxxx****b-Lxxxi****b-Lxxxii****b-Lxxxiii****b-Lxxxiv****b-Lxxxv****b-Lxxxvi****b-Lxxxviii****b-Lxxxix****b-xC****b-xCi****b-xCii****b-xCiii****b-xCiv****b-xCv****b-xCvi****b-xCvii****b-xCviii****b-xCix**



[085] In some embodiments, R^W is selected from halo, cyano, CF_3 , CHF_2 , $OCHF_2$, Me, Et, $CH(Me)_2$, $CHMeEt$, n-propyl, t-butyl, OMe, OEt, OPh, O-fluorophenyl, O-difluorophenyl, O-methoxyphenyl, O-tolyl, O-benzyl, SMe, SCF_3 , $SCHF_2$, SEt, CH_2CN , NH_2 , $NHMe$, $N(Me)_2$, $NHEt$, $N(Et)_2$, $C(O)CH_3$, $C(O)Ph$, $C(O)NH_2$, SPh, SO_2 -(amino-pyridyl), SO_2NH_2 , SO_2Ph , SO_2NHPh , SO_2 -N-morpholino, SO_2 -N-pyrrolidyl, N-pyrrolyl, N-morpholino, 1-piperidyl, phenyl, benzyl, (cyclohexyl-methylamino)methyl, 4-Methyl-2,4-dihydro-pyrazol-3-one-2-yl, benzimidazol-2-yl, furan-2-yl, 4-methyl-4H-[1,2,4]triazol-3-yl, 3-(4'-chlorophenyl)-[1,2,4]oxadiazol-5-yl, $NHC(O)Me$, $NHC(O)Et$, $NHC(O)Ph$, or $NHSO_2Me$

[086] In some embodiments, X and R^X , taken together, is Me, Et, halo, CN, CF_3 , OH, OMe, OEt, $SO_2N(Me)$ (fluorophenyl), SO_2 -(4-methyl-piperidin-1-yl), or SO_2 -N-pyrrolidinyl.

[087] According to another embodiment, the present invention provides compounds of formula **IVA**, formula **IVB**, or formula **IVC**:



[088] In one embodiment compounds of formula **IVA**, formula **IVB**, and formula **IVC** have y occurrences of substituent $X-R^X$, wherein y is 0-4. Or, y is 1. Or, y is 2.

[089] In one embodiment, the present invention provides compounds of formula **IVA**, formula **IVB**, and formula **IVC**, wherein X is a bond and R^X is hydrogen.


[090] In one embodiment, the present invention provides compounds of formula **IVB**, and formula **IVC**, wherein ring A₂ is an optionally substituted, saturated, unsaturated, or aromatic seven membered ring with 0-3 heteroatoms selected from O, S, or N. Exemplary rings include azepanyl, 5,5-dimethyl azepanyl, etc.

[091] In one embodiment, the present invention provides compounds of formula **IVB** and **IVC**, wherein ring A₂ is an optionally substituted, saturated, unsaturated, or aromatic six membered ring with 0-3 heteroatoms selected from O, S, or N. Exemplary rings include piperidinyl, 4,4-dimethylpiperidinyl, etc.

[092] In one embodiment, the present invention provides compounds of formula **IVB** and **IVC**, wherein ring A₂ is an optionally substituted, saturated, unsaturated, or aromatic five membered ring with 0-3 heteroatoms selected from O, S, or N.

[093] In one embodiment, the present invention provides compounds of formula **IVB** and **IVC**, wherein ring A₂ is an optionally substituted five membered ring with one nitrogen atom, e.g., pyrrolyl or pyrrolidinyl.

[094] According to one embodiment of formula **IVA**, the following compound of formula **VA-1** is provided:



VA-1

wherein each of WR^{W2} and WR^{W4} is independently selected from hydrogen, CN, CF₃, halo, C1-C6 straight or branched alkyl, 3-12 membered cycloaliphatic, phenyl, C5-C10 heteroaryl or C3-C7 heterocyclic, wherein said heteroaryl or heterocyclic has up to 3 heteroatoms selected from O, S, or N, wherein said WR^{W2} and WR^{W4} is independently and optionally substituted with up to three substituents selected from -OR', -CF₃, -OCF₃, SR', S(O)R', SO₂R', -SCF₃, halo, CN, -COOR', -COR', -O(CH₂)₂N(R')(R'), -O(CH₂)N(R')(R'), -CON(R')(R'), -(CH₂)₂OR', -(CH₂)OR', CH₂CN, optionally substituted phenyl or phenoxy, -N(R')(R'), -NR'C(O)OR', -NR'C(O)R', -(CH₂)₂N(R')(R'), or -(CH₂)N(R')(R'); and

WR^{W5} is selected from hydrogen, -OH, NH₂, CN, CHF₂, NHR', N(R')₂, -NHC(O)R', -NHC(O)OR', NHSO₂R', -OR', CH₂OH, CH₂N(R')₂, C(O)OR', SO₂NHR', SO₂N(R')₂, or

CH₂NHC(O)OR'. Or, WR^{W4} and WR^{W5} taken together form a 5-7 membered ring containing 0-3 three heteroatoms selected from N, O, or S, wherein said ring is optionally substituted with up to three WR^W substituents.

[095] In one embodiment, compounds of formula VA-1 have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0.

[096] In one embodiment, the present invention provides compounds of formula VA-1, wherein X is a bond and R^X is hydrogen.

[097] In one embodiment, the present invention provides compounds of formula VA-1, wherein:

each of WR^{W2} and WR^{W4} is independently selected from hydrogen, CN, CF₃, halo, C1-C6 straight or branched alkyl, 3-12 membered cycloaliphatic, or phenyl, wherein said WR^{W2} and WR^{W4} is independently and optionally substituted with up to three substituents selected from -OR', -CF₃, -OCF₃, -SCF₃, halo, -COOR', -COR', -O(CH₂)₂N(R')(R'), -O(CH₂)N(R')(R'), -CON(R')(R'), -(CH₂)₂OR', -(CH₂)OR', optionally substituted phenyl, -N(R')(R'), -NC(O)OR', -NC(O)R', -(CH₂)₂N(R')(R'), or -(CH₂)N(R')(R'); and

WR^{W5} is selected from hydrogen, -OH, NH₂, CN, NHR', N(R')₂, -NHC(O)R', -NHC(O)OR', NHSO₂R', -OR', CH₂OH, C(O)OR', SO₂NHR', or CH₂NHC(O)O-(R').

[098] In one embodiment, the present invention provides compounds of formula VA-1, wherein:

WR^{W2} is a phenyl ring optionally substituted with up to three substituents selected from -OR', -CF₃, -OCF₃, SR', S(O)R', SO₂R', -SCF₃, halo, CN, -COOR', -COR', -O(CH₂)₂N(R')(R'), -O(CH₂)N(R')(R'), -CON(R')(R'), -(CH₂)₂OR', -(CH₂)OR', CH₂CN, optionally substituted phenyl or phenoxy, -N(R')(R'), -NR'C(O)OR', -NR'C(O)R', -(CH₂)₂N(R')(R'), or -(CH₂)N(R')(R');

WR^{W4} is C1-C6 straight or branched alkyl; and

WR^{W5} is OH.

[099] In one embodiment, each of WR^{W2} and WR^{W4} is independently selected from CF₃ or halo. In one embodiment, each of WR^{W2} and WR^{W4} is independently selected from optionally substituted hydrogen, C1-C6 straight or branched alkyl. In certain embodiments, each of WR^{W2} and WR^{W4} is independently selected from optionally substituted n-propyl,

isopropyl, n-butyl, sec-butyl, t-butyl, 1,1-dimethyl-2-hydroxyethyl, 1,1-dimethyl-2-(ethoxycarbonyl)-ethyl, 1,1-dimethyl-3-(t-butoxycarbonyl-amino) propyl, or n-pentyl.

[0100] In one embodiment, each of WR^{W2} and WR^{W4} is independently selected from optionally substituted 3-12 membered cycloaliphatic. Exemplary embodiments of such cycloaliphatic include cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, adamantyl, [2.2.2.]bicyclo-octyl, [2.3.1.] bicyclo-octyl, or [3.3.1]bicyclo-nonyl.

[0101] In certain embodiments WR^{W2} is hydrogen and WR^{W4} is C1-C6 straight or branched alkyl. In certain embodiments, WR^{W4} is selected from methyl, ethyl, propyl, n-butyl, sec-butyl, or t-butyl.

[0102] In certain embodiments WR^{W4} is hydrogen and WR^{W2} is C1-C6 straight or branched alkyl. In certain embodiments, WR^{W2} is selected from methyl, ethyl, propyl, n-butyl, sec-butyl, t-butyl, or n-pentyl.

[0103] In certain embodiments each of WR^{W2} and WR^{W4} is C1-C6 straight or branched alkyl. In certain embodiments, each of WR^{W2} and WR^{W4} is selected from methyl, ethyl, propyl, n-butyl, sec-butyl, t-butyl, or pentyl.

[0104] In one embodiment, WR^{W5} is selected from hydrogen, CHF_2 , NH_2 , CN , NHR' , $N(R')_2$, $CH_2N(R')_2$, $-NHC(O)R'$, $-NHC(O)OR'$, $-OR'$, $C(O)OR'$, or SO_2NHR' . Or, WR^{W5} is $-OR'$, e.g., OH .

[0105] In certain embodiments, WR^{W5} is selected from hydrogen, NH_2 , CN , CHF_2 , $NH(C1-C6 \text{ alkyl})$, $N(C1-C6 \text{ alkyl})_2$, $-NHC(O)(C1-C6 \text{ alkyl})$, $-CH_2NHC(O)O(C1-C6 \text{ alkyl})$, $-NHC(O)O(C1-C6 \text{ alkyl})$, $-OH$, $-O(C1-C6 \text{ alkyl})$, $C(O)O(C1-C6 \text{ alkyl})$, $CH_2O(C1-C6 \text{ alkyl})$, or SO_2NH_2 . In another embodiment, WR^{W5} is selected from $-OH$, OMe , NH_2 , $-NHMe$, $-N(Me)_2$, $-CH_2NH_2$, CH_2OH , $NHC(O)OMe$, $NHC(O)OEt$, CN , CHF_2 , $-CH_2NHC(O)O(t\text{-butyl})$, $-O\text{-(ethoxyethyl)}$, $-O\text{-(hydroxyethyl)}$, $-C(O)OMe$, or $-SO_2NH_2$.

[0106] In one embodiment, compound of formula VA-1 has one, preferably more, or more preferably all, of the following features:

- i) WR^{W2} is hydrogen;
- ii) WR^{W4} is C1-C6 straight or branched alkyl or monocyclic or bicyclic aliphatic; and
- iii) WR^{W5} is selected from hydrogen, CN , CHF_2 , NH_2 , $NH(C1-C6 \text{ alkyl})$, $N(C1-C6 \text{ alkyl})_2$, $-NHC(O)(C1-C6 \text{ alkyl})$, $-NHC(O)O(C1-C6 \text{ alkyl})$, $-CH_2C(O)O(C1-C6 \text{ alkyl})$, $-OH$, $-O(C1-C6 \text{ alkyl})$, $C(O)O(C1-C6 \text{ alkyl})$, or SO_2NH_2 .

[0107] In one embodiment, compound of formula VA-1 has one, preferably more, or more preferably all, of the following features:

- i) WR^{W2} is halo, C1-C6 alkyl, CF_3 , CN, or phenyl optionally substituted with up to 3 substituents selected from C1-C4 alkyl, $-O(C1-C4 \text{ alkyl})$, or halo;
- ii) WR^{W4} is CF_3 , halo, C1-C6 alkyl, or C6-C10 cycloaliphatic; and
- iii) WR^{W5} is OH, NH_2 , $NH(C1-C6 \text{ alkyl})$, or $N(C1-C6 \text{ alkyl})$.


[0108] In one embodiment, $X-R^X$ is at the 6-position of the quinolinyl ring. In certain embodiments, $X-R^X$ taken together is C1-C6 alkyl, $-O-(C1-C6 \text{ alkyl})$, or halo.

[0109] In one embodiment, $X-R^X$ is at the 5-position of the quinolinyl ring. In certain embodiments, $X-R^X$ taken together is -OH.

[0110] In another embodiment, the present invention provides compounds of formula VA-1, wherein WR^{W4} and WR^{W5} taken together form a 5-7 membered ring containing 0-3 three heteroatoms selected from N, O, or S, wherein said ring is optionally substituted with up to three WR^W substituents.

[0111] In certain embodiments, WR^{W4} and WR^{W5} taken together form an optionally substituted 5-7 membered saturated, unsaturated, or aromatic ring containing 0 heteroatoms. In other embodiments, WR^{W4} and WR^{W5} taken together form an optionally substituted 5-7 membered ring containing 1-3 heteroatoms selected from N, O, or S. In certain other embodiments, WR^{W4} and WR^{W5} taken together form an optionally substituted saturated, unsaturated, or aromatic 5-7 membered ring containing 1 nitrogen heteroatom. In certain other embodiments, WR^{W4} and WR^{W5} taken together form an optionally substituted 5-7 membered ring containing 1 oxygen heteroatom.

[0112] In another embodiment, the present invention provides compounds of formula V-A-2:



V-A-2

wherein:

Y is CH_2 , $C(O)O$, $C(O)$, or $S(O)_2$;

m is 0-4; and

X, R^X, W, and R^W are as defined above.

[0113] In one embodiment, compounds of formula VA-2 have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0114] In one embodiment, Y is C(O). In another embodiment, Y is C(O)O. Or, Y is S(O)₂. Or, Y is CH₂.

[0115] In one embodiment, m is 1 or 2. Or, m is 1. Or, m is 0.


[0116] In one embodiment, W is a bond.

[0117] In another embodiment, R^W is C1-C6 aliphatic, halo, CF₃, or phenyl optionally substituted with C1-C6 alkyl, halo, cyano, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.

Exemplary embodiments of WR^W include methyl, ethyl, propyl, tert-butyl, or 2-ethoxyphenyl.

[0118] In another embodiment, R^W in Y-R^W is C1-C6 aliphatic optionally substituted with N(R'')₂, wherein R'' is hydrogen, C1-C6 alkyl, or two R'' taken together form a 5-7 membered heterocyclic ring with up to 2 additional heteroatoms selected from O, S, or NR'. Exemplary such heterocyclic rings include pyrrolidinyl, piperidyl, morpholinyl, or thiomorpholinyl.

[0119] In another embodiment, the present invention provides compounds of formula V-A-3:



V-A-3

wherein:

Q is W;

R^Q is R^W;

m is 0-4;

n is 0-4; and

X, R^X, W, and R^W are as defined above.

[0120] In one embodiment, compounds of formula VA-3 have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.


[0121] In one embodiment, n is 0-2.

[0122] In another embodiment, m is 0-2. In one embodiment, m is 0. In one embodiment, m is 1. Or, m is 2.

[0123] In one embodiment, QR^Q taken together is halo, CF₃, OCF₃, CN, C1-C6 aliphatic, O-C1-C6 aliphatic, O-phenyl, NH(C1-C6 aliphatic), or N(C1-C6 aliphatic)₂, wherein said aliphatic and phenyl are optionally substituted with up to three substituents selected from C1-C6 alkyl, O-C1-C6 alkyl, halo, cyano, OH, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, SOR', SO₂R', -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.

[0124] Exemplary QR^Q include methyl, isopropyl, sec-butyl, hydroxymethyl, CF₃, NMe₂, CN, CH₂CN, fluoro, chloro, OEt, OMe, SMe, OCF₃, OPh, C(O)OMe, C(O)O-iPr, S(O)Me, NHC(O)Me, or S(O)₂Me.

[0125] In another embodiment, the present invention provides compounds of formula V-A-4:



V-A-4

wherein X, R^X, and R^W are as defined above.


[0126] In one embodiment, compounds of formula VA-4 have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0127] In one embodiment, R^W is C1-C12 aliphatic, C5-C10 cycloaliphatic, or C5-C7 heterocyclic ring, wherein said aliphatic, cycloaliphatic, or heterocyclic ring is optionally substituted with up to three substituents selected from C1-C6 alkyl, halo, cyano, oxo, OH, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally

replaced with $-\text{CO}-$, $-\text{CONR}'-$, $-\text{CO}_2-$, $-\text{OCO}-$, $-\text{NR}'\text{CO}_2-$, $-\text{O}-$, $-\text{NR}'\text{CONR}'-$, $-\text{OCONR}'-$, $-\text{NR}'\text{CO}-$, $-\text{S}-$, $-\text{NR}'-$, $-\text{SO}_2\text{NR}'-$, $\text{NR}'\text{SO}_2-$, or $-\text{NR}'\text{SO}_2\text{NR}'-$. In another embodiment, R' above is C1-C4 alkyl.

[0128] Exemplary R^{W} includes methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, t-butyl, n-pentyl, vinyl, cyanomethyl, hydroxymethyl, hydroxyethyl, hydroxybutyl, cyclohexyl, adamantyl, or $-\text{C}(\text{CH}_3)_2\text{-NHC}(\text{O})\text{O-T}$, wherein T is C1-C4 alkyl, methoxyethyl, or tetrahydrofuranylmethyl.

[0129] In another embodiment, the present invention provides compounds of formula V-A-5:



wherein:

m is 0-4; and

X , R^{X} , W , R^{W} , and R' are as defined above.

[0130] In one embodiment, compounds of formula V-A-5 have y occurrences of X-R^{X} , wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.


[0131] In one embodiment, m is 0-2. Or, m is 1. Or, m is 2.

[0132] In another embodiment, both R' are hydrogen. Or, one R' is hydrogen and the other R' is C1-C4 alkyl, e.g., methyl. Or, both R' are C1-C4 alkyl, e.g., methyl.

[0133] In another embodiment, m is 1 or 2, and R^{W} is halo, CF_3 , CN, C1-C6 aliphatic, O-C1-C6 aliphatic, or phenyl, wherein said aliphatic and phenyl are optionally substituted with up to three substituents selected from C1-C6 alkyl, O-C1-C6 alkyl, halo, cyano, OH, or CF_3 , wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with $-\text{CO}-$, $-\text{CONR}'-$, $-\text{CO}_2-$, $-\text{OCO}-$, $-\text{NR}'\text{CO}_2-$, $-\text{O}-$, $-\text{NR}'\text{CONR}'-$, $-\text{OCONR}'-$, $-\text{NR}'\text{CO}-$, $-\text{S}-$, $-\text{NR}'-$, $-\text{SO}_2\text{NR}'-$, $\text{NR}'\text{SO}_2-$, or $-\text{NR}'\text{SO}_2\text{NR}'-$. In another embodiment, R' above is C1-C4 alkyl.

[0134] Exemplary embodiments of R^{W} include chloro, CF_3 , OCF_3 , methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, methoxy, ethoxy, propyloxy, or 2-ethoxyphenyl.

[0135] In another embodiment, the present invention provides compounds of formula V-A-6:



V-A-6

wherein:

ring B is a 5-7 membered monocyclic or bicyclic, heterocyclic or heteroaryl ring optionally substituted with up to n occurrences of $-Q-R^Q$, wherein n is 0-4, and Q and R^Q are as defined above; and

Q, R^Q , X, R^X , W, and R^W are as defined above.

[0136] In one embodiment, compounds of formula V-A-6 have y occurrences of $X-R^X$, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.


[0137] In one embodiment, m is 0-2. Or, m is 0. Or m is 1.

[0138] In one embodiment, n is 0-2. Or, n is 0. Or, n is 1.

[0139] In another embodiment, ring B is a 5-7 membered monocyclic, heterocyclic ring having up to 2 heteroatoms selected from O, S, or N, optionally substituted with up to n occurrences of $-Q-R^Q$. Exemplary heterocyclic rings include N-morpholinyl, N-piperidinyl, 4-benzoyl-piperazin-1-yl, pyrrolidin-1-yl, or 4-methyl-piperidin-1-yl.

[0140] In another embodiment, ring B is a 5-6 membered monocyclic, heteroaryl ring having up to 2 heteroatoms selected from O, S, or N, optionally substituted with up to n occurrences of $-Q-R^Q$. Exemplary such rings include benzimidazol-2-yl, 5-methyl-furan-2-yl, 2,5-dimethyl-pyrrol-1-yl, pyridine-4-yl, indol-5-yl, indol-2-yl, 2,4-dimethoxy-pyrimidin-5-yl, furan-2-yl, furan-3-yl, 2-acyl-thien-2-yl, benzothiophen-2-yl, 4-methyl-thien-2-yl, 5-cyano-thien-2-yl, 3-chloro-5-trifluoromethyl-pyridin-2-yl.

[0141] In another embodiment, the present invention provides compounds of formula V-B-1:

**V-B-1**

wherein:

one of Q_1 and Q_3 is $N(WR^W)$ and the other of Q_1 and Q_3 is selected from O, S, or $N(WR^W)$;

Q_2 is $C(O)$, $CH_2-C(O)$, $C(O)-CH_2$, CH_2 , CH_2-CH_2 , CF_2 , or CF_2-CF_2 ;

m is 0-3; and


X , W , R^X , and R^W are as defined above.

[0142] In one embodiment, compounds of formula **V-B-1** have y occurrences of $X-R^X$, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0143] In one embodiment, Q_3 is $N(WR^W)$; exemplary WR^W include hydrogen, C1-C6 aliphatic, $C(O)C1-C6$ aliphatic, or $C(O)OC1-C6$ aliphatic.

[0144] In another embodiment, Q_3 is $N(WR^W)$, Q_2 is $C(O)$, CH_2 , CH_2-CH_2 , and Q_1 is O.

[0145] In another embodiment, the present invention provides compounds of formula **V-B-2**:

**V-B-2**

wherein:

R^{W1} is hydrogen or C1-C6 aliphatic;

each of R^{W3} is hydrogen or C1-C6 aliphatic; or

both R^{W3} taken together form a C3-C6 cycloalkyl or heterocyclic ring having up to two heteroatoms selected from O, S, or NR' , wherein said ring is optionally substituted with up to two WR^W substituents;

m is 0-4; and


X, R^X, W, and R^W are as defined above.

[0146] In one embodiment, compounds of formula **V-B-2** have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0147] In one embodiment, WR^{W1} is hydrogen, C1-C6 aliphatic, C(O)C1-C6 aliphatic, or C(O)OC1-C6 aliphatic.

[0148] In another embodiment, each R^{W3} is hydrogen, C1-C4 alkyl. Or, both R^{W3} taken together form a C3-C6 cycloaliphatic ring or 5-7 membered heterocyclic ring having up to two heteroatoms selected from O, S, or N, wherein said cycloaliphatic or heterocyclic ring is optionally substituted with up to three substituents selected from WR^{W1}. Exemplary such rings include cyclopropyl, cyclopentyl, optionally substituted piperidyl, etc.

[0149] In another embodiment, the present invention provides compounds of formula **V-B-3**:



V-B-3

wherein:

Q₄ is a bond, C(O), C(O)O, or S(O)₂;

R^{W1} is hydrogen or C1-C6 aliphatic;


m is 0-4; and

X, W, R^W, and R^X are as defined above.

[0150] In one embodiment, compounds of formula **V-B-3** have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0.

[0151] In one embodiment, Q₄ is C(O). Or Q₄ is C(O)O. In another embodiment, R^{W1} is C1-C6 alkyl. Exemplary R^{W1} include methyl, ethyl, or t-butyl.

[0152] In another embodiment, the present invention provides compounds of formula **V-B-4**:

**V-B-4**

wherein:

m is 0-4; and


X, R^X, W, and R^W are as defined above.

[0153] In one embodiment, compounds of formula **V-B-4** have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0154] In one embodiment, m is 0-2. Or, m is 0. Or, m is 1.

[0155] In another embodiment, said cycloaliphatic ring is a 5-membered ring. Or, said ring is a six-membered ring.

[0156] In another embodiment, the present invention provides compounds of formula **V-B-5**:

**V-B-5**

wherein:

ring A₂ is a phenyl or a 5-6 membered heteroaryl ring, wherein ring A₂ and the phenyl ring fused thereto together have up 4 substituents independently selected from WR^W;


m is 0-4; and

X, W, R^W and R^X are as defined above.

[0157] In one embodiment, compounds of formula **V-B-5** have y occurrences of X-R^X, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0158] In one embodiment, ring A₂ is an optionally substituted 5-membered ring selected from pyrrolyl, furanyl, thienyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, thiadiazolyl, oxadiazolyl, or triazolyl.

[0159] In one embodiment, ring A₂ is an optionally substituted 5-membered ring selected from pyrrolyl, pyrazolyl, thiadiazolyl, imidazolyl, oxazolyl, or triazolyl. Exemplary such rings include:



wherein said ring is optionally substituted as set forth above.

[0160] In another embodiment, ring A₂ is an optionally substituted 6-membered ring. Exemplary such rings include pyridyl, pyrazinyl, or triazinyl. In another embodiment, said ring is an optionally pyridyl.


[0161] In one embodiment, ring A₂ is phenyl.

[0162] In another embodiment, ring A₂ is pyrrolyl, pyrazolyl, pyridyl, or thiadiazolyl.

[0163] Exemplary W in formula V-B-5 includes a bond, C(O), C(O)O or C1-C6 alkylene.

[0164] Exemplary R^W in formula V-B-5 include cyano, halo, C1-C6 aliphatic, C3-C6 cycloaliphatic, aryl, 5-7 membered heterocyclic ring having up to two heteroatoms selected from O, S, or N, wherein said aliphatic, phenyl, and heterocyclic are independently and optionally substituted with up to three substituents selected from C1-C6 alkyl, O-C1-C6 alkyl, halo, cyano, OH, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.

[0165] In one embodiment, the present invention provides compounds of formula V-B-5-a:



V-B-5-a

wherein:

G_4 is hydrogen, halo, CN, CF_3 , CHF_2 , CH_2F , optionally substituted C1-C6 aliphatic, aryl-C1-C6 alkyl, or a phenyl, wherein G_4 is optionally substituted with up to 4 WR^W substituents; wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with $-CO-$, $-CONR'-$, $-CO_2-$, $-OCO-$, $-NR'CO_2-$, $-O-$, $-NR'CONR'-$, $-OCONR'-$, $-NR'CO-$, $-S-$, $-NR'-$, $-SO_2NR'-$, $NR'SO_2-$, or $-NR'SO_2NR'-$. ;

G_5 is hydrogen or an optionally substituted C1-C6 aliphatic;

wherein said indole ring system is further optionally substituted with up to 3 substituents independently selected from WR^W .

[0166] In one embodiment, compounds of formula V-B-5-a have y occurrences of $X-R^X$, wherein y is 0-4. In one embodiment, y is 0. Or, y is 1. Or, y is 2.

[0167] In one embodiment, G_4 is hydrogen. Or, G_5 is hydrogen.

[0168] In another embodiment, G_4 is hydrogen, and G_5 is C1-C6 aliphatic, wherein said aliphatic is optionally substituted with C1-C6 alkyl, halo, cyano, or CF_3 , and wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with $-CO-$, $-CONR'-$, $-CO_2-$, $-OCO-$, $-NR'CO_2-$, $-O-$, $-NR'CONR'-$, $-OCONR'-$, $-NR'CO-$, $-S-$, $-NR'-$, $-SO_2NR'-$, $NR'SO_2-$, or $-NR'SO_2NR'-$. In another embodiment, R' above is C1-C4 alkyl.

[0169] In another embodiment, G_4 is hydrogen, and G_5 is cyano, methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, t-butyl, cyanomethyl, methoxyethyl, $CH_2C(O)OMe$, $(CH_2)_2-NHC(O)O-tert-butyl$, or cyclopentyl.


[0170] In another embodiment, G_5 is hydrogen, and G_4 is halo, C1-C6 aliphatic or phenyl, wherein said aliphatic or phenyl is optionally substituted with C1-C6 alkyl, halo, cyano, or CF_3 , wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with $-CO-$, $-CONR'-$, $-CO_2-$, $-OCO-$, $-NR'CO_2-$, $-O-$, $-NR'CONR'-$, $-OCONR'-$, $-NR'CO-$, $-S-$, $-NR'-$, $-SO_2NR'-$, $NR'SO_2-$, or $-NR'SO_2NR'-$. In another embodiment, R' above is C1-C4 alkyl.

[0171] In another embodiment, G₅ is hydrogen, and G₄ is halo, CF₃, ethoxycarbonyl, t-butyl, 2-methoxyphenyl, 2-ethoxyphenyl, (4-C(O)NH(CH₂)₂-NMe₂)-phenyl, 2-methoxy-4-chloro-phenyl, pyridine-3-yl, 4-isopropylphenyl, 2,6-dimethoxyphenyl, sec-butylaminocarbonyl, ethyl, t-butyl, or piperidin-1-ylcarbonyl.

[0172] In another embodiment, G₄ and G₅ are both hydrogen, and the nitrogen ring atom of said indole ring is substituted with C1-C6 aliphatic, C(O)(C1-C6 aliphatic), or benzyl, wherein said aliphatic or benzyl is optionally substituted with C1-C6 alkyl, halo, cyano, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.

[0173] In another embodiment, G₄ and G₅ are both hydrogen, and the nitrogen ring atom of said indole ring is substituted with acyl, benzyl, C(O)CH₂N(Me)C(O)CH₂NHMe, or ethoxycarbonyl.

[0174] In another embodiment, the present invention provides compounds of formula I':



or pharmaceutically acceptable salts thereof,

wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and Ar¹ is as defined above for compounds of formula I'.

[0175] In one embodiment, each of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, and Ar¹ in compounds of formula I' is independently as defined above for any of the embodiments of compounds of formula I.

[0176] Representative compounds of the present invention are set forth below in Table 1 below.

[0177] Table 1

Cmpd No.	Name
1	N-[5-(5-chloro-2-methoxy-phenyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
2	N-(3-methoxy-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
3	N-[2-(2-methoxyphenoxy)-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
4	N-(2-morpholinophenyl)-4-oxo-1H-quinoline-3-carboxamide
5	N-[4-(2-hydroxy-1,1-dimethyl-ethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
6	N-[3-(hydroxymethyl)-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
7	N-(4-benzoylamino-2,5-diethoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
8	N-(3-amino-4-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
9	4-oxo-N-(3-sulfamoylphenyl)-1H-quinoline-3-carboxamide
10	1,4-dihydro-N-(2,3,4,5-tetrahydro-1H-benzo[b]azepin-8-yl)-4-oxoquinoline-3-carboxamide
11	4-oxo-N-[2-[2-(trifluoromethyl)phenyl]phenyl]-1H-quinoline-3-carboxamide
12	N-[2-(4-dimethylaminophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
13	N-(3-cyano-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
14	[5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2- <i>tert</i> -butyl-phenyl]aminoformic acid methyl ester
15	N-(2-methoxy-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
16	4-oxo-N-(2-propylphenyl)-1H-quinoline-3-carboxamide
17	N-(5-amino-2-propoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
18	N-(9H-fluoren-1-yl)-4-oxo-1H-quinoline-3-carboxamide
19	4-oxo-N-(2-quinolyl)-1H-quinoline-3-carboxamide
20	N-[2-(2-methylphenoxy)phenyl]-4-oxo-1H-quinoline-3-carboxamide
21	4-oxo-N-[4-(2-pyridylsulfamoyl)phenyl]-1H-quinoline-3-carboxamide
22	4-Oxo-1,4-dihydro-quinoline-3-carboxylic acid N-(1',2'-dihydrospiro[cyclopropane-1,3'-[3H]indol]-6'-yl)-amide
23	N-[2-(2-ethoxyphenyl)-5-hydroxy-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
24	4-oxo-N-(3-pyrrolidin-1-ylsulfonylphenyl)-1H-quinoline-3-carboxamide

25	N-[2-(3-acetylamino-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
26	4-oxo-N-[2-(1-piperidyl)phenyl]-1H-quinoline-3-carboxamide
27	N-[1-[2-[methyl-(2-methylaminoacetyl)-amino]acetyl]-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
28	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid 2-methoxyethyl ester
29	1-isopropyl-4-oxo-N-phenyl-1H-quinoline-3-carboxamide
30	[2-isopropyl-5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid methyl ester
31	4-oxo-N-(p-tolyl)-1H-quinoline-3-carboxamide
32	N-(5-chloro-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
33	N-(1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
34	N-[4-(1,1-diethylpropyl)-2-fluoro-5-hydroxy-phenyl]-4-hydroxy-quinoline-3-carboxamide
35	1,4-dihydro-N-(2,3,4,5-tetrahydro-5,5-dimethyl-1H-benzo[b]azepin-8-yl)-4-oxoquinoline-3-carboxamide
36	N-(2-isopropylphenyl)-4-oxo-1H-quinoline-3-carboxamide
37	N-(1H-indol-7-yl)-4-oxo-1H-quinoline-3-carboxamide
38	N-[2-(1H-indol-2-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
39	[3-[(2,4-dimethoxy-3-quinolyl)carbonylamino]-4- <i>tert</i> -butyl-phenyl]aminoformic acid <i>tert</i> -butyl ester
40	N-[2-(2-hydroxyethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
41	N-(5-amino-2-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
42	N-[2-[[3-chloro-5-(trifluoromethyl)-2-pyridyl]oxy]phenyl]-4-oxo-1H-quinoline-3-carboxamide
43	N-[2-(3-ethoxyphenyl)-5-hydroxy-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
44	N-(2-methylbenzothiazol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
45	N-(2-cyano-3-fluoro-phenyl)-4-oxo-1H-quinoline-3-carboxamide
46	N-[3-chloro-5-(2-morpholinoethylsulfonylamino)phenyl]-4-oxo-1H-quinoline-3-carboxamide
47	N-[4-isopropyl-2-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
48	N-(5-chloro-2-fluoro-phenyl)-4-oxo-1H-quinoline-3-carboxamide
49	N-[2-(2,6-dimethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide

50	4-oxo-N-(2,4,6-trimethylphenyl)-1H-quinoline-3-carboxamide
51	6-[(4-methyl-1-piperidyl)sulfonyl]-4-oxo-N-(5- <i>tert</i> -butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
52	N-[2-(<i>m</i> -tolyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
53	4-oxo-N-(4-pyridyl)-1H-quinoline-3-carboxamide
54	4-oxo-N-(8-thia-7,9-diazabicyclo[4.3.0]nona-2,4,6,9-tetraen-5-yl)-1H-quinoline-3-carboxamide
55	N-(3-amino-2-methoxy-5- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
56	1,4-dihydro-N-(1,2,3,4-tetrahydro-6-hydroxynaphthalen-7-yl)-4-oxoquinoline-3-carboxamide
57	N-[4-(3-ethyl-2,6-dioxo-3-piperidyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
58	N-[3-amino-4-(trifluoromethoxy)phenyl]-4-oxo-1H-quinoline-3-carboxamide
59	N-[2-(5-isopropyl-2-methoxy-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
60	[4-isopropyl-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid <i>tert</i> -butyl ester
61	N-(2,3-dimethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
62	4-oxo-N-[3-(trifluoromethoxy)phenyl]-1H-quinoline-3-carboxamide
63	N-[2-(2,4-difluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
64	4-oxo-N-(2-oxo-1,3-dihydrobenzoimidazol-5-yl)-1H-quinoline-3-carboxamide
65	4-oxo-N-[5-(3-pyridyl)-1H-indol-6-yl]-1H-quinoline-3-carboxamide
66	N-(2,2-difluorobenzo[1,3]dioxol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
67	6-ethyl-4-hydroxy-N-(1H-indol-6-yl)quinoline-3-carboxamide
68	3-[2-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]benzoic acid methyl ester
69	N-(3-amino-4-isopropyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
70	4-oxo-N-[2-(4-pyridyl)phenyl]-1H-quinoline-3-carboxamide
71	3-[2-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]benzoic acid isopropyl ester
72	N-(2-ethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
73	4-oxo-N-(2-phenyl-3H-benzoimidazol-5-yl)-1H-quinoline-3-carboxamide
74	4-oxo-N-[5-(trifluoromethyl)-2-pyridyl]-1H-quinoline-3-carboxamide

75	4-oxo-N-(3-quinolyl)-1H-quinoline-3-carboxamide
76	N-[2-(3,4-difluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
77	N-(5-fluoro-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
78	4-oxo-N-(2-sulfamoylphenyl)-1H-quinoline-3-carboxamide
79	N-[2-(4-fluoro-3-methyl-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
80	N-(2-methoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
81	4-oxo-N-(3-propionylaminophenyl)-1H-quinoline-3-carboxamide
82	N-(4-diethylamino-2-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
83	N-[2-(3-cyanophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
84	N-(4-methyl-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
85	N-[2-(3,4-dichlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
86	N-[4-[2-(aminomethyl)phenyl]phenyl]-4-oxo-1H-quinoline-3-carboxamide
87	4-oxo-N-(3-phenoxyphenyl)-1H-quinoline-3-carboxamide
88	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid tert-butyl ester
89	N-(2-cyano-5-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
90	4-oxo-N-(2- <i>tert</i> -butylphenyl)-1H-quinoline-3-carboxamide
91	N-(3-chloro-2,6-diethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
92	N-[2-fluoro-5-hydroxy-4-(1-methylcyclohexyl)-phenyl]-4-oxo-1H-quinoline-3-carboxamide
93	N-[2-(5-cyano-2-thienyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
94	N-(5-amino-2-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
95	N-(2-cyanophenyl)-4-oxo-1H-quinoline-3-carboxamide
96	N-[3-(cyanomethyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
97	N-[2-(2,4-dimethoxypyrimidin-5-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
98	N-(5-dimethylamino-2-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
99	4-oxo-N-(4-pentylphenyl)-1H-quinoline-3-carboxamide

100	N-(1H-indol-4-yl)-4-oxo-1H-quinoline-3-carboxamide
101	N-(5-amino-2-isopropyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
102	N-[2-[3-(4-chlorophenyl)-1,2,4-oxadiazol-5-yl]phenyl]-4-oxo-1H-quinoline-3-carboxamide
103	6-fluoro-N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
104	N-(2-methyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
105	1,4-dihydro-N-(3,4-dihydro-2H-benzo[b][1,4]oxazin-6-yl)-4-oxoquinoline-3-carboxamide
106	N-(2-cyano-4,5-dimethoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
107	7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1,2,3,4-tetrahydroisoquinoline-2-carboxylic acid tert-butyl ester
108	4,4-dimethyl-7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1,2,3,4-tetrahydroquinoline-1-carboxylic acid tert-butyl ester
109	N-(1-acetyl-2,3,4,5-tetrahydro-5,5-dimethyl-1H-benzo[b]azepin-8-yl)-1,4-dihydro-4-oxoquinoline-3-carboxamide
110	N-[4-(cyanomethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
111	4-oxo-N-[2-(trifluoromethyl)phenyl]-1H-quinoline-3-carboxamide
112	6-ethoxy-4-hydroxy-N-(1H-indol-6-yl)quinoline-3-carboxamide
113	N-(3-methyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
114	[4-(2-ethoxyphenyl)-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid tert-butyl ester
115	N-[2-(2-furyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
116	5-hydroxy-N-(1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
117	N-(3-dimethylamino-4-isopropyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
118	N-[2-(1H-indol-5-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
119	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid ethyl ester
120	N-(2-methoxy-5-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
121	N-(3,4-dichlorophenyl)-4-oxo-1H-quinoline-3-carboxamide
122	N-(3,4-dimethoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
123	N-[2-(3-furyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
124	6-fluoro-4-oxo-N-(5-tert-butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide

125	N-(6-ethyl-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
126	N-[3-hydroxy-4-[2-(2-methoxyethoxy)-1,1-dimethyl-ethyl]-phenyl]-4-oxo-1H-quinoline-3-carboxamide
127	[5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2- <i>tert</i> -butyl-phenyl]aminoformic acid ethyl ester
128	1,6-dimethyl-4-oxo-N-(2-phenylphenyl)-1H-quinoline-3-carboxamide
129	[2-ethyl-5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid methyl ester
130	4-hydroxy-N-(1H-indol-6-yl)-5,7-bis(trifluoromethyl)quinoline-3-carboxamide
131	N-(3-amino-5-chloro-phenyl)-4-oxo-1H-quinoline-3-carboxamide
132	N-(5-acetylamino-2-ethoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
133	N-[3-chloro-5-[2-(1-piperidyl)ethylsulfonylamino]phenyl]-4-oxo-1H-quinoline-3-carboxamide
134	N-[2-(4-methylsulfinylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
135	N-(2-benzo[1,3]dioxol-5-ylphenyl)-4-oxo-1H-quinoline-3-carboxamide
136	N-(2-hydroxy-3,5-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
137	6-[(4-fluorophenyl)-methyl-sulfamoyl]-N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
138	N-[2-(3,5-difluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
139	N-[2-(2,4-dichlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
140	N-(4-cyclohexylphenyl)-4-oxo-1H-quinoline-3-carboxamide
141	[2-methyl-5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid ethyl ester
142	4-oxo-N-(2-sec-butylphenyl)-1H-quinoline-3-carboxamide
143	N-(2-fluoro-5-hydroxy-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
144	N-(3-hydroxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
145	6-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1H-indole-4-carboxylic acid ethyl ester
146	4-oxo-N-(1,7,9-triazabicyclo[4.3.0]nona-2,4,6,8-tetraen-5-yl)-1H-quinoline-3-carboxamide
147	N-[2-(4-fluorophenoxy)-3-pyridyl]-4-oxo-1H-quinoline-3-carboxamide
148	4-oxo-N-[5-(1-piperidylcarbonyl)-1H-indol-6-yl]-1H-quinoline-3-carboxamide
149	N-(3-acetylamino-4-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide

150	4-oxo-N-[4-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]phenyl]-1H-quinoline-3-carboxamide
151	N-[2-(4-methyl-2-thienyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
152	4-oxo-N-(2-oxo-3H-benzooxazol-6-yl)-1H-quinoline-3-carboxamide
153	N-[4-(1,1-diethyl-2,2-dimethyl-propyl)-2-fluoro-5-hydroxy-phenyl]-4-hydroxy-quinoline-3-carboxamide
154	N-[3,5-bis(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
155	4-oxo-N-(2-pyridyl)-1H-quinoline-3-carboxamide
156	4-oxo-N-[2-[2-(trifluoromethoxy)phenyl]phenyl]-1H-quinoline-3-carboxamide
157	N-(2-ethyl-5-methylamino-phenyl)-4-oxo-1H-quinoline-3-carboxamide
158	4-oxo-N-(5-phenyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
159	[7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]tetralin-1-yl]aminoformic acid methyl ester
160	N-(3-amino-4-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
161	N-[3-(2-ethoxyethoxy)-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
162	N-(6-methoxy-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
163	N-[5-(aminomethyl)-2-(2-ethoxyphenyl)-phenyl]-4-oxo-1H-quinoline-3-carboxamide
164	4-oxo-N-[3-(trifluoromethyl)phenyl]-1H-quinoline-3-carboxamide
165	4-oxo-N-(4-sulfamoylphenyl)-1H-quinoline-3-carboxamide
166	4-[2-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]benzoic acid methyl ester
167	N-(3-amino-4-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
168	4-oxo-N-(3-pyridyl)-1H-quinoline-3-carboxamide
169	N-(1-methyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
170	N-(5-chloro-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
171	N-[2-(2,3-dichlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
172	N-(2-(benzo[b]thiophen-2-yl)phenyl)-1,4-dihydro-4-oxoquinoline-3-carboxamide
173	N-(6-methyl-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
174	N-[2-(5-acetyl-2-thienyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide

175	4-Oxo-1,4-dihydro-quinoline-3-carboxylic acid N-(1'-Acetyl-1',2'-dihydrospiro[cyclopropane-1,3'-3H-indol]-6'-yl)-amide
176	4-oxo-N-[4-(trifluoromethoxy)phenyl]-1H-quinoline-3-carboxamide
177	N-(2-butoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
178	4-oxo-N-[2-(2- <i>tert</i> -butylphenoxy)phenyl]-1H-quinoline-3-carboxamide
179	N-(3-carbamoylphenyl)-4-oxo-1H-quinoline-3-carboxamide
180	N-(2-ethyl-6-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
181	4-oxo-N-[2-(<i>p</i> -tolyl)phenyl]-1H-quinoline-3-carboxamide
182	N-[2-(4-fluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
183	7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1,2,3,4-tetrahydroquinoline-1-carboxylic acid <i>tert</i> -butyl ester
184	N-(1H-indol-6-yl)-4-oxo-2-(trifluoromethyl)-1H-quinoline-3-carboxamide
185	N-(3-morpholinosulfonylphenyl)-4-oxo-1H-quinoline-3-carboxamide
186	N-(3-cyclopentyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
187	N-(1-acetyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
188	6-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1H-indole-5-carboxylic acid ethyl ester
189	N-(4-benzyloxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
190	N-[2-(3-chloro-4-fluoro-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
191	4-oxo-N-(5-quinolyl)-1H-quinoline-3-carboxamide
192	N-(3-methyl-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
193	N-(2,6-dimethoxy-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
194	N-(4-cyanophenyl)-4-oxo-1H-quinoline-3-carboxamide
195	N-(5-methyl-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
196	N-[5-(3,3-dimethylbutanoylamino)-2- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
197	4-oxo-N-[6-(trifluoromethyl)-3-pyridyl]-1H-quinoline-3-carboxamide
198	N-(4-fluorophenyl)-4-oxo-1H-quinoline-3-carboxamide
199	N-[2-(<i>o</i> -tolyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide

200	1,4-dihydro-N-(1,2,3,4-tetrahydro-1-hydroxynaphthalen-7-yl)-4-oxoquinoline-3-carboxamide
201	N-(2-cyano-3-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
202	N-[2-(5-chloro-2-methoxy-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
203	N-(1-benzyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
204	N-(4,4-dimethylchroman-7-yl)-4-oxo-1H-quinoline-3-carboxamide
205	N-[2-(4-methoxyphenoxy)-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
206	N-[2-(2,3-dimethylphenoxy)-3-pyridyl]-4-oxo-1H-quinoline-3-carboxamide
207	2-[6-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1H-indol-3-yl]acetic acid ethyl ester
208	N-[4-(2-adamantyl)-5-hydroxy-2-methyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
209	N-[4-(hydroxymethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
210	2,4-dimethoxy-N-(2-phenylphenyl)-quinoline-3-carboxamide
211	N-(2-methoxy-5- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
212	N-[3-(3-methyl-5-oxo-1,4-dihydropyrazol-1-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
213	N-[2-(2,5-dichlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
214	N-(3-methylsulfonylaminophenyl)-4-oxo-1H-quinoline-3-carboxamide
215	4-oxo-N-phenyl-1H-quinoline-3-carboxamide
216	N-(3H-benzoimidazol-2-yl)-4-oxo-1H-quinoline-3-carboxamide
217	N-(1H-indazol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
218	6-fluoro-N-[2-fluoro-5-hydroxy-4-(1-methylcyclohexyl)-phenyl]-4-oxo-1H-quinoline-3-carboxamide
219	4-oxo-N-pyrazin-2-yl-1H-quinoline-3-carboxamide
220	N-(2,3-dihydroxy-4,6-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
221	[5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2-propyl-phenyl]aminoformic acid methyl ester
222	N-(3-chloro-2-cyano-phenyl)-4-oxo-1H-quinoline-3-carboxamide
223	N-[2-(4-methylsulfanylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
224	4-oxo-N-[4-[2-[(2,2,2-trifluoroacetyl)aminomethyl]phenyl]phenyl]-1H-quinoline-3-carboxamide

225	[2-isopropyl-5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid ethyl ester
226	4-oxo-N-(4-propylphenyl)-1H-quinoline-3-carboxamide
227	N-[2-(3H-benzoimidazol-2-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
228	N-[2-(hydroxy-phenyl-methyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
229	N-(2-methylsulfanylphenyl)-4-oxo-1H-quinoline-3-carboxamide
230	N-(2-methyl-1H-indol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
231	3-[4-hydroxy-2-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-5- <i>tert</i> -butyl-phenyl]benzoic acid methyl ester
232	N-(5-acetylamino-2-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
233	N-(1-acetylinolin-6-yl)-4-oxo-1H-quinoline-3-carboxamide
234	4-oxo-N-[5-(trifluoromethyl)-1H-indol-6-yl]-1H-quinoline-3-carboxamide
235	N-(6-isopropyl-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
236	4-oxo-N-[4-(trifluoromethyl)phenyl]-1H-quinoline-3-carboxamide
237	N-[5-(2-methoxyphenyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
238	7'-[(4-oxo-1H-quinolin-3-ylcarbonyl)amino]-spiro[piperidine-4,4'(1'H)-quinoline], 2',3'-dihydro-carboxylic acid <i>tert</i> -butyl ester
239	[4-isopropyl-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid methyl ester
240	N-(2-benzyloxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
241	4-oxo-N-(8-quinolyl)-1H-quinoline-3-carboxamide
242	N-(5-amino-2,4-dichloro-phenyl)-4-oxo-1H-quinoline-3-carboxamide
243	N-(5-acetylamino-2-isopropyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
244	4-oxo-N-(6,7,8,9-tetrahydro-5H-carbazol-2-yl)-1H-quinoline-3-carboxamide
245	N-[2-(2,4-dichlorophenoxy)phenyl]-4-oxo-1H-quinoline-3-carboxamide
246	N-(3,4-dimethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
247	4-oxo-N-[2-(2-phenoxyphenyl)phenyl]-1H-quinoline-3-carboxamide
248	N-(3-acetylamino-4-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
249	[4-ethyl-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid methyl ester

250	N-(5-acetylamino-2-methoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
251	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid isobutyl ester
252	N-(2-benzoylphenyl)-4-oxo-1H-quinoline-3-carboxamide
253	4-oxo-N-[2-[3-(trifluoromethoxy)phenyl]phenyl]-1H-quinoline-3-carboxamide
254	6-fluoro-N-(5-fluoro-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
255	N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-6-pyrrolidin-1-ylsulfonyl-1H-quinoline-3-carboxamide
256	N-(1H-benzotriazol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
257	N-(4-fluoro-3-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
258	N-indolin-6-yl-4-oxo-1H-quinoline-3-carboxamide
259	4-oxo-N-(3-sec-butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
260	N-(5-amino-2- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
261	N-[2-(3,4-dimethylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
262	1,4-dihydro-N-(3,4-dihydro-3-oxo-2H-benzo[b][1,4]thiazin-6-yl)-4-oxoquinoline-3-carboxamide
263	N-(4-bromo-2-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
264	N-(2,5-diethoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
265	N-(2-benzylphenyl)-4-oxo-1H-quinoline-3-carboxamide
266	N-[5-hydroxy-4- <i>tert</i> -butyl-2-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
267	4-oxo-N-(4-phenoxyphenyl)-1H-quinoline-3-carboxamide
268	4-oxo-N-(3-sulfamoyl-4- <i>tert</i> -butyl-phenyl)-1H-quinoline-3-carboxamide
269	[4-isopropyl-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid ethyl ester
270	N-(2-cyano-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
271	N-(3-amino-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
272	N-[3-(2-morpholinoethylsulfonylamino)-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
273	[7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]tetralin-1-yl]aminoformic acid <i>tert</i> -butyl ester
274	4-oxo-6-pyrrolidin-1-ylsulfonyl-N-(5- <i>tert</i> -butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide

275	4-benzyloxy-N-(3-hydroxy-4- <i>tert</i> -butyl-phenyl)-quinoline-3-carboxamide
276	N-(4-morpholinosulfonylphenyl)-4-oxo-1H-quinoline-3-carboxamide
277	N-[2-(3-fluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
278	4-oxo-N-[2-[3-(trifluoromethyl)phenyl]phenyl]-1H-quinoline-3-carboxamide
279	N-[2-(2-methylsulfanylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
280	4-oxo-N-(6-quinolyl)-1H-quinoline-3-carboxamide
281	N-(2,4-dimethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
282	N-(5-amino-2-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
283	N-[2-(3-methoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
284	N-(1H-indazol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
285	N-[2-(2,3-difluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
286	1,4-dihydro-N-(1,2,3,4-tetrahydronaphthalen-5-yl)-4-oxoquinoline-3-carboxamide
287	N-[2-fluoro-5-hydroxy-4-(1-methylcyclohexyl)-phenyl]-5-hydroxy-4-oxo-1H-quinoline-3-carboxamide
288	N-(5-fluoro-2-methoxycarbonyloxy-3- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
289	N-(2-fluoro-4-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
290	N-[2-(3-isopropylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
291	N-(2-chloro-5-hydroxy-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
292	N-(5-chloro-2-phenoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
293	4-oxo-N-[2-(1H-pyrrol-1-yl)phenyl]-1H-quinoline-3-carboxamide
294	N-(1H-indol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
295	4-oxo-N-(2-pyrrolidin-1-ylphenyl)-1H-quinoline-3-carboxamide
296	2,4-dimethoxy-N-(2- <i>tert</i> -butylphenyl)-quinoline-3-carboxamide
297	N-[2-(2,5-dimethyl-1H-pyrrol-1-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
298	[2-ethyl-5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid ethyl ester
299	4-oxo-N-(1,2,3,4-tetrahydroquinolin-7-yl)-1H-quinoline-3-carboxamide

300	N-(4,4-dimethyl-1,2,3,4-tetrahydroquinolin-7-yl)-4-oxo-1H-quinoline-3-carboxamide
301	N-[4-(4-methyl-4H-1,2,4-triazol-3-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
302	N-[2-[4-(hydroxymethyl)phenyl]phenyl]-4-oxo-1H-quinoline-3-carboxamide
303	N-(2-acetyl-1,2,3,4-tetrahydroisoquinolin-7-yl)-4-oxo-1H-quinoline-3-carboxamide
304	[4-(2-ethoxyphenyl)-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenylmethyl]aminoformic acid tert-butyl ester
305	N-[2-(4-methoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
306	N-[2-(3-ethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
307	N-[2-(3-chlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
308	N-[2-(cyanomethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
309	N-(3-isoquinolyl)-4-oxo-1H-quinoline-3-carboxamide
310	4-oxo-N-(4-sec-butylphenyl)-1H-quinoline-3-carboxamide
311	N-[2-(5-methyl-2-furyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
312	N-[2-(2,4-dimethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
313	N-[2-(2-fluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
314	N-(2-ethyl-6-isopropyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
315	N-(2,6-dimethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
316	N-(5-acetylamino-2-tert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
317	N-(2,6-dichlorophenyl)-4-oxo-1H-quinoline-3-carboxamide
318	4-oxo-N-[3-[2-(1-piperidyl)ethylsulfonylamino]-5-(trifluoromethyl)phenyl]-1H-quinoline-3- carboxamide
319	6-fluoro-N-(2-fluoro-5-hydroxy-4-tert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
320	4-oxo-N-(2-tert-butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
321	N-[2-(4-benzoylpiperazin-1-yl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
322	N-(2-ethyl-6-sec-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
323	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid methyl ester
324	N-(4-butylphenyl)-4-oxo-1H-quinoline-3-carboxamide

325	N-(2,6-diethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
326	N-[2-(4-methylsulfonylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
327	N-[5-(2-ethoxyphenyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
328	N-(3-acetylphenyl)-4-oxo-1H-quinoline-3-carboxamide
329	N-[2-(o-tolyl)benzooxazol-5-yl]-4-oxo-1H-quinoline-3-carboxamide
330	N-(2-chlorophenyl)-4-oxo-1H-quinoline-3-carboxamide
331	N-(2-carbamoylphenyl)-4-oxo-1H-quinoline-3-carboxamide
332	N-(4-ethynylphenyl)-4-oxo-1H-quinoline-3-carboxamide
333	N-[2-[4-(cyanomethyl)phenyl]phenyl]-4-oxo-1H-quinoline-3-carboxamide
334	7'-[(4-oxo-1H-quinolin-3-ylcarbonyl)amino]-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro- carboxylic acid tert-butyl ester
335	N-(2-carbamoyl-5-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
336	N-(2-butylphenyl)-4-oxo-1H-quinoline-3-carboxamide
337	N-(5-hydroxy-2,4-ditert-butyl-phenyl)-N-methyl-4-oxo-1H-quinoline-3-carboxamide
338	N-(3-methyl-1H-indol-4-yl)-4-oxo-1H-quinoline-3-carboxamide
339	N-(3-cyano-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
340	N-(3-methylsulfonylamino-4-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
341	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid neopentyl ester
342	N-[5-(4-isopropylphenyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
343	N-[5-(isobutylcarbamoyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
344	N-[2-(2-ethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
345	6-fluoro-4-hydroxy-N-(1H-indol-6-yl)quinoline-3-carboxamide
346	4-oxo-N-phenyl-7-(trifluoromethyl)-1H-quinoline-3-carboxamide
347	N-[5-[4-(2-dimethylaminoethylcarbamoyl)phenyl]-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
348	N-[2-(4-ethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
349	4-oxo-N-(2-phenylsulfonylphenyl)-1H-quinoline-3-carboxamide

350	N-(1-naphthyl)-4-oxo-1H-quinoline-3-carboxamide
351	N-(5-ethyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
352	2-[6-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1H-indol-3-yl]ethylaminoformic acid tert-butyl ester
353	[3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-4- <i>tert</i> -butyl-phenyl]aminoformic acid tert-butyl ester
354	N-[2-[(cyclohexyl-methyl-amino)methyl]phenyl]-4-oxo-1H-quinoline-3-carboxamide
355	N-[2-(2-methoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
356	N-(5-methylamino-2-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
357	N-(3-isopropyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
358	6-chloro-4-hydroxy-N-(1H-indol-6-yl)quinoline-3-carboxamide
359	N-[3-(2-dimethylaminoethylsulfonylamino)-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
360	N-[4-(difluoromethoxy)phenyl]-4-oxo-1H-quinoline-3-carboxamide
361	N-[2-(2,5-dimethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
362	N-(2-chloro-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
363	N-[2-(2-fluoro-3-methoxy-phenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
364	N-(2-methyl-8-quinolyl)-4-oxo-1H-quinoline-3-carboxamide
365	N-(2-acetylphenyl)-4-oxo-1H-quinoline-3-carboxamide
366	4-oxo-N-[2-[4-(trifluoromethyl)phenyl]phenyl]-1H-quinoline-3-carboxamide
367	N-[2-(3,5-dichlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
368	N-(3-amino-4-propoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
369	N-(2,4-dichloro-6-cyano-phenyl)-4-oxo-1H-quinoline-3-carboxamide
370	N-(3-chlorophenyl)-4-oxo-1H-quinoline-3-carboxamide
371	4-oxo-N-[2-(trifluoromethylsulfanyl)phenyl]-1H-quinoline-3-carboxamide
372	N-[2-(4-methyl-1-piperidyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
373	N-indan-4-yl-4-oxo-1H-quinoline-3-carboxamide
374	4-hydroxy-N-(1H-indol-6-yl)-2-methylsulfanyl-quinoline-3-carboxamide

375	1,4-dihydro-N-(1,2,3,4-tetrahydronaphthalen-6-yl)-4-oxoquinoline-3-carboxamide
376	4-oxo-N-(2-phenylbenzooxazol-5-yl)-1H-quinoline-3-carboxamide
377	6,8-difluoro-4-hydroxy-N-(1H-indol-6-yl)quinoline-3-carboxamide
378	N-(3-amino-4-methoxy-phenyl)-4-oxo-1H-quinoline-3-carboxamide
379	N-[3-acetylamino-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
380	N-(2-ethoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
381	4-oxo-N-(5- <i>tert</i> -butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
382	[5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2-propyl-phenyl]aminoformic acid ethyl ester
383	N-(3-ethyl-1H-indol-6-yl)-4-oxo-1H-quinoline-3-carboxamide
384	N-[2-(2,5-difluorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
385	N-[2-(2,4-difluorophenoxy)-3-pyridyl]-4-oxo-1H-quinoline-3-carboxamide
386	N-(3,3-dimethylindolin-6-yl)-4-oxo-1H-quinoline-3-carboxamide
387	N-[2-methyl-3-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
388	4-oxo-N-[2-[4-(trifluoromethoxy)phenyl]phenyl]-1H-quinoline-3-carboxamide
389	N-(3-benzylphenyl)-4-oxo-1H-quinoline-3-carboxamide
390	N-[3-(aminomethyl)-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
391	N-[2-(4-isobutylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
392	N-(6-chloro-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
393	N-[5-amino-2-(2-ethoxyphenyl)-phenyl]-4-oxo-1H-quinoline-3-carboxamide
394	1,6-dimethyl-4-oxo-N-phenyl-1H-quinoline-3-carboxamide
395	N-[4-(1-adamantyl)-2-fluoro-5-hydroxy-phenyl]-4-hydroxy-quinoline-3-carboxamide
396	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid tetrahydrofuran-3-ylmethyl ester
397	4-oxo-N-(4-phenylphenyl)-1H-quinoline-3-carboxamide
398	4-oxo-N-[2-(<i>p</i> -tolylsulfonylamino)phenyl]-1H-quinoline-3-carboxamide
399	N-(2-isopropyl-5-methylamino-phenyl)-4-oxo-1H-quinoline-3-carboxamide


400	N-(6-morpholino-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
401	N-[2-(2,3-dimethylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
402	4-oxo-N-(5-phenyl-2-pyridyl)-1H-quinoline-3-carboxamide
403	N-[2-fluoro-5-hydroxy-4-(1-methylcyclooctyl)-phenyl]-4-hydroxy-quinoline-3-carboxamide
404	N-[5-(2,6-dimethoxyphenyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
405	N-(4-chlorophenyl)-4-oxo-1H-quinoline-3-carboxamide
406	6-[(4-fluorophenyl)-methyl-sulfamoyl]-4-oxo-N-(5- <i>tert</i> -butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
407	N-(2-fluoro-5-hydroxy-4- <i>tert</i> -butyl-phenyl)-5-hydroxy-4-oxo-1H-quinoline-3-carboxamide
408	N-(3-methoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
409	N-(5-dimethylamino-2-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
410	4-oxo-N-[2-(4-phenoxyphenyl)phenyl]-1H-quinoline-3-carboxamide
411	7-chloro-4-oxo-N-phenyl-1H-quinoline-3-carboxamide
412	6-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1H-indole-7-carboxylic acid ethyl ester
413	4-oxo-N-(2-phenoxyphenyl)-1H-quinoline-3-carboxamide
414	N-(3H-benzoimidazol-5-yl)-4-oxo-1H-quinoline-3-carboxamide
415	N-(3-hydroxy-4- <i>tert</i> -butyl-phenyl)-4-methoxy-quinoline-3-carboxamide
416	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid propyl ester
417	N-(2-(benzo[b]thiophen-3-yl)phenyl)-1,4-dihydro-4-oxoquinoline-3-carboxamide
418	N-(3-dimethylaminophenyl)-4-oxo-1H-quinoline-3-carboxamide
419	N-(3-acetylaminophenyl)-4-oxo-1H-quinoline-3-carboxamide
420	2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propanoic acid ethyl ester
421	N-[5-methoxy-4- <i>tert</i> -butyl-2-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
422	N-(5,6-dimethyl-3H-benzoimidazol-2-yl)-4-oxo-1H-quinoline-3-carboxamide
423	N-[3-(2-ethoxyethyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide
424	N-[2-(4-chlorophenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide

425	N-(4-isopropylphenyl)-4-oxo-1H-quinoline-3-carboxamide
426	N-(4-chloro-5-hydroxy-2- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
427	5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-1,2,3,4-tetrahydroisoquinoline-2-carboxylic acid <i>tert</i> -butyl ester
428	N-(3-hydroxy-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
429	N-[3-amino-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
430	N-(2-isopropyl-6-methyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
431	N-(3-aminophenyl)-4-oxo-1H-quinoline-3-carboxamide
432	N-[2-(4-isopropylphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
433	N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
434	N-(2,5-dimethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
435	N-[2-(2-fluorophenoxy)-3-pyridyl]-4-oxo-1H-quinoline-3-carboxamide
436	N-[2-(3,4-dimethoxyphenyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
437	N-benzo[1,3]dioxol-5-yl-4-oxo-1H-quinoline-3-carboxamide
438	N-[5-(difluoromethyl)-2,4-ditert-butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
439	N-(4-methoxyphenyl)-4-oxo-1H-quinoline-3-carboxamide
440	N-(2,2,3,3-tetrafluoro-2,3-dihydrobenzo[b][1,4]dioxin-6-yl)-1,4-dihydro-4-oxoquinoline-3-carboxamide
441	N-[3-methylsulfonylamino-5-(trifluoromethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
442	4-oxo-N-[3-(1-piperidylsulfonyl)phenyl]-1H-quinoline-3-carboxamide
443	4-oxo-N-quinoxalin-6-yl-1H-quinoline-3-carboxamide
444	5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2- <i>tert</i> -butyl-benzoic acid methyl ester
445	N-(2-isopropenylphenyl)-4-oxo-1H-quinoline-3-carboxamide
446	N-(1,1-dioxobenzothiophen-6-yl)-4-oxo-1H-quinoline-3-carboxamide
447	N-(3-cyanophenyl)-4-oxo-1H-quinoline-3-carboxamide
448	4-oxo-N-(4- <i>tert</i> -butylphenyl)-1H-quinoline-3-carboxamide
449	N-(<i>m</i> -tolyl)-4-oxo-1H-quinoline-3-carboxamide

450	N-[4-(1-hydroxyethyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
451	N-(4-cyano-2-ethyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
452	4-oxo-N-(4-vinylphenyl)-1H-quinoline-3-carboxamide
453	N-(3-amino-4-chloro-phenyl)-4-oxo-1H-quinoline-3-carboxamide
454	N-(2-methyl-5-phenyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
455	N-[4-(1-adamantyl)phenyl]-4-oxo-1H-quinoline-3-carboxamide
456	4-oxo-N-[3-(trifluoromethylsulfanyl)phenyl]-1H-quinoline-3-carboxamide
457	N-(4-morpholinophenyl)-4-oxo-1H-quinoline-3-carboxamide
458	N-[3-(2-hydroxyethoxy)-4- <i>tert</i> -butyl-phenyl]-4-oxo-1H-quinoline-3-carboxamide
459	N-(<i>o</i> -tolyl)-4-oxo-1H-quinoline-3-carboxamide
460	[2-methyl-2-[4-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]-propyl]aminoformic acid butyl ester
461	4-oxo-N-(2-phenylphenyl)-1H-quinoline-3-carboxamide
462	N-(3-dimethylamino-4-propyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
463	N-(4-ethylphenyl)-4-oxo-1H-quinoline-3-carboxamide
464	5-hydroxy-N-(5-hydroxy-2,4-ditert-butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
465	[5-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-2- <i>tert</i> -butyl-phenylmethyl]aminoformic acid tert-butyl ester
466	N-(2,6-diisopropylphenyl)-4-oxo-1H-quinoline-3-carboxamide
467	N-(2,3-dihydrobenzofuran-5-yl)-4-oxo-1H-quinoline-3-carboxamide
468	1-methyl-4-oxo-N-phenyl-1H-quinoline-3-carboxamide
469	4-oxo-N-(2-phenylphenyl)-7-(trifluoromethyl)-1H-quinoline-3-carboxamide
470	4-oxo-N-(4-phenylsulfanylphenyl)-1H-quinoline-3-carboxamide
471	[3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]-4-propyl-phenyl]aminoformic acid methyl ester
472	[4-ethyl-3-[(4-oxo-1H-quinolin-3-yl)carbonylamino]phenyl]aminoformic acid ethyl ester
473	1-isopropyl-4-oxo-N-(2- <i>tert</i> -butylphenyl)-1H-quinoline-3-carboxamide
474	N-(3-methyl-2-oxo-3H-benzooxazol-5-yl)-4-oxo-1H-quinoline-3-carboxamide

475	N-(2,5-dichloro-3-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
476	N-(2-cyano-5-hydroxy-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
477	N-(5-fluoro-2-pyridyl)-4-oxo-1H-quinoline-3-carboxamide
478	4-oxo-N-(3- <i>tert</i> -butyl-1H-indol-6-yl)-1H-quinoline-3-carboxamide
479	N-(1H-indol-6-yl)-5-methoxy-4-oxo-1H-quinoline-3-carboxamide
480	1-ethyl-6-methoxy-4-oxo-N-phenyl-1H-quinoline-3-carboxamide
481	N-(2-naphthyl)-4-oxo-1H-quinoline-3-carboxamide
482	[7-[(4-oxo-1H-quinolin-3-yl)carbonylamino]tetralin-1-yl]aminoformic acid ethyl ester
483	N-[2-fluoro-5-hydroxy-4-(1-methylcycloheptyl)-phenyl]-4-hydroxy-quinoline-3-carboxamide
484	N-(3-methylamino-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide
485	N-(3-dimethylamino-4- <i>tert</i> -butyl-phenyl)-4-oxo-1H-quinoline-3-carboxamide

[0178] In another embodiment, the present invention provides compounds useful as intermediates in the synthesis of compounds of formula I. In one embodiment, such compounds have formula **A-I**:



A-I;

or a salt thereof;

wherein:

G_1 is hydrogen, R' , $C(O)R'$, $C(S)R'$, $S(O)R'$, $S(O)_2R'$, $Si(CH_3)_2R'$, $P(O)(OR')_3$, $P(S)(OR')_3$, or $B(OR')_2$;

G_2 is halo, CN, CF_3 , isopropyl, or phenyl wherein said isopropyl or phenyl is optionally substituted with up to 3 substituents independently selected from WR^W , wherein W and R^W are as defined above for formula I and embodiments thereof;

G₃ is an isopropyl or a C3-C10 cycloaliphatic ring, wherein said G₃ is optionally substituted with up to 3 substituents independently selected from WR^W, wherein W and R^W are as defined above for formula I and embodiments thereof; provided that when G₁ is methoxy, G₃ is tert-butyl, then G₂ is not 2-amino-4-methoxy-5-*tert*-butyl-phenyl.

[0179] In one embodiment, the present invention provides compounds of formula A-I, provided that when G₂ and G₃ each is t-butyl, then G₁ is not hydrogen.

[0180] In another embodiment:

G₁ is hydrogen;

G₂ is halo or isopropyl, wherein said isopropyl is optionally substituted with up to 3 substituents independently selected from R'; and

G₃ is an isopropyl or a C3-C10 cycloaliphatic ring, wherein said G₃ is optionally substituted with up to 3 substituents independently selected from R'.

[0181] In another embodiment:

G₁ is hydrogen;

G₂ is halo, preferably fluoro; and

G₃ is a C3-C10 cycloaliphatic ring, wherein said G₃ is optionally substituted with up to 3 substituents independently selected from methyl, ethyl, propyl, or butyl.

[0182] In another embodiment:

G₁ is hydrogen;

G₂ is CN, halo, or CF₃; and

G₃ is an isopropyl or a C3-C10 cycloaliphatic ring, wherein said G₃ is optionally substituted with up to 3 substituents independently selected from R'.

[0183] In another embodiment:

G₁ is hydrogen;

G₂ is phenyl is optionally substituted with up to 3 substituents independently selected from -OC1-C4 alkyl, CF₃, halo, or CN; and

G₃ is an isopropyl or a C3-C10 cycloaliphatic ring, wherein said G₃ is optionally substituted with up to 3 substituents independently selected from R'.

[0184] Exemplary G_3 include optionally substituted cyclopentyl, cyclohexyl, cycloheptyl, or adamantyl. Or, G_3 is C3-C8 branched aliphatic chain. Exemplary G_3 include isopropyl, t-butyl, 3,3-diethyl-prop-3-yl, or 3,3-diethyl-2,2-dimethyl-prop-3-yl.


[0185] In another embodiment:

G_1 is hydrogen;

G_2 is t-butyl; and

G_3 is a t-butyl.

[0186] In another embodiment, the present invention provides a compound of formula **A-II**:



A-II;

or a salt thereof, wherein:

G_4 is hydrogen, halo, CN, CF_3 , CHF_2 , CH_2F , optionally substituted C1-C6 aliphatic, aralkyl, or a phenyl ring optionally substituted with up to 4 WR^W substituents;

G_5 is hydrogen or an optionally substituted C1-C6 aliphatic;

provided that both, G_4 and G_5 , are not simultaneously hydrogen;

wherein said indole ring system is further optionally substituted with up to 3 substituents independently selected from WR^W .

[0187] In one embodiment, G_4 is hydrogen. Or, G_5 is hydrogen.

[0188] In another embodiment, G_4 is hydrogen, and G_5 is C1-C6 aliphatic, wherein said aliphatic is optionally substituted with C1-C6 alkyl, halo, cyano, or CF_3 , and wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with $-CO-$, $-CONR'-$, $-CO_2-$, $-OCO-$, $-NR'CO_2-$, $-O-$, $-NR'CONR'-$, $-OCONR'-$, $-NR'CO-$, $-S-$, $-NR'-$, $-SO_2NR'-$, $NR'SO_2-$, or $-NR'SO_2NR'-$. In another embodiment, R' above is C1-C4 alkyl.

[0189] In another embodiment, G_4 is hydrogen, and G_5 is cyano, methyl, ethyl, propyl, isopropyl, butyl, sec-butyl, t-butyl, cyanomethyl, methoxyethyl, $CH_2C(O)OMe$, $(CH_2)_2-NHC(O)O-tert-But$, or cyclopentyl.

[0190] In another embodiment, G_5 is hydrogen, and G_4 is halo, C1-C6 aliphatic or phenyl, wherein said aliphatic or phenyl is optionally substituted with C1-C6 alkyl, halo,

cyano, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.

[0191] In another embodiment, G₅ is hydrogen, and G₄ is halo, ethoxycarbonyl, t-butyl, 2-methoxyphenyl, 2-ethoxyphenyl, 4-C(O)NH(CH₂)₂-NMe₂, 2-methoxy-4-chloro-phenyl, pyridine-3-yl, 4-isopropylphenyl, 2,6-dimethoxyphenyl, sec-butylaminocarbonyl, ethyl, t-butyl, or piperidin-1-ylcarbonyl.

[0192] In a related embodiment of formula A-II, the nitrogen ring atom of said indole ring is substituted with C1-C6 aliphatic, C(O)(C1-C6 aliphatic), or benzyl, wherein said aliphatic or benzyl is optionally substituted with C1-C6 alkyl, halo, cyano, or CF₃, wherein up to two methylene units of said C1-C6 aliphatic or C1-C6 alkyl is optionally replaced with -CO-, -CONR'-, -CO₂-, -OCO-, -NR'CO₂-, -O-, -NR'CONR'-, -OCONR'-, -NR'CO-, -S-, -NR'-, -SO₂NR'-, NR'SO₂-, or -NR'SO₂NR'-. In another embodiment, R' above is C1-C4 alkyl.


[0193] In another embodiment the nitrogen ring atom of said indole ring is substituted with acyl, benzyl, C(O)CH₂N(Me)C(O)CH₂NHMe, or ethoxycarbonyl.

[0194] 4. General Synthetic Schemes


[0195] Compounds of the present invention are readily prepared by methods known in the art. Illustrated below are exemplary methods for the preparation of compounds of the present invention.

[0196] The scheme below illustrates the synthesis of acid precursors of the compounds of the present invention.

[0197] Synthesis of Acid Precursors P-IV-A, P-IV-B or P-IV-C:




[0198] Synthesis of Acid Precursors P-IV-A, P-IV-B or P-IV-C:




a) AcONH_4 ; b) $\text{EtOCHC}(\text{CO}_2\text{Et})_2$, 130°C ; c) Ph_2O , ΔT ; d) I_2 , EtOH ; e) NaOH .

[0199] Synthesis of Acid Precursors **P-IV-A**, **P-IV-B** or **P-IV-C**




[0200] POCl_3 ; b) $\text{R}'\text{ONa}$; c) $n\text{-BuLi}$, ClCO_2Et ; d) NaOH

[0201] Synthesis of Amine Precursor **P-III-A**:




$(\text{CH}_3)_2\text{SO}_4$; b) $\text{K}_3\text{Fe}(\text{CN})_6$, NaOH , H_2O ; c) HNO_3 , H_2SO_4 ; d) RCOCH_3 , MeOH , NH_3 ; e) H_2 , Raney Ni

[0202] Synthesis of Amine Precursor **P-IV-A**:




[0203] HNO_3 , HOAc; b) $\text{Na}_2\text{S}_2\text{O}_4$, THF/ H_2O ; c) H_2 , Pd/C.

[0204] Synthesis of Amine Precursor **P-V-A-1**:




[0205] KNO_3 , H_2SO_4 ; b) NaNO_2 , H_2SO_4 - H_2O ; c) $\text{NH}_4\text{CO}_2\text{H}$, Pd-C; d) $\text{R}'\text{X}$; e) $\text{NH}_4\text{CO}_2\text{H}$, Pd-C

[0206] Synthesis of Amine Precursor **P-V-A-1**:




a) SO_2Cl_2 , $\text{R}_2 = \text{Cl}$; b) R_2OH , $\text{R}_2 = \text{alkyl}$; c) NBS, $\text{R}_1 = \text{Br}$; d) ClCO_2R , TEA; e) HNO_3 , H_2SO_4 ; f) base; g) $\text{ArB}(\text{OH})_2$, $\text{R}_1 = \text{Br}$; h) $[\text{H}]$; i) $\text{R}'\text{X}$, $\text{R}_1 = \text{Br}$; j) $\text{ClCF}_2\text{CO}_2\text{Me}$; k) $[\text{H}]$; l) $[\text{H}]$.

[0207] Synthesis of Amine Precursor **P-V-A-1**:




[0208] KNO_3 ; b) $[\text{H}]$; c) KNO_3 ; d) AcCl ; e) $[\text{H}]$; f) i) NaNO_2 ; ii) H_2O ; g) HCl

[0209] Synthesis of Amine Precursor **P-V-A-1**:




[0210] HNO₃, H₂SO₄; b) [H]; c) protection; d) R'CHO; e) deprotection; f) [H]; g) Na₂S, S, H₂O; h) nitration; i) (BOC)₂O; j) [H]; k) RX; l) [H]; PG= protecting group

[0211] Synthesis of Amine Precursors **P-V-A-1** or **P-V-A-2**:




[0212] a) Br_2 ; b) $\text{Zn}(\text{CN})_2$, $\text{Pd}(\text{PPh})_3$; c) $[\text{H}]$; d) BH_3 ; e) $(\text{BOC})_2\text{O}$; f) $[\text{H}]$; g) H_2SO_4 , H_2O ; h) $\text{R}'\text{X}$; i) $[\text{H}]$; j) LiAlH_4


[0213] Synthesis of Amine Precursors **P-V-A-1** or **P-V-A-2**:



(i) NaNO_2 , HCl ; ii) Na_2SO_3 , CuSO_4 , HCl ; b) NH_4Cl ; c) $[\text{H}]$


[0214] Synthesis of Amine Precursors P-V-A-1:

a) CHCl_2OMe ; b) $\text{KNO}_3, \text{H}_2\text{SO}_4$; c) Deoxo-Fluor; d) Fe

[0215] Synthesis of Amine Precursors P-V-A-3:


Ar = Aryl or Heteroaryl

a) Nitration; b) $\text{ArB}(\text{OH})_2, \text{Pd}$; c) BH_3 ; d) $(\text{BOC})_2\text{O}$

[0216] Synthesis of Amine Precursors P-V-B-1:

a) AcCl ; b) DEAD ; c) AlCl_3 ; d) NaOH


[0217] Synthesis of Amine Precursors P-V-B-1:




a) ClCH_2COCl ; b) $[\text{H}]$; c) protection; d) $[\text{H}]$

PG= protecting group

[0218] Synthesis of Amine Precursors P-V-B-1:




a) $\text{HSCH}_2\text{CO}_2\text{H}$; b) $[\text{H}]$


[0219] Synthesis of Amine Precursors **P-V-B-2**:

a) AlCl_3 ; b) $[\text{H}]$; c) i) $\text{R}_1\text{R}_2\text{CHCOCH}_2\text{CH}_2\text{Cl}$; ii) NaBH_4 ; d) NH_2OH ; e) DIBAL-H; f) nitration;
 g) protection; h) $[\text{H}]$

PG= protecting group


[0220] Synthesis of Amine Precursors **P-V-B-3**:

[0223] Synthesis of Amine Precursors V-B-5:




a) NaNO₂, HCl, SnCl₂·2H₂O, H₂O; b) RCH₂COR, AcOH, EtOH; c) H₃PO₄, toluene; d) H₂, Pd-C, EtOH

[0224] Synthesis of Amine Precursors P-V-B-5:




a) NaNO₂, HCl, SnCl₂·2H₂O, H₂O; b) RCH₂COH, AcOH, EtOH; c) H₃PO₄, toluene; d) H₂, Pd-C, EtOH

[0225] Synthesis of Amine Precursors P-V-B-5:




a) RX (X=Br, I), zinc triflate, TBAI, DIEA, toluene; b) H₂, Raney Ni, EtOH or H₂, Pd-C, EtOH or SnCl₂·2H₂O, EtOH; c) ClSO₂NCO, DMF, CH₃CN; d) Me₂NH, H₂CO, AcOH; e) MeI, DMF, THF, H₂O; f) MNu (M= Na, K, Li; Nu= nucleophile)


[0226] Synthesis of Amine Precursors P-V-B-5:



a) HNO₃, H₂SO₄; b) Me₂NCH(OMe)₂, DMF; c) H₂, Raney Ni, EtOH

[0227] Synthesis of Amine Precursors **P-V-B-5**:




a) When PG= SO₂Ph: PhSO₂Cl, Et₃N, DMAP, CH₂Cl₂; When PG= Ac: AcCl, NaHCO₃, CH₂Cl₂;
 b) When R= RCO: (RCO)₂O, AlCl₃, CH₂Cl₂; When R=Br: Br₂, AcOH; c) HBr or HCl; d) KNO₃, H₂SO₄; e) MnO₂, CH₂Cl₂ or DDQ, 1,4-dioxane; f) H₂, Raney Ni, EtOH.


[0228] Synthesis of Amine Precursors **P-V-B-5**:

a) NBS, DMF; b) KNO₃, H₂SO₄; c) HC≡CSiMe₃, Pd(PPh₃)₂Cl₂, CuI, Et₃N, Toluene, H₂O; d) CuI, DMF; e) H₂, Raney Ni, MeOH

[0229] Synthesis of Amine Precursors P-V-A-3 and P-V-A-6:


Ar= Aryl or heteroaryl

a) ArB(OH)₂, Pd(PPh₃)₄, K₂CO₃, H₂O, THF or ArB(OH)₂, Pd₂(dba)₃, P(tBu)₃, KF, THF**[0230] Synthesis of Amine Precursors P-V-A-4:**R= CN, CO₂Et; a) MeI, NaOtBu, DMF; b) HCO₂K, Pd-C, EtOH or HCO₂NH₄, Pd-C, EtOH**[0231] Synthesis of Amine Precursors P-V-A-4:**a) ArBr, Pd(OAc)₂, PS-PPh₃, K₂CO₃, DMF**[0232] Synthesis of Amine Precursors P-V-B-4:**




a) H_2 , Pd-C, MeOH

[0233] Synthesis of Amine Precursors P-V-B-4:




a) NaBH_4 , MeOH; b) H_2 , Pd-C, MeOH; c) NH_2OH , Pyridine; d) H_2 , Pd-C, MeOH; e) Boc_2O , Et_3N , MeOH

[0234] Synthesis of Compounds of Formula I:




a) $\text{Ar}_1\text{R}_7\text{NH}$, coupling reagent, base, solvent. Examples of conditions used: HATU, DIEA; BOP, DIEA, DMF; HBTU, Et_3N , CH_2Cl_2 ; PFPTFA, pyridine.

[0235] Synthesis of Compounds of Formula I':




R⁵ = aliphatic: a) R⁵X (X= Br, I), Cs₂CO₃, DMF

[0236] Synthesis of Compounds of formula V-B-5:




a) NaOH, THF; b) HNR₂, HATU, DIEA, DMF

[0237] Synthesis of Compounds of formula V-B-5:




WR^w = aryl or heteroaryl: a) ArB(OH)₂, (dppf)PdCl₂, K₂CO₃, DMF

[0238] Synthesis of Compounds of Formula V-A-2 & V-A-5:




a) SnCl₂·2H₂O, EtOH; b) PG= BOC: TFA, CH₂Cl₂; c) CH₂O, NaBH₃CN, CH₂Cl₂, MeOH; d) RXCl, DIEA, THF or RXCl, NMM, 1,4-dioxane or RXCl, CH₂Cl₂, DMF; e) R'R''NH, LiClO₄, CH₂Cl₂, iPrOH

[0239] Synthesis of compounds of formula V-B-2:




a) When PG = BOC: TFA, CH₂Cl₂; When PG = Ac: NaOH or HCl, EtOH or THF

[0240] Synthesis of compounds of formula V-A-2:




a) When PG = BOC: TFA, CH₂Cl₂



a) When PG = BOC: TFA, CH₂Cl₂; b) ROCOCl, Et₃N, DMF

[0241] Synthesis of compounds of formula V-A-4:



a) When PG = BOC: TFA, CH₂Cl₂; b) When R^w = CO₂R: ROCOCl, DIEA, MeOH

[0242] In the schemes above, the radical R employed therein is a substituent, e.g., R^w as defined hereinabove. One of skill in the art will readily appreciate that synthetic routes suitable for various substituents of the present invention are such that the reaction conditions and steps employed do not modify the intended substituents.

[0243] 5. Uses, Formulation and Administration

[0244] *Pharmaceutically acceptable compositions*

[0245] As discussed above, the present invention provides compounds that are useful as modulators of ABC transporters and thus are useful in the treatment of disease, disorders or conditions such as cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome,

53568-20

Perlizaues-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebular ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease.

[0246] Accordingly, in another aspect of the present invention, pharmaceutically acceptable compositions are provided, wherein these compositions comprise any of the compounds as described herein, and optionally comprise a pharmaceutically acceptable carrier, adjuvant or vehicle. In certain embodiments, these compositions optionally further comprise one or more additional therapeutic agents.

[0247] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative or a prodrug thereof. According to the present invention, a pharmaceutically acceptable derivative or a prodrug includes, but is not limited to, pharmaceutically acceptable salts, esters, salts of such esters, or any other adduct or derivative which upon administration to a patient in need thereof is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof.

[0248] As used herein, the term "pharmaceutically acceptable salt" refers to those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. A "pharmaceutically acceptable salt" means any non-toxic salt or salt of an ester of a compound of this invention that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound of this invention or an inhibitorily active metabolite or residue thereof.

[0249] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge, *et al.* describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of

an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0250] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. Remington's Pharmaceutical Sciences, Sixteenth Edition, E. W. Martin (Mack Publishing Co., Easton, Pa., 1980) discloses various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable

composition, its use is contemplated to be within the scope of this invention. Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such a propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[0251] *Uses of Compounds and Pharmaceutically Acceptable Compositions*

[0252] In yet another aspect, the present invention provides a method of treating a condition, disease, or disorder implicated by ABC transporter activity, e.g., CFTR. In certain embodiments, the present invention provides a method of treating a condition, disease, or disorder implicated by a deficiency of the ABC transporter activity, the method comprising administering a composition comprising a compound of formula (I) to a subject, preferably a mammal, in need thereof.

[0253] In certain embodiments, the present invention provides a method of treating cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia,

abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease, comprising the step of administering to said mammal an effective amount of a composition comprising a compound of the present invention.

[0254] According to an alternative preferred embodiment, the present invention provides a method of treating cystic fibrosis comprising the step of administering to said mammal a composition comprising the step of administering to said mammal an effective amount of a composition comprising a compound of the present invention.

[0255] According to the invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of one or more of cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease,

Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease.

[0256] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route of administration effective for treating or lessening the severity of one or more of cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker syndrome, COPD, dry-eye disease, or Sjogren's disease.

[0257] In one embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in a patient.

[0258] In certain embodiments, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who exhibit residual CFTR activity in the apical membrane of respiratory and non-respiratory epithelia. The presence of residual CFTR activity at the epithelial surface can be readily detected using methods known in the art, e.g., standard electrophysiological, biochemical, or

histochemical techniques. Such methods identify CFTR activity using *in vivo* or *ex vivo* electrophysiological techniques, measurement of sweat or salivary Cl⁻ concentrations, or *ex vivo* biochemical or histochemical techniques to monitor cell surface density. Using such methods, residual CFTR activity can be readily detected in patients heterozygous or homozygous for a variety of different mutations, including patients homozygous or heterozygous for the most common mutation, $\Delta F508$.

[0259] In another embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients who have residual CFTR activity induced or augmented using pharmacological methods or gene therapy. Such methods increase the amount of CFTR present at the cell surface, thereby inducing a hitherto absent CFTR activity in a patient or augmenting the existing level of residual CFTR activity in a patient.

[0260] In one embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients within certain genotypes exhibiting residual CFTR activity, e.g., class III mutations (impaired regulation or gating), class IV mutations (altered conductance), or class V mutations (reduced synthesis) (Lee R. Choo-Kang, Pamela L., Zeitlin, *Type I, II, III, IV, and V cystic fibrosis Transmembrane Conductance Regulator Defects and Opportunities of Therapy*; Current Opinion in Pulmonary Medicine 6:521 – 529, 2000). Other patient genotypes that exhibit residual CFTR activity include patients homozygous for one of these classes or heterozygous with any other class of mutations, including class I mutations, class II mutations, or a mutation that lacks classification.

[0261] In one embodiment, the compounds and compositions of the present invention are useful for treating or lessening the severity of cystic fibrosis in patients within certain clinical phenotypes, e.g., a moderate to mild clinical phenotype that typically correlates with the amount of residual CFTR activity in the apical membrane of epithelia. Such phenotypes include patients exhibiting pancreatic sufficiency or patients diagnosed with idiopathic pancreatitis and congenital bilateral absence of the vas deferens, or mild lung disease.

[0262] The exact amount required will vary from subject to subject, depending on the species, age, and general condition of the subject, the severity of the infection, the particular agent, its mode of administration, and the like. The compounds of the invention are preferably

formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts. The term "patient", as used herein, means an animal, preferably a mammal, and most preferably a human.

[0263] The pharmaceutically acceptable compositions of this invention can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, as an oral or nasal spray, or the like, depending on the severity of the infection being treated. In certain embodiments, the compounds of the invention may be administered orally or parenterally at dosage levels of about 0.01 mg/kg to about 50 mg/kg and preferably from about 1 mg/kg to about 25 mg/kg, of subject body weight per day, one or more times a day, to obtain the desired therapeutic effect.

[0264] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0265] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0266] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0267] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered compound form is accomplished by dissolving or suspending the compound in an oil vehicle. Injectable depot forms are made by forming microcapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0268] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are

solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[0269] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar--agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0270] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0271] The active compounds can also be in microencapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as

sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such a magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[0272] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms are prepared by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[0273] As described generally above, the compounds of the invention are useful as modulators of ABC transporters. Thus, without wishing to be bound by any particular theory, the compounds and compositions are particularly useful for treating or lessening the severity of a disease, condition, or disorder where hyperactivity or inactivity of ABC transporters is implicated in the disease, condition, or disorder. When hyperactivity or inactivity of an ABC transporter is implicated in a particular disease, condition, or disorder, the disease, condition, or disorder may also be referred to as a "ABC transporter-mediated disease, condition or disorder". Accordingly, in another aspect, the present invention provides a method for treating or lessening the severity of a disease, condition, or disorder where hyperactivity or inactivity of an ABC transporter is implicated in the disease state.

[0274] The activity of a compound utilized in this invention as a modulator of an ABC transporter may be assayed according to methods described generally in the art and in the Examples herein.

[0275] It will also be appreciated that the compounds and pharmaceutically acceptable compositions of the present invention can be employed in combination therapies, that is, the compounds and pharmaceutically acceptable compositions can be administered concurrently with, prior to, or subsequent to, one or more other desired therapeutics or medical procedures. The particular combination of therapies (therapeutics or procedures) to employ in a combination regimen will take into account compatibility of the desired therapeutics and/or procedures and the desired therapeutic effect to be achieved. It will also be appreciated that the therapies employed may achieve a desired effect for the same disorder (for example, an inventive compound may be administered concurrently with another agent used to treat the same disorder), or they may achieve different effects (e.g., control of any adverse effects). As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated".

[0276] In one embodiment, the additional agent is selected from a mucolytic agent, bronchodilator, an anti-biotic, an anti-infective agent, an anti-inflammatory agent, a CFTR modulator other than a compound of the present invention, or a nutritional agent.

[0277] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

[0278] The compounds of this invention or pharmaceutically acceptable compositions thereof may also be incorporated into compositions for coating an implantable medical device, such as prostheses, artificial valves, vascular grafts, stents and catheters. Accordingly, the present invention, in another aspect, includes a composition for coating an implantable device comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. In still

another aspect, the present invention includes an implantable device coated with a composition comprising a compound of the present invention as described generally above, and in classes and subclasses herein, and a carrier suitable for coating said implantable device. Suitable coatings and the general preparation of coated implantable devices are described in US Patents 6,099,562; 5,886,026; and 5,304,121. The coatings are typically biocompatible polymeric materials such as a hydrogel polymer, polymethyldisiloxane, polycaprolactone, polyethylene glycol, polylactic acid, ethylene vinyl acetate, and mixtures thereof. The coatings may optionally be further covered by a suitable topcoat of fluorosilicone, polysaccharides, polyethylene glycol, phospholipids or combinations thereof to impart controlled release characteristics in the composition.

[0279] Another aspect of the invention relates to modulating ABC transporter activity in a biological sample or a patient (e.g., *in vitro* or *in vivo*), which method comprises administering to the patient, or contacting said biological sample with a compound of formula I or a composition comprising said compound. The term "biological sample", as used herein, includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof.

[0280] Modulation of ABC transporter activity, e.g., CFTR, in a biological sample is useful for a variety of purposes that are known to one of skill in the art. Examples of such purposes include, but are not limited to, the study of ABC transporters in biological and pathological phenomena; and the comparative evaluation of new modulators of ABC transporters.

[0281] In yet another embodiment, a method of modulating activity of an anion channel *in vitro* or *in vivo*, is provided comprising the step of contacting said channel with a compound of formula (I). In preferred embodiments, the anion channel is a chloride channel or a bicarbonate channel. In other preferred embodiments, the anion channel is a chloride channel.

[0282] According to an alternative embodiment, the present invention provides a method of increasing the number of functional ABC transporters in a membrane of a cell, comprising the step of contacting said cell with a compound of formula (I). The term "functional ABC transporter" as used herein means an ABC transporter that is capable of transport activity. In preferred embodiments, said functional ABC transporter is CFTR.

[0283] According to another preferred embodiment, the activity of the ABC transporter is measured by measuring the transmembrane voltage potential. Means for measuring the voltage potential across a membrane in the biological sample may employ any of the known methods in the art, such as optical membrane potential assay or other electrophysiological methods.

[0284] The optical membrane potential assay utilizes voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J 69(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol 4(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today 4(9): 431-439).

[0285] These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, DiSBAC₂(3), and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged DiSBAC₂(3) to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission can be monitored using VIPRTM II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

[0286] In another aspect the present invention provides a kit for use in measuring the activity of a ABC transporter or a fragment thereof in a biological sample *in vitro* or *in vivo* comprising (i) a composition comprising a compound of formula (I) or any of the above embodiments; and (ii) instructions for a) contacting the composition with the biological sample and b) measuring activity of said ABC transporter or a fragment thereof. In one embodiment, the kit further comprises instructions for a) contacting an additional composition with the biological sample; b) measuring the activity of said ABC transporter or a fragment thereof in the presence of said additional compound, and c) comparing the activity of the ABC transporter in the presence of the additional compound with the density of the ABC transporter


in the presence of a composition of formula (I). In preferred embodiments, the kit is used to measure the density of CFTR.

[00287] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLES


[00286] Example 1:

[00287] General scheme to prepare Acid Moities:



a) 140-150 °C; b) PPA, POCl₃, 70 °C or diphenyl ether, 220 °C; c) i) 2N NaOH ii) 2N HCl

[00288] Specific example: 2-Phenylaminomethylene-malonic acid diethyl ester



A mixture of aniline (25.6 g, 0.28 mol) and diethyl 2-(ethoxymethylene)malonate (62.4 g, 0.29 mol) was heated at 140-150 °C for 2 h. The mixture was cooled to room temperature and dried under reduced pressure to afford 2-phenylaminomethylene-malonic acid diethyl ester as a solid, which was used in the next step without further purification. ¹H NMR (*d*-DMSO) δ 11.00 (d, 1H), 8.54 (d, *J* = 13.6 Hz, 1H), 7.36-7.39 (m, 2H), 7.13-7.17 (m, 3 H), 4.17-4.33 (m, 4H), 1.18-1.40 (m, 6H).

[00289] 4-Hydroxyquinoline-3-carboxylic acid ethyl ester


A 1 L three-necked flask fitted with a mechanical stirrer was charged with 2-phenylaminomethylene-malonic acid diethyl ester (26.3 g, 0.1 mol), polyphosphoric acid (270 g) and phosphoryl chloride (750 g). The mixture was heated to about 70 °C and stirred for 4 h. The

mixture was cooled to room temperature, and filtered. The residue was treated with aqueous Na_2CO_3 solution, filtered, washed with water and dried. 4-Hydroxyquinoline-3-carboxylic acid ethyl ester was obtained as a pale brown solid (15.2 g, 70 %). The crude product was used in next step without further purification.

[00290] A-1; 4-Oxo-1,4-dihydroquinoline-3-carboxylic acid


4-Hydroxyquinoline-3-carboxylic acid ethyl ester (15 g, 69 mmol) was suspended in sodium hydroxide solution (2N, 150 mL) and stirred for 2 h under reflux. After cooling, the mixture was filtered, and the filtrate was acidified to pH 4 with 2N HCl. The resulting precipitate was collected via filtration, washed with water and dried under vacuum to give 4-oxo-1,4-dihydroquinoline-3-carboxylic acid (**A-1**) as a pale white solid (10.5 g, 92 %). ^1H NMR (d -DMSO) δ 15.34 (s, 1 H), 13.42 (s, 1 H), 8.89 (s, 1 H), 8.28 (d, $J = 8.0$ Hz, 1H), 7.88 (m, 1H), 7.81 (d, $J = 8.4$ Hz, 1H), 7.60 (m, 1H).

[00291] Specific Example: A-2; 6-Fluoro-4-hydroxy-quinoline-3-carboxylic acid



6-Fluoro-4-hydroxy-quinoline-3-carboxylic acid (**A-2**) was synthesized following the general scheme above starting from 4-fluoro-phenylamine. Overall yield (53 %). ^1H NMR ($\text{DMSO}-d_6$) δ 15.2 (br s, 1 H), 8.89 (s, 1 H), 7.93-7.85 (m, 2 H), 7.80-7.74 (m, 1 H); ESI-MS 207.9 m/z (MH^+).

[00292] Example 2:



[00293] 2-Bromo-5-methoxy-phenylamine

A mixture of 1-bromo-4-methoxy-2-nitro-benzene (10 g, 43 mmol) and Raney Ni (5 g) in ethanol (100 mL) was stirred under H₂ (1 atm) for 4 h at room temperature. Raney Ni was filtered off and the filtrate was concentrated under reduced pressure. The resulting solid was purified by column chromatography to give 2-bromo-5-methoxy-phenylamine (7.5 g, 86 %).

[00294] 2-[(2-Bromo-5-methoxy-phenylamino)-methylene]-malonic acid diethyl ester

A mixture of 2-bromo-5-methoxy-phenylamine (540 mg, 2.64 mmol) and diethyl 2-(ethoxymethylene)malonate (600 mg, 2.7 mmol) was stirred at 100 °C for 2 h. After cooling, the reaction mixture was recrystallized from methanol (10 mL) to give 2-[(2-bromo-5-methoxy-phenylamino)-methylene]-malonic acid diethyl ester as a yellow solid (0.8 g, 81 %).

[00295] 8-Bromo-5-methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester


2-[(2-Bromo-5-methoxy-phenylamino)-methylene]-malonic acid diethyl ester (9 g, 24.2 mmol) was slowly added to polyphosphoric acid (30 g) at 120 °C. The mixture was stirred at this temperature for additional 30 min and then cooled to room temperature. Absolute ethanol (30 mL) was added and the resulting mixture was refluxed for 30 min. The mixture was basified with aqueous sodium bicarbonate at 25 °C and extracted with EtOAc (4 x 100 mL). The organic layers were combined, dried and the solvent evaporated to give 8-bromo-5-methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (2.3 g, 30 %).

[00296] 5-Methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester

A mixture of 8-bromo-5-methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (2.3 g, 7.1 mmol), sodium acetate (580 mg, 7.1 mmol) and 10 % Pd/C (100 mg) in glacial acetic acid (50 ml) was stirred under H₂ (2.5 atm) overnight. The catalyst was removed via filtration, and the reaction mixture was concentrated under reduced pressure. The resulting oil was dissolved in CH₂Cl₂ (100 mL) and washed with aqueous sodium bicarbonate solution and water. The organic layer was dried, filtered and concentrated. The crude product was purified by column chromatography to afford 5-methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester as a yellow solid (1 g, 57 %).

[00297] A-4; 5-Methoxy-4-oxo-1, 4-dihydro-quinoline-3-carboxylic acid

A mixture of 5-methoxy-4-oxo-1, 4-dihydro-quinoline-3-carboxylic acid ethyl ester (1 g, 7.1 mmol) in 10% NaOH solution (50 mL) was heated to reflux overnight and then cooled to room temperature. The mixture was extracted with ether. The aqueous phase was separated and acidified with conc. HCl solution to pH 1-2. The resulting precipitate was collected by filtration to give 5-methoxy-4-oxo-1, 4-dihydro-quinoline-3-carboxylic acid (**A-4**) (530 mg, 52 %). ¹H NMR (DMSO) δ : 15.9 (s, 1 H), 13.2 (br, 1 H), 8.71 (s, 1 H), 7.71 (t, J = 8.1 Hz, 1 H), 7.18 (d, J = 8.4 Hz, 1 H), 6.82 (d, J = 8.4 Hz, 1 H), 3.86 (s, 3 H); ESI-MS 219.9 m/z (MH⁺).

[00298] Example 3:**[0288] Sodium 2-(mercapto-phenylamino-methylene)-malonic acid diethyl ester**

To a suspension of NaH (60% in mineral oil, 6 g, 0.15 mol) in Et₂O at room temperature was added dropwise, over a 30 minutes period, ethyl malonate (24 g, 0.15 mol). Phenyl isothiocyanate (20.3 g, 0.15 mol) was then added dropwise with stirring over 30 min. The mixture was refluxed for 1 h and then stirred overnight at room temperature. The solid was separated, washed with anhydrous ether (200 mL), and dried under *vacuum* to yield sodium 2-(mercapto-phenylamino-methylene)-malonic acid diethyl ester as a pale yellow powder (46 g, 97 %).

[00299] 2-(Methylsulfanyl-phenylamino-methylene)-malonic acid diethyl ester


Over a 30 min period, methyl iodide (17.7 g, 125 mmol) was added dropwise to a solution of sodium 2-(mercapto-phenylamino-methylene)-malonic acid diethyl ester (33 g, 104 mmol) in DMF (100 mL) cooled in an ice bath. The mixture was stirred at room temperature for 1 h, and then poured into ice water (300 mL). The resulting solid was collected via filtration, washed with water and dried to give 2-(methylsulfanyl-phenylamino-methylene)-malonic acid diethyl ester as a pale yellow solid (27 g, 84 %).

[00300] 4-Hydroxy-2-methylsulfanyl-quinoline-3-carboxylic acid ethyl ester

A mixture of 2-(methylsulfanyl-phenylamino-methylene)-malonic acid diethyl ester (27 g, 87 mmol) in 1,2-dichlorobenzene (100 mL) was heated to reflux for 1.5 h. The solvent was removed under reduced pressure and the oily residue was triturated with hexane to afford a pale yellow solid that was purified by preparative HPLC to yield 4-hydroxy-2-methylsulfanyl-quinoline-3-carboxylic acid ethyl ester (8 g, 35 %).

[00301] A-16; 2-Methylsulfanyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid

4-Hydroxy-2-methylsulfanyl-quinoline-3-carboxylic acid ethyl ester (8 g, 30 mmol) was heated under reflux in NaOH solution (10%, 100 mL) for 1.5 h. After cooling, the mixture was acidified with concentrated HCl to pH 4. The resulting solid was collected via filtration, washed with water (100 mL) and MeOH (100 mL) to give 2-methylsulfanyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid (**A-16**) as a white solid (6 g, 85 %). ¹H NMR (CDCl₃) δ 16.4 (br s, 1 H), 11.1 (br s, 1 H), 8.19 (d, *J* = 8 Hz, 1H), 8.05 (d, *J* = 8 Hz, 1H), 7.84 (t, *J* = 8, 8 Hz, 1H), 7.52 (t, *J* = 8 Hz, 1H), 2.74 (s, 3H); ESI-MS 235.9 m/z (MH⁺).

[00302] Example 4:

a) PPh_3 , Et_3N , CCl_4 , $\text{CF}_3\text{CO}_2\text{H}$; b) diethyl malonate; c) $T \sim 200^\circ\text{C}$; d) 10% NaOH

[00303] 2,2,2-Trifluoro-N-phenyl-acetimidoyl chloride

A mixture of Ph_3P (138.0 g, 526 mmol), Et_3N (21.3 g, 211 mmol), CCl_4 (170 mL) and TFA (20 g, 175 mmol) was stirred for 10 min in an ice-bath. Aniline (19.6 g, 211 mmol) was dissolved in CCl_4 (20 mL) was added. The mixture was stirred at reflux for 3 h. The solvent was removed under vacuum and hexane was added. The precipitates (Ph_3PO and Ph_3P) were filtered off and washed with hexane. The filtrate was distilled under reduced pressure to yield 2,2,2-trifluoro-N-phenyl-acetimidoyl chloride (19 g), which was used in the next step without further purification.

[00304] 2-(2,2,2-Trifluoro-1-phenylimino-ethyl)-malonic acid diethyl ester


To a suspension of NaH (3.47 g, 145 mmol, 60 % in mineral oil) in THF (200 mL) was added diethyl malonate (18.5 g, 116 mmol) at 0°C . The mixture was stirred for 30 min at this temperature and 2,2,2-trifluoro-N-phenyl-acetimidoyl chloride (19 g, 92 mmol) was added at 0°C . The reaction mixture was allowed to warm to room temperature and stirred overnight. The mixture was diluted with CH_2Cl_2 , washed with saturated sodium bicarbonate solution and brine. The combined organic layers were dried over Na_2SO_4 , filtered and concentrated to provide 2-(2,2,2-trifluoro-1-phenylimino-ethyl)-malonic acid diethyl ester, which was used directly in the next step without further purification.

[00305] 4-Hydroxy-2-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester

2-(2,2,2-Trifluoro-1-phenylimino-ethyl)-malonic acid diethyl ester was heated at 210°C for 1 h with continuous stirring. The mixture was purified by column chromatography (petroleum ether) to yield 4-hydroxy-2-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester (12 g, 24 % over 3 steps).

[00306] A-15; 4-Hydroxy-2-trifluoromethyl-quinoline-3-carboxylic acid

A suspension of 4-hydroxy-2-trifluoromethyl-quinoline-3-carboxylic acid ethyl ester (5 g, 17.5 mmol) in 10% aqueous NaOH solution was heated at reflux for 2 h. After cooling, dichloromethane was added and the aqueous phase was separated and acidified with concentrated HCl to pH 4. The resulting precipitate was collected via filtration, washed with water and Et₂O to provide 4-hydroxy-2-trifluoromethyl-quinoline-3-carboxylic acid (**A-15**) (3.6 g, 80 %). ¹H NMR (DMSO-*d*₆) δ 8.18-8.21 (d, *J* = 7.8 Hz, 1 H), 7.92-7.94 (d, *J* = 8.4 Hz, 1 H), 7.79-7.83 (t, *J* = 14.4 Hz, 1 H), 7.50-7.53 (t, *J* = 15 Hz, 1 H); ESI-MS 257.0 m/z (MH⁺).

[00307] Example 5:

a) CH₃C(O)ONH₄, toluene; b) EtOCHC(CO₂Et)₂, 130 °C; c) Ph₂O; d) I₂, EtOH; e) NaOH

[00308] 3-Amino-cyclohex-2-enone

A mixture of cyclohexane-1,3-dione (56.1 g, 0.5 mol) and AcONH₄ (38.5 g, 0.5 mol) in toluene was heated at reflux for 5 h with a Dean-stark apparatus. The resulting oily layer was separated and concentrated under reduced pressure to give 3-amino-cyclohex-2-enone (49.9 g, 90 %), which was used directly in the next step without further purification.

[00309] 2-[(3-Oxo-cyclohex-1-enylamino)-methylene]-malonic acid diethyl

ester

A mixture of 3-amino-cyclohex-2-enone (3.3 g, 29.7 mmol) and diethyl 2-(ethoxymethylene)malonate (6.7 g, 31.2 mmol) was stirred at 130 °C for 4 h. The reaction mixture was concentrated under reduced pressure and the resulting oil was purified by column chromatography (silica gel, ethyl acetate) to give 2-[(3-oxo-cyclohex-1-enylamino)-methylene]-malonic acid diethyl ester (7.5 g, 90 %).

[00310] 4,5-Dioxo-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid ethyl ester

A mixture of 2-[(3-oxo-cyclohex-1-enylamino)-methylene]-malonic acid diethyl ester (2.8 g, 1 mmol) and diphenylether (20 mL) was refluxed for 15 min. After cooling, *n*-hexane (80 mL) was added. The resulting solid was isolated via filtration and recrystallized from methanol to give 4,5-dioxo-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid ethyl ester (1.7 g 72 %).


[00311] 5-Hydroxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester

To a solution of 4,5-dioxo-1,4,5,6,7,8-hexahydro-quinoline-3-carboxylic acid ethyl ester (1.6 g, 6.8 mmol) in ethanol (100 mL) was added iodine (4.8 g, 19 mmol). The mixture was refluxed for 19 h and then concentrated under reduced pressure. The resulting solid was washed with ethyl acetate, water and acetone, and then recrystallized from DMF to give 5-hydroxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (700 mg, 43 %).

[00312] A-3; 5-Hydroxy-4-oxo-1, 4-dihydro-quinoline-3-carboxylic acid

A mixture of 5-hydroxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid ethyl ester (700 mg, 3 mmol) in 10% NaOH (20 ml) was heated at reflux overnight. After cooling, the mixture was extracted with ether. The aqueous phase was separated and acidified with conc. HCl to pH 1-2. The resulting precipitate was collected via filtration to give 5-hydroxy-4-oxo-1, 4-dihydro-quinoline-3-carboxylic acid (A-3) (540 mg, 87 %). ¹H NMR (DMSO-*d*₆) δ 13.7 (br, 1 H), 13.5 (br, 1 H), 12.6 (s, 1 H), 8.82 (s, 1 H), 7.68 (t, *J* = 8.1 Hz, 1 H), 7.18 (d, *J* = 8.4 Hz, 1 H), 6.82 (d, *J* = 8.4 Hz, 1 H); ESI-MS 205.9 *m/z* (MH⁺).

[00313] Example 6:



a) POCl_3 ; b) MeONa ; c) $n\text{-BuLi}$, ClCO_2Et ; d) NaOH

2,4-Dichloroquinoline

A suspension of quinoline-2,4-diol (15 g, 92.6 mmol) in POCl_3 was heated at reflux for 2 h. After cooling, the solvent was removed under reduced pressure to yield 2,4-dichloroquinoline, which was used without further purification.

[00314] 2,4-Dimethoxyquinoline

To a suspension of 2,4-dichloroquinoline in MeOH (100 mL) was added sodium methoxide (50 g). The mixture was heated at reflux for 2 days. After cooling, the mixture was filtered. The filtrate was concentrated under reduced pressure to yield a residue that was dissolved in water and extracted with CH_2Cl_2 . The combined organic layers were dried over Na_2SO_4 and concentrated to give 2,4-dimethoxyquinoline as a white solid (13 g, 74 % over 2 steps).

[00315] Ethyl 2,4-dimethoxyquinoline-3-carboxylate

To a solution of 2,4-dimethoxyquinoline (11.5 g, 60.8 mmol) in anhydrous THF was added dropwise $n\text{-BuLi}$ (2.5 M in hexane, 48.6 mL, 122 mmol) at 0 °C. After stirring for 1.5 h at 0 °C, the mixture was added to a solution of ethyl chloroformate in anhydrous THF and stirred at 0 °C for additional 30 min and then at room temperature overnight. The reaction mixture was poured into water and extracted with CH_2Cl_2 . The organic layer was dried over Na_2SO_4 and concentrated under vacuum. The resulting residue was purified by column chromatography (petroleum ether / EtOAc = 50 / 1) to give ethyl 2, 4-dimethoxyquinoline-3-carboxylate (9.6 g, 60 %).

[00316] A-17; 2,4-Dimethoxyquinoline-3-carboxylic acid

Ethyl 2,4-dimethoxyquinoline-3-carboxylate (1.5 g, 5.7 mmol) was heated at reflux in NaOH solution (10 %, 100 mL) for 1 h. After cooling, the mixture was acidified with concentrated HCl to pH 4. The resulting precipitate was collected via filtration and washed with water and ether to give 2,4-dimethoxyquinoline-3-carboxylic acid (**A-17**) as a white solid (670 mg, 50 %). ¹H NMR (CDCl₃) δ 8.01-8.04 (d, *J*=12 Hz, 1 H), 7.66-7.76 (m, 2 H), 7.42-7.47 (t, *J*= 22 Hz, 2 H), 4.09 (s, 3 H). 3.97 (s, 3 H); ESI-MS 234.1 m/z (MH⁺).

[00317] Commercially available acids


Acid	Name
A-5	6,8-Difluoro-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-6	6-[(4-Fluoro-phenyl)-methyl-sulfamoyl]-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-7	6-(4-Methyl-piperidine-1-sulfonyl)-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-8	4-Oxo-6-(pyrrolidine-1-sulfonyl)-1,4-dihydro-quinoline-3-carboxylic acid
A-10	6-Ethyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-11	6-Ethoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-12	4-Oxo-7-trifluoromethyl-1,4-dihydro-quinoline-3-carboxylic acid
A-13	7-Chloro-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-14	4-Oxo-5,7-bis-trifluoromethyl-1,4-dihydro-quinoline-3-carboxylic acid
A-20	1-Methyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-21	1-Isopropyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-22	1,6-Dimethyl-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-23	1-Ethyl-6-methoxy-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid
A-24	6-Chloro-4-oxo-1,4-dihydro-quinoline-3-carboxylic acid

[00318] Amine Moieties

[00319] N-1 Substituted 6-aminoindoles


[00320] **Example 1:**

[00321] General Scheme:



a) RX (X = Cl, Br, I), K₂CO₃, DMF or CH₃CN; b) H₂, Pd-C, EtOH or SnCl₂·2H₂O, EtOH.

[00322] Specific example:



[00323] **1-Methyl-6-nitro-1H-indole**


To a solution of 6-nitroindole (4.05 g 25 mmol) in DMF (50 mL) was added K₂CO₃ (8.63 g, 62.5 mmol) and MeI (5.33 g, 37.5 mmol). After stirring at room temperature overnight, the mixture was poured into water and extracted with ethyl acetate. The combined organic layers were dried over Na₂SO₄ and concentrated under *vacuum* to give the product 1-methyl-6-nitro-1H-indole (4.3 g, 98 %).

[00324] **B-1; 1-Methyl-1H-indol-6-ylamine**

A suspension of 1-methyl-6-nitro-1H-indole (4.3 g, 24.4 mmol) and 10% Pd-C (0.43 g) in EtOH (50 mL) was stirred under H₂ (1 atm) at room temperature overnight. After filtration, the filtrate was concentrated and acidified with HCl-MeOH (4 mol/L) to give 1-methyl-1H-indol-6-ylamine hydrochloride salt (**B-1**) (1.74 g, 49 %) as a grey powder. ¹H NMR (DMSO-*d*₆): δ 9.10 (s, 2 H),


7.49 (d, $J = 8.4$ Hz, 1 H), 7.28 (d, $J = 2.0$ Hz, 1H), 7.15(s, 1 H), 6.84 (d, $J = 8.4$ Hz, 1 H), 6.38 (d, $J = 2.8$ Hz, 1H), 3.72 (s, 3 H); ESI-MS 146.08 m/z (MH^+).

[00325] Other examples:



[00326] **B-2; 1-Benzyl-1H-indol-6-ylamine**


1-Benzyl-1H-indol-6-ylamine (**B-2**) was synthesized following the general scheme above starting from 6-nitroindole and benzyl bromide. Overall yield (~ 40 %). HPLC ret. time 2.19 min, 10–99 % CH_3CN , 5 min run; ESI-MS 223.3 m/z (MH^+).



[00327] **B-3; 1-(6-Amino-indol-1-yl)-ethanone**

1-(6-Amino-indol-1-yl)-ethanone (**B-3**) was synthesized following the general scheme above starting from 6-nitroindole and acetyl chloride. Overall yield (~ 40 %). HPLC ret. time 0.54 min, 10–99 % CH_3CN , 5 min run; ESI-MS 175.1 m/z (MH^+).

[00328] **Example 2:**



[00329] {[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid ethyl ester

To a stirred solution of (*tert*-butoxycarbonyl-methyl-amino)-acetic acid (37 g, 0.2 mol) and Et₃N (60.6 g, 0.6 mol) in CH₂Cl₂ (300 mL) was added isobutyl chloroformate (27.3 g, 0.2 mmol) dropwise at −20 °C under argon. After stirring for 0.5 h, methylamino-acetic acid ethyl ester hydrochloride (30.5 g, 129 mmol) was added dropwise at −20 °C. The mixture was allowed to warm to room temperature (c.a. 1 h) and quenched with water (500 mL). The organic layer was separated, washed with 10 % citric acid solution, dried over Na₂SO₄, filtered and concentrated. The residue was purified by column chromatography (petroleum ether / EtOAc 1:1) to give {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid ethyl ester (12.5 g, 22 %).

[00330] {[2-(*tert*-Butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid

A suspension of {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid ethyl ester (12.3 g, 42.7 mmol) and LiOH (8.9 g, 214 mmol) in H₂O (20 mL) and THF (100 mL) was stirred overnight. Volatile solvent was removed under *vacuum* and the residue was extracted with ether (2 x 100 mL). The aqueous phase was acidified to pH 3 with dilute HCl solution, and then extracted with CH₂Cl₂ (2 x 300 mL). The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under *vacuum* to give {[2-(*tert*-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid as a colorless oil (10 g, 90 %). ¹H NMR (CDCl₃) δ

53568-20

7.17 (br s, 1 H), 4.14-4.04 (m, 4 H), 3.04-2.88 (m, 6 H), 1.45-1.41 (m, 9 H); ESI-MS 282.9 m/z (M+Na⁺).


[00331] Methyl-({methyl-[2-(6-nitro-indol-1-yl)-2-oxo-ethyl]-carbamoyl}-methyl)-carbamic acid tert-butyl ester

To a mixture of {[2-(tert-butoxycarbonyl-methyl-amino)-acetyl]-methyl-amino}-acetic acid (13.8g, 53 mmol) and TFFH (21.0g, 79.5 mmol) in anhydrous THF (125 mL) was added DIEA (27.7 mL, 159 mmol) at room temperature under nitrogen. The solution was stirred at room temperature for 20 min. A solution of 6-nitroindole (8.6g, 53 mmol) in THF (75 mL) was added and the reaction mixture was heated at 60 °C for 18 h. The solvent was evaporated and the crude mixture was re-partitioned between EtOAc and water. The organic layer was separated, washed with water (x 3), dried over Na₂SO₄ and concentrated. Diethyl ether followed by EtOAc was added. The resulting solid was collected via filtration, washed with diethyl ether and air dried to yield methyl-({methyl-[2-(6-nitro-indol-1-yl)-2-oxo-ethyl]-carbamoyl}-methyl)-carbamic acid *tert*-butyl ester (6.42 g, 30 %). ¹H NMR (400 MHz, DMSO-d₆) δ 1.37 (m, 9H), 2.78 (m, 3H), 2.95 (d, J = 1.5 Hz, 1H), 3.12 (d, J = 2.1 Hz, 2H), 4.01 (d, J = 13.8 Hz, 0.6H), 4.18 (d, J = 12.0 Hz, 1.4H), 4.92 (d, J = 3.4 Hz, 1.4H), 5.08 (d, J = 11.4 Hz, 0.6H), 7.03 (m, 1H), 7.90 (m, 1H), 8.21 (m, 1H), 8.35 (d, J = 3.8 Hz, 1H), 9.18 (m, 1H); HPLC ret. time 3.12 min, 10-99 % CH₃CN, 5 min run; ESI-MS 405.5 m/z (MH⁺).

[00332] B-26; ({[2-(6-Amino-indol-1-yl)-2-oxo-ethyl]-methyl-carbamoyl}-methyl)-methyl-carbamic acid tert-butyl ester

A mixture of methyl-({methyl-[2-(6-nitro-indol-1-yl)-2-oxo-ethyl]-carbamoyl}-methyl)-carbamic acid *tert*-butyl ester (12.4 g, 30.6 mmol), SnCl₂·2H₂O (34.5g, 153.2 mmol) and DIEA (74.8 mL, 429 mmol) in ethanol (112 mL) was heated to 70 °C for 3 h. Water and EtOAc were added and the mixture was filtered through a short plug of CeliteTM. The organic layer was separated, dried over Na₂SO₄ and concentrated to yield ({[2-(6-Amino-indol-1-yl)-2-oxo-ethyl]-methyl-carbamoyl}-methyl)-methyl-carbamic acid *tert*-butyl ester (**B-26**) (11.4 g, quant.). HPLC ret. time 2.11 min, 10-99 % CH₃CN, 5 min run; ESI-MS 375.3 m/z (MH⁺).

[00333] 2-Substituted 6-aminoindoles

[00334] Example 1:**[00335] B-4-a; (3-Nitro-phenyl)-hydrazine hydrochloride salt**

3-Nitrophenylamine (27.6 g, 0.2 mol) was dissolved in a mixture of H_2O (40 mL) and 37% HCl (40 mL). A solution of NaNO_2 (13.8 g, 0.2 mol) in H_2O (60 mL) was added at 0 °C, followed by the addition of $\text{SnCl}_2 \cdot \text{H}_2\text{O}$ (135.5 g, 0.6 mol) in 37% HCl (100 mL) at that temperature. After stirring at 0 °C for 0.5 h, the solid was isolated via filtration and washed with water to give (3-nitro-phenyl)-hydrazine hydrochloride salt (**B-4-a**) (27.6 g, 73 %).

[00336] 2-[(3-Nitro-phenyl)-hydrazono]-propionic acid ethyl ester

(3-Nitro-phenyl)-hydrazine hydrochloride salt (**B-4-a**) (30.2 g, 0.16 mol) and 2-oxo-propionic acid ethyl ester (22.3 g, 0.19 mol) was dissolved in ethanol (300 mL). The mixture was stirred at room temperature for 4 h. The solvent was evaporated under reduced pressure to give 2-[(3-nitro-phenyl)-hydrazono]-propionic acid ethyl ester, which was used directly in the next step.

[00337] B-4-b; 4-Nitro-1H-indole-2-carboxylic acid ethyl ester and 6-Nitro-1H-indole-2-carboxylic acid ethyl ester


2-[(3-Nitro-phenyl)-hydrazono]-propionic acid ethyl ester from the preceding step was dissolved in toluene (300 mL). PPA (30 g) was added. The mixture was heated at reflux overnight and then cooled to room temperature. The solvent was removed to give a mixture of 4-nitro-1H-

indole-2-carboxylic acid ethyl ester and 6-nitro-1H-indole-2-carboxylic acid ethyl ester (**B-4-b**) (15 g, 40 %).

[00338] B-4; 2-Methyl-1H-indol-6-ylamine

To a suspension of LiAlH_4 (7.8 g, 0.21 mol) in THF (300 mL) was added dropwise a mixture of 4-nitro-1H-indole-2-carboxylic acid ethyl ester and 6-nitro-1H-indole-2-carboxylic acid ethyl ester (**B-4-b**) (6g, 25.7 mmol) in THF (50 mL) at 0 °C under N_2 . The mixture was heated at reflux overnight and then cooled to 0 °C. H_2O (7.8 mL) and 10 % NaOH (7.8 mL) were added to the mixture at 0 °C. The insoluble solid was removed via filtration. The filtrate was dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography to afford 2-methyl-1H-indol-6-ylamine (**B-4**) (0.3 g, 8 %). ^1H NMR (CDCl_3) δ 7.57 (br s, 1 H), 7.27 (d, $J = 8.8$ Hz, 1 H), 6.62 (s, 1 H), 6.51-6.53 (m, 1 H), 6.07 (s, 1 H), 3.59-3.25 (br s, 2 H), 2.37 (s, 3H); ESI-MS 147.2 m/z (MH^+).

[00339] Example 2:



[00340] 6-Nitro-1H-indole-2-carboxylic acid and 4-Nitro-1H-indole-2-carboxylic acid

A mixture of 4-nitro-1H-indole-2-carboxylic acid ethyl ester and 6-nitro-1H-indole-2-carboxylic acid ethyl ester (**B-4-b**) (0.5 g, 2.13 mmol) in 10 % NaOH (20 mL) was heated at reflux overnight and then cooled to room temperature. The mixture was extracted with ether. The aqueous phase was separated and acidified with HCl to pH 1-2. The resulting solid was isolated via filtration to give a mixture of 6-nitro-1H-indole-2-carboxylic acid and 4-nitro-1H-indole-2-carboxylic acid (0.3 g, 68 %).

[00341] 6-Nitro-1H-indole-2-carboxylic acid amide and 4-Nitro-1H-indole-2-carboxylic acid amide

A mixture of 6-nitro-1H-indole-2-carboxylic acid and 4-nitro-1H-indole-2-carboxylic acid (12 g, 58 mmol) and SOCl₂ (50 mL, 64 mmol) in benzene (150 mL) was refluxed for 2 h. The benzene and excessive SOCl₂ was removed under reduced pressure. The residue was dissolved in CH₂Cl₂ (250 mL). NH₄OH (21.76 g, 0.32 mol) was added dropwise at 0 °C. The mixture was stirred at room temperature for 1 h. The resulting solid was isolated via filtration to give a crude mixture of 6-nitro-1H-indole-2-carboxylic acid amide and 4-nitro-1H-indole-2-carboxylic acid amide (9 g, 68 %), which was used directly in the next step.

[00342] 6-Nitro-1H-indole-2-carbonitrile and 4-Nitro-1H-indole-2-carbonitrile


A mixture of 6-nitro-1H-indole-2-carboxylic acid amide and 4-nitro-1H-indole-2-carboxylic acid amide (5 g, 24 mmol) was dissolved in CH₂Cl₂ (200 mL). Et₃N (24.24 g, 0.24 mol) was added, followed by the addition of (CF₃CO)₂O (51.24 g, 0.24 mol) at room temperature. The mixture was stirred for 1 h and poured into water (100 mL). The organic layer was separated. The aqueous layer was extracted with EtOAc (100 mL x 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure. The crude residue was purified by column chromatography to give a mixture of 6-nitro-1H-indole-2-carbonitrile and 4-nitro-1H-indole-2-carbonitrile (2.5 g, 55 %).

[00343] B-5; 6-Amino-1H-indole-2-carbonitrile

A mixture of 6-nitro-1H-indole-2-carbonitrile and 4-nitro-1H-indole-2-carbonitrile (2.5 g, 13.4 mmol) and Raney Ni (500 mg) in EtOH (50 mL) was stirred at room temperature under H₂ (1

atm) for 1 h. Raney Ni was filtered off. The filtrate was evaporated under reduced pressure and purified by column chromatography to give 6-amino-1H-indole-2-carbonitrile (**B-5**) (1 g, 49 %). ^1H NMR (DMSO- d_6) δ 12.75 (br s, 1 H), 7.82 (d, $J = 8$ Hz, 1 H), 7.57 (s, 1 H), 7.42 (s, 1 H), 7.15 (d, $J = 8$ Hz, 1 H); ESI-MS 158.2 m/z (MH^+).

[00344] **Example 3:**



[00345] **2,2-Dimethyl-N-*o*-tolyl-propionamide**

To a solution of *o*-tolylamine (21.4 g, 0.20 mol) and Et_3N (22.3 g, 0.22 mol) in CH_2Cl_2 was added 2,2-dimethyl-propionyl chloride (25.3 g, 0.21 mol) at 10 °C. The mixture was stirred overnight at room temperature, washed with aq. HCl (5%, 80 mL), saturated NaHCO_3 solution and brine, dried over Na_2SO_4 and concentrated under *vacuum* to give 2,2-dimethyl-N-*o*-tolyl-propionamide (35.0 g, 92 %).

[00346] **2-*tert*-Butyl-1H-indole**

To a solution of 2,2-dimethyl-N-*o*-tolyl-propionamide (30.0 g, 159 mmol) in dry THF (100 mL) was added dropwise *n*-BuLi (2.5 M, in hexane, 190 mL) at 15 °C. The mixture was stirred

overnight at 15 °C, cooled in an ice-water bath and treated with saturated NH₄Cl solution. The organic layer was separated and the aqueous layer was extracted with ethyl acetate. The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and concentrated in *vacuum*. The residue was purified by column chromatography to give 2-*tert*-butyl-1H-indole (23.8 g, 88 %).

[00347] 2-*tert*-Butyl-2,3-dihydro-1H-indole

To a solution of 2-*tert*-butyl-1H-indole (5.0 g, 29 mmol) in AcOH (20 mL) was added NaBH₄ at 10 °C. The mixture was stirred for 20 min at 10 °C, treated dropwise with H₂O under ice cooling, and extracted with ethyl acetate. The combined organic layers were dried over anhydrous Na₂SO₄, filtered, and concentrated under *vacuum* to give a mixture of starting material and 2-*tert*-butyl-2,3-dihydro-1H-indole (4.9 g), which was used directly in the next step.

[00348] 2-*tert*-Butyl-6-nitro-2,3-dihydro-1H-indole

To a solution of the mixture of 2-*tert*-butyl-2,3-dihydro-1H-indole and 2-*tert*-butyl-1H-indole (9.7 g) in H₂SO₄ (98%, 80 mL) was slowly added KNO₃ (5.6 g, 55.7 mmol) at 0 °C. The reaction mixture was stirred at room temperature for 1 h, carefully poured into cracked ice, basified with Na₂CO₃ to pH~8 and extracted with ethyl acetate. The combined extracts were washed with brine, dried over anhydrous Na₂SO₄ and concentrated under *vacuum*. The residue was purified by column chromatography to give 2-*tert*-butyl-6-nitro-2,3-dihydro-1H-indole (4.0 g, 32 % over 2 steps).

[00349] 2-*tert*-Butyl-6-nitro-1H-indole

To a solution of 2-*tert*-butyl-6-nitro-2,3-dihydro-1H-indole (2.0 g, 9.1 mmol) in 1,4-dioxane (20 mL) was added DDQ at room temperature. After refluxing for 2.5 h, the mixture was filtered and the filtrate was concentrated under *vacuum*. The residue was purified by column chromatography to give 2-*tert*-butyl-6-nitro-1H-indole (1.6 g, 80 %).


[00350] B-6; 2-*tert*-Butyl-1H-indol-6-ylamine

To a solution of 2-*tert*-butyl-6-nitro-1H-indole (1.3 g, 6.0 mmol) in MeOH (10 mL) was added Raney Ni (0.2 g). The mixture was stirred at room temperature under H₂ (1 atm) for 3 h. The

reaction mixture was filtered and the filtrate was concentrated. The residue was washed with petroleum ether to give 2-*tert*-butyl-1H-indol-6-ylamine (**B-6**) (1.0 g, 89 %). ¹H NMR (DMSO-*d*₆) δ 10.19 (s, 1 H), 6.99 (d, *J* = 8.1 Hz, 1 H), 6.46 (s, 1 H), 6.25 (dd, *J* = 1.8, 8.1 Hz, 1 H), 5.79 (d, *J* = 1.8 Hz, 1 H), 4.52 (s, 2 H), 1.24 (s, 9 H); ESI-MS 189.1 *m/z* (MH⁺).

[00351] 3-Substituted 6-aminoindoles

[00352] **Example 1:**



[00353] **N-(3-Nitro-phenyl)-N'-propylidene-hydrazine**

Sodium hydroxide solution (10 %, 15 mL) was added slowly to a stirred suspension of (3-nitro-phenyl)-hydrazine hydrochloride salt (**B-4-a**) (1.89 g, 10 mmol) in ethanol (20 mL) until pH 6. Acetic acid (5 mL) was added to the mixture followed by propionaldehyde (0.7 g, 12 mmol). After stirring for 3 h at room temperature, the mixture was poured into ice-water and the resulting precipitate was isolated via filtration, washed with water and dried in air to obtain N-(3-nitro-phenyl)-N'-propylidene-hydrazine, which was used directly in the next step.

[00354] **3-Methyl-4-nitro-1H-indole and 3-Methyl-6-nitro-1H-indole**


A mixture of N-(3-nitro-phenyl)-N'-propylidene-hydrazine dissolved in 85 % H₃PO₄ (20 mL) and toluene (20 mL) was heated at 90-100 °C for 2 h. After cooling, toluene was removed under reduced pressure. The resultant oil was basified with 10 % NaOH to pH 8. The aqueous layer

was extracted with EtOAc (100 mL x 3). The combined organic layers were dried, filtered and concentrated under reduced pressure to afford a mixture of 3-methyl-4-nitro-1H-indole and 3-methyl-6-nitro-1H-indole (1.5 g, 86 % over two steps), which was used directly in the next step.

[00355] B-7; 3-Methyl-1H-indol-6-ylamine

A mixture of 3-methyl-4-nitro-1H-indole and 3-methyl-6-nitro-1H-indole (3 g, 17 mol) and 10 % Pd-C (0.5 g) in ethanol (30 mL) was stirred overnight under H₂ (1 atm) at room temperature. Pd-C was filtered off and the filtrate was concentrated under reduced pressure. The residue was purified by column chromatography to give 3-methyl-1H-indol-6-ylamine (**B-7**) (0.6 g, 24 %). ¹H NMR (CDCl₃) δ 7.59 (br s, 1 H), 7.34 (d, *J* = 8.0 Hz, 1 H), 6.77 (s, 1H), 6.64 (s, 1 H), 6.57 (m, 1 H), 3.57 (br s, 2 H), 2.28 (s, 3H); ESI-MS 147.2 m/z (MH⁺).

[00356] Example 2:



[00357] 6-Nitro-1H-indole-3-carbonitrile


To a solution of 6-nitroindole (4.86 g 30 mmol) in DMF (24.3 mL) and CH₃CN (243 mL) was added dropwise a solution of ClSO₂NCO (5 mL, 57 mmol) in CH₃CN (39 mL) at 0 °C . After addition, the reaction was allowed to warm to room temperature and stirred for 2 h. The mixture was poured into ice-water, basified with sat. NaHCO₃ solution to pH 7-8 and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated to give 6-nitro-1H-indole-3-carbonitrile (4.6 g, 82 %).

[00358] B-8; 6-Amino-1H-indole-3-carbonitrile

A suspension of 6-nitro-1H-indole-3-carbonitrile (4.6 g, 24.6 mmol) and 10% Pd-C (0.46 g) in EtOH (50 mL) was stirred under H₂ (1 atm) at room temperature overnight. After filtration, the filtrate was concentrated and the residue was purified by column chromatography (Pet. Ether / EtOAc = 3 / 1) to give 6-amino-1H-indole-3-carbonitrile (**B-8**) (1 g, 99 %) as a pink powder. ¹H

NMR (DMSO- d_6) δ 11.51 (s, 1 H), 7.84 (d, $J = 2.4$ Hz, 1 H), 7.22 (d, $J = 8.4$ Hz, 1 H), 6.62 (s, 1H), 6.56 (d, $J = 8.4$ Hz, 1 H), 5.0 (s, 2H); ESI-MS 157.1 m/z (MH^+).

[00359] Example 3:



[00360] Dimethyl-(6-nitro-1H-indol-3-ylmethyl)-amine

A solution of dimethylamine (25 g, 0.17 mol) and formaldehyde (14.4 mL, 0.15 mol) in acetic acid (100 mL) was stirred at 0 °C for 30 min. To this solution was added 6-nitro-1H-indole (20 g, 0.12 mol). After stirring for 3 days at room temperature, the mixture was poured into 15% aq. NaOH solution (500 mL) at 0 °C. The precipitate was collected via filtration and washed with water to give dimethyl-(6-nitro-1H-indol-3-ylmethyl)-amine (23 g, 87 %).

[00361] B-9-a; (6-Nitro-1H-indol-3-yl)-acetonitrile


To a mixture of DMF (35 mL) and MeI (74.6 g, 0.53 mol) in water (35 mL) and THF (400 mL) was added dimethyl-(6-nitro-1H-indol-3-ylmethyl)-amine (23 g, 0.105 mol). After the reaction mixture was refluxed for 10 min, potassium cyanide (54.6 g, 0.84 mol) was added and the mixture was kept refluxing overnight. The mixture was then cooled to room temperature and filtered. The filtrate was washed with brine (300 mL x 3), dried over Na_2SO_4 , filtered and concentrated. The residue was purified by column chromatography to give (6-nitro-1H-indol-3-yl)-acetonitrile (**B-9-a**) (7.5 g, 36 %).

[00362] B-9; (6-Amino-1H-indol-3-yl)-acetonitrile

A mixture of (6-nitro-1H-indol-3-yl)-acetonitrile (**B-9-a**) (1.5 g, 74.5 mmol) and 10 % Pd-C (300 mg) in EtOH (50 mL) was stirred at room temperature under H_2 (1 atm) for 5 h. Pd-C was

removed via filtration and the filtrate was evaporated to give (6-amino-1H-indol-3-yl)-acetonitrile (**B-9**) (1.1 g, 90 %). ^1H NMR ($\text{DMSO-}d_6$) δ 10.4 (br s, 1 H), 7.18 (d, $J = 8.4$ Hz, 1 H), 6.94 (s, 1H), 6.52 (s, 1 H), 6.42 (dd, $J = 8.4, 1.8$ Hz, 1 H), 4.76(s, 2 H), 3.88 (s, 2 H); ESI-MS 172.1 m/z (MH^+).

[00363] Example 4:




[00364] [2-(6-Nitro-1H-indol-3-yl)-ethyl]-carbamic acid *tert*-butyl ester


To a solution of (6-nitro-1H-indol-3-yl)-acetonitrile (**B-9-a**) (8.6 g, 42.8 mmol) in dry THF (200 mL) was added a solution of 2 M borane-dimethyl sulfide complex in THF (214 mL, 0.43 mol) at 0 °C. The mixture was heated at reflux overnight under nitrogen. The mixture was then cooled to room temperature and a solution of $(\text{Boc})_2\text{O}$ (14 g, 64.2 mmol) and Et_3N (89.0 mL, 0.64 mol) in THF was added. The reaction mixture was kept stirring overnight and then poured into ice-water. The organic layer was separated and the aqueous phase was extracted with EtOAc (200 x 3 mL). The combined organic layers were washed with water and brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The crude was purified by column chromatography to give [2-(6-nitro-1H-indol-3-yl)-ethyl]-carbamic acid *tert*-butyl ester (5 g, 38 %).

[00365] **B-10; [2-(6-Amino-1H-indol-3-yl)-ethyl]-carbamic acid *tert*-butyl ester**

A mixture of [2-(6-nitro-1H-indol-3-yl)-ethyl]-carbamic acid *tert*-butyl ester (5 g, 16.4 mmol) and Raney Ni (1 g) in EtOH (100 mL) was stirred at room temperature under H_2 (1 atm) for 5 h. Raney Ni was filtered off and the filtrate was evaporated under reduced pressure. The crude product was purified by column chromatography to give [2-(6-amino-1H-indol-3-yl)-ethyl]-carbamic acid *tert*-butyl ester (**B-10**) (3 g, 67 %). ^1H NMR ($\text{DMSO-}d_6$) δ 10.1 (br s, 1 H), 7.11 (d, $J = 8.4$ Hz, 1 H), 6.77-6.73 (m, 2 H), 6.46 (d, $J = 1.5$ Hz, 1 H), 6.32 (dd, $J = 8.4, 2.1$ Hz, 1 H), 4.62 (s, 2 H), 3.14-3.08 (m, 2 H), 2.67-2.62 (m, 2 H), 1.35 (s, 9H); ESI-MS 275.8 m/z (MH^+).

[00366] Example 5:**[00367] General Scheme:**

a) RX (X=Br,I), zinc triflate, TBAI, DIEA, toluene; b) H₂, Raney Ni, EtOH or SnCl₂·2H₂O, EtOH.

[00368] Specific example:**[00369] 3-*tert*-Butyl-6-nitro-1H-indole**


To a mixture of 6-nitroindole (1 g, 6.2 mmol), zinc triflate (2.06 g, 5.7 mmol) and TBAI (1.7 g, 5.16 mmol) in anhydrous toluene (11 mL) was added DIEA (1.47 g, 11.4 mmol) at room temperature under nitrogen. The reaction mixture was stirred for 10 min at 120 °C, followed by addition of *t*-butyl bromide (0.707 g, 5.16 mmol). The resulting mixture was stirred for 45 min at 120 °C. The solid was filtered off and the filtrate was concentrated to dryness and purified by column chromatography on silica gel (Pet.Ether./EtOAc 20:1) to give 3-*tert*-butyl-6-nitro-1H-indole as a yellow solid (0.25 g, 19 %). ¹H NMR (CDCl₃) δ 8.32 (d, *J* = 2.1 Hz, 1H), 8.00 (dd, *J* = 2.1, 14.4 Hz, 1H), 7.85 (d, *J* = 8.7 Hz, 1H), 7.25 (s, 1H), 1.46 (s, 9H).

[00370] B-11; 3-*tert*-Butyl-1H-indol-6-ylamine

A suspension of 3-*tert*-butyl-6-nitro-1H-indole (3.0 g, 13.7 mmol) and Raney Ni (0.5 g) in ethanol was stirred at room temperature under H₂ (1 atm) for 3 h. The catalyst was filtered off and the


filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel (Pet.Ether. / EtOAc 4 : 1) to give 3-*tert*-butyl-1H-indol-6-ylamine (**B-11**) (2.0 g, 77.3%) as a gray solid. ^1H NMR (CDCl_3): δ 7.58 (m, 2H), 6.73 (d, $J = 1.2$ Hz, 1H), 6.66 (s, 1H), 6.57 (dd, $J = 0.8, 8.6$ Hz, 1H), 3.60 (br s, 2H), 1.42 (s, 9H).

[00371] Other examples:




[00372] **B-12; 3-Ethyl-1H-indol-6-ylamine**

3-Ethyl-1H-indol-6-ylamine (**B-12**) was synthesized following the general scheme above starting from 6-nitroindole and ethyl bromide. Overall yield (42 %). HPLC ret. time 1.95 min, 10–99 % CH_3CN , 5 min run; ESI-MS 161.3 m/z (MH^+).




[00373] **B-13; 3-Isopropyl-1H-indol-6-ylamine**

3-Isopropyl-1H-indol-6-ylamine (**B-13**) was synthesized following the general scheme above starting from 6-nitroindole and isopropyl iodide. Overall yield (17 %). HPLC ret. time 2.06 min, 10–99 % CH_3CN , 5 min run; ESI-MS 175.2 m/z (MH^+).




[00374] **B-14; 3-sec-Butyl-1H-indol-6-ylamine**

3-*sec*-Butyl-1H-indol-6-ylamine (**B-14**) was synthesized following the general scheme above starting from 6-nitroindole and 2-bromobutane. Overall yield (20 %). HPLC ret. time 2.32 min, 10–99 % CH₃CN, 5 min run; ESI-MS 189.5 m/z (MH⁺).




[00375] B-15; 3-Cyclopentyl-1H-indol-6-ylamine

3- Cyclopentyl -1H-indol-6-ylamine (**B-15**) was synthesized following the general scheme above starting from 6-nitroindole and iodo-cyclopentane. Overall yield (16 %). HPLC ret. time 2.39 min, 10–99 % CH₃CN, 5 min run; ESI-MS 201.5 m/z (MH⁺)..



[00376] B-16; 3-(2-Ethoxy-ethyl)-1H-indol-6-ylamine


3-(2-Ethoxy-ethyl)-1H-indol-6-ylamine (**B-16**) was synthesized following the general scheme above starting from 6-nitroindole and 1-bromo-2-ethoxy-ethane. Overall yield (15 %). HPLC ret. time 1.56 min, 10–99 % CH₃CN, 5 min run; ESI-MS 205.1 m/z (MH⁺).



[00377] B-17; (6-Amino-1H-indol-3-yl)-acetic acid ethyl ester

(6-Amino-1H-indol-3-yl)-acetic acid ethyl ester (**B-17**) was synthesized following the general scheme above starting from 6-nitroindole and iodo-acetic acid ethyl ester. Overall yield (24 %). HPLC ret. time 0.95 min, 10–99 % CH₃CN, 5 min run; ESI-MS 219.2 m/z (MH⁺).

[00378] 4-Substituted 6-aminoindole



[00379] 2-Methyl-3,5-dinitro-benzoic acid

To a mixture of HNO₃ (95%, 80 mL) and H₂SO₄ (98%, 80 mL) was slowly added 2-methylbenzoic acid (50 g, 0.37 mol) at 0 °C. After addition, the reaction mixture was stirred for 1.5 h while keeping the temperature below 30 °C, poured into ice-water and stirred for 15 min. The resulting precipitate was collected via filtration and washed with water to give 2-methyl-3,5-dinitro-benzoic acid (70 g, 84 %).

[00380] 2-Methyl-3,5-dinitro-benzoic acid ethyl ester


A mixture of 2-methyl-3,5-dinitro-benzoic acid (50 g, 0.22 mol) in SOCl₂ (80 mL) was heated at reflux for 4 h and then was concentrated to dryness. CH₂Cl₂ (50 mL) and EtOH (80 mL) were added. The mixture was stirred at room temperature for 1 h, poured into ice-water and extracted with EtOAc (3 x 100 mL). The combined extracts were washed with sat. Na₂CO₃ (80 mL), water (2 x 100 mL) and brine (100 mL), dried over Na₂SO₄ and concentrated to dryness to give 2-methyl-3,5-dinitro-benzoic acid ethyl ester (50 g, 88 %).


[00381] 2-(2-Dimethylamino-vinyl)-3,5-dinitro-benzoic acid ethyl ester

A mixture of 2-methyl-3,5-dinitro-benzoic acid ethyl ester (35 g, 0.14 mol) and dimethoxymethyl-dimethyl-amine (32 g, 0.27 mol) in DMF (200 mL) was heated at 100 °C for 5 h. The mixture was poured into ice-water. The precipitate was collected via filtration and washed with water to give 2-(2-dimethylamino-vinyl)-3,5-dinitro-benzoic acid ethyl ester (11.3 g, 48 %).

[00382] B-18; 6-Amino-1H-indole-4-carboxylic acid ethyl ester

A mixture of 2-(2-dimethylamino-vinyl)-3,5-dinitro-benzoic acid ethyl ester (11.3 g, 0.037 mol) and SnCl₂ (83 g, 0.37 mol) in ethanol was heated at reflux for 4 h. The mixture was concentrated to dryness and the residue was poured into water and basified with sat. Na₂CO₃ solution to pH 8. The precipitate was filtered off and the filtrate was extracted with ethyl acetate (3 x 100 mL). The combined extracts were washed with water (2 x 100 mL) and brine (150 mL), dried over Na₂SO₄ and concentrated to dryness. The residue was purified by column chromatography on silica gel to give 6-amino-1H-indole-4-carboxylic acid ethyl ester (**B-18**) (3 g, 40 %). ¹H NMR (DMSO-*d*₆) δ 10.76 (br s, 1 H), 7.11-7.14 (m, 2 H), 6.81-6.82 (m, 1 H), 6.67-6.68 (m, 1 H), 4.94 (br s, 2 H), 4.32-4.25 (q, *J* = 7.2 Hz, 2 H), 1.35-1.31 (t, *J* = 7.2, 3 H). ESI-MS 205.0 m/z (MH⁺).

[00383] 5-Substituted 6-aminoindoles**[00384] Example 1:****[00385] General Scheme:****[00386] Specific example:**



1-Fluoro-5-methyl-2,4-dinitro-benzene

To a stirred solution of HNO₃ (60 mL) and H₂SO₄ (80 mL), cooled in an ice bath, was added 1-fluoro-3-methylbenzene (27.5g, 25 mmol) at such a rate that the temperature did not rise over 35 °C. The mixture was allowed to stir for 30 min at room temperature and poured into ice water (500 mL). The resulting precipitate (a mixture of the desired product and 1-fluoro-3-methyl-2,4-dinitrobenzene, approx. 7:3) was collected via filtration and purified by recrystallization from 50 mL isopropyl ether to give 1-fluoro-5-methyl-2,4-dinitrobenzene as a white solid (18 g, 36 %).


[00387] [2-(5-Fluoro-2,4-dinitro-phenyl)-vinyl]-dimethyl-amine

A mixture of 1-fluoro-5-methyl-2,4-dinitrobenzene (10 g, 50 mmol), dimethoxymethyl-dimethylamine (11.9 g, 100 mmol) and DMF (50 mL) was heated at 100 °C for 4 h. The solution was cooled and poured into water. The red precipitate was collected via filtration, washed with water adequately and dried to give [2-(5-fluoro-2,4-dinitro-phenyl)-vinyl]-dimethyl-amine (8 g, 63 %).

[00388] B-20; 5-Fluoro-1H-indol-6-ylamine


A suspension of [2-(5-fluoro-2,4-dinitro-phenyl)-vinyl]-dimethyl-amine (8 g, 31.4 mmol) and Raney Ni (8 g) in EtOH (80 mL) was stirred under H₂ (40 psi) at room temperature for 1 h. After filtration, the filtrate was concentrated and the residue was purified by chromatography (Pet.Ether/ EtOAc = 5 / 1) to give 5-fluoro-1H-indol-6-ylamine (**B-20**) as a brown solid (1 g, 16 %). ¹H NMR (DMSO-*d*₆) δ 10.56 (br s, 1 H), 7.07 (d, *J* = 12 Hz, 1 H), 7.02 (m, 1H), 6.71 (d, *J* = 8 Hz, 1H), 6.17 (s, 1H), 3.91 (br s, 2H); ESI-MS 150.1 m/z (MH⁺).

[00389] Other examples:




[00390] B-21; 5-Chloro-1H-indol-6-ylamine

5-Chloro-1H-indol-6-ylamine (**B-21**) was synthesized following the general scheme above starting from 1-chloro-3-methyl-benzene. Overall yield (7 %). ^1H NMR (CDCl_3) δ 7.85 (br s, 1 H), 7.52 (s, 1 H), 7.03 (s, 1H), 6.79 (s, 1H), 6.34 (s, 1H), 3.91 (br s, 2H); ESI-MS 166.0 m/z (MH^+).

**[00391] B-22; 5-Trifluoromethyl-1H-indol-6-ylamine**

5-Trifluoromethyl-1H-indol-6-ylamine (**B-22**) was synthesized following the general scheme above starting from 1-methyl-3-trifluoromethyl-benzene. Overall yield (2 %). ^1H NMR ($\text{DMSO}-d_6$) 10.79 (br s, 1 H), 7.55 (s, 1 H), 7.12 (s, 1 H), 6.78 (s, 1 H), 6.27 (s, 1 H), 4.92 (s, 2 H); ESI-MS 200.8 m/z (MH^+).

[00392] Example 2:**[00393] 1-Benzenesulfonyl-2,3-dihydro-1H-indole**

To a mixture of DMAP (1.5 g), benzenesulfonyl chloride (24 g, 136 mmol) and 2,3-dihydro-1H-indole (14.7 g, 124 mmol) in CH_2Cl_2 (200 mL) was added dropwise Et_3N (19 g, 186 mmol) in an ice-water bath. After addition, the mixture was stirred at room temperature overnight, washed with water, dried over Na_2SO_4 and concentrated to dryness under reduced pressure to provide 1-benzenesulfonyl-2,3-dihydro-1H-indole (30.9 g, 96 %).

[00394] 1-(1-Benzenesulfonyl-2,3-dihydro-1H-indol-5-yl)-ethanone

To a stirring suspension of AlCl_3 (144 g, 1.08 mol) in CH_2Cl_2 (1070 mL) was added acetic anhydride (54 mL). The mixture was stirred for 15 minutes. A solution of 1-benzenesulfonyl-2,3-dihydro-1H-indole (46.9 g, 0.18 mol) in CH_2Cl_2 (1070 mL) was added dropwise. The mixture was stirred for 5 h and quenched by the slow addition of crushed ice. The organic layer was separated and the aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were washed with saturated aqueous NaHCO_3 and brine, dried over Na_2SO_4 and concentrated under *vacuum* to yield 1-(1-benzenesulfonyl-2,3-dihydro-1H-indol-5-yl)-ethanone (42.6 g, 79 %).

[00395] 1-Benzenesulfonyl-5-ethyl-2,3-dihydro-1H-indole

To magnetically stirred TFA (1600 mL) was added at 0 °C sodium borohydride (64 g, 1.69 mol) over 1 h. To this mixture was added dropwise a solution of 1-(1-benzenesulfonyl-2,3-dihydro-1H-indol-5-yl)-ethanone (40 g, 0.13 mol) in TFA (700 mL) over 1 h. The mixture was stirred overnight at 25 °C, diluted with H_2O (1600 mL), and basified with sodium hydroxide pellets at 0 °C. The organic layer was separated and the aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to give 1-benzenesulfonyl-5-ethyl-2,3-dihydro-1H-indole (16.2 g, 43 %).

[00396] 5-Ethyl-2,3-dihydro-1H-indole

A mixture of 1-benzenesulfonyl-5-ethyl-2,3-dihydro-1H-indole (15 g, 0.05 mol) in HBr (48%, 162 mL) was heated at reflux for 6 h. The mixture was basified with sat. NaOH solution to pH 9 and extracted with ethyl acetate. The organic layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to give 5-ethyl-2,3-dihydro-1H-indole (2.5 g, 32 %).

[00397] 5-Ethyl-6-nitro-2,3-dihydro-1H-indole

To a solution of 5-ethyl-2,3-dihydro-1H-indole (2.5 g, 17 mmol) in H_2SO_4 (98%, 20 mL) was slowly added KNO_3 (1.7 g, 17 mmol) at 0 °C. After addition, the mixture was stirred at 0 - 10

°C for 10 min, carefully poured into ice, basified with NaOH solution to pH 9 and extracted with ethyl acetate. The combined extracts were washed with brine, dried over Na₂SO₄ and concentrated to dryness. The residue was purified by column chromatography on silica gel to give 5-ethyl-6-nitro-2,3-dihydro-1H-indole (1.9 g, 58 %).


[00398] 5-Ethyl-6-nitro-1H-indole

To a solution of 5-ethyl-6-nitro-2,3-dihydro-1H-indole (1.9 g, 9.9 mmol) in CH₂Cl₂ (30 mL) was added MnO₂ (4 g, 46 mmol). The mixture was stirred at room temperature for 8 h. The solid was filtered off and the filtrate was concentrated to dryness to give crude 5-ethyl-6-nitro-1H-indole (1.9 g, quant.).

[00399] B-23; 5-Ethyl-1H-indol-6-ylamine

A suspension of 5-ethyl-6-nitro-1H-indole (1.9 g, 10 mmol) and Raney Ni (1 g) was stirred under H₂ (1 atm) at room temperature for 2 h. The catalyst was filtered off and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel to give 5-ethyl-1H-indol-6-ylamine (**B-23**) (760 mg, 48 %). ¹H NMR (CDCl₃) δ 7.90 (br s, 1H), 7.41 (s, 1H), 7.00 (s, 1H), 6.78 (s, 2H), 6.39 (s, 1H), 3.39 (br s, 2H), 2.63 (q, *J* = 7.2 Hz, 2H), 1.29 (t, *J* = 6.9 Hz, 3H); ESI-MS 161.1 m/z (MH⁺).

[00400] Example 3:



[00401] 2-Bromo-4-tert-butylphenylamine

To a solution of 4-*tert*-butyl-phenylamine (447 g, 3 mol) in DMF (500 mL) was added dropwise NBS (531 g, 3 mol) in DMF (500 mL) at room temperature. Upon completion, the reaction mixture was diluted with water and extracted with EtOAc. The organic layer was washed with water, brine, dried over Na₂SO₄ and concentrated. The crude product was directly used in the next step without further purification.

[00402] 2-Bromo-4-*tert*-butyl-5-nitro-phenylamine

2-Bromo-4-*tert*-butyl-phenylamine (162 g, 0.71 mol) was added dropwise to H₂SO₄ (410 mL) at room temperature to yield a clear solution. This clear solution was then cooled down to -5 to -10 °C. A solution of KNO₃ (82.5 g, 0.82 mol) in H₂SO₄ (410 mL) was added dropwise while the temperature was maintained between -5 to -10 °C. Upon completion, the reaction mixture was poured into ice / water and extracted with EtOAc. The combined organic layers were washed with 5% Na₂CO₃ and brine, dried over Na₂SO₄ and concentrated. The residue was purified by a column chromatography (EtOAc / petroleum ether 1 / 10) to give 2-bromo-4-*tert*-butyl-5-nitro-phenylamine as a yellow solid (152 g, 78 %).

[00403] 4-*tert*-Butyl-5-nitro-2-trimethylsilanylethynyl-phenylamine

To a mixture of 2-bromo-4-*tert*-butyl-5-nitro-phenylamine (27.3 g, 100 mmol) in toluene (200 mL) and water (100 mL) was added Et₃N (27.9 mL, 200 mmol), Pd(PPh₃)₂Cl₂ (2.11 g, 3 mmol), CuI (950 mg, 0.5 mmol) and trimethylsilyl acetylene (21.2 mL, 150 mmol) under a nitrogen atmosphere. The reaction mixture was heated at 70 °C in a sealed pressure flask for 2.5 h., cooled down to room temperature and filtered through a short plug of Celite. The filter cake was washed with EtOAc. The combined filtrate was washed with 5% NH₄OH solution and water, dried over Na₂SO₄ and concentrated. The crude product was purified by column chromatography (0 – 10 % EtOAc / petroleum ether) to provide 4-*tert*-butyl-5-nitro-2-trimethylsilanylethynyl-phenylamine as a brown viscous liquid (25 g, 81 %).

[00404] 5-*tert*-Butyl-6-nitro-1H-indole


To a solution of 4-*tert*-butyl-5-nitro-2-trimethylsilanylethynyl-phenylamine (25 g, 86 mmol) in DMF (100 mL) was added CuI (8.2 g, 43 mmol) under a nitrogen atmosphere. The mixture was heated at 135 °C in a sealed pressure flask overnight, cooled down to room temperature and

filtered through a short plug of Celite. The filter cake was washed with EtOAc. The combined filtrate was washed with water, dried over Na₂SO₄ and concentrated. The crude product was purified by column chromatography (10 – 20 % EtOAc / Hexane) to provide 5-*tert*-butyl-6-nitro-1H-indole as a yellow solid (12.9 g, 69 %).

[00405] B-24; 5-*tert*-Butyl-1H-indol-6-ylamine

Raney Ni (3 g) was added to 5-*tert*-butyl-6-nitro-1H-indole (14.7 g, 67 mmol) in methanol (100 mL). The mixture was stirred under hydrogen (1 atm) at 30 °C for 3 h. The catalyst was filtered off. The filtrate was dried over Na₂SO₄ and concentrated. The crude dark brown viscous oil was purified by column chromatography (10 – 20 % EtOAc / petroleum ether) to give 5-*tert*-butyl-1H-indol-6-ylamine (**B-24**) as a gray solid (11 g, 87 %). ¹H NMR (300 MHz, DMSO-d₆) δ 10.3 (br s, 1H), 7.2 (s, 1H), 6.9 (m, 1H), 6.6 (s, 1H), 6.1 (m, 1H), 4.4 (br s, 2H), 1.3 (s, 9H).

[00406] Example 4:



[00407] 5-Methyl-2,4-dinitro-benzoic acid


To a mixture of HNO₃ (95 %, 80 mL) and H₂SO₄ (98 %, 80 mL) was slowly added 3-methylbenzoic acid (50 g, 0.37 mol) at 0 °C. After addition, the mixture was stirred for 1.5 h while maintaining the temperature below 30 °C. The mixture was poured into ice-water and stirred for 15 min. The precipitate was collected via filtration and washed with water to give a mixture of 3-methyl-2,6-dinitro-benzoic acid and 5-methyl-2,4-dinitro-benzoic acid (70 g, 84 %). To a solution of this mixture in EtOH (150 mL) was added dropwise SOCl₂ (53.5 g, 0.45 mol). The mixture was heated at reflux for 2 h and concentrated to dryness under reduced pressure. The residue was dissolved in EtOAc (100 mL) and extracted with 10% Na₂CO₃ solution (120 mL). The organic layer was found to contain 5-methyl-2,4-dinitro-benzoic acid ethyl ester while the aqueous layer contained 3-methyl-2,6-dinitro-benzoic acid. The organic layer was washed with brine (50 mL), dried over Na₂SO₄ and concentrated to dryness to provide 5-methyl-2,4-dinitro-benzoic acid ethyl ester (20 g, 20 %).

[00408] 5-(2-Dimethylamino-vinyl)-2,4-dinitro-benzoic acid ethyl ester

A mixture of 5-methyl-2,4-dinitro-benzoic acid ethyl ester (39 g, 0.15 mol) and dimethoxymethyl-dimethylamine (32 g, 0.27 mol) in DMF (200 mL) was heated at 100 °C for 5 h. The mixture was poured into ice water. The precipitate was collected via filtration and washed with water to afford 5-(2-dimethylamino-vinyl)-2,4-dinitro-benzoic acid ethyl ester (15 g, 28 %).

[00409] B-25; 6-Amino-1H-indole-5-carboxylic acid ethyl ester

A mixture of 5-(2-dimethylamino-vinyl)-2,4-dinitro-benzoic acid ethyl ester (15 g, 0.05 mol) and Raney Ni (5 g) in EtOH (500 mL) was stirred under H₂ (50 psi) at room temperature for 2 h. The catalyst was filtered off and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel to give 6-amino-1H-indole-5-carboxylic acid ethyl ester (**B-25**) (3 g, 30 %). ¹H NMR (DMSO-*d*₆) δ 10.68 (s, 1 H), 7.99 (s, 1 H), 7.01-7.06 (m, 1 H), 6.62 (s, 1 H), 6.27-6.28 (m, 1 H), 6.16 (s, 2 H), 4.22 (q, *J* = 7.2 Hz, 2 H), 1.32-1.27 (t, *J* = 7.2 Hz, 3 H).

[00410] Example 5:**1-(2,3-Dihydro-indol-1-yl)-ethanone**

To a suspension of NaHCO₃ (504 g, 6.0 mol) and 2,3-dihydro-1H-indole (60 g, 0.5 mol) in CH₂Cl₂ (600 mL) cooled in an ice-water bath, was added dropwise acetyl chloride (78.5 g, 1.0

mol). The mixture was stirred at room temperature for 2 h. The solid was filtered off and the filtrate was concentrated to give 1-(2,3-dihydro-indol-1-yl)-ethanone (82 g, 100 %).

[00411] 1-(5-Bromo-2,3-dihydro-indol-1-yl)-ethanone

To a solution of 1-(2,3-dihydro-indol-1-yl)-ethanone (58.0 g, 0.36 mol) in acetic acid (3000 mL) was added Br₂ (87.0 g, 0.54 mol) at 10 °C. The mixture was stirred at room temperature for 4 h. The precipitate was collected via filtration to give crude 1-(5-bromo-2,3-dihydro-indol-1-yl)-ethanone (100 g, 96 %), which was used directly in the next step.

[00412] 5-Bromo-2,3-dihydro-1H-indole

A mixture of crude 1-(5-bromo-2,3-dihydro-indol-1-yl)-ethanone (100 g, 0.34 mol) in HCl (20 %, 1200 mL) was heated at reflux for 6 h. The mixture was basified with Na₂CO₃ to pH 8.5-10 and then extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel to give 5-bromo-2,3-dihydro-1H-indole (37 g, 55 %).

[00413] 5-Bromo-6-nitro-2,3-dihydro-1H-indole

To a solution of 5-bromo-2,3-dihydro-1H-indole (45 g, 0.227 mol) in H₂SO₄ (98 %, 200 mL) was slowly added KNO₃ (23.5 g, 0.23 mol) at 0 °C. After addition, the mixture was stirred at 0 – 10 °C for 4 h, carefully poured into ice, basified with Na₂CO₃ to pH 8 and extracted with ethyl acetate. The combined organic extracts were washed with brine, dried over Na₂SO₄ and concentrated to dryness. The residue was purified by column chromatography on silica gel to give 5-bromo-6-nitro-2,3-dihydro-1H-indole (42 g, 76 %).


[00414] 5-Bromo-6-nitro-1H-indole

To a solution of 5-bromo-6-nitro-2,3-dihydro-1H-indole (20 g, 82.3 mmol) in 1,4-dioxane (400 mL) was added DDQ (30 g, 0.13 mol). The mixture was stirred at 80 °C for 2 h. The solid was filtered off and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel to afford 5-bromo-6-nitro-1H-indole (7.5 g, 38 %).

[00415] B-27; 5-Bromo-1H-indol-6-ylamine

A mixture of 5-bromo-6-nitro-1H-indole (7.5 g, 31.1 mmol) and Raney Ni (1 g) in ethanol was stirred under H₂ (1 atm) at room temperature for 2 h. The catalyst was filtered off and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel to give 5-bromo-1H-indol-6-ylamine (**B-27**) (2 g, 30 %). ¹H NMR (DMSO-*d*₆) δ 10.6 (s, 1 H), 7.49 (s, 1 H), 6.79-7.02 (m, 1 H), 6.79 (s, 1 H), 6.14-6.16 (m, 1 H), 4.81 (s, 2 H).

[00416] 7-Substituted 6-aminoindole



[00417] 3-Methyl-2,6-dinitro-benzoic acid

To a mixture of HNO₃ (95 %, 80 mL) and H₂SO₄ (98 %, 80 mL) was slowly added 3-methylbenzoic acid (50 g, 0.37 mol) at 0 °C. After addition, the mixture was stirred for 1.5 h while maintaining the temperature below 30 °C. The mixture was poured into ice-water and stirred for 15 min. The precipitate was collected via filtration and washed with water to give a mixture of 3-methyl-2,6-dinitro-benzoic acid and 5-methyl-2,4-dinitro-benzoic acid (70 g, 84 %). To a solution of this mixture in EtOH (150 mL) was added dropwise SOCl₂ (53.5 g, 0.45 mol). The mixture was heated to reflux for 2 h and concentrated to dryness under reduced pressure. The residue was dissolved in EtOAc (100 mL) and extracted with 10% Na₂CO₃ solution (120 mL). The organic layer was found to contain 5-methyl-2,4-dinitro-benzoic acid ethyl ester. The aqueous layer was acidified with HCl to pH 2 ~ 3 and the resulting precipitate was collected via filtration, washed with water and dried in air to give 3-methyl-2,6-dinitro-benzoic acid (39 g, 47 %).

[00418] 3-Methyl-2,6-dinitro-benzoic acid ethyl ester


A mixture of 3-methyl-2,6-dinitro-benzoic acid (39 g, 0.15 mol) and SOCl_2 (80 mL) was heated at reflux for 4 h. The excess SOCl_2 was removed under reduced pressure and the residue was added dropwise to a solution of EtOH (100 mL) and Et_3N (50 mL). The mixture was stirred at 20 °C for 1 h and concentrated to dryness. The residue was dissolved in EtOAc (100 mL), washed with Na_2CO_3 (10 %, 40 mL x 2), water (50 mL x 2) and brine (50 mL), dried over Na_2SO_4 and concentrated to give 3-methyl-2,6-dinitro-benzoic acid ethyl ester (20 g, 53 %).

[00419] 3-(2-Dimethylamino-vinyl)-2,6-dinitro-benzoic acid ethyl ester

A mixture of 3-methyl-2,6-dinitro-benzoic acid ethyl ester (35 g, 0.14 mol) and dimethoxymethyl-dimethylamine (32 g, 0.27 mol) in DMF (200 mL) was heated at 100 °C for 5 h. The mixture was poured into ice water and the precipitate was collected via filtration and washed with water to give 3-(2-dimethylamino-vinyl)-2,6-dinitro-benzoic acid ethyl ester (25 g, 58 %).

[00420] B-19; 6-Amino-1H-indole-7-carboxylic acid ethyl ester

A mixture of 3-(2-dimethylamino-vinyl)-2,6-dinitro-benzoic acid ethyl ester (30 g, 0.097 mol) and Raney Ni (10 g) in EtOH (1000 mL) was stirred under H_2 (50 psi) for 2 h. The catalyst was filtered off, and the filtrate was concentrated to dryness. The residue was purified by column chromatography on silica gel to give 6-amino-1H-indole-7-carboxylic acid ethyl ester (**B-19**) as an off-white solid (3.2 g, 16 %). ^1H NMR ($\text{DMSO}-d_6$) δ 10.38 (s, 1 H), 7.44-7.41 (d, $J = 8.7$ Hz, 1 H), 6.98 (t, 1 H), 6.65 (s, 2 H), 6.50-6.46 (m, 1 H), 6.27-6.26 (m, 1 H), 4.43-4.36 (q, $J = 7.2$ Hz, 2 H), 1.35 (t, $J = 7.2$ Hz, 3 H).

[00421] Phenols**[00422] Example 1:**

[00423] 2-tert-Butyl-5-nitroaniline

To a cooled solution of sulfuric acid (90 %, 50 mL) was added dropwise 2-*tert*-butyl-phenylamine (4.5 g, 30 mmol) at 0 °C. Potassium nitrate (4.5 g, 45 mmol) was added in portions at 0 °C. The reaction mixture was stirred at 0-5 °C for 5 min, poured into ice-water and then extracted with EtOAc three times. The combined organic layers were washed with brine and dried over Na₂SO₄. After removal of solvent, the residue was purified by recrystallization using 70 % EtOH - H₂O to give 2-*tert*-butyl-5-nitroaniline (3.7 g, 64 %). ¹H NMR (400 MHz, CDCl₃) δ 7.56 (dd, J = 8.7, 2.4 Hz, 1H), 7.48 (d, J = 2.4 Hz, 1H), 7.36 (d, J = 8.7 Hz, 1H), 4.17 (s, 2H), 1.46 (s, 9H); HPLC ret. time 3.27 min, 10-99 % CH₃CN, 5 min run; ESI-MS 195.3 m/z (MH⁺).

[00424] C-1-a; 2-tert-Butyl-5-nitrophenol


To a mixture of 2-*tert*-butyl-5-nitroaniline (1.94 g, 10 mmol) in 40 mL of 15 % H₂SO₄ was added dropwise a solution of NaNO₂ (763 mg, 11.0 mmol) in water (3 mL) at 0 °C. The resulting mixture was stirred at 0-5 °C for 5 min. Excess NaNO₂ was neutralized with urea, then 5 mL of H₂SO₄-H₂O (v/v 1:2) was added and the mixture was refluxed for 5 min. Three additional 5 mL aliquots of H₂SO₄-H₂O (v/v 1:2) were added while heating at reflux. The reaction mixture was cooled to room temperature and extracted with EtOAc twice. The combined organic layers were washed with brine and dried over MgSO₄. After removal of solvent, the residue was purified by column chromatography (0-10 % EtOAc – Hexane) to give 2-*tert*-butyl-5-nitrophenol (**C-1-a**) (1.2 g, 62 %). ¹H NMR (400 MHz, CDCl₃) δ 7.76 (dd, J = 8.6, 2.2 Hz, 1H), 7.58 (d, J = 2.1 Hz, 1H), 7.43 (d, J = 8.6 Hz, 1H), 5.41 (s, 1H), 1.45 (s, 9H); HPLC ret. time 3.46 min, 10-99 % CH₃CN, 5 min run.

[00425] C-1; 2-tert-Butyl-5-aminophenol. To a refluxing solution of 2-*tert*-butyl-5-nitrophenol (**C-1-a**) (196 mg, 1.0 mmol) in EtOH (10 mL) was added ammonium formate (200 mg, 3.1 mmol), followed by 140 mg of 10% Pd-C. The reaction mixture was refluxed for additional 30 min, cooled to room temperature and filtered through a plug of Celite. The filtrate was concentrated to dryness and purified by column chromatography (20-30% EtOAc-Hexane) to give 2-*tert*-butyl-5-aminophenol (**C-1**) (144 mg, 87 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.76 (s, 1H), 6.74 (d, J = 8.3 Hz, 1H), 6.04 (d, J = 2.3 Hz, 1H), 5.93 (dd, J = 8.2, 2.3 Hz, 1H),

4.67 (s, 2H), 1.26 (s, 9H); HPLC ret. time 2.26 min, 10-99 % CH₃CN, 5 min run; ESI-MS 166.1 m/z (MH⁺).


[00426] Example 2:

[00427] General scheme:



a) RX (X = Br, I), K₂CO₃ or Cs₂CO₃, DMF; b) HCO₂NH₄ or HCO₂K, Pd-C, EtOH

[00428] Specific example:



[00429] 1-*tert*-Butyl-2-methoxy-4-nitrobenzene


To a mixture of 2-*tert*-butyl-5-nitrophenol (**C-1-a**) (100 mg, 0.52 mmol) and K₂CO₃ (86 mg, 0.62 mmol) in DMF (2 mL) was added CH₃I (40 μ L, 0.62 mmol). The reaction mixture was stirred at room temperature for 2 h, diluted with water and extracted with EtOAc. The combined organic layers were washed with brine and dried over MgSO₄. After filtration, the filtrate was evaporated to dryness to give 1-*tert*-butyl-2-methoxy-4-nitrobenzene (82 mg, 76 %) that was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.77 (t, J = 4.3 Hz, 1H), 7.70 (d, J = 2.3 Hz, 1H), 7.40 (d, J = 8.6 Hz, 1H), 3.94 (s, 3H), 1.39 (s, 9H).

[00430] C-2; 4-*tert*-Butyl-3-methoxyaniline

To a refluxing solution of 1-*tert*-butyl-2-methoxy-4-nitrobenzene (82 mg, 0.4 mmol) in EtOH (2 mL) was added potassium formate (300 mg, 3.6 mmol) in water (1 mL), followed by 10% Pd-C (15 mg). The reaction mixture was refluxed for additional 60 min, cooled to room temperature and filtered through Celite. The filtrate was concentrated to dryness to give 4-*tert*-butyl-3-


methoxyaniline (**C-2**) (52 mg, 72 %) that was used without further purification. HPLC ret. time 2.29 min, 10-99 % CH₃CN, 5 min run; ESI-MS 180.0 m/z (MH⁺).

[00431] Other examples:




[00432] **C-3; 3-(2-Ethoxyethoxy)-4-*tert*-butylbenzenamine**

3-(2-Ethoxyethoxy)-4-*tert*-butylbenzenamine (**C-3**) was synthesized following the general scheme above starting from 2-*tert*-butyl-5-nitrophenol (**C-1-a**) and 1-bromo-2-ethoxyethane. ¹H NMR (400 MHz, CDCl₃) δ 6.97 (d, J = 7.9 Hz, 1H), 6.17 (s, 1H), 6.14 (d, J = 2.3 Hz, 1H), 4.00 (t, J = 5.2 Hz, 2H), 3.76 (t, J = 5.2 Hz, 2H), 3.53 (q, J = 7.0 Hz, 2H), 1.27 (s, 9H), 1.16 (t, J = 7.0 Hz, 3H); HPLC ret. time 2.55 min, 10-99 % CH₃CN, 5 min run; ESI-MS 238.3 m/z (MH⁺).



[00433] **C-4; 2-(2-*tert*-Butyl-5-aminophenoxy)ethanol**

2-(2-*tert*-Butyl-5-aminophenoxy)ethanol (**C-4**) was synthesized following the general scheme above starting from 2-*tert*-butyl-5-nitrophenol (**C-1-a**) and 2-bromoethanol. HPLC ret. time 2.08 min, 10-99 % CH₃CN, 5 min run; ESI-MS 210.3 m/z (MH⁺).

[00434] Example 3:**[00435] N-(3-Hydroxy-phenyl)-acetamide and acetic acid 3-formylamino-phenyl ester**

To a well stirred suspension of 3-amino-phenol (50 g, 0.46 mol) and NaHCO_3 (193.2 g, 2.3 mol) in chloroform (1 L) was added dropwise chloroacetyl chloride (46.9 g, 0.6 mol) over a period of 30 min at 0 °C. After the addition was complete, the reaction mixture was refluxed overnight and then cooled to room temperature. The excess NaHCO_3 was removed via filtration. The filtrate was poured into water and extracted with EtOAc (300 x 3 mL). The combined organic layers were washed with brine (500 mL), dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give a mixture of N-(3-hydroxy-phenyl)-acetamide and acetic acid 3-formylamino-phenyl ester (35 g, 4:1 by NMR analysis). The mixture was used directly in the next step.

[00436] N-[3-(3-Methyl-but-3-enyloxy)-phenyl]-acetamide

A suspension of the mixture of N-(3-hydroxy-phenyl)-acetamide and acetic acid 3-formylamino-phenyl ester (18.12 g, 0.12 mol), 3-methyl-but-3-en-1-ol (8.6 g, 0.1 mol), DEAD (87 g, 0.2 mol) and Ph_3P (31.44 g, 0.12 mol) in benzene (250 mL) was heated at reflux overnight and then cooled to room temperature. The reaction mixture was poured into water and the organic layer was separated. The aqueous phase was extracted with EtOAc (300 x 3 mL). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 and concentrated. The

residue was purified by column chromatography to give N-[3-(3-methyl-but-3-enyloxy)-phenyl]-acetamide (11 g, 52 %).

[00437] N-(4,4-Dimethyl-chroman-7-yl)-acetamide


A mixture of N-[3-(3-methyl-but-3-enyloxy)-phenyl]-acetamide (2.5 g, 11.4 mmol) and AlCl_3 (4.52 g, 34.3 mmol) in fluoro-benzene (50 mL) was heated at reflux overnight. After cooling, the reaction mixture was poured into water. The organic layer was separated and the aqueous phase was extracted with EtOAc (40 x 3 mL). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 and concentrated under *vacuum*. The residue was purified by column chromatography to give N-(4,4-dimethyl-chroman-7-yl)-acetamide (1.35 g, 54 %).

[00438] C-5; 3,4-Dihydro-4,4-dimethyl-2H-chromen-7-amine

A mixture of N-(4,4-dimethyl-chroman-7-yl)-acetamide (1.35 g, 6.2 mmol) in 20 % HCl solution (30 mL) was heated at reflux for 3 h and then cooled to room temperature. The reaction mixture was basified with 10 % aq. NaOH to pH 8 and extracted with EtOAc (30 x 3 mL). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 and concentrated to give 3,4-dihydro-4,4-dimethyl-2H-chromen-7-amine (**C-5**) (1 g, 92 %). ^1H NMR ($\text{DMSO}-d_6$) δ 6.87 (d, $J = 8.4$ Hz, 1 H), 6.07 (dd, $J = 8.4, 2.4$ Hz, 1 H), 5.87 (d, $J = 2.4$ Hz, 1 H), 4.75 (s, 2 H), 3.99 (t, $J = 5.4$ Hz, 2 H), 1.64 (t, $J = 5.1$ Hz, 2 H), 1.15 (s, 6 H); ESI-MS 178.1 m/z (MH^+).


[00439] Example 4:

[00440] General scheme:



X = F, Cl; a) ROH, H₂SO₄ or MeSO₃H, CH₂Cl₂; b) R'CO₂Cl, Et₃N, 1,4-dioxane or CHCl₃; c) HNO₃, H₂SO₄ or KNO₃, H₂SO₄ or HNO₃, AcOH; d) piperidine, CH₂Cl₂; e) HCO₂NH₄, Pd-C, EtOH or SnCl₂.2H₂O, EtOH or H₂, Pd-C, MeOH.

[00441] Specific example



[00442] 2-tert-Butyl-4-fluorophenol

4-Fluorophenol (5g, 45 mmol) and *tert*-butanol (5.9 mL, 63 mmol) were dissolved in CH₂Cl₂ (80 mL) and treated with concentrated sulfuric acid (98 %, 3 mL). The mixture was stirred at room temperature overnight. The organic layer was washed with water, neutralized with NaHCO₃,

dried over MgSO_4 and concentrated. The residue was purified by column chromatography (5-15 % EtOAc - Hexane) to give 2-*tert*-butyl-4-fluorophenol (3.12 g, 42 %). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.32 (s, 1H), 6.89 (dd, $J = 11.1, 3.1$ Hz, 1H), 6.84-6.79 (m, 1H), 6.74 (dd, $J = 8.7, 5.3$ Hz, 1H), 1.33 (s, 9H).

[00443] 2-*tert*-Butyl-4-fluorophenyl methyl carbonate

To a solution of 2-*tert*-butyl-4-fluorophenol (2.63g, 15.7 mmol) and NEt_3 (3.13 mL, 22.5 mmol) in dioxane (45 mL) was added methyl chloroformate (1.27 mL, 16.5 mmol). The mixture was stirred at room temperature for 1 h. The precipitate was removed via filtration. The filtrate was then diluted with water and extracted with ether. The ether extract was washed with water and dried over MgSO_4 . After removal of solvent, the residue was purified by column chromatography to give 2-*tert*-butyl-4-fluorophenyl methyl carbonate (2.08g, 59 %). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.24 (dd, $J = 8.8, 5.4$ Hz, 1H), 7.17-7.10 (m, 2H), 3.86 (s, 3H), 1.29 (s, 9H).

[00444] 2-*tert*-Butyl-4-fluoro-5-nitrophenyl methyl carbonate (C-7-a) and 2-*tert*-butyl-4-fluoro-6-nitrophenyl methyl carbonate (C-6-a)

To a solution of 2-*tert*-butyl-4-fluorophenyl methyl carbonate (1.81g, 8 mmol) in H_2SO_4 (98 %, 1 mL) was added slowly a cooled mixture of H_2SO_4 (1 mL) and HNO_3 (1 mL) at 0 °C. The mixture was stirred for 2 h while warming to room temperature, poured into ice and extracted with diethyl ether. The ether extract was washed with brine, dried over MgSO_4 and concentrated. The residue was purified by column chromatography (0-10 % EtOAc - Hexane) to give 2-*tert*-butyl-4-fluoro-5-nitrophenyl methyl carbonate (C-7-a) (1.2 g, 55 %) and 2-*tert*-butyl-4-fluoro-6-nitrophenyl methyl carbonate (C-6-a) (270 mg, 12 %). 2-*tert*-Butyl-4-fluoro-5-nitrophenyl methyl carbonate (C-7-a): ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.24 (d, $J = 7.1$ Hz, 1H), 7.55 (d, $J = 13.4$ Hz, 1H), 3.90 (s, 3H), 1.32 (s, 9H). 2-*tert*-butyl-4-fluoro-6-nitrophenyl methyl carbonate (C-6-a): ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.04 (dd, $J = 7.6, 3.1$ Hz, 1H), 7.69 (dd, $J = 10.1, 3.1$ Hz, 1H), 3.91 (s, 3H), 1.35 (s, 9H).


[00445] 2-*tert*-Butyl-4-fluoro-5-nitrophenol

To a solution of 2-*tert*-butyl-4-fluoro-5-nitrophenyl methyl carbonate (**C-7-a**) (1.08 g, 4 mmol) in CH₂Cl₂ (40 mL) was added piperidine (3.94 mL, 10 mmol). The mixture was stirred at room temperature for 1 h and extracted with 1N NaOH (3x). The aqueous layer was acidified with 1N HCl and extracted with diethyl ether. The ether extract was washed with brine, dried (MgSO₄) and concentrated to give 2-*tert*-butyl-4-fluoro-5-nitrophenol (530 mg, 62 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.40 (s, 1H), 7.49 (d, *J* = 6.8 Hz, 1H), 7.25 (d, *J* = 13.7 Hz, 1H), 1.36 (s, 9H).

[00446] C-7; 2-*tert*-Butyl-5-amino-4-fluorophenol


To a refluxing solution of 2-*tert*-butyl-4-fluoro-5-nitrophenol (400 mg, 1.88 mmol) and ammonium formate (400 mg, 6.1 mmol) in EtOH (20 mL) was added 5 % Pd-C (260 mg). The mixture was refluxed for additional 1 h, cooled and filtered through Celite. The solvent was removed by evaporation to give 2-*tert*-butyl-5-amino-4-fluorophenol (**C-7**) (550 mg, 83 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.83 (br s, 1H), 6.66 (d, *J* = 13.7 Hz, 1H), 6.22 (d, *J* = 8.5 Hz, 1H), 4.74 (br s, 2H), 1.26 (s, 9H); HPLC ret. time 2.58 min, 10-99 % CH₃CN, 5 min run; ESI-MS 184.0 m/z (MH⁺).

[00447] Other examples:



[00448] C-10; 2-*tert*-Butyl-5-amino-4-chlorophenol


2-*tert*-Butyl-5-amino-4-chlorophenol (**C-10**) was synthesized following the general scheme above starting from 4-chlorophenol and *tert*-butanol. Overall yield (6 %). HPLC ret. time 3.07 min, 10-99 % CH₃CN, 5 min run; ESI-MS 200.2 m/z (MH⁺).



[00449] C-13; 5-Amino-4-fluoro-2-(1-methylcyclohexyl)phenol


5-Amino-4-fluoro-2-(1-methylcyclohexyl)phenol (**C-13**) was synthesized following the general scheme above starting from 4-fluorophenol and 1-methylcyclohexanol. Overall yield (3 %).

HPLC ret. time 3.00 min, 10-99 % CH₃CN, 5 min run; ESI-MS 224.2 m/z (MH⁺).




[00450] C-19; 5-Amino-2-(3-ethylpentan-3-yl)-4-fluoro-phenol

5-Amino-2-(3-ethylpentan-3-yl)-4-fluoro-phenol (**C-19**) was synthesized following the general scheme above starting from 4-fluorophenol and 3-ethyl-3-pentanol. Overall yield (1 %).




[00451] C-20; 2-Admantyl-5-amino-4-fluoro-phenol

2-Admantyl-5-amino-4-fluoro-phenol (**C-20**) was synthesized following the general scheme above starting from 4-fluorophenol and adamantan-1-ol.




[00452] C-21; 5-Amino-4-fluoro-2-(1-methylcycloheptyl)phenol


5-Amino-4-fluoro-2-(1-methylcycloheptyl)phenol (**C-21**) was synthesized following the general scheme above starting from 4-fluorophenol and 1-methyl-cycloheptanol.

**[00453] C-22; 5-Amino-4-fluoro-2-(1-methylcyclooctyl)phenol**

5-Amino-4-fluoro-2-(1-methylcyclooctyl)phenol (**C-22**) was synthesized following the general scheme above starting from 4-fluorophenol and 1-methyl-cyclooctanol.


**[00454] C-23; 5-Amino-2-(3-ethyl-2,2-dimethylpentan-3-yl)-4-fluoro-phenol**

5-Amino-2-(3-ethyl-2,2-dimethylpentan-3-yl)-4-fluoro-phenol (**C-23**) was synthesized following the general scheme above starting from 4-fluorophenol and 3-ethyl-2,2-dimethyl-pentan-3-ol.

[00455] Example 5:**[00456] C-6; 2-*tert*-Butyl-4-fluoro-6-aminophenyl methyl carbonate**

To a refluxing solution of 2-*tert*-butyl-4-fluoro-6-nitrophenyl methyl carbonate (250 mg, 0.92 mmol) and ammonium formate (250 mg, 4 mmol) in EtOH (10 mL) was added 5 % Pd-C (170 mg). The mixture was refluxed for additional 1 h, cooled and filtered through Celite. The solvent was removed by evaporation and the residue was purified by column chromatography (0-15 %, EtOAc – Hexane) to give 2-*tert*-butyl-4-fluoro-6-aminophenyl methyl carbonate (**C-6**) (60 mg, 27 %). HPLC ret. time 3.35 min, 10-99 % CH₃CN, 5 min run; ESI-MS 242.0 m/z (MH⁺).

[00457] Example 6:



[00458] Carbonic acid 2,4-di-*tert*-butyl-phenyl ester methyl ester

Methyl chloroformate (58 mL, 750 mmol) was added dropwise to a solution of 2,4-di-*tert*-butylphenol (103.2g, 500 mmol), Et₃N (139 mL, 1000 mmol) and DMAP (3.05g, 25 mmol) in dichloromethane (400 mL) cooled in an ice-water bath to 0 °C. The mixture was allowed to warm to room temperature while stirring overnight, then filtered through silica gel (approx. 1L) using 10% ethyl acetate – hexanes (~ 4 L) as the eluent. The combined filtrates were concentrated to yield carbonic acid 2,4-di-*tert*-butyl-phenyl ester methyl ester as a yellow oil (132 g, quant.). ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.35 (d, J = 2.4 Hz, 1H), 7.29 (dd, J = 8.5, 2.4 Hz, 1H), 7.06 (d, J = 8.4 Hz, 1H), 3.85 (s, 3H), 1.30 (s, 9H), 1.29 (s, 9H).

[00459] Carbonic acid 2,4-di-*tert*-butyl-5-nitro-phenyl ester methyl ester and Carbonic acid 2,4-di-*tert*-butyl-6-nitro-phenyl ester methyl ester

To a stirring mixture of carbonic acid 2,4-di-*tert*-butyl-phenyl ester methyl ester (4.76 g, 18 mmol) in conc. sulfuric acid (2 mL), cooled in an ice-water bath, was added a cooled mixture of sulfuric acid (2 mL) and nitric acid (2 mL). The addition was done slowly so that the reaction temperature did not exceed 50 °C. The reaction was allowed to stir for 2 h while warming to room temperature. The reaction mixture was then added to ice-water and extracted into diethyl ether. The ether layer was dried (MgSO₄), concentrated and purified by column chromatography (0 – 10% ethyl acetate – hexanes) to yield a mixture of carbonic acid 2,4-di-*tert*-butyl-5-nitro-phenyl ester methyl ester and carbonic acid 2,4-di-*tert*-butyl-6-nitro-phenyl ester methyl ester as a pale yellow solid (4.28 g), which was used directly in the next step.

[00460] 2,4-Di-*tert*-butyl-5-nitro-phenol and 2,4-Di-*tert*-butyl-6-nitro-phenol

The mixture of carbonic acid 2,4-di-*tert*-butyl-5-nitro-phenyl ester methyl ester and carbonic acid 2,4-di-*tert*-butyl-6-nitro-phenyl ester methyl ester (4.2 g, 12.9 mmol) was dissolved in MeOH (65 mL) and KOH (2.0g, 36 mmol) was added. The mixture was stirred at room temperature for 2 h. The reaction mixture was then made acidic (pH 2-3) by adding conc. HCl and partitioned between water and diethyl ether. The ether layer was dried (MgSO₄), concentrated and purified by column chromatography (0 – 5 % ethyl acetate – hexanes) to provide 2,4-di-*tert*-butyl-5-nitro-phenol (1.31 g, 29 % over 2 steps) and 2,4-di-*tert*-butyl-6-nitro-phenol. 2,4-Di-*tert*-butyl-5-nitro-phenol: ¹H NMR (400 MHz, DMSO-*d*₆) δ 10.14 (s, 1H, OH), 7.34 (s, 1H), 6.83 (s, 1H), 1.36 (s, 9H), 1.30 (s, 9H). 2,4-Di-*tert*-butyl-6-nitro-phenol: ¹H NMR (400 MHz, CDCl₃) δ 11.48 (s, 1H), 7.98 (d, J = 2.5 Hz, 1H), 7.66 (d, J = 2.4 Hz, 1H), 1.47 (s, 9H), 1.34 (s, 9H).

[00461] C-9; 5-Amino-2,4-di-*tert*-butyl-phenol


To a refluxing solution of 2,4-di-*tert*-butyl-5-nitro-phenol (1.86 g, 7.4 mmol) and ammonium formate (1.86 g) in ethanol (75 mL) was added Pd-5% wt. on activated carbon (900 mg). The reaction mixture was stirred at reflux for 2 h, cooled to room temperature and filtered through Celite. The Celite was washed with methanol and the combined filtrates were concentrated to yield 5-amino-2,4-di-*tert*-butyl-phenol as a grey solid (1.66 g, quant.). ¹H NMR (400 MHz,

DMSO-*d*₆) δ 8.64 (s, 1H, OH), 6.84 (s, 1H), 6.08 (s, 1H), 4.39 (s, 2H, NH₂), 1.27 (m, 18H); HPLC ret. time 2.72 min, 10-99 % CH₃CN, 5 min run; ESI-MS 222.4 m/z (MH⁺).

[00462] C-8; 6-Amino-2,4-di-*tert*-butyl-phenol

A solution of 2,4-di-*tert*-butyl-6-nitro-phenol (27 mg, 0.11 mmol) and SnCl₂·2H₂O (121 mg, 0.54 mmol) in EtOH (1.0 mL) was heated in microwave oven at 100 °C for 30 min. The mixture was diluted with EtOAc and water, basified with sat. NaHCO₃ and filtered through Celite. The organic layer was separated and dried over Na₂SO₄. Solvent was removed by evaporation to provide 6-amino-2,4-di-*tert*-butyl-phenol (**C-8**), which was used without further purification. HPLC ret. time 2.74 min, 10-99 % CH₃CN, 5 min run; ESI-MS 222.5 m/z (MH⁺).

[00463] Example 7:



[00464] 4-*tert*-butyl-2-chloro-phenol

To a solution of 4-*tert*-butyl-phenol (40.0 g, 0.27 mol) and SO₂Cl₂ (37.5 g, 0.28 mol) in CH₂Cl₂ was added MeOH (9.0 g, 0.28 mol) at 0 °C. After addition was complete, the mixture was stirred overnight at room temperature and then water (200 mL) was added. The resulting solution was extracted with ethyl acetate. The combined organic layers were dried over anhydrous Na₂SO₄, filtered and concentrated under *vacuum*. The residue was purified by column chromatography (Pet. Ether / EtOAc, 50:1) to give 4-*tert*-butyl-2-chloro-phenol (47.0 g, 95 %).

[00465] 4-*tert*-Butyl-2-chlorophenyl methyl carbonate

To a solution of 4-*tert*-butyl-2-chlorophenol (47.0 g, 0.25 mol) in dichloromethane (200 mL) was added Et₃N (50.5 g, 0.50 mol), DMAP (1 g) and methyl chloroformate (35.4 g, 0.38 mol) at 0 °C. The reaction was allowed to warm to room temperature and stirred for additional 30 min. The reaction mixture was washed with H₂O and the organic layer was dried over Na₂SO₄ and concentrated to give 4-*tert*-butyl-2-chlorophenyl methyl carbonate (56.6 g, 92 %), which was used directly in the next step.

[00466] 4-*tert*-Butyl-2-chloro-5-nitrophenyl methyl carbonate


4-*tert*-Butyl-2-chlorophenyl methyl carbonate (36.0 g, 0.15 mol) was dissolved in conc. H₂SO₄ (100 mL) at 0 °C. KNO₃ (0.53 g, 5.2 mmol) was added in portions over 25 min. The reaction was stirred for 1.5 h and poured into ice (200 g). The aqueous layer was extracted with dichloromethane. The combined organic layers were washed with aq. NaHCO₃, dried over Na₂SO₄ and concentrated under *vacuum* to give 4-*tert*-butyl-2-chloro-5-nitrophenyl methyl carbonate (41.0 g), which was used without further purification.

[00467] 4-*tert*-Butyl-2-chloro-5-nitro-phenol

Potassium hydroxide (10.1 g, 181 mmol) was added to 4-*tert*-butyl-2-chloro-5-nitrophenyl methyl carbonate (40.0 g, 139 mmol) in MeOH (100 mL). After 30 min, the reaction was acidified with 1N HCl and extracted with dichloromethane. The combined organic layers were combined, dried over Na₂SO₄ and concentrated under *vacuum*. The crude residue was purified by column chromatography (Pet. Ether / EtOAc, 30:1) to give 4-*tert*-butyl-2-chloro-5-nitro-phenol (23.0 g, 68 % over 2 steps).

[00468] C-11; 4-*tert*-Butyl-2-chloro-5-amino-phenol

To a solution of 4-*tert*-butyl-2-chloro-5-nitro-phenol (12.6 g, 54.9 mmol) in MeOH (50 mL) was added Ni (1.2 g). The reaction was shaken under H₂ (1 atm) for 4 h. The reaction mixture was filtered and the filtrate was concentrated. The residue was purified by column chromatography (P.E. / EtOAc, 20:1) to give 4-*tert*-butyl-2-chloro-5-amino-phenol (C-11) (8.5 g, 78 %). ¹H NMR (DMSO-*d*₆) δ 9.33 (s, 1 H), 6.80 (s, 1 H), 6.22 (s, 1 H), 4.76 (s, 1 H), 1.23 (s, 9 H); ESI-MS 200.1 m/z (MH⁺).

[00469] Example 8:**2-Admantyl-4-methyl-phenyl ethyl carbonate**

Ethyl chloroformate (0.64 mL, 6.7 mmol) was added dropwise to a solution of 2-admantyl-4-methylphenol (1.09 g, 4.5 mmol), Et_3N (1.25 mL, 9 mmol) and DMAP (catalytic amount) in dichloromethane (8 mL) cooled in an ice-water bath to 0 °C. The mixture was allowed to warm to room temperature while stirring overnight, then filtered and the filtrate was concentrated. The residue was purified by column chromatography (10-20 % ethyl acetate – hexanes) to yield 2-admantyl-4-methyl-phenyl ethyl carbonate as a yellow oil (1.32 g, 94 %).

[00470] 2-Admantyl-4-methyl-5-nitrophenyl ethyl carbonate

To a cooled solution of 2-admantyl-4-methyl-phenyl ethyl carbonate (1.32 g, 4.2 mmol) in H_2SO_4 (98 %, 10 mL) was added KNO_3 (510 mg, 5.0 mmol) in small portions at 0 °C. The mixture was stirred for 3 h while warming to room temperature, poured into ice and then extracted with dichloromethane. The combined organic layers were washed with NaHCO_3 and brine, dried over MgSO_4 and concentrated to dryness. The residue was purified by column chromatography (0-10 % EtOAc - Hexane) to yield 2-admantyl-4-methyl-5-nitrophenyl ethyl carbonate (378 mg, 25 %).


[00471] 2-Admantyl-4-methyl-5-nitrophenol

To a solution of 2-admantyl-4-methyl-5-nitrophenyl ethyl carbonate (378 mg, 1.05 mmol) in CH_2Cl_2 (5 mL) was added piperidine (1.0 mL). The solution was stirred at room temperature for 1 h, adsorbed onto silica gel under reduced pressure and purified by flash chromatography on silica gel (0-15 %, EtOAc - Hexanes) to provide 2-admantyl-4-methyl-5-nitrophenol (231 mg, 77 %).

[00472] C-12; 2-Admantyl-4-methyl-5-aminophenol

To a solution of 2-admantyl-4-methyl-5-nitrophenol (231 mg, 1.6 mmol) in EtOH (2 mL) was added Pd- 5% wt on carbon (10 mg). The mixture was stirred under H_2 (1 atm) overnight and then filtered through Celite. The filtrate was evaporated to dryness to provide 2-admantyl-4-methyl-5-aminophenol (**C-12**), which was used without further purification. HPLC ret. time 2.52 min, 10-99 % CH_3CN , 5 min run; ESI-MS 258.3 m/z (MH^+).

[00473] Example 9:



2-tert-Butyl-4-bromophenol

To a solution of 2-tert-butylphenol (250g, 1.67 mol) in CH_3CN (1500 mL) was added NBS (300 g, 1.67 mol) at room temperature. After addition, the mixture was stirred at room temperature

overnight and then the solvent was removed. Petroleum ether (1000 mL) was added, and the resulting white precipitate was filtered off. The filtrate was concentrated under reduced pressure to give the crude 2-*tert*-butyl-4-bromophenol (380 g), which was used without further purification.

[00474] Methyl (2-*tert*-butyl-4-bromophenyl) carbonate

To a solution of 2-*t*-butyl-4-bromophenol (380 g, 1.67 mol) in dichloromethane (1000 mL) was added Et₃N (202 g, 2 mol) at room temperature. Methyl chloroformate (155 mL) was added dropwise to the above solution at 0 °C. After addition, the mixture was stirred at 0 °C for 2 h., quenched with saturated ammonium chloride solution and diluted with water. The organic layer was separated and washed with water and brine, dried over Na₂SO₄, and concentrated to provide the crude methyl (2-*tert*-butyl-4-bromophenyl) carbonate (470 g), which was used without further purification.

[00475] Methyl (2-*tert*-butyl-4-bromo-5-nitrophenyl) carbonate

Methyl (2-*tert*-butyl-4-bromophenyl) carbonate (470 g, 1.67 mol) was dissolved in conc. H₂SO₄ (1000 ml) at 0 °C. KNO₃ (253 g, 2.5 mol) was added in portions over 90 min. The reaction mixture was stirred at 0 °C for 2 h and poured into ice-water (20 L). The resulting precipitate was collected via filtration and washed with water thoroughly, dried and recrystallized from ether to give methyl (2-*tert*-butyl-4-bromo-5-nitrophenyl) carbonate (332 g, 60 % over 3 steps).

[00476] C-14-a; 2-*tert*-Butyl-4-bromo-5-nitro-phenol

To a solution of methyl (2-*tert*-butyl-4-bromo-5-nitrophenyl) carbonate (121.5 g, 0.366 mol) in methanol (1000 mL) was added potassium hydroxide (30.75 g, 0.549 mol) in portions. After addition, the mixture was stirred at room temperature for 3 h and acidified with 1N HCl to pH 7. Methanol was removed and water was added. The mixture was extracted with ethyl acetate and the organic layer was separated, dried over Na₂SO₄ and concentrated to give 2-*tert*-butyl-4-bromo-5-nitro-phenol (**C-14-a**) (100 g, 99 %).

[00477] 1-*tert*-Butyl-2-(benzyloxy)-5-bromo-4-nitrobenzene

To a mixture of 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) (1.1 g, 4 mmol) and Cs₂CO₃ (1.56 g, 4.8 mmol) in DMF (8 mL) was added benzyl bromide (500 μ L, 4.2 mmol). The mixture was stirred at room temperature for 4 h, diluted with H₂O and extracted twice with EtOAc. The combined organic layers were washed with brine and dried over MgSO₄. After removal of solvent, the residue was purified by column chromatography (0-5 % EtOAc - Hexane) to yield 1-*tert*-butyl-2-(benzyloxy)-5-bromo-4-nitrobenzene (1.37 g, 94 %). ¹H NMR (400 MHz, CDCl₃) 7.62 (s, 1H), 7.53 (s, 1H), 7.43 (m, 5H), 5.22 (s, 2H), 1.42 (s, 9H).

[00478] 1-*tert*-Butyl-2-(benzyloxy)-5-(trifluoromethyl)-4-nitrobenzene


A mixture of 1-*tert*-butyl-2-(benzyloxy)-5-bromo-4-nitrobenzene (913 mg, 2.5 mmol), KF (291 mg, 5 mmol), KBr (595 mg, 5 mmol), CuI (570 mg, 3 mmol), methyl chlorodifluoroacetate (1.6 mL, 15 mmol) and DMF (5 mL) was stirred at 125 °C in a sealed tube overnight, cooled to room temperature, diluted with water and extracted three times with EtOAc. The combined organic layers were washed with brine and dried over anhydrous MgSO₄. After removal of the solvent, the residue was purified by column chromatography (0-5 % EtOAc - Hexane) to yield 1-*tert*-butyl-2-(benzyloxy)-5-(trifluoromethyl)-4-nitrobenzene (591 mg, 67 %). ¹H NMR (400 MHz, CDCl₃) 7.66 (s, 1H), 7.37 (m, 5H), 7.19 (s, 1H), 5.21 (s, 2H), 1.32 (s, 9H).

[00479] C-14; 5-Amino-2-*tert*-butyl-4-trifluoromethyl-phenol


To a refluxing solution of 1-*tert*-butyl-2-(benzyloxy)-5-(trifluoromethyl)-4-nitrobenzene (353 mg, 1.0 mmol) and ammonium formate (350 mg, 5.4 mmol) in EtOH (10 mL) was added 10% Pd-C (245 mg). The mixture was refluxed for additional 2 h, cooled to room temperature and filtered through Celite. After removal of solvent, the residue was purified by column chromatography to give 5-Amino-2-*tert*-butyl-4-trifluoromethyl-phenol (**C-14**) (120 mg, 52 %). ¹H NMR (400 MHz, CDCl₃) δ 7.21 (s, 1H), 6.05 (s, 1H), 1.28 (s, 9H); HPLC ret. time 3.46 min, 10-99 % CH₃CN, 5 min run; ESI-MS 234.1 m/z (MH⁺).

[00480] Example 10:

[00481] General scheme:



a) ArB(OH)_2 , K_2CO_3 , $\text{Pd(PPh}_3)_4$, H_2O , DMF or ArB(OH)_2 , $(\text{dppf})\text{PdCl}_2$, K_2CO_3 , EtOH; b) H_2 , Raney Ni, MeOH or HCO_2NH_4 , Pd-C, EtOH or $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$.



[00483] 2-tert-Butyl-4-(2-ethoxyphenyl)-5-nitrophenol


To a solution of 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) (8.22 g, 30 mmol) in DMF (90 mL) was added 2-ethoxyphenyl boronic acid (5.48 g, 33 mmol), potassium carbonate (4.56 g, 33 mmol), water (10 mL) and $\text{Pd(PPh}_3)_4$ (1.73 g, 1.5 mmol). The mixture was heated at 90 °C for 3 h under nitrogen. The solvent was removed under reduced pressure. The residue was partitioned between water and ethyl acetate. The combined organic layers were washed with water and brine, dried and purified by column chromatography (petroleum ether - ethyl acetate, 10:1) to afford 2-*tert*-butyl-4-(2-ethoxyphenyl)-5-nitrophenol (9.2 g, 92 %). $^1\text{H NMR}$ ($\text{DMSO-}d_6$) δ 10.38 (s, 1 H), 7.36 (s, 1 H), 7.28 (m, 2 H), 7.08 (s, 1 H), 6.99 (t, 1 H, $J = 7.35$ Hz), 6.92 (d, 1 H, $J = 8.1$ Hz), 3.84 (q, 2 H, $J = 6.6$ Hz), 1.35 (s, 9 H), 1.09 (t, 3 H, $J = 6.6$ Hz); ESI-MS 314.3 m/z (MH^+).

[00484] C-15; 2-tert-Butyl-4-(2-ethoxyphenyl)-5-aminophenol

To a solution of 2-*tert*-butyl-4-(2-ethoxyphenyl)-5-nitrophenol (3.0 g, 9.5 mmol) in methanol (30 mL) was added Raney Ni (300 mg). The mixture was stirred under H_2 (1 atm) at room temperature for 2 h. The catalyst was filtered off and the filtrate was concentrated. The residue was purified by column chromatography (petroleum ether – ethyl acetate, 6:1) to afford 2-*tert*-


butyl-4-(2-ethoxyphenyl)-5-aminophenol (**C-15**) (2.35 g, 92 %). ^1H NMR ($\text{DMSO}-d_6$) δ 8.89 (s, 1H), 7.19 (t, 1H, $J = 4.2$ Hz), 7.10 (d, 1H, $J = 1.8$ Hz), 7.08 (d, 1H, $J = 1.8$ Hz), 6.94 (t, 1H, $J = 3.6$ Hz), 6.67 (s, 1 H), 6.16 (s, 1 H), 4.25 (s, 1 H), 4.00 (q, 2H, $J = 6.9$ Hz), 1.26 (s, 9H), 1.21 (t, 3 H, $J = 6.9$ Hz); ESI-MS 286.0 m/z (MH^+).

[00485] Other examples:



[00486] **C-16; 2-*tert*-Butyl-4-(3-ethoxyphenyl)-5-aminophenol**


2-*tert*-Butyl-4-(3-ethoxyphenyl)-5-aminophenol (**C-16**) was synthesized following the general scheme above starting from 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) and 3-ethoxyphenyl boronic acid. HPLC ret. time 2.77 min, 10-99 % CH_3CN , 5 min run; ESI-MS 286.1 m/z (MH^+).



[00487] **C-17; 2-*tert*-Butyl-4-(3-methoxycarbonylphenyl)-5-aminophenol (C-17)**

2-*tert*-Butyl-4-(3-methoxycarbonylphenyl)-5-aminophenol (**C-17**) was synthesized following the general scheme above starting from 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) and 3-(methoxycarbonyl)phenylboronic acid. HPLC ret. time 2.70 min, 10-99 % CH_3CN , 5 min run; ESI-MS 300.5 m/z (MH^+).

[00488] **Example 11;**



[00489] 1-*tert*-Butyl-2-methoxy-5-bromo-4-nitrobenzene

To a mixture of 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) (1.5 g, 5.5 mmol) and Cs_2CO_3 (2.2 g, 6.6 mmol) in DMF (6 mL) was added methyl iodide (5150 μL , 8.3 mmol). The mixture was stirred at room temperature for 4 h, diluted with H_2O and extracted twice with EtOAc. The combined organic layers were washed with brine and dried over MgSO_4 . After removal of solvent, the residue was washed with hexane to yield 1-*tert*-butyl-2-methoxy-5-bromo-4-nitrobenzene (1.1 g, 69 %). ^1H NMR (400 MHz, CDCl_3) δ 7.58 (s, 1H), 7.44 (s, 1H), 3.92 (s, 3H), 1.39 (s, 9H).


[00490] 1-*tert*-Butyl-2-methoxy-5-(trifluoromethyl)-4-nitrobenzene

A mixture of 1-*tert*-butyl-2-methoxy-5-bromo-4-nitrobenzene (867 mg, 3.0 mmol), KF (348 mg, 6 mmol), KBr (714 mg, 6 mmol), CuI (684 mg, 3.6 mmol), methyl chlorodifluoroacetate (2.2 mL, 21.0 mmol) in DMF (5 mL) was stirred at 125 $^\circ\text{C}$ in a sealed tube overnight, cooled to room temperature, diluted with water and extracted three times with EtOAc. The combined organic layers were washed with brine and dried over anhydrous MgSO_4 . After removal of the solvent, the residue was purified by column chromatography (0-5 % EtOAc - Hexane) to yield 1-*tert*-butyl-2-methoxy-5-(trifluoromethyl)-4-nitrobenzene (512 mg, 61 %). ^1H NMR (400 MHz, CDCl_3) δ 7.60 (s, 1H), 7.29 (s, 1H), 3.90 (s, 3H), 1.33 (s, 9H).

[00491] C-18; 1-*tert*-Butyl-2-methoxy-5-(trifluoromethyl)-4-aminobenzene

To a refluxing solution of 1-*tert*-butyl-2-methoxy-5-(trifluoromethyl)-4-nitrobenzene (473 mg, 1.7 mmol) and ammonium formate (473 mg, 7.3 mmol) in EtOH (10 mL) was added 10% Pd-C (200 mg). The mixture was refluxed for 1 h, cooled and filtered through Celite. The solvent was removed by evaporation to give 1-*tert*-butyl-2-methoxy-5-(trifluoromethyl)-4-aminobenzene (**C-18**) (403 mg, 95 %). ¹H NMR (400 MHz, CDCl₃) δ 7.19 (s, 1H), 6.14 (s, 1H), 4.02 (bs, 2H), 3.74 (s, 3H), 1.24 (s, 9H).


[00492] **Example 12:**



[00493] **C-27; 2-*tert*-Butyl-4-bromo-5-amino-phenol**

To a solution of 2-*tert*-butyl-4-bromo-5-nitrophenol (**C-14-a**) (12 g, 43.8 mmol) in MeOH (90 mL) was added Ni (2.4 g). The reaction mixture was stirred under H₂ (1 atm) for 4 h. The mixture was filtered and the filtrate was concentrated. The crude product was recrystallized from ethyl acetate and petroleum ether to give 2-*tert*-butyl-4-bromo-5-amino-phenol (**C-27**) (7.2 g, 70 %). ¹H NMR (DMSO-*d*₆) δ 9.15 (s, 1 H), 6.91 (s, 1 H), 6.24 (s, 1 H), 4.90 (br s, 2 H), 1.22 (s, 9 H); ESI-MS 244.0 m/z (MH⁺).


[00494] **Example 13:**



[00495] **C-24; 2,4-Di-*tert*-butyl-6-(N-methylamino)phenol**

A mixture of 2,4-di-*tert*-butyl-6-amino-phenol (**C-9**) (5.08 g, 23 mmol), NaBH₃CN (4.41 g, 70 mmol) and paraformaldehyde (2.1 g, 70 mmol) in methanol (50 mL) was stirred at reflux for 3 h. After removal of the solvent, the residue was purified by column chromatography (petroleum ether – EtOAc, 30:1) to give 2,4-di-*tert*-butyl-6-(*N*-methylamino)phenol (**C-24**) (800 mg, 15 %). ¹HNMR (DMSO-*d*₆) δ 8.67 (s, 1 H), 6.84 (s, 1 H), 5.99 (s, 1 H), 4.36 (q, *J* = 4.8 Hz, 1H), 2.65 (d, *J* = 4.8 Hz, 3 H), 1.23 (s, 18 H); ESI-MS 236.2 *m/z* (MH⁺).

[00496] Example 14:



[00497] 2-Methyl-2-phenyl-propan-1-ol

To a solution of 2-methyl-2-phenyl-propionic acid (82 g, 0.5 mol) in THF (200 mL) was added dropwise borane-dimethyl sulfide (2M, 100 mL) at 0-5 °C. The mixture was stirred at this temperature for 30 min and then heated at reflux for 1 h. After cooling, methanol (150 mL) and

water (50 mL) were added. The mixture was extracted with EtOAc (100 mL $\times$ 3), and the combined organic layers were washed with water and brine, dried over Na₂SO₄ and concentrated to give 2-methyl-2-phenyl-propan-1-ol as an oil (70 g, 77 %).

[00498] 2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-benzene

To a suspension of NaH (29 g, 0.75 mol) in THF (200 mL) was added dropwise a solution of 2-methyl-2-phenyl-propan-1-ol (75 g, 0.5 mol) in THF (50 mL) at 0 °C. The mixture was stirred at 20 °C for 30 min and then a solution of 1-bromo-2-methoxy-ethane (104 g, 0.75 mol) in THF (100 mL) was added dropwise at 0 °C. The mixture was stirred at 20 °C overnight, poured into water (200 mL) and extracted with EtOAc (100 mL $\times$ 3). The combined organic layers were washed with water and brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (silica gel, petroleum ether) to give 2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-benzene as an oil (28 g, 27 %).

[00499] 1-[2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-4-nitro-benzene

To a solution of 2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-benzene (52 g, 0.25 mol) in CHCl₃ (200 mL) was added KNO₃ (50.5 g, 0.5 mol) and TMSCl (54 g, 0.5 mol). The mixture was stirred at 20 °C for 30 min and then AlCl₃ (95 g, 0.7 mol) was added. The reaction mixture was stirred at 20 °C for 1 h and poured into ice-water. The organic layer was separated and the aqueous layer was extracted with CHCl₃ (50 mL $\times$ 3). The combined organic layers were washed with water and brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (silica gel, petroleum ether) to obtain 1-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-4-nitro-benzene (6 g, 10 %).

[00500] 4-[2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenylamine

A suspension of 1-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-4-nitro-benzene (8.1 g, 32 mmol) and Raney Ni (1 g) in MeOH (50 mL) was stirred under H₂ (1 atm) at room temperature for 1 h. The catalyst was filtered off and the filtrate was concentrated to obtain 4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenylamine (5.5 g, 77 %).

[00501] 4-[2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenylamine

To a solution of 4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenylamine (5.8 g, 26 mmol) in H₂SO₄ (20 mL) was added KNO₃ (2.63 g, 26 mmol) at 0 °C. After addition was complete, the mixture was stirred at this temperature for 20 min and then poured into ice-water. The mixture was extracted with EtOAc (50 mL × 3). The combined organic layers were washed with water and brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (petroleum ether – EtOAc, 100:1) to give 4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenylamine (5 g, 71 %).

[00502] N-{4-[2-(2-Methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenyl}-acetamide

To a suspension of NaHCO₃ (10 g, 0.1 mol) in dichloromethane (50 mL) was added 4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenylamine (5 g, 30 mmol) and acetyl chloride (3 mL, 20 mmol) at 0-5 °C. The mixture was stirred overnight at 15 °C and then poured into water (200 mL). The organic layer was separated and the aqueous layer was extracted with dichloromethane (50 mL × 2). The combined organic layers were washed with water and brine, dried over Na₂SO₄, and concentrated to dryness to give N-{4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenyl}-acetamide (5.0 g, 87 %).

[00503] N-{3-Amino-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}-acetamide

A mixture of N-{4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-3-nitro-phenyl}-acetamide (5 g, 16 mmol) and Raney Ni (1 g) in MeOH (50 mL) was stirred under H₂ (1 atm) at room temperature 1 h. The catalyst was filtered off and the filtrate was concentrated. The residue was purified by column chromatography (petroleum ether – EtOAc, 100:1) to give N-{3-amino-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}-acetamide (1.6 g, 35 %).

[00504] N-{3-Hydroxy-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}-acetamide


To a solution of N-{3-amino-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}-acetamide (1.6 g, 5.7 mmol) in H₂SO₄ (15 %, 6 mL) was added NaNO₂ at 0-5 °C. The mixture was stirred at this temperature for 20 min and then poured into ice water. The mixture was extracted with

EtOAc (30 mL $\times$ 3). The combined organic layers were washed with water and brine, dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography (petroleum ether – EtOAc, 100:1) to give N-{3-hydroxy-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}- acetamide (0.7 g, 38 %).

[00505] C-25; 2-(1-(2-Methoxyethoxy)-2-methylpropan-2-yl)-5-aminophenol

A mixture of N-{3-hydroxy-4-[2-(2-methoxy-ethoxy)-1,1-dimethyl-ethyl]-phenyl}- acetamide (1 g, 3.5 mmol) and HCl (5 mL) was heated at reflux for 1 h. The mixture was basified with Na_2CO_3 solution to pH 9 and then extracted with EtOAc (20 mL $\times$ 3). The combined organic layers were washed with water and brine, dried over Na_2SO_4 and concentrated to dryness. The residue was purified by column chromatography (petroleum ether – EtOAc, 100:1) to obtain 2-(1-(2-methoxyethoxy)-2-methylpropan-2-yl)-5-aminophenol (**C-25**) (61 mg, 6 %). ^1H NMR (CDCl_3) δ 9.11 (br s, 1 H), 6.96-6.98 (d, $J = 8$ Hz, 1 H), 6.26-6.27 (d, $J = 4$ Hz, 1 H), 6.17-6.19 (m, 1 H), 3.68-3.69 (m, 2 H), 3.56-3.59 (m, 4 H), 3.39 (s, 3 H), 1.37 (s, 6 H); ESI-MS 239.9 m/z (MH^+).

[00506] Example 15:



[00507]

[00508] 4,6-di-tert-Butyl-3-nitrocyclohexa-3,5-diene-1,2-dione

To a solution of 3,5-di-tert-butylcyclohexa-3,5-diene-1,2-dione (4.20 g, 19.1 mmol) in acetic acid (115 mL) was slowly added HNO_3 (15 mL). The mixture was heated at 60 $^\circ\text{C}$ for 40 min before it was poured into H_2O (50 mL). The mixture was allowed to stand at room temperature

for 2 h, then was placed in an ice bath for 1 h. The solid was collected and washed with water to provide 4,6-di-*tert*-butyl-3-nitrocyclohexa-3,5-diene-1,2-dione (1.2 g, 24 %). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 6.89 (s, 1H), 1.27 (s, 9H), 1.24 (s, 9H).

[0001] 4,6-Di-*tert*-butyl-3-nitrobenzene-1,2-diol

In a separatory funnel was placed THF/ H_2O (1:1, 400 mL), 4,6-di-*tert*-butyl-3-nitrocyclohexa-3,5-diene-1,2-dione (4.59 g, 17.3 mmol) and $\text{Na}_2\text{S}_2\text{O}_4$ (3 g, 17.3 mmol). The separatory funnel was stoppered and was shaken for 2 min. The mixture was diluted with EtOAc (20 mL). The layers were separated and the organic layer was washed with brine, dried over MgSO_4 and concentrated to provide 4,6-di-*tert*-butyl-3-nitrobenzene-1,2-diol (3.4 g, 74 %), which was used without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.24 (s, 1H), 8.76 (s, 1H), 6.87 (s, 1H), 1.35 (s, 9H), 1.25 (s, 9H).


[0002] C-26; 4,6-Di-*tert*-butyl-3-aminobenzene-1,2-diol

To a solution of 4,6-di-*tert*-butyl-3-nitrobenzene-1,2-diol (1.92 g, 7.2 mmol) in EtOH (70 mL) was added Pd-5% wt. on carbon (200 mg). The mixture was stirred under H_2 (1 atm) for 2 h. The reaction was recharged with Pd-5% wt. on carbon (200 mg) and stirred under H_2 (1 atm) for another 2 h. The mixture was filtered through Celite and the filtrate was concentrated and purified by column chromatography (10-40 % ethyl acetate – hexanes) to give 4,6-di-*tert*-butyl-3-aminobenzene-1,2-diol (C-26) (560 mg, 33 %). ^1H NMR (400 MHz, CDCl_3) δ 7.28 (s, 1H), 1.42 (s, 9H), 1.38 (s, 9H).


[0003] Anilines

[0004] Example 1:

[0005] General scheme




[00509] Specific example:



[00510] **D-1; 4-Chloro-benzene-1,3-diamine**


A mixture of 1-chloro-2,4-dinitro-benzene (100 mg, 0.5 mmol) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (1.12 g, 5 mmol) in ethanol (2.5 mL) was stirred at room temperature overnight. Water was added and then the mixture was basified to pH 7-8 with saturated NaHCO_3 solution. The solution was extracted with ethyl acetate. The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated to yield 4-chloro-benzene-1,3-diamine (**D-1**) (79 mg, quant.). HPLC ret. time 0.38 min, 10-99 % CH_3CN , 5 min run; ESI-MS 143.1 m/z (MH^+)

[00511] Other examples:




[00512] **D-2; 4,6-Dichloro-benzene-1,3-diamine**

4,6-Dichloro-benzene-1,3-diamine (**D-2**) was synthesized following the general scheme above starting from 1,5-dichloro-2,4-dinitro-benzene. Yield (95 %). HPLC ret. time 1.88 min, 10-99 % CH_3CN , 5 min run; ESI-MS 177.1 m/z (MH^+).




[00513] **D-3; 4-Methoxy-benzene-1,3-diamine**

4-Methoxy-benzene-1,3-diamine (**D-3**) was synthesized following the general scheme above starting from 1-methoxy-2,4-dinitro-benzene. Yield (quant.). HPLC ret. time 0.31 min, 10-99 % CH₃CN, 5 min run.



[00514] D-4; 4-Trifluoromethoxy-benzene-1,3-diamine

4-Trifluoromethoxy-benzene-1,3-diamine (**D-4**) was synthesized following the general scheme above starting from 2,4-dinitro-1-trifluoromethoxy-benzene. Yield (89 %). HPLC ret. time 0.91 min, 10-99 % CH₃CN, 5 min run; ESI-MS 193.3 m/z (MH⁺).




[00515] D-5; 4-Propoxybenzene-1,3-diamine

4-Propoxybenzene-1,3-diamine (**D-5**) was synthesized following the general scheme above starting from 5-nitro-2-propoxy-phenylamine. Yield (79 %). HPLC ret. time 0.54 min, 10-99 % CH₃CN, 5 min run; ESI-MS 167.5 m/z (MH⁺).


[00516] Example 2:

[00517] General scheme



a) HNO₃, H₂SO₄; b) SnCl₂·2H₂O, EtOH or H₂, Pd-C, MeOH

[00518] Specific example:




[00519] 2,4-Dinitro-propylbenzene

A solution of propylbenzene (10 g, 83 mmol) in conc. H_2SO_4 (50 mL) was cooled at 0 °C for 30 min, and a solution of conc. H_2SO_4 (50 mL) and fuming HNO_3 (25 mL), previously cooled to 0 °C, was added in portions over 15 min. The mixture was stirred at 0 °C for additional 30 min, and then allowed to warm to room temperature. The mixture was poured into ice (200 g) - water (100 mL) and extracted with ether (2 x 100 mL). The combined extracts were washed with H_2O (100 mL) and brine (100 mL), dried over MgSO_4 , filtered and concentrated to afford 2,4-dinitro-propylbenzene (15.6 g, 89 %). ^1H NMR (CDCl_3 , 300 MHz) δ 8.73 (d, $J = 2.2$ Hz, 1H), 8.38 (dd, $J = 8.3$, $J = 2.2$, 1H), 7.6 (d, $J = 8.5$ Hz, 1H), 2.96 (dd, 2H), 1.73 (m, 2H), 1.06 (t, $J = 7.4$ Hz, 3H).

[00520] D-6; 4-Propyl-benzene-1,3-diamine


To a solution of 2,4-dinitro-propylbenzene (2.02 g, 9.6 mmol) in ethanol (100 mL) was added SnCl_2 (9.9 g, 52 mmol) followed by conc. HCl (10 mL). The mixture was refluxed for 2 h, poured into ice-water (100 mL), and neutralized with solid sodium bicarbonate. The solution was further basified with 10% NaOH solution to pH ~ 10 and extracted with ether (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over MgSO_4 , filtered, and concentrated to provide 4-propyl-benzene-1,3-diamine (**D-6**) (1.2 g, 83 %). No further purification was necessary for use in the next step; however, the product was not stable for an extended period of time. ^1H NMR (CDCl_3 , 300 MHz) δ 6.82 (d, $J = 7.9$ Hz, 1H), 6.11 (dd, $J = 7.5$, $J = 2.2$ Hz, 1H), 6.06 (d, $J = 2.2$ Hz, 1H), 3.49 (br s, 4H, NH_2), 2.38 (t, $J = 7.4$ Hz, 2H), 1.58 (m, 2H), 0.98 (t, $J = 7.2$ Hz, 3H); ESI-MS 151.5 m/z (MH^+).

[00521] Other examples:




[00522] D-7; 4-Ethylbenzene-1,3-diamine

4-Ethylbenzene-1,3-diamine (**D-7**) was synthesized following the general scheme above starting from ethylbenzene. Overall yield (76 %).



[00523] D-8; 4-Isopropylbenzene-1,3-diamine

4-Isopropylbenzene-1,3-diamine (**D-8**) was synthesized following the general scheme above starting from isopropylbenzene. Overall yield (78 %).




[00524] D-9; 4-*tert*-Butylbenzene-1,3-diamine

4-*tert*-Butylbenzene-1,3-diamine (**D-9**) was synthesized following the general scheme above starting from *tert*-butylbenzene. Overall yield (48 %). ^1H NMR (400 MHz, CDCl_3) δ 7.01 (d, J = 8.3 Hz, 1H), 6.10 (dd, J = 2.4, 8.3 Hz, 1H), 6.01 (d, J = 2.4 Hz, 1H), 3.59 (br, 4H), 1.37 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 145.5, 145.3, 127.6, 124.9, 105.9, 104.5, 33.6, 30.1; ESI-MS 164.9 m/z (MH^+).


[00525] Example 3:

[00526] General scheme



a) KNO_3 , H_2SO_4 ; b) (i) HNO_3 , H_2SO_4 ; (ii) Na_2S , S , H_2O ; c) Boc_2O , NaOH , THF ; d) H_2 , Pd-C , MeOH

[00527] Specific example:



[00528] **4-*tert*-Butyl-3-nitro-phenylamine**


To a mixture of 4-*tert*-butyl-phenylamine (10.0 g, 67.01 mmol) dissolved in H_2SO_4 (98 %, 60 mL) was slowly added KNO_3 (8.1 g, 80.41 mmol) at 0 °C. After addition, the reaction was allowed to warm to room temperature and stirred overnight. The mixture was then poured into ice-water and basified with sat. NaHCO_3 solution to pH 8. The mixture was extracted several times with CH_2Cl_2 . The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography (petroleum ether - EtOAc , 10:1) to give 4-*tert*-butyl-3-nitro-phenylamine (10 g, 77 %).

[00529] **(4-*tert*-Butyl-3-nitro-phenyl)-carbamic acid *tert*-butyl ester**


A mixture of 4-*tert*-butyl-3-nitro-phenylamine (4.0 g, 20.6 mmol) and Boc_2O (4.72 g, 21.6 mmol) in NaOH (2N, 20 mL) and THF (20 mL) was stirred at room temperature overnight. THF was removed under reduced pressure. The residue was dissolved in water and extracted with CH_2Cl_2 . The organic layer was washed with NaHCO_3 and brine, dried over Na_2SO_4 and concentrated to afford (4-*tert*-butyl-3-nitro-phenyl)-carbamic acid *tert*-butyl ester (4.5 g, 74 %).

[00530] D-10; (3-Amino-4-*tert*-butyl-phenyl)-carbamic acid *tert*-butyl ester


A suspension of (4-*tert*-butyl-3-nitro-phenyl)-carbamic acid *tert*-butyl ester (3.0 g, 10.19 mol) and 10% Pd-C (1 g) in MeOH (40 mL) was stirred under H₂ (1 atm) at room temperature overnight. After filtration, the filtrate was concentrated and the residue was purified by column chromatograph (petroleum ether - EtOAc, 5:1) to give (3-amino-4-*tert*-butyl-phenyl)-carbamic acid *tert*-butyl ester (**D-10**) as a brown oil (2.5 g, 93 %). ¹H NMR (CDCl₃) δ 7.10 (d, *J* = 8.4 Hz, 1 H), 6.92 (s, 1 H), 6.50-6.53 (m, 1 H), 6.36 (s, 1 H), 3.62 (br s, 2 H), 1.50 (s, 9 H), 1.38 (s, 9 H); ESI-MS 528.9 m/z (2M+H⁺).

[00531] Other examples:**[00532] D-11; (3-Amino-4-isopropyl-phenyl)-carbamic acid *tert*-butyl ester**

(3-Amino-4-isopropyl-phenyl)-carbamic acid *tert*-butyl ester (**D-11**) was synthesized following the general scheme above starting from isopropylbenzene. Overall yield (56 %).

**[00533] D-12; (3-Amino-4-ethyl-phenyl)-carbamic acid *tert*-butyl ester**


(3-Amino-4-ethyl-phenyl)-carbamic acid *tert*-butyl ester (**D-12**) was synthesized following the general scheme above starting from ethylbenzene. Overall yield (64 %). ¹H NMR (CD₃OD, 300 MHz) δ 6.87 (d, *J* = 8.0 Hz, 1H), 6.81 (d, *J* = 2.2 Hz, 1H), 6.63 (dd, *J* = 8.1, *J* = 2.2, 1H), 2.47 (q, *J* = 7.4 Hz, 2H), 1.50 (s, 9H), 1.19 (t, *J* = 7.4 Hz, 3H); ESI-MS 237.1 m/z (MH⁺).



[00534] D-13; (3-Amino-4-propyl-phenyl)-carbamic acid *tert*-butyl ester

(3-Amino-4-propyl-phenyl)-carbamic acid *tert*-butyl ester (**D-13**) was synthesized following the general scheme above starting from propylbenzene. Overall yield (48 %).

[00535] Example 4:



[00536] (3-Amino-4-*tert*-butyl-phenyl)-carbamic acid benzyl ester

A solution of 4-*tert*-butylbenzene-1,3-diamine (**D-9**) (657 mg, 4 mmol) and pyridine (0.39 mL, 4.8 mmol) in CH₂Cl₂ / MeOH (12 / 1, 8 mL) was cooled to 0 °C, and a solution of benzyl chloroformate (0.51 mL, 3.6 mmol) in CH₂Cl₂ (8 mL) was added dropwise over 10 min. The mixture was stirred at 0 °C for 15 min, then warmed to room temperature. After 1 h, the mixture was washed with 1M citric acid (2 x 20 mL), saturated aqueous sodium bicarbonate (20 mL), dried (Na₂SO₄), filtered and concentrated in vacuo to afford the crude (3-amino-4-*tert*-butyl-phenyl)-carbamic acid benzyl ester as a brown viscous gum (0.97 g), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.41-7.32 (m, 6H), 7.12 (d, *J* = 8.5 Hz, 1H), 6.89 (br s, 1H), 6.57 (dd, *J* = 2.3, 8.5 Hz, 1H), 5.17 (s, 2H), 3.85 (br s, 2H), 1.38 (s, 9H); ¹³C NMR (100 MHz, CDCl₃, rotameric) δ 153.3 (br), 145.3, 136.56, 136.18, 129.2, 128.73, 128.59, 128.29, 128.25, 127.14, 108.63 (br), 107.61 (br), 66.86, 33.9, 29.7; ESI-MS 299.1 *m/z* (MH⁺).

[00537] (4-*tert*-Butyl-3-formylamino-phenyl)-carbamic acid benzyl ester

A solution of (3-amino-4-*tert*-butyl-phenyl)-carbamic acid benzyl ester (0.97 g, 3.25 mmol) and pyridine (0.43 mL, 5.25 mmol) in CH₂Cl₂ (7.5 mL) was cooled to 0 °C, and a solution of formic-acetic anhydride (3.5 mmol, prepared by mixing formic acid (158 µL, 4.2 mmol, 1.3 equiv) and acetic anhydride (0.32 mL, 3.5 mmol, 1.1 eq.) neat and ageing for 1 hour) in CH₂Cl₂ (2.5 mL) was added dropwise over 2 min. After the addition was complete, the mixture was allowed to warm to room temperature, whereupon it deposited a precipitate, and the resulting slurry was stirred overnight. The mixture was washed with 1 M citric acid (2 x 20 mL), saturated aqueous sodium bicarbonate (20 mL), dried (Na₂SO₄), and filtered. The cloudy mixture deposited a thin bed of solid above the drying agent, HPLC analysis showed this to be the desired formamide. The filtrate was concentrated to approximately 5 mL, and diluted with hexane (15 mL) to precipitate further formamide. The drying agent (Na₂SO₄) was slurried with methanol (50 mL), filtered, and the filtrate combined with material from the CH₂Cl₂ / hexane recrystallisation. The resultant mixture was concentrated to afford (4-*tert*-butyl-3-formylamino-phenyl)-carbamic acid benzyl ester as an off-white solid (650 mg, 50 % over 2 steps). ¹H and ¹³C NMR (CD₃OD) show the product as a rotameric mixture. ¹H NMR (400 MHz, CD₃OD, rotameric) δ 8.27 (s, 1H-a), 8.17 (s, 1H-b), 7.42-7.26 (m, 8H), 5.17 (s, 1H-a), 5.15 (s, 1H-b), 4.86 (s, 2H), 1.37 (s, 9H-a), 1.36 (s, 9H-b); ¹³C NMR (100 MHz, CD₃OD, rotameric) δ 1636.9, 163.5, 155.8, 141.40, 141.32, 139.37, 138.88, 138.22, 138.14, 136.4, 135.3, 129.68, 129.65, 129.31, 129.24, 129.19, 129.13, 128.94, 128.50, 121.4 (br), 118.7 (br), 67.80, 67.67, 35.78, 35.52, 31.65, 31.34; ESI-MS 327.5 m/z (MH⁺).

[00538] *N*-(5-Amino-2-*tert*-butyl-phenyl)-formamide

A 100 mL flask was charged with (4-*tert*-butyl-3-formylamino-phenyl)-carbamic acid benzyl ester (650 mg, 1.99 mmol), methanol (30 mL) and 10% Pd-C (50 mg), and stirred under H₂ (1 atm) for 20 h. CH₂Cl₂ (5 mL) was added to quench the catalyst, and the mixture then filtered through Celite, and concentrated to afford *N*-(5-amino-2-*tert*-butyl-phenyl)-formamide as an off-white solid (366 mg, 96 %). Rotameric by ¹H and ¹³C NMR (DMSO-*d*₆). ¹H NMR (400 MHz, DMSO-*d*₆, rotameric) δ 9.24 (d, *J* = 10.4 Hz, 1H), 9.15 (s, 1H), 8.23 (d, *J* = 1.5 Hz, 1H), 8.06 (d, *J* = 10.4 Hz, 1H), 7.06 (d, *J* = 8.5 Hz, 1H), 7.02 (d, *J* = 8.5 Hz, 1H), 6.51 (d, *J* = 2.5 Hz, 1H), 6.46 (dd, *J* = 2.5, 8.5 Hz, 1H), 6.39 (dd, *J* = 2.5, 8.5 Hz, 1H), 6.29 (d, *J* = 2.5 Hz, 1H), 5.05 (s,


2H), 4.93 (s, 2H), 1.27 (s, 9H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$, rotameric) δ 164.0, 160.4, 147.37, 146.74, 135.38, 135.72, 132.48, 131.59, 127.31, 126.69, 115.15, 115.01, 112.43, 112.00, 33.92, 33.57, 31.33, 30.92; ESI-MS 193.1 m/z (MH^+).

[00539] D-14; 4-*tert*-butyl-*N*³-methyl-benzene-1,3-diamine


A 100 mL flask was charged with *N*-(5-amino-2-*tert*-butyl-phenyl)-formamide (340 mg, 1.77 mmol) and purged with nitrogen. THF (10 mL) was added, and the solution was cooled to 0 °C. A solution of lithium aluminum hydride in THF (4.4 mL, 1M solution) was added over 2 min. The mixture was then allowed to warm to room temperature. After refluxing for 15 h, the yellow suspension was cooled to 0 °C, quenched with water (170 μL), 15 % aqueous NaOH (170 μL), and water (510 μL) which were added sequentially and stirred at room temperature for 30 min. The mixture was filtered through Celite, and the filter cake washed with methanol (50 mL). The combined filtrates were concentrated in *vacuo* to give a gray-brown solid, which was partitioned between chloroform (75 mL) and water (50 mL). The organic layer was separated, washed with water (50 mL), dried (Na_2SO_4), filtered, and concentrated to afford 4-*tert*-butyl-*N*³-methyl-benzene-1,3-diamine (**D-14**) as a brown oil which solidified on standing (313 mg, 98 %). ^1H NMR (400 MHz, CDCl_3) δ 7.01 (d, J = 8.1 Hz, 1H), 6.05 (dd, J = 2.4, 8.1 Hz, 1H), 6.03 (d, J = 2.4 Hz, 1H), 3.91 (br s, 1H), 3.52 (br s, 2H), 2.86 (s, 3H), 1.36 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 148.4, 145.7, 127.0, 124.3, 103.6, 98.9, 33.5, 31.15, 30.31; ESI-MS 179.1 m/z (MH^+).

[00540] Example 5:

[00541] General scheme:



[00542] Specific example:



[00543] 2,4-Dinitro-propylbenzene

A solution of propylbenzene (10 g, 83 mmol) in conc. H_2SO_4 (50 mL) was cooled at 0 °C for 30 mins, and a solution of conc. H_2SO_4 (50 mL) and fuming HNO_3 (25 mL), previously cooled to 0 °C, was added in portions over 15 min. The mixture was stirred at 0 °C for additional 30 min. and then allowed to warm to room temperature. The mixture was poured into ice (200 g) -water (100 mL) and extracted with ether (2 x 100 mL). The combined extracts were washed with H_2O (100 mL) and brine (100 mL), dried over MgSO_4 , filtered and concentrated to afford 2,4-dinitro-propylbenzene (15.6 g, 89 %). ^1H NMR (CDCl_3 , 300 MHz) δ 8.73 (d, $J = 2.2$ Hz, 1H), 8.38 (dd, $J = 8.3, 2.2$ Hz, 1H), 7.6 (d, $J = 8.5$ Hz, 1H), 2.96 (m, 2H), 1.73 (m, 2H), 1.06 (t, $J = 7.4$ Hz, 3H).

[00544] 4-Propyl-3-nitroaniline

A suspension of 2,4-dinitro-propylbenzene (2 g, 9.5 mmol) in H_2O (100 mL) was heated near reflux and stirred vigorously. A clear orange-red solution of polysulfide (300 mL (10 eq.)), previously prepared by heating sodium sulfide nanohydrate (10.0 g), sulfur powder (2.60 g) and H_2O (400 mL), was added dropwise over 45 mins. The red-brown solution was heated at reflux for 1.5 h. The mixture was cooled to 0 °C and then extracted with ether (2 x 200 mL). The combined organic extracts were dried over MgSO_4 , filtered, and concentrated under reduced pressure to afford 4-propyl-3-nitroaniline (1.6 g, 93 %), which was used without further purification.

[00545] (3-Nitro-4-propyl-phenyl)-carbamic acid tert-butyl ester

4-Propyl-3-nitroaniline (1.69 g, 9.4 mmol) was dissolved in pyridine (30 mL) with stirring. Boc anhydride (2.05 g, 9.4 mmol) was added. The mixture was stirred and heated at reflux for 1 h before the solvent was removed *in vacuo*. The oil obtained was re-dissolved in CH₂Cl₂ (300 mL) and washed with water (300 mL) and brine (300 mL), dried over Na₂SO₄, filtered, and concentrated. The crude oil that contained both mono- and *bis*-acylated nitro products was purified by column chromatography (0–10 % CH₂Cl₂ - MeOH) to afford (3-nitro-4-propyl-phenyl)-carbamic acid *tert*-butyl ester (2.3 g, 87 %).


[00546] Methyl-(3-nitro-4-propyl-phenyl)-carbamic acid *tert*-butyl ester

To a solution of (3-nitro-4-propyl-phenyl)-carbamic acid *tert*-butyl ester (200 mg, 0.71 mmol) in DMF (5 mL) was added Ag₂O (1.0 g, 6.0 mmol) followed by methyl iodide (0.20 mL, 3.2 mmol). The resulting suspension was stirred at room temperature for 18 h and filtered through a pad of Celite. The filter cake was washed with CH₂Cl₂ (10 mL). The filtrate was concentrated *in vacuo*. The crude oil was purified by column chromatography (0–10 % CH₂Cl₂ - MeOH) to afford methyl-(3-nitro-4-propyl-phenyl)-carbamic acid *tert*-butyl ester as a yellow oil (110 mg, 52 %). ¹H NMR (CDCl₃, 300 MHz) δ 7.78 (d, *J* = 2.2 Hz, 1H), 7.42 (dd, *J* = 8.2, 2.2 Hz, 1H), 7.26 (d, *J* = 8.2 Hz, 1H), 3.27 (s, 3H), 2.81 (t, *J* = 7.7 Hz, 2H), 1.66 (m, 2H), 1.61 (s, 9H), 0.97 (t, *J* = 7.4 Hz, 3H).


[00547] D-15; (3-Amino-4-propyl-phenyl)-methyl-carbamic acid *tert*-butyl ester

To a solution of methyl-(3-nitro-4-propyl-phenyl)-carbamic acid *tert*-butyl ester (110 mg, 0.37 mmol) in EtOAc (10 mL) was added 10% Pd-C (100 mg). The resulting suspension was stirred at room temperature under H₂ (1 atm) for 2 days. The progress of the reaction was monitored by TLC. Upon completion, the reaction mixture was filtered through a pad of Celite. The filtrate was concentrated *in vacuo* to afford (3-Amino-4-propyl-phenyl)-methyl-carbamic acid *tert*-butyl ester (**D-15**) as a colorless crystalline compound (80 mg, 81 %). ESI-MS 265.3 m/z (MH⁺).

[00548] Other examples:




[00549] **D-16; (3-Amino-4-ethyl-phenyl)-methyl-carbamic acid *tert*-butyl ester**
 (3-Amino-4-ethyl-phenyl)-methyl-carbamic acid *tert*-butyl ester (**D-16**) was synthesized following the general scheme above starting from ethylbenzene. Overall yield (57 %).



[00550] **D-17; (3-Amino-4-isopropyl-phenyl)-methyl-carbamic acid *tert*-butyl ester**
 (3-Amino-4-isopropyl-phenyl)-methyl-carbamic acid *tert*-butyl ester (**D-17**) was synthesized following the general scheme above starting from isopropylbenzene. Overall yield (38 %).

[00551] **Example 6:**



[00552] **2'-Ethoxy-2,4-dinitro-biphenyl**

A pressure flask was charged with 2-ethoxyphenylboronic acid (0.66 g, 4.0 mmol), KF (0.77 g, 13 mmol), Pd₂(dba)₃ (16 mg, 0.02 mmol), and 2,4-dinitro-bromobenzene (0.99 g, 4.0 mmol) in THF (5 mL). The vessel was purged with argon for 1 min followed by the addition of tri-*tert*-butylphosphine (0.15 ml, 0.48 mmol, 10 % solution in hexanes). The reaction vessel was purged with argon for additional 1 min., sealed and heated at 80 °C overnight. After cooling to room temperature, the solution was filtered through a plug of Celite. The filter cake was rinsed with CH₂Cl₂ (10 mL), and the combined organic extracts were concentrated under reduced pressure to provide the crude product 2'-ethoxy-2,4-dinitro-biphenyl (0.95 g, 82%). No further purification was performed. ¹H NMR (300 MHz, CDCl₃) δ 8.75 (s, 1H), 8.43 (d, *J* = 8.7 Hz, 1H), 7.60 (d, *J* = 8.4 Hz, 1H), 7.40 (t, *J* = 7.8 Hz, 1H), 7.31 (d, *J* = 7.5 Hz, 1H), 7.08 (t, *J* = 7.5 Hz, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), 3.44 (q, *J* = 6.6 Hz, 2H), 1.24 (t, *J* = 6.6 Hz, 3H); HPLC ret. time 3.14 min, 10-100 % CH₃CN, 5 min gradient.

[00553] 2'-Ethoxy-2-nitrobiphenyl-4-yl amine

A clear orange-red solution of polysulfide (120 ml, 7.5 eq.), previously prepared by heating sodium sulfide monohydrate (10 g), sulfur (1.04 g) and water (160 ml), was added dropwise at 90 °C over 45 minutes to a suspension of 2'-ethoxy-2,4-dinitro-biphenyl (1.2 g, 4.0 mmol) in water (40 ml). The red-brown solution was heated at reflux for 1.5 h. The mixture was cooled to room temperature, and solid NaCl (5 g) was added. The solution was extracted with CH₂Cl₂ (3 x 50 mL), and the combined organic extracts was concentrated to provide 2'-ethoxy-2-nitrobiphenyl-4-yl amine (0.98 g, 95 %) that was used in the next step without further purification. ¹H NMR (300 MHz, CDCl₃) δ 7.26 (m, 2H), 7.17 (d, *J* = 2.7 Hz, 1H), 7.11 (d, *J* = 7.8 Hz, 1H), 7.00 (t, *J* = 6.9 Hz, 1H), 6.83 (m, 2H), 3.91 (q, *J* = 6.9 Hz, 2H), 1.23 (t, *J* = 7.2 Hz, 3H); HPLC ret. time 2.81 min, 10-100 % CH₃CN, 5 min gradient; ESI-MS 259.1 m/z (MH⁺).

[00554] (2'-Ethoxy-2-nitrobiphenyl-4-yl)-carbamic acid *tert*-butyl ester


A mixture of 2'-ethoxy-2-nitrobiphenyl-4-yl amine (0.98 g, 4.0 mmol) and Boc₂O (2.6g, 12 mmol) was heated with a heat gun. Upon the consumption of the starting material as indicated by TLC, the crude mixture was purified by flash chromatography (silica gel, CH₂Cl₂) to provide (2'-ethoxy-2-nitrobiphenyl-4-yl)-carbamic acid *tert*-butyl ester (1.5 g, 83 %). ¹H NMR (300 MHz, CDCl₃) δ 7.99 (s, 1H), 7.55 (d, *J* = 8.4 Hz, 1H), 7.25 (m, 3H), 6.99 (t, *J* = 7.5 Hz, 1H),

6.82 (m, 2H), 3.88 (q, $J = 6.9$ Hz, 2H), 1.50 (s, 9 H), 1.18 (t, $J = 6.9$ Hz, 3H); HPLC ret. time 3.30 min, 10-100 % CH₃CN, 5 min gradient.

[00555] D-18; (2'-ethoxy-2-aminobiphenyl-4-yl)-carbamic acid *tert*-butyl ester

To a solution of NiCl₂·6H₂O (0.26 g, 1.1 mmol) in EtOH (5 mL) was added NaBH₄ (40 mg, 1.1 mmol) at -10 °C. Gas evolution was observed and a black precipitate was formed. After stirring for 5 min, a solution of 2'-ethoxy-2-nitrobiphenyl-4-yl)carbamic acid *tert*-butyl ester (0.50 g, 1.1 mmol) in EtOH (2 mL) was added. Additional NaBH₄ (80 mg, 60 mmol) was added in 3 portions over 20 min. The reaction was stirred at 0 °C for 20 min followed by the addition of NH₄OH (4 mL, 25% aq. solution). The resulting solution was stirred for 20 min. The crude mixture was filtered through a short plug of silica. The silica cake was flushed with 5% MeOH in CH₂Cl₂ (10 mL), and the combined organic extracts was concentrated under reduced pressure to provide (2'-ethoxy-2-aminobiphenyl-4-yl)-carbamic acid *tert*-butyl ester (**D-18**) (0.36 g, quant.), which was used without further purification. HPLC ret. time 2.41 min, 10-100 % CH₃CN, 5 min gradient; ESI-MS 329.3 m/z (MH⁺).

[00556] Example 7:




[00557] D-19; N-(3-Amino-5-trifluoromethyl-phenyl)-methanesulfonamide

A solution of 5-trifluoromethyl-benzene-1,3-diamine (250 mg, 1.42 mmol) in pyridine (0.52 mL) and CH₂Cl₂ (6.5 mL) was cooled to 0 °C. Methanesulfonyl chloride (171 mg, 1.49 mmol) was slowly added at such a rate that the temperature of the solution remained below 10 °C. The mixture was stirred at ~ 8 °C and then allowed to warm to room temperature after 30 min. After stirring at room temperature for 4 h, reaction was almost complete as indicated by LCMS analysis. The reaction mixture was quenched with sat. aq. NH₄Cl (10 mL) solution, extracted

with CH_2Cl_2 (4 x 10 mL), dried over Na_2SO_4 , filtered, and concentrated to yield N-(3-amino-5-trifluoromethyl-phenyl)-methanesulfonamide (**D-19**) as a reddish semisolid (0.35 g, 97 %), which was used without further purification. $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 6.76 (m, 1H), 6.70 (m, 1H), 6.66 (s, 1H), 3.02 (s, 3H); ESI-MS 255.3 m/z (MH^+).

[00558] Cyclic amines

[00559] Example 1:



[00560] 7-Nitro-1,2,3,4-tetrahydro-quinoline

To a mixture of 1,2,3,4-tetrahydro-quinoline (20.0 g, 0.15 mol) dissolved in H_2SO_4 (98 %, 150 mL), KNO_3 (18.2 g, 0.18 mol) was slowly added at 0 °C. The reaction was allowed to warm to room temperature and stirred over night. The mixture was then poured into ice-water and basified with sat. NaHCO_3 solution to pH 8. After extraction with CH_2Cl_2 , the combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography (petroleum ether – EtOAc, 10:1) to give 7-nitro-1,2,3,4-tetrahydro-quinoline (6.6 g, 25 %).

[00561] 7-Nitro-3,4-dihydro-2H-quinoline-1-carboxylic acid *tert*-butyl ester


A mixture of 7-nitro-1,2,3,4-tetrahydro-quinoline (4.0 g, 5.61 mmol), Boc_2O (1.29 g, 5.89 mmol) and DMAP (0.4 g) in CH_2Cl_2 was stirred at room temperature overnight. After diluted with water, the mixture was extracted with CH_2Cl_2 . The combined organic layers were washed with NaHCO_3 and brine, dried over Na_2SO_4 and concentrated to provide crude 7-nitro-3,4-dihydro-

2H-quinoline-1-carboxylic acid *tert*-butyl ester that was used in the next step without further purification.

[00562] DC-1; *tert*-Butyl 7-amino-3,4-dihydroquinoline-1(2H)-carboxylate

A suspension of the crude 7-nitro-3,4-dihydro-2H-quinoline-1-carboxylic acid *tert*-butyl ester (4.5 g, 16.2 mol) and 10% Pd-C (0.45 g) in MeOH (40 mL) was stirred under H₂ (1 atm) at room temperature overnight. After filtration, the filtrate was concentrated and the residue was purified by column chromatography (petroleum ether – EtOAc, 5:1) to give *tert*-butyl 7-amino-3,4-dihydroquinoline-1(2H)-carboxylate (**DC-1**) as a brown solid (1.2 g, 22 % over 2 steps). ¹H NMR (CDCl₃) δ 7.15 (d, *J* = 2 Hz, 1 H), 6.84 (d, *J* = 8 Hz, 1 H), 6.36-6.38 (m, 1 H), 3.65-3.68 (m, 2 H), 3.10 (br s, 2 H), 2.66 (t, *J* = 6.4 Hz, 2 H), 1.84-1.90 (m, 2 H), 1.52 (s, 9 H); ESI-MS 496.8 m/z (2M+H⁺).

[00563] Example 2:



[00564] 3-(2-Hydroxy-ethyl)-1,3-dihydro-indol-2-one

A stirring mixture of oxindole (5.7 g, 43 mmol) and Raney nickel (10 g) in ethane-1,2-diol (100 mL) was heated in an autoclave. After the reaction was complete, the mixture was filtered and

the excess of diol was removed under *vacuum*. The residual oil was triturated with hexane to give 3-(2-hydroxy-ethyl)-1,3-dihydro-indol-2-one as a colorless crystalline solid (4.6 g, 70 %).

[00565] 1,2-Dihydro-3-spiro-1'-cyclopropyl-1*H*-indole-2-one

To a solution of 3-(2-hydroxy-ethyl)-1,3-dihydro-indol-2-one (4.6 g, 26 mmol) and triethylamine (10 mL) in CH₂Cl₂ (100 mL) was added MsCl (3.4 g, 30 mmol) dropwise at -20 °C. The mixture was then allowed to warm up to room temperature and stirred overnight. The mixture was filtered and the filtrate was concentrated under *vacuum*. The residue was purified by column chromatography to give crude 1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole-2-one as a yellow solid (2.5 g), which was used directly in the next step.

[00566] 1,2-Dihydro-3-spiro-1'-cyclopropyl-1*H*-indole

To a solution of 1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole-2-one (2.5 g crude) in THF (50 mL) was added LiAlH₄ (2 g, 52 mmol) portionwise. After heating the mixture to reflux, it was poured into crushed ice, basified with aqueous ammonia to pH 8 and extracted with EtOAc. The combined organic layers were washed with brine, dried over Na₂SO₄ and concentrated to give the crude 1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole as a yellow solid (about 2 g), which was used directly in the next step.

[00567] 6-Nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole

To a cooled solution (-5 °C to -10 °C) of NaNO₃ (1.3 g, 15.3 mmol) in H₂SO₄ (98 %, 30 mL) was added 1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole (2 g, crude) dropwise over a period of 20 min. After addition, the reaction mixture was stirred for another 40 min and poured over crushed ice (20 g). The cooled mixture was then basified with NH₄OH and extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure to yield 6-nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole as a dark gray solid (1.3 g)

[00568] 1-Acetyl-6-nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole


NaHCO₃ (5 g) was suspended in a solution of 6-nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole (1.3 g, crude) in CH₂Cl₂ (50 mL). While stirring vigorously, acetyl chloride (720 mg) was

added dropwise. The mixture was stirred for 1 h and filtered. The filtrate was concentrated under *vacuum*. The residue was purified by flash column chromatography on silica gel to give 1-acetyl-6-nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole (0.9 g, 15 % over 4 steps).

[00569] DC-2; 1-Acetyl-6-amino-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole

A mixture of 1-acetyl-6-nitro-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole (383 mg, 2 mmol) and Pd-C (10 %, 100 mg) in EtOH (50 mL) was stirred at room temperature under H₂ (1 atm) for 1.5 h. The catalyst was filtered off and the filtrate was concentrated under reduced pressure. The residue was treated with HCl / MeOH to give 1-acetyl-6-amino-1,2-dihydro-3-spiro-1'-cyclopropyl-1*H*-indole (**DC-2**) (300 mg, 90 %) as a hydrochloride salt.

[00570] Example 3:



[00571] 3-Methyl-but-2-enoic acid phenylamide

A mixture of 3-methyl-but-2-enoic acid (100 g, 1 mol) and SOCl₂ (119 g, 1 mol) was heated at reflux for 3 h. The excess SOCl₂ was removed under reduced pressure. CH₂Cl₂ (200 mL) was added followed by the addition of aniline (93 g, 1.0 mol) in Et₃N (101 g, 1 mol) at 0 °C. The mixture was stirred at room temperature for 1 h and quenched with HCl (5%, 150 mL). The aqueous layer was separated and extracted with CH₂Cl₂. The combined organic layers were

washed with water (2x100 mL) and brine (100 mL), dried over Na₂SO₄ and concentrated to give 3-methyl-but-2-enoic acid phenylamide (120 g, 80 %).

[00572] 4,4-Dimethyl-3,4-dihydro-1H-quinolin-2-one

AlCl₃ (500 g, 3.8 mol) was carefully added to a suspension of 3-methyl-but-2-enoic acid phenylamide (105 g, 0.6 mol) in benzene (1000 mL). The reaction mixture was stirred at 80 °C overnight and poured into ice-water. The organic layer was separated and the aqueous layer was extracted with ethyl acetate (250 mL x 3). The combined organic layers were washed with water (200 mL x 2) and brine (200 mL), dried over Na₂SO₄ and concentrated to give 4,4-dimethyl-3,4-dihydro-1H-quinolin-2-one (90 g, 86 %).

[00573] 4,4-Dimethyl-1,2,3,4-tetrahydro-quinoline

A solution of 4,4-dimethyl-3,4-dihydro-1H-quinolin-2-one (35 g, 0.2 mol) in THF (100 mL) was added dropwise to a suspension of LiAlH₄ (18 g, 0.47 mol) in THF (200 mL) at 0 °C. After addition, the mixture was stirred at room temperature for 30 min and then slowly heated to reflux for 1 h. The mixture was then cooled to 0 °C. Water (18 mL) and NaOH solution (10 %, 100 mL) were carefully added to quench the reaction. The solid was filtered off and the filtrate was concentrated to give 4,4-dimethyl-1,2,3,4-tetrahydro-quinoline.

[00574] 4,4-Dimethyl-7-nitro-1,2,3,4-tetrahydro-quinoline

To a mixture of 4,4-dimethyl-1,2,3,4-tetrahydro-quinoline (33 g, 0.2 mol) in H₂SO₄ (120 mL) was slowly added KNO₃ (20.7 g, 0.2 mol) at 0 °C. After addition, the mixture was stirred at room temperature for 2 h, carefully poured into ice water and basified with Na₂CO₃ to pH 8. The mixture was extracted with ethyl acetate (3 x 200 mL). The combined extracts were washed with water and brine, dried over Na₂SO₄ and concentrated to give 4, 4-dimethyl-7-nitro-1, 2, 3, 4-tetrahydro-quinoline (21 g, 50 %).

[00575] 4,4-Dimethyl-7-nitro-3,4-dihydro-2H-quinoline-1-carboxylic acid tert-butyl ester


A mixture of 4,4-dimethyl-7-nitro-1,2,3,4-tetrahydro-quinoline (25 g, 0.12 mol) and Boc₂O (55 g, 0.25 mol) was stirred at 80 °C for 2 days. The mixture was purified by silica gel

chromatography to give 4,4-dimethyl-7-nitro-3,4-dihydro-2H-quinoline-1-carboxylic acid *tert*-butyl ester (8 g, 22 %).

[00576] DC-3; *tert*-Butyl 7-amino-3,4-dihydro-4,4-dimethylquinoline-1(2H)-carboxylate

A mixture of 4,4-dimethyl-7-nitro-3,4-dihydro-2H-quinoline-1 carboxylic acid *tert*-butyl ester (8.3 g, 0.03 mol) and Pd-C (0.5 g) in methanol (100 mL) was stirred under H₂ (1 atm) at room temperature overnight. The catalyst was filtered off and the filtrate was concentrated. The residue was washed with petroleum ether to give *tert*-butyl 7-amino-3,4-dihydro-4,4-dimethylquinoline-1(2H)-carboxylate (**DC-3**) (7.2 g, 95 %). ¹H NMR (CDCl₃) δ 7.11-7.04 (m, 2 H), 6.45-6.38 (m, 1 H), 3.71-3.67 (m, 2 H), 3.50-3.28 (m, 2 H), 1.71-1.67 (m, 2 H), 1.51 (s, 9 H), 1.24 (s, 6 H).

[00577] Example 4:



[00578] 1-Chloro-4-methylpentan-3-one

Ethylene was passed through a solution of isobutyryl chloride (50 g, 0.5 mol) and AlCl₃ (68.8 g, 0.52 mol) in anhydrous CH₂Cl₂ (700 mL) at 5 °C. After 4 h, the absorption of ethylene ceased, and the mixture was stirred at room temperature overnight. The mixture was poured into cold diluted HCl solution and extracted with CH₂Cl₂. The combined organic phases were washed

with brine, dried over Na_2SO_4 , filtered and concentrated to give the crude 1-chloro-4-methylpentan-3-one, which was used directly in the next step without further purification.

[00579] 4-Methyl-1-(phenylamino)-pentan-3-one

A suspension of the crude 1-chloro-4-methylpentan-3-one (about 60 g), aniline (69.8 g, 0.75 mol) and NaHCO_3 (210 g, 2.5 mol) in CH_3CN (1000 mL) was heated at reflux overnight. After cooling, the insoluble salt was filtered off and the filtrate was concentrated. The residue was diluted with CH_2Cl_2 , washed with 10% HCl solution (100 mL) and brine, dried over Na_2SO_4 , filtered and concentrated to give the crude 4-methyl-1-(phenylamino)-pentan-3-one.

[00580] 4-Methyl-1-(phenylamino)-pentan-3-ol

At $-10\text{ }^\circ\text{C}$, NaBH_4 (56.7 g, 1.5 mol) was gradually added to a mixture of the crude 4-methyl-1-(phenylamino)-pentan-3-one (about 80 g) in MeOH (500 mL). After addition, the reaction mixture was allowed to warm to room temperature and stirred for 20 min. The solvent was removed and the residue was repartitioned between water and CH_2Cl_2 . The organic phase was separated, washed with brine, dried over Na_2SO_4 , filtered and concentrated. The resulting gum was triturated with ether to give 4-methyl-1-(phenylamino)-pentan-3-ol as a white solid (22 g, 23 %).

[00581] 5,5-Dimethyl-2,3,4,5-tetrahydro-1H-benzo[b]azepine

A mixture of 4-methyl-1-(phenylamino)-pentan-3-ol (22 g, 0.11 mol) in 98% H_2SO_4 (250 mL) was stirred at $50\text{ }^\circ\text{C}$ for 30 min. The reaction mixture was poured into ice-water basified with sat. NaOH solution to pH 8 and extracted with CH_2Cl_2 . The combined organic phases were washed with brine, dried over Na_2SO_4 , filtered and concentrated. The residue was purified by column chromatography (petroleum ether) to afford 5,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[b]azepine as a brown oil (1.5 g, 8 %).

[00582] 5,5-Dimethyl-8-nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine

At $0\text{ }^\circ\text{C}$, KNO_3 (0.76 g, 7.54 mmol) was added portionwise to a solution of 5,5-dimethyl-2,3,4,5-tetrahydro-1H-benzo[b]azepine (1.1 g, 6.28 mmol) in H_2SO_4 (15 mL). After stirring 15 min at this temperature, the mixture was poured into ice water, basified with sat. NaHCO_3 to pH 8 and

extracted with EtOAc. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated to give crude 5,5-dimethyl-8-nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine (1.2 g), which was used directly in the next step without further purification.


[00583] 1-(5,5-dimethyl-8-nitro-2,3,4,5-tetrahydrobenzo[b]azepin-1-yl)ethanone

Acetyl chloride (0.77 mL, 11 mmol) was added to a suspension of crude 5,5-dimethyl-8-nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine (1.2 g, 5.45 mmol) and NaHCO₃ (1.37 g, 16.3 mmol) in CH₂Cl₂ (20 mL). The mixture was heated at reflux for 1 h. After cooling, the mixture was poured into water and extracted with CH₂Cl₂. The organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography to afford 1-(5,5-dimethyl-8-nitro-2,3,4,5-tetrahydrobenzo[b]azepin-1-yl)ethanone (1.05 g, 64 % over two steps).

[00584] DC-4; 1-(8-Amino-2,3,4,5-tetrahydro-5,5-dimethylbenzo[b]azepin-1-yl)ethanone

A suspension of 1-(5,5-dimethyl-8-nitro-2,3,4,5-tetrahydrobenzo[b]azepin-1-yl)ethanone (1.05 g, 40 mmol) and 10% Pd-C (0.2 g) in MeOH (20 mL) was stirred under H₂ (1 atm) at room temperature for 4 h. After filtration, the filtrate was concentrated to give 1-(8-amino-2,3,4,5-tetrahydro-5,5-dimethylbenzo[b]azepin-1-yl)ethanone as a white solid (**DC-4**) (880 mg, 94 %). ¹H NMR (CDCl₃) δ 7.06 (d, *J* = 8.0 Hz, 1 H), 6.59 (dd, *J* = 8.4, 2.4 Hz, 1 H), 6.50 (br s, 1H), 4.18-4.05 (m, 1H), 3.46-3.36 (m, 1H), 2.23 (s, 3H), 1.92-1.85 (m, 1H), 1.61-1.51 (m, 3H), 1.21 (s, 3H), 0.73 (t, *J* = 7.2 Hz, 3 H); ESI-MS 233.0 m/z (MH⁺).

[00585] Example 5:



[00586] Spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl

A mixture of spiro[1H-indene-1,4'-piperidin]-1'-carboxylic acid, 2,3-dihydro-3-oxo-, 1,1-dimethylethyl ester (9.50 g, 31.50 mmol) in saturated HCl/MeOH (50 mL) was stirred at 25 °C overnight. The solvent was removed under reduced pressure to yield an off-white solid (7.50 g). To a solution of this solid in dry CH₃CN (30 mL) was added anhydrous K₂CO₃ (7.85 g, 56.80 mmol). The suspension was stirred for 5 min, and benzyl bromide (5.93 g, 34.65 mmol) was added dropwise at room temperature. The mixture was stirred for 2 h, poured into cracked ice and extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄ and concentrated under *vacuum* to give crude spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl (7.93 g, 87 %), which was used without further purification.

[00587] Spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl, oxime

To a solution of spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl (7.93 g, 27.25 mmol) in EtOH (50 mL) were added hydroxylamine hydrochloride (3.79 g, 54.50 mmol) and anhydrous sodium acetate (4.02 g, 49.01 mmol) in one portion. The mixture was refluxed for 1 h, and then cooled to room temperature. The solvent was removed under reduced pressure and 200 mL of

water was added. The mixture was extracted with CH₂Cl₂. The combined organic layers were dried over Na₂SO₄ and concentrated to yield spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl, oxime (7.57 g, 91 %), which was used without further purification.

[00588] 1,2,3,4-Tetrahydroquinolin-4-spiro-4'-(N'-benzyl-piperidine)

To a solution of spiro[1H-indene-1,4'-piperidin]-3(2H)-one, 1'-benzyl, oxime (7.57 g, 24.74 mmol) in dry CH₂Cl₂ (150 mL) was added dropwise DIBAL-H (135.7 mL, 1M in toluene) at 0 °C. The mixture was stirred at 0 °C for 3 h, diluted with CH₂Cl₂ (100 mL), and quenched with NaF (20.78 g, 495 mmol) and water (6.7 g, 372 mmol). The resulting suspension was stirred vigorously at 0 °C for 30 min. After filtration, the residue was washed with CH₂Cl₂. The combined filtrates were concentrated under *vacuum* to give an off-brown oil that was purified by column chromatography on silica gel (CH₂Cl₂ – MeOH, 30:1) to afford 1,2,3,4-tetrahydroquinolin-4-spiro-4'-(N'-benzyl-piperidine) (2.72 g, 38 %).

[00589] 1,2,3,4-Tetrahydroquinolin-4-spiro-4'-piperidine

A suspension of 1,2,3,4-Tetrahydroquinolin-4-spiro-4'-(N'-benzyl-piperidine) (300 mg, 1.03 mmol) and Pd(OH)₂-C (30 mg) in MeOH (3 mL) was stirred under H₂ (55 psi) at 50 °C overnight. After cooling, the catalyst was filtered off and washed with MeOH. The combined filtrates were concentrated under reduced pressure to yield 1,2,3,4-tetrahydroquinolin-4-spiro-4'-piperidine as a white solid (176 mg, 85 %), which was used without further purification.

[00590] 7'-Nitro-spiro[piperidine-4,4'(1'H)-quinoline], 2',3'-dihydro-carboxylic acid *tert*-butyl ester

KNO₃ (69.97 mg, 0.69 mmol) was added portion-wise to a suspension of 1,2,3,4-tetrahydroquinolin-4-spiro-4'-piperidine (133 mg, 0.66 mmol) in 98% H₂SO₄ (2 mL) at 0 °C. After the addition was complete, the reaction mixture was allowed to warm to room temperature and stirred for additional 2 h. The mixture was then poured into cracked ice and basified with 10% NaOH to pH~ 8. Boc₂O (172 mg, 0.79 mmol) was added dropwise and the mixture was stirred at room temperature for 1 h. The mixture was then extracted with EtOAc and the combined organic layers were dried over Na₂SO₄, filtered and concentrated to yield crude 7'-

nitro-spiro[piperidine-4,4'(1'H)-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester (230 mg), which was used in the next step without further purification.


[00591] 7'-nitro-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro-carboxylic acid *tert*-butyl ester

Acetyl chloride (260 mg, 3.30 mmol) was added dropwise to a suspension of 7'-nitro-spiro[piperidine-4,4'(1'H)-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester (230 mg) and NaHCO₃ (1.11 g, 13.17 mmol) in MeCN (5 mL) at room temperature. The reaction mixture was refluxed for 4 h. After cooling, the suspension was filtered and the filtrate was concentrated. The residue was purified by column chromatography (petroleum ether – EtOAc, 10:1) to provide 7'-nitro-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester (150 mg, 58 % over 2 steps)

[00592] DC-5; 7'-Amino-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester

A suspension of 7'-nitro-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester (150 mg, 0.39 mmol) and Raney Ni (15 mg) in MeOH (2 mL) was stirred under H₂ (1 atm) at 25 °C overnight. The catalyst was removed via filtration and washed with MeOH. The combined filtrates were dried over Na₂SO₄, filtered, and concentrated to yield 7'-amino-spiro[piperidine-4,4'(1'H)-1-acetyl-quinoline], 2',3'-dihydro- carboxylic acid *tert*-butyl ester (**DC-5**) (133 mg, 96 %).

[00593] Example 7:



[00594] 2-(2,4-Dinitrophenylthio)-acetic acid


Et₃N (1.5 g, 15 mmol) and mercapto-acetic acid (1 g, 11 mmol) were added to a solution of 1-chloro-2,4-dinitrobenzene (2.26 g, 10 mmol) in 1,4-dioxane (50 mL) at room temperature. After

stirring at room temperature for 5 h, H₂O (100 mL) was added. The resulting suspension was extracted with ethyl acetate (100 mL x 3). The ethyl acetate extract was washed with water and brine, dried over Na₂SO₄ and concentrated to give 2-(2,4-dinitrophenylthio)-acetic acid (2.3 g, 74 %), which was used without further purification.

[00595] DC-7; 6-Amino-2H-benzo[b][1,4]thiazin-3(4H)-one

A solution of 2-(2,4-dinitrophenylthio)-acetic acid (2.3 g, 9 mmol) and tin (II) chloride dihydrate (22.6 g, 0.1 mol) in ethanol (30 mL) was refluxed overnight. After removal of the solvent under reduced pressure, the residual slurry was diluted with water (100 mL) and basified with 10 % Na₂CO₃ solution to pH 8. The resulting suspension was extracted with ethyl acetate (3 x 100 mL). The ethyl acetate extract was washed with water and brine, dried over Na₂SO₄, and concentrated. The residue was washed with CH₂Cl₂ to yield 6-amino-2H-benzo[b][1,4]thiazin-3(4H)-one (**DC-7**) as a yellow powder (1 g, 52 %). ¹H NMR (DMSO-*d*₆) δ 10.24 (s, 1 H), 6.88 (d, 1 H, *J* = 6 Hz), 6.19-6.21 (m, 2H), 5.15 (s, 2 H), 3.28 (s, 2 H); ESI-MS 181.1 m/z (MH⁺).

[00596] Example 7:



[00597] N-(2-Bromo-5-nitrophenyl)acetamide

Acetic anhydride (1.4 mL, 13.8 mmol) was added dropwise to a stirring solution of 2-bromo-5-nitroaniline (3 g, 13.8 mmol) in glacial acetic acid (30 mL) at 25 °C. The reaction mixture was stirred at room temperature overnight, and then poured into water. The precipitate was collected

via filtration, washed with water and dried under *vacuum* to provide N-(2-bromo-5-nitrophenyl)acetamide as an off white solid (3.6 g, 90 %).

[00598] N-(2-Bromo-5-nitrophenyl)-N-(2-methylprop-2-enyl)acetamide


At 25 °C, a solution of 3-bromo-2-methylpropene (3.4 g, 55.6 mmol) in anhydrous DMF (30 mL) was added dropwise to a solution of N-(2-bromo-5-nitrophenyl)acetamide (3.6 g, 13.9 mmol) and potassium carbonate (3.9 g, 27.8 mmol) in anhydrous DMF (50 mL). The reaction mixture was stirred at 25 °C overnight. The reaction mixture was then filtered and the filtrate was treated with sat. Na₂CO₃ solution. The organic layer was separated and the aqueous layer was extracted with EtOAc. The combined organic extracts were washed with water and brine, dried over MgSO₄, filtered and concentrated under *vacuum* to provide N-(2-bromo-5-nitrophenyl)-N-(2-methylprop-2-enyl)acetamide as a golden solid (3.1 g, 85 %). ESI-MS 313 m/z (MH⁺).

[00599] 1-(3,3-Dimethyl-6-nitroindolin-1-yl)ethanone

A solution of N-(2-bromo-5-nitrophenyl)-N-(2-methylprop-2-enyl)acetamide (3.1 g, 10.2 mmol), tetraethylammonium chloride hydrate (2.4 g, 149 mmol), sodium formate (1.08 g, 18 mmol), sodium acetate (2.76 g, 34.2 mmol) and palladium acetate (0.32 g, 13.2 mmol) in anhydrous DMF (50 mL) was stirred at 80 °C for 15 h under N₂ atmosphere. After cooling, the mixture was filtered through Celite. The Celite was washed with EtOAc and the combined filtrates were washed with sat. NaHCO₃. The separated organic layer was washed with water and brine, dried over MgSO₄, filtered and concentrated under reduced pressure to provide 1-(3,3-dimethyl-6-nitroindolin-1-yl)ethanone as a brown solid (2.1 g, 88%).

[00600] DC-8; 1-(6-Amino-3,3-dimethyl-2,3-dihydro-indol-1-yl)-ethanone

10% Pd-C (0.2 g) was added to a suspension of 1-(3,3-dimethyl-6-nitroindolin-1-yl)ethanone (2.1 g, 9 mmol) in MeOH (20 mL). The reaction was stirred under H₂ (40 psi) at room temperature overnight. Pd-C was filtered off and the filtrate was concentrated under *vacuum* to give a crude product, which was purified by column chromatography to yield 1-(6-amino-3,3-dimethyl-2,3-dihydro-indol-1-yl)-ethanone (**DC-8**) (1.3 g, 61 %).

[00601] Example 8:**[00602] 2,3,4,5-Tetrahydro-1H-benzo[b]azepine**

DIBAL (90 mL, 90 mmol) was added dropwise to a solution of 4-dihydro-2H-naphthalen-1-one oxime (3 g, 18 mmol) in dichloromethane (50 mL) at 0 °C. The mixture was stirred at this temperature for 2 h. The reaction was quenched with dichloromethane (30 mL), followed by treatment with NaF (2 g, 0.36 mol) and H₂O (5 mL, 0.27 mol). Vigorous stirring of the resulting suspension was continued at 0 °C for 30 min. After filtration, the filtrate was concentrated. The residue was purified by flash column chromatography to give 2,3,4,5-tetrahydro-1H-benzo[b]azepine as a colorless oil (1.9 g, 70 %).

[00603] 8-Nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine

At -10 °C, 2,3,4,5-tetrahydro-1H-benzo[b]azepine (1.9 g, 13 mmol) was added dropwise to a solution of KNO₃ (3 g, 30 mmol) in H₂SO₄ (50 mL). The mixture was stirred for 40 min, poured over crushed ice, basified with aq. ammonia to pH 13, and extracted with EtOAc. The combined organic phases were washed with brine, dried over Na₂SO₄ and concentrated to give 8-nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine as a black solid (1.3 g, 51 %), which was used without further purification.

[00604] 1-(8-Nitro-2,3,4,5-tetrahydro-benzo[b]azepin-1-yl)-ethanone


Acetyl chloride (1 g, 13 mmol) was added dropwise to a mixture of 8-nitro-2,3,4,5-tetrahydro-1H-benzo[b]azepine (1.3 g, 6.8 mmol) and NaHCO₃ (1 g, 12 mmol) in CH₂Cl₂ (50 mL). After

stirring for 1 h, the mixture was filtered and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2 , washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by column chromatography to give 1-(8-nitro-2,3,4,5-tetrahydro-benzo[b]azepin-1-yl)-ethanone as a yellow solid (1.3 g, 80 %).

[00605] DC-9; 1-(8-Amino-2,3,4,5-tetrahydro-benzo[b]azepin-1-yl)-ethanone

A mixture of 1-(8-nitro-2,3,4,5-tetrahydro-benzo[b]azepin-1-yl)-ethanone (1.3 g, 5.4 mmol) and Pd-C (10 %, 100 mg) in EtOH (200 mL) was stirred under H_2 (1 atm) at room temperature for 1.5 h. The mixture was filtered through a layer of Celite and the filtrate was concentrated to give 1-(8-amino-2,3,4,5-tetrahydro-benzo[b]azepin-1-yl)-ethanone (**DC-9**) as a white solid (1 g, 90 %). ^1H NMR (CDCl_3) δ 7.01 (d, J = 6.0 Hz, 1 H), 6.56 (dd, J = 6.0, 1.8 Hz, 1 H), 6.50 (d, J = 1.8 Hz, 1 H), 4.66-4.61 (m, 1 H), 3.50 (br s, 2 H), 2.64-2.55 (m, 3 H), 1.94-1.91 (m, 5 H), 1.77-1.72 (m, 1 H), 1.32-1.30 (m, 1 H); ESI-MS 204.1 m/z (MH^+).

[00606] Example 9:



[00607] 6-Nitro-4H-benzo[1,4]oxazin-3-one

At 0 °C, chloroacetyl chloride (8.75 mL, 0.11 mol) was added dropwise to a mixture of 4-nitro-2-aminophenol (15.4 g, 0.1 mol), benzyltrimethylammonium chloride (18.6 g, 0.1 mol) and NaHCO_3 (42 g, 0.5 mol) in chloroform (350 mL) over a period of 30 min. After addition, the reaction mixture was stirred at 0 °C for 1 h, then at 50 °C overnight. The solvent was removed under reduced pressure and the residue was treated with water (50 mL). The solid was collected

via filtration, washed with water and recrystallized from ethanol to provide 6-nitro-4H-benzo[1,4]oxazin-3-one as a pale yellow solid (8 g, 41 %).

[00608] 6-Nitro-3,4-dihydro-2H-benzo[1,4]oxazine


A solution of $\text{BH}_3 \cdot \text{Me}_2\text{S}$ in THF (2 M, 7.75 mL, 15.5 mmol) was added dropwise to a suspension of 6-nitro-4H-benzo[1,4]oxazin-3-one (0.6 g, 3.1 mmol) in THF (10 mL). The mixture was stirred at room temperature overnight. The reaction was quenched with MeOH (5 mL) at 0 °C and then water (20 mL) was added. The mixture was extracted with Et_2O and the combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated to give 6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine as a red solid (0.5 g, 89 %), which was used without further purification.

[00609] 4-Acetyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine

Under vigorous stirring at room temperature, acetyl chloride (1.02 g, 13 mmol) was added dropwise to a mixture of 6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine (1.8 g, 10 mmol) and NaHCO_3 (7.14 g, 85 mmol) in CH_2Cl_2 (50 mL). After addition, the reaction was stirred for 1 h at this temperature. The mixture was filtered and the filtrate was concentrated under *vacuum*. The residue was treated with Et_2O : hexane (1:2, 50 mL) under stirring for 30 min and then filtered to give 4-acetyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine as a pale yellow solid (2 g, 90 %).

[00610] DC-10; 4-Acetyl-6-amino-3,4-dihydro-2H-benzo[1,4]oxazine

A mixture of 4-acetyl-6-nitro-3,4-dihydro-2H-benzo[1,4]oxazine (1.5 g, 67.6 mmol) and Pd-C (10 %, 100 mg) in EtOH (30 mL) was stirred under H_2 (1 atm) overnight. The catalyst was filtered off and the filtrate was concentrated. The residue was treated with HCl / MeOH to give 4-acetyl-6-amino-3,4-dihydro-2H-benzo[1,4]oxazine hydrochloride (**DC-10**) as an off-white solid (1.1 g, 85 %). ^1H NMR ($\text{DMSO}-d_6$) δ 10.12 (br s, 2H), 8.08 (br s, 1H), 6.90-7.03 (m, 2 H), 4.24 (t, $J = 4.8$ Hz, 2 H), 3.83 (t, $J = 4.8$ Hz, 2H), 2.23 (s, 3 H); ESI-MS 192.1 m/z (MH^+).

[00611] Example 10:[00612] **1,2,3,4-Tetrahydro-7-nitroisoquinoline hydrochloride**

1,2,3,4-Tetrahydroisoquinoline (6.3 mL, 50.0 mmol) was added dropwise to a stirred ice-cold solution of concentrated H_2SO_4 (25 mL). KNO_3 (5.6 g, 55.0 mmol) was added portionwise while maintaining the temperature below 5 °C. The mixture was stirred at room temperature overnight, carefully poured into an ice-cold solution of concentrated NH_4OH , and then extracted three times with CHCl_3 . The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated. The resulting dark brown oil was taken up into EtOH, cooled in an ice bath and treated with concentrated HCl. The yellow precipitate was collected via filtration and recrystallized from methanol to give 1,2,3,4-tetrahydro-7-nitroisoquinoline hydrochloride as yellow solid (2.5 g, 23 %). ^1H NMR (400 MHz, DMSO- d_6) δ 9.86 (s, 2H), 8.22 (d, J = 1.6 Hz, 1H), 8.11 (dd, J = 8.5, 2.2 Hz, 1H), 7.53 (d, J = 8.5 Hz, 1H), 4.38 (s, 2H), 3.38 (s, 2H), 3.17-3.14 (m, 2H); HPLC ret. time 0.51 min, 10-99 % CH_3CN , 5 min run; ESI-MS 179.0 m/z (MH^+).

[00613] ***tert*-Butyl 3,4-dihydro-7-nitroisoquinoline-2(1H)-carboxylate**


A mixture of 1,2,3,4-Tetrahydro-7-nitroisoquinoline (2.5 g, 11.6 mmol), 1,4-dioxane (24 mL), H_2O (12 mL) and 1N NaOH (12 mL) was cooled in an ice-bath, and Boc_2O (2.8 g, 12.8 mmol) was added. The mixture was stirred at room temperature for 2.5 h, acidified with a 5% KHSO_4 solution to pH 2-3, and then extracted with EtOAc. The organic layer was dried over MgSO_4 and concentrated to give *tert*-butyl 3,4-dihydro-7-nitroisoquinoline-2(1H)-carboxylate (3.3 g, quant.), which was used without further purification. ^1H NMR (400 MHz, DMSO- d_6) δ 8.13 (d, J = 2.3 Hz, 1H), 8.03 (dd, J = 8.4, 2.5 Hz, 1H), 7.45 (d, J = 8.5 Hz, 1H), 4.63 (s, 2H), 3.60-3.57 (m, 2H),

2.90 (t, $J = 5.9$ Hz, 2H), 1.44 (s, 9H); HPLC ret. time 3.51 min, 10-99 % CH₃CN, 5 min run; ESI-MS 279.2 m/z (MH⁺).

[00614] **DC-6; *tert*-Butyl 7-amino-3,4-dihydroisoquinoline-2(1H)-carboxylate**
Pd(OH)₂ (330.0 mg) was added to a stirring solution of *tert*-butyl 3,4-dihydro-7-nitroisoquinoline-2(1H)-carboxylate (3.3 g, 12.0 mmol) in MeOH (56 mL) under N₂ atmosphere. The reaction mixture was stirred under H₂ (1 atm) at room temperature for 72 h. The solid was removed by filtration through Celite. The filtrate was concentrated and purified by column chromatography (15–35 % EtOAc - Hexanes) to provide *tert*-butyl 7-amino-3,4-dihydroisoquinoline-2(1H)-carboxylate (**DC-6**) as a pink oil (2.0 g, 69 %). ¹H-NMR (400 MHz, DMSO-d₆) δ 6.79 (d, $J = 8.1$ Hz, 1H), 6.40 (dd, $J = 8.1, 2.3$ Hz, 1H), 6.31 (s, 1H), 4.88 (s, 2H), 4.33 (s, 2H), 3.48 (t, $J = 5.9$ Hz, 2H), 2.58 (t, $J = 5.9$ Hz, 2H), 1.42 (s, 9H); HPLC ret. time 2.13 min, 10-99 % CH₃CN, 5 min run; ESI-MS 249.0 m/z (MH⁺).

[00615] **Other amines**

[00616] **Example 1:**



[00617] 4-Bromo-3-nitrobenzonitrile

To a solution of 4-bromobenzonitrile (4.0 g, 22 mmol) in conc. H_2SO_4 (10 mL) was added dropwise at 0 °C nitric acid (6 mL). The reaction mixture was stirred at 0 °C for 30 min, and then at room temperature for 2.5 h. The resulting solution was poured into ice-water. The white precipitate was collected via filtration and washed with water until the washings were neutral. The solid was recrystallized from an ethanol/water mixture (1:1, 20 mL) twice to afford 4-bromo-3-nitrobenzonitrile as a white crystalline solid (2.8 g, 56 %). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.54 (s, 1H), 8.06 (d, $J = 8.4$ Hz, 1H), 7.99 (d, $J = 8.4$ Hz, 1H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 150.4, 137.4, 136.6, 129.6, 119.6, 117.0, 112.6; HPLC ret. time 1.96 min, 10-100 % CH_3CN , 5 min gradient; ESI-MS 227.1 m/z (MH^+).

[00618] 2'-Ethoxy-2-nitrobiphenyl-4-carbonitrile

A 50 mL round-bottom flask was charged with 4-bromo-3-nitrobenzonitrile (1.0 g 4.4 mmol), 2-ethoxyphenylboronic acid (731 mg, 4.4 mmol), $\text{Pd}_2(\text{dba})_3$ (18 mg, 0.022 mmol) and potassium fluoride (786 mg, 13.5 mmol). The reaction vessel was evacuated and filled with argon. Dry THF (300 mL) was added followed by the addition of $\text{P}(t\text{-Bu})_3$ (0.11 mL, 10% wt. in hexane). The reaction mixture was stirred at room temperature for 30 min., and then heated at 80 °C for 16 h. After cooling to room temperature, the resulting mixture was filtered through a Celite pad and concentrated. 2'-Ethoxy-2-nitrobiphenyl-4-carbonitrile was isolated as a yellow solid (1.12 g, 95%). ^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 8.51 (s, 1H), 8.20 (d, $J = 8.1$ Hz, 1H), 7.68 (d, $J = 8.4$ Hz, 1H), 7.41 (t, $J = 8.4$ Hz, 1H), 7.37 (d, $J = 7.5$ Hz, 1H), 7.08 (t, $J = 7.5$ Hz, 1H), 7.03 (d, $J = 8.1$ Hz, 1H), 3.91 (q, $J = 7.2$ Hz, 2H), 1.12 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) δ 154.9, 149.7, 137.3, 137.2, 134.4, 131.5, 130.4, 128.4, 125.4, 121.8, 117.6, 112.3, 111.9, 64.1, 14.7; HPLC ret. time 2.43 min, 10-100 % CH_3CN , 5 min gradient; ESI-MS 269.3 m/z (MH^+).

[00619] 4-Aminomethyl-2'-ethoxy-biphenyl-2-ylamine


To a solution of 2'-ethoxy-2-nitrobiphenyl-4-carbonitrile (500 mg, 1.86 mmol) in THF (80 mL) was added a solution of $\text{BH}_3\cdot\text{THF}$ (5.6 mL, 10% wt. in THF, 5.6 mmol) at 0 °C over 30 min. The reaction mixture was stirred at 0 °C for 3 h and then at room temperature for 15 h. The reaction solution was chilled to 0 °C, and a $\text{H}_2\text{O}/\text{THF}$ mixture (3 mL) was added. After being agitated at room temperature for 6 h, the volatiles were removed under reduced pressure. The residue was

dissolved in EtOAc (100 mL) and extracted with 1N HCl (2 x 100 mL). The aqueous phase was basified with 1N NaOH solution to pH 1 and extracted with EtOAc (3 x 50 mL). The combined organic layers were washed with water (50 mL), dried over Na₂SO₄, filtered, and evaporated. After drying under vacuum, 4-aminomethyl-2'-ethoxy-biphenyl-2-ylamine was isolated as a brown oil (370 mg, 82 %). ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.28 (dt, *J* = 7.2 Hz, *J* = 1.8 Hz, 1H), 7.09 (dd, *J* = 7.2 Hz, *J* = 1.8 Hz, 1H), 7.05 (d, *J* = 7.5 Hz, 1H), 6.96 (dt, *J* = 7.2 Hz, *J* = 0.9 Hz, 1H), 6.83 (d, *J* = 7.5 Hz, 1H), 6.66 (d, *J* = 1.2 Hz, 1H), 6.57 (dd, *J* = 7.5 Hz, *J* = 1.5 Hz, 1H), 4.29 (s, 2H), 4.02 (q, *J* = 6.9 Hz, 2H), 3.60 (s, 2H), 1.21 (t, *J* = 6.9 Hz, 3H); HPLC ret. time 1.54 min, 10-100 % CH₃CN, 5 min gradient; ESI-MS 243.3 m/z (MH⁺).

[00620] E-1; (2-Amino-2'-ethoxy-biphenyl-4-ylmethyl)carbamic acid *tert*-butyl ester

A solution of Boc₂O (123 mg, 0.565 mmol) in 1,4-dioxane (10 mL) was added over a period of 30 min. to a solution of 4-aminomethyl-2'-ethoxy-biphenyl-2-ylamine (274 mg, 1.13 mmol) in 1,4-dioxane (10 mL). The reaction mixture was stirred at room temperature for 16 h. The volatiles were removed on a rotary evaporator. The residue was purified by flash chromatography (silica gel, EtOAc – CH₂Cl₂, 1:4) to afford (2-Amino-2'-ethoxy-biphenyl-4-ylmethyl)carbamic acid *tert*-butyl ester (**E-1**) as a pale yellow oil (119 mg, 31 %). ¹H NMR (300 MHz, DMSO-*d*₆) δ 7.27 (m, 2H), 7.07 (dd, *J* = 7.2 Hz, *J* = 1.8 Hz, 1H), 7.03 (d, *J* = 7.8 Hz, 1H), 6.95 (dt, *J* = 7.2 Hz, *J* = 0.9 Hz, 1H), 6.81 (d, *J* = 7.5 Hz, 1H), 6.55 (s, 1H), 6.45 (dd, *J* = 7.8 Hz, *J* = 1.5 Hz, 1H), 4.47 (s, 2H), 4.00 (q, *J* = 7.2 Hz, 2H), 1.38 (s, 9H), 1.20 (t, *J* = 7.2 Hz, 3H); HPLC ret. time 2.34 min, 10-100 % CH₃CN, 5 min gradient; ESI-MS 343.1 m/z (MH⁺).

[00621] Example 2:



[00622] 2-Bromo-1-*tert*-butyl-4-nitrobenzene

To a solution of 1-*tert*-butyl-4-nitrobenzene (8.95 g, 50 mmol) and silver sulfate (10 g, 32 mmol) in 50 mL of 90% sulfuric acid was added dropwise bromine (7.95 g, 50 mmol). Stirring was continued at room temperature overnight, and then the mixture was poured into dilute sodium hydrogen sulfite solution and was extracted with EtOAc three times. The combined organic layers were washed with brine and dried over MgSO₄. After filtration, the filtrate was concentrated to give 2-bromo-1-*tert*-butyl-4-nitrobenzene (12.7 g, 98 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.47 (d, J = 2.5 Hz, 1H), 8.11 (dd, J = 8.8, 2.5 Hz, 1H), 7.63 (d, J = 8.8 Hz, 1H), 1.57 (s, 9H); HPLC ret. time 4.05 min, 10-100 % CH₃CN, 5 min gradient.


[00623] 2-*tert*-Butyl-5-nitrobenzonitrile

To a solution of 2-bromo-1-*tert*-butyl-4-nitrobenzene (2.13 g, 8.2 mmol) and Zn(CN)₂ (770 mg, 6.56 mmol) in DMF (10 mL) was added Pd(PPh₃)₄ (474 mg, 0.41 mmol) under a nitrogen atmosphere. The mixture was heated in a sealed vessel at 205 °C for 5 h. After cooling to room temperature, the mixture was diluted with water and extracted with EtOAc twice. The combined organic layers were washed with brine and dried over MgSO₄. After removal of solvent, the residue was purified by column chromatography (0-10 % EtOAc-Hexane) to give 2-*tert*-butyl-5-nitrobenzonitrile (1.33 g, 80 %). ¹H NMR (400 MHz, CDCl₃) δ 8.55 (d, J = 2.3 Hz, 1H), 8.36 (dd, J = 8.8, 2.2 Hz, 1H), 7.73 (d, J = 8.9 Hz, 1H), 1.60 (s, 9H); HPLC ret. time 3.42 min, 10-100 % CH₃CN, 5 min gradient.

[00624] E-2; 2-*tert*-Butyl-5-aminobenzonitrile

To a refluxing solution of 2-*tert*-butyl-5-nitrobenzonitrile (816 mg, 4.0 mmol) in EtOH (20 mL) was added ammonium formate (816 mg, 12.6 mmol), followed by 10% Pd-C (570 mg). The reaction mixture was refluxed for additional 90 min, cooled to room temperature and filtered through Celite. The filtrate was concentrated to give 2-*tert*-butyl-5-aminobenzonitrile (**E-2**) (630 mg, 91 %), which was used without further purification. HPLC ret. time 2.66 min, 10-99 % CH₃CN, 5 min run; ESI-MS 175.2 m/z (MH⁺).

[00625] Example 3:



[00626] (2-*tert*-Butyl-5-nitrophenyl)methanamine

To a solution of 2-*tert*-butyl-5-nitrobenzonitrile (612 mg, 3.0 mmol) in THF (10 mL) was added a solution of $\text{BH}_3\cdot\text{THF}$ (12 mL, 1M in THF, 12.0 mmol) under nitrogen. The reaction mixture was stirred at 70 °C overnight and cooled to 0 °C. Methanol (2 mL) was added followed by the addition of 1N HCl (2 mL). After refluxing for 30 min, the solution was diluted with water and extracted with EtOAc. The aqueous layer was basified with 1N NaOH and extracted with EtOAc twice. The combined organic layers were washed with brine and dried over Mg_2SO_4 . After removal of solvent, the residue was purified by column chromatography (0-10 % MeOH - CH_2Cl_2) to give (2-*tert*-butyl-5-nitrophenyl)methanamine (268 mg, 43 %). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.54 (d, $J = 2.7$ Hz, 1H), 7.99 (dd, $J = 8.8, 2.8$ Hz, 1H), 7.58 (d, $J = 8.8$ Hz, 1H), 4.03 (s, 2H), 2.00 (t, $J = 2.1$ Hz, 2H), 1.40 (s, 9H); HPLC ret. time 2.05 min, 10-100 % CH_3CN , 5 min gradient; ESI-MS 209.3 m/z (MH^+).

[00627] *tert*-Butyl 2-*tert*-butyl-5-nitrobenzylcarbamate


A solution of (2-*tert*-butyl-5-nitrophenyl)methanamine (208 mg, 1 mmol) and Boc_2O (229 mg, 1.05 mmol) in THF (5mL) was refluxed for 30 min. After cooling to room temperature, the solution was diluted with water and extracted with EtOAc. The combined organic layers were washed with brine and dried over MgSO_4 . After filtration, the filtrate was concentrated to give *tert*-butyl 2-*tert*-butyl-5-nitrobenzylcarbamate (240 mg, 78 %), which was used without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 8.26 (d, $J = 2.3$ Hz, 1H), 8.09 (dd, $J = 8.8, 2.5$ Hz, 1H), 7.79 (t, $J = 5.9$ Hz, 1H), 7.68 (d, $J = 8.8$ Hz, 1H), 4.52 (d, $J = 6.0$ Hz, 2H), 1.48 (s, 18H); HPLC ret. time 3.72 min, 10-100 % CH_3CN , 5 min gradient.

[00628] E-4; *tert*-Butyl 2-*tert*-butyl-5-aminobenzylcarbamate

To a solution of *tert*-butyl 2-*tert*-butyl-5-nitrobenzylcarbamate (20 mg, 0.065 mmol) in 5% AcOH-MeOH (1 mL) was added 10% Pd-C (14 mg) under nitrogen atmosphere. The mixture

was stirred under H₂ (1 atm) at room temperature for 1 h. The catalyst was removed via filtration through Celite, and the filtrate was concentrated to give *tert*-butyl 2-*tert*-butyl-5-aminobenzylcarbamate (**E-4**), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.09 (d, J = 8.5 Hz, 1H), 6.62 (d, J = 2.6 Hz, 1H), 6.47 (dd, J = 8.5, 2.6 Hz, 1H), 4.61 (br s, 1H), 4.40 (d, J = 5.1 Hz, 2H), 4.15 (br s, 2H), 1.39 (s, 9H), 1.29 (s, 9H); HPLC ret. time 2.47 min, 10-100 % CH₃CN, 5 min gradient; ESI-MS 279.3 m/z (MH⁺).

[00629] Example 4:



[00630] 2-*tert*-Butyl-5-nitrobenzoic acid


A solution of 2-*tert*-butyl-5-nitrobenzonitrile (204 mg, 1 mmol) in 5 mL of 75% H₂SO₄ was microwaved at 200 °C for 30 min. The reaction mixture was poured into ice, extracted with EtOAc, washed with brine and dried over MgSO₄. After filtration, the filtrate was concentrated to give 2-*tert*-butyl-5-nitrobenzoic acid (200 mg, 90 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.36 (d, J = 2.6 Hz, 1H), 8.24 (dd, J = 8.9, 2.6 Hz, 1H), 7.72 (d, J = 8.9 Hz, 1H) 1.51 (s, 9H); HPLC ret. time 2.97 min, 10-100 % CH₃CN, 5 min gradient.

[00631] Methyl 2-*tert*-butyl-5-nitrobenzoate

To a mixture of 2-*tert*-butyl-5-nitrobenzoic acid (120 mg, 0.53 mmol) and K₂CO₃ (147 mg, 1.1 mmol) in DMF (5.0 mL) was added CH₃I (40 μL, 0.64 mmol). The reaction mixture was stirred at room temperature for 10 min, diluted with water and extracted with EtOAc. The combined organic layers were washed with brine and dried over MgSO₄. After filtration, the filtrate was concentrated to give methyl 2-*tert*-butyl-5-nitrobenzoate, which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.20 (d, J = 2.6 Hz, 1H), 8.17 (t, J = 1.8 Hz, 1H), 7.66 (d, J = 8.6 Hz, 1H), 4.11 (s, 3H), 1.43 (s, 9H).

[00632] E-6; Methyl 2-*tert*-butyl-5-aminobenzoate

To a refluxing solution of 2-*tert*-butyl-5-nitrobenzoate (90 mg, 0.38 mmol) in EtOH (2.0 mL) was added potassium formate (400 mg, 4.76 mmol) in water (1 mL), followed by the addition of 20 mg of 10% Pd-C. The reaction mixture was refluxed for additional 40 min, cooled to room temperature and filtered through Celite. The filtrate was concentrated to give methyl 2-*tert*-butyl-5-aminobenzoate (**E-6**) (76 mg, 95 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.24 (d, J = 8.6 Hz, 1H), 6.67 (dd, J = 8.6, 2.7 Hz, 1H), 6.60 (d, J = 2.7 Hz, 1H), 3.86 (s, 3H), 1.34 (s, 9H); HPLC ret. time 2.19 min, 10-99 % CH₃CN, 5 min run; ESI-MS 208.2 m/z (MH⁺).

[00633] Example 5:**[00634] 2-*tert*-Butyl-5-nitrobenzene-1-sulfonyl chloride**

A suspension of 2-*tert*-butyl-5-nitrobenzenamine (0.971 g, 5 mmol) in conc. HCl (5 mL) was cooled to 5-10 °C and a solution of NaNO₂ (0.433 g, 6.3 mmol) in H₂O (0.83 mL) was added dropwise. Stirring was continued for 0.5 h, after which the mixture was vacuum filtered. The filtrate was added, simultaneously with a solution of Na₂SO₃ (1.57 g, 12.4 mmol) in H₂O (2.7 mL), to a stirred solution of CuSO₄ (0.190 g, 0.76 mmol) and Na₂SO₃ (1.57 g, 12.4 mmol) in HCl (11.7 mL) and H₂O (2.7 mL) at 3-5 °C. Stirring was continued for 0.5 h and the resulting precipitate was filtered off, washed with water and dried to give 2-*tert*-butyl-5-nitrobenzene-1-

sulfonyl chloride (0.235 g, 17 %). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.13 (d, $J = 2.5$ Hz, 1H), 8.36 (dd, $J = 8.9, 2.5$ Hz, 1H), 7.88 (d, $J = 8.9$ Hz, 1H), 1.59 (s, 9H).


[00635] 2-*tert*-Butyl-5-nitrobenzene-1-sulfonamide

To a solution of 2-*tert*-butyl-5-nitrobenzene-1-sulfonyl chloride (100 mg, 0.36 mmol) in ether (2 mL) was added aqueous NH_4OH (128 μL , 3.6 mmol) at 0 °C. The mixture was stirred at room temperature overnight, diluted with water and extracted with ether. The combined ether extracts were washed with brine and dried over Na_2SO_4 . After removal of solvent, the residue was purified by column chromatography (0-50 % EtOAc-Hexane) to give 2-*tert*-butyl-5-nitrobenzene-1-sulfonamide (31.6 mg, 34 %).

[00636] E-7; 2-*tert*-Butyl-5-aminobenzene-1-sulfonamide

A solution of 2-*tert*-butyl-5-nitrobenzene-1-sulfonamide (32 mg, 0.12 mmol) and $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (138 mg, 0.61 mmol) in EtOH (1.5 mL) was heated in microwave oven at 100 °C for 30 min. The mixture was diluted with EtOAc and water, basified with sat. NaHCO_3 and filtered through Celite. The organic layer was separated from water and dried over Na_2SO_4 . Solvent was removed by evaporation to provide 2-*tert*-butyl-5-aminobenzene-1-sulfonamide (**E-7**) (28 mg, 100 %), which was used without further purification. HPLC ret. time 1.99 min, 10-99 % CH_3CN , 5 min run; ESI-MS 229.3 m/z (MH^+).

[00637] Example 6:




[00638] E-8; (2-*tert*-Butyl-5-aminophenyl)methanol

To a solution of methyl 2-*tert*-butyl-5-aminobenzoate (159 mg, 0.72 mmol) in THF (5 mL) was added dropwise LiAlH_4 (1.4 mL, 1M in THF, 1.4 mmol) at 0 °C. The reaction mixture was

refluxed for 2 h, diluted with H₂O and extracted with EtOAc. The combined organic layers were washed with brine and dried over MgSO₄. After filtration, the filtrate was concentrated to give (2-*tert*-butyl-5-aminophenyl)methanol (**E-8**) (25 mg, 20 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 7.17 (d, *J* = 8.5 Hz, 1H), 6.87 (d, *J* = 2.6 Hz, 1H), 6.56 (dd, *J* = 8.4, 2.7 Hz, 1H), 4.83 (s, 2H), 1.36 (s, 9H).

[00639] Example 7:



[00640] 1-Methyl-pyridinium monomethyl sulfuric acid salt

Methyl sulfate (30 mL, 39.8 g, 0.315 mol) was added dropwise to dry pyridine (25.0 g, 0.316 mol) added dropwise. The mixture was stirred at room temperature for 10 min, then at 100 °C for 2 h. The mixture was cooled to room temperature to give crude 1-methyl-pyridinium monomethyl sulfuric acid salt (64.7 g, quant.), which was used without further purification.

[00641] 1-Methyl-2-pyridone

A solution of 1-methyl-pyridinium monomethyl sulfuric acid salt (50 g, 0.243 mol) in water (54 mL) was cooled to 0 °C. Separate solutions of potassium ferricyanide (160 g, 0.486 mol) in water (320 mL) and sodium hydroxide (40 g, 1.000 mol) in water (67 mL) were prepared and added dropwise from two separatory funnels to the well-stirred solution of 1-methyl-pyridinium monomethyl sulfuric acid salt, at such a rate that the temperature of reaction mixture did not rise above 10 °C. The rate of addition of these two solutions was regulated so that all the sodium hydroxide solution had been introduced into the reaction mixture when one-half of the potassium

Ferric Cyanide solution had been added. After addition was complete, the reaction mixture was allowed to warm to room temperature and stirred overnight. Dry sodium carbonate (91.6 g) was added, and the mixture was stirred for 10 min. The organic layer was separated, and the aqueous layer was extracted with CH₂Cl₂ (100 mL × 3). The combined organic layers were dried and concentrated to yield 1-methyl-2-pyridone (25.0 g, 94 %), which was used without further purification.

[00642] 1-Methyl-3,5-dinitro-2-pyridone

1-Methyl-2-pyridone (25.0 g, 0.229 mol) was added to sulfuric acid (500 mL) at 0 °C. After stirring for 5 min., nitric acid (200 mL) was added dropwise at 0 °C. After addition, the reaction temperature was slowly raised to 100 °C, and then maintained for 5 h. The reaction mixture was poured into ice, basified with potassium carbonate to pH 8 and extracted with CH₂Cl₂ (100 mL × 3). The combined organic layers were dried over Na₂SO₄ and concentrated to yield 1-methyl-3,5-dinitro-2-pyridone (12.5 g, 28 %), which was used without further purification.


[00643] 2-Isopropyl-5-nitro-pyridine

To a solution of 1-methyl-3,5-dinitro-2-pyridone (8.0 g, 40 mmol) in methyl alcohol (20 mL) was added dropwise 3-methyl-2-butanone (5.1 mL, 48 mmol), followed by ammonia solution in methyl alcohol (10.0 g, 17%, 100 mmol). The reaction mixture was heated at 70 °C for 2.5 h under atmospheric pressure. The solvent was removed under *vacuum* and the residual oil was dissolved in CH₂Cl₂, and then filtered. The filtrate was dried over Na₂SO₄ and concentrated to afford 2-isopropyl-5-nitro-pyridine (1.88 g, 28 %).

[00644] E-9; 2-Isopropyl-5-amino-pyridine

2-Isopropyl-5-nitro-pyridine (1.30 g, 7.82 mmol) was dissolved in methyl alcohol (20 mL), and Raney Ni (0.25 g) was added. The mixture was stirred under H₂ (1 atm) at room temperature for 2 h. The catalyst was filtered off, and the filtrate was concentrated under *vaccum* to give 2-isopropyl-5-amino-pyridine (**E-9**) (0.55 g, 52 %). ¹H NMR (CDCl₃) δ 8.05 (s, 1 H), 6.93-6.99 (m, 2 H), 3.47 (br s, 2 H), 2.92-3.02 (m, 1 H), 1.24-1.26 (m, 6 H). ESI-MS 137.2 m/z (MH⁺).

[00645] Example 8:



[00646] Phosphoric acid 2,4-di-*tert*-butyl-phenyl ester diethyl ester

To a suspension of NaH (60% in mineral oil, 6.99 g, 174.7 mmol) in THF (350 mL) was added dropwise a solution of 2,4-di-*tert*-butylphenol (35 g, 169.6 mmol) in THF (150 mL) at 0 °C. The mixture was stirred at 0 °C for 15 min and then phosphorochloridic acid diethyl ester (30.15 g, 174.7 mmol) was added dropwise at 0 °C. After addition, the mixture was stirred at this temperature for 15 min. The reaction was quenched with sat. NH_4Cl (300 mL). The organic layer was separated and the aqueous phase was extracted with Et_2O (350 mL $\times$ 2). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 and concentrated under *vacuum* to give crude phosphoric acid 2,4-di-*tert*-butyl-phenyl ester diethyl ester as a yellow oil (51 g, contaminated with some mineral oil), which was used directly in the next step.

[00647] 1,3-Di-*tert*-butyl-benzene

To NH₃ (liquid, 250 mL) was added a solution of phosphoric acid 2,4-di-*tert*-butyl-phenyl ester diethyl ester (51 g, crude from last step, about 0.2 mol) in Et₂O (anhydrous, 150 mL) at -78 °C under N₂ atmosphere. Lithium metal was added to the solution in small pieces until a blue color persisted. The reaction mixture was stirred at -78 °C for 15 min and then quenched with sat. NH₄Cl solution until the mixture turned colorless. Liquid NH₃ was evaporated and the residue was dissolved in water, extracted with Et₂O (300 mL × 2). The combined organic phases were dried over Na₂SO₄ and concentrated to give crude 1,3-di-*tert*-butyl-benzene as a yellow oil (30.4 g, 94 % over 2 steps, contaminated with some mineral oil), which was used directly in next step.

[00648] 2,4-Di-*tert*-butyl-benzaldehyde and 3,5-di-*tert*-butyl-benzaldehyde

To a stirred solution of 1,3-di-*tert*-butyl-benzene (30 g, 157.6 mmol) in dry CH₂Cl₂ (700 mL) was added TiCl₄ (37.5 g, 197 mmol) at 0 °C, and followed by dropwise addition of MeOCHCl₂ (27.3 g, 236.4 mmol). The reaction was allowed to warm to room temperature and stirred for 1 h. The mixture was poured into ice-water and extracted with CH₂Cl₂. The combined organic phases were washed with NaHCO₃ and brine, dried over Na₂SO₄ and concentrated. The residue was purified by column chromatography (petroleum ether) to give a mixture of 2,4-di-*tert*-butyl-benzaldehyde and 3,5-di-*tert*-butyl-benzaldehyde (21 g, 61 %).

[00649] 2,4-Di-*tert*-butyl-5-nitro-benzaldehyde and 3,5-di-*tert*-butyl-2-nitro-benzaldehyde

To a mixture of 2,4-di-*tert*-butyl-benzaldehyde and 3,5-di-*tert*-butyl-benzaldehyde in H₂SO₄ (250 mL) was added KNO₃ (7.64 g, 75.6 mmol) in portions at 0 °C. The reaction mixture was stirred at this temperature for 20 min and then poured into crushed ice. The mixture was basified with NaOH solution to pH 8 and extracted with Et₂O (10 mL × 3). The combined organic layers were washed with water and brine and concentrated. The residue was purified by column chromatography (petroleum ether) to give a mixture of 2,4-di-*tert*-butyl-5-nitro-benzaldehyde and 3,5-di-*tert*-butyl-2-nitro-benzaldehyde (2:1 by NMR) as a yellow solid (14.7 g, 82 %). After further purification by column chromatography (petroleum ether), 2,4-di-*tert*-butyl-5-nitro-benzaldehyde (2.5 g, contains 10% 3,5-di-*tert*-butyl-2-nitro-benzaldehyde) was isolated.

[00650] 1,5-Di-*tert*-butyl-2-difluoromethyl-4-nitro-benzene and 1,5-Di-*tert*-butyl-3-difluoromethyl-2-nitro-benzene


2,4-Di-*tert*-butyl-5-nitro-benzaldehyde (2.4 g, 9.11 mmol, contaminated with 10% 3,5-di-*tert*-butyl-2-nitro-benzaldehyde) in neat deoxofluor solution was stirred at room temperature for 5 h. The reaction mixture was poured into cooled sat. NaHCO₃ solution and extracted with dichloromethane. The combined organics were dried over Na₂SO₄, concentrated and purified by column chromatography (petroleum ether) to give 1,5-di-*tert*-butyl-2-difluoromethyl-4-nitro-benzene (1.5 g) and a mixture of 1,5-di-*tert*-butyl-2-difluoromethyl-4-nitro-benzene and 1,5-di-*tert*-butyl-3-difluoromethyl-2-nitro-benzene (0.75 g, contains 28 % 1,5-di-*tert*-butyl-3-difluoromethyl-2-nitro-benzene).

[00651] E-10; 1,5-Di-*tert*-butyl-2-difluoromethyl-4-amino-benzene

To a suspension of iron powder (5.1 g, 91.1 mmol) in 50% acetic acid (25 ml) was added 1,5-di-*tert*-butyl-2-difluoromethyl-4-nitro-benzene (1.3 g, 4.56 mmol). The reaction mixture was heated at 115 °C for 15 min. Solid was filtered off was washed with acetic acid and CH₂Cl₂. The combined filtrate was concentrated and treated with HCl/MeOH. The precipitate was collected via filtration, washed with MeOH and dried to give 1,5-Di-*tert*-butyl-2-difluoromethyl-4-amino-benzene HCl salt (**E-10**) as a white solid (1.20 g, 90 %). ¹H NMR (DMSO-*d*₆) δ 7.35-7.70 (t, *J* = 53.7 Hz, 1 H), 7.56 (s, 1 H), 7.41 (s, 1 H), 1.33-1.36 (d, *J* = 8.1 Hz, 1H); ESI-MS 256.3 *m/z* (MH⁺).

[00652] Example 9

[00653] General scheme:



A) Pd(PPh₃)₄, K₂CO₃, H₂O, THF; **B)** Pd₂(dba)₃, P(*t*Bu)₃, KF, THF

[00654] Method A

In a 2-dram vial, 2-bromoaniline (100 mg, 0.58 mmol) and the corresponding aryl boronic acid (0.82 mmol) were dissolved in THF (1 mL). H₂O (500 µL) was added followed by K₂CO₃ (200 mg, 1.0 mmol) and Pd(PPh₃)₄ (100 mg, 0.1 mmol). The vial was purged with argon and sealed. The vial was then heated at 75 °C for 18 h. The crude sample was diluted in EtOAc and filtered through a silica gel plug. The organics were concentrated *via* Savant Speed-vac. The crude amine was used without further purification.

[00655] Method B


In a 2-dram vial, the corresponding aryl boronic acid (0.58 mmol) was added followed by KF (110 mg, 1.9 mmol). The solids were suspended in THF (2 mL), and then 2-bromoaniline (70 µL, 0.58 mmol) was added. The vial was purged with argon for 1 min. P(^tBu)₃ (100 µL, 10% sol. in hexanes) was added followed by Pd₂(dba)₃ (900 µL, 0.005 M in THF). The vial was purged again with argon and sealed. The vial was agitated on an orbital shaker at room temperature for 30 min and heated in a heating block at 80 °C for 16 h. The vial was then cooled to 20 °C and the suspension was passed through a pad of Celite. The pad was washed with EtOAc (5 mL). The organics were combined and concentrated under vacuum to give a crude amine that was used without further purification.

[00656] The table below includes the amines made following the general scheme above.

Product	Name	Method
F-1	4'-Methyl-biphenyl-2-ylamine	A
F-2	3'-Methyl-biphenyl-2-ylamine	A
F-3	2'-Methyl-biphenyl-2-ylamine	A
F-4	2',3'-Dimethyl-biphenyl-2-ylamine	A
F-5	(2'-Amino-biphenyl-4-yl)-methanol	A
F-6	N*4'*,N*4'*-Dimethyl-biphenyl-2,4'-diamine	B
F-7	2'-Trifluoromethyl-biphenyl-2-ylamine	B
F-8	(2'-Amino-biphenyl-4-yl)-acetonitrile	A

F-9	4'-Isobutyl-biphenyl-2-ylamine	A
F-10	3'-Trifluoromethyl-biphenyl-2-ylamine	B
F-11	2-Pyridin-4-yl-phenylamine	B
F-12	2-(1H-Indol-5-yl)-phenylamine	B
F-13	3',4'-Dimethyl-biphenyl-2-ylamine	A
F-14	4'-Isopropyl-biphenyl-2-ylamine	A
F-15	3'-Isopropyl-biphenyl-2-ylamine	A
F-16	4'-Trifluoromethyl-biphenyl-2-ylamine	B
F-17	4'-Methoxy-biphenyl-2-ylamine	B
F-18	3'-Methoxy-biphenyl-2-ylamine	B
F-19	2-Benzo[1,3]dioxol-5-yl-phenylamine	B
F-20	3'-Ethoxy-biphenyl-2-ylamine	B
F-21	4'-Ethoxy-biphenyl-2-ylamine	B
F-22	2'-Ethoxy-biphenyl-2-ylamine	B
F-23	4'-Methylsulfanyl-biphenyl-2-ylamine	B
F-24	3',4'-Dimethoxy-biphenyl-2-ylamine	B
F-25	2',6'-Dimethoxy-biphenyl-2-ylamine	B
F-26	2',5'-Dimethoxy-biphenyl-2-ylamine	B
F-27	2',4'-Dimethoxy-biphenyl-2-ylamine	B
F-28	5'-Chloro-2'-methoxy-biphenyl-2-ylamine	B
F-29	4'-Trifluoromethoxy-biphenyl-2-ylamine	B
F-30	3'-Trifluoromethoxy-biphenyl-2-ylamine	B
F-31	4'-Phenoxy-biphenyl-2-ylamine	B
F-32	2'-Fluoro-3'-methoxy-biphenyl-2-ylamine	B
F-33	2'-Phenoxy-biphenyl-2-ylamine	B
F-34	2-(2,4-Dimethoxy-pyrimidin-5-yl)-phenylamine	B
F-35	5'-Isopropyl-2'-methoxy-biphenyl-2-ylamine	B
F-36	2'-Trifluoromethoxy-biphenyl-2-ylamine	B
F-37	4'-Fluoro-biphenyl-2-ylamine	B
F-38	3'-Fluoro-biphenyl-2-ylamine	B


F-39	2'-Fluoro-biphenyl-2-ylamine	B
F-40	2'-Amino-biphenyl-3-carbonitrile	B
F-41	4'-Fluoro-3'-methyl-biphenyl-2-ylamine	B
F-42	4'-Chloro-biphenyl-2-ylamine	B
F-43	3'-Chloro-biphenyl-2-ylamine	B
F-44	3',5'-Difluoro-biphenyl-2-ylamine	B
F-45	2',3'-Difluoro-biphenyl-2-ylamine	B
F-46	3',4'-Difluoro-biphenyl-2-ylamine	B
F-47	2',4'-Difluoro-biphenyl-2-ylamine	B
F-48	2',5'-Difluoro-biphenyl-2-ylamine	B
F-49	3'-Chloro-4'-fluoro-biphenyl-2-ylamine	B
F-50	3',5'-Dichloro-biphenyl-2-ylamine	B
F-51	2',5'-Dichloro-biphenyl-2-ylamine	B
F-52	2',3'-Dichloro-biphenyl-2-ylamine	B
F-53	3',4'-Dichloro-biphenyl-2-ylamine	B
F-54	2'-Amino-biphenyl-4-carboxylic acid methyl ester	B
F-55	2'-Amino-biphenyl-3-carboxylic acid methyl ester	B
F-56	2'-Methylsulfanyl-biphenyl-2-ylamine	B
F-57	N-(2'-Amino-biphenyl-3-yl)-acetamide	B
F-58	4'-Methanesulfinyl-biphenyl-2-ylamine	B
F-59	2',4'-Dichloro-biphenyl-2-ylamine	B
F-60	4'-Methanesulfonyl-biphenyl-2-ylamine	B
F-61	2'-Amino-biphenyl-2-carboxylic acid isopropyl ester	B
F-62	2-Furan-2-yl-phenylamine	B
F-63	1-[5-(2-Amino-phenyl)-thiophen-2-yl]-ethanone	B
F-64	2-Benzo[b]thiophen-2-yl-phenylamine	B
F-65	2-Benzo[b]thiophen-3-yl-phenylamine	B
F-66	2-Furan-3-yl-phenylamine	B
F-67	2-(4-Methyl-thiophen-2-yl)-phenylamine	B
F-68	5-(2-Amino-phenyl)-thiophene-2-carbonitrile	B

[00657] Example 10:**[00658] Ethyl 2-(4-nitrophenyl)-2-methylpropanoate**


Sodium t-butoxide (466 mg, 4.85 mmol) was added to DMF (20 mL) at 0 °C. The cloudy solution was re-cooled to 5 °C. Ethyl 4-nitrophenylacetate (1.0 g, 4.78 mmol) was added. The purple slurry was cooled to 5 °C and methyl iodide (0.688 mL, 4.85 mmol) was added over 40 min. The mixture was stirred at 5-10 °C for 20 min, and then re-charged with sodium t-butoxide (466 mg, 4.85 mmol) and methyl iodide (0.699 mL, 4.85 mmol). The mixture was stirred at 5-10 °C for 20 min and a third charge of sodium t-butoxide (47 mg, 0.48 mmol) was added followed by methyl iodide (0.057 mL, 0.9 mmol). Ethyl acetate (100 mL) and HCl (0.1 N, 50 mL) were added. The organic layer was separated, washed with brine and dried over Na₂SO₄. After filtration, the filtrate was concentrated to provide ethyl 2-(4-nitrophenyl)-2-methylpropanoate (900 mg, 80 %), which was used without further purification.

[00659] G-1; Ethyl 2-(4-aminophenyl)-2-methylpropanoate

A solution of ethyl 2-(4-nitrophenyl)-2-methylpropanoate (900 mg, 3.8 mmol) in EtOH (10 mL) was treated with 10% Pd-C (80 mg) and heated to 45 °C. A solution of potassium formate (4.10 g, 48.8 mmol) in H₂O (11 mL) was added over a period of 15 min. The reaction mixture was stirred at 65 °C for 2 h and then treated with additional 300 mg of Pd/C. The reaction was stirred for 1.5 h and then filtered through Celite. The solvent volume was reduced by approximately 50 % under reduced pressure and extracted with EtOAc. The organic layers were dried over Na₂SO₄ and the solvent was removed under reduced pressure to yield ethyl 2-(4-aminophenyl)-2-methylpropanoate (**G-1**) (670 mg, 85 %). ¹H NMR (400 MHz, CDCl₃) δ 7.14 (d, J = 8.5 Hz, 2H), 6.65 (d, J = 8.6 Hz, 2H), 4.10 (q, J = 7.1 Hz, 2H), 1.53 (s, 6H), 1.18 (t, J = 7.1 Hz, 3H).

[00660] Example 11:**[00661] G-2; 2-(4-Aminophenyl)-2-methylpropan-1-ol**

A solution of ethyl 2-(4-aminophenyl)-2-methylpropanoate (30 mg, 0.145 mmol) in THF (1 mL) was treated with LiAlH_4 (1M solution in THF, 0.226 mL, 0.226 mmol) at 0 °C and stirred for 15 min. The reaction was treated with 0.1N NaOH, extracted with EtOAc and the organic layers were dried over Na_2SO_4 . The solvent was removed under reduced pressure to yield 2-(4-aminophenyl)-2-methylpropan-1-ol (G-2), which was used without further purification: ^1H NMR (400 MHz, CDCl_3) δ 7.17 (d, J = 8.5 Hz, 2H), 6.67 (d, J = 8.5 Hz, 2H), 3.53 (s, 2H), 1.28 (s, 6H).

[00662] Example 12:**[00663] 2-methyl-2-(4-nitrophenyl)propanenitrile**

A suspension of sodium *tert*-butoxide (662 mg, 6.47 mmol) in DMF (20 mL) at 0 °C was treated with 4-nitrophenylacetonitrile (1000 mg, 6.18 mmol) and stirred for 10 min. Methyl iodide (400 μL , 6.47 mmol) was added dropwise over 15 min. The solution was stirred at 0-10 °C for 15 min and then at room temperature for additional 15 min. To this purple solution was added sodium *tert*-butoxide (662 mg, 6.47 mmol) and the solution was stirred for 15 min. Methyl iodide (400 μL , 6.47 mmol) was added dropwise over 15 min and the solution was stirred

overnight. Sodium *tert*-butoxide (192 mg, 1.94 mmol) was added and the reaction was stirred at 0 °C for 10 minutes. Methyl iodide (186 μ L, 2.98 mmol) was added and the reaction was stirred for 1h. The reaction mixture was then partitioned between 1N HCl (50 mL) and EtOAc (75 mL). The organic layer was washed with 1 N HCl and brine, dried over Na₂SO₄ and concentrated to yield 2-methyl-2-(4-nitrophenyl)propanenitrile as a green waxy solid (1.25 g, 99 %). ¹H NMR (400 MHz, CDCl₃) δ 8.24 (d, J = 8.9 Hz, 2H), 7.66 (d, J = 8.9 Hz, 2H), 1.77 (s, 6H).

[00664] 2-Methyl-2-(4-nitrophenyl)propan-1-amine

To a cooled solution of 2-methyl-2-(4-nitrophenyl)propanenitrile (670 mg, 3.5 mmol) in THF (15 mL) was added BH₃ (1M in THF, 14 mL, 14 mmol) dropwise at 0 °C. The mixture was warmed to room temperature and heated at 70 °C for 2 h. 1N HCl solution (2 mL) was added, followed by the addition of NaOH until pH > 7. The mixture was extracted with ether and ether extract was concentrated to give 2-methyl-2-(4-nitrophenyl)propan-1-amine (610 mg, 90 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.20 (d, J = 9.0 Hz, 2H), 7.54 (d, J = 9.0 Hz, 2H), 2.89 (s, 2H), 1.38 (s, 6H).

[00665] *tert*-Butyl 2-methyl-2-(4-nitrophenyl)propylcarbamate


To a cooled solution of 2-methyl-2-(4-nitrophenyl)propan-1-amine (600 mg, 3.1 mmol) and 1N NaOH (3 mL, 3 mmol) in 1,4-dioxane (6 mL) and water (3 mL) was added Boc₂O (742 mg, 3.4 mmol) at 0 °C. The reaction was allowed to warm to room temperature and stirred overnight. The reaction was made acidic with 5% KHSO₄ solution and then extracted with ethyl acetate. The organic layer was dried over MgSO₄ and concentrated to give *tert*-butyl 2-methyl-2-(4-nitrophenyl)propylcarbamate (725 mg, 80 %), which was used without further purification. ¹H NMR (400 MHz, CDCl₃) δ 8.11 (d, J = 8.9 Hz, 2H), 7.46 (d, J = 8.8 Hz, 2H), 3.63 (s, 2H), 1.31-1.29 (m, 15H).

[00666] G-3; *tert*-Butyl 2-methyl-2-(4-aminophenyl)propylcarbamate

To a refluxing solution of *tert*-butyl 2-methyl-2-(4-nitrophenyl)propylcarbamate (725 mg, 2.5 mmol) and ammonium formate (700 mg, 10.9 mmol) in EtOH (25 mL) was added Pd-5%wt on carbon (400 mg). The mixture was refluxed for 1 h, cooled and filtered through Celite. The filtrate was concentrated to give *tert*-butyl 2-methyl-2-(4-aminophenyl)propylcarbamate (G-3)

(550 mg, 83 %), which was used without further purification. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 6.99 (d, $J = 8.5$ Hz, 2H), 6.49 (d, $J = 8.6$ Hz, 2H), 4.85 (s, 2H), 3.01 (d, $J = 6.3$ Hz, 2H), 1.36 (s, 9H), 1.12 (s, 6H); HPLC ret. time 2.02 min, 10-99 % CH_3CN , 5 min run; ESI-MS 265.2 m/z (MH^+).

[00667] Example 13:




[00668] 7-Nitro-1,2,3,4-tetrahydro-naphthalen-1-ol

7-Nitro-3,4-dihydro-2H-naphthalen-1-one (200 mg, 1.05 mmol) was dissolved in methanol (5 mL) and NaBH_4 (78 mg, 2.05 mmol) was added in portions. The reaction was stirred at room temperature for 20 min and then concentrated and purified by column chromatography (10-50 % ethyl acetate - hexanes) to yield 7-nitro-1,2,3,4-tetrahydro-naphthalen-1-ol (163 mg, 80 %). ^1H NMR (400 MHz, CD_3CN) δ 8.30 (d, $J = 2.3$ Hz, 1H), 8.02 (dd, $J = 8.5, 2.5$ Hz, 1H), 7.33 (d, $J = 8.5$ Hz, 1H), 4.76 (t, $J = 5.5$ Hz, 1H), 2.96-2.80 (m, 2H), 2.10-1.99 (m, 2H), 1.86-1.77 (m, 2H); HPLC ret. time 2.32 min, 10-99 % CH_3CN , 5 min run.

[00669] H-1; 7-Amino-1,2,3,4-tetrahydro-naphthalen-1-ol

7-nitro-1,2,3,4-tetrahydro-naphthalen-1-ol (142 mg, 0.73 mmol) was dissolved in methanol (10 mL) and the flask was flushed with N_2 (g). 10% Pd-C (10 mg) was added and the reaction was stirred under H_2 (1 atm) at room temperature overnight. The reaction was filtered and the filtrate concentrated to yield 7-amino-1,2,3,4-tetrahydro-naphthalen-1-ol (**H-1**) (113 mg, 95 %). HPLC ret. time 0.58 min, 10-99 % CH_3CN , 5 min run; ESI-MS 164.5 m/z (MH^+).

[00670] Example 14:



[00671] 7-Nitro-3,4-dihydro-2H-naphthalen-1-one oxime

To a solution of 7-nitro-3,4-dihydro-2H-naphthalen-1-one (500 mg, 2.62 mmol) in pyridine (2 mL) was added hydroxylamine solution (1 mL, ~50% solution in water). The reaction was stirred at room temperature for 1 h, then concentrated and purified by column chromatography (10-50 % ethyl acetate - hexanes) to yield 7-nitro-3,4-dihydro-2H-naphthalen-1-one oxime (471 mg, 88 %). HPLC ret. time 2.67 min, 10-99 % CH₃CN, 5 min run; ESI-MS 207.1 m/z (MH⁺).

[00672] 1,2,3,4-Tetrahydro-naphthalene-1,7-diamine


7-Nitro-3,4-dihydro-2H-naphthalen-1-one oxime (274 mg, 1.33 mmol) was dissolved in methanol (10 mL) and the flask was flushed with N₂ (g). 10 % Pd-C (50 mg) was added and the reaction was stirred under H₂ (1 atm) at room temperature overnight. The reaction was filtered and the filtrate was concentrated to yield 1,2,3,4-tetrahydro-naphthalene-1,7-diamine (207 mg, 96 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 6.61-6.57 (m, 2H), 6.28 (dd, J = 8.0, 2.4 Hz, 1H), 4.62 (s, 2H), 3.58 (m, 1H), 2.48-2.44 (m, 2H), 1.78-1.70 (m, 2H), 1.53-1.37 (m, 2H).

[00673] H-2; (7-Amino-1,2,3,4-tetrahydro-naphthalen-1-yl)-carbamic acid tert-butyl ester

To a solution of 1,2,3,4-tetrahydro-naphthalene-1,7-diamine (154 mg, 0.95 mmol) and triethylamine (139 μL, 1.0 mmol) in methanol (2 mL) cooled to 0 °C was added di-*tert*-butyl dicarbonate (207 mg, 0.95 mmol). The reaction was stirred at 0 °C and then concentrated and purified by column chromatography (5-50 % methanol - dichloromethane) to yield (7-amino-

1,2,3,4-tetrahydro-naphthalen-1-yl)-carbamic acid tert-butyl ester (**H-2**) (327 mg, quant.). HPLC ret. time 1.95 min, 10-99 % CH₃CN, 5 min run; ESI-MS 263.1 m/z (MH⁺).

[00674] Example 15:



[00675] N-(2-Bromo-benzyl)-2,2,2-trifluoro-acetamide

To a solution of 2-bromobenzylamine (1.3 mL, 10.8 mmol) in methanol (5 mL) was added ethyl trifluoroacetate (1.54 mL, 21.6 mmol) and triethylamine (1.4 mL, 10.8 mmol) under a nitrogen atmosphere. The reaction was stirred at room temperature for 1 h. The reaction mixture was then concentrated under vacuum to yield N-(2-bromo-benzyl)-2,2,2-trifluoro-acetamide (3.15g, quant.). HPLC ret. time 2.86 min, 10-99 % CH₃CN, 5 min run; ESI-MS 283.9 m/z (MH⁺).

[00676] I-1; N-(4'-Amino-biphenyl-2-ylmethyl)-2,2,2-trifluoro-acetamide

A mixture of N-(2-bromo-benzyl)-2,2,2-trifluoro-acetamide (282 mg, 1.0 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (284 mg, 1.3 mmol), Pd(OAc)₂ (20 mg, 0.09 mmol) and PS-PPh₃ (40 mg, 3 mmol / g, 0.12 mmol) was dissolved in DMF (5 mL) and 4M K₂CO₃ solution (0.5 mL) was added. The reaction was heated at 80 °C overnight. The mixture was filtered, concentrated and purified by column chromatography (0-50 % ethyl acetate - hexanes) to yield N-(4'-amino-biphenyl-2-ylmethyl)-2,2,2-trifluoro-acetamide (**I-1**) (143 mg, 49 %). HPLC ret. time 1.90 min, 10-99 % CH₃CN, 5 min run; ESI-MS 295.5 m/z (MH⁺).

[00677] Commercially available amines

<u>Amine</u>	Name
J-1	2-methoxy-5-methylbenzenamine
J-2	2,6-diisopropylbenzenamine
J-3	pyridin-2-amine
J-4	4-pentylbenzenamine
J-5	isoquinolin-3-amine
J-6	aniline
J-7	4-phenoxybenzenamine
J-8	2-(2,3-dimethylphenoxy)pyridin-3-amine
J-9	4-ethynylbenzenamine
J-10	2-sec-butylbenzenamine
J-11	2-amino-4,5-dimethoxybenzonitrile
J-12	2- <i>tert</i> -butylbenzenamine
J-13	1-(7-amino-3,4-dihydroisoquinolin-2(1H)-yl)ethanone
J-14	4-(4-methyl-4H-1,2,4-triazol-3-yl)benzenamine
J-15	2'-Aminomethyl-biphenyl-4-ylamine
J-16	1H-Indazol-6-ylamine
J-17	2-(2-methoxyphenoxy)-5-(trifluoromethyl)benzenamine
J-18	2- <i>tert</i> -butylbenzenamine
J-19	2,4,6-trimethylbenzenamine
J-20	5,6-dimethyl-1H-benzo[d]imidazol-2-amine
J-21	2,3-dihydro-1H-inden-4-amine
J-22	2-sec-butyl-6-ethylbenzenamine
J-23	quinolin-5-amine
J-24	4-(benzyloxy)benzenamine
J-25	2'-Methoxy-biphenyl-2-ylamine
J-26	benzo[c][1,2,5]thiadiazol-4-amine
J-27	3-benzylbenzenamine
J-28	4-isopropylbenzenamine
J-29	2-(phenylsulfonyl)benzenamine

J-30	2-methoxybenzenamine
J-31	4-amino-3-ethylbenzonitrile
J-32	4-methylpyridin-2-amine
J-33	4-chlorobenzenamine
J-34	2-(benzyloxy)benzenamine
J-35	2-amino-6-chlorobenzonitrile
J-36	3-methylpyridin-2-amine
J-37	4-aminobenzonitrile
J-38	3-chloro-2,6-diethylbenzenamine
J-39	3-phenoxybenzenamine
J-40	2-benzylbenzenamine
J-41	2-(2-fluorophenoxy)pyridin-3-amine
J-42	5-chloropyridin-2-amine
J-43	2-(trifluoromethyl)benzenamine
J-44	(4-(2-aminophenyl)piperazin-1-yl)(phenyl)methanone
J-45	1H-benzo[d][1,2,3]triazol-5-amine
J-46	2-(1H-indol-2-yl)benzenamine
J-47	4-Methyl-biphenyl-3-ylamine
J-48	pyridin-3-amine
J-49	3,4-dimethoxybenzenamine
J-50	3H-benzo[d]imidazol-5-amine
J-51	3-aminobenzonitrile
J-52	6-chloropyridin-3-amine
J-53	o-toluidine
J-54	1H-indol-5-amine
J-55	[1,2,4]triazolo[1,5-a]pyridin-8-amine
J-56	2-methoxypyridin-3-amine
J-57	2-butoxybenzenamine
J-58	2,6-dimethylbenzenamine
J-59	2-(methylthio)benzenamine

J-60	2-(5-methylfuran-2-yl)benzenamine
J-61	3-(4-aminophenyl)-3-ethylpiperidine-2,6-dione
J-62	2,4-dimethylbenzenamine
J-63	5-fluoropyridin-2-amine
J-64	4-cyclohexylbenzenamine
J-65	4-Amino-benzenesulfonamide
J-66	2-ethylbenzenamine
J-67	4-fluoro-3-methylbenzenamine
J-68	2,6-dimethoxypyridin-3-amine
J-69	4- <i>tert</i> -butylbenzenamine
J-70	4- <i>sec</i> -butylbenzenamine
J-71	5,6,7,8-tetrahydronaphthalen-2-amine
J-72	3-(Pyrrolidine-1-sulfonyl)-phenylamine
J-73	4-Adamantan-1-yl-phenylamine
J-74	3-amino-5,6,7,8-tetrahydronaphthalen-2-ol
J-75	benzo[d][1,3]dioxol-5-amine
J-76	5-chloro-2-phenoxybenzenamine
J-77	N1-tosylbenzene-1,2-diamine
J-78	3,4-dimethylbenzenamine
J-79	2-(trifluoromethylthio)benzenamine
J-80	1H-indol-7-amine
J-81	3-methoxybenzenamine
J-82	quinolin-8-amine
J-83	2-(2,4-difluorophenoxy)pyridin-3-amine
J-84	2-(4-aminophenyl)acetonitrile
J-85	2,6-dichlorobenzenamine
J-86	2,3-dihydrobenzofuran-5-amine
J-87	p-toluidine
J-88	2-methylquinolin-8-amine
J-89	2- <i>tert</i> -butylbenzenamine

J-90	3-chlorobenzenamine
J-91	4- <i>tert</i> -butyl-2-chlorobenzenamine
J-92	2-Amino-benzenesulfonamide
J-93	1-(2-aminophenyl)ethanone
J-94	m-toluidine
J-95	2-(3-chloro-5-(trifluoromethyl)pyridin-2-yloxy)benzenamine
J-96	2-amino-6-methylbenzonitrile
J-97	2-(prop-1-en-2-yl)benzenamine
J-98	4-Amino-N-pyridin-2-yl-benzenesulfonamide
J-99	2-ethoxybenzenamine
J-100	naphthalen-1-amine
J-101	Biphenyl-2-ylamine
J-102	2-(trifluoromethyl)-4-isopropylbenzenamine
J-103	2,6-diethylbenzenamine
J-104	5-(trifluoromethyl)pyridin-2-amine
J-105	2-aminobenzamide
J-106	3-(trifluoromethoxy)benzenamine
J-107	3,5-bis(trifluoromethyl)benzenamine
J-108	4-vinylbenzenamine
J-109	4-(trifluoromethyl)benzenamine
J-110	2-morpholinobenzenamine
J-111	5-amino-1H-benzo[d]imidazol-2(3H)-one
J-112	quinolin-2-amine
J-113	3-methyl-1H-indol-4-amine
J-114	pyrazin-2-amine
J-115	1-(3-aminophenyl)ethanone
J-116	2-ethyl-6-isopropylbenzenamine
J-117	2-(3-(4-chlorophenyl)-1,2,4-oxadiazol-5-yl)benzenamine
J-118	N-(4-amino-2,5-diethoxyphenyl)benzamide
J-119	5,6,7,8-tetrahydronaphthalen-1-amine

J-120	2-(1H-benzo[d]imidazol-2-yl)benzenamine
J-121	1,1-Dioxo-1H-1 λ 6*-benzo[b]thiophen-6-ylamine
J-122	2,5-diethoxybenzenamine
J-123	2-isopropyl-6-methylbenzenamine
J-124	tert-butyl 5-amino-3,4-dihydroisoquinoline-2(1H)-carboxylate
J-125	2-(2-aminophenyl)ethanol
J-126	(4-aminophenyl)methanol
J-127	5-methylpyridin-2-amine
J-128	2-(pyrrolidin-1-yl)benzenamine
J-129	4-propylbenzenamine
J-130	3,4-dichlorobenzenamine
J-131	2-phenoxybenzenamine
J-132	Biphenyl-2-ylamine
J-133	2-chlorobenzenamine
J-134	2-amino-4-methylbenzonitrile
J-135	(2-aminophenyl)(phenyl)methanone
J-136	aniline
J-137	3-(trifluoromethylthio)benzenamine
J-138	2-(2,5-dimethyl-1H-pyrrol-1-yl)benzenamine
J-139	4-(Morpholine-4-sulfonyl)-phenylamine
J-140	2-methylbenzo[d]thiazol-5-amine
J-141	2-amino-3,5-dichlorobenzonitrile
J-142	2-fluoro-4-methylbenzenamine
J-143	6-ethylpyridin-2-amine
J-144	2-(1H-pyrrol-1-yl)benzenamine
J-145	2-methyl-1H-indol-5-amine
J-146	quinolin-6-amine
J-147	1H-benzo[d]imidazol-2-amine
J-148	2-o-tolylbenzo[d]oxazol-5-amine
J-149	5-phenylpyridin-2-amine

J-150	Biphenyl-2-ylamine
J-151	4-(difluoromethoxy)benzenamine
J-152	5- <i>tert</i> -butyl-2-methoxybenzenamine
J-153	2-(2- <i>tert</i> -butylphenoxy)benzenamine
J-154	3-aminobenzamide
J-155	4-morpholinobenzenamine
J-156	6-aminobenzo[d]oxazol-2(3H)-one
J-157	2-phenyl-3H-benzo[d]imidazol-5-amine
J-158	2,5-dichloropyridin-3-amine
J-159	2,5-dimethylbenzenamine
J-160	4-(phenylthio)benzenamine
J-161	9H-fluoren-1-amine
J-162	2-(4-aminophenyl)-1,1,1,3,3,3-hexafluoropropan-2-ol
J-163	4-bromo-2-ethylbenzenamine
J-164	4-methoxybenzenamine
J-165	3-(Piperidine-1-sulfonyl)-phenylamine
J-166	quinoxalin-6-amine
J-167	6-(trifluoromethyl)pyridin-3-amine
J-168	3-(trifluoromethyl)-2-methylbenzenamine
J-169	(2-aminophenyl)(phenyl)methanol
J-170	aniline
J-171	6-methoxypyridin-3-amine
J-172	4-butylbenzenamine
J-173	3-(Morpholine-4-sulfonyl)-phenylamine
J-174	2,3-dimethylbenzenamine
J-175	aniline
J-176	Biphenyl-2-ylamine
J-177	2-(2,4-dichlorophenoxy)benzenamine
J-178	pyridin-4-amine
J-179	2-(4-methoxyphenoxy)-5-(trifluoromethyl)benzenamine


J-180	6-methylpyridin-2-amine
J-181	5-chloro-2-fluorobenzenamine
J-182	1H-indol-4-amine
J-183	6-morpholinopyridin-3-amine
J-184	aniline
J-185	1H-indazol-5-amine
J-186	2-[(Cyclohexyl-methyl-amino)-methyl]-phenylamine
J-187	2-phenylbenzo[d]oxazol-5-amine
J-188	naphthalen-2-amine
J-189	2-aminobenzonitrile
J-190	N1,N1-diethyl-3-methylbenzene-1,4-diamine
J-191	aniline
J-192	2-butylbenzenamine
J-193	1-(4-aminophenyl)ethanol
J-194	2-amino-4-methylbenzamide
J-195	quinolin-3-amine
J-196	2-(piperidin-1-yl)benzenamine
J-197	3-Amino-benzenesulfonamide
J-198	2-ethyl-6-methylbenzenamine
J-199	Biphenyl-4-ylamine
J-200	2-(o-tolyloxy)benzenamine
J-201	5-amino-3-methylbenzo[d]oxazol-2(3H)-one
J-202	4-ethylbenzenamine
J-203	2-isopropylbenzenamine
J-204	3-(trifluoromethyl)benzenamine
J-205	2-amino-6-fluorobenzonitrile
J-206	2-(2-aminophenyl)acetonitrile
J-207	2-(4-fluorophenoxy)pyridin-3-amine
J-208	aniline
J-209	2-(4-methylpiperidin-1-yl)benzenamine

J-210	4-fluorobenzenamine
J-211	2-propylbenzenamine
J-212	4-(trifluoromethoxy)benzenamine
J-213	3-aminophenol
J-214	2,2-difluorobenzo[d][1,3]dioxol-5-amine
J-215	2,2,3,3-tetrafluoro-2,3-dihydrobenzo[b][1,4]dioxin-6-amine
J-216	N-(3-aminophenyl)acetamide
J-217	1-(3-aminophenyl)-3-methyl-1H-pyrazol-5(4H)-one
J-218	5-(trifluoromethyl)benzene-1,3-diamine
J-219	5-tert-butyl-2-methoxybenzene-1,3-diamine
J-220	N-(3-amino-4-ethoxyphenyl)acetamide
J-221	N-(3-Amino-phenyl)-methanesulfonamide
J-222	N-(3-aminophenyl)propionamide
J-223	N1,N1-dimethylbenzene-1,3-diamine
J-224	N-(3-amino-4-methoxyphenyl)acetamide
J-225	benzene-1,3-diamine
J-226	4-methylbenzene-1,3-diamine
J-227	1H-indol-6-amine
J-228	6,7,8,9-tetrahydro-5H-carbazol-2-amine
J-229	1H-indol-6-amine
J-230	1H-indol-6-amine
J-231	1H-indol-6-amine
J-232	1H-indol-6-amine
J-233	1H-indol-6-amine
J-234	1H-indol-6-amine
J-235	1H-indol-6-amine
J-236	1H-indol-6-amine
J-237	1H-indol-6-amine
J-238	1H-indol-6-amine
J-239	1-(6-Amino-2,3-dihydro-indol-1-yl)-ethanone

J-240	5-Chloro-benzene-1,3-diamine
-------	------------------------------

[00678] Amides (Compounds of formula I)


[00679] General scheme:



a) Ar₁R₇NH₂, coupling reagent, base, solvent. Examples of conditions used:

HATU, DIEA, DMF; BOP, DIEA, DMF; HBTU, Et₃N, CH₂Cl₂; PFP-TFA, pyridine

[00680] Specific example:



[00681] **215; 4-Oxo-N-phenyl-1H-quinoline-3-carboxamide**

To a solution of 4-hydroxy-quinoline-3-carboxylic acid (**A-1**) (19 mg, 0.1 mmol), HATU (38 mg, 0.1mmol) and DIEA (34.9 μ L, 0.2mmol) in DMF (1 mL) was added aniline (18.2 μ L, 0.2 mmol) and the reaction mixture was stirred at room temperature for 3 h. The resulting solution was filtered and purified by HPLC (10-99 % CH₃CN / H₂O) to yield 4-oxo-N-phenyl-1H-quinoline-3-carboxamide (**215**) (12 mg, 45 %). ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.97 (s, 1H), 12.50 (s, 1H), 8.89 (s, 1H), 8.34 (dd, J = 8.1, 1.1 Hz, 1H), 7.83 (t, J = 8.3 Hz, 1H), 7.75 (m, 3H),

7.55 (t, J = 8.1 Hz, 1H), 7.37 (t, J = 7.9 Hz, 2H), 7.10 (t, J = 6.8 Hz, 1H); HPLC ret. time 3.02 min, 10-99 % CH₃CN, 5 min run; ESI-MS 265.1 m/z (MH⁺).

[00682] The table below lists other examples synthesized by the general scheme above.

Compound of formula I	Acid	Amine
2	A-1	C-2
3	A-1	J-17
4	A-1	J-110
5	A-1	G-2
6	A-1	E-8
7	A-1	J-118
8	A-1	D-7
9	A-1	J-197
11	A-1	F-7
12	A-1	F-6
13	A-1	E-2
15	A-1	J-56
16	A-1	J-211
18	A-1	J-161
19	A-1	J-112
20	A-1	J-200
21	A-1	J-98
23	A-1	C-15
24	A-1	J-72
25	A-1	F-57
26	A-1	J-196
29	A-21	J-208
31	A-1	J-87
32	A-1	B-21
33	A-1	J-227
34	A-1	C-19
36	A-1	J-203
37	A-1	J-80

38	A-1	J-46
39	A-17	D-10
40	A-1	J-125
42	A-1	J-95
43	A-1	C-16
44	A-1	J-140
45	A-1	J-205
47	A-1	J-102
48	A-1	J-181
49	A-1	F-25
50	A-1	J-19
51	A-7	B-24
52	A-1	F-2
53	A-1	J-178
54	A-1	J-26
55	A-1	J-219
56	A-1	J-74
57	A-1	J-61
58	A-1	D-4
59	A-1	F-35
60	A-1	D-11
61	A-1	J-174
62	A-1	J-106
63	A-1	F-47
64	A-1	J-111
66	A-1	J-214
67	A-10	J-236
68	A-1	F-55
69	A-1	D-8
70	A-1	F-11
71	A-1	F-61
72	A-1	J-66
73	A-1	J-157
74	A-1	J-104
75	A-1	J-195
76	A-1	F-46

77	A-1	B-20
78	A-1	J-92
79	A-1	F-41
80	A-1	J-30
81	A-1	J-222
82	A-1	J-190
83	A-1	F-40
84	A-1	J-32
85	A-1	F-53
86	A-1	J-15
87	A-1	J-39
88	A-1	G-3
89	A-1	J-134
90	A-1	J-18
91	A-1	J-38
92	A-1	C-13
93	A-1	F-68
95	A-1	J-189
96	A-1	B-9
97	A-1	F-34
99	A-1	J-4
100	A-1	J-182
102	A-1	J-117
103	A-2	C-9
104	A-1	B-4
106	A-1	J-11
107	A-1	DC-6
108	A-1	DC-3
109	A-1	DC-4
110	A-1	J-84
111	A-1	J-43
112	A-11	J-235
113	A-1	B-7
114	A-1	D-18
115	A-1	F-62
116	A-3	J-229

118	A-1	F-12
120	A-1	J-1
121	A-1	J-130
122	A-1	J-49
123	A-1	F-66
124	A-2	B-24
125	A-1	J-143
126	A-1	C-25
128	A-22	J-176
130	A-14	J-233
131	A-1	J-240
132	A-1	J-220
134	A-1	F-58
135	A-1	F-19
136	A-1	C-8
137	A-6	C-9
138	A-1	F-44
139	A-1	F-59
140	A-1	J-64
142	A-1	J-10
143	A-1	C-7
144	A-1	J-213
145	A-1	B-18
146	A-1	J-55
147	A-1	J-207
150	A-1	J-162
151	A-1	F-67
152	A-1	J-156
153	A-1	C-23
154	A-1	J-107
155	A-1	J-3
156	A-1	F-36
160	A-1	D-6
161	A-1	C-3
162	A-1	J-171
164	A-1	J-204

165	A-1	J-65
166	A-1	F-54
167	A-1	J-226
168	A-1	J-48
169	A-1	B-1
170	A-1	J-42
171	A-1	F-52
172	A-1	F-64
173	A-1	J-180
174	A-1	F-63
175	A-1	DC-2
176	A-1	J-212
177	A-1	J-57
178	A-1	J-153
179	A-1	J-154
180	A-1	J-198
181	A-1	F-1
182	A-1	F-37
183	A-1	DC-1
184	A-15	J-231
185	A-1	J-173
186	A-1	B-15
187	A-1	B-3
188	A-1	B-25
189	A-1	J-24
190	A-1	F-49
191	A-1	J-23
192	A-1	J-36
193	A-1	J-68
194	A-1	J-37
195	A-1	J-127
197	A-1	J-167
198	A-1	J-210
199	A-1	F-3
200	A-1	H-1
201	A-1	J-96

202	A-1	F-28
203	A-1	B-2
204	A-1	C-5
205	A-1	J-179
206	A-1	J-8
207	A-1	B-17
208	A-1	C-12
209	A-1	J-126
210	A-17	J-101
211	A-1	J-152
212	A-1	J-217
213	A-1	F-51
214	A-1	J-221
215	A-1	J-136
216	A-1	J-147
217	A-1	J-185
218	A-2	C-13
219	A-1	J-114
220	A-1	C-26
222	A-1	J-35
223	A-1	F-23
224	A-1	I-1
226	A-1	J-129
227	A-1	J-120
228	A-1	J-169
229	A-1	J-59
230	A-1	J-145
231	A-1	C-17
233	A-1	J-239
234	A-1	B-22
235	A-1	E-9
236	A-1	J-109
240	A-1	J-34
241	A-1	J-82
242	A-1	D-2
244	A-1	J-228

245	A-1	J-177
246	A-1	J-78
247	A-1	F-33
250	A-1	J-224
252	A-1	J-135
253	A-1	F-30
254	A-2	B-20
255	A-8	C-9
256	A-1	J-45
257	A-1	J-67
259	A-1	B-14
261	A-1	F-13
262	A-1	DC-7
263	A-1	J-163
264	A-1	J-122
265	A-1	J-40
266	A-1	C-14
267	A-1	J-7
268	A-1	E-7
270	A-1	B-5
271	A-1	D-9
273	A-1	H-2
274	A-8	B-24
276	A-1	J-139
277	A-1	F-38
278	A-1	F-10
279	A-1	F-56
280	A-1	J-146
281	A-1	J-62
283	A-1	F-18
284	A-1	J-16
285	A-1	F-45
286	A-1	J-119
287	A-3	C-13
288	A-1	C-6
289	A-1	J-142

290	A-1	F-15
291	A-1	C-10
292	A-1	J-76
293	A-1	J-144
294	A-1	J-54
295	A-1	J-128
296	A-17	J-12
297	A-1	J-138
301	A-1	J-14
302	A-1	F-5
303	A-1	J-13
304	A-1	E-1
305	A-1	F-17
306	A-1	F-20
307	A-1	F-43
308	A-1	J-206
309	A-1	J-5
310	A-1	J-70
311	A-1	J-60
312	A-1	F-27
313	A-1	F-39
314	A-1	J-116
315	A-1	J-58
317	A-1	J-85
319	A-2	C-7
320	A-1	B-6
321	A-1	J-44
322	A-1	J-22
324	A-1	J-172
325	A-1	J-103
326	A-1	F-60
328	A-1	J-115
329	A-1	J-148
330	A-1	J-133
331	A-1	J-105
332	A-1	J-9

333	A-1	F-8
334	A-1	DC-5
335	A-1	J-194
336	A-1	J-192
337	A-1	C-24
338	A-1	J-113
339	A-1	B-8
344	A-1	F-22
345	A-2	J-234
346	A-12	J-6
348	A-1	F-21
349	A-1	J-29
350	A-1	J-100
351	A-1	B-23
352	A-1	B-10
353	A-1	D-10
354	A-1	J-186
355	A-1	J-25
357	A-1	B-13
358	A-24	J-232
360	A-1	J-151
361	A-1	F-26
362	A-1	J-91
363	A-1	F-32
364	A-1	J-88
365	A-1	J-93
366	A-1	F-16
367	A-1	F-50
368	A-1	D-5
369	A-1	J-141
370	A-1	J-90
371	A-1	J-79
372	A-1	J-209
373	A-1	J-21
374	A-16	J-238
375	A-1	J-71

376	A-1	J-187
377	A-5	J-237
378	A-1	D-3
380	A-1	J-99
381	A-1	B-24
383	A-1	B-12
384	A-1	F-48
385	A-1	J-83
387	A-1	J-168
388	A-1	F-29
389	A-1	J-27
391	A-1	F-9
392	A-1	J-52
394	A-22	J-170
395	A-1	C-20
397	A-1	J-199
398	A-1	J-77
400	A-1	J-183
401	A-1	F-4
402	A-1	J-149
403	A-1	C-22
405	A-1	J-33
406	A-6	B-24
407	A-3	C-7
408	A-1	J-81
410	A-1	F-31
411	A-13	J-191
412	A-1	B-19
413	A-1	J-131
414	A-1	J-50
417	A-1	F-65
418	A-1	J-223
419	A-1	J-216
420	A-1	G-1
421	A-1	C-18
422	A-1	J-20


423	A-1	B-16
424	A-1	F-42
425	A-1	J-28
426	A-1	C-11
427	A-1	J-124
428	A-1	C-1
429	A-1	J-218
430	A-1	J-123
431	A-1	J-225
432	A-1	F-14
433	A-1	C-9
434	A-1	J-159
435	A-1	J-41
436	A-1	F-24
437	A-1	J-75
438	A-1	E-10
439	A-1	J-164
440	A-1	J-215
441	A-1	D-19
442	A-1	J-165
443	A-1	J-166
444	A-1	E-6
445	A-1	J-97
446	A-1	J-121
447	A-1	J-51
448	A-1	J-69
449	A-1	J-94
450	A-1	J-193
451	A-1	J-31
452	A-1	J-108
453	A-1	D-1
454	A-1	J-47
455	A-1	J-73
456	A-1	J-137
457	A-1	J-155
458	A-1	C-4

459	A-1	J-53
461	A-1	J-150
463	A-1	J-202
464	A-3	C-9
465	A-1	E-4
466	A-1	J-2
467	A-1	J-86
468	A-20	J-184
469	A-12	J-132
470	A-1	J-160
473	A-21	J-89
474	A-1	J-201
475	A-1	J-158
477	A-1	J-63
478	A-1	B-11
479	A-4	J-230
480	A-23	J-175
481	A-1	J-188
483	A-1	C-21
484	A-1	D-14
B-26-I	A-1	B-26
B-27-I	A-1	B-27
C-27-I	A-1	C-27
D-12-I	A-1	D-12
D-13-I	A-1	D-13
D-15-I	A-1	D-15
D-16-I	A-1	D-16
D-17-I	A-1	D-17
DC-10-I	A-1	DC-10
DC-8-I	A-1	DC-8
DC-9-I	A-1	DC-9

[00683] Indoles


[00684] **Example 1:**

[00685] General Scheme:



[00686]

Specific example:



[00687]

188-I; 6-[(4-Oxo-1H-quinolin-3-yl)carbamoyl]-1H-indole-5-carboxylic acid

A mixture of 6-[(4-oxo-1H-quinolin-3-yl)carbamoyl]-1H-indole-5-carboxylic acid ethyl ester (**188**) (450 mg, 1.2 mmol) and 1N NaOH solution (5 mL) in THF (10 mL) was heated at 85 °C overnight. The reaction mixture was partitioned between EtOAc and water. The aqueous layer was acidified with 1N HCl solution to pH 5, and the precipitate was filtered, washed with water and air dried to yield 6-[(4-oxo-1H-quinolin-3-yl)carbamoyl]-1H-indole-5-carboxylic acid (**188-I**) (386 mg, 93 %). ¹H-NMR (400 MHz, DMSO-*d*₆) δ 12.92-12.75 (m, 2H), 11.33 (s, 1H), 8.84 (s, 1H), 8.71 (s, 1H), 8.30 (dd, *J* = 8.1, 0.9 Hz, 1H), 8.22 (s, 1H), 7.80-7.72 (m, 2H), 7.49 (t, *J* = 8.0 Hz, 1H), 7.41 (t, *J* = 2.7 Hz, 1H), 6.51 (m, 1H); HPLC ret. time 2.95 min, 10-99 % CH₃CN, 5 min run; ESI-MS 376.2 *m/z* (MH⁺).


[00688]

343; N-[5-(Isobutylcarbamoyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide

To a solution of 6-[(4-oxo-1H-quinolin-3-yl)carbamoyl]-1H-indole-5-carboxylic acid (**188-I**) (26 mg, 0.08 mmol), HATU (38 mg, 0.1 mmol) and DIEA (35 μL, 0.2 mmol) in DMF (1 mL) was added isobutylamine (7 mg, 0.1 mmol) and the reaction mixture was stirred at 65 °C

overnight. The resulting solution was filtered and purified by HPLC (10-99 % CH₃CN / H₂O) to yield the product, N-[5-(isobutylcarbamoyl)-1H-indol-6-yl]-4-oxo-1H-quinoline-3-carboxamide (**343**) (20 mg, 66 %). ¹H-NMR (400 MHz, DMSO-*d*₆) δ 12.66 (d, *J* = 7.4 Hz, 1H), 12.42 (s, 1H), 11.21 (s, 1H), 8.81 (d, *J* = 6.6 Hz, 1H), 8.47 (s, 1H), 8.36 (t, *J* = 5.6 Hz, 1H), 8.30 (d, *J* = 8.4 Hz, 1H), 7.79 (t, *J* = 7.9 Hz, 1H), 7.72-7.71 (m, 2H), 7.51 (t, *J* = 7.2 Hz, 1H), 7.38 (m, 1H), 6.48 (m, 1H), 3.10 (t, *J* = 6.2 Hz, 2H), 1.88 (m, 1H), 0.92 (d, *J* = 6.7 Hz, 6H); HPLC ret. time 2.73 min, 10-99 % CH₃CN, 5 min run; ESI-MS 403.3 *m/z* (MH⁺).

[00689] Another example:




[00690] **148; 4-Oxo-N-[5-(1-piperidylcarbonyl)-1H-indol-6-yl]-1H-quinoline-3-carboxamide**

4-Oxo-N-[5-(1-piperidylcarbonyl)-1H-indol-6-yl]-1H-quinoline-3-carboxamide (**148**) was synthesized following the general scheme above, coupling the acid (**188-I**) with piperidine. Overall yield (12 %). HPLC ret. time 2.79 min, 10-99 % CH₃CN, 5 min run; ESI-MS 415.5 *m/z* (MH⁺).

[00691] **Example 2:**

[00692] General scheme:



[00693] Specific example:

DEMANDES OU BREVETS VOLUMINEUX

**LA PRÉSENTE PARTIE DE CETTE DEMANDE OU CE BREVETS
COMPREND PLUS D'UN TOME.**

CECI EST LE TOME __1__ DE __2__

NOTE: Pour les tomes additionels, veuillez contacter le Bureau Canadien des Brevets.

JUMBO APPLICATIONS / PATENTS

**THIS SECTION OF THE APPLICATION / PATENT CONTAINS MORE
THAN ONE VOLUME.**


THIS IS VOLUME __1__ OF __2__

NOTE: For additional volumes please contact the Canadian Patent Office.


53568-20

CLAIMS:


1. A compound which is B-24

**B-24**


- 5 2. A process for the preparation of Compound B-24

**B-24**

comprising contacting 4-*tert*-butylaniline with N-bromosuccinimide to produce 2-bromo-4-*tert*-butylaniline


**2-bromo-4-*tert*-butylaniline**

contacting 2-bromo-4-*tert*-butylaniline with a mixture of KNO₃ and sulfuric acid to produce 2-bromo-4-*tert*-butyl-5-nitroaniline

**2-bromo-4-*tert*-butyl-5-nitroaniline**


53568-20

contacting 2-bromo-4-*tert*-butyl-5-nitroaniline with trimethylsilylacetylene, Pd(PPh₃)Cl₂, CuI and triethylamine to produce 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline




5 **4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline**

contacting 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline with CuI to produce 5-*tert*-butyl-6-nitro-1H-indole



5-*tert*-butyl-6-nitro-1H-indole

10 contacting 5-*tert*-butyl-6-nitro-1H-indole with Raney Ni in the presence of hydrogen gas to produce 6-amino-5-*tert*-butyl-1H-indole



6-amino-5-*tert*-butyl-1H-indole (B-24).

3. The process according to claim 2, wherein 4-*tert*-butylaniline is contacted with
15 N-bromosuccinimide in the presence of a solvent.

4. The process according to claim 3, wherein the solvent is DMF.

53568-20

5. The process according to claim 2, wherein 2-bromo-4-*tert*-butyl-5-nitroaniline is contacted with trimethylsilylacetylene, Pd(PPh₃)Cl₂, CuI and triethylamine in the presence of a solvent.
6. The process according to claim 5, wherein the solvent is a mixture of toluene
5 and water.
7. The process according to claim 6, wherein the ratio of toluene to water in the mixture is about 2:1.
8. The process according to claim 2, wherein 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline is contacted with CuI in the presence of a solvent.
- 10 9. The process according to claim 8, wherein the solvent is DMF.
10. The process according to claim 9, wherein 4-*tert*-butyl-5-nitro-2-((trimethylsilyl)ethynyl)aniline is contacted with CuI at a temperature from about 120°C to about 153°C.

