



(86) Date de dépôt PCT/PCT Filing Date: 1999/10/06
(87) Date publication PCT/PCT Publication Date: 2000/04/13
(45) Date de délivrance/Issue Date: 2012/10/02
(85) Entrée phase nationale/National Entry: 2001/04/05
(86) N° demande PCT/PCT Application No.: NL 1999/000618
(87) N° publication PCT/PCT Publication No.: 2000/020555
(30) Priorité/Priority: 1998/10/06 (EP PCT/EP98/06496)

(51) Cl.Int./Int.Cl. *C12N 15/80* (2006.01),
C12N 1/15 (2006.01), *C12N 15/55* (2006.01),
C12N 15/56 (2006.01), *C12N 15/57* (2006.01),
C12N 9/16 (2006.01), *C12N 9/24* (2006.01),
C12N 9/36 (2006.01), *C12N 9/38* (2006.01),
C12N 9/42 (2006.01), *C12N 9/58* (2006.01)

(72) Inventeurs/Inventors:
EMALFARB, MARK AARON, US;
BURLINGAME, RICHARD PAUL, US;
OLSON, PHILIP TERRY, US;
SINITSYN, ARKADY PANTELEIMONOVICH, RU;
...

(73) Propriétaire/Owner:

(54) Titre : SYSTÈME DE TRANSFORMATION DANS LE DOMAINE DES HOTES FONGIQUES FILAMENTEUX
(54) Title: TRANSFORMATION SYSTEM IN THE FIELD OF FILAMENTOUS FUNGAL HOSTS

(57) **Abrégé/Abstract:**

A novel transformation system in the field of filamentous fungal hosts for expressing and secreting heterologous proteins or polypeptides is described. The invention also covers a process for producing large amounts of polypeptide or protein in an economical manner. The system comprises a transformed or transfected fungal strain of the genus *Chrysosporium*, more particularly of *Chrysosporium lucknowense* and mutants or derivatives thereof. It also covers transformants containing *Chrysosporium* coding sequences, as well as expression-regulating sequences of *Chrysosporium* genes.

(72) **Inventeurs(suite)/Inventors(continued):** PARRICHE, MARTINE, FR; BOUSSON, JEAN CHRISTOPHE, FR;
PYNNONEN, CHRISTINE MARIE, US; PUNT, PETER JAN, NL; VAN ZEIJL, CORNELIA MARIA JOHANNA, NL

(73) **Propriétaires(suite)/Owners(continued):**DYADIC INTERNATIONAL (USA), INC., US

(74) **Agent:** MCCARTHY TETRAULT LLP

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/80, 15/55, 15/56, 9/42, 1/15, 9/58, 15/57, 9/16, 9/38, 9/24	A3	(11) International Publication Number: WO 00/20555 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/NL99/00618 (22) International Filing Date: 6 October 1999 (06.10.99) (30) Priority Data: PCT/EP98/06496 6 October 1998 (06.10.98) EP (71) Applicant (for all designated States except US): AARL, INC. [US/US]; Suite 404, 140 Intracoastal Pointe Drive, Jupiter, FL 33477 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): EMALFARB, Mark, Aaron [US/US]; 193 Spyglass Court, Jupiter, FL 33477 (US). BURLINGAME, Richard, Paul [US/US]; 808 North 9th Street, Manitowoc, WI 54220 (US). OLSON, Philip, Terry [US/US]; 3220 Waldo Boulevard, Manitowoc, WI 54220 (US). SINITSYN, Arkady Panteleimonovich [RU/RU]; 7-761 Dovzenko Street, Moscow, 119550 (RU). PARRICHE, Martine [FR/FR]; 3, rue Jean Gayral, F-31200 Toulouse (FR). BOUSSON, Jean, Christophe [FR/FR]; 19 A, rue de Cèdres, F-31400 Toulouse (FR). PYNNONEN, Christine, Marie [US/US]; 808 North 9th Street, Manitowoc, WI 54220 (US). PUNT, Peter, Jan [NL/NL]; Boekdrukkersgilde 1, NL-3994 XT Houten (NL).	VAN ZEIJL, Cornelia, Maria, Johanna [NL/NL]; Henri Dunantlaan 24, NL-3451 EP Vleuten-De Meern (NL). (74) Agent: DE BRUIJN, Leendert, C.; Nederlandsch Octrooi- bureau, Scheveningseweg 82, P.O. Box 29720, NL-2502 LS The Hague (NL). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 5 October 2000 (05.10.00)	
(54) Title: TRANSFORMATION SYSTEM IN THE FIELD OF FILAMENTOUS FUNGAL HOSTS: IN <i>CHRYSO Sporium</i> (57) Abstract A novel transformation system in the field of filamentous fungal hosts for expressing and secreting heterologous proteins or polypeptides is described. The invention also covers a process for producing large amounts of polypeptide or protein in an economical manner. The system comprises a transformed or transfected fungal strain of the genus <i>Chrysosporium</i> , more particularly of <i>Chrysosporium lucknowense</i> and mutants or derivatives thereof. It also covers transformants containing <i>Chrysosporium</i> coding sequences, as well as expression-regulating sequences of <i>Chrysosporium</i> genes.		

TRANSFORMATION SYSTEM IN THE FIELD OF FILAMENTOUS FUNGAL HOSTS

Summary of the invention

5 The subject invention pertains to a novel transformation system in the field of filamentous fungal hosts for expressing and secreting heterologous proteins or polypeptides. The invention also covers a process for producing large amounts of polypeptide in an economical manner. The system comprises a transformed or transfected fungal strain of the genus *Chrysosporium*, more particularly of *Chrysosporium lucknowense* and mutants or derivatives thereof. It also covers transformants containing *Chrysosporium* coding sequences. Novel mutant *Chrysosporium* strains are disclosed as
10 are novel enzymes derived therefrom.

Background to the invention.

A number of hosts for gene expression and methods of transformation have been disclosed in the prior art. Bacteria are often mentioned e.g. *Escherichia coli*. *E. coli* is however a micro-
15 organism incapable of secretion of a number of proteins or polypeptides and as such is undesirable as host cell for production of protein or polypeptide at the industrial level. An additional disadvantage for *E. coli*, which is valid also for bacteria in general, is that prokaryotes cannot provide additional modifications required for numerous eukaryotic proteins or polypeptides to be produced in an active form. Glycosylation of proteins and proper folding of proteins are examples of processing required to
20 ensure an active protein or polypeptide is produced. To ensure such processing one can sometimes use mammalian cells; however, the disadvantage of such cells is that they are often difficult to maintain and require expensive media. Such transformation systems are therefore not practical for production of proteins or polypeptides at the industrial level. They may be cost efficient for highly priced pharmaceutical compounds requiring relatively low amounts, but certainly not for industrial enzymes.

25 A number of fungal expression systems have been developed e.g. *Aspergillus niger*, *Aspergillus awamori*, *Aspergillus nidulans*, *Trichoderma reesei*. A number of others have been suggested but for various reasons have not found wide-spread acceptance or use. In general terms the ideal host must fulfil a large number of criteria:

- The ideal host must be readily fermented using inexpensive medium.
- 30 - The ideal host should use the medium efficiently.
- The ideal host must produce the polypeptide or protein in high yield, i.e. must exhibit high protein to biomass ratio.
- The ideal host should be capable of efficient secretion of the protein or polypeptide.
- The ideal host must enable ease of isolation and purification of the desired protein or polypeptide.
- 35 - The ideal host must process the desired protein or polypeptide such that it is produced in an active form not requiring additional activation or modification steps.

- The ideal host should be readily transformed.
- The ideal host should allow a wide range of expression regulatory elements to be used thus ensuring ease of application and versatility.
- The ideal host should allow use of easily selectable markers that are cheap to use.
- 5 - The ideal host should produce stable transformants.
- The ideal host should allow cultivation under conditions not detrimental to the expressed protein or polypeptide e.g. low viscosity, low shear.

Fungal systems that have not yet found widespread use are described e.g. in US Patent 5,578,463 by Berka et al suggesting *Neurospora*, *Podospora*, *Endothia*, *Mucor*, *Cochoibolus* and
 10 *Pyricularia* together with *Aspergillus* and *Trichoderma*. However only illustrations of transformation and expression are provided for *Aspergillus* and *Trichoderma* and no details are provided for any of the other suggested hosts.

1 WO 96/02563 and US patents 5,602,004, 5,604,129 and 5,695,985 to Novo Nordisk describe the drawbacks of *Aspergillus* and *Trichoderma* systems and suggests cultivation conditions
 15 for other fungi may be more suited to large scale protein production. The only examples provided for any transformed cultures are those of *Myceliophthora thermophila*, *Acremonium alabamense*, *Thielavia terrestris* and *Sporotrichum cellulophilum* strains. The *Sporotrichum* strain is reported to lyse and produce green pigment under fermentation conditions not leading to such results for the other strains. A non-sporulating mutant of *Thielavia terrestris* is described as being the organism of choice
 20 by virtue of its morphology. However it is also stated that the protoplasting efficiency of *Thielavia* and *Acremonium* (whereby the *Acremonium* strain used was the imperfect state of the *Thielavia* strain used) is low and that hygromycin was not useful as a selection marker. A large number of others are suggested as being potentially useful by virtue of their morphology but no transformation thereof is described. The suggested strains are *Corynascus*, *Thermoascus*, *Chaetomium*, *Ctenomyces*,
 25 *Scytalidium* and *Talaromyces*. The transformed hosts are mentioned as only producing low levels of the introduced *Humicola* xylanase with *Thielavia* producing the lowest amount; however, the information is ambiguous and could actually infer *Thielavia* was the best embodiment. The nomenclature of this reference is based on the ATCC names of Industrial Fungi of 1994. Thus it is apparent no high degree of heterologous expression was achieved and in fact no positive correlation
 30 could be derived between the postulated morphology and the degree of expression. If any correlation could be made, it was more likely to be negative. According to the 1996 ATCC fungal classification *Sporotrichum thermophilum* ATCC 20493 is a *Myceliophthora thermophila* strain. Currently the strain is still identified as *Myceliophthora thermophila*. The unpredictability of the art is apparent from these recent disclosures.

35 Also Allison et al (*Curr. Genetics* 21:225-229, 1992) described transformation of *Humicola grisea* var. *thermoidea* using the lithium acetate method and a *Humicola* enzyme-encoding sequence,

but no report of expression of heterologous protein from such a strain has been provided.

In 1997 a patent issued to Hawaii Biotechnology Group for transformed *Neurospora* for expression of mammalian peptide such as chymosin. The transformation of auxotrophic *Neurospora crassa* occurred with spheroplasts. Endogenous transcriptional regulatory regions were introduced and
5 cotransformation was carried out. Nothing is mentioned concerning other hosts and other transformation protocols. Nothing is apparent from the disclosure concerning the degree of expression. It is doubtful whether the degree of expression is high, as immunotechniques (which are useful for detecting small amounts of protein) are the only techniques used to illustrate the presence of the protein. No actual isolation of the protein is disclosed.

10 WO 97/26330 of Novo Nordisk suggests a method of obtaining mutants of filamentous fungal parent cells having an improved property for production of heterologous polypeptide. The method comprises first finding a specific altered morphology followed by assessing whether a transformant produces more heterologous polypeptide than the parent. The method is illustrated only for strains of *Fusarium* A3/5 and *Aspergillus oryzae*. The method is suggested to be applicable for
15 *Aspergillus*, *Trichoderma*, *Thielavia*, *Fusarium*, *Neurospora*, *Acremonium*, *Tolytlocadium*, *Humicola*, *Scytalidium*, *Myceliophthora* or *Mucor*. As stated above the unpredictability in the art and also the unpredictability of the method of the cited application do not provide a generally applicable teaching with a reasonable expectation of success.

20 Detailed description of the invention.

We now describe an alternative fungal expression system with the simplicity of use of the above-mentioned *Aspergilli* and *Trichoderma* fulfilling the above requirements. The new system has not been taught or suggested in the prior art. The new system according to the invention provides the additional advantages that transformation rates are higher than those for the frequently used
25 *Trichoderma reesei* system. In addition the culture conditions offer the additional bonus of being advantageous for the expressed polypeptide. Where reference is made in this specification and in the appending claims to "polypeptides" or "polypeptides of interest" as the products of the expression system of the invention, this term also comprise proteins, i.e polypeptides having a particular function and/or secondary and/or tertiary structure.

30 The pH of the culture medium can be neutral or alkaline thus no longer subjecting the produced protein or polypeptide to aggressive and potentially inactivating acid pH. It is also possible to culture at acid pH such as pH 4 for cases where the protein or polypeptide is better suited to an acidic environment. Suitably culture can occur at a pH between 4.0-10.0. A preference however exists for neutral to alkaline pH as the host strain exhibits better growth at such pH, e.g. between 6 and 9.
35 Growth at alkaline temperature which can be from pH 8 up and can even be as high as 10 is also a good alternative for some cases. Also the cultivation temperature of such host strains is advantageous

to the stability of some types of produced polypeptide. The cultivation temperature is suitably at a temperature of 25-43°C. A temperature in the range from 40°C down to 23°C or 30°C is also advantageously applied. Clearly such conditions are of particular interest for production of mammalian polypeptides. The selected temperature will depend on cost effectiveness of the cultivation and sensitivity of the polypeptide or cultivation strain. The conditions will be determined by the skilled person without undue burden on a case-by-case basis, as is common in the art.

It has also been ascertained that the biomass to viscosity relation and the amount of protein produced is exceedingly favourable for the host according to the invention. Comparisons have been carried out with *Trichoderma longibrachiatum* (formerly also known as *Trichoderma reesei*) and with *Aspergillus niger*. *Trichoderma longibrachiatum* gave 2.5-5 g/l biomass, *Aspergillus niger* gave 5-10 g/l biomass and the host according to the invention gave 0.5-1 g/l biomass under their respective optimised conditions. This thus offers 5-10 fold improvement over the commercially used strains. These commercial strains are strains which themselves are considered in the art to be high producers of proteins and they are successfully used for commercial protein production. They have been cultured under their optimal conditions developed and run viably in large-scale commercial fermenters. The same strains were used to illustrate enormous improvement in viscosity values for cultures of the host according to the invention. At the end of the fermentation process *Trichoderma longibrachiatum* gave a value of 200-600 cP (Centipoise), *Aspergillus niger* gave a value of 1500-2000 cP and the host according to the invention gave a value below 10 cP. This thus provides at least 20-200 fold improvement for viscosity values over the commercially used strains. A quite surprising further aspect was that the protein levels determined for the host cells according to the invention were much higher than for the commercial *Aspergilli* and *Trichoderma reesei* strains, even with the above mentioned surprisingly low biomass and viscosity levels. In summary an easy to use versatile improved transformation system and expression system with improved culturing conditions has hereby been introduced. The strains according to the invention produce surprisingly higher protein levels under these improved conditions and in addition they do such in a shorter fermenter time.

The subject invention is directed at mutant *Chrysosporium* strains comprising a nucleic acid sequence encoding a heterologous protein or polypeptide, said nucleic acid sequence being operably linked to an expression regulating region and optionally a secretion signal encoding sequence and/or a carrier protein encoding sequence. Preferably a recombinant strain according to the invention will secrete the polypeptide of interest. This will avoid the necessity of disrupting the cell in order to isolate the polypeptide of interest and also minimise the risk of degradation of the expressed product by other components of the host cell.

Chrysosporium can be defined by morphology consistent with that disclosed in Barnett and Hunter 1972, Illustrated Genera of Imperfect Fungi, 3rd Edition of Burgess Publishing Company. Other sources providing details concerning classification of fungi of the genus *Chrysosporium* are

known e.g. Sutton Classification (Van Oorschot, C.A.N. (1980) "A revision of *Chrysosporium* and allied genera" in Studies in Mycology No. 20 of the CBS in Baarn The Netherlands p 1-36). CBS is one of the depository institutes of the Budapest Treaty. According to these teachings the genus *Chrysosporium* falls within the family *Moniliaceae* which belongs to the order *Hyphomycetales*. The

5 criteria that can be used are the following:

1. Signs of *Hyphomycetales* order:

Conidia are produced directly on mycelium, on separate sporogenous cells or on distinct conidiophores.

2. Signs of *Moniliaceae* family:

10 Both conidia and conidiophores (if present) are hyaline or brightly coloured; conidiophores are single or in loose clusters.

3. Signs of *Chrysosporium* Corda 1833 genus:

Colonies are usually spreading, white, sometimes cream-coloured, pale brown or yellow, felty and/or powdery. Hyphae are mostly hyaline and smooth-walled, with irregular, more or less

15 orthotopic branching. Fertile hyphae exhibit little or no differentiation. Conidia are terminal and lateral, thallic, borne all over the hyphae, sessile or on short protrusions or side branches, subhyaline or pale yellow, thin- or thick-walled, subglobose, clavate, pyriform, obovoid, 1-celled, rarely 2-celled, truncate. Intercalary conidia are sometimes present, are solitary, occasionally catenate, subhyaline or pale yellow, broader than the supporting hyphae, normally 1-celled, truncate at both

20 ends. Chlamydospores are occasionally present.

Another source providing information on fungal nomenclature is ATCC (US) at their website. CBS also has a website providing relevant information. VKM in Moscow is also a reliable source of such information. All these institutions can provide teaching on the distinguishing characteristics of a *Chrysosporium*.

25 Strains defined as being of *Myceliophthora thermophila* are not considered to define *Chrysosporium* strains according to the definition of the invention. In the past there has been considerable confusion over the nomenclature of some *Myceliophthora* strains. Preferably the *Chrysosporium* according to the invention are those which are clearly distinguishable as such and cannot be confused with *Myceliophthora*, *Sporotrichum* or *Phanerochaete chrysosporium*.

30 The following strains are defined as *Chrysosporium* but the definition of *Chrysosporium* is not limited to these strains: *C. botryoides*, *C. carmichaelii*, *C. crassitunicatum*, *C. europae*, *C. evolceannui*, *C. farinicola*, *C. fastidium*, *C. filiforme*, *C. georgiae*, *C. globiferum*, *C. globiferum* var. *articulatum*, *C. globiferum* var. *niveum*, *C. hirundo*, *C. hispanicum*, *C. holmii*, *C. indicum*, *C. inops*, *C.*

keratinophilum, *C. kreiselii*, *C. kuzurovianum*, *C. lignorum*, *C. lobatum*, *C. lucknowense*, *C. lucknowense* Garg 27K, *C. medium*, *C. medium* var. *spissescens*, *C. mephiticum*, *C. merdarium*, *C. merdarium* var. *roseum*, *C. minor*, *C. pannicola*, *C. parvum*, *C. parvum* var. *crescens*, *C. pilosum*, *C. pseudomerdarium*, *C. pyriformis*, *C. queenslandicum*, *C. sigleri*, *C. sulfureum*, *C. synchronum*, *C. tropicum*, *C. undulatum*, *C. vallenarense*, *C. vespertilium*, *C. zonatum*.

C. lucknowense forms one of the species of *Chrysosporium* that have raised particular interest as it has provided a natural high producer of cellulase proteins (WO 98/15633 and related US 5,811,381). The characteristics of this *Chrysosporium lucknowense* are:

Colonies attain 55 mm diameter on Sabouraud glucose agar in 14 days, are cream-coloured, felty and fluffy; dense and 3-5 mm high; margins are defined, regular, and fimbriate; reverse pale yellow to cream-coloured. Hyphae are hyaline, smooth- and thin-walled, little branched. Aerial hyphae are mostly fertile and closely septate, about 1-3.5 mm wide. Submerged hyphae are infertile, about 1-4.5 mm wide, with the thinner hyphae often being contorted. Conidia are terminal and lateral, mostly sessile or on short, frequently conical protrusions or short side branches. Conidia are solitary but in close proximity to one another, 1-4 conidia developing on one hyphal cell, subhyaline, fairly thin- and smooth-walled, mostly subglobose, also clavate obovoid, 1-celled, 2.5 x 11 x 1.5-6 mm, with broad basal scars (1-2 mm). Intercalary conidia are absent. Chlamydospores are absent. ATCC 44006, CBS 251.72, CBS 143.77 and CBS 272.77 are examples of *Chrysosporium lucknowense* strains and other examples are provided in WO 98/15633 and US 5,811,381.

A further strain was isolated from this species with an even higher production capacity for cellulases. This strain is called C1 by its internal notation and was deposited with the International Depository of the All Russian Collection of micro-organisms of the Russian Academy of Sciences Bakrushina Street 8, Moscow, Russia 113184 on August 29, 1996, as a deposit according to the Budapest Treaty and was assigned Accession Number VKM F-3500D. It is called *Chrysosporium lucknowense* Garg 27K. The characteristics of the C1 strain are as follows:

Colonies grow to about 55-66 mm diameter in size on potato-dextrose agar in about 7 days; are white-cream-coloured, felty, 2-3 mm high at the center; margins are defined, regular, fimbriate; reverse pale, cream-coloured. Hyphae are hyaline, smooth- and thin-walled, little branched. Aerial hyphae are fertile, septate, 2-3 mm wide. Submerged hyphae are infertile. Conidia are terminal and lateral; sessile or on short side branches; absent; solitary, but in close proximity to one another, hyaline, thin- and smooth-walled, subglobose, clavate or obovoid, 1-celled, 4-10 mm. Chlamydospores are absent. Intercalary conidia are absent.

The method of isolation of the C1 strain is described in WO 98/15633 and US 5,811,381. Strains exhibiting such morphology are included within the definition of *Chrysosporium* according to the invention. Also included within the definition of *Chrysosporium* are strains derived from *Chrysosporium* predecessors including those that have mutated somewhat either naturally or by induced

mutagenesis. In particular the invention covers mutants of *Chrysosporium* obtained by induced mutagenesis, especially by a combination of irradiation and chemical mutagenesis.

For example strain C1 was mutagenised by subjecting it to ultraviolet light to generate strain UV13-6. This strain was subsequently further mutated with N-methyl-N'-nitro-N-nitrosoguanidine to generate strain NG7C-19. The latter strain in turn was subjected to mutation by ultraviolet light resulting in strain UV18-25. During this mutation process the morphological characteristics have varied somewhat in culture in liquid or on plates as well as under the microscope. With each successive mutagenesis the cultures showed less of the fluffy and felty appearance on plates that are described as being characteristic of *Chrysosporium*, until the colonies attained a flat and matted appearance. A brown pigment observed with the wild type strain in some media was also less prevalent in mutant strains. In liquid culture the mutant UV18-25 was noticeably less viscous than the wild type strain C1 and the mutants UV13-6 and NG7C-19. While all strains maintained the gross microscopic characteristics of *Chrysosporium*, the mycelia became narrower with each successive mutation and with UV18-25 distinct fragmentation of the mycelia could be observed. This mycelial fragmentation is likely to be the cause of the lower viscosity associated with cultures of UV18-25. The ability of the strains to sporulate decreased with each mutagenic step. The above illustrates that for a strain to belong to the genus *Chrysosporium* there is some leeway from the above morphological definition. At each mutation step production of cellulase and extracellular proteins has in addition also increased, while several mutations resulted in decrease of protease expression. Criteria with which fungal taxonomy can be determined are available from CBS, VKMF and ATCC for example.

In particular the anamorph form of *Chrysosporium* has been found to be suited for the production application according to the invention. The metabolism of the anamorph renders it extremely suitable for a high degree of expression. A teleomorph should also be suitably as the genetic make-up of the anamorphs and teleomorphs is identical. The difference between anamorph and teleomorph is that one is the asexual state and the other is the sexual state. The two states exhibit different morphology under certain conditions.

It is preferable to use non-toxic *Chrysosporium* strains of which a number are known in the art as this will reduce risks to the environment upon large scale production and simplify production procedures with the concomitant reduction in costs.

An expression-regulating region is a DNA sequence recognised by the host *Chrysosporium* strain for expression. It comprises a promoter sequence operably linked to a nucleic acid sequence encoding the polypeptide to be expressed. The promoter is linked such that the positioning vis-à-vis the initiation codon of the sequence to be expressed allows expression. The promoter sequence can be constitutive or inducible. Any expression regulating sequence or combination thereof capable of permitting expression of a polypeptide from a *Chrysosporium* strain is envisaged. The expression regulating sequence is suitably a fungal expression-regulating region e.g. an ascomycete regulating

region. Suitably the fungal expression regulating region is a regulating region from any of the following genera of fungi: *Aspergillus*, *Trichoderma*, *Chrysosporium*, *Hansenula*, *Mucor*, *Pichia*, *Neurospora*, *Tolypocladium*, *Rhizomucor*, *Fusarium*, *Penicillium*, *Saccharomyces*, *Talaromyces* or alternative sexual forms thereof like *Emericella*, *Hypocrea* e.g. the cellobiohydrolase promoter from *Trichoderma*, glucoamylase promoter from *Aspergillus*, glyceraldehyde phosphate dehydrogenase promoter from *Aspergillus*, alcohol dehydrogenase A and alcohol dehydrogenase R promoter of *Aspergillus*, TKA amylase promoter from *Aspergillus*, phosphoglycerate and cross-pathway control promoters of *Neurospora*, aspartic proteinase promoter of *Rhizomucor miehei*, lipase promoter of *Rhizomucor miehei* and beta-galactosidase promoter of *Penicillium canescens*. An expression regulating sequence from the same genus as the host strain is extremely suitable, as it is most likely to be specifically adapted to the specific host. Thus preferably the expression regulating sequence is one from a *Chrysosporium* strain.

We have found particular strains of *Chrysosporium* to express proteins in extremely large amounts and natural expression regulating sequences from these strains are of particular interest. These strains are internally designated as *Chrysosporium* strain C1, strain UV13-6, strain NG7C-19 and strain UV18-25. They have been deposited in accordance with the Budapest Treaty with the All Russian Collection (VKM) depository institute in Moscow. Wild type C1 strain was deposited in accordance with the Budapest Treaty with the number VKM F-3500 D, deposit date 29-08-1996, C1 UV13-6 mutant was deposited with number VKM F-3632 D, and deposit date 02-09-1998, C1 NG7c-19 mutant was deposited with number VKM F-3633 D and deposit date 02-09-1998 and C1 UV18-25 mutant was deposited with number VKM F-3631 D and deposit date 02-09-1998.

Preferably an expression-regulating region enabling high expression in the selected host is applied. This can also be a high expression-regulating region derived from a heterologous host, such as are well known in the art. Specific examples of proteins known to be expressed in large quantities and thus providing suitable expression regulating sequences for the invention are without being limited thereto hydrophobin, protease, amylase, xylanase, pectinase, esterase, beta-galactosidase, cellulase (e.g. endo-glucanase, cellobiohydrolase) and polygalacturonase. The high production has been ascertained in both solid state and submerged fermentation conditions. Assays for assessing the presence or production of such proteins are well known in the art. The catalogues of Sigma and Megazyme for example provide numerous examples. Megazyme is located at Bray Business Park, Bray, County Wicklow in Ireland. Sigma Aldrich has many affiliates world wide e.g. USA P.O. Box 14508 St. Louis Missouri. For cellulase we refer to commercially available assays such as CMCase assays, endovisco-metric assays, Avicelase assays, beta-glucanase assays, RBBCMCase assays, Cellazyme C assays. Alternatives are well known to a person skilled in the art and can be found from general literature concerning the subject and such information is considered incorporated herein by reference. By way of example we refer to "Methods in Enzymology Volume 1, 1955 right through to

volumes 297-299 of 1998. Suitably a *Chrysosporium* promoter sequence is applied to ensure good recognition thereof by the host.

We have found that heterologous expression-regulating sequences work as efficiently in *Chrysosporium* as native *Chrysosporium* sequences. This allows well known constructs and vectors to be used in transformation of *Chrysosporium* as well as offering numerous other possibilities for constructing vectors enabling good rates of expression in this novel expression and secretion host. For example standard *Aspergillus* transformation techniques can be used as described for example by Christiansen et al in Bio/Technol. 6:1419-1422 (1988). Other documents providing details of *Aspergillus* transformation vectors, e.g. US patents 4,816,405, 5,198,345, 5,503,991, 5,364,770 and 5,578,463, EP-B-215.594 (also for *Trichoderma*) and their contents are incorporated by reference. As extremely high expression rates for cellulase have been ascertained for *Chrysosporium* strains, the expression regulating regions of such proteins are particularly preferred. We refer for specific examples to the previously mentioned deposited *Chrysosporium* strains.

A nucleic acid construct comprising a nucleic acid expression regulatory region from *Chrysosporium*, preferably from *Chrysosporium lucknowense* or a derivative thereof forms a separate embodiment of the invention as does the mutant *Chrysosporium* strain comprising such operably linked to a polypeptide to be expressed. Suitably such a nucleic acid construct will be an expression regulatory region from *Chrysosporium* associated with cellulase or xylanase expression, preferably cellobiohydrolase expression, more specifically expression of a 55 kDa cellobiohydrolase. The *Chrysosporium* promoter sequences of an endoglucanase of 25 kDa (C1-EG5) and of an endoglucanase of 43 kDa (C1-EG6), wherein the molecular weights are determined according to SDS PAGE (with the molecular weights according to amino acid sequence data being 21.9 kDa and 39.5 kDa), are provided by way of example. Thus, the *Chrysosporium* promoter sequences of hydrophobin, protease, amylase, xylanase, esterase, pectinase, beta-galactosidase, cellulase (e.g. endoglucanase, cellobiohydrolase) and polygalacturonase are considered to also fall within the scope of the invention. Any of the promoters or regulatory regions of expression of enzymes disclosed in Table A or B can be suitably employed. The nucleic acid sequence according to the invention can suitably be obtained from a *Chrysosporium* strain according to the invention, such strain being defined elsewhere in the description. The manner in which promoter sequences can be determined are numerous and well known in the art. Nuclease deletion experiments of the region upstream of the ATG codon at the beginning of the relevant gene will provide such sequence. Also for example analysis of consensus sequences can lead to finding a gene of interest. Using hybridisation and amplification techniques one skilled in the art can readily arrive at the corresponding promoter sequences.

The promoter sequences of C1 endoglucanases were identified in this manner, by cloning the corresponding genes, and are given in SEQ ID No.'s 2 (EG5) and 1 (EG6), respectively. Other preferred promoters according to the invention are the 55 kDa cellobiohydrolase (CBH1) promoter

-10-

and the 30kDa xylanase (XylF) promoters, as the enzymes are expressed at high level by their own promoters. The corresponding promoter sequences can be identified in a straightforward manner by cloning as described below for the endoglucanase promoters, using the partial sequence information given in SEQ ID No. 4 (for CBH1) and SEQ ID No. 5 (for XylF), respectively. The promoters of the carbohydrate-degrading enzymes of *Chrysosporium*, especially C1 promoters, can advantageously be used for expressing desired polypeptides in a host organism, especially a fungal or other microbial host organism. Promoter sequences having at least 60%, preferably at least 70%, most preferably at least 80% nucleotide sequence identity with the sequence given in SEQ ID No's 1 and 2, or with the sequences found for other *Chrysosporium* genes, are part of the present invention.

For particular embodiments of the recombinant strain and the nucleic acid sequence according to the invention we also refer to the examples. We also refer for the recombinant strains to prior art describing high expression promoter sequences in particular those providing high expression in fungi e.g. such as are disclosed for *Aspergillus* and *Trichoderma*. The prior art provides a number of expression regulating regions for use in *Aspergillus* e.g. US 5,252,726 of Novo and US 5,705,358 of Unilever.

The hydrophobin gene is a fungal gene that is highly expressed. It is thus suggested that the promoter sequence of a hydrophobin gene, preferably from *Chrysosporium*, may be suitably applied as expression regulating sequence in a suitable embodiment of the invention. *Trichoderma reesei* and *Trichoderma harzianum* gene sequences for hydrophobin have been disclosed for example in the prior art as well as a gene sequence for *Aspergillus fumigatus* and *Aspergillus nidulans* and the relevant sequence information is hereby incorporated by reference (Munoz et al, *Curr. Genet.* 1997, 32(3);225-230; Nakari-Setälä T. et al, *Eur. J. Biochem.* 1996 15:235 (1-2):248-255, M. Parta et al, *Infect. Immun.* 1994 62 (10) : 4389-4395 and Stringer M.A. et al. *Mol. Microbiol.* 1995 16(1) :33-44). Using this sequence information a person skilled in the art can obtain the expression regulating sequences of *Chrysosporium* hydrophobin genes without undue experimentation following standard techniques as suggested already above. A recombinant *Chrysosporium* strain according to the invention can comprise a hydrophobin-regulating region operably linked to the sequence encoding the polypeptide of interest.

An expression regulating sequence can also additionally comprise an enhancer or silencer. There are also well known in the prior art and are usually located some distance away from the promoter. The expression regulating sequences can also comprise promoters with activator binding sites and repressor binding sites. In some cases such sites may also be modified to eliminate this type of regulation. Filamentous fungal promoters in which creA sites are present have been described. Such creA sites can be mutated to ensure the glucose repression normally resulting from the presence of the non-mutated sites is eliminated. Gist-Brocades' WO 94-13820 illustrates this principle. Use of such a promoter enables production of the polypeptide encoded by the nucleic acid sequence regulated

by the promoter in the presence of glucose. The same principle is also apparent from WO 97/09438. These promoters can be used either with or without their creA sites. Mutants in which the creA sites have been mutated can be used as expression regulating sequences in a recombinant strain according to the invention and the nucleic acid sequence it regulates can then be expressed in the presence of glucose. Such *Chrysosporium* promoters ensure derepression in an analogous manner to that illustrated in WO 97/09438. The identity of creA sites is known from the prior art. Alternatively, it is possible to apply a promoter with CreA binding sites that have not been mutated in a host strain with a mutation elsewhere in the repression system e.g. in the creA gene itself, so that the strain can, notwithstanding the presence of creA binding sites, produce the protein or polypeptide in the presence of glucose.

Terminator sequences are also expression-regulating sequences and these are operably linked to the 3' terminus of the sequence to be expressed. Any fungal terminator is likely to be functional in the host *Chrysosporium* strain according to the invention. Examples are *A. nidulans* trpC terminator (1), *A. niger* alpha-glucosidase terminator (2), *A. niger* glucoamylase terminator (3), *Mucor miehei* carboxyl protease terminator (US 5,578,463) and the *Trichoderma reesei* cellobiohydrolase terminator. Naturally *Chrysosporium* terminator sequences will function in *Chrysosporium* and are suitable e.g. EG6 terminator.

A suitable recombinant *Chrysosporium* strain according to the invention has the nucleic acid sequence to be expressed operably linked to a sequence encoding the amino acid sequence defined as signal sequence. A signal sequence is an amino acid sequence which when operably linked to the amino acid sequence of the expressed polypeptide allows secretion thereof from the host fungus. Such a signal sequence may be one normally associated with the heterologous polypeptide or may be one native to the host. It can also be foreign to both host and the polypeptide. The nucleic acid sequence encoding the signal sequence must be positioned in frame to permit translation of the signal sequence and the heterologous polypeptide. Any signal sequence capable of permitting secretion of a polypeptide from a *Chrysosporium* strain is envisaged. Such a signal sequence is suitably a fungal signal sequence, preferably an ascomycete signal sequence.

Suitable examples of signal sequences can be derived from yeasts in general or any of the following specific genera of fungi: *Aspergillus*, *Trichoderma*, *Chrysosporium*, *Pichia*, *Neurospora*, *Rhizomucor*, *Hansenula*, *Humicola*, *Mucor*, *Tolypocladium*, *Fusarium*, *Penicillium*, *Saccharomyces*, *Talaromyces* or alternative sexual forms thereof like *Emericella*, *Hypocrea*. Signal sequences that are particularly useful are often natively associated with the following proteins a cellobiohydrolase, an endoglucanase, a beta-galactosidase, a xylanase, a pectinase, an esterase, a hydrophobin, a protease or an amylase. Examples include amylase or glucoamylase of *Aspergillus* or *Humicola* (4), TAKA amylase of *Aspergillus oryzae*, alpha-amylase of *Aspergillus niger*, carboxyl peptidase of *Mucor* (US 5,578,463), a lipase or proteinase from *Rhizomucor miehei*, cellobiohydrolase of *Trichoderma* (5),

beta-galactosidase of *Penicillium canescens* and alpha mating factor of *Saccharomyces*.

Alternatively the signal sequence can be from an amylase or subtilisin gene of a strain of *Bacillus*. A signal sequence from the same genus as the host strain is extremely suitable as it is most likely to be specifically adapted to the specific host thus preferably the signal sequence is a signal sequence of *Chrysosporium*. We have found particular strains of *Chrysosporium* to excrete proteins in extremely large amounts and naturally signal sequences from these strains are of particular interest. These strains are internally designated as *Chrysosporium* strain C1, strain UV13-6, strain NG7C-19 and strain UV18-25. They have been deposited in accordance with the Budapest Treaty as described elsewhere in this description. Signal sequences from filamentous fungi, yeast and bacteria are useful. Signal sequences of non-fungal origin are also considered useful, particularly bacterial, plant and mammalian.

A recombinant *Chrysosporium* strain according to any of the embodiments of the invention can further comprise a selectable marker. Such a selectable marker will permit easy selection of transformed or transfected cells. A selectable marker often encodes a gene product providing a specific type of resistance foreign to the non-transformed strain. This can be resistance to heavy metals, antibiotics and biocides in general. Prototrophy is also a useful selectable marker of the non-antibiotic variety. Non-antibiotic selectable markers can be preferred where the protein or polypeptide of interest is to be used in food or pharmaceuticals with a view to speedier or less complicated regulatory approval of such a product. Very often the GRAS indication is used for such markers. A number of such markers are available to the person skilled in the art. The FDA e.g. provides a list of such. Most commonly used are selectable markers selected from the group conferring resistance to a drug or relieving a nutritional defect e.g the group comprising amdS (acetamidase), hph (hygromycin phosphotransferase), pyrG (orotidine-5'-phosphate decarboxylase), trpC (anthranilate synthase), argB (ornithine carbamoyltransferase), sC (sulphate adenylyltransferase), bar (phosphinothricin acetyltransferase), glufosinate resistance, niaD (nitrate reductase), a bleomycin resistance gene, more specifically Sh ble, sulfonyleurea resistance e.g. acetolactate synthase mutation ilv1. Selection can also be carried out by virtue of cotransformation where the selection marker is on a separate vector or where the selection marker is on the same nucleic acid fragment as the polypeptide-encoding sequence for the polypeptide of interest.

As used herein the term heterologous polypeptide is a protein or polypeptide not normally expressed and secreted by the *Chrysosporium* host strain used for expression according to the invention. The polypeptide can be of plant or animal (vertebrate or invertebrate) origin e.g. mammalian, fish, insect, or micro-organism origin, with the proviso it does not occur in the host strain. A mammal can include a human. A micro-organism comprises viruses, bacteria, archaeobacteria and fungi i.e. filamentous fungi and yeasts. Bergey's Manual for Bacterial Determinology provides adequate lists of bacteria and archaeobacteria. For pharmaceutical purposes quite often a preference

will exist for human proteins thus a recombinant host according to the invention forming a preferred embodiment will be a host wherein the polypeptide is of human origin. For purposes such as food production suitably the heterologous polypeptide will be of animal, plant or algal origin. Such embodiments are therefore also considered suitable examples of the invention. Alternative
5 embodiments that are useful also include a heterologous polypeptide of any of bacterial, yeast, viral, archaeobacterial and fungal origin. Fungal origin is most preferred.

A suitable embodiment of the invention will comprise a heterologous nucleic acid sequence with adapted codon usage. Such a sequence encodes the native amino acid sequence of the host from which it is derived, but has a different nucleic acid sequence, i.e. a nucleic acid sequence in which
10 certain codons have been replaced by other codons encoding the same amino acid but which are more readily used by the host strain being used for expression. This can lead to better expression of the heterologous nucleic acid sequence. This is common practice to a person skilled in the art. This adapted codon usage can be carried out on the basis of known codon usage of fungal vis-à-vis non-fungal codon usage. It can also be even more specifically adapted to codon usage of *Chrysosporium*
15 itself. The similarities are such that codon usage as observed in *Trichoderma*, *Humicola* and *Aspergillus* should enable exchange of sequences of such organisms without adaptation of codon usage. Details are available to the skilled person concerning the codon usage of these fungi and are incorporated herein by reference.

The invention is not restricted to the above mentioned recombinant *Chrysosporium* strains,
20 but also covers a recombinant *Chrysosporium* strain comprising a nucleic acid sequence encoding a homologous protein for a *Chrysosporium* strain, said nucleic acid sequence being operably linked to an expression-regulating region and said recombinant strain expressing more of said protein than the corresponding non-recombinant strain under the same conditions. In the case of homologous polypeptide of interest such is preferably a neutral or alkaline enzyme like a hydrolase, a protease or a
25 carbohydrate degrading enzyme as already described elsewhere. The polypeptide may also be acidic. Preferably the recombinant strain will express the polypeptide in greater amounts than the non-recombinant strain. All comments mentioned vis-à-vis the heterologous polypeptide are also valid (mutatis mutandis) for the homologous polypeptide cellulase.

Thus the invention also covers genetically engineered *Chrysosporium* strains wherein the
30 sequence that is introduced can be of *Chrysosporium* origin. Such a strain can, however, be distinguished from natively occurring strains by virtue of for example heterologous sequences being present in the nucleic acid sequence used to transform or transfect the *Chrysosporium*, by virtue of the fact that multiple copies of the sequence encoding the polypeptide of interest are present or by virtue of the fact that these are expressed in an amount exceeding that of the non-engineered strain under
35 identical conditions or by virtue of the fact that expression occurs under normally non-expressing conditions. The latter can be the case if an inducible promoter regulates the sequence of interest

contrary to the non-recombinant situation or if another factor induces the expression than is the case in the non-engineered strain. The invention as defined in the preceding embodiments is not intended to cover naturally occurring *Chrysosporium* strains. The invention is directed at strains derived through engineering either using classical genetic technologies or genetic engineering methodologies.

- 5 All the recombinant strains of the invention can comprise a nucleic acid sequence encoding a heterologous protein selected from carbohydrate-degrading enzymes (cellulases, xylanases, mannanases, mannosidases, pectinases, amylases, e.g. glucoamylases, α -amylases, alpha- and beta-galactosidases, α - and β -glucosidases, β -glucanases, chitinases, chitanases), proteases (endoproteases, amino-proteases, amino-and carboxy-peptidases), other hydrolases (lipases, esterases, phytases),
 10 oxidoreductases (catalases, glucose-oxidases) and transferases (transglycosylases, transglutaminases, isomerases and invertases).

Table A: pH range where enzymes retain activity and/or stability

Sample	pH range retaining > 50% enzymatic activity			pH range retaining > 70% enzymatic activity			Stability (20h, 50°C) % from max
	CMC ase	RBB-CMC ase	Other sub-strates	CMC ase	RBB-CMC ase	Other sub-strates	pH 7.5/8
30 Kd protease (alkaline) 30 kD	-	-	12.5	-	-	12.0	-
Xyl (alkaline)	-	-	10.0	-	-	8.5	80
51 kD Xyl	-	-	8.0	-	-	7.5	-
60 kD Xyl	-	-	9.5	-	-	9.0	85
45 kD endo	7.0	8.0	-	6.5	7.0	-	75
55 kD endo	8.0	8.0	-	7.0	7.0	-	55
25 kD(21.8 kD*)endo	7.5	10.0	-	6.5	9.0	-	80
43 kD(39.6 kD*)endo	8.0	8.0	-	7.2	7.2	-	-
45 kD α , β -Gal/ β -Gluc	-	-	6.8	-	-	5.7	-
48 kD CBH with β -Gluc traces	5.2	7.5	8.0	5.0	6.8	-	-
55 kD CBH	8.0	9.0	-	7.4	8.5	-	70
65 kD PGU	-	-	8.0	-	-	7.3	-
90 kD protease	-	-	9.0	-	-	9.0	-
100 kD esterase	-	-	9.0	-	-	9.0	-

- 15 * molecular weights (by MALDI)

Note: * all other molecular weights by SDS PAGE
 * enzymes were taken in equal protein contents
 * xyl = xylanase
 * endo = endoglucanase
 * gal = galactosidase
 * gluc = glucosidase
 * CBN = cellbiohydrolase
 * PGU = polygalacturonase

Table B. Activities of enzymes isolated from ultrafiltrate from 18-25 strain toward different substrates (pH 5), units/mg protein

Sample	pI	CMC	RBB-CMC	CMC-41	FP	CMC (visc)	b-Glu can	pNP-a-G	pNP-b-G	Cellobiose	Avicel	MUF-cellobioside	MUF-lactoside	MUF-xyloside	Lactose	Xylan	Polygalacturonic acid	MUF-glucoside	Galactomannan	pNP-a-galactoside	pNP-b-galactoside	Dyed casein**	pNP butyrate
		50oC	40oC	40oC	50oC	40oC	50oC	40oC	40oC	40oC	40oC	40oC	40oC	40oC	40oC	50oC	50oC	40oC	40oC	40oC	40oC	50oC	60oC
30 kD protease	8.9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0.4	0
30 kD Xyl	9.1	0.1	2	0.1	0.16	0.1	0	0	0	-	0	0	0	0	-	25	0	0	0	0	-	0	0
51 kD Xyl	8.7	0.1	4.2	-	0.19	-	0	0	0	-	0	0	0	0	-	19	0	0	0	0	-	0	0
60 kD Xyl	4.7	0	-	-	0	-	0	0	0	-	0	0	0.02	0.04	-	163	0	0	0	0	0	0	0
45 kD endo	6	51	86	7.6	0.2	47	36	-	0	-	0.5	0	0	0	-	1	-	0	1.8	0	-	0	0
55 kD endo	4.9	47	94	7.7	0.3	39	25	-	0	-	0.5	0	0	0	-	0	-	0	0.4	0	-	0	0
25 kD (21.8 kD*) endo	4.1	19	15	3.9	0.3	11	3.8	-	0	0	0.05	0	0	-	0	0.03	0	-	0	0	0	0	0
43 kD (39.6 kD*) endo	4.2	0.43	0.2	0.1	0	0.2	0.2	-	0	0	0	0	0	-	0	0	0	-	0	0	0	0	0
45 kD a,b-Gal/b-Gluc	4.2	0	0	0	0	0.01	0.01	0	0.4	0.06	0	0	0	-	0.01	0	0.1	0.1	0.2	0.2	0.3	0	1.7
46 kD CBH with b-Gluc traces+ glucono-d-lactone	4.4	0.67	1.3	1.2	0.4	0.8	0.77	0	1.7	0.08	0	0.2	0.36	-	0	0	0.1	0.4	0	0	0	0	2.3
									0				0.36										
55 kD CBH with b-Gluc traces + glucono-d-lactone	4.4	0.7	0.16	0.27	0.4	0.1	0.1	-	0.05	0.08	0.46	0.2	0.7	-	0	0.1	0	-	0	0	0	0	0
												0.14	0.6										
65 kD PGU	4.4	0	0	0	0	0	0	-	0	0	0	0	0	-	0	0	1	-	0	0	0	0	0
90 kD protease	4.2	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	0.01	-
100 kD esterase	4.5	0	0	0	0	0	0	-	0	0	0	0	0	-	0	0	0	0	0	0	0	0	0.8

*molecular weights (by MALDI)

** activity toward dyed casein was expressed in arbitrary units/mg

The most interesting products to be produced according to invention are cellulases, xylanases, pectinases, lipases and proteases, wherein cellulases and xylanases cleave beta-1,4-bonds, and cellulases comprise endoglucanases, cellobiohydrolases and beta-glucosidases. These proteins are extremely useful in various industrial processes known in the art. Specifically for cellulases we refer
5 e.g. to WO 98/15633 describing cellobiohydrolases and endoglucanases of use. The contents of said application are hereby incorporated by reference. We also refer to Tables A and B providing further details of interesting *Chrysosporium* proteins.

It was found according to the invention, that *Chrysosporium* mutants can be made that have reduced expression of protease, thus making them even more suitable for the production of
10 proteinaceous products, especially if the proteinaceous product is sensitive to protease activity. Thus the invention also involves a mutant *Chrysosporium* strain which produces less protease than non-mutant *Chrysosporium* strain, for example less than *C. lucknowense* strain C1 (VKM F-3500 D). In particular the protease activity of such strains is less than half the amount, more in particular less than 30% of the amount produced by C1 strain. The decreased protease activity can be
15 measured by known methods, such as by measuring the halo formed on skim milk plates or BSA degradation.

An embodiment of the invention that is of particular interest is a recombinant *Chrysosporium* according to the invention wherein the nucleic acid sequence encoding the polypeptide of interest encodes a polypeptide that is inactivated or unstable at acid pH i.e. pH below 6, even below
20 pH 5,5, more suitably even below pH 5 and even as low as or lower than pH 4. This is a particularly interesting embodiment, as the generally disclosed fungal expression systems are not cultured under conditions that are neutral to alkaline, but are cultured at acidic pH. Thus the system according to the invention provides a safe fungal expression system for proteins or polypeptides that are susceptible to being inactivated or are unstable at acid pH.

Quite specifically a recombinant strain as defined in any of the embodiments according to the invention, wherein the nucleic acid sequence encoding the polypeptide of interest encodes a protein or polypeptide exhibiting optimal activity and/or stability at a pH above 5, preferably at neutral or alkaline pH (i.e. above 7) and/or at a pH higher than 6, is considered a preferred
25 embodiment of the invention. More than 50%, more than 70% and even more than 90% of optimal activities at such pH values are anticipated as being particularly useful embodiments. A polypeptide expressed under the cultivation conditions does not necessarily have to be active at the cultivation conditions, in fact it can be advantageous for it to be cultured under conditions under which it is inactive as its active form could be detrimental to the host. This is the case for proteases for example. What is however required is for the protein or polypeptide to be stable under the cultivation
30 conditions. The stability can be thermal stability. It can also be stability against specific compositions or chemicals, such as are present for example in compositions or processes of production or

application of the polypeptide or protein of interest. LAS in detergent compositions comprising cellulases or lipases, etc. is an example of a chemical often detrimental to proteins. The time periods of use in applications can vary from short to long exposure so stability can be over a varying length of time varying per application. The skilled person will be able to ascertain the correct conditions on a case by case basis. One can use a number of commercially available assays to determine the optimal activities of the various enzymatic products. The catalogues of Sigma and Megazyme for example show such. Specific examples of tests are mentioned elsewhere in the description. The manufacturers provide guidance on the application.

We have surprisingly found that a *Chrysosporium* strain that can be suitably used to transform or transfect with the sequence of interest to be expressed is a strain exhibiting relatively low biomass. We have found that *Chrysosporium* strains having a biomass two to five times lower than that of *Trichoderma reesei* when cultured to a viscosity of 200-600 cP at the end of fermentation and exhibiting a biomass of 10 to 20 times lower than that of *Aspergillus niger* when cultured to a viscosity of 1500-2000 cP under corresponding conditions, i.e. their respective optimal cultivation conditions can provide a high level of expression. This level of expression far exceeds that of the two commercial reference strains at a much lower biomass and at much lower viscosity. This means that the yield of expression of such *Chrysosporium* strains will be appreciably higher than from *Aspergillus niger* and *Trichoderma reesei*. Such a transformed or transfected *Chrysosporium* strain forms a suitable embodiment of the invention.

We find a biomass of 0,5-1,0 g/l for *Chrysosporium* strain C1(18-25) as opposed to 2,5-5,0 g/l for *Trichoderma reesei* and 5-10 g/l of *Aspergillus niger* under the above described conditions. In the Examples we provide details of this process.

In a suitable embodiment a recombinant *Chrysosporium* strain according to the invention produces protein or polypeptide in at least the amount equivalent to the production in moles per liter of cellulase by the strain UV13-6 or C-19, and most preferably at least equivalent to or higher than that of the strain UV18-25 under the corresponding or identical conditions, i.e. their respective optimal cultivation conditions.

Unexpectedly we have also found that expression and secretion rates are exceedingly high when using a *Chrysosporium* strain exhibiting the mycelial morphology of strain UV18-25 i.e. fragmented short mycelia. Thus a recombinant strain according to the invention will preferably exhibit such morphology. The invention however also covers non-recombinant strains or otherwise engineered strains of *Chrysosporium* exhibiting this novel and inventive characteristic. Also covered by the invention is a recombinant *Chrysosporium* strain in any of the embodiments described according to the invention further exhibiting reduced sporulation in comparison to C1, preferably below that of strain UV13-6, preferably below that of NG7C-19, preferably below that of UV18-25 under equivalent fermenter conditions. Also covered by the invention is a recombinant *Chrysosporium*

strain in any of the embodiments described according to the invention further exhibiting at least the amount of protein production ratio to biomass in comparison to C1, preferably in comparison to that of any of strains UV13-6, NG7C-19 and UV18-25 under equivalent fermenter conditions. The invention however also covers non-recombinant strains or otherwise engineered strains of *Chrysosporium* exhibiting this novel and inventive characteristic as such or in combination with any of the other embodiments.

Another attractive embodiment of the invention also covers a recombinant *Chrysosporium* strain exhibiting a viscosity below that of strain NG7C-19, preferably below that of UV18-25 under corresponding or identical fermenter conditions. The invention however also covers non-recombinant strains or otherwise engineered strains of *Chrysosporium* exhibiting this novel and inventive characteristic as such or in combination with any of the other embodiments. We have determined that the viscosity of a culture of UV18-25 is below 10 cP opposed to that of *Trichoderma reesei* being of the order 200-600 cP, with that of *Aspergillus niger* being of the order 1500-2000 cP under their respective optimal culture conditions at the end of fermentation. The process used for such determination is provided in the examples.

Viscosity can be assessed in many cases by visual monitoring. The fluidity of the substance can vary to such a large extent that it can be nearly solid, sauce like or liquid. Viscosity can also readily be ascertained by Brookfield rotational viscometry, use of kinematic viscosity tubes, falling ball viscometer or cup type viscometer. The yields from such a low viscosity culture are higher than from the commercial known higher viscosity cultures per time unit and per cell.

The processing of such low viscosity cultures according to the invention is advantageous in particular when the cultures are scaled up. The subject *Chrysosporium* strains with the low viscosity perform very well in cultures as large as up to 150,000 liter cultures. Thus any culture size up to 150,000 litres provides a useful embodiment of the invention. Any other conventional size of fermentation should be carried out well with the strains according to the invention. The reasoning behind this is that problems can arise in large scale production with the formation of aggregates that have mycelia that are too dense and/or are unevenly distributed. The media as a result cannot be effectively utilised during the culture thus leading to an inefficient production process in particular in large scale fermentations i.e. over 150,000 liters. Aeration and mixing become problematic leading to oxygen and nutrient starvation and thus reduced concentration of productive biomass and reduced yield of polypeptide during the culture and/or can result in longer fermentation times. In addition high viscosity and high shear are not desirable in commercial fermentation processes and in current commercial processes they are the production limiting factors. All these negative aspects can be overcome by the *Chrysosporium* host according to the invention which exhibits much better characteristics than *Trichoderma reesei*, *Aspergillus niger* and *Aspergillus oryzae* that are commercially used in this respect i.e. exhibits better protein production levels and viscosity properties

and biomass figures.

A *Chrysosporium* strain selected from C1, UV13-6, NG7C-19 and UV18-25 illustrates various aspects of the invention exceedingly well. The invention however also covers recombinant strains or otherwise engineered strains of *Chrysosporium* derived from the four deposited strains that also exhibit any of the novel and inventive characteristics as such or in combination. The deposit data for these strains have been presented elsewhere in the description. The invention also covers recombinant strains or otherwise engineered strains of *Chrysosporium* derived from the four deposited strains that also exhibit any of the novel and inventive characteristics as such or in combination. A *Chrysosporium* strain according to the invention also comprises a strain exhibiting under the corresponding culture conditions a biomass at least twice as low as that of *Trichoderma reesei*, suitably even more up to 5 times lower than that of *Trichoderma reesei*, specifically of a *Trichoderma reesei* exhibiting a viscosity of 200-600 cP as disclosed under the conditions of the examples. A *Chrysosporium* strain according to the invention also comprises a strain producing the polypeptide in at least the amount in moles per liter of cellulase by the strain C1, UV13-6, NG7C-19 or UV18-25 under the corresponding or identical conditions.

Chrysosporium strains according to the invention are further preferred if they exhibit optimal growth conditions at neutral to alkaline pH and temperatures of 25-43°C. A preference can exist for neutral and even for alkaline pH. Such production conditions are advantageous to a number of polypeptides and proteins, in particular those susceptible to attack by acidic pH or those that are inactive or unstable at low temperatures. It is however also an embodiment of the invention to include *Chrysosporium* strains that can be cultured at acidic pH as this can be useful for certain proteins and polypeptides. A suitable acidic pH lies from 7.0. An acidic pH lower than 6.5 is envisaged as providing a good embodiment of the invention. A pH around 5.0-7.0 is also a suitable embodiment. A neutral pH can be 7.0 or around 7 e.g. 6.5-7.5. As stated elsewhere the pH of optimal interest depends on a number of factors that will be apparent to the person skilled in the art. A pH higher than 7.5 is alkaline, suitably between 7.5-9.0 can be used.

When comparing data of strains according to the invention with other strains perhaps having other optimal conditions (e.g. *Aspergillus* and *Trichoderma*) for viscosity measurements, biomass determination or protein production comparisons should be made using the relevant optimal conditions for the relevant strain. This will be obvious to the person skilled in the art.

A *Chrysosporium* strain according to any of the above-mentioned embodiments of the invention, said strain further exhibiting production of one or more of the fungal enzymes selected from the carbohydrate-degrading enzymes, proteases, other hydrolases, oxidoreductase, and transferases mentioned above is considered a particularly useful embodiment of the invention. The most interesting products are specifically cellulases, xylanases, pectinases, lipases and proteases. Also useful as embodiment of the invention however is a *Chrysosporium* strain exhibiting production of

one or more fungal enzymes that exhibit neutral or alkaline optimal stability and/or activity, preferably alkaline optimal stability and/or activity, said enzyme being selected from carbohydrate-degrading enzymes, hydrolases and proteases, preferably hydrolases and carbohydrate-degrading enzymes. In the case of non-recombinant *Chrysosporium*, such enzymes are suitably other than cellulase as disclosed
5 in WO 98/15633. Enzymes of particular interest are xylanases, proteases, esterases, alpha galactosidases, beta-galactosidases, beta-glucanases and pectinases. The enzymes are not limited to the aforementioned. The comments vis-à-vis stability and activity elsewhere in the description are valid here also.

The invention also covers a method of producing a polypeptide of interest, said method
10 comprising culturing a *Chrysosporium* strain in any of the embodiments according to the invention under conditions permitting expression and preferably secretion of the polypeptide and recovering the subsequently produced polypeptide of interest.

Where protein or polypeptide is mentioned, variants and mutants e.g. substitution, insertion or deletion mutants of naturally occurring proteins are intended to be included that exhibit the activity
15 of the non-mutant. The same is valid vis-à-vis the corresponding nucleic acid sequences. Processes such as gene shuffling, protein engineering and directed evolution site directed mutagenesis and random mutagenesis are processes through which such polypeptides, variants or mutants can be obtained. US 5,223,409, US 5,780,279 and US 5,770,356 provide teaching of directed evolution. Using this process a library of randomly mutated gene sequences created for example by gene
20 shuffling via error prone PCR occurs in any cell type. Each gene has a secretion region and an immobilising region attached to it such that the resulting protein is secreted and stays fixed to the host surface. Subsequently conditions are created that necessitate the biological activity of the particular polypeptide. This occurs for a number of cycles ultimately leading to a final gene with the desired characteristics. In other words a speeded up directed process of evolution. US 5,763,192 also
25 describes a process for obtaining DNA, RNA, peptides, polypeptides or protein by way of synthetic polynucleotide coupling stochastically generated sequences, introduction thereof into a host followed by selection of the host cell with the corresponding predetermined characteristic.

Standard cloning and protein or polypeptide isolation techniques can be used to arrive at the required sequence information. Parts of known sequences can be used as probes to isolate other
30 homologues in other genera and strains. The nucleic acid sequence encoding a particular enzyme activity can be used to screen a *Chrysosporium* library for example. A person skilled in the art will realise which hybridisation conditions are appropriate. Conventional methods for nucleic acid hybridisation construction of libraries and cloning techniques are described in Sambrook et al (Eds) (1989) In "Molecular Cloning. A Laboratory Manual" Cold Spring Harbor, Press Plainview, New
35 York, and Ausubel et al (Eds) "Current Protocols in Molecular Biology" (1987) John Wiley and Sons, New York. The relevant information can also be derived from later handbooks and patents, as well as

from various commercially available kits in the field.

In an alternative embodiment, said method comprises culturing a strain according to the invention under conditions permitting expression and preferably secretion of the protein or polypeptide or precursor thereof and recovering the subsequently produced polypeptide and optionally
5 subjecting the precursor to additional isolation and purification steps to obtain the polypeptide of interest. Such a method may suitably comprise a cleavage step of the precursor into the polypeptide or precursor of interest. The cleavage step can be cleavage with a Kex-2 like protease, any basic amino acid paired protease or Kex-2 for example when a protease cleavage site links a well secreted protein carrier and the polypeptide of interest. A person skilled in the art can readily find Kex-2-like protease
10 sequences as consensus sequence details for such are available and a number of alternatives have already been disclosed e.g. furin.

Suitably in a method for production of the polypeptide according to any of the embodiments of the invention the cultivation occurs at pH higher than 5, preferably 5-10, more preferably 6-9. Suitably in such a method the cultivation occurs at a temperature between 25-43 °C, preferably 30-
15 40°C. The *Chrysosporium* strain used in the method according to the invention is quite suitably a recombinant *Chrysosporium* strain according to any of the embodiments disclosed. The method according to the invention in such a case can further be preceded by the step of production of a recombinant *Chrysosporium* strain according to the invention. The selection of the appropriate conditions will depend on the nature of the polypeptide to be expressed and such selection lies well
20 within the realm of normal activity of a person skilled in the art.

The method of production of a recombinant *Chrysosporium* strain according to the invention is also part of the subject invention. The method comprises stably introducing a nucleic acid sequence encoding a heterologous or homologous polypeptide into a *Chrysosporium* strain, said nucleic acid sequence being operably linked to an expression regulating region, said introduction
25 occurring in a manner known per se for transforming filamentous fungi. As stated above numerous references hereof are available and a small selection has been cited. The information provided is sufficient to enable the skilled person to carry out the method without undue burden. The method comprises introduction of a nucleic acid sequence comprising any of the nucleic acid elements described in the various embodiments of the recombinant *Chrysosporium* according to the invention
30 as such or in combination.

By way of example the introduction can occur using the protoplast transformation method. The method is described in the examples. Alternative protoplast or spheroplast transformation methods are known and can be used as have been described in the prior art for other filamentous fungi. Details of such methods can be found in many of the cited references and are thus incorporated
35 by reference. A method according to the invention suitably comprises using a non-recombinant strain of *Chrysosporium* according to the invention as starting material for introduction of the desired

sequence encoding the polypeptide of interest.

The subject invention also covers a method of producing *Chrysosporium* enzyme, said method comprising culturing a *Chrysosporium* strain according to any of the embodiments of the invention as described above in or on a cultivation medium at pH higher than 5, preferably 5-10, more preferably 6-9, suitably 6-7.5, 7.5-9 as examples of neutral and alkaline pH ranges.

The subject invention also covers such a method using a cultivation medium at a temperature between 25-43 °C, preferably 30-40°C. The combination of preferred pH and temperature is an especially preferred embodiment of the method of producing *Chrysosporium* enzyme according to the invention.

More in general the invention further covers a method of producing enzymes exhibiting neutral or alkaline optimal activity and/or stability, preferably alkaline optimal activity and/or stability. The preferred ranges vis-à-vis pH and optimal activity as well as assays with which to determine such have been provided elsewhere in the description. The enzyme should be selected from carbohydrate-degrading enzymes, proteases, other hydrolases, oxidoreductases, and transferases, as described above, said method comprising cultivating a host cell transformed or transfected with the corresponding enzyme-encoding nucleic acid sequence. Suitably such an enzyme will be a *Chrysosporium* enzyme. A suitable method such as this comprises production specifically of cellulase, xylanase, pectinase, lipase and protease, wherein cellulase and xylanase cleave β -1,4-bonds and cellulase comprises endoglucanase, cellobiohydrolase and β -glucosidase. The method according to the invention can comprise cultivating any *Chrysosporium* host according to the invention comprising nucleic acid encoding such aforementioned enzymes. Suitably the production of non-recombinant *Chrysosporium* hosts according to the invention is directed at production of carbohydrate degrading enzymes, hydrolases and proteases. In such a case the enzyme is suitably other than a cellulase. Suitable examples of products to be produced are given in Tables A and B. Methods of isolating are analogous to those described in WO 98/15633 and are incorporated by reference.

The enzymes produced by *Chrysosporium* strains according to the invention are also covered by the invention. Enzymes of *Chrysosporium* origin as can be isolated from non-recombinant *Chrysosporium* strains according to the invention are also covered. They exhibit the aforementioned stability, activity characteristics. Suitably they are stable in the presence of LAS. In particular proteases with pI 4-9.5, proteases with a MW of 25-95 kD, xylanases with pI between 4.0 and 9.5, xylanases with MW between 25 and 65 kD, endoglucanases with a pI between 3.5 and 6.5, endoglucanases with MW of 25-55 kDa, β -glucosidases, α,β -galactosidases with a pI of 4-4.5, β -glucosidases, α,β -galactosidases with a MW of 45-50 kDa, cellobiohydrolases of pI 4-5, cellobiohydrolases of MW 45-60 kDa, e.g. a MW of 55 kD and pI 4.4, polygalacturonases, with a pI of 4.0-5.0 polygalacturonase of 60-70 kDa, e.g. 65 kDa, esterases with a pI 4-5, and esterases with a MW of 95-105 kDa with the afore-mentioned stability, activity characteristics are claimed. The

molecular weights (MW) are those determined by SDS-PAGE. The non-recombinant i.e. natively occurring enzyme is other than cellulase as disclosed in WO 98/15633. An enzyme as disclosed in WO 98/15633 is excluded. Enzymes according to the invention are represented by the enzymes of Table B. Enzymes with combinations of the pI values and molecular weights mentioned above are also covered.

The invention is also concerned with the (over)production of non-protein products by the mutant (recombinant) strains of the invention. Such non-protein products include primary metabolites such as organic acids, amino acids, and secondary such as antibiotics, e.g. penicillins and cephalosporins. These products are the result of combinations of biochemical pathways, involving several fungal genes of interest. Fungal primary and secondary metabolites and procedures for producing these metabolites in fungal organisms are well known in the art. Examples of the production of primary metabolites have been described by Matthey M., The Production of Organic Acids, *Current Reviews in Biotechnology*, 12, 87-132 (1992). Examples of the production of secondary metabolites have been described by Penalva et al. The Optimization of Penicillin Biosynthesis in Fungi, *Trends in Biotechnology* 16, 483-489 (1998).

EXAMPLES

EXAMPLES OF BIOMASS AND VISCOSITY DETERMINATIONS

The following operating parameter data ranges have been determined for fungal fermentations using three different fungal organisms. The three fungal organisms compared are: *Trichoderma longibrachiatum* (formerly *T. reesei*), *Aspergillus niger* and *Chrysosporium lucknowense* (UV18-25).

Viscosity:

Viscosity is determined on a Brookfield LVF viscometer using the small sample adapter and spindle number 31.

Turn the water-circulating pump on 5 minutes prior to viscometer use to equilibrate the water jacket. The water bath temperature should be 30°C.

Obtain a fresh sample of fermentation broth and place 10 ml of the broth in the small sample spindle. Select the spindle speed to give a reading in the range 10-80. Wait four (4) minutes and take the reading from the viscometer scale. Multiply the reading by the factor given below to get the viscosity in centipoise (cP).

Spindle Speed	Multiplication Factor
6	50
12	25
30	10
60	5

The following viscosity ranges have been determined for fermentations using the specified fungal

organism using the above procedure:

	Viscosity in cP
<i>T. longibrachiatum</i>	200 - 600
<i>A. niger</i>	1,500 - 2,000
5 <i>C. lucknowense</i> (UV18-25)	LT 10

Biomass:

Biomass is determined by the following procedure:

Preweigh 5.5 cm filter paper (Whatman 54) in an aluminium weighing dish.

- 10 Filter 5.0 ml whole broth through the 5.5 cm paper on a Buchner funnel, wash the filter cake with 10 ml deionised water, place the washed cake and filter in a weighing pan and dry overnight at 60°C. Finish drying at 100°C for 1 hour, then place in desiccator to cool.

Measure the weight of dried material. Total biomass (g/l) is equal to the difference between the initial and final weights multiplied by 200.

- 15 The following biomass ranges have been determined for fermentations using the specified fungal organism using the above procedure:

	Biomass in g/l
<i>T. longibrachiatum</i>	2.5 - 5
<i>A. niger</i>	5 - 10
20 <i>C. lucknowense</i> (UV18-25)	0.5 - 1

Protein:

Protein levels were determined using the BioRad Assay Procedure from Sigma Company. Protein levels were highest for the *Chrysosporium*.

- 25 The data presented above represent values determined 48 hours into the fermentation process until fermentation end; All values of *Aspergilli* and *Trichoderma* are for commercially relevant fungal organisms and reflect actual commercial data.

- 30 A fungal strain such as *C. lucknowense* (UV18-25) has the advantage that the low viscosity permits the use of lower power input and/or shear the in the fermentation to meet oxygen demands for those cases where shear stress on the product may be detrimental to productivity due to physical damage of the product molecule. The lower biomass production at high protein production indicates a more efficient organism in the conversion of fermentation media to product. Thus the *Chrysosporium* provides better biomass and viscosity data whilst also delivering at least as much protein, and in fact a lot more protein than the two commercially used systems which obviously are better than for typically
35 deposited *Aspergillus* or *Trichoderma reesei* strains in general public collections.

The high protein production with low biomass concentration produced by *C. lucknowense*

(UV18-25) would allow development of fermentation conditions with higher multiples of increase in biomass, if increasing biomass results in increased productivity, for the desired product before reaching limiting fermentation conditions. The present high levels of biomass and viscosity produced by the *T. longibrachiatum* and *A. niger* organisms restrict the increase of biomass as the present levels of biomass and viscosity are near limiting practical fermentation conditions.

EXAMPLES OF TRANSFORMATION COMPARING CHRYSOSPORIUM, TRICHODERMA AND TOLYPOCLADIUM GEODES

Two untransformed *Chrysosporium* C1 strains and one *Trichoderma reesei* reference strain were tested on two media (Gs pH 6,8 and Pridham agar, PA, pH 6,8). To test the antibiotic resistance level spores were collected from 7 day old PDA plates. Selective plates were incubated at 32°C and scored after 2,4 and 5 days. It followed that the C-1 strains NG7C-19 and UV18-25 clearly have a low basal resistance level both to phleomycin and hygromycin. This level is comparable to that for a reference *T. reesei* commonly used laboratory strain. Thus there is clear indication these two standard fungal selectable markers can be used well in *Chrysosporium* strains. Problems with other standard fungal selectable markers should not be expected.

Selection of Sh-ble (phleomycin-resistance) transformed *Chrysosporium* strains was successfully carried out at 50 µg/ml. This was also the selection level used for *T. reesei* thus showing that differential selection can be easily achieved in *Chrysosporium*. The same comments are valid for transformed strains with hygromycin resistance at a level of 150 µg/ml.

Table C

	Gs (pH 6.8)			Pridham Agar (PA,pH 6.8)		
	NG7C-19	UV18-25	T.r.11D5	NG7C-19	UV18-25	T.r.11D5
Phleomycin	7.5 µg/ml	10 µg/ml	5-7.5 µg/ml	2.5 µg/ml	10 µg/ml	2.5 µg/ml
Hygromycin	7.5 - 10 µg/ml	10 µg/ml	10 µg/ml	15 µg/ml	25 µg/ml	15 µg/ml

The protoplast transformation technique was used on *Chrysosporium* based on the most generally applied fungal transformation technology. All spores from one 90mm PDA plate were recovered in 8ml IC1 and transferred into a shake flask of 50ml IC1 medium for incubation for 15 hours at 35°C and 200 rpm. After this the culture was centrifuged, the pellet was washed in MnP, brought back into solution in 10ml MnP and 10mg/ml Caylase C₃ and incubated for 30 minutes at 35°C with agitation (150 rpm).

The solution was filtered and the filtrate was subjected to centrifugation for 10 minutes at 3500 rpm. The pellet was washed with 10 ml MnPCa^{2+} . This was centrifuged for 10 minutes at 25°C. Then 50 microlitres of cold MPC was added. The mixture was kept on ice for 30 minutes whereupon 2,5 ml PMC was added. After 15 minutes at room temperature 500 microlitres of the treated protoplasts were mixed to 3 ml of MnR Soft and immediately plated out on a MnR plate containing phleomycin or hygromycin as selection agent. After incubation for five days at 30°C transformants were analysed (clones become visible after 48 hours). Transformation efficiency was determined using 10 microgrammes of reference plasmid pAN8-1¹⁹. The results are presented in the following Table D.

10

Table D: Transformation efficiency (using 10 µg of reference plasmid pAN8-1)

	T. reesei	NG7C-19	UV18-25
Viability	$10^6/200 \mu\text{l}$	$5 \cdot 10^6/200 \mu\text{l}$	$5 \cdot 10^6/200 \mu\text{l}$
Transformants Per 200 µl	2500	10^4	10^4
Transformants per 10^6 viable cells	2500	2000	2000

The results show that the *Chrysosporium* transformants viability is superior to that of *Trichoderma*. The transformability of the strains is comparable and thus the number of transformants obtained in one experiment lies 4 times higher for *Chrysosporium* than for *T. reesei*. Thus the *Chrysosporium* transformation system not only equals the commonly used *T. reesei* system, but even outperforms it. This improvement can prove especially useful for vectors that are less transformation efficient than pAN8-1. Examples of such less efficient transformation vectors are protein carrier vectors for production of non-fungal proteins which generally yield 10 times fewer transformants.

A number of other transformation and expression vectors were constructed with homologous *Chrysosporium* protein encoding sequences and also with heterologous protein encoding sequences for use in transformation experiments with *Chrysosporium*. The vector maps are provided in the figures 6-11.

The homologous protein to be expressed was selected from the group of cellulases produced by *Chrysosporium* and consisted of endoglucanase 6 which belongs to family 6 (MW 43 kDa) and the heterologous protein was endoglucanase 3 which belongs to family 12 (MW 25 kDa) of *Penicillium*.

pF6g comprises *Chrysosporium* endoglucanase 6 promoter fragment linked to endoglucanase 6 signal sequence in frame with the endoglucanase 6 open reading frame followed by the endoglucanase 6 terminator sequence. Transformant selection is carried out by using cotransformation

with a selectable vector.

pUT1150 comprises *Trichoderma reesei* cellobiohydrolase promoter linked to endoglucanase 6 signal sequence in frame with the endoglucanase 6 open reading frame followed by the *T. reesei* cellobiohydrolase terminator sequence. In addition this vector carries a second expression
5 cassette with a selection marker i.e. the phleomycin resistance gene (Sh-ble gene).

pUT1152 comprises *Aspergillus nidulans* glyceraldehyde-3-phosphate dehydrogenase A promoter linked to endoglucanase 6 signal sequence in frame with the endoglucanase 6 open reading frame followed by the *A. nidulans* anthranilate synthase (trpC) terminator sequence. In addition this vector carries a second expression cassette with a selection marker i.e. the phleomycin resistance gene
10 (Sh-ble gene).

pUT1155 comprises *A. nidulans* glyceraldehyde-3-phosphate dehydrogenase A promoter linked to *Trichoderma reesei* cellobiohydrolase signal sequence in frame with the carrier protein Sh-ble which in turn is linked in frame to the endoglucanase 6 open reading frame followed by the *A. nidulans* trpC terminator sequence. This vector uses the technology of the carrier protein fused to the
15 protein of interest which is known to very much improve the secretion of the protein of interest.

pUT1160 comprises *Aspergillus nidulans* glyceraldehyde-3-phosphate dehydrogenase A promoter linked to *Trichoderma reesei* cellobiohydrolase signal sequence in frame with the carrier protein Sh-ble which in turn is linked in frame to the endoglucanase 3 open reading frame of *Penicillium* followed by the *A. nidulans* trpC terminator sequence.

20 pUT1162 comprises *Trichoderma reesei* cellobiohydrolase promoter linked to endoglucanase 3 signal sequence in frame with the endoglucanase 3 open reading frame of *Penicillium* followed by the *T. reesei* cellobiohydrolase terminator sequence. In addition this vector carries a second expression cassette with a selection marker i.e. the phleomycin resistance gene (Sh-ble gene).

25 **Table E: Comparative transformations**

Vector	Strain	Transformation	No of transf.	Tested in liquid culture
PUT1150	UV18-25 <i>T. geodes</i>	selection phleo	285	5
		selection phleo	144	5
PUT1152	UV18-25 <i>T. geodes</i>	cotransformation pAN8.1	398	5
		cotransformation pAN8.1	45	4
PF6g	UV18-25 <i>T. geodes</i>	cotransformation pAN8.1	252	6
		cotransformation pAN8.1	127	5
PUT1162	UV18-25 <i>T. geodes</i>	selection phleo Not done yet	>400	

Table E shows the results of transformation of both *Chrysosporium* UV18-25 and *Tolypocladium geodes*. The transformation protocol used is described in the section for heterologous transformation.

EXAMPLES OF HETEROLOGOUS AND HOMOLOGOUS EXPRESSION OF CHRYSO-SPORIUM TRANSFORMANTS

C1 strains (NG7C-19 and/or UV18-25) have been tested for their ability to secrete various heterologous proteins: a bacterial protein (*Streptoalloteichus hindustanus* phleomycin-resistance protein, Sh ble), a fungal protein (*Trichoderma reesei* xylanase II, XYN2) and a human protein (the human lysozyme, HLZ).

The details of the process are as follows:

[1] C1 secretion of *Streptoalloteichus hindustanus* phleomycin-resistance protein (Sh ble).

C1 strains NG7C-19 and UV18-25 have been transformed by the plasmid pUT720¹. This vector presents the following fungal expression cassette:

- *Aspergillus nidulans* glyceraldehyde-3-phosphate dehydrogenase (*gpdA*) promoter²
- A synthetic *Trichoderma reesei* cellobiohydrolase I (*cbh1*) signal sequence^{1,3}
- *Streptoalloteichus hindustanus* phleomycin-resistance gene *Sh ble*⁴
- *Aspergillus nidulans* tryptophan-synthase (*trpC*) terminator⁵

The vector also carries the beta-lactamase gene (*bla*) and *E. coli* replication origin from plasmid pUC18⁶. The detailed plasmid map is provided in figure 2.

C1 protoplasts were transformed according to Durand et al.⁷ adapted to C1 (media & solutions composition is given elsewhere): All spores from one 90mm PDA plate of untransformed C1 strain were recovered in 8ml IC1 and transferred into a shake flask with 50ml IC1 medium for incubation 15 hours at 35°C and 150 rpm. Thereupon, the culture was spun down, the pellet washed in MnP, resolved in 10ml MnP + 10mg/ml Caylase C₃, and incubated 30 min at 35°C with agitation (150 rpm). The solution was filtrated and the filtrate was centrifuged 10 min at 3500 rpm. The pellet was washed with 10ml MnPCa²⁺. This was spun down 10min at 3500 rpm and the pellet was taken up into 1ml MnPCa²⁺. 10µg of pUT720 DNA were added to 200µl of protoplast solution and incubated 10min at room temperature (~20°C). Then, 50µl of cold MPC was added. The mixture was kept on ice for 30min whereupon 2.5ml PMC was added. After 15min at room temperature 500µl of the treated protoplasts were mixed to 3ml of MnR Soft and immediately plated out on a MnR plate containing phleomycin (50µg/ml at pH6.5) as selection agent. After 5 days incubation at 30°C, transformants were analysed (clones start to be visible after 48 hours).

The Sh ble production of C1 transformants (phleomycin-resistant clones) was analysed as

follows: Primary transformants were toothpicked to GS+phleomycin (5µg/ml) plates and grown for 5 days at 32°C for resistance verification. Each validated resistant clone was subcloned onto GS plates. Two subclones per transformant were used to inoculate PDA plates in order to get spores for liquid culture initiation. The liquid cultures in IC1 were grown 5 days at 27°C (shaking 200 rpm). Then, the cultures were centrifuged (5000g, 10min.) and 500µl of supernatant were collected. From these samples, the proteins were precipitated with TCA and resuspended in Western Sample Buffer to 4 mg/ml of total proteins (Lowry Method ⁸). 10µl (about 40µg of total proteins) were loaded on a 12% acrylamide/SDS gel and run (BioRad Mini Trans-Blot system). Western blotting was conducted according to BioRad instructions (Schleicher & Schull 0.2µm membrane) using rabbit anti-Sh ble antiserum (Cayla Cat. Ref. #ANTI-0010) as primary antibody.

The results are shown in Figure 1 and Table F:

Table F: Sh ble estimated production levels in C1

	Estimated Sh ble quantity on the Western blot	Estimated Sh ble concentration in the production media
Untransformed NG7C-19	Not detectable	
NG7C-19::720 clone 4-1	25 ng	0.25 mg/l
NG7C-19::720 clone 5-1	25 ng	0.25 mg/l
NG7C-19::720 clone 2-2	250 ng	2.5 mg/l
Untransformed UV18-25	Not detectable	
UV18-25::720 clone 1-2	500 ng	5 mg/l
UV18-25::720 clone 3-1	250 ng	2.5 mg/l

These data show that:

- 1) The heterologous transcription/translation signals from pUT720 are functional in *Chrysosporium*.
- 2) The heterologous signal sequence of pUT720 is functional in *Chrysosporium*.
- 3) *Chrysosporium* can be used a host for the secretion of an heterologous bacterial protein.

[2] C1 secretion of the human lysozyme (HLZ).

C1 strains NG7C-19 and UV18-25 have been transformed by the plasmid pUT970G ⁹. This vector presents the following fungal expression cassette:

- *Aspergillus nidulans* glyceraldehyde-3-phosphate dehydrogenase (*gpdA*) promoter ²
- A synthetic *Trichoderma reesei* cellobiohydrolase I (*cbhI*) signal sequence ^{1,3}
- *Streptoalloteichus hindustanus* phleomycin-resistance gene *Sh ble* ⁴ used as carrier-protein

10

- *Aspergillus niger* glucoamylase (*glaA2*) hinge domain cloned from plasmid pAN56-2 ^{11, 12}
- A linker peptide (LGERK) featuring a KEX2-like protease cleavage site ¹
- A synthetic human lysozyme gene (*hlz*) ¹⁰
- 5 - *Aspergillus nidulans* tryptophan-synthase (*trpC*) terminator ⁵

The vector also carries the beta-lactamase gene (*bla*) and *E. coli* replication origin from plasmid pUC18 ⁶. The detailed plasmid map is provided in figure 3.

C1 protoplasts were transformed with plasmid pUT970G following the same procedure already described in example 1. The fusion protein (Sh ble :: GAM hinge :: HLZ) is functional with respect to
10 the phleomycin-resistance thus allowing easy selection of the C1 transformants. Moreover, the level of phleomycin resistance correlates roughly with the level of *hlz* expression.

The HLZ production of C1 transformants (phleomycin-resistant clones) was analysed by lysozyme-activity assay as follow: Primary transformants were toothpicked to GS+phleomycin (5µg/ml) plates (resistance verification) and also on LYSO plates (HLZ activity detection by clearing
15 zone visualisation ^{1, 10}). Plates were grown for 5 days at 32°C. Each validated clone was subcloned onto LYSO plates. Two subclones per transformant were used to inoculate PDA plates in order to get spores for liquid culture initiation. The liquid cultures in IC1 were grown 5 days at 27°C (shaking 180 rpm). Then, the cultures were centrifuged (5000g, 10min.). From these samples, lysozyme activity was measured according to Mörsky et al. ¹³.

20

Table G: Active HLZ production levels in C1

	Active HLZ concentration in culture media
Untransformed NG7C-19	0 mg/l
NG7C-19::970G clone 4	4 mg/l
NG7C-19::970G clone 5	11 mg/l
Untransformed UV18-25	0 mg/l
UV18-25::970G clone 1	8 mg/l
UV18-25::970G clone 2	4 mg/l
UV18-25::970G clone 3	2 mg/l
UV18-25::970G clone 2	2.5 mg/l

These data show that:

- 1) Points 1 & 2 from example 1 are confirmed.
- 25 2) Sh ble is functional in *Chrysosporium* as resistance-marker.

3) *Sh ble* is functional in *Chrysosporium* as carrier-protein.

4) The KEX2-like protease cleavage site is functional in *Chrysosporium* (otherwise HLZ wouldn't be active).

5) *Chrysosporium* can be used as host for the secretion of a heterologous mammalian protein.

[3] C1 secretion of *Trichoderma reesei* xylanase II (XYN2).

C1 strain UV18-25 has been transformed by the plasmids pUT1064 and pUT1065.

pUT1064 presents the two following fungal expression cassettes:

10 The first cassette allows the selection of phleomycin-resistant transformants:

- *Neurospora crassa* cross-pathway control gene 1 (*cpc-1*) promoter ¹⁴
- *Streptoalloteichus hindustanus* phleomycin-resistance gene *Sh ble* ⁴
- *Aspergillus nidulans* tryptophan-synthase (*trpC*) terminator ⁵

The second cassette is the xylanase production cassette:

- 15
- *T. reesei* strain TR2 *cbh1* promoter ¹⁵
 - *T. reesei* strain TR2 *xyn2* gene (including its signal sequence) ¹⁶
 - *T. reesei* strain TR2 *cbh1* terminator ¹⁵

The vector also carries an *E. coli* replication origin from plasmid pUC19 ⁶. The plasmid detailed map is provided in figure 4.

20 pUT1065 presents the following fungal expression cassette:

- *A. nidulans* glyceraldehyde-3-phosphate dehydrogenase (*gpdA*) promoter ²
- A synthetic *T. reesei* cellobiohydrolase I (*cbh1*) signal sequence ^{1,3}
- *S. hindustanus* phleomycin-resistance gene *Sh ble* ⁴ used as carrier-protein ¹⁰
- A linker peptide (SGERK) featuring a KEX2-like protease cleavage site ¹
- 25 - *T. reesei* strain TR2 *xyn2* gene (without signal sequence) ¹⁶
- *A. nidulans* tryptophan-synthase (*trpC*) terminator ⁵

The vector also carries the beta-lactamase gene (*bla*) and an *E. coli* replication origin from plasmid pUC18 ⁶. The plasmid detailed map is provided in figure 5.

30 C1 protoplasts were transformed with plasmid pUT1064 or pUT1065 following the same procedure already described in example 1. The fusion protein in plasmid pUT1065 (*Sh ble* :: XYN2) is functional with respect to the phleomycin-resistance thus allowing easy selection of the C1 transformants. Moreover, the level of phleomycin resistance correlates roughly with the level of *xyn2* expression. In pUT1064, *xyn2* was cloned with its own signal sequence.

35 The xylanase production of C1 transformants (phleomycin-resistant clones) was analysed by xylanase-activity assay as follow: Primary transformants were toothpicked to GS+phleomycin (5µg/ml) plates (resistance verification) and also on XYLAN plates (xylanase activity detection by

clearing zone visualisation ¹⁷). Plates were grown for 5 days at 32°C. Each validated clone was subcloned onto XYLAN plates. Two subclones per transformant were used to inoculate PDA plates in order to get spores for liquid culture initiation. The liquid cultures in IC1+ 5g/l KPhtalate were grown 5 days at 27°C (shaking 180 rpm). Then, the cultures were centrifuged (5000g, 10min.). From these samples, xylanase activity was measured by DNS Technique according to Miller et al. ¹⁸

Table H: Active XYN2 production levels in C1 (best producers)

	Active xylanase II concentration in culture media	Xylanase II specific activity in culture media
Untransformed UV18-25	3.9 U./ml	3.8 U./mg total prot.
UV18-25::1064 clone 7-1	4.7 U./ml	4.7 U./mg total prot.
UV18-25::1064 clone 7-2	4.4 U./ml	4.3 U./mg total prot.
UV18-25::1065 clone 1-1	29.7 U./ml	25.6 U./mg total prot.
UV18-25::1065 clone 1-2	30.8 U./ml	39.4U./mg total prot.

These data show that:

- 1) Points 1 to 4 from example 2 are confirmed.
- 2) C1 can be used as host for the secretion of a heterologous fungal protein.

[4] We also illustrate data from expression of transformed UV18-25 wherein the table I shows the results for the plasmids with which transformation was carried out. The Table shows good expression levels for endoglucanase and cellobiohydrolase using heterologous expression regulating sequences and signal sequences but also with homologous expression regulating sequences and signal sequences. The details of the various plasmids can be derived elsewhere in the description and from the figures. The production occurs at alkaline pH at a temperature of 35°C.

Table I: Expression data of transformed UV18-25 strain

Culture	Total proteins	CMCase		β-glucanase		pH value
	mg/ml	u/ml	u/mg	u/ml	u/mg	
*UV 18-25	100%	100%	100%	100%	100%	7.90
1150-23	94%	105%	111%	140%	149%	7.90
-30	96%	105%	110%	145%	151%	8.10
1152-3	94%	112%	120%	147%	156%	7.85
-4	100%	105%	105%	132%	132%	7.90
1160-2	69%	81%	118%	90%	131%	7.90
-4	73%	72%	98%	83%	114%	8.35
-1	92%	95%	103%	120%	130%	8.45
1162-1	102%	105%	103%	145%	142%	8.20
-11	112%	109%	98%	115%	103%	8.20
F6g-20	104%	102%	98%	130%	125%	7.90
-25	-	-	-	-	-	-

Culture conditions (shake flask): 88h, 35°C, 230 rpm

* all above figures are in relative % to parent UV18-25 strain

5

Appendix to the Examples: Media

Transformation media:

10	Mandels Base:	MnP Medium :	
	KH ₂ PO ₄	2.0 g/l	Mandels Base with
	(NH ₄) ₂ SO ₄	1.4 g/l	Peptone 1 g/l
	MgSO ₄ .7H ₂ O	0.3 g/l	MES 2 g/l
	CaCl ₂	0.3 g/l	Sucrose 100 g/l
15	Oligoelements	1.0 ml/l	Adjust pH to 5
	MnR	MnP CA²⁺ :	
	MnP+sucrose	130 g/l	MnP Medium +
	Yeast extract	2.5 g/l	CaCl ₂ 2H ₂ O 50 mM
	Glucose	2.5 g/l	Adjust pH to 6.5
20	Agar	15 g/l	
	MnR Soft :	MnR with only 7.5 g/l of agar.	

	MPC :		
	CaCl ₂	50 mM	pH 5.8
	MOPS	10 mM	
5	PEG	40%	
<u>For selection and culture</u>			
	GS :		
	Glucose	10 g/l	
10	Biosoyase	5 g/l	[Merieux]
	Agar	15 g/l	pH should be 6.8
	PDA :		
	Potato Dextrose Agar	39 g/l	[Difco]
15			pH should be 5.5
	MPG :		
	Mandels Base with		
	K.Phtalate	5 g/l	
	Glucose	30 g/l	
20	Yeast extract	5 g/l	

The regeneration media (MnR) supplemented with 50 µg/ml phleomycin or 100-150 µg/ml hygromycin is used to select transformants. GS medium, supplemented with 5 µg/ml phleomycin is used to confirm antibiotic resistance.

- 25 PDA is a complete medium for fast growth and good sporulation. Liquid media are inoculated with 1/20th of spore suspension (all spores from one 90mm PDA plate in 5ml 0.1% Tween). Such cultures are grown at 27°C in shake flasks (200 rpm).

ISOLATION AND CHARACTERISATION OF CI PROTEINS

- 30 The process for obtaining various proteins is described as are a number of characteristics of the proteins. The tables A, B and J provide details of purification scheme and activities. Isolation occurs from the *Chrysosporium* culture filtrate using DEAE-Toyopearl ion exchange chromatography analogously to the method described in WO 98/15633, which is incorporated herein by reference. The non-bound fraction (F 60-31 CF) obtained from this chromatography was purified using Macro Prep
- 35 Q ion exchange chromatography after equilibration to pH 7.6. The non-bound fraction (NBNB) was pooled and bound proteins were eluted in 0-1 M NaCl gradient. The NBNB fraction provided major protein bands of 19, 30, 35 and 46 kD and a minor one of 51 kD. In 0-1 M NaCl gradient protein peaks were eluted from various fractions. 39-41 included 28, 36 and 60 kD proteins, 44-48 included

28, 45 and 66 kD as major protein bands with 33, 36, 55, 60 and 67 kD proteins, the 49-51 fraction gave 30, 36, 56 and 68 kD proteins and the 52-59 fraction included major 33 and 55 kD proteins and minor 28 and 36 kD proteins. The pooled NBNB fraction was further purified by hydrophobic chromatography on Phenyl Superose. The NBNB fraction was equilibrated with 0,03M Na-phosphate
 5 buffer pH 7,0 containing 1,2 M $(\text{NH}_4)_2\text{SO}_4$ and applied to a column. Adsorbed proteins were eluted in 1,2-0,6 M $(\text{NH}_4)_2\text{SO}_4$ gradient. Thus homogeneous xylanase with MW 30 and 51 kD and pI 9.1 and 8.7 respectively were obtained as was a 30 kD protease with pI 8,9.

The xylanases did not possess MUF cellobiase activity and are thus true xylanases. The alkaline 30 kD xylanase (pI 9.1) possessed high activity within a very broad pH range from 5-8
 10 maintaining 65% of maximum activity at pH 9-10; it is a member of the xylanase F family; its partial nucleotide and amino acid sequences are depicted in SEQ ID No. 5. The partial amino acid sequence depicted corresponds to about amino acids 50-170 from the N terminus of the mature protein. Xylanases according to invention have at least 60%, preferably at least 70%, most preferably at least 80% sequence identity of the partial amino acid sequence of SEQ ID No. 5. The corresponding
 15 xylanase promoter, which is a preferred embodiment of the invention, can be identified using the partial nucleotide sequence of SEQ ID No. 5. The 51 kD xylanase (pI 8,7) possessed maximum activity at pH 6 and retained at least 70% of its activity at pH 7,5 and it retained at least 50% of its activity at pH 8,0. It was not very stable with only 15% activity at pH 5,5 and 4% at pH 7,5. The Michaelis constant toward birch xylan was 4,2 g/l for 30kD xylanase and 3,4 g/l for 51 kD xylanase.
 20 Temperature optimum was high and equal to 70°C for both xylanases.

The 30 kD protease activity measured towards proteins of the NBNB fraction appeared to be equal to $0,4 \times 10^3$ units/ml at 50°C and pH 7,90 kD. The fraction exhibited activity toward dyed casein of 0,4 arbitrary units/mg (pH 7). Addition of urea as chaotropic agent resulted in 2-3 times increase of protease activity. The effect of the protease on xylanase activity was significant. Only 30%
 25 xylanase activity remained at pH 10,3 and 50°C after 30 minutes of incubation. At pH 8 95% of the xylanase activity remained. LAS addition resulted in a dramatic decrease of xylanase activity at pH 8 and 10,3 with only 50% xylanase activity after 10 minutes of incubation with or without protease inhibitor PMSF. The 30 kD protease was alkaline with pH optimum at pH 10-11. The activity is inhibited by phenylmethylsulfonyl fluoride (PMSF) and not by iodoacetic acid, pepstatin A and
 30 EDTA which characterises it as a serine type protease. The protease is not active towards C1 proteins at neutral pH and 50°C without chaotropic agents. Increase of pH and the addition of chaotropic agents such as LAS, SDS and urea significantly increase proteolysis.

The 39-41 fraction was purified by hydrophobic chromatography on plenol superose. Fractions were equilibrated with 0,03M Na phosphate buffer pH 7,2 containing 1,5 M $(\text{NH}_4)_2\text{SO}_4$ and
 35 applied to a column. Adsorbed proteins were eluted in 1,5-0 M $(\text{NH}_4)_2\text{SO}_4$ gradient. Thus homogenous xylanase with MW 60 kD and pI 4,7 was obtained. This xylanase possessed activities

04-12-2000

BO 42057

NL 009900618
PCT/NL77/00010

-36-

04. 12. 2000

(42)

towards xylan, MUF-cellobioside, MUF-xyloside and MUF-lactoside. This xylanase probably belongs to family 10 (family F). This xylanase was stable at pH from 5 to 8 during 24 hours and retained more than 80% activity at 50°C. It retained 70% activity at pH 5-7 at 60°C. It kept 80% activity during 5 hours and 35% during 24 hours at 50°C and pH 9. At pH 10 60% activity was retained at 50°C and 0,5 hours of incubation. After 5 hours of incubation at pH 8 and 60°C 45% activity was found decreasing to 0 after 24 hours. It had a pH optimum within the pH range of 6-7 and kept 70% activity at pH 9 and 50% of its activity at pH 9,5. The Michaelis constant toward birch xylan was 0,5 g/l. Temperature optimum was high and equal to 80°C.

Fraction 44-48 was then purified by chromatofocusing on Mono P. A pH gradient from 7,63-5,96 was used for the elution of the proteins. As a result 45 kD endoglucanase was isolated with a pI of 6. The 45 kD endo had maximum activity at pH 5 toward CMC and at pH 5-7 toward RBB-CMC. The 45 kD endo retained 70% of its maximal activity toward CMC at pH 6,5 and 70% of its maximal activity toward RBB-CMC was retained at pH 7,0; 50% of its maximal activity toward CMC was retained at pH 7 and 50% of its maximal activity toward RBB-CMC was retained at pH 8. The Michaelis constant toward CMC was 4,8 g/l. Temperature optimum was high and equal to 80°C. Other proteins 28, 33, 36, 55, 60 and 66 kD were eluted mixed together.

Fraction 52-58 was purified by chromatofocusing on Mono P too with a pH gradient 7,6-4,5. Individual 55 kD endoglucanase with pI 4,9 was obtained. The 55 kD endo was neutral. It has a broad pH optimum from 4,5-6 and 70% activity was retained at pH 7,0 both for CMC and RBB-CMC and 50% activity was retained at pH 8 for both CMC and RBB-CMC. The Michaelis constant toward CMC was 1 g/l. Temperature optimum was high and equal to 80°C. A number of fractions also held proteins with MW of 28, 33 and 36 kD.

45, 48 and 100 kD proteins were isolated from bound DEAE Toyopearl fraction of F 60-8 UF conc of *Chrysosporium* culture from fractions 50-53 using Macro Prep Q chromatography.

Fraction 50-53 was equilibrated with 0.03 M imidazole HCL buffer, pH 5.75 and was applied to a column and the adsorbed proteins were eluted in 0,1-0,25 M NaCl gradient for 4 h. As a result 45 kD (pI 4.2), 48 kD (pI 4.4) and 100 kD (pI 4.5) proteins were isolated in homogenous states (figure 17).

The 45 kD is supposedly a alpha beta-galactosidase by virtue of its activity toward p-nitrophenyl alpha-galactoside and p-nitrophenyl beta-galactoside. The pH optimum was 4,5 70% activity was maintained at pH 5,7 and 50% of its activity was retained at pH 6,8. The temperature optimum was 60°C (figure 18).

The 48 kD protein was a cellobiohydrolase having high activity toward p-nitrophenyl beta-glucoside and also activities toward MUF cellobioside, MUF lactoside and p-nitrophenyl butyrate. The 48 kD protein had a pH optimum of 5 toward CMC and 5-6 toward RBB-CMC (figure 19).

The 100 kD protein with pI 4,5 possessed activity only toward p-nitrophenyl butyrate. It is

04-12-2000

BO 42057

NL 009900618
PCT/NL99/00618

-37-

probably an esterase but is not a feruloyl esterase as it had no activity against methyl ester of ferulic acid. It had neutral/alkaline pH optimum (pH 8-9) and optimal temperature of 55-60°C (fig's 20, 21).

The 90 kD protease with pI 4,2 was isolated from the bound fraction and the activity measured towards proteins of the NBNB fraction appeared to be equal to 12×10^{-3} units/ml at 50°C and pH 7,90 kD. The fraction exhibited activity toward dyed casein of 0,01 arbitrary units/mg (pH 7). Addition of urea as chaotropic agent resulted in 2-3 fold increase of protease activity as did addition of LAS at both pH 7 and 9 (50°C). The 90 kD protease was neutral with pH optimum at pH 8. The activity is inhibited by phenylmethylsulfonyl fluoride (PMSF) and not by iodoacetic acid, pepstatin A and EDTA which characterises it as a serine type protease.

Also isolated from the bound fraction were 43 kD endoglucanase with pI 4.2 (fraction 33-37) and 25 kD endoglucanase with pI 4.1 (fraction 39-43), 55 kD cellobiohydrolase with pI 4.9 (fraction 39-43) and 65 kD polygalacturonase with pI 4.4 (fraction 39-43). The endoglucanases did not possess activity towards avicel or MUF cellobioside and possessed high activity toward MC, RBB-CMC, CMC41, beta-glucan and endoglucanase. The 25 kD endo did not produce glucose from CMC and the 43 kD endo did. No glucose was formed from avicel. The pH optimum for the 43 kD protein was 4,5 with 70% maximum activity maintained at pH 7.2 and 50% at pH 8. The 43 kD endo kept 70% activity at pH 5 and 6 during 25 hours of incubation. It kept only 10% at pH 7 during this incubation period. The 25 kD endo had pH optimum of activity at pH 5 toward CMC and broad pH optimum of activity toward RBB-CMC with 70% of the maximum activity being kept at pH 9 and with 50% of the maximum activity being at pH 10. It kept 100% activity at pH 5 and 6 and 80% at pH 7, 8, 8.6 and 9.6 during 120 hours of incubation. The 25 kD endo had a temperature optimum of activity at 70°C. The 43 kD endo had a temperature optimum of activity at 60°C. The Michaelis constants towards CMC were 62 and 12,7 g/l for 25 and 43 kD endo respectively. The polygalacturonase is a pectinase. The Michaelis constant toward PGA was 3.8 g/l. The pH optimum of PGU activity is within pH range 5-7 and T optimum within 50-65°C.

Genes encoding *C. lucknowense* proteins were obtained using PCR and characterised by sequence analysis. The corresponding full genes were obtained by screening (partial) gene libraries using the isolated PCR fragments. The full gene of the 43 kD endoglucanase (EG6, Family 6) of the C1 strain has been cloned, sequenced and analysed (including 2.5 kb promoter region and 0.5 kb terminator region). Its nucleotide and amino acid sequences are depicted in SEQ ID No. 1. Predicted molecular weight of the mature protein is 39,427 Da and predicted pI is 4.53, which values correspond well with the measured values. Protein alignment analysis with other glycosyl hydrolases of the family 6.2 shows that C1-EG6 does not include a cellulose-binding domain (CBD) Homology analysis using SwissProt SAMBA software (Smith & Waterman algorithm, Gap penalty 12/2, alignment 10, Blossum62 matrix) shows that C1-EG6 has 51.6% identity with *Fusarium oxysporum* EG-B (over 376 amino acids), 51.0% identity with *Agaricus bisporus* CBH3 (over 353 amino acids), and 50.7%

AMENDED SHEET

04-12-2000

BO 42057

NL 009900618
PCT/NL99/00618

-38-

identity with *Trichoderma reesei* CBH2 (over 367 amino acids). The putative signal sequence runs Met 1 to Arg 28. The promoter contains several potential CreA binding sites, so it is very likely that this promoter would be subject to glucose repression in a fungal strain with working CreA regulation.

Similarly, the full gene of the 25 kD endoglucanase (EG5, Family 45) of the C1 strain has been cloned, sequenced and analysed (including 3.3 kb promoter region and 0.7 kb terminator region). The nucleotide and amino acid sequences are depicted in SEQ ID No. 2. Predicted molecular weight of the mature protein is 21,858 Da and predicted pI is 4.66, which values correspond well with the measured values. This is the smallest fungal endoglucanase known to date. Protein alignment analysis with other glycosyl hydrolases of the family 45 shows that C1-EG5 does not include a cellulose-binding domain (CBD), nor a cohesin/dockerin domain. Homology analysis using NCBI-BLASTP2 software (Gap penalty 11/1, alignment 10, Blosun62 matrix) shows that the closest homologous protein to C1-EG5 is *Fusarium oxysporum* EG-K with 63% identity. The putative signal sequence runs Met 1 to Ala 18. The promoter contains many potential CreA binding sites, so it is very likely that this promoter would be subject to glucose repression in a fungal strain with working CreA regulation.

Furthermore, an additional endoglucanase was found by PCR based on family 12 cellulases homology analysis. The partial nucleotide and amino acid sequence of this additional endoglucanase (EG3, Family 12) is given in SEQ ID No. 3.

The 55kD protein was a cellobiohydrolase (referred to herein as CBH1) with activity against MUF-cellobioside, MUF lactoside, FP and avicel, also against p-nitrophenyl β -glucoside, cellobiose and p-nitrophenyl lactoside. Its activity toward MUF cellobioside is inhibited by cellobiose. The inhibition constant 0,4 mM was determined. The Michaelis constant toward MUF cellobioside was 0,14 mM, toward MUF lactoside was 4 mM and toward CMC was 3,6 g/l. The pH optimum is rather broad from 4,5 to 7. 50% of maximum activity toward CMC and 80% activity toward RBB-CMC is kept at pH 8. 70-80% activity within pH 5-8 is kept during 25 hours of incubation. The temperature optimum is 60-70°C. CBH1 is probably a member of the cellobiohydrolase family 7; its partial nucleotide and amino acid sequences are depicted in SEQ ID No. 4. The partial amino acid sequence depicted corresponds to about amino acids 300-450 from the N terminus of the mature protein. A cellobiohydrolase according to the invention has at least 60%, preferably at least 70%, most preferably at least 80% sequence identity of the partial amino acid sequence of SEQ ID No. 4. The corresponding CBH promoter, which is a preferred embodiment of the invention, can be identified using the partial nucleotide sequence of SEQ ID No. 4. A synergistic effect was observed between 25 kD endo and 55 kD CBH during avicel hydrolysis. Synergism coefficient was maximal at the ratio of 25 kD endo to 55 kD CBH 80:20. The K_{syn} was 1,3 at its maximum.


AMENDED SHEET

Tables A, B and J illustrate the details of the above.

Table J

Purification scheme

Purification of F 60-8 (UF-conc) and F 60-31 CF samples.



04-12-2000

BO 42057

NL 009900618
PCT/NL99/00618

-40-

The expression level of five main *Chrysosporium* genes was studied by Northern analysis. Various strains of *C. lucknowense* were grown in rich medium containing pharmedia with cellulose and lactose (medium 1) or rich medium containing pharmedia and glucose (medium 2) at 33°C. After 48 h, mycelium was harvested and RNA was isolated. The RNA was hybridised with 5 different probes: EG5, EG6, EG3, XylF and CBH. After exposure, the Northern blots were stripped and hybridised again with a probe for ribosomal L3 as a control for the amount of mRNA on the blot. Most strains showed very high response for CBH and high response for XylF in medium 1; in medium 2, half of the strain showed high response for all genes, and the other half showed low response. The order of expression strength was deducted from these data as CBH > XylF > EG5 > EG3 > EG6.

10 Tables A, B and J illustrate the details of the above.

Description of the figures

Figure 1 is a Western blot as described in the Examples

Figure 2 is a pUT720 map

15 Figure 3 is a pUT970G map

Figure 4 is a pUT1064 map

Figure 5 is a pUT1065 map

Figure 6 is a pF6g map

Figure 7 is a pUT1150 map

20 Figure 8 is a pUT1152 map

Figure 9 is a pUT1155 map

Figure 10 is a pUT1160 map

Figure 11 is a pUT1162 map

25 Figure 12: Ion exchange chromatography on Macro Prep Q of NB-fraction after DEAE-Toyopearl of F-60-31 CF sample.

Figure 13: pH dependencies of activity of enzymes from NB fractions of F-60-31 CF sample.

Figure 14: Stability of enzymes from NB fraction of F-60-31 CF sample at pH 5.5 and 7.5 (60°C).

Figure 15: pH stability at 60°C and 50°C of 60 kD Xyl (pI 4.7) from NB fraction of F-60-31 sample.

Figure 16: Temperature dependencies of enzymes from NB fraction of F-60-31 sample.

30 Figure 17: Ion exchange chromatography on Macro Prep Q of bound fractions 50-53 after DEAE-Toyopearl of F-60-8 sample.

Figure 18: pH and temperature dependencies of α -galactosidase activity of F-60-43, UF - conc.

Figure 19: pH dependencies of activity of 48 kD CBH (pI 4.4) from bound fractions of F-60-8 UF - conc.

35 Figure 20: Temperature dependencies of activity towards *p*-nitrophenyl butyrate of F-60-8 UF - conc.

Figure 21: pH dependencies of activity towards *p*-nitrophenyl butyrate of F-60-8 UF - conc.

Figure 22: pH courses of activities of 30kD (pI 8.9) and 90 kD (pI 4.2) proteases toward C1 proteins (50°C, 30 min. incubation).

Figure 23: Effect of 30 kD (pI 8.9) "alkaline" protease on xylanase activity of the NBNB-fraction (Macro Prep Q) of F 60-31 CF at 50°.

5 Figure 24: Effect of 90 kD (pI 4.2) "neutral" protease on CMCase activity of the proteins in the bound fraction #44-45 (DEAE-Toyopearl) of F 60-8 UV-conc sample at 50°C.

Figure 25: Complete hydrolysis of polygalacturonic acid by 65 kD polygalacturonase (pI 4.4): 50°C, pH 4.5; concentration of PGA = 5 g/l, concentration of protein = 0.1 g/l.

Figure 26: pH- and temperature dependencies of polygalacturonase activity of F-60-43 UF-conc.

10 Figure 27: Inhibition of activity toward MUF-cellobioside by cellobiose for 55 kD CBH (pI 4.4): pH 4.5, 40°C.

Figure 28: Synergistic effect between 25 kD Endo (pI 4.1) and 55 kD CBH (pI 4.4) toward avicel (40°C, pH 5, 25 min).

15 Figure 29: Complete hydrolysis of CMC (a) and avicel (b) by the enzymes isolated from bound fractions of F-60-8 UF-conc. sample (50°C, pH 5): concentration of CMC and avicel = 5 g/l, concentration of 25 kD Endo = 0.01 g/l, concentration of 43 kD Endo = 0.02 g/l; 1-25 kD Endo (pI 4.1), 2-43 kD Endo (pI 4.2).

20 Figure 30: Complete hydrolysis of CMC (1) and avicel (2) by 55 kD CBH (pI 4.4) without (a) and with (b) glucono- δ -lactone (50°C, pH 4.5): concentration of CMC and avicel = 5 g/l, concentration of protein = 0.1 g/l, concentration of glucono- δ -lactone = 5 g/l.

Figure 31: pH-Dependence is of CMCase and RBB-CMCase activities of the enzymes isolated from F-60-8 UF-conc. sample: 1-25 kD Endo (pI 4.1), 2-43 kD Endo (pI 4.2).

Figure 32: pH-Dependencies of CMCase and RBB-CMCase activities of 55 kD CBH (pI 4.4).

25 Figure 33: Temperature dependencies of CMCase activity (pH 4.5) of the enzymes isolated from bound fractions of F-60-8 UF-conc. sample: 1-55 kD CBH (pI 4.4), 2-25 kD Endo (pI 4.1), 3-43 kD Endo (pI 4.2).

Figure 34: pH-stability (50°C) of the enzymes isolated from bound fractions of F-60-8 UF-conc. sample: 1-55 kD CBH (pI 4.4), 2-25 kD Endo (pI 4.1), 3-43 kD Endo (pI 4.2).

30 Figure 35: Adsorption of the enzymes isolated from bound fractions of F-60-8 UF-conc. sample.

References

1. Calmels T.P., Martin F., Durand H., and Tiraby G. (1991) *Proteolytic events in the processing of secreted proteins in fungi*. J Biotechnol 17(1): p. 51-66.
2. Punt P.J., Dingemans M.A., Jacobs-Meijnsing B.J., Pouwels P.H., and van den Hondel C.A. (1988) *Isolation and characterization of the glyceraldehyde-3-phosphate dehydrogenase gene of Aspergillus nidulans*. Gene 69(1): p. 49-57.

04-12-2000

BO 42057

NL 009900618
PCT/NL77/00010

-42-

3. Shoemaker S., Schweickart V., Ladner M., Gelfand D., Kwok S., Myambo K., and Innis M. (1983) *Molecular cloning of exo-cellobiohydrolase I derived from Trichoderma reesei strain L27*. Bio/Technology Oct.:691-696.
4. Drocourt D., Calmels T., Reynes J.P., Baron M., and Tiraby G. (1990) *Cassettes of the Streptoalloteichus hindustanus ble gene for transformation of lower and higher eukaryotes to phleomycin resistance*. Nucleic Acids Res 18(13): p. 4009.
5. Mullaney E.J., Hamer J.E., Roberti K.A., Yelton M.M., and Timberlake W.E. (1985) *Primary structure of the trpC gene from Aspergillus nidulans*. Mol Gen Genet 199(1): p. 37-45.
6. Yanisch-Perron C., Vieira J., and Messing J. (1987) *Improved M13 phage cloning vectors and host strains: nucleotide sequences of the M13mp18 and pUC19 vectors*. Gene 33:103-119.
7. Durand H., Baron M., Calmels T., and Tiraby G. (1988) *Classical and molecular genetics applied to Trichoderma reesei for the selection of improved cellulolytic industrial strains*, in *Biochemistry and genetics of cellulose degradation*, J.P. Aubert, Editor. Academic Press. p. 135-151.
8. Lowry O.H., Rosebrough N.J., Farr A.L., and Randall R.J. (1951) *Protein measurements with the folin phenol reagent*. J. Biol. Chem. 7:193-265.
9. Parriche M., Bousson J.C., Baron M., and Tiraby G. *Development of heterologous protein secretion systems in filamentous fungi*. in *3rd European Conference on Fungal Genetics*. 1996. Münster, Germany.
10. Baron M., Tiraby G., Calmels T., Parriche M., and Durand H. (1992) *Efficient secretion of human lysozyme fused to the Sh ble phleomycin resistance protein by the fungus Tolypocladium geodes*. J Biotechnol 24(3): p. 253-266.
11. Jeenes D.J., Marczinke B., MacKenzie D.A., and Archer D.B. (1993) *A truncated glucoamylase gene fusion for heterologous protein secretion from Aspergillus niger*. FEMS Microbiol. Lett. 107(2-3): p. 267-271.
12. Stone P.J., Makoff A.J., Parish J.H., and Radford A. (1993) *Cloning and sequence-analysis of the glucoamylase gene of neurospora-crassa*. Current Genetics 24(3): p. 205-211.
13. Mörsky P. (1983) *Turbidimetric determination of lysozyme with Micrococcus lysodeikticus cells: Reexamination of reaction conditions*. Analytical Biochem. 128:77-85.
14. Paluh J.L., Orbach M.J., Legerton T.L., and Yanofsky C. (1988) *The cross-pathway control gene of Neurospora crassa, cpc-1, encodes a protein similar to GCN4 of yeast and the DNA-binding domain of the oncogene v-jun-encoded protein*. Proc Natl Acad Sci U S A 85(11): p. 3728-32.
15. Nakari T., Onnela M.L., Ilmen M., Nevalainen K., and Penttilä M. (1994) *Fungal promoters active in the presence of glucose*, Patent #WO 94/04673, Alko.
16. Torronen A., Mach R.L., Messner R., Gonzalez R., Kalkkinen N., Harkki A., and Kubicek C.P. (1992) *The two major xylanases from Trichoderma reesei: characterization of both enzymes and genes*. Biotechnology (N Y) 10(11): p. 1461-5.

Chrysosporium sequences

SEQ ID No. 1:
C1-EG6 "43kD" (Family 6) was obtained by PCR based on "43kD Endo" protein sequencing and family 6 cellulases homology analysis.

-2508	GGATCCACACCTACCATACCGGATAGTATGCTACCCAAGTGACATAGG	-2461	creA
	GTTGGTAAAGTAATACGAGAACTCAGAGAGCACTGCCCATATGGCTCGCCAATGACCTCA	-2401	
	AGTGCCAGGTCAGCTTTGCGAGACAGACCTGAGCGCGTCGGATGTGTGACATGGAACGCG	-2341	
	CCGGATCGCCTTGTGATTAAATTATAGGGAAGTAGCGAGGAAGGTTTCAGCAATTGACGT	-2281	
	GAGCGTACATTAAAGCTGTATGATTTCAGGAAGACGAGCCATGGACCAAGTTTCAAGGC	-2221	
	TGAATGGCTTGACGACTTAAGCACCCGAACGAGGAATGAAGAATGAAGAAGTGGGGATCA	-2161	
	TTCTGGCCCCCTCTCGTATGTCGAGTGTAAAGAAAGCGGTTCTACGGAGGACCTAAAGA	-2101	
	GCTCCAAATTTGCTCTGTGAGCTTAAGCCACATATCTCAAGATGAATACATGTCAGGCAT	-2041	
	AGTCACCCCTGATCTTGTTCATCAGTCCACACACTTTTCAGTTCAGCATGTTGATTCCTCA	-1981	
	TCCATATCACTTTCATTAATCTCTTTATGTCTTGGTCAAGACTCCAAGAAACCGAT	-1921	
	AGGTGAGCATCGGTGAGGCTCCCTCAAGGTACCAAGTAGCCATCATCACCGAGTCTGG	-1861	
	GAATGGCGCCGTGCCCGATCTGAGTCTCCAACTCCACGGTACGACGACGACGTCACA	-1801	
	TTGACGCACCAACGTTGAACAAGCAGAGAGGACACGCTTTGTCTACGCGAATCCTGGCAC	-1741	
	TGGATGGAGACGCGTGTGAGCAGGTTTCCGGAACCATGACGGCCTGTCCGGCTTCTCGA	-1681	
	ACAAAGAAAGTGAACACAAAAGAACCGAAACCGAAACGAGGCACGGCATCGACGACCG	-1621	
	GATTGTCCACGGGGACCTCGGCCAGTCAAGCGTTGCCCTGGCCGTGAGTCCCTGGCGA	-1561	
	CGGGGATTCAGCACATCTCACGTTATAGCGACCTCATCCCCCTTCCGCTTGTGCGGTC	-1501	
	GTGCTCCGTGCCGAGTACCCAGGCGTCCGGGCGCTTTAGCCGGGCGGAATCAGAGTC	-1441	creA
	AAGATGCGGCCGAATTGGACGGCAGACGAAGTTTCGTAGAGGTCATGATCGGCACTGAC	-1381	
	GACACCCACCCCTGCGTGATCCCGTGCCCTGGGCTGGGAATTGCCGGCTAATAATCTAC	-1321	
	GGCTTAATAGATATGCACCTTTGCACGCGGTGCAGATAAATAAGCTGTGGTTTCAAAACT	-1261	
	GGCCTCCGTAATTTACCCACCACTGCCGCTTAGCGCCGGGACCTGAGTCTTGGGAGTGC	-1201	
	GCGGAGCGGACCCACCTCGGGTTAGCGTACACACGACGGCTGCATGCGGGATGCCCGG	-1141	creA
	TGCATGGCTTCATAGTACGACAGACCGTCAAGTCCAAATCTGGGTGATGCTTGATGAG	-1081	creA
	ATGACAGCGAGCCCGTCGGCGCACCCCGGCTATGCATCGCGAATTGACAACACTCTCA	-1021	
	GCTCTATTGCGACCCATCGGATAAAAGAAAGAAAGAAAATGGACCTTGAGTACGGGCGT	-961	
	CAGAAACCAAAAAAATCTCGGAACCAATAATGTGGGCATGGCCGGGTGAACGACCG	-901	
	CTACTCCCGGTTCCCTTCTTCGAAACAGAACGCTACAGAGGGTTTCTGGTTTGTCAA	-841	
	GAGTTCGGAGGTCCTCTGCTCCGCGAATGCGTGGTGAAACCCACAGCAGCCATTGTTC	-781	
	GCATGCGTGGCGGACCGTTAGCCGCTGATCGACATGGCGAGCTTCCACCTCAGACCTGG	-721	creA
	AGCAGACGGTTGCGAGGAGCAAGGGCTGCCCTCCCCCTGACGTCGGACCCCAATGACT	-661	
	TCCCCAAACGGGACATCGAGGGTCGTGCATGATGGTGAAAGTAGTTGCAGTATGGAA	-601	
	GTACCCCGGGTTGCCAGGAACCGTTGTTCCGGCCCCCCCCACATTTTCTCTGCCCATGTCAA	-541	
	CTGTGTGTCGTCGAGAGTTTCTGGCTCCGGCCCCCCCCGTCCAATTCCCTAACGGGACCGC	-481	creA

GGGGCATCGCCTGTAACTAATTCCAATGAAGCCGGATATGAGGAGGGAGATTGGATC	-421	area
TGGCAAGCCAGCCATTCTGCTGCGATCGGCACTCGTCCGTACGCCCGCAGTCCATATCCC	-361	
CAAGGCAACTGCTCGGCGGCTCAAGTCTTCTCGGAACGTCCAGCCCGAAGGCGCGC	-301	
GCCAGCACCGCCCTATGTTCTCTGATTGCGATCCTCGATCTCCAGAGACGGGTCACTCG	-241	
CCTCGAGGACGGTGCAGGGCATCGGCTTCGCTTAGAGTCCGGGCTGTGTGTGGTC	-181	
AAGGGAGAAGGCGGCGGCCAAGGTGCTCTCGGCGCACTACCCATCGCCTTTACCC	-121	
CCCTCCCCCAGTATATAAAGATGGCCATCGTCTCCTCGTCTGTTGGGAAGAAAGGA	-61	transc.ini.
TCTCTGACCATGCACCACAGCCTAGCTCTAACCCAGCTTGTGCTGTGTGTTGCCCAGC	-1	
ATGAAGTTCGTGAGTCCGCCACCCCTGGCGTTTCGCCCGCCACGGCCCTCGCTGGCCCTCG	+60	Putative
M K F V Q S A T L A F A A T A L A A P S	20	Signal Seq
CGCACGACTCCCCCAGAAGCCCCCGCAGGCCTCGGCGGCTGCGGTGCGCGTGACGCTC	+120	
R T T P Q K P R Q A S A G C A S A V T L	40	
GATGCCAGCACCAACGTGTTCAGCAGTACACGCTGCACCCCAACAACCTTCTACCGTGCC	+180	
D A S T N V F Q Q Y T L H P N N F Y R A	60	
GAGGTCGAGGCTGCCGCGAGGCCATCTCCGACTCGGCGCTGGCCGAGAAAGGCCGCAAG	+240	
E V E A A A E A I S D S A L A E K A R K	80	
GTCGCCGACGTCGGTACCTTCCTGTGGCTCGACACCATCGAGAACATTGGCCGGCTGGAG	+300	
V A D V G T F L W L D T I E N I G R L E	100	
CCCCGCTCGAGGACGTGCCCTGCGAGAACATCGTGGTCTCGTCACTACGACCTCCCG	+360	
P A L E D V P C E N I V G L V I Y D L P	120	
GGCCGTGACTGCGGGCCAAGGCCTCCAACGGCGAGCTCAAGGTCGGCGAGCTCGACAGG	+420	
G R D C A A K A S N G E L K V G E L D R	140	
TACAAGACCGAGTACATCGACAGtgagttaacccttctgtggcccccttctttcccccgag	+480	Intron 1
Y K T E Y I D	147	
agagcgtctggttgagtggggtgtgagagagaaaaatggggcgagcttaaaagactgacgt	+540	
gttggctcgagAGATCGCCGAGATCTCAAGGCCCACTCCAACACGGCCTTCGCCCTCG	+600	
K I A E I L K A H S N T A F A L	163	
TCATCGAGCCGACTCGCTCCCCAACCTGGTCAACCAATAGCGACCTGCAGACGTGCCAGC	+660	
V I E P D S L P N L V T N S D L Q T C Q	183	

AGAGCGCTTCGGGTACCGGAGGGTGTGCGCTATGCCCTCAAGCAGCTCAACCTCCCCA	+720	Intron 2
Q S A S G Y R E G V A Y A L K Q L N L P	203	
ACGTGGTCATGTACATGCGATGCCGGCCACGGTGGCTGGCTGGGACGCCAACCTCA	+780	
N V V M Y I D A G H G G W L G W D A N L	223	
AGCCCGGCGCCAGAGCTCGCCAGCGTCTACAAGTCTGCTGGTTCCGCCCTCGCAAGTCC	+840	
K P G A Q E L A S V Y K S A G S P S Q V	243	
GCGGTATCTCCACCAACGTGGCTGGTTGGAACGCCCTGgtaagacactctatgtccccctc	+900	
R G I S T N V A G W N A W	256	
gtcgggtcaatggcgagcggaatggcgtgaaatgcatgggtgaccttbtgatcttttccc	+960	
cctcctatagGGACCAGGAGCCCGGTGAGTTCTCGGACGCCCTCGGATGCCCAGTACAACA	+1020	
D Q E P G E F S D A S D A Q Y N	272	
AGTGCCAGAACGAGAAGATCTACATCAACACCTTTGGCGCTGAGCTCAAGTCTGCCGGCA	+1080	
K C Q N E K I Y I N T F G A E L K S A G	292	
TGCCCAACCAAGCCATCATCGACACTGGCCGCAACGGTGTCAACCGGTCTCCGCGACGAGT	+1140	
M P N H A I I D T G R N G V T G L R D E	312	
GGGTGACTGGTGCAACGTCAACGGCGCGCTTCGGTGTGCGCCCGACTGCCAACACTG	+1200	
W G D W C N V N G A G F G V R P T A N T	332	
GCGACGAGCTCGCCGACGCCCTTCGTGTGGTCAAGCCCGTGGCGAGTCCGACGGCACCA	+1260	
G D E L A D A F V W V K P G G E S D G T	352	
GCGACTCGTGGCGCGCTACGACAGCTTCTGCGGCAAGCCCGACGCCCTTCAAGCCCA	+1320	
S D S S A A R Y D S F C G K P D A F K P	372	
GCCCCGAGCCGGTACCTGGAACCAAGCCCTACTTCGAGATGCTCCTCAAGAACGCCAACC	+1380	
S P E A G T W N Q A Y F E M L L K N A N	392	
CGTCCTTCTAAGTCTCCTCGACGGCTTCTGTCTCAGTCGCTCTGACGGTGGTGTGCTGG	+1440	t49
P S F *	395	
TGGTGCCCCCTGCTCCTGTCTGTCTCCGCGGGAGGGAGGCAACGAAATGAAGTC	+1500	t109

-3309
AATCACGAGAAGCTAATTGGGCTCTATAGTATCCGACAAAGATGACCCAGAGCGAGATTGA
GGATCTCGAGGGAACCTGAAGCAGAGCAGCAACAACGACACAGCCTCTCCGCGACCT
GCTCGACAAGATTCCCGATGGCCTCTCTCGCGGCAACAACAATCCAAGCTGACGATAT
CCAGAGCAACGCGAGCGCAGATGGAGAACCTGAGCGTCTCGCCGCGGAACCCGA
GGAGCTGACCAGATACGTCCAGGAAGTGTCCGTGAGATCATGCCGCCATCAAGTTCCA
TGACCAGCTTCTCCAGGACATCTCGAGGCCATCGACAAGATCCCGTGTGCCCAAGAT
TGTGGAGCAGCTGGAGGAGCAGATGTCCATCTTTGTATTCAGATCATGGCCCCGTTCTGT
GGTTCCGCTTATCGAGCAGATCAAGAACGAGCTCGGACTGGCTCCAGCGAGATCATCCA
GAGCAGCAGGGCTGAGCAGCACAACGTCTTTGAGGACGACAACGCCACCGACCCGACTCA
CTCGATGTTGGCCAAGGACCACCTTGTAGTAACGTAAGCCGACCCCTAATCAGAAGCTCGCA
TGTAAGAAATTGAGTTAGACTGACGCGACTTGTTTCCCGTCTCTGTAGATCCTCAACGAGAT
CGCGGTCCGCGCCTCCAAAGTCTGTCTCTGGTCTGTTGACAAGATCATTTACGGAGTGTCCA
GGACGATGACAGCTCGACGTGGACCGCCTGCTTGACAAGATCATTTACGGAGTGTCCA
CCATCCCGCGAGCGCACCATGGGCCCTGAGGGCGCTCCGAGGGCCGGAGCTCATCTT
CAACATGTCGCGAGTGTGGAGGACATGAGCGACGGGCAGCGGACGAGTACCGGGG
CAAGCTGAGCCGCGAGGAGTCGAGAGAGGCGACAACACCGGAGGGCCAGCACGACTG
CGGCCACGGCTGCGGGGCAAGCTCAAGATGCACAAGAACTTCGGAAACGAGCGCCCCA
GACGGTAGAGGACCAGATCGCGGGCGCCGCGGAGGCCATCATGGAGGCGTCAAGCA
GGGCCCTGTCAGAGCCGTGCAGAACGCCCGCGCCAGGAGTCGTCGGAGAGCAGCGG
CCTGGGTGGTTTCATCAGCAGCGTCGCGGGCGGCTCTCTGGCGGCGCTCAAGAGGA
CGAGACAGAGTCGTACCAAGCCGGCGCGCACCGAGACGGCGGTACACGCAGACCAC

GACCGAGTACGGCTACTCCGAGGCCGCTACGGCCAGGCCCAGTACACGGAGACGCAGTA	-1981	
CGCGGGCGGCGGCGGCCGAGCGAGTACCGCCGTACGAGCAGCGCGAGGATGATGA	-1921	
CGCCCGGTCCAGAGCTACGGATACACGGAACAGCGCACCGAGACGCGCTACGACAGCTA	-1861	
CTCGGTGCTATGGCGCCGCGAGAGACACAGCAGCTATGGCGCGCGCGCAGCGCGAG	-1801	
CGAATACATTCTAGTCTCCAGCAGAGTAGCTACCGTGGCAGCGCTATGGCAGTGGGTA	-1741	
CGTCTGTCGTGATGAAGAGAGAGCAGCGGCTATGGAAGTGGTTACGGTCGTCGTGATGA	-1681	
AGAGGAGAGTGGTGGTTATGGTGGCGCTATGGCCCGCTCAGGAAGAAGAGAGTAGCAG	-1621	
CTATGGAAGCGGTTATGGTCTGTCGTCTGATGAAGAGAGCGCGGTTATGGTGGTGG	-1561	
CTACGGCCCGCTCAGGAAGAAGAGAGTAGCGGCTATGGAAGTGGTTACGGTCGTCGTGA	-1501	
TGAAGAAGGAGCGCGGTTATGGTGGTGGCTACGGCCCGCTCATGAGGAAGAGAGCAG	-1441	
TGGTTACGGCAGCGGCTATGGTCTGTCGCTATGAAGAGGCGGCTACGGCAGTGG	-1381	
TTACGGCCCGCGGCAACGACGAGGAGGAAGAGGAGTGGCGACGCCGAGGTGGG	-1321	creA
TTACTAGGGTGAACCTCTCCGGCCGGTCTTGTGTGAACCTTGTCTGTCATGGGCAG	-1261	
GACCGGTGCATCATGAACAGGACGGTGGCTGTGTTTTTTTTTCTCGGGTCTTGATTG	-1201	
TTTTGTTGAATCTCCCTTTTCGAGGATACGAGCTCTCTCGGGACGAATAGATGAAGCAA	-1141	nit2
TCTGACAGATTTGCTCTCAAAAAGAACTGATATCTCTCCACCATGCATGTATGTACA	-1081	
TTACATACATTATCCCCCTCCACTGGATTGACAAACGGAAGCAATGGCGCTGATTC	-1021	
AAGAACCATCAGGGCTGTCTATGGCTTGTGTCGCGTGGCGCGGTGACGCCCACTAT	-961	
GACTCTCTGGGCAGCGGCAACTGGTGCCAGATATATTAATCCGGGCATAGCGCATAT	-901	creA
CTTCCCTTGATTTGTAGAGTACTAGTACACTAACCCCTTCTCCACATGGGCCACTGTTT	-841	nit2
GGTAGATCTGCCCCAAGTGCAAGTGCGGGGGGCCAACTAGTAATATCCTCCGCTC	-781	creA
TCCCGAGTGGCGGACTAACCGTCAATTGCTCCAGAGGCTTGCACTCTATCGCAGGCCCT	-721	nit2
TTCCAATAAGGATGGGCGTTTCGGCGGTGATGATCCCGTCTGTCGGGCATACGGGGAG	-661	creA
GGTAGATAGAAATAACGACGCTGGTGTGTTGGAGAGGGGAGGACTATTAGGGGAGG	-601	creA nit2
GAAATACAGGGCAGGGGTGAGACGGGTGACGTTCCGGCGGAACCTCGCGCTTGTCAAA	-541	
CAAGCAGCCCTGTTAGTTGCTCTAGACTAGTGTACATACATACATATGTACATACTGTA	-481	
TGTACTGCACATACTTTAACTTGGTCTTCCCTGTGAGCCGCCAGGAACATCACAACTGC	-421	
AAGCGGAAAAGCCCCCATATACGGGGCGGCTTGTGCGGATGGCTCCCCCTTCGGAACGG	-361	
GTCTGACTCCGAGGATTTTACCCTGCTTCACTTGGGTATTCGCGATGGCCTGTTCAACC	-301	
CTTCCCCCTGGCCGAACCGTTTCTTGGCTCGATCCTAGTGTACACTACACTACTCGTAGAC	-241	
TGCCCTGCCCGACGATCCGCGGAACGCGGAGTGTGAGTGGAGACGGCGCGGCTG	-181	creA
ATGTCGTGTAATTAAATATATAAGTGAGAGTGTGTTTGTGACTGCCCGGTTCTGCTAGT	-121	TATA box
TGAAGGGAAGTTCGATGCTCTCTGCTGCTGCTCTCTGCTGCTCTCTGCTGCGCATCCTCC	-61	
ATCCGTCCGCCTTTGATAACCCGCTCCCGACTCAGTCAAGACGACGCATACTTGGCACC	-1	
ATGCATCTCTCCGCCACCAACCGGTTCTCGCCCTCCCGGCCCTGGCCCTGGCCCAAGCTC	+60	Put .SS
M H L S A T T G F L A L P A L A L A Q L	20	
TCGGGCAGCGGCCAGACGACCCGGTACTGGGACTGTGCAAGCCGAGCTGCGCCTGGCCC	+120	
S G S G Q T T R Y W D C C K P S C A W P	40	

GGCAAGGGCCCCCTCGTCTCCGGTGCAGGCCTGCGACAAGACGACAACCCGCTCAACGAC G K G P S S P V Q A C D K N D N P L N D	+180 60	
GGCGGCTCCACCCGGTCCGGTGCAGCGGGCGGCGCCTACATGTGCTCCTCCCAG G G S T R S G C D A G G S A Y M C S S Q	+240 80	
AGCCCCCTGGCCGTCAGCGACGAGCTGTCTGTACGGCTGGGGCGCCGTCAAGCTCGCCGGC S P W A V S D E L S Y G W A A V K L A G	+300 100	
AGCTCCGAGTCGCGAGTGGTGTGCGCCTGTCTACGAGCTGACCTTACACAGCGGCCGGTC S S E S Q W C C A C Y E L T F T S G P V	+360 120	
GCGGGCAAGAAGATGATTGTGCAGGCGACCAACACCGGTGGCGACCTGGGCGACAAACCAC A G K K M I V Q A T N T G G D L G D N H	+420 140	
TTTGACCTGGCCGtgagttggcctcccccttctccccggaccgctcagattagatgagatta F D L A	+480 Intron 1 144	
gactttgctcgtaaatcgggtccaagattcccttgactgaccaaaacatcatcaggga	+540	
gATCCCCGGTGGCGGTGTCGGTATTTTCAACGgtaagctggtgcccccgaccctcccc I P G G G V G I F N	+600 Intron 2 154	
ggacccctccccctttctccagcgagccgagttgggatacgcgagatcgagaaactcac	+660	
acaacttctctctcgacagCCTGCACCGACAGTACGGCGCTCCCCCGAACGGCTGGGGC A C T D Q Y G A P P N G W G	+720 168	
GACCGCTACGGCGCATCCATTCCAAGGAAGAGTGCGAATCCTTCCGGAGGCCCTCAAG D R Y G G I H S K E E C E S F P E A L K	+780 188	
CCCGGCTGCAACTGGCGCTTCGACTGgtacgttgctttgacataccggaacccaattcct P G C N W R F D W	+840 Intron 3 197	
ccaacccccccccctttctccccaaactccggggtagtcggaatgtcgcgactgaccct	+900	
atttcagGTTCCAAACGCCGACAAACCGTCGGTCACCTTCCAGGAGGTGGCCCTGCCCGT F Q N A D N P S V T F Q E V A C P	+960 214	

CGGAGCTCACGTCCAAGACGGGCTGCTCCCGTTAAGAGGGAAGAGAGGGGCTGGAAGGA	+1020	t25
S E L T S K S G C S R *	225	
CCGAAAGATTCAACCCTCTGCTCCTGCTGGGAAGCTCGGGCGGAGTGTGAAACTGGTG	+1080	t85
AAATATTGTGGCACACAAGCTACTACAGTCCTCGCTCGCCGTCCGGCTAACTAGCCTTG	+1140	t145
CTGCGGATCTGTCCATCTTCGGTCCGAACGTGTCCGTTGCTGTTTTGGCTCGGTGCCTCAT	+1200	t205
CCTCTCCC AACCTAGTCAAGAATGAATCGTGAGAGAGGCTGAGAGAGATAAGATCGACTT	+1260	t265
CAGAAATCCAGGGTTGAAAGCAAATAAAAAAATTCCTGTGGATGAATATCTCGTGATGC	+1320	poly ^M
AACGACCCCTCTAGGAAACCTTGACGAAATTTGCTGACGGCAAAATCTTCAAAGACTCGT	+1380	t385
TAAACGGTCGCCCGTAGTGGTCTGTGCCCCCAATCCGTTTGTGTTGAAATGACATTGCC	+1440	t445
CGTAACGCCGGACTCATATCAACTGCGTACCGAAAGCCAATCCCTCCCCAACACGCCCT	+1500	t505
CTCTAATAAGCTCTCCAAACAAGACCTCTTGAGACAGAAAATAACGCCAGATGCTGAGG	+1560	t565
ACTTGACAAGCCGGGGGGGGGGCTTGTCAAGTGCAAAAACTTGCCCATTTTCATGC	+1620	t625
TGGTATCAAAAAAACA AAAAAAAAAAACAATTTCAAGTCGCGGATGCCCATTTACAT	+1680	t685
TGCTTGGGTGGCCCAATAGAAACTTGCAACACGTCAGTGTCACTTGCACGCCCTTGG	+1737	t742

SEQ ID No. 3:

C1-EG3 (Family 12) gene fragment obtained by PCR based on family 12 cellulases homology analysis.

GAATTGGGGATTACGAGCTAATGATCTGGtcagtttttttctttctttctttctttctttctcctgttttatattcttta 100
g d y e l m i w
tccattgcttgccctctttccttaaccctgctgactctcttcttgtcaatgatactgtaataaggctggcgagatttcggcgacgtctacctcatcggc 200
L A R F G D V Y P I G
TCGTCCCAGGGCCACGTCAACGTGGCGGCCAGGACTGGGAGCTGTGGACGGGCTTCAANGGNAACATCGGGTCTACAGCTTCGTAGCGCCCCANCCCC 299
S Q G H V N V A G Q D W E L W T G F X G N M R V Y S F V A P X P
CGCAACAGNTTCAGCGCCAACGTCAAGGACTTCTTCAACTATCTCCAGTCCAAACCAGGGCTTCCCGGCAGCAGCCAATACTTCTCAAgtaaggagacga 400
r n x f s a n v k d f f n y l q s n q g f p a s s q y l l n?
gatctcgaacagcataccatatatgctgcggtacaaagtgcactaaccccccttttttcccgttcgcagctCTCCAGTTCCGGCACTG 487
? ? ? ?

WO 00/20555

PCT/NL99/00618

-50-

SERQ ID No. 4:

Chrysosporium cellobiohydrolase CBH1

```
TTTNGGGCGCGTCTTACTCCTACCTTGACCGTGATCGGCCAGTCGCGCTGCGAGGGCG
1  ? G A V L L L P C T V I G Q S R C E G

ACTCGTGGGGGTACCTACAGCACCGACCGCTATGCCGGCATCTGCCACCCCGACGGAT
61 D S C G G T Y S T D R Y A G I C D P D G

GCGACTTCAACTCGTACCGCCAGGGCAACAAGACCTTCTACGGCAAGGCATGACGGTCG
121 C D F N S Y R Q G N K T F Y G K G M T V

ACACGACCAAGAAGATCAGGTGTCGTACCCAGTTCCTCAAGAACTCGCCGCGAGCTCT
181 D T T K K I T V V T Q F L K N S A G E L

CCGAGATCAAGCGGTTCTACGTCCAGAACGGCAAGGTCTATCCCCAACTCCGAGTCCACCA
241 S E I K R F Y V Q N G K V I P N S E S T

TCCCGGGCGTCGAGGGCAACTCCATCACCCAGGACTGGTGCGACCGCCAGAAGCCGCCT
301 I P G V E G N S I T Q D W C D R Q K A A

TCGGCGACGTGACCGACTTCCAGACAAGGCGGCATGCTCCAGATGGGCAAGGCCCTCG
361 F G D V T D F Q D K G G M V Q M G K A L

CGGGGCCCATGGTCCTCGTCATGTCCATATGGGACGACCCAGCCAGTNAACA
421 A G P M V L V M S I W D D H A S ? -
```

WO 00/20555

PCT/NL99/00618

-51-

SEQ ID No. 5:

Chrysosporium xylanase F

```
1      TGACCTTCTCCTCCTTCTCCCGAACAAATAATAGATAATTACGAGCCGGTTCGAGGCTGAC
61     ATTGCGCGATTCTAGCGAGCCGCAATCAATTCAACTTTGCCAACGCCGACGCGGTTGTC
      S R N Q F N F A N A D A V V
120    AACTTTGCCAGGCCAACGGCAAGCTCATCCGGGGCCACACCCTCCTCTGGCACTCTCAG
      N F A Q A N G K L I R G H T L L W H S Q
180    CTGCCGAGTGGTGCAAGACATCAACGACCGCAACACCTTGACCCAGGTCATCGAGAAC
      L P Q W V Q N I N D R N T L T Q V I E N
240    CAGGTCACCAACCCTTGTCACCTCGCTACAAGGCAAGATCCTCCACTGGGACGTCGTTAAC
      H V T T L V T R Y K G K I L H W D V V N
300    GAGATCTTTGCCGAGGACGGCTCGCTCCGCGACAGCGTCTTCAGCCGCGTCCTCGGCGAG
      E I F A E D G S L R D S V F S R V L G E
360    GACTTTGTGGCATCGCCTTCCGGCGCGCGCGCGCGATCCCAACGCCAAGCTCTAC
      D F V G I A F R A A R A A D P N A K L Y
420    ATCAACGACTACAGTCGACA
      I N D Y R S T -
```

SEQUENCE LISTING

<110> Emalfarb, Mark Aaron

<120> Transformation system in the field of filamentous fungal hosts:
in Chrysosporium

<130> 065355-286935

<140> 2,345,356

<141> 1999-10-06

<150> PCT/EP98/06496

<151> 1998-10-06

<160> 16

<170> PatentIn version 3.5

<210> 1

<211> 4376

<212> DNA

<213> Chrysosporium

<220>

<221> CDS

<222> (2509) .. (2949)

<220>

<221> Intron

<222> (2950) .. (3059)

<220>

<221> CDS

<222> (3060) .. (3386)

<220>

<221> Intron

<222> (3387) .. (3479)

<220>

<221> CDS

<222> (3480) .. (3899)

<400> 1

```

ggatccacac ctaccatacc ggatagtatg ctaccaagt gacatagggt tggtaaagta      60
atacgagaac tcagagagca ctgcccataat ggctcgccaa tgacctcaag tgccagggtca    120
gctttgcgag acagacctga gcgcgtcgga tgtgtgacat ggaacgcgcc ggatcgctt      180

```

gttgattaat tataggggaag tagcgaggaa ggtttcagca attgacgtga gcgtacatta	240
aaagctgtat gatttcagga agacgagcca tggaccaggt ttcaaggctg aatggcttga	300
cgacttaagc accgaacgag gaatgaaaga atgaaaagtg ggggatcatt ctggcccctc	360
ctcgatatgtc gagtggttaa gaaggcgggt ctacggagga cctaaagagc tccaatttgc	420
tctgttgagc ttaagccaca tatctcaaga tgaatacatg tcaggcatag tcaccctgat	480
cttgttcatc agtccacaca cttttcagtt cagcatgttg attcctcatc catatcactt	540
tccattacta tctctttatg tccttggtca agactccaag gaaccgatag gtgagcatcg	600
gtgaggctcc ctcaaggtag caaagtagcc atcatcaccg aggtctggga atggcgccgt	660
gcccgatctg agtcctccaa ctccacggta cgacgacagc acgtcacatt gacgcaccac	720
ggttgaacaa gcagagaggg acacgtcttg ctacgcgaat cctggcactg gatggagacg	780
cgtgtgagca ggtttccgga accatgacgg cctgggccgg cttctcgaac aaagaagtgg	840
aacacaaaaa gaaccgaaac ggaaacgcag gcacggcatc gacgaccgga ttgtcccacg	900
gggacctcgg ccagtcaagc gttgccctgg ccgtcagctc cctggcgacg gggattcagc	960
acatctcacg ttataggcga cctcatcccc cttccgtctt gtgcggtcgt tgctccgtgc	1020
cgagtacca ggcgtgccgg ggcctttagc cggggcgga tcagagtcaa gatgcggccg	1080
aattggacgg cagacgaagt ttcgtagagg gtcatgatcg gcactgacga caccacccc	1140
tgcgtgatcc cgtggccctg ggctgggaat tgccggctaa taatctacgg cttaatagat	1200
atgcactttg cacgcggtgc agataaataa gctgtggttt caaacactgg cctccgtact	1260
ttaccacca actgccgctt agcgccggga cctgagtctt gggagtgcgc ggagcggcag	1320
ccacctcggg ttagcgtaca cacgacggct gcatgcgggg atgccgcgtg catggcttca	1380
tagtgtacga cagaccgtca agtccaaatc tgggtgatgc ttgatgagat gacagcgagc	1440
cccgtcggcg gcaccccggc tatgcatcgc gaattgacaa cactctcagc tctattgcga	1500
cccatcggat aaaagaagaa gaaaaaatg gaccttgagt acgggcgtca gaaacaaaa	1560
aaaaactccg gaaccaaata tgtcgggcat ggccgggggtg aacgaccgct actccccgtt	1620
cccttcttcg caaacagaac gctacagagg gttttctggt ttgtcaaaga gttcggaggt	1680
cctctgctcc gcgaatgcgt ggtgaacca ccagcagcca ttgttcttgc atgcgtggcg	1740
gaccgttagc cgctgatcga catggcgagc tccccacctc agacctggag cagacggttg	1800

-54-

cgaggagcaa ggggctgccc tccccctgac ggtcggaccc caatgacttc cccaaacggg	1860
gacatcgagg gtcgtgcatg atgggtgaaa gtagttgcag tatgggaagt accccgggtt	1920
gccaggaacc gttgttcggc cccccacatt ttctctctgc catgtcaact gtgtgtcgtt	1980
cgagagttcc tggctccggc cccccgtcca attccctaac gggaccgcgg ggcategcct	2040
gtaactaact tccaaatgaa gccggatatg agggagggag attggatctg gcaagccagc	2100
cattcgctgc gatcggcact cgtccgtcag ccccgagtc catatcccca aaggcaactg	2160
ctcggcgcgg ctcaagtctt cttcggaacg tccagcccga aggcgcgcgc cagcaccggc	2220
cctatgttcc tgattgcgat cctcgatctc cagagacggg tcacctcgcc tcgaggacgg	2280
tgcaggggca tcggcttcgc ttcctagagc tccgggctgt gtgtggtcaa ggggagaagg	2340
cggcggcgc aaggtgcgtc tcggcgcact caccatcgc ctttaccccc ctcccccca	2400
gtatataaaa gatggccatc gtctcctcgt ctgcttgga agaaaggatc tctcgaccat	2460
gcaccacagc ctagctctaa cccagcttgt cgtgtgttgt tgcccagc atg aag ttc	2517
Met Lys Phe	
1	
gtg cag tcc gcc acc ctg gcg ttc gcc gcc acg gcc ctc gct gcg ccc	2565
Val Gln Ser Ala Thr Leu Ala Phe Ala Ala Thr Ala Leu Ala Ala Pro	
5 10 15	
tcg cgc acg act ccc cag aag ccc cgc cag gcc tcg gcg ggc tgc gcg	2613
Ser Arg Thr Thr Pro Gln Lys Pro Arg Gln Ala Ser Ala Gly Cys Ala	
20 25 30 35	
tcg gcc gtg acg ctc gat gcc agc acc aac gtg ttc cag cag tac acg	2661
Ser Ala Val Thr Leu Asp Ala Ser Thr Asn Val Phe Gln Gln Tyr Thr	
40 45 50	
ctg cac ccc aac aac ttc tac cgt gcc gag gtc gag gct gcc gcc gag	2709
Leu His Pro Asn Asn Phe Tyr Arg Ala Glu Val Glu Ala Ala Ala Glu	
55 60 65	
gcc atc tcc gac tcg gcg ctg gcc gag aag gcc cgc aag gtc gcc gac	2757
Ala Ile Ser Asp Ser Ala Leu Ala Glu Lys Ala Arg Lys Val Ala Asp	
70 75 80	
gtc ggt acc ttc ctg tgg ctc gac acc atc gag aac att ggc cgg ctg	2805
Val Gly Thr Phe Leu Trp Leu Asp Thr Ile Glu Asn Ile Gly Arg Leu	
85 90 95	
gag ccc gcg ctc gag gac gtg ccc tgc gag aac atc gtg ggt ctc gtc	2853
Glu Pro Ala Leu Glu Asp Val Pro Cys Glu Asn Ile Val Gly Leu Val	
100 105 110 115	

-55-

atc tac gac ctc ccg ggc cgt gac tgc gcg gcc aag gcc tcc aac ggc Ile Tyr Asp Leu Pro Gly Arg Asp Cys Ala Ala Lys Ala Ser Asn Gly 120 125 130	2901
gag ctc aag gtc ggc gag ctc gac agg tac aag acc gag tac atc gac Glu Leu Lys Val Gly Glu Leu Asp Arg Tyr Lys Thr Glu Tyr Ile Asp 135 140 145	2949
agtgagttaa ccctttgtgg ccccttcttt tcccccgaga gagcgtctgg ttgagtgggg	3009
ttgtgagaga gaaaatgggg cgagcttaaa gactgacgtg ttggctcgca gag atc Glu Ile	3065
gcc gag atc ctc aag gcc cac tcc aac acg gcc ttc gcc ctc gtc atc Ala Glu Ile Leu Lys Ala His Ser Asn Thr Ala Phe Ala Leu Val Ile 150 155 160 165	3113
gag ccc gac tcg ctc ccc aac ctg gtc acc aat agc gac ctg cag acg Glu Pro Asp Ser Leu Pro Asn Leu Val Thr Asn Ser Asp Leu Gln Thr 170 175 180	3161
tgc cag cag agc gct tcc ggc tac cgc gag ggt gtc gcc tat gcc ctc Cys Gln Gln Ser Ala Ser Gly Tyr Arg Glu Gly Val Ala Tyr Ala Leu 185 190 195	3209
aag cag ctc aac ctc ccc aac gtg gtc atg tac atc gat gcc ggc cac Lys Gln Leu Asn Leu Pro Asn Val Val Met Tyr Ile Asp Ala Gly His 200 205 210	3257
ggt ggc tgg ctc ggc tgg gac gcc aac ctc aag ccc ggc gcc cag gag Gly Gly Trp Leu Gly Trp Asp Ala Asn Leu Lys Pro Gly Ala Gln Glu 215 220 225	3305
ctc gcc agc gtc tac aag tct gct ggt tcg ccc tcg caa gtc cgc ggt Leu Ala Ser Val Tyr Lys Ser Ala Gly Ser Pro Ser Gln Val Arg Gly 230 235 240 245	3353
atc tcc acc aac gtg gct ggt tgg aac gcc tgg taagacactc tatgtccccc Ile Ser Thr Asn Val Ala Gly Trp Asn Ala Trp 250 255	3406
tcgtcgggtca atggcgagcg gaatggcgtg aaatgcatgg tgctgacctt tgatcttttc	3466
cccctcctat agg gac cag gag ccc ggt gag ttc tcg gac gcc tcg gat Asp Gln Glu Pro Gly Glu Phe Ser Asp Ala Ser Asp 260 265	3515
gcc cag tac aac aag tgc cag aac gag aag atc tac atc aac acc ttt Ala Gln Tyr Asn Lys Cys Gln Asn Glu Lys Ile Tyr Ile Asn Thr Phe 270 275 280	3563

-56-

ggc gct gag ctc aag tct gcc ggc atg ccc aac cac gcc atc atc gac 3611
 Gly Ala Glu Leu Lys Ser Ala Gly Met Pro Asn His Ala Ile Ile Asp
 285 290 295 300

act ggc cgc aac ggt gtc acc ggt ctc cgc gac gag tgg ggt gac tgg 3659
 Thr Gly Arg Asn Gly Val Thr Gly Leu Arg Asp Glu Trp Gly Asp Trp
 305 310 315

tgc aac gtc aac ggc gcc ggc ttc ggt gtg cgc ccg act gcc aac act 3707
 Cys Asn Val Asn Gly Ala Gly Phe Gly Val Arg Pro Thr Ala Asn Thr
 320 325 330

ggc gac gag ctc gcc gac gcc ttc gtg tgg gtc aag ccc ggt ggc gag 3755
 Gly Asp Glu Leu Ala Asp Ala Phe Val Trp Val Lys Pro Gly Gly Glu
 335 340 345

tcc gac ggc acc agc gac tcg tcg gcg gcg cgc tac gac agc ttc tgc 3803
 Ser Asp Gly Thr Ser Asp Ser Ser Ala Ala Arg Tyr Asp Ser Phe Cys
 350 355 360

ggc aag ccc gac gcc ttc aag ccc agc ccc gag gcc ggt acc tgg aac 3851
 Gly Lys Pro Asp Ala Phe Lys Pro Ser Pro Glu Ala Gly Thr Trp Asn
 365 370 375 380

cag gcc tac ttc gag atg ctc ctc aag aac gcc aac ccg tcc ttc taa 3899
 Gln Ala Tyr Phe Glu Met Leu Leu Lys Asn Ala Asn Pro Ser Phe
 385 390 395

gctcctcgac ggcttcttgc tgtcagtcgc tctgacgggtg gtgtgctggt ggtgccctg 3959

ctcctgctgc tgctgctccg cggggagggg aggcaacgaa aatgaagtcc tgcttcaaaa 4019

caaaacagaa acaagcgagg cgcggtgcaa tggctcgtgcg ttcgtctttt ttcattgttcc 4079

cttctagtgt agtagtttga tagtcgtaca taaggggttt cagaaccgtc tctctgtctc 4139

ggtctttttg cgagttgttg cgactcgtga ttatggcctt tgttgctcgt tgcggcagag 4199

tagaaccaca gcgtgttggg gtagcagctt gctccgtagg acgtagggaa acaacctgag 4259

actctggaat tgcagtcagc ctgcgtcgcc cctctaggaa acgaagggga gaaccagtag 4319

tggctgcagc ttacaaacgc gagcatgggtg aacatctccg agaaaaggga gggatcc 4376

<210> 2

<211> 147

<212> PRT

<213> Chrysosporium

<400> 2

-57-

Met Lys Phe Val Gln Ser Ala Thr Leu Ala Phe Ala Ala Thr Ala Leu
 1 5 10 15

Ala Ala Pro Ser Arg Thr Thr Pro Gln Lys Pro Arg Gln Ala Ser Ala
 20 25 30

Gly Cys Ala Ser Ala Val Thr Leu Asp Ala Ser Thr Asn Val Phe Gln
 35 40 45

Gln Tyr Thr Leu His Pro Asn Asn Phe Tyr Arg Ala Glu Val Glu Ala
 50 55 60

Ala Ala Glu Ala Ile Ser Asp Ser Ala Leu Ala Glu Lys Ala Arg Lys
 65 70 75 80

Val Ala Asp Val Gly Thr Phe Leu Trp Leu Asp Thr Ile Glu Asn Ile
 85 90 95

Gly Arg Leu Glu Pro Ala Leu Glu Asp Val Pro Cys Glu Asn Ile Val
 100 105 110

Gly Leu Val Ile Tyr Asp Leu Pro Gly Arg Asp Cys Ala Ala Lys Ala
 115 120 125

Ser Asn Gly Glu Leu Lys Val Gly Glu Leu Asp Arg Tyr Lys Thr Glu
 130 135 140

Tyr Ile Asp
 145

<210> 3
 <211> 109
 <212> PRT
 <213> Chrysosporium

<400> 3

Glu Ile Ala Glu Ile Leu Lys Ala His Ser Asn Thr Ala Phe Ala Leu
 1 5 10 15

Val Ile Glu Pro Asp Ser Leu Pro Asn Leu Val Thr Asn Ser Asp Leu
 20 25 30

-58-

Gln Thr Cys Gln Gln Ser Ala Ser Gly Tyr Arg Glu Gly Val Ala Tyr
 35 40 45

Ala Leu Lys Gln Leu Asn Leu Pro Asn Val Val Met Tyr Ile Asp Ala
 50 55 60

Gly His Gly Gly Trp Leu Gly Trp Asp Ala Asn Leu Lys Pro Gly Ala
 65 70 75 80

Gln Glu Leu Ala Ser Val Tyr Lys Ser Ala Gly Ser Pro Ser Gln Val
 85 90 95

Arg Gly Ile Ser Thr Asn Val Ala Gly Trp Asn Ala Trp
 100 105

<210> 4
 <211> 139
 <212> PRT
 <213> Chrysosporium

<400> 4

Asp Gln Glu Pro Gly Glu Phe Ser Asp Ala Ser Asp Ala Gln Tyr Asn
 1 5 10 15

Lys Cys Gln Asn Glu Lys Ile Tyr Ile Asn Thr Phe Gly Ala Glu Leu
 20 25 30

Lys Ser Ala Gly Met Pro Asn His Ala Ile Ile Asp Thr Gly Arg Asn
 35 40 45

Gly Val Thr Gly Leu Arg Asp Glu Trp Gly Asp Trp Cys Asn Val Asn
 50 55 60

Gly Ala Gly Phe Gly Val Arg Pro Thr Ala Asn Thr Gly Asp Glu Leu
 65 70 75 80

Ala Asp Ala Phe Val Trp Val Lys Pro Gly Gly Glu Ser Asp Gly Thr
 85 90 95

-59-

Ser Asp Ser Ser Ala Ala Arg Tyr Asp Ser Phe Cys Gly Lys Pro Asp
 100 105 110

Ala Phe Lys Pro Ser Pro Glu Ala Gly Thr Trp Asn Gln Ala Tyr Phe
 115 120 125

Glu Met Leu Leu Lys Asn Ala Asn Pro Ser Phe
 130 135

<210> 5
 <211> 5046
 <212> DNA
 <213> Chrysosporium

<220>
 <221> CDS
 <222> (3310)..(3741)

<220>
 <221> Intron
 <222> (3742)..(3850)

<220>
 <221> CDS
 <222> (3851)..(3880)

<220>
 <221> Intron
 <222> (3881)..(3987)

<220>
 <221> CDS
 <222> (3988)..(4116)

<220>
 <221> Intron
 <222> (4117)..(4217)

<220>
 <221> CDS
 <222> (4218)..(4304)

<400> 5

gcttaggaga atcacgagaa gctaattggg ctctatagta tccgacaaga tgacccagag 60
 cgagattgag gatctcgagg gaaccctgaa gcagagcagc aacaacgaca ccagcctcct 120
 ccgcgacctg ctcgacaaga ttcccgatgg cctcctcggc ggcaacaaca aatccaagct 180

ggacgatatc cagagcaacg cgcaggccgc gcagatggag aacctgagcg tctcgccgcg	240
ggaacccgag gagctgacca gatacgtcca ggaagtgttc cgtcagatca tgcccgccat	300
caagttccat gaccagcttc tccaggacat ctcggaggcc atcgacaaga tcccggtgct	360
gccaagatt gtggagcagc tggaggagca gatgtccatc tttgtattcc agatcatggc	420
cccgttcgtg gttccgctta tcgagcagat caagaacgag ctcgcgactg gctccagcga	480
gatcatccag agcagcaggg ctgagcagca caacgtcttt gaggacgaca acgccaccga	540
cccgactcac tcgatgttgg ccaaggacca ctttagtaac gtaaagccga ccctaatacag	600
aagctcgcat gtagaattga gttagactga cgcgacttgt ttcccgtctc tgtagatcct	660
caacgagatc ggcggtcgcg ccgcctccaa ggtcgtctcc tgggtcgtcc cgcagctcat	720
ggaggcctgg gacgatgaca gcgtcgacgt ggaccgcctg cttgacaaga tcatttacgg	780
agtgttccac catcccgcgc agcgcaccat gggccctgag ggggcgtccg agggccggga	840
gctcatcttc aacatggtgc gcgagtgggt ggaggacatg agcgacgggc agcgcgacga	900
gtaccggggc aagctgagcc gcgagggagt cgagagaggc gacaaccacc gcgagggcca	960
gcacgactgc ggccacggct gcggggggcaa gctcaagatg cacaagaact tccggaacga	1020
ggcgccccag acggtagagg accagatcgc gggcgccgcc gcggaggcca tcatgggagg	1080
cgtcaagcag ggctgtcgc aggccgtgca gaacgccgcc ggccgccagg agtcgtcga	1140
gagcagcggc ctgggtgggt tcatcagcag cgtcgcgggc ggctcctgg gcggcgccct	1200
caagagggac gagacagagt cgtaccaggc cggcggccgc accgaggacg gcgggtacac	1260
gcagaccacg accgagtacg gctactccgg aggccgctac ggccaggccc agtacacgga	1320
gacgcagtac ggcggcggcg gcggcgggcg cagcgagtac cgccgctacg agcagcgca	1380
ggatgatgac ggccgggtcc agagctacgg atacacggaa cagcgcaccg agacgcgcta	1440
cgacagctac tcgggtggct atggcgggcg cgaggagacc agcagctatg gcggcgggcg	1500
cagcgcgagc gaatacatte gtagctccca gcagagtagc tacggtggca gcggctatgg	1560
cagtgggtac ggtcgtcgtg atgaagaaga gagcagcggc tatggaagtg gttacggtcg	1620
tcgtgatgaa gaggagagtg gtggttatgg tggcggtat ggccgccgtc aggaagaaga	1680
gagtagcagc tatggaagcg gttatggtcg tcgtcgtgat gaagaagaga gcggcggtta	1740
tgggtgggtggc tacggccgcc gtcaggaaga agagagtagc ggctatggaa gtggttacgg	1800

-61-

tcgtcgtgat gaagaaggga gcggcgggta tgggtgggtggc tacggccgcc gtcattgagga	1860
agagagcagt ggttacggca gcggttatgg tcgtcgccat gaagaggagg gcggtggcta	1920
cggcagtggt tacggccgcc ggcgcaacga cgaggaggaa gaggaggatg gcggacgccg	1980
gaggtggggg tactaggggtg aactcttccg gccgggtctct tgttgtgaac cttgctgttg	2040
catgggcagg accggtgcat catgaacagg acggtgcgct gtgttttttt tttctcgggg	2100
tcttgattgt ttgttgaatc tcccttttcg aggatacgag ctctctcggg gacgaataga	2160
tgaaggcaat ctgacagatt tgctctcaaa aaaagactga tatctcttcc accatgcact	2220
gtatgtacat tacatacatt atccccctcc actggattcg cacaacggaa agcaatggcg	2280
cgctgattca agaaccatca gggctgtcat tggcttgttt tgtgccgtgg ccgcggtgac	2340
gccactatg actctctggg caggcggcaa ctgggtgcca gatataataa tccggggcat	2400
agcgcatatc ttccttgatt tgtagagtac tagtacacta acccccttct ccacatgggg	2460
ccactgttcg gtagatctgc ccgaagtgca agtgcggggg gggccaaact aggtaatatc	2520
ctcccgtct cccgagtgcg cggactaacc gtcattgctc ccagaggctt gcactctatc	2580
gcaggccttt tccaataagg atggggcggt cggcgggtgat gatgccggtc gtgcggggca	2640
tacggggagg gtagatagaa aataacgacg ctggtgtttt ggagagggga gggggactat	2700
taggggaggg aaatacaggg gcaggggggt agacgggtga cgttccggcg gaacctcgcg	2760
cttgtcaaac aagcagccct gttaggttgc tctagactag tgtacataca tacatatgta	2820
catactgtat gtactgcaca tactttaact tgggtgcttcc ctgtgagccg ccaggaacat	2880
cacaactgca agcggaaaag gcccataata cggggcggct tgtcgggatg gctccccct	2940
tcggaacggg tctgacttcc gaggatttta cctgcttcat ttgggtattc tgcgatggcc	3000
tgttcaaccc ttcccctggc cgaaccgttt cttggctcga tcctagtgtg cactacacta	3060
ctcgtagact gcctgcccga cgatccgcgg gaacgggcca ggagtgtgga gtggagacgg	3120
gcggcgggtga tgcgtgttaa ttaaataatat aagtgagagt gttttttgac tgccccgggt	3180
tctggtagtt gaagggaagt tcgatgctct ctgctgtcgt cgctctcgtc gctctcgtcg	3240
gcattctcca tccgtccgcc tttgataacc cgctccccga ctcagtcaag acgacgcata	3300
cttggcacc atg cat ctc tcc gcc acc acc ggg ttc ctc gcc ctc ccg gcc	3351
Met His Leu Ser Ala Thr Thr Gly Phe Leu Ala Leu Pro Ala	
1 5 10	

ctg gcc ctg gcc cag ctc tcg ggc agc ggc cag acg acc cgg tac tgg	3399
Leu Ala Leu Ala Gln Leu Ser Gly Ser Gly Gln Thr Thr Arg Tyr Trp	
15 20 25 30	
gac tgc tgc aag ccg agc tgc gcc tgg ccc ggc aag ggc ccc tcg tct	3447
Asp Cys Cys Lys Pro Ser Cys Ala Trp Pro Gly Lys Gly Pro Ser Ser	
35 40 45	
ccg gtg cag gcc tgc gac aag aac gac aac ccg ctc aac gac ggc ggc	3495
Pro Val Gln Ala Cys Asp Lys Asn Asp Asn Pro Leu Asn Asp Gly Gly	
50 55 60	
tcc acc cgg tcc ggc tgc gac gcg ggc ggc agc gcc tac atg tgc tcc	3543
Ser Thr Arg Ser Gly Cys Asp Ala Gly Gly Ser Ala Tyr Met Cys Ser	
65 70 75	
tcc cag agc ccc tgg gcc gtc agc gac gag ctg tcg tac ggc tgg gcg	3591
Ser Gln Ser Pro Trp Ala Val Ser Asp Glu Leu Ser Tyr Gly Trp Ala	
80 85 90	
gcc gtc aag ctc gcc ggc agc tcc gag tcg cag tgg tgc tgc gcc tgc	3639
Ala Val Lys Leu Ala Gly Ser Ser Glu Ser Gln Trp Cys Cys Ala Cys	
95 100 105 110	
tac gag ctg acc ttc acc agc ggg ccg gtc gcg ggc aag aag atg att	3687
Tyr Glu Leu Thr Phe Thr Ser Gly Pro Val Ala Gly Lys Lys Met Ile	
115 120 125	
gtg cag gcg acc aac acc ggt ggc gac ctg ggc gac aac cac ttt gac	3735
Val Gln Ala Thr Asn Thr Gly Gly Asp Leu Gly Asp Asn His Phe Asp	
130 135 140	
ctg gcc gtgagttgcc tccccctctc cccggaccgc tcagattaga tgagattaga	3791
Leu Ala	
ctttgctcgt aaatcgggtcc aagattccct tgactgacca acaaacatca tacgggcag	3850
atc ccc ggt ggc ggt gtc ggt att ttc aac ggtaagctgg tgcccccgga	3900
Ile Pro Gly Gly Gly Val Gly Ile Phe Asn	
145 150	
ccccccccg gaccctccc ctttttctc cagcgagccg agttgggata gccgagatcg	3960
agaactcaca caactttctct ctcgaca gcc tgc acc gac cag tac ggc gct ccc	4014
Ala Cys Thr Asp Gln Tyr Gly Ala Pro	
155 160	
ccg aac ggc tgg ggc gac cgc tac ggc ggc atc cat tcc aag gaa gag	4062
Pro Asn Gly Trp Gly Asp Arg Tyr Gly Gly Ile His Ser Lys Glu Glu	
165 170 175	

-63-

tgc gaa tcc ttc ccg gag gcc ctc aag ccc ggc tgc aac tgg cgc ttc 4110
 Cys Glu Ser Phe Pro Glu Ala Leu Lys Pro Gly Cys Asn Trp Arg Phe
 180 185 190 195

gac tgg tacgttgctt tgacataccg gaacccaatt cctccaaccc cccccctttt 4166
 Asp Trp

ctcccccaac tccgggggta gtcggaatgt cgcgactgac cctatttcag g ttc caa 4223
 Phe Gln

aac gcc gac aac ccg tcg gtc acc ttc cag gag gtg gcc tgc ccg tcg 4271
 Asn Ala Asp Asn Pro Ser Val Thr Phe Gln Glu Val Ala Cys Pro Ser
 200 205 210 215

gag ctc acg tcc aag agc ggc tgc tcc cgt taa gagggaagag agggggctgg 4324
 Glu Leu Thr Ser Lys Ser Gly Cys Ser Arg
 220 225

aaggaccgaa agattcaacc tctgctcctg ctggggaagc tcgggcgcga gtgtgaaact 4384

ggtgtaaata ttgtggcaca cacaagctac tacagtccgt ctgcgccgtcc ggctaactag 4444

ccttgctgcg gatctgtcca tcttcgggtcc gaactgtccg ttgctgtttt ggctcgggtgc 4504

ctcatcttct cccaacctag tcaagaatga atcgtgagag aggctgagag agataagatc 4564

gacttcagaa atccagggtt gaaagcaata aaaaaaattc ctgtgggatg aatatctcgt 4624

gatgcaacga ccctcctagg aaaccttgac gaaatttgct gacggcaaatt tcttcaaaga 4684

ctcgттаacc ggtcgcccggt agtgggtcctg ttgccccaat ccgttttgtgt tgaaatgaca 4744

ttgcgcgtaa cgccggactc atatcaactg cgtaccgaaa gccaatccct ccccaaacac 4804

gccctctcta ataagctctc ccaaacaaga cctcttgaga cagaaaatac gccagatgc 4864

tgaggacttg acaagccggg gggggggggg ggcttgtcaa gtgcaaaaac ttgcccattt 4924

catgctggta tcaaaaaaac aaaaaaaaaa aaaaacattt caagtcgcgg atgccccatt 4984

tacattgctt gcgtgcgcca atagaaactt gcaacacgtc agtgtcatct tgcacgcctt 5044

gg 5046

<210> 6

<211> 144

<212> PRT

<213> Chrysosporium

<400> 6

-64-

Met His Leu Ser Ala Thr Thr Gly Phe Leu Ala Leu Pro Ala Leu Ala
1 5 10 15

Leu Ala Gln Leu Ser Gly Ser Gly Gln Thr Thr Arg Tyr Trp Asp Cys
20 25 30

Cys Lys Pro Ser Cys Ala Trp Pro Gly Lys Gly Pro Ser Ser Pro Val
35 40 45

Gln Ala Cys Asp Lys Asn Asp Asn Pro Leu Asn Asp Gly Gly Ser Thr
50 55 60

Arg Ser Gly Cys Asp Ala Gly Gly Ser Ala Tyr Met Cys Ser Ser Gln
65 70 75 80

Ser Pro Trp Ala Val Ser Asp Glu Leu Ser Tyr Gly Trp Ala Ala Val
85 90 95

Lys Leu Ala Gly Ser Ser Glu Ser Gln Trp Cys Cys Ala Cys Tyr Glu
100 105 110

Leu Thr Phe Thr Ser Gly Pro Val Ala Gly Lys Lys Met Ile Val Gln
115 120 125

Ala Thr Asn Thr Gly Gly Asp Leu Gly Asp Asn His Phe Asp Leu Ala
130 135 140

<210> 7
<211> 10
<212> PRT
<213> Chrysosporium

<400> 7

Ile Pro Gly Gly Gly Val Gly Ile Phe Asn
1 5 10

<210> 8
<211> 43
<212> PRT
<213> Chrysosporium

-65-

<400> 8

Ala Cys Thr Asp Gln Tyr Gly Ala Pro Pro Asn Gly Trp Gly Asp Arg
 1 5 10 15

Tyr Gly Gly Ile His Ser Lys Glu Glu Cys Glu Ser Phe Pro Glu Ala
 20 25 30

Leu Lys Pro Gly Cys Asn Trp Arg Phe Asp Trp
 35 40

<210> 9

<211> 28

<212> PRT

<213> Chrysosporium

<400> 9

Phe Gln Asn Ala Asp Asn Pro Ser Val Thr Phe Gln Glu Val Ala Cys
 1 5 10 15

Pro Ser Glu Leu Thr Ser Lys Ser Gly Cys Ser Arg
 20 25

<210> 10

<211> 487

<212> DNA

<213> Chrysosporium

<220>

<221> CDS

<222> (7)..(30)

<220>

<221> misc_feature

<222> (61)..(61)

<223> n is a, c, g, or t

<220>

<221> CDS

<222> (168)..(386)

<220>

<221> misc_feature

<222> (260)..(260)

<223> n is a, c, g, or t

<220>

-66-

<221> misc_feature
 <222> (263)..(263)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (295)..(295)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (308)..(308)
 <223> n is a, c, g, or t

<400> 10

gaattc ggg gat tac gag cta atg atc tgg tcagtttttt ttttcttttt 50
 Gly Asp Tyr Glu Leu Met Ile Trp
 1 5

tcttttcttc ncttttcttt tcttttcctt tctcctgttt tattttctta tccattgctt 110

cgccctcttt ccttaaccct gctgactctc tcttcttgtc aatgatactg taatagg 167

ctg gcg aga ttc ggc gac gtc tac ccc atc ggc tcg tcc cag ggc cac 215
 Leu Ala Arg Phe Gly Asp Val Tyr Pro Ile Gly Ser Ser Gln Gly His
 10 15 20

gtc aac gtg gcc ggc cag gac tgg gag ctg tgg acg ggc ttc aan ggn 263
 Val Asn Val Ala Gly Gln Asp Trp Glu Leu Trp Thr Gly Phe Xaa Gly
 25 30 35 40

aac atg cgg gtc tac agc ttc gta gcg ccc anc ccc cgc aac agn ttc 311
 Asn Met Arg Val Tyr Ser Phe Val Ala Pro Xaa Pro Arg Asn Xaa Phe
 45 50 55

agc gcc aac gtc aag gac ttc ttc aac tat ctc cag tcc aac cag ggc 359
 Ser Ala Asn Val Lys Asp Phe Phe Asn Tyr Leu Gln Ser Asn Gln Gly
 60 65 70

ttc ccg gcc agc agc caa tac ctt ctc aagtaaggag acgagatctc 406
 Phe Pro Ala Ser Ser Gln Tyr Leu Leu
 75 80

gaacagcata ccatatatgc gtgcggtaca agtgcactaa cccctttttt ttcccgttcg 466

cagtcttcca gttcggcact g 487

<210> 11
 <211> 8
 <212> PRT
 <213> Chrysosporium

-67-

<400> 11

Gly Asp Tyr Glu Leu Met Ile Trp
 1 5

<210> 12

<211> 73

<212> PRT

<213> Chrysosporium

<220>

<221> misc_feature

<222> (31)..(31)

<223> The 'Xaa' at location 31 stands for Lys or Asn.

<220>

<221> misc_feature

<222> (43)..(43)

<223> The 'Xaa' at location 43 stands for Asn, Ser, Thr, or Ile.

<220>

<221> misc_feature

<222> (47)..(47)

<223> The 'Xaa' at location 47 stands for Arg or Ser.

<400> 12

Leu Ala Arg Phe Gly Asp Val Tyr Pro Ile Gly Ser Ser Gln Gly His
 1 5 10 15

Val Asn Val Ala Gly Gln Asp Trp Glu Leu Trp Thr Gly Phe Xaa Gly
 20 25 30

Asn Met Arg Val Tyr Ser Phe Val Ala Pro Xaa Pro Arg Asn Xaa Phe
 35 40 45

Ser Ala Asn Val Lys Asp Phe Phe Asn Tyr Leu Gln Ser Asn Gln Gly
 50 55 60

Phe Pro Ala Ser Ser Gln Tyr Leu Leu
 65 70

<210> 13

<211> 472

<212> DNA

<213> Chrysosporium

-68-

<220>
 <221> misc_feature
 <222> (4)..(4)
 <223> n is a, c, g, or t

<220>
 <221> CDS
 <222> (6)..(467)

<220>
 <221> misc_feature
 <222> (468)..(468)
 <223> n is a, c, g, or t

<400> 13

tttng ggc gcc gtc tta ctc cta cct tgc acc gtg atc ggc cag tcg cgc 50
 Gly Ala Val Leu Leu Leu Pro Cys Thr Val Ile Gly Gln Ser Arg
 1 5 10 15

tgc gag ggc gac tcg tgc ggc ggt acc tac agc acc gac cgc tat gcc 98
 Cys Glu Gly Asp Ser Cys Gly Gly Thr Tyr Ser Thr Asp Arg Tyr Ala
 20 25 30

ggc atc tgc gac ccc gac gga tgc gac ttc aac tcg tac cgc cag ggc 146
 Gly Ile Cys Asp Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly
 35 40 45

aac aag acc ttc tac ggc aag ggc atg acg gtc gac acg acc aag aag 194
 Asn Lys Thr Phe Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys
 50 55 60

atc acg gtc gtc acc cag ttc ctc aag aac tcg gcc ggc gag ctc tcc 242
 Ile Thr Val Val Thr Gln Phe Leu Lys Asn Ser Ala Gly Glu Leu Ser
 65 70 75

gag atc aag cgg ttc tac gtc cag aac ggc aag gtc atc ccc aac tcc 290
 Glu Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro Asn Ser
 80 85 90 95

gag tcc acc atc ccg ggc gtc gag ggc aac tcc atc acc cag gac tgg 338
 Glu Ser Thr Ile Pro Gly Val Glu Gly Asn Ser Ile Thr Gln Asp Trp
 100 105 110

tgc gac cgc cag aag gcc gcc ttc ggc gac gtg acc gac ttc cag gac 386
 Cys Asp Arg Gln Lys Ala Ala Phe Gly Asp Val Thr Asp Phe Gln Asp
 115 120 125

aag ggc ggc atg gtc cag atg ggc aag gcc ctc gcg ggg ccc atg gtc 434
 Lys Gly Gly Met Val Gln Met Gly Lys Ala Leu Ala Gly Pro Met Val
 130 135 140

-69-

ctc gtc atg tcc ata tgg gac gac cac gcc agt naaca
 Leu Val Met Ser Ile Trp Asp Asp His Ala Ser
 145 150

472

<210> 14
 <211> 154
 <212> PRT
 <213> Chrysosporium

<400> 14

Gly Ala Val Leu Leu Leu Pro Cys Thr Val Ile Gly Gln Ser Arg Cys
 1 5 10 15

Glu Gly Asp Ser Cys Gly Gly Thr Tyr Ser Thr Asp Arg Tyr Ala Gly
 20 25 30

Ile Cys Asp Pro Asp Gly Cys Asp Phe Asn Ser Tyr Arg Gln Gly Asn
 35 40 45

Lys Thr Phe Tyr Gly Lys Gly Met Thr Val Asp Thr Thr Lys Lys Ile
 50 55 60

Thr Val Val Thr Gln Phe Leu Lys Asn Ser Ala Gly Glu Leu Ser Glu
 65 70 75 80

Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro Asn Ser Glu
 85 90 95

Ser Thr Ile Pro Gly Val Glu Gly Asn Ser Ile Thr Gln Asp Trp Cys
 100 105 110

Asp Arg Gln Lys Ala Ala Phe Gly Asp Val Thr Asp Phe Gln Asp Lys
 115 120 125

Gly Gly Met Val Gln Met Gly Lys Ala Leu Ala Gly Pro Met Val Leu
 130 135 140

Val Met Ser Ile Trp Asp Asp His Ala Ser
 145 150

<210> 15

-70-

<211> 440
 <212> DNA
 <213> Chrysosporium

<220>
 <221> CDS
 <222> (78) .. (440)

<400> 15

tgaccttctc ctccttctcc cgaacaataa tagataatta cgagccgggtt cgaggctgac 60

attgcgcgat tctagcg agc cgc aat caa ttc aac ttt gcc aac gcc gac 110
 Ser Arg Asn Gln Phe Asn Phe Ala Asn Ala Asp
 1 5 10

gcg gtt gtc aac ttt gcc cag gcc aac ggc aag ctc atc cgc ggc cac 158
 Ala Val Val Asn Phe Ala Gln Ala Asn Gly Lys Leu Ile Arg Gly His
 15 20 25

acc ctc ctc tgg cac tct cag ctg ccg cag tgg gtg cag aac atc aac 206
 Thr Leu Leu Trp His Ser Gln Leu Pro Gln Trp Val Gln Asn Ile Asn
 30 35 40

gac cgc aac acc ttg acc cag gtc atc gag aac cac gtc acc acc ctt 254
 Asp Arg Asn Thr Leu Thr Gln Val Ile Glu Asn His Val Thr Thr Leu
 45 50 55

gtc act cgc tac aag ggc aag atc ctc cac tgg gac gtc gtt aac gag 302
 Val Thr Arg Tyr Lys Gly Lys Ile Leu His Trp Asp Val Val Asn Glu
 60 65 70 75

atc ttt gcc gag gac ggc tcg ctc cgc gac agc gtc ttc agc cgc gtc 350
 Ile Phe Ala Glu Asp Gly Ser Leu Arg Asp Ser Val Phe Ser Arg Val
 80 85 90

ctc ggc gag gac ttt gtc ggc atc gcc ttc cgc gcc gcc cgc gcc gcc 398
 Leu Gly Glu Asp Phe Val Gly Ile Ala Phe Arg Ala Ala Arg Ala Ala
 95 100 105

gat ccc aac gcc aag ctc tac atc aac gac tac agg tcg aca 440
 Asp Pro Asn Ala Lys Leu Tyr Ile Asn Asp Tyr Arg Ser Thr
 110 115 120

<210> 16
 <211> 121
 <212> PRT
 <213> Chrysosporium

<400> 16

-71-

Ser Arg Asn Gln Phe Asn Phe Ala Asn Ala Asp Ala Val Val Asn Phe
1 5 10 15

Ala Gln Ala Asn Gly Lys Leu Ile Arg Gly His Thr Leu Leu Trp His
20 25 30

Ser Gln Leu Pro Gln Trp Val Gln Asn Ile Asn Asp Arg Asn Thr Leu
35 40 45

Thr Gln Val Ile Glu Asn His Val Thr Thr Leu Val Thr Arg Tyr Lys
50 55 60

Gly Lys Ile Leu His Trp Asp Val Val Asn Glu Ile Phe Ala Glu Asp
65 70 75 80

Gly Ser Leu Arg Asp Ser Val Phe Ser Arg Val Leu Gly Glu Asp Phe
85 90 95

Val Gly Ile Ala Phe Arg Ala Ala Arg Ala Ala Asp Pro Asn Ala Lys
100 105 110

Leu Tyr Ile Asn Asp Tyr Arg Ser Thr
115 120

Claims

1. A mutant *Chrysosporium* strain comprising a nucleic acid sequence encoding a polypeptide of interest, said nucleic acid sequence being operably linked to an expression-regulating region selected from the group consisting of a promoter sequence associated with cellulase expression, xylanase expression, or *gpdA* expression; and optionally a secretion signal sequence, said mutant strain expressing said polypeptide of interest at a higher level than the corresponding non-mutant strain under the same conditions.
2. The mutant *Chrysosporium* strain according to claim 1, said mutant being obtained by recombinant methods comprising stable introduction of at least one heterologous nucleic acid sequence selected from heterologous polypeptide-encoding nucleic acid sequences, heterologous signal sequences and heterologous expression-regulating sequences.
3. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest is a homologous polypeptide which is expressed at a higher level than in the corresponding non-mutant strain under the same conditions.
4. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest is selected from carbohydrate-degrading enzymes, proteases, lipases, esterases, other hydrolases, oxidoreductases and transferases.
5. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest is selected from fungal enzymes allowing production or overproduction of primary metabolites, organic acids, secondary metabolites, and antibiotics.
6. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest is inactivated at a pH below 6.
7. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest exhibits at a pH above 6 at least one of optimal activity or stability.

8. The mutant *Chrysosporium* strain according to claim 7, wherein said polypeptide of interest at a pH above 6 also exhibits at least one of more than 70% of its activity or more than 70% stability.
9. The mutant *Chrysosporium* strain according to claim 1, wherein said polypeptide of interest at a pH above 6 exhibits at least one of more than 70% of its activity or more than 70% stability.
10. The mutant *Chrysosporium* strain according to claim 1, comprising a heterologous signal sequence.
11. The mutant *Chrysosporium* strain according to claim 1, wherein said secretion signal sequence is a fungal signal sequence.
12. The mutant *Chrysosporium* strain according to claim 11, wherein the fungal signal sequence is a signal sequence of a cellulase, β -galactosidase, xylanase, pectinase, esterase, protease, amylase, polygalacturonase or hydrophobin.
13. The mutant *Chrysosporium* strain according to claim 1, further comprising a selectable marker.
14. The mutant *Chrysosporium* strain according to claim 13, wherein the selectable marker confers resistance to a drug or relieves a nutritional defect.
15. The mutant *Chrysosporium* strain according to claim 1, comprising a heterologous expression-regulating region.
16. The mutant *Chrysosporium* strain according to claim 1, comprising a fungal expression-regulating region.
17. The mutant *Chrysosporium* strain according to claim 16, wherein the expression-

regulating region comprises an inducible promoter.

18. The mutant *Chrysosporium* strain according to claim 16, wherein the expression-regulating region comprises a high expression promoter.

19. The mutant *Chrysosporium* strain according to claim 1, said mutant being obtained by mutagenesis steps, the steps including at least one step chosen from the group consisting of UV irradiation and chemical mutagenesis.

20. The mutant *Chrysosporium* strain according to claim 19, wherein the mutagenesis steps comprise a first UV irradiation step, a N-methyl-N'-nitro-N-nitrosoguanidine treatment step, and a second UV irradiation step.

21. The mutant *Chrysosporium* strain according to claim 1, wherein said mutant *Chrysosporium* strain is a mutant *Chrysosporium lucknowense* strain.

22. The mutant *Chrysosporium* strain according to claim 21, wherein said mutant is or is a mutant of a strain selected from the group consisting of *C. lucknowense* strain C1 (VKM F-3500 D), UV13-6 (VKM F-3632 D), NG7C-19 (VKM F-3633 D), and UV18-25 (VKM F-3631 D).

23. The mutant *Chrysosporium* strain according to claim 1, wherein, when *Trichoderma reesei* and said *Chrysosporium* strain are cultured under conditions optimal for each strain, when the *Trichoderma* culture attains a viscosity of 200-600 cP, said *Chrysosporium* strain exhibits a biomass of less than half that of the *Trichoderma*.

24. The mutant *Chrysosporium* strain according to claim 1, said strain producing at least the amount of cellulase in moles per liter as produced by any of the *Chrysosporium lucknowense* mutant strains C1 (VKM F-3500 D), UV13-6 (VKM F-3632 D), NG7C-19 (VKM F-3633 D), and UV18-25 (VKM F-3631 D).

25. The mutant *Chrysosporium* strain according to claim 1, said strain producing less protease than produced by the *Chrysosporium lucknowense* strain C1 (VKM F-3500 D).

-75-

26. The mutant *Chrysosporium* strain according to claim 25, wherein the strain produces less than half the amount of protease produced by the C1 strain.


Fig 1


Fig 2


Fig 3


Fig 4


Fig 5


Fig 6


Fig 7


Fig 8


Fig 9




Fig 10

Fig 11


12/35

Fig 12


13/35

Fig 13a


45kD endo (pI 6.0)

**Fig 13b**


55 kD endo (pI 4.9)

**Fig 13c**


30kD Xyl (pI 9.1)

**Fig 13d**

51kD Xyl (pI 8.7)

**Fig 13e**

60 kD Xyl (pI 4.7)




Xylanase


14/35

Fig 14a


45 kD Endo (pI 6.0)

**Fig 14b**


55 kD Endo (pI 4.9)



**Fig 14c**

30 kD Xyl (pI 9.1)






**Fig 14d**

51 kD Xyl (pI 8.7)








16/35

Fig 16a**Fig 16b****Fig 16c****Fig 16d****Fig 16e**


17/35




18/35

Fig 18a*Fig 18b*


19/35

Fig 19


20/35







21/35









22/35




23/35

Fig 23a*Fig 23b**Fig 23c**Fig 23d*



24/35




25/35

Fig 25

26/35

Fig 26a*Fig 26b*

27/35



28/35










Fig 28

Fig 29a*Fig 29b**Fig 29c**Fig 29d*

30/35



31/35

Fig 31b









Fig 31a



32/35



33/35

Fig 33a*Fig 33b**Fig 33c*

34/35




Fig 34a*Fig 34b**Fig 34c*

Fig 35