

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200124649 B2**
(10) Patent No. **783545**

(54) Title
Universal procedure for refolding recombinant proteins

(51)⁶ International Patent Classification(s)
C07K 001/14

(21) Application No: 200124649 (22) Application Date: 2000 .12 .28

(87) WIPO No: W001/55174

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/177836	2000 .01 .25	US
60/178368	2000 .01 .27	US
60/210292	2000 .06 .08	US
60/210306	2000 .06 .08	US

(43) Publication Date : 2001 .08 .07

(43) Publication Journal Date : 2001 .10 .18

(44) Accepted Journal Date : 2005 .11 .10

(71) Applicant(s)
Oklahoma Medical Research Foundation

(72) Inventor(s)
xinli Lin

(74) Agent/Attorney
PIZZEYS,GPO Box 1374,BRISBANE QLD 4001

(56) Related Art
WEI C. ET AL; BIOCHEM J. (1999) 340, 345-351

AU 200124649

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55174 A3

- (51) International Patent Classification⁷: C07K 1/113
- (21) International Application Number: PCT/US00/35632
- (22) International Filing Date:
28 December 2000 (28.12.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/177,836 25 January 2000 (25.01.2000) US
60/178,368 27 January 2000 (27.01.2000) US
60/210,292 8 June 2000 (08.06.2000) US
60/210,306 8 June 2000 (08.06.2000) US
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- (71) Applicant: OKLAHOMA MEDICAL RESEARCH FOUNDATION [US/US]; 825 N.E. Thirteenth Street, Oklahoma City, OK 73104 (US).
- (72) Inventor: LIN, Xinli; 1201 Charlton, Edmond, OK 73003 (US).
- (74) Agents: PABST, Patrea, L. et al.; Holland & Knight LLP, One Atlantic Center, Suite 2000, 1201 West Peachtree Street, N.E., Atlanta, GA 30309-3400 (US).
- (88) Date of publication of the international search report:
7 March 2002
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: UNIVERSAL PROCEDURE FOR REFOLDING RECOMBINANT PROTEINS

WO 01/55174 A3

(57) Abstract: A universal folding method that has been demonstrated to be effective in refolding a variety of very different proteins expressed in bacteria as inclusion bodies has been developed. Representative proteins that can be dissolved and refolded in biologically active form, with the native structure, are shown in Table I. The method has two key steps to unfold and then refold the proteins expressed in the inclusion bodies. The first step is to raise the pH of the protein solution in the presence of denaturing agents to pH greater than 9, preferably 10. The protein solution may be maintained at the elevated pH for a period of up to about 24 hours, or the pH immediately decreased slowly, in increments of about 0.2 pH units/24 hours, until the solution reaches a pH of about 8.0, or both steps used. In the preferred embodiment, purified inclusion bodies are dissolved in 8 M urea, 0.1 M Tris, 1 mM glycine, 1 mM EDTA, 10 mM beta-mercaptoethanol, 10 mM dithiothreitol (DTT), 1 mM reduced glutathione (GSH), 0.1 mM oxidized glutathione (GSSG), pH 10. The absorbance at 280 nm (OD280) of the protein solution is 5.0. This solution is rapidly diluted into 20 volumes of 20 mM Tris base. The resulting solution is adjusted to pH 9.0 with 1 M HCl and is kept at 4 C for 24 hrs. The pH is adjusted to pH 8.8 and the solution is kept at 4 C for another 24 hrs. This process is repeated until the pH is adjusted to 8.0. After 24 hrs at pH 8.0, the refolded proteins can be concentrated by ultrafiltration and applied to a gel filtration column for purification.

UNIVERSAL PROCEDURE FOR REFOLDING RECOMBINANT PROTEINS

Background of the Invention

5 The present invention is generally in the field of methods for manufacture of recombinant proteins, and especially in the field of refolding of recombinant proteins expressed in the inclusion bodies of procaryotic expression systems such as *E. coli*.

 Expression of recombinant proteins with natural biological activity
10 and structure, referred to as "proteomics", becomes increasingly important with the completion of genomic sequencing for several organisms and the near completion of human genome sequencing. One aspect of proteomics is to express large amounts of protein for structural and functional studies, as well as for commercial applications. The least expensive and most efficient
15 way to express recombinant proteins is to express the proteins in *E. coli*. Proteins are expressed either intracellularly or secreted into the periplasmic spaces. In the former case, the proteins are often deposited in inclusion bodies, especially if the protein has disulfide bonds.

 However, one of the problems in expressing mammalian proteins in
20 *E. coli* is that most of the expressed proteins form insoluble inclusion bodies. While this problem can be circumvented by using various mammalian or insect expression systems, growing *E. coli* is faster and less expensive compared to mammalian and insect cultures. Moreover, some proteins are toxic to the host when expressed in their native forms, thus expression as
25 insoluble inclusion bodies is the only way to obtain large quantities of recombinant proteins. Importantly, high levels of expression can be achieved for most proteins. 400 to 600 mg of inclusion bodies per liter of bacterial culture can routinely be achieved, with up to 9,700 mg/L having been reported using this method (Jeong KL; Lee SY, 1999. *Appl. Environ. Microbiol.* 65:3027-32). Inclusion bodies can be easily purified to greater
30 than 90% with a simple freeze/thaw and detergent washing procedure.

 Inclusion bodies appear as dense cytoplasmic granules when the cells are observed under a light microscope. Typically, the cells will be lysed by

mechanical disruption of the cells, followed by centrifugation for 30 min at 4700 g. Inclusion bodies will sediment at low g forces and can be separated from many other intracellular proteins. Further purification can be done by washing the pellet with the buffer used during the cell disruption, or by centrifuging the resuspended pellet in 40-50% glycerol.

Many extracellular proteins of eukaryotes contain disulfide bonds. Proteins having multiple disulfide bonds may form non-native disulfide bonds during folding from the reduced species. Further folding is then blocked unless the incorrect disulfide bond is cleaved by reduction with an external thiol or by attack from a protein thiol. Eukaryotic organisms that secrete disulfide containing proteins also machinery for ensuring proper disulfide bond formation. A distinct disadvantage of expression of recombinant proteins in prokaryotes as inclusion bodies is that the proteins are not obtained in their native state, and typically are not functionally active. A variety of methods have been used to re-solubilize the proteins and refold them to reform active protein. Dissolution of the pelleted recombinant protein usually requires the use of denaturants such as 7 M guanidine hydrochloride or 8 M urea. The amount of aggregation may continue to increase with time if the protein is allowed to remain in the denaturant (Kelley and Winkler, "Folding of Eukaryotic Proteins Produced in *Escherichia coli*" Genetic Engineering 12, 1-19 at p. 6 (1990)). Removal of the denaturant from the solubilized inclusion bodies by dialysis or desalting columns will cause the protein to precipitate under conditions where the native protein needs to be refolded. A misfolded protein solution can also have a very low specific activity in biological assays.

Although there are many reports of expression and refolding of various proteins in *E. coli* as inclusion bodies, one of the misconceptions in protein refolding is that a unique refolding method has to be developed for each individual protein (see Kelley and Winkler at p. 6). Another misconception is that most of the mammalian proteins cannot be refolded from inclusion bodies. (for review, see: Rudolph R., Lilie H., 1996. *FASEB J* 10:49-56; Lilie H, Schwarz E, Rudolph R. 1998. *Curr Opin Biotechnol*



9:497-501). Because published works are mostly "success" stories in refolding inclusion bodies from *E. coli*, it is impossible to get a general idea about what percentage of mammalian proteins can be purified using this procedure.

- 5 There are probably more refolding methods than refolded proteins reported in the literature (for review, see Rudolph R., Lilie, H. 1996, *FASEB J* 10:49-56; Lilie, H., Schwarz, E., Rudolph, R. 1998, *Curr Opin. Biotechnol.* 9:497-501; Huxtable, *et al.*, *Protein Expression and Purification* 12:305-314 (1998); Tichy, *et al.*, *Protein Expression and Purification* 4:59-63 (1993)).
- 10 Different chaperones, detergents, and chaotrophs have been used to help refolding. In addition, pH, ionic strength, temperature, buffer formulation, and reducing/oxidation reagents can all effect refolding. It would be prohibitive to test all these conditions for refolding large amounts of proteins, as required for studies in proteomics or structural genomics.
- 15 A single simplified procedure to refold most of the proteins that are expressed in recombinant systems, especially those which form inclusion bodies in systems such as *E. coli*, is therefore needed.
- It is therefore an object of the present invention to provide a "universal" method for refolding of proteins, especially recombinant proteins, especially
- 20 recombinant proteins present in inclusion bodies in bacterial hosts.

Summary of the Invention

- A universal folding method that has been demonstrated to be effective in refolding a variety of very different proteins expressed in bacteria as
- 25 inclusion bodies has been developed. According to a first broad form of the invention there is provided a method for refolding of a recombinant protein comprising maintaining the pH of 9.0 or greater, in the presence of one or more chaotrophic and reducing agents, and decreasing the pH of the solution gradually over a period of at least about 24 hours to a pH of about 8.0 to
- 30 induce renaturation of at least a protein so that it qualitatively exhibits a biological activity and structure characteristic of the protein.

According to a second broad form of the invention there is provided a method for refolding of a recombinant protein comprising maintaining the protein at a pH of 9.0 or greater, in the presence of one or more chaotropic and reducing agents, and decreasing the pH of the solution gradually over a period of at least about 24 hrs to a physiological pH to induce renaturation of at least a portion of the protein so that it qualitatively exhibits a biological activity and structure characteristic of the protein. Representative proteins that can be dissolved and refolded in biologically active form, with the native structure, are shown in Table I. The method has two key steps to unfold and then refold the proteins expressed in the inclusion bodies. The first step is to raise the pH of the protein solution in the presence of denaturing agents to pH greater than 9, preferably 10. The protein solution may be maintained at the elevated pH for a period of up to about 24 hours, or the pH immediately decreased slowly, in increments of about 0.2 pH units/24 hours, until the solution reaches a pH of about 8.0, or both steps used. In the preferred



embodiment, purified inclusion bodies are dissolved in 8 M urea, 0.1 M Tris, 1 mM glycine, 1 mM EDTA, 10 mM beta-mercaptoethanol, 10 mM dithiothreitol (DTT), 1 mM reduced glutathione (GSH), 0.1 mM oxidized glutathione (GSSG), pH 10. the absorbance at 280 nm (OD280) of the protein solution is 5.0. This solution is rapidly diluted into 20 volumes of 20 mM Tris base. The resulting solutin is adjusted to pH 9.0 with 1 M HCl and is kept at 4°C for another 24 hrs. The pH is adjusted to pH 8.8 and the solution is kept at 4°C for another 24 hrs. This process is repeated until the pH is adjusted to 8.0. After 24 hr at pH 8.0, the refolded proteins can be concentrated by ultrafiltration and applied to a gel filtration column for purification.

Preferably the pH of the solution is reduced more quickly at the higher pH and more gradually near the physiological pH range of the protein.

Detailed Description of the Invention

Expression of Recombinant Proteins

Recombinant proteins are typically expressed in a suitable host, for example, a prokaryotic expression system such as *E.coli* or other type of bacteria, using a standard expression vector like a plasmid, bacteriophage or even naked DNA, and the protein expressed from the plasmid or DNA integrated the host chromosome. Suitable bacterial strains are commercially available or can be obtained from the American Type Culture Collection, Rockville, MD. (the "ATCC").

Suitable vectors can be obtained from any number of sources, including the ATCC. These need a promoter to insure that the DNA is expressed in the host, and may include other regulatory sequences. The vector may also include means for detection, such as an antibiotic resistance marker, green fluorescent protein tag, or antigen tag to facilitate in purification of the recombinant protein.

Once the DNA encoding the protein to be purified is introduced into the host, the host is cultured under appropriate conditions until sufficient amounts of recombinant protein are obtained.

Refolding/purification methods

Once the protein has been expressed in the maximum amount, it must be separated and purified from the bacterial host. The protein is isolated generally by lysing the cells, for example, by suspending in detergent, adding
5 lysozyme, and then freezing (for example, by suspending cells in 20 ml of TN/1% Triton™ X-100, adding 10 mg lysozyme and freezing at -20°C overnight), thawing and adding DNAase to degrade all of the bacterial DNA, then washing the resulting precipitate in a buffered solution. The precipitate is then dissolved in an appropriate solution as discussed below, for refolding.

10 The isolated protein is then refolded. There are several critical aspects of an universal refolding method.

(i) **High pH refolding.** Most published procedures refold proteins using reducing chaotrophs (such as 8 M urea) at a physiological pH, usually pH 7.4 to 8.0. This usually produces large quantities of precipitation
15 or aggregation, making refolding either impossible or with a very low yield. It has been found that some proteins cannot be refolded at physiological pH, but can be refolded when initial refolding pH is high (at least pH 9.0, although higher pH, such as pH 10, may be desirable). This strategy was initially inspired by the fact that pepsinogen can be reversibly
20 denatured/renatured between pH 8.0 and 9.0. It is postulated that at high pH (such as pH 9.0), proteins can obtain some secondary structures, allowing it to be refolded more efficiently when acidity of the refolding solution is lowered to the biological pH. Later it was found that even for the proteins that could be refolded at a physiological pH, high pH refolding resulted in a
25 better yield. In addition, high pH refolding is an excellent method for preventing initial large scale precipitation.

(ii) **Non-denaturing chaotroph concentration.** It has been shown in several labs that non-denaturing concentrations of chaotrophic reagents, such as 0.5 to 1.0 M of urea, guanidine hydrochloride, and L-
30 arginine, can be used to assist refolding and stabilize refolded proteins (Rudolph et al. 1996, *FASEB J* 10:49-56). The preferred concentration of chaotrophic reagents is 0.4 M, although the concentration may range from 0

to 4 M. A moderate concentration of urea in the refolding/purification procedures has not been found to have a denaturing effect on proteins.

(iii) **Reducing/oxidation reagents.** Inclusion bodies for mammalian proteins containing disulfide bonds need to be dissolved in the presence of reducing reagents. Representative reducing agents include beta-mercaptoethanol, in a range of from 0.1 mM to 100 mM, preferably 10 mM; DTT, in a range of from 0.1 mM to 10 mM, preferably 10 mM; reduced glutathion (GSH), in a range from 0.1 mM to 10 mM, preferably 1 mM; and oxidized glutathion (GSSG), in a range from 0.1 mM to 10 mM, preferably 1 mM. Beta-mercaptoethanol is a preferred reducing reagent. In addition, dithiothreitol and/or reduced/oxidized glutathion (GSH, GSSG) can also be included to facilitate "oxido-shuffling" of wrongly folded, intermediate disulfide bonds.

(iv) **pH control** It is important that the protein solution remain at an elevated condition long enough to refold the protein. This is preferably achieved by decreasing the pH slowly, in 0.2 pH unit increments per 24 hours. In this method, the protein solution is adjusted to a high pH, preferably at least 9.0 or higher, to 10 or less preferably 11. The protein is preferably maintained at each pH for at least 24 hours, although comparable effects can be achieved with shorter periods of time, for example, for a period of three, six, nine, twelve, eighteen or twenty hours, most preferably at least twelve hours. The pH can be adjusted by addition of an acid or by dialysis or dilution into a lower pH. Addition of the acid is preferred.

The four conditions discussed above are considered the most essential aspects of the basic protocol for the "universal" refolding procedure.

Table 1. Expression, refolding, and purification of different proteins from *E. coli*

Name	From	Organism	Refold	Purification	Ref.
Pepsinogen	Full-length	Porcine	Yes	Yes	Lin, et al, 1989
Pepsinogen	N and C	Porcine	Yes	Yes	Lin, et al, 1992
	Domain				Lin, et al, 1993
Rhizopus-	full-length	Fungus	Yes	Yes	Chen, et al, 1991
Pepsinogen					Lin, et al, 1992
Thermopsin	full-length	Archae	No	No	None
Thermopsin	fusion	Archae	Yes	Partial	Lin, Liu, Tang, 92
Cathepsin D	full-length	human	No	No	None
					low yield
Pregnancy	full-length	Bovine	No	No	None
Specific Ant		Ovine			
HIV protease	full-length	HIV	Yes	Yes	Lin, et al, 1995
					Ermolief, et al, 97
SAP	full-length	Yeast	Yes	Yes	Lin, et al, 1993
					Koelsch, et al, 98
Streptokinase	full-length	bacteria	Yes	Yes	Wang, et al, 1998
Plasminogen	cat-domain	human	Yes	Yes	Wang, et al, 2000
Cadosin A	full-length	plant	Yes	Yes	Faro, et al, 1999

Nepsin 1	full-length	human	No	No	Koelsch, et al, 00
Memapsin 2	full-length	human	Yes	Yes	Lin, et al, 2000
Memapsin 1	full-length	human	Yes	Yes	
PreS	partial	HBV	Yes	Yes	
unc-76	full-length	C. elegans	Yes	Yes	
odc-1	full-length	C. elegans	Yes	Yes	
ceb-10	full-length	C. elegans	Yes	Yes	
ppp-1	full-length	C. elegans	No	No	

References for Table 1:

- Lin, X., Wong, R.N.S., and Tang, J. (1989) "Synthesis, purification, and active site mutagenesis of recombinant porcine pepsinogen". *J. Biol. Chem.* 264:4482-4489.
- Lin, X.L., Lin, Y.-Z., Koelsch, G., Gustichina, A., Wlodawer, A., and Tang, J. (1992) "Enzymic activities of two-chain pepsinogen, two-chain pepsin, and the amino-terminal lobe of pepsinogen". *J. Biol. Chem.* 267:17257-17263.
- Lin, X., Loy, J.A., Sussman, F., and Tang, J. (1993) "Conformational instability of the N- and C-terminal lobes of porcine pepsin in neutral and alkaline solutions". *Procl. Sci.* 2:1383-1390.
- Chen, Z., Koelsch, G., Han, H.-P., Wang, X.-J., Lin, X., Hartsuck, J.A., and Tang, J. (1991) "Recombinant rhizopuspepsinogen", *J. Biol. Chem.* 266:11718-11725.
- Lin, Y.-Z., Fusek, M., Lin, X.L., Hartsuck, J.A., Kezdy, F.J., and Tang, J. (1992) "pH Dependence of kinetic parameters of pepsin, rhizopuspepsin, and their active-site hydrogen bond mutants". *J. Biol. Chem.* 267:18413-18418.
- Lin, X.L., Liu, M.T., and Tang, J. (1992) "Heterologous expression of thermopsis, a heat stable acid proteinase". *Enzyme Microb. Technol.* 14:696-701.

- Lin, Y.-Z., Lin, X., Hong, L., Foundling, S., Heinrichson, R.L., Thaisrivongs, S., Leelanit, W., Raterman, D., Shah, M., Dunn, B.M., and Tang, J. (1995) "Effect of point mutations on the kinetics and the inhibition of human immunodeficiency virus type 1 protease: relationship to drug resistance". *Biochemistry* 34:1143-1152.
- Ermoloff, J., Lin, X., and Tang, J. (1997) Kinetic properties of sequinavir-resistant mutants of human immunodeficiency virus type 1 protease and their implications in drug resistance *in vivo*. *Biochemistry* 36:12364-12370.
- Lin, X., Tang, J., Koelsch, G., Monod, M., and Foundling, S. (1993) "Recombinant cardiotropin, an extracellular aspartic protease from yeast *Candida tropicalis*". *J. Biol. Chem.* 268 20143-20147.
- Koelsch, G., Tang, J., Monod, M., Foundling, S.L., Lin, X. (1998) "Primary substrate specificities of secreted aspartic proteases of *Candida albicans*". *Adv. Exp. Med. Biol.* 436:335-338.
- Wang, X., Lin, X., Lowy, J.A., Tang, J., Zhang, X.C. (1998) "Crystal structure of the catalytic domain of human plasmin complexed with streptokinase". *Science*. 281:1662-1665.
- Wang, X., Terzyan, S., Tang, J., Loy, J., Lin, X., and Zhang, X. (2000) "Human plasminogen catalytic domain undergoes a novel conformational change upon activation". *J. Mol. Biol.* (in press).
- Faro, C., Ramalho-Santos, M., Vieira, M., Mendes, A., Simoes, I., Andrade, R., Verissimo, P., Lin, X., Tang, J., Pires, E. (1999) "Cloning and Characterization of cDNA Encoding Cardosin A, an RGD-containing Plant Aspartic Proteinase," *J. Biol. Chem.* 274(40):28724-28729.
- Lin, X., Koelsch, G., Wu, S., Downs, D., Dashti, A., and Tang, J. (2000) "Human aspartic protease memapsin 2 cleaves the -secretase site of -amyloid precursor protein. *Proc. Natl. Acc. Sci.* 97(4):1456-1460.
- Ghosh, A.K., Shin, D., Downs, D., Koelsch, G., Lin, X., Ermoloff, J., Tang, J. (2000) "Design of potent inhibitors form human brain memapsin 2 (-secretase)" *J. Amer. Chem. Soc.*, 122:3522-3523.

The present invention will be further understood by reference to the following non-limiting examples.

Example 1: Preferred Method for Refolding Recombinant Proteins

A. Reagents:

- 5 ZB media: 10 g N-Z-Amine A, 5 g NaCl/L
- LB media: 10 g Tryptone, 5 g Yeast extract, 10 g NaCl/L, pH 7.5
- 8 M urea: 8 M urea, 0.1 M TRIS™, 1 mM glycine, 1 mM EDTA, pH 10
- TN buffer: 0.05 mM TRIS™, 0.15 M NaCl, pH 7.5

B. Expression of the Recombinant Protein:

- 10 1. Expression plasmids should be transfected into an appropriate host, such as the BL21(DE3) strain of *E. coli* and plated on ZB/Ampicillin plates, which selects for the desired recombinant organisms. A single colony from each construct is inoculated into 100 ml of ZB/ampicillin media and grown 16 h at 37°C.
- 15 2. Inoculate 20 ml of the overnight culture into 1 L of LB/ampicillin, and shake at 37°C till OD₆₀₀ reaches 0.4-0.6. Add IPTG to 0.5 mM, then shake for 3 h.
- 3. Centrifuge and resuspend cells in 20 ml of TN/1% Triton™ X-100. Add 10 mg lysozyme and freeze at -20°C overnight.
- 20 4. Thaw the frozen cells, add 20 µl 1 M MgSO₄ 100 µg DNAase, and stir until the bacterial DNA is completely dissolved.
- 5. Add 250 ml of TN/1% Triton and stir for 2-4h. Centrifuge and repeat the Triton wash one more time.
- 6. Dissolve the pellet in 10 ml of 8 M urea solution, add beta-
- 25 Mercaptoethanol to 100 mM. This solution can be ultracentrifuged, and is then ready for refolding.

C. Refolding and purification:

- The OD₂₈₀ of the solution containing the inclusion bodies is adjusted to 5.0 with the 8 M urea solution. The final solution contains the following
- 30 reducing reagents:
- 10 mM beta-Mercaptoethanol

10 mM DTT (Dithiothreitol)
 1 mM reduced glutathion (GSH)
 0.1 mM oxidized glutathion (GSSG)

The final pH of the solution is 10.0.

5 1. The above solution is rapidly diluted into 20 volumes of 20 mM TRISTM base, the pH is adjusted to 9.0, and then slowly adjusted to 8.0 with 1 M HCl, by adjusting pH to 8.8 for twenty four hours, then 8.6 for twenty four hours, etc., until the pH is 8.0. Alternatively, the proteins can be refolded using dialysis. The OD₂₈₀ of the 8 M urea solution is adjusted to 0.5, and dialyzed
 10 against 20 volumes of TRISTM base. The pH of the solution is again slowly adjusted to 8.0.

2. The refolded material is then concentrated by ultrafiltration, and separated by gel filtration, for example, on a SEPHACRYLTM S-300 column equilibrated with 20 mM TRISTM, HCl, 0.4 M urea, pH 8.0.

15 3. The S-300 fractions can be checked by running a non-reduced SDS-PAGE. The wrongly refolded protein runs at a very high molecular weight, while folded proteins run at a normal molecular weight.

4. The refolded peak from the S-300 column can be further purified with a FPLC Resource-QTM or Resource-STM column, which is equilibrated with
 20 20 mM TRISTM-HCl (HEPES buffer for Resource-STM), 0.4 M urea, pH 8.0. The enzyme is eluted from the column with a linear gradient of NaCl. Table I lists all of the proteins which have been refolded directly or as inclusion bodies.

Example 2. Expression and Refolding of memapsin 2.

25 Pro-memapsin 2 was PCR amplified and cloned into the *Bam*HI site of a pET11a vector. The resulting vector expresses pro-memapsin 2 having a sequence from Ala-8p to Ala 326. Two expression vectors, pET11-memapsin 2-T1 (hereafter T1) and pET11-memapsin 2-T2 (hereafter T2), were constructed. In both vectors, the N-terminal 15 residues of the expressed recombinant
 30 proteins are derived from the expression vector. Pro-memapsin 2 residues start at residue Ala-16. The two recombinant pro-memapsin 2s have different C-

terminal lengths. Clone T1 ends at Thr- 454 and clone T2 ends at Ala-419. The T1 construct contains a C-terminal extension from the T2 construct but does not express any of the predicted transmembrane domain.

The T1 and T2 expression vectors were separately transfected into *E. coli* strain BL21(DE3). The procedures for the culture of transfected bacteria, induction for synthesis of recombinant proteins and the recovery and washing of inclusion bodies containing recombinant proteins are essentially as described by Lin et al., 1994 Methods in Enzymology 214, 195-224. Basically, the inclusion bodies are washed with 1% (v/v) Triton X-100 and 0.15 M NaCl in 0.1 M TrisTM HCl, pH 7.4, the insoluble protein is dissolved in a solution containing 8 M urea, 0.05 M cyclohexylaminopropanesulfonic acid, 10 mM 2-mercaptoethanol, 10 mM DTT (Dithiothreitol), 1 mM reduced glutathion (GSH), 0.1 mM oxidized glutathion (GSSG), 1 mM glycine, and 1 mM ethylenediaminetetraacetic acid (EDTA), pH 10.5, to a protein concentration of about 5 mg/ml. This solution is added dropwise to 20 vol of rapidly stirred 20 mM Tris base. The pH of the diluted solution is readjusted to 9.0 with 1 M HCl and kept at 4 °C for 24 hr. The pH is then adjusted to 8.8 with 1 M HCl, and kept at 4 °C for 24 hr again. The process is repeated until the pH is adjusted to 8.0. Three different refolding methods have produced satisfactory results.

(i) *The rapid dilution method.*

Pro-memapsin 2 in 8 M urea/10 mM DTT (Dithiothreitol), 1 mM reduced glutathion (GSH), 0.1 mM oxidized glutathion (GSSG), with OD_{280nm} = 5 was rapidly diluted into 20 volumes of 20 mM-Tris base. The solution was slowly adjusted into pH 8 with 1 M HCl. The refolding solution was then kept at 4° C for 24 to 48 hours before proceeding with purification.

(ii) *The reverse dialysis method*

An equal volume of 20 mM TRISTM, 0.5 mM oxidized/1.25 mM reduced glutathione, pH 9.0 is added to rapidly stirred pro-memapsin 2 in 8 M urea/10 mM beta-mercaptoethanol with OD_{280 nm} = 5. The process is repeated three more times with 1 hour intervals. The resulting solution is then dialyzed against

sufficient volume of 20 mM TRISTM base so that the final urea concentration is 0.4 M. The pH of the solution is then slowly adjusted to 8.0 with 1 M HCl.

iii. The preferred method for refolding.

- Inclusion bodies are dissolved in 8 M urea, 0.1 M TRISTM, 1 mM Glycine, 1 mM EDTA, 100 mM beta-mercaptoethanol, pH 10.0. The OD₂₈₀ of the inclusion bodies are adjusted to 5.0 with the 8 M urea solution without beta-mercaptoethanol. The final solution contains the following reducing reagents: 10 mM beta-mercaptoethanol, 10 mM DTT (Dithiothreitol), 1 mM reduced glutathione, and 0.1 M oxidized glutathione. The final pH of the solution is 10.0.
- 5 The above solution is rapidly diluted into 20 volumes of 20 mM TRISTM base, the pH is adjusted to 9.0, and the resulting solution is kept at 4 °C for 16 hr. The solution is equilibrated to room temperature in 6 hr, and the pH is adjusted to 8.5. The solution is returned to 4 °C again for 18 hr.


- 15 The solution is again equilibrated to room temperature in 6 hr, and the pH is adjusted to 8.0. The solution is returned to 4 °C again for 4 to 7 days.

Purification of recombinant pro-memapsin 2-T1

- The refolded material is concentrated by ultrafiltration, and separated on a SEPHACRYLTM S-300 column equilibrated with 20 mM TRISTMHCl, 0.4 M urea, pH 8.0. The refolded peak (second peak) from the S-300 column can be
- 20 further purified with a FPLC RESOURCE-QTM column, which is equilibrated with 20 mM TRISTM-HCl, 0.4 M urea, pH 8.0. The enzyme is eluted from the column with a linear gradient of NaCl. The refolded peak from S-300 can also be activated before further purification. For activation, the fractions are mixed with equal volume 0.2 M Sodium Acetate, 70% glycerol, pH 4.0. The mixture is
- 25 incubated at 22 °C for 18 hr, and then dialyzed twice against 20 volumes of 20 mM Bis-TRISTM, 0.4 M urea, pH 6.0. The dialyzed materials are then further purified on a FPLC RESOURCE-QTM column equilibrated with 20 Bis-TRISTM, 0.4 M urea, pH 6.0. The enzyme is eluted with a linear gradient of NaCl.

- 30 Modifications and variation of these methods are intended to come within the scope of the appended claims.

In the specification the term 'comprising' shall be understood to have a broad meaning similar to the term 'including' and will be understood to imply the inclusion of a stated integer or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps. This definition also applies to variations on the term 'comprising' such as 'comprise' and 'comprises'.



THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method for refolding of a recombinant protein comprising maintaining the protein at pH of 9.0 or greater, in the presence of one or more chaotropic and
 5 reducing agents, and
 decreasing the pH of the solution gradually over a period of at least about 24 hrs to a pH of about 8.0 to induce renaturation of at least a portion of the protein so that it qualitatively exhibits a biological activity and structure characteristic of the protein.
 10
2. A method for refolding of a recombinant protein comprising maintaining the protein at a pH of 9.0 or greater, in the presence of one or more chaotropic and reducing agents, and
 decreasing the pH of the solution gradually over a period of at least about 24
 15 hrs to a physiological pH to induce renaturation of at least a portion of the protein so that it qualitatively exhibits a biological activity and structure characteristic of the protein.
3. The method of claim 2 wherein the pH is decreased to pH 8.0.
4. The method of claim 2 wherein the pH is decreased to pH 7.4.
5. The method of any one of the preceding claims wherein the pH is decreased in increments equivalent to 0.2 pH units per 24 hours.
6. The method of any one of the preceding claims wherein the protein is maintained at a pH of greater than 9.0 for a period of at least 24 hours.
7. The method of any one of the preceding claims wherein the pH is
 30 decreased by addition of acid.

8. The method of any one of claims 1 to 6 wherein the pH is decreased by dilution or dialysis into a solution of a lower pH.

9. The method of any one of the preceding claims wherein one or more
5 the chaotropic and reducing reagents are selected from the group consisting of urea, guanidine hydrochloride, L-arginine, beta-mercaptoethanol, DTT, reduced glutathione, and oxidized glutathione.

10. The method of any one of the preceding claims wherein the chaotropic and reducing reagents are selected from the group consisting of between 0.4 and 1.0
10 M urea, 0.1 mM to 100 mM beta-mercaptoethanol, 0.1 mM to 100 mM DTT, 0.1 mM to 10 mM reduced glutathione, and 0.1 mM to 10 mM oxidized glutathione.

11. The method of any one of the preceding claims wherein the protein is
15 first extracted from bacterial inclusion bodies.

12. The method of claim 11 wherein the bacteria is *E. coli*.

13. The method of claim 11 or claim 12 wherein the inclusion bodies are
20 dissolved at a final pH of between 9.0 and 10.0.

14. The method of claim 11 or claim 12 wherein the inclusion bodies are dissolved at a final pH of greater than 9.0.

15. The method of claim 11 or claim 12 wherein the inclusion bodies are
25 dissolved at a final pH of greater than 10.0.

16. The method of any one of the preceding claims, wherein the protein is maintained at a pH above about 10.0.

30

17. The method of any one of the preceding claims, wherein the protein is maintained at a pH above about 11.0.

18. The method of any one of the preceding claims, wherein the protein is maintained at a pH above about 12.0.

5 19. The method of any one of the preceding claims wherein the pH is decreased over a period of at least 36 hours.

20. The method of any one of the preceding claims, wherein the pH of the solution is reduced more quickly at the higher pH and more gradually nearer the
10 physiological pH range of the protein.

21. The method of any one of the preceding claims comprising the additional step of separating protein species which exhibit biological activity from inactive, or wrongly folded, protein species.

15

22. A method for refolding, recombinant proteins substantially as hereinbefore described with reference to any one of examples 1 or 2.



20 23. The method of any one of claims 11 – 15 wherein the inclusion bodies are dissolved in 8 M urea, 0.1 M Tris, 1 mM glycine, 1 mM EDTA, 10 mM beta-mercaptoethanol, 10 mM dithiothreitol, 1 mM reduced glutathione, 0.1 mM oxidized glutathione, pH 10.



25 24. The method of claim 23 further comprising adjusting absorbance at 280 nm of the solution to 5.0; and rapidly diluting the solution into 20 volumes of 20 mM Tris base before the step of decreasing the pH of the solution.

30

DATED THIS ELEVENTH DAY OF JULY 2005

OKLAHOMA MEDICAL RESEARCH FOUNDATION

BY

PIZZEYS PATENT AND TRADE MARK ATTORNEYS