

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



WIPO | PCT



(10) International Publication Number
WO 2012/166534 A1

(43) International Publication Date
6 December 2012 (06.12.2012)

- (51) **International Patent Classification:**
A01N 33/24 (2006.01) *A61K 31/13* (2006.01)
- (21) **International Application Number:**
PCT/US2012/039401
- (22) **International Filing Date:**
24 May 2012 (24.05.2012)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
61/490,929 27 May 2011 (27.05.2011) US
- (71) **Applicant (for all designated States except US):** GENO LLC [US/US]; 2941 Oxbow Circle, Cocoa, FL 32926 (US).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** WATTS, Tammy [US/US]; 305 Fox Trail Street, Palm Bay, FL 32909 (US). ROSCIGNO, Robert, F. [US/US]; 7415 Aquarina Beach Drive, Unit 403, Melbourne Beach, FL 32951 (US). FINE, David, H. [US/US]; P.O. Box 321610, Cocoa Beach, FL 32932 (US). RUBIN, Lewis, J. [US/US]; 5550 Caminito Genio, La Jolla, CA 92037 (US).
- (74) **Agents:** FOX, Harold H. et al.; Steptoe & Johnson LLP, 1330 Connecticut Avenue, NW, Washington, DC 20036 (US).

- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** METHOD OF DETERMINING VASOREACTIVITY USING INHALED NITRIC OXIDE

(57) **Abstract:** A method for treating pulmonary arterial hypertension in a patient can include determining whether a patient with pulmonary arterial hypertension has an enlarged right ventricle or right ventricular dysfunction, where the patient has previously been administered a prostanoid and weaning the patient off of the prostanoid if the patient does not have an enlarged right ventricle or right ventricular dysfunction.



WO 2012/166534 A1

**METHOD OF DETERMINING VASOREACTIVITY USING INHALED NITRIC
OXIDE**

CLAIM OF PRIORITY

[0001] This application claims the benefit of prior U.S. Provisional Application No.
5 61/490,929 filed on May 27, 2011, which is incorporated by reference in its entirety.

TECHNICAL FIELD

[0002] This description relates to methods of determining vasoreactivity using inhaled nitric oxide.

BACKGROUND

- 10 [0003] Nitric oxide (NO), also known as nitrosyl radical, is a free radical that is an important signalling molecule. For example, NO can cause smooth muscles in blood vessels to relax, thereby resulting in vasodilation and increased blood flow through the blood vessel. These effects can be limited to small biological regions since NO can be highly reactive with a lifetime of a few seconds and can be quickly metabolized in the body.
- 15 [0004] Generally, nitric oxide (NO) can be inhaled or otherwise delivered to an individual's lungs. Providing a dose of NO can result in localized pulmonary vasodilation. Typically, the NO gas can be supplied in a bottled gaseous form diluted in another gas, for example, nitrogen gas (N₂). Great care should be taken to prevent the presence of even trace amounts of oxygen (O₂) in the tank of NO gas because the NO, in the presence of O₂,
20 can be oxidized to nitrogen dioxide (NO₂). Unlike NO, the part per million levels of NO₂ gas can be highly toxic if inhaled and can form nitric and nitrous acid in the lungs.

SUMMARY

- [0005] In one aspect, a method for treating pulmonary arterial hypertension in a patient can include determining whether a patient with pulmonary arterial hypertension has
25 an enlarged right ventricle or right ventricular dysfunction.

[0006] In some embodiments, the patient has previously been administered a prostanoid. A prostanoid can include a prostaglandin, a prostaglandin derivative or a thromboxane. A prostanoid can include prostacyclin, epoprostenol, trepostinil, iloprost or beraprost. A prostanoid can be administered intravenously, subcutaneously, orally (e.g.

liquid, tablet or capsule) or by inhalation. A patient can be administered a prostanoid as part of a treatment course. A course can include administration of the prostanoid at least once a day, at least once a week or at least once a month. A course can last for at least three months, at least six months, at least one year, at least two years, or more than two years.

5 [0007] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

10 [0008] In some embodiments, determining if the patient has an enlarged right ventricle or right ventricular dysfunction can include administering a vasoactive agent to the patient. A vasoactive agent can cause vasculature to dilate, increasing the internal diameter of the vasculature. In some embodiments, a vasoactive agent can be inhaled nitric oxide.

[0009] In some embodiments, administering inhaled nitric oxide can include
15 communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[0010] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO_2 or N_2O_4 to NO can be used as determined by a person of skill in the art. For example, a reducing agent can include a
20 hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II) , Mo(VI) , NaI , Ti(III) or Cr(III) . A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An
25 antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[0011] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

[0012] In some embodiments, administering nitric oxide can further include
30 communicating the air flow to the outlet through the surface-active material.

[0013] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

[0014] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[0015] In some embodiments, determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[0016] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.

[0017] In some embodiments, determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include calculating a change in at least one indicator of the hemodynamics of the patient between before the inhaled nitric oxide was administered and after the inhaled nitric oxide was administered. For example, an at least one indicator of the hemodynamics of the patient can be monitored before administration of a vasoactive agent and then the at least one indicator of the hemodynamics of the patient can be monitored after administration of a vasoactive agent. The difference between the indicator of the hemodynamics of the patient can be determined or calculated based on at least those two time points. A change in at least one indicator of the hemodynamics of the patient can be a negative value, zero or a positive value.

[0018] In some embodiments, the at least one indicator of hemodynamics can include pulmonary vascular resistance and cardiac output. In some embodiments,

determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include determining the presence of an increase in cardiac output after administration of nitric oxide which correlates with a decrease in pulmonary vascular resistance after administration of nitric oxide. For example, a minimal decrease in

5 pulmonary vascular resistance can correlate with a minimal increase in cardiac output.

Accordingly, a moderate decrease in pulmonary vascular resistance can correlate with a moderate increase in cardiac output, and a marked decrease in pulmonary vascular resistance can correlate with a marked increase in cardiac output. A minimal decrease in pulmonary vascular resistance can be a decrease of at least zero mmHg·min/l and at most

10 0.2 mmHg·min/l, a moderate decrease in pulmonary vascular resistance can be a decrease of at least 0.2 mmHg·min/l and at most 1.0 mmHg·min/l, and a marked decrease can be of at least 1.0 mmHg·min/l. A pulmonary vascular resistance can also increase. A minimal increase in cardiac output can be at least 0.0 l/min to at most 0.1 l/min, a moderate increase in cardiac output can be at least 0.1 l/min to at most 0.3 l/min, a marked increase in cardiac output can be at least 0.3 l/min. Also, cardiac output can decrease.

[0019] In some embodiments, a decrease of at least 0.2 mmHg·min/l in pulmonary vascular resistance can correlate with an increase of at least 0.1 l/min in cardiac output.

[0020] In some embodiments, a statistically significant decrease in pulmonary vascular resistance can correlate with a statistically significant increase in cardiac output.

20 Statistical significance can be independent of the magnitude of change (i.e. minimal, moderate, marked). Statistical significance can be determined using any of a variety of statistical tests known to those of skill in the art, including but not limited to, chi-square test, student's t-test, z- test, Fisher's exact test, runs test, Kolmogorov-Smirnov test, Mann-Whitney U test, Wald-Wolfowitz runs, Kruskal-Wallis test, Jonckheere-Terpstra test,

25 McNemar test, Wilcoxon sign test, Friedman test, Kendall's W test and Cochran Q test.

[0021] In some embodiments, an increase in cardiac output which correlates with a decrease in pulmonary vascular resistance can indicate that an enlarged right ventricle or right ventricular dysfunction may not be present. A decrease in pulmonary vascular resistance can occur as a result of vasodilation. Therefore, as pulmonary vascular resistance decreases, a correlative increase in cardiac output can be expected, particularly when the

30 right ventricle is functioning properly. On the other hand, a decrease in pulmonary vascular resistance without a correlative cardiac output can indicate that the right ventricle is unable

to respond in the change in the hemodynamics and can suggest an enlarged right ventricle or right ventricular dysfunction.

[0022] In some embodiments, the method can include weaning the patient off of the prostanoid if the patient does not have an enlarged right ventricle or right ventricular dysfunction. Weaning can mean the decreasing the amount of prostanoid administered over time to a patient during a given unit of time (e.g. per day, per week, per month), most preferably, decreasing the amount to zero.

[0023] In another aspect, a method for treating pulmonary arterial hypertension in a patient can include administering a vasoactive agent to a patient with pulmonary arterial hypertension. The patient may have previously been administered a prostanoid. A prostanoid can include a prostaglandin, a prostaglandin derivative or a thromboxane. A prostanoid can include prostacyclin, epoprostenol, trepostinil, iloprost or beraprost. A prostanoid can be administered intravenously, subcutaneously, orally (e.g. liquid, tablet or capsule) or by inhalation. A patient can be administered a prostanoid as part of a treatment course. A course can include administration of the prostanoid at least once a day, at least once a week or at least once a month. A course can last for at least three months, at least six months, at least one year, at least two years, or more than two years.

[0024] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

[0025] In some embodiments, a method can include determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent. A vasoactive agent can cause vasculature to dilate, increasing the internal diameter of the vasculature. In some embodiments, a vasoactive agent can be inhaled nitric oxide.

[0026] In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[0027] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO_2 or N_2O_4 to NO can be used as

determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[0028] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

[0029] In some embodiments, administering nitric oxide can further include communicating the air flow to the outlet through the surface-active material.

[0030] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

[0031] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[0032] In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[0033] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.

[0034] In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can further include calculating a change in at least one indicator of the hemodynamics of the patient between before the vasoactive was administered and after the vasoactive was administered. For example, an at least one indicator of the hemodynamics of the patient can be monitored before administration of a vasoactive agent and then the at least one indicator of the hemodynamics of the patient can be monitored after administration of a vasoactive agent. The difference between the indicator of the hemodynamics of the patient can be determined or calculated based on at least those two time points. A change in at least one indicator of the hemodynamics of the patient can be a negative value, zero or a positive value.

[0035] In some embodiments, the at least one indicator of hemodynamics can include mean pulmonary arterial pressure. In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can further include determining the presence of a change in mean pulmonary arterial pressure of greater than 5 mm Hg, greater than 10 mm Hg between, greater than 12 mm Hg or greater than 15 mm Hg before the inhaled nitric oxide was administered and after the inhaled nitric oxide was administered.

[0036] In some embodiments, the method can further included weaning the patient off of the prostanoid if the patient demonstrates an acute vasodilator response to the vasoactive agent. Weaning can mean the decreasing the amount of prostanoid administered over time to a patient during a given unit of time (e.g. per day, per week, per month), most preferably, decreasing the amount to zero.

[0037] In another aspect, a method for monitoring a patient with pulmonary arterial hypertension can include performing a vasoreactivity challenge test.

[0038] In another aspect, a method for monitoring a patient with pulmonary arterial hypertension can include administering inhaled nitric oxide to the patient with pulmonary arterial hypertension.

[0039] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years

old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

[0040] In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle
5 can include an inlet, an outlet, and/or a surface-active material.

[0041] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[0042] In some embodiments, a surface-active material can include a reducing
10 agent. Any appropriate reducing agent that can convert NO₂ or N₂O₄ to NO can be used as determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic
15 acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[0043] In some embodiments an inlet can be configured to receive the flow of
20 nitrogen dioxide.

[0044] In some embodiments, administering nitric oxide can further include communicating the air flow to the outlet through the surface-active material.

[0045] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient
25 temperature.

[0046] In some embodiments, a method can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure
30 or cardiac output. Hemodynamics can be anything relating to the movement, flow or

circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[0047] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering inhaled nitric oxide to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering inhaled nitric oxide to the patient.

[0048] In another aspect, a method for diagnosing right ventricular dysfunction can include administering a vasoactive agent to a patient. A patient can be suspected of having right ventricular dysfunction. A patient can have been diagnosed with pulmonary arterial hypertension.

[0049] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

[0050] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[0051] In some embodiments, a vasoactive agent can be nitric oxide, more specifically, inhaled nitric oxide. In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[0052] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO_2 or N_2O_4 to NO can be used as determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II) , Mo(VI) , NaI , Ti(III) or Cr(III) . A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. An antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An

antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[0053] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

5 [0054] In some embodiments, administering nitric oxide can further include communicating the air flow to the outlet through the surface-active material.

[0055] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

10 [0056] In some embodiments, a method can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or
15 circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[0057] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the
20 patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.

[0058] In some embodiments, a method can include determining whether the patient has right ventricular dysfunction.

[0059] In some embodiments, the at least one indicator of hemodynamics can
25 include pulmonary vascular resistance and cardiac output. In some embodiments, determining whether the patient has right ventricular dysfunction can further include determining the presence of an increase in cardiac output after administration of nitric oxide which correlates with a decrease in pulmonary vascular resistance after administration of nitric oxide. For example, a minimal decrease in pulmonary vascular resistance can
30 correlate with a minimal increase in cardiac output. Accordingly, a moderate decrease in pulmonary vascular resistance can correlate with a moderate increase in cardiac output, and

a marked decrease in pulmonary vascular resistance can correlate with a marked increase in cardiac output. A minimal decrease in pulmonary vascular resistance can be a decrease of at least zero mmHg·min/l and at most 0.2 mmHg·min/l, a moderate decrease in pulmonary vascular resistance can be a decrease of at least 0.2 mmHg·min/l and at most 1.0

5 mmHg·min/l, and a marked decrease can be of at least 1.0 mmHg·min/l. A pulmonary vascular resistance can also increase. A minimal increase in cardiac output can be at least 0.0 l/min to at most 0.1 l/min, a moderate increase in cardiac output can be at least 0.1 l/min to at most 0.3 l/min, a marked increase in cardiac output can be at least 0.3 l/min. Also, cardiac output can decrease.

10 **[0060]** In some embodiments, a decrease of at least 0.2 mmHg·min/l in pulmonary vascular resistance can correlate with an increase of at least 0.1 l/min in cardiac output.

[0061] In some embodiments, a statistically significant decrease in pulmonary vascular resistance can correlate with a statistically significant increase in cardiac output. Statistical significance can be independent of the magnitude of change (i.e. minimal,
15 moderate, marked). Statistical significance can be determined using any of a variety of statistical tests known to those of skill in the art, including but not limited to, chi-square test, student's t-test, z- test, Fisher's exact test, runs test, Kolmogorov-Smirnov test, Mann-Whitney U test, Wald-Wolfowitz runs, Kruskal-Wallis test, Jonckheere-Terpstra test, McNemar test, Wilcoxon sign test, Friedman test, Kendall's W test and Cochran Q test.

20 **[0062]** In some embodiments, an increase in cardiac output which correlates with a decrease in pulmonary vascular resistance can indicate that an enlarged right ventricle or right ventricular dysfunction may not be present. A decrease in pulmonary vascular resistance can occur as a result of vasodilation. Therefore, as pulmonary vascular resistance decreases, a correlative increase in cardiac output can be expected, particularly when the
25 right ventricle is functioning properly. On the other hand, a decrease in pulmonary vascular resistance without a correlative cardiac output can indicate that the right ventricle is unable to respond in the change in the hemodynamics and can suggest an enlarged right ventricle or right ventricular dysfunction.

[0063] Other features will become apparent from the following detailed description,
30 taken in conjunction with the accompanying drawings, which illustrate by way of example, the features of the various embodiments.

BRIEF DESCRIPTION OF THE DRAWINGS

[0064] FIG. 1 is a diagram of a NO delivery system.

DETAILED DESCRIPTION

Introduction

5 [0065] Pulmonary arterial hypertension (PAH), a rare progressive disorder, can carry a poor prognosis. PAH can be associated with significant morbidity and mortality, having a historical survival rate rarely exceeding 5 years. Patients with PAH can be classified based on a system developed by the World Health Organization (WHO), which places patients in one of four classes based on aetiology. (Nauser, et al., "Diagnosis and
10 Treatment of Pulmonary Hypertension," Am Fam Physician, 2001; 63: 1789-98, 1800, which is incorporated by reference in its entirety). Patients can also be classified using a Functional classification system. (Nauser, 2001).

[0066] The use of parenteral prostacyclin analogs, the first medications approved for the treatment of PAH, has resulted in substantial improvement in survival for PAH
15 WHO functional class III and IV patients. However, treatment with prostacyclin analogs can be costly and complex to administer, can require continuous infusion, and can be associated with frequent side effects (e.g., flushing, jaw pain, diarrhea) and a potential for severe or life-threatening sequelae (e.g., catheter-related infections and sepsis). Also, significant, potentially life-threatening, hemodynamic compromise may occur due to abrupt
20 cessation of the infusion. The complications associated with parenteral prostacyclin therapy can include reduced quality of life, increased hospitalizations, morbidity and in certain cases, can be life threatening.

[0067] Traditionally, initiating treatment with parenteral prostacyclin therapies can be a life-long requirement with no opportunity to consider alternative therapies that may be
25 less invasive and less cumbersome. However, in the past decade significant advances in the treatment of PAH, along with a better understanding of the pathogenesis of this debilitating disease, suggest alternative patient management strategies may be available.

[0068] It has been suggested that PAH can become a "fixed" defect in the pulmonary vasculature with decreased reactivity to various vasodilators. The exact
30 mechanism of decreased responsiveness to acute vasodilator testing is unknown. The propensity for select patients with PAH to have normalization of pulmonary hemodynamics

following long-term administration of parenteral prostacyclin therapy is also unknown. However, some patients have demonstrated a normalization of pulmonary hemodynamics following long-term administration of parental prostacyclin therapy.

[0069] For example, one case study involving four WHO class IV patients reported successful transition from chronic parenteral prostacyclin infusion to less complex and less invasive treatment following normalization of pulmonary pressures and restored pulmonary vasoreactivity (improving to WHO class I-II). (Kim et al., 2003).

[0070] Following this single center experience, additional successes have been reported, documenting a total of 63 transitions from prostanoid therapy to alternative therapies. (Sidhu, et al., 2011; Johnson, et al., 2007; Steiner, et al., 2006; Diaz-Guzman, et al., 2008; Suleman, et al., 2004; Camacho, et al., 2006). Table 1 summarizes the available literature from parenteral prostacyclin therapy to oral agents.

[0071] Predictors of successful transition remain difficult to identify because of the variety of detail reported in the literature, the retrospective nature of the experiences, the population baseline characteristics, and the inconsistent methodologies for collecting data across the reported single center experiences, for example. This can make it difficult to draw any conclusions from the reports. Additionally, the transition process can be a complex process that can require careful monitoring and close follow-up.

[0072] However, common criteria has been noted for selecting patients for transition from prostanoid therapy to alternative therapies. The common criteria can include that a patient is considered “clinically stable.” A patient can be considered “clinically stable” if the patient is on stable doses of prostanoid analogs, has no evidence of overt right-sided heart failure, and/or is in WHO classes I-III. While classes I-III have shown evidence of being capable of transition,

Table 1 Transition From Parenteral Prostacyclins to Oral Agents: Summary of Available Literature								
	Kim, et al.	Sulenan and Frost	Steiner, et al.	Johnson, et al	Diaz-Guzman	Sidhu	Camacho	
# of Patients	4	23	22	15	21	7	5	
% Success	4 (100%)	11 (47.8%)	7 (31.8%)	10 (66.6%)	15 (71.4%)	6	5 (100%)	
Design	Retrospective	Prospective	Prospective	Retrospective	Retrospective	Retrospective	Prospective	
Wean Criteria	Normalization of PAP	In frequent increase PGI2 dose, FC II, III no heart failure	Stable FC, <10% decrease in 6MWD and no PGI2 dose changes	Stable FC, no increase in PGI2, oral therapy for 3 months	Stable FC I, II, or III, stable 6MWD, stable prostanoid dose, no right heart failure, oral therapy at least 2 months	Stable FC I,II	Stable, complications associated with PGI Tx	
Mean PGI2 Duration (mos.)								
Success/Failed	68.4	35.2/83.4	41/36	45.1	26/16	14	57	
Mean max PGI2 Dose (ng/kg/min)								
Success/Failed	40.5/NR	25.9/40.1	25.5/37.67	23.2/NR	18/15.5	17/NR	35/NR (+ 2 on Ilprost)	
At wean Success/Failed								
Mean 6MWD	N/A	361.8	317.9/294	420	427.7/469.1	405	465	
mPAP	21	65.8/84.1	38/56	35.8	38.2/42.8		N/A	
CI (L/min/m ² /CO)	CI-2.85	CO-4.4/4.4	N/A	CO4.9	CI-2.1/1.9	CI-	N/A	
PVR (units)	N/A	N/A	N/A	6.2	10.3/17.5		N/A	
Mean FU mo.	11	9.6	17.7	29.9	27.3	41.3	17.6	
AE transition	None	2 late failures, no deaths	4 deaths; 2 Success 2 Failures	2 late failures, no deaths	2 deaths; 1 Success- pneumonia 1 Failure	1 late failure	None	
NR = Not Reported								

Sidhu, et al. has also reported the successful transition of functional class III-IV patients from prostanoid therapy to an alternative therapy. However, it is not clear if the functional class documented by Sidhu was determined at time of diagnosis or following prostanoid therapy.

5 **[0073]** Normalization of pulmonary pressures, restoration of pulmonary vasoreactivity, and/or clinical response to treatment can have significant implications regarding a patient's condition. These factors may prove to be valuable measures of treatment effect. These factors may also prove valuable for evaluating the extent or reversal of beneficial pulmonary vascular remodelling, which can be helpful for evaluating a PAH
10 patient. Consequently, measuring the hemodynamic response to a vasodilator, for example inhaled nitric oxide, may indicate possible pulmonary vascular remodelling. The measurement can then be used to support a change in patient management for carefully selected patients, for instance, in support of transitioning a patient from prostanoid therapy to alternative therapies. Ultimately, this information may result in individual patient
15 management strategies that pose less risk, improve quality of life, and cost less, thus imparting a positive effect on clinical outcome.

[0074] Inhaled nitric oxide (NO) can be administered during right heart catheterization (RHC) to evaluate and characterize pulmonary hemodynamics following long term parenteral prostacyclin therapy and can provide information that may influence
20 the choice of treatment strategies for these patients. These treatment strategies may involve changing vasodilator therapy, eligibility for lung and/or heart transplantation, or changes in prognosis. Inhaled nitric oxide can be a potent, selective pulmonary vasodilator agent and has been well studied up to 80 ppm. It is expected that inhaled nitric oxide as a diagnostic agent during RHC may be useful for facilitating patient management strategies and provide
25 clinical information beneficial to patient care.

[0075] In sum, despite the promising reports of successful transition from continuous infusion therapy, patient selection and factors that may predict success remain unclear. Therefore, there remains a need for methods of evaluating PAH patients, patients who are undergoing parenteral prostacyclin therapy, and patients who are candidates for
30 transitioning from parenteral prostacyclin therapy to an alternative therapy.

Transition of Patients from Prostanoid Therapy

[0076] Pulmonary arterial hypertension (PAH) is a rare disease of the pulmonary vessels that currently has no cure. PAH can eventually affect the right ventricle (RV). Right ventricular performance can be key in explaining symptoms, pathophysiology, and survival in PAH. At least one report has that ~70% of PAH deaths can be attributed to RV failure. (D'Alonzo, et al., 1991). Current therapies can remain palliative, improving symptoms, hemodynamics and longevity, while morbidity and mortality in this population can remain very high.

[0077] Several case studies have reported the successful wean from epoprostenol, a prostacyclin, to oral agents. (Kim, et al., 2003; Johnson, et al., 2007). These studies have also demonstrated normalization of hemodynamics, with two reported cases where patient's exhibited return of vasoreactivity after treatment with epoprostenol. A search of the literature revealed 63 reports of experiences with the transition of patients from prostanoid therapy which shared the following common criteria: 1) normalization/stable hemodynamics, 2) stable dose of prostanoid therapy, 3) stable Functional Class and 4) no clinical signs of heart failure, (JVD, edema, etc.).

[0078] Among these experiences there are instances where pulmonary hemodynamics had improved. This may support a theory that some degree of positive remodeling occurs, possibly more frequently than initially thought. The same trend was noted in the data review of the MUST study, although the positive remodeling phenomenon alone does not seem to be a reliable characteristic to determine the success or failure of the wean/transition process.

[0079] Reports of improved hemodynamics can be found in the literature, which can support the positive effect of epoprostenol, as in the instance of Barst, et al. (1996). This is of particular interest because the findings correlated with the results of a six mile walk test (6MWD), as well. This study, a 12-week open-label, prospective, randomized, multicenter study of 81 patients in which the active group, ($n=41$) demonstrated an improvement in hemodynamics treatment with average decline of 8% in the mean pulmonary arterial pressure and a 21% decline in pulmonary vascular resistance.

[0080] McLaughlin, et al. (1998) reported similar findings with an improvement in symptoms, hemodynamic measures and a reduction in pulmonary vascular resistance with

epoprostenol therapy in a prospective study of 27 consecutive patients. Twenty-six of the 27 patients demonstrated signs of overall improvement, where pulmonary vascular resistance declined by 53%, cardiac output increased by 67%, and mean pulmonary artery pressure declined 22%.

5 [0081] Despite these findings, the improved clinical outcomes for this population and the availability of several pulmonary-specific vasodilators, the median survival for patients with no treatment can be 2.8 years with 1-, 3-, and 5-year survival rates of 68%, 48%, and 34%, respectively. Continuous prostanoid therapy may have improved median survival at 1-, 3-, and 5-years 97-99%, 63-71% and 56%, respectively. (McLaughlin, 2002
10 et al., Barst, 2006, et al.). Right ventricular (RV) dysfunction can be the main cause of death in this population and has been estimated to be responsible for at least 70% of all PAH deaths.

Right Ventricle

[0082] Normally, the right ventricle (RV) can perform against very little resistance,
15 can be a thin walled compliant chamber. The RV can be more adaptive to sudden increases in volume and can be less tolerant to sudden increases in afterload. Capable of maintaining a relatively stable cardiac output, the RV can accommodate significant variations in volume but can be poorly adapted to generating high pressure, particularly in situations where there is a sudden increase in afterload. A sudden increase in afterload can follow massive
20 pulmonary emboli.

[0083] In pulmonary hypertension, RV dysfunction and failure can be the result of the increased longstanding pressure overload of the RV. It may not be the degree of pulmonary hypertension (pulmonary artery pressure) that correlates with symptoms or survival, but RV mass, RV size and/or right atrial pressure that can reflect functional status
25 and can be strong predictors of survival. (Voelkel, et al., 2006). In the PAH population, the 6-minute walk test (6MWT or 6MWD) can be used as a measure of outcome and can correlate better with RV function than pulmonary artery pressure. (ATS Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories, "ATS Statement: Guidelines for the Six-Minute Walk Test," Am J Respir Crit Care Med, Vol. 166, pp. 111-
30 117, 2002, which incorporated by reference in its entirety). However, the 6-minute walk test may not provide information on RV-pulmonary vascular function.

[0084] In PAH, the increase in afterload can manifest as elevated pulmonary vascular resistance. The RV can initially adapt to the increased pressure with myocardial hypertrophy minimizing wall stress through Laplace Law. This initial compensation can lead to eventual decompensation, chamber dilatation, heart failure and/or left ventricular diastolic dysfunction. Abnormal RV function can be reversed with effective medical therapy as can be seen with stabilization and improved exercise capacity. (Hinderliter et al., 1997).

[0085] Prostacyclin therapy can be intended to target the pulmonary vascular bed, by decreasing pulmonary pressures. Therapy can be implemented at very low doses of prostacyclin and continually up-titrated to achieve the established target dose or until dose limiting side effects occur. The idea that continuous up-titration, regardless of clinical status, can be required is under question. Continuous up-titration can lead to excessively high cardiac output and even high output heart failure. A better option may be to conduct frequent reassessment of symptoms, exercise capacity and periodic hemodynamic evaluations.

[0086] Prostacyclin therapies can restore the vasodilator/vasoconstrictor balance and can improve pulmonary hemodynamics, exercise capacity and/or survival. However, this complicated therapy may not be without its own associated risks, and in many instances, the opportunity to transition to alternative less complicated therapies can also have risks. Rapid deterioration with no response to resumption of prostacyclin therapy can occur.

[0087] Criteria for the selection of patients that may be considered suitable for wean or transition can include two recommendations for patient selection: 1) stable/normalization of hemodynamics and/or 2) no clinical evidence of heart failure (Table 1). There may be support that, following prostacyclin therapy, pulmonary pressure can improve and pulmonary vascular resistance (PVR) can decrease.

[0088] The second of the common patient selection criteria, clinical evidence of heart failure, has previously been evaluated during clinical assessment for the presence of jugular vein distension (JVD), peripheral edema and presence of a third heart sound at the time of wean. The echocardiography data available did not seem to correlate reliably with the clinical presentation. Some cases presented did have clinical documentation of signs of heart failure, but the ECHO was not supportive of these findings. No recent catheterization

data found supports the clinical findings, which may give credence to assessing RV systolic function.

[0089] The increase in PVR can lead to increased RV afterload. The fixed nature of this pressure overload may restrict the ability of the RV to augment cardiac output (CO).

5 Cardiac catheterization can be the gold standard for the diagnosis and management of PH and RV dysfunction. However, it can provide the only direct measurement of right sided pressures and direct measurement of RV afterload, disease severity and determining prognostic markers (RAP, CO and mPAP). ('Alonzo et al., 1991).

[0090] The characterization of stable hemodynamics and RV function can provide
10 an indication of the effect of afterload reduction and the RV-pulmonary vascular relationship following long term treatment. This can help to guide and optimize patient management. In the presence of normal RV function and pulmonary hypertension, there can be minimal influence on RV ejection fraction or cardiac output (Bhorade, 1999, et al.). Identification of the RV-pulmonary vascular relationship may also allow for the detection
15 and progression of arteriopathy and the subsequent decline of RV function earlier than current diagnostic methods.

[0091] Vasodilator challenge following therapy, particularly parenteral prostacyclin therapy, can be useful as a diagnostic agent. For example, vasodilator challenge can provide information to guide medical management. Following inhalation of nitric oxide, a
20 reduction in PVR can allow for assessment of the RV hemodynamic response, the assessment of RV function and/or the inotropic effect of prostacyclin therapy. The information obtained from a vasodilator challenge can aid in the determination of the stability of hemodynamics, the RV-pulmonary vascular response to afterload reduction, the preservation of RV function.

25 **[0092]** Repeat vasoreactivity testing (i.e. vasodilator challenge) can be a method suitable for the identification and optimization of PAH therapy (e.g. medication adjustments) and/or characterization of the RV and response to a reduction in afterload. This information can provide significant clinical information regarding the RV-pulmonary vascular relationship. Right ventricular function can be determined and a measure of
30 outcomes information regarding the function of the RV can aid in early detection of heart failure and consequently failure to transition from a parental prostacyclin. Therefore, right

ventricular function can guide treatment decisions and the optimization of patient management.

Study Results

Objectives/Hypothesis

- 5 [0093] An aim of the project was to collect and review the existing data from centers experienced with the practice of transition from parenteral prostacyclins to an alternative therapy. The review was conducted to determine if a population could be characterized based on the hemodynamic response using inhaled nitric oxide during right heart catheterization. A primary objective of the project was to determine the correlation, if
10 any, between the return of pulmonary vascular reactivity following parenteral prostacyclin therapy and the ability of the patient to transition from parenteral prostacyclins to an alternative therapy. Secondary objectives included the exploration of clinical characteristics of a patient population with a documented wean or transition from parenteral prostacyclins, as well as evaluating any change in clinical impression.

15 Method

- [0094] To date seven sites have contributed to the data set collected for a total of 47 baseline visits, and 246 follow-up visits. There are five failed weans and seven late failures. Data was available to review for 40 cases entered. No statistical analysis was performed and reviews completed were descriptive in order to identify any shared characteristics in the
20 population either as a success, failure or late failure.

- [0095] Due to the retrospective nature of the study, the data sets collected for each case at baseline through follow-up are not complete and statistical interpretation is limited. However, the summary review of the data entered for the MUST Study, in addition to the reported cases in the literature identifying experiences with successful transition from
25 parenteral prostanoid, may provide a sufficient amount of information that will guide the pivotal Phase 3 study in this development program.

- [0096] This data reviewed collectively with published literature has identified a link between the return of a degree of vasoreactivity and the suitability for wean or transition from parenteral prostacyclin therapy and the characterization of the population that have a
30 reported wean/transition.

[0097] Three cases, where nitric oxide re-challenge data was available at the time of subject wean, are represented in Table 2.

Table 2 Nitric oxide challenge at time of wean						
PAPm	NO PAPm	CO	NO CO	PVR	NO PVR	IV Prost
45	49	3.7	5.2	10	8.3	Yes
54	48	7.3	7.8	10.2	3.8	No
75	70	4.77	4.77	14.6	14.4	No

[0098] As can be seen, there was only a modest decrease in the PAPm for all three cases. The modest decreases failed to meet the criteria of acute vasodilator response,

5 defined as a fall in PAPm of > 10 mm Hg to < 40 mm Hg. Of interest was the effect of inhaled nitric oxide in the case of the failed transition, a marked increase in the CO and a minimal decrease in PVR suggested the RV compliance had been affected because the increase in CO did not correlate with the decrease in PVR. Review of the five failure to wean cases and four of the seven late failures showed the presence of right ventricular
 10 enlargement in four of the five failure to wean cases (with one blank) and in three of the four reviewed late failure cases. In these cases, the clinical findings did not coorelate with the ECHO data.

[0099] The lack of correlation between the clinical findings and the ECHO data prompted further review of the data. Baseline hemodynamics, wean hemodynamics and
 15 echocardiography data were reviewed for all the failed transitions. Table 3 provides a summary of the available data. A trend was noted among the cases that failed to wean or were identified as late failures having to return to prostacyclin therapy. In each case, where echocardiography data was available and right ventricular enlargement was indicated and/or right ventricular dysfunction was indicated, a failed wean or late failure was common. This
 20 data was not considered conclusive due to the small sample size and missing data.

Table 3 Hemodynamics Failures													
BL PAPm	BL NO PAPm	BL CO	BL NO CO	BL PVR	BL NO PVR	WN PAPm	WN NO PAPm	WN CO	WN NO CO	WN PVR	WN NO PVR	IV Prost	Prost Date
Baseline-Time of Diagnosis						Time of Wean							
60		3.77		13		34		5.74		3.8		Yes	3/14/05
37	32	2.31	2.86	6.5	3.2	26		2.91		5.2		Yes	No wean
53	42	4.2	4.4	9.2	6	49		5.8		6.3		Yes	No wean
58	56	5.8	5.8	8	7.2	47		8.53		3.9		Yes	No wean
56	52	4.3	6.5	6.98	6.3	52		5.49		7.84		Yes	No wean
72		4.24		12.5		46		5.47		7.25		Yes	No wean
34	35	5.2	5.6	5.77	4.49							Yes	5/22/03
70	61	4.28	5.15	13.1							Exp.	Yes	7/23/01
46	48	3.6	5.3	10.3	7.7	45	49	3.7	5.2	10	8.3	Yes	4/4/05
51	61	5.4	7									Yes	8/7/07
50	50	3.5	4.2	10.9	9							Yes	7/5/02
						32		3.76		6.38		Yes	1/1/97

Table 4 Hemodynamics Success												
BL PAPm	BL NO PAPm	BL CO	BL NO CO	BL PVR	BL NO PVR	WN PAPm	WN NO PAPm	WN CO	WN NO CO	WN PVR	WN NO PVR	IV Prost
<i>Baseline-Time of Diagnosis</i>						<i>Time of Wean</i>						
54	40	3	4.02	13.3	5.97	26		7.14		3.2		No
51						18		8.8		1.1		No
57		1.9		27.4		30		3.92		5.6		No
60		2.4		19.2		36		5.37		5.4		No
46		3.1		4.5		45		5.9		6.27		No
						37		5.62		4.2		No
43	40	4.9	4.8	7.55	6.66			6.9		1.87		No
40		2.1		11.18		15		5.86		0.85		No
33	26	4.71	5.4	5.7	3.41	32		4.09		6.6		No
51		6.2		6.4		53		5.7		7.6		No
57		6.4		12		85		4.05		20		No
59	50	5.7	7.2	9.5	6.3	61		4.76		11.5		No
90		6.8		12.1		72		7.63		7.9		No
42				7.71		37	41	6.55		4.58	3.11	No
58		58		2.5		4.1		18				No
58	52	4.5	4.2	16.48	10.7	54	48	7.3	7.8	10.2	3.8	No
75	70	4.77	4.77	14.6	14.4	75	70	4.77	4.77	14.6	14.4	No

Vasoreactive Methods

[00100] Therefore, in one aspect, a method can include vasoreactivity testing to provide a medical professional with information regarding the status of a patient with a pulmonary condition.

5 [00101] In one aspect, a method for treating pulmonary arterial hypertension in a patient can include determining whether a patient with pulmonary arterial hypertension has an enlarged right ventricle or right ventricular dysfunction.

[00102] In some embodiments, the patient has previously been administered a prostanoid. A prostanoid can include a prostaglandin, a prostaglandin derivative or a
10 thromboxane. A prostanoid can include prostacyclin, epoprostenol, trepostinil, iloprost or beraprost. A prostanoid can be administered intravenously, subcutaneously, orally (e.g. liquid, tablet or capsule) or by inhalation. A patient can be administered a prostanoid as part of a treatment course. A course can include administration of the prostanoid at least once a day, at least once a week or at least once a month. A course can last for at least three
15 months, at least six months, at least one year, at least two years, or more than two years.

[00103] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years
20 old). In still other embodiments, a patient can be an adult (at least 18 years old).

[00104] In some embodiments, determining if the patient has an enlarged right ventricle or right ventricular dysfunction can include administering a vasoactive agent to the patient. A vasoactive agent can cause vasculature to dilate, increasing the internal diameter of the vasculature. In some embodiments, a vasoactive agent can be inhaled nitric oxide.

25 [00105] In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[00106] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO₂ or N₂O₄ to NO can be used as
30 determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as

Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An
5 antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[00107] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

[00108] In some embodiments, administering nitric oxide can further include
10 communicating the air flow to the outlet through the surface-active material.

[00109] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

[00110] In some embodiments, a vasoactive agent can be administered repeatedly. In
15 some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[00111] In some embodiments, determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include
20 measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[00112] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.

[00113] In some embodiments, determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include calculating a change in at least

one indicator of the hemodynamics of the patient between before the inhaled nitric oxide was administered and after the inhaled nitric oxide was administered. For example, an at least one indicator of the hemodynamics of the patient can be monitored before administration of a vasoactive agent and then the at least one indicator of the hemodynamics of the patient can be monitored after administration of a vasoactive agent. The difference between the indicator of the hemodynamics of the patient can be determined or calculated based on at least those two time points. A change in at least one indicator of the hemodynamics of the patient can be a negative value, zero or a positive value.

[00114] In some embodiments, the at least one indicator of hemodynamics can include pulmonary vascular resistance and cardiac output. In some embodiments, determining whether the patient has an enlarged right ventricle or right ventricular dysfunction can further include determining the presence of an increase in cardiac output after administration of nitric oxide which correlates with a decrease in pulmonary vascular resistance after administration of nitric oxide. For example, a minimal decrease in pulmonary vascular resistance can correlate with a minimal increase in cardiac output. Accordingly, a moderate decrease in pulmonary vascular resistance can correlate with a moderate increase in cardiac output, and a marked decrease in pulmonary vascular resistance can correlate with a marked increase in cardiac output. A minimal decrease in pulmonary vascular resistance can be a decrease of at least zero mmHg·min/l and at most 0.2 mmHg·min/l, a moderate decrease in pulmonary vascular resistance can be a decrease of at least 0.2 mmHg·min/l and at most 1.0 mmHg·min/l, and a marked decrease can be of at least 1.0 mmHg·min/l. A pulmonary vascular resistance can also increase. A minimal increase in cardiac output can be at least 0.0 l/min to at most 0.1 l/min, a moderate increase in cardiac output can be at least 0.1 l/min to at most 0.3 l/min, a marked increase in cardiac output can be at least 0.3 l/min. Also, cardiac output can decrease.

[00115] In some embodiments, a decrease of at least 0.2 mmHg·min/l in pulmonary vascular resistance can correlate with an increase of at least 0.1 l/min in cardiac output.

[00116] In some embodiments, a statistically significant decrease in pulmonary vascular resistance can correlate with a statistically significant increase in cardiac output. Statistical significance can be independent of the magnitude of change (i.e. minimal, moderate, marked). Statistical significance can be determined using any of a variety of statistical tests known to those of skill in the art, including but not limited to, chi-square

test, student's t-test, z- test, Fisher's exact test, runs test, Kolmogorov-Smirnov test, Mann-Whitney U test, Wald-Wolfowitz runs, Kruskal-Wallis test, Jonckheere-Terpstra test, McNemar test, Wilcoxon sign test, Friedman test, Kendall's W test and Cochran Q test.

[00117] In some embodiments, an increase in cardiac output which correlates with a decrease in pulmonary vascular resistance can indicate that an enlarged right ventricle or right ventricular dysfunction may not be present. A decrease in pulmonary vascular resistance can occur as a result of vasodilation. Therefore, as pulmonary vascular resistance decreases, a correlative increase in cardiac output can be expected, particularly when the right ventricle is functioning properly. On the other hand, a decrease in pulmonary vascular resistance without a correlative cardiac output can indicate that the right ventricle is unable to respond in the change in the hemodynamics and can suggest an enlarged right ventricle or right ventricular dysfunction.

[00118] In some embodiments, the method can include weaning the patient off of the prostanoid if the patient does not have an enlarged right ventricle or right ventricular dysfunction. Weaning can mean the decreasing the amount of prostanoid administered over time to a patient during a given unit of time (e.g. per day, per week, per month), most preferably, decreasing the amount to zero.

[00119] In another aspect, a method for treating pulmonary arterial hypertension in a patient can include administering a vasoactive agent to a patient with pulmonary arterial hypertension. The patient may have previously been administered a prostanoid. A prostanoid can include a prostaglandin, a prostaglandin derivative or a thromboxane. A prostanoid can include prostacyclin, epoprostenol, trepostinil, iloprost or beraprost. A prostanoid can be administered intravenously, subcutaneously, orally (e.g. liquid, tablet or capsule) or by inhalation. A patient can be administered a prostanoid as part of a treatment course. A course can include administration of the prostanoid at least once a day, at least once a week or at least once a month. A course can last for at least three months, at least six months, at least one year, at least two years, or more than two years.

[00120] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

[00121] In some embodiments, a method can include determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent. A vasoactive agent can cause vasculature to dilate, increasing the internal diameter of the vasculature. In some embodiments, a vasoactive agent can be inhaled nitric oxide.

5 [00122] In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[00123] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO_2 or N_2O_4 to NO can be used as
10 determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II) , Mo(VI) , NaI , Ti(III) or Cr(III) . A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used
15 depending on the activities and properties as determined by a person of skill in the art. An antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[00124] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

20 [00125] In some embodiments, administering nitric oxide can further include communicating the air flow to the outlet through the surface-active material.

[00126] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

25 [00127] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[00128] In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can include monitoring an at least one
30 indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least

one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

- 5 **[00129]** Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.
- 10 **[00130]** In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can further include calculating a change in at least one indicator of the hemodynamics of the patient between before the vasoactive was administered and after the vasoactive was administered. For example, an at least one indicator of the hemodynamics of the patient can be monitored before administration of a
- 15 vasoactive agent and then the at least one indicator of the hemodynamics of the patient can be monitored after administration of a vasoactive agent. The difference between the indicator of the hemodynamics of the patient can be determined or calculated based on at least those two time points. A change in at least one indicator of the hemodynamics of the patient can be a negative value, zero or a positive value.
- 20 **[00131]** In some embodiments, the at least one indicator of hemodynamics can include mean pulmonary arterial pressure. In some embodiments, determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent can further include determining the presence of a change in mean pulmonary arterial pressure of greater than 5 mm Hg, greater than 10 mm Hg between, greater than 12 mm Hg or greater than 15
- 25 mm Hg before the inhaled nitric oxide was administered and after the inhaled nitric oxide was administered.
- [00132]** In some embodiments, the method can further included weaning the patient off of the prostanoid if the patient demonstrates an acute vasodilator response to the vasoactive agent. Weaning can mean the decreasing the amount of prostanoid administered
- 30 over time to a patient during a given unit of time (e.g. per day, per week, per month), most preferably, decreasing the amount to zero.

[00133] In another aspect, a method for monitoring a patient with pulmonary arterial hypertension can include performing a vasoreactivity challenge test.

[00134] In another aspect, a method for monitoring a patient with pulmonary arterial hypertension can include administering inhaled nitric oxide to the patient with pulmonary
5 arterial hypertension.

[00135] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years
10 old). In still other embodiments, a patient can be an adult (at least 18 years old).

[00136] In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[00137] In some embodiments, a vasoactive agent can be administered repeatedly. In
15 some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[00138] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO_2 or N_2O_4 to NO can be used as determined by a person of skill in the art. For example, a reducing agent can include a
20 hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II) , Mo(VI) , NaI , Ti(III) or Cr(III) . A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An
25 antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[00139] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

[00140] In some embodiments, administering nitric oxide can further include
30 communicating the air flow to the outlet through the surface-active material.

[00141] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

[00142] In some embodiments, a method can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[00143] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering inhaled nitric oxide to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering inhaled nitric oxide to the patient.

[00144] In another aspect, a method for diagnosing right ventricular dysfunction can include administering a vasoactive agent to a patient. A patient can be suspected of having right ventricular dysfunction. A patient can have been diagnosed with pulmonary arterial hypertension.

[00145] In some embodiments, a patient can be a neonate (i.e. less than one month old), an infant (at least one month to one year old) or a toddler (at least one year to three years old). In other embodiments, a patient can be a child (at least three years to 18 years old). More specifically, a patient can be a young child (at least three years to seven years old). In still other embodiments, a patient can be an adult (at least 18 years old).

[00146] In some embodiments, a vasoactive agent can be administered repeatedly. In some embodiments, a vasoactive agent can be administered at least twice, at least three times, at least four times, at least five times, at least ten times or greater than ten times.

[00147] In some embodiments, a vasoactive agent can be nitric oxide, more specifically, inhaled nitric oxide. In some embodiments, administering inhaled nitric oxide can include communicating an air flow including nitrogen dioxide through a receptacle. The receptacle can include an inlet, an outlet, and/or a surface-active material.

[00148] In some embodiments, a surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO₂ or N₂O₄ to NO can be used as determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. An antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[00149] In some embodiments an inlet can be configured to receive the flow of nitrogen dioxide.

[00150] In some embodiments, administering nitric oxide can further include communicating the air flow to the outlet through the surface-active material.

[00151] In some embodiments, administering nitric oxide can further include converting the gaseous nitrogen dioxide to nitric oxide, most preferably, at ambient temperature.

[00152] In some embodiments, a method can further include monitoring an at least one indicator of hemodynamics of the patient. Monitoring at least one indicator can include measuring a hemodynamic property. For example, a hemodynamic property or an at least one indicator can include pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output. Hemodynamics can be anything relating to the movement, flow or circulation of blood. An indicator of hemodynamics can include, but is not limited to, a pressure, a volume or a rate.

[00153] Monitoring an at least one indicator can include monitoring at any time point during a method. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient before administering a vasoactive agent to the patient. In some embodiments, monitoring can include monitoring the at least one indicator of the hemodynamics of the patient after administering a vasoactive agent to the patient.

[00154] In some embodiments, a method can include determining whether the patient has right ventricular dysfunction.

[00155] In some embodiments, the at least one indicator of hemodynamics can include pulmonary vascular resistance and cardiac output. In some embodiments, determining whether the patient has right ventricular dysfunction can further include determining the presence of an increase in cardiac output after administration of nitric oxide which correlates with a decrease in pulmonary vascular resistance after administration of nitric oxide. For example, a minimal decrease in pulmonary vascular resistance can correlate with a minimal increase in cardiac output. Accordingly, a moderate decrease in pulmonary vascular resistance can correlate with a moderate increase in cardiac output, and a marked decrease in pulmonary vascular resistance can correlate with a marked increase in cardiac output. A minimal decrease in pulmonary vascular resistance can be a decrease of at least zero mmHg·min/l and at most 0.2 mmHg·min/l, a moderate decrease in pulmonary vascular resistance can be a decrease of at least 0.2 mmHg·min/l and at most 1.0 mmHg·min/l, and a marked decrease can be of at least 1.0 mmHg·min/l. A pulmonary vascular resistance can also increase. A minimal increase in cardiac output can be at least 0.0 l/min to at most 0.1 l/min, a moderate increase in cardiac output can be at least 0.1 l/min to at most 0.3 l/min, a marked increase in cardiac output can be at least 0.3 l/min. Also, cardiac output can decrease.

[00156] In some embodiments, a decrease of at least 0.2 mmHg·min/l in pulmonary vascular resistance can correlate with an increase of at least 0.1 l/min in cardiac output.

[00157] In some embodiments, a statistically significant decrease in pulmonary vascular resistance can correlate with a statistically significant increase in cardiac output. Statistical significance can be independent of the magnitude of change (i.e. minimal, moderate, marked). Statistical significance can be determined using any of a variety of statistical tests known to those of skill in the art, including but not limited to, chi-square test, student's t-test, z- test, Fisher's exact test, runs test, Kolmogorov-Smirnov test, Mann-Whitney U test, Wald-Wolfowitz runs, Kruskal-Wallis test, Jonckheere-Terpstra test, McNemar test, Wilcoxon sign test, Friedman test, Kendall's W test and Cochran Q test.

[00158] In some embodiments, an increase in cardiac output which correlates with a decrease in pulmonary vascular resistance can indicate that an enlarged right ventricle or right ventricular dysfunction may not be present. A decrease in pulmonary vascular resistance can occur as a result of vasodilation. Therefore, as pulmonary vascular resistance decreases, a correlative increase in cardiac output can be expected, particularly when the

right ventricle is functioning properly. On the other hand, a decrease in pulmonary vascular resistance without a correlative cardiac output can indicate that the right ventricle is unable to respond in the change in the hemodynamics and can suggest an enlarged right ventricle or right ventricular dysfunction.

5 [00159] Regardless of the method, in some embodiments, a vasoactive agent can include nitric oxide, most preferably, inhaled nitric oxide. When delivering nitric oxide (NO) for therapeutic use to a mammal, it can be important to avoid delivery of nitrogen dioxide (NO₂) to the mammal. Nitrogen dioxide (NO₂) can be formed by the oxidation of nitric oxide (NO) with oxygen (O₂). The rate of formation of nitrogen dioxide (NO₂) can be
10 proportional to the oxygen (O₂) concentration multiplied by the square of the nitric oxide (NO) concentration.

[00160] A NO delivery system can convert nitrogen dioxide to nitric oxide. A NO delivery system can include a surface-active material. A surface-active material can include a reducing agent. Any appropriate reducing agent that can convert NO₂ or N₂O₄ to NO can
15 be used as determined by a person of skill in the art. For example, a reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). A reducing agent can be an antioxidant. An antioxidant can be an aqueous solution of an antioxidant. A antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used
20 depending on the activities and properties as determined by a person of skill in the art. An antioxidant can be used dry or wet. A surface-active material can be coated with a reducing agent or coated with an aqueous solution of a reducing agent.

[00161] The system can employ a surface-active material coated with a reducing agent, for example an antioxidant, as a simple and effective mechanism for making the
25 conversion. More particularly, NO₂ can be converted to NO by passing the dilute gaseous NO₂ over a surface-active material including a reducing agent, e.g. an antioxidant, such that the reducing agent reacts with the NO₂. As an example, when the aqueous antioxidant is ascorbic acid (that is, vitamin C), the reaction can be quantitative at ambient temperatures.

[00162] One example of a surface-active material can be silica gel. Another example
30 of a surface-active material that can be used is cotton. The surface-active material may be or may include a substrate capable of retaining a liquid, for example, water. A surface-active material can include a relatively large surface area. A surface-active material can

also allow a fluid to pass through it. Another type of surface-active material that has a large surface area that is capable of absorbing moisture also may be used.

[00163] Fig. 1 illustrates a cartridge 100 for generating NO by converting NO₂ to NO. The cartridge 100, which may be referred to as a cartridge, a converter, a NO generation cartridge, a GENO cartridge, or a GENO cylinder or Nitrosyl™ Primary Cartridge, can include an inlet 105 and an outlet 110. Screen and/or glass wool 115 can be located at the inlet 105 and/or the outlet 110. The remainder of the cartridge 100 can be filled with a surface-active material 120 that is coated with a reducing agent. The surface-active material can be soaked with a saturated solution including a reducing agent to coat the surface-active material. The saturated solution can be, for example, an antioxidant in water. The screen and/or glass wool 115 can also be soaked with the saturated solution before being inserted into the cartridge 100. The antioxidant can be ascorbic acid.

[00164] In a general process for converting NO₂ to NO, a gas flow (e.g. air flow) having NO₂ can be received through the inlet 105. The gas flow can be fluidly communicated to the outlet 110 through the surface-active material 120 coated with the aqueous reducing agent, e.g. antioxidant. As long as the surface-active material remains moist and the reducing agent may not been used up in the conversion, the general process can be effective at converting NO₂ to NO at ambient temperature.

[00165] The inlet 105 also may receive a gas flow, e.g. air flow, having NO₂, for example, from source of NO₂. A source of NO₂ can include a pressurized bottle of NO₂, which also may be referred to as a tank of NO₂. The inlet 105 also may receive a gas flow with NO₂ in nitrogen (N₂), air, or oxygen (O₂). The conversion can occur over a wide concentration range. Experiments have been carried out at concentrations in a gas including from about 2 ppm NO₂ to 100 ppm NO₂, and even to over 1000 ppm NO₂. In one example, a cartridge that was approximately 6 inches long and had a diameter of 1.5-inches was packed with silica gel that had first been soaked in a saturated aqueous solution of ascorbic acid. The moist silica gel was prepared using ascorbic acid (i.e., vitamin C) designated as A.C.S reagent grade 99.1% pure from Aldrich Chemical Company and silica gel from Fischer Scientific International, Inc., designated as S8 32-1, 40 of Grade of 35 to 70 sized mesh. Other sizes of silica gel also can be effective. For example, silica gel having an eighth-inch diameter could also work.

[00166] The surface active material, e.g. silica gel, can be moistened with a solution including a reducing agent, e.g. a saturated solution including a reducing agent. For example, a saturated solution of ascorbic acid in water; more specifically, the saturated solution can be a saturated solution that had been prepared by mixing 35% by weight ascorbic acid in water, stirring, and straining the water/ascorbic acid mixture through the silica gel, followed by draining. The conversion of NO₂ to NO can proceed well when the surface active material (e.g. silica gel) coated with a reducing agent (e.g. ascorbic acid) is moist. The conversion of NO₂ to NO may not proceed well with a reducing agent alone, for example, in an aqueous solution of ascorbic acid alone.

[00167] The cartridge can be filled with the wet silica gel/reducing agent. For example, a cartridge filled with the wet silica gel/ascorbic acid was able to convert 1000 ppm of NO₂ in air to NO at a flow rate of 150 ml per minute, quantitatively, non-stop for over 12 days. A wide variety of flow rates and NO₂ concentrations have been successfully tested, ranging from only a few ml per minute to flow rates of up to 5,000 ml per minute. Any appropriate reducing agent that can convert NO₂ or N₂O₄ to NO can be used as determined by a person of skill in the art. For example, the reducing agent can include a hydroquinone, glutathione, thiols, nitrites, and/or one or more reduced metal salts such as Fe(II), Mo(VI), NaI, Ti(III) or Cr(III). The reducing agent can be an antioxidant. The antioxidant can be an aqueous solution of an antioxidant. The antioxidant can be ascorbic acid, alpha tocopherol, or gamma tocopherol. Any appropriate antioxidant can be used depending on the activities and properties as determined by a person of skill in the art. The antioxidant can be used dry or wet.

[00168] The antioxidant/surface-active material GENO cartridge may be used for inhalation therapy. In one such example, the GENO cartridge can be used as a NO₂ scrubber for NO inhalation therapy that delivers NO from a pressurized bottle source. The GENO cartridge can be used to remove any NO₂ that chemically forms during inhalation therapy. This GENO cartridge can be used to help ensure that no harmful levels of NO₂ are inadvertently inhaled by the patient.

[00169] The GENO cartridge may be used to supplement or replace some or all of the safety devices used during inhalation therapy in conventional NO inhalation therapy. For example, one type of safety device can warn of the presence of NO₂ in air when the concentration of NO₂ exceeds a preset or predetermined limit, usually 1 part per million or

greater of NO₂. Such a safety device may be unnecessary when a GENO cartridge is positioned in a NO delivery system just prior to the patient breathing the NO laden air. The GENO cartridge can convert any NO₂ to NO just prior to the patient breathing the NO laden gas, which can make a device to warn of the presence of NO₂ in gas unnecessary.

5 [00170] Alternatively or additionally, a NO₂ removal cartridge can be inserted just before the attachment of the delivery system to the patient to further enhance safety and help ensure that all traces of the toxic NO₂ have been removed. The NO₂ removal cartridge may be a GENO cartridge used to remove any trace amounts of NO₂. Alternatively, the NO₂ removal cartridge may include heat-activated alumina. A cartridge with heat-activated
10 alumina, such as supplied by Fisher Scientific International, Inc., designated as A505-212, of 8-14 sized mesh can be effective at removing low levels of NO₂ from an air or oxygen stream, and yet can let NO gas pass through without loss. Activated alumina, and other high surface area materials like it, can be used to scrub NO₂ from a NO inhalation line.

[00171] The various embodiments described are provided by way of illustration only
15 and should not be construed to limit the claimed invention. Those skilled in the art will readily recognize various modifications and changes that may be made to the claimed invention without following the example embodiments and applications illustrated and described herein, and without departing from the true spirit and scope of the claimed invention, which is set forth in the claims.

20 References

1. Barst, R.J., et al. A comparison of continuous intravenous epoprostenol (prostacyclin) with conventional therapy for primary pulmonary hypertension. The Primary Pulmonary Hypertension Study Group. N Engl J Med 334, 296-302 (1996).
2. Barst, R.J., et al Long-term outcome in pulmonary arterial hypertension patients
25 treated with subcutaneous treprostinil. Eur Respir J. 28: 1195-1203 (2006).
3. Barst, R.J., et al. Updated evidence-based treatment algorithm in pulmonary arterial hypertension. J Am Coll Cardiol. 54, S78-84. (2009).
4. Bhorade, S., et al. Response to inhaled nitric oxide in patients with acute right heart syndrome. Am J Respir Crit Care Med 159, 571-579 (1999).

5. Bloch, K.D., Ichinose, F., Roberts, J.D., Jr. & Zapol, W.M. Inhaled NO as a therapeutic agent. *Cardiovasc Res* 75, 339-348 (2007).
6. Bogaard, H.J., et al. Chronic Pulmonary Artery Pressure Elevation Is Insufficient to Explain Right Heart Failure. *Circulation* 120, 1951-1960 (2009).
- 5 7. Flox Camacho, Á., et al. Transition From Prostacyclin to Bosentan in Five Patients With Severe Pulmonary Hypertension: The Switch Is Possible. *Revista Española de Cardiología* 59, 737-739 (2006).
8. Chakinala, M. Changing the Prognosis of Pulmonary Arterial Hypertension: Impact of Medical Therapy. Vol. 26 409-416 (Medscape, 2005).
- 10 9. Champion, H.C., Michelakis, E.D. & Hassoun, P.M. Comprehensive Invasive and Noninvasive Approach to the Right Ventricle-Pulmonary Circulation Unit: State of the Art and Clinical and Research Implications. *Circulation* 120, 992-1007 (2009).
10. Chin, K.M. & Rubin, L.J. Pulmonary arterial hypertension. *J Am Coll Cardiol*. 51, 1527-1538. (2008).
- 15 11. D'Alonzo, G.E., *et al.* Survival in patients with primary pulmonary hypertension. Results from a national prospective registry. *Ann Intern Med* **115**, 343-349 (1991).
12. Di Salvo, T., Mathier, M., Semigran, M. & Dec, G. Preserved right ventricular ejection fraction predicts exercise capacity and survival in advanced heart failure. *J Am Coll Cardiol* 25, 1143-1153 (1995).
- 20 13. Diaz-Guzman, E., Heresi, G.A., Dweik, R.A. & Minai, O.A. Long-term experience after transition from parenteral prostanoids to oral agents in patients with pulmonary hypertension. *Respir Med* 102, 681-689 (2008).
14. Diller, G.-P., Dimopoulos, K., Kafka, H., Ho, S.Y. & Gatzoulis, M.A. Model of chronic adaptation: right ventricular function in Eisenmenger syndrome. *European Heart Journal Supplements* 9, H54-H60 (2007).
- 25 15. Damman, K., et al. Increased Central Venous Pressure Is Associated With Impaired Renal Function and Mortality in a Broad Spectrum of Patients With Cardiovascular Disease. *J Am Coll Cardiol* 53, 582-588 (2009).

16. Gaine, S.P. & Rubin, L.J. Primary pulmonary hypertension. *Lancet*. **352**, 719-725. (1998).
17. Ghofrani, H.A., Wilkins, M.W. & Rich, S. Uncertainties in the Diagnosis and Treatment of Pulmonary Arterial Hypertension. *Circulation* 118, 1195-1201 (2008).
- 5 18. Gomberg-Maitland, M., et al. Transition from intravenous epoprostenol to intravenous treprostinil in pulmonary hypertension. *Am J Respir Crit Care Med* 172, 1586-1589 (2005).
19. Greyson, C.R. Pathophysiology of right ventricular failure. *Critical Care Medicine* 36, S57-S65 10.1097/1001.CCM.0000296265.0000252518.0000296270 (2008).
- 10 20. Haddad, E., Lowson, S.M., Johns, R.A. & Rich, G.F. Use of inhaled nitric oxide perioperatively and in intensive care patients. *Anesthesiology* 92, 1821-1825 (2000).
21. Haddad, F., Hunt, S.A., Rosenthal, D.N. & Murphy, D.J. Right Ventricular Function in Cardiovascular Disease, Part I: Anatomy, Physiology, Aging, and Functional Assessment of the Right Ventricle. *Circulation* 117, 1436-1448 (2008).
- 15 22. Haddad, F., Doyle, R., Murphy, D.J. & Hunt, S.A. Right Ventricular Function in Cardiovascular Disease, Part II: Pathophysiology, Clinical Importance, and Management of Right Ventricular Failure. *Circulation* 117, 1717-1731 (2008).
23. Handoko, M.L., et al. Perspectives on novel therapeutic strategies for right heart failure in pulmonary arterial hypertension: lessons from the left heart. *European Respiratory Review* 19, 72-82 (2010).
- 20 24. Hemnes, A.R. & Kawut, S.M. The right ventricle in pulmonary hypertension: from dogma to data. *Am J Respir Crit Care Med* 182, 586-588 (2010).
- 25 25. Hinderliter, A.L., et al. Effects of Long-term Infusion of Prostacyclin (Epoprostenol) on Echocardiographic Measures of Right Ventricular Structure and Function in Primary Pulmonary Hypertension. *Circulation* 95, 1479-1486 (1997).
26. Hoeper, M.M., Markevych, I., Spiekerkoetter, E., Welte, T. & Niedermeyer, J. Goal-oriented treatment and combination therapy for pulmonary arterial hypertension. *European Respiratory Journal* 26, 858-863 (2005).

27. Humbert, M., et al. Survival in patients with idiopathic, familial, and anorexigen-associated pulmonary arterial hypertension in the modern management era. *Circulation* 122, 156-163 (2010).
28. Inglessis, I., et al. Hemodynamic effects of inhaled nitric oxide in right ventricular myocardial infarction and cardiogenic shock. *J Am Coll Cardiol* 44, 793-798 (2004).
29. Johnson, R.F., Loyd, J.E., Mullican, A.L., Fink, C.A. & Robbins, I.M. Long-term follow-up after conversion from intravenous epoprostenol to oral therapy with bosentan or sildenafil in 13 patients with pulmonary arterial hypertension. *J Heart Lung Transplant* 26, 363-369 (2007).
30. Krasuski, R.A., Wang, A., Harrison, J.K., Tapson, V.F. & Bashore, T.M. The response to inhaled nitric oxide in patients with pulmonary artery hypertension is not masked by baseline vasodilator use. *Am Heart J* 150, 725-728 (2005).
31. Kim, N.H., Channick, R.N. & Rubin, L.J. Successful withdrawal of long-term epoprostenol therapy for pulmonary arterial hypertension. *Chest*. 124, 1612-1615. (2003).
32. Krasuski, R.A., et al. Inhaled nitric oxide selectively dilates pulmonary vasculature in adult patients with pulmonary hypertension, irrespective of etiology. *Journal of the American College of Cardiology* 36, p2204-2211 (2000).
33. Kevin, L.G. & Barnard, M. Right ventricular failure. *Continuing Education in Anaesthesia, Critical Care & Pain* 7, 89-94 (2007).
34. Lang, I.M. Management of acute and chronic RV dysfunction. *European Heart Journal Supplements* 9, H61-H67 (2007).
35. Lepore, J.J., et al. Hemodynamic effects of sildenafil in patients with congestive heart failure and pulmonary hypertension: Combined administration with inhaled nitric oxide. *Chest* 127, 1647-1653 (2005).
36. Leci, E., et al. Prognostic Parameters For Survival Of Patients With Pulmonary Arterial Hypertension Associated To Connective Tissue Diseases Treated With Targeted Therapy. *Am. J. Respir. Crit. Care Med*. 181, A4844- (2010).

37. Loh, E., Stamler, J.S., Hare, J.M., Loscalzo, J. & Colucci, W.S. Cardiovascular effects of inhaled nitric oxide in patients with left ventricular dysfunction. *Circulation*. 90, 2780-2785. (1994).
38. Lopez-Candales, A., et al. Right ventricular dyssynchrony in patients with pulmonary hypertension is associated with disease severity and functional class. *Cardiovascular Ultrasound* 3, 23 (2005).
39. McLaughlin, V.V., Genthner, D.E., Panella, M.M. & Rich, S. Reduction in pulmonary vascular resistance with long-term epoprostenol (prostacyclin) therapy in primary pulmonary hypertension. *N Engl J Med*. 338, 273-277. (1998).
40. Moraes, D.L., Colucci, W.S. & Givertz, M.M. Secondary Pulmonary Hypertension in Chronic Heart Failure : The Role of the Endothelium in Pathophysiology and Management. *Circulation* 102, 1718-1723 (2000).
41. Matthews, J.C., Koelling, T.M., Pagani, F.D. & Aaronson, K.D. The Right Ventricular Failure Risk Score: A Pre-Operative Tool for Assessing the Risk of Right Ventricular Failure in Left Ventricular Assist Device Candidates. *J Am Coll Cardiol* 51, 2163-2172 (2008).
42. Naeije, R. Pulmonary Hypertension and Right Heart Failure in Chronic Obstructive Pulmonary Disease. *Proc Am Thorac Soc* 2, 20-22 (2005).
43. Natori, S., et al. Inhaled nitric oxide modifies left ventricular diastolic stress in the presence of vasoactive agents in heart failure. *Am J Respir Crit Care Med* 167, 895-901 (2003).
44. Pilcher, D.V., et al. High central venous pressure is associated with prolonged mechanical ventilation and increased mortality after lung transplantation. *J Thorac Cardiovasc Surg* 129, 912-918 (2005).
45. Price, L.C., Wort, S.J., Finney, S.J., Marino, P.S. & Brett, S.J. Pulmonary vascular and right ventricular dysfunction in adult critical care: current and emerging options for management: a systematic literature review, (2010).
46. Rich, G.F. Management of the Patient with Pulmonary Hypertension and Right Ventricular Failure. *ASA Refresher Courses in Anesthesiology* 33, 203-212 (2005).

47. Rich, S., et al. The short-term effects of digoxin in patients with right ventricular dysfunction from pulmonary hypertension. *Chest* 114, 787-792 (1998).
48. Raffy, O., et al. Clinical Significance of the Pulmonary Vasodilator Response During Short-term Infusion of Prostacyclin in Primary Pulmonary Hypertension. *Circulation* 93, 484-488 (1996).
5. 49. Roeleveld, R.J., et al. Effects of epoprostenol on right ventricular hypertrophy and dilatation in pulmonary hypertension. *Chest* 125, 572-579 (2004).
- 10 50. Safdar, Z. Outcome of pulmonary hypertension subjects transitioned from intravenous prostacyclin to oral bosentan. *Respiratory medicine* 103, 1688-1692 (2009).
51. Sánchez, M.A.G. & Torbicki, A. Foreword: right ventricular function and pulmonary hypertension. *European Heart Journal Supplements* 9, H3-H4 (2007).
- 15 52. Schattke, S., et al. Early right ventricular systolic dysfunction in patients with systemic sclerosis without pulmonary hypertension: a Doppler Tissue and Speckle Tracking echocardiography study. *Cardiovascular Ultrasound* 8, 3 (2010).
53. Sidhu MS, A.J., Saggar Rajan, Saggar Rajeev, Belperio J, Ardehali A, Ross DJ. TRANSITION FROM PROSTANOID TO COMBINATION ORAL THERAPY FOR PULMONARY ARTERIAL HYPERTENSION. *E-Journal of Cardiology* 1, 8 (2011).
- 20 54. Simon, M.A. & Pinsky, M.R. Right Ventricular Dysfunction and Failure in Chronic pressure Overload. *Cardiology Research and Practice* 2011, 1 (2011).
55. Skoro-Sajer, N., et al. Pulmonary vascular reactivity and prognosis in patients with chronic thromboembolic pulmonary hypertension: a pilot study. *Circulation* 119, 298-305 (2009).
- 25 56. Steiner, M.K., et al. Conversion to bosentan from prostacyclin infusion therapy in pulmonary arterial hypertension: a pilot study. *Chest*. 130, 1471-1480. (2006).
57. Suleman, N. & Frost, A.E. Transition from epoprostenol and treprostinil to the oral endothelin receptor antagonist bosentan in patients with pulmonary hypertension. *Chest*. 126, 808-815. (2004).

58. Triantafyllou, K., Kranidis, A., Karabinos, E., Grassos, H. & Babalis, D. Clinical implications of the echocardiographic evaluation of right ventricular function on the long axis using newer techniques. *Hellenic J Cardiol* 51, 42-48 (2010).
 59. Tayyareci, Y., et al. Early detection of right ventricular systolic dysfunction by using myocardial acceleration during isovolumic contraction in patients with mitral stenosis. *Eur J Echocardiogr* 9, 516-521 (2008).
 60. Voelkel, N.F., et al. Right ventricular function and failure: report of a National Heart, Lung, and Blood Institute working group on cellular and molecular mechanisms of right heart failure. *Circulation* 114, 1883-1891 (2006).
 61. van Wolferen, S.A., et al. Right ventricular reverse remodelling after sildenafil in pulmonary arterial hypertension. *Heart* 92, 1860-1861 (2006).
 62. Vlahakes, G.J. Management of Pulmonary Hypertension and Right Ventricular Failure: Another Step Forward. *Ann Thorac Surg* 61, 1051-1052 (1996).
 63. Zakir, R.M., Al-Dehneh, A., Maher, J., Saric, M. & Berkowitz, R.L. Right ventricular failure in patients with preserved ejection fraction and diastolic dysfunction: an underrecognized clinical entity. *Congest Heart Fail* 13, 164-169 (2007).
 64. Gaine, S.P. & Rubin, L.J. Primary pulmonary hypertension. *Lancet*. 352, 719-725. (1998).
- 20 **[00172]** Each of the references above is incorporated by reference in its entirety.

What is claimed:

1. A method for treating pulmonary arterial hypertension in a patient, comprising:
administering a vasoactive agent to a patient with pulmonary arterial hypertension;
determining whether the patient has an enlarged right ventricle or right ventricular
5 dysfunction; and
weaning the patient off of the prostanoid if the patient does not have an enlarged
right ventricle or right ventricular dysfunction.
2. The method of claim 1, wherein the vasoactive agent is inhaled nitric oxide.
- 10 3. The method of claim 1 or 2, wherein determining whether the patient has an
enlarged right ventricle or right ventricular dysfunction includes monitoring an at least one
indicator of hemodynamics of the patient before and after administering the vasoactive
agent to a patient.
- 15 4. The method of any one of claims 1-3, wherein determining whether the patient has
an enlarged right ventricle or right ventricular dysfunction includes calculating a change in
the at least one indicator of the hemodynamics of the patient between before the vasoactive
agent was administered and after the vasoactive agent was administered.
- 20 5. The method of claim 3 or 4, wherein the at least one indicator of hemodynamics
includes pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output.
6. The method of any one of claims 3-5, wherein the at least one indicator of
25 hemodynamics includes pulmonary vascular resistance and cardiac output, and wherein
determining whether the patient has an enlarged right ventricle or right ventricular
dysfunction further includes determining the presence of an increase in cardiac output after
administration of the vasoactive agent which correlates with a decrease in pulmonary
vascular resistance after administration of the vasoactive agent, such that an increase in
30 cardiac output which correlates with a decrease in pulmonary vascular resistance indicates
that an enlarged right ventricle or right ventricular dysfunction is not present.

7. The method of any one of claims 1-6, wherein administering a vasoactive agent includes communicating an air flow including nitrogen dioxide through a receptacle, the receptacle including an inlet, an outlet, a surface-active material and a reducing agent, wherein the inlet is configured to receive the flow of nitrogen dioxide; and communicating
5 the air flow to the outlet through the surface-active material.

8. A method for treating pulmonary arterial hypertension in a patient, comprising:
administering a vasoactive agent to a patient with pulmonary arterial hypertension, wherein the patient has previously been administered a prostanoid;
10 determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent; and
weaning the patient off of the prostanoid if the patient demonstrates an acute vasodilator response to the vasoactive agent.

15 9. The method of claim 8, wherein the vasoactive agent is inhaled nitric oxide.

10. The method of claim 8 or 9, wherein determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent includes monitoring an at least one indicator of hemodynamics of the patient.

20 11. The method of claim 10, wherein monitoring includes monitoring the at least one indicator of the hemodynamics of the patient before and after administering the vasoactive agent to the patient.

25 12. The method of claim 10 or 11, wherein the at least one indicator of hemodynamics includes pulmonary vascular resistance, mean pulmonary arterial pressure or cardiac output.

13. The method of any one of claims 8-12, wherein determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent includes calculating a
30 change in at least one indicator of the hemodynamics of the patient between before the vasoactive agent was administered and after the vasoactive agent was administered.

14. The method of any one of claims 10-13, wherein the at least one indicator of hemodynamics includes mean pulmonary arterial pressure, and wherein determining whether the patient demonstrates an acute vasodilator response to the vasoactive agent further includes determining the presence of a change in mean pulmonary arterial pressure of greater than 10 mm Hg between before the vasoactive agent was administered and after the vasoactive agent was administered.

15. The method of any one of claims 8-14, wherein administering a vasoactive agent includes communicating an air flow including nitrogen dioxide through a receptacle, the receptacle including an inlet, an outlet, a surface-active material and a reducing agent, wherein the inlet is configured to receive the flow of nitrogen dioxide; and communicating the air flow to the outlet through the surface-active material.

16. A method for monitoring a patient with pulmonary arterial hypertension, comprising:

administering inhaled nitric oxide to a patient with pulmonary arterial hypertension, wherein administering inhaled nitric oxide includes

communicating an air flow including nitrogen dioxide through a receptacle, the receptacle including an inlet, an outlet, a surface-active material and a reducing agent, wherein the inlet is configured to receive the flow of nitrogen dioxide;

communicating the air flow to the outlet through the surface-active material;

and

converting the gaseous nitrogen dioxide to nitric oxide; and
monitoring an at least one indicator of hemodynamics of the patient.

17. The method of claim 16, wherein the reducing agent is an antioxidant.

18. The method of claim 16 or 17, wherein the reducing agent comprises ascorbic acid, alpha tocopherol or gamma tocopherol.

19. The method of any one of claims 16-18, wherein the reducing agent is included on the surface-active material.

20. The method of any one of claims 16-19, wherein the reducing agent is coated on the surface-active material.

21. The method of any one of claims 16-20, wherein the surface-active material
5 comprises a substrate that retains water.

22. The method of any one of claims 16-21, wherein the surface-active material comprises a silica gel.

10 23. A method for diagnosing right ventricular dysfunction, comprising:
administering a vasoactive agent to a patient suspected of having right ventricular
dysfunction;
monitoring an at least one indicator of hemodynamics of the patient; and
determining whether the patient has right ventricular dysfunction.

15 24. The method of claim 23, wherein the vasoactive agent is inhaled nitric oxide.

25. The method of claim 23 or 24, wherein the at least one indicator of hemodynamics
includes pulmonary vascular resistance and cardiac output, and wherein determining
20 whether the patient has right ventricular dysfunction further includes determining the
presence of an increase in cardiac output after administration of the vasoactive agent which
correlates with a decrease in pulmonary vascular resistance after administration of the
vasoactive agent, such that an increase in cardiac output which correlates with a decrease in
pulmonary vascular resistance indicates that the patient does not have right ventricular
25 dysfunction.

26. The method of any one of claims 23-25, wherein administering a vasoactive agent
includes communicating an air flow including nitrogen dioxide through a receptacle, the
receptacle including an inlet, an outlet, a surface-active material and a reducing agent,
30 wherein the inlet is configured to receive the flow of nitrogen dioxide; and communicating
the air flow to the outlet through the surface-active material.




FIGURE 1

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/39401

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01N 33/24; A61K 31/13 (2012.01)

USPC - 514/645

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC - 514/645

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
USPC - 424/718; 514/645; 544/297; 546/335; 548/391; 560/194 (see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

*** Please see extra sheet ***

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Oudiz et al. Cardiac Catheterization in Pulmonary Arterial Hypertension: An Updated Guide to Proper Use. Advances in Pulmonary Hypertension: Official Journal of the Pulmonary Hypertension Association, 2005, Vol 4, No 3, pp 15-25; especially pg 22, col 1, para 6 and col 2, para 4-5; pg 24, col 1, para 1-3	1-3, 23-25
Y		8-11, 16-18
Y	US 2009/0197922 A1 (Maitland et al.) 06 August 2009 (06.08.2009) para [0011], [0022], [0026]	8-11
Y	WO 2011/063335 A1 (Fine et al.) 26 May 2011 (26.05.2011) para [0004], [0006], [0018]-[0019], [0031]-[0033]	16-18
A,P	US 2012/0093948 A1 (Fine et al.) 19 April 2012 (19.04.2012) entire document	1-3, 8-11, 16-18, 23-25

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

01 August 2012 (01.08.2012)

Date of mailing of the international search report

17 AUG 2012

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/39401

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4-7, 12-15, 19-22 and 26
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 12/39401

*** Databases: WEST (PGPB, USPT, USOC, EPAB, JPAB); Google, Google Scholar

*** Search Terms Used: Geno, Watts, Roscigno, Fine, Rubin, vasoactive, vasodilator, nitric oxide, pulmonary, arterial, hypertension, prostanoid, prostaglandin, thromboxane, prostacyclin, epoprostenol, trepostinil, iloprost, orberaprost, vascular remodeling, hemodynamic, ventricle, enlarged, dysfunction, monitoring, monitor, indicator, indicating, tocopherol, cardiac output, vascular resistance, arterial pressure, cardiac output,