




(86) **Date de dépôt PCT/PCT Filing Date:** 2015/06/30
(87) **Date publication PCT/PCT Publication Date:** 2016/01/07
(45) **Date de délivrance/Issue Date:** 2022/12/06
(85) **Entrée phase nationale/National Entry:** 2016/12/29
(86) **N° demande PCT/PCT Application No.:** IL 2015/050670
(87) **N° publication PCT/PCT Publication No.:** 2016/001918
(30) **Priorité/Priority:** 2014/07/01 (US62/019,439)

(51) **Cl.Int./Int.Cl.** **A61F 13/42** (2006.01),
A61F 13/15 (2006.01), **A61L 15/16** (2006.01),
A61L 15/56 (2006.01)
(72) **Inventeurs/Inventors:**
HERSHCOVITZ-FUNDOIANO, YIFAT, IL;
PARDO-YISSAR, VERED, IL;
KESSARY SHOHAM, HADAR, IL
(73) **Propriétaire/Owner:**
COMMON SENSE LTD., IL
(74) **Agent:** RIDOUT & MAYBEE LLP

(54) **Titre : COMPOSITION DE DIAGNOSTIC POUR IDENTIFIER UN LIQUIDE AMNIOTIQUE**
(54) **Title: DIAGNOSTIC COMPOSITION FOR IDENTIFYING AMNIOTIC FLUID**



(57) **Abrégé/Abstract:**

An indicator composition for detecting amniotic fluid in vaginal secretion comprises a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein the composition is devoid of a wetting agent. An article for detecting amniotic fluid in a biological fluid secreted from a person comprises an absorbent material for absorbing the biological fluid, and the indicator composition. A kit comprises the article and instructions for use. A method for diagnosing amniotic fluid leakage in a biological fluid secreted from a person comprises positioning the article to receive the biological fluid secreted from the person, contacting the article with vaginal secretion; and viewing the article, wherein a color change indicates the presence of amniotic fluid.

ABSTRACT

An indicator composition for detecting amniotic fluid in vaginal secretion comprises a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein the composition is devoid of a wetting agent. An article for detecting amniotic fluid in a biological fluid secreted from a person comprises an absorbent material for absorbing the biological fluid, and the indicator composition. A kit comprises the article and instructions for use. A method for diagnosing amniotic fluid leakage in a biological fluid secreted from a person comprises positioning the article to receive the biological fluid secreted from the person, contacting the article with vaginal secretion; and viewing the article, wherein a color change indicates the presence of amniotic fluid.

DIAGNOSTIC COMPOSITION FOR IDENTIFYING AMNIOTIC FLUID

FIELD OF THE INVENTION

The present invention relates to the field of medical diagnostics and more
5 specifically, to identification of amniotic fluid.

BACKGROUND OF THE INVENTION

Premature rupture of the amniotic sac is a common disorder that happens to occur
in around 10% of pregnant women. Failure to treat premature rupture may cause intra-
10 amnion infection, disruption in fetal lung development and even perinatal death.

Several approaches for detecting amniotic leakage are known in the art, including
chromatographic detection of a color change in a diagnostic article following contact with
vaginal secretion.

A number of devices involving panty shields, swabs and the like, with indicators
15 aimed to identify amniotic fluid were developed, for example, in U.S. Patent Nos.
5,897,834; 6,126,597 and 6,149,590. These devices can be worn, or otherwise used, by the
user and whenever there is a secretion it is absorbed and further detected.

US Patent No. 6,921,647 and Continuation-in-Parts thereof, including, US Patent
Nos. 7,314,752; 7,541,177 and 7,947,467 disclose secretion-monitoring article for
20 identifying a secreted biological fluid, including, amniotic fluid, comprising: a body that
includes an absorbent material and an indicator system having at least one pH determining
member. The pH determining member includes a chemical composition which reacts with
fluids containing urea differently than with fluids that do not contain urea, in a manner
such that the former can be distinguished from the latter while avoiding false positive
25 indications. The pH determining member may further, or alternatively, include a
hydrophobic chemical composition that reacts with fluids based on the buffer capacity
thereof, such that fluids having different buffer capacities can be distinguished.

It has been also proposed to detect the rupture of the amniotic sac by employing
immunochemical analysis of the proteins in the amniotic fluid, such as, alpha-fetoprotein,
30 prolactin, fibronectin, and insulin-like growth-factor binding protein 1. For example, US
Patent No. 8,114,610 discloses a method of detecting amniotic fluid in a sample

comprising a vaginal secretion of a pregnant woman using a combination of PAMG-1-specific monoclonal antibodies.

There is an unmet need for an improved diagnostic system that can differentiate between amniotic fluid and an interfering biological fluid, such as, urine, with minimal false positive results, while providing a reliable indication which is stable over time, and while reducing the amount of time required to get the reliable result.

SUMMARY OF THE INVENTION

The present invention discloses an indicator composition for detecting amniotic fluid in vaginal secretion comprising: a plurality of polymers; a plasticizer; a surfactant; an ion-balance reagent; an acid and an indicator agent, wherein said composition is devoid of a wetting agent. The present invention further discloses diagnostic articles for identifying amniotic fluid and methods of using same.

Advantageously, the articles of the invention may operate right after contacting the vaginal secretion. Specifically, the articles and diagnostic compositions disclosed herein do not require any drying time and/or do not require waiting for urine evaporation. Thus, detection of amniotic fluid (or other fluids) by the articles and compositions disclosed herein is a one-step process, namely, upon contact of liquid (amniotic fluid or urine) with an article comprising the diagnostic composition, detection can be carried out on the article as is, right after being in contact with the fluids. Without wishing to be bound by any theory or mechanism, it is assumed that this advantage is rendered by the unique combination of the ingredients in the diagnostic formulations of the invention, or may be specifically attributed to the lack of wetting agent, the presence of an aromatic carboxylic acid, the presence of an hydrophilic non-ionic polymer as surfactant, the presence of a tri-ester non-cyclic plasticizer or the combination thereof, together with the rest of the ingredients.

Furthermore, the articles of the invention enable initial self-detection and do not require the patient to visit health-care professional in order to distinguish between positive and false-positives. Accordingly, using the article of the invention saves a significant amount of time required to determine if amniotic fluid is actually leaking.

In some embodiments, there is provided an indicator composition for detecting amniotic fluid in vaginal secretion comprising: a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent.

5 In some embodiments, said aromatic carboxylic acid comprises salicylic acid, benzoic acid or a combination thereof.

In some embodiments, the aromatic carboxylic acid is salicylic acid.

In some embodiments, the hydrophilic surfactant is a non-ionic surfactant.

In some embodiments, the hydrophilic surfactant is a polymer.

10 In some embodiments, the hydrophilic surfactant comprises a block copolymer surfactant.

In some embodiments, the indicator agent is negatively charged.

In some embodiments, the ion-balance reagent is a quaternary ammonium salt.

In some embodiments, the non-cyclic plasticizer is a tri-ester plasticizer.

15 In some embodiments, there is provided an indicator composition for detecting amniotic fluid in vaginal secretion consisting of a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent.

20 In some embodiments, there is provided an article for detecting amniotic fluid in vaginal secretion comprising: an absorbent material for absorbing a biological fluid secreted from a person, and an indicator composition, comprising: a plurality of polymers; a non-cyclic plasticizer; an hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent.

25 In some embodiments, the absorbent material is selected from the group consisting of a swab, gauze, panty shield, hygienic napkin, diaper and interlabial absorbent structure.

In some embodiments, there is provided a method for diagnosing amniotic fluid leakage in vaginal secretion comprising:

(a) positioning a secretion-monitoring article to receive a biological fluid secreted from the subject, the article comprising an indicator composition comprising: a plurality of polymers; a non-cyclic plasticizer; an hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent;

(b) contacting said article with vaginal secretion; and

(c) viewing the article, wherein a color change indicates the presence of amniotic fluid.

In some embodiments, said color change is a dark stain relative to background of the secretion monitoring article. In some embodiments, the dark stain is a dark blue stain or a dark green stain.

In some embodiments, said viewing is performed within less than 10 minutes after said contacting. In some embodiments, said viewing is performed within less than 9 minutes after said contacting. In some embodiments, said viewing is performed within less than 8 minutes after said contacting. In some embodiments, said viewing is performed within less than 7 minutes after said contacting. In some embodiments, said viewing is performed within less than 6 minutes after said contacting. In some embodiments, said viewing is performed within less than 5 minutes after said contacting. In some embodiments, said viewing is performed within less than 4 minutes after said contacting. In some embodiments, said viewing is performed within less than 3 minutes after said contacting. In some embodiments, said viewing is performed within less than 2 minutes after said contacting.

In some embodiments, said indicator composition is consisting of a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent.

In some embodiments, there is provided a kit comprising the article for detecting amniotic fluid in vaginal secretion, and instructions for use.

In some embodiments, the kit further comprises a color scale presenting the color expected to appear upon contact of the article with amniotic fluid and the color expected to appear upon contact of the article with urine and any non-amniotic fluid.

Further embodiments, features, advantages and the full scope of applicability of the present invention will become apparent from the detailed description and drawings given herein after. However, it should be understood that the detailed description, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE FIGURES

Exemplary embodiments are illustrated in referenced figures. It is intended that the embodiments and figures disclosed herein are to be considered illustrative rather than restrictive. The figures are listed below.

Fig. 1 shows the relative specificity of the articles disclosed herein (including ALM1 which is also termed "AL-SENSE Blue") at the following time points: 0-2 min., 5 min., 10 min., 20 min., 30 min 1 hr. and 2 hrs.

Fig. 2 exhibits the results obtained from contacting urine samples (pH 7) with one articles encompassing candidate composition - ALM1 and test compositions 3Mar1 and 3Mar2.

Fig. 3A demonstrates a photo of an absorbent article containing the ALM1 formulation 30 min. following contact with amniotic fluid.

Fig. 3B demonstrates a photo of an absorbent article containing the ALM1 formulation 30 min. following contact with urine.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides an indicator composition, a diagnostic article comprising same, and methods of use thereof for obtaining an immediate, clear and reliable identification of amniotic fluid leakage.

The terms "amniotic fluid leakage" and "premature rupture of the amniotic sac" are interchangeable and relate to the spontaneous rupture of the membranes (the 'amniotic sac' is often termed 'fetal membrane') 24 or more hours before the onset of labor. About 30-50% of premature ruptures occur before the 37th week of pregnancy. As amniotic fluid

leakage is associated with a significant increase in the risk of an intrauterine infection and disturbance of development of the fetal lung system, definitive and reliable diagnosis of the rupture is material. Intrauterine penetration of such infections increases both maternal and perinatal morbidity and mortality by about ten percent. Immediate diagnosis of a
5 rupture at 38 to 40 weeks of pregnancy is crucial, since once detected delivery should be induced as soon as possible. The rupture diagnosis is also important before 37 weeks of pregnancy because it enables prevention of intra-amnion infection and the stimulation of fetal lung development.

Due to the severe consequences of amniotic fluid leakage, pregnant women
10 undergo severe stress and often go to a health-care professional upon secretion of any liquid from the vicinity of the vagina. The health-care professional looks for the presence of amniotic fluid by checking the pH of the vaginal secretions - amniotic fluid is having a pH of between 6.7 and 7.5. Since pregnant women often have urinary incontinence and since urine typically has a pH of between 5.0 and 8.0, if only pH is checked, a false
15 positive result may occur: urine being identified as amniotic fluid. Consequently, it is necessary that a vaginal secretion be examined using a microscope for the presence of a fern-shaped pattern indicative of amniotic fluid.

As the time between the fluid secretion and the arrival at the health-care professional may be long, there is often no evidence of amniotic fluid upon examination.
20 The secretion may mistakenly assumed to be urine, which may result with tragic consequences. On the other hand, the healthcare professional may decide to err on the side of caution, misdiagnosing the secretion of urine as amniotic fluid leading to an unnecessary hospitalization and patient stress.

Various diagnostic indicators are known, however, an obstacle of many is that they
25 often provide "false positives" due to changes in pH on drying, interfering biological fluids and repetitive cycles of drying/wetting. The "false positive" readings can be stressful and time consuming to the user. A device that minimizes these "false positive" readings is disclosed herein. The present invention is an improvement over the prior art, providing a diagnostic composition which is more accurate, reliable and convenient for the
30 user. The diagnostic composition of the present invention comprises an indicator agent that identifies the pH associated with the fluid, a plurality of polymers, a tri-ester non-

cyclic plasticizer, hydrophilic non-ionic polymer surfactant, an ion-balance reagent and at least one aromatic carboxylic acid, wherein said composition is devoid of a wetting agent.

As used herein and in the claims the term “plurality” means at least two.

In some embodiments, there is provided an indicator composition for detecting
5 amniotic fluid in vaginal secretion comprising: a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent.

In some embodiments there is provided an indicator composition for detecting amniotic fluid in vaginal secretion comprising: a plurality of polymers; a tri-ester non-cyclic plasticizer; a hydrophilic non-ionic polymer surfactant; an aromatic carboxylic
10 acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent.

In some embodiments, the indicator composition is consisting of a plurality of polymers; a non-cyclic plasticizer; a hydrophilic surfactant; an aromatic carboxylic acid;
15 an ion-balance reagent and an indicator agent.

In some embodiments, the indicator composition is consisting of a plurality of polymers; a tri-ester non-cyclic plasticizer; a hydrophilic non-ionic polymer surfactant; an aromatic carboxylic acid; an ion-balance reagent and an indicator agent.

In some embodiments there is provided an article for detecting amniotic fluid in
20 vaginal secretion comprising: an absorbent material for absorbing a biological fluid secreted from a person, and an indicator composition comprising a plurality of polymers; a non-cyclic plasticizer; a hydrophobic surfactant; an aromatic acid; an ion-balance reagent and an indicator agent, wherein said composition is devoid of a wetting agent.

In some embodiments there is provided a method for diagnosing amniotic fluid
25 leakage in vaginal secretion comprising:

(a) positioning a secretion-monitoring article to receive a biological fluid secreted from the subject, the article comprising: an absorbent material and an indicator composition comprising a plurality of polymers, an aromatic acid, a non-cyclic plasticizer, a hydrophobic surfactant, an ion-balance reagent and an indicator
30 agent, wherein said composition is devoid of a wetting agent;

- (b) contacting said article with vaginal secretion; and
- (c) viewing the article, wherein a color change indicates the presence of amniotic fluid.

In some embodiments, contacting the article comprises removing release tapes
5 covering adhesive strips, thereby exposing the adhesive strip, and placing the article in the
crotch portion of an undergarment, thereby attaching said article to said crotch portion of
the undergarment.

In some embodiments, viewing the color change comprises detecting dark staining
on the article, wherein the dark staining is indicative of the presence of amniotic fluid.

10 In some embodiments, the dark staining is blue staining. In some embodiments, the
dark staining is green staining.

The compositions described herein may be prepared by methods well known to
those skilled in the art.

In some embodiments, said plurality of polymers comprises a first polymer and a
15 second polymer, the hydrophobicity of said first polymer is higher than the hydrophobicity
of said second polymer.

In some embodiments, said plurality of polymers comprises a first polymer and a
second polymer having similar hydrophobicity. The terms "hydrophobic" and
"hydrophilic" polymers refer on the amount of water vapor absorbed by the polymers at
20 100% relative humidity. Thus, "hydrophobic polymers" typically refer to polymers that
absorb only up to 1 wt. % water at 100% relative humidity, while moderately hydrophilic
polymers absorb 1-10% wt. % water and hydrophilic polymers are capable of absorbing
more than 10 wt. % of water. In some embodiments, the polymers of the invention are
water-insoluble polymers. In some embodiments, each of the polymers comprises
25 cellulose, derivatives thereof or copolymers thereof, including, without limitation,
cellulose acetate, cellulose acetate butyrate, cellulose acetate phthalate, cellophane,
cellulose nitrate, cellulose propionate, cellulose propionate butyrate, cellulose diacetate,
cellulose triacetate, cellulose ethers, and carboxymethyl cellulose, and any copolymers and
any blends of the aforementioned polymers.

30 In some embodiments, each polymer comprises cellulose, derivatives thereof or
copolymers thereof.

In some embodiments, the polymers of the invention are cellulose esters comprising partially acetylated cellulose polymeric chains, including, but not limited to, cellulose acetate in which substantially each one of positions 2, 3 and 6 of the monomeric glucose units of the cellulose polymer, consists of either free hydroxyl or acetate. In some
5 embodiments, the polymers are cellulose esters comprising substantially acetylated cellulose polymeric chains, including, but not limited to, cellulose triacetate. In some embodiments, the polymers are cellulose esters comprising partially esterified cellulose polymeric chains, wherein the esterification comprises partial acetylation and partial butyration, including, but not limited to, cellulose acetate butyrate, in which substantially
10 each one of positions 2, 3 and 6 of the monomeric glucose units of the cellulose polymer consists of either free hydroxyl, acetate or butyrate. In some embodiments, the polymers are cellulose esters comprising partially esterified cellulose polymeric chains, wherein the esterification comprises partial acetylation and partial propionylation, including, but not limited to, cellulose acetate propionate, in which substantially each one of positions 2, 3
15 and 6 of the monomeric glucose units of the cellulose polymer consists of either free hydroxyl, acetate or propionate.

Without wishing to be bound by any theory or mechanism of action, the less hydrophobic polymer, is more hydrophilic thereby improves the penetration of the fluid to be tested, into the panty liner. In addition, the more hydrophobic polymer provides
20 improved adhesion of the indicator composition to the absorbent carrier (e.g. the fabric) thereby minimizing or even avoiding leaching of the formulation components.

In some embodiments, said plurality of polymers comprises at least one polymer selected from the group of cellulose acetate and cellulose acetate butyrate.

In some embodiments, said plurality of polymers comprises a first polymer, such
25 as, cellulose acetate butyrate, and a second polymer, such as, cellulose acetate, the hydrophobicity of said first polymer is higher than the hydrophobicity of said second polymer.

In some embodiments, the plasticizer is a low molecular weight plasticizer. Typically, a low molecular weight plasticizer is a plasticizer having a molecular weight of
30 less than 600 gr/mol. In some embodiments, the plasticizer is having a molecular weight of less than 500 gr/mol. In some embodiments, the plasticizer is having a molecular

weight of less than 400 gr/mol. In some embodiments, the plasticizer is having a molecular weight of less than 300 gr/mol.

Suitable low molecular weight plasticizers include alkyl and aryl phosphates, such as, tributyl phosphate, trioctyl phosphate, tricresyl phosphate, and triphenyl phosphate; 5 alkyl citrate and citrate esters, such as, trimethyl citrate, triethyl citrate, tributyl citrate, acetyl triethyl citrate, and trihexyl citrate; dialkyladipates such as dioctyladipate (DOA; also referred to as bis(2-ethylhexyl)adipate), diethyl adipate, di(2-methylethyl)adipate, and dihexyladipate; dialkyl tartrates such as diethyl tartrate and dibutyl tartrate; dialkylsebacates such as diethyl sebacate, dipropylsebacate and dinonylsebacate; dialkyl 10 succinates such as diethyl succinate and dibutyl succinate; alkyl glycolates, alkyl glycerolates, glycol esters and glycerol esters such as glycerol diacetate, glycerol triacetate (triacetin), glycerol monolactatediacetate, methyl phthalyl ethyl glycolate, butyl phthalyl butyl glycolate, ethylene glycol diacetate, ethylene glycol dibutyrate, triethylene glycol diacetate, triethylene glycol dibutyrate and triethylene glycol dipropionate; acetylated 15 monoglycerides and mixtures thereof.

In some embodiments, the plasticizers comprise low molecular weight plasticizers, including, but not limited to, alkyl citrate and citrate esters, such as, trimethyl citrate, triethyl citrate, tributyl citrate, acetyl triethyl citrate, and trihexyl citrate, acetylated monoglycerides and mixtures thereof.

20 In some embodiments, the non-cyclic plasticizer is a tri-ester plasticizer. In some embodiments, the low molecular weight plasticizers comprise a triester structure. Such plasticizers include, but are not limited to, alkyl citrate and citrate esters, such as, trimethyl citrate, triethyl citrate, tributyl citrate, acetyl triethyl citrate and trihexyl citrate. In some embodiments, the plasticizer is triethyl citrate. In some embodiments, the plasticizer is 25 other than dibutyl phthalate. In some embodiments, the plasticizer is other than Hexamoll Dinch (1,2-cyclohexane dicarboxylic acid diisononyl ester).

In some embodiments, the non-cyclic plasticizer comprises triethyl citrate. In some embodiments, the amount of the surfactant is within the range of 1% to 10% (wt/wt), relative to the total weight of the solid components.

The term “solid components” or “total solids” as used herein refer to the all ingredients within each formulation, in the absence of solvent, in particular, water and/or acetone.

The term “surfactant” as used herein refers to one or more surfactants. In some
5 embodiments, surfactants refer to a plurality of surfactants.

In some embodiments, the surfactant comprises a hydrophilic surfactant. In some
embodiments, the hydrophilic surfactant is a polymer. In some embodiments, the
surfactant comprises a block copolymer surfactant. In some embodiments, the polymeric
surfactant comprises a tri-block copolymer surfactant. In some embodiments, the
10 polymeric surfactant comprises poly(ethylene oxide) and/or copolymers thereof. In some
embodiments, the polymeric surfactant comprises poly(propylene oxide) and/or
copolymers thereof. In some embodiments, the block copolymer surfactant comprises
copolymer of poly(ethylene oxide) and poly(propylene oxide) and/or copolymers thereof.

In some embodiments, the hydrophilic surfactant is a non-ionic surfactant. In some
15 embodiments, the non-ionic surfactant comprise a non-ionic polymeric surfactant. In some
embodiments, the surfactant is selected from the group of poloxamer, alkyl poly(ethylene
oxide), copolymers of poly(ethylene oxide) and poly(propylene oxide), octylglucoside,
decylmaltoside, cetyl alcohol, oleyl alcohol, cocamidemonoethanolamine (cocamide
MEA), cocamidediethanolamine (cocamide DEA), cocamidetriethanolamine (cocamide
20 TEA), polysorbate, such as, Tween® (e.g. Tween 20® or Tween 80®), lecithin and
combinations thereof.

In some embodiments, the non-ionic polymeric surfactant is selected from the
group of poly(ethylene oxide) and copolymers of poly(ethylene oxide), poly(propylene
oxide), polysorbate, such as, Tween® (e.g. Tween 20® or Tween 80®) and combinations
25 thereof.

In some embodiments, the surfactant comprises poloxamer, polysorbate and a
combination thereof.

In some embodiments, the surfactant comprises a poloxamer. In some
embodiments, the surfactant comprises poloxamer 407.

30 In some embodiments, the surfactant comprises a polysorbate. In some
embodiments, the non-ionic surfactant is other than ionic surfactants, such as, aliquot 336.

The term “poloxamer” as used herein refers to nonionic triblock copolymers comprising a central hydrophobic chain of polyoxypropylene (poly(propylene oxide)) and two hydrophilic chains of polyoxyethylene (poly(ethylene oxide)). Poloxamers are commonly termed "P" followed by three digits, the first two times 100 refer to the estimated molecular mass of the polyoxypropylene core, and the last digit times 10 is the percentage polyoxyethylene content. For example, P407 refers to a poloxamer with a polyoxypropylene molecular mass of 4,000 g/mol and 70% polyoxyethylene content.

The term “polysorbate” as used herein generally refers to polyoxyethylene derivative of sorbitan monolaurate. Common commercial brand names for polysorbates include Tween®, such as, Tween 20® or Tween 80®. The number following the polyoxyethylene part refers to the total number of oxyethylene groups found in the molecule. The number following the polysorbate part is related to the type of fatty acid associated with the polyoxyethylene sorbitan part of the molecule. Monolaurate is indicated by 20, monopalmitate is indicated by 40, monostearate by 60 and monooleate by 80.

Without wishing to be bound by any theory or mechanism, the non-ionic polymeric surfactant distributes within the urine and as a result retention time of urine is reduced thereby avoiding false positive results from urine.

In some embodiments, the at least one carboxylic acid comprises at least one unsubstituted or substituted carbocyclic or heterocyclic carboxylic acid. In some embodiments, the at least one carboxylic acid comprises at least one of unsubstituted or substituted aromatic or heteroaromatic carboxylic acid. In some embodiments, the at least one carboxylic acid comprises at least one unsubstituted or substituted aromatic carboxylic acid. In some embodiments, the at least one substituted carboxylic acid is selected from the group consisting of salicylic acid, benzoic acid, phthalic acid, terphthalic acid, isophthalic acid, α -naphthoic acid, β -naphthoic acid, isomers of toluic acid, isomers of halobenzoic acid, such as, but not limited to *p*-fluorobenzoic acid and *m*-chlorobenzoic acid, isomers of nitrobenzoic acids, isomers of trifluoromethylbenzoic acid, isomers of aminobenzoic acid, such as, but not limited to, anthranilic acid, isomers of anisic acid, *m*-hydroxybenzoic acid and *p*-hydroxybenzoic acid. Each possibility represents as separate embodiment of the present invention.

The term “isomer” as used herein refers to a positional isomer, denoting a relative location in which two or more substituents are located on the aromatic ring.

In some embodiments, the acid is an aromatic carboxylic acid. In some embodiments, the aromatic carboxylic acid is in the form of Ar-COOH. In some
 5 embodiments, Ar is selected from the group consisting of aryl and heteroaryl rings, wherein Ar is unsubstituted or substituted. Non limiting examples of substitutions include halogen, haloalkyl, C₁-C₄ alkyl, aryl, heterocyclyl, hydroxy, alkoxy, aryloxy, alkylaryloxy, heteroaryloxy, oxo, cycloalkyl, phenyl, heteroaryl, naphthyl, amino, alkylamino, arylamino, heteroarylamino, dialkylamino, diarylamino, alkylarylamine,
 10 alkylheteroarylamine, arylheteroarylamine, acyl, acyloxy, nitro, carboxy, carbamoyl, carboxamide, cyano, sulfonyl, sulfonylamino, sulfinyl, sulfinylamino, thiol, C₁ to C₁₀ alkylthio, arylthio, or C₁ to C₁₀ alkylsulfonyl groups. Any substituent can be unsubstituted or further substituted with any one of these aforementioned substituents. Each possibility represents as separate embodiment of the present invention.

15 The terms “aryl” and “Ar” as used herein alone or as part of another group are interchangeable and denote an aromatic ring system containing from 6-14 ring carbon atoms. The aryl ring can be a monocyclic, bicyclic, tricyclic and the like. Non-limiting examples of aryl groups are phenyl, naphthyl, including, 1-naphthyl and 2-naphthyl, and the like. The aryl group may be unsubstituted or substituted through available carbon
 20 atoms with one or more groups defined hereinabove for alkyl.

The term “heteroaryl” used herein alone or as part of another group denotes a heteroaromatic system containing at least one heteroatom ring atom selected from nitrogen, sulfur and oxygen. The heteroaryl generally contains 5 or more ring atoms. The heteroaryl group can be monocyclic, bicyclic, tricyclic and the like. Also included in this
 25 expression are the benzoheterocyclic rings and other fused heterocyclic ring systems. If nitrogen is a ring atom, the present invention also contemplates the *N*-oxides of the nitrogen containing heteroaryls. Non-limiting examples of heteroaryls include thienyl, benzothienyl, 1-naphthothienyl, thianthrenyl, furyl, benzofuryl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, isoindolyl, indazolyl,
 30 purinyl, isoquinolyl, quinolyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, carbolinyl, thiazolyl, oxazolyl, isothiazolyl, isoxazolyl and the like. The

heteroaryl group may be substituted through available atoms with one or more groups defined hereinabove for alkyl.

The terms “heterocyclic ring” or “heterocyclyl” used herein alone or as part of another group denote a five-membered to eight-membered rings that have 1 to 4 heteroatoms, such as oxygen, sulfur and/or nitrogen, in particular nitrogen, either alone or in conjunction with sulfur or oxygen ring atoms. These five-membered to eight-membered rings can be saturated, fully unsaturated or partially unsaturated. Preferred heterocyclic rings include piperidinyl, pyrrolidinyl, pyrrolinyl, pyrazolinyl, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, pyranyl, thiopyranyl, piperazinyl, indolinyl, dihydrofuranyl, tetrahydrofuranyl, dihydrothiophenyl, tetrahydrothiophenyl, dihydropyranyl, tetrahydropyranyl, and the like. The heterocyclyl group can be unsubstituted or substituted through available atoms with one or more groups defined hereinabove for alkyl.

The term “halogen” used herein alone or as part of another group includes chloro, fluoro, bromo, and iodo.

In some embodiments, the at least one carboxylic acid comprises at least one carboxylic acid having a pKa smaller than 3. In some embodiments, the at least one carboxylic acid comprises at least one carboxylic acid having a water solubility of less than 100 gr/L. In some embodiments, the at least one carboxylic acid comprises at least one carboxylic acid having a water solubility of less than 10 gr/L. In some embodiments, the at least one carboxylic acid comprises at least one carboxylic acid having a water solubility of less than 5 gr/L.

In some embodiments, the aromatic carboxylic acid is selected from the group consisting of salicylic acid, benzoic acid and the like and combinations thereof. Each possibility represents as separate embodiment of the present invention.

In some embodiments, the aromatic carboxylic acid is salicylic acid.

In some embodiments, the aromatic carboxylic acid is other than non-aromatic carboxylic acids, such as, lactic acid, oxalic acid or citric acid. In some embodiments, the at least one carboxylic acid is other than citric acid.

Without being bound by any theory or mechanism, the aromatic carboxylic acid creates a complex, and may function as a trap, with the ammonium ions within the urine.

This interaction results in reduction of interference from urine, including urine with pH level of 7.0 units, and even higher, and therefore improves indicator specificity.

In some embodiments, the ion-balance reagent is in an amount of about 0.1% to 2%. In some embodiments, the ion-balance reagent is present in an amount of about 0.1% to 1%; and the indicator agent is present in an amount of about 0.2% to 0.7%. In some
5 embodiments, the ion-balance reagent is a quaternary ammonium ion. In some
embodiments, the ion-balance reagent is a quaternary ammonium salt.

In some embodiments, the ion-balance reagent is selected from the group consisting of: di(long-chain alkyl)dimethyl ammonium chloride, *N*-methyl-*N,N*-bis(long-
10 chain alkanoyloxyethyl)-*N*-(2-hydroxyethyl) ammonium methylsulfate,
vinylbenzyltrimethylammonium chloride, methyl trioctyl ammonium chloride
tricaprylmethyl ammonium chloride, tridodecylmethyl ammonium chloride and
cetyltrimethyl ammonium chloride. Each possibility represents as separate embodiment of
the present invention.

15 Any suitable ion-balance reagent may be used, including, tridodecylmethyl
ammonium chloride (TDMAC; CAS 7173-54-8) or cetyltrimethyl ammonium chloride
(CTAC; CAS 112-02-7). In some embodiments, the ion balance reagent is a combination
of ion-balance reagents. In some embodiments, the ion-balance reagent is TDMAC
(Tridodecylmethylammonium chloride).

20 In some embodiments, the indicator is selected from the group of chemical
compounds having negatively charged functional groups. In some embodiments, the
indicator is a weak acid. In some embodiments, the indicator is selected from the group
consisting of nitrazine yellow, thymol blue, bromthymol blue, xylene blue, bromoxylene
blue, phenol red, m-cresol purple, chlorophenol red, bromcresol purple, alizarin, neutral
25 red, and cresol red. Each possibility represents as separate embodiment of the present
invention. A list of other suitable indicators can be found, for example, in U.S. Patent No.
5,897,834. It is clear to one skilled in the art that the indicators specifically mentioned
herein are just examples and any suitable indicators may be used.

In some embodiments, the indicator agent is selected from the group consisting of:
30 cresol red, alizarin, bromocresol purple, chlorophenol red, nitrazine yellow, bromthymol

blue, bromoxylene blue, neutral red, phenol red, thymol blue, xylene blue and m-cresol purple. Each possibility represents as separate embodiment of the present invention.

In some embodiments, the indicator agent is negatively charged.

In some embodiments, the indicator agent is a weak acid.

5 In some embodiments, the indicator agent comprises nitrazine yellow.

In some embodiments, the diagnostic composition is associated with an absorbent material, thereby forming a diagnostic article for absorbing the fluids secreted from a pregnant woman. The diagnostic article can be implemented using many devices and methods. In some embodiments, the article is implemented in a manner that can be easily
10 used by non-skilled personnel, specifically a user. The body of the article comprising the absorbent material can be supplied to the user, for example, in the form of a pad, gauze, a swab, a fiber ball, but most preferably, as a sanitary napkin, diaper, panty shield, and interlabial structure.

In some embodiments, the article further comprises a means for mounting the
15 article to facilitate the collection of the secreted biological fluid. An example of a mounting means includes, but is not limited to, adhesive strips associated with the article. In some embodiments, the article further comprises release tapes, each covering each of the adhesive strips. Covering as used herein also means protecting (e.g. from drying out). Typically, the user removes the release tapes to expose the adhesive strips of the article
20 and places the article in the crotch portion of their undergarment, thereby preventing the article from moving out of position during regular use. Types of adhesive compounds that can be used are well known in the art.

In some embodiments, the diagnostic composition provides a visible indication of amniotic fluid. In some embodiments, the visual indication is stable for about a week.

25 The terms “stable indication” and “irreversible indication” are interchangeably used herein to describe an indication, typically a color indication, that once obtained remains sufficiently unaltered for the time required for clinical examination by a professional. Preferably, the color change is stable for at least 48 hours, more preferably at least 72 hours, and in some embodiments, preferably the color change is stable for about a
30 week.

In some embodiments, the absorbent material is selected from the group consisting of a swab, gauze, panty shield, hygienic napkin, diaper and interlabial absorbent structure. Each possibility represents as separate embodiment of the present invention.

In some embodiments, there is provided a kit comprising the article for detecting
5 amniotic fluid in vaginal secretion, and instructions for use of the article in order to detect amniotic fluid in vaginal secretion.

In some embodiments, the kit further comprises a color scale presenting the color expected to appear upon contact of the article with amniotic fluid and the color expected to appear upon contact of the article with urine.

10 In some embodiments, the kit comprises a plurality of article for detecting amniotic fluid in vaginal secretion.

In some embodiments, each article within the kit is separately wrapped.

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These
15 Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitation.

20

EXAMPLES

EXAMPLE 1: Clinical Study

Samples of urine and amniotic fluid were collected from 83 pregnant women, according to the following distribution: urine and amniotic fluid were collected from 8
25 women, urine only was collected from 63 women, and amniotic fluid was collected from 12 women.

The inclusion criteria for subject participation in the study are pregnant women:

- (i). Pregnant women 18 years to 50 years of age.
- (ii). Subject is ready to sign the informed consent form.
- 30 (iii). Subject arrives at delivery room undergo amniotomy.

The exclusion criteria for subject participation in this study are:

- (iv). Subject suffers from bleeding.
- (v). Subject is unable or unwilling to cooperate with study procedures.
- (vi). Subject is currently participating in another clinical study.

5

Subjects arriving at the delivery room were requested to collect a random urine sample in a clean container. Amniotic fluid samples were collected, by the clinician, during the amniotomy procedure. Each fresh urine and amniotic fluid sample in the study was tested using the test devices (panty liners) each comprising candidate formulation or a reference formulation, as follows:

10

1. Dipstick for pH and other urine parameters (using Combur-Test® strips, Roche Diagnostics).
2. 100µl of amniotic fluid was dropped onto each device candidate and reference devices (AL-SENSE and AmniScreen).
- 15 3. 400µl of urine was dropped onto each device candidate (a panty liner containing one of the eight candidate formulations) and each reference device (a panty liner containing the AmniScreen or Al-Sense formulations).
4. The study coordinator stirred each panty liner in a way that will mimic the "real life" usage.
- 20 5. For the amniotic fluid and urine samples, the coordinator recorded the result status (positive or negative) immediately (0-2 min), and after 5 min, 10 min, 15 min, 20 min, 30 min, 60 min and 120 min, based on the color observed on the tested pads. A positive result is concluded when the yellow panty liner show a blue, green stain of any size or shape, even if as a light color. A negative result is concluded when the panty
- 25 liner color remained yellow.

Eight types of device candidates (Table 1) and two reference devices, AL-SENSE and AmniScreen (Table 2), were used. The ingredients in each candidate and reference formulations were dissolved in 50 ml acetone and 2 ml water.

30

Table 1: Formulations of candidate devices

Agent	ALM1/ALM1G [§]		ALM2/ALM2G [§]		ALM3/ALM3G [§]		ALM4/ALM4G [§]	
	Wt. (g)	wt/wt(%)	Wt. (g)	wt/wt(%)	Wt. (g)	wt/wt(%)	Wt. (g)	wt/wt(%)
Cellulose acetate	2	44.6	2	45.7	2	45.7	2	45.8
Cellulose acetate-butyrates	0.8	17.9	0.8	18.3	0.85	19.4	0.8	18.3
Triethyl citrate	1	22.3	1	22.8	1	22.8	1	22.9
TDMAC	0.053	1.2	0.053	1.2	0.053	1.2	0.046	1.1
Poloxamer	0.3	6.7						
Salicylic acid	0.3	6.7			0.25	5.71		
Benzoic acid			0.3	6.7			0.3	6.86
Tween 80			0.2	6.7	0.2	4.57	0.2	4.58
Nitrazine yellow	0.024	0.54	0.024	0.55	0.024	0.55	0.024	0.55
Total	4.48	100	4.38	100	4.38	100	4.37	100

[§]G = contains glue between the layers as a part of the manufacturing process.

5 Table 2: AL-SENSE and AmniScreen Compositions

Ingredient	AL-Sense	AmniScreen	Units
Cellulose Acetate	1.6	1.5	Gr
Cellulose acetate butyrate	0.8		Gr
Dibutyl phthalate		1.07	MI
Hexamoll Dinch	0.8		MI
TDMAC	0.0453		Gr
Aliquat 336		0.23	MI
2-ethoxyethanol		1.5	MI
Citric acid	0.0112		Gr
Nitrazine Yellow	0.0091	0.024	Gr

The values of the AmniScreen reference device in terms of sensitivity and specificity were as follows: efficacy of 84% specificity and 96% sensitivity. AmniScreen provides final reading after 30 minutes. Significance level of findings was determined to be equal to, or lower than, 5%. All statistical analyses were performed using SAS v9.3 (SAS®, SAS Institute Cary, NC USA) software.

Results (stains) were detected at different time points. While accurate detection with 100% sensitivity was obtained by articles comprising the indicator composition of the invention within the first 2 minutes after contact with amniotic fluid (Table 3), it was required to wait 30 minutes and 10 minutes when using AmniScreen and AL-SENSE,

respectively, in order to avoid false positive readings (i.e. staining by urine). This is because in AmniScreen and AL-SENSE a stain is obtained from non-specific staining (by urine), which requires some time (30 minutes in AmniScreen and 10 minutes in AL-SENSE) to fade.

5 The results from the different tests, based on the pads color, were designated "Positive", "Negative". The term "Positive" means a blue stain, which appeared upon the yellow background of the panty liner device. The term "Negative" means that a stain did not appear, or appeared and faded back to yellow during detection time. A stain formed in a urine test was considered as false positive and when a stain does not appear in a urine
10 test it was true negative. Similarly, when a stain did not appear in a test of amniotic fluid it was referred to as false negative.

Tables 3-4 and Figure 1 present the detected results read from the candidate devices following exposure to amniotic fluid (Table 3) or to urine (Table 4). Efficacy evaluation was obtained by comparing the sensitivity, i.e. the proportion of "Positive" readings upon
15 exposure of the pads to amniotic fluid, and the specificity, i.e. the proportion of "Negative" upon exposure of the pads to urine, from the various devices (pads). The resulting sensitivities and specificities are presented in Tables 3 and 4 along with the respective two-sided 90% exact Confidence Intervals (CI) compared with McNemar's test. For each candidate device agreement between the candidate device and the AmniScreen
20 was calculated together with 95% exact confidence intervals. Agreements in amniotic fluid, in urine and altogether were evaluated as the percentage of equal diagnoses (positive-positive negative-negative) between pairs of devices. The devices diagnosis agreement levels were cross-tabulated against those of the AmniScreen overall agreement levels and compared using McNemar's test. As shown in Table 3 all candidate devices had
25 a sensitivity of 100%, i.e. all candidates exhibited a positive reading (blue stain) following exposure to amniotic fluid. Furthermore, in all candidate pads the color change (blue stain formation on a yellow background) occurred immediately or by no longer than two minutes after contacting the test sample (i.e. 0-2 min) and remained stable for the duration of the two-hour follow-up. Regarding specificity, candidate pads ALM1 and ALM1G
30 demonstrated superiority over the remaining candidate pads (Table 4 and Figure 1). These two candidates exhibited 84% specificity at 5 minutes and even better specificity values at 10 minutes with no fluctuations: Specificity of the ALM1 (and ALM1G) candidate at 5

minutes was 84.5% (60/71) and at 10 minutes 87.2% (62/71). Thus, ALM1 and ALM1G meet the goal of good accuracy in a short time span.

Tables 5 and 6 present the sensitivity and specificity, respectively, of five selected candidate pads, at 5 and 10 minutes along with the results of AL-SENSE and AmniScreen with lower two-sided 90% exact confidence bounds, based on the values presented in Tables 3 and 4. According to Table 5, the sensitivity of ALM1 at 5 and 10 minutes is 100% (namely, all amniotic fluid samples that were contacted with ALM1, resulted in a positive detection – blue stain). In terms of performance, it may be stated that for this study, the value of 86.09% has 95% confidence. Stated otherwise, the value of 86.09% refers to the assertion that may be made given the sample size and the sensitivity data. The same applies to AmniScreen. AL-SENSE has a 90% sensitivity, which is lower than that of ALM1 and AmniScreen.

Table 3: Readings of candidate devices upon exposure to amniotic fluid

Candidate device	0-2m		5m		10m		15m		20m		30m		60m		120m	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
ALM1																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM2																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM3																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM4																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM1G																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM2G																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM3G																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0
ALM4G																
Positive	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0	20	100.0

‘%’ -% sensitivity; ‘N’ - the number of samples tested; ‘m’ – minutes.

According to Table 6, specificity of ALM1 at 5 minutes (84.51%) is better than that of AmniScreen (77.46%; $p=0.0588$) with a significant superiority at 10 minutes (87.32%) relative to AmniScreen ($p=0.0082$) and AL-SENSE (74.65%, $p=0.0067$). Specificities of AmniScreen and AL-SENSE were not significantly different ($p=0.5637$).

Specificity, as used herein refers, to correct negative reading, namely, non-staining of a composition upon contact thereof with a urine sample.

The clear and sharp detection with ALM1 is further demonstrated in Figures 3A and 3B. Figure 3A presents the dark stain obtained after contacting the composition of ALM1 absorbed into a panty shield with amniotic fluid (the dark stain is surrounded by a dashed line and an arrow is pointing at the stain). The dark stain presented in Figure 3A is clearer and much more pronounced compared to the stains observed after contacting the reference compositions, namely, AL-SENSE and AmniScreen, with amniotic fluid. Figure 3B presents a corresponding composition of ALM1 absorbed into a panty shield after being in contact with urine. As can be seen, no staining is observed indicating the absence of amniotic fluid.

10

Table 4: Readings of the candidate devices when exposed to urine, over time

	0-2m		5m		10m		15m		20m		30m		60m		120m	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
ALM1																
Negative	58	81.7	60	84.5	62	87.3	62	87.3	62	87.3	64	90.1	64	90.1	64	90.1
Positive	13	18.3	11	15.5	9	12.7	9	12.7	9	12.7	7	9.9	7	9.9	7	9.9
ALM2																
Negative	53	74.6	57	80.3	60	84.5	60	84.5	61	85.9	62	87.3	62	87.3	62	87.3
Positive	18	25.4	14	19.7	11	15.5	11	15.5	10	14.1	9	12.7	9	12.7	9	12.7
ALM3																
Negative	52	73.2	56	78.9	60	84.5	61	85.9	61	85.9	64	90.1	64	90.1	64	90.1
Positive	19	26.8	15	21.1	11	15.5	10	14.1	10	14.1	7	9.9	7	9.9	7	9.9
ALM4																
Negative	57	80.3	60	84.5	60	84.5	61	85.9	60	84.5	61	85.9	61	85.9	61	85.9
Positive	14	19.7	11	15.5	11	15.5	10	14.1	11	15.5	10	14.1	10	14.1	10	14.1
ALM1G																
Negative	58	81.7	60	84.5	62	87.3	63	88.7	63	88.7	65	91.5	65	91.5	65	91.5
Positive	13	18.3	11	15.5	9	12.7	8	11.3	8	11.3	6	8.5	6	8.5	6	8.5
ALM2G																
Negative	56	78.9	59	83.1	61	85.9	61	85.9	61	85.9	63	88.7	63	88.7	63	88.7
Positive	15	21.1	12	16.9	10	14.1	10	14.1	10	14.1	8	11.3	8	11.3	8	11.3
ALM3G																
Negative	56	78.9	59	83.1	61	85.9	62	87.3	63	88.7	64	90.1	64	90.1	64	90.1
Positive	15	21.1	12	16.9	10	14.1	9	12.7	8	11.3	7	9.9	7	9.9	7	9.9
ALM4G																
Negative	56	78.9	58	81.7	60	84.5	60	84.5	61	85.9	63	88.7	63	88.7	63	88.7
Positive	15	21.1	13	18.3	11	15.5	11	15.5	10	14.1	8	11.3	8	11.3	8	11.3

‘%’ - % sensitivity; ‘N’ - the number of samples tested; ‘m’ – minutes.

Table 5: Sensitivity of the candidate and reference pads

Device	n/N	Sensitivity (%)	Lower 90% CL
ALM1 5 min	20/20	100.0%	86.09%
ALM1 10 min	20/20	100.0%	86.09%
AL-Sense 10 min	18/20	90.0%	71.74%
AmniScreen 30 min	20/20	100.0%	86.09%

Table 6: Specificity of the candidate and reference pads

Device	n/N	Specificity (%)	Lower 90% CL	P-Value vs. AmniScreen	P-Value vs. AL-SENSE
ALM1 5 min	60/71	84.51%	75.66%	0.0588	0.0196
ALM1 10 min	62/71	87.32%	78.92%	0.0082	0.0067
AL-SENSE 10 min	53/71	74.65%	64.76%	0.5637	NR
AmniScreen 30 min	55/71	77.46%	67.81%	NR	0.5637

5 The clear and sharp detection with ALM1 is further demonstrated in Figures 3A and 3B. Figure 3A presents the dark stain obtained immediately after contacting the composition of ALM1 absorbed into a panty shield with amniotic fluid (the dark stain is surrounded by a dashed line and an arrow is pointing at the stain). The dark stain presented in Figure 3A is clearer and much more pronounced compared to the stains observed after

10 contacting the reference compositions, namely, AL-SENSE and AmniScreen, with amniotic fluid. Figure 3B presents a corresponding composition of ALM1 absorbed into a panty shield after being in contact with urine. As can be seen, no staining is observed indicating the absence of amniotic fluid.

15 Table 7 presents cross-tabulation readings of ALM1 and AL-SENSE along with the reading of AmniScreen - all readings were obtained on the same sample.

20 Table 8 displays the total percent agreement level of the candidate and the reference (AL-SENSE and AmniScreen) devices on positive and negative cases and overall with two-sided 95% exact confidence intervals for each. These values represent are the percent of cases in which the devices gave the same reading in amniotic fluid (positive cases; N=20), in urine (negative cases; N=71) and overall (together in both urine and amniotic fluid; N=91). As can be seen in the positive cases, i.e. amniotic fluid, ALM1 readings at 5 and 10 minutes correspond to AmniScreen readings. Thus, the agreement between ALM1

and AmniScreen in diagnosis of amniotic fluid is 100% for both time points, where agreement between ALM1 and AL-SENSE was only 90% (Table 5). In urine, ALM1 at 5 minutes agrees with AmniScreen in 98.2% of the cases and at 10 minutes in all (100%) cases where the AmniScreen provided a negative reading. AmniScreen demonstrated six false positive readings at 5 minutes and seven false positive readings at 10 minutes while ALM1 correctly read negative reading at 5 minutes and 10 minutes (Table 6).

Table 7: Cross tabulation of candidate, AL-Sense versus AmniScreen

Test Sample		AmniScreen			
		Negative		Positive	
		N	%	N	%
Amniotic Fluid	<u>ALM1 5m</u>				
	Positive	.	.	20	100.0
	<u>ALM1 10m</u>				
	Positive	.	.	20	100.0
Urine	<u>AL-SENSE</u>				
	Negative	.	.	2	10.0
	Positive	.	.	18	90.0
	<u>ALM1 5m</u>				
	Negative	54	98.2	6	37.5
	Positive	1	1.8	10	62.5
	<u>ALM1 10m</u>				
	Negative	55	100.0	7	43.8
	Positive	.	.	9	56.3
	<u>AL-SENSE</u>				
Overall	Negative	48	87.3	5	31.3
	Positive	7	12.7	11	68.8
	<u>ALM1 5m</u>				
	Negative	54	98.2	6	16.7
	Positive	1	1.8	30	83.3
	<u>ALM1 10m</u>				
	Negative	55	100.0	7	19.4
	Positive	.	.	29	80.6
	<u>AL-SENSE</u>				
	Negative	48	87.3	7	19.4
	Positive	7	12.7	29	80.6

In general, there was a disagreement between ALM1 at 5 min and 10min and AmniScreen in seven cases out of which ALM1 gave the correct reading in six (6) cases at 5 min and in all seven (7) cases at the 10 min evaluation point. The differences were not statistically significant at 5 minutes ($p=0.0588$, Table 8) but were found statistically significant at 10 min ($p=0.0082$, Table 8). Stated otherwise, as evident from Table 7, the

aforementioned disagreement is only on the negative cases (i.e., urine) where at 5 min ALM1 gave the correct reading in 6 of those 7 discrepant cases and in all 7 cases at the 10 min evaluation point. Disagreement between AL-SENSE and AmniScreen was greater - 14 cases, of which the AmniScreen was correct in 9 of the cases. Despite the better accuracy, AmniScreen and AL-Sense were found to be similar ($p=1.00$, Table 8) with respect to percent of positive and negative readings yet less effective than ALM1.

Table 8: Agreement of ALM1 and AL-SENSE with AmniScreen

	Device	n/N	Agreement Percent	Lower 95% CL	Upper 95% CL	P-Value McNemar's Test
Amniotic Fluid	ALM1 5 min	20/20	100.0%	83.16%	100.0%	-
	ALM1 10 min	20/20	100.0%	83.16%	100.0%	-
	AL-Sense	18/20	90.0%	68.30%	99.98%	-
Urine	ALM1 5 min	64/71	90.14%	80.74%	95.94%	-
	ALM1 10 min	64/71	90.14%	80.74%	95.94%	-
	AL-Sense	59/71	83.10%	72.34%	90.95%	-
Overall	ALM1 5 min	84/91	92.31%	84.79%	96.85%	0.0588
	ALM1 10 min	84/91	92.31%	84.79%	96.85%	0.0082
	AL-Sense	77/91	84.62%	75.54%	91.33%	1.000

10 **EXAMPLE 2: Clinical Study with Articles Lacking Surfactants and Acids**

Two formulations, 3Mar1 and 3Mar2, were prepared, containing all of the ingredients of the ALM1 formulation aside from poloxamer 407 or salicylic acid (Table 9).

Table 9: Formulations Lacking Surfactants or Acids

Ingredient	ALM1	3Mar1	3Mar2
Cellulose Acetate	2.00	2.50	2.50
Cellulose acetate butyrate	0.80	1.00	1.00
Triethyl citrate	1.00	1.00	1.00
TDMAC	0.053	0.053	0.053
Poloxamer 407	0.30	0.30	
Salicylic acid	0.30		0.30
Nitrazine Yellow	0.024	0.024	0.024

Urine samples (400 µl) having pH 7.0 were dripped on articles containing the 3Mar1 and 3Mar2 formulations and on an article containing the ALM1 formulation.

5 As shown in Figure 2, following contact with urine each of the articles containing the 3Mar1 or the 3Mar2 formulations produced a false positive result, namely, a dark distinctive (blue) stain formed at the center of each of the articles. However, the ALM1 article provided a true negative result as it did not produce any stain upon contacting the urine sample.

10 The results indicate that surfactant(s) and acid(s) are required for obtaining a reliable detection.

EXAMPLE 3: Feasibility of Wearing Study

15 A primary efficacy analysis with an article which includes the diagnostic composition ALM1 (also termed hereinafter "AL SENSE Blue") was conducted on a total of 43 women who were compliant with the study protocol and had valid data for the analysis. The analysis consists of a comparison between the patient's reading of the AL-SENSE Blue result and the clinical diagnosis.

20 Table 10 presents the subject's reading of the AL-SENSE Blue device versus the clinical diagnosis (sterile speculum Pooling test, if result was negative, then a microscopic Ferning test and pH test were performed).

Table 10: Subject's reading vs. clinical diagnosis

Clinical Diagnosis Frequency (N)	AL-SENSE Blue Subject's reading		
	Positive	Negative	Total
Negative	1	16	17
Positive	25	1	26
Total	26	17	43

The sensitivity observed in this study is 96.15% (25/26) with a 95% exact binomial confidence interval of 80.36% - 99.90% and the specificity is 94.12% (16/17) with a 95% exact binomial confidence interval of 71.31% - 99.85%. Since the Success Criteria were 91% for the sensitivity and 73% for the specificity, it is concluded that the study was successful.

In a Secondary Efficacy study the women (all participants) were requested to complete a questionnaire. The first three questions were rated on a scale of 1 through 5, where 1 represents a poor result and 5 a very satisfactory result. The last two questions relate to patient's comprehension of the results reading. Tables 11-14 present in a descriptive manner the distributions of response gathered.

Table 11: Responses to Question 1

Is the Blue-Green Stain on the Indicator Strip Clear to Distinguish? (N=35)		
Response	Frequency	%
Slightly Clear	1	2.86
Somewhat Clear	10	28.57
Clear	17	48.57
Very Clear	7	20

Table 12: Responses to Question 2:

Is the pad comfortable for use? (N=40)		
Response	Frequency	%
Quite Comfortable	24	60
Very Comfortable	16	40

Table 13: Responses to Question 3

Did you experience any discomfort related to use of the pad? (N=41)		
Response	Frequency	%
Indifferent	17	41.46
I was quite comfortable with the AL-SENSE Blue	11	26.83
I experienced no discomfort related to the AL-SENSE Blue	13	31.71

5 Table 14: Responses to Question 4

According to the color of the used indicator strip and to the instructions leaflet, do you understand the cause of your wetness sensation? (N=41)		
Response	Frequency	%
Amniotic fluid leakage	23	53.66
Urine leakage	0	0
Nothing	7	17.07
I don't know	11	29.27

Of the 35 women that responded to question 1, 97.14% (34/35) responded that the Blue or green stain on the yellow strip was somewhat clear, clear or very clear to distinguish.

10 Of the 40 women that responded to question 2, 100% (40/40) responded the pad was quite or very comfortable for use.

Of the 41 women that responded to question 3, no one claimed that they had experienced even slight discomfort relating to the use of the AL-SENSE Blue.

Of the 41 women that responded to question 4, 24 marked their reading as positive. 95.83% of these women (23/24) stated that from the strip they had understood that the cause of their wetness was due to amniotic fluid leakage, all of them actually marked their reading as positive.

5 'Time to Reading' of the tested article was improved relative to the reference articles. Specifically, reading of the liner color was detected and recorded at 2, 5, 10, 15, 20 and 30 minutes (Table 15). In 100% (43/43) of the cases the final reading was already present from 2 minutes after exposition. In contrast, the Time to Reading of the reference article AmniScreen is 30 minutes while the Time to Reading of the reference article AL-SENSE is at least 10
10 minutes.

Table 15: Listing of the readings over time

Subject Number	Patient result 0-2 min	Patient result 5 min	patient result 10 min	Patient result 15 min	Patient result 20 min	Patient result 30 min
9200	Positive	Positive	Positive	Positive	Positive	Positive
9201	Positive	Positive	Positive	Positive	Positive	Positive
9202	Positive	Positive	Positive	Positive	Positive	Positive
9203	Negative	Negative	Negative	Negative	Negative	Negative
9204	Positive	Positive	Positive	Positive	Positive	Positive
9205	Positive	Positive	Positive	Positive	Positive	Positive
9206	Negative	Negative	Negative	Negative	Negative	Negative
9207	Positive	Positive	Positive	Positive	Positive	Positive
9208	Positive	Positive	Positive	Positive	Positive	Positive
9209	Positive	Positive	Positive	Positive	Positive	Positive
9210	Positive	Positive	Positive	Positive	Positive	Positive
9212	Negative	Negative	Negative	Negative	Negative	Negative
9213	Negative	Negative	Negative	Negative	Negative	Negative
9214	Negative	Negative	Negative	Negative	Negative	Negative
9215	Negative	Negative	Negative	Negative	Negative	Negative
9217	Positive	Positive	Positive	Positive	Positive	Positive
9218	Positive	Positive	Positive	Positive	Positive	Positive
9219	Positive	Positive	Positive	Positive	Positive	Positive
9222	Positive	Positive	Positive	Positive	Positive	Positive
9223	Positive	Positive	Positive	Positive	Positive	Positive
9224	Positive	Positive	Positive	Positive	Positive	Positive
9226	Negative	Negative	Negative	Negative	Negative	Negative
9227	Positive	Positive	Positive	Positive	Positive	Positive
9228	Negative	Negative	Negative	Negative	Negative	Negative
9229	Negative	Negative	Negative	Negative	Negative	Negative
9230	Positive	Positive	Positive	Positive	Positive	Positive

9231	Positive	Positive	Positive	Positive	Positive	Positive
9232	Positive	Positive	Positive	Positive	Positive	Positive
9233	Negative	Negative	Negative	Negative	Negative	Negative
9234	Negative	Negative	Negative	Negative	Negative	Negative
9237	Positive	Positive	Positive	Positive	Positive	Positive
9238	Negative	Negative	Negative	Negative	Negative	Negative
9239	Negative	Negative	Negative	Negative	Negative	Negative
9240	Negative	Negative	Negative	Negative	Negative	Negative
9241	Negative	Negative	Negative	Negative	Negative	Negative
9242	Positive	Positive	Positive	Positive	Positive	Positive
9243	Negative	Negative	Negative	Negative	Negative	Negative
9244	Negative	Negative	Negative	Negative	Negative	Negative
9246	Positive	Positive	Positive	Positive	Positive	Positive
9247	Positive	Positive	Positive	Positive	Positive	Positive
9248	Positive	Positive	Positive	Positive	Positive	Positive
7189	Positive	Positive	Positive	Positive	Positive	Positive
7190	Positive	Positive	Positive	Positive	Positive	Positive

In summary of the primary and secondary studies, the AL-SENSE Blue subject reading sensitivity is 96.15%, which is very good for a home use test. Furthermore, the performance altogether (sensitivity, specificity, time of reading etc.) of AL-SENSE Blue is better than that of the reference articles AL-SENSE and AmniScreen.

5

EXAMPLE 4: Article Preparation

Article preparation initially involved preparing each of the candidate and reference (including 3Mar1 and 3Mar2) formulations in the form of solutions. For this purpose, the solid part of each formulation, excluding Nitrazine yellow, was dissolved in 50 ml acetone. Nitrazine yellow was dissolved in 2 ml water and then added to the acetone solution.

The resulting solutions were sprayed on a substrate material. As a result, the substrate material acquired a faint yellow color, which gave a suitable background for the detection of a color change due to interaction with vaginal fluids, particularly, for the detection of a blue stain as a proper result of interaction with amniotic fluid. Each substrate was then integrated with a pre-formed panty liner, thereby obtaining a distinct diagnostic (candidate or reference) panty liner.

The foregoing description of the specific embodiments will so fully reveal the general nature of the invention that others can, by applying current knowledge, readily modify and/or adapt for various applications such specific embodiments without undue experimentation and without departing from the generic concept, and, therefore, such
5 adaptations and modifications should and are intended to be comprehended within the meaning and range of equivalents of the disclosed embodiments. It is to be understood that the phraseology or terminology employed herein is for the purpose of description and not of limitation. The means, materials, and steps for carrying out various disclosed functions may take a variety of alternative forms without departing from the invention.

WE CLAIM:

1. An indicator composition for detecting amniotic fluid in vaginal secretion comprising:

a plurality of polymers;

a non-cyclic plasticizer;

a hydrophilic surfactant;

an aromatic carboxylic acid;

an ion-balance reagent and

an indicator agent,

wherein said composition is devoid of a wetting agent.
2. The indicator composition of claim 1, wherein said aromatic carboxylic acid comprises salicylic acid, benzoic acid or a combination thereof.
3. The indicator composition of claim 1, wherein said aromatic carboxylic acid is salicylic acid.
4. The indicator composition of claim 1, wherein said hydrophilic surfactant is a non-ionic surfactant.
5. The indicator composition of claim 1, wherein said hydrophilic surfactant is a polymer.
6. The indicator composition of claim 5, wherein said hydrophilic surfactant comprises a block copolymer surfactant.
7. The indicator composition of claim 1, wherein the indicator agent is negatively charged.
8. The indicator composition of claim 1, wherein the ion-balance reagent is a quaternary ammonium salt.

9. The indicator composition of claim 1, wherein the non-cyclic plasticizer is a tri-ester plasticizer.
10. An article for detecting amniotic fluid in a biological fluid secreted from a person, comprising:

an absorbent material for absorbing the biological fluid secreted from the person, and

the indicator composition of any one of claims 1-9.
11. The article of claim 10, wherein the absorbent material is selected from the group consisting of a swab, gauze, panty shield, hygienic napkin, diaper and interlabial absorbent structure.
12. A method for diagnosing amniotic fluid leakage in a biological fluid secreted from a person, comprising:

positioning a secretion-monitoring article to receive the biological fluid secreted from the person, the article comprising the indicator composition of claim 1;

contacting said article with vaginal secretion; and

viewing the article, wherein a color change indicates the presence of amniotic fluid.
13. The method of claim 12, wherein said color change is a dark stain relative to background of the secretion monitoring article.
14. The method of claim 13, wherein the dark stain is a dark blue stain or a dark green stain.
15. The method of claim 14, wherein said viewing is performed within less than 10 minutes after said contacting.
16. The method of claim 15, wherein said viewing is performed within less than 5 minutes after said contacting.

17. A kit comprising the article of claim 10 for detecting the presence of the amniotic fluid in a biological fluid secreted from a person, and instructions for use.
18. The kit of claim 17, further comprises a color scale presenting the color expected to appear upon contact of the article with the amniotic fluid.
19. The indicator composition of claim 1, wherein the plurality of polymers are cellulose acetate and cellulose acetate butyrate;

the non-cyclic plasticizer is triethyl citrate;

the hydrophilic surfactant is a poloxamer;

the aromatic carboxylic acid is salicylic acid;

the ion-balance reagent is tridodecylmethyl ammonium chloride; and

the indicator agent is nitrazine yellow.
20. The indicator composition of claim 19, wherein the indicator composition comprises 44.6 wt% cellulose acetate, 17.9 wt% cellulose acetate butyrate, 22.3 wt% triethyl citrate, 6.7 wt% poloxamer, 6.7 wt% salicylic acid, 1.2 wt% tridodecylmethyl ammonium chloride, and 0.54 wt% nitrazine yellow.
21. An indicator composition for detecting amniotic fluid in vaginal secretion comprising:

at least one polymer selected from cellulose, a cellulose ester, and a copolymer thereof;

a plasticizer, wherein the plasticizer has a molecular weight less than or equal to 600 g/mol;

a surfactant, wherein the surfactant is a poloxamer;

an acid represented by a structural formula that includes a ring moiety selected from an unsubstituted or substituted C₆-C₁₄ aromatic ring moiety, an unsubstituted or substituted C₃-C₈ carbocyclic ring moiety, an unsubstituted or substituted 5- to 8-membered

heterocyclic ring moiety having 1 to 4 heteroatoms selected from nitrogen, sulfur and oxygen, or an unsubstituted or substituted 5- to 14-membered heteroaromatic ring moiety having 1 to 4 heteroatoms selected from nitrogen, sulfur, and oxygen;

an ion-balance reagent, wherein the ion-balance reagent is a quaternary ammonium salt;
and

an indicator agent.

22. The indicator composition of claim 21, wherein the ring moiety is an unsubstituted C₆-C₁₄ aromatic ring moiety.
23. The indicator composition of claim 21, wherein the ring moiety is a substituted C₆-C₁₄ aromatic ring moiety.
24. The indicator composition of claim 21, wherein the acid is selected from salicylic acid, benzoic acid and combinations thereof.
25. The indicator composition of claim 21, wherein the acid is salicylic acid.
26. The indicator composition of any one of claims 21-25, wherein the surfactant comprises poloxamer 407.
27. The indicator composition of any one of claims 21-26, wherein the molecular weight of the plasticizer is less than 300 g/mol.
28. The indicator composition of any one of claims 21-27, wherein the plasticizer is triethyl citrate.
29. The indicator composition of any one of claims 21-28, comprising a plurality of polymers, wherein each polymer of the plurality of polymers is independently a cellulose ester.
30. The indicator composition of any one of claims 21-29, wherein the polymer is independently selected from cellulose acetate, cellulose acetate butyrate, cellophane,

- cellulose nitrate, cellulose propionate, cellulose propionate butyrate, cellulose diacetate, cellulose triacetate, cellulose ethers, carboxymethyl cellulose, and any copolymers thereof.
31. The indicator composition of any one of claims 21-30, wherein the polymer is cellulose acetate butyrate.
 32. The indicator composition of any one of claims 21-31, wherein the ion-balance reagent is selected from di(long-chain alkyl)dimethyl ammonium chloride, N-methyl-N,N-bis(long-chain alkanoyloxyethyl)-N-(2-hydroxyethyl) ammonium methylsulfate, vinylbenzyltrimethylammonium chloride, methyl trioctyl ammonium chloride, tricaprylmethyl ammonium chloride, tridodecylmethyl ammonium chloride and cetyltrimethyl ammonium chloride.
 33. The indicator composition of any one of claims 21-32, wherein the indicator agent is negatively charged.
 34. The indicator composition of any one of claims 21-32, wherein the indicator agent is a weak acid.
 35. The indicator composition of any one of claims 21-32, wherein the indicator agent is selected from cresol red, alizarin, bromocresol purple, chlorophenol red, nitrazine yellow, bromothymol blue, bromoxylene blue, neutral red, phenol red, thymol blue, xylenol blue and m-cresol purple.
 36. An article for detecting amniotic fluid in vaginal secretion comprising an absorbent material associated with the indicator composition of any one of claims 1-9 and 19-35.
 37. A method for detecting amniotic fluid leakage in vaginal secretion comprising:
 - a. contacting the indicator composition of any one of claims 21-35 or the article of claim 36 with a biological fluid secreted from a person; and
 - b. viewing the indicator composition or the article, wherein a color change indicates the presence of amniotic fluid.

1/3




Fig. 1

2/3




Fig. 2

3/3




Fig. 3A




Fig. 3B

