




(86) Date de dépôt PCT/PCT Filing Date: 1992/01/15	(51) Cl.Int. ⁵ /Int.Cl. ⁵ A61K 31/19, A61K 31/375, A61K 31/355, A61K 31/20
(87) Date publication PCT/PCT Publication Date: 1992/09/17	(72) Inventeur/Inventor: Martin, Alain, US
(45) Date de délivrance/Issue Date: 2002/07/30	(73) Propriétaire/Owner: WARNER-LAMBERT COMPANY, US
(85) Entrée phase nationale/National Entry: 1993/08/19	(74) Agent: MACRAE & CO.
(86) N° demande PCT/PCT Application No.: US 1992/000249	
(87) N° publication PCT/PCT Publication No.: 1992/015292	
(30) Priorité/Priority: 1991/03/01 (663,500) US	

- (54) Titre : COMPOSITIONS THERAPEUTIQUES DE PROTECTION ET DE REANIMATION DE CELLULES DE MAMMIFERES ET METHODE POUR LEUR PREPARATION ET UTILISATION
- (54) Title: THERAPEUTIC COMPOSITIONS TO PROTECT AND RESUSCITATE MAMMALIAN CELLS AND METHODS FOR PREPARING AND USING SAME

(57) **Abrégé/Abstract:**

The present invention pertains to therapeutic compositions for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells. In one embodiment, the therapeutic composition comprises (a) pyruvate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. In another embodiment, the therapeutic composition comprises (a) pyruvate, (b) lactate, and (c) a mixture of saturated and unsaturated fatty acids. In yet another embodiment, the therapeutic composition comprises (a) an antioxidant, and (b) a mixture of saturated and unsaturated fatty acids. In still yet another embodiment, the therapeutic composition comprises (a) lactate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. The therapeutic compositions may be utilized in a wide variety of topical and ingestible pharmaceutical products. This invention also relates to methods for preparing and using the therapeutic compositions and the topical and ingestible pharmaceutical products in which the therapeutic compositions may be used.





INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/20, 31/375, 31/355 // (A61K 31/20, 31:19, 31:07) (A61K 31/20, 31:19, 31:015) (A61K 31/475, 31:20, 31:19) (A61K 31/355, 31:20, 31:19)	A1	(11) International Publication Number: 2104461 (43) International Publication Date: 17 September 1992 (17.09.92)
(21) International Application Number: PCT/US92/00249 (22) International Filing Date: 15 January 1992 (15.01.92) (30) Priority data: 663,500 1 March 1991 (01.03.91) US (71) Applicant: WARNER-LAMBERT COMPANY [US/US]; 201 Tabor Road, Morris Plains, NJ 07950 (US). (72) Inventor: MARTIN, Alain ; 31 Country Club Drive, Ringoes, NJ 08851 (US). (74) Agents: BULLITT, Richard, S. et al.; Warner-Lambert Company, 201 Tabor Road, Morris Plains, NJ 07950 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), MC (European patent), NL (European patent), SE (European patent). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: THERAPEUTIC COMPOSITIONS TO PROTECT AND RESUSCITATE MAMMALIAN CELLS AND METHODS FOR PREPARING AND USING SAME (57) Abstract The present invention pertains to therapeutic compositions for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells. In one embodiment, the therapeutic composition comprises (a) pyruvate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. In another embodiment, the therapeutic composition comprises (a) pyruvate, (b) lactate, and (c) a mixture of saturated and unsaturated fatty acids. In yet another embodiment, the therapeutic composition comprises (a) an antioxidant, and (b) a mixture of saturated and unsaturated fatty acids. In still yet another embodiment, the therapeutic composition comprises (a) lactate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. The therapeutic compositions may be utilized in a wide variety of topical and ingestible pharmaceutical products. This invention also relates to methods for preparing and using the therapeutic compositions and the topical and ingestible pharmaceutical products in which the therapeutic compositions may be used.		

2104461

- 1 -

5

10

**THERAPEUTIC COMPOSITIONS
TO PROTECT AND RESUSCITATE MAMMALIAN CELLS
AND METHODS FOR PREPARING AND USING SAME**

15

BACKGROUND OF THE INVENTION

20

1. Field of the Invention

25

30

35

This invention pertains to therapeutic compositions for protecting and resuscitating mammalian cells. In one embodiment, the therapeutic composition comprises (a) pyruvate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. In another embodiment, the therapeutic composition comprises (a) pyruvate, (b) lactate, and (c) a mixture of saturated and unsaturated fatty acids. In yet another embodiment, the therapeutic composition comprises (a) an antioxidant and (b) a mixture of saturated and unsaturated fatty acids. In still yet another embodiment, the therapeutic composition comprises (a) lactate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. This invention also pertains to methods for preparing and using the therapeutic compositions and the topical and

SUBSTITUTE SHEET

2104461

- 2 -

ingestible pharmaceutical products in which the therapeutic compositions may be used.

5

2. Description of the Prior Art

Mammalian cells are continuously exposed to activated oxygen species such as superoxide (O_2^-),
10 hydrogen peroxide (H_2O_2), hydroxyl radical ($OH^\cdot$), and singlet oxygen ($^1O_2$). In vivo, these reactive oxygen intermediates are generated by cells in response to aerobic metabolism, catabolism of drugs and other xenobiotics, ultraviolet and x-ray radiation, and the
15 respiratory burst of phagocytic cells (such as white blood cells) to kill invading bacteria. Hydrogen peroxide, for example, is produced during respiration of most living organisms especially by stressed and injured cells.

20

These active oxygen species can injure cells. An important example of such damage is lipid peroxidation which involves the oxidative degradation of unsaturated lipids. Lipid peroxidation is highly
25 detrimental to membrane structure and function and can cause numerous cytopathological effects. Cells defend against lipid peroxidation by producing radical scavengers such as superoxide dismutase, catalase, and peroxidase. Injured cells have a decreased ability to
30 produce radical scavengers. Excess hydrogen peroxide can react with DNA to cause backbone breakage, produce mutations, and alter and liberate bases. Hydrogen peroxide can also react with pyrimidines to open the
35 of pyrimidines to hydrogen bond to complementary bases, Hallaender et al. (1971). Such oxidative biochemical injury can result in the loss of cellular membrane

SUBSTITUTE SHEET

2104461

- 3 -

integrity, reduced enzyme activity, changes in transport kinetics, changes in membrane lipid content, and leakage of potassium ions, amino acids, and other cellular material.

5

The production of reactive oxygen intermediates has been suggested to cause many skin, tissue, and organ disorders such as atherosclerosis, arthritis, cytotoxicity, skin inflammation, photoaging, wrinkling, actinic keratosis, tumor formation, cancer, hypertension, Parkinson's disease, lung disease, and heart disease. The role of active oxygen radicals in promoting tumors has been proposed based on the findings that (a) tumor promoters increase the level of oxygen radicals, (b) many free radical generating systems promote tumors, and (c) certain antioxidants inhibit the biochemical effects of tumor promoters.

In vitro, reactive oxygen intermediates can be generated in cellular culture media by autooxidation and photooxidation of media components. During excision and storage, transplant organs can suffer oxidative injuries which result in the loss of cellular membrane integrity and shorten the usable life of the organ.

When cells are stressed by oxidative injury, a resuscitation step is necessary to recondition the cells. Antioxidants have been shown to inhibit damage associated with active oxygen species. For example, pyruvate and other alpha-ketoacids have been reported to react rapidly and stoichiometrically with hydrogen peroxide to protect cells from cytolytic effects, O'Donnell-Tormey et al., J. Exp. Med., 165, pp. 500-514 (1987).

SUBSTITUTE SHEET

2104461

- 4 -

United States patents nos. 3,920,835, 3,984,556, and 3,988,470, all issued to Van Scott et al., disclose methods for treating acne, dandruff, and palmar keratosis, respectively, which consist of applying to the affected area a topical composition comprising from about 1% to about 20% of a lower aliphatic compound containing from two to six carbon atoms selected from the group consisting of *alpha*-hydroxyacids, *alpha*-ketoacids and esters thereof, and 3-hydroxybutyric acid in a pharmaceutically acceptable carrier. The aliphatic compounds include pyruvic acid and lactic acid.

United States patents nos. 4,105,783 and 4,197,316, both issued to Yu et al., disclose a method and composition, respectively, for treating dry skin which consists of applying to the affected area a topical composition comprising from about 1% to about 20% of a compound selected from the group consisting of amides and ammonium salts of *alpha*-hydroxyacids, *beta*-hydroxyacids, and *alpha*-ketoacids in a pharmaceutically acceptable carrier. The compounds include the amides and ammonium salts of pyruvic acid and lactic acid.

United States patent no. 4,234,599, issued to Van Scott et al., discloses a method for treating actinic and nonactinic skin keratoses which consists of applying to the affected area a topical composition comprising an effective amount of a compound selected from the group consisting of *alpha*-hydroxyacids, *beta*-hydroxyacids, and *alpha*-ketoacids in a pharmaceutically acceptable carrier. The acidic compounds include pyruvic acid and lactic acid.

United States patent no. 4,294,852, issued to Wildnauer et al., discloses a composition for treating skin which comprises the *alpha*-hydroxyacids, *beta*-

SUBSTITUTE SHEET

- 5 -

hydroxyacids, and α -ketoacids disclosed above by Van Scott et al. in combination with C_3 - C_8 aliphatic alcohols.

5 United States patent no. 4,663,166, issued to Veech, discloses an electrolyte solution which comprises a mixture of L-lactate and pyruvate in a ratio from 20:1 to 1:1, respectively, or a mixture of D-beta-hydroxybutyrate and acetoacetate, in a ratio
10 from 6:1 to 0.5:1, respectively.

Sodium pyruvate has been reported to reduce the number of erosions, ulcers, and hemorrhages on the gastric mucosa in guinea pigs and rats caused by
15 acetylsalicylic acid. The analgesic and antipyretic properties of acetylsalicylic acid were not impaired by sodium pyruvate, Puschmann, Arzneimittelforschung, 33, pp. 410-415 and 415-416 (1983).

20 Pyruvate has been reported to exert a positive inotropic effect in stunned myocardium, which is a prolonged ventricular dysfunction following brief periods of coronary artery occlusions which does not produce irreversible damage, Mentzer et al., Ann. Surg., 209, pp. 629-633 (1989).
25

Pyruvate has been reported to produce a relative stabilization of left ventricular pressure and work parameter and to reduce the size of infarctions.
30 Pyruvate improves resumption of spontaneous beating of the heart and restoration of normal rates and pressure development, Bunger et al., J. Mol. Cell. Cardiol., 18, pp. 423-438 (1986), Mochizuki et al., J. Physiol. (Paris), 76, pp. 805-812 (1980), Regitz et al., Cardiovasc. Res., 15, pp. 652-658 (1981),
35 Giannelli et al., Ann. Thorac. Surg., 21, pp. 386-396 (1976).

2104461

- 6 -

Sodium pyruvate has been reported to act as an antagonist to cyanide intoxicification (presumably through the formation of a cyanohydrin) and to protect against the lethal effects of sodium sulfide and to retard the onset and development of functional, morphological, and biochemical measures of acrylamide neuropathy of axons, Schwartz et al., Toxicol. Appl. Pharmacol., 50, pp. 437-442 (1979), Sabri et al., Brain Res., 483, pp. 1-11 (1989).

A chemotherapeutic cure of advanced L1210 leukemia has been reported using sodium pyruvate to restore abnormally deformed red blood cells to normal. The deformed red blood cells prevented adequate drug delivery to tumor cells, Cohen, Cancer Chemother. Pharmacol., 5, pp. 175-179 (1981).

Primary cultures of heterotopic tracheal transplant exposed *in vivo* to 7, 12-dimethylbenz(a)anthracene were reported to be successfully maintained in enrichment medium supplemented with sodium pyruvate along with cultures of interleukin-2 stimulated peripheral blood lymphocytes, and plasmacytomas and hybridomas, pig embryos, and human blastocysts, Shacter, J. Immunol. Methods, 99, pp. 259-270 (1987), Marchok et al., Cancer Res., 37, pp. 1811-1821 (1977), Davis, J. Reprod. Fertil. Suppl., 33, pp. 115-124 (1985), Okamoto et al., No To Shinkei, 38, pp. 593-598 (1986), Cohen et al., J. In Vitro Fert. Embryo Transfer, 2, pp. 59-64 (1985).

United States patents nos. 4,158,057, 4,351,835, 4,415,576, and 4,645,764, all issued to Stanko, disclose methods for preventing the accumulation of fat in the liver of a mammal due to the ingestion of alcohol, for controlling weight in a

SUBSTITUTE SHEET

2104461

- 7 -

mammal, for inhibiting body fat while increasing protein concentration in a mammal, and for controlling the deposition of body fat in a living being, respectively. The methods comprise administering to
5 the mammal a therapeutic mixture of pyruvate and dihydroxyacetone, and optionally riboflavin. United States patent no. 4,548,937, issued to Stanko, discloses a method for controlling the weight gain of a mammal which comprises administering to the mammal a
10 therapeutically effective amount of pyruvate, and optionally riboflavin. United States patent no. 4,812,479, issued to Stanko, discloses a method for controlling the weight gain of a mammal which comprises administering to the mammal a therapeutically effective
15 amount of dihydroxyacetone, and optionally riboflavin and pyruvate.

Rats fed a calcium-oxalate lithogenic diet including sodium pyruvate were reported to develop
20 fewer urinary calculi (stones) than control rats not given sodium pyruvate, Ogawa et al., Hinyokika Kyo, 32, pp. 1341-1347 (1986).

United States patent no. 4,521,375, issued to
25 Houlsby, discloses a method for sterilizing surfaces which come into contact with living tissue. The method comprises sterilizing the surface with aqueous hydrogen peroxide and then neutralizing the surface with pyruvic acid.

30
United States patent no. 4,416,982, issued to Tauda et al., discloses a method for decomposing hydrogen peroxide by reacting the hydrogen peroxide with a phenol or aniline derivative in the presence of
35 peroxidase.

SUBSTITUTE SHEET

2104461

- 8 -

United States patent no. 4,696,917, issued to Lindstrom et al., discloses an irrigation solution which comprises Eagle's Minimum Essential Medium with Earle's salts, chondroitin sulfate, a buffer solution, 2-mercaptoethanol, and a pyruvate. The irrigation solution may optionally contain ascorbic acid and α -tocopherol. United States patent no. 4,725,586, issued to Lindstrom et al., discloses an irrigation solution which comprises a balanced salt solution, chondroitin sulfate, a buffer solution, 2-mercaptoethanol, sodium bicarbonate or dextrose, a pyruvate, a sodium phosphate buffer system, and cystine. The irrigation solution may optionally contain ascorbic acid and γ -tocopherol.

15

United States patent no. 3,887,702 issued to Baldwin, discloses a composition for treating fingernails and toenails which consists essentially of soybean oil or sunflower oil in combination with Vitamin E.

20

United States patent no. 4,847,069, issued to Bissett et al., discloses a photoprotective composition comprising (a) a sorbohydroxamic acid, (b) an anti-inflammatory agent selected from steroidal anti-inflammatory agents and a natural anti-inflammatory agent, and (c) a topical carrier. Fatty acids may be present as an emollient. United States patent no. 4,847,071, issued to Bissett et al., discloses a photoprotective composition comprising (a) a tocopherol or tocopherol ester radical scavenger, (b) an anti-inflammatory agent selected from steroidal anti-inflammatory agents and a natural anti-inflammatory agent, and (c) a topical carrier. United States patent no. 4,847,072, issued to Bissett et al., discloses a topical composition comprising not more than 25% tocopherol sorbate in a topical carrier.

25

30

35

SUBSTITUTE SHEET

2104461

- 9 -

United States patent no. 4,533,637, issued to Yamane et al., discloses a culture medium which comprises a carbon source, a nucleic acid source precursor, amino acids, vitamins, minerals, a lipophilic nutrient, and serum albumin, and cyclodextrins. The lipophilic substances include unsaturated fatty acids and lipophilic vitamins such as Vitamin A, D, and E. Ascorbic acid may also be present.

United Kingdom patent application no. 2,196,348A, to Kovar et al., discloses a synthetic culture medium which comprises inorganic salts, monosaccharides, amino acids, vitamins, buffering agents, and optionally sodium pyruvate adding magnesium hydroxide or magnesium oxide to the emulsion. The oil phase may include chicken fat.

United States patent no. 4,284,630, issued to Yu et al., discloses a method for stabilizing a water-in-oil emulsion which comprises adding magnesium hydroxide or magnesium oxide to the emulsion. The oil phase may include chicken fat.

Preparation-H has been reported to increase the rate of wound healing in artificially created rectal ulcers. The active ingredients in Preparation-H are skin respiratory factor and shark oil, Subramanyam et al., Digestive Diseases and Sciences, 29, pp. 829-832 (1984).

The addition of sodium pyruvate to bacterial and yeast systems has been reported to inhibit hydrogen peroxide production, enhance growth, and protect the systems against the toxicity of reactive oxygen intermediates. The optimum ratio of unsaturated to

2104461 - 10 -

5 saturated fatty acids contained within chicken fat enhanced membrane repair and reduced cytotoxicity. The antioxidants glutathione and thioglycollate reduced the injury induced by oxygen radical species, Martin, Ph.D. thesis, (1987-89).

10 While the above therapeutic compositions are reported to inhibit the production of reactive oxygen intermediates, none of the above compositions are entirely satisfactory. None of the compositions has the ability to simultaneously decrease cellular levels of hydrogen peroxide production, increase cellular resistance to cytotoxic agents, increase rates of cellular proliferation, and increase cellular viability to protect and resuscitate mammalian cells. The present invention provides such improved therapeutic compositions without the disadvantages characteristic of previously known compositions. This invention also relates to methods for preparing and using the therapeutic compositions and the topical and ingestible pharmaceutical products in which the therapeutic compositions may be used.

25 SUMMARY OF THE INVENTION

30 The present invention pertains to therapeutic compositions for preventing and reducing injury to mammalian cells and increasing the resuscitation rate of injured mammalian cells. In one embodiment, the therapeutic composition comprises (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids wherein the

35

2104461

- 11 -

fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

5 In another embodiment, the invention pertains to a therapeutic composition which comprises (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

15

In yet another embodiment, the invention pertains to a therapeutic composition which comprises (a) an antioxidant, and (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

20

25 In still yet another embodiment, the invention pertains to a therapeutic composition which comprises (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

30

35 The therapeutic compositions may be utilized in a wide variety of topical and ingestible pharmaceutical products. This invention also relates to methods for preparing and using the therapeutic compositions and the topical and ingestible

SUBSTITUTE SHEET

2104461

- 12 -

pharmaceutical products in which the therapeutic compositions may be used.

5

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1 depicts in bar graph format the viability of U937 monocytic cells following exposure of the cells to various antioxidants (Examples 1-5).

FIGURE 2 depicts in bar graph format the viability of U937 monocytic cells following exposure of the cells to various combinations of antioxidants (Examples 6-13).

FIGURE 3 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells following exposure of the cells to various antioxidants (Examples 14-18).

FIGURE 4 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells following exposure of the cells to various combinations of antioxidants (Examples 19-26).

FIGURE 5 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells following exposure of the cells to various combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids (Examples 27-32).

FIGURE 6 depicts in bar graph format the levels of hydrogen peroxide produced by epidermal keratinocytes following exposure of the cells to

SUBSTITUTE SHEET

- 13 -

various antioxidants with and without a mixture of saturated and unsaturated fatty acids (Examples 33-42).

FIGURE 7 depicts in bar graph format the levels of hydrogen peroxide produced by epidermal keratinocytes following exposure of the cells to various combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids (Examples 43-52).

10

DETAILED DESCRIPTION OF THE INVENTION

Applicant has discovered therapeutic compositions for preventing and reducing injury to mammalian cells and increasing the resuscitation rate of injured mammalian cells. Cells treated with the therapeutic compositions of the present invention show decreased levels of hydrogen peroxide production, increased resistance to cytotoxic agents, increased rates of proliferation, and increased viability. Cellular cultures containing the therapeutic compositions showed enhanced differentiation and proliferation over control cultures and rapidly formed attachments or tight junctions between the cells to form an epidermal sheet.

The term "injured cell" as used herein means a cell which has (a) injured membranes so that transport through the membranes is diminished resulting in an increase in toxins and normal cellular wastes inside the cell and a decrease in nutrients and other components necessary for cellular repair inside the cell, (b) an increase in concentration of oxygen radicals inside the cell because of the decreased ability of the cell to produce antioxidants and

2104461

- 14 -

enzymes, and (c) damaged DNA, RNA, and ribosomes which must be repaired or replaced before normal cellular functions can be resumed. The term "resuscitation" of injured mammalian cells as used herein means the reversal of cytotoxicity, the stabilization of the cellular membrane, an increase in the proliferation rate of the cell, and/or the normalization of cellular functions such as the secretion of growth factors, hormones, and the like. The term "cytotoxicity" as used herein means a condition caused by a cytotoxic agent that injures the cell. Injured cells do not proliferate because injured cells expend all energy on cellular repair. Aiding cellular repair promotes cellular proliferation.

15

Epidermal keratinocytic cells and monocytic cells have multiple oxygen generating mechanisms and the degree to which each type of mechanism functions differs in each type of cell. In monocytes, for example, the respiratory bursting process is more pronounced than in epidermal keratinocytes. Hence, the components in the therapeutic compositions of the present invention may vary depending upon the types of cells involved in the condition being treated.

25

In a first embodiment, the therapeutic composition for treating mammalian cells, preferably epidermal keratinocytes, comprises (a) pyruvate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids.

30

While not wishing to be bound by theory, applicant believes that pyruvate can be transported inside a cell where it can act as an antioxidant to neutralize oxygen radicals in the cell. Pyruvate can also be used inside the cell in the citric acid cycle to provide energy to increase cellular viability, and

35

SUBSTITUTE SHEET

- 15 -

as a precursor in the synthesis of important biomolecules to promote cellular proliferation. In addition, pyruvate can be used in the multifunction oxidase system to reverse cytotoxicity. Antioxidants, especially lipid-soluble antioxidants, can be absorbed into the cell membrane to neutralize oxygen radicals and thereby protect the membrane. The combination of pyruvate inside the cell and an antioxidant in the cellular membrane functions in a synergistic manner to reduce hydrogen peroxide production in the cell to levels lower than can be achieved by use of either type of component alone.

The saturated and unsaturated fatty acids in the present invention are those fatty acids required for the resuscitation of mammalian cells. Hence, the fatty acids in the therapeutic composition are readily available for the repair of injured cells. Cells injured by oxygen radicals need to produce unsaturated fatty acids to repair cellular membranes. However, the production of unsaturated fatty acids by cells requires oxygen. Thus, the injured cell needs high levels of oxygen to produce unsaturated fatty acids and at the same time needs to reduce the level of oxygen within the cell to reduce oxidative injury. By providing the cell with the unsaturated fatty acids needed for repair, the need of the cell for unsaturated fatty acids is reduced and the need for high oxygen levels is also reduced. The presence of mixtures of saturated and unsaturated fatty acids in the therapeutic composition significantly enhances the ability of pyruvate and the antioxidant to inhibit reactive oxygen production. By stabilizing the cellular membrane, unsaturated fatty acids also improve membrane function and enhance pyruvate transport into the cell. Hence, the three components in the therapeutic composition function together in a synergistic manner to prevent

2104461

- 16 -

and reduce injury to mammalian cells and increase the resuscitation rate of injured mammalian cells.

5 In a second embodiment, the therapeutic composition for treating mammalian cells, preferably epidermal keratinocytes, comprises (a) pyruvate, (b) lactate, and (c) a mixture of saturated and unsaturated fatty acids. In this embodiment, lactate is employed instead of an antioxidant. Antioxidants react with, 10 and neutralize, oxygen radicals after the radicals are already formed. Lactate, on the other hand, is a component in the cellular feedback mechanism and inhibits the respiratory bursting process to suppress the production of active oxygen species. The 15 combination of pyruvate to neutralize active oxygen species and lactate to suppress the respiratory bursting process functions in a synergistic manner to reduce hydrogen peroxide production in the cell to levels lower than can be achieved by use of either type 20 of component alone. The presence of mixtures of saturated and unsaturated fatty acids in the therapeutic composition significantly enhances the ability of pyruvate and lactate to inhibit reactive oxygen production. Hence, the three components in the 25 therapeutic composition in this embodiment function together in a synergistic manner to protect and resuscitate mammalian cells.

30 In a third embodiment, the therapeutic composition for treating mammalian cells, preferably epidermal keratinocytes, comprises (a) an antioxidant, and (b) a mixture of saturated and unsaturated fatty acids. The presence of mixtures of saturated and unsaturated fatty acids in the therapeutic composition 35 in this embodiment significantly enhances the ability of the antioxidant to inhibit reactive oxygen production. The combination of an antioxidant to

SUBSTITUTE SHEET

2104461

- 17 -

neutralize active oxygen species and fatty acids to rebuild cellular membranes and reduce the need of the cell for oxygen functions in a synergistic manner to reduce hydrogen peroxide production in the cell to
5 levels lower than can be achieved by either type of component alone. Hence, the components in the therapeutic composition in this embodiment function together in a synergistic manner to protect and resuscitate mammalian cells.

10

In a fourth embodiment, the therapeutic composition for treating mammalian cells, preferably monocytes, comprises (a) lactate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty
15 acids. In this embodiment, lactate is employed because the respiratory bursting process is more pronounced in monocytes than in epidermal keratinocytes. The combination of lactate to suppress the respiratory bursting process and an antioxidant to neutralize
20 active oxygen species functions in a synergistic manner to reduce hydrogen peroxide production in the cell to levels lower than can be achieved by either component alone. The presence of mixtures of saturated and unsaturated fatty acids in the therapeutic composition
25 in this embodiment significantly enhances the ability of lactate and the antioxidant to inhibit reactive oxygen production. Hence, the three components in the therapeutic composition in this embodiment function together in an unexpected synergistic manner to protect
30 and resuscitate mammalian cells.

Accordingly, the combination of ingredients set out in the above embodiments functions together in an enhanced manner to prevent and reduce injury to
35 mammalian cells and increase the resuscitation rate of injured mammalian cells. The therapeutic effect of the combination of the components in each of the above

SUBSTITUTE SHEET

2104461 - 18 -

embodiments is markedly greater than that expected by the mere addition of the individual therapeutic components. Hence, applicant's therapeutic compositions for treating mammalian cells have the ability to decrease intracellular levels of hydrogen peroxide production, increase cellular resistance to cytotoxic agents, increase rates of cellular proliferation, and increase cellular viability.

The cells which may be treated with the therapeutic compositions in the present invention are mammalian cells. Although applicant will describe the present therapeutic compositions as useful for treating mammalian epidermal keratinocytes and mammalian monocytes, applicant contemplates that all mammalian cells which may be protected or resuscitated by applicant's therapeutic compositions may be used in the present invention. Keratinocytes are representative of normal mammalian cells and are the fastest proliferating cells in the body. The correlation between the reaction of keratinocytes to injury and therapy and that of mammalian cells in general is very high. Monocytes are representative of specialized mammalian cells such as the white blood cells in the immune system and the organ cells in liver, kidney, heart, and brain. The mammalian cells may be treated *in vivo* and *in vitro*.

Epidermal keratinocytes are the specialized epithelial cells of the epidermis which synthesize keratin, a scleroprotein which is the principal constituent of epidermis, hair, nails, horny tissue, and the organic matrix of the enamel of teeth. Mammalian epidermal keratinocytes constitute about 95% of the epidermal cells and together with melanocytes form the binary system of the epidermis. In its various successive stages, epidermal keratinocytes are

- 19 -

also known as basal cells, prickle cells, and granular cells.

Monocytes are mononuclear phagocytic leukocytes which undergo respiratory bursting and are involved in reactive oxygen mediated damage within the epidermis. Leukocytes are white blood cells or corpuscles which may be classified into two main groups: granular leukocytes (granulocytes) which are leukocytes with abundant granules in the cytoplasm and nongranular leukocytes (nongranulocytes) which are leukocytes without specific granules in the cytoplasm and which include the lymphocytes and monocytes. Phagocyte cells are cells which ingest microorganisms or other cells and foreign particles. Monocytes are also known as large mononuclear leukocytes, and hyaline or transitional leukocytes.

Pyruvic acid (2-oxopropanoic acid, α -ketopropionic acid, CH_3COCOOH) or pyruvate is a fundamental intermediate in protein and carbohydrate metabolism and in the citric acid cycle. The citric acid cycle (tricarboxylic acid cycle, Krebs's cycle) is the major reaction sequence which executes the reduction of oxygen to generate adenosine triphosphate (ATP) by oxidizing organic compounds in respiring tissues to provide electrons to the transport system. Acetyl coenzyme A ("active acetyl") is oxidized in this process and is thereafter utilized in a variety of biological processes and is a precursor in the biosynthesis of many fatty acids and sterols. The two major sources of acetyl coenzyme A are derived from the metabolism of glucose and fatty acids. Glycolysis consists of a series of transformations wherein each glucose molecule is transformed in the cellular cytoplasm into two molecules of pyruvic acid. Pyruvic acid may then enter the mitochondria where it is

SUBSTITUTE SHEET

2104461

- 20 -

oxidized by coenzyme A in the presence of enzymes and cofactors to acetyl coenzyme A. Acetyl coenzyme A can then enter the citric acid cycle.

5 In muscle, pyruvic acid (derived from glycogen) is reduced to lactic acid during exertion. Lactic acid is reoxidized and partially retransformed to glycogen during rest. Pyruvate can also act as an antioxidant to neutralize oxygen radicals in the cell
10 and can be used in the multifunction oxidase system to reverse cytotoxicity.

 The pyruvate in the present invention may be selected from the group consisting of pyruvic acid,
15 pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof. In general, the pharmaceutically acceptable salts of pyruvic acid may be alkali salts and alkaline earth salts. Preferably, the pyruvate is selected from the group consisting of pyruvic acid,
20 lithium pyruvate, sodium pyruvate, potassium pyruvate, magnesium pyruvate, calcium pyruvate, zinc pyruvate, manganese pyruvate, and mixtures thereof. More preferably, the pyruvate is selected from the group of salts consisting of sodium pyruvate, potassium
25 pyruvate, magnesium pyruvate, calcium pyruvate, zinc pyruvate, manganese pyruvate, and mixtures thereof. Most preferably, the pyruvate is sodium pyruvate.

 The amount of pyruvate present in the
30 therapeutic compositions of the present invention is a therapeutically effective amount. A therapeutically effective amount of pyruvate is that amount of pyruvate necessary to prevent and reduce injury to mammalian cells and increase the resuscitation rate of injured
35 mammalian cells. The exact amount of pyruvate is a matter of preference subject to such factors as the type of condition being treated as well as the other

2104461

- 21 -

ingredients in the composition. In a preferred embodiment, pyruvate is present in the therapeutic composition in an amount from about 10% to about 50%, preferably from about 20% to about 45%, and more preferably from about 25% to about 40%, by weight of the therapeutic composition.

L-Lactic acid ((*S*)-2-hydroxypropanoic acid, (+) *alpha*-hydroxypropionic acid, $\text{CH}_3\text{CHOHCOOH}$) or lactate occurs in small quantities in the blood and muscle fluid of mammals. Lactic acid concentration increases in muscle and blood after vigorous activity. Lactate is a component in the cellular feedback mechanism and inhibits the natural respiratory bursting process of cells thereby suppressing the production of oxygen radicals.

The lactate in the present invention may be selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof. In general, the pharmaceutically acceptable salts of lactic acid may be alkali salts and alkaline earth salts. Preferably, the lactate is selected from the group consisting of lactic acid, lithium lactate, sodium lactate, potassium lactate, magnesium lactate, calcium lactate, zinc lactate, manganese lactate, and mixtures thereof. More preferably, the lactate is selected from the group consisting of lactic acid, sodium lactate, potassium lactate, magnesium lactate, calcium lactate, zinc lactate, manganese lactate, and mixtures thereof. Most preferably, the lactate is lactic acid.

The amount of lactate present in the therapeutic compositions of the present invention is a therapeutically effective amount. A therapeutically effective amount of lactate is that amount of lactate

SUBSTITUTE SHEET

- 22 -

necessary to prevent and reduce injury to mammalian cells and increase the resuscitation rate of injured mammalian cells. For an ingestible composition, a therapeutically effective amount of lactate is that amount necessary to suppress the respiratory bursting process of white blood cells to protect and resuscitate the mammalian cells. In general, a therapeutically effective amount of lactate in an ingestible composition is from about 5 to about 10 times the amount of lactate normally found in serum. The exact amount of lactate is a matter of preference subject to such factors as the type of condition being treated as well as the other ingredients in the composition. In a preferred embodiment, lactate is present in the therapeutic composition in an amount from about 10% to about 50%, preferably from about 20% to about 45%, and more preferably from about 25% to about 40%, by weight of the therapeutic composition.

Antioxidants are substances which inhibit oxidation or suppress reactions promoted by oxygen or peroxides. Antioxidants, especially lipid-soluble antioxidants, can be absorbed into the cellular membrane to neutralize oxygen radicals and thereby protect the membrane. The antioxidants useful in the present invention may be selected from the group consisting of Vitamin A (retinol), Vitamin A₂ (3, 4-didehydroretinol), all forms of carotene such as alpha-carotene, beta-carotene (beta, beta-carotene), gamma-carotene, delta-carotene, Vitamin C (ascorbic acid, L-ascorbic acid), all forms of tocopherol such as Vitamin E (alpha-tocopherol, 3,4-dihydro-2,5,7,8-tetramethyl-2-(4,8,12-trimethyltri-decyl)-2H-1-benzopyran-6-ol), beta-tocopherol, gamma-tocopherol, and delta-tocopherol, and mixtures thereof. Preferably, the antioxidant is selected from the group of lipid-soluble antioxidants consisting of Vitamin A,

SUBSTITUTE SHEET

2104461

- 23 -

beta-carotene, Vitamin E, and mixtures thereof. More preferably, the antioxidant is Vitamin E.

5 The amount of antioxidant present in the therapeutic compositions of the present invention is a therapeutically effective amount. A therapeutically effective amount of antioxidant is that amount of antioxidant necessary to prevent and reduce injury to mammalian cells and increase the resuscitation rate of
10 injured mammalian cells. The exact amount of antioxidant is a matter of preference subject to such factors as the type of condition being treated as well as the other ingredients in the composition. In a preferred embodiment, the antioxidant is present in the
15 therapeutic composition in an amount from about 10% to about 50%, preferably from about 20% to about 45%, and more preferably from about 25% to about 40%, by weight of the therapeutic composition.

20 The mixture of saturated and unsaturated fatty acids in the present invention are those fatty acids required for the repair of mammalian cellular membranes. Hence, the fatty acids are readily incorporated into the cell and are immediately
25 available for the repair of injured cells. By providing the cell with the unsaturated fatty acids needed for repair, the need of the cell for unsaturated fatty acids is reduced and the need for high oxygen levels is also reduced. Accordingly, the presence of
30 the mixtures of saturated and unsaturated fatty acids in the therapeutic compositions significantly enhances the ability of pyruvate, lactate, and the antioxidant to inhibit reactive oxygen production.

35 Fatty acids are carboxylic acid compounds found in animal and vegetable fat and oil. Fatty acids are classified as lipids and are composed of chains of

SUBSTITUTE SHEET

2104461

- 24 -

alkyl groups containing from 4 to 22 carbon atoms and 0-3 double bonds and characterized by a terminal carboxyl group, -COOH. Fatty acids may be saturated or unsaturated and may be solid, semisolid, or liquid.

5 The most common saturated fatty acids are butyric acid (C_4), lauric acid (C_{12}), palmitic acid (C_{16}), and stearic acid (C_{18}). Unsaturated fatty acids are usually derived from vegetables and consist of alkyl chains containing from 16 to 22 carbon atoms and 0-3

10 double bonds with the characteristic terminal carboxyl group. The most common unsaturated fatty acids are oleic acid, linoleic acid, and linolenic acid (all C_{18} acids).

15 In general, the mixture of saturated and unsaturated fatty acids required for the repair of mammalian cellular membranes in the present invention may be derived from animal fats and waxes. Cells produce the chemical components and the energy required

20 for cellular viability and store excess energy in the form of fat. Fat is adipose tissue stored between organs of the body to furnish a reserve supply of energy. The preferred animal fats and waxes have a fatty acid composition similar to that of human fat and

25 the fat contained in human breast milk. The preferred animal fats and waxes may be selected from the group consisting of human fat, chicken fat, cow fat (defined herein as a bovine domestic animal regardless of sex or age), sheep fat, horse fat, pig fat, and whale fat.

30 The more preferred animal fats and waxes may be selected from the group consisting of human fat and chicken fat. The most preferred animal fat is human fat. Mixtures of other fats and waxes, such as vegetable waxes and synthetic waxes, which have a fatty

35 acid composition similar to that of animal fats and waxes, and preferably to that of human fats and waxes, may also be employed. The mixture of saturated and

SUBSTITUTE SHEET

- 25 -

unsaturated fatty acids may also be derived from shark liver oil.

5 In a preferred embodiment, the mixture of saturated and unsaturated fatty acids has composition similar to that of human fat and comprises the following fatty acids: butyric acid, caproic acid, caprylic acid, capric acid, lauric acid, myristic acid, myristoleic acid, palmitic acid, palmitoleic acid, 10 stearic, oleic acid, linoleic acid, linolenic acid, arachidic acid, and gaddoleic acid. Preferably, butyric acid, caproic acid, caprylic acid, capric acid, lauric acid, myristic acid, myristoleic acid, palmitic acid, palmitoleic acid, stearic, oleic acid, linoleic acid, 15 linolenic acid, arachidic acid, and gaddoleic acid are present in the mixture in about the following percentages by weight, respectively (carbon chain number and number of unsaturations are shown parenthetically, respectively): 0.2%-0.4% (C_4), 0.1% (C_6), 0.3%-0.8% (C_8), 2.2%-3.5% (C_{10}), 0.9%-5.5% (C_{12}), 2.8%-8.5% (C_{14}), 0.1%-0.6% ($C_{14:1}$), 23.2%-24.6% (C_{16}), 1.8%-3.0% ($C_{16:1}$), 6.9%-9.9% (C_{18}), 36.0%-36.5% ($C_{18:1}$), 20%-20.6% ($C_{18:2}$), 7.5-7.8% ($C_{18:3}$), 1.1%-4.9% (C_{20}), and 3.3%-6.4% ($C_{20:1/2}$).

25

In another preferred embodiment, the mixture of saturated and unsaturated fatty acids is chicken fat comprising the following fatty acids: lauric acid, myristic acid, myristoleic acid, pentadecanoic acid, 30 palmitic acid, palmitoleic acid, margaric acid, margaroleic acid, stearic, oleic acid, linoleic acid, linolenic acid, arachidic acid, and gaddoleic acid. Preferably, lauric acid, myristic acid, myristoleic acid, pentadecanoic acid, palmitic acid, palmitoleic acid, margaric acid, margaroleic acid, stearic, oleic acid, 35 linoleic acid, linolenic acid, arachidic acid, and gaddoleic acid are present in the mixture in about

SUBSTITUTE SHEET

2104461 - 26 -

the following percentages by weight, respectively: 0.1%
(C₁₂), 0.8% (C₁₄), 0.2% (C_{14:1}), 0.1% (C₁₅), 25.3%
(C₁₆), 7.2% (C_{16:1}), 0.1% (C₁₇), 0.1% (C_{17:1}), 6.5%
(C₁₈), 37.7% (C_{18:1}), 20.6% (C_{18:2}), 0.8% (C_{18:3}), 0.2%
5 (C₂₀), and 0.3% (C_{20:1}).

The above fatty acids and percentages thereof
present in the fatty acid mixture are given as an
example. The exact type of fatty acid present in the
10 fatty acid mixture and the exact amount of fatty acid
employed in the fatty acid mixture may be varied in
order to obtain the result desired in the final product
and such variations are now within the capabilities of
those skilled in the art without the need for undue
15 experimentation.

The amount of fatty acids present in the
therapeutic compositions of the present invention is a
therapeutically effective amount. A therapeutically
20 effective amount of fatty acids is that amount of fatty
acids necessary to prevent and reduce injury to
mammalian cells and increase the resuscitation rate of
injured mammalian cells. The exact amount of fatty
acids employed is subject to such factors as the type
25 and distribution of fatty acids employed in the
mixture, the type of condition being treated, and the
other ingredients in the composition. In a preferred
embodiment, the fatty acids are present in the
therapeutic composition in an amount from about 10% to
30 about 50%, preferably from about 20% to about 45%, and
more preferably from about 25% to about 40%, by weight
of the therapeutic composition.

The present invention extends to methods for
35 making the therapeutic compositions. In general, a
therapeutic composition is made by forming an admixture
of the components of the composition. In one

SUBSTITUTE SHEET

- 27 -

embodiment, a therapeutic composition is made by forming an admixture of (a) a pyruvate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids. In a second embodiment, a
5 therapeutic composition is made by forming an admixture of (a) a pyruvate, (b) a lactate, and (c) a mixture of saturated and unsaturated fatty acids. In a third embodiment, a therapeutic composition is made by forming an admixture of (a) an antioxidant, and (b) a
10 mixture of saturated and unsaturated fatty acids. In a fourth embodiment, a therapeutic composition is made by forming an admixture of (a) a lactate, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids.

15

For some applications, the admixture may be formed in a solvent such as water. If necessary, the pH of the solvent is adjusted to a range from about 3.5 to about 8.0, and preferably from about 4.5 to about
20 7.5, and more preferably about 6.0 to about 7.4. The admixture is then sterile filtered. Other ingredients may also be incorporated into the therapeutic composition as dictated by the nature of the desired composition as well known by those having ordinary
25 skill in the art. The ultimate therapeutic compositions are readily prepared using methods generally known in the pharmaceutical arts.

The present invention extends to methods for
30 employing the therapeutic compositions in vivo and in vitro. In general, a therapeutic composition is employed by contacting the therapeutic composition with mammalian cells.

35 In one embodiment, the invention is directed at a method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate

2104461

- 28 -

of injured mammalian cells, which comprises the steps of (A) providing a therapeutic composition which comprises (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells, and (B) contacting the therapeutic composition with the mammalian cells.

In a second embodiment, the invention is directed at a method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of (A) providing a therapeutic composition which comprises (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells, and (B) contacting the therapeutic composition with the mammalian cells.

In a third embodiment, the invention is directed at a method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of (A) providing a therapeutic composition which comprises (a) an antioxidant, and (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells, and (B)

SUBSTITUTE SHEET

2104461

- 29 -

contacting the therapeutic composition with the mammalian cells.

5 In a fourth embodiment, the invention is directed at a method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of (A) providing a therapeutic composition which comprises (a) lactate selected from the group
10 consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, (b) an antioxidant, and (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
15 injured mammalian cells, and (B) contacting the therapeutic composition with the mammalian cells.

The therapeutic compositions of the present invention may be utilized in a wide variety of topical
20 and ingestible pharmaceutical products. The therapeutic compositions may be used alone, may be used in combination with medicaments useful for treating injured mammalian cells to prepare augmented pharmaceutical compositions, and may be used in
25 combination with medicaments which are cytotoxic to mammalian cells to prepare cytoprotective pharmaceutical compositions.

When the therapeutic compositions are used to
30 treat mammalian cells in combination a medicament, the therapeutic composition may be selected from the group consisting of:

- (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
35 salts of pyruvic acid, and mixtures thereof;
(b) an antioxidant; and

SUBSTITUTE SHEET

2104461

- 30 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
10 salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

15 (3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

20 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and
25 unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

In a preferred embodiment, the therapeutic
30 composition for treating mammalian cells, preferably epidermal keratinocytes, in combination a medicament may be selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
35 salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

SUBSTITUTE SHEET

- 31 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

15 (3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

20

In a more preferred embodiment, the therapeutic composition for treating mammalian cells, preferably epidermal keratinocytes, in combination a medicament may be selected from the group consisting of:

25

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

30

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

35

2104461

- 32 -

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

5 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

10 In a most preferred embodiment, the therapeutic composition for treating mammalian cells, preferably epidermal keratinocytes, in combination a medicament comprises:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
15 salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for membrane repair and
20 stabilization.

In another preferred embodiment, the therapeutic composition for treating mammalian cells, preferably monocytes, in combination a medicament
25 comprises:

(a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

30 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

35 In one embodiment, the therapeutic compositions may be utilized by themselves to protect, that is, to prevent and reduce injury to mammalian

- 33 -

cells, and increase the resuscitation rate of injured mammalian cells. In a second embodiment, the therapeutic compositions may be combined with a medicament which is useful for treating injured mammalian cells to form augmented pharmaceutical compositions having an enhanced ability to prevent and reduce injury to mammalian cells and further increase the resuscitation rate of injured mammalian cells. In a third embodiment, the therapeutic compositions may be combined with a medicament which is cytotoxic to cells to form cytoprotective pharmaceutical compositions to prevent and reduce injury to mammalian cells from the cytotoxic medicament, increase the resuscitation rate of injured mammalian cells, and thereby increase the therapeutic effect of the cytotoxic medicament. In a fourth embodiment, the therapeutic compositions, without the mixture of saturated and unsaturated fatty acids, may be combined with a medicament which is cytotoxic to cells to form prophylactic cytoprotective pharmaceutical compositions for preventing and reducing injury to mammalian cells from the cytotoxic medicament and thereby increasing the therapeutic effect of the cytotoxic medicament. In a fifth embodiment, therapeutic compositions comprising pyruvate may be combined with a medicament which is cytotoxic to cells to form preventative cytoprotective pharmaceutical compositions for preventing and reducing injury to mammalian cells from the cytotoxic medicament and thereby increasing the therapeutic effect of the cytotoxic medicament. In a sixth embodiment, the components of the therapeutic compositions may be utilized in wound healing pharmaceutical compositions to increase the resuscitation rate of injured mammalian cells.

35

In the first embodiment, the therapeutic compositions may be utilized by themselves in topical

SUBSTITUTE SHEET

2104461

- 34 -

products, ingestible products, and tissue culture medium to protect mammalian cells and increase the resuscitation rate of injured mammalian cells. For example, the therapeutic compositions may be used in
5 topical skin care products to protect and increase the resuscitation rate of skin tissue such as in the treatment of various dermatological disorders such as hyperkeratosis, photo-aging, and sunburn photoreactive processes. The topical therapeutic compositions may
10 also be used orally in the form of a mouth wash or spray to protect and accelerate the healing of injured oral tissue such as mouth sores and burns. The topical therapeutic compositions may further be used in ophthalmological preparations such as eye care products
15 to neutralize hydrogen peroxide used in the cleaning of contact lenses. The topical therapeutic compositions may in addition be used in anorectal creams and suppositories to treat such conditions as pruritus ani, proctitis, anal fissures, and hemorrhoids.

20

The therapeutic compositions may also be used in ingestible products to protect and increase the resuscitation rate of erosions, stomach ulcers, and hemorrhages in the gastric mucosa. Other ingestible
25 therapeutic products include: heart medication to improve regional ventricular function and restore normal heart rate and pressure functions; lung medication to suppress generalized polymorphonuclear cell activation; liver medication to suppress
30 lipogenesis of alcoholic origin and prevent hepatic steatosis; kidney medication to suppress urinary calculi (kidney stones); detoxification medication to antagonize heavy metal poisoning, cyanide poisoning, sodium sulfide poisoning, other types of poisoning;
35 and reduce and neutralize the production of oxygen radicals which produces injury to tissue.

SUBSTITUTE SHEET

- 35 -

The therapeutic compositions of the present invention may also be used in tissue culture media and organ transplant media to prevent and reduce injury to mammalian cells and increase the resuscitation rate of injured mammalian cells. Tissue cultures and transplant organs encounter reactive oxygen species generated in the culture media by the injured cells. Organs particularly susceptible to oxidative damage during transport and transplantation due to reperfusion injury following ischemia are corneas, livers, hearts, and kidneys. The therapeutic compositions may be useful to abrogate reperfusion injury to such transplant organs.

In the second embodiment, the therapeutic compositions may be combined with a medicament which is useful for treating injured mammalian cells to form topical and ingestible augmented pharmaceutical compositions. In this embodiment, the combination of the therapeutic compositions of the present invention and the medicament useful for treating injured mammalian cells provides an augmented pharmaceutical composition having an enhanced ability to prevent and reduce injury to mammalian cells and further increase the resuscitation rate of injured mammalian cells. The tissue damage associated with many diseases and conditions such as autoimmune disease and benign and malignant skin growths is believed to be caused by the production of cellular produced active oxygen species. Combination of the therapeutic compositions of the present invention and medicaments useful for treating such diseases and conditions may suppress such reactive oxygen-linked tissue injury. For example, the therapeutic compositions may be used in topical augmented pharmaceutical compositions in combination with medicaments such as anti-inflammatories, antibacterials, antiseptics, burn relief medications,

SUBSTITUTE SHEET

2104461

- 36 -

sun burn medications, acne preparations, insect bite and sting medications, wound cleansers and wound dressings, to protect and further enhance the resuscitation rate of the injured mammalian cells. The
5 therapeutic compositions may also be used in ingestible augmented pharmaceutical compositions in combination with medicaments used to treat injured mammalian cells such as stroke medications, autoimmune disease medications, arthritis medications, ulcer medications,
10 and cancer medications, to protect and further enhance the resuscitation rate of the injured mammalian cells.

In a specific embodiment, the invention is directed at an augmented pharmaceutical composition
15 having an enhanced ability to prevent and reduce injury to mammalian cells which comprises:

(A) a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group
20 consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are
25 those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

30 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are
35 those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

SUBSTITUTE SHEET

- 37 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a medicament useful for treating injured mammalian cells.

15

In a preferred embodiment, the augmented pharmaceutical composition comprises the therapeutic compositions of the present invention and a medicament useful for treating injured mammalian cells selected from the group consisting of anti-inflammatories and wound cleansers and wound dressings. In a more preferred embodiment, the augmented pharmaceutical composition comprises a medicament selected from the group consisting of wound cleansers and wound dressings.

20

25

In the third embodiment, the therapeutic compositions may be combined with a medicament which is cytotoxic to cells to form topical and ingestible cytoprotective pharmaceutical compositions. In this embodiment, the combination of the therapeutic compositions of the present invention and the medicament cytotoxic to cells provides a cytoprotective pharmaceutical composition having an ability to prevent and reduce injury to mammalian cells from a cytotoxic medicament, increase the resuscitation rate of injured mammalian cells, and thereby increase the dose of the

30

35

2104461

- 38 -

cytotoxic medicament. Medicaments taken on a long term regimen tend to cause liver, kidney, tissue, and other toxicity problems. In addition, certain cytotoxic medicaments, such as potent chemotherapeutic medicaments used to treat malignant tissues, are believed to stimulate release of significant amounts of reactive oxygen species by mammalian tissues which can cause oxidative injury. Combination of the therapeutic compositions of the present invention with such cytotoxic medicaments may inhibit induction of reactive oxygen production while simultaneously decreasing side effects of such medicaments. By decreasing the side effects of such medicaments, the dosage levels of the medicaments may be increased thereby increasing the therapeutic effect of the medicaments. For example, the therapeutic compositions may be used in topical cytoprotective pharmaceutical compositions in combination with cytotoxic medicaments such as epithelial cell cohesiveness reducers such as tretinoin (Retin A), dermatological abrasants, and anti-inflammatory, to protect and enhance the resuscitation rate of the injured mammalian cells. The therapeutic compositions may also be used in ingestible cytoprotective pharmaceutical compositions in combination with medicaments that cause cytotoxic side effects such as anti-tumor, anti-viral, and antibacterial medicaments including the lipid regulating agents gemfibrozil and lovastatin, centrally acting anticholinesterases such as tacrine, chemotherapeutic medicaments such as the anthracycline antibiotic doxorubicin, gastric irritants such as acetylsalicylic acid and ibuprofen, to protect and enhance the resuscitation rate of the injured mammalian cells.

35

In a specific embodiment, the invention is directed at a cytoprotective pharmaceutical composition

SUBSTITUTE SHEET

2104461

- 39 -

for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises:

5 (A) a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

20 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

25 (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

30 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a medicament having cytotoxic properties.

SUBSTITUTE SHEET

2104461 - 40 -

In a preferred embodiment, the medicament having cytotoxic properties in the cytoprotective pharmaceutical compositions is selected from the group consisting of gemfibrozil, lovastatin, tacrine, and doxorubicin. In a more preferred embodiment, the medicament having cytotoxic properties in the cytoprotective pharmaceutical compositions is selected from the group consisting of gemfibrozil and tacrine.

In a most preferred embodiment, the medicament having cytotoxic properties is gemfibrozil. Gemfibrozil (LOPID) is a lipid regulating agent which lowers elevated serum lipids primarily by decreasing serum triglyceride with a variable reduction in total serum cholesterol. Lovastatin (MEVACOR) is a cholesterol lowering agent which inhibits the enzymatic biosynthesis of cholesterol. Tacrine (1,2,3,4-tetrahydro-9-acridinamine) is a centrally active anticholinesterase useful as a cognition activator. Tacrine has undergone clinical trials for use in treating severe Alzheimer's disease (presenile dementia). Doxorubicin (adriamycin) is a cytotoxic anthracycline antibiotic reported to produce regression in disseminated neoplastic conditions such as in various leukemias, tumors, neuroblastomas, sarcomas, and carcinomas.

In a fourth embodiment, the therapeutic compositions, without the mixture of saturated and unsaturated fatty acids, may be combined with a medicament, which is cytotoxic to cells, to form a prophylactic cytoprotective pharmaceutical composition. These prophylactic compositions may be used to protect mammalian cells, which have not been injured by the medicament having cytotoxic properties and which do not require resuscitation, and to thereby increase the therapeutic effect of the cytotoxic medicament.

SUBSTITUTE SHEET

2104461

- 41 -

In a specific embodiment, the invention is directed at a prophylactic cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises:

(A) a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

(B) a medicament having cytotoxic properties.

25

In a preferred embodiment, the therapeutic composition in the prophylactic cytoprotective pharmaceutical composition is selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant; and

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

35

2104461

- 42 -

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof.

5 In a more preferred embodiment, the therapeutic composition in the prophylactic cytoprotective pharmaceutical composition comprises:

 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
10 salts of pyruvic acid, and mixtures thereof; and
 (b) an antioxidant.

 In another preferred embodiment, the therapeutic composition in the prophylactic
15 cytoprotective pharmaceutical composition comprises:

 (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and
 (b) an antioxidant.

20

 In another form of the fourth embodiment, the therapeutic compositions of the present invention, without the mixture of saturated and unsaturated fatty acids, may be combined in an immediate release form
25 with an anticancer medicament having cytotoxic properties in a timed release form to provide a timed release prophylactic cytoprotective pharmaceutical composition. In this embodiment, the timed release composition releases the therapeutic composition
30 substantially immediately and releases the cytotoxic chemotherapeutic medicament after a suitable period of time, for example 15 minutes after releasing the therapeutic composition, to selectively protect non-cancerous cells in the presence of cancerous cells
35 against the cytotoxic chemotherapeutic medicament. Cancer cells, unlike normal cells or benign tumor cells, exhibit the properties of invasion and

2104461

- 43 -

metastasis and are highly anaplastic. Because cancerous cells have a rapid metabolism, cancerous cells will rapidly consume the protective therapeutic composition and will not be protected by the therapeutic compositions when the chemotherapeutic medicament is subsequently released. Non-cancerous cells which do not have such a rapid metabolism will not rapidly consume the therapeutic compositions and will be protected when the chemotherapeutic medicament is subsequently released.

In a specific embodiment, the invention is directed at a timed release prophylactic cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises:

(A) a therapeutic composition in an immediate release form selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(3) (a) an antioxidant; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

(B) an anticancer medicament having cytotoxic properties in timed release form;

SUBSTITUTE SHEET

2104461

- 44 -

wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have
5 substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition.

A suitable or sufficient period of time is
10 that period of time wherein the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition. The period of
15 time should not be so long that the non-cancerous cells substantially metabolize the therapeutic composition and are unprotected. The exact time is subject to such factors as the type and quantity of therapeutic composition employed, the medicament having cytotoxic properties used, and the type of cancerous cells and
20 non-cancerous cells being treated. Thus, the period of time may be varied in order to obtain the result desired and such variations are within the capabilities of those skilled in the art without the need for undue experimentation.

25 In an alternative embodiment, the timed release prophylactic cytoprotective pharmaceutical composition may further comprise a mixture of saturated and unsaturated fatty acids in the therapeutic
30 composition in immediate release form wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

In a fifth embodiment, therapeutic
35 compositions comprising pyruvate may be combined with a medicament, which is cytotoxic to cells, to form a preventative cytoprotective pharmaceutical composition.

SUBSTITUTE SHEET

- 45 -

These preventative compositions may be used to protect mammalian cells, which have not been injured by the medicament having cytotoxic properties and which do not require resuscitation, and to thereby increase the therapeutic effect of the cytotoxic medicament.

In a specific embodiment, the invention is directed at a preventative cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises:

(A) a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) a medicament having cytotoxic properties.

In another form of the fifth embodiment, therapeutic compositions comprising pyruvate may be combined in an immediate release form with an anticancer medicament having cytotoxic properties in a timed release form to provide a timed release preventative cytoprotective pharmaceutical composition. In this embodiment, the timed release composition releases the therapeutic composition substantially immediately and releases the cytotoxic chemotherapeutic medicament after a suitable period of time, for example 15 minutes after releasing the therapeutic composition, to selectively protect non-cancerous cells in the presence of cancerous cells against the cytotoxic chemotherapeutic medicament.

In a specific embodiment, the invention is direct at a timed release preventative cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian cells in the presence of

2104461

- 46 -

cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises:

5 (A) a therapeutic composition in an immediate release form comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) an anticancer medicament having cytotoxic properties in timed release form;

10 wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition
15 and the non-cancerous cells have not substantially metabolized the therapeutic composition.

In a sixth embodiment, the components in the therapeutic compositions may be utilized in wound
20 healing products to increase the resuscitation rate of injured mammalian cells. In this embodiment, the mixture of saturated and unsaturated fatty acids is employed with another component in the therapeutic compositions. Not all components of the therapeutic
25 composition may be required in the wound healing product to heal a wound. For example, if the wound is not undergoing oxidative injury, the therapeutic composition may not require an antioxidant. For other types of wounds, the components in the therapeutic
30 composition may vary depending upon the type of wound being treated.

In a specific embodiment, the invention is directed at a wound healing pharmaceutical composition
35 to increase the resuscitation rate of injured mammalian cells which comprises:

SUBSTITUTE SHEET

2104461

- 47 -

(A) a first wound healing component comprising a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

5 (B) a second wound healing component selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

10 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) an antioxidant.

15 In a preferred embodiment, the second wound healing component in the wound healing pharmaceutical composition is selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

20 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof.

25 In a more preferred embodiment, the second wound healing component in the wound healing pharmaceutical composition comprises:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof.

30 The present invention extends to methods for making the augmented pharmaceutical compositions, cytoprotective pharmaceutical compositions, prophylactic cytoprotective pharmaceutical compositions, timed release prophylactic cytoprotective pharmaceutical compositions, preventative

SUBSTITUTE SHEET

2104461

- 48 -

cytoprotective pharmaceutical compositions, timed
release preventative cytoprotective pharmaceutical
compositions, and wound healing pharmaceutical
compositions. In general, the pharmaceutical
5 compositions are made by forming an admixture of the
therapeutic components in the compositions. For
example, an augmented pharmaceutical composition is
made by forming an admixture of the therapeutic
compositions and the medicament which is useful for
10 treating injured mammalian cells. A cytoprotective
pharmaceutical composition is made by forming an
admixture of the therapeutic compositions and the
medicament having cytotoxic properties. A prophylactic
cytoprotective pharmaceutical composition, including
15 the timed release form, is made by forming an admixture
of the prophylactic therapeutic components and the
medicament having cytotoxic properties. A preventative
cytoprotective pharmaceutical composition, including
the timed release form, is made by forming an admixture
20 of the preventative therapeutic components and the
medicament having cytotoxic properties. A wound
healing pharmaceutical composition is made by forming
an admixture of the wound healing components.

25 The present invention also extends to methods
for employing the augmented pharmaceutical
compositions, the cytoprotective pharmaceutical
compositions, the prophylactic cytoprotective
pharmaceutical compositions, the timed release
30 prophylactic cytoprotective pharmaceutical
compositions, the preventative cytoprotective
pharmaceutical compositions, the timed release
preventative cytoprotective pharmaceutical
compositions, and the wound healing pharmaceutical
35 compositions.

SUBSTITUTE SHEET

- 49 -

In general, a pharmaceutical composition is employed by contacting the pharmaceutical composition with mammalian cells. In a specific embodiment, the invention is directed at a method for augmenting the therapeutic effect of a medicament useful for treating injured mammalian cells which comprises the steps of:

(A) providing a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are

SUBSTITUTE SHEET

2104461 - 50 -

those fatty acids required for the resuscitation of injured mammalian cells;

(B) providing a medicament useful for treating injured cells; and

5 (C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to augment the therapeutic effect of the medicament useful for treating injured mammalian cells.

10

In another specific embodiment, the invention is directed at a method for protecting mammalian cells from a medicament having cytotoxic properties which comprises the steps of:

15 (A) providing a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

20

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

25

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

30

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

35

(3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are

SUBSTITUTE SHEET

- 51 -

those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(B) providing a medicament having cytotoxic properties; and

15 (C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells from the medicament having cytotoxic properties.

20 In another embodiment, the invention is directed at a method for protecting mammalian cells prophylactically from a medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition selected from the group consisting of:

25 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

30 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant; and

35 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

SUBSTITUTE SHEET

2104461

- 52 -

(b) an antioxidant; and

(B) providing a medicament having cytotoxic properties; and

5 (C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells prophylactically from the medicament having cytotoxic properties.

10 In another embodiment, the invention is directed at a method for prophylactically protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps
15 of:

(A) providing a therapeutic composition in an immediate release form selected from the group consisting of:

20 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and
25

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(3) (a) an antioxidant; and

30 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant;

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and
35

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently

SUBSTITUTE SHEET

- 53 -

to mammalian cells to selectively protect non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties;

5 wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition
10 and the non-cancerous cells have not substantially metabolized the therapeutic composition.

 The therapeutic compositions of the present invention may also be administered to cells prior to
15 the administration of a cytotoxic chemotherapeutic medicament, for example 15 minutes prior to administering the chemotherapeutic medicament, to selectively protect non-cancerous cells in the presence of cancerous cells against a cytotoxic chemotherapeutic
20 medicament. Because cancerous cells have a rapid metabolism, cancerous cells will rapidly consume the protective therapeutic composition and will not be protected by the therapeutic compositions when the chemotherapeutic medicament is subsequently
25 administered.

 In a specific embodiment, the invention is direct at a method for selectively protecting non-cancerous mammalian cells prophylactically in the
30 presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

(A) administering to mammalian cells a therapeutic composition to prevent and reduce injury to the
35 mammalian cells selected from the group consisting of:

SUBSTITUTE SHEET

2104461

- 54 -

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

5 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
10 salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant;

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

15 (b) an antioxidant; and

(B) waiting a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic
20 composition; and

(C) administering the cytotoxic anticancer medicament to the mammalian cells to treat the cancerous cells which are unprotected by the therapeutic composition and the non-cancerous cells
25 which are protected by the therapeutic composition to thereby increase the therapeutic effect of the anticancer medicament.

A suitable period of time is that period of
30 time wherein the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition. The period of time should not be so long that the non-cancerous cells substantially
35 metabolize the therapeutic composition and are unprotected. The exact time is subject to such factors as the type and quantity of therapeutic composition

SUBSTITUTE SHEET

- 55 -

employed, the medicament having cytotoxic properties used, and the type of cancerous cells and non-cancerous cells being treated. Thus, the period of time may be varied in order to obtain the result desired and such variations are within the capabilities of those skilled in the art without the need for undue experimentation.

In another embodiment, the invention is directed at a method for protecting mammalian cells preventatively from a medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) providing a medicament having cytotoxic properties; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells preventatively from the medicament having cytotoxic properties.

In another embodiment, the invention is directed at a method for preventatively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition in an immediate release comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently

SUBSTITUTE SHEET

2104461

- 56 -

to mammalian cells to selectively protect non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties;

5 wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition
10 and the non-cancerous cells have not substantially metabolized the therapeutic composition.

 In another embodiment, the invention is directed at a method for selectively protecting non-
15 cancerous mammalian cells preventatively in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

 (A) administering to mammalian cells a therapeutic
20 composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

 (B) waiting a period of time sufficient such that the cancerous cells have substantially metabolized the
25 therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition; and

 (C) administering the cytotoxic anticancer medicament to the mammalian cells to treat the
30 cancerous cells which are unprotected by the therapeutic composition and the non-cancerous cells which are protected by the therapeutic composition to thereby increase the therapeutic effect of the anticancer medicament.

35

 In another embodiment, the invention is directed at a method for increasing the healing rate of

- 57 -

a wound. In this embodiment, not all components of the therapeutic composition may be required to heal the wound. For example, if the wound is not undergoing oxidative injury, the therapeutic composition may not
5 require an antioxidant. For other wounds, the components in the therapeutic composition may vary depending upon the type of wound being treated. In a specific embodiment, the invention is directed at a method for increasing the healing rate of a wound in a
10 mammal which comprises:

(A) providing a wound healing component selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
15 salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(c) an antioxidant;

20 (d) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(e) a combination of pyruvate selected from
25 the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

30 (f) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and an antioxidant;

(g) a combination of pyruvate selected from
35 the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids

SUBSTITUTE SHEET

2104461

- 58 -

wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (h) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and an antioxidant;

10 (i) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

15 (j) a combination of an antioxidant and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(B) contacting the wound healing component from step (A) with the wound to increase the healing rate of the wound.

20

In a preferred embodiment, the wound healing component may be selected from the group consisting of:

25 (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(c) an antioxidant; and

30 (d) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

35 In another preferred embodiment, the wound healing component may be selected from the group consisting of:

SUBSTITUTE SHEET

- 59 -

(e) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and lactate selected from the group consisting of
5 lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(f) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof,
10 and an antioxidant;

(g) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids
15 wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(h) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof,
20 and an antioxidant;

(i) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids
25 wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(j) a combination of an antioxidant and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required
30 for the resuscitation of injured mammalian cells.

In another preferred embodiment, the wound healing component may be selected from the group consisting of:

35 (g) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof,

SUBSTITUTE SHEET

2104461

- 60 -

and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(i) a combination of lactate selected from
5 the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

10

The types of wounds which may be healed using the compositions of the present invention are those which result from an injury which causes epidermal damage, chronic ulcers, gastric ulcers, burns, and
15 donor site wounds. Such wounds include ophthalmic wounds, such as those which result from corneal ulcers, radialkeratotomy, corneal transplants, epikeratophakia and other surgically induced wounds in the eye, and cutaneous wounds such as burn wounds, donor site wounds
20 from skin transplants and ulcers (cutaneous, decubitis, venous stasis, and diabetic). In addition, dermatological wounds such as psoriasis, sunburn, and skin rashes may also be treated with the compositions of the present invention. The compositions may be
25 applied to the wound site either topically or internally depending on the type of wound.

Methods for increasing the rate of wound healing comprise contacting the wound healing
30 composition with the wound to increase the healing rate of the wound. Preferably, the method comprises topically administering the compositions of the present invention directly to a wound site. The composition is maintained in contact with the wound for a period of
35 time sufficient to increase the rate of cell growth at the wound site.

SUBSTITUTE SHEET

2104461

- 61 -

In another embodiment, the invention is directed at a method for preserving mammalian cells in a culture medium which comprises the steps of:

5 (A) providing a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

20 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

25 (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

30 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

SUBSTITUTE SHEET

2104461

- 62 -

(B) providing mammalian cells in a culture medium;
and

(C) contacting the therapeutic composition from
step (A) with the mammalian cells in the culture medium
5 from step (B).

In another embodiment, the invention is
directed at a cytoprotective composition for reversing
the cytotoxic effects of ascorbic acid on mammalian
10 cells which comprises ascorbic acid and a
therapeutically cytoprotective effective amount of
pyruvate. The invention also extends to a method for
reversing the cytotoxic effects of ascorbic acid on
mammalian cells which comprises administering to the
15 mammalian cells ascorbic acid and a therapeutically
cytoprotective effective amount of pyruvate. A
therapeutically effective amount of pyruvate is that
amount of pyruvate which will reverse the cytotoxic
effects of ascorbic acid on mammalian cells. In
20 general, ascorbic acid and pyruvate are present in the
cytoprotective composition in a ratio by weight from
about 1:10 to about 10:1, preferably from about 1:5 to
about 5:1, and more preferably from about 1:2 to about
2:1, respectively.

25 Once prepared, the inventive therapeutic
compositions may be stored for future use or may be
formulated in effective amounts with pharmaceutically
acceptable carriers to prepare a wide variety of
30 pharmaceutical compositions. Examples of
pharmaceutically acceptable carriers are pharmaceutical
appliances, topical vehicles (non-oral and oral), and
ingestible vehicles.

35 Examples of pharmaceutical appliances are
sutures, staples, gauze, bandages, burn dressings,
artificial skins, liposome or micell formulations,

- 63 -

microcapsules, aqueous vehicles for soaking gauze dressings, and the like, and mixtures thereof. Non-oral topical compositions employ non-oral topical vehicles, such as creams, gels formulations, foams, ointments and sprays, salves, and films, which are intended to be applied to the skin or body cavity and are not intended to be taken by mouth. Oral topical compositions employ oral vehicles, such as mouthwashes, rinses, oral sprays, suspensions, and dental gels, which are intended to be taken by mouth but are not intended to be ingested. Ingestible compositions employ ingestible or partly ingestible vehicles such as confectionery bulking agents which include hard and soft confectionery such as lozenges, tablets, toffees, nougats, suspensions, chewy candies, and chewing gums.

In one form of the invention, the therapeutic composition is incorporated into a pharmaceutical appliance which may be in the form of sutures, staples, gauze, bandages, burn dressings, artificial skins, liposome or micell formulations, microcapsules, aqueous vehicles for soaking gauze dressings, and the like, and mixtures thereof. A variety of traditional ingredients may optionally be included in the pharmaceutical composition in effective amounts such as buffers, preservatives, tonicity adjusting agents, antioxidants, polymers for adjusting viscosity or for use as extenders, and excipients, and the like. Specific illustrative examples of such traditional ingredients include acetate and borate buffers; thimerosal, sorbic acid, methyl and propyl paraben and chlorobutanol preservatives; sodium chloride and sugars to adjust the tonicity; and excipients such as mannitol, lactose and sucrose. Other conventional pharmaceutical additives known to those having ordinary skill in the pharmaceutical arts may also be used in the pharmaceutical composition.

2104461

- 64 -

In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be employed
5 in the pharmaceutical appliance. These amounts are readily determined by those skilled in the art without the need for undue experimentation. The exact amount of the therapeutic composition employed is subject to such factors as the type and concentration of the
10 therapeutic composition and the type of pharmaceutical appliance employed. Thus, the amount of therapeutic composition may be varied in order to obtain the result desired in the final product and such variations are within the capabilities of those skilled in the art
15 without the need for undue experimentation. In a preferred embodiment, the pharmaceutical composition will comprise the therapeutic composition in an amount from about 0.1% to about 5%, by weight of the pharmaceutical composition. In a more preferred
20 embodiment, the pharmaceutical composition will comprise the therapeutic composition in an amount from about 0.1% to about 3%, by weight of the pharmaceutical composition. In a most preferred embodiment, the pharmaceutical composition will comprise the
25 therapeutic composition in an amount from about 0.1% to about 1%, by weight of the pharmaceutical composition.

The present invention extends to methods for making the pharmaceutical compositions. In general, a
30 pharmaceutical composition is made by contacting a therapeutically effective amount of a therapeutic composition with a pharmaceutical appliance and the other ingredients of the final desired pharmaceutical composition. The therapeutic composition may be in a
35 solvent and may be absorbed onto a pharmaceutical appliance.

2104461

- 65 -

Other ingredients will usually be incorporated into the composition as dictated by the nature of the desired composition as well known by those having ordinary skill in the art. The ultimate
5 pharmaceutical compositions are readily prepared using methods generally known in the pharmaceutical arts.

In another form of the invention, the therapeutic composition is incorporated into a non-oral
10 topical vehicle which may be in the form of a cream, gel, foam, ointment, spray, and the like. Typical non-toxic non-oral topical vehicles known in the pharmaceutical arts may be used in the present invention. The preferred non-oral topical vehicles are
15 water and pharmaceutically acceptable water-miscible organic solvents such as ethyl alcohol, isopropyl alcohol, propylene glycol, glycerin, and the like, and mixtures of these solvents. Water-alcohol mixtures are particularly preferred and are generally employed in a
20 weight ratio from about 1:1 to about 20:1, preferably from about 3:1 to about 20:1, and most preferably from about 3:1 to about 10:1, respectively.

The non-oral topical therapeutic compositions
25 may also contain conventional additives employed in those products. Conventional additives include humectants, emollients, lubricants, stabilizers, dyes, and perfumes, providing the additives do not interfere with the therapeutic properties of the therapeutic
30 composition.

Suitable humectants useful in the non-oral topical therapeutic compositions include glycerin, propylene glycol, polyethylene glycol, sorbitan,
35 fructose, and the like, and mixtures thereof. Humectants, when employed, may be present in amounts

SUBSTITUTE SHEET

2104461

- 66 -

from about 10% to about 20%, by weight of the topical therapeutic composition.

5 The coloring agents (colors, colorants) useful in the non-oral topical therapeutic composition are used in amounts effective to produce the desired color. These coloring agents include pigments which may be incorporated in amounts up to about 6% by weight of the non-oral topical therapeutic composition. A
10 preferred pigment, titanium dioxide, may be incorporated in amounts up to about 2%, and preferably less than about 1%, by weight of the non-oral topical therapeutic composition. The coloring agents may also include natural food colors and dyes suitable for food,
15 drug and cosmetic applications. These coloring agents are known as F.D. & C. dyes and lakes. The materials acceptable for the foregoing uses are preferably water-soluble. Illustrative nonlimiting examples include the indigoid dye known as F.D. & C. Blue No. 2, which is the
20 disodium salt of 5,5-indigotindisulfonic acid. Similarly, the dye known as F.D. & C. Green No. 1 comprises a triphenylmethane dye and is the monosodium salt of 4-[4-(N-ethyl-p-sulfoniumbenzylamino) diphenylmethylene]-[1-(N-ethyl-N-p-sulfoniumbenzyl)-
25 delta-2,5-cyclohexadieneimine]. A full recitation of all F.D. & C. coloring agents and their corresponding chemical structures may be found in the Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Edition, in volume 5 at pages 857-884, which text is incorporated
30 herein by reference.

 In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be admixed
35 with a non-oral topical vehicle to form a topical therapeutic composition. These amounts are readily determined by those skilled in the art without the need

SUBSTITUTE SHEET

- 67 -

for undue experimentation. In a preferred embodiment, the non-oral topical therapeutic compositions will comprise the therapeutic composition in an amount from about 0.1% to about 10% and a non-oral topical vehicle in a quantity sufficient to bring the total amount of composition to 100%, by weight of the non-oral topical therapeutic composition. In a more preferred embodiment, the non-oral topical therapeutic compositions will comprise the therapeutic composition in an amount from about 0.1% to about 5%, and in a most preferred embodiment, the non-oral topical therapeutic compositions will comprise the therapeutic composition in an amount from about 0.1% to about 2%, and a non-oral topical vehicle in a quantity sufficient to bring the total amount of composition to 100%, by weight of the non-oral topical therapeutic composition.

The present invention extends to methods for preparing the non-oral topical therapeutic compositions. In such a method, the non-oral topical therapeutic composition is prepared by admixing a therapeutically effective amount of the therapeutic composition of the present invention and a non-oral topical vehicle. The final compositions are readily prepared using standard methods and apparatus generally known by those skilled in the pharmaceutical arts. The apparatus useful in accordance with the present invention comprises mixing apparatus well known in the pharmaceutical arts, and therefore the selection of the specific apparatus will be apparent to the artisan.

In another form of the invention, the therapeutic composition is incorporated into an oral topical vehicle which may be in the form of a mouthwash, rinse, oral spray, suspension, dental gel, and the like. Typical non-toxic oral vehicles known in the pharmaceutical arts may be used in the present

2104461

- 68 -

invention. The preferred oral vehicles are water, ethanol, and water-ethanol mixtures. The water-ethanol mixtures are generally employed in a weight ratio from about 1:1 to about 20:1, preferably from about 3:1 to about 20:1, and most preferably from about 3:1 to about 10:1, respectively. The pH value of the oral vehicle is generally from about 4 to about 7, and preferably from about 5 to about 6.5. An oral topical vehicle having a pH value below about 4 is generally irritating to the oral cavity and an oral vehicle having a pH value greater than about 7 generally results in an unpleasant mouth feel.

The oral topical therapeutic compositions may also contain conventional additives normally employed in those products. Conventional additives include a fluorine providing compound, a sweetening agent, a flavoring agent, a coloring agent, a humectant, a buffer, and an emulsifier, providing the additives do not interfere with the therapeutic properties of the therapeutic composition.

The coloring agents and humectants, and the amounts of these additives to be employed, set out above as useful in the non-oral topical therapeutic composition may be used in the oral topical therapeutic composition.

Fluorine providing compounds may be fully or slightly water soluble and are characterized by their ability to release fluoride ions or fluoride containing ions in water and by their lack of reaction with other components in the composition. Typical fluorine providing compounds are inorganic fluoride salts such as water-soluble alkali metal, alkaline earth metal, and heavy metal salts, for example, sodium fluoride, potassium fluoride, ammonium fluoride, cuprous

- 69 -

fluoride, zinc fluoride, stannic fluoride, stannous fluoride, barium fluoride, sodium fluorosilicate, ammonium fluorosilicate, sodium fluorozirconate, sodium monofluorophosphate, aluminum mono- and di-
5 fluorophosphates and fluorinated sodium calcium pyrophosphate. Alkali metal fluorides, tin fluoride and monofluorophosphates, such as sodium and stannous fluoride, sodium monofluorophosphate and mixtures thereof, are preferred.

10

The amount of fluorine providing compound present in the present oral topical therapeutic composition is dependent upon the type of fluorine providing compound employed, the solubility of the
15 fluorine compound, and the nature of the final oral therapeutic composition. The amount of fluorine providing compound used must be a nontoxic amount. In general, the fluorine providing compound when used will be present in an amount up to about 1%, preferably from
20 about 0.001% to about 0.1%, and most preferably from about 0.001% to about 0.05%, by weight of the oral topical therapeutic composition.

When sweetening agents (sweeteners) are used,
25 those sweeteners well known in the art, including both natural and artificial sweeteners, may be employed. The sweetening agent used may be selected from a wide range of materials including water-soluble sweetening agents, water-soluble artificial sweetening agents,
30 water-soluble sweetening agents derived from naturally occurring water-soluble sweetening agents, dipeptide based sweetening agents, and protein based sweetening agents, including mixtures thereof. Without being limited to particular sweetening agents, representative
35 categories and examples include:

(a) water-soluble sweetening agents such as monosaccharides, disaccharides and polysaccharides such

2104461

- 70 -

as xylose, ribose, glucose (dextrose), mannose, galactose, fructose (levulose), sucrose (sugar), maltose, invert sugar (a mixture of fructose and glucose derived from sucrose), partially hydrolyzed starch, corn syrup solids, dihydrochalcones, monellin, steviosides, and glycyrrhizin, and mixtures thereof;

(b) water-soluble artificial sweeteners such as soluble saccharin salts, i.e., sodium or calcium saccharin salts, cyclamate salts, the sodium, ammonium or calcium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide, the potassium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide (Acesulfame-K), the free acid form of saccharin, and the like;

(c) dipeptide based sweeteners, such as L-aspartic acid derived sweeteners, such as L-aspartyl-L-phenylalanine methyl ester (Aspartame) and materials described in United States patent no. 3,492,131, L-alpha-aspartyl-N-(2,2,4,4-tetramethyl-3-thietanyl)-D-alanin-amide hydrate (Alitame), methyl esters of L-aspartyl-L-phenylglycerine and L-aspartyl-L-2,5-dihydrophenyl-glycine, L-aspartyl-2,5-dihydro-L-phenylalanine; L-aspartyl-L-(1-cyclohexen)-alanine, and the like;

(d) water-soluble sweeteners derived from naturally occurring water-soluble sweeteners, such as chlorinated derivatives of ordinary sugar (sucrose), e.g., chlorodeoxysugar derivatives such as derivatives of chlorodeoxysucrose or chlorodeoxygalactosucrose, known, for example, under the product designation of Sucralose; examples of chlorodeoxysucrose and chlorodeoxygalacto-sucrose derivatives include but are not limited to: 1-chloro-1'-deoxysucrose; 4-chloro-4-deoxy-alpha-D-galacto-pyranosyl-alpha-D-fructofuranoside, or 4-chloro-4-deoxygalactosucrose; 4-chloro-4-deoxy-alpha-D-galacto-pyranosyl-1-chloro-1-deoxy-beta-D-fructo-furanoside, or 4,1'-dichloro-4,1'-

SUBSTITUTE SHEET

2104461

- 71 -

dideoxygalactosucrose; 1',6'-dichloro-1',6'-
dideoxysucrose; 4-chloro-4-deoxy-alpha-D-galacto-
pyranosyl-1,6-dichloro-1,6-dideoxy-beta-D-fructo-
furanoside, or 4,1',6'-trichloro-4,1',6'-
5 trideoxygalacto-sucrose; 4,6-dichloro-4,6-dideoxy-
alpha-D-galacto-pyranosyl-6-chloro-6-deoxy-beta-D-
fructofuranoside, or 4,6,6'-trichloro-4,6,6'-
trideoxygalactosucrose; 6,1',6'-trichloro-6,1',6'-
trideoxysucrose; 4,6-dichloro-4,6-dideoxy-alpha-D-
10 galacto-pyranosyl-1,6-dichloro-1,6-di-deoxy-beta-D-
fructofuranoside, or 4,6,1',6'-tetrachloro-4,6,1',6'-
tetraideoxygalacto-sucrose; and 4,6,1',6'-tetrachloro-
4,6,1',6'-tetraideoxy-sucrose; and
(e) protein based sweeteners such as
15 thaumaococcus danielli (Thaumatococcus daniellii).

In general, an effective amount of sweetening
agent is utilized to provide the level of sweetness
desired in the particular oral topical therapeutic
20 composition, and this amount will vary with the
sweetener selected and the final oral therapeutic
product desired. The amount of sweetener normally
present is in the range from about 0.0025% to about
90%, by weight of the oral topical therapeutic
25 composition, depending upon the sweetener used. The
exact range of amounts for each type of sweetener is
well known in the art and is not the subject of the
present invention.

30 The flavoring agents (flavors, flavorants)
which may be used include those flavors known to the
skilled artisan, such as natural and artificial
flavors. Suitable flavoring agents include mints, such
as peppermint, citrus flavors such as orange and lemon,
35 artificial vanilla, cinnamon, various fruit flavors,
both individual and mixed, and the like.

2104461

- 72 -

5 The amount of flavoring agent employed in the oral topical therapeutic composition is normally a matter of preference subject to such factors as the type of final oral therapeutic composition, the individual flavor employed, and the strength of flavor desired. Thus, the amount of flavoring may be varied in order to obtain the result desired in the final product and such variations are within the capabilities of those skilled in the art without the need for undue experimentation. The flavoring agents, when used, are generally utilized in amounts that may, for example, range in amounts from about 0.05% to about 6%, by weight of the oral topical therapeutic composition.

15 Sutable buffer solutions useful in the non-oral topical therapeutic compositions include citric acid-sodium citrate solution, phosphoric acid-sodium phosphate solution, and acetic acid-sodium acetate solution in amounts up to about 1%, and preferably from about 0.05% to about 0.5% by weight of the oral topical therapeutic composition.

25 In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be admixed with an oral topical vehicle to form a topical therapeutic composition. These amounts are readily determined by those skilled in the art without the need for undue experimentation. In a preferred embodiment, the oral topical therapeutic compositions will comprise the therapeutic composition in an amount from about 0.1% to about 10% and a oral topical vehicle in a quantity sufficient to bring the total amount of composition to 100%, by weight of the oral topical therapeutic composition. In a more preferred embodiment, the oral topical therapeutic compositions will comprise the therapeutic composition in an amount

2104461

- 73 -

from about 0.1% to about 5%, and in a most preferred embodiment, the oral topical therapeutic compositions will comprise the therapeutic composition in an amount from about 0.1% to about 2%, and a oral topical vehicle in a quantity sufficient to bring the total amount of composition to 100%, by weight of the oral topical therapeutic composition.

The present invention extends to methods for preparing the oral topical therapeutic compositions. In such a method, the oral topical therapeutic composition is prepared by admixing a therapeutically effective amount of the therapeutic composition of the present invention and an oral topical vehicle. The final compositions are readily prepared using standard methods and apparatus generally known by those skilled in the pharmaceutical arts. The apparatus useful in accordance with the present invention comprises mixing apparatus well known in the pharmaceutical arts, and therefore the selection of the specific apparatus will be apparent to the artisan.

In a preferred embodiment, an oral topical therapeutic composition is made by first dissolving coloring agents, sweetening agents, and similar additives in water. The therapeutic composition is then admixed with the aqueous solution. Then sufficient water or ethanol, or mixtures of water and ethanol, are added to the solution with mixing until the final solution volume is reached. In a more preferred embodiment, the therapeutic composition is added to the solution as the final ingredient. The final oral topical therapeutic compositions are readily prepared using methods generally known in the pharmaceutical arts.

SUBSTITUTE SHEET

2104461

- 74 -

The oral therapeutic composition may also be in the form of dental gel. As used herein, the term "gel" means a solid or semisolid colloid which contains considerable quantities of water. The colloid particles in a gel are linked together in a coherent meshwork which immobilizes the water contained inside the meshwork.

The dental gel compositions of the present invention may contain the conventional additives set out above for oral topical therapeutic compositions such as mouthwashes, rinses, oral sprays, and suspensions and, in addition, may contain additional additives such as a polishing agent, a desensitizing agent, and the like, providing the additional additives do not interfere with the therapeutic properties of the therapeutic composition.

In a dental gel composition, the oral vehicle generally comprises water, typically in an amount from about 10% to about 90%, by weight of the dental gel composition. Polyethylene glycol, propylene glycol, glycerin, and mixtures thereof may also be present in the vehicle as humectants or binders in amounts from about 18% to about 30%, by weight of the dental gel composition. Particularly preferred oral vehicles comprise mixtures of water with polyethylene glycol or water with glycerin and polypropylene glycol.

The dental gels of the present invention include a gelling agent (thickening agent) such as a natural or synthetic gum or gelatin. Gelling agents such as hydroxyethyl cellulose, methyl cellulose, glycerin, carboxypolymethylene, and gelatin and the like, and mixtures thereof may be used. The preferred gelling agent is hydroxyethyl cellulose. Gelling agents may be used in amounts from about 0.5% to about

SUBSTITUTE SHEET

2104461

- 75 -

5%, and preferably from about 0.5% to about 2%, by weight of the dental gel composition.

5 The dental gel compositions of the present invention may also include a polishing agent. In clear gels, a polishing agent of colloidal silica and/or alkali metal aluminosilicate complexes is preferred since these materials have refractive indices close to the refractive indices of the gelling systems commonly
10 used in dental gels. In non-clear gels, a polishing agent of calcium carbonate or calcium dihydrate may be used. These polishing agents may be used in amounts up to about 75%, and preferably in amounts up to about 50%, by weight of the dental gel composition.

15

The dental gel may also contain a desensitizing agent such as a combination of citric acid and sodium citrate. Citric acid may be used in an amount from about 0.1% to about 3%, and preferably from
20 about 0.2% to about 1%, by weight, and sodium citrate may be used in an amount from about 0.3% to about 9%, and preferably from about 0.6% to about 3%, by weight of the dental gel composition.

25

In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be admixed into the dental gel compositions. These amounts are readily determined by those skilled in the art without
30 the need for undue experimentation. In a preferred embodiment, the dental gel compositions will comprise the therapeutic composition in an amount from about 0.1% to about 10% and an oral topical vehicle in a quantity sufficient to bring the total amount of
35 composition to 100%, by weight of the dental gel composition. In a more preferred embodiment, the dental gel compositions will comprise the therapeutic

2104461

- 76 -

composition in an amount from about 0.1% to about 5%,
and in a most preferred embodiment, the dental gel
compositions will comprise the therapeutic composition
in an amount from about 0.1% to about 2%, and an oral
5 topical vehicle in a quantity sufficient to bring the
total amount of composition to 100%, by weight of the
dental gel composition.

The present invention extends to methods for
10 preparing the therapeutic dental gel compositions. In
such a method, the dental gel composition is prepared
by admixing a therapeutically effective amount of the
therapeutic composition of the present invention and an
oral topical vehicle. The final compositions are
15 readily prepared using methods generally known by those
skilled in the dental and pharmaceutical arts. The
apparatus useful in accordance with the present
invention comprises mixing apparatus well known in the
pharmaceutical arts, and therefore the selection of the
20 specific apparatus will be apparent to the artisan.

In a preferred embodiment, a therapeutic
dental gel composition is made by first dispersing a
gelling agent in a humectant or water, or a mixture of
25 both, then admixing to the dispersion an aqueous
solution of the water-soluble additives such as the
fluorine providing compound, sweeteners and the like,
then adding the polishing agent, and lastly admixing
the flavoring agent and the therapeutic composition.
30 The final gel mixture is then tubed or otherwise
packaged. The liquids and solids in a gel product are
proportioned to form a creamy or gelled mass which is
extrudable from a pressurized container or from a
collapsible tube. The final therapeutic compositions
35 are readily prepared using methods generally known in
the pharmaceutical arts.

SUBSTITUTE SHEET

- 77 -

In yet another form of the invention, the therapeutic composition is incorporated into an ingestible vehicle. The ingestible vehicle may be a confectionery bulking agent in the form of lozenges, tablets, toffees, nougats, suspensions, chewy candies, chewing gums, and the like. The pharmaceutically acceptable carriers may be prepared from a wide range of materials including, but not limited to, diluents, binders and adhesives, lubricants, disintegrants, coloring agents, bulking agents, flavoring agents, sweetening agents and miscellaneous materials such as buffers and adsorbents that may be needed in order to prepare a particular therapeutic confection.

The preparation of confectionery formulations is historically well known and has changed little through the years. Confectionery items have been classified as either "hard" confectionery or "soft" confectionery. The therapeutic compositions of the present invention can be incorporated into confectionery compositions by admixing the inventive composition into conventional hard and soft confections.

As used herein, the term confectionery material means a product containing a bulking agent selected from a wide variety of materials such as sugar, corn syrup, and in the case of sugarless bulking agents, sugar alcohols such as sorbitol and mannitol and mixtures thereof. Confectionery material may include such exemplary substances as lozenges, tablets, toffee, nougat, suspensions, chewy candy, chewing gum and the like. The bulking agent is present in a quantity sufficient to bring the total amount of composition to 100%. In general, the bulking agent will be present in amounts up to about 99.98%, preferably in amounts up to about 99.9%, and more

2104461

- 78 -

preferably in amounts up to about 99%, by weight of the ingestible therapeutic composition.

Lozenges are flavored medicated dosage forms intended to be sucked and held in the mouth. Lozenges may be in the form of various shapes such as flat, circular, octagonal and biconvex forms. The lozenge bases are generally in two forms: hard boiled candy lozenges and compressed tablet lozenges.

10

Hard boiled candy lozenges may be processed and formulated by conventional means. In general, a hard boiled candy lozenge has a base composed of a mixture of sugar and other carbohydrate bulking agents kept in an amorphous or glassy condition. This amorphous or glassy form is considered a solid syrup of sugars generally having from about 0.5% to about 1.5% moisture. Such materials normally contain up to about 92% corn syrup, up to about 55% sugar and from about 0.1% to about 5% water, by weight of the final composition. The syrup component is generally prepared from corn syrups high in fructose, but may include other materials. Further ingredients such as flavoring agents, sweetening agents, acidulants, coloring agents and the like may also be added.

25

Boiled candy lozenges may also be prepared from non-fermentable sugars such as sorbitol, mannitol, and hydrogenated corn syrup. Typical hydrogenated corn syrups are Lycasin, a commercially available product manufactured by Roquette Corporation, and Hystar, a commercially available product manufactured by Lonza, Inc. The candy lozenges may contain up to about 95% sorbitol, a mixture of sorbitol and mannitol in a ratio from about 9.5:0.5 up to about 7.5:2.5, and hydrogenated corn syrup up to about 55%, by weight of the solid syrup component.

35

- 79 -

Boiled candy lozenges may be routinely prepared by conventional methods such as those involving fire cookers, vacuum cookers, and scraped-surface cookers also referred to as high speed atmospheric cookers.

Fire cookers involve the traditional method of making a boiled candy lozenge base. In this method, the desired quantity of carbohydrate bulking agent is dissolved in water by heating the agent in a kettle until the bulking agent dissolves. Additional bulking agent may then be added and cooking continued until a final temperature of 145° C. to 156° C. is achieved. The batch is then cooled and worked as a plastic-like mass to incorporate additives such as flavors, colorants and the like.

A high-speed atmospheric cooker uses a heat-exchanger surface which involves spreading a film of candy on a heat exchange surface, the candy is heated to 165° C. to 170° C. in a few minutes. The candy is then rapidly cooled to 100° C. to 120° C. and worked as a plastic-like mass enabling incorporation of the additives, such as flavors, colorants and the like.

In vacuum cookers, the carbohydrate bulking agent is boiled to 125° C. to 132° C., vacuum is applied and additional water is boiled off without extra heating. When cooking is complete, the mass is a semi-solid and has a plastic-like consistency. At this point, flavors, colorants, and other additives are admixed in the mass by routine mechanical mixing operations.

35

The optimum mixing required to uniformly mix the flavoring agents, coloring agents and other

2104461

- 80 -

additives during conventional manufacturing of boiled
candy lozenges is determined by the time needed to
obtain a uniform distribution of the materials.
Normally, mixing times of from 4 to 10 minutes have
5 been found to be acceptable.

Once the boiled candy lozenge has been
properly tempered, it may be cut into workable portions
or formed into desired shapes. A variety of forming
10 techniques may be utilized depending upon the shape and
size of the final product desired. A general
discussion of the composition and preparation of hard
confections may be found in H.A. Lieberman,
Pharmaceutical Dosage Forms: Tablets, Volume 1 (1980),
15 Marcel Dekker, Inc., New York, N.Y. at pages 339 to
469, which disclosure is incorporated herein by
reference.

The apparatus useful in accordance with the
20 present invention comprises cooking and mixing
apparatus well known in the confectionery manufacturing
arts, and therefore the selection of the specific
apparatus will be apparent to the artisan.

25 In contrast, compressed tablet confections
contain particulate materials and are formed into
structures under pressure. These confections generally
contain sugars in amounts up to about 95%, by weight of
the composition, and typical tablet excipients such as
30 binders and lubricants as well as flavoring agents,
coloring agents and the like.

In addition to hard confectionery materials,
the lozenges of the present invention may be made of
35 soft confectionery materials such as those contained in
nougat. The preparation of soft confections, such as
nougat, involves conventional methods, such as the

SUBSTITUTE SHEET

2104461

- 81 -

combination of two primary components, namely (1) a high boiling syrup such as a corn syrup, hydrogenated starch hydrolysate or the like, and (2) a relatively light textured frappe, generally prepared from egg albumin, gelatin, vegetable proteins, such as soy derived compounds, sugarless milk derived compounds such as milk proteins, and mixtures thereof. The frappe is generally relatively light, and may, for example, range in density from about 0.5 to about 0.7 grams/cc.

The high boiling syrup, or "bob syrup" of the soft confectionery is relatively viscous and has a higher density than the frappe component, and frequently contains a substantial amount of carbohydrate bulking agent such as a hydrogenated starch hydrolysate. Conventionally, the final nougat composition is prepared by the addition of the "bob syrup" to the frappe under agitation, to form the basic nougat mixture. Further ingredients such as flavoring agents, additional carbohydrate bulking agent, coloring agents, preservatives, medicaments, mixtures thereof and the like may be added thereafter also under agitation. A general discussion of the composition and preparation of nougat confections may be found in B.W. Minifie, Chocolate, Cocoa and Confectionery: Science and Technology, 2nd edition, AVI Publishing Co., Inc., Westport, Conn. (1980), at pages 424-425, which disclosure is incorporated herein by reference.

The procedure for preparing the soft confectionery involves known procedures. In general, the frappe component is prepared first and thereafter the syrup component is slowly added under agitation at a temperature of at least about 65° C., and preferably at least about 100° C. The mixture of components is continued to be mixed to form a uniform mixture, after

2104461

- 82 -

which the mixture is cooled to a temperature below 80° C., at which point, the flavoring agent may be added. The mixture is further mixed for an additional period until it is ready to be removed and formed into
5 suitable confectionery shapes.

The ingestible therapeutic compositions may also be in the form of a pharmaceutical suspension. Pharmaceutical suspensions of this invention may be
10 prepared by conventional methods long established in the art of pharmaceutical compounding. Suspensions may contain adjunct materials employed in formulating the suspensions of the art. The suspensions of the present invention can comprise:

15 (a) preservatives such as butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), benzoic acid, ascorbic acid, methyl paraben, propyl paraben, tocopherols, and the like, and mixtures thereof. Preservatives are generally present in
20 amounts up to about 1%, and preferably from about 0.05% to about 0.5%, by weight of the suspension;

(b) buffers such as citric acid-sodium citrate, phosphoric acid-sodium phosphate, and acetic acid-sodium acetate in amounts up to about 1%, and
25 preferably from about 0.05% to about 0.5%, by weight of the suspension;

(c) suspending agents or thickeners such as cellulosics like methylcellulose, carrageenans like alginic acid and its derivatives, xanthan gums,
30 gelatin, acacis, and microcrystalline cellulose in amounts up to about 20%, and preferably from about 1% to about 15%, by weight of the suspension;

(d) antifoaming agents such as dimethyl polysiloxane in amounts up to about 0.2%, and
35 preferably from about 0.01% to about 0.1%, by weight of the suspension;

2104461

- 83 -

(e) sweetening agents such as those sweeteners well known in the art, including both natural and artificial sweeteners. Sweetening agents such as monosaccharides, disaccharides and polysaccharides such as xylose, ribose, glucose (dextrose), mannose, galactose, fructose (levulose), sucrose (sugar), maltose, invert sugar (a mixture of fructose and glucose derived from sucrose), partially hydrolyzed starch, corn syrup solids, dihydrochalcones, monellin, steviosides, glycyrrhizin, and sugar alcohols such as sorbitol, mannitol, maltitol, hydrogenated starch hydrolysates and mixtures thereof may be utilized in amounts up to about 60%, and preferably from about 20% to about 50%, by weight of the suspension. Water-soluble artificial sweeteners such as soluble saccharin salts, i.e., sodium or calcium saccharin salts, cyclamate salts, the sodium, ammonium or calcium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide, the potassium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide (Acesulfame-K), the free acid form of saccharin, and the like may be utilized in amounts from about 0.001% to about 5%, by weight of the suspension;

(f) flavoring agents such as those flavors well known to the skilled artisan, such as natural and artificial flavors and mints, such as peppermint, menthol, citrus flavors such as orange and lemon, artificial vanilla, cinnamon, various fruit flavors, both individual and mixed and the like may be utilized in amounts from about 0.5% to about 5%, by weight of the suspension;

(g) coloring agents such as pigments which may be incorporated in amounts up to about 6%, by weight of the suspension. A preferred pigment, titanium dioxide, may be incorporated in amounts up to about 2%, and preferably less than about 1%, by weight of the suspension. The coloring agents may also

2104461

- 84 -

include natural food colors and dyes suitable for food, drug and cosmetic applications. These colorants are known as F.D.& C. dyes and lakes. The materials acceptable for the foregoing uses are preferably water-soluble. Such dyes are generally present in amounts up to about 0.25%, and preferably from about 0.05% to about 0.2%, by weight of the suspension;

(h) decolorizing agents such as sodium metabisulfite, ascorbic acid and the like may be incorporated into the suspension to prevent color changes due to aging. In general, decolorizing agents may be used in amounts up to about 0.25%, and preferably from about 0.05% to about 0.2%, by weight of the suspension; and

(i) solubilizers such as alcohol, propylene glycol, polyethylene glycol, and the like may be used to solubilize the flavoring agents. In general, solubilizing agents may be used in amounts up to about 10%, and preferably from about 2% to about 5%, by weight of the suspension.

The pharmaceutical suspensions of the present invention may be prepared as follows:

(A) admix the thickener with water heated from about 40° C. to about 95° C., preferably from about 40° C. to about 70° C., to form a dispersion if the thickener is not water soluble or a solution if the thickener is water soluble;

(B) admix the sweetening agent with water to form a solution;

(C) admix the therapeutic composition with the thickener-water admixture to form a uniform thickener-therapeutic composition;

(D) combine the sweetener solution with the thickener-therapeutic composition and mix until uniform; and

SUBSTITUTE SHEET

- 85 -

(E) admix the optional adjunct materials such as coloring agents, flavoring agents, decolorants, solubilizers, antifoaming agents, buffers and additional water with the mixture of step (D) to form
5 the suspension.

The ingestible therapeutic compositions of this invention may also be in chewable form. To achieve acceptable stability and quality as well as
10 good taste and mouth feel in a chewable formulation several considerations are important. These considerations include the amount of active substance per tablet, the flavoring agent employed, the degree of compressibility of the tablet and the organoleptic
15 properties of the composition.

Chewable therapeutic candy is prepared by procedures similar to those used to make soft confectionery. In a typical procedure, a boiled sugar-corn syrup blend is formed to which is added a frappe
20 mixture. The boiled sugar-corn syrup blend may be prepared from sugar and corn syrup blended in parts by weight ratio of about 90:10 to about 10:90. The sugar-corn syrup blend is heated to temperatures above about
25 120° C. to remove water and to form a molten mass. The frappe is generally prepared from gelatin, egg albumin, milk proteins such as casein, and vegetable proteins such as soy protein, and the like, which is added to a gelatin solution and rapidly mixed at ambient
30 temperature to form an aerated sponge like mass. The frappe is then added to the molten candy mass and mixed until homogeneous at temperatures between about 65° C. and about 120° C.

35 The ingestible therapeutic composition of the instant invention can then be added to the homogeneous mixture as the temperature is lowered to about 65° C.-

2104461

- 86 -

95° C. whereupon additional ingredients can then be added such as flavoring agents and coloring agents. The formulation is further cooled and formed into pieces of desired dimensions.

5

A general discussion of the lozenge and chewable tablet forms of confectionery may be found in H.A. Lieberman and L. Lachman, Pharmaceutical Dosage Forms: Tablets Volume 1, Marcel Dekker, Inc., New York, N.Y. at pages 289 to 466, which disclosure is incorporated herein by reference.

In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be admixed into the hard and soft confectionery products. These amounts are readily determined by those skilled in the art without the need for undue experimentation. In a preferred embodiment, the ingestible therapeutic composition will comprise the therapeutic composition in an amount from about 0.1% to about 10% and an ingestible vehicle, that is a pharmaceutically acceptable carrier, in a quantity sufficient to bring the total amount of composition to 100%, by weight the ingestible therapeutic composition. In a more preferred embodiment, the ingestible composition will comprise the therapeutic composition in an amount from about 0.1% to about 5%, and in a most preferred embodiment, the ingestible composition will comprise the therapeutic composition in an amount from about 0.1% to about 2%, and an ingestible vehicle in a quantity sufficient to bring the total amount of composition to 100%, by weight the ingestible therapeutic composition.

35

The present invention extends to methods of making the ingestible therapeutic compositions. In

- 87 -

such methods, an ingestible therapeutic composition is prepared by admixing a therapeutically effective amount of the therapeutic composition with a pharmaceutically-acceptable carrier. The apparatus useful in accordance with the present invention comprises mixing and heating apparatus well known in the confectionery arts, and therefore the selection of the specific apparatus will be apparent to the artisan. The final ingestible therapeutic compositions are readily prepared using methods generally known in the confectionery arts.

The therapeutic compositions may also be incorporated into chewing gums. In this form of the invention, the chewing gum composition contains a gum base, a bulking agent, the inventive therapeutic composition, and various additives.

The gum base employed will vary greatly depending upon various factors such as the type of base desired, the consistency of gum desired and the other components used in the composition to make the final chewing gum product. The gum base may be any water-insoluble gum base known in the art, and includes those gum bases utilized for chewing gums and bubble gums. Illustrative examples of suitable polymers in gum bases include both natural and synthetic elastomers and rubbers. For example, those polymers which are suitable as gum bases include, without limitation, substances of vegetable origin such as chicle, crown gum, nispero, rosadinha, jelutong, perillo, niger gutta, tunu, balata, gutta-percha, lechi-capsi, sorva, gutta kay, mixtures thereof and the like. Synthetic elastomers such as butadiene-styrene copolymers, polyisobutylene, isobutylene-isoprene copolymers, polyethylene, mixtures thereof and the like are particularly useful.

2104461

- 88 -

The gum base may include a non-toxic vinyl polymer, such as polyvinyl acetate and its partial hydrolysate, polyvinyl alcohol, and mixtures thereof. When utilized, the molecular weight of the vinyl polymer may range from about 2,000 up to and including about 94,000.

The amount of gum base employed will vary greatly depending upon various factors such as the type of base used, the consistency of the gum desired and the other components used in the composition to make the final chewing gum product. In general, the gum base will be present in amounts from about 5% to about 94%, by weight of the final chewing gum composition, and preferably in amounts from about 15% to about 45%, and more preferably in amounts from about 15% to about 35%, and most preferably in amounts from about 20% to about 30%, by weight of the final chewing gum composition.

The gum base composition may contain conventional elastomer solvents to aid in softening the elastomer base component. Such elastomer solvents may comprise terpinene resins such as polymers of alpha-pinene or beta-pinene, methyl, glycerol or pentaerythritol esters of rosins or modified rosins and gums, such as hydrogenated, dimerized or polymerized rosins or mixtures thereof. Examples of elastomer solvents suitable for use herein include the pentaerythritol ester of partially hydrogenated wood or gum rosin, the pentaerythritol ester of wood or gum rosin, the glycerol ester of wood rosin, the glycerol ester of partially dimerized wood or gum rosin, the glycerol ester of polymerized wood or gum rosin, the glycerol ester of tall oil rosin, the glycerol ester of wood or gum rosin and the partially hydrogenated wood or gum rosin and the partially hydrogenated methyl

SUBSTITUTE SHEET

- 89 -

ester of wood or rosin, mixtures thereof, and the like. The elastomer solvent may be employed in amounts from about 5% to about 75%, by weight of the gum base, and preferably from about 45% to about 70%, by weight of the gum base.

A variety of traditional ingredients may be included in the gum base in effective amounts such as plasticizers or softeners such as lanolin, palmitic acid, oleic acid, stearic acid, sodium stearate, potassium stearate, glyceryl triacetate, glyceryl lecithin, glyceryl monostearate, propylene glycol monostearate, acetylated monoglyceride, glycerine, mixtures thereof, and the like may also be incorporated into the gum base to obtain a variety of desirable textures and consistency properties. Waxes, for example, natural and synthetic waxes, hydrogenated vegetable oils, petroleum waxes such as polyurethane waxes, polyethylene waxes, paraffin waxes, microcrystalline waxes, fatty waxes, sorbitan monostearate, tallow, propylene glycol, mixtures thereof, and the like may also be incorporated into the gum base to obtain a variety of desirable textures and consistency properties. These traditional additional materials are generally employed in amounts up to about 30%, by weight of the gum base, and preferably in amounts from about 3% to about 20%, by weight of the gum base.

The gum base may include effective amounts of mineral adjuvants such as calcium carbonate, magnesium carbonate, alumina, aluminum hydroxide, aluminum silicate, talc, tricalcium phosphate, dicalcium phosphate and the like as well as mixtures thereof. These mineral adjuvants may serve as fillers and textural agents. These fillers or adjuvants may be used in the gum base in various amounts. Preferably

2104461

- 90 -

the amount of filler when used will be present in an amount up to about 60%, by weight of the chewing gum base.

5 The chewing gum base may additionally include the conventional additives of coloring agents, antioxidants, preservatives and the like. For example, titanium dioxide and other dyes suitable for food, drug and cosmetic applications, known as F.D. & C. dyes, may
10 be utilized. An antioxidant such as butylated hydroxytoluene (BHT), butylated hydroxyanisole (BHA), propyl gallate, and mixtures thereof, may also be included. Other conventional chewing gum additives known to one having ordinary skill in the chewing gum
15 art may also be used in the chewing gum base.

 The gum composition may include effective amounts of conventional additives selected from the group consisting of sweetening agents (sweeteners),
20 plasticizers, softeners, emulsifiers, waxes, fillers, bulking agents, mineral adjuvants, flavoring agents (flavors, flavorings), coloring agents (colorants, colorings), antioxidants, acidulants, thickeners, mixtures thereof and the like. Some of these additives
25 may serve more than one purpose. For example, in sugarless gum compositions, the sweetener, e.g., sorbitol or other sugar alcohol or mixtures thereof, may also function as a bulking agent. Similarly, in sugar containing gum compositions, the sugar sweetener
30 can also function as a bulking agent.

 The plasticizers, softeners, mineral adjuvants, colorants, waxes and antioxidants discussed above as being suitable for use in the gum base may
35 also be used in the gum composition. Examples of other conventional additives which may be used include emulsifiers, such as lecithin and glyceryl

SUBSTITUTE SHEET

- 91 -

monostearate, thickeners, used alone or in combination with other softeners, such as methyl cellulose, alginates, carrageenan, xanthan gum, gelatin, carob, tragacanth, locust bean, and carboxy methyl cellulose, acidulants such as malic acid, adipic acid, citric acid, tartaric acid, fumaric acid, and mixtures thereof, and fillers, such as those discussed above under the category of mineral adjuvants. The fillers when used may be utilized in an amount up to about 60%, by weight of the gum composition.

Bulking agents (carriers, extenders) suitable for use in chewing gums include sweetening agents selected from the group consisting of monosaccharides, disaccharides, poly-saccharides, sugar alcohols, and mixtures thereof; polydextrose; maltodextrins; minerals, such as calcium carbonate, talc, titanium dioxide, dicalcium phosphate, and the like. Bulking agents may be used in amounts up to about 90%, by weight of the final gum composition, with amounts from about 40% to about 70%, by weight of the gum composition being preferred, with from about 50% to about 65%, by weight, being more preferred and from about 55% to about 60%, by weight of the chewing gum composition, being most preferred.

The sweetening agent used may be selected from a wide range of materials including water-soluble sweeteners, water-soluble artificial sweeteners, water-soluble sweeteners derived from naturally occurring water-soluble sweeteners, dipeptide based sweeteners, and protein based sweeteners, including mixtures thereof. Without being limited to particular sweeteners, representative categories and examples include:

(a) water-soluble sweetening agents such as monosaccharides, disaccharides and polysaccharides such

2104461

- 92 -

as xylose, ribulose, glucose (dextrose), mannose, galactose, fructose (levulose), sucrose (sugar), maltose, invert sugar (a mixture of fructose and glucose derived from sucrose), partially hydrolyzed starch, corn syrup solids, dihydrochalcones, monellin, steviosides, glycyrrhizin, and sugar alcohols such as sorbitol, mannitol, maltitol, hydrogenated starch hydrolysates and mixtures thereof;

(b) water-soluble artificial sweeteners such as soluble saccharin salts, i.e., sodium or calcium saccharin salts, cyclamate salts, the sodium, ammonium or calcium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide, the potassium salt of 3,4-dihydro-6-methyl-1,2,3-oxathiazine-4-one-2,2-dioxide (Acesulfame-K), the free acid form of saccharin, and the like;

(c) dipeptide based sweeteners, such as L-aspartic acid derived sweeteners, such as L-aspartyl-L-phenylalanine methyl ester (Aspartame) and materials described in United States patent no. 3,492,131, L-alpha-aspartyl-N-(2,2,4,4-tetramethyl-3-thietanyl)-D-alanin-amide hydrate (Alitame), methyl esters of L-aspartyl-L-phenylglycerine and L-aspartyl-L-2,5-dihydrophenyl-glycine, L-aspartyl-2,5-dihydro-L-phenylalanine; L-aspartyl-L-(1-cyclohexen)-alanine, and the like;

(d) water-soluble sweeteners derived from naturally occurring water-soluble sweeteners, such as chlorinated derivatives of ordinary sugar (sucrose), known, for example, under the product designation of Sucralose; and

(e) protein based sweeteners such as thaumaococcus danielli (Thaumatin I and II).

35 In general, an effective amount of sweetener
is utilized to provide the level of bulk and/or
sweetness desired, and this amount will vary with the

- 93 -

sweetener selected. This amount of sweetener will normally be present in amounts from about 0.0025% to about 90%, by weight of the gum composition, depending upon the sweetener used. The exact range of amounts for each type of sweetener is well known in the art and is not the subject of the present invention. The amount of sweetener ordinarily necessary to achieve the desired level of sweetness is independent from the flavor level achieved from flavor oils.

10

Preferred sugar based-sweeteners are sugar (sucrose), corn syrup and mixtures thereof. Preferred sugarless sweeteners are the sugar alcohols, artificial sweeteners, dipeptide based sweeteners and mixtures thereof. Preferably, sugar alcohols are used in the sugarless compositions because these sweeteners can be used in amounts which are sufficient to provide bulk as well as the desired level of sweetness. Preferred sugar alcohols are selected from the group consisting of sorbitol, xylitol, maltitol, mannitol, and mixtures thereof. More preferably, sorbitol or a mixture of sorbitol and mannitol is utilized. The gamma form of sorbitol is preferred. An artificial sweetener or dipeptide based sweetener is preferably added to the gum compositions which contain sugar alcohols.

The coloring agents useful in the gum compositions are used in amounts effective to produce the desired color. These coloring agents include pigments which may be incorporated in amounts up to about 6% by weight of the gum composition. A preferred pigment, titanium dioxide, may be incorporated in amounts up to about 2%, and preferably less than about 1% by weight of the composition. The colorants may also include natural food colors and dyes suitable for food, drug and cosmetic applications. These colorants are known as F.D. & C. dyes and lakes. The materials

2104461

- 94 -

acceptable for the foregoing uses are preferably water-soluble. Illustrative nonlimiting examples include the indigoid dye known as F.D. & C. Blue No. 2, which is the disodium salt of 5,5-indigotindisulfonic acid. Similarly, the dye known as F.D. & C. Green No. 1 comprises a triphenylmethane dye and is the monosodium salt of 4-[4-(N-ethyl-p-sulfoniumbenzylamino)diphenylmethylene]-[1-(N-ethyl-N-p-sulfoniumbenzyl)-delta-2,5-cyclohexadieneimine]. A full recitation of all F.D. & C. colorants and their corresponding chemical structures may be found in the Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Edition, in volume 5 at pages 857-884, which text is incorporated herein by reference.

15

Suitable oils and fats usable in gum compositions include partially hydrogenated vegetable or animal fats, such as coconut oil, palm kernel oil, beef tallow, lard, and the like. These ingredients when used are generally present in amounts up to about 7%, by weight, and preferably up to about 3.5%, by weight of the gum composition.

In accordance with this invention, therapeutically effective amounts of the therapeutic compositions of the present invention may be admixed into a chewing gum. These amounts are readily determined by those skilled in the art without the need for undue experimentation. In a preferred embodiment, the final chewing gum composition will comprise the therapeutic composition in an amount from about 0.1% to about 10% and a chewing gum composition in a quantity sufficient to bring the total amount of composition to 100%, by weight of the chewing gum composition. In a more preferred embodiment, the final chewing gum composition will comprise the therapeutic composition in an amount from about 0.1% to about 5%, and in a most

2104461

- 95 -

preferred embodiment, the final chewing gum composition will comprise the therapeutic composition in an amount from about 0.1% to about 2%, and a chewing gum composition in a quantity sufficient to bring the total
5 amount of composition to 100%, by weight of the chewing gum composition.

The present invention extends to methods of making the therapeutic chewing gum compositions. The
10 therapeutic compositions may be incorporated into an otherwise conventional chewing gum composition using standard techniques and equipment known to those skilled in the art. The apparatus useful in accordance with the present invention comprises mixing and heating
15 apparatus well known in the chewing gum manufacturing arts, and therefore the selection of the specific apparatus will be apparent to the artisan.

For example, a gum base is heated to a
20 temperature sufficiently high enough to soften the base without adversely effecting the physical and chemical make up of the base. The optimum temperatures utilized may vary depending upon the composition of the gum base used, but such temperatures are readily determined by
25 those skilled in the art without undue experimentation.

The gum base is conventionally melted at temperatures that range from about 60° C. to about 120° C. for a period of time sufficient to render the
30 base molten. For example, the gum base may be heated under these conditions for a period of about thirty minutes just prior to being admixed incrementally with the remaining ingredients of the base such as the plasticizer, fillers, the bulking agent and/or
35 sweeteners, the softener and coloring agents to plasticize the blend as well as to modulate the hardness, viscoelasticity and formability of the base.

2104461 - 96 -

The chewing gum base is then blended with the therapeutic composition of the present invention which may have been previously blended with other traditional ingredients. Mixing is continued until a uniform
5 mixture of gum composition is obtained. Thereafter the gum composition mixture may be formed into desirable chewing gum shapes.

In a specific embodiment, the invention is
10 directed at a therapeutic pharmaceutical composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises:

(A) a therapeutically effective amount of a
15 therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

20 (b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

25 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
30 salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

35 (3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are

SUBSTITUTE SHEET

2104461

- 97 -

those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a pharmaceutically acceptable carrier.

15 The pharmaceutically acceptable carrier may be selected from the group consisting of pharmaceutical appliances, topical vehicles, and ingestible vehicle.

20 In another specific embodiment, the invention is directed at a method for preparing a therapeutic pharmaceutical composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of :

25 (A) providing a therapeutically effective amount of a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

30 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

35 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

2104461

- 98 -

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

5 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

10 (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
15 salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
20 injured mammalian cells; and

(B) providing a pharmaceutically acceptable carrier; and

(C) admixing the therapeutic composition from step (A) and the pharmaceutically acceptable carrier
25 from step (B) to form a therapeutic pharmaceutical composition.

Throughout this application, various publications have been referenced. The disclosures in
30 these publications are incorporated herein by reference in order to more fully describe the state of the art.

The present invention is further illustrated by the following examples which are not intended to
35 limit the effective scope of the claims. All parts and percentages in the examples and throughout the

SUBSTITUTE SHEET

- 99 -

specification and claims are by weight of the final composition unless otherwise specified.

EXAMPLES 1-26

5

These examples demonstrate a comparison of the viability of U937 monocytic cells after exposure of the cells to various antioxidants and combinations of antioxidants. These examples also demonstrate a comparison of the levels of hydrogen peroxide produced by U937 monocytic cells and mammalian epidermal keratinocytes after exposure of the cells to various antioxidants and combinations of antioxidants.

15

Mammalian epidermal keratinocytes and monocytes were employed to examine the ability of various antioxidants to reduce levels of hydrogen peroxide in these cells. Hydrogen peroxide was measured after the cells were exposed to ultraviolet light in the wavelength range from 290 to 320 nm (UV-B) or to the inflammatory compound 12-O-tetradecanoyl-phorbol-13-acetate (TPA). Sodium pyruvate was tested at various concentrations to determine the effect of concentrations of this antioxidant on the hydrogen peroxide production by epidermal cells and monocytes. Magnesium pyruvate, calcium pyruvate, zinc pyruvate, and combinations of sodium pyruvate with ascorbic acid, lactic acid, and Vitamin E were then tested to determine the effect of these salts and combinations of antioxidants on the hydrogen peroxide production by epidermal cells and monocytes.

35

Mammalian epidermal keratinocytes were isolated by trypsinization of epithelial sheets and grown in modified basal MCDB 153 medium supplemented with epidermal growth factor, bovine pituitary extract, and hydrocortisone. Cells were maintained in a

SUBSTITUTE SHEET

2104461

- 100 -

humidified incubator with 5% carbon dioxide at 37 °C. Keratinocytes were seeded in 60 mm culture dishes at a cell density of 3×10^5 cells per dish and the cultures were exposed to 1 M.E.D. dose of ultraviolet-B light (100 mJ/cm²) or treated with 100 ng/ml of TPA.

U937 monocytic cells are a cultured cell line grown in RPMI media with 10% fetal calf serum. Cells were maintained in a 60 mm culture dish at 5% carbon dioxide at 37 °C. at a seeding density not exceeding 1×10^6 cells per dish.

Sodium pyruvate, lactic acid, ascorbic acid, and Vitamin E were dissolved in distilled water. The concentrations of the sodium pyruvate solutions prepared were 1 mM, 10 mM, 50 mM, 100 mM, and 200 mM. The concentrations of the lactic acid solutions prepared were 1.0%, 0.1%, and 0.05%. The concentrations of the ascorbic acid solutions prepared were 1.0%, 0.1%, 0.05%, and 0.025%. The concentrations of the Vitamin E solutions prepared were 1 U, 10 U, 50 U, and 100 U. The test solutions were adjusted to a pH value of 7.4 with 1.0N sodium hydroxide solution and then sterile filtered. The appropriate concentration of test solution or combination of test solutions was added to the cells immediately prior to exposure of the cells to ultraviolet light-B or TPA [100ng/ml]. Stock solutions were prepared so that the vehicle did not constitute more than 1% of the total volume of the culture media.

Intracellular hydrogen peroxide production by mammalian epidermal keratinocytes and U937 monocytes was measured using dichlorofluorescein diacetate (DCFH-DA, Molecular Probes, Eugene, Ore.). DCFH-DA is a non-polar non-fluorescent compound that readily diffuses into cells where it is hydrolyzed to the polar non-

SUBSTITUTE SHEET

2104461

- 101 -

fluorescent derivative DCFH which then becomes trapped within the cells. In the presence of intracellular hydrogen peroxide, DCFH is oxidized to the highly fluorescent compound DCF. Hence, cellular fluorescence intensity is directly proportional to the level of intracellular hydrogen peroxide produced. Cellular fluorescence intensity can be monitored by fluorimetry and by flow cytometry.

10 Mammalian epidermal keratinocytes and U937 cultured monocytes (1×10^6 per dish) were incubated at 37°C . with 5 μM of DCFH-DA. Production of hydrogen peroxide was measured using a Coulter Profile analytical flow cytometer. Linear and log
15 intensity of green fluorescence data was collected. For each analysis, a quantity of 10,000 to 20,000 events was accumulated. Optical alignment for the instrument was performed daily. Coefficients of variation for forward angle light scatter and
20 integrated green fluorescence were generally less than two. Each analysis was repeated three times and the quantitation of fluorescence was expressed in terms of femtomoles (fmol , 10^{-15} moles) of DCF oxidized per cell, which is a direct measure of the intracellular
25 hydrogen peroxide produced. Alternatively, in the saturated and unsaturated fatty acid examples in examples 27-52, fluorimetry was used to assess the DCF oxidation per cell.

30 The viability of the U937 monocytic cells after exposure of the cells to various antioxidants for 24 hours was measured. The viability of the cells was determined by exposing the cells to the dye propidium iodide. Permeable cell membranes which absorbed the
35 dye were not considered viable. The viability of the cells was represented as the percentage of cells that excluded propidium iodide. FIGURE 1 depicts in bar

SUBSTITUTE SHEET

2104461 - 102 -

graph format the viability of U937 monocytic cells after exposure of the cells to no antioxidant (Example 1, control), to sodium pyruvate (Example 2), to ascorbic acid (Example 3), to lactic acid (Example 4), and to Vitamin E (Example 5). FIGURE 2 depicts in bar graph format the viability of U937 monocytic cells after exposure of the cells to various combinations of antioxidants. Specifically, the viability of U937 monocytic cells was measured after exposure to no antioxidant (Example 6, control), to ascorbic acid and lactic acid (Example 7), to ascorbic acid and Vitamin E (Example 8), to sodium pyruvate and ascorbic acid (Example 9), to sodium pyruvate and lactic acid (Example 10), to sodium pyruvate and Vitamin E (Example 11), to lactic acid and Vitamin E (Example 12), and to sodium pyruvate, ascorbic acid, and lactic acid (Example 13).

FIGURE 1 shows that ascorbic acid is cytotoxic to monocytes at concentrations as low as 0.25%. FIGURE 2 shows that the cytotoxicity of ascorbic acid was reversed by the addition of 10 mM of sodium pyruvate. FIGURES 1 and 2 show that the viability rate of 15% to 20% of the cells when treated with ascorbic acid was increased to 95% to 98% upon addition of sodium pyruvate. Lactic acid and Vitamin E did not reverse the cytotoxicity of ascorbic acid.

Sodium pyruvate was then tested at various concentrations to determine the effect of concentrations of this antioxidant on the hydrogen peroxide production by epidermal cells and monocytes. Mammalian epidermal keratinocytes and monocytes were exposed to (a) 1 M.E.D. dose of ultraviolet light-B and (b) 100 ng/ml of 12-O-tetradecanoylphorbol-13-acetate (TPA) in the presence of sodium pyruvate at the

SUBSTITUTE SHEET

- 103 -

following concentrations: 200 mM, 100 mM, 50 mM, 10 mM, 1 mM.

5 The optimum concentration of sodium pyruvate to reduce the hydrogen peroxide production by epidermal cells and monocytes was found to be 10 mM. Concentrations of sodium pyruvate of 50 mM and above were cytotoxic to both epidermal keratinocytes and monocytes.

10

 Magnesium pyruvate, calcium pyruvate, zinc pyruvate, ascorbic acid, lactic acid, and Vitamin E, and combinations of sodium pyruvate with ascorbic acid, lactic acid, and Vitamin E were then tested to
15 determine the effect of these salts and combinations of antioxidants on the hydrogen peroxide production by epidermal cells and monocytes. The following test solutions were prepared.

20

(a) sodium pyruvate [10 mM];
(b) zinc salt [10 mM];
(c) magnesium salt [10 mM];
(d) calcium salt [10 mM];
(e) sodium pyruvate [10 mM] and ascorbic acid
25 [0.025%];

25

(f) sodium pyruvate [10 mM] and lactic acid
 [0.05%];
(g) sodium pyruvate [10 mM], lactic acid,
 [0.05%], and ascorbic acid

30

[0.025%];

(h) lactic acid [1.0%, 0.1%, and 0.05%];
(i) ascorbic acid [1.0%, 0.1%, 0.05%, and
 0.025%];
(j) Vitamin E [1 U, 10 U, 50 U, and 100 U];

35

and

(k) vehicle solvent controls.

2104461

- 104 -

There was no significant difference among the zinc, magnesium, and calcium salts of pyruvic acid on the hydrogen peroxide production by epidermal cells and monocytes. The zinc and calcium salts of pyruvic acid induced differentiation of keratinocytes. For convenience, the sodium salt was used in subsequent tests.

The optimum concentration of lactic acid to reduce the hydrogen peroxide production by epidermal cells and monocytes was found to be 0.05%. The optimum concentration of ascorbic acid was found to be 0.025%. The higher concentrations of both of these compounds were found to be cytotoxic to both types of cells. The optimum concentration of Vitamin E was found to be 50 U.

FIGURE 3 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells after exposure of the cells to no antioxidant (Example 14, control), to sodium pyruvate (Example 15), to ascorbic acid (Example 16), to lactic acid (Example 17), and to Vitamin E (Example 18). Sodium pyruvate and Vitamin E significantly reduced the hydrogen peroxide production by monocytes.

FIGURE 4 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells after exposure of the cells to various combinations of antioxidants. Specifically, the levels of hydrogen peroxide produced by U937 monocytic cells were measured after exposure to no antioxidant (Example 19, control), to ascorbic acid and lactic acid (Example 20), to ascorbic acid and Vitamin E (Example 21), to sodium pyruvate and ascorbic acid (Example 22), to sodium pyruvate and lactic acid (Example 23), to sodium pyruvate and Vitamin E

SUBSTITUTE SHEET

2104461

- 105 -

(Example 24), to lactic acid and Vitamin E (Example 25), and to sodium pyruvate, ascorbic acid, and lactic acid (Example 26). The combination of lactic acid (0.05%) and Vitamin E (50 U) significantly
5 reduced the hydrogen peroxide production by monocytes.

The morphological alterations in epidermal keratinocytes were observed in control cultures and in cultures exposed to ultraviolet-B. Cells in the layer
10 closest to the dermis are basal keratinocytes. These cells proliferate and migrate into the spinous and granular layers of the epidermis where the cells begin to differentiate. The differentiation pattern results in cells enucleating and forming cornified envelopes at
15 the uppermost portion of the epidermis, the stratum corneum. The differentiation of keratinocytes is controlled by the levels of calcium, magnesium, and other elements in the medium. Cells in culture systems promoting differentiation appear as an epidermal sheet
20 forming attachments or tight junctions with each other. Keratinocytes that become nonadherent or float in the media were considered responding to a cytotoxic event.

The following morphological alterations in
25 the mammalian epidermal keratinocytes were observed for the following control cultures:

10 mM Sodium Pyruvate: Tight junctions of cells were formed and the proliferation rate of the cells was
30 higher than the rate of the control cells.

0.025% Ascorbic Acid: Cells were floating in a cytotoxic response to ascorbic acid.

35 0.025% Ascorbic acid and 10 mM Sodium Pyruvate: Few tight junctions of cells were observed and cells

2104461

- 106 -

appeared similar to the cells in the sodium pyruvate culture.

5 0.05% Lactic Acid: Cells appeared dramatically altered as an epidermal sheet and as flat granular cells.

10 0.05% Lactic Acid and 10 mM Sodium Pyruvate: Cells formed an epidermal sheet but appeared smaller than the cell in the lactic acid culture.

15 50 U Vitamin E: Cells appeared the same as the cells in the control culture.

20 50 U Vitamin E and 10 mM Sodium Pyruvate: Cells increased in number and changed in appearance resembling the cells in the sodium pyruvate culture.

25 The following morphological alterations in the mammalian epidermal keratinocytes were observed for the corresponding cultures exposed to ultraviolet light-B, 100 mJoules, for 24 hours:

30 10 mM Sodium Pyruvate: Cells proliferated more rapidly than the cells in the control culture.

35 0.025% Ascorbic Acid: Cells were nonadherent and floating in a cytotoxic response to ascorbic acid greater than the cytotoxic response of the corresponding cells without ultraviolet-B light exposure.

40 0.05% Lactic Acid: Cells formed an epidermal sheet and were more granular than cells in the control culture without ultraviolet-B light exposure.

35

SUBSTITUTE SHEET

- 107 -

50 U Vitamin E: Cell growth was inhibited but cells appeared similar to cells in the control culture without ultraviolet-B light exposure.

- 5 50 U Vitamin E and 10 mM Sodium Pyruvate: Cells appeared similar to cells in the control culture and proliferated to a greater extent than cells in the control cultures without ultraviolet-B light exposure.
- 10 Morphological alterations in the U937 monocytic cell line were also observed for control cultures and cultures exposed to ultraviolet light-B, 100 mJoules, for 24 hours. The following compounds and combination of compounds, at the concentrations set out
- 15 below, significantly inhibited the levels of hydrogen peroxide produced by U937 monocytic cells

- Sodium pyruvate at 10 mM and 50 mM;
Vitamin E at 50 U and 100 U; and
- 20 Lactic acid at 0.05% and Vitamin E at 50 U.

EXAMPLES 27-52

- 25 These examples demonstrate a comparison of the levels of hydrogen peroxide produced by U937 monocytic cells and epidermal keratinocytes after exposure of the cells to various combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids.

- 30 Mammalian epidermal keratinocytes and U937 monocytic cells and the test solutions of sodium pyruvate, lactic acid, ascorbic acid, and Vitamin E were prepared as describe above for Examples 1-26.
- 35 Intracellular hydrogen peroxide production by the mammalian epidermal keratinocytes and U937 monocytes was also measured as described above.

SUBSTITUTE SHEET

2104461

- 108 -

A mixture of fatty acids derived from chicken fat was prepared for addition to the cultured cells by mixing 0.1% of the chicken fat with the culture media. At the temperature of the culture media, 37 °C., the chicken fat was miscible. This chicken fat mixture was added to cultures of cells prior to exposure of the cells to ultraviolet-B light or TPA treatment.

As set out in examples 1-26, mammalian epidermal keratinocytes and monocytes were exposed to (a) 1 M.E.D. dose of ultraviolet light-B and (b) 100 ng/ml of 12-O-tetradecanoylphorbol-13-acetate in the presence of various antioxidants and combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids [0.1%, 0.5%, and 1.0% chicken fat].

FIGURE 5 depicts in bar graph format the levels of hydrogen peroxide produced by U937 monocytic cells after exposure of the cells to various combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids. Specifically, the levels of hydrogen peroxide produced by U937 monocytic cells were measured after exposure to lactic acid and Vitamin E without fatty acids (Example 27) and with fatty acids (Example 28), to ascorbic acid and lactic acid without fatty acids (Example 29) and with fatty acids (Example 30), and to ascorbic acid and Vitamin E without fatty acids (Example 31) and with fatty acids (Example 32). The ability of the combinations of lactic acid and Vitamin E, ascorbic acid and lactic acid, and ascorbic acid and Vitamin E to reduce the hydrogen peroxide production by monocytes was increased in the presence of fatty acids. The most effective combination to reduce the hydrogen peroxide production of monocytes

SUBSTITUTE SHEET

2104461

- 109 -

was lactic acid (0.05%) and Vitamin E (50 E) in the presence of a mixture of saturated and unsaturated fatty acids (0.5%).

5 FIGURE 6 depicts in bar graph format the levels of hydrogen peroxide produced by epidermal keratinocytes after exposure of the cells to various antioxidants with and without a mixture of saturated and unsaturated fatty acids. Specifically, the levels
10 of hydrogen peroxide produced by epidermal keratinocytes were measured after exposure to no antioxidant without fatty acids (Example 33, control) and with fatty acids (Example 34), to sodium pyruvate without fatty acids (Example 35) and with fatty acids
15 (Example 36), to ascorbic acid without fatty acids (Example 37) and with fatty acids (Example 38), to lactic acid without fatty acids (Example 39) and with fatty acids (Example 40), and to Vitamin E without fatty acids (Example 41) and with fatty acids
20 (Example 42). The ability of sodium pyruvate and Vitamin E to reduce the hydrogen peroxide production by epidermal keratinocytes was increased in the presence of fatty acids. The most effective combinations to reduce the hydrogen peroxide production of epidermal
25 keratinocytes were sodium pyruvate in combination with a mixture saturated and unsaturated fatty acids and Vitamin E in combination with a mixture of saturated and unsaturated fatty acids.

30 FIGURE 7 depicts in bar graph format the levels of hydrogen peroxide produced by epidermal keratinocytes after exposure of the cells to various combinations of antioxidants with and without a mixture of saturated and unsaturated fatty acids.
35 Specifically, the levels of hydrogen peroxide produced by epidermal keratinocytes were measured after exposure to no antioxidant without fatty acids (Example 43,

SUBSTITUTE SHEET

2104461

- 110 -

control) and with fatty acids (Example 44), to sodium pyruvate and ascorbic acid without fatty acids (Example 45) and with fatty acids (Example 46), to sodium pyruvate and lactic acid without fatty acids (Example 47) and with fatty acids (Example 48), to sodium pyruvate and Vitamin E without fatty acids (Example 49) and with fatty acids (Example 50), and to ascorbic acid and Vitamin E without fatty acids (Example 51) and with fatty acids (Example 52). The ability of all combinations of antioxidants to reduce the hydrogen peroxide production by epidermal keratinocytes was increased in the presence of fatty acids. In order of potency, the most effective combinations to reduce the hydrogen peroxide production of epidermal keratinocytes were sodium pyruvate and Vitamin E, sodium pyruvate and lactic acid, and Vitamin E, each in combination with a mixture of saturated and unsaturated fatty acids (0.5%).

Because of the cytotoxicity of cells towards ascorbic acid described above, the ascorbic acid combinations without sodium pyruvate were not considered significantly different from the control test solution.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modifications are intended to be included within the scope of the following claims.

2104461

- 111 -

I claim:

1. A therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises:
- 5 (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;
- 10 (b) an antioxidant; and
- (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.
- 15 2. The composition according to claim 1, wherein the mammalian cells comprise epidermal keratinocytes.
- 20 3. The composition according to claim 1, wherein the pyruvate is selected from the group consisting of pyruvic acid, lithium pyruvate, sodium pyruvate, potassium pyruvate, magnesium pyruvate, calcium pyruvate, zinc pyruvate, manganese pyruvate, and mixtures thereof.
- 25 4. The composition according to claim 3, wherein the pyruvate is sodium pyruvate.
- 30 5. The composition according to claim 1, wherein the antioxidant is selected from the group consisting of retinol, 3, 4-didehydroretinol, alpha-carotene, beta-carotene, gamma-carotene, delta-carotene, ascorbic acid, alpha-tocopherol, beta-tocopherol, gamma-tocopherol, delta-tocopherol, and mixtures thereof.
- 35

SUBSTITUTE SHEET

2104461 - 112 -

6. The composition according to claim 5,
wherein the antioxidant is *alpha*-tocopherol.

7. The composition according to claim 1,
5 wherein the mixture of saturated and unsaturated fatty
acids comprises lauric acid, myristic acid, myristoleic
acid, pentadecanoic acid, palmitic acid, palmitoleic
acid, margaric acid, margaroleic acid, stearic, oleic
acid, linoleic acid, linolenic acid, arachidic acid,
10 and gaddoleic acid.

8. The composition according to claim 7,
wherein the mixture of saturated and unsaturated fatty
acids comprises human fat, chicken fat, cow fat, sheep
15 fat, horse fat, pig fat, and whale fat.

9. The composition according to claim 1,
wherein pyruvate is present in the therapeutic
composition in an amount from about 10% to about 50%,
20 by weight of the therapeutic composition.

10. The composition according to claim 1,
wherein the antioxidant is present in the therapeutic
composition in an amount from about 10% to about 50%,
25 by weight of the therapeutic composition.

11. The composition according to claim 1,
wherein the mixture of saturated and unsaturated fatty
acids is present in the therapeutic composition in an
30 amount from about 10% to about 50%, by weight of the
therapeutic composition.

12. A therapeutic composition for preventing
and reducing injury to mammalian cells, and increasing
35 the resuscitation rate of injured mammalian cells,
which comprises:

SUBSTITUTE SHEET

2104461

- 113 -

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

5 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured
10 mammalian cells.

13. The composition according to claim 12, wherein the mammalian cells comprise epidermal keratinocytes.
15

14. The composition according to claim 12, wherein the pyruvate is selected from the group consisting of pyruvic acid, lithium pyruvate, sodium pyruvate, potassium pyruvate, magnesium pyruvate, calcium pyruvate, zinc pyruvate, manganese pyruvate, and mixtures thereof.
20

15. The composition according to claim 14, wherein the pyruvate is sodium pyruvate.
25

16. The composition according to claim 12, wherein the lactate is selected from the group consisting of lactic acid, lithium lactate, sodium lactate, potassium lactate, magnesium lactate, calcium lactate, zinc lactate, manganese lactate, and mixtures thereof.
30

17. The composition according to claim 16, wherein the lactate is lactic acid.
35

2104461

- 114 -

18. The composition according to claim 12, wherein the mixture of saturated and unsaturated fatty acids comprises lauric acid, myristic acid, myristoleic acid, pentadecanoic acid, palmitic acid, palmitoleic acid, margaric acid, margaroleic acid, stearic, oleic acid, linoleic acid, linolenic acid, arachidic acid, and gaddoleic acid.

19. The composition according to claim 18, wherein the mixture of saturated and unsaturated fatty acids comprises human fat, chicken fat, cow fat, sheep fat, horse fat, pig fat, and whale fat.

20. The composition according to claim 12, wherein pyruvate is present in the therapeutic composition in an amount from about 10% to about 50%, by weight of the therapeutic composition.

21. The composition according to claim 12, wherein lactate is present in the therapeutic composition in an amount from about 10% to about 50%, by weight of the therapeutic composition.

22. The composition according to claim 12, wherein the mixture of saturated and unsaturated fatty acids is present in the therapeutic composition in an amount from about 10% to about 50%, by weight of the therapeutic composition.

23. A therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises:

(a) an antioxidant; and

2104461

- 115 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

5

24. The composition according to claim 23, wherein the mammalian cells comprise epidermal keratinocytes.

10

25. The composition according to claim 23, wherein the antioxidant is selected from the group consisting of retinol, 3, 4-didehydroretinol, alpha-carotene, beta-carotene, gamma-carotene, delta-carotene, ascorbic acid, alpha-tocopherol, beta-tocopherol, gamma-tocopherol, delta-tocopherol, and mixtures thereof.

15

26. The composition according to claim 23, wherein the mixture of saturated and unsaturated fatty acids comprises lauric acid, myristic acid, myristoleic acid, pentadecanoic acid, palmitic acid, palmitoleic acid, margaric acid, margaroleic acid, stearic, oleic acid, linoleic acid, linolenic acid, arachidic acid, and gaddoleic acid.

20

27. The composition according to claim 26, wherein the mixture of saturated and unsaturated fatty acids comprises human fat, chicken fat, cow fat, sheep fat, horse fat, pig fat, and whale fat.

25

28. The composition according to claim 23, wherein the antioxidant is present in the therapeutic composition in an amount from about 10% to about 50%, by weight of the therapeutic composition.

30

35

2104461

- 116 -

29. The composition according to claim 23, wherein the mixture of saturated and unsaturated fatty acids is present in the therapeutic composition in an amount from about 10% to about 50%, by weight of the
5 therapeutic composition.

30. A therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells,
10 which comprises:

(a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

15 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

20 31. The composition according to claim 30, wherein the mammalian cells comprise monocytes.

32. The composition according to claim 30, wherein the lactate is selected from the group
25 consisting of lactic acid, lithium lactate, sodium lactate, potassium lactate, magnesium lactate, calcium lactate, zinc lactate, manganese lactate, and mixtures thereof.

30 33. The composition according to claim 32, wherein the lactate is lactic acid.

34. The composition according to claim 30, wherein the antioxidant is selected from the group
35 consisting of retinol, 3, 4-didehydroretinol, alpha-carotene, beta-carotene, gamma-carotene, delta-carotene, ascorbic acid, alpha-tocopherol, beta-

SUBSTITUTE SHEET

2104461

- 117 -

tocopherol, gamma-tocopherol, delta-tocopherol, and mixtures thereof.

35. The composition according to claim 34,
5 wherein the antioxidant is alpha-tocopherol.

36. The composition according to claim 30,
wherein the mixture of saturated and unsaturated fatty
acids comprises lauric acid, myristic acid, myristoleic
10 acid, pentadecanoic acid, palmitic acid, palmitoleic
acid, margaric acid, margaroleic acid, stearic, oleic
acid, linoleic acid, linolenic acid, arachidic acid,
and gaddoleic acid.

37. The composition according to claim 36,
wherein the mixture of saturated and unsaturated fatty
acids comprises human fat, chicken fat, cow fat, sheep
fat, horse fat, pig fat, and whale fat.

38. The composition according to claim 30,
wherein lactate is present in the therapeutic
composition in an amount from about 10% to about 50%,
by weight of the therapeutic composition.

39. The composition according to claim 30,
wherein the antioxidant is present in the therapeutic
composition in an amount from about 10% to about 50%,
by weight of the therapeutic composition.

40. The composition according to claim 30,
wherein the mixture of saturated and unsaturated fatty
acids is present in the therapeutic composition in an
amount from about 10% to about 50%, by weight of the
therapeutic composition.

35

2104461

- 118 -

41. A method for preparing a therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps
5 of admixing the following ingredients:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

15 42. A method for preparing a therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of admixing the following ingredients:

20 (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
25 salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

30

43. A method for preparing a therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps
35 of admixing the following ingredients:

(a) an antioxidant; and

SUBSTITUTE SHEET

2104461

- 119 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

5

44. A method for preparing a therapeutic composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of admixing the following ingredients:

10

(a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

15

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells.

20

45. A method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of:

(A) providing a therapeutic composition which comprises:

25

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

30

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) contacting the therapeutic composition with the mammalian cells.

35

2104461

- 120 -

46. A method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of:

5 (A) providing a therapeutic composition which comprises:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

10 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

15 (B) contacting the therapeutic composition with the mammalian cells.

20 47. A method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of:

(A) providing a therapeutic composition which comprises:

(a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

30 (B) contacting the therapeutic composition with the mammalian cells.

48. A method for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises the steps of:

2104461

- 121 -

(A) providing a therapeutic composition which comprises:

(a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) contacting the therapeutic composition with the mammalian cells.

49. An augmented pharmaceutical composition having an enhanced ability to prevent and reduce injury to mammalian cells which comprises:

(A) a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

SUBSTITUTE SHEET

2104461

- 122 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a medicament useful for treating injured mammalian cells.

15

50. The augmented pharmaceutical composition according to claim 49, wherein the medicament is selected from the group consisting of anti-inflammatory, antibacterials, antiseptics, burn relief medications, sun burn medications, acne preparations, insect bite and sting medications, wound cleansers and wound dressings, stroke medications, autoimmune disease medications, arthritis medications, ulcer medications, and cancer medications.

25

51. A cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises:

30 (A) a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

35

(b) an antioxidant; and

2104461

- 123 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

15 (3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

20 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a medicament having cytotoxic properties.

30 52. The cytoprotective pharmaceutical composition according to claim 51, wherein the medicament having cytotoxic properties is selected from the group consisting of epithelial cell cohesiveness reducers, dermatological abrasants, anti-inflammatory, lipid regulating agents, centrally acting anticholinesterases, chemotherapeutic drugs, and

35 gastric irritants.

SUBSTITUTE SHEET

2104461

- 124 -

53. The cytoprotective pharmaceutical composition according to claim 52, wherein the medicament having cytotoxic properties is selected from the group consisting of gemfibrozil, lovastatin, tacrine, and doxorubicin.

54. A prophylactic cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises:

(A) a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

(B) a medicament having cytotoxic properties.

55. A timed release prophylactic cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises:

2104461

- 125 -

(A) a therapeutic composition in an immediate release form selected from the group consisting of:

5 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

10 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(3) (a) an antioxidant; and

15 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

(B) an anticancer medicament having cytotoxic properties in timed release form;

20 wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition
25 and the non-cancerous cells have not substantially metabolized the therapeutic composition.

56. A preventative cytoprotective pharmaceutical composition for preventing and reducing
30 injury to mammalian cells from a medicament having cytotoxic properties which comprises:

(A) a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and
35 mixtures thereof; and

(B) a medicament having cytotoxic properties.

SUBSTITUTE SHEET

2104461

- 126 -

57. A timed release preventative cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises:

(A) a therapeutic composition in an immediate release form comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) an anticancer medicament having cytotoxic properties in timed release form;

wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition.

58. A wound healing pharmaceutical composition to increase the resuscitation rate of injured mammalian cells which comprises:

(A) a first wound healing component comprising a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a second wound healing component selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) an antioxidant.

- 127 -

59. A method for preparing an augmented pharmaceutical composition having an enhanced ability to prevent and reduce injury to mammalian cells which comprises the steps of:

(A) providing a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

2104461

- 128 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (B) providing a medicament useful for treating injured mammalian cells; and

(C) admixing the therapeutic composition from step (A) with the medicament from step (B) to prepare the augmented pharmaceutical composition.

10

60. A method for preparing a cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises the steps of:

15 (A) providing a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

20 (b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

25 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

30 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

35 (3) (a) an antioxidant; and

- 129 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(B) providing a medicament having cytotoxic properties; and

15 (C) admixing the therapeutic composition from step (A) with the medicament from step (B) to prepare the cytoprotective pharmaceutical composition.

20 61. A method for preparing a prophylactic cytoprotective pharmaceutical composition for preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises the steps of:

25 (A) providing a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

30 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

35 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant; and

2104461

- 130 -

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant;

5 (B) providing a medicament having cytotoxic properties; and

(C) admixing the therapeutic composition from step (A) with the medicament from step (B) to prepare the prophylactic cytoprotective pharmaceutical
10 composition.

62. A method for preparing a timed release prophylactic cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian
15 cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition in an immediate release form selected from the group
20 consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

25 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
30 salts of lactic acid, and mixtures thereof; and

(3) (a) an antioxidant; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

35 (b) an antioxidant;

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and

- 131 -

(C) admixing the therapeutic composition from step (A) with the medicament from step (B) to prepare the timed release prophylactic cytoprotective pharmaceutical composition;

5 wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition
10 and the non-cancerous cells have not substantially metabolized the therapeutic composition.

63. A method for preparing a preventative cytoprotective pharmaceutical composition for
15 preventing and reducing injury to mammalian cells from a medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic
20 acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(B) providing a medicament having cytotoxic properties; and

(C) admixing the therapeutic composition from
25 step (A) with the medicament from step (B) to prepare the timed release preventative cytoprotective pharmaceutical composition.

64. A method for preparing a timed release
30 preventative cytoprotective pharmaceutical composition for selectively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

2104461

- 132 -

(A) providing a therapeutic composition in an immediate release form comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and

(C) admixing the therapeutic composition from step (A) with the medicament from step (B) to prepare the timed release preventative cytoprotective pharmaceutical composition;

wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition.

65. A method for preparing a wound healing pharmaceutical composition to increase the resuscitation rate of injured mammalian cells which comprises the steps of:

(A) providing a first wound healing component comprising a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) providing a second wound healing component selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(c) an antioxidant; and

2104461

- 133 -

(C) admixing the first wound healing component from step (A) with the second wound healing component from step (B) to prepare the wound healing pharmaceutical composition.

5

66. A method for augmenting the therapeutic effect of a medicament useful for treating injured mammalian cells which comprises the steps of:

10 (A) providing a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

15 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

20 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

25 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

30 (b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

35 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

SUBSTITUTE SHEET

2104461

- 134 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (B) providing a medicament useful for treating injured cells; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to augment the therapeutic effect of
10 the medicament useful for treating injured mammalian cells.

67. A method for protecting mammalian cells from a medicament having cytotoxic properties which
15 comprises the steps of:

(A) providing a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
20 salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
25 injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and
30

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
35 injured mammalian cells;

(3) (a) an antioxidant; and

SUBSTITUTE SHEET

2104461

- 135 -

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(B) providing a medicament having cytotoxic properties; and

15 (C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells from the medicament having cytotoxic properties.

20 68. A method for protecting mammalian cells prophylactically from a medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition selected from the group consisting of:

25 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

30 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

35 (3) (a) an antioxidant; and

2104461

- 136 -

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

5 (B) providing a medicament having cytotoxic properties; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells
10 prophylactically from the medicament having cytotoxic properties.

69. A method for prophylactically protecting non-cancerous mammalian cells in the presence of
15 cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition in an immediate release form selected from the group
20 consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

25 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable
30 salts of lactic acid, and mixtures thereof; and

(3) (a) an antioxidant; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

35 (b) an antioxidant;

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and

SUBSTITUTE SHEET

- 137 -

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to prophylactically protect non-cancerous mammalian cells in the presence of cancerous mammalian cells from the anticancer medicament having cytotoxic properties;

wherein the therapeutic composition is released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition.

70. A method for protecting mammalian cells preventatively from a medicament having cytotoxic properties which comprises the steps of:

(A) providing a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(B) providing a medicament having cytotoxic properties; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently to mammalian cells to protect the mammalian cells preventatively from the medicament having cytotoxic properties.

71. A method for preventatively protecting non-cancerous mammalian cells in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which comprises the steps of:

SUBSTITUTE SHEET

2104461

- 138 -

(A) providing a therapeutic composition in an immediate release comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;
5 and

(B) providing an anticancer medicament having cytotoxic properties in timed release form; and

(C) administering the therapeutic composition from step (A) and the medicament from step (B) concurrently
10 to mammalian cells to preventatively protect non-cancerous mammalian cells in the presence of cancerous mammalian cells from the anticancer medicament having cytotoxic properties;

wherein the therapeutic composition is
15 released substantially immediately and the anticancer medicament is released after a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially
20 metabolized the therapeutic composition.

72. A method for selectively protecting non-cancerous mammalian cells prophylactically in the presence of cancerous mammalian cells from an
25 anticancer medicament having cytotoxic properties which comprises the steps of:

(A) administering to mammalian cells a therapeutic composition to prevent and reduce injury to the mammalian cells selected from the group consisting of:

30 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

(b) an antioxidant;

(2) (a) pyruvate selected from the group
35 consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

SUBSTITUTE SHEET

- 139 -

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(3) (a) an antioxidant;

5 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(b) an antioxidant; and

10 (B) waiting a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition; and

15 (C) administering the cytotoxic anticancer medicament to the mammalian cells to treat the cancerous cells which are unprotected by the therapeutic composition and the non-cancerous cells which are protected by the therapeutic composition to thereby increase the therapeutic effect of the
20 anticancer medicament.

73. The method according to claim 72, wherein the medicament having cytotoxic properties is doxorubicin.

25

74. A method for selectively protecting non-cancerous mammalian cells preventatively in the presence of cancerous mammalian cells from an anticancer medicament having cytotoxic properties which
30 comprises the steps of:

(A) administering to mammalian cells a therapeutic composition comprising pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof; and

2104461

- 140 -

(B) waiting a period of time sufficient such that the cancerous cells have substantially metabolized the therapeutic composition and the non-cancerous cells have not substantially metabolized the therapeutic composition; and

(C) administering the cytotoxic anticancer medicament to the mammalian cells to treat the cancerous cells which are unprotected by the therapeutic composition and the non-cancerous cells which are protected by the therapeutic composition to thereby increase the therapeutic effect of the anticancer medicament.

75. A method for increasing the healing rate of a wound in a mammal which comprises the steps of:

(A) providing a wound healing component selected from the group consisting of:

(a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(c) an antioxidant;

(d) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(e) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

2104461

- 141 -

(f) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and an antioxidant;

5 (g) a combination of pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required
10 for the resuscitation of injured mammalian cells;

(h) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and an antioxidant;

15 (i) a combination of lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof, and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required
20 for the resuscitation of injured mammalian cells; and

(j) a combination of an antioxidant and a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

25 (B) contacting the wound healing component from step (A) with the wound to increase the healing rate of the wound.

76. A method for preserving mammalian cells
30 in a culture medium which comprises the steps of:

(A) providing a therapeutic composition selected from the group of consisting of:

(1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable
35 salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

2104461

- 142 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

5 (2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

10 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

15 (3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

20 (4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and

25 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

(B) providing mammalian cells in a culture medium; and

30 (C) contacting the therapeutic composition from step (A) with the mammalian cells in the culture medium from step (B).

- 143 -

5 77. A cytoprotective composition for reversing the cytotoxic effects of ascorbic acid on mammalian cells which comprises ascorbic acid and a therapeutically cytoprotective effective amount of pyruvate.

10 78. The cytoprotective composition according to claim 77, wherein ascorbic acid and pyruvate are present in the composition in a ratio by weight from about 1:10 to about 10:1, respectively.

15 79. A method for reversing the cytotoxic effects of ascorbic acid on mammalian cells which comprises administering to the mammalian cells ascorbic acid and a therapeutically cytoprotective effective amount of pyruvate.

20 80. The method according to claim 79, wherein ascorbic acid and pyruvate are present in the method in a ratio by weight from about 1:10 to about 10:1, respectively.

25 81. A therapeutic pharmaceutical composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which comprises:

(A) a therapeutically effective amount of a therapeutic composition selected from the group consisting of:

30 (1) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

35 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells;

SUBSTITUTE SHEET

2104461

- 144 -

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

5 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
10 injured mammalian cells;

(3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of
15 injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

(b) an antioxidant; and
20 (c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and

(B) a pharmaceutically acceptable carrier.
25

82. The pharmaceutical composition according to claim 81, wherein the pharmaceutically acceptable carrier is a pharmaceutical appliance.

30 83. The pharmaceutical composition according to claim 81, wherein the pharmaceutically acceptable carrier is a topical vehicle.

84. The pharmaceutical composition according to claim 81, wherein the pharmaceutically acceptable
35 carrier is an ingestible vehicle.

- 145 -

85. A method for preparing a therapeutic pharmaceutical composition for preventing and reducing injury to mammalian cells, and increasing the resuscitation rate of injured mammalian cells, which
5 comprises the steps of :

(A) providing a therapeutically effective amount of a therapeutic composition selected from the group consisting of:

(1) (a) pyruvate selected from the group
10 consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

(b) an antioxidant; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are
15 those fatty acids required for the resuscitation of injured mammalian cells;

(2) (a) pyruvate selected from the group consisting of pyruvic acid, pharmaceutically acceptable salts of pyruvic acid, and mixtures thereof;

20 (b) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof; and

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are
25 those fatty acids required for the resuscitation of injured mammalian cells;

(3) (a) an antioxidant; and

(b) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are
30 those fatty acids required for the resuscitation of injured mammalian cells; and

(4) (a) lactate selected from the group consisting of lactic acid, pharmaceutically acceptable salts of lactic acid, and mixtures thereof;

35 (b) an antioxidant; and


2104461

- 146 -

(c) a mixture of saturated and unsaturated fatty acids wherein the fatty acids are those fatty acids required for the resuscitation of injured mammalian cells; and


5 (B) providing a pharmaceutically acceptable carrier; and

(C) admixing the therapeutic composition from step (A) and the pharmaceutically acceptable carrier from step (B) to form a therapeutic pharmaceutical
10 composition.




TREATMENT
FIG. 1

2 / 7



TREATMENT
FIG. 2


3 / 7



TREATMENT
FIG. 3

2104461


4 / 7



TREATMENT
FIG. 4

2104461


5 / 7



TREATMENT
FIG. 5

2104461

6 / 7



TREATMENT
FIG. 6

7 / 7

