



(51) International Patent Classification:

A61K 35/28 (2015.01) *A61P 39/00* (2006.01)
A61P 9/00 (2006.01) *G01N 33/15* (2006.01)

(21) International Application Number:

PCT/US2017/061031

(22) International Filing Date:

10 November 2017 (10.11.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/421,048 11 November 2016 (11.11.2016) US

(71) Applicants: **LONGEVERON LLC** [US/US]; 1951 NW 7th Avenue, Suite 300, Miami, Florida 33136 (US).
UNIVERSITY OF MIAMI [US/US]; 1951 NW 7th Avenue, Miami, Florida 33136 (US).

(72) Inventors: **HARE, Joshua M.**; c/o Longeveron LLC, 1951 NW 7th Avenue, Suite 300, Miami, Florida 33136 (US).
LANDIN, Ana Marie; c/o University of Miami, 1951 NW 7th Avenue, Miami, Florida 33136 (US).

(74) Agent: **HANSEN, Christine M.** et al.; Buchanan Ingersoll & Rooney PC, P.O. Box 1404, Alexandria, Virginia 22313-1404 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: METHODS OF USING HUMAN MESENCHYMAL STEM CELLS TO EFFECT CELLULAR AND HUMORAL IMMUNITY

(57) Abstract: The present invention provides a method of treating subjects with non-ischemic dilated cardiomyopathy by administering a therapeutically effective amount of an isolated population of allogeneic human mesenchymal stem cells. The present invention also provides a method of treating subjects with symptoms of aging frailty by administering a therapeutically effective amount of an isolated population of allogeneic human mesenchymal stem cells.

-1-

Methods of Using Human Mesenchymal Stem Cells to Effect Cellular and Humoral Immunity

FIELD OF THE INVENTION

[0001] The present invention relates to methods of administering therapeutically effective amounts of isolated populations of allogeneic mesenchymal stem cells to effect cellular and humoral immunity in subjects suffering from non-ischemic dilated cardiomyopathy. The present invention also relates to methods of administering therapeutically effective amounts of isolated populations of allogeneic mesenchymal stem cells to effect cellular and humoral immunity in subjects suffering from symptoms of aging frailty.

BACKGROUND OF THE INVENTION

Aging Frailty and Inflammaging

[0002] Aging frailty poses a very concerning problem for the overall health and well-being of individuals and is characterized as a syndrome of multisystem physiological dysregulation. Aging frailty is a geriatric syndrome characterized by weakness, low physical activity, slowed motor performance, exhaustion, and unintentional weight loss. *See* Yao, X. *et al.*, *Clinics in Geriatric Medicine* 27(1):79-87 (2011). Furthermore, there are many studies showing a direct correlation between aging frailty and inflammation. *See* Hubbard, R.E. *et al.*, *Biogerontology* 11(5):635-641 (2010).

[0003] Immunosenescence is characterized by a low grade, chronic systemic inflammatory state known as inflammaging. *See* Franceschi, C. *et al.*, *Annals of the New York Academy of Sciences* 908:244-254 (2000). This heightened inflammatory state or chronic inflammation found in aging and aging frailty leads to immune dysregulation and a complex remodeling of both innate and adaptive immunity. In immunosenescence, the T cell and B cell repertoire is skewed resulting in an increase in CD8⁺ T effector memory cells re-expressing CD45ra (TEMRA) and in the CD19⁺ late/exhausted memory B cells, and a decrease in the CD8⁺ Naïve T cells, and in the switched memory B cells (CD27⁺). *See* Blomberg, B.B. *et al.*, *Immunologic Research* 57(1-3):354-360 (2013); Colonna-Romano, G. *et al.*, *Mechanisms of Ageing and Development* 130(10):681-690 (2009); and Koch S. *et al.*, *Immunity & Ageing*: 5:6 (2008). This shift in the T cell and B cell repertoire results in a

-2-

refractory or less efficient immune status. It is well known that age-associated increase in systemic inflammation (TNF- α , IL-6, IL-8, INF γ and CRP) induces impaired B cell function.

[0004] Inflammaging has received considerable attention because it proposes a link between immune changes and a number of diseases and conditions (such as aging frailty) common in old age. Circulating inflammatory mediators such as cytokines and acute phase proteins are markers of the low-grade inflammation observed to increase with aging. These pro-inflammatory cytokines (*e.g.*, TNF- α , IL-6) impair the capacity of B cells to make protective antibodies to exogenous antigens and vaccines. This impaired B cell response is measured by reduced class switch recombination (CSR) which is the ability of immunoglobulins to switch isotype from IgM to a secondary isotype (IgG, IgA, or IgE). Immunoglobulin isotype switching is crucial for a proper immune response as the effector functions differ in each isotype. A key player in CSR and somatic hypermutation (SHM) is the enzyme, activation-induced cytidine deaminase (AID), encoded by the *Aicda* gene. AID's basic function in CSR and SHM is to initiate breaks in the DNA by converting cytosines to uracils in the switch and variable regions of immunoglobulins. E47, encoded by the *Tcf2a* (E2A) gene, is a transcription factor belonging to the class I basic helix loop helix (bHLH) proteins, also known as E proteins. Without E47 expression, the B cell specific transcription factors EBF1 (early B cell factor) and Pax-5 (paired box protein) are not expressed. Both E47 and Pax-5 are key transcription factors in early development for the B cell lineage and mature B cell function. *See Hagman J. et al., Immunity 27(1):8-10 (2007); Horcher M. et al., Immunity 14(6):779-790 (2001); Riley R.L. et al., Seminars in Immunology 17(5):330-336 (2005).* The Pax-5 gene encodes the B cell lineage specific activator protein (BSAP) that is expressed at all stages of B cell differentiation, but not in terminally differentiated B cells. Pax-5 controls B cell commitment by repressing B lineage inappropriate genes and activating B cell specific genes making Pax-5 the B cell gatekeeper and is exclusively expressed in the B lymphoid lineage from the committed pro-B cell to the mature B cell stage. The B cell specific transcription factor, Pax-5, is not only highly important in early B cell development and B cell lineage commitment, it is also involved in CSR.

[0005] It has also been shown in humans that the amount of TNF- α made: (1) depends on the amount of system inflammation and (2) impairs the ability of the same B cells to be stimulated with mitogens or antigens. *See Frasca, D. et al., Journal of Immunology*

-3-

188(1):279-286 (2012). Thus, the immune response in subjects suffering from aging frailty is impaired for a number of reasons.

Non-Ischemic Dilated Cardiomyopathy (NIDCM)

[0006] Non-ischemic dilated cardiomyopathy (NIDCM) is a progressive disorder with no current cure, often culminating in heart transplantation. See Felker, G.M. *et al.*, *The New England Journal of Medicine* 342:1077-84 (2000) and Kirklin, J.K. *et al.*, *J. Heart Lung Transplant.* 35:407-412 (2016). NIDCM is a disorder with a major component of immune dysregulation as an underlying etiology. See Efthimiadis, I. *et al.*, *Hippokratia* 15:335-342 (2011) and Meng, X. *et al.*, *Nature Reviews Cardiology* 13:167-79 (2016). Cell-based therapy for heart disease is a promising new treatment strategy undergoing evaluation, with a major challenge and opportunity in developing allogeneic therapy. See Patel, A.N. *et al.*, *Lancet* 387:2412-21 (2016); Assmus, B. *et al.*, *The New England Journal of Medicine* 355:1222-32 (2006); Hare, J.M. *et al.*, *J. Am. Coll. Cardiol.* 54:2277-86 (2009); Heldman, A.W. *et al.*, *JAMA* 311:62-73 (2014); Perin, E.C. *et al.*, *JAMA* 307:1717-26 (2012); Golpanian, S. *et al.*, *Physiol. Rev.* 96:1127-68 (2016); and Hare, J.M. *et al.*, *JAMA* 308:2369-79 (2012).

Mesenchymal Stem Cells

[0007] Mesenchymal stem cells are multipotent cells able to migrate to sites of injury, while also being immunoprivileged by not detectably expressing major histocompatibility complex class II (MHC-II) molecules, and expressing MHC-I molecules at low levels. See Le Blanc, K. *et al.*, *Lancet* 371(9624):1579-1586 (2008) and Klyushnenkova E. *et al.*, *J. Biomed. Sci.* 12(1):47-57 (2005). As such, allogeneic mesenchymal stem cells hold great promise for therapeutic and regenerative medicine, and have been repeatedly shown to have a high safety and efficacy profile in clinical trials for multiple disease processes. See Hare, J.M. *et al.*, *Journal of the American College of Cardiology* 54(24):2277-2286 (2009); Hare, J.M. *et al.*, *Tex. Heart Inst. J.* 36(2):145-147 (2009); and Lalu, M.M. *et al.*, *PloS One* 7(10):e47559 (2012). They have also been shown to not undergo malignant transformation after transplantation into patients. See Togel F. *et al.*, *American Journal of Physiology Renal Physiology* 289(1):F31-F42 (2005). Treatment with mesenchymal stem cells has been shown to ameliorate severe graft-versus-host disease, protect against ischemic acute renal failure, contribute to pancreatic islet and renal glomerular repair in diabetes, reverse fulminant hepatic failure, regenerate damaged lung tissue, attenuate sepsis, and reverse remodeling and improve cardiac function after myocardial infarction. See Le Blanc K. *et al.*, *Lancet* 371(9624):1579-1586 (2008); Hare, J.M. *et al.*, *Journal of the American College of Cardiology* 54(24):2277-

2286 (2009); Togel F. *et al.*, *American Journal of Physiology Renal Physiology* 289(1):F31-F42 (2005); Lee R.H. *et al.*, *PNAS* 103(46):17438-17442 (2006); Parekkadan, B. *et al.*, *PloS One* 2(9):e941 (2007); Ishizawa K. *et al.*, *FEBS Letters* 556(1-3):249-252 (2004); Nemeth K. *et al.*, *Nature Medicine* 15(1):42-49 (2009); Iso Y. *et al.*, *Biochem. Biophys. Res. Comm.* 354(3):700-706 (2007); Schuleri K.H. *et al.*, *Eur. Hearth J.* 30(22):2722-2732 (2009); and Heldman A.W. *et al.*, *JAMA* 311(1):62-73 (2014). Furthermore, mesenchymal stem cells are also a potential source of multiple cell types for use in tissue engineering. See Gong Z. *et al.*, *Methods in Mol. Bio.* 698:279-294 (2011); Price, A.P. *et al.*, *Tissue Engineering Part A* 16(8):2581-2591 (2010); and Togel F. *et al.*, *Organogenesis* 7(2):96-100 (2011).

[0008] Mesenchymal stem cells have immuno-modulatory capacity. They control inflammation and the cytokine production of lymphocytes and myeloid-derived immune cells without evidence of immunosuppressive toxicity and are hypo-immunogenic. See Bernardo M.E. *et al.*, *Cell Stem Cell* 13(4):392-402 (2013).

[0009] Mesenchymal stem cells also have the capacity to differentiate not only into cells of mesodermal origin, but into cells of endodermal and ectodermal origin. See Le Blanc K. *et al.*, *Exp. Hematol.* 31(10):890-896 (2003). For example, *in vitro*, mesenchymal stem cells cultured in airway growth media differentiate to express lung-specific epithelial markers, *e.g.*, surfactant protein-C, Clara cell secretory protein, and thyroid transcription factor-1. See Jiang Y. *et al.*, *Nature* 418(6893):41-49 (2002) and Kotton D.N. *et al.*, *Development* 128(24):5181-5188 (2001).

[00010] *In vivo* studies have shown that human mesenchymal stem cells undergo site-specific differentiation into various cell types, including myocytes and cardiomyocytes, when transplanted into fetal sheep. See Airey J.A. *et al.*, *Circulation* 109(11):1401-1407 (2004). These mesenchymal stem cells can persist for as long as 13 months in multiple tissues after transplantation in non-immunosuppressed immunocompetent hosts. Other *in vivo* studies using rodents, dogs, goats, and baboons similarly demonstrate that human mesenchymal stem cells xenografts do not evoke lymphocyte proliferation or systemic allo-antibody production in the recipient. See Klyushnenkova E. *et al.*, *J. Biomed. Sci.* 12(1):47-57 (2005); Aggarwal S. *et al.*, *Blood* 105(4):1815-22 (2005); Augello A. *et al.*, *Arthritis and Rheumatism* 56(4):1175-86 (2007); Bartholomew A. *et al.*, *Exp Hematol.* 30(1):42-48. (2002); Dokic J. *et al.*, *European Journal of Immunology* 43(7):1862-72 (2013); Gerdoni E. *et al.*, *Annals of Neurology* 61(3):219-227 (2007); Lee S.H. *et al.*, *Respiratory Research* 11:16 (2010); Urban V.S. *et al.*, *Stem Cells* 26(1):244-253 (2008); Yang H. *et al.*, *PloS One* 8(7):e69129 (2013);

-5-

Zappia E. *et al.*, *Blood* 106(5):1755-1761 (2005); Bonfield T.L. *et al.*, *American Journal of Physiology Lung Cellular and Molecular Physiology* 299(6):L760-70 (2010); Glenn J.D. *et al.*, *World Journal of Stem Cells*. 6(5):526-39 (2014); Guo K. *et al.*, *Frontiers in Cell and Developmental Biology* 2:8 (2014); Puissant B. *et al.*, *British Journal of Haematology* 129(1):118-129 (2005); and Sun L. *et al.*, *Stem Cells* 27(6):1421-32 (2009). Taken as a whole, these repeated finding of allogeneic safety and efficacy solidify the notion for using mesenchymal stem cells as an allograft for successful tissue regeneration.

[00011] However, despite being a safe therapeutic agent, mesenchymal stem cells are reported in the literature to exert a suppressive effect on antibody production as well as proliferation and maturation of B cells. See Uccelli, A. *et al.*, *Trends in Immunology* 28(5):219-226 (2007). Mesenchymal stem cells are also reported to inhibit the generation and function of antigen presenting cells. See Hoogduijn M.J. *et al.*, *Int. Immunopharmacology* 10(12):1496-1500 (2010). Finally, mesenchymal stem cells are reported to suppress CD4⁺ and CD8⁺ T cell proliferation. See Ghannam S. *et al.*, *Stem Cell Res. & Ther.* 1:2 (2010). Finally, some preclinical data indicates a higher risk of immunological clearance with administration of allogeneic mesenchymal stem cells as compared to the administration of autologous mesenchymal stem cells. See, e.g., Huang, X.P. *et al.*, *Circulation* 122:2419-29 (2010).

SUMMARY

[00012] Surprisingly, despite the reports of mesenchymal stem cells having a suppressive effect on aspects of the immune system, the present inventors discovered a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof. The present inventors also discovered a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof. The present inventors identified several biomarkers specific for non-ischemic dilated cardiomyopathy and aging frailty including, but not limited to, the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺,

-6-

CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁻), the CD4⁺:CD8⁺ T cell ratio, and the TNF- α concentration in serum.

[00013] One aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells (CD19⁺, CD27⁻, IgD⁻) in a sample of the subject's serum decreases by at least 25% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00014] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻) in a sample of the subject's serum increases by at least 100% as compared to the number of switched memory B cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00015] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum decreases by at least 30% as compared to the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00016] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of early activated T-cells (CD3⁺, CD69⁺) in a sample of the subject's serum decreases by at least 30% as compared to the number of early activated T-cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

-7-

[00017] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of chronic activated T-cells ($CD3^+$, $CD25^+$) in a sample of the subject's serum decreases by at least 70% as compared to the number of chronic activated T-Cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00018] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of Temra cells ($CD45RA^+$, $CCR7^-$) in a sample of the subject's serum decreases by at least 40% as compared to the number of Temra cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00019] Another aspect of the invention relates to a method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the TNF- α concentration in a sample of the subject's serum decreases by at least 80% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the non-ischemic dilated cardiomyopathy.

[00020] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-) in a sample of the subject's serum decreases by at least 10% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00021] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) in a sample of the

-8-

subject's serum increases by at least 75% as compared to the number of switched memory B cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00022] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum decreases by at least 60% as compared to the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00023] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of early activated T-cells (CD3⁺, CD69⁺) in a sample of the subject's serum decreases by at least 30% as compared to the number of early activated T-cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00024] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of chronic activated T-cells (CD3⁺, CD25⁺) in a sample of the subject's serum decreases by at least 75% as compared to the number of chronic activated T-Cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00025] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of Temra cells (CD45RA⁺, CCR7⁺) in a sample of the subject's serum decreases by at least 20% as compared to the number of Temra cells in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00026] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the TNF- α concentration in a sample of the subject's serum decreases by at least 50% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00027] Another aspect of the invention relates to a method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the CD4⁺:CD8⁺ T cell ratio in a sample of the subject's serum increases by at least 100% as compared to the CD4⁺:CD8⁺ T cell ratio in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, thereby treating the symptoms of aging frailty.

[00028] In one embodiment of the invention, the subject is a human. In another embodiment of the invention, the subject is a human who exhibits inflammaging.

[00029] In one embodiment of the invention, the mesenchymal stem cells are bone marrow-derived mesenchymal stem cells. In one embodiment of the invention, the mesenchymal stem cells do not express STRO-1. In another embodiment of the invention, the mesenchymal stem cells do not express CD45. In another embodiment of the invention, the mesenchymal stem cells do not express fibroblast surface markers or have a fibroblast morphology. In another embodiment of the invention, the mesenchymal stem cells are not genetically manipulated.

[00030] In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered in a single dose. In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered in multiple doses, *e.g.*, two or more doses. In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered at least yearly.

[00031] In one embodiment, the isolated population of allogeneic mesenchymal stem cells is administered systemically. In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered by infusion or direct injection. In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered intravenously, intraarterially, intramuscularly,

-10-

intraperitoneally, subcutaneously, intradermally, orally, transendocardially, or intranasally. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered intravenously. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered intramuscularly.

[00032] In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 20×10^6 mesenchymal stem cells. In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 100×10^6 mesenchymal stem cells. In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 200×10^6 mesenchymal stem cells.

[00033] In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells are obtained from a human donor and wherein a step of MHC matching of the human donor to the subject is not employed prior to the administration of the isolated population of allogeneic mesenchymal stem cells to the subject.

[00034] Another aspect of the invention relates to a method of evaluating cellular and humoral immunity status in a subject, comprising:

(1) obtaining a serum sample from a subject selected for evaluation based on a determination that the subject was previously in need of treatment of non-ischemic dilated cardiomyopathy and the subject had been administered an initial dose of an isolated population of allogeneic human mesenchymal stem cells;

(2) performing one or more assays configured to detect a non-ischemic dilated cardiomyopathy marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), and serum $TNF-\alpha$ by introducing the serum sample obtained from the subject into an assay instrument which (i) contacts the serum sample with one or more antibodies which specifically bind for detection the biomarker(s) which are assayed, and (ii) generates one or more assay results indicating of binding of each biomarker which is assayed to a respective antibody to provide one or more assay results;

(3) correlating the assay result(s) generated by the assay instrument to the immunity status of the subject, wherein the correlating step comprises assigning a likelihood of one or more future changes in immune status to the subject based on the assay result(s); and

-11-

(4) treating the subject based on the predetermined subpopulation of individuals to which the subject is assigned, wherein the treatment comprises administration of one or more additional doses of an isolated population of allogeneic human mesenchymal stem cells.

[00035] Another aspect of the invention relates to a method of evaluating cellular and humoral immunity status in a subject, comprising:

(1) obtaining a serum sample from a subject selected for evaluation based on a determination that the subject was previously in need of treatment of symptoms of aging frailty and the subject had been administered an initial dose of an isolated population of allogeneic human mesenchymal stem cells;

(2) performing one or more assays configured to detect aging frailty marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), the $CD4^+$: $CD8^+$ T cell ratio, and serum $TNF-\alpha$ by introducing the serum sample obtained from the subject into an assay instrument which (i) contacts the serum sample with one or more antibodies which specifically bind for detection the biomarker(s) which are assayed, and (ii) generates one or more assay results indicating of binding of each biomarker which is assayed to a respective antibody to provide one or more assay results;

(3) correlating the assay result(s) generated by the assay instrument to the immunity status of the subject, wherein the correlating step comprises assigning a likelihood of one or more future changes in immune status to the subject based on the assay result(s); and

(4) treating the subject based on the predetermined subpopulation of individuals to which the subject is assigned, wherein the treatment comprises administration of one or more additional doses of an isolated population of allogeneic human mesenchymal stem cells.

[00036] In one embodiment of the invention, one or more future changes in immune status comprise one or more of an increase in the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), a decrease in the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), an increase in the number of B-cells expressing intracellular $TNF-\alpha$, an increase in the number of early activated T-cells ($CD3^+$, $CD69^+$), an increase in the number of chronic activated T-cells ($CD3^+$, $CD25^+$), an increase in the number of Temra cells ($CD45RA^+$, $CCR7^-$), a decrease in the $CD4^+$: $CD8^+$ T cell ratio, and an increase in serum $TNF-\alpha$.

[00037] Another aspect of the invention relates to an *in vitro* method of determining efficacy of treatment of non-ischemic dilated cardiomyopathy in a subject comprising:

-12-

determining the levels of one or more biomarkers selected from the group consisting of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), and the $TNF-\alpha$ concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if

(1) the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-) decreases by at least 25% as compared to the number of exhausted B cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(2) the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) increases by at least 100% as compared to the number of switched memory B cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(3) the number of B-cells expressing intracellular $TNF-\alpha$ decreases by at least 30% as compared to the number of B-cells expressing intracellular $TNF-\alpha$ prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(4) the number of early activated T-cells ($CD3^+$, $CD69^+$) decreases by at least 30% as compared to the number of early activated T-cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(5) the number of chronic activated T-cells ($CD3^+$, $CD25^+$) decreases by at least 70% as compared to the number of chronic activated T-Cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(6) the number of Temra cells ($CD45RA^+$, $CCR7^-$) decreases by at least 40% as compared to the number of Temra cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells, and/or

(7) the $TNF-\alpha$ concentration in a sample of the subject's serum decreases by at least 80% as compared to the $TNF-\alpha$ concentration in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells.

[00038] Another aspect of the invention relates to an *in vitro* method of determining efficacy of treatment of symptoms of aging frailty in a subject comprising: determining the

-13-

levels of one or more biomarkers selected from the group consisting of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), the $CD4^+$: $CD8^+$ T cell ratio, and the $TNF-\alpha$ concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if

(1) the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-) decreases by at least 10% as compared to the number of exhausted B cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(2) the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) increases by at least 75% as compared to the number of switched memory B cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(3) the number of B-cells expressing intracellular $TNF-\alpha$ decreases by at least 60% as compared to the number of B-cells expressing intracellular $TNF-\alpha$ prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(4) the number of early activated T-cells ($CD3^+$, $CD69^+$) decreases by at least 30% as compared to the number of early activated T-cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(5) the number of chronic activated T-cells ($CD3^+$, $CD25^+$) decreases by at least 75% as compared to the number of chronic activated T-Cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(6) the number of Temra cells ($CD45RA^+$, $CCR7^-$) decreases by at least 20% as compared to the number of Temra cells prior to administration of the population of isolated allogeneic human mesenchymal stem cells,

(7) the $TNF-\alpha$ concentration in a sample of the subject's serum decreases by at least 50% as compared to the $TNF-\alpha$ concentration in a sample of the subject's serum prior to administration of the population of isolated allogeneic human mesenchymal stem cells, and/or

(8) the $CD4^+$: $CD8^+$ T cell ratio in a sample of the subject's serum increases by at least 100% as compared to the $CD4^+$: $CD8^+$ T cell ratio in a sample of the subject's serum

prior to administration of the population of isolated allogeneic human mesenchymal stem cells.

[00039] Another aspect of the invention relates to the use *in vitro* of the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁻), and the TNF- α concentration in serum for determining whether treatment for non-ischemic dilated cardiomyopathy is efficacious.

[00040] Another aspect of the invention relates to the use *in vitro* of the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁻), the CD4⁺:CD8⁺ T cell ratio, and the TNF- α concentration in serum for determining whether treatment for aging frailty is efficacious.

BRIEF DESCRIPTION OF THE FIGURES

[00041] FIG. 1 is a Consort Diagram for a randomized, double-blinded, placebo-controlled study investigating the use of allogeneic mesenchymal stem cells for the treatment of older individuals with frailty.

[00042] FIG. 2 provides baseline patient characteristics for subjects enrolled in the randomized, double-blinded, placebo-controlled study investigating the use of allogeneic mesenchymal stem cells for the treatment of older individuals with frailty.

[00043] FIG. 3 shows a decrease in TNF- α in older subjects administered allogeneic human mesenchymal stem cells.

[00044] FIG. 4 shows a decrease in early activated T cells expressing CD69 in older subjects administered allogeneic human mesenchymal stem cells.

[00045] FIG. 5 shows a decrease in the numbers of chronic/late activated T cells expressing CD25 in older subjects administered allogeneic human mesenchymal stem cells.

[00046] FIG. 6 shows measurements of CD8⁺ T cells in older subjects administered allogeneic human mesenchymal stem cells.

[00047] FIG. 7 shows an increase in the ratio of CD4⁺:CD8⁺ T cells in older subjects administered allogeneic human mesenchymal stem cells.

[00048] FIG. 8 shows an increased immunosenescence score in patients with aging frailty who received two infusions of human mesenchymal stem cells.

-15-

[00049] FIG. 9 shows a correlation between baseline serum TNF- α and baseline B cells expressing IC TNF- α .

[00050] FIG. 10 provides the % of B cells expressing IC TNF- α in Young Control BL, Low TNF:IC TNF BL, and High TNF:IC TNF BL.

[00051] FIG. 11 provides the % of Switch Memory B cells in Young Control BL, Low TNF:SwMem B cells BL, and High TNF:SwMem B cells BL.

[00052] FIG. 12 provides the % of Exhausted B cells in Young Control BL, Low TNF:Exh B cells BL, and High TNF:Exh B cells BL.

[00053] FIG. 13 provides the % of Temra cells in Young Control BL, Low TNF:Temra cells BL, and High TNF:Temra cells BL.

[00054] FIG. 14 provides the CD4⁺/CD8⁺ T cell ratio in Young Control BL, Low TNF:CD4:CD8 BL, and High TNF:CD4:CD8 BL.

[00055] FIG. 15 provides the absolute change of serum TNF- α in pg/ml in Placebo Control, Low TNF- α , and High TNF- α .

[00056] FIG. 16 shows a correlation between baseline serum TNF- α and the change in serum TNF- α .

[00057] FIG. 17 shows the absolute change of B cells expressing IC TNF- α in Placebo Control, Low TNF- α , and High TNF- α .

[00058] FIG. 18 shows a correlation between serum TNF- α and change in B cells expressing IC TNF- α .

[00059] FIG. 19 shows the absolute change of % of Switch Memory B cells in Placebo Control, Low TNF- α , and High TNF- α .

[00060] FIG. 20 shows the absolute change of % of Exhausted B cells in Placebo Control, Low TNF- α , and High TNF- α .

[00061] FIG. 21 shows the absolute change of % of Temra T cells in Placebo Control, Low TNF- α , and High TNF- α .

[00062] FIG. 22 shows a correlation between the change in serum TNF- α and the change in Temra T cells.

[00063] FIG. 23 shows the absolute change in the CD4⁺/CD8⁺ T cell ratio in Placebo Control, Low TNF- α , and High TNF- α .

[00064] FIG. 24 shows a correlation between the change in CD4⁺/CD8⁺ T cell ratio and the change in Temra T cells.

DETAILED DESCRIPTION

[00065] In certain embodiments, the present invention is directed to methods of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof. The examples demonstrate that *in vivo* administration of isolated populations of allogeneic human mesenchymal stem cells result in an increase in the percentage of switched memory B cells and a decrease in exhausted B cells in subjects. The examples also demonstrate that *in vivo* administration of isolated allogeneic human mesenchymal stem cells results in an improvement in the CD4⁺:CD8⁺ T cell ratio in subjects. Also, as shown in the examples, the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁺), and the TNF- α concentration in serum is reduced in subjects having received infusions of allogeneic human mesenchymal stem cells. From these unexpected results, the present inventors determined that isolated allogeneic human mesenchymal stem cells favorably altered several immunologic markers typically elevated in chronic inflammation. Restoration of immune competence has clinical relevance in subjects who are of higher risk for co-morbid infectious disease.

[00066] In other embodiments, the present invention is directed to methods of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof. The examples demonstrate that *in vivo* administration of isolated populations of allogeneic human mesenchymal stem cells result in an increase in the percentage of switched memory B cells and a decrease in exhausted B cells in subjects. The examples also demonstrate that *in vivo* administration of isolated allogeneic human mesenchymal stem cells results in an improvement in the CD4⁺:CD8⁺ T cell ratio in subjects. Also, as shown in the examples, the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁺), and the TNF- α concentration in serum is reduced in subjects having received infusions of allogeneic human mesenchymal stem cells. From these unexpected results, the present inventors determined that isolated allogeneic human mesenchymal stem cells are effective at reducing inflammaging, a prevalent feature in aging frailty.

Definitions

[00067] Embodiments may be practiced without the theoretical aspects presented. Moreover, the theoretical aspects are presented with the understanding that the embodiments are not bound by any theory presented.

[00068] Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

[00069] The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, the singular forms “a”, “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms “including”, “includes”, “having”, “has”, “with”, or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term “comprising.”

[00070] Recitation of ranges of values herein are merely intended to serve as a shorthand method of referring individually to each separate value falling within the range, unless otherwise indicated herein, and each separate value is incorporated into the specification as if it were individually recited herein. The use of any and all examples, or exemplary language (*e.g.*, “such as”) provided herein, is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention unless otherwise claimed. No language in the specification should be construed as indicating any non-claimed element as essential to the practice of the invention.

[00071] The term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system. For example, “about” can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, “about” can mean a range of $\pm 10\%$ of the referenced value.

Dosage, Duration and Subjects

[00072] “A therapeutically effective amount” means an amount that stimulates a B- or T-cell dependent immune response. Such a response is characterized by the ability to elicit

-18-

significant levels of IgG and opsonic activity. The dosage and number of doses (*e.g.*, single or multiple dose) administered to the subject will vary depending upon a variety of factors, including the route of administration, patient conditions and characteristics (sex, age, body weight, health, size), extent of symptoms, concurrent treatments, frequency of treatment and the effect desired, and the like.

[00073] In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered as a single dose. In another embodiment, the isolated population of allogeneic mesenchymal stem cells is administered in multiple doses, *e.g.*, two or more doses. In other embodiments, the isolated population of allogeneic mesenchymal stem cells is administered at least yearly.

[00074] In another embodiment of the invention, the administration of the isolated population of allogeneic mesenchymal stem cells is repeated, such as at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18 months after the first administration of the isolated population of allogeneic mesenchymal stem cells, or repeated between 2-4, 2-6, 2-8, 2-10, 3-4, 3-6, 3-8, 3-10, 4-6, 4-8, 4-10, 6-8, 6-10, 6-12, or 12-18 months after the first administration of the isolated population of allogeneic mesenchymal stem cells.

[00075] In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 1×10^6 , 2×10^6 , 5×10^6 , 10×10^6 , 20×10^6 , 30×10^6 , 40×10^6 , 50×10^6 , 60×10^6 , 70×10^6 , 80×10^6 , 90×10^6 , 100×10^6 , 110×10^6 , 120×10^6 , 130×10^6 , 140×10^6 , 150×10^6 , 160×10^6 , 170×10^6 , 180×10^6 , 190×10^6 , 200×10^6 , 300×10^6 , 400×10^6 , 500×10^6 , or 10×10^7 mesenchymal stem cells. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 20×10^6 mesenchymal stem cells. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 100×10^6 mesenchymal stem cells. In yet a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 200×10^6 mesenchymal stem cells. In further embodiments, the isolated population of allogeneic mesenchymal stem cells is administered at a dose of from about $1-400 \times 10^6$, $10-400 \times 10^6$, $100-400 \times 10^6$, $20-200 \times 10^6$, $20-400 \times 10^6$, $0.1-5 \times 10^6$, $0.1-10 \times 10^6$, $0.1-100 \times 10^6$, $1-50 \times 10^6$, $1-100 \times 10^6$, $0.01-10 \times 10^6$ or $0.01-100 \times 10^6$ mesenchymal stem cells.

[00076] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to increase the ratio of $CD4^+ : CD8^+$ T cells in a subject, such as to increase the ratio of $CD4^+ : CD8^+$ T cells by at least

-19-

two-, three-, four-, five-, or six-fold as compared to the ratio prior to administration of the isolated population of allogeneic mesenchymal stem cells, *i.e.*, an increase in the ratio of CD4⁺:CD8⁺ T cells by at least 100%, 200%, 300%, 400%, or 500%.

[00077] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to increase the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻) in a subject, such as to increase the number of switched memory B cells by at least two-, three-, four-, or five-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells, *i.e.*, an increase in the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻) by at least 100%, 200%, 300%, or 400%. In other embodiments, the number of switched memory B cells as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells increases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%.

[00078] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of B cells expressing intracellular TNF- α in a subject, such as to decrease the number by at least two-, three-, four-, five-, or six-fold as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of B cells expressing intracellular TNF- α as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, 50%, or 60%.

[00079] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of exhausted B cells (CD19⁺, CD27⁻, IgD⁻) in a subject, such as to decrease the number of exhausted B cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of exhausted B cells as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%.

[00080] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of early activated T-cells (CD3⁺, CD69⁺) in a subject, such as to decrease the number of early

-20-

activated T-cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of early activated T-cells as compared to the number of such T cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, or 50%.

[00081] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of chronic activated T-cells ($CD3^+$, $CD25^+$) in a subject, such as to decrease the number of chronic activated T-cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of chronic activated T-cells as compared to the number of such T cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%.

[00082] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of Temra cells ($CD45RA^+$, $CCR7^-$) in a subject, such as to decrease the number of Temra cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of Temra cells as compared to the number of such Temra cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, or 50%.

[00083] In some embodiments, the therapeutically effective amount of the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the TNF- α concentration in a sample of the subject's serum, such as to decrease the TNF- α concentration by at least two- or three-fold as compared to the TNF- α concentration prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the concentration of TNF- α decreases by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%, as compared to the concentration of TNF- α prior to administration of the isolated population of allogeneic mesenchymal stem cells.

[00084] Other aspects of the invention include methods for evaluating the cellular and humoral immunity status in subjects to determine if the subjects would benefit from the administration of an isolated population of allogeneic mesenchymal stem cells and treating a

-21-

predetermined subpopulation of subjects based on biomarker data. For example, one or more assays could be performed to detect a non-ischemic dilated cardiomyopathy marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), and serum $TNF-\alpha$. Similarly, one or more assays could be performed to detect an aging frailty marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), the $CD4^+$: $CD8^+$ T cell ratio, and serum $TNF-\alpha$.

[00085] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to increase the ratio of $CD4^+$: $CD8^+$ T cells in a subject, such as to increase the ratio of $CD4^+$: $CD8^+$ T cells by at least two-, three-, four-, five-, or six-fold as compared to the ratio prior to administration of the isolated population of allogeneic mesenchymal stem cells, *i.e.*, an increase in the ratio of $CD4^+$: $CD8^+$ T cells by at least 100%, 200%, 300%, 400%, or 500%.

[00086] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to increase the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) in a subject, such as to increase the number of switched memory B cells by at least two-, three-, four-, or five-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells, *i.e.*, an increase in the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) by at least 100%, 200%, 300%, or 400%. In other embodiments, the number of switched memory B cells as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells increases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%.

[00087] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of B cells expressing intracellular $TNF-\alpha$ in a subject, such as to decrease the number by at least two-, three-, four-, five-, or six-fold as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of B cells expressing intracellular $TNF-\alpha$ as compared to the

-22-

number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, 50%, or 60%.

[00088] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-) in a subject, such as to decrease the number of exhausted B cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of exhausted B cells as compared to the number of such B cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%.

[00089] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of early activated T-cells ($CD3^+$, $CD69^+$) in a subject, such as to decrease the number of early activated T-cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of early activated T-cells as compared to the number of such T cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, or 50%.

[00090] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of chronic activated T-cells ($CD3^+$, $CD25^+$) in a subject, such as to decrease the number of chronic activated T-cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of chronic activated T-cells as compared to the number of such T cells prior to administration of the isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%.

[00091] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the number of Temra cells ($CD45RA^+$, $CCR7^-$) in a subject, such as to decrease the number of Temra cells by at least two- or three-fold as compared to the number prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the number of Temra cells as compared to the number of such Temra cells prior to administration of the

isolated population of allogeneic mesenchymal stem cells decreases by at least 10%, 20%, 30%, 40%, or 50%.

[00092] In some embodiments, treatment is demonstrated to be efficacious when the isolated population of allogeneic mesenchymal stem cells is sufficient to decrease the TNF- α concentration in a sample of the subject's serum, such as to decrease the TNF- α concentration by at least two- or three-fold as compared to the TNF- α concentration prior to administration of the isolated population of allogeneic mesenchymal stem cells. In other embodiments, the concentration of TNF- α decreases by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%, as compared to the concentration of TNF- α prior to administration of the isolated population of allogeneic mesenchymal stem cells.

[00093] "Administering" a composition may be accomplished by oral administration, injection, infusion, parenteral, intravenous, mucosal, sublingual, intramuscular, intradermal, intranasal, intraperitoneal, intraarterial, subcutaneous absorption or by any method in combination with other known techniques. In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered systemically. In another embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered by infusion or direct injection. In one embodiment of the invention, the isolated population of allogeneic mesenchymal stem cells is administered intramuscularly, intravenously, intraarterially, intraperitoneally, subcutaneously, intradermally, orally, transendocardially, or intranasally. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered intramuscularly. In a further embodiment, the isolated population of allogeneic mesenchymal stem cells is administered intravenously.

[00094] The term "subject" as used herein includes, but is not limited to, humans and non-human vertebrates such as wild, domestic, and farm animals. In some embodiments, the term refers to humans, such as elderly humans ≥ 65 years of age, or elderly humans 60-95 years of age. In some embodiments, the human subject exhibits symptoms of aging frailty. In some embodiments, the human subject exhibits inflammaging.

[00095] The term "allogeneic" refers to a cell that is of the same animal species but genetically different in one or more genetic loci as the animal that becomes the "recipient host." This usually applies to cells transplanted from one animal to another non-identical animal of the same species.

-24-

[00096] As used herein, the phrase “in need thereof” means that the subject has been identified as having a need for the particular method or treatment. In some embodiments, the identification can be by any means of diagnosis. In any of the methods and treatments described herein, the subject can be in need thereof. In some embodiments, the subject is in an environment or will be traveling to an environment in which a particular disease, disorder, or condition is prevalent.

[00097] Cells are referred to herein as being positive or negative for certain markers. For example, a cell can be negative for CD45, which can also be referred to as CD45⁻. The superscript notation “⁻” refers to a cell that is negative for the marker linked to the superscript. In contrast a marker with the “⁺” refers to a cell that is positive for that marker. For example, a cell that is referenced as “CD8⁺” is positive for CD8. A “+” can also be used to reference the marker as positive. A “-” can also be used to reference the marker as negative.

[00098] As used herein, the term “stem cell” refers to a cell from the embryo, fetus, or adult that has, under certain conditions, the ability to reproduce itself for long periods or, in the case of adult stem cells, throughout the life of the organism. It also can give rise to specialized cells that make up the tissues and organs of the body.

[00099] Mesenchymal stem cells are the formative pluripotent blast cells found *inter alia* in bone marrow, blood, dermis, and periosteum that are capable of differentiating into any kind of the specific types of mesenchymal or connective tissues (*i.e.*, the tissues of the body that support the specialized elements; particularly adipose, osseous, cartilaginous, elastic, and fibrous connective tissues) depending upon various influences from bioactive factors, such as cytokines.

[000100] Certain methods of isolating and/or purifying mesenchymal stem cells have been described herein and are known in the art. In some embodiments, mesenchymal stem cells are isolated from bone marrow of adult humans. In some embodiments, the cells are passed through a density gradient to eliminate undesired cell types. The cells can be plated and cultured in appropriate media. In some embodiments, the cells are cultured for at least one day or about three to about seven days, and removing non-adherent cells. The adherent cells can then be plated and expanded.

[000101] Other methods for isolating and culturing stem cells are also known. Placenta is an excellent readily available source for mesenchymal stem cells. Moreover, mesenchymal stem cells can be derivable from adipose tissue and bone marrow stromal cells are speculated

-25-

to be present in other tissues. While there are dramatic qualitative and quantitative differences in the organs from which adult stem cells can be derived, the initial differences between the cells may be relatively superficial and balanced by the similar range of plasticity they exhibit.

[000102] Homogeneous human mesenchymal stem cell compositions are provided which serve as the progenitors for all mesenchymal cell lineages. Mesenchymal stem cells are identified by specific cell surface markers which are identified with unique monoclonal antibodies. The homogeneous mesenchymal stem cell compositions are obtained by positive selection of adherent marrow or periosteal cells which are free of markers associated with either hematopoietic or differentiated mesenchymal cells. These isolated mesenchymal cell populations display epitopic characteristics associated with only mesenchymal stem cells, have the ability to regenerate in culture without differentiating, and have the ability to differentiate into specific mesenchymal lineages when either induced *in vitro* or placed *in vivo* at a site of inflammation.

[000103] In order to obtain the human mesenchymal stem cells for the compositions, methods, and kits disclosed herein, pluripotent mesenchymal stem cells are separated from other cells in the bone marrow or other mesenchymal stem cell source. Bone marrow cells may be obtained from iliac crest, femora, tibiae, spine, rib, or other medullary spaces. Other spaces of human mesenchymal stem cells include embryonic yolk sac, placenta, umbilical cord, fetal and adolescent skin, and blood.

[000104] In some embodiments, the human mesenchymal stem cells are identified by the absence of markers. For example, human mesenchymal stem cells useful in the invention include those that are negative for STRO-1 and/or negative for CD45. Similarly, human mesenchymal stem cells useful in the invention include those that do not express fibroblast surface markers or have a fibroblast morphology.

Methods of Enhancing Immune Responses

[000105] As discussed above, the present invention is directed to a method of enhancing a subject's cellular or humoral immune response, comprising administering to the subject therapeutically effective amounts of an isolated population of human mesenchymal stem cells. In some embodiments of the invention, the mesenchymal stem cells are not genetically manipulated. In some embodiments of the invention, the mesenchymal stem cells are obtained from a human donor and wherein a step of MHC matching of the human donor to

-26-

the subject is not employed prior to the administration of the isolated population of human mesenchymal stem cells.

[000106] Compositions for use in the invention may be formulated using any suitable method. Formulation of cells with standard pharmaceutically acceptable carriers and/or excipients may be carried out using routine methods in the pharmaceutical art. The exact nature of a formulation will depend upon several factors including the cells to be administered and the desired route of administration. Suitable types of formulation are fully described in Remington's Pharmaceutical Sciences, 19th Edition, Mack Publishing Company, Eastern Pennsylvania, USA.

[000107] Compositions may be prepared together with a physiologically acceptable carrier or diluent. Typically, such compositions are prepared as liquid suspensions of cells. The cells may be mixed with an excipient which is pharmaceutically acceptable and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, of the like and combinations thereof

[000108] In addition, if desired, the pharmaceutical compositions of the invention may contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, and/or adjuvants which enhance effectiveness. In one embodiment of the invention, the adjuvant comprises human serum albumin (HSA).

[000109] One suitable carrier or diluent is PlasmaLyte ATM. This is a sterile, nonpyrogenic isotonic solution for intravenous administration. Each 100 mL contains 526 mg of Sodium Chloride, USP (NaCl); 502 mg of Sodium Gluconate (C₆H₁₁NaO₇); 368 mg of Sodium Acetate Trihydrate, USP (C₂H₃NaO₂·3H₂O); 37 mg of Potassium Chloride, USP (KCl); and 30 mg of Magnesium Chloride, USP (MgCl₂·6H₂O). It contains no antimicrobial agents. The pH is adjusted with sodium hydroxide. The pH is 7.4 (6.5 to 8.0).

[000110] In one embodiment of the invention the mesenchymal stem cells are not genetically manipulated. In another embodiment of the invention, the mesenchymal stem cells are cryopreserved. For example, the mesenchymal stem cells can be suspended in cryoprotectant consisting of Hespan® (6% hetastarch in 0.9% sodium chloride) supplemented with 2% HSA and 5% DMSO and then aliquoted into cryopreservation containers for placement in vapor phase nitrogen freezers. In another embodiment, the mesenchymal stem cells may be provided in PlasmaLyte ATM supplemented with 1% HSA.

EXAMPLES

-27-

Example 1

Treatment of Chronic Non-Ischemic Dilated Cardiomyopathy

[000111] Briefly, thirty-seven patients were randomized to receive either allogeneic or autologous human mesenchymal stem cells in a 1:1 ratio. Patients were recruited between December 2011 and July 2015 at the University of Miami Hospital. Patients (age: 55.8 ± 11.2 ; 32% female) received human mesenchymal stem cells (100 million) by transendocardial stem cell injection (TESI) in ten left ventricular sites by NOGA Catheter. Treated patients were evaluated at baseline, 30 days, 3-, 6- and 12-months for immune biomarkers.

Patient Population

[000112] Patient eligibility was determined after confirmation of diagnosis of NIDCM with an EF of less than 40% and either a left ventricular end diastolic diameter greater than 5.9 cm in male subjects and greater than 5.6 cm in female subjects or a left ventricular end diastolic volume index $> 125 \text{ mL/m}^2$. Thirty-seven patients were randomized to either autologous human mesenchyme stem cells (hMSCs) or allogeneic hMSCs in a ratio of 1:1. Thirty-four patients received study injection; 16 and 18 patients in the auto- and allo-hMSC, respectively. Three patients did not receive the study injection. One patient withdrew consent before treatment. Another patient was recruited but did not receive treatment due to automatic implantable cardioverter-defibrillator placement ($n=1$), and one patient died before treatment ($n=1$). The mean age of injected participants was 55.8 ± 11.2 , 29% were female, and 35% were Hispanic. The mean years of NIDCM diagnosis before the TESI was 6.1 ± 6.2 years for allo and 6.9 ± 7.3 years for auto patients ($p=0.5$ between groups). Fifty percent of patients had NYHA class II symptoms, mean baseline global EF was $26.5 \pm 9.64\%$, mean 6MWT was $422 \pm 86.8 \text{ M}$, and median baseline MLHFQ scores were 36 (IQR 18.0, 64.0).

Study Procedures and Timeline

[000113] Baseline assessments included chemistry and hematology laboratories, echocardiography, and chest, abdominal and pelvic computed tomography scans.

Human Mesenchymal Stem Cells (hMSCs) for Cell Therapy

[000114] All allogeneic and autologous human mesenchymal stem cells were manufactured at the University of Miami ISCI. See, e.g., Golpanian, S. *et al.*, *Physiol. Rev.* 96:1127-68 (2016) and Mushtaq, M. *et al.*, *J. Cardio. Trans. Res.* 7:769-80 (2014). Allogeneic human mesenchymal stem cells were derived from Caucasian male donors mean age 25.4 ± 3.3 years and were between 80 to 90% viable at the time of TESI. The auto-

-28-

hMSCs were from 11 males with a mean age of 58.0 ± 9.9 and six females with a mean age of 55.0 ± 12.4 years.

Transendocardial Stem Cell Injection

[000115] Injection sites were selected to prioritize safety of the TESI procedure and to distribute sites throughout the accessible myocardial territories. Considerations for site selection included avoidance of the ventricular apex, and optimization of catheter stability prior to needle extension.

Immune Monitoring

[000116] Calculated panel reactive antibodies (cPRA) were measured at baseline and at six-months using Luminex 200. Serum TNF- α was measured using human TNF- α ELISA high sensitivity kit (eBiosciences). Lymphocytes were stained for T-cell markers of activation. Late/exhausted T-cells, B-cell subsets (Switched memory and Late/Exhausted B-cells) and TNF- α by B-cells. All samples were acquired using the LSR-Fortessa-HTS analyzer (BD Pharmingen) and analyzed with the FlowJo V10 software, *i.e.*, all lymphocyte populations were assessed by flow cytometry and expressed as a percent from gated (B-cell parameters gated on CD19⁺ and T-cell parameters gated on CD3⁺).

[000117] cPRA results showed that 67% of allo and 92% of auto recipients had no reaction to low cPRA (0-20% cPRA). Twenty-seven % of allo and 8% of auto had a moderate cPRA (21-79% cPRA), and one subject (7%) receiving allo MSCs had a high cPRA response (+80% cPRA).

[000118] Elevated baseline levels of TNF- α decreased from baseline to 6-months in both groups (allo: -10.6 ± 1.6 pg/ml, $p < 0.0001$, auto: -6.8 ± 1.4 pg/ml, $p < 0.0001$, Table 1; between group, $p = 0.05$). Temra T-cells (exhausted T-cell phenotype) were reduced in both groups with a greater decrease in allo (allo: $-15.9 \pm 5.4\%$, $p < 0.0001$, auto: $-9.3 \pm 3.3\%$, $p < 0.0001$, Table 1; between group, $p = 0.0111$; Table 1). Suppressed %Switch Memory B-cells (predictive biomarker for antibody response) at baseline were significantly increased at six-months in both groups with a greater improvement in allo (allo: $+10.2 \pm 4.9\%$, $p < 0.0001$ vs. auto: $+4.3 \pm 3.9\%$, $p = 0.0014$, between group, $p < 0.0001$; Table 1). Finally, intracellular TNF- α expression in B23 cells was also decreased at 6-months relative to baseline in both groups (allo: $-11.2 \pm 3.3\%$, $p < 0.0001$, auto: $-8.5 \pm 3.0\%$, $p < 0.0001$; between group, $p = 0.174$; Table 1). In contrast, Late/exhausted B-cells decreased significantly in both groups (allo: $-5.4 \pm 1.03\%$, $p < 0.0001$, auto: $-5.9 \pm 2.5\%$, $p = 0.003$; between group difference, $p = 0.57$; Table 1). Early T-cell activation (allo: $-5.57 \pm 1.03\%$, $p < 0.0001$, auto: $-2.92 \pm 1.5\%$, $p = 0.02$; between

-29-

group, $p=0.08$; Table 1) decreased to similar degrees in both group. Whereas late/chronic T-cell activation did not significantly decrease in either group (allo: $-2.3 \pm 1.3\%$, $p=0.4$, and auto: $-3.4 \pm 2.7\%$, $p=0.7$).

Table 1. Effects of Human Mesenchymal Stem Cells on Cellular and Humoral Immunity at 6-months.

Immune Biomarkers	Allogeneic		Autologous	
	Baseline	6-Months	Baseline	6-Months
Serum TNF- α (pg/ml)	13.5 ± 1.5	$2.3 \pm 0.2^{**,\dagger}$	11.8 ± 0.9	$4.8 \pm 0.8^{**}$
%Early T-cell Activation (CD3 ⁺ , CD69 ⁺)	13.4 ± 4.4	$7.9 \pm 3.3^{**}$	13.3 ± 4.8	$10.6 \pm 5.4^*$
%Late/Chronic T-cell activation (CD3 ⁺ , CD25 ⁺)	8.9, IQR (5.9, 10.4)	3.4, IQR (3.0, 5.0)*	8.1, IQR (5.2, 14.1)	4.4, IQR (3.6, 11.1)
% Temra (CD3 ⁺ , CD45RA ⁺ , CCR7)	33.7 ± 7.6	$17.9 \pm 5.2^{**,\dagger}$	30.1 ± 10.8	$21.3 \pm 7.5^{**}$
%Late/Exhausted B-cells (CD19 ⁺ , CD27 ⁺ , IgD ⁺)	19.3, IQR (17.9, 28.6)	14.7, IQR (14.2, 17.3)** [†]	20.0, IQR (17.7, 32.0)	15.5, IQR (14.0, 19.0)*
%Switched Memory B-cells (CD19 ⁺ , CD27 ^{high} , IgD ⁺)	10.0 ± 3.6	$20.2 \pm 3.1^{**,\dagger}$	9.6 ± 3.1	$14.1 \pm 4.5^*$
% B-cells expressing intracellular TNF- α	32.1 ± 7.3	$20.4 \pm 6.3^{**}$	28.2 ± 6.8	$19.7 \pm 4.8^{**}$

* indicates $p \leq 0.05$ within group; ** indicates $p \leq 0.001$ within group; [†] indicates between group $p \leq 0.5$; [‡] indicates between group $p \leq 0.001$.

Example 2

Treatment of Symptoms of Aging Frailty – Study 1

-30-

[000119] Thirty subjects aged 60 years of age or older who met the study inclusion/exclusion criteria for frailty by the Canadian Study of Health and Aging (CSHA) were randomized. Ten subjects received 100 million MSCs, ten subjects received 200 million MSCs, and ten subjects received placebo. Subjects were followed for one year for safety and efficacy. Assessments included at least the following: a study of inflammatory biomarkers and a review of immunologic effects. FIG. 1 is a Consort Diagram. FIG. 2 provides baseline patient characteristics. FIGs. 3-7 illustrate the impact of allogeneic human MSCs on immune biomarkers.

[000120] Figure 3 shows a decrease TNF- α in subjects administered human MSCs. Both early and late/chronic T cell activation decreases after allogeneic MSC treatment. FIG. 4 shows a decrease in early activated T cells expressing CD69 in subjects administered human MSCs. FIG. 5 shows a decrease in the numbers of chronic/late activated T cells expressing CD25 in subjects administered human MSCs. FIG. 7 shows an increase in the ratio of CD4⁺/CD8⁺ T cells in subjects administered human MSCs.

Example 3

Treatment of Symptoms of Aging Frailty – Study 2

[000121] Fifteen aged subjects were included in an open-label study and administered human mesenchymal stem cells in two separate infusions. Subjects were evaluated for changes in various immune biomarkers to determine a response to the injected human mesenchymal stem cells. These immune biomarkers included the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7⁻), the CD4⁺:CD8⁺ T cell ratio, and the TNF- α concentration in serum. Subjects were evaluated 6 months after the first injection and 1, 3, and 6 months after the second injection. Provided in Tables 2-9 are the measured values for each of the immune biomarkers discussed above. Also provided in Tables 2-9 are Scores assigned to each of the immune biomarkers assessed in each patient at different time-points that indicate whether an increase or decrease was observed in the immune biomarker measured. Finally, in Table 10 are the total Scores for each patient that includes different time-points as well as averages among the patients. As can be observed in the last row of Table 10 and in FIG. 8, the immunosenescence Score improves in subjects with aging frailty after the first injection of

-31-

human mesenchymal stem cells and continues to stay in an improved state after a second injection of human mesenchymal stem cells.

Table 2. Effects of Human Mesenchymal Stem Cells on Switched Memory B Cells

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	6.61	21.2	14.2	16.5	22.8	21.3	1	2	2	2	2	2
2	21.3	41	24.9	24	25.4	30.8	2	4	3	3	3	3
3	5.34	9.31	7.14	16.6	7.58	9.81	1	1	1	2	1	1
4	10.5	17	30.1	36.6	18.7	40.3	1	2	3	4	2	2
5	10.7	24	25.	33.3	28.8	31.5	1	3	3	3	3	3
6	7.51	14.6	1.7	14.7	19.7	19.8	1	2	2	2	2	2
7	7.32	20.7	12.3	13.7	21.2	27.1	1	2	2	2	2	2
8	7.69	9.89	8.31	11.7	26.4	21.5	1	1	1	2	2	2
9	10.1	17.8	15.8	15.3	25.8	35.9	1	2	2	2	3	2
10	21.7	35.3	23.1	23.6	23.8	25.7	2	4	2	3	3	3
15	9.25	13.5	22.3	28.1	28.1	29.7	1	2	2	3	3	3

Improvement Scale: No Change=0; Improved $>0 \leq 1$ fold=1; Improved $>1 \leq 2$ fold=2; Improved $>2 \leq 3$ fold=3; Improved >3 fold=4; Worsened $>0 \leq 1$ fold=-1; Worsened $>1 \leq 2$ fold=-2; Worsened $>2 \leq 3$ fold=-3; and Worsened >3 fold=-4.

Table 3. Effects of Human Mesenchymal Stem Cells on %B-Cells Expressing Intracellular TNF- α

-32-

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	23.4	9.65	15.6	9.59	5.35	8.21	1	3	2	2	4	3
2	34.8	9.33	10	9.83	6.88	6.71	1	3	2	3	3	3
3	14.3	7.62	40.4	21.2	15.7	8.02	2	3	1	2	3	3
4	14.3	7.63	13.7	12	5.25	3.63	2	2	2	2	4	4
5	19.5	10.9	9.52	8.89	8.33	7.59	2	2	2	3	3	3
6	37.7	1.22	5.57	4	4.04	4.33	1	4	4	4	4	4
7	13.2	2.71	4.1	3.67	4.42	2.52	2	4	4	4	4	4
8	30.8	4.66	6.82	7.01	7.05	7.42	1	4	3	3	3	3
9	11.1	6.94	11.2	5.76	7.79	5.1	2	3	2	4	3	4
10	13.5	4.2	3.55	3.87	3.72	3.42	2	4	4	4	4	4
15	7.05	4.75	5.75	3.84	6.46	2.04	3	4	4	4	4	4

Table 4. Effects of Human Mesenchymal Stem Cells on Exhausted B Cells

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	23.2	40.4	15.7	9.88	10.7	15.6	1	1	2	3	3	2
2	26.8	27.6	6.59	3.33	6.67	7.51	1	1	4	4	4	4

-33-

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
3	7.63	18.2	15.7	15.7	9.01	14.2	4	2	2	2	3	2
4	29.1	14.8	11.9	7.12	9.86	10.1	1	2	3	4	3	3
5	53.3	15.2	19.2	8.39	13.6	12.9	1	2	2	4	2	3
6	34.6	14.1	9.47	9.09	8.01	5.26	1	2	3	3	4	4
7	44.1	51.9	66.5	57.2	43.1	40.3	1	1	1	1	1	1
8	9.23	7.1	12.6	11.6	8.05	12	3	4	3	3	4	3
9	15.4	17.8	15.6	7.85	9.51	17.8	2	2	2	4	3	2
10	12.5	24.6	7.17	11.9	13.4	7.42	2	2	4	3	2	4
15	32.4	26.6	24.4	16.2	11.7	12.4	1	1	2	2	3	3

Table 5. Effects of Human Mesenchymal Stem Cells on Temra Cells

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	30.1	42.6	14.6	17.4	19.4	15.4	1	1	2	2	1	2
2	25.6	27.8	25.2	27	26.4	37.1	1	1	1	1	1	1
3	7.55	2.85	14.4	13.3	10.1	11.8	3	4	2	2	2	2
4	15.7	3.06	15	12.7	9.5	12.5	2	4	2	2	3	2

-34-

Patient #	BL	6-Mo	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
5	10	2.85	9.86	8.43	11	10.4	2	4	2	3	2	2
6	39.5	18.2	21.6	16.4	20	18.1	1	2	1	2	1	2
7	13.3	14.2	11.4	10.3	15	17.7	2	2	2	2	2	2
8	31.5	12.2	11.8	7.09	11.6	33.3	1	2	2	3	2	1
9	16.1	11.1	24	25.1	34.3	38.8	2	2	1	1	1	1
10	7.15	4.89	12.2	19	9.52	13.2	3	4	2	2	3	2
15	15.2	13.3	30.2	40.6	46.1	34.9	2	2	1	1	1	1

-35-

Table 6. Effects of Human Mesenchymal Stem Cells on Early Activation of T Cells (CD69)

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	16	16.6	12.98	8.03	8.74	6.07	2	2	2	3	3	4
2	5.8	7.5	10.31	13.82	10.77	12.21	4	4	3	2	3	2
3	29	13	14.76	3.36	4.83	9.14	1	2	2	4	4	3
4	65	30	7.87	5	4.33	17.07	1	1	3	4	4	2
5	4.4	11.7	13.15	3.42	20.88	6.75	4	2	2	4	2	4
6	42	22.4	13.23	13.56	8.6	13.29	1	2	2	2	3	2
7	25.8	22.1	13.73	10.26	10.87	37.35	1	2	2	3	3	1
8	28.3	17.1	4.22	5.1	4.93	12.32	1	2	4	4	4	2
9	4.4	6.2	20.12	7.63	12.06	11.9	4	4	2	3	2	2
10	19.4	17.9	14.18	13.53	6.22	13.85	2	2	2	2	4	2
15	10.9	6.18	14.18	12.79	14.2	13.1	3	4	2	2	2	2

Table 7. Effects of Human Mesenchymal Stem Cells on Late/Chronic Activation of T Cells (CD25)

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	1.7	2.9	5.74	8.32	10.98	7.63	4	4	3	2	2	2
2	2.2	3	4.25	4.69	3.84	4.7	4	4	3	3	4	3
3	20	5	10.83	4.93	14.32	5.87	1	3	2	3	1	3
4	53	3.2	4.54	5.88	7.44	6.97	1	4	3	3	2	2
5	10	1.9	5.69	7.9	7.91	8.9	2	4	3	2	2	2

-36-

Patient #	BL	6-Mo.	2 nd Inject . BL	2 nd Inject . 1-Mo.	2 nd Inject . 3-Mo.	2 nd Inject . 6-Mo.	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
6	16	1.4	5.69	6.95	6.69	2.53	1	4	3	2	2	4
7	0.9	2.3	7.35	8.38	6.77	8.19	4	4	2	2	2	2
8	20.5	3.7	6.81	9.41	10.74	4.26	1	4	2	2	2	3
9	2.1	1.3	3	4.47	3.88	3.82	4	4	4	3	4	4
10	6.6	2.9	11.61	5.29	3.88	5.08	2	4	2	3	4	3
15	1.4	1.11	11.61	2.21	3.14	2.11	4	4	2	4	4	4

Table 8. Effects of Human Mesenchymal Stem Cells on Ratio of CD4⁺:CD8⁺ T Cells

Patient #	BL Ratio	6-Mo. Ratio	2 nd Inject . BL Ratio	2 nd Inject . 1-Mo. Ratio	2 nd Inject . 3-Mo. Ratio	2 nd Inject . 6-Mo. Ratio	BL Score	6-Mo. Score	2 nd Inject . BL Score	2 nd Inject . 1-Mo. Score	2 nd Inject . 3-Mo. Score	2 nd Inject . 6-Mo. Score
1	0.77	0.69	0.48	1.03	0.68	0.66	1	1	1	1	1	1
2	2.17	2.48	0.98	1.16	1.48	0.96	2	2	1	1	2	1
3	1.66	1.94	2.24	1.95	2.98	2.32	2	2	2	2	3	2
4	0.96	3.43	6.49	2.18	3.67	4.59	1	3	4	2	3	4
5	2.34	5.32	5.05	4.27	4.45	3.79	2	4	4	4	4	3
6	0.94	3.47	4.38	3.54	3.50	4.42	1	3	4	2	3	4
7	0.51	1.60	0.95	0.92	0.92	0.97	1	2	1	1	1	1
8	0.83	2.26	2.49	2.49	2.58	0.21	1	2	2	2	2	1
9	1.57	2.68	1.56	2.48	1.25	0.67	2	2	2	2	1	1
10	3.02	4.76	3.97	3.26	4.66	2.65	3	4	3	3	4	2
15	0.68	1.94	0.84	0.62	0.47	1.10	1	2	1	1	1	1

Table 9. Effects of Human Mesenchymal Stem Cells on Serum TNF- α Concentration

Patient #	BL	6-Mo.	6-Mo. Score
1	3.5	2.4	2
2	3.6	2.2	2
3	2.8	2.3	2

-37-

Patient #	BL	6-Mo.	6-Mo. Score
4	2.8	0.9	2
5	8	6.8	0
6	3.2	1.1	2
7	3.5	0.14	2
8	7.3	0	2
9	6.8	4.1	2
10	5.7	2.8	2
11	6.3	2.7	2
12	4.1	2.2	2
13	3.5	0.6	2
14	9.2	3.8	2
15	6.3	1.2	2

Table 10. Total Immunosenescence Scores of Patients

Patient #	BL	6-Mo.	2 nd Inject. BL	2 nd Inject. 1-Mo. Score	2 nd Inject. 3-Mo. Score	2 nd Inject. 60-Mo. Score
1	11	14	14	15	16	16
2	15	19	17	17	20	17
3	14	17	12	17	17	16
4	9	18	20	21	21	21
5	14	21	18	23	18	20
6	7	19	19	17	19	22
7	12	17	14	15	15	14
8	9	19	17	19	19	16
9	17	19	15	19	17	18
10	16	24	19	20	24	20
15	15	19	14	17	18	18
Average	12.64	18.73	16.27	18.18	18.55	18

WHAT IS CLAIMED IS:

1. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells (CD19⁺, CD27⁻, IgD⁻) in a sample of the subject's serum decreases by at least 25% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.
2. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻) in a sample of the subject's serum increases by at least 100% as compared to the number of switched memory B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.
3. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum decreases by at least 30% as compared to the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.
4. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of early activated T-cells (CD3⁺, CD69⁺) in a sample of the subject's serum decreases by at least 30% as compared to the number of early activated T-cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.
5. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of chronic activated T-cells (CD3⁺, CD25⁺) in a sample of the subject's serum decreases by at least 70% as

-39-

compared to the number of chronic activated T-Cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.

6. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of Temra cells (CD45RA⁺, CCR7⁺) in a sample of the subject's serum decreases by at least 40% as compared to the number of Temra cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.

7. A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the TNF- α concentration in a sample of the subject's serum decreases by at least 80% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.

8. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells (CD19⁺, CD27⁺, IgD⁺) in a sample of the subject's serum decreases by at least 10% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

9. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁺) in a sample of the subject's serum increases by at least 75% as compared to the number of switched memory B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

10. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal

-40-

stem cells to a subject in need thereof, wherein the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum decreases by at least 60% as compared to the number of B-cells expressing intracellular TNF- α in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

11. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of early activated T-cells (CD3⁺, CD69⁺) in a sample of the subject's serum decreases by at least 30% as compared to the number of early activated T-cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

12. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of chronic activated T-cells (CD3⁺, CD25⁺) in a sample of the subject's serum decreases by at least 75% as compared to the number of chronic activated T-Cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

13. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of Temra cells (CD45RA⁺, CCR7⁻) in a sample of the subject's serum decreases by at least 20% as compared to the number of Temra cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

14. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the TNF- α concentration in a sample of the subject's serum decreases by at least 50% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

-41-

15. A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the $CD4^+ : CD8^+$ T cell ratio in a sample of the subject's serum increases by at least 100% as compared to the $CD4^+ : CD8^+$ T cell ratio in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.
16. The method of any one of claims 1-7, wherein the subject is a human.
17. The method of any one of claims 8-15, wherein the subject exhibits inflammaging.
18. The method of any one of claims 1-17, wherein the mesenchymal stem cells are bone marrow-derived mesenchymal stem cells.
19. The method of any one of claims 1-18, wherein the mesenchymal stem cells do not express STRO-1.
20. The method of any one of claims 1-8, wherein the mesenchymal stem cells do not express CD45.
21. The method of any one of claims 1-20, wherein the mesenchymal stem cells do not express fibroblast surface markers or have a fibroblast morphology.
22. The method of any one of claims 1-21, wherein the mesenchymal stem cells are not genetically manipulated.
23. The method of any one of claims 1-22, wherein the isolated population of allogeneic mesenchymal stem cells is administered in a single dose.
24. The method of any one of claims 1-22, wherein the isolated population of allogeneic mesenchymal stem cells is administered in two or more doses.
25. The method of any one of claims 1-22, wherein the isolated population of allogeneic mesenchymal stem cells is administered at least yearly.
26. The method of any one of claims 1-25, wherein the isolated population of allogeneic mesenchymal stem cells is administered systemically.
27. The method of any one of claims 1-26, wherein the isolated population of allogeneic mesenchymal stem cells is administered by infusion or direct injection.
28. The method of any one of claims 1-27, wherein the isolated population of allogeneic mesenchymal stem cells is administered intravenously, intraarterially, or intraperitoneally.
29. The method of claim 28, wherein the isolated population of allogeneic mesenchymal stem cells is administered intravenously.

-42-

30. The method of any one of claims 1-25, wherein the isolated population of allogeneic mesenchymal stem cells is administered intramuscularly, intravenously, intraarterially, intraperitoneally, subcutaneously, intradermally, orally, tranendocardially, or intranasally.
31. The method of claim 30, wherein the isolated population of allogeneic mesenchymal stem cells is administered intramuscularly.
32. The method of any one of claims 1-31, wherein the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 20×10^6 mesenchymal stem cells.
33. The method of any one of claims 1-31, wherein the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 100×10^6 mesenchymal stem cells.
34. The method of any one of claims 1-31, wherein the isolated population of allogeneic mesenchymal stem cells is administered at a dose of about 200×10^6 mesenchymal stem cells.
35. The method of any one of claims 1-34, wherein the isolated population of allogeneic mesenchymal stem cells are obtained from a human donor and wherein a step of MHC matching of the human donor to the subject is not employed prior to the administration of the isolated population of allogeneic mesenchymal stem cells to the subject.
36. A method of evaluating cellular and humoral immunity status in a subject, comprising:
- (1) obtaining a serum sample from a subject selected for evaluation based on a determination that the subject was previously in need of treatment of non-ischemic dilated cardiomyopathy and said subject had been administered an initial dose of an isolated population of allogeneic human mesenchymal stem cells;
 - (2) performing one or more assays configured to detect a non-ischemic dilated cardiomyopathy marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), and serum $TNF-\alpha$ by introducing the serum sample obtained from the subject into an assay instrument which (i) contacts the serum sample with one or more antibodies which specifically bind for detection the biomarker(s) which are assayed, and (ii) generates one or more assay results indicating of binding of each biomarker which is assayed to a respective antibody to provide one or more assay results;
 - (3) correlating the assay result(s) generated by the assay instrument to the immunity status of the subject, wherein said correlating step comprises assigning a likelihood of one or more future changes in immune status to the subject based on the assay result(s); and

-43-

(4) treating the subject based on the predetermined subpopulation of individuals to which the subject is assigned, wherein the treatment comprises administration of one or more additional doses of an isolated population of allogeneic human mesenchymal stem cells.

37. A method of evaluating cellular and humoral immunity status in a subject, comprising:

(1) obtaining a serum sample from a subject selected for evaluation based on a determination that the subject was previously in need of treatment of symptoms of aging frailty and said subject had been administered an initial dose of an isolated population of allogeneic human mesenchymal stem cells;

(2) performing one or more assays configured to detect aging frailty marker selected from the group of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular $TNF-\alpha$, early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), the $CD4^+$: $CD8^+$ T cell ratio, and serum $TNF-\alpha$ by introducing the serum sample obtained from the subject into an assay instrument which (i) contacts the serum sample with one or more antibodies which specifically bind for detection the biomarker(s) which are assayed, and (ii) generates one or more assay results indicating of binding of each biomarker which is assayed to a respective antibody to provide one or more assay results;

(3) correlating the assay result(s) generated by the assay instrument to the immunity status of the subject, wherein said correlating step comprises assigning a likelihood of one or more future changes in immune status to the subject based on the assay result(s); and

(4) treating the subject based on the predetermined subpopulation of individuals to which the subject is assigned, wherein the treatment comprises administration of one or more additional doses of an isolated population of allogeneic human mesenchymal stem cells.

38. The method of claims 36 or 37, wherein said one or more future changes in immune status comprise one or more of an increase in the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), a decrease in the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), an increase in the number of B-cells expressing intracellular $TNF-\alpha$, an increase in the number of early activated T-cells ($CD3^+$, $CD69^+$), an increase in the number of chronic activated T-cells ($CD3^+$, $CD25^+$), an increase in the number of Temra cells ($CD45RA^+$, $CCR7^-$), a decrease in the $CD4^+$: $CD8^+$ T cell ratio, and an increase in serum $TNF-\alpha$.

-44-

39. An *in vitro* method of determining efficacy of treatment of non-ischemic dilated cardiomyopathy in a subject comprising: determining the levels of one or more biomarkers selected from the group consisting of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), B-cells expressing intracellular TNF- α , early activated T-cells (CD3⁺, CD69⁺), chronic activated T-cells (CD3⁺, CD25⁺), Temra cells (CD45RA⁺, CCR7⁻), and the TNF- α concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if

(1) the number of exhausted B cells (CD19⁺, CD27⁻, IgD⁻) decreases by at least 25% as compared to the number of exhausted B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(2) the number of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻) increases by at least 100% as compared to the number of switched memory B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(3) the number of B-cells expressing intracellular TNF- α decreases by at least 30% as compared to the number of B-cells expressing intracellular TNF- α prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(4) the number of early activated T-cells (CD3⁺, CD69⁺) decreases by at least 30% as compared to the number of early activated T-cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(5) the number of chronic activated T-cells (CD3⁺, CD25⁺) decreases by at least 70% as compared to the number of chronic activated T-Cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(6) the number of Temra cells (CD45RA⁺, CCR7⁻) decreases by at least 40% as compared to the number of Temra cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells, and/or

(7) the TNF- α concentration in a sample of the subject's serum decreases by at least 80% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells.

40. An *in vitro* method of determining efficacy of treatment of symptoms of aging frailty in a subject comprising: determining the levels of one or more biomarkers selected from the

-45-

group consisting of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-), switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-), B-cells expressing intracellular TNF- α , early activated T-cells ($CD3^+$, $CD69^+$), chronic activated T-cells ($CD3^+$, $CD25^+$), Temra cells ($CD45RA^+$, $CCR7^-$), the $CD4^+$: $CD8^+$ T cell ratio, and the TNF- α concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if

(1) the number of exhausted B cells ($CD19^+$, $CD27^-$, IgD^-) decreases by at least 10% as compared to the number of exhausted B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(2) the number of switched memory B cells ($CD19^+$, $CD27^{high}$, IgD^-) increases by at least 75% as compared to the number of switched memory B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(3) the number of B-cells expressing intracellular TNF- α decreases by at least 60% as compared to the number of B-cells expressing intracellular TNF- α prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(4) the number of early activated T-cells ($CD3^+$, $CD69^+$) decreases by at least 30% as compared to the number of early activated T-cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(5) the number of chronic activated T-cells ($CD3^+$, $CD25^+$) decreases by at least 75% as compared to the number of chronic activated T-Cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(6) the number of Temra cells ($CD45RA^+$, $CCR7^-$) decreases by at least 20% as compared to the number of Temra cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,

(7) the TNF- α concentration in a sample of the subject's serum decreases by at least 50% as compared to the TNF- α concentration in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, and/or

(8) the $CD4^+$: $CD8^+$ T cell ratio in a sample of the subject's serum increases by at least 100% as compared to the $CD4^+$: $CD8^+$ T cell ratio in a sample of the subject's serum

prior to administration of said population of isolated allogeneic human mesenchymal stem cells.

41. Use *in vitro* of the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7), and the TNF- α concentration in serum for determining whether treatment for non-ischemic dilated cardiomyopathy is efficacious.

42. Use *in vitro* of the levels of exhausted B cells (CD19⁺, CD27⁻, IgD⁻), the levels of switched memory B cells (CD19⁺, CD27^{high}, IgD⁻), the levels of B-cells expressing intracellular TNF- α , the levels of early activated T-cells (CD3⁺, CD69⁺), the levels of chronic activated T-cells (CD3⁺, CD25⁺), the levels of Temra cells (CD45RA⁺, CCR7), the CD4⁺:CD8⁺ T cell ratio, and the TNF- α concentration in serum for determining whether treatment for aging frailty is efficacious.




FIG. 1

2/24

Characteristic	Treatment Group			Total n=30
	Allo-100M n=10	Allo-200M n=10	Placebo n=10	
Sex				
Male	6 (60%)	6 (60%)	6 (60%)	18 (60%)
Female	4 (40%)	4 (40%)	4 (40%)	12 (40%)
Race				
American Indian/Alaskan Native	0 (0.0%)	1 (10.0%)	0 (0.0%)	1 (3.3%)
Asian	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Hawaiian/Pacific Islander	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Black or African American	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
White	10 (100.0%)	9 (90.0%)	10 (100.0%)	29 (96.7%)
More than One Race	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Other, Specify	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Unknown	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Not Answered	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Age at infusion in years				
Mean	75 ± 7.4	75.3 ± 8.4	75.3 ± 6.8	75.5 ± 7.3
Infusion Status				
Canadian Clinical Frailty Scale	10 (100.0%)	10 (100.0%)	10 (100.0%)	30 (100.0%)
4	5 (50.0%)	7 (70.0%)	5 (50.0%)	17 (56.7%)
5	3 (30.0%)	1 (10.0%)	5 (50.0%)	9 (30.0%)
6	2 (20.0%)	2 (20.0%)	0 (0.0%)	4 (13.3%)
7	0 (0.0%)	0 (0.0%)	0 (0.0%)	0 (0.0%)
Mini Mental State Examination				
Mean	29.3 ± 0.8	28.5 ± 1.1	29.5 ± 1.0	29.1 ± 1.0
Hemoglobin Level (g/dL)				
Mean	14.1 ± 1.2	13.5 ± 1.3	14.3 ± 1.2	14.0 ± 1.3
White Blood Cell Count (cells/mm³)				
Mean	7160.0 ± 2437.8	6600.0 ± 1303.8	7070.0 ± 2214.6	6943.3 ± 1989.1
AST (U/L)				
Mean	24.5 ± 7.6	20.7 ± 3.6	29.3 ± 11.1	24.8 ± 8.5
Six Minute Walk Test Distance (meters)				
Mean	345.9 ± 103.4	390.6 ± 148.9	385.8 ± 83.1	374.1 ± 112.9
FEV1 (Liters)				
Mean	2.5 ± 0.7	2.3 ± 0.9	2.3 ± 0.5	2.4 ± 0.7

FIG. 2

3/24




FIG. 3

4/24




FIG. 4

5/24




FIG. 5

6/24




FIG. 6

7/24




FIG. 7

8/24





FIG. 8

9/24

Correlation between Baseline Serum TNF-α and Baseline B cell Expressing IC TNF-α




$R^2 = 0.16, p=0.02$

FIG. 9

10/24

Baseline B cells Expressing IC TNF-α




††† p≤0.0001 compared to young control;
†† p≤0.0001 compared to low TNF-α.

FIG. 10

11/24

Baseline Switch Memory B Cells



††† p<0.0001 compared to young control.

FIG. 11

12/24

Baseline Exhausted B cells




FIG. 12 † p≤0.05 compared to young control.




FIG. 13




FIG. 14





FIG. 15


Correlation between Baseline Serum TNF- α and Change in Serum TNF- α



$R^2 = 0.65, p < 0.0001$

FIG. 16


Change in B cells Expressing IC TNF- α



***p<0.0001 compared to baseline;
† p<0.05 compared to placebo.

FIG. 17


Correlation between Serum TNF- α and Change in B Cells Expressing IC TNF- α



$R^2 = 0.12, p=0.053$


FIG. 18

Change in Switch Memory B Cells




* $p \leq 0.05$ compared to baseline;
*** $p \leq 0.0001$ compared to baseline;

FIG. 19



*p $\leq$ 0.05 compared to baseline.


FIG. 20



***p≤0.0001 compared to baseline;
†p≤0.005 compared to placebo;
‡p≤0.05 compared to low TNF-α.

FIG. 21


Correlation between Change in Serum TNF-α and Change in Temra T Cells



Absolute Change of Serum TNF-α pg/ml
 $R^2 = 0.13$, $p=0.04$

FIG. 22


Change in Immune Risk Phenotype (CD4:CD8 ratio)



*p≤0.05 compared to baseline.

FIG. 23

Correlation between Change in CD4:CD8 ratio and Change in Temra T cells



$R^2 = 0.09$, $p=0.07$

FIG. 24

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2017/061031

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-7, 16(completely); 18-35(partially)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2017/061031

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K35/28 A61P9/00 A61P39/00 G01N33/15
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K A61P G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, EMBASE, CHEM ABS Data, WPI Data, BIOSIS, Sequence Search

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2016/067285 A1 (DE LA ROSA OLGA [ES] ET AL) 10 March 2016 (2016-03-10) the whole document	1-7,16, 18-25, 27-35
X	US 2012/269785 A1 (WOODS ERIK JOHN [US] ET AL) 25 October 2012 (2012-10-25) paragraphs [0002], [0012], [0013], [0020], [0023], [0024], [0052], [0053]; claims 1,4,14	1-7,16, 18-35
X	WO 2014/203267 A2 (KASIAK RES PVT LTD [IN]) 24 December 2014 (2014-12-24) claims 1,8-12 ----- -/-	1-7,16, 18-35



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 February 2018

Date of mailing of the international search report

30/04/2018

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Greif, Gabriela

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/061031

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ICHIM THOMAS E ET AL: "Placental mesenchymal and cord blood stem cell therapy for dilated cardiomyopathy", REPRODUCTIVE BIOMEDICINE ON,, vol. 16, no. 6, 1 June 2008 (2008-06-01), pages 898-905, XP009180675, ISSN: 1472-6491, DOI: 10.1016/S1472-6483(10)60159-9 abstract page 902, right-hand column, paragraph 2 - page 903, left-hand column, paragraph 1 page 989, right-hand column, paragraph 1 - page 990, left-hand column, paragraph 1 -----	1-7,16, 22,24-26
X	PSALTIS P J ET AL: "Reparative Effects of Allogeneic Mesenchymal Precursor Cells Delivered Transendocardially in Experimental Nonischemic Cardiomyopathy", JACC: CARDIOVASCULAR INTERVENTIONS, ELSEVIER, AMSTERDAM, NL, vol. 3, no. 9, 1 September 2010 (2010-09-01), pages 974-983, XP027290326, ISSN: 1936-8798 [retrieved on 2010-09-01] the whole document -----	1-7,25, 26,30,31
X	JOSHUA M. HARE ET AL: "Comparison of Allogeneic vs Autologous Bone Marrow-Derived Mesenchymal Stem Cells Delivered by Transendocardial Injection in Patients With Ischemic Cardiomyopathy", JAMA THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION, vol. 308, no. 22, 12 December 2012 (2012-12-12), page 2369, XP055254332, US ISSN: 0098-7484, DOI: 10.1001/jama.2012.25321 the whole document -----	1-7,16, 18-35
X,P	WO 2017/123667 A2 (CARDIOCELL LLC [US]) 20 July 2017 (2017-07-20) the whole document ----- -/--	1-7

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2017/061031

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BLOMBERG BONNIE B ET AL: "Age effects on mouse and human B cells", IMMUNOLOGIC RESEARCH, HUMANA PRESS, INC, US, vol. 57, no. 1, 8 November 2013 (2013-11-08), pages 354-360, XP035952301, ISSN: 0257-277X, DOI: 10.1007/S12026-013-8440-9 [retrieved on 2013-11-08] the whole document -----	1-7,16, 18-35

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2017/061031

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2016067285	A1	10-03-2016	AU 2014230014 A1 29-10-2015 CA 2906299 A1 18-09-2014 EP 2972368 A2 20-01-2016 HK 1219775 A1 13-04-2017 JP 2016516985 A 09-06-2016 SG 11201507521U A 29-10-2015 US 2016067285 A1 10-03-2016 WO 2014140362 A2 18-09-2014
US 2012269785	A1	25-10-2012	AU 2011295954 A1 11-04-2013 BR 112013004917 A2 20-09-2016 CA 2809272 A1 08-03-2012 CN 103180435 A 26-06-2013 EP 2611906 A1 10-07-2013 JP 2013536860 A 26-09-2013 JP 2016185972 A 27-10-2016 US 2012052049 A1 01-03-2012 US 2012269785 A1 25-10-2012 US 2012269786 A1 25-10-2012 US 2012269787 A1 25-10-2012 US 2017014454 A1 19-01-2017 WO 2012030968 A1 08-03-2012
WO 2014203267	A2	24-12-2014	NONE
WO 2017123667	A2	20-07-2017	NONE

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-7, 16(completely); 18-35(partially)

A method of treating non-ischemic dilated cardiomyopathy in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells (CD19⁺, CD27⁺, IgD⁺) in a sample of the subject's serum decreases by at least 25% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said non-ischemic dilated cardiomyopathy.

2. claims: 8-15, 17(completely); 18-35(partially)

A method of treating symptoms of aging frailty in a subject, comprising administering a therapeutically effective amount of a population of isolated allogeneic human mesenchymal stem cells to a subject in need thereof, wherein the number of exhausted B cells (CD19⁺, CD27⁺, IgD⁺) in a sample of the subject's serum decreases by at least 10% as compared to the number of exhausted B cells in a sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells, thereby treating said symptoms of aging frailty.

3. claims: 36-38

A method of evaluating cellular and humoral immunity status in a subject, comprising:(1) obtaining a serum sample from a subject selected for evaluation based on a determination that the subject was previously in need of treatment of non-ischemic dilated cardiomyopathy or treatment of symptoms of aging frailty and said subject had been administered an initial dose of an isolated population of allogeneic human mesenchymal stem cells:(2) performing one or more assays configured to detect a non-ischemic dilated cardiomyopathy marker selected from the group of exhausted B cells (CD19⁺, CD27⁺, IgD⁺), switched memory B cells (CD19⁺, CD27^{high}, IgD⁺), B-cells expressing intracellular TNF- α , early activated T-cells (CD3⁺, CD69⁺), chronic activated T-cells (CD3⁺, CD25⁺), Temra cells (CD45RA⁺, CCR7⁺), and serum TNF- α by introducing the serum sample obtained from the subject into an assay instrument which (i) contacts the serum sample with one or more antibodies which specifically bind for detection the biomarker(s) which are assayed, and (ii) generates one or more assay results indicating of binding of each biomarker which is assayed to a respective antibody to provide one or more assay results;(3)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

correlating the assay result(s) generated by the assay instrument to the immunity status of the subject, wherein said correlating step comprises assigning a likelihood of one or more future changes in immune status to the subject based on the assay result(s); and(4) treating the subject based on the predetermined subpopulation of individuals to which the subject is assigned, wherein the treatment comprises administration of one or more additional doses of an isolated population of allogeneic human mesenchymal stem cells

4. claims: 39, 41

An in vitro method of determining efficacy of treatment of non-ischemic dilated cardiomyopathy in a subject comprising: determining the levels of one or more biomarkers selected from the group consisting of exhausted B cells (CD19⁺ , CD27⁺, IgD⁺), switched memory B cells (CD19⁺ , CD27^{high} , IgD⁻), B-cells expressing intracellular TNF- α , early activated T-cells (CD3⁺ , CD69⁺), chronic activated T-cells (CD3⁺, CD25⁺), Temra cells (CD45RA⁺ , CCR7⁻). and the TNF- γ concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if(1) the number of exhausted B cells (CD19⁺ , CD27⁺, IgD⁺) decreases by at least 25% as compared to the number of exhausted B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,(2) the number of switched memory B cells (CD19⁺ , CD27^{high} , IgD⁻) increases by at least 100% as compared to the number of switched memory B cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells.(3) the number of B-cells expressing intracellular TNF- γ decreases by at least 30% as compared to the number of B-cells expressing intracellular TNF- α prior to administration of said population of isolated allogeneic human mesenchymal stem cells,(4) the number of early activated T-cells (CD3⁺ , CD69⁺) decreases by at least 30% as compared to the number of early activated T-cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,(5) the number of chronic activated T-cells (CD3⁺, CD25⁺) decreases by at least 70% as compared to the number of chronic activated T-Cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells,(6) the number of Temra cells (CD45RA⁺ , CCR7⁻) decreases by at least 40% as compared to the number of Temra cells prior to administration of said population of isolated allogeneic human mesenchymal stem cells. and/or(7) the TNF- α concentration in a sample of the subject's serum decreases by at least 80% as compared to the TNF- α concentration in a

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

sample of the subject's serum prior to administration of said population of isolated allogeneic human mesenchymal stem cells

5. claims: 40, 42

An in vitro method of determining efficacy of treatment of symptoms of aging frailty in a subject comprising: determining the levels of one or more biomarkers selected from exhausted B cells (CD191 , CD27- , IgD-), switched memory B cells (CD19+ , CD27high , IgD-), B-cells expressing intracellular TNF-a, early activated T-cells (CD3+ , CD69+), chronic activated T-cells (CD3+ , CD25+), Temra cells (CD45RA+ . CCR7-), the CD4+ :CD8+ T cell ratio, and the TNF-a concentration in serum obtained from the subject before and after administration of a population of isolated allogeneic human mesenchymal stem cells to the subject, and comparing the levels of the one or more biomarkers in the serum obtained before and after administration of the population of isolated human mesenchymal stem cells, wherein treatment is efficacious if (1) the number of exhausted B cells (CD19+ , CD27', IgD-) decreases by at least 100% as compared to prior to administration of said human mesenchymal stem cells,(2) the number of switched memory B cells (CD19+ , CD27high , IgD-) increases by at least 75% as compared to prior to administration of said mesenchymal stem cells,(3) the number of B-cells expressing intracellular TNF-a decreases by at least 60% as compared to prior to administration of said mesenchymal stem cells,(4) the number of early activated T-cells (CD3+ , CD69+) decreases by at least 30% as compared to prior to administration of said human mesenchymal stem cells,(5) the number of chronic activated T-cells (CD3+ , CD25+) decreases by at least 75% as compared to prior to administration of the mesenchymal stem cells,(6) the number of Temra cells (CD45RA+, CCR7") decreases by at least 20% as compared to prior to administration of said population of isolated allogeneic human mesenchymal stem cells,(7) the TNF-a concentration in a sample of the subject's serum decreases by at least 50% as compared to prior to administration of said population of isolated allogeneic human mesenchymal stem cells, and/or(8) the CD4+:CD8+ T cell ratio in a sample of the subject's serum increases by at least 100% as compared to prior to administration of said population of isolated allogeneic human mesenchymal stem cells.
