



US 20230192802A1

(19) **United States**(12) **Patent Application Publication**
ARNETT et al.(10) **Pub. No.: US 2023/0192802 A1**(43) **Pub. Date: Jun. 22, 2023**(54) **BTNL3 PROTEINS, NUCLEIC ACIDS AND
ANTIBODIES AND USES THEREOF**(71) Applicant: **Amgen Inc.**, Thousand Oaks, CA (US)(72) Inventors: **Heather ARNETT**, Seattle, WA (US);
Sabine Schadow ESCOBAR,
Sammamish, WA (US); **Ryan Michael**
SWANSON, Seattle, WA (US); **Joanne**
Louise VINEY, Belmont, MA (US)(73) Assignee: **Amgen Inc.**, Thousand Oaks, CA (US)(21) Appl. No.: **17/893,998**(22) Filed: **Aug. 23, 2022****Related U.S. Application Data**(62) Division of application No. 16/256,951, filed on Jan.
24, 2019, now Pat. No. 11,459,370, which is a divi-
sion of application No. 14/407,904, filed on Dec. 12,
2014, now abandoned, filed as application No. PCT/
US2013/051097 on Jul. 18, 2013.(60) Provisional application No. 61/673,639, filed on Jul.
19, 2012.**Publication Classification**(51) **Int. Cl.****C07K 14/705** (2006.01)**A61K 39/00** (2006.01)**A61K 39/395** (2006.01)**C07K 16/46** (2006.01)(52) **U.S. Cl.**CPC **C07K 14/705** (2013.01); **C07K 14/70503**
(2013.01); **A61K 39/0013** (2013.01); **A61K**
39/3955 (2013.01); **C07K 16/46** (2013.01);
C07K 2319/30 (2013.01); **A61K 38/00**
(2013.01)

(57)

ABSTRACT

The invention provides novel BTNL3 proteins, including multimers, fragments, fusion proteins, and variants. In addition, antibodies that can bind to BTNL3 proteins and nucleic acids encoding BTNL3 proteins are provided. Methods of making BTNL3 proteins using such nucleic acids are also provided. Uses for BTNL3 proteins, and agonists or antagonists thereof, are described.

Specification includes a Sequence Listing.





FIGURE 1






Figure 3




Figure 4




FIGURE 5




FIGURE 6

**BTNL3 PROTEINS, NUCLEIC ACIDS AND
ANTIBODIES AND USES THEREOF****CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a divisional of U.S. application Ser. No. 16/256,951, filed on Jan. 24, 2019, which is a divisional of pending U.S. application Ser. No. 14/407,904, filed on Dec. 12, 2014; which is a 371 of International Patent Application No. PCT/US13/51097, filed Jul. 18, 2013; which claims the benefit of U.S. Provisional Application No. 61/673,639 filed Jul. 19, 2012, each of which is incorporated by reference herein.

INCORPORATION BY REFERENCE

[0002] The present application contains a Sequence Listing, which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. The XML copy of the Sequence Listing, which was created on Mar. 10, 2023, is named A-1722-US04-DIV_ST26.XML and is 33,754 bytes in size.

FIELD

[0003] This invention relates to a butyrophilin-like protein and fragments, variants, and derivatives thereof, nucleic acids encoding such proteins, antibodies that bind to these proteins, and agonists and antagonists of these proteins. Pharmaceutical compositions containing such molecules and uses for such molecules or compositions containing them are also contemplated.

BACKGROUND

[0004] Modulation of an immune or inflammatory response may be valuable in various therapeutic settings. Downmodulation of an immune or inflammatory response may be desirable in treatments for various kinds of autoimmune or inflammatory diseases. Enhancement of any immune response may be valuable to, for example, amplify a response to a particular antigen, such as an antigen contained in a vaccine and/or an antigen preferentially expressed on a cancer cell, a cell mediating a fibrotic disease, or a pathogen. Thus, molecules capable of modulating an immune or inflammatory response are potentially of therapeutic value in a variety of pathological settings. The present invention provides therapeutic agents to diagnose and treat diseases characterized by inappropriate, inadequate, and/or abnormal inflammation and/or immune responses. Some of these agents can stimulate an immune response. Others can inhibit inflammation and/or immune responses.

SUMMARY

[0005] The invention provides BTNL3 proteins and variants thereof, nucleic acids encoding them, and antibodies that bind to them. More specifically, the BTNL3 proteins described herein are multimeric proteins and or fusion proteins that can be isolated and/or soluble proteins. BTNL3 proteins can be attached to a surface and/or expressed on the surface of a cell. Also provided are uses for BTNL3 proteins and for antagonists and agonists of BTNL3, including anti-BTNL3 antibodies.

[0006] In one embodiment, the invention encompasses an isolated multimeric BTNL3 protein, optionally a soluble protein, or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface comprising (a) a polypeptide having an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9 and (b) a second polypeptide having an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequences of the polypeptides of (a) and (b) with amino acids 18-236 of SEQ ID NO:2 or with amino acids 18-166 of SEQ ID NO:9 is at least 50, 60, 70, 80, or 100 amino acids long, wherein the BTNL3 protein is at least a dimer, a trimer, or a tetramer, wherein the BTNL3 protein has been produced by a non-human host cell, and wherein the multimeric BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. In a slightly different embodiment, the invention provides isolated multimeric BTNL3 protein, optionally a soluble protein, comprising (a) a polypeptide having an amino acid sequence at least 90% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9, and (b) a second polypeptide having an amino acid sequence at least 90% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequences of the polypeptides of (a) and (b) with amino acids 18-236 of SEQ ID NO:2 or with amino acids 18-166 of SEQ ID NO:9 is at least 50, 60, 70, 80, or 100 amino acids long, wherein the BTNL3 protein has a molecular weight that is at least about two, three, or four times as large as that of a monomeric polypeptide of (a) and/or at least about five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of a monomeric polypeptide of (a), wherein the BTNL3 protein has been produced by a non-human host cell, and wherein the multimeric BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. Alternatively or in addition in either of these embodiments, the polypeptides of (a) and (b) can comprise a sequence having not more than 20, 15, 12, 10, 8, or 5 insertions, deletions, or substitutions of a single amino acid relative to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9. In some embodiments the multimeric BTNL3 protein can be no more than three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of a monomeric polypeptide of (a). The multimeric BTNL3 protein in either of these embodiments can also be at least a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octomer, a nonamer, a decamer, and/or a higher order multimer, which also means that the multimeric BTNL3 protein can be a dimer, a trimer, tetramer, a pentamer, a hexamer, a heptamer, an octomer, a nonamer, a decamer, and/or a higher order multimer. In some embodiments, the multimeric BTNL3 protein can be no more than a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octomer, a nonamer, or a decamer. The multimeric BTNL3 protein can comprise the amino acid sequence from amino acid 18, 19, 20, 21, 22, or 23 to 234, 235, 236, 237, or 238 of SEQ ID NO:2 or the amino acid sequence from amino acid 18, 19, 20, 21, 22, or 23 to 164, 165, 166, 167, or 168

of SEQ ID NO:9. In some embodiments, the amino acid sequences of the polypeptides of (a) and (b) do not comprise amino acids 237 to 259 of SEQ ID NO:2 or amino acids 167 to 189 of SEQ ID NO:9, and in some embodiments these amino acid sequences may comprise an additional amino acid sequence, which is, for example, an amino acid sequence of an Fc polypeptide. Such Fc polypeptides can comprise (i) the amino acid sequence of a native human Fc region or (ii) an amino acid sequence that is substantially similar to that of the native human Fc region having not more than 15, not more than 10, or not more than 5 insertions, deletions, or substitutions of a single amino acid relative to the amino acid sequence of a native human Fc region. The native human Fc may be of the IgG, including IgG1, IgG2, IgG3, or IgG4, IgA, IgD, IgM, or IgE isotype. The multimeric BTNL3 protein can be a homodimer, homotrimer, homotetramer, a homopentamer, a homohexamer, a homoheptamer, a homooctamer, a homononamer, a homodecamer, a higher order homomultimer, a heteromultimer, or a mixture of species. The multimeric BTNL3 protein can have a molecular weight that is: approximately 4 times as large as the molecular weight of the polypeptide of (a) or (b); approximately the sum of two times the molecular weight of the polypeptide of (a) plus two times the molecular weight of the polypeptide of (b); approximately the sum of three times the molecular weight of the polypeptide of (a) plus the molecular weight of the polypeptide of (b); or approximately the sum of the molecular weight of the polypeptide of (a) plus three times the molecular weight of the polypeptide of (b). Nucleic acids encoding such multimeric BTNL3 proteins are also provided, as well as vectors comprising these nucleic acids and host cells containing the vectors and/or the nucleic acids.

[0007] In another embodiment, the invention provides a BTNL3 fusion protein comprising (a) a first polypeptide comprising an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence of the BTNL3 fusion protein with amino acids 18-236 is SEQ ID NO:2 or with amino acids 18 to 166 of SEQ ID NO:9 is at least 80 amino acids long, and (b) a second polypeptide that has a different amino acid sequence from that of the first polypeptide and does not comprise a fragment of the sequence from amino acid 237 to 466 of SEQ ID NO:2 that is at least 20 amino acids long, wherein the BTNL3 fusion protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The fusion protein can be an isolated and/or a soluble protein. The second polypeptide can be an Fc polypeptide, wherein the Fc polypeptide has an amino acid sequence that is identical or substantially similar to an amino acid sequence of a native human Fc region and contains not more than 5, 10, 15, or 20 insertions, deletions, or substitutions of a single amino acid relative to the native human Fc region. The native human Fc region can be of the IgG, including IgG1, IgG2, IgG3, or IgG4, IgA, IgD, IgE, or IgM isotype. The BTNL3 fusion protein can comprise amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9. The BTNL3 fusion protein can comprise an amino acid sequence that is substantially similar to SEQ ID NO:7, wherein the amino acid sequence comprises not more than 5, 10, 15, or 20 insertions, deletions, or substitutions of a single amino acid relative to SEQ ID NO:7, and/or the BTNL3 fusion

protein can comprise SEQ ID NO:7. The BTNL3 fusion protein can be at least a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, or a decamer. In some embodiments the BTNL3 fusion protein can be not more than a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, or a decamer. The BTNL3 fusion protein can comprise a linker. In some embodiments the BTNL3 fusion protein does not comprise a linker. Such a BTNL3 fusion protein can be a multimer, wherein the multimer can have a molecular weight at least about four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric BTNL3 fusion protein. In another aspect, some such multimers can have a molecular weight that is no more than about four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric BTNL3 fusion protein.

[0008] Nucleic acids encoding such BTNL3 fusion proteins are also provided, as well as vectors comprising these nucleic acids and host cells containing the vectors and/or the nucleic acids.

[0009] In another embodiment, the invention provides a BTNL3 protein, optionally a soluble protein, or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface, comprising the amino acid sequence of a fragment of SEQ ID NO:2 extending from about position 25-131 of SEQ ID NO:2, 6, or 9, or a variant thereof comprising no more than 5 or 10 insertions, deletions, or substitutions of a single amino acid relative to amino acids 25-131 of SEQ ID NO:2, 6, or 9, wherein the BTNL3 protein does not also comprise the amino acid sequence of a fragment of SEQ ID NO:2 extending from about position 132 to 236 of SEQ ID NO:2 or a variant thereof comprising no more than 20, 15, 10, 10, or 5 insertions, deletions, or substitutions of a single amino acid relative to amino acids 132 to 236 of SEQ ID NO:2, and wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The BTNL3 protein can be made in a non-human or mammalian host cell. The BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface may comprise no more than 5 insertions, deletions or substitutions of a single amino acid relative to amino acids 25 to 131 of SEQ ID NO:2, 6, or 9. Or, in another aspect, the amino acid sequence of the soluble BTNL3 protein can be at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 25 to 131 of SEQ ID NO:2. The soluble BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be at least a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, or a decamer. Such a BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can also be a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, a decamer, a higher order multimer, or a mixture of these species. In another aspect, the soluble BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface may be no more than a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, or a decamer. Such a soluble BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be a multimer, wherein the

multimer has a molecular weight at least about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric soluble BTNL3 protein. In another aspect, such a soluble BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be a multimer that has a molecular weight that is no more than about three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric soluble BTNL3 protein. Such a soluble BTNL3 protein or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can further comprise another polypeptide, such as, for example, an Fc fragment of an antibody and/or a linker. Nucleic acids encoding such soluble BTNL3 proteins are also provided, as well as vectors comprising these nucleic acids and host cells containing the vectors and/or the nucleic acids.

[0010] Alternatively, a BTNL3 protein, optionally a soluble protein, or a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface, can comprise the amino acid sequence of a fragment of SEQ ID NO:2 or 6 extending from about position 132 to 236 of SEQ ID NO:2 or 6, or a variant thereof, comprising no more than 20, 15, 10, or 5 insertions, deletions, or substitutions of a single amino acid relative to amino acids 132 to 236 of SEQ ID NO:2 or 6, wherein the BTNL3 protein does not also comprise the amino acid sequence of a fragment of SEQ ID NO:2 extending from position 25 to 131 of SEQ ID NO:2 or 6 or a variant thereof comprising no more than 10 insertions, deletions, or substitutions of a single amino acid relative to amino acids 25 to 131 of SEQ ID NO:2 or 6, and wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The BTNL3 protein can be made in a non-human or mammalian host cell. Such a soluble BTNL3 protein or such a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be at least a dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, a decamer, or a higher order multimer. In another aspect, the soluble BTNL3 protein or the BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface may be no more than dimer, a trimer, a tetramer, a pentamer, a hexamer, a heptamer, an octamer, a nonamer, a decamer. Such a BTNL3 protein or such a BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be a multimer, wherein the multimer has a molecular weight at least about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric soluble BTNL3 protein. In addition, the soluble BTNL3 protein or the BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface can be a multimer, wherein the multimer has a molecular weight no more than about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric soluble BTNL3 protein. Nucleic acids encoding such BTNL3 fusion proteins are also

provided, as well as vectors comprising these nucleic acids and host cells containing the vectors and/or the nucleic acids.

[0011] In a further embodiment, there is provided a BTNL3 fusion protein encoded by a DNA, wherein the DNA comprises the following: (a) a DNA, which encodes a polypeptide, wherein the DNA (i) consists of the nucleotide sequence from nucleotide 52, 55, 58, or 61 to 696, 699, 702, 705, or 708 of SEQ ID NO:1 or nucleotide 52, 55, 58, or 61 to 486, 489, 492, 495, or 498 of SEQ ID NO:8; or (ii) hybridizes under stringent conditions to the DNA of (i); and (b) another DNA that does not hybridize to a polynucleotide consisting of the sequence of SEQ ID NO:1 or SEQ ID NO:8 and encodes another polypeptide in frame with the polypeptide encoded by the polynucleotide of (a); wherein the fusion protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The BTNL3 fusion protein can comprise a linker sequence and can be an isolated and/or soluble protein. Such a BTNL3 fusion protein can be a multimer, wherein the multimer has a molecular weight at least about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric BTNL3 fusion protein. In a further aspect, such a BTNL3 fusion protein can be a multimer, wherein the multimer has a molecular weight of no more than about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times as large as that of the monomeric BTNL3 fusion protein. Nucleic acids encoding such BTNL3 fusion proteins are also provided, as well as vectors comprising these nucleic acids and host cells containing the vectors and/or the nucleic acids.

[0012] Any of the BTNL3 proteins discussed above or below can be isolated and/or soluble and can comprise multimers or aggregated species, which comprise multiple molecules of a

[0013] BTNL3 protein. The molecular weight of the monomeric BTNL3 protein species contained in the multimer or aggregate can be measured by gel electrophoresis under reducing conditions or by size exclusion chromatography (SEC) done under reducing conditions. The molecular weight of the multimeric or aggregated species can be measured by gel electrophoresis and/or SEC done under non-reducing conditions. In some embodiments the multimer or aggregate has a molecular weight that is at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 times that of the monomeric species. In some embodiments the multimer or aggregate has a molecular weight that is no more than about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, or 16 times that of the monomeric species. The monomeric BTNL3 protein of such a multimer or aggregate comprises (a) a polypeptide containing the amino acid sequence from amino acid from amino acid 18, 19, 20, 21, 22, or 23 to 234, 235, 236, 237, or 238 of SEQ ID NO:2 or from amino acid 18, 19, 20, 21, 22, or 23 to 164, 165, 166, 167, or 168 of SEQ ID NO:9; (b) a polypeptide having an amino acid sequence at least 90%, 95%, 97% or 99% identical to amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence of the polypeptide of (b) with amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9 is at least 50, 60, 70, 80, 90, or 100 amino acids long; or (c) a polypeptide having a sequence like that of amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9 except that it can contain no

more than 20, 15, 10, or 5 insertions, deletions, or substitutions of a single amino acid relative to amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9.

[0014] In another aspect, described herein is an isolated variant BTNL3 protein comprising a polypeptide comprising an amino acid sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or to amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, wherein the protein does not comprise the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, and wherein the protein can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ IN NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody.

[0015] Also described herein is an isolated DNA encoding any of the BTNL3 proteins described herein, including the multimeric, fusion, and variant BTNL3 proteins, wherein the DNA does not include exon sequences.

[0016] In another aspect, there is provided a DNA encoding a fusion protein comprising a BTNL3 protein and another polypeptide, wherein the DNA comprises: (a) a DNA (i) that consists of the DNA sequence from nucleotide 52, 55, 58, or 61 to 696, 699, 702, 705, or 708 of SEQ ID NO:1 or nucleotide or 52, 55, 58, or 61 to 486, 489, 492, 495, or 498 of SEQ ID NO:8; or (ii) that hybridizes under stringent conditions to the DNA of (i); and (b) another DNA that does not hybridize to a DNA consisting of the sequence of SEQ ID NO:1 or SEQ ID NO:8 and encodes a another polypeptide in frame with the polypeptide encoded by the DNA of (a); wherein the fusion protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. Vectors containing these DNAs and host cells containing the vectors and/or the DNAs are also contemplated.

[0017] The invention provides a method of making any of the BTNL3 proteins discussed above, including the multimeric BTNL3 protein, the BTNL3 fusion proteins, and the soluble or variant BTNL3 protein or the BTNL3 protein attached to a surface such as a microbead or expressed on a cell surface, comprising culturing a host cell containing nucleic acids encoding the BTNL3 protein in a medium under conditions suitable for expression of the DNA and recovering the expressed BTNL3 protein from the cell mass or the culture medium. The DNAs can be introduced into the host cell.

[0018] In still another aspect, a method of treating a patient having an autoimmune or inflammatory disease is provided, which comprises administering to the patient a therapeutically effective dose of any one of the following: (1) any BTNL3 protein comprising the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, (2) a variant thereof which comprises an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, or which comprises an amino acid sequence that has no more than 5, 10, 15, or 20 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the BTNL3 variant protein can inhibit the proliferation of a T cell stimulated by an immobilized

anti-CD3 antibody; or (3) a BTNL3 agonist such as an agonistic anti-BTNL3 antibody or an anti-idiotypic antibody to an anti-BTNL3 antibody. Alternatively, such a treatment could be performed ex vivo. As a further alternative, the BTNL3 protein or agonist could be attached to a small particle for in vivo or ex vivo administration such as administration on a nanoparticle. This method would include the use of any of the multimeric BTNL3 proteins, the BTNL3 fusion proteins, or the BTNL3 proteins discussed above and below for practicing the method with the exception of those BTNL3 variant proteins that can antagonize the inhibition of T cell proliferation by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7. The autoimmune or inflammatory disease can be arthritis including rheumatoid arthritis, juvenile idiopathic arthritis, and psoriatic arthritis, Addison's disease, asthma, polyglandular endocrinopathy syndromes, systemic lupus erythematosus, chronic active hepatitis, thyroiditis, lymphocytic adenohypophysitis, premature ovarian failure, idiopathic phyoparathyroidism, pernicious anemia, glomerulonephritis, autoimmune neutropenia, Goodpasture's syndrome, myasthenia gravis, scleroderma, primary Sjogren's syndrome, polymyositis, autoimmune hemolytic anemia, an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, psoriasis, dermatitis, sarcoidosis, multiple sclerosis, chronic obstructive pulmonary disease, asthma, type 1 and type 2 diabetes, transplantation-related conditions such as graft rejection or graft versus host disease, gout and related inflammatory crystal deposition diseases, or a fibrotic disease, such as atherosclerosis, chronic obstructive pulmonary disease (COPD), cirrhosis, scleroderma, kidney transplant fibrosis, kidney allograft nephropathy, pulmonary fibrosis, including idiopathic pulmonary fibrosis, autoimmune thrombocytopenic purpura, pemphigus vulgaris, acute rheumatic fever, mixed essential cryoglobulinemia, and warm autoimmune hemolytic anemia, among many others.

[0019] In a further aspect, a method for inhibiting T cell proliferation is provided, which comprises adding to the T cell (1) any BTNL3 protein comprising the amino acid sequence from amino acid 18, 19, 20, 21, 22, or 23 to 234, 235, 236, 237, or 238 of SEQ ID NO:2 or from amino acid 18, 19, 20, 21, 22, or 23 to 164, 165, 166, 167, or 168 of SEQ ID NO:9 or (2) a variant thereof which comprises an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9 or which comprises an amino acid sequence that has no more than 5, 10, 15, or 20 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the BTNL3 protein can inhibit the proliferation of T cells stimulated by an immobilized anti-CD3 antibody. This method encompasses the use of soluble multimeric BTNL3 protein, the BTNL3 fusion proteins, or the soluble BTNL3 protein discussed above and below to inhibit T cell proliferation. This inhibition of T cell proliferation can occur in vitro, ex vivo, or in vivo.

[0020] In another aspect, a method for expanding regulatory T ("T reg") cells is provided, which comprises adding to a T cell (1) a BTNL3 protein comprising the amino acid sequence from amino acid 18, 19, 20, 21, 22, or 23 to 234, 235, 236, 237, or 238 of SEQ ID NO:2 or from amino acid 18, 19, 20, 21, 22, or 23 to 164, 165, 166, 167, or 168 of SEQ

ID NO:9 or (2) a variant thereof which comprises an amino acid sequence at least 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9 or which comprises an amino acid sequence that has no more than 5, 10, 15, or 20 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the BTNL3 protein or the variant BTNL3 protein can inhibit the proliferation of T cells stimulated by an immobilized anti-CD3 antibody. This method encompasses the use of a BTNL3 protein attached to a surface such as a plate or a bead or expressed on a cell surface, a soluble BTNL3 protein, which can be multimeric, or a BTNL3 fusion protein or soluble BTNL3 protein discussed above and below to expand T reg cells. This expansion of T reg cells can occur in vitro, ex vivo, or in vivo. The method comprises contacting a population of T cells with an effective amount of a BTNL3 protein, fusion protein, or agonist as described herein. "Expansion" of T reg cells means that the ratio of T reg cells (CD3⁺ FOXP3⁺) to T cells as a whole (CD3⁺ FOXP3) becomes greater. This ratio can be determined by FACS analysis using antibodies to detect cell proteins, a method well known in the art. See, e.g., Swanson et al. (2013), *J. Immunol.* 190: 2027-2035, the relevant portions of which are incorporated herein by reference.

[0021] Another embodiment includes a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective dose of an anti-BTNL3 antibody, wherein the anti-BTNL3 antibody agonizes the inhibition of proliferation of T cells by a native BTNL3 protein, wherein the anti-BTNL3 antibody binds to a protein consisting of the amino acid sequence of amino acids 18-236 of SEQ ID NO:2.

[0022] Another embodiment includes a method for treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective dose of an anti-BTNL3 antibody, wherein the anti-BTNL3 antibody can bind to a protein consisting of the amino acid sequence of amino acids 18 to 236 of SEQ ID NO:2. In some embodiments, the anti-BTNL3 antibody can bind to a cell surface BTNL3 protein and induce an intracellular signaling cascade via the B30.2 domain of BTNL3, for example, in neutrophils expressing BTNL3. Such intracellular signaling can inhibit proliferation of the cells expressing BTNL3 or, in some cases, cause cell death. The autoimmune or inflammatory diseases include, for example, those that are characterized by excess neutrophil activity such as congestive obstructive pulmonary disease (COPD), asthma, inflammatory bowel disease, including ulcerative colitis and Crohn's disease, and gout and related inflammatory crystal deposition diseases.

[0023] A further method includes a treatment for a cancer patient comprising administering to the patient a therapeutically effective amount of an antibody that binds to a BTNL3 protein consisting of amino acids 18 to 236 of SEQ ID NO:2 or to a BTNL3 protein at least 90% or 95% identical to amino acids 18 to 236 of SEQ ID NO:2. In some embodiments, such an anti-BTNL3 antibody can bind to a human BTNL3 protein as described herein with an equilibrium dissociation constant (K_D) of no more than 10^{-8} , 10^{-9} , 10^{-10} , or 10^{-11} M. In some embodiments, such an anti-BTNL3 antibody can bind to a BTNL3 protein comprising

an amino acid sequence at least 90%, 95%, 97%, 98%, or 100% identical to SEQ ID NO:2 and/or SEQ ID NO:9, wherein the alignment window with SEQ ID NO:2 and/or 9 is at least 50, 60, 70, 80, or 100 amino acids long and, optionally, wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. In some embodiments, an anti-BTNL3 antibody can bind to a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 and/or 9. This antibody can be an antagonistic anti-BTNL3 antibody that antagonizes the suppression of T cell proliferation by BTNL3. Alternatively, the antibody may bind to BTNL3 expressed on a cancer cell, thereby signaling the cancer cell so as to inhibit its proliferation or, in some cases cause cell death. The cancer can be, for example, acute or chronic leukemias, lymphoma, non-Hodgkin's lymphoma, Hodgkin's disease, lymphocytic leukemias, lymphocytic or cutaneous lymphomas, carcinomas, adenocarcinomas, sarcomas, thymomas, neoplasms of the mediastinum, breast cancer, prostate cancer, cancers of the head and neck, lung cancer, non-small cell lung cancer, small cell lung cancer, various kinds of skin cancer, cancer of the bladder, malignant gliomas, cancer of the esophagus, cancer of the stomach, cancer of the pancreas, hepatobiliary neoplasms, cancer of the small intestine, colon, or rectum, cancer of the kidney or ureter, testicular cancer, cancer of the urethra or penis, gynecologic tumors, ovarian cancer, sarcomas of the bone, cancers of the endocrine system, cutaneous melanoma, intraocular melanoma, neoplasms of the central nervous system, and plasma cell neoplasms. In some embodiments, the antibody can be an agonistic antibody that binds to BTNL3 expressed on the cancer cells, causing intracellular signaling via the B30.2 domain.

[0024] Further provided is a method of treating a cancer patient comprising administering to the patient a therapeutically effective amount of a variant BTNL3 protein comprising a polypeptide comprising an amino acid sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, wherein the variant BTNL3 protein does not comprise the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, and wherein the variant BTNL3 protein can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The cancer can be an adenocarcinoma.

[0025] In another aspect, provided herein is a method of treating a patient infected with a pathogen comprising administering to the patient a therapeutically effective amount of an antibody that binds to a BTNL3 protein, the amino acid sequence of which consists of amino acids 18-236 of SEQ ID NO:2 and/or amino acids 18-166 of SEQ ID NO:9, or to a BTNL3 protein, the amino acid sequence of which consists of a sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 and/or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long. The antibody can be an antagonistic antibody that can antagonize the inhibition

by a BTNL3 protein comprising the amino acid sequence of SEQ IN NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody.

[0026] In a further aspect, described herein is a method of treating a patient infected with a pathogen comprising administering to the patient a therapeutically effective amount of a variant BTNL3 protein comprising a polypeptide comprising an amino acid sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, wherein the variant BTNL3 protein does not comprise the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, and wherein the variant BTNL3 protein can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ IN NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody.

[0027] In another aspect, described herein is a method for treating a patient having an autoimmune or inflammatory condition comprising the following steps: (a) removing T cells from the patient; (b) stimulating the T cells with a combination of proteins comprising an anti-CD3 antibody and a BTNL3 protein, wherein the BTNL3 protein comprises (i) the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, (ii) an amino acid sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, or (iii) an amino acid sequence that has no more than 20 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9; (c) harvesting the stimulated T cells; and (d) returning the harvested T cells to the patient; wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The BTNL3 protein can be the BTNL3 protein of (b)(iii), wherein the amino acid sequence has no more than 5 or 10 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9. The BTNL3 protein can be the BTNL3 protein of (b)(i). The autoimmune or inflammatory condition can be selected from the group consisting of arthritis including rheumatoid arthritis, juvenile idiopathic arthritis, and psoriatic arthritis, Addison's disease, asthma, polyglandular endocrinopathy syndromes, systemic lupus erythematosus, chronic active hepatitis, thyroiditis, lymphocytic adenohypophysitis, premature ovarian failure, idiopathic hypoparathyroidism, pernicious anemia, glomerulonephritis, autoimmune neutropenia, Goodpasture's syndrome, myasthenia gravis, scleroderma, primary Sjogren's syndrome, polymyositis, autoimmune hemolytic anemia, an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, psoriasis, dermatitis, sarcoidosis, multiple sclerosis, chronic obstructive pulmonary disease, asthma, type 1 and type 2 diabetes, transplantation-related conditions such as graft rejection or graft versus host disease, gout and related inflammatory crystal deposition diseases, or a fibrotic disease, such as atherosclerosis,

chronic obstructive pulmonary disease (COPD), cirrhosis, scleroderma, kidney transplant fibrosis, kidney allograft nephropathy, pulmonary fibrosis, including idiopathic pulmonary fibrosis, autoimmune thrombocytopenic purpura, pemphigus vulgaris, acute rheumatic fever, mixed essential cryoglobulinemia, and warm autoimmune hemolytic anemia, among many others.

[0028] A method is provided for vaccinating a patient, for example against a cancer or a pathogen, which comprises administering to the patient an antigen and (a) an antagonistic antibody that binds to (i) a BTNL3 protein, the amino acid sequence of which consists of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, or (ii) a BTNL3 protein, the amino acid sequence of which consists of a sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long; or (b) a variant BTNL3 protein comprising a polypeptide comprising an amino acid sequence at least 90% or 95% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, wherein the variant BTNL3 protein does not comprise the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, and wherein the variant BTNL3 protein can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ IN NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. The antigen can be a cancer antigen or an antigen that can elicit an immune response against the pathogen. The antagonistic antibody can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ IN NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. In such methods the antagonistic anti-BTNL3 antibody or the variant BTNL3 protein can be administered before, concurrently with, or after the antigen. Contemplated here are vaccines comprising an antagonistic anti-BTNL3 antibody or a variant BTNL3 protein and an antigen.

BRIEF DESCRIPTION OF THE FIGURES

[0029] FIG. 1: The domain structures of the human proteins that are part of the butyrophilin-like (BTNL) protein family (BTNL2, BTNL3, BTNL8, BTNL9, ERMAL, and MOG), as well as the general structures of the related butyrophilin and B7 families, are diagrammed. Each oval or circle represents a protein domain. The domains are indicated as follows: , immunoglobulin variable region-like (IgV-like) domain; , immunoglobulin constant region-like (IgC-like) domain; , transmembrane domain; and , B30.2 domain. Regions not identified with a particular domain structure are indicated by a horizontal line.

[0030] FIG. 2: This figure indicates relative amount of BTNL3 mRNA present in various primary human immune cells derived from the blood of healthy donors. The vertical axis indicates the intensity value for expression of BTNL3 mRNA generated using ROSETTA RESOLVER®. The various cell types tested are indicated along the x axis as follows: 1, peripheral blood mononuclear cells; 2, T cells; 3, CD4⁺ T cells; 4, CD8⁺ T cells; 5, B cells; 6, monocytes; 7,

macrophages; 8, neutrophils; 9, eosinophils; 10, natural killer (NK) cells. The horizontal line plus error bar in lane 8 indicates the average intensity value obtained for BTNL3 expression in neutrophils plus the standard deviation. Methods are described in detail in Example 1.

[0031] FIG. 3: This figure shows the amount of expression of BTNL3 as measured by next generation RNA sequencing of RNA extracted from normal human tissues identified on the X axis as follows: 1, adrenal, not otherwise specified; 2, artery, aorta; 3, blood; 4, bone marrow, not otherwise specified; 5, brain, anterior cingulate; 6, brain, cerebellum; 7, brain, frontal cortex; 8, brain, not otherwise specified; 9, breast; 10, bronchus; 11, ascending colon; 12, colon, not otherwise specified; 13, sigmoid colon; 14, duodenum; 15, esophagus; 16, abdominal fat; 17, heart, not otherwise specified; 18, heart, ventricle; 19, heart, left ventricle; 20, islet cell; 21, kidney, cortex; 22, kidney, medulla; 23, kidney, not otherwise specified; 24, liver; 25, lung; 26, mesenteric lymph node; 27, lymph node, not otherwise specified; 28, skeletal muscle, not otherwise specified; 29, optic nerve; 30, ovary; 31, head of pancreas; 32, pancreas, not otherwise specified; 33, peripheral blood mononuclear cells; 34, prostate; 35, rectum; 36, retina; 37, skin, not otherwise specified; 38, small intestine, duodenum; 39, small intestine, ileum; 40, small intestine, jejunum; 41, spinal cord, cervical; 42, spinal cord, lumbar; 43, spinal cord, thoracic; 44, spleen; 45, stomach, body; 46, stomach, not otherwise specified; 47, stomach, pylorus; 48, testis; 49, thyroid gland; 50, tongue; and 51, urinary bladder, detrusor muscle. The y axis indicates the amount of expression, shown as the number of BTNL3 transcripts per cell.

[0032] FIG. 4: This figure shows the amount of expression of BTNL3 as measured by next generation RNA sequencing analysis of samples of various human disease tissues. The x axis indicates the nature of each sample as follows: 1, metastatic adenocarcinoma; 2, adenocarcinoma, metastatic colon cancer; 3, adenocarcinoma, metastatic gastric cancer; 4, adenocarcinoma, moderately differentiated; 5, adenocarcinoma, not otherwise specified; 6, adenocarcinoma, poorly differentiated; 7, small cell carcinoma, not otherwise specified; 8, small cell carcinoma, oat cell; 9, carcinoma, undifferentiated; 10, type II diabetes sample; 11, acute myelogenous leukemia; 12, acute myeloid leukemia; 13, malignant melanoma; and 14, malignant, metastatic melanoma. The y axis indicates the amount of expression, shown as the number of BTNL3 transcript per cell.

[0033] FIG. 5: The y axis indicates the levels of proliferation of mouse CD4⁺ T cells that have been activated, in some cases, with a protein immobilized on a microtiter dish, sometimes in the presence of various other proteins in solution. The lanes marked along the x axis represent samples treated as follows: lane 1, no immobilized activating protein or other protein used; lane 2, immobilized anti-CD3 antibody used to activate the cells, and human IgG included in the reaction; lane 3, immobilized anti-CD3 antibody used to activate the cells, and recombinant B7-2.Fc included in the reaction; lane 4, immobilized anti-CD3 antibody used to activate the cells, and recombinant BTNL2.Fc included in the reaction; lane 5, immobilized anti-CD3 antibody used to activate the cells, and recombinant BTNL3.Fc included in the reaction; lane 6, immobilized human IgG used to activate the cells, and recombinant BTNL2.Fc

included in the reaction; and lane 7, immobilized human IgG used to activate the cells, and recombinant BTNL3.Fc included in the reaction.

[0034] FIG. 6: In this figure, each black dot reflects data on RNA expression of a particular sequence in murine CD4⁺ T cells, where the expression of the sequence is significantly up- or down-regulated by the stimulus the cells were subjected to. Sequences whose expression was unaffected by the stimulus are not represented in the figure. The x axis indicates the logio of the signal intensity average of both the unstimulated and stimulated for each particular sequence detected in the array analysis. The y axis indicates the logio of the ratio of expression after 24 hours of stimulation compared to expression of same sequence in unstimulated cells cultured for 24 hours. The stimulations used were as follows: left, stimulation with an anti-CD3 antibody alone; middle, stimulation with an anti-CD3 antibody plus a murine BTNL2.Fc fusion protein; and right, stimulation with an anti-CD3 antibody plus a human BTNL3.Fc fusion protein.

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

- [0035]** SEQ ID NO:1: Nucleotide sequence encoding human BTNL3 coding region.
- [0036]** SEQ ID NO:2: Amino acid sequence of human BTNL3.
- [0037]** SEQ ID NO:3: Amino acid sequence of an IgK signal sequence.
- [0038]** SEQ ID NO:4: Amino acid sequence of a signal sequence for human growth hormone.
- [0039]** SEQ ID NO:5: Full length nucleotide sequence encoding human BTNL3.Fc fusion protein.
- [0040]** SEQ ID NO:6: Full length amino acid sequence of human BTNL3 fusion protein.
- [0041]** SEQ ID NO:7: Amino acid sequence of mature BTNL3.Fc (without signal sequence).
- [0042]** SEQ ID NO:8: Nucleotide sequence encoding a splice variant of BTNL3.
- [0043]** SEQ ID NO:9: Amino acid sequence encoded by SEQ ID NO:8.
- [0044]** SEQ ID NO:10: Peptide linker.
- [0045]** SEQ ID NO:11: Peptide linker.
- [0046]** SEQ ID NO:12: Peptide linker.
- [0047]** SEQ ID NO:13: Peptide linker.
- [0048]** SEQ ID NO:14: Peptide linker.
- [0049]** SEQ ID NO:15: Peptide linker.
- [0050]** SEQ ID NO:16: Peptide linker.
- [0051]** SEQ ID NO:17: Peptide linker.
- [0052]** SEQ ID NO:18: Peptide linker.

DETAILED DESCRIPTION

[0053] The invention provides uses for BTNL3 proteins or antagonists or agonists of a BTNL3 protein, such as anti-BTNL3 antibodies and/or variant forms of a BTNL3 protein. The invention provides BTNL3 proteins, including multimers, fusion proteins, and variants thereof, and uses for such proteins, as well as nucleic acids encoding all of the above. BTNL3 proteins can alter T cell function by attenuating T cell activation, proliferation, and cytokine production. Such effects can lead to effective treatments of T cell-mediated autoimmune or inflammatory diseases such as inflammatory bowel diseases and fibrotic disorders, among a number of

others. Antagonists of BTNL3 can function to prevent or antagonize BTNL3 from inhibition of T cell activation, proliferation, and cytokine secretion, thus, leading to an overall increase in T cell activation. Such effects can be useful for treating diseases such as cancer or for enhancing the efficacy of a vaccine. Further, agonists of BTNL3 may be able to alter immune cell function, for example, by clustering the BTNL3 protein without blocking its function. Such agonists may mediate intracellular signaling into cells expressing BTNL3 via the B30.2 domain that may modulate the activity of such cells by, for example, increasing or decreasing proliferation and/or cytokine secretion. Since BTNL3 is expressed on neutrophils and some kinds of cancer cells, such intracellular signaling may modulate the course of diseases mediated by such cells.

Definitions

[0054] The biological activity of a BTNL3 antagonist that can “antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody” can be determined by doing a T cell proliferation assay as described in Example 4 and adding the antagonist to some of the samples in a controlled fashion so as to determine whether the antagonist affects the activity of a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7. By a similar test, the activity of an agonistic antibody that increases the inhibition of T cell proliferation by T cells can be assayed.

[0055] An “antibody,” as meant herein, comprises a heavy chain variable region of an immunoglobulin and/or a light chain variable region of an immunoglobulin. An antibody may be a full length, tetrameric antibody comprising a light chain variable region (V_L), a light chain constant region (C_L), a heavy chain variable region (V_H), a first heavy chain constant region (C_{H1}), a hinge region, a second heavy chain constant region (C_{H2}), and a third heavy chain constant region (C_{H3}), such as an IgG, IgA, IgD, IgM, or IgE antibody. Alternatively, an antibody can be a fragment such as a Fab fragment or, optionally, a recombinant fragment, such as an scFv fragment. Single domain antibodies comprising a single variable region, either a V_H or V_L region, are also antibodies as meant herein. Single domain antibodies are described in US Patent Appin. Publication US 2006/0062784, the portions of which describe single domain antibodies are hereby incorporated by reference. Further, various forms of monovalent (including single chain antibodies such as scFvs, Fabs, scFv-Fcs, domain antibodies, and various formats described, for example, in International Application WO 2009/089004 and U.S. Pat. No. 5,837,821, the descriptive portions of which are incorporated herein by reference) and multivalent molecules (such as F(ab)2’s and those described, for example, in International Application WO 2009/089004 and U.S. Pat. No. 5,837,821, the descriptive portions of which are incorporated herein by reference) are encompassed within the meaning of “antibody.”

[0056] It is said in multiple places herein that a multimeric species of a protein has a molecular weight “at least about” two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times that of a monomeric species of the protein. While the meaning of this is plain, this phrase is specifically meant to include species that are about four, five, six, etc. times larger than a monomer and not only combinations of such species with

larger species or only larger species. Similarly, it is said in multiple places that a multimer is “at least” a dimer, a trimer, a tetramer, etc. This phrase is specifically meant to include species that are dimers, trimers, tetramers, etc. and not only the combination of the stated species with larger species or only larger species. Further, it is said in multiple places herein that a multimeric species has a molecular weight that is “no more than” two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen, or sixteen times that of a monomeric species of the protein. While the meaning of this is plain, this phrase is specifically meant to include species that are about four, five, six, etc. times larger than a monomer and not only combinations of such species with smaller species or only smaller species. Similarly, it is said in multiple places that a multimer is “no more than” a dimer, a trimer, a tetramer, etc. This phrase is specifically meant to include species that are dimers, trimers, tetramers, etc. and not only the combination of the stated species with smaller species or only smaller species.

[0057] “BTNL3 proteins,” as meant herein, includes full length human, BTNL3 proteins and fragments and/or variants thereof, which includes proteins encoded by naturally-occurring allelic variants of the BTNL3 gene, as well as recombinantly-produced BTNL3 proteins, that is, proteins produced in host cells such as non-human or mammalian host cells, which may contain some sequence changes relative to naturally-occurring BTNL3 proteins. Proteins comprising soluble fragments of a full length human BTNL3 protein that include most or the entire extracellular region and do not include the transmembrane region are specifically contemplated. Such proteins can be attached to a surface such as a bead or a microtiter plate.

[0058] A patient is receiving “concurrent” treatment with two or more molecules, which may be, for example, therapeutics or components of a vaccine, when the patient receives the two or more molecules during the same general timeframe, optionally at the very same time.

[0059] For example, if a patient were dosed with one molecule daily on an ongoing basis and were also dosed with another molecule once a month on an ongoing basis, the patient would be receiving these two molecules concurrently. Similarly, a patient dosed with two different molecules, each administered every two weeks, but not on the same day, would be receiving concurrent treatment with the two molecules. Further, a patient receiving one molecule on an ongoing basis once per week and another molecule once per day for only three days would be receiving concurrent treatment for a short period of time with these two molecules.

[0060] A “scFv” is a single chain antibody comprising a heavy chain variable region (V_H) and a light chain variable region (V_L) and not comprising a constant region of an antibody. In some embodiments scFv’s can also comprise a linker of variable length between the heavy and light chain variable regions. Although an scFv can be fused to other amino acid sequences, the portion of a protein referred to as an scFv preferably does not comprise any substantial amount of amino acid sequence other than a V_H region, a V_L region, and, optionally, a linker joining these sequences.

[0061] An “Fc polypeptide,” as meant herein, is a fragment of an antibody heavy chain comprising a C_{H2} and a C_{H3} domain and all or part of a hinge region or a variant of such a fragment. An Fc polypeptide does not comprise a C_{H1} domain or a V_H domain. See e.g. Kuby, *Immunology*, Second Edition, p.110-11, W.H. Freeman and Co., New York (1994).

An Fc polypeptide can be of the IgA, IgD, IgE, IgG, or IgA isotype, including IgG1, IgG2, IgG3, IgG4 or other subtypes. An Fc polypeptide can be multimeric, in some cases dimeric. Such multimeric Fc polypeptides are referred to herein as “Fc regions.” Some Fc polypeptides, for example an IgM or IgA Fc polypeptides, can form higher order multimers. A single polypeptide chain within a multimeric Fc region is referred to as an “Fc polypeptide.” Variants of Fc polypeptides, as meant herein, may comprise from 1 to 30 (including specifically, no more than 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, etc.) insertions, deletions, or substitutions of a single amino acid relative to a naturally-occurring Fc polypeptide. A naturally occurring or “native” Fc polypeptide has a sequence that occurs in nature in a living organism, for example, a human or a mouse Fc polypeptide. Thus, a “native human” Fc polypeptide has an amino acid sequence that is found in a naturally occurring human Fc polypeptide. Guidance as to where variations can be tolerated without affecting function can be found in the art. For example, alterations of amino acid residues identified in U.S. Pat. No. 5,807,706 and International Application WO 2009/089004, the relevant portions of which are incorporated herein by reference, may be used to encourage heterodimer formation as compared to homodimer formation. Similarly, alterations to the Fc polypeptide that do not prevent binding of the neonatal Fc receptor, FcRn, are encompassed within the alterations that can occur in Fc variants as meant herein. Binding of an Fc polypeptide to FcRn can be ascertained at about pH 6 using a Biacore instrument, such as a Biacore 3000. Human FcRn can be coupled to a CM5 chip using standard chemistry. The Fc-containing protein can be part of the mobile phase, and the response can be measured in resonance units. Alterations of Fc polypeptides are described in, for example, International Patent Application Publication WO 97/34631, the relevant portions of which are incorporated herein by reference. Alternatively, comparisons of, for example, IgG sequences within and between species can locate highly conserved amino acids, which would suggest to one of skill in the art that alteration of those amino acids may affect structure and/or function. Numerous alignments of sequences of hinge, C_H2 and C_H3 regions (which together form the Fc region) are available in, for example, Kabat et al., *Sequences of Immunological Interest*, National Institutes of Health, Publication No. 91-3242, 1991, the relevant portions of which are incorporated herein by reference. On the other hand, amino acids which vary among various IgGs are sites at which variation is likely to be tolerated without effect on function. Similarly, Fc variants that have other desired properties, such as increased or decreased effector functions, including antibody dependent cellular cytotoxicity and/or C1q binding, which leads to complement fixation, are encompassed within what is meant by Fc variants.

[0062] The term “full length antibody” refers to a molecule similar in structure to most naturally-occurring antibodies, for example, an IgG antibody, that is, containing two entire heavy chains and two entire light chains. See e.g. Kabat et al., *supra* or Kuby, *Immunology*, Second Edition, p.109-32, W.H. Freeman and Co., New York (1994) for discussion of the structure of naturally-occurring antibodies. The portions of these references describing the structure of full length antibodies are incorporated herein by reference. Also included among “full length antibodies” are antibodies similar in structure to the naturally-occurring dromedary antibodies that contain only two complete heavy chains

(often with an unusually long CDR3 region) and no light chains Muldermans et al. (2001), *J. Biotechnol.* 74:277-302; Desmyter et al. (2001), *J. Biol. Chem.* 276:26285-26290. The portions of these references describing the structure of these dromedary antibodies are incorporated herein by reference.

[0063] A “multimeric” protein, such as a multimeric BTNL3 protein, is a protein comprising more than one polypeptide chain. The term “multimer” encompasses terms such as “dimer,” “trimer,” or “tetramer,” which specify how many polypeptide chains the multimer contains. A “homomultimer” consists of two or more copies of the same polypeptide chain and does not contain any different polypeptide chains. Similarly, a “homodimer” consists of two copies of the same polypeptide chain, a “homotrimer” consists of three copies of the same polypeptide chain, etc. A “heteromultimer” contains at least two different polypeptide chains. If the heteromultimer has three or more polypeptide chains, some of them can be identical to each other as long as at least one is different from the others. When a protein is said to be “at least a trimer,” it is meant that it is a trimer or a higher order multimer. Similar meanings would be ascribed to “at least a tetramer,” “at least a pentamer,” etc. Similarly, a multimeric protein that is “no more than” a trimer is a trimer or a dimer.

[0064] A “Fab fragment” is an antibody fragment comprising a light chain comprising a V_L and C_L region and a portion of a heavy chain comprising a V_H and a C_H1 region. A Fab fragment does not comprise a C_H2 or C_H3 region. See e.g., Kuby, *Immunology*, Second Edition, pp.110-11 W.H. Freeman and Co., New York (1994) for a discussion of what Fab fragments are. Different kinds of Fab fragments may contain either no hinge region, a portion of a hinge region, or an entire hinge region.

[0065] An “scFv-Fc,” as used herein, is a recombinant protein that is a fusion of an scFv with an Fc polypeptide. See Li et al. (2000), *Cancer Immunol. Immunother.* 49:243-252; Powers et al. (2001), *J. Immunol. Methods* 251:123-135; Gilliland et al. (1996), *Tissue Antigens* 47:1-20. An scFv-Fc can be dimeric.

[0066] As meant herein, “non-reducing conditions” are conditions under which disulfide bridges within a protein will not be reduced such that the bridges will be destroyed.

[0067] A “recombinant” protein or antibody is one resulting from the process of genetic engineering. The term “genetic engineering” refers to a recombinant DNA or RNA method used to create a cell that expresses a gene at elevated levels or at lowered levels, or expresses a mutant form of the gene. In other words, the cell has been transfected, transformed or transduced with a recombinant polynucleotide molecule, and thereby altered so as to cause the cell to alter expression of a desired polypeptide.

[0068] Soluble secreted proteins and proteins expressed on the cell surface often comprise an N-terminal “signal sequence,” which is a hydrophobic sequence that mediates insertion of the protein through the membrane bounding the endoplasmic reticulum (ER) in a eukaryotic cell. Type I transmembrane proteins also comprise signal sequences. “Signal sequences,” as meant herein are amino-terminal hydrophobic sequences which are usually enzymatically removed following the insertion of part or all of the protein through the ER membrane into the lumen of the ER. Thus, it is known in the art that a signal sequence can be present as part of a precursor form of a secreted or transmembrane

protein, but will generally be absent from the mature form of the protein. When a protein is said to comprise a signal sequence, it is to be understood that, although a precursor form of the protein does contain the signal sequence, a mature form of the protein will likely not contain the signal sequence. Signal sequences contain a residue adjacent to and immediately upstream from the cleavage site (position —1) and another residue at position —3, which are important for this enzymatic cleavage. Nielsen et al. (1997), *Protein Eng.* 10(1):1-6; von Heijne (1983), *Eur. J. Biochem.* 133:17-21; von Heijne (1985), *J. Mol. Biol.* 184:99-105, the portions of which describe signal sequences and how to identify them are incorporated herein by reference. Signal sequences can be identified as described by Nielsen et al. (supra). Examples of signal peptides or sequences that are functional in mammalian host cells include the following: the signal sequence for interleukin-7 (IL-7) described in U.S. Pat. No. 4,965,195; the signal sequence for interleukin-2 receptor described in Cosman et al. ((1984), *Nature* 312:768); the interleukin-4 receptor signal peptide described in EP Patent 0 367 566; the type I interleukin-1 receptor signal sequence described in U.S. Pat. No. 4,968,607; the type II interleukin-1 receptor signal peptide described in EP Patent 0 460 846; the signal sequence of human IgK (which is METDTLLLVLLLWVPGSTG; SEQ ID NO:3); and the signal sequence of human growth hormone (MATGSRTSLLAFLGLCLPWLQEGSA; SEQ ID NO:4). The relevant portions of these references are incorporated herein by reference. Many other signal sequences are known in the art.

[0069] An “immunoglobulin-like” (Ig-like) domain, as meant herein, is distinguished mainly by its tertiary structure. See e.g. Bork et al. (1994), *J. Mol. Biol.* 242: 309-20; Hunkapiller and Hood (1989), *Adv. Immunol.* 44: 1-63; Williams and Barclay (1988), *Ann. Rev. Immunol.* 6: 381-405. However, variable and constant immunoglobulin-like domains do contain a handful of highly conserved amino acids that occur at conserved positions within their primary amino acid sequence. See e.g. Kabat et al. (1991), *Sequences of Proteins of Immunological Interest*, U.S. Dept. of Health and Human Services, Public Health Service, National Institutes of Health, NIH Publication No. 91-3242. Such conserved amino acids in variable regions and in C_H1 and C_H2 constant regions are discussed in detail in, e.g., Harpaz and Chothia (1994), *J. Mol. Biol.* 238: 528-39 and Williams and Barclay (1988), *Ann. Rev. Immunol.* 6: 381-405. The portions of these references that discuss such conserved residues are incorporated herein by reference. The presence of such highly conserved amino acids or conservative variants thereof occurring in the proper spacing can indicate the presence of an IgC-like or IgV-like domain.

[0070] The percent identity of two amino acid or two nucleic acid sequences can be determined by comparing sequence information using the computer program GAP, i.e., Genetics Computer Group (GCG; Madison, Wis.) Wisconsin package version 10.0 program, GAP (Devereux et al. (1984), *Nucleic Acids Res.* 12: 387-95). The preferred default parameters for the GAP program includes: (1) The GCG implementation of a unary comparison matrix (containing a value of 1 for identities and 0 for non-identities) for nucleotides, and the weighted amino acid comparison matrix of Gribskov and Burgess, ((1986) *Nucleic Acids Res.* 14: 6745) as described in *Atlas of Polypeptide Sequence and Structure*, Schwartz and Dayhoff, eds., National Biomedical Research Foundation, pp. 353-358 (1979) or other com-

parable comparison matrices; (2) a penalty of 8 for each gap and an additional penalty of 2 for each symbol in each gap for amino acid sequences, or a penalty of 50 for each gap and an additional penalty of 3 for each symbol in each gap for nucleotide sequences; (3) no penalty for end gaps; and (4) no maximum penalty for long gaps.

[0071] In connection with comparisons to determine sequence identity of polynucleotides or polypeptides, what is meant by an “alignment window” is the portion of the polynucleotide or polypeptide that is matched, partially or wholly, with another polynucleotide or polypeptide by the computer program GAP (Devereux et al. (1984), *Nucleic Acids Res.* 12: 387-95) using the parameters stated herein. For example, when a polypeptide of 20 amino acids is aligned with a considerably longer protein and the first 10 amino acids match the longer protein exactly while the last 10 amino acids do not match the longer protein at all, the alignment window is 10 amino acids. If, on the other hand, the first and last amino acids of the 20 amino acid polypeptide match the longer protein, and eight other matches are scattered between, the alignment window is 20 amino acids long. However, long stretches in either aligned strand without identical or conservatively substituted amino acids or identical nucleotides of at least, for example, 25 amino acids or 75 nucleotides constitute an endpoint of an alignment window, as meant herein. Alignment windows for a comparison of sequences can be at least about 25, 50, 60, 75, 80, 90, 100, 150, 200, 225, 300, 400, 450, 500, or 600 amino acids or nucleotides in length.

[0072] Two polypeptide or nucleotide sequences are considered “substantially similar” when they are at least 90% identical as determined using the GAP program as described above and have similar biological activity. In the case of a BTNL3 protein or variant thereof as described herein, the biological activity to be tested in determining whether two sequences are substantially similar can be the ability to inhibit the proliferation of T cells activated by an immobilized anti-CD3 antibody or to antagonize such inhibition for certain antagonistic BTNL3 variant proteins.

The BTNL Family

[0073] BTNL3 has been placed within the butyrophilin-like (BTNL) family of proteins based on its domain structure. See, e.g., Arnett et al. (2008), *Current Immunology Reviews* 4: 43-52 and Arnett et al. (2009), *Cytokine* 46: 370-75. The human proteins in the BTNL family include BTNL2, BTNL3, BTNL8, BTNL9, ERMAP, and MOG, and the domain structures of these proteins are shown diagrammatically in FIG. 1. As is apparent from FIG. 1, BTNL2 is the only member of the family having four immunoglobulin-like (Ig-like) domains in its extracellular region (two IgV-like and two IgC-like domains). MOG and ERMAP each have only one Ig-like domain. BTNL3, BTNL8, and BTNL9 also have one extracellular domain that is clearly an IgV-like or Ig-like domain and another domain with many characteristics similar to an Ig-like domain, although these domains are not actually classified as Ig-like. All BTNL family members have a transmembrane domain. BTNL2 and MOG have short intracellular regions, whereas BTNL3, BTNL8, BTNL9, and ERMAP have longer intracellular

	Percent identity between human members of BTNL family of proteins				
	BTNL2	BTNL8	BTNL9	ERMAP	MOG
BTNL3	35%	74%	43%	41%	32%

As shown in FIG. 1, BTNL3, BTNL8, and BTNL9 have similar domain structures. The amino acid sequence of BTNL3 is much more similar to that of BTNL8 than to those of other BTNL proteins.

[0075] Beyond levels of sequence identity, certain sites within the BTNL protein family are highly conserved as shown in Table 2 below, which is an alignment of all six human BTNL-like proteins. Beneath the alignment is a consensus sequence (labeled “ALL”). If the consensus amino acid(s) occurs in all proteins in which the amino acid sequence spans the portion of the alignment in which the amino acid occurs, it is shown in bold. If it occurs in all but one of the proteins in which the sequence spans that portion of the alignment, it appears in regular font. If a site has in all cases one of two or more amino acids, each of which are conservative variations of the other, these amino acids are listed below that position in bold font. If a site has in all but one sequence spanning that portion of the alignment one of two or more amino acids, each of which are conservative variations of the other, these amino acids are listed below that position in regular font. The numbering above the alignments in Table 2 is the numbering of SEQ ID NO:2, which is the full length amino acid sequence of human BTNL3, including the signal sequence, which ends at position 17 of SEQ ID NO:2.

Alignment and consensus sequence of BTNL proteins

[illegible]

TABLE 2-continued

Alignment and consensus sequence of BTNL proteins					
	119				168
BTNL3	QIYDEEATWE	LRVAALGSLP	LISIVGYVDG	GIQLLCLSSG	WFPQPTAKWK
BTNL8	QSYYQKATWE	LQVSALGSVP	LISITGYVDR	DIQLLCQSSG	WFPRTAKWK
BTNL9	DNFSGEALWE	LEVAGLGSDP	HLSLEGFKEG	GIQLRLRSSG	WYPKPKVQWR
ERMAP	GNLSKEDTVI	LQVAA...P	SV.....		GSLSPSA...
MOG	HSYQEEAAME	LKVED...P	FY.....		W.VSPGV...
BTNL2	DDVYQEASLD	LKVVG LGSSP	LITVEGQEDG	EMQPMCSDG	WFPQPHVPWR
ALL	EA E L V P				W P A V
	D				
	169				216
BTNL3	GPQGQDLSSD	SRANA.DGYS	LYDVEISIIV	QENA.GSILC	SIHLAEQSHE
BTNL8	GPQGQDLSTD	SRTNR.DMHG	LFDVEISLTV	QENA.GSISC	SMRHAHLSRE
BTNL9	DHQQGCLPPE	FEAIVWDAQD	LFSLETSVVV	RAGALSNVSV	SIQNLLLSQK
ERMAP	VA	LAVILPVLVL	LIMVCLCLIW	KQRRAKEKLL	
MOG	LV	LLAVLPVLLL	QITVGLVFLC	LQYRLRGKLR	
BTNL2	DMEGKTIPSS	SQALTQGSHG	LHFVQTLRLV	TNISAVDVTC	SISIPFLGEE
ALL			LF V L V A I		
			L L V L V V		
			Y I I L		
			A F		
	217				254
BTNL3	VESKVLIGET	FFQ.PSPWR..	LASILLGL	LCGALCGVVM	GM
BTNL8	VESRVQIGDT	FFE.PISWH..	LATKVLGI	LCCGLFFGIV	GL
BTNL9	KELVVQIADV	FVPGASAWKS	AFVATLPLLL	VLAALALGVL	RKQRRSREKL
ERMAP	YEHVTEVDNL	L.....			SDHAKE....
MOG	AE.IENLHRT	F.....			DPHFLRVPCW
BTNL2	KIATFSLSES	..RMTFLWKT	LLVWGLLLAV	AVGLPRKRS~	~~~~~
ALL	E I				
	V				
	L				
	255				297
BTNL3	IIVFFKSK..	GKIQA	ELDWRRKHGQ	AELRDARKHA	VEVTLDPETA
BTNL8	KIFFSKFQ..	WKIQA	ELDWRRKHGQ	AELRDARKHA	VEVTLDPETA
BTNL9	RKQAEKRQEK	LTALEKLQT	ELDWRRAEGQ	AEWRAAQKYA	VDVTLDPASA
ERMAP	KGKLEKAYKK	LRSELK....	...LKRAAAN	SGWRRARLHF	VAVTLDPDPA
MOG	KITLFLVIVPV	LGPLVALIIC	YNWLHRRLAG	QFLEELLFHL	EALSG~~~~
BTNL2	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ALL	K F		RR GQ	R AR HA V	VTLDPETA
	R L		K AN	LQ L LS DS	
	A		H	F	
	298				344
BTNL3	HPKLCVS.DL	KTVTHRKAPO	.EVPHSEKRF	TRKSVVAS.Q	GFQAGKHYWE
BTNL8	HPKLCVS.DL	KTVTHRKAPO	.EVPHSEKRF	TRKSVVAS.Q	SFQAGKHYWE
BTNL9	HPSLEVSSEDG	KSVSSRGAPP	GPAPGHPQRF	SEQTCALSLE	RFSAGRHYWE
ERMAP	HPKLILSDQ	RCV.RLGDRR	QVPDPNPQRF	DFVVSILGSE	YFTTGCHYWE
MOG	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
BTNL2	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ALL	HPKL VS D	KTV HR APQ	VP KRF	T KS VAS	F AGKHYWE
	L	S R R	A Q	S QT AL	R
				I	
	345				394
BTNL3	VDVGQNVGWY	VGVCRDDVD	GKNNVTLSNP	NGYWVLRLTT	EHLYFTFNPH
BTNL8	VDGGHNKRWR	VGVCRDDVD	RKEYVTLSPD	HGYWVLRLNG	EHLYFTLNPR
BTNL9	VHVGRSRWF	LGACLAAPR	A.GPARLSPA	AGYWVLGLWN	GCEYFVLAPH
ERMAP	VYVGDKTKWI	LGVCSESVSR	.KGKVTASPA	NGHWLLRQSR	GNEYEALTSP
MOG	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
BTNL2	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ALL	V VGQN RW	VGVC D V R	K VTLSP	NGYWVLRL	H LF L PH
	HR K	L A E	A A	H	N F R
	RK				
	395				444
BTNL3	FISLPPSTPP	TRVGVFLDYE	GGTISFFNTN	DQSLIYTLT	CQFEGLLRPY
BTNL8	FISVFPRTPP	TKIGVFLDYE	CGTISFFNIN	DQSLIYT.LT	CRFEGLLRPY
BTNL9	RVALTLRVPP	RRLGVFLDYE	AGELSFENVS	DGSHIFTFHD	.TFSGALCAY
ERMAP	QTSFRLKEPP	RCVGIFLDYE	AGVISFYNTV	NKSHIFTF.T	HNFSGPLRPF
MOG	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
BTNL2	~~~~~	~~~~~	~~~~~	~~~~~	~~~~~
ALL	SL R PP	RVGVFLDYE	G ISFFN	DQS IYT T	QF G LRPY

TABLE 2-continued

Alignment and consensus sequence of BTNL proteins																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																					
	V	K		KI	I		L	Y		K	F	N		F																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
	F				L							R																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
	445													466																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
BTNL3	IQHAMYD	EE		KGTP	IFICPV	SWG	~	~	~	~	~	~	~																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
BTNL8	IEYPSYN	EQ		NGTP	IVICPV	TQESKEASW	Q	R	A	S	A	I	P	E	T	S	N	S	E	S	S	Q	A	T	T																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
BTNL9	FRPRAHDGGE			HPDPL	TICPL	P	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.</

One of skill in the art will appreciate that the consensus sequence among these proteins reflects features that may be important for the structures or functions of these proteins. Given their varying expression patterns, it is likely that these proteins do not have identical functions, and, thus, it is unlikely that all amino acids important for the function of each protein would be conserved within the family. However, many of the conserved amino acids may be important to maintain the proper structure, which is, of course, necessary for function. At many sites one of two or more amino acids that are conservative variations of each other occurs in all or most members of the BTNL family. One of skill in the art would understand that such conservative variations in BTNL3 would likely not adversely affect function. For example, at position 36 of SEQ ID NO:2 (which has the same numbering as the alignment of Table 2 above), various members of the BTNL family have one of three different hydrophobic amino acids, alanine (BTNL3, BTNL8, and ERMAP), isoleucine (BTNL2), or valine (BTNL3 and MOG). One of skill in the art would understand that a change from valine to alanine or isoleucine at this position of the BTNL3 amino acid sequence would be unlikely to affect function. Similar considerations would apply at all of the sites where conservative variations occur within the family. Thus, a BTNL3 protein, as meant herein, includes proteins comprising SEQ ID NO:2, or a fragment thereof, wherein the sequence may be altered by conservative variation at a site where conservative variation occurs among members of the BTNL family and wherein the protein can inhibit the proliferation of T cells as measured by the method described in the examples below. Such sites include, for example, positions 3, 16, 20, 28, 30, 32, 34, 35, 36, 38, 42, 48, 53, 55, 63, 64, 66, 67, 68, 69, 72, 79, 81, 82, 87, 88, 89, 91, 97, 98, 100, 101, 104, 112, 116, 128, 165, 189, 191, 195, 197, 201, 204, 223, 255, 258, 272, 276, 277, 283, 284, 287, 290, 291, 303, 308, 311, 316, 318, 323, 326, 328, 329, 331, 332, 340, 349, 350, 352, 355, 357, 360, 369, 371, 375, 386, 391, 394, 398, 401, 406, 407, 409, 418, 421, 426, 430, 436, and 444 of SEQ ID NO:2. Further, variations may also be tolerated at other sites within BTNL3 without effect on

function. For example conservative substitutions at non-conserved positions would be unlikely to affect function.

[0076] Thus, a BTNL3 protein, as meant herein, includes proteins that (1) have naturally-occurring polymorphisms or recombinantly-introduced amino acid changes, (2) are at least 90%, 95%, 96%, 97%, 98% or 99% identical to SEQ ID NO:2 and/or to amino acids 18-236 of SEQ ID NO:2 and/or to amino acids 18-166 of SEQ ID NO:9, and (3) retain the ability to attenuate T cell proliferation as measured by the methods described herein or act as an inhibitor of native BTNL3. Some such polymorphisms may enhance the ability of a BTNL3 protein to inhibit T cell proliferation and/or may make a BTNL3 protein easier to produce in a commercial production process. Other such polymorphisms may produce an inhibitor of native BTNL3. These polymorphisms can occur at sites within BTNL3 that are not conserved such as, for example, position 22, 25, 26, 27, 29, 37, 39, 41, 43, 45, 46, 47, and any other site shown to be nonconserved in Table 2.

[0077] The expression patterns and biological functions of the BTNL proteins have been explored to some extent in some cases, but not in others. ERMAP is expressed on the surface of red blood cells and has not been assigned a specific biological function. MOG is a component of the myelin sheath. Neither ERMAP nor MOG is thought to play a role in the immune system, although antibodies to MOG are often detected in patients with multiple sclerosis. BTNL1 is homologous to MOG, and BTNL1 is found in cow's milk. It has been hypothesized that human consumption of cow's milk may lead to the development of antibodies to BTNL1 that cross-react with human MOG, thus leading to autoimmune diseases such as multiple sclerosis. See Guggenmos et al. (2004), J. Immunol. 172: 61-68. BTNL2 has been shown to inhibit T cell proliferation and cytokine secretion, but not B cell proliferation. Thus, BTNL2 is thought to act as a negative co-regulator of T-cell mediated events. See, e.g., U.S. Pat. No. 7,244,822, the relevant portions of which are incorporated herein by reference. A polymorphism in BTNL2 has been clearly linked to sarcoidosis, suggesting that BTNL2 may play a role in either initiating or mediating or contributing or responding to this disease. Valentonyte et

al. (2005), *Nature Genetics* 37(4): 357-64. More tentative associations have been drawn between various BTNL2 polymorphisms and ulcerative colitis, rheumatoid arthritis, spontaneous inclusion body myositis, systemic lupus erythematosus, type I diabetes, tuberculoid leprosy, and antigen-specific IgE responsiveness. Arnett et al. (2009), *Cytokine* 46: 370-75.

[0078] Levels of RNAs encoding the various BTNL proteins in various cell types and tissues have been reported. BTNL3 is expressed in ileum, small intestine, colon, liver, lung, bone marrow, and spleen. Arnett et al. (2009), *Cytokine* 46: 370-375. Among various cell types associated with immune function that have been tested for BTNL3 expression, BTNL3 RNA is expressed predominantly in neutrophils. Arnett et al. (2009), *Cytokine* 46: 370-75. The expression of BTNL3 RNA in cells involved in immune function suggests that BTNL3 may play a role in immune function, either by driving the inflammatory response or in dampening the response following a flare.

BTNL3 Protein

[0079] The instant invention encompasses secreted, soluble versions of BTNL3, which can, optionally, be immobilized by attachment to a surface such as a nanoparticle, a bead, or a microtiter plate, as well as versions comprising a transmembrane domain that can be expressed on a cell surface. Such BTNL3 proteins can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. Some variant BTNL3 proteins also have this activity, although variants that antagonize this inhibition are also contemplated. Such proteins can be isolated, that is, be part of a purified protein preparation in which the BTNL3 protein constitutes at least 80% or at least 90% of the protein present in the preparation. The invention further includes BTNL3 proteins encoded by the BTNL3 nucleic acids described below. A BTNL3 protein, as meant herein, encompasses a protein comprising the amino acid sequence of SEQ ID NO:2, 6, 7, or 9, as well as fragments, derivatives, and variants thereof, including fusion proteins and multimers, as discussed above and below. The amino acid sequences of SEQ ID NO:2, 6, and 9 include a signal sequence starting at position 1 and ending at a position from about position 15 to about position 20, optionally at position 17. Thus, the amino acid sequence of the mature BTNL3 begins at a position from about 16 to about position 21 of SEQ ID NO:2, 6, or 9. Optionally, the mature amino acid sequence of BTNL3 begins at position 18 of SEQ ID NO:2, 6, or 9.

[0080] The signal sequence of BTNL3 is followed by an Ig-like domain extending from about position 25 to about position 131 of SEQ ID NO:2, 6, or 9. The following region, from about position 132 to about position 236 of SEQ ID NO:2 or 6, aligns with IgC-like domains in BTNL2, but lacks some of the characteristic sequence features commonly found in a IgC1-like domain. See, e.g., Williams and Barclay (1988), *Ann. Rev. Immunol.* 6: 381-405; Peach et al. (1995), *J. Biol. Chem.* 270(36): 21181-21187. This domain is substantially shortened in the amino acid sequence of SEQ ID NO:9, which is the amino acid sequence encoded by a splice variant of BTNL3. The transmembrane domain of BTNL3 begins at about position 237 of SEQ ID NO:2 or about position 167 of SEQ ID NO: 9 and ends at about position 259 of SEQ ID NO:2 or about position 189 of SEQ ID NO:9. The intracellular portion of BTNL3 begins at about position 260 and ends at position 466 of SEQ ID NO:2

or begins at about position 190 and ends at position 432 of SEQ ID NO:9. The intracellular region contains a B30.2 domain extending from about position 288 to about position 445 of SEQ ID NO:2 or about position 254 to position 411 of SEQ ID NO:9. A B30.2 domain is a globular domain of approximately 170 amino acids. Henry et al. (supra) discuss B30.2 domains in some detail and provide an alignment of a number of B30.2 domains and a consensus sequence derived from the alignment. The portions of Henry et al. (1998), *Mol. Biol. Evol.* 15(12): 1696-1705 that show (by sequence comparison) and explain what a B30.2 domain are incorporated herein by reference. B30.2 domains are also found in BTNL9, BTNL8, and ERMAP, all of which are members of the butyrophilin-like family of proteins, as discussed herein. The alignment of BTNL proteins in Table 2 above from about position 288-445 exhibits a high degree of homology, certainly higher than is observed between the more disparate collections of proteins containing B30.2 domains aligned by Henry et al. supra.

[0081] Similarly, the protein having the amino acid sequence of SEQ ID NO:9 includes a signal sequence starting at position 1 and ending at a position from about position 15 to about position 20, optionally at position 17. Thus, the amino acid sequence of the mature protein begins at a position from about 16 to about position 21 of SEQ ID NO:2. Optionally, the mature amino acid sequence begins at position 18 of SEQ ID NO:9. This is followed by an IgV-like domain extending from about 25 to 131 of SEQ ID NO:9. The majority of the following domain present in SEQ ID NO:2 is missing in SEQ ID NO:9. The transmembrane domain begins at about position 167 of SEQ ID NO:9 and extends to about position 189 of SEQ ID NO:9. Beginning at about 10 amino acids beyond the transmembrane domain, SEQ ID NO:9 includes a stretch of 36 amino acids not found in SEQ ID NO:2. The B30.2 domain begins at about position 254 of SEQ ID NO:9 and ends at about position 411 of SEQ ID NO:9.

[0082] BTNL3 proteins, as meant herein, include hetero- and homo-multimers comprising at least two BTNL3 proteins. In some embodiments, biologically active multimers can be homomultimers. The size of such homomultimers can be determined by polyacrylamide gel electrophoresis under non-reducing conditions or by size exclusion chromatography. The size of the monomeric BTNL3 protein contained in such multimers can be determined by polyacrylamide gel electrophoresis of the multimer under reducing conditions. Such conditions would be expected to break disulfide bridges and interfere with non-covalent interactions such as hydrogen bonds or charge interactions. Thus, multimers held together by disulfide bonds or non-covalent interactions would be expected to be reduced to monomers under reducing conditions. In some embodiments, the size of the biologically active BTNL3 homomultimer can be at least about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen or sixteen times the size of the monomeric BTNL3 protein. In some embodiments, the size of such a homomultimer can be no more than about two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, thirteen, fourteen, fifteen or sixteen times the size of the monomeric BTNL3 protein.

[0083] BTNL3 proteins, as meant herein, also include proteins encoded by splice variants of the full length BTNL3 mRNA. The cDNA coding sequence for BTNL3 extends from position 1-1401 of SEQ ID NO:1, the last three

nucleotides being a stop codon. The coding sequence starts within the first exon. The most common splice variant contains eight exons: exon 1, encoding amino acids 1-16 of SEQ ID NO:2; exon 2, encoding amino acids 17-132 of SEQ ID NO:2; exon 3, encoding amino acids 133-224 of SEQ ID NO:2; exon 4, encoding amino acids 225-262 of SEQ ID NO:2; exon 5, encoding amino acids 263-269 of SEQ ID NO:2; exon 6, encoding amino acids 270-278 of SEQ ID NO:2; exon 7, encoding amino acids 279-287 of SEQ ID NO:2; and exon 8, encoding amino acids 288-466. However, alternative splicing can generate different messages including different exons. One of these was cloned by Shibui and coworkers. Shibui et al. (1999), *J. Hum. Genet.* 44: 249-252. The amino acid sequence reported in Shibui et al. from clone ColF4100 is incorporated by reference herein. The cDNA sequence encoding this amino acid sequence from GenBank Accession no. AB020625 is also incorporated herein by reference. The protein encoded by this splice variant lacks a portion of SEQ ID NO:2 encoded by the third and fourth exons. Hence, the encoded protein is missing the majority of the second Ig-like domain of mature BTNL3 protein. Compare SEQ ID NO:2 to SEQ ID NO:9. Moreover, the intracellular region of SEQ ID NO:9 includes sequences not found in SEQ ID NO:2 (residues 200-235 of SEQ ID NO:9). Both the insertion and deletion of extra amino acids in this splice variant are a result from the use of non-canonical splice sites. SEQ ID NO:9, like SEQ ID NO:2, contains a signal sequence ending at a residue from 16 to 21 of SEQ ID NO:9, optionally at residue 17.

[0084] Other splice variants might be formed. For example, a splice variant missing either of exons 2 or 3, that is one of exons encoding either of the two extracellular domains, is contemplated. Similarly, splice variants missing any of exons 5, 6, 7, or 8 are contemplated. BTNL3 proteins, as meant herein, can be encoded by splice variants that are missing various exons. Splice variants can, in addition, use cryptic splice sites.

[0085] In some embodiments, a BTNL3 protein can be a soluble fragment of the full length transmembrane protein comprising SEQ ID NO:2 or SEQ ID NO:9, or a variant thereof. In some embodiments, a BTNL3 protein comprises a fragment of BTNL3 comprising the immunoglobulin-like domain extending from residue 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 of SEQ ID NO:2 or SEQ ID NO:9 to residue 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, or 140 of SEQ ID NO:2 or SEQ ID NO:9. Such embodiments may or may not include all of the following domain extending from residue 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, or 140 of SEQ ID NO:2 to residue 230, 231, 232, 233, 234, 235, 236, 237, or 238 of SEQ ID NO:2. For example, SEQ ID NO:9 lacks the amino acids present at positions 158 to 227 of SEQ ID NO:2. Variants lacking other portions of this domain are also contemplated. In further embodiments, a BTNL3 protein can comprise a fragment extending from residue 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, or 140 of SEQ ID NO:2 to residue 220, 221, 222, 223, 234, 235, 236, 237, 238, 239, 230, 231, 232, 233, 234, 235, 236, 237, or 238 of SEQ ID NO:2. Such embodiments may or may not include the preceding domain extending from residue extending from residue 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 of SEQ ID NO:2

to residue 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, or 140 of SEQ ID NO:2. A BTNL3 protein can comprise a fragment which includes most or all of the extracellular region of BTNL3. Such a protein can comprise an amino acid sequence extending from residue 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 of SEQ ID NO:2 to residue 220, 221, 222, 223, 234, 235, 236, 237, 238, 239, 230, 231, 232, 233, 234, 235, 236, 237, or 238 of SEQ ID NO:2, optionally from about residue 18 to about residue 236 of SEQ ID NO:2. Alternatively, a soluble BTNL3 protein can comprise a fragment that includes most or all of the extracellular region extending from residue 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 of SEQ ID NO:9 to residue 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170 of SEQ ID NO:9. All of these fragments can contain variations relative to SEQ ID NO:2 or SEQ ID NO:9 and can contain a defined number of substitutions, insertions, or deletions of a single amino acid relative to SEQ ID NO:2 or SEQ ID NO:9 as discussed below. These embodiments can inhibit the proliferation of T cells stimulated by an immobilized anti-CD3 antibody. Some BTNL3 protein variants that do not comprise the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 may antagonize the inhibition of the proliferation of T cells stimulated by an immobilized anti-CD3 antibody.

[0086] The invention encompasses epitopes of BTNL3 proteins that are useful for generating antibodies, which are referred to herein as immunogenic fragments. Immunogenic fragments are preferably at least 10 amino acids long and can comprise contiguous amino acids from SEQ ID NO:2 or SEQ ID NO:9. Such epitopes can span regions of a BTNL3 protein encoded by a splice junction, which may have the advantage of specific binding to proteins encoded by specific splice variants. In some embodiments the epitope is located within the extracellular region of BTNL3, which extends from about amino acid position 18 to about position 236 of SEQ ID NO:2 or about residue 18 to about residue 166 of SEQ ID NO:9. An epitope can be partially or completely within the first immunoglobulin variable region-like domain, which extends from about amino acid position 25 to position 131 of SEQ ID NO:2 or SEQ ID NO:9. Alternatively, an epitope can be partially or completely within the following domain, which extends from about amino acid position 132 to 236 of SEQ ID NO:2. In SEQ ID NO:9, this domain, which is substantially shortened, extends from about position 132 to 166 of SEQ ID NO:9. This shortened domain can also contain part or all of an epitope to which an antibody binds.

[0087] A BTNL3 protein, as meant herein, may contain one or more insertions, deletions, or substitutions of a single amino acid relative to SEQ ID NO:2 or SEQ ID NO:9 or to one of the fragments or portions of SEQ ID NO:2 or SEQ ID NO:9 discussed above. In some embodiments, a BTNL3 protein contains not more than 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 substitutions, insertions, or deletions of a single amino acid relative to SEQ ID NO:2 or SEQ ID NO:9 or relative to one of the fragments of SEQ ID NO:2 or SEQ ID NO:9 discussed above. Such BTNL3 protein variants within the scope of the invention can retain the ability to attenuate T cell proliferation or can act as an

inhibitor or antagonist of this attenuation of T cell proliferation by unaltered BTNL3 protein as assayed by the methods described herein.

[0088] In some embodiments the substitutions can be conservative amino acid substitutions. Examples of conservative amino acid substitutions, unlikely to affect biological activity, include the following: alanine for serine, valine for isoleucine, aspartate for glutamate, threonine for serine, alanine for glycine, alanine for threonine, serine for asparagine, alanine for valine, serine for glycine, tyrosine for phenylalanine, alanine for proline, lysine for arginine, aspartate for asparagine, leucine for isoleucine, leucine for valine, alanine for glutamate, aspartate for glycine, and these changes in the reverse. See e.g. Neurath et al., *The Proteins*, Academic Press, New York (1979), the relevant portions of which are incorporated herein by reference. Further, an exchange of one amino acid within a group for another amino acid within the same group is a conservative substitution, where the groups are the following: (1) alanine, valine, leucine, isoleucine, methionine, norleucine, and phenylalanine; (2) histidine, arginine, lysine, glutamine, and asparagine; (3) aspartate and glutamate; (4) serine, threonine, alanine, tyrosine, phenylalanine, tryptophan, and cysteine; and (5) glycine, proline, and alanine. As discussed above, sites at which conservative substitutions exist when BTNL family members are aligned (as in Table 2) are sites where conservative substitutions are particularly unlikely to affect function. Similarly, highly conserved residues, for example residues 2, 40, 57, and 65, among many others, of SEQ ID NO:2, are likely to be less tolerant of substitution.

[0089] A BTNL3 protein or variant thereof can be at least 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO:2 or SEQ ID NO:9, wherein the alignment window is at least 50, 60, 75, 80, 90, or 100 amino acids long, and wherein the BTNL3 protein can inhibit the proliferation of T cells activated with an immobilized anti-CD3 antibody. As discussed above, sequence mismatches with other human BTNL family members can guide one of skill in the art as to where modifications in the sequence of SEQ ID NO:2 or SEQ ID NO:9 can be made without affecting function. In some embodiments, the insertions, deletions, or substitutions can occur at, or adjacent to, residues that are not conserved among human BTNL family members. See Table 2. In some embodiments, these alterations occur at (in the case of deletions or substitutions) or adjacent to (in the case of insertions) one or more of the following non-conserved residues of SEQ ID NO:2: 22, 25-27, 29, 37, 39, 41, 43, 45-47, 49, 50, 56, 58-62, 71, 73-76, 80, 86, 93, 94, 96, 102, 103, 105-107, 109, 111, 113, 115, 117-123, 126, 127, 130, 132-137, 139-158, 160-162, 164, 166-187, 190, 192-194, 196, 198-200, 202-203, 205-217, 219-222, and/or 224-236. Alterations at the analogous sites in SEQ ID NO:9, where such analogous sites exist, are also contemplated. Alternatively, a BTNL3 protein can contain not more than 1, 2, 3, 4, 5, 6, 8, 10, 15, 20, 25, or 30 amino acid substitutions, deletions, or insertions relative to SEQ ID NO:2 or SEQ ID NO:9. The proteins described above are BTNL3 proteins or variants thereof as meant herein as long as they can inhibit the proliferation of T cells activated by an immobilized anti-CD3 antibody or antagonize such inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7.

[0090] BTNL3 proteins may be glycosylated to varying degrees or not glycosylated. As an illustration, a BTNL3

protein may comprise one or more N- or O-linked glycosylation sites in addition to those already found in a protein comprising SEQ ID NO:2 or SEQ ID NO:9. SEQ ID NO:2 contains two potential N-linked glycosylation sites at positions 103 and 368. SEQ ID NO:9 contains the potential N-linked glycosylation sites at position 103 and 334. The potential N-linked glycosylation site at position 103 of SEQ ID NO:2 or 9 is also present in BTNL8, the BTNL protein most closely related to BTNL3. Thus, this glycosylation site may play a role in BTNL3 function or structure. One of skill in the art would be aware that asparagine residues that are part of the sequence Asn Xxx Ser/Thr (where Xxx is any amino acid except proline) can serve as sites of addition for N-glycans. In addition, there are many serine and threonine residues that may serve as O-linked glycosylation sites. Glycosylation may increase in vivo half life or alter biological activity. Variants of BTNL3 proteins also include proteins comprising one, two, three, four, five, six, seven, eight, nine, or ten more N- and/or O-linked glycosylation sites than are present in SEQ ID NO:2 as long as the resulting protein can inhibit the proliferation of T cells. Variant BTNL3 proteins also include those that have one, two, three, four, or five fewer N- and/or O-linked glycosylation sites than are present in SEQ ID NO:2 as long as they can inhibit the proliferation of T cells activated with an immobilized anti-CD3 antibody or can inhibit the ability of unmutated versions of BTNL3 protein to do so.

[0091] BTNL3 proteins, as meant herein, can be fusion proteins comprising at least one BTNL3 polypeptide, which can comprise an amino acid sequence that is a variant and/or a fragment of SEQ ID NO:2 or SEQ ID NO:9 (as explained above), and at least one other moiety. The other moiety can be a polypeptide other than a BTNL3 protein. The other moiety can also be a non-protein moiety such as, for example, a polyethylene glycol (PEG) moiety or a cytotoxic, cytostatic, luminescent, and/or radioactive moiety.

[0092] Attachment of PEG has been shown to increase the in vivo half life of at least some proteins. Moreover, cytotoxic, cytostatic, luminescent, and/or radioactive moieties have been fused to antibodies for diagnostic or therapeutic purposes, for example, to locate, to inhibit proliferation of, or to kill cells to which the antibodies can bind. Similarly, BTNL3 proteins fused to such moieties can be used to locate, to inhibit proliferation of, or to kill cells that BTNL3 can bind to, for example, cells involved in immune response. Among such cytotoxic, cytostatic, luminescent, and/or radioactive moieties are, for example, maytansine derivatives (such as DM1), enterotoxins (such as a Staphylococcal enterotoxin), iodine isotopes (such as iodine-125), technetium isotopes (such as Tc-99m), cyanine fluorochromes (such as Cy5.5.18), ribosome-inactivating proteins (such as bouganin, gelonin, or saporin-S6), and calicheamicin, a cytotoxic substance that is part of a product marketed under the trademark MYLOTARG™ (Wyeth-Ayerst).

[0093] A variety of polypeptides other than BTNL3 can be fused to a BTNL3 polypeptide for a variety of purposes such as, for example, to increase in vivo half life of the protein, to facilitate identification, isolation and/or purification of the protein, to increase the activity of the protein, and to promote oligomerization of the protein.

[0094] Many polypeptides can facilitate identification and/or purification of a recombinant fusion protein of which they are a part. Examples include polyarginine, polyhistidine, or HAT™ (Clontech), which is a naturally-occurring sequence

of non-adjacent histidine residues that possess a high affinity for immobilized metal ions. BTNL3 proteins comprising these polypeptides can be purified by, for example, affinity chromatography using immobilized nickel or TALON™ resin (Clontech), which comprises immobilized cobalt ions. See e.g. Knol et al. (1996), *J. Biol. Chem.* 27(26): 15358-15366. Polypeptides comprising polyarginine allow effective purification by ion exchange chromatography. Other useful polypeptides include, for example, the antigenic identification peptides described in U.S. Pat. No. 5,011,912 and in Hopp et al. (1988), *Bio/Technology* 6:1204. One such peptide is the FLAG® peptide, which is highly antigenic and provides an epitope reversibly bound by a specific monoclonal antibody, enabling rapid assay and facile purification of expressed recombinant fusion protein. A murine hybridoma designated 4E11 produces a monoclonal antibody that binds the FLAG® peptide in the presence of certain divalent metal cations, as described in U.S. Pat. No. 5,011,912. The 4E11 hybridoma cell line has been deposited with the American Type Culture Collection under Accession No. HB 9259. Monoclonal antibodies that bind the FLAG® peptide can be used as affinity reagents to recover a polypeptide purification reagent that comprises the FLAG® peptide. Other suitable protein tags and affinity reagents are: 1) those described in GST-Bind™ system (Novagen), which utilizes the affinity of glutathione-S-transferase fusion proteins for immobilized glutathione; 2) those described in the T7-TAG® affinity purification kit (Novagen), which utilizes the affinity of the amino terminal 11 amino acids of the T7 gene 10 protein for a monoclonal antibody; or 3) those described in the STREP-TAG® system (Novagen), which utilizes the affinity of an engineered form of streptavidin for a protein tag. Some of the above-mentioned protein tags, as well as others, are described in Sassenfeld (1990), *TIBTECH* 8: 88-93, Brewer et al., in *Purification and Analysis of Recombinant Proteins*, pp.239-266, Seetharam and Sharma (eds.), Marcel Dekker, Inc. (1991), and Brewer and Sassenfeld, in *Protein Purification Applications*, pp. 91-111, Harris and Angal (eds.), Press, Inc., Oxford England (1990). The portions of these references that describe protein tags are incorporated herein by reference. Further, fusions of two or more of the tags described above, such as, for example, a fusion of a FLAG tag and a polyhistidine tag, can be fused to a BTNL3 protein of the invention.

[0095] Recombinant fusion proteins comprising polypeptides other than BTNL3 may have other kinds of unique advantages, such as, for example, a propensity to form dimers, trimers, or higher order multimers, an increased in vivo half-life, and/or an increased biological activity. A “higher order multimer” when used in conjunction with, for example, “dimer,” means a multimer containing more than two polypeptide chains. When used in a phrase like “a trimer or a higher order multimer,” the higher order multimer contains more than three polypeptide chains. Thus, a higher order multimer is one that has more polypeptide chains than the multimer it is compared to. Techniques for preparing fusion proteins are known, and are described, for example, in International Application WO 99/31241 and in Cosman et al. ((2001). *Immunity* 14: 123-133).

[0096] As an illustration, a polypeptide that comprises an Fc region of an antibody, optionally an IgG antibody, or a substantially similar protein, can be fused to a BTNL3 polypeptide or fragment thereof. An Fc region of an antibody is a polypeptide comprising the most or all of hinge

plus the C_H2, and the C_H3 domains from an antibody or immunoglobulin domains substantially similar to these. For discussion, see Hasemann and Capra, *Immunoglobulins: Structure and Function*, in William E Paul, ed., *Fundamental Immunology*, Second Edition, 212-213 (1989). The Fc fragment can be a human IgG Fc, such as an IgG1, IgG2, IgG3, or IgG4 Fc. An Fc fragment can be a native human or animal Fc fragment. Truncated forms of Fc regions, that is, forms missing some portion of the hinge, C_H2, and/or C_H3 domains, that promote dimerization can also be used. Other portions of antibodies and other immunoglobulin isotypes can be used. Recombinant fusion proteins comprising Fc regions of antibodies are likely to form dimers or higher order multimers. Fusion proteins comprising various portions of antibody-derived proteins have been described by Ashkenazi et al. ((1991) *Proc. Natl. Acad. Sci. USA* 88:10535-39), Byrn et al. ((1990), *Nature* 344: 677-70), Hollenbaugh and Aruffo (in *Current Protocols in Immunology*, Suppl. 4, pp. 10.19.1-10.19.11 (1992)), Baum et al. ((1994), *EMBO J.* 13: 3992-4001) and in U.S. Pat. No. 5,457,035 and International Application WO 93/10151, the relevant portions of which are incorporated herein by reference. In some embodiments, an altered Fc region can have the advantage of having a lower affinity for Fc receptors compared to a wild type Fc region. This can be an advantage because it may lessen the lysis of cells to which such recombinant fusion proteins bind by immune effector cells. Such alterations to the Fc region are described in U.S. Pat. No. 5,457,035 and International Patent Application WO 93/10151, the relevant portions of which are incorporated herein by reference. Example 2 below describes the production of a fusion protein containing the extracellular region of human BTNL3 and the Fc region of a human IgG1 antibody.

[0097] Alternatively, a BTNL3 fusion protein can comprise all or part of a BTNL3 polypeptide as described herein plus a non-Fc polypeptide that has the effect of increasing the half life of the fusion protein such as, for example, a serum albumin protein, optionally a human protein, or a fragment thereof, or a protein that binds to serum albumin or a fragment thereof. In some embodiments this additional polypeptide could be all or part of a fibronectin protein, which is, optionally a human protein.

[0098] A recombinant fusion protein comprising a BTNL3 protein can comprise a polypeptide comprising a leucine zipper. Among known leucine zipper sequences are sequences that promote dimerization and sequences that promote trimerization. See e.g. Landschulz et al. (1988), *Science* 240: 1759-64, the relevant portions of which are incorporated herein by reference. Leucine zippers comprise a repetitive heptad repeat, often with four or five leucine residues interspersed with other amino acids. Use and preparation of leucine zippers are well known in the art. A BTNL3 fusion protein can comprise one or more peptide linkers. Generally, a peptide linker is a stretch of amino acids that serves to link plural polypeptides to form multimers and provides the flexibility or rigidity required for the desired function of the linked portions of the protein. Typically, a peptide linker is between about 1 and 30 amino acids in length. Examples of peptide linkers include, but are not limited to, —Gly-Gly—, GGGGS (SEQ ID NO:10), (GGGGS)_n (SEQ ID NO:11), GKSSGSGSESKS (SEQ ID NO:12), GSTSGSGKSSEGKKG (SEQ ID NO:13), GSTSGSGKSSEGSGSTKG (SEQ ID NO:14), GSTSGSGKSSEGKKG (SEQ ID NO:15), GSTSGSGKPGSGEG-

STKG (SEQ ID NO:16), EGKSSGSGSESKEF (SEQ ID NO:17) and GGGGSGGGGSGGGGS (SEQ ID NO:18). Linking moieties are described, for example, in Huston, J. S., et al., *Proc. Natl. Acad. Sci.* 85: 5879-83 (1988), Whitlow, M., et al., *Protein Engineering* 6: 989-95 (1993), Newton, D. L., et al., *Biochemistry* 35: 545-53 (1996), and U.S. Pat. Nos. 4,751,180 and 4,935,233. The relevant portions of these references, that is, the portions describing linkers, are incorporated herein by reference.

[0099] Alternatively, described herein are BTNL3 variant proteins that can antagonize the inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7 of proliferation of a T cell stimulated by an immobilized anti-CD3 antibody. Such BTNL3 variants can comprise an amino acid sequence at least about 90%, 95%, or 98% identical to amino acids 18-236 of SEQ ID NO:2 or 18-166 of SEQ ID NO:9 and/or can comprise an amino acid sequence containing not more than 20, 15, 10, 8 or 5 insertions, deletions, or substitutions of a single amino acid relative to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9. Such variants do not contain an amino acid sequence identical to amino acids 18-236 of SEQ ID NO:2 or amino acid 18-166 of SEQ ID NO:9.

[0100] A recombinant BTNL3 fusion protein can comprise a BTNL3 protein that lacks its normal signal sequence and has, instead, a different signal sequence replacing it. The choice of a signal sequence depends on the type of host cells in which the recombinant protein is to be produced, and a different signal sequence can replace the native signal sequence. Examples of signal sequences that are functional in mammalian host cells include the following: the signal sequence for interleukin-7 (IL-7) described in U.S. Pat. No. 4,965,195; the signal sequence of human IgK (which is METDTLLWVLLWVPGSTG; SEQ ID NO:3); the signal sequence for human growth hormone (that is, MATGSRTSLLLAFGLLCLPWLEQSGA; SEQ ID NO:4); the signal sequence for interleukin-2 receptor described in Cosman et al. ((1984), *Nature* 312: 768); the interleukin-4 receptor signal peptide described in EP Patent 0 367 566; the type I interleukin-1 receptor signal peptide described in U.S. Pat. No. 4,968,607; and the type II interleukin-1 receptor signal peptide described in EP Patent 0 460 846. The portions of these references describing these signal sequences are incorporated herein by reference.

BTNL3 Nucleic Acids

[0101] The invention encompasses isolated nucleic acids, including, for example, DNAs and RNAs that encode the BTNL3 proteins described herein, which include proteins comprising the amino acid sequence of SEQ ID NO:2 or SEQ ID NO:9 and fragments and/or variants thereof. Such nucleic acids include DNAs that do not include exon sequences. These nucleic acids are useful for, inter alia, producing recombinant proteins and detecting the presence of BTNL3 nucleic acids in tissue samples, e.g., for diagnostic uses. Such nucleic acids can be genomic DNA or cDNA. As is known in the art, a cDNA sequence does not include exon sequence that can be present in genomic DNA. The nucleic acids can comprise an uninterrupted open reading frame encoding a BTNL3 protein. Nucleic acid molecules of the invention include DNA and RNA in both single-stranded and double-stranded form, as well as the corresponding complementary sequences. An "isolated nucleic acid" is a nucleic acid that has been separated from

adjacent genetic sequences present in the genome of the organism from which the nucleic acid was isolated, in the case of nucleic acids isolated from naturally-occurring sources. In the case of nucleic acids synthesized chemically, such as oligonucleotides, or enzymatically from a template, such as polymerase chain reaction (PCR) products or cDNAs, it is understood that the nucleic acids resulting from such processes are isolated nucleic acids. An isolated nucleic acid molecule refers to a nucleic acid molecule in the form of a separate fragment or as a component of a larger nucleic acid construct such as a vector.

[0102] Further, the invention encompasses fragments of a nucleic acid encoding a BTNL3 protein that can serve (1) as probes for detecting BTNL3 nucleic acids by a number of methods well known in the art, e.g., Southern and northern blotting, dot blotting, colony hybridizations, hybridization to an array, etc., (2) as polymerase chain reaction (PCR) primers to amplify BTNL3 nucleic acids, or (3) as a means to regulate expression of BTNL3 nucleic acids, e.g., through inhibition of expression with antisense nucleic acids (including peptide nucleic acids), ribozymes, triple helix-forming molecules, or interfering RNAs. DNAs that encode any of these RNAs are also BTNL3 nucleic acids as meant herein. PCR primers can comprise, in addition to BTNL3 nucleic acid sequences, other sequences such as restriction enzyme cleavage sites that facilitate the use of the amplified nucleic acid. PCR is described, for example, in the following references: Saiki et al. (1988), *Science* 239: 487-91; *PCR Technology*, Erlich, ed., Stockton Press, (1989). As explained below, PCR can be useful to detect over- or under-expression of BTNL3 mRNAs, and PCR primers can be taken from various parts of the BTNL3 gene and can also be selected to distinguish between different splice variants. Antisense RNAs (and DNAs encoding them), DNAs, or synthetic nucleotides and their use to regulate expression are well known in the art and are described in, e.g. Izant and Weintraub (1984), *Cell* 36(4): 1007-15; Izant and Weintraub (1985), *Science* 229(4711): 345-52; Harel-Bellan et al. (1988), *J. Exp. Med.* 168(6): 2309-18; Sarin et al. (1988), *Proc. Natl. Acad. Sci. USA* 85(20): 7448-51; Zon (1988), *Pharm. Res.* 5(9): 539-49; Harel-Bellan et al. (1988), *J. Immunol.* 140(7): 2431-35; Marcus-Sekura et al. (1987), *Nucleic Acids Res.* 15(14): 5749-63; Gambari (2001), *Curr. Pharm. Des.* 7(17): 1839-62; and Lemaitre et al. (1987), *Proc. Natl. Acad. Sci. USA* 84(3): 648-52. The portions of these references describing techniques of modulating gene expression using nucleic acids are incorporated by reference herein. Similarly, interfering RNAs (and DNAs encoding them) and their use to inhibit expression of selected genes are well known in the art and described in, e.g., Fjose et al. (2001), *Biotechnol. Ann. Rev.* 7: 31-57; Boshier and Labouesse (2000), *Nature Cell Biol.* 2: E31-E36. The relevant portions of these references are incorporated herein by reference. Further, ribozymes or DNazymes can be targeted to cleave specific RNAs and thus used to inhibit gene expression as described in, e.g., Lewin and Hauswirth (2001), *Trends Mol. Med.* 7(5): 221-28; Menke and Hobom (1997), *Mol. Biotechnol.* 8(1): 17-33; Norris et al. (2000), *Adv. Exp. Med. Biol.* 465: 293-301; Sioud (2001), *Curr. Mol. Med.* 1(5): 575-88; and Santiago and Khachigian (2001), *J. Mol. Med.* 79(12): 695-706. The portions of these references describing these methods of modulating gene expression are incorporated by reference herein. Such nucleic acids that can regulate BTNL3 expression can find

use in in vivo or in vitro studies of BTNL3 function or as therapeutics, optionally, as gene therapy agents.

[0103] The present invention also includes nucleic acids, for example, DNAs, comprising the sequence of SEQ ID NO:1 or SEQ ID NO:8 or a fragment thereof or nucleic acids that hybridize under moderately stringent conditions, and optionally highly stringent conditions, to nucleic acids comprising the nucleotide sequence of SEQ ID NO:1 or SEQ ID NO:8, which are BTNL3 cDNAs, wherein the nucleic acid encodes a protein that can inhibit the proliferation of T cells activated with an immobilized anti-CD3 antibody. Hybridization techniques are well known in the art and are described by Sambrook, J., E. F. Fritsch, and T. Maniatis (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., chapters 9 and 11, (1989)) and *Current Protocols in Molecular Biology* (F. M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4 (1995)), the relevant portions of which are incorporated by reference herein. Generally, moderately stringent conditions for filter hybridizations include hybridization in about 50% formamide, 6×SSC at a temperature from about 42° C. to 55° C. and washing at about 60° C. in 0.5×SSC, 0.1% SDS. Generally, highly stringent conditions are defined as hybridization conditions as above, but with washing at approximately 68° C. in 0.2×SSC, 0.1% SDS. SSPE (1×SSPE is 0.15 M NaCl, 10 mM NaH₂PO₄, and 1.26 mM EDTA, pH 7.4) can be substituted for SSC (1×SSC is 0.15 M NaCl and 15 mM sodium citrate) in the hybridization and wash buffers; washes, optionally at least two washes, are performed for 15 minutes after hybridization is complete.

[0104] It should be understood that the wash temperature and wash salt concentration can be adjusted as necessary to achieve a desired degree of stringency by applying the basic principles that govern hybridization reactions and duplex stability, as known to those skilled in the art and described further below (see e.g., Sambrook et al., supra). When nucleic acids of known sequence are hybridized, the hybrid length can be determined by aligning the sequences of the nucleic acids (for example, using GAP) and identifying the region or regions of optimal sequence complementarity. Somewhat different conditions than those described above can apply for hybrids of relatively short lengths. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5 to 10° C. less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, T_m (degrees C.)=2(# of A+T bases)+4(# of G+C bases). For hybrids above 18 base pairs in length, T_m (degrees C.)=81.5+16.6 (log₁₀ [Na⁺])+0.41(% G+C)=(600/N), where N is the number of bases in the hybrid, and [Na⁺] is the concentration of sodium ions in the hybridization buffer. Each such hybridizing nucleic acid can have a length that is at least 15 nucleotides, or at least 18 nucleotides, or at least 20 nucleotides, or at least 25 nucleotides, or at least 30 nucleotides, or at least 40 nucleotides, or at least 50 nucleotides, or at least 100 nucleotides. Sambrook et al., supra.

[0105] BTNL3 nucleic acids, such as cDNAs, include nucleic acids comprising the following polynucleotides: (1) all or a fragment of SEQ ID NO:1 or SEQ ID NO:8, wherein the fragment encodes a BTNL3 protein that can inhibit proliferation of T cells; (2) a polynucleotide including nucleotide sequences at least 80%, 85%, 90%, 95%, 97%,

98%, 99%, 99.5%, or 99.7% identical to SEQ ID NO:1 or SEQ ID NO:8, wherein the alignment window is at least 100, 125, 150, 175, 200, 225, 250, 300, 400, 500, 600, 800, 1000, or 1200 nucleotides long and wherein the sequence encodes a BTNL3 protein that can inhibit the proliferation of T cells activated with an immobilized anti-CD3 antibody or can antagonize such inhibition; (3) fragments of SEQ ID NO:1 or SEQ ID NO:8 or substantially similar sequences that are useful for detecting or amplifying nucleic acids encoding the BTNL3 proteins of the invention or for regulating the expression of BTNL3 mRNAs and/or proteins; (4) a polynucleotide that comprises not more than 1, 2, 3, 4, 6, 8, 10, 15, 20, 25, or 30 alteration(s) of a single nucleotide relative to SEQ ID NO:1 or SEQ ID NO:8, wherein an alteration can be an insertion, deletion or substitution of a single nucleotide, and wherein the polynucleotide encodes a BTNL3 protein that can inhibit the proliferation of T cells activated with an immobilized anti-CD3 antibody or can serve as an antagonist of such inhibition; and (5) a polynucleotide that encodes a BTNL3 protein as described herein, which includes fragments, derivatives and variants of a human BTNL3 protein.

Methods of Making BTNL3 Proteins and Anti-BTNL3 Antibodies

[0106] BTNL3 proteins, including multimeric BTNL3 proteins or variant BTNL3 proteins, or anti-BTNL3 antibodies (or anti-idiotypic antibodies) can be made as follows. A nucleic acid, for example a DNA, that encodes a BTNL3 protein or an anti-BTNL3 antibody, as described herein, can be introduced into a vector, which can be introduced into a host cell, for example a non-human host cell, which can be a mammalian cell. Vectors and host cells comprising nucleic acids, such as DNAs, encoding a BTNL3 protein or an anti-BTNL3 antibody are encompassed by the invention. The host cell containing the nucleic acids encoding a BTNL3 protein or an anti-BTNL3 antibody can be cultured under conditions such that the BTNL3 protein or the anti-BTNL3 antibody can be expressed. The expressed BTNL3 protein or anti-BTNL3 antibody can then be obtained from the medium in which the cells are cultured or from the cells and purified by any of the many appropriate means known in the art. In addition, genetic engineering methods for the production of BTNL3 proteins include the expression of the polynucleotide molecules in cell free expression systems, in cellular hosts, in tissues, and in animal models, according to known methods.

[0107] The vector can include a selectable marker and an origin of replication, for propagation in a host. The vector can further include suitable transcriptional or translational regulatory sequences, such as those derived from mammalian, microbial, viral, or insect genes, operably linked to the nucleic acid encoding the BTNL3 protein or the anti-BTNL3 antibody. Examples of such regulatory sequences include transcriptional promoters, operators, or enhancers, mRNA ribosomal binding sites, and appropriate sequences that control transcription and translation. Nucleotide sequences are operably linked when the regulatory sequence functionally relates to the DNA encoding the target protein. Thus, a promoter nucleotide sequence is operably linked to a nucleic sequence if the promoter nucleotide sequence directs the transcription of the anti-BTNL3 antibody-encoding or BTNL3 protein-encoding sequence. If the BTNL3 protein is a fusion protein, a nucleic acid sequence encoding a portion

of the fusion protein, for example, a signal sequence, can be part of a vector, and a nucleic acid encoding an anti-BTNL3 antibody or a BTNL3 protein can be inserted into the vector such that a protein comprising the added signal sequence plus the anti-BTNL3 antibody or BTNL3 protein is encoded by the vector.

[0108] Suitable host cells for expression of BTNL3 proteins or anti-BTNL3 antibodies include human and non-human cells including prokaryotic cells such as bacteria, yeast cells, plant cells, insect cells, and higher eukaryotic cells, including mammalian cells. The regulatory sequences in the vector will be chosen such that they are operable in the host cell. Suitable prokaryotic host cells include bacteria of the genera *Escherichia*, *Bacillus*, and *Salmonella*, as well as members of the genera *Pseudomonas*, *Streptomyces*, and *Staphylococcus*. For expression in prokaryotic cells, for example, in *E. coli*, the polynucleotide molecule encoding a BTNL3 protein or anti-BTNL3 antibody preferably includes an N-terminal methionine residue to facilitate expression of the recombinant polypeptide. The N-terminal methionine may optionally be cleaved from the expressed polypeptide. Suitable yeast host cells include cells from genera including *Saccharomyces*, *Pichia*, and *Kluyveromyces*, including species such as *S. cerevisiae* and *P. pastoris*. A suitable system for expression in an insect host cell is described, for example, in the review by Luckow and Summers ((1988), *BioTechnology* 6: 47), the relevant portions of which are incorporated herein by reference. Suitable mammalian host cells include the COS-7 line of monkey kidney cells (Gluzman et al. (1981), *Cell* 23: 175-182), baby hamster kidney (BHK) cells, Chinese hamster ovary (CHO) cells (Puck et al. (1958), *PNAS USA* 60: 1275-1281), CV-1 (Fischer et al. (1970), *Int. J. Cancer* 5: 21-27), 293 cells from human kidney (American Type Culture Collection (ATCC®) catalog no. CRL-10852™), and human cervical carcinoma cells (HELA) (ATCC® CCL 2). The relevant portions of the references referred to in this paragraph are incorporated herein by reference.

[0109] Expression vectors for use in cellular hosts generally comprise one or more phenotypic selectable marker genes. Such genes encode, for example, a protein that confers antibiotic resistance or that supplies an auxotrophic requirement. A wide variety of such vectors are readily available from commercial sources. Examples include pGEM® vectors (Promega), pSPORT™ vectors, and pPROEXTM vectors (Invitrogen, Life Technologies, Carlsbad, Calif.), Bluescript vectors (Stratagene), and pQE vectors (Qiagen). Yeast vectors will often contain an origin of replication sequence from a 2 μ yeast plasmid, an autonomously replicating sequence (ARS), a promoter region, sequences for polyadenylation, sequences for transcription termination, and a selectable marker gene. Vectors replicable in both yeast and *E. coli* (termed shuttle vectors) may also be used. In addition to the above-mentioned features of yeast vectors, a shuttle vector will also include sequences for replication and selection in *E. coli*. Direct secretion of the target polypeptides expressed in yeast hosts may be accomplished by the inclusion of nucleotide sequence encoding the yeast a-factor leader sequence at the 5' end of the BTNL3- or antibody-encoding nucleotide sequence. Brake (1989), *Biotechnology* 13: 269-280.

[0110] Examples of suitable expression vectors for use in mammalian host cells include pcDNA3.1/Hygro+ (Invitrogen), pDC409 (McMahan et al. (1991), *EMBO J.* 10:

2821-2832), and pSVL (Pharmacia Biotech). Expression vectors for use in mammalian host cells can include transcriptional and translational control sequences derived from viral genomes. Commonly used promoter sequences and enhancer sequences that can be used to express BTNL3 RNA include, but are not limited to, those derived from human cytomegalovirus (CMV), Adenovirus 2, Polyoma virus, and Simian virus 40 (SV40). Methods for the construction of mammalian expression vectors are disclosed, for example, in Okayama and Berg ((1982) *Mol. Cell. Biol.* 2:161-170), Cosman et al. ((1986) *Mol. Immunol.* 23:935-941), Cosman et al. ((1984) *Nature* 312: 768-771), EP-A-0367566, and WO 91/18982. The relevant portions of these references are incorporated herein by reference.

Anti-BTNL3 Antibodies

[0111] Antibodies that bind specifically to the BTNL3 proteins described herein, anti-idiotypic antibodies that bind to anti-BTNL3 antibodies and uses of these antibodies are described herein. An anti-BTNL3 antibody can bind to a polypeptide, wherein the amino acid sequence of the polypeptide consists of the amino acids 18 to 236 of SEQ ID NO:2 or amino acids 18 to 166 of SEQ ID NO:9. As used herein, specific binding of an epitope on a BTNL3 protein by a first antibody means that the first antibody can be displaced from the BTNL3 protein by another antibody that competes with the first antibody, but not by other anti-BTNL3 antibodies that do not compete with the first antibody for binding. Numerous competitive binding assays are known in the art.

[0112] Typically competition of antibodies for binding can be evaluated by a fluorescence activated cell sorting (FACS) assay. The antibody of interest can be biotinylated. The biotinylated antibodies are combined with cells known to express the antigen to which the antibodies bind. If the biotinylated antibodies bind to the cells as expected, a shift in fluorescence intensity should be observed using a secondary consisting of streptavidin conjugated to a fluorochrome. Pre-incubation of the cells with an unlabeled version of the same antibody should completely eliminate the observed shift in fluorescence. Pre-incubation with a different unlabeled antibody may completely or partially eliminate the fluorescence shift or have no effect. In the later case, one would conclude that the unlabeled antibody does not compete with the labeled antibody. In the former case, the antibodies do compete for binding, and, as meant herein, one would conclude that the epitopes are either fully or partially overlapping. Alternatively, the unlabeled antibody may compete by altering the conformation of the epitope bound by the labeled antibody. Among the antibodies contemplated are those that compete, either fully or partially, with any specifically provided anti-BTNL3 antibody.

[0113] In some embodiments, an anti-BTNL3 antibody can bind to a human BTNL3 protein with an equilibrium dissociation constant (KD) of no more than 10^{-8} , 10^{-9} , 10^{-10} , or 10^{-11} M. A KD is commonly determined by use of a specialized instrument that can detect interactions between proteins in real time using surface plasmon resonance, such as those sold by Biacore Life Sciences, Piscataway, N.J. In some embodiments, an anti-BTNL3 antibody can bind to a BTNL3 protein comprising an amino acid sequence at least 90%, 95%, 97%, 98%, or 100% identical to SEQ ID NO:2 and/or SEQ ID NO:9, wherein the alignment window with SEQ ID NO:2 and/or 9 is at least 50, 60, 70, 80, or 100

amino acids long and, optionally, wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody or can antagonize such inhibition by a BTNL3 protein comprising the amino acid sequence of SEQ ID NO:7. In some embodiments, an anti-BTNL3 antibody can bind to a protein comprising a fragment of the amino acid sequence of SEQ ID NO:2 and/or 9.

[0114] In addition, the impact of an anti-BTNL3 antibody on activation of anti-CD3-activated T cells in the presence or absence of a BTNL3 protein may provide additional useful information about the functional properties of an antibody. Particularly, antibodies that prevent or inhibit BTNL3 from exerting its usual role of inhibiting the proliferation of anti-CD3-activated T cells are contemplated. Alternatively, antibodies that agonize or enhance the inhibition of proliferation of anti-CD3-activated T cells by BTNL3 are contemplated. Thus, the antibodies can be either antagonists or agonists of BTNL3. Since BTNL3 contains an intracellular B30.2 domain, it may transduce a signal into the cell on which it is expressed when its extracellular domain is engaged by an agonist, such as a naturally occurring ligand or an agonistic antibody. This kind of agonistic antibody can be distinct from antibodies that enhance the inhibition of proliferation of T cells. The invention includes monoclonal antibodies, each of which binds to a particular epitope of BTNL3, and monoclonal antibodies that compete with these for binding. The antibodies can be human, humanized, or chimeric antibodies and can be in various formats.

[0115] Epitopes on BTNL3 protein may comprise contiguous amino acids and also may comprise non-contiguous amino acids. Epitopes can be identified by methods known in the art, including the use of protein fragment or peptide libraries, alanine scanning, epitope extraction, epitope excision, or X-ray crystallography. See e.g. Leinonen et al. (2002), *Clin. Chem.* 48(12): 2208-16; Kroger et al. (2002), *Biosens. Bioelectron.* 17(11-12): 937-44; Zhu et al. (2001), *Biochem. Biophys. Res. Commun.* 282(4): 921-27; Obungu et al. (2009), *Biochemistry* 48: 7251-60. The relevant portions of these references, i.e., the portions describing methods of epitope mapping, are incorporated herein by reference. In addition, especially in view of the fact that BTNL3 includes some domains of well known tertiary structure, such as an IgV-like domain, tertiary structures of BTNL3 generated by analogy to known tertiary structures can be used as an aid to determining epitopes.

[0116] Antibodies can be polyclonal or monoclonal antibodies and can be produced by methods well known in the art. See, for example, *Monoclonal Antibodies, Hybridomas: A New Dimension in Biological Analyses*, Kennet et al. (eds.), Plenum Press, New York (1980); and *Antibodies: A Laboratory Manual*, Harlow and Land (eds.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (1988); Kohler and Milstein (1980) *Proc. Natl. Acad. Sci., USA*, 77: 2197; Kozbor et al. (1984), *J. Immunol.* 133: 3001-3005 (describing the human B-cell hybridoma technique); Cole et al., *Monoclonal Antibodies And Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96 (1985) (which describes EBV-hybridoma technique); Kuby, *Immunology*, Second Edition, p.162-64, W.H. Freeman and Co., New York (1994); the relevant portions of these references are incorporated herein by reference. Hybridoma cell lines that produce monoclonal antibodies specific for the BTNL3 proteins as described

herein are also contemplated herein. Such hybridoma lines can be produced and identified by conventional techniques. The hybridoma producing an anti-BTNL3 antibody can be cultivated in vitro or in vivo. Further, anti-BTNL3 antibodies can be produced in other cultured cells, including, for example, Chinese hamster ovary (CHO), HeLa, VERO, BHK, Cos, MDCK, 293, 3T3, myeloma (e.g. NSO, NSI), or WI38 cells, yeast cells, insect cells, and bacterial cells, including, for example, *Escherichia coli*. Such antibodies can be produced by introducing nucleic acids, such as DNAs, encoding the antibodies plus nucleic acids to enable expression of these nucleic acids into desired host cells. The antibodies can then be produced by culturing the cells into which these nucleic acids have been introduced. Monoclonal antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD and any subclass thereof, such as, for example, IgG1, IgG2, IgG3, or IgG4.

[0117] Anti-BTNL3 antibodies can be full-length tetrameric antibodies comprising two heavy chains and two light chains, like those found in most mammalian species. Such a full length antibody can be of the IgA, IgD, IgG, IgM, or IgE isotype. If it is an IgG antibody, it can be an IgG1, IgG2, IgG3, or IgG4 antibody. Alternatively, anti-BTNL3 antibodies can be single chain antibodies comprising a heavy and a light chain variable region and, optionally, also one or more constant region-like domain (U.S. Pat. No. 4,946,778; Bird et al. (1988), *Science* 242: 423-26; Huston et al. (1988), *Proc. Natl. Acad. Sci. USA* 85: 5879-83), dimeric or multivalent antibodies (see e.g. Lantto et al. (2002), *J. Gen. Virol.* 83: 2001-05; Hudson and Souriau (2001), *Expert Opin. Biol. Ther.* 1(5): 845-55), tetrameric antibodies (see e.g. Janeway et al., *Immunobiology: The Immune System in Health and Disease*, Fifth Edition, Part II, Ch. 3, Garland Publishing (2001)), chimeric antibodies (Hudson and Souriau, supra; Boulianne et al. (1984), *Nature* 312:643-46; Morrison et al. (1984), *Proc. Natl. Acad. Sci. USA* 81: 6851-55; Takeda et al. (1985), *Nature* 314: 452-54; Neuberger et al. (1985), *Nature* 314: 268-70), fully human antibodies produced in a non-human transgenic mammal (described in e.g., U.S. Pat. No. 6,150,584) or by in vitro selection (US Patent Application 2002/0058033) or humanized antibodies (Morrison et al., supra; Takeda et al., supra; Boulianne et al., supra). Further, antibodies can be "matured" by in vitro selection schemes to yield an antibody with altered properties such as, for example, a higher affinity for the epitope to which it binds. See e.g. Jackson et al. (1995), *J. Immunol.* 154(7): 3310-19; Pini and Bracci (2000), *Curr. Protein Pept. Sci.* 1(2): 155-69; Ellmark et al. (2002), *Mol. Immunol.* 39(5-6): 349; O'Connell et al. (2002), *J. Mol. Biol.* 321(1): 49-56; Huls et al. (2001), *Cancer Immunol. Immunother.* 50: 163-71; Hudson and Souriau, supra; Adams and Schier (1999), *J. Immunol. Methods* 231(1-2): 249-60; Schmitz et al. (2000), *Placenta* 21 Suppl. A: S106-12. Alternatively, fragments of antibodies such as, for example, Fab fragments, F(ab')₂ fragments, or single chain Fv fragments (scFv's) that can bind specifically to a BTNL3 protein of the invention are also encompassed by what is meant herein as an anti-BTNL3 antibody. See Kuby, supra, pp.109-112 and Janeway et al., supra, for discussion of Fab and Fv fragments. The invention also encompasses anti-idiotypic antibodies that bind specifically to antibodies that bind specifically to BTNL3 proteins and that mimic the effects of BTNL3 proteins. Such anti-idiotypic antibodies find the same uses as BTNL3 proteins.

Methods for generating anti-idiotypic antibodies are well known in the art. See e.g. Kuby et al., *supra*, at 371-72. Various kinds of recombinant and non-recombinant bispecific antibodies that can bind specifically to a BTNL3 protein of the invention and another protein are also contemplated. Various kinds of bispecific antibodies and methods for making them are described in e.g. U.S. Pat. Nos. 4,474,893, 6,060,285, and 6,106,833.

[0118] The anti-BTNL3 antibodies can be multimeric antibodies, including full-length, tetrameric, bispecific antibodies containing two complete heavy chains and two complete light chains or multimeric monovalent antibodies containing, for example, a heavy chain plus a light chain plus an Fc region. Such multimeric antibodies can contain certain mutations in their Fc region that facilitate the formation of heterodimers. Such antibodies and mutations are described in International Patent Publication No. International Application WO 2009/089004 and US Application 2007/0105199, the portions of which describe such antibodies and mutations are incorporated by reference herein. The Fc regions in such antibodies can have native human sequences or sequences native to other species. Alternatively or in addition, the Fc regions of such antibodies can contain mutations in their Fc regions that either increase or decrease effector function by increasing or decreasing the affinity of various Fc receptors for the Fc region. Some such Fc alterations are discussed in U.S. Pat. No. 5,457,035 and International Patent Application Publication No. WO 93/10151, the relevant portions of which are incorporated herein by reference.

[0119] Anti-BTNL3 antibodies can also be conjugated to moieties can be used to locate, to inhibit proliferation of, or to kill cells that antibody can bind to. Among such cytotoxic, cytostatic, luminescent, and/or radioactive moieties are, for example, maytansine derivatives (such as DM1), enterotoxins (such as a Staphylococcal enterotoxin), iodine isotopes (such as iodine-125), technetium isotopes (such as Tc-99m), cyanine fluorochromes (such as Cy5.5.18), ribosome-inactivating proteins (such as bouganin, gelonin, or saporin-S6), and calicheamicin, a cytotoxic substance that is part of a product marketed under the trademark MYLOTARG™ (Wyeth-Ayerst).

[0120] An antibody may contain only a single heavy or light chain variable region, optionally fused to another portion of an antibody as described in US Patent Application 2004/058820, the portions of which describe these single domain antibodies are incorporated herein by reference.

[0121] Some naturally-occurring antibodies, which have been found in camels and llamas, are dimers consisting of two heavy chains and include no light chains. Muldermans et al., 2001, *J. Biotechnol.* 74:277-302; Desmyter et al., 2001, *J. Biol. Chem.* 276:26285-90, the portions of which describe the structures of these antibodies are incorporated herein by reference. Anti-BTNL3 antibodies having this structure are among the anti-BTNL3 antibodies of the invention.

[0122] Anti-BTNL3 antibodies can have a variety of activities and uses. Anti-BTNL3 antibodies may be antagonistic antibodies that block or inhibit the biological function of BTNL3, for example by blocking or antagonizing BTNL3-dependent inhibition of T cell proliferation, which can be assayed by the methods described in the Examples herein. The antibodies can also be agonistic antibodies. In some embodiments, agonistic antibodies can bind to the

BTNL3 counterstructure and mimic BTNL3 binding to inhibit T cell activation or proliferation. Agonistic antibodies may also be anti-idiotypic antibodies that bind to anti-BTNL3 antibodies and/or also bind to the BTNL3 counterstructure and mimic the activity of

[0123] BTNL3, that is, they inhibit the proliferation of T cells as described herein. Such agonistic anti-BTNL3 antibodies can be used for the same uses as a BTNL3 protein. Anti-BTNL3 antibodies can be agonistic or antagonistic by, for example, stabilizing or disrupting the BTNL3 protein, possibly in combination with other proteins, on the cell surface. For example, an agonistic antibody may enhance the activity of BTNL3 by stabilizing the transmembrane form of BTNL3, or by stabilizing the interaction of a BTNL3 protein with other BTNL3 proteins or different proteins on the surface of a cell or the clustering of BTNL3 protein on the surface of cells. Further, an antagonistic antibody may inhibit BTNL3 activity by destabilizing the transmembrane form of BTNL3, or by destabilizing interactions among multiple molecules of BTNL3 or interactions of BTNL3 with other proteins, on the cell surface of cells. Agonistic anti-BTNL3 antibodies can also bind transmembrane forms of BTNL3, causing it to transduce a biological signal into the cell on which it is expressed. Such a signal, may, for example, decrease or increase the immune function of an immune cell, e.g., a neutrophil, expressing BTNL3. In some embodiments, an agonistic anti-BTNL3 antibody may cause a decrease in proliferation or cell death of cells expressing BTNL3, such as cancer cells or neutrophils. An antagonistic anti-BTNL3 antibody can be used to enhance an immune response. Hence, antagonistic anti-BTNL3 antibodies can find use, for example, in the treatment of cancer or in a vaccine, for example in a vaccine to induce a response to a cancer-specific antigen.

[0124] Anti-BTNL3 antibodies can be conjugated to a cytotoxic, cytostatic, luminescent, and/or radioactive moiety. Such antibodies can be used to locate, to inhibit proliferation of, or to kill cells that express BTNL3. Since some cancer cells express BTNL3, such antibody conjugates can be used to treat such cancers. Similarly, since neutrophils express BTNL3, such antibody conjugates can be used to treat conditions characterized by excessive numbers of neutrophils or excessive neutrophil activity, for example, asthma, inflammatory bowel disease including Crohn's disease and ulcerative colitis, COPD, and gout and related inflammatory crystal deposition diseases. Among such cytotoxic, cytostatic, luminescent, and/or radioactive moieties are, for example, maytansine derivatives (such as DM1), enterotoxins (such as a Staphylococcal enterotoxin), iodine isotopes (such as iodine-125), technetium isotopes (such as Tc-99m), cyanine fluorochromes (such as Cy5.5.18), ribosome-inactivating proteins (such as bouganin, gelonin, or saporin-S6), and calicheamicin, a cytotoxic substance that is part of a product marketed under the trademark MYLOTARG™ (Wyeth-Ayerst).

[0125] The antibodies of the invention can also be used in assays to detect the presence of the BTNL3 proteins of the invention, either in vitro or in vivo. The antibodies also can be employed in purifying BTNL3 proteins of the invention by immunoaffinity chromatography.

Agonists and Antagonists of BTNL3 Polypeptides

[0126] In addition to antagonistic or agonistic antibodies, other antibody-related molecules that can bind specifically

BTNL3 proteins, such as affibodies (Ronnmark et al. (2002), *J. Immunol. Methods* 261(1-2): 199-211, the portion of which describes affibodies is incorporated by reference herein) and the biologically active peptides described in International Application WO 00/24782 (the portions of which describe these peptides are incorporated herein by reference) that can bind specifically to BTNL3 and inhibit the biological activity of BTNL3 proteins are encompassed by the invention. Further, BTNL3 antagonists include the nucleic acids described above that are useful for modulating expression of BTNL3 protein and/or mRNA, such as, for example, interfering RNAs (or DNAs that encode them) or antisense RNAs or DNAs.

[0127] Antagonists further include proteins that comprise amino acid sequences selected in vitro to bind to BTNL3 or its receptor and that can, optionally, interfere with the interaction of BTNL3 and its receptor. In some embodiments such proteins can be variant forms of BTNL3 that lack the normal activity of BTNL3 and essentially act as decoys. Alternatively, such proteins can be BTNL3 agonists that promote or mimic the biological function of BTNL3. Proteins that bind to BTNL3 or its receptor can be screened for their ability to interfere with the interaction of BTNL3 with its receptor, or, alternatively, a selection can be designed to obtain such proteins directly. Alternatively, a BTNL3 antagonist can be a variant BTNL3 protein that interferes with the biological activity of the wild type BTNL3 protein, for example, a soluble, biologically inactive BTNL3 variant that blocks the interaction of wild type BTNL3 with its receptor.

[0128] Proteins may be selected by a number of methods such as, for example, phage display or display of the surface of a bacterium. See e.g. Parmley and Smith (1989), *Adv. Exp. Med. Biol.* 251: 215-218; Luzzago et al. (1995), *Biotechnol. Annu. Rev.* 1: 149-83; Lu et al. (1995), *Biotechnology (NY)* 13(4): 366-372. In these methods, each member of a library of binding domains can be displayed on individual phage particles or bacterial cells, and bacteria or phage that bind to a protein of interest under chosen conditions can be selected. Nucleic acids encoding the selected binding domains can be obtained by growing the selected phage or bacteria and isolating nucleic acids from them.

[0129] Alternatively, a protein can be selected entirely in vitro. For example, each individual polypeptide in a library of potential binding domains can be attached to nucleic acids encoding it, and those that bind to the protein of interest under chosen conditions can be selected. Since the polypeptides are attached to nucleic acids encoding them, subsequent operations, such as amplifying, cloning, or sequencing nucleic acids encoding effective binding domains are facilitated. Various schemes for such selections are known in the art, including antibody-ribosome-mRNA particles, ribosome display, covalent RNA-peptide fusions, or covalent DNA-RNA-peptide fusions. He and Taussig (1997), *Nucleic Acids. Res.* 25(24): 5132-5134; Hanes and Pluckthun (1997), *Proc. Natl. Acad. Sci.* 94: 4937-4942; Roberts and Szostak (1997), *Proc. Natl. Acad. Sci.* 94: 12297-12302; Lohse and Wright (2001), *Curr. Opin. Drug Discov. Devel.* 4(2): 198-204; Kurz et al. (2000), *Nucleic Acids Res.* 28(18): E83; Liu et al. (2000), *Methods Enzymol.* 318: 268-93; Nemoto et al. (1997), *FEBS Lett.* 414(2): 405-08; US Patent 6,261,804; International Applications WO 00/32823; and WO 00/34784. The portions of these publications that describe how such selections can be done are

incorporated by reference herein. Such proteins can be selected to be antagonists or agonists.

Therapeutic Uses

[0130] It is demonstrated herein that a BTNL3.Fc fusion protein can inhibit proliferation of activated T cells. See Example 4; FIG. 5. BTNL3 also reduces the activation state of CD4⁺ T cells in response to an immobilized anti-CD3 antibody as reflected in a changed gene expression profile. See Example 5; FIG. 6. Also, it is shown herein that BTNL3 is relatively highly expressed on neutrophils and some kinds of cancer cells. FIGS. 2 and 4. In some situations, polymorphic forms of BTNL3 expressed on neutrophils could possibly alter immune homeostasis. Hence, BTNL3, or a molecule with the ability to agonize the BTNL3 pathway, may be useful as a therapeutic to treat autoimmune or inflammatory diseases that are mediated by T cells or neutrophils. These diseases include, for example, arthritis including rheumatoid arthritis, juvenile idiopathic arthritis, and psoriatic arthritis, Addison's disease, asthma, polyglandular endocrinopathy syndromes, systemic lupus erythematosus, chronic active hepatitis, thyroiditis, lymphocytic adenohypophysitis, premature ovarian failure, idiopathic hypoparathyroidism, pernicious anemia, glomerulonephritis, autoimmune neutropenia, Goodpasture's syndrome, myasthenia gravis, scleroderma, primary Sjogren's syndrome, polymyositis, autoimmune hemolytic anemia, an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, psoriasis, dermatitis, sarcoidosis, multiple sclerosis, chronic obstructive pulmonary disease, asthma, type 1 and type 2 diabetes, transplantation-related conditions such as graft rejection or graft versus host disease, gout and related inflammatory crystal deposition diseases, or a fibrotic disease, such as atherosclerosis, chronic obstructive pulmonary disease (COPD), cirrhosis, scleroderma, kidney transplant fibrosis, kidney allograft nephropathy, pulmonary fibrosis, including idiopathic pulmonary fibrosis, autoimmune thrombocytopenic purpura, pemphigus vulgaris, acute rheumatic fever, mixed essential cryoglobulinemia, and warm autoimmune hemolytic anemia, among many others.

[0131] BTNL3 antagonists that block or inhibit the BTNL3 pathway can find use in oncology settings. As explained above, BTNL3 antagonists can include, for example, anti-BTNL3 antibodies or variant forms of a BTNL3 protein that interfere with binding of endogenous BTNL3 proteins expressed on a cell membrane and their receptors. As shown below, BTNL3 is relatively highly expressed in certain cancer cells. Example 3; FIG. 4.

[0132] This suggests that the ability of BTNL3 to inhibit activation of T cells might play a role in the ability of cancer cells to evade elimination by the immune system.

[0133] An antibody that binds to either BTNL3 or its receptor and can block or inhibit the interaction between these molecules can be used as a therapeutic to treat cancer. Other antagonists of BTNL3 described above could also be used. Some of the various cancers that might be treated with a BTNL3 pathway blocker include acute or chronic leukemias, lymphoma, non-Hodgkin's lymphoma, Hodgkin's disease, lymphocytic leukemias, lymphocytic or cutaneous lymphomas, carcinomas, adenocarcinomas, sarcomas, thymomas, neoplasms of the mediastinum, breast cancer, prostate cancer, cancers of the head and neck, lung cancer, non-small cell lung cancer, small cell lung cancer, various kinds of skin cancer, cancer of the bladder, malignant

gliomas, cancers of the gastrointestinal system including cancer of the esophagus, stomach, small intestine, colon, or rectum, cancer of the pancreas, adenocarcinomas, hepatobiliary neoplasms, cancer of the kidney or ureter, testicular cancer, cancer of the urethra or penis, gynecologic tumors, ovarian cancer, sarcomas of the bone, cancers of the endocrine system, cutaneous melanoma, intraocular melanoma, neoplasms of the central nervous system, and plasma cell neoplasms, among many other cancers.

[0134] Similarly, BTNL3 antagonists, including anti-BTNL3 antibodies, can be used as a therapeutic to treat infection by a wide variety of pathogens. This may be particularly important where a pathogen can suppress T cell response. Pathogenic infections that can be treated with a BTNL3 antagonist include, without limitation, infections by viruses, bacteria, fungi, and various eukaryotic pathogens including malaria, i.e., members of the species *Plasmodium*, the species *Leishmania*, nematodes, and helminths, e.g., the species *Ascaris*, among many others. In some embodiments, the BTNL3 antagonist can be administered before, concurrently with, or after administration of an antigen that is all or some portion of the pathogen or is similar enough to all or some portion of the pathogen that it can elicit an immune response to the pathogen.

[0135] A BTNL3 antagonist can also find use as an agent to make a vaccine more effective. BTNL3 could be used with a vaccine to induce a response against any antigen. Among these antigens are antigens that are highly expressed on cancer cells, such as cells from the cancers mentioned above. Among these cancer antigens are the following human proteins: WT1, MUC1, LMP2, EGFRvIII, HER-2/neu, MAGE-A3, NY-ESO-1, PSMA, GM2/GD2 synthase, CEA, MLANA/MART1, gp100, survivin, prostate-specific antigen (PSA), telomerase reverse transcriptase (hTERT), sarcoma translocation breakpoints, EPHA2, prostatic acid phosphatase (PAP), melanoma inhibitor of apoptosis (ML-IAP), α -fetoprotein (AFP), epithelial cell adhesion molecule (EpCAM), ERG, NA17.A2 peptide (VLPDVFIRC), paired box 3 (PAX3), anaplastic lymphoma kinase (ALK), androgen receptor, cyclin B1, polysialic acid, rho-related GTP-binding protein RhoC, v-myc myelocytomatosis viral related oncogene (MYCN), TRP-2, GD3 ganglioside, fucosyl GM1, mesothelin, prostate stem cell antigen (PSCA), MAGE-A1, CYP1B1, PLAC1, GM3, BORIS, tetranectin (TN), ETV6-AML1 (especially peptides including the breakpoint), NY-BR-1, RGS5, SART3, STn, carbonic anhydrase IX, PAX5, proacrosin binding protein sp32 precursor (OY-TES-1), sperm protein 17 (Sp17), LCK, high molecular weight melanoma-associated antigen (HMWMAA, also known as melanoma chondroitin sulfate proteoglycan), AKAP-4, SSX2, XAGE-1, B7H3 (also known as CD276), legumain, TIE2, prostate-associated gene 4 protein (PAGE-4), vascular endothelial growth factor receptor 2 (VEGFR2), protamine 2 (also known as MAD-CT-1), glomulin (also known as FAP), PDGFR- β , SSX2, SSX5, Fos-related antigen 1, CD20, integrin α v β 3, 5T4 oncofetal antigen, CA IX, CD5, CD19, CD22 (also known as Siglec-2), CD30 (also known as TNFRSF1), CD33 (also known as Siglec-3), CD40, CD44V6, CD55, CD56 (also known as NCAM), CTLA-4 (also known as CD152), EGFR, GD2, HER2, HLA-DR10 (MHC II), IGF1R, IL-6, sialyl Lewis Y, TAG-72, TAL6, TRAILR2, VEGF, CD52 (also known as CAM-PATH-1), CD4, CD73, CA125 (also known as MUC16), CD66e, CD80 (also known as B7-1), PDGFR β , prostate

specific membrane antigen (PSMA, also known as glutamate carboxypeptidase 2, among many other names). Cancer antigens also include the human herpes virus 4 protein LMP2, the human papillomavirus proteins E6 and E7, and the glycosphingolipid globo H (as described in Gilewski et al. (2001), *Proc. Natl. Acad. Sci.* 98(6): 3270-3275, the portions of which describe globo H are incorporated herein by reference), the α 4 subunit of the α 4 β 1 and α 4 β 7 integrins, the α 4 β 7 integrin, BAFF, APRIL, CD2, CD3, CD20, CD52, CD73, CD80, CD86, the C₅ complement protein, IgE, IL-1 β , IL-5, IL-6R, IL-12, IL-23, and tumor necrosis factor α (TNF α).

[0136] When used to make a vaccine more effective, a BTNL3 antagonist, for example, an anti-BTNL3 antibody, can be administered before, concurrently with, or after an antigen. The antigen can be a part or all of a cancer antigen as described above. An antigen can also be part or all of a pathogen, or a molecule similar to part or all of a pathogen, part or all of a molecule that is expressed specifically on the surface of any cell that mediates a disease, a soluble antigen secreted by cells, or a molecule that can elicit an immune response to a pathogen.

[0137] A molecule that agonizes intracellular signaling through BTNL3, can serve as a treatment for a cancer where the cancer cells express BTNL3. Similarly, an inflammatory disease mediated by immune cells expressing BTNL3, for example neutrophils, can be treated with a molecule that agonizes intracellular signaling through BTNL3. Such an agonist can be an anti-BTNL3 antibody, and the intracellular signaling may be mediated through the B30.2 domain of BTNL3.

[0138] Thus, molecules that can bind to BTNL3 expressed on a neutrophil, thereby downregulating or inhibiting proliferation of the neutrophil, can be useful treatments in conditions where there is an excess of neutrophils or excessive neutrophil activity, for example, in congestive obstructive pulmonary disease (COPD), asthma, inflammatory bowel disease, including ulcerative colitis and Crohn's disease, and gout and related inflammatory crystal deposition diseases. Such molecules can include anti-BTNL3 antibodies.

[0139] On the other hand, molecules that encourage neutrophil proliferation can be useful in patients with neutropenia, who are susceptible to infection. Such patients include patients that have received chemotherapy or radiation for cancer, patients that have been exposed to toxic chemicals or drugs, or patients that have certain autoimmune diseases. Molecules that bind to BTNL3 on the surface of a neutrophil in such a way as to prevent intracellular signals of downregulation or inhibition of proliferation can be useful treatments for such patients. Such molecules can include anti-BTNL3 antibodies. Further, molecules that act as decoys, thereby preventing the interaction of BTNL3 expressed on a neutrophil with a ligand that ordinarily binds to BTNL3 thereby downregulating or inhibiting proliferation of the neutrophil on which BTNL3 is expressed, can be treatments for such conditions. Such molecules can include a BTNL3 protein or variants thereof.

[0140] In addition, an antibody conjugated to a cytotoxic, cytostatic, luminescent, and/or radioactive moiety as described above can be used to treat a cancer in which the cancer cells express BTNL3 or can be used to treat an inflammatory or autoimmune disease characterized by increased numbers of neutrophils, for example, asthma,

COPD, inflammatory bowel disease, including ulcerative colitis and Crohn's disease, and gout and related inflammatory crystal deposition diseases.

[0141] "Treatment" of any disease mentioned herein encompasses an alleviation of at least one symptom of the disease, a reduction in the severity of the disease, or the delay or prevention of disease progression to more serious symptoms that may, in some cases, accompany the disease or lead to at least one other disease. Treatment need not mean that the disease is totally cured. A useful therapeutic agent needs only to reduce the severity of a disease, reduce the severity of one or more symptoms associated with the disease or its treatment, or delay the onset of more serious symptoms or a more serious disease that can occur with some frequency following the treated condition. For example, if the disease is an inflammatory bowel disease, a therapeutic agent may reduce the number of distinct sites of inflammation in the gut, the total extent of the gut affected, reduce pain and/or swelling, reduce symptoms such as diarrhea, constipation, or vomiting, and/or prevent perforation of the gut. A patient's condition can be assessed by standard techniques such as an x-ray performed following a barium enema or enteroclysis, endoscopy, colonoscopy, and/or a biopsy. Suitable procedures vary according to the patient's condition and symptoms.

[0142] A "therapeutically effective amount" of a drug used to treat a disease is an amount that can reduce the severity of a disease, reduce the severity of one or more symptoms associated with the disease or its treatment, or delay the onset of more serious symptoms or a more serious disease that can occur with some frequency following the treated condition.

[0143] The invention encompasses a method of treating inflammatory diseases such as arthritis including rheumatoid arthritis, juvenile idiopathic arthritis, and psoriatic arthritis, Addison's disease, asthma, polyglandular endocrinopathy syndromes, systemic lupus erythematosus, chronic active hepatitis, thyroiditis, lymphocytic adenohypophysitis, premature ovarian failure, idiopathic hypoparathyroidism, pernicious anemia, glomerulonephritis, autoimmune neutropenia, Goodpasture's syndrome, myasthenia gravis, scleroderma, primary Sjogren's syndrome, polymyositis, autoimmune hemolytic anemia, an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, psoriasis, dermatitis, sarcoidosis, multiple sclerosis, chronic obstructive pulmonary disease, asthma, type 1 and type 2 diabetes, transplantation-related conditions such as graft rejection or graft versus host disease, gout and related inflammatory crystal deposition diseases, or a fibrotic disease, such as atherosclerosis, chronic obstructive pulmonary disease (COPD), cirrhosis, scleroderma, kidney transplant fibrosis, kidney allograft nephropathy, pulmonary fibrosis, including idiopathic pulmonary fibrosis, autoimmune thrombocytopenic purpura, pemphigus vulgaris, acute rheumatic fever, mixed essential cryoglobulinemia, and warm autoimmune hemolytic anemia, among many others.

[0144] Such treatment involves using a therapeutically effective amount of a BTNL3 protein, or an agonistic antibody that binds to BTNL3 or its receptor, for a time sufficient to induce a sustained improvement over baseline of an indicator that reflects the severity of a particular disorder or the severity of symptoms caused by the disorder or to delay or prevent the onset of a more serious disease that follows the treated condition in some or all cases. The

treatments of the invention may be used before, after, or during other treatments for the disorder in question that are commonly used, or they may be used without other treatments. For example, Crohn's disease and ulcerative colitis are commonly treated with sulfasalazine, 5-aminosalicylic acid, or corticosteroids. These treatments may be used before, during, or after treatments using a BTNL3 protein or an agonist or antagonist thereof.

[0145] Similarly, cancer is often treated with chemotherapeutic agents and such agents can be used along with the BTNL3 antagonist therapeutics, such as anti-BTNL3 antibodies, described herein. In some embodiments, the BTNL3 antagonist can be administered before, concurrently with, or after a chemotherapeutic agent. Chemotherapeutic agents include, for example, the following therapeutics: alkylating agents (e.g. busulfan, temozolomide, cyclophosphamide, lomustine (CCNU), methyllomustine, streptozotocin, cis-diamminedi-chloroplatinum, aziridinybenzo-quinone, and thiotepea); inorganic ions (e.g. cisplatin and carboplatin); nitrogen mustards (e.g. melphalan hydrochloride, ifosfamide, chlorambucil, and mechlorethamine HCl); nitrosoureas (e.g. carmustine (BCNU)); anti-neoplastic antibiotics (e.g. adriamycin (doxorubicin), daunomycin, mitomycin C, daunorubicin, idarubicin, mithramycin, and bleomycin); plant derivatives (e.g. vincristine, vinblastine, vinorelbine, paclitaxel, docetaxel, vindesine, VP-16, and VM-26); antimetabolites (e.g. methotrexate with or without leucovorin, 5-fluorouracil with or without leucovorin, 5-fluorodeoxyuridine, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycytosine, gemcitabine, and fludarabine); podophyllotoxins (e.g. etoposide, irinotecan, and topotecan); as well as actinomycin D, dacarbazine (DTIC), mAMSA, procarbazine, hexamethylmelamine, pentamethylmelamine, L-asparaginase, and mitoxantrone, among many known in the art. See e.g. *Cancer: Principles and Practice of Oncology*, 4th Edition,

[0146] DeVita et al., eds., J. B. Lippincott Co., Philadelphia, Pa. (1993), the relevant portions of which are incorporated herein by reference.

[0147] Cancer is also commonly treated by a variety of methods with a variety of forms of radiation. It is contemplated herein that the administration of a BTNL3 antagonist to a cancer patient can occur before, concurrently with, or after radiation treatment or any other anti-neoplastic treatment.

[0148] For autoimmune or inflammatory conditions, T cells can be removed from a patient, for example, through apheresis, and stimulated ex vivo using BTNL3, optionally plus other proteins, such that the T cells attain a regulatory or inhibitory phenotype. The T cells can then be transferred back into the patient. To stimulate the T cells to attain a regulatory or inhibitory phenotype, they can be incubated in the presence of a surface, for example, with beads or in microtiter plate well, that is coated with human T cell agonistic anti-CD3 antibody, rBTNL3.Fc, and rB7-1.Fc or rB7-2.Fc in the presence of TGF-beta and IL-2. Alternatively, the surface could be coated with a combination of proteins that includes rBTNL3 or BTNL3.Fc plus an agonistic anti-CD3 antibody or a combination that includes only these proteins. In one embodiment, the agonistic anti-CD3 antibody, rBTNL3.Fc, and rB7-1.Fc or rB7-2.Fc can be, for example, at a molecular weight ratio of 2:10:2.5. The TGF-beta and IL-2 can be at appropriate concentrations, such as, for example, from about 0.05 to 5 ng/ml for

TGF-beta and from about 0.5 to 10 ng/ml for IL-2. This can program the T cells to become inhibitory or regulatory, resulting in an "expansion" of regulatory T ("T reg") cells. "Expansion" of T reg cells means that the ratio of T reg cells (CD3⁺ FOXP3⁺) to T cells as a whole (CD3⁺ FOXP3⁻) becomes greater. This ratio can be determined by FACS analysis using antibodies to detect cell proteins, a method well known in the art. See, e.g., Swanson et al. (2013), J. Immunol. 190: 2027-2035, the relevant portions of which are incorporated herein by reference. The T cells can be incubated in such a setting for, e.g., three to seven days and then harvested and delivered back to the same patient. Optionally, the T cells can be incubated about three, four, five, six, or seven days. In some embodiments, the T cells can also be rested, i.e., cultured in the presence of, for example, IL-2, without T cell receptor or costimulatory stimulus, and then restimulated as explained above one to four more times. The autoimmune or inflammatory conditions treatable with such a regime include, for example, arthritis including rheumatoid arthritis, juvenile idiopathic arthritis, and psoriatic arthritis, Addison's disease, asthma, polyglandular endocrinopathy syndromes, systemic lupus erythematosus, chronic active hepatitis, thyroiditis, lymphocytic adenohypophysitis, premature ovarian failure, idiopathic hypoparathyroidism, pernicious anemia, glomerulonephritis, autoimmune neutropenia, Goodpasture's syndrome, myasthenia gravis, scleroderma, primary Sjogren's syndrome, polymyositis, autoimmune hemolytic anemia, an inflammatory bowel disease, such as Crohn's disease or ulcerative colitis, psoriasis, dermatitis, sarcoidosis, multiple sclerosis, chronic obstructive pulmonary disease, asthma, type 1 and type 2 diabetes, transplantation-related conditions such as graft rejection or graft versus host disease, gout and related inflammatory crystal deposition diseases, or a fibrotic disease, such as atherosclerosis, chronic obstructive pulmonary disease (COPD), cirrhosis, scleroderma, kidney transplant fibrosis, kidney allograft nephropathy, pulmonary fibrosis, including idiopathic pulmonary fibrosis, autoimmune thrombocytopenic purpura, pemphigus vulgaris, acute rheumatic fever, mixed essential cryoglobulinemia, and warm autoimmune hemolytic anemia, among many others.

[0149] Any of the above-described therapeutic agents can be administered in the form of a composition, that is, with one or more additional components such as a physiologically acceptable carrier, excipient, or diluent. For example, a composition may comprise a soluble BTNL3 protein, an anti-BTNL3 antibody, or a BTNL3 agonist or antagonist as described herein plus a buffer, an antioxidant such as ascorbic acid, a low molecular weight polypeptide (such as those having less than 10 amino acids), a protein, amino acids, carbohydrates such as glucose, sucrose, or dextrans, chelating agent such as EDTA, glutathione, and/or other stabilizers, excipients, and/or preservatives. The composition may be formulated as a liquid or a lyophilizate. Further examples of components that may be employed in pharmaceutical formulations are presented in Remington's Pharmaceutical Sciences, 16th Ed., Mack Publishing Company, Easton, Pa., (1980), the relevant portions of which are incorporated herein by reference.

[0150] Compositions comprising therapeutic molecules described above can be administered by any appropriate means including, but not limited to, parenteral, topical, oral, nasal, vaginal, rectal, or pulmonary (by inhalation) admin-

istration. If injected, the composition(s) can be administered intra-articularly, intravenously, intraarterially, intramuscularly, intraperitoneally, or subcutaneously by bolus injection or continuous infusion. Localized administration, that is, at the site of disease, is contemplated, as are transdermal delivery and sustained release from implants, skin patches, or suppositories. Delivery by inhalation includes, for example, nasal or oral inhalation, use of a nebulizer, inhalation in aerosol form, and the like. Administration via a suppository inserted into a body cavity can be accomplished, for example, by inserting a solid form of the composition in a chosen body cavity and allowing it to dissolve. Other alternatives include eye drops, oral preparations such as pills, lozenges, syrups, and chewing gum, and topical preparations such as lotions, gels, sprays, and ointments. In most cases, therapeutic molecules that are polypeptides can be administered topically or by injection or inhalation.

[0151] The therapeutic molecules described above can be administered at any dosage, frequency, and duration that can be effective to treat the condition being treated. The dosage depends on the molecular nature of the therapeutic molecule and the nature of the disorder being treated. Treatment may be continued as long as necessary to achieve the desired results. The periodicity of treatment may or may not be constant throughout the duration of the treatment. For example, treatment may initially occur at weekly intervals and later occur every other week. Treatments having durations of days, weeks, months, or years are encompassed by the invention. Treatment may be discontinued and then restarted. Maintenance doses may be administered after an initial treatment.

[0152] Dosage may be measured as milligrams per kilogram of body weight (mg/kg) or as milligrams per square meter of skin surface (mg/m²) or as a fixed dose, irrespective of height or weight. These are standard dosage units in the art. A person's skin surface area is calculated from her height and weight using a standard formula. For example, a therapeutic BTNL3 protein or an antibody that binds to BTNL3 or its receptor can be administered at a dose of from about 0.05 mg/kg to about 10 mg/kg or from about 0.1 mg/kg to about 1.0 mg/kg. Alternatively, a dose of from about 1 mg to about 500 mg can be administered. Or a dose of about 5 mg, 10 mg, 15 mg, 20 mg, 25 mg, 30 mg, 35 mg, 40 mg, 45 mg, 50 mg, 55 mg, 60 mg, 100 mg, 200 mg, or 300 mg can be administered.

[0153] The invention is described below with reference to specific examples. These examples are not meant to limit the invention in any way. It is understood for purposes of this disclosure, that various changes and modifications may be made to the invention that are well within the scope of the invention. Numerous other changes may be made which will readily suggest themselves to those skilled in the art and which are encompassed in the spirit of the invention disclosed herein and as defined in the appended claims.

EXAMPLES

Example 1: Expression of mRNA Encoding Human BTNL3 Protein in Human Immune Cells

[0154] The following experiments were done in order to gather information on expression of mRNA encoding human BTNL3 in primary human immune cells.

[0155] Primary immune cells were isolated from whole blood or leukopaks via various commercially available

selection methods from Stem Cell Sciences (Palo Alto, California) or Miltenyi Biotech (Germany). For example, the EASYSEP® Human T cell enrichment kit and/or the CD4⁺ T cell enrichment kit (both from Stem Cell Sciences), was used to isolate CD4⁺ T cells, while monocytes were isolated using the Miltenyi Monocyte Isolation Kit II. Such cell separations using such commercially available reagents are routine in the art. Macrophages were obtained through the ex vivo maturation of negatively-selected monocytes for seven days. Each isolated cell population was analyzed by fluorescence activated cell sorting (FACS) to determine whether the isolated cell population was expressing the expected cell surface proteins. RNA was isolated and assessed by Affymetrix array (Affymetrix GENECHIP™ HG-U1333 Plus 2.0). Data normalization and analysis for human BTNL3 transcript detection was performed using ROSETTA RESOLVER® software (Rosetta Biosoftware, Cambridge, Mass., USA).

[0156] The results of these analyses are shown in FIG. 2. The cells type tested included peripheral blood mononuclear cells, T cells, CD4⁺ T cells, CD8⁺ T cells, B cells, monocytes, macrophages, neutrophils, eosinophils, and NK cells. Among cell types tested, neutrophils (lane 8 in FIG. 2) expressed the highest amounts of BTNL3.

Example 2: Preparation of Human BTNL3.Fc

[0157] The following describes how a fusion protein containing the extracellular region of human BTNL3, a linker, and the Fc portion of a human IgG1 antibody was made. A cDNA in an appropriate vector was constructed encoding the extracellular domain of human BTNL3, that is amino acids 1 to 236 of SEQ ID NO:2, fused to a linker plus a human IgG1 Fc fragment. SEQ ID NO:5 provides the sequence of this cDNA, and SEQ ID NO:6 provides the amino acid sequence of the BTNL3.Fc protein encoded by this cDNA, including the signal sequence. The mature BTNL3.Fc protein sequence, that is, lacking the signal sequence, is provided in SEQ ID NO:7. Cos PKB cells were transfected with the BTNL3.Fc mammalian expression construct using LIPOFECTAMINE™ 2000 (Invitrogen) and cultured in complete Dulbecco's Modified Eagle Medium (DMEM) with 0.5% Low Ig Serum. These methods are described in detail by Ettahdieh et al., OVEREXPRESSION OF PROTEIN KINASE B α ENHANCES RECOMBINANT PROTEIN EXPRESSION IN TRANSIENT SYSTEMS in *Animal Cell Technology: From Target to Market: Proceedings of the 17th ESACT Meeting*, Tylosand, Sweden, Jun. 10-14, 2001, Vol. 1, Lindner-Olsson et al., eds., pp. 31-35, Springer, 2001. The portions of this reference describing how to make a recombinant protein are incorporated herein by reference. Seven days post transfection, supernatants were harvested, and the BTNL3.Fc protein was purified by Protein A column chromatography (MABSE-LECT™ SuRe column, GE Healthcare).

[0158] The size of the resulting protein was determined by gel electrophoresis and size exclusion chromatography (SEC). It was expected that the protein would be dimeric because of the presence of the Fc region, which causes dimerization due to noncovalent interactions between C_H3 regions and disulfide bonds formed between hinge regions. If the protein were a dimer, its expected molecular weight would likely be in excess of about 120 kD due to variable glycosylation. Similarly, its expected molecular weight under reducing conditions, in which it would be expected to be monomeric, would be at least about 60 kD. On a reducing

gel, a number of bands were observed, most of which were between about 50 and 64 kD. The observed molecular weight of the main peak by SEC under non-reducing conditions was about 140 kD, although both higher and lower molecular weight species were also observed. These data indicate that the protein is at least dimeric under non-reducing conditions and that it likely also forms higher order multimers.

Example 3: Expression of BTNL3 in Human Normal and Tumor Tissue

[0159] To determine the level of expression of BTNL3 at the RNA level in human normal and tumor tissue, RNA was isolated and assessed by next-generation sequencing analysis. In particular, the TruSeq® (Illumina, Inc., San Diego, Calif.) reagents were used to prepare samples and reactions that were processed in a HiSeq® 2000 (Illumina, Inc., San Diego, Calif.) sequencing system. Data was analyzed to quantify the number of BTNL3 RNAs per cell. The results of this analysis for normal human tissues are shown in FIG. 3. Little or no BTNL3 expression was detected in most tissues, but higher levels of expression were detected in samples from colon and small intestine. Thus, BTNL3, like BTNL2, is expressed primarily in the gut.

[0160] Similar methods were used to determine levels of BTNL3 RNA expression in various human tumor tissue samples. These results are shown in FIG. 4. Moderate levels of BTNL3 expression were detected in several different adenocarcinoma samples, but not in all tumor tissues. In addition, moderate and high levels of BTNL3 expression were detected in some samples from type II diabetes patients, suggesting that BTNL3 could play a role in mediating or responding to this disease in some cases.

Example 4: In Vitro Analysis of Marine CD4⁺ T Cell Proliferation

[0161] The following experiment was done to determine the effects of a human BTNL3.Fc fusion protein on the proliferation of mouse CD4⁺ T cells in vitro.

[0162] A single cell splenocyte suspension, which was prepared from spleens harvested from at least five female C57BL/6 mice per experiment, was used to purify CD4⁺ T cells with the mouse EASYSEP™ CD4⁺ negative selection kit (Stem Cell Sciences). Purity of CD4⁺ T cells was greater than 90% as assessed by FACS analysis. Tissue culture-treated microtiter plates were coated with variable concentrations of an anti-mouse CD3 monoclonal antibody (Clone 2C11, BD Biosciences Pharmingen, San Diego, Calif., USA) and the Fc fusion protein indicated in the brief description of FIG. 5. Each well was equalized for total protein content using human IgG (Sigma). All coatings/dilutions were done in PBS. Plates were incubated overnight. Wells were then washed with PBS, and then 1-2×10⁵ purified CD4⁺ splenocytes/well were added. Proliferation of the CD4⁺ T cells was determined by incorporation of ³H-thymidine (1 μ Ci/well) during the last 6 hours of the 72 hour culture. Human IgG was used as a negative control. As positive controls, mouse BTNL2.Fc, which had been previously shown to inhibit proliferation of T cells, and human B7-2-Fc (R & D Biosystems), a known positive costimulator of T cells, were also included.

[0163] The results are shown in FIG. 5. Lane 1 in FIG. 5 represents a negative control assay containing no immobi-

lized protein and no additional protein. Lanes 2 and 3 show results from positive control assays containing immobilized anti-mouse CD3 antibody plus either human IgG (lane 2) or a human B7-2-Fc protein. Lane 4 shows results from an assay containing immobilized anti-mouse CD3 antibody plus mouse BTNL2.Fc, a negative costimulatory molecule. Lane 5 shows results of an assay containing immobilized anti-mouse CD3 antibody plus human BTNL3.Fc. Lanes 6 and 7 show results from an assay using immobilized human IgG plus either mouse BTNL2.Fc (lane 6) or human BTNL3.Fc (lane 7). These data confirm the stimulatory effect of human B7-2-Fc and the inhibitory effect of mouse BTNL2.Fc on mouse T cell proliferation and indicate that human BTNL3.Fc can inhibit mouse T cell proliferation. These data suggest that a human BTNL3.Fc fusion protein can have effects on murine T cells similar to those observed using a murine BNTL2.Fc fusion protein. Furthermore, these data suggest that human BTNL3 can interact with a receptor on murine T cells, despite the fact that a close murine homolog of human BTNL3 has not been isolated

Example 5: Effects of BTNL3.Fc on Gene Expression of Mouse CD4+ T Cells

[0164] In a standard anti-CD3 proliferation assay, mouse CD4+ T cells were isolated as described above and stimulated with an immobilized anti-CD3 antibody plus no additional protein, immobilized mouse BTNL2.Fc, or immobilized human BTNL3.Fc. Proliferation was measured by ³H-thymidine incorporation. After 24 hours of stimulation, RNA was isolated from the cells and analyzed by Affymetrix cDNA chips to determine levels of expression of a large variety of sequences. Results are shown in FIG. 6 and Table 4 below. Stimulation of murine CD4+ cells with an immobilized anti-CD3 antibody upregulated the expression many immunoregulatory and inflammatory genes as expected. Inclusion of murine BTNL2.Fc or human BTNL3.Fc dampened this response.

[0165] In FIG. 6, a black dot represents the relative amount of expression of a particular sequence observed in stimulated as compared to unstimulated cells. Sequences with no significant difference in expression between stimulated and un-stimulated cells are not represented in FIG. 6.

In FIG. 6, the left panel shows data from cells stimulated with an anti-CD3 antibody alone, the middle panel shows data from cells stimulated with an anti-CD3 antibody plus a murine BTNL2.Fc fusion protein, and the right panel shows data from cells stimulated with an anti-CD3 antibody plus a human BTNL3.Fc fusion protein. These data show that the expression of many genes is significantly up- or down-regulated by 24 hours of stimulation with anti-CD3 antibody. However, addition of either murine BTNL2.Fc or human BTNL3.Fc substantially reduces this number of genes. These data confirm that BTNL2.Fc is a negative costimulator and indicate that BTNL3.Fc is also a negative costimulator. Thus, these results indicate that both molecules inhibit the activation of murine CD4+ T cells by anti-CD3 antibody.

[0166] Table 4 below provides additional numeric data from this experiment. Out of 45,060 sequences assayed, expression of many (26,530) sequences was not changed in a statistically significant manner by stimulation with an anti-CD3 antibody alone. However, expression of 8,631 and 9,899 sequences was significantly up- or down-regulated, respectively, upon stimulation with anti-CD3 antibody alone. These numbers were substantially decreased when either murine BTNL2.Fc or human BTNL3.Fc was added. Hence, both BTNL2 and BTNL3 inhibit the activation of T cells.

TABLE 4

Effects of BTNL3 on the Gene Expression Signature of anti-CD3 Antibody				
Growth Conditions	# Data Points			Total # Data Points
	Unchanged	Up-Regulated (p ≤ 0.01)	Down-Regulated (p ≤ 0.01)	
Anti-CD3 24 h/No Stim 24 h	26,530	8,631	9,899	45,060
Anti-CD3 + rBTNL2 Fc 24 h/No Stim 24 h	40,752	2,056	2,252	45,060
Anti-CD3 + rBTNL3 Fc 24 h/No Stim 24 h	39,161	2,831	3,068	45,060

SEQUENCE LISTING

Sequence total quantity: 23
SEQ ID NO: 1 moltype = DNA length = 1401
FEATURE Location/Qualifiers
source 1..1401
mol_type = other DNA
organism = Homo sapiens
CDS 1..1401
SEQUENCE: 1
atgggtttttg tgcctatctt gggtctcagt ttctacgagc tgggtgcagg acagtggcaa 60
gtcactggac cgggcaagtt tgtccaggcc ttggtggggg aggacgccgt gttctcctgc 120
tccctctttc ctgagaccag tgcagaggct atggaagtgc ggttcttcag gaatcagttc 180
catgctgtgg tccactctca cagagatggg gaagactggg aatctaagca gatgccacag 240
tatacaggga gaactgagtt tgtgaaggac tccattgcag gggggcgtgt ctctctaagg 300
ctaaaaaaca tcaactccctc ggacatcgcc ctgtatgggt getgggtcag ttcccagatt 360
tacgatgagg agggccactg ggagctgcgg gtggcagcac tgggtcactc tcctctcatt 420
tccatcgtgg gatattgtga cggagggtatc cagtactctc gctgtcctc aggetgggtc 480
ccccagccca cagccaagtg gaaagggtcca caaggacagg atttgtcttc agactccaga 540
gcaaatgcag atgggtacag cctgtatgat gtggagatct ccattatagt ccaggaaaat 600
gctgggagca tattgtgttc catccacctt gctgagcaga gtcattgaggt ggaatccaag 660
gtattgatag gagagacgtt ttccagccc tcaccttggc gctggcttc tattttactc 720
gggttactct gtgggtccct gtgtggtgtt gtcattggga tgataattgt tttcttcaaa 780

-continued

```

tccaaagggg aaatccaggc ggaactggac tggagaagaa agcacggaca ggcagaattg 840
agagacgccc ggaaacacgc agtggagggt actctggatc cagagacggc tcacccgaag 900
ctctgcgttt ctgatctgaa aactgtaacc catagaaaaa ctccccagga ggtgcctcac 960
tctgagaaga gatttacaag gaagagtgtg gtggcttctc agggtttcca agcagggaaa 1020
cattactggg aggtggagct gggacaaaat gtagggtggt atgtgggagt gtgtcgggat 1080
gacgtagaca gggggaagaa caatgtgact ttgtctccca acaatgggta ttgggtcctc 1140
agactgacaa cagaacattt gtatttcaca ttcaatcccc attttatcag cctccccccc 1200
agcacccttc ctacacgagt aggggtcttc ctggactatg aggggtgggac catctccttc 1260
ttcaatacaa atgaccagtc ccttatttat accctgctga catgtcagtt tgaaggcttg 1320
ttgagaccct atatccagca tgcgatgtat gacgaggaaa aggggactcc catattcata 1380
tgtccagctg cctgggggatg a 1401

```

```

SEQ ID NO: 2      moltype = AA length = 466
FEATURE          Location/Qualifiers
source           1..466
                 mol_type = protein
                 organism = Homo sapiens

```

```

SEQUENCE: 2
MAFVLILVLS FYELVSGQWQ VTGPGKFVQA LVGEDAVFSC SLFPETSAEA MEVRFFRNQF 60
HAVVHLYRDG EDWESKQMPQ YRGRTEFVKD SIAGGRVSLR LKNITPSDIG LYGCWFSSQI 120
YDEEATWELR VAALGSLPLI SIVGYVDGGI QLLCLSSGWF PQPTAKWKP QGQDLSSDSR 180
ANADGYSLYD VEISIIQEN AGSILCSIHL AEQSHEVESK VLIGETFPQP SPWRLASILL 240
GLLCGALCGV VMGMIIFFFK SKGKIQAELD WRRKHQAEL RDARKHAVEV TLPETAHPK 300
LCVSLDKTVT HRKAPQEVPH SEKRFRKSV VASQGFQAGK HYWEVDVQON VGWYVGVCRD 360
DVDRGKNNVT LSPNNGYWVL RLTEHLYPT FNPFIISLP STPPTRVGVF LDYEGGTISF 420
FNTNDQSLIY TLLTCQPEGL LRPYIQHAMY DEEKGTPIFI CPVSWG 466

```

```

SEQ ID NO: 3      moltype = AA length = 20
FEATURE          Location/Qualifiers
source           1..20
                 mol_type = protein
                 organism = Homo sapiens

```

```

SEQUENCE: 3
METDTLLLVW LLLWVPGSTG 20

```

```

SEQ ID NO: 4      moltype = AA length = 26
FEATURE          Location/Qualifiers
source           1..26
                 mol_type = protein
                 organism = Homo sapiens

```

```

SEQUENCE: 4
MATGSRTSL LAFGLLCLPW LQEGSA 26

```

```

SEQ ID NO: 5      moltype = DNA length = 1437
FEATURE          Location/Qualifiers
misc_feature     1..1437
                 note = Fusion Protein
source           1..1437
                 mol_type = other DNA
                 organism = synthetic construct
CDS              1..1437

```

```

SEQUENCE: 5
atggcttttg tgetcatttt ggttctcagt ttctacgagc tgggtgcagg acagtggcaa 60
gtcactggac cgggcaagtt tgtccaggcc ttggtggggg aggacgccgt gttctcctgc 120
tccctctttc ctgagaccag tgcagaggct atggaagtgc ggttcttcag gaatcagttc 180
catgtctgtg tccaccteta cagagatggg gaagactggg aatctaagca gatgccacag 240
tatcgaggga gaactgagtt tgtgaaggac tccattgcag gggggcgctg ctctctaagg 300
ctaaaaaaca tcaactccctc ggacatcggc ctgtatgggt gctgggtcag ttcccagatt 360
tacgatgagg agggccacctg ggagctgcgg gtggcagcac tgggctcact tcctctcatt 420
tccatcgtag gatattgtga cggaggtatc cagttactct gcctgtcctc aggtcgggtc 480
ccccagccca cagccaagtg gaaaggtcca caaggacagg atttgtcttc agactccaga 540
gcaaatgcag atgggtacag cctgtatgat gtggagatct ccattatagt ccaggaaaat 600
gctgggagca tatgtgttc catccacctt gctgagcaga gtcatgaggt ggaatccaag 660
gtattgatag gagagacggt ttccagccc tcaccttggc gcctggctgg aggtggaggc 720
tccggaggtg gaggttcogg ttgaggtgga tccgacaaaa ctacacatg tccaccgtgc 780
ccagcacctg aactcctcggg gggaccgtca gtttctctc tcccccaaa acccaaggac 840
accctcatga tctcccgagc ccttgaggtc acatgcgttg tgggtggacgt gagccacgaa 900
gacctgagg tcaagttcaa ctggtacgtg gacggcgctg aggtgcataa tgccaagaca 960
aagccgcggg aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg 1020
caccaggact ggctgaatgg caaggagtac aagtgcagg tctccaacaa agccctccca 1080
gcccccatcg agaaaaacat ctccaaagcc aaagggcagc ccgagaacc acaggtgtac 1140
accctgcccc cataccggga tgagctgacc aagaaccagg tcagcctgac ctgcctgggc 1200
aaaggcttct atccacgca catcgccgtg gagggggaga gcaatgggca gccggagaa 1260
aactacaaga ccacgctcct cgtgctggac tccgacggct ccttcttctc ctacagcaag 1320
ctcacctggg acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat 1380
gaggctctgc acaaccacta caccagaaag agcctctccc tgtctccggg taaatga 1437

```

-continued

SEQ ID NO: 6 moltype = AA length = 478
 FEATURE Location/Qualifiers
 REGION 1..478
 note = Synthetic Construct
 source 1..478
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 6

MAFVLILVLS	FYELVSGQWQ	VTGPGKFVQA	LVGEDAVFSC	SLFPETSAEA	MEVRFFRNQF	60
HAVVHLYRDG	EDWESKQMPQ	YRGRTEFVKD	SIAGGRVSLR	LKNITPSDIG	LYGCWFSSQI	120
YDEEATWELR	VAALGSLPLI	SIVGYVDGGI	QLLCLSSGWF	PQPTAKWKGP	QGQDLSSDSR	180
ANADGYSLYD	VEISIIQVEN	AGSILCSIHL	AEQSHEVESK	VLIGETFFQP	SPWRLAGGGG	240
SGGGGSGGGG	SDKTHTCPPC	PAPELLGGPS	VFLFPPKPKD	TLMISRTPEV	TCVVVDVSHE	300
DPEVKFNWYV	DGVEVHNAKT	KPREEQYNST	YRVVSVLTVL	HQDWLNGKEY	KCKVSNKALP	360
APIEKTISKA	KGPQREPQVY	TLPPSRDELT	KNQVSLTCLV	KGFYPSDIAV	EWESNGQPEN	420
NYKTTTPVLD	SDGSFFLYSK	LTVDKSRWQQ	GNVFSCSV MH	EALHNHYTQK	SLSLSPGK	478

SEQ ID NO: 7 moltype = AA length = 461
 FEATURE Location/Qualifiers
 REGION 1..461
 note = Fusion Protein
 source 1..461
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 7

QWQVTGPGKF	VQALVGEDAV	FSCSLFPETS	AEAMEVRFFR	NQFHAVVHLY	RDGEDWESKQ	60
MPQYRGRTEF	VKDSIAGGRV	SLRLKNITPS	DIGLYGCWFS	SQIYDEEATW	ELRVAALGSL	120
PLISIVGYVD	GGIQLLCLSS	GWFPQPTAKW	KGPQGQDLSS	DSRANADGYS	LYDVEISIIV	180
QENAGSILCS	IHLAEQSHEV	ESKVLIGETF	FQPSPWRLAG	GGGSGGGGSG	GGGSDKTHTC	240
PENAPPELLG	GPSVFLPPPK	PKDTLMISRT	PEVTCVVVDV	SHEDPEVKFN	WYVDGVEVHN	300
AKTKPREEQY	NSTYRVVSVL	TVLHQDWLNG	KEYKCKVSNK	ALPAPIEKTI	SKAKGQPREP	360
QVYTLPPSRD	ELTKNQVSLT	CLVKGFYPSD	IAVEWESNGQ	PENNYKTPP	VLDSDGSFFL	420
YSKLTVDKSR	WQQGNVFS	CS VMHEALHNHY	TQKSLSLSPG	K		461

SEQ ID NO: 8 moltype = DNA length = 1299
 FEATURE Location/Qualifiers
 source 1..1299
 mol_type = other DNA
 organism = Homo sapiens
 CDS 1..1299

SEQUENCE: 8

atggcttttg	tgtctcatttt	ggttctcagt	ttctacgagc	tgggtgcagg	acagtgccaa	60
gtcactggac	cgggcaagtt	tgtccaggcc	ttgggtgggg	aggacgccgt	gttctcctgc	120
tccctctttc	ctgagaccag	tgcagaggct	atggaagtgc	ggttcttcag	gaatcagttc	180
catgctgtgg	tccacctcta	cagagatggg	gaagactggg	aatctaagca	gatgccacag	240
tatcgaggga	gaactgagtt	tgtgaaggac	tccattgcag	gggggcgtgt	ctctctaagg	300
ctaaaaaaca	tcactccctc	ggacatcgcc	ctgtatgggt	gctgggtcag	ttcccagatt	360
tacgatgagg	aggccacctg	ggagctgcgg	gtggcagcac	tgggtcact	tcctctcatt	420
tccatcgctg	gatatgttga	cggaggtatc	cagttactct	gcctgtcctc	attccagccc	480
tcaccttgcc	gcctggcttc	tattttactc	gggttactct	gtgtgtccct	gtgtgtgtgt	540
gtcatgggga	tgataattgt	ttcttcaaaa	tccaaaggga	aaatccaggc	ggaactgggt	600
atgtgtcatg	tcctgagcct	cccacacatg	gttctcccg	gtccctccct	gateccacagt	660
ttgagcctct	ggacgaccct	ggctgcaggc	tggacaggaa	gcaccgactg	gagaagaaa	720
cacggacagg	cagaattgag	agacgcccgg	aaacacgcag	tggaggtgac	tctggatcca	780
gagacggctc	accggaagct	ctgcgtttct	gatctgaaaa	ctgtaaccca	tagaaaagct	840
ccccaggagg	tgcctcactc	tgagaagaga	tttacaagga	agagtgtggt	ggcttctcag	900
gggtttccaag	cagggaaaaca	ttactgggag	gtggacgtgg	gacaaaatgt	agggtgggtat	960
gtgggagtg	gtcgggatga	cgtagacagg	gggaagaaca	atgtgacttt	gtctcccaac	1020
aatgggtatt	gggtccctcag	actgacaaca	gaacatttgt	atttcacatt	caatccccat	1080
tttatcagcc	tccccccag	caccctcctc	acacgagtag	gggtcttcct	ggactatgag	1140
gggtgggacca	tctccttctt	caatacaaat	gaccagtcct	ttattttatac	cctgctgaca	1200
tgtagctttg	aaggcttggt	gagaccctat	atccagcatg	cgatgtatga	cgaggaaaag	1260
gggactccca	tattcatatg	tccagtgtcc	tggggatga			1299

SEQ ID NO: 9 moltype = AA length = 432
 FEATURE Location/Qualifiers
 source 1..432
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 9

MAFVLILVLS	FYELVSGQWQ	VTGPGKFVQA	LVGEDAVFSC	SLFPETSAEA	MEVRFFRNQF	60
HAVVHLYRDG	EDWESKQMPQ	YRGRTEFVKD	SIAGGRVSLR	LKNITPSDIG	LYGCWFSSQI	120
YDEEATWELR	VAALGSLPLI	SIVGYVDGGI	QLLCLSSFQ	SPWRLASILL	GLLCGALCGV	180
VMGMIIVFFK	SGKIQAELG	MCHVLSLPHM	VLPGPSLIHS	LSLWTTLAAG	WTGSTDWRRK	240
HQAELRDAR	KHAVEVTLDP	ETAHPKLCVS	DLKTVTHRKA	PQEVPHSEKR	FTRKSVVASQ	300

-continued

GFQAGKHYWE	VDVGQNVGWY	VGVCRDDVDR	GKNNVTLSFN	NGYWVLRLLT	EHLYFTFNPH	360
FISLPPSTPP	TRVGVFLDYE	GGTISFFNTN	DQSLIYTLLT	CQFEGLLRPY	IQHAMYDEEK	420
GTPIFICPVS	WG					432
SEQ ID NO: 10	moltype = AA length = 5					
FEATURE	Location/Qualifiers					
REGION	1..5					
	note = Linker					
source	1..5					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 10						
GGGGS						5
SEQ ID NO: 11	moltype = AA length = 6					
FEATURE	Location/Qualifiers					
REGION	1..6					
	note = Linker					
source	1..6					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 11						
GGGGSN						6
SEQ ID NO: 12	moltype = AA length = 12					
FEATURE	Location/Qualifiers					
REGION	1..12					
	note = Linker					
source	1..12					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 12						
GKSSGSGSES KS						12
SEQ ID NO: 13	moltype = AA length = 14					
FEATURE	Location/Qualifiers					
REGION	1..14					
	note = Linker					
source	1..14					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 13						
GSTSGSGKSS EGKG						14
SEQ ID NO: 14	moltype = AA length = 18					
FEATURE	Location/Qualifiers					
REGION	1..18					
	note = Linker					
source	1..18					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 14						
GSTSGSGKSS EGGSTKG						18
SEQ ID NO: 15	moltype = AA length = 14					
FEATURE	Location/Qualifiers					
REGION	1..14					
	note = Linker					
source	1..14					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 15						
GSTSGSGKSS EGKG						14
SEQ ID NO: 16	moltype = AA length = 18					
FEATURE	Location/Qualifiers					
REGION	1..18					
	note = Linker					
source	1..18					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 16						
GSTSGSGKPG SGGSTKG						18
SEQ ID NO: 17	moltype = AA length = 14					
FEATURE	Location/Qualifiers					

-continued

REGION	1..14	
	note = Linker	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 17		
EGKSSGSGSE SKEF		14
SEQ ID NO: 18	moltype = AA length = 15	
FEATURE	Location/Qualifiers	
REGION	1..15	
	note = Linker	
source	1..15	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 18		
GGGSGGGGS GGGGS		15
SEQ ID NO: 19	moltype = AA length = 500	
FEATURE	Location/Qualifiers	
source	1..500	
	mol_type = protein	
	organism = Homo sapiens	
SEQUENCE: 19		
MALMLSLVLS LLKLGSGQWQ VFGPDKPVQA LVGEDAAFSC FLSPKTNAEA MEVRFRRGQF		60
SSVHLYRDG KDQPFMQMPQ YQGRTKLVKD SIAEGRISLR LENITVLDAG LYGCRISQS		120
YYQKAIWELQ VSALGSVPLI SITGYVDRDI QLLCQSSGWF PRPTAKWKGP QGQDLSTDSR		180
TNRDMHGLFD VEISLTQEN AGSISCSMRH AHLREVESR VQIGDTFFEP ISWHLATKVL		240
GILCCGLFFG IVGLKIFFSK FQWKIQAELE WRRKHGQAELE RDARKHAVEV TLDPETAHPK		300
LCVSDLKTVT HRKAPQEVPH SEKRFRTRKSV VASQSFQAGK HYWEVDGGHN KRWRVGVCRD		360
DVDRRKEYVT LSPDHGYWVL RLNGEHLFT LNPRFISVFP RTPPTKIGVF LDYECGTISF		420
FNINDQSLIY TLTCTFEGLL RPYIEYPSYN EQNGTPIVIC PVTQESEKEA SWQRASAIPE		480
TSNSESQA TTPFLPRGEM		500
SEQ ID NO: 20	moltype = AA length = 535	
FEATURE	Location/Qualifiers	
source	1..535	
	mol_type = protein	
	organism = Homo sapiens	
SEQUENCE: 20		
MVDLSVSPDS LKPVSLTSSL VFLMHLLLQ PGEPSSEVKV LGPEYPILAL VGEEVEFPCH		60
LWPQLDAQQM EIRWFRSQTF NVVHLYQEQQ ELPGRQMPAF RNRTKLVKDD IAYGSVVLQL		120
HSIIPSDKGT YGCRFHSDFN SGEALWELEV AGLGSDPHLS LEGFKEGGIQ LRLRSSGWYP		180
KPKVQWRDHQ GQCLPPEFEA IVWDAQDLFS LETSVVVRAG ALSNVSVSIQ NLLLSQKKEL		240
VVQIADVFPV GASAWKSAFV ATPLPLLVLA ALALGVLRKQ RRSREKLKQ AEKRQEKLT		300
ELEKLQTELD WRAEGQAEW RAAQKYAVDV TLDPASAHPS LEVSEDGKSV SSRGAPPGPA		360
PGHPQRFSEQ TCALSLEFRS AGRHYWEVHV GRRSRWFLGA CLAAVPRAGP ARLSPAAGYW		420
VLGLWNGCEY FVLAPHRVAL TLRVPPRRLG VFLDYEAGEL SFFNVSDGSH IFTFHDFTSG		480
ALCAYFRPRA HDGGEHPDPL TICPLPVRGT GVPEENDSDT WLQPYEPADP ALDWW		535
SEQ ID NO: 21	moltype = AA length = 475	
FEATURE	Location/Qualifiers	
source	1..475	
	mol_type = protein	
	organism = Homo sapiens	
SEQUENCE: 21		
MEMASSAGSW LSGCLIPLVF LRLSVHVS GH AGDAGKFHVA LLGGTAELLC PLSLWPGTVP		60
KEVRWLRSPP PQRSQAVHIF RDGKDQDEDL MPEYKGRTVL VRDAQEGSVT LQILDVRLD		120
QGSYRCLIQV GNLSKEDTVI LQVAAPSVGS LSPSAVALAV ILPVLVLLIM VCLCLIWKQR		180
RAKEKLLYEH VTEVDNLLSD HAKEKGKHLK AVKKLRSELK LKRAAANS GW RRARLHFVAV		240
TLDPDTAHPK LILSEDQRCV RLGD RRQVPV DNPQRFDV V SILGSEYFTT GCHYWEVYVG		300
DKTKWILGVC SESVSRKGKV TASPANGHWL LRQSRGNEYE ALTSPQTSFR LKEPPRCVGI		360
FLDYEAGVIS FYNVTNKSHI FTFTHNFSGP LRPFFEPCLH DGGKNTAPLV ICSELHKSEE		420
SIVPRPEGKG HANGDVS LKV NSSLLPPKAP ELKDIILSLP PDLGPALQEL KAPSF		475
SEQ ID NO: 22	moltype = AA length = 252	
FEATURE	Location/Qualifiers	
source	1..252	
	mol_type = protein	
	organism = Homo sapiens	
SEQUENCE: 22		
MASLSRPSLP SCLCSFLLLL LLQVSSSYAG QFRVIGPRHP IRALVGDEVE LPCRISPGKN		60
ATGMEVGN YR PPF SRVHLY R NGKDQDQDQ APEYRGRT EL LKDAIGEGKV TLRIRNVRF		120
DEGGFTCFER DHSYQEEAAM ELKVEDPFYV VSPGVLVLLA VLPVLLQIT VGLVFLCLQY		180
RLRGKLR AEI ENLHRTDPH FLRVPCWKIT LFVIVPV LGP LVALIICYNW LHRRLAGQFL		240
EELLPHLEAL SG		252

-continued

SEQ ID NO: 23	moltype = AA length = 282				
FEATURE	Location/Qualifiers				
source	1..282				
	mol_type = protein				
	organism = Homo sapiens				
SEQUENCE: 23					
YAEATLVVRN	ASAESVSCLV	HNPVLTEEGK	SVISLPEKLQ	TELASLKVNG	PSQPILVRVG 60
EDIQLTCYLS	PKANAQSM EV	RWDRSHRYPA	VHVYMDGDHV	AGEQMAEYRG	RTVLVSDAID 120
EGRLTLQILS	ARPSDDGQYR	CLFEKDDVYQ	EASLDLKVVG	LGSSPLITVE	GQEDGEMQPM 180
CSSDGFPPQP	HVPWRDMEGK	TIPSSSQALT	QGSHGLFHVQ	TLLRVTNISA	VDVTCISISIP 240
FLGEEKIATF	SLSESRTFL	WKTLLVWGLL	LAVAVGLPRK	RS	282

- 1-52. (canceled)
53. A method of treating a patient having an autoimmune or inflammatory disease comprising administering to the patient a therapeutically effective dose of a BTNL3 protein comprising
- (a) the amino acid sequence of amino acids 18-236 of SEQ ID NO:2 or the amino acid sequence of amino acids 18-166 of SEQ ID NO:9,
 - (b) an amino acid sequence at least 90% identical to amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9, wherein the alignment window of the amino acid sequence with amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9 is at least 80 amino acids long, or
 - (c) an amino acid sequence that has no more than 20 insertions, deletions, or substitutions of a single amino acid relative to the sequence of amino acids 18-236 of SEQ ID NO:2 or amino acids 18-166 of SEQ ID NO:9,

- wherein the BTNL3 protein can inhibit the proliferation of a T cell stimulated by an immobilized anti-CD3 antibody.
54. The method of claim 53, wherein the autoimmune or inflammatory disease is selected from the group consisting of systemic lupus erythematosus, rheumatoid arthritis, an inflammatory bowel disease, a transplantation-related condition, Crohn's disease, ulcerative colitis, psoriasis, sarcoidosis, asthma, or a fibrotic disease.
55. The method of claim 54, wherein the autoimmune or inflammatory disease is Crohn's disease.
56. The method of claim 54, wherein the autoimmune or inflammatory disease is ulcerative colitis.
57. The method of claim 54, wherein the autoimmune or inflammatory disease is a fibrotic disease.
- 58-77. (canceled)

* * * * *