




- (51) **International Patent Classification:**
A61K 39/395 (2006.01) C07K 16/28 (2006.01)
A61P 35/00 (2006.01)
- (21) **International Application Number:**
PCT/EP2010/005245
- (22) **International Filing Date:**
26 August 2010 (26.08.2010)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
09011047.9 28 August 2009 (28.08.2009) EP
- (71) **Applicant (for all designated States except US):** F. **HOFFMANN-LA ROCHE AG** [CH/CH]; Grenzacherstrasse 124, CH-4070 Basel (CH).
- (72) **Inventors; and**
- (75) **Inventors/Applicants (for US only):** **BOSENMAIER, Birgit** [DE/DE]; Horst-Wolfram-Geissler-Weg 10, 82229 Seefeld (DE). **FERTIG, Georg** [DE/DE]; St.-Klara-Strasse 2, 82377 Penzberg (DE). **KRELL, Hans-Willi** [AT/DE]; Zugspitzstrasse 14a, 82377 Penzberg (DE). **LAMMERS, Reiner** [DE/DE]; Pfizerstrasse 32, 72070 Tuebingen (DE). **LIFKE, Alexander** [DE/DE]; An der Freiheit 26, 82377 Penzberg (DE). **NIEDERFELLNER, Gerhard** [DE/DE]; Dorfstrasse 36b, 82386 Oberhausen, Germany (DE).
- (74) **Agent:** **BURGER, Alexander**; Roche Diagnostics GmbH, Patent Department (TR-E), Postfach 11 52 (DE).
- (81) **Designated States (unless otherwise indicated, for every kind of national protection available):** AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States (unless otherwise indicated, for every kind of regional protection available):** ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
- of inventorship (Rule 4.17(iv))
- Published:**
- with international search report (Art. 21(3))
 - with sequence listing part of description (Rule 5.2(a))

(54) **Title:** ANTIBODIES AGAINST CDCP1 FOR THE TREATMENT OF CANCER

Fig. 1



(57) **Abstract:** The present disclosure relates to antibodies against human CDCP1 binding to the same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer.

Antibodies against CDCP1 for the treatment of cancer

The present invention relates to antibodies against human CDCP1 binding to the same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer.

Background of the Invention

5 Human CDCP1 ((CUB domain containing protein 1, B345, CD318, SIMA135, TRASK; SEQ ID NO: 1 and variants with mutation R525Q (i.e. replacement of Arginine (R) with Glutamine (Q) at amino acid position 525 of SEQ ID NO: 1) and/or mutation G709D (i.e. replacement of Glycine (G) with Aspartic acid (D) at amino acid position 709 of SEQ ID NO: 1)) is a transmembrane protein containing
10 three extracellular CUB domains. This protein is found to be overexpressed in colon and lung cancers. Its expression level is correlated with the metastatic ability of carcinoma cells. It has been shown to be tyrosine phosphorylated in a cancer cell line. (WO 2002/004508; Scherl-Mostageer, M. et al., *Oncogene* 20 (2001) 4402-8; Hooper, J.D., et al., *Oncogene* 22 (2003) 1783-94; Perry, S.E. et al., *FEBS Lett.* 15 581 (2007) 1137-42; Brown, T.A., *J. Biol. Chem.* 279 (2004) 14772-14783; Ota, T., et al., *Nat. Genet.* 36 (2004) 40-45). Alternatively spliced transcript variants encoding distinct isoforms have been reported.

WO 2002/004508 refers to CDCP1 as tumor associated antigen B345. WO 2004/074481 relates to CDCP1 as glycoprotein antigen SIMA135 expressed in
20 metastatic tumor cells. WO 2005/042102 relates to CDCP1 as protein involved in ovarian cancer. WO 2007/ 005502 relates to methods and compositions for treating diseases targeting CDCP1.

US 2004/0053343 (and Conze, T. et al., *Ann. N. Y. Acad. Sci.* 996 (2003) 222-6 and Buhning, H.J. et al., *Stem Cells* 22 (2004) 334-43) relates to CDCP1 antibodies
25 for identifying and/ or certain cell stem cell populations.

Summary of the Invention

One aspect of the invention is an antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer.

The invention further comprises an antibody characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer, and further characterized in that

- 5 a) the heavy chain variable domain is SEQ ID NO: 7 and the light chain variable domain is SEQ ID NO: 8,
b) the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or
c) the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or
10 d) the heavy chain variable domain is SEQ ID NO: 31 and the light chain variable domain is SEQ ID NO: 32, or
or a humanized version thereof.

15 The invention further comprises an antibody characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer, and further characterized in that

- 20 a) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region of SEQ ID NO: 14, or
b) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22, or
25 c) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.
30

The invention further comprises an antibody according to the invention, characterized in that said antibody is of human IgG1 subclass.

Another aspect of the invention is a pharmaceutical composition comprising an antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551), for the treatment of cancer.

5 Another aspect of the invention is the use of an antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) in the preparation of a medicament for the treatment of cancer

10 Another aspect of the invention is a method of treatment of a patient suffering from cancer by administering an antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) to said patient in the need of such treatment.

The invention further comprises an antibody according to the invention, characterized in that

15 a) the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or
b) the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or
c) the heavy chain variable domain is SEQ ID NO: 31 and the light chain
20 variable domain is SEQ ID NO: 32, or
or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

25 a) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region of SEQ ID NO: 14, or
b) the heavy chain variable domain comprises a CDRH1 region of SEQ ID
30 NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22, or

5 c) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.

10 Preferably said antibody is characterized in that said antibody is of human IgG1 subclass. The invention further comprises a pharmaceutical composition comprising said antibody. The invention further comprises the use of said antibody according to the preparation of a medicament for the treatment of cancer. The invention further comprises a method of treatment of a patient suffering from cancer by administering said antibody to said patient in the need of such treatment.

15 The invention provides nucleic acid encoding the antibody according to the invention. The invention further provides expression vectors containing nucleic acid according to the invention capable of expressing said nucleic acid in a prokaryotic or eukaryotic host cell, and host cells containing such vectors for the recombinant production of an antibody according to the invention.

The invention further comprises a prokaryotic or eukaryotic host cell comprising a vector according to the invention.

20 The invention further comprises a method for the production of a recombinant antibody according to the invention, characterized by expressing a nucleic acid according to the invention in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant. The invention further comprises the antibody obtained by such a recombinant method.

25 It has now surprisingly been found that the antibodies specifically binding to CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) are especially useful for the treatment of cancer compared to CDCP1 antibodies binding to other epitopes of CDCP1 like e.g. CUB1 (Deposition No. DSM ACC2569).

Detailed Description of the Invention

30 The CUB4 antibody refers to the deposited antibody with the Deposition No. DSM ACC2551 from DE 10242146 (EP 1 396 501, US 7,541,030) with the heavy chain variable domain (VH) of SEQ ID NO: 7 and the light chain variable domain (VL)

of SEQ ID NO: 8. Said CUB4 antibody is specifically binding to human CDCP1. (The Deposition of No. DSM ACC2551 (DSMZ) was made by Eberhard-Karls-University Tübingen, Universitätsklinikum Tübingen, Geissweg 3, 72076 Tübingen).

- 5 The invention comprises an antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer.

The invention further comprises an antibody according to the invention, characterized in that

- 10 the heavy chain variable domain is SEQ ID NO: 7 and the light chain variable domain is SEQ ID NO: 8,
or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

- 15 the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or
or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

- 20 the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or
or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

- 25 the heavy chain variable domain is SEQ ID NO: 31 and the light chain variable domain is SEQ ID NO: 32, or
or a humanized version thereof.

The invention further comprises an antibody according to the invention, characterized in that

5 the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region of SEQ ID NO: 14, or

The invention further comprises an antibody according to the invention, characterized in that

10 the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22, or

The invention further comprises an antibody according to the invention, characterized in that

15 the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.

20 The term “antibody” encompasses the various forms of antibody structures including but not being limited to whole antibodies and antibody fragments. The antibody according to the invention is preferably a humanized antibody, chimeric antibody, or further genetically engineered antibody as long as the characteristic properties according to the invention are retained. “Antibody fragments” comprise
25 a portion of a full length antibody, preferably the variable domain thereof, or at least the antigen binding site thereof. Examples of antibody fragments include diabodies, single-chain antibody molecules, and multispecific antibodies formed from antibody fragments. scFv antibodies are, e.g., described in Huston, J.S., Methods in Enzymol. 203 (1991) 46-88. In addition, antibody fragments comprise
30 single chain polypeptides having the characteristics of a V_H domain, namely being able to assemble together with a V_L domain, or of a V_L domain binding to CDCP1, namely being able to assemble together with a V_H domain to a functional antigen binding site and thereby providing the properties of an antibody according to the

invention. The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of a single amino acid composition. The term "humanized antibody" or "humanized version of an antibody" refers to antibodies in which the framework and/or "complementary determining regions" (CDR) have been modified to comprise the CDR (e.g. CDR3) of an immunoglobulin of different species as compared to that of the parent immunoglobulin. In a preferred embodiment, a mouse CDR (e.g. CDR3) is grafted into the framework region of a human antibody to prepare the "humanized antibody" (see, e.g., Riechmann, L. et al., Nature 332 (1988) 323-327; and Neuberger, M. S. et al., Nature 314 (1985) 268-270).

The terms "monoclonal antibody" or "monoclonal antibody composition" as used herein refer to a preparation of antibody molecules of a single amino acid composition.

The term "chimeric antibody" refers to a monoclonal antibody comprising a variable region, i.e., binding region, from mouse and at least a portion of a constant region derived from a different source or species, usually prepared by recombinant DNA techniques. Chimeric antibodies comprising a mouse variable region and a human constant region are especially preferred. Such rat/human chimeric antibodies are the product of expressed immunoglobulin genes comprising DNA segments encoding rat immunoglobulin variable regions and DNA segments encoding human immunoglobulin constant regions. Other forms of "chimeric antibodies" encompassed by the present invention are those in which the class or subclass has been modified or changed from that of the original antibody. Such "chimeric" antibodies are also referred to as "class-switched antibodies." Methods for producing chimeric antibodies involve conventional recombinant DNA and gene transfection techniques now well known in the art (see, e.g., Morrison, S.L. et al., Proc. Natl. Acad. Sci. USA 81 (1984) 6851-6855; US 5,202,238 and US 5,204,244).

Human CDCP1 ((CUB domain containing protein 1, B345, CD318, SIMA135, TRASK; SEQ ID NO: 1 and variants with mutation R525Q (i.e. replacement of Arginine (R) with Glutamine (Q) at amino acid position 525 of SEQ ID NO: 1) and/or mutation G709D (i.e. replacement of Glycine (G) with Aspartic acid (D) at amino acid position 709 of SEQ ID NO: 1)) is a transmembrane protein containing three extracellular CUB domains. This protein is found to be overexpressed in colon and lung cancers. Its expression level is correlated with the metastatic ability

of carcinoma cells. It has been shown to be tyrosine phosphorylated in a cancer cell line. (WO 2002/004508; Scherl-Mostageer, M., et al., *Oncogene* 20 (2001) 4402-8; Hooper, J.D. et al., *Oncogene* 22 (2003) 1783-94; Perry, S.E. et al., *FEBS Lett.* 581 (2007) 1137-42; Brown, T.A., *J. Biol. Chem.* 279 (2004) 14772-14783; Ota, T. et al., *Nat. Genet.* 36 (2004) 40-45). Alternatively spliced transcript variants encoding distinct isoforms have been reported.

The term "Kabat numbering" or "numbering according to Kabat" or "EU index" unless otherwise stated, is defined as the numbering of the residues in, e.g., an IgG antibody using the EU index as in Kabat, et al. (*Sequences of Proteins of Immunological Interest*, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991)).

As used herein, "specifically binding to human CDCP1" refers to an antibody specifically binding to the human CDCP1 antigen. The binding affinity is of KD-value of 1.0×10^{-8} mol/l or lower, preferably of a KD-value of 1.0×10^{-9} mol/l or lower. The binding affinity is determined with a standard binding assay, such as surface plasmon resonance technique (Biacore®). Thus an "antibody specifically binding to human CDCP1" as used herein refers to an antibody binding to human CDCP1 antigen with a binding affinity of KD 1.0×10^{-8} mol/l or lower (e.g. of KD 1.0×10^{-8} mol/l - 1.0×10^{-13} mol/l, preferably of KD 1.0×10^{-9} mol/l - 1.0×10^{-12} mol/l).

The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or a humanized version thereof.

The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or a humanized version thereof.

The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

the heavy chain variable domain is SEQ ID NO: 31 and the light chain variable domain is SEQ ID NO: 32, or a humanized version thereof.

The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

5 the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region of SEQ ID NO: 14.

10 The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region of
15 SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22.

The invention further comprises an antibody specifically binding to human CDCP1, characterized in that

20 the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.

25 The term "epitope" denotes a protein determinant of human CDCP1 capable of specifically binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually epitopes have specific three dimensional structural characteristics, as well as specific charge characteristics. Conformational and nonconformational epitopes are distinguished in that the binding to the former but not the latter is lost in the
30 presence of denaturing solvents.

The term “binding to the same epitope as CUB4 (Deposition No. DSM ACC2551)” as used herein refers to an anti-CDCP1 antibody of the invention that binds to the same epitope on CDCP1 to which the antibody CUB4 (Deposition No. DSM ACC2551) binds. The epitope binding property of an anti-CDCP1 antibody of the present invention may be determined using techniques known in the art. The CDCP1 antibody is measured at 25 °C by Surface Plasmon Resonance (SPR) in an in vitro competitive binding inhibition assay to determine the ability of antibody CUB4 (Deposition No. DSM ACC2551) to inhibit binding of the test antibody to CDCP1 (see Fig 2). Binding of antibodies binding to the same epitope as CUB4 is inhibited and no binding signal is detected after addition of the test antibody. (e.g. 100 seconds after the injection time (= 0 seconds) of the test antibody the binding signal is not higher than the signal at the time of injection; the signal is measured in RU (Relative Units)) (e.g. CDCP1-004, CDCP1-012, CDCP1-015, see Figure 3a). Binding of antibodies binding to a different epitope as CUB4 is not inhibited and a binding signal is detected after addition of the test antibody (e.g. CUB1 and CUB3, see Figure 3b) (e.g. 100 seconds after the injection time (= 0 seconds) the binding signal is higher than the signal at the time of injection; the signal is measured in RU (Relative Units)). This can be investigated by a BIAcore assay (Pharmacia Biosensor AB, Uppsala, Sweden) as described. e.g. in Example 2.

The “variable domain” (variable domain of a light chain (VL), variable domain of a heavy chain (VH)) as used herein denotes each of the pair of light and heavy chain domains which are involved directly in binding the antibody to the antigen. The variable light and heavy chain domains have the same general structure and each domain comprises four framework (FR) regions whose sequences are widely conserved, connected by three “hypervariable regions” (or complementary determining regions, CDRs). The framework regions adopt a β -sheet conformation and the CDRs may form loops connecting the β -sheet structure. The CDRs in each chain are held in their three-dimensional structure by the framework regions and form together with the CDRs from the other chain the antigen binding site. The antibody’s heavy and light chain CDR3 regions play a particularly important role in the binding specificity/affinity of the antibodies according to the invention and therefore provide a further object of the invention.

The term “antigen-binding portion of an antibody” when used herein refer to the amino acid residues of an antibody which are responsible for antigen-binding. The antigen-binding portion of an antibody comprises amino acid residues from the

“complementary determining regions” or “CDRs”. The term “antigen-binding portion” of an antibody of the invention may contain six complementarity determining regions (CDRs) which contribute in varying degrees to the affinity of the binding site for antigen. There are three heavy chain variable domain CDRs (CDRH1, CDRH2 and CDRH3) and three light chain variable domain CDRs (CDRL1, CDRL2 and CDRL3). The extent of CDR and framework regions (FRs) is determined by comparison to a compiled database of amino acid sequences in which those regions have been defined according to variability among the sequences.

“Framework” or “FR” regions are those variable domain regions other than the hypervariable region residues as herein defined. Therefore, the light and heavy chain variable domains of an antibody comprise from N- to C-terminus the domains FR1, CDR1, FR2, CDR2, FR3, CDR3, and FR4. CDR and FR regions are determined according to the standard definition of Kabat et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991) and/or those residues from a “hypervariable loop”.

The terms “nucleic acid” or “nucleic acid molecule”, as used herein, are intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, but preferably is double-stranded DNA.

The term “amino acid” as used within this application denotes the group of naturally occurring carboxy α -amino acids comprising alanine (three letter code: ala, one letter code: A), arginine (arg, R), asparagine (asn, N), aspartic acid (asp, D), cysteine (cys, C), glutamine (gln, Q), glutamic acid (glu, E), glycine (gly, G), histidine (his, H), isoleucine (ile, I), leucine (leu, L), lysine (lys, K), methionine (met, M), phenylalanine (phe, F), proline (pro, P), serine (ser, S), threonine (thr, T), tryptophan (trp, W), tyrosine (tyr, Y), and valine (val, V).

The antibody according to the invention is characterized in that the constant region is of human origin, and is preferably of human IgG1 subclass. The constant region includes the heavy chain and light chain constant region of an antibody. The heavy chain constant region comprises in N-terminal to C-terminal direction an antibody constant heavy chain domain 1 (CH1), an antibody hinge region (HR), an antibody heavy chain constant domain 2 (CH2), and an antibody heavy chain constant domain 3 (CH3),; and optionally an antibody heavy chain constant domain 4 (CH4)

in case of an antibody of the subclass IgE. The light chain constant region comprises an antibody light chain constant domain (CL). The antibody light chain constant domain (CL) can be κ (kappa) or λ (lambda). Such constant chains are well known in the state of the art and e.g. described by Kabat, E., A., (see e.g. Johnson, G. and Wu, T., T., *Nucleic Acids Res.* 28 (2000) 214-218). For example, a useful human heavy chain constant region of IgG1 subclass comprises an amino acid sequence of SEQ ID NO: 3. For example, a useful human light chain constant region comprises an amino acid sequence of a kappa-light chain constant region of SEQ ID NO: 4; another useful human light chain constant region comprises an amino acid sequence of a lambda-light chain constant region of SEQ ID NO: 5.

The "Fc part" of an antibody is not involved directly in binding of an antibody to an antigen, but exhibit various effector functions. A "Fc part of an antibody" is a term well known to the skilled artisan and defined on the basis of papain cleavage of antibodies. Depending on the amino acid sequence of the constant region of their heavy chains, antibodies or immunoglobulins are divided in the classes: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses (isotypes), e.g. IgG1, IgG2, IgG3, and IgG4, IgA1, and IgA2. According to the heavy chain constant regions the different classes of immunoglobulins are called α , δ , ϵ , γ , and μ , respectively.

The Fc part of an antibody is directly involved in ADCC (antibody-dependent cell-mediated cytotoxicity) and CDC (complement-dependent cytotoxicity) based on complement activation, C1q binding and Fc receptor binding. Complement activation (CDC) is initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. While the influence of an antibody on the complement system is dependent on certain conditions, binding to C1q is caused by defined binding sites in the Fc part. Such binding sites are known in the state of the art and described e.g. by Boackle, R.J. et al., *Nature* 282 (1979) 742-743; Lukas, T.J. et al., *J. Immunol.* 127 (1981) 2555-2560; Brunhouse, R. and Cebra, J.J., *Mol. Immunol.* 16 (1979) 907-917; Burton, D.R. et al., *Nature* 288 (1980) 338-344; Thommesen, J.E. et al., *Mol. Immunol.* 37 (2000) 995-1004; Idusogie, E.E. et al., *J. Immunol.* 164 (2000) 4178-4184; Hezareh, M. et al., *J. Virology* 75 (2001) 12161-12168; Morgan, A. et al., *Immunology* 86 (1995) 319-324; EP 0307434. Such binding sites are e.g. L234, L235, D270, N297, E318, K320, K322, P331 and P329 (numbering according to EU index of Kabat, E.A., see below). Antibodies of subclass IgG1, IgG2 and IgG3 usually show complement activation and C1q and

C3 binding, whereas IgG4 do not activate the complement system and do not bind C1q and C3.

The antibody according to the invention comprises a Fc part derived from human origin and preferably all other parts of the human constant regions. As used herein the term “Fc part derived from human origin” denotes a Fc part which is either a Fc part of a human antibody of the subclass IgG1, IgG2, IgG3 or IgG4, preferably a Fc part from human IgG1 subclass, a mutated Fc part from human IgG1 subclass (preferably with a mutation on L234A + L235A), a Fc part from human IgG4 subclass or a mutated Fc part from human IgG4 subclass (preferably with a mutation on S228P). Mostly preferred are the human heavy chain constant regions of human IgG1 subclass (see e.g. of SEQ ID NO: 3), of human IgG1 subclass with mutations L234A and L235A, of human IgG4 subclass (see e.g. of SEQ ID NO: 6), or of human IgG4 subclass with mutation S228P.

The term “antibody-dependent cellular cytotoxicity (ADCC)” refers to lysis of human target cells by an antibody according to the invention in the presence of effector cells. ADCC is measured preferably by the treatment of a preparation of CDCP1 expressing cells with an antibody according to the invention in the presence of effector cells such as freshly isolated PBMC or purified effector cells from buffy coats, like monocytes or natural killer (NK) cells or a permanently growing NK cell line.

The term “complement-dependent cytotoxicity (CDC)” denotes a process initiated by binding of complement factor C1q to the Fc part of most IgG antibody subclasses. Binding of C1q to an antibody is caused by defined protein-protein interactions at the so called binding site. Such Fc part binding sites are known in the state of the art (see above). Such Fc part binding sites are, e.g., characterized by the amino acids L234, L235, D270, N297, E318, K320, K322, P331, and P329 (numbering according to EU index of Kabat). Antibodies of subclass IgG1, IgG2, and IgG3 usually show complement activation including C1q and C3 binding, whereas IgG4 does not activate the complement system and does not bind C1q and/or C3.

Cell-mediated effector functions of monoclonal antibodies can be enhanced by engineering their oligosaccharide component as described in Umana, P. et al., Nature Biotechnol. 17 (1999) 176-180, and US 6,602,684. IgG1 type antibodies, the most commonly used therapeutic antibodies, are glycoproteins that have a

conserved N-linked glycosylation site at Asn297 in each CH2 domain. The two complex biantennary oligosaccharides attached to Asn297 are buried between the CH2 domains, forming extensive contacts with the polypeptide backbone, and their presence is essential for the antibody to mediate effector functions such as antibody dependent cellular cytotoxicity (ADCC) (Lifely, M.R. et al., *Glycobiology* 5 (1995) 813-822; Jefferis, R. et al., *Immunol. Rev.* 163 (1998) 59-76; Wright, A. and Morrison, S.L., *Trends Biotechnol.* 15 (1997) 26-32). Umana, P., et al. *Nature Biotechnol.* 17 (1999) 176-180 and WO 99/54342 showed that overexpression in Chinese hamster ovary (CHO) cells of $\beta(1,4)$ -N-acetylglucosaminyltransferase III (“GnTIII”), a glycosyltransferase catalyzing the formation of bisected oligosaccharides, significantly increases the in vitro ADCC activity of antibodies. Alterations in the composition of the Asn297 carbohydrate or its elimination affect also binding to Fc γ R and C1q (Umana, P. et al., *Nature Biotechnol.* 17 (1999) 176-180; Davies, J. et al., *Biotechnol. Bioeng.* 74 (2001) 288-294; Mimura, Y. et al., *J. Biol. Chem.* 276 (2001) 45539-45547; Radaev, S. et al., *J. Biol. Chem.* 276 (2001) 16478-16483; Shields, R.L. et al., *J. Biol. Chem.* 276 (2001) 6591-6604; Shields, R.L. et al., *J. Biol. Chem.* 277 (2002) 26733-26740; Simmons, L.C. et al., *J. Immunol. Methods* 263 (2002) 133-147).

Methods to enhance cell-mediated effector functions of monoclonal antibodies are reported e.g. in WO 2005/044859, WO 2004/065540, WO2007/031875, Umana, P. et al., *Nature Biotechnol.* 17 (1999) 176-180, WO 99/154342, WO 2005/018572, WO 2006/116260, WO 2006/114700, WO 2004/065540, WO 2005/011735, WO 2005/027966, WO 1997/028267, US 2006/0134709, US 2005/0054048, US 2005/0152894, WO 2003/035835 and WO 2000/061739 or e.g. in Niwa, R. et al., *J. Immunol. Methods* 306 (2005) 151-160; Shinkawa, T. et al., *J. Biol. Chem.* 278 (2003) 3466-3473; WO 03/055993 and US 2005/0249722.

Therefore in one embodiment of the invention, the antibody according to the invention is glycosylated (if it comprises an Fc part of IgG1 or IgG3 subclass) with a sugar chain at Asn297 whereby the amount of fucose within said sugar chain is 65% or lower (Numbering according to Kabat). In another embodiment is the amount of fucose within said sugar chain is between 5% and 65%, preferably between 20% and 40%. “Asn297” according to the invention means amino acid asparagine located at about position 297 in the Fc region. Based on minor sequence variations of antibodies, Asn297 can also be located some amino acids (usually not more than ± 3 amino acids) upstream or downstream of position 297, i.e. between

position 294 and 300. In one embodiment the glycosylated antibody according to the invention the IgG subclass is of human IgG1 subclass, of human IgG1 subclass with the mutations L234A and L235A or of IgG3 subclass. In a further embodiment the amount of N-glycolylneuraminic acid (NGNA) is 1% or less and/or the amount of N-terminal alpha-1,3-galactose is 1 % or less within said sugar chain. The sugar chains show preferably the characteristics of N-linked glycans attached to Asn297 of an antibody recombinantly expressed in a CHO cell.

The term “the sugar chains show characteristics of N-linked glycans attached to Asn297 of an antibody recombinantly expressed in a CHO cell” denotes that the sugar chain at Asn297 of the antibody according to the invention has the same structure and sugar residue sequence except for the fucose residue as those of the same antibody expressed in unmodified CHO cells, e.g. as those reported in WO 2006/103100.

The term “NGNA” as used within this application denotes the sugar residue N-glycolylneuraminic acid.

Glycosylation of human IgG1 or IgG3 occurs at Asn297 as core fucosylated biantennary complex oligosaccharide glycosylation terminated with up to two Gal residues. Human constant heavy chain regions of the IgG1 or IgG3 subclass are reported in detail by Kabat, E.A., et al., Sequences of Proteins of Immunological Interest, 5th ed., Public Health Service, National Institutes of Health, Bethesda, MD (1991), and by Brueggemann, M., et al., J. Exp. Med. 166 (1987) 1351-1361; Love, T.W. et al., Methods Enzymol. 178 (1989) 515-527. These structures are designated as G0, G1 (α -1,6- or α -1,3-), or G2 glycan residues, depending from the amount of terminal Gal residues (Raju, T.S., Bioprocess Int. 1 (2003) 44-53). CHO type glycosylation of antibody Fc parts is e.g. described by Routier, F.H., Glycoconjugate J. 14 (1997) 201-207. Antibodies which are recombinantly expressed in non-glycomodified CHO host cells usually are fucosylated at Asn297 in an amount of at least 85%. The modified oligosaccharides of the antibody may be hybrid or complex. Preferably the bisected, reduced/not-fucosylated oligosaccharides are hybrid. In another embodiment, the bisected, reduced/not-fucosylated oligosaccharides are complex.

According to the invention “amount of fucose” means the amount of said sugar within the sugar chain at Asn297, related to the sum of all glycostructures attached to Asn297 (e.g. complex, hybrid and high mannose structures) measured by

MALDI-TOF mass spectrometry and calculated as average value (see e.g. WO 2008/077546). The relative amount of fucose is the percentage of fucose-containing structures related to all glycostructures identified in an N-Glycosidase F treated sample (e.g. complex, hybrid and oligo- and high-mannose structures, resp.)
5 by MALDI-TOF.

The antibodies according to the invention are preferably produced by recombinant means. Such methods are widely known in the state of the art and comprise protein expression in prokaryotic and eukaryotic cells with subsequent isolation of the antibody polypeptide and usually purification to a pharmaceutically acceptable
10 purity. For the protein expression, nucleic acids encoding light and heavy chains or fragments thereof are inserted into expression vectors by standard methods. Expression is performed in appropriate prokaryotic or eukaryotic host cells like CHO cells, NS0 cells, SP2/0 cells, HEK293 cells, COS cells, yeast, or E.coli cells, and the antibody is recovered from the cells (supernatant or cells after lysis).

15 Recombinant production of antibodies is well-known in the state of the art and described, for example, in the review articles of Makrides, S.C., *Protein Expr. Purif.* 17 (1999) 183-202; Geisse, S. et al., *Protein Expr. Purif.* 8 (1996) 271-282; Kaufman, R.J., *Mol. Biotechnol.* 16 (2000) 151-160; Werner, R.G., *Drug Res.* 48 (1998) 870-880.

20 The antibodies may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. Purification is performed in order to eliminate other cellular components or other contaminants, e.g. other cellular nucleic acids or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis, and others well known in the
25 art (see Ausubel, F., et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987)).

Expression in NS0 cells is described by, e.g., Barnes, L.M. et al., *Cytotechnology* 32 (2000) 109-123; and Barnes, L.M. et al., *Biotech. Bioeng.* 73 (2001) 261-270. Transient expression is described by, e.g., Durocher, Y. et al., *Nucl. Acids. Res.* 30
30 (2002) E9. Cloning of variable domains is described by Orlandi, R. et al., *Proc. Natl. Acad. Sci. USA* 86 (1989) 3833-3837; Carter, P. et al., *Proc. Natl. Acad. Sci. USA* 89 (1992) 4285-4289; and Norderhaug, L. et al., *J. Immunol. Methods* 204 (1997) 77-87. A preferred transient expression system (HEK 293) is described by

Schlaeger, E.J., and Christensen, K., in *Cytotechnology* 30 (1999) 71-83 and by Schlaeger, E. J., in *J. Immunol. Methods* 194 (1996) 191-199.

5 The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, enhancers and polyadenylation signals.

10 Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading frame. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

20 The monoclonal antibodies are suitably separated from the culture medium by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography. DNA and RNA encoding the monoclonal antibodies are readily isolated and sequenced using conventional procedures. The hybridoma cells can serve as a source of such DNA and RNA. Once isolated, the DNA may be inserted into expression vectors, which are then transfected into host cells such as HEK 293 cells, CHO cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of recombinant monoclonal antibodies in the host cells.

30 As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that all progeny may not be precisely identical in DNA content, due to deliberate or inadvertent mutations. Variant progeny that have the same function or

biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

5 The term "transformation" as used herein refers to process of transfer of a vectors/nucleic acid into a host cell. If cells without formidable cell wall barriers are used as host cells, transfection is carried out e.g. by the calcium phosphate precipitation method as described by Graham, F.L. and van der Eb, A.J., *Virology* 10 52 (1973) 456-467. However, other methods for introducing DNA into cells such as by nuclear injection or by protoplast fusion may also be used. If prokaryotic cells or cells which contain substantial cell wall constructions are used, e.g. one method of transfection is calcium treatment using calcium chloride as described by Cohen, S.N. et al., *PNAS* 69 (1972) 2110-2114.

15 As used herein, "expression" refers to the process by which a nucleic acid is transcribed into mRNA and/or to the process by which the transcribed mRNA (also referred to as transcript) is subsequently being translated into peptides, polypeptides, or proteins. The transcripts and the encoded polypeptides are collectively referred to as gene product. If the polynucleotide is derived from genomic DNA, expression in a eukaryotic cell may include splicing of the mRNA.

20 A "vector" is a nucleic acid molecule, in particular self-replicating, which transfers an inserted nucleic acid molecule into and/or between host cells. The term includes vectors that function primarily for insertion of DNA or RNA into a cell (e.g., chromosomal integration), replication of vectors that function primarily for the replication of DNA or RNA, and expression vectors that function for transcription and/or translation of the DNA or RNA. Also included are vectors that provide more than one of the functions as described.

25 An "expression vector" is a polynucleotide which, when introduced into an appropriate host cell, can be transcribed and translated into a polypeptide. An "expression system" usually refers to a suitable host cell comprised of an expression vector that can function to yield a desired expression product.

30 One aspect of the invention is a pharmaceutical composition comprising an antibody according to the invention. Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a pharmaceutical composition. A further aspect of the invention is a method for the manufacture of a pharmaceutical composition comprising an antibody according to the invention. In

another aspect, the present invention provides a composition, e.g. a pharmaceutical composition, containing an antigen binding protein according to the present invention, formulated together with a pharmaceutical carrier.

5 Said antibodies specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) have been turned out to be especially useful for the treatment of cancer compared with other anti-CDCP1 antibodies as e.g. CUB1 antibody (deposited antibody with the Deposition No. DSM ACC2569 from DE 10242146 (EP 1 396 501, US 7,541,030)).

10 Therefore one aspect of the invention is said pharmaceutical composition for the treatment of cancer.

Another aspect of the invention is the use of an antibody according to the invention for the manufacture of a medicament for the treatment of cancer.

15 Another aspect of the invention is a method of treatment of a patient suffering from cancer by administering an antibody according to the invention to said patient in the need of such treatment.

20 As used herein, "pharmaceutical carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g. by injection or infusion).

25 A composition of the present invention can be administered by a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. To administer a compound of the invention by certain routes of administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation. For example, the compound may be administered to a subject in an appropriate carrier, for example, liposomes, or a diluent. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutical carriers include sterile aqueous solutions or dispersions
30 and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art.

The phrases “parenteral administration” and “administered parenterally” as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intra-arterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intra-articular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

The term “cancer” as used herein may be, for example, lung cancer, non small cell lung (NSCL) cancer, bronchioloalviolar cell lung cancer, bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, uterine cancer, ovarian cancer, rectal cancer, cancer of the anal region, stomach cancer, gastric cancer, colon cancer, breast cancer, uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, mesothelioma, hepatocellular cancer, biliary cancer, neoplasms of the central nervous system (CNS), spinal axis tumors, brain stem glioma, glioblastoma multiforme, astrocytomas, schwannomas, ependymomas, medulloblastomas, meningiomas, squamous cell carcinomas, pituitary adenoma, lymphoma, lymphocytic leukemia, including refractory versions of any of the above cancers, or a combination of one or more of the above cancers. Preferably such cancer is a breast cancer, ovarian cancer, cervical cancer, lung cancer or prostate cancer and more preferably lung cancer. Preferably such cancers are further characterized by CDCP1 expression or overexpression. More preferably such cancers are further characterized by overexpression.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be

brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

5 Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

10 Actual dosage levels of the active ingredients in the pharmaceutical compositions of the present invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, 15 the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

20 The composition must be sterile and fluid to the extent that the composition is deliverable by syringe. In addition to water, the carrier preferably is an isotonic buffered saline solution.

25 Proper fluidity can be maintained, for example, by use of coating such as lecithin, by maintenance of required particle size in the case of dispersion and by use of surfactants. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol or sorbitol, and sodium chloride in the composition.

30 The following examples, sequence listing and figures are provided to aid the understanding of the present invention, the true scope of which is set forth in the appended claims. It is understood that modifications can be made in the procedures set forth without departing from the spirit of the invention.

Description of the Sequence Listing

	SEQ ID NO:1	human CDCP1
	SEQ ID NO:2	extracellular-domain-(ECD)-comprising fragment of human CDCP1):
5	SEQ ID NO:3	IgG1 constant heavy chain region from human origin
	SEQ ID NO:4	kappa constant light chain region from human origin
	SEQ ID NO:5	lambda constant light chain region from human origin
	SEQ ID NO:6	IgG4 constant heavy chain region from human origin
10	SEQ ID NO: 7	heavy chain variable domain VH, CUB4 (Deposition No. DSM ACC2551)
	SEQ ID NO: 8	light chain variable domain VL, CUB4 (Deposition No. DSM ACC2551)
	SEQ ID NO: 9	heavy chain CDRH1 , Mab CDCP1-004
	SEQ ID NO: 10	heavy chain CDRH2, Mab CDCP1-004
15	SEQ ID NO: 11	heavy chain CDRH3, Mab CDCP1-004
	SEQ ID NO: 12	light chain CDRL1, Mab CDCP1-004
	SEQ ID NO: 13	light chain CDRL2, Mab CDCP1-004
	SEQ ID NO: 14	light chain CDRL3, Mab CDCP1-004
	SEQ ID NO: 15	heavy chain variable domain VH, Mab CDCP1-004
20	SEQ ID NO: 16	light chain variable domain VL, Mab CDCP1-004
	SEQ ID NO: 17	heavy chain CDRH1, Mab CDCP1-012
	SEQ ID NO: 18	heavy chain CDRH2, Mab CDCP1-012
	SEQ ID NO: 19	heavy chain CDRH3, Mab CDCP1-012
	SEQ ID NO: 20	light chain CDRL1, Mab CDCP1-012
25	SEQ ID NO: 21	light chain CDRL2, Mab CDCP1-012
	SEQ ID NO: 22	light chain CDRL3, Mab CDCP1-012
	SEQ ID NO: 23	heavy chain variable domain VH, Mab CDCP1-012
	SEQ ID NO: 24	light chain variable domain VL, Mab CDCP1-012
	SEQ ID NO: 25	heavy chain CDRH1, Mab CDCP1-015
30	SEQ ID NO: 26	heavy chain CDRH2, Mab CDCP1-015
	SEQ ID NO: 27	heavy chain CDRH3, Mab CDCP1-015
	SEQ ID NO: 28	light chain CDRL1, Mab CDCP1-01510
	SEQ ID NO: 29	light chain CDRL2, Mab CDCP1-015
	SEQ ID NO: 30	light chain CDRL3, Mab CDCP1-015
35	SEQ ID NO: 31	heavy chain variable domain VH, Mab CDCP1-015
	SEQ ID NO: 32	light chain variable domain VL, Mab CDCP1-015

Description of the Figures

5 **Figure 1** In vivo tumor growth inhibition in human lung cancer H322M xenograft of anti CDCP1 antibodies CUB4 and CUB1.

10 **Figure 2** Schematic Assay format of Surface Plasmon Resonance (SPR-)technology (BIAcore®) with immobilized CUB4 antibody, extracellular domain (ECD) of CDCP1, and further anti-CDCP1 test antibodies (e.g. CUB1 and CUB3).

15 **Figure 3** Biacore sensogram of immobilized CUB4 antibody, extracellular domain (ECD) of CDCP1 and anti-CDCP1 antibodies CUB1 and CUB3 (x-axis = time; y-axis = response in relative units (RU))

Fig 3a: The binding of CDCP1-004, CDCP1-012 and CDCP1-015 antibodies to the same epitope on CDCP1 as CUB4 antibody is shown.

20 **Fig 3b:** The binding of CUB1 and CUB3 antibodies to another epitope on CDCP1 than CUB4 antibody is shown.

Figure 4 Binding of the CDCP1 antibodies according to the invention to human CDCP1 Extracellular Domain (CDCP1-ECD) and subdomains of human CDCP1-ECD

25 **Figure 5** Stimulation of CDCP1 phosphorylation in HCT 116 cells

Figure 6 Stimulation of CDCP1 internalisation and phosphorylation in MDAMB-231 cells

Example 1

30 **In vivo tumor growth inhibition of anti-CDCP1 antibody CUB4 and in comparison to CUB1 antibody**

Study name: CDCP1_PZ_H322M_002

The present in vivo study was performed to compare the efficacy of anti-CDCP1 specific antibodies CUB4 with antibodies binding another epitope, as e.g. CUB1

5 H322M non small cell lung cancer cells were originally obtained from the NCI collection and were deposited after expansion in the Roche cell bank, Penzberg. Tumor cell line was routinely cultured in RPMI 1640 medium supplemented with 10% fetal bovine serum and 2mM L-glutamine at 37°C in a water-saturated atmosphere at 5% CO₂. Passage 4 was used for cell transplantation.

The human non small cell lung cancer cell line H322M was subcutaneously inoculated (5x10⁶ cells) with matrigel into the right flank of the mice.

10 Animal treatment started at the day of randomisation 17 days after cell transplantation. Antibodies were administered i.p. q7d until study termination day 62 at the indicated dosage of 10 mg/kg. Also the corresponding vehicle was administered on the same days. The administration volume was 10 ml/kg.

Groups:

Treatment Group 1: Vehicle

Treatment Group 2: murine CUB4 10 mg/kg i.p.;

15 Treatment Group 3: murine CUB1 10 mg/kg i.p.;

TGI (Tumor growth inhibition in%)

20 The following tables show the values of the tumor growth inhibition, which is calculated as $100 - \frac{\text{Average}(T_{\text{treatment}}[\text{day } x] - T_{\text{treatment}}[\text{baseline}])}{\text{Average}(T_{\text{reference}}[\text{day } x] - T_{\text{reference}}[\text{baseline}])} * 100$ for each group and time point based on means and medians, respectively. The reference for the TGI calculations was chosen to be group 'Vehicle'. Results are also shown in Figure 1.

Table 1: Tumor growth inhibition (TGI)

Group	Compound	Treatment schedule	Dose (mg/kg)	Median tumor volume at staging day 17 (mm ³)	Median tumor volume at day 62, (mm ³)	TGI %
1	vehicle	1x/week i.p.	-	175.7	926.3	-
2	CUB4	1x/week i.p.	10mg/kg	174.7	311.8	82
3	CUB1	1x/week i.p.	10mg/kg	172.8	414.1	68

The CUB4 antibody (Deposition No. DSM ACC2551) surprisingly shows a clearly higher tumor growth inhibition than CUB1 antibody (deposited antibody with the Deposition No. DSM ACC2569 from DE 10242146 (EP 1 396 501, US 7,541,030)).

Analogously the in vivo tumor growth inhibition of the anti-CDCP1 antibodies according to the invention CDCP1_004, CDCP1_012 and CDCP1_015 can be determined.

Example 2

10 Epitope binding assay (Biacore)

To determine the epitope regions of different test anti-CDCP1 antibodies, CUB4 (Deposition No. DSM ACC2551) was immobilized on the surface of a CM5 biosensorchip using amine-coupling chemistry. Flow cells were activated with a 1:1 mixture of 0.1 M N-hydroxysuccinimide and 0.4 M 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide at a flow rate of 5 μ l/min. Anti-CDCP1 antibodies CUB4 (Deposition No. DSM ACC2551) was injected in sodium acetate, pH 4.5-5.0 at 10-30 μ g/ml, 12 minutes, which resulted in a surface density of approx 15000 RU. Surfaces were blocked with an injection of 1 M ethanolamine/HCl pH 8.5. Soluble ECD of human CDCP1 (SEQ. ID NO.:2) (analyte 1) and anti-CDCP1 antibodies (analyte 2) were diluted in PBST + 0.8M NaCl and injected at a flow rate of 30 μ l/min. The contact time (association phase) was 150-200 sec for the ECD of human CDCP1 at a concentration of 250 nM-500nM and 300 sec for the anti-CDCP1 antibody at a concentration of 100 nM.

Then the chip surface was washed with PBST + 0.8M NaCl for 3 min (dissociation phase). All interactions were performed at exactly 25°C (standard temperature). A regeneration solution of 10 mM Glycine, pH 2.0 was injected for 60-150 sec to remove any non-covalently bound protein after each binding cycle. Signals were detected at a detection rate of one signal per second.

To determine whether an anti-CDCP1 antibody binds to the same or a different epitope of human CDCP1 as CUB4, the ECD of human CDCP1 was injected and bound by the immobilized antibodies. Shortly after the binding, antibodies with unknown epitope were injected. Test anti-CDCP1-antibodies which do not show an increase of the binding signal after injection (i.e. e.g. 100 seconds after the injection time (= 0 seconds) of the test antibody, the binding signal is not higher than the signal at the time of injection; the signal is measured in RU (Relative Units)), are binding to the same epitope as CUB4. Test anti-CDCP1-antibodies which show an increase of the binding signal are binding to a different epitope as CUB4 (e.g. 100 seconds after the injection time (= 0 seconds) the binding signal is higher than the signal at the time of injection; the signal is measured in RU (Relative Units)).

Results of the triple determinations are shown in Figure 3 a and 3 b. Antibodies CUB3, CUB1 (see Fig 3b) were found to bind at different epitopes as CUB4, whereas anti-CDCP1 antibodies CDCP1-004, CDCP1-012 and CDCP1-015 were identified as binding to the same epitope as CUB4 (see Table 2 and Fig 3a, all triple experiments). Antibodies CUB3 and CUB1 each were found to bind to another epitope as CUB4 and are different from one another (CUB1 \neq CUB3). (see Table 2 and Fig 3b, all triple experiments). Control experiments with CUB4 and dilution buffer as negative controls did not show an increase in the binding signal.

CUB1 showed no sandwich complex formation with preformed complex consisting of amine coupled Mab CUB3 and vice versa. (data not shown).

CDCP1-004, CDCP1-012 and CDCP1-015 showed a sandwich complex formation with both preformed complexes consisting of Mab CUB1 or Mab CUB3, respectively and antigen. (data not shown)

Table 2 (see also Figure 3):

Antibodies binding to the same epitope as CUB4	Antibodies binding to the different epitope as CUB4
Murine CUB4	Murine CUB1
Chimeric CUB4	Murine CUB3
Murine CDCP1-004	
Murine CDCP1-012	
Murine CDCP1-015	

In a separate Biacore experiment the binding constants and affinities of anti-CDCP1 antibodies to immobilized hCDCP1-ECD have been determined as follows:

5

Table 3:

Mab<hCDCP1>	antigene	k_a [1/Ms]	k_d [1/s]	K_D[M]	t(1/2) [min]
Mab CUB4	hum. CDCP1 ECD	9.9E+05	5.2E-04	5.2E-10	22.4
Mab CDCP1-015	hum. CDCP1 ECD	4.6E+05	2.2E-04	4.7E-10	52.8
Mab CUB1	hum. CDCP1 ECD	1.4E+06	1.9E-02	1.4E-08	0.6
Mab CUB3	hum. CDCP1 ECD	5.4E+04	3.1E-02	5.7E-07	0.4

k_a: association rate constant, k_d dissociation rate constant, K_D dissociation equilibrium constant (binding affinity) , t ½ : halflife time of complex,

10

Example 3

Antigene specific ELISA

Soluble CDCP1 extracellular Domain (CDCP1-ECD) (SEQ ID No: 2), as well as Short1 (huCDCP1_SH1_ECD aa1-216)_SBP) and Short5 (hu_CDCP1_SH5_ECD (aa 1-361)_SBP), which comprehend amino acids 1-216 of CDCP1 extracellular domain and 1-361 of CDCP1 extracellular domain, respectively, each fused to streptavidine Binding Protein (SBP) (SEQ ID No: 2), were captured on a

15

streptavidine plate. To define optimal binding of the antibodies to SBP-CDCP1-ECD, SBP-CDCP1 Short1 and SBP-CDCP1-Short5, 96 well polystyrene plates (Roche, streptavidin-coated, ID- No. 1989685) have been coated with pure or 1:4 diluted (dissolved in Dulbecco's Modified Eagle's Medium, PAN Biotech, containing 10% of Fetal Bovine Serum, Pan Biotech ID-No 3302-P251116) HEK293 supernatant. For the standard coating, SBP-CDCP1-Short5 containing HEK293 supernatant was diluted (1:4) contrary to undiluted SBP-CDCP1-Short1 and SBP-CDCP1-ECD supernatant and incubated overnight at 2-8°C (60µl). Intensive washing of the microtiter plate is necessary to remove remaining unbound SBP-CDCP1-ECD, SBP-CDCP1-Short1 and SBP-CDCP1-Short5.

The coated wells were blocked by performing an one hour blocking step using blocking reagent for ELISA (Roche 11112589) 250µl/well.

Anti-CDCP1 antibodies CUB4 (Deposition No. DSM ACC2551) and CDCP1_004, CDCP1_012 and CDCP1_015 were tested using a 1:500 dilution (0,1µg antibody diluted in 50µl PBS containing 1% BSA Fraction V, Sigma A3059). 50µl per well for each sample was incubated for 60 minutes at room temperature. After intensive washing using PBS-T (0,05% Tween 20 in PBS, Fluka #08057) 50µl of goat anti-mouse IgG antibodies coupled with HRP (BioRad #1706516, dilution 1:1000) were added and incubated for 1 hour at room temperature. After intensive washing the binding of the antibodies was detected with BM Blue POD Substrate (Roche 11484281001) 50µl. Absorbance at 370nm/492nm was measured using a standard photometer.

The binding region of the antibodies CDCP1-004, CDCP1-012, CDCP1-015 and CUB4 is located within the aa 1 and 216 in the extracellular domain of CDCP1. As a consequence all these antibodies also recognize the complete ECD of CDCP1 and a construct containing aa 1-361 (short 5) (see Figure 4).

Example 4

Stimulation of CDCP1 internalisation and phosphorylation in MDAMB-231 or HCT 116 cells

7x10⁵ per 6 well MDAMB-231 cells were cultured in DMEM + L-Glutamine, +Pyruvat (Gibco, 41966) 10% FCS (PAN Biotech ID-No 3302-P251116) 1% MEM Non Essentials Amino Acids (PAA, P0832100) over night. MDA-MB-231 cells were treated with 10µg/ of the different antibodies for 10 minutes and 5 hours:

5 mouse CUB4 (Deposition No. DSM ACC2551) and antibodies CDCP1_004, CDCP1_012 and CDCP1_015. HCT 116 cells were treated with 20µg/ of the different antibodies for 10 minutes and 5 hours: mouse CUB4 (Deposition No. DSM ACC2551), mouse CUB1 and mouse CUB3. Cells were lysed with ice cooled freshly prepared RIPA-Lysis buffer (Thermo Scientific, #89901) containing 10 1mM PMSF in Ethanol, 10µg/ml Aprotinin, 0,4 mM Orthovanadat). After 15 minutes on ice cell lysates were centrifuged 20 minutes at 13000 rpm. The lysates were separated on SDS-PAGE by standard protocol and transferred to nitrocellulose by Western blotting. Western Blots were detected by an anti-CDCP1 10 antibody (cell signaling #4115), an anti-phospho-CDCP1 antibody or PY 4G10 in the case of HCT116.

Phosphorylation level of CDCP1 present in untreated HCT116 cells is unchanged at time point 10 minutes or 5 hours. CUB4 mediates a stronger phosphorylation of 15 CDCP1 after 10 min incubation of the cell compared to CUB1. Inhibition of phosphorylation and down modulation (data not shown) of CDCP1 is much more pronounced by CUB4 compared to CUB1. CUB3 incubation leads to weak stimulation of CDCP1 phosphorylation and can not mediated downmodulation (data not shown) of the protein. Results are shown in Figure 5a and b.

20 Expression level of CDCP1 present in untreated MDA-MB231 cells is unchanged at time point 10 minutes or 5 hours. CUB4 incubation as well as incubation of the cells with CDCP1-004, CDCP1-012 and CDCP1-0015 antibody for 5 hours leads at least to partial degradation of the CDCP1 protein. Phosphorylation of CDCP1 is 25 hard to detect in the untreated cell. Treatment of cells for 10 min with CUB4 or the antibodies CDCP1-004, CDCP1-012 and CDCP1-0015 leads to an increase of CDCP1 phosphorylation on tyrosine 734. After 5 hour treatment with these antibodies as a consequence of the CDCP1 down modulation the phosphorylation of tyrosine 734 in CDCP1 barely to detect. Results are shown in Figure 6a to c.

Patent Claims

1. An antibody specifically binding to human CDCP1 characterized in binding to same epitope as CUB4 (Deposition No. DSM ACC2551) for the treatment of cancer.
- 5 2. The antibody according to claim 1, characterized in that
 - a) the heavy chain variable domain is SEQ ID NO: 7 and the light chain variable domain is SEQ ID NO: 8,
 - b) the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or
 - 10 c) the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or
 - d) the heavy chain variable domain is SEQ ID NO: 31 and the light chain variable domain is SEQ ID NO: 32, oror a humanized version thereof.
- 15 3. The antibody according to claims 1, characterized in that
 - a) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region of SEQ ID NO: 14, or
 - 20 b) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22, or
 - 25 c) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region

of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.

4. The antibody according to any one of claims 1 to 3, characterized in that said antibody is of human IgG1 subclass.
- 5 5. A pharmaceutical composition comprising the antibody according to any one of claims 1 to 4, for the treatment of cancer
6. Use of the antibody according to any one of claims 1 to 4, for the preparation of a medicament for the treatment of cancer
7. A method of treatment of a patient suffering from cancer by administering
10 the antibody according to any one of claims 1 to 4 to said patient in the need of such treatment.
8. An antibody specifically binding to human CDCP1 characterized in that
 - a) the heavy chain variable domain is SEQ ID NO: 15 and the light chain variable domain is SEQ ID NO: 16, or
 - 15 b) the heavy chain variable domain is SEQ ID NO: 23 and the light chain variable domain is SEQ ID NO: 24, or
 - c) the heavy chain variable domain is SEQ ID NO: 31 and the light chain variable domain is SEQ ID NO: 32, oror a humanized version thereof.
- 20 9. The antibody according to claim 8, characterized in that
 - a) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 9, a CDRH2 region of SEQ ID NO: 10, and a CDRH3 region of SEQ ID NO: 11, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 12, a CDRL2 region of SEQ ID NO: 13, and a CDRL3 region
25 of SEQ ID NO: 14, or
 - b) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 17, a CDRH2 region of SEQ ID NO: 18, and a CDRH3 region of SEQ ID NO: 19, and the light chain variable domain comprises a CDRL1 region

of SEQ ID NO: 20, a CDRL2 region of SEQ ID NO: 21, and a CDRL3 region of SEQ ID NO: 22, or

c) the heavy chain variable domain comprises a CDRH1 region of SEQ ID NO: 25, a CDRH2 region of SEQ ID NO: 26, and a CDRH3 region of SEQ ID NO: 27, and the light chain variable domain comprises a CDRL1 region of SEQ ID NO: 28, a CDRL2 region of SEQ ID NO: 29, and a CDRL3 region of SEQ ID NO: 30.

10. The antibody according to any one of claims 8 to 9, characterized in that said antibody is of human IgG1 subclass.
11. A pharmaceutical composition comprising the antibody according to claims 8 to 10.
12. Use of the antibody according to any one of claims 8 to 10, for the preparation of a medicament for the treatment of cancer
13. A method of treatment of a patient suffering from cancer by administering the antibody according to any one of claims 8 to 10 to said patient in the need of such treatment.
14. Nucleic acid encoding the antibody according to claims 8 to 10.
15. Expression vectors containing a nucleic acid according claim 14, capable of expressing said nucleic acid in a prokaryotic or eukaryotic host cell.
16. A prokaryotic or eukaryotic host cell comprising a vector according to claim 15.
17. A method for the production of a recombinant antibody according to claims 8 to 10, characterized by expressing a nucleic acid according to claim 14 in a prokaryotic or eukaryotic host cell and recovering said antibody from said cell or the cell culture supernatant.

Fig. 1




Fig. 2

Assay format (1a)




Fig. 3a




Fig. 3b




Fig 4




Figure 5a

Stimulation of the phosphorylation of CDCP1 in HCT116




Figure 5b




Figure 6a

Stimulation of CDCP1 internalisation and phosphorylation in MDAMB-231




Figure 6b





Figure 6c



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/005245

A. CLASSIFICATION OF SUBJECT MATTER
 INV. A61K39/395 A61P35/00
 ADD. C07K16/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 A61K A61P C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
 EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	JP 2007 112734 A (CHUGAI PHARMACEUTICAL CO LTD) 10 May 2007 (2007-05-10)	2-17
Y	paragraph [0006] - paragraph [0011]; claims 1-19; figure 1; examples 1-2	1-7, 12, 13
X	NEZU JUN-ICHI ET AL: "Identification of CDCP1 as a novel molecular target of anti-cancer therapeutic antibody targeting prostate cancer", PROCEEDINGS OF THE ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH; 98TH ANNUAL MEETING OF THE AMERICAN-ASSOCIATION-FOR- CANCER-RESEARCH, NEW YORK, NY, US; LOS ANGELES, CA, USA, vol. 48, 1 April 2007 (2007-04-01), page 153, XP008115766, ISSN: 0197-016X * abstract	1-17

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

<p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>	<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>
--	--

Date of the actual completion of the international search	Date of mailing of the international search report
12 November 2010	19/11/2010

Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Siaterli, Maria
--	---

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2010/005245

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SIVA AMARA C ET AL: "Targeting CUB domain-containing protein 1 with a monoclonal antibody inhibits metastasis in a prostate cancer model.", CANCER RESEARCH 15 MAY 2008, vol. 68, no. 10, 15 May 2008 (2008-05-15), pages 3759-3766, XP002559247, ISSN: 1538-7445 page 3759, right-hand column, paragraph 2 page 3760, right-hand column, paragraph 3 - paragraph 4; figures 4,5 page 3763, left-hand column, paragraph 3 - paragraph 4	1-17
X	WO 2007/005502 A2 (APPLERA CORP [US]; DOMON BRUNO [US]; BIRSE CHARLES E [US]; LEE CANDY []) 11 January 2007 (2007-01-11) paragraph [0308] - paragraph [0332] paragraph [0357] - paragraph [0377] paragraph [0382]; figures 2,11-13	1-17
X	EP 1 396 501 A1 (UNIV EBERHARD KARLS [DE]) 10 March 2004 (2004-03-10) cited in the application	8-11, 14-17
Y	paragraph [0161] - paragraph [0163]; claims 1-6	1-7,12, 13
X	WO 02/04504 A1 (BIOWINDOW GENE DEV INC [CN]; MAO YUMIN [CN]; XIE YI [CN]) 17 January 2002 (2002-01-17) claim 9	1-17
X	WO 2004/074481 A1 (NOVARTIS AG [CH]; NOVARTIS PHARMA GMBH [AT]; SCRIPPS RESEARCH INST [US]) 2 September 2004 (2004-09-02) example XII	1-17
X	WO 2008/133851 A1 (ALEXION PHARMA INC [US]; BOWDISH KATHERINE S [US]; XIN HONG [US]; YANT) 6 November 2008 (2008-11-06) claims 1-48; examples 4,7-10	1-17

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2010/005245

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-7(partially)

antibody CUB4, with a VH of SEQ ID NO 7 and a VL of SEQ ID NO 8, or humanized versions thereof for the treatment of cancer

2. claims: 1-17(partially)

antibody CDCP1-012 with a VH of SEQ ID NO 23 and a VL of SEQ ID NO 24 or or comprising the CDR's with SEQ ID NO 17-22 & humanized versions thereof,

3. claims: 1-17(partially)

antibody CDCP1-015 with a VH of SEQ ID NO 31 and a VL of SEQ ID NO 32 or comprising the CDR's with SEQ ID NO 25-30 & humanized versions thereof

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2010/005245

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
JP 2007112734	A	10-05-2007	NONE	
WO 2007005502	A2	11-01-2007	NONE	
EP 1396501	A1	10-03-2004	AT 462728 T	15-04-2010
			DE 10242146 A1	18-03-2004
			DK 1396501 T3	21-06-2010
			EP 2206729 A1	14-07-2010
			ES 2344982 T3	13-09-2010
			PT 1396501 E	17-05-2010
			SI 1396501 T1	30-07-2010
			US 2009181033 A1	16-07-2009
			US 2004053343 A1	18-03-2004
			US 2010179306 A1	15-07-2010
WO 0204504	A1	17-01-2002	AU 9362201 A	21-01-2002
			CN 1329061 A	02-01-2002
WO 2004074481	A1	02-09-2004	AT 473279 T	15-07-2010
			AU 2004213568 A1	02-09-2004
			BR PI0407680 A	01-03-2006
			CA 2516366 A1	02-09-2004
			CN 1761750 A	19-04-2006
			EP 1597367 A1	23-11-2005
			JP 2007523836 T	23-08-2007
			MX PA05008794 A	10-03-2006
			PT 1597367 E	02-09-2010
WO 2008133851	A1	06-11-2008	US 2008008719 A1	10-01-2008