



US00PP29645P3

(12) **United States Plant Patent**
Randag

(10) **Patent No.:** **US PP29,645 P3**

(45) **Date of Patent:** **Sep. 4, 2018**

(54) **CALLA LILY PLANT NAMED ‘NEW YORK’**

(50) Latin Name: *Zantedeschia sprengeri*
Varietal Denomination: **New York**

(71) Applicant: **Sande B.V.**, 't Zand (NL)

(72) Inventor: **Cecilius Jan-Jochem Randag**, 't Zand (NL)

(73) Assignee: **Sande B.V.**, 't Zand (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 44 days.

(21) Appl. No.: **15/530,234**

(22) Filed: **Dec. 14, 2016**

(65) **Prior Publication Data**
US 2017/0181363 P1 Jun. 22, 2017

Related U.S. Application Data
(60) Provisional application No. 62/386,953, filed on Dec. 17, 2015.

(51) **Int. Cl.**
A01H 5/02 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./490**

(58) **Field of Classification Search**
USPC Plt./263.1, 490
See application file for complete search history.

Primary Examiner — Susan McCormick Ewoldt
Assistant Examiner — Karen M Redden
(74) *Attorney, Agent, or Firm* — The Webb Law Firm

(57) **ABSTRACT**

‘New York’ is a new variety of calla lily having inflorescences with a pink-colored spathe that produces 5-25 inflorescences per tuber. The inflorescence stems are about 27-47 cm in height, and the leaves are green with whitish spots and have a leathery texture.

3 Drawing Sheets

1

Botanical classification: *Zantedeschia sprengeri*.
Varietal denomination: ‘NEW YORK’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct cultivar of calla lily, botanically known as *Zantedeschia sprengeri* and hereinafter referred to by the cultivar name ‘New York’.

‘New York’ is a product of a planned breeding program which had the objective of creating *Zantedeschia* hybrids for pot flower production in a wide range of colors with a large, classic flower shape. The breeding program began in 1989, and the new cultivar is a seedling selected from the crossing of a *Zantedeschia sprengeri* selection referred to as Z060032-937 (female parent, unpatented) with a *Zantedeschia sprengeri* selection referred to as Z071598-949 (male parent, unpatented). ‘New York’ was selected in 2011 by the inventor in ‘t Zand, The Netherlands as one flowering plant within the progeny of the stated cross.

The first act of asexual reproduction of ‘New York’ by tissue culture was performed by the inventor in September of 2011 in ‘t Zand, The Netherlands. Subsequent asexual reproductions by tissue culture at the same location have demonstrated that the combination of characteristics as herein disclosed for the new cultivar are retained and reproduced true to type through successive generations of asexual reproduction.

The following observations, measurements and comparisons describe plants grown in 12-18 cm pots at 12-14 weeks after planting in ‘t Zand, The Netherlands under greenhouse conditions, which approximate those generally used in horticultural practice. Color references are made to The 2001 R.H.S. Colour Chart of The Royal Horticultural Society of London, except where general color terms of ordinary significance are used.

2

The present invention has not been evaluated under all possible environmental conditions. The phenotype may vary with variations in environment without a change in the genotype of the plant.

The following traits have been repeatedly observed and determined to be basic characteristics of ‘New York’ which, in combination, distinguish this calla lily as a new and distinct cultivar:

- 1. Pink-colored spathes;
- 2. Compact plant architecture; and
- 3. High inflorescence production.

Further, when compared to calla lily plant named ‘Paco’ (U.S. Plant Pat. No. 24,246), the upper surface and lower surface spathe color of ‘Paco’ varies between Purple Group RHS N79B and N79D, the upper surface spathe color of ‘New York’ varies between Red-Purple Group RHS N57A and 58B and the lower surface spathe color of ‘New York’ varies between Red-Purple Group RHS 58B and 63A.

Table 1 provides differences of ‘New York’ to its parents, *Zantedeschia sprengeri* selections Z060032-937 and Z071598-949.

TABLE 1

‘New York’	Differences
Z060032-937	Spathe color Varies between Red-Purple Group RHS 58A to 58C Plant height 35-60 cm
Z071598-949	Spathe color Varies between Red-Purple Group RHS 68A to 71D Plant height 25-40 cm

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying photographic drawings illustrate the new variety at 12-14 weeks after planting, with the colors being as nearly true as is possible with color illustrations of this type:

FIG. 1 is a photograph of a group of potted plants of the new variety;

FIG. 2 is a photograph of a close-up view of the new variety; and

FIG. 3 is a photograph of a side view of the new variety. 5

DESCRIPTION OF THE PLANT

The Plant

Size:

Height of the leaf canopy above the soil.—20-40 cm.

Height of top of inflorescence above the soil.—24-47 cm.

Diameter.—8-12 cm.

Form: Erect.

Number of inflorescences per tuber:

Size 14-16 cm in diameter.—5-7 inflorescences.

Size 16-18 cm in diameter.—7-11 inflorescences.

Size 18-20 cm in diameter.—12-18 inflorescences.

Size 20-22 cm in diameter.—15-25 inflorescences. 10

Branches:

Number.—From 3-8.

Color.—Yellow-Green Group RHS 144A.

Leaves:

Size.—Width: 6-11 cm. Length: 12-24 cm.

Number per plant.—27-39.

Shape.—Lanceolate.

Color.—Upper surface: Yellow-Green Group RHS 146A to 147A, with transparent White Group RHS 155A and N155A spots. Lower surface: Yellow-Green Group RHS 146A to 147B, with transparent White Group RHS 155A and N155A spots. 15

Apex.—Apiculate.

Base.—Decurrent.

Margin.—Undulate; having a color of Yellow-Green Group RHS 146A.

Veins.—Configuration: Pinnate. Color: Yellow-Green Group RHS 146A to 146B.

Surface quality.—Leathery.

Petiole.—Length: 8-16 cm. Average diameter: 10 mm.

Color: The base is White Group RHS 155A and the upper part is Green Group RHS 143A to 143B. 20

Roots:

Color.—White.

Branching.—Similar to other *Zantedeschia* varieties.

The Inflorescence and Seeds

Spathe:

Size.—Length (measured from base opening to tip): 6.5-9 cm. Width: 3-7 cm. Height: 7-10 cm.

Color.—Upper surface: Varies between Red-Purple Group RHS N57A and 58B. Lower surface: Varies between Red-Purple Group RHS 58B and 63A.

Vein color.—Varies between Red-Purple Group RHS 59A and 59B.

Shape.—Cupped.

Apex.—Apiculate.

Base.—Obtuse.

Texture.—Smooth (upper and lower surfaces).

Spadix:

Size.—Length: 25-46 mm. Diameter: 5-9 mm.

Color.—Yellow Group RHS 12A.

Position relative to spathe.—Upright.

Peduncle:

Size.—Length: 20-37 cm. Diameter: 6-8 mm.

Color.—The upper part is Yellow-Green Group RHS 144A, and the lower part is Green-White Group RHS 157C.

Texture.—Smooth.

Reproductive organs:

Location of female organs.—Basal position of the spadix.

Location of male organs.—Upper position of the spadix.

Perianth.—Conspicuous.

Stamens.—Presence: Not visible before pollen release.

Number: More than 20. Pollen amount: Abundant and similar to 'Picasso' (U.S. Plant Pat. No. 15,282), but not quantifiable due to minute pollen size. Color: Between White Group RHS 155A and 155B.

Anther.—Shape: Round. Length: Less than 1 mm. Color: Yellow Group RHS 12A.

Filament.—Length: Less than 1 mm. Color: Between White Group RHS 155A and 155B.

Pistil.—Number: 15 to 30. Length beyond perianth: About 1 mm.

Stigma.—Shape: Round. Size: Less than 1 mm.

Style.—Length: Less than 1 mm. Color: Between Red-Purple Group RHS 58A and Greyed-Purple Group RHS 186B.

Ovaries.—Texture: Smooth. Length: Between 1 to 10 mm. Width: Between 1 to 10 mm. Color: Top color is between Yellow Group RHS 6A and 7A, bottom color is from White Group RHS 155B to Green-White Group RHS 157C. 25

Seeds:

Length.—4-7 mm.

Width.—3-6 mm.

Color.—Between Yellow-Green Group RHS 150A and 151A.

Amount.—Between 1-60.

General

Disease resistance: No unusual susceptibility to diseases noticed to date.

Pest resistance: No unusual susceptibility to pests noticed to date.

Heat tolerance: No heat tolerance.

Hardiness: Not hardy.

Flowering: Tubers planted in a greenhouse in The Netherlands in February produce inflorescences in April. Tubers planted outdoors in The Netherlands in May produce inflorescences in July. In general, the time from planting to first flowering takes eight weeks. However, tubers stored for a longer period before planting may flower earlier—up to five weeks after planting.

Lastingness: About two weeks on the plant.

Fragrance: None.

I claim:

1. A new and distinct variety of calla lily plant named 'New York' as herein described and illustrated. 30

* * * * *



Fig. 1




Fig. 2



Fig. 3