



(72) GANEMO, AGNETA, SE

(71) YAMANOUCHI EUROPE B.V., NL

(51) Int.Cl.⁶ A61K 31/19, A61K 9/06, A61K 31/045

(30) 1997/11/25 (97203681.8) EP

(54) **UTILISATION D'UN MELANGE DE DIOL ET D'ACIDE ALPHA-HYDROXYLE POUR TRAITER DES HYPERKERATOSES**

(54) **USE OF A MIXTURE OF A DIOL AND AN ALPHA-HYDROXY ACID FOR THE TREATMENT OF HYPERKERATOTIC SKIN DISEASES**

(57) L'invention concerne l'utilisation de compositions contenant un mélange constitué par un diol et par un acide alpha-hydroxylé dans un véhicule semi-occlusif pour le traitement localisé d'hyperkératoses.

(57) The use of compositions containing a mixture of a diol and an alpha-hydroxy acid in a semi-occluding vehicle for the topical treatment of hyperkeratotic skin diseases is provided.

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/19, 9/06 // (A61K 31/19, 31:045)	A1	(11) International Publication Number: WO 99/26617 (43) International Publication Date: 3 June 1999 (03.06.99)
(21) International Application Number: PCT/EP98/07759 (22) International Filing Date: 25 November 1998 (25.11.98) (30) Priority Data: 97203681.8 25 November 1997 (25.11.97) EP (71) Applicant (for all designated States except US): YAMANOUCI EUROPE B.V. [NL/NL]; Elisabethhof 19, P.O. Box 108, NL-2350 AC Leiderdorp (NL). (72) Inventor; and (75) Inventor/Applicant (for US only): GÅNEMO, Agneta [SE/SE]; Gånarp 2517, S-266 92 Munka Ljungby (SE). (74) Common Representative: YAMANOUCI EUROPE B.V.; Dept. of Legal Affairs, Patents & Trademarks, Elisabethhof 19, P.O. Box 108, NL-2350 AC Leiderdorp (NL).	(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: USE OF A MIXTURE OF A DIOL AND AN ALPHA-HYDROXY ACID FOR THE TREATMENT OF HYPERKERATOTIC SKIN DISEASES		
(57) Abstract The use of compositions containing a mixture of a diol and an alpha-hydroxy acid in a semi-occluding vehicle for the topical treatment of hyperkeratotic skin diseases is provided.		

5 **USE OF A MIXTURE OF A DIOL AND AN ALPHA-HYDROXY ACID FOR THE
TREATMENT OF HYPERKERATOTIC SKIN DISEASES**

The present invention relates to the use of a combination of a diol and an alpha-hydroxy acid for the topical treatment of hyperkeratotic skin diseases.

10

BACKGROUND OF THE INVENTION

Chronic hyperkeratotic skin diseases are characterised by a disturbance in the keratinisation process and include among other diseases psoriasis, hyperkeratotic eczema and ichthyosis.

15

Psoriasis is characterised by reddened flaking skin lesions. The disease, has as a rule, a chronic course. Topical agents for the treatment thereof are often smeary and/or discolouring, e.g. dithranol and tar. Steroid preparations are cosmetically attractive but have important side-effects. Emollient preparations containing e.g. salicylic acid and urea in a cream or ointment base have a certain peeling effect but are not separately sufficiently active.

20

Hyperkeratotic eczema is characterised by thickened scaling skin often with chaps (rhagades).

Lamellar ichthyosis is a rare, congenital genodermatosis characterised by generalised hyperkeratosis, intense dryness of the skin, and large scales especially over the extremities (H. Traupe, ed. (1989) *The Ichthyoses: A guide to clinical diagnosis, genetic counseling and therapy*, 1st ed., Springer Verlag, Berlin). Though usually not life-threatening, the disease can be very disfiguring and causes considerable distress to the sufferers throughout life. The treatment of lamellar ichthyosis involves oral vitamin A derivatives (retinoids) combined with adjuvant topical treatment with different emollient formulations containing e.g. urea, propylene glycol and alpha-hydroxy acids (AHA). However, the therapeutic results, like in other hyperkeratotic skin diseases, are often disappointing and associated with side-effects.

25

30

E.J. Van Scott et al. (1974) *Arch. Dermatol.* 110: 586-590 pioneered in using alpha-hydroxy acids against ichthyotic conditions. Further E.J. Van Scott et al. (1984) *J. Am. Acad. Dermatol.* 11: 867-879 described a keratolytic gel (5-10% alpha-hydroxy acid in a

35

vehicle consisting of water, ethanol and propylene glycol (4:4:2), which however fell in
5 oblivion since it was irritating and not suitable for whole body application.

In EP-0292495 compositions, consisting essentially of a mixture of propylene
glycol, urea and optionally lactic acid, were disclosed. The compositions after topical
application to the skin were said to have a beneficial effect on hyperkeratotic skin diseases.
Simple application of the said mixture to the skin is possible, but in practice proved to be
10 very difficult to achieve effectively. The mixture is fluid and slow to dry (evaporate) and the
use of plasters and/or bandages is deemed necessary. This procedure is considered to be too
inconvenient for general use. Besides that, due to the high concentrations of propylene
glycol (40-80%) and urea (5-20%), side-effects, such as irritation, may occur, which also are
a serious draw-back for daily application of such preparations during a prolonged period of
15 time.

US Patent 4,105,783 discloses compositions containing a product, prepared by
reacting, in aqueous or alcoholic aqueous solution, one or more of an alpha-or a beta-
hydroxy acid and a base selected from the group consisting of ammonium hydroxide and an
organic alkylamine, in a total amount of from 1 to 20% for the topical treatment of dry skin
20 disorders, such as psoriasis and ichthyosis. The bases are successfully used to raise the pH of
the compositions containing the hydroxy acids without compromising the therapeutic
activity of the active ingredients, thereby reducing the skin irritation. Although it was stated
that the reaction products, viz. the ammonium salts and amides, thus formed, need no
isolation procedure and can be directly incorporated into the therapeutic composition, good
25 manufacturing and clinical practices nowadays require a careful assessment of the active
ingredient both qualitatively and quantitatively and extensive pharmacological and
toxicological testing of the reaction product formed.

There thus still exists a need for an effective product based on well-known and well-
defined active ingredients or a method which can be topically applied to hyperkeratotic skin
30 without causing serious side-effects, such as irritation.

SUMMARY OF THE INVENTION

5

The object of the present invention is to provide the use of compositions containing a combination of a diol and an alpha-hydroxy acid in a vehicle, which has semi-occluding properties, for topical treatment of hyperkeratotic skin diseases.

10

DETAILED DESCRIPTION OF THE INVENTION

It has now been found that compositions containing a combination of a diol and an alpha-hydroxy acid in a semi-occluding vehicle on topical application to the skin of patients, having a hyperkeratotic skin disease, are effective and show a considerable reduction of side-effects, such as irritation, as compared with compositions, containing the same amounts of the diol and the alpha-hydroxy acid in a non-occluding vehicle, such as a gel. In addition thereto the compositions of the invention show a remarkably greater efficacy than compositions comprising either the diol or the alpha-hydroxy acid in the same amount.

Examples of diols which may be used are propylene glycol, butylene glycol, pentanediol and hexylene glycol. There is a preference for propylene glycol and hexylene glycol, which may be used in a concentration of up to 40% and preferably from 10-20%.

Examples of the alpha-hydroxy acids or derivatives thereof are lactic acid, citric acid, glycolic acid, glucuronic acid, galacturonic acid, pyruvic acid etc, but preferably lactic acid is used. The alpha-hydroxy acid may be used in an amount of up to 10%, but preferably up to 5% is used.

The semi-occluding vehicles to be used are characterised by the fact that in in vitro and in vivo tests an occluding activity is observed, which is lower than for white soft paraffin, which is considered to be an occluding vehicle. The lower limit for the occluding activity is determined by values which are obtained for preparations, known to be non-occluding. E.g. in the in vitro test as disclosed in example 7 preparations having an occlusion factor below about 70 are considered to be non-occluding preparations. A proper in vivo test, wherein the effectiveness of topical preparations in restoring a cutaneous barrier function of diseased skin is assessed, is disclosed in example 8. Examples of such semi-occluding vehicles are fatty creams, containing a certain amount of occluding fats, such as

white soft paraffin, or creams, containing an aqueous suspension of solid lipid nanoparticles,
5 e.g. wherein the solid lipid is hard paraffin, having a melting point range from 54 to 57°C.

The combination of only 5% lactic acid and 20% propylene glycol in a fatty cream
base is readily acceptable by most patients and has proved to be much more effective and
less irritating than monotherapy with either of the two compounds in the same or even at
twice these concentrations. The astonishingly potent effect of the mixture of the diol and the
10 alpha-hydroxy acid in the semi-occluding vehicle suggests that the ingredients act
synergistically in reverting hyperkeratosis.

The compositions according to the invention can be used in several stages of the
treatment of hyperkeratotic skin diseases. In the pretreatment of psoriatic plaques before
initiation of treatment with e.g. corticosteroids the compositions have proven to be equally
15 good as the standard product, which is 5% of salicylic acid in white soft paraffin. However,
the cosmetic properties of the compositions according to the invention were considered by
the patients to be superior over those of the standard product. The treatment of lamellar
ichthyosis normally consists of systemic therapy with retinoids combined with adjuvant
topical treatment. The compositions according to the invention have proven to be
20 advantageously used in such adjuvant topical treatment. For less severe hyperkeratotic skin
disorders the compositions according to the invention can be used without systemic therapy.

Although the foregoing invention has been described in some detail by way of
illustration and example for purposes of clarity and understanding, it will be readily apparent
to those of ordinary skill in the art in the light of the teachings of this invention that certain
25 changes and modifications may be made thereto without departing from the spirit and the
scope of the appended claims.

The following examples further illustrate the invention.

5

EXAMPLES

5

Example 1

lactic acid	5%
propylene glycol	20%
fatty cream base*	75%

10

*: The fatty cream base consists of 3% polyoxyethylene 1000 monocetyl ether (Cetomacrogol® 1000), 6% cetostearyl alcohol, 18% liquid paraffin, 42% white soft paraffin, preservatives q.s., buffers q.s. and water up to 100%.

15

Example 2

lactic acid	5%
propylene glycol	20%
nanoparticles suspension*	75%

20

*: The nanoparticles suspension consists of 30% solid paraffin (melting point range 54-57°C), 5% polyoxyethylene 1000 monocetyl ether and 65% water.

Example 3

25

lactic acid	5%
propylene glycol	20%
nanoparticles cream*	75%

30

*: The nanoparticles cream consists of 2% cetyl dimethicone copolyol, 10% isopropyl myristate, 16.67% cyclomethicone, 0.33% sodium chloride, 10% solid paraffin, 1.67 polyoxyethylene 1000 monocetyl ether, buffers q.s., preservatives q.s. and water up to 100%.

Example 4

5	lactic acid	5%
	propylene glycol	20%
	cream*	65%

10 *: The cream consists of 32.5% cetostearyl alcohol, 5% polyoxyethylene 1000 monocetyl ether and 62.5% water.

Example 5

	lactic acid	5%
	propylene glycol	20%
15	nanoparticles suspension/ cream (2.5/1)*	65%

20 *: The vehicle consists of a mixture of the nanoparticles suspension of example 2 and the cream of example 4.

Example 6

25 Ten patients (7 females; age 2-38 years) with severe lamellar ichthyosis, two of whom had ongoing treatment with oral acitretin (50-75 mg/day) were recruited after informed consent. The patients were instructed to apply the formulation according to example 1 twice daily to the legs and refrain from any other topical medication at that side. Signs of scaling, hyperkeratosis and xerosis were separately scored on a scale ranging from 0 (feature not present) to 4 (feature present to a large extent) and the sums of the scores for these features (max.: 12) were calculated for each patient before and after changing therapy. Use of the formulation according to example 1 resulted after 1 month in a reduction of the mean total score from 9.6 ± 2.2 to 3.5 ± 2.3 ($p < 0.001$; paired t-test). All patients reported increased shredding of scales after a few days of treatment, followed within 1-2 weeks by the appearance of a much smoother, almost normal looking skin. Nine patients considered the effect "better than ever obtained before" and later choose to extend the treatment to the rest of the body with equally good

30

results. Hence one patient was able to reduce her acitretin dosage from 75 mg to 25 mg
 5 daily without deterioration of skin symptoms. Some patients experienced a transient skin
 irritation after applying the cream. No other side effects were noted.

Example 7

A vessel in the form of a beaker was used. The vessel had a diameter of 5.5. cm
 10 and a height of 7 cm, and had been designed to receive on top a closing standard
 laboratory paper filter (TVN, sold by Schut, the Netherlands, surface 23.8 cm²). The test
 was performed by placing 50 g of distilled water in the vessel, closing the vessel with the
 paper filter on the upper surface of which 200 mg of the preparation to be tested were
 evenly distributed, and placing the closed vessel for a period of 72 hours in a stove at
 15 33°C and 58% RH. All other conditions having been kept equal, the weight loss of water
 from the vessel (water flux) after 72 hours exclusively depended on the occlusivity of the
 preparation tested.

The occlusion factor F of the tested preparation was calculated according to the
 equation:

$$20 \quad F = 100 \{(A-B)/A\}$$

wherein A is the water flux through the uncovered filter and B the water flux through the
 filter when covered by the tested preparation.

All preparations were tested in triplicate, the maximal deviation between the
 results of one preparation being 10%. The following table presents the means of the
 25 occlusion factors F found.

Vehicle of example	Occlusion Factor F
1	91.7
2	78.5
4	73.0
5	87.0

Example 8

5 Hospitalised female patients (8) with chronic constitutional eczema were included
in a study to assess the effectiveness of various topical preparations in restoring a
cutaneous barrier function by means of Trans Epidermal Water Loss measurements. The
flexor aspect of the right forearm was used throughout the study. The included sites (area
in between a distance of about 5 cm from wrist to elbow) were examined to be classified
10 as light, moderate or severely affected. The patients were not allowed to use any topical
medication 24 hours before commencement of the study. The experiments were
performed in a separate room with constant humidity and temperature. These parameters
were checked before the start of the experiments. The patients were quietly seated for at
least 10 minutes prior to the start of the measurements. The measurements were carried
15 out using a Servo Med EP1 Evaporimeter (Servo Med, Stockholm, Sweden), calibrated
according to the manufacturers instructions.

6 areas of about 3x3 cm were indicated on the forearm of each patient. 0.5 ml of the
preparations were applied on the indicated spots For each patient the preparations were
applied at the same spot. After massaging for 0.5 minutes with a fingertip and a period of
20 2 minutes, the remaining of the preparations was discarded with a wooden scraper
followed by gently wiping with a piece of tissue. The sixth spot was left untreated to
allow control measurements. Initial TEWL-values were determined on each spot before
application of the dressings (t=0 values). Accordingly, the indicated spots were measured
at 0.5, 1, 1.5, 2; 3; 4; 6 and 8 hours after treatment. All measurements were conducted in
25 triplicate.

The results showed that white soft paraffin is always the best preparation with respect to
prevention of TEWL and can be considered as an occluding preparation. The vehicle of
example 1 resulted in a lower normalisation of the skin barrier function than white soft
paraffin, but the preparation can be considered to be semi-occluding.

CLAIM

5

1. Use of a combination of a diol and an alpha-hydroxy acid in a vehicle for the preparation of a medicament for the topical treatment of hyperkeratotic skin diseases, characterised in that the vehicle has semi-occluding properties.

10