



(43) International Publication Date
24 December 2014 (24.12.2014)

(51) International Patent Classification:

A61K 36/75 (2006.01) A61K 36/487 (2006.01)
A61K 36/12 (2006.01) A61K 36/539 (2006.01)
A61K 36/185 (2006.01) A61K 36/605 (2006.01)
A61K 36/19 (2006.01) A61K 36/62 (2006.01)
A61K 36/23 (2006.01) A61K 36/73 (2006.01)
A61K 36/232 (2006.01) A61P 3/04 (2006.01)
A61K 36/38 (2006.01) A61P 3/10 (2006.01)
A61K 36/44 (2006.01)

(21) International Application Number:

PCT/US2014/042534

(22) International Filing Date:

16 June 2014 (16.06.2014)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/836,843 19 June 2013 (19.06.2013) US

(71) Applicants: ACCESS BUSINESS GROUP INTERNATIONAL LLC [US/US]; 7575 Fulton Street East, Ada, MI 49355 (US). THE REGENTS OF THE UNIVERSITY OF COLORADO [US/US]; 1800 Grant Street, 8th Floor, Denver, CO 80203 (US).

(72) Inventors: RANA, Jatinder; 5708 Tall Timber Street SE, Grand Rapids, MI 49546 (US). RANDOLPH, Russell, Keith; 6909 E. Shorecrest Drive, Anaheim, CA 92807 (US). SCHOLTEN, Jeffrey; 476 Crahen Ave SE, Grand Rapids, MI 49525 (US). LE, Myphuong, Thi; 5353

Laredo Street, Denver, CO 80239 (US). JOHNSON, Richard, J.; 15857 E Aberdeen Avenue, Centennial, CO 80016 (US). LANASPA GARCIA, Miguel, Angel; 2908 Ironton Street, Denver, CO 80238 (US).

(74) Agent: CILELLA, Magdalena, O.; Brinks Gilson & Li-one, P.O. Box 10087, Chicago, IL 60610 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: PLANT-BASED INHIBITORS OF KETOHEXOKINASE FOR THE SUPPORT OF WEIGHT MANAGEMENT

Overview of Extraction Process




FIG. 1

(57) Abstract: A composition for inhibiting ketohexokinase, for example, ketohexokinase-C (KHK-C) activity, may include a plant extract exhibiting at least IC50 (i.e., 50% KHK-C inhibition at a concentration in the range of from about 0.1 µg/mL to about 1000 µg/mL. The composition may be in a form suitable for oral ingestion. A method for inhibiting KHK-C activity in a subject may include administering a plant extract that exhibits at least 50% KHK-C inhibition at a concentration from about 0.1 µg/mL to about 1000 µg/mL. The administering may be done to treat or prevent at least one of sugar addiction, obesity, or metabolic syndrome. The administering may be done to provide a diminished craving in the subject from at least one member selected from the group consisting of craving of sugar, fructose, fructose-containing sugars, carbohydrates, and combinations thereof. The subject may be pre-diabetic, diabetic and or insulin resistant.

WO 2014/204853 A1


PLANT-BASED INHIBITORS OF KETOHEXOKINASE FOR THE SUPPORT OF WEIGHT MANAGEMENT

Background

[0001] The present disclosure relates generally to inhibitors of ketohexokinase and, more particularly, to plant-based inhibitors of ketohexokinase and the use of such plant-based inhibitors for the support of weight management.

[0002] The intake of added sugars, especially sucrose and high fructose corn syrup (HFCS), has increased markedly over the last century in developed countries around the world. Epidemiological studies strongly associate the consumption of dietary sugar with the incidence of metabolic syndrome and obesity. Experimentally, the administration of fructose to rats has been shown to induce all features of metabolic syndrome, weight gain, and increased body fat.

[0003] Ketohexokinase (KHK) is an enzyme found in the liver, the renal cortex, and the small intestine that is involved in the metabolism of fructose in the body. KHK catalyzes the phosphorylation of fructose by adenosine triphosphate (ATP) to produce fructose-1-phosphate and adenosine diphosphate (ADP) according to the following reaction:



Fructose-1-phosphate is then metabolized by aldolase B to generate various substrates. The phosphorylation of fructose consumes ATP and generates ADP.

[0004] Fructose is distinct from other sugars in that it causes transient intracellular ATP depletion in the liver prior to generating energy. This occurs with regularly ingested oral doses of fructose, even in humans. The mechanism may be due to the rapid phosphorylation of fructose by KHK. It is believed that such rapid phosphorylation of fructose is possible because KHK does not have a negative feedback system like hexokinases (e.g., glucokinase), which catalyze the phosphorylation of hexoses (e.g., glucose). KHK consumes ATP rapidly, resulting in activation of adenosine monophosphate (AMP) deaminase and the generation of uric acid, which increases in both hepatocytes and transiently in the circulation. ATP depletion by KHK is critical for fatty liver formation.

Summary

[0005] In one example, a composition for inhibiting ketohexokinase, for example, for inhibiting ketohexokinase-C (KHK-C) activity, may include a plant extract exhibiting at least IC₅₀ (i.e., at least 50% KHK-C inhibition at a concentration from about 0.1 µg/mL to about 1000 µg/mL). The plant extract may be obtained from a plant from a genus selected from the group consisting of *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, *Diospyros*, *Andrographis*, *Nymphaea*, *Chloroxylon*, *Petroselinum*, *Morus*, *Pteris*, *Garcinia*, and *Malus*. The plant extract may be obtained from a plant selected from the group consisting of *Angelica archangelica*, *Cratoxylum prunifolium*, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*, *Andrographis paniculata*, *Nymphaea lotus*, *Chloroxylon swietenia*, *Petroselinum crispum*, *Morus alba*, *Pteris wallichiana*, *Garcinia mangostana*, and *Malus domestica*. The plant extract may include a compound selected from the group consisting of Osthol, Cratoxyarborenone E, gamma-Mangostin, Osthenol, a Polyketide type molecule, 4-Hydroxy-Derricin, Isobavachalcone, Methoxy-isobavachalcone, Oroxylin A, 5,7-Dimethoxy-8-prenylcoumarin, Apigenin 7-glucuronide, 3',4',5,7-THMethoxy3'-O-β-D-Xylopyranoside, Swietenocoumarin B, Apiin, Mulberrin, Flavaspidic acid AB, Mangostin, Phloretin, and combinations thereof. The composition may be in a form suitable for oral ingestion.

[0006] In another example, a method for inhibiting KHK-C activity in a subject may include administering a plant extract that exhibits at least IC₅₀ (i.e., 50% KHK-C inhibition at a concentration from about 0.1 µg/mL to about 1000 µg/mL). The administering may be done to treat or prevent at least one of sugar addiction, obesity, or metabolic syndrome. The administering may be done to provide a diminished craving in the subject from at least one member selected from the group consisting of sugar, fructose, fructose-containing sugars, carbohydrates, and combinations thereof. The subject may be pre-diabetic, diabetic and/or insulin resistant.

[0007] These and other features and advantages of the invention will become apparent upon consideration of the following detailed description of the presently preferred embodiments, viewed in conjunction with the appended drawings.

Brief Description of the Drawings

[0008] Figure 1 depicts an overview of the extraction process described in the application;

[0009] Figure 2 depicts an HPLC fingerprint profile for *Angelica archangelica* (Wild Celery);

[0010] Figure 3 depicts an HPLC fingerprint profile for *Myrica cerifera* (Bayberry);

[0011] Figure 4 depicts an HPLC fingerprint profile for *Scutellaria baicalensis* (Skullcap);

[0012] Figure 5 depicts an HPLC fingerprint profile for *Petroselinum crispum* (Garden Parsley);

[0013] Figure 6 depicts an HPLC fingerprint profile for *Garcinia mangostana* (Mangosteen);

[0014] Figure 7 depicts an HPLC fingerprint profile for *Psoralea corylifolia* (Malay tea); and

[0015] Figure 8 depicts an HPLC fingerprint profile for *Morus alba* (Mulberry).

Detailed Description

[0016] Throughout this disclosure, the terms “ketoheokinase” (KHK) and “fructokinase” may be used interchangeably and may refer to ketoheokinase-C (KHK-C), ketoheokinase-A (KHK-A), or combinations thereof.

[0017] As used herein, the term “administration” of a compound refers to introducing or delivering the compound to a subject to perform its intended function. The administration may be carried out by any suitable route such as

orally, intranasally, parenterally (intravenously, intramuscularly, intraperitoneally, or subcutaneously), rectally, or topically.

[0018] As used herein, the term “effective amount” or “therapeutically effective amount” refers to an amount effective at dosages and for periods of time sufficient to achieve a desired result.

[0019] As used herein, the term “subject” refers to any animal (e.g., a mammal) including, but not limited to, humans, non-human primates, rodents, and the like, to which a compound may be administered.

[0020] As used herein, the term “carrier” refers to a composition that aids in maintaining one or more plant extracts in a soluble and homogeneous state in a form suitable for administration, which is nontoxic and which does not interact with other components in a deleterious manner.

[0021] Unless indicated otherwise, all proportions and percentages recited throughout this disclosure are by weight.

[0022] A KHK-C inhibitor may be administered to a subject to inhibit KHK-C activity within the body. Such inhibition of KHK-C activity may effectively reduce the metabolism or absorption of fructose within the body. The fructose present within the body may be derived from fructose-containing sugars (e.g., fructose, sucrose, or high fructose corn syrup), sugars that can be converted to fructose within the body (e.g., sorbitol), glucose, carbohydrates (e.g., starches), or any other source. Absorption of fructose and increased KHK-C activity may contribute to a variety of conditions (e.g., obesity, metabolic syndrome, renal disease, pre-diabetes, diabetes, adenosine triphosphate (ATP) depletion, monocyte chemotactic protein-1 (MCP-1) production, insulin resistance, or intrarenal uric acid production). Inhibition of KHK-C activity may be beneficial for supporting weight management (e.g., by reducing the absorption of fructose and the associated caloric intake).

[0023] Inhibition of KHK-C activity may effectively reduce the craving for fructose from any source. A craving for fructose may result in repeated sugar intake, which may contribute to obesity, metabolic syndrome, or other conditions. Reducing the craving for fructose may be beneficial for supporting weight

management (e.g., by reducing the consumption of fructose and the associated caloric intake).

[0024] In one example, a composition for inhibiting KHK-C activity may include a plant extract. The plant extract may exhibit at least IC₅₀ (i.e., at least 50% KHK-C inhibition at a concentration from about 0.1 µg/mL to about 1000 µg/mL). In another example, the plant extract may exhibit at least 50% KHK-C inhibition at a concentration of less than about 50 µg/mL. Alternatively, the plant extract may exhibit at least 50% KHK-C inhibition at a concentration of less than about 30 µg/mL, less than about 10 µg/mL, or less than about 2 µg/mL. Preferably, the plant extract may reduce expression of a KHK-C gene or the activity of a KHK-C polypeptide by at least about 10%, preferably at least about 50%, more preferably at least about 75%, at least about 90%, or at least about 100% relative to the absence of the plant extract. The composition may be suitable for administration to a subject to support weight management. In one example, the composition may be administered to a subject to treat or prevent at least one of sugar addiction, obesity, diabetes, insulin resistance and metabolic syndrome. In one example, the composition may be administered to provide a diminished craving in the subject for at least one of sugar, fructose, fructose-containing sugars, carbohydrates, and combinations thereof. In one example, the subject may be diabetic.

[0025] The plant extract may include any suitable plant extract capable of inhibiting KHK-C activity. The plant extract may be present in the composition in an amount suitable to inhibit KHK-C activity in a subject. In one example, the plant extract may be obtained from a plant from a genus selected from the group consisting of *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, *Diospyros*, *Andrographis*, *Nymphaea*, *Chloroxylon*, *Petroselinum*, *Morus*, *Pteris*, *Garcinia*, and *Malus*. The plant extract may be obtained from a plant from a genus selected from the group consisting of *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, and *Diospyros*.


[0026] In one example, the plant extract may be obtained from a plant selected from the group consisting of *Angelica archangelica*, *Cratoxylum*

prunifolium, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*, *Andrographis paniculata*, *Nymphaea lotus*, *Chloroxylon swietenia*, *Petroselinum crispum*, *Morus alba*, *Pteris wallichiana*, *Garcinia mangostana*, and *Malus domestica*. The plant extract may be obtained from a plant selected from the group consisting of *Angelica archangelica*, *Cratoxylum prunifolium*, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*.


[0027] In one example, the plant extract may include two or more plant extracts each independently obtained from a plant from a genus selected from the group consisting of, *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, *Diospyros*, *Andrographis*, *Nymphaea*, *Chloroxylon*, *Petroselinum*, *Morus*, *Pteris*, *Garcinia*, and *Malus*. The two or more plant extracts each independently may be obtained from a plant selected from the group consisting of, *Angelica archangelica*, *Cratoxylum prunifolium*, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*, *Andrographis paniculata*, *Nymphaea lotus*, *Chloroxylon swietenia*, *Petroselinum crispum*, *Morus alba*, *Pteris wallichiana*, *Garcinia mangostana*, and *Malus domestica*.

[0028] The composition for inhibiting KHK-C activity may include one or more compounds which may function as active ingredients. The compound may be a component of the plant extract. For example, the compound may include a phytochemical present in the plant from which the plant extract is obtained. The compound may be at least partially responsible for the inhibition of KHK-C activity exhibited by the plant extract. The compound may include any compound capable of inhibiting KHK-C activity. In one example, the compound may be selected from the group consisting of Osthol, Cratoxyarborenone E, gamma-Mangostin, Osthenol, a Polyketide type molecule, 4-Hydroxy-Derricin, Isobavachalcone, Methoxy-isobavachalcone, Oroxylin A, 5,7-Dimethoxy-8-prenylcoumarin, Apigenin 7-glucuronide, 3',4',5,7-THMethoxy3'-O-β-D-Xylopyranoside, Swietenocoumarin B, Apiin, Mulberrin, Flavaspidic acid AB, Mangostin, Phloretin, and combinations thereof. The compound may be selected from the group consisting of Osthol, Cratoxyarborenone E, gamma-Mangostin,


Osthenol, a Polyketide type molecule, 4-Hydroxy-Derricin, Isobavachalcone, Methoxy-isobavachalcone, Oroxylin A, 5,7-Dimethoxy-8-prenylcoumarin, and combinations thereof. In one example, the compound may include a flavonoid, a polyphenol, or a combination thereof. The flavonoid may be a derivative of a phenyl-benzopyrone compound (e.g., 2-phenyl-1,4-benzopyrone, 3-phenyl-1,4-benzopyrone, or 4-phenyl-1,2-benzopyrone). In one example, the compound may include a prenylated side chain. In one example, the compound may include at least one of the functional groups I, II, or III, shown below:



I



II



III

[0029] The plant extract may be commercially obtained from various sources. The plant extract may be obtained using any suitable extraction technique. Generally, any part of a plant may be used to produce the plant extract including, but not limited to, the root, the stem, the leaf, the flower, the fruit, and the fruit pod. One or more parts of the plant may be extracted to yield the plant extract. In this regard, one or more parts of the plant may be collected and milled. Thereafter, the milled material may be extracted with a suitable solvent. The solvent may be removed in a concentration step. For example, the

extracted material may be screened or filtered to create a supernatant and a cake. The cake may be pressed to remove a substantial portion of the liquid, which may be added to the supernatant. The cake then may be dehydrated and used as a fiber source. The supernatant may be distilled to remove the solvent, or a portion thereof, to form a plant extract liquid concentrate. The removed solvent may be recycled. The concentrate may be dried (e.g., by spray drying) to provide a dried plant extract. The dried plant extract may be assayed and/or standardized as described herein.

[0030] The solvent may include alcohol, water, or a combination thereof. Exemplary alcoholic solvents may include, but are not limited to, C1-C7 alcohols (e.g., methanol, ethanol, propanol, isopropanol, and butanol), hydro-alcohols, or mixtures of alcohol and water (e.g., hydroethanol), polyhydric alcohols (e.g., propylene glycol and butylene glycol), and fatty alcohols. Any of these alcoholic solvents may be used in the form of a mixture. In one example, the plant extract is extracted using ethanol, water, or a combination thereof (e.g., a mixture of about 95% ethanol and about 5% water).

[0031] In one example, the plant extract may be obtained using an organic solvent extraction technique. In another example, solvent sequential fractionation may be used to obtain the plant extract. Total hydro-ethanolic extraction techniques also may be used to obtain the plant extract. Generally, this is referred to as a lump-sum extraction. The plant extract generated in the process may include a broad variety of phytochemicals present in the extracted material. The phytochemicals may be fat soluble or water soluble. Following collection of the extract solution, the solvent may be evaporated, resulting in the extract.

[0032] Total ethanol extraction also may be used. This technique uses ethanol as the solvent. This extraction technique may generate a plant extract that includes fat soluble and/or lipophilic compounds in addition to water soluble compounds.

[0033] Another example of an extraction technique that may be used to obtain the plant extract is supercritical fluid carbon dioxide extraction (SFE). In

this extraction procedure, the material to be extracted may not be exposed to any organic solvents. Rather, carbon dioxide may be used as the extraction solvent, with or without a modifier, in super-critical conditions (>31.3°C and > 73.8 bar). Those of skill in the art will appreciate that temperature and pressure conditions can be varied to obtain the best yield of extract. This technique may generate an extract of fat soluble and/or lipophilic compounds, similar to a total hexane and ethyl acetate extraction technique.

[0034] The plant extract may be standardized to a specified amount of a particular compound. For example, the plant extract may be standardized to a specified amount of an active ingredient or phytochemical.

[0035] The amount of plant extract present in the KHK-C inhibiting composition may depend upon several factors, including the desired level of KHK-C inhibition, the KHK-C inhibiting level of a particular plant extract or component thereof, and other factors. Preferably, the plant extract may be present in an amount of from about 0.005 weight percent to about 50 weight percent based on the weight of the total composition.

[0036] The KHK-C inhibiting composition may include one or more acceptable carriers. The carrier may aid in enabling incorporation of the plant extract into a KHK-C inhibiting composition having a suitable form for administration to a subject. A wide number of acceptable carriers are known in the art, and the carrier may be any suitable carrier. The carrier may be suitable for administration to animals, including humans, and may be able to act as a carrier without substantially affecting the desired activity of the plant extract and/or any active ingredient. The carrier may be selected based upon the desired administration route and dosage form of the composition. For example, the composition may be suitable for use in a variety of dosage forms, such as liquid form and solid form. In one example, the composition may be provided as a gel, a syrup, a slurry, or a suspension. In one example, the composition may be provided in a liquid dosage form such as a drink shot or a liquid concentrate. In one example, the composition may be provided in a solid dosage form, such as a tablet, a pill, a capsule, a dragée, or a powder. The composition, in liquid or

solid dosage form, may be in a food delivery form that is suitable for incorporation into food for delivery. Examples of suitable carriers for use in solid forms (particularly tablet and capsule forms) may include, but are not limited to, organic and inorganic inert carrier materials such as gelatin, starch, magnesium stearate, talc, gums, silicon dioxide, stearic acid, cellulose, and the like. The carrier may be substantially inert.

[0037] In one example, silicified microcrystalline cellulose may be used as a carrier. Silicified microcrystalline cellulose is a physical mixture of microcrystalline cellulose and colloidal silicon dioxide. One suitable form of silicified microcrystalline cellulose may include Prosolve 90 available from Penwest of Patterson, N.J. Silicon dioxide, in addition to that provided by the silicified microcrystalline cellulose, may be added to the composition as a processing aid. For example, silicon dioxide may be included as a glidant to improve the flow of powder during compression in the manufacturing of solid dosage units, such as tablets.

[0038] The KHK-C inhibiting composition may include other inert ingredients, such as lubricants and/or glidants. Lubricants may ease the handling of tablets during manufacturing, such as during ejection from dies. Glidants may improve powder flow during tablet compression. Stearic acid may be used as an acceptable lubricant/glidant.

[0039] The KHK-C inhibiting composition may be made in a solid dosage form, such as tablets and capsules. This form may provide a product that can be easily transported with an individual to a place of eating, such as a restaurant, and taken prior to consumption of a foodstuff. The composition may be formulated into dosage units that contain suitable amounts of the plant extract and/or active ingredient to permit an individual to determine an appropriate number of units to take based upon appropriate parameters, such as body weight, foodstuff size, or carbohydrate (e.g., sugar) content.

[0040] In one example, the KHK-C inhibiting composition may be provided in a solid dosage form (e.g., tablets or caplets) individually including from about 50 mg to about 2 g of the plant extract. The compound may be administered

such that a dosage of the plant extract is from about 150 mg per day to about 2 g per day. The compound may be administered as a single dose or in multiple doses. In one example, the compound may be administered in up to three doses per day. For example, the compound may be administered prior to meals.

[0041] The dosage may be selected to provide a level of inhibitory effect in a single unit that may be effective for some individuals and/or some foodstuffs, while also allowing for relatively simple dosage increases to provide other levels of inhibitory effects that may be effective for other individuals and/or other foodstuffs.

[0042] In one example, the KHK-C inhibiting composition may be in a form adapted for oral ingestion. The form may be configured as a single dosage form intended to provide a specified dose of the plant extract. For example, the single dosage form may be a pill, a tablet, a capsule, or a drink shot. The single dosage form may include from about 50 mg to about 2 g of the plant extract.


[0043] In one example, the carrier may include saline, buffered saline, dextrose, or water. The carrier may include suitable excipients or auxiliaries to facilitate processing of the active compounds into preparations suitable for administration to a subject. The composition may be administered by any suitable route including oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Oral dosage forms may include tablets, pills, dragées, capsules, liquids, gels, syrups, slurries, suspensions, and the like.

[0044] Certain embodiments relate to a method for inhibiting KHK-C activity in a subject comprising administering a plant extract that exhibits at least 50% KHK-C inhibition at a concentration of from about 0.1 $\mu\text{g}/\text{mL}$ to about 1000 $\mu\text{g}/\text{mL}$. The administering is done to treat or prevent at least one of sugar addiction, obesity, or metabolic syndrome. The administering is also done to provide a diminished craving in the subject from at least one member selected from the group consisting of craving of sugar, fructose, fructose-containing

sugars, carbohydrates, and combinations thereof. The subject may be pre-diabetic, diabetic, and or insulin resistant.

Examples

[0045] The plant extracts identified below with reference to Table 1 are evaluated for inhibitory properties in a cell-free KHK-C model assay system. Each plant extract demonstrates meaningful inhibitory activity against KHK-C (i.e., a 50% inhibitory activity concentration in the low μM range, concentrations that are feasible within the body following oral consumption of low milligram doses). Interestingly, a number of the plant extracts possess a prenylated side chain (e.g., an isoprenyl, a geranyl, or a 1,1-dimethylallyl moiety) as part of their natural molecular backbones. The following structure illustrates one example of a compound having such prenylated side chains.



[0046] The plant extracts are screened using a 96-well high throughput enzymatic KHK assay that utilizes recombinant proteins. Recombinant proteins of human KHK-C and KHK-A are produced using the Profinity eXact fusion-tag system available from Bio-Rad Laboratories, Hercules, CA. KHK-C and KHK-A activity is assayed using a 3-step reaction. Fructose is broken down by fructokinase into fructose-1-phosphate. The ADP generated is coupled with p-enolpyruvate to generate pyruvate. The pyruvate is then coupled with NADH and broken down into NAD⁺ and lactate by lactate dehydrogenase. A Synergy 2 multi-mode microplate reader, available from BioTek Instruments, Inc., Winooski, VT, is used to measure the decrease in NADH using absorbance at 340 nm ($A_{340\text{nm}}$).

[0047] For the screening of the plant extracts, the KHK-C enzymatic assay is measured at 37°C and uses 50 mM piperazine-N,N'-bis(2-ethanesulfonic acid) (PIPES), 6 mM MgCl₂, 100 mM KCl, 5 mM ATP, 2 mM phosphoenolpyruvate, 0.3 mM NADH, 10 U of pyruvate kinase, 10 U of lactate dehydrogenase, 75 ng of KHK-C, and 1 mM fructose in a total reaction volume of 200 µl. The KHK-A enzymatic assay uses the same reaction conditions except that 30 mM fructose and 50 ng/µl KHK-A are used. The mastermix (without fructose) is incubated for 5 min at 37°C. The mixture is then added to a 96-well plate containing 10 µl of the plant extracts and then incubated for 15 min at 37°C. Fructose is added to the reactions, except for the negative controls, and A_{340nm} data is collected every minute for 1 hr. The change in absorbance during the first 30 minutes is calculated for each sample. The change in absorbance is calculated as the difference between A_{340nm} at 0 min and A_{340nm} at 30 min according to the following equation:

$$\Delta A_{340nm} = A_{340nm}(0 \text{ min}) - A_{340nm}(30 \text{ min})$$

The samples are then adjusted for the negative control by calculating the difference between ΔA_{340nm} of the respective sample and ΔA_{340nm} of the negative control according to the following equation:

$$\text{Adj } \Delta A_{340nm} = \Delta A_{340nm} \text{ Sample} - \Delta A_{340nm} \text{ Negative Control}$$

The percentage inhibition of KHK-C is calculated using the following formula:

$$\frac{\text{Adjusted } \Delta A_{340nm} \text{ Positive Control} - \text{Adjusted } \Delta A_{340nm} \text{ Sample}}{\text{Adjusted } \Delta A_{340nm} \text{ Positive Control}} \times 100$$

4-(hydroxymercuri) benzoic acid sodium salt is used as a positive inhibitory control of both KHK-C and KHK-A. Using this procedure, a 50% KHK inhibitory activity concentration (IC₅₀) is calculated for each plant extract.






[0048] Samples 1-16 shown in Table 1 are screened using the procedure described above. Each sample is an extract from the plant of the listed genus and species and exhibits the 50% KHK inhibitory activity concentration (IC₅₀) recited in Table 1.







Table 1






Sample No.	Plate ID	Well ID	Genus	Species	Description	IC ₅₀ (μM)
1	UCO-02	G10	<i>Angelica</i>	<i>archangelica</i>	European medicinal plant	0.53
2	UCO-02	C08	<i>Cratoxylum</i>	<i>prunifolium</i>	tree from SE Asia	1.40
3	UCO-04	F02	<i>Cratoxylum</i>	<i>prunifolium</i>	tree from SE Asia	1.97
4	UCO-05	G10	<i>Angelica</i>	<i>archangelica</i>	European medicinal plant	5.97
5	UCO-02	B08	<i>Myrica</i>	<i>cerifera</i>	medicinal plant from SE N America	6.20
6	UCO-10	B11	<i>Psoralea</i>	<i>corylifolia</i>	Ayurvedic medicinal plant	6.20
7	UCO-03	E01	<i>Scutellaria</i>	<i>baicalensis</i>	Chinese medicinal plant	7.70
8	UCO-01	G07	<i>Diospyros</i>	<i>attenuata</i>	tree from E Africa	9.40
9	UCO-04	F03	<i>Andrographis</i>	<i>paniculata</i>	Chinese medicinal plant	24.10
10	UCO-01	D10	<i>Nymphaea</i>	<i>lotus</i>	plant from SE Asia and E Africa	26.00
11	UCO-13	G6	<i>Chloroxylon</i>	<i>swietenia</i>	threatened species	26.00
12	UCO-06	E01	<i>Petroselinum</i>	<i>crispum</i>	edible	40.60
13	UCO-03	D07	<i>Morus</i>	<i>alba</i>	edible	44.20
14	UCO-02	F08	<i>Pteris</i>	<i>wallichiana</i>	fern from SE Asia	44.90
15	FTL plate 8	B09	<i>Garcinia</i>	<i>mangostana</i>	tree from SE Asia	30ug/mL
16	FTL plate 9	E10	<i>Malus</i>	<i>domestica</i>	apple tree	9ug/mL


[0049] Table 2 shows a phytochemical present in each of samples 1-17, including the structure of each phytochemical.

Table 2

Sample No.	Phytochemical	CAS	Structure
1	Osthol	484-12-8	
2	Cratoxyarborenone E		
3	gamma-Mangostin	31271-07-5	
4	Osthenol	484-14-0	
5	Polyketide type molecule		

6	4-Hydroxy-Derricin, Isobavachalcone	20784-50-3	
7	Methoxy-Isobavachalcone		
8	Oroxylin A	480-11-5	
9	5,7-Dimethoxy-8-prenylcoumarin	17245-25-9	
10	Apigenin 7-glucuronide	29741-09-1	
11	3',4',5,7-THMethoxy3'-O-β-D-Xylopyranoside	93373-16-1	

12	Swietenocoumarin B	64652-23-9	 The chemical structure of Swietenocoumarin B is a complex polycyclic molecule. It features a central coumarin core with multiple oxygen atoms and a side chain containing a benzene ring with a hydroxyl group and a lactone ring.
13	Apiin	26544-34-3	 The chemical structure of Apiin is a complex polycyclic molecule. It features a central coumarin core with multiple oxygen atoms and a side chain containing a benzene ring with a hydroxyl group and a lactone ring.
14	Mulberrin	62949-79-5	 The chemical structure of Mulberrin is a complex polycyclic molecule. It features a central coumarin core with multiple oxygen atoms and a side chain containing a benzene ring with a hydroxyl group and a lactone ring.
15	Flavaspidic acid AB	3761-64-6	 The chemical structure of Flavaspidic acid AB is a complex polycyclic molecule. It features a central coumarin core with multiple oxygen atoms and a side chain containing a benzene ring with a hydroxyl group and a lactone ring.
16	Mangostin		 The chemical structure of Mangostin is a complex polycyclic molecule. It features a central coumarin core with multiple oxygen atoms and a side chain containing a benzene ring with a hydroxyl group and a lactone ring.

17	Phloretin	
----	-----------	--

EXAMPLES

[0050] Example 1: Method of Extraction: Preparation of three fractions, hydrophilic, lipophilic and mixed/combination fractions for biological in vitro high through-put screening.

[0051] Reagents / solutions

[0052] General Chemistry Laboratory supply and standard equipment.

[0053] Deionized water (DI). HPLC grade or equivalent.

[0054] Chloroform, (Trichloromethane), ACS grade. Fisher Scientific #C298-4 or equivalent.

[0055] Methanol, Optima grade, Fisher Scientific #A456-4 or equivalent.

[0056] Plant Materials: All plant materials used in this study were obtained from Applicants' farms in the form of dry powders.

[0057] Figure 1 shows a general schematic diagram of the extraction procedures described below.

[0058] A. Preparation of Hydrophilic Fractions:

[0059] Approximately 50 g, to the nearest 0.01 g, of the powdered botanical was weighed into a wide mouth 500 mL Erlenmeyer flask. A stir bar was added and 300 mL of methanol was poured into the flask. The flask was loosely covered using aluminum foil. The flask was placed on a magnetic stir plate and stirred for 12 hours (minimum) using a slow / medium stir rate. The

samples were kept out of direct light. Next, the flask was removed from the stir plate and sonicated for one hour, with occasional swirling, at room temperature. The sample solution was then filtered through GF/A Filter Paper directly into a 500 mL round flat bottom boiling flask. The filter paper was scraped and botanical residue was collected from the filter paper onto an aluminum weigh boat (or foil). The sample was dried at room temperature in a hood for at least 12 hours and the residue stored in an appropriate container. Next, using a graduated cylinder, 100 mL aliquot of this sample solution was pipetted out in an Erlenmeyer flask and capped with aluminum foil and stored in a refrigerator. The sample was then properly identified. This solution was then used for the preparation of a combination fraction in "Section C" below of this procedure.

[0060] Using a rotary evaporator, the remaining solvent was evaporated in the round bottom boiling flask. The volume of the solvent was reduced to less than 10 mL. The concentrated extract (still in liquid form) was then transferred, using a glass pipette, to a pre-weighed scintillation vial (weigh without cap). Methanol was used for further dilution for transfer purposes as needed.

[0061] The vial was then placed under nitrogen evaporator to reduce volume to as minimum as possible (using slow stream of nitrogen). Recommended water bath temperature should be around 40°C. The vial was then removed from the nitrogen evaporator and placed in a vacuum desiccator until dry (approximately 12 hours). The final dry weight (without cap) (check for constant weight) of the fraction was recorded in the scintillation vial and the weight of the fraction (by difference) was also recorded.

[0062] B. Preparation of Lipophilic Fractions:

[0063] Approximately 50g, to the nearest 0.01g, of the powdered botanical was weighed into a wide mouth 500 mL Erlenmeyer flask. A stir bar was added and 300 mL of chloroform was poured into the flask. The flask opening was covered using aluminum foil. The flask was then placed on a magnetic stir plate and stirred for 12 hours (minimum) using a slow / medium stir rate. The samples were kept out of direct light. The flask was then removed from the stir plate and

the sample was sonicated for one hour, with occasional swirling, at room temperature. Next, the sample solution was filtered through GF/A Filter Paper directly into a 500 mL round, flat bottom boiling flask. Using a graduated cylinder, a 100 mL aliquot of this solution was removed and stored in a foiled Erlenmeyer flask in a refrigerator for further step. This solution was then used for the preparation of a combination fraction in "Section C" of this procedure.

[0064] Using a rotary evaporator, the remaining solvent in the round bottom boiling flask was evaporated. The volume of the solvent was reduced to less than 10 mL. The concentrated extract, (still in a liquid form) was then transferred using a glass pipette, to a pre-weighed scintillation vial (weigh vial without cap). Chloroform was used for further dilution and transfer purposes as needed.

[0065] Next, to prepare a lipophilic fraction, the vial was placed under a nitrogen evaporator to reduce volume to as minimum as possible (using slow stream of nitrogen). Recommended water bath temperature should be around 40°C. The vial was then removed from the nitrogen evaporator and placed in a vacuum desiccator until dry (approximately 12 hours). The final dry weight (without cap) (check for constant weight) of the fraction was recorded in the scintillation vial and the weight of the fraction (by difference) was also recorded.

[0066] C. Preparation of a Mixed / Combination Fractions:

[0067] The two 100 mL aliquots of hydrophilic and lipophilic solutions saved during Preparations A and B above were combined. Specifically, a 100 mL aliquot of the sample solution from "Section A" above and 100 mL aliquot of the solution from "Section B" were combined into a 500 mL round bottom boiling flask and mixed well. Using a rotary evaporator, the solvent in the sample concentrate was evaporated. The volume of the solvent was reduced to less than 10 mL. The concentrated extract (still in liquid form) was transferred using a glass pipette to a pre-weighed scintillation vial (weigh without cap). A mixture of chloroform / methanol (1/1 v/v) was used for further dilution for transfer purposes as needed.

[0068] Next, the vial was placed under a nitrogen evaporator and the volume was reduced to minimum as possible (using slow stream of nitrogen). Recommended water bath temperature should be around 40 °C. The vial was then removed from the nitrogen evaporator and placed in a desiccator until dry (approximately 12 hours) and cooled to room temperature. The final dry weight (without cap) (check for constant weight) of the fraction in the scintillation vial (by difference) was recorded.

[0069] The sample was then properly identified.

[0070] The vial was stored in a refrigerator for future use.

[0071] **D. HPLC Methodology:**

[0072] Materials and Instrumentation:

[0073] All solvents were HPLC grade and purchased from Fisher Scientific. HPLC separation was achieved using an Agilent Technologies, Santa Clara, CA. HP1100 System equipped with Photodiode-array detection and Chemstation software using Waters C 18 4um NovaPak column (250 X 4.6mm) Part No 0528401. Botanical samples were fingerprinted with 0.2% ortho-phosphoric acid (OPA) v/v with deionized (DI) water and acetonitrile (ACN) elution gradient as outlined in the Table 1.

[0074] TABLE 1 (HPLC Conditions):

HPLC Gradient:		
Time	0.2% o-Phosphoric acid (OPA) in DI water	ACN (Acetonitrile)
0.0	92	8
12	90	10
14	88	12
26	76	24
34	60	40
37	60	40
40	92	8
42	92	8

Flow Rate:	1.0 mL/min
Column Temperature:	Ambient
Injection Volume:	10 µL

Detection Wavelengths:	210-370nm
Integration:	Peak area
Run Time:	42 minutes

[0075] Sample Preparation:

[0076] Approximately 300 mg of dry powdered sample of botanical extracts was weighted, to the nearest 0.1 mg, into a 50 mL volumetric flask. About 40 mL of 80/20 Methanol in DI Water (diluent) was added and the mixture was shaken well to dissolve. The flask was then placed in a sonic bath and sonicated for 10 minutes. The mixture was then cooled to room temperature, diluted to volume with diluent and mixed thoroughly. The sample solution was then filtered with a disposable syringe through a 0.45 micron PVDF filter into an HPLC auto sampler vial.

[0077] Results:

[0078] Typical HPLC fingerprint profiles of the tested botanicals are shown in FIGURES 2-8.

[0079] Specifically, an HPLC fingerprint profile for *Angelica archangelica* (Wild Celery) is shown in Figure 2; an HPLC fingerprint profile for *Myrica cerifera* (Bayberry) is shown in Figure 3; an HPLC fingerprint profile for *Scutellaria baicalensis* (Skullcap) is shown in Figure 4; an HPLC fingerprint profile for *Petroselinum crispum* (Garden Parsley) is shown in Figure 5; an HPLC fingerprint profile for *Garcinia mangostana* (Mangosteen) is shown in Figure 6; an HPLC fingerprint profile for *Psoralea corylifolia* (Malay tea) is shown in Figure 7; and an HPLC fingerprint profile for *Morus alba* (Mulberry) is shown in Figure 8.

[0080] While the present invention has been described with reference to specific exemplary embodiments, it will be evident that various modifications and changes may be made to these embodiments without departing from the spirit and scope of the invention. It is the following claims, including all equivalents, which are intended to define the spirit and scope of the invention.

We claim:

1. A composition for inhibiting ketohexokinase-C (KHK-C) activity comprising a plant extract exhibiting at least IC₅₀, wherein at least 50% KHK-C inhibition occurs at a concentration from about 0.1 µg/mL to about 1000 µg/mL.
2. The composition of claim 1, wherein the plant extract is obtained from a plant from a genus selected from the group consisting of *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, *Diospyros*, *Andrographis*, *Nymphaea*, *Chloroxylon*, *Petroselinum*, *Morus*, *Pteris*, *Garcinia*, and *Malus*.
3. The composition of any of claims 1-2, wherein the plant extract is obtained from a plant selected from the group consisting of *Angelica archangelica*, *Cratoxylum prunifolium*, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*, *Andrographis paniculata*, *Nymphaea lotus*, *Chloroxylon swietenia*, *Petroselinum crispum*, *Morus alba*, *Pteris wallichiana*, *Garcinia mangostana*, and *Malus domestica*.
4. The composition of any of claims 1-3, wherein the plant extract is obtained from a plant from a genus selected from the group consisting of *Angelica*, *Cratoxylum*, *Myrica*, *Psoralea*, *Scutellaria*, and *Diospyros*.
5. The composition of claim 4, wherein the plant extract is obtained from a plant selected from the group consisting of *Angelica archangelica*, *Cratoxylum prunifolium*, *Myrica cerifera*, *Psoralea corylifolia*, *Scutellaria baicalensis*, and *Diospyros attenuata*.
6. The composition of any of claims 1-5, wherein the plant extract comprises a compound selected from the group consisting of Osthol, Cratoxyarborenone E, gamma-Mangostin, Osthonol, a Polyketide type molecule, 4-Hydroxy-Derricin, Isobavachalcone, Methoxy isobavachalcone, Oroxylin A, 5,7-Dimethoxy-8-

prenylcoumarin, Apigenin 7-glucuronide, 3',4',5,7-THMethoxy3'-O- β -D-Xylopyranoside, Swietenocoumarin B, Apiin, Mulberrin, Flavaspidic acid AB, Mangostin, Phloretin, and combinations thereof.



7. The composition of any of claims 1-6, wherein the plant extract comprises a compound having a prenylated side chain.


8. The composition of any of claims 1-7, wherein an amount of the plant extract in the composition is between about 0.005 weight percent and about 50 weight percent.

9. The composition of any of claims 1-8, wherein the composition is in a form suitable for oral ingestion.

10. The composition of claim 9, wherein the form is selected from the group consisting of a pill, a tablet, a capsule, a caplet, a dragée, a powder, a liquid, a gel, a syrup, a slurry, and a suspension.

11. The composition of any of claims 1-10, wherein the plant extract comprises at least one of the following functional groups I, II, or III:






III


12. A method for inhibiting KHK-C activity in a subject comprising administering a plant extract that exhibits at least 50% KHK-C inhibition at a concentration of from about 0.1 $\mu\text{g/mL}$ to about 1000 $\mu\text{g/mL}$.
13. The method of claim 12, wherein the administering is done to treat or prevent at least one of sugar addiction, obesity, or metabolic syndrome.
14. The method of any of claims 12-13, wherein the administering is done to provide a diminished craving in the subject from at least one member selected from the group consisting of sugar, fructose, fructose-containing sugars, carbohydrates, and combinations thereof.
15. The method of any of claims 12-14, wherein the subject is pre-diabetic.
16. The method of any of claims 12-14, wherein the subject is diabetic.
17. The method of any of claims 12-14, wherein the subject is insulin resistant.

Overview of Extraction Process




Powdered Plants are Extracted & Concentrated Using a Standardized Process


FIG. 1



Angelica archangelica (Wild Celery)
FIG. 2




Myrica cerifera (Bayberry)
FIG. 3




Scutellaria baicalensis (Skullcap)


FIG. 4



Petroselinum crispum (Garden Parsley)
FIG. 5




Garcinia mangostana (Mangosteen)
FIG. 6



Psoralea corylifolia (Malay tea)

FIG. 7



Morus alba (Mulberry)

FIG. 8

INTERNATIONAL SEARCH REPORT

International application No PCT/US2014/042534

A. CLASSIFICATION OF SUBJECT MATTER		
INV.	A61K36/75	A61K36/12
	A61K36/232	A61K36/38
	A61K36/605	A61K36/62
		A61K36/185
		A61K36/44
		A61K36/73
		A61K36/19
		A61K36/487
		A61P3/04
		A61K36/23
		A61K36/539
		A61P3/10
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, BIOSIS, EMBASE, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P	EP 2 679 229 A1 (BIOACTIVE FOOD GMBH [DE]) 1 January 2014 (2014-01-01) claims; examples	1-3,6-17
X	----- WO 2010/100653 A2 (LAILA NUTRACEUTICALS [IN]; GOKARAJU GANGA RAJU [IN]; GOKARAJU RAMA RAJ) 10 September 2010 (2010-09-10) claims; examples	1-3,6-17
X	----- US 2006/251749 A1 (JIA QI [US] ET AL) 9 November 2006 (2006-11-09) claims; examples	1-17
	----- -/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search	Date of mailing of the international search report	
13 August 2014	26/08/2014	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Friederich, Martin	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2014/042534

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>DATABASE WPI Week 201263 Thomson Scientific, London, GB; AN 2012-E65777 XP002728441, & KR 2012 0031800 A (CHOI S W) 4 April 2012 (2012-04-04) abstract</p> <p style="text-align: center;">-----</p>	1-3,6-17
X	<p>DATABASE WPI Week 200940 Thomson Scientific, London, GB; AN 2009-K16841 XP002728442, & JP 2009 125030 A (SHIROTA Y) 11 June 2009 (2009-06-11) abstract</p> <p style="text-align: center;">-----</p>	1-3,6-17
X	<p>RAMMOHAN SUBRAMANIAN ET AL: "A bitter plant with a sweet future? A comprehensive review of an oriental medicinal plant:", PHYTOCHEMISTRY REVIEWS, KLUWER ACADEMIC PUBLISHERS, DO, vol. 11, no. 1, 2 October 2011 (2011-10-02), pages 39-75, XP035015554, ISSN: 1572-980X, DOI: 10.1007/S11101-011-9219-Z the whole document</p> <p style="text-align: center;">-----</p>	1-3,6-17
X	<p>Bandna Devi ET AL: "MORUS ALBA LINN: A PHYTOPHARMACOLOGICAL REVIEW", 1 January 2013 (2013-01-01), XP055134293, Retrieved from the Internet: URL:http://www.ijppsjournal.com/Vol15Suppl2 /6747.pdf [retrieved on 2014-08-12] the whole document</p> <p style="text-align: center;">-----</p>	1-3,6-17
X,P	<p>CHOPRA BHAWNA ET AL: "Psoralea corylifoliaL. (Buguchi) - Folklore to modern evidence: Re", FITOTERAPIA, vol. 90, 4 July 2013 (2013-07-04), pages 44-56, XP028739851, ISSN: 0367-326X, DOI: 10.1016/J.FITOTE.2013.06.016 the whole document</p> <p style="text-align: center;">-----</p>	1-17
	-/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2014/042534

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	TAO X ET AL: "Using Chinese natural products for diabetes mellitus drug discovery and development", EXPERT OPINION ON DRUG DISCOVERY, INFORMA HEALTHCARE, LONDON, GB, vol. 2, no. 7, 1 July 2007 (2007-07-01), pages 977-986, XP009140069, ISSN: 1746-0441, DOI: 10.1517/17460441.2.7.977 table 5	1-17
X	----- EVA KEMPS ET AL: "Non-food odorants reduce chocolate cravings", APPETITE, ACADEMIC PRESS, NEW YORK, NY, US, vol. 58, no. 3, 2 March 2012 (2012-03-02), pages 1087-1090, XP028419563, ISSN: 0195-6663, DOI: 10.1016/J.APPET.2012.03.002 [retrieved on 2012-03-09] table 1	1-3,6-17
X	----- BLUM ET AL: "Genotrim(TM), a DNA-customized nutrigenomic product, targets genetic factors of obesity: Hypothesizing a dopamine-glucose correlation demonstrating reward deficiency syndrome (RDS)", MEDICAL HYPOTHESES, EDEN PRESS, PENRITH, US, vol. 68, no. 4, 25 January 2007 (2007-01-25), pages 844-852, XP005744405, ISSN: 0306-9877, DOI: 10.1016/J.MEHY.2006.08.041 page 849, column 1, last paragraph -----	1,2,6-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No PCT/US2014/042534

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 2679229	A1	01-01-2014	NONE

WO 2010100653	A2	10-09-2010	AU 2010220058 A1 21-07-2011
			CA 2751227 A1 10-09-2010
			CN 102292093 A 21-12-2011
			EP 2391374 A2 07-12-2011
			JP 2012516842 A 26-07-2012
			KR 20110122158 A 09-11-2011
			US 2011280951 A1 17-11-2011
			WO 2010100653 A2 10-09-2010

US 2006251749	A1	09-11-2006	BR PI0608796 A2 26-01-2010
			CN 101217968 A 09-07-2008
			EP 1881839 A2 30-01-2008
			JP 5443754 B2 19-03-2014
			JP 2008540551 A 20-11-2008
			KR 20080016609 A 21-02-2008
			KR 20130125411 A 18-11-2013
			US 2006251749 A1 09-11-2006
			US 2011223267 A1 15-09-2011
			WO 2006122160 A2 16-11-2006

KR 20120031800	A	04-04-2012	NONE

JP 2009125030	A	11-06-2009	JP 5450952 B2 26-03-2014
			JP 2009125030 A 11-06-2009

摘要

一种抑制己酮糖激酶的组合，例如，己酮糖激酶-C (KHK-C) 活性，可以包括一种具有至少如下 IC₅₀ (例如，浓度范围在约 0.1 μg/mL 至约 1000 μg/mL 范围内的 50% KHK-C 抑制) 的植物提取物。组合物可以是适于口服的形式。抑制受试者中 KHK-C 活性的方法包括施用一种植物提取物，其具有在约 0.1 μg/mL 至约 1000 μg/mL 浓度的至少 50% KHK-C 抑制。施用可以治疗或预防糖成瘾、肥胖或代谢综合征中的至少一种。施用可以减少受试者对选自以下组中至少一个成员的渴求，该组包括糖、果糖、含果糖的糖、碳水化合物、以及它们的组合。受试者可以是糖尿病前期、糖尿病和 / 或胰岛素抵抗。

