

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 200041853 B2**
(10) Patent No. **774738**

(54) Title
(E)-styryl sulfone anticancer agents

(51) ⁶ International Patent Classification(s)
A61K 031/10 C07C 321/20
C07C 317/10

(21) Application No: 200041853 (22) Application Date: 2000.03.31

(87) WIPO No: W000/59495

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/127683	1999.04.02	US
60/143975	1999.07.15	US

(43) Publication Date : 2000.10.23
(43) Publication Journal Date : 2001.01.04
(44) Accepted Journal Date : 2004.07.08

(71) Applicant(s)
Temple University - of The Commonwealth System of Higher Education

(72) Inventor(s)
E. Premkumar Reddy; M. V. Ramana Reddy

(74) Agent/Attorney
Davies Collison Cave, Level 15, 1 Nicholson Street, MELBOURNE VIC 3000

(56) Related Art
CA 101: 23063
CA 125: 327911
CA 120: 210378



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A61K 31/10, C07C 317/10, 321/20		A1	(11) International Publication Number: WO 00/59495
			(43) International Publication Date: 12 October 2000 (12.10.00)
(21) International Application Number: PCT/US00/08565		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 31 March 2000 (31.03.00)			
(30) Priority Data: 60/127,683 2 April 1999 (02.04.99) US 60/143,975 15 July 1999 (15.07.99) US			
(71) Applicant (for all designated States except US): TEMPLE UNIVERSITY - OF THE COMMONWEALTH SYSTEM OF HIGHER EDUCATION [US/US]; Broad Street and Montgomery Avenue, Philadelphia, PA 19122 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): REDDY, E., Premkumar [US/US]; 547 Atterbury Road, Villanova, PA 19085 (US). REDDY, M., V., Ramana [IN/US]; 921 St. Joseph Drive, Upper Darby, PA 19082 (US).			
(74) Agent: MONACO, Daniel, A.; Seidel, Gonda, Lavorgna & Monaco, P.C., Two Penn Center Plaza, Suite 1800, Philadelphia, PA 19102 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: (E)-STYRYL SULFONE ANTICANCER AGENTS			
(57) Abstract			
(E)-styryl benzyldisulfones of formula (I) are useful as anticancer agents wherein R₁, R₂, R₃, and R₄ are independently selected from the group consisting of hydrogen, fluoro, chloro, iodo, bromo, C1-C6 alkyl, C1-C4 alkoxy, nitro, cyano and trifluoromethyl, with the proviso that (a) R₁, R₂, and R₃ not all hydrogen when R₄ is 2-chloro or 4-chloro; (b) when R₁ and R₃ are hydrogen and R₂ is 4-bromo or 4-chloro, then R₄ may not be 4-chloro, 4-fluoro or 4-bromo; (c) when R₁ and R₃ are hydrogen and R₂ is 4-fluoro, then R₄ may not be 4-fluoro or 4-bromo; (d) when R₁ is hydrogen, and R₄ is 2-fluoro, the R₂ and R₃ may not be 4-fluoro; and (e) when R₁ is hydrogen and R₃ is 4-hydrogen, 4-chloro, 4-bromo, 4-methyl or 4-methoxy, and R₄ is 2-hydrogen, 2-chloro, or 2-fluoro; then R₂ may not be 4-hydrogen, 4-chloro, 4-fluoro, or 4-bromo.			

(E)-STYRYL SULFONE ANTICANCER AGENTS

5

Field of the Invention

10 The invention relates to compositions and methods for the treatment of cancer.

Background of the Invention

15 Extracellular signals received at transmembrane receptors are relayed into the cells by the signal transduction pathways (Pelech *et al.*, *Science* **257**:1335 (1992)) which have been implicated in a wide array of physiological processes such as induction of cell proliferation, differentiation or apoptosis (Davis *et al.*, *J. Biol. Chem.* **268**:14553 (1993)). The Mitogen Activated Protein Kinase (MAPK) cascade is a major signaling system by which cells transduce extracellular cues into intracellular responses (Nishida *et al.*, *Trends Biochem. Sci.* **18**:128 (1993); Blumer *et al.*, *Trends Biochem. Sci.* **19**:236 (1994)). Many steps of this cascade are

20

- 2 -

conserved, and homologous for MAP kinases have been discovered in different species.

In mammalian cells, the Extracellular-Signal-Regulated Kinases (ERKs), ERK-1 and ERK-2 are the archetypal and best-studied members of the MAPK family, which all have the unique feature of being activated by phosphorylation on threonine and tyrosine residues by an upstream dual specificity kinase (Posada *et al.*, *Science* **255**:212 (1992); Biggs III *et al.*, *Proc. Natl. Acad. Sci. USA* **89**:6295 (1992); Garner *et al.*, *Genes Dev.* **6**:1280 (1992)).

Recent studies have identified an additional subgroup of MAPKs, known as c-Jun NH2-terminal kinases 1 and 2 (JNK-1 and JNK-2), that have different substrate specificities and are regulated by different stimuli (Hibi *et al.*, *Genes Dev.* **7**:2135 (1993)). JNKs are members of the class of stress-activated protein kinases (SPKs). JNKs have been shown to be activated by treatment of cells with UV radiation, pro-inflammatory cytokines and environmental stress (Derijard *et al.*, *Cell* **102**:5 (1994)). The activated JNK binds to the amino terminus of the c-Jun protein and increases the protein's transcriptional activity by phosphorylating it at ser63 and ser73 (Adler *et al.*, *Proc. Natl. Acad. Sci. USA* **89**:5341 (1992); Kwok *et al.*, *Nature* **370**:223 (1994)).

Analysis of the deduced primary sequence of the JNKs indicates that they are distantly related to ERKs (Davis, *Trends Biochem. Sci.* **19**:470 (1994)). Both ERKs and JNKs are phosphorylated on Tyr and Thr in response to external stimuli resulting in their activation (Davis, *Trends Biochem. Sci.* **19**:470 (1994)). The phosphorylation (Thr and Tyr) sites, which play a critical role in their activation are conserved between ERKs and JNKs (Davis, *Trends Biochem. Sci.* **19**:470 (1994)). However, these sites of phosphorylation are located within distinct dual phosphorylation motifs: Thr-Pro-Tyr (JNK) and Thr-Glu-Tyr (ERK). Phosphorylation of MAPKs and JNKs by an external signal often involves the activation of protein tyrosine kinases (PTKs) (Gille *et al.*, *Nature*

358:414 (1992)), which constitute a large family of proteins encompassing several growth factor receptors and other signal transducing molecules.

Protein tyrosine kinases are enzymes which catalyze a well defined chemical reaction: the phosphorylation of a tyrosine residue
5 (Hunter *et al.*, *Annu Rev Biochem* 54:897 (1985)). Receptor tyrosine kinases in particular are attractive targets for drug design since blockers for the substrate domain of these kinases is likely to yield an effective and selective antiproliferative agent. The potential use of protein tyrosine kinase blockers as antiproliferative agents was recognized as early as
10 1981, when quercetin was suggested as a PTK blocker (Graziani *et al.*, *Eur. J. Biochem.* 135:583-589 (1983)).

The best understood MAPK pathway involves extracellular signal-regulated kinases which constitute the Ras/Raf/MEK/ERK kinase cascade (Boudewijn *et al.*, *Trends Biochem. Sci.* 20, 18 (1995)). Once this
15 pathway is activated by different stimuli, MAPK phosphorylates a variety of proteins including several transcription factors which translocate into the nucleus and activate gene transcription. Negative regulation of this pathway could arrest the cascade of these events.

What are needed are new anticancer chemotherapeutic
20 agents which target receptor tyrosine kinases and which arrest the Ras/Raf/MEK/ERK kinase cascade. Oncoproteins in general, and signal transducing proteins in particular, are likely to be more selective targets for chemotherapy because they represent a subclass of proteins whose activities are essential for cell proliferation, and because their activities are
25 greatly amplified in proliferative diseases.

What is also needed are new anticancer therapeutics which are highly selective in the killing of tumor cells, but not normal cells.

Summary of the Invention

30 In one embodiment the present invention provides compounds, compositions and methods for the treatment of cancer and other


proliferative diseases. The biologically active compounds are in the form of (E)-styryl benzylsulfones.

In another embodiment, the present invention provides compounds which are highly selective in killing tumor cells but not normal cells.

In yet another embodiment, the present invention provides novel polymers prepared by polymerization of (E)-styryl benzylsulfones.

In still yet another embodiment, the present invention provides intermediates useful for the preparation of compounds having anticancer activity. The intermediates comprise (E)-styryl benzylsulfonyl acetic acids.

10 According to one embodiment of the invention, novel compounds are provided according to formula I:



wherein:

15 R₁, R₂, R₃, and R₄ are independently selected from the group consisting of hydrogen; fluoro; chloro; bromo; C1-C6 alkyl; C1-C6 alkoxy; nitro; cyano; and trifluoromethyl;

with the proviso that

- 20 (a) R₁, R₂, R₃, and R₄ may not all be hydrogen;
- (b) when R₁, R₂, and R₃ are hydrogen, then R₄ may not be:
- (i) 2- or 4-chloro or 4-fluoro;
 - (ii) 2-nitro, 3-nitro or 4-nitro;
 - (iii) 4-methoxy or 4-ethoxy; or
 - (iv) 4-methyl;
- (c) when R₁ and R₃ are hydrogen and R₂ is 4-chloro, then R₄ may not be 4-chloro, 4-fluoro, 4-bromo, 4-nitro, 4-isopropyl or 4-ethoxy;

(d) when R_1 and R_3 are hydrogen and R_2 is 4-fluoro, then R_4 may not be 4-fluoro, 4-bromo, or 4-chloro

(e) when R_1 and R_3 are hydrogen and R_2 is 4-nitro, then R_4 may not be 4-chloro, 4-nitro, 4-bromo, 4-fluoro, 4-methyl, or 4-methoxy;

5 (f) when R_1 and R_3 are hydrogen and R_2 is 4-methyl, R_4 may not be 4-chloro, 4-bromo, 4-fluoro, 4-methyl or 2-chloro;

(g) when R_1 and R_3 are hydrogen and R_2 is 4-bromo, then R_4 may not be 4-fluoro, 4-bromo or 4-chloro;

10 (h) when R_1 and R_2 are hydrogen, then R_3 and R_4 may not be 2, 4-dichloro, 2,3-dimethoxy or 3,4-dimethoxy;

(i) when R_1 is hydrogen, then R_2 , R_3 and R_4 may not all be fluoro; and

15 (j) when R_1 is hydrogen and R_3 is 2-fluoro, then R_2 and R_4 may not both be selected from the group consisting of 4-chloro, 4-bromo, and 4-fluoro.


According to a preferred embodiment of the invention, novel compounds are provided according to formula I wherein R_1 , R_2 , R_3 , and R_4 are independently selected from the group consisting of hydrogen, chloro, fluoro, bromo, nitro, cyano and trifluoromethyl. According to a more preferred embodiment, R_1 , R_2 , R_3 , and R_4 are independently selected from the group consisting of hydrogen, chloro, fluoro and bromo; most preferably hydrogen, chloro and fluoro.

20 In a further preferred embodiment, novel compounds are provided according to formula I wherein (1) at least one of R_1 and R_2 is other than hydrogen and is located at the 2-, 3- and/or 4- position of the phenyl ring to which it is attached, and is preferably selected from chloro and fluoro, most preferably chloro; and/or (2) wherein at least one of R_3 and R_4 is other than hydrogen and is located at the 2- and/or 4- position of the phenyl ring to which it is attached, and is preferably selected from chloro and fluoro. In other preferred embodiments wherein at least one of R_1 and R_2 is other than hydrogen, and at least one of R_3 and R_4 is other than

- 6 -

hydrogen, (i) R_2 is 4-halogen or 4-cyano, and R_4 is 4-nitro; or (ii) R_2 is 4-C1-C6 alkoxy, and R_4 is 4-nitro or 4-halogen. R_1 and R_3 are preferably hydrogen in these embodiments.

- In another embodiment of the invention, a pharmaceutical composition is provided comprising a pharmaceutically acceptable carrier and one or more compounds of formula II:



wherein


- R_1 , R_2 , R_3 , and R_4 are independently selected from the group consisting of hydrogen; fluoro; chloro; bromo; C1-C6 alkyl; C1-C6 alkoxy; nitro; cyano; and trifluoromethyl;

with the proviso that

- (a) R_1 , R_2 , and R_3 are not all hydrogen when R_4 is 2-chloro or 4-chloro;
- (b) when R_1 and R_3 are hydrogen and R_2 is 4-bromo or 4-chloro, then R_4 may not be 4-chloro, 4-fluoro or 4-bromo;
- (c) when R_1 and R_3 are hydrogen and R_2 is 4-fluoro, R_4 may not be 4-fluoro or 4-bromo;
- (d) when R_1 is hydrogen, and R_4 is 2-fluoro, then R_2 and R_3 may not be 4-fluoro;
- (e) when R_1 is hydrogen and R_3 is 4-hydrogen, 4-chloro, 4-bromo, 4-methyl or 4-methoxy, and R_4 is 2-hydrogen, 2-chloro or 2-fluoro; then R_2 may not be 4-hydrogen, 4-chloro, 4-fluoro or 4-bromo.

According to a related invention, novel compounds are provided according to formula III:

- 7 -



wherein

R_1 , R_2 , R_3 , and R_4 are independently selected from the group consisting of hydrogen; fluoro; chloro; bromo; iodo; C1-C6 alkyl; C1-C6 alkoxy; nitro; cyano; and trifluoromethyl;

5 provided at least one of R_1 , R_2 , R_3 , and R_4 is iodo.

According to a preferred embodiment, at least one of R_1 and R_2 in formula III is other than hydrogen and is located at the 2- or 4-position of the phenyl ring to which it is attached; and at least one of R_3 and R_4 is other than hydrogen and is located at the 2- or 4-position of the phenyl ring to which it is attached. According to a further preferred embodiment, R_2 and R_4 in formula III are hydrogen, and R_1 and R_3 are located at the 4-position of the respective phenyl rings to which they are attached. According to a further preferred embodiment, one of R_1 or R_3 is selected from the group consisting of chloro, fluoro, bromo and nitro, the other of R_1 or R_3 being iodo.

15 A pharmaceutical composition is also provided comprising a pharmaceutically acceptable carrier and one or more compounds of formula III above, wherein R_1 , R_2 , R_3 , and R_4 are defined as above for formula III.

20 Where R_1 , R_2 , R_3 or R_4 is an alkyl or alkoxy group in any compound of formulae I, II or III, the carbon chain may be branched or straight, with straight being preferred. Preferably, the alkyl and alkoxy groups comprise C1-C3 alkyl and C1-C4 alkoxy, most preferably methyl and methoxy.

- 8 -

According to another embodiment of the invention, a method of treating an individual for a proliferative disorder, particularly cancer, is provided, comprising administering to said individual an effective amount of a compound according to formula II or III, alone or in combination with a pharmaceutically acceptable carrier.

In another embodiment of the invention, a method of inhibiting growth of tumor cells in an individual afflicted with cancer is provided comprising administering to said individual an effective amount of a compound according to formula II or III, alone or in combination with a pharmaceutically acceptable carrier.

In another embodiment, a method of inducing apoptosis of cancer cells, more preferably tumor cells, in an individual afflicted with cancer is provided, comprising administering to said individual an effective amount of a compound according to formula II or III, alone or in combination with a pharmaceutically acceptable carrier.

In yet another embodiment of the present invention, benzyl sulfones having the structural formula II or III, may be utilized as monomers in the synthesis of a new class of polymers having pendant benzylsulfone groups.

The present invention also provides a series of substituted benzylsulfonyl acetic acid compounds having structural formula V, below. The substituted benzylsulfonyl acetic acid compounds are useful as intermediates in the synthesis of novel (E)-styryl benzylsulfone compounds of formula I, according to Method A, below.

Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

- 8a -



The reference to any prior art in this specification is not, and should not be taken as, an acknowledgment or any form of suggestion that that prior art forms part of the common general knowledge in Australia.

5

Detailed Description of the Invention

According to the present invention, certain (E)-styryl benzylsulfone derivatives selectively kill various tumor cell types without killing normal cells. With wishing to be bound by any theory, it is believed that the compounds affect the MAPK signal transduction pathway,

10



thereby affecting tumor cell growth and viability. This cell growth inhibition is associated with regulation of the ERK and JNK types of MAPK. Without wishing to be bound by any theory, the styryl sulfones of the present invention may block the phosphorylating capacity of ERK-2.

5 The compounds of the invention have been shown to inhibit the proliferation of tumor cells by inducing cell death. The compounds are believed effective against a broad range of tumor types, including but not limited to the following: breast, prostate, ovarian, lung, colorectal, brain (i.e., glioma) and renal. The compounds are also believed effective against
10 leukemic cells. The compounds do not kill normal cells in concentrations at which tumor cells are killed.

 The compounds are also useful in the treatment of non-cancer proliferative disorders, including but not limited to the following: hemangiomatosis in new born, secondary progressive multiple sclerosis,
15 chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid formation, Pagets Disease of the bone, fibrocystic disease of the breast, Peronies and Duputren's fibrosis, restenosis and cirrhosis.

 Treatment of this broad range of tumor cells with the styryl
20 benzylsulfone compounds of the invention leads to inhibition of cell proliferation and induction of apoptotic cell death. In breast tumors, the effect is observed for estrogen receptor (ER) positive as well as estrogen receptor negative cells.

 Tumor cells treated with the compounds of the invention
25 accumulate in the G2/M phase of the cell cycle. As the cells exit the G2/M phase, they appear to undergo apoptosis. Treatment of normal cells with the styryl sulfones does not result in apoptosis.

Synthesis of (E)- Styryl Benzylsulfones

 The styryl benzylsulfones are characterized by cis-trans
30 isomerism resulting from the presence of one or more double bonds. The

- 10 -


compounds are named according to the Cahn-Ingold-Prelog system, the IUPAC 1974 Recommendations, Section E: Stereochemistry, in *Nomenclature of Organic Chemistry*, John Wiley & Sons, Inc., New York, NY, 4th ed., 1992, p. 127-138. Stearic relations around a double bond are

5 designated as "Z" or "E".

(E)-styryl benzylsulfones are prepared by Knoevenagel condensation of aromatic aldehydes with benzylsulfonyl acetic acids. The procedure is described by Reddy *et al.*, *Acta. Chim. Hung.* **115**:269 (1984); Reddy *et al.*, *Sulfur Letters* **13**:83 (1991); Reddy *et al.*, *Synthesis* **322**
10 (1984); and Reddy *et al.*, *Sulfur Letters* **7**:43 (1987), the entire disclosures of which are incorporated herein by reference.

The (E)-styryl benzylsulfones can be prepared according to either of the following Methods A and B:

- 11 -



- 12 -


METHOD A

A benzyl thioacetic acid V formed by the reaction of sodium thioglycollate and a benzyl chloride IV. The benzyl thioacetic acid V is oxidized with 30% hydrogen peroxide to give a corresponding benzylsulfonyl acetic acid VI. Condensation of VI with an aromatic aldehyde VII via a Knoevenagel reaction in the presence of benzylamine and glacial acetic acid yields the (E)-styryl benzylsulfone I, II or III.

METHOD B

A benzylthioacetic acid V is formed by the reaction of the appropriate sodium benzylthiolate VIII with chloroacetic acid. Oxidation of V to the corresponding benzylsulfonyl acetic acid VI and subsequent Knoevenagel condensation with aldehyde VII is carried out as in Method A.


Substituted benzylsulfonyl acetic acid compounds Va, Vb, Vc, and Vd according to formula V were prepared by reacting the corresponding benzyl chloride with thioglycolic acid under basic conditions (Method A). These compounds are novel intermediates.



No.	Compound	R ₁	R ₂	M.P. (°C)
Va	4-nitrobenzylsulfonyl acetic acid	H	4-NO ₂	165-166
Vb	4-trifluoromethylbenzylsulfonyl acetic acid	H	4-CF ₃	164-165
Vc	2,4-dichlorobenzylsulfonyl acetic acid	2-Cl	4-Cl	165-166
Vd	3,4-dichlorobenzylsulfonyl acetic acid	3-Cl	4-Cl	132-134

- 13 -

- (E)-Styryl benzylsulfones may be utilized as monomers in the synthesis of polymers X having pendant aryl and benzylsulfone groups. The polymerization of styryl benzylsulfones defined according to formula IX below into formula X polymers is accomplished by heating the formula IX compound above 250°C in the presence of a free radical initiator. The
- 5 IX compound above 250°C in the presence of a free radical initiator. The initiator may comprise benzoyl peroxide, for example:



- 14 -

The degree of polymerization in the polymer of formula X, "x", may range from about 10 to about 150, providing an oligomer or polymer of from 5,000 to 50,000 daltons. Other degrees of polymerization are also contemplated.

5 R₁, R₂, R₃, and R₄ in the monomer of formula IX, and in the polymer of formula X, are independently selected from the group consisting of hydrogen; halogen, i.e., fluoro, chloro, bromo and iodo, most preferably fluoro, chloro and bromo; C1-C6 alkyl; C1-C6 alkoxy; nitro; cyano; and trifluoromethyl.

10 The (E)-styryl benzylsulfones may be derivatized with a chemical group to permit conjugation to a carrier molecule, for the purpose of raising antibodies to the styryl sulfones. Suitable derivatizing chemistries are well-known to those skilled in the art. Preferably, the derivative comprises a carboxylic acid derivative. The carrier may comprise any molecule sufficiently large to be capable of generating an immune
15 response in an appropriate host animal. One such preferred carrier is keyhole limpet haemocyanin (KLH).

Therapeutic Administration

The (E)-styryl benzylsulfones of the invention may be administered in the form of a pharmaceutical composition, in combination
20 with a pharmaceutically acceptable carrier. The active ingredient in such formulations may comprise from 0.1 to 99.99 weight percent. By "pharmaceutically acceptable carrier" is meant any carrier, diluent or excipient which is compatible with the other ingredients of the formulation and to deleterious to the recipient.

25 The compounds of the invention may be administered to individuals (mammals, including animals and humans) afflicted with cancer. The compounds are also useful in the treatment of non-cancer proliferative disorders, that is, proliferative disorders which are characterized by benign indications. Such disorders may also be known as "cytoproliferative" or
30 "hyperproliferative" in that cells are made by the body at an atypically

elevated rate. Such disorders include, but are not limited to, the following:
hemangiomas in new born, secondary progressive multiple sclerosis,
chronic progressive myelodegenerative disease, neurofibromatosis,
ganglioneuromatosis, keloid formation, Pagets Disease of the bone,
5 fibrocystic disease of the breast, Peronies and Duputren's fibrosis,
restenosis and cirrhosis.

The compounds may be administered by any route, including
oral and parenteral administration. Parenteral administration includes, for
example, intravenous, intramuscular, intraarterial, intraperitoneal,
10 intranasal, rectal, intravaginal, topical or subcutaneous administration. The
active agent is preferably administered with a pharmaceutically acceptable
carrier selected on the basis of the selected route of administration and
standard pharmaceutical practice.

The active agent may be formulated into dosage forms
15 according to standard practices in the field of pharmaceutical preparations.
See Gennaro Alphonso, ed., *Remington's Pharmaceutical Sciences*, 18th
Ed., (1990) Mack Publishing Co., Easton, PA. Suitable dosage forms may
comprise, for example, tablets, capsules, solutions, parenteral solutions,
troches, suppositories, or suspensions.

20 For parenteral administration, the active agent may be mixed
with a suitable carrier or diluent such as water, an oil (particularly a
vegetable oil), ethanol, saline solution, aqueous dextrose (glucose) and
related sugar solutions, glycerol, or a glycol such as propylene glycol or
polyethylene glycol. Solutions for parenteral administration preferably
25 contain a water soluble salt of the active agent. Stabilizing agents,
antioxidizing agents and preservatives may also be added. Suitable
antioxidizing agents include sulfite, ascorbic acid, citric acid and its salts,
and sodium EDTA. Suitable preservatives include benzalkonium chloride,
methyl- or propyl-paraben, and chlorbutanol. The composition for
30 parenteral administration may take the form of an aqueous or nonaqueous
solution, dispersion, suspension or emulsion.

- 16 -

For oral administration, the active agent may be combined with one or more solid inactive ingredients for the preparation of tablets, capsules, pills, powders, granules or other suitable oral dosage forms. For example, the active agent may be combined with at least one excipient
5 such as fillers, binders, humectants, disintegrating agents, solution retarders, absorption accelerators, wetting agents, absorbents or lubricating agents. According to one tablet embodiment, the active agent may be combined with carboxymethylcellulose calcium, magnesium stearate, mannitol and starch, and then formed into tablets by conventional tableting
10 methods.

The specific dose of compound according to the invention to obtain therapeutic benefit will, of course, be determined by the particular circumstances of the individual patient including, the size, weight, age and sex of the patient, the nature and stage of the disease, the aggressiveness
15 of the disease, and the route of administration. For example, a daily dosage of from about 0.05 to about 50 mg/kg/day may be utilized. Higher or lower doses are also contemplated.

Examples

General Procedure for Synthesis (E)-Styryl Benzylsulfones

20 **Part A.** To a solution of (8g, 0.2 mol) sodium hydroxide in methanol (200 ml), thioglycolic acid (0.1 mol) is added slowly and the precipitate formed is dissolved by stirring the contents of the flask. Then an appropriately substituted or unsubstituted benzyl chloride (0.1 mol) is added stepwise and the reaction mixture is refluxed for 2-3 hours. The
25 cooled contents are poured onto crushed ice and neutralized with dilute hydrochloric acid (200 ml). The resulting corresponding benzylthioacetic acid (0.1 mol) is subjected to oxidation with 30% hydrogen peroxide (0.12 mol) in glacial acetic acid (125 ml) by refluxing for 1 hour. The contents are

- 17 -

cooled and poured onto crushed ice. The separated solid is recrystallized from hot water to give the corresponding pure benzylsulfonylacetic acid.

Part B. A mixture of the benzylsulfonyl acetic acid (10 mmol), an appropriately substituted or unsubstituted aromatic aldehyde (10 mmol), and benzylamine (200 μ l) in glacial acetic acid (12 ml) is refluxed for 2-3 hours. The contents are cooled and treated with cold ether (50 ml). Any product precipitated out is separated by filtration. The filtrate is diluted with more ether and washed successively with a saturated solution of sodium bicarbonate (20 ml), sodium bisulfite (20 ml), dilute hydrochloric acid (20 ml) and finally with water (35 ml). Evaporation of the dried ethereal layer yields styryl benzylsulfones as a solid material.

In each of the following examples, the substituted benzylsulfonyl acetic acid was made according to Part A of the General Procedure. All the styryl benzylsulfone compounds were recrystallized from 2-propanol and the purity was checked by thin layer chromatography.

Example 1

(E)-4-Fluorostyryl-4-trifluoromethylbenzylsulfone

A solution of 4-trifluorobenzylsulfonylacetic acid (10 mmol) and 4-fluorobenzaldehyde (10mmol) was subjected to the General Procedure, Part B. The title compound melting point 166-168°C, was obtained in 82% yield.

Example 2

(E)-4-Chlorostyryl-4-trifluoromethylbenzylsulfone

A solution of 4-trifluoromethylbenzylsulfonylacetic acid (10 mmol) and 4-chlorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 164-168°C, was obtained in 88% yield.

Example 3**(E)-4-Bromostyryl-4-trifluoromethylbenzylsulfone**

5 A solution of 4-trifluoromethylbenzylsulfonylacetic acid (10 mmol) and 4-bromobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 181-183°C, was obtained in 85% yield.

Example 4**(E)-4-Fluorostyryl-2,4-dichlorobenzylsulfone**

10 A solution of 2,4-dichlorobenzylsulfonyl acid (10 mmol) and 4-fluorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 146-148°C, was obtained in 78% yield.

Example 5**(E)-4-Chlorostyryl-2,4-dichlorobenzylsulfone**

15 A solution of 2,4-dichlorobenzylsulfonylacetic acid (10 mmol) and 4-chlorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 148-149°C, was obtained in 84% yield.

Example 6**20 (E)-4-Fluorostyryl-3,4-dichlorobenzylsulfone**

A solution of 3,4-dichlorobenzylsulfonylacetic acid (10 mmol) and 4-fluorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 120-122°C, was obtained in 82% yield.

- 19 -

Example 7**(E)-4-Chlorostyryl-3,4-dichlorobenzylsulfone**

5 A solution of 3,4-dichlorobenzylsulfonylacetic acid (10 mmol) and 4-chlorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 149-151°C, was obtained in 86% yield.

Example 8**(E)-4-Bromostyryl-3,4-dichlorobenzylsulfone**

10 A solution of 3,4-dichlorobenzylsulfonylacetic acid (10 mmol) and 4-bromobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 154-155°C, was obtained in 84% yield.

Example 9**(E)-4-Fluorostyryl-4-nitrobenzylsulfone**

15 A solution of 4-nitrobenzylsulfonylacetic acid (10 mmol) and 4-fluorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 160-161°C, was obtained in 76% yield.

Example 10**20 (E)-4-Fluorostyryl-4-cyanobenzylsulfone**

A solution of 4-cyanobenzylsulfonylacetic acid (10 mmol) and 4-fluorobenzaldehyde (10 mmol) was subjected to the General Procedure Part B. The title compound, melting point 150-151°C, was obtained in 82% yield.

- 20 -

Example 11**(E)-4-Chlorostyryl-4-cyanobenzylsulfone**

5 A solution of 4-cyanobenzylsulfonyl acetic acid (10 mmol) and 4-chlorobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 173-177°C, was obtained in 86% yield.

Example 12**(E)-4-Bromostyryl-4-cyanobenzylsulfone**

10 A solution of 4-cyanobenzylsulfonyl acetic acid (10 mmol) and 4-bromobenzaldehyde (10 mmol) was subjected to the General Procedure, Part B. The title compound, melting point 183-184°C, was obtained in 77% yield.

Example 13**(E)-3,4-Difluorostyryl-4-chlorobenzylsulfone**

15 A solution of 4-chlorobenzylsulfonyl acetic acid (10 mmol) and 3,4 difluorobenzaldehyde was subjected to the General Procedure, Part B. The title compound, melting point 204-205°C, was obtained in 73% yield.

Example 14**(E)-3-Chloro-4-fluorostyryl-4-chlorobenzylsulfone**

20 A solution of 4-chlorobenzylsulfonylacetic acid (10 mmol) and 3-chloro-4-fluorobenzaldehyde was subjected to the General Procedure, Part B. The title compound, melting point 181-183°C, was obtained in 78% yield.

Example 15**25 (E)-2-Chloro-4-fluorostyryl-4-chlorobenzylsulfone**

A solution of 4-chlorobenzylsulfonylacetic acid (10 mmol) and 2-chloro-4-fluorobenzaldehyde was subjected to the General Procedure,

- 21 -

Part B. The title compound, melting point 149-150°C, was obtained in 68% yield.

Example 16**(E)-2,4-Dichlorostyryl-4-chlorobenzylsulfone**

5 A solution of 4-chlorobenzylsulfonyl acetic acid (10 mmol) and 2,4-dichlorobenzaldehyde was subjected to the General Procedure, Part B. The title compound, melting point 164-165°C, was obtained in 78% yield.

Example 17**(E)-3,4-Dichlorostyryl-4-chlorobenzylsulfone**

10 A solution of 4-chlorobenzylsulfonyl acetic acid (10 mmol) and 3,4-dichlorobenzaldehyde (10 mmol) was subjected to the General procedure, Part B. The title compound, melting point 170-171°C, was obtained in 73% yield.

Example 18**(E)-2,3-Dichlorostyryl-4-chlorobenzylsulfone**

15 A solution of 4-chlorobenzylsulfonyl acetic acid (10 mmol) and 2,3-dichlorobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 170-171°C, was obtained in 72% yield.

Example 19**(E)-4-Fluorostyryl-4-iodobenzylsulfone**

20 A solution of 4-iodobenzylsulfonyl acetic acid (10 mmol) and 4-fluorobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 171-173°C, was obtained in 98%
25 yield. (1H NMR, CDCl₃) δ 4.27 (s, CH₂), 6.60 (d, = CH, J = 15.7 Hz), 7.18-7.80 (m, 9H, Aro + = CH).

- 22 -

Example 20**(E)-4-Iodostyryl-4-fluorobenzylsulfone**

A solution of 4-fluorobenzylsulfonyl acetic acid (10 mmol) and 4-iodobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 168-170°C, was obtained in 58% yield.

Example 21**(E)-4-Iodostyryl-4-chlorobenzylsulfone**

A solution of 4-chlorobenzylsulfonyl acetic acid (10 mmol) and 4-iodobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 181-182°C, was obtained in 70% yield. (1H NMR, CDCl₃) δ 4.27 (s, CH₂), 6.60 (d, =CH, J = 15.7 Hz), 7.18-7.80 (m, 9H, Aro + =CH).

Example 22**(E)-4-Iodostyryl-4-bromobenzylsulfone**

A solution of 4-bromobenzylsulfonyl acetic acid (10 mmol) and 4-iodobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 201-203°C, was obtained in 71% yield.

Example 23**(E)-4-Chlorostyryl-4-iodobenzylsulfone**

A solution of 4-iodobenzylsulfonyl acetic acid (10 mmol) and 4-chlorobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 200-202°C, was obtained in 86% yield. (1H NMR, CDCl₃) δ 4.27 (s, CH₂), 6.60 (d, =CH, J = 15.7 Hz), 7.18-7.80 (m, 9H, Aro + =CH).

- 23 -

Example 24**(E)-4-Bromostyryl-4-iodobenzylsulfone**

5 A solution of 4-iodobenzylsulfonyl acetic acid (10 mmol) and 4-bromobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 217-219°C, was obtained in 88% yield.

Example 25**(E)-2-Nitrostyryl-4-iodobenzylsulfone**

10 A solution of 4-iodobenzylsulfonyl acetic acid (10 mmol) and 2-nitrobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 227-229°C, was obtained in 62% yield.

Example 26**(E)-4-Nitrostyryl-4-iodobenzylsulfone**

15 A solution of 4-iodobenzylsulfonyl acetic acid (10 mmol) and 4-nitrobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 227-228°C, was obtained in 62% yield.

Example 27

20 **(E)-4-Iodostyryl-4-methoxybenzylsulfone**

A solution of 4-methoxybenzylsulfonyl acetic acid (10 mmol) and 4-iodobenzaldehyde (10 mmol) was subjected to the General Procedure, part B. The title compound, melting point 201-203°C, was obtained in 56% yield.

25

Example 28**(E)-4-Iodostyryl-2,4-dichlorobenzylsulfone**

A solution of 2,4-dichlorobenzylsulfonyl acetic acid (10 mmol) and 4-iodobenzaldehyde (10 mmol) was subjected to the General Procedure, part

B. The title compound, melting point 181-182°C, was obtained in 60% yield.

The following additional compounds in Table 1 were prepared according to the same synthetic methods (M.P. = melting point):

5

Table 1

Ex.	M.P. (°C)	Yield (%)	Compound
29	134-136	55	(E)-2-nitrostyryl-4-fluorobenzylsulfone
30	170-173	64	(E)-3-nitrostyryl-4-fluorobenzylsulfone
31	151-152	61	(E)-4-nitrostyryl-4-fluorobenzylsulfone
10 32	96-98	54	(E)-2-trifluoromethylstyryl-4-fluorobenzylsulfone
33	117-119	55	(E)-3-trifluoromethylstyryl-4-fluorobenzylsulfone
34	125-128	73	(E)-4-trifluoromethylstyryl-4-fluorobenzylsulfone
35	108-112	52	(E)-2-trifluoromethyl-4-fluorostyryl-4-fluorobenzylsulfone
36	128-132	58	(E)-2-nitrostyryl-4-chlorobenzylsulfone
15 37	156-157	60	(E)-3-nitrostyryl-4-chlorobenzylsulfone
38	189-191	61	(E)-4-nitrostyryl-4-chlorobenzylsulfone
39	100-101	55	(E)-2-trifluoromethylstyryl-4-chlorobenzylsulfone
40	155-157	58	(E)-3-trifluoromethylstyryl-4-chlorobenzylsulfone
41	164-166	59	(E)-4-trifluoromethylstyryl-4-chlorobenzylsulfone
20 42	115-117	63	(E)-2-trifluoromethyl-4-fluorostyryl-4-chlorobenzylsulfone
43	169-171	63	(E)-3-methyl-4-fluorostyryl-4-chlorobenzylsulfone
44	136-138	57	(E)-2-nitrostyryl-2,4-dichlorobenzylsulfone
45	136-138	57	(E)-2-trifluoromethyl-4-fluorostyryl-2,4-dichlorobenzylsulfone
46	131-132	63	(E)-2-nitrostyryl-4-bromobenzylsulfone
25 47	168-170	56	(E)-3-nitrostyryl-4-bromobenzylsulfone
48	205-207	67	(E)-4-nitrostyryl-4-bromobenzylsulfone
49	102-104	57	(E)-2-trifluoromethylstyryl-4-bromobenzylsulfone
50	160-161	55	(E)-3-trifluoromethylstyryl-4-fluorobenzylsulfone

- 25 -

5	51	174-175	62	(E)-4-trifluoromethylstyryl-4-bromoyanobenzylsulfone
	52	167-168	63	(E)-2-nitrostyryl-4-cyanobenzylsulfone
	53	192-193	62	(E)-3-nitrostyryl-4-cyanobenzylsulfone
	54	219-220	66	(E)-4-nitrostyryl-4-cyanobenzylsulfone
	55	182-184	70	(E)-4-fluorostyryl-4-methylbenzylsulfone
10	56	191-192	70	(E)-4-bromostyryl-4-methylbenzylsulfone
	57	128-130	51	(E)-2-nitrostyryl-4-methylbenzylsulfone
	58	201-203	56	(E)-3-nitrostyryl-4-methylbenzylsulfone
	59	194-195	57	(E)-4-nitrostyryl-4-methylbenzylsulfone
	60	148-149	60	(E)-4-fluorostyryl-4-methoxybenzylsulfone
15	61	176-177	66	(E)-4-chlorostyryl-4-methoxybenzylsulfone
	62	179-181	60	(E)-4-bromostyryl-4-methoxybenzylsulfone
	63	127-129	57	(E)-2-nitrostyryl-4-methoxybenzylsulfone
	64	153-155	59	(E)-3-nitrostyryl-4-methoxybenzylsulfone
	65	179-181	56	(E)-4-nitrostyryl-4-methoxybenzylsulfone
30	66	176-177	66	(E)-4-chlorostyryl-4-nitrobenzylsulfone
	67	199-200	60	(E)-4-fluorostyryl-4-nitrobenzylsulfone

Effect of (E)-Styryl Benzylsulfones on Breast, and Prostate Tumor Cell Lines

20 A. Cells.

The effect of the (E)-styryl benzylsulfones on normal fibroblasts and on tumor cells of breast, and prostate origin was examined utilizing one or more of the following cell lines: breast tumor cell lines MCF-7 and BT-20; prostate tumor cell line DU-145; colorectal carcinoma cell line DLD-1; non-small cell lung carcinoma cell line H157; and NIH/3T3 and HFL cells. MCF-7 is estrogen-responsive, while BT-20 is an estrogen-unresponsive cell line. NIH/3T3 and HFL are normal murine and human fibroblasts, respectively. MCF-7, BT-20, DLD-1 and H157 were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum supplemented with penicillin and streptomycin. DU145 was cultured in RPMI with 10%

- 26 -

fetal bovine serum containing penicillin and streptomycin. NIH3T3 and HFL cells were grown in DMEM containing 10% calf serum supplemented with penicillin and streptomycin. All cell cultures were maintained at 37°C in a humidified atmosphere of 5% CO₂.


5 B. Treatment with (E)-Styryl Sulfones and Viability Assay

Cells were treated with test compound at 2.5 mM concentration and cell viability was determined after 96 hours by the Trypan blue exclusion method. The results are set forth in Table 2. Activity for each compound is reported as a range of cell induced death (% Death) with the lowest activity in the range of 5-10% and the highest being above 80%.

10

Normal cells HFL and NIH 3T3 were treated with the same compounds in Table 2 under the same conditions of concentration and time. The normal cells displayed 5% growth inhibition but no appreciable cell death.

Table 2
Effect of (E)-styryl benzylsulfones on tumor cells



	Ex	R ₁	R ₂	R ₃	R ₄	Tumor cell type			H157	BT20
						MCF-7	DU145	DLD-1		
5	1	4-CF ₃	H	4-F	H	+++	+++	+++	++	+++
	2	4-CF ₃	H	4-Cl	H	++	++	++	ND	++
	3	4-CF ₃	H	4-Br	H	ND	+	+	ND	++
	4	2-Cl	4-Cl	4-F	H	++	++	ND	+	++
	5	2-Cl	4-Cl	4-Cl	H	+++	+	++	++	ND
10	6	3-Cl	4-Cl	4-F	H	++++	++++	ND	ND	++++
	7	3-Cl	4-Cl	4-Cl	H	ND	-	ND	ND	++
	8	3-Cl	4-Cl	4-Br	H	ND	-	-	ND	-
	9	4-NO ₂	H	4-F	H	++	++	-	++	++
	10	4-CN	H	4-F	H	+++	++	ND	+++	ND
15	11	4-CN	H	4-Cl	H	+	+	ND	+	ND
	12	4-CN	H	4-Br	H	-	-	-	+	ND
	13	4-Cl	H	3-F	4-F	-	-	-	-	ND
	14	4-Cl	H	3-Cl	4-F	-	-	-	-	ND
	15	4-Cl	H	2-Cl	4-F	++++	++++	++++	++++	ND
20	16	4-Cl	H	2-Cl	4-Cl	++++	+++	++++	ND	ND
	17	4-Cl	H	3-Cl	4-Cl	+	+	+	ND	ND
	18	4-Cl	H	2-Cl	3-Cl	+	+	+	ND	ND
	19	4-I	H	4-F	H	ND	++++	++++	++++	++++
	20	4-F	H	4-I	H	ND	ND	ND	ND	ND
25	21	4-Cl	H	4-I	H	ND	++	++	++	++
	22	4-Br	H	4-I	H	ND	++	++	++	++

- 28 -

5	23	4-I	H	4-Cl	H	ND	++	++	++	+++
	24	4-I	H	4-Br	H	ND	ND	ND	ND	ND
	25	4-I	H	2-NO ₂	H	ND	++	++	++	++
	26	4-I	H	4-NO ₂	H	ND	ND	ND	ND	ND
	27	4-CH ₃ O	H	4-I	H	ND	ND	ND	ND	ND
10	28	4-Cl	2-Cl	4-I	H	ND	ND	ND	ND	ND
	29	4-F	H	2-NO ₂	H	ND	+	+	+	+
	30	4-F	H	3-NO ₂	H	ND	+	+	+	+
	31	4-F	H	4-NO ₂	H	ND	++++	++++	++++	++++
	32	4-F	H	2-CF ₃	H	ND	+	+	+	+
15	33	4-F	H	3-CF ₃	H	ND	+	+	+	+
	34	4-F	H	4-CF ₃	H	ND	+	+	+	+
	35	4-F	H	2-CF ₃	4-F	ND	+	+	+	+
	36	4-Cl	H	2-NO ₂	H	ND	+	+	+	+
	37	4-Cl	H	3-NO ₂	H	ND	+	+	+	+
20	38	4-Cl	H	4-NO ₂	H	ND	++++	++++	++++	++++
	39	4-Cl	H	2-CF ₃	H	ND	+	+	+	+
	40	4-Cl	H	3-CF ₃	H	ND	+	+	+	+
	41	4-Cl	H	4-CF ₃	H	ND	+	+	+	+
	42	4-Cl	H	4-F	2-CF ₃	ND	+	+	+	+
25	43	4-Cl	H	4-F	3-CH ₃	ND	+	+	+	+
	44	4-Cl	2-Cl	2-NO ₂	H	ND	+	+	+	+
	45	4-Cl	2-Cl	4-F	2-CF ₃	ND	+++	+++	+++	+++
	46	4-Br	H	2-NO ₂	H	ND	+	+	+	+
	47	4-Br	H	3-NO ₂	H	ND	+	+	+	+
30	48	4-Br	H	4-NO ₂	H	ND	++++	++++	++++	++++
	49	4-Br	H	2-CF ₃	H	ND	+	+	+	+
	50	4-Br	H	3-CF ₃	H	ND	+	+	+	+
	51	4-Br	H	4-CF ₃	H	ND	+	+	+	+
	52	4-CN	H	2-NO ₂	H	ND	+	+	+	+
35	53	4-CN	H	3-NO ₂	H	ND	+	+	+	+
	54	4-CN	H	4-NO ₂	H	ND	++++	++++	++++	++++
	55	4-CH ₃	H	4-F	H	ND	+	+	+	+
	56	4-CH ₃	H	4-Br	H	ND	+	+	+	+

- 29 -

5	57	4-CH ₃	H	2-NO ₂	H	ND	+	+	+	+
	58	4-CH ₃	H	3-NO ₂	H	ND	+	+	+	+
	59	4-CH ₃	H	4-NO ₂	H	ND	+	+	+	+
	60	4-CH ₃ O	H	4-F	H	ND	++++	++++	++++	++++
	61	4-CH ₃ O	H	4-Cl	H	ND	++++	++++	++++	++++
10	62	4-CH ₃ O	H	4-Br	H	ND	++++	++++	++++	++++
	63	4-CH ₃ O	H	2-NO ₂	H	ND	+	+	+	+
	64	4-CH ₃ O	H	3-NO ₂	H	ND	+	+	+	+
	65	4-CH ₃ O	H	4-NO ₂	H	ND	++++	++++	++++	++++
	66	4-NO ₂	H	4-Cl	H	ND	+	+	+	+
	67	4-NO ₂	H	4-F	H	ND	+	+	+	+

Cell death:

15	0%	=	-	50-60%	=	++++
	5-10%	=	+	Above 80%	=	+++++
	10-15%	=	++	ND	=	not done
	40-50%	=	+++			

Example 68**Conjugation of (E)-4-Fluorostyryl 4-chlorobenzylsulfone to Keyhole Limpet Haemocyanin**

20 A carboxylic acid derivative of (E)-4-fluorostyryl 4-chlorobenzylsulfone was synthesized by mixing 4-chlorobenzyl sulfonyl acetic acid (10 mmol), 4-fluorobenzaldehyde (10mmol), glacial acetic acid (15 ml) and piperidine (0.5 ml) at room temperature (22°C) over a magnetic stirrer for 5 hours. The mixture was then diluted with ether and the ethereal

25 layer was washed with water. Evaporation of the ethereal layer yielded a semisolid material which on treating with 2-propanol gave a white solid. Recrystallization with 2-propanol gave 2-(4-chlorobenzyl sulfonyl)-3-(4-fluorophenyl)propenoic acid as white crystals, (yield 32%), m.p 111-112°C.

30 The above carboxylic acid derivative (10 mM) was made into an active ester by treatment with 10 mM 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC) and 10 mM N-hydroxysuccinamide (NHS), and then cross-linked to KLH by mixing with 1

- 30 -


ml of a KLH water solution containing 500 mg KLH. The mixture was stirred at room temperature for 5-6 hours. The KLH conjugate was then separated by passing the mixture through a size exclusion column (PD 10, Pharmacia). The conjugate was then used to inject rabbits for raising
5 antibodies.

All references cited with respect to synthetic, preparative and analytical procedures are incorporated herein by reference.

The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof and,
10 accordingly, reference should be made to the appended claims, rather than to the foregoing specification, as indication the scope of the invention.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound of the formula:



wherein:

R_1 and R_2 are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, C1-C6 alkyl, C1-C6 alkoxy, nitro, cyano and trifluoromethyl;

R_3 and R_4 are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, nitro, cyano and trifluoromethyl;

at least one of R_1 and R_2 is other than hydrogen; and

at least one of R_3 and R_4 is other than hydrogen;

with the proviso that

(a) when R_1 and R_3 are hydrogen and R_2 is 4-chloro, then R_4 may not be 4-chloro, 4-fluoro, 4-bromo, or 4-nitro;

(b) when R_1 and R_3 are hydrogen and R_2 is 4-fluoro or 4-bromo, then R_4 may not be 4-fluoro, 4-bromo, or 4-chloro;

(c) when R_1 and R_3 are hydrogen and R_2 is 4-nitro, then R_4 may not be 4-chloro, 4-nitro, 4-bromo, or 4-fluoro;

(d) when R_1 and R_3 are hydrogen and R_2 is 4-methyl, then R_4 may not be 4-chloro, 4-bromo, 4-fluoro, or 2-chloro;

(e) when R_1 is hydrogen, then R_2 , R_3 and R_4 may not all be fluoro;

(f) when R₁ is hydrogen and R₃ is 2-fluoro, then R₂ and R₄ may not both be selected from the group consisting of 4-chloro, 4-bromo and 4-fluoro; and

(g) the compound is not (*E*)-2-chloro-4-fluorostyryl-4-fluorobenzyl sulfone, (*E*)-2-chloro-4-fluorostyryl-4-chlorobenzyl sulfone, (*E*)-2-chloro-4-fluorostyryl-4-bromobenzyl sulfone, (*E*)-2,4-dichlorostyryl-4-fluorobenzyl sulfone, (*E*)-2,4-dichlorostyryl-4-chlorobenzyl sulfone, (*E*)-2,4-dichlorostyryl-4-bromobenzyl sulfone, (*E*)-2-chloro-4-bromostyryl-4-fluorobenzyl sulfone, (*E*)-2-chloro-4-bromostyryl-4-chlorobenzyl sulfone, or (*E*)-2-chloro-4-bromostyryl-4-bromobenzyl sulfone.

2. A compound according to claim 1 wherein at least one of R₁ and R₂ is located at the 2-, 3- or 4- position of the phenyl ring to which it is attached; and


wherein at least one of R₃ and R₄ is located at the 2- or 4- position of the phenyl ring to which it is attached.

3. A compound according to claim 2 selected from the group consisting of (*E*)-4-fluorostyryl-4-trifluoromethylbenzylsulfone, (*E*)-2-trifluoromethyl-4-fluorostyryl-2,4-dichlorobenzylsulfone, (*E*)-4-fluorostyryl-4-nitrobenzylsulfone and (*E*)-4-fluorostyryl-4-cyanobenzylsulfone.

4. A compound according to claim 2 wherein at least one of R₁ and R₂ is chloro and at least one of R₃ and R₄ is chloro or fluoro.

5. A compound according to claim 4 selected from the group consisting of (*E*)-4-fluorostyryl-3,4-dichlorobenzylsulfone, and (*E*)-4-fluorostyryl-2,4-dichlorobenzylsulfone.

6. A compound according to claim 2 wherein R_2 is 4-halogen or 4-cyano, and R_4 is 4-nitro.
7. A compound according to claim 6 selected from the group consisting of (*E*)-4-nitrostyryl-4-fluorobenzylsulfone, (*E*)-4-nitrostyryl-4-bromobenzylsulfone and (*E*)-4-nitrostyryl-4-cyanobenzylsulfone.
8. A compound according to claim 2 wherein R_2 is 4-C1-C6 alkoxy, and R_4 is 4-halogen or 4-nitro.
9. A compound according to claim 8 selected from the group consisting of (*E*)-4-fluorostyryl-4-methoxybenzylsulfone, (*E*)-4-chlorostyryl-4-methoxybenzylsulfone, (*E*)-4-bromostyryl-4-methoxybenzylsulfone and (*E*)-4-nitrostyryl-4-methoxybenzylsulfone.
10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of the formula:



wherein:

R_1 and R_2 are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, C1-C6 alkyl, C1-C6 alkoxy, nitro, cyano and trifluoromethyl;

R_3 and R_4 are independently selected from the group consisting of hydrogen, fluoro, chloro, bromo, nitro, cyano and trifluoromethyl;

at least one of R₁ and R₂ is other than hydrogen; and

at least one of R₃ and R₄ is other than hydrogen;

with the proviso that

(a) when R₁ and R₃ are hydrogen and R₂ is 4-bromo or 4-chloro, R₄ may not be 4-chloro, 4-fluoro, or 4-bromo;

(b) when R₁ and R₃ are hydrogen and R₂ is 4-fluoro, R₄ may not be 4-fluoro or 4-bromo;

(c) when R₁ is hydrogen and R₄ is 2-fluoro, then R₂ and R₃ may not be 4-fluoro;

(d) when R₁ is hydrogen and R₃ is 4-hydrogen, 4-chloro, or 4-bromo, and R₄ is 2-hydrogen, 2-chloro or 2-fluoro, then R₂ may not be 4-hydrogen, 4-chloro, 4-fluoro, or 4-bromo; and

(e) the compound is not (*E*)-2-chloro-4-fluorostyryl-4-fluorobenzyl sulfone, (*E*)-2-chloro-4-fluorostyryl-4-chlorobenzyl sulfone, (*E*)-2-chloro-4-fluorostyryl-4-bromobenzyl sulfone, (*E*)-2, 4-difluorostyryl-4-chlorobenzyl sulfone, or (*E*)-2,4-difluorostyryl-4-bromobenzyl sulfone.

11. A composition according to claim 10 wherein the compound is selected from the group consisting of (*E*)-4-fluorostyryl-4-trifluoromethylbenzylsulfone, (*E*)-2-trifluoromethyl-4-fluorostyryl-2,4-dichlorobenzylsulfone, (*E*)-4-fluorostyryl-4-cyanobenzylsulfone, (*E*)-4-fluorostyryl-3,4-dichlorobenzylsulfone, (*E*)-4-fluorostyryl-2,4-dichlorobenzylsulfone, (*E*)-4-fluorostyryl-4-nitrobenzylsulfone, (*E*)-4-chlorostyryl-4-nitrobenzylsulfone, (*E*)-4-nitrostyryl-4-fluorobenzylsulfone, (*E*)-4-nitrostyryl-4-bromobenzylsulfone and (*E*)-4-nitrostyryl-4-cyanobenzylsulfone, (*E*)-4-fluorostyryl-4-methoxybenzylsulfone, (*E*)-4-chlorostyryl-4-methoxybenzylsulfone, (*E*)-4-bromostyryl-4-methoxybenzylsulfone and (*E*)-4-nitrostyryl-4-methoxybenzylsulfone.

12. A method of treating an individual for a proliferative disorder comprising administering to said individual an effective amount of a pharmaceutical composition according to claim 10.

13. A method according to claim 12 wherein the proliferative disorder is selected from the group consisting of hemangiomatosis in new born, secondary progressive multiple sclerosis, chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid formation, Pagets Disease of the bone, fibrocystic disease of the breast, Peronies and Duputren's fibrosis, restenosis and cirrhosis.


14. A method according to claim 12 wherein the proliferative disorder is cancer.

15. A method according to claim 14 wherein the cancer is selected from the group consisting of ovarian, breast, prostate, lung, renal, colorectal and brain cancers, or the cancer is a leukemia.

16. A method of inducing apoptosis of tumor cells in an individual afflicted with cancer comprising administering to said individual an effective amount of a pharmaceutical composition according to claim 10.

17. A method according to claim 16 wherein the tumor cells are selected from the group consisting of ovarian, breast, prostate, lung, renal, colorectal and brain tumors.

18. A compound useful as an intermediate in the synthesis of a compound according to claim 1, said intermediate compound having the formula



wherein:

R₁ and R₂ are both halogen; or

R₁ is hydrogen and R₂ is CF₃;

or a salt thereof.

19. A compound according to claim 18 wherein R₁ and R₂ are both chloro.

20. A compound according to claim 36 selected from the group consisting of 4-trifluoromethylbenzylsulfonyl acetic acid, 2,4-dichlorobenzylsulfonyl acetic acid and 3,4-dichlorobenzylsulfonyl acetic acid, and salts thereof.

21. A method of treating an individual for a proliferative disorder, said method comprising administering to said individual an effective amount of a compound according to any one of claims 1 to 9.

22. A method according to claim 21 wherein the proliferative disorder is selected from the group consisting of hemangiomatosis in new born, secondary progressive multiple sclerosis, chronic progressive myelodegenerative disease, neurofibromatosis, ganglioneuromatosis, keloid formation, Pagets Disease of the bone, fibrocystic disease of the breast, Peronies and Duputren's fibrosis, restenosis and cirrhosis.

23. A method according to claim 21 wherein the proliferative disorder is cancer.

24. A method according to claim 23 wherein the cancer is selected from the group consisting of ovarian, breast, prostate, lung, renal, colorectal and brain cancers, or the cancer is a leukemia.

25. A method of inducing apoptosis of tumor cells in an individual afflicted with cancer, said method comprising administering to said individual an effective amount of a compound according to any one of claims 1 to 9.

26. A method according to claim 25 wherein the tumor cells are selected from the group consisting of ovarian, breast, prostate, lung, renal, colorectal and brain tumors.

27. A method of inhibiting growth of tumor cells in an individual afflicted with cancer, said method comprising administering to said individual an effective amount of a compound according to any one of claims 1 to 9.

28. The method according to claim 27 wherein the tumor cells are selected from the group consisting of ovarian, breast, prostate, lung, colorectal, renal and brain tumors.

29. Use of a compound according to any one of claims 1 to 9 in the manufacture of a medicament to treat a proliferative disorder in an individual.

30. Use of a compound according to any one of claims 1 to 9 in the manufacture of a medicament to induce apoptosis of tumor cells in an individual.

31. Use of a compound according to any one of claims 1 to 9 in the manufacture of a medicament to inhibit growth of tumor cells in an individual.

5

10 Patent Attorneys for the Applicant