



(51) International Patent Classification:

C12N 5/071 (2010.01) C12N 15/117 (2010.01)

C12N 5/10 (2006.01) C12Q 1/68 (2018.01)

C12N 15/113 (2010.01)

(21) International Application Number:

PCT/US2019/028108

(22) International Filing Date:

18 April 2019 (18.04.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/659,281

18 April 2018 (18.04.2018)

US

(71) Applicant: THE TRUSTEES OF COLUMBIA UNIVERSITY IN THE CITY OF NEW YORK

[US/US]; 412 Low Memorial Library, 535 West 116th Street, New York, NY 10027 (US).

(72) Inventors: **HIRANO, Michio**; 120 Cabrini Blvd, Apt. 126, New York, NY 10033 (US). **AKMAN, Hasan, O.**; 454 Jefferson Avenue, Haworth, NJ 07641 (US). **LOPEZ-GOMEZ, Carlos**; 128 Fort Washington Avenue, Apt. 4A, New York, NY 10032 (US).

(74) Agent: **CARNEY, Bonnie, K.** et al.; Leason Ellis LLP, One Barker Avenue, Fifth Floor, White Plains, NY 10601 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(54) Title: GENE THERAPY FOR DISEASES CAUSED BY UNBALANCED NUCLEOTIDE POOLS INCLUDING MITOCHONDRIAL DNA DEPLETION SYNDROMES




FIG. 1

(57) Abstract: The invention relates generally to a method of treatment for a human genetic disease, such as diseases characterized by unbalanced nucleotide pools, e.g., mitochondrial DNA depletion syndromes, and more specifically, thymidine kinase 2 (TK2) deficiency, using gene therapy. The gene therapy may involve administration of one or more constructs, such as a viral vector, containing a nucleic acid encoding a functional protein. The functional protein may correspond to a nuclear gene. For treatment of TK2 deficiency, the gene therapy may involve administration of one or more constructs, such as a viral vector, containing a nucleic acid encoding a functional TK2 enzyme. The treatment may also involve the administration of pharmacological therapy in conjunction with the gene therapy. The treatment protocols of the disclosure, such as those involving gene therapy alone or in combination with pharmacological therapy, can be used to treat, prevent, and/or cure various other disorders of unbalanced nucleoside pools, especially those found in mitochondrial DNA depletion syndrome.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

GENE THERAPY FOR DISEASES CAUSED BY UNBALANCED NUCLEOTIDE POOLS INCLUDING MITOCHONDRIAL DNA DEPLETION SYNDROMES

CROSS-REFERENCE TO RELATED APPLICATIONS

5 This present application claims priority to U.S. Provisional Patent Application Ser. No. 62/659,281 filed April 18, 2018, which is incorporated herein by reference in its entirety.

STATEMENT OF GOVERNMENT SUPPORT

 This invention was made with government support under HD080642 awarded by NIH.
10 The government has certain rights in the invention.

FIELD OF THE INVENTION

 The invention relates to the field of gene therapy for the treatment of genetic diseases, including human diseases characterized by unbalanced nucleotide pools. Examples of such
15 diseases are mitochondrial DNA depletion syndromes, such as thymidine kinase 2 (TK2) deficiency.

BACKGROUND OF THE INVENTION

 Mitochondrial diseases are clinically heterogeneous diseases due to defects of the
20 mitochondrial respiratory chain (RC) and oxidative phosphorylation, the biochemical pathways that convert energy in electrons into adenosine triphosphate (ATP). The respiratory chain is comprised of four multi-subunit enzymes (complexes I-IV) that transfer electrons to generate a proton gradient across the inner membrane of mitochondria. The flow of protons down this concentration gradient through complex V subsequently drives ATP synthesis (DiMauro and
25 Schon 2003; DiMauro and Hirano 2005). Coenzyme Q₁₀ (CoQ₁₀) is an essential molecule that shuttles electrons from complexes I and II to complex III. The respiratory chain is unique in eukaryotic, *e.g.*, mammalian, cells by virtue of being controlled by two genomes: one composed of mitochondrial DNA (mtDNA) and another composed of nuclear DNA (nDNA). As a consequence, mutations in either genome can cause mitochondrial diseases.

30 Mitochondrial DNA depletion syndrome (MDS), which is a subgroup of mitochondrial disease, is a frequent cause of severe childhood encephalomyopathy characterized molecularly by reduction of mitochondrial DNA (mtDNA) copy number in tissues and insufficient synthesis of mitochondrial RC complexes (Hirano, et al. 2001). Mutations in several nuclear genes have been identified as causes of infantile MDS, including TK2, DGUOK, POLG,

POLG2, SCLA25A4, MPV17, RRM2B, SUCLA2, SUCLG1, TYMP, OPA1, and C10orf2 (PEO1). (Bourdon, et al. 2007; Copeland 2008; Elpeleg, et al. 2005; Mandel, et al. 2001; Naviaux and Nguyen 2004; Ostergaard, et al. 2007; Saada, et al. 2003; Sarzi, et al. 2007; Spinazzola, et al, 2006). In addition, mutations in these nuclear genes can also cause multiple
5 deletions of mtDNA with or without mtDNA depletion (Béhin, et al. 2012; Garone, et al. 2012; Longley, et al. 2006; Nishino, et al. 1999; Paradas, et al. 2012; Ronchi, et al. 2012; Spelbrink, et al. 2001; Tyynismaa, et al. 2012; Tyynismaa, et al. 2009; Van Goethem, et al. 2001).

Most mitochondrial diseases affect multiple body organs and are typically fatal in childhood or early adult life. There are no proven effective treatments or cures for
10 mitochondrial diseases, only supportive therapies. There remains a need for compositions and methods for treating and preventing mitochondrial diseases, such as MDS.

SUMMARY OF THE INVENTION

The present disclosure relates to compositions and methods that can be used to treat a
15 subject (e.g., a mammalian subject, such as a human subject) that has or is at risk of developing a disease characterized by unbalanced nucleotide pools. Exemplary pathologies that may be treated, prevented, and/or cured using the compositions and methods of the disclosure are mitochondrial diseases, including mitochondrial DNA depletion syndromes, such as deficiencies in a gene selected from TK2, encoding thymidine kinase 2 protein; DGUOK,
20 encoding deoxyguanosine kinase protein; RRM2B, encoding p53R2, p53 inducible subunit of ribonucleotide reductase; TYMP, encoding thymidine phosphorylase; SUCLA2, encoding succinate-CoA ligase ADP-forming beta subunit; SUCLG1, encoding succinate-CoA ligase ADP-forming alpha subunit; MPV17, encoding mitochondrial inner membrane protein MPV17; and POLG, encoding DNA polymerase gamma, catalytic subunit.

25 Using the compositions and methods of the disclosure, a subject (e.g., a mammalian subject, such as a human subject) that has or is at risk of developing a disease described above may be administered a composition containing a transgene encoding one or more of the foregoing proteins. The composition may be a vector, for example, a viral vector, such as an adeno-associated virus (AAV) vector. In some embodiments, the subject is administered a
30 second composition containing a transgene encoding one or more of the foregoing proteins. The second composition may be a vector, for example, a viral vector, such as an AAV vector. In some embodiments, the subject is additionally administered one or more subsequent compositions, such as a third, fourth, fifth, sixth, seventh, eighth, ninth, and/or tenth composition, containing a transgene encoding one or more of the foregoing proteins. The one

or more subsequent compositions may each be, independently, a vector, for example, a viral vector, such as an AAV vector.

In a first aspect, the disclosure features a method of treating, preventing, and/or curing a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof.

5 The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding thymidine kinase 2 (TK2), deoxyguanosine kinase (dGK), thymidine phosphorylase (TP), p53 inducible small subunit of ribonucleotide reductase (p53R2), succinyl-CoA ligase ADP-forming subunit beta (SUCLA2), succinyl-CoA ligase GDP-forming subunit alpha (SUCLG1), mitochondrial inner membrane protein MPV17
10 (MPV17), and/or DNA polymerase subunit gamma (POLG).

In a further aspect, the disclosure features a method of restoring enzyme activity in a subject having a disease or disorder characterized by unbalanced nucleotide pools. The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or
15 POLG.

In an additional aspect, the disclosure features a method of alleviating one or more symptoms associated with a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof. The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding TK2, dGK, TP, p53R2,
20 SUCLA2, SUCLG1, MPV17, and/or POLG.

In any of the aspects of this invention, the disclosure also provides a composition as described herein for use in a method as described herein. The disclosure also provides the use of a composition as described herein for the manufacture of a medicament for a method as described herein. The transgene may encode TK2, dGK, TP, p53R2, SUCLA2, SUCLG1,
25 MPV17, and/or POLG. For example, the transgene may encode TK2, dGK, TP, p53R2, SUCLA2, SUCLG1 and/or MPV17.

As part of the foregoing aspects, the disclosure therefore also provides a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for use in treating, preventing, and/or curing a disease or disorder characterized by
30 unbalanced nucleotide pools. Also provided is a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for use in restoring enzyme activity in a subject having a disease or disorder characterized by unbalanced nucleotide pools. Furthermore provided is a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for use in alleviating one or

more symptoms associated with a disease or disorder characterized by unbalanced nucleotide pools.

As part of the foregoing aspects, the disclosure also provides the use of a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for the manufacture of a medicament for treating, preventing, and/or curing a disease or disorder characterized by unbalanced nucleotide pools. Also provided is the use of a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for the manufacture of a medicament for restoring enzyme activity in a subject having a disease or disorder characterized by unbalanced nucleotide pools. Furthermore provided is the use of a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG for the manufacture of a medicament for alleviating one or more symptoms associated with a disease or disorder characterized by unbalanced nucleotide pools.

In some embodiments of any of the foregoing aspects, the disease or disorder is a mitochondrial disease, such as a mitochondrial DNA depletion syndrome (MDS). In some embodiments, the MDS is a myopathic MDS characterized by one or more mutations in an endogenous gene encoding TK2. In some embodiments, the MDS is an encephalomyopathic form characterized by one or more mutations in an endogenous gene encoding SUCLA2. In some embodiments, the MDS is a neurogastrointestinal encephalopathic form characterized by one or more mutations in an endogenous gene encoding TP. In some embodiments, the MDS is a hepatopathic form characterized by one or more mutations in an endogenous gene encoding dGK, MPV17, and/or POLG.

In some embodiments, the transgene encodes TK2. The TK2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from

1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by way of one or more conservative amino acid
 5 substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid
 10 sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 85%, 86%, 87%,
 15 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some
 20 embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TK2 is at least
 25 at least 70% identical to coding sequence of the transgene that encodes TK2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TK2 has the nucleic acid
 30 sequence of SEQ ID NO: 17.

In some embodiments, the transgene encodes dGK. The dGK may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In

some embodiments, the dGK has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding dGK is at least at least 70% identical to coding sequence of the transgene that encodes dGK (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%,

82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding dGK has the nucleic acid sequence of SEQ ID NO: 18.

5 In some embodiments, the transgene encodes TP. The TP may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5).
10 In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).
15 20

In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or
25 30

100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TP is at least at least 70% identical to coding sequence of the transgene that encodes TP (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TP has the nucleic acid sequence of SEQ ID NO: 19.

In some embodiments, the transgene encodes p53R2. The p53R2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%,

99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding p53R2 is at least at least 70% identical to coding sequence of the transgene that encodes p53R2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding p53R2 has the nucleic acid sequence of SEQ ID NO: 20.

In some embodiments, the transgene encodes SUCLA2. The SUCLA2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments,

the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

5 In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some
10 embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid
15 sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of
20 SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding SUCLA2 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLA2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%,
25 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLA2 has the nucleic acid sequence of SEQ ID NO: 21.

In some embodiments, the transgene encodes SUCLG1. The SUCLG1 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO:
30 11 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid

sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that differs
5 from SEQ ID NO: 11 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the SUCLG1 has an amino acid sequence that differs from SEQ ID NO: 11 by way of one or
10 more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic
15 acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is
20 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO:
25 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding
30 SUCLG1 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLG1 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding

sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLG1 has the nucleic acid sequence of SEQ ID NO: 22.

In some embodiments, the transgene encodes MPV17. The MPV17 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 13 (e.g.,
5 an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of
10 SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more amino acid substitutions, insertions, and/or deletions, such
15 as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or
20 more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%,
25 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%
30 identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that

is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding
5 MPV17 is at least at least 70% identical to coding sequence of the transgene that encodes MPV17 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding MPV17
10 has the nucleic acid sequence of SEQ ID NO: 23.

In some embodiments, the transgene encodes POLG. The POLG may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In
15 some embodiments, the POLG has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is
20 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23,
25 24, 25, or more, conservative amino acid substitutions). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

30 In some embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some

embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene
5 encoding POLG has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid
10 sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding POLG is at least at least 70% identical to coding sequence of the transgene that encodes POLG (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%,
15 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding POLG has the nucleic acid sequence of SEQ ID NO: 24.

In another aspect, the disclosure features a method of treating, preventing, and/or curing
20 TK2 deficiency in a subject in need thereof. The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding TK2.

In a further aspect, the disclosure features a method of restoring TK2 enzyme activity in a subject in need thereof. The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding TK2.

25 In another aspect, the disclosure features a method of alleviating one or more symptoms associated with a TK2 deficiency in a subject in need thereof. The method includes administering to the subject a therapeutically effective amount of a composition containing a transgene encoding TK2.

As part of these aspects, the disclosure also provides a composition containing a
30 transgene encoding TK2 for use in treating, preventing, and/or curing TK2 deficiency. Also provided is a composition containing a transgene encoding TK2 for use in restoring TK2 enzyme activity in a subject having a TK2 deficiency. Furthermore provided is a composition containing a transgene encoding TK2 for use in alleviating one or more symptoms associated with a TK2 deficiency. Typically the subject has a TK2 deficiency.

As part of these aspects, the disclosure also provides the use of a composition containing a transgene encoding TK2 for the manufacture of a medicament for treating, preventing, and/or curing TK2 deficiency. Also provided is the use of a composition containing a transgene encoding TK2 for the manufacture of a medicament for restoring TK2 enzyme
5 activity in a subject having a TK2 deficiency. Furthermore provided is the use of a composition containing a transgene encoding TK2 for the manufacture of a medicament for alleviating one or more symptoms associated with a TK2 deficiency. Typically the subject has a TK2 deficiency.

In some embodiments of any of the preceding three aspects, the subject has or is at risk
10 of developing a MDS. The TK2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 1 (e.g.,
15 an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino
20 acid sequence that differs from SEQ ID NO: 1 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by
25 way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is
30 at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 85% identical to the

nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TK2 is at least at least 70% identical to coding sequence of the transgene that encodes TK2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TK2 has the nucleic acid sequence of SEQ ID NO: 17.

In some embodiments of any of the above aspects of the disclosure, the composition comprises a vector, such as a viral vector. The viral vector may be, for example, an AAV, adenovirus, lentivirus, retrovirus, poxvirus, baculovirus, herpes simplex virus, vaccinia virus, or a synthetic virus (e.g., a chimeric virus, mosaic virus, or pseudotyped virus, and/or a virus that contains a foreign protein, synthetic polymer, nanoparticle, or small molecule).

In some embodiments, the viral vector is an AAV, such as an AAV1 (i.e., an AAV containing AAV1 inverted terminal repeats (ITRs) and AAV1 capsid proteins), AAV2 (i.e., an AAV containing AAV2 ITRs and AAV2 capsid proteins), AAV3 (i.e., an AAV containing AAV3 ITRs and AAV3 capsid proteins), AAV4 (i.e., an AAV containing AAV4 ITRs and AAV4 capsid proteins), AAV5 (i.e., an AAV containing AAV5 ITRs and AAV5 capsid proteins), AAV6 (i.e., an AAV containing AAV6 ITRs and AAV6 capsid proteins), AAV7 (i.e., an AAV containing AAV7 ITRs and AAV7 capsid proteins), AAV8 (i.e., an AAV containing AAV8 ITRs and AAV8 capsid proteins), AAV9 (i.e., an AAV containing AAV9 ITRs and AAV9 capsid proteins), AAVrh74 (i.e., an AAV containing AAVrh74 ITRs and AAVrh74 capsid proteins), AAVrh.8 (i.e., an AAV containing AAVrh.8 ITRs and AAVrh.8 capsid proteins), or AAVrh.10 (i.e., an AAV containing AAVrh.10 ITRs and AAVrh.10 capsid proteins).

In some embodiments, the viral vector is a pseudotyped AAV, containing ITRs from one AAV serotype and capsid proteins from a different AAV serotype. In some embodiments, the pseudotyped AAV is AAV2/9 (i.e., an AAV containing AAV2 ITRs and AAV9 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/8 (i.e., an AAV containing AAV2 ITRs and AAV8 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/1 (i.e., an AAV containing AAV2 ITRs and AAV1 capsid proteins).

In some embodiments, the AAV contains a recombinant capsid protein, such as a capsid protein containing a chimera of one or more of capsid proteins from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10.

In certain embodiments, the viral vector is AAV9. For example, the composition may comprise AAV9 comprising a nucleic acid sequence comprising a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG. In certain embodiments, the composition comprises AAV9 comprising a nucleic acid sequence comprising a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1 or MPV17. In a particular embodiment, the composition comprises AAV9 comprising a nucleic acid sequence comprising a transgene encoding TK2.

In certain embodiments, the viral vector is an AAV and the transgene is TK2. For example, the composition may comprise a recombinant AAV (rAAV), such as AAV9, comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein.

In some embodiments of any of the above aspects of the disclosure, the composition is a liposome, vesicle, synthetic vesicle, exosome, synthetic exosome, dendrimer, or nanoparticle.

In some embodiments of any of the above aspects of the disclosure, the transgene is operably linked to a promoter that induces expression of the transgene in a muscle cell. The promoter may be, for example, a chicken beta actin promoter, cytomegalovirus (CMV) promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter, Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial natriuretic factor promoter.

In some embodiments of any of the above aspects of the disclosure, the transgene is operably linked to an enhancer that induces expression of the transgene in a muscle cell. Exemplary enhancers that may be used in conjunction with the compositions and methods of the disclosure are a CMV enhancer, a myocyte enhancer factor 2 (MEF2) enhancer, and a MyoD enhancer.

In some embodiments, the composition is administered to the subject as soon as, or immediately after, the subject is diagnosed as having a deficiency in an endogenous gene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG. In some embodiments, the composition is administered to the subject as soon as, or immediately after,
5 the subject is diagnosed as having a deficiency in an endogenous gene encoding TK2.

In some embodiments, the method further includes administering to the subject a therapeutically effective amount of a second composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG.

In some embodiments, the transgene of the second composition encodes TK2. The
10 TK2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least
15 90% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that differs from
20 SEQ ID NO: 1 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by way of one or more
25 conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding TK2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO:
30 2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene of the second composition encoding TK2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g.,

a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TK2 is at least at least 70% identical to coding sequence of the transgene that encodes TK2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TK2 has the nucleic acid sequence of SEQ ID NO: 17.

In some embodiments, the transgene of the second composition encodes dGK. The dGK may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or

more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding dGK has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene of the second composition encoding dGK has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding dGK is at least at least 70% identical to coding sequence of the transgene that encodes dGK (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding dGK has the nucleic acid sequence of SEQ ID NO: 18.

In some embodiments, the transgene of the second composition encodes TP. The TP may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid

sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding TP has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene of the second composition encoding TP has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TP is at least at least 70% identical to coding sequence of the transgene that encodes TP (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TP has the nucleic acid sequence of SEQ ID NO: 19.

In some embodiments, the transgene of the second composition encodes p53R2. The p53R2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding p53R2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene of the second composition encoding p53R2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of

SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding p53R2 is at least at least 70% identical to coding sequence of the transgene that encodes p53R2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding p53R2 has the nucleic acid sequence of SEQ ID NO: 20.

In some embodiments, the transgene of the second composition encodes SUCLA2. The SUCLA2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding SUCLA2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID

NO: 10). In some embodiments, the transgene of the second composition encoding SUCLA2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding SUCLA2 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLA2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLA2 has the nucleic acid sequence of SEQ ID NO: 21.

In some embodiments, the transgene of the second composition encodes SUCLG1. The SUCLG1 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that differs from SEQ ID NO: 11 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In

some embodiments, the SUCLG1 has an amino acid sequence that differs from SEQ ID NO: 11 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding SUCLG1 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene of the second composition encoding SUCLG1 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding SUCLG1 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLG1 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLG1 has the nucleic acid sequence of SEQ ID NO: 22.

In some embodiments, the transgene of the second composition encodes MPV17. The MPV17 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence

that is at least 90% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments, the transgene of the second composition encoding MPV17 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene of the second composition encoding MPV17 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding MPV17 is at least at least 70% identical to coding sequence of the transgene that encodes MPV17 (e.g., a nucleic acid sequence that is 70%, 71%,

72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding MPV17 has the nucleic acid sequence of SEQ ID NO: 23.

5 In some embodiments, the transgene of the second composition encodes POLG. The POLG may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

25 In some embodiments, the transgene of the second composition encoding POLG has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene of the second composition encoding POLG has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG has a nucleic acid sequence

that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding POLG is at least at least 70% identical to coding sequence of the transgene that encodes POLG (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding POLG has the nucleic acid sequence of SEQ ID NO: 24.

In some embodiments, the second composition is a vector, such as a viral vector. The viral vector may be, for example, an AAV, adenovirus, lentivirus, retrovirus, poxvirus, baculovirus, herpes simplex virus, vaccinia virus, or a synthetic virus (e.g., a chimeric virus, mosaic virus, or pseudotyped virus, and/or a virus that contains a foreign protein, synthetic polymer, nanoparticle, or small molecule).

In some embodiments, the second composition is an AAV, such as an AAV1 (i.e., an AAV containing AAV1 inverted terminal repeats (ITRs) and AAV1 capsid proteins), AAV2 (i.e., an AAV containing AAV2 ITRs and AAV2 capsid proteins), AAV3 (i.e., an AAV containing AAV3 ITRs and AAV3 capsid proteins), AAV4 (i.e., an AAV containing AAV4 ITRs and AAV4 capsid proteins), AAV5 (i.e., an AAV containing AAV5 ITRs and AAV5 capsid proteins), AAV6 (i.e., an AAV containing AAV6 ITRs and AAV6 capsid proteins), AAV7 (i.e., an AAV containing AAV7 ITRs and AAV7 capsid proteins), AAV8 (i.e., an AAV containing AAV8 ITRs and AAV8 capsid proteins), AAV9 (i.e., an AAV containing AAV9 ITRs and AAV9 capsid proteins), AAVrh74 (i.e., an AAV containing AAVrh74 ITRs and AAVrh74 capsid proteins), AAVrh.8 (i.e., an AAV containing AAVrh.8 ITRs and AAVrh.8 capsid proteins), or AAVrh.10 (i.e., an AAV containing AAVrh.10 ITRs and AAVrh.10 capsid proteins).

In some embodiments, the second composition is a pseudotyped AAV, containing ITRs from one AAV serotype and capsid proteins from a different AAV serotype. In some embodiments, the pseudotyped AAV is AAV2/9 (i.e., an AAV containing AAV2 ITRs and AAV9 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/8 (i.e., an

AAV containing AAV2 ITRs and AAV8 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/1 (i.e., an AAV containing AAV2 ITRs and AAV1 capsid proteins).

In some embodiments, the second composition is an AAV that contains a recombinant capsid protein, such as a capsid protein containing a chimera of one or more of capsid proteins from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10.

In some embodiments, the second composition is a liposome, vesicle, synthetic vesicle, exosome, synthetic exosome, dendrimer, or nanoparticle.

In some embodiments, the transgene of the second composition is operably linked to a promoter that induces expression of the transgene in a muscle cell. The promoter may be, for example, a chicken beta actin promoter, cytomegalovirus (CMV) promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter, Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial natriuretic factor promoter.

In some embodiments, the transgene of the second composition is operably linked to an enhancer that induces expression of the transgene in a muscle cell. Exemplary enhancers that may be used in conjunction with the compositions and methods of the disclosure are a CMV enhancer, a myocyte enhancer factor 2 (MEF2) enhancer, and a MyoD enhancer.

In some embodiments, the second composition is administered to the subject after administration of the first composition to the subject. The second composition may be administered to the subject, for example, within one or more days or weeks of administration of the first composition to the subject. In some embodiments, the second composition is administered to the subject at least one month after administration of the first composition to the subject. In some embodiments, administration of the first composition continues while the second composition is administered to the subject.

In some embodiments, the method further includes administering to the subject a therapeutically effective amount of a third composition containing a pharmacological agent. The pharmacological agent may be, for example, deoxycytidine (dC), deoxythymidine (dT), deoxyadenosine (dA), deoxyguanosine (dG), deoxycytidine monophosphate (dCMP), deoxythymidine monophosphate (TMP), deoxyadenosine monophosphate (dAMP), deoxyguanosine monophosphate (dGMP), or a mixture thereof. In some embodiments, the pharmacological agent is dC, dT, or a mixture thereof.

In some embodiments, the third composition is administered to the subject after administration of the first composition to the subject. The third composition may be administered to the subject, for example, within one or more days or weeks of administration of the first composition to the subject. In some embodiments, the third composition is administered to the subject at least one month after administration of the first composition to the subject.

In some embodiments, the third composition is administered to the subject after administration of the second composition to the subject. The third composition may be administered to the subject, for example, within one or more days or weeks of administration of the second composition to the subject. In some embodiments, the third composition is administered to the subject at least one month after administration of the second composition to the subject.

In some embodiments, administration of the first composition and administration of the second composition continue while the third composition is administered to the subject.

In some embodiments, the third composition is administered to the subject in an amount of from about 100 mg/kg/day to about 1,000 mg/kg/day (e.g., about 100 mg/kg/day, 105 mg/kg/day, 110 mg/kg/day, 115 mg/kg/day, 120 mg/kg/day, 125 mg/kg/day, 130 mg/kg/day, 135 mg/kg/day, 140 mg/kg/day, 145 mg/kg/day, 150 mg/kg/day, 155 mg/kg/day, 160 mg/kg/day, 165 mg/kg/day, 170 mg/kg/day, 175 mg/kg/day, 180 mg/kg/day, 185 mg/kg/day, 190 mg/kg/day, 195 mg/kg/day, 200 mg/kg/day, 205 mg/kg/day, 210 mg/kg/day, 215 mg/kg/day, 220 mg/kg/day, 225 mg/kg/day, 230 mg/kg/day, 235 mg/kg/day, 240 mg/kg/day, 245 mg/kg/day, 250 mg/kg/day, 255 mg/kg/day, 260 mg/kg/day, 265 mg/kg/day, 270 mg/kg/day, 275 mg/kg/day, 280 mg/kg/day, 285 mg/kg/day, 290 mg/kg/day, 295 mg/kg/day, 300 mg/kg/day, 305 mg/kg/day, 310 mg/kg/day, 315 mg/kg/day, 320 mg/kg/day, 325 mg/kg/day, 330 mg/kg/day, 335 mg/kg/day, 340 mg/kg/day, 345 mg/kg/day, 350 mg/kg/day, 355 mg/kg/day, 360 mg/kg/day, 365 mg/kg/day, 370 mg/kg/day, 375 mg/kg/day, 380 mg/kg/day, 385 mg/kg/day, 390 mg/kg/day, 395 mg/kg/day, 400 mg/kg/day, 405 mg/kg/day, 410 mg/kg/day, 415 mg/kg/day, 420 mg/kg/day, 425 mg/kg/day, 430 mg/kg/day, 435 mg/kg/day, 440 mg/kg/day, 445 mg/kg/day, 450 mg/kg/day, 455 mg/kg/day, 460 mg/kg/day, 465 mg/kg/day, 470 mg/kg/day, 475 mg/kg/day, 480 mg/kg/day, 485 mg/kg/day, 490 mg/kg/day, 495 mg/kg/day, 500 mg/kg/day, 505 mg/kg/day, 510 mg/kg/day, 515 mg/kg/day, 520 mg/kg/day, 525 mg/kg/day, 530 mg/kg/day, 535 mg/kg/day, 540 mg/kg/day, 545 mg/kg/day, 550 mg/kg/day, 555 mg/kg/day, 560 mg/kg/day, 565 mg/kg/day, 570 mg/kg/day, 575 mg/kg/day, 580 mg/kg/day, 585 mg/kg/day, 590 mg/kg/day, 595 mg/kg/day, 600

mg/kg/day, 605 mg/kg/day, 610 mg/kg/day, 615 mg/kg/day, 620 mg/kg/day, 625 mg/kg/day, 630 mg/kg/day, 635 mg/kg/day, 640 mg/kg/day, 645 mg/kg/day, 650 mg/kg/day, 655 mg/kg/day, 660 mg/kg/day, 665 mg/kg/day, 670 mg/kg/day, 675 mg/kg/day, 680 mg/kg/day, 685 mg/kg/day, 690 mg/kg/day, 695 mg/kg/day, 700 mg/kg/day, 705 mg/kg/day, 710 mg/kg/day, 715 mg/kg/day, 720 mg/kg/day, 725 mg/kg/day, 730 mg/kg/day, 735 mg/kg/day, 740 mg/kg/day, 745 mg/kg/day, 750 mg/kg/day, 755 mg/kg/day, 760 mg/kg/day, 765 mg/kg/day, 770 mg/kg/day, 775 mg/kg/day, 780 mg/kg/day, 785 mg/kg/day, 790 mg/kg/day, 795 mg/kg/day, 800 mg/kg/day, 805 mg/kg/day, 810 mg/kg/day, 815 mg/kg/day, 820 mg/kg/day, 825 mg/kg/day, 830 mg/kg/day, 835 mg/kg/day, 840 mg/kg/day, 845 mg/kg/day, 850 mg/kg/day, 855 mg/kg/day, 860 mg/kg/day, 865 mg/kg/day, 870 mg/kg/day, 875 mg/kg/day, 880 mg/kg/day, 885 mg/kg/day, 890 mg/kg/day, 895 mg/kg/day, 900 mg/kg/day, 905 mg/kg/day, 910 mg/kg/day, 915 mg/kg/day, 920 mg/kg/day, 925 mg/kg/day, 930 mg/kg/day, 935 mg/kg/day, 940 mg/kg/day, 945 mg/kg/day, 950 mg/kg/day, 955 mg/kg/day, 960 mg/kg/day, 965 mg/kg/day, 970 mg/kg/day, 975 mg/kg/day, 980 mg/kg/day, 985 mg/kg/day, 990 mg/kg/day, 995 mg/kg/day, or 1,000 mg/kg/day).

In some embodiments, the third composition is administered to the subject in an amount of from about 200 mg/kg/day to about 800 mg/kg/day (e.g., about 200 mg/kg/day, 205 mg/kg/day, 210 mg/kg/day, 215 mg/kg/day, 220 mg/kg/day, 225 mg/kg/day, 230 mg/kg/day, 235 mg/kg/day, 240 mg/kg/day, 245 mg/kg/day, 250 mg/kg/day, 255 mg/kg/day, 260 mg/kg/day, 265 mg/kg/day, 270 mg/kg/day, 275 mg/kg/day, 280 mg/kg/day, 285 mg/kg/day, 290 mg/kg/day, 295 mg/kg/day, 300 mg/kg/day, 305 mg/kg/day, 310 mg/kg/day, 315 mg/kg/day, 320 mg/kg/day, 325 mg/kg/day, 330 mg/kg/day, 335 mg/kg/day, 340 mg/kg/day, 345 mg/kg/day, 350 mg/kg/day, 355 mg/kg/day, 360 mg/kg/day, 365 mg/kg/day, 370 mg/kg/day, 375 mg/kg/day, 380 mg/kg/day, 385 mg/kg/day, 390 mg/kg/day, 395 mg/kg/day, 400 mg/kg/day, 405 mg/kg/day, 410 mg/kg/day, 415 mg/kg/day, 420 mg/kg/day, 425 mg/kg/day, 430 mg/kg/day, 435 mg/kg/day, 440 mg/kg/day, 445 mg/kg/day, 450 mg/kg/day, 455 mg/kg/day, 460 mg/kg/day, 465 mg/kg/day, 470 mg/kg/day, 475 mg/kg/day, 480 mg/kg/day, 485 mg/kg/day, 490 mg/kg/day, 495 mg/kg/day, 500 mg/kg/day, 505 mg/kg/day, 510 mg/kg/day, 515 mg/kg/day, 520 mg/kg/day, 525 mg/kg/day, 530 mg/kg/day, 535 mg/kg/day, 540 mg/kg/day, 545 mg/kg/day, 550 mg/kg/day, 555 mg/kg/day, 560 mg/kg/day, 565 mg/kg/day, 570 mg/kg/day, 575 mg/kg/day, 580 mg/kg/day, 585 mg/kg/day, 590 mg/kg/day, 595 mg/kg/day, 600 mg/kg/day, 605 mg/kg/day, 610 mg/kg/day, 615 mg/kg/day, 620 mg/kg/day, 625 mg/kg/day, 630 mg/kg/day, 635 mg/kg/day, 640 mg/kg/day, 645 mg/kg/day, 650 mg/kg/day, 655 mg/kg/day, 660 mg/kg/day, 665 mg/kg/day, 670

mg/kg/day, 675 mg/kg/day, 680 mg/kg/day, 685 mg/kg/day, 690 mg/kg/day, 695 mg/kg/day, 700 mg/kg/day, 705 mg/kg/day, 710 mg/kg/day, 715 mg/kg/day, 720 mg/kg/day, 725 mg/kg/day, 730 mg/kg/day, 735 mg/kg/day, 740 mg/kg/day, 745 mg/kg/day, 750 mg/kg/day, 755 mg/kg/day, 760 mg/kg/day, 765 mg/kg/day, 770 mg/kg/day, 775 mg/kg/day, 780
5 mg/kg/day, 785 mg/kg/day, 790 mg/kg/day, 795 mg/kg/day, or 800 mg/kg/day).

In some embodiments, the third composition is administered to the subject in an amount of from about 250 mg/kg/day to about 400 mg/kg/day (e.g., about 250 mg/kg/day, 255 mg/kg/day, 260 mg/kg/day, 265 mg/kg/day, 270 mg/kg/day, 275 mg/kg/day, 280 mg/kg/day, 285 mg/kg/day, 290 mg/kg/day, 295 mg/kg/day, 300 mg/kg/day, 305 mg/kg/day, 310
10 mg/kg/day, 315 mg/kg/day, 320 mg/kg/day, 325 mg/kg/day, 330 mg/kg/day, 335 mg/kg/day, 340 mg/kg/day, 345 mg/kg/day, 350 mg/kg/day, 355 mg/kg/day, 360 mg/kg/day, 365 mg/kg/day, 370 mg/kg/day, 375 mg/kg/day, 380 mg/kg/day, 385 mg/kg/day, 390 mg/kg/day, 395 mg/kg/day, or 400 mg/kg/day).

In some embodiments, the third composition is administered to the subject once daily,
15 twice daily, three times daily, four times daily, five times daily, or six times daily. The third composition may be administered to the subject orally in admixture with cow's milk, human breast milk, infant formula, and/or water.

In some embodiments, the first composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal,
20 subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular, inhalation, perfusion, lavage, and/or oral administration.

In some embodiments, the second composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular,
25 inhalation, perfusion, lavage, and/or oral administration.

In some embodiments, the third composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular, inhalation, perfusion, lavage, and/or oral administration.

30 In some embodiments, the subject is a mammal, such as a human. In some embodiments, the subject is a pediatric human subject, such as a human subject of from about 1 month to about 12 years of age (e.g., a human subject of from about 1 month to about 1 year, 2 years, 3 years, 4 years, 5 years, 6 years, 7 years, 8 years, 9 years, 10 years, 11 years, or 12 years of age).

In a further aspect, the disclosure features a composition containing a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG.

In some embodiments of the preceding aspect, the transgene encodes TK2. The TK2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 1). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TK2 has an amino acid sequence that differs from SEQ ID NO: 1 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding TK2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments of the preceding aspect, the transgene encoding TK2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 has a

nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 2). In some embodiments, the transgene encoding TK2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TK2 is at least at least 70% identical to coding sequence of the transgene that encodes TK2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TK2 has the nucleic acid sequence of SEQ ID NO: 17.

In some embodiments of the preceding aspect, the transgene encodes dGK. The dGK may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 3 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 3). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the dGK has an amino acid sequence that differs from SEQ ID NO: 3 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding dGK has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO:

4). In some embodiments of the preceding aspect, the transgene encoding dGK has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 4 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 4). In some embodiments, the transgene encoding dGK is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding dGK is at least at least 70% identical to coding sequence of the transgene that encodes dGK (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding dGK has the nucleic acid sequence of SEQ ID NO: 18.

In some embodiments of the preceding aspect, the transgene encodes TP. The TP may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 5 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 5). In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the TP has an amino acid sequence that differs from SEQ ID NO: 5 by way of one or more conservative

amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding TP has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g.,
 5 a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments of the preceding aspect, the transgene encoding TP has a nucleic acid sequence
 10 that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is
 15 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 6 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 6). In some embodiments, the transgene encoding TP is codon
 20 optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding TP is at least at least 70% identical to coding sequence of the transgene that encodes TP (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding
 25 sequence). In some embodiments, the nucleic acid sequence of the transgene encoding TP has the nucleic acid sequence of SEQ ID NO: 19.

In some embodiments of the preceding aspect, the transgene encodes p53R2. The p53R2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%,
 30 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino

acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 7 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 7). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the p53R2 has an amino acid sequence that differs from SEQ ID NO: 7 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding p53R2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments of the preceding aspect, the transgene encoding p53R2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 8 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 8). In some embodiments, the transgene encoding p53R2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding p53R2 is at least at least 70% identical to coding sequence of the transgene that encodes p53R2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence

of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding p53R2 has the nucleic acid sequence of SEQ ID NO: 20.

In some embodiments of the preceding aspect, the transgene encodes SUCLA2. The SUCLA2 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 9 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 9). In some embodiments, the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the SUCLA2 has an amino acid sequence that differs from SEQ ID NO: 9 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments of the preceding aspect, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100%

identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 10 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 10). In some embodiments, the transgene encoding SUCLA2 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding SUCLA2 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLA2 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLA2 has the nucleic acid sequence of SEQ ID NO: 21.

In some embodiments of the preceding aspect, the transgene encodes SUCLG1. The SUCLG1 may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 11 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 11). In some embodiments, the SUCLG1 has an amino acid sequence that differs from SEQ ID NO: 11 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the SUCLG1 has an amino acid sequence that differs from SEQ ID NO: 11 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO:

12 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments of the preceding aspect, the transgene encoding SUCLG1 has a

5 nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic

10 acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the transgene encoding SUCLG1 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 12 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 12). In some embodiments, the

15 transgene encoding SUCLG1 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding SUCLG1 is at least at least 70% identical to coding sequence of the transgene that encodes SUCLG1 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical

20 to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding SUCLG1 has the nucleic acid sequence of SEQ ID NO: 22.

In some embodiments of the preceding aspect, the transgene encodes MPV17. The MPV17 may have an amino acid sequence that is at least 85% identical to the amino acid

25 sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical

30 to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 13 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 13). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more amino acid substitutions,

insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the MPV17 has an amino acid sequence that differs from SEQ ID NO: 13 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding MPV17 has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments of the preceding aspect, the transgene encoding MPV17 has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 14 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 14). In some embodiments, the transgene encoding MPV17 is codon optimized to increase efficiency. In some embodiments, the nucleic acid of the transgene encoding MPV17 is at least at least 70% identical to coding sequence of the transgene that encodes MPV17 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the transgene encoding MPV17 has the nucleic acid sequence of SEQ ID NO: 23.

In some embodiments of the preceding aspect, the transgene encodes POLG. The POLG may have an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 85%, 86%, 87%, 88%, 89%,

90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 15 (e.g., an amino acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the amino acid sequence of SEQ ID NO: 15). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more amino acid substitutions, insertions, and/or deletions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, amino acid substitutions, insertions, and/or deletions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions). In some embodiments, the POLG has an amino acid sequence that differs from SEQ ID NO: 15 by way of one or more conservative amino acid substitutions, such as by from 1 to 10, 1 to 15, 1 to 20, 1 to 25, or more, conservative amino acid substitutions (e.g., by 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, or more, conservative amino acid substitutions).

In some embodiments of the preceding aspect, the transgene encoding POLG has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments of the preceding aspect, the transgene encoding POLG has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 90% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 16 (e.g., a nucleic acid sequence that is 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of SEQ ID NO: 16). In some embodiments, the transgene encoding POLG is codon optimized to increase efficiency. In some embodiments, the nucleic

acid of the transgene encoding POLG is at least at least 70% identical to coding sequence of the transgene that encodes POLG (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence of the coding sequence). In some embodiments, the nucleic acid sequence of the

In some embodiments of the preceding aspect, the composition is a vector, such as a viral vector. The viral vector may be, for example, an AAV, adenovirus, lentivirus, retrovirus, poxvirus, baculovirus, herpes simplex virus, vaccinia virus, or a synthetic virus (e.g., a chimeric virus, mosaic virus, or pseudotyped virus, and/or a virus that contains a foreign protein, synthetic polymer, nanoparticle, or small molecule).

In some embodiments of the preceding aspect, the viral vector is an AAV, such as an AAV1 (i.e., an AAV containing AAV1 inverted terminal repeats (ITRs) and AAV1 capsid proteins), AAV2 (i.e., an AAV containing AAV2 ITRs and AAV2 capsid proteins), AAV3 (i.e., an AAV containing AAV3 ITRs and AAV3 capsid proteins), AAV4 (i.e., an AAV containing AAV4 ITRs and AAV4 capsid proteins), AAV5 (i.e., an AAV containing AAV5 ITRs and AAV5 capsid proteins), AAV6 (i.e., an AAV containing AAV6 ITRs and AAV6 capsid proteins), AAV7 (i.e., an AAV containing AAV7 ITRs and AAV7 capsid proteins), AAV8 (i.e., an AAV containing AAV8 ITRs and AAV8 capsid proteins), AAV9 (i.e., an AAV containing AAV9 ITRs and AAV9 capsid proteins), AAVrh74 (i.e., an AAV containing AAVrh74 ITRs and AAVrh74 capsid proteins), AAVrh.8 (i.e., an AAV containing AAVrh.8 ITRs and AAVrh.8 capsid proteins), or AAVrh.10 (i.e., an AAV containing AAVrh.10 ITRs and AAVrh.10 capsid proteins).

In some embodiments of the preceding aspect, the viral vector is a pseudotyped AAV, containing ITRs from one AAV serotype and capsid proteins from a different AAV serotype. In some embodiments, the pseudotyped AAV is AAV2/9 (i.e., an AAV containing AAV2 ITRs and AAV9 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/8 (i.e., an AAV containing AAV2 ITRs and AAV8 capsid proteins). In some embodiments, the pseudotyped AAV is AAV2/1 (i.e., an AAV containing AAV2 ITRs and AAV1 capsid proteins).

In some embodiments of the preceding aspect, the AAV contains a recombinant capsid protein, such as a capsid protein containing a chimera of one or more of capsid proteins from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10.

In some embodiments of the preceding aspect, the composition is a liposome, vesicle, synthetic vesicle, exosome, synthetic exosome, dendrimer, or nanoparticle.

In some embodiments of the preceding aspect, the transgene is operably linked to a promoter that induces expression of the transgene in a muscle cell. The promoter may be, for example, a chicken beta actin promoter, cytomegalovirus (CMV) promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter, Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial natriuretic factor promoter.

In some embodiments of the preceding aspect, the transgene is operably linked to an enhancer that induces expression of the transgene in a muscle cell. Exemplary enhancers that may be used in conjunction with the compositions and methods of the disclosure are a CMV enhancer, a myocyte enhancer factor 2 (MEF2) enhancer, and a MyoD enhancer.

In another aspect, the disclosure features a kit containing the composition of the preceding aspect. The kit may further contain a package insert, such as a package insert instructing a user of the kit to administer the composition to a subject in accordance with the method of any of the above aspects or embodiments of the disclosure. The kit may further contain a pharmacological agent selected from the group consisting of dC, dT, dA, dG, dCMP, TMP, dAMP, dGMP, and mixtures thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

For the purpose of illustrating the invention, there are depicted in drawings certain embodiments of the invention. However, the invention is not limited to the precise arrangements and instrumentalities of the embodiments depicted in the drawings.

In the drawings, certain abbreviations are used. “WT” means wild-type mice which are Tk2⁺. “Het” means mice heterozygous for the TK2 gene. “Mut” means Tk2^{-/-} mice. “P#” means post-natal day #. “AAV2-hTK2” refers to an AAV2 vector containing a human TK2 transgene. “AAV9-hTK2” refers to an AAV9 vector containing a human TK2 transgene. “dC” refers to deoxycytidine. “dT” refers to deoxythymidine. “IV” refers to intravenous administration. “vc” refers to a quantity of vector genome copies.

FIG. 1 is a schematic of a recombinant adeno-associated virus (AAV) vector comprising human TK2 cDNA.

FIG. 2 shows the survival curves of the following mice (curves from left to right): Tk2^{-/-} mice untreated (n=13); Tk2^{-/-} mice administered milk only (n=6); Tk2^{-/-} mice treated with

AAV9-hTK2 at 4.2×10^{10} vc by IV at postnatal day 1 (n=5); *Tk2*^{-/-} mice treated with 520 mg of dC+dT (n=11); and *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1 (n=10). Survival of untreated *Tk2*⁺ (wild-type) is shown by the top horizontal line.

FIG. 3 shows graphs of weight versus time in the following male and female mice:
 5 *Tk2*⁺ treated with 520 mg of dC+dT; *Tk2*^{-/-} mice treated with 520 mg of dC+dT; *Tk2*⁺ treated AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1; and *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{10} vc by IV at postnatal day 1.

FIG. 4 shows graphs of the result of grip strength test of the fore limbs (bar test) and both fore and hind limbs (grid test) normalized by mouse weight at postnatal day 29 and
 10 postnatal day 60 of male and female mice including untreated *Tk2*⁺, *Tk2*⁺ treated with AAV9-hTK2 and *Tk2*^{-/-} mice treated with AAV9-hTK2.

FIG. 5 shows graphs of the results of mice motor function as measured by accelerating rotarod performance test at postnatal day 29 and postnatal day 60 of male and female mice including *Tk2*⁺ treated with AAV9-hTK2 and *Tk2*^{-/-} mice treated with AAV9-hTK2.

FIG. 6 shows graphs of the activity of TK2 enzyme measured as pmol/min/mg of protein at postnatal day 29 in brain, liver, muscle and kidney tissue of the following mice:
 15 untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2 at 4.2×10^{11} vc; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc; *Tk2*⁺ treated with 520 mg of dC+dT; and *Tk2*^{-/-} mice treated with 520 mg of dC+dT.

FIG. 7 shows graphs of mRNA expression of human *TK2* measure as a percent of mouse *TK2* mRNA in brain, liver, muscle and kidney tissue of *Tk2*⁺ treated with AAV9-hTK2
 20 at one month, two months and six months.

FIG. 8 shows graphs of showing mtDNA copy number (mtDNA /nDNA as a percent of untreated *Tk2*⁺) at postnatal day 29 in brain, liver, kidney, heart, muscle, and intestine tissue
 25 of the following mice: untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2; mice which are heterozygous for the *Tk2* gene treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2; and *Tk2*^{-/-} mice treated with 520 mg of dC+dT.

FIG. 9 shows graphs of showing mtDNA copy number (mtDNA /nDNA as a percent of untreated *Tk2*⁺) at postnatal day 60 in brain, liver, kidney, heart, muscle, and intestine tissue
 30 of the following mice: untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2; mice which are heterozygous for the *Tk2* gene treated with AAV9-hTK2; and *Tk2*^{-/-} mice treated with AAV9-hTK2.

FIG. 10 shows graphs of respiratory chain enzyme activity at postnatal day 29 in brains of the following mice: untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2; mice which are

heterozygous for the *Tk2* gene treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2; and *Tk2*^{-/-} mice treated with 520 mg of dC+dT.

FIG. 11 shows graphs of respiratory chain enzyme levels (oxphos steady-state levels normalized to vinculin) at postnatal day 29 in brains of the following mice: untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2; and *Tk2*^{-/-} mice treated with 520 mg of dC+dT.

FIG. 12 shows images of stained (H & E) kidneys of AAV9-hTK2 treated mutant and wild-type mice at postnatal day 96.

FIG. 13 shows graphs of creatinine and BUN index in AAV9-hTK2 treated mutant and wild-type mice in a survival cohort and a postnatal day 60 cohort.

FIG. 14 shows the survival curve of the following mice (curves from left to right): *Tk2*^{-/-} mice untreated (n=13); *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{10} vc by IV at postnatal day 1 (n=5); *Tk2*^{-/-} mice treated with 520 mg of dC+dT (n=11); *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1 (n=10); mice co-treated with AAV9-hTK2 at 2.1×10^{11} at day 1 and AAV2-hTK2 at 1.05×10^{11} at day 29 (n=7); and mice under co-treatment with AAV9-hTK2 at day 1, AAV2-hTK2 at day 29 and supplemented with 520 mg/kg/day of oral dC+dT from day 21 (n=7). Survival of untreated *Tk2*⁺ (wild-type) is shown by the top horizontal line.

FIG. 15 shows weight versus time in the following male mice: *Tk2*⁺ untreated; *Tk2*⁺ treated with 520 mg of dC+dT; *Tk2*^{-/-} mice treated with 520 mg of dC+dT; *Tk2*⁺ treated AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1 (“AAV9”); *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{10} vc by IV at postnatal day 1; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 2.1×10^{11} vc by IV at postnatal day 1 and AAV2-hTK2 at 1.05×10^{11} vc by IV at postnatal day 29 (“AAV9 + AAV2”); and *Tk2*^{-/-} mice treated with AAV9-hTK2 at 2.1×10^{11} vc by IV at postnatal day 1, AAV2-hTK2 at 1.05×10^{11} vc by IV at postnatal day 29 and 520 mg/kg/day of oral dC+dT from day 21 (“AAV9 + AAV2 + dCdT”).

FIG. 16 shows weight versus time in the following female mice: *Tk2*⁺ untreated; *Tk2*⁺ treated with 520 mg of dC+dT; *Tk2*^{-/-} mice treated with 520 mg of dC+dT; *Tk2*⁺ treated AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{11} vc by IV at postnatal day 1 (“AAV9”); *Tk2*^{-/-} mice treated with AAV9-hTK2 at 4.2×10^{10} vc by IV at postnatal day 1; *Tk2*^{-/-} mice treated with AAV9-hTK2 at 2.1×10^{11} vc by IV at postnatal day 1 and AAV2-hTK2 at 1.05×10^{11} vc by IV at postnatal day 29 (“AAV9 + AAV2”); and *Tk2*^{-/-} mice treated with AAV9-hTK2 at 2.1×10^{11} vc by IV at postnatal day 1,

AAV2-hTK2 at 1.05×10^{11} vc by IV at postnatal day 29 and 520 mg/kg/day of oral dC+dT from day 21 (AAV9 + AAV2+ dCdT”).

FIG. 17 shows graphs of the result of grip strength test of the fore limbs (bar test) and both fore and hind limbs (grid test) normalized by mouse weight at postnatal day 60 of male and female mice including untreated *Tk2*⁺, *Tk2*⁺ treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2, and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

FIG. 18 shows graphs of the result of grip strength test of the fore limbs (bar test) and both fore and hind limbs (grid test) normalized by mouse weight at postnatal day 90 of male and female mice including untreated *Tk2*⁺, *Tk2*⁺ treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2, and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

FIG. 19 shows graphs of the results of mice motor function as measured by accelerating rotarod performance test at postnatal day 60 and postnatal day 90 of male and female mice including untreated *Tk2*⁺, *Tk2*⁺ treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2, and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

FIG. 20 shows graphs of showing mtDNA copy number (mtDNA /nDNA as a percent of untreated *Tk2*⁺) at postnatal day 60 in brain, liver, muscle, heart, kidney, and intestine tissue of the following mice: untreated *Tk2*⁺; *Tk2*^{-/-} mice treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2; and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

FIG. 21 depicts the results from the protein test in a urine dip stick using urine from mice at age 21-29 days and age 60 days. All *Tk2*^{-/-} mice were treated with AAV9-hTK2 at day 1, while 3 out of the four *Tk2*^{-/-} mice were treated with AAV2-hTK2 at day 29.

FIG. 22 shows graphs of BUN index in mice at day 60 and mice in an endpoint cohort including untreated *Tk2*⁺, *Tk2*⁺ treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2, *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2, and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

FIG. 23 shows graphs of the activity of TK2 enzyme measured as pmol/min/mg of protein at postnatal day 60 in brain, liver, muscle and kidney tissue of the following mice: untreated *Tk2*⁺; *Tk2*⁺ treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2; *Tk2*^{-/-} mice treated with AAV9-hTK2 and AAV2-hTK2; and *Tk2*^{-/-} mice treated with AAV9-hTK2, AAV2-hTK2 and dC+dT.

DETAILED DESCRIPTION OF THE INVENTION

The current invention is based, at least in part, upon the surprising discovery that mitochondrial DNA depletion syndromes, including TK2 deficiency, can be treated, prevented and/or cured via gene therapy by delivery of nucleic acids encoding for a functional protein, such as a functional protein encoded by a nuclear gene. Exemplary nucleic acids that can be used in conjunction with the compositions and methods of the disclosure encode thymidine kinase 2 (TK2), deoxyguanosine kinase (dGK), thymidine phosphorylase (TP), p53 inducible small subunit of ribonucleotide reductase (p53R2), succinyl-CoA ligase ADP-forming subunit beta (SUCLA2), succinyl-CoA ligase GDP-forming subunit alpha (SUCLG1), mitochondrial inner membrane protein MPV17 (MPV17), and/or DNA polymerase subunit gamma (POLG). Such nucleic acids can be delivered by way of one or more viral vectors, such as by way of an adeno-associated virus (AAV), among others described herein. The gene therapy can be administered alone or in combination with pharmacological therapy, such as deoxynucleoside or deoxyribonucleoside monophosphate therapy. The sections that follow describe the compositions and methods of the disclosure in further detail.

Definitions

The terms used in this specification generally have their ordinary meanings in the art, within the context of this invention and the specific context where each term is used. Certain terms are discussed below, or elsewhere in the specification, to provide additional guidance to the practitioner in describing the methods of the invention and how to use them. Moreover, it will be appreciated that the same thing can be said in more than one way. Consequently, alternative language and synonyms may be used for any one or more of the terms discussed herein, nor is any special significance to be placed upon whether or not a term is elaborated or discussed herein. Synonyms for certain terms are provided. A recital of one or more synonyms does not exclude the use of the other synonyms. The use of examples anywhere in the specification, including examples of any terms discussed herein, is illustrative only, and in no way limits the scope and meaning of the invention or any exemplified term. Likewise, the invention is not limited to its preferred embodiments.

The term “about” or “approximately” means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, *i.e.*, the limitations of the measurement system, *i.e.*, the degree of precision required for a particular purpose, such as a pharmaceutical formulation. For example, “about” can mean within 1 or more than 1 standard deviations, per the practice in the art. Alternatively, “about” can mean a range of up to 20%, preferably up to 10%, more

preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated, the term
5 “about” meaning within an acceptable error range for the particular value should be assumed.

The terms “mutant mouse”, “TK2 knockin mouse” and “*Tk2*^{-/-}” refer to the homozygous Tk2 H126N knock-in mutant (*Tk2*^{-/-}) mouse model, described in Akman, *et al.* 2008. The terms “wild-type mouse”, “WT”, and “*Tk2*⁺” will be used interchangeably as well. In some cases, mice which are heterozygous for the *Tk2* gene (“*Tk2*^{+/-}”) are used for comparison and would
10 be expected to behave the same as the wild-type mice. These mice are designated “Het.”

The term “subject” as used in this application refers to animals in need of therapeutic or prophylactic treatment. Subjects include mammals, such as canines, felines, rodents, bovine, equines, porcines, ovines, and primates. Thus, the invention can be used in veterinary medicine, *e.g.*, to treat companion animals, farm animals, laboratory animals in zoological
15 parks, and animals in the wild. The invention is particularly desirable for human medical applications.

The term “patient” as used in this application means a human subject. In some embodiments of the present invention, the “patient” is known or suspected of having a disease or disorder characterized by unbalanced nucleotide pools, mitochondrial disease,
20 mitochondrial DNA depletion syndrome, or TK2 deficiency.

The phrase “therapeutically effective amount” is used herein to mean an amount sufficient to cause an improvement in a clinically significant condition in the subject, or delays or minimizes or mitigates one or more symptoms associated with the disease or disorder, or results in a desired beneficial change of physiology in the subject.

25 The terms “treat”, “treatment”, and the like refer to a means to slow down, relieve, ameliorate or alleviate at least one of the symptoms of the disease or disorder, or reverse the disease or disorder after its onset.

The terms “prevent”, “prevention”, and the like refer to acting prior to overt disease or disorder onset, to prevent the disease or disorder from developing or minimize the extent of the
30 disease or disorder, or slow its course of development.

The term “cure” and the like means to heal, to make well, or to restore to good health or to allow a time without recurrence of disease so that the risk of recurrence is small.

The term “in need thereof” would be a subject known or suspected of having or being at risk of having a disease or disorder characterized by unbalanced nucleotide pools, mitochondrial disease, mitochondrial DNA depletion syndrome, or TK2 deficiency.

5 The term “agent” as used herein means a substance that produces or is capable of producing an effect and would include, but is not limited to, chemicals, pharmaceuticals, biologics, small organic molecules, antibodies, nucleic acids, peptides, and proteins.

The terms “deoxynucleoside” or “dN” as used herein means deoxythymidine or dT, deoxycytidine or dC, deoxyadenosine or dA, and deoxyguanosine or dG. The full length name and common abbreviation for each will be used interchangeably. Such deoxynucleosides also
10 include physiologically functional derivatives of the deoxynucleosides.

The term “deoxyribonucleoside monophosphate” as used herein means thymidine-5'-monophosphate or (TMP) or 2'-deoxycytidine-5'-monophosphate (dCMP), deoxyadenosine monophosphate or dAMP, and deoxyguanosine monophosphate or dGMP. The full length name and common abbreviation for each will be used interchangeably.

15 The term “pharmacological agent” as used herein means deoxynucleosides or dNs and/or deoxyribonucleoside monophosphate or dNMPs, alone or in a composition or a pharmaceutical composition.

The terms “pharmacological therapy” or “pharmacological treatment” as used herein means the administration of dNs and/or dNMPs as disclosed herein and as fully described in
20 co-owned U.S. Application Serial Nos.15/082,207 (hereinafter “the ‘207 application”) and 15/736,092 (hereinafter the ‘092 application”).

As used herein, the term “physiologically functional derivative” refers to a compound (*e.g.*, a drug precursor) that is transformed *in vivo* to yield a deoxynucleoside. The transformation may occur by various mechanisms (*e.g.*, by metabolic or chemical processes),
25 such as, for example, through hydrolysis in blood. Prodrugs are such derivatives, and a discussion of the use of prodrugs is provided by T. Higuchi and W. Stella, “Pro-drugs as Novel Delivery Systems,” Vol. 14 of the A.C.S. Symposium Series, and in *Bioreversible Carriers in Drug Design*, ed. Edward B. Roche, American Pharmaceutical Association and Pergamon Press, 1987.

30 As used herein “an adverse effect” is an unwanted reaction caused by the administration of a drug. In most cases, the administration of the gene and pharmacological therapy caused no adverse effects.

As used herein, the term “carrier” refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered, and includes any and all solvents, dispersion media,

vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art.

The term "pharmaceutically-acceptable" refers to molecular entities and compositions
5 that do not produce an allergic or similar untoward reaction when administered to a host, such as gastric upset, dizziness and the like, when administered to a human, and approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans.

"Isolated nucleic acid molecule" means a DNA or RNA of genomic, mRNA, cDNA, or
10 synthetic origin or some combination thereof which is not associated with all or a portion of a polynucleotide in which the isolated polynucleotide is found in nature or is linked to a polynucleotide to which it is not linked in nature. For purposes of this disclosure, it should be understood that "a nucleic acid molecule comprising" a particular nucleotide sequence does not encompass intact chromosomes. Isolated nucleic acid molecules "comprising" specified
15 nucleic acid sequences may include, in addition to the specified sequences, coding sequences for up to ten or even up to twenty or more other proteins or portions or fragments thereof, or may include operably linked regulatory sequences that control expression of the coding region of the recited nucleic acid sequences, and/or may include vector sequences.

The phrase "control sequences" refers to DNA sequences necessary for the expression
20 of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to use promoters, polyadenylation signals, and enhancers.

A nucleic acid is "operably linked" when it is placed into a functional relationship with
25 another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked"
30 means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

As used herein, the expressions "cell," "cell line," and "cell culture" are used interchangeably and all such designations include progeny. Thus, the words "transformants" and "transformed cells" include the primary subject cell and cultures derived therefrom without regard for the number of transfers. It is also understood that not all progeny will have precisely identical DNA content, due to deliberate or inadvertent mutations. Mutant progeny that have the same function or biological activity as screened for in the originally transformed cell are included. Where distinct designations are intended, it will be clear from the context.

In some aspects, the invention provides isolated adeno-associated vectors (AAVs). As used herein with respect to AAVs, the term "isolated" refers to an AAV that has been isolated from its natural environment (*e.g.*, from a host cell, tissue, or subject) or artificially produced. Isolated AAVs may be produced using recombinant methods. Such AAVs are referred to herein as "recombinant AAVs". Recombinant AAVs (rAAVs) preferably have tissue-specific targeting capabilities, such that a transgene of the rAAV will be delivered specifically to one or more predetermined tissue(s). The AAV capsid is an important element in determining these tissue-specific targeting capabilities.

Methods for obtaining recombinant AAVs having a desired capsid protein have been described (See, for example, U.S. Patent No. 7,906,111). A number of different AAV capsid proteins have been described, for example, those disclosed in Gao, *et al.*, *J. Virology* 78(12):6381-6388 (June 2004); Gao, *et al.*, *Proc Natl Acad Sci USA* 100(10):6081-6086 (May 13, 2003); and U.S. Patent No. 7,906,111; U.S. Patent No. 8,999,678. For the desired packaging of the presently described constructs and methods, the AAV9 vector and capsid, or the AAV2 vector and capsid, is preferred. However, it is noted that other suitable AAVs such as rAAVrh.8 and rAAVrh.10, or other similar vectors may be adapted for use in the present invention. Typically the methods involve culturing a host cell which contains a nucleic acid sequence encoding an AAV capsid protein or fragment thereof; a functional rep gene; a recombinant AAV vector composed of AAV inverted terminal repeats (ITRs) and a transgene; and sufficient helper functions to permit packaging of the recombinant AAV vector into the AAV capsid proteins.

The components to be cultured in the host cell to package a rAAV vector in an AAV capsid may be provided to the host cell in trans. Alternatively, any one or more of the required components (*e.g.*, recombinant AAV vector, rep sequences, cap sequences, and/or helper functions) may be provided by a stable host cell which has been engineered to contain one or more of the required components using methods known to those of skill in the art. Most suitably, such a stable host cell will contain the required component(s) under the control of an

inducible promoter. However, the required component(s) may be under the control of a constitutive promoter. In still another alternative, a selected stable host cell may contain selected component(s) under the control of a constitutive promoter and other selected component(s) under the control of one or more inducible promoters. For example, a stable host
5 cell may be generated which is derived from 293 cells (which contain E1 helper functions under the control of a constitutive promoter), but which contain the rep and/or cap proteins under the control of inducible promoters.

The recombinant AAV vector, rep sequences, cap sequences, and helper functions for producing the rAAV may be delivered to the packaging host cell using any appropriate genetic
10 element (vector). The selected genetic element may be delivered by any suitable method, including those described herein. See, *e.g.*, Fisher *et al*, *J. Virology* 70:520-532 (1993) and U.S. Patent No. 5,478,745.

In some embodiments, recombinant AAVs may be produced using the triple transfection method (*e.g.*, as described in detail in U.S. Patent No. 6,001,650). Typically, the
15 recombinant AAVs are produced by transfecting a host cell with a recombinant AAV vector (comprising a transgene) to be packaged into AAV particles, an AAV helper function vector, and an accessory function vector. An AAV helper function vector encodes the "AAV helper function" sequences (*i.e.*, rep and cap), which function in trans for productive AAV replication and encapsidation. Preferably, the AAV helper function vector supports efficient AAV vector
20 production without generating any detectable wild-type AAV virions (*i.e.*, AAV virions containing functional rep and cap genes). Non-limiting examples of vectors suitable for use with the present invention include pHLP19, described in U.S. Patent No. 6,001,650 and pRep6cap6 vector, described in U.S. Patent No. 6,156,303, the entirety of both incorporated by reference herein. The accessory function vector encodes nucleotide sequences for non-AAV
25 derived viral and/or cellular functions upon which AAV is dependent for replication (*i.e.*, "accessory functions"). The accessory functions include those functions required for AAV replication, including, without limitation, those moieties involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of cap expression products, and AAV capsid assembly. Viral-based accessory functions can be
30 derived from any of the known helper viruses such as adenovirus, herpesvirus (other than herpes simplex virus type-1), and vaccinia virus.

As used herein, the terms "AAV1," "AAV2," "AAV3," "AAV4," and the like refer to AAV vectors containing ITRs from AAV1, AAV2, AAV3, or AAV4, respectively, as well as capsid proteins from AAV1, AAV2, AAV3, or AAV4, respectively. The terms "AAV2/1,"

"AAV2/8," "AAV2/9," and the like refer to pseudotyped AAV vectors containing ITRs from AAV2 and capsid proteins from AAV1, AAV8, or AAV9, respectively.

With respect to transfected host cells, the term "transfection" is used to refer to the uptake of foreign DNA by a cell, and a cell has been "transfected" when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art. See, *e.g.*, Graham *et al.*, *Virology* 52:456 (1973), Sambrook *et al.*, *Molecular Cloning, a Laboratory Manual*, Cold Spring Harbor Laboratories, New York (1989), Davis *et al.*, *Basic Methods in Molecular Biology*, Elsevier (1986), and Chu *et al.*, *Gene* 13:197 (1981). Such techniques can be used to introduce one or more exogenous nucleic acids, such as a nucleotide integration vector and other nucleic acid molecules, into suitable host cells.

A "host cell" refers to any cell that harbors, or is capable of harboring, a substance of interest. Often a host cell is a mammalian cell. A host cell may be used as a recipient of an AAV helper construct, an AAV minigene plasmid, an accessory function vector, or other transfer DNA associated with the production of recombinant AAVs. The term includes the progeny of the original cell which has been transfected. Thus, a "host cell" as used herein may refer to a cell which has been transfected with an exogenous DNA sequence. It is understood that the progeny of a single parental cell may not necessarily be completely identical in morphology or in genomic or total DNA complement as the original parent, due to natural, accidental, or deliberate mutation.

With respect to cells, the term "isolated" refers to a cell that has been isolated from its natural environment (*e.g.*, from a tissue or subject). The term "cell line" refers to a population of cells capable of continuous or prolonged growth and division *in vitro*. Often, cell lines are clonal populations derived from a single progenitor cell. It is further known in the art that spontaneous or induced changes can occur in karyotype during storage or transfer of such clonal populations. Therefore, cells derived from the cell line referred to may not be precisely identical to the ancestral cells or cultures, and the cell line referred to includes such variants. As used herein, the terms "recombinant cell" refers to a cell into which an exogenous DNA segment, such as DNA segment that leads to the transcription of a biologically-active polypeptide or production of a biologically active nucleic acid such as an RNA, has been introduced.

The term "vector" includes any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, artificial chromosome, virus, or virion, which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

In some embodiments, useful vectors are contemplated to be those vectors in which the nucleic acid segment to be transcribed is positioned under the transcriptional control of a promoter. A "promoter" refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrases "operatively positioned," "operatively linked," "under control," or "under transcriptional control" means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

The term "expression vector" or "expression construct" or "construct" means any type of genetic construct containing a nucleic acid in which part or all of the nucleic acid encoding sequence is capable of being transcribed. In some embodiments, expression includes transcription of the nucleic acid, for example, to generate a biologically-active polypeptide product or inhibitory RNA from a transcribed gene.

As used herein, the term "TK2" refers to a gene encoding thymidine kinase 2, or the corresponding protein product. The terms "TK2" and "thymidine kinase 2" include wild-type forms of the TK2 gene or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion constructs, among others) of wild-type TK2 proteins and nucleic acids encoding the same. Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or more) to any of the amino acid sequences of a wild-type TK2 protein (e.g., SEQ ID NO: 1), provided that the TK2 variant retains the function of a wild-type TK2. Additionally, the terms "TK2" and "thymidine kinase 2" may refer to fusion proteins, or nucleic acids encoding the same, in which the TK2 is operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The term "TK2" may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary TK2 amino acid sequence (SEQ ID NO: 1):

MGAFQCRPSSDKEQEKEKKSVCVEGNASGKTTCLEFFSNATDVEVLTEPVSKWRNVGRHNPLGLMY
HD
ASRWGLTLQTYVQLTMLDRHTRPQVSSVRLMERSIHSARYIFVENLYRSGKMPEVDYVVLSEWFDWIL
RN
MDVSVDLIVYLRTNPETCYQRLKKRCREEEKVIPLEYLEAIHHLHEEWLIKSLFPMAAPVLVIEADHH
M
ERMLELFEQNRDRILTPENRKHCP

Exemplary TK2 nucleic acid sequence (SEQ ID NO: 2):

GCCAAGTTATGGGTGCGTTCTGCCAGCGTCTTAGCAGTGATAAAGAACAGGAAAAAGAGAAAAATCAGT
GATCTGTGTGCGAGGGCAATATTGCAAGTGGGAAGACGACATGCCTGGAATTCTTCTCCAACGCGACAGAC
GTGAGAGTGTTAACGGAGCCTGTGTCCAAGTGGAGAAATGTCCGTGGCCACAATCCTCTGGGCTGATGT
ACCACGATGCCCTCTCGCTGGGGTCTTACGCTACAGACTTATGTGCAGCTCACCATGCTGGACAGGCATAC
5 TCGTCCTCAGGTGTCTGTACGGTTGATGGAGAGGTCGATTACAGCGCAAGATACATTTTTGTAGAA
AACCTGTATAGAAGTGGGAAGATGCCAGAAGTGGACTATGTAGTTCTGTGGAATGGTTTACTGGATCT
TGAGGAACATGGACGTGTCTGTTGATTTGATAGTTTACCTTCGGACCAATCCTGAGACTTGTACCAGAG
GTTAAAGAAGAGATGCAGGGAAGAGGAGAAGGTCATTCCGCTGGAATACCTGGAAGCAATTCACCATCTC
CATGAGGAGTGGCTCATCAAAGGCAGCCTTTTCCCCATGGCAGCCCCTGTTCTGGTGATTGAGGCTGACC
10 ACCACATGGAGAGGATGTTAGAACTCTTTGAACAAAATCGGGATCGAATATTAACCTCCAGAGAATCGGAA
GCATTGCCCATAGGAGGCAAAAGGTCTATGGCTCATGTCTGAAAAATGCCTGCTGCTGCCAAGTTAGCTA
TTGGGAGCAATCTGGAAAACTTGTCTCCAGGAGGGCTTTGTGTCTGGCCAGCTTGATTTTCCATAATGGT
CTCATCTCCTTTGCTAGTGTCTTTGTCTATGCGTCTCTGGCCCTCGTGGGTAAATGACAAACGGGACCAAT
GGGTTTGCCAAGCCCTTTGCTGTTGCGAGCCTCACATTCCCCGGTGCCTCTCCCATGGCTTTGTGTCTGC
15 TGAGTCGCTCTCATGAAGCCCTTAGGGAGAGCACCTGTTGTGTGCTGACACCACGCTGGAGCTGTGTAC
CAATCGTCTCAGCCTTCATTAGGAGGCCGAGGTAGGAGTCTTATATCCCAGGTGAGGAATTTGAAGCTCA
GAAAGTTGAGGGGCTCCCCAGAGGTACACAGCCTGTGTGCAGTGGAGCTGGCACCATTGAGACTTTCA
GCCGACTCAGCAACTTTCCCTTGCCCTGGGCTGCCTCCTCCTGAGAGCTGTTCCCCACCGCCCTGCCTCT
TCCGGTTGGAGGCTCTCATGTCTCTTTGGGGAGAGCTGGCAGTGTGCGGAGCTGATAACATTTTCCCAAT
20 ATTGAGCAGTTCCTCAAGGACAGTCAGCATTTCTAGACTTCCACAAAATTATGCTGCATTTGGCTGGAGCC
CGGTGTTTCAGTGGTTTCCCTGCCCCGAGGTGCTGCGAGCCCATCTACCACATCTTCATGTGGACATTGAG
ATTCACATGCTGGCTCCTGAAGGGTGTCTAGTCTCCTTGGTGATTAAAGTCTGCTTGAAGTGTGCCAA
CTCCATGTGAGGGAAGTGTCTTTTGGTGCCTGGCTGGTTTGGCCAGAGCCAAGCTGGGGCAAGGGGCAGC
CAGCCCTGGCTTCCAAGGCTCCCGTACTGTCTGTGTCTTGTATAAGGAGCTTTGCTCTTGGAAATTACTG
25 AAAGTCTGTGGCCCAAGAGAGAGACACAAGTGGCCTTAAGTCTTTTTGAAGTGTTATTTTCATCCAGGGAA
ATGCCTCGAGCCATAGAGCCTGAAATCATCTTTGTTGGCTCAGAAAATACCTTAGCTTCACTCAGCTGGA
CTGCATTGAAGGCGAGGCTGCCCCCTGGATCAAGCAGAAAAACAAGAGAAAGAAAGAACGTTCCCTTTGGG
GATAGTCTGGAAGTTGGGATTTGCAAATAAAGGCTCTGGAAGCATTGCTGGTCTGAAGCTTTGGAGGT
GGGCAGAGAGAGCTTCAAGAAGACTAGATGCAAACCTGGAAAGGATTAAGGCTCAACTCTGGAGAAACA
30 GGCCACAGCCTCTCAGAGCAGCTGTTGGCTGTAAATAGAGGTAGCAAGGCCGCTCCCAGGCCCTGTGAG
TGTGGGCACCTGTGCATGCAATGCTCCGACTCTGCAGAGGTGCCAAGTGCCCTGCTGGGCCAGTCCCAG
AGAGTTAGGAAGTCAAGGCCTGCAACTCCTGGTTCTTCTGTTTGGACCAGTTCTTGTGCCATTGGCAGG
ATGAGAGGCAGCAGCCAGGCGGGAGCTGTGTCTAGCAGCACCTGTAGCCCACGTGCTGCTAATTAGCTGG
AAAAGTGGCGAAGGCAGAACCTTTGCTACCAGGAATCTTGACATGTGGGTCTGTCTTTGAGAAATTGTA
35 AATGAACAGTCCAATATTTCTCTCGGCTCATTTTGACATCCATTTTGGGGAAATGTGATTTCTCTCT
CTTTTTTTTTTTTTTTTTTTTGCCTAAGGCACAATCTCAAGAGGTCTGAGACCACGTACCCATGATTTTT
TTCTTGCTCTGTGATACCAATAACTCCTTCACCTAAGCCCTGTTGTTGATTTTGAAGTGCTTCTTAGGC
CAGTGATTTTAGCTTCTGCCAGCTGCTTTTGCCAGTAGATTAACGTGTTTTTATTTTTCAAACCTCCGTGT
TTCTTAACGTGGAGTGTATGGGTCTAAGAGAGCCTGTGTCTCCTGCCTTCCACCTTGGAGAGGAGGC
40 TGGACGCATCAGCAGTGGCCAGGGCAGGTGCAAAATCTCCAGCCTAGAGACCACACCTGAAACGGCTG
AAGCCAGCTTGACAAGGGCTGTGTCCCTCTGCGGCAGGCAGAGCTGGTGGGGGCAGGGGTCACAGAGC

AGTCATAGACACCATGGACCAGGGCAGGAGAAGGGCAGATGGCACATGGGCACAACAGGGCCTTGTCCTT
 AGAGCACTGGGGGGTCATGGCTGGGAGGGGCATGGCAGGGGCTGGCATCCCTGTAGAGCCAGAGGGGCCA
 CCCAGGGCAGTGACATTCCAGATATGTTGGGCTCACCTCATCCTTGCTGTGAGACTGGAGTTCCATGGGG
 ACATGAAGTCAGTACACCGCAGAGCTGCTCAGCTGCTCTACCTCTCGCTGACTTTTTTGTGACATATA
 5 CATTTCCTTTCAATTAGCATTATTTTACGCTTTTATTTAAGCTTTTTTGACAGTACATGTAAATATATGA
 TTATAACCATTAAAAATACCTTATGTACCTGGTTTTTTTTTGAAACTAGATAGAAATATATTTATCTTT
 TACATAAAAGAAGTGTGTAGTGGGTGGTCCAGGGCTTTGTGGTGGCTTAGTGGCCATCGGGGTCCCAGG
 CTCTTCCCACCTTTCTCTTTTGGGTCCACCTCTTGGCTCCTGGCTTCCTTCCTCTGGGGCCTGACTGTCC
 AGGATGGCAGCTGGAGCTCCTGCTCTGGGCACGGTTGGCATCAGAGCCCACTGCTCCCCATCCACTCTTT
 10 AATCCAGATGCTGGTAATGTCCCTTTCCAAGGCATAACTAAGATAAGCTGGAAGGTTCTTACAGAGGCTTT
 GCTAGAGGCCCTGGGAGGTGGGGGAAAGGCAAGAGGGCAGTGCCACCCATAGACCGGGTCACATGACC
 TGGCATCAGCGCCGTGGGGTCCCTTTGGGCTTCACCTCCCTCTCCCTCTGCCCCACGTCATCCCACTT
 CATCTCACCTCACCATTTCCCATCCCGTACCGTAATGTCCCTTGCAAGGCCTAACTCTGTGAGGAATGA
 AACCACCTCATCCTTGTTCCAATAACATGAGTGACCGAATTTACAAACGAGTGAATGTGGACCTCTGGAA
 15 ACATTACCCAGCTTGTTTTTACTTTTCACTTTCTCTTTCTGCCCCCTTTCATTTCCGTAGGAGCCCTTTAC
 CTAGATGAGAAGTGTCCCCCGCTGGGGAATATATCAGTCAGAACAATCTTCCTGCAGACATGCACCAT
 TAGACCCGAGTGACGGTGGTGCCATTTAAACCTCAGAGCAGGTAAAAGGTGGTCTGAAACCTGTCTACC
 CACAGGGTGCTATGGAATCTGAATCACTTCTTTTTTCTAGAGCCCTGGGGTGGGGAGCTCCCTCAAGTG
 TTCACATGTGTGTGGAATGAGGAACACCCATCTCCTTGGCCCTCTCCACCTGAAGAGTTAGTTATTTAAA
 20 ATAATTGGCAAGCTCTTGCAATGTCAGTCATCCATTGTTTCAAGATGGAATAGCAATAATACATCCCTGG
 CTGCCCTGGGCTTGGCCAGGATTACTCACTGAAGGCCTCAGGGTTACTGGCACACACTTTCTTTTCTTAA
 TAATCCCATCCCCTCAGCTTTTCTAAGGCTAGAGTGAATTTTCGTGTTCTTTAGTTTACATAAGATGGTG
 AACTTGGCAAAAGCTATCATTAAACAGAAGCTAAGAGAAAGCCTATGTCGTGGAATCCAGAATGGGTATT
 GCCATTCACTGCTGTCCACAGAAGCTGTCTTGAATTTCTTTCTGTGTCTTTCTTTTTTTTTCTTTAAGA
 25 CTGTTGTTTACCAGACTGGGCTCTGTGGAACACAGGIGTCCTGGGAGATGGTTAATCATTACAAAATATT
 GGTAACAATCTAAAGATGCATACATAAGAGAGTGGTCAAATAAACCATTTTCCATTCA

As used herein, the term "dGK" refers to a gene encoding deoxyguanosine kinase, or
 the corresponding protein product. The terms "dGK" and "deoxyguanosine kinase" include
 wild-type forms of the dGK gene or protein, as well as variants (e.g., splice variants,
 30 truncations, concatemers, and fusion constructs, among others) of wild-type dGK proteins and
 nucleic acids encoding the same. Examples of such variants are proteins having at least 70%
 sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%,
 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%,
 98%, 99%, or 99.9% identity, or more) to any of the amino acid sequences of a wild-type dGK
 35 protein (e.g., SEQ ID NO: 3), provided that the dGK variant retains the function of a wild-type
 dGK. Additionally, the terms "dGK" and "deoxyguanosine kinase" may refer to fusion
 proteins, or nucleic acids encoding the same, in which the dGK is operably linked to another
 polypeptide, half-life-modifying agent, or therapeutic agent. The term "dGK" may refer to the

protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary dGK amino acid sequence (SEQ ID NO: 3):

MAAGRLFLSRLRAPFSSMAKSPLEGVSSSRGLHAGRGPRRLSIEGNIAVGKSTFVKLLTKTYPEWHVATE
 5 PVATWQNIQAAGTQKACTAQLGNLLDMMYREPARWSYTFQTFSLRLKVLQLEPFPEKLLQARKPVQIF
 ERLHFEALMNIPVLVLDVNDDFSEEVTKQEDLMREVNTFVKNL

Exemplary dGK nucleic acid sequence (SEQ ID NO: 4):

AACGGTGCCTGGAGCGAGTGAGCAGCGATACCTAGGGCGGAAGTGCTCTCGGCGGAAGTGATCGCTGTG
 TGAATCGTGGGTGGGATGGCCGCGGGCCGCTCTTTCTAAGTCGGCTTCGAGCACCCCTTCAGTTCATGG
 10 CCAAGAGCCCACTCGAGGGCGTTTCTCTCCAGAGGCCTGCACGCGGGGCGGGGCCCCGAAGGCTCTC
 CATCGAAGGCAACATTGCTGTGGGAAAGTCCACGTTTGTGAAGTTACTCACGAAAACCTACCCAGAATGG
 CACGTAGCTACAGAACCTGTAGCAACATGGCAGAATATCCAGGCTGCTGGCACCCAAAAAGCCTGCACTG
 CCAAAGTCTTGAAAACTTGCTGGATATGATGTACCGGAGCCAGCACGATGGTCTACACATTCCAGAC
 ATTTTCCTTTTGTAGCCGCTGAAAGTACAGCTGGAGCCCTTCCCTGAGAACTCTTACAGGCCAGGAAG
 15 CCAGTACAGATCTTTGAGAGGCTCCACTTTGAGGCTCTGATGAACATTCCAGTGCTGGTGTGGATGTCA
 ATGATGATTTTTCTGAGGAAGTAACCAAACAAGAAGACCTCATGAGAGAGGTAAACACCTTTGTAAAGAA
 TCTGTAACCAATACCATGAAGTTCAGGCTGTGATCTGGGCTCCCTGACTTTCTGAAGCTAGAAAAATGTT
 GTGTCTCCCAACCACCTTTCCATCCCCAGCCCCCTCTCATCCCTGGAGCACTCTGCCGCTCAAGAGCTGGT
 TTGTTAATTATGTTAGACTTTGCCATTGTTTTCTTTGTACCTGAAGCATTTTGAATAAAGTTTACT
 20 TAAGTTATGCTTGTTTTTCTAAAAAATAAAGTTTACT

As used herein, the term "p53R2" refers to a gene encoding p53 inducible subunit of ribonucleotide reductase, or the corresponding protein product. The terms "p53R2" and "p53 inducible subunit of ribonucleotide reductase" include wild-type forms of the p53R2 gene or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion
 25 constructs, among others) of wild-type p53R2 proteins and nucleic acids encoding the same. Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or more) to any of the amino acid sequences of a wild-type p53R2 protein (e.g., SEQ ID NO: 5),
 30 provided that the p53R2 variant retains the function of a wild-type p53R2. Additionally, the terms "p53R2" and "p53 inducible subunit of ribonucleotide reductase" may refer to fusion proteins, or nucleic acids encoding the same, in which the p53R2 is operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The term "p53R2" may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated
 35 by one of skill in the art.

Exemplary p53R2 amino acid sequence (SEQ ID NO: 5):

MLLLRLPPHRSHASPLDCKLQDRCKCYSPRSGQACPPALAAWLRRRCERRGGRPRGGRRKELTLGLRPA

RCSAPGPAKDDAWRPQAGRSSSDTNESEIKSNEEPLLRKSSRRFVIFPIQYPDIWKMYKQAQASFWTAE
 VDLKDLPHWNKLKADEKYFISHILAFFAASDGIVNENLVERFSQEVQVPEARCFYGFQILIENVHSEMY
 SLLIDTYIRDPKKREFLFNAIETMPYVKKKADWALRWIADRKSTFGERVVAFAAVEGVFFSGSFAAIFWL
 KKRGLMPGLTFSNELISRDEGLHCDFACLMFQYLVNKPSEERVREIIVDAVKIEQEFLTEALPVGLIGMN
 5 CILMKQYIEFVADRLLEVLFSGVSKVFAQENPFDFMENISLEGKTNFFEKRVSSEYQRFVMAETTDNVFTLD
 ADF

Exemplary p53R2 nucleic acid sequence (SEQ ID NO: 6):

ATGAGGTAATGTGTGCTGTTGCCGCTTCCCCCTCACCGCAGTCACGCCAGCCCCTTAGATTGCAAGTTGC
 AGGACCGCTGTAGGAAATGTTATTCGCCGCGGTGAGGACAGGCCTGTCCGCCCGCCCTCGCCGAGCCTG
 10 GCTTCGTCGTTGCGAGCGCCGGGAGGCCGTCCCCGGGAGGGCGGAGGAAGGAGCTGACTTTGGGTTTG
 CGTCCCGCTCGCTGCTGTGCCCCGGGGCCAGCCAAGGACGACGCTTGGAGGCCTCAGGCCGGGAGATCAT
 CTTCAGACACCAACGAAAGTGAAATAAAGTCAAATGAAGAGCCACTCCTAAGAAAGAGTTCTCGCCGGTT
 TGTCTCTTTTCCAATCCAGTACCCTGATATTTGGAAAATGTATAAACAGGCACAGGCTTCCTTCTGGACA
 GCAGAAGAGGTCGACTTATCAAAGGATCTCCCTCACTGGAACAAGCTTAAAGCAGATGAGAAGTACTTCA
 15 TCTCTCACATCTTAGCCTTTTTTGCAGCCAGTGATGGAATTGTAAATGAAAATTTGGTGGAGCGCTTTAG
 TCAGGAGGTGCAGGTTCCAGAGGCTCGCTGTTTCTATGGCTTTCAAATTCTCATCGAGAATGTTCACTCA
 GAGATGTACAGTTTGTGATAGACACTTACATCAGAGATCCCAAGAAAAGGGAATTTTTATTTAATGCAA
 TTGAAACCATGCCCTATGTTAAGAAAAAAGCAGATTGGGCCTTGCGATGGATAGCAGATAGAAAATCTAC
 TTTTGGGGAAAGAGTGGTGGCCTTTGCTGCTGTAGAAGGAGTTTTCTTCTCAGGATCTTTTGTGCTATA
 20 TTCTGGCTAAAGAAGAGAGGTCTTATGCCAGGACTCACTTTTTCCAATGAACTCATCAGCAGAGATGAAG
 GACTTCACTGTGACTTTGCTTGCCTGATGTTCCAATACTTAGTAAATAAGCCTTCAGAAGAAAGGGTCAG
 GGAGATCATTGTTGATGCTGTCAAAATTGAGCAGGAGTTTTTAACAGAAGCCTTGCCAGTTGGCCTCATT
 GGAATGAATTGCATTTTGATGAAACAGTACATTGAGTTGTAGCTGACAGATTACTTGTGGAACCTGGAT
 TCTCAAAGGTTTTTCAGGCAGAAAATCCTTTTGATTTTATGGAAAACATTTCTTTAGAAGGAAAAACAAA
 25 TTTCTTTGAGAAACGAGTTTCAGAGTATCAGCGTTTTGCAGTTATGGCAGAAACCACAGATAACGTCCTTC
 ACCTTGGATGCAGATTTTTAAAAAACCTCTCGTTTTAAACTCTATAAACTTGTCAATTGGTAAATAGTAG
 TCTATTTTCTCTGCTTAAAAAAATTTTAAGTATATCCTTTAAAGGACTGGGGGTTTGTCAAAAGGAA
 ATCCAAAACCTATTCTAAACAATTTGCATTTATATAATTTTCTGTTTAAACAACAGAGTGTGACCTAAA
 TGCTTTTGTCTGTCACTGAAATAAAGATGGCATTATGTGGTTAAGAGCATGGGGCGAGGGGTGACACA
 30 TGAGTCTAAGGTTCTGCCCTTACTCCAGTGTGTGACCTTGGCAAGTCAGTTAATCTTGGTAAACCTCGG
 TGTACTTATCTTTAAATGGGAGTAATAGTAGGTCCTAAATTCATAGAGTGGATATTAGGATTAGGATGC
 AAAATAAATGCTTAACCAACACTACTACTGTTAGCACCACCTACTAATTATCATTGATAATATTA
 TTGCAATGATGTTGTAATAAAATACTCTCATTTCTTAAATAAATTGTGATTCTAGGTCCTAGGATCTAG
 AATTAGATCTTTGTATTTTAAATGCTTAGGGGAAGAATATAAGTATCTCCTTAAAAAGAACATAATTCTC
 35 ATTCACGCAAGAATAAGTTCTTTGAATTCCTTAGTATGTAGTGAAGAAAATTTAGTTGTTAGTTGCTTTG
 GGAAGCCTACTTATGGAGTGGAAACCAGGAGGTTATCATGGTAGTTGACCTTATAAGAAAAATGATTCTT
 CTTCAGAAATTAACATAACTATTGCCAGATTTAGCTCTGGAATGTTTGAATCAGGCTAGAATAGCA
 TTTTCCAAAGAAATATTCTAAGAGCTATTAGCTCCTCTAGATATTTTTTGGGGGAAAAAGGGGATTCTGT
 GGTGAGATGAGTTTGGGAAATGCTGAACACTTCATTCTCTTTAGCAAGTACAGTCAGTACATCAAAGAC
 40 TGAGCAGTTGAGTTGACATAAATTTATCTCGCCCTGCATATTTCCCAACATACTTAACACAGATGTTTTT
 TACCTGTTAACATCTCACCCAGCTAGTGTTCTCAGAACAAAGATTGGAAAAAGCTGGCCGAGAACCATT

TATACATAGAGGAAGGGCTTATGGACTGAGAAAGGGAGAACATGGTAGGGATTATTGAATCATTTCAAAT
 TTATACCAGCCTGAATAGTGTACCAGCAATTGACTTAGGCTGTGTTTCTTTATGGTTTTAAACTCTTGA
 GCTGTTATAAGAGATAGTTCTTTTAATGTGACTATGCAACATGATAGCCAATGGTGAGGGAAAAGGAGGT
 TTCTCTAGAAGAGTCTGATGAAAGGCCGGGAACCAAGGTTTTTGAGAAGTCTGCCCTATTTATTTTTAG
 5 TAAGTATCAAGAGGTAGCCTGAGCCTAGTTAGAGTTAGACCTGTCTTTGGATGAAGAAGTCTTAATACTG
 AAATACTGAATTTTTAATACATTATTATTTGGTATTCTGTATACCCCTTCAAGCAGTTGTTTCCCATTCC
 CAACAACTGTACTTTATACAATTCTGGATGCTAAACTTAGAGATTTTCTCTTGCATAAATTTGGCT
 CCATTCTTTCCATAACAATCTAATCAAACTGGGAGTTCTCAAGTGAATGCAAAAGGAGCAGGCCATAAC
 TTTATTTGTTAGATACACTGTCAGAACTTGAGATCTTTTGGCCTATGATAATACCATTAAATTTTGCAT
 10 TGCTTCAGTTTGCCAAGTGTTTTTACATCATCTCATTGATCTCAAAACAGCTTGACAGAGCAACTGTTA
 TTGAAATATTACAGATGGAAGAATGAGGCTCAGGGAAGTTAAATGACTTGCCAAGATCTGCTCATCGT
 CACTGTCTGTACAGTATTTTTTTTTAGAGGTTGTAATGTCTCAGATTTAGTCCTTTACCATCTATGTTGA
 TTTGCTTTTGCTATTTTCTCATTAAATGAATATACTTTAAATATATATATTAAAGTATCAAAATATAGA
 GAGACATTTGAACTGTATTACAGTAATATGTTTAAAGATATTTATATATTGCCATACAAAACTTAACAT
 15 TTAAACTGATAATATCTGTAATGACATCAGAATGAAAGAAAAAAATTGTACAGTGTATATTCCTTTGT
 TTTGAATCCAACTCTTTTTTCATAGTAATGACAGATGCCTTAATGTGAAGCTTATTTATAATAGCAATAA
 ACCTAACTGGATTTGGATGAAGAAGTCTTAATACTGACATACTGGATTTTAAATGCACTGGTTTGTATT
 TGGTATTCTATCTCTTTTTCCAGGCCTCCAGGTTGCACATTTATTTATTATGTTCAATACTTTGGTTCTT
 AGTTCTTAAAGAATCAAGAAGTTGTGTAATCTTTTAAAAATATTATCTTGACAGATAAAGAAAAAATTAA
 20 GAGTGTGTTTACAACCTGTTTTCTCTTTTTTACAGTACATGTATTTAAATCATTGCTATAATAAAGTTAAG
 TTCATTAGGAATATAAAAACTTGCACTTCTATGATAGATTGCATTTATTTAAAAATGTTTCATTGTATCAC
 ATAGAAATATGGCCAGGAAGGACTTGAGAAGACAGTTTGATCCATTGCTTTTAGACAGGACTGGGTTTTG
 CTGTCCAATTATATACAATAATAGTTTTTCTTACAACCTAAGCTGGCCCCAGCCTTGCTCTTGATATTAATA
 CATGAAATTTTTATAATTGTCTCATTGTCTCATTAGAAACATCCATATTTTTCTGCTTTTTCTATTGCC
 25 ATTTTTTATTTGTGCATGAATTGATTATTGAGAAAATGTAGCAGTTGCATATTTAAAAATTAATCATTT
 TGCATTTTACATTTAAATATGCTAACATCACTGTATAGAATTCCCAAATTTTCAATTTGTAGATACTGAAC
 TAAGGGCTAATGTCAGGAGCTGATTTTAAATGATAAAGCTGCAGATGGGCTAAATAAAGCCAAATTAAT
 CCTACAATCAGGTATTATGTTTTTAAACCAAGTTGAGTGAATTGGTAGTGGACTTGGGAAATCTTCCCCA
 GCAGAATCTGGATGAATGGCACAGAATTGAAATCTCTTTGTTTCCCACCATTTCCTTTAAGTGCTCTGC
 30 TCCTTTGTAAAAAGTTAAAGATTTGAAAGAGAATCTCATATTTCCCGAGGCATTAGGAAGAAAGGATTTAA
 TCCCTTCAATTTGGGGCTTAATCTTGTTTTAAAAAATGTAAGTGAAGATGGAAGGCTGGAGAGAATGATT
 GCTTTTTGTACAGTTAAATAAGGTCACAATATTCTTACATACTTTGTTTTACAACCTGTGTTTTCATTTTT
 TCAATGTCTGGCCATTTAGCAAAGTTATTTACTATTTACTGTGTACATAGAAAGCTTTATTATGTGTGG
 TGTATCTAAATTTTTTTTGCTGAAATACATTATGGTCAATCAAGCCAAGCCTGCATGTACAGAATTTGTT
 35 TTTTTTCAAATAAATTAGTTGTTTTCTTATTTTTTTGGCTTAGTATGTTGAAATAAACTATGGTATCTT
 CATCATTTTGTACATTTTCTTTTTGAGGAAGGTTTCTTTATAAGTGCAAGGGCTACCCTAATAAAGGAAT
 GTATATACTTACT

As used herein, the term "TP" refers to a gene encoding thymidine phosphorylase, or
 the corresponding protein product. The terms "TP" and "thymidine phosphorylase" include
 40 wild-type forms of the TP gene or protein, as well as variants (e.g., splice variants, truncations,

concatemers, and fusion constructs, among others) of wild-type TP proteins and nucleic acids encoding the same. Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or more) to any of the amino acid sequences of a wild-type TP protein (e.g., SEQ ID NO: 7), provided that the TP variant retains the function of a wild-type TP. Additionally, the terms “TP” and “thymidine phosphorylase” may refer to fusion proteins, or nucleic acids encoding the same, in which the TP is operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The term “TP” may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary TP amino acid sequence (SEQ ID NO: 7):

MAALMTPGTGAPPAPGDFSGEGSQGLPDPSPPEPKQLPELIRMKRDGGRLSEADIRGFVAAVVNGSAQGAQ
IGAMLMAIRLRGMDLEETSVLTQALAQSGQQLWEPAWRQQLVDKHSTGGVGDQVSLVLAPALAAACGCKV
PMISGRGLGHTGGTLDKLESIPGFNVIQSPEQMQLLDQAGCCIVGQSEQLVPADGILYAARDVTATVDS
LPLITASILSKKLVEGLSALVVDVKFGGAAVFPNQEQARELAKTLVGVGASLGLRVAAALTAMDKPLGRC
VGHALEVEEALLCMDGAGPPDLRDLVTTLGGALLWLSGHAGTQAQGAARVAAALDDGSALGRFERMLAAQ
GVDPGLARALCSGSPAERRQLLPRAREQEELLAPADGTVELVRALPLALVLHELGAAGRSRAGEPLRLGVG
AELLVDVGQRLRRGTPWLRVHRDGPALSGPQSRALQEALVLSDRAPFAAPSPFAELVLPQQ

Exemplary TP nucleic acid sequence (SEQ ID NO: 8):

CGACTGCCGAGCTCCGCCCTCCAGGCGGCCCCACCCGCTGCCGTCTGGGGCGCCGCCGCCGCCGCCGCC
GGCAGTGGACCGCTGTGCGCAACCCTGAACCCTACGGTCCCGACCCGCGGGCGAGGCCGGGTACCTGGG
CTGGGATCCGGAGCAAGCGGGCGAGGGCAGCGCCCTAAGCAGGCATCCCCGAGGCCCGGAGCGATGGCA
GCCTTGATGACCCCGGAACCGGGGCCCCACCCGCGCTGGTGACTTCTCCGGGAAGGGAGCCAGGGAC
TTCCCGACCTTCGCCAGAGCCCAAGCAGCTCCCGAGCTGATCCGCATGAAGCGAGACGGAGGCCGCT
GAGCGAAGCGGACATCAGGGGCTTCGTGGCCGCTGTGGTGAATGGGAGCGCGCAGGGCGCACAGATCGGG
GCCATGCTGATGGCCATCCGACTTCGGGGCATGGATCTGGAGGAGACCTCGGTGCTGACCCAGGCCCTGG
CTCAGTCGGGACAGCAGCTGGAGTGGCCAGAGGCCTGGCGCCAGCAGCTTGTGGACAAGCATTCCACAGG
GGGTGTGGGTGACAAGGTCAGCCTGGTCTCGCACCTGCCCTGGCGGCATGTGGCTGCAAGGTGCCAATG
ATCAGCGGACGTGGTCTGGGGCACACAGGAGGCACCTTGGATAAGCTGGAGTCTATTCTGGATTCAATG
TCATCCAGAGCCAGAGCAGATGCAAGTGCTGCTGGACCAGGCGGGCTGCTGTATCGTGGGTGACAGTGA
GCAGCTGGTTCTGCGGACGGAATCCTATATGCAGCCAGAGATGTGACAGCCACCGTGGACAGCCTGCCA
CTCATCACAGCCTCCATTCTCAGTAAGAACTCGTGGAGGGGCTGTCCGCTCTGGTGGTGGACGTTAAGT
TCGGAGGGGGCCCGCTCTTCCCCAACCCAGGAGCAGGCCCGGGAGCTGGCAAAGACGCTGGTTGGCGTGGG
AGCCAGCCTAGGGCTTCGGGTGCGGCAGCGCTGACCGCCATGGACAAGCCCTGGGTGCTGCGTGGGC
CACGCCCTGGAGGTGGAGGAGGCGCTGCTCTGCATGGACGGCGCAGGCCCGCCAGACTTAAGGGACCTGG
TCACCACGCTCGGGGGCGCCCTGCTCTGGCTCAGCGGACACGCGGGGACTCAGGCCAGGGCGCTGCCCG
GGTGGCCGCGGCGCTGGACGACGGCTCGGCCCTTGGCCGCTTCGAGCGGATGCTGGCGGCGCAGGGCGTG

GATCCCGGTCTGGCCCGAGCCCTGTGCTCGGGAAGTCCCGCAGAACGCCGGCAGCTGCTGCCTCGCGCCC
 GGGAGCAGGAGGAGCTGCTGGCGCCCGCAGATGGCACCGTGGAGCTGGTCCGGGCGCTGCCGCTGGCGCT
 GGTGCTGCACGAGCTCGGGGCGGGCGCAGCCGCGCTGGGGAGCCGCTCCGCTGGGGGTGGGCGCAGAG
 CTGCTGGTTCGACGTGGGTCTAGAGGCTGCGCCGTGGGACCCCTGGCTCCGCTGCACCGGGACGGCCCCG
 5 CGCTCAGCGGCCCCGAGAGCCGCGCCCTGCAGGAGGCGCTCGTACTCTCCGACCGCGCGCCATTGCGCCG
 CCCCTCGCCCTTCGCAGAGCTCGTTCTGCGCCCGCAGCAATAAAGCTCCTTTGCCGCGAAAAA

As used herein, the term "SUCLA2" refers to a gene encoding succinate-CoA ligase ADP-forming beta subunit, or the corresponding protein product. The terms "SUCLA2" and "succinate-CoA ligase ADP-forming beta subunit" include wild-type forms of the SUCLA2
 10 gene or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion constructs, among others) of wild-type SUCLA2 proteins and nucleic acids encoding the same. Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or
 15 more) to any of the amino acid sequences of a wild-type SUCLA2 protein (e.g., SEQ ID NO: 9), provided that the SUCLA2 variant retains the function of a wild-type SUCLA2. Additionally, the terms "SUCLA2" and "succinate-CoA ligase ADP-forming beta subunit" may refer to fusion proteins, or nucleic acids encoding the same, in which the SUCLA2 is operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The
 20 term "SUCLA2" may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary SUCLA2 amino acid sequence (SEQ ID NO: 9):

MAASMFYGRVLAVATLRNHRPRTAQRAAAQVLGSSGLFNNHGLQVQQQQRNLSLHEYMSMELLQEAGVS
 VPKGYVAKSPDEAYAIKKLGSKDVVIKAQVLGGRGKGTFFESGLKGGVKIVFSPEEAKAVSSQMIGKKL
 25 FTKQTGEKGRICNQVLVCERKYPREYYFAITMERSFQGPVLIGSSHGGVNIEDVAAESPEAIKEPIDI
 EEGIKKEQALQLAQKMGFPPNIVESAAENMVKLYSLFLKYDATMIEINPMVEDSDGAVLCMDAKINFDSN
 SAYRQKKIFDLQDWTQEDERDKDAKANLNYIGLDGNIGCLVNGAGLAMATMDIIKLHGGTPANFLDVGG
 GATVHQVTEAFKLITSDKKVLAILVNIFGGIMRCDVIAQGIVMAVKDLEIKIPVVRLQGTRVDDAKALI
 ADSGLKILACDDLDEAARMVVKLSEIVTLAKQAHVDVKFQLPI

30 Exemplary SUCLA2 nucleic acid sequence (SEQ ID NO: 10):

CCGCCTGTCGCTGTGCGCTGCGCGCGGCGCCGAGGGGACGGGGTCCGACTCAGAAATGGCGGCCTCCA
 TGTTCTACGGCAGGCTAGTGGCCGTGGCCACCCTTCGGAACCACCGGCCTCGGACGGCCAGCGGGCTGC
 TGCTCAGGTTCTGGGAAGTTCTGGATTGTTTAAATAACCATGGACTCCAAGTACAGCAGCAACAGCAAAGG
 AATCTCTCACTACATGAATACATGAGTATGGAATTATTGCAAGAAGCTGGTGTCTCCGTTCCCAAAGGAT
 35 ATGTGGCAAAGTCACCAGATGAAGCTTATGCAATTGCCAAAAAATTAGGTTCAAAAGATGTCGTGATAAA
 GGCACAGGTTTTAGCTGGTGGTAGAGGAAAAGGAACATTTGAAAGTGGCCTCAAAGGAGGAGTGAAGATA
 GTTTTCTCTCCAGAAGAAGCAAAAGCTGTTTCTTCACAAATGATTGGGAAAAAATTGTTTACCAAGCAAA

CGGGAGAAAAGGGCAGAATATGCAATCAAGTATTGGTCTGTGAGCGAAAAATATCCCAGGAGAGAATACTA
 CTTTGAATAACAATGGAAAGGTCATTTCAAGGTCCTGTATTAATAGGAAGTTCACATGGTGGTGTCAAC
 ATTGAAGATGTTGCTGCTGAGTCTCCTGAAGCAATAATTAAGAACCTATTGATATTGAAGAAGGCATCA
 AAAAGGAACAAGCTCTCCAGCTTGACAGAAAGATGGGATTTCACCTAATATTGTGGAATCAGCAGCAGA
 5 AAACATGGTCAAGCTTTACAGCCTTTTTCTGAAATACGATGCAACCATGATAGAAATAAATCCAATGGTG
 GAAGATTGATGGAGCTGTATTGTGTATGGATGCAAAGATCAATTTTGACTCTAATTCAGCCTATCGCC
 AAAAGAAAATCTTTGATCTACAGGACTGGACCCAGGAAGATGAAAGGGACAAAGATGCTGCTAAGGCAAA
 TCTCAACTACATTGGCCTCGATGGAAATATAGGCTGCCTAGTAAATGGTGCTGGTTTGGCTATGGCCACA
 ATGGATATAATAAACTTCATGGAGGGACTCCAGCCAACTTCCTTGATGTTGGTGGTGGTGTACAGTCC
 10 ATCAAGTAACAGAAGCATTAAAGCTTATCACTTCAGATAAAAAGGTAAGTGGCTATTCTGGTCAACATTTT
 TGGAGGAATCATGCGCTGTGATGTTATTGCACAGGGTATAGTCATGGCAGTAAAAGACTTGAAATTAAA
 ATACCTGTTGTGGTACGGTTACAAGGTACACGAGTCGATGATGCTAAGGCACTGATAGCGGACAGTGGAC
 TTAATACTTTGCTTGTGATGACTTGGAATGAAGCTGCTAGAATGGTTGTAAAGCTCTCTGAAATAGTGAC
 CTTAGCGAAGCAAGCACATGTGGATGTGAAATTTCAAGTTGCCAATATGATCTGAAAACCCAGTGGATGGC
 15 TGAAGGTGTTAAATGTGCTATAATCATTAAAGAATACTGTGTTCTGTGTTATTGTTCTTTTTCTTTTAGT
 GTGTGGAGATTGTAATTGCCATCTAGGCACACAAACATTTAAAAGGATTTGGACTGCATTTAATTGTACC
 ATTCAGAATGGACTGTTTGTACGAAGCATGTATAATGCAGTTATCTTCTTTCTTTGTGCGACCCAGTCT
 TTTTGTCTTCTCTACAAAACGTAACCTTGCAATTTGCCAGTTTATTATTGTTGGATACAAAGTCTTCAT
 TGATAAGAGTCCATAAAATAAGATAAAATACGAAGATAAAGCTTTATTCTTTAGTGTTAAATACAGTATA
 20 TCTAATAACTAGCCTCATTAGTAGAGCAGTATATTAACAATGTTTTATGTAAAAAGTGTATTCTTCA
 GCACCAAATACATGATAAATGTATCAATCACTATTTATAAACAGAGCTTTCAAACACTCCTCAGAATATT
 CTTCTAAGTATTTTGATGAAGTAACTTTGTAATTATTTGAACATTGTTTTAATCATTAGGAAACACTGAT
 TAACTGCAAGTCTTCATGATTCTGTCATATTAAGAAACACCTGTAGGTTTGCTTCAAATAAAGGCATATA
 TACCAAGGACTTACAGACAAAATTAAGAATGTCAATTTAAGTTAATAAAAAATCTCCCAATATGAAAAAA
 25 AAAAAAAAAA

As used herein, the term "SUCLG1" refers to a gene encoding succinate-CoA ligase
 ADP-forming alpha subunit, or the corresponding protein product. The terms "SUCLG1" and
 "succinate-CoA ligase ADP-forming alpha subunit" include wild-type forms of the SUCLG1
 gene or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion
 30 constructs, among others) of wild-type SUCLG1 proteins and nucleic acids encoding the same.
 Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%,
 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,
 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or
 more) to any of the amino acid sequences of a wild-type SUCLG1 protein (e.g., SEQ ID NO:
 35 11), provided that the SUCLG1 variant retains the function of a wild-type SUCLG1.
 Additionally, the terms "SUCLG1" and "succinate-CoA ligase ADP-forming alpha subunit"
 may refer to fusion proteins, or nucleic acids encoding the same, in which the SUCLG1 is
 operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The

term “SUCLG1” may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary SUCLG1 amino acid sequence (SEQ ID NO: 11):

MTATLAAAADIATMVSGSSGLAAARLLSRSFLLPQNGIRHCSYASRQHLYVDKNTKIICQGFTGKQGT
 5 HSQQALEYGTKLVGGTTPGKGGQTHLGLPVFNTVKEAKEQTGATASVIYVPPPFAAAAINEAIEAEIPLV
 VCITEGIPQQDMVRVKHKLRLQEKTRLIGPNCPGVINPGECKIGIMPGHIHKKGRIGIVSRSGTLTYEAV
 HQTQVGLGQSLCVGIGGDPFNGTDFIDCLEIFLNDSETEGIIILIGEIGNAEENAAEFLKQHNSGPNK
 PVVSFIAGLTAPPGRMRMGHAGAIAGGKGGAKEKISALQSAGVVVMSPAQLGTTIYKEFEKRKML

Exemplary SUCLG1 nucleic acid sequence (SEQ ID NO: 12):

10 GTCATTGGCGTATGACCGCAACCCTTGCCGCTGCCGCTGACATCGCTACCATGGTCTCCGGCAGCAGCGG
 CCTCGCCGCCGCCGCTCTCCTGTCGCGCAGCTTCCTCCTGCCGCGAATGGAATTCGGCATTGTTCTTAC
 ACAGCTTCTCGGCAACATCTCTATGTTGATAAAAATACAAAGATTATTTGCCAGGGTTTCACTGGCAAAC
 AGGGCACCTTTACAGCCAGCAGGCATTGGAATATGGCACCAAACTCGTTGGAGGAACCACTCCAGGGAA
 AGGAGGCCAGACACATCTGGGCTTACCTGTCTTTAATACTGTGAAGGAGGCCAAAGAACAGACAGGAGCA
 15 ACGGCTTCTGTCATTTATGTTCTCCGCTTTTGCTGCTGCTGCCATTAATGAAGCTATTGAGGCAGAAA
 TTCCCTTGGTTGTGTATCACTGAAGGAATTCCCCAGCAGGACATGGTACGAGTCAAGCACAAACTGCT
 GCGCCAGGAAAAGACAAGGCTAATTGGGCCCACTGCCCTGGAGTCATCAATCCTGGAGAATGTAAATT
 GGCATCATGCCTGGCCATATTCACAAAAAAGGAAGGATTGGCATTGTGTCCAGATCTGGCACCTGACTT
 ATGAAGCAGTTCACCAACAACGCAAGTTGGATTGGGGCAGTCTTTGTGCGTTGGCATTGGAGGTGATCC
 20 TTTTAATGGAACAGATTTTATTGACTGCCTCGAAATCTTTTGAACGATTCTGCCACAGAAGGCATCATA
 TTGATTGGTGAAATTGGTGGTAATGCAGAAGAGAATGCTGCAGAATTTTGAAGCAACATAATTCAGGTC
 CAAATTCCAAGCCTGTAGTGTCTTCATTGCTGGTTTAACTGCTCCTCCTGGGAGAAGAATGGGTCATGC
 CGGGGCAATTATTGCTGGAGGAAAAGGTGGAGCTAAAGAGAAGATCTCTGCCCTTCAGAGTGCAGGAGTT
 GTGGTCAGTATGTCTCCTGCACAGCTGGGAACCACGATCTACAAGGAATTTGAAAAGAGGAAGATGCTAT
 25 GAAAGAAAAAATTCCTAAACTGTGGAATGGATCACGTAGACATGTAACCCAGCAGCAGTTTGCTT
 CTGTTGTCCACTGATTAATCAGCCTATGTGCCTGACACTGGTCTTGAGTACAAGTGAAGCCAAAACAA
 GGTGGAAGATGCTCTGAATTAAGATGTTTTACCACATTGTATTACAGAGACAGCCAATAAATCTACTAT
 TTGATTTCAA

As used herein, the term “MPV17” refers to a gene encoding mitochondrial inner
 30 membrane protein MPV17, or the corresponding protein product. The terms “MPV17” and
 “mitochondrial inner membrane protein MPV17” include wild-type forms of the MPV17 gene
 or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion
 constructs, among others) of wild-type MPV17 proteins and nucleic acids encoding the same.
 Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%,
 35 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%,
 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or
 more) to any of the amino acid sequences of a wild-type MPV17 protein (e.g., SEQ ID NO:
 13), provided that the MPV17 variant retains the function of a wild-type MPV17. Additionally,

the terms “MPV17” and “mitochondrial inner membrane protein MPV17” may refer to fusion proteins, or nucleic acids encoding the same, in which the MPV17 is operably linked to another polypeptide, half-life-modifying agent, or therapeutic agent. The term “MPV17” may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary MPV17 amino acid sequence (SEQ ID NO: 13):

MALWRAYQRALAAHPWKVQVLTAGSLMGLGDIISQQLVERRGLQEHQGRGLTMVSLGCGFVGPVVGWY
KVLDRFIPGTTKVDALKKMLLDQGGFAPCFLGCFPLVLVGALNGLSAQDNWAKLQRDYPDALITNYYLWPA
VQLANFYLVPLHYRLAVVQCVAVIWNSYLSWKAHRL

Exemplary MPV17 nucleic acid sequence (SEQ ID NO: 14):

AGTTCCTAGGCCAGCCTGTACGTGGGAGGGAGGCTCGGCGCTCAGGAAGCATGGCACTCTGGCGGGCAT
ACCAGCGGGCCCTGGCCGCTCACCCGTGGAAAGTACAGGTCCTGACAGCTGGGTCCCTGATGGGCTGGG
TGACATTATCTACAGCAGCTGGTGGAGAGGCGGGGTCTGCAGGAACACCAGAGAGGCCGGACTCTGACC
ATGGTGTCCCTGGGCTGTGGCTTTGTGGGCCCTGTGGTAGGAGGCTGGTACAAGGTTTTGGATCGGTTCA
TCCCTGGCACCACCAAAGTGGATGCACTGAAGAAGATGTTGTTGGATCAGGGGGGCTTTGCCCCGTGTTT
TCTAGGCTGCTTTCTCCCACTGGTAGGGGCACCTTAATGGACTGTGAGCCAGGACAACCTGGGCCAACTA
CAGCGGGATTATCCTGATGCCCTTATCACCACCTACTATCTATGGCCTGCTGTGCAGTTAGCCAACTTCT
ACCTGGTCCCCCTTCATTACAGGTTGGCCGTTGTCCAATGTGTTGCTGTTATCTGGAACCTCTACCTGTC
CTGGAAGGCACATCGGCTCTAAGCCTGCCTCACTCCATCGTTTCCACCTTGCACTGATGCAGCTTGACCC
TGGAACGGTCAGACAACCTCCTCAAAGTGGGCATACCAGTTTCCACGGGGTTGGGTGGCCGGTCAGAGCT
TAAGAGGACTAGCACCCCTGCAATGCCCTCTTCACTCTAAATGTACACTGACTGCTTTAGAGCCCTTGA
TAATAGTCTTATTTCCACCACATACTAGGCACTCCATAAATATCTGTTGAACCTTCATGACCTTATCAAC
TTTACACCCATATCCCAGCAAATGCCACTCATCCCCACTCTTCATAGACACATTTGTTACTCTAACCCCTG
CCTAGGCTTCTGTAGCTCCAGCTCTTTAGAGACTCCCGGAACCCCTTTATATGGTGCCTCAGTAAATATG
TTATTAAATATGTAATCCGGAA

As used herein, the term “POLG” refers to a gene encoding DNA polymerase gamma, catalytic subunit, or the corresponding protein product. The terms “POLG” and “DNA polymerase gamma, catalytic subunit” include wild-type forms of the POLG gene or protein, as well as variants (e.g., splice variants, truncations, concatemers, and fusion constructs, among others) of wild-type POLG proteins and nucleic acids encoding the same. Examples of such variants are proteins having at least 70% sequence identity (e.g., 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91 %, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.9% identity, or more) to any of the amino acid sequences of a wild-type POLG protein (e.g., SEQ ID NO: 15), provided that the POLG variant retains the function of a wild-type POLG. Additionally, the terms “POLG” and “DNA polymerase gamma, catalytic subunit” may refer to fusion proteins, or nucleic acids encoding the same, in which the POLG is operably linked to another polypeptide, half-life-

modifying agent, or therapeutic agent. The term “POLG” may refer to the protein or the gene encoding this protein, depending upon the context, as will be appreciated by one of skill in the art.

Exemplary POLG amino acid sequence (SEQ ID NO: 15):

5 MSRLLRKVAGATVGGPVPAPGRWVSSVPSADPSDGQRRRQQQQQQQQQQQPQQPQVLSSEGGQLR
HNPLDIQMLSRGLHEQIFGQGGEMPGEAAVRRSVEHLQKHGLWGQPAVPLPDVELRLPPLYGDNLDQHFR
LLAQKQSLPYLEAANLLLQAQLPKPPAWAWAEGWTRYGPEGEAVPVAIPEERALVFDVEVCLAEGTCPT
LAVAISSPSAWYSWCSQRLVEERYSWTSQSPADLIPLEVPTGASSPTQRDWQEQLVVGHNVSFDRAHIRE
QYLIQGSRMRFDLTMSMHMAISGLSSFQSRSLWIAAQKGKHKVQPPTKQGQKSQRKARRGPAISSWDWLDI
10 SSVNSLAEVHRLYVGGPPELEKEPRELFVKGTMKDIRENFQDLMQYCAQDVWATHEVFQQQLPLFLERCPH
PVTLAGMLEMGVSYLPVNQNWERYLAEAQGTYEELQREMKKSLMDLANDACQLLSGERYKEDPWLWDLEW
DLQEFKQKKAKVKKEPATASKLPIEGAGAPGDPMDQEDLGPCSEEEEFQQQVDMARACLQKLKGTTELLP
KRPQHLPGHPGWYRKLCPRLLDDPAWTPGPSLLSLQMRVTPKLMALTWDGFPLHYSERHGWGYLVPGRDN
LAKLPTGTTLESAGVVCYPYRAIESLYRKHCLEQGKQQLMPQEAGLAEEFLTDNSAIWQTVEELDYLEVE
15 AEAKMENLRAAVPGQPLALTARGGPKDTQPSYHHGNGPYNDVDIPGCWFFKLPHKDGNSCNVGSPPFAKDF
LPKMEDGTLQAGPGGASGPRALEINKMISFWRNAHKRISSQM VVWLPRSALPRAVIRHPDYDEEGLYGAI
LPQVVTAGTITRRAVEPTWLTASNARPDVRGSELKAMVQAPPGYTLVGADVDSQELWIAAVLGDAHFAGM
HGCTAFGWMTLQGRKSRGTDLHSKTATTVGISREHAKIFNYGRIYGAGQPFAERLLMQFNHRLTQQEAAE
KAQQMYAATKGLRWYRLSDEGEWLVRLENLVPDRTEGGWISLQDLRKVQRETARKSQWKKWEVVAERAWK
20 GGTESEMFNKLESIAATSDIPRTPVLGCCISRLEPSAVQEEFMTSRVNWVQSSAVDYLHMLVAMKWLF
EEFAIDGRFCISIHDEVRYLVREEDRYRAALALQITNLLTRCMFAYKLGLNDLPQSVAFFSAVDIDRCLR
KEVTMDCKTPSNPTGMERRYGIPQGEALDIYQIIELTKGSLEKRSQPGP

Exemplary POLG nucleic acid sequence (SEQ ID NO: 16):

GCGGACCGGCGGGGTGGAGGCCACACGCTACCCCGAGGCTGCGTAGGCCGCGCGAAGGGGGACGCCGTGC
25 CGTGGGCCTGGGGTCGGGGGAGCAGCAGACCGGGAAGCACCGATTGGGGTGAAGGCAGGCATGGTCAA
ACCCATTTCACTGACAGGAGAGCAGAGACAGGACGTGTCTCTCTCCACGTCTTCCAGCCAGTAAAGAAG
CCAAGCTGGAGCCCAAAGCCAGGTGTCTGACTCCAGCGTGGGGGTCCCTGCACCAACCATGAGCCGCC
TGCTCTGGAGGAAGGTGGCCGCGCCACCGTCGGGCCAGGGCCGGTTCCAGCTCCGGGGCGCTGGGTCTC
CAGCTCCGTCCCGCGTCCGACCCAGCGACGGGCAGCGCGGCGGCAGCAGCAGCAGCAGCAGCAGCAG
30 CAGCAGCAACAGCAGCCTCAGCAGCCGAAGTGCTATCCTCGGAGGGCGGGCAGCTGCGGCACAACCCAT
TGGACATCCAGATGCTCTCGAGAGGGCTGCACGAGCAAATCTTCGGGCAAGGAGGGGAGATGCTGGCGA
GGCCGCGGTGCGCCGAGCGTCGAGCACCTGCAGAAGCACGGGCTCTGGGGGCAGCCAGCCGTGCCCTTG
CCCAGCTGGAGCTGCGCCTGCCGCCCTCTACGGGGACAACCTGGACCAGCACTCCGCCTCTGGCCC
AGAAGCAGAGCCTGCCCTACCTGGAGGCGGCCAACTTGCTGTTGCAGGGCCAGCTGCCCCGAAGCCCCC
35 GGCTTGGGCCTGGGCGGAGGGCTGGACCCGGTACGGCCCCGAGGGGGAGGCCGTACCCGTGGCCATCCCC
GAGGAGCGGGCCCTGGTGTTCGACGTGGAGGTCTGCTTGGCAGAGGGAACCTGCCCCACATTGGCGGTGG
CCATATCCCCCTCGGCCTGGTATTCCTGGTGCAGCCAGCGGCTGGTGAAGAGCGTTACTCTTGGACCAG
CCAGCTGTGCGCGGCTGACCTCATCCCCCTGGAGGTCCCTACTGGTGCCAGCAGCCCCACCCAGAGAGAC
TGGCAGGAGCAGTTAGTGGTGGGGCACAATGTTTCTTTGACCGAGCTCATATCAGGGAGCAGTACCTGA
40 TCCAGGGTTCCCGCATGCGTTTCTTGACACCATGAGCATGCACATGGCCATCTCAGGGCTAAGCAGCTT

CCAGCGCAGTCTGTGGATAGCAGCCAAGCAGGGCAAACACAAGGTCCAGCCCCCACAAGCAAGGCCAG
AAGTCCCAGAGGAAAGCCAGAAGAGGGCCAGCGATCTCATCTGGGACTGGCTGGACATCAGCAGTGTCA
ACAGTCTGGCAGAGGTGCACAGACTTTATGTAGGGGGCCCTCCCTTAGAGAAGGAGCCTCGAGAACTGTT
TGTGAAGGGCACCATGAAGGACATTCGTGAGAACTTCCAGGACCTGATGCAGTACTGTGCCCAGGACGTG
5 TGGGCCACCCATGAGGTTTTCCAGCAGCAGCTACCGCTCTTCTTGGAGAGGTGTCCCCACCCAGTGA
TGGCCGGCATGCTGGAGATGGGTGTCTCCTACCTGCCTGTCAACCAGAACTGGGAGCGTTACCTGGCAGA
GGCACAGGGCACTTATGAGGAGCTCCAGCGGGAGATGAAGAAGTCGTTGATGGATCTGGCCAATGATGCC
TGCCAGCTGCTCTCAGGAGAGAGGTACAAAGAAGACCCCTGGCTCTGGGACCTGGAGTGGGACCTGCAAG
AATTTAAGCAGAAGAAAGCTAAGAAGGTGAAGAAGGAACAGCCACAGCCAGCAAGTTGCCCCATCGAGGG
10 GGCTGGGGCCCCCTGGTGATCCCATGGATCAGGAAGACCTCGGGCCCTGCAGTGAGGAGGAGGAGTTTCAA
CAAGATGTTCATGCCCCGCGCTGCTTGCAGAAGCTGAAGGGGACCACAGAGCTCCTGCCAAGCGGCCCC
AGCACCTTCTCGACACCCCTGGATGGTACCGGAAGCTCTGCCCCGGCTAGACGACCCCTGCATGGACCCC
GGGCCCCAGCCTCCTCAGCCTGCAGATGCGGGTCACACCTAAACTCATGGCACTTACCTGGGATGGCTTC
CCTCTGCACTACTCAGAGCGTCATGGCTGGGGCTACTTGGTGCTGGGCGCGGGACAACCTGGCCAAGC
15 TGCCGACAGGTACCAACCTGGAGTCAGCTGGGGTGGTCTGCCCCCTACAGAGCCATCGAGTCCCTGTACAG
GAAGCACTGTCTCGAACAGGGGAAGCAGCAGCTGATGCCCCAGGAGGCCGCTGGCGGAGGAGTTCTTG
CTCACTGACAATAGTGCCATATGGCAAACGCTAGAAGAACTGGATTACTTAGAAGTGGAGGCTGAGGCCA
AGATGGAGAACTTGCGAGCTGCAGTGCCAGGTCAACCCCTAGCTCTGACTGCCCCGTGGTGGCCCCAAGGA
CACCCAGCCCAGCTATCACCATGGCAATGGACCTTACAACGACGTGGACATCCCTGGCTGCTGGTTTTTC
20 AAGCTGCCTCACAAGGATGGTAATAGCTGTAATGTGGGAAGCCCCCTTTGCCAAGGACTTCTGCCAAGA
TGGAGGATGGCACCCCTGCAGGCTGGCCCAGGAGGTGCCAGTGGGCCCCGTGCTCTGGAATCAACAAAAT
GATTTCTTTCTGGAGGAACGCCCATAAACGTATCAGCTCCCAGATGGTGGTGTGGCTGCCCAGGTGAGCT
CTGCCCCGTGCTGTGATCAGGCACCCCGACTATGATGAGGAAGGCCTCTATGGGGCCATCCTGCCCCAAG
TGGTGA CTGCCGGCACCATCACTCGCCGGGCTGTGGAGCCACATGGCTCACCGCCAGCAATGCCCGGCC
25 TGACCGAGTAGGCAGTGAGTTGAAAGCCATGGTGCAGGCCCCACCTGGCTACACCCCTTGTGGGTGCTGAT
GTGGACTCCCAAGAGCTGTGGATTGCAGCTGTGCTTGGAGACGCCCACTTTGCCGGCATGCATGGCTGCA
CAGCCTTTGGGTGGATGACACTGCAGGGCAGGAAGAGCAGGGGCACTGATCTACACAGTAAGACAGCCAC
TACTGTGGGCATCAGCCGTGAGCATGCCAAAATCTTCAACTACGGCCGCATCTATGGTGCTGGGCAGCCC
TTTGCTGAGCGCTTACTAATGCAGTTTAAACCACCGGCTCACACAGCAGGAGGCAGCTGAGAAGGCCAGC
30 AGATGTACGCTGCCACCAAGGGCCTCCGCTGGTATCGGCTGTGCGATGAGGGCGAGTGGCTGGTGAGGGA
GTTGAACCTCCCACTGGACAGGACTGAGGGTGGCTGGATTTCCCTGCAGGATCTGCGCAAGGTCCAGAGA
GAAACTGCAAGGAAGTCACAGTGAAGAAGTGGGAGGTGGTTGCTGAACGGGCATGGAAGGGGGGCACAG
AGTCAGAAATGTTCAATAAGCTTGAGAGCATTGCTACGTCTGACATACCACGTACCCCGGTGCTGGGCTG
CTGCATCAGCCGAGCCCTGGAGCCCTCGGCTGTCCAGGAAGAGTTTATGACCAGCCGTGTGAATTGGGTG
35 GTACAGAGCTCTGCTGTTGACTACTTACACCTCATGCTTGTGGCCATGAAGTGGCTGTTTGAAGAGTTTG
CCATAGATGGGCGCTTCTGCATCAGCATCCATGACGAGGTTTCGCTACCTGGTGCGGGAGGAGGACCGCTA
CCGCGCTGCCCTGGCCTTGCAGATACCAACCTCTTGACCAGGTGCATGTTTGCCTACAAGCTGGGTCTG
AATGACTTGCCCCAGTCAGTCGCCTTTTTTCAGTGCAGTCGATATTGACCGGTGCCTCAGGAAGGAAGTGA
CCATGGATTGTA AAAACCCCTTCCAACCCAACTGGGATGGAAAGGAGATACGGGATTCCCCAGGTGAAGC
40 GCTGGATATTTACCAGATAATTGAACTCACCAAAGGCTCCTTGGAAAAACGAAGCCAGCCTGGACCATAG
CACTGCCTGGAGGCTCTGTATTTGCTCCCGTGGAGCTTCATCGGGGTGGTGCAGGCTCCCAAACCTCAGGC

5

10

Mitochondrial DNA Depletion Syndrome Treatment and Prevention

20

30

35

nucleosides to generate deoxythymidine monophosphate (TMP) and deoxycytidine monophosphate (dCMP), which in turn, are converted to deoxynucleotide triphosphates (dNTPs) required for mitochondrial DNA synthesis. Autosomal recessive *TK2* mutations cause devastating neuromuscular weakness with severe depletion of mitochondrial DNA (mtDNA) in infants and children, as well as progressive external ophthalmoplegia with mtDNA multiple deletions in adults. Many patients cannot walk and require some type of mechanical ventilation and feeding tube. The central nervous system is variably involved in these disorders, with symptoms that include seizures, encephalopathy, cognitive impairment, and hearing loss. Less than 7% of patients live more than 42 years.

Based on clinical and molecular genetics findings of patients thus diagnosed, three disease presentations were identified: i) infantile-onset (≤ 1 year-old) myopathy with onset of weakness in the first year of life with severe mtDNA depletion and early mortality; ii) childhood-onset (>1 -11 years-old) myopathy with severe mtDNA depletion; and iii) late-onset myopathy (≥ 12 years-old) with mild weakness at onset and slow progression to loss of ambulation, respiratory insufficiency, or both, often with chronic progressive external ophthalmoparesis in adolescence or adulthood in association with mtDNA multiple deletions, reduced mtDNA copy number, or both. See Garone, *et al.* 2018.

Attempts to study the pathogenesis and test therapies for TK2 deficiency using cultured fibroblasts from patients have been unsuccessful, because the replicating cells failed to manifest mtDNA depletion. In contrast, a homozygous Tk2 H126N knock-in mutant (*Tk2*^{-/-}) mouse model, manifests a phenotype that is strikingly similar to the human infantile encephalomyopathy caused by *TK2* mutations, characterized by onset at age 10 days with decreased ambulation, unstable gait, coarse tremor, growth retardation, and depletion of mitochondrial DNA (mtDNA) progressing rapidly to early death at age 14 to 16 days, which is a time period analogous to the human infantile-onset disease (Akman, *et al.* 2008; Dorado, *et al.* 2011). This mouse model has been used to show the administration of oral dC/dT prolonged delayed the onset of clinical symptoms of TK2 deficiency and prolonged the lives of the mice by two- to three-fold ('092 application). It was also used to show the administration of oral dCMP/TMP delayed clinical symptoms and prolonged life ('207 application).

Exemplary findings from AAV vectors encoding TK2 in mouse models of MDS

As shown herein using a mouse model of Tk2 deficiency, the disease can be cured by replacing the mutant gene with a normal gene that encodes a functional Tk2 protein. An exemplary gene therapy described herein involves the use of an AAV (e.g., an AAV2 or AAV9

virus) containing a transgene encoding TK2. It has been discovered that administration of an AAV9 virus expressing human TK2 (referred to as “AAV9-hTK2” in the Examples provided below) prolonged the lifespan of TK2 mutant mice up to a maximum lifespan of about 4 months and an average of about 3 months. This was more than double the lifespan of the oral dC/dT pharmacological-treated TK2 mutant mice, which was an average of 43 days, and five times longer than the untreated TK2 mutant mice which was 18 days (see, e.g., Example 2). Weight also increased from about 55% over mice with dC/dT treatment alone and about 80% over the weight of wild type mice (see, e.g., Example 3).

Muscle strength (normalized to weight) and motor function assessed by Rotarod test was performed at postnatal day 60 and showed no differences between the treated TK2 mutant mice and untreated wild-type mice (see, e.g., Examples 3 and 4). Wild-type mice showed no adverse side-effects related to the therapy and were overall healthy.

Activity and levels of OXPHOS enzyme complexes in brain of treated mutant mice were similar to that showed in untreated and treated wild-type mice (see, e.g., Examples 7 and 8).

Additionally, treated mutant mice never showed head tremor, characteristic of untreated and dC/dT treated *Tk2*^{-/-} mice, confirming that rescue of mitochondrial function in central nervous system prevents pathology.

Mitochondrial DNA was also rescued by the AAV9 administration in most tissue of the mutant mice (see, e.g., Example 6).

Treatment with 4.2×10^{11} vc of AAV9-hTK2 at P1 does not efficiently transduce kidney cells, as previously observed in other studies. Consequently, TK2 activity in treated mutant mice was very poor compared to wild-type mice and there were signs of mtDNA depletion as soon as post-natal day 29. Histology and biochemical analysis showed compromised kidney function, which may be the underlying condition leading to early death of *Tk2*^{-/-} mice (see, e.g., Example 8).

However, when the TK2 mutant mice were administered with a combination of the AAV9-hTK2 and an AAV2 virus expressing human *TK2* gene (AAV2-hTK2), they survived significantly longer despite having lower levels of total viral dose. The mutant mice treated with both constructs grew as much as the mice treated with only AAV9-hTK2 and had slightly higher levels of mtDNA than mice treated with only AAV9-hTK2. They had equal growth and strength and motor coordination as the mutant mice treated with AAV9-hTK2 only as well as a decrease of protein in their urine and lower levels of index BUN (see, e.g., Example 9).

Additionally, when the mutant mice were treated with both AAV9-hTK2 and AAV2-

hTK2 were further administered oral dC+dT in water, they survived even longer and have increased growth and mtDNA levels in the liver and kidney and decreased protein in their urine (see, e.g., Example 10).

Additionally, the TK2 mutant mice are treated with the oral dC/dT as described in the
5 '092 application or the oral dCMP/TMP as described in '207 application, starting at an early age, e.g., postnatal day 4, and subsequently administered the AAV9-hTK2 vector, alone or in combination with the AAV2-hTK2 vector, at a later age, ranging from one month to about two months, with about 21 days being ideal. The AAV2-hTK2 vector can be administered simultaneously with the AAV9-hTK2 vector or subsequently. These mutant mice survived
10 longer than the mutant mice administered only the AAV9-hTK2 and have increased growth and strength, and mtDNA and RCEs are restored in all tissues including kidney (see, e.g., Examples 11 and 12).

Methods of treating, preventing, and/or curing mitochondrial diseases in patients

15 Patients who would benefit from the administration of the described gene therapy include those diagnosed with a deficiency in one or more of TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG. In these patients, compositions containing a nucleic acid encoding one or more of the deficient proteins (e.g., viral vectors, such as AAV vectors, containing such nucleic acids) may be administered to the patient. These compositions may be
20 administered alone or in combination with pharmacological agents, such as deoxynucleosides (e.g., dC and dT or mixtures thereof), or deoxyribonucleoside monophosphates (e.g., dCMP and TMP or mixtures thereof). For example, the subject may have a mitochondrial DNA depletion syndrome and the treatment comprises administering (i) a composition containing a viral vector (e.g. AAV, such as an AAV that targets at least muscle tissue, optionally at least
25 muscle tissue and CNS, for example AAV9) containing a nucleic acid encoding one or more of the deficient proteins, and (ii) a composition comprising deoxynucleosides (e.g., dC and dT or mixtures thereof), or deoxyribonucleoside monophosphates (e.g., dCMP and TMP or mixtures thereof). In one embodiment, the mitochondrial DNA depletion syndrome is TK2 deficiency and the viral vector (e.g. AAV, such as an AAV that targets at least muscle tissue,
30 optionally at least muscle tissue and CNS, for example AAV9) comprises a nucleic acid encoding TK2. In a particular embodiment, the mitochondrial DNA depletion syndrome is TK2 deficiency and the viral vector is AAV9 comprising a nucleic acid encoding TK2.

In some embodiments, the present disclosure provides methods of treating, preventing, curing, and/or reducing the severity or extent of TK2 deficiency by administering to a subject

in need thereof a therapeutically effective amount of a composition, such as a viral vector (e.g., an AAV), comprising a nucleic acid encoding TK2. In some embodiments, the viral vector is an AAV, such as AAV9 or AAV2. In some embodiments, the composition (e.g., viral vector, such as an AAV) comprising a nucleic acid encoding TK2 is administered as soon as TK2
5 deficiency is diagnosed or suspected. In some embodiments, the amount of AAV comprising the transgene administered is about 4.2×10^{11} or 4.2×10^{10} genome or vector or vector copies.

The disclosure also provides methods of treating, preventing, curing, and/or reducing the severity or extent of TK2 deficiency by administering to a subject in need thereof a therapeutically effective amount of a first composition (e.g., viral vector, such as AAV)
10 containing a nucleic acid encoding TK2 and further comprising administering to the subject a therapeutically effective amount of a second composition (e.g., viral vector, such as AAV) containing a nucleic acid encoding TK2. In some embodiments, the first and second AAV are each independently an AAV2 or AAV9 vector encoding TK2. In some embodiments, the first AAV vector is administered prior to the second AAV vector. In some embodiments, the first
15 composition (e.g., AAV vector) is administered as soon as TK2 deficiency is diagnosed or suspected, and the second composition (e.g., AAV vector) is administered at a time point after the first composition. In some embodiments, the second composition (e.g., AAV vector) is administered within days of the first composition (e.g., AAV vector). In some embodiments, the second composition (e.g., AAV vector) is administered within weeks of the first
20 composition (e.g., AAV vector). In some embodiments, the second composition (e.g., AAV vector) is administered months after the first composition (e.g., AAV vector). In some embodiments, the first composition (e.g., AAV vector) and the second composition (e.g., AAV vector) are administered simultaneously at any given time point. In some embodiments, the two compositions (e.g., AAV vectors) are present within the same larger composition, and in
25 some embodiments, the two are separate compositions.

In some embodiments, the method further includes further administering a therapeutically effective amount of a pharmacological agent, such as dC and/or dT. The pharmacological agent may be administered at the same time that the first composition (e.g., viral vector, such as an AAV vector) is administered or at a time point after the first
30 composition is administered. In some embodiments, the pharmacological agent, such as dC and/or dT, is administered within days of the first composition. In some embodiments, the pharmacological agent, such as dC and/or dT, is administered within weeks of the first composition, and in some embodiments, the pharmacological agent, such as dC and/or dT, is administered months after the first composition. In some embodiments, the first composition,

second composition, and the pharmacological agent are administered simultaneously at any given time point.

In some embodiments, the first composition (e.g., viral vector, such as an AAV, encoding a transgene described herein, for example TK2) is administered as soon as TK2
5 deficiency is known or suspected. The pharmacological agent (e.g., dC and/or dT) may be administered starting from about 7 days to about 35 days after the administration of the first composition, starting from about 14 days to about 28 days after the administration of the first composition, starting from about 14 days to about 21 days after the administration of the first composition, or starting from about 21 days after the administration of the first composition.
10 The second composition (e.g., viral vector, such as an AAV, encoding a transgene described herein, for example TK2) may be administered starting from about 14 days to about 45 days after the administration of the first composition, starting from about 21 days to about 35 days after the administration of the first composition, starting from about 21 days to about 30 days after the administration of the first composition, or starting from about 29 days after the administration of the first composition. In some embodiments, administration of the pharmacological agent (e.g., dC and/or dT) continues throughout the subject's lifetime.

In some embodiments, the disclosure provides methods of treating, preventing, curing, and/or reducing the severity or extent of TK2 deficiency by administering to a subject in need thereof a therapeutically effective amount of a pharmacological agent (e.g., dC and/or dT) and
20 further includes administering to the subject a therapeutically effective amount of a composition containing a viral vector, such as an AAV vector (e.g. rAAV), containing a nucleic acid encoding TK2. For example, the composition may contain an AAV2 or AAV9 containing a nucleic acid encoding TK2. In certain embodiments, the AAV is AAV9. In some embodiments, the pharmacological agent is administered prior to the AAV vector. In some
25 embodiments, the pharmacological agent is administered as soon as TK2 deficiency is diagnosed or suspected, and the AAV vector is administered at a time point after. In some embodiments, the AAV vector is administered within days of the pharmacological agent, and in some embodiments, the AAV vector is administered within weeks of the pharmacological agent. In some embodiments, the AAV vector is administered months after the
30 pharmacological agent. In some embodiments, the pharmacological agent and the AAV vector are administered simultaneously at any given time point. In some embodiments, a second AAV vector (e.g., AAV2 or AAV9) encoding TK2 is also administered. The first and second AAV vector may be the same or different. For example, the first AAV vector may be an AAV2 vector (i.e., containing AAV2 ITRs and AAV2 capsid proteins) and the second AAV vector

may be an AAV9 vector (e.g., containing AAV9 ITRs and AAV9 capsid proteins). In some embodiments, one or both of the first and second vectors is a pseudotyped AAV vector containing AAV2 ITRs and capsid proteins from a different AAV serotype (e.g., an AAV2/1, AAV2/8, or AAV2/9 vector). In some embodiments, the first AAV vector is administered
5 simultaneously or later in time relative to administration of the second AAV vector. The first and second vectors may be administered in the same composition or in different compositions.

In some embodiments, the disclosure provides methods of treating, preventing and/or reducing the severity or extent of TK2 deficiency by administering to a subject in need thereof a therapeutically effective amount of a pharmacological agent containing dCMP and/or TMP
10 and additionally administering to the subject a therapeutically effective amount of a composition containing an AAV vector (e.g., an AAV2 or AAV9) containing a nucleic acid encoding TK2. In some embodiments, the dCMP and/or TMP is administered prior to administration of the AAV vector. In some embodiments, the dCMP and/or TMP is administered as soon as TK2 deficiency is diagnosed or suspected, and the AAV vector is
15 administered at a time point after. In some embodiments, the AAV vector is administered within days of dCMP and/or TMP administration, and in some embodiments, the AAV vector is administered within weeks of the dCMP and/or TMP. In some embodiments, the AAV vector is administered months after the dCMP and/or TMP. In some embodiments, the dCMP and/or TMP and the AAV vector are administered simultaneously at any time point. In some
20 embodiments, an additional AAV vector (e.g., an AAV2 or AAV9 vector) containing a nucleic acid encoding TK2 is also administered to the subject. The second AAV vector can be administered simultaneously or later in time relative to administration of the first AAV vector. The first and second AAV vectors may be administered as part of the same composition or as different composition.

25 All of the aforementioned methods can be used to restore function of a dysfunctional TK2 protein in a subject with TK2 deficiency.

A parallel defect of dGK, due to autosomal recessive mutations in *DGUOK* with deficiencies in dGMP and dAMP, causes mtDNA depletion typically manifesting as early childhood-onset hepatocerebral disease (Mandel, *et al.* 2001). These patients would benefit
30 from the administration of an AAV vector comprising the nucleic acid that encodes dGK and at least one deoxypurine, dG or dA, or mixtures thereof. Similarly, patients having deficiencies of TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG may be treated using the compositions and methods of the disclosure, for example, by providing such a patient with a therapeutically effective amount of a composition (e.g., a viral vector, such as an AAV vector)

containing a transgene encoding the deficient protein. The composition may be provided to the patient alone or in combination with a pharmacological agent, such as a deoxynucleoside or deoxynucleoside monophosphate described herein.

For example, other forms of MDS, as well as other disorders related to unbalanced
 5 nucleotide pools, can be treated by administration of compositions (e.g., viral vectors, such as AAV vectors) containing nucleic acids encoding one or more of TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG. The composition may be administered alone or in combination with a pharmacological agent, such as a deoxynucleosides (e.g., dA, dG, dC, or dT, or mixture thereof) or deoxynucleoside monophosphate (e.g., dAMP, dGMP, dCMP, or
 10 TMP or mixture thereof). These disorders include, without limitation, deficiencies related to *RRM2B* (encoding p53R2, the p53 inducible small subunit of ribonucleotide reductase, RNR) and mutations in *TYMP* (encoding thymidine phosphorylase, TP) which cause mitochondrial neurogastrointestinal encephalomyopathy (MNGIE). Additional nuclear genes that disrupt mitochondrial dNTP pools include but are not limited to *SUCLA2*, *SUCLG1* and *MPV17*.

15

Assessing the condition of a patient having a mitochondrial disease

Patients that exhibit the phenotype discussed above for TK2 deficiency including the most typical presentation of progressive muscle disease characterized by generalized hypotonia, proximal muscle weakness, loss of previously acquired motor skills, poor feeding,
 20 and respiratory difficulties, can be tested to definitively diagnose the disease.

If the clinical presentation is highly suspicious for mtDNA depletion syndrome, molecular genetic testing using a panel of genes known to cause mtDNA depletion syndrome should be performed (Chanprasert, *et al.* 2012). The *TK2* gene is the only gene in which mutations are known to cause TK2-related mitochondrial DNA depletion syndrome.
 25 This testing can include a sequence analysis of the entire coding and exon/intron junction regions of *TK2* for sequence variants and deletion/duplication. If compound heterozygous or homozygous deleterious mutations are identified in the sequence analysis, the diagnosis of TK2 deficiency is confirmed, and thus, the subject would benefit from the therapy described herein. If sequence analysis does not identify two compound heterozygous or homozygous deleterious
 30 mutations, deletion/duplication analysis should be considered to determine and/or confirm a TK2 deficiency diagnosis.

Further tests to determine and/or confirm a TK2 deficiency diagnosis may include testing serum creatine kinase (CK) concentration, electromyography, histopathology on skeletal muscle, mitochondrial DNA (mtDNA) content (copy number), and electron transport

chain (ETC) activity in skeletal muscle. If one or more of the following is found in these tests, the TK2 deficiency is determined and/or confirmed. Elevated CK concentration as compared to healthy controls can indicate TK2 deficiency. A skeletal muscle biopsy can be performed, and then a mtDNA content analysis in skeletal muscle performed. If the skeletal muscle biopsy shows prominent variance in fiber size, variable sarcoplasmic vacuoles, variable increased connective tissue, and ragged red fibers as well as increased succinate dehydrogenase (SDH) activity and low to absent cytochrome c oxidase (COX) activity, and mtDNA copy number is severely reduced (typically less than 20% of age- and tissue-matched healthy controls), a diagnosis of TK2 deficiency can be determined and/or confirmed (Chanprasert, *et al.* 2012).

Additionally, TK2 deficiency is inherited in an autosomal recessive manner. Thus, a sibling of an affected patient can be tested as early as possible after birth to diagnose the disease.

In all of these examples, gene therapy alone or in combination with deoxynucleoside therapy or deoxyribonucleoside monophosphate therapy should be started as soon as possible after a diagnosis of TK2 deficiency.

Recombinant AAV Vectors

"Recombinant AAV (rAAV) vectors" described herein generally include a transgene (*e.g.*, encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG). The transgene is flanked by 5' and 3' ITRs, and may be operably linked to one or more regulatory elements in a manner that permits transgene transcription, translation, and/or expression in a cell of a target tissue. Such regulatory elements may include a promoter or enhancer, such as the chicken beta actin promoter or cytomegalovirus enhancer, among others described herein. The recombinant AAV genome is generally encapsidated by capsid proteins (*e.g.*, from the same AAV serotype as that from which the ITRs are derived or from a different AAV serotype from that which the ITRs are derived). The AAV vector may then be delivered to a selected target cell. In some embodiments, the transgene is a nucleic acid sequence, heterologous to the vector sequences, which encodes a polypeptide, protein, functional RNA molecule (*e.g.*, miRNA, miRNA inhibitor) or other gene product of interest (*e.g.*, TK2). Components of exemplary AAV vectors that may be used in conjunction with the compositions and methods of the disclosure are described below.

Any AAV serotype or combination of AAV serotype can be used in the methods and compositions of the present invention. Because the methods and compositions of the present invention are for the treatment and cure of mitochondrial disorders, AAV serotypes that target

at least muscle, or at least muscle and the central nervous system can be used in some embodiments and include but are not limited to AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, and AAV9.

In some embodiments, AAV9 serotype, which has a wide tropism, is used.

5 *Components of AAV vectors*

The AAV vectors described herein may contain cis-acting 5' and 3' ITRs (See, *e.g.*, Carter, in "Handbook of Parvoviruses", ed., P. Tijsser, CRC Press, pp. 155 168 (1990)). The ITR sequences are typically about 145 bp in length. Preferably, substantially the entire sequences encoding the ITRs are used in the molecule, although some degree of minor
10 modification of these sequences is permissible. (See, *e.g.*, texts such as Sambrook *et al.*, (1989) and Fisher *et al.*, (1996)). An example of such a molecule is a "cis-acting" plasmid containing the transgene, in which the selected transgene sequence and associated regulatory elements are flanked by the 5' and 3' AAV ITR sequences. The AAV ITR sequences may be obtained from any known AAV, including presently identified mammalian AAV types.

15 In addition to the elements identified above for recombinant AAV vectors, the vector may also include conventional control elements which are operably linked to the transgene in a manner which permits its transcription, translation and/or expression in a cell transfected with the plasmid vector or infected with the virus produced by the invention. As used herein, "operably linked" sequences include both expression control sequences that are contiguous
20 with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (*i.e.*, Kozak consensus sequence);
25 sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. A great number of expression control sequences, including promoters which are native, constitutive, inducible and/or tissue-specific, are known in the art and may be utilized.

As used herein, a nucleic acid sequence (*e.g.*, coding sequence) and regulatory
30 sequences are said to be operably linked when they are covalently linked in such a way as to place the expression or transcription of the nucleic acid sequence under the influence or control of the regulatory sequences. If it is desired that the nucleic acid sequences be translated into a functional protein, two DNA sequences are said to be operably linked if induction of a promoter in the 5' regulatory sequences results in the transcription of the coding sequence and if the

nature of the linkage between the two DNA sequences does not (1) result in the introduction of a frame-shift mutation, (2) interfere with the ability of the promoter region to direct the transcription of the coding sequences, or (3) interfere with the ability of the corresponding RNA transcript to be translated into a protein. Thus, a promoter region would be operably
5 linked to a nucleic acid sequence if the promoter region were capable of effecting transcription of that DNA sequence such that the resulting transcript might be translated into the desired protein or polypeptide. Similarly two or more coding regions are operably linked when they are linked in such a way that their transcription from a common promoter results in the expression of two or more proteins having been translated in frame. In some embodiments,
10 operably linked coding sequences yield a fusion protein. In some embodiments, operably linked coding sequences yield a functional RNA (*e.g.*, shRNA, miRNA).

For nucleic acids encoding proteins, a polyadenylation sequence generally is inserted following the transgene sequences and before the 3' AAV ITR sequence. An rAAV construct useful in the present invention may also contain an intron, desirably located between the
15 promoter/enhancer sequence and the transgene. One possible intron sequence is derived from SV-40, and is referred to as the SV-40 T intron sequence.

Another vector element that may be used is an internal ribosome entry site (IRES). An IRES sequence is used to produce more than one polypeptide from a single gene transcript. An IRES sequence would be used to produce a protein that contain more than one polypeptide
20 chains. Selection of these and other common vector elements are conventional and many such sequences are available (see, *e.g.*, Sambrook *et al*, and references cited). Such a motif may be useful, for example, for instances in which multiple genes or portions thereof are expressed from the same AAV vector.

The precise nature of the regulatory sequences needed for gene expression in host cells
25 may vary between species, tissues or cell types, but shall in general include, as necessary, 5' non-transcribed and 5' non-translated sequences involved with the initiation of transcription and translation respectively, such as a TATA box, capping sequence, CAAT sequence, enhancer elements, and the like. Especially, such 5' non-transcribed regulatory sequences will include a promoter region that includes a promoter sequence for transcriptional control of the
30 operably joined gene. Regulatory sequences may also include enhancer sequences or upstream activator sequences as desired. The vectors may optionally include 5' leader or signal sequences.

Examples of constitutive promoters include, without limitation, a chicken beta actin promoter, a retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with a RSV

enhancer), a cytomegalovirus (CMV) promoter (optionally with a CMV enhancer), a SV40 promoter, a dihydrofolate reductase promoter, a 13-actin promoter, a phosphoglycerol kinase (PGK) promoter, and an EFla promoter (Invitrogen).

Inducible promoters allow regulation of gene expression and can be regulated by
5 exogenously supplied compounds, environmental factors such as temperature, or the presence of a specific physiological state, *e.g.*, acute phase, a particular differentiation state of the cell, or in replicating cells only. Inducible promoters and inducible systems are available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech and Ariad. Examples of inducible promoters regulated by exogenously supplied promoters include a zinc-
10 inducible sheep metallothionine (MT) promoter, a dexamethasone (Dex)-inducible mouse mammary tumor virus (MMTV) promoter, a T7 polymerase promoter system (WO 98/10088); a ecdysone insect promoter (No *et al.*, *Proc. Natl. Acad. Sci. USA* 93:3346-3351 (1996)), a tetracycline-repressible system (Gossen *et al.*, *Proc. Natl. Acad. Sci. USA* 89:5547-5551 (1992)), a tetracycline-inducible system (Gossen *et al.*, *Science* 268:1766-1769 (1995), a
15 RU486-inducible system (Wang *et al.*, *Nat. Biotech.* 15:239-243 (1997) and Wang *et al.*, *Gene Ther.* 4:432-441 (1997)) and a rapamycin-inducible system (Magari *et al.*, *J. Clin. Invest.* 100:2865-2872 (1997)). Still other types of inducible promoters which may be useful in this context are those which are regulated by a specific physiological state, *e.g.*, temperature, acute phase, a particular differentiation state of the cell, or in replicating cells only.

20 In another embodiment, a native promoter, or fragment thereof, for the transgene will be used. The native promoter may be preferred when it is desired that expression of the transgene should mimic the native expression. The native promoter may be used when expression of the transgene must be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific transcriptional stimuli. In a further embodiment,
25 other native expression control elements, such as enhancer elements, polyadenylation sites or Kozak consensus sequences may also be used to mimic the native expression.

In some embodiments, the regulatory sequences impart tissue-specific gene expression capabilities. In some cases, the tissue-specific regulatory sequences bind tissue-specific transcription factors that induce transcription in a tissue specific manner.

30 In some embodiments, one or more bindings sites for one or more of miRNAs are incorporated in a transgene of a rAAV vector, to inhibit the expression of the transgene in one or more tissues of a subject harboring the transgenes. The miRNA target sites in the mRNA may be in the 5' UTR, the 3' UTR or in the coding region. Typically, the target site is in the 3' UTR of the mRNA. Furthermore, the transgene may be designed such that multiple miRNAs

regulate the mRNA by recognizing the same or multiple sites. The presence of multiple miRNA binding sites may result in the cooperative action of multiple RISCs and provide highly efficient inhibition of expression. The target site sequence may comprise a total of 5-100, 10-60, or more nucleotides. The target site sequence may comprise at least 5 nucleotides of the sequence of a target gene binding site.

For example, a 3'UTR site which would inhibit the expression of the transgene in the liver can be incorporated into a transgene. This would be beneficial for transgenes which encode therapeutic proteins which are toxic to the liver as most of the virus administered (approximately 60 to 90%) is eventually found in the liver. Thus suppressing the therapeutic gene expression in liver relieves the burden from liver cells.

In some embodiments, the AAV vector will be modified to be a self-complementing AAV. A self-complementing AAV carries complementary sequence of the transgene (i.e., a double copy of the transgene). Self complementation makes the gene more stable after it enters the cell.

15

Transgene Coding Sequences

Nucleic acid sequences of transgenes described herein may be designed based on the knowledge of the specific composition (e.g., viral vector) that will express the transgene. For example, one type of transgene sequence includes a reporter sequence, which upon expression produces a detectable signal. In another example, the transgene encodes a therapeutic protein or therapeutic functional RNA. In another example, the transgene encodes a protein or functional RNA that is intended to be used for research purposes, e.g., to create a somatic transgenic animal model harboring the transgene, e.g., to study the function of the transgene product. In another example, the transgene encodes a protein or functional RNA that is intended to be used to create an animal model of disease. Appropriate transgene coding sequences will be apparent to the skilled artisan.

In embodiments of the current invention the transgenes would encode a functional protein including but not limited to TK2, dGK, p53R2, TP, SUCLA2, SUCLG1, POLG1, and a mitochondrial membrane protein MPV17.

The human *TK2* gene (Gene ID: 7084) can be used to obtain a transgene encoding a functional TK2 protein. The human *DGUOK* (Gene ID: 1716) can be used to obtain a transgene encoding a functional dGK protein. The human *TYMP* gene (Gene ID: 1890) can be used to obtain a transgene encoding a functional TP protein. The human RRM2B gene (Gene ID: 50584) can be used to obtain a transgene encoding a functional p53R2 protein. The human

SUCLA2 gene (Gene ID: 8803) can be used to obtain a transgene encoding a functional *SUCLA2* protein. The human *SUCLG1* gene (Gene ID: 8802) can be used to obtain a transgene encoding a functional *SUCLG1* protein. The human *POLG* gene (Gene ID: 5428) can be used to obtain a transgene encoding a functional *POLG1* protein. The human *MPV17* gene (Gene ID: 4538) can be used to obtain a transgene encoding a functional mitochondrial membrane protein MPV17.

Codon optimization of transgene coding sequences

Codon optimization of the transgene coding sequences can increase the efficiency of the gene therapy. Thus, in some embodiments, a nucleic acid that is at least 70% identical to the coding sequence of the transgene encoding the therapeutic protein (e.g., a nucleic acid sequence that is 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical to the nucleic acid sequence) is used.

Codon optimization tools are known in the art.

Exemplary codon optimized nucleic acids are as follows.

Exemplary codon optimized TK2 nucleic acid (SEQ ID NO: 17):

ATGGGAGCATTGTCAGCGACCTAGTTCCGATAAAGAACAGGAGAAAGAGAAGAAGAGCGTCATCTG
TGTGGAAGGTAAACATCGCTAGTGGCAAACGACCTGTTTGGAATTTTCTCTAATGCCACCGATGTGG
AGGTGCTTACAGAGCCTGTTAGTAAATGGCGAAATGTTTCGAGGACATAATCCGCTGGGACTGATGTAT
CACGATGCCTCCAGATGGGGACTGACCCTTCAGACTTATGTCCAACCTACAATGCTTGACAGGCACAC
TAGACCGCAGGTTAGCAGCGTGAGGCTGATGGAGCGCTCAATTCCTCTGCCAGGTATATATTCGTTG
AGAACCTCTACCGATCTGGTAAGATGCCTGAGGTAGATTACGTAGTGCTGAGTGAATGGTTTGACTGG
ATTCTTCGCAACATGGATGTATCAGTGGATCTCATTGTTTACCTTAGGACTAACCCGGAGACGTGTTA
TCAGCGCTTGAAAAAGCGCTGCCGGAAGAGGAAAAAGTGATTCTCTGGAATACTTGGAAGCTATTC
ATCACCTTCACGAAGAGTGGCTGATAAAAGGTTCCCTGTTTCCCATGGCGGCCCCCGTGCTTGTCATA
GAAGCGGATCACCACATGGAAAGGATGCTCGAACTTTTCGAGCAAAATCGAGACCGGATTCTCACGCC
GGAGAATCGAAAACATTGCCCATAG

Exemplary codon optimized dGK nucleic acid (SEQ ID NO: 18):

ATGGCTGCAGGGAGGCTTTTTTGTCTAGATTGAGGGCACCTTTTAGCTCCATGGCGAAGTCTCCGCT
TGAGGGGGGTGAGTAGCAGCCGGGGGCTCCATGCTGGCAGAGGGCCCCGGAGGCTGTCTATTGAGGGGA
ATATTGCAGTTGGTAAGAGCACGTTTGTAAATTGCTCACGAAGACCTATCCAGAATGGCACGTTGCT
ACAGAACCAGTAGCTACATGGCAAAACATTCAAGCAGCAGGCACACAGAAGGCTTGCACTGCACAGTC
TCTCGGAAACCTGCTCGATATGATGTACAGAGAGCCCGCCGGTGGTCATACAGTTTCAGACCTTCA
GCTTTCTTTCTCGGTTGAAGGTACAGTTGGAGCCCTTCCAGAGAAGTTGCTTCAAGCACGGAAACCT

GTACAGATATTCGAACGATTGCATTTTCGAGGCGCTGATGAACATACCTGTGCTTGTTCCTGGATGTGAA
CGACGATTTTTCCGAGGAAGTTACTAAGCAAGAGGACCTCATGCGGGAGGTAAACACGTTTCGTTAAGA
ACCTGTAA

Exemplary codon optimized TP nucleic acid (SEQ ID NO: 19):

5 ATGGCAGCATTGATGACACCAGGAACGGGGCTCCGCCTGCGCCTGGCGATTTTTTCAGGAGAGGGGAG
TCAAGGTCTGCCGGACCCTTCACCGGAACCCAAACAGCTTCCCGAGCTTATTAGAATGAAGCGAGATG
GGGGTCGCCTCTCTGAGGCTGACATACGCGGCTTTGTTGCGGCGGTTCGTCAACGGCTCAGCACAAGGT
GCGCAAATAGGAGCGATGCTGATGGCTATCAGGCTGAGGGGAATGGATTTGGAAGAGACCAGTGTGCT
GACTCAAGCGCTGGCACAGAGCGGCCAGCAGCTTGAATGGCCAGAAGCTTGGCGCCAACAATTGGTTCG
10 AATAAAGCTCAACCGGGGGCGTCGGGGACAAAGTCTCCCTTGTACTGGCACCTGCTCTGGCTGCGTGC
GGATGTAAGGTTCCCTATGATTAGCGGGCGGGGCTTGGGACATACGGGAGGAACGTTGGACAAACTCGA
ATCCATCCCTGGTTTCAACGTGATTACAGAGCCCTGAACAAATGCAAGTACTCCTGGATCAGGCAGGCT
GCTGTATTGTTGGCCAAAGCGAACAACCTCGTGCCTGCGGATGGTATCCTCTATGCTGCCAGGGATGTG
ACCGCCACAGTTGACTCCCTCCCGCTGATAACAGCGTCAATTTTGTAGTAAGAAGCTCGTCGAAGGGCT
15 CAGCGCTCTTGTGGTTGATGTAAAATTTGGGGGGCGCAGTATTCCCCAACAGGAACAAGCACGGG
AACTGGCAAAAACCTGGTTGGTGTGGGCGCTTCACTGGGCTGAGAGTTGCCGCTGCCTTGACCGCC
ATGGATAAGCCGCTCGGTAGGTGCGTAGGTGCTGCACTTGAAGTAGAAGAAGCCCTTTTGTGCATGGA
TGGGGCCGGTCCCCCTGACTTGCAGGATCTCGTAACCACGCTTGGCGGCGCGTTGCTTTGGCTTTCTG
GACACGCTGGTACCCAAGCGCAAGGGGCGAGCAAGAGTCGCAGCAGCGCTTGATGACGGATCAGCTCTT
20 GGGCGATTTGAAAGAATGCTTGCAGCTCAGGGCGTAGATCCAGGGCTGGCGCGGGCCCTTTGCTCAGG
TTCTCCAGCAGAGCGCCGACAGCTCCTTCCAGGGCGCGAGAGCAGGAAGAATTGCTGGCCCCGGCTG
ATGGTACCGTCGAACCTCGTACGGGCTTTGCCGCTGGCTCTTGTGTTTGCACGAACCTGGGGGCTGGGCGG
AGTCGCGCCGGTGAACCTCTCAGACTCGGTGTGGGTGCGGAGCTCCTCGTTGACGTCGGGCAACGACT
TCGAGGGGAACCCCTTGGCTTAGGGTACACAGGGACGGGCCAGCACTCAGCGGCCCTCAGTCCAGGG
25 CCGTTCAAGAAGCTCTCGTGCTGAGTGATCGAGCGCCTTTTGCCGCTCCCTCACCATTGCTGAATTG
GTATTGCCACCCAGCAGTAA

Exemplary codon optimized p53R2 nucleic acid (SEQ ID NO: 20):

ATGCTGCTTCTCAGACTGCCACCCACCGCAGTCATGCTTCCCCACTTGACTGTAACTTCA
AGATAGGTGCCGGAAGTGTTACTACCCCGATCAGGACAAGCGTGCCCTCCTGCGTTGGCGG
30 CCGCCTGGCTTCGGCGGTGTGAACGCCGAGGAGTTCGGCCGAGGGGTGGGCGACGCAAGGAA
CTGACCCTTGGCCTGCGCCCTGCACGGTGTTCAGCTCCCGTCTGCCAAGGATGATGCATG
GAGACCTCAGGCAGGTGCTCCTCCTCAGATACCAACGAGAGTGAAATTAAATCTAATGAAG
AACCCCTTCTTCGCAAGAGTAGTCGCAGGTTCGTGATATTTCCCATACAATATCCTGATATC
TGGAAAATGTACAAGCAAGCCCAAGCATCATTCTGGACCGCAGAAGAGGTTGATCTCAGTAA
35 GGACCTCCACACTGGAACAACTGAAAGCCGACGAAAAATATTTTCATATCCCATATACTTG
CTTTCTTTGCAGCCTCAGATGGAATTGTGAATGAAACTTGGTAGAGCGCTTCTCTCAAGAG

GTCCAAGTTCCTGAAGCACGATGCTTTTACGGATTTCAAATCCTGATAGAAAATGTACATTC
 CGAAATGTATTCTTGCTTATAGACACGTATATCAGAGATCCCCAAAAAAGGGAGTTCTTGT
 TCAATGCGATCGAGACCATGCCATATGTTAAGAAAAAAGCCGATTGGGCCCTGAGGTGGATA
 GCTGATAGGAAGTCTACATTTGGGGAACGCGTTGTTGCCCTCGCCGCCGTCGAGGGAGTCTT
 5 CTTCTCCGGATCCTTCGCTGCTATCTTTTGGCTGAAAAAACGCGGCTTGATGCCTGGTTTGA
 CCTTTAGCAATGAGTTGATATCACGGGATGAAGGGCTGCATTGTGACTTTGCCCTGCCTTATG
 TTCCAGTACTTGGTGAACAAACCGTCTGAGGAACGGGTAAAGGGAAATCATAGTGGATGCAGT
 AAAAATAGAGCAGGAGTTCCTTACTGAAGCGCTCCCCGTAGGGCTCATTGGCATGAACTGCA
 TTCTTATGAAGCAGTACATCGAGTTCGTGGCCGATAGACTTTTGGTGGAGCTCGGGTTCTCA
 10 AAGGTTTTTTCAGGCTGAAAAATCCTTTTGATTTTATGGAGAACATATCTCTGGAGGGCAAGAC
 TAACTTTTTTGA AAAAAGAGTATCCGAGTATCAAAGGTTTGCCGTGATGGCAGAAACAACAG
 ATAATGTCTTCACACTTGATGCGGATTTTTAA

Exemplary codon optimized SUCLA2 nucleic acid (SEQ ID NO: 21):

ATGGCGGCTTCTATGTTCTATGGACGCTTGGTCGCCGTTGCCACGTTGCGGAATCACAGGCCCCGAAC
 15 TGCACAAAGGGCTGCCGCTCAGGTGCTTGAAGCAGCGGATTGTTCAACAATCACGGCCTCCAAGTTC
 AGCAACAGCAGCAGAGAAACCTCTCTCTGCATGAGTATATGAGTATGGAACCTGCTCCAGGAAGCAGGG
 GTGTCTGTACCAAAGGGCTACGTAGCAAAAAGCCCCGACGAGGCGTACGCCATAGCTAAAAAACTGGG
 AAGTAAGGACGTAGTTATCAAAGCACAGGTTCTGGCCGGTGGCCGCGGTAAGGGAACCTTCGAGAGTG
 GTTTGAAAGGCGGTGTCAAGATCGTATTTAGTCCCGAGGAGGCCAAGGCTGTAAGTTCCCAGATGATC
 20 GGCAAAAAATTGTTCAAAAGCAAACCGCGCAAAAGGGTAGAATTTGCAACCAAGTACTTGTCTGCGA
 AAGAAAGTATCCGAGAAGGGAGTATTATTTGCTATAACCATGGAAAGATCATTCCAAGGTCCCGTGC
 TCATCGGCAGTTCACACGGGGGCGTTAATATAGAGGATGTCGCTGCAGAGTCTCCGAGAGCCATAATT
 AAGGAGCCTATAGACATCGAGGAAGGCATTAAAAAAGAACAAGCCTTGCAACTGGCACAAAAAATGGG
 TTTTCTCCGAACATCGTCGAGAGTGCCGCAGAAAATATGGTGAACTGTACAGCCTGTTTTTGAAGT
 25 ACGATGCGACAATGATAGAAATTAATCCGATGGTCGAGGACTCAGATGGAGCCGTGCTTTGTATGGAC
 GCCAAAATTAACCTTCGATTCAAACAGCGCTTATCGACAGAAGAAAATTTTCGATTTGCAAGATTGGAC
 CCAAGAGGATGAGCGAGACAAAGATGCCGCCAAAGCAAATCTCAATTATATAGGACTCGATGGTAATA
 TTGGATGTTTGGTCAATGGTGCCGGTCTCGCGATGGCAACTATGGATATAATCAAGTTGCATGGTGGG
 ACTCCCGCCAACTTTCTTGACGTCGGGGGAGGCGCCACCGTTACCAAGTTACGGAGGCTTTCAAAC
 30 CATAACCTCTGACAAGAAGGTGCTGGCGATTCTTGTAATATCTTTGGTGGCATTATGCGGTGTGATG
 TTATTGCCAGGGAATAGTAATGGCTGTCAAAGATTTGGAGATTAAATACCCGTCGTGGTGC GGCTC
 CAAGGTACTCGAGTAGATGATGCTAAGGCTCTGATTGCCGACAGCGGGCTGAAGATTCTGGCATGTGA
 TGATTTGGACGAGGCAGCGAGAATGGTCTGAAGCTCTCAGAAATAGTAACTTTGGCGAAACAGGCTC
 ATGTCGATGTAAAGTTCCAACCTTCCTATATGA

35 Exemplary codon optimized SUCLG1 nucleic acid (SEQ ID NO: 22):

ATGACCGCTACGCTTGC GGCTGCAGCCGACATCGCGACAATGGTCTCTGGAAGTTCAGGCTTGGCCGC
 GGCTCGGCTTCTGAGTAGAAGTTTCTTGTGTCGCGAAAACGGTATCAGGCACTGCTCCTACACCGCCA
 GTAGACAACACCTTTATGTTGATAAGAACACGAAGATAATTTGCCAAGGATTACCCGGTAAACAGGGG
 ACCTTCCATTCCCAGCAGGCACTGGAATATGGAATAAATTGGTAGGCGGCACTACGCCTGGGAAGGG
 5 TGGCCAAACACATCTTGGTCTTCCCGTTTTTAACACAGTGAAGGAGGCTAAAGAACAAACGGGGGCAA
 CGGCTAGCGTTATCTATGTCCCACCCCGTTTGGCGCTGCAGCCATAAATGAGGCGATTGAAGCCGAG
 ATCCCGCTTGTAGTCTGCATAACGGAGGGAATTCCGCAACAAGACATGGTGCAGATAAAGCACAAGCT
 TCTTCGACAGGAAAAACAAGACTGATAGGTCCGAATTGTCTTGGCGTAATTAACCCCGGTGAATGCA
 AAATCGGAATTATGCCGGGACATATTCACAAAAAAGGCCGAATCGGGATAGTCAGCAGATCAGGCACC
 10 TTGACATACGAAGCGGTTACCAGACTACGCAAGTCGGTTTGGGACAAAGCCTTTGCGTCGGTATCGG
 TGGCGACCCATTTAACGGGACGGATTTCAATTGACTGTCTGGAGATCTTCCTCAACGATTCCGCGACAG
 AAGGAATCATTTTGTATAGGAGAAATAGGCGGGAACGCCGAAGAGAATGCAGCGGAATTCTCAAACAA
 CATAACTCTGGCCCTAATAGTAAGCCGGTGGTATCATTATAGCCGGTCTTACAGCGCCGCCGGGTGCG
 CAGAATGGGCCACGCAGGTGCAATTATAGCGGGGGGCAAGGGAGGTGCCAAAGAAAAGATATCCGCTC
 15 TTCAAAGCGCAGGTGTAGTAGTTAGTATGAGTCCAGCTCAACTGGGCACAACCATCTACAAAGAATTT
 GAAAAGCGGAAAATGCTTTGA

Exemplary codon optimized MPV17 nucleic acid (SEQ ID NO: 23):

ATGGCCCTCTGGAGAGCTTACCAACGAGCCTTGGCCGCGCACCCCTTGAAGGTACAGGTCTTGACCGC
 CGGATCTTTGATGGGTCTTGGAGATATTATTTCTCAACAGTTGGTTGAACGGCGAGGCCTCCAGGAAC
 20 ACCAAAGGGGGAGAACCCCTTACAATGGTAAGTCTGGGTTGCGGATTCGTGCGGCCTGTCTGGGGGGA
 TGGTATAAGGTTCTGGATCGCTTTATACCGGGGACCACCAAGGTCGATGCCTTGAAGAAGATGTTGCT
 GGATCAGGGAGGATTGCTCCGTGTTTTTTGGGATGTTTTCTGCCACTTGTTGGGGCTCTGAACGGAC
 TGTCCGCGCAGGATAACTGGGCGAAGTTGCAACGCGACTACCCAGACGCCCTGATAACAAATTACTAT
 CTCTGGCCAGCAGTTCAGTTGGCCAATTTTTACCTGGTACCTCTCCACTATCGCCTTGCTGTAGTACA
 25 GTGTGTCGCCGTCATCTGGAACCTACCTTTTCATGGAAGGCTCATAGATTGTAA

Exemplary codon optimized POLG nucleic acid (SEQ ID NO: 24):

ATGTCCAGGCTGCTCTGGCGGAAGGTCGAGGCGCCACTGTCTGGTCCAGGACCAGTTCCAGCTCCCGG
 TCGCTGGGTGAGCAGCAGCGTGCCAGCTAGCGATCCGAGCGACGGTCAGCGGCGACGGCAACAACAGC
 AGCAACAGCAACAACAGCAACAACAACAGCCCCAGCAGCCTCAGGTTCTCAGTTCCGAGGGTGGCCAA
 30 CTCCGACACAACCCACTGGATATACAGATGCTCTCCGCGGTCTCCACGAACAAATATTTCGACAAAG
 GGGTGAGATGCCGGGAGAGGCTGCGGTGAGGCGCAGTGTAGAATCTCCAGAAACACGGGTTGTGGG
 GCCAACCGGCCGTTCTCTCCCCGATGTTGAACTGCGGCTTCCACCTCTCTACGGTGATAATCTGGAC
 CAGCACTTTAGACTGCTCGCTCAAAAGCAGAGTCTCCCTTACCTGGAAGCCGCTAACCTCCTGCTCCA
 AGCCCAATTGCCCCCTAAACCGCCAGCCTGGGCTTGGGCGGAGGGATGGACGAGGTATGGACCCGAAG
 35 GGGAGGCTGTGCCAGTTGCTATACCAGAGGAACGCGCTCTGGTTTTTCGACGTAGAGGTTTGTCTCGCG
 GAGGGAACTTGTCTTACACTGGCTGTAGCAATTTCCCTTCAGCCTGGTACAGCTGGTGTCTCAGAG
 ATTGGTGGAAGAAAGGTATAGCTGGACTAGCCAGCTGAGTCCCGCGGACCTCATTCCACTTGAGGTAC

CCACCGGGGCGTCAAGCCCAACTCAGAGGGACTGGCAGGAACAATTGGTAGTTGGGCATAATGTGAGT
TTTGACAGGGGCTCATATCCGCGAACAGTATCTTATCCAGGGCTCTAGAATGCGATTCCCTTGACACGAT
GAGCATGCACATGGCAATCAGCGGACTTAGTTCCCTTTCAGAGGTCATTGTGGATTGCAGCCAAGCAGG
GAAAGCATAAGGTCCAACCCCGACAAAACAAGGTCAGAAATCCCAGAGAAAAGCCCGGCGAGGCCCC
5 GCCATCAGTTCCCTGGGATTGGTTGGATATCAGTAGTGTGAATAGCCTTGCTGAGGTGCATCGCCTGTA
TGTGGGTGGACCCCCACTTGAGAAAGAGCCTAGGGAGCTCTTCGTCAAAGGCACCATGAAGGATATTA
GAGAGAATTTTCAAGATCTCATGCAATACTGCGCACAAGACGTATGGGCAACGCATGAGGTCTTTCAA
CAGCAACTCCCCCTCTTTTTTGAACGATGTCCACATCCCGTCACTCTTGCTGGGATGTTGGAAATGGG
TGTAAGTTATTTGCCAGTCAATCAAAATTGGGAGAGATACTTGGCTGAAGCGCAGGGTACATATGAGG
10 AACTTCAGCGAGAAATGAAAAAAGTCTTATGGATTGGCCAATGACGCCTGCCAGCTGCTTTCCGGT
GAGCGGTACAAAGAAGATCCATGGCTTTGGGATCTCGAATGGGATTTGCAGGAATTTAAACAGAAAAA
GGCCAAGAAGGTCAAGAAAGAGCCAGCTACAGCCTCAAAGCTCCCTATAGAGGGAGCGGGAGCACCGG
GTGATCCGATGGATCAAGAGGATTTGGGACCTTGCTCCGAAGAGGAGGAATTCCAACAAGATGTAATG
GCAAGGGCCTGCCTGCAAAAGCTCAAAGGGACAACAGAACTCTTGCCCAAGAGGCCTCAACATCTGCC
15 CGGCCATCCAGGTTGGTATCGCAAACCTCTGTCCAAGGCTGGACGATCCCGCCTGGACCCCGGGGCCCT
CCCTTCTGAGTCTGCAGATGAGAGTGACACCTAAGCTGATGGCACTTACTTGGGATGGGTTCCCTCTT
CACTATTCAGAGAGACACGGGTGGGGATATCTTGTCCCAGGTGCAAGGGACAATCTGGCGAAGCTTCC
CACAGGAACTACCTTGAGAGTGCGGGCGTAGTATGTCCTTATCGAGCCATAGAAAGTCTGTATAGAA
AGCATTGCCTTGAACAAGGCAAACAACAACCTCATGCCTCAGGAAGCCGGCCTCGCTGAAGAATTTCTT
20 CTTACTGATAACTCTGCTATCTGGCAAACAGTGGAGGAACTGGATTACCTCGAAGTCGAGGCCGAAGC
GAAGATGGAAAAATTTGCGGGCGGCGGTCCCAGGTCAGCCCCTTGCTTTGACAGCACGGGGGGGCCCTA
AGGATACCCAGCCCAGCTATCATCACGGGAACGGTCCATATAATGATGTTGATATACCTGGTTGTTGG
TTTTTTAAACTTCCTCATAAAGATGGCAATTCATGTAACGTTGGATCCCCATTTCGCAAAGGACTTCCT
CCCCAAAATGGAGGATGGGACGTTGCAAGCAGGTCCCGGTGGAGCTTCTGGGCCGCGAGCCCTGGAAA
25 TAAATAAGATGATTAGCTTCTGGAGGAATGCACACAAGCGCATTTCTTCACAGATGGTAGTGTGGCTG
CCTCGGAGTGCTTTGCCAGGGCTGTGATCAGACACCCGATTATGATGAAGAGGGACTGTACGGGGC
AATATTGCCCCAGGTGGTTACGGCTGGTACTATTACCCGCCGGGCAGTTGAGCCGACCTGGCTGACAG
CATCTAATGCCAGGCCTGATCGCGTGGGTTCTGAACTTAAAGCAATGGTCCAAGCTCCGCCTGGATAC
ACGCTTGTGCGCGCGGACGTGGACTCCCAGGAACCTTGGATAGCGGCTGTCCTTGGCGATGCACATTT
30 TGCAGGGATGCACGGGTGCACGGCTTTTGGCTGGATGACACTTCAGGGGAGGAAATCAAGGGGGACCG
ACCTGCACTCCAAGACCGCGACAACGGTAGGAATCTCAAGGGAACACGCTAAAATTTTCAATTATGGG
AGAATATATGGTGCCGGTCAACCGTTCGCTGAGCGCCTTCTGATGCAGTTTAACCATAGGCTGACACA
GCAGGAGGCAGCGGAGAAGGCGCAGCAGATGTACGCCGCAACTAAGGGTCTCAGATGGTATCGCCTCT
CAGACGAGGGCGAGTGGCTTGTCCGGGAATTGAACCTCCCGGTCGATCGGACGGAAGGTGGTTGGATT
35 AGTCTTCAGGATCTCAGAAAGGTGCAGCGGGAAACAGCACGCAAGTCTCAGTGAAGAAATGGGAGGT
GGTCGCCGAGCGGGCATGGAAGGGTGAACAGAATCCGAGATGTTTAACAAGCTGGAGAGCATAGCAA
CTAGTGACATCCCTAGGACACCGGTCTTGGGATGTTGCATTAGCCGGGCACTCGAGCCATCTGCCGTA

CAGGAAGAATTCATGACGTCACGGGTCAACTGGGTTGTCCAGTCTTCAGCCGTCGATTATCTGCATTT
 GATGCTGGTCGCGATGAAGTGGCTCTTTCGAAGAGTTTGCTATAGATGGCCGATTTTGCATCTCTATTC
 ACGATGAGGTGCGCTACCTGGTAAGGGAAGAAGACCGATACCGAGCCGCCCTCGCCCTTCAGATAACA
 AATCTGCTTACCAGGTGTATGTTTGCATACAAGCTGGGGTTGAACGACCTTCCCCAGTCCGTCGCTTT
 5 CTTTTCAGCTGTTGATATAGATCGCTGCCTGAGAAAAGAGGTTACGATGGACTGCAAAACGCCCTCAA
 ACCCCACTGGTATGGAGCGCAGATATGGCATCCCCCAAGGAGAAGCCCTCGACATATACCAGATAATT
 GAGCTCACGAAGGGCAGCCTGGAGAAGAGATCCCAACCTGGCCCATAG

Exemplary Recombinant AAV Compositions

10 The current disclosure provides for compositions containing a recombinant AAV
 containing a nucleic acid sequence that encodes a functional protein. Such proteins include,
 without limitation, TK2, dGK, p53R2, TP, SUCLA2, SUCLG1, POLG1, and a mitochondrial
 membrane protein MPV17. In some embodiments, the rAAV comprises a chicken beta-actin
 promoter. In some embodiments, the rAAV further comprises a CMV enhancer. In some
 15 embodiments, the rAAV further comprises ITRs. In some embodiments, the rAAV further
 comprises a 3'UTR. In some embodiments, the rAAV is AAV 2 or AAV9. In some
 embodiments, the transgene encodes a functional TK2 enzyme.

An exemplary rAAV-hTK2 vector comprising the human TK2 transgene, a chicken
 beta-actin promoter, a CMV IE enhancer, a BGH 3'UTR, a poly A signal, and a 5' and 3' ITR
 20 is shown in FIG. 1.

Both an rAAV9-hTK2 and an rAAV2-hTK2 vector were used in the Examples. These
 constructs have the same ITR but different capsid protein sequences.

In some embodiments, the composition further comprises a pharmaceutical carrier.

In some embodiments, the composition comprises more than one rAAV comprising a
 25 transgene that encodes a functional protein including but not limited to TK2, dGK, p53R2, TP,
 SUCLA2, SUCLG1, POLG1, and a mitochondrial membrane protein MPV17.

Routes of Administration and Dosing

30 The current invention provides rAAV vectors for use in methods of treating, preventing,
 and/or curing a disease or disorder characterized by unbalanced nucleotide pools, and/or
 restoring function of a protein that is dysfunctional in a disease or disorder characterized by
 unbalanced nucleotide pools, and/or alleviating in a subject at least one of the symptoms
 associated with unbalanced nucleotide pools. In some embodiments, methods involve
 administration of a rAAV vector that encodes one or more therapeutic peptides, polypeptides,

shRNAs, microRNAs, or antisense nucleotides, in a pharmaceutically-acceptable carrier to the subject in an amount and for a period of time sufficient to treat, prevent and/or cure the disease or disorder characterized by unbalanced nucleotide pools in the subject having or suspected of having such a disorder.

5 Diseases or disorders characterized by unbalanced nucleotide pools that can be treated, prevented and/or cured by the method of the current invention include, but are not limited to, those characterized by mutations in the following genes: *TK2*; *DGUOK*; *TYMP*; *RRM2B*; *SUCLA2*; *SUCLG1*; and *MPV17*. The proteins encoded by these genes can be restored by the method of the invention and include but are not limited to TK2, dGK, p53R2, TP, SUCLA2,
10 SUCLG1, POLG1, and a mitochondrial membrane protein MPV17.

The rAAVs may be delivered to a subject in compositions according to any appropriate methods known in the art. The rAAV, preferably suspended in a physiologically compatible carrier (*e.g.*, in a composition), may be administered to a subject. In certain embodiments, compositions may comprise a rAAV alone, or in combination with one or more other viruses
15 (*e.g.*, a second rAAV encoding having one or more different transgenes). In one embodiment, a composition can comprise an rAAV9 vector comprising a nucleic acid sequence comprising a transgene encoding a functional protein including but not limited to TK2, dGK, p53R2, TP, SUCLA2, SUCLG1, POLG1, and a mitochondrial membrane protein MPV17.

In a further embodiment, a composition can comprise an rAAV2 vector comprising a
20 nucleic acid sequence comprising a transgene encoding a functional protein including but not limited to TK2, dGK, p53R2, TP, SUCLA2, SUCLG1, POLG1, and a mitochondrial membrane protein MPV17.

In yet a further embodiment, a composition can comprise an rAAV9 vector comprising a nucleic acid sequence comprising a transgene encoding a functional TK2. In a further
25 embodiment, a composition can comprise an rAAV2 vector comprising a nucleic acid sequence comprising a transgene encoding a functional TK2. In a further embodiment, a composition can comprise an rAAV9 vector comprising a nucleic acid sequence comprising a transgene encoding TK2 and an rAAV2 vector comprising a nucleic acid sequence comprising a transgene encoding TK2.

30 Suitable carriers may be readily selected by one of skill in the art in view of the indication for which the rAAV is directed. For example, one suitable carrier includes saline, which may be formulated with a variety of buffering solutions (*e.g.*, phosphate buffered saline). Other exemplary carriers include sterile saline, lactose, sucrose, calcium phosphate, gelatin,

dextran, agar, pectin, peanut oil, sesame oil, and water. The selection of the carrier is not a limitation of the present invention.

Optionally, the compositions of the invention may contain, in addition to the rAAV and carrier(s), other conventional pharmaceutical ingredients, such as preservatives, or chemical stabilizers. Suitable exemplary preservatives include chlorobutanol, potassium sorbate, sorbic acid, sulfur dioxide, propyl gallate, the parabens, ethyl vanillin, glycerin, phenol, and parachlorophenol. Suitable chemical stabilizers include gelatin and albumin.

In some embodiments, rAAV compositions are formulated to reduce aggregation of AAV particles in the composition, particularly where high rAAV concentrations are present (*e.g.*, about 10^{13} GC/ml or more). Methods for reducing aggregation of rAAVs are well known in the art and, include, for example, addition of surfactants, pH adjustment, and salt concentration adjustment (see, *e.g.*, Wright, *et al.*, *Molecular Therapy* 12:171-178 (2005)).

Formulation of pharmaceutically-acceptable excipients and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens. Typically, these formulations may contain at least about 0.1% of the active ingredient or more, although the percentage of the active ingredient(s) may, of course, be varied and may conveniently be between about 1 or 2% and about 70% or 80% or more of the weight or volume of the total formulation. Naturally, the amount of active ingredient in each therapeutically-useful composition may be prepared in such a way that a suitable dosage will be obtained in any given unit dose of the compound. Factors such as solubility, bioavailability, biological half-life, route of administration, product shelf life, as well as other pharmacological considerations will be contemplated by one skilled in the art of preparing such pharmaceutical formulations, and as such, a variety of dosages and treatment regimens may be desirable.

The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms. In many cases the form is sterile and fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (*e.g.*, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils.

Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

For administration of an injectable aqueous solution, for example, the solution may be suitably buffered, if necessary, and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion. Some variation in dosage will necessarily occur depending on the condition of the host. The person responsible for administration will, in any event, determine the appropriate dose for the individual host.

Sterile injectable solutions are prepared by incorporating the active rAAV in the required amount in the appropriate solvent with various of the other ingredients enumerated herein, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

In addition to the methods of delivery described above, the following techniques are also contemplated as alternative methods of delivering the rAAV compositions to a host. Sonophoresis (*i.e.*, ultrasound) has been used and described in U.S. Patent No. 5,656,016 as a device for enhancing the rate and efficacy of drug permeation into and through the circulatory system. Other drug delivery alternatives contemplated are intraosseous injection (U.S. Patent No. 5,779,708), microchip devices (U.S. Patent No. 5,797,898), ophthalmic formulations, transdermal matrices (U.S. Patent Nos. 5,770,219 and 5,783,208) and feedback-controlled delivery (U.S. Patent No. 5,697,899).

rAAVS are administered in sufficient amounts to transfect the cells of a desired tissue and to provide sufficient levels of gene transfer and expression without undue adverse effects. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the selected tissue (*e.g.*, intracerebral administration, intrathecal administration), intravenous, oral, inhalation (including intranasal and intratracheal delivery), intraocular, intravenous, intramuscular, subcutaneous, intradermal, and other parental routes of administration. Routes of administration may be combined, if desired. The administration regimen depends on several factors, including the serum or tissue turnover rate of the therapeutic composition, the level of symptoms, and the accessibility of the target cells in the biological matrix. Preferably, the administration regimen delivers sufficient therapeutic composition to effect improvement in the target disease state, while simultaneously minimizing undesired side effects. Accordingly, the amount of biologic delivered depends in part on the particular therapeutic composition and the severity of the condition being treated.

The present invention provides stable pharmaceutical compositions comprising rAAV virions. The compositions remain stable and active even when subjected to freeze/thaw cycling and when stored in containers made of various materials, including glass.

Appropriate doses will depend on the subject being treated (*e.g.*, human or nonhuman primate or other mammal), age and general condition of the subject to be treated, the severity of the condition being treated, the mode of administration of the rAAV virions, among other factors. An appropriate effective amount can be readily determined by one of skill in the art.

The dose of rAAV virions required to achieve a desired effect or "therapeutic effect," *e.g.*, the units of dose in vector genomes/per kilogram of body weight (vg/kg), will vary based on several factors including, but not limited to: the route of rAAV administration; the level of gene or RNA expression required to achieve a therapeutic effect; the specific disease or disorder being treated; and the stability of the gene or RNA product. One of skill in the art can readily determine a rAAV virion dose range to treat a subject having a particular disease or disorder based on the aforementioned factors, as well as other factors that are well known in the art. An effective amount of the rAAV is generally in the range of from about 10 μ l to about 100 ml of solution containing from about 10^9 to 10^{16} genome copies per subject. Other volumes of solution may be used. The volume used will typically depend, among other things, on the size of the subject, the dose of the rAAV, and the route of administration. For example, for intrathecal or intracerebral administration a volume in range of 1 μ l to 10 μ l or 10 μ l to 100 μ l may be used. For intravenous administration a volume in range of 10 μ l to 100 μ l, 100 μ l to 1 ml, 1 ml to 10 ml, or more may be used. In some cases, a dosage between about 10^{10} to 10^{12}

rAAV genome copies per subject is appropriate. In certain embodiments, 10^{12} rAAV genome copies per subject is effective to target desired tissues. In some embodiments the rAAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , 10^{14} , or 10^{15} genome copies per subject. In some embodiments the rAAV is administered at a dose of 10^{10} , 10^{11} , 10^{12} , 10^{13} , or 10^{14} genome copies per kg.

Thus, a "therapeutically effective amount" will fall in a relatively broad range that can be determined through clinical trials. For example, for *in vivo* injection, *i.e.*, injection directly to the subject, a therapeutically effective dose will be on the order of from about 10^5 to 10^{16} of the rAAV virions, more preferably 10^8 to 10^{14} rAAV virions. For *in vitro* transduction, an effective amount of rAAV virions to be delivered to cells will be on the order of 10^5 to 10^{13} , preferably 10^8 to 10^{13} of the rAAV virions. If the composition comprises transduced cells to be delivered back to the subject, the amount of transduced cells in the pharmaceutical compositions will be from about 10^4 to 10^{10} cells, more preferably 10^5 to 10^8 cells. The dose, of course, depends on the efficiency of transduction, promoter strength, the stability of the message and the protein encoded thereby, etc. Effective dosages can be readily established by one of ordinary skill in the art through routine trials establishing dose response curves.

Dosage treatment may be a single dose schedule or a multiple dose schedule to ultimately deliver the amount specified above. Moreover, the subject may be administered as many doses as appropriate. Thus, the subject may be given, *e.g.*, 10^5 to 10^{16} rAAV virions in a single dose, or two, four, five, six or more doses that collectively result in delivery of, *e.g.*, 10^5 to 10^{16} rAAV virions. One of skill in the art can readily determine an appropriate number of doses to administer.

Pharmaceutical compositions will thus comprise sufficient genetic material to produce a therapeutically effective amount of the protein of interest, *i.e.*, an amount sufficient to reduce or ameliorate symptoms of the disease state in question or an amount sufficient to confer the desired benefit. Thus, rAAV virions will be present in the subject compositions in an amount sufficient to provide a therapeutic effect when given in one or more doses. The rAAV virions can be provided as lyophilized preparations and diluted in the virion-stabilizing compositions for immediate or future use. Alternatively, the rAAV virions may be provided immediately after production and stored for future use.

The pharmaceutical compositions will also contain a pharmaceutically acceptable excipient or carriers. Such excipients include any pharmaceutical agent that does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity. Pharmaceutically acceptable excipients

include, but are not limited to, liquids such as water, saline, glycerol and ethanol. Pharmaceutically acceptable salts can be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary
5 substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles. A thorough discussion of pharmaceutically acceptable excipients is available in *Remington's Pharmaceutical Sciences* and *U.S. Pharmacopeia: National Formulary*, Mack Publishing Company, Easton, PA (1984).

Formulations of therapeutic and diagnostic agents may be prepared by mixing with
10 acceptable carriers, excipients, or stabilizers in the form of, *e.g.*, lyophilized powders, slurries, aqueous solutions or suspensions.

Toxicity and therapeutic efficacy of the therapeutic compositions, administered alone or in combination with another agent, can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, *e.g.*, for determining the LD₅₀ (the dose
15 lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index (LD₅₀/ED₅₀). In particular aspects, therapeutic compositions exhibiting high therapeutic indices are desirable. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably
20 within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration.

Determination of the appropriate dose is made by the clinician, *e.g.*, using parameters or factors known or suspected in the art to affect treatment. Generally, the dose begins with an
25 amount somewhat less than the optimum dose and it is increased by small increments thereafter until the desired or optimum effect is achieved relative to any negative side effects. Important diagnostic measures include those of symptoms of, *e.g.*, the inflammation or level of inflammatory cytokines produced. In general, it is desirable that a biologic that will be used is derived from the same species as the animal targeted for treatment, thereby minimizing any
30 immune response to the reagent.

A preferred route of administration of the AAVs is intravenously.

A preferred dose ranges from about 1×10^{10} to about 8×10^{11} , from about 2×10^{10} to about 6×10^{11} , from about 4×10^{10} to about 4×10^{11} genome or viral copy (vc) total administration. A preferred dose is about 4×10^{11} genome or viral copy (vc) total administration of rAAV.

If more than one rAAV is used a preferred total dose of vector ranges from about 1×10^{10} to about 6×10^{11} , from about 2×10^{10} to about 5×10^{11} , from about 1×10^{10} to about 4×10^{11} genome or viral copy (vc) total administration. A preferred dose of total vector is about 3×10^{11} . The AAV can be administered in equal amounts, *e.g.*, ratio of 50/50, or in or in ratios of about 5/95,
5 10/90, 15/85, 20/80, 25/75, 30/70, 35/65, 40/60, 45/55, 55/45, 60/40, 65/35, 70/30, 75/25, 80/20, 85/15, 90/10, and 95/5.

Doses can be adjusted to optimize the effects in the subject. Additionally, a subject can be monitored for improvement of their condition prior to increasing the dosage. A subject's response to the therapeutic administration of the rAAV can be monitored by observing a
10 subject's muscle strength and control, and mobility as well as changes in height and weight. If one or more of these parameters increase after the administration, the treatment can be continued. If one or more of these parameters stays the same or decreases, the dosage can be increased.

15 Pharmacological Therapy - Methods of Administration and Dosing

The present invention encompasses the administration of deoxynucleosides, and/or deoxyribonucleoside monophosphates, in combination with the administration of a composition containing a transgene encoding a protein described herein.

Most preferred methods of administration are oral, intrathecal and parental including
20 intravenous. The deoxynucleosides or deoxyribonucleoside monophosphates must be in the appropriate form for administration of choice.

Deoxynucleosides or deoxyribonucleoside monophosphates are easily dissolved in liquid are easily dissolved in liquid (such as water, formula or milk) whereas the free acid form does not readily dissolve in liquid.

25 Such pharmaceutical compositions comprising one of more deoxynucleosides or deoxyribonucleoside monophosphates for administration may comprise a therapeutically effective amount of the deoxynucleosides or deoxyribonucleoside monophosphates and a pharmaceutically acceptable carrier. Such pharmaceutical carriers can be sterile liquids, such as saline solutions in water and oils, including those of petroleum, animal, vegetable, or
30 synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil, and the like. A saline solution is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel,

sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents.

Oral administration is a preferred method of administration. The deoxynucleosides or deoxyribonucleoside monophosphates can be added to any form of liquid a patient would consume including but not limited to, milk, both cow's and human breast, infant formula, and water.

Additionally, pharmaceutical compositions adapted for oral administration may be capsules, tablets, powders, granules, solutions, syrups, suspensions (in non-aqueous or aqueous liquids), or emulsions. Tablets or hard gelatin capsules may comprise lactose, starch or derivatives thereof, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, stearic acid or salts thereof. Soft gelatin capsules may comprise vegetable oils, waxes, fats, semi-solid, or liquid polyols. Solutions and syrups may comprise water, polyols, and sugars. An active agent intended for oral administration may be coated with or admixed with a material that delays disintegration and/or absorption of the active agent in the gastrointestinal tract. Thus, the sustained release may be achieved over many hours and if necessary, the active agent can be protected from degradation within the stomach. Pharmaceutical compositions for oral administration may be formulated to facilitate release of an active agent at a particular gastrointestinal location due to specific pH or enzymatic conditions.

In order to overcome any issue of the deoxynucleosides or deoxyribonucleoside monophosphates crossing the blood/brain barrier, intrathecal administration is an additional preferred form of administration (Galbiati, *et al.* 2006; Gotz, *et al.* 2008). Intrathecal administration involves injection of the drug into the spinal canal, more specifically the subarachnoid space such that it reaches the cerebrospinal fluid. This method is commonly used for spinal anesthesia, chemotherapy, and pain medication. Intrathecal administration can be performed by lumbar puncture (bolus injection) or by a port-catheter system (bolus or infusion). The catheter is most commonly inserted between the laminae of the lumbar vertebrae and the tip is threaded up the thecal space to the desired level (generally L3-L4). Intrathecal formulations most commonly use water, and saline as excipients but EDTA and lipids have been used as well.

A further preferred form of administration is parenteral including intravenous administration. Pharmaceutical compositions adapted for parenteral administration, including intravenous administration, include aqueous and non-aqueous sterile injectable solutions or suspensions, which may contain anti-oxidants, buffers, bacteriostats, and solutes that render

the compositions substantially isotonic with the blood of the subject. Other components which may be present in such compositions include water, alcohols, polyols, glycerine, and vegetable oils. Compositions adapted for parental administration may be presented in unit-dose or multi-dose containers, such as sealed ampules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of a sterile carrier, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules, and tablets. Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include: Water for Injection USP; aqueous vehicles such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Additionally, since some patients may be receiving enteral nutrition by the time the deoxynucleoside or deoxyribonucleoside monophosphates treatment begins, the dNs or dNMPs can be administered through a gastronomy feeding tube or other enteral nutrition means.

Further methods of administration include mucosal, such as nasal, sublingual, vaginal, buccal, or rectal; or transdermal administration to a subject.

Pharmaceutical compositions adapted for nasal and pulmonary administration may comprise solid carriers such as powders, which can be administered by rapid inhalation through the nose. Compositions for nasal administration may comprise liquid carriers, such as sprays or drops. Alternatively, inhalation directly through into the lungs may be accomplished by inhalation deeply or installation through a mouthpiece. These compositions may comprise aqueous or oil solutions of the active ingredient. Compositions for inhalation may be supplied in specially adapted devices including, but not limited to, pressurized aerosols, nebulizers or insufflators, which can be constructed so as to provide predetermined dosages of the active ingredient.

Pharmaceutical compositions adapted for rectal administration may be provided as suppositories or enemas. Pharmaceutical compositions adapted for vaginal administration may be provided as pessaries, tampons, creams, gels, pastes, foams or spray formulations.

Pharmaceutical compositions adapted for transdermal administration may be provided as discrete patches intended to remain in intimate contact with the epidermis of the recipient over a prolonged period of time.

The deoxynucleoside therapy comprises the administration of one or more deoxynucleosides chosen from the group consisting of deoxythymidine (dT), deoxycytidine (dC), deoxyadenosine (dA) and deoxyguanosine (dG).

A skilled practitioner can determine which deoxynucleosides are beneficial based upon the deficiency. It is also within the skill of the art for the practitioner to determine if mixtures of the deoxynucleosides should be administered and in what ratio. If two deoxynucleosides are to be administered, they can be in a ratio of 50/50 of each deoxynucleoside, *e.g.*, dC and dT, or in ratios of about 5/95, 10/90, 15/85, 20/80, 25/75, 30/70, 35/65, 40/60, 45/55, 55/45, 60/40, 65/35, 70/30, 75/25, 80/20, 85/15, 90/10, and 95/5.

By way of example, dT and dC are administered in mixture of equal amounts for TK2 deficiency.

The deoxyribonucleoside monophosphates therapy comprises the administration of one or more deoxyribonucleoside monophosphates chosen from the group consisting of TMP, dCMP, dAMP, and dGMP. If two deoxyribonucleoside monophosphates, ratios can be determined.

By way of example, TMP and dCMP are administered in mixture of equal amounts for TK2 deficiency.

Selection of a therapeutically effective dose will be determined by the skilled artisan considering several factors, which will be known to one of ordinary skill in the art. Such factors include the particular form of the deoxynucleoside or deoxyribonucleoside monophosphate, and its pharmacokinetic parameters such as bioavailability, metabolism, and half-life, which will have been established during the usual development procedures typically employed in obtaining regulatory approval for a pharmaceutical compound. Further factors in considering the dose include the condition or disease to be treated or the benefit to be achieved in a normal individual, the body mass of the patient, the route of administration, whether the administration is acute or chronic, concomitant medications, and other factors well known to affect the efficacy of administered pharmaceutical agents. Thus, the precise dose should be decided according to the judgment of the person of skill in the art, and each patient's circumstances, and according to standard clinical techniques.

A preferred dose ranges from about 100 mg/kg/day to about 1,000 mg/kg/day. A further preferred dose ranges from about 200 mg/kg/day to about 800 mg/kg/day. A further preferred dose ranges from about 250 mg/kg/day to about 400 mg/kg/day. These dosage amounts are of individual deoxynucleosides or deoxyribonucleoside monophosphates or of a composition with a mixture of more than one deoxynucleosides, *e.g.*, dT and dC or

deoxyribonucleoside monophosphates. For example, a dose can comprise 400 mg/kg/day of dT alone. In a further example, a dose can comprise a mixture of 200 mg/kg/day of dT and 200 mg/kg/day of dC. In a further example, a dose can comprise 400 mg/kg/day of a mixture of dT and dC.

5 Administration of the deoxynucleosides or deoxyribonucleoside monophosphates can be once a day, twice a day, three times a day, four times a day, five times a day, up to six times a day, preferably at regular intervals. For example, when the deoxynucleosides are administered four times daily, doses would be at 8:00 AM, 12:00 PM, 4:00 PM, and 8:00 PM.

Doses can also be lowered if being administered intravenously or intrathecally.
10 Preferred dose ranges for such administration are from about 50 mg/kg/day to about 500 mg/kg/day.

As shown in the '092 application, doses can be adjusted to optimize the effects in the subject. For example, the deoxynucleosides can be administered at 100 mg/kg/day to start, and then increased over time to 200 mg/kg/day, to 400 mg/kg/day, to 800 mg/kg/day, up to 1000
15 mg/kg/day, depending upon the subject's response and tolerability.

A subject can be monitored for improvement of their condition prior to increasing the dosage. A subject's response to the therapeutic administration of the deoxynucleosides or deoxyribonucleoside monophosphates can be monitored by observing a subject's muscle strength and control, and mobility as well as changes in height and weight. If one or more of
20 these parameters increase after the administration, the treatment can be continued. If one or more of these parameters stays the same or decreases, the dosage of the deoxynucleosides can be increased.

As shown in the prior patent applications, the pharmacological therapy is well tolerated. Any observed adverse effects were minor and were mostly diarrhea, abdominal bloating and
25 other gastrointestinal manifestations. A subject can also be monitored for any adverse effects, such as gastrointestinal intolerance, *e.g.*, diarrhea. If one or more adverse effects are observed after administration, then the dosage can be decreased. If no such adverse effects are observed, then the dosage can be increased. Additionally, once a dosage is decreased due to the observation of an adverse effect, and the adverse effect is no longer observed, the dosage can
30 be increased.

The deoxynucleosides or deoxyribonucleoside monophosphates can also be co-administered with other agents. Such agents would include therapeutic agents for treating the symptoms of the particular form of MDS. In particular, for TK2 deficiency, the dT and dC can be co-administered with an inhibitor of ubiquitous nucleoside catabolic enzymes, including but

not limited to enzyme inhibitors such as tetrahydrouridine (inhibitor of cytidine deaminase) and immucillin H (inhibitor of purine nucleoside phosphorylase) and tipiracil (inhibitor of thymidine phosphorylase). Such inhibitors are known and used in the treatment of some cancers.

5

Kits

The present invention also provides kits comprising the components of the combinations of the invention in kit form. A kit of the present invention includes one or more components including, but not limited to, viral vectors (e.g., AAV vectors) and/or
10 pharmacological agents (e.g., deoxynucleosides and/or nucleoside monophosphates) described herein. Kits may further include a pharmaceutically acceptable carrier, as discussed herein. The viral vector or pharmacological agent can be formulated as a pure composition or in combination with a pharmaceutically acceptable carrier, in a pharmaceutical composition.

In some embodiments, a kit includes an AAV vector containing a transgene described
15 herein in one container (e.g., in a sterile glass or plastic vial).

In some embodiments, a kit includes an AAV vector containing a transgene described herein in one container (e.g., in a sterile glass or plastic vial) and a second AAV vector encoding a transgene described herein in another container (e.g., in a sterile glass or plastic vial).

In some embodiments, a kit includes an AAV vector containing a transgene described
20 herein in one container (e.g., in a sterile glass or plastic vial) and a pharmacological agent in another container (e.g., in a sterile glass or plastic vial).

In some embodiments, a kit includes an AAV2 vector containing a transgene described herein in one container (e.g., in a sterile glass or plastic vial) and a pharmacological agent in another container (e.g., in a sterile glass or plastic vial).

25 In some embodiments, a kit includes an AAV9 vector containing a transgene described herein in one container (e.g., in a sterile glass or plastic vial) and a pharmacological agent in another container (e.g., in a sterile glass or plastic vial).

In some embodiments, a kit includes an AAV vector encoding TK2, dGK, p53R2, TP, SUCLA2, SUCLG1, POLG1, and/or a mitochondrial membrane protein MPV17, or a
30 pharmaceutical composition thereof in one container (e.g., in a sterile glass or plastic vial).

If the kit includes one or more pharmaceutical compositions for parenteral administration to a subject, the kit can include a device for performing such administration. For example, the kit can include one or more hypodermic needles or other injection devices as discussed above.

The kit can include a package insert including information concerning the pharmaceutical compositions and dosage forms in the kit. Generally, such information aids patients and physicians in using the enclosed pharmaceutical compositions and dosage forms effectively and safely. For example, the following information regarding a combination of the invention may be supplied in the insert: pharmacokinetics, pharmacodynamics, clinical studies, efficacy parameters, indications and usage, contraindications, warnings, precautions, adverse reactions, overdosage, proper dosage and administration, how supplied, proper storage conditions, references, manufacturer/distributor information and patent information.

10 Exemplary embodiments of the invention include:

1. A method of preventing or treating thymidine kinase 2 (TK2) deficiency in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein.
- 15 2. The method of embodiment 1, wherein the rAAV is chosen from the group consisting of AAV9 and AAV2.
3. The method of embodiment 1, wherein the subject is a mammal.
4. The method of embodiment 1, wherein the subject is a human.
5. The method of embodiment 1, wherein the composition is a pharmaceutical composition.
- 20 6. The method of embodiment 1, wherein the composition is the first composition comprising a first rAAV, wherein the first rAAV is AAV9 and further comprising administering a therapeutically effective amount of a second composition comprising a second rAAV comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein, wherein the second rAAV is AAV2 and is administered at a time point after the administration of the first rAAV.
- 25 7. The method of embodiment 6, wherein the second composition is a pharmaceutical composition.
8. The method of embodiment 6, wherein the first composition is administered as soon as the subject is diagnosed with TK2 deficiency or suspected of having TK2 deficiency, and is continued to be administered during the administration of the second composition.
- 30 9. The method of embodiment 6, wherein the second composition is administered within days of the first composition.

10. The method of embodiment 6, wherein the second composition is administered within weeks of the first composition.
11. The method of embodiment 6, wherein the second composition is administered at least a month after the first composition.
- 5 12. The method of embodiment 6, further comprising administering a therapeutically effective amount of a third composition comprising a pharmacological agent chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), and mixtures thereof.
13. The method of embodiment 12, wherein the third composition is administered at the same time point as the second composition.
- 10 14. The method of embodiment 12, wherein the third composition is administered at a later time point as the second composition.
15. The method of embodiment 12, wherein the therapeutically effective amount of the third composition is between about 100 mg/kg/day and about 1000 mg/kg/day.
16. The method of embodiment 12, wherein the therapeutically effective amount of the
15 third composition is between about 200 mg/kg/day and about 800 mg/kg/day.
17. The method of embodiment 12, wherein the therapeutically effective amount of the third composition is between about 250 mg/kg/day and about 400 mg/kg/day.
18. The method of embodiment 12, wherein the third composition is administered once daily, twice daily, three times daily, four times daily, five times daily or six times daily.
- 20 19. The method of embodiment 12, wherein the third composition is administered orally, intrathecally, enterally, or intravenously.
20. The method of embodiment 19, wherein the third composition is administered orally and comprises deoxynucleoside mixed with cow's milk, human breast milk, infant formula or water.
- 25 21. A method of preventing or treating thymidine kinase 2 (TK2) deficiency in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a first composition comprising a pharmacological agent, and further administering a therapeutically effective amount of a second composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding
30 a functional TK2 protein.
22. The method of embodiment 21, wherein the pharmacological agent is chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), and mixtures thereof; and deoxycytidine monophosphate (dCMP), deoxythymidine monophosphate (dTMP), and mixtures thereof.

23. The method of embodiment 21, wherein the rAAV is AAV9.
24. The method of embodiment 21, wherein the second composition is a pharmaceutical composition.
25. The method of embodiment 21, wherein the first composition is administered as soon as the subject is diagnosed with TK2 deficiency or suspected of having TK2 deficiency, and is continued to be administered during the administration of the second composition.
26. The method of embodiment 21, wherein the second composition is administered within days of the first composition.
27. The method of embodiment 21, wherein the second composition is administered within weeks of the first composition.
28. The method of embodiment 21, wherein the second composition is administered at least a month after the first composition.
29. The method of embodiment 21, wherein the therapeutically effective amount of the first composition is between about 100 mg/kg/day and about 1000 mg/kg/day.
30. The method of embodiment 21, wherein the therapeutically effective amount of the first composition is between about 200 mg/kg/day and about 800 mg/kg/day.
31. The method of embodiment 21, wherein the therapeutically effective amount of the first composition is between about 250 mg/kg/day and about 400 mg/kg/day.
32. The method of embodiment 21, wherein the first composition is administered once daily, twice daily, three times daily, four times daily, five times daily or six times daily.
33. The method of embodiment 21, wherein the first composition is administered orally, intrathecally, enterally, or intravenously.
34. The method of embodiment 33, wherein the first composition is administered orally and comprises the pharmacological agent mixed with cow's milk, human breast milk, infant formula or water.
35. The method of embodiment 21, further comprising the administration of a third composition comprising a second rAAV comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein wherein the second rAAV is AAV2.
36. A method of restoring thymidine kinase 2 enzyme activity in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein.

37. The method of embodiment 36, wherein the rAAV is chosen from the group consisting of AAV9 and AAV2.
38. The method of embodiment 36, wherein the subject is a mammal.
39. The method of embodiment 36, wherein the subject is a human.
- 5 40. The method of embodiment 36, wherein the composition is a pharmaceutical composition.
41. The method of embodiment 36, wherein the composition is the first composition comprising a first rAAV, wherein the first rAAV is AAV9 and further comprising administering a therapeutically effective amount of a second composition comprising a second
10 rAAV comprising
a nucleic acid sequence comprising a transgene encoding a functional TK2 protein, wherein the second rAAV is AAV2 and is administered at a time point after the administration of the first rAAV.
42. The method of embodiment 41, wherein the second composition is a pharmaceutical
15 composition.
43. The method of embodiment 41, wherein the first composition is administered as soon as the subject is diagnosed with TK2 deficiency or suspected of having TK2 deficiency, and is continued to be administered during the administration of the second composition.
44. The method of embodiment 41, wherein the second composition is administered within
20 days of the first composition.
45. The method of embodiment 41, wherein the second composition is administered within weeks of the first composition.
46. The method of embodiment 41, wherein the second composition is administered at least a month after the first composition.
- 25 47. The method of embodiment 41, further comprising administering a therapeutically effective amount of a third composition comprising a pharmacological agent chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), and mixtures thereof.
48. The method of embodiment 47, wherein the third composition is administered at the same time point as the second composition.
- 30 49. The method of embodiment 47, wherein the third composition is administered at a later time point as the second composition.
50. The method of embodiment 47, wherein the therapeutically effective amount of the third composition is between about 100 mg/kg/day and about 1000 mg/kg/day.

51. The method of embodiment 47, wherein the therapeutically effective amount of the third composition is between about 200 mg/kg/day and about 800 mg/kg/day.
52. The method of embodiment 47, wherein the therapeutically effective amount of the third composition is between about 250 mg/kg/day and about 400 mg/kg/day.
- 5 53. The method of embodiment 47, wherein the third composition is administered once daily, twice daily, three times daily, four times daily, five times daily or six times daily.
54. The method of embodiment 47, wherein the third composition administered orally, intrathecally, enterally, or intravenously.
55. The method of embodiment 54, wherein the third composition is administered orally
10 and comprises deoxynucleoside mixed with cow's milk, human breast milk, infant formula or water.
56. A method of restoring thymidine kinase 2 enzyme activity in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a first composition comprising a pharmacological agent, and further administering a therapeutically
15 effective amount of a second composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein.
57. The method of embodiment 56, wherein the pharmacological agent is chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), and mixtures thereof; and
20 deoxycytidine monophosphate (dCMP), deoxythymidine monophosphate (dTMP), and mixtures thereof.
58. The method of embodiment 56, wherein the rAAV is AAV9.
59. The method of embodiment 56, wherein the second composition is a pharmaceutical composition.
- 25 60. The method of embodiment 56, wherein the first composition is administered as soon as the subject is diagnosed with TK2 deficiency or suspected of having TK2 deficiency, and is continued to be administered during the administration of the second composition.
61. The method of embodiment 56, wherein the second composition is administered within days of the first composition.
- 30 62. The method of embodiment 56, wherein the second composition is administered within weeks of the first composition.
63. The method of embodiment 56, wherein the second composition is administered at least a month after the first composition.

64. The method of embodiment 56, wherein the therapeutically effective amount of the first composition is between about 100 mg/kg/day and about 1000 mg/kg/day.
65. The method of embodiment 56, wherein the therapeutically effective amount of the first composition is between about 200 mg/kg/day and about 800 mg/kg/day.
- 5 66. The method of embodiment 56, wherein the therapeutically effective amount of the first composition is between about 250 mg/kg/day and about 400 mg/kg/day.
67. The method of embodiment 56, wherein the first composition is administered once daily, twice daily, three times daily, four times daily, five times daily or six times daily.
68. The method of embodiment 56, wherein the first composition administered orally,
10 intrathecally, enterally, or intravenously.
69. The method of embodiment 56, wherein the first composition is administered orally and comprises the pharmacological agent mixed with cow's milk, human breast milk, infant formula or water.
70. The method of embodiment 56, further comprising the administration of a third
15 composition comprising a second rAAV comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene encoding a functional TK2 protein wherein the second rAAV is AAV2.
71. A method of preventing or treating a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof comprising administering to the subject a
20 therapeutically effective amount of a composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene chosen from the group consisting of TK2; DGUOK; TYMP; RRM2B; SUCLA2; SUCLGJ; and MPV17.
72. A method of preventing or treating a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof comprising administering to the subject a
25 therapeutically effective amount of a first composition comprising a pharmacological agent, and further administering a therapeutically effective amount of a second composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a
transgene chosen from the group consisting of TK2; DGUOK; TYMP; RRM2B; SUCLA2;
30 SUCLGJ; and MPV17.
73. The method of embodiment 72, wherein the pharmacological agent is chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), deoxyadenosine (dA), deoxyguanosine (dG), and mixtures thereof; and deoxycytidine monophosphate (dCMP),

deoxythymidine monophosphate (dTMP), deoxyadenosine monophosphate (dAMP), deoxyguanosine monophosphate (dGMP) and mixtures thereof.

74. A method of restoring enzyme activity in a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof comprising administering to the
5 subject a therapeutically effective amount of a composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene chosen from the group consisting of TK2; DGUOK; TYMP; RRM2B; SUCLA2; SUCLGJ; and MPV17.

75. A method of restoring enzyme activity in a disease or disorder characterized by
10 unbalanced nucleotide pools in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a first composition comprising a pharmacological agent, and further administering a therapeutically effective amount of a second composition comprising a recombinant adeno-associated vector (rAAV) comprising a nucleic acid sequence comprising a transgene chosen from the group consisting of TK2;
15 DGUOK; TYMP; RRM2B; SUCLA2; SUCLGJ; and MPV17.

76. The method of embodiment 75, wherein the pharmacological agent is chosen from the group consisting of deoxycytidine (dC), deoxythymidine (dT), deoxyadenosine (dA), deoxyguanosine (dG), and mixtures thereof; and deoxycytidine monophosphate (dCMP), deoxythymidine monophosphate (dTMP), deoxyadenosine monophosphate (dAMP),
20 deoxyguanosine monophosphate (dGMP) and mixtures thereof.

EXAMPLES

The present invention may be better understood by reference to the following non-limiting examples, which are presented in order to more fully illustrate the preferred
25 embodiments of the invention. They should in no way be construed to limit the broad scope of the invention.

Example 1- Materials and Methods

Mouse Model of TK2 Deficiency

30 A homozygous *Tk2* H126N knock-in mutant (*Tk2*^{-/-}) mouse that manifests a phenotype strikingly similar to the human infantile encephalomyopathy has been previously reported (Akman, *et al.* 2008). Between postnatal day 10 and 13, *Tk2*^{-/-} mice rapidly develop fatal encephalomyopathy characterized by decreased ambulation, unstable gait, coarse tremor, growth retardation, and rapid progression to early death at age 14 to 16 days. Molecular and

biochemical analyses of the mouse model demonstrated that the pathogenesis of the disease is due to loss of enzyme activity and ensuing dNTP pool imbalances with decreased dTTP levels in brain and both dTTP and dCTP levels in liver, which, in turn, produces mtDNA depletion and defects of respiratory chain enzymes containing mtDNA-encoded subunits, most prominently in the brain and spinal cord.

All experiments were performed according to a protocol approved by the Institutional Animal Care and Use Committee of the Columbia University Medical Center, and were consistent with the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Mice were housed and bred according to international standard conditions, with a 12-hour light, 12-hour dark cycle, and sacrificed at 4, 13, and 29 days of age.

Organs (brain, spinal cord, liver, heart, kidney, quadriceps muscle, lung, and gastrointestinal tract) were removed and either frozen in the liquid phase of isopentane, pre-cooled near its freezing point (-160°C) with dry ice or fixed in 10% neutral buffered formalin and embedded in paraffin using standard procedures. Paraffin embedded tissue were then stained with hematoxylin and eosin (H&E) for morphological study or processed for immunostaining studies with GFAP, COX I, or complex I. Both heterozygous and homozygous wild type mice were considered as control group ($Tk2^{+}$) since no clinical and biochemical difference were previously described (Akman, *et al.* 2008; Dorado, *et al.* 2011).

hTK2 Construct

The vector comprising the human TK2 gene (pAAVsc CB6 PI TK2 is shown in FIG. 1). Key components of the construct include the human TK2 cDNA as well as 5' ITR, CMV IE enhancer, chicken beta actin promoter, BGH 3' UTR, Poly A signal, GRE, and 3' ITR.

This vector was encapsulated in both AAV9 and AAV2 capsids for the subsequent experiments.

Gene Therapy Administration

Based upon previous results using an AAV9-GFP vector in wild-type mice and analyzing the distribution of the vector showed that AAV9 virus does not target the brain when administered I.P. (Results not shown.) Based upon this work, the administration of the AAV9-hTK2 construct was done by intravenous (IV) injection.

The AAV9-hTK2 was administered via IV with a retro-orbital injection of a total volume of 35 μl of PBS 1X containing about 4.2×10^{11} or 4.2×10^{10} genome or vector copies at postnatal day 1.

Alternatively, AAV9-hTK2 was administered via IV with a retro-orbital injection of 35 μ L of PBS 1X containing about 2.1×10^{11} vector copies at postnatal day 1, followed by administration of 100 μ L of AAV2-hTK2 via IV with a tail vein injection containing about 1.05×10^{11} vector copies at day 29 and with and without supplementation with 520 mg/kg/day of oral dC+dT (in drinking water) from day 21 assuming a water consumption of 4 ml/day/mouse.

Pharmacological Treatment Administration and Experimental Plan

Deoxycytidine (dC) and deoxythymidine (dT) are administered in 50 μ l of Esbilac milk formula for small pets (Pet-Ag) by daily oral gavage to Tk2^{-/-} and aged matched control wild-type (Tk2⁺) using a dose of 520 mg/kg/day, from postnatal day 4 to 29 days. At age postnatal 21 days, mice are separated from the mother and the treatment was continued by administration of dC and dT in drinking water assuming a water consumption of 4 ml/day/mouse. A negative control group of untreated Tk2 mutant and control wild-type mice are weighed and observed closely for comparison.

Deoxycytidine monophosphate (dCMP) and deoxythymidine monophosphate (TMP) (Hongene Biotech, Inc.) are administered in 50 μ l of Esbilac milk formula for small pets (Pet-Ag) by daily oral gavage to Tk2 H126N knock-in mice (Tk2^{-/-}) and aged-matched control wild-type (Tk2⁺) using 2 doses, 200 mg/kg/day and 400 mg/kg/day, from postnatal day 4 to 29 days. At age 29 days, mice are separated from the mother and the treatment is continued by administration of dCMP and TMP in drinking water using a dose of 400 mg/kg/day.

Phenotype assessment

Body weight was assessed daily, since it has been previously observed that incapacity of gaining weight is the first sign of disease (Akman, *et al.* 2008).

To define the degree of safety and efficacy of the gene therapy, survival time, age-at-onset of disease, type and severity of symptoms, occurrence of side effects, and proportion of treatment termination due to adverse events in treated and untreated Tk2 mice were compared. General behavior, survival time, and body weights of the mice were assessed daily beginning at postnatal day 4.

TK2 Enzyme Measurement

The activity of TK2 enzyme was measured as previously described in Franzolin, *et al.* 2006.

mtDNA Copy Number Measurement

mtDNA copy number was measured as previously described in Spinazzola, *et al.* 2006/

Mitochondrial respiratory chain enzyme activities

Mitochondrial RC enzymes analysis was performed in cerebrum tissue as previously described (DiMauro, *et al.* 1987; Birch-Machin, *et al.* 1994; Quinzii, *et al.* 2013).

Statistical methods

5 Data are expressed as the mean $\pm$ SD of at least 3 experiments per group. Gehan-Breslow-Wilcoxon test was used to compare the survival proportion of each group of mice. A p-value of <0.05 was considered to be statistically significant.

Example 2 - Administration of AAV9-hTK2 Resulted in Prolonged Lifespan

10 The mutant mice described in Example 1 were administered via IV with a retro-orbital injection at postnatal day 1 with the AAV9-hTK2 construct (containing about 4.2×10^{11} or 4.2×10^{10} genome or vector copies) also described in Example 1.

Treatment with AAV9-hTK2 resulted in a dose-dependent response with administration of 4.2×10^{10} vc prolonging lifespan up to 3-fold (average of 39 days), similar to
15 520 mg/kg/day of dC+dT therapy. Administration of 4.2×10^{11} vc prolonged lifespan up to 6-fold (average of 89 days) with a maximum of 129 days. See FIG. 2

Example 3 – Administration of AAV9-hTK2 Resulted in Increased Growth, Strength and Motor Function in Mutant Mice

The mice described in Example 1 were treated as described in Example 1 with the
20 AAV9-hTK2 construct also described in Example 1.

The *Tk2*^{-/-} mice treated with the lower dose of AAV9-hTK2 (4.2×10^{10} vc) grew at the same rate as those *Tk2*^{-/-} mice treated with 520 mg/kg/day of dC+dT. They also grew at the same rate as *Tk2*⁺ mice until postnatal day 20, when they reached a plateau.

The *Tk2*^{-/-} mice treated with the higher dose of AAV9-hTK2 (4.2×10^{11} vc) grew at the
25 same rate as *Tk2*⁺ mice until P30.

No differences were found between untreated *Tk2*⁺, and *Tk2*⁺ treated with the higher dose of AAV9-hTK2 (4.2×10^{11} vc). See FIG. 3.

The *Tk2*^{-/-} mice treated with 4.2×10^{11} vc at P1 showed no differences in strength compared to untreated or AAV9-hTK2 treated wild-type mice. Strength was normalized by
30 weight and measured at P29 and P60. See FIG. 4.

Rotarod testing, an outcome measure believed to model motor behavior defects, *i.e.* motor phenotypes, observed in human patients for assessing the motor function, was performed at postnatal day 29 and 60, and showed no differences between the treated mutant mice and treated wild-type mice, although two mutant, one male and one female, could not perform the

test because of weakness and lack of balance. See FIG. 5.

Wild-type mice treated with the AAV9-hTK2 showed no adverse side-effects related to the therapy and were overall healthy.

The results regarding strength and motor function appear to rule out myopathy in the mutant mice and suggest that reduced weight showed in *Tk2*^{-/-} may be caused by a non-muscle related health condition.

Example 4 - Treatment with AAV9-Tk2 Restored TK2 Enzyme Activity in Most Tissues

Using the methods described in Example 1, the activity of the TK2 enzyme was measured at P29 and shown to be restored in mutant mice treated with 4.2×10^{11} vc of AAV9-hTK2 at P1 in brain and liver to levels similar to those found in untreated wild-type mice. In muscle, AAV9-hTK2 increased 40-fold the TK2 activity as compared to untreated wild-type mice. However, the administration of the AAV9-hTK2 showed a poor efficiency restoring TK2 activity in kidney. See FIG. 6.

15

Example 5 - Treatment with AAV9-hTk2 Increased the Expression of hTK2 in Most Tissues

Human *TK2* was measured as a percentage with respect of mRNA expression of mouse *TK2* as described in Example 1 in wild-type mice treated with AAV9-hTK2 (4×10^{11} vc) at 1 month, 2 months, and 6 months in brain, liver, muscle, and kidney tissue.

Human *TK2* was expressed in brain, liver and muscle above 100% of the expression of mouse *TK2* on target tissue, even in wild-type mice after 6 months of treatment. In contrast, expression of human *TK2* in kidney is very low 2 months after the treatment. See FIG. 7.

Example 6 - Treatment with AAV9-hTK2 Rescued mtDNA in Most Tissues

mtDNA copy number was measured as described in Example 1. At postnatal day 29, *Tk2*^{-/-} mice treated with AAV9-hTK2 (4×10^{11} vc) at postnatal day 1 had signs of severe mtDNA depletion only in kidney tissue and mtDNA was partially rescued in brain, heart, liver, intestine and muscle (FIG. 8).

At postnatal day 60, there was very mild mtDNA depletion (65-80% of untreated wild-type mice) in liver, heart, muscle and intestine tissue of AAV9-hTK treated mutant mice. mtDNA in brain tissue of treated mutant mice was 87% that of untreated wild-type mice. This showed a rescue of mtDNA in most tissues of AAV9-hTK2 treated *Tk2*^{-/-} mice. There was severe mtDNA depletion in kidney tissue of the same treated mutant mice. See FIG. 9.

Example 7 - Treatment with AAV9-hTK2 Ameliorated Biochemical Abnormalities in Brain

Respiratory chain enzyme (RCE) activities and protein levels were measured as described in Example 1.

RCE activity in the brain at postnatal day 29 was completely restored in the brains of *Tk2*^{-/-} mice treated with AAV9-hTK2 (4×10^{11} vc) (FIG. 10). Only a slight reduction in complex III activity could be observed in AAV9-hTK2 treated *Tk2*^{-/-} mice compared to untreated wild-type mice (FIG. 11).

Example 8 - AAV9-hTK2 Does Not Rescue Kidney Function

Using the methods in Example 1, kidney tissue of mutant mice treated with AAV9-hTK2 at 4.2×10^{11} vc at their time of death at postnatal day 96 was stained with SDH and Cox. Kidney tissue of age matched wild-type mice treated with AAV9-hTK2 at 4.2×10^{11} vc were stained as well. SDH (blue) staining showed no co-localization with Cox (brown) staining indicating Cox deficiency in the kidney tissue of the mutant mice and likely dysfunctional mitochondria (FIG. 12).

Creatinine and blood urea nitrogen (BUN) index were assessed in both mutant mice treated with AAV9-hTK2 at 4.2×10^{11} vc and wild-type treated with AAV9-hTK2 at 4.2×10^{11} vc both in the survival cohort and at postnatal day 60. Both creatinine and BUN were higher in the mutant mice in both cohorts suggesting kidney dysfunction (FIG. 13).

Example 9 – Administration of AAV9-hTK2 in Combination with AAV2-hTK2 Increases Survival Over the Use of AAV9-hTK2 Alone

Mutant mice as described in Example 1 were administered AAV9-hTK2 at 2.1×10^{11} vc via retro-orbital injection of a total volume of 35 μ l as also described in Example 1 at postnatal day 1. Wild-type mice are also treated at postnatal day 1.

At postnatal day 21, one third of these treated mutant and wild-type mice were supplemented with 520 mg/kg/day of oral dC+dT (in drinking water). At postnatal day 29 two thirds of the mice (all of those supplemented with dC+dT and half of the remaining mice) are administered with AAV2-hTK (Example 1) at 1.05×10^{11} vc, via tail vein injection in a total volume of 100 μ l.

The mutant mice treated with only the AAV9-hTK2 survived an average of 89 days as consistent with the results in Example 2. The mutant mice who received both AAV9-hTK2 and AAV2-hTK2 lived significantly longer, about 120 days (FIG. 14). Despite having an

overall lower dose of vector copies (3.15×10^{11} vc in total versus 4.2×10^{11} vc), mutant mice treated with both constructs grew as much as mutant mice treated only with AAV9-hTK2 (FIG.s 15 and 16) and showed equal strength and motor coordination (FIGs. 17, 18 and 19). At day 60, mutant mice treated with both AAV9-hTK2 and AAV2-hTK2 showed slightly higher levels of mtDNA than those treated only with AAV9-hTK2 (FIG. 20). Mice treated with a combination of AAV9-hTK2 and AAV2-hTK2 also showed a decrease in the content of protein in the urine (FIG. 21) and slightly lower levels of the index BUN (FIG. 22) as compared to mice treated only with AAV9-hTK2, as well as higher Tk2 activity in liver (FIG. 23).

The combined data on mtDNA, protein in the urine and BUN showed that kidney dysfunction can be delayed in mice co-treated with AAV2 or AAV2+dN which also increases lifespan.

Example 10 – Administration of Supplemental Nucleosides Enhances the Effects of the Gene Therapy

Half of the mutant and wild-type mice treated with both the AAV9-hTK2 and the AAV2-hTK2 in Example 9 were further administered oral dC + dT at 520 mg/kg/day in drinking water as described in Example 1. It is assumed the mice ingest 4 ml a day per mouse.

The mutant mice who were given the supplemental nucleosides survive significantly longer, with a survival of more than 180 days (FIG. 14). They also had increased growth (FIGs. 15 and 16), and equal strength and motor coordination than mice treated with either AAV9-hTK2 or a combination of AAV9-hTK2 and AAV2-hTK2. Mice treated with the combination of AAV9-hTK2 and AAV2-hTK2 and further supplemented with dC+dT showed higher levels of mtDNA in liver.

Example 11 – Administration of AAV9-hTK2 in Combination with dC+dT Increase Survival Over the Use of AAV9-hTK2 Alone

Mutant mice as described in Example 1 are administered oral dC+dT (260 or 520mg/kg/day each in milk) from postnatal day 4). At postnatal day 21, half the mice are administered with AAV9-hTK2 as described in Example 1.

The mutant mice given only the dC+dT treatment survived a mean of 31 and 40 days with the 260 and 520 mg/kg/day dose, respectively. The mutant mice who received the subsequent AAV9-hTK2 survive significantly longer.

Of the mice that received the AAV9-hTK2 half of these mice then are administered

AAV2-hTK2 at postnatal day 30. These mice who received the dC+dT and the AAV9-hTK2 and the AAV2-hTK2 survive significantly longer than the mice receiving only the dC+dT and AAV9-hTK2.

5 Example 12 – Administration of AAV9-hTK2 in Combination with dCMP + TMP Increase Survival Over the Use of AAV9-hTK2 Alone

 Mutant mice as described in Example 1 are administered dCMP + TMP (200 mg/kg/day or 400 mg/kg/day each in milk) from postnatal day 4. At postnatal day 21, half the mice are administered with AAV9-hTK2 as described in Example 1.

10 The mutant mice given only the dCMP + TMP treatment survive a mean of 35 and 44 days with the 200 and 400 mg/kg/day dose, respectively. The mutant mice who received the subsequent AAV9-hTK2 survive significantly longer.

 Of the mice that received the AAV9-hTK2 half of these mice then are administered AAV2-hTK2 at postnatal day 30. These mice who received the dCMP + TMP and the AAV9-
15 hTK2 and the AAV2-hTK2 survive significantly longer than the mice receiving only the dC+dT and AAV9-hTK2.

REFERENCES

- Akman, *et al.* (2008) Thymidine kinase 2 (H126N) knock in mice show the essential role of balanced deoxynucleotide pools for mitochondrial DNA maintenance. *Hum. Mol. Genet.* 17:2433-2440
- 5 Béhin, *et al.* (2012) Adult cases of mitochondrial DNA depletion due to TK2 defect An expanding spectrum. *Neurology* 78:644-648
- Birch-Machin, *et al.* (1994) An evaluation of the measurement of the activities of complexes I-IV in the respiratory chain of human skeletal muscle mitochondria. *Biochem Med Metab Biol* 51:35-42
- 10 Bourdon, *et al.* (2007) Mutation of RRM2B, encoding p53-controlled ribonucleotide reductase (p53R2), causes severe mitochondrial DNA depletion. *Nature Genetics* 39:776-780
- 15 Chanprasert, *et al.* (2012) TK2-Related Mitochondrial DNA Depletion Syndrome, Myopathic Form. *GeneReviews® Internet*, December 6, 2012
- Copeland (2008) Inherited mitochondrial diseases of DNA replication. *Ann. Rev. Med.* 59:131-146
- 20 DiMauro, *et al.* (1987) Cytochrome c oxidase deficiency in Leigh syndrome. *Ann. Neurol.* 22:498-506
- 25 DiMauro, Schon. (2003) Mitochondrial respiratory-chain diseases. *New England Journal of Medicine* 348:2656-2668
- DiMauro, Hirano. (2005) Mitochondrial encephalomyopathies: an update. *Neuromuscul. Disord.* 15:276-286
- 30 Dorado, *et al.* (2011) Onset and organ specificity of Tk2 deficiency depends on Tk1 down-regulation and transcriptional compensation. *Hum. Mol. Genet.* 20:155-64

- Elpeleg, *et al.* (2005) Deficiency of the ADP-forming succinyl-CoA synthase activity is associated with encephalomyopathy and mitochondrial DNA depletion. *Am. J. Hum. Genet.* 76:1081-1086
- 5 Franzolin, *et al.* (2006) Bromovinyl-deoxyuridine: a selective substrate for mitochondrial thymidine kinase in cell extracts. *Biochem. Biophys. Res. Commun.* 344(1):30-6
- Galbiati, *et al.* (2006) New mutations in TK2 gene associated with mitochondrial DNA depletion. *Pediatr. Neurol.* 34:177-185
- 10 Garone, *et al.* (2012). MPV17 Mutations Causing Adult-Onset Multisystemic Disorder With Multiple Mitochondrial DNA Deletions. *Arch Neurol* 69:1648-1651
- Garone, *et al.* (2018) Retrospective Natural History of Thymidine Kinase 2 Deficiency. *J. Med.*
15 *Genetics* 55:515-21
- Gotz, *et al.* (2008) Thymidine kinase 2 defects can cause multi-tissue mtDNA depletion syndrome. *Brain* 131:2841-2850
- 20 Hirano, *et al.* (2001) Defects of intergenomic communication: autosomal disorders that cause multiple deletions and depletion of mitochondrial DNA. *Semin. Cell. Develop. Biol.* 12:417-427
- Longley, *et al.* (2006). Mutant POLG2 disrupts DNA polymerase gamma subunits and causes
25 progressive external ophthalmoplegia. *Am J Hum Genet.* 78:1026-1034
- Mandel, *et al.* (2001) The deoxyguanosine kinase gene is mutated in individuals with depleted hepatocerebral mitochondrial DNA. *Nature Genet.* 29:337-341
- 30 Naviaux, Nguyen. (2004) POLG mutations associated with Alpers' syndrome and mitochondrial DNA depletion. *Ann. Neurol.* 55:706-712
- Nishino, *et al.* (1999). Thymidine phosphorylase gene mutations in MNGIE, a human mitochondrial disorder. *Science* 283:689-692.

- Ostergaard, *et al.* (2007) Deficiency of the alpha subunit of succinate-coenzyme A ligase causes fatal infantile lactic acidosis with mitochondrial DNA depletion. *Am. J. Hum. Genet.* 81: 383-387
- Paradas, *et al.* (2012) TK2 mutation presenting as indolent myopathy. *Neurology* 29:504-506
- 5 Quinzii, *et al.* (2013) Tissue-specific oxidative stress and loss of mitochondria in CoQ-deficient Pdss2 mutant mice. *FASEB J.* 27:612-621
- Ronchi, *et al.* (2012). Next-generation sequencing reveals DGUOK mutations in adult patients
10 with mitochondrial DNA multiple deletions. *Brain* 135:3404-3415.
- Saada, *et al.* (2003) Mitochondrial deoxyribonucleoside triphosphate pools in thymidine kinase 2 deficiency. *Biochem. Biophys. Res. Commun.* 310:963-966
- 15 Sarzi, *et al.* (2007) Twinkle helicase (PEO1) gene mutation causes mitochondrial DNA depletion. *Ann. Neurol.* 62: 579-587
- Spelbrink, *et al.* (2001). Human mitochondrial DNA deletions associated with mutations in the gene encoding Twinkle, a phage T7 gene 4-like protein localized in mitochondria. *Nature*
20 *Genet.* 28:223-231
- Spinazzola, *et al.* (2006) MPV17 encodes an inner mitochondrial membrane protein and is mutated in infantile hepatic mitochondrial DNA depletion. *Nature Genet.* 38:570-575
- 25 Tynismaa, *et al.* (2012) Thymidine kinase 2 mutations in autosomal recessive progressive external ophthalmoplegia with multiple mitochondrial DNA deletions. *Hum. Mol. Genet.* 21:66-75
- Tynismaa, *et al.* (2009). A heterozygous truncating mutation in RRM2B causes autosomal-
30 dominant progressive external ophthalmoplegia with multiple mtDNA deletions. *Am. J. Hum. Genet.* 85: 290-295
- Van Goethem, *et al.* (2001) Mutation of POLG is associated with progressive external ophthalmoplegia characterized by mtDNA deletions. *Nature Genet.* 28:211-212.

CLAIMS:

1. A method of treating, preventing, and/or curing a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof, the method comprising administering
5 to the subject a therapeutically effective amount of a composition comprising a transgene encoding thymidine kinase 2 (TK2), deoxyguanosine kinase (dGK), thymidine phosphorylase (TP), p53 inducible small subunit of ribonucleotide reductase (p53R2), succinyl-CoA ligase ADP-forming subunit beta (SUCLA2), succinyl-CoA ligase GDP-forming subunit alpha (SUCLG1), mitochondrial inner membrane protein MPV17 (MPV17), and/or DNA
10 polymerase subunit gamma (POLG).
2. A method of restoring enzyme activity in a subject having a disease or disorder characterized by unbalanced nucleotide pools, the method comprising administering to the subject a therapeutically effective amount of a composition comprising a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG.
- 15 3. A method of alleviating one or more symptoms associated with a disease or disorder characterized by unbalanced nucleotide pools in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG.
- 20 4. The method of any one of claims 1-3, wherein the disease or disorder is a mitochondrial disease.
5. The method of claim 4, wherein the mitochondrial disease is a mitochondrial DNA depletion syndrome (MDS).
6. The method of claim 5, wherein the MDS is a myopathic MDS characterized by one or
25 more mutations in an endogenous gene encoding TK2.
7. The method of claim 5, wherein the MDS is an encephalomyopathic form characterized by one or more mutations in an endogenous gene encoding SUCLA2.
8. The method of claim 5, wherein the MDS is a neurogastrointestinal encephalopathic form characterized by one or more mutations in an endogenous gene encoding TP.

9. The method of claim 5, wherein the MDS is a hepatopathic form characterized by one or more mutations in an endogenous gene encoding dGK, MPV17, and/or POLG.
10. The method of any one of claims 1-9, wherein the transgene encodes TK2.
11. The method of claim 10, wherein the TK2 has an amino acid sequence that is at least
5 85% identical to the amino acid sequence of SEQ ID NO: 1.
12. The method of claim 11, wherein the TK2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 1.
13. The method of claim 12, wherein the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1.
- 10 14. The method of claim 13, wherein the TK2 has the amino acid sequence of SEQ ID NO: 1.
15. The method of any one of claims 10-14, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2.
16. The method of claim 15, wherein the transgene has a nucleic acid sequence that is at
15 least 85% identical to the nucleic acid sequence of SEQ ID NO: 2.
17. The method of claim 16, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2.
18. The method of claim 17, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 2.
- 20 19. The method of any one of claims 1-9, wherein the transgene encodes dGK.
20. The method of claim 19, wherein the dGK has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 3.
21. The method of claim 20, wherein the dGK has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 3.
- 25 22. The method of claim 21, wherein the dGK has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 3.

23. The method of claim 22, wherein the dGK has the amino acid sequence of SEQ ID NO: 3.
24. The method of any one of claims 19-23, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 4.
- 5 25. The method of claim 24, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 4.
26. The method of claim 25, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 4.
- 10 27. The method of claim 26, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 4.
28. The method of any one of claims 1-9, wherein the transgene encodes TP.
29. The method of claim 28, wherein the TP has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 5.
30. The method of claim 29, wherein the TP has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 5.
- 15 31. The method of claim 30, wherein the TP has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 5.
32. The method of claim 31, wherein the TP has the amino acid sequence of SEQ ID NO: 5.
- 20 33. The method of any one of claims 28-32, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 6.
34. The method of claim 33, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 6.
35. The method of claim 34, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 6.
- 25 36. The method of claim 35, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 6.

37. The method of any one of claims 1-9, wherein the transgene encodes p53R2.
38. The method of claim 37, wherein the p53R2 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 7.
39. The method of claim 38, wherein the p53R2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 7.
40. The method of claim 39, wherein the p53R2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 7.
41. The method of claim 40, wherein the p53R2 has the amino acid sequence of SEQ ID NO: 7.
42. The method of any one of claims 37-41, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 8.
43. The method of claim 42, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 8.
44. The method of claim 43, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 8.
45. The method of claim 44, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 8.
46. The method of any one of claims 1-9, wherein the transgene encodes SUCLA2.
47. The method of claim 46, wherein the SUCLA2 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 9.
48. The method of claim 47, wherein the SUCLA2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 9.
49. The method of claim 48, wherein the SUCLA2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 9.
50. The method of claim 49, wherein the SUCLA2 has the amino acid sequence of SEQ ID NO: 9.

51. The method of any one of claims 46-50, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 10.
52. The method of claim 51, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 10.
- 5 53. The method of claim 52, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 10.
54. The method of claim 53, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 10.
55. The method of any one of claims 1-9, wherein the transgene encodes SUCLG1.
- 10 56. The method of claim 55, wherein the SUCLG1 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 11.
57. The method of claim 56, wherein the SUCLG1 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 11.
58. The method of claim 57, wherein the SUCLG1 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 11.
- 15 59. The method of claim 58, wherein the SUCLG1 has the amino acid sequence of SEQ ID NO: 11.
60. The method of any one of claims 55-59, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 12.
- 20 61. The method of claim 60, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 12.
62. The method of claim 61, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 12.
63. The method of claim 62, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 12.
- 25 64. The method of any one of claims 1-9, wherein the transgene encodes MPV17.

65. The method of claim 64, wherein the MPV17 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 13.
66. The method of claim 65, wherein the MPV17 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 13.
- 5 67. The method of claim 66, wherein the MPV17 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 13.
68. The method of claim 67, wherein the MPV17 has the amino acid sequence of SEQ ID NO: 13.
69. The method of any one of claims 64-68, wherein the transgene has a nucleic acid
10 sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 14.
70. The method of claim 69, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 14.
71. The method of claim 70, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 14.
- 15 72. The method of claim 71, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 14.
73. The method of any one of claims 1-9, wherein the transgene encodes POLG.
74. The method of claim 73, wherein the POLG has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 15.
- 20 75. The method of claim 74, wherein the POLG has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 15.
76. The method of claim 75, wherein the POLG has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 15.
77. The method of claim 76, wherein the POLG has the amino acid sequence of SEQ ID
25 NO: 15.
78. The method of any one of claims 73-77, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 16.

79. The method of claim 78, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 16.
80. The method of claim 79, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 16.
- 5 81. The method of claim 80, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 16.
82. A method of treating, preventing, and/or curing TK2 deficiency in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising a transgene encoding TK2.
- 10 83. A method of restoring TK2 enzyme activity in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising a transgene encoding TK2.
84. A method of alleviating one or more symptoms associated with a TK2 deficiency in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a composition comprising a transgene encoding TK2.
- 15 85. The method of any one of claims 82-84, wherein the subject has or is at risk of developing a MDS.
86. The method of any one of claims 82-85, wherein the TK2 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1.
- 20 87. The method of claim 86, wherein the TK2 has an amino acid sequence that is at least 90% identical to the amino acid sequence of SEQ ID NO: 1.
88. The method of claim 87, wherein the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1.
89. The method of claim 88, wherein the TK2 has the amino acid sequence of SEQ ID NO: 1.
- 25 90. The method of any one of claims 82-89, wherein the transgene has a nucleic acid sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2.

91. The method of claim 90, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 2.
92. The method of claim 91, wherein the transgene has a nucleic acid sequence that is at least 95% identical to the nucleic acid sequence of SEQ ID NO: 2.
- 5 93. The method of claim 92, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 2.
94. The method of any one of claims 1-93, wherein the composition is a vector.
95. The method of claim 94, wherein the vector is a viral vector.
96. The method of claim 95, wherein the viral vector is selected from the group consisting
10 of adeno-associated virus (AAV), adenovirus, lentivirus, retrovirus, poxvirus, baculovirus, herpes simplex virus, vaccinia virus, and a synthetic virus.
97. The method of claim 96, wherein the synthetic virus is chimeric virus, mosaic virus, or pseudotyped virus, and/or comprises a foreign protein, synthetic polymer, nanoparticle, or small molecule.
- 15 98. The method of claim 96, wherein the viral vector is an AAV.
99. The method of claim 98, wherein the AAV is an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10 serotype.
100. The method of claim 99, wherein the AAV is an AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, and AAV9.
- 20
101. The method of claim 98, wherein the viral vector is a pseudotyped AAV.
102. The method of claim 101, wherein the pseudotyped AAV is AAV2/9.
103. The method of claim 101, wherein the pseudotyped AAV is AAV2/8.
104. The method of claim 98, wherein the AAV comprises a recombinant capsid protein.
- 25 105. The method of any one of claims 1-93, wherein the composition is a liposome, vesicle, synthetic vesicle, exosome, synthetic exosome, dendrimer, or nanoparticle.

106. The method of any one of claims 1-105, wherein the transgene is operably linked to a promoter that induces expression of the transgene in a muscle cell.
107. The method of claim 106, wherein the promoter is a chicken beta actin promoter, cytomegalovirus (CMV) promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter, Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial natriuretic factor promoter.
108. The method of any one of claims 1-107, wherein the transgene is operably linked to an enhancer that induces expression of the transgene in a muscle cell.
109. The method of claim 108, wherein the enhancer is a CMV enhancer, a myocyte enhancer factor 2 (MEF2) enhancer, or a MyoD enhancer.
110. The method of any one of claims 1-109, wherein the composition is administered to the subject as soon as, or immediately after, the subject is diagnosed as having a deficiency in an endogenous gene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG, optionally wherein the composition is administered to the subject as soon as, or immediately after, the subject is diagnosed as having a deficiency in an endogenous gene encoding TK2.
111. The method of any one of claims 1-110, the method further comprising administering to the subject a therapeutically effective amount of a second composition comprising a transgene encoding TK2, dGK, TP, p53R2, SUCLA2, SUCLG1, MPV17, and/or POLG.
112. The method of claim 111, wherein the second composition is a vector.
113. The method of claim 112, wherein the vector is a viral vector.
114. The method of claim 113, wherein the viral vector is selected from the group consisting of adeno-associated virus (AAV), adenovirus, lentivirus, retrovirus, poxvirus, baculovirus, herpes simplex virus, vaccinia virus, and a synthetic virus.
115. The method of claim 114, wherein the synthetic virus is chimeric virus, mosaic virus, or pseudotyped virus, and/or comprises a foreign protein, synthetic polymer, nanoparticle, or small molecule.

116. The method of claim 114, wherein the viral vector is an AAV.
117. The method of claim 116, wherein the AAV is an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10 serotype.
118. The method of claim 116, wherein the viral vector is a pseudotyped AAV.
- 5 119. The method of claim 118, wherein the pseudotyped AAV is AAV2/9.
120. The method of claim 118, wherein the pseudotyped AAV is AAV2/8.
121. The method of claim 116, wherein the AAV comprises a recombinant capsid protein.
122. The method of claim 111, wherein the second composition is a liposome, vesicle, synthetic vesicle, exosome, synthetic exosome, dendrimer, or nanoparticle.
- 10 123. The method of any one of claims 110-121, wherein the transgene comprised by the second composition is operably linked to a promoter that induces expression of the transgene in a muscle cell.
124. The method of claim 122, wherein the promoter is a chicken beta actin promoter, CMV promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter,
- 15 Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial natriuretic factor promoter.
125. The method of any one of claims 110-124, wherein the transgene comprised by the second composition is operably linked to an enhancer that induces expression of the transgene
- 20 in a muscle cell.
126. The method of claim 125, wherein the enhancer is a CMV enhancer, a MEF2 enhancer, or a MyoD enhancer.
127. The method of any one of claims 110-126, wherein the second composition is administered to the subject after administration of the first composition to the subject.
- 25 128. The method of claim 127, wherein the second composition is administered to the subject within one or more days or weeks of administration of the first composition to the subject.

129. The method of claim 128, wherein the second composition is administered to the subject at least one month after administration of the first composition to the subject.
130. The method of any one of claims 110-129, wherein administration of the first composition continues while the second composition is administered to the subject.
- 5 131. The method of any one of claims 1-130, the method further comprising administering to the subject a therapeutically effective amount of a third composition comprising a pharmacological agent.
132. The method of claim 131, wherein the pharmacological agent is selected from the group consisting of deoxycytidine (dC), deoxythymidine (dT), deoxyadenosine (dA),
10 deoxyguanosine (dG), deoxycytidine monophosphate (dCMP), deoxythymidine monophosphate (TMP), deoxyadenosine monophosphate (dAMP), deoxyguanosine monophosphate (dGMP), and mixtures thereof.
133. The method of claim 132, wherein the pharmacological agent is selected from the group consisting of dC, dT, and a mixture thereof.
- 15 134. The method of any one of claims 131-133, wherein the third composition is administered to the subject after administration of the first composition to the subject.
135. The method of claim 134, wherein the third composition is administered to the subject within one or more days or weeks of administration of the first composition to the subject.
136. The method of claim 135, wherein the third composition is administered to the subject
20 at least one month after administration of the first composition to the subject.
137. The method of any one of claims 131-136, wherein the third composition is administered to the subject after administration of the second composition to the subject.
138. The method of claim 137, wherein the third composition is administered to the subject within one or more days or weeks of administration of the second composition to the subject.
- 25 139. The method of claim 138, wherein the third composition is administered to the subject at least one month after administration of the second composition to the subject.

140. The method of any one of claims 131-139, wherein administration of the first composition and administration of the second composition continue while the third composition is administered to the subject.

141. The method of any one of claims 131-140, wherein the third composition is administered to the subject in an amount of from about 100 mg/kg/day to about 1,000 mg/kg/day.

142. The method of claim 141, wherein the third composition is administered to the subject in an amount of from about 200 mg/kg/day to about 800 mg/kg/day.

143. The method of claim 142, wherein the third composition is administered to the subject in an amount of from about 250 mg/kg/day to about 400 mg/kg/day.

144. The method of any one of claims 131-143, wherein the third composition is administered to the subject once daily, twice daily, three times daily, four times daily, five times daily, or six times daily.

145. The method of any one of claims 131-143, wherein the third composition is administered to the subject orally in admixture with cow's milk, human breast milk, infant formula, or water.

146. The method of any one of claims 1-145, wherein the first composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular, inhalation, perfusion, lavage, and/or oral administration.

147. The method of any one of claims 1-146, wherein the second composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular, inhalation, perfusion, lavage, and/or oral administration.

148. The method of any one of claims 1-147, wherein the third composition is administered to the subject by way of intravenous, intrathecal, intradermal, transdermal, parenteral, intramuscular, intranasal, subcutaneous, percutaneous, intratracheal, intraperitoneal, intraarterial, intravascular, inhalation, perfusion, lavage, and/or oral administration.

149. The method of any one of claims 1-148, wherein the subject is a mammal.
150. The method of claim 149, wherein the subject is a human.
151. The method of claim 150, wherein the subject is a pediatric human subject.
152. A recombinant viral vector comprising a transgene encoding TK2, dGK, TP, p53R2,
5 SUCLA2, SUCLG1, MPV17, and/or POLG.
153. The vector of claim 152, wherein the transgene encodes TK2.
154. The vector of claim 153, wherein the TK2 has an amino acid sequence that is at least 85% identical to the amino acid sequence of SEQ ID NO: 1.
155. The vector of claim 154, wherein the TK2 has an amino acid sequence that is at least
10 90% identical to the amino acid sequence of SEQ ID NO: 1.
156. The vector of claim 155, wherein the TK2 has an amino acid sequence that is at least 95% identical to the amino acid sequence of SEQ ID NO: 1.
157. The vector of claim 156, wherein the TK2 has the amino acid sequence of SEQ ID NO: 1.
158. The vector of any one of claims 153-157, wherein the transgene has a nucleic acid
15 sequence that is at least 70% identical to the nucleic acid sequence of SEQ ID NO: 2.
159. The vector of claim 158, wherein the transgene has a nucleic acid sequence that is at least 85% identical to the nucleic acid sequence of SEQ ID NO: 2.
160. The vector of claim 159, wherein the transgene has a nucleic acid sequence that is at
20 least 95% identical to the nucleic acid sequence of SEQ ID NO: 2.
161. The vector of claim 160, wherein the transgene has the nucleic acid sequence of SEQ ID NO: 2.
162. The vector of any one of claims 152-161, wherein the vector is selected from the group consisting of adeno-associated virus (AAV), adenovirus, lentivirus, retrovirus,
25 poxvirus, baculovirus, herpes simplex virus, vaccinia virus, and a synthetic virus.

163. The vector of claim 162, wherein the synthetic virus is chimeric virus, mosaic virus, or pseudotyped virus, and/or comprises a foreign protein, synthetic polymer, nanoparticle, or small molecule.
164. The vector of claim 163, wherein the viral vector is an AAV.
- 5 165. The vector of claim 164, wherein the AAV is an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAVrh74, AAVrh.8, or AAVrh.10 serotype.
166. The vector of claim 165, wherein the AAV is an AAV1, AAV2, AAV4, AAV5, AAV6, AAV7, AAV8, and AAV9.
167. The vector of claim 164, wherein the viral vector is a pseudotyped AAV.
- 10 168. The vector of claim 167, wherein the pseudotyped AAV is AAV2/9.
169. The vector of claim 167, wherein the pseudotyped AAV is AAV2/8.
170. The vector of claim 165, wherein the AAV comprises a recombinant capsid protein.
171. The vector of any one of claims 152-170, wherein the transgene is operably linked to a promoter that induces expression of the transgene in a muscle cell.
- 15 172. The vector of claim 171, wherein the promoter is a chicken beta actin promoter, cytomegalovirus (CMV) promoter, myosin light chain-2 promoter, alpha actin promoter, troponin 1 promoter, Na⁺/Ca²⁺ exchanger promoter, dystrophin promoter, creatine kinase promoter, alpha7 integrin promoter, brain natriuretic peptide promoter, alpha B-crystallin/small heat shock protein promoter, alpha myosin heavy chain promoter, or atrial
- 20 natriuretic factor promoter.
173. The vector of any one of claims 152-172, wherein the transgene is operably linked to an enhancer that induces expression of the transgene in a muscle cell.
174. The vector of claim 173, wherein the enhancer is a CMV enhancer, a myocyte enhancer factor 2 (MEF2) enhancer, or a MyoD enhancer.
- 25 175. A kit comprising the vector of any one of claims 152-174 and a package insert instructing a user of the kit to administer the vector to a subject in accordance with the method of any one of claims 1-158.

176. The kit of claim 175, wherein the kit further comprises a pharmacological agent selected from the group consisting of dC, dT, dA, dG, dCMP, TMP, dAMP, dGMP, and mixtures thereof.

FIG. 1





FIG. 2







FIG. 3

FIG. 4







FIG. 5

FIG. 6







FIG. 6 (cont)

FIG. 7







FIG. 8

FIG. 8 (cont)







FIG. 9

FIG. 9 (cont)







FIG. 10

FIG. 10 (cont)






FIG. 11






FIG. 11 (cont)

FIG. 12



Tk2^{-/-} AAV9-hTK2 4.2x10¹¹ vc
at time of death (P96)



Tk2⁺ AAV9-hTK2 4.2x10¹¹ vc
Age matched (P96)




FIG. 13

FIG. 13 (cont)




FIG. 14




FIG. 15




FIG. 16




FIG. 17

Grip test (strength) at P60




FIG. 18





FIG. 19

Rotarod (motor function)



mtDNA copy number

FIG. 20




FIG. 20 (cont)




FIG. 21

Comparison of different time points





FIG. 22

BUN levels in plasma







FIG. 23

FIG. 23 (cont)



02196839.TXT
SEQUENCE LISTING

<110> The Trustees of Columbia University in the City of New York
<120> Gene Therapy for Diseases Caused by Unbalanced Nucleotide Pools
including Mitochondrial DNA Depletion Syndromes
<130> 01001/006458-W00
<140> Filed Herewith
<141> 2019-04-18
<150> 62/659,281
<151> 2018-04-18
<160> 24
<170> PatentIn version 3.5
<210> 1
<211> 234
<212> PRT
<213> Homo sapiens
<400> 1

Met Gly Ala Phe Cys Gln Arg Pro Ser Ser Asp Lys Glu Gln Glu Lys
1 5 10 15

Glu Lys Lys Ser Val Ile Cys Val Glu Gly Asn Ile Ala Ser Gly Lys
20 25 30

Thr Thr Cys Leu Glu Phe Phe Ser Asn Ala Thr Asp Val Glu Val Leu
35 40 45

Thr Glu Pro Val Ser Lys Trp Arg Asn Val Arg Gly His Asn Pro Leu
50 55 60

Gly Leu Met Tyr His Asp Ala Ser Arg Trp Gly Leu Thr Leu Gln Thr
65 70 75 80

Tyr Val Gln Leu Thr Met Leu Asp Arg His Thr Arg Pro Gln Val Ser
85 90 95

Ser Val Arg Leu Met Glu Arg Ser Ile His Ser Ala Arg Tyr Ile Phe
100 105 110

Val Glu Asn Leu Tyr Arg Ser Gly Lys Met Pro Glu Val Asp Tyr Val
115 120 125

Val Leu Ser Glu Trp Phe Asp Trp Ile Leu Arg Asn Met Asp Val Ser
130 135 140

Val Asp Leu Ile Val Tyr Leu Arg Thr Asn Pro Glu Thr Cys Tyr Gln
145 150 155 160

Arg Leu Lys Lys Arg Cys Arg Glu Glu Glu Lys Val Ile Pro Leu Glu
165 170 175

Tyr Leu Glu Ala Ile His His Leu His Glu Glu Trp Leu Ile Lys Gly
180 185 190

Ser Leu Phe Pro Met Ala Ala Pro Val Leu Val Ile Glu Ala Asp His
195 200 205

His Met Glu Arg Met Leu Glu Leu Phe Glu Gln Asn Arg Asp Arg Ile
210 215 220

Leu Thr Pro Glu Asn Arg Lys His Cys Pro
225 230

<210> 2
<211> 4678
<212> DNA
<213> Homo sapiens

<400> 2
gccaagttat ggggtgcgttc tgccagcgtc ctagcagtga taaagaacag gaaaaagaga 60
aaaaatcagt gatctgtgtc gagggcaata ttgcaagtgg gaagacgaca tgcctggaat 120
tcttctccaa cgcgacagac gtcgaggtgt taacggagcc tgtgtccaag tggagaaatg 180
tccgtggcca caatcctctg ggcctgatgt accacgatgc ctctcgctgg ggtcttacgc 240
tacagactta tgtgcagctc accatgctgg acaggcatac tcgtcctcag gtgtcatctg 300
tacggttgat ggagaggtcg attcacagcg caagatacat tttttagtaa aacctgtata 360

02196839.TXT

gaagtgggaa gatgccagaa gtggactatg tagttctgtc ggaatggttt gactggatct	420
tgaggaacat ggacgtgtct gttgatttga tagtttacct tcggaccaat cctgagactt	480
gttaccagag gttaaagaag agatgcaggg aagaggagaa ggtcattccg ctggaatacc	540
tggaagcaat tcaccatctc catgaggagt ggctcatcaa aggcagcctt ttcccatgg	600
cagcccctgt tctggtgatt gaggctgacc accacatgga gaggatgtta gaactctttg	660
aacaaaatcg ggatcgaata ttaactccag agaatcgga gcattgcca taggaggcaa	720
aaggtctatg gctcatgtct gaaaaatgcc tgctgctgcc aagttagcta ttgggagcaa	780
tctggaaaaa cttgctccca ggagggcttt gtgtctggcc agcttgattt tcctaattgg	840
ctcatctcct ttgctagtgt ctttgtcatg cgtctctggc cctcgtgggt aaatgacaaa	900
cgggaccaat gggtttgcca agccctttgc tgttcgcagc ctcacattcc cccggtgcct	960
ctcccatggc tttgtgctgc tgagtcgctc tcatgaagcc cttagggaga gcacctgttg	1020
tgtgcctgac accacgtgg agctgtgtac caatcgtctc agccttcatt aggaggccga	1080
ggtaggagtc ttatatccca ggtgaggaat ttgaagctca gaaaggttga ggggctcccc	1140
agaggtcaca cagcctgtgt gcagtggagc tggcaccatt cagactttca gccgactcag	1200
caactttccc ttgccctggg ctgcctcctc ctgagagctg ttccccaccg ccctgcctct	1260
tccggttggg ggctctcatg tctctttggg gagagctggc agtgtgcgga gctgataaca	1320
ttttcccaat attgagcagt tcccaaggac agtcagcatt tctagacttc cacaaaatta	1380
tgctgcattt ggctggagcc cgggtgtcag tggtttccct gcccagggtc gctgcagccc	1440
catctaccac atcttcatgt ggacattgag attcacatgc tggctcctga aggggtgctca	1500
gtctccttgg tgattaaggt cctgcttgaa ctgctgcaa ctccatgtca gggaagtcgc	1560
ttttggtgcc tggctgggtt gccagagcc aagctggggc aaggggcagc cagccctggc	1620
ttccaaggct cccgtactgt ctgtgtcctt gtataaggag ctttgctctt ggaattactg	1680
aaagtctgtg gccaagaga gagacacaag tggccttaag tctttttgaa gtgttatttc	1740
atccagggaa atgcctcgag ccatagagcc tgaaatcatc tttgttggct cagaaaatac	1800
cttagcttca ctcagctgga ctgcattgaa ggcgaggctg ccccttggat caagcagaaa	1860
acaagagaaa gaaagaacgt tccctttggg gatagtctgg aaagtggga tttgcaaata	1920

02196839.TXT

aaggctctgg aagcattgct ggtcctgaag ctttggagggt gggcagagag agcttcaaga	1980
agactagatg caaaccttg aaaggattaa ggctcaactc tggagaaaca ggccacagcc	2040
tctcagagca gctgttggct gtaaatagag gtagcaaggc cgctcccagg cccctgtgag	2100
tgtgggcacc tgtgcatgca atgctccgac tctgcagagg tgccaagtgc ccctgctggg	2160
ccagtcccag agagttagga agtcaaggcc tgcaactcct ggttcttcct gtttggacca	2220
gttcttgtgc cattggcagg atgagaggca gcagccaggc gggagctgtg tctagcagca	2280
cctgtagccc acgtgctgct aattagctgg aaaactggcg aaggcagaac ctttgctacc	2340
aggaatcttg acatgtgggg tctgtctttg agaatttgta aatgaacagt ccaatatctc	2400
ctctcggctc attttgcaca tccatctttg gggaaatgtg atttctctct cttttttttt	2460
tttttttttt gcctaaggca caatctcaag aggtcctgag accacgtacc catgattttt	2520
ttcttgctct gtgataccca ataactcctt cacctaagcc ctgttggtga ttttgaagtg	2580
cttcctaggc cagtgatttt agcttctgcc agctgctttt gccagtagat taacgtgttt	2640
ttatctttca aactccgtgt ttcctaactg ggagtgtatg ggtctaagag agcctgctgt	2700
cctccctgcc ttccacctg gagaggaggc tggacgcac agcagtggcc agggcaggctc	2760
gcaaaatctc ccagcctaga gaccacacct gaaacggctg aagccagctt gcacaagggc	2820
tgctgtccct ctgcggcagg cagagctggg gggggcaggg gtcacagagc agtcatagac	2880
accatggacc agggcaggag aagggcagat ggcacatggg cacaacaggg ccttgtcctt	2940
agagcactgg ggggtcatgg ctgggagggg catggcaggg gctggcatcc ctgtagagcc	3000
agaggggcca cccagggcag tgacattcca gatatgttgg gctcacctca tccttgctgt	3060
gagactggag ttccatgggg acatgaagtc agtacaccgc agagctgctc agctgctcta	3120
cctctcgctg acttttttgt tgcacatata ctttttcttt caattagcat ttatttcagc	3180
ttttatttaa gctttttgac agtacatgta aaatatatga ttataaccat ttaaaaatac	3240
cttatgtacc tgggtttttt tggaactag atagaaatat atttatcttt tacataaaag	3300
aagtgtgtag tgggggtggc cagggtcttg tgggtggctta gtggccatcg ggggtcccagg	3360
ctcttccac ctttctcttt tgggtccacc tcttggctcc tggcttcctt cctctggggc	3420
ctgactgtcc aggatggcag ctggagctcc tgctctgggc acggttggca tcagagccca	3480

02196839.TXT

ctgctcccca tccactcttt aatccagatg ctggtaatgt ccctttccaa ggcataacta	3540
agataagctg gaaggttctt acgaggcttt gctagaggcc ctgggaggtg gggggaaagg	3600
caagagggca gtgcccaccc atagaccggg tcacatgacc tggcatcagc gccgtggggt	3660
cccttgggct tcacctccct ctcccccttg ccccccacgtc atcccaccct catcctcacc	3720
ctcaccattc ccatcccgtc cgtaatgtc ccttgcaagg cctaactctg tgaggaatga	3780
aaccaccta tccttgttcc aataacatga gtgaccgaat ttacaaacga gtgaatgtgg	3840
acctctggaa acattacca gcttgttttt acttttctact ttctctttct gcccctttca	3900
tttccgtagg agccctttac ctagatgaga agtgtccccc cgcctgggga atatatcagt	3960
cagaacaatc ttctgcaga catgcaccat tagaccgag tgacggtggt gccatttaaa	4020
cctcagagca ggtaaaagggt ggtcctgaaa cctgtctacc cacagggtgc tatggaatct	4080
gaatcattc ttttttcta gagccctggg gtggggagct ccctcaagtg ttcacatgtg	4140
tgtggaatga ggaacaccca tctccttggc cctctccacc ctgaagagtt agttattaaa	4200
ataattggca agctcttgca aatgtcagtc atccattgtt cagaatggaa tagcaataat	4260
acatccctgg ctgccctggg cttggccagg attactcact gaaggcctca gggttactgg	4320
cacacacttt cttttcctaa taatcccatc ccctcagctt tcctaaggct agagtgaatt	4380
tcgtgttcct ttagtttaca taagatggtg aacttggcaa aagctatcat taaacagaag	4440
ctaagagaaa gcctatgtcg tggaatccag aatgggtatt gccattcact gctgtccaca	4500
gaagctgtct tgaatttctt tctgtgtctt ttcttttttt ttctttaaga ctgttgttta	4560
ccagactggg ctctgtggaa cacaggtgtc ctgggagatg gttaatcatt acaaaatatt	4620
ggtaacaatc taaagatgca tacataagag agtgggtcaaa taaaccattt tccattca	4678

<210> 3
 <211> 183
 <212> PRT
 <213> Homo sapiens

<400> 3

Met	Ala	Ala	Gly	Arg	Leu	Phe	Leu	Ser	Arg	Leu	Arg	Ala	Pro	Phe	Ser
1				5					10					15	

02196839.TXT

Ser Met Ala Lys Ser Pro Leu Glu Gly Val Ser Ser Ser Arg Gly Leu
20 25 30

His Ala Gly Arg Gly Pro Arg Arg Leu Ser Ile Glu Gly Asn Ile Ala
35 40 45

Val Gly Lys Ser Thr Phe Val Lys Leu Leu Thr Lys Thr Tyr Pro Glu
50 55 60

Trp His Val Ala Thr Glu Pro Val Ala Thr Trp Gln Asn Ile Gln Ala
65 70 75 80

Ala Gly Thr Gln Lys Ala Cys Thr Ala Gln Ser Leu Gly Asn Leu Leu
85 90 95

Asp Met Met Tyr Arg Glu Pro Ala Arg Trp Ser Tyr Thr Phe Gln Thr
100 105 110

Phe Ser Phe Leu Ser Arg Leu Lys Val Gln Leu Glu Pro Phe Pro Glu
115 120 125

Lys Leu Leu Gln Ala Arg Lys Pro Val Gln Ile Phe Glu Arg Leu His
130 135 140

Phe Glu Ala Leu Met Asn Ile Pro Val Leu Val Leu Asp Val Asn Asp
145 150 155 160

Asp Phe Ser Glu Glu Val Thr Lys Gln Glu Asp Leu Met Arg Glu Val
165 170 175

Asn Thr Phe Val Lys Asn Leu
180

<210> 4

<211> 879

<212> DNA

<213> Homo sapiens

<400> 4

aacggtgcgc tggagcgagt gaggcagcgat acctagggcg gaagtgcctc cggcggaagt

60

02196839.TXT

gatcgctgtg tgaatcgtgg gtgggatggc cgcgggccgc ctcttttctaa gtcggcttcg	120
agcacccttc agttccatgg ccaagagccc actcgagggc gtttcctcct ccagaggcct	180
gcacgcgggg cgcgggcccc gaaggctctc catcgaaggc aacattgctg tgggaaagtc	240
cacgtttgtg aagttactca cgaaaactta cccagaatgg cacgtagcta cagaacctgt	300
agcaacatgg cagaatatcc aggctgctgg caccacaaaa gcctgcactg cccaaagtct	360
tggaaacttg ctggatatga tgtaccggga gccagcacga tggtcctaca cattccagac	420
attttccttt ttgagccgcc tgaaagtaca gctggagccc ttccctgaga aactcttaca	480
ggccaggaag ccagtacaga tctttgagag gctccacttt gaggctctga tgaacattcc	540
agtgctggtg ttggatgtca atgatgattt ttctgaggaa gtaaccaaac aagaagacct	600
catgagagag gtaaacacct ttgtaaagaa tctgtaacca ataccatgaa gttcaggctg	660
tgatctgggc tccctgactt tctgaagcta gaaaaatgtt gtgtctcca accaccttc	720
catccccagc ccctctcatc cctggagcac tctgccgctc aagagctggt ttgttaatta	780
ttgttagact ttgccattgt tttcttttgt acctgaagca ttttgaaaat aaagtttact	840
taagttatgc ttgtttttct aaaaaaaaaa aaaaaaaaaa	879

<210> 5
 <211> 423
 <212> PRT
 <213> Homo sapiens

<400> 5

Met	Leu	Leu	Leu	Arg	Leu	Pro	Pro	His	Arg	Ser	His	Ala	Ser	Pro	Leu
1				5					10					15	

Asp	Cys	Lys	Leu	Gln	Asp	Arg	Cys	Arg	Lys	Cys	Tyr	Ser	Pro	Arg	Ser
			20					25						30	

Gly	Gln	Ala	Cys	Pro	Pro	Ala	Leu	Ala	Ala	Ala	Trp	Leu	Arg	Arg	Cys
		35					40					45			

Glu	Arg	Arg	Gly	Gly	Arg	Pro	Arg	Gly	Gly	Arg	Arg	Lys	Glu	Leu	Thr
	50					55					60				

02196839.TXT

Leu Gly Leu Arg Pro Ala Arg Cys Ser Ala Pro Gly Pro Ala Lys Asp
65 70 75 80

Asp Ala Trp Arg Pro Gln Ala Gly Arg Ser Ser Ser Asp Thr Asn Glu
85 90 95

Ser Glu Ile Lys Ser Asn Glu Glu Pro Leu Leu Arg Lys Ser Ser Arg
100 105 110

Arg Phe Val Ile Phe Pro Ile Gln Tyr Pro Asp Ile Trp Lys Met Tyr
115 120 125

Lys Gln Ala Gln Ala Ser Phe Trp Thr Ala Glu Glu Val Asp Leu Ser
130 135 140

Lys Asp Leu Pro His Trp Asn Lys Leu Lys Ala Asp Glu Lys Tyr Phe
145 150 155 160

Ile Ser His Ile Leu Ala Phe Phe Ala Ala Ser Asp Gly Ile Val Asn
165 170 175

Glu Asn Leu Val Glu Arg Phe Ser Gln Glu Val Gln Val Pro Glu Ala
180 185 190

Arg Cys Phe Tyr Gly Phe Gln Ile Leu Ile Glu Asn Val His Ser Glu
195 200 205

Met Tyr Ser Leu Leu Ile Asp Thr Tyr Ile Arg Asp Pro Lys Lys Arg
210 215 220

Glu Phe Leu Phe Asn Ala Ile Glu Thr Met Pro Tyr Val Lys Lys Lys
225 230 235 240

Ala Asp Trp Ala Leu Arg Trp Ile Ala Asp Arg Lys Ser Thr Phe Gly
245 250 255

Glu Arg Val Val Ala Phe Ala Ala Val Glu Gly Val Phe Phe Ser Gly
260 265 270

02196839.TXT

Ser Phe Ala Ala Ile Phe Trp Leu Lys Lys Arg Gly Leu Met Pro Gly
275 280 285

Leu Thr Phe Ser Asn Glu Leu Ile Ser Arg Asp Glu Gly Leu His Cys
290 295 300

Asp Phe Ala Cys Leu Met Phe Gln Tyr Leu Val Asn Lys Pro Ser Glu
305 310 315 320

Glu Arg Val Arg Glu Ile Ile Val Asp Ala Val Lys Ile Glu Gln Glu
325 330 335

Phe Leu Thr Glu Ala Leu Pro Val Gly Leu Ile Gly Met Asn Cys Ile
340 345 350

Leu Met Lys Gln Tyr Ile Glu Phe Val Ala Asp Arg Leu Leu Val Glu
355 360 365

Leu Gly Phe Ser Lys Val Phe Gln Ala Glu Asn Pro Phe Asp Phe Met
370 375 380

Glu Asn Ile Ser Leu Glu Gly Lys Thr Asn Phe Phe Glu Lys Arg Val
385 390 395 400

Ser Glu Tyr Gln Arg Phe Ala Val Met Ala Glu Thr Thr Asp Asn Val
405 410 415

Phe Thr Leu Asp Ala Asp Phe
420

<210> 6
<211> 4913
<212> DNA
<213> Homo sapiens

<400> 6
atgaggtaaa tgttgctgtt gcgtcttccc cctcaccgca gtcacgccag cccgttagat 60
tgcaagttgc aggaccgctg taggaaatgt tattcgccgc ggtcaggaca ggcctgtccg 120
cccgccctcg ccgcagcctg gcttcgtcgt tgcgagcgcc ggggaggccg tccccgggga 180

02196839.TXT

gggcggagga	aggagctgac	tttgggtttg	cgtcccgctc	gctgctctgc	cccggggcca	240
gccaaggacg	acgcttggag	gcctcaggcc	gggagatcat	cttcagacac	caacgaaagt	300
gaaataaagt	caaatgaaga	gccactccta	agaaagagtt	ctcgccgggt	tgtcatcttt	360
ccaatccagt	accctgatat	ttggaaaatg	tataaacagg	cacaggcttc	cttctggaca	420
gcagaagagg	tcgacttata	aaaggatctc	cctcactgga	acaagcttaa	agcagatgag	480
aagtacttca	tctctcacat	cttagccttt	tttgcagcca	gtgatggaat	tgtaaatgaa	540
aatttggtgg	agcgctttag	tcaggaggtg	caggttccag	aggctcgctg	tttctatggc	600
tttcaaattc	tcacgcagaa	tgttcactca	gagatgtaca	gtttgctgat	agacacttac	660
atcagagatc	ccaagaaaag	ggaattttta	tttaatgcaa	ttgaaaccat	gccctatgtt	720
aagaaaaaag	cagattgggc	cttgcgatgg	atagcagata	gaaaatctac	ttttggggaa	780
agagtgggtg	cctttgctgc	tgtagaagga	gttttcttct	caggatcttt	tgctgctata	840
ttctggctaa	agaagagagg	tcttatgcca	ggactcactt	tttccaatga	actcatcagc	900
agagatgaag	gacttcactg	tgactttgct	tgccctgatgt	tccaatactt	agtaaataag	960
ccttcagaag	aaagggtcag	ggagatcatt	gttgatgctg	tcaaaattga	gcaggagttt	1020
ttaacagaag	ccttgccagt	tggcctcatt	ggaatgaatt	gcattttgat	gaaacagtac	1080
attgagtttg	tagctgacag	attactttgt	gaacttggat	tctcaaaggt	ttttcaggca	1140
gaaaatcctt	ttgattttat	ggaaaacatt	tctttagaag	gaaaaacaaa	tttctttgag	1200
aaacgagttt	cagagtatca	gcgttttgca	gttatggcag	aaaccacaga	taacgtcttc	1260
accttggatg	cagattttta	aaaaacctct	cgtttttaaa	ctctataaac	ttgtcattgg	1320
taaatagtag	tctattttcc	tctgcttaaa	aaaaatttta	agtatatcct	ttaaaggact	1380
ggggggtttg	tcaaaaggaa	atccaaaacc	tattctaaac	aatttgcat	tatataattt	1440
tcctgtttta	caacaagagt	gtgacctaaa	tgcttttgct	ttgtcactga	aataaaagat	1500
ggcattatgt	ggttaagagc	atggggcgag	gggtcagaca	tgagtctaag	gttctgccct	1560
tactccagtg	tgtgaccctt	ggcaagtcag	ttaatcttgg	taaacctcgg	tgtacttatc	1620
tttaaaatgg	gagtaatagt	aggtcctaaa	ttcatagagt	ggatattagg	attaggatgc	1680
aaaaataaat	gcttaaccaa	cactactact	gttagcacca	ctactaatta	tcattcattg	1740

02196839.TXT

ataatattaa ttgcaatgat gttgtaataa aatactctca tttccttaaa ataattgtga	1800
ttctaggtcc taggatctag aattagatct ttgtatTTTT aatgcttagg ggaagaatat	1860
aagtatctcc ttaaaaagaa cataattctc attcacgcaa gaataagttc tttgaattcc	1920
ttagtatgta gtgaagaaaa tttagttggt agttgctttg ggaagcctac ttatggagtg	1980
gaaaccagga ggttatcatg gtagttgacc ttataagaaa aatgattctt cttcagaaat	2040
taaaaacata actattgccca gatttagctc tggaatgttt agaatcaggc tagaatagca	2100
ttttccaaag aatattctaa gagctattag ctctctaga tatttttttg ggggaaaaag	2160
gggattctgt ggtcagatga gtttgggaaa tgctgaacac ttcattcttc tttagcaagt	2220
acagtcagta catcaaagac tgagcagttc agtggtagat aaatttatct cgccctgcat	2280
attcccaaca tacttaacac agatgttttt tacctgttaa catctcacc agctagtgtt	2340
cctcagaaca aagattggaa aaagctggcc gagaaccatt tatacataga ggaagggctt	2400
atggactgag aaaggagaa catggtaggg attattgaat catttcaa at ttataaccagc	2460
ctgaatagtg taccagcaat tgacttaggc tgtgtttctt tatggtttta aaactcttga	2520
gctgttataa gagatagttc ttttaatgtg actatgcaac atgatagcca atggtgaggg	2580
aaaaggaggt ttctctagaa gagtctgatg aaaggccggg aaccaagggt tttgagaagt	2640
ctgcccctat ttatttttag taagtatcaa gaggtagcct gagcctagtt agagttagac	2700
ctgtctttgg atgaagaagt cttactactg aaatactgaa tttttaatac attattattt	2760
ggtattctgt atacccttc aagcagttgt ttcccattcc caacaaactg tactttatac	2820
aattctggat gctaaaactt agagattttc tctttgcata aattttggct ccattctttc	2880
cataacaatc taatcaaac tgggagttct caagtgaatg caaaaggagc aggccataac	2940
tttatttggt agatacactg tcagaaactt gagatctttt ggcctatgat aataccatta	3000
atttttgcat tgcttcagtt tgccaagtgt ttttacatca tctcatttga tctcaaaaca	3060
gcttgacaga gcaactgtta ttgaaatatt acagatggaa agaatgaggc tcagggaagt	3120
taaatgactt ggccaagatc tgctcatcgt cactgtctgt acagtatttt ttttagagg	3180
ttgtaatgtc tcagatttag tcctttacca tctatgttga tttgcttttg tctatttcct	3240
cattaattga atatacttta aatatatata ttaaagtatc aaaatataga gagacatttg	3300

02196839.TXT

aactgtattc aggtaatatg tttaaagata tttatatatt gccatacaaa aacttaacat	3360
ttaaaactga taatatctgt aatgacatca gaatgaaaga aaaaaaattg tacagtgtat	3420
attcctttgt tttgaatcca aatctttttc ataggtaatg acagatgcct taatgtgaag	3480
cttatttata atagcaataa acctaactgg atttggatga agaagtctta atactgacat	3540
actggatttt taatgcactg gtttgttatt tgggtattcta tctctttttc caggcctcca	3600
ggttgacacat ttatttatta tgttcaatac tttggttctt agttcttaaa gaatcaagaa	3660
gttgtgtaat cttttaaaaa tattatcttg cagataaaga aaaaaattaa gagtgtgttt	3720
acaactgttt tctctttttt acagtacatg tatttaaadc attgctataa taaagttaag	3780
ttcattagga atataaaaac ttgcagttct atgatagatt gcatttatta aaaatgtttc	3840
attgtatcac atagaaatat ggccaggaag gacttgagaa gacagtttga tccattgctt	3900
ttagacagga ctgggttttg ctgtccaatt atatacaata atagtttttc ttacaactaa	3960
gctggcccca gccttgtctt gatattaata catgaaattt ttataattgt ctcatgtctt	4020
catttagaaa catccatatt tttctgcttt ttctattgcc attttttatt tgtgcatgaa	4080
ttgattattg agaaaatgta gcagtttgca tatttaaaaa ttaatcattt tgcattttac	4140
atttaaatat gctaacatca ctgtcataga attcccaaatt ttcatttgta gataactgaac	4200
taagggctaa tgtcaggagc tgatttttaa tgataaagct gcagatgggc taaataaaaag	4260
ccaaattaat cctacaatca ggtattatgt ttttaaacca agttgagtga attggtagt	4320
gacttgggaa atcttcccca gcagaatctg gatgaatggc acagaattga aatctctttg	4380
tttcccacca tttcccttta agtgctctgc tcctttgtaa aaagttaaag atttgaaaga	4440
gaatctcata ttcccagggc attaggaaga aaggatttaa tcccttcaat ttggggctta	4500
atcttgttta aaaaaatgta agtgaagatg gaaggctgga gagaatgatt gctttttgta	4560
cagttaaata aggtcacaat attcttacat actttgtttt acaactgtgt tttcattttt	4620
tcaaagtgtc ggccatttag caaagttatt tactatttac tgtgtacata gaaagcttta	4680
ttatgtgtgg tgtatctaaa ttttttttgc tgaaatacat tatgggtcaat caagccaagc	4740
ctgcatgtac agaatttggt tttttttcaa ataaattagt tgttttctta tttttttggc	4800
ttagtatgtt gaaataaact atggtatctt catcattttg tacatttcct ttttgaggaa	4860

ggtttcttta taagtgaag ggctacccta ataaaggaat gtatatactt act

4913

<210> 7

<211> 482

<212> PRT

<213> Homo sapiens

<400> 7

Met Ala Ala Leu Met Thr Pro Gly Thr Gly Ala Pro Pro Ala Pro Gly
1 5 10 15

Asp Phe Ser Gly Glu Gly Ser Gln Gly Leu Pro Asp Pro Ser Pro Glu
20 25 30

Pro Lys Gln Leu Pro Glu Leu Ile Arg Met Lys Arg Asp Gly Gly Arg
35 40 45

Leu Ser Glu Ala Asp Ile Arg Gly Phe Val Ala Ala Val Val Asn Gly
50 55 60

Ser Ala Gln Gly Ala Gln Ile Gly Ala Met Leu Met Ala Ile Arg Leu
65 70 75 80

Arg Gly Met Asp Leu Glu Glu Thr Ser Val Leu Thr Gln Ala Leu Ala
85 90 95

Gln Ser Gly Gln Gln Leu Glu Trp Pro Glu Ala Trp Arg Gln Gln Leu
100 105 110

Val Asp Lys His Ser Thr Gly Gly Val Gly Asp Lys Val Ser Leu Val
115 120 125

Leu Ala Pro Ala Leu Ala Ala Cys Gly Cys Lys Val Pro Met Ile Ser
130 135 140

Gly Arg Gly Leu Gly His Thr Gly Gly Thr Leu Asp Lys Leu Glu Ser
145 150 155 160

Ile Pro Gly Phe Asn Val Ile Gln Ser Pro Glu Gln Met Gln Val Leu
165 170 175

Leu Asp Gln Ala Gly Cys Cys Ile Val Gly Gln Ser Glu Gln Leu Val
 180 185 190

Pro Ala Asp Gly Ile Leu Tyr Ala Ala Arg Asp Val Thr Ala Thr Val
 195 200 205

Asp Ser Leu Pro Leu Ile Thr Ala Ser Ile Leu Ser Lys Lys Leu Val
 210 215 220

Glu Gly Leu Ser Ala Leu Val Val Asp Val Lys Phe Gly Gly Ala Ala
 225 230 235 240

Val Phe Pro Asn Gln Glu Gln Ala Arg Glu Leu Ala Lys Thr Leu Val
 245 250 255

Gly Val Gly Ala Ser Leu Gly Leu Arg Val Ala Ala Ala Leu Thr Ala
 260 265 270

Met Asp Lys Pro Leu Gly Arg Cys Val Gly His Ala Leu Glu Val Glu
 275 280 285

Glu Ala Leu Leu Cys Met Asp Gly Ala Gly Pro Pro Asp Leu Arg Asp
 290 295 300

Leu Val Thr Thr Leu Gly Gly Ala Leu Leu Trp Leu Ser Gly His Ala
 305 310 315 320

Gly Thr Gln Ala Gln Gly Ala Ala Arg Val Ala Ala Ala Leu Asp Asp
 325 330 335

Gly Ser Ala Leu Gly Arg Phe Glu Arg Met Leu Ala Ala Gln Gly Val
 340 345 350

Asp Pro Gly Leu Ala Arg Ala Leu Cys Ser Gly Ser Pro Ala Glu Arg
 355 360 365

Arg Gln Leu Leu Pro Arg Ala Arg Glu Gln Glu Glu Leu Leu Ala Pro
 370 375 380

Ala Asp Gly Thr Val Glu Leu Val Arg Ala Leu Pro Leu Ala Leu Val
 385 390 395 400

Leu His Glu Leu Gly Ala Gly Arg Ser Arg Ala Gly Glu Pro Leu Arg
 405 410 415

Leu Gly Val Gly Ala Glu Leu Leu Val Asp Val Gly Gln Arg Leu Arg
 420 425 430

Arg Gly Thr Pro Trp Leu Arg Val His Arg Asp Gly Pro Ala Leu Ser
 435 440 445

Gly Pro Gln Ser Arg Ala Leu Gln Glu Ala Leu Val Leu Ser Asp Arg
 450 455 460

Ala Pro Phe Ala Ala Pro Ser Pro Phe Ala Glu Leu Val Leu Pro Pro
 465 470 475 480

Gln Gln

<210> 8
 <211> 1679
 <212> DNA
 <213> Homo sapiens

<400> 8
 cgactgccga gctccgccct ccaggcggcc ccacccgcct gccgtcctgg ggcgccgccg 60
 ccccgccgcc ggcagtggac cgctgtgctg gaaccctgaa ccctacggtc ccgacccgcg 120
 ggcgaggccg ggtacctggg ctgggatccg gagcaagcgg gcgagggcag cgccctaagc 180
 aggcattccc gcaggcccgg agcgatggca gccttgatga ccccggaac cggggcccca 240
 cccgcgcctg gtgacttctc cggggaaggg agccaggagac ttcccgaccc ttcgccagag 300
 cccaagcagc tcccggagct gatccgcatg aagcgagacg gaggccgcct gagcgaagcg 360
 gacatcaggg gcttcgtggc cgctgtggtg aatgggagcg cgcagggcgc acagatcggg 420
 gccatgctga tggccatccg acttcggggc atggatctgg aggagacctc ggtgctgacc 480

02196839.TXT

caggccctgg	ctcagtcggg	acagcagctg	gagtggccag	aggcctggcg	ccagcagctt	540
gtggacaagc	attccacagg	gggtgtgggt	gacaaggcca	gcctggtcct	cgcacctgcc	600
ctggcggcat	gtggctgcaa	ggtgccaatg	atcagcggac	gtggctctggg	gcacacagga	660
ggcaccttgg	ataagctgga	gtctattcct	ggattcaatg	tcattccagag	cccagagcag	720
atgcaagtgc	tgctggacca	ggcgggctgc	tgtatcgtgg	gtcagagtga	gcagctgggt	780
cctgcggacg	gaatcctata	tgcagccaga	gatgtgacag	ccaccgtgga	cagcctgcc	840
ctcatcacag	cctccattct	cagtaagaaa	ctcgtggagg	ggctgtccgc	tctgggtggtg	900
gacgttaagt	tcggaggggc	cgcctcttct	cccaaccagg	agcaggcccg	ggagctggca	960
aagacgctgg	ttggcgtggg	agccagccta	gggcttcggg	tcgcggcagc	gctgaccgcc	1020
atggacaagc	ccctgggtcg	ctgcgtgggc	cacgccctgg	aggctggagga	ggcgtgctc	1080
tgcatggacg	gcgcaggccc	gccagactta	agggacctgg	tcaccacgct	cgggggcgcc	1140
ctgctctggc	tcagcggaca	cgcggggact	caggcccagg	gcgctgccc	ggcggccgcg	1200
gcgctggacg	acggctcggc	ccttgccgcg	ttcagcggga	tgctggcggc	gcagggcgtg	1260
gatcccggtc	tggcccgagc	cctgtgctcg	ggaagtcccg	cagaacgccg	gcagctgctg	1320
cctcgcgccc	gggagcagga	ggagctgctg	gcgcccgcag	atggcaccgt	ggagctggtc	1380
cgggcgctgc	cgctggcgct	ggtgctgcac	gagctcgggg	ccgggcgcag	ccgcgctggg	1440
gagccgctcc	gcctgggggt	gggcgcagag	ctgctggctg	acgtgggtca	gaggctgcgc	1500
cgtgggaccc	cctggctccg	cgtgcaccgg	gacggccccc	cgctcagcgg	cccgcagagc	1560
cgcgccctgc	aggaggcgct	cgtactctcc	gaccgcgcgc	cattcgccgc	cccctcgccc	1620
ttcgagagc	tcgttctgcc	gccgcagcaa	taaagctcct	ttgccgcgaa	aaaaaaaaa	1679

<210> 9
 <211> 463
 <212> PRT
 <213> Homo sapiens

<400> 9

Met	Ala	Ala	Ser	Met	Phe	Tyr	Gly	Arg	Leu	Val	Ala	Val	Ala	Thr	Leu
1				5					10					15	

02196839.TXT

Arg Asn His Arg Pro Arg Thr Ala Gln Arg Ala Ala Ala Gln Val Leu
20 25 30

Gly Ser Ser Gly Leu Phe Asn Asn His Gly Leu Gln Val Gln Gln Gln
35 40 45

Gln Gln Arg Asn Leu Ser Leu His Glu Tyr Met Ser Met Glu Leu Leu
50 55 60

Gln Glu Ala Gly Val Ser Val Pro Lys Gly Tyr Val Ala Lys Ser Pro
65 70 75 80

Asp Glu Ala Tyr Ala Ile Ala Lys Lys Leu Gly Ser Lys Asp Val Val
85 90 95

Ile Lys Ala Gln Val Leu Ala Gly Gly Arg Gly Lys Gly Thr Phe Glu
100 105 110

Ser Gly Leu Lys Gly Gly Val Lys Ile Val Phe Ser Pro Glu Glu Ala
115 120 125

Lys Ala Val Ser Ser Gln Met Ile Gly Lys Lys Leu Phe Thr Lys Gln
130 135 140

Thr Gly Glu Lys Gly Arg Ile Cys Asn Gln Val Leu Val Cys Glu Arg
145 150 155 160

Lys Tyr Pro Arg Arg Glu Tyr Tyr Phe Ala Ile Thr Met Glu Arg Ser
165 170 175

Phe Gln Gly Pro Val Leu Ile Gly Ser Ser His Gly Gly Val Asn Ile
180 185 190

Glu Asp Val Ala Ala Glu Ser Pro Glu Ala Ile Ile Lys Glu Pro Ile
195 200 205

Asp Ile Glu Glu Gly Ile Lys Lys Glu Gln Ala Leu Gln Leu Ala Gln
210 215 220

02196839.TXT

Lys Met Gly Phe Pro Pro Asn Ile Val Glu Ser Ala Ala Glu Asn Met
225 230 235 240

Val Lys Leu Tyr Ser Leu Phe Leu Lys Tyr Asp Ala Thr Met Ile Glu
245 250 255

Ile Asn Pro Met Val Glu Asp Ser Asp Gly Ala Val Leu Cys Met Asp
260 265 270

Ala Lys Ile Asn Phe Asp Ser Asn Ser Ala Tyr Arg Gln Lys Lys Ile
275 280 285

Phe Asp Leu Gln Asp Trp Thr Gln Glu Asp Glu Arg Asp Lys Asp Ala
290 295 300

Ala Lys Ala Asn Leu Asn Tyr Ile Gly Leu Asp Gly Asn Ile Gly Cys
305 310 315 320

Leu Val Asn Gly Ala Gly Leu Ala Met Ala Thr Met Asp Ile Ile Lys
325 330 335

Leu His Gly Gly Thr Pro Ala Asn Phe Leu Asp Val Gly Gly Gly Ala
340 345 350

Thr Val His Gln Val Thr Glu Ala Phe Lys Leu Ile Thr Ser Asp Lys
355 360 365

Lys Val Leu Ala Ile Leu Val Asn Ile Phe Gly Gly Ile Met Arg Cys
370 375 380

Asp Val Ile Ala Gln Gly Ile Val Met Ala Val Lys Asp Leu Glu Ile
385 390 395 400

Lys Ile Pro Val Val Val Arg Leu Gln Gly Thr Arg Val Asp Asp Ala
405 410 415

Lys Ala Leu Ile Ala Asp Ser Gly Leu Lys Ile Leu Ala Cys Asp Asp
420 425 430

02196839.TXT

Leu Asp Glu Ala Ala Arg Met Val Val Lys Leu Ser Glu Ile Val Thr
 435 440 445

Leu Ala Lys Gln Ala His Val Asp Val Lys Phe Gln Leu Pro Ile
 450 455 460

<210> 10
 <211> 2182
 <212> DNA
 <213> Homo sapiens

<400> 10
 ccgcctgtcg cctgtgcgcc tgcgcgcggc gccgagggga cgggggtccga ctcagaaatg 60
 gcggcctcca tgttctacgg caggctagtg gccgtggcca cccttcggaa ccaccggcct 120
 cggacggccc agcgggctgc tgctcaggtt ctgggaagtt ctggattggt taataaccat 180
 ggactccaag tacagcagca acagcaaagg aatctctcac tacatgaata catgagtatg 240
 gaattattgc aagaagctgg tgtctccgtt cccaaaggat atgtggcaaa gtcaccagat 300
 gaagcttatg caattgcaa aaaattaggt tcaaaagatg tcgtgataaa ggcacaggtt 360
 ttagctggtg gtagaggaaa aggaacattt gaaagtggcc tcaaaggagg agtgaagata 420
 gttttctctc cagaagaagc aaaagctggt tcttcacaaa tgattgggaa aaaattgttt 480
 accaagcaaa cgggagaaaa gggcagaata tgcaatcaag tattggtctg tgagcgaaaa 540
 tatcccagga gagaatacta ctttgcaata acaatggaaa ggatcatttca aggtcctgta 600
 ttaataggaa gttcacatgg tgggtgtcaac attgaagatg ttgctgctga gtctcctgaa 660
 gcaataatta aagaacctat tgatattgaa gaaggcatca aaaaggaaca agctctccag 720
 cttgcacaga agatgggatt tccacctaatt attgtggaat cagcagcaga aaacatggtc 780
 aagctttaca gcctttttct gaaatacgat gcaaccatga tagaaataaa tccaatgggtg 840
 gaagattcag atggagctgt attgtgtatg gatgcaaaga tcaattttga ctctaattca 900
 gcctatcgcc aaaagaaaat ctttgatcta caggactgga cccaggaaga tgaaaggac 960
 aaagatgctg ctaaggcaaa tctcaactac attggcctcg atggaaatat aggctgccta 1020
 gtaaagtgtg ctggtttggc tatggccaca atggatataa taaaacttca tggagggact 1080
 ccagccaact tccttgatgt tgggtggtggt gctacagtcc atcaagtaac agaagcattt 1140

02196839.TXT

aagcttatca cttcagataa aaaggtactg gctattctgg tcaacathtt tggaggaatc	1200
atgcgctgtg atgttattgc acagggtata gtcattggcag taaaagactt ggaaattaaa	1260
atacctgttg tggtagcgtt acaaggtaca cgagtcgatg atgctaaggc actgatagcg	1320
gacagtggac ttaaaatact tgcttgtgat gacttggatg aagctgctag aatggttgta	1380
aagctctctg aaatagtgac ctttagcgaag caagcacatg tggatgtgaa atttcagttg	1440
ccaatatgat ctgaaaaccc agtggatggc tgaagggtgtt aaatgtgcta taatcattaa	1500
gaatactgtg ttctgtgtta ttgttctttt tctttttagt gtgtggagat tgtaattgcc	1560
atctaggcac acaaacatht aaaaggatht ggactgcatt taattgtacc attcagaatg	1620
gactgtttgt acgaagcatg tataatgcag ttatcttctt tcttttgtcg cagccagtct	1680
tttttgcttc tcctacaaaa cgtaacttgc aatttgccag tttattattg ttggatacaa	1740
agttcttcat tgataagagt cctataaata agataaatac gaagataaag ctttattctt	1800
tagtgtaaaa atacagtata tctaataact agcctcatta gtagagcagt atattaaaac	1860
aatgttttat gtaaaaagtg tttatcttca gcaccaaata catgataaat gtatcaatca	1920
ctatttataa acagagcttt caaacactcc tcagaatatt cttctaagta ttttgatgaa	1980
gtaactttgt aattatttga acattgtttt aatcattagg aaacactgat taactgcaag	2040
tcttcatgat tctgtcatat taagaaacac ctgtaggttt gcttcaaata aaggcatata	2100
taccaaggac ttacagacaa aattaagaat gtcaatttaa gttaataaaaa atctcccaat	2160
atgaaaaaaaa aaaaaaaaaa aa	2182

<210> 11
 <211> 346
 <212> PRT
 <213> Homo sapiens

<400> 11

Met	Thr	Ala	Thr	Leu	Ala	Ala	Ala	Ala	Asp	Ile	Ala	Thr	Met	Val	Ser
1				5					10					15	

Gly	Ser	Ser	Gly	Leu	Ala	Ala	Ala	Arg	Leu	Leu	Ser	Arg	Ser	Phe	Leu
			20					25					30		

02196839.TXT

Leu Pro Gln Asn Gly Ile Arg His Cys Ser Tyr Thr Ala Ser Arg Gln
35 40 45

His Leu Tyr Val Asp Lys Asn Thr Lys Ile Ile Cys Gln Gly Phe Thr
50 55 60

Gly Lys Gln Gly Thr Phe His Ser Gln Gln Ala Leu Glu Tyr Gly Thr
65 70 75 80

Lys Leu Val Gly Gly Thr Thr Pro Gly Lys Gly Gly Gln Thr His Leu
85 90 95

Gly Leu Pro Val Phe Asn Thr Val Lys Glu Ala Lys Glu Gln Thr Gly
100 105 110

Ala Thr Ala Ser Val Ile Tyr Val Pro Pro Pro Phe Ala Ala Ala Ala
115 120 125

Ile Asn Glu Ala Ile Glu Ala Glu Ile Pro Leu Val Val Cys Ile Thr
130 135 140

Glu Gly Ile Pro Gln Gln Asp Met Val Arg Val Lys His Lys Leu Leu
145 150 155 160

Arg Gln Glu Lys Thr Arg Leu Ile Gly Pro Asn Cys Pro Gly Val Ile
165 170 175

Asn Pro Gly Glu Cys Lys Ile Gly Ile Met Pro Gly His Ile His Lys
180 185 190

Lys Gly Arg Ile Gly Ile Val Ser Arg Ser Gly Thr Leu Thr Tyr Glu
195 200 205

Ala Val His Gln Thr Thr Gln Val Gly Leu Gly Gln Ser Leu Cys Val
210 215 220

Gly Ile Gly Gly Asp Pro Phe Asn Gly Thr Asp Phe Ile Asp Cys Leu
225 230 235 240

02196839.TXT

Glu Ile Phe Leu Asn Asp Ser Ala Thr Glu Gly Ile Ile Leu Ile Gly
245 250 255

Glu Ile Gly Gly Asn Ala Glu Glu Asn Ala Ala Glu Phe Leu Lys Gln
260 265 270

His Asn Ser Gly Pro Asn Ser Lys Pro Val Val Ser Phe Ile Ala Gly
275 280 285

Leu Thr Ala Pro Pro Gly Arg Arg Met Gly His Ala Gly Ala Ile Ile
290 295 300

Ala Gly Gly Lys Gly Gly Ala Lys Glu Lys Ile Ser Ala Leu Gln Ser
305 310 315 320

Ala Gly Val Val Val Ser Met Ser Pro Ala Gln Leu Gly Thr Thr Ile
325 330 335

Tyr Lys Glu Phe Glu Lys Arg Lys Met Leu
340 345

<210> 12
<211> 1270
<212> DNA
<213> Homo sapiens

<400> 12
gtcattggcg tatgaccgca acccttgccg ctgccgctga catcgctacc atggtctccg 60
gcagcagcgg cctcgccgcc gcccgctctc tgtcgcgcag cttcctcctg ccgcagaatg 120
gaattcggca ttgttcttac acagcttctc ggcaacatct ctatgttgat aaaaatacaa 180
agattatattg ccagggtttc actggcaaac agggcacctt tcacagccag caggcattgg 240
aatatggcac caaactcggt ggaggaacca ctccagggaaggaggccag acacatctgg 300
gcttacctgt ctttaatact gtgaaggagg ccaaagaaca gacaggagca acggcttctg 360
tcatttatgt tcctccgcct ttgctgctg ctgccattaa tgaagctatt gaggcagaaa 420
ttcccttggt tgtgtgtatc actgaaggaa ttccccagca ggacatggta cgagtcaagc 480
acaaactgct gcgccaggaa aagacaaggc taattgggcc caactgccct ggagtcatca 540

02196839.TXT

```

atcctggaga atgtaaaatt ggcatcatgc ctggccatat tcacaaaaaa ggaaggattg      600
gcatttgtgc cagatctggc accctgactt atgaagcagt tcaccaaaca acgcaagttg      660
gattggggca gtctttgtgc gttggcattg gaggtgatcc ttttaatgga acagatttta      720
ttgactgcct cgaaatcttt ttgaacgatt ctgccacaga aggcatcata ttgattggtg      780
aaattggtgg taatgcagaa gagaatgctg cagaatTTTT gaagcaacat aattcaggtc      840
caaattccaa gcctgtagtg tccttcattg ctggtttaac tgctcctcct gggagaagaa      900
tgggtcatgc cggggcaatt attgctggag gaaaagggtg agctaaagag aagatctctg      960
cccttcagag tgcaggagtt gtggtcagta tgtctcctgc acagctggga accacgatct     1020
acaaggaatt tgaaaagagg aagatgctat gaaagaaaaa aaaaattcct aaaactgtgg     1080
aatggatcac gtagacatgt aaccagcag cagtttgctt ctgttgcca ctgattaatc     1140
agcctatgtg cctgacactg gtcttgcaat acaactggaa gccaaaacaa ggtggaagat     1200
gtcctgaatt aagatgtttt caccacattg tattacagag acagccaata aatctactat     1260
ttgatttcaa                                     1270

```

<210> 13
 <211> 176
 <212> PRT
 <213> Homo sapiens

<400> 13

Met Ala Leu Trp Arg Ala Tyr Gln Arg Ala Leu Ala Ala His Pro Trp
 1 5 10 15

Lys Val Gln Val Leu Thr Ala Gly Ser Leu Met Gly Leu Gly Asp Ile
 20 25 30

Ile Ser Gln Gln Leu Val Glu Arg Arg Gly Leu Gln Glu His Gln Arg
 35 40 45

Gly Arg Thr Leu Thr Met Val Ser Leu Gly Cys Gly Phe Val Gly Pro
 50 55 60

Val Val Gly Gly Trp Tyr Lys Val Leu Asp Arg Phe Ile Pro Gly Thr
 65 70 75 80

Thr Lys Val Asp Ala Leu Lys Lys Met Leu Leu Asp Gln Gly Gly Phe
 85 90 95

Ala Pro Cys Phe Leu Gly Cys Phe Leu Pro Leu Val Gly Ala Leu Asn
 100 105 110

Gly Leu Ser Ala Gln Asp Asn Trp Ala Lys Leu Gln Arg Asp Tyr Pro
 115 120 125

Asp Ala Leu Ile Thr Asn Tyr Tyr Leu Trp Pro Ala Val Gln Leu Ala
 130 135 140

Asn Phe Tyr Leu Val Pro Leu His Tyr Arg Leu Ala Val Val Gln Cys
 145 150 155 160

Val Ala Val Ile Trp Asn Ser Tyr Leu Ser Trp Lys Ala His Arg Leu
 165 170 175

<210> 14
 <211> 1002
 <212> DNA
 <213> Homo sapiens

<400> 14
 agttcctagg ccagcctgtc acgtgggagg gaggctcggc gctcaggaag catggcactc 60
 tggcgggcat accagcgggc cctggccgct caccctgtga aagtacaggt cctgacagct 120
 gggtcctga tgggcctggg tgacattatc tcacagcagc tgggtggagag gcgggggtctg 180
 caggaacacc agagaggccg gactctgacc atgggtgtccc tgggctgtgg ctttgtgggc 240
 cctgtggtag gaggctggta caaggttttg gatcggttca tccctggcac caccaaagtg 300
 gatgcactga agaagatggt gttggatcag gggggctttg ccccggtgtt tctaggctgc 360
 tttctccac tggtaggggc acttaatgga ctgtcagccc aggacaactg ggccaaacta 420
 cagcgggatt atcctgatgc cttatcacc aactactatc tatggcctgc tgtgcagtta 480
 gccaaacttct acctgggtccc ctttcattac aggttggccg ttgtccaatg tgttgctgtt 540
 atctggaact cctacctgtc ctggaaggca catcggctct aagcctgcct cactccatcg 600

02196839.TXT

tttccacctt gcagtgatgc agcttgaccc tggaacgggtc agacaacctc ctcaaagtgg	660
gcataccagt ttccacggggg ttgggttgcc ggtcagagct taagaggact agcacctgc	720
aatgccctc ttactctaa aatgtacact gactgcttta gagcccttga taatagtctt	780
attcccacca catactaggc actccataaa tatctgttga accttcatga ctttatcaac	840
tttacacca tatcccagca aatgccactc atccccactc ttcatagaca catttggtac	900
tctaaccctg cctaggcttc ttgtagctcc agctctttag agactcccgg aaccctttat	960
atggtgcctc agtaaatatg ttattaaata tgtaatccgg aa	1002

<210> 15
 <211> 1239
 <212> PRT
 <213> Homo sapiens

<400> 15

Met Ser Arg Leu Leu Trp Arg Lys Val Ala Gly Ala Thr Val Gly Pro	
1 5 10 15	

Gly Pro Val Pro Ala Pro Gly Arg Trp Val Ser Ser Ser Val Pro Ala	
20 25 30	

Ser Asp Pro Ser Asp Gly Gln Arg Arg Arg Gln Gln Gln Gln Gln Gln	
35 40 45	

Gln Gln Gln Gln Gln Gln Gln Pro Gln Gln Pro Gln Val Leu Ser Ser	
50 55 60	

Glu Gly Gly Gln Leu Arg His Asn Pro Leu Asp Ile Gln Met Leu Ser	
65 70 75 80	

Arg Gly Leu His Glu Gln Ile Phe Gly Gln Gly Gly Glu Met Pro Gly	
85 90 95	

Glu Ala Ala Val Arg Arg Ser Val Glu His Leu Gln Lys His Gly Leu	
100 105 110	

Trp Gly Gln Pro Ala Val Pro Leu Pro Asp Val Glu Leu Arg Leu Pro	
115 120 125	

02196839.TXT

Pro Leu Tyr Gly Asp Asn Leu Asp Gln His Phe Arg Leu Leu Ala Gln
130 135 140

Lys Gln Ser Leu Pro Tyr Leu Glu Ala Ala Asn Leu Leu Leu Gln Ala
145 150 155 160

Gln Leu Pro Pro Lys Pro Pro Ala Trp Ala Trp Ala Glu Gly Trp Thr
165 170 175

Arg Tyr Gly Pro Glu Gly Glu Ala Val Pro Val Ala Ile Pro Glu Glu
180 185 190

Arg Ala Leu Val Phe Asp Val Glu Val Cys Leu Ala Glu Gly Thr Cys
195 200 205

Pro Thr Leu Ala Val Ala Ile Ser Pro Ser Ala Trp Tyr Ser Trp Cys
210 215 220

Ser Gln Arg Leu Val Glu Glu Arg Tyr Ser Trp Thr Ser Gln Leu Ser
225 230 235 240

Pro Ala Asp Leu Ile Pro Leu Glu Val Pro Thr Gly Ala Ser Ser Pro
245 250 255

Thr Gln Arg Asp Trp Gln Glu Gln Leu Val Val Gly His Asn Val Ser
260 265 270

Phe Asp Arg Ala His Ile Arg Glu Gln Tyr Leu Ile Gln Gly Ser Arg
275 280 285

Met Arg Phe Leu Asp Thr Met Ser Met His Met Ala Ile Ser Gly Leu
290 295 300

Ser Ser Phe Gln Arg Ser Leu Trp Ile Ala Ala Lys Gln Gly Lys His
305 310 315 320

Lys Val Gln Pro Pro Thr Lys Gln Gly Gln Lys Ser Gln Arg Lys Ala
325 330 335

Arg Arg Gly Pro Ala Ile Ser Ser Trp Asp Trp Leu Asp Ile Ser Ser
 340 345 350

Val Asn Ser Leu Ala Glu Val His Arg Leu Tyr Val Gly Gly Pro Pro
 355 360 365

Leu Glu Lys Glu Pro Arg Glu Leu Phe Val Lys Gly Thr Met Lys Asp
 370 375 380

Ile Arg Glu Asn Phe Gln Asp Leu Met Gln Tyr Cys Ala Gln Asp Val
 385 390 395 400

Trp Ala Thr His Glu Val Phe Gln Gln Gln Leu Pro Leu Phe Leu Glu
 405 410 415

Arg Cys Pro His Pro Val Thr Leu Ala Gly Met Leu Glu Met Gly Val
 420 425 430

Ser Tyr Leu Pro Val Asn Gln Asn Trp Glu Arg Tyr Leu Ala Glu Ala
 435 440 445

Gln Gly Thr Tyr Glu Glu Leu Gln Arg Glu Met Lys Lys Ser Leu Met
 450 455 460

Asp Leu Ala Asn Asp Ala Cys Gln Leu Leu Ser Gly Glu Arg Tyr Lys
 465 470 475 480

Glu Asp Pro Trp Leu Trp Asp Leu Glu Trp Asp Leu Gln Glu Phe Lys
 485 490 495

Gln Lys Lys Ala Lys Lys Val Lys Lys Glu Pro Ala Thr Ala Ser Lys
 500 505 510

Leu Pro Ile Glu Gly Ala Gly Ala Pro Gly Asp Pro Met Asp Gln Glu
 515 520 525

Asp Leu Gly Pro Cys Ser Glu Glu Glu Glu Phe Gln Gln Asp Val Met
 530 535 540

Ala Arg Ala Cys Leu Gln Lys Leu Lys Gly Thr Thr Glu Leu Leu Pro
 545 550 555 560

Lys Arg Pro Gln His Leu Pro Gly His Pro Gly Trp Tyr Arg Lys Leu
 565 570 575

Cys Pro Arg Leu Asp Asp Pro Ala Trp Thr Pro Gly Pro Ser Leu Leu
 580 585 590

Ser Leu Gln Met Arg Val Thr Pro Lys Leu Met Ala Leu Thr Trp Asp
 595 600 605

Gly Phe Pro Leu His Tyr Ser Glu Arg His Gly Trp Gly Tyr Leu Val
 610 615 620

Pro Gly Arg Arg Asp Asn Leu Ala Lys Leu Pro Thr Gly Thr Thr Leu
 625 630 635 640

Glu Ser Ala Gly Val Val Cys Pro Tyr Arg Ala Ile Glu Ser Leu Tyr
 645 650 655

Arg Lys His Cys Leu Glu Gln Gly Lys Gln Gln Leu Met Pro Gln Glu
 660 665 670

Ala Gly Leu Ala Glu Glu Phe Leu Leu Thr Asp Asn Ser Ala Ile Trp
 675 680 685

Gln Thr Val Glu Glu Leu Asp Tyr Leu Glu Val Glu Ala Glu Ala Lys
 690 695 700

Met Glu Asn Leu Arg Ala Ala Val Pro Gly Gln Pro Leu Ala Leu Thr
 705 710 715 720

Ala Arg Gly Gly Pro Lys Asp Thr Gln Pro Ser Tyr His His Gly Asn
 725 730 735

Gly Pro Tyr Asn Asp Val Asp Ile Pro Gly Cys Trp Phe Phe Lys Leu
 740 745 750

Pro His Lys Asp Gly Asn Ser Cys Asn Val Gly Ser Pro Phe Ala Lys
 755 760 765

Asp Phe Leu Pro Lys Met Glu Asp Gly Thr Leu Gln Ala Gly Pro Gly
 770 775 780

Gly Ala Ser Gly Pro Arg Ala Leu Glu Ile Asn Lys Met Ile Ser Phe
 785 790 795 800

Trp Arg Asn Ala His Lys Arg Ile Ser Ser Gln Met Val Val Trp Leu
 805 810 815

Pro Arg Ser Ala Leu Pro Arg Ala Val Ile Arg His Pro Asp Tyr Asp
 820 825 830

Glu Glu Gly Leu Tyr Gly Ala Ile Leu Pro Gln Val Val Thr Ala Gly
 835 840 845

Thr Ile Thr Arg Arg Ala Val Glu Pro Thr Trp Leu Thr Ala Ser Asn
 850 855 860

Ala Arg Pro Asp Arg Val Gly Ser Glu Leu Lys Ala Met Val Gln Ala
 865 870 875 880

Pro Pro Gly Tyr Thr Leu Val Gly Ala Asp Val Asp Ser Gln Glu Leu
 885 890 895

Trp Ile Ala Ala Val Leu Gly Asp Ala His Phe Ala Gly Met His Gly
 900 905 910

Cys Thr Ala Phe Gly Trp Met Thr Leu Gln Gly Arg Lys Ser Arg Gly
 915 920 925

Thr Asp Leu His Ser Lys Thr Ala Thr Thr Val Gly Ile Ser Arg Glu
 930 935 940

His Ala Lys Ile Phe Asn Tyr Gly Arg Ile Tyr Gly Ala Gly Gln Pro
 945 950 955 960

02196839.TXT

Phe Ala Glu Arg Leu Leu Met Gln Phe Asn His Arg Leu Thr Gln Gln
965 970 975

Glu Ala Ala Glu Lys Ala Gln Gln Met Tyr Ala Ala Thr Lys Gly Leu
980 985 990

Arg Trp Tyr Arg Leu Ser Asp Glu Gly Glu Trp Leu Val Arg Glu Leu
995 1000 1005

Asn Leu Pro Val Asp Arg Thr Glu Gly Gly Trp Ile Ser Leu Gln
1010 1015 1020

Asp Leu Arg Lys Val Gln Arg Glu Thr Ala Arg Lys Ser Gln Trp
1025 1030 1035

Lys Lys Trp Glu Val Val Ala Glu Arg Ala Trp Lys Gly Gly Thr
1040 1045 1050

Glu Ser Glu Met Phe Asn Lys Leu Glu Ser Ile Ala Thr Ser Asp
1055 1060 1065

Ile Pro Arg Thr Pro Val Leu Gly Cys Cys Ile Ser Arg Ala Leu
1070 1075 1080

Glu Pro Ser Ala Val Gln Glu Glu Phe Met Thr Ser Arg Val Asn
1085 1090 1095

Trp Val Val Gln Ser Ser Ala Val Asp Tyr Leu His Leu Met Leu
1100 1105 1110

Val Ala Met Lys Trp Leu Phe Glu Glu Phe Ala Ile Asp Gly Arg
1115 1120 1125

Phe Cys Ile Ser Ile His Asp Glu Val Arg Tyr Leu Val Arg Glu
1130 1135 1140

Glu Asp Arg Tyr Arg Ala Ala Leu Ala Leu Gln Ile Thr Asn Leu
1145 1150 1155

Leu Thr Arg Cys Met Phe Ala Tyr Lys Leu Gly Leu Asn Asp Leu
 1160 1165 1170

Pro Gln Ser Val Ala Phe Phe Ser Ala Val Asp Ile Asp Arg Cys
 1175 1180 1185

Leu Arg Lys Glu Val Thr Met Asp Cys Lys Thr Pro Ser Asn Pro
 1190 1195 1200

Thr Gly Met Glu Arg Arg Tyr Gly Ile Pro Gln Gly Glu Ala Leu
 1205 1210 1215

Asp Ile Tyr Gln Ile Ile Glu Leu Thr Lys Gly Ser Leu Glu Lys
 1220 1225 1230

Arg Ser Gln Pro Gly Pro
 1235

<210> 16
 <211> 4452
 <212> DNA
 <213> Homo sapiens

<400> 16
 gcggaccggc cgggtggagg ccacacgcta ccccgaggct gcgtaggccg cgcgaagggg 60
 gacgccgtgc cgtgggcctg gggtcggggg agcagcagac cgggaagcac cgatttgggg 120
 tggaaggcag gcatggtcaa acccatttca ctgacaggag agcagagaca ggacgtgtct 180
 ctctccacgt cttccagcca gtaaaagaag ccaagctgga gcccaaagcc aggtgttctg 240
 actcccagcg tgggggtccc tgcaccaacc atgagccgcc tgctctggag gaaggtggcc 300
 ggcgccaccg tcgggccagg gccggttcca gctccggggc gctgggtctc cagctccgtc 360
 cccgcgtccg accccagcga cgggcagcgg cggcggcagc agcagcagca gcagcagcag 420
 cagcagcaac agcagcctca gcagccgcaa gtgctatcct cggagggcgg gcagctgcgg 480
 cacaacccat tggacatcca gatgctctcg agagggtgc acgagcaa atcttcgggcaa 540
 ggaggggaga tgcctggcga ggccgcggtg cgccgcagcg tcgagcacct gcagaagcac 600

02196839.TXT

gggctctggg	ggcagccagc	cgtgcccttg	cccgacgtgg	agctgcgcct	gccgcccctc	660
tacggggaca	acctggacca	gcacttccgc	ctcctggccc	agaagcagag	cctgccctac	720
ctggaggcgg	ccaacttgct	gttgcaggcc	cagctgcccc	cgaagcccc	ggcttgggcc	780
tgggcggagg	gctggacccg	gtacggcccc	gagggggagg	ccgtaccctg	ggccatcccc	840
gaggagcggg	ccctggtgtt	cgacgtggag	gtctgcttgg	cagagggaac	ttgccccaca	900
ttggcgggtg	ccatatcccc	ctcggcctgg	tattcctggg	gcagccagcg	gctggtggaa	960
gagcgttact	cttggaccag	ccagctgtcg	ccggctgacc	tcatccccct	ggaggtccct	1020
actggtgcca	gcagccccac	ccagagagac	tggcaggagc	agttagtggg	ggggcacaat	1080
gtttcctttg	accgagctca	tatcaggagg	cagtacctga	tccagggttc	ccgcatgcgt	1140
ttcctggaca	ccatgagcat	gcacatggcc	atctcagggc	taagcagctt	ccagcgcagt	1200
ctgtggatag	cagccaagca	gggcaaacac	aaggtccagc	ccccacaaa	gcaaggccag	1260
aagtcccaga	ggaaagccag	aagaggccca	gcgatctcat	cctgggactg	gctggacatc	1320
agcagtgtca	acagtctggc	agaggtgcac	agactttatg	tagggggggc	tcccttagag	1380
aaggagcctc	gagaactgtt	tgtgaagggc	accatgaagg	acattcgtga	gaacttccag	1440
gacctgatgc	agtactgtgc	ccaggacgtg	tgggccaccc	atgaggtttt	ccagcagcag	1500
ctaccgctct	tcttggagag	gtgtccccac	ccagtgactc	tggccggcat	gctggagatg	1560
ggtgtctcct	acctgcctgt	caaccagaac	tgggagcggt	acctggcaga	ggcacagggc	1620
acttatgagg	agctccagcg	ggagatgaag	aagtcgttga	tggatctggc	caatgatgcc	1680
tgccagctgc	tctcaggaga	gaggtacaaa	gaagaccctt	ggctctggga	cctggagtgg	1740
gacctgcaag	aatttaagca	gaagaaagct	aagaagggtga	agaaggaacc	agccacagcc	1800
agcaagttgc	ccatcgaggg	ggctggggcc	cctggtgatc	ccatggatca	ggaagacctc	1860
ggcccctgca	gtgaggagga	ggagtttcaa	caagatgtca	tggccccgcg	ctgcttgcat	1920
aagctgaagg	ggaccacaga	gctcctgccc	aagcggcccc	agcaccttcc	tggacaccct	1980
ggatggtacc	ggaagctctg	ccccggcta	gacgaccctg	catggacccc	gggccccagc	2040
ctcctcagcc	tgcagatgcg	ggtcacacct	aaactcatgg	cacttacctg	ggatggcttc	2100
cctctgcact	actcagagcg	tcatggctgg	ggctacttgg	tgcctgggcg	gcgggacaac	2160

02196839.TXT

ctggccaagc tgccgacagg taccaccctg gagtcagctg ggggtggtctg cccctacaga	2220
gccatcgagt ccctgtacag gaagcactgt ctcgaacagg ggaagcagca gctgatgccc	2280
caggaggccg gcctggcgga ggagttcctg ctcactgaca atagtgccat atggcaaacg	2340
gtagaagaac tggattactt agaagtggag gctgaggcca agatggagaa cttgcgagct	2400
gcagtgccag gtcaaccctt agctctgact gcccgtgggtg gcccgaagga caccagccc	2460
agctatcacc atggcaatgg accttacaac gacgtggaca tccctggctg ctggtttttc	2520
aagctgcctc acaaggatgg taatagctgt aatgtgggaa gcccctttgc caaggacttc	2580
ctgccaaga tggaggatgg caccctgcag gctggcccag gaggtgccag tgggccccgt	2640
gctctggaaa tcaacaaaat gattttctttc tggaggaacg cccataaacg tatcagctcc	2700
cagatggtgg tgtggctgcc caggtcagct ctgccccgtg ctgtgatcag gcaccccgac	2760
tatgatgagg aaggcctcta tggggccatc ctgccccaaag tgggtgactgc cggcaccatc	2820
actcgccggg ctgtggagcc cacatggctc accgccagca atgcccggcc tgaccgagta	2880
ggcagtgagt tgaaagccat ggtgcaggcc ccacctggct acacccttgt ggggtgctgat	2940
gtggactccc aagagctgtg gattgcagct gtgcttggag acgcccactt tgccggcatg	3000
catggctgca cagcctttgg gtggatgaca ctgcagggca ggaagagcag gggcactgat	3060
ctacacagta agacagccac tactgtgggc atcagccgtg agcatgcca aatcttcaac	3120
tacggccgca tctatggtgc tgggcagccc tttgctgagc gcttactaat gcagtttaac	3180
caccggctca cacagcagga ggcagctgag aaggcccagc agatgtacgc tgccaccaag	3240
ggcctccgct ggtatcggct gtcggatgag ggcgagtggc tgggtgaggga gttgaacctc	3300
ccagtggaca ggactgaggg tggctggatt tccctgcagg atctgcgcaa ggtccagaga	3360
gaaactgcaa ggaagtcaca gtggaagaag tgggaggtgg ttgctgaacg ggcatggaag	3420
gggggcacag agtcagaaat gttcaataag cttgagagca ttgctacgtc tgacatacca	3480
cgtaccccggtg tgctgggctg ctgcatcagc cgagccctgg agccctcggc tgtccaggaa	3540
gagtttatga ccagccgtgt gaattgggtg gtacagagct ctgctgttga ctacttacac	3600
ctcatgcttg tggccatgaa gtggctgttt gaagagtttg ccatagatgg gcgcttctgc	3660
atcagcatcc atgacaggt tgcgtacctg gtgcgggagg aggaccgcta ccgcgtgcc	3720

02196839.TXT

ctggccttgc agatcaccaa cctcttgacc aggtgcatgt ttgcctacaa gctgggtctg	3780
aatgacttgc cccagtcagt cgcctttttc agtgcagtcg atattgaccg gtgcctcagg	3840
aaggaagtga ccatggattg taaaaccct tccaaccaa ctgggatgga aaggagatac	3900
gggattcccc aggggtgaagc gctggatatt taccagataa ttgaactcac caaaggctcc	3960
ttggaaaaac gaagccagcc tggaccatag cactgcctgg aggctctgta tttgctcccg	4020
tggagcttca tcgggggtggt gcaggctccc aaactcaggc tttcagctgt gctttttgca	4080
aaagggttgc cctaaggcca gccatttttc agtagcagga cctgccaaga agattccttc	4140
taactgaagg tgcagttgaa ttcagtgggt tcagaaccaa gatgccaaca tcggtgtgga	4200
ctacaggaca aggggcattg ttgcttggtg ggtaaaaatg aagcagaagc cccaaagttc	4260
acattaactc aggcatttca tttatTTTTT ccttttcttc ttggctgggt ctttgttctg	4320
tccccatgc tctgatgcag tgccctagaa ggggaaagaa ttaatgctct aacgtgataa	4380
acctgctcca aggcagtgga aataaaaaaga aggaaaaaaa agactctatc ttctcaaaaa	4440
aaaaaaaaaa aa	4452

<210> 17
 <211> 705
 <212> DNA
 <213> Homo sapiens

<400> 17	
atgggagcat tttgccagcg acctagtacc gataaagaac aggagaaaga gaagaagagc	60
gtcatctgtg tggaaggtaa catcgctagt ggcaaaacga cctgtttgga atttttctct	120
aatgccaccg atgtggaggt gcttacagag cctgttagta aatggcgaaa tgttcgagga	180
cataatccgc tgggactgat gtatcacgat gcctccagat ggggactgac ctttcagact	240
tatgtccaac ttacaatgct tgacaggcac actagaccgc aggttagcag cgtgaggctg	300
atggagcgct caattcactc tgccaggtat atattcgttg agaacctcta ccgatctggt	360
aagatgcctg aggtagatta cgtagtgtg agtgaatggt ttgactggat tcttcgcaac	420
atggatgtat cagtggatct cattgtttac cttaggacta acccgagac gtgttatcag	480
cgcttgaaaa agcgctgccg ggaagaggaa aaagtgattc ctctggaata cttggaagct	540
attcatcacc ttcacgaaga gtggctgata aaaggttccc tgtttcccat ggcggtcccc	600

02196839.TXT

gtgcttgtca tagaagcgga tcaccacatg gaaaggatgc tcgaactttt cgagcaaaat	660
cgagaccgga ttctcacgcc ggagaatcga aaacattgcc catag	705

<210> 18
 <211> 552
 <212> DNA
 <213> Homo sapiens

<400> 18 atggctgcag ggaggctttt tttgtctaga ttgagggcac ctttttagctc catggcgaag	60
tctccgcttg aggggggtgag tagcagccgg gggctccatg ctggcagagg gccccggagg	120
ctgtctattg aggggaatat tgcagttggg aagagcacgt ttgttaaatt gctcacgaag	180
acctatccag aatggcacgt tgctacagaa ccagtagcta catggcaaaa cattcaagca	240
gcaggcacac agaaggcttg cactgcacag tctctcggaa acctgctcga tatgatgtac	300
agagagcccg cccggtgggc atacacgttt cagaccttca gctttctttc tcggttgaag	360
gtacagttgg agcccttccc agagaagttg cttcaagcac ggaaacctgt acagatattc	420
gaacgattgc atttcgaggc gctgatgaac atacctgtgc ttgttctgga tgtgaacgac	480
gatttttccg aggaagttac taagcaagag gacctcatgc gggaggtaaa cacgttcgtt	540
aagaacctgt aa	552

<210> 19
 <211> 1449
 <212> DNA
 <213> Homo sapiens

<400> 19 atggcagcat tgatgacacc aggaacgggg gctccgcctg cgcctggcga tttttcagga	60
gaggggagtc aaggtctgcc ggacccttca ccggaaccca aacagcttcc cgagcttatt	120
agaatgaagc gagatggggg tcgcctctct gaggctgaca tacgcggctt tgttgccggc	180
gtcgtcaacg gctcagcaca aggtgcgcaa ataggagcga tgctgatggc tatcaggctg	240
aggggaatgg atttggaaga gaccagtgtg ctgactcaag cgctggcaca gagcggccag	300
cagcttgaat ggccagaagc ttggcgccaa caattggctg ataaacactc aaccgggggc	360
gtcggggaca aagtctccct tgtactggca cctgctctgg ctgcgtgcgg atgtaaggtt	420

02196839.TXT

cctatgatta gcgggcgggg cttgggacat acgggaggaa cgttggacaa actcgaatcc	480
atccctgggt tcaacgtgat tcagagccct gaacaaatgc aagtactcct ggatcaggca	540
ggctgctgta ttgttggcca aagcgaacaa ctctgtcctg cggatgggtat cctctatgct	600
gccagggatg tgaccgccac agttgactcc ctcccgtga taacagcgtc aattttgagt	660
aagaagctcg tcgaagggtc cagcgtcttt gtggttgatg taaaatttgg gggggccgca	720
gtattcccca accaggaaca agcacgggaa ctggcaaaaa ccctggtttg tgtgggcgct	780
tcactgggcc tgagagttgc cgctgccttg accgccatgg ataagccgct cggtaggtgc	840
gtaggtcatg cacttgaagt agaagaagcc cttttgtgca tggatggggc cgggtccccct	900
gacttgcggg atctcgtaac cacgcttggc ggcgcgttgc tttggctttc tggacacgct	960
ggtaccaag cgcaaggggc agcaagagtc gcagcagcgc ttgatgacgg atcagctctt	1020
gggcgatttg aaagaatgct tgcaagctcag ggcgtagatc cagggtctggc gcgggcccctt	1080
tgctcaggtt ctccagcaga gcgccgacag ctctttccca gggcgcgaga gcaggaagaa	1140
ttgctggccc cggctgatgg taccgtcgaa ctctgtacggg ctttgccgct ggctcttggt	1200
ttgcacgaac tgggggctgg gcggagtcgc gccggtgaac ctctcagact cgggtgtgggt	1260
gcggagctcc tcgttgacgt cgggcaacga cttcgcaggg gaacccttg gcttagggta	1320
cacaggagacg ggccagcact cagcggccct cagtccaggg cccttcaaga agctctcgtg	1380
ctgagtgatc gagcgccttt tgccgctccc tcaccatttg ctgaattggt attgccaccc	1440
cagcagtaa	1449

<210> 20
 <211> 1272
 <212> DNA
 <213> Homo sapiens

<400> 20	
atgctgcttc tcagactgcc accccaccgc agtcatgctt cccacttga ctgtaaactt	60
caagataggt gccggaagtg ttactcacc ccatcaggac aagcgtgccc tcctgcgttg	120
gcggccgcct ggcttcggcg gtgtgaacgc cgaggaggtc ggccgagggg tgggcgacgc	180
aaggaactga cccttggcct gcgccctgca cgggtgttcag ctcccgttcc tgccaaggat	240

02196839.TXT

gatgcatgga gacctcaggc aggtcgctcc tcctcagata ccaacgagag tgaaattaaa	300
tctaataaag aaccccttct tcgcaagagt agtcgcaggt tcgtgatatt ccccatataa	360
tatcctgata tctggaaaat gtacaagcaa gcccaagcat cattctggac cgcagaagag	420
gttgatctca gtaaggacct cccacactgg aacaaactga aagccgacga aaaatatttc	480
atatcccata tacttgcttt ctttgcagcc tcagatggaa ttgtgaatga aaacttggtg	540
gagcgcttct ctcaagaggt ccaagttcct gaagcacgat gcttttacgg atttcaaata	600
ctgatagaaa atgtacattc cgaaatgtat tccttgctta tagacacgta tatcagagat	660
ccccaaaaaa gggagttcct gttcaatgcg atcgagacca tgccatatgt taagaaaaaa	720
gccgattggg ccctgaggtg gatagctgat aggaagtcta ctttgggga acgcgttggt	780
gccttcgccg ccgtcgaggg agtcttcttc tccggatcct tcgtgctat cttttggctg	840
aaaaaacgcg gcttgatgcc tggtttgacc tttagcaatg agttgatata acgggatgaa	900
gggctgcatt gtgactttgc ctgccttatg ttccagtact tggatgaaca accgtctgag	960
gaacgggtaa gggaaatcat agtggatgca gtaaaaatag agcaggagtt cttactgaa	1020
gcgctccccg tagggctcat tggcatgaac tgcatctta tgaagcagta catcgagttc	1080
gtggccgata gacttttggg ggagctcggg ttctcaaagg tttttcaggc tgaaaatcct	1140
tttgatttta tggagaacat atctctggag ggcaagacta acttttttga aaaaagagta	1200
tccgagtata aaaggtttgc cgtgatggca gaaacaacag ataatgtctt cacacttgat	1260
gcggattttt aa	1272

<210> 21
 <211> 1392
 <212> DNA
 <213> Homo sapiens

<400> 21	
atggcggctt ctatgttcta tggacgcctg gtcgccgttg ccacgttgcg gaatcacagg	60
ccccgaactg cacaaagggc tgccgctcag gtgcttggaa gcagcggatt gttcaacaat	120
cacggcctcc aagttcagca acagcagcag agaaacctct ctctgcatga gtatatgagt	180
atggaactgc tccaggaagc aggggtgtct gtaccaaagg gctacgtagc aaaaagcccc	240
gacgaggcgt acgcatagc taaaaaactg ggaagtaagg acgtagttat caaagcacag	300

02196839.TXT

gttctggccg gtggccgcgg taagggaacc ttcgagagtg gtttgaaagg cgggtgtcaag	360
atcgtattta gtcccaggga ggccaaggct gtaagttccc agatgatcgg caaaaaattg	420
ttcaciaaagc aaaccggcga aaagggtaga atttgcaacc aagtacttgt ctgcgaaaga	480
aagtatccga gaaggagta ttatttcgct ataaccatgg aaagatcatt ccaaggtccc	540
gtgctcatcg gcagttcaca cgggggcgtt aatatagagg atgtcgctgc agagtctccg	600
gaggccataa ttaaggagcc tatagacatc gaggaaggca ttaaaaaaga acaagccttg	660
caactggcac aaaaaatggg ttttcctccg aacatcgctc agagtgccgc agaaaatatg	720
gtgaaactgt acagcctggt tttgaagtac gatgcgacaa tgatagaaat taatccgatg	780
gtcgaggact cagatggagc cgtgctttgt atggacgcca aaattaactt cgattcaaac	840
agcgcttatt gacagaagaa aattttcgat ttgcaagatt ggacccaaga ggatgagcga	900
gacaaagatg ccgccaaagc aaatctcaat tatataggac tcgatggtaa tattggatgt	960
ttggtcaatg gtgccggtct cgcgatggca actatggata taatcaagtt gcatggtggg	1020
actcccgcca actttcttga cgtcggggga ggcgccaccg ttcaccaagt tacggaggct	1080
ttcaaactca taacctctga caagaagggt ctggcgattc ttgtaaatat ctttgggtggc	1140
attatgcggt gtgatgttat tgcccaggga atagtaatgg ctgtcaaaga tttggagatt	1200
aaaatacccg tcgtgggtgc gctccaaggt actcgagtag atgatgctaa ggctctgatt	1260
gccgacagcg ggctgaagat tctggcatgt gatgatattg acgaggcagc gagaatggtc	1320
gtgaagctct cagaaatagt aactttggcg aaacaggctc atgtcgatgt aaagttccaa	1380
cttcctatat ga	1392

<210> 22
 <211> 1041
 <212> DNA
 <213> Homo sapiens

<400> 22	
atgaccgcta cgcttgccgc tgcagccgac atcgcgacaa tggctctctgg aagttcaggc	60
ttggccgcgg ctccgcttct gagtagaagt ttcttggtgc cgcaaacgg tatcaggcac	120
tgctcctaca ccgccagtag acaacacctt tatgttgata agaacacgaa gataatttgc	180

02196839.TXT

caaggattca ccggttaaaca ggggaccttc cattcccagc aggcactgga atatggaact	240
aaattggttag gcggcactac gcctgggaag ggtggccaaa cacatcttgg tcttcccgtt	300
tttaacacag tgaaggaggc taaagaacaa acggggggcaa cggctagcgt tatctatgtc	360
ccacccccgt ttgccgtgc agccataaat gaggcgattg aagccgagat cccgcttgta	420
gtctgcataa cggagggaat tccgcaacaa gacatggtgc gagtaaagca caagcttctt	480
cgacaggaaa aaacaagact gataggtccg aattgtcctg gcgtaattaa ccccggtgaa	540
tgcaaaatcg gaattatgcc gggacatatt cacaaaaaag gccgaatcgg gatagtcagc	600
agatcaggca ccttgacata cgaagcggtt caccagacta cgcaagtcgg tttgggacaa	660
agcctttgcg tcggtatcgg tggcgacca tttaacggga cggatttcat tgactgtctg	720
gagatcttcc tcaacgattc cgcgacagaa ggaatcattt tgataggaga aataggcggg	780
aacgccgaag agaatgcagc ggaattcctc aaacaacata actctggccc taatagtaag	840
ccggtggtat cattcatagc cggctttaca gcgccgccgg gtcgcagaat gggccacgca	900
ggtgcaatta tagcggggggg caaggaggt gccaaagaaa agatatccgc tcttcaaagc	960
gcaggtgtag tagttagtat gagtccagct caactgggca caaccatcta caaagaattt	1020
gaaaagcgga aaatgctttg a	1041

<210> 23
 <211> 531
 <212> DNA
 <213> Homo sapiens

<400> 23	
atggccctct ggagagctta ccaacgagcc ttggccgcgc acccttggaa ggtacaggtc	60
ttgaccgccg gatctttgat gggctttgga gatattatct ctcaacagtt ggttgaacgg	120
cgaggcctcc aggaacacca aagggggaga acccttaca tggtaagtct gggttgcgga	180
ttcgtcgggc ctgtcgtggg gggatggtat aaggttctgg atcgctttat accggggacc	240
accaaggtcg atgccttgaa gaagatgttg ctggatcagg gaggattcgc tccgtgtttt	300
ttgggatgtt ttctgccact tgttggggct ctgaacggac tgtccgcgca ggataactgg	360
gcgaagtgc aacgcgacta cccagacgcc ctgataacaa attactatct ctggccagca	420
gttcagttgg ccaattttta cctggtacct ctccactatc gccttgctgt agtacagtgt	480

gtcgccgtca tctggaactc atacctttca tggaaggctc atagattgta a 531

<210> 24

<211> 3720

<212> DNA

<213> Homo sapiens

<400> 24

atgtccaggc tgctctggcg gaaggctgca ggcgccactg tcggtccagg accagttcca 60
gctcccggtc gctgggtgag cagcagcgtg ccagctagcg atccgagcga cggtcagcgg 120
cgacggcaac aacagcagca acagcaacaa cagcaacaac aacagcccca gcagcctcag 180
gttctcagtt ccgagggtgg ccaactccga cacaaccac tggaataca gatgctctcc 240
cgcggtctcc acgaacaaat attcggacaa gggggtgaga tgccgggaga ggctgcggtc 300
aggcgcagtg tagaacatct ccagaaacac gggttgtggg gccaacccgc cgttcctctc 360
cccgatgttg aactgcggct tccacctctc tacggtgata atctggacca gcactttaga 420
ctgctcgtc aaaagcagag tctcccttac ctggaagccg ctaacctcct gtcceaagcc 480
caattgcccc ctaaaccgcc agcctgggct tgggcggagg gatggacgag gtatggaccc 540
gaaggggagg ctgtgccagt tgctatacca gaggaacgcg ctctggtttt cgacgtagag 600
gtttgtctcg cggagggaac ttgtcctaca ctggctgtag caatttcccc ttcagcctgg 660
tacagctggg gctctcagag attggtggaa gaaaggata gctggactag ccagctgagt 720
cccgcgacc tcattccact tgaggtaccc accggggcgt caagcccaac tcagagggac 780
tggcaggaac aattggtagt tgggcataat gtgagttttg acagggtca tatccgcgaa 840
cagtatctta tccagggtc tagaatgcga ttccttgaca cgatgagcat gcacatggca 900
atcagcggac ttagttcctt tcagaggtca ttgtggattg cagccaagca gggaaagcat 960
aaggtccaac ccccgacaaa acaaggtcag aaatcccaga gaaaagcccg gcgaggcccc 1020
gccatcagtt cctgggattg gttggatatc agtagtgtga atagccttgc tgagggtgat 1080
cgcctgtatg tgggtggacc ccacttgag aaagagccta gggagctctt cgtcaaaggc 1140
accatgaagg atattagaga gaattttcaa gatctcatgc aatactgcgc acaagacgta 1200
tgggcaacgc atgaggtctt tcaacagcaa ctccccctt ttttgaacg atgtccacat 1260

02196839.TXT

cccgctcactc ttgctgggat gttggaaatg ggtgtaagtt atttgccagt caatcaaaat	1320
tgggagagat acttggctga agcgcagggt acatatgagg aacttcagcg agaaatgaaa	1380
aaaagtctta tggatttggc caatgacgcc tgccagctgc tttccggtga gcggtacaaa	1440
gaagatccat ggctttggga tctcgaatgg gatttgcagg aatttaaaca gaaaaaggcc	1500
aagaaggatca agaaagagcc agctacagcc tcaaagctcc ctatagaggg agcgggagca	1560
ccgggtgatc cgatggatca agaggatttg ggaccttgct ccgaagagga ggaattccaa	1620
caagatgtaa tggcaagggc ctgcctgcaa aagctcaaag ggacaacaga actcttgccc	1680
aagaggcctc aacatctgcc cggccatcca ggttggatat gcaaactctg tccaaggctg	1740
gacgatcccg cctggacccc ggggccctcc cttctgagtc tgcatagag agtgacacct	1800
aagctgatgg cacttacttg ggatgggttc cctcttcact attcagagag acacgggttg	1860
ggatatcttg tcccaggtcg aagggacaat ctggcgaagc ttcccacagg aactaccttg	1920
gagagtgcgg gcgtagtatg tccttatcga gccatagaaa gtctgtatag aaagcattgc	1980
cttgaacaag gcaaacaaca actcatgcct caggaagccg gcctcgctga agaatttctt	2040
cttactgata actctgctat ctggcaaaca gtggagggaac tggattacct cgaagtcgag	2100
gccgaagcga agatggaaaa tttgcgggcg gcggtcccag gtcagcccct tgctttgaca	2160
gcacggggggg gccctaagga taccagccc agctatcatc acgggaacgg tccatataat	2220
gatgttgata tacctggttg ttggtttttt aaacttcctc ataaagatgg caattcatgt	2280
aacgttggat ccccatctgc aaaggacttc ctccccaaaa tggaggatgg gacgttgcaa	2340
gcaggtcccg gtggagcttc tgggccgcga gccctggaaa taaataagat gattagcttc	2400
tggaggaatg cacacaagcg catttcttca cagatggtag tgtggctgcc tcggagtgct	2460
ttgcccaggg ctgtgatcag acaccccgat tatgatgaag agggactgta cggggcaata	2520
ttgccccagg tggttacggc tggtactatt acccgccggg cagttgagcc gacctggctg	2580
acagcatcta atgccaggcc tgatcgcgtg ggttctgaac ttaaagcaat ggtccaagct	2640
ccgcctggat acacgcttgt cggcgcggac gtggactccc aggaactttg gatagcggct	2700
gtccttggcg atgcacattt tgcagggatg cacgggtgca cggtttttgg ctggatgaca	2760
cttcagggga ggaaatcaag ggggaccgac ctgcactcca agaccgcgac aacggtagga	2820

02196839.TXT

atctcaaggg aacacgctaa aatttttcaat tatgggagaa tatatgggtgc cgggtcaaccg	2880
ttcgctgagc gccttctgat gcagtttaac cataggctga cacagcagga ggacgcggag	2940
aaggcgcagc agatgtacgc cgcaactaag ggtctcagat ggtatcgcct ctacagacgag	3000
ggcgagtggc ttgtccggga attgaacctc ccggtcgatc ggacggaagg tggttggatt	3060
agtcttcagg atctcagaaa ggtgcagcgg gaaacagcac gcaagtctca gtggaagaaa	3120
tgggaggtgg tcgccgagcg ggcatggaag ggtggaacag aatccgagat gttaacaag	3180
ctggagagca tagcaactag tgacatccct aggacaccgg tcttgggatg ttgcattagc	3240
cgggcactcg agccatctgc cgtacaggaa gaattcatga cgtcacgggt caactgggtt	3300
gtccagtctt cagccgtcga ttatctgcat ttgatgctgg tcgcgatgaa gtggctcttc	3360
gaagagtttg ctatagatgg ccgattttgc atctctattc acgatgaggt gcgctacctg	3420
gtaagggaag aagaccgata ccgagccgcc ctgcgcccttc agataacaaa tctgcttacc	3480
aggtgtatgt ttgcatacaa gctgggggttg aacgaccttc cccagtccgt cgctttcttt	3540
tcagctgttg atatagatcg ctgcctgaga aaagagggtta cgatggactg caaacgccc	3600
tcaaacccca ctggtatgga gcgcagatat ggcatcccc aaggagaagc cctcgacata	3660
taccagataa ttgagctcac gaagggcagc ctggagaaga gatcccaacc tggcccatag	3720