



Office de la Propriété

Intellectuelle  
du Canada

Un organisme  
d'Industrie Canada

Canadian  
Intellectual Property  
Office

An agency of  
Industry Canada

CA 2464107 A1 2003/06/05

(21) 2 464 107

(12) **DEMANDE DE BREVET CANADIEN**  
**CANADIAN PATENT APPLICATION**

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2002/11/19  
(87) Date publication PCT/PCT Publication Date: 2003/06/05  
(85) Entrée phase nationale/National Entry: 2004/04/19  
(86) N° demande PCT/PCT Application No.: EP 2002/012957  
(87) N° publication PCT/PCT Publication No.: 2003/045919  
(30) Priorité/Priority: 2001/11/26 (01811143.5) EP

(51) Cl.Int.<sup>7</sup>/Int.Cl.<sup>7</sup> C07D 211/94, C08F 2/38, C08K 5/3492,  
C07D 493/10, C08K 5/3435, C09K 15/20, C07C 239/20,  
C07D 401/14

(71) **Demandeur/Applicant:**  
CIBA SPECIALTY CHEMICALS HOLDING INC., CH


(72) **Inventeurs/Inventors:**  
FREY, MARKUS, CH;  
RAST, VALERIE, CH

(74) **Agent:** FETHERSTONHAUGH & CO.

(54) Titre : PROCÉDÉ DE SYNTHÈSE D'AMINES ETHERS A PARTIR D'AMINOXYDES SECONDAIRES  
(54) Title: PROCESS FOR THE SYNTHESIS OF AMINE ETHERS FROM SECONDARY AMINO OXIDES AND USES

**(57) Abrégé/Abstract:**

Amine ethers of sterically hindered amines are obtained in good yield from the corresponding N-oxyl hindered amine precursor by reaction with a hydrocarbon in the presence of an organic hydroperoxide and an iodide. The products of present process find utility as polymerization regulators and/or light stabilizers for organic material.



## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number  
**WO 2003/045919 A3**

(51) International Patent Classification<sup>7</sup>: **C07D 211/94**, 401/14, 493/10, C07C 239/20, C09K 15/20, C08K 5/3435, 5/3492, C08F 2/38

(74) Common Representative: **CIBA SPECIALTY CHEMICALS HOLDING INC.**; Patentabteilung, Klybeckstrasse 141, CH-4057 Basel (CH).

(21) International Application Number:  
PCT/EP2002/012957

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:  
19 November 2002 (19.11.2002)

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language:  
English

Published:

— with international search report

(26) Publication Language:  
English

(88) Date of publication of the international search report:  
29 April 2004

(30) Priority Data:  
01811143.5 26 November 2001 (26.11.2001) EP

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(71) Applicant (for all designated States except US): **CIBA SPECIALTY CHEMICALS HOLDING INC.** [CH/CH]; Klybeckstrasse 141, CH-4057 Basel (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FREY, Markus** [CH/CH]; Alte Saline 12, CH-4310 Rheinfelden (CH). **RAST, Valérie** [CH/CH]; St.-Johanns-Parkweg 18, CH-4056 Basel (CH).

WO 2003/045919 A3

(54) Title: PROCESS FOR THE SYNTHESIS OF AMINE ETHERS FROM SECONDARY AMINO OXIDES AND USES

(57) Abstract: Amine ethers of sterically hindered amines are obtained in good yield from the corresponding N-oxyl hindered amine precursor by reaction with a hydrocarbon in the presence of an organic hydroperoxide and an iodide. The products of present process find utility as polymerization regulators and/or light stabilizers for organic material.

PROCESS FOR THE SYNTHESIS OF AMINE ETHERS FROM SECONDARY AMINO  
OXIDES

The instant invention pertains to a process for preparing amine ethers, e.g. N-hydrocarbyloxy substituted hindered amine compounds, by the reaction of the corresponding N-oxyl intermediate with a hydrocarbon in presence of an organic hydroperoxide and an iodide catalyst.

4-Hydroxy-1-oxyl-2,2,6,6-tetramethylpiperidine and 4-oxo-1-oxyl-2,2,6,6-tetramethylpiperidine are described as scavengers for some carbon centered radicals (S. Nigam et al., J. Chem. Soc., Trans. Faraday Soc., 1976, (72), 2324 and by K.-D. Asmus et al., Int. J. Radiat. Biol., 1976, (29), 211).

D. H. R. Barton et al., Tetrahedron, 1996, (52), 10301 describe the formation of some N-alkoxy-2,2,6,6-tetramethylpiperidine derivatives in the reaction of hydrocarbons with iron(II) and iron(III) species, hydrogen peroxide and various coadditives in the presence of N-oxyl-2,2,6,6-tetramethylpiperidine (TEMPO).

United States Patent No 5,374,729 describes a process for the preparation of N-methoxy derivatives of hindered amines from the reaction of the corresponding N-oxyl compound with methyl radicals produced from dimethyl sulfoxide by decomposing aqueous hydrogen peroxide in presence of a metal salt or by thermal decomposition of di-tert.butyl peroxide.

United States Patent No. 4,921,962 describes a process for the formation of N-hydrocarbyloxy derivatives of sterically hindered amines in which a hindered amine or N-oxyl substituted hindered amine is reacted with a hydrocarbon solvent in the presence of a hydroperoxide and a molybdenum catalyst.

It has now been found that N-hydrocarbyloxy substituted sterically hindered amines can most suitably be prepared from the N-oxyl intermediate and a hydrocarbon in presence of an organic hydroperoxide and an iodide catalyst. The process of the invention uses only catalytic quantities of iodide and does not require high temperatures.

Thus, present invention pertains to a process for the preparation of an amine ether


- 2 -

of a sterically hindered amine by reacting a corresponding sterically hindered aminoxide with an aliphatic hydrocarbon compound, characterized in that the reaction is carried out in the presence of an organic hydroperoxide and an iodide, which is preferably used in a catalytic amount.

The aliphatic hydrocarbon compound may be any compound selected from alkane, alkene, alkyne, or cyclic or polycyclic analogues thereof, and optionally may be substituted, e.g. by aryl, halogen, alkoxy etc., provided that an aliphatic CH (or CH<sub>2</sub>, CH<sub>3</sub>) moiety is contained.

Advantageously, the process of the invention is carried out in the absence of a copper or a copper compound, preferably in the absence of any heavy metal or heavy metal compound. Heavy metal is to be understood as transition metal or any metal of higher molecular weight than calcium. Metal compounds, the presence of which is advantageously to be avoided in the present process, include any form like salts, complexes, solutions and dispersions thereof. The amounts of these compounds to be tolerated within the process of the invention are preferably well below the catalytic level, e.g. below 0.0001 molar equivalent per mole of nitroxyl moiety, more preferably within or below the ppm-level (up to 1000 parts by weight of heavy metal per 1 million parts by weight of total reaction mixture).

Preferred is a process for the preparation of an amine ether of the formula A



wherein



a is 1 or 2;

when a is 1, E is E'

when a is 2, E is L;


E' is C<sub>1</sub>-C<sub>36</sub> alkyl; C<sub>3</sub>-C<sub>18</sub> alkenyl; C<sub>2</sub>-C<sub>18</sub> alkinyl; C<sub>5</sub>-C<sub>18</sub> cycloalkyl; C<sub>5</sub>-C<sub>18</sub> cycloalkenyl; a radical of a saturated or unsaturated aliphatic bicyclic or tricyclic hydrocarbon of 7 to 12 carbon atoms; C<sub>2</sub>-C<sub>7</sub>alkyl or C<sub>3</sub>-C<sub>7</sub>alkenyl substituted by halogen, C<sub>1</sub>-C<sub>8</sub>alkoxy or phenoxy;

$C_4$ - $C_{12}$ heterocycloalkyl;  $C_4$ - $C_{12}$ heterocycloalkenyl;  $C_7$ - $C_{15}$  aralkyl or  $C_4$ - $C_{12}$ heteroaralkyl, each of which is unsubstituted or substituted by  $C_1$ - $C_4$  alkyl or phenyl; or  $E'$  is a radical of formula (VII) or (VIII)



$\text{Ar}$  is  $C_6$ - $C_{10}$ aryl or  $C_5$ - $C_9$ heteroaryl;

$X$  is phenyl, naphthyl or biphenyl, which are substituted by 1, 2, 3 or 4 D and optionally further substituted by  $\text{NO}_2$ , halogen, amino, hydroxy, cyano, carboxy,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylamino or di( $C_1$ - $C_4$ alkyl)amino;

D is a group  , a group  $\text{C}(\text{O})-\text{G}_{13}$  or a group  $\text{C}(\text{O})-\text{G}_9-\text{C}(\text{O})-\text{G}_{13}$ ;

$G_1$  and  $G_2$ , independently of each other, are hydrogen, halogen,  $\text{NO}_2$ , cyano,  $-\text{CONR}_5\text{R}_6$ ,  $-(\text{R}_9)\text{COOR}_4$ ,  $-\text{C}(\text{O})-\text{R}_7$ ,  $-\text{OR}_8$ ,  $-\text{SR}_8$ ,  $-\text{NHR}_8$ ,  $-\text{N}(\text{R}_{18})_2$ , carbamoyl, di( $C_1$ - $C_{18}$ alkyl)carbamoyl,  $-\text{C}(\text{=NR}_5)(\text{NHR}_6)$ ,  $C_1$ - $C_{18}$ alkyl;  $C_3$ - $C_{18}$ alkenyl;  $C_3$ - $C_{18}$ alkinyl,  $C_7$ - $C_9$ phenylalkyl,  $C_3$ - $C_{12}$ cycloalkyl or  $C_2$ - $C_{12}$ heterocycloalkyl;  $C_1$ - $C_{18}$ alkyl or  $C_3$ - $C_{18}$ alkenyl or  $C_3$ - $C_{18}$ alkinyl or  $C_7$ - $C_9$ phenylalkyl,  $C_3$ - $C_{12}$ cycloalkyl or  $C_2$ - $C_{12}$ heterocycloalkyl substituted by OH, halogen,  $\text{NO}_2$ , amino, cyano, carboxy,  $\text{COOR}_{21}$ ,  $\text{C}(\text{O})-\text{R}_{22}$ ,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylamino or di( $C_1$ - $C_4$ alkyl)amino or a group  $-\text{O}-\text{C}(\text{O})-\text{R}_7$ ;  $C_2$ - $C_{18}$ alkyl which is interrupted by at least one O atom and/or  $\text{NR}_5$  group; or are  $C_6$ - $C_{10}$ aryl; or phenyl or naphthyl which are substituted by  $C_1$ - $C_4$ alkyl,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ alkylthio, halogen, cyano, hydroxy, carboxy,  $\text{COOR}_{21}$ ,  $\text{C}(\text{O})-\text{R}_{22}$ ,  $C_1$ - $C_4$ alkylamino or di( $C_1$ - $C_4$ alkyl)amino; or  $G_1$  and  $G_2$  together with the linking carbon atom form a  $C_3$ - $C_{12}$ cycloalkyl radical;

$G_5$  and  $G_6$  are independently of each other H or  $\text{CH}_3$ ;

$G_9$  is  $C_1$ - $C_{12}$ alkylene or a direct bond;

$G_{13}$  is  $C_1$ - $C_{18}$ alkyl;


$G_{14}$  is  $C_1$ - $C_{18}$ alkyl,  $C_5$ - $C_{12}$ cycloalkyl, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms, an acyl radical of a cyclo-aliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, or acyl radical of an aromatic acid containing 7 to 15 carbon atoms;

$G_{55}$  is H,  $\text{CH}_3$  or phenyl;

$G_{66}$  is  $-\text{CN}$  or a group of the formula  $-\text{COOR}_4$  or  $-\text{CONR}_5\text{R}_6$  or  $-\text{CH}_2-\text{O}-\text{G}_{14}$ ;

L is alkylene of 1 to 18 carbon atoms, cycloalkylene of 5 to 8 carbon atoms, cycloalkenylene of 5 to 8 carbon atoms, alkenylene of 3 to 18 carbon atoms, alkylene of 1 to 12 carbon atoms substituted by phenyl or by phenyl substituted by alkyl of 1 to 4 carbon atoms; or is alkylene of 4 to 18 carbon atoms interrupted by COO and/or phenylene;

T' is tertiary C<sub>4</sub>-C<sub>18</sub>alkyl or phenyl, each of which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>; or T' is C<sub>5</sub>-C<sub>12</sub>cycloalkyl; C<sub>5</sub>-C<sub>12</sub>cycloalkyl which is interrupted by at least one O or -NR<sub>18</sub>-; a polycyclic alkyl radical having 7-18 carbon atoms, or the same radical which is interrupted by at least one O or -NR<sub>18</sub>-; or T' is -C(G<sub>1</sub>)(G<sub>2</sub>)-T"; or C<sub>1</sub>-C<sub>18</sub>alkyl



or C<sub>5</sub>-C<sub>12</sub>cycloalkyl substituted by ;

T" is hydrogen, halogen, NO<sub>2</sub>, cyano, or is a monovalent organic radical comprising 1-50 carbon atoms;

or T" and T' together form a divalent organic linking group completing, together with the hindered amine nitrogen atom and the quaternary carbon atom substituted by G<sub>1</sub> and G<sub>2</sub>, an optionally substituted five- or six-membered ring structure;

and

R<sub>4</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl, phenyl, an alkali metal cation or a tetraalkylammonium cation;

R<sub>5</sub> and R<sub>6</sub> are hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>2</sub>-C<sub>18</sub>alkyl which is substituted by hydroxy or, taken together, form a C<sub>2</sub>-C<sub>12</sub>alkylene bridge or a C<sub>2</sub>-C<sub>12</sub>-alkylene bridge interrupted by O or/and NR<sub>18</sub>;

R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>6</sub>-C<sub>10</sub>aryl;

R<sub>8</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>2</sub>-C<sub>18</sub>hydroxyalkyl;

R<sub>9</sub> is C<sub>1</sub>-C<sub>12</sub>alkylene or a direct bond;

R<sub>18</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl or phenyl, which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>;


R<sub>21</sub> is hydrogen, a alkali metal atom or C<sub>1</sub>-C<sub>18</sub>alkyl; and

R<sub>22</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl;

which process comprises

- 5 -

reacting a N-oxyl amine of formula B




with a compound of formula IV or V



in the presence of an organic hydroperoxide and a catalytic amount of an iodide.

More specifically, present invention pertains to a process for the preparation of an amine ether of the formula A




wherein

a is 1 or 2;

when a is 1, E is  $E'$

when a is 2, E is  $L$ ;

$E'$  is  $C_1-C_{36}$  alkyl;  $C_3-C_{18}$  alkenyl;  $C_2-C_{18}$  alkinyl;  $C_5-C_{18}$  cycloalkyl;  $C_5-C_{18}$  cycloalkenyl; a radical of a saturated or unsaturated aliphatic bicyclic or tricyclic hydrocarbon of 7 to 12 carbon atoms;  $C_2-C_7$  alkyl or  $C_3-C_7$  alkenyl substituted by halogen;  $C_7-C_{15}$  aralkyl or  $C_7-C_{15}$  aralkyl substituted by  $C_1-C_4$  alkyl or phenyl; or  $E'$  is a radical of formula (VII)



X is phenyl, naphthyl or biphenyl, which are substituted by 1, 2, 3 or 4 D and optionally further substituted by  $NO_2$ , halogen, amino, hydroxy, cyano, carboxy,  $C_1-C_4$  alkoxy,  $C_1-C_4$  alkylthio,  $C_1-C_4$  alkylamino or di( $C_1-C_4$  alkyl)amino;

- 6 -

D is a group  , a group C(O)-G<sub>13</sub> or a group C(O)-G<sub>9</sub>-C(O)-G<sub>13</sub>;

G<sub>1</sub> and G<sub>2</sub>, independently of each other, are hydrogen, halogen, NO<sub>2</sub>, cyano, -CONR<sub>5</sub>R<sub>6</sub>, -(R<sub>9</sub>)COOR<sub>4</sub>, -C(O)-R<sub>7</sub>, -OR<sub>8</sub>, -SR<sub>8</sub>, -NHR<sub>8</sub>, -N(R<sub>18</sub>)<sub>2</sub>, carbamoyl, di(C<sub>1</sub>-C<sub>18</sub>alkyl)carbamoyl, -C(=NR<sub>5</sub>)(NHR<sub>6</sub>), C<sub>1</sub>-C<sub>18</sub>alkyl; C<sub>3</sub>-C<sub>18</sub>alkenyl; C<sub>3</sub>-C<sub>18</sub>alkinyl, C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl; C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>3</sub>-C<sub>18</sub>alkenyl or C<sub>3</sub>-C<sub>18</sub>alkinyl or C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl substituted by OH, halogen, NO<sub>2</sub>, amino, cyano, carboxy, COOR<sub>21</sub>, C(O)-R<sub>22</sub>, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylamino or di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino or a group -O-C(O)-R<sub>7</sub>; C<sub>2</sub>-C<sub>18</sub>alkyl which is interrupted by at least one O atom and/or NR<sub>5</sub> group; or are C<sub>6</sub>-C<sub>10</sub>aryl; or phenyl or naphthyl which are substituted by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, halogen, cyano, hydroxy, carboxy, COOR<sub>21</sub>, C(O)-R<sub>22</sub>, C<sub>1</sub>-C<sub>4</sub>alkylamino or di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino; or G<sub>1</sub> and G<sub>2</sub> together with the linking carbon atom form a C<sub>3</sub>-C<sub>12</sub>cycloalkyl radical;


G<sub>5</sub> and G<sub>6</sub> are independently of each other H or CH<sub>3</sub>;

G<sub>9</sub> is C<sub>1</sub>-C<sub>12</sub>alkylene or a direct bond;

G<sub>13</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl;

L is alkylene of 1 to 18 carbon atoms, cycloalkylene of 5 to 8 carbon atoms, cycloalkenylene of 5 to 8 carbon atoms, alkenylene of 3 to 18 carbon atoms, alkylene of 1 to 12 carbon atoms substituted by phenyl or by phenyl substituted by alkyl of 1 to 4 carbon atoms;

T' is tertiary C<sub>4</sub>-C<sub>18</sub>alkyl or phenyl, each of which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>; or T' is C<sub>5</sub>-C<sub>12</sub>cycloalkyl; C<sub>5</sub>-C<sub>12</sub>cycloalkyl which is interrupted by at least one O or -NR<sub>18</sub>-; a polycyclic alkyl radical having 7-18 carbon atoms, or the same radical which is interrupted by at least one O or -NR<sub>18</sub>-; or T' is -C(G<sub>1</sub>)(G<sub>2</sub>)-T"; or C<sub>1</sub>-C<sub>18</sub>alkyl



or C<sub>5</sub>-C<sub>12</sub>cycloalkyl substituted by

T" is hydrogen, halogen, NO<sub>2</sub>, cyano, or is a monovalent organic radical comprising 1-50 carbon atoms;

or T" and T' together form a divalent organic linking group completing, together with the hindered amine nitrogen atom and the quaternary carbon atom substituted by G<sub>1</sub> and G<sub>2</sub>, an

- 7 -

optionally substituted five- or six-membered ring structure;  
and

$R_4$  is hydrogen,  $C_1$ - $C_{18}$ alkyl, phenyl, an alkali metal cation or a tetraalkylammonium cation;  
 $R_5$  and  $R_6$  are hydrogen,  $C_1$ - $C_{18}$ alkyl,  $C_2$ - $C_{18}$ alkyl which is substituted by hydroxy or, taken together, form a  $C_2$ - $C_{12}$ alkylene bridge or a  $C_2$ - $C_{12}$ -alkylene bridge interrupted by O or/and  $NR_{18}$ ;

$R_7$  is hydrogen,  $C_1$ - $C_{18}$ alkyl or  $C_6$ - $C_{10}$ aryl;

$R_8$  is hydrogen,  $C_1$ - $C_{18}$ alkyl or  $C_2$ - $C_{18}$ hydroxyalkyl;

$R_9$  is  $C_1$ - $C_{12}$ alkylene or a direct bond;


$R_{18}$  is  $C_1$ - $C_{18}$ alkyl or phenyl, which are unsubstituted or substituted by halogen, OH,  $COOR_{21}$  or  $C(O)-R_{22}$ ;

$R_{21}$  is hydrogen, a alkali metal atom or  $C_1$ - $C_{18}$ alkyl; and

$R_{22}$  is  $C_1$ - $C_{18}$ alkyl;

which process comprises

reacting a N-oxyl amine of formula B




with a hydrocarbon of formula IV or V




in the presence of an organic hydroperoxide and a catalytic amount of an iodide.

In particular, present invention pertains to a process for the synthesis of a hindered amine of formula I or II



- 8 -




wherein


$G_1$ ,  $G_2$ ,  $G_3$  and  $G_4$  independently of each other are  $C_1$ - $C_{18}$ alkyl;  $C_3$ - $C_{18}$ alkenyl;  $C_3$ - $C_{18}$ alkinyl;  $C_1$ - $C_{18}$ alkyl or  $C_3$ - $C_{18}$ alkenyl or  $C_3$ - $C_{18}$ alkinyl substituted by OH, halogen or a group  $-O-C(O)-R_5$ ;  $C_2$ - $C_{18}$ alkyl which is interrupted by at least one O atom and/or  $NR_5$  group; or are  $C_3$ - $C_{12}$ cycloalkyl; or  $C_6$ - $C_{10}$ aryl; or  $G_1$  and  $G_2$  and/or  $G_3$  and  $G_4$  together with the linking carbon atom form a  $C_3$ - $C_{12}$ cycloalkyl radical;

$a$  is 1 or 2;

when  $a$  is 1,  $E$  is  $E'$ , wherein  $E'$  is  $C_1$ - $C_{36}$  alkyl;  $C_2$ - $C_{18}$  alkenyl;  $C_2$ - $C_{18}$  alkinyl;  $C_5$ - $C_{18}$  cycloalkyl;  $C_5$ - $C_{18}$  cycloalkenyl; a radical of a saturated or unsaturated aliphatic bicyclic or tricyclic hydrocarbon of 7 to 12 carbon atoms;  $C_2$ - $C_7$ alkyl or  $C_3$ - $C_7$ alkenyl substituted by halogen;  $C_7$ - $C_{15}$  aralkyl or  $C_7$ - $C_{15}$  aralkyl substituted by  $C_1$ - $C_4$  alkyl or phenyl; or  $E'$  is a radical of formula (VII)



$X$  is phenyl, naphthyl or biphenyl, which are substituted by 1, 2, 3 or 4 D and optionally further substituted by  $NO_2$ , halogen, amino, hydroxy, cyano, carboxy,  $C_1$ - $C_4$ alkoxy,  $C_1$ - $C_4$ alkylthio,  $C_1$ - $C_4$ alkylamino or di( $C_1$ - $C_4$ alkyl)amino;

$D$  is a group , a group  $C(O)-G_{13}$  or a group  $C(O)-G_9-C(O)-G_{13}$ ;

when  $a$  is 2,  $E$  is  $L$ ;

$G_5$  and  $G_6$  are independently of each other H or  $CH_3$ ;

$G_9$  is  $C_1$ - $C_{12}$ alkylene or a direct bond;


$G_{13}$  is  $C_1$ - $C_{18}$ alkyl;

$L$  is alkylene of 1 to 18 carbon atoms, cycloalkylene of 5 to 8 carbon atoms, cycloalkenylene of 5 to 8 carbon atoms, alkenylene of 3 to 18 carbon atoms, alkylene of 1 to 12 carbon atoms substituted by phenyl or by phenyl substituted by alkyl of 1 to 4 carbon atoms;

T is a divalent organic radical required to complete formula I to form, together with the hindered amine nitrogen atom and the two quaternary carbon atoms substituted by G<sub>1</sub> and G<sub>2</sub> or G<sub>3</sub> and G<sub>4</sub>, a five- or six-membered ring structure;

T<sub>1</sub> is hydrogen, halogen, NO<sub>2</sub>, cyano, -(R<sub>9</sub>)COOR<sub>4</sub>, -(R<sub>9</sub>)C(O)-R<sub>7</sub>, -OR<sub>8</sub>, unsubstituted C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>2</sub>-C<sub>18</sub>alkenyl, C<sub>2</sub>-C<sub>18</sub>alkynyl, C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl; or T<sub>1</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>2</sub>-C<sub>18</sub>alkenyl, C<sub>2</sub>-C<sub>18</sub>alkynyl, C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl, which is substituted by NO<sub>2</sub>, halogen, hydroxy, cyano, carboxy, C<sub>1</sub>-C<sub>6</sub>alkanoyl, C<sub>1</sub>-C<sub>12</sub>alkoxy; or phenyl, naphthyl, which are unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, halogen, cyano, hydroxy, carboxy; or T<sub>1</sub> is a residue -CH<sub>2</sub>-O-R<sub>10</sub> or -CH<sub>2</sub>-NR<sub>18</sub>-R<sub>10</sub> or -C(=CH<sub>2</sub>)-R<sub>11</sub> or -C(=O)-R<sub>12</sub>;

T<sub>2</sub> is tertiary C<sub>4</sub>-C<sub>18</sub>alkyl or phenyl, which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>; or T<sub>2</sub> is C<sub>5</sub>-C<sub>12</sub>cycloalkyl; C<sub>5</sub>-C<sub>12</sub>cycloalkyl which is interrupted by at least one O; a polycyclic alkyl radical having 7-18 carbon atoms or the same radical which is interrupted by at least one O atom; or T<sub>2</sub> is -C(G<sub>1</sub>)(G<sub>2</sub>)-T<sub>1</sub>; or



R<sub>4</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl, phenyl, an alkali metal cation or a tetraalkylammonium cation;

R<sub>5</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>6</sub>-C<sub>10</sub>aryl

R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or phenyl;

R<sub>8</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>2</sub>-C<sub>18</sub>hydroxyalkyl;

R<sub>9</sub> is C<sub>1</sub>-C<sub>12</sub>alkylene or a direct bond;

R<sub>10</sub> is hydrogen, formyl, C<sub>2</sub>-C<sub>18</sub>alkylcarbonyl, benzoyl, C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>5</sub>-C<sub>12</sub>cycloalkyl, C<sub>5</sub>-C<sub>12</sub>cycloalkyl interrupted by O or NR<sub>18</sub>, or is benzyl or phenyl which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>;

R<sub>11</sub> is OH, C<sub>1</sub>-C<sub>18</sub>alkoxy, benzyloxy, O-C(O)-(C<sub>1</sub>-C<sub>18</sub>)alkyl, N(R<sub>18</sub>)<sub>2</sub>, or a group C(O)R<sub>25</sub>;

R<sub>12</sub> is OH, O(alkali-metal), C<sub>1</sub>-C<sub>18</sub>alkoxy, benzyloxy, N(R<sub>18</sub>)<sub>2</sub>;



R<sub>18</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>2</sub>-C<sub>18</sub>hydroxyalkyl;

R<sub>21</sub> is hydrogen, a alkali metal atom or C<sub>1</sub>-C<sub>18</sub>alkyl; and

R<sub>22</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl;

R<sub>25</sub> is OH, C<sub>1</sub>-C<sub>18</sub>alkoxy, benzyloxy, N(R<sub>18</sub>)<sub>2</sub>;

which process comprises  
reacting a N-oxyl hindered amine of formula III or IIIa



with a hydrocarbon of formula IV or V



in the presence of an organic hydroperoxide and a catalytic amount of an iodide.

In the context of the description of the present invention, the term alkyl comprises, for example, methyl, ethyl and the isomers of propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl and dodecyl. Examples of aryl-substituted alkyl (aralkyl) are benzyl,  $\alpha$ -methylbenzyl or cumyl. Examples of alkoxy are methoxy, ethoxy, propoxy, butoxy, octyloxy etc.. Examples of alkenyl are vinyl and especially allyl. Examples of alkylene including alkylidene are ethylene, n-propylene or 1,2-propylene.

Some examples of cycloalkyl are cyclobutyl, cyclopentyl, cyclohexyl, methylcyclopentyl, dimethylcyclopentyl and methylcyclohexyl.

Examples of aryl are phenyl and naphthyl. Examples of substituted aryl are methyl-, dimethyl-, trimethyl-, methoxy- or phenyl-substituted phenyl.

Some examples of an aliphatic carboxylic acid are acetic, propionic, butyric, stearic acid. An example of a cycloaliphatic carboxylic acid is cyclohexanoic acid. An example of an aromatic carboxylic acid is benzoic acid. An example of a phosphorus-containing acid is methylphosphonic acid. An example of an aliphatic dicarboxylic acid is malonyl, maleoyl or


- 11 -

succinyl, or sebacic acid. An example of a residue of an aromatic dicarboxylic acid is phthaloyl.

A group heterocycloalkyl or heterocycloalkenyl embraces one or two heteroatoms, and a group heteroaryl from one to four heteroatoms, the heteroatoms being preferably selected from the group consisting of nitrogen, sulfur and oxygen. Some examples of heterocycloalkyl are tetrahydrofuryl, pyrrolidinyl, piperazinyl and tetrahydrothienyl. Some examples of heteroaryl are furyl, thienyl, pyrrolyl, pyridyl and pyrimidinyl.  $C_2$ - $C_{12}$ heterocycloalkyl is typically oxirane, 1,4-dioxane, tetrahydrofuran,  $\gamma$ -butyrolactone,  $\varepsilon$ -caprolactam, oxirane, aziridine, diaziridine, pyrrole, pyrrolidine, thiophen, furan, pyrazole, imidazole, oxazole, oxazolidine, thiazole, pyran, thiopyran, piperidine or morpholine.


An example of a monovalent silyl radical is trimethylsilyl.

Polycyclic alkyl radicals which may also be interrupted by at least one oxygen or nitrogen atom are for example adamantane, cubane, twistane, norbornane, bycyclo[2.2.2]octane bycyclo[3.2.1]octane, hexamethylentetramine (urotropine) or a group



Acyl radicals of monocarboxylic acids are, within the definitions, a residue of the formula  $-CO-R'$ , wherein  $R'$  may stand inter alia for an alkyl, alkenyl, cycloalkyl or aryl radical as defined. Preferred acyl radicals include acetyl, benzoyl, acryloyl, methacryloyl, propionyl, butyryl, valeroyl, hexanoyl, heptanoyl, octanoyl, nonanoyl, decanoyl, undecanoyl, dodecanoyl, pentadecanoyl, stearoyl. Polyacyl radicals of polyvalent acids are of the formula  $(-CO)_n-R'$ , wherein  $n$  is the valency, e.g. 2, 3, 4, 5 or 6. Some preferred examples for such residues are given elsewhere.

In preferred products of the instant process,  $E'$  is selected from the group consisting of




$-CH_2CH=CH_2$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-}(C_1\text{-}C_{12})\text{alkyl}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-}(C_6\text{-}C_{10})\text{aryl}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-}(C_1\text{-}C_{12})\text{alkoxy}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-phenoxy}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-}N\text{-di}(C_1\text{-}C_{12})\text{alkyl}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}CO\text{-}NH(C_1\text{-}C_{12})\text{alkyl}$ ,  $(C_1\text{-}C_{12})\text{alkyl-}CR_{30}\text{-}C(O)\text{-}N\text{-di}(C_1\text{-}C_{12})\text{alkyl}$

- 12 -

$\text{CR}_{30}\text{-CO-NH}_2$ ,  $-\text{CH}_2\text{CH=CH-CH}_3$ ,  $-\text{CH}_2\text{-C(CH}_3\text{)=CH}_2$ ,  $-\text{CH}_2\text{-CH=CH-phenyl}$ ,  $-\text{CH}_2\text{-C}\equiv\text{CH}$ ,

$(\text{C}_1\text{-C}_{12})\text{alkyl-}\text{CR}_{30}\text{-CN}$ ,



, wherein

$\text{R}_{30}$  is hydrogen or  $\text{C}_1\text{-C}_{12}\text{alkyl}$ ;

the aryl groups are phenyl or naphthyl, which are unsubstituted or substituted with  $\text{C}_1\text{-C}_{12}\text{alkyl}$ , halogen,  $\text{C}_1\text{-C}_{12}\text{alkoxy}$ , formyl,  $\text{C}_2\text{-C}_{12}\text{alkylcarbonyl}$ , glycidyloxy, OH, -COOH or -COOC $\text{C}_1\text{-C}_{12}\text{alkyl}$ . More preferably  $\text{E}'$  is selected from the group consisting of -CH<sub>2</sub>-phenyl, CH<sub>3</sub>CH-phenyl, (CH<sub>3</sub>)<sub>2</sub>C-phenyl, (C<sub>5</sub>-C<sub>6</sub>cycloalkyl)<sub>2</sub>CCN, (CH<sub>3</sub>)<sub>2</sub>CCN, -CH<sub>2</sub>CH=CH<sub>2</sub>, CH<sub>3</sub>CH-CH=CH<sub>2</sub> (C<sub>1</sub>-C<sub>8</sub>alkyl)CR<sub>30</sub>-C(O)-phenyl, (C<sub>1</sub>-C<sub>8</sub>)alkyl- $\text{CR}_{30}\text{-C(O)-(C}_1\text{-C}_8\text{)alkoxy}$ , (C<sub>1</sub>-C<sub>8</sub>)alkyl- $\text{CR}_{30}\text{-C(O)-(C}_1\text{-C}_8\text{)alkyl}$ , (C<sub>1</sub>-C<sub>8</sub>)alkyl- $\text{CR}_{30}\text{-C(O)-N-di(C}_1\text{-C}_8\text{)alkyl}$ , (C<sub>1</sub>-C<sub>8</sub>)alkyl- $\text{CR}_{30}\text{-C(O)-NH(C}_1\text{-C}_8\text{)alkyl}$ , (C<sub>1</sub>-C<sub>8</sub>)alkyl- $\text{CR}_{30}\text{-C(O)-NH}_2$ , (C<sub>1</sub>-C<sub>12</sub>)alkyl- $\text{CR}_{30}\text{-CN}$ , wherein  $\text{R}_{30}$  is hydrogen or (C<sub>1</sub>-C<sub>8</sub>)alkyl.

$\text{G}_1$  and  $\text{G}_2$  and/or  $\text{G}_3$  and  $\text{G}_4$  forming, together with the linking carbon atom, a C<sub>3</sub>-C<sub>12</sub>cycloalkyl radical, preferably form a C<sub>5</sub>-C<sub>12</sub>cycloalkyl radical, especially cyclopentylene, cyclohexylene or cycloheptylene.

$\text{G}_1$ ,  $\text{G}_2$ ,  $\text{G}_3$  and  $\text{G}_4$  independently are preferably alkyl of 1 to 4 carbon atoms, or the adjacent radicals  $\text{G}_1$  and  $\text{G}_2$  and/or  $\text{G}_3$  and  $\text{G}_4$  together are pentamethylene. More preferably,  $\text{G}_1$ ,  $\text{G}_2$ ,  $\text{G}_3$  and  $\text{G}_4$  independently are methyl or ethyl or propyl, especially methyl or ethyl. In the products most preferred,  $\text{G}_1$  and  $\text{G}_3$  are each methyl while  $\text{G}_2$  and  $\text{G}_4$  independently are methyl, ethyl or propyl.

T usually is an organic linking group containing 2-500 carbon atoms and forming, together with the carbon atoms it is directly connected to and the nitrogen atom, a substituted, 5-, 6 or 7-membered cyclic ring structure; T is preferably a C<sub>2</sub>-C<sub>500</sub>hydrocarbon optionally containing 1-200 hetero atoms selected from nitrogen, oxygen, phosphorus, sulfur,

silicon and halogen, T therein can be part of a 6-membered cyclic ring structure. More


preferably, T is an organic linking group of the formula  $E_1$    $E_2$  (VI), wherein

$E_2$  is  $-\text{CO-}$  or  $-(\text{CH}_2)_b-$ , while b is 0, 1 or 2;

$E_1$  is a carbon atom carrying the two residues  $R_{24}$  and  $R_{25}$ , or is  $>\text{N-}R_{25}$ , or is oxygen, and  $R_{24}$  and  $R_{25}$  are hydrogen or an organic residue, characterized in that the linking group T in total contains 2-500 carbon atoms and forms, together with the carbon atoms it is directly connected to it and the nitrogen atom, a substituted, 5-, 6 or 7-membered cyclic ring structure, or wherein  $R_{24}$  and  $R_{25}$  together are  $=\text{O}$  or wherein  $R_{24}$  is hydrogen and  $R_{25}$  is hydrogen or hydroxy. T is most preferably 2-hydroxy-1,3-propanediyl or 2-oxo-1,3-propanediyl.

Preferred products of the formula (I) are those wherein  $G_1$ ,  $G_2$ ,  $G_3$  and  $G_4$ , independently of each other, are methyl, ethyl, phenyl or  $\text{COOR}_4$ ;

E is a carbon centered radical formed from a  $\text{C}_7\text{-C}_{11}$ phenylalkane or a  $\text{C}_6\text{-C}_{10}$ pyridylalkane; or  $\text{C}_5\text{-C}_{12}$ cycloalkane; or  $\text{C}_5\text{-C}_{12}$ cycloalkene; or an oxacyclohexane or oxycyclohexene; or  $\text{C}_3\text{-C}_8$ alkene; or  $\text{C}_3\text{-C}_8$ alkene substituted by phenoxy; or a benzene which is substituted by  $\text{C}_1\text{-C}_4$ alkyl and a further substituent selected from  $\text{C}_1\text{-C}_4$ alkoxy, glycidyl or glycidyloxy; or E is a radical of formula (VIII)



Ar is  $\text{C}_6\text{-C}_{10}$ aryl or  $\text{C}_5\text{-C}_9$ heteroaryl;

$G_{14}$  is  $\text{C}_1\text{-C}_4$ alkyl or an acyl radical of an aliphatic carboxylic acid containing 2 to 4 carbon atoms or benzoyl;

$G_{55}$  is H,  $\text{CH}_3$  or phenyl;

$G_{66}$  is  $-\text{CN}$  or a group of the formula  $-\text{COOR}_4$  or  $-\text{CH}_2\text{-O-}G_{14}$ ;

$R_4$  is hydrogen or  $\text{C}_1\text{-C}_8$ alkyl;

L is a carbon centered radical formed from propane, butane, pentane, 2,2-dimethyl-propane, xylene; and

- 14 -





T is phenylene or an organic linking group of the formula  $E_1 E_2$  (VI), wherein

$E_2$  is  $-CO-$  or  $-(CH_2)_b-$ , while  $b$  is 0, 1 or 2;


$E_1$  is a carbon atom carrying the two residues  $R_{24}$  and  $R_{25}$ , or is  $>N-R_{25}$ , or is oxygen, and  $R_{24}$  and  $R_{25}$  are hydrogen or an organic residue, characterized in that the linking group T in total contains 2-500 carbon atoms and forms, together with the carbon atoms it is directly connected to it and the nitrogen atom, a substituted, 5-, 6 or 7-membered cyclic ring structure, or wherein  $R_{24}$  and  $R_{25}$  together are  $=O$  or wherein  $R_{24}$  is hydrogen and  $R_{25}$  is hydrogen or hydroxy;

or  $E_1$  and  $E_2$  together are 1,2-phenylene.

The product of formula A most preferably corresponds to one of the formulae



- 15 -



wherein

$G_1$ ,  $G_2$ ,  $G_3$  and  $G_4$  independently of each other are  $C_1$ - $C_{18}$ alkyl;  $C_3$ - $C_{18}$ alkenyl;  $C_3$ - $C_{18}$ alkinyl;  $C_1$ - $C_{18}$ alkyl or  $C_3$ - $C_{18}$ alkenyl or  $C_3$ - $C_{18}$ alkinyl substituted by OH, halogen or a group  $-O-C(O)-R_5$ ;  $C_2$ - $C_{18}$ alkyl which is interrupted by O;  $C_5$ - $C_{12}$ cycloalkyl; or phenyl; or  $G_1$  and  $G_2$  and/or  $G_3$  and  $G_4$  together with the linking carbon atom form a  $C_5$ - $C_{12}$ cycloalkyl radical;

$Z_1$  is O or  $NR_8$ ;

$R_8$  is hydrogen, OH,  $C_1$ - $C_{18}$ alkyl,  $C_3$ - $C_{18}$ alkenyl,  $C_3$ - $C_{18}$ alkinyl,  $C_1$ - $C_{18}$ alkyl,  $C_3$ - $C_{18}$ alkenyl,  $C_3$ - $C_{18}$ alkinyl which are substituted by one or more OH, halogen or a group  $-O-C(O)-R_5$ ,  $C_2$ - $C_{18}$ alkyl which is interrupted by at least one O atom and/or  $NR_5$  group,  $C_3$ - $C_{12}$ cycloalkyl or  $C_6$ - $C_{10}$ aryl,  $C_7$ - $C_9$ phenylalkyl,  $C_5$ - $C_{10}$ heteroaryl,  $-C(O)-C_1-C_{18}$ alkyl,  $-O-C_1-C_{18}$ alkyl or  $-COOC_1-C_{18}$ alkyl;

$Q$  is a direct bond or a divalent radical  $CR_9R_{10}$ ,  $CR_9R_{10}-CR_{11}R_{12}$ ,  $CR_9R_{10}CR_{11}R_{12}CR_{13}R_{14}$ ,  $C(O)$  or  $CR_9R_{10}C(O)$ ;

$R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$  and  $R_{14}$  are independently hydrogen, phenyl, or  $C_1$ - $C_{18}$ alkyl;

$T$  is  $CH_2-C(R_{24})(R_{25})-CH_2$ , wherein  $R_{24}$  and  $R_{25}$  together are  $=O$  or independently are H, OH or an organic residue, characterized in that the linking group  $T$  in total contains 2-500 carbon atoms and optionally 1-200 hetero atoms selected from, oxygen, phosphorus, sulfur, silicon, halogen and tertiary nitrogen.

The sterically hindered aminoxides, also referred to as N-oxyl educts for the instant process

which include compounds of formulae B, III or IIIa, are largely known in the art; they may be prepared by oxidation of the corresponding N-H hindered amine with a suitable oxygen donor, e.g. by the reaction of the corresponding N-H hindered amine with hydrogen peroxide and sodium tungstate as described by E. G. Rozantsev et al., in *Synthesis*, 1971, 192; or with tert-butyl hydroperoxide and molybdenum (VI) as taught in United States Patent No. 4,691,015, or obtained in analogous manner.

The preferred amount of hydrocarbon for the instant process depends to some extent on the relative number of reactive hydrogens on the hydrocarbon reactant and the hindered amine nitroxyl compound. The reaction is typically carried out with a ratio of 1 to 100 moles of hydrocarbon per mole of nitroxyl moiety with the preferred ratio being 1 to 50 moles per mole of nitroxyl moiety, and the most preferred ratio being 1 to 30 moles of hydrocarbon per mole of nitroxyl moiety.

The preferred amount of organic hydroperoxide is 1 to 20 moles per mole of nitroxyl moiety, with the more preferred amount being 1 to 5 moles of peroxide per mole of nitroxyl moiety and the most preferred amount being 1 to 3 moles of peroxide per mole of nitroxyl moiety.

The organic hydroperoxide used in the process of present invention can be of the formula R-OOH, wherein R usually is a hydrocarbon containing 1-18 carbon atoms. The organic hydroperoxide preferably is a peroxyalcohol containing 3-18 carbon atoms. R is often aliphatic, preferably C<sub>1</sub>-C<sub>12</sub>alkyl. Most preferred organic hydroperoxide is tert.butyl hydroperoxide.


The preferred amount of iodide catalyst is from about 0.0001 to 0.5, especially 0.0005 to 0.1 molar equivalent per mole of nitroxyl moiety, with a ratio of 0.001 to 0.05 moles of iodide per mole of nitroxyl moiety being the most preferred.

The reaction is preferably run at 0° to 100°C; more preferably at 20° to 100°C, especially in the range 20-80°C.

More specifically, the instant process involves the reaction of a mixture of 1 to 100 moles of the hydrocarbon, e.g. of formula IV or V, 1 to 20 moles of organic hydroperoxide, and 0.001 mmoles to 0.5 moles of iodide catalyst per mole of N-oxyl compound, such as the compound

of formula B (1 mmol is 0.001 mol). Preferably, the molar ratio of iodide catalyst per mole of N-oxyl compound is in the range from 1:100 to 1:100000, especially 1:300 to 1:100000.

E is preferably a carbon centered radical formed from a C<sub>7</sub>-C<sub>11</sub>phenylalkane or a C<sub>6</sub>-C<sub>10</sub>pyridylalkane; or C<sub>5</sub>-C<sub>12</sub>cycloalkane; or C<sub>5</sub>-C<sub>12</sub>cycloalkene; or an oxacyclohexane or oxycyclohexene; or C<sub>3</sub>-C<sub>8</sub>alkene; or C<sub>3</sub>-C<sub>8</sub>alkene substituted by phenoxy; or a benzene which is substituted by C<sub>1</sub>-C<sub>4</sub>alkyl and a further substituent selected from C<sub>1</sub>-C<sub>4</sub>alkoxy, glycidyl or glycidyloxy; or E is a radical of formula (VIII)



Ar is C<sub>6</sub>-C<sub>10</sub>aryl or C<sub>5</sub>-C<sub>9</sub>heteroaryl;

G<sub>14</sub> is C<sub>1</sub>-C<sub>4</sub>alkyl or an acyl radical of an aliphatic carboxylic acid containing 2 to 4 carbon atoms or benzoyl;

G<sub>55</sub> is H, CH<sub>3</sub> or phenyl;

G<sub>66</sub> is -CN or a group of the formula -COOR<sub>4</sub> or -CH<sub>2</sub>-O-G<sub>14</sub>;

R<sub>4</sub> is hydrogen or C<sub>1</sub>-C<sub>8</sub>alkyl;

L is a carbon centered radical formed from propane, butane, pentane, 2,2-dimethyl-propane, xylene.

Important are those educts, which are pure hydrocarbons.

The educt hydrocarbon, such as compound of formula IV or V, may serve two functions both as reactant and as solvent for the reaction. The reaction can also be carried out using an inert organic or inorganic solvent. A mixture of products may result if the hydrocarbon contains non-equivalent carbon-hydrogen bonds which are reactive in the instant process. For example, cyclohexane can give only one product whereas isopentane can give three distinct reaction products.

Usually the hydrocarbon reactand, e.g. compound of formula IV or V, reacts with its most active aliphatic carbon-hydrogen bond.

A solvent may be used, especially if the hydrocarbon, such as the compound of of formula IV or V, is a solid at the temperature of the reaction or if the catalyst is not very soluble in the

hydrocarbon. Inert solvents should have less active carbon-hydrogen bonds; typical inert solvents are acetonitrile, aromatic hydrocarbons like benzene, chlorobenzene,  $\text{CCl}_4$ , alcohols (e.g. methanol, ethanol, ethylene glycol, ethylene glycol monomethyl ether), or, especially for reactions with activated hydrocarbons like alkylated aromatics or alkenes, also alkanes like hexane, decane etc., or mixtures thereof. Inorganic solvents such as water are possible as well. The reaction can be carried out in one liquid phase or in separate phases.

Good results can be achieved when phase transfer catalysts such as quaternary ammonium or phosphonium salts are used. For example, quaternary ammonium or phosphonium halogenides such as chlorides or bromides may be employed for this purpose. The structure of the ammonium or phosphonium cation is less important; usually, quaternary ammonium or phosphonium cations contain 4 hydrocarbon residues bonded to the central nitrogen or phosphorus atom, which may be, for example, alkyl, phenylalkyl or phenyl groups. Some readily available materials are tetra- $\text{C}_1\text{-C}_{12}$ alkylated.

The iodide catalyst may be selected from any iodide compound, including organic and inorganic iodide compounds. Examples are alkaline or alkaline earth metal iodides, or onium iodides such as ammonium or phosphonium or sulfonium iodides. Suitable metal iodides are, *inter alia*, those of lithium, sodium, potassium, magnesium or calcium.

Especially good results can be achieved when onium iodides are used which are soluble in organic solvents. Suitable onium iodides embrace quaternary ammonium, phosphonium or sulfonium iodides. The structure of the onium cation is less important provided the solubility in organic solvents is high enough; the latter can be increased by increasing the hydrophobicity of the hydrocarbon residues attached to the onium cation. Some readily available materials are tetra- $\text{C}_1\text{-C}_{12}$ alkylated ammonium iodides and/or the following compounds:

Tetrabutylammonium iodide;

Tetraoctylammonium iodide;

Tetra(hexamethylbenzyl)ammonium iodide;

Tetradodecylammonium iodide;

Tetrahexylammonium iodide;

Di-octadecyl-dimethyl-ammonium iodide;

Hexadecyl-benzyl-dimethyl-ammonium iodide;

Tributyl-methyl-ammonium iodide<sup>A)</sup>;

Di-tetradecyl-dimethyl-ammonium iodide;  
Trioctyl-propyl-ammonium iodide;  
Octyl-benzyl-dimethyl ammonium iodide;  
Trioctylmethylammonium iodide<sup>B)</sup>;  
Hexadecylpyridinium iodide;  
Dioctyl-dimethyl-ammonium iodide;  
Octyl-trimethylammonium iodide;  
Tetraethyl ammonium iodide;  
Dioctyl-methyl sulfonium iodide;  
Tetraphenylphosphonium iodide;  
Triphenyl-isopropyl phosphonium iodide;  
Triphenylethylphosphonium iodide;  
Triphenylhexyl phosphonium iodide;  
Tetrabutyl phosphonium iodide;  
Tributyl-hexadecyl phosphonium iodide;  
Tetraoctyl phosphonium iodide;  
Triphenylmethyl phosphonium iodide;  
Diphenyl-dimethyl-phosphonium iodide;  
Tetraethylphosphonium iodide;  
Phenyl-trimethyl-phosphonium iodide;  
Triphenyl-(CH<sub>2</sub>CO<sub>2</sub>CH<sub>3</sub>)phosphonium iodide;  
Triphenylbenzylphosphonium iodide.

A) iodide form of ALIQUAT® 175

B) iodide form of ALIQUAT® 336

In a preferred embodiment, the iodide catalyst functions the same time as a phase transfer catalyst, e.g. when a quaternary ammonium or phosphonium iodide such as tetrabutylammoniumiodide is used as catalyst. These compounds are known, many are commercially available.

The onium iodides can be generated from any other onium salt (e.g., hydroxide, sulfate, hydrogensulfate, fluoride, acetate, chloride, cyanide, bromide, nitrate, nitrite, perchlorate etc.) via insitu anion exchange using a watersoluble inorganic iodide such as alkaline or alkaline earth metal iodides, other iodine containing salts or elemental iodine. For example,

- 20 -

commercial onium chlorides of the ALIQUAT® series may conveniently be brought into the above iodide form by in situ anion exchange.

The onium iodides can be bound to an organic or inorganic polymer backbone, rendering a homogeneous or heterogenous catalytic system.

Preferably, the pH of the aqueous phase, if present, is held between 7 and 11, especially between 9 and 10, most preferably at 9 during the reaction.

Preferred are quaternary ammonium or phosphonium iodides, especially tetraalkyl ammonium iodides.

The instant process can be run in air or in an inert atmosphere such a nitrogen or argon. The instant process can be run under atmospheric pressure as well as under reduced or elevated pressure. Elevated pressure can especially be useful in reactions with a hydrocarbon, which is gaseous under atmospheric pressure and the reaction temperature; in this case, pressure/temperature conditions are advantageous where the hydrocarbon forms a liquid phase or is at least partially dissolved in a suitable solvent.

There are several variations of the instant process. One variation involves the addition of a solution of organic hydroperoxide to a mixture of the N-oxyl hindered amine, the hydrocarbon and cosolvent (if used), and catalyst which has been brought to the desired temperature for reaction. The proper temperature may be maintained by controlling the rate of peroxide addition and/or by using a heating or cooling bath. After the hydroperoxide is added, the reaction mixture is conveniently stirred till the starting N-oxyl, e.g. compound of formula III, has disappeared or is no longer being converted to the desired product, e.g. compound of formula I and/or II. The reaction can be monitored by methods known in the art such as UV-Vis spectroscopy, thin layer chromatography, gas chromatography or liquid chromatography. Additional portions of catalyst can be added while the reaction is in progress. After the initial hydroperoxide charge has been added to the reaction mixture, more hydroperoxide can be added dropwise to bring the reaction to completion.

A second variation of the instant process is to simultaneously add separate solutions of the hydroperoxide and the nitroxyl compound to a mixture of the hydrocarbon, cosolvent (if used) and catalyst. The nitroxyl compound may be dissolved in water or the

- 21 -


alcohol solvent used in the reaction. Some of the nitroxyl compound may be introduced into the reaction mixture prior to starting the peroxide addition, and all of the nitroxyl compound should be added prior to completing the peroxide addition.

Another variation of the instant process involves the simultaneous addition of separate solutions of the hydroperoxide and of the aqueous or alcohol solution of the catalyst to a mixture of the nitroxyl compound, hydrocarbon, and cosolvent (if used). Some of the metal may be introduced into the reaction mixture prior to starting the peroxide addition.


Still another variation of the instant process is the simultaneous addition of separate solutions of the hydroperoxide, of the aqueous or alcohol solution of the nitroxyl compound, and of an aqueous or alcohol solution of the catalyst to the hydrocarbon and cosolvent (if used). A portion of the nitroxyl compound and/or catalyst may be introduced into the reaction mixture prior to starting the hydroperoxide addition. All of the nitroxyl compound should be added prior to completing the hydroperoxide addition.

At the end of the reaction, the residual hydroperoxide should be carefully decomposed prior to the isolation of any products.

Examples for compounds which can be obtained advantageously with the process of present invention are those of formulae 1-28:




(1)




(2)


- 22 -




(3)




(4)



(5)




(6)




(7)


- 23 -




(8)



(9)




(10)




(11)


- 24 -



(12)







(13)




(14)


- 25 -




- 26 -




(19)




(20)



(21)




(22)




(23)


- 27 -




(24)



(25)




(26)



(27)

- 28 -



wherein in formulas (1) to (15):

m is 0 or 1;

R<sub>1</sub> is hydrogen, hydroxyl or hydroxymethyl;

R<sub>2</sub> is hydrogen, alkyl of 1 to 12 carbon atoms or alkenyl of 2 to 12 carbon atoms;

n is 1 to 4;

when n is 1,

R<sub>3</sub> is alkyl of 1 to 18 carbon atoms, alkoxy carbonyl alkylene carbonyl of 4 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, glycidyl, 2,3-dihydroxypropyl, 2-hydroxy or 2-(hydroxymethyl) substituted alkyl of 3 to 12 carbon atoms which alkyl is interrupted by oxygen, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms, an acyl radical of a cycloaliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, or acyl radical of an aromatic acid containing 7 to 15 carbon atoms;

when n is 2,

R<sub>3</sub> is alkylene of 2 to 18 carbon atoms, a divalent acyl radical of an aliphatic or unsaturated aliphatic dicarboxylic or dicarbamic acid containing 2 to 18 carbon atoms, a divalent acyl radical of a cycloaliphatic dicarboxylic or dicarbamic acid containing 7 to 12 carbon atoms, or a divalent acyl radical of an aromatic dicarboxylic acid containing 8 to 15 carbon atoms;

when n is 3,

$R_3$  is a trivalent acyl radical of an aliphatic or unsaturated aliphatic tricarboxylic acid containing 6 to 18 carbon atoms, or a trivalent acyl radical of an aromatic tricarboxylic acid containing 9 to 15 carbon atoms;

when n is 4,

$R_3$  is a tetravalent acyl radical of an aliphatic or unsaturated aliphatic tetracarboxylic acid, especially 1,2,3,4-butanetetracarboxylic acid, 1,2,3,4-but-2-enetetracarboxylic acid, 1,2,3,5-pantanetetracarboxylic acid and 1,2,4,5-pantanetetracarboxylic acid, or  $R_3$  is a tetravalent acyl radical of an aromatic tetracarboxylic acid containing 10 to 18 carbon atoms;

p is 1 to 3,

$R_4$  is hydrogen, alkyl of 1 to 18 carbon atoms or acyl of 2 to 6 carbon atoms;

when p is 1,

$R_5$  is hydrogen, alkyl of 1 to 18 carbon atoms, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms; an acyl radical of a cycloaliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, an acyl radical of an aromatic carboxylic acid containing 7 to 15 carbon atoms, or  $R_4$  and  $R_5$  together are  $-(CH_2)_5CO-$ , phthaloyl or a divalent acyl radical of maleic acid;

when p is 2,

$R_5$  is alkylene of 2 to 12 carbon atoms, a divalent acyl radical of an aliphatic or unsaturated aliphatic dicarboxylic or dicarbamic acid containing 2 to 18 carbon atoms, a divalent acyl radical of a cycloaliphatic dicarboxylic or dicarbamic acid containing 7 to 12 carbon atoms, or a divalent acyl radical of an aromatic dicarboxylic acid containing 8 to 15 carbon atoms;

when p is 3,

$R_5$  is a trivalent acyl radical of an aliphatic or unsaturated aliphatic tricarboxylic acid containing 6 to 18 carbon atoms, or a trivalent acyl radical of an aromatic tricarboxylic acid containing 9 to 15 carbon atoms;

when n is 1,

- 30 -

$R_6$  is alkoxy of 1 to 18 carbon atoms, alkenyloxy of 2 to 18 carbon atoms,  $-NHalkyl$  of 1 to 18 carbon atoms or  $-N(alkyl)_2$  of 2 to 36 carbon atoms,

when  $n$  is 2,

$R_6$  is alkylenedioxy of 2 to 18 carbon atoms, alkenylenedioxy of 2 to 18 carbon atoms,  $-NH-alkylene-NH-$  of 2 to 18 carbon atoms or  $-N(alkyl)-alkylene-N(alkyl)-$  of 2 to 18 carbon atoms, or  $R_6$  is 4-methyl-1,3-phenylenediamino,

when  $n$  is 3,

$R_6$  is a trivalent alkoxy radical of a saturated or unsaturated aliphatic triol containing 3 to 18 carbon atoms,


when  $n$  is 4,

$R_6$  is a tetravalent alkoxy radical of a saturated or unsaturated aliphatic tetraol containing 4 to 18 carbon atoms,

$R_7$  and  $R_8$  are independently chlorine, alkoxy of 1 to 18 carbon atoms,  $-O-T_1$ , amino substituted by 2-hydroxyethyl,  $-NH(alkyl)$  of 1 to 18 carbon atoms,  $-N(alkyl)T_1$  with alkyl of 1 to 18 carbon atoms, or  $-N(alkyl)_2$  of 2 to 36 carbon atoms,

$R_9$  is oxygen, or  $R_9$  is nitrogen substituted by either hydrogen, alkyl of 1 to 12 carbon atoms or  $T_1$


$T_1$  is



$R_{10}$  is hydrogen or methyl,

$q$  is 2 to 8,

$R_{11}$  and  $R_{12}$  are independently hydrogen or the group  $T_2$



$R_{13}$  is hydrogen, phenyl, straight or branched alkyl of 1 to 12 carbon atoms, alkoxy of 1 to 12 carbon atoms, straight or branched alkyl of 1 to 4 carbon atoms substituted by phenyl, cycloalkyl of 5 to 8 carbon atoms, cycloalkenyl of 5 to 8 carbon atoms, alkenyl of 2 to 12 carbon atoms, glycidyl, allyloxy, straight or branched hydroxyalkyl of 1 to 4 carbon atoms, or silyl or silyloxy substituted three times independently by hydrogen, by phenyl, by alkyl of 1 to 4 carbon atoms or by alkoxy of 1 to 4 carbon atoms;

$R_{14}$  is hydrogen or silyl substituted three times independently by hydrogen, by phenyl, by alkyl of 1 to 4 carbon atoms or by alkoxy of 1 to 4 carbon atoms;

$d$  is 0 or 1;

$h$  is 0 to 4;

$k$  is 0 to 5;

$x$  is 3 to 6;

$y$  is 1 to 10;


$z$  is an integer such that the compound has a molecular weight of 1000 to 4000 amu, e.g.  $z$  may be from the range 3-10;

- 32 -

$R_{15}$  is morpholino, piperidino, 1-piperizinyl, alkylamino of 1 to 8 carbon atoms, especially branched alkylamino of 3 to 8 carbon atoms such as tert-octylamino,  $-N(alkyl)T_1$  with alkyl of 1 to 8 carbon atoms, or  $-N(alkyl)_2$  of 2 to 16 carbon atoms,

$R_{16}$  is hydrogen, acyl of 2 to 4 carbon atoms, carbamoyl substituted by alkyl of 1 to 4 carbon atoms, s-triazinyl substituted once by chlorine and once by  $R_{15}$ , or s-triazinyl substituted twice by  $R_{15}$  with the condition that the two  $R_{15}$  substituents may be different;

$R_{17}$  is chlorine, amino substituted by alkyl of 1 to 8 carbon atoms or by  $T_1$ ,  $-N(alkyl)T_1$  with alkyl of 1 to 8 carbon atoms,  $-N(alkyl)_2$  of 2 to 16 carbon atoms, or the group  $T_3$



$R_{18}$  is hydrogen, acyl of 2 to 4 carbon atoms, carbamoyl substituted by alkyl of 1 to 4 carbon atoms, s-triazinyl substituted twice by  $-N(alkyl)_2$  of 2 to 16 carbon atoms or s-triazinyl substituted twice by  $-N(alkyl)T_1$  with alkyl of 1 to 8 carbon atoms;

in formulas (16) to (28),  $R_1$ ,  $R_2$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{13}$ ,  $R_{14}$ ,  $d$ ,  $h$ ,  $k$ ,  $m$ ,  $q$ , and  $T_1$  have the same meanings as in formulas (1) to (15);

$R_{19}$  is hydrogen, alkyl of 1 to 18 carbon atoms, alkenyl of 2 to 18 carbon atoms, glycidyl, 2,3-dihydroxypropyl, 2-hydroxy or 2-(hydroxymethyl) substituted alkyl of 3 to 12 carbon atoms which alkyl is interrupted by oxygen, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms, an acyl radical of a cycloaliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, or acyl radical of an aromatic acid containing 7 to 15 carbon atoms;

$R_{20}$  is alkylene of 2 to 18 carbon atoms, a divalent acyl radical of an aliphatic or unsaturated aliphatic dicarboxylic or dicarbamic acid containing 2 to 18 carbon atoms, a

divalent acyl radical of a cycloaliphatic dicarboxylic or dicarbamic acid containing 7 to 12 carbon atoms, or a divalent acyl radical of an aromatic dicarboxylic acid containing 8 to 15 carbon atoms;

$R_{21}$  is hydrogen, alkyl of 1 to 18 carbon atoms or acyl of 2 to 6 carbon atoms;

$R_{22}$  is hydrogen, alkyl of 1 to 18 carbon atoms, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms, an acyl radical of a cycloaliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, an acyl radical of an aromatic carboxylic acid containing 7 to 15 carbon atoms, or  $R_4$  and  $R_5$  together are  $-(CH_2)_5CO-$ , phthaloyl or a divalent acyl radical of maleic acid;

$R_{23}$  is hydrogen, alkyl of 1 to 4 carbon atoms or acyl of 2 to 6 carbon atoms;

$R_{24}$  is alkylene of 2 to 18 carbon atoms, a divalent acyl radical of an aliphatic or unsaturated aliphatic dicarboxylic or dicarbamic acid containing 2 to 18 carbon atoms, a divalent acyl radical of a cycloaliphatic dicarboxylic or dicarbamic acid containing 7 to 12 carbon atoms, or a divalent acyl radical of an aromatic dicarboxylic acid containing 8 to 15 carbon atoms;

$R_{25}$  is alkoxy of 1 to 18 carbon atoms, alkenyloxy of 2 to 18 carbon atoms,  $-NHalkyl$  of 1 to 18 carbon atoms or  $-N(alkyl)_2$  of 2 to 36 carbon atoms,

$R_{26}$  is alkyleneedioxy of 2 to 18 carbon atoms, alkenyleneedioxy of 2 to 18 carbon atoms,  $-NH-alkylene-NH-$  of 2 to 18 carbon atoms or  $-N(alkyl)-alkylene-N(alkyl)-$  of 3 to 18 carbon atoms.

$E$  is a carbon centered radical formed preferably from a  $C_7-C_{11}$ phenylalkane, especially toluene, ethylbenzene, isopropylbenzene; or  $C_5-C_{12}$ cycloalkane, especially cyclohexene; or  $C_5-C_{12}$ cycloalkene, especially cyclohexene; or  $C_3-C_8$ alkene, especially propene; or a benzene which is substituted by  $C_1-C_4$ alkyl and a further substituent selected from  $C_1-C_4$ alkoxy, glycidyl or glycidyloxy.

$L$  is a carbon centered radical formed preferably from propane, butane, pentane, 2,2-dimethyl-propane, xylene, diethylbenzene.

Preferably, the reaction site in the compound E-H or H-L-H is an activated carbon-hydrogen bond, whose carbon, for example, is linked to an electron pushing functional group or a functional group able to stabilize the radical formed after cleavage of the carbon-hydrogen bond. Electron withdrawing groups, if present in E-H or H-L-H, are preferably not directly linked to the reactive site.

Products of the present process can be employed with advantage for stabilizing organic material against the damaging effect of light, oxygen and/or heat, especially for stabilizing synthetic organic polymers or compositions containing them. They are notable for high thermal stability, substrate compatibility and good persistence in the substrate.

The compounds made by the instant process are particularly effective in the stabilization of polymer compositions against harmful effects of light, oxygen and/or heat; they are also useful as initiators or regulators for radical polymerization processes which provide homopolymers, random copolymers, block copolymers, multiblock copolymers, graft copolymers and the like, at enhanced rates of polymerization and enhanced monomer to polymer conversions.

Of particular interest is the use of products of the present process as stabilizers in synthetic organic polymers, for example a coating or a bulk polymer or article formed therefrom, especially in thermoplastic polymers and corresponding compositions as well as in coating compositions. Thermoplastic polymers of most importance in present compositions are polyolefins and their copolymers, thermoplastic polyolefin (TPO), thermoplastic polyurethane (TPU), thermoplastic rubber (TPR), polycarbonate, such as in item 19 above, and blends, such as in item 28 above. Of utmost importance are polyethylene (PE), polypropylene (PP), polycarbonate (PC) and polycarbonate blends such as PC/ABS blends, as well as in acid or metal catalyzed coating compositions.

In general the products of present invention may be added to the material to be stabilized in amounts of from 0.1 to 10 %, preferably from 0.01 to 5 %, in particular from 0.01 to 2 % (based on the material to be stabilized). Particular preference is given to the use of the novel compounds in amounts of from 0.05 to 1.5 %, especially from 0.1 to 0.5 %. Where compounds of present invention are used as flame retardants, dosages are usually higher,

e.g. 0.1 to 25 % by weight, mainly 0.1 to 10 % by weight of the organic material to be stabilized and protected against inflammation.

Used in polymerizable compositions as a polymerization regulator or initiator, preferably the regulator/initiator compound is present in an amount of from 0.01 mol-% to 30 mol-%, more preferably in an amount of from 0.1 mol-% to 20 mol-% and most preferred in an amount of from 0.5 mol-% to 10 mol-% based on the monomer or monomer mixture.

The following examples are for illustrative purposes only and are not to be construed to limit the instant invention in any manner whatsoever. Percentages given are usually percent by weight if not otherwise indicated. Abbreviations used:

min. minutes;

HPLC high pressure liquid chromatography;

GC gas chromatography;

Bu butyl;

Ph phenyl;

Me methyl;

Oct octyl;

Hex hexyl;

Et ethyl;


Bz benzyl;

Py 1-pyridinium;

TEMPO 2,2,6,6-tetramethylpiperidine-N-oxide;

eq. equivalent (of nitroxide, if not otherwise indicated).

**Example 1:** Preparation of the compound of formula




To a stirred mixture of 5g (32 mmol) 2,2,6,6-tetramethylpiperidine-N-oxide (TEMPO), 34 g (320 mmol) of ethylbenzene and 0.12 g (0.32 mmol) of tetrabutylammoniumiodide, 6.2 g (48 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 30 minutes. The temperature is maintained at 60°C for 25 minutes until all of the TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 61 g of an aqueous solution of Na<sub>2</sub>SO<sub>3</sub> (10%) until the disappearance of excess t-butylhydroperoxide. The aqueous phase


is then separated and washed with ethylbenzene. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered, and the solvent is distilled off on a rotary-evaporator. The crude product is purified by flash-chromatography (silica gel, hexane : ethylacetate 9 : 1), yielding 5 g (60 % of theory) of a yellow oil. Analysis required for  $\text{C}_{17}\text{H}_{27}\text{NO}$  (261.41): C 78.11%, H 10.41%, N 5.36%; found: C 78.04%, H 10.46%, N 5.26%.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 0.66 (broad s, 3H), 1.03-1.52 (m, 15H), 1.48 (d,  $J = 8\text{Hz}$ , 3H), 4.78 (q,  $J = 8\text{Hz}$ , 1H), 7.21-7.33 (m, 5H).

**Example 2:** Example 1 is repeated except that 32 mmol of 2,2,6,6-Tetramethylpiperidine-N-oxide are replaced by the equivalent amount of 2,2,6,6-Tetramethylpiperidine-4-one-N-oxide,

yielding a compound of formula




**Example 3:** Preparation of a compound of formula




A stirred mixture of 0.5g (3.2 mmol) TEMPO, 1.14 g (6.4 mmol) of 2-(4-ethyl-phenoxyethyl)-oxirane, 0.0118 g (0.032 mmol) of tetrabutylammoniumiodide and 0.62 g (4.8 mmol) of t-butylhydroperoxid (70% aqueous solution) is brought to 60°C. The temperature is maintained at 60°C for 4 hours until all of the TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 20g of a 10% aqueous  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with ethylbenzene. The combined organic phases are passed through a plug of silica gel, washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 0.9 g of a colorless oil. Quantitative HPLC-analysis reveals a product-concentration of 65% w/w, corresponding to an overall yield of 54.8%.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ),  $\delta$  (ppm; 2-(4-Ethyl-phenoxyethyl)-oxirane not shown): 0.63 (broad s, 3H), 1.01-1.56 (m, 15H), 1.45 (d,  $J = 8\text{Hz}$ , 3H), 2.75-2.76 (m, 1H), 2.89-2.91 (m, 1H), 3.34-3.36 (m, 1H), 3.95-3.99 (m, 1H), 4.17-4.21 (m, 1H), 4.73 (q,  $J = 8\text{Hz}$ , 1H), 6.84-6.88 (m, 2H), 7.21-7.26 (m, 2H).

Example 4: Preparation of the compound of formula




To a stirred mixture of 5g (32 mmol) TEMPO, 39.1 g (320 mmol) of phenetole and 0.12 g (0.32 mmol) of tetrabutylammoniumiodide, 12.37 g (96 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 60 minutes. The temperature is maintained at 60°C for 21 hours until all TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 121 g of a 10% aqueous Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over MgSO<sub>4</sub>, filtered, and the solvent is distilled off on a rotary-evaporator. The crude product is purified by flash-chromatography (silica gel, Hexane / Ethylacetate 9 / 1), yielding 4.6 g (51.8 % of theory) of a slightly yellow oil. Analysis required for C<sub>17</sub>H<sub>27</sub>NO<sub>2</sub> (277.41): C 73.61%, H 9.81%, N 5.05%; found: C 73.15%, H 9.89%, N 4.95%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm): 1.13 (s, 3H), 1.16 (s, 3H), 1.19 (s, 6H), 1.30-1.69 (m, 6H), 1.47 (d, J = 8Hz, 3H), 5.58 (q, J = 8Hz, 1H), 6.92-6.96 (m, 1H), 7.01-7.03 (m, 2H), 7.24-7.28 (m, 2H).

Example 5: Preparation of




To a stirred mixture of 50 mmol 4-propoxy-2,2,6,6-tetramethylpiperidine-1-oxyl, 41.1 g (500 mmol) of cyclohexene and 0.18 g (0.5 mmol) of tetrabutylammoniumiodide, 7.4 g (58 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 55°C within 30 minutes. The reaction mixture is cooled down to 25°C and stirred with 63 g of an aqueous 20% Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are passed through a plug of silica gel and washed with brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator. The crude product is purified by distillation, yielding the title product.

Example 6: Preparation of


by Hydrogenation of the

Product of Example 5


A mixture of 4 mmol) of the product of Example 5 and 0.2 g Pd on charcoal (10%) in 10 ml of methanol is hydrogenated at 25°C and 4 bar of hydrogen. Filtration and evaporation of the solvent yields the title product as a slightly orange oil.

Example 7: Preparation of the compound of the formula To a stirred mixture of 5.5 g (35

mmol) TEMPO, 10.5 g (70 mmol) of phenylacetic acid methyl ester and 0.13 g (0.35 mmol) of tetrabutylammoniumiodide, 6.75 g (52.5 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 25 minutes. The temperature is maintained at 60°C for 46 hours. The reaction mixture is cooled down to 25°C and stirred with 66 g of a 10% aqueous Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with ethylbenzene. The combined organic phases are washed with brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator. The crude product is purified by flash-chromatography (silica gel, hexane : ethylacetate 9 : 1), yielding 6 g (56 % of theory) of the title product as a white crystalline solid, mp 85°C - 87°C. Analysis required for C<sub>18</sub>H<sub>27</sub>NO<sub>3</sub> (305.42): C 70.79%, H 8.91%, N 4.59%; found: C 70.60%, H 9.13%, N 4.53%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm): 0.72 (s, 3H), 1.07 (s, 3H), 1.14 (s, 3H), 1.23 (s, 3H), 1.28 - 1.58 (m, 6H), 3.65 (s, 3H), 5.21 (s, 1H), 7.27 - 7.35 (m, 3H), 7.43 - 7.45 (d-like, 2H).

Example 8: Preparation of the compound of the formula

To a stirred mixture of 6.8 g (32 mmol) of 2,6-diethyl-2,3,6-trimethyl-piperidin-4-one-N-oxide, 34 g (320 mmol) of ethylbenzene and 0.12 g (0.32 mmol) of tetrabutylammoniumiodide, 6.2 g (48 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 30 minutes. The temperature is maintained at 60°C for 13 hours, after which another 6.2 g of t-butylhydroperoxid and 0.12 g of tetrabutylammoniumiodide are added. The temperature is maintained at 60°C for another 24 hours, cooled down to 25°C and stirred with 120 g of a 10% aqueous Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with ethylbenzene. The combined organic phases are washed with brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator. The crude product is purified by flash-chromatography (silica gel, hexane : Ethylacetate 9 : 1), yielding the title product as a yellow oil. Analysis required for C<sub>20</sub>H<sub>31</sub>NO<sub>2</sub> (317.48): C 75.67%, H 9.84%, N 4.41%; found: C 74.01%, H 9.76%, N 4.30%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm, O-CH only): 4.83 (p-like, 1H).


Example 9: Preparation of the compound of the formula

To a stirred mixture of 6.4 g (25 mmol) of 3,3,8,8,10,10-hexamethyl-1,5-dioxa-9-aza-spiro[5.5]undecane-N-oxide, 8.9 g (50 mmol) of 2-(4-ethyl-phenoxy)methyl-oxirane and 0.09 g (0.25 mmol) of tetrabutylammoniumiodide, 3.4 g (37.5 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 30 minutes. The temperature is maintained at 60°C for 17.6 hours. The reaction mixture is cooled down to 25°C and stirred with 47g of an aqueous 10% Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator, yielding 12.2 g of a brownish oil partially crystallizing at low temperature. The title product is obtained as an off-white solid, mp 106°C - 110°C. Analysis required for C<sub>25</sub>H<sub>39</sub>NO<sub>5</sub> (433.59): C 69.25%, H 9.07%, N 3.23%; found: C 68.24%, H 9.04%, N 2.87%. <sup>1</sup>H-

- 40 -


NMR ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 0.63 (br s, 3H), 0.93 (br s, 3H), 0.95 (br s, 3H), 1.14 (br s, 3H), 1.30 (br s, 6H), 1.45 - 1.48 (m, 4H), 1.53 - 1.60 (m, 1H), 2.05 - 2.09 (d-like, 1H), 2.16 - 2.20 (d-like, 1H), 2.75 - 2.76 (m, 1H), 2.89 - 2.91 (m, 1H), 3.34 - 3.36 (m, 1H), 3.45 (s, 4H), 3.94 - 3.99 (m, 1H), 4.18 - 4.21 (m, 1H), 4.74 (q,  $J = 8\text{Hz}$ , 1H), 6.84 - 6.87 (d-like, 2H), 7.22 - 7.25 (d-like, 2H).

Example 10: Preparation of the compound of the formula




A stirred mixture of 1.42 g (2.5 mmol) of N,N'-dibutyl-6-chloro-N,N'-bis-(2,2,6,6-tetramethylpiperidin-4-yl-N-oxide)-[1,3,5]-triazine-2,4-diamine, 4.2 g (50 mmol) cyclohexane, 0.018 g (0.05 mmol) tetrabutylammoniumiodide and 1.93 g (15 mmol) t-butylhydroperoxid (70% aqueous solution) is brought to 68°C. The temperature is maintained at 68°C for 22 hours. The reaction mixture is cooled down to 25°C and stirred with 18.9 g of an aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 1.1 g of a reddish solid. Purification by flash-chromatography (silica gel, hexane : ethylacetate 9 : 1) yields the title product as a white solid, mp 86°C - 90°C. Analysis required for  $\text{C}_{41}\text{H}_{74}\text{ClN}_7\text{O}_2$  (732.55): C 67.23%, H 10.18%, Cl 4.84%, N 13.38%; found: C 67.16%, H 10.08%, Cl 4.91%, N 12.86%.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 0.88 - 0.96 (m, 6H), 1.05 - 1.4 (m, 42H), 1.45 - 1.60 (m, 6H), 1.63 - 1.80 (m, 8H), 2.0 - 2.1 (m, 4H), 3.25 - 3.35 (m, 4H), 3.55 - 3.65 (m, 2H), 4.9 - 5.1 (m, 2H).

Example 11: Preparation of the compound of the formula




To a stirred mixture of 8 g (35 mmol) of propionic acid-2,2,6,6-tetramethylpiperidin-4-yl-N-oxide ester, 29.5 g (350 mmol) cyclohexane and 0.13 g (0.35 mmol) of tetrabutylammoniumiodide, 13.5 g (105 mmol) of t-butylhydroperoxid (70% aqueous solution) are added at 60°C within 20 minutes. The temperature is maintained at 60°C for 2.8 hours. The reaction mixture is cooled down to 25°C and stirred with 132 g of an aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 10 g of a reddish oil. Purification by flash-chromatography (silica gel, hexane : ethylacetate 9 : 1) yields the title product as a yellowish oil. Analysis required for  $\text{C}_{18}\text{H}_{33}\text{NO}_3$  (311.47): C 69.41%, H 10.68%, N 4.50%; found: C 69.32%, H 10.57%, N 4.40%.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 1.09 (t,  $J$  = 8Hz, 3H), 1.10 - 1.26 (m, 17H), 1.52 - 1.57 (m, 3H), 1.74 - 1.84 (m, 4H), 2.03 - 2.05 (m, 2H), 2.28 (q,  $J$  = 8Hz, 2H), 3.56 - 3.62 (m, 1H), 4.98 - 5.06 (m, 1H).

Example 12: Preparation of the compound




To a stirred mixture of 8.95 g (30 mmol) 8,10-diethyl-3,3,7,8,10-pentamethyl-1,5-dioxa-9-aza-spiro[5.5]undecane-N-oxide, 24.6 g (300 mmol) cyclohexene and 0.11 g (0.3 mmol) tetrabutylammoniumiodide are added at 65°C within 20 minutes 5.8 g (45 mmol) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 65°C for 15 minutes until all of the N-oxide has reacted. The reaction mixture is cooled down to 25°C and stirred with 57 g of an aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 10.5 g (92% of theory) of a slightly orange oil. Purification by Flash-Chromatography (silica gel, Hexane / Ethylacetate 8 / 2) affords 9.7 g (85% of theory) the title compound as a viscous, colourless oil. Analysis required for  $\text{C}_{23}\text{H}_{41}\text{NO}_3$  (379.58): C 72.78%, H 10.89%, N 3.69%; found: C 72.61%, H 10.65%, N 3.66%.

Example 13: Preparation of the compound




To a stirred mixture of 9.1 g (30 mmol) 8,10-Diethyl-3,3,7,8,10-pentamethyl-1,5-dioxa-9-aza-spiro[5.5]undecane-N-oxide, 31.9 g (300 mmol) Ethylbenzene and 0.11 g (0.3 mmol) Tetrabutylammoniumiodide are added at 60°C within 25 minutes 5.8 g (45 mmol) t-Butylhydroperoxid (70% aqueous solution). The temperature is maintained at 65°C for 15 minutes until all of the N-oxide has reacted. The reaction mixture is cooled down to 25°C and stirred with 57 g of an aqueous 10% Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-Butylhydroperoxide. The aqueous phase is then separated and washed with Ethylbenzene. The combined organic phases are washed with Brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator, yielding 12.4 g (102% of theory) of a slightly yellow oil. Purification by Flash-Chromatography (silica gel, Hexane / Ethylacetate 9.5 / 0.5) affords 10 g (82.6 % of theory) of the title compound as a viscous, colourless oil. Analysis required for C<sub>25</sub>H<sub>41</sub>NO<sub>3</sub> (403.61): C 74.40%, H 10.24%, N 3.47%; found: C 74.29%, H 10.47%, N 3.36%.

Example 14: Preparation of the compound of Example 1 with the catalyst Bu<sub>4</sub>NI generated in situ from Bu<sub>4</sub>NCl / NaI; yield determination by HPLC.




To a stirred mixture of 0.5g (3.2 mmol) 2,2,6,6-tetramethylpiperidine-N-oxide (TEMPO), 3.8 g (35.6 mmol) ethylbenzene, 0.0092 g (0.032 mmol) tetrabutylammoniumchloride and 0.0048g (0.032mmol) sodium iodide dissolved in 1ml water are added at 50°C 0.62 g (4.8 mmol) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 50°C for 80 minutes, after which a sample is withdrawn and analyzed by quantitative HPLC. The yield is 78%.

Example 15: Preparation of the compound of Example 12 using immobilized onium iodide. This allows the catalyst be filtered off after the reaction.



To a stirred mixture of 8.95 g (30 mmol) 8,10-diethyl-3,3,7,8,10-pentamethyl-1,5-dioxa-9-aza-spiro[5.5]undecane-N-oxide, 24.6 g (300 mmol) cyclohexene and 0.3 g (0.3 mmol) tributylmethylammonium iodide bound to polystyrene (1meq iodide / g) are added at 70°C within 35 minutes 5.8 g (45 mmol) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 70°C for 18.5 hours until all of the nitroxide has reacted. The reaction mixture is cooled down to 25°C and the catalyst filtered off. The filtrate is stirred with 57 g of an aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 10.7 g (94% of theory) of the title product as a slightly orange oil.


Example 16: Preparation of the compound of Example 9



To a stirred mixture of 0.769 g (3 mmol) 3,3,8,8,10,10-hexamethyl-1,5-dioxa-9-aza-spiro[5.5]undecane-N-oxide, 1.6 g (9 mmol, 3eq) 2-(4-ethyl-phenoxy)methyl-oxirane, 0.046 g (0.3 mmol, 0.1eq) biphenyl (internal standard) and 0.03 mmol (0.01eq) onium iodide are added at 60°C 0.579 g (4.5 mmol, 1.5eq) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 60°C. Samples are withdrawn and analyzed by quantitative HPLC.

Using  $\text{Bu}_4\text{NI}$  as onium iodide yields 82% of theory after 22 h (nitroxide conversion: 97%).

Good results are also achieved when the amount of 2-(4-ethyl-phenoxy)methyl-oxirane is reduced to 2, 1.5 or 1 eq.; or when using 1 eq. of 2-(4-ethyl-phenoxy)methyl-oxirane, the catalyst is replaced by the equivalent amount of  $\text{Ph}_4\text{PI}$  or  $\text{Oct}_3\text{MeNI}$ , or the amount of  $\text{Bu}_4\text{NI}$  is increased to 0.15 mmol (0.05 eq.).

Example 17: Preparation of the compound

CAS Regno 117174-66-0

To a stirred mixture of 0.829 g (3 mmol) benzoic acid-2,2,6,6-tetramethyl-piperidin-4-yl-N-oxid ester, 2.53 g (30 mmol, 10eq) cyclohexane, 0.046 g (0.3 mmol, 0.1eq) biphenyl (internal standard) and 0.03 mmol (0.01eq) onium iodide are added at 60°C 0.579 g (4.5 mmol, 1.5eq) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained constant. Samples are withdrawn after 22 h and analyzed by quantitative HPLC. Results are given in the tables below:

Table: Yield and nitroxide conversion after 22 h reaction at various temperatures

Catalyst	Reaction Temperature	Product yield [%]	Nitroxide conversion [%]	
Bu <sub>4</sub> NI	60°C	33	38	
Oct <sub>3</sub> MeNI	60°C	31	35	
Bu <sub>4</sub> NI	70°C	43	48	
Bu <sub>4</sub> NI	80°C	46	52	

Good results are also achieved when the amount onium iodide catalyst or the amount of tert.butyl hydroperoxide is doubled.

**Table:** Product yield and nitroxide conversion after 22 h reaction at 80°C and using 9 mmol (3 eq.) of tert.butyl hydroperoxide

Catalyst	Product yield [%]	Nitroxide conversion [%]
Bu <sub>4</sub> NI	63	69
Oct <sub>4</sub> NI	59	67
Hexadecyl <sub>4</sub> NI	59	68
Dodecyl <sub>4</sub> NI	58	67
Hex <sub>4</sub> NI	58	68
Octadecyl <sub>2</sub> Me <sub>2</sub> NI	57	64
HexadecylBzMe <sub>2</sub> NI	57	63
Tetradecyl <sub>2</sub> Me <sub>2</sub> NI	56	63
Oct <sub>3</sub> PrNI	56	65
OctBzMe <sub>2</sub> NI	56	63
Oct <sub>3</sub> MeNI	54	63
HexadecylPyl	54	59
Oct <sub>2</sub> Me <sub>2</sub> NI	53	62
OctMe <sub>3</sub> NI	52	57
Et <sub>4</sub> N	38	42
Oct <sub>2</sub> MeSI	12	17
Ph <sub>4</sub> PI	74	88
Ph <sub>3</sub> iPrPI	71	87
Ph <sub>3</sub> EtPI	63	74
Ph <sub>3</sub> HexPI	61	71
Bu <sub>4</sub> PI	61	68
Bu <sub>3</sub> HexadecylPI	61	68
Oct <sub>4</sub> PI	58	66
Ph <sub>3</sub> MePI	57	65
Ph <sub>2</sub> Me <sub>2</sub> PI	51	56
Et <sub>4</sub> PI	46	50
PhMe <sub>3</sub> PI	39	44
Ph <sub>3</sub> (CH <sub>2</sub> CO <sub>2</sub> Me)PI	36	35
Ph <sub>3</sub> BzPI	34	40


Abbreviations: *Me* methyl, *Et* ethyl, *Pr* n-propyl, *iPr* iso-propyl, *Bu* n-butyl, *Hex* n-hexyl, *Oct* n-octyl, *Ph* phenyl, *Bz* benzyl, *Py* 1-pyridinium

Using a wide variety of catalysts, the present process effectively converts the N-oxide into the desired product, yielding only low levels of by-products.

Example 18: Preparation of the compound of Example 17 using  $\text{Ph}_4\text{PI}$  as catalyst

To a stirred mixture of 8.3 g (30 mmol) benzoic acid-2,2,6,6-tetramethyl-piperidin-4-yl-N-oxide ester, 25.4 g (300 mmol) cyclohexane and 0.14 g (0.3 mmol) tetraphenylphosphonium iodide are added at 80°C within 30 minutes 11.6 g (90 mmol) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 80°C for 19.3 hours. The reaction mixture is cooled down to 25°C and stirred with aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is then separated and washed with cyclohexane. The combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 9 g of a red oil. Purification by flash-chromatography (silica gel, hexane / ethylacetate 9 / 1) affords 6.8 g (63% of theory) of the product as a viscous, colorless oil. Analysis required for  $\text{C}_{22}\text{H}_{33}\text{NO}_3$  (359.51): C 73.50%, H 9.25%, N 3.90%; found: C 72.68%, H 9.39%, N 3.85%.

Example 19: Preparation of the compound




CAS Regno 264224-73-9

To a stirred mixture of 7.7 g (45 mmol) triacetoneamine-N-oxide, 37.3 g (450 mmol) cyclohexene and 0.17 g (0.45 mmol) tetrabutylammonium iodide are added at 60°C within 1 hour 17.4 g (135 mmol) t-butylhydroperoxid (70% aqueous solution). The temperature is maintained at 60°C for 21.7 hours. After further addition of catalyst (0.24g, 0.45 mmol trioctylmethylammonium iodide) and t-butylhydroperoxide (17.4g, 135 mmol) the temperature is maintained another 24 hours. The reaction mixture is then cooled down to 25°C and stirred with aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-butylhydroperoxide. The aqueous phase is separated and washed with cyclohexane. The

- 47 -


combined organic phases are washed with brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator, yielding 11.7 g of an orange oil. Purification by flash-chromatography (silica gel, hexane / ethylacetate 9 / 1) affords the title product as a colorless oil. Analysis required for  $\text{C}_{15}\text{H}_{25}\text{NO}_2$  (251.37): C 71.67%, H 10.02%, N 5.57%; found: C 71.33%, H 10.03%, N 5.78%.

Example 20: Preparation of the compound



To a stirred mixture of 5g (32 mmol) TEMPO, 52.5 g (320 mmol) 2-Phenylethylacetate and 0.12 g (0.32 mmol) Tetrabutylammoniumiodide are added at 60°C within 25 minutes 12.37 g (96 mmol) t-Butylhydroperoxid (70% aqueous solution). The temperature is maintained at 60°C for 18.67 hours until all of the TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 121 g of an aqueous 10%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess t-Butylhydroperoxide. The aqueous phase is then separated and washed with Ethylbenzene. The combined organic phases are washed with Brine, dried over  $\text{MgSO}_4$ , filtered and the solvent distilled off on a rotary-evaporator. The crude product is purified by flash-chromatography (silica gel, Hexane / Ethylacetate 9 / 1), yielding the title product as a colorless oil. Analysis for  $\text{C}_{19}\text{H}_{29}\text{NO}_3$  (319.45): C 71.44%, H 9.15%, N 4.38%; found: C 71.36%, H 9.20%, N 4.21%.  $^1\text{H-NMR}$  ( $\text{CDCl}_3$ ),  $\delta$  (ppm): 0.66 (broad s, 3H), 1.08-1.60 (m, 15H), 1.95 (s, 3H), 4.23-4.30 (m, 1H), 4.57-4.61 (m, 1H), 4.91 (t,  $J = 8\text{Hz}$ , 1H), 7.28-7.37 (m, 5H).


Example 21: Preparation of the compound



To a stirred mixture of 7.8 g (50 mmol) TEMPO, 41.1 g (500 mmol) Cyclohexene and 0.18 g (0.5 mmol) Tetrabutylammoniumiodide are added at 55°C within 30 minutes 7.4 g (58 mmol) t-Butylhydroperoxid (70% aqueous solution). The reaction mixture is cooled down to 25°C and stirred with 63 g of an aqueous 20%  $\text{Na}_2\text{SO}_3$  solution until the disappearance of excess

t-Butylhydroperoxide. The aqueous phase is then separated and washed with Cyclohexane. The combined organic phases are passed through a plug of silica gel and washed with Brine, dried over  $MgSO_4$ , filtered and the solvent distilled off on a rotary-evaporator. The crude product is purified by distillation, yielding 8 g (67.4 % of theory) of an orange oil (bp 62°C-65°C / 0.04 mbar). Analysis required for  $C_{15}H_{27}NO$  (237.39): C 75.90%, H 11.46%, N 5.90%; found: C 75.69%, H 11.99%, N 5.75%.  $^1H$ -NMR ( $CDCl_3$ ),  $\delta$  (ppm): 1.13-2.07 (m, 24H), 4.24 (br s, 1H), 5.77-5.81 (m, 1H), 5.91-5.95 (m, 1H).


Example 22: Hydrogenation of the product of Example 21




A mixture of 0.95 g (4 mmol) 1-(Cyclohex-2-enyloxy)-2,2,6,6-tetramethyl-piperidine and 0.2 g Pd on charcoal (10%) in 10 ml Methanol is hydrogenated at 25°C and 4 bar Hydrogen. Filtration and evaporation of the solvent yields the title product as a slightly orange oil. Analysis for  $C_{15}H_{29}NO$  (239.40): C 75.26%, H 12.21%, N 5.85%; found: C 74.53%, H 12.07%, N 5.90%.  $^1H$ -NMR ( $CDCl_3$ ),  $\delta$  (ppm): 1.12-1.39 (m, 19H), 1.40-1.65 (m, 7H), 1.74 (br s, 1H), 2.04 (br s, 1H), 3.58 (m, 1H).

Example 23: Hydrogenation of the crude product of Example 21


A mixture of the crude product from example 21 (10.87 g, 91.6% of theory) and 2.4 g Pd on charcoal (10%) in 120 ml Methanol is hydrogenated as described in example 22. Filtration and evaporation of the solvent yields 6.8 g of a slightly yellow oil. Analysis required for  $C_{15}H_{29}NO$  (239.40): C 75.26%, H 12.21%, N 5.85%; found: C 74.53%, H 12.07%, N 5.90%.  $^1H$ -NMR ( $CDCl_3$ ),  $\delta$  (ppm): 1.12-1.39 (m, 19H), 1.40-1.65 (m, 7H), 1.74 (br s, 1H), 2.04 (br s, 1H), 3.58 (m, 1H).

Example 24: Preparation of the compound

To a stirred mixture of 7.3 g (32 mmol) Propionic acid-2,2,6,6-tetramethylpiperidin-4-yl-N-oxide ester, 26.3 g (320 mmol) Cyclohexene and 0.12 g (0.32 mmol) Tetrabutylammoniumiodide are added at 55°C within 25 minutes 6.2 g (48 mmol) t-Butylhydroperoxid (70% aqueous solution). The temperature is maintained at 55°C for 5 minutes until all of the TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 61 g of an aqueous 10% Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-Butylhydroperoxide. The aqueous phase is then separated and washed with Cyclohexane. The combined organic phases are passed through a plug of silica gel and washed with Brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator, yielding 8.7 g (87.9% of theory) of the above product as a slightly orange oil. Analysis required for C<sub>18</sub>H<sub>31</sub>NO<sub>3</sub> (309.45): C 69.87%, H 10.10%, N 4.53%; found: C 69.36%, H 10.03%, N 4.45%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm): 1.12 (t, J = 8Hz, 3H), 1.20-1.26 (m, 12H), 1.52-1.58 (m, 4H), 1.73-2.1 (m, 6H), 2.29 (q, J = 8Hz, 2H), 4.23 (m, 1H), 5.05 (m, 1H), 5.79-5.82 (m, 1H), 5.90-5.94 (m, 1H).

Example 25: Hydrogenation of the product of Example 24


A mixture of CG40-1201 (1g, 3.19 mmol) and 0.17 g Pd on charcoal (10%) in 30 ml Hexane is hydrogenated as described in example 6. Filtration and evaporation of the solvent yields 0.9 g (90.6% of theory) of a slightly yellow oil. Analysis required for C<sub>18</sub>H<sub>33</sub>NO<sub>3</sub> (311.47): C 69.41%, H 10.68%, N 4.50%; found: C 69.20%, H 10.76%, N 4.42%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm): 1.09 (t, J = 8Hz, 3H), 1.10 - 1.26 (m, 17H), 1.52 - 1.57 (m, 3H), 1.74 - 1.84 (m, 4H), 2.03 - 2.05 (m, 2H), 2.28 (q, J = 8Hz, 2H), 3.56 - 3.62 (m, 1H), 4.98 - 5.06 (m, 1H).

Example 26: Preparation of the compound

To a stirred mixture of 14.2 g (25 mmol) of N,N'-Dibutyl-6-chloro-N,N'-bis-(2,2,6,6-tetramethyl-piperidin-4-yl-N-oxide)-[1,3,5]-triazine-2,4-diamine, 41 g (500 mmol) Cyclohexene and 0.18 g (0.5 mmol) Tetrabutylammoniumiodide are added at 57°C within 30 minutes 9.7 g (75 mmol) t-Butylhydroperoxid (70% aqueous solution). The temperature is maintained at 57°C for 5 minutes until all of the TEMPO has reacted. The reaction mixture is cooled down to 25°C and stirred with 63 g of an aqueous 10% Na<sub>2</sub>SO<sub>3</sub> solution until the disappearance of excess t-Butylhydroperoxide. The aqueous phase is then separated and washed with Cyclohexane. The combined organic phases are washed with Brine, dried over MgSO<sub>4</sub>, filtered and the solvent distilled off on a rotary-evaporator, yielding 14.5 g (79.6% of theory) of a slightly yellow solid. Crystallization from Acetone / Hexane yields 12.2 g (67%) of a white solid, mp 83°C - 87°C. Analysis required for C<sub>41</sub>H<sub>70</sub>ClN<sub>7</sub>O<sub>2</sub> (728.51): C 67.60%, H 9.69%, Cl 4.87%, N 13.46%; found: C 67.27%, H 9.63%, Cl 4.97%, N 13.34%. <sup>1</sup>H-NMR (CDCl<sub>3</sub>), δ (ppm): 0.89 - 0.96 (m, 6H), 1.22 - 1.32 (m, 26H), 1.49 - 1.56 (m, 12H), 1.73 - 1.78 (m, 8H), 1.89 - 2.04 (m, 6H), 3.31 - 3.32 (m, 4H), 4.24 - 4.26 (m, 2H), 4.99 - 5.06 (m, 2H), 5.80 - 5.83 (m, 2H), 5.92 - 6.02 (m, 2H).

## WHAT IS CLAIMED IS:

1. Process for the preparation of an amine ether of a sterically hindered amine by reacting a corresponding sterically hindered aminoxide with an aliphatic hydrocarbon compound, characterized in that the reaction is carried out in the presence of an organic hydroperoxide and an iodide.
2. Process of claim 1 for the preparation of an amine ether of a sterically hindered amine by reacting a corresponding sterically hindered aminoxide with a hydrocarbon compound, characterized in that the reaction is carried out in the presence of an organic hydroperoxide and a catalytic amount of an iodide.
3. Process of claim 1, wherein the amine ether is of the formula A





wherein

a is 1 or 2;

when a is 1, E' is E

when a is 2, E' is L;

E is C<sub>1</sub>-C<sub>36</sub> alkyl; C<sub>3</sub>-C<sub>18</sub> alkenyl; C<sub>2</sub>-C<sub>18</sub> alkinyl; C<sub>5</sub>-C<sub>18</sub> cycloalkyl; C<sub>5</sub>-C<sub>18</sub> cycloalkenyl; a radical of a saturated or unsaturated aliphatic bicyclic or tricyclic hydrocarbon of 7 to 12 carbon atoms; C<sub>2</sub>-C<sub>7</sub>alkyl or C<sub>3</sub>-C<sub>7</sub>alkenyl substituted by halogen, C<sub>1</sub>-C<sub>8</sub>alkoxy or phenoxy; C<sub>4</sub>-C<sub>12</sub>heterocycloalkyl; C<sub>4</sub>-C<sub>12</sub>heterocycloalkenyl; C<sub>7</sub>-C<sub>15</sub> aralkyl or C<sub>4</sub>-C<sub>12</sub>heteroaralkyl, each of which is unsubstituted or substituted by C<sub>1</sub>-C<sub>4</sub> alkyl or phenyl; or E is a radical of formula (VII) or (VIII)



Ar is C<sub>6</sub>-C<sub>10</sub>aryl or C<sub>5</sub>-C<sub>9</sub>heteroaryl;

X is phenyl, naphthyl or biphenyl, which is substituted by 1, 2, 3 or 4 D and optionally further substituted by NO<sub>2</sub>, halogen, amino, hydroxy, cyano, carboxy, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylamino or di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino;

D is a group , a group C(O)-G<sub>13</sub> or a group C(O)-G<sub>9</sub>-C(O)-G<sub>13</sub>;

G<sub>1</sub> and G<sub>2</sub>, independently of each other, are hydrogen, halogen, NO<sub>2</sub>, cyano, -CONR<sub>5</sub>R<sub>6</sub>, -(R<sub>9</sub>)COOR<sub>4</sub>, -C(O)-R<sub>7</sub>, -OR<sub>8</sub>, -SR<sub>8</sub>, -NHR<sub>8</sub>, -N(R<sub>18</sub>)<sub>2</sub>, carbamoyl, di(C<sub>1</sub>-C<sub>18</sub>alkyl)carbamoyl, -C(=NR<sub>5</sub>)(NHR<sub>6</sub>), C<sub>1</sub>-C<sub>18</sub>alkyl; C<sub>3</sub>-C<sub>18</sub>alkenyl; C<sub>3</sub>-C<sub>18</sub>alkinyl, C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl; C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>3</sub>-C<sub>18</sub>alkenyl or C<sub>3</sub>-C<sub>18</sub>alkinyl or C<sub>7</sub>-C<sub>9</sub>phenylalkyl, C<sub>3</sub>-C<sub>12</sub>cycloalkyl or C<sub>2</sub>-C<sub>12</sub>heterocycloalkyl substituted by OH, halogen, NO<sub>2</sub>, amino, cyano, carboxy, COOR<sub>21</sub>, C(O)-R<sub>22</sub>; C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, C<sub>1</sub>-C<sub>4</sub>alkylamino or di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino or a group -O-C(O)-R<sub>7</sub>; C<sub>2</sub>-C<sub>18</sub>alkyl which is interrupted by at least one O atom and/or NR<sub>5</sub> group; or are C<sub>6</sub>-C<sub>10</sub>aryl; or phenyl or naphthyl which are substituted by C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkoxy, C<sub>1</sub>-C<sub>4</sub>alkylthio, halogen, cyano, hydroxy, carboxy, COOR<sub>21</sub>, C(O)-R<sub>22</sub>, C<sub>1</sub>-C<sub>4</sub>alkylamino or di(C<sub>1</sub>-C<sub>4</sub>alkyl)amino; or G<sub>1</sub> and G<sub>2</sub> together with the linking carbon atom form a C<sub>3</sub>-C<sub>12</sub>cycloalkyl radical;

G<sub>5</sub> and G<sub>6</sub> are independently of each other H or CH<sub>3</sub>;

G<sub>9</sub> is C<sub>1</sub>-C<sub>12</sub>alkylene or a direct bond;

G<sub>13</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl;

G<sub>14</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>5</sub>-C<sub>12</sub>cycloalkyl, an acyl radical of an aliphatic or unsaturated aliphatic carboxylic or carbamic acid containing 2 to 18 carbon atoms, an acyl radical of a cyclo-aliphatic carboxylic or carbamic acid containing 7 to 12 carbon atoms, or acyl radical of an aromatic acid containing 7 to 15 carbon atoms;

G<sub>55</sub> is H, CH<sub>3</sub> or phenyl;


G<sub>66</sub> is -CN or a group of the formula -COOR<sub>4</sub> or -CONR<sub>5</sub>R<sub>6</sub> or -CH<sub>2</sub>-O-G<sub>14</sub>;

L is alkylene of 1 to 18 carbon atoms, cycloalkylene of 5 to 8 carbon atoms, cycloalkenylene of 5 to 8 carbon atoms, alkenylene of 3 to 18 carbon atoms, alkylene of 1 to 12 carbon atoms substituted by phenyl or by phenyl substituted by alkyl of 1 to 4 carbon atoms; or is alkylene of 4 to 18 carbon atoms interrupted by COO and/or phenylene;

T' is tertiary C<sub>4</sub>-C<sub>18</sub>alkyl or phenyl, each of which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>; or T' is C<sub>5</sub>-C<sub>12</sub>cycloalkyl; C<sub>5</sub>-C<sub>12</sub>cycloalkyl which is interrupted by at

- 53 -

least one O or -NR<sub>18</sub>-; a polycyclic alkyl radical having 7-18 carbon atoms, or the same radical which is interrupted by at least one O or -NR<sub>18</sub>-; or T' is -C(G<sub>1</sub>)(G<sub>2</sub>)-T"; or C<sub>1</sub>-C<sub>18</sub>alkyl



or C<sub>5</sub>-C<sub>12</sub>cycloalkyl substituted by ;

T" is hydrogen, halogen, NO<sub>2</sub>, cyano, or is a monovalent organic radical comprising 1-50 carbon atoms;

or T" and T' together form a divalent organic linking group completing, together with the hindered amine nitrogen atom and the quaternary carbon atom substituted by G<sub>1</sub> and G<sub>2</sub>, an optionally substituted five- or six-membered ring structure;  
and

R<sub>4</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl, phenyl, an alkali metal cation or a tetraalkylammonium cation;  
R<sub>5</sub> and R<sub>6</sub> are hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl, C<sub>2</sub>-C<sub>18</sub>alkyl which is substituted by hydroxy or, taken together, form a C<sub>2</sub>-C<sub>12</sub>alkylene bridge or a C<sub>2</sub>-C<sub>12</sub>-alkylene bridge interrupted by O or/and NR<sub>18</sub>;

R<sub>7</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>6</sub>-C<sub>10</sub>aryl;

R<sub>8</sub> is hydrogen, C<sub>1</sub>-C<sub>18</sub>alkyl or C<sub>2</sub>-C<sub>18</sub>hydroxyalkyl;


R<sub>9</sub> is C<sub>1</sub>-C<sub>12</sub>alkylene or a direct bond;

R<sub>18</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl or phenyl, which are unsubstituted or substituted by halogen, OH, COOR<sub>21</sub> or C(O)-R<sub>22</sub>;



R<sub>21</sub> is hydrogen, a alkali metal atom or C<sub>1</sub>-C<sub>18</sub>alkyl; and

R<sub>22</sub> is C<sub>1</sub>-C<sub>18</sub>alkyl;

the aminoxide is of formula B



and the hydrocarbon is of formula IV or V




wherein E, G<sub>1</sub>, G<sub>2</sub>, L, T and T' are as defined for formula A.


4. Process according to claim 1, wherein the organic hydroperoxide used in the process of present invention is a peroxyalcohol containing 3-18 carbon atoms, especially tert.butyl-hydroperoxide.
5. Process according to claim 1, wherein 1 to 100 moles of the hydrocarbon, 1 to 20 moles of organic hydroperoxide, and 0.001 mmoles to 0.5 moles of iodide catalyst are used per mole of aminoxide.
6. Process according to claim 1, which is carried out in the absence of copper or a copper compound.
7. Process according to claim 1, wherein the hydrocarbon is used in excess and serves both as reactant and as solvent for the reaction and/or wherein a further inert organic or inorganic solvent is used.
8. Process according to any of claims 1 to 7, wherein the reaction is carried out in the presence of a phase transfer catalyst.
9. Process according to claim 8, wherein the catalyst is selected from alkaline or alkaline earth metal iodides, ammonium iodides and phosphonium iodides, especially from tetraalkyl ammonium iodides, tetraphenylphosphonium iodide and triphenylalkylphosphonium iodides.
10. Process according to claim 3, wherein in the formulae A and B T and T' together are an organic linking group containing 2-500 carbon atoms and 0-200 hetero atoms selected from oxygen, phosphorus, sulfur, silicon, halogen and nitrogen as tertiary nitrogen, and forming, together with the carbon atoms it is directly connected to and the nitrogen atom, an optionally substituted, 5-, 6 or 7-membered cyclic ring structure.
11. Process according to claim 1, wherein the aliphatic hydrocarbon compound contains an ethylenic double bond, and the product is subsequently hydrogenated.
12. Use of an organic hydroperoxide together with an iodide and an aliphatic hydrocarbon compound for the preparation of a sterically hindered amine ether from its N-oxyl precursor.
13. A compound of the formula a, b c or d:

- 55 -


a)




b)



c)



d)



14. Use of a sterically hindered amine ether obtained according to any of claims 1-10 as a stabilizer for organic material against degradation by light, oxygen and/or heat, or as a polymerization regulator.