



(51) International Patent Classification:

A61P 31/18 (2006.01) C07K 16/10 (2006.01)

(21) International Application Number:

PCT/US2019/040342

(22) International Filing Date:

02 July 2019 (02.07.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/693,642 03 July 2018 (03.07.2018) US
 62/810,191 25 February 2019 (25.02.2019) US

(71) Applicant: GILEAD SCIENCES, INC. [US/US]; 333 Lakeside Drive, Foster City, California 94404 (US).

(72) Inventors: BALAKRISHNAN, Mini; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). CARR, Brian A.; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). HUNG, Magdeleine S.; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). KANWAR, Manu; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). PACE,

Craig S.; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). REHDER, Doug; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). SCHENAUER, Matthew Robert; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). SERAFINI, Loredana; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). STEPHENSON, Heather Theresa; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). THOMSEN, Nathan D.; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). YU, Helen; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US). ZHANG, Xue; c/o Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US).

(74) Agent: WAHLSTEN, Jennifer L. et al.; Gilead Sciences, Inc., 333 Lakeside Drive, Foster City, California 94404 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

(54) Title: ANTIBODIES THAT TARGET HIV GP120 AND METHODS OF USE




FIG. 1

(57) Abstract: Antibodies that bind to HIV gp120 and neutralize HIV are disclosed. Also disclosed are methods of using such antibodies alone or in combination with other therapeutic agents to treat or prevent HIV infection.

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

ANTIBODIES THAT TARGET HIV GP120 AND METHODS OF USECross Reference to Related Applications

[0001] This application claims the benefit under 35 U.S.C. § 119(e) of U.S. provisional application no. 62/693,642, filed on July 3, 2018 and U.S. provisional application no. 62/810,191, filed on February 25, 2019, which are hereby incorporated herein by reference in their entireties for all purposes.

Sequence Listing

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on June 10, 2019, is named 1232_P2F_SL.txt and is 899,216 bytes in size.

Field

[0003] This disclosure relates to antibodies and antigen-binding fragments thereof for the treatment and/or prevention of human immunodeficiency virus (HIV) infection.

Background

[0004] Human immunodeficiency virus (HIV) infection and related diseases are a major public health problem worldwide. Most currently approved therapies for HIV infection target the viral reverse transcriptase, protease enzymes, and integrase. Yet resistance of HIV to these existing drugs, long-term toxicity, and lack of patient adherence to daily dosing regimens have been associated with these therapies. Therefore, it is important to discover and develop new anti-HIV antibodies with advantageous properties suitable for therapeutic uses.

[0005] WO 2012/158948 describes human anti-HIV antibodies derived from memory B cells of HIV-infected donors, which are capable of inhibiting infection by HIV-1 species from a plurality of clades. Anti-HIV antibodies are also disclosed *e.g.*, in WO 2005/058963, WO 2013/090644, WO 2014/063059 and EP 0690132B1. The therapeutic use of the antibodies may be limited due to their intra-patient viral coverage, pharmacokinetics, polyspecificity, and other properties. Accordingly, there is a need for novel anti-HIV antibodies for therapeutic uses.

Summary

[0006] The present disclosure provides compositions for treating or preventing HIV. More specifically, provided herein are antibodies that bind human immunodeficiency

virus (HIV) envelope (Env) glycoprotein gp120 (gp120). This disclosure provides anti-HIV antibodies and antigen-binding fragments thereof, including broadly neutralizing anti-HIV antibodies and antigen-binding fragments thereof, pharmaceutical compositions containing such antibodies and fragments thereof, and methods for using these antibodies and fragments thereof in the treatment and prevention of HIV infection.

[0007] In one aspect, this disclosure provides an antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120. The antibody or antigen-binding fragment thereof comprises a heavy chain variable region (VH) comprising VH complementary determining regions (CDRs) and a light chain variable region (VL) comprising VL CDRs. In some embodiments, the VH CDRs and VL CDRs have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 159, 138, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 160, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 161, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 162, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 163, 139, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 138, 164, 140, 141, and 142, respectively; SEQ ID NOs.: 159, 138, 164, 140, 141, and 142, respectively; SEQ ID NOs.: 137, 138, 139, 140, 165, and 142, respectively; SEQ ID NOs.: 137, 138, 139, 140, 166, and 142, respectively; SEQ ID NOs.: 137, 138, 139, 140, 167, and 142, respectively; SEQ ID NOs.: 137, 138, 139, 140, 168, and 142, respectively; SEQ ID NOs.: 137, 138, 154, 140, 141, and 142, respectively, or SEQ ID NOs.: 137, 138, 139, 570, 141, and 142, respectively. In some cases, the antibody or antigen-binding fragment thereof comprises in framework region 3 (FR3) of the VH at position corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO: 453 or SEQ ID NO: 627. In some, the VH CDRs and VL CDRs have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein the antibody or antigen-binding fragment thereof comprises in framework region 3 (FR3) of the VH at position corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO: 627. In some cases, the antibody or antigen-binding fragment thereof comprises a FR3 of the VH comprising the following amino acid sequence:
 RVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 628) or
 RVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In certain embodiments, the antibody or antigen-binding fragment thereof comprises a FR3

of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In some, the VH CDRs and VL CDRs have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein the antibody or antigen-binding fragment thereof comprises a FR3 of the VH comprising the following amino acid sequence: RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629).

[0008] In another aspect, the VH CDRs and VL CDRs have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively; or SEQ ID NOs.: 153, 138, 139, 140, 141, and 142, respectively. In certain cases, the VH of this antibody has one or more of: histidine at position 3, serine at position 5, glutamine at position 72, tyrosine at position 76, valine at position 82c, isoleucine at position 89 (position numbering according to Kabat). In certain cases, the VL of this antibody has one or more of: arginine at position 14, alanine at position 60, valine at position 83, and isoleucine at position 98 (position numbering according to Kabat). In some cases, the antibody or antigen-binding fragment thereof comprises in framework region 3 (FR3) of the VH at position corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO: 453 or SEQ ID NO: 627. In some cases, the antibody or antigen-binding fragment thereof comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 628) or RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In certain embodiments, the antibody or antigen-binding fragment thereof comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises in framework region 3 (FR3) of the VH at position corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO: 627. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In

certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629), and comprises a VL comprising the amino acid sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477, and comprises a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629), and comprises a VL comprising the amino acid sequence set forth in SEQ ID NO: 278.

[0009] The foregoing antibodies may further comprise a VH with one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, or 18) of the following amino acids at the indicated positions (position numbering according to Kabat): valine at position 5, glutamic acid at position 10, lysine at position 12, lysine at position 23, asparagine at position 28, arginine at position 30, tyrosine at position 32, threonine at position 68, methionine at position 69, histidine at position 72, phenylalanine at position 76, alanine at position 78, serine at position 82a, arginine at position 82b, threonine at position 89, tyrosine at position 99, glutamine at position 105, or methionine at position 108. In

certain embodiments, the antibody may further comprise a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72 and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprise a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72 and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprise a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72, phenylalanine a position 74a and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73, phenylalanine a position 76 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72 and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73 and tyrosine at position 98, wherein the amino acid positions are with respect

to SEQ ID NO: 477). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72, phenylalanine a position 74a and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73, phenylalanine a position 76 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477).

[0010] In some embodiments, the VL comprises one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) of the following amino acids at the indicated positions (position numbering according to Kabat): arginine at position 18, lysine at position 39, proline at position 40, threonine at position 56, serine at position 65, threonine at position 72, serine at position 76, serine at position 77, threonine at position 99, glycine at position 99, asparagine at position 103, or isoleucine at position 106. In other embodiments, the VL comprises one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) of the following amino acids at the indicated positions (position numbering according to Kabat): arginine at position 18, alanine at position 19, serine at position 65, threonine or histidine at position 72, lysine at position 74, serine at position 76, serine at position 77, phenylalanine at position 98, or glycine at position 99. In certain embodiments, the VL comprises an alanine at position 19 (Kabat numbering). In yet other embodiments, the VH comprises one or more of the following amino acids at the indicated positions (position numbering according to Kabat): histidine at position 72, phenylalanine at position 76, or phenylalanine at position 74a. In other embodiments, the VL comprises one or more (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8) of the following amino acids at the indicated positions (position numbering according to Kabat): arginine at position 18, alanine at position 19, serine at position 65, threonine at position 72, serine at position 76, serine at position 77, phenylalanine at position 98, or glycine at position 99. In certain embodiments, the antibody or antigen-binding fragment thereof

comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72, phenylalanine at position 76, and phenylalanine at position 74a, and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73, phenylalanine a position 76 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a VL with an alanine at position 19 (Kabat numbering). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72, phenylalanine at position 76, and phenylalanine at position 74a, and tyrosine

at position 99, and comprises a VL with the following amino acids at the indicated positions (position numbering according to Kabat): alanine at position 19.

[0011] In certain embodiments, the VL comprises an amino acid sequence set forth in any one of SEQ ID NOs.: 332 to 342. In some cases, the antibody comprises a human IgG1 Fc region. In certain embodiments, the human IgG1 Fc region is IgG1m17 (SEQ ID NO: 348).

[0012] The foregoing antibody or antigen-binding fragment thereof further comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): (i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330; (ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434; (iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434; (iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434; (v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or (vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434. In certain embodiments, the antibody or antigen-binding fragment thereof further comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and further comprises a human IgG1 Fc region. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and further comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434. In certain embodiments, antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least

96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278 and further comprises a human IgG1 Fc region. In certain embodiments, antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278 and further comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434.

[0013] In certain embodiments, the antibody comprises a human kappa light chain constant region. In some cases, the human kappa light chain constant region is Km3 (SEQ ID NO:351). In a certain embodiment, the human kappa light chain constant region is Km3 (SEQ ID NO: 351). In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and further comprises the human kappa light chain constant region Km3 (SEQ ID NO: 351). In certain embodiments, antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278 and further comprises the human kappa light chain constant region Km3 (SEQ ID NO: 351). In certain embodiments, antibody or antigen-binding

fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278 and further comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434, and the human kappa light chain constant region Km3 (SEQ ID NO: 351).

[0014] In some embodiments, the antibody or antigen-binding fragment has improved, extended, enhanced or increased serum half-life in a mammal (*e.g.*, in a non-human primate, in a human) compared to other anti-HIV antibodies, such as Antibody A. In some embodiments, the antibody or antigen-binding fragment has a serum half-life in a human of at least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer. In some embodiments, the antibody or antigen-binding fragment has improved, enhanced or increased killing potency of HIV-infected cells compared to other anti-HIV antibodies, such as Antibody A. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and has improved, extended, enhanced or increased serum half-life in a mammal (*e.g.*, in a non-human primate, in a human) compared to other anti-HIV antibodies, such as Antibody A. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and has improved, enhanced or increased killing potency of HIV-infected cells compared to other anti-HIV antibodies, such as Antibody A.

[0015] In another aspect, the disclosure provides an antibody that binds to HIV-1 Envelope glycoprotein gp120. The antibody comprises a VH comprising VH CDRs 1-3

and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively; or SEQ ID NOs.: 153, 138, 154, 140, 141, and 142, respectively. The antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): (i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330; (ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434; (iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434; (iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434; (v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or (vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein the antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434.

[0016] In certain embodiments, the antibody comprises a light chain comprising an alanine at position 19 (Kabat numbering). In some embodiments, the antibody comprises in framework region 3 (FR3) of the VH at positions corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO:453 or SEQ ID NO: 627. In certain embodiments, the antibody comprises in framework region 3 (FR3) of the VH at positions corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO: 627. In some embodiments, the antibody comprises a FR3 of the VH comprising the following amino acid sequence: RVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 628) or RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In some embodiments, the antibody comprises a FR3 of the VH comprising the following amino acid sequence: RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629). In some

embodiments, the antibody comprises an amino acid sequence set forth in any one of SEQ ID NOs.: 332 to 342. In some cases, the antibody comprises a VH and VL having the amino acid sequence set forth in SEQ ID NOs.: 182 and 223, respectively. In some cases, the antibody comprises a VH and VL having the amino acid sequence set forth in SEQ ID NOs.: 220 and 276, respectively. In certain embodiments, the antibody comprises a VH and VL having the amino acid sequence set forth in SEQ ID NOs.: 477 and 278, respectively. In other embodiments, the human IgG1 Fc region is IgG1m17 (SEQ ID NO: 348). In some embodiments, the antibody comprises a human kappa light chain constant region. In certain cases, the human kappa light chain constant region is Km3 (SEQ ID NO: 351).

[0017] In some embodiments, the antibody or antigen-binding fragment has improved, extended, enhanced or increased serum half-life in a mammal (*e.g.*, in a non-human primate, in a human) compared to other anti-HIV antibodies, such as Antibody A and/or Antibody B. In some embodiments, the antibody or antigen-binding fragment has a serum half-life in a human of at least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer. In some embodiments, the antibody has improved, increase, or enhanced killing potency of HIV-infected cells compared to other anti-HIV antibodies such as Antibody A and/or Antibody B. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein the antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): leucine at position 428, and serine at position 434, and has improved, extended, enhanced or increased serum half-life in a mammal (*e.g.*, in a non-human primate, in a human) compared to other anti-HIV antibodies, such as Antibody A and/or Antibody B. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein the antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, and

has improved, enhanced or increased killing potency of HIV-infected cells compared to other anti-HIV antibodies, such as Antibody A and/or Antibody B.

[0018] In yet another aspect, the disclosure provides an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region (VH) and a light chain variable region (VL), wherein the VH and VL comprise the amino acid sequences set forth, respectively: (1) SEQ ID NOs.: 184 and 223; (2) SEQ ID NOs.: 185 and 223; (3) SEQ ID NOs.: 182 and 225; (4) SEQ ID NOs.: 185 and 225; (5) SEQ ID NOs.: 186 and 223; (6) SEQ ID NOs.: 187 and 223; (7) SEQ ID NOs.: 188 and 223; (8) SEQ ID NOs.: 189 and 223; (9) SEQ ID NOs.: 190 and 223; (10) SEQ ID NOs.: 191 and 223; (11) SEQ ID NOs.: 192 and 223; (12) SEQ ID NOs.: 193 and 223; (13) SEQ ID NOs.: 194 and 223; (14) SEQ ID NOs.: 195 and 223; (15) SEQ ID NOs.: 196 and 223; (16) SEQ ID NOs.: 197 and 223; (17) SEQ ID NOs.: 198 and 223; (18) SEQ ID NOs.: 199 and 223; (19) SEQ ID NOs.: 200 and 223; (20) SEQ ID NOs.: 201 and 223; (21) SEQ ID NOs.: 202 and 223; (22) SEQ ID NOs.: 203 and 223; (23) SEQ ID NOs.: 204 and 223; (24) SEQ ID NOs.: 205 and 223; (25) SEQ ID NOs.: 206 and 223; (26) SEQ ID NOs.: 207 and 223; (27) SEQ ID NOs.: 208 and 223; (28) SEQ ID NOs.: 209 and 223; (29) SEQ ID NOs.: 182 and 226; (30) SEQ ID NOs.: 182 and 227; (31) SEQ ID NOs.: 182 and 229; (32) SEQ ID NOs.: 182 and 230; (33) SEQ ID NOs.: 182 and 231; (34) SEQ ID NOs.: 182 and 232; (35) SEQ ID NOs.: 182 and 233; (36) SEQ ID NOs.: 182 and 234; (37) SEQ ID NOs.: 182 and 235; (38) SEQ ID NOs.: 182 and 236; (39) SEQ ID NOs.: 182 and 237; (40) SEQ ID NOs.: 182 and 238; (41) SEQ ID NOs.: 182 and 239; (42) SEQ ID NOs.: 182 and 240; (43) SEQ ID NOs.: 182 and 241; (44) SEQ ID NOs.: 182 and 242; (45) SEQ ID NOs.: 182 and 243; (46) SEQ ID NOs.: 182 and 244; (47) SEQ ID NOs.: 182 and 245; (48) SEQ ID NOs.: 182 and 246; (49) SEQ ID NOs.: 182 and 247; (50) SEQ ID NOs.: 182 and 248; (51) SEQ ID NOs.: 182 and 249; (52) SEQ ID NOs.: 182 and 250; (53) SEQ ID NOs.: 182 and 251; (54) SEQ ID NOs.: 182 and 252; (55) SEQ ID NOs.: 182 and 253; (56) SEQ ID NOs.: 210 and 238; (57) SEQ ID NOs.: 211 and 238; (58) SEQ ID NOs.: 212 and 238; (59) SEQ ID NOs.: 210 and 240; (60) SEQ ID NOs.: 211 and 240; (61) SEQ ID NOs.: 212 and 240; (62) SEQ ID NOs.: 213 and 223; (63) SEQ ID NOs.: 214 and 223; (64) SEQ ID NOs.: 215 and 223; (65) SEQ ID NOs.: 216 and 223; (66) SEQ ID NOs.: 217 and 223; (67) SEQ ID NOs.: 218 and 223; (68) SEQ ID NOs.: 182 and 254; (69) SEQ ID NOs.: 213 and 254; (70) SEQ ID NOs.: 214 and 254; (71) SEQ ID NOs.: 215 and 254; (72) SEQ ID NOs.: 216 and 254; (73) SEQ ID NOs.: 217 and 254; (74) SEQ ID NOs.: 218 and 254; (75) SEQ ID NOs.: 182 and 255; (76)

SEQ ID NOs.: 213 and 255; (77) SEQ ID NOs.: 214 and 255; (78) SEQ ID NOs.: 215 and 255; (79) SEQ ID NOs.: 216 and 255; (80) SEQ ID NOs.: 217 and 255; (81) SEQ ID NOs.: 218 and 255; (82) SEQ ID NOs.: 182 and 256; (83) SEQ ID NOs.: 213 and 256; (84) SEQ ID NOs.: 214 and 256; (85) SEQ ID NOs.: 215 and 256; (86) SEQ ID NOs.: 216 and 256; (87) SEQ ID NOs.: 217 and 256; (88) SEQ ID NOs.: 218 and 256; (89) SEQ ID NOs.: 182 and 257; (90) SEQ ID NOs.: 213 and 257; (91) SEQ ID NOs.: 214 and 257; (92) SEQ ID NOs.: 215 and 257; (93) SEQ ID NOs.: 216 and 257; (94) SEQ ID NOs.: 217 and 257; (95) SEQ ID NOs.: 218 and 257; (96) SEQ ID NOs.: 182 and 258; (97) SEQ ID NOs.: 213 and 258; (98) SEQ ID NOs.: 214 and 258; (99) SEQ ID NOs.: 215 and 258; (100) SEQ ID NOs.: 216 and 258; (101) SEQ ID NOs.: 217 and 258; (102) SEQ ID NOs.: 218 and 258; (103) SEQ ID NOs.: 182 and 259; (104) SEQ ID NOs.: 213 and 259; (105) SEQ ID NOs.: 214 and 259; (106) SEQ ID NOs.: 215 and 259; (107) SEQ ID NOs.: 216 and 259; (108) SEQ ID NOs.: 217 and 259; (109) SEQ ID NOs.: 218 and 259; (110) SEQ ID NOs.: 182 and 260; (111) SEQ ID NOs.: 182 and 261; (112) SEQ ID NOs.: 182 and 262; (113) SEQ ID NOs.: 182 and 263; (114) SEQ ID NOs.: 182 and 264; (115) SEQ ID NOs.: 182 and 265; (116) SEQ ID NOs.: 182 and 266; (117) SEQ ID NOs.: 182 and 267; (118) SEQ ID NOs.: 182 and 268; (119) SEQ ID NOs.: 182 and 269; (120) SEQ ID NOs.: 182 and 270; (121) SEQ ID NOs.: 182 and 271; (122) SEQ ID NOs.: 182 and 272; (123) SEQ ID NOs.: 219 and 273; (124) SEQ ID NOs.: 191 and 274; (125) SEQ ID NOs.: 182 and 275; (126) SEQ ID NOs.: 220 and 277; (127) SEQ ID NOs.: 182 and 278; (128) SEQ ID NOs.: 182 and 279; (129) SEQ ID NOs.: 182 and 280; (130) SEQ ID NOs.: 182 and 281; (131) SEQ ID NOs.: 182 and 282; (132) SEQ ID NOs.: 221 and 228; (133) SEQ ID NOs.: 221 and 283; (134) SEQ ID NOs.: 182 and 284; (135) SEQ ID NOs.: 221 and 285; (136) SEQ ID NOs.: 182 and 286; (137) SEQ ID NOs.: 221 and 287; (138) SEQ ID NOs.: 221 and 288; (139) SEQ ID NOs.: 221 and 289; (140) SEQ ID NOs.: 182 and 290; (141) SEQ ID NOs.: 221 and 291; (142) SEQ ID NOs.: 182 and 292; (143) SEQ ID NOs.: 221 and 293; (144) SEQ ID NOs.: 221 and 294; (145) SEQ ID NOs.: 221 and 295; (146) SEQ ID NOs.: 182 and 296; (147) SEQ ID NOs.: 221 and 297; (148) SEQ ID NOs.: 182 and 298; (149) SEQ ID NOs.: 221 and 299; (150) SEQ ID NOs.: 221 and 300; (151) SEQ ID NOs.: 221 and 301; (152) SEQ ID NOs.: 182 and 302; (153) SEQ ID NOs.: 221 and 303; (154) SEQ ID NOs.: 182 and 304; (155) SEQ ID NOs.: 221 and 305; (156) SEQ ID NOs.: 182 and 306; (157) SEQ ID NOs.: 182 and 307; (158) SEQ ID NOs.: 182 and 308; (159) SEQ ID NOs.: 182 and 309; (160) SEQ ID NOs.: 220 and 310; (161) SEQ ID NOs.: 220 and 311; (162) SEQ ID NOs.: 182 and 228; (163) SEQ ID NOs.: 465 and 276; (164) SEQ ID NOs.: 466

and 276; (166) SEQ ID NOs.: 182 and 479; (167) SEQ ID NOs.: 465 and 479; (168) SEQ ID NOs.: 466 and 479; (169) SEQ ID NOs.: 182 and 480; (170) SEQ ID NOs.: 465 and 480; (171) SEQ ID NOs.: 466 and 480; (172) SEQ ID NOs.: 182 and 481; (173) SEQ ID NOs.: 182 and 482; (174) SEQ ID NOs.: 465 and 482; (175) SEQ ID NOs.: 466 and 482; (176) SEQ ID NOs.: 182 and 483; (177) SEQ ID NOs.: 182 and 484; (178) SEQ ID NOs.: 465 and 484; (179) SEQ ID NOs.: 466 and 484; (180) SEQ ID NOs.: 182 and 485; (181) SEQ ID NOs.: 182 and 486; (182) SEQ ID NOs.: 465 and 486; (183) SEQ ID NOs.: 466 and 486; (184) SEQ ID NOs.: 182 and 487; (185) SEQ ID NOs.: 182 and 488; (186) SEQ ID NOs.: 465 and 488; (187) SEQ ID NOs.: 466 and 488; (188) SEQ ID NOs.: 182 and 489; (189) SEQ ID NOs.: 465 and 489; (190) SEQ ID NOs.: 466 and 489; (191) SEQ ID NOs.: 182 and 491; (192) SEQ ID NOs.: 465 and 491; (193) SEQ ID NOs.: 466 and 491; (194) SEQ ID NOs.: 182 and 492; (195) SEQ ID NOs.: 465 and 492; (196) SEQ ID NOs.: 466 and 492; (197) SEQ ID NOs.: 182 and 493; (198) SEQ ID NOs.: 182 and 494; (199) SEQ ID NOs.: 465 and 494; (200) SEQ ID NOs.: 466 and 494; (201) SEQ ID NOs.: 182 and 277; (202) SEQ ID NOs.: 465 and 277; (203) SEQ ID NOs.: 466 and 277; (204) SEQ ID NOs.: 182 and 495; (205) SEQ ID NOs.: 465 and 495; (206) SEQ ID NOs.: 466 and 495; (207) SEQ ID NOs.: 182 and 496; (208) SEQ ID NOs.: 465 and 496; (209) SEQ ID NOs.: 466 and 496; (210) SEQ ID NOs.: 182 and 497; (211) SEQ ID NOs.: 465 and 497; (212) SEQ ID NOs.: 466 and 497; (213) SEQ ID NOs.: 182 and 498; (214) SEQ ID NOs.: 182 and 499; (215) SEQ ID NOs.: 465 and 499; (216) SEQ ID NOs.: 466 and 499; (217) SEQ ID NOs.: 182 and 500; (218) SEQ ID NOs.: 182 and 501; (219) SEQ ID NOs.: 465 and 501; (220) SEQ ID NOs.: 466 and 501; (221) SEQ ID NOs.: 182 and 502; (222) SEQ ID NOs.: 182 and 503; (223) SEQ ID NOs.: 182 and 504; (224) SEQ ID NOs.: 182 and 505; (225) SEQ ID NOs.: 182 and 506; (226) SEQ ID NOs.: 182 and 507; (227) SEQ ID NOs.: 182 and 508; (228) SEQ ID NOs.: 182 and 509; (229) SEQ ID NOs.: 182 and 510; (230) SEQ ID NOs.: 182 and 511; (231) SEQ ID NOs.: 182 and 512; (232) SEQ ID NOs.: 182 and 513; (233) SEQ ID NOs.: 182 and 514; (234) SEQ ID NOs.: 182 and 515; (235) SEQ ID NOs.: 467 and 223; (236) SEQ ID NOs.: 468 and 223; (237) SEQ ID NOs.: 469 and 223; (238) SEQ ID NOs.: 470 and 223; (239) SEQ ID NOs.: 471 and 223; (240) SEQ ID NOs.: 472 and 223; (241) SEQ ID NOs.: 473 and 223; (242) SEQ ID NOs.: 474 and 223; (243) SEQ ID NOs.: 475 and 223; (244) SEQ ID NOs.: 476 and 223; (245) SEQ ID NOs.: 182 and 516; (246) SEQ ID NOs.: 182 and 276; (247) SEQ ID NOs.: 182 and 569; (248) SEQ ID NOs.: 477 and 223; (249) SEQ ID NOs.: 477 and 278; (250) SEQ ID NOs.: 477 and 292; or (251) SEQ ID NOs.: 478 and 276.

[0019] In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 275, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 278, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 223, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 292, respectively. In certain embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 465 and 276, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 466 and 276, respectively. In certain embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 491, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 465 and 491, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 466 and 491, respectively. In certain embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 493, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 220 and 276, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 516, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 276, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 569, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 477 and 223, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 477 and 278, respectively. In some embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 477 and 292, respectively. In other embodiments, the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 478 and 276, respectively.

[0020] In some embodiments, the antibody or antigen-binding fragment thereof comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid

sequence selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569. In some embodiments, the antibody or antigen-binding fragment thereof comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278.

[0021] In some embodiments, the antibody further comprises a human IgG1 Fc region. In some instances, the human IgG1 Fc region is IgG1m17 (SEQ ID NO:348). In certain embodiments, the antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering): (i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330; (ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434; (iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434; (iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434; (v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or (vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434. In some embodiments, the antibody comprises a human kappa light chain constant region. In certain cases, the human kappa light chain constant region is Km3 (SEQ ID NO: 351).

[0022] In some embodiments, the antibody or antigen-binding fragment has improved, extended, enhanced or increased serum half-life in a mammal (*e.g.*, in a non-human primate, in a human) compared to other anti-HIV antibodies, such as Antibody A and/or Antibody B. In some embodiments, the antibody or antigen-binding fragment has a serum half-life in a human of at least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer. In some embodiments, the antibody or antigen-binding fragment has improved, enhanced, or

increased killing potency of HIV-infected cells compared to other anti-HIV antibodies such as Antibody A and/or Antibody B.

[0023] In another aspect, the disclosure provides an antibody comprising a heavy chain and a light chain, wherein the heavy chain and the light chain comprise any of the amino acid sequences set forth in Table X and XI, respectively.

[0024] In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 49, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 100, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 42 and 101, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 103, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 117, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 517 and 101, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 518 and 101, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 542, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 517 and 542, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 518 and 542, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 544, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 567, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 568, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 529 and 49, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 529 and 103, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 529 and 117, respectively. In some embodiments, the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 530 and 101, respectively. In some embodiments, antibody comprises a heavy chain (HC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid

sequence selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and a light chain (LC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 48-136 and 531-567. In certain embodiments, antibody comprises a heavy chain (HC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to the amino acid sequence set forth in SEQ ID NO: 529 and a light chain (LC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 103. In some embodiments, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or more, N-linked glycosylation sites in the VL are sialylated. In some embodiments, the N-linked glycosylation sites in the VL have a sialic acid occupancy (*e.g.*, a glycan comprising one or two terminal sialic acid residues) of at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or more.

[0025] In a related aspect, provided is an antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody or antigen-binding fragment thereof comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: (i) SEQ ID NOs.: 159, 138, 139, 140, 141, and 142, respectively; (ii) SEQ ID NOs.: 137, 160, 139, 140, 141, and 142, respectively; (iii) SEQ ID NOs.: 137, 161, 139, 140, 141, and 142, respectively; (iv) SEQ ID NOs.: 137, 162, 139, 140, 141, and 142, respectively; (v) SEQ ID NOs.: 137, 163, 139, 140, 141, and 142, respectively; (vi) SEQ ID NOs.: 137, 138, 164, 140, 141, and 142, respectively; (vii) SEQ ID NOs.: 159, 138, 164, 140, 141, and 142, respectively; (viii) SEQ ID NOs.: 137, 138, 139, 140, 165, and 142, respectively; (ix) SEQ ID NOs.: 137, 138, 139, 140, 166, and 142, respectively; (x) SEQ ID NOs.: 137, 138, 139, 140, 167, and 142, respectively; (xi) SEQ ID NOs.: 137, 138, 139, 140, 168, and 142, respectively; (xii) SEQ ID NOs.: 137, 138, 154, 140, 141, and 142, respectively, or (xiii) SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and wherein at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or

more, N-linked glycosylation sites in the VL are sialylated. In certain embodiments, is an antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody or antigen-binding fragment thereof comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and wherein at least 50%, at least 60%, at least 70%, least 80%, at least 85%, at least 90%, or more, N-linked glycosylation sites in the VL are sialylated. In certain embodiments, is an antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody or antigen-binding fragment thereof comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, wherein comprises a VH with the following amino acids at the indicated positions (position numbering according to Kabat): asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 72, phenylalanine at position 76, and phenylalanine at position 74a, and tyrosine at position 99 (*e.g.*, asparagine at position 28, arginine at position 30, tyrosine at position 32, histidine at position 73, phenylalanine a position 76 and tyrosine at position 98, wherein the amino acid positions are with respect to SEQ ID NO: 477), and wherein at least 50%, at least 60%, at least 70%, least 80%, at least 85%, at least 90%, or more, N-linked glycosylation sites in the VL are sialylated. In some embodiments, the N-linked glycosylation sites in the VL have a sialic acid occupancy (*e.g.*, one or two terminal sialic acid residues) of at least 40%, at least 50%, at least 60%, at least 70%, least 80%, at least 85%, at least 90%, or more. In some embodiments, the asparagine at VL amino acid position 72 according to Kabat numbering (N72) is sialylated. In some embodiments, the sialylated N-linked glycosylation sites in the VL comprise from 1 to 5 sialic acid residues, *e.g.*, from 1 to 4 sialic acid residues, *e.g.*, from 1 to 3 sialic acid residues, *e.g.*, from 1 to 2 sialic acid residues. In some embodiments, the VL are sialylated with N-acetylneuraminic acid (NANA). In some embodiments, the sialic acid residues are present in biantennary structures. In some embodiments, the sialic acid residues are present in complex N-linked glycan structures. In some embodiments, the sialic acid residues are present in hybrid N-linked glycan structures.

[0026] In a further aspect, provided is a bispecific antibody comprising: a first antigen binding arm that binds to gp120, the first antigen binding arm comprising: (i) the VH CDRs 1-3 and the VL CDRs 1-3; or (ii) the VH and the VL of any one or claims 1 to 63; and a second antigen binding arm binding to a second antigen. In certain embodiments, is a bispecific antibody comprising: a first antigen binding arm that binds to gp120, the first antigen binding arm comprising the VH CDRs 1-3 and the VL CDRs 1-3 as set forth in SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and a second antigen binding arm binding to a second antigen. In certain embodiments, is a bispecific antibody comprising: a first antigen binding arm that binds to gp120, the first antigen binding arm comprising the VH and the VL comprising the amino acid sequences set forth in SEQ ID NOs: 477 and 278, respectively, and a second antigen binding arm binding to a second antigen. In some embodiments, the second antigen is selected from the group consisting of CD3, FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16); CD89, CCR5, CD4, gp41, killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3), killer cell lectin like receptor C1 (KLRC1), killer cell lectin like receptor C2 (KLRC2), killer cell lectin like receptor C3 (KLRC3), killer cell lectin like receptor C4 (KLRC4), killer cell lectin like receptor D1 (KLRD1), killer cell lectin like receptor K1 (KLRK1), natural cytotoxicity triggering receptor 3 (NCR3 or NKp30), natural cytotoxicity triggering receptor 2 (NCR2 or NK-p44), natural cytotoxicity triggering receptor 1 (NCR1 or NK-p46), CD226 (DNAM-1), cytotoxic and regulatory T cell molecule (CRTAM or CD355), signaling lymphocytic activation molecule family member 1 (SLAMF1), CD48 (SLAMF2), lymphocyte antigen 9 (LY9 or SLAMF3), CD244 (2B4 or SLAMF4), CD84 (SLAMF5), SLAM family member 6 (SLAMF6 or NTB-A), SLAM family member 7 (SLAMF7 or CRACC), CD27 (TNFRSF7), semaphorin 4D (SEMA4D or CD100), and CD160 (NK1), and a second epitope of gp120.

[0027] The disclosure also provides a pharmaceutical composition comprising an antibody or antigen-binding fragment described herein, and a pharmaceutically acceptable carrier.

[0028] In certain embodiments, the pharmaceutical composition further comprises a second agent (*e.g.*, one or more additional agents) for treating an HIV infection. In some cases, the pharmaceutical composition further comprises a latency reversing agent (LRA) or an immunostimulatory agent, *e.g.*, an agonist of a toll-like receptor (TLR), *e.g.*, an agonist of TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, and/or TLR10. In some embodiments, the LRA is a TLR7 agonist or a TLR8 agonist. In certain instances, the TLR7 agonist is selected from the group consisting of vesatolimod, imiquimod, and resiquimod. In some embodiments, the pharmaceutical composition further comprises an antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV. In some embodiments, the pharmaceutical composition further comprises a second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, wherein the second antibody or antigen-binding fragment thereof does not compete with the antibody or antigen-binding fragment, as described herein, for binding to gp120. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, competes with or comprises VH and VL variable domains of a broadly neutralizing antibody (bNAb) against HIV. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 selected from the group consisting of: (i) third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan; (ii) second variable loop (V2) and/or Env trimer apex; (iii) gp120/gp41 interface; or (iv) silent face of gp120. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722, PGT-121.60, PGT-121.66, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137, PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the second variable loop (V2) and/or Env trimer apex and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PG9, PG16, PGC14, PGG14, PGT-142, PGT-143, PGT-144, PGT-145, CH01, CH59, PGDM1400, CAP256, CAP256-

VRC26.08, CAP256-VRC26.09, CAP256-VRC26.25, PCT64-24E and VRC38.01. In some embodiments, the second antibody or antigen-binding fragment binds to an epitope or region of gp120 in the gp120/gp41 interface and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PGT-151, CAP248-2B, 35O22, 8ANC195, ACS202, VRC34 and VRC34.01. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp120 silent face and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC-PG05 and SF12. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER). In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER) and competes with or comprises VH and VL regions from an antibody selected from the group consisting of 10E8, 10E8v4, 10E8-5R-100cF, 4E10, DH511.11P, 2F5, 7b2, and LN01. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp41 fusion peptide and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC34 and ACS202. In some embodiments, the second or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of PGT121.60 or PGT121.66. In certain cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of SEQ ID NO: 443 and/or SEQ ID NO: 447. In other cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO: 454 and the VL within SEQ ID NO: 455. In yet other cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO: 454 and the VL within SEQ ID NO: 456.

[0029] In another aspect, the disclosure provides nucleic acids, nucleotides, or polynucleotides encoding an antibody or antigen-binding fragment disclosed herein. In some embodiments, the nucleic acid or nucleic acids comprise DNA, cDNA or mRNA. In some embodiments, the nucleic acid or nucleic acids encode a VH selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at

least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 572-581; and encode a VL selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 582-595. In some embodiments, the nucleic acid or nucleic acids encode a HC selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 596-605; and encode a LC selected from the group consisting of SEQ ID NOs: 48-136 and 531-567 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 606-619. In another aspect, the disclosure provides an expression vector or expression vectors comprising the nucleic acid or nucleic acids operably linked to a regulatory sequence. In some embodiments, the expression vector or expression vectors comprise a plasmid vector or a viral vector. Further provided are pharmaceutical compositions comprising the nucleic acid or nucleic acids, or the expression vector or expression vector, as described herein, and a pharmaceutically acceptable carrier. Further provided are lipid nanoparticles comprising the nucleic acid or nucleic acids, or the expression vector or expression vector, as described herein.

[0030] In yet another aspect, the disclosure provides a host cell, or population of host cells, comprising the nucleic acid or nucleic acids, or the expression vector or expression vectors, described herein. In some embodiments, the cell or population of cells comprises a eukaryotic cell. In some embodiments, the cell or population of cells comprises a mammalian cell, a human cell, a hamster cell, an insect cell, a plant cell or a yeast cell. In some embodiments, the mammalian cell is a Chinese Hamster Ovary (CHO) cell or a human cell, *e.g.*, a human embryonic kidney cell or a human B-cell. In some embodiments, the cell predominantly sialylates N-linked glycosylation sites in the variable domains (Fv) of the expressed antigen binding molecules, *e.g.*, expressed antibodies or antigen binding fragments. In some embodiments, the cell sialylates at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least

80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the variable domains (Fv) of expressed antibodies or antigen-binding fragments. In some embodiments, the cell sialylates at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the VL of expressed antibodies or antigen-binding fragments. In some embodiments, the asparagine at VL amino acid position 72 according to Kabat numbering (N72) is sialylated. In some embodiments, the sialylated N-linked glycosylation sites in the VL comprise from 1 to 5 sialic acid residues, *e.g.*, from 1 to 4 sialic acid residues, *e.g.*, from 1 to 3 sialic acid residues, *e.g.*, from 1 to 2 sialic acid residues. In some embodiments, the VL are sialylated with N-acetylneuraminic acid (NANA). In some embodiments, the sialic acid residues are present in biantennary structures. In some embodiments, the sialic acid residues are present in complex N-linked glycan structures. In some embodiments, the sialic acid residues are present in hybrid N-linked glycan structures.

[0031] In yet another aspect, provided herein are antigen-binding fragments of the antibodies described herein. In some embodiments, the antigen-binding fragments are selected from the group consisting of a scFv, sc(Fv)₂, Fab, F(ab)₂, Fab', F(ab')₂, Facb or Fv fragment. Further provided is a chimeric antigen receptor (CAR) including an antigen-binding antibody fragment as described herein. In certain embodiments, the CAR is expressed on a T-cell, a B-cell, a macrophage or a NK cell. Further provided is a CAR T-cell including a CAR as described herein. In certain embodiments, the T-cell is a CD4⁺ T-cell, a CD8⁺ T-cell, or a combination thereof. In certain embodiments, the cell is administered to a subject. In certain embodiments, the cell is autologous. In certain embodiments, the cell is allogeneic.

[0032] In yet another aspect, provided herein is a method of producing an antibody or antigen-binding fragment thereof described herein. The method involves culturing the host cell in a cell culture and isolating the antibody or antigen-binding fragment from the cell culture. In certain cases, the method further involves formulating the antibody or antigen-binding fragment into a sterile pharmaceutical composition suitable for administration to a human subject.

[0033] In another aspect, the disclosure provides a method of treating or preventing HIV in a human subject in need thereof. The method involves administering to the subject an

effective amount of an antibody or antigen-binding fragment thereof, or a pharmaceutical composition described herein.

[0034] In some embodiments, the method further comprises administering to the subject a second agent (*e.g.*, one or more additional agents) for treating an HIV infection. In some cases, the method comprises administering to the subject a TLR7 agonist. In certain instances, the TLR7 agonist is selected from the group consisting of vesatolimod, imiquimod, and resiquimod. In some embodiments, the method further comprises administering to the subject an antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV. In some embodiments, the method further comprises administering a second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, wherein the second antibody or antigen-binding fragment thereof does not compete with the antibody or antigen-binding fragment, as described herein, for binding to gp120. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, competes with or comprises VH and VL variable domains of a broadly neutralizing antibody (bNAb) against HIV. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 selected from the group consisting of: (i) third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan; (ii) second variable loop (V2) and/or Env trimer apex; (iii) gp120/gp41 interface; or (iv) silent face of gp120. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722, GS-9722, PGT-121.60, PGT-121.66, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137, PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the second variable loop (V2) and/or Env trimer apex and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PG9, PG16, PGC14, PGG14, PGT-142, PGT-143, PGT-144, PGT-145, CH01, CH59, PGDM1400, CAP256,

CAP256-VRC26.08, CAP256-VRC26.09, CAP256-VRC26.25, PCT64-24E and VRC38.01. In some embodiments, the second antibody or antigen-binding fragment binds to an epitope or region of gp120 in the gp120/gp41 interface and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PGT-151, CAP248-2B, 35O22, 8ANC195, ACS202, VRC34 and VRC34.01. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp120 silent face and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC-PG05 and SF12. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER). In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER) and competes with or comprises VH and VL regions from an antibody selected from the group consisting of 10E8, 10E8v4, 10E8-5R-100cF, 4E10, DH511.11P, 2F5, 7b2, and LN01. In some embodiments, the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp41 fusion peptide and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC34 and ACS202. In some embodiments, the second or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of PGT121.60 or PGT121.66. In certain cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of SEQ ID NO: 443 and/or SEQ ID NO: 447. In other cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO: 454 and the VL within SEQ ID NO: 455. In yet other cases, the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO: 454 and the VL within SEQ ID NO: 456. In some embodiments, the antibody or antigen-binding fragments, as described herein, are co-administered to a human subject with an anti-HIV vaccine. In various embodiments, the anti-HIV vaccine comprises a viral vaccine. In certain embodiments, the viral vaccine is from a virus selected from the group consisting of an arenavirus, an adenovirus, a poxvirus, and a rhabdovirus.

[0035] In another aspect, the disclosure relates to a method of inhibiting HIV in a human subject in need thereof. The method involves administering to the subject an effective

amount of an antibody or antigen-binding fragment thereof, or a pharmaceutical composition described herein.

[0036] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the exemplary methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present application, including definitions, will control. The materials, methods, and examples are illustrative only and not intended to be limiting.

Brief Description of the Drawings

[0037] Figure 1 illustrates the results of an ADCC reporter assay conducted on the antibody A-1 stress panel. The sample stressed at 37 °C in pH 5.9 formulation buffer for six weeks showed a large reduction in activity relative to other samples.

[0038] Figure 2 illustrates kinetics of W74a oxidation over time as measured in the stress panel. Diamond: Antibody A-1, 25°C, pH 5.9. Open circle: Antibody A-1, 37°C, pH 5.9. Open triangle: Antibody A-1, 37°C, pH 7.4. The degree of oxidation is greatest in the pH 5.9 sample stressed at 37°C for 6 weeks, suggesting that W74a oxidation may be the source of potency loss observed in this condition. In addition to the significant oxidation at heavy chain W74a observed in pH 5.9 conditions, a steady percentage of deamidation at light chain position N26 was observed on the constructs coming out of cell culture and increased further at pH 7.4 incubation conditions.

[0039] Figure 3 illustrates kinetics of N26 deamidation over time as measured in the stress panel (include oxidation to aspartic acid, isoaspartic acid, and aspartyl succinimide intermediate). Diamond: Antibody A-1, 25°C, pH 5.9. Open circle: Antibody A-1, 37°C, pH 5.9. Open triangle: Antibody A-1, 37°C, pH 7.4. The degree of deamidation was greatest at the pH 7.4 sample stressed at 37°C for 6 weeks.

[0040] Figure 4 illustrates a dot plot representation of the neutralization profile of seven mAb variants. Antibodies were screened against a panel of 152 patient-derived HIV-1 pseudotyped with Env from subtype B plasma viral clones (n=133) and isolates (n=19). Each dot represents neutralization IC₉₅ for one virus. In parentheses (Breadth/Median

IC95). Breadth represents % viruses neutralized with an IC95 $\leq$ 50 mg/mL. Median IC95 values calculated using viruses with IC95 $\leq$ 50 mg/mL. (1) Antibody A-1 (89%/2.66 μ g/mL); (2) 1.1.90-1 (86%/2.59 μ g/mL); (3) 1.1.64-1 (92%/2.25 μ g/mL); (4) 1.1.10-1 (86%/1.93 μ g/mL); (5) 1.52.1-1 (83%/3.66 μ g/mL); (6) 1.52.90 (78%/4.42 μ g/mL); (7) 1.1.138-1 (82%/2.59 μ g/mL).

[0041] Figure 5 illustrates a dot plot representation of the neutralization profile of three mAbs. Antibodies were screened against a panel of 142 HIV-1 pseudotyped with Env from subtype B plasma isolates. In parentheses (Breadth/Median IC95), defined the same as for Figure 4. Each dot represents neutralization IC95 for one virus. (1) Antibody A (87%/1.72 μ g/mL); (2) Antibody A-1 (87%/1.09 μ g/mL); (3) 1.52.64-1 (86%/2.0 μ g/mL).

[0042] Figure 6 illustrates that mutations in the IgG1 Fc that enhance effector cell killing activity (*e.g.*, aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330 according to EU number (DEAL)) can shorten serum half-life *in vivo*. Such shortened serum half-life can be partially or wholly reversed by also incorporating mutations in the IgG1 Fc that enhance FcRn binding (*e.g.*, leucine at position 428, and serine at position 434 according to EU numbering (LS)). Depicted are illustrative dose normalized pharmacokinetic profiles for PGT121-WT (circle), PGT121-DEAL (triangle), PGT121.60 (square), PGT121-LS (diamond), and A-1 (solid circle) dosed IV to naïve cynomolgus monkeys (n=3). Each symbol is the measured mean ($\pm$ SD) serum concentration.

[0043] Figure 7 illustrates pharmacokinetic profiles for Antibody A (triangle), Antibody A-1 Lot 14 (circle), Antibody A-1 Lot 22 (open triangle), Antibody A-1 Lot 3 (open circle), Antibody A-1 Lot 10 (square), and Antibody A-1 Lot 7 (open square) following intravenous (IV) dosing to naïve male cynomolgus monkeys (n=3). Each symbol is the measured mean ($\pm$ SD) serum concentration.

[0044] Figure 8 illustrates mean serum ($\pm$ SD) concentration-time profiles of three lots of 1.52.64-1 following IV administrations to naïve male and female cynomolgus monkeys (n=3). Lot 4 (open square) was administered at 0.5 mg/kg slow IV bolus, while Lot 18-PP21 (open circle) and Lot 14525-32 (circle) were administered at 30 mg/kg via a 30 minute IV infusion. Each symbol is the measured mean ($\pm$ SD) serum concentration.

Detailed Description

[0045] This disclosure provides antibodies that target human immunodeficiency virus (HIV). The antibodies described herein bind to HIV envelope (Env) protein gp120 (gp120). In some embodiments, these are HIV neutralizing antibodies. In certain embodiments, these antibodies broadly neutralize HIV.

[0046] HIV-1 is the main family of HIV and accounts for 95% of all infections worldwide. HIV-2 is mainly seen in a few West African countries. HIV viruses are divided into specific groups, M, N, O and P, of which M is the “major” group and responsible for majority of HIV/AIDS globally. Based on their genetic sequence, Group M is further subdivided into subtypes (also called clades) with prevalence in distinct geographical locations.

[0047] A Group M “subtype” or “clade” is a subtype of HIV-1 group M defined by genetic sequence data. Examples of Group M subtypes include Subtypes A-K. Some of the subtypes are known to be more virulent or are resistant to different medications. There are also “circulating recombinant forms” or CRFs derived from recombination between viruses of different subtypes, which are each given a number. CRF12_{BF}, for example, is a recombination between subtypes B and F. Subtype A is common in West Africa. Subtype B is the dominant form in Europe, the Americas, Japan, Thailand, and Australia. Subtype C is the dominant form in Southern Africa, Eastern Africa, India, Nepal, and parts of China. Subtype D is generally only seen in Eastern and central Africa. Subtype E has never been identified as a nonrecombinant, only recombined with subtype A as CRF01_{AE}. Subtype F has been found in central Africa, South America and Eastern Europe. Subtype G (and the CRF02_{AG}) have been found in Africa and central Europe. Subtype H is limited to central Africa. Subtype I was originally used to describe a strain that is now accounted for as CRF04_{cpx}, with the cpx for a “complex” recombination of several subtypes. Subtype J is primarily found in North, Central and West Africa, and the Caribbean. Subtype K is limited to the Democratic Republic of Congo and Cameroon. These subtypes are sometimes further split into sub-subtypes such as A1 and A2 or F1 and F2. In 2015, the strain CRF19, a recombinant of subtype A, subtype D, and subtype G, with a subtype D protease was found to be strongly associated with rapid progression to AIDS in Cuba.

[0048] This disclosure provides neutralizing antibodies (*e.g.*, broadly neutralizing Abs) that target the gp120 polypeptide on the surface of HIV-infected cells. Without being bound to any hypothesis, neutralizing antibodies against viral envelope proteins may

provide adaptive immune defense against HIV-1 exposure by blocking the infection of susceptible cells. Broad neutralization indicates that the antibodies can neutralize HIV-1 isolates from different clades. Thus, the antibodies encompassed by this disclosure have cross-clade binding activity.

HIV envelope glycoprotein gp120

[0049] Envelope glycoprotein gp120 (or gp120) is a 120 kDa glycoprotein that is part of the outer layer of HIV. It presents itself as viral membrane spikes consisting of three molecules of gp120 linked together and anchored to the membrane by gp41 protein. Gp120 is essential for viral infection as it facilitates HIV entry into the host cell through its interaction with cell surface receptors. These receptors include DC-SIGN, Heparan Sulfate Proteoglycan, the CD4 receptor, C-C motif chemokine receptor 5 (CCR5) and C-X-C motif chemokine receptor 4 (CXCR4). Binding to CD4 on helper T-cells induces the start of a cascade of conformational changes in gp120 and gp41 that lead to the fusion of the virus with the host cell membrane.

[0050] Gp120 is encoded by the HIV *env* gene. The *env* gene encodes a gene product of around 850 amino acids. The primary *env* product is the protein gp160, which gets cleaved to gp120 (about 480 amino acids) and gp41 (about 345 amino acids) in the endoplasmic reticulum by the cellular protease furin.

[0051] The amino acid sequence of an exemplary gp160 polypeptide of HIV clone WITO is provided below (the V3 hypervariable loop is boldened and the N332 potential N-linked glycosylation site is boldened and underlined):

```
MKVMGTTKKNYQHLWRWGIMLLGMLMMSSAAEQLWVTVYYGVPVWREANTTLFCASDAKA
YDTEVHNVWATHACVPTDPNPQEVVMGNVTEDFNMWKNMVEQMHEDIISLWDQSLKPC
VKLTPLCVTLHCTNVTISSTNGSTANVTMREEMKNCSFNNTTIVIRDKIQKEYALFYKLD
IVPIEGKNTNTSYRLINCNTSVITQACPKVSFEPIPIHYCAPAGFAILKCNNKTFNGKG
PCRNVTSTVQCTHGIKPVVSTQLLLNGSLAEEDIIRSENFTNNGKNIIVQLKEPVKINC
TRPGNNTRRSINIGPGRAFYATGAIIGDIRKAHCNISTEQWNNNTLTQIVDKLREQFGNK
TIIIFNQSSGGDPEVVMHTFNCGGEFFYCNSTQLEFNSTWFNNGTSTWNSTADNITLPCRI
KQVINMWQEVGKAMYAPPPIRGQIDCSSNITGLILTRDGGSNSSQNETFRPGGGMKDNW
RSELYKYKVKIEPLGIAPTRAKRRVVQREKRAVTLGAVFLGFLGAAGSTMGAASLTLT
VQARLLLSGIVQQSNLLRAIEAQQHMLQLTVWGIKQLQARVLAIERYLKDQQLLGIWG
CSGKLICTTTVPWNTSWSNKSVDYIWNMTWMQWEREIDNYTGFIYTLIEESQNQQEKN
ELELLELDKWASLWNWFNITNWLWYIKLFIMIIGGLVGLRIVCAVLSIVNRVRQGYSP
SFQTRLNPNRGPDRPEETEGEGGERDRDRSARLVNGFLAIIWDDLRLSLCLFSYHRLRDL
LLIVARVVEILGRRGWEILKYWWNLLKYWSQELKNSAVSLLNVTATAVAEGTDRVIEIV
QRAVRAILHIPTRIRQGFERALL (SEQ ID NO: 343)
```

[0052] The amino acid sequence of an exemplary gp120 polypeptide is provided below (the V3 hypervariable loop is boldened and the N332 potential N-linked glycosylation site is boldened and underlined):

AEQLWVTVYYGVPVWREANTTLFCASDAKAYDTEVHNWATHACVPTDPNPQEVVMGNV
TEDFNMWKNMVEQMHEDIISLWDQSLKPCVKLTPLCVTLHCTNVTISSTNGSTANVTM
REEMKNCSFNNTTIVIRDKIQKEYALFYKLDIVPIEGKNTNTSYRLINCNTSVITQACPK
VSFEPIPIHYCAPAGFAILKCNNKTFNGKGPCRNVSTVQCTHGIKPVVSTQLLLNGSLA
EEDIIIRSENFTNNGKNIIVQLKEPVKIN**CTRPGNNTRRSINIGPGRAFYATGAIIGDI**
RKAHCNISTEQWNNLTQIVDKLREQFGNKTIIFNQSSGGDPEVVMHTFNCGGEFFYCN
STQLENSTWFNNGTSTWNSTADNITLPCRIKQVINMWQEVGKAMYAPPPIRGQIDCSSNI
TGLILTRDGGSNSSQNETFRPGGGMKDNWRSELYKYKVVKIEPLGIAPTRAKRRVVQR
EKR (SEQ ID NO: 344)

[0053] The amino acid sequence of another exemplary gp120 polypeptide (see, bioafrica.net/proteomics/ENV-GP120prot.html) is provided below (the V3 hypervariable loop is boldened and the N332 potential N-linked glycosylation site is boldened and underlined):

TEKLWVTVYY GVPVWKEATT TLFCASDAKA YDTEVHNVWA THACVPTDPN
PQEVVLVNVT ENFNMWKNM VEQMHEDIIS LWDQSLKPCV KLTPLCVSLK
CTDLKNDTNT NSSSGRMIME KGEIKNCSFN ISTSIRGKVQ KEYAFFYKLD
IIPIDNDTTS YKLTSCNTSV ITQACPKVSF EPIPIHYCAP AGFAILKCNN
KTFNGTGPCT NVSTVQCTHG IRPVVSTQLL LNGSLAEEEV VIRSVNFTDN
AKTIIIVQLNT SVEIN**CTRPN NNTRKRIRIQ RGPGRAFVTI GKIGNMRQAH**
CNISRAKWNN TLKQIASKLR EQFGNNKTII FKQSSGGDPE IVTHSFNCGG
EFFYCNSTQL FNSTWFNSTW STEGSNNTEG SDTITLPCRI KQIINMWQKV
GKAMYAPPIS GQIRCSSNIT GLLLTRDGGN SNESEIFRP GGDMRDNR
SELYKYKVVK IEPLGVAPTK AKRRVVQREK R (SEQ ID NO: 345)

[0054] Genomic diversity among independent human immunodeficiency virus type 1 (HIV-1) isolates, to a lesser degree among sequential isolates from the same patients, and even within a single patient isolate is a well-known feature of HIV-1. Although this sequence heterogeneity is distributed throughout the genome, most of the heterogeneity is located in the *env* gene. Comparison of predicted amino acid sequences from several different isolates has shown that sequence heterogeneity is clustered in five hypervariable regions (designated V1 through V5) of the surface glycoprotein, gp120. The V3 region, although only 35 amino acids long, exhibits considerable sequence variability. In spite of this variability, the V3 region includes determinants that mediate interactions with CD4⁺ cells. The increase in gp120 variability results in higher levels of viral replication, suggesting an increase in viral fitness in individuals infected by diverse HIV-1 variants. Without being bound to theory, the higher levels of viral replication may be due to host immune response pressure (*e.g.*, immune response escape) and/or to

adaptation to each individual host to maximize the rate of virus replication. Variability in potential N-linked glycosylation sites (PNGSs) also result in increased viral fitness. PNGSs allow for the binding of long-chain carbohydrates to the high variable regions of gp120. Thus, the number and precise location of PNGSs in *env* might affect the fitness of the virus, or the replication capacity of each virus variant, by providing more or less sensitivity to host immune responses, particularly the neutralizing antibodies.

[0055] A consensus sequence of the V3 region of gp120 (Milich et al., *J. Virol.*, 67(9):5623-5634 (1993)) is provided below:

CTRPNNNTRKSIHIGPGRAFYTGTGEIIGDIRQAH (SEQ ID NO: 346).

[0056] Antibody variants described herein bind to the CD4 binding site (CD4bs) of HIV gp120. The CD4 binding site (CD4bs) involves structurally conserved sites located within the β 1- α 1, loop D, β 20- β 21 (bridging sheet) and β 24- α 5 of gp120, which determine the CD4 binding and are involved in the epitopes of CD4bs-directed antibodies (Qiao, *et al.*, *Antiviral Res.* 2016 Aug;132:252-61). The CD4bs of gp120 forms conformational epitopes recognized by anti-CD4bs antibodies involving one or more amino acid residues selected from Thr278, Asp279, Ala281, Thr283, Asp368, Trp427, Glu460, Ser461, Glu462, Leu452, Leu453 and Arg476. The amino acid residues and position numbering is with reference to HXB2 subtype B HIV-1 isolate, which corresponds to residues 1-511 of NCBI Ref Seq No. NP_057856.1, provided below. Residues Thr278, Asp279, Asn280, Ala281, Thr283, Asp368, Trp427, Leu452, Leu453, Gly459, Glu464, Ser465, Glu466, Ile467, Gly472, Gly473 and Arg476, which can contribute to the gp120 CD4bs, are boldened and underlined:

MRVKEKYQHLWRWGWRWGTMLLGMLMICSATEKLWVTVYYGVPVWKEATTTLFCASDAK
 AYDTEVHNWATHACVPTDPNPQEVVLNVNVTENFNMWKNDMVEQMHEDIISLWDQSLKP
 CVKLTPLCVSLKCTDLKNDTNTNSSSGRMIMEKGEIKNCSFNISTSIRGKVQKEYAFFY
 KLDIIPIDNDTTSYKLTSCNTSVITQACPKVSFEPIPIHYCAPAGFAILKCNNKTFNGT
 GPCTNVSTVQCTHGIRPVVSTQLLLNGSLAEEEVVIRSVNF**TDNAKT**IIVQLNTSVEIN
 CTRPNNNTRKRIRIQRGPGRAVFTIGKIGNMRQAHCNISRAKWNNTLKQIASKLREQFG
 NNKTIIFKQSSGG**D**PEIVTHSFNCGGEFFYCNSTQLFNSTWFNSTWSTEGSNNTGSDT
 ITLPCRIKQIINM**W**QKVGKAMYAPPISGQIRCSSNITG**LLL**TRDG**G**NSNN**ESEI**FRPG**G**
GDMRDNWRSELYKVKVVKIEPLGVAPTKAKRRVVQREKR (SEQ ID NO: 571).

[0057] Tridimensional models depicting amino acid residues contributing to the gp120 CD4bs are provided, *e.g.*, in Canducci, *et al.*, *Retrovirology*. 2009 Jan 15;6:4; Falkowska, *et al.*, *J. Virol.* 2012 Apr;86(8):4394-403; and Li, *et al.*, *J. Virol.* 2012

Oct;86(20):11231–41; Gristick, *et al.*, *Nat Struct Mol Biol.* 2016 Oct;23(10):906-915; Kwon, *et al.*, *Nat Struct Mol Biol.* 2015 Jul;22(7):522-31; Liu, *et al.*, *Nat Struct Mol Biol.* 2017 Apr;24(4):370-378; Chen, *et al.*, *Science.* 2009 Nov 20;326(5956):1123-7 and Lyumkis, *et al.*, *Science.* 2013 Dec 20;342(6165):1484-90. In some embodiments, the antibody variants described herein compete with anti-CD4bs antibodies b12, CH103, 1NC9, 12A12, VRC01, VRC07-523, N6, 3BNC117, NIH45-46 and/or PGV04 (VRC-PG04) for binding to gp120 CD4bs. In some embodiments, the antibody variants described herein bind to an overlapping or identical epitope to the epitope bound by anti-CD4bs antibodies b12, CH103, 1NC9, 12A12, VRC01, VRC07-523, N6, 3BNC117, NIH45-46 and/or PGV04 (VRC-PG04).

Anti-gp120 Antibodies

[0058] This disclosure provides anti-gp120 antibodies. In certain embodiments, these antibodies bind to HIV-1 antigens expressed on a cell surface and eliminate or kill the infected cell.

[0059] In certain embodiments, these antibodies are neutralizing antibodies (*e.g.*, monoclonal) that target HIV-1. A “neutralizing antibody” is one that neutralizes the ability of HIV to initiate and/or perpetuate an infection in a host and/or in target cells *in vitro*. The disclosure provides neutralizing monoclonal human antibodies, wherein the antibody recognizes an antigen from HIV, *e.g.*, a gp120 polypeptide. In certain embodiments, a “neutralizing antibody” may inhibit the entry of HIV-1 virus, *e.g.*, SF162 and/or JR-CSF, with a neutralization index >1.5 or >2.0 (Kostrikis LG *et al.*, *J. Virol.*, 70(1): 445-458 (1996)).

[0060] In some embodiments, these antibodies are broadly neutralizing antibodies (*e.g.*, monoclonal) that target HIV-1. By “broadly neutralizing antibodies” are meant antibodies that neutralize more than one HIV-1 virus species (from diverse clades and different strains within a clade) in a neutralization assay. A broadly neutralizing antibody may neutralize at least 2, 3, 4, 5, 6, 7, 8, 9 or more different strains of HIV-1, the strains belonging to the same or different clades. In some embodiments, a broad neutralizing antibody may neutralize multiple HIV-1 species belonging to at least 2, 3, 4, 5, or 6 different clades. In certain embodiments, the inhibitory concentration of the antibody may be less than about 0.0001 µg/mL, less than about 0.001 µg/mL, less than about 0.01 µg/mL, less than about 0.1 µg/mL, less than about 0.5 µg/mL, less than about 1.0 µg/mL, less than about 5 µg/mL, less than about 10 µg/mL, less than about 25

µg/mL, less than about 50 µg/mL, or less than about 100 µg/mL to neutralize about 50% of the input virus in the neutralization assay.

[0061] In certain embodiments, these antibodies show broad and potent activity and fall within the group of highly active agonistic anti-CD4 binding site antibodies (HAADs). Such antibodies mimic binding of the host receptor CD4 protein to gp120. In certain embodiments, the antibodies or antigen-binding fragments thereof comprise in their heavy chain variable region tryptophan at position 50; asparagine at position 58; arginine at position 71; and tryptophan at position 100 (position numbering according to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise in their light chain variable region tryptophan or phenylalanine at position 67; and glutamic acid at position 96 (position numbering according to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according to Kabat). In certain instances, the light chain variable region includes an N-linked glycosylation site in framework region 3. In certain embodiments, the antibodies or antigen-binding fragments thereof comprise in their heavy chain variable region tryptophan at position 50; asparagine at position 58; arginine at position 71; and tryptophan at position 100; and comprise in their light chain variable region tryptophan or phenylalanine at position 67; and glutamic acid at position 96 (position numbering according to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise in their heavy chain variable region tryptophan at position 50; asparagine at position 58; arginine at position 71; and tryptophan at position 100; and comprise in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and further comprise in their heavy chain variable region tryptophan at position 50; asparagine at position 58; arginine at position 71; and tryptophan at position 100; and comprise in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and further comprise in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according

to Kabat). In certain embodiments, the antibodies or antigen-binding fragments thereof comprise VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and further comprise in their heavy chain variable region tryptophan at position 50; asparagine at position 58; arginine at position 71; and tryptophan at position 100; and comprise in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according to Kabat) and in their light chain variable region tryptophan at position 67 and glutamic acid at position 96 (position numbering according to Kabat).

[0062] Exemplary HAADs include the antibodies disclosed herein as well as those disclosed in Scheid et al., *Science*, 333:1633-1637 (2011); and West et al., *Proc. Natl. Acad. Sci. USA*, E2083-E2090 (2012). Studies have shown that Antibody A and Antibody B are of the same B cell lineage from one patient and differ at four amino acid positions in their light chain variable regions and at ten amino acid positions in their heavy chain variable regions (Scheid et al., 2011). The exemplary antibodies include but are not limited to Antibody A, Antibody B, and an antibody comprising the heavy chain of Antibody A and the light chain of Antibody B.

[0063] Table I provides the complementarity determining regions (CDRs) of the heavy chain variable region and the light chain variable region of Antibody A and Antibody B according to the Kabat, Chothia, and IMGT definitions.

[0064] Table I. CDRs of Antibody A and Antibody B

AntibodyA CDR	Kabat	Chothia	IMGT
VH-CDR1	DYFIH (SEQ ID NO: 137)	GYNIRDY (SEQ ID NO: 143)	GYNIRDYF (SEQ ID NO: 149)
VH-CDR2	WINPKTGQPN NPRQFQG (SEQ ID NO: 138)	PKTG (SEQ ID NO: 144)	INPKTGQP (SEQ ID NO: 150)
VH-CDR3	QRSDYWDFDV (SEQ ID NO: 139)	RSDYWDFD (SEQ ID NO: 145)	ARQRSDYWDFDV (SEQ ID NO: 151)
VL-CDR1	QANGYLN (SEQ ID NO: 140)	NGY (SEQ ID NO: 146)	GY (SEQ ID NO: 152)
VL-CDR2	DGSKLER (SEQ ID NO: 141)	DGS (SEQ ID NO: 147)	DGS (SEQ ID NO: 147)
VL-CDR3	QVYEF (SEQ ID NO: 142)	YE (SEQ ID NO: 148)	QVYEF (SEQ ID NO: 142)

Antibody B CDR	Kabat	Chothia	IMGT
VH-CDR1	DHFIH (SEQ ID NO: 153)	GYKISDH (SEQ ID NO: 155)	GYKISDHF (SEQ ID NO: 157)
VH-CDR2	WINPKTGQPN NPRQFQG	PKTG	INPKTGQP

	(SEQ ID NO: 138)	(SEQ ID NO: 144)	(SEQ ID NO: 150)
VH-CDR3	QRSDFWDFDV (SEQ ID NO: 154)	RSDFWDFD (SEQ ID NO: 156)	ARQRSDFWDFDV (SEQ ID NO: 158)
VL-CDR1	QANGYLN (SEQ ID NO: 140)	NGY (SEQ ID NO: 146)	GY (SEQ ID NO: 152)
VL-CDR2	DGSKLER (SEQ ID NO: 141)	DGS (SEQ ID NO: 147)	DGS (SEQ ID NO: 147)
VL-CDR3	QVYEF (SEQ ID NO: 142)	YE (SEQ ID NO: 148)	QVYEF (SEQ ID NO: 142)

[0065] The complementarity determining regions (CDRs) of exemplary antibodies of the present application are provided below: the CDRs according to the Kabat definition (Tables II and V), Chothia definition (Tables III and VI), and IMGT definition (Tables IV and VII). Antibodies comprising the CDRs listed below are encompassed by the present application.

[0066] In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments thereof of this disclosure in addition to including the six CDRs of Antibody A or Antibody B according to the Kabat, Chothia, or IMGT definitions provided below also include tryptophan (W) or phenylalanine (F) at Kabat position 74a, aspartic acid (D) at Kabat position 74b, phenylalanine (F) at Kabat position 74c, and aspartic acid (D) at Kabat position 74d; *i.e.*, the WDFD (SEQ ID NO: 453) or the FDFD (SEQ ID NO: 627) sequence in framework region 3 of their VH or heavy chain domain. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments thereof of this disclosure in addition to including the six CDRs of Antibody A, also include phenylalanine (F) at Kabat position 74a, aspartic acid (D) at Kabat position 74b, phenylalanine (F) at Kabat position 74c, and aspartic acid (D) at Kabat position 74d; *i.e.*, the FDFD (SEQ ID NO: 627) sequence in framework region 3 of their VH or heavy chain domain. Crystallographic studies have shown that framework region 3 at VH Kabat position numbers 74a, 74b, 74c and 74d form part of the paratope of the herein described antibody variants, directly contacting the antigen target, gp120. *See, e.g.*, Lee, *et al.*, *Immunity* (2017) 46(4): 690–702 (Figure 1G, identifying residue W71d); Klein, *et al.*, *Cell*. (2013) 153(1):126-38 (Figures 4 and 5); and Zhou, *et al.*, (2013) *Immunity* (2013) 39 245-258 (Table 1); ribbon diagrams of crystallized structures of 5V8L, 5V8M, 4JPV and 4LSV can be viewed at rcsb.org.

Table II. CDR Definitions (Kabat) of Antibodies						
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3	
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFMH (SEQ ID NO: 159)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFIH (SEQ ID NO: 137)	WINPKWGQPNPRQFQG (SEQ ID NO: 160)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFIH (SEQ ID NO: 137)	WINPKGQPNPRQFQG (SEQ ID NO: 161)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFIH (SEQ ID NO: 137)	WINPKAGQPNPRQFQG (SEQ ID NO: 162)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFIH (SEQ ID NO: 137)	WINPKHGQPNPRQFQG (SEQ ID NO: 163)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 164)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	
DYFMH (SEQ ID NO: 159)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 164)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)	

SUBSTITUTE SHEET (RULE 26)

Table II. CDR Definitions (Kabat) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DASKLER (SEQ ID NO: 165)	QVYEF (SEQ ID NO: 142)
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSNLER (SEQ ID NO: 166)	QVYEF (SEQ ID NO: 142)
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLET (SEQ ID NO: 167)	QVYEF (SEQ ID NO: 142)
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DASNLER (SEQ ID NO: 168)	QVYEF (SEQ ID NO: 142)
DHFIH (SEQ ID NO: 153)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 139)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 154)	QANGYLN (SEQ ID NO: 140)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)
DYFIH (SEQ ID NO: 137)	WINPKTGQPNPRQFQG (SEQ ID NO: 138)	QRSDYWDFDV (SEQ ID NO: 154)	QATGYLN (SEQ ID NO: 570)	DGSKLER (SEQ ID NO: 141)	QVYEF (SEQ ID NO: 142)

Table III. CDR Definitions (Chothia) of Antibodies						
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3	
GYNIRDY (SEQ ID NO: 143)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKWG (SEQ ID NO: 169)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKGG (SEQ ID NO: 170)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKAG (SEQ ID NO: 171)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKHG (SEQ ID NO: 172)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKTG (SEQ ID NO: 144)	RTDYWDFD (SEQ ID NO: 173)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 143)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DAS (SEQ ID NO: 174)	YE (SEQ ID NO: 148)	
GYNIRDY (SEQ ID NO: 459)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)	
GYNISDY (SEQ ID NO:)	PKTG (SEQ ID NO:)	RSDYWDFD (SEQ ID NO:)	NGY (SEQ ID NO:)	DGS (SEQ ID NO:)	YE (SEQ ID NO:)	

Table III. CDR Definitions (Chothia) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
460)	NO: 144)	145)	NO: 146)	NO: 147)	NO: 148)
GYNIRDH (SEQ ID NO: 461)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)
GYNIRDY (SEQ ID NO: 143)	PKTG (SEQ ID NO: 144)	RSDFWDFD (SEQ ID NO: 156)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)

Table IV. CDR Definitions (IMGT) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
GYNIRDYF (SEQ ID NO: 149)	INPKTGQP (SEQ ID NO: 150)	ARQRSYWDYDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GYNIRDYF (SEQ ID NO: 149)	INPKWGQP (SEQ ID NO: 175)	ARQRSYWDYDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GYNIRDYF (SEQ ID NO: 149)	INPKGGQP (SEQ ID NO: 176)	ARQRSYWDYDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GYNIRDYF (SEQ ID NO: 149)	INPKAGQP (SEQ ID NO: 177)	ARQRSYWDYDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GYNIRDYF (SEQ ID NO: 149)	INPKHGQP (SEQ ID NO: 178)	ARQRSYWDYDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GYNIRDYF	INPKTGQP	ARQRTDYWDYDFDV	GY	DGS	QVYEF

Table IV. CDR Definitions (IMGT) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
(SEQ ID NO: 149)	(SEQ ID NO: 150)	(SEQ ID NO: 179)	(SEQ ID NO: 152)	(SEQ ID NO: 147)	(SEQ ID NO: 142)
GYNIRDYF	INPKTGQP	ARQRSYWDVDV	GY	DAS	QVYEF
(SEQ ID NO: 149)	(SEQ ID NO: 150)	(SEQ ID NO: 151)	(SEQ ID NO: 152)	(SEQ ID NO: 180)	(SEQ ID NO: 142)
GYKIRDYF	INPKTGQP	ARQRSYWDVDV	GY	DGS	QVYEF
(SEQ ID NO: 462)	(SEQ ID NO: 150)	(SEQ ID NO: 151)	(SEQ ID NO: 152)	(SEQ ID NO: 147)	(SEQ ID NO: 142)
GYNISDYF	INPKTGQP	ARQRSYWDVDV	GY	DGS	QVYEF
(SEQ ID NO: 463)	(SEQ ID NO: 150)	(SEQ ID NO: 151)	(SEQ ID NO: 152)	(SEQ ID NO: 147)	(SEQ ID NO: 142)
GYNIRDHF	INPKTGQP	ARQRSYWDVDV	GY	DGS	QVYEF
(SEQ ID NO: 464)	(SEQ ID NO: 150)	(SEQ ID NO: 151)	(SEQ ID NO: 152)	(SEQ ID NO: 147)	(SEQ ID NO: 142)
GYNIRDYF	INPKTGQP	ARQRSYWDVDV	GY	DGS	QVYEF
(SEQ ID NO: 149)	(SEQ ID NO: 150)	(SEQ ID NO: 158)	(SEQ ID NO: 152)	(SEQ ID NO: 147)	(SEQ ID NO: 142)

Table V. CDR Definitions (Kabat) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
DYFIH	WINPKTGQPNPRQFG	QRSYWDVDV	QANGYLN	DGSKLER	QVYEF
(SEQ ID NO: 137)	(SEQ ID NO: 138)	(SEQ ID NO: 139)	(SEQ ID NO: 140)	(SEQ ID NO: 141)	(SEQ ID NO: 142)
DHFIH	WINPKTGQPNPRQFG	QRSYWDVDV	QANGYLN	DGSKLER	QVYEF
(SEQ ID NO: 153)	(SEQ ID NO: 138)	(SEQ ID NO: 139)	(SEQ ID NO: 140)	(SEQ ID NO: 141)	(SEQ ID NO: 142)

Table VI. CDR Definitions (Chothia) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
GYNIRDY (SEQ ID NO: 143)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)
GKIRDH (SEQ ID NO: 457)	PKTG (SEQ ID NO: 144)	RSDYWDFD (SEQ ID NO: 145)	NGY (SEQ ID NO: 146)	DGS (SEQ ID NO: 147)	YE (SEQ ID NO: 148)

Table VII. CDR Definitions (IMGT) of Antibodies					
VH-CDR1	VH-CDR2	VH-CDR3	VL-CDR1	VL-CDR2	VL-CDR3
GYNIRDYF (SEQ ID NO: 149)	INPKTGQP (SEQ ID NO: 150)	ARQRSDYWDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)
GKIRDHF (SEQ ID NO: 458)	INPKTGQP (SEQ ID NO: 150)	ARQRSDYWDFDV (SEQ ID NO: 151)	GY (SEQ ID NO: 152)	DGS (SEQ ID NO: 147)	QVYEF (SEQ ID NO: 142)

[0067] Encompassed by the present application are anti-gp120 antibodies or gp120-binding fragments thereof that include the six CDRs of each of the antibodies disclosed herein (*see, e.g.*, Tables I- VII). In certain embodiments, one or more of these anti-gp120 antibodies or gp120-binding fragments thereof also include tryptophan (W) or phenylalanine (F) at Kabat position 74a, aspartic acid (D) at Kabat position 74b, phenylalanine (F) at Kabat position 74c, and aspartic acid (D) at Kabat position 74d. It is to be understood that this disclosure also encompasses anti-gp120 antibodies or gp120-binding fragments thereof comprising the CDRs according to any other CDR definition (*e.g.*, Honegger definition, enhanced Chothia definition, AbM definition, contact definition, *see, e.g.*, www.bioinf.org.uk/abs/#cdrdef) of the anti-HIV antibodies disclosed herein. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have improved killing ability of HIV-1 infected target CD4 T cells compared to Antibody A and/or Antibody B. In certain embodiments, antibodies comprising VH and VL comprising the amino acid sequences set forth in SEQ ID NOs.: 477 and 278, respectively, or HC and LC comprising the amino acid sequences set forth in SEQ ID NOs.: 529 and 103, respectively, have improved killing ability of HIV-1 infected target CD4 T cells compared to Antibody A and/or Antibody B. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 2 µg/mL in ADCC assays of NK cell mediated killing of HIV-infected cells (*e.g.*, HIV-1-infected cells). In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 1.5 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 1.0 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 0.85 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 0.75 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 0.5 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.05 to 0.3 µg/mL. In certain instances, the anti-gp120 antibodies or gp120-binding fragments disclosed herein have an EC₅₀ of 0.07 to 0.2 µg/mL.

[0068] The amino acid sequences of the heavy chain variable region (VH) of and light chain variable region (VL) of exemplary antibodies of the presentation application are provided in Tables VIII and IX, respectively. The amino acid sequences of the VH and

VL of controls used in some assays of this disclosure (*e.g.*, Antibody C and Antibody D) are also included.

Table VIII. VH Sequences		
SEQ ID NO	Name	Heavy Chain Variable Region (VH) Amino Acid Sequence
181	C	QVRLSQSGGQMKKPGDSMRISCRASGYEFINCPINWIRLAPGKRPEW MGWMKPRWGAVSYARQLQGRVTMTRDMYSETAFLELRSLTSDDTAVY FCTRGKYCTARDYYNWDFEHWGQGTPTVTVSS
182	A-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
183	D-1	QVRLSQSGGQMKKPGDSMRISCRASGYEFINCPINWIRLAPGKRPEW MGWMKPRHGAVSYARQLQGRVTMTRDMYSETAFLELRSLTSDDTAVY FCTRGKYCTARDYYNWDFEHWGQGTPTVTVSS
184	1v2-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
185	1.2.1-1	QVQLLQSGAEVKKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTMVTVSS
186	1.3.1-1	QVSLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
187	1.4.1-1	QVQLVQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
188	1.5.1-1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
189	1.6.1-1	QVQLLQSGAEVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
190	1.7.1-1	QVQLLQSGAEVKKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
191	1.8.1-1	QVQLVQSGAEVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS
192	1.9.1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSYWDVDFVWGSGTQVTVSS

Table VIII. VH Sequences		
SEQ ID NO	Name	Heavy Chain Variable Region (VH) Amino Acid Sequence
193	1.10.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFMHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
194	1.11.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFMHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSAYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
195	1.15.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKWGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
196	1.16.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKGGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
197	1.17.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKAGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
198	1.18.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKHGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
199	1.19.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVTLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
200	1.20.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVTMTTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
201	1.21.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRDASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
202	1.22.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASTFSFYMDLKALRSDDTAVY FCARQQRSDYWDFDVWGSQTQVTVSS
203	1.24.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFSMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
204	1.25.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLSRLRSDD TAVYFCARQQRSDYWDFDVWGSQTQVTVSS
205	1.26.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TATYFCARQQRSDYWDFDVWGSQTQVTVSS
206	1.27.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRTDYWDFDVWGSQTQVTVSS
207	1.28.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TATYFCARQRTDYWDFDVWGSQTQVTVSS

Table VIII. VH Sequences		
SEQ ID NO	Name	Heavy Chain Variable Region (VH) Amino Acid Sequence
208	1.29.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSGTQVTVSS
209	1.30.1-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSGTMVTVSS
210	1.12.15-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSGTQVTVSS
211	1.13.15-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFMHWVRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSGTQVTVSS
212	1.14.15-1	QVQLLQSGAAVTKPGASVRVSCEASGYNIRDYFMHWVRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSAYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGSGTQVTVSS
213	1.31.1-1	QVQLVQSGAEVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TATYFCARQQRSDYWDFDVWGSGTQVTVSS
214	1.32.1-1	QVQLVQSGAEVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRTDYWDFDVWGSGTQVTVSS
215	1.33.1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVTLTRHASWDFDTFSFYMDLKALRSDD TATYFCARQQRSDYWDFDVWGSGTQVTVSS
216	1.34.1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVTLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRTDYWDFDVWGSGTQVTVSS
217	1.35.1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFMHWVRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVTMTRHASWDFDTFSFYMDLSRLRSDD TATYFCARQRTDYWDFDVWGQGTMTVTVSS
218	1.36.1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFMHWVRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVTMTRHASWDFDTFSAYMDLSRLRSDD TATYFCARQRTDYWDFDVWGQGTMTVTVSS
219	1.37.51-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTYSFYMDLSRLRSDD TAVYFCARQQRSDYWDFDVWGQGTMTVTVSS
220	B-1	QVHLSQSGAAVTKPGASVRVSCEASGYKISDHFHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRQASWDFDTYSFYMDLKAVRSDD TAIYFCARQQRSDYWFDFDVWGSGTQVTVSS
221	1.41.5-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQQRSDYWDFDVWGQGTMTVTVSS
465	2.2.1-1	QVHLSQSGAAVTKPGASVRVSCEASGYKIRDHFHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKAVRSDD TAIYFCARQQRSDYWDFDVWGSGTQVTVSS
466	2.3.1-1	QVHLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKAVRSDD TAIYFCARQQRSDYWDFDVWGSGTQVTVSS

Table VIII. VH Sequences		
SEQ ID NO	Name	Heavy Chain Variable Region (VH) Amino Acid Sequence
467	1.42.1-1	QVHLLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
468	1.43.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
469	1.44.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYKIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
470	1.45.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNISDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
471	1.46.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDHFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
472	1.47.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRQASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
474	1.49.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKAVRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
475	1.50.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAIYFCARQRSDYWDFDVWGSGTQVTVSS
476	1.51.1-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYDFDVWGSGTQVTVSS
477	1.52.64-1	QVQLSQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKALRSDD TAVYFCARQRSDYWDFDVWGSGTQVTVSS
478	2.4.1-1	QVHLSQSGAAVTKPGASVRVSCEASGYKIRDHFIHWWRQAPGQGLQW VGWINPKTGQPN NPRQFQGRVSLTRHASWDFDTFSFYMDLKAVRSDD TAIYFCARQRSDYWDFDVWGSGTQVTVSS

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
222	C	EIVLTQSPGTLSSLSPGETAIISCRTSQYGSLAWYQQRPGQAP RLVIYSGSTRAAGIPDRFSGSRWGPDYNLTI SNLESGDFGVY YCQQYEFFGQGTKVQVDIK
223	A-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
224	D-1	SLTQSPGTLSSLSPGETAIISCRTSQYGSLAWYQQRPGQAPRL VIYSGSTRAAGIPDRFSGSRWGPDYNLTI SNLESGDFGVYYC QQYEFFGQGTKVQVDIK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
225	1.1.2-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTKVDIK
226	1.1.3-1	EIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
227	1.1.4-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
228	1.1.5-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
229	1.1.6-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDASKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
230	1.1.7-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSNLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
231	1.1.8-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDASNLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
232	1.1.9-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLETGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
233	1.1.10-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYTLTINNLPEDIATYFC QVYEFVVPGTRLDLK
234	1.1.11-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYTLTINNLPEDIATYFC QVYEFVVPGTRLDLK
235	1.1.12-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTISLQPEDATYFC QVYEFVVPGTRLDLK
236	1.1.13-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYTLTISLQPEDATYFC QVYEFVVPGTRLDLK
237	1.1.14-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGSGTDFTFTINNLPEDIATYFC QVYEFVVPGTRLDLK
238	1.1.15-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFFVVPGTRLDLK
239	1.1.16-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVGPTRLDLK
240	1.1.17-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFFGPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
241	1.1.18-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVSPGTRLDLK
242	1.1.19-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVTPGTRLDLK
243	1.1.20-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTNLDLK
244	1.1.21-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVPGTNLDLK
245	1.1.22-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVSPGTNLDLK
246	1.1.23-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVTPGTNLDLK
247	1.1.24-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRVDLK
248	1.1.25-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTNVDLK
249	1.1.26-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDIK
250	1.1.27-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRVDIK
251	1.1.28-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTNVDIK
252	1.1.29-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVPGTNVDIK
253	1.1.30-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVTPGTNVDIK
254	1.1.31-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYTLTISSLQPEDIATYFC QVYEFVVPGTNLDLK
255	1.1.32-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYTLTISSLQPEDIATYFC QVYEFVTPGTRLDLK
256	1.1.33-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSNLERGVPSRFSGRRWGQEYTLTISSLQPEDIATYFC QVYEFVVPGTNLDIK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
257	1.1.34-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSNLERGVPSRFSGRRWGQEYTLTISSLQPEDATYFC QVYEFVTPGTRLDIK
258	1.1.35-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQKPGKAPKL LIYDGSNLETGVPSRFSGSRWGQEYTLTISSLQPEDATYFC QVYEFVGPNTLDIK
259	1.1.36-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQKPGKAPKL LIYDGSNLETGVPSRFSGSRWGQEYTLTISSLQPEDATYFC QVYEFVTPGNTLDIK
260	1.1.37-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYTFITNNLQPEDATYFC QVYEFVVPGTRLDLK
261	1.1.38-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYSLTINNLPEDATYFC QVYEFVVPGTRLDLK
262	1.1.39-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYSFTINNLPEDATYFC QVYEFVVPGTRLDLK
263	1.1.40-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYALTINNLPEDATYFC QVYEFVVPGTRLDLK
264	1.1.41-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYAFITINNLPEDATYFC QVYEFVVPGTRLDLK
265	1.1.42-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHITINNLPEDATYFC QVYEFVVPGTRLDLKR
266	1.1.43-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHFTINNLPEDATYFC QVYEFVVPGTRLDLK
267	1.1.44-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYQLTINNLPEDATYFC QVYEFVVPGTRLDLK
268	1.1.45-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYQFTINNLPEDATYFC QVYEFVVPGTRLDLK
269	1.1.46-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLKINNLPEDATYFC QVYEFVVPGTRLDLK
270	1.1.47-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNFKINNLPEDATYFC QVYEFVVPGTRLDLK
271	1.1.48-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLAINNLPEDATYFC QVYEFVVPGTRLDLK
272	1.1.49-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNFAINNLPEDATYFC QVYEFVVPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
273	1.37.51-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLETGVPSRFSGSRWGQEYTLTINNLPEDIATYFC QVYEFFGPGTRLDLK
274	1.8.52-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYTLTINNLPEDIATYFC QVYEFVVPGTRLDLK
275	1.1.54-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYTLTINNLPEDIATYFC QVYEFVVPGTRLDLK
276	B-1	DIQMTQSPSSLSARVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRWRWGQEYNLTINNLPEDVATYFC QVYEFIVPGTRLDLK
277	2.1.2-1	DIQMTQSPSSLSARVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRWRWGQEYHLTINNLPEDVATYFC QVYEFIVPGTRLDLK
278	1.1.64-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
279	1.1.67-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYNLTINNLPEDIATYFC QVYEFFGPGTRLDLK
280	1.1.72-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYHLTINNLPEDIATYFC QVYEFFGPGTRLDLK
281	1.1.75-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYHLTINNLPEDIATYFC QVYEFVVPGTRLDLK
282	1.1.78-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYHLTINNLPEDIATYFC QVYEFFGPGTRLDLK
283	1.41.81-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYNLTINNLPEDIATYFC QVYEFFGPGTRLDLK
284	1.1.82-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDVATYFC QVYEFVVPGTRLDLK
285	1.41.83-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDVATYFC QVYEFVVPGTRLDLK
286	1.1.84-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDVATYFC QVYEFFGPGTRLDLK
287	1.41.85-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDVATYFC QVYEFFGPGTRLDLK
288	1.41.86-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRWRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
289	1.41.87-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFFGPGTRLDLK
290	1.1.88-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDATYFC QVYEFVVPGTRLDLK
291	1.41.89-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDATYFC QVYEFVVPGTRLDLK
292	1.1.90-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDATYFC QVYEFFGPGTRLDLK
293	1.41.91-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYNLTISLQPEDATYFC QVYEFFGPGTRLDLK
294	1.41.92-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHLTINNLPEDIATYFC QVYEFVVPGTRLDLK
295	1.41.93-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHLTINNLPEDIATYFC QVYEFFGPGTRLDLK
296	1.1.94-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFVVPGTRLDLK
297	1.41.95-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFVVPGTRLDLK
298	1.1.96-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFFGPGTRLDLK
299	1.41.97-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFFGPGTRLDLK
300	1.41.98-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHLTINNLPEDIATYFC QVYEFVVPGTRLDLK
301	1.41.99-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYHLTINNLPEDIATYFC QVYEFFGPGTRLDLK
302	1.1.100-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFVVPGTRLDLK
303	1.41.101-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFVVPGTRLDLK
304	1.1.102-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISLQPEDATYFC QVYEFFGPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
305	1.41.103-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYHLTISSLQPEDVATYFC QVYEFFGPGTRLDLK
306	1.1.110-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGTRRGQDYIFSINNLPEDIATYFC QVYEFVVPGTRLDLK
307	1.1.111-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRFGQDYILTIINNLPEDIATYFC QVYEFVVPGTRLDLK
308	1.1.112-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRFGQKYQLSINNLPEDIATYFC QVYEFVVPGTRLDLK
309	1.1.113-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRFGQDYILTIINNLPEDIATYFC QVYEFVVPGTRLDLK
310	2.1.3-1	DIQMTQSPSSLSARVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRFGQDYILTIINNLPEDVATYFC QVYEFIVPGTRLDLK
311	2.1.4-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRFGQDYILTIINNLPEDVATYFC QVYEFIVPGTRLDLK
479	3.1.8-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYNLTINNLPEDVATYFC QVYEFIVPGTRLDLK
480	3.1.9-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYNLTINNLPEDVATYFC QVYEFFGPGTRLDLK
481	1.1.115-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYILTIINNLPEDIATYFC QVYEFVVPGTRLDLK
482	3.1.10-1	DIQMTQSPSSLSARVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYILTIINNLPEDVATYFC QVYEFIVPGTRLDLK
483	1.1.116-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYILTIINNLPEDIATYFC QVYEFVVPGTRLDLK
484	3.1.11-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYILTIINNLPEDVATYFC QVYEFIVPGTRLDLK
485	1.1.117-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYILTIINNLPEDIATYFC QVYEFFGPGTRLDLK
486	3.1.12-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYILTIINNLPEDVATYFC QVYEFFGPGTRLDLK
487	1.1.118-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYILTISSLQPEDVATYFC QVYEFFGPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
488	3.1.13-1	DIQMTQSPSSLSARVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGSRWGQEYILTISLQPEDVATYFC QVYEFFGPGTRLDLK
489	3.1.14-1	DIQMTQSPSSLSARVGDTVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYTLTINNLPEDVATYFC QVYEFIVPGTRLDLK
491	3.1.5-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYTLTINNLPEDVATYFC QVYEFIVPGTRLDLK
492	3.1.15-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYTLTINNLPEDVATYFC QVYEFFGPGTRLDLK
493	1.1.119-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGSRWGQEYTLTISLQPEDVATYFC QVYEFFGPGTRLDLK
494	3.1.7-1	DIQMTQSPSSLSARVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGSRWGQEYTLTISLQPEDVATYFC QVYEFFGPGTRLDLK
495	3.1.16-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYHLTINNLPEDVATYFC QVYEFIVPGTRLDLK
496	3.1.17-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQEYHLTINNLPEDVATYFC QVYEFFGPGTRLDLK
497	3.1.18-1	DIQMTQSPSSLSARVGDRATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGSRWGQEYHLTISLQPEDVATYFC QVYEFFGPGTRLDLK
498	1.1.120-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQDYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
499	3.1.19-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRWGQDYILTINNLPEDVATYFC QVYEFIVPGTRLDLK
500	1.1.121-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRFGQEYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
501	3.1.20-1	DIQMTQSPSSLSARVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPARFSGRRFGQEYILTINNLPEDVATYFC QVYEFIVPGTRLDLK
502	1.1.122-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYVLTINNLPEDIATYFC QVYEFVVPGTRLDLK
503	1.1.123-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYLLTINNLPEDIATYFC QVYEFVVPGTRLDLK
504	1.1.124-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYMLTINNLPEDIATYFC QVYEFVVPGTRLDLK

Table IX. VL Sequences		
SEQ ID NO	Name	Light Chain Variable Region (VL) Amino Acid Sequence
505	1.1.125-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYALTINNLPEDIATYFC QVYEFVVPGTRLDLK
506	1.1.126-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYSLTINNLPEDIATYFC QVYEFVVPGTRLDLK
507	1.1.127-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYFLTINNLPEDIATYFC QVYEFVVPGTRLDLK
508	1.1.128-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGTRWGQEYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
509	1.1.129-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
510	1.1.130-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
511	1.1.131-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGTRWGQDYILTINNLPEDIATYFC QVYEFVVPGTRLDLK
512	1.1.132-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
513	1.1.133-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
514	1.1.134-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDVATYFC QVYEFVVPGTRLDLK
515	1.1.135-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFIVPGTRLDLK
569	1.1.138-1	DIQMTQSPSSLSASVGDVTITCQATGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYNLTINNLPEDIATYFC QVYEFVVPGTRLDLK
516	1.1.104-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKL LIYDGSKLERGVPSRFSGRRWGQEYTLTINNLPEDIATYFC QVYEFFGPGTRLDLK

[0069] In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and a VL that is at least 80%, at least

85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569. In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a VH selected from the group consisting of SEQ ID NOs: 181-221 and 465-478, and a VL selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569.

[0070] When comparing polynucleotide and polypeptide sequences, two sequences are said to be “identical” if the sequence of nucleotides or amino acids in the two sequences is the same when aligned for maximum correspondence, as described below. Comparisons between two sequences are typically performed by comparing the sequences over a comparison window to identify and compare local regions of sequence similarity. A “comparison window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned.

[0071] Alignment of sequences for comparison may be conducted using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. This program embodies several alignment schemes described in the following references: Dayhoff, M.O. (1978) A model of evolutionary change in proteins - Matrices for detecting distant relationships. In Dayhoff, M.O. (ed.) Atlas of Protein Sequence and Structure, National Biomedical Research Foundation, Washington DC Vol. 5, Suppl. 3, pp. 345-358; Hein J. (1990) Unified Approach to Alignment and Phylogenies pp. 626-645 Methods in Enzymology vol. 183, Academic Press, Inc., San Diego, CA; Higgins, D.G. and Sharp, P.M. (1989) CABIOS 5: 151-153; Myers, E.W. and Muller W. (1988) CABIOS 4:11-17; Robinson, E.D. (1971) Comb. Theor 77: 105; Santou, N. Nes, M. (1987) Mol. Biol. Evol. 4:406-425; Sneath, P.H.A. and Sokal, R.R. (1973) Numerical Taxonomy - the Principles and Practice of Numerical Taxonomy, Freeman Press, San Francisco, CA; Wilbur, W.J. and Lipman, D.J. (1983) Proc. Natl. Acad., Sci. USA 80:726-730.

[0072] Alternatively, alignment of sequences for comparison may be conducted by the local identity algorithm of Smith and Waterman (1981) Add. APL. Math 2:482, by the identity alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443, by the search for similarity methods of Pearson and Lipman (1988) Proc. Natl. Acad. Sci.

USA 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), 575 Science Dr., Madison, WI), or by inspection.

[0073] One example of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) Nucl. Acids Res. 25:3389-3402 and Altschul et al. (1990) J. Mol. Biol. 215:403-410, respectively. BLAST and BLAST 2.0 can be used, for example with the parameters described herein, to determine percent sequence identity for the polynucleotides and polypeptides described herein. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information.

[0074] In one illustrative example, cumulative scores can be calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a word length (W) of 11, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) Proc. Natl. Acad. Sci. USA 89: 10915) alignments, (B) of 50, expectation (E) of 10, M=5, N=-4 and a comparison of both strands.

[0075] For amino acid sequences, a scoring matrix can be used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T and X determine the sensitivity and speed of the alignment.

[0076] In one approach, the “percentage of sequence identity” is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or

less, usually 5 to 15 percent, or 10 to 12 percent, as compared to the reference sequences (which does not comprise additions or deletions) for alignment of the two sequences.

The percentage is calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residues occur in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (i.e., the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

[0077] Encompassed by this disclosure are anti-gp120 antibodies or gp120-binding fragments thereof that include the VH of any of antibodies disclosed herein. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the VH of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.90-1, Antibody 2.2.1-1, Antibody 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, Antibody 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1, Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody 2-1. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the VH of Antibody 1.52.64-1.

[0078] Encompassed by this disclosure are anti-gp120 antibodies or gp120-binding fragments thereof that include the VL of any of the antibodies disclosed above. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the VL of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.1.90-1, Antibody 2.2.1-1, 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1, Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody B-1-1. Also encompassed are anti-gp120 antibodies or gp120-binding fragments thereof that include the VH and VL of any of the antibodies disclosed herein. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the VH and VL of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.1.90-1, Antibody 2.2.1-1, Antibody 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, Antibody 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1, Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody B-1. Also encompassed by this disclosure are the antibodies comprising the CDRs of any of the foregoing VL and/or VH sequences.

[0079] In certain instances, the anti-gp120 antibodies or gp120-binding fragments thereof comprises in addition to the VH amino acid sequence of any of the antibodies disclosed herein, a heavy chain constant region comprising an amino acid sequence below with 0 to 10 (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 437);

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 438);

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPDVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 439);

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 440);

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPDVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 441);

or

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT
 VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTLRVVSILTTLVHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPLVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTKQKSLSLSPGK (SEQ ID NO: 442).

[0080] In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments thereof comprises the VH amino acid sequence set forth in SEQ ID NO: 477 and a heavy chain constant region comprising an amino acid sequence set forth in SEQ ID NO: 438 with 0 to 10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions.

[0081] The amino acid sequences of the heavy chain and light chain of exemplary antibodies of the present application are shown in Tables X and XI, respectively. The amino acid sequence of the heavy and light chain of control antibodies used in a number of the assays of this disclosure (*e.g.*, Antibody C and Antibody D-1) are also included.

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
1	C	QVRLSQSGGQMKKPGDSMRISCRASGYEFINCPINWIRLAPGKRPEWMGWMKPRWGAVSYA RQLQGRVTMTRDMYSETAFLELRSLTSDDTAVYFCTRGRKYCTARDYYNWD FEHWGQGT PVT VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLG GPSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQ GNVFSCSVLHEALHNHYTQKSLSLSPGK
2	A-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
3	A	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGGP SVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHNHYTQKSLSLSPGK
4	D-1	QVRLSQSGGQMKKPGDSMRISCRASGYEFINCPINWIRLAPGKRPEWMGWMKPRHGAVSYA RQLQGRVTMTRDMYSETAFLELRSLTSDDTAVYFCTRGRKYCTARDYYNWD FEHWGQGT PVT VSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQ SSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLA GPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYN STYRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEM TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQ GNVFSCSVLHEALHSHYTQKSLSLSPGK
5	1v2-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
6	1.2.1 -1	QVQLLQSGAEVKKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTMVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
7	1.3.1 -1	QVSLQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
8	1.4.1 -1	QVQLVQSGAAVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
9	1.5.1 -1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
10	1.6.1 -1	QVQLLQSGAEVTKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
11	1.7.1 -1	QVQLLQSGAEVKKPGASVRVSCEASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
12	1.8.1 -1	QVQLVQSGAEVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
13	1.9.1 -1	QVQLVQSGAEVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
14	1.10. 1-1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFMHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
15	1.11. 1-1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFMHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSAYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
16	1.15. 1-1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKWGPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
17	1.16. 1-1	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKGGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
18	1.17. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKAGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
19	1.18. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKHGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
20	1.19. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
21	1.20. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTMTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
22	1.21. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRDASWDFDTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
23	1.22. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASTFSFYMDLKALRSDDTAVYFCARQRSYDWFDFVWGSGTQVTVSSAST KGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS LSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFL FPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVV SVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTKNQVS LTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSC SVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
24	1.24. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
25	1.25. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKLRLSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
26	1.26. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTATYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
27	1.27. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRTDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
28	1.28. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTATYFCARQRTDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
29	1.29. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGQGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
30	1.30.1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTMVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
31	1.12.15-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
32	1.13.15-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFMHWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
33	1.14.15-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFMHWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSAYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
34	1.31.1-1	QVQLVQSGAEVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
35	1.32.1-1	QVQLVQSGAEVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQRTDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
36	1.33. 1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTLTRHASWDFDFTSFYMDLKALRSDDTATYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
37	1.34. 1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTLTRHASWDFDFTSFYMDLKALRSDDTAVYFCARQRTDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
38	1.35. 1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFMHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTMTRHASWDFDFTSFYMDLSRLRSDDTATYFCARQRTDYWDFDVWGQGTMTVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
39	1.36. 1-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFMHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVTMTRHASWDFDFTSFAYMDLSRLRSDDTATYFCARQRTDYWDFDVWGQGTMTVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
40	1.37. 51-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTSFYMDLSRLRSDDTAVYFCARQQRSDYWDFDVWGQGTMTVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
41	A-2	QVQLVQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
42	B-1	QVHLSQSGAAVTKPGASVRVSCASGYKISDHFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRQASWDFDFTSFYMDLKAVRSDDTAIYFCARQQRSDYWDFDVWGSQTQVTVS

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
		SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
43	A-3	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
44	A-4	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
45	A-5	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
46	A-6	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTLRVVSILTTLVHQDWLNGKEYKCKVSNKALPAPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPLVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
47	1.41.5-1	QVQLVQSGAEVKKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
517	2.2.1-1	QVHLSQSGAAVTKPGASVRVSCASGYKIRDHFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDFTFSFYMDLKA VRSDDTAIYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
		DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
518	2.3.1 -1	QVHLSQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA VRSDDTAIYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
519	1.42. 1-1	QVHLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
520	1.43. 1-1	QVQLLSQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
521	1.44. 1-1	QVQLLSQSGAAVTKPGASVRVSCASGYKIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
522	1.45. 1-1	QVQLLSQSGAAVTKPGASVRVSCASGYNISDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
523	1.46. 1-1	QVQLLSQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKA LRSDDTAVYFCARQ RSDYWDFDVWGSGTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK

Table X. Heavy Chain Sequences		
SEQ ID NO	Name	Heavy Chain Amino Acid Sequence
		NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYTQKSLSLSPGK
524	1.47. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRQASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
526	1.49. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
527	1.50. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
528	1.51. 1-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
529	1.52. 64-1	QVQLLQSGAAVTKPGASVRVSCASGYNIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK
530	2.4.1 -1	QVHLSQSGAAVTKPGASVRVSCASGYKIRDYFIHWWRQAPGQGLQWVGWINPKTGQPNNP RQFQGRVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCARQQRSDYWDFDVWGSQTQVTVS SASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST YRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTK NQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGN VFSCSVLHEALHSHYTQKSLSLSPGK

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
48	C	EIVLTQSPGTLSSLSPGETAII SCRTSQYGS LAWYQQRPGQAPRLVIYSGSTRAAGIPDRFS GSRWGPDYNTIISNLESGDFGVYYCQQYEFFGQGTKVQVDIKRTVAAPSVFIFPPSDEQLK SGTASVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYE KHKVYACEVTHQGLSSPVTKSFNRGEC
49	A-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
50	D-1	SLTQSPGTLSSLSPGETAII SCRTSQYGS LAWYQQRPGQAPRLVIYSGSTRAAGIPDRFSGS RWGPDYNTIISNLESGDFGVYYCQQYEFFGQGTKVQVDIKRTVAAPSVFIFPPSDEQLKSG TASVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHK KVYACEVTHQGLSSPVTKSFNRGEC
51	1.1.3-1	EIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
52	1.1.4-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
53	1.1.5-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
54	1.1.6-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDASKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
55	1.1.7-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSNLERGVP SRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
56	1.1.8-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDASNLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
57	1.1.9-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLETGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
58	1.1.10-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
59	1.1.11-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYTLTINNLPEDIATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
60	1.1.12-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTISLQPEDATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA SVVCLLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
61	1.1.13-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQQRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYTLTISLQPEDATYFCQVYEFVVPGTRLDLKRVAAPSVFIFPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
62	1.1.14-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS GSGTDFTFITINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
63	1.1.15-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
64	1.1.16-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
65	1.1.17-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
66	1.1.18-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
67	1.1.19-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
68	1.1.20-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTNLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
69	1.1.21-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTNLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
70	1.1.22-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTNLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
71	1.1.23-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTNLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
72	1.1.24-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRVDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
73	1.1.25-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTNVDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
74	1.1.26-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
75	1.1.27-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVPGTRVDIKRTVAAPSVFI FPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
76	1.1.28-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTNVDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
77	1.1.29-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTNVDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
78	1.1.30-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTNVDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
79	1.1.31-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
80	1.1.32-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
81	1.1.33-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSNLERGVPSRFSGR RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
82	1.1.34-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSNLERGVPSRFSGR RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
83	1.1.35-1	DIQMTQSPSSLSASVGDVITITCQANGYLNWYQKPGKAPKLLIYDGSNLETGVPSRFSGS RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
84	1.1.36-1	DIQMTQSPSSLSASVGDVITITCQANGYLNWYQKPGKAPKLLIYDGSNLETGVPSRFSGS RWGQEYTLTISLQPEDIAITFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
85	1.1.37-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTFITINNLPEDIATYFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
86	1.1.38-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYSLTINNLPEDIATYFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
87	1.1.39-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYSFTINNLPEDIATYFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
88	1.1.40-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYALTINNLPEDIATYFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
89	1.1.41-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYAFITINNLPEDIATYFCQVYEFVVPGTNLDIKRTVAAPSVFI FPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
90	1.1.42-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHFTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
91	1.1.43-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHFTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
92	1.1.44-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYQFTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
93	1.1.45-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYQFTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
94	1.1.46-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLKINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
95	1.1.47-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNFKINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
96	1.1.48-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLAINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
97	1.1.49-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNFAINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
98	1.37.51-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLETGVPSRFSGS RWGQEYTLTINNLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
99	1.8.52-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
100	1.1.54-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYTLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
101	B-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYNLTINNLPEDVATYFCQVYEFIVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
102	2.1.2-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYHFTINNLPEDVATYFCQVYEFIVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
103	1.1.64-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
104	1.1.67-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
105	1.1.72-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
106	1.1.75-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
107	1.1.78-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
108	1.41.81-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
109	1.1.82-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
110	1.41.83-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
111	1.1.84-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
112	1.41.85-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
113	1.41.86-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
114	1.41.87-1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
115	1.1.88-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
116	1.41.89-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFVPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
117	1.1.90-1	DIQMTQSPSSLSASVGDRTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYNLTISSLQPEDATYFCQVYEFFGPGTRLDLKRVAAPSVFI FPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
118	1.41.91-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
119	1.41.92-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHNTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
120	1.41.93-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHNTINNLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
121	1.1.94-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
122	1.41.95-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
123	1.1.96-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
124	1.41.97-1	DIQMTQSPSSLSASVGDRVTITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
125	1.41.98-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHNTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
126	1.41.99-1	DIQMTQSPSSLSASVGDTATITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYHNTINNLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
127	1.1.100-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
128	1.41.101-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
129	1.1.102-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
130	1.41.103-1	DIQMTQSPSSLSASVGDRATITCQANGYLNWYQQKPGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYHNTISSLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
131	1.1.110-1	DIQMTQSPSSLSASVGDTVTITCQANGYLNWYQQRRGKAPKLLIYDGSKLERGVPSRFSGT RRGQDYIFSIINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
132	1.1.111-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RFGQDYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
133	1.1.112-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RFGQKYQLSINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
134	1.1.113-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RFGQDYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
135	2.1.3-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RFGQDYILTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
136	2.1.4-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RFGQDYILTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
531	3.1.8-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYNLTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
532	3.1.9-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYNLTINNLPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
533	1.1.115-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
534	3.1.10-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYILTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
535	1.1.116-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
536	3.1.11-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYILTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
537	1.1.117-1	DIQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYILTINNLPEDIATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
538	3.1.12-1	DIQMTQSPSSLSARVGDVTITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYILTINNLPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYSLSSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
539	1.1.118-1	DIQMTQSPSSLSASVGDRTATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYILTISLQPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
540	3.1.13-1	DIQMTQSPSSLSARVGDRAITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGS RWGQEYILTISLQPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
541	3.1.14-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYTLTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
542	3.1.5-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYTLTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
543	3.1.15-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYTLTINNLPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
544	1.1.119-1	DIQMTQSPSSLSASVGDRAITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGS RWGQEYTLTISLQPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
545	3.1.7-1	DIQMTQSPSSLSARVGDRAITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGS RWGQEYTLTISLQPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
546	3.1.16-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYHILTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
547	3.1.17-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYHILTINNLPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
548	3.1.18-1	DIQMTQSPSSLSARVGDRAITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGS RWGQEYHILTISLQPEDVATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
549	1.1.120-1	DIQMTQSPSSLSASVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQDYILTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
550	3.1.19-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQDYILTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
551	1.1.121-1	DIQMTQSPSSLSASVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RFGQEYILTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
552	3.1.20-1	DIQMTQSPSSLSARVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RFGQEYILTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
553	1.1.122-1	DIQMTQSPSSLSASVGDVITITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYVLTINNLPEDVATYFCQVYEFVPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
554	1.1.123 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYLLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
555	1.1.124 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYMLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
556	1.1.125 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYALTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
557	1.1.126 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYSLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
558	1.1.127 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYFLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
559	1.1.128 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGT RWGQEYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
560	1.1.129 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RRGQEYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
561	1.1.130 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RYGQEYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
562	1.1.131 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGT RWGQDYILTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
563	1.1.132 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
564	1.1.133 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPARFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
565	1.1.134 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDVATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
566	1.1.135 -1	DIQMTQSPSSLSASVGDATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA SVVCLNNFYFPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
568	1.1.138 -1	DIQMTQSPSSLSASVGDATITCQATGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEYNLTINNLPEDIATYFCQVYEFVVPGTRLDLKRTVAAPSVFI FPPSDEQLKSGTA

Table XI. Light Chain Sequences		
SEQ ID NO	Name	Light Chain Amino Acid Sequence
		SVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC
567	1.1.104-1	DIQMTQSPSSLSASVGDITATITCQANGYLNWYQRRGKAPKLLIYDGSKLERGVPSRFSGR RWGQEQYTLTINNLPEDATYFCQVYEFFGPGTRLDLKRTVAAPSVFIFPPSDEQLKSGTA SVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLTLSKADYEKHKV YACEVTHQGLSSPVTKSFNRGEC

[0082] In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a heavy chain (HC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and a light chain (LC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 48-136 and 531-567. In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a HC selected from the group consisting of SEQ ID NOs: 1-47 and 517-530, and a LC selected from the group consisting of SEQ ID NOs: 48-136 and 531-567. In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a heavy chain (HC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to the amino acid sequence set forth in SEQ ID NO: 529 and a light chain (LC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to the amino acid sequence set forth in SEQ ID NO: 103. In some embodiments, the anti-gp120 antibodies or gp120-binding fragments described herein have a HC with the amino acid sequence set forth in SEQ ID NO: 529, and a LC with the amino acid sequence set forth in SEQ ID NO: 103.

[0083] Encompassed by this disclosure are anti-gp120 antibodies or gp120-binding fragments thereof that include the heavy chain of any of the antibodies disclosed herein. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the heavy chain of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.1.90-1, Antibody 2.2.1-1, Antibody 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, Antibody 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1,

Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody B-1. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the heavy chain of Antibody 1.52.64-1.

[0084] Encompassed by this disclosure are anti-gp120 antibodies or gp120-binding fragments thereof that include the light chain of any of the antibodies disclosed herein. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the light chain of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.1.90-1, Antibody 2.2.1-1, Antibody 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, Antibody 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1, Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody B-1. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the light chain of Antibody 1.52.64-1.

[0085] Also encompassed are anti-gp120 antibodies or gp120-binding fragments thereof that include the heavy and light chain of any of the antibodies disclosed herein. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the heavy and light chains of any one of Antibody A-1, Antibody 1.1.64-1, Antibody 1.1.90-1, Antibody 2.2.1-1, Antibody 2.3.1-1, Antibody 3.1.5-1, Antibody 2.2.5-1, Antibody 2.3.5-1, Antibody 1.1.119-1, Antibody 1.1.104-1, Antibody 1.52.64-1, Antibody 2.4.1-1, Antibody 1.1.54-1, or Antibody B-1. In certain embodiments, the anti-gp120 antibody or gp120-binding fragment thereof includes the heavy and light chains of Antibody 1.52.64-1.

[0086] Encompassed by this disclosure are anti-gp120 antibodies or gp120-binding fragments thereof that include any of the VH and/or VL amino acid substitutions shown above.

[0087] In some embodiments, the variable heavy chain of any of the anti-gp120 antibodies of this disclosure is linked to a heavy chain constant region comprising a CH1 domain and a hinge region. In some embodiments, the variable heavy chain of any of the anti-gp120 antibodies of this disclosure is linked to a heavy chain constant region comprising a CH3 domain. In certain embodiments, the variable heavy chain of any of the anti-gp120 antibodies of this disclosure is linked to a heavy chain constant region comprising a CH1 domain, hinge region, and CH2 domain from IgG4 and a CH3 domain (*e.g.*, from IgG1, IgG2, IgG3, or IgG4). In some instances, the variable heavy chain of any of the anti-gp120 antibodies of this disclosure is linked to a heavy chain constant region comprising a CH1 domain, hinge region, CH2 domain, and a CH3 domain from

IgG1, IgG2, IgG3, or IgG4. In certain embodiments, the variable heavy chain of any of the anti-gp120 antibodies of this disclosure is linked to a heavy chain constant region comprising a CH1 domain, CH2 domain, and a CH3 domain from IgG1 (*e.g.*, human IgG1, *e.g.*, IgG1m3 allotype) and an IgG3 hinge region (*e.g.*, an “open” IgG3 hinge region designated “IgG3 C-” in WO 2017/096221 (*see, e.g.*, Fig. 2A of this PCT publication)). This IgG3 hinge region is expected to exhibit improved Fab arm flexibility and the ability to span over a 200 Å distance that is sufficient for intra-trimeric interactions. In certain embodiments, such a chimeric antibody contains one or more additional mutations in the heavy chain constant region that increase the stability of the chimeric antibody. In certain embodiments, the heavy chain constant region includes substitutions that modify the properties of the antibody (*e.g.*, increase effector function, improve pharmacokinetics, increase or decrease Fc receptor binding, increase or decrease antibody glycosylation, increase or decrease binding to C1q, increase half-life).

[0088] In certain embodiments, the anti-gp120 antibody is an IgG antibody (*e.g.*, IgG1, IgG2, IgG3, IgG4). In one embodiment, the antibody is human IgG1. In another embodiment, the antibody is human IgG2. In some embodiments, the antibody has a chimeric heavy chain constant region (*e.g.*, having the CH1, hinge, and CH2 regions of human IgG4 and CH3 region of human IgG1). In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and the antibody is human IgG1. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and the antibody is human IgG1.

[0089] IgG antibodies exist in various allotypes and isoallotypes. In certain embodiments, antibodies of the present disclosure include an IgG1 heavy chain having an allotype of G1m1; nG1m2; G1m3; G1m17,1; G1m17,1,2; G1m3,1; or G1m17. Each of these allotypes or isoallotypes is characterized by the following amino acid residues at the indicated positions within the IgG1 heavy chain constant region (Fc) (EU

numbering): G1m1: D356, L358; nG1m1: E356, M358; G1m3: R214, E356, M358, A431; G1m17,1: K214, D356, L358, A431; G1m17,1,2: K214, D356, L358, G431; G1m3,1: R214, D356, L358, A431; and G1m17: K214, E356, M358, A431. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and the antibody has an IgG1 heavy chain having an allotype of G1m1; nG1m2; G1m3; G1m17,1; G1m17,1,2; G1m3,1; or G1m17. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and the antibody has an IgG1 heavy chain having an allotype of G1m1; nG1m2; G1m3; G1m17,1; G1m17,1,2; G1m3,1; or G1m17.

[0090] In one embodiment, any of the VHs of an anti-gp120 antibody disclosed herein is directly linked to, or linked via an intervening amino acid sequence (*e.g.*, a G-S linker), to a wild type IgG1m3 sequence provided below (representative allotype-determining residues are indicated in bold).

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
VVTVPSSSLGTQTYICNVNHKPSNTKVDK**K**VEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL
MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE
YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPS**REEM**TKNQVSLTCLVKGFYPSDIAVEWESNG
QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVME**A**LHNHYTQKSLSLSPGK (SEQ
ID NO: 347).

[0091] In another embodiment, any of the VHs of an anti-gp120 antibody disclosed herein is directly linked to, or linked via an intervening amino acid sequence (*e.g.*, a G-S linker), to a wild type IgG1m17 sequence provided below (representative allotype-determining residues are indicated in bold).

[0092] IgG1m17:

ASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSS
VVTVPSSSLGTQTYICNVNHKPSNTKVDK**K**VEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTL
MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKE

YKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSR**EE**MTKNQVSLTCLVKGFYPSDIAVEWESNG
 QPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQSVME**AL**HNHYTQKSLSLSPGK (SEQ
 ID NO: 348).

[0093] In certain embodiments, a VH of an anti-gp120 antibody disclosed herein is directly linked to, or linked via an intervening amino acid sequence (*e.g.*, a G-S linker), to a IgG1m17 sequence with 1 to 10 (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions in SEQ ID NO:348 (*e.g.*, substitutions made to improve effector function and/or to increase half-life). Exemplary amino acid substitutions in the Fc region (of *e.g.*, IgG1 such as IgG1m17) include S239D, I332E, G236A, A330L, M428L, N434S; S239D, I332E, G236A, A330L; S239D, I332E M428L, N434S; S239D, I332E, A330L, M428L, N434S; F243L, R292P, Y300L, V305I, P396L, M428L, N434S; and S239D, I332E, G236A, A330L.

[0094] In certain embodiments, the anti-gp120 antibody is a human IgG1/human kappa antibody. In some embodiments, antibodies of this disclosure comprise a kappa light chain having an allotype selected from Km1; Km1,2; or Km3. Each of these allotypes is characterized by the following amino acid residues at the indicated positions within the light chain (EU numbering): Km1: V153, L191; Km1,2: A153, L191; and Km3: A153, V191. . In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and comprises a kappa light chain having an allotype selected from Km1; Km1,2; or Km3. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and comprises a kappa light chain having an allotype Km3. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and is a human IgG1/human kappa antibody, such as an human IgG1/Km3. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at

least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and is a human IgG1/human kappa antibody, such as an human IgG1/Km3.

[0095] In certain embodiments, an anti-gp120 antibody of this disclosure comprises a human kappa light chain comprising one of the following amino acid sequences, in which representative allotype-determining residues are indicated in bold:

Km1:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN**VL**QSGNSQESVTEQDSKDYSTYS
LSSTLTLSKADYEEKHK**LY**ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 349);

Km1,2:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN**AL**QSGNSQESVTEQDSKDYSTYS
LSSTLTLSKADYEEKHK**LY**ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 350); or

Km3:

RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN**AL**QSGNSQESVTEQDSKDYSTYS
LSSTLTLSKADYEEKHK**Y**ACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 351).

[0096] In one embodiment, an anti-gp120 antibody of this disclosure comprises a human kappa light chain, Km3. In a specific embodiment, a VL of an anti-gp120 antibody disclosed herein is directly linked to, or linked via an intervening amino acid sequence (*e.g.*, a G-S linker), to a wild type human Km3 sequence (SEQ ID NO:351). In certain embodiments, the VL of an anti-gp120 antibody disclosed herein is directly linked to, or linked via an intervening amino acid sequence (*e.g.*, a G-S linker), to a mutant human Km3 sequence having 1 to 5 (*i.e.*, 1, 2, 3, 4, 5) amino acid substitutions within SEQ ID NO:351.

[0097] In certain embodiments, the anti-gp120 antibody is a human IgG1/human lambda antibody. Each individual human includes between seven and eleven different lambda light chain genes, which encode light chains selected from Lambda1, Lambda2, Lambda3, Lambda4, Lambda5, Lambda6, and Lambda7. In certain embodiments, antibodies of the present disclosure comprise a lambda light chain selected from Lambda1, Lambda2, Lambda3, Lambda4, Lambda5, Lambda6, and Lambda7. In some embodiments, an antibody described herein comprises a lambda light chain comprising one of the following amino acid sequences, in which representative lambda-determining residues are indicated in bold:

Lambda1:

GQPKA**NP**TVTLFPPSSEELQANKATLVCL**I**SDFYPGAVTVAWKAD**GSPVK**AGVETTKPSKQSNNKYA
ASSYLSLTPEQWKSH**RSYSCQ**VTHEGSTVEKTVAP**TECS** (SEQ ID NO: 352);

Lambda2:

GQPKA**AP**SVTLFPPSSEELQANKATLVCL**I**SDFYPGAVTVAWKAD**SSPVK**AGVETTTPSKQSNNKYA
ASSYLSLTPEQWKSH**RSYSCQ**VTHEGSTVEKTVAP**TECS** (SEQ ID NO: 353);

Lambda3:

GQPKA**AP**SVTLFPPSSEELQANKATLVCL**I**SDFYPGAVTVAWKAD**SSPAK**AGVETTTPSKQSNNKYA
ASSYLSLTPEQWKSH**KSYSCQ**VTHEGSTVEKTVAP**TECS** (SEQ ID NO: 354);

orLambda7:

GQPKA**AP**SVTLFPPSSEELQANKATLVCL**V**SDFYPGAVTVAWKAD**GSPVKV**GVETTKPSKQSNNKYA
ASSYLSLTPEQWKSH**RSYSCR**VTHEGSTVEKTVAP**AECS** (SEQ ID NO: 355).

[0098] In one embodiment, the anti-gp120 antibody is a human IgG1m17/human Km3 antibody. The constant regions (light and/or heavy) can include 1 to 10 (*e.g.*, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions (*e.g.*, substitutions made to improve effector function and/or to increase half-life). In some embodiments, the antibodies are afucosylated. In some embodiments, the antibodies comprise one or more tags. In certain embodiments, the one or more tags comprise an avidin tag. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and is a human IgG1m17/human Km3 antibody. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and is a human IgG1m17/human Km3 antibody, wherein the heavy chain constant region includes 1 to 10 amino acid substitutions. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively and is a human IgG1m17/human Km3 antibody, wherein the heavy chain constant region includes the following amino acid substitutions compared to SEQ ID

NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and is a human IgG1/human kappa antibody, such as an human IgG1/Km3, wherein the heavy chain constant region includes the following amino acid substitutions compared to SEQ ID NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, these substitutions improve effector function. In certain embodiments, these substitutions increase half-life. In certain embodiments, these substitutions improve effector function and improve half-life.

[0099] In certain embodiments, the antibody that binds to gp120 comprises an amino acid sequence of a VH of an anti-gp120 antibody disclosed herein and of a VL of an anti-gp120 antibody disclosed herein. Exemplary VH and VL amino acid sequences of an anti-gp120 antibody include the sequences set forth in SEQ ID NOs: 182 and 223, respectively; SEQ ID NOs: 182 and 275, respectively; SEQ ID NOs: 182 and 278, respectively; SEQ ID NOs: 182 and 292, respectively; SEQ ID NOs: 220 and 276, respectively; SEQ ID NOs: 465 and 276, respectively; SEQ ID NOs: 466 and 276, respectively; SEQ ID NOs: 182 and 491, respectively; SEQ ID NOs: 465 and 491, respectively; SEQ ID NOs: 466 and 491, respectively; SEQ ID NOs: 182 and 493, respectively; SEQ ID NOs: 182 and 516, respectively; SEQ ID NOs: 182 and 276, respectively; SEQ ID NOs: 182 and 569, respectively; SEQ ID NOs: 477 and 223, respectively; SEQ ID NOs: 477 and 278, respectively; SEQ ID NOs: 477 and 292, respectively; and SEQ ID NOs: 478 and 276, respectively. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs: 477 and 278, respectively. In certain embodiments, each of these antibodies are human IgG1m17/human Km3 antibodies. In certain embodiments, these antibodies comprise the amino acid sequence set forth in SEQ ID NO: 348 and/or 351. In some instances, these antibodies include up to 10 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10) amino acid substitutions (e.g., substitutions made to improve effector function and/or to increase half-life) within SEQ ID NO: 348 and/or 351, respectively. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3,

wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351 with 1 to 10 amino acid sequence substitutions within SEQ ID NO: 348 and/or 351. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351 with 1 to 10 amino acid sequence substitutions within SEQ ID NO: 348. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351, with the following amino acid substitutions in SEQ ID NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and is a IgGm17/human Km3 antibody. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and the antibody comprises a human kappa light chain comprising the amino acid sequence set forth in SEQ ID NO: 351 and a IgG1 heavy chain having an allotype with the amino acid sequence set forth in SEQ ID NO: 348. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351 with 1 to 10 amino acid sequence substitutions within SEQ ID NO: 348 and/or 351. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351 with 1 to 10 amino acid sequence substitutions within SEQ ID NO: 348. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and comprises the amino acid sequence set forth in SEQ ID NOs: 348 and 351, with the following amino acid substitutions in SEQ ID NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid

sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and is a IgGm17/human Km3 antibody. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and the antibody comprises a human kappa light chain comprising the amino acid sequence set forth in SEQ ID NO: 351 and a IgG1 heavy chain having an allotype with the amino acid sequence set forth in SEQ ID NO: 348.

[0100] In certain embodiments, the antibody that binds to gp120 comprises an amino acid sequence of a heavy chain of an anti-gp120 antibody disclosed herein and a light chain of an anti-gp120 antibody disclosed herein. Exemplary heavy chain and light chain sequences of an anti-gp120 antibody include the sequences set forth in SEQ ID NOs: 2 and 49, respectively; SEQ ID NOs: 2 and 100, respectively; SEQ ID NOs: 42 and 101, respectively; SEQ ID NOs: 2 and 103, respectively; SEQ ID NOs: 517 and 101, respectively; SEQ ID NOs: 518 and 101, respectively; SEQ ID NOs: 2 and 542, respectively; SEQ ID NOs: 517 and 542, respectively; SEQ ID NOs: 2 and 117, respectively; SEQ ID NOs: 518 and 542, respectively; SEQ ID NOs: 2 and 544, respectively; SEQ ID NOs: 2 and 567, respectively; SEQ ID NOs: 2 and 568, respectively; SEQ ID NOs: 529 and 49, respectively; SEQ ID NOs: 529 and 103, respectively; SEQ ID NOs: 529 and 117, respectively; and SEQ ID NOs: 530 and 101, respectively. In certain embodiments, the antibody that binds to gp120 comprises a heavy chain with the amino acid sequence set forth in SEQ ID NO: 529 and a light chain with the amino acid sequence set forth in SEQ ID NO: 103.

[0101] Antibodies or antigen-binding fragments described herein can be made, for example, by preparing and expressing nucleic acids that encode the amino acid sequences of the antibody.

Multispecific Antibodies

[0102] In another aspect, this disclosure provides multispecific antibodies. Multispecific antibodies are antibodies which binds two or more different epitopes (*e.g.*, bispecific antibodies, trivalent antibodies, tetravalent antibodies). The anti-gp120 antibodies described above can be comprised as part of multispecific antibodies. The multispecific antibodies may have binding sites to at least one other antigen or one other epitope that is not bound by the anti-gp120 antibody binding site of the multispecific antibody. The anti-gp120 comprising multispecific antibody can include a dimerization domain and three or more (*e.g.*, three, four, five, six) antigen binding sites. An exemplary dimerization domain comprises (or consists of) an Fc region. An anti-gp120 comprising

multispecific antibody can comprise (or consist of) three to about eight (i.e., three, four, five, six, seven, eight) antigen binding sites. The multispecific antibody optionally comprises at least one polypeptide chain (*e.g.*, two polypeptide chains, three polypeptide chains), wherein the polypeptide chain(s) comprise three or more variable domains. For instance, the polypeptide chain(s) may comprise, *e.g.*, VD1-(X1)_n-VD2-(X2)_n-Fc, or VD1-(X1)_n-VD2-(X2)_n-VD3-(X3)_n-Fc, wherein VD1 is a first variable domain, VD2 is a second variable domain, VD3 is a third variable domain Fc is a polypeptide chain of an Fc region, X1, X2, and X3 represent an amino acid or peptide spacer, and n is 0 or 1. In certain instances, the variable domains may each be an scFv. Multispecific antibodies can be readily produced by recombinant expression of nucleic acid encoding the polypeptide chains of the antibody.

Bispecific Antibodies

[0103] In one aspect, the multispecific antibody is a bispecific antibody. Bispecific antibodies are antibodies that have binding specificities for two different epitopes. A bispecific antibody has two “arms.” One arm of the bispecific antibody binds one epitope and the other arm another epitope. In one embodiment, one arm of the bispecific antibody binds a first antigen and the other arm of the bispecific antibody binds a second antigen. In another embodiment, the two arms of the bispecific antibody bind to two different epitopes of the same antigen (*e.g.*, gp120).

[0104] In one aspect, this disclosure provides a bispecific antibody that specifically binds to gp120 and specifically binds to a second antigen. In certain embodiments, the second antigen is a triggering molecule on a leukocyte so as to focus and localize cellular defense mechanisms to the infected cell. In some cases, the second antigen is a T-cell receptor molecule (*e.g.*, CD3, CD4); Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16); CD89; an HIV-1 antigen (*e.g.*, gp41); CCR5; a KIR family member, such as killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3); an NKG2 family receptor such as, killer cell lectin like receptor C1 (KLRC1), killer cell lectin like receptor C2 (KLRC2), killer cell lectin like receptor C3 (KLRC3), killer cell lectin like receptor C4 (KLRC4), killer cell lectin like receptor D1

(KLRD1), killer cell lectin like receptor K1 (KLRK1); a natural cytotoxicity triggering receptor, such as natural cytotoxicity triggering receptor 3 (NCR3 or NKp30), natural cytotoxicity triggering receptor 2 (NCR2 or NK-p44), natural cytotoxicity triggering receptor 1 (NCR1 or NK-p46), CD226 (DNAM-1), cytotoxic and regulatory T cell molecule (CRTAM or CD355); a SLAM family member, such as signaling lymphocytic activation molecule family member 1 (SLAMF1), CD48 (SLAMF2), lymphocyte antigen 9 (LY9 or SLAMF3), CD244 (2B4 or SLAMF4), CD84 (SLAMF5), SLAM family member 6 (SLAMF6 or NTB-A), SLAM family member 7 (SLAMF7 or CRACC); CD27 (TNFRSF7), semaphorin 4D (SEMA4D or CD100), or CD160 (NK1). In certain embodiments, the second arm of the bispecific antibody binds a different epitope of gp120.

[0105] In a further embodiment, a bispecific antibody molecule of this disclosure includes a dual-variable-domain antibody (DVD-Ig), where each light chain and heavy chain contains two variable domains in tandem through a short peptide linkage (Wu et al., Generation and Characterization of a Dual Variable Domain Immunoglobulin (DVD-IgTM) Molecule, In: Antibody Engineering, Springer Berlin Heidelberg (2010)). In some embodiments, the bispecific antibody is a chemically-linked bispecific (Fab')₂ fragment. In other embodiments, the bispecific antibody comprises a Tandab (i.e., a fusion of two single chain diabodies resulting in a tetravalent bispecific antibody that has two binding sites for each of the target antigens). In certain embodiments, the bispecific antibody is a flexibody, which is a combination of scFvs with a diabody resulting in a multivalent molecule. In yet another embodiment, the bispecific antibody comprises a “dock and lock” molecule, based on the “dimerization and docking domain” in Protein Kinase A, which, when applied to Fabs, can yield a trivalent bispecific binding protein consisting of two identical Fab fragments linked to a different Fab fragment. In another instance, the bispecific antibodies of this disclosure comprise a “Scorpion molecule,” comprising, *e.g.*, two scFvs fused to both termini of a human Fab-arm. In yet another embodiment, the bispecific antibody of this disclosure comprises a diabody.

[0106] Exemplary classes of bispecific antibodies include but are not limited to IgG-like molecules with complementary CH3 domains to force heterodimerization; IgG fusion molecules, wherein full length IgG antibodies are fused to extra Fab fragment or parts of Fab fragment; Fc fusion molecules, wherein single chain Fv molecules or stabilized diabodies are fused to heavy-chain constant-domains, Fc-regions or parts thereof; Fab fusion molecules, wherein different Fab-fragments are fused together; recombinant IgG-

like dual targeting molecules, wherein the two sides of the molecule each contain the Fab fragment or part of the Fab fragment of at least two different antibodies; scFv- and diabody-based and heavy chain antibodies (*e.g.*, domain antibodies, nanobodies) wherein different single chain Fv molecules or different diabodies or different heavy-chain antibodies (*e.g.* domain antibodies, nanobodies) are fused to each other or to another protein or carrier molecule.

[0107] Examples of Fab fusion bispecific antibodies include but are not limited to F(ab)₂ (Medarex/AMGEN), Dual-Action or Bis-Fab (Genentech), Dock-and-Lock (DNL) (ImmunoMedics), Bivalent Bispecific (Biotecnol) and Fab-Fv (UCB-Celltech).

[0108] Examples of scFv-, diabody-based and domain antibodies include but are not limited to Bispecific T Cell Engager (BITE) (Micromet, Tandem Diabody (Tandab) (Affimed), Dual Affinity Retargeting Technology (DART) (MacroGenics), Single-chain Diabody (Academic), TCR-like Antibodies (AIT, ReceptorLogics), Human Serum Albumin ScFv Fusion (Merrimack) and COMBODY (Epigen Biotech), dual targeting nanobodies (Ablynx), and dual targeting heavy chain only domain antibodies.

Antigen-Binding Fragments

[0109] This disclosure encompasses antigen-binding fragments of the anti-gp120 antibodies disclosed herein. Antigen-binding antibody fragments (*e.g.*, scFv, sc(Fv)₂, Fab, F(ab)₂, Fab', F(ab')₂, Facb, and Fv) may be prepared, *e.g.*, by recombinant methods or by proteolytic digestion of intact antibodies. For example, antibody fragments can be obtained by treating the whole antibody with an enzyme such as papain, pepsin, or plasmin. Papain digestion of whole antibodies produces F(ab)₂ or Fab fragments; pepsin digestion of whole antibodies yields F(ab')₂ or Fab'; and plasmin digestion of whole antibodies yields Facb fragments.

[0110] Alternatively, antibody fragments can be produced recombinantly. For example, nucleic acids encoding the antibody fragments of interest can be constructed, introduced into an expression vector, and expressed in suitable host cells. See, *e.g.*, Co, M.S. et al., *J. Immunol.*, 152:2968-2976 (1994); Better, M. and Horwitz, A.H., *Methods in Enzymology*, 178:476-496 (1989); Plueckthun, A. and Skerra, A., *Methods in Enzymology*, 178:476-496 (1989); Lamoyi, E., *Methods in Enzymology*, 121:652-663 (1989); Rousseaux, J. et al., *Methods in Enzymology*, (1989) 121:663-669 (1989); and Bird, R.E. et al., *TIBTECH*, 9:132-137 (1991)). Antibody fragments can be expressed in and secreted from *E. coli*, thus allowing the facile production of large amounts of these

fragments. Antibody fragments can be isolated from the antibody phage libraries. Alternatively, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab)₂ fragments (Carter et al., *Bio/Technology*, 10:163-167 (1992)). According to another approach, F(ab')₂ fragments can be isolated directly from recombinant host cell culture. Fab and F(ab')₂ fragment with increased *in vivo* half-life comprising a salvage receptor binding epitope residues are described in U.S. Pat. No. 5,869,046.

Minibodies

[0111] Also encompassed by this disclosure are minibodies that bind gp120. Minibodies include diabodies, single chain (scFv), and single-chain (Fv)₂ (sc(Fv)₂).

[0112] A “diabody” is a bivalent minibody constructed by gene fusion (*see, e.g.*, Holliger, P. et al., *Proc. Natl. Acad. Sci. U. S. A.*, 90:6444-6448 (1993); EP 404,097; WO 93/11161). Diabodies are dimers composed of two polypeptide chains. The VL and VH domain of each polypeptide chain of the diabody are bound by linkers. The number of amino acid residues that constitute a linker can be between 2 to 12 residues (*e.g.*, 3-10 residues or five or about five residues). The linkers of the polypeptides in a diabody are typically too short to allow the VL and VH to bind to each other. Thus, the VL and VH encoded in the same polypeptide chain cannot form a single-chain variable region fragment, but instead form a dimer with a different single-chain variable region fragment. As a result, a diabody has two antigen-binding sites.

[0113] An scFv is a single-chain polypeptide antibody obtained by linking the VH and VL with a linker (*see e.g.*, Huston et al., *Proc. Natl. Acad. Sci. U. S. A.*, 85:5879-5883 (1988); and Plickthun, “The Pharmacology of Monoclonal Antibodies” Vol.113, Ed Resenburger and Moore, Springer Verlag, New York, pp.269-315, (1994)). The order of VHs and VLs to be linked is not particularly limited, and they may be arranged in any order. Examples of arrangements include: [VH] linker [VL]; or [VL] linker [VH]. The H chain V region and L chain V region in an scFv may be derived from any anti-gp120 antibody or antigen-binding fragment thereof described herein.

[0114] An sc(Fv)₂ is a minibody in which two VHs and two VLs are linked by a linker to form a single chain (Hudson, et al., *J. Immunol. Methods*, (1999), 231: 177-189). An sc(Fv)₂ can be prepared, for example, by connecting scFvs with a linker. The sc(Fv)₂ of the present disclosure include antibodies preferably in which two VHs and two VLs are arranged in the order of: VH, VL, VH, and VL ([VH] linker [VL] linker [VH] linker

[VL]), beginning from the N terminus of a single-chain polypeptide; however the order of the two VHs and two VLs is not limited to the above arrangement, and they may be arranged in any order. Examples of arrangements are listed below:

[VL] linker [VH] linker [VH] linker [VL]
 [VH] linker [VL] linker [VL] linker [VH]
 [VH] linker [VH] linker [VL] linker [VL]
 [VL] linker [VL] linker [VH] linker [VH]
 [VL] linker [VH] linker [VL] linker [VH]

[0115] Normally, three linkers are required when four antibody variable regions are linked; the linkers used may be identical or different. There is no particular limitation on the linkers that link the VH and VL regions of the minibodies. In some embodiments, the linker is a peptide linker. Any arbitrary single-chain peptide comprising about three to 25 residues (*e.g.*, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18) can be used as a linker. Examples of such peptide linkers include: Ser; Gly Ser; Gly Gly Ser; Ser Gly Gly; Gly Gly Gly Ser (SEQ ID NO: 427); Ser Gly Gly Gly (SEQ ID NO: 428); Gly Gly Gly Gly Ser (SEQ ID NO: 429); Ser Gly Gly Gly Gly (SEQ ID NO: 430); Gly Gly Gly Gly Gly Ser (SEQ ID NO: 431); Ser Gly Gly Gly Gly Gly (SEQ ID NO: 432); Gly Gly Gly Gly Gly Gly Ser (SEQ ID NO: 433); Ser Gly Gly Gly Gly Gly Gly (SEQ ID NO: 434); (Gly Gly Gly Gly Ser)_n (SEQ ID NO: 435), wherein n is an integer of one or more; and (Ser Gly Gly Gly Gly)_n (SEQ ID NO: 436), wherein n is an integer of one or more.

[0116] In certain embodiments, the linker is a synthetic compound linker (chemical cross-linking agent). Examples of cross-linking agents that are available on the market include N-hydroxysuccinimide (NHS), disuccinimidylsuberate (DSS), bis(sulfosuccinimidyl)suberate (BS3), dithiobis(succinimidylpropionate) (DSP), dithiobis(sulfosuccinimidylpropionate) (DTSSP), ethyleneglycol bis(succinimidylsuccinate) (EGS), ethyleneglycol bis(sulfosuccinimidylsuccinate) (sulfo-EGS), disuccinimidyl tartrate (DST), disulfosuccinimidyl tartrate (sulfo-DST), bis[2-(succinimidooxycarbonyloxy)ethyl]sulfone (BSOCOES), and bis[2-(sulfosuccinimidooxycarbonyloxy)ethyl]sulfone (sulfo-BSOCOES).

[0117] The amino acid sequence of the VH or VL in the minibodies may include modifications such as substitutions, deletions, additions, and/or insertions. For example, the modification may be in one or more of the CDRs of the anti-gp120 antibody or antigen-binding fragment thereof. In certain embodiments, the modification involves one, two, or three amino acid substitutions in one or more CDRs of the VH and/or VL

domain of the anti-gp120 minibody. Such substitutions are made to improve the binding and/or functional activity of the anti-gp120 minibody. In other embodiments, one, two, or three amino acids of the CDRs of the anti-gp120 antibody or antigen-binding fragment thereof may be deleted or added as long as there is gp120 binding and/or functional activity when VH and VL are associated.

[0118] In some embodiments, the antibodies and antigen-binding fragments thereof, described herein, do not comprise a signal peptide. In some embodiments, the antibodies and antigen-binding fragments thereof, described herein, comprise an N-terminal signal peptide. The signal peptide can be an endogenous signal peptide (*e.g.*, from a native or wild-type immunoglobulin protein), or from a heterologous polypeptide (*e.g.*, a non-immunoglobulin protein). In some embodiments, the heterologous signal peptide is from a secreted protein, *e.g.*, a serum protein, an immunoglobulin or a cytokine. In some embodiments, the signal peptide is from a serum albumin signal peptide (*e.g.*, having the amino acid sequence KWTFTISLLFLFSSAYS (SEQ ID NO: 620). In some embodiments, the signal peptide is comprises a sequence selected from the group consisting of MDPKGSLSWRILLFLSLAFELSYG (SEQ ID NO: 621), MSVPTQVLGLLLLWLT DARC (SEQ ID NO: 622), METDTLLLWVLLLWVPGSTG (SEQ ID NO: 623), MKWVTFISLLFLFSSAYS (SEQ ID NO: 624), MRCLAEFLGLLVLWIPGAIG (SEQ ID NO: 625), and MDPKGSLSWRILLFLSLAFELSYG (SEQ ID NO: 626). The signal peptide can be designed to be cleaved off, *e.g.*, after secretion from the cell, to form a mature fusion protein. A modified human serum albumin signal peptide to secrete proteins in cells that can find use in expressing the present fusion proteins is described, *e.g.*, in Attallah, et al., *Protein Expr Purif.* (2017) 132:27-33. Additional guidance for selection of signal peptide sequences for use in expressing the herein described antibodies and antigen-binding fragments thereof are described, *e.g.*, in Kober, et al., *Biotechnol Bioeng.* (2013) 110(4):1164-73; Gibson, et al., *Biotechnol Bioeng.* 2017 Sep;114(9):1970-1977; Lin, et al., *Biotechnol J.* 2017 Sep;12(9). doi: 10.1002/biot.201700268 (PMID 28727292); Ramezani, et al., *Protein Expr Purif.* 2017 Jul;135:24-32; and Haryadi, et al., *PLoS One.* 2015 Feb 23;10(2):e0116878. As appropriate, the heavy chain and the light chain, or antigen-binding fragments thereof, can have the same or different signal peptides when expressed as individual proteins.

Fc Modifications

[0119] In certain embodiments, the antibodies of this disclosure include one or more

amino acid sequence modifications in the heavy chain constant region (Fc) as compared to the IgG1m17 amino acid sequence (i.e., SEQ ID NO: 348). In certain embodiments, the antibodies of this disclosure include one or more amino acid sequence modifications in the heavy chain constant region (Fc) as compared to other anti-HIV-antibodies such as Antibody A or Antibody B. In some embodiments, these modifications increase stability or increase binding affinity of the modified antibody as compared to Antibody A or Antibody B. In certain embodiments, these modifications increase stability or increase effector function of the modified antibody as compared to Antibody A or Antibody B. In some embodiments, certain of these modifications, improve the pharmacokinetics of the antibody as compared to Antibody A or Antibody B. In certain embodiments, certain of these modifications, increase half-life of the antibody as compared to Antibody A or Antibody B. In other embodiments, certain of these modifications, increase antibody effector function and improve the pharmacokinetics of the antibody as compared to Antibody A or Antibody B. In other embodiments, certain of these modifications, increase antibody effector function and increase half-life of the antibody as compared to the Antibody A or Antibody B. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a heavy chain constant region with one or more amino acid sequence modifications as compared to SEQ ID NO: 348. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and comprises a heavy chain constant region with one or more amino acid sequence modifications as compared to SEQ ID NO: 348. In some embodiments, these substitutions improve effector function. In some embodiments, these substitutions increase half-life. In some embodiments, these substitutions improve effector function and increase half-life.

[0120] In certain embodiments, the one or more modifications are selected from the following Fc amino acid substitutions (EU numbering) or combinations thereof: L234F; L235E; G236A; S239D; F243L; D265E; D265A; S267E; H268F; R292P; N297Q; N297A; S298A; S324T; I332E; S239D; A330L; L234F; L235E; P331S; F243L; Y300L; V305I; P396L; S298A; E333A; K334A; E345R; L235V; F243L; R292P; Y300L; P396L; M428L; E430G; N434S; G236A, S267E, H268F, S324T, and I332E; G236A, S239D, and I332E; S239D, A330L, I332E; L234F, L235E, and P331S; F243L, R292P, Y300L, V305I, and P396L; G236A, H268F, S324T, and I332E; S239D, H268F, S324T, and I332E; S298A, E333A, and K334A; L235V, F243L, R292P, Y300L, and P396L; S239D, I332E; S239D, S298A, and I332E; G236A, S239D, I332E, M428L, and N434S; G236A, S239D, A330L, I332E, M428L, and N434S; S239D, I332E, G236A and A330L; M428L and N434S; M428L, N434S; G236A, S239D, A330L, and I332E; and G236A and I332E. In certain embodiments, one, two, three, four, or more amino acid substitutions are introduced into a Fc region to alter (*e.g.*, increase) the effector function of the antibody. For example, these substitutions are located at positions selected from the group consisting of amino acid residues 236, 239, 330 and 332 (according to EU numbering). These positions can be replaced with a different amino acid residue such that the antibody has an improved effector function. In certain embodiments, the antibody comprises a VH comprising VH CDRs 1-3 and a VL comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a heavy chain constant region with the following modifications (EU numbering) compared to SEQ ID NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, the anti-gp120 antibodies or gp120-binding fragments have a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278, and comprises a heavy chain constant region with the following modifications (EU numbering) compared to SEQ ID NO: 348: S239D, I332E, G236A, A330L, M428L, N434S. In certain embodiments, the antibody comprises a VH and VL comprising the amino acid sequences set forth in: SEQ ID NOs.: 477 and 278, respectively, and comprises a heavy chain constant region with the following modifications (EU numbering) compared to SEQ ID NO: 348: S239D, I332E, G236A,

A330L, M428L, N434S. In some embodiments, these substitutions improve effector function. In some embodiments, these substitutions increase half-life. In some embodiments, these substitutions improve effector function and increase half-life.

[0121] In certain instances, the antibodies of the present application comprise mutations that increase or enhance effector function by enhancing the binding of the Fc to activating FcγRs. In some instances, the antibodies of the present application comprise mutations that increase the pharmacokinetic half-life of the antibody.

[0122] Mutations that increase the half-life of an antibody are known in the art. In one embodiment, the constant region of an antibody described herein comprises a methionine to tyrosine substitution at position 252 (EU numbering), a serine to threonine substitution at position 254 (EU numbering), and a threonine to glutamic acid substitution at position 256 (EU numbering). *See, e.g.*, U.S. Patent No. 7,658,921. This type of mutant, designated as a “YTE mutant” exhibits a four-fold increased half-life relative to wild-type versions of the same antibody (Dall’Acqua et al., *J Biol Chem*, 281: 23514-24 (2006); Robbie et al., *Antimicrob Agents Chemother*, 57(12):6147-6153 (2013)). In certain embodiments, an antibody comprises an IgG constant domain comprising one, two, three or more amino acid substitutions of amino acid residues at positions 251-257, 285-290, 308-314, 385-389, and 428-436 (EU numbering). In other embodiments, an antibody described herein comprises T250Q and M428L (EU numbering) mutations. In other embodiments, an antibody described herein (*e.g.*, Duobodies®) comprises H433K and N434F (EU numbering) mutations.

Conjugated Antibodies

[0123] Any of the antibodies disclosed herein may be conjugated antibodies which are bound to various molecules including macromolecular substances such as polymers (*e.g.*, polyethylene glycol (PEG), polyethylenimine (PEI) modified with PEG (PEI-PEG), polyglutamic acid (PGA) (N-(2-Hydroxypropyl) methacrylamide (HPMA) copolymers), hyaluronic acid, radioactive materials (*e.g.*, ⁹⁰Y, ¹³¹I, ¹²⁵I, ³⁵S, ³H, ¹²¹In, ⁹⁹Tc), fluorescent substances (*e.g.*, fluorescein and rhodamine), luminescent substances (*e.g.*, luminol), Qdots, haptens, enzymes (*e.g.*, glucose oxidase), metal chelates, biotin, avidin, and drugs.

[0124] In some embodiments, the antibodies or antigen-binding fragments thereof described herein are conjugated to a cytotoxic agent, *e.g.*, for delivery to and killing of an HIV infected cell. In various embodiments, the cytotoxic agent is a

small organic compound or an inhibitory nucleic acid, *e.g.*, a short-inhibitory RNA (siRNA), a microRNA (miRNA). In some embodiments, the antibodies or antigen-binding fragments thereof described herein are conjugated to a cytotoxic agent selected from the group consisting of monomethyl auristatin E (MMAE), monomethyl auristatin F (MMAF), a calicheamicin, ansamitocin, maytansine or an analog thereof (*e.g.*, mertansine/emtansine (DM1), ravtansine/soravtansine (DM4)), an anthracycline (*e.g.*, doxorubicin, daunorubicin, epirubicin, idarubicin), pyrrolbenzodiazepine (PBD) DNA cross-linking agent SC-DR002 (D6.5), duocarmycin, a microtubule inhibitors (MTI) (*e.g.*, a taxane, a vinca alkaloid, an epothilone), a pyrrolbenzodiazepine (PBD) or dimer thereof, a duocarmycin (A, B1, B2, C1, C2, D, SA, CC-1065), and a *Pseudomonas* exotoxin.

[0125] The above-described conjugated antibodies can be prepared by performing chemical modifications on the antibodies or the lower molecular weight forms thereof described herein. Methods for modifying antibodies are well known in the art (*e.g.*, US 5,057,313 and US 5,156,840).

Nucleic Acids

[0126] This disclosure also provides a polynucleotide or polynucleotides encoding an antibody or antigen-binding fragment described herein, vectors comprising such polynucleotides, and host cells (*e.g.*, mammalian cells including hamster cells or human cells, plant cells, yeast cells, bacterial cells, including *E. coli* cells) comprising such polynucleotides or expression vectors. Provided herein are polynucleotides comprising nucleotide sequence(s) encoding any of the antibodies provided herein, as well as vector(s) comprising such polynucleotide sequences, *e.g.*, expression vectors for their efficient expression in host cells, *e.g.*, mammalian cells.

[0127] In another aspect, this disclosure provides polynucleotides or nucleic acid molecules encoding an antibody or antigen-binding fragment thereof according to the present invention. In some embodiments, the nucleic acid molecules encode an antibody light chain (or a fragment thereof) or an antibody heavy chain (or a fragment thereof), or both of the present application. In other embodiments, the nucleic acid is a DNA, a cDNA, or an mRNA. In some other embodiments, the nucleic acid molecule is codon-optimized to enhance expression in a host cell.

[0128] In one aspect, this disclosure provides polynucleotides comprising nucleotide sequences encoding the VH, VL, or VH and VL of the antibodies or antigen-binding

fragments which bind to gp120. In certain instances, the VH and VL have the amino acids set forth respectively in SEQ ID NOs.: 182 and 275; 182 and 278; 182 and 279; 182 and 280; 182 and 281; 182 and 282; 182 and 292; 182 and 304; 182 and 307; 182 and 309; 220 and 310; or 220 and 311.

[0129] In another aspect, provided herein are polynucleotides comprising a nucleotide sequence encoding the CDRs, light chain, or heavy chain of an antibody described herein. The polynucleotides can comprise nucleotide sequences encoding a light chain or light chain variable domain comprising the VL CDRs of antibodies described herein (see, *e.g.*, Tables above). The polynucleotides can comprise nucleotide sequences encoding a heavy chain or heavy chain variable domain comprising the VH CDRs of antibodies described herein (see, *e.g.*, Tables above). In one embodiment, a polynucleotide described herein encodes a variable light chain or light chain with the VL-CDRs comprising the amino acid sequence set forth in SEQ ID NOs: 140, 141, and 142, respectively. In another embodiment, a polynucleotide described herein encodes a variable heavy chain or heavy chain with VH CDRs comprising the amino acid sequence set forth in SEQ ID NOs: 137, 138, and 139, respectively. In one embodiment, a polynucleotide described herein encodes a VL domain comprising the amino acid sequence set forth in SEQ ID NO: 275, 278, 279, 280, 281, 282, 292, 304, 307, 309, 310 or 311. In another embodiment, a polynucleotide described herein encodes a VH domain comprising the amino acid sequence set forth in SEQ ID NO: 182 or 220. In yet another embodiment, a polynucleotide described herein encodes a light chain comprising the amino acid sequence set forth in SEQ ID NO: 49, 100, 101, 103, 104, 105, 106, 107, 117, 129, 132, 134, 135, or 136. In another embodiment, a polynucleotide described herein encodes a heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 2 or 42. In one embodiment, a polynucleotide described herein encodes a VL domain comprising the amino acid sequence set forth in SEQ ID NO: 278. In another embodiment, a polynucleotide described herein encodes a VH domain comprising the amino acid sequence set forth in SEQ ID NO: 477. In yet another embodiment, a polynucleotide described herein encodes a light chain comprising the amino acid sequence set forth in SEQ ID NO: 103. In another embodiment, a polynucleotide described herein encodes a heavy chain comprising the amino acid sequence set forth in SEQ ID NO: 529.

[0130] In some embodiments, the nucleic acid or nucleic acids encode a VH selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and having at least

80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 572-581; and encode a VL selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 582-595.

[0131] In some embodiments, the nucleic acid or nucleic acids encode a HC selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 596-605; and encode a LC selected from the group consisting of SEQ ID NOs: 48-136 and 531-567 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 606-619.

[0132] In some embodiments, the nucleic acid molecule or molecules are codon-biased to enhance expression in a desired host cell, *e.g.*, in human cells, mammalian cells, yeast cells, plant cells, insect cells, or bacterial cells, *e.g.*, *E. coli* cells. Accordingly, provided are polynucleotides encoding an antibody or antigen-binding fragment, as described herein, wherein the polynucleotides are codon-biased, comprise replacement heterologous signal sequences, and/or have mRNA instability elements eliminated. Methods to generate codon-biased nucleic acids can be carried out by adapting the methods described in, *e.g.*, U.S. Patent Nos. 5,965,726; 6,174,666; 6,291,664; 6,414,132; and 6,794,498. Preferred codon usage for expression of the an antibody or antigen-binding fragments in desired host cells is provided, *e.g.*, at kazusa.or.jp/codon/; and genscript.com/tools/codon-frequency-table.

[0133] Illustrative polynucleotides encoding the VH and the VL of the anti-gp120 antibodies and antigen-binding fragments described herein, codon-biased for improved expression in a mammalian host cell, are provided in Tables XII and XIII. Illustrative polynucleotides encoding the HC and the LC of the anti-gp120 antibodies and antigen-

binding fragments described herein, codon-biased for improved expression in a mammalian host cell, are provided in Tables XIV and XV.

[0134] As appropriate, in certain embodiments, the 3'-end of the polynucleotide or polynucleotides encoding the antibodies or antigen-binding fragments described herein, comprise multiple tandem stop codons, *e.g.*, two or more tandem TAG ("amber"), TAA ("ochre") or TGA ("opal" or "umber") stop codons. The multiple tandem stop codons can be the same or different. In embodiments where the polynucleotide is an mRNA, the 3'-end of the polynucleotide can comprise a poly-A tail.

[0135] Also encompassed by this disclosure are polynucleotides encoding an anti-gp120 antibody or antigen-binding fragment thereof, an anti-CD3 antibody or antigen-binding fragment thereof, an anti-CD16 antibody or antigen-binding fragment thereof, or an anti-CD89 antibody or antigen-binding fragment thereof that are optimized, *e.g.*, by codon optimization, replacement with heterologous signal sequences, and elimination of mRNA instability elements. Methods to generate optimized nucleic acids can be carried out by adapting the methods described in, *e.g.*, U.S. Patent Nos. 5,965,726; 6,174,666; 6,291,664; 6,414,132; and 6,794,498.

[0136] In some embodiments, the one or more polynucleotides encoding the antibodies or antigen-binding fragments, described herein, are formulated or encapsulated in a lipid nanoparticle (LNP). As used herein, the term "lipid nanoparticle" refers to one or more spherical nanoparticles with an average diameter of between about 10 to about 1000 nanometers, and which comprise a solid lipid core matrix that can solubilize lipophilic molecules. In certain embodiments, the lipid core is stabilized by surfactants (*e.g.*, emulsifiers), and can comprise one or more of triglycerides (*e.g.*, tristearin), diglycerides (*e.g.*, glycerol behenate), monoglycerides (*e.g.*, glycerol monostearate), fatty acids (*e.g.*, stearic acid), steroids (*e.g.*, cholesterol), and waxes (*e.g.*, cetyl palmitate), including combinations thereof. Lipid nanoparticles are described, for example, in Petrilli et al., *Curr Pharm Biotechnol.* 15:847-55, 2014; and U.S. Patent Nos. 6,217,912; 6,881,421; 7,402,573; 7,404,969; 7,550,441; 7,727,969; 8,003,621; 8,691,750; 8,871,509; 9,017,726; 9,173,853; 9,220,779; 9,227,917; and 9,278,130, each of which is incorporated by reference in its entirety. LNP-encapsulated mRNA molecules encoding a broadly neutralizing antibody are described, *e.g.*, in Pardi, *et al.*, *Nat Commun.* (2017) 8:14630. In certain embodiments, the one or more polynucleotides encoding the antibodies or antigen-binding fragments, described herein, are formulated or encapsulated in an LNP comprised of an ionizable cationic

lipid/phosphatidylcholine/cholesterol/PEG-lipid, *e.g.*, in molar ratios of about 50:10:38.5:1.5 mol mol⁻¹, respectively.

TABLE XII - POLYNUCLEOTIDES ENCODING HEAVY CHAIN VARIABLE REGIONS (VH)	
SEQ ID NO:	Polynucleotide sequence encoding VH
572	CAGGTGCAGTTGTTGCAGTCTGGCGCCGCTGTTACAAAGCCTGGCGCTTCTGTTAGAGTGTCCTGCGAGGCCCTCCGGCTACAACAT CAGAGACTACTTCAATCCACTGGTGGCGGCAGGCTCCAGGACAGGGATTGCAATGGTGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCTTAGACAGTTCCAGGGCAGAGTGTCCTGACCCAGACACGCCCTCTTGGGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCCTGAGATCCGACGATACCGCCGTGTAATCTGCGCCAGACAGAGAAAGCGACTACTGGGATTTTCGATGTGTGGGGCTC TGGCACCCCAAGTGACCGTGTCTCT
573	CAGGTGCAGTGTGTCAGTCTGGCGCCGCTGTGACAAAACAGGCGCTTCTGTGCGGGTGTCCTGCGAGGCCAGCGGCTACAACAT CCGGGACTACTTCAATCACTGGTGGCGCCAGGCCCCCTGGACAGGACTGCAGTGGTGGATGGATCAACCCCAAGACCGGCCAGC CCAAACAACCCAGACAGTTCCAGGGCAGAGTGTCCTGACCCAGACACGCCAGCTGGGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCCTGCGGAGCGACGATACCGCCGTGTAATCTGCGCCAGACAGAGAAAGCGACTACTGGGATTTTCGACGTGTGGGGCAG CGGCACCCCAAGTGACCGTGTCTCT
574	CAGGTGCAGTTGTTGCAGTCTGGCGCCGCTGTTACAAAGCCTGGCGCTTCTGTTAGAGTGTCCTGCGAGGCCCTCCGGCTACAACAT CAGAGACTACTTCAATCCACTGGTGGCGGCAGGCTCCAGGACAGGGATTGCAATGGTGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCTTAGACAGTTCCAGGGCAGAGTGTCCTGACCCAGACACGCCCTCTTTCGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCCTGAGATCCGACGATACCGCCGTGTAATCTGCGCCAGACAGAGAAAGCGACTACTGGGACTTCGATGTGTGGGGCTC TGGCACCCCAAGTGACCGTGTCTCT
575	CAGGTGCAGTGTGTCAGTCTGGCGCCGCTGTGACAAAACAGGCGCTTCTGTGCGGGTGTCCTGCGAGGCCAGCGGCTACAACAT CCGGGACTACTTCAATCACTGGTGGCGCCAGGCCCCCTGGACAGGACTGCAGTGGTGGATGGATCAACCCCAAGACCGGCCAGC CCAAACAACCCAGACAGTTCCAGGGCAGAGTGTCCTGACCCAGACACGCCAGCTTCGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCCTGCGGAGCGACGATACCGCCGTGTAATCTGCGCCAGACAGAGAAAGCGACTACTGGGATTTTCGACGTGTGGGGCAG CGGCACCCCAAGTGACCGTGTCTCT
576	CAGGTCCACTTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCTGGCGCTTCTGTGAGAGTGTCCTGCGAGGCCCTTCGGCTACAAGAT CCGGGACCACTTTATCCACTGGTGGCGACAGGCTCCAGGACAGGGATTGCAATGGTGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCTTAGACAGTTCCAGGGCAGAGTGTCCTGACCCAGACACGCCCTCTTGGGACTTCGACACCTTCAGCTTCTACATGGAC

TABLE XII - POLYNUCLEOTIDES ENCODING HEAVY CHAIN VARIABLE REGIONS (VH)	
SEQ ID NO:	Polynucleotide sequence encoding VH
	CTGAAGGCCGTGGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGAGATCCGACTACTGGGATTTTCGATGTGTGGGGCTC TGGCACCCAAAGTGACCGTGTCCTCT
577	CAGGTCACCTGTCTCAATCTGGCGCCGCTGTACAAAACAGGCGCCTCTGTTAGAGTGTCTTTCGAGGCCAGCGGCTACAAGAT CAGGGACCACTTTTATTCACCTGGTGGCGCCAGGCTCCAGGACAGGGATTCGAGTGGGTCGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCCAGACAGTTCCAGGGCAGAGTGTCTCTGACAAGACACGCCAGCTGGGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCGTGGGAGCGACGACACCGCCATCTATTTTGGCCAGACAGAGAAGCGACTACTGGGATTTTCGATGTGTGGGGCAG CGGCACCCAAAGTGACAGTCTCTTCT
578	CAGGTCACCTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCTGGCGCTTCTGTTCAGAGTGTCTTTCGAGGCCCTCTGGCTACAAGAT CCGGGACCACTTTTATCCACTGGTGGCGACAGGCTCCAGGACAGGGATTCGAGTGGGTCGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCTTAGACAGTTCCAGGGCAGAGTGTCCCTGACCAGACACGCCCTCTTTCGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCGTGGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGAGATCCGACTACTGGGACTTCGATGTGTGGGGCTC TGGCACCCAAAGTGACCGTGTCCTCT
579	CAGGTCACCTGTCTCAATCTGGCGCCGCTGTACAAAACAGGCGCCTCTGTTCAGAGTGTCTTTCGAGGCCAGCGGCTACAAGAT CAGGGACCACTTTTATTCACCTGGTGGCGCCAGGCTCCAGGACAGGGACTTCGAGTGGGTCGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCCAGACAGTTCCAGGGCAGAGTGTCTCTGACAAGACACGCCAGCTTCGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCGTGGGAGCGACGACACCGCCATCTATTTTGGCCAGACAGAGAAGCGACTACTGGGATTTTCGATGTGTGGGGCAG CGGCACCCAAAGTGACAGTCTCTTCT
580	CAGGTCACCTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCTGGCGCTTCTGTTCAGAGTGTCTTTCGAGGCCCTCCGGCTACAACAT CCGGGACTACTTTTATCCACTGGTGGCGCCAGGCTCCAGGACAGGGATTCGAGTGGGTCGGATGGATCAACCTTAAGACCGGCCAGC CTAACAAACCTTAGACAGTTCCAGGGCAGAGTGTCCCTGACCAGACACGCCCTCTTTCGACTTCGACACCTTCAGCTTCTACATGGAC CTGAAGGCCGTGGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGAGATCCGACTACTGGGATTTTCGATGTGTGGGGCTC TGGCACCCAAAGTGACCGTGTCCTCT
581	CAGGTCACCTGTCTCAATCTGGCGCCGCTGTACAAAACAGGCGCCTCTGTTCAGAGTGTCTTTCGAGGCCAGCGGCTACAACAT CCGGGACTACTTTTATTCACCTGGTGGCGCCAGGCTCCAGGACAGGGACTTCGAGTGGGTCGGATGGATCAACCTTAAGACCGGCCAGC

TABLE XII - POLYNUCLEOTIDES ENCODING HEAVY CHAIN VARIABLE REGIONS (VH)	
SEQ ID NO:	Polynucleotide sequence encoding VH
	CTAACAAACCCAGACAGTTCAGGGCAGAGTGTCTCTGACAAAGACAGCCAGCTGGGACTTCGACACCTTCAGCTTCTACATGGACCTGAAGGCCGTGCGAGCGACGACACCGCCATCTATTTTGGCCAGACAGAGAAAGCGACTACTGGGATTCGATGTGTGGGGCAGCGGCACCCAAAGTGACAGTCTCTTCT

TABLE XIII - POLYNUCLEOTIDES ENCODING LIGHT CHAIN VARIABLE REGIONS (VL)	
SEQ ID NO:	Polynucleotide sequence encoding VL
582	GACATCCAGATGACCCAGAGCCCTTCTCTTTATCCGTAGCGTGGCGATACCGTGACCATCACATGCCAAGCTAACGGCTACCTCAACTGGTACCAGCAGCGGAGGGGAAAGCCCCCAAGCTGCTGATCTACGACGGCTCCAAGCTGGAGAGGGAGTGCCTTCCCGGTTCAGCGGAAGAGGTGGGACAAGAATACAATTAAACCATCAACAATTTACAGCCCCGAGGACATCGCTACTTCTGCCAAGTTTACGAGTTCGTGGTGCCCGGCACCTCGTCTGGATCTGAAG
583	GACATCCAGATGACCCAGAGCCCTAGCAGCCTGAGCGCCAGCGTGGCGATACCGTGACCATTACCTGCCAGGCCAACGGCTACCTGAACTGGTATCAGCAGCGGAGAGGCAAGGCCCCCAAGCTGCTGATCTACGACGGCAGCAAGCTGGAAAGAGGCGTGCCAGCAGATTCAGCGGCAGAAGATGGGGCCAGGAGTACAACCTGACCATCAACAACCTGCAGCCCCGAGGATATGCCACATACTTTTGCCAGGTGTACGAGTTCGTGGTGCCCGGCACACGGCTGGACCTGAAA
584	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCTCTGTGGCGATACCGGTACCATCACCTGTGAGGCCAACGGCTACCTGAACTGGTATCAGCAGAGAAAGAGGCAAGGCCCCCAAGCTGCTGATCTACGACGGCTCCAAGCTGGAAAGAGGCGTGCCCTCTCGGTTCCTCTGGCAGAAGATGGGGCCAAAGAGTACAACCTGACCATCAACAACCTGCAGCCTGAGGATATGCCACATACTTTTGCCAGGTGTACGAGTTCGTGGTGCCCTGGCACAAGACTGGACCTGAAG
585	GATATTCAGATGACACAGAGCCCCAGTAGCCTGAGCGCCAGCGTGGCGACACCGCAACCATCACCTGTGAGGCCAACGGCTATCTGAACTGGTATCAACAGAGGAGGGGCAAGGCCCCCAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGCGTTCCAGCCGCTTCAGCGGCAGGAGGTGGGGCCAGGAGTACAACCTTACAATCAACAACCTGCAGCCCCGAGGACATCGCCACCTATTTCTGCCAAGTTTACGAGTTCGTGGTGCCCGGCACACAGGCTGGACCTGAAG

TABLE XIII - POLYNUCLEOTIDES ENCODING LIGHT CHAIN VARIABLE REGIONS (VL)	
SEQ ID NO:	Polynucleotide sequence encoding VL
586	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCCTCTGTGGCGACAGAGCTACCATCACCTGTTCAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAAGAGCGTGCCCTCCAGAT TCTCCGGCTCTAGATGGGGCCAAAGAGTACAACCTGACCATCTCCAGCCTCCAGCCTGAGGATATGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAG
587	GATATTTCAGATGACACAGAGCCCCAGTAGCTGAGCGCCAGCGTGGCGACAGAGCAACCATCACCTGTTCAGGCCAACGGCTATCT GAACTGGTATCAACAGAGAAAGGGCAAGGCCCTAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGCGTTCCAGCGCGCT TCAGCGGCTCAAGGTGGGGCCAGGAGTACAACCTTACAATCTCATCCCTGCAGCCCCGAGGACATGCCACCTATTTCTGCCAAGTT TACGAGTTCTTCGGACCCGGCACACAGGCTGGACCTGAAG
588	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCAGAGTGGCGACACCGTGACAATCACCTGTTCAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAAGAGCGTGCCCGCTAGAT TCTCCGGCAGAAGATGGGGCCAAAGAGTACAACCTGACCATCAACAACCTGCAGCCTGAGGACGTGGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGTGCCCGGCACACAGACTGGACCTGAAG
589	GATATTTCAGATGACACAGAGCCCCAGTAGCTGAGCGCCCGCTGGCGACACCGTGACCATCACCTGTTCAGGCCAACGGCTATCT GAACTGGTATCAACAGAGAGAGGGGCAAGGCCCTAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGCGTTCCCGCACGCT TCAGCGGCAGGAGGTGGGGCCAGGAGTACAACCTTACAATCAACAACCTGCAGCCCCGAGGACGTGCCACCTATTTCTGCCAAGTT TACGAGTTCTTCGTGCCCGGCACACAGGCTGGACCTGAAG
590	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCCTCTGTGGCGATACCGTACCATCACCTGTTCAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAAGAGCGTGCCCTCTCGGT TCTCTGGCAGAAGATGGGGCCAAAGAGTACAACCTGACCATCAACAACCTGCAGCCTGAGGATATGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAG
591	GACATCCAGATGACACAGAGCCCCAGTAGCTGTGCCAGCGTGGAGATACCGCCACAATTACCTGTTCAGGCCAACGGCTACCT GAACTGGTATCAGCAGCGGAGAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCAGCAAGCTGGAAAAGAGCGTGCCAGCAGAT TCAGCGGCAGAAGATGGGGCCAAAGAGTACAACCTGACCATCAACAACCTGCAGCCTGAGGATATGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAG

TABLE XIII - POLYNUCLEOTIDES ENCODING LIGHT CHAIN VARIABLE REGIONS (VL)	
SEQ ID NO:	Polynucleotide sequence encoding VL
592	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCTCTGTGGCGACAGAGCTACCATCACCTGTCTAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAAGAGGCGTGCCCTCCAGAT TCTCCGGCTCTAGATGGGGCCAAAGAGTACACCTGACCATCTCTAGCCTGCAGCCTGAGGATATGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAG
593	GACATCCAGATGACACAGAGCCCTAGCAGCCTGTCTGCCCTCTGTGGCGATAGAGCCACAATCACCTGTCTAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCAGCAAACTGGAAAAGAGGCGTGCCAAAGCAGAT TCAGCGGCTCTAGATGGGGCCAAAGAGTACACCTGACCATCTCTAGCCTGCAGCCTGAGGATATGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAA
594	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCAGAGTGGCGGATACCGCTACCATCACCTGTCTAGGCCAACGGCTACCT GAACTGGTATCAGCAGAGAAAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAAGAGGCGTGCCCGCTAGAT TCTCCGGCAGAAGATGGGGCCAAAGAGTACACCTGACCATCAACAACCTGCAGCCTGAGGACGTGGCCACATACTTTTGCCAGGTG TACGAGTTCTTCGTGCCCGGCACACAGACTGGACCTGAAG
595	GATATTCAGATGACACAGAGCCCCAGTAGCCTGAGCGCCCGCTGGCGGACACCGCGACCATCACCTGTCTAGGCCAACGGCTATCT GAACTGGTATCAACAGAGGAGGGGCAAGGCCCTCAAGCTCTGATATACGACGGCAGCAAGCTGGAGAGGGGCGTTCCCGCACGCT TCAGCGGCAGGAGGTGGGGCCAGGAGTACACCTTACAATCAACAACCTGCAGCCCCGAGGACGTGCCACCTATTCTTGCCAAAGTT TACGAGTTCTTCGTGCCCGGCACACAGGTGGACCTGAAG

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
596	CAGGTGCAGTTGTTGCAGTCTGGCGCCGCTGTTTACAAAAGCCTGGCGCTTCTGTTAGAGTGTCTCGAGGGCTCCGGCTACAACATC AGAGACTACTTTCATCCACTGGTGGCGGCAGGCTCCAGGACAGGATGGATGCAATGGTGGATGGATCAACCTTAAGACCGGCCAGCCT AACAAACCTTAGACAGTTCCAGGCGAGAGTGTCCCTGACACAGACGCCCTCTTGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCCTGAGATCCGACGATACCGCGCTGTACTTCTGGCCAGACAGAGAACGCACTACTGGGATTTTCTGATGTGTGGGCTCTGGC ACCAAAGTGACCGTGTCTCTGCTTCTACCAAGGACCCCTGTGTCTCCCTCTGGCTCCCTCCAGCAAGTCTACCTCTGTTGGAACC GCTGCTTGGGCTGCTTGGTCAAGGATTAATTTCTGAGCCCTGTGACAGTGTCTCTGGAACCTCTGGTGTCTGACCTCCGGCGTGCAC ACATTTCCAGCTGTGTGAGTCTCCGGCTGTACTCTCTGTCTGTGACCGTGCCTCTAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACACAAAGCCTTCCAAACACAAAGGTGGACAAAGTGGAAACCAAGTCTTGGACAAAGACCCACACCTGT CCTCCATGTCTGTCTCAGAACTGTGTGGTGGCGCCGATGTCTTCTGTCTCCCTCCAAAGCCTAAGGACACCCCTGATGATCTCTCGG ACCCCTGAAAGTGACCTGCGTGGTGGATGTCTACGAGGATCCCGAAGTGAAGTCAATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACGCAAGACCAAGCTTAGAGAGGAACAGTACAACTCCACCTACAGAGTGTGTCCGTGTGACCGTGTGACCCAGGATTTGG CTGAACGGCAAGAGTACAAAGTGAAGTGTCCAAAGGCCCTGCTCTGCTGAGGAAAAGACCAATCTTAAGGCTAAGGGCCAG CCTCGGAGCCTCAGGTTTACACACTGCTCTCATCTCGGGAAGAGATGACCAAGAACAGGTGTCACTGACCTGCCTCGTGAAGGGC TTCTACCCCTTCCGATATCGCGTGAATGGAGTCCAAATGGCCAGCCTGAGAACAACTACAAAGACAAACCCCTCTCTGTGTGGACTCC GACGGCTCAATTTCTTCTGTACTCCAAAGCTGACAGTGGACAAAGTCTCGGTGGCAGCAGGGCAACGTGTTCTCTTGTAGTGTGCTGCAC GAGGCCCTGCACCTCCACTATATACCCAGAAAGTCTCTGTCTGAGCCCCGGCAAA
597	CAGGTGCAGTGTGTCAGTCTGGCGCCGCTGTGACAAAACCAAGGCGCTTCTGTGCGGGTGTCTCGAGGGCCAGCGGCTACAACATC CGGGACTACTTTCATTCACTGGTGGCGCCAGGCCCCCTGGACAGGACTGCAGTGGTGGATGGATCAACCCCAAGACCGGCCAGCCCC AACAAACCCAGACAGTTCAGGGCAGAGTGTCCCTGACACAGACAGCCAGCTGGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCCTGCGGAGCGACGATACCGCGCTGTACTTCTGGCCAGACAGAGAAAGCGACTACTGGGATTTTCTGACCGTGTGGGCGACGGC ACCAAAGTGACCGTGTCTGTAGCACCAAGGGCCCCAGCGTGTCTCCCTCTGGCCCCCTAGCAGCAAGAGCAATCTGGCGGAACA GCGGCCCTGGGCTGCCCTGTGAAGGACTACTTTTCCGAGCCGCTGACCGTGTCTGGAACTCTGGCGCTCTGACAAGCGGCGTGCAC ACCTTTCCAGCCGTGTGACAGCAGCGGCTGTACTCTCTGAGCAGCGTCTGTGACAGTGTGCGGAGCAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACACAAAGCCAGCAACACAAAGGTGGACAAAGTGGAAACCAAGAGTGTGACAGTGTGACAGCAACCCACCTGT CCCCCTGTCTTGGCCCCGAACTGTGGTGGCTGGCCCTGACGTGTCTCTGTTCCTCCCAAGCCCAAGGACACCCCTGATGATCAGCCGG ACCCCCGAAAGTGACCTGCGTGGTGGATGTCTCCACGAGGACCCCTGAAAGTGAAGTTCATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACGCAAGACCAAGCTTAGAGAGGAACAGTACAAACAGCACCTACCGGGTGGTGTCCGTGCTGACAGTGTGCTGCACCCAGGACTGG

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
	CTGAACGGGAAAAGAGTACAAAGTGCAAGGTGTCCAAACAAGGCCCTGCCCTGCCCCGAGGAAAAGACCATCTTAAGGCCAAGGGACAG CCCCGCGAGCCCCAGGTGTACACACTGCCTCCAAAGCCGGGAAAGAGATGACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC TTCTACCCCAAGCGACATTTGCCGTGGAATGGGAGAGCAACGGCCAGCCCGAGAACAACTACAGACCAACCCCCCTGTGCTGGACAGC GACGGCTCAATTTCTTCTGTACAGCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCGGCAACGTGTTAGCTGTAGCGTGTTCAT GAGGCTCTGCACAGCCACTACACGCAGAAAGGCCCTCTCCCTGTCTCCGGGTAAA
598	CAGGTGCAGTTGTTGCAGTCTGGCGCCGCTGTTACAAAAGCCCTGGCGCTTCTGTTAGAGTGTCTCGAGGGCCCTCCGGCTACAACATC AGAGACTACTTTCATCCACTGGTGGCGGCAGGCTCCAGGACAGGGATTGCAATGGTCCGATGGATCAACCCCTAAGACCGGCCAGCCT AACAAACCTTAGACAGTTCAGGGCAGAGTGTCCCTGACCAGACAGCCCTCTTTCGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCCTGAGATCCGACGATACCGCGTGTACTTCTGCGCAGACAGAGAACGCACTACTGGGACTTCGATGTGTGGGCTCTGGC ACCAAAGTGACCGTGTCTCTGCTTCTACCAAGGGACCCCTGTGTTCCCTCTGGCTCCCTCCAGCAAGTCTACCTCTGTGGAAACC GCTGCTCTGGGCTGCCCTGGTCAAGGATTACTTTCTGAGCCTGTGACAGTGTCTGGAACTCTGGTGTCTGACCTCCGGCGTGCAC ACATTTCCAGCTGTGCTGCAGTCTCCGGCTGTACTCTGCTCTGCTGACCCGTGCCCTCTAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACCAAGCCTTCCAAACCAAGGTGGACAAGAGTGGAAACCAAGTCTCGCAGCAAGACCCACACCTGT CCTCCATGTCCTGCTCCAGAACTGTGGTGGCGCCCGATGTTCTTCTGTTCCCTCCAAAGCCTAAGGACACCCCTGATGATCTCTCGG ACCCCTGAAAGTGACCTGCGTGGTGGATGTGCTCACGAGGATCCCGAAGTGAAAGTTCATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACCCAAAGACCAAGCCTAGAGAGGAAACAGTACAACTCCACCTACAGAGTGGTGTCCGTGCTGACCGTGTGCAACAGGATTGG CTGAACGGCAAAAGAGTACAAAGTGCAAGGTGTCCAAACAAGGCCCTGCCCTGAGGAAAAGACCATCTCTAAGGCTAAGGGCCAG CCTCGGAGCCTCAGGTTTACACACTGCCCTCCATCTCGGGAAGAGATGACCAAGAACAGGTGTCACTGACCTGCCCTCGTGAAGGGC TTCTACCCCTTCGATATCGCGTGGAAATGGGAGTCCAAATGGCCAGCCTGAGAACAACTACAGACAAACCCCTCCCTGTGCTGGACTCC GACGGCTCAATTTCTTCTGTACTCCAAAGCTGACAGTGGACAAGTCTCGGTGGCAGCGGCAACGTGTTCTCTTGTAGTGTGCTGCAC GAGGCCCTGCACCTCCCACTATACCCAGAAAGTCTCTGCTGAGCCCCCGGCAAA
599	CAGGTGCAGCTGCTGCAGTCTGGCGCCGCTGTGACAAAAACAGGGCGCTTCTGTGCGGGTGTCTCGAGGGCCAGCGGCTACAACATC CGGGACTACTTTCATTTCACTGGTGGCGCCAGGCCCTTGACAGGGACTGGAGTGGTGGATGGATCAACCCCAAGACCGGCCAGCCCC AACAAACCCAGACAGTTCAGGGCAGAGTGTCCCTGACCAGACACGCCAGCTTCGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCCTGCGGAGCGACGATACCGCGTGTACTTCTGGCCAGACAGAGAGCACTACTGGGATTTGACCTGTGGGCGAGCGGC ACCAAAGTGACCGTGTACTGTGTAGCACCAAGGGCCCGAGCGTGTTCCTCTGGCCCTTAGCAGCAAGAGCACATCTGGCGGAACA GCCGCCCTGGGCTGCCCTCGTGAAGGACTACTTTCCCGAGCCCGTGACCGTGTCTCTGGAACTCTGGCGCTCTGACAAGCGGCGTGCAC

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
	ACCTTTCCAGCCGTGCTGCAGAGCAGCGGGCTGTACTCTCTGAGCAGCGTCTGTGACAGTGCCTCAGCAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACACACAAGCCAGCAACACAAAGGTGGACAAAGAGTGGAAACCCAAAGAGTGGCAAGACCCACACCTGT CCCCCTTGCTCCTGCCCCGAACTGCTGGCTGGCTGACGTGTTCTGTTCCCTCCAAAGCCAAAGGACACCCCTGATGATCAGCCCGG ACCCCGAAAGTGACCTGCGTGGTGGATGTCTCCACGAGGACCCCTGAAAGTGAAGTTCATTTGTTACGTGGACGGCGTGGAAAGTG CACAAACGCCAAGACCAAGCCTAGAGAGGAAACAGTACAAACAGCACCTACCCGGTGGTGTCCGTGCTGACAGTGTGACCCAGGACTGG CTGAACGGCAAGAGTACAAAGTGAAGTGTCCAAACAAAGGCCCTGCCCTGCCGAGGAAAAGACCATCTCTAAGGCCAAGGGACAG CCCCAGAGCCCAAGGTGTACACACTGCCCTCCAAAGCCGGGAAAGAGTACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC TTCTACCCCAAGCAGACATTTGCCGTGGAATGGGAGAGCAACGGCCAGACCCGAGAACAACTACAAAGACCAACCCCTGTGCTGGACAGC GACGGCTCAATTTCTTCTGTACAGCAAGCTGACCGTGGACAAAGTCCCGTGGCAGCAGGGCAACGTGTTCAAGCTGTAGCGTGTTCGAT GAGGCTCTGCACAGCCACTACACGCAGAAAGACCTCTCCCTGTCTCCGGGTAAAA
600	CAGGTCCACTTTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCTGGCGCTTCTGTGACAGTGTCTTGGAGGGCCTCTGGCTACAAGATC CGGGACCACCTTTATCCACTGGTGGCGACAGAGCTCCAGGACAGGGATTGAGTGGGTCCGATGGATCAACCTTAAGACCGGCCAGCCT AACAAACCTAGACAGTTCAGGGCAGAGTGTCCCTGACACAGACAGCCTCTTGGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGCGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGAGATCCGACTACTGGGATTTTCGATGTGTGGGGCTCTGGC ACCCAAGTGACCGTGTCTCTGTCTTCTACCAAGGGACCCCTCTGTGTTCCCTCTGGCTCCTTCCAGCAAGTCTACCTCTGGTGGAAACC GCTGCTCTGGGCTGCCGTGGTCAAGGATTAATTTCTGAGCCTGTGACAGTGTCTGGAACTCTGGTGTCTGACCTCCGGCGTGCAC ACATTTCCAGCTGTGCTGCAGTCTCCGGCTGTACTCTGTCTCTGTGCTGACCCGTGCCCTTCTAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACCAAGCCTTCCAAACACAAAGGTGGACAAAGAGTGGAAACCAAGTCTCGGACAAAGACCCACACCTGT CCTCCATGTCTGCTCCAGAACTGCTGGGTGGCTGGCGCCCGATGTCTTCTGTTCCTCCAAAGCCTAAGGACACCCCTGATGATCTCTCGG ACCCTGAAAGTGACCTGCGTGGTGGATGTGTCTACGAGGATCCCGAAAGTGAAGTTCATTTGTTACGTGGACGGCGTGGAAAGTG CACAAACGCCAAGCCTAGAGAGGAAACAGTACAACTCCACCTACAGAGTGGTGTCCGTGCTGACCGTGTGACCCAGGATTTGG CTGAACGGCAAGAGTACAAAGTGAAGTGTCCAAACAAAGGCCCTGCCCTGCTGCTGAGGAAAAGACCATCTCTAAGGCTAAGGGCCAG CCTCGGAGCCTCAGGTTTACACACTGCCCTCCATCTCGGGAAAGATGACCAAGAACAGGTGTCACTGACCTGCCCTCGTGAAGGGC TTCTACCCCTTCGATATCGCGTGGAAATGGGAGTCCCAATGGCCAGCCTGAGAACAACTACAAAGACCAACCCCTCCTGTGCTGGACTCC GACGGCTCAATTTCTTCTGTACTCCAAAGCTGACAGTGGACAAAGTCTCGGTGGCAGCAGGGCAACGTGTTCTCTTGTAGTGTGCTGCAC GAGGCCCTGCACCTCCCACTATACCCAGAAAGTCCCTGTCTCTGTCCCTCTGGCAAA

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
601	CAGGTCCACCTGTCTCAATCTGGCGCCGCTGTATTACAAAACCAGGCGCCTCTGTAGAGTGTCTTGGAGGCCAGCGGCTACAAGATC AGGGACCACCTTTATTCACTGGTGGCGCCAGGCTCCAGGACAGGACTTCAATGGTCCGATGGATCAACCCTAAGACCGGCCAGCCT AACAAACCCAGACAGTTCCAGGCGAGAGTGTCTGTGACAAAGACAGCCAGCTGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGGGAGCGACGACACCGGCATCTATTTTGGCCAGACAGAGAGCGACTACTGGGATTTTCGATGTGTGGGCGACGCGC ACCCAAAGTGACAGTCTTTCTGTAGCACCAAGGGCCCGAGCGTGTCCCTCTGGCCCTAGCAGCAAGAGACATCTGGCGGAACA GCCGCCCTGGGCTGCCCTGTGAAGGACTACTTTCCCGAGCCCGTGACCGTGTCTCTGGAACTCTGGCGCTCTGACAAGCGCGCTGCAC ACCTTTCCAGCCGTGTGACAGAGCGGCGCTGTACTCTGTAGCAGCGTGTGACAGTGCAGAGCTCTCTGGGCAACCCACACCTGT TACATCTGCAACGTGAACACAAAGCCAGCAACACAAAGGTGGACAAAGAGTGGAAACCCAAAGAGCTGGACAAAGACCCACACCTGT CCCCCTGTCTGCCCGGAACTGTGGCTGGCGCTGACGTGTCTGTCTCCCGCAAGCCCAAGGACACCCCTGATGATCAGCCCGG ACCCCGGAAGTGACCTGCGTGGTGGATGTCTCCAGGAGCCCTGAAAGTGAAGTCAATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACCCAGCAAGCCTAGAGAGGAACAGTACAAACAGACCTACCCGGTGTCTCCGTGTGACAGTGTGACAGGCTGTGACAGGACTGG CTGAACGGCAAGAGTACAAAGTGAAGGTGTCCAAAGGCCCTGCCCTGCCGAGGAAAAGACCATCTTAAGGCCAAGGGACAG CCCCGCGAGCCCGAGGTGTACACACTGCGCTCCAAAGCCGGGAAGAGATGACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC TTCTACCCAGCGACATTTGCCGTGGAATGGGAGAGCAACGGCCAGCCCGAGAACAACTACAAAGACCAACCCCTGTGCTGGACAGC GACGGCTCAATTTCTTCTGTACAGCAAGCTGACCGTGGACAAAGTCCCGGTGGCAGCAGGGCAACGTGTTCAGCTGTAGCGTGTTCAT GAGGCTCTGCACAGCCACTACACGCAAGAGCCCTCTCCCTGTCTCCGGGTAAA
602	CAGGTCCACCTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCCTGGCGCTTCTGTGACAGTGTCTTGGAGGCCCTCTGGCTACAAGATC CGGGACCACCTTTATCCACTGGTGGCGACAGGCTCCAGGACAGGATTTGCAGTGGTCCGATGGATCAACCCTAAGACCGGCCAGCCT AACAAACCCTAGACAGTTCCAGGCGAGAGTGTCCCTGACACAGACAGCCCTTTTCGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGAGATCCGACTACTGGGACTTCGATGTGTGGGCTCTGGC ACCCAAAGTGACCGTGTCTGCTTCTACCAAGGGACCCCTGTGTCTCCCTCTGGCTCCCTCCAGCAAGTCTACCTCTGGTGGAAAC GCTGTCTTGGGCTGCCCTGGTCAAGGATTAATTTCTGTAGCCCTGTGACAGTGTCTCTGGAACTCTGGTGTCTGACCTCCGGCGCTGCAC ACATTTCCAGCTGTGCTGCAGTCTCCGGCTGTACTCTGTCTGTCTGTGACCGCTGTCCCTCTAGCTCTCTGGGCAACCCAGAC TACATCTGCAACGTGAACCAAGCCTTCCAAACCAAGGTGGACAAAGAGTGGAAACCCAAAGTCTCTGGCAAGACCCACACCTGT CCTCCATGTCTGTGCTCCAGAACTGTGGCTGGCGCCCGATGTCTTTCTGTTCCTCCAAAGCCTTAAGGACACCCCTGATGATCTCTCGG ACCCCTGAAGTGACCTGCGTGGTGGATGTCTCACAGGATCCCGGAAGTGAAGTCAATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACCCAGCAAGCCTAGAGAGGAACAGTACAACTCCACCTACAGAGTGGTGTCCGTGCTGACCGTGTGCAACCGAGGATTTGG

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
	CTGAACGGGAAAAGAGTACAAAGTGCAAGGTGTCCAAACAAGGCCCTGCCCTGCTCTGAGGAAAAGACCAATCTCTAAGGCTAAGGGCCAG CCTCGCGAGCCTCAGGTTTACACACTGCCTCCATCTCGGGAAGAGATGACCAAGAACAGAGTGTCACCTGACCTGCCTCGTGAAGGGC TTCTACCCCTTCCGATATCGCCGTGGAATGGAGTCCAAATGGCCAGCCTGAGAACAACTACAGACAAACCCCTCCTGTGCTGGACTCC GACGGCTCAATCTTCTCTACTCCAAAGCTGACAGTGGACAAGTCTCGGTGGCAGCAGGGCAACGTGTTCTCTTTGTAGTGTGCTGCAC GAGGCCCTGCACCTCCACTATATACCCAGAAAGTCCCTGTCTCTGTCCTCCCTGGCAAA
603	CAGGTCCACCTGTCTCAATCTGGCGCCGCTGTACAAAACCAAGGCCCTCTGTTAGAGTGTCTTGGAGGGCCAGCGGCTACAAGATC AGGGACCACCTTTATTCACTGGTGGCGCCAGGCTCCAGGACAGGGACTTCAATGGTGGATGGATCAACCCTAAGACCGGCCAGCCT AACAAACCCAGACAGTTCCAGGGCAGAGTGTCTGTACAAGACACGCCAGCTTCGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGGGAGCGACGACACCGCCATCTAATTTTGGCCAGACAGAGAAGCGACTACTGGGATTTTCGATGTGTGGGCGACGGGC ACCAAAGTGACAGTCTCTTCTGTAGCACCAAGGGCCCGAGCGTGTCTCCCTCTGGCCCCCTAGCAGCAAGAGCACATCTGGCGGAACA GCGGCCCTGGGCTGCCCTCGTGAAGGACTACTTTTCCGAGCCCGTGACCGGTCTCTGGGCTCTGACAAGCGGCGTGCAC ACCTTTCCAGCCGTGTGCAGAGCAGCGGCTGTACTCTCTGAGCAGCGTCTGTGACAGTCCCCAGCAGCTCTCTGGGACCCAGACC TACATCTGCAACGTGAACCAAGCCAGCAACCAAGGTGGACAAGAGTGGAAACCAAGAGCTGCGACAAGACCCACACCTGT CCCCCTGTCTTCTGCCCCGAACTGTGGTGGCTGGCCCTGACGTGTTCTGTTCCCCCAAGCCCAAGGACACCTGATGATCAGCCGG ACCCCGGAAGTGACCTGCGTGGTGGATGTCTCCACAGGACCCCTGAAAGTGAAGTTCATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACCCAAAGCCCTAGAGAGGAAACAGTACAACAGCACCTACCGGGTGGTGTCCGTGCTGACAGTGTGCTGCCAGGACTGG CTGAACGGCAAGAGTACAAAGTGCAAGGTGTCCAAACAAGGCCAGAGATGACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC CCCCGAGCCCCAGGTGTACACACTGCCCTCCAAAGCCGGGAAGAGATGACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC TTCTACCCCGAGCGACATTTGCCGTGGAATGGGAGAGCAACGGCCAGCCCGAGAACAACTACAAAGACCAACCCCTGTGCTGGACAGC GACGGCTCAATCTTCTGTACAGCAAGCTGACCGTGGACAAGTCCCGGTGGCAGGCGAACGTGTTCAAGCTGTAGCGTGTGTCAT GAGGCTCTGCACAGCCACTACACGCAGAAAGCCCTCTCCCTGTCTCCCGGGTAAA
604	CAGGTCCACCTGTCTCAATCTGGCGCCGCTGTGACAAAAGCCCTGGCGCTTCTGTGACAGTGTCTTGGAGGGCCCTCCGGCTACAACATC CGGGACTACTTTTATCCACTGGTGGCGGCAGGCTCCAGGACAGGGATTTGCAATGGGTGGATGGATCAACCCTAAGACCGGCCAGCCT AACAAACCCTAGACAGTTCCAGGGCAGAGTGTCCCTGACCAGACACGCCCTCTTGGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGGGAGCGACGACACCGCTATCTACTTTTGGCCAGACAGATCCGACTACTGGGATTTTCGATGTGTGGGCTCTGGC ACCAAAGTGACCGTGTCTGCTTCTACCAAGGGACCCCTGTGTCTCCCTCTGGCTCCCTCCAGCAAGTCTACCTCTGTTGGTGAACC GCTGCTCTGGGCTGCCCTGGTCAAGGATTACTTTCTGTAGCCCTGTGACAGTGTCTTGGAACTCTGGTGTCTGTGACCTCCGGCGTGCAC

TABLE XIV - POLYNUCLEOTIDES ENCODING HEAVY CHAIN (HC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING HC
	ACATTTCCAGCTGTGCTGAGTCTCTCCGGCTGTACTCTCTGTCTCTGTCTGTCGACCGTGCCTTTCTAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACACACAAGCCTTCAAACACAAAGGTGGACAAAGAGTGGAAACCCAGTCTTGCGACAAAGACCCACACCTGT CCTCCATGTCTGCTCCAGAACTGTGCTGGCTGGCCCGATGTCTTCTGTTCCTCCAAAGCCTAAGGACACCCCTGATGATCTCTCTCGG ACCCCTGAAAGTGACCTGCGTGGTGGATGTCTCACGAGGATCCCGAAGTGAAGTTCATTTGTTACGTGGACGGCGTGGAAAGTG CACAAACGCCAAGACCAAGCCTAGAGAGGAACAGTACAACTCCACCTACAGAGTGGTGTCCGTGCTGACCCGTGCTGCACCCAGGATTTGG CTGAACGGCAAGAGTACAAAGTGAAGTGTCCAAACAAGGCCCTGCCCTGCTGAGGAAAAGACCAATCTCTAAGGCTAAGGGCCAG CCTCGGAGCCTCAGGTTTACACACTGCCCTCCATCTCGGGAAGAGATGACCAAGAACAGGTGTCACTGACCTGCCCTCGTGAAGGGC TTTACCCCTTCCGATATCGCCGTGGAATGGGAGTCCCAATGGCCAGCCTGAGAAACAATACAAAGACAAACCCCTCCTGTGCTGGACTCC GACGGCTCAATTTCTTCTGTACTCCAAGCTGACAGTGGACAAGTCTCGGTGGCAGCGGCAACGTGTTCTCTTGTAGTGTGCTGCAC GAGGCCCTGCACCTCCACTATATACCCAGAAAGTCCCTGTCTCTGTCCCTTGGCAAA
605	CAGGTCCACCTGTCTCAATCTGGCGCCGCTGTATACAAAACACAGGCGCTCTGTTAGAGTGTCTTGGAGGGCCAGCGGTACAAACATC CGGACTACTTTTATTCACTGGTGGCGCCAGGCTCCAGGACAGGACTTCAATGGTCCGATGGATCAACCTTAAGACCGGCCAGCCT AAACAACCCAGACAGTTCAGGGCAGAGTGTCTGTACAAGACAGCCAGCTGGACTTCGACACCTTCAGCTTCTACATGGACCTG AAGGCCGTGCGGAGCGACGACACCGCCATCTATTTTTCGCCAGACAGAGAAAGCGACTACTGGGATTTTCGATGTGTGGGCGACGGGC ACCAAAGTGACAGTCTCTTCTGTAGCACCAAGGGCCCGAGCGTGTTCCTCTGGCCCCCTAGCAGCAAGAGACATCTGGCGGAACA GCCGCCCTGGGCTGCCCTCGTGAAGGACTACTTTTCCGAGCCCGTGACCGTGTCTGGAACTCTGGCGCTCTGACAAGCGCGCTGCAC ACCTTTCCAGCCGTGCTGCAGAGCAGCGGCTGTACTCTGTAGCAGCGTCTGTGACAGTCCCCAGCAGCTCTCTGGGCACCCAGACC TACATCTGCAACGTGAACCAAGCCAGCAACACAAAGGTGGACAAAGAGTGGAAACCAAGAGCTGCGACAAAGACCCACACCTGT CCCCCTTGTCTTGTCTGCCCCGAACTGTGGTGGCTGGCCCTGACAGTGTTCCTGTTCCTCCCAAGCCCAAGGACACCCCTGATGATCAGCCGG ACCCCGAAGTGACCTGCGTGGTGGATGTGTCCACGAGGACCCCTGAAAGTGAAGTTCATTTGGTACGTGGACGGCGTGGAAAGTG CACAAACGCCAAGCCTAGAGAGGAACAGTACAAACAGACCTACCCGGGTGGTGTCCGTGCTGACAGTGTGTCACCCAGGACTGG CTGAACGGCAAGAGTACAAAGTGAAGGTGTCCAAACAAGGCCCTGCCCTGAGGAAAAGACCAATCTCTAAGGCCAAGGGACAG CCCCGCGAGCCCGAGGTGTACACACTGCCCTCCAAAGCCGGGAAGAGATGACCAAGAACAGGTGTCCCTGACCTGTCTCGTGAAGGC TTTCTACCCCGAGCGACATTTGCCGTGGAATGGGAGAGCAACGGCCAGCCCGAGAAACAATACAAAGACCAACCCCTCCTGTGCTGGACAGC GACGGCTCAATTTCTTCTGTACAGCAAGCTGACCGTGGACAAGTCCCGGTGGCAGCGGCAACGTGTTCTAGCTGTAGCGTGTGTCAT GAGGCTCTGCACAGCCACTACACGCAGAAAGAGCCCTCTCCCTGTCTCCCGGTAAA

TABLE XV - POLYNUCLEOTIDES ENCODING LIGHT CHAIN (LC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING LC
606	GACATCCAGATGACCCAGAGCCCTTCTCTTTATCCGCTAGCGTGGCGGATACCGTGACCATCACATGCCAAGCTAACGGCTACCTC AACTGGTATCAGCAGCGGAGGGGAAAAGCCCCCAAGCTGCTGATCTACGACGGCTCCAAAGCTGGAGGGGAGTGCCCTCCCGGTTTC AGCGGAAGGAGGTGGGGACAAGAATACAAATTAACCATCAAAATTTACAGCCGAGGACATCGCTACCTACTTCTGCCAAGTTTAC GAGTTCGTGGTGCCCGGCACTCGTCTGGATCTGAAAGAGGACCGTGGCGGCCCTCCGTGTTCACTCTTCCCTTCCGACGAGCAG CTGAAAGTCCGGCACCGCTCCGTGGTGTGTTTACTGAACAACCTTCTACCTCGTGAGGCCAAGGTGCAAGTGAAGTGGACAACGCT TTACAGTCCGGCAACTCCCAAGAATCCGTGACCCGAGCAAGATAGCAAGGACTCCACCTACTCCCTCTCCAGCACTTTAACTTTATCC AAGCCGACTACGAGAAGCACAAAGGTGTACGCTGTGAGGTGACCCACCAAGGTCTGTCTCTCCCTCCGTGACAAAGTCTTCAATCGG GGCGAGTGT
607	GACATCCAGATGACCCAGAGCCCTAGCAGCCTGAGCGCGAGCGTGCGGATACCGTGACCAATTACCTGCCAGGCCAACGGCTACCTG AACTGGTATCAGCAGCGGAGAGGCAAGGCCCAAGCTGCTGATCTACGACGGCAGCAAGCTGGAAGAGGGCGTGCCACGAGATTTC AGCGGCAGAAAGATGGGCGCAGGAGTACAACCTGACCATCAACAACCTGACAGCCGAGGATATCGCCACATACTTTTGCCAGGTGTAC GAGTTCGTGGTGCCCGGCACACGGCTGGACCTGAAACGTACGGTGGCTGCACCATCTGTCTTCACTCTTCCGCCATCTGATGAGCAG TTGAAATCTGAACTGCCCTCTGTTGTGCTGCTGAATACTTCTATCCAGAGGCCCAAGTACAGTGGAAAGTGGATAACGCC CTCCAATCGGGTAACTCCAGGAGAGTGTACAGAGCAGCAAGCAAGCACTACAGCCTCAGCAGCACCTGACGCTGAGC AAAGCAGACTACGAGAAACACAAAAGTCTACGCGCTGCGAAAGTCAACCATCAGGGCCTGAGCTCGCCCGTCAAAAAGAGCTTCAACAGG GGAGAGTGT
608	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGTGCGCTGTGGCGGATACCGGTACCAATCACCTGTGAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAAGAGGCAAGGCCCTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAGAGGGCGTGCCCTCTCGGTTTC TCTGGCAGAAAGATGGGCGCAAGAGTACAACCTGACCATCAACAACCTGACAGCCTGAGGATATCGCCACATACTTTTGCCAGGTGTAC GAGTTCGTGGTGGCCTGGCAAGACTGGACCTGAAAGAACCTGAAAGAACCTGGCCGCTCTTCCGTGTTCACTCTTCCACCATCTGACGAGCAG CTGAAAGTCTGGCACCGCTTCTGTGCTGTGCTGCTGAAACAACTTCTACCTCGGGAAGCCAAAGTGCAGTGAAGTGGACAATGCC CTGCAGTCCGGCAACTCCCAAGAGTCTGTGACCGAGCAGGACTCCAAGGACTCTACCTACAGCCTGTCTCTCCACACTGACCCCTGTCT AAGGCCGACTACGAGAAGCACAAAGGTGTACGCGCTGCGAAAGTGAACCATCAGGGGACTGTCTAGCCCCCGTGACCAAGTCTTCAACAGA GGCGAGTGT
609	GATATTGAGATGACACAGAGCCCCAGTAGCCTGAGCGCCAGCGTGGGCGACACCGCAACCAATCACCTGTGAGGCCAACGGCTATCTG AACTGGTATCAACAGAGGAGGGGCAAGGCCCTCAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGGCGTTCCACAGCCGCTTC

TABLE XV - POLYNUCLEOTIDES ENCODING LIGHT CHAIN (LC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING LC
	AGCGGCAGGAGGTGGGGCCAGGAGTACAAACCTTACAAATCAAAACCTGAGCCGAGGACATGCCACCTATTTCTGCCAAGTTTAC GAGTTCTGTGGTGGCCCGCACAGGCTGACCTGAAAGGGACCGTGGCCGCCCGCCAGCGTGTTCATCTTCCCTCCAGCGACGAGCAG CTGAAAGTCTGGCACCGCCAGCGTGGTGTGGCTGTGAACAACTTACCCCGGAGGCCAAGGTGCAGTGGAAAGGTGGACAAACGCC CTGCAGAGCGGCAACAGCCAGGAGAGCGTGACCGAGGAGGACTCCAAAGGACAGCACCTACAGCCTGAGCAGCACCTGACCCCTGAGC AAGGCCGACTACGAGAAACAAAGGTGTACCGCTGCGAGGTGACCCACAGGGGACTGTCTAGCCCCCTGACCAAGAGCTTCAACCCGG GGCGAGTGC
610	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCCTCTGTGGGCGACAGAGCTACCATCACCTGTCAAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAAGGGCAAGGCCCTAAAGTGTGATCTACGACGGCTCCAAACTGGAAAGAGGGCGTCCCTCCAGATTTC TCCGGCTCTAGATGGGGCCAAAGATACAACTGACCATCTCCAGCCTCCAGGATATCGCCACATATCTTTTGGCCAGGTGTAC GAGTTCTTCGGCCCTGGCACACAGACTGGACCTGAAAGAGAAACAGTGGCCGCTCCTTCGCTGTTCATCTTCCCACCATCTGACGAGCAG CTGAAAGTCTGGCACCCGCTTCTGTCTGTGGCTGCTGAAACAACTTACCCCTCGGGAAGCCCAAGGTGCAGTGGAAAGGTGGACAAATGCT CTCCAGTCCGGCAACTCCAAAGATCTGTGACCGAGCAGGACTCCAAGGACTCTACCTACAGCCTGTCTCCACACTGACCCCTGTCT AAGGCCGACTACGAGAAACAAAGGTGTACCGCTGCGAAAGTGACCCATCAGGGACTGTCTAGCCCCCTGACCAAGTCTTCAACACAGA GGCGAGTGT
611	GATATTCAGATGACACAGAGCCCCAGTAGCCTGAGGGCCAGCGTGGGCGACAGAGCAACCATCACCTGTCAAGGCCAACGGCTATCTG AACTGGTATCAACAGAGAAAGGGCAAGGCCCGCCAAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGCGGTCCAGAGCCGCTTC AGCGGCTCAAGGTGGGCGCAGGAGTACAACTTACAATCTCATCCCTGACGCCCGAGGACATCGCCACCTATTTCTGCCAAGTTTAC GAGTTCTTCGACCCCGCACAGGCTGGACCTGAAAGCGGACCGTGGCCGCCCCCAGCGTGTTCATCTTCCCTCCAGCGACGAGCAG CTGAAAGTCTGGCACCCCGCAGCGTGGTGTGGCTGCTGAAACAACTTACCCCGGAGGCCAAGGTGCAGTGGAAAGGTGGACAAACGCC CTGCAGAGCGGCAACAGCCAGGAGCGTGACCGAGGACTCCAAAGGACAGCACTACAGCCTGAGCAGCACCTGACCCCTGAGC AAGGCCGACTACGAGAAACAAAGGTGTACCGCTGCGAGGTGACCCACAGGGGACTGTCTAGCCCCCTGACCAAGAGCTTCAACCCGG GGCGAGTGC
612	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCAGAGTGGGCGACACCGTGACAAATCACCTGTCAAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAAGGGCAAGGCCCGCCAAAGCTGCTGATCTACGACGGCTCCAAACTGGAAAGAGGGCGTCCCGCTAGATTTC TCCGGCAGAAAGATGGGGCCAAAGATACAACTGACCATCAACAACTTGAGCGCTGAGGACGTGGCCACATATCTTTTGGCCAGGTGTAC GAGTTCTATCTGTGGCCCGCACAGACTGGACCTGAAAGAGAAACAGTTGCCGCTCCTTCCGCTGTTCATCTTCCCACCTTCCGACGAGCAG

TABLE XV - POLYNUCLEOTIDES ENCODING LIGHT CHAIN (LC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING LC
	CTGAAAGTCTGGCACAGCTTCTGTCTGTGTGCTGCTGAAACAACTTTACCTCTGGGAAAGCCAAAGGTGCA GTGGAAGTGGACAAATGCC CTGCAGTCCGGCAACTCCAAAGAGTCTGTACCGAGAGGACTCCAAAGGACTTACCTACAGCCTGTCTCCACACACTGACCCCTGTCT AAGGCCGACTACGAGAAAGCACAAAGGTGTACGCCCTGCGAAAGTGACCCATCAGGGACTGTCTAGCCCCGTGACCAAGTCTTCAACACAGA GCGGAGTGT
613	GATATTCAGATGACACAGAGCCCCAGTAGCCTGAGCGCCCGCTGGGCGACACCGTGACCATCACCTGTCAAGGCCAACGGCTATCTG AACTGGTATCAACAGAGGAGGGGCAAGGCCCAAGCTCTGATATACGACGGCAGCAAGCTGGAGAGGGGCGTTCCCGCACGCTTC AGCGCAGGAGGTGGGGCCAGGAGTACAACTTACAAATCAACAACTTGACGCCCGAGGACGTGCCACCTATTTCTGCCCAAGTTTAC GAGTTCATCTGTGCCCGGCACAGGCTGGACCTGAAAGGGACCGTGGCGGCCCAAGCGTGTTCATCTTCCCTCCAGCGACGAGCAG CTGAAAGTCTGGCACCGCCAGCGTGGTGTGCTGAAACAACTTCTACCCCGCGAGGCCAAAGGTGCAGTGGAAAGTGGACAAACGCC CTGCAGAGCGGCAACAGCCAGGAGAGCGTGACCGAGGAGACTCCAAAGGACAGCACCTACAGCCTGAGCAGCACCCCTGACCCCTGAGC AAGGCCGACTACGAGAAAGCACAAAGGTGTACGCCCTGCGAGGTGACCCACAGGGACTGTCTAGCCCCGTGACCAAGAGCTTCAACCCGG GCGGAGTGC
614	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGTGCTGTGGCGGATACCGGTACCATCACCTGTCAAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAAGAGGCAAGGCCCTAAAGCTGTGATCTACGACGGCTCCAAACTGGAAGAGGGCGTGCCCTCTCGGTTTC TCTGGCAGAAAGATGGGGCCAAAGATACACCTGTGACCATCAACAACTTGACGCTGAGGATATCGCCACATATCTTTTGGCAGGTGTAC GAGTTCCTTCGGCCCTGGCACAGACTGGACCTGAAAGAGAAACAGTGGCGGCTCCTTCCGTGTTCATCTTCCACCATCTGACGAGCAG CTGAAAGTCTGGCACCGCTTCTGTCTGTGCTGCTGAAACAACTTCTACCCCTCGGGAGGCCAAGGTGCAGTGGAAAGTGGACAAATGCC CTGCAGTCCGGCAACTCCAAAGAGTCTGTGACCGGAGGAGACTCCAAAGGACTCTACCTACAGCCTGTCTCCACACTGACCCCTGTCT AAGGCCGACTACGAGAAAGCACAAAGGTGTACGCCCTGCGAAAGTGACCCATCAGGGACTGTCTAGCCCCGTGACCAAGTCTTCAACACAGA GCGGAGTGT
615	GACATCCAGATGACACAGAGCCCTAGCAGCCTGTCTGCCAGCGTGGGAGATACCGCCACAAATTAACCTGTCAAGGCCAACGGCTACCTG AACTGGTATCAGCAGCGGAGAGGCAAGGCCCTAAAGCTGTGATCTACGACGGCAGCAAGCTGGAAGAGGGCGTGCCACGAGATTTC AGCGGCAGAAAGATGGGGCCAAAGATACACCTGTGACCATCAACAACTTGACGCTGAGGATATTCGCCACATATCTTTTGGCAGGTGTAC GAGTTCCTTCGGCCCTGGCACAGACTGGACCTGAAAGAGAAACAGTGGCGGCTCCTAGCGTGTTCATCTTCCACCTTCCGACGAGCAG CTGAAAGTCTGGCACAGCCTCTGTCTGTGCTGCTGAAACAACTTCTACCCCGAGAGAACCAAGGTGCAGTGGAAAGTGGACAAACGCC CTGCAGAGCGGCAATAGCCAAAGAGAGCGGTGACCGGAGGAGCAAGGACTCTACCTACTCTCTGAGGAGCACCCCTGACACTGAGC

TABLE XV - POLYNUCLEOTIDES ENCODING LIGHT CHAIN (LC)

SEQ ID No :	POLYNUCLEOTIDE SEQUENCE ENCODING LC
616	AAGCCGACTACGAGAAGCACAAAAGTGACGCCCTGCGAAGTGACCCACGAGGCCCTTTCTAGCCCTGTGACCAAGAGCTTCAACCGG GGCGAGTGT GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCGCTCTGTGGCGACAGAGTACCATCACCTGTGAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAGAGGCAAGGCCCTTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAGAGGCGTGCCCTCCAGATTCTC TCCGGCTCTAGATGGGGCCAAGAGTACACCCCTGACCATCTCTAGCCTGCAGCCTGAGGATATCGCCACATACTTTTGCAGGTGTAC GAGTTCTTCGGCCCTGGCACCAAGACTGGACCTGAAGAGAACAGTGGCCGCTCCTTCGTGTTTCATCTTCCACCATCTGACGAGCAG CTGAAGTCTGGCACCGCTTCTGTGCTGCTGAACAACTTCTACCCCTCGGAAGCCAAAGTGCAGTGAAGGTGGACAATGCC CTGCAGTCCGGCAACTCCCAAGAGTCTGTGACCGAGCAGGACTCCAAGGACTTACCTACAGCCTGTCTCCACACTGACCCCTGTCT AAGCCGACTACGAGAAGCACAAAGGTGTACGCCCTGCGAAGTGACCCATCAGGGACTGTCTAGCCCCGTGACCAAAGTCTTCAACAGA GGCGAGTGT
617	GACATCCAGATGACACAGAGCCCTAGCAGCCTGTCTGCGCTCTGTGGCGATAGAGCCACAATCACCTGTGAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAGAGGCAAGGCCCTTAAGCTGCTGATCTACGACGGCAGCAAACTGGAAGAGGCGTGCCAAAGCAGATTCTC AGCGGCTCTAGATGGGGCCAAGAGTACACCCCTGACCATCTCTAGCCTGCAGCCTGAGGATATCGCCACATACTTTTGCAGGTGTAC GAGTTCTTCGGCCCTGGCACCAAGACTGGACCTGAAACGTACGGTGGCTGCACCATCTGTCTTTCATCTTCCGCCATCTGATGAGCAG TTGAAATCTGAAACTGCCCTCTGTTGTGCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTACAGTGAAGGTGGATAACGCC CTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCCTACAGCCTCAGCAGCACCTGACGCTGAGC AAAGCAGACTACGAGAACAACAAAAGTCTAGCCCTGCGAAGTGACCCATCAGGGCCTGAGCTCGCCCCGTCACAAAGAGCTTCAACAGG GGAGAGTGT
618	GACATCCAGATGACCCAGTCTCCATCCTCTCTGTCTGCCAGAGTGGCGGATACCGTACCATCACCTGTGAGGCCAACGGCTACCTG AACTGGTATCAGCAGAGAAGAGGCAAGGCCCTTAAGCTGCTGATCTACGACGGCTCCAAACTGGAAGAGGCGTGCCCGCTAGATTCTC TCCGGCAGAAGATGGGGCCAAGAGTACACCCCTGACCATCAACAACTGAGCCTGAGGACGTGGCCACATACTTTTGCAGGTGTAC GAGTTCTATCGTGCCCGGCACCAAGACTGGACCTGAAGAGAACAGTTGCCGCTCCTTCGTGTTTCATCTTCCACCTTCCGACGAGCAG CTGAAGTCTGGCACAGCTTCTGTGCTGCTGCTGAACAACTTCTACCCCTCGGAAGCCAAAGTGCAGTGAAGGTGGACAATGCC CTGCAGTCCGGCAACTCCCAAGAGTCTGTGACCGAGCAGGACTCCAAGGACTTACCTACAGCCTGTCTCCACACTGACCCCTGTCT AAGCCGACTACGAGAAGCACAAAGGTGTACGCCCTGCGAAGTGACCCATCAGGGACTGTCTAGCCCCGTGACCAAAGTCTTCAACAGA GGCGAGTGT

TABLE XV - POLYNUCLEOTIDES ENCODING LIGHT CHAIN (LC)	
SEQ ID NO:	POLYNUCLEOTIDE SEQUENCE ENCODING LC
619	GATATTGAGATGACACAGAGAGCCCCAGTAGCCTGAGCGCCCCGCTGGGCGGACACCCGCGACCATCACCTGTGAGGCCCAACGGCTATCTG AACTGGTATCAACAGAGAGGGGCAAGCCCCCAAGCTCCTGATATACGACGGCAGCAAGCTGGAGAGGGCGGTTCCCGCACGCTTC AGCGGCAGGAGGTGGGGCCAGGAGTACACCCCTTACAATCAACAAACCTGACGCCGAGGACGTGCCACCTATTCTGCCAAGTTTAC GAGTTCATCGTGGCCCGGCACACAGGCTGGACCTGAAAGCGGACCGTGGCCGCCGCCAGCGTGTTCATCTTCCCTCCAGCGACGAGCAG CTGAAAGTCTGGCACCGCCAGCGTGGTGTGCCTGCTGAACAACCTTCTACCCCGCGAGGCCCAAGGTGCAGTGAAGGTGGACAACGCC CTGCAGAGCGGCAACAGCCAGGAGCGGTGACCCGAGCAGGACTCCAAGGACAGCACCTACAGCCTGAGCAGCACCTGACCCCTGAGC AAGCCGACTACGAGAAGCACAAAGGTGTACGCCCTGGGAGGTGACCCACACAGGGACTGTCTAGCCCCGCTGACCAAGAGCTTCAACCCGG GGCGAGTGC

Vectors and Host Cells

[0137] This disclosure also encompasses vectors comprising a nucleic acid(s) disclosed herein. A vector can be of any type, for example, a recombinant vector such as an expression vector. Vectors include, but are not limited to, plasmids, cosmids, bacterial artificial chromosomes (BAC) and yeast artificial chromosomes (YAC) and vectors derived from bacteriophages or plant or animal (including human) viruses. Vectors can comprise an origin of replication recognized by the proposed host cell and in the case of expression vectors, promoter and other regulatory regions recognized by the host cell. In additional embodiments, a vector comprises a polynucleotide encoding an antibody of the disclosure operably linked to a promoter and optionally additional regulatory elements. Certain vectors are capable of autonomous replication in a host into which they are introduced (*e.g.*, vectors having a bacterial origin of replication can replicate in bacteria). Other vectors can be integrated into the genome of a host upon introduction into the host, and thereby are replicated along with the host genome. Vectors include, but are not limited to, those suitable for recombinant production of the antibodies disclosed herein.

[0138] The choice of the vector is dependent on the recombinant procedures followed and the host used. Introduction of vectors into host cells can be effected by *inter alia* calcium phosphate transfection, virus infection, DEAE-dextran-mediated transfection, lipofectamine transfection or electroporation. Vectors may be autonomously replicating or may replicate together with the chromosome into which they have been integrated. In certain embodiments, the vectors contain one or more selection markers. The choice of the markers may depend on the host cells of choice. These include, but are not limited to, kanamycin, neomycin, puromycin, hygromycin, zeocin, thymidine kinase gene from Herpes simplex virus (HSV-TK), and dihydrofolate reductase gene from mouse (*dhfr*). Vectors comprising one or more nucleic acid molecules encoding the antibodies described herein, operably linked to one or more nucleic acid molecules encoding proteins or peptides that can be used to isolate the antibodies, are also covered by the disclosure. These proteins or peptides include, but are not limited to, glutathione-S-transferase, maltose binding protein, metal-binding polyhistidine, green fluorescent protein, luciferase and beta-galactosidase.

[0139] In other embodiments, the vector that is used is pcDNA™3.1+ (ThermoFisher, MA).

[0140] The disclosure also provides host cells comprising a nucleic acid or a vector described herein. Any of a variety of host cells can be used. In one embodiment, a host cell is a prokaryotic cell, for example, *E. coli*. In another embodiment, a host cell is a eukaryotic cell, for example, a yeast cell, a plant cell (*e.g.*, a tobacco plant cell), or a mammalian cell, such as a Chinese Hamster Ovary (CHO) cell (*e.g.*, CHO-S, [®], CHO-K1, CHO-K1a, CHO DG44, ExpiCHO™), COS cells, BHK cells, NSO cells or Bowes melanoma cells. Examples of human host cells are, *inter alia*, HeLa, 911, AT1080, A549, 293 and HEK293 (*e.g.*, HEK293E, HEK293T, Expi293™) cells. In addition, antibodies (*e.g.*, scFv's) can be expressed in a yeast cell such as *Pichia* (see, *e.g.*, Powers *et al.*, *J Immunol Methods*. 251:123-35 (2001)), *Hansenula*, or *Saccharomyces*. Antibody production in transgenic tobacco plants and cultured plant cells is described, *e.g.*, in Sacks, *et al.*, *Plant Biotechnol J.* (2015) 13(8):1094-105; Klimyuk, *et al.*, *Curr Top Microbiol Immunol.* (2014) 375:127-54 and Cramer, *et al.*, *Curr Top Microbiol Immunol.* (1999) 240:95-118.

[0141] In some embodiments, the host cell predominantly sialylates N-linked glycosylation sites with the variable regions of an immunoglobulin antigen binding domain. In some embodiments, the polynucleotides encoding an antibody or antigen-binding fragment thereof, as described herein, are expressed in a host cell that sialylates at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the variable domains (Fv, particularly VL) of expressed antibodies or antigen-binding fragments thereof. In some embodiments, the cell sialylates at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the VL of expressed antibodies or antigen-binding fragments. In some embodiments, the N-linked glycosylation sites in the VL have a sialic acid occupancy (*e.g.*, a glycan comprising one or two terminal sialic acid residues) of at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or more. As used herein, "occupancy" refers to the percentage of the time that a glycan is attached at a predicted amino acid glycosylation site. In some embodiments, the asparagine at VL amino acid position 72 according to Kabat numbering (N72) is sialylated. In some embodiments, the sialylated N-linked glycosylation sites in the VL comprise from 1 to 5 sialic acid residues, *e.g.*, from 1 to 4 sialic acid residues, *e.g.*, from 1 to 3 sialic acid residues, *e.g.*, from 1 to 2 sialic acid

residues. Human and hamster host cells predominantly sialylate with N-acetylneuraminic acid (NANA). In some embodiments, the VL are sialylated or predominantly sialylated with N-acetylneuraminic acid (NANA). Mouse host cells predominantly sialylate with N-glycolylneuraminic acid (NGNA). In some embodiments, the VL are sialylated or predominantly sialylated with N-acetylneuraminic acid (NGNA). In some embodiments, the sialic acid residues are present in biantennary structures. In some embodiments, the sialic acid residues are present in complex N-linked glycan structures (*e.g.*, can contain almost any number of the other types of saccharides, including more than the original two N-acetylglucosamines). In some embodiments, the sialic acid residues are present in hybrid N-linked glycan structures (*e.g.*, can contain mannose residues on one side of the branch, while on the other side a N-acetylglucosamine initiates a complex branch).

[0142] The term “nucleic acid molecule” refers to a polymeric form of nucleotides and includes both sense and anti-sense strands of RNA, cDNA, genomic DNA, and synthetic forms and mixed polymers of the above. As used herein, the term nucleic acid molecule may be interchangeable with the term polynucleotide. In some embodiments, a nucleotide refers to a ribonucleotide, deoxynucleotide or a modified form of either type of nucleotide, and combinations thereof. The terms also include, but are not limited to, single- and double-stranded forms of DNA. In addition, a polynucleotide, *e.g.*, a cDNA or mRNA, may include either or both naturally occurring and modified nucleotides linked together by naturally occurring and/or non-naturally occurring nucleotide linkages. The nucleic acid molecules may be modified chemically or biochemically or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those of skill in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analogue, internucleotide modifications such as uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages (*e.g.*, phosphorothioates, phosphorodithioates, etc.), pendent moieties (*e.g.*, polypeptides), intercalators (*e.g.*, acridine, psoralen, etc.), chelators, alkylators, and modified linkages (*e.g.*, alpha anomeric nucleic acids, etc.). The above term is also intended to include any topological conformation, including single-stranded, double-stranded, partially duplexed, triplex, hairpinned, circular and padlocked conformations. A reference to a nucleic acid sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a particular sequence should be understood to encompass

its complementary strand, with its complementary sequence. The term also includes codon-optimized nucleic acids.

[0143] The term “operably linked” refers to two or more nucleic acid sequence elements that are usually physically linked and are in a functional relationship with each other. For instance, a promoter is operably linked to a coding sequence if the promoter is able to initiate or regulate the transcription or expression of a coding sequence, in which case, the coding sequence should be understood as being “under the control of” the promoter.

[0144] A “substitution,” as used herein, denotes the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

[0145] An “isolated” nucleic acid refers to a nucleic acid molecule that has been separated from a component of its natural environment. An isolated nucleic acid includes a nucleic acid molecule contained in cells that ordinarily contain the nucleic acid molecule, but the nucleic acid molecule is present extrachromosomally or at a chromosomal location that is different from its natural chromosomal location. “Isolated nucleic acid encoding an antibody or fragment thereof” refers to one or more nucleic acid molecules encoding antibody heavy and light chains (or fragments thereof), including such nucleic acid molecule(s) in a single vector or separate vectors, and such nucleic acid molecule(s) present at one or more locations in a host cell.

[0146] The term “vector,” as used herein, refers to a nucleic acid molecule capable of propagating another nucleic acid to which it is linked. The term includes the vector as a self-replicating nucleic acid structure as well as the vector incorporated into the genome of a host cell into which it has been introduced. Some vectors are suitable for delivering the nucleic acid molecule or polynucleotide of the present application. Certain vectors are capable of directing the expression of nucleic acids to which they are operatively linked. Such vectors are referred to herein as expression vectors.

[0147] The terms “host cell,” “host cell line,” and “host cell culture” are used interchangeably and refer to cells into which exogenous nucleic acid has been introduced, including the progeny of such cells. Host cells include “transformants” and “transformed cells,” which include the primary transformed cell and progeny derived therefrom without regard to the number of passages. Progeny may not be completely identical in nucleic acid content to a parent cell, but may contain mutations. Mutant progeny that have the same function or biological activity as screened or selected for in the originally transformed cell are included herein.

[0148] A polynucleotide “variant,” as the term is used herein, is a polynucleotide that typically differs from a polynucleotide specifically disclosed herein in one or more substitutions, deletions, additions and/or insertions. Such variants may be naturally occurring or may be synthetically generated, for example, by modifying one or more of the polynucleotide sequences of the invention and evaluating one or more biological activities of the encoded polypeptide as described herein and/or using any of a number of techniques well known in the art.

[0149] The term “variant” may also refer to any naturally occurring or engineered molecule comprising one or more nucleotide or amino acid mutations.

[0150] Further provided is a chimeric antigen receptor (CAR) including an antigen-binding antibody fragment as described herein. In certain embodiments, the CAR is expressed on a T-cell or a NK cell. Further provided is a CAR T-cell including a CAR as described herein. In certain embodiments, the T-cell is a CD4⁺ T-cell, a CD8⁺ T-cell, or a combination thereof. In certain embodiments, the cell is administered to a subject. In certain embodiments, the cell is autologous. In certain embodiments, the cell is allogeneic.

Methods of Producing Antibodies

[0151] Monospecific antibodies that bind to gp120 and bispecific antibodies that bind to gp120 and human CD3 (*e.g.*, human CD3 ϵ or human CD3 δ) or to gp120 and CD89 can be produced by any method known in the art for the synthesis of antibodies, for example, by chemical synthesis or by recombinant expression techniques.

[0152] Methods of making monospecific antibodies are very well known in the art. Methods of making bispecific antibodies are described, for example, in U.S. Pat. Nos. 5,731,168; 5,807,706; 5,821,333; and U.S. Appl. Publ. Nos. 2003/020734 and 2002/0155537. Bispecific tetravalent antibodies, and methods of making them are described, *e.g.*, in WO 02/096948 and WO 00/44788, the disclosures of both of which are herein incorporated by reference in its entirety. In addition, other publications relating to making bispecific antibodies include WO 91/00360, WO 92/08802, WO92/05793, and WO 93/17715; Tutt et al., *J. Immunol.* 147:60-69 (1991); U.S. Pat. Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819 and 9,212,230; and Kostelny et al., *J. Immunol.* 148:1547-1553 (1992).

[0153] Another exemplary method of making bispecific antibodies is by the knobs-into-holes technology (Ridgway et al., *Protein Eng.*, 9:617-621 (1996); WO 2006/028936). The mispairing problem of Ig heavy chains that is a chief drawback for making bispecific antibodies is reduced in this technology by mutating selected amino acids forming the interface of the CH3 domains in IgG. At positions within the CH3 domain at which the two heavy chains interact directly, an amino acid with a small side chain (hole) is introduced into the sequence of one heavy chain and an amino acid with a large side chain (knob) into the counterpart interacting residue location on the other heavy chain. In some instances, antibodies of the disclosure have immunoglobulin chains in which the CH3 domains have been modified by mutating selected amino acids that interact at the interface between two polypeptides so as to preferentially form a bispecific antibody. The bispecific antibodies can be composed of immunoglobulin chains of the same subclass or different subclasses. In one instance, a bispecific antibody that binds to gp120 and CD3 comprises a T366W (EU numbering) mutation in the “knobs chain” and T366S, L368A, Y407V (9EU numbering) mutations in the “hole chain.” In certain embodiments, an additional interchain disulfide bridge is introduced between the CH3 domains by, e.g., introducing a Y349C mutation into the “knobs chain” and a E356C mutation or a S354C mutation into the “hole chain.” In certain embodiments, R409D, K370E mutations are introduced in the “knobs chain” and D399K, E357K mutations in the “hole chain.” In other embodiments, Y349C, T366W mutations are introduced in one of the chains and E356C, T366S, L368A, Y407V mutations in the counterpart chain. In some embodiments, Y349C, T366W mutations are introduced in one chain and S354C, T366S, L368A, Y407V mutations in the counterpart chain. In some embodiments, Y349C, T366W mutations are introduced in one chain and S354C, T366S, L368A, Y407V mutations in the counterpart chain. In yet other embodiments, Y349C, T366W mutations are introduced in one chain and S354C, T366S, L368A, Y407V mutations in the counterpart chain (all EU numbering).

[0154] Another exemplary method of making bispecific antibodies is by using the Bispecific T-cell Engagers (BiTEs®) platform. BiTEs are made by genetically fusing a first scFv (e.g., an scFv that binds gp120) to a second scFv (e.g., an scFv that binds human CD3) via flexible peptide linker (e.g., GGGGS (SEQ ID NO: 429)). See, e.g., Staerz et al., *Nature*, 314:628-631 (1985); Mack et al., *PNAS*, 92:7021-7025 (1995); Huehls et al., *Immunol. Cell Biol.*, 93:290-296 (2015).

- [0155] Another exemplary method of making bispecific antibodies is by using the Dual-Affinity Re-targeting (DART) platform. This technology is based on the diabody format of Holliger et al. (*PNAS*, 90:6444-6448 (1993)) and further improved for stability and optimal pairing of the VH and VL chains (Johnson et al., *J Mol. Biol.*, 399:436-449 (2010); Sung et al., *J Clin Invest.*, 125(11): 4077-4090 (2015)).
- [0156] Yet another exemplary method of making bispecific antibodies is by using the Trifunctional Hybrid Antibodies platform - Triomab®. This platform employs a chimeric construction made up of half of two full-length antibodies of different isotypes, mouse IgG2a and rat IgG2b. This technology relies on species-preferential heavy/light chain pairing associations. See, Lindhofer et al., *J Immunol.*, 155:219-225 (1995).
- [0157] A further exemplary method of making bispecific antibodies is by using the TandAb® platform. This technology is based on the diabody concept but are designed as a single polypeptide chain VH1-VL2-VH2-VL1 comprising short linkers to prevent intra-chain pairing. Head-to-tail dimerization of this single chain results in the formation of a tetravalent homodimer (Kipriyanov et al., *J Mol. Biol.*, 293:41-56 (1999)).
- [0158] Yet another method for making bispecific antibodies is the CrossMab technology. CrossMab are chimeric antibodies constituted by the halves of two full-length antibodies. For correct chain pairing, it combines two technologies: (i) the knob-into-hole which favors a correct pairing between the two heavy chains; and (ii) an exchange between the heavy and light chains of one of the two Fabs to introduce an asymmetry which avoids light-chain mispairing. See, Ridgway et al., *Protein Eng.*, 9:617-621 (1996); Schaefer et al., *PNAS*, 108:11187-11192 (2011). CrossMabs can combine two or more antigen-binding domains for targeting two or more targets or for introducing bivalency towards one target such as the 2:1 format.
- [0159] The antibodies of this disclosure may be produced in bacterial or eukaryotic cells. Antibodies can also be produced in eukaryotic cells such as transformed cell lines (*e.g.*, CHO-based or CHO-origin cell lines (*e.g.*, CHO-S, CHO DG44, ExpiCHO™, CHOZN® ZFN-modified GS-/- CHO cell line, CHO-K1, CHO-K1a), 293E, 293T, COS, NIH3T3). In addition, antibodies (including antibody fragments, *e.g.*, Fabs, scFv's) can be expressed in a yeast cell such as *Pichia* (see, *e.g.*, Powers et al., *J Immunol Methods*, 251:123-35 (2001)), *Hansenula*, or *Saccharomyces*. In one embodiment, the antibodies described herein are produced in a CHO cell line, *e.g.*, a CHO-S, CHO DG44, ExpiCHO™, CHOZN®, CHO-K1 or CHO-K1a cell line. To produce the antibody of interest, a polynucleotide encoding the antibody is constructed, introduced into an

expression vector, and then expressed in suitable host cells. Standard molecular biology techniques are used to prepare the recombinant expression vector, transfect the host cells, select for transformants, culture the host cells, and recover the antibody.

[0160] If the antibody is to be expressed in bacterial cells (*e.g.*, *E. coli*), the expression vector should have characteristics that permit amplification of the vector in the bacterial cells. Additionally, when *E. coli* such as JM109, DH5 α , HB101, or XL1-Blue is used as a host, the vector must have a promoter, for example, a lacZ promoter (Ward *et al.*, 341:544-546 (1989), araB promoter (Better *et al.*, *Science*, 240:1041-1043 (1988)), or T7 promoter that can allow efficient expression in *E. coli*. Examples of such vectors include, for example, M13-series vectors, pUC-series vectors, pBR322, pBluescript, pCR-Script, pGEX-5X-1 (Pharmacia), "QIAexpress system" (QIAGEN), pEGFP, and pET (when this expression vector is used, the host is preferably BL21 expressing T7 RNA polymerase). The expression vector may contain a signal sequence for antibody secretion. For production into the periplasm of *E. coli*, the *pelB* signal sequence (Lei *et al.*, *J. Bacteriol.*, 169:4379 (1987)) may be used as the signal sequence for antibody secretion. For bacterial expression, calcium chloride methods or electroporation methods may be used to introduce the expression vector into the bacterial cell.

[0161] If the antibody is to be expressed in animal cells such as CHO, CHO-S, CHO DG44, CHOZN®, ExpiCHO™, CHO-K1, CHO-K1a, COS, and NIH3T3 cells, the expression vector includes a promoter necessary for expression in these cells, for example, an SV40 promoter (Mulligan *et al.*, *Nature*, 277:108 (1979)), MMLV-LTR promoter, EF1 α promoter (Mizushima *et al.*, *Nucleic Acids Res.*, 18:5322 (1990)), or CMV promoter. In addition to the nucleic acid sequence encoding the immunoglobulin or domain thereof, the recombinant expression vectors may carry additional sequences, such as sequences that regulate replication of the vector in host cells (*e.g.*, origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see *e.g.*, U.S. Pat. Nos. 4,399,216, 4,634,665 and 5,179,017). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin, or methotrexate, on a host cell into which the vector has been introduced. Examples of vectors with selectable markers include pMAM, pDR2, pBK-RSV, pBK-CMV, pOPRSV, and pOP13.

[0162] In one embodiment, antibodies are produced in mammalian cells. Exemplary mammalian host cells for expressing an antibody include Chinese Hamster Ovary (CHO) cells, including, *e.g.*, CHO-S, CHO DG44, ExpiCHO™, CHOZN®, CHO-K1 or CHO-

K1a cells) (including *dhfr*⁻ CHO cells, described in Urlaub and Chasin (1980) *Proc. Natl. Acad. Sci. USA* 77:4216-4220, used with a DHFR selectable marker, *e.g.*, as described in Kaufman and Sharp (1982) *Mol. Biol.* 159:601-621), human embryonic kidney 293 cells (*e.g.*, 293, 293E, 293T), COS cells, NIH3T3 cells, human B-cells, lymphocytic cell lines, *e.g.*, NS0 myeloma cells and SP2 cells, and a cell from a transgenic animal, *e.g.*, a transgenic mammal. For example, in some embodiments, the cell is a mammary epithelial cell.

[0163] In an exemplary system for antibody expression, recombinant expression vectors encoding the antibody heavy chain and the antibody light chain of an antibody of this disclosure are introduced into *dhfr*⁻ CHO cells by calcium phosphate-mediated transfection. In a specific embodiment, the *dhfr*⁻ CHO cells are cells of the DG44 cell line, such as DG44i (*see, e.g.*, Derouaz et al., *Biochem Biophys Res Commun.*, 340(4):1069-77 (2006)). Within the recombinant expression vectors, the antibody heavy and light chain genes are each operatively linked to enhancer/promoter regulatory elements (*e.g.*, derived from SV40, CMV, adenovirus and the like, such as a CMV enhancer/AdMLP promoter regulatory element or an SV40 enhancer/AdMLP promoter regulatory element) to drive high levels of transcription of the genes. The recombinant expression vectors also carry a *DHFR* gene, which allows for selection of CHO cells that have been transfected with the vector using methotrexate selection/amplification. The selected transformant host cells are cultured to allow for expression of the antibody heavy and light chains and the antibody is recovered from the culture medium.

[0164] Antibodies can also be produced by a transgenic animal. For example, U.S. Pat. No. 5,849,992 describes a method of expressing an antibody in the mammary gland of a transgenic mammal. A transgene is constructed that includes a milk-specific promoter and nucleic acids encoding the antibody of interest and a signal sequence for secretion. The milk produced by females of such transgenic mammals includes, secreted-therein, the antibody of interest. The antibody can be purified from the milk, or for some applications, used directly. Animals are also provided comprising one or more of the nucleic acids described herein.

[0165] The antibodies of the present disclosure can be isolated from inside or outside (such as medium) of the host cell and purified as substantially pure and homogenous antibodies. Methods for isolation and purification commonly used for antibody purification may be used for the isolation and purification of antibodies, and are not limited to any particular method. Antibodies may be isolated and purified by

appropriately selecting and combining, for example, column chromatography, filtration, ultrafiltration, salting out, solvent precipitation, solvent extraction, distillation, immunoprecipitation, SDS-polyacrylamide gel electrophoresis, isoelectric focusing, dialysis, and recrystallization. Chromatography includes, for example, affinity chromatography, ion exchange chromatography, hydrophobic chromatography, gel filtration, reverse-phase chromatography, and adsorption chromatography (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed Daniel R. Marshak et al., Cold Spring Harbor Laboratory Press, 1996). Chromatography can be carried out using liquid phase chromatography such as HPLC and FPLC. Columns used for affinity chromatography include protein A column and protein G column. Examples of columns using protein A column include Hyper D, POROS, and Sepharose FF (GE Healthcare Biosciences). The present disclosure also includes antibodies that are highly purified using these purification methods.

Pharmaceutical Compositions

[0166] This disclosure also includes pharmaceutical compositions comprising an antibody described herein, or a polynucleotide encoding an antibody described herein, and a pharmaceutically acceptable diluent, carrier or excipient. In certain embodiments, the pharmaceutical composition comprises a therapeutically effective amount of the antibody or polynucleotide.

[0167] Various pharmaceutically acceptable diluents, carriers, and excipients, and techniques for the preparation and use of pharmaceutical compositions will be known to those of skill in the art in light of the present disclosure. Illustrative pharmaceutical compositions and pharmaceutically acceptable diluents, carriers, and excipients are also described in Remington: The Science and Practice of Pharmacy 20th Ed. (Lippincott, Williams & Wilkins 2003); Loyd V. Allen Jr (Editor), "Remington: The Science and Practice of Pharmacy," 22nd Edition, 2012, Pharmaceutical Press; Brunton, Knollman and Hilal-Dandan, "Goodman and Gilman's The Pharmacological Basis of Therapeutics," 13th Edition, 2017, McGraw-Hill Education / Medical; McNally and Hastedt (Editors), "Protein Formulation and Delivery, 2nd Edition, 2007, CRC Press; Banga, "Therapeutic Peptides and Proteins: Formulation, Processing, and Delivery Systems," 3rd Edition, 2015, CRC Press; Lars Hovgaard, Frokjaer and van de Weert (Editors), "Pharmaceutical Formulation Development of Peptides and Proteins," 2nd Edition, 2012, CRC Press; Carpenter and Manning (Editors), "Rational Design of Stable Protein Formulations: Theory and Practice," 2002, Springer (Pharmaceutical

Biotechnology (Book 13)); Meyer (Editor), "Therapeutic Protein Drug Products: Practical Approaches to Formulation in the Laboratory, Manufacturing, and the Clinic, 2012, Woodhead Publishing; and Shire, "Monoclonal Antibodies: Meeting the Challenges in Manufacturing, Formulation, Delivery and Stability of Final Drug Product, 2015, Woodhead Publishing.

[0168] In some embodiments, each carrier, diluent or excipient is "acceptable" in the sense of being compatible with the other ingredients of the pharmaceutical composition and not injurious to the subject. Often, the pharmaceutically acceptable carrier is an aqueous pH-buffered solution. Some examples of materials which can serve as pharmaceutically-acceptable carriers, diluents or excipients include: sterile water; buffers, *e.g.*, phosphate-buffered saline; sugars, such as lactose, glucose, trehalose and sucrose; starches, such as corn starch and potato starch; cellulose, and its derivatives, such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients, such as cocoa butter and suppository waxes; oils, such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols, such as propylene glycol; polyols, such as glycerin, sorbitol, mannitol and polyethylene glycol; esters, such as ethyl oleate and ethyl laurate; agar; buffering agents, such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol; phosphate buffer solutions; amino acids (*e.g.*, charged amino acids, including without limitation, aspartate, asparagine, glutamate, glutamine, histidine, lysine); and other non-toxic compatible substances employed in pharmaceutical formulations. Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

[0169] The formulation of and delivery methods of pharmaceutical compositions will generally be adapted according to the site and the disease to be treated. Exemplary formulations include, but are not limited to, those suitable for parenteral administration, *e.g.*, intravenous, intra-arterial, intramuscular, or subcutaneous administration, including formulations encapsulated in micelles, liposomes or drug-release capsules (active agents incorporated within a biocompatible coating designed for slow-release); ingestible formulations; formulations for topical use, such as creams, ointments and gels; and other formulations such as inhalants, aerosols and sprays.

Methods of Use

[0170] This disclosure provides methods for treating or preventing an HIV infection or a related disease or disorder in a subject in need thereof (*e.g.*, a human subject), comprising providing to a subject in need thereof an effective amount of an antibody or antibodies described herein, or a polynucleotide encoding the antibody or antibodies. As used herein, the term “effective amount” in the context of the administration of a therapy to a subject refers to the amount of a therapy that achieves a desired prophylactic or therapeutic effect. The polynucleotide may be present in a vector, *e.g.*, a viral vector. In some embodiments, the related disease or disorder is caused by infection with HIV. In other embodiments, it is acquired immune deficiency syndrome (AIDS). In certain embodiments, the subject is a virologically suppressed HIV-infected mammal, while in other embodiments, the subject is a treatment-naïve HIV-infected mammal. In certain embodiments, a treatment-naïve subject has a viral load between 10^3 and 10^5 copies/ml, and in certain embodiments, a virologically suppressed subject has a viral load < 50 copies/ml. In another embodiment, the subject is a mammal, *e.g.*, a human. In certain embodiments, the subject has been diagnosed with an HIV, *e.g.*, HIV-1 or HIV-2, infection or a related disease or disorder, *e.g.*, AIDS, or is considered at risk for developing an HIV, *e.g.*, HIV-1 or HIV-2, infection or a related disease or disorder, *e.g.*, AIDS. Subjects at risk for HIV-related diseases or disorders include patients who have come into contact with an infected person or who have been exposed to HIV in some other way. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of HIV-related disease or disorder, such that a disease or disorder is prevented or, alternatively, delayed in its progression. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody

or antigen-binding fragment thereof comprises a VH sequence set forth in SEQ ID NO: 477 and a VL sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a heavy chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 529 and a light chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 103. In certain embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain sequence set forth in SEQ ID NO: 529 and a light chain sequence set forth in SEQ ID NO: 103.

[0171] Also provided are methods for preventing or inhibiting an increase in HIV virus titer, virus replication, virus proliferation or an amount of an HIV viral DNA, HIV proviral DNA, or HIV viral protein in a subject (*e.g.*, a human subject). In one embodiment, the method comprises providing to the subject in need thereof an amount of an antibody or antibodies (or their antigen-binding fragments) described herein, or a polynucleotide encoding the antibody or antibodies (or their antigen-binding fragments), effective to prevent an increase in HIV titer, virus replication, or an amount of an HIV protein of one or more HIV strains or isolates in the subject. In certain embodiments, the method further comprises measuring an amount of HIV viral or proviral DNA or protein at one or more time points, *e.g.*, before and after the subject is provided with an antibody or antibodies of the present disclosure. Methods and biomarkers for determining an amount of HIV viral or proviral DNA or protein in a subject are known and available in the art, and described for example, in Siliciano, J.D. et al., *Curr Opin HIV AIDS*, 5(6):491-7 (2010), and Rouzioux, C. et al., *Curr Opin HIV AIDS*, 8(3):170-5 (2013).

[0172] In certain aspect, an antibody or antibodies of the present disclosure may be used in, for example, methods of inhibiting certain viruses such as HIV isolates described herein, prophylactic inhibiting or preventing infections of certain viruses such as HIV isolates described herein, detection of certain viruses such as HIV isolates described herein in a sample, inhibiting certain viruses such as HIV isolates described herein, or diagnosis of certain viruses such as HIV isolates described herein.

[0173] For *in vivo* treatment of mammalian subject, *e.g.*, humans, the subject may be administered or provided a pharmaceutical composition comprising an antibody or antibodies described herein. When used for *in vivo* therapy, an antibody or antibodies described herein are typically administered or provided to the patient in therapeutically effective amounts (*i.e.*, amounts that eliminate or reduce the patient's viral burden and/or viral reservoir). The antibodies are administered or provided to a mammalian subject, *e.g.*, a human, in accord with known methods, such as, but not limited to, intravenous administration, *e.g.*, as a bolus or by continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intraarticular, intrasynovial, intrathecal, oral, topical, or inhalation routes. The antibodies may be administered parenterally, when possible, at the target cell site, or intravenously. In one embodiment, administration of the antibody or antibodies to the subject is via an intravenous route. In another embodiment, administration of the antibody or antibodies to the subject is via a subcutaneous route. In additional embodiments, pharmaceutical compositions of the disclosure are administered to a subject systemically, parenterally, or locally.

[0174] In certain embodiments, the present disclosure provides a method for treating an HIV infection, comprising administering to a human subject in need thereof a therapeutically effective amount of an antibody or antibodies disclosed herein. In some embodiments, the present disclosure provides a method for preventing an HIV infection, comprising administering to a human subject in need thereof a therapeutically effective amount of an antibody or antibodies disclosed herein.

Combination Therapy

[0175] In certain embodiments, this disclosure provides a method for treating (*e.g.*, including long-term or extended suppression) or preventing an HIV infection in a human subject having, or at risk of having, the HIV infection. The method comprises administering to the human subject a therapeutically effective amount of an antibody or antibodies disclosed herein, or a pharmaceutical composition thereof, in combination with a therapeutically effective amount of one or more (*e.g.*, one, two, three, one or two, or one to three) additional therapeutic agents. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139,

140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-binding fragment thereof comprises a VH sequence set forth in SEQ ID NO: 477 and a VL sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a heavy chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 529 and a light chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 103. In certain embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain sequence set forth in SEQ ID NO: 529 and a light chain sequence set forth in SEQ ID NO: 103. In one embodiment, a method for treating an HIV infection in a human subject having or at risk of having the infection is provided, the method comprising administering to the human subject a therapeutically effective amount of an antibody or antibodies disclosed herein, or a pharmaceutically acceptable salt thereof, in combination with a therapeutically effective amount of one or more (*e.g.*, one, two, three, one or two, or one to three) additional therapeutic agents. In some embodiments, after one or more administrations of the antibody or antigen-binding fragments thereof, optionally with one or more additional therapeutic agents, the subject does not exhibit symptoms of HIV or AIDS in the absence of anti-retroviral treatment (ART) for at least 6 months, at least 1 year, at least 2 years, at least 3 years, or more. In some embodiments, after one or more administrations of the binding molecule, the subject has a viral load of copies/ml blood of less than 500, *e.g.*, less than 400, less than 300, less than 200, less than 100, less than 50, in the absence of anti-retroviral treatment (ART) for at least 6 months, at least 1 year, at least 2 years, at least 3 years, or more.

[0176] Multiple clinical studies have now shown that treatment of HIV infected individuals with single broadly neutralizing antibodies (bNAbs) leads to temporary suppression of sensitive viruses, followed by rapid outgrowth of resistant viruses – many of which appear to be rare pre-existing viral variants.

[0177] Antibody A and Antibody B were previously shown to neutralize 96% of 118 cross-clade viruses tested *in vitro* (Scheid et al., Science, 333: 1633-1637 (2011)). The clinical trials showed that many HIV infected patients receiving the antibody treatment exhibited rare and pre-existing resistant clones, even when their plasma HIV isolates appeared to be sensitive to the antibody (Caskey et al., Nature, 522:487-491 (2016); Scheid et al., Nature, 535:556-560 (2016)). These results suggested that Antibody A may be broad when tested against HIV isolates collected from different patients (inter-patient breadth), yet it may not neutralize 100% of viral isolates within individual patients (intra-patient breadth).

[0178] An antibody known as 10-1074, part of the PGT121 lineage and taken from the same donor and with similar neutralizing breadth, has also been tested in clinical trials (Mouquet et al., PNAS, 109:E3268-3277 (2012); Caskey et al., Nature Medicine, 23:185-191 (2017)). 10-1074 was originally shown to neutralize approximately 66% of 60 viruses tested at an IC₅₀ below 50 µg/mL (Mouquet et al., PNAS (supra)). The 10-1074 trials showed that in many patients received 10-1074 therapy, there were resistant clones, even when the plasma HIV isolates appeared to be sensitive to the antibody (Caskey et al. Nature Medicine (supra)). This data suggests that most patients may harbor rare pre-existing viral variants that are resistant to 10-1074. These 10-1074 resistance variants showed correlated cross-resistance to PGT121, consistent with close evolutionary relationship between 10-1074 and PGT121. However, nearly all of the resistant viruses isolated during the 10-1074 clinical trial were sensitive to neutralization by Antibody A (Caskey et al. (supra)). This data suggests that combination antibody therapy, using complementary bNAbs, may allow for more complete intra-patient viral coverage.

[0179] The bNAb combinations may achieve complete intra-patient viral coverage. In some embodiments, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and another anti-HIV bNAbs antibody (*i.e.*, a neutralizing antibody that neutralizes multiple HIV-1 viral strains). Various bNAbs are known in the art and may be used in this invention. Examples include, but are not limited to, those

described in U.S. Patent No. 8673307, 9,493,549, 9,783,594, WO2014/063059, WO2012/158948, WO2015/117008, and PCT/US2015/41272, and WO2017/096221, including antibodies 12A12, 12A21, NIH45-46, bANC131, 8ANC134, IB2530, INC9, 8ANC195, 8ANC196, 10-259, 10-303, 10-410, 10-847, 10-996, 10-1074, 10-1121, 10-1130, 10-1146, 10-1341, 10-1369, and 10-1074GM. Additional examples include those described in Klein et al., *Nature*, 492(7427): 118-22 (2012), Horwitz et al., *Proc Natl Acad Sci U S A*, 110(41): 16538-43 (2013), Scheid, et al., *Science*, 333 : 1633-1637 (2011), Scheid, et al., *Nature*, 458:636-640 (2009), Eroshkin et al, *Nucleic Acids Res.*, 42 (Database issue):D1 133-9 (2014), Mascola et al., *Immunol Rev.*, 254(l):225-44 (2013), such as 2F5, 4E10, M66.6, CAP206-CH12, 10E81 (all of which bind the MPER of gp41); PG9, PG16, CH01-04 (all of which bind V1V2-glycan), 2G12 (which binds to outer domain glycan); b12, HJ16, CH103-106, VRC01-03, VRC-PG04, 04b, VRC-CH30-34, 3BNC62, 3BNC89, 3BNC91, 3BNC95, 3BNC104, 3BNC176, and 8ANC131 (all of which bind to the CD4 binding site).

[0180] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAb)) that binds to an epitope or region of gp120 selected from the group consisting of: (i) third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan; (ii) second variable loop (V2) and/or Env trimer apex; (iii) gp120/gp41 interface; or (iv) silent face of gp120. The foregoing epitopes or regions of gp120 bound by broadly neutralizing antibodies are described, *e.g.*, in McCoy, *Retrovirology* (2018) 15:70; Sok and Burton, *Nat Immunol.* 2018 19(11):1179-1188; Possas, et al., *Expert Opin Ther Pat.* 2018 Jul;28(7):551-560; and Stephenson and Barouch, *Curr HIV/AIDS Rep* (2016) 13:31-37, which are hereby incorporated herein by reference in their entirety for all purposes.

[0181] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAb) that binds to an epitope or region of gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722, PGT-121.60, PGT-121.66, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137,

PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03. Additional broadly neutralizing antibodies that bind to gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and which can be used as the second antibody or antigen-binding fragment thereof are described, *e.g.*, in WO 2012/030904; WO 2014/063059; WO 2016/149698; WO 2017/106346; WO 2018/075564, WO 2018/125813 and WO 2018/237148, which are hereby incorporated herein by reference in their entireties for all purposes.

[0182] In some embodiments, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and another anti-HIV antibody (*e.g.*, GS-9722, PGT-121.60, PGT-121.66, PGT-121, PGT-122, PGT-123, PGT-124, PGT-133, or PGT-134) having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies from Tables 1 and 2 of US2017/0190763A1. These improved or optimized versions of PGT121 have enhanced drug-like-properties, reduced immunogenicity, enhanced ADCC, and suitable pharmacokinetic properties. Such antibodies were shown to bind to the HIV envelope glycoprotein expressed on the surface of virion or infected cells, and mediating both direct neutralization of the virus as well as potent NK, Monocyte and PBMC killing of these cells. This property allows the antibodies to treat HIV infections by neutralizing the virus, and also kill and eliminate latently HIV infected cells in infected individuals, potentially leading to a sterilizing cure for HIV.

[0183] In one embodiment, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of the antibody having the sequences below:

Heavy Chain (VH underlined)	<u>QMQLQESGPGLVKPSETLSLTCSVSGASISDSYWSWIRRRSPGKGLEWIGY</u> <u>VHKSGDTNYPNPSLKSRLVHLSLDTSKNQVSLSLTGVTAAADSGKYCARTLH</u> <u>GRRYIGIVAFNEWFTYFYMDVWGTGTQVTVSSASTKGPSVFPLAPSSKSTS</u> GGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVT VPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGP DVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAK GQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENN YKTTTPVLDSGDSFFLYSKLTVDKSRWQQGNVFCFVLSHESLYTQS LSLSPGK (SEQ ID NO: 443)
--------------------------------	---

Heavy CDR1 Kabat	DSYWS (SEQ ID NO: 444)
Heavy CDR2 Kabat	YVHKSGDTNYPNPSLKS (SEQ ID NO: 445)
Heavy CDR3 Kabat	TLHGRRYIGIVAFNEWFTYFYMDV (SEQ ID NO: 446)
Light Chain (VL underlined)	<u>SDISVAPGETARISCGEKSLSRAVQWYQHRAGQAPSLIIYNNQDRPSGIP</u> <u>ERFSGSPDSRPGTTATLTITSVEAGDEADYYCHIWDSRVPTKWVFGGGTTL</u> <u>TVLGQPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADSSP</u> VKAGVETTTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEK TVAPTECS (SEQ ID NO: 447)
Light CDR1 Kabat	GEKSLSRAVQ (SEQ ID NO: 448)
Light CDR2 Kabat	NNQDRPS (SEQ ID NO: 449)
Light CDR3 Kabat	HIWDSRVPTKWV (SEQ ID NO: 450)

[0184] In one embodiment, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of other additional anti-HIV antibodies such as those disclosed in US2017/0190763. In certain embodiments, the additional anti-HIV antibodies comprise an antibody comprising the VH (or heavy) and the VL (or light) chains provided below:

[0185] Heavy Chain (VH underlined):

QMQLQESGPGPLVKPSETLSLTCSVSGASISDSYWSWIRRSPPGKLEWIGYVHKSGDTNYPNPSLK
SRVHLSLDTSKNQVLSLTGVTAADSGKYCARTLHGRRYIGIVAFNEWFTYFYMDVWGTTGTQV
TVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS
GLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVF
LFPPKPKDITLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSV
LTVLHQDWLNGKEYKCKVSNKALPLPEEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV
KGFPYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSCSVLHEALH
SHYTQKSLSLSPGK (SEQ ID NO: 454)

[0186] Light Chain (VL underlined):

SDISVAPGETARISCGEKSLSRAVQWYQHRAGQAPSLIIYNNQDRPSGIPERFSGSPDY
RPGTTATLTITSVEAGDEADYYCHIWDSRVPTKWVFGGGTTLTVLGQPKAAPSVTLFPPS
SEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTTPSKQSNKYAASSYLSLT
PEQWKSHRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 455)

[0187] In one embodiment, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of another anti-HIV antibody, the heavy

chain of which has the amino acid sequence set forth in SEQ ID NO:40 and the light chain of which has the sequence provided below:

[0188] Light Chain (VL underlined):

SDISVAPGETARISCGEKSLSRAVQWYQHRAGQAPSLIIYNNQDRPSGIPERFSGSPDF
RP GTTATLTITSVEAGDEADYYCHIWDSRVPTKWVFGGGTTLTVLGQPKAAPSVTLFP
SEELQANKATLVCLISDFYPGAVTVAWKADSSPVKAGVETTTPSKQSNKYAASSYLSLT
PEQWKSHRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 456)

[0189] In one embodiment, the combination therapy includes an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of any of the antibodies disclosed herein and an antibody having the same CDRs, VH, VL, VH and VL, heavy, light, or heavy and light chains of the antibody described below:

Clone Designation	PGT121.42 hlgG1/hLambda
Heavy Chain (VH underlined)	<u>QMQLQESGPGLVKPSETLSLTCSVSGASISDSYWSWIRRRSPGKGLEWIGY</u> <u>VHKSGDTNYPNPSLKS RVHLSLDT SKNQVSLSLSSVTAADSGKYCARTLHG</u> <u>RRIYGIVAFNEWFTYFYMDVWGKGTVTVSSASTKGPSVFPLAPSSKSTSG</u> <u>GTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTV</u> <u>PSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPD</u> <u>VFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTK</u> <u>PREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPLPEEKTISKAKG</u> <u>QPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNY</u> <u>KTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCVSLHEALHSHYTQKSL</u> <u>SLSPGK</u> (SEQ ID NO: 451)
Heavy CDR1 Kabat	DSYWS (SEQ ID NO: 444)
Heavy CDR2 Kabat	YVHKSGDTNYPNPSLKS (SEQ ID NO: 445)
Heavy CDR3 Kabat	TLHGRRYIGIVAFNEWFTYFYMDV (SEQ ID NO: 446)
Light Chain (VL underlined)	<u>SDISVAPGETARISCGEKSLSRAVQWYQHRAGQAPSLIIYNNQDRPSGIP</u> <u>ERFSGSPDSPFGTTATLTITSVEAGDEADYYCHIWDSRVPTKWVFGGGTTL</u> <u>TVLGQPKAAPSVTLFPSSSEELQANKATLVCLISDFYPGAVTVAWKADSSP</u> <u>VKAGVETTTPSKQSNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEK</u> <u>TVAPTECS</u> (SEQ ID NO: 452)
Light CDR1 Kabat	GEKSLGSRAVQ (SEQ ID NO: 448)
Light CDR2 Kabat	NNQDRPS (SEQ ID NO: 449)
Light CDR3 Kabat	HIWDSRVPTKWV (SEQ ID NO: 450)

[0190] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of gp120 in the second variable loop (V2) and/or Env trimer apex and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PG9, PG16, PGC14, PGG14, PGT-142, PGT-143, PGT-144, PGT-145, CH01, CH59, PGDM1400, CAP256, CAP256-

VRC26.08, CAP256-VRC26.09, CAP256-VRC26.25, PCT64-24E and VRC38.01.

Additional broadly neutralizing antibodies that bind to gp120 in the second variable loop (V2) and/or Env trimer apex and which can be used as the second antibody or antigen-binding fragment thereof are described, *e.g.*, in WO 2010/107939; WO 2012/030904; WO 2018/075564 and WO 2018/125813, which are hereby incorporated herein by reference in their entireties for all purposes.

[0191] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of gp120 in the gp120/gp41 interface and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PGT-151, CAP248-2B, 35O22, 8ANC195, ACS202, VRC34 and VRC34.01. Additional broadly neutralizing antibodies that bind to gp120 in the gp120/gp41 interface and which can be used as the second antibody or antigen-binding fragment thereof are described, *e.g.*, in WO 2011/038290; WO 2012/030904 and WO2017/079479, which are hereby incorporated herein by reference in their entireties for all purposes.

[0192] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of the gp120 silent face and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC-PG05 and SF12. *See, e.g.*, Schoofs, *et al.*, “Broad and Potent Neutralizing Antibodies Recognize the Silent Face of the HIV Envelope,” *Immunity* (2019) May 14. pii: S1074-7613(19)30194-3 (PMID 31126879).

[0193] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of gp41 in the membrane proximal region (MPER). Additional broadly neutralizing antibodies that bind to gp41 in the MPER and which can be used as the second antibody or antigen-binding fragment thereof are described, *e.g.*, in WO 2011/034582; WO 2011/038290; WO 2011/046623 and WO 2013/070776, which are hereby incorporated herein by reference in their entireties for all purposes.

[0194] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of gp41 in the membrane proximal region (MPER) and competes with or comprises VH and VL regions from an antibody selected from the group consisting of 10E8, 10E8v4, 10E8-5R-100cF, 4E10, DH511.11P, 2F5, 7b2, and LN01.

[0195] In some embodiments, the antibodies or antigen-binding fragments thereof, described herein, are combined or co-administered with a second antibody or antigen-binding fragment thereof (*e.g.*, a second non-competing broadly neutralizing antibody (bNAbs)) that binds to an epitope or region of the gp41 fusion peptide and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC34 and ACS202.

[0196] Additional broadly neutralizing antibodies which can be used as a second therapeutic agent in a combination therapy are described, *e.g.*, in U.S. Patent Nos. 8,673,307; 9,493,549; 9,783,594; and WO 2012/154312; WO2012/158948; WO 2013/086533; WO 2013/142324; WO2014/063059; WO 2014/089152, WO 2015/048462; WO 2015/103549; WO 2015/117008; WO2016/014484; WO 2016/154003; WO 2016/196975; WO 2016/149710; WO2017/096221; WO 2017/133639; WO 2017/133640, which are hereby incorporated herein by reference in their entireties for all purposes. Additional examples include those described in Sajadi, et al., Cell. (2018) 173(7):1783-1795; Sajadi, et al., J Infect Dis. (2016) 213(1):156-64; Klein et al., Nature, 492(7427): 118-22 (2012), Horwitz et al., Proc Natl Acad Sci U S A, 110(41): 16538-43 (2013), Scheid, et al., Science, 333 : 1633-1637 (2011), Scheid, et al., Nature, 458:636-640 (2009), Eroshkin et al, Nucleic Acids Res., 42 (Database issue):D1133-9 (2014), Mascola et al., Immunol Rev., 254(1):225-44 (2013), such as 2F5, 4E10, M66.6, CAP206-CH12, 10E8, 10E8v4, 10E8-5R-100cF, DH511.11P, 7b2, and LN01 (all of which bind the MPER of gp41); PG9, PG16, CH01-04 (all of which bind V1V2-glycan), 2G12 (which binds to outer domain glycan), which are hereby incorporated herein by reference in their entireties for all purposes.

[0197] Exemplary VH and VL amino acid sequences of an anti-gp120 antibody of this disclosure that are used in the combination therapy include the sequences set forth in SEQ ID NOs: 182 and 275, respectively; SEQ ID NOs: 182 and 278, respectively; SEQ ID NOs: 182 and 279, respectively; SEQ ID NOs: 182 and 280, respectively; SEQ ID

NOs: 182 and 281, respectively; SEQ ID NOs: 182 and 282, respectively; SEQ ID NOs: 182 and 292, respectively; SEQ ID NOs: 182 and 304, respectively; SEQ ID NOs: 182 and 307, respectively; SEQ ID NOs: 182 and 309, respectively; SEQ ID NOs: 182 and 310, respectively; SEQ ID NOs: 220 and 310, respectively; SEQ ID NOs: 477 and 223, respectively; SEQ ID NOs: 477 and 278, respectively; SEQ ID NOs: 477 and 292, respectively; and SEQ ID NOs: 220 and 311, respectively. In certain embodiments, the VH and VL amino acid sequences of an anti-gp120 antibody used in the combination therapy are the sequences set forth in SEQ ID NOs: 477 and 278, respectively. In certain embodiments, the arm of the bispecific antibody that binds to gp120 comprises an amino acid sequence of a heavy chain of an anti-gp120 antibody disclosed herein. In certain embodiments, the arm of the bispecific antibody that binds to gp120 comprises an amino acid sequence of a light chain of an anti-gp120 antibody disclosed herein. Exemplary heavy chain and light chain sequences of an anti-gp120 antibody of this disclosure that are used in the combination therapy include the sequences set forth in SEQ ID NOs: 2 and 49, respectively; SEQ ID NOs: 2 and 100, respectively; SEQ ID NOs: 42 and 101, respectively; SEQ ID NOs: 2 and 103, respectively; SEQ ID NOs: 2 and 104, respectively; SEQ ID NOs: 2 and 105, respectively; SEQ ID NOs: 2 and 106, respectively; SEQ ID NOs: 2 and 107, respectively; SEQ ID NOs: 2 and 117, respectively; SEQ ID NOs: 2 and 129, respectively; SEQ ID NOs: 2 and 132, respectively; SEQ ID NOs: 2 and 134, respectively; SEQ ID NOs: 2 and 569, respectively; SEQ ID NOs: 42 and 135, respectively; SEQ ID NOs: 529 and 49, respectively; SEQ ID NOs: 529 and 103, respectively; SEQ ID NOs: 529 and 117, respectively; and SEQ ID NOs: 42 and 136, respectively. In certain embodiments, the heavy chain and light chain sequences of an anti-gp120 antibody used in the combination therapy are the sequences set forth in SEQ ID NOs: 529 and 103, respectively.

[0198] In one embodiment, pharmaceutical compositions comprising an antibody disclosed herein, or a pharmaceutical composition thereof, in combination with one or more (*e.g.*, one, two, three, one or two, or one to three) additional therapeutic agents, and a pharmaceutically acceptable carrier, diluent, or excipient are provided.

[0199] In certain embodiments, the present disclosure provides a method for treating an HIV infection, comprising administering to a patient in need thereof a therapeutically effective amount of an antibody disclosed herein, or a pharmaceutical composition thereof, in combination with a therapeutically effective amount of one or more additional therapeutic agents which are suitable for treating an HIV infection.

[0200] In certain embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with one, two, three, four, or more additional therapeutic agents. In certain embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with two additional therapeutic agents. In other embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with three additional therapeutic agents. In further embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with four additional therapeutic agents. The one, two, three, four, or more additional therapeutic agents can be different therapeutic agents selected from the same class of therapeutic agents, and/or they can be selected from different classes of therapeutic agents.

[0201] In certain embodiments, an antibody disclosed herein is administered with one or more additional therapeutic agents. Co-administration of an antibody disclosed herein with one or more additional therapeutic agents generally refers to simultaneous or sequential administration of a compound disclosed herein and one or more additional therapeutic agents, such that therapeutically effective amounts of the antibody disclosed herein and the one or more additional therapeutic agents are both present in the body of the patient. When administered sequentially, the combination may be administered in two or more administrations.

[0202] Co-administration includes administration of unit dosages of the antibodies disclosed herein before or after administration of unit dosages of one or more additional therapeutic agents. For example, the antibody disclosed herein may be administered within seconds, minutes, or hours of the administration of the one or more additional therapeutic agents. In some embodiments, a unit dose of an antibody disclosed herein is administered first, followed within seconds or minutes by administration of a unit dose of one or more additional therapeutic agents. Alternatively, a unit dose of one or more additional therapeutic agents is administered first, followed by administration of a unit dose of an antibody disclosed herein within seconds or minutes. In other embodiments, a unit dose of an antibody disclosed herein is administered first, followed, after a period of hours (*e.g.*, 1-12 hours), by administration of a unit dose of one or more additional therapeutic agents. In yet other embodiments, a unit dose of one or more additional therapeutic agents is administered first, followed, after a period of hours (*e.g.*, 1-12 hours), by administration of a unit dose of an antibody disclosed herein.

[0203] In certain embodiments, an antibody disclosed herein is combined with one or more additional therapeutic agents in a unitary dosage form for simultaneous administration to a patient, for example as a solid dosage form for oral administration.

[0204] In certain embodiments, an antibody of this disclosure is formulated as a liquid, which may optionally contain an additional therapeutic agent(s) useful for treating HIV. In certain embodiments, the liquid can contain another active ingredient for treating HIV, such as another anti-HIV antibody or antigen-binding fragment thereof, a HIV protease inhibitor, a HIV non-nucleoside or non-nucleotide inhibitor of reverse transcriptase, a HIV nucleoside or nucleotide inhibitor of reverse transcriptase, a HIV integrase inhibitor, a HIV non-catalytic site (or allosteric) integrase inhibitor, pharmacokinetic enhancer, and combinations thereof.

[0205] In some embodiments, the additional therapeutic agent is a latency reversing agent (LRA), *e.g.*, an agonist of a toll-like receptor (TLR), *e.g.*, an agonist of TLR1 (NCBI Gene ID: 7096), TLR2 (NCBI Gene ID: 7097), TLR3 (NCBI Gene ID: 7098), TLR4 (NCBI Gene ID: 7099), TLR5 (NCBI Gene ID: 7100), TLR6 (NCBI Gene ID: 10333), TLR7 (NCBI Gene ID: 51284), TLR8 (NCBI Gene ID: 51311), TLR9 (NCBI Gene ID: 54106), and/or TLR10 (NCBI Gene ID: 81793). In some embodiments, the LRA is a TLR7 agonist. In other embodiments, the additional therapeutic agent is a latency reversing agent (LRA), *e.g.*, a TLR8 agonist. Examples of TLR agonists include but are not limited to Vesatolimod. Additional examples include but are not limited to the compounds described in U.S. Patent No. 8,367,670 and the compounds described in U.S. Patent Application Publication No. 2016/0289229. In one embodiment, the antibody of the present invention may be combined with TLR7 agonist such as Vesatolimod. In another embodiment, the antibody of the present invention may be combined with TLR8 agonist, *e.g.*, GS-9688. In one embodiment, the additional therapeutic agent is a TLR modulator. TLR modulators may include modulators of TLR1, TLR2, TLR3, TLR4, TLR5, TLR6, TLR7, TLR8, TLR9, TLR10, TLR11, TLR12, and TLR13. Examples of TLR3 modulators include rintatolimod, poly-ICLC, RIBOXXON®, Apoxsim, RIBOXXIM®, IPH-33, MCT-465, MCT-475, and ND-1.1. Examples of TLR7 modulators include GS-9620, GSK-2245035, imiquimod, resiquimod, DSR-6434, DSP-3025, IMO-4200, MCT-465, MEDI-9197, 3M-051, SB-9922, 3M-052, Limtop, TMX-30X, TMX-202, RG-7863, RG-7795, and the compounds disclosed in US20100143301 (Gilead Sciences), US20110098248 (Gilead Sciences), and US20090047249 (Gilead Sciences). Examples of TLR8 modulators include GS-9688,

motolimod, resiquimod, 3M-051, 3M-052, MCT-465, IMO-4200, VTX-763, VTX-1463, and the compounds disclosed in US20140045849 (Janssen), US20140073642 (Janssen), WO2014/056953 (Janssen), WO2014/076221 (Janssen), WO2014/128189 (Janssen), US20140350031 (Janssen), WO2014/023813 (Janssen), US20080234251 (Array Biopharma), US20080306050 (Array Biopharma), US20100029585 (Ventirx Pharma), US20110092485 (Ventirx Pharma), US20110118235 (Ventirx Pharma), US20120082658 (Ventirx Pharma), US20120219615 (Ventirx Pharma), US20140066432 (Ventirx Pharma), US20140088085 (Ventirx Pharma), US20140275167 (Novira Therapeutics), and US20130251673 (Novira Therapeutics). Examples of TLR9 modulators include BB-001, BB-006, CYT-003, IMO-2055, IMO-2125, IMO-3100, IMO-8400, IR-103, IMO-9200, agatolimod, DIMS-9054, DV-1079, DV-1179, AZD-1419, leftolimod (MGN-1703), litemimod, and CYT-003-QbG10.

[0206] In some embodiments, the additional therapeutic agent is an agonist of DExD/H-box helicase 58 (DDX58; *a.k.a.*, RIG-I, RIG1, RIGI, RLR-1, SGMRT2; NCBI Gene ID: 23586). An illustrative RIG-I agonist is KIN1148, described by Hemann, *et al.*, *J Immunol* May 1, 2016, 196 (1 Supplement) 76.1. Additional RIG-I agonists are described, *e.g.*, in Elion, *et al.*, *Cancer Res.* (2018) 78(21):6183-6195; and Liu, *et al.*, *J Virol.* (2016) 90(20):9406-19. RIG-I agonists are commercially available, *e.g.*, from Invivogen (invivogen.com).

[0207] In certain embodiments, such formulations are suitable for once daily dosing.

[0208] In some embodiments, the additional therapeutic agent may be an anti-HIV agent. In some instances, the additional therapeutic agent can be HIV protease inhibitors, HIV non-nucleoside or non-nucleotide inhibitors of reverse transcriptase, HIV nucleoside or nucleotide inhibitors of reverse transcriptase, HIV integrase inhibitors, HIV non-catalytic site (or allosteric) integrase inhibitors, HIV entry inhibitors, HIV maturation inhibitors, HIV capsid inhibitors, HIV Tat or Rev inhibitors, immunomodulators (*e.g.*, immunostimulators), immunotherapeutic agents, antibody-drug conjugates, gene modifiers, gene editors (such as CRISPR/Cas9, zinc finger nucleases, homing nucleases, synthetic nucleases, TALENs), cell therapies (such as chimeric antigen receptor T-cell, CAR-T, and engineered T-cell receptors, TCR-T, autologous T-cell therapies), latency reversing agents, compounds that target the HIV capsid, immune-based therapies, phosphatidylinositol 3-kinase (PI3K) inhibitors, HIV antibodies, bispecific antibodies and “antibody-like” therapeutic proteins, HIV p17 matrix protein inhibitors, IL-13 antagonists, peptidyl-prolyl cis-trans isomerase A modulators, protein disulfide

isomerase inhibitors, complement C5a receptor antagonists, DNA methyltransferase inhibitor, HIV vif gene modulators, Vif dimerization antagonists, HIV-1 viral infectivity factor inhibitors, TAT protein inhibitors, HIV-1 Nef modulators, Hck tyrosine kinase modulators, mixed lineage kinase-3 (MLK-3) inhibitors, HIV-1 splicing inhibitors, Rev protein inhibitors, integrin antagonists, nucleoprotein inhibitors, splicing factor modulators, COMM domain containing protein 1 modulators, HIV ribonuclease H inhibitors, retrocyclin modulators, CDK-9 inhibitors, dendritic ICAM-3 grabbing nonintegrin 1 inhibitors, HIV GAG protein inhibitors, HIV POL protein inhibitors, Complement Factor H modulators, ubiquitin ligase inhibitors, deoxycytidine kinase inhibitors, cyclin dependent kinase inhibitors, proprotein convertase PC9 stimulators, ATP dependent RNA helicase DDX3X inhibitors, reverse transcriptase priming complex inhibitors, G6PD and NADH-oxidase inhibitors, pharmacokinetic enhancers, HIV gene therapy, HIV vaccines, and combinations thereof.

[0209] In some embodiments, the additional therapeutic agent is selected from the group consisting of combination drugs for HIV, other drugs for treating HIV, HIV protease inhibitors, HIV reverse transcriptase inhibitors, HIV integrase inhibitors, HIV non-catalytic site (or allosteric) integrase inhibitors, HIV entry (fusion) inhibitors, HIV maturation inhibitors, latency reversing agents, capsid inhibitors, immune-based therapies, PI3K inhibitors, HIV antibodies, and bispecific antibodies, and “antibody-like” therapeutic proteins, and combinations thereof.

Combination Drugs

[0210] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV combination drug. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 477 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-

binding fragment thereof comprises a VH sequence set forth in SEQ ID NO: 477 and a VL sequence set forth in SEQ ID NO: 278. In certain embodiments, the antibody or antigen-binding fragment thereof comprises VH CDRs and VL CDRs having the sequences set forth in: SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and comprises a heavy chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 529 and a light chain that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, such as 100%, identical to an amino acid sequence set forth in SEQ ID NO: 103. In certain embodiments, the antibody or antigen-binding fragment thereof comprises a heavy chain sequence set forth in SEQ ID NO: 529 and a light chain sequence set forth in SEQ ID NO: 103. Examples of combination drugs that can be employed with an antibody of this disclosure include ATRIPLA® (efavirenz, tenofovir disoproxil fumarate, and emtricitabine); COMPLERA® (EVIPLERA®; rilpivirine, tenofovir disoproxil fumarate, and emtricitabine); STRIBILD® (elvitegravir, cobicistat, tenofovir disoproxil fumarate, and emtricitabine); TRUVADA® (tenofovir disoproxil fumarate and emtricitabine; TDF+FTC); DESCOVY® (tenofovir alafenamide and emtricitabine); ODEFSEY® (tenofovir alafenamide, emtricitabine, and rilpivirine); GENVOYA® (tenofovir alafenamide, emtricitabine, cobicistat, and elvitegravir); darunavir, tenofovir alafenamide hemifumarate, emtricitabine, and cobicistat; efavirenz, lamivudine, and tenofovir disoproxil fumarate; lamivudine and tenofovir disoproxil fumarate; tenofovir and lamivudine; tenofovir alafenamide and emtricitabine; tenofovir alafenamide hemifumarate and emtricitabine; tenofovir alafenamide hemifumarate, emtricitabine, and rilpivirine; tenofovir alafenamide hemifumarate, emtricitabine, cobicistat, and elvitegravir; COMBIVIR® (zidovudine and lamivudine; AZT+3TC); EPZICOM® (LIVEXA®; abacavir sulfate and lamivudine; ABC+3TC); KALETRA® (ALUVIA®; lopinavir and ritonavir); TRIUMEQ® (dolutegravir, abacavir, and lamivudine); TRIZIVIR® (abacavir sulfate, zidovudine, and lamivudine; ABC+AZT+3TC); atazanavir and cobicistat; atazanavir sulfate and cobicistat; atazanavir sulfate and ritonavir; darunavir and cobicistat; dolutegravir and rilpivirine; dolutegravir and rilpivirine hydrochloride; dolutegravir, abacavir sulfate, and lamivudine; lamivudine, nevirapine, and zidovudine; raltegravir and lamivudine; doravirine, lamivudine, and tenofovir disoproxil fumarate; doravirine, lamivudine, and tenofovir disoproxil; dolutegravir + lamivudine, lamivudine + abacavir + zidovudine, lamivudine + abacavir,

lamivudine + tenofovir disoproxil fumarate, lamivudine + zidovudine + nevirapine, lopinavir + ritonavir, lopinavir + ritonavir + abacavir + lamivudine, lopinavir + ritonavir + zidovudine + lamivudine, tenofovir + lamivudine, and tenofovir disoproxil fumarate + emtricitabine + rilpivirine hydrochloride, lopinavir, ritonavir, zidovudine and lamivudine; Vacc-4x and romidepsin; and APH-0812.

Other HIV Drugs

[0211] Examples of other drugs for treating HIV that can be combined with an antibody of this disclosure include acemannan, alisporivir, BanLec, deferiprone, Gamimmune, metenkefalin, naltrexone, Prolastin, REP 9, RPI-MN, VSSP, H1viral, SB-728-T, 1,5-dicaffeoylquinic acid, rHIV7-shl-TAR-CCR5RZ, AAV-eCD4-Ig gene therapy, MazF gene therapy, BlockAide, ABX-464, AG-1105, APH-0812, BIT-225, CYT-107, HGTV-43, HPH-116, HS-10234, IMO-3100, IND-02, MK-1376, MK-2048, MK-4250, MK-8507, MK-8591, NOV-205, PA-1050040 (PA-040), PGN-007, SCY-635, SB-9200, SCB-719, TR-452, TEV-90110, TEV-90112, TEV-90111, TEV-90113, RN-18, Immuglo, and VIR-576.

HIV Protease Inhibitors

[0212] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV protease inhibitor. Examples of HIV protease inhibitors that can be combined with an antibody of this disclosure include amprenavir, atazanavir, brecanavir, darunavir, fosamprenavir, fosamprenavir calcium, indinavir, indinavir sulfate, lopinavir, nelfinavir, nelfinavir mesylate, ritonavir, saquinavir, saquinavir mesylate, tipranavir, DG-17, TMB-657 (PPL-100), T-169, BL-008, MK-8122, TMB-607, and TMC-310911.

HIV Reverse Transcriptase Inhibitors

[0213] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a non-nucleoside or non-nucleotide inhibitor. Examples of HIV non-nucleoside or non-nucleotide inhibitors of reverse transcriptase that can be combined with an antibody of this disclosure include dapivirine, delavirdine, delavirdine mesylate, doravirine, efavirenz, etravirine, lentinan, nevirapine, rilpivirine, ACC-007, AIC-292, KM-023, PC-1005, and elsulfavirine (VM-1500.).

[0214] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV nucleoside or nucleotide inhibitor. Examples of HIV nucleoside or nucleotide inhibitors of reverse transcriptase that can be combined with an

antibody of this disclosure include adefovir, adefovir dipivoxil, azvudine, emtricitabine, tenofovir, tenofovir alafenamide, tenofovir alafenamide fumarate, tenofovir alafenamide hemifumarate, tenofovir disoproxil, tenofovir disoproxil fumarate, tenofovir disoproxil hemifumarate, VIDEX® and VIDEX EC® (didanosine, ddl), abacavir, abacavir sulfate, alovudine, apricitabine, censavudine, didanosine, elvucitabine, festinavir, fosalvudine tidoxil, CMX-157, dapivirine, doravirine, etravirine, OCR-5753, tenofovir disoproxil orotate, fozivudine tidoxil, lamivudine, phosphazid, stavudine, zalcitabine, zidovudine, rovafovir etalafenamide (GS-9131), GS-9148, MK-8504, MK-8591, MK-858, VM-2500 and KP-1461.

HIV Integrase Inhibitors

[0215] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV integrase inhibitor. Examples of HIV integrase inhibitors that can be combined with an antibody of this disclosure include elvitegravir, curcumin, derivatives of curcumin, chicoric acid, derivatives of chicoric acid, 3,5-dicaffeoylquinic acid, derivatives of 3,5-dicaffeoylquinic acid, aurintricarboxylic acid, derivatives of aurintricarboxylic acid, caffeic acid phenethyl ester, derivatives of caffeic acid phenethyl ester, tyrphostin, derivatives of tyrphostin, quercetin, derivatives of quercetin, raltegravir, dolutegravir, JTK-351, bictegravir, AVX-15567, cabotegravir (long-acting injectable), diketo quinolin-4-1 derivatives, integrase-LEDGF inhibitor, ledgins, M-522, M-532, NSC-310217, NSC-371056, NSC-48240, NSC-642710, NSC-699171, NSC-699172, NSC-699173, NSC-699174, stilbenedisulfonic acid, T-169, VM-3500 and cabotegravir.

[0216] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a HIV non-catalytic site, or allosteric, integrase inhibitor (NCINI). Examples of HIV non-catalytic site, or allosteric, integrase inhibitors (NCINI) that can be combined with an antibody of this disclosure include CX-05045, CX-05168, and CX-14442.

HIV Entry Inhibitors

[0217] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV entry inhibitor. Examples of HIV entry (fusion) inhibitors that can be combined with an antibody of this disclosure include cenicriviroc, CCR5 inhibitors, gp41 inhibitors, CD4 attachment inhibitors, gp120 inhibitors, and CXCR4 inhibitors.

[0218] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a CCR5 inhibitor. Examples of CCR5 inhibitors that can be combined with an antibody of this disclosure include aplaviroc, vicriviroc, maraviroc, cenicriviroc, leronlimab (PRO-140), adaptavir (RAP-101), nifeviroc (TD-0232), anti-GP120/CD4 or CCR5 bispecific antibodies, B-07, MB-66, polypeptide C25P, TD-0680, and vMIP (Haimipu).

[0219] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a gp41 inhibitor. Examples of gp41 inhibitors that can be combined with an antibody of this disclosure include albuvirtide, enfuvirtide, BMS-986197, enfuvirtide biobetter, enfuvirtide biosimilar, HIV-1 fusion inhibitors (P26-Bapc), ITV-1, ITV-2, ITV-3, ITV-4, PIE-12 trimer and sifuvirtide.

[0220] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a CD4 attachment inhibitor. Examples of CD4 attachment inhibitors that can be combined with an antibody of this disclosure include ibalizumab and CADA analogs.

[0221] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a gp120 inhibitor. Examples of gp120 inhibitors that can be combined with an antibody of this disclosure include Radha-108 (receptol) 3B3-PE38, BanLec, bentonite-based nanomedicine, fostemsavir tromethamine, IQP-0831, and BMS-663068.

[0222] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a CXCR4 inhibitor. Examples of CXCR4 inhibitors that can be combined with an antibody of this disclosure include plerixafor, ALT-1188, N15 peptide, and vMIP (Haimipu).

[0223] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a HIV maturation inhibitor. Examples of HIV maturation inhibitors that can be combined with an antibody of this disclosure include BMS-955176, GSK-3640254 and GSK-2838232.

Latency Reversing Agents

[0224] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a latency reversing agent (LRA). Examples of latency reversing agents that can be combined with an antibody of this disclosure include toll-like receptor (TLR) agonists (including TLR7 agonists, *e.g.*, GS-9620 and TLR8

agonists, *e.g.*, GS-9688), histone deacetylase (HDAC) inhibitors, proteasome inhibitors such as velcade, protein kinase C (PKC) activators, Smyd2 inhibitors, BET-bromodomain 4 (BRD4) inhibitors, ionomycin, IAP antagonists (inhibitor of apoptosis proteins, such as APG-1387, LBW-242), SMAC mimetics (including TL32711, LCL161, GDC-0917, HGS1029, AT-406), PMA, SAHA (suberanilohydroxamic acid, or suberoyl, anilide, and hydroxamic acid), NIZ-985, IL-15 modulating antibodies (including IL-15, IL-15 fusion proteins and IL-15 receptor agonists, *e.g.*, ALT-803), JQ1, disulfiram, amphotericin B, and ubiquitin inhibitors such as largazole analogs, APH-0812, and GSK-343. Examples of HDAC inhibitors include romidepsin, vorinostat, and panobinostat. Examples of PKC activators include indolactam, prostratin, ingenol B, and DAG-lactones.

Toll-Like Receptor (TLR) Agonists

[0225] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with an agonist of a toll-like receptor (TLR), *e.g.*, an agonist of TLR1 (NCBI Gene ID: 7096), TLR2 (NCBI Gene ID: 7097), TLR3 (NCBI Gene ID: 7098), TLR4 (NCBI Gene ID: 7099), TLR5 (NCBI Gene ID: 7100), TLR6 (NCBI Gene ID: 10333), TLR7 (NCBI Gene ID: 51284), TLR8 (NCBI Gene ID: 51311), TLR9 (NCBI Gene ID: 54106), and/or TLR10 (NCBI Gene ID: 81793). Example TLR7 agonists that can be co-administered include without limitation AL-034, DSP-0509, GS-9620 (vesatolimod), LHC-165, TMX-101 (imiquimod), GSK-2245035, resiquimod, DSR-6434, DSP-3025, IMO-4200, MCT-465, MEDI-9197, 3M-051, SB-9922, 3M-052, Limtop, TMX-30X, TMX-202, RG-7863, RG-7854, RG-7795, and the compounds disclosed in US20100143301 (Gilead Sciences), US20110098248 (Gilead Sciences), and US20090047249 (Gilead Sciences), US20140045849 (Janssen), US20140073642 (Janssen), WO2014/056953 (Janssen), WO2014/076221 (Janssen), WO2014/128189 (Janssen), US20140350031 (Janssen), WO2014/023813 (Janssen), US20080234251 (Array Biopharma), US20080306050 (Array Biopharma), US20100029585 (Ventirx Pharma), US20110092485 (Ventirx Pharma), US20110118235 (Ventirx Pharma), US20120082658 (Ventirx Pharma), US20120219615 (Ventirx Pharma), US20140066432 (Ventirx Pharma), US20140088085 (Ventirx Pharma), US20140275167 (Novira Therapeutics), and US20130251673 (Novira Therapeutics). An TLR7/TLR8 agonist that can be co-administered is NKTR-262, telratolimod and BDB-001. Example TLR8 agonists that can be co-administered include without limitation E-6887, IMO-4200, IMO-8400, IMO-9200, MCT-465, MEDI-9197,

motolimod, resiquimod, GS-9688, VTX-1463, VTX-763, 3M-051, 3M-052, and the compounds disclosed in US20140045849 (Janssen), US20140073642 (Janssen), WO2014/056953 (Janssen), WO2014/076221 (Janssen), WO2014/128189 (Janssen), US20140350031 (Janssen), WO2014/023813 (Janssen), US20080234251 (Array Biopharma), US20080306050 (Array Biopharma), US20100029585 (Ventirx Pharma), US20110092485 (Ventirx Pharma), US20110118235 (Ventirx Pharma), US20120082658 (Ventirx Pharma), US20120219615 (Ventirx Pharma), US20140066432 (Ventirx Pharma), US20140088085 (Ventirx Pharma), US20140275167 (Novira Therapeutics), and US20130251673 (Novira Therapeutics). Example TLR9 agonists that can be co-administered include without limitation AST-008, cobitolimod, CMP-001, IMO-2055, IMO-2125, litenimod, MGN-1601, BB-001, BB-006, IMO-3100, IMO-8400, IR-103, IMO-9200, agatolimod, DIMS-9054, DV-1079, DV-1179, AZD-1419, lefitolimod (MGN-1703), CYT-003, CYT-003-QbG10, tilsitolimod and PUL-042. Examples of TLR3 agonist include rintatolimod, poly-ICLC, RIBOXXON®, Apoxxim, RIBOXXIM®, IPH-33, MCT-465, MCT-475, and ND-1.1. Examples of TLR4 agonist include G-100, and GSK-1795091.

Histone Deacetylase (HDAC) Inhibitors

[0226] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with an inhibitor of a histone deacetylase, *e.g.*, histone deacetylase 9 (HDAC9, HD7, HD7b, HD9, HDAC, HDAC7, HDAC7B, HDAC9B, HDAC9FL, HDRP, MITR; Gene ID: 9734). Examples of HDAC inhibitors include without limitation, abexinostat, ACY-241, AR-42, BEBT-908, belinostat, CKD-581, CS-055 (HBI-8000), CUDC-907 (fimepinostat), entinostat, givinostat, mocetinostat, panobinostat, pracinostat, quisinostat (JNJ-26481585), resminostat, ricolinostat, romidepsin, SHP-141, valproic acid (VAL-001), vorinostat, tinostamustine, remetinostat, entinostat.

Capsid Inhibitors

[0227] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a capsid inhibitor. Examples of capsid inhibitors that can be combined with an antibody of this disclosure include capsid polymerization inhibitors or capsid disrupting compounds, HIV nucleocapsid p7 (NCp7) inhibitors such as azodicarbonamide, HIV p24 capsid protein inhibitors, GS-6207, AVI-621, AVI-101, AVI-201, AVI-301, and AVI-CAN1-15 series.

Immune-based Therapies

[0228] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an immune-based therapy. Examples of immune-based therapies that can be combined with an antibody of this disclosure include toll-like receptors (TLR) modulators (*e.g.*, agonists) such as TLR1, TLR 2, TLR 3, TLR 4, TLR 5, TLR 6, TLR 7, TLR 8, TLR 9, TLR 10, TLR 11, TLR 12, and/or TLR 13 agonists; programmed cell death protein 1 (PD-1) modulators; programmed death-ligand 1 (PD-L1) modulators; IL-15 agonists (*e.g.*, ALT-803); DermaVir; interleukin-7; plaquenil (hydroxychloroquine); proleukin (aldesleukin, IL-2); interferon alfa; interferon alfa-2b; interferon alfa-n3; pegylated interferon alfa; interferon gamma; hydroxyurea; mycophenolate mofetil (MPA) and its ester derivative mycophenolate mofetil (MMF); ribavirin; rintatolimod, polymer polyethyleneimine (PEI); gepon; IL-12; WF-10; VGV-1; MOR-22; BMS-936559; CYT-107, interleukin-15/Fc fusion protein, AM-0015, ALT-803, NIZ-985, NKTR-255, normferon, peginterferon alfa-2a, peginterferon alfa-2b, recombinant interleukin-15, RPI-MN, GS-9620, GS-9688, STING modulators, RIG-I modulators, NOD2 modulators, SB-9200, and IR-103.

[0229] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a TLR agonist. Examples of TLR agonists include without limitation: vesatolimod (GS-9620), lefitolimod, tilsetolimod, rintatolimod, DSP-0509, AL-034, G-100, cobitolimod, AST-008, motolimod, GSK-1795091, GSK-2245035, VTX-1463, GS-9688, LHC-165, BDB-001, RG-7854, and telratolimod.

Immune Checkpoint Receptor Protein Modulators

[0230] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with one or more blockers or inhibitors of inhibitory immune checkpoint proteins or receptors and/or with one or more stimulators, activators or agonists of one or more stimulatory immune checkpoint proteins or receptors. Blockade or inhibition of inhibitory immune checkpoints can positively regulate T-cell or NK cell activation and prevent immune escape of infected cells. Activation or stimulation of stimulatory immune check points can augment the effect of immune checkpoint inhibitors in infective therapeutics. In various embodiments, the immune checkpoint proteins or receptors regulate T cell responses (*e.g.*, reviewed in Xu, et al., J Exp Clin Cancer Res. (2018) 37:110). In various embodiments, the immune checkpoint proteins or receptors regulate NK cell responses (*e.g.*, reviewed in Davis, et al., Semin

Immunol. (2017) 31:64–75 and Chiossone, et al., Nat Rev Immunol. (2018) 18(11):671-688).

[0231] Examples of immune checkpoint proteins or receptors include without limitation CD27, CD70; CD40, CD40LG; CD47, CD48 (SLAMF2), transmembrane and immunoglobulin domain containing 2 (TMIGD2, CD28H), CD84 (LY9B, SLAMF5), CD96, CD160, MS4A1 (CD20), CD244 (SLAMF4); CD276 (B7H3); V-set domain containing T cell activation inhibitor 1 (VTCN1, B7H4); V-set immunoregulatory receptor (VSIR, B7H5, VISTA); immunoglobulin superfamily member 11 (IGSF11, VSIG3); natural killer cell cytotoxicity receptor 3 ligand 1 (NCR3LG1, B7H6); HERV-H LTR-associating 2 (HHLA2, B7H7); inducible T cell co-stimulator (ICOS, CD278); inducible T cell costimulator ligand (ICOSLG, B7H2); TNF receptor superfamily member 4 (TNFRSF4, OX40); TNF superfamily member 4 (TNFSF4, OX40L); TNFRSF8 (CD30), TNFSF8 (CD30L); TNFRSF10A (CD261, DR4, TRAILR1), TNFRSF9 (CD137), TNFSF9 (CD137L); TNFRSF10B (CD262, DR5, TRAILR2), TNFRSF10 (TRAIL); TNFRSF14 (HVEM, CD270), TNFSF14 (HVEM); CD272 (B and T lymphocyte associated (BTLA)); TNFRSF17 (BCMA, CD269), TNFSF13B (BAFF); TNFRSF18 (GITR), TNFSF18 (GITRL); MHC class I polypeptide-related sequence A (MICA); MHC class I polypeptide-related sequence B (MICB); CD274 (CD274, PDL1, PD-L1); programmed cell death 1 (PDCD1, PD1, PD-1); cytotoxic T-lymphocyte associated protein 4 (CTLA4, CD152); CD80 (B7-1), CD28; nectin cell adhesion molecule 2 (NECTIN2, CD112); CD226 (DNAM-1); Poliovirus receptor (PVR) cell adhesion molecule (PVR, CD155); PVR related immunoglobulin domain containing (PVRIG, CD112R); T cell immunoreceptor with Ig and ITIM domains (TIGIT); T cell immunoglobulin and mucin domain containing 4 (TIMD4; TIM4); hepatitis A virus cellular receptor 2 (HAVCR2, TIMD3, TIM3); galectin 9 (LGALS9); lymphocyte activating 3 (LAG3, CD223); signaling lymphocytic activation molecule family member 1 (SLAMF1, SLAM, CD150); lymphocyte antigen 9 (LY9, CD229, SLAMF3); SLAM family member 6 (SLAMF6, CD352); SLAM family member 7 (SLAMF7, CD319); UL16 binding protein 1 (ULBP1); UL16 binding protein 2 (ULBP2); UL16 binding protein 3 (ULBP3); retinoic acid early transcript 1E (RAET1E; ULBP4); retinoic acid early transcript 1G (RAET1G; ULBP5); retinoic acid early transcript 1L (RAET1L; ULBP6); lymphocyte activating 3 (CD223); killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR, CD158E1); killer cell lectin like receptor C1 (KLRC1, NKG2A, CD159A); killer cell

lectin like receptor K1 (KLRK1, NKG2D, CD314); killer cell lectin like receptor C2 (KLRC2, CD159c, NKG2C); killer cell lectin like receptor C3 (KLRC3, NKG2E); killer cell lectin like receptor C4 (KLRC4, NKG2F); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3); killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1); killer cell lectin like receptor D1 (KLRD1); and SLAM family member 7 (SLAMF7).

[0232] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with one or more blockers or inhibitors of one or more T-cell inhibitory immune checkpoint proteins or receptors. Illustrative T-cell inhibitory immune checkpoint proteins or receptors include without limitation CD274 (CD274, PDL1, PD-L1); programmed cell death 1 ligand 2 (PDCD1LG2, PD-L2, CD273); programmed cell death 1 (PDCD1, PD1, PD-1); cytotoxic T-lymphocyte associated protein 4 (CTLA4, CD152); CD276 (B7H3); V-set domain containing T cell activation inhibitor 1 (VTCN1, B7H4); V-set immunoregulatory receptor (VSIR, B7H5, VISTA); immunoglobulin superfamily member 11 (IGSF11, VSIG3); TNFRSF14 (HVEM, CD270), TNFSF14 (HVEM); CD272 (B and T lymphocyte associated (BTLA)); PVR related immunoglobulin domain containing (PVRIG, CD112R); T cell immunoreceptor with Ig and ITIM domains (TIGIT); lymphocyte activating 3 (LAG3, CD223); hepatitis A virus cellular receptor 2 (HAVCR2, TIMD3, TIM3); galectin 9 (LGALS9); killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR, CD158E1); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3); and killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1). In various embodiments, the FLT3L-Fc fusion proteins, homodimers, heterodimers, polynucleotides, vectors, LNPs and/or pharmaceutical compositions, as described herein, are combined with one or more agonist or activators of one or more T-cell stimulatory immune checkpoint proteins or receptors. Illustrative T-cell stimulatory immune checkpoint proteins or receptors include without limitation CD27, CD70, CD40, CD40LG; inducible T cell costimulator (ICOS, CD278); inducible T cell costimulator

ligand (ICOSLG, B7H2); TNF receptor superfamily member 4 (TNFRSF4, OX40); TNF superfamily member 4 (TNFSF4, OX40L); TNFRSF9 (CD137), TNFSF9 (CD137L); TNFRSF18 (GITR), TNFSF18 (GITRL); CD80 (B7-1), CD28; nectin cell adhesion molecule 2 (NECTIN2, CD112); CD226 (DNAM-1); CD244 (2B4, SLAMF4), Poliovirus receptor (PVR) cell adhesion molecule (PVR, CD155). See, *e.g.*, Xu, et al., *J Exp Clin Cancer Res.* (2018) 37:110.

[0233] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with one or more blockers or inhibitors of one or more NK-cell inhibitory immune checkpoint proteins or receptors. Illustrative NK-cell inhibitory immune checkpoint proteins or receptors include without limitation killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR, CD158E1); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2); killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3); killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1); killer cell lectin like receptor C1 (KLRC1, NKG2A, CD159A); and killer cell lectin like receptor D1 (KLRD1, CD94). In various embodiments, the FLT3L-Fc fusion proteins, homodimers, heterodimers, polynucleotides, vectors, LNPs and/or pharmaceutical compositions, as described herein, are combined with one or more agonist or activators of one or more NK-cell stimulatory immune checkpoint proteins or receptors. Illustrative NK-cell stimulatory immune checkpoint proteins or receptors include without limitation CD16, CD226 (DNAM-1); CD244 (2B4, SLAMF4); killer cell lectin like receptor K1 (KLRK1, NKG2D, CD314); SLAM family member 7 (SLAMF7). See, *e.g.*, Davis, et al., *Semin Immunol.* (2017) 31:64–75; Fang, et al., *Semin Immunol.* (2017) 31:37-54; and Chiossone, et al., *Nat Rev Immunol.* (2018) 18(11):671-688.

[0234] In some embodiments, the one or more immune checkpoint inhibitors comprises a proteinaceous (*e.g.*, antibody or fragment thereof, or antibody mimetic) inhibitor of PD-L1 (CD274), PD-1 (PDCD1) or CTLA4. In some embodiments, the one or more immune checkpoint inhibitors comprises a small organic molecule inhibitor of PD-L1 (CD274), PD-1 (PDCD1) or CTLA4.

[0235] Examples of inhibitors of CTLA4 that can be co-administered include without limitation ipilimumab, tremelimumab, BMS-986218, AGEN1181, AGEN1884, BMS-986249, MK-1308, REGN-4659, ADU-1604, CS-1002, BCD-145, APL-509, JS-007,

BA-3071, ONC-392, AGEN-2041, JHL-1155, KN-044, CG-0161, ATOR-1144, PBI-5D3H5, BPI-002, as well as multi-specific inhibitors FPT-155 (CTLA4/PD-L1/CD28), PF-06936308 (PD-1/CTLA4), MGD-019 (PD-1/CTLA4), KN-046 (PD-1/CTLA4), MEDI-5752 (CTLA4/PD-1), XmAb-20717 (PD-1/CTLA4), and AK-104 (CTLA4/PD-1).

[0236] Examples of inhibitors of PD-L1 (CD274) or PD-1 (PDCD1) that can be co-administered include without limitation pembrolizumab, nivolumab, cemiplimab, pidilizumab, AMP-224, MEDI0680 (AMP-514), spartalizumab, atezolizumab, avelumab, durvalumab, BMS-936559, CK-301, PF-06801591, BGB-A317 (tislelizumab), GLS-010 (WBP-3055), AK-103 (HX-008), AK-105, CS-1003, HLX-10, MGA-012, BI-754091, AGEN-2034, JS-001 (toripalimab), JNJ-63723283, genolimzumab (CBT-501), LZM-009, BCD-100, LY-3300054, SHR-1201, SHR-1210 (camrelizumab), Sym-021, ABBV-181, PD1-PIK, BAT-1306, (MSB0010718C), CX-072, CBT-502, TSR-042 (dostarlimab), MSB-2311, JTX-4014, BGB-A333, SHR-1316, CS-1001 (WBP-3155, KN-035, IBI-308 (sintilimab), HLX-20, KL-A167, STI-A1014, STI-A1015 (IMC-001), BCD-135, FAZ-053, TQB-2450, MDX1105-01, GS-4224, GS-4416, INCB086550, MAX10181, as well as multi-specific inhibitors FPT-155 (CTLA4/PD-L1/CD28), PF-06936308 (PD-1/CTLA4), MGD-013 (PD-1/LAG-3), FS-118 (LAG-3/PD-L1) MGD-019 (PD-1/CTLA4), KN-046 (PD-1/CTLA4), MEDI-5752 (CTLA4/PD-1), RO-7121661 (PD-1/TIM-3), XmAb-20717 (PD-1/CTLA4), AK-104 (CTLA4/PD-1), M7824 (PD-L1/TGF β -EC domain), CA-170 (PD-L1/VISTA), CDX-527 (CD27/PD-L1), LY-3415244 (TIM3/PDL1), and INBRX-105 (4-1BB/PDL1).

[0237] In some embodiments, the small molecule inhibitor of CD274 or PDCD1 is selected from the group consisting of GS-4224, GS-4416, INCB086550 and MAX10181. In some embodiments, the small molecule inhibitor of CTLA4 comprises BPI-002.

[0238] In various embodiments, the antibodies or antigen-binding fragments as described herein are combined with anti-TIGIT antibodies, such as BMS-986207, RG-6058, AGEN-1307

TNF Receptor Superfamily (TNFRSF) Member Agonists or Activators

[0239] In various embodiments, the antibodies or antigen-binding fragments as described herein are combined with an agonist of one or more TNF receptor superfamily (TNFRSF) members, *e.g.*, an agonist of one or more of TNFRSF1A (NCBI Gene ID: 7132), TNFRSF1B (NCBI Gene ID: 7133), TNFRSF4 (OX40, CD134; NCBI Gene ID:

7293), TNFRSF5 (CD40; NCBI Gene ID: 958), TNFRSF6 (FAS, NCBI Gene ID: 355), TNFRSF7 (CD27, NCBI Gene ID: 939), TNFRSF8 (CD30, NCBI Gene ID: 943), TNFRSF9 (4-1BB, CD137, NCBI Gene ID: 3604), TNFRSF10A (CD261, DR4, TRAILR1, NCBI Gene ID: 8797), TNFRSF10B (CD262, DR5, TRAILR2, NCBI Gene ID: 8795), TNFRSF10C (CD263, TRAILR3, NCBI Gene ID: 8794), TNFRSF10D (CD264, TRAILR4, NCBI Gene ID: 8793), TNFRSF11A (CD265, RANK, NCBI Gene ID: 8792), TNFRSF11B (NCBI Gene ID: 4982), TNFRSF12A (CD266, NCBI Gene ID: 51330), TNFRSF13B (CD267, NCBI Gene ID: 23495), TNFRSF13C (CD268, NCBI Gene ID: 115650), TNFRSF16 (NGFR, CD271, NCBI Gene ID: 4804), TNFRSF17 (BCMA, CD269, NCBI Gene ID: 608), TNFRSF18 (GITR, CD357, NCBI Gene ID: 8784), TNFRSF19 (NCBI Gene ID: 55504), TNFRSF21 (CD358, DR6, NCBI Gene ID: 27242), and TNFRSF25 (DR3, NCBI Gene ID: 8718).

[0240] Example anti-TNFRSF4 (OX40) antibodies that can be co-administered include without limitation, MEDI6469, MEDI6383, MEDI0562 (tavolixizumab), MOXR0916, PF-04518600, RG-7888, GSK-3174998, INCAGN1949, BMS-986178, GBR-8383, ABBV-368, and those described in WO2016179517, WO2017096179, WO2017096182, WO2017096281, and WO2018089628.

[0241] Example anti-TNFRSF5 (CD40) antibodies that can be co-administered include without limitation RG7876, SEA-CD40, APX-005M and ABBV-428.

[0242] In some embodiments, the anti-TNFRSF7 (CD27) antibody varlilumab (CDX-1127) is co-administered.

[0243] Example anti-TNFRSF9 (4-1BB, CD137) antibodies that can be co-administered include without limitation urelumab, utomilumab (PF-05082566), AGEN2373 and ADG-106.

[0244] Example anti-TNFRSF18 (GITR) antibodies that can be co-administered include without limitation, MEDI1873, FPA-154, INCAGN-1876, TRX-518, BMS-986156, MK-1248, GWN-323, and those described in WO2017096179, WO2017096276, WO2017096189, and WO2018089628. In some embodiments, an antibody, or fragment thereof, co-targeting TNFRSF4 (OX40) and TNFRSF18 (GITR) is co-administered. Such antibodies are described, *e.g.*, in WO2017096179 and WO2018089628.

Bi-and Tri-Specific Natural Killer (NK)-Cell Engagers

[0245] In various embodiments, the antibodies or antigen-binding fragments as described herein, are combined with a bi-specific NK-cell engager (BiKE) or a tri-

specific NK-cell engager (TriKE) (*e.g.*, not having an Fc) or bi-specific antibody (*e.g.*, having an Fc) against an NK cell activating receptor, *e.g.*, CD16A, C-type lectin receptors (CD94/NKG2C, NKG2D, NKG2E/H and NKG2F), natural cytotoxicity receptors (NKp30, NKp44 and NKp46), killer cell C-type lectin-like receptor (NKp65, NKp80), Fc receptor FcγR (which mediates antibody-dependent cell cytotoxicity), SLAM family receptors (*e.g.*, 2B4, SLAMF6 and SLAMF7), killer cell immunoglobulin-like receptors (KIR) (KIR-2DS and KIR-3DS), DNAM-1 and CD137 (41BB). Illustrative anti-CD16 bi-specific antibodies, BiKEs or TriKEs that can be co-administered include AFM26 (BCMA/CD16A) and AFM-13 (CD16/CD30). As appropriate, the anti-CD16 binding bi-specific molecules may or may not have an Fc. BiKEs and TriKEs are described, *e.g.*, in Felices, et al., *Methods Mol Biol.* (2016) 1441:333–346; Fang, et al., *Semin Immunol.* (2017) 31:37-54. Examples of a trispecific NK cell engager (TRiKE) include OXS-3550, and CD16-IL-15-B7H3 TriKe.

Phosphatidylinositol 3-kinase (PI3K) Inhibitors

[0246] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a PI3K inhibitor. Examples of PI3K inhibitors that can be combined with an antibody of this disclosure include idelalisib, alpelisib, buparlisib, CAI orotate, copanlisib, duvelisib, gedatolisib, neratinib, panulisib, perifosine, pictilisib, pilaralisib, puquitinib mesylate, rigosertib, rigosertib sodium, sonolisib, taselisib, AMG-319, AZD-8186, BAY-1082439, CLR-1401, CLR-457, CUDC-907, DS-7423, EN-3342, GSK-2126458, GSK-2269577, GSK-2636771, INCB-040093, LY-3023414, MLN-1117, PQR-309, RG-7666, RP-6530, RV-1729, SAR-245409, SAR-260301, SF-1126, TGR-1202, UCB-5857, VS-5584, XL-765, and ZSTK-474.

alpha-4/beta-7 antagonists

[0247] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an alpha-4/beta-7 antagonist. Examples of Integrin alpha-4/beta-7 antagonists that can be combined with an antibody of this disclosure include PTG-100, TRK-170, abrilumab, etrolizumab, carotegrast methyl, and vedolizumab.

[0248] Examples of HIV antibodies, bispecific antibodies, and “antibody-like” therapeutic proteins that can be combined with an antibody of this disclosure include DARTs®, DUOBODIES®, BITES®, XmAbs®, TandAbs®, Fab derivatives, bNAbs (broadly neutralizing HIV-1 antibodies), BMS-936559, TMB-360, and those targeting HIV gp120 or gp41, antibody-Recruiting Molecules targeting HIV, anti-CD63

monoclonal antibodies, anti-GB virus C antibodies, anti-GP120/CD4, CCR5 bispecific antibodies, anti-nef single domain antibodies, anti-Rev antibody, camelid derived anti-CD18 antibodies, camelid-derived anti-ICAM-1 antibodies, DCVax-001, gp140 targeted antibodies, gp41-based HIV therapeutic antibodies, human recombinant mAbs (PGT-121), ibalizumab, Immuglo, MB-66. Examples of those targeting HIV in such a manner include bavituximab, UB-421, C2F5, 2G12, C4E10, C2F5+C2G12+C4E10, 8ANC195, 3-BNC-117, 3BNC117-LS, 3BNC60, D1D2, 10-1074, 10-1074-LS, GS-9722, DH411-2, BG18, PGT145, PGT121, PGT122, PGT-151, PGT-133, PGT-134, PGT-135, PGT-128, MDX010 (ipilimumab), DH511, DH511-2, N6, N6LS, N49P6, N49P7, N49P7.1, N49P9, N49P11, N60P1.1, N60P25.1, N60P2.1, N60P31.1, N60P22, NIH 45-46, PG9, PG16, 2Dm2m, 4Dm2m, 6Dm2m, PGDM1400, MDX010 (ipilimumab), VRC01, VRC-01-LS, A32, 7B2, 10E8, VRC-07-523, VRC07-523LS, 10E8VLS, 3810109, 10E8v4, IMC-HIV, iMabm36, eCD4-Ig, IOMA, CAP256-VRC26.25, DRVIA7, VRC-HIVMAB080-00-AB, VRC-HIVMAB060-00-AB, P2G12, VRC07 and SF12. Examples of HIV bispecific and trispecific antibodies include MGD014, TMB-bispecific, SAR-441236, VRC-01/PGDM-1400/10E8v4, 10E8.4/iMab, 10E8v4/PGT121-VRC01. Example of *in vivo* delivered bnABs such as AAV8-VRC07; mRNA encoding anti-HIV antibody VRC01; and engineered B-cells encoding 3BNC117 (Hartweiger *et al.*, *J. Exp. Med.* (2019), 1301).

Pharmacokinetic Enhancers

[0249] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a pharmacokinetic enhancer. Examples of pharmacokinetic enhancers that can be combined with an antibody of this disclosure include cobicistat and ritonavir.

Additional Therapeutic Agents

[0250] Examples of additional therapeutic agents that can be combined with an antibody of this disclosure include the compounds disclosed in WO 2004/096286 (Gilead Sciences), WO 2006/015261 (Gilead Sciences), WO 2006/110157 (Gilead Sciences), WO 2012/003497 (Gilead Sciences), WO 2012/003498 (Gilead Sciences), WO 2012/145728 (Gilead Sciences), WO 2013/006738 (Gilead Sciences), WO 2013/159064 (Gilead Sciences), WO 2014/100323 (Gilead Sciences), US 2013/0165489 (University of Pennsylvania), US 2014/0221378 (Japan Tobacco), US 2014/0221380 (Japan Tobacco), WO 2009/062285 (Boehringer Ingelheim), WO 2010/130034 (Boehringer Ingelheim), WO 2013/006792 (Pharma Resources), US 20140221356 (Gilead Sciences), US

20100143301 (Gilead Sciences) and WO 2013/091096 (Boehringer Ingelheim).

HIV Vaccines

[0251] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with an HIV vaccine. In various embodiments, the HIV vaccine elicits a T-cell response. Illustrative vaccines that can be combined with the herein described antibodies and fragments thereof include without limitation viral vectored vaccines (e.g., arenaviruses, adenoviruses, poxviruses, rhabdoviruses) as well as nucleic acid-based vaccines (e.g., DNA, RNA and self-replicating RNA). In some embodiments, the anti-HIV vaccine comprises one or more polypeptide vaccine immunogens. Examples of HIV vaccines that can be combined with an antibody of this disclosure include peptide vaccines, recombinant subunit protein vaccines, live vector vaccines, DNA vaccines, CD4-derived peptide vaccines, vaccine combinations, adenoviral vector vaccines, Chimp adenoviral vaccines (e.g., ChAdOX1, ChAd68, ChAd3 etc), Coxsackieviruses based vaccines, Gorilla adenovirus vaccines, arenavirus vaccines (LCMV, Pichinde), measles virus based vaccine, Varicella-zoster virus based vaccine, Human parainfluenza virus 3 (PIV3) based vaccines, poxvirus based vaccine (modified vaccinia virus Ankara (MVA), the NYVAC, and the ALVAC strains); rhabdovirus-based vaccines, such as VSV and marabavirus; alphavirus-based vaccines, such as semliki forest virus, venezuelan equine encephalitis virus and sindbis virus; (see Lauer, *Clinical and Vaccine Immunology*, (2017), DOI: 10.1128/CDVI.00298-16); LNP formulated mRNA based therapeutic vaccines; LNP-formulated self-replicating RNA/self-amplifying RNA vaccines, rgp120 (AIDSVAX), ALVAC HIV (vCP1521)/AIDSVAX B/E (gp120) (RV144), monomeric gp120 HIV-1 subtype C vaccine, Remune, ITV-1, Contre Vir, Ad5-ENVA-48, DCVax-001 (CDX-2401), Vacc-4x, Vacc-C5, VAC-3S, multiclade DNA recombinant adenovirus-5 (rAd5), rAd5 gag-pol env A/B/C vaccine, Pennvax-G, Pennvax-GP, Pennvax-G/MVA-CMDR, HIV-TriMix-mRNA vaccine, HIV-LAMP-vax, Ad35, Ad35-GRIN, NAcGM3/VSSP ISA-51, poly-ICLC adjuvanted vaccines, TatImmune, GTU-multiHIV (FIT-06), gp140[delta]V2.TV1+MF-59, rVSVIN HIV-1 gag vaccine, SeV-Gag vaccine, AT-20, DNK-4, ad35-Grin/ENV, TBC-M4, HIVAX, HIVAX-2, NYVAC-HIV-PT1, NYVAC-HIV-PT4, DNA-HIV-PT123, rAAV1-PG9DP, GOVX-B11, GOVX-B21, TVI-HIV-1, Ad-4 (Ad4-env Clade C+Ad4-mGag), Paxvax, EN41-UGR7C, EN41-FPA2, PreVaxTat, AE-H, MYM-V101, CombiHIVvac, ADVAX, MYM-V201, MVA-CMDR, DNA-Ad5 gag/pol/nef/nev (HVTN505), MVATG-17401, ETV-01, CDX-1401, rcAD26.MOS1.HIV-Env, Ad26.Mod.HIV

vaccine, Ad26.Mod.HIV + MVA mosaic vaccine + gp140, AGS-004, AVX-101, AVX-201, PEP-6409, SAV-001, ThV-01, TL-01, TUTI-16, VGX-3300, IHV-001, and virus-like particle vaccines such as pseudovirion vaccine, CombiVICHvac, LFn-p24 B/C fusion vaccine, GTU-based DNA vaccine, HIV gag/pol/nef/env DNA vaccine, anti-TAT HIV vaccine, conjugate polypeptides vaccine, dendritic-cell vaccines (*e.g.*, such as DermaVir), gag-based DNA vaccine, GI-2010, gp41 HIV-1 vaccine, HIV vaccine (PIKA adjuvant), I i-key/MHC class II epitope hybrid peptide vaccines, ITV-2, ITV-3, ITV-4, LIPO-5, multiclade Env vaccine, MVA vaccine, Pennvax-GP, pp71-deficient HCMV vector HIV gag vaccine, recombinant peptide vaccine (HIV infection), NCI, rgp160 HIV vaccine, RNative HIV vaccine, SCB-703, Tat Oyi vaccine, TBC-M4, therapeutic HIV vaccine, UBI HIV gp120, Vacc-4x + romidepsin, variant gp120 polypeptide vaccine, rAd5 gag-pol env A/B/C vaccine, DNA.HTI and MVA.HTI, VRC-HIVDNA016-00-VP + VRC-HIVADV014-00-VP, INO-6145, JNJ-9220, gp145 C.6980; eOD-GT8 60mer based vaccine, PD-201401, env (A, B, C, A/E)/gag (C) DNA Vaccine, gp120 (A,B,C,A/E) protein vaccine, PDPHV-201401, Ad4-EnvCN54, EnvSeq-1 Envs HIV-1 vaccine (GLA-SE adjuvanted), HIV p24gag pri, me-boost plasmid DNA vaccine, arenavirus vector-based vaccines (Vaxwave, TheraT), MVA-BN HIV-1 vaccine regimen, UBI HIV gp120, mRNA based prophylactic vaccines, and TBL-1203HI.

Birth control (contraceptive) combination therapy

[0252] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a birth control or contraceptive regimen. Therapeutic agents used for birth control (contraceptive) that can be combined with an antibody of this disclosure include cyproterone acetate, desogestrel, dienogest, drospirenone, estradiol valerate, ethinyl Estradiol, ethynodiol, etonogestrel, levomefolate, levonorgestrel, lynestrenol, medroxyprogesterone acetate, mestranol, mifepristone, misoprostol, nomegestrol acetate, norelgestromin, norethindrone, noretynodrel, norgestimate, ormeloxifene, seggestersone acetate, ulipristal acetate, and any combinations thereof.

[0253] In one embodiment, an antibody disclosed herein, or a pharmaceutically acceptable salt thereof, is combined with one, two, three, four or more additional therapeutic agents selected from ATRIPLA® (efavirenz, tenofovir disoproxil fumarate, and emtricitabine); COMPLERA® (EVIPLERA®; rilpivirine, tenofovir disoproxil fumarate, and emtricitabine); STRIBILD® (elvitegravir, cobicistat, tenofovir disoproxil fumarate, and emtricitabine); TRUVADA® (tenofovir disoproxil fumarate and emtricitabine; TDF +FTC); DESCOVY® (tenofovir alafenamide and emtricitabine);

ODEFSEY® (tenofovir alafenamide, emtricitabine, and rilpivirine); GENVOYA® (tenofovir alafenamide, emtricitabine, cobicistat, and elvitegravir); adefovir; adefovir dipivoxil; cobicistat; emtricitabine; tenofovir; tenofovir disoproxil; tenofovir disoproxil fumarate; tenofovir alafenamide; tenofovir alafenamide hemifumarate; TRIUMEQ® (dolutegravir, abacavir, and lamivudine); dolutegravir, abacavir sulfate, and lamivudine; raltegravir; raltegravir and lamivudine; maraviroc; enfuvirtide; ALUVIA® (KALETRA®; lopinavir and ritonavir); COMBIVIR® (zidovudine and lamivudine; AZT+3TC); EPZICOM® (LIVEXA®; abacavir sulfate and lamivudine; ABC+3TC); TRIZIVIR® (abacavir sulfate, zidovudine, and lamivudine; ABC+AZT+3TC); rilpivirine; rilpivirine hydrochloride; atazanavir sulfate and cobicistat; atazanavir and cobicistat; darunavir and cobicistat; atazanavir; atazanavir sulfate; dolutegravir; elvitegravir; ritonavir; atazanavir sulfate and ritonavir; darunavir; lamivudine; prolastin; fosamprenavir; fosamprenavir calcium efavirenz; etravirine; nelfinavir; nelfinavir mesylate; interferon; didanosine; stavudine; indinavir; indinavir sulfate; tenofovir and lamivudine; zidovudine; nevirapine; saquinavir; saquinavir mesylate; aldesleukin; zalcitabine; tipranavir; amprenavir; delavirdine; delavirdine mesylate; Radha-108 (receptol); lamivudine and tenofovir disoproxil fumarate; efavirenz, lamivudine, and tenofovir disoproxil fumarate; phosphazid; lamivudine, nevirapine, and zidovudine; abacavir; and abacavir sulfate.

[0254] In some embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with an HIV nucleoside or nucleotide inhibitor of reverse transcriptase and an HIV non-nucleoside inhibitor of reverse transcriptase. In another specific embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with an HIV nucleoside or nucleotide inhibitor of reverse transcriptase, and an HIV protease inhibiting compound. In an additional embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with an HIV nucleoside or nucleotide inhibitor of reverse transcriptase, an HIV non-nucleoside inhibitor of reverse transcriptase, and a pharmacokinetic enhancer. In certain embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with at least one HIV nucleoside inhibitor of reverse transcriptase, an integrase inhibitor, and a pharmacokinetic enhancer. In another embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with two HIV nucleoside or nucleotide inhibitors of reverse transcriptase.

[0255] In a certain embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with abacavir sulfate, tenofovir, tenofovir disoproxil, tenofovir disoproxil fumarate, tenofovir disoproxil hemifumarate, tenofovir alafenamide, or tenofovir alafenamide hemifumarate.

[0256] In another embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with tenofovir, tenofovir disoproxil, tenofovir disoproxil fumarate, tenofovir alafenamide, or tenofovir alafenamide hemifumarate.

[0257] In yet another embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with a first additional therapeutic agent selected from the group consisting of abacavir sulfate, tenofovir, tenofovir disoproxil, tenofovir disoproxil fumarate, tenofovir alafenamide, and tenofovir alafenamide hemifumarate, and a second additional therapeutic agent selected from the group consisting of emtricitabine and lamivudine.

[0258] In another embodiment, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with a first additional therapeutic agent selected from the group consisting of tenofovir, tenofovir disoproxil, tenofovir disoproxil fumarate, tenofovir alafenamide, and tenofovir alafenamide hemifumarate, and a second additional therapeutic agent, wherein the second additional therapeutic agent is emtricitabine.

[0259] In some embodiments, an antibody disclosed herein, or a pharmaceutical composition thereof, is combined with a first additional therapeutic agent (a contraceptive) selected from the group consisting of cyproterone acetate, desogestrel, dienogest, drospirenone, estradiol valerate, ethinyl Estradiol, ethynodiol, etonogestrel, levomefolate, levonorgestrel, lynestrenol, medroxyprogesterone acetate, mestranol, mifepristone, misoprostol, nomegestrol acetate, norelgestromin, norethindrone, noretynodrel, norgestimate, ormeloxifene, segestersone acetate, ulipristal acetate, and any combinations thereof.

Gene Therapy and Cell Therapy

[0260] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a gene or cell therapy regimen. Gene therapy and cell therapy include without limitation the genetic modification to silence a gene; genetic approaches to directly kill the infected cells; the infusion of immune cells designed to replace most of the patient's own immune system to enhance the immune response to infected cells, or activate the patient's own immune system to kill infected cells, or find and kill the

infected cells; genetic approaches to modify cellular activity to further alter endogenous immune responsiveness against the infection. Examples of dendritic cell therapy include AGS-004. CCR5 gene editing agents include SB-728T. CCR5 gene inhibitors include Cal-1. In some embodiments, C34-CCR5/C34-CXCR4 expressing CD4-positive T-cells are co-administered with the herein described antibodies or antigen-binding fragments thereof. In some embodiments, the antibodies or antigen-binding fragments are co-administered with AGT-103-transduced autologous T-cell therapy or AAV-eCD4-Ig gene therapy.

Gene Editors

[0261] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a gene editor, *e.g.*, an HIV targeted gene editor. In various embodiments, the genome editing system can be selected from the group consisting of: a CRISPR/Cas9 complex, a zinc finger nuclease complex, a TALEN complex, a homing endonucleases complex, and a meganuclease complex. An illustrative HIV targeting CRISPR/Cas9 system includes without limitation EBT-101.

CAR-T-cell therapy

[0262] In some embodiments, the antibodies or antigen-binding fragments described herein can be co-administered with a population of immune effector cells engineered to express a chimeric antigen receptor (CAR), wherein the CAR comprises an HIV antigen binding domain. The HIV antigen include an HIV envelope protein or a portion thereof, gp120 or a portion thereof, a CD4 binding site on gp120, the CD4-induced binding site on gp120, N glycan on gp120, the V2 of gp120, the membrane proximal region on gp41. The immune effector cell is a T-cell or an NK cell. In some embodiments, the T-cell is a CD4⁺ T-cell, a CD8⁺ T-cell, or a combination thereof. Cells can be autologous or allogeneic. Examples of HIV CAR-T include VC-CAR-T, CMV-N6-CART, anti-CD4 CART-cell therapy, autologous hematopoietic stem cells genetically engineered to express a CD4 CAR and the C46 peptide.

TCR-T-cell therapy

[0263] In certain embodiments, the antibodies or antigen-binding fragments described herein are combined with a population of TCR-T-cells. TCR-T-cells are engineered to target HIV derived peptides present on the surface of virus-infected cells.

B-cell therapy

[0264] In certain embodiments, the antibodies or antigen-binding fragments described

herein are combined with a population of B cells genetically modified to express broadly neutralizing antibodies, such as 3BNC117 (Hartweger, *et al.*, *J. Exp. Med.* 2019, 1301, Moffett, *et al.*, *Sci. Immunol.* 4, eaax0644 (2019) 17 May 2019).

Kits

[0265] This disclosure also encompasses kits comprising one or more antibodies or antigen binding fragments, described herein, or conjugates thereof. In one instance, provided herein is a pharmaceutical pack or kit comprising one or more containers (*e.g.*, vials, ampules) filled with one or more of the ingredients of the pharmaceutical compositions described herein, such as one or more antibodies provided herein. In some instances, the kits contain a pharmaceutical composition described herein. In one embodiment, kits comprising an antibody disclosed herein, or a pharmaceutical composition thereof, in combination with one or more (*e.g.*, one, two, three, one or two, or one to three) additional therapeutic agents (such as those disclosed above) are provided.

[0266] In some embodiments, the kits comprise one or more unitary doses of the antibodies or antigen-binding fragments, or the polynucleotide or polynucleotides, in one or more containers. In some embodiments, the kits comprise one or more unitary doses of the antibodies or antigen-binding fragments and a second agent (*e.g.*, one or more additional agents) for treating an HIV infection in separate containers. In some embodiments, the kits further comprise one or more unitary doses of a toll-like receptor (TLR) agonist. In some embodiments, the TLR agonist is a TLR7 agonist or a TLR8 agonist. In some embodiments, the TLR7 agonist is selected from the group consisting of vesatolimod, imiquimod, and resiquimod. In some embodiments, the kits comprise one or more unitary doses of the antibodies or antigen-binding fragments, as described herein, and one or more unitary doses of a second, third or fourth anti-HIV antibody, or antigen-binding fragments thereof, wherein the second, third or fourth anti-HIV antibodies, or antigen-binding fragments thereof, bind to epitopes or regions of gp120 selected from the group consisting of: (i) third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan; (ii) second variable loop (V2) and/or Env trimer apex; (iii) gp120/gp41 interface; or (iv) silent face of gp120. In some embodiments, the second anti-HIV antibody or antigen-binding fragment thereof, binds to the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan. In some embodiments, the second anti-HIV antibody competes with or comprises VH and VL regions from an antibody selected from the group

consisting of GS-9722, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137, PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03. In some embodiments, the second anti-HIV antibody or antigen binding fragments thereof competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722 and PGT-121. In some embodiments, the kits comprise two or more unitary doses, wherein the unitary doses are the same. In some embodiments, the kits comprise two or more unitary doses, wherein the unitary doses are different.

[0267] Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

Examples

[0268] The following examples are provided to illustrate the various embodiments and are not to be interpreted as limiting the scope of the present application. To the extent that specific materials are mentioned, it is merely for purposes of illustration and is not intended to limit the present application. One skilled in the art can develop equivalent means or reactants without the exercise of inventive capacity and without departing from the scope of the present application.

Example 1: ADCC Activity of Antibody A

[0269] ADCC of HIV-infected target CD4⁺ T cells by the antibodies were assayed *in vitro* using HIV-infected CEM.NKr.CCR5⁺Luc⁺ cells and primary human NK effector cells from independent healthy donors.

[0270] The study included both PGT121-sensitive and PGT121-resistant viruses and antibodies having modifications to the Fc (Fc-modified) of Antibody A. Table 1 summarizes the killing potency and efficacy of Antibodies A, A-1, A-2, A-3, A-4, A-5 and A-6, when assayed in the presence of 5 mg/mL of human serum IgG and using primary human NK cells from three independent human donors and CEM.NKr.CCR5⁺Luc⁺ cells infected with viral isolates 92US712 or 92US657.

[0271] Table 1. ADCC activity

92US712-infected cells								
	ID:	A	A-2	A-1	A-3	A-4	A-5	A-6
E _{max} (%)	NK Donor 1	48	76	77	77	78	79	68
	NK Donor 2	7	60	62	59	61	61	54
	NK Donor 3	27	60	62	67	64	66	51
EC ₅₀ (μg/mL)	NK Donor 1	2.23	0.18	0.19	0.07	0.20	0.19	0.27
	NK Donor 2	>100	0.08	0.09	0.08	0.13	0.08	0.54
	NK Donor 3	3.63	0.20	0.16	0.20	0.40	0.09	0.18
92US657-infected cells								
	ID:	A	A-2	A-1	A-3	A-4	A-5	A-6
E _{max} (%)	NK Donor 1	2	59	58	49	52	58	54
	NK Donor 2	0	51	50	49	53	61	53
	NK Donor 3	0	56	52	56	50	61	52
EC ₅₀ (μg/mL)	NK Donor 1	>100	0.54	0.74	0.68	0.60	0.57	3.93
	NK Donor 2	>100	0.81	0.55	0.67	1.21	0.83	1.21
	NK Donor 3	>100	1.13	0.37	1.39	1.64	0.76	3.21

EC₅₀ noted as >100 μg/mL for dose responses with E_{max} < 10%

[0272] The Fc-modified antibodies exhibited increased killing of HIV-1-infected target CD4 T cells compared to Antibody A *in vitro* by primary human NK cells from independent donors and target cells infected with different viral isolates (Table 1). Antibody A-mediated minimal killing (E_{max} <10%) with primary NK cells from some donors, while with NK cells from other donors killing was detectable. Compared to Antibody A, the Fc-modified antibodies exhibited increased potency (EC₅₀) and maximum killing (E_{max}) of HIV-1-infected cells, as observed in ADCC assays performed with primary human NK cells from three independent healthy donors (Table 1). The increase in potency observed ranged from about 10- to 40-fold with donors where Antibody A was active. A panel of 22 infected target cell cultures was generated by infecting CEM.NKr.CCR5⁺Luc⁺ cells with 22 unique viral clones resistant to neutralization (*e.g.*, infected cell killing) by PGT121.60 (*see*, WO 2017/106346). ADCC activity and breadth of Antibody A-1 and Antibody PGT121.60 were evaluated against this panel of infected target cells using primary human NK effector cells from healthy donors in the absence of competing serum IgG. 86% (19/22) of the infected target cell cultures resistant to ADCC by PGT121.60 were killed by Antibody A-1 (E_{max} >30%).

Antibody A-1 mediated ADCC of cells infected with HIV strains that were resistant to PGT121.60. The results of this assessment are summarized in Table 2.

[0273] Table 2. Infected cell killing of PGT121.60 resistant by Antibody A-1 and Antibody PGT121.60. Numbers depict ADCC Emax (%) average from two donors.

Virus	ADCC Emax (%)	
	PGT121.60	Antibody A-1
VS001	1.0	26.8
VS002	2.0	22.9
VS003	3.0	44.4
VS004	4.0	31.7
VS007	9	45
VS008	22	60
VS010	10	69
VS011	8.0	34.6
VS017	9.0	40.5
VS023	10.0	0.8
VS026	11.0	31.3
VS029	12.0	1.5
VS030	13.0	39.3
VS032	14.0	29.1
VS033	15.0	31.1
VS034	16.0	40.6
VS038	17.0	36.1
VS042	18.0	39.6
VS044	19.0	31.4
VS046	20.0	41.9
VS049	21.0	7.8
VS052	22.0	34.0

[0274] Antibody-dependent cellular cytotoxicity was also evaluated using HIV-infected primary CD4⁺ T cells as target cells and autologous primary NK cells, monocytes and neutrophils as effector cells.

[0275] The NK cells, monocytes and CD4⁺ T cells were isolated from PBMCs obtained from healthy donors, while neutrophils were isolated from whole blood from healthy

donors. Total CD4⁺ T cells were spininfected in the absence of T-cell activation to maintain low cell surface antigen expression levels and potentially mimic antigen expression levels on latently infected CD4⁺ T cells. Viral isolates used were 8176 and 92US076 (antibody A neutralization sensitive) and 8398 (antibody A neutralization resistant). Assays were performed in the presence of 1 mg/ml nonspecific human serum IgG which compete with effector mAbs for FcγR binding. Antibody-dependent killing was measured by the reduction in p24⁺ CD4 T cells using flow cytometry.

[0276] The killing AUC, EC₅₀ (μg/mL) and Emax (%) values are tabulated in Table 3-11.

[0277] Table 3. Killing AUC by NK cells

Virus	Donor	AUC (NK)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	49	108	103	168
	3594	18	142	139	205
302076	0117	26	116	78	18
	3594	28	76	101	26
8398	0117	20	4	0	203
	3594	0	6	15	211

[0278] Table 4. Killing EC₅₀ by NK cells

Virus	Donor	EC ₅₀ (NK)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	>100	1.101	0.949	0.246
	3594	>100	0.518	1.350	0.051
302076	0117	>100	1.701	7.602	100.000
	3594	>100	2.613	3.114	72.050
8398	0117	>100	>100	>100	0.339
	3594	>100	>100	>100	0.220

[0279] Table 5. Killing Emax by NK cells

Virus	Donor	Emax (NK)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	<20	56	54	65
	3594	<20	62	69	62
302076	0117	<20	68	69	20
	3594	<20	49	64	45
8398	0117	<20	<20	<20	79
	3594	<20	<20	<20	81

[0280] Table 6. Killing AUC by monocytes

Virus	Donor	AUC (Monocytes)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	0	83	87	122
	3594	17	141	159	157
302076	0117	24	54	61	24
	3594	138	166	158	108
8398	0117	0	0	4	53
	3594	0	13	4	186

[0281] Table 7. Killing EC50 by monocytes

Virus	Donor	EC50 (Monocytes)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	>100	0.402	0.490	0.232
	3594	>100	0.309	0.202	0.010
302076	0117	10.570	6.514	7.236	>100
	3594	0.006	0.062	0.088	0.019
8398	0117	>100	>100	>100	0.728
	3594	>100	>100	>100	0.201

[0282] Table 8. Killing Emax by monocytes

Virus	Donor	Emax (Monocytes)			
		A	A-1	1.52.64-1	PGT121.60
8176	0117	<20	34	35	44
	3594	<20	47	57	40
302076	0117	24	46	47	20
	3594	33	49	54	30
8398	0117	<20	<20	<20	25
	3594	<20	<20	<20	67

[0283] Table 9. Killing AUC by neutrophils

Virus	Donor	AUC (Neutrophils)			
		A	A-1	1.52.64-1	PGT121.60
8176	92132	41	74	89	125
	92602	21	47	45	71

[0284] Table 10. Killing EC50 by neutrophils

Virus	Donor	EC50 (Neutrophils)			
		A	A-1	1.52.64-1	PGT121.60
8176	92132	>100	0.231	0.307	0.012
	92602	>100	1.097	0.199	0.232

[0285] Table 11. Killing Emax by neutrophils

Virus	Donor	Emax (Neutrophils)			
		A	A-1	1.52.64-1	PGT121.60
8176	92132	<20	29	37	34
	92602	<20	23	<20	25

[0286] The results presented in Tables 3-11 demonstrate that, consistent with NK-mediated ADCC of CEM cells, the Fc-engineered mAbs (1.52.64-1, A-1 and PGT121.60) also exhibited increased killing of HIV-infected primary CD4 T cells by NK cells, monocytes and neutrophils compared to Antibody A.

Example 2: Antibody Campaign

[0287] The sequences of Antibody A and Antibody B were compared to the human germline, revealing several mutations, insertions and deletions both inside and outside of the CDRs. Briefly, a contiguous region of germline mismatch in heavy chain framework region 3 (HC FR3) was identified at position 72-78 of the heavy chain (HC). Four amino acid insertions were identified between position 74 and 75 in HC FR3. A germline deletion was identified in CDR L1 at positions 27-30 of the light chain (LC). A contiguous region of germline mismatch was identified in light chain framework region 3 (LC FR3) at position 65-77. A N72 linked consensus glycosylation motif was identified in LC FR3 at position 72-74. A germline deletion in CDR L3 was identified at position 92-95. Two residues that are highly conserved in human IgG light chains (F98 and G99) were mutated in both Antibody A and Antibody B.

[0288] Mass spectrometry studies of ExpiCHO expressed Antibody A were conducted to determine whether there was glycosylation at LC position 72-74. Accelerated stress and potency assays were conducted to see if there were any chemical liabilities (*e.g.*

oxidation, deamidation etc.) present in antibody A or its variants. Due to the high degree of somatic hypermutation, T-cell epitope mapping of the primary sequence was conducted to identify potentially immunogenic motifs. Additionally, an iterative protein engineering campaign was conducted in order to generate new antibodies without the N72 glycosylation motif and/or with a closer overall match to the human germline. Without being bound to any theories, this campaign may yield new antibodies that have desired properties including but not limited to a reduced risk of immunogenicity, HIV neutralization potency and breadth equal to or better than Antibody A or Antibody B, and/or improved biophysical and development properties.

[0289] Table 12 provides the SEQ ID NOs of the VH and VL CDRs (according to the Kabat definition) of the anti-gp120 antibodies disclosed herein.

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
A-1	137	138	139	140	141	142
A	137	138	139	140	141	142
1v2-1	137	138	139	140	141	142
1.2.1-1	137	138	139	140	141	142
1.1.2-1	137	138	139	140	141	142
1.2.2-1	137	138	139	140	141	142
1.3.1-1	137	138	139	140	141	142
1.4.1-1	137	138	139	140	141	142
1.5.1-1	137	138	139	140	141	142
1.6.1-1	137	138	139	140	141	142
1.7.1-1	137	138	139	140	141	142
1.8.1-1	137	138	139	140	141	142
1.9.1-1	137	138	139	140	141	142
1.10.1-1	159	138	139	140	141	142
1.11.1-1	159	138	139	140	141	142
1.15.1-1	137	160	139	140	141	142
1.16.1-1	137	161	139	140	141	142
1.17.1-1	137	162	139	140	141	142
1.18.1-1	137	163	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
1.19.1-1	137	138	139	140	141	142
1.20.1-1	137	138	139	140	141	142
1.21.1-1	137	138	139	140	141	142
1.22.1-1	137	138	139	140	141	142
1.24.1-1	137	138	139	140	141	142
1.25.1-1	137	138	139	140	141	142
1.26.1-1	137	138	139	140	141	142
1.27.1-1	137	138	164	140	141	142
1.28.1-1	137	138	164	140	141	142
1.29.1-1	137	138	139	140	141	142
1.30.1-1	137	138	139	140	141	142
1.1.3-1	137	138	139	140	141	142
1.1.4-1	137	138	139	140	141	142
1.1.5-1	137	138	139	140	141	142
1.1.6-1	137	138	139	140	165	142
1.1.7-1	137	138	139	140	166	142
1.1.8-1	137	138	139	140	168	142
1.1.9-1	137	138	139	140	167	142
1.1.10-1	137	138	139	140	141	142
1.1.11-1	137	138	139	140	141	142
1.1.12-1	137	138	139	140	141	142
1.1.13-1	137	138	139	140	141	142
1.1.14-1	137	138	139	140	141	142
1.1.15-1	137	138	139	140	141	142
1.1.16-1	137	138	139	140	141	142
1.1.17-1	137	138	139	140	141	142
1.1.18-1	137	138	139	140	141	142
1.1.19-1	137	138	139	140	141	142
1.1.20-1	137	138	139	140	141	142
1.1.21-1	137	138	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
1.1.22-1	137	138	139	140	141	142
1.1.23-1	137	138	139	140	141	142
1.1.24-1	137	138	139	140	141	142
1.1.25-1	137	138	139	140	141	142
1.1.26-1	137	138	139	140	141	142
1.1.27-1	137	138	139	140	141	142
1.1.28-1	137	138	139	140	141	142
1.1.29-1	137	138	139	140	141	142
1.1.30-1	137	138	139	140	141	142
1.12.15-1	137	138	139	140	141	142
1.13.15-1	137	138	139	140	141	142
1.14.15-1	137	138	139	140	141	142
1.12.17-1	137	138	139	140	141	142
1.13.17-1	137	138	139	140	141	142
1.14.17-1	137	138	139	140	141	142
1.31.1-1	137	138	139	140	141	142
1.32.1-1	137	138	139	140	141	142
1.33.1-1	137	138	139	140	141	142
1.34.1-1	137	138	164	140	141	142
1.35.1-1	159	138	164	140	141	142
1.36.1-1	159	138	164	140	141	142
1.1.31-1	137	138	139	140	141	142
1.31.31-1	137	138	139	140	141	142
1.32.31-1	137	138	139	140	141	142
1.33.31-1	137	138	139	140	141	142
1.34.31-1	137	138	164	140	141	142
1.35.31-1	159	138	164	140	141	142
1.36.31-1	159	138	164	140	141	142
1.1.32-1	137	138	139	140	141	142
1.31.32-1	137	138	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
1.32.32-1	137	138	139	140	141	142
1.33.32-1	137	138	139	140	141	142
1.34.32-1	137	138	164	140	141	142
1.35.32-1	159	138	164	140	141	142
1.36.32-1	159	138	164	140	141	142
1.1.33-1	137	138	139	140	166	142
1.31.33-1	137	138	139	140	166	142
1.32.33-1	137	138	139	140	166	142
1.33.33-1	137	138	139	140	166	142
1.34.33-1	137	138	164	140	166	142
1.35.33-1	159	138	164	140	166	142
1.36.33-1	159	138	164	140	166	142
1.1.34-1	137	138	139	140	166	142
1.31.34-1	137	138	139	140	166	142
1.32.34-1	137	138	139	140	166	142
1.33.34-1	137	138	139	140	166	142
1.34.34-1	137	138	164	140	166	142
1.35.34-1	159	138	164	140	166	142
1.36.34-1	159	138	164	140	166	142
1.1.35-1	137	138	139	140	166	142
1.31.35-1	137	138	139	140	166	142
1.32.35-1	137	138	139	140	166	142
1.33.35-1	137	138	139	140	166	142
1.34.35-1	137	138	164	140	166	142
1.35.35-1	159	138	164	140	166	142
1.36.35-1	159	138	164	140	166	142
1.1.36-1	137	138	139	140	166	142
1.31.36-1	137	138	139	140	166	142
1.32.36-1	137	138	139	140	166	142
1.33.36-1	137	138	139	140	166	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
1.34.36-1	137	138	164	140	166	142
1.35.36-1	159	138	164	140	166	142
1.36.36-1	159	138	164	140	166	142
1.1.37-1	137	138	139	140	141	142
1.1.38-1	137	138	139	140	141	142
1.1.39-1	137	138	139	140	141	142
1.1.40-1	137	138	139	140	141	142
1.1.41-1	137	138	139	140	141	142
1.1.42-1	137	138	139	140	141	142
1.1.43-1	137	138	139	140	141	142
1.1.44-1	137	138	139	140	141	142
1.1.45-1	137	138	139	140	141	142
1.1.46-1	137	138	139	140	141	142
1.1.47-1	137	138	139	140	141	142
1.1.48-1	137	138	139	140	141	142
1.1.49-1	137	138	139	140	141	142
1.37.51-1	137	138	139	140	141	142
1.8.52-1	137	138	139	140	141	142
1.1.54-1	137	138	139	140	141	142
A-2	137	138	139	140	141	142
B-1	153	138	154	140	141	142
2.1.2-1	153	138	154	140	141	142
1.1.64-1	137	138	139	140	141	142
1.1.67-1	137	138	139	140	141	142
1.1.72-1	137	138	139	140	141	142
1.1.75-1	137	138	139	140	141	142
1.1.78-1	137	138	139	140	141	142
A-3	137	138	139	140	141	142
A-4	137	138	139	140	141	142
A-5	137	138	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
A-6	137	138	139	140	141	142
1.41.5-1	137	138	139	140	141	142
1.41.81-1	137	138	139	140	141	142
1.1.82-1	137	138	139	140	141	142
1.41.83-1	137	138	139	140	141	142
1.1.84-1	137	138	139	140	141	142
1.41.85-1	137	138	139	140	141	142
1.41.86-1	137	138	139	140	141	142
1.41.87-1	137	138	139	140	141	142
1.1.88-1	137	138	139	140	141	142
1.41.89-1	137	138	139	140	141	142
1.1.90-1	137	138	139	140	141	142
1.41.91-1	137	138	139	140	141	142
1.41.92-1	137	138	139	140	141	142
1.41.93-1	137	138	139	140	141	142
1.1.94-1	137	138	139	140	141	142
1.41.95-1	137	138	139	140	141	142
1.1.96-1	137	138	139	140	141	142
1.41.97-1	137	138	139	140	141	142
1.41.98-1	137	138	139	140	141	142
1.41.99-1	137	138	139	140	141	142
1.1.100-1	137	138	139	140	141	142
1.41.101-1	137	138	139	140	141	142
1.1.102-1	137	138	139	140	141	142
1.41.103-1	137	138	139	140	141	142
1.1.110-1	137	138	139	140	141	142
1.1.111-1	137	138	139	140	141	142
1.1.112-1	137	138	139	140	141	142
1.1.113-1	137	138	139	140	141	142
2.1.3-1	153	138	154	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
2.1.4-1	153	138	154	140	141	142
2.2.1-1	137	138	139	140	141	142
2.3.1-1	153	138	139	140	141	142
3.1.8-1	137	138	139	140	141	142
2.2.8-1	137	138	139	140	141	142
2.3.8-1	153	138	139	140	141	142
3.1.9-1	137	138	139	140	141	142
2.2.9-1	137	138	139	140	141	142
2.3.9-1	153	138	139	140	141	142
1.1.115-1	137	138	139	140	141	142
3.1.10-1	137	138	139	140	141	142
2.2.10-1	137	138	139	140	141	142
2.3.10-1	153	138	139	140	141	142
1.1.116-1	137	138	139	140	141	142
3.1.11-1	137	138	139	140	141	142
2.2.11-1	137	138	139	140	141	142
2.3.11-1	153	138	139	140	141	142
1.1.117-1	137	138	139	140	141	142
3.1.12-1	137	138	139	140	141	142
2.2.12-1	137	138	139	140	141	142
2.3.12-1	153	138	139	140	141	142
1.1.118-1	137	138	139	140	141	142
3.1.13-1	137	138	139	140	141	142
2.2.13-1	137	138	139	140	141	142
2.3.13-1	153	138	139	140	141	142
3.1.14-1	137	138	139	140	141	142
2.2.14-1	137	138	139	140	141	142
2.3.14-1	153	138	139	140	141	142
3.1.5-1	137	138	139	140	141	142
2.2.5-1	137	138	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
2.3.5-1	153	138	139	140	141	142
3.1.15-1	137	138	139	140	141	142
2.2.15-1	137	138	139	140	141	142
2.3.15-1	153	138	139	140	141	142
1.1.119-1	137	138	139	140	141	142
3.1.7-1	137	138	139	140	141	142
2.2.7-1	137	138	139	140	141	142
2.3.7-1	153	138	139	140	141	142
3.1.2-1	137	138	139	140	141	142
2.2.2-1	137	138	139	140	141	142
2.3.2-1	153	138	139	140	141	142
3.1.16-1	137	138	139	140	141	142
2.2.16-1	137	138	139	140	141	142
2.3.16-1	153	138	139	140	141	142
3.1.17-1	137	138	139	140	141	142
2.2.17-1	137	138	139	140	141	142
2.3.17-1	153	138	139	140	141	142
3.1.18-1	137	138	139	140	141	142
2.2.18-1	137	138	139	140	141	142
2.3.18-1	153	138	139	140	141	142
1.1.120-1	137	138	139	140	141	142
3.1.19-1	137	138	139	140	141	142
2.2.19-1	137	138	139	140	141	142
2.3.19-1	153	138	139	140	141	142
1.1.121-1	137	138	139	140	141	142
3.1.20-1	137	138	139	140	141	142
2.2.20-1	137	138	139	140	141	142
2.3.20-1	153	138	139	140	141	142
1.1.122-1	137	138	139	140	141	142
1.1.123-1	137	138	139	140	141	142

Table 12. SEQ ID NOs of the VH and VL CDRs of Antibodies						
Antibody Name	VHCDR1	VHCDR2	VHCDR3	VLCDR1	VLCDR2	VLCDR3
1.1.124-1	137	138	139	140	141	142
1.1.125-1	137	138	139	140	141	142
1.1.126-1	137	138	139	140	141	142
1.1.127-1	137	138	139	140	141	142
1.1.128-1	137	138	139	140	141	142
1.1.129-1	137	138	139	140	141	142
1.1.130-1	137	138	139	140	141	142
1.1.131-1	137	138	139	140	141	142
1.1.132-1	137	138	139	140	141	142
1.1.133-1	137	138	139	140	141	142
1.1.134-1	137	138	139	140	141	142
1.1.135-1	137	138	139	140	141	142
1.1.138-1	137	138	139	570	141	142
1.42.1-1	137	138	139	140	141	142
1.43.1-1	137	138	139	140	141	142
1.44.1-1	137	138	139	140	141	142
1.45.1-1	137	138	139	140	141	142
1.46.1-1	153	138	139	140	141	142
1.47.1-1	137	138	139	140	141	142
1.49.1-1	137	138	139	140	141	142
1.50.1-1	137	138	139	140	141	142
1.51.1-1	137	138	154	140	141	142
1.1.104-1	137	138	139	140	141	142
3-1	137	138	139	140	141	142
1.52.1-1	137	138	139	140	141	142
1.52.64-1	137	138	139	140	141	142
1.52.90	137	138	139	140	141	142
2.4.1-1	153	138	139	140	141	142

[0290] Table 13 provides the SEQ ID NOs of the VH, VL, heavy and light chains of the anti-gp120 antibodies disclosed herein.

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
B	181	222	1	48
A-1	182	223	2	49
A	182	223	3	49
C-1	183	224	4	50
1v2-1	184	223	5	49
1.2.1-1	185	223	6	49
1.1.2-1	182	225	2	50
1.2.2-1	185	225	6	50
1.3.1-1	186	223	7	49
1.4.1-1	187	223	8	49
1.5.1-1	188	223	9	49
1.6.1-1	189	223	10	49
1.7.1-1	190	223	11	49
1.8.1-1	191	223	12	49
1.9.1-1	192	223	13	49
1.10.1-1	193	223	14	49
1.11.1-1	194	223	15	49
1.15.1-1	195	223	16	49
1.16.1-1	196	223	17	49
1.17.1-1	197	223	18	49
1.18.1-1	198	223	19	49
1.19.1-1	199	223	20	49
1.20.1-1	200	223	21	49
1.21.1-1	201	223	22	49
1.22.1-1	202	223	23	49
1.24.1-1	203	223	24	49
1.25.1-1	204	223	25	49
1.26.1-1	205	223	26	49

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.27.1-1	206	223	27	49
1.28.1-1	207	223	28	49
1.29.1-1	208	223	29	49
1.30.1-1	209	223	30	49
1.1.3-1	182	226	2	51
1.1.4-1	182	227	2	52
1.1.5-1	182	228	2	53
1.1.6-1	182	229	2	54
1.1.7-1	182	230	2	55
1.1.8-1	182	231	2	56
1.1.9-1	182	232	2	57
1.1.10-1	182	233	2	58
1.1.11-1	182	234	2	59
1.1.12-1	182	235	2	60
1.1.13-1	182	236	2	61
1.1.14-1	182	237	2	62
1.1.15-1	182	238	2	63
1.1.16-1	182	239	2	64
1.1.17-1	182	240	2	65
1.1.18-1	182	241	2	66
1.1.19-1	182	242	2	67
1.1.20-1	182	243	2	68
1.1.21-1	182	244	2	69
1.1.22-1	182	245	2	70
1.1.23-1	182	246	2	71
1.1.24-1	182	247	2	72
1.1.25-1	182	248	2	73
1.1.26-1	182	249	2	74
1.1.27-1	182	250	2	75
1.1.28-1	182	251	2	76
1.1.29-1	182	252	2	77

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.1.30-1	182	253	2	78
1.12.15-1	210	238	31	63
1.13.15-1	211	238	32	63
1.14.15-1	212	238	33	63
1.12.17-1	210	240	31	65
1.13.17-1	211	240	32	65
1.14.17-1	212	240	33	65
1.31.1-1	213	223	34	49
1.32.1-1	214	223	35	49
1.33.1-1	215	223	36	49
1.34.1-1	216	223	37	49
1.35.1-1	217	223	38	49
1.36.1-1	218	223	39	49
1.1.31-1	182	254	2	79
1.31.31-1	213	254	34	79
1.32.31-1	214	254	35	79
1.33.31-1	215	254	36	79
1.34.31-1	216	254	37	79
1.35.31-1	217	254	38	79
1.36.31-1	218	254	39	79
1.1.32-1	182	255	2	80
1.31.32-1	213	255	34	80
1.32.32-1	214	255	35	80
1.33.32-1	215	255	36	80
1.34.32-1	216	255	37	80
1.35.32-1	217	255	38	80
1.36.32-1	218	255	39	80
1.1.33-1	182	256	2	81
1.31.33-1	213	256	34	81
1.32.33-1	214	256	35	81
1.33.33-1	215	256	36	81

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.34.33-1	216	256	37	81
1.35.33-1	217	256	38	81
1.36.33-1	218	256	39	81
1.1.34-1	182	257	2	82
1.31.34-1	213	257	34	82
1.32.34-1	214	257	35	82
1.33.34-1	215	257	36	82
1.34.34-1	216	257	37	82
1.35.34-1	217	257	38	82
1.36.34-1	218	257	39	82
1.1.35-1	182	258	2	83
1.31.35-1	213	258	34	83
1.32.35-1	214	258	35	83
1.33.35-1	215	258	36	83
1.34.35-1	216	258	37	83
1.35.35-1	217	258	38	83
1.36.35-1	218	258	39	83
1.1.36-1	182	259	2	84
1.31.36-1	213	259	34	84
1.32.36-1	214	259	35	84
1.33.36-1	215	259	36	84
1.34.36-1	216	259	37	84
1.35.36-1	217	259	38	84
1.36.36-1	218	259	39	84
1.1.37-1	182	260	2	85
1.1.38-1	182	261	2	86
1.1.39-1	182	262	2	87
1.1.40-1	182	263	2	88
1.1.41-1	182	264	2	89
1.1.42-1	182	265	2	90
1.1.43-1	182	266	2	91

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.1.44-1	182	267	2	92
1.1.45-1	182	268	2	93
1.1.46-1	182	269	2	94
1.1.47-1	182	270	2	95
1.1.48-1	182	271	2	96
1.1.49-1	182	272	2	97
1.37.51-1	219	273	40	98
1.8.52-1	191	274	12	99
1.1.54-1	182	275	2	100
A-2	182	223	41	49
B-1	220	276	42	101
2.1.2-1	220	277	42	102
1.1.64-1	182	278	2	103
1.1.67-1	182	279	2	104
1.1.72-1	182	280	2	105
1.1.75-1	182	281	2	106
1.1.78-1	182	282	2	107
A-3	182	223	43	49
A-4	182	223	44	49
A-5	182	223	45	49
A-6	182	223	46	49
1.41.5-1	221	228	47	53
1.41.81-1	221	283	47	108
1.1.82-1	182	284	2	109
1.41.83-1	221	285	47	110
1.1.84-1	182	286	2	111
1.41.85-1	221	287	47	112
1.41.86-1	221	288	47	113
1.41.87-1	221	289	47	114
1.1.88-1	182	290	2	115
1.41.89-1	221	291	47	116

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.1.90-1	182	292	2	117
1.41.91-1	221	293	47	118
1.41.92-1	221	294	47	119
1.41.93-1	221	295	47	120
1.1.94-1	182	296	2	121
1.41.95-1	221	297	47	122
1.1.96-1	182	298	2	123
1.41.97-1	221	299	47	124
1.41.98-1	221	300	47	125
1.41.99-1	221	301	47	126
1.1.100-1	182	302	2	127
1.41.101-1	221	303	47	128
1.1.102-1	182	304	2	129
1.41.103-1	221	305	47	130
1.1.110-1	182	306	2	131
1.1.111-1	182	307	2	132
1.1.112-1	182	308	2	133
1.1.113-1	182	309	2	134
2.1.3-1	220	310	42	135
2.1.4-1	220	311	42	136
2.2.1-1	465	276	517	101
2.3.1-1	466	276	518	101
3.1.8-1	182	479	2	531
2.2.8-1	465	479	517	531
2.3.8-1	466	479	518	531
3.1.9-1	182	480	2	532
2.2.9-1	465	480	517	532
2.3.9-1	466	480	518	532
1.1.115-1	182	481	2	533
3.1.10-1	182	482	2	534
2.2.10-1	465	482	517	534

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
2.3.10-1	466	482	518	534
1.1.116-1	182	483	2	535
3.1.11-1	182	484	2	536
2.2.11-1	465	484	517	536
2.3.11-1	466	484	518	536
1.1.117-1	182	485	2	537
3.1.12-1	182	486	2	538
2.2.12-1	465	486	517	538
2.3.12-1	466	486	518	538
1.1.118-1	182	487	2	539
3.1.13-1	182	488	2	540
2.2.13-1	465	488	517	540
2.3.13-1	466	488	518	540
3.1.14-1	182	489	2	541
2.2.14-1	465	489	517	541
2.3.14-1	466	489	518	541
3.1.5-1	182	491	2	542
2.2.5-1	465	491	517	542
2.3.5-1	466	491	518	542
3.1.15-1	182	492	2	543
2.2.15-1	465	492	517	543
2.3.15-1	466	492	518	543
1.1.119-1	182	493	2	544
3.1.7-1	182	494	2	545
2.2.7-1	465	494	517	545
2.3.7-1	466	494	518	545
3.1.2-1	182	277	2	102
2.2.2-1	465	277	517	102
2.3.2-1	466	277	518	102
3.1.16-1	182	495	2	546
2.2.16-1	465	495	517	546

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
2.3.16-1	466	495	518	546
3.1.17-1	182	496	2	547
2.2.17-1	465	496	517	547
2.3.17-1	466	496	518	547
3.1.18-1	182	497	2	548
2.2.18-1	465	497	517	548
2.3.18-1	466	497	518	548
1.1.120-1	182	498	2	549
3.1.19-1	182	499	2	550
2.2.19-1	465	499	517	550
2.3.19-1	466	499	518	550
1.1.121-1	182	500	2	551
3.1.20-1	182	501	2	552
2.2.20-1	465	501	517	552
2.3.20-1	466	501	518	552
1.1.122-1	182	502	2	553
1.1.123-1	182	503	2	554
1.1.124-1	182	504	2	555
1.1.125-1	182	505	2	556
1.1.126-1	182	506	2	557
1.1.127-1	182	507	2	558
1.1.128-1	182	508	2	559
1.1.129-1	182	509	2	560
1.1.130-1	182	510	2	561
1.1.131-1	182	511	2	562
1.1.132-1	182	512	2	563
1.1.133-1	182	513	2	564
1.1.134-1	182	514	2	565
1.1.135-1	182	515	2	566
1.1.138-1	182	569	2	568
1.42.1-1	467	223	519	49

Table 13. SEQ ID NOs of VH, VL, heavy chains (HC) and light chains (LC) of anti-gp120 antibodies				
Antibody Name	VH	VL	HC	LC
1.43.1-1	468	223	520	49
1.44.1-1	469	223	521	49
1.45.1-1	470	223	522	49
1.46.1-1	471	223	523	49
1.47.1-1	472	223	524	49
1.49.1-1	474	223	526	49
1.50.1-1	475	223	527	49
1.51.1-1	476	223	528	49
1.1.104-1	182	516	2	567
3-1	182	276	2	101
1.52.1-1	477	223	529	49
1.52.64-1	477	278	529	103
1.52.90	477	292	529	117
2.4.1-1	478	276	530	101

Example 3: Mass Spectrometry Analysis

[0291] Antibody A-1 was transiently expressed in ExpiCHO cells and protein-A purified using standard methods. The sample was denatured and reduced by using 4 M guanidine hydrochloride and 50 mM DTT (final concentrations) and heating for 20 minutes at 60°C. The sample was desalted online as reduced heavy and light chains were separated on a BEH C4 reverse phase chromatography column prior to infusion into the source of a Waters Synapt G2Si hybrid time-of-flight mass spectrometer. Multiply-charged protein peak packets were deconvoluted used the Maximum Entropy deconvolution algorithm. Results show that the Antibody A light chain was glycosylated. The observed light chain mass spectrum reveals the presence of a G0-glycan modification with additional glycan-associated mass heterogeneity. This observation is consistent with the presence of an N72 consensus glycosylation motif in the Antibody A VL domain (NLT), and with previous crystal structures of Antibody A showing glycosylation at this position (Zhou et al., *Immunity*, 39:245-258 (2013); Klein et al., *Cell*, 153:126-138 (2013)).

Example 4: Accelerated Stress-Induced Potency Loss

[0292] To identify chemical liabilities, an accelerated thermal stability study (stress panel) was performed for A-1. The antibody was stressed at pH 5.9 at 25° C and 37° C (formulation like stress) and at pH 7.4 at 37° C (mock physiological-like stress). Samples were pulled and frozen at T0, in addition to 2, 4, and 6 weeks. Select samples were screened for stress-induced potency loss prior to implementation of other methods. The potency assay employed for the stressed A-1 samples was an ADCC reporter assay which uses a reporter cell that expresses luciferase when the FcγRIIIa receptors on its cell surface are tethered via a functional mAb's Fc and Fab domains to a target cell. The target cell in the assay expresses the HIV Env glycoprotein to which the A-1 Fab binds. Luciferase stoichiometrically converts excess luminescent substrate producing light measured in the assay. Response curves are indicative of antibody potency.

[0293] As shown in Figure 1, the most significant potency losses for A-1 occurred in the pH 5.9 conditions. We next conducted peptide mapping on the stress panel to identify the stress-induced chemical modification leading to loss of activity at pH 5.9. We additionally conducted peptide maps on the pH 7.4 stressed samples to identify modifications that might be prone to occur under physiological-like conditions.

[0294] The antibody A-1 stress panel samples were denatured, reduced, and alkylated with iodoacetamide prior digestion with the endoproteinase Lys-C. Protein digests were subsequently analyzed by reverse phase LC-MS/MS on a Thermo Q-Exactive HF mass spectrometer. Peptide maps were analyzed using Thermo Pepfinder and Xcalibur softwares, while ion lists were further analyzed in Microsoft Excel. Since our ADCC reporter data suggested the most significant potency losses at the pH 5.9 conditions we searched the ion lists for modifications occurring over time but that were unique to the pH 5.9 conditions. The most significant stress-induced, time dependent modification unique to the pH 5.9 conditions was oxidation of tryptophan 76 in the mAb heavy chain observed on the peptide T₅₅GQPNNPRQFQGRVSLTRHASWDFDTFSFYMDLK₈₈ (T55-K88) (SEQ ID NO: 630) as additions of oxygen (+15.99 Da) and further conversion to kynurenine (+3.99 Da). For relative quantification of these conversions the peak intensities from the Pepfinder ion list outputs for the two oxidized variants were summed and then compared to the sum of all modified and non-modified T55-K88 peptide peak intensities. The resultant summed oxidized peptide outputs for the various stress conditions are presented in Figure 2. It was on the basis of these studies that we

identified W74a (Kabat, FR3 insertion) oxidation as a potential risk to pharmaceutical stability of antibody A-1.

[0295] In addition to the significant oxidation at heavy chain W74a observed in pH 5.9 conditions, approximately 8-9% deamidation at light chain position N26 was observed on the constructs at T0 and increased further at pH 7.4 incubation conditions. The percentage of deamidation reported reflect the combination of asparagine deamidated to aspartic acid (+0.98 Da), isoaspartic acid (+0.98), and aspartyl succinimide (-17.03 Da) and were observed on light chain peptide
D₁IQMTQSPSSLSASVGDVTITCQANGYLNWYQRRGK₃₈ (SEQ ID NO: 631).
The results are depicted in Figure 3.

[0296] Although it is part of the antibody framework, heavy chain residue, W74a is found within an unusual framework insertion loop that forms part of the antibody paratope, and thus directly contacts the HIV gp120 (Lee *et al.* 2017. *Immunity* 46: 690-702). Light chain residue N26 is part of an NG deamidation risk motif in CDR1 that is formed by an unusual germline deletion in antibody A-1. Like W74a, N26 forms part of the paratope and is predicted to make contact with elements of HIV gp120. Based on available structural models, we next designed a panel of 15 mutants designed to remove the W74a oxidation site and the N26 deamidation motif. The mutations were screened in HIV neutralization assays (*see*, Example 10) to identify a variant that removed W74a, but had minimal impact on neutralization potency or breadth of antibody A-1.

Example 5: T-cell Epitope Mapping

[0297] To assess the immunogenicity and identify immunodominant T-cell epitopes, the Antitope Epi-Screen T-cell Epitope Mapping Assay was used to screen overlapping 15-mer peptides covering the entire Antibody A LC and HC Fv sequence. The background donor response (n=50 donors) of the assay was 8%, and responses >10% would be considered as positive in this assay. The T-cell epitope mapping results on Antibody A HC and LC identified a single peptide, GDTVITTCQANGYLN (SEQ ID NO: 320), containing a putative T-cell epitope- with a donor response rate of 18% in the Antibody A light chain.

[0298] Computational prediction of the core 9-mer using the antitope iTope algorithm identified VTITCQANG (SEQ ID NO: 321) as the potential MHCII binding 9-mer core within the peptide, with residue V19 being the P1 anchor position. The C-terminus of this epitope overlaps with non-germline residues in CDR L1 that are known to contact

the gp120 antigen as observed in co-crystal structures. To avoid disrupting antigen binding via germline reversion of CDR L1, this epitope was removed by introducing the LC V19A mutation at the P1 anchor position.

Example 6: Antibody Characterization

[0299] Scanning and combinatorial mutagenesis were used to generate additional antibodies to assess the biophysical and functional impact of germline reversions and glycan removal on Antibody A. Single point ELISA assays at the EC₅₀ concentration for Antibody A-1 were conducted for each of three unique HIV gp120 antigens in 384 well format and normalized to plate controls. DSF assays were conducted in parallel to assess the impact of mutations on Fab melting temperature (T_m). The results are shown in Table 14.

[0300] Table 14. Antibody characterization by ELISA and DSF

Antibody	ELISA gp120 Bal (A ₄₅₀)	ELISA gp120 SHIV SF162 P3 (A ₄₅₀)	ELISA gp120 CAAN (A ₄₅₀)	ELISA BSA (A ₄₅₀)	DSF Fab T _m (°C)
A-1	0.91	0.94	1.06	0.06	79.69
1.2.2-1	0.91	0.97	0.80	0.06	79.65
1.2.1-1	0.93	1.02	1.01	0.07	79.35
1.3.1-1	0.97	1.13	0.94	0.06	77.85
1.4.1-1	1.04	1.19	0.96	0.06	78.90
1.5.1-1	0.92	1.15	1.07	0.06	82.02
1.6.1-1	0.95	1.10	0.90	0.06	78.27
1.7.1-1	0.98	1.08	1.04	0.06	79.56
1.8.1-1	0.86	0.94	0.90	0.06	78.77
1.9.1-1	1.08	1.23	0.98	0.06	82.40
1.10.1-1	0.95	1.03	0.94	0.06	78.23
1.11.1-1	0.93	0.57	0.75	0.09	71.14
1.15.1-1	0.87	1.24	1.38	0.06	71.14
1.16.1-1	0.92	0.90	0.87	0.07	72.77
1.17.1-1	0.84	0.31	0.70	0.06	71.27

Antibody	ELISA gp120 Bal (A₄₅₀)	ELISA gp120 SHIV SF162 P3 (A₄₅₀)	ELISA gp120 CAAN (A₄₅₀)	ELISA BSA (A₄₅₀)	DSF Fab T_m (°C)
1.18.1-1	0.97	1.36	1.18	0.07	71.39
1.19.1-1	0.85	0.96	0.93	0.06	80.03
1.20.1-1	0.89	0.91	0.80	0.06	77.78
1.21.1-1	0.87	0.82	0.77	0.07	78.28
1.22.1-1	1.03	0.78	0.87	0.06	77.15
1v2-1	1.03	1.29	1.00	0.08	79.28
1.24.1-1	0.98	1.09	0.90	0.06	75.02
1.25.1-1	1.07	1.03	1.16	0.06	77.16
1.26.1-1	0.81	0.94	1.00	0.06	78.54
1.27.1-1	0.86	0.86	0.86	0.07	75.28
1.28.1-1	1.05	0.99	0.99	0.06	76.03
1.29.1-1	1.03	0.98	1.08	0.06	80.04
1.30.1-1	0.81	1.03	1.05	0.06	79.17
1.1.2-1	0.95	1.01	0.93	0.06	78.54
1.1.3-1	0.97	0.97	1.00	0.07	79.29
1.1.4-1	0.96	1.07	1.09	0.06	79.42
1.1.5-1	0.90	0.93	0.90	0.06	78.41
1.1.6-1	0.91	0.41	0.46	0.07	78.92
1.1.7-1	0.95	1.02	1.00	0.06	78.89
1.1.8-1	1.01	0.37	0.50	0.06	79.40
1.1.9-1	0.98	1.09	0.93	0.07	77.81
1.1.10-1	0.94	1.02	1.10	0.06	78.89
1.1.11-1	0.92	0.93	1.03	0.06	78.64
1.1.12-1	1.05	1.13	1.02	0.06	78.52
1.1.13-1	0.94	0.97	1.11	0.06	78.77

Antibody	ELISA gp120 Bal (A ₄₅₀)	ELISA gp120 SHIV SF162 P3 (A ₄₅₀)	ELISA gp120 CAAN (A ₄₅₀)	ELISA BSA (A ₄₅₀)	DSF Fab T _m (°C)
1.1.14-1	Low Yield/No Expression				
1.1.15-1	0.93	0.79	0.79	0.06	79.02
1.1.16-1	1.06	1.05	0.99	0.06	80.77
1.1.17-1	1.02	1.07	0.90	0.06	85.20
1.1.18-1	0.92	0.94	0.92	0.06	80.66
1.1.19-1	1.03	1.12	1.05	0.06	80.36
1.1.20-1	0.94	1.01	0.95	0.07	77.90
1.1.21-1	0.98	1.03	0.99	0.06	81.78
1.1.22-1	0.88	1.04	1.02	0.06	80.53
1.1.23-1	0.90	1.07	0.95	0.06	81.66
1.1.24-1	0.88	0.97	0.98	0.06	80.15
1.1.25-1	0.90	1.01	0.92	0.07	78.65
1.1.26-1	0.96	1.09	1.07	0.06	79.03
1.1.27-1	0.97	1.10	0.99	0.06	77.78
1.1.28-1	0.98	1.09	0.93	0.06	77.65
1.1.29-1	0.91	0.90	1.01	0.06	79.17
1.1.30-1	0.90	0.98	0.86	0.06	80.67
1.12.15-1	0.69	0.53	0.63	0.06	78.41
1.13.15-1	0.70	0.52	0.60	0.06	76.41
1.14.15-1	Low Yield/No Expression				
1.12.17-1	0.90	0.75	0.84	0.06	82.67
1.13.17-1	1.02	0.69	0.78	0.06	79.04
1.14.17-1	0.89	0.24	0.43	0.10	73.15
1.31.1-1	0.82	0.87	0.92	0.06	79.67
1.32.1-1	0.93	0.91	0.83	0.06	75.03
1.33.1-1	0.88	0.96	0.98	0.06	83.28

Antibody	ELISA gp120 Bal (A ₄₅₀)	ELISA gp120 SHIV SF162 P3 (A ₄₅₀)	ELISA gp120 CAAN (A ₄₅₀)	ELISA BSA (A ₄₅₀)	DSF Fab T _m (°C)
1.34.1-1	0.83	0.79	0.90	0.07	79.15
1.35.1-1	0.84	0.80	0.87	0.07	75.77
1.36.1-1	1.13	0.59	0.75	0.18	69.63
1.1.31-1	0.87	0.87	0.96	0.06	78.64
1.31.31-1	0.98	0.97	1.02	0.07	79.15
1.32.31-1	0.95	0.66	0.90	0.07	75.01
1.33.31-1	0.92	0.89	1.18	0.07	82.44
1.34.31-1	0.91	0.77	0.92	0.06	78.64
1.35.31-1	1.05	0.68	0.99	0.07	75.39
1.36.31-1	Low Yield/No Expression				
1.1.32-1	0.93	0.89	1.04	0.06	81.07
1.31.32-1	0.98	0.91	1.04	0.06	81.40
1.32.32-1	0.89	0.70	0.90	0.08	77.52
1.33.32-1	0.98	0.98	1.13	0.07	85.42
1.34.32-1	0.93	0.69	0.97	0.06	81.78
1.35.32-1	0.96	0.58	1.01	0.06	77.27
1.36.32-1	0.96	0.18	0.50	0.07	71.26
1.1.33-1	0.87	0.84	0.94	0.07	78.40
1.31.33-1	0.89	0.74	0.86	0.06	78.27
1.32.33-1	0.93	0.71	0.68	0.06	74.27
1.33.33-1	1.09	0.97	1.06	0.07	81.78
1.34.33-1	1.10	0.82	1.04	0.06	77.15
1.35.33-1	0.90	0.65	0.91	0.06	74.64
1.36.33-1	Low Yield/No Expression				
1.1.34-1	0.90	0.83	0.96	0.06	79.02
1.31.34-1	0.76	0.82	0.86	0.06	79.27

Antibody	ELISA gp120 Bal (A ₄₅₀)	ELISA gp120 SHIV SF162 P3 (A ₄₅₀)	ELISA gp120 CAAN (A ₄₅₀)	ELISA BSA (A ₄₅₀)	DSF Fab T _m (°C)
1.32.34-1	0.81	0.71	0.86	0.06	77.15
1.33.34-1	0.95	0.82	0.95	0.06	82.66
1.34.34-1	0.89	0.70	0.88	0.06	80.03
1.35.34-1	1.00	0.58	0.85	0.06	77.27
1.36.34-1	1.01	0.15	0.43	0.07	71.01
1.1.35-1	0.96	0.72	0.75	0.08	77.65
1.31.35-1	0.90	0.61	0.77	0.06	78.14
1.32.35-1	0.89	0.50	0.59	0.06	74.46
1.33.35-1	1.03	0.71	0.81	0.07	82.03
1.34.35-1	0.94	0.56	0.77	0.06	78.02
1.35.35-1	0.88	0.42	0.67	0.06	74.51
1.36.35-1	0.89	0.16	0.36	0.06	66.88
1.1.36-1	0.93	0.72	0.88	0.06	79.02
1.31.36-1	1.06	0.93	0.90	0.06	79.40
1.32.36-1	0.92	0.62	0.64	0.06	75.51
1.33.36-1	0.94	0.88	0.82	0.06	83.15
1.34.36-1	0.94	0.77	0.71	0.07	77.69
1.35.36-1	0.85	0.56	0.67	0.06	75.39
1.36.36-1	1.12	0.14	0.34	0.06	68.38

[0301] The results indicated that some germline reversions and combinatorial modifications affected gp120 binding and/or Fv thermal stability. Based on these data, multiple rounds of engineering were conducted. The V19A mutation (which may remove the predicted T-cell epitope shown above) and mutations made at light chain position N72 (Kabat numbering) (which may remove the N72-linked Fv glycan) were combined with other mutations in order to identify an antibody with improved functional

and biophysical properties. The resulting antibodies were characterized by expression titer analysis, polyspecificity analysis, and/or HIV neutralization assays.

Example 7: Expression Titer Analysis of Antibodies Without the Glycosylation Motif

[0302] When expressing and purifying protein for the ELISA and DSF screening campaign, reduced expression titer was observed for antibodies lacking the glycosylation motif. Further mutations were generated to identify antibodies with improved protein expression.

[0303] The antibodies were expressed in Expi293F™ cells using Expifectamine™293 system following manufacturer's protocol (ThermoFisher Scientific, MA). Transfection was carried out in 30 ml scale in 50 ml SeptaVent™ (Optimum Processing, CA). Briefly, 30 µg total of heavy and light chain (ratio of HC:LC is 2:3) expressing plasmids were used per transfection. Diluted DNA in OptiMEM was added to diluted Expifectamine™293 reagent to allow complex formation. After 20 minutes incubation at room temperature, the reagent DNA complex was added to 28 mL of cells seeded at 2.5 million/mL. Culture was incubated at 37°C in 8% CO₂ with shaking at 250 rpm for four days. Clarified supernatant was harvested by centrifugation at 500x g for 15mins. Antibodies were purified by Hamilton STAR Liquid handler (Hamilton, NV) using Phytips (PhyNexus, CA) pre-packed with 160µL MaSelectSuRe resin (GE Healthcare, NJ). Each of the 30 mL transfected volume was purified using 3 Phytips. After capture of the antibody, the resin was washed with 1 x PBS prior to elution with 100mM NaAcetate pH3.5. The eluted sample was neutralized with 1/10th volume of 1M Tris pH8.0. Samples were stored at 4°C overnight. The elution plate was centrifuged at 1000x g for 10 minutes to remove precipitate if any. Concentration of the clarified elution was determined by measuring its absorbance at A280. Titer of each of the antibodies is expressed as follows (mg/L): [concentration (mg/mL) x volume of elution (mL) * 1000]/ 30 mL. Glycosylation site mutations and expression titer are summarized in Table 15.

[0304] Table 15. Glycosylation site mutations and expression titer

Antibody	Titer (mg/L)
A-1	243
1.1.10-1	148

Antibody	Titer (mg/L)
1.1.37-1	104
1.1.38-1	113
1.1.39-1	93
1.1.40-1	133
1.1.41-1	104
1.1.42-1	158
1.1.43-1	124
1.1.44-1	70
1.1.45-1	77
1.1.46-1	136
1.1.47-1	45
1.1.48-1	65
1.1.49-1	27

[0305] The results in Table 15 show that all antibodies lacking the “NLT” glycosylation consensus motif exhibited reduced expression titer. This suggests that removal of the N72-linked glycan may have a negative effect on protein expression. The results also show that the L73F germline reversion, systematically reduces expression titer. Among the mutations tested, N72H, N72T and T74K had the highest expression titer and were carried forward for further analysis.

Example 8: Mammalian Display

[0306] To identify mutations that eliminate the Fab glycan while maintaining binding to HIV Env, improve expression titer, and/or reduce polyspecificity, a combinatorial light chain mutation library was designed and constructed using a set of trimer oligoes (GenScript) varied at 6 sites, including R65, W67, E70, N72, L73, and T74. The synthesized light chain library harboring ~18,000 antibodies was sub-cloned into a modified pcDNA5/FRT vector (Invitrogen), containing the Antibody A heavy chain fused with a human PDGFR transmembrane domain at the C-terminus.

[0307] To display the antibodies in stably transfected cells, the constructed expression vector was co-transfected with pOG44 to Flp-In -CHO cells following the manufacturer's instructions (R758-07, Invitrogen). The transfected cells were selected and then maintained in hygromycin supplemented culture media. Antibody display and binding to HIV Env were analyzed by FACS following anti-human IgG (Fcγ specific) and HIV BG505.SOSSIP (*J. Virol.*, 89(10):5318-29 (2015)) staining. Cells collected after FACS sorting were expanded for DNA extraction and subsequent PCR-sequencing analysis to identify recovered mutations. More than one hundred clones were picked for sequencing before and after FACS sorting. The sequences recovered from two consecutive rounds of FACS sorts were next examined.

[0308] Results showed that antibodies with a sequence of TRRGQQYNLT (SEQ ID NO: 332), RRWGQNYNFT (SEQ ID NO: 333), TRRGQDYIFS (SEQ ID NO: 334), RRRGQDYILA (SEQ ID NO: 335), RRRGQNYTFT (SEQ ID NO: 336), RRRGQDYILT (SEQ ID NO: 337), TRFGQNYSLQ (SEQ ID NO: 338), or TRRGQNYTLA (SEQ ID NO: 339), TRRGQQYTLP (SEQ ID NO: 340), TRRGQDYILA (SEQ ID NO: 341), or SRFGQKYQLS (SEQ ID NO: 342) in the LC FR3 region had desirable expression levels and retain binding affinity to HIV BG505.SOSSIP. The mutations in SEQ ID NO:334, SEQ ID NO:337 and SEQ ID NO: 342 were incorporated into Antibodies 1.1.110-1, 1.1.111-1, 1.1.113-1, 2.1.3-1, 2.1.4-1 and 1.1.112-1.

Example 9: Polyspecificity Assessment

[0309] Polyspecificity of therapeutic antibodies may adversely affect pharmacokinetic properties and present potential safety concerns. It has been shown that Antibody A was polyreactive to double-stranded DNA and lipopolysaccharide in a four-antigen panel ELISA assay (*Science*, 333(6049):1633–1637 (2011)). The polyspecificity risk of antibodies evaluated herein were tested in multiple assays including anti-nuclear antibody (*Genes Immun.*, 13(5): 399–410 (2012)), anti-cardiolipin (*Hum Antibodies*, 14(3-4): 59–67 (2005)), anti-baculoviral particle ELISA (*Proc. Natl. Acad. Sci. USA*, 114(5):944-949 (2017)), and FACS-based HEK-293 and HEp2 cell binding assays (*J. Virol.*, 88(21):12669-82 (2014)). To compare polyspecificity, Antibody C and Antibody D, two polyspecific bNAbs (*J. Virol.*, 88(21):12669-82 (2014)), were used as positive controls; and a clinical sample of Rituximab (Myoderm Medical Supply) was used as a benchmark for low risk of polyspecificity. The tested articles were diluted to 1 μM in ELISA assays, and OD₄₅₀ values were normalized to control (no antibody) to calculate fold

change. In cell binding assays, HEK293 or HEp2 cells were permeabilized and then incubated with serially diluted tested articles. The stained samples were FACS analyzed, and MFI (mean fluorescence intensity) was normalized to anti-human IgG-Fc γ secondary antibody only stained control. The relative binding signals were plotted against antibody concentrations, and fitted to non-linear response curve. Non-specific cell binding of each tested antibody was represented by binding AUC (area under curve).

[0310] Three single mutants with the N72 glycan removed (via point mutagenesis) show the highest expression titer (Table 5). In order to evaluate their contributions to A-1 polyspecificity, the mutants were tested in anti-nuclear antibody (ANA) and anti-cardiolipin ELISA assays as described above. The results of two independent assays are shown in Table 16. These results suggest that removal of the N72 glycan may lead to increases in polyspecificity. Among mutations tested, the N72T and N72H mutation show the lowest polyspecificity scores.

[0311] Table 16: Polyspecificity Assessment

mAb (1 μ M)	ANA		Anti-Cardiolipin	
	(Normalized OD ₄₅₀)		(Normalized OD ₄₅₀)	
C	24.5	26.5	15.6	17.2
A-1	2.3	2.6	1.6	1.5
1.1.10-1	8.3	8.2	2.5	2.2
1.1.42-1	4.0	3.7	2.3	2.0
1.1.46-1	11.8	10.1	4.4	3.7
Rituximab	1.5		1.3	

[0312] Antibodies with N72T, V19A and other mutations selected based on the functional analysis presented in Table 14 and Table 23 were tested in ANA and anti-cardiolipin ELISA assays for polyspecificity assessment. The results of these analyses are shown in Table 17.

[0313] Table 17. Polyspecificity Assessment of antibodies with N72T Mutation

mAb (1 μ M)	ANA		Anti-Cardiolipin	
	(Normalized OD ₄₅₀)		(Normalized OD ₄₅₀)	
A	1.9		1.2	
A-1	2.7		1.4	

mAb (1 μM)	ANA (Normalized OD₄₅₀)	Anti-Cardiolipin (Normalized OD₄₅₀)
1.1.10-1	5.7	1.7
1.33.32-1	4.9	7.7
1.1.54-1	3.0	1.6
1.37.51-1	4.0	6.2
1.8.52-1	5.6	1.7
C	22.5	4.4
D-1	14.6	7.7
Rituximab	1.3	1.0

[0314] The results in Table 17 show that all antibodies lacking the N72 glycan exhibited increased polyspecificity compared to Antibody A-1. Antibody 1.1.54, which contains the N72T and the V19A mutation, exhibited reduced polyspecificity compared to Antibody 1.1.10, which contained the N72T mutation alone. This suggests that the V19A mutation, which was introduced to remove a T-cell epitope, may have unexpected benefits in reducing the polyspecificity of the antibodies disclosed herein.

[0315] In order to identify antibodies with decreased polyspecificity, a 32-member combinatorial panel comprised of 5 sets of mutations to Antibody A-1 (Table 18) was tested in ANA, anti-cardiolipin ELISA, HEK293 and HEp2 binding assays as described above.

[0316] Table 18. Mutations Used to Generate a 32 Member Combinatorial Library

Mutation Set	HC mutations	LC mutations
Set 1	None	V19A
Set 2	None	N72H
Set 3	None	V98F, V99G
Set 4	None	T18R, R65S, N76S, N77S
Set 5	L5V, A10E, T12K, E23K, S105Q, Q108M	R39K, R40P

[0317] The results of the assays were summarized and compared using polyspecificity scores (P-scores) that were calculated as ratio of each tested antibody to rituximab in each assay (Table 19). The average P-score values were used to rank the risk of polyspecificity of the tested antibodies. To statistically analyze the contribution of each mutation in the combinatorial dataset, pairwise comparisons were done for each

combinatorial antibody in the presence or absence of the mutation sets listed in Table 19.

In the context of the 32 member combinatorial antibody panel tested herein, 16 independent comparisons were conducted for each of the five mutation sets tested.

[0318] Table 19. Polyspecificity Scores (P-score) of Combinatorial Antibodies

mAb	ANA		Anti-Cardiolipin		HEK293 Binding		HEp2 Binding	Mean	stdev
A-1	2.8	3.2	1.3	1.5	3.7	2.2	3.5	2.6	1
1.1.17-1	1.8	1.8	1.2	1.1	2.5	1.6	2.7	1.8	0.5
1.1.42-1	n/a	4.3	n/a	1.9	6	n/a	5	4.3	1.5
1.1.64-1	2	1.9	1.2	1.1	2.4	1.6	2.5	1.8	0.5
1.1.67-1	2.3	2.5	1.2	1.4	2.8	1.9	2.6	2.1	0.6
1.1.72-1	n/a	4.7	n/a	1.4	4	n/a	3.9	3.5	1.2
1.1.75-1	3.2	3.3	1.7	1.7	3.8	2.3	4.4	2.9	1
1.1.78-1	3.4	3.5	1.8	1.4	2.9	2.6	3.4	2.7	0.8
1.41.5-1	n/a	5	n/a	3.7	7.2	n/a	4.9	5.2	1.3
1.41.81-1	4.3	4.5	2.7	3.4	6.1	2	4.6	3.9	1.3
1.1.82-1	3	2.9	1.2	1.2	3.7	1.6	2.9	2.4	0.9
1.41.83-1	4.2	4.5	4.1	5.3	4.9	2.1	3.9	4.1	0.9
1.1.84-1	2.2	2.4	1.3	1.2	2.3	1.7	2	1.9	0.5
1.41.85-1	3.3	3.5	3.3	3.2	3.7	1.6	3.2	3.1	0.6
1.41.86-1	5.1	6.2	3	3.8	4.1	1.8	4	4	1.3
1.41.87-1	3.3	3.4	3.5	3.7	3.2	1.9	2.9	3.1	0.6
1.1.88-1	1.6	1.8	1.2	1.2	3.7	1.5	2.4	1.9	0.8
1.41.89-1	1.9	1.9	1.9	2.2	6.3	1.7	4.4	2.9	1.6
1.1.90-1	1.5	1.2	1.2	0.9	3.7	1	2.3	1.7	0.9
1.41.91-1	2.9	3.1	3.6	6.2	2.5	2.1	2.9	3.3	1.2
1.41.92-1	n/a	15.7	n/a	11.2	5.8	n/a	5.4	9.5	4.2
1.41.93-1	n/a	9.5	n/a	9.8	5.1	n/a	5.4	7.4	2.2
1.1.94-1	6.2	7.7	4.7	4.9	3.4	3.2	3.3	4.8	1.6
1.41.95-1	10.2	14.8	8.9	9.2	6.5	3.8	5.4	8.4	3.4
1.1.96-1	4.7	6.1	2.4	2.7	2.4	2.8	2.8	3.4	1.3
1.41.97-1	5.4	6.5	5.4	7.3	4.5	3.9	4.1	5.3	1.2
1.41.98-1	5.9	8	4.6	5.9	3.9	3.3	4	5.1	1.5
1.41.99-1	6.4	6.9	6.1	9.2	7.8	3.2	4.6	6.3	1.9
1.1.100-1	4.6	4.2	2	2.1	4.7	1.7	2.7	3.1	1.2
1.41.101-1	5.6	7.2	5.5	6.8	8.7	3.2	4.8	6	1.7
1.1.102-1	3.3	4	1.8	1.7	4.5	2.8	2.6	3	1
1.41.103-1	3.6	4.6	5.9	5.3	6.8	4.4	4	5	1
C	17.6	14.7	3.8	8.3	13.1	8.6	6.9	10.4	4.5
D-1	8.7	10.4	1.9	3.3	7.3	7.1	4	6.1	2.8
Rituximab	1	1	1	1	1	1	1.1	1	0

[0319] For each of the 16 pairwise combinations, the average P-score across the seven assays shown in Table 19 was compared using a paired T-test. The results showed an increase in polyspecificity due to introduction of the light chain N72H mutation as well as due to the introduction of the Set 5 mutations. The results showed a decrease in polyspecificity due to introduction of the light chain V19A or the V98F+V99G mutations. A modest but not statistically significant decrease in polyspecificity was observed upon introduction of the Set 4 mutations. Consistent with this statistical analysis, the antibody with the lowest average polyspecificity score was Antibody 1.1.90, which incorporated the V19A mutation, the V98F+V99G mutations, and the Set 4 mutations.

[0320] Antibodies A-1 and B-1 were next compared in polyspecificity assays. Additionally, antibodies with the following mutations were tested in various combinations: N72T, N72H, V19A, V98F+V99G, the Set 4 mutations, or the mutations identified in SEQ ID NO: 37. The antibodies were tested in baculoviral particle (BVP) ELISA and the results are summarized in Table 20. Test articles were assayed at 1 μ M concentration in duplicate in each experiment and the BVP score was calculated as a ratio of OD₄₅₀ to no mAb background.

[0321] Table 20. BVP Scores

mAb	mean	SD	n
A	10.9	1.1	2
A-1	9.4	0.7	6
1.1.10-1	13.9	n/a	1
1.1.42-1	9.7	1.1	2
1.1.111-1	22.1	2.0	4
1.1.113-1	11.9	n/a	1
1.1.90-1	2.5	0.6	2
B-1	2.7	0.9	3
2.1.2-1	47.1	n/a	1
2.1.3-1	8.2	n/a	1
2.1.4-1	7.7	n/a	1
D-1	29.2	9.7	4
C	44.2	10.2	4
Rituximab	4.0	0.5	4

[0322] The results in Table 20 show that Antibody B-1 exhibited reduced polyspecificity compared to Antibody A-1. Like Antibody A, removal of the N72 glycan using the N72H mutation in Antibody 1.1.42 or Antibody 2.1.2 resulted in an increase in

polyspecificity. Incorporation of the mutations discovered via mammalian display into Antibody 1.1.111 may increase polyspecificity, while incorporation of the same mutations into Antibody 2.1.3 may reduce polyspecificity compared to the N72H mutation. Adding the V19A mutation (*e.g.* Antibody 1.1.113 or Antibody 2.1.4) may systematically lower the polyspecificity in both cases.

[0323] An additional panel of 96 antibodies was generated to identify antibodies with improved neutralization breadth and potency, and ideally lacking the N72 linked glycan. This panel tested the effects of the set 1, 3 and 4 mutations (Table 18) as well as various N72 mutations and mutations derived from mammalian display in the context of antibody variable domains derived from Antibody A or combining elements of both Antibody A and Antibody B. The library also included a scanning mutagenesis campaign, where each amino acid differing between Antibody A and Antibody B was tested individually in the context of Antibody A. The panel was produced using high throughput methods and assayed using a high throughput BVP ELISA normalized for variations in sample concentration. The results of this assay are shown in Table 21.

[0324] Table 21: BVP Scores

mAb	Normalized BVP Score
A-1	5.5
3-1	3.7
2.2.1-1	5.5
2.3.1-1	3.5
1.1.64-1	7.2
3.1.8-1	3.8
2.2.8-1	13.7
2.3.8-1	10.8
1.1.67-1	5.0
3.1.9-1	8.2
2.2.9-1	15.7
2.3.9-1	9.0
1.1.115-1	15.0
3.1.10-1	6.0
2.2.10-1	25.8
2.3.10-1	13.6
1.1.116-1	27.1
3.1.11-1	14.6
2.2.11-1	34.4
2.3.11-1	19.1
1.1.117-1	18.7
3.1.12-1	15.2

mAb	Normalized BVP Score
2.2.12-1	35.7
2.3.12-1	24.3
1.1.118-1	13.1
3.1.13-1	19.4
2.2.13-1	15.2
2.3.13-1	12.5
1.1.10-1	10.8
3.1.14-1	12.5
2.2.14-1	26.4
2.3.14-1	6.0
1.1.54-1	8.3
3.1.5-1	7.0
2.2.5-1	13.2
2.3.5-1	6.3
1.1.104-1	7.8
3.1.15-1	8.1
2.2.15-1	25.3
2.3.15-1	12.2
1.1.119-1	12.9
3.1.7-1	8.4
2.2.7-1	15.5
2.3.7-1	9.1
1.1.42-1	7.7
3.1.2-1	7.4
2.2.2-1	26.0
2.3.2-1	11.3
1.1.75-1	14.6
3.1.16-1	13.4
2.2.16-1	33.9
2.3.16-1	18.0
1.1.78-1	10.4
3.1.17-1	14.4
2.2.17-1	39.6
2.3.17-1	25.6
1.1.102-1	7.4
3.1.18-1	11.9
2.2.18-1	26.5
2.3.18-1	15.4
1.1.120-1	14.2
3.1.19-1	14.1
2.2.19-1	32.1
2.3.19-1	24.4
1.1.121-1	8.4
3.1.20-1	6.7

mAb	Normalized BVP Score
2.2.20-1	22.7
2.3.20-1	13.2
1.1.122-1	12.5
1.1.123-1	10.0
1.1.124-1	8.9
1.1.125-1	8.6
1.1.126-1	9.8
1.1.127-1	13.9
1.1.128-1	12.0
1.1.129-1	16.8
1.1.130-1	13.8
1.1.131-1	8.3
1.1.111-1	17.1
B-1	7.0
1.1.90-1	7.2
1.1.132-1	12.3
1.1.133-1	12.1
1.1.134-1	7.1
1.1.135-1	11.3
1.42.1-1	13.9
1.43.1-1	12.6
1.44.1-1	22.2
1.45.1-1	1.8
1.46.1-1	10.0
1.47.1-1	8.8
1.49.1-1	11.9
1.50.1-1	8.0
1.51.1-1	8.1
A-1	8.1

[0325] The results for two separate production runs of Antibody A-1, shown in the first and last rows of Table 21, had BVP scores between 5.5 and 8.1. Antibodies incorporating mutations at N72 had systematically higher BVP scores, which was consistent with the above results demonstrating that removing the N72 linked glycan may lead to increased polyspecificity. Selected antibodies lacking N72 and incorporating the Antibody B light chain or heavy chains (or mutants derived from these chains), such as Antibodies 3.1.10-1, 2.3.14-1, 1.1.54-1, 3.1.5-1, and 2.3.5-1, did not show increased BVP ELISA scores compared to Antibody A-1.

[0326] An additional panel of 12 antibodies with N72 linked glycan or lacking the glycan was produced to further evaluate the role of the glycan in polyspecificity. Some

antibodies were produced in both Expi293 and CHO-S cells. Mutations that may decrease polyspecificity, identified in the above assays, were incorporated into this panel. The results of this assay are shown in Table 22. Antibodies retaining the light chain N72 linked glycosylation motif had relatively lower BVP scores than antibodies lacking the N72 linked glycosylation motif.

[0327] Table 22: BVP Scores. N $\geq$ 3 for each antibody

N72 linked glycan	Name	Cell line	BVP Score
Yes	A-1	Expi293	2.2
		CHO-S	7.7
	1.1.64-1	Expi293	2.0
		CHO-S	6.6
	1.52.64-1	Expi293	2.4
		CHO-S	2.6
	1.1.90-1	Expi293	1.7
	2.2.101	Expi293	12.8
No	2.4.1-1	Expi293	5.2
	2.3.1-1	Expi293	5.3
	1.1.104-1	ExpiCHO	8.8
	1.1.119-1	ExpiCHO	6.0
	3.1.5-1	ExpiCHO	3.3
		CHO-S	12.0
	2.2.5-1	Expi293	32.5
	2.3.5-1	Expi293	3.6

Example 10: HIV Neutralization Assay

[0328] To assess the breadth of antigen recognition for antibodies, HIV neutralization assays were conducted using a variety of virus isolates and clones. HIV neutralization potency (expressed as IC₅₀ in μ g/mL) of the antibodies were measured in the CEM-NKr-CCR5-Luc reporter cell based assay (Trkola et al., (1999), J. Virol., 73(11):8966-74) against a panel of replication competent subtype B viruses that included isolates and clones amplified from patient plasma samples (NIH AIDS Reagent Program) and the lab adapted strain HIV-1 BaL.

[0329] Table 23. HIV Neutralization Potency

Antibody	<i>Virus Neutralization Potency (µg/mL)</i>			
	CHO77	Bal	92US657	8320
A-1	0.12	0.16	1.31	0.31
1.2.2-1	0.12	0.23	1.38	0.70
1.3.1-1	0.08	0.26	1.84	0.37
1.4.1-1	0.04	0.21	1.35	0.28
1.5.1-1	0.14	0.26	0.85	0.50
1.6.1-1	0.10	0.39	1.32	0.62
1.7.1-1	0.10	0.51	1.10	0.54
1.8.1-1	0.09	0.04	1.88	0.59
1.9.1-1	0.17	0.20	1.39	0.50
1.15.1-1	0.10	0.17	1.42	0.31
1.18.1-1	0.10	0.12	1.78	0.33
1.21.1-1	0.17	0.11	2.51	0.59
1.22.1-1	0.95	>20	11.3	13.2
1v2-1	0.07	0.08	0.88	0.28
1.25.1-1	0.12	0.17	1.46	0.23
1.26.1-1	0.11	0.06	0.93	0.38
1.27.1-1	0.13	0.54	0.56	0.58
1.28.1-1	0.10	0.11	1.63	0.38
1.29.1-1	0.09	0.14	1.33	0.41
1.30.1-1	0.06	0.15	0.93	0.39
1.1.2-1	0.14	0.20	0.62	0.78
1.1.4-1	0.12	0.10	1.44	0.35
1.1.5-1	0.12	0.21	1.93	0.63
1.1.10-1	0.07	0.10	0.60	0.33
1.1.11-1	0.13	0.09	1.17	0.39
1.1.12-1	0.12	0.06	1.64	0.50
1.1.13-1	0.09	0.19	1.43	0.45
1.1.17-1	0.10	0.08	1.27	0.52
1.1.19-1	0.12	0.09	0.90	0.34
1.1.26-1	0.13	0.07	1.49	0.45
1.1.27-1	0.10	0.12	1.25	0.53
1.14.15-1	>20	>20	>20	>20
1.33.1-1	0.17	0.11	0.95	0.77
1.33.32-1	0.08	0.09	0.83	0.35
1.34.32-1	0.11	0.31	1.47	0.54
1.36.35-1	0.32	>20	5.37	3.82
1.36.36-1	0.30	>20	2.07	3.84

[0330] Some antibodies displayed no loss of function in the ELISA assays (Table 14), but exhibited reduced potency in HIV neutralization assays (Table 23). Several antibodies showed either no change in virus neutralization activity or exhibited small gains in neutralization potency.

[0331] Table 24. HIV Neutralization Potency on Antibodies Incorporating Mutations

Antibody	<i>Virus Neutralization Potency (ug/mL)</i>					
	BaL	92US727	92HT593	92US657	92US712	302076
A	0.063	7.81	0.019	2.63	0.104	0.183
A-1	0.037	6.08	0.013	2.160	0.085	0.145
1.1.10-1	0.021	>20	0.013	1.83	0.088	0.131
1.1.42-1	0.030	>20	0.046	2.14	0.078	0.157
1.33.32-1	0.063	>20	0.010	1.39	0.062	0.077
1.1.54-1	0.062	11.4	0.011	1.50	0.076	0.114
1.37.51-1	0.053	>20	0.014	2.58	0.149	0.112
1.8.52-1	0.076	>20	0.015	2.81	0.141	0.202

[0332] As shown in Table 24, antibodies lacking the N72 glycan exhibited reduced potency in neutralizing the 92US727 virus. Antibody 1.1.54-1 (V19A +N72T) showed increased neutralization potency for the 92US727 virus compared to Antibody 1.1.10-1 (which contained N72T). This suggests that, combined with the N72T mutation, V19A may reduce polyspecificity and improve neutralization potency on select viruses.

[0333] Table 25. HIV Neutralization Potency on Antibodies Using an Expanded Panel of Viruses

Virus	<i>Virus Neutralization Potency (ug/mL)</i>			
	A-1	1.1.54-1	1.37.51-1	1.1.42-1
7467	0.07	0.06	0.06	0.07
302076	0.10	0.08	0.06	0.09
CH058	0.10	0.14	0.09	0.1
92US712	0.11	0.09	0.13	0.09
92HT593	0.12	0.06	0.12	0.05
7015	0.14	0.14	0.10	0.11
BaL	0.15	0.13	0.82	0.07
RHPA	0.16	0.16	0.09	0.15
1489	0.16	0.22	0.26	0.21
WITO	0.26	0.22	0.11	0.07
8176	0.28	0.18	0.18	0.16
8318	0.38	0.31	0.37	0.24
7576	0.46	0.32	0.34	0.4
8339	0.52	0.32	0.29	0.39
7051	0.61	0.52	0.49	0.29
8089	0.67	0.57	1.14	0.73
8106	1.03	0.82	0.97	0.92

Virus	<i>Virus Neutralization Potency (ug/mL)</i>			
	A-1	1.1.54-1	1.37.51-1	1.1.42-1
8359	1.56	1.19	1.51	1.2
92US657	1.81	3.45	3.39	2.97
92US727	1.82	9.65	50.33	33.7
8117	1.93	1.12	0.92	1.17
CH077	2.45	1.63	2.29	1.80
CH106	2.71	3.05	2.8	1.30
REJO	2.88	1.93	3.04	2.82
THRO	3.1	1.81	1.28	2.33
1413	3.84	2.8	2.45	2.88
8320	4.07	2.61	4.09	2.55
7103	4.82	3.03	2.28	3.23
8134	5.08	7.19	6.97	8.71
7141	5.35	8.93	51.0	23.2
8110	7.09	5.29	4.14	5.97
7714	8.96	6.78	4.33	8.71
1003	23.0	15.35	12.6	17.7
7595	24.6	18.9	16.17	18.02
8339	>200	>200	>200	>200
8398	>200	>200	>200	>200
7406	>200	>200	>200	>200
7552	>200	>200	>200	>200
7007	>200	>200	>200	>200

[0334] Among antibodies profiled in this assay, Antibody 1.1.54 containing the N72T and the V19A mutations, exhibited the highest neutralization potency (Table 25).

[0335] Table 26. HIV Neutralization Potency of Select Antibodies

Virus	A-1	1.1.110-1	1.1.111-1	1.1.112-1	1.1.90-1	1.1.64-1
CH106	1.29	1.33	1.42	1.31	1.69	0.99
1413	2.04	3.69	2.79	2.05	3.55	2.35
BaL	0.05	0.11	0.13	0.12	0.13	0.08
92HT593	0.34	0.50	0.52	0.56	0.68	0.35
92US657	0.70	0.69	0.79	0.74	1.17	0.79
302076	0.13	0.25	0.18	0.28	0.16	0.15
7015	0.68	0.77	0.73	0.75	1.03	0.60
8117	1.13	1.35	1.09	1.45	1.14	1.13
8339	>50	>50	>50	31.4	>50	>50
92US727	1.62	>50	35.19	30.3	3.80	0.79

Virus	A-1	1.1.110-1	1.1.111-1	1.1.112-1	1.1.90-1	1.1.64-1
92US712	0.08	0.06	0.06		0.07	0.06
7141	2.10	>50	27.09		2.41	1.99
7007	>50	>50	>50		>50	>50
8134	0.36	1.29	0.87		1.05	0.29
8318	0.15	0.23	0.19		0.21	0.23
7596	15.9	14.9	11.6		11.54	16.19
7103	0.95	1.00	0.69		1.13	1.01
8359	0.40	0.35	0.41		0.58	0.48
8110	2.63	3.85	2.19		1.65	2.62
7595	4.38	3.97	3.59		3.71	4.48
REJO	0.03	0.03	0.05		0.07	0.04
CH058	0.04	0.04	0.04		0.05	0.03
CH077	0.03	0.03	0.05		0.10	0.03
RHPA	0.19	0.20	0.16		0.16	0.09
WITO	0.12	0.08	0.06		0.07	0.13
7714	4.20	2.65	3.78		3.76	7.48
THRO	6.84	6.77	6.20		7.42	9.60
CH040	4.86	6.86	16.1		9.57	18.59
8089	0.32	1.30	0.60		0.47	0.32
1489	0.09	0.08	0.10		0.24	0.13
7467	0.03	0.03	0.04		0.06	0.03
7352	>50	>50	>50		>50	40.1
8320	0.28	0.26	0.25		0.39	0.37
1003	1.25	1.35	1.17		1.92	1.25
1012	1.24	1.65	1.43		1.40	1.40
8398	>50	>50	>50		>50	>50
7576	0.38	0.50	0.32		0.36	0.41
7051	0.21	0.25	0.19		0.19	0.23
8176	0.18	0.18	0.19		0.24	0.24
8106	0.57	0.61	0.61		0.89	1.08
7552	27.24	>50	>50		>50	16.1

[0336] HIV neutralization potency was tested on select antibodies identified via mammalian display (Antibodies 1.1.110, 1.1.111 and 1.1.112) and those that showed reduced polyspecificity (Antibodies 1.1.90 and 1.1.64). Loss of potency was observed against viruses 92US727 and 7141 for the antibodies identified via mammalian display (Table 26).

[0337] Table 27. HIV Neutralization Results for Select Antibodies

Virus	B-1	2.1.3-1	2.1.4-1	A-1	1.1.54-1	1.1.90-1	1.1.111-1	1.1.113-1
92US712	0.10	0.20	0.11	0.10	0.07	0.09	0.08	0.07
8318	0.11	0.24	0.21	0.11	0.32	0.57	0.48	0.14
92HT593	0.35	0.30	0.33	0.41	0.26	0.39	0.29	0.24
RHPA	0.09	0.07	0.05	0.12	0.03	0.15	0.08	0.05
7051	0.41	0.42	0.46	0.52	0.23	0.44	0.32	0.41
8320	0.27	0.28	0.23	0.42	0.29	0.34	0.30	0.28
7576	0.24	0.49	0.55	0.34	0.48	0.41	0.26	0.28
7015	1.12	0.89	0.59	1.17	0.40	0.41	0.48	0.44
92US657	0.53	0.42	0.38	0.52	0.45	0.48	0.40	0.40
CH106	1.51	1.01	0.91	0.91	0.90	1.00	0.86	0.59
1413	2.51	2.78	2.16	2.17	1.72	1.94	2.33	3.67
7141	1.67	13.76	6.78	2.45	1.61	2.24	29.81	27.85
92US727	2.13	14.34	7.27	2.01	11.18	2.17	>50	>50
7595	1.24	1.35	1.19	1.73	1.62	1.76	1.31	1.78
CH040	0.14	0.47	0.36	0.57	>50	22.06	3.71	0.15
THRO	5.32	6.56	4.97	6.30	5.47	10.62	8.58	9.39
7552	26.02	>50	>50	16.55	6.22	27.82	>50	>50
8339	>50	>50	>50	>50	>50	>50	>50	>50
7007	>50	>50	>50	>50	>50	>50	>50	>50
8398	>50	>50	>50	>50	>50	>50	>50	>50

[0338] The HIV neutralization results shown in in Table 27 suggest that removal of the N72 glycan (2.1.3-1, 2.1.4-1, 1.1.54-1, 1.1.111-1, and 1.1.113-1) may result in loss of neutralization sensitivity for select viruses (*i.e.*, 7141, 92US727) compared to antibodies retaining the N72 glycan (Antibodies B-1, A-1, 1.1.90-1).

[0339] The antibodies tested in the high throughput BVP ELISA shown in Table 21 were examined in an HIV neutralization assay against 4 viruses. The results of this study are shown in Table 28.

[0340] Table 28. HIV Neutralization Results

mAb	Neutralization IC ₅₀ Values (μg/mL) for the Virus Indicated			
	92US712	92US727	7141	VS001
A-1	0.13	3.8	3	0.18
3-1	0.11	3.92	4.03	0.1
2.2.1-1	0.05	~6-11	1.61	0.1
2.3.1-1	0.13	~4-12	2.65	0.2
1.1.64-1	0.05	0.55	0.64	0.05
3.1.8-1	0.12	1.15	3.51	0.11
2.2.8-1	0.07	3.38	1.6	0.07
2.3.8-1	0.07	1.72	1.64	0.15
1.1.67-1	0.1	8.1	3.76	0.15
3.1.9-1	0.11	3.54	4.35	0.16
2.2.9-1	0.06	13.1	1.75	0.15
2.3.9-1	0.09	11.1	2.56	0.22
1.1.115-1	0.08	>50	10.1	0.75
3.1.10-1	0.34	>50	44.8	2.24
2.2.10-1	0.05	>50	4.31	0.79
2.3.10-1	0.09	>50	15.3	0.94
1.1.116-1	0.12	>36	14.6	0.44
3.1.11-1	0.07	>44	8.7	0.57
2.2.11-1	0.04	>50	2.65	0.4
2.3.11-1	0.07	>50	11.2	0.83
1.1.117-1	0.09	>50	16.9	3.03
3.1.12-1	0.11	>41	24.5	2.41
2.2.12-1	0.04	>50	7.08	2.1
2.3.12-1	0.09	>50	23.8	1.92
1.1.118-1	0.18	>50	20.2	2.4
3.1.13-1	0.1	>21	13.2	0.56
2.2.13-1	0.08	>50	3.8	1.48
2.3.13-1	0.14	>38	15.5	1.5
1.1.10-1	0.12	45	5.43	0.17
3.1.14-1	0.12	>27	6.42	0.19
2.2.14-1	0.06	>50	2.26	0.21
2.3.14-1	0.11	35.3	5.23	0.21
1.1.54-1	0.16	>50	7.08	0.32
3.1.5-1	0.14	>17	9.8	0.23
2.2.5-1	0.05	>50	2.34	0.13
2.3.5-1	0.05	>50	3.52	0.19
1.1.104-1	0.12	>50	5.13	0.45

mAb	Neutralization IC50 Values (µg/mL) for the Virus Indicated			
	92US712	92US727	7141	VS001
3.1.15-1	0.12	>33	9.1	0.49
2.2.15-1	0.03	>50	1.99	0.37
2.3.15-1	0.1	>50	5.93	0.85
1.1.119-1	0.08	>50	6.92	0.39
3.1.7-1	0.09	>50	5.65	0.3
2.2.7-1	0.04	>50	1.58	0.27
2.3.7-1	0.07	>50	5.08	0.47
1.1.42-1	0.29	>50	>50	0.9
3.1.2-1	0.09	>50	21.9	0.73
2.2.2-1	0.04	>50	2.06	0.44
2.3.2-1	0.1	>50	14.6	0.88
1.1.75-1	0.08	>50	11.3	0.5
3.1.16-1	0.29	>43	39	2.15
2.2.16-1	0.05	>50	3.38	0.78
2.3.16-1	0.09	>50	10.2	0.72
1.1.78-1	0.1	>50	29.5	3.3
3.1.17-1	0.1	>50	25.5	2.21
2.2.17-1	0.05	>50	6.28	4.38
2.3.17-1	0.08	>50	31.3	4.54
1.1.102-1	0.14	>50	22.7	3
3.1.18-1	0.24	>9.8	>9.8	3.52
2.2.18-1	0.09	>34	5.04	2.51
2.3.18-1	0.16	>23	>23	2.37
1.1.120-1	0.2	>32.9	17.8	0.88
3.1.19-1	0.1	>50	23.9	0.57
2.2.19-1	0.06	>50	4.83	1.04
2.3.19-1	0.09	>50	12	0.48
1.1.121-1	0.1	>37	28.7	1.3
3.1.20-1	0.06	>50	22.1	0.95
2.2.20-1	0.05	>50	4.01	1.25
2.3.20-1	0.08	>50	15.9	1.23
1.1.122-1	0.13	>42.4	6.79	0.32
1.1.123-1	0.09	>43.8	11.2	0.23
1.1.124-1	0.14	>25.4	9.87	0.3
1.1.125-1	0.12	>50	7.69	0.37
1.1.126-1	0.12	>50	6.49	0.35
1.1.127-1	0.16	>23	5.78	0.18
1.1.128-1	0.17	>28.3	9.5	0.67
1.1.129-1	0.15	>29.7	>29.7	2.09
1.1.130-1	0.21	>27.3	16.7	1.67
1.1.131-1	0.15	>31.9	16.4	0.57
1.1.111-1	0.13	>50	>50	3.24
B-1	0.13	2.57	1.43	0.29

mAb	Neutralization IC50 Values (µg/mL) for the Virus Indicated			
	92US712	92US727	7141	VS001
1.1.90-1	0.12	8.72	2.65	0.2
1.1.132-1	0.12	2.26	3.39	0.14
1.1.133-1	0.14	2.27	3.38	0.16
1.1.134-1	0.14	4.23	3.15	0.22
1.1.135-1	0.22	5.61	4	0.31
1.42.1-1	0.13	0.98	2.34	0.11
1.43.1-1	0.08	3.35	2.42	0.1
1.44.1-1	0.06	~7-12	1.43	0.13
1.45.1-1	9.8	>50	>50	27.6
1.46.1-1	0.17	1.68	2.44	0.24
1.47.1-1	0.11	1.17	2.48	0.24
1.49.1-1	0.14	2.41	2.98	0.26
1.50.1-1	0.08	0.92	2.54	0.14
1.51.1-1	0.1	2.67	3.5	0.22
A-1	0.13	4.8	2.9	0.2

[0341] All antibodies carrying mutations at position N72 exhibited loss of function on the 92US727 virus. The antibody with the highest median potency was Antibody 1.1.64. Some antibodies show improvements in median neutralization potency compared to Antibody A-1. Among the antibodies containing the N72T mutation with reduced BVP ELISA scores, Antibody 2.3.5 also exhibited increased potency in the HIV neutralization assays.

[0342] Additional neutralization assays were conducted on a panel of variants designed to remove the heavy chain W74a oxidation motif and the light chain N26 deamidation motif in the A-1 variable domain. The results of the assessment are shown in Tables 29 and 30. The results show that many variants exhibited loss of function, while select variants retained potency more similar to A-1.

[0343] Table 29: HIV neutralization results for A-1 variants designed to remove chemical liabilities

		Virus							
Liability Addressed	Antibody Name	VS003	VS046	VS001	VS026	VS052	US657	VS043	VS004
None	A-1	0.03	0.03	0.08	0.26	0.38	0.86	1.47	3.42
Light Chain N26	1.1.136-1	0.03	0.04	0.20	0.76	0.38	0.92	1.57	>100
	1.1.137-1	0.03	0.05	0.16	0.43	0.35	0.78	1.65	>100
	1.1.138-1	0.06	0.04	0.18	0.35	0.46	0.54	1.41	39.24
	1.1.139-1	0.03	0.05	0.36	0.94	0.51	0.54	1.37	>100
	1.1.140-1	0.03	0.04	0.17	0.56	0.47	0.51	2.82	94.28
	1.1.141-1	0.02	0.04	0.25	0.69	0.51	0.38	1.30	>100
	1.1.142-1	0.02	0.03	0.20	0.48	0.56	0.44	1.42	>100
	1.1.143-1	0.05	0.05	0.40	3.85	0.92	0.82	2.05	89.88
	1.1.144-1	0.03	0.05	0.38	1.07	0.66	0.38	1.29	>100
Heavy Chain W74a	1.52.1-1	0.05	0.06	0.21	0.48	0.54	0.69	1.39	9.90
	1.53.1-1	0.06	0.05	0.18	0.49	0.57	0.72	1.44	14.10
	1.54.1-1	0.08	0.06	0.55	0.59	0.75	1.27	1.94	36.15
	1.55.1-1	0.08	0.07	0.49	0.83	1.08	1.39	2.48	34.00
	1.56.1-1	0.09	0.07	0.49	0.72	0.87	1.28	1.51	21.80
	1.57.1-1	0.06	0.09	0.65	0.91	1.21	2.43	2.47	56.68

[0344] Table 30: HIV neutralization results for selected A-1 variants on an expanded viral panel

Avg. IC50 ($\mu\text{g/mL}$)	Antibody A-1	1.1.136-1	1.1.137-1	1.1.138-1	1.1.140-1	1.1.142-1	1.52.1-1	1.53.1-1
US657(1)	0.864	0.92	0.78	0.55	0.55	0.51	0.44	0.73
VS003	0.027	0.03	0.03	0.06	0.03	0.02	0.04	0.06
VS046	0.029	0.04	0.05	0.04	0.04	0.03	0.05	0.05
VS001	0.080	0.20	0.16	0.18	0.17	0.20	0.21	0.18
VS026	0.259	0.76	0.43	0.35	0.56	0.48	0.48	0.50
VS052	0.381	0.38	0.35	0.45	0.47	0.56	0.54	0.56
VS043	1.466	1.57	1.65	1.41	2.82	1.42	1.39	1.44
VS004	3.419	>100	>100	39.24	>100	>100	9.90	14.10
US657(2)	0.336	0.30	0.31	0.42	0.46	0.59	0.82	0.65
VS002	0.024	0.02	0.03	0.03	0.03	0.02	0.05	0.05
VS011	0.147	0.19	0.18	0.25	0.25	0.37	0.33	0.34
VS017	0.443	2.14	1.75	2.02	2.13	2.87	3.01	3.03
VS030	1.614	2.49	2.86	2.81	3.28	3.61	4.86	3.54
VS032	0.256	0.27	0.28	0.25	0.20	0.17	0.27	0.25
VS033	0.207	0.15	0.20	0.19	0.09	0.08	0.27	0.11
VS034	0.358	0.54	0.40	0.45	0.23	0.33	0.28	0.45
VS038	0.210	0.22	0.18	0.12	0.28	0.34	0.40	0.18
VS039	0.121	0.10	0.07	0.07	0.05	0.07	0.17	0.15
VS042	3.171	3.37	2.23	1.73	2.01	1.72	4.54	3.42
VS044	0.312	0.73	0.39	0.36	0.06	0.07	0.16	0.12
VS053	0.530	1.83	0.88	1.52	1.61	1.17	1.07	0.87

[0345] Next, select A-1 variants were profiled for impact of the mutations on neutralization breadth and potency, using a PhenoSense™ HIV Neutralization Assay

(Monogram Biosciences; *see also*, Richman, *et al.*, *Proc Natl Acad Sci USA*. (2003) 100(7):4144-9, Whitcomb, *et al.*, *Antimicrob Agents Chemother*. (2007) 51(2):566-75), using reporter viruses pseudotyped with patient virus-derived envelopes (Table 31). The panel comprised 152 Env vectors in total (one vector per patient), with 133 clonal vectors and 19 isolates (representing the quasispecies in the sampled plasma). Briefly, the pseudoviruses were incubated with 5-fold serial dilutions of the antibody for 1 hour at 37°C and then used to infect U87 cells expressing CD4, CCR5 and CXCR4 (CD4+/CCR5+/CXCR4+/U87). The ability of an antibody to neutralize HIV infectivity was assessed by measuring luciferase activity 72 hours post incubation of cells with virus. Virus and antibody controls were employed to monitor plate to plate performance within a run and to allow for comparison of runs over time. All test antibodies were screened against a control panel of viruses consisting of HIV-1 NL4.3 (CXCR4-tropic), JRCSF (CCR5-tropic), and MLV (non-HIV specificity control). A broadly neutralizing HIV+ plasma sample served as the antibody control. While some mutations had more subtle impact on activity, inducing either a slight reduction or slight gain in activity, other mutations induced a notable loss of neutralization breadth (Table 31 and Figure 4).

[0346] Table 31: Neutralization potency and breadth of select antibodies were profiled against 152 patient-derived subtype B viruses obtained from pre-ART plasma.

Name	Breadth ^a	Median IC ₉₅ , μg/mL ^b	Geomean IC ₉₅ , μg/mL ^b
A-1	89 %	2.66	3.06
1.1.90-1	86 %	2.59	3.01
1.1.64-1	92 %	2.25	2.70
1.1.10-1	86 %	1.93	2.53
1.52.1-1	83 %	3.66	4.33
1.52.90-1	78 %	4.42	4.56
1.1.138-1	82 %	2.59	2.96

a Breadth represents % viruses neutralized with an IC₉₅ ≤ 50 μg/mL

b Median and Geomean IC₉₅ values calculated using only viruses with IC₉₅ ≤ 50 μg/mL

[0347] The neutralization data shown above was combined with the results of the polyspecificity screening (Example 9) and immunogenicity screening (Example 11) in order to design a pane of 12 lead variants. The panel of 12 Expi293-expressed antibodies tested in the BVP ELISA shown in Table 22 were next examined in an HIV neutralization assay against an expanded panel of viruses. The results for antibodies

retaining the N72-linked glycan are shown in Table 32, while the results for antibodies lacking the N72-linked glycan are shown in Table 33.

[0348] Table 32: HIV Neutralization Potency of 7 antibodies retaining the N72-linked glycan tested using an expanded panel of viruses

Virus	<i>Virus Neutralization Potency ($\mu\text{g/mL}$)</i>						
	A-1	1.1.64-1	1.52.64-1	1.1.90-1	2.2.1-1	2.4.1-1	2.3.1-1
1003	1.79	2.97	3.29	1.17	1.61	3.62	1.87
1012	1.41	1.41	1.71	2.92	3.70	3.38	1.55
1413	5.31	4.22	7.79	2.07	4.13	5.04	5.28
1489	0.12	0.16	0.17	0.22	0.26	0.16	0.18
7015	0.53	0.65	0.98	0.63	0.47	0.54	0.63
7051	0.79	2.05	0.82	0.83	0.97	1.18	0.52
7103	0.76	1.56	1.09	1.75	1.72	1.31	0.94
7141	2.32	1.82	30.02	2.66	3.93	4.82	1.62
7467	0.04	0.06	0.07	0.05	0.03	0.05	0.06
7552	12.04	>80	50.05	>80	39.62	37.05	20.69
7576	0.69	0.39	1.46	1.42	0.70	1.40	0.92
7595	4.88	1.71	7.34	10.46	9.72	5.06	8.50
7714	4.20	ND	ND	ND	ND	ND	ND
8106	2.72	3.31	8.15	4.54	2.46	4.25	3.12
8110	3.64	6.74	7.04	3.16	3.62	5.14	3.48
8117	0.86	1.77	2.18	1.47	0.94	2.58	2.08
8134	1.57	0.60	3.66	3.78	1.52	1.05	2.27
8176	0.26	0.66	0.34	0.27	0.30	0.33	0.30
8318	0.27	0.33	0.42	0.39	0.36	0.40	0.31
8320	2.20	2.89	3.82	3.22	3.88	6.69	2.88
302076	0.08	0.08	0.11	0.08	0.07	0.13	0.10
92HT593	0.27	0.37	0.25	0.22	0.17	0.37	0.29
92US657	0.38	0.42	0.48	0.40	0.25	0.34	0.36
92US712	0.06	0.06	0.13	0.05	0.05	0.07	0.06
92US727	2.43	1.12	5.47	8.61	4.84	13.74	2.02
CHO77	0.04	0.04	0.12	0.06	0.02	0.08	0.03

Virus	<i>Virus Neutralization Potency ($\mu\text{g/mL}$)</i>						
	A-1	1.1.64-1	1.52.64-1	1.1.90-1	2.2.1-1	2.4.1-1	2.3.1-1
REJO	0.01	0.02	0.02	0.04	0.01	0.02	0.02
THRO	2.80	3.87	3.11	2.04	2.17	4.17	3.96
VS001	0.06	0.06	0.11	0.10	0.08	0.18	0.07
VS004	2.66	2.43	4.74	3.21	2.84	6.08	1.92
VS017	0.70	0.91	2.01	1.08	0.84	3.91	0.88
VS026	0.11	0.17	0.23	0.20	0.13	0.27	0.13
VS030	2.04	2.60	7.98	7.40	2.51	5.20	6.38
VS039	0.12	0.09	0.27	0.14	0.03	0.11	0.06
VS042	2.53	2.33	4.44	1.88	0.89	2.00	1.11
VS043	0.79	0.85	1.30	1.11	1.01	1.16	0.97
VS044	0.18	0.16	0.38	0.26	0.13	0.25	0.16
VS046	0.04	0.04	0.05	0.03	0.03	0.05	0.03
VS049	39.54	32.19	>100	>100	2.89	>100	4.86
VS052	0.28	0.45	0.61	0.64	0.24	0.50	0.33

[0349] Table 33: HIV Neutralization potency of 5 antibodies lacking the N72 glycan using an expanded panel of viruses

Virus	<i>Virus Neutralization Potency ($\mu\text{g/mL}$)</i>				
	1.1.104-1	1.1.119-1	3.1.5-1	2.2.5-1	2.3.5-1
1003	1.44	0.91	1.02	0.78	1.06
1012	2.87	1.06	1.19	0.98	1.63
1413	4.34	4.52	4.30	3.62	3.79
1489	0.28	0.17	0.16	0.11	0.14
7015	0.67	1.08	1.16	0.95	1.89
7051	1.32	1.89	1.27	3.51	4.46
7103	1.82	1.16	0.91	0.78	1.05
7141	6.57	4.98	3.31	1.75	2.41
7467	0.06	0.19	0.12	0.07	0.09
7552	>80	>80	>80	>80	>80
7576	1.10	0.60	0.55	0.58	0.61
7595	8.47	3.07	3.56	2.94	3.07

Virus	<i>Virus Neutralization Potency (μg/mL)</i>				
	1.1.104-1	1.1.119-1	3.1.5-1	2.2.5-1	2.3.5-1
7714	ND	1.93	1.50	0.80	1.47
8106	2.80	1.44	1.45	0.94	1.44
8110	4.19	2.37	1.81	1.10	1.82
8117	1.22	1.66	1.02	0.75	1.06
8134	6.07	3.44	2.78	1.74	1.52
8176	0.16	0.27	0.30	0.20	0.33
8318	0.32	0.42	0.36	0.33	0.41
8320	4.34	1.47	1.33	1.21	1.48
302076	0.10	0.08	0.06	0.06	0.06
92HT593	0.28	0.15	0.16	0.11	0.17
92US657	0.36	0.30	0.27	0.15	0.24
92US712	0.04	0.03	0.06	0.04	0.05
92US727	10.55	43.70	11.98	16.74	9.32
CHO77	0.03	0.02	0.04	0.02	0.04
REJO	0.02	0.03	0.02	0.01	0.02
THRO	4.27	2.80	2.70	1.22	2.34
VS001	0.34	0.37	0.10	0.10	0.13
VS004	4.80	6.02	1.55	3.22	2.77
VS017	2.33	2.75	1.03	0.77	0.70
VS026	0.19	0.14	0.10	0.06	0.10
VS030	7.90	4.96	6.86	4.22	6.23
VS039	0.10	0.13	0.12	0.06	0.15
VS042	2.33	2.19	1.94	1.21	1.56
VS043	1.49	0.81	0.54	0.41	0.79
VS044	0.28	0.20	0.16	0.12	0.17
VS046	0.03	0.03	0.03	0.03	0.04
VS049	>100	>100	20.37	10.85	36.15
VS052	0.27	0.39	0.31	0.20	0.31

[0350] The results in Tables 32 and 33 show that all 12 antibody variants tested have similar virus neutralization potency values on the expanded panel of viruses. A subset of

variants were also profiled for neutralization breadth and potency via the phenosense neutralization assay, using a panel of 141 reporter viruses pseudotyped with subtype B patient virus-derived envelopes (Table 34 and Figure 5). Each envelope vector comprised isolate sampled from one patient. The variants exhibited comparable neutralization potency and breadth.

[0351] Table 34: Neutralization Activity of mAb Variants Profiled Against HIV-1 Pseudotyped with Env from Subtype B Plasma Isolates

	A	A-1	1.52.64-1
Breadth ^a	87 %	87 %	86 %
Median IC₉₅, µg/mL ^b	1.72	1.93	2.00
Geomean IC₉₅, µg/mL ^b	2.26	2.23	2.94

^a Breadth represents % viruses neutralized with an IC₉₅ ≤ 50 µg/mL

^b Median and Geomean IC₉₅ values calculated using only viruses with IC₉₅ ≤ 50 µg/mL

Example 11: *In vitro* Whole Molecule T-cell Proliferation and IL2 Release

[0352] Host anti-drug-antibody (ADA) responses can negatively impact the efficacy and pharmacokinetics of therapeutic antibodies and the resulting immune complexes may present safety concerns (Pratt KP. 2018. Antibodies. 7:19, Krishna M and Nadler SG. 2016. Front. Immunol. 7:21). As a result, *in vitro* T-cell proliferation and IL2 release assays such as the EpiScreen™ (Abzena Ltd.) have been developed to assess the overall immunogenic risk of biotherapeutics. The EpiScreen™ measures biotherapeutic induced IL2 release via the Enzyme Linked Immunosorbent Spot (ELISpot) assay and T-cell proliferation via 3H-thymidine incorporation in CD8+ T-cell depleted primary PMBC cultures obtained from 50 donors selected to represent HLA allotypes expressed among the world population. A highly immunogenic protein such as keyhole limpet hemocyanin (KLH) will induce both IL2 release and T-cell proliferation in >80% of donors, approved biotherapeutics such as Alemtuzumab and Infliximab with high rates of clinical immunogenicity will induce response rates in 25%-40% of donors, while biotherapeutics with low immunogenic risk typically show donor response rates ≤10%. Donor response rates in the EpiScreen™ have been shown to correlate with clinical ADA rates (Baker and Jones 2007. Curr. Opin. Drug Discov. Devel. 10: 219-227).

[0353] Table 35 shows the results of the EpiScreen™ assay for a panel of anti gp120 bNAbs that were transiently expressed in Expi293 cells and purified using protein A and size exclusion chromatography. Also shown are the A33 antibody and KLH as positive controls. In contrast to the immunogenic positive control proteins, many of the anti-gp120 antibodies tested, including A-1, show unusually high T-cell proliferation rates, but have relatively low IL2 release rates. This data suggests that in the absence of target, A-1 and other anti-gp120 bNAbs may directly stimulate ³H-thymidine incorporation in primary human PBMCs *in vitro* via an unknown mechanism. This unknown mechanisms, hereafter referred to as “off-target activity” could present safety liabilities if it translated *in-vivo*.

[0354] Table 35: EpiScreen™ results for 10 Expi293™ expressed anti-gp120 Abs tested on PBMCs from 50 donors.

Ab Name	Lot #	Expression Cell Line	³ H-thymidine incorporation (%)	IL2 Release (%)	³ H + IL2 (%)
E	3	Expi293™	16	6	4
A-1	3	Expi293™	32	6	6
F	4	Expi293™	60	10	10
G	4	Expi293™	60	8	8
H	4	Expi293™	50	14	12
I	3	Expi293™	56	16	10
J	4	Expi293™	6	14	4
K	4	Expi293™	16	6	2
E-7	2	Expi293™	14	10	2
E-6	2	Expi293™	10	8	2
A33	NA	NA	20	22	14
KLH	NA	NA	90	82	80

Response rates (%) indicate the percent of 50 donors showing responses in the indicated assays.

[0355] The heavy and light chains of antibodies E, F, G, H, I, J, K, L, L-1, E-6 and E-7 are provided in Table 36.

TABLE 36 - COMPARISON / CONTROL ANTIBODIES		
Ab Name	Heavy Chain (HC) Amino Acid Sequence	Light Chain Amino Acid Sequence
E	EVQLVESGGGLVKAGGSLILSCGVSNFRI SAHTMNVRRVPVGGGLEWVASISTST YRDYADAVKGRFTVSRDDLEDFVYLQMHKMRVEDTAIYYCARKGSDRLSDNDPDA WGPGLVTVSPASTKGPVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSNWSGA LTSVHTFPAVLQSSGLYSLSVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMI SRTPEVTCVAVDVSHEDPEVK FNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVMHEALHNHYTQKSLS LSPGK (SEQ ID NO: 632)	DVMTQSPSTLSASVGDITITCRASQSIET WLAWYQQKPGKAPKLLIYKASTLKTGVPSRF SGSGSGTEFTLTISGLQDDEDFATYHCQHYAG YSATFGQGRVEIKRTVAAPSVFIFPPSDEQ LKSGTASVVCCLNNFYPREAKVQWKVDNALQ SGNSQESVTEQDSKSDSTYLSSTLTLSKADY EKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 643)
F	EVQLVQSGTQMKEPGASVTI SCVTSGYEFVEILINWVRQVPGRGLEWGMWNPGRGG GVNVARQFQGVMTDRVYRDTAYLTLSGLTSGDTAKYFCVGRGRSCCGGRRHCNGA DCFNWDFAHWGQGLTVIVSPASTKGPVFLAPSSKSTSGGTAALGCLVKDYFPEP VTVSNWSGALTSGVHTFPAVLQSSGLYSLSVTVPSSSLGTQTYI CNVNHKPSNT KVDKKVEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMI SRTPEVTCVAVD VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKC KVSNAKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVLHEALH SHYTQKSLSLSPGK (SEQ ID NO: 633)	YIGVTSQSPAILSVSLGERVTLSCKTSQLTP RHLVWHRQKGGQAPSLVMTGTSERASGIPDR FIGSGSGTDFTLITITRLEAEDFAVYQCQCLE AFGQGTKEIKRTVAAPSVFIFPPSDEQLKS GTASVVCCLNNFYPREAKVQWKVDNALQSGN SQESVTEQDSKSDSTYLSSTLTLSKADYEKH KVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 644)
G	EVQLVQSGTQMKEPGASVTI SCVTSGYEFVEILINWVRQVPGRGLEWGMWNPGRGG GVNVARQFQGVMTDRVYRDTAYLTLSGLTSGDTAKYFCVGRKSCCAGRRFCGPT DCYNWDFAHWGQGLTVIVSPASTKGPVFLAPSSKSTSGGTAALGCLVKDYFPEP VTVSNWSGALTSGVHTFPAVLQSSGLYSLSVTVPSSSLGTQTYI CNVNHKPSNT KVDKKVEPKSCDKTHTCPPCPAPELGGPSVFLFPPKPKDTLMI SRTPEVTCVAVD VSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKC KVSNAKALPLPEEKTI SKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFSVLHEALH SHYTQKSLSLSPGK (SEQ ID NO: 634)	EIVLTQSPGTLSSLSPGETAII SCRTSQYGS AWYQQRPQGAPRLVIYSGSTRAAGI PDRFSG SRWGPDYNLITISNLESGDFGVYQCQYEFFG QGTQVQVDIKRTVAAPSVFIFPPSDEQLKSG TASVVCCLNNFYPREAKVQWKVDNALQSGNS QESVTEQDSKSDSTYLSSTLTLSKADYEKH VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 645)
H	QVRLSQSGGQMKKPGDSMRI SCRASGYEFINCPINWIRLAPGRPEWGMWMPKPRGG AVSYARQLQGRVTMTDMYSETAFLELRSLTSDDTAVYFCTRGKYCTARDYYNWD EHWGQGTPTVTSSASTKGPVFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSNWS GALTSGVHTFPAVLQSSGLYSLSVTVPSSSLGTQTYI CNVNHKPSNTKVDKKVE	EIVLTQSPGTLSSLSPGETAII SCRTSQYGS AWYQQRPQGAPRLVIYSGSTRAAGI PDRFSG SRWGPDYNLITISNLESGDFGVYQCQYEFFG QGTQVQVDIKRTVAAPSVFIFPPSDEQLKSG

TABLE 36 – COMPARISON / CONTROL ANTIBODIES		
Ab Name	Heavy Chain (HC) Amino Acid Sequence	Light Chain Amino Acid Sequence
	PKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDITLMI SRTPEVTCVVVDVSHEDPE VKFNWYVDGVEVHNKATKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKAL PLPEEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESN GQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSHSHYTTQKS LSLSPGK (SEQ ID NO: 635)	TASVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSYSLSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 646)
I	RVQLVESGGGVVQPGKSVRLSCVVSDFPFESKYPMYWVRQAPGKGLEWVAALISGDAW HVVYSNSVQGRFLVSRDNLVNTLYLEMNLSLKIEDTAVYRCARMFQESGPPRLDRWS GRNYYSYSGMDVWGQGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF PEPVTISWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHP SNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDITLMI SRTPEVTCV VVDVSHEDPEVKFNWYVDGVEVHNKATKPREEQYNSTYRVVSVLTVLHQDWLNGKE YKCVSNKALPLPEEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPS SDIAVEWESNGQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSH ALSHYTTQKSLSLSPGK (SEQ ID NO: 636)	DIVMTQTPLSLSVTPGPASISCKSSSESLRQ SNGKTSLYWYRQKPGQSPQQLLVFEVSNRFSG VSDRFVSGSGGTFTLRI SRVEAEDVGFYFC MQSKDFPLTFGGGTQVVDLKRITVAAPSVFIFP PSDEQLKSGTASVCLLNFFYPREAKVQWKV DNALQSGNSQESVTEQDSKDSYSLSTLTLS SKADYEKHKVYACEVTHQGLSSPVTKSFNRG EC (SEQ ID NO: 647)
J	QEQLVESGGGVVQPGGSLRLSCLASGFTTHKYGMHWVRQAPGKGLEWVALISDDGM RKYHSDSMWGRVTISRDNKNTLYLQFSLLKVEDTAMFFCAREAGGPIWHDDVKYY DFNDGYNYHYMDVWGKGTITVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKD YFPEPVTISWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHP KPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDITLMI SRTPEVT CVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEQYNSTYRVVSVLTVLHQDWLNG KEYKCVSNKALPLPEEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGF YPSDIAVEWESNGQPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCFVLSH ALSHYTTQKSLSLSPGK (SEQ ID NO: 637)	QSALTQPASVSGSPGQTITISCNGTSSDVG FDSVWYQQSPGKAPKVMVFDVSHRPSGISN RFGSGKSGNTASLTISGLHIEDEGDYFCSSL TDRSHRIFGGGTQVTVLQGPKAAPSVTLFPP SSEELQANKATLVCLISDFYPGAVTVAAKAD SSPVKAGVETTTTPSKQSNKKYAAASSYLSLTP EQWKSRRSYSCQVTHEGSTVEKTVAPTECS (SEQ ID NO: 648)
K	QVQLVQSGAEVKKPGSSVKVSKASGNSFNSHVDHWVRQATGQGLEWMGMWSHEGD KTGLAQKFQGRVTITRDGASTVYMEIRGLTADDTAIYCLTGSKHRLRDYFLYNE YGPNYEEWGDYLA TLVDWGHGTA VTVSSASTKGPSVFPLAPSSKSTSGGTAALGCL VKDYFPEPVTISWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICN VNHKPSNTKVDKKVEPKSCDKTHTCPPCPAPELLAGPDVFLFPPKPKDITLMI SRT PEVT CVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEQYNSTYRVVSVLTVLHQDW LNGKEYKCVSNKALPLPEEKTIISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLV	EVVITQSPFLFPLVTPGEAASLSCKSHSLQH STGANLYAWYLRPGQTPTLLIHLATHRASG VPDRFSGSGGTFTLKI SRVESDDVGTYYC MQGLHSPVTFGGGTQVTVLQGPKAAPSVTLFPP PSDEQLKSGTASVCLLNFFYPREAKVQWKV DNALQSGNSQESVTEQDSKDSYSLSTLTLS SKADYEKHKVYACEVTHQGLSSPVTKSFNRG EC (SEQ ID NO: 649)

TABLE 36 – COMPARISON / CONTROL ANTIBODIES		
Ab Name	Heavy Chain (HC) Amino Acid Sequence	Light Chain Amino Acid Sequence
L	KGFYPSDIAVEWESNGQPENNYKTTTPPVLSDSGSFFLYSKLTVDKSRWQQGNVFSC SVLHEALHSHYTQKSLSPGK (SEQ ID NO: 638)	EIVLTQSPGTLSSLSPGETAIIISCRTSQYGS AWYQQRPQGAPRLVIYSGSTRAAGIPDRFSG SRWGPDYNLITISNLESGDFGVYCCQYEFFG QGTKVQVDIKRTVAAPSVFIFPPSDEQLKSG TASVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 650)
L-1	QVQLVQSGGQMKKPGESMRI SCRASGYEFIDCTLNWIRLAPGKRPEWMGWLKPRGG AVNYARPLQGRVTMTRDVSDTAFLERSLTVDDTAVFCTRGKNCNDYNWDFEHWG RGTPVIVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQYICNVNHKPSNTKVDKKVEPKSC DKHTCCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVAVDVSHEDPEVKFN WYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPI EKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPE NNYKTTTPPVLSDSGSFFLYSKLTVDKSRWQQGNVFSCSVMEALHNHYYTQKSLSL PGK (SEQ ID NO: 639)	EIVLTQSPGTLSSLSPGETAIIISCRTSQYGS AWYQQRPQGAPRLVIYSGSTRAAGIPDRFSG SRWGPDYNLITISNLESGDFGVYCCQYEFFG QGTKVQVDIKRTVAAPSVFIFPPSDEQLKSG TASVCLLNFFYPREAKVQWKVDNALQSGNS QESVTEQDSKDSSTYSLSSTLTLSKADYEKHK VYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 651)
E-6	EVQLVESGGGLVKAGGSLILSCGVSNFRISAHTTMNWRRVPGGGLWVASISTST YRDYADAVKGRFTVSRDDEDFVYLQMHKMRVEDTAIYYCARKGSDRLSDNDPFD WGPSTVTVSPASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGA LTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVAVDVSHEDPEVK FNWYVDGVEVHNAKTKPREEQYNSTLRVVSILTVLHQDWLNGKEYKCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTTPVLSDSGSFFLYSKLTVDKSRWQQGNVFSCSVLHEALHSHYYTQKSL LSPGK (SEQ ID NO: 641)	DVAVMTQSPSTLSASVGDITITITCRASQSIET WLAWYQQKPGKAPKLLIYKASTLKTGVPSRF SGSGSGTEFTLTISGLQFDDFATYHCQHYAG YSATFGQGRTRVEIKRTVAAPSVFIFPPSDEQ LKSGETASVCLLNFFYPREAKVQWKVDNALQ SGNSQESVTEQDSKDSSTYSLSSTLTLSKADY EKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 652)

TABLE 36 - COMPARISON / CONTROL ANTIBODIES		
Ab Name	Heavy Chain (HC) Amino Acid Sequence	Light Chain Amino Acid Sequence
E-7	EVQLVESGGGLVKAGGSLILSCGVSNFRI SAHTMNWRRVPGGGLEWVASISTST YRDYADAVKGRFTVSRDDLEDFVYLQMHKMRVEDTAIYYCARKGSDRLSDNDPFD WGPGTVTVSPASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGA LTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVDKKVEPK SCDKTHTCPPCPAPELVGGPSVFLPPKPKDTLMI SRTPEVTCVAVDVSHPEDPEVK FNWYVDGVEVHNAKTKPREEQYNSTLRVSVLTVLHQDWLNGKEYCKVSNKALPA PIEKTISKAKGQPREPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQ PENNYKTTPLPLDSDGSFFFLYSKLTITVDKSRWQQGNVFCSVLHEALHSHYITQKSL LSPGK (SEQ ID NO: 642)	DVVMTQSPSTLSASVGDITITICRASQSIET WLAWYQQKPGKAPKLLIYKASTLKTGVPSRF SGSGSGTEFTLTISGLQFDDDFATYHCQHYAG YSATFGQGTREIKRTVAAPSVFIFPPSDEQ LKSGTASVVCCLNNFYFREAKVQWKVDNALQ SGNSQESVTEQDSKDSSTYSLSSTLTLSKADY EKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO: 653)

[0356] To better understand the unexpected off-target activity of A-1, we conducted a second EpiScreen assay on A-1 and variants described herein lacking the N72 linked glycosylation motif in the A-1 light chain. To determine if expression-host-dependent N72-linked glycan composition changes (Example 14, below) might impact the off-target activity of A-1, proteins for the second EpiScreen assay were expressed in the ExpiCHO™ cell line rather than the Expi293™ cell line. The results of this EpiScreen™ assay are shown in Table 37. Antibody A-1 expressed in the ExpiCHO cell line showed lower T-cell proliferation rates (16%) than A-1 expressed in Expi293 cells (32%) suggesting that the expression cell line and associated N72-linked glycan composition changes may have an impact on the putative off-target activity observed in the EpiScreen™. Unexpectedly, all variants of antibody A-1 lacking the N72-linked glycosylation site in the antibody light chain showed much higher T-cell proliferation rates. The results suggest that the composition of the N72-linked Fab glycan may play a role in modulating the off-target T-cell proliferation activity, but that removal of the N72-linked Fab glycan potentiates the off-target activity.

[0357] Table 37: EpiScreen™ results for 7 ExpiCHO™ expressed anti-gp120 Abs tested on PBMCs from 50 donors.

Name	Lot #	Expression Cell Line	³ H-thymidine incorporation (%)	IL2 Release (%)	³ H + IL2 (%)
A-1	6	ExpiCHO™	16	6	4
1.1.10-1	2	ExpiCHO™	32	6	6
1.1.42-1	2	ExpiCHO™	60	10	10
1.33.32-1	3	ExpiCHO™	60	8	8
1.1.54-1	2	ExpiCHO™	50	14	12
1.37.51-1	2	ExpiCHO™	56	16	10
1.8.52-1	2	ExpiCHO™	6	14	4
A33	NA	NA	16	6	2
KLH	NA	NA	14	10	2

Response rates (%) indicate the percent of 50 donors showing responses in the indicated assays.

[0358] Since the EpiScreen™ assay measures ³H-thymidine incorporation in primary PBMC cultures, it is possible that in the absence of IL2-release, the off-target activities

observed for A-1 and variants thereof could involve proliferation of any cell type present in the PBMCs (*e.g.*, B-cell proliferation instead of T-cell proliferation). To determine if ExpiCHO™ derived A-1 and a variant thereof lacking the N72-glycan were stimulating proliferation of T-cells, we next conducted an EpiScreen™ assay using either CD8+ T-cell depleted PBMCs or CD8+ and CD4+ T-cell depleted PBMCs taken from the same 10 donors. For a negative control, we selected Expi293 derived antibody L, which had previously shown low donor response rates in the EpiScreen™ (*see, e.g.*, WO 2017/106346). The results of this assay are shown in Table 38. The results clearly show that ³H-thymidine incorporation rates are reduced in the absence of CD4+ T-cells. This data shows that the off-target activity observed for A-1 and variants thereof is dependent on the presence of T-cells. As HIV infects and establishes a latent reservoir in T-cells, off-target anti-gp120 antibody induced T-cell proliferation could potentially expand the HIV-1 reservoir, and would thus be undesirable as part of an HIV cure strategy intended to deplete the HIV-1 reservoir.

[0359] Table 38: EpiScreen™ results for 3 anti-gp120 Abs tested on PBMCs from 10 donors in the presence (+CD4) or absence (-CD4) of CD4+ T-cells.

Name	Lot #	Expression Cell Line	³ H-thymidine incorporation +CD4 (%)	³ H-thymidine incorporation -CD4 (%)
A-1	6	ExpiCHO™	40	0
1.1.54-1	2	ExpiCHO™	40	10
L	3	Expi293™	0	0

Response rates (%) indicate the percent of 10 donors showing responses in the indicated assays.

[0360] As described later in Example 15, the molecular composition of the A-1 N72-linked light chain glycan and resulting pharmacokinetics can change dramatically depending on the expression host and resulting sialylation content of the light chain N72-linked Fab glycan. Based on the results of the EpiScreen assays reported in Tables 35 to 38, we hypothesized that the molecular composition of the A-1 N72-linked light chain glycan might impact the observed off-target T-cell proliferation activity described herein. To test this idea, we next conducted a 10 donor EpiScreen measuring T-cell proliferation using either ExpiCHO or CHO-S derived A-1 or variants thereof. As described in Examples 14 and 15, CHO-S derived A-1 has significantly higher N72-glycan sialylation content than Expi293™ or ExpiCHO™ derived material. The results of this EpiScreen

are shown in Table 39. Antibodies A-1 and 1.1.90-1 expressed in the CHO-S cell line showed no off-target T-cell proliferation. Although the number of donors in this screen was small, this data suggested that the A-1 expression cell line and associated N72-linked light chain glycan composition could modulate not just pharmacokinetics, but also modulate the observed off-target activity in the EpiScreen™ assay.

[0361] Table 39: EpiScreen™ T-cell proliferation results for 6 ExpiCHO™ or CHO-S expressed anti-gp120 Abs tested on PBMCs from 10 donors.

Name	Lot #	Expression Cell Line	³ H-thymidine incorporation (%)
1.1.54-1	6	ExpiCHO™	40
L-1	2	CHO-S	20
1.1.111-1	2	CHO-S	30
1.1.90-1	3	CHO-S	0
B-1	2	CHO-S	20
A-1	18	CHO-S	0
Exenatide	NA	NA	40
KLH	NA	NA	60

Response rates (%) indicate the percent of 50 donors showing responses in the indicated assays.

[0362] Based on the preliminary results shown in Table 39, we next conducted a 50 donor EpiScreen™ on a panel of 7 anti-gp120 antibodies including A-1 and variants thereof expressed in ExpiCHO or CHO-S cell lines. The results of this screen are shown in Table 40 and show that A-1 demonstrates very low T-cell proliferation and IL2 release when generated using CHO-S cell lines that incorporate a high level of sialic acid into the N72-linked light chain glycosylation site (see examples 14-15). The results further demonstrate that selected variants of A-1 have further reduced T-cell proliferation rates when produced and tested in the same manner.

[0363] Table 40: EpiScreen™ results for 7 anti-gp120 Abs tested on PBMCs from 50 donors.

Name	Lot #	Expression Cell Line	³ H -thymidine incorporation (%)	IL2 Release (%)	³ H + IL2 (%)
------	-------	----------------------	---	-----------------	--------------------------

1.1.64-1	5	CHO-S	36	6	2
A-1	27	CHO-S	10	0	0
1.52.64-1	3	CHO-S	4	4	0
3.1.5-1	3	CHO-S	28	6	4
2.3.5-1	3	CHO-S	28	2	0
1.1.10-1	3	ExpiCHO™	22	10	4
L-1	7	CHO-S	12	4	2
Exenatide	NA	NA	38	20	10
KLH	NA	NA	98	94	92

Response rates (%) indicate the percent of 50 donors showing responses in the indicated assays.

Example 12: *In vitro* Binding Assays

[0364] The pharmacokinetics (PK) and pharmacodynamics (PD) of antibody therapeutics is mediated by specific binding to target proteins via the variable domains and/or by binding to Fc-gamma receptors (FcγR) on innate immune cells, neonatal Fc-receptor (FcRn) on endothelial cells and circulating complement protein C1q (Nimmerjahn and Ravetch. 2008. Nat. Rev. Immunol. 8:34-47, Rogers et al. 2014. Immunol. Res. 59:203-210, Kuo TT and Aveson VG. 2011. MAbs 3:422-430). Genetic engineering of the antibody variable domain or Fc domain can impact binding to these receptors and influence PK and PD. We thus assessed the relative affinity of selected antibodies described herein using a variety of common *in vitro* binding assays including surface plasmon resonance (SPR) and enzyme linked immunosorbent assay (ELISA).

[0365] The *in vitro* binding dissociation constants (KD) of selected antibodies described herein for human and cynomolgus macaque (cyno) Fc binding receptors (FcγRs, FcRn) were determined using the Biacore 4000 surface plasmon resonance (SPR) biosensor, and either C1 or CM4 sensor chips (GE Healthcare). Biotinylated human FcRn was purchased from Immunitrack. Biotinylated cynomolgus macaque FcRn and human FcγRIIIB-NA1 and FcγRIIIB-NA2 were purchased from Acro Biosystems. Human FcγRIIA-167H, FcγRIIA-167R, FcγRIIA-176F, FcγRIIA-176V, FcγRIIB/C, FcγRI, and cynomolgus macaque FcγRI, FcγRIIA, FcγRIIB and FcγRIII were purchased from R&D systems.

[0366] For human FcRn binding assays, 600 RU of streptavidin was amine coupled to a C1 sensor chip using standard NHS/EDC coupling. The immobilization buffer was

PBS+0.005% Tween 20, pH 7.4. Streptavidin was prepared at 50 µg/ml in 10 mM NaAc pH 4.5. Activation, coupling, and blocking steps were run for 10 minutes, each at 10 µl/min. Biotinylated human FcRn was captured to about 20 relative units (RU). mAb samples A-1, A and 1.52.64-1 were tested for binding to the FcRn surface using a two-fold concentration series up to 1 µM. Data were collected at pH 6.0 and pH 7.4 in triplicate. The response data at steady-state were fit to a simple binding isotherm.

[0367] Human FcγRIIA and FcγRIIIA were amine coupled at 4 different densities (about 100 RU, about 250 RU, about 375 RU and about 725 RU) on a CM4 sensor chip. The three mAb samples were tested for binding in PBS pH 7.4 + Tween20 (0.005%) running buffer in a 2-fold dilution series up to 1 µM. Each mAb concentration series was tested twice over each of the 4 receptor densities surfaces generating 8 data sets for each interaction. The response data at steady-state were fit to a simple binding isotherm.

[0368] Human FcRIIB/C was amine coupled to a CM4 sensor chip at three different levels (50, 400 and 800 RU). The three mAbs were tested using 2 µM as the highest concentration in a two-fold dilution series. The concentration series was run in triplicate for each antibody across the low, medium and high density receptor surfaces. The response data at steady-state were fit to a simple binding isotherm.

[0369] To determine human FcγRIIIB binding affinities, each test antibody was amine coupled to a CM4 sensor chip at two densities (about 100 RU and about 800 RU). Human FcγRIIIB samples were tested for binding using a two-fold concentration series up to 0.5 µM. The response data at steady-state were fit to a simple binding isotherm.

[0370] To determine human FcγRI binding affinities, each test antibody was amine coupled to a CM4 sensor chip at two densities (about 100 RU and about 800 RU). Human FcγRI was tested for binding using a two-step titration series (3 nM and 30 nM). Responses were fit to a simple kinetic model.

[0371] To determine cynomolgus macaque FcRn binding affinities, 600 RU of streptavidin was amine coupled to a C1 sensor chip using standard NHS/EDC coupling. The immobilization buffer was PBS+0.005% Tween 20, pH 7.4. Streptavidin was prepared at 50 µg/ml in 10 mM NaAc pH 4.5. Activation, coupling, and blocking steps were run for 10 minutes, each at 10 µl/min. Biotinylated cyno FcRn was captured to about 20 RU. Antibodies were tested for binding to the FcRn surface using a two-fold concentration series up to 1 µM. Data were collected at pH 6.0 and pH 7.4 in triplicate. The response data at steady-state were fit to a simple binding isotherm.

[0372] To determine cynomolgus macaque FcγRIIA, FcγRIIB, FcγRIII and FcγRI binding affinities each test antibody was amine coupled to a CM4 sensor chip at two densities (about 100 RU and about 800 RU). Cyno FcγRIIA and FcγRIIB were tested in a two-fold concentration series up to 1 μM. FcγRIII was tested in a two-fold concentration up to 500 nM. Cyno FcγRI was tested for binding using a two-step titration (3 nM and 30 nM). The response data for FcγRIIA, FcγRIIB, FcγRIII at steady-state were fit to a simple binding isotherm. Responses for FcγRI were fit to a simple kinetic model.

[0373] The full set of binding constants determined by surface plasmon resonance (SPR) are shown in Table 41. The data shows that variants of antibody A with genetically engineered Fc domains have enhanced binding affinity to both human and cyno FcγR and FcRn proteins.

[0374] Table 41: Fc Receptor Binding Constants (KD) Determined by SPR

Fc receptor type-allele	A	A-1	1.52.64-1
Human FcγRI	0.107 ± 0.040 nM	0.002 ± 0.002 nM	0.0012 ± 0.0005 nM
Cyno FcγRI	0.038 ± 0.016 nM	0.005 ± 0.004 nM	0.005 ± 0.003 nM
Human FcγRIIA-167H	1.8 ± 0.5 μM	131 ± 22 nM	221 ± 21 nM
Human FcγRIIA-167R	3 ± 1 μM	130 ± 9 nM	199 ± 21 nM
Cyno FcγRIIA	2000 ± 1000 nM	1100 ± 80 nM	1180 ± 60 nM
Human FcγRIIB	11 ± 0.8 μM	1.6 ± 0.2 μM	1.9 ± 0.2 μM
Cyno FcγRIIB	895 ± 50 nM	240 ± 9 nM	280 ± 32 nM
Human FcγRIIIA – 176V	670 ± 40 nM	59 ± 4 nM	67 ± 6 nM
Human FcγRIIIA – 176F	2.3 ± 0.6 μM	52 ± 4 nM	63 ± 5 nM
Human FcγRIIIB – NA1	2000 ± 1000 nM	59 ± 9 nM	64 ± 15 nM
Human FcγRIIIB – NA2	1500 ± 400 nM	56 ± 18 nM	55 ± 14 nM
Cyno FcγRIII	200 ± 70 nM	7.1 ± 0.7 nM	8 ± 2 nM
Human FcRn pH 7.4	42 ± 1 μM	1.7 ± 0.3 μM	1.12 ± 0.08 μM
Human FcRn pH 6.0	485 ± 43 nM	38 ± 3 nM	49 ± 5 nM
Cyno FcRn pH 7.4	12.8 ± 0.4 μM	4.3 ± 0.4 μM	5.3 ± 0.2 μM
Cyno FcRn pH 6.0	1100 ± 100 nM	16 ± 2 nM	22 ± 2 nM

[0375] A dose response binding ELISA was conducted to determine the relative C1q binding affinity of antibodies described herein. To conduct the assay, a 384-well Maxisorp plate was coated with 25 μ L of antibody solution at 5 μ g/mL in PBS pH 7.4 overnight at 4 °C. Plates were then blocked with 75 μ L of 1% BSA in PBS for 2 hours and washed 4 times with PBS + 0.05% Tween 20 (PBST). Next, 25 μ L of a three-fold serial dilution of human C1q protein in PBS+5% BSA was added to the plates. Plates were incubated with shaking at 600 rpm for one hour, washed 4 times with PBST and then 25 μ L of anti-C1q-HRP conjugated polyclonal antibody was added in PBS+5% BSA. Plates were incubated with shaking at 600 rpm for 15 minutes, washed 8 times with PBST and then developed using 3,3',5,5'-Tetramethylbenzidine (TMB) substrate and quenched with HCl. Absorbance at 450nm was read using a spectramax m5 plate-reader and EC50 values were determined using a 4-parameter dose response fit.

[0376] The average EC50 values for the C1q binding ELISA were calculated from three independent assays and are shown in Table 42.

Table 42: C1q Binding EC50 values Determined by ELISA (n=3 assays)

Antibody	C1q Binding EC50 (nM)
A	2.2 $\pm$ 1.2 nM
A-1	> 100 nM
1.52.64-1	> 100 nM

[0377] The results show that Fc engineered variants of antibody A have significantly reduced C1q binding affinity.

[0378] A dose response binding ELISA was conducted to determine the relative gp120 binding affinities of the antibodies described herein. To conduct the assay, a 384 well Maxisorp plate was coated with 25 μ L of 5 μ g/mL gp120 and incubated overnight at 4°C. The plate was washed 4 times with PBS 0.05% Tween 20 and blocked with 75 μ L of PBS 5% BSA for 1hr at room temperature while shaking at 600rpm. After blocking, the wells were aspirated and 25 μ L of a 3-fold serial dilution of primary antibody was added and incubated at room temperature for 1hr with shaking at 600rpm. The plate was then washed 4 times with PBS 0.05% Tween 20 and 25 μ L of goat anti-human IgG (H+L) HRP secondary antibody diluted 1/10,000 in PBS 1% BSA was added and incubated at room temperature, shaking at 600 rpm for 30mins. Next, the plate was washed 4 times

with PBS 0.05% Tween 20 and 25 μ l fresh TMB substrate was added. The plate was developed for 90secs with shaking at 600rpm and before being quenched with 25 μ l 1M HCl. The absorbance was read at A450 on a Spectramax m5 plate reader.

[0379] The average EC₅₀ values were calculated from three independent ELISA assays and are shown in Table 43.

[0380] Table 43: gp120 binding EC₅₀ values determined by ELISA

gp120 protein	A	A-1	1.52.64-1
Bal	0.05 $\pm$ 0.02 nM	0.06 $\pm$ 0.01 nM	0.07 $\pm$ 0.02 nM
CAAN	1.84 $\pm$ 0.22 nM	2.17 $\pm$ 0.56 nM	3.79 $\pm$ 1.26 nM
REJO	2.21 $\pm$ 0.44 nM	2.37 $\pm$ 0.68 nM	4.02 $\pm$ 0.45 nM

[0381] The results suggest that all antibodies tested bind HIV gp120 protein with similar affinities.

Example 13: Effects of Fc Mutations on Serum Half-Life

[0382] In this example, IgG1 Fc mutations that enhance effector cell killing and/or that enhance FcRn binding were evaluated for effects on serum half-life. The data are consistent with the conclusion that mutations in the IgG1 Fc that enhance effector cell killing activity (*e.g.*, aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330 according to EU number (DEAL)) can shorten serum half-life *in vivo*. Such shortened serum half-life can be partially or wholly recovered by also incorporating mutations in the IgG1 Fc that enhance FcRn binding (*e.g.*, leucine at position 428, and serine at position 434 according to EU numbering (LS)).

[0383] PGT121-WT, PGT121-DEAL, PGT121.60, PGT121-LS (described, *e.g.*, in WO 2017/106346), and A-1 from the present application were administered to cynomolgus macaque monkeys (Covance, TX) at 10 mg/kg or 0.5 mg/kg (A-1) via a single intravenous (IV) injection to characterize their basic pharmacokinetic (PK) profiles. Serum samples collected from monkeys were analyzed using a bioanalytical method of sufficient selectivity and sensitivity to determine serum concentration-time profiles and calculate the mean serum PK parameters by non-compartmental PK analysis (NCA). The bioanalytical method utilized clade B gp120 antigen (Immune-tech, CA) as a capture reagent and biotin conjugated goat anti-human IgG antibody (Southern

Biotech, AL) as a secondary reagent, with SULFO-TAG labeled Streptavidin (MesoScale Discovery, MD) for electrochemical detection.

[0384] The measured serum concentration versus time profiles of PGT121-WT, PGT121-DEAL, PGT121.60, PGT121-LS, and A-1, depicted in Figure 6, were used to calculate the mean ($\pm$ SD) PK parameters depicted in Table 44.

[0385] Table 44: Pharmacokinetic parameters of PGT121-WT, PGT121-DEAL, PGT121.60, PGT121-LS, and A-1 after IV administration in naïve cynomolgus monkeys (n=3).

Test Article	Dose (mg/kg IV)	AUC _{0-∞} (day*ug/mL)	Cl (mL/day/kg)	V _d (mL/kg)	t _{1/2} (day)
PGT121-WT	10	1510 $\pm$ 470	7.0 $\pm$ 1.9	105 $\pm$ 17.6	10.6 $\pm$ 1.3
PGT121-DEAL	10	1020 $\pm$ 167	9.9 $\pm$ 1.5	109 $\pm$ 20	7.7 $\pm$ 1.3
PGT121.60	10	1490 $\pm$ 377	7.0 $\pm$ 1.9	96 $\pm$ 19	9.7 $\pm$ 0.8
PGT121-LS	10	3540 $\pm$ 463	2.9 $\pm$ 0.4	82 $\pm$ 11	19.9 $\pm$ 2.1
A-1	0.5	70 $\pm$ 7.0	7.2 $\pm$ 0.7	91 $\pm$ 14	8.7 $\pm$ 0.8

[0386] The PK analysis showed that inclusion of the Fc mutations (DEAL) to PGT121-WT negatively impacted the PK by increasing the clearance (Cl) to 9.9 $\pm$ 1.5 mL/day/kg for PGT121-DEAL relative to 7.0 $\pm$ 1.9 mL/day/kg for PGT121-WT and reduced the half-life (t_{1/2}) to 7.7 $\pm$ 1.3 days versus 10.6 $\pm$ 1.3 days for PGT121-WT. Inclusion of the FcRn binding mutations (LS) to antibodies with an Fc that contains the DEAL mutations (PGT121.60 and A-1), resulted in Cl values of 7.0 $\pm$ 1.9 and 7.2 $\pm$ 0.7 mL/day/kg and t_{1/2} values of 9.7 $\pm$ 0.8 and 8.7 $\pm$ 0.8 days, respectively, which are comparable to the PK of PGT121-WT. While inclusion of LS alone to PGT121-WT reduced the Cl to 2.9 $\pm$ 0.4 mL/day/kg and increased the t_{1/2} to 19.9 $\pm$ 2.1 days for PGT121-LS. The PK analysis support that introduction of the Fc-enhancing mutations DEAL reduces antibody PK (likely due to enhanced FcγR binding), which can be recovered by inclusion of the LS FcRn binding mutations.

Example 14. Light Chain Fab Glycan Profile Assessments

[0387] Two techniques were used to isolate and analyze the light chain Fab glycan profiles in the absence of potentially interfering heavy chain Fc glycans. The primary goal of these experiments was to understand the relative percentage of light chain glycans terminating with one or more sialic acid groups (referred to as percent sialylation hereafter). The first approach (“method 1”) was reverse phase mass spectrometry of the reduced, intact light chain. In this technique, observed mass shifts in the deconvoluted mass spectrum are assigned to the glycan structure known from biosynthetic N-glycan pathways to correspond to the mass shift. Relative quantification of the sialylated forms is obtained by summing the deconvoluted peak heights for the sialylated species and dividing this value by the total of all sialylated and non-sialylated peak heights. A second method (“method 2”) to quantify the sialylation on the light chain fab glycans relied on selective enzymatic release of the Fc glycans (under purely aqueous conditions) prior to isolations of the remaining protein and release of the remaining light chain Fab glycans. The separate aliquots corresponding to the Fc and Fab glycans are then fluorescently labeled (Waters RapiFluor) and analyzed, identified, and quantified by HILIC chromatography. The percent Fab sialylation values for multiple antibodies described herein and analyzed by one of these techniques are shown below in Tables 45A and 45B.

[0388] Table 45A. Light chain Fab glycan assessment of antibody A-1

Lot	14	10	7
Expression System	CHO-S	Tuna293™	ExpiCHO™
<u>Glycan ID¹</u>	% Peak Area	% Peak Area	% Peak Area
Unknown Peaks ²	6.47	3.50	4.16
G0-GlcNAc	0.00	4.20	4.49
G0	0.91	55.97	52.43
G0F	0.00	1.25	1.52
G1F-GlcNAc	0.00	11.77	0.00
Man5	0.88	4.02	0.00
G1(a)	0.44	6.50	9.54
G1(b)	0.46	0.43	6.92
G1-GlcNAc	0.00	0.00	11.17
G1F(a)	0.27	0.00	0.00

Lot	14	10	7
Expression System	CHO-S	Tuna293™	ExpiCHO™
<u>Glycan ID¹</u>	% Peak Area	% Peak Area	% Peak Area
G1F(b)	0.00	0.00	0.31
G1S	<u>0.00</u>	<u>1.88</u>	<u>0.87</u>
G2	0.00	7.42	8.59
G2F	6.14	0.00	0.00
G2S(a)	<u>0.28</u>	<u>1.79</u>	<u>0.00</u>
G2S(b)	<u>38.65</u>	<u>1.30</u>	<u>0.00</u>
G2FS	<u>0.67</u>	<u>0.00</u>	<u>0.00</u>
G2S2	<u>39.13</u>	<u>0.00</u>	<u>0.00</u>
G2FS2	<u>0.69</u>	<u>0.00</u>	<u>0.00</u>
G2S + 2 GlcNAc	<u>2.60</u>	<u>0.00</u>	<u>0.00</u>
G2S2+2GlcNAc(a)	<u>0.82</u>	<u>0.00</u>	<u>0.00</u>
G2S2+2GlcNAc(b)	<u>1.60</u>	<u>0.00</u>	<u>0.00</u>
Sum Sialylated Glycans³	84.44	4.97	0.87

¹Identification and peak percentages derived from selective fab glycan (VL) release, labeling, and hydrophilic interaction liquid chromatography (HILIC) method. All identification are based on observed monoisotopic masses and known biosynthetic pathways, however isomeric variants are possible for some entries.

²Unknown, system, and reagent peak totals.

³Sum of sialylated glycans; sum of identified N-glycans terminating in one or more sialic acid (N-acetylneuraminic acid) residues (underlined).

[0389] Table 45B. Light chain Fab glycan assesement comparing Antibodies A-1 and 1.52.64-1

Name	Lot #	Cell Line	Percent Sialylation	Method
A-1	5	Expi293™	67/52	1/2
	7	ExpiCHO™	1/1	1/2
	10	Tuna293™	5	2
	14	CHO-S	84	2
	22	CHO-origin	73	1
1.52.64-1	18-PP21	CHO-origin	49	1
	14525-02	CHO-origin	83	1

¹Reduced Light Chain LC/MS

²Selective Fab Glycan Release, Labeling, and HILIC Chromatography

Example 15: Effects of Fv mutations and Fv-glycosylation profiles on antibody pharmacokinetics

[0390] Antibody A and several engineered antibodies described herein were administered to cynomolgus macaque monkeys to characterize their pharmacokinetic (PK) profiles. In certain cases, Antibody A-1 variants were transiently or stably produced in different expression cell lines to assess the impact of N72-linked Fab glycan sialylation on PK. Percent Fab glycan sialylation was determined using LCMS as described in Example 14. Serum samples collected from monkeys were analyzed using a bioanalytical method of sufficient selectivity and sensitivity to determine serum concentration-time profiles and mean serum PK parameters by non-compartmental PK analysis (NCA). The bioanalytical method utilized clade B gp120 antigen (Immune-tech, CA) as a capture reagent and biotin conjugated goat anti-human IgG antibody (Southern Biotech, AL) as a secondary reagent, with SULFO-TAG labeled Streptavidin (MesoScale Discovery, MD) for electrochemical detection.

[0391] The *in vivo* disposition of Antibody A and several engineered variants transiently expressed in different cell lines was characterized after a single intravenous (IV) administration in three (n=3) naïve male cynomolgus monkeys per group (Covance, TX). The measured mean $\pm$ standard deviation (SD) serum concentration-time profiles is depicted in Figure 7. The pharmacokinetic analysis of Antibody A transiently expressed in Expi293™ (ThermoFisher Scientific, MA) dosed at 0.5 mg/kg IV showed clearance (Cl) values of 17.9 ± 1.0 and corresponding half-life ($t_{1/2}$) of 8.9 ± 1.7 days which were comparable to Antibody A-1 Lot 3, expressed in Expi293™ under similar conditions, with a Cl of 18.7 ± 2.3 mL/day/kg and $t_{1/2}$ of 7.6 ± 0.3 days (Table 46).

[0392] Antibodies with variable domain Fab glycans containing low sialic acid or high mannose may have altered PK (Liu L. 2015. J. Pharm. Sci. 104:1866-1884). Glycan compositions can be altered as a result of protein expression conditions, therefore the *in vivo* disposition of A-1 was evaluated using additional transiently expressed lots characterized for their % Fab glycan sialylation content, namely CHO-S (Lot 14), CHO-origin (Lot 22) (Sigma-Aldrich, MO), and Tuna293™ (Lot 10) (LakePharma, CA), and ExpiCHO™ (Lot 7) (ThermoFisher Scientific, MA). Antibodies were characterized after a single IV dose of 0.5 mg/kg (Lot 14, 22, and 10) or 5.0 mg/kg (Lot 7) in naïve male cynomolgus monkeys (Covance, TX). The measured mean ($\pm$ SD) serum concentration-time profiles of each lot of Antibody A-1 are depicted in Figure 7. Lot 7 was dose normalized for direct comparison. The pharmacokinetic analysis of the tested Antibody

A-1 lots showed variable PK based on % Fab sialylation content (Table 46). Antibody A-1 Lot 14 with 84% Fab glycan sialylation had the lowest clearance (Cl) value of 7.2 ± 0.7 mL/day/kg, while the Cl was progressively faster with Antibody A-1 Lot 22 (73%) with a Cl of 10.7 ± 1.7 , Antibody A-1 Lot 3 with a Cl of 18.7 ± 2.3 mL/day/kg, Antibody A-1 Lot 10 (5%) with a Cl of 68.7 ± 19.8 mL/day/kg, and Antibody A-1 Lot 7 (<1%) with a Cl of 120 ± 46.7 mL/day/kg. The data supports protein expression conditions can impact Fab glycan composition and resultant PK.

[0393] Table 46: Pharmacokinetics of antibody A and several engineered variants after IV administration in naïve male cynomolgus monkeys (n=3).

Test Article	Expression System	Lot	IV Dose (mg/kg)	Cl (mL/day/kg)	% Fab Sialylation
A	Expi293 TM	5	0.5	17.9 ± 1.0	ND
A-1	Expi293 TM	3	0.5	18.7 ± 2.3	ND
A-1	CHO-S	14	0.5	7.2 ± 0.7	84
A-1	CHO-origin	22	0.5	10.7 ± 1.7	73
A-1	Tuna293 TM	10	0.5	68.7 ± 20	5
A-1	ExpiCHO TM	7	5	120 ± 47	<1
1.1.54-1	ExpiCHO TM	3	5	12 ± 1	ND
1.37.51-1	ExpiCHO TM	3	5	15 ± 12	ND

ND = not determined

[0394] To evaluate the impact of protein modifications aimed to remove the variable domain N72-linked glycan and polyspecificity, in vivo PK of 1.1.54-1 and 1.37.51-1 (two antibodies without the N72-linked glycan removed) was evaluated. Both antibodies were transiently expressed in the ExpiCHOTM mammalian cell expression system under similar conditions which resulted in reduced PK of A-1 (Lot 7, above). Antibodies were characterized after a single IV bolus dose of 5 mg/kg to three naïve male cynomolgus monkeys (Covance, TX). The PK analysis (Table 46) demonstrated that 1.1.54-1 and 1.37.51-1 were comparable in Cl (12 ± 1 and 15 ± 12 mL/day/kg, respectively), yet significantly improved over A-1 Lot 7 (Cl of 120 ± 47 mL/day/kg), supporting that protein modifications which remove the variable domain N72-linked glycan can improve the PK of the antibody variants described herein. Removing the glycan did not achieve the same clearance as the highly sialylated lots, supporting that the N72-linked glycan may be present to reduce non-specific protein interactions.

[0395] The PK of 1.52.64-1 (Lot 4) derived from transient expression in CHO-S, or 1.52.64-1 from a stable pool of CHO-origin cells (Lot 18-PP21) or from a clonally selected CHO-origin cell line (Lot 14525-32) was studied following a single IV administration in naïve male and female cynomolgus monkeys (n=3). The mean $\pm$ SD serum concentration-time profiles for days 0-14 are presented in Figure 8. Results of the NCA are depicted in Table 47. 1.52.64-1 (Lot 4) contained approximately 75% Fab sialylation. 1.52.64-1 Lot 4 dosed at 0.5 mg/kg IV slow bolus resulted in a cynomolgus monkey clearance of 7.8 ± 0.6 mL/day/kg; equivalent to A-1 Lot 14 (7.2 ± 0.7 mL/day/kg) expressed in CHO-S under similar conditions.

[0396] Table 47: Pharmacokinetic parameters of three lots of 1.52.64-1 following IV administrations in naïve male and female cynomolgus monkeys (n=3).

Expression system	Lot	IV Dose (mg/kg)	Cl (mL/day/kg)	% Fab Sialylation
CHO-S transient transfection	Lot 4	0.5	7.8 ± 0.6	75
CHO-origin stable pool	Lot 18-PP21	30	20.8 ± 9.5	49
CHO-origin stable clone	Lot 14525-32	30	7.9 ± 1.3	84

[0397] 1.52.64-1 Lot 18-PP21 yielded material with approximately 49% Fab sialylation while Lot 14525-32 yielded material with approximately 84% Fab sialylation from the CHO-origin stable expression system. 1.52.64-1 Lot 18-PP21 and Lot 14525-32 were administered via a 30 minute IV infusion at 30 mg/kg. PK analysis revealed that Lot 18-PP21 had reduced exposure relative to Lot 14525-32 due to the increased clearance of 20.8 ± 9.5 mL/day/kg compared to 7.9 ± 1.3 mL/day/kg, respectively. The increased clearance is consistent with the reduced % Fab glycan sialylation (49% vs 84%). The totality of the preclinical PK assessments demonstrate that antibody A variants containing a Fab glycan structure require controlled protein production conditions to yield antibodies with high Fab glycan sialylation (*e.g.* $\geq 75\%$) that will achieve desirable antibody pharmacokinetics.

Example 16: Selection of High Sialylation Cell Lines

[0398] In view of the foregoing data and analyses, we isolated cell lines to produce highly sialylated antibody. To accomplish this, cell line development (CLD) was biased towards identification of cell lines that express highly sialylated anti-gp120 antibodies, as described herein. Briefly, the CHO-based development cell line was transfected with a vector encoding the heavy host and light chains of antibody variants described herein. Multiple stable pools were assessed for bioreactor performance and product quality (including % sialylation). Stable pools expressing antibody having a high level of sialylation (*e.g.*, at least about 75 % sialylated) were selected for clone generation. In order to further bias clonal cell line isolation towards higher sialylation, clonal cell lines generated from the parent stable pool with the highest % sialylation (approximately 95 % sialylated) were over-represented throughout the clone generation workflow. Multiple clonal cell lines were assessed for bioreactor performance and product quality (including % sialylation) and a clonal cell line expressing highly sialylated antibody (>85 %) was selected as the lead cell line for master cell bank (MCB) manufacturing.

OTHER EMBODIMENTS

[0399] While the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims.

Claims

1. An antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody or antigen-binding fragment thereof comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in:

- (i) SEQ ID NOs.: 159, 138, 139, 140, 141, and 142, respectively;
- (ii) SEQ ID NOs.: 137, 160, 139, 140, 141, and 142, respectively;
- (iii) SEQ ID NOs.: 137, 161, 139, 140, 141, and 142, respectively;
- (iv) SEQ ID NOs.: 137, 162, 139, 140, 141, and 142, respectively;
- (v) SEQ ID NOs.: 137, 163, 139, 140, 141, and 142, respectively;
- (vi) SEQ ID NOs.: 137, 138, 164, 140, 141, and 142, respectively;
- (vii) SEQ ID NOs.: 159, 138, 164, 140, 141, and 142, respectively;
- (viii) SEQ ID NOs.: 137, 138, 139, 140, 165, and 142, respectively;
- (ix) SEQ ID NOs.: 137, 138, 139, 140, 166, and 142, respectively;
- (x) SEQ ID NOs.: 137, 138, 139, 140, 167, and 142, respectively;
- (xi) SEQ ID NOs.: 137, 138, 139, 140, 168, and 142, respectively;
- (xii) SEQ ID NOs.: 137, 138, 154, 140, 141, and 142, respectively; or
- (xiii) SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively; and

optionally includes in framework region 3 (FR3) of the VH at position corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO:453 or SEQ ID NO: 627.

2. The antibody or antigen-binding fragment thereof of claim 1, wherein the VH comprises one or more of the following amino acids at the indicated positions (position numbering according to Kabat): valine at position 5, glutamic acid at position 10, lysine at position 12, lysine at position 23, asparagine at position 28, arginine at position 30, tyrosine at position 32, threonine at position 68, methionine at position 69, histidine at position 72, phenylalanine at position 76, alanine at position 78, serine at position 82a, arginine at position 82b, threonine at position 89, tyrosine at position 99, glutamine at position 105, or methionine at position 108.

3. The antibody or antigen-binding fragment thereof of claim 1 or 2, comprising FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASWDFDTSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 628) or
RVSLTRHASFDFDTSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629).

4. The antibody or antigen-binding fragment thereof of any one of claims 1 to 3, wherein the VL comprises one or more of the following amino acids at the indicated positions (position numbering according to Kabat): arginine at position 18, lysine at position 39, proline at position 40, threonine at position 56, serine at position 65, threonine at position 72, serine at position 76, serine at position 77, threonine at position 99, glycine at position 99, asparagine at position 103, or isoleucine at position 106.

5. The antibody or antigen-binding fragment thereof of any one of claims 1 to 4, wherein the VL comprises one or more of the following amino acids at the indicated positions (position numbering according to Kabat): arginine at position 18, alanine at position 19, serine at position 65, threonine or histidine at position 72, lysine at position 74, serine at position 76, serine at position 77, phenylalanine at position 98, or glycine at position 99.

6. The antibody or antigen-binding fragment thereof of any one of claims 1 to 5, wherein the VL comprises an amino acid sequence set forth in any one of SEQ ID NOs.: 332 to 342.

7. The antibody or antigen-binding fragment thereof of any one of claims 1 to 6, comprising a human IgG1 Fc region.

8. The antibody or antigen-binding fragment thereof of claim 7, wherein the human IgG1 Fc region is IgG1m17 (SEQ ID NO: 348).

9. The antibody or antigen-binding fragment thereof of any one of claims 1 to 8, comprising a human IgG1 Fc region comprising (position numbered according to EU numbering):

(i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330;

(ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434;

(iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434;

(iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434;

(v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or

(vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434.

10. The antibody or antigen-binding fragment thereof of any one of claims 1 to 7, comprising a human kappa light chain constant region.

11. The antibody or antigen-binding fragment thereof of claim 10, wherein the human kappa light chain constant region is Km3 (SEQ ID NO:351).

12. The antibody or antigen-binding fragment thereof of any one of claims 1 to 11, wherein the antibody or antigen-binding fragment has greater killing potency of HIV-infected cells than Antibody A.

13. The antibody or antigen-binding fragment thereof of any one of claims 1 to 12, wherein the antibody or antigen-binding fragment has an improved, extended, enhanced or increased serum half-life in a human than Antibody A.

14. The antibody or antigen-binding fragment thereof of any one of claims 1 to 13, wherein the antibody or antigen-binding fragment has a serum half-life in a human of at least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer.

15. An antibody that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in:

(i) SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively; or

(ii) SEQ ID NOs.: 153, 138, 154, 140, 141, and 142, respectively,

wherein the antibody comprises a human IgG1 Fc region comprising (position numbered according to EU numbering):

(i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330;

(ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434;

(iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434;

(iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434;

(v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or

(vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434.

16. The antibody of claim 15, wherein the antibody comprises a light chain comprising an alanine at position 19 (Kabat numbering).

17. The antibody of any one of claims 15 to 16, comprising in framework region 3 (FR3) of the VH at positions corresponding to 74a, 74b, 74c, and 74d (Kabat numbering) the amino acid sequence set forth in SEQ ID NO:453 or SEQ ID NO: 627.

18. The antibody of any one of claims 15 to 17, comprising a FR3 of the VH comprising the following amino acid sequence:

RVSLTRHASWDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 628) or
RVSLTRHASFDFDTFSFYMDLKALRSDDTAVYFCAR (SEQ ID NO: 629).

19. The antibody of any one of claims 15 to 18, wherein the antibody comprises a VL amino acid sequence set forth in any one of SEQ ID NOs.: 332 to 342.

20. The antibody of any one of claims 15 to 19, wherein the human IgG1 Fc region is IgG1m17 (SEQ ID NO:348).

21. The antibody of any one of claims 15 to 20, wherein the antibody comprises a human kappa light chain constant region.

22. The antibody of claim 21, wherein the human kappa light chain constant region is Km3 (SEQ ID NO:351).

23. The antibody of any one of claims 15 to 22, wherein the antibody has greater killing potency of HIV-infected cells than Antibody A and/or Antibody B.

24. The antibody or antigen-binding fragment thereof of any one of claims 15 to 23, wherein the antibody or antigen-binding fragment has an improved, extended, enhanced or increased serum half-life in a human than Antibody A and/or Antibody B.

25. The antibody or antigen-binding fragment thereof of any one of claims 15 to 24, wherein the antibody or antigen-binding fragment has a serum half-life in a human of at least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer.

26. An antibody or an antigen-binding fragment thereof, the antibody or antigen-binding fragment thereof comprising a heavy chain variable region (VH) and a light chain variable region (VL), wherein the VH and VL comprise the amino acid sequences set forth, respectively, below:

- (1) SEQ ID NOs.: 184 and 223;
- (2) SEQ ID NOs.: 185 and 223;
- (3) SEQ ID NOs.: 182 and 225;
- (4) SEQ ID NOs.: 185 and 225;
- (5) SEQ ID NOs.: 186 and 223;
- (6) SEQ ID NOs.: 187 and 223;
- (7) SEQ ID NOs.: 188 and 223;
- (8) SEQ ID NOs.: 189 and 223;
- (9) SEQ ID NOs.: 190 and 223;
- (10) SEQ ID NOs.: 191 and 223;
- (11) SEQ ID NOs.: 192 and 223;
- (12) SEQ ID NOs.: 193 and 223;
- (13) SEQ ID NOs.: 194 and 223;
- (14) SEQ ID NOs.: 195 and 223;
- (15) SEQ ID NOs.: 196 and 223;
- (16) SEQ ID NOs.: 197 and 223;
- (17) SEQ ID NOs.: 198 and 223;
- (18) SEQ ID NOs.: 199 and 223;
- (19) SEQ ID NOs.: 200 and 223;
- (20) SEQ ID NOs.: 201 and 223;

- (21) SEQ ID NOs.: 202 and 223;
- (22) SEQ ID NOs.: 203 and 223;
- (23) SEQ ID NOs.: 204 and 223;
- (24) SEQ ID NOs.: 205 and 223;
- (25) SEQ ID NOs.: 206 and 223;
- (26) SEQ ID NOs.: 207 and 223;
- (27) SEQ ID NOs.: 208 and 223;
- (28) SEQ ID NOs.: 209 and 223;
- (29) SEQ ID NOs.: 182 and 226;
- (30) SEQ ID NOs.: 182 and 227;
- (31) SEQ ID NOs.: 182 and 229;
- (32) SEQ ID NOs.: 182 and 230;
- (33) SEQ ID NOs.: 182 and 231;
- (34) SEQ ID NOs.: 182 and 232;
- (35) SEQ ID NOs.: 182 and 233;
- (36) SEQ ID NOs.: 182 and 234;
- (37) SEQ ID NOs.: 182 and 235;
- (38) SEQ ID NOs.: 182 and 236;
- (39) SEQ ID NOs.: 182 and 237;
- (40) SEQ ID NOs.: 182 and 238;
- (41) SEQ ID NOs.: 182 and 239;
- (42) SEQ ID NOs.: 182 and 240;
- (43) SEQ ID NOs.: 182 and 241;
- (44) SEQ ID NOs.: 182 and 242;
- (45) SEQ ID NOs.: 182 and 243;
- (46) SEQ ID NOs.: 182 and 244;
- (47) SEQ ID NOs.: 182 and 245;
- (48) SEQ ID NOs.: 182 and 246;
- (49) SEQ ID NOs.: 182 and 247;
- (50) SEQ ID NOs.: 182 and 248;
- (51) SEQ ID NOs.: 182 and 249;
- (52) SEQ ID NOs.: 182 and 250;
- (53) SEQ ID NOs.: 182 and 251;
- (54) SEQ ID NOs.: 182 and 252;
- (55) SEQ ID NOs.: 182 and 253;

- (56) SEQ ID NOs.: 210 and 238;
- (57) SEQ ID NOs.: 211 and 238;
- (58) SEQ ID NOs.: 212 and 238;
- (59) SEQ ID NOs.: 210 and 240;
- (60) SEQ ID NOs.: 211 and 240;
- (61) SEQ ID NOs.: 212 and 240;
- (62) SEQ ID NOs.: 213 and 223;
- (63) SEQ ID NOs.: 214 and 223;
- (64) SEQ ID NOs.: 215 and 223;
- (65) SEQ ID NOs.: 216 and 223;
- (66) SEQ ID NOs.: 217 and 223;
- (67) SEQ ID NOs.: 218 and 223;
- (68) SEQ ID NOs.: 182 and 254;
- (69) SEQ ID NOs.: 213 and 254;
- (70) SEQ ID NOs.: 214 and 254;
- (71) SEQ ID NOs.: 215 and 254;
- (72) SEQ ID NOs.: 216 and 254;
- (73) SEQ ID NOs.: 217 and 254;
- (74) SEQ ID NOs.: 218 and 254;
- (75) SEQ ID NOs.: 182 and 255;
- (76) SEQ ID NOs.: 213 and 255;
- (77) SEQ ID NOs.: 214 and 255;
- (78) SEQ ID NOs.: 215 and 255;
- (79) SEQ ID NOs.: 216 and 255;
- (80) SEQ ID NOs.: 217 and 255;
- (81) SEQ ID NOs.: 218 and 255;
- (82) SEQ ID NOs.: 182 and 256;
- (83) SEQ ID NOs.: 213 and 256;
- (84) SEQ ID NOs.: 214 and 256;
- (85) SEQ ID NOs.: 215 and 256;
- (86) SEQ ID NOs.: 216 and 256;
- (87) SEQ ID NOs.: 217 and 256;
- (88) SEQ ID NOs.: 218 and 256;
- (89) SEQ ID NOs.: 182 and 257;
- (90) SEQ ID NOs.: 213 and 257;

- (91) SEQ ID NOs.: 214 and 257;
- (92) SEQ ID NOs.: 215 and 257;
- (93) SEQ ID NOs.: 216 and 257;
- (94) SEQ ID NOs.: 217 and 257;
- (95) SEQ ID NOs.: 218 and 257;
- (96) SEQ ID NOs.: 182 and 258;
- (97) SEQ ID NOs.: 213 and 258;
- (98) SEQ ID NOs.: 214 and 258;
- (99) SEQ ID NOs.: 215 and 258;
- (100) SEQ ID NOs.: 216 and 258;
- (101) SEQ ID NOs.: 217 and 258;
- (102) SEQ ID NOs.: 218 and 258;
- (103) SEQ ID NOs.: 182 and 259;
- (104) SEQ ID NOs.: 213 and 259;
- (105) SEQ ID NOs.: 214 and 259;
- (106) SEQ ID NOs.: 215 and 259;
- (107) SEQ ID NOs.: 216 and 259;
- (108) SEQ ID NOs.: 217 and 259;
- (109) SEQ ID NOs.: 218 and 259;
- (110) SEQ ID NOs.: 182 and 260;
- (111) SEQ ID NOs.: 182 and 261;
- (112) SEQ ID NOs.: 182 and 262;
- (113) SEQ ID NOs.: 182 and 263;
- (114) SEQ ID NOs.: 182 and 264;
- (115) SEQ ID NOs.: 182 and 265;
- (116) SEQ ID NOs.: 182 and 266;
- (117) SEQ ID NOs.: 182 and 267;
- (118) SEQ ID NOs.: 182 and 268;
- (119) SEQ ID NOs.: 182 and 269;
- (120) SEQ ID NOs.: 182 and 270;
- (121) SEQ ID NOs.: 182 and 271;
- (122) SEQ ID NOs.: 182 and 272;
- (123) SEQ ID NOs.: 219 and 273;
- (124) SEQ ID NOs.: 191 and 274;
- (125) SEQ ID NOs.: 182 and 275;

- (126) SEQ ID NOs.: 220 and 277;
- (127) SEQ ID NOs.: 182 and 278;
- (128) SEQ ID NOs.: 182 and 279;
- (129) SEQ ID NOs.: 182 and 280;
- (130) SEQ ID NOs.: 182 and 281;
- (131) SEQ ID NOs.: 182 and 282;
- (132) SEQ ID NOs.: 221 and 228;
- (133) SEQ ID NOs.: 221 and 283;
- (134) SEQ ID NOs.: 182 and 284;
- (135) SEQ ID NOs.: 221 and 285;
- (136) SEQ ID NOs.: 182 and 286;
- (137) SEQ ID NOs.: 221 and 287;
- (138) SEQ ID NOs.: 221 and 288;
- (139) SEQ ID NOs.: 221 and 289;
- (140) SEQ ID NOs.: 182 and 290;
- (141) SEQ ID NOs.: 221 and 291;
- (142) SEQ ID NOs.: 182 and 292;
- (143) SEQ ID NOs.: 221 and 293;
- (144) SEQ ID NOs.: 221 and 294;
- (145) SEQ ID NOs.: 221 and 295;
- (146) SEQ ID NOs.: 182 and 296;
- (147) SEQ ID NOs.: 221 and 297;
- (148) SEQ ID NOs.: 182 and 298;
- (149) SEQ ID NOs.: 221 and 299;
- (150) SEQ ID NOs.: 221 and 300;
- (151) SEQ ID NOs.: 221 and 301;
- (152) SEQ ID NOs.: 182 and 302;
- (153) SEQ ID NOs.: 221 and 303;
- (154) SEQ ID NOs.: 182 and 304;
- (155) SEQ ID NOs.: 221 and 305;
- (156) SEQ ID NOs.: 182 and 306;
- (157) SEQ ID NOs.: 182 and 307;
- (158) SEQ ID NOs.: 182 and 308;
- (159) SEQ ID NOs.: 182 and 309;
- (160) SEQ ID NOs.: 220 and 310;

(161) SEQ ID NOs.: 220 and 311;
(162) SEQ ID NOs.: 182 and 228;
(163) SEQ ID NOs.: 465 and 276;
(164) SEQ ID NOs.: 466 and 276;
(165) SEQ ID NOs.: 182 and 479;
(166) SEQ ID NOs.: 465 and 479;
(167) SEQ ID NOs.: 466 and 479;
(168) SEQ ID NOs.: 182 and 480;
(169) SEQ ID NOs.: 465 and 480;
(170) SEQ ID NOs.: 466 and 480;
(171) SEQ ID NOs.: 182 and 481;
(172) SEQ ID NOs.: 182 and 482;
(173) SEQ ID NOs.: 465 and 482;
(174) SEQ ID NOs.: 466 and 482;
(175) SEQ ID NOs.: 182 and 483;
(176) SEQ ID NOs.: 182 and 484;
(177) SEQ ID NOs.: 465 and 484;
(178) SEQ ID NOs.: 466 and 484;
(179) SEQ ID NOs.: 182 and 485;
(180) SEQ ID NOs.: 182 and 486;
(181) SEQ ID NOs.: 465 and 486;
(182) SEQ ID NOs.: 466 and 486;
(183) SEQ ID NOs.: 182 and 487;
(184) SEQ ID NOs.: 182 and 488;
(185) SEQ ID NOs.: 465 and 488;
(186) SEQ ID NOs.: 466 and 488;
(187) SEQ ID NOs.: 182 and 489;
(188) SEQ ID NOs.: 465 and 489;
(189) SEQ ID NOs.: 466 and 489;
(190) SEQ ID NOs.: 182 and 491;
(191) SEQ ID NOs.: 465 and 491;
(192) SEQ ID NOs.: 466 and 491;
(193) SEQ ID NOs.: 182 and 492;
(194) SEQ ID NOs.: 465 and 492;
(195) SEQ ID NOs.: 466 and 492;

(196) SEQ ID NOs.: 182 and 493;
(197) SEQ ID NOs.: 182 and 494;
(198) SEQ ID NOs.: 465 and 494;
(199) SEQ ID NOs.: 466 and 494;
(200) SEQ ID NOs.: 182 and 277;
(201) SEQ ID NOs.: 465 and 277;
(202) SEQ ID NOs.: 466 and 277;
(203) SEQ ID NOs.: 182 and 495;
(204) SEQ ID NOs.: 465 and 495;
(205) SEQ ID NOs.: 466 and 495;
(206) SEQ ID NOs.: 182 and 496;
(207) SEQ ID NOs.: 465 and 496;
(208) SEQ ID NOs.: 466 and 496;
(209) SEQ ID NOs.: 182 and 497;
(210) SEQ ID NOs.: 465 and 497;
(211) SEQ ID NOs.: 466 and 497;
(212) SEQ ID NOs.: 182 and 498;
(213) SEQ ID NOs.: 182 and 499;
(214) SEQ ID NOs.: 465 and 499;
(215) SEQ ID NOs.: 466 and 499;
(216) SEQ ID NOs.: 182 and 500;
(217) SEQ ID NOs.: 182 and 501;
(218) SEQ ID NOs.: 465 and 501;
(219) SEQ ID NOs.: 466 and 501;
(220) SEQ ID NOs.: 182 and 502;
(221) SEQ ID NOs.: 182 and 503;
(222) SEQ ID NOs.: 182 and 504;
(223) SEQ ID NOs.: 182 and 505;
(224) SEQ ID NOs.: 182 and 506;
(225) SEQ ID NOs.: 182 and 507;
(226) SEQ ID NOs.: 182 and 508;
(227) SEQ ID NOs.: 182 and 509;
(228) SEQ ID NOs.: 182 and 510;
(229) SEQ ID NOs.: 182 and 511;
(230) SEQ ID NOs.: 182 and 512;

- (231) SEQ ID NOs.: 182 and 513;
- (232) SEQ ID NOs.: 182 and 514;
- (233) SEQ ID NOs.: 182 and 515;
- (234) SEQ ID NOs.: 467 and 223;
- (235) SEQ ID NOs.: 468 and 223;
- (236) SEQ ID NOs.: 469 and 223;
- (237) SEQ ID NOs.: 470 and 223;
- (238) SEQ ID NOs.: 471 and 223;
- (239) SEQ ID NOs.: 472 and 223;
- (240) SEQ ID NOs.: 473 and 223;
- (241) SEQ ID NOs.: 474 and 223;
- (242) SEQ ID NOs.: 475 and 223;
- (243) SEQ ID NOs.: 476 and 223;
- (244) SEQ ID NOs.: 182 and 516;
- (245) SEQ ID NOs.: 182 and 276;
- (246) SEQ ID NOs.: 182 and 569;
- (247) SEQ ID NOs.: 477 and 223;
- (248) SEQ ID NOs.: 477 and 278;
- (249) SEQ ID NOs.: 477 and 292; or
- (250) SEQ ID NOs.: 478 and 276.

27. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 275, respectively.

28. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 278, respectively.

29. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 223, respectively.

30. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 292, respectively.

31. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 465 and 276, respectively.

32. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 466 and 276, respectively.

33. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 491, respectively.

34. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 465 and 491, respectively.

35. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 466 and 491, respectively.

36. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 493, respectively.

37. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 220 and 276, respectively.

38. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 182 and 516, respectively.

39. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 477 and 278, respectively.

40. The antibody or antigen-binding fragment of claim 26, wherein the VH and VL comprise the amino acid sequence set forth in SEQ ID NOs.: 478 and 276, respectively.

41. The antibody or antigen-binding fragment thereof of any one of claims 1 to 40, comprising a VH that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and a VL that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569.

42. The antibody or antigen-binding fragment thereof of any one of claims 26 to 41, comprising a human IgG1 Fc region.

43. The antibody or antigen-binding fragment thereof of claim 42, wherein the human IgG1 Fc region is IgG1m17 (SEQ ID NO:348).

44. The antibody or antigen-binding fragment thereof of any one of claims 26 to 43, comprising a human IgG1 Fc region comprising (position numbered according to EU numbering):

(i) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330;

(ii) aspartic acid at position 239, glutamic acid at position 332, leucine at position 428, and serine at position 434;

(iii) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 428, and serine at position 434;

(iv) aspartic acid at position 239, glutamic acid at position 332, leucine at position 330, leucine at position 428, and serine at position 434;

(v) aspartic acid at position 239, glutamic acid at position 332, alanine at position 236, leucine at position 330, leucine at position 428, and serine at position 434; or

(vi) leucine at position 243, proline at position 292, leucine at position 300, isoleucine at position 305, leucine at position 396, leucine at position 428, and serine at position 434.

45. The antibody or antigen-binding fragment thereof of any one of claims 26 to 44, wherein the antibody comprises a human kappa light chain constant region.

46. The antibody or antigen-binding fragment thereof of claim 45, wherein the human kappa light chain constant region is Km3 (SEQ ID NO:351).

47. The antibody or antigen-binding fragment thereof of any one of claims 26 to 46, wherein the antibody or antigen-binding fragment has greater killing potency of HIV-infected cells than Antibody A and/or Antibody B.

48. The antibody or antigen-binding fragment thereof of any one of claims 26 to 47, wherein the antibody or antigen-binding fragment has an improved, extended, enhanced or increased serum half-life in a human than Antibody A and/or Antibody B.

49. The antibody or antigen-binding fragment thereof of any one of claims 26 to 48, wherein the antibody or antigen-binding fragment has a serum half-life in a human of at

least about 3 days, *e.g.*, at least about 4 days, at least about 5 days, at least about 6 days, at least about 7 days, at least about 8 days, at least about 9 days, at least about 10 days, at least about 12 days, at least about 14 days, at least about 16 days, at least about 18 days, at least about 20 days, at least about 21 days, at least about 24 days, at least about 28 days, at least about 30 days, or longer.

50. An antigen-binding fragment of any one of claims 1 to 49, comprising a scFv, sc(Fv)₂, Fab, F(ab)₂, Fab', F(ab')₂, Facb or Fv fragment.

51. An antibody, the antibody comprising a heavy chain and a light chain, wherein the heavy chain and the light chain comprise the amino acid sequences set forth, respectively, below:

- (1) SEQ ID NOs.: 2 and 49;
- (2) SEQ ID NOs.: 5 and 49;
- (3) SEQ ID NOs.: 6 and 49;
- (4) SEQ ID NOs.: 2 and 50;
- (5) SEQ ID NOs.: 6 and 50;
- (6) SEQ ID NOs.: 7 and 49;
- (7) SEQ ID NOs.: 8 and 49;
- (8) SEQ ID NOs.: 9 and 49;
- (9) SEQ ID NOs.: 10 and 49;
- (10) SEQ ID NOs.: 11 and 49;
- (11) SEQ ID NOs.: 12 and 49;
- (12) SEQ ID NOs.: 13 and 49;
- (13) SEQ ID NOs.: 14 and 49;
- (14) SEQ ID NOs.: 15 and 49;
- (15) SEQ ID NOs.: 16 and 49;
- (16) SEQ ID NOs.: 17 and 49;
- (17) SEQ ID NOs.: 18 and 49;
- (18) SEQ ID NOs.: 19 and 49;
- (19) SEQ ID NOs.: 20 and 49;
- (20) SEQ ID NOs.: 21 and 49;
- (21) SEQ ID NOs.: 22 and 49;
- (22) SEQ ID NOs.: 23 and 49;
- (23) SEQ ID NOs.: 24 and 49;
- (24) SEQ ID NOs.: 25 and 49;

- (25) SEQ ID NOs.: 26 and 49;
- (26) SEQ ID NOs.: 27 and 49;
- (27) SEQ ID NOs.: 28 and 49;
- (28) SEQ ID NOs.: 29 and 49;
- (29) SEQ ID NOs.: 30 and 49;
- (30) SEQ ID NOs.: 2 and 51;
- (31) SEQ ID NOs.: 2 and 52;
- (32) SEQ ID NOs.: 2 and 53;
- (33) SEQ ID NOs.: 2 and 54;
- (34) SEQ ID NOs.: 2 and 55;
- (35) SEQ ID NOs.: 2 and 56;
- (36) SEQ ID NOs.: 2 and 57;
- (37) SEQ ID NOs.: 2 and 58;
- (38) SEQ ID NOs.: 2 and 59;
- (39) SEQ ID NOs.: 2 and 60;
- (40) SEQ ID NOs.: 2 and 61;
- (41) SEQ ID NOs.: 2 and 62;
- (42) SEQ ID NOs.: 2 and 63;
- (43) SEQ ID NOs.: 2 and 64;
- (44) SEQ ID NOs.: 2 and 65;
- (45) SEQ ID NOs.: 2 and 66;
- (46) SEQ ID NOs.: 2 and 67;
- (47) SEQ ID NOs.: 2 and 68;
- (48) SEQ ID NOs.: 2 and 69;
- (49) SEQ ID NOs.: 2 and 70;
- (50) SEQ ID NOs.: 2 and 71;
- (51) SEQ ID NOs.: 2 and 72;
- (52) SEQ ID NOs.: 2 and 73;
- (53) SEQ ID NOs.: 2 and 74;
- (54) SEQ ID NOs.: 2 and 75;
- (55) SEQ ID NOs.: 2 and 76;
- (56) SEQ ID NOs.: 2 and 77;
- (57) SEQ ID NOs.: 2 and 78;
- (58) SEQ ID NOs.: 31 and 63;
- (59) SEQ ID NOs.: 32 and 63;

- (60) SEQ ID NOs.: 33 and 63;
- (61) SEQ ID NOs.: 31 and 65;
- (62) SEQ ID NOs.: 32 and 65;
- (63) SEQ ID NOs.: 33 and 65;
- (64) SEQ ID NOs.: 34 and 49;
- (65) SEQ ID NOs.: 35 and 49;
- (66) SEQ ID NOs.: 36 and 49;
- (67) SEQ ID NOs.: 37 and 49;
- (68) SEQ ID NOs.: 38 and 49;
- (69) SEQ ID NOs.: 39 and 49;
- (70) SEQ ID NOs.: 2 and 79;
- (71) SEQ ID NOs.: 34 and 79;
- (72) SEQ ID NOs.: 35 and 79;
- (73) SEQ ID NOs.: 36 and 79;
- (74) SEQ ID NOs.: 37 and 79;
- (75) SEQ ID NOs.: 38 and 79;
- (76) SEQ ID NOs.: 39 and 79;
- (77) SEQ ID NOs.: 2 and 80;
- (78) SEQ ID NOs.: 34 and 80;
- (79) SEQ ID NOs.: 35 and 80;
- (80) SEQ ID NOs.: 36 and 80;
- (81) SEQ ID NOs.: 37 and 80;
- (82) SEQ ID NOs.: 38 and 80;
- (83) SEQ ID NOs.: 39 and 80;
- (84) SEQ ID NOs.: 2 and 81;
- (85) SEQ ID NOs.: 34 and 81;
- (86) SEQ ID NOs.: 35 and 81;
- (87) SEQ ID NOs.: 36 and 81;
- (88) SEQ ID NOs.: 37 and 81;
- (89) SEQ ID NOs.: 38 and 81;
- (90) SEQ ID NOs.: 39 and 81;
- (91) SEQ ID NOs.: 2 and 82;
- (92) SEQ ID NOs.: 34 and 82;
- (93) SEQ ID NOs.: 35 and 82;
- (94) SEQ ID NOs.: 36 and 82;

- (95) SEQ ID NOs.: 37 and 82;
- (96) SEQ ID NOs.: 38 and 82;
- (97) SEQ ID NOs.: 39 and 82;
- (98) SEQ ID NOs.: 2 and 83;
- (99) SEQ ID NOs.: 34 and 83;
- (100) SEQ ID NOs.: 35 and 83;
- (101) SEQ ID NOs.: 36 and 83;
- (102) SEQ ID NOs.: 37 and 83;
- (103) SEQ ID NOs.: 38 and 83;
- (104) SEQ ID NOs.: 39 and 83;
- (105) SEQ ID NOs.: 2 and 84;
- (106) SEQ ID NOs.: 34 and 84;
- (107) SEQ ID NOs.: 35 and 84;
- (108) SEQ ID NOs.: 36 and 84;
- (109) SEQ ID NOs.: 37 and 84;
- (110) SEQ ID NOs.: 38 and 84
- (111) SEQ ID NOs.: 39 and 84;
- (112) SEQ ID NOs.: 2 and 85;
- (113) SEQ ID NOs.: 2 and 86;
- (114) SEQ ID NOs.: 2 and 87;
- (115) SEQ ID NOs.: 2 and 88;
- (116) SEQ ID NOs.: 2 and 89;
- (117) SEQ ID NOs.: 2 and 90;
- (118) SEQ ID NOs.: 2 and 91;
- (119) SEQ ID NOs.: 2 and 92;
- (120) SEQ ID NOs.: 2 and 93;
- (121) SEQ ID NOs.: 2 and 94;
- (122) SEQ ID NOs.: 2 and 95;
- (123) SEQ ID NOs.: 2 and 96;
- (124) SEQ ID NOs.: 2 and 97;
- (125) SEQ ID NOs.: 40 and 98;
- (126) SEQ ID NOs.: 12 and 99;
- (127) SEQ ID NOs.: 2 and 100;
- (128) SEQ ID NOs.: 41 and 49;
- (129) SEQ ID NOs.: 42 and 101;

- (130) SEQ ID NOs.: 42 and 102;
- (131) SEQ ID NOs.: 2 and 103;
- (132) SEQ ID NOs.: 2 and 104;
- (133) SEQ ID NOs.: 2 and 105;
- (134) SEQ ID NOs.: 2 and 106;
- (135) SEQ ID NOs.: 2 and 107;
- (136) SEQ ID NOs.: 43 and 49;
- (137) SEQ ID NOs.: 44 and 49;
- (138) SEQ ID NOs.: 45 and 49;
- (139) SEQ ID NOs.: 46 and 49;
- (140) SEQ ID NOs.: 47 and 53;
- (141) SEQ ID NOs.: 47 and 108;
- (142) SEQ ID NOs.: 2 and 109;
- (143) SEQ ID NOs.: 47 and 110;
- (144) SEQ ID NOs.: 2 and 111;
- (145) SEQ ID NOs.: 47 and 112;
- (146) SEQ ID NOs.: 47 and 113;
- (147) SEQ ID NOs.: 47 and 114;
- (148) SEQ ID NOs.: 2 and 115;
- (149) SEQ ID NOs.: 47 and 116;
- (150) SEQ ID NOs.: 2 and 117;
- (151) SEQ ID NOs.: 47 and 118;
- (152) SEQ ID NOs.: 47 and 119;
- (153) SEQ ID NOs.: 47 and 120;
- (154) SEQ ID NOs.: 2 and 121;
- (155) SEQ ID NOs.: 47 and 122;
- (156) SEQ ID NOs.: 2 and 123;
- (157) SEQ ID NOs.: 47 and 124;
- (158) SEQ ID NOs.: 47 and 125;
- (159) SEQ ID NOs.: 47 and 126;
- (160) SEQ ID NOs.: 2 and 127;
- (161) SEQ ID NOs.: 47 and 128;
- (162) SEQ ID NOs.: 2 and 129;
- (163) SEQ ID NOs.: 47 and 130;
- (164) SEQ ID NOs.: 2 and 131;

- (165) SEQ ID NOs.: 2 and 132;
- (166) SEQ ID NOs.: 2 and 133;
- (167) SEQ ID NOs.: 2 and 134;
- (168) SEQ ID NOs.: 42 and 135;
- (169) SEQ ID NOs.: 42 and 136;
- (170) SEQ ID NOs.: 517 and 101;
- (171) SEQ ID NOs.: 518 and 101;
- (172) SEQ ID NOs.: 2 and 531;
- (173) SEQ ID NOs.: 517 and 531;
- (174) SEQ ID NOs.: 518 and 531;
- (175) SEQ ID NOs.: 2 and 532;
- (176) SEQ ID NOs.: 517 and 532;
- (177) SEQ ID NOs.: 518 and 532;
- (178) SEQ ID NOs.: 2 and 533;
- (179) SEQ ID NOs.: 2 and 534;
- (180) SEQ ID NOs.: 517 and 534;
- (181) SEQ ID NOs.: 518 and 534;
- (182) SEQ ID NOs.: 2 and 535;
- (183) SEQ ID NOs.: 2 and 536;
- (184) SEQ ID NOs.: 517 and 536;
- (185) SEQ ID NOs.: 518 and 536;
- (186) SEQ ID NOs.: 2 and 537;
- (187) SEQ ID NOs.: 2 and 538;
- (188) SEQ ID NOs.: 517 and 538;
- (189) SEQ ID NOs.: 518 and 538;
- (190) SEQ ID NOs.: 2 and 539;
- (191) SEQ ID NOs.: 2 and 540;
- (192) SEQ ID NOs.: 517 and 540;
- (193) SEQ ID NOs.: 518 and 540;
- (194) SEQ ID NOs.: 2 and 541;
- (195) SEQ ID NOs.: 517 and 541;
- (196) SEQ ID NOs.: 518 and 541;
- (197) SEQ ID NOs.: 2 and 542;
- (198) SEQ ID NOs.: 517 and 542;
- (199) SEQ ID NOs.: 518 and 542;

- (200) SEQ ID NOs.: 2 and 543;
- (201) SEQ ID NOs.: 517 and 543;
- (202) SEQ ID NOs.: 518 and 543;
- (203) SEQ ID NOs.: 2 and 544;
- (204) SEQ ID NOs.: 2 and 545;
- (205) SEQ ID NOs.: 517 and 545;
- (206) SEQ ID NOs.: 518 and 545;
- (207) SEQ ID NOs.: 2 and 102;
- (208) SEQ ID NOs.: 517 and 102;
- (209) SEQ ID NOs.: 518 and 102;
- (210) SEQ ID NOs.: 2 and 546;
- (211) SEQ ID NOs.: 517 and 546;
- (212) SEQ ID NOs.: 518 and 546;
- (213) SEQ ID NOs.: 2 and 547;
- (214) SEQ ID NOs.: 517 and 547;
- (215) SEQ ID NOs.: 518 and 547;
- (216) SEQ ID NOs.: 2 and 548;
- (217) SEQ ID NOs.: 517 and 548;
- (218) SEQ ID NOs.: 518 and 548;
- (219) SEQ ID NOs.: 2 and 549;
- (220) SEQ ID NOs.: 2 and 550;
- (221) SEQ ID NOs.: 517 and 550;
- (222) SEQ ID NOs.: 518 and 550;
- (223) SEQ ID NOs.: 2 and 551;
- (224) SEQ ID NOs.: 2 and 552;
- (225) SEQ ID NOs.: 517 and 552;
- (226) SEQ ID NOs.: 518 and 552;
- (227) SEQ ID NOs.: 2 and 553;
- (228) SEQ ID NOs.: 2 and 554;
- (229) SEQ ID NOs.: 2 and 555;
- (230) SEQ ID NOs.: 2 and 556;
- (231) SEQ ID NOs.: 2 and 557;
- (232) SEQ ID NOs.: 2 and 558;
- (233) SEQ ID NOs.: 2 and 559;
- (234) SEQ ID NOs.: 2 and 560;

- (235) SEQ ID NOs.: 2 and 561;
- (236) SEQ ID NOs.: 2 and 562;
- (237) SEQ ID NOs.: 2 and 563;
- (238) SEQ ID NOs.: 2 and 564;
- (239) SEQ ID NOs.: 2 and 565;
- (240) SEQ ID NOs.: 2 and 566;
- (241) SEQ ID NOs.: 519 and 49;
- (242) SEQ ID NOs.: 520 and 49;
- (243) SEQ ID NOs.: 521 and 49;
- (244) SEQ ID NOs.: 522 and 49;
- (245) SEQ ID NOs.: 523 and 49;
- (246) SEQ ID NOs.: 524 and 49;
- (247) SEQ ID NOs.: 526 and 49;
- (248) SEQ ID NOs.: 527 and 49;
- (249) SEQ ID NOs.: 528 and 49;
- (250) SEQ ID NOs.: 2 and 567;
- (251) SEQ ID NOs.: 2 and 568;
- (252) SEQ ID NOs.: 2 and 101;
- (253) SEQ ID NOs.: 529 and 49;
- (254) SEQ ID NOs.: 529 and 103;
- (255) SEQ ID NOs.: 529 and 117; or
- (256) SEQ ID NOs.: 530 and 101.

52. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 49, respectively.

53. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 100, respectively.

54. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 42 and 101, respectively.

55. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 103, respectively.

56. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 117, respectively.

57. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 517 and 101, respectively.

58. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 518 and 101, respectively.

59. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 542, respectively.

60. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 517 and 542, respectively.

61. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 518 and 542, respectively.

62. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 544, respectively.

63. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 2 and 567, respectively.

64. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 529 and 103, respectively.

65. The antibody of claim 51, wherein the heavy chain and light have the amino acid sequence set forth in SEQ ID NOs.: 530 and 101, respectively.

66. The antibody or antigen-binding fragment thereof of any one of claims 1 to 65, comprising a heavy chain (HC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and a light chain (LC) that is at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 48-136 and 531-567.

67. The antibody or antigen-binding fragment thereof of any one of claims 1 to 66, wherein at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or more, N-linked glycosylation sites in the VL are sialylated.

68. An antibody or an antigen-binding fragment thereof that binds to human immunodeficiency virus-1 (HIV-1) Envelope glycoprotein gp120, the antibody or antigen-binding fragment thereof comprising (i) a heavy chain variable region (VH) comprising VH complementary determining regions 1-3 (CDRs 1-3) and (ii) a light chain variable region (VL) comprising VL CDRs 1-3, wherein the VH CDRs 1-3 and VL CDRs 1-3 have the sequences set forth in:

- (i) SEQ ID NOs.: 159, 138, 139, 140, 141, and 142, respectively;
- (ii) SEQ ID NOs.: 137, 160, 139, 140, 141, and 142, respectively;
- (iii) SEQ ID NOs.: 137, 161, 139, 140, 141, and 142, respectively;
- (iv) SEQ ID NOs.: 137, 162, 139, 140, 141, and 142, respectively;
- (v) SEQ ID NOs.: 137, 163, 139, 140, 141, and 142, respectively;
- (vi) SEQ ID NOs.: 137, 138, 164, 140, 141, and 142, respectively;
- (vii) SEQ ID NOs.: 159, 138, 164, 140, 141, and 142, respectively;
- (viii) SEQ ID NOs.: 137, 138, 139, 140, 165, and 142, respectively;
- (ix) SEQ ID NOs.: 137, 138, 139, 140, 166, and 142, respectively;
- (x) SEQ ID NOs.: 137, 138, 139, 140, 167, and 142, respectively;
- (xi) SEQ ID NOs.: 137, 138, 139, 140, 168, and 142, respectively;
- (xii) SEQ ID NOs.: 137, 138, 154, 140, 141, and 142, respectively, or
- (xiii) SEQ ID NOs.: 137, 138, 139, 140, 141, and 142, respectively, and wherein

at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, or more, N-linked glycosylation sites in the VL are sialylated.

69. The antibody or antigen-binding fragment of any one of claims 67 to 68, wherein the asparagine at VL amino acid position 72 according to Kabat numbering (N72) is sialylated.

70. The antibody or antigen-binding fragment of any one of claims 67 to 69, wherein the sialylated N-linked glycosylation sites in the VL comprise from 1 to 5 sialic acid residues, *e.g.*, from 1 to 4 sialic acid residues, *e.g.*, from 1 to 3 sialic acid residues, *e.g.*, from 1 to 2 sialic acid residues.

71. The antibody or antigen-binding fragment thereof of any one of claims 67 to 70, wherein the VL are sialylated with N-acetylneuraminic acid (NANA).

72. The antibody or antigen-binding fragment thereof of any one of claims 67 to 71, wherein the sialic acid residues are present in biantennary structures.

73. The antibody or antigen-binding fragment thereof of any one of claims 67 to 72, wherein the sialic acid residues are present in complex N-linked glycan structures.

74. The antibody or antigen-binding fragment thereof of any one of claims 67 to 72, wherein the sialic acid residues are present in hybrid N-linked glycan structures.

75. The antibody or antigen-binding fragment thereof of any one of claims 67 to 74, wherein the glycans are terminally sialylated.

76. A bispecific antibody comprising:
a first antigen binding arm that binds to gp120, the first antigen binding arm comprising:

- i. the VH CDRs 1-3 and the VL CDRs 1-3; or
 - ii. the VH and the VL of any one of claims 1 to 75; and
- a second antigen binding arm binding to a second antigen.

77. The bi-specific antibody of claim 76, wherein the second antigen is selected from the group consisting of CD3, FcγRI (CD64), FcγRII (CD32), FcγRIII (CD16); CD89, CCR5, CD4, gp41, killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, three Ig domains and long cytoplasmic tail 1 (KIR3DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 1 (KIR2DL1), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 2 (KIR2DL2), killer cell immunoglobulin like receptor, two Ig domains and long cytoplasmic tail 3 (KIR2DL3), killer cell lectin like receptor C1 (KLRC1), killer cell lectin like receptor C2 (KLRC2), killer cell lectin like receptor C3 (KLRC3), killer cell lectin like receptor C4 (KLRC4), killer cell lectin like receptor D1 (KLRD1), killer cell lectin like receptor K1 (KLRK1), natural cytotoxicity triggering receptor 3 (NCR3 or NKp30), natural cytotoxicity triggering receptor 2 (NCR2 or NK-p44), natural cytotoxicity triggering receptor 1 (NCR1 or NK-p46), CD226 (DNAM-1), cytotoxic and regulatory T cell molecule (CRTAM or CD355), signaling lymphocytic activation molecule family member 1 (SLAMF1), CD48 (SLAMF2), lymphocyte antigen 9 (LY9 or SLAMF3), CD244 (2B4 or SLAMF4), CD84 (SLAMF5), SLAM family member 6 (SLAMF6 or NTB-A), SLAM family member 7 (SLAMF7 or CRACC), CD27 (TNFRSF7), semaphorin 4D (SEMA4D or CD100), and CD160 (NK1), and a second epitope of gp120.

78. A pharmaceutical composition comprising the antibody or antigen-binding fragment of any one of claims 1-77, and a pharmaceutically acceptable carrier.

79. The pharmaceutical composition of claim 78, further comprising a second agent for treating an HIV infection.

80. The pharmaceutical composition of any one of claims 78 to 79, further comprising a toll-like receptor (TLR) agonist.

81. The pharmaceutical composition of any one of claims 78 to 80, wherein the TLR agonist is a TLR2 agonist, a TLR3 agonist, a TLR7 agonist, a TLR8 agonist or a TLR9 agonist

82. The pharmaceutical composition of any one of claims 78 to 80, further comprising a TLR7 agonist.

83. The pharmaceutical composition of claim 82, wherein the TLR7 agonist is selected from the group consisting of vesatolimod, imiquimod, and resiquimod.

84. The pharmaceutical composition of claim 78, further comprising an antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV.

85. The pharmaceutical composition of any one of claims 78 to 84, further comprising a second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, wherein the second antibody or antigen-binding fragment thereof does not compete with the antibody or antigen-binding fragment of any one of claims 1-70 for binding to gp120.

86. The pharmaceutical composition of claim 85, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, competes with or comprises VH and VL variable domains of a broadly neutralizing antibody (bNAb) against HIV.

87. The pharmaceutical composition of any one of claims 85 to 86, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 selected from the group consisting of:

- i. third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan;
- ii. second variable loop (V2) and/or Env trimer apex;
- iii. gp120/gp41 interface; or
- iv. silent face of gp120.

88. The pharmaceutical composition of any one of claims 85 to 87, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137, PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03.

89. The pharmaceutical composition of any one of claims 85 to 87, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the second variable loop (V2) and/or Env trimer apex and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PG9, PG16, PGC14, PGG14, PGT-142, PGT-143, PGT-144, PGT-145, CH01, CH59, PGDM1400, CAP256, CAP256-VRC26.08, CAP256-VRC26.09, CAP256-VRC26.25, PCT64-24E and VRC38.01.

90. The pharmaceutical composition of any one of claims 85 to 87, wherein the second antibody or antigen-binding fragment binds to an epitope or region of gp120 in the gp120/gp41 interface and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PGT-151, CAP248-2B, 35O22, 8ANC195, ACS202, VRC34 and VRC34.01.

91. The pharmaceutical composition of any one of claims 85 to 87, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp120 silent face and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC-PG05 and SF12.

92. The pharmaceutical composition of any one of claims 85 to 86, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER).

93. The pharmaceutical composition of claim 92, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER) and competes with or

comprises VH and VL regions from an antibody selected from the group consisting of 10E8, 10E8v4, 10E8-5R-100cF, 4E10, DH511.11P, 2F5, 7b2, and LN01.

94. The pharmaceutical composition of any one of claims 85 to 86, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp41 fusion peptide and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC34 and ACS202.

95. The pharmaceutical composition of claim 84, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of PGT121.60 or PGT121.66.

96. The pharmaceutical composition of claim 84, wherein the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO:454 and the VL within SEQ ID NO:455.

97. The pharmaceutical composition of claim 84, wherein the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO:454 and the VL within SEQ ID NO:456.

98. A nucleic acid or nucleic acids encoding the antibody or antigen-binding fragment of any one of claims 1-77.

99. The nucleic acid or nucleic acids of claim 98, wherein the nucleic acid or nucleic acids comprise DNA, cDNA or mRNA.

100. The nucleic acid or nucleic acids of any one of claims 98 to 99, encoding a VH selected from the group consisting of SEQ ID NOs: 181-221 and 465-478 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 572-581; and encoding a VL selected from the group consisting of SEQ ID NOs: 222-311, 479-516 and 569 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 582-595.

101. The nucleic acid or nucleic acids of any one of claims 98 to 99, encoding a HC selected from the group consisting of SEQ ID NOs: 1-47 and 517-530 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 596-605; and encoding a LC selected from the group consisting of SEQ ID NOs: 48-136 and 531-567 and having at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, identical to a nucleic acid sequence selected from the group consisting of SEQ ID NOs: 606-619.

102. An expression vector or expression vectors comprising the nucleic acid or nucleic acids of any one of claims 98 to 101 operably linked to a regulatory sequence.

103. The expression vector or expression vectors of claim 102, wherein the expression vector or expression vectors comprise a plasmid vector or a viral vector.

104. A pharmaceutical composition comprising the nucleic acid or nucleic acids of any one of claims 98 to 101, or the expression vector or expression vectors of any one of claims 102 to 103, and a pharmaceutically acceptable carrier.

105. A lipid nanoparticle (LNP) comprising the nucleic acid or nucleic acids of any one of claims 98 to 101, or the expression vector or expression vectors of any one of claims 102 to 103.

106. A chimeric antigen receptor (CAR) comprising an antigen-binding fragment according to claim 50.

107. A CAR T-cell comprising the CAR of claim 106.

108. A host cell or population of cells comprising the nucleic acid or nucleic acids of any one of claims 98 to 101, or the expression vector or expression vectors of any one of claims 102 to 103.

109. The host cell or population of cells of claim 108, wherein the cell or population of cells comprises a eukaryotic cell.

110. The cell or population of cells of any one of claims 108 to 109, wherein the cell or population of cells comprises a mammalian cell, an insect cell, a plant cell or a yeast cell.

111. The cell or population of cells of any one of claims 108 to 110, wherein the mammalian cell is a Chinese Hamster Ovary (CHO) cell.

112. The cell or population of cells of any one of claims 108 to 110, wherein the mammalian cell is a human cell.

113. The cell or population of cells of claim 112, wherein the cell is a human embryonic kidney cell or a human B-cell.

114. The cell or population of cells of any one of claims 108 to 113, wherein the cell predominantly sialylates N-linked glycosylation sites in the variable domains (Fv) of expressed antibodies or antigen binding fragments.

115. The cell or population of cells of claim 114, wherein at least 50%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the variable domains (Fv) of expressed antibodies or antigen binding fragments are sialylated.

116. The cell or population of cells of any one of claims 114 to 115, wherein at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, or more, N-linked glycosylation sites in the VL are sialylated.

117. The cell or population of cells of any one of claims 114 to 116, wherein the asparagine at VL amino acid position 72 according to Kabat numbering (N72) is sialylated.

118. The cell or population of cells of any one of claims 114 to 117, wherein the sialylated N-linked glycosylation sites in the VL comprise from 1 to 5 sialic acid residues, *e.g.*, from 1 to 4 sialic acid residues, *e.g.*, from 1 to 3 sialic acid residues, *e.g.*, from 1 to 2 sialic acid residues.

119. The cell or population of cells of any one of claims 114 to 118, wherein the VL are sialylated with N-acetylneuraminic acid (NANA).

120. The cell or population of cells of any one of claims 114 to 119, wherein the sialic acid residues are present in biantennary structures.

121. The cell or population of cells of any one of claims 114 to 120, wherein the sialic acid residues are present in complex N-linked glycan structures.

122. The cell or population of cells of any one of claims 114 to 121, wherein the sialic acid residues are present in hybrid N-linked glycan structures.

123. The cell or population of cells of any one of claims 114 to 122, wherein the glycans are terminally sialylated.

124. A method of producing an antibody or antigen-binding fragment thereof, the method comprising:

culturing the host cell of any one of claims 108 to 123 in a cell culture; and

isolating the antibody or antigen-binding fragment from the cell culture.

125. The method of claim 124, further comprising formulating the antibody or antigen-binding fragment into a sterile pharmaceutical composition suitable for administration to a human subject.

126. A method of treating or preventing HIV in a human subject in need thereof, the method comprising administering to the subject an effective amount of the antibody or antigen-binding fragment thereof of any one of claims 1-77, or the pharmaceutical composition of any one of claims 78-97.

127. The method of claim 126, further comprising administering to the subject a second agent for treating an HIV infection.

128. The method of any one of claims 126 to 127, further comprising administering to the subject a toll-like receptor (TLR) agonist.

129. The method of any one of claims 126 to 128, wherein the TLR agonist is a TLR2 agonist, a TLR3 agonist, a TLR7 agonist, a TLR8 agonist or a TLR9 agonist.

130. The method of any one of claims 127 to 129, further comprising administering to the subject a TLR7 agonist.

131. The method of claim 130, wherein the TLR7 agonist is selected from the group consisting of vesatolimod, imiquimod, and resiquimod.

132. The method of any one of claims 126 to 131, further comprising administering to the human subject an antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV.

133. The method of any one of claims 127 to 132, further comprising administering to the human subject an effective amount of a second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, wherein the second antibody or antigen-binding fragment thereof does not compete with the antibody or antigen-binding fragment of any one of claims 1-70 for binding to gp120.

134. The method of claim 133, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, competes with or comprises VH and VL variable domains of a broadly neutralizing antibody (bNAb) against HIV.

135. The method of any one of claims 133 to 134, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 selected from the group consisting of:

- i. third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan;
- ii. second variable loop (V2) and/or Env trimer apex;
- iii. gp120/gp41 interface; or
- iv. silent face of gp120.

136. The method of any one of claims 133 to 135, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the third variable loop (V3) and/or high mannose patch comprising a N332 oligomannose glycan and competes with or comprises VH and VL regions from an antibody selected from the group consisting of GS-9722, PGT-121, PGT-122, PGT-123, PGT-124, PGT-125, PGT-126, PGT-128, PGT-130, PGT-133, PGT-134, PGT-135, PGT-136, PGT-137, PGT-138, PGT-139, 10-1074, VRC24, 2G12, BG18, 354BG8, 354BG18, 354BG42, 354BG33, 354BG129, 354BG188, 354BG411, 354BG426, DH270.1, DH270.6, PGDM12, VRC41.01, PGDM21, PCDN-33A, BF520.1 and VRC29.03.

137. The method of any one of claims 133 to 135, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp120 in the second variable loop (V2) and/or Env trimer apex and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PG9, PG16, PGC14, PGG14, PGT-142, PGT-143, PGT-144, PGT-145, CH01, CH59, PGDM1400, CAP256, CAP256-VRC26.08, CAP256-VRC26.09, CAP256-VRC26.25, PCT64-24E and VRC38.01.

138. The method of any one of claims 133 to 135, wherein the second antibody antigen-binding fragment binds to an epitope or region of gp120 in the gp120/gp41 interface and competes with or comprises VH and VL regions from an antibody selected from the group consisting of PGT-151, CAP248-2B, 35O22, 8ANC195, ACS202, VRC34 and VRC34.01.

139. The method of any one of claims 133 to 135, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp120 silent face and competes with or comprises VH and VL regions from antibody selected from the group consisting of VRC-PG05 and SF12.

140. The method of any one of claims 133 to 135, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER).

141. The method of any one of claims 133 to 134, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of gp41 in the membrane proximal region (MPER) and competes with or comprises VH and VL regions from an antibody selected from the group consisting of 10E8, 10E8v4, 10E8-5R-100cF, 4E10, DH511.11P, 2F5, 7b2, and LN01.

142. The method of any one of claims 133 to 134, wherein the second antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV, binds to an epitope or region of the gp41 fusion peptide and competes with or comprises VH and VL regions from an antibody selected from the group consisting of VRC34 and ACS202.

143. The method of claim 132, wherein the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH and VL of PGT121.60 or PGT121.66.

144. The method of claim 132, wherein the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO:454 and the VL within SEQ ID NO:455.

145. The method of claim 132, wherein the antibody or antigen-binding fragment thereof that binds, inhibits, and/or neutralizes HIV comprises the VH within SEQ ID NO:454 and the VL within SEQ ID NO:456.

146. The method of any one of claims 126 to 145, further comprising administering to the human subject an anti-HIV vaccine.

147. The method of claim 146, wherein the anti-HIV vaccine comprises a viral vaccine.

148. The method of any one of claims 146 to 147, wherein the viral vaccine is from a virus selected from the group consisting of an arenavirus, an adenovirus, a poxvirus, and a rhabdovirus.

149. A method of inhibiting HIV in a human subject in need thereof, the method comprising administering to the subject an effective amount of the antibody or antigen-binding fragment thereof of any one of claims 1-77, or the pharmaceutical composition of any one of claims 78-97.




FIG. 1

2/8




FIG. 2




FIG. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2019/040342

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61P31/18 C07K16/10
ADD. A61K39/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61P A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, Sequence Search, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2013/192589 A1 (CALIFORNIA INST OF TECHN [US]) 27 December 2013 (2013-12-27) sequences 9,10	1-149
X	WO 2016/196740 A1 (UNIV ROCKEFELLER [US]) 8 December 2016 (2016-12-08) sequences 39-47 ----- -/--	1-149



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

25 November 2019

Date of mailing of the international search report

10/12/2019

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Scheffzyk, Irmgard

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2019/040342

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	J. F. SCHEID ET AL: "Sequence and Structural Convergence of Broad and Potent HIV Antibodies That Mimic CD4 Binding: supporting online material", SCIENCE, NIH AUTHOR MANUSCRIPT, vol. 333, no. 6049, 14 July 2011 (2011-07-14), pages 1-54, XP055400143, US ISSN: 0036-8075, DOI: 10.1126/science.1207227 cited in the application the whole document -----	1-149

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2019/040342

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013192589 A1	27-12-2013	US 2014017234 A1	16-01-2014
		US 2018230203 A1	16-08-2018
		WO 2013192589 A1	27-12-2013

WO 2016196740 A1	08-12-2016	US 2018179299 A1	28-06-2018
		WO 2016196740 A1	08-12-2016
