




(86) **Date de dépôt PCT/PCT Filing Date:** 2011/01/06
(87) **Date publication PCT/PCT Publication Date:** 2011/07/14
(45) **Date de délivrance/Issue Date:** 2016/01/05
(85) **Entrée phase nationale/National Entry:** 2012/06/20
(86) **N° demande PCT/PCT Application No.:** US 2011/020416
(87) **N° publication PCT/PCT Publication No.:** 2011/085128
(30) **Priorité/Priority:** 2010/01/07 (US61/293,128)

(51) **Cl.Int./Int.Cl.** **C07D 277/28** (2006.01),
A61K 31/421 (2006.01), **A61K 31/426** (2006.01),
A61K 31/496 (2006.01), **A61K 31/5377** (2006.01),
A61P 35/00 (2006.01), **C07D 233/64** (2006.01),
C07D 263/32 (2006.01)

(72) **Inventeurs/Inventors:**
VERNIER, JEAN-MICHEL, US;
MAY, JOHN, US;
O'CONNOR, PATRICK, US;
RIPKA, WILLIAM, US;
PINKERTON, ANTHONY, US;
BOUNAUD, PIERRE-YVES, US;
HOPKINS, STEPHANIE, US

(73) **Propriétaire/Owner:**

(54) **Titre : INHIBITEURS DE LA VOIE HEDGEHOG**
(54) **Title: HEDGEHOG INHIBITORS**



(57) **Abrégé/Abstract:**

Described herein are compounds, pharmaceutical compositions and methods for the inhibition of Hedgehog signaling. Said compounds, pharmaceutical compositions and methods have utility in the treatment of human and veterinary disease and disorders.



(73) **Propriétaires(suite)/Owners(continued):**SELEXAGEN THERAPEUTICS, INC., US

(74) **Agent:** GOWLING LAFLEUR HENDERSON LLP

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 July 2011 (14.07.2011)

PCT

(10) International Publication Number
WO 2011/085128 A1

(51) International Patent Classification:

A01N 43/78 (2006.01) *A61K 31/425* (2006.01)

(21) International Application Number:

PCT/US2011/020416

(22) International Filing Date:

6 January 2011 (06.01.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/293,128 7 January 2010 (07.01.2010) US

(71) Applicant (for all designated States except US): **SELEX-AGEN THERAPEUTICS, INC.** [US/US]; 4653 Carmel Mountain Road, Suite 308, San Diego, CA 92130 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **VERNIER, Jean-Michel** [FR/US]; 5150 Sea Mist Court, San Diego, CA 92121 (US). **MAY, John** [US/US]; 14169 Caminito Vis-tana, San Diego, CA 92130 (US). **O'CONNOR, Patrick** [US/US]; 3586 Torrey View Court, San Diego, CA 92130 (US). **RIPKA, William** [US/US]; 10819 Red Rock Road, San Diego, CA 92131 (US). **PINKERTON, Anthony** [US/US]; 7613 Mona Lane, San Diego, CA 92130 (US). **BOUNAUD, Pierre-Yves** [FR/US]; 4610 Janet Place, San Diego, CA 92115 (US). **HOPKINS, Stephanie** [US/US]; 12716 Tustin Street, Poway, CA 92064 (US).(74) Agents: **CLARK, David** et al.; Wilson Sonsini Goodrich & Rosati, 650 Page Mill Road, Palo Alto, CA 94304-1050 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

Published:

- with international search report (Art. 21(3))

(54) Title: HEDGEHOG INHIBITORS

Alkaline Phosphatase Activity
(units/ mg protein)

(57) Abstract: Described herein are compounds, pharmaceutical compositions and methods for the inhibition of Hedgehog signaling. Said compounds, pharmaceutical compositions and methods have utility in the treatment of human and veterinary disease and disorders.

WO 2011/085128 A1


HEDGEHOG INHIBITORS

BACKGROUND OF THE INVENTION

[0002] Described herein are compounds, pharmaceutical compositions and methods for the inhibition of Hedgehog signaling. Said compounds, pharmaceutical compositions and methods have utility in the treatment of human and veterinary disease and disorders.

SUMMARY OF THE INVENTION

[0003] One embodiment provides a compound having the structure of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-

alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -
CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl,
aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-
linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -
CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -
NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

10 **[0004]** Another embodiment provides a compound having the structure of Formula (I)
wherein X is -S-, -O-.

[0005] Another embodiment provides a compound having the structure of Formula (I)
wherein G¹ and G² can not both be hydrogen.

15 **[0006]** Another embodiment provides a compound having the structure of Formula (I)
wherein:

X is -S-, -O-, -N(H)- or -N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

20 G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃, and wherein G¹ and G² can
not both be hydrogen;

R¹ is H or C₁-C₃ alkyl;

25 R² is selected from halogen, -CN, alkyl, aryl, -O-aryl, -O-heteroaryl, -CH₂-aryl,
-CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, -NH-heteroaryl, -NH-alkyl, -CH₂-NH-
alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked
heterocycle), N-linked heterocycle, and C-linked heterocycle;

30 R⁴ is selected from alkoxy, -CN, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-
aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -
CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H,
and -CO₂alkyl; and

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl,
aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-
linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -

CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -
 NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
 N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂.

[0007] Another embodiment provides a compound having the structure of Formula (I)

5 wherein n is 0.

[0008] Another embodiment provides a compound having the structure of Formula (I)

wherein n is 1.

[0009] Another embodiment provides a compound having the structure of Formula (I)

wherein G² is H.

10 [0010] Another embodiment provides a compound having the structure of Formula (I)

wherein G² is H and G¹ is alkyl.

[0011] Another embodiment provides a compound having the structure of Formula (I)

wherein X is -S-.

[0012] Another embodiment provides a compound having the structure of Formula (I)

15 wherein R² is -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), or -CH₂-(C-linked heterocycle).

[0013] Another embodiment provides a compound having the structure of Formula (I)

wherein R² is halogen, -CN, -alkyl, or -CF₃.

[0014] Another embodiment provides a compound having the structure of Formula (I)

20 wherein R⁴ is -SO₂Me or -OMe.

[0015] Another embodiment provides a compound having the structure of Formula (I)

wherein R² is halogen and R⁴ is -SO₂Me.

[0016] Another embodiment provides a compound having the structure of Formula (I)

wherein R² is halogen and R⁴ is -OMe.

25 [0017] Another embodiment provides a compound having the structure of Formula (I)


wherein n is 0; G² is H; G¹ is alkyl; X is -S-; R² is halogen and R⁴ is -OMe.

[0018] Another embodiment provides a compound having the structure of Formula (I)

wherein n is 0; G² is H; G¹ is alkyl; X is -S-; R² is halogen and R⁴ is -SO₂Me.

[0019] One embodiment provides a pharmaceutical composition comprising a compound of

30 Formula (I), or a stereoisomer, hydrate, solvate or pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable excipient, wherein the compound of Formula (I) has the following structure:



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;


R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[0020] One embodiment provides a method of inhibiting the Hedgehog pathway in a cell comprising contacting the cell with an inhibitory concentration of a compound of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;


R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[0021] Another embodiment provides the method wherein the cell is characterized by a patched loss-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a smoothened gain-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a constitutively active smoothened

phenotype. Another embodiment provides the method wherein the cell is characterized by expression of *Gli*.

[0022] One embodiment provides a method of inhibiting the activity of smoothened protein in a cell comprising contacting the smoothened protein with an inhibitory concentration of a compound of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;


R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -

NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[0023] Another embodiment provides the method wherein the cell is characterized by a
5 patched loss-of-function phenotype. Another embodiment provides the method wherein the
cell is characterized by a smoothened gain-of-function phenotype. Another embodiment
provides the method wherein the cell is characterized by a constitutively active smoothened
phenotype. Another embodiment provides the method wherein the cell is characterized by
expression of *Gli*.

10 [0024] One embodiment provides a method of inhibiting the transcriptional activity of *Gli*
transcription factor in a cell comprising contacting the cell with an inhibitory concentration
of a compound of Formula (I):



15 or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt
thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

20 G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -
O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl -
25 NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked
heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked
heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -
SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -


NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -
CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl,
aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-
linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -
CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -
NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

10 [0025] Another embodiment provides the method wherein the cell is characterized by a
patched loss-of-function phenotype. Another embodiment provides the method wherein the
cell is characterized by a smoothened gain-of-function phenotype. Another embodiment
provides the method wherein the cell is characterized by a constitutively active smoothened
phenotype. Another embodiment provides the method wherein the cell is characterized by
15 expression of *Gli*.

[0026] One embodiment provides a method of inhibiting *Gli*-mediated gene transcription in
a cell comprising contacting the cell with an inhibitory concentration of a compound of
Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt
thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R^2 is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;


R^4 is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R^3 , R^5 and R^6 are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R^7 is H or C₁-C₃ alkyl.

[0027] Another embodiment provides the method wherein the cell is characterized by a patched loss-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a smoothened gain-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a constitutively active smoothened phenotype. Another embodiment provides the method wherein the cell is characterized by expression of *Gli*.

[0028] One embodiment provides a method of treating a human disease or disorder mediated by Hedgehog pathway comprising administering to a patient a therapeutically effective amount of a composition comprising a compound of Formula (I), or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein the compound of Formula (I) has the following structure:



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

5 n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

10 R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl, -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

15 R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;


20 R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

25 **[0029]** Another embodiment provides the method wherein the disease or disorder is a proliferative disease. Another embodiment provides the method wherein the proliferative disease is selected from colon cancer, lung cancer, pancreatic cancer, gastric cancer, prostate cancer, and hepatocellular carcinoma. Another embodiment provides the method wherein the proliferative disease is selected from basal cell carcinoma, breast cancer, bone
30 sarcoma, soft tissue sarcoma, chronic myeloid leukemia, acute myeloid leukemia, hematological cancer, medulloblastoma, rhabdomyosarcoma, neuroblastoma, pancreatic cancer, breast carcinoma, meningioma, glioblastoma, astrocytoma, melanoma, stomach cancer, esophageal cancer, biliary tract cancer, prostate cancer, small cell lung cancer, non-

small cell lung cancer, glial cell cancer, multiple myeloma, colon cancer, neuroectodermal tumor, neuroendocrine tumor, mastocytoma and Gorlin syndrome. Another embodiment provides the method wherein the proliferative disease is basal cell carcinoma.

[0030] One embodiment provides a method of treating a veterinary disease or disorder mediated by Hedgehog pathway comprising administering to a subject a therapeutically effective amount of a composition comprising a compound of Formula (I), or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein the compound of Formula (I) has the following structure:



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-

linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -
 CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -
 NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
 N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

5 R^7 is H or C₁-C₃ alkyl.

[0031] Another embodiment provides a method of treating a veterinary disease or disorder wherein the disease or disorder is a proliferative disease selected from mast cell tumors or osteosarcoma.

10

BRIEF DESCRIPTION OF THE DRAWINGS

[0033] The novel features of the invention are set forth with particularity in the appended
 15 claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:


[0034] Figure 1 shows the dose-reponse of cyclopamine, a positive control, in the alkaline
 20 phosphatase assay described herein.

[0035] Figure 2 shows the dose-reponse of the compound of Example 4 in the alkaline phosphatase assay described herein.

DETAILED DESCRIPTION OF THE INVENTION

Heterocyclic Benzamide Hedgehog Inhibitors

25 [0036] One embodiment provides a compound having the structure of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl, -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[0037] Another embodiment provides a compound having the structure of Formula (I) wherein X is -S-, -O-.

[0038] Another embodiment provides a compound having the structure of Formula (I) wherein G¹ and G² can not both be hydrogen.

[0039] Another embodiment provides a compound having the structure of Formula (I) wherein:

X is -S-, -O-, -N(H)- or -N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃, and wherein G¹ and G² can not both be hydrogen;

R^1 is H or C_1 - C_3 alkyl;

R^2 is selected from halogen, -CN, alkyl, aryl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R^4 is selected from alkoxy, -CN, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl; and

R^3 , R^5 and R^6 are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂.

[0040] Another embodiment provides a compound having the structure of Formula (I) wherein n is 0.

[0041] Another embodiment provides a compound having the structure of Formula (I) wherein n is 1.

[0042] Another embodiment provides a compound having the structure of Formula (I) wherein G^2 is H.

[0043] Another embodiment provides a compound having the structure of Formula (I) wherein G^2 is H and G^1 is alkyl.

[0044] Another embodiment provides a compound having the structure of Formula (I) wherein X is -S-.

[0045] Another embodiment provides a compound having the structure of Formula (I) wherein R^2 is -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), or -CH₂-(C-linked heterocycle).

[0046] Another embodiment provides a compound having the structure of Formula (I) wherein R^2 is halogen, -CN, -alkyl, or -CF₃.

[0047] Another embodiment provides a compound having the structure of Formula (I) wherein R^4 is -SO₂Me or -OMe.

[0048] Another embodiment provides a compound having the structure of Formula (I) wherein R^2 is halogen and R^4 is $-\text{SO}_2\text{Me}$.





[0049] Another embodiment provides a compound having the structure of Formula (I) wherein R^2 is halogen and R^4 is $-\text{OMe}$.







5 [0050] Another embodiment provides a compound having the structure of Formula (I) wherein n is 0; G^2 is H; G^1 is alkyl; X is $-\text{S}-$; R^2 is halogen and R^4 is $-\text{OMe}$.







[0051] Another embodiment provides a compound having the structure of Formula (I) wherein n is 0; G^2 is H; G^1 is alkyl; X is $-\text{S}-$; R^2 is halogen and R^4 is $-\text{SO}_2\text{Me}$.



10 [0052] In certain specific embodiments, the compounds of Formula (I) have the structures shown in Table 1.

Table 1


Example	Structure
1	 2,4-dichloro-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide
2	 3,4,5-trimethoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide
3	 4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide
4	 2-chloro-4-(methylsulfonyl)-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide

5	 4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)-2-(morpholinomethyl)benzamide
6	 2-chloro-<i>N</i>-(4-(2-methyloxazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide
7	 2-chloro-<i>N</i>-(3-chloro-4-(2-methylthiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide
8	 2-chloro-4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide
9	 2-chloro-<i>N</i>-methyl-4-(methylsulfonyl)-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide
10	 2-chloro-<i>N</i>-(4-(thiazol-4-yl)phenyl)benzamide


11	 2-chloro-4-(methylsulfonyl)-N-(4-(thiazol-4-yl)phenyl)benzamide
12	 2-chloro-N-(4-(2-(2-hydroxypropan-2-yl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide
13	 2-chloro-N-(4-(2-(hydroxymethyl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide
14	 2-chloro-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide
15	 2-(methylsulfonyl)-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide
16	 4-methoxy-2-methyl-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide

17	 2-fluoro-4-methoxy-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide
18	 4-cyano-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide


[0053] In further embodiments, a compound of Formula (I) is selected from the structures shown below as examples 19-193.




Example 19




Example 20




example 21




example 22




example 23




example 24




example 25




example 26




example 27




example 28




example 29




example 30




example 31




example 32




example 33




example 34




example 35




example 36




example 37






















example 38






















example 39


















































Example 184




Example 185




Example 186




Example 187




Example 188




Example 189




Example 190



Example 191



Example 192



Example 193

Definitions

[0054] As used herein and in the appended claims, the singular forms "a", "and", and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a compound" includes a plurality of such compounds, and reference to "the

cell" includes reference to one or more cells (or to a plurality of cells) and equivalents thereof known to those skilled in the art, and so forth. When ranges are used herein for physical properties, such as molecular weight, or chemical properties, such as chemical formulae, all combinations and subcombinations of ranges and specific embodiments
5 therein are intended to be included. The term "about" when referring to a number or a numerical range means that the number or numerical range referred to is an approximation within experimental variability (or within statistical experimental error), and thus the number or numerical range may vary between 1% and 15% of the stated number or numerical range. The term "comprising" (and related terms such as "comprise" or
10 "comprises" or "having" or "including") is not intended to exclude that in other certain embodiments, for example, an embodiment of any composition of matter, composition, method, or process, or the like, described herein, may "consist of" or "consist essentially of" the described features.

[0055] "Amino" refers to the -NH_2 radical.

15 [0056] "Cyano" refers to the -CN radical.

[0057] "Nitro" refers to the -NO_2 radical.

[0058] "Oxa" refers to the -O- radical.

[0059] "Oxo" refers to the =O radical.

[0060] "Imino" refers to the =NH radical.

20 [0061] "Thioxo" refers to the =S radical.

[0062] "Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to fifteen carbon atoms (*e.g.*, $\text{C}_1\text{-C}_{15}$ alkyl). In certain embodiments, an alkyl comprises one to thirteen carbon atoms (*e.g.*, $\text{C}_1\text{-C}_{13}$ alkyl). In certain embodiments, an alkyl comprises one
25 to eight carbon atoms (*e.g.*, $\text{C}_1\text{-C}_8$ alkyl). In other embodiments, an alkyl comprises one to five carbon atoms (*e.g.*, $\text{C}_1\text{-C}_5$ alkyl). In other embodiments, an alkyl comprises one to four carbon atoms (*e.g.*, $\text{C}_1\text{-C}_4$ alkyl). In other embodiments, an alkyl comprises one to three carbon atoms (*e.g.*, $\text{C}_1\text{-C}_3$ alkyl). In other embodiments, an alkyl comprises one to two carbon atoms (*e.g.*, $\text{C}_1\text{-C}_2$ alkyl). The alkyl is attached to the rest of the molecule by a
30 single bond, for example, methyl (Me), ethyl (Et), *n*-propyl, 1-methylethyl (*iso*-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, and the like. Unless stated otherwise specifically in the specification, an alkyl group is optionally substituted by one or more of the following substituents: halo, cyano, nitro, oxo, thioxo,

trimethylsilanyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)N(R^a)_2$,
 $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1
 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl,
 fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl,
 5 heteroaryl or heteroarylalkyl. Unless stated otherwise in the specification, an alkyl group is
 optionally a fluorinated or perfluorinated alkyl group, such as CF_3 , CF_2CF_3 , CH_2F , CHF_2 ,
 CH_2CF_3 and the like.

[0063] "Alkenyl" refers to a straight or branched hydrocarbon chain radical group
 consisting solely of carbon and hydrogen atoms, containing at least one double bond, and
 10 having from two to twelve carbon atoms. In certain embodiments, an alkenyl comprises
 two to eight carbon atoms. In other embodiments, an alkenyl comprises two to four carbon
 atoms. The alkenyl is attached to the rest of the molecule by a single bond, for example,
 ethenyl (*i.e.*, vinyl), prop-1-enyl (*i.e.*, allyl), but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and
 the like. Unless stated otherwise specifically in the specification, an alkenyl group is
 15 optionally substituted by one or more of the following substituents: halo, cyano, nitro, oxo,
 thioxo, trimethylsilanyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$, $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)N(R^a)_2$,
 $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1
 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl,
 fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl,
 20 heteroaryl or heteroarylalkyl.

[0064] "Alkynyl" refers to a straight or branched hydrocarbon chain radical group
 consisting solely of carbon and hydrogen atoms, containing at least one triple bond, having
 from two to twelve carbon atoms. In certain embodiments, an alkynyl comprises two to
 eight carbon atoms. In other embodiments, an alkynyl has two to four carbon atoms. The
 25 alkynyl is attached to the rest of the molecule by a single bond, for example, ethynyl,
 propynyl, butynyl, pentynyl, hexynyl, and the like. Unless stated otherwise specifically in
 the specification, an alkynyl group is optionally substituted by one or more of the following
 substituents: halo, cyano, nitro, oxo, thioxo, trimethylsilanyl, $-OR^a$, $-SR^a$, $-OC(O)-R^a$,
 $-N(R^a)_2$, $-C(O)R^a$, $-C(O)OR^a$, $-C(O)N(R^a)_2$, $-N(R^a)C(O)OR^a$, $-N(R^a)C(O)R^a$, $-N(R^a)S(O)_tR^a$
 30 (where t is 1 or 2), $-S(O)_tOR^a$ (where t is 1 or 2) and $-S(O)_tN(R^a)_2$ (where t is 1 or 2) where
 each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl,
 aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

[0065] "Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, for example, methylene, ethylene, propylene, *n*-butylene, and the like. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond. The points of attachment of the alkylene chain to the rest of the molecule and to the radical group can be through one carbon in the alkylene chain or through any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkylene chain is optionally substituted by one or more of the following substituents: halo, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, thioxo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2) and -S(O)_tN(R^a)₂ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, carbocyclyl, carbocyclylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl.

[0066] "Alkenylene" or "alkenylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group, consisting solely of carbon and hydrogen, containing at least one double bond and having from two to twelve carbon atoms, for example, ethenylene, propenylene, *n*-butenylene, and the like. The alkenylene chain is attached to the rest of the molecule through a double bond or a single bond and to the radical group through a double bond or a single bond. The points of attachment of the alkenylene chain to the rest of the molecule and to the radical group can be through one carbon or any two carbons within the chain. Unless stated otherwise specifically in the specification, an alkenylene chain is optionally substituted by one or more of the following substituents: halo, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, thioxo, trimethylsilyl, -OR^a, -SR^a, -OC(O)-R^a, -N(R^a)₂, -C(O)R^a, -C(O)OR^a, -C(O)N(R^a)₂, -N(R^a)C(O)OR^a, -N(R^a)C(O)R^a, -N(R^a)S(O)_tR^a (where t is 1 or 2), -S(O)_tOR^a (where t is 1 or 2) and -S(O)_tN(R^a)₂ (where t is 1 or 2) where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0067] "Aryl" refers to a radical derived from an aromatic monocyclic or multicyclic hydrocarbon ring system by removing a hydrogen atom from a ring carbon atom. The aromatic monocyclic or multicyclic hydrocarbon ring system contains only hydrogen and carbon from six to eighteen carbon atoms, where at least one of the rings in the ring system is fully unsaturated, *i.e.*, it contains a cyclic, delocalized $(4n+2)$ π -electron system in accordance with the Hückel theory. Aryl groups include, but are not limited to, groups such as phenyl, fluorenyl, and naphthyl. Unless stated otherwise specifically in the specification, the term "aryl" or the prefix "ar-" (such as in "aralkyl") is meant to include aryl radicals optionally substituted by one or more substituents independently selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-OC(O)-R^a$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl (optionally substituted with one or more halo groups), aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0068] "Aralkyl" refers to a radical of the formula $-R^c$ -aryl where R^c is an alkylene chain as defined above, for example, benzyl, diphenylmethyl and the like. The alkylene chain part of the aralkyl radical is optionally substituted as described above for an alkylene chain. The aryl part of the aralkyl radical is optionally substituted as described above for an aryl group.

[0069] "Aralkenyl" refers to a radical of the formula $-R^d$ -aryl where R^d is an alkenylene chain as defined above. The aryl part of the aralkenyl radical is optionally substituted as described above for an aryl group. The alkenylene chain part of the aralkenyl radical is optionally substituted as defined above for an alkenylene group.

[0070] "Aralkynyl" refers to a radical of the formula $-R^e$ -aryl, where R^e is an alkynylene chain as defined above. The aryl part of the aralkynyl radical is optionally substituted as

described above for an aryl group. The alkynylene chain part of the aralkynyl radical is optionally substituted as defined above for an alkynylene chain.

[0071] "Carbocyclyl" refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which includes fused or bridged ring systems, having from three to fifteen carbon atoms. In certain embodiments, a carbocyclyl comprises three to ten carbon atoms. In other embodiments, a carbocyclyl comprises five to seven carbon atoms. The carbocyclyl is attached to the rest of the molecule by a single bond. Carbocyclyl is optionally saturated, (*i.e.*, containing single C-C bonds only) or unsaturated (*i.e.*, containing one or more double bonds or triple bonds.) A fully saturated carbocyclyl radical is also referred to as "cycloalkyl." Examples of monocyclic cycloalkyls include, *e.g.*, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. An unsaturated carbocyclyl is also referred to as "cycloalkenyl." Examples of monocyclic cycloalkenyls include, *e.g.*, cyclopentenyl, cyclohexenyl, cycloheptenyl, and cyclooctenyl. Polycyclic carbocyclyl radicals include, for example, adamantyl, norbornyl (*i.e.*, bicyclo[2.2.1]heptanyl), norbornenyl, decalanyl, 7,7-dimethyl-bicyclo[2.2.1]heptanyl, and the like. Unless otherwise stated specifically in the specification, the term "carbocyclyl" is meant to include carbocyclyl radicals that are optionally substituted by one or more substituents independently selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, oxo, thioxo, cyano, nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-SR^a$, $-R^b-OC(O)-R^a$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where *t* is 1 or 2), $-R^b-S(O)_tOR^a$ (where *t* is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where *t* is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0072] "Carbocyclalkyl" refers to a radical of the formula $-R^c$ -carbocyclyl where R^c is an alkylene chain as defined above. The alkylene chain and the carbocyclyl radical is optionally substituted as defined above.

[0073] "Halo" or "halogen" refers to bromo, chloro, fluoro or iodo substituents.

[0074] "Fluoroalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more fluoro radicals, as defined above, for example, trifluoromethyl, difluoromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, and the like. The alkyl part of the fluoroalkyl radical is optionally substituted as defined above for an alkyl group.

[0075] "Heterocyclyl" refers to a stable 3- to 18-membered non-aromatic ring radical that comprises two to twelve carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, and includes fused or bridged ring systems. The heteroatoms in the heterocyclyl radical is optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized.

The heterocyclyl radical is partially or fully saturated. The heterocyclyl is attached to the rest of the molecule through any atom of the ring(s). Examples of such heterocyclyl radicals include, but are not limited to, dioxolanyl, thienyl[1,3]dithianyl,

decahydroisoquinolyl, imidazolyl, imidazolidinyl, isothiazolidinyl, isoxazolidinyl, morpholyl, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidinyl, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidinyl, quinuclidinyl, thiazolidinyl, tetrahydrofuryl, trithianyl, tetrahydropyranyl, thiomorpholyl, thiamorpholyl, 1-oxo-thiomorpholyl, and 1,1-dioxo-thiomorpholyl.

Unless stated otherwise specifically in the specification, the term "heterocyclyl" is meant to include heterocyclyl radicals as defined above that are optionally substituted by one or more substituents selected from alkyl, alkenyl, alkynyl, halo, fluoroalkyl, oxo, thioxo, cyano,

nitro, optionally substituted aryl, optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclalkyl, optionally substituted heterocyclyl, optionally substituted heterocyclalkyl, optionally substituted heteroaryl, optionally substituted heteroarylalkyl, $-R^b-OR^a$, $-R^b-SR^a$, $-R^b-OC(O)-R^a$, $-R^b-N(R^a)_2$, $-R^b-C(O)R^a$, $-R^b-C(O)OR^a$, $-R^b-C(O)N(R^a)_2$, $-R^b-O-R^c-C(O)N(R^a)_2$, $-R^b-N(R^a)C(O)OR^a$, $-R^b-N(R^a)C(O)R^a$, $-R^b-N(R^a)S(O)_tR^a$ (where t is 1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl,

heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

5 [0076] "*N*-heterocyclyl" or "N-attached heterocyclyl" refers to a heterocyclyl radical as defined above containing at least one nitrogen and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a nitrogen atom in the heterocyclyl radical. An *N*-heterocyclyl radical is optionally substituted as described above for heterocyclyl radicals. Examples of such *N*-heterocyclyl radicals include, but are not
10 limited to, 1-morpholinyl, 1-piperidinyl, 1-piperazinyl, 1-pyrrolidinyl, pyrazolidinyl, imidazolinyl, and imidazolidinyl.

[0077] "*C*-heterocyclyl" or "C-attached heterocyclyl" refers to a heterocyclyl radical as defined above containing at least one heteroatom and where the point of attachment of the heterocyclyl radical to the rest of the molecule is through a carbon atom in the heterocyclyl
15 radical. A *C*-heterocyclyl radical is optionally substituted as described above for heterocyclyl radicals. Examples of such *C*-heterocyclyl radicals include, but are not limited to, 2-morpholinyl, 2- or 3- or 4-piperidinyl, 2-piperazinyl, 2- or 3-pyrrolidinyl, and the like.

[0078] "Heterocyclylalkyl" refers to a radical of the formula $-R^c$ -heterocyclyl where R^c is an alkylene chain as defined above. If the heterocyclyl is a nitrogen-containing
20 heterocyclyl, the heterocyclyl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heterocyclylalkyl radical is optionally substituted as defined above for an alkylene chain. The heterocyclyl part of the heterocyclylalkyl radical is optionally substituted as defined above for a heterocyclyl group.

[0079] "Heteroaryl" refers to a radical derived from a 3- to 18-membered aromatic ring
25 radical that comprises two to seventeen carbon atoms and from one to six heteroatoms selected from nitrogen, oxygen and sulfur. As used herein, the heteroaryl radical is a monocyclic, bicyclic, tricyclic or tetracyclic ring system, wherein at least one of the rings in the ring system is fully unsaturated, *i.e.*, it contains a cyclic, delocalized $(4n+2)$ π -electron system in accordance with the Hückel theory. Heteroaryl includes fused or bridged ring
30 systems. The heteroatom(s) in the heteroaryl radical is optionally oxidized. One or more nitrogen atoms, if present, are optionally quaternized. The heteroaryl is attached to the rest of the molecule through any atom of the ring(s). Examples of heteroaryls include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzindolyl, 1,3-benzodioxolyl,

benzofuranyl, benzooxazolyl, benzo[d]thiazolyl, benzothiadiazolyl,
 benzo[b][1,4]dioxepinyl, benzo[b][1,4]oxazinyl, 1,4-benzodioxanyl, benzonaphthofuranyl,
 benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl,
 benzofuranonyl, benzothienyl (benzothiophenyl), benzothieno[3,2-d]pyrimidinyl,
 5 benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, carbazolyl, cinnolinyl,
 cyclopenta[d]pyrimidinyl, 6,7-dihydro-5H-cyclopenta[4,5]thieno[2,3-d]pyrimidinyl,
 5,6-dihydrobenzo[h]quinazolinyl, 5,6-dihydrobenzo[h]cinnolinyl, 6,7-dihydro-5H-
 benzo[6,7]cyclohepta[1,2-c]pyridazinyl, dibenzofuranyl, dibenzothiophenyl, furanyl,
 furanonyl, furo[3,2-c]pyridinyl, 5,6,7,8,9,10-hexahydrocycloocta[d]pyrimidinyl,
 10 5,6,7,8,9,10-hexahydrocycloocta[d]pyridazinyl,
 5,6,7,8,9,10-hexahydrocycloocta[d]pyridinyl, isothiazolyl, imidazolyl, indazolyl, indolyl,
 indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indolizinyl, isoxazolyl,
 5,8-methano-5,6,7,8-tetrahydroquinazolinyl, naphthyridinyl, 1,6-naphthyridinonyl,
 oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl,
 15 5,6,6a,7,8,9,10,10a-octahydrobenzo[h]quinazolinyl, 1-phenyl-1H-pyrrolyl, phenazinyl,
 phenothiazinyl, phenoxazinyl, phthalazinyl, pteridinyl, purinyl, pyrrolyl, pyrazolyl,
 pyrazolo[3,4-d]pyrimidinyl, pyridinyl, pyrido[3,2-d]pyrimidinyl, pyrido[3,4-d]pyrimidinyl,
 pyrazinyl, pyrimidinyl, pyridazinyl, pyrrolyl, quinazolinyl, quinoxalinyl, quinolinyl,
 isoquinolinyl, tetrahydroquinolinyl, 5,6,7,8-tetrahydroquinazolinyl,
 20 5,6,7,8-tetrahydrobenzo[4,5]thieno[2,3-d]pyrimidinyl,
 6,7,8,9-tetrahydro-5H-cyclohepta[4,5]thieno[2,3-d]pyrimidinyl,
 5,6,7,8-tetrahydropyrido[4,5-c]pyridazinyl, thiazolyl, thiadiazolyl, triazolyl, tetrazolyl,
 triazinyl, thieno[2,3-d]pyrimidinyl, thieno[3,2-d]pyrimidinyl, thieno[2,3-c]pyridinyl, and
 thiophenyl (*i.e.* thienyl). Unless stated otherwise specifically in the specification, the term
 25 "heteroaryl" is meant to include heteroaryl radicals as defined above which are optionally
 substituted by one or more substituents selected from alkyl, alkenyl, alkynyl, halo,
 fluoroalkyl, haloalkenyl, haloalkynyl, oxo, thioxo, cyano, nitro, optionally substituted aryl,
 optionally substituted aralkyl, optionally substituted aralkenyl, optionally substituted
 aralkynyl, optionally substituted carbocyclyl, optionally substituted carbocyclylalkyl,
 30 optionally substituted heterocyclyl, optionally substituted heterocyclylalkyl, optionally
 substituted heteroaryl, optionally substituted heteroarylalkyl, -R^b-OR^a, -R^b-SR^a,
 -R^b-OC(O)-R^a, -R^b-N(R^a)₂, -R^b-C(O)R^a, -R^b-C(O)OR^a, -R^b-C(O)N(R^a)₂,
 -R^b-O-R^c-C(O)N(R^a)₂, -R^b-N(R^a)C(O)OR^a, -R^b-N(R^a)C(O)R^a, -R^b-N(R^a)S(O)_tR^a (where t is

1 or 2), $-R^b-S(O)_tOR^a$ (where t is 1 or 2) and $-R^b-S(O)_tN(R^a)_2$ (where t is 1 or 2), where each R^a is independently hydrogen, alkyl, fluoroalkyl, cycloalkyl, cycloalkylalkyl, aryl, aralkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, each R^b is independently a direct bond or a straight or branched alkylene or alkenylene chain, and R^c is a straight or
5 branched alkylene or alkenylene chain, and where each of the above substituents is unsubstituted unless otherwise indicated.

[0080] "*N*-heteroaryl" refers to a heteroaryl radical as defined above containing at least one nitrogen and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a nitrogen atom in the heteroaryl radical. An *N*-heteroaryl radical is
10 optionally substituted as described above for heteroaryl radicals.

[0081] "*C*-heteroaryl" refers to a heteroaryl radical as defined above and where the point of attachment of the heteroaryl radical to the rest of the molecule is through a carbon atom in the heteroaryl radical. A *C*-heteroaryl radical is optionally substituted as described above for heteroaryl radicals.


[0082] "Heteroarylalkyl" refers to a radical of the formula $-R^c$ -heteroaryl, where R^c is an alkylene chain as defined above. If the heteroaryl is a nitrogen-containing heteroaryl, the heteroaryl is optionally attached to the alkyl radical at the nitrogen atom. The alkylene chain of the heteroarylalkyl radical is optionally substituted as defined above for an alkylene chain. The heteroaryl part of the heteroarylalkyl radical is optionally substituted as defined
15 above for a heteroaryl group.

[0083] The compounds, or their pharmaceutically acceptable salts may contain one or more asymmetric centers and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. When the compounds described herein contain
25 olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both *E* and *Z* geometric isomers (*e.g.*, *cis* or *trans*). Likewise, all possible isomers, as well as their racemic and optically pure forms, and all tautomeric forms are also intended to be included.

[0084] A "stereoisomer" refers to the relationship between two or more compounds made
30 up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not superimposable. The term "enantiomer" refers to two stereoisomers that are nonsuperimposable mirror images of one another. It is contemplated

that the various stereoisomers of the compounds disclosed herein, and mixtures thereof, are within the scope of the present disclosure and specifically includes enantiomers.

[0085] A "tautomer" refers to a compound wherein a proton shift from one atom of a molecule to another atom of the same molecule is possible. The compounds presented herein may exist as tautomers. In solutions where tautomerization is possible, a chemical equilibrium of the tautomers will exist. The exact ratio of the tautomers depends on several factors, including temperature, solvent, and pH. Some examples of tautomeric equilibrium are shown below.



[0086] "Optional" or "optionally" means that a subsequently described event or circumstance may or may not occur and that the description includes instances when the event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution.


General Methods for the Synthesis of Heterocyclic Benzamide Hedgehog Inhibitors

[0087] Synthetic Schemes 1-5 illustrate general methods for the synthesis of benzamide hedgehog inhibitors. The generic substituents illustrated in Schemes 1-5 are defined by Formula (I) as previously presented herein.


[0088] Scheme 1 illustrates the synthesis of benzamide hedgehog inhibitors. Methyl 2-iodo-4-methylsulfonylbenzoate is subjected to hydrolysis of the ester followed by coupling with a heterocycle substituted aniline gives the advanced benzamide intermediate. A variety of cross coupling reactions (such as those described by D. A. Evans, et al,

Tetrahedron Letters, 1998, 39, 2937-2940; D. M. T. Chan, et al, Tetrahedron Lett., 1998, 39, 2933-2936; P. Y. S. Lam, et al, Tetrahedron Lett., 1998, 39, 2941-2944; Y.-C. Wong, et al, Org. Lett., 2006, 8, 5613-5616; S. A. Weissman, D. Zewge, C. Chen, J. Org. Chem., 2005, 70, 1508-1510; M. McLaughlin, Org. Lett., 2005, 7, 4875-4878; A. Cwik, Z. Hell, F. Figueras, Org. Biomol. Chem., 2005, 3, 4307-4309; R. Kuwano, M. Yokogi, Org. Lett., 2005, 7, 945-947 among others) can be performed on the halobenzamide or on the borylated benzamide (available by the procedure of T. Ishiyama et al, J. Org. Chem., 1995, 60, 7508-7510) to give the desired benzamide hedgehog inhibitor.

10


Scheme 1

Scheme 2




[0089] Scheme 2 illustrates the synthesis of benzamide hedgehog inhibitors. Coupling of the acid with a 4-iodoaniline gives the 4-iodobenzamide. A palladium catalyzed cross coupling reaction between thiazole boronic acid or a thiazole stannane and the 4-iodobenzamide gives the benzamide hedgehog inhibitor.

Scheme 3




[0090] Scheme 3 illustrates the synthesis of benzamide hedgehog inhibitors. Methyl 3-iodo-4-substituted benzoate derivatives, or 3-boryl derivatives prepared by the method of T. Ishiyama et al, J. Org. Chem., 1995, 60, 7508-7510, are subjected to cross coupling reactions (such as those described by D. A. Evans, et al, Tetrahedron Letters, 1998, 39, 2937-2940; D. M. T. Chan, et al, Tetrahedron Lett., 1998, 39, 2933-2936; P. Y. S. Lam, et al, Tetrahedron Lett., 1998, 39, 2941-2944; Y.-C. Wong, et al, Org. Lett., 2006, 8, 5613-

5616; S. A. Weissman, D. Zewge, C. Chen, J. Org. Chem., 2005, 70, 1508-1510; M. McLaughlin, Org. Lett., 2005, 7, 4875-4878; A. Cwik, Z. Hell, F. Figueras, Org. Biomol. Chem., 2005, 3, 4307-4309; R. Kuwano, M. Yokogi, Org. Lett., 2005, 7, 945-947 among others) to give functionalized benzoates. Hydrolysis of the ester and coupling with a
 5 functionalized aniline affords the benzamide hedgehog inhibitor.

Scheme 4

[0091] Scheme 4 illustrates the general synthesis of the aniline intermediate that can be coupled to the appropriate carboxylic acid to form the desired product.

Scheme 5

[0092] Scheme 5 further illustrates the synthesis of thiazole based Hedgehog inhibitors based on amide bond formation with hexafluorophosphate (o-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium (HATU).

15 The Hedgehog Pathway

[0093] Since the late 1990's, scientists have unraveled several of the complex processes by which normal cells become cancer cells and developed a deeper knowledge of the heterogeneous nature of tumors. The mutational events which result in aberrant growth factor signaling in bulk tumor cells has led to the theory of "oncogene addiction", that
 20 ascribes cancer cell proliferation and survival to a dependence upon the activation of certain

pathways or on the activity of oncogenic proteins within these pathways. More recently, researchers have found small populations of tumor cells with “stem cell” like characteristics, commonly referred to as cancer stem cells, within human primary tumor samples. These newly described cancer stem cells replicate more slowly, are more resistant to conventional chemotherapy, and their survival appears to be a major contributor to tumor re-growth following surgery and/or chemotherapy. In contrast to bulk tumor cells, cancer stem cells appear to be more reliant on embryonic pathways for their proliferation and survival traits.

[0094] The Hedgehog Pathway: Several key signaling pathways (e.g. Hedgehog, Notch, Wnt) are involved in most processes essential to the normal development of an embryo. The Hedgehog pathway was initially discovered in *Drosophila* by Dr. Eric Wieschaus and Dr. Christiane Nusslein-Volhard, and is a major regulator for cell differentiation, tissue polarity and cell proliferation. It is also becoming clear that the Hedgehog pathway may play a crucial role in tumorigenesis when reactivated in adult tissues through either mutation or other mechanisms. It is thought that the Hedgehog pathway is an important driver of tumorigenesis in at least 1/3rd of all types of cancer.

[0095] Oncogenic mutations in the Hedgehog pathway have been found in basal cell carcinoma and medulloblastoma, and Hh over expression is associated with at least pancreatic, colon, gastric, liver and prostate cancer. The estimated incidence of cancers with ligand dependent activation of Hh in the US is > 200,000 cases annually and approximately 10-fold higher worldwide, see Table 2.

TABLE 2 Hh Pathway Over Expression in Solid Tumors

Tumor	2008 New US Cases (Deaths)	Hh Pathway Expression (% Total)	References
Colon	108,070 (49,960)	92,940 (86%)	Douard et al, Surgery 136, 665-670 (2006)
Lung	215,020 (161,840)	53,755* (25%-50%)	Watkins et al, Nature 422, 313-317 (2003)
Pancreas	37,680 (34,290)	18,840 (50%)	Thayer et al, Nature 425, 851-855. (2003)
Gastric	21,500 (10,880)	13,760 (64%)	Ma et al, Carcinogenesis 26, 1698-1705 (2005)
Hepatocellular	21,370 (18,410)	10,685 (50%)	Huang et al Carcinogenesis 27, 1334-1340 (2006)
Prostate	186,320 (28,660)	55,896 (30%)	Sanchez et al, PNAS 101, 12561-12566 (2004)
Total	589,960	245,876 (41%)	

[0096] More is becoming known about the role of cancer stem cells in the recurrence and spread of cancer. Control of the self-renewal and differentiation processes in cancer stems
5 cells is thought to be regulated by embryonic pathways including Hedgehog. Growing evidence suggests that these pathways are deregulated in several cases, leading to abnormal cellular expansion and the formation of cancer.


Hedgehog Pathway Signaling

[0097] Human Sonic Hedgehog protein (SHh) is synthesized as a 45 kDa precursor protein
10 that undergoes autocleavage to yield a 20 kDa fragment that is responsible for normal Hedgehog pathway signaling. At the cell surface that Hedgehog signal is thought to be relayed through the 12 transmembrane domain protein, Patched (Ptc) and the 7 transmembrane domain protein, Smoothened (Smo). In normal adult cells, Ptc serves as a negative regulatory of Smo activity. The binding of SHh to Ptc inhibits the normal
15 inhibitory effect of Ptc on Smo allowing Smo to transduce the SHh signal across the plasma membrane. The signal cascade initiated by Smo results in the activation of Gli transcription factors that migrate to the nucleus where they control target transcription factors effecting

cell growth and differentiation in embryonic cells and where uncontrolled activation in adult cells is associated with malignancies.

Methods of Inhibiting Hedgehog Signaling

[0098] One embodiment provides a method of inhibiting the Hedgehog pathway in a cell comprising contacting the cell with an inhibitory concentration of a compound of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;


R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -

NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[0099] Another embodiment provides the method wherein the cell is characterized by a
5 patched loss-of-function phenotype. Another embodiment provides the method wherein the
cell is characterized by a smoothened gain-of-function phenotype. Another embodiment
provides the method wherein the cell is characterized by a constitutively active smoothened
phenotype. Another embodiment provides the method wherein the cell is characterized by
expression of *Gli*.

10 [00100] One embodiment provides a method of inhibiting the activity of smoothened
protein in a cell comprising contacting the smoothened protein with an inhibitory
concentration of a compound of Formula (I):



15 or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt
thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

20 G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -
O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl -
25 NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked
heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked
heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -
SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -


NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -
CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl,
aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-
linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -
CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -
NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -
N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

10 **[00101]** Another embodiment provides the method wherein the cell is characterized
by a patched loss-of-function phenotype. Another embodiment provides the method
wherein the cell is characterized by a smoothened gain-of-function phenotype. Another
embodiment provides the method wherein the cell is characterized by a constitutively active
smoothened phenotype. Another embodiment provides the method wherein the cell is
15 characterized by expression of *Gli*.

[00102] One embodiment provides a method of inhibiting the transcriptional activity
of *Gli* transcription factor in a cell comprising contacting the cell with an inhibitory
concentration of a compound of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt
thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R^2 is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;


R^4 is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R^3 , R^5 and R^6 are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R^7 is H or C₁-C₃ alkyl.

[00103] Another embodiment provides the method wherein the cell is characterized by a patched loss-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a smoothened gain-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a constitutively active smoothened phenotype. Another embodiment provides the method wherein the cell is characterized by expression of *Gli*.

[00104] One embodiment provides a method of inhibiting *Gli*-mediated gene transcription in a cell comprising contacting the cell with an inhibitory concentration of a compound of Formula (I):



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

5 G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

10 R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

15 R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

20 R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and


R⁷ is H or C₁-C₃ alkyl.

[00105] Another embodiment provides the method wherein the cell is characterized by a patched loss-of-function phenotype. Another embodiment provides the method
25 wherein the cell is characterized by a smoothened gain-of-function phenotype. Another embodiment provides the method wherein the cell is characterized by a constitutively active smoothened phenotype. Another embodiment provides the method wherein the cell is characterized by expression of *Gli*.

Methods of Treatment

30 [00106] One embodiment provides a method of treating a human disease or disorder mediated by Hedgehog pathway comprising administering to a patient a therapeutically effective amount of a composition comprising a compound of Formula (I), or a

stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein the compound of Formula (I) has the following structure:



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl - NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;


R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.

[00107] Another embodiment provides the method wherein the disease or disorder is a proliferative disease. Another embodiment provides the method wherein the proliferative disease is selected from colon cancer, lung cancer, pancreatic cancer, gastric cancer, prostate cancer, and hepatocellular carcinoma. Another embodiment provides the method wherein the proliferative disease is selected from basal cell carcinoma, breast cancer, bone sarcoma, soft tissue sarcoma, chronic myeloid leukemia, acute myeloid leukemia, hematological cancer, medulloblastoma, rhabdomyosarcoma, neuroblastoma, pancreatic cancer, breast carcinoma, meningioma, glioblastoma, astrocytoma, melanoma, stomach cancer, esophageal cancer, biliary tract cancer, prostate cancer, small cell lung cancer, non-small cell lung cancer, glial cell cancer, multiple myeloma, colon cancer, neuroectodermal tumor, neuroendocrine tumor, mastocytoma and Gorlin syndrome. Another embodiment provides the method wherein the proliferative disease is basal cell carcinoma.

[00108] One embodiment provides a method of treating a veterinary disease or disorder mediated by Hedgehog pathway comprising administering to a subject a therapeutically effective amount of a composition comprising a compound of Formula (I), or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein the compound of Formula (I) has the following structure:



or a stereoisomer, tautomer, hydrate, solvate or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -O-, -N(H)- or —N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, 2 or 3;

G¹ is hydrogen, halogen, C₁-C₆ alkyl, C₃-C₆ cycloalkyl, -CN, -CF₃, or aryl;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

R¹ is H or C₁-C₃ alkyl;

R² is selected from hydrogen, halogen, -CN, alkyl, -CF₃, aryl, -O-alkyl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, SO₂-alkyl -

NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, and C-linked heterocycle;

R⁴ is selected from hydrogen, halogen, alkyl, alkoxy, -CN, -CF₃, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, and -CO₂alkyl;

R³, R⁵ and R⁶ are each independently selected from hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), and -N(alkyl)CON(alkyl)₂; and

R⁷ is H or C₁-C₃ alkyl.


[00109] Another embodiment provides a method of treating a veterinary disease or disorder wherein the disease or disorder is a proliferative disease selected from mast cell tumors or osteosarcoma.

EXAMPLES


I. Chemical Synthesis

Synthesis of intermediates

[00110] Intermediate 1: 3-chloro-(2-methylthiazol-4-yl)aniline



[00111] Step 1: 3-chloro-4-(bromoacetyl)nitrobenzene



[00112] A stirring solution of 3-chloro-4-acetyl-nitrobenzene (1.685 g, 8.44 mmol) (prepared as described in *J. Med. Chem.* **2005**, 48, 6066) and paratoluenesulfonic acid (2.41 g, 12.66 mmol) in acetonitrile (100 mL) was treated with NBS (1.503 g, 8.44 mmol) and heated to 80°C for 23h. Another gram of NBS was added after 21h. The reaction was cooled


to room temperature and concentrated in vacuo. The residue was taken up in EtOAc and the organic layer was washed with water (2x), brine, and dried over MgSO₄, filtered, and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 0-40% EtOAc/hexane yielded 554 mg of the titled compound as a waxy solid (23% yield): ¹H

NMR (400 MHz, DMSO-*d*₆) δ 4.95 (s, 2H), 8.09 (d, 1H), 8.35 (dd, 1H), 8.44 (d, 1H).


Step 2: 3-chloro-(2-methylthiazol-4-yl)aniline

[00113] A solution of 3-chloro-4-(bromoacetyl)nitrobenzene (546 mg, 1.96 mmol) and thioacetamide (162 mg, 2.15 mmol) in absolute EtOH (8 mL) was stirred at 85°C for 1h and cooled to room temperature. Solvent was evaporated in vacuo. A suspension of SnCl₂·2H₂O (1.46 g, 6.47 mmol) and conc. HCl (2 mL) in absolute EtOH (12 mL) was heated to reflux and the resulting clear solution was added to the residue. The reaction mixture was stirred at reflux for 1h, cooled to room temperature and poured into a solution of KOH (4.4 g) in water (50 mL) at 0°C. The aqueous layer was extracted with EtOAc (3x) and the combined organics were dried over MgSO₄, filtered, and concentrated in vacuo to give 438 mg of the titled product as a dark yellow oil (quant. yield): ¹H NMR (400 MHz, DMSO-*d*₆) δ 2.70 (s, 3H), 5.61 (broad s, 2H), 6.59 (dd, 1H), 6.69 (d, 1H), 7.58 (d, 1H), 7.62 (s, 1H); [M+H]⁺ *m/z* 225.

[00114] Intermediate 2: 2-chloro-(4-methylsulfonyl)-*N*-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide and 2-chloro-(4-methylsulfonyl)-*N*-(4-(thiazol-4-yl)phenyl)benzamide




[00115] Step 1: (2-ethoxycarbonylthiazol-4-yl)nitrobenzene



[00116] A solution of 4-bromoacetyl-nitrobenzene (1.8 g, 7.37 mmol) and ethyl thiooxamate (982 mg, 7.37 mmol) in absolute EtOH (20 mL) was stirred at 80°C for 19h, cooled to room temperature, and poured into saturated aqueous sodium carbonate. The resulting precipitate was filtered, washed with water and dried in vacuo to give 1.826 g of

the titled product as a yellow solid (89% yield): ^1H NMR (400 MHz, DMSO-*d*6) δ 1.40 (t, 3H), 4.46 (q, 2H), 8.32 (d, 2H), 8.38 (d, 2H), 8.89 (s, 1H).

Step 2: (2-ethoxycarbonylthiazol-4-yl)aniline and 4-(4-aminophenyl)thiazole



5 [00117] A suspension of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (4.01 g, 17.79 mmol) and conc. HCl (5.4 mL) in absolute EtOH (25 mL) was heated to reflux and the resulting clear solution was added to (2-ethoxycarbonylthiazol-4-yl)nitrobenzene (1.5 g, 5.39 mmol). The reaction mixture was stirred at reflux for 2h, cooled to room temperature and poured into a solution of KOH (12.1 g) in water (140 mL) at 0°C. The aqueous layer was extracted with EtOAc (3x) and the
10 combined organics were dried over MgSO_4 , filtered through a pad of silica, and concentrated in vacuo to give 540 mg of a (2:1) mixture of titled products as a dark yellow oil.

[00118] Step 3: 2-chloro-(4-methylsulfonyl)-*N*-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide and 2-chloro-(4-methylsulfonyl)-*N*-(4-(thiazol-4-yl)phenyl)benzamide
15 (1:1 mixture)


[00119] To a stirring solution of (2-ethoxycarbonylthiazol-4-yl)aniline/4-(4-aminophenyl)thiazole mixture (540 mg, 2.17 mmol) in DCM (10 mL) was added 2-chloro-4-methylsulfonylbenzoic acid (510 mg, 2.17 mmol) and DMAP (27 mg, 0.217 mmol). The reaction mixture was stirred for 5 min. then EDCI.HCl (500 mg, 2.60 mmol) was added.
20 After stirring for 17h, the reaction mixture was partitioned between water and EtOAc. The organic layer was separated, washed with water (2x) and brine, dried over MgSO_4 , filtered, and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 10-70% EtOAc/hexane yielded 520 mg of a (1:1) mixture of titled compounds as a yellow solid.

25 2-chloro-(4-methylsulfonyl)-*N*-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide:


$[\text{M}+\text{H}^+]^+ m/z$ 465.

2-chloro-(4-methylsulfonyl)-*N*-(4-(thiazol-4-yl)phenyl)benzamide: $[\text{M}+\text{H}^+]^+ m/z$ 393.

[00120] Intermediate 3: 4-methoxy-2-(morpholinomethyl)benzoic acid




[00121] Step 1: Methyl 2-(bromomethyl)-4-methoxybenzoate




[00122] To a solution of methyl 4-methoxy-2-methylbenzoate (1 g, 5.6 mmol) in CCl₄ (25 mL) was added dropwise N-bromosuccinimide (1.1 g, 6.2 mmol) previously dissolved in CCl₄ (5 mL) and a catalytic amount of benzoyl peroxide. The mixture was refluxed for 2 hours, cooled to room temperature and poured onto iced water. The aqueous mixture was extracted with DCM (3x), and the combined organics were dried over MgSO₄, filtered, and concentrated in vacuo to provide 2.1 g of the titled product as light yellow solid (~ 100% yield): [M+H⁺] *m/z* 260.

[00123] Step 2: methyl 4-methoxy-2-(morpholinomethyl)benzoate




[00124] To a solution of methyl 2-(bromomethyl)-4-methoxybenzoate (960 mg, 3.7 mmol) in DCM (18.5 mL) was added morpholine (0.68 mL, 7.8 mmol). The reaction mixture was stirred at room temperature overnight and poured onto iced water. The aqueous mixture was extracted with DCM (3x), and the combined organics were dried over MgSO₄, filtered, and concentrated in vacuo to provide a yellow oil. The crude material was purified by flash chromatography on silica using a gradient of EtOAc in hexane (0 to 50%) as eluant. The title product was obtained as a clear oil (490 mg, 50%)

[00125] Step 3: 4-methoxy-2-(morpholinomethyl)benzoic acid



[00126] To a solution of methyl 2-(bromomethyl)-4-methoxybenzoate (160 mg, 0.6 mmol) was dissolved in EtOH (5 mL) and a aqueous solution of sodium hydroxide was added (1N, 6.1 mL, 6.1 mmol). The resulting mixture was stirred at room temperature over night and concentrated under reduced pressure to produce a white solid. The compound
5 was used as is for the next step. LC-MS (M+H): 252.09


[00127] Intermediate 4: 4-(2-methyloxazol-4-yl)aniline




[00128] 2-methyl-4-(4-nitrophenyl)oxazole (J. Heterocyclic Chemistry, 1981, 885)(1.12 g, 5.48 mmol) was added to a refluxing solution of SnCl₂·2H₂O (6.33 g, 28.1 mmol), conc HCL (10 mL) and EtOH (20 mL). The reaction is stirred for 30 min then cooled to room temperature and poured into a solution of 24g of KOH in 100 mL of water. The resulting mixture was cooled in an ice bath and stirred for an additional 30 min. The product was collected by filtration as a yellow solid (1.35 g, 83%). LC-MS (M+H): 191

General Procedures:

Method A




[00129] Example 1: 2,4-dichloro-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide




[00130] To a solution of 4-(2-methylthiazol-4-yl)aniline (0.05 g, 0.26 mmol) and 2,4-dichlorobenzoic acid (0.05g, 0.26 mmol) in dimethylformamide (20 mL) and diisopropylethylamine (0.067g, 0.09 mL, 0.52 mmol) at room temperature was added HATU (0.148g, 0.39 mmol) in one portion. The reaction was stirred at room temperature for 4 hours, then poured in to water (200 mL) and stirred for 20 minutes. The product
25 (0.094g, 100%) as a light yellow solid was collected via filtration. LC-MS (M+H): 364. ¹H NMR (400 MHz, dmso-d₆): 10.67 (s, 1H), 7.95 (d, 2H), 7.87 (s, 1H), 7.80-7.77 (m, 3H), 7.69-7.58 (m, 2H), 2.74 (s, 3H).

[00131] Examples 2, 3, 4, 5 and 6 were prepared according to the Method A (see table 3)

Method B

[00132] Example 7: 2-chloro-N-(3-chloro-4-(2-methylthiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide:




[00133] To a stirring mixture of 3-chloro-(2-methylthiazol-4-yl)aniline (435 mg, 1.936 mmol), 2-chloro-4-methylsulfonylbenzoic acid (545 mg, 2.32 mmol) and DMAP (237 mg, 1.936 mmol) in DCM (8 mL) was added EDCI.HCl (445 mg, 2.32 mmol). After stirring overnight, the reaction mixture was partitioned between water and EtOAc. The organic


layer was separated, washed with saturated aqueous sodium bicarbonate, saturated aqueous ammonium chloride and brine, dried over MgSO₄, filtered, and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 20-80% EtOAc/hexane yielded 543 mg of the titled compound as a light yellow solid (63% yield).

Alternatively, some compounds were purified by precipitation: The DCM was removed from the reaction mixture and DMF and 0.1M NH₄Cl (compound 18) or 1N HCl (compounds 14,15,16,17) were added which caused the products to crash out of solution.

[00134] Examples 8, 14, 15, 16, 17, and 18 were prepared according to the Method B (see table 3)


Method C

[00135] Example 9: 2-chloro-N-methyl-4-(methylsulfonyl)-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide




[00136] To a stirring solution of 2-chloro-4-(methylsulfonyl)-*N*-(4-(2-methylthiazol-4-yl)phenyl)benzamide (50 mg, 0.123 mmol) in THF (1 mL) under nitrogen atmosphere was added 1M solution of *t*BuOK in *t*BuOH (0.14 mL) dropwise. After stirring for 15 min, CH₃I (0.009 mL, 0.14 mmol) was added and the reaction mixture was stirred at 60°C for 1h, before partitioning between water and EtOAc. The organic layer was isolated, washed with brine, and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 40-100% EtOAc/hexane yielded 40 mg of the titled compound as an off-white solid (77% yield).

Method D




[00137] Example 10: 2-chloro-*N*-(4-(thiazol-4-yl)phenyl)benzamide




[00138] A stirring suspension of 2-chloro-(4-methylsulfonyl)-*N*-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide and 2-chloro-(4-methylsulfonyl)-*N*-(4-(thiazol-4-yl)phenyl)benzamide (1:1 mixture) (100 mg, 0.215 mmol) in THF (1 mL) under nitrogen atmosphere was cooled to 0°C and treated with 1M LAH solution in THF (0.32 mL) dropwise. The reaction mixture was stirred at 0°C for 1h, quenched with 0.64 mL of 1N aqueous HCl, and partitioned between water and EtOAc. The aqueous layer was isolated, extracted with EtOAc (2x), and the combined organics were dried over MgSO₄, filtered and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 0-80% EtOAc/hexane yielded 12 mg of the titled compound as a yellow solid (18% yield).

Method E




[00139] Example 11 and 12: 2-chloro-N-(4-(2-(hydroxymethyl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide and 2-chloro-N-(4-(2-(2-hydroxypropan-2-yl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide




[00140] A stirring suspension of 2-chloro-(4-methylsulfonyl)-N-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide and 2-chloro-(4-methylsulfonyl)-N-(4-(thiazol-4-yl)phenyl)benzamide (1:1 mixture) (100 mg, 0.215 mmol) in THF (1 mL) under nitrogen atmosphere was cooled to 0°C and treated with 1.4M CH_3MgBr solution in toluene/THF (0.77 mL) dropwise. The reaction mixture was stirred at 0°C for 1h, quenched with 1 mL of 1N aqueous HCl, and partitioned between water and EtOAc. The aqueous layer was isolated, extracted with EtOAc (2x), and the combined organics were dried over MgSO_4 , filtered and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 0-20% $\text{CH}_3\text{CN}/\text{DCM}$ yielded 38 mg of 2-chloro-N-(4-(2-(hydroxymethyl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide as a white solid and 34 mg of 2-chloro-N-(4-(2-(2-hydroxypropan-2-yl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide as an off-white solid.

Method F



[00141] Example 13: 2-chloro-N-(4-(2-(hydroxymethyl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide














[00142] A stirring suspension of 2-chloro-(4-methylsulfonyl)-N-(4-(2-ethoxycarbonylthiazol-4-yl)phenyl)benzamide and 2-chloro-(4-methylsulfonyl)-N-(4-(thiazol-4-yl)phenyl)benzamide (1:1 mixture) (50 mg, 0.107 mmol) in THF (1 mL) under nitrogen atmosphere was treated with 2M LiBH₄ solution in THF (0.11 mL) dropwise. The reaction mixture was stirred at 60°C for 1h, quenched with water, and partitioned between water and EtOAc. The aqueous layer was isolated, extracted with EtOAc (3x), and the combined organics were dried over MgSO₄, filtered and adsorbed on silica. Purification on silica by flash chromatography using a gradient of 0-20% CH₃CN/DCM then 10% MeOH/DCM yielded 5 mg of the titled product as a white solid.






[00143] Compounds in Table 3 were prepared using the indicated method.


Table 3

Example	Structure	Synthesis Method	¹ H NMR (400 MHz), δ (ppm)	Activity Ranking	[M+H] ⁺
1	2,4-dichloro-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide	A	(DMSO- <i>d</i> ₆) 10.67 (s, 1H), 7.95 (d, 2H), 7.87 (s, 1H), 7.80-7.77 (m, 3H), 7.69-7.58 (m, 2H), 2.74 (s, 3H).	A	365

2	 3,4,5-trimethoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	A	(DMSO- <i>d</i> ₆) 10.23 (s, 1H), 7.96 (d, 2H), 7.87-7.80 (m, 3H), 7.32 (s, 2H), 3.90 (s, 6H), 3.75 (s, 3H), 2.74 (s, 3H).	C	385
3	 4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	A	(DMSO- <i>d</i> ₆) 10.21 (s, 1H), 8.00 (d, 2H), 7.93 (d, 2H), 7.88-7.87 (m, 3H), 7.09 (d, 2H), 3.87 (s, 3H), 2.73 (s, 3H).	A	326
4	 2-chloro-4-(methylsulfonyl)-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	A	(DMSO- <i>d</i> ₆) 10.82 (s, 1H), 8.17-8.16 (m, 1H), 8.05-7.90 (m, 5H), 7.79 (d, 2H), 3.38 (s, 3H), 2.75 (s, 3H).	A	407
5	 4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)-2-(morpholinomethyl)benzamide	A	(DMSO- <i>d</i> ₆) 11.71 (s, 1H), 8.08 (d, 1H), 7.92 (d, 2H), 7.76 (d, 2H), 7.29 (s, 1H), 7.01 (d, 1H), 6.76 (s, 1H), 3.89 (s, 3H), 3.83 (s, 4H), 2.83 (s, 3H), 2.62 (s, 4H)	A	424
6	 2-chloro-<i>N</i>-(4-(2-methyloxazol-4-yl)phenyl)-(methylsulfonyl)benzamide	A	(DMSO- <i>d</i> ₆) 10.81 (s, 1H), 8.45 (a, 1H), 8.15 (s, 1H), 8.04 (d, 1H), 8.02 (d, 1H), 7.77 (m, 4H), 2.49 (s, 3H)	B	390
7	 2-chloro-<i>N</i>-(3-chloro-4-(2-methylthiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide	B	(DMSO- <i>d</i> ₆) 2.75 (s, 3H), 3.38 (s, 3H), 7.69 (dd, 1H), 7.95 (m, 3H), 8.04 (m, 2H), 8.17 (d, 1H), 11.0 (s, 1H)	C	441

8	 2-chloro-4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 2.74 (s, 3H), 3.86 (s, 3H), 7.06 (dd, 1H), 7.17 (d, 1H), 7.58 (d, 1H), 7.79 (d, 2H), 7.87 (s, 1H), 7.93 (d, 2H), 10.5 (s, 1H)	A	359
9	 2-chloro-<i>N</i>-methyl-4-(methylsulfonyl)-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	C	(DMSO- <i>d</i> ₆) 2.69 (s, 3H), 3.25 (s, 3H), 3.46 (s, 3H), 7.34 (d, 2H), 7.72 (d, 1H), 7.76 (dd, 1H), 7.83 (d, 2H), 7.88 (d, 1H), 7.94 (s, 1H)	D	421
10	 2-chloro-<i>N</i>-(4-(thiazol-4-yl)phenyl)benzamide	D	(DMSO- <i>d</i> ₆) 7.49-7.55 (m, 2H), 7.63 (m, 2H), 7.83 (d, 2H), 8.01 (d, 2H), 8.12 (d, 1H), 9.21 (d, 1H), 10.6 (s, 1H)	B	315
11	 2-chloro-4-(methylsulfonyl)-<i>N</i>-(4-(thiazol-4-yl)phenyl)benzamide	E	(DMSO- <i>d</i> ₆) 3.38 (s, 3H), 7.81 (d, 2H), 7.94 (d, 1H), 8.03 (m, 3H), 8.16 (dd, 2H), 9.22 (d, 1H), 10.8 (s, 1H)	B	393
12	 2-chloro-<i>N</i>-(4-(2-(2-hydroxypropan-2-yl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide	E	(DMSO- <i>d</i> ₆) 1.59 (s, 6H), 3.38 (s, 3H), 6.04 (s, 1H), 7.79 (d, 2H), 7.91-7.97 (m, 4H), 8.02 (dd, 1H), 8.15 (d, 1H), 10.8 (s, 1H)	B	451

13	 2-chloro-<i>N</i>-(4-(2-(hydroxymethyl)thiazol-4-yl)phenyl)-4-(methylsulfonyl)benzamide	F	(DMSO- <i>d</i> ₆) 3.38 (s, 3H), 4.82 (broad s, 2H), 6.15 (broad s, 1H), 7.79 (d, 2H), 7.92-8.02 (m, 5H), 8.16 (d, 1H), 10.8 (s, 1H)	C	423
14	 2-chloro-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 10.63 (s, 1), 7.96 (d, 2H), 7.87 (s, 1H), 7.64 (d, 2H), 7.56-7.49 (m, 4H), 2.74 (s, 3H)	A	329
15	 2-(methylsulfonyl)-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 10.77 (s, 1), 8.06 (d, 1H), 7.96 (d, 2H), 7.81 (m, 2H), 7.79-7.76 (m, 4H), 3.42 (s, 3H), 2.74 (s, 3H)	D	373
16	 4-methoxy-2-methyl-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 10.27 (s, 1), 7.93 (d, 2H), 7.85 (s, 1H), 7.83 (d, 2H), 7.50 (d, 1H), 6.90 (m, 2H), 3.82 (s, 3H), 2.74 (s, 3H), 2.43 (s, 3H)	A	339
17	 2-fluoro-4-methoxy-<i>N</i>-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 10.30 (s, 1), 7.94 (d, 2H), 7.87 (s, 1H), 7.81 (d, 2H), 7.68 (t, 1H), 6.95 (d, 1H), 6.92 (d, 1H), 3.87 (s, 3H), 2.74 (s, 3H)	A	343


18	 4-cyano-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide	B	(DMSO- <i>d</i> ₆) 10.6 (s, 1H), 8.16 (d, 2H), 8.08 (d, 2H), 7.98 (d, 2H), 7.89-7.86 (m, 3H), 2.74 (s, 3H)	B	320
----	---	---	--	---	-----

Note: Activity Ranking refers to the percent activity compared to control alkaline phosphatase activity at 500 nM as determined by modified method A. Rank: A = 0 to 25% control activity; B = 26 to 50% control activity; C = 51 to 75% control activity; and D = 76 to 100% control activity.


5

[00144] Example 19: 4-methoxy-2-((4-methylpiperazin-1-yl)methyl)-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide

10




[00145] The example 19 is synthesized using the chemistry described for example 5 according to the following synthetic scheme.




15

[00146] Example 20: 2-((2-hydroxyethylamino)methyl)-4-methoxy-N-(4-(2-methylthiazol-4-yl)phenyl)benzamide




- 5 [00147] The example 20 is synthesized using the chemistry described for example 5 according to the following synthetic scheme.




10


[00148] The following compound are synthesized using 4-(2-methylthiazol-4-yl)aniline (commercially available) coupled with the appropriate carboxylic acid using method A or B.




example 21




example 22




example 23



example 24




example 25




example 26


[00149] The following compounds are synthesized using intermediate 1 coupled with the appropriate carboxylic acid using method A or B.




example 27




example 28




example 29




example 30




example 31




example 32




example 33




example 34




example 35




example 36



example 37




example 38




example 39

[00150] Intermediate 5: 4-(2-(trifluoromethyl)thiazol-4-yl)aniline




[00151] The compound is made according to the following scheme using 2,2,2-trifluoroethanethioamide.




5

[00152] The following examples are synthesized using intermediate 5 coupled with the appropriate carboxylic acid using method A or B.




10

[00153] Intermediate 6: 3-chloro-4-(2-(trifluoromethyl)thiazol-4-yl)aniline




[00154] The compound is made according to the following scheme using 2,2,2-trifluoroethanethioamide (see intermediate 1).




5

[00155] The following examples are synthesized using intermediate 6 coupled with the appropriate carboxylic acid using method A or B.




10


[00156] Intermediate 7: 4-(2-ethylthiazol-4-yl)aniline




[00157] The compound is made according to the following scheme using propanethioamide




5 **[00158]** The following examples are synthesized using intermediate 7 coupled with the appropriate carboxylic acid using method A or B:



[00159] Intermediate 8: 3-chloro-4-(2-ethylthiazol-4-yl)aniline








[00160] The compound is made according to the following scheme using propanethioamide (see intermediate 1).




5


[00161] The following examples are synthesized using intermediate 8 coupled with the appropriate carboxylic acid using method A or B.



10 **[00162]** Intermediate 9: 4-(2-isopropylthiazol-4-yl)aniline




[00163] The compound is made according to the following scheme using 2-methylpropanethioamide.




5


[00164] The following examples are synthesized using intermediate 9 coupled with the appropriate carboxylic acid using method A or B.



10 **[00165]** Intermediate 10: 3-chloro-4-(2-isopropylthiazol-4-yl)aniline




[00166] The compound is made according to the following scheme using 2-methylpropanethioamide (see intermediate 1).




5

[00167] The following examples are synthesized using intermediate 10 coupled with the appropriate carboxylic acid using method A or B.




10

[00168] Intermediate 11: 4-(2-cyclopropylthiazol-4-yl)aniline




[00169] The compound is made according to the following scheme using cyclopropanecarbothioamide.




5

[00170] The following examples are synthesized using intermediate 11 coupled with the appropriate carboxylic acid using method A or B.




10

[00171] Intermediate 12: 3-chloro-4-(2-cyclopropylthiazol-4-yl)aniline








[00172] The compound is made according to the following scheme using cyclopropanecarbothioamide (see intermediate 1).




5


[00173] The following examples are synthesized using intermediate 12 coupled with the appropriate carboxylic acid using method A or B.



10 **[00174]** Intermediate 13: 4-(2-phenylthiazol-4-yl)aniline




[00175] The compound is made according to the following scheme using phenylcarbothioamide.




5


[00176] The following examples are synthesized using intermediate 13 coupled with the appropriate carboxylic acid using method A or B.



10 **[00177]** Intermediate 14: 3-chloro-4-(2-phenylthiazol-4-yl)aniline




[00178] The compound is made according to the following scheme using phenylcarbothioamide (see intermediate 1).




5

[00179] The following examples are synthesized using intermediate 14 coupled with the appropriate carboxylic acid using method A or B.




10

[00180] The examples 180 and 182 are prepared according to the following scheme.




Journal of Pharmaceutical Sciences
(1969), 58(7), 852-7

[00181] The examples 181 and 184 are prepared according to the following scheme.




Journal of Heterocyclic Chemistry
(1989), 26(2), 269-75

[00182] The Example 183 is prepared according to the following scheme.




5


[00183] The Examples 185-187 are prepared according to the following scheme.



[00184] The Examples 188 and 189 are prepared according the following scheme using the appropriate carboxylic acid.



[00185] The Example 190-193 are prepared according to the following scheme using the appropriate carboxylic acid.



II. Biological Evaluation

[00186] The ability of compounds described in the current invention to inhibit hedgehog pathway signaling was determined from the following cell differentiation assays (Method A, or Modified Method A as described below) in which cellular alkaline phosphatase activity is assessed in the presence of control vehicle or test compound, as previously described in the literature (Wu et al., Chemistry & Biology, 11: 1229-1238, 2004; Couve-Privat, et al., Cancer Research, 64: 3559-3565, 2004; Dwyer et al., J. Biological Chemistry, 282: 8959-8968, 2007) with modifications as described below.

[00187] Method A: Mouse C3H10T1/2 (CCL-226TM) or M2-10B4 (CRL-1972TM) cells obtained from the American Type Tissue Culture Collection (Maryland, USA) were cultured to 60-80% confluence in Dulbecco's modified Eagle's Medium (C3H10T1/2 cells) containing heat-inactivated 10% fetal bovine serum or RPMI-1640 media (M2-10B4 cells) containing heat-inactivated 10% fetal bovine serum. Cell cultures were maintained in 10 U/mL penicillin, 100 µg/mL streptomycin and 2 mM glutamine. Cells were then trypsinized, counted and plated into 96-well microtiter plates prior to incubation overnight at 37 °C in 5% CO₂. The following morning, control vehicle (DMSO) or compound

dissolved in 100% DMSO were serially added to individual wells, 30 minutes prior to the addition of Control Buffer or Recombinant Mouse Sonic Hedgehog (Shh-N, CF, 461-SH-025/CF, R&D Systems, Minnesota, USA) to final concentration of 2 $\mu\text{g/mL}$. The 96-well microtiter plates were then incubated at 37 °C in 5% CO₂ for 72 hours before being assayed for alkaline phosphatase activity using p-nitrophenyl phosphate (pNPP, Anaspec, California, USA). Briefly, after 72 hours incubation, cell culture media was carefully aspirated from the wells of the 96-well microtiter plates and cells gently washed with phosphate-buffered saline, pH 7.4 (PBS). Following removal of PBS, cells were lysed in 50 μL RIPA lysis buffer and alkaline phosphatase activity assayed following addition of 50 μL pNPP reaction mixture for 30 minutes during which reagents were mixed by gently shaking of the plates. Absorbance was subsequently measured at 405 nm, and the concentration of test compound required to cause 50% inhibition of sonic hedgehog stimulated alkaline phosphatase activity (IC₅₀) calculated from the dose-response curve. Similarly, an IC₉₀ was determined. At least triplicate determinations for each individual test compound concentration were made and data plotted as mean + standard deviation relative to the DMSO control vehicle. Figure 1 shows the dose-reponse of cyclopamine, a positive control, in the assay described above. Figure 2 shows the dose-reponse of the compound of Example 4 in the assay described above. The compound of Example 4 was found to have an IC₅₀ of 63 nM and an IC₉₀ of 300 nM.


[00188] Modified Method A: An additional method to assess inhibition of hedgehog pathway signaling was also applied in which 100 nM of the smoothened agonist, purmorphamine (Stemgent, California) was added to confluent C3H10T1/2 cells instead of recombinant sonic hedgehog protein and 72 hours after co-incubation with compounds alkaline phosphatase activity was assayed using the alkaline phosphatase assay kit from BioAssay Systems (Haywood, California).

[00189] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

CLAIMS

We claim:

1. A compound having the structure of Formula (I):



or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is -S-, -N(H)- or -N(R¹)-;

Y is halogen, C₁-C₃ alkyl, -CN, or -CF₃;

n is 0, 1, or 2;

G¹ is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃;

G² is hydrogen, halogen, C₁-C₃ alkyl, -CN, or -CF₃, and wherein G¹ and G² can not both be hydrogen;

R¹ is H or C₁-C₃ alkyl;


R² is halogen, -CN, alkyl, aryl, -O-aryl, -O-heteroaryl, -CH₂-aryl, -CH₂-heteroaryl, -NH-aryl, -SO₂-aryl, -NH-heteroaryl, -NH-alkyl, -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), -CH₂-(C-linked heterocycle), N-linked heterocycle, or C-linked heterocycle;

R⁴ is alkoxy, -CN, -SO₂-alkyl, -SO₂NH₂, -NHSO₂-alkyl, -NHSO₂-aryl, -NHCO-alkyl, -NHCO-aryl, -NHCONH-alkyl, -NHCONH-aryl, -CONH₂, -CONH-alkyl, -CONH-aryl, -CON(alkyl)₂, -CON(aryl)₂, -CO₂H, or -CO₂alkyl; and

R³, R⁵ and R⁶ are each independently hydrogen, halogen, -CN, alkyl, aryl, heteroaryl, C-linked heterocycle, -O-alkyl, -O-aryl, -O-heteroaryl, N-linked heterocycle, -NH-alkyl, -N(alkyl)₂, -NH-aryl, -NHheteroaryl, -CO₂H, -CO₂alkyl, -SO₂alkyl, -SO₂NH₂, -SO₂NHalkyl, -SO₂N(alkyl)₂, -NHSO₂alkyl, -NHSO₂aryl, -NHCONH-alkyl, -NHCON(alkyl)₂, -N(alkyl)CONH₂, -N(alkyl)CONH(alkyl), or -N(alkyl)CON(alkyl)₂; and R⁷ is H or C₁-C₃ alkyl.

2. The compound of claim 1 wherein n is 0.

3. The compound of claim 1 wherein n is 1.
4. The compound of claim 1 wherein G^2 is H.
5. The compound of claim 1 wherein G^2 is H and G^1 is C_1 - C_3 alkyl.
6. The compound of claim 1 wherein X is -S-.
7. The compound of claim 1 wherein R^2 is -CH₂-NH-alkyl, -CH₂-N(alkyl)₂, -CH₂-(N-linked heterocycle), or -CH₂-(C-linked heterocycle).
8. The compound of claim 1 wherein R^2 is halogen, -CN, or -alkyl.
9. The compound of claim 1 wherein R^4 is -SO₂Me or -OMe.
10. The compound of claim 1 wherein R^2 is halogen and R^4 is -SO₂Me.
11. The compound of claim 1 wherein R^2 is halogen and R^4 is -OMe.
12. The compound of claim 1 wherein n is 0; G^2 is H; G^1 is C_1 - C_3 alkyl; X is -S-; R^2 is halogen and R^4 is -OMe.
13. The compound of claim 1 wherein n is 0; G^2 is H; G^1 is C_1 - C_3 alkyl; X is -S-; R^2 is halogen and R^4 is -SO₂Me.
14. The compound of claim 1 wherein the compound of Formula (I) is:



15. A pharmaceutical composition comprising the compound of any one of claims 1 to 14 and at least one pharmaceutically acceptable excipient.
16. An *in vitro* method of inhibiting the Hedgehog pathway in a cell comprising contacting the cell with an inhibitory concentration of the compound of any one of claims 1 to 14.
17. An *in vitro* method of inhibiting the activity of smoothened protein in a cell comprising contacting the smoothened protein with an inhibitory concentration the compound of any one of claims 1 to 14.
18. An *in vitro* method of inhibiting the transcriptional activity of *Gli* transcription factor in a cell comprising contacting the cell with an inhibitory concentration of the compound of any one of claims 1 to 14.

19. An *in vitro* method of inhibiting *Gli*-mediated gene transcription in a cell comprising contacting the cell with an inhibitory concentration of the compound of any one of claims 1 to 14.
20. The method of any one of claims 16-19, wherein the cell is characterized by a patched loss-of-function phenotype.
21. The method of any one of claims 16-19, wherein the cell is characterized by a smoothened gain-of-function phenotype.
22. The method of any one of claims 16-19, wherein the cell is characterized by a constitutively active smoothened phenotype.
23. The method of any one of claims 16-19, wherein the cell is characterized by expression of *Gli*.
24. Use of the compound of any one of claims 1 to 14 for treating a human disease or disorder mediated by Hedgehog pathway.
25. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for treating a human disease or disorder mediated by Hedgehog pathway.
26. The compound of any one of claims 1 to 14 for use in treating a human disease or disorder mediated by Hedgehog pathway.
27. The use of claim 24 or 25, or the compound for use of claim 26, wherein the disease or disorder is a proliferative disease.
28. The use or compound for use of claim 27, wherein the proliferative disease is colon cancer, lung cancer, pancreatic cancer, gastric cancer, prostate cancer, or hepatocellular carcinoma.
29. The use or compound for use of claim 27, wherein the proliferative disease is basal cell carcinoma, breast cancer, bone sarcoma, soft tissue sarcoma, chronic myeloid leukemia, acute myeloid leukemia, hematological cancer, medulloblastoma, rhabdomyosarcoma, neuroblastoma, pancreatic cancer, breast carcinoma, meningioma, glioblastoma, astrocytoma, melanoma, stomach cancer, esophageal cancer, biliary tract cancer, prostate cancer, small cell lung cancer, non-small cell lung cancer, glial cell cancer, multiple myeloma, colon cancer, neuroectodermal tumor, neuroendocrine tumor, mastocytoma or Gorlin syndrome.
30. The use or compound for use of claim 29, wherein the proliferative disease is basal cell carcinoma.

31. Use of the compound of any one of claims 1 to 14 for treating a veterinary disease or disorder mediated by Hedgehog pathway.
32. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for treating a veterinary disease or disorder mediated by Hedgehog pathway.
33. The compound of any one of claims 1 to 14 for use in treating a veterinary disease or disorder mediated by Hedgehog pathway.
34. The use of claim 31 or 32, or the compound for use of claim 33, wherein the disease or disorder is a proliferative disease and the proliferative disease is mast cell tumors or osteosarcoma.
35. Use of the compound of any one of claims 1 to 14 for inhibiting the Hedgehog pathway in a cell.
36. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for inhibiting the Hedgehog pathway in a cell.
37. Use of the compound of any one of claims 1 to 14 for inhibiting the activity of smoothened protein in a cell.
38. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for inhibiting the activity of smoothened protein in a cell.
39. Use of the compound of any one of claims 1 to 14 for inhibiting the transcriptional activity of *Gli* transcription factor in a cell.
40. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for inhibiting the transcriptional activity of *Gli* transcription factor in a cell.
41. Use of the compound of any one of claims 1 to 14 for inhibiting *Gli*-mediated gene transcription in a cell.
42. Use of the compound of any one of claims 1 to 14 in the manufacture of a medicament for inhibiting *Gli*-mediated gene transcription in a cell.
43. The compound of any one of claims 1 to 14 for use in inhibiting the Hedgehog pathway in a cell.
44. The compound of any one of claims 1 to 14 for use in inhibiting the activity of smoothened protein in a cell.
45. The compound of any one of claims 1 to 14 for use in inhibiting the transcriptional activity of *Gli* transcription factor in a cell.

46. The compound of any one of claims 1 to 14 for use in inhibiting *Gli*-mediated gene transcription in a cell.
47. The use of any one of claims 35-42 or the compound for use of any one of claims 43-46, wherein the cell is characterized by a patched loss-of-function phenotype.
48. The use of any one of claims 35-42 or the compound for use of any one of claims 43-46, wherein the cell is characterized by a smoothened gain-of-function phenotype.
49. The use of any one of claims 35-42 or the compound for use of any one of claims 43-46, wherein the cell is characterized by a constitutively active smoothened phenotype.
50. The use of any one of claims 35-42 or the compound for use of any one of claims 43-46, wherein the cell is characterized by expression of *Gli*.
51. The pharmaceutical composition of claim 15 for use in treating a human disease or disorder mediated by Hedgehog pathway.
52. Use of the pharmaceutical composition of claim 15 in the manufacture of a medicament for treating a human disease or disorder mediated by Hedgehog pathway.
53. Use of the pharmaceutical composition of claim 15 for treating a human disease or disorder mediated by Hedgehog pathway.

1/2

FIGURE 1




FIGURE 2

