



US00PP16500P2

(12) **United States Plant Patent**  
**Uchneat**

(10) **Patent No.:** **US PP16,500 P2**

(45) **Date of Patent:** **May 2, 2006**

(54) **DOUBLE *IMPATIENS* PLANT NAMED  
'BALOLEPURP'**

(52) **U.S. Cl.** ..... **Plt./317**

(58) **Field of Classification Search** ..... **Plt./317**  
See application file for complete search history.

(50) Latin Name: *Impatiens walleriana*  
Varietal Denomination: **Balolepurp**

(75) Inventor: **Michael S. Uchneat**, Geneva, IL (US)

*Primary Examiner*—Anne Marie Grunberg  
*Assistant Examiner*—Annette H Para

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

(74) *Attorney, Agent, or Firm*—Wood, Phillips, Katz, Clark  
& Mortimer

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 54 days.

(57) **ABSTRACT**

A new and distinct Double *Impatiens* plant named  
'Balolepurp' characterized by its fully double dark purple-  
colored flowers, medium green-colored foliage, compact,  
upright and mounded growth habit, and good basal branching  
character.

(21) Appl. No.: **11/011,315**

(22) Filed: **Dec. 13, 2004**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

**1 Drawing Sheet**

**1**

**2**

Latin Name of genus and species of plant claimed:  
*Impatiens walleriana*.  
Variety denomination: 'Balolepurp'.

Plants of the new cultivar differ from plants of the female  
parent primarily in flower form and color and from the male  
parent primarily in foliage color.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of Double *Impatiens* plant botanically known as *Impatiens*  
*walleriana* and hereinafter referred to by the cultivar name  
'Balolepurp'.

Of the many commercially available Double *Impatiens*  
cultivars known to the inventor, the most similar to the new  
cultivar is the *Impatiens walleriana* variety 'Balfiepuna',  
U.S. Plant Pat. No. 13,611. However, in side-by-side com-  
parisons conducted at West Chicago, Ill., plants of the new  
cultivar differed from plants of 'Balfiepuna' in the following  
characteristics:

The new cultivar originated in a controlled breeding  
program during April 2000 at Elburn, Ill. The objective of  
the breeding program was the development of new *Impa-*  
*tiens* cultivars with numerous, fully double flowers, excel-  
lent basal branching, and upright compact growth habit.

1. Plants of the new cultivar are shorter than plants of  
'Balfiepuna'.
2. The leaves of the new cultivar are smaller than the  
leaves of 'Balfiepuna'.

The female (seed) parent of the new cultivar was the  
proprietary *Impatiens walleriana* selection designated 3177-  
1-1-2, not patented, characterized by its small, single,  
salmon-colored flowers. The male (pollen) parent of the new  
cultivar was the proprietary *Impatiens walleriana* selection  
designated 12865-2, not patented, characterized by its dark  
green-colored foliage. The new cultivar was discovered and  
selected by the inventor as a single flowering plant within  
the progeny of the stated cross-pollination during September  
2001 at Elburn, Ill. Asexual reproduction of the new cultivar  
by terminal stem cuttings since September 2001 at West  
Chicago, Ill., has demonstrated that the new cultivar repro-  
duces true to type with all characteristics, as herein  
described, firmly fixed and retained through successive  
generations of such asexual propagation.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs show, as nearly true as it  
is reasonably possible to make the same in color illustrations  
of this type, typical flower and foliage characteristics of the  
new cultivar. Colors in the photographs differ slightly from  
the color values cited in the detailed description, which  
accurately describe the colors of 'Balolepurp'. The plants  
were grown for 8 weeks in a greenhouse at West Chicago,  
Ill.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of 'Balolepurp'.

FIG. 2 illustrates a close-up view of an individual flower  
of 'Balolepurp'.

**SUMMARY OF THE INVENTION**

The following characteristics of the new cultivar have  
been repeatedly observed and can be used to distinguish  
'Balolepurp' as a new and distinct cultivar of Double *Impa-*  
*tiens* plant:

**DETAILED BOTANICAL DESCRIPTION**

1. Fully double purple-colored flowers.
2. Medium green-colored foliage.
3. A good basal branching character.
4. A compact, upright and mounded growth habit.

The new cultivar has not been observed under all possible  
environmental conditions to date. Accordingly, it is possible  
that the phenotype may vary somewhat with variations in the  
environment, such as temperature, light intensity, and day  
length without, however, any variance in phenotype.

The chart used in the identification of colors described  
herein is the R.H.S. Colour Chart of The Royal Horticultural

Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined on Jul. 20, 2004 between 10:00 and 11:45 a.m. under natural light conditions.

The following descriptions and measurements describe plants produced from cuttings taken from stock plants and grown in a double polycarbonate-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown at West Chicago, Ill. in 10 cm pots for 8 weeks using a soilless growth medium. Greenhouse temperatures were maintained at approximately 65°–75° F. during the day and approximately 55°–62° F. during the night. Greenhouse light levels were maintained at approximately 4,000–6,000 footcandles during the day.

Botanical classification: *Impatiens walleriana* cultivar Balolepurp.

Parentage:

*Female (seed) parent*.—Proprietary *Impatiens walleriana* selection designated 3177-1-1-2, not patented.

*Male (pollen) parent*.—Proprietary *Impatiens walleriana* selection designated 12865-2, not patented.

Propagation:

*Type cutting*.—Terminal stem.

*Time to initiate roots*.—Approximately 7–14 days.

*Time to produce a rooted cutting*.—Approximately 21 days.

*Root description*.—Fine, fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Crop time*.—Approximately 6–8 weeks from planting of rooted cutting to finish.

*Growth habit*.—Compact with good basal branching.

*Form*.—Upright and mounded.

*Size*.—Height: Approximately 12.9 cm. Width (area of plant spread): Approximately 24.9 cm in width (plant spread).

*Branch*.—Quantity: Approximately 3 per plant. Strength: Strong. Length: Approximately 12 cm. Diameter: Approximately 6.8 mm. Texture: Glabrous. Color: 144A. Internode length: Approximately 1.3 cm.

*Foliage*.—Quantity of leaves per branch: Approximately 11. Fragrance: None. Type: Simple. Arrangement: Alternate. Aspect: At right angle to stem. Shape: Ovate. Apex: Cuspidate. Base: Attenuate. Margin: Crenate/ciliate. Venation pattern: Pinnate, arcuate. Size of mature foliage: Length: Approximately 1.1 cm. Width: Approximately 8.9 mm. Texture of both surfaces: Glabrous. Color of upper surface of mature foliage: Darker than 137A, with veins and mid-vein of 144A. Color of lower surface of mature foliage: Closest to 138B with veins and mid-vein of 144C. Petiole length: Approximately 5.5 mm. Petiole diameter: Approximately 0.6 mm. Petiole texture: Both surfaces are glabrous. Petiole color: 144C.

Flowering description:

*Flowering habit*.—Freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year round in greenhouse environment.

*Lastingness of individual bloom*.—Approximately 7–10 days.

*Quantity of flowers*.—Approximately 28 flowers and 20 buds per plant at any one time.

Flower description:

*Flower bud rate of opening*.—Generally it takes from 3–6 days for buds to progress from first color to fully open flowers.

*Mature flower buds (just before opening)*.—Shape: Elliptic. Length: Approximately 1.1 cm. Diameter: Approximately 1 cm. Color: N74D.

*Flower description*.—Type: Fully double. Fragrance: None. Shape: Spherical. Diameter: Approximately 3.2 cm. Depth: Approximately 1.3 cm. Borne: Above foliage facing upward or outward. Flowers are not persistent or fragrant.

*Petals*.—Quantity per flower: Approximately 29. Shape: Obovate. Margin: Entire. Apex: Retuse. Base: Attenuate. Texture: Glabrous. Appearance: Iridescent. Size of outermost petals: Length: Approximately 1.8 cm. Width: 1.6 cm. Size of innermost petals: Length: 1.2 cm. Width: 1 cm.

*Flower color*.—Upper surface of petals when fully open: Slightly darker than N74A. Lower surface of petals when fully open: Closest to N74D. Petal color fading with age to: 83A.

*Calyx*.—Quantity of sepals: Three, with the lower sepal modified into a spur. Sepal apex: Acuminate. Sepal texture: Both surfaces are glabrous. Lateral sepal shape: Lanceolate. Lateral sepal apex: Acute. Lateral sepal length: Approximately 3 mm. Lateral sepal width: Approximately 1 mm. Lateral sepal color: Both surfaces are 144C. Lower sepal shape: Ovate. Lower sepal apex: Acute. Lower sepal base forms spur. Lower sepal length: Approximately 7 mm. Lower sepal width: Approximately 7 mm. Lower sepal, color of both surfaces: 154D with N144D at tip.

*Spur*.—Quantity: One per flower. Length: Approximately 1.1 cm. Diameter at base: Approximately 2 mm. Diameter at tip: Approximately 1 mm. Texture: Glabrous. Color: 154D with overlay of 71B.

*Peduncles*.—Length: Approximately 2 cm. Diameter: Approximately 0.4 mm. Texture: Glabrous. Strength: Strong. Angle to stem: Acute. Color: 144C.

*Reproductive organs*.—None observed.

Seed and fruit development: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of Double *Impatiens* plant named 'Balolepurp', substantially as herein shown and described.

\* \* \* \* \*



FIG. 1




FIG. 2