

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 October 2007 (11.10.2007)

PCT

(10) International Publication Number
WO 2007/113256 A1

(51) International Patent Classification:

C07D 417/06 (2006.01) **C07D 417/14** (2006.01)
C07D 401/06 (2006.01) **A61K 31/506** (2006.01)
C07D 239/48 (2006.01) **A61K 31/5377** (2006.01)
C07D 403/06 (2006.01) **A61P 31/18** (2006.01)
C07D 513/04 (2006.01)

(21) International Application Number:

PCT/EP2007/053113

(22) International Filing Date: 30 March 2007 (30.03.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

06112045.7 30 March 2006 (30.03.2006) EP

(71) Applicant (for all designated States except US): **TI-BOTEC PHARMACEUTICALS LTD.** [IE/IE]; Eastgate Village, Eastgate, Little Island, Co Cork (IE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GUILLEMONT, Jérôme, Emile, Georges** [FR/FR]; 51bis, route de Muids, F-27430 Andé (FR). **MORDANT, Céline, Isabelle** [FR/FR]; 52 Avenue Victor Segoffin, Batiment B4, F-31400 Toulouse (FR). **SCHMITT, Benoit, Antoine** [FR/BE]; Lange Heistraat 60, B-2800 Mechelen (BE).

(74) Agent: **WANTE, Dirk**; Generaal De Wittelaan L 11B 3, B-2800 Mechelen (BE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: HIV INHIBITING 5-(HYDROXYMETHYLENE AND AMINOMETHYLENE) SUBSTITUTED PYRIMIDINES

(57) Abstract: This invention concerns 5-(hydroxymethylene and aminomethylene)pyrimidine derivatives having HIV (Human Immunodeficiency Virus) replication inhibiting properties, to the preparation thereof and to pharmaceutical compositions comprising these compounds.

WO 2007/113256 A1

HIV INHIBITING 5-(HYDROXYMETHYLENE AND AMINOMETHYLENE)
SUBSTITUTED PYRIMIDINES

5 This invention concerns 5-(hydroxymethylene and aminomethylene)pyrimidine derivatives having HIV (Human Immunodeficiency Virus) replication inhibiting properties, to the preparation thereof and to pharmaceutical compositions comprising these compounds.


10 Resistance of the HIV virus against currently available HIV drugs continues to be a major cause of therapy failure. This has led to the introduction of combination therapy of two or more anti-HIV agents usually having a different activity profile. Significant progress was made by the introduction of HAART therapy (Highly Active
15 Anti-Retroviral Therapy), which has resulted in a significant reduction of morbidity and mortality in HIV patient populations treated therewith. HAART involves various combinations of nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs) and protease inhibitors (PIs). But even these multidrug therapies do not completely eliminate HIV and long-term treatment often
20 leads to multidrug resistance. In many cases, resistant virus is carried over to newly infected individuals, resulting in limited therapy options for these drug-naïve patients.

Therefore there is a continued need for new combinations of active ingredients that are effective against HIV. New types of anti-HIV agents, differing in chemical structure and activity profile are useful in new types of combination therapy. Finding such
25 active ingredients therefore is a highly desirable goal to achieve.

The present invention is aimed at providing particular novel series of pyrimidine derivatives having HIV replication inhibiting properties. WO 99/50250, WO 00/27825 and WO 01/85700 disclose certain substituted aminopyrimidines having HIV
30 replication inhibiting properties.

The compounds of the invention differ from prior art compounds as regards their chemical structure, as well as their pharmacological profile. It has been found that the introduction of certain substituents in the 5-position of specifically substituted
35 pyrimidines results in compounds the compounds not only acting favorably in terms of their capability to inhibit the replication of Human Immunodeficiency Virus (HIV), but also by their improved ability to inhibit the replication of mutant strains, in particular strains which have become resistant to one or more known NNRTI drugs, which strains are referred to as drug or multidrug resistant HIV strains.

Thus in one aspect, the present invention concerns compounds of formula



the stereochemically isomeric forms thereof, the pharmaceutically acceptable addition
 5 salts thereof, the pharmaceutically acceptable hydrates or solvates thereof, the N-oxides thereof, wherein

each **R**¹ independently is hydrogen; aryl; formyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyl;
 C₁₋₆alkyloxycarbonyl;

- 10 **R**², **R**³, **R**⁷ and **R**⁸ independently are hydrogen; hydroxy; halo; C₃₋₇cycloalkyl;
 C₁₋₆alkyloxy; carboxyl; C₁₋₆alkyloxycarbonyl; cyano; nitro; amino; mono- or
 di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; -C(=O)R¹⁰; C₁₋₆alkyl
 optionally substituted with halo, cyano or -C(=O)R¹⁰; C₂₋₆alkenyl optionally
 15 substituted with halo, cyano or -C(=O)R¹⁰; C₂₋₆alkynyl optionally substituted with
 halo, cyano or -C(=O)R¹⁰;

R⁴ and **R**⁹ independently are hydroxy; halo; C₃₋₇cycloalkyl; C₁₋₆alkyloxy; carboxyl;
 C₁₋₆alkyloxycarbonyl; formyl; cyano; nitro; amino; mono- or di(C₁₋₆alkyl)amino;
 polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; -C(=O)R¹⁰; cyano; -S(=O)_rR¹⁰;
 -NH-S(=O)_rR¹⁰; -NHC(=O)H; -C(=O)NHNH₂; -NHC(=O)R¹⁰; Het; -Y-Het;
 20 C₁₋₁₂alkyl optionally substituted with halo, cyano, amino, mono- or
 di(C₁₋₆alkyl)-amino, -C(=O)-R¹⁰, Het or with C₁₋₆alkyloxy; C₂₋₁₂alkenyl optionally
 substituted with halo, cyano, amino, mono- or di(C₁₋₆alkyl)amino, -C(=O)-R¹⁰, Het
 or with C₁₋₆alkyloxy; C₂₋₁₂alkynyl optionally substituted with halo, cyano, amino,
 mono- and di(C₁₋₆alkyl)amino, -C(=O)-R¹⁰, Het or with C₁₋₆alkyloxy;

- 25 **R**⁵ is C₁₋₆alkyl optionally substituted with Ar or with Het; C₂₋₆alkenyl optionally
 substituted with Ar or with Het; C₂₋₆alkynyl optionally substituted with Ar or with Het;
 C₃₋₇cycloalkyl; Ar; Het;

R⁶ is H, C₁₋₆alkyl, Het;

Y is -OR¹¹, -NR¹²R¹³;

each **R**¹⁰ independently is C₁₋₆alkyl, amino, mono- or di(C₁₋₆alkyl)amino or polyhalo-C₁₋₆alkyl;

R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy, phenyl or pyridyl;

5 **R**¹² is hydrogen or C₁₋₆alkyl;

R¹³ is hydrogen or C₁₋₆alkyl; or **R**¹² and **R**¹³ taken together with the nitrogen atom to which they are substituted form pyrrolidinyl; piperidinyl; morpholinyl; piperazinyl; piperazinyl optionally substituted with C₁₋₆alkyl or C₁₋₆alkylcarbonyl; imidazolyl;

X is -NR¹-, -O-, -C(=O)-, -CH₂-, -CHOH-, -S-, -S(=O)_r-;

10 each **Y** independently is -NR¹-, -O-, -C(=O)-, -S-, -S(=O)_r-;
each *r* independently is 1 or 2;

Het is a 5- or 6-membered completely unsaturated ring wherein one, two, three or four ring members are hetero atoms, each independently selected from nitrogen, oxygen and sulfur, and wherein the remaining ring members are carbon atoms; and, where possible, any nitrogen ring member may optionally be substituted with C₁₋₆alkyl; which 5- or 6-membered ring may optionally be annelated with a benzene, thiophene or pyridine ring; and wherein any ring carbon atom, including any carbon of an optionally annelated benzene, thiophene or pyridine ring, may, each independently, optionally be substituted with a substituent selected from halo, hydroxy, mercapto, cyano, C₁₋₆alkyl, hydroxyC₁₋₄alkyl, carboxyC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkyloxycarbonylC₁₋₄alkyl, cyanoC₁₋₄alkyl, mono- and di(C₁₋₄alkyl)aminoC₁₋₄alkyl, Het¹C₁₋₄alkyl, arylC₁₋₄alkyl, polyhaloC₁₋₄alkyl, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, aryl-C₂₋₄alkenyl, C₁₋₄alkyloxy, -OCONH₂, polyhaloC₁₋₄alkyloxy, aryloxy, amino, mono- and di-C₁₋₄alkylamino, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, 4-C₁₋₆alkylpiperazinyl, C₁₋₄alkylcarbonylamino, formyl, C₁₋₄alkylcarbonyl, C₁₋₄alkyloxy-carbonyl, aminocarbonyl, mono- and diC₁₋₄alkylaminocarbonyl, aryl, Het¹;

Het¹ is pyridyl, thienyl, furanyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, quinolinyl, benzothienyl, benzofuranyl; which each may be optionally substituted with one or two C₁₋₄alkyl radicals;

each **aryl** independently is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono and di(C₁₋₆alkyl)-aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, phenylC₁₋₆alkyloxy,

C₁₋₆alkyloxycarbonyl, aminosulfonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, phenyl, Het and -Y-Het.

5 Het is a completely unsaturated ring, meaning that the ring has the maximum number of double bonds. One type of Het rings are the aromatic heterocycles. In one embodiment Het as a whole, i.e. including the optionally annelated benzene, thiophene or pyridine ring may optionally be substituted with one, two or three of the substituents mentioned above or hereinafter. In particular, Het contains no more than two oxygen or sulfur ring atoms, more in particular no more than one oxygen or sulfur ring atom.

10

In one embodiment, each Het independently is pyridyl, thienyl, furanyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, quinolinyl, benzothienyl, benzofuranyl, benzoxazolyl, benzothiazolyl, imidazothiazolyl; which each may optionally be substituted with one or two substituents each independently
15 selected from C₁₋₆alkyl; halo; hydroxy; cyano; C₁₋₆alkyloxy; C₂₋₁₂alkenyl substituted with halo, hydroxy or with cyano.

As used hereinbefore or hereinafter C₁₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon
20 atoms such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methylpropyl, t.butyl; C₁₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the group defined for C₁₋₄alkyl and 1-pentyl, 2-pentyl, 1-hexyl, 2-hexyl, 3-hexyl, 2-methylbutyl, 3-methylpentyl and the like; C₁₋₂alkyl defines methyl or ethyl; C₃₋₇cycloalkyl is generic
25 to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. Preferred amongst C₁₋₆alkyl are C₁₋₄alkyl or C₁₋₂alkyl. Preferred amongst C₃₋₇cycloalkyl are cyclopentyl and cyclohexyl.

The term "C₂₋₆alkenyl" as a group or part of a group defines straight and branched
30 chained hydrocarbon radicals having saturated carbon-carbon bonds and at least one double bond, and having from 2 to 6 carbon atoms, such as, for example, ethenyl (or vinyl), 1-propenyl, 2-propenyl (or allyl), 1-butenyl, 2-butenyl, 3-butenyl, 2-methyl-2-propenyl, 1-pentenyl, 2-pentenyl, 3-pentenyl, 2-methyl-1-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 2-methyl-2-pentenyl, 1,2-dimethyl-
35 1-butenyl and the like. Preferred are C₂₋₆alkenyls having one double bond. Of interest amongst C₂₋₆alkenyl radicals are the C₂₋₄alkyl radicals. The term "C₃₋₆alkenyl" is as C₂₋₆alkenyl but is limited to unsaturated hydrocarbon radicals having from 3 to 6 carbon atoms. In the instances where a C₃₋₆alkenyl is linked to a heteroatom, the carbon

atom linked to the heteroatom by preference is saturated. The term "C₂₋₁₂alkenyl" is as C₂₋₆alkenyl but has from 2 to 12 carbon atoms and includes the C₂₋₆alkenyl radicals and the higher homologs such as 1-heptenyl, 2-heptenyl, 3-heptenyl, 1-methyl-1-hexenyl, 1,2-dimethyl-1-pentenyl, 2-methyl-1-hexenyl, 2-ethyl-2-pentenyl, 3-propyl-2-hexenyl, 1-octenyl, 2-octenyl, 1-nonenyl, 1-decenyl, 1-undecenyl, 1-dodecenyl and the like. Preferred amongst C₂₋₁₂alkenyl are the C₂₋₆alkenyl radicals.

The term "C₂₋₆alkynyl" as a group or part of a group defines straight and branched chained hydrocarbon radicals having saturated carbon-carbon bonds and at least one triple bond, and having from 2 to 6 carbon atoms, such as, for example, ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, 2-methyl-2-propynyl, 2-pentyne, 3-pentyne, 2-hexynyl, 3-hexynyl, 4-hexynyl, 2-methyl-2-butyne, 2-methyl-2-pentyne and the like. Preferred are C₂₋₆alkynyls having one triple bond. Of interest amongst C₂₋₆alkynyl radicals are the C₂₋₄alkynyl radicals. The term "C₃₋₆alkynyl" is as C₂₋₆alkynyl but is limited to unsaturated hydrocarbon radicals having from 3 to 6 carbon atoms. In the instances where a C₃₋₆alkynyl is linked to a heteroatom, the carbon atom linked to the heteroatom by preference is saturated. The term "C₂₋₁₂alkynyl" is as C₂₋₆alkynyl but has from 2 to 12 carbon atoms and includes the C₂₋₆alkynyl radicals and the higher homologs such as 1-heptyne, 2-heptyne, 1-octynyl, 2-octynyl, 1-nonyne, 1-decynyl, 1-undecynyl, 1-dodecynyl and the like. Preferred amongst C₂₋₁₂alkynyl are the C₂₋₆alkynyl radicals.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom.

The terms carboxyl, carboxy or hydroxycarbonyl refer to a group -COOH.

The term "halo" is generic to fluoro, chloro, bromo or iodo.

The term "polyhaloC₁₋₆alkyl" as a group or part of a group, e.g. in polyhaloC₁₋₆alkoxy, is defined as mono- or polyhalo substituted C₁₋₆alkyl, in particular C₁₋₆alkyl substituted with up to one, two, three, four, five, six, or more halo atoms, such as methyl or ethyl with one or more fluoro atoms, for example, difluoromethyl, trifluoromethyl, trifluoro-ethyl. Preferred is trifluoromethyl. Also included are perfluoroC₁₋₆alkyl groups, which are C₁₋₆alkyl groups wherein all hydrogen atoms are replaced by fluoro atoms, e.g. pentafluoroethyl. In case more than one halogen atom is attached to an alkyl group within the definition of polyhaloC₁₋₆alkyl, the halogen atoms may be the same or different.

Any of the heterocycles mentioned in the definitions of Het is meant to comprise any isomer such as for example oxadiazole may be 1,2,4-oxadiazole, 1,3,4-oxadiazole, or 1,2,3-oxadiazole; likewise for thiadiazole which may be 1,2,4-thiadiazole,
5 1,3,4-thia-diazole, or 1,2,3-thiadiazole; similarly, imidazole may be 1H-imidazole or 3H-imidazole.

Whenever a radical occurs in the definition of the compounds of formula (I) or in any of the subgroups specified herein, said radical independently is as specified above in
10 the definition of the compounds of formulas (I) or in the more restricted definitions as specified hereinafter.

It should also be noted that the radical positions on any molecular moiety used in the definitions may be anywhere on such moiety as long as it is chemically stable. For
15 instance pyridine includes 2-pyridine, 3-pyridine and 4-pyridine; pentyl includes 1-pentyl, 2-pentyl and 3-pentyl.

When any variable (e.g. halogen, C₁₋₆alkyl, aryl, Het, etc.) occurs more than one time in any moiety, each definition is independent.
20

Any limited definitions of the radicals specified herein are meant to be applicable to the group of compounds of formula (I) as well as to any subgroup defined or mentioned herein.

25 Lines drawn from substituents into ring systems indicate that the bond may be attached to any of the suitable ring atoms.

The term "compounds of formula (I)", or any similar terms such as "compounds of the invention" and the like, is meant to also comprise any *N*-oxide forms of the compounds
30 of formula (I), which are compounds of formula (I) wherein one or several nitrogen atoms are oxidized to the *N*-oxide form.

The pharmaceutically acceptable addition salts that the compounds of the present invention are able to form can conveniently be prepared using the appropriate acids,
35 such as, for example, inorganic acids such as hydrohalic acids, e.g. hydrochloric or hydrobromic acid, sulfuric, hemisulphuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, aspartic, dodecyl-sulphuric, heptanoic, hexanoic, nicotinic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, citric, methanesulfonic, ethanesulfonic,

benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-amino-salicylic, pamoic and the like acids. Conversely said acid addition salt forms can be converted by treatment with an appropriate base into the free base form.

- 5 The compounds of formula (I) containing acidic protons may be converted into their pharmaceutically acceptable metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g.
- 10 primary, secondary and tertiary aliphatic and aromatic amines such as methylamine, ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethyl-amine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-*n*-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, the benzathine,
- 15 *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.
- 20 The invention also comprises the hydrates and solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

It will be appreciated that some of the compounds of formula (I) and the addition salts

25 thereof may contain one or more centers of chirality and exist as stereochemically isomeric forms. Of special interest are those compounds of formula (I), which are stereochemically pure.

The term "stereochemically isomeric forms" as used hereinbefore defines all the

30 possible stereoisomeric forms, which the compounds of formula (I) and the addition salts thereof may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and

35 their salts or solvates substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Thus, when a compound of formula (I) is for instance specified as (E), this means that the compound is substantially free of the (Z) isomer. In particular,

stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-configuration.

Compounds having double bonds can have an E (entgegen) or Z (zusammen)
5 -stereochemistry at said double bond. The terms *cis*, *trans*, R, S, E and Z are well known to a person skilled in the art.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included
10 within the scope of the present invention.

The present invention is also intended to include any isotopes of atoms present in the compounds of the invention. For example, isotopes of hydrogen include tritium and deuterium and isotopes of carbon include C-13 and C-14.
15

Whenever used hereinabove or hereinafter, the terms “compounds of formula (I)”, “the present compounds”, “the compounds of the present invention” or any equivalent terms, and similarly, the terms “subgroups of compounds of formula (I)”, “subgroups of the present compounds”, “subgroups of the compounds of the present invention” or
20 any equivalent terms, are meant to include the compounds of general formula (I), or subgroups of the compounds of general formula (I), as well as their salts and stereoisomers.

Whenever mention is made hereinbefore or hereinafter that substituents can be selected
25 each independently out of a list of definitions, such as for example for R⁸ and R⁹, any possible combinations are intended to be included, which are chemically possible or which lead to molecules of such chemical stability that they can be processed in standard pharmaceutical procedures.

30 Embodiment A of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein R¹ is hydrogen.


Embodiment B of the present invention comprises those compounds of formula (I) or
35 any of the subgroups of compounds of formula (I), such as those of embodiment A, wherein

(a) R², R³, R⁷ and R⁸ independently are hydrogen; hydroxy; halo; C₁₋₆alkyl; C₃₋₇cyclo-alkyl; C₁₋₆alkyloxy; carboxyl; C₁₋₆alkyloxycarbonyl; cyano; nitro; amino; mono- or di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; -C(=O)R¹⁰;

- (b) R^2 , R^3 , R^7 and R^8 independently are hydrogen; hydroxy; halo; C_{1-6} alkyl; C_{1-6} alkyloxy; carboxyl; C_{1-6} alkyloxycarbonyl; cyano; nitro; amino; mono- or di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl; $-C(=O)R^{10}$;
- (c) R^2 , R^3 , R^7 and R^8 independently are hydrogen; hydroxy; halo; C_{1-6} alkyl; C_{1-6} alkyloxy; cyano; amino; mono- or di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl;
- (d) R^2 , R^3 , R^7 and R^8 independently are hydrogen; halo; C_{1-6} alkyl; cyano;
- (e) R^2 , R^3 , R^7 and R^8 independently are hydrogen; halo; C_{1-6} alkyl; cyano;
- (f) R^2 and R^3 are hydrogen and R^7 and R^8 independently are hydrogen; halo; cyano.
- 10 Embodiment C of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A or B, wherein
- (a) R^4 and R^9 independently are hydroxy; halo; C_{1-6} alkyloxy; carboxyl; C_{1-6} alkyloxy-carbonyl; formyl; cyano; amino; mono- or di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl; $-C(=O)R^{10}$; Het; -Y-Het; C_{1-12} alkyl optionally substituted with halo, cyano, amino, mono- or di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het; C_{2-12} alkenyl optionally substituted with halo, cyano, amino, mono- or di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het; C_{2-12} alkynyl optionally substituted with halo, cyano, amino, mono- and di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het;
- 15 (b) R^4 and R^9 independently are hydroxy; halo; C_{1-6} alkyloxy; carboxyl; C_{1-6} alkyloxy-carbonyl; formyl; cyano; amino; mono- or di(C_{1-6} alkyl)amino; polyhalo C_{1-6} alkyl; $-C(=O)R^{10}$; Het; -Y-Het; C_{1-12} alkyl optionally substituted with halo, cyano, amino, mono- and di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het; C_{2-12} alkenyl optionally substituted with halo, cyano, amino, mono- or di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het; C_{2-12} alkynyl optionally substituted with halo, cyano, amino, mono- or di(C_{1-6} alkyl)amino, $-C(=O)-R^{10}$, Het; and wherein each Het in particular is independently selected from thienyl, furanyl, oxazolyl, thiazolyl, optionally substituted with halo, C_{1-6} alkyl, cyano, carboxyl, $-C(=O)-R^{10}$;
- 20 (c) R^4 and R^9 independently are hydroxy; halo; C_{1-6} alkyloxy; carboxyl; C_{1-6} alkyloxy-carbonyl; cyano; amino; mono- or di(C_{1-6} alkyl)amino; $-C(=O)R^{10}$; Het; -Y-Het; C_{1-6} alkyl optionally substituted with cyano, $-C(=O)-R^{10}$, Het; C_{2-6} alkenyl optionally substituted with cyano, $-C(=O)-R^{10}$, Het; C_{2-6} alkynyl optionally substituted with cyano, $-C(=O)-R^{10}$, Het; and wherein each Het in particular is independently selected from thienyl, furanyl, oxazolyl, thiazolyl, optionally substituted with halo, C_{1-6} alkyl, cyano, carboxyl, $-C(=O)-R^{10}$;
- 25 (d) R^4 and R^9 independently are halo; carboxyl; C_{1-6} alkyloxycarbonyl; cyano; $-C(=O)R^{10}$; Het; -Y-Het; C_{1-6} alkyl optionally substituted with cyano, $-C(=O)-R^{10}$, Het; C_{2-12} alkenyl optionally substituted with cyano, $-C(=O)-R^{10}$, Het; and wherein
- 30
- 35

- each Het in particular is independently selected from thienyl, furanyl, oxazolyl, thiazolyl, optionally substituted with halo, C₁₋₆alkyl, cyano, carboxyl, -C(=O)-R¹⁰;
- (e) R⁴ and R⁹ independently are cyano; -C(=O)R¹⁰; Het; C₁₋₆alkyl optionally substituted with cyano, -C(=O)-R¹⁰, Het; C₂₋₆alkenyl optionally substituted with cyano, -C(=O)-R¹⁰, Het; and wherein each Het in particular is independently thienyl or furanyl, each optionally substituted with cyano, -C(=O)-R¹⁰;
- (f) R⁴ and R⁹ independently are cyano; C₁₋₆alkyl substituted with cyano; C₂₋₆alkenyl substituted with cyano.
- Embodiment D of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, or C, wherein
- (a) R⁵ is C₁₋₆alkyl substituted with Ar or with Het, C₂₋₆alkenyl, C₂₋₆alkynyl optionally substituted with Het, C₃₋₇cycloalkyl, Ar, Het;
- R⁶ is H, C₁₋₆alkyl, Het;
- Y is OR¹¹, NR¹²R¹³;
- R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy, phenyl or pyridyl;
- R¹² is hydrogen or C₁₋₆alkyl;
- R¹³ is hydrogen or C₁₋₆alkyl; or R¹² and R¹³ taken together with the nitrogen atom to which they are substituted form imidazolyl;
- (b) R⁵ is C₂₋₆alkenyl, C₂₋₆alkynyl substituted with Het, Ar, Het;
- R⁶ is H, C₁₋₆alkyl, Het;
- Y is OR¹¹, NR¹²R¹³;
- R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy;
- R¹² is hydrogen or C₁₋₆alkyl;
- R¹³ is hydrogen;
- wherein in (a) or (b) Ar and Het may be as defined for the compounds of formula (I) or (I'), or any subgroup thereof or more in particular
- Ar is phenyl optionally substituted with C₁₋₆alkyloxy; and/or
- (i) Het is pyridyl, thiazolyl, imidazolyl, benzothiazolyl, imidazothiazolyl, each optionally substituted with C₁₋₆alkyl, halo; or more in particular
- (ii) Het is pyridyl, thiazolyl, imidazolyl, each optionally substituted with C₁₋₆alkyl, halo; or more in particular
- (iii) Het is pyridyl, thiazolyl the latter optionally substituted with C₁₋₆alkyl.

Of particular interest are compounds of formula (I) or any of the subgroups of compounds of formula (I) wherein the radical



Embodiment E of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C or D, wherein

- 5 each R¹⁰ independently is C₁₋₆alkyl, amino, mono- or di(C₁₋₆alkyl)amino.

Embodiment F of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C, D or E, wherein

- 10 X is -NR¹-, -O-, -S-, -S(=O)_r-;

X is -NR¹-, -O-;

X is -NR¹-;

X is -NH-;

- 15 Embodiment G of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C, D, E or F, wherein
each Y independently is -NR¹-, -O-, -S-, -S(=O)_r-; or each Y independently is -NR¹-.


- 20 Embodiment H of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C, D, E, F or G, wherein each r independently is 2.

- Embodiment I of the present invention comprises those compounds of formula (I) or
25 any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C, D, E, F, G and H, wherein each Het independently is pyridyl, thienyl, furanyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, quinolinyl, benzothienyl, benzofuranyl; which each may optionally be substituted with one or two substituents each independently selected from C₁₋₆alkyl, halo, hydroxy,
30 cyano, C₁₋₆alkyloxy, C₂₋₁₂alkenyl substituted with halo, hydroxy or with cyano.

- Embodiment J of the present invention comprises those compounds of formula (I) or any of the subgroups of compounds of formula (I), such as those of embodiments A, B, C, D, E, F, G, H and I, wherein each aryl independently is phenyl or phenyl substituted
35 with one, two or three substituents each independently selected from those mentioned above or in particular from:

- (a) halo, hydroxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, hydroxyC₁₋₆alkyl, amino-C₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, phenylC₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, aminosulfonyl, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, phenyl, Het or -Y-Het; or
- (b) halo, hydroxy, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)-aminoC₁₋₆alkyl, C₁₋₆alkyloxy, phenylC₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, cyano, polyhaloC₁₋₆alkyl, aminocarbonyl.

10 One embodiment of the present invention concerns compounds of formula



the pharmaceutically acceptable addition salts or stereochemically isomeric forms thereof, wherein X, Y, R¹, R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are as defined above.

- 15 In a particular embodiment, R⁹ in the compounds of formula (I) or (I'), or any subgroup thereof, is -CH₂-CH₂-CN, -CH=CH-CN, or -C≡C-CN. Of interest are those compounds wherein R⁹ is the (E)-isomer of -CH=CH-CN.

Another embodiment relates to those compounds of formula (I) or (I'), or any subgroup thereof, wherein one or more of the following restrictions apply:

- 20 (i) each R¹ independently is hydrogen, aryl, formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl;
- (ii) R⁴ is hydroxy, halo, C₁₋₆alkyl, carboxyl, cyano, -C(=O)R¹⁰, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl;
- 25 (iii) X is -NR¹-, -O-, -S-, -S(=O)_r-;
- (iv) R⁷ is H, C₁₋₆alkyl, halo;
- (v) R⁸ is H, C₁₋₆alkyl, halo;
- (vi) R⁵ is C₁₋₆alkyl substituted with Ar or with Het; C₂₋₆alkenyl; C₂₋₆alkynyl optionally substituted with Het; C₃₋₇cycloalkyl; Ar; Het;
- 30 R⁶ is H, C₁₋₆alkyl, Het;

- Y is -OR¹¹, -NR¹²R¹³;
 R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy,
 phenyl or pyridyl;
 R¹² is hydrogen or C₁₋₆alkyl;
 5 R¹³ is hydrogen or C₁₋₆alkyl; or R¹² and R¹³ taken together with the nitrogen atom
 to which they are substituted form imidazolyl;
- (vii) each aryl is phenyl or phenyl substituted with one, two, three, four or five
 substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl,
 hydroxyc₁₋₆alkyl, aminoc₁₋₆alkyl, mono and di(C₁₋₆alkyl)aminoc₁₋₆alkyl,
 10 C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl,
 C₁₋₆alkyl-thio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy,
 aminocarbonyl.
- Another embodiment relates to those compounds of formula (I) or (I'), or any subgroup
 15 thereof, wherein one or more of the following restrictions apply:
- (i) R⁹ is -CH₂-CH₂-CN or -CH=CH-CN ; or in particular wherein R⁹ is
 -CH=CH-CN;
- (ii) R¹ is hydrogen, formyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl;
 20 (ii-a) R¹ is hydrogen, C₁₋₆alkyl;
 (ii-b) R¹ is hydrogen, methyl;
 (ii-c) R¹ is hydrogen;
- (iii) R⁴ is cyano, aminocarbonyl; or wherein (iii-a) R² is cyano.
- (iv) X is -NR¹-, -O-;
 25 (iv-a) X is -NR¹-,
 (iv-b) X is -NH-, -N(C₁₋₄alkyl)-, -O-;
 (iv-c) X is -NH-;
- (v) R⁷ is H, C₁₋₆alkyl, halo; (v-a) R⁷ is H, C₁₋₄alkyl, halo; (v-b) R⁷ is C₁₋₄alkyl.
- (vi) R⁸ is H, C₁₋₆alkyl, halo; (v-a) R⁸ is H, C₁₋₄alkyl, halo; (v-b) R⁸ is C₁₋₄alkyl;
- 30 (vii) R⁵ is C₂₋₆alkenyl, C₂₋₆alkynyl substituted with Het, Ar, Het;
 R⁶ is H, C₁₋₆alkyl, Het;
 Y is OR¹¹, NR¹²R¹³;
 R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy;
 R¹² is hydrogen or C₁₋₆alkyl;
 35 R¹³ is hydrogen.


Still other subgroups of the compounds of formula (I) or (I') are those compounds of formula (I) or (I'), or any subgroup thereof, wherein

(a) R^{10} is hydrogen, C_{1-4} alkyl; or wherein (b) R^{10} is hydrogen or C_{1-2} alkyl.


Other subgroups of the compounds of formula (I) or (I') are those compounds of formula (I) or (I'), or any subgroup thereof, wherein

- 5 (a) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono and di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro,
10 polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl.
- (b) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono and di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, trifluoromethyl,
15 trifluoromethoxy, aminocarbonyl.
- (c) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino- C_{1-6} alkyl, mono and di(C_{1-6} alkyl)amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, cyano, nitro, trifluoromethyl.
- 20 (d) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro, trifluoromethyl.


- The compounds of formula (I-a), which are compounds of formula (I) wherein Y is OH
25 and R^6 is hydrogen, can be prepared by reacting a pyrimidine aldehyde of formula (II) with an organo-metal compound ($M-R^5$). The thus obtained compounds of formula (I-a) can be converted to the corresponding compounds of formula (I-b), which are compounds of formula (I) wherein Y is OR^{11} , wherein R^{11} is other than hydrogen. The group R^{11} can be introduced by an ether forming reaction such as an O-alkylation
30 reaction with a reagent W^1-R^{11} , wherein W^1 is a leaving group such as halo, in particular chloro, bromo or iodo, or a sulfate or azide group. M in $M-R^5$ is a metal such as an alkali metal, in particular Li, Na or K, or a magnesium derivative such as a Grignard type of reagent ($M-R^5$ is halo-Mg- R^5). These reactions are typically conducted in a reaction-inert solvent such as an ether (THF, diethylether, dioxan) or a
35 halogenated hydrocarbon (CH_2Cl_2 , $CHCl_3$).



- The compounds of formula (I) wherein Y is OR¹¹ and R⁵ and R⁶ have the same meaning, can be prepared by reacting a pyrimidine carboxylic ester of formula (III) with an organo-metal compound (M-R⁵, which is as specified above), thus obtaining
- 5 compounds of formula (I-c), which are compounds of formula (I) wherein Y is OH and R⁵ and R⁶ have the same meaning. The carboxylic ester in (III), i.e. the group –COOR, can be a C₁₋₆ alkyl ester, e.g. a methyl, ethyl or propyl ester. The compounds of formula (I-c) can be converted to the corresponding compounds of formula (I-d), which are compounds of formula (I) wherein Y is OR¹¹, wherein R¹¹ is other than hydrogen,
- 10 and R⁵ and R⁶ have the same meaning, by an ether forming reaction as described above. These reactions are typically conducted in a reaction-inert solvent such as the solvents mentioned in relation to the preparation of (I-a) and (I-b).




The compounds of formula (I) wherein Y is OH (hereafter represented by (I-e)) can be prepared from a keto derivative (IV) by a reaction with an organo-metal compound (M-R⁶) as described above in the conversion of (III) to (I-c). The keto derivative (IV) is obtained as from a compound (I-a) by an oxidation reaction, in particular with a mild oxidant such as MnO₂. The compounds (I-e) are converted to compounds (I-f), which are compounds of formula (I) wherein Y is OR¹¹ and wherein R¹¹ is other than hydrogen, by an ether forming reaction as described above. Also these reactions typically are conducted in a reaction-inert solvent (e.g. as the solvents mentioned above).




The intermediates of formula (IV), the salts and possible stereoisomers thereof, are novel compounds and constitute a further aspect of the present invention. Salts of the intermediates (IV) in particular are acid-addition salts, more in particular pharmaceutically acceptable acid-addition salts, such as the salts mentioned above in relation to the compounds of formula (I).


The compounds of formula (I) wherein Y is NR¹¹R¹², herein represented by (I-g), can be prepared by reacting a compound of formula (I-c) with a leaving group introducing agent such as POCl₃ or SOCl₂, thus obtaining intermediates (V), which are converted to the corresponding amines with R¹¹R¹²NH. The group W in (IV) represents a leaving group, which in particular is halo, preferably chloro or bromo, or a tosyl, mesyl or similar. Also these reactions are typically conducted in a reaction-inert solvent (such as the solvents mentioned above).



- The intermediates (II) are prepared by first halogenating a starting material of formula (VI), which can be prepared as described in WO-03/016306. Other leaving groups can be introduced by replacing the halo group using suitable reagents. The thus obtained intermediates (VII) are then converted to the corresponding intermediates (II), which have a group $-CHO$ in the 5-position of the pyrimidine moiety by reaction of (VII) with pressurized CO gas in the presence of sodium formate and a suitable catalyst, e.g. dichlorobis(triphenylphosphine)-palladium(II).
- 5



The ester intermediates (III) can be obtained by first reacting intermediates (VII) with pressurized CO gas in the presence of ethanol or a similar lower alkanol and a suitable catalyst, e.g. dichlorobis(triphenylphosphine)-palladium(II).



5

The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a tertiary nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzene-carbo-peroxy acid or halo substituted benzenecarboxy peroxy acid, e.g. 3-chlorobenzenecarboxy peroxy acid, peroxyalkanoic acids, e.g. peroxyacetic acid,

10

alkylhydroperoxides, e.g. tert.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

5

The compounds of formula (I) may further be converted into each other using art-known functional group transformation reactions. Compounds of formula (I) wherein R^2 or R^3 is hydrogen, can be converted into a compounds of formula (I) wherein one or more of R^2 , R^3 , R^7 or R^8 represents halo, by reaction with a suitable halo-introducing agent, e.g. *N*-chlorosuccinimide or *N*-bromosuccinimide, in the presence of a suitable solvent, e.g. acetic acid. Compounds of formula (I) wherein R^1 represents C_{1-6} alkyloxycarbonyl, can be converted into a compound of formula (I) wherein R^1 represents hydrogen, by reaction with a suitable base, such as for example sodium hydroxide or methoxide. Where R^1 is t.butyloxycarbonyl, the corresponding compounds wherein R^1 is hydrogen are prepared by treatment with trifluoroacetic acid.

10

15

20

25

30

Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereo-chemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.

35

An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

The compounds of formula (I) have antiretroviral properties (reverse transcriptase inhibiting properties), in particular against Human Immunodeficiency Virus (HIV), which is the aetiological agent of Acquired Immune Deficiency Syndrome (AIDS) in

humans. The HIV virus preferentially infects human T-4 cells and destroys them or changes their normal function, particularly the coordination of the immune system. As a result, an infected patient has an ever-decreasing number of T-4 cells, which moreover behave abnormally. Hence, the immunological defence system is unable to combat infections and neoplasms and the HIV infected subject usually dies by opportunistic infections such as pneumonia, or by cancers. Other conditions associated with HIV infection include thrombocytopaenia, Kaposi's sarcoma and infection of the central nervous system characterized by progressive demyelination, resulting in dementia and symptoms such as, progressive dysarthria, ataxia and disorientation. HIV infection further has also been associated with peripheral neuropathy, progressive generalized lymphadenopathy (PGL) and AIDS-related complex (ARC).

The present compounds also show activity against (multi) drug resistant HIV strains, in particular (multi) drug resistant HIV-1 strains, more in particular the present compounds show activity against HIV strains, especially HIV-1 strains that have acquired resistance to one or more art-known non-nucleoside reverse transcriptase inhibitors. Art-known non-nucleoside reverse transcriptase inhibitors are those non-nucleoside reverse transcriptase inhibitors other than the present compounds and known to the person skilled in the art, in particular commercial non-nucleoside reverse transcriptase inhibitors. The present compounds also have little or no binding affinity to human α -1 acid glycoprotein; human α -1 acid glycoprotein does not or only weakly affect the anti HIV activity of the present compounds.

Due to their antiretroviral properties, particularly their anti-HIV properties, especially their anti-HIV-1-activity, the compounds of formula (I), the pharmaceutically acceptable addition salts and stereochemically isomeric forms thereof, are useful in the treatment of individuals infected by HIV and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the enzyme reverse transcriptase. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic Central Nervous System diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

The compounds of the present invention may therefore be used as medicines against above-mentioned conditions. Said use as a medicine or method of treatment comprises

the administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1. In particular, the compounds of formula (I) may be used in the manufacture of a medicament for the treatment or the prevention of HIV infections.

5

In further aspect of this invention, there is provided a method of treating warm-blooded animals, including humans, suffering from conditions associated with viral infection, in particular HIV infection, said method comprising the administration to said warm-blooded animals, including humans, an anti-virally effective amount of a compound of formula (I) as specified herein. Furthermore there is provided a method of preventing the development of conditions associated with viral infection, in particular HIV infection, in warm-blooded animals, including humans, said method comprising the administration to said warm-blooded animals, including humans, an anti-virally effective amount of a compound of formula (I) as specified herein.

10

15

The present invention also provides compositions for treating viral infections comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent.

20

The compounds of the present invention may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable

25

30

35

solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are
5 intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the
10 administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

The compounds of the present invention may also be administered via inhalation or
15 insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder. Any system developed for the delivery of solutions, suspensions or dry powders via oral or nasal inhalation or insufflation are suitable for the administration of
20 the present compounds.

To aid solubility of the compounds of formula (I), suitable ingredients, e.g. cyclo-dextrins, may be included in the compositions. Appropriate cyclodextrins are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the
25 hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C_{1-6} alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxy C_{1-6} alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl; carboxy- C_{1-6} alkyl, particularly carboxymethyl or carboxy-ethyl; C_{1-6} alkylcarbonyl, particularly acetyl. Especially noteworthy as complexants and/or solubilizers are
30 β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxypropyl- β -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two
35 cyclo-dextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree

(D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared spectroscopy (IR). Depending on the technique used, slightly different values may be
5 obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

Other suitable compositions for oral or rectal administration comprise particles consisting of a solid dispersion comprising a compound of formula (I) and one or more
10 appropriate pharmaceutically acceptable water-soluble polymers.

The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the compound of formula (I) and the water-soluble polymer, wherein one component is
15 dispersed more or less evenly throughout the other component or components (in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermo-dynamics, such a
20 solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may
25 be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.

The term "a solid dispersion" also comprises dispersions, which are less homogenous
30 throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid dispersion" also relates to a system having domains or small regions wherein amorphous, microcrystalline or crystalline compound of formula (I), or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or
35 less evenly in another phase comprising water-soluble polymer, or compound of formula (I), or a solid solution comprising compound of formula (I) and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by

some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

5 Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

The solution-evaporation process comprises the following steps :

- a) dissolving the compound of formula (I) and the water-soluble polymer in an appropriate solvent, optionally at elevated temperatures;
- 10 b) heating the solution resulting under point a), optionally under vacuum, until the solvent is evaporated. The solution may also be poured onto a large surface so as to form a thin film, and evaporating the solvent therefrom.

15 In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer followed by evaporating the solvent from the resulting droplets at elevated temperatures.

20 The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

- a) mixing a compound of formula (I) and an appropriate water-soluble polymer,
- b) optionally blending additives with the thus obtained mixture,
- c) heating and compounding the thus obtained blend until one obtains a homogenous melt,
- 25 d) forcing the thus obtained melt through one or more nozzles; and
- e) cooling the melt until it solidifies.

30 The terms "melt" and "melting" are not only meant to refer to the transition from a solid state to a liquid state, but also to refer to the transition to a glassy state or a rubbery state, in which it is possible for one component of the mixture to get embedded more or less homogeneously into the other. In particular cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

35 After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved. The solid dispersion product may be milled or ground to particles having a particle size of less than 600 μm , preferably less than 400 μm and most preferably less than 125 μm .

The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

- 5 The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkyl-celluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali
10 metal salts of carboxyalkylcelluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivates, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans, tragacanth, agar-agar, gum arabic, guar gum and xanthan gum, polyacrylic acids and the salts thereof, polymethacrylic acids and the
15 salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

- 20 Also one or more cyclodextrins can be used as water-soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins known in the art, more particularly α , β or γ cyclodextrins or the pharmaceutically
25 acceptable derivatives thereof.

- Substituted cyclodextrins which can be used to prepare the above described particles include polyethers described in U.S. Patent 3,459,731. Further substituted
30 cyclo-dextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C_{1-6} alkyl, hydroxy C_{1-6} alkyl, carboxy- C_{1-6} alkyl or C_{1-6} alkyloxy-carbonyl C_{1-6} alkyl or mixed ethers thereof. In particular such substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy
groups is replaced by C_{1-3} alkyl, hydroxy C_{2-4} alkyl or carboxy C_{1-2} alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxy-
35 methyl or carboxyethyl.

Of particular utility are the β -cyclodextrin ethers, e.g. dimethyl- β -cyclodextrin as described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl β -cyclodextrin and hydroxyethyl β -cyclodextrin, being

examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between β -cyclodextrin and propylene oxide and may have a M.S. value of about 0.125 to 10, e.g. about 0.3 to 3. Another type of substituted cyclodextrins that can be used are the sulfobutylcyclodextrins.

The ratio of the compound of formula (I) over the water soluble polymer may vary widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) over cyclodextrin range from about 1/10 to 10/1. More interesting ratios range from about 1/5 to 5/1.

It may further be convenient to formulate the compounds of formula (I) in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm. Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound. Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.

Yet another way of formulating the compounds of formula (I) involves a pharmaceutical composition whereby the compounds of formula (I) are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration. These beads comprise a central, rounded or spherical core, a coating film of a hydrophilic polymer and a compound of formula (I) and optionally a seal-coating layer. Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including

scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.

Those of skill in the treatment of HIV-infection could determine the effective daily amount from the test results presented here. In general it is contemplated that an effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, more preferably from 0.1 mg/kg to 10 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines and are not intended to limit the scope or use of the invention to any extent.

The compounds of formula (I) can be used alone or in combination with other therapeutic agents, such as anti-virals, antibiotics, immunomodulators or vaccines for the treatment of viral infections. They may also be used alone or in combination with other prophylactic agents for the prevention of viral infections. The present compounds may be used in vaccines and methods for protecting individuals against viral infections over an extended period of time. The compounds may be employed in such vaccines either alone or together with other compounds of this invention or together with other anti-viral agents in a manner consistent with the conventional utilization of reverse transcriptase inhibitors in vaccines. Thus, the present compounds may be combined with pharmaceutically acceptable adjuvants conventionally employed in vaccines and administered in prophylactically effective amounts to protect individuals over an extended period of time against HIV infection.

Also, the combination of one or more additional antiretroviral compounds and a compound of formula (I) can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I), and (b) one or more additional antiretroviral compounds, as a combined preparation for simultaneous,

separate or sequential use in anti-HIV treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. Said other antiretroviral compounds may be any known antiretroviral compounds such as suramine, pentamidine, thymopentin, castanospermine, dextran (dextran sulfate),

5 foscarnet-sodium (trisodium phosphono formate); nucleoside reverse transcriptase inhibitors (NRTIs), e.g. zidovudine (AZT), didanosine (ddI), zalcitabine (ddC), lamivudine (3TC), stavudine (d4T), emtricitabine (FTC), abacavir (ABC), amdoxovir (DAPD), elvucitabine (ACH-126,443), AVX 754 ((-)-dOTC), fozivudine tidoxil (FZT), phosphazide, HDP-990003, KP-1461, MIV-210, racivir (PSI-5004), UC-781 and the

10 like; non-nucleoside reverse transcriptase inhibitors (NNRTIs) such as delavirdine (DLV), efavirenz (EFV), nevirapine (NVP), dapivirine (TMC120), etravirine (TMC125), rilpivirine (TMC278), DPC-082, (+)-Calanolide A, BILR-355, and the like; nucleotide reverse transcriptase inhibitors (NtRTIs), e.g. tenofovir ((R)-PMPA) and tenofovir disoproxil fumarate (TDF), and the like; nucleotide-competing reverse

15 transcriptase inhibitors (NcRTIs), e.g. NcRTI-1 and the like; inhibitors of trans-activating proteins, such as TAT-inhibitors, e.g. RO-5-3335, BI-201, and the like; REV inhibitors; protease inhibitors e.g. ritonavir (RTV), saquinavir (SQV), lopinavir (ABT-378 or LPV), indinavir (IDV), amprenavir (VX-478), TMC126, nelfinavir (AG-1343), atazanavir (BMS 232,632), darunavir (TMC114), fosamprenavir

20 (GW433908 or VX-175), brecanavir (GW-640385, VX-385), P-1946, PL-337, PL-100, tipranavir (PNU-140690), AG-1859, AG-1776, Ro-0334649 and the like; entry inhibitors which comprise fusion inhibitors (e.g. enfuvirtide (T-20)), attachment inhibitors and co-receptor inhibitors, the latter comprise the CCR5 antagonists (e.g. ancriviroc, CCR5mAb004, maraviroc (UK-427,857), PRO-140, TAK-220, TAK-652,

25 vicriviroc (SCH-D, SCH-417,690)) and CXR4 antagonists (e.g. AMD-070, KRH-27315), examples of entry inhibitors are PRO-542, TNX-355, BMS-488,043, BlockAide/CR™, FP 21399, hNM01, nonakine, VGV-1; a maturation inhibitor for example is PA-457; inhibitors of the viral integrase e.g. MK-0518, JTK-303 (GS-9137), BMS-538,158; ribozymes; immunomodulators; monoclonal antibodies;

30 gene therapy; vaccines; siRNAs; antisense RNAs; microbicides; Zinc-finger inhibitors.

The compounds of the present invention may also be administered in combination with immunomodulators (e.g., bropirimine, anti-human alpha interferon antibody, IL-2, methionine enkephalin, interferon alpha, and naltrexone) with antibiotics (e.g.,

35 pentamidine isothiorate) cytokines (e.g. Th2), modulators of cytokines, chemokines or modulators of chemokines, chemokine receptors (e.g. CCR5, CXCR4), modulators chemokine receptors, or hormones (e.g. growth hormone) to ameliorate, combat, or eliminate HIV infection and its symptoms. Such combination therapy in different

formulations, may be administered simultaneously, sequentially or independently of each other. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately.

5

The compounds of the present invention may also be administered in combination with modulators of the metabolism following application of the drug to an individual. These modulators include compounds that interfere with the metabolism at cytochromes, such as cytochrome P450. It is known that several isoenzymes exist of cytochrome P450, one of which is cytochrome P450 3A4. Ritonavir is an example of a modulator of metabolism via cytochrome P450. Such combination therapy in different formulations, may be administered simultaneously, sequentially or independently of each other. Alternatively, such combination may be administered as a single formulation, whereby the active ingredients are released from the formulation simultaneously or separately. Such modulator may be administered at the same or different ratio as the compound of the present invention. Preferably, the weight ratio of such modulator vis-à-vis the compound of the present invention (modulator:compound of the present invention) is 1:1 or lower, more preferable the ratio is 1:3 or lower, suitably the ratio is 1:10 or lower, more suitably the ratio is 1:30 or lower.

20


Although the present invention focuses on the use of the present compounds for preventing or treating HIV infections, the present compounds may also be used as inhibitory agents for other viruses which depend on similar reverse transcriptases for obligatory events in their life cycle.

25


The following examples are intended to illustrate the present invention and not to limit its scope thereto.




Examples












30 Example 1: Preparation of compound 1:









n-BuLi 1,6M (3.04 mmol) was added dropwise to a solution of 2-bromopyrine (3.04 mmol) in THF (5 ml) at -78°C . The mixture was stirred at -78°C for 1h. Then intermediate 1 (0.761 mmol) was added dropwise at -78°C and this mixture was stirred at -78°C for 1h and at room temperature overnight. Water was added and the mixture was extracted with ethyl acetate. The organic layer was dried over magnesium sulfate, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 98/2/0.1 to $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 92/8/0.4 ; Kromasil 5 μm). The pure fractions were collected and the solvent evaporated. Yield : 0.085 g of compound 2 (17 %, melting point : 244°C).

Table 1


Compound No.	R	Phys. Data and stereo-chemistry
1		(E) 239 °C
2		(E) 244 °C
3		(E) 248°C

Compound No.	R	Phys. Data and stereo-chemistry
		30%
4		(E) 250°C 9%
5		(E) 236°C 14%
6		(E) nd°C 14%
7		(E) nd°C 5%
8		(E) 180°C 10%
9		(E) 215°C 25%
10		(E) 160°C 12%
11		(E) 240°C 6%
12		(E) nd°C 14%
13		(E) 200°C 12%
14		(E) 156°C 13%

Compound No.	R	Phys. Data and stereo-chemistry
15		(E) 178°C 13%
16		(E) 190°C 7%
17		(E) 190°C 7%
18		(E) 218°C 11%
19		A2* (E) 236 °C
20		A1* (E) 236 °C

* enantiomeric separation by chiral chromatography

Example 2: Preparation of intermediate 2 :




- Manganese dioxide (0.012 mol) was added to a solution of compound 2 (0.00062 mol) in CH_2Cl_2 (20 ml) and the mixture was stirred at room temperature for 24h then filtered over celite. The celite was washed with $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{THF}$ and the filtrate was evaporated. The residue was purified by column chromatography over silica gel

(eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}$ 99/1 ; SiO_2 10 μm). The pure fractions were collected and the solvent evaporated. Yielding : 0.060 g of intermediate 2 (20%, melting point : $>250^\circ\text{C}$)


Example 3: Preparation of tertiary alcohols :

5 Method A :



n-BuLi 1,6M (3,08mmol) was added dropwise to a solution of thiazole (3.08 mmol) in THF (5 ml) at -78°C . The mixture was stirred at -78°C for 1h. Then intermediate 2 (0.761 mmol) was added dropwise at -78°C then the mixture was stirred at -78°C for 1h and at room temperature overnight. Water and an NH_4Cl 10% solution were added and the mixture was extracted with $\text{CH}_2\text{Cl}_2/\text{THF}$. The organic layer was dried over magnesium sulfate, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 99/1/0.1 to $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 95/5/0.5 ; Kromasil 5 μm). The pure fractions were collected and the solvent evaporated. Yield : 0.044 g of compound 5 (11 %).







Method B :




Commercial magnesium reagent (3,5eq) was added to a solution of intermediate 2 (1 eq) in THF (15ml) at 0°C , the mixture was stirred for 1h at 0°C and then at room

temperature overnight. Water and an NH_4Cl 10% solution were added and the mixture was extracted with $\text{CH}_2\text{Cl}_2/\text{THF}$. The organic layer was dried over magnesium sulfate, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel. The pure fractions were collected and the solvent

5 evaporated, yielding the compounds 21-27.

Compound No.	R	Method	Phys. Data and stereo-chemistry
21		A	(E) 239 °C 11%
22		B	(E) 130°C 28%
23		B	(E) nd°C 14%
24		B	(E) 141°C 43%
25	CH_3	B	(E) 240°C 37%
26		B	(E) 152°C 32%
27		B	(E) 140°C 26%

Example 4: Preparation of thiazole derivatives:

Thionyl chloride (1ml) was added dropwise to a solution of Compound 2 (0,60 mmol) in CH_2Cl_2 (5 ml) at 0°C . The mixture was stirred at 0°C for 3h and then at room temperature overnight. The mixture was evaporated to dryness and the resulting solid (Intermediate 4) filtered off and rinsed with Et_2O . This intermediate was used in the next reactions with no further purification.

10 Method A

Morpholine (1.2 ml) was added dropwise to a solution of Intermediate 4 (0.30 mmol) in THF (15 ml) at room temperature. The mixture was stirred at room temperature overnight. The mixture was poured in water and K_2CO_3 10 % and extracted with $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{THF}$. The organic layer was dried over magnesium sulfate, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 98/2/0.2 to $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 94/6/0.6 ; Kromasil $5\mu\text{m}$). The pure fractions were collected and the solvent evaporated. The residue was crystallized from Et_2O . Yield: 0.047 g of compound 29 (28 %, melting point : 220°C)

20

Method B

A mixture of compound 2 (0.00028 mol, 1eq), 1,1'-Carbonylbisimidazol (0,0031mol, 1.5eq) and methylamine (0,0031mol, 1.5eq) in THF (2 ml) was stirred 18 hours. The mixture was poured in water and K_2CO_3 10 % and extracted with $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{THF}$. The organic layer was dried over magnesium sulfate, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 99/1/0.1 to $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 94/6/0.6 ; Kromasil $5\mu\text{m}$). The pure fractions were collected and the solvent evaporated. The

25







residue was crystallized from $i\text{Pr}_2\text{O}$. Yield : 0.040 g of compound 30 (36 %, melting point : 165°C)


Method C

- 5 1,2-Ethanediol (8.02mmol, 2eq) was added dropwise to a solution of compound 6 (4.01 mmol, 1eq) and triethylamine (8.02mmol, 2eq) in THF (4 ml) at room temperature. The mixture was stirred at room temperature overnight, then heated to reflux 24 hours. The mixture was poured in water and K_2CO_3 10 % and extracted with $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{THF}$. The organic layer was dried over magnesium sulfate, filtered and
- 10 the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 99/1/0.1 to $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ 95/5/0.5 ; Kromasil $5\mu\text{m}$). The pure fractions were collected and the solvent evaporated. The residue was crystallized from $i\text{Pr}_2\text{O}$. Yield : 0.048 g of compound 31 (22 %, melting point : 120°C)

15

The following table list compounds which were prepared according to the procedures described in the above example.

Compound No.	R	Method	Phys. Data and stereo-chemistry
28		A	(E) 215°C 26%
29		A	(E) 220°C 28%
30		B	(E) 165°C 36%
31		C	(E) 120°C 22%
32		C	(E) 210°C 24%
33		C	(E) nd°C 18%

Compound No.	R	Method	Phys. Data and stereo-chemistry
34		C	(E) nd°C 13%

nd means non-determined

Formulation examples

Capsules

- 5 A compound of formula (I) is dissolved in organic solvent such as ethanol, methanol or methylene chloride, preferably, a mixture of ethanol and methylene chloride. Polymers such as polyvinylpyrrolidone copolymer with vinyl acetate (PVP-VA) or hydroxyl-propylmethylcellulose (HPMC), typically 5 mPa.s, are dissolved in organic solvents such as ethanol, methanol methylene chloride. Suitably the polymer is
- 10 dissolved in ethanol. The polymer and compound solutions are mixed and subsequently spray dried. The ratio of compound/polymer is selected from 1/1 to 1/6. Intermediate ranges can be 1/1.5 and 1/3. A suitable ratio can be 1/6. The spray-dried powder, a solid dispersion, is subsequently filled in capsules for administration. The drug load in
- 15 one capsule ranges between 50 and 100 mg depending on the capsule size used.

Film-coated Tablets

Preparation of Tablet Core

- A mixture of 100 g of a compound of formula (I), 570 g lactose and 200 g starch is mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and
- 20 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture is sieved, dried and sieved again. Then there is added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole is mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

- To a solution of 10 g methylcellulose in 75 ml of denaturated ethanol there is added a solution of 5 g of ethylcellulose in 150 ml of dichloromethane. Then there is added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol is molten and dissolved in 75 ml of dichloromethane. The latter solution is added to the
- 30 former and then there is added 2.5 g of magnesium octadecanoate, 5 g of polyvinyl-pyrrolidone and 30 ml of concentrated color suspension and the whole is homogenized. The tablet cores are coated with the thus obtained mixture in a coating apparatus.

Antiviral spectrum:

Because of the increasing emergence of drug resistant HIV strains, the present compounds were tested for their potency against clinically isolated HIV strains

5 harboring several mutations. These mutations are associated with resistance to reverse transcriptase inhibitors and result in viruses that show various degrees of phenotypic cross-resistance to the currently commercially available drugs such as for instance AZT and delavirdine.

10 The antiviral activity of the compound of the present invention has been evaluated in the presence of wild type HIV and HIV mutants bearing mutations at the reverse transcriptase gene. The activity of the compounds is evaluated using a cellular assay which was performed according to the following procedure.

15 The human T-cell line MT4 is engineered with Green Fluorescent Protein (GFP) and an HIV-specific promoter, HIV-1 long terminal repeat (LTR). This cell line is designated MT4 LTR-EGFP, and can be used for the in vitro evaluation of anti-HIV activity of investigational compounds. In HIV-1 infected cells, the Tat protein is produced which upregulates the LTR promotor and finally leads to stimulation of the GFP reporter
20 production, allowing to measure ongoing HIV-infection fluorometrically.

Analogously, MT4 cells are engineered with GFP and the constitutional cytomegalovirus (CMV) promotor. This cell line is designated MT4 CMV-EGFP, and can be used for the in vitro evaluation of cytotoxicity of investigational compounds. In
25 this cell line, GFP levels are comparably to those of infected MT4 LTR-EGFP cells. Cytotoxic investigational compounds reduce GFP levels of mock-infected MT4 CMV-EGFP cells.

Effective concentration values such as 50% effective concentration (EC₅₀) can be
30 determined and are usually expressed in μM . An EC₅₀ value is defined as the concentration of test compound that reduces the fluorescence of HIV-infected cells by 50%. The 50% cytotoxic concentration (CC₅₀ in μM) is defined as the concentration of test compound that reduces fluorescence of the mock-infected cells by 50%. The ratio of CC₅₀ to EC₅₀ is defined as the selectivity index (SI) and is an indication of the
35 selectivity of the anti-HIV activity of the inhibitor. The ultimate monitoring of HIV-1 infection and cytotoxicity is done using a scanning microscope. Image analysis allows very sensitive detection of viral infection. Measurements are done before cell necrosis,

which usually takes place about five days after infection, in particular measurements are performed three days after infection.

The columns IIIB, L100I, etc. in the table list the pEC₅₀ values against various strains

5 IIIB, L100I, etc.

Strain IIIB is wild type HIV strain.


“MDR” refers to a strain that contains mutations L100I, K103N, Y181C, E138G, V179I, L2214F, V278V/I and A327A/V in HIV reverse transcriptase.

Compound No	IIIB	pSi (IIIB)	MDR	L100I + K103N	K103N + Y181C	Y181C
1	8.96	3.99	7.09	8.82	8.49	8.44
2	8.46	3.63	6.09	8.46	8.49	8.07
4	8.75	3.87	6.45	8.46	8.24	8.27
5	8.39	3.78	6.19	8.07	7.73	8.07
7	7.94	3.15	5.79	7.83	7.32	7.67
8	8.41	3.53	6.68	8.37	8.02	7.84
9	7.77	2.94	5.18	6.80	6.12	6.61
10	8.52	3.50	6.84	8.58	8.39	8.42
21	7.74	2.90	5.48	7.63	7.07	7.25
6	8.43	3.36	5.65	7.73	7.02	7.61
11	8.59	3.78	6.08	8.43	8.14	8.22
12	8.91	3.34	6.37	8.86	8.38	8.44
13	8.44	3.84	6.27	7.20	7.41	7.71
14	7.90	2.95	5.20	7.15	7.00	7.12
15	8.17	3.26	6.33	8.11	7.74	7.75
22	7.75	2.97	5.66	7.51	7.03	7.21
28	7.98	3.37	5.80	7.79	7.18	7.70
16	7.60	3.00	5.14	7.20	7.02	7.05
18	7.74	3.14	5.66	7.51	6.97	7.10
25	7.75	2.79	6.40	7.85	7.62	7.74
26	7.00	2.05	5.12	6.33	6.24	6.66

Compound No	IIIB	pSi (IIIB)	MDR	L100I + K103N	K103N + Y181C	Y181C
27	7.70	2.78	6.27	7.84	7.54	7.70
20	8.55	3.63	5.97	8.50	8.37	8.13
19	8.64	3.61	7.15	8.82	8.44	8.40
30	8.43	3.30	5.89	7.41	7.64	7.71
31	8.52	3.57	5.82	8.04	7.70	8.06
32	8.48	3.62	6.07	8.41	7.75	7.93
33	8.62	3.82	6.81	7.62	7.50	7.86
34	7.96	3.36	5.81	6.58	6.59	7.03

ClaimsClaims

1. A compound of formula



5 a stereochemically isomeric form thereof, a pharmaceutically acceptable addition salt thereof, a pharmaceutically acceptable hydrate or solvate thereof, a N-oxide thereof, wherein

each **R¹** independently is hydrogen; aryl; formyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyl;

10 C₁₋₆alkyloxycarbonyl;

R², **R³**, **R⁷** and **R⁸** independently are hydrogen; hydroxy; halo; C₃₋₇cycloalkyl; C₁₋₆alkyloxy; carboxyl; C₁₋₆alkyloxycarbonyl; cyano; nitro; amino; mono- or di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; -C(=O)R¹⁰; C₁₋₆alkyl optionally substituted with halo, cyano or -C(=O)R¹⁰; C₂₋₆alkenyl optionally substituted with halo, cyano or -C(=O)R¹⁰; C₂₋₆alkynyl optionally substituted with halo, cyano or -C(=O)R¹⁰;

R⁴ and **R⁹** independently are hydroxy; halo; C₃₋₇cycloalkyl; C₁₋₆alkyloxy; carboxyl; C₁₋₆alkyloxycarbonyl; formyl; cyano; nitro; amino; mono- or di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl; polyhaloC₁₋₆alkyloxy; -C(=O)R¹⁰; cyano; -S(=O)_rR¹⁰; -NH-S(=O)_rR¹⁰; -NHC(=O)H; -C(=O)NHNH₂; -NHC(=O)R¹⁰; Het; -Y-Het; C₁₋₁₂alkyl optionally substituted with halo, cyano, amino, mono- or di(C₁₋₆alkyl)-amino, -C(=O)-R¹⁰, Het or with C₁₋₆alkyloxy; C₂₋₁₂alkenyl optionally substituted with halo, cyano, amino, mono- or di(C₁₋₆alkyl)amino, -C(=O)-R¹⁰, Het or with C₁₋₆alkyloxy; C₂₋₁₂alkynyl optionally substituted with halo, cyano, amino, mono- and di(C₁₋₆alkyl)amino, -C(=O)-R¹⁰, Het or with C₁₋₆alkyloxy;

R⁵ is C₁₋₆alkyl optionally substituted with Ar or with Het; C₂₋₆alkenyl optionally substituted with Ar or with Het; C₂₋₆alkynyl optionally substituted with Ar or with Het; C₃₋₇cycloalkyl; Ar; Het;

R⁶ is H, C₁₋₆alkyl, Het;

Y is -OR¹¹, -NR¹²R¹³;

each **R**¹⁰ independently is C₁₋₆alkyl, amino, mono- or di(C₁₋₆alkyl)amino or polyhalo-C₁₋₆alkyl;

R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy, phenyl or pyridyl;

R¹² is hydrogen or C₁₋₆alkyl;

R¹³ is hydrogen or C₁₋₆alkyl; or **R**¹² and **R**¹³ taken together with the nitrogen atom to which they are substituted form pyrrolidinyl; piperidinyl; morpholinyl; piperazinyl; piperazinyl optionally substituted with C₁₋₆alkyl or C₁₋₆alkylcarbonyl; imidazolyl;

X is -NR¹-, -O-, -C(=O)-, -CH₂-, -CHOH-, -S-, -S(=O)_r-;

each **Y** independently is -NR¹-, -O-, -C(=O)-, -S-, -S(=O)_r-;

each *r* independently is 1 or 2;


Het is a 5- or 6-membered completely unsaturated ring wherein one, two, three or four ring members are hetero atoms, each independently selected from nitrogen, oxygen and sulfur, and wherein the remaining ring members are carbon atoms; and, where possible, any nitrogen ring member may optionally be substituted with C₁₋₆alkyl; which 5- or 6-membered ring may optionally be annelated with a benzene, thiophene or pyridine ring; and wherein any ring carbon atom, including any carbon of an optionally annelated benzene, thiophene or pyridine ring, may, each independently, optionally be substituted with a substituent selected from halo, hydroxy, mercapto, cyano, C₁₋₆alkyl, hydroxyC₁₋₄alkyl, carboxyC₁₋₄alkyl, C₁₋₄alkyloxyC₁₋₄alkyl, C₁₋₄alkyloxycarbonylC₁₋₄alkyl, cyanoC₁₋₄alkyl, mono- and di(C₁₋₄alkyl)aminoC₁₋₄alkyl, Het¹C₁₋₄alkyl, arylC₁₋₄alkyl, polyhaloC₁₋₄alkyl, C₃₋₇cycloalkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, aryl-C₂₋₄alkenyl, C₁₋₄alkyloxy, -OCONH₂, polyhaloC₁₋₄alkyloxy, aryloxy, amino, mono- and di-C₁₋₄alkylamino, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, 4-C₁₋₆alkylpiperazinyl, C₁₋₄alkylcarbonylamino, formyl, C₁₋₄alkylcarbonyl, C₁₋₄alkyloxy-carbonyl, aminocarbonyl, mono- and diC₁₋₄alkylaminocarbonyl, aryl, Het¹;

Het¹ is pyridyl, thienyl, furanyl, oxazolyl, isoxazolyl, imidazolyl, pyrazolyl, thiazolyl, thiadiazolyl, oxadiazolyl, quinolinyl, benzothienyl, benzofuranyl; which each may be optionally substituted with one or two C₁₋₄alkyl radicals;

each **aryl** independently is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono and

di(C₁₋₆alkyl)-aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, phenylC₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, aminosulfonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, aminocarbonyl, phenyl, Het and -Y-Het.

- 5 2. A compound according to claim 1 wherein the compound is of formula



wherein X, Y, R¹, R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are as defined in claim 1.

3. A compound according to claims 1 or 2 wherein R¹ is hydrogen.
- 10 4. A compound according to claims 1 or 3 wherein R², R³, R⁷ and R⁸ independently are hydrogen; hydroxy; halo; C₁₋₆alkyl; C₁₋₆alkyloxy; cyano; amino; mono- or di(C₁₋₆alkyl)amino; polyhaloC₁₋₆alkyl.
- 15 5. A compound according to claims 1 to 4 wherein R⁴ and R⁹ independently are cyano; C₁₋₆alkyl substituted with cyano; C₂₋₆alkenyl substituted with cyano.
6. A compound according to any one of claims 1 to 5 wherein R⁵ is C₁₋₆alkyl substituted with Ar or with Het, C₂₋₆alkenyl, C₂₋₆alkynyl optionally substituted with
- 20 Het, C₃₋₇cycloalkyl, Ar, Het;
R⁶ is H, C₁₋₆alkyl, Het;
Y is OR¹¹, NR¹²R¹³;
R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy, phenyl or pyridyl;
- 25 R¹² is hydrogen or C₁₋₆alkyl;
R¹³ is hydrogen or C₁₋₆alkyl; or R¹² and R¹³ taken together with the nitrogen atom to which they are substituted form imidazolyl.
7. A compound according to any one of claims 1 to 5 wherein R⁵ is C₂₋₆alkenyl,
- 30 C₂₋₆alkynyl substituted with Het, Ar, Het;

R⁶ is H, C₁₋₆alkyl, Het;

Y is OR¹¹, NR¹²R¹³;

R¹¹ is hydrogen or C₁₋₆alkyl optionally substituted with hydroxy, C₁₋₆alkyloxy;

R¹² is hydrogen or C₁₋₆alkyl;

5 R¹³ is hydrogen.

8. A compound according to any one of claims 1 to 7 wherein each Het independently is pyridyl, thienyl, thiazolyl, furanyl, each of which may be optionally substituted with a radical selected from C₁₋₆alkyl.

10

9. A compound according to any one of claims 1 to 8 wherein each aryl independently may be phenyl optionally substituted with C₁₋₆alkyl. amino, mono- or diC₁₋₆alkyl-amino, C₁₋₆alkyloxy, aminosulfonyl, Het, the latter more in particular being thiadiazolyl

15

10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredient a therapeutically effective amount of a compound as claimed in any one of claims 1 to 9.

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/053113

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D417/06 C07D401/06 C07D239/48 C07D403/06 C07D513/04
C07D417/14 A61K31/506 A61K31/5377 A61P31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 2003/171374 A1 (KUKLA MICHAEL JOSEPH ET AL) 11 September 2003 (2003-09-11) cited in the application abstract table 1; compounds 2,3,14,27 -----	1-10
Y	WO 2004/065378 A (WARNER-LAMBERT COMPANY LLC; BIWERSI, CATHLIN, MARIE; MCNAMARA, DENNIS,) 5 August 2004 (2004-08-05) * claim 11: page 76, line 15; page 78, line 7 * claim 14 ----- -/--	1-10

☒ Further documents are listed in the continuation of Box C.☒ See patent family annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

6 July 2007

Date of mailing of the international search report

17/07/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

MATES VALDIVIELSO, J

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2007/053113

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>LUDOVICI D W ET AL: "Evolution of anti-HIV drug candidates. Part 3: Diarylpyrimidine (DAPY) analogues" BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, OXFORD, GB, vol. 11, 2001, pages 2235-2239, XP002329859 ISSN: 0960-894X the whole document -----</p>	1,10

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2007/053113

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2003171374 A1	11-09-2003	AR 028423 A1	07-05-2003
		AU 783981 B2	12-01-2006
		AU 5633301 A	20-11-2001
		CA 2406562 A1	15-11-2001
		WO 0185700 A2	15-11-2001
		EP 1282606 A2	12-02-2003
		JP 2003535828 T	02-12-2003
WO 2004065378 A	05-08-2004	BR 0406809 A	27-12-2005
		CA 2512646 A1	05-08-2004
		EP 1590341 A1	02-11-2005
		JP 2006516561 T	06-07-2006
		MX PA05007503 A	21-09-2005