



(22) Date de dépôt/Filing Date: 1994/05/26

(41) Mise à la disp. pub./Open to Public Insp.: 1995/01/05

(45) Date de délivrance/Issue Date: 2005/08/02

(62) Demande originale/Original Application: 2 166 059

(30) Priorités/Priorities: 1993/06/25 (08/082,693) US;
1993/10/27 (08/144,084) US; 1994/05/11 (08/237,466) US

(51) Cl.Int.⁷/Int.Cl.⁷ C07D 211/34, A61K 31/445

(72) Inventeurs/Inventors:

KRAUSS, RICHARD CARL, US;
STROM, ROBERT MICHAEL, US;
SCORTICHINI, CAREY LEE, US;
KRUPER, WILLIAM JOHN, US;
WOLF, RICHARD APPLEBY, US;
CARR, ALBERT ANTHONY, US;
RUDISILL, DUANE ERIC, US;

...


(73) Propriétaire/Owner:

MERRELL PHARMACEUTICALS INC., US

(74) Agent: OSLER, HOSKIN & HARCOURT LLP

(54) Titre : NOUVEAUX INTERMEDIAIRES POUR LA PREPARATION DE DERIVES DE 4-DIPHENYLMETHYL/DIPHENYLMETHOXY PIPERIDINE, ANTIHISTAMINIQUES

(54) Title: NOVEL INTERMEDIATES FOR THE PREPARATION OF ANTIHISTAMINIC 4-DIPHENYLMETHYL/DIPHENYLMETHOXY PIPERIDINE DERIVATIVES




(57) Abrégé/Abstract:

The present invention is related to novel intermediates and processes which are useful in the preparation of certain antihistaminic piperidine derivatives of formula (I): (see above formula) wherein W represents -C(=O)- or -CH(OH)-; R₁ represents hydrogen or hydroxy; R₂ represents hydrogen; R₁ and R₂ taken together form a second bond between the carbon atoms bearing R₁ and R₂; n is an integer of from 1 to 5; m is an integer 0 or 1; R₃ is -COOH or -COOalkyl wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched; each of A is hydrogen or hydroxy; and pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R₁ and R₂ are taken together to form a second bond between the carbon atoms bearing R₁ and R₂ or where R₁ represented hydroxy, m is an integer 0.

(72) Inventeurs(suite)/Inventors(continued): PANZONE, GIANBATTISTA, IT; HAY, DAVID ALLEN, US;
WU, WEISHI WILSON, US

ABSTRACT

The present invention is related to novel intermediates and processes which are useful in the preparation of certain antihistaminic piperidine derivatives of formula (I):



wherein

W represents $-C(=O)-$ or $-CH(OH)-$;

R_1 represents hydrogen or hydroxy;

R_2 represents hydrogen;

R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;

n is an integer of from 1 to 5;

m is an integer 0 or 1;

R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

each of A is hydrogen or hydroxy; and

pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.

-1-

5

1 NOVEL INTERMEDIATES FOR THE PREPARATION OF ANTIHISTAMINIC 4-
DIPHENYLMETHYL/DIPHENYLMETHOXY PIPERIDINE DERIVATIVES

BACKGROUND OF THE INVENTION


15

20 The present invention is related to novel intermediates
which are useful in the preparation of certain piperidine
derivatives which are useful as antihistamines, antiallergy
agents and bronchodilators [United States Patent No.
4,254,129, March 3, 1981, United States Patent No.
25 4,254,130, March 3, 1981, United States Patent No.
4,285,958, April 25, 1981 and United States Patent No.
4,550,116, Oct. 29, 1985].

These antihistaminic piperidine derivatives can be
30 described by the following formula:

35

- 2 -



wherein

- W represents $-\text{C}(=\text{O})-$ or $-\text{CH}(\text{OH})-$;
- 15 R_1 represents hydrogen or hydroxy;
- R_2 represents hydrogen;
- R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R_3 is $-\text{COOH}$ or $-\text{COOalkyl}$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;
- each of A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual
- 25 optical isomers thereof,
- with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.


30

SUMMARY OF THE INVENTION

The present invention provides novel intermediates useful for the preparation of certain antihistaminic

35 piperidine derivatives of formula (I)

- 3 -



wherein

- W represents $-\text{C}(=\text{O})-$ or $-\text{CH}(\text{OH})-$;
- 15 R_1 represents hydrogen or hydroxy;
- R_2 represents hydrogen; or
- R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R_3 is $-\text{COOH}$ or $-\text{COOalkyl}$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;
- each of A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual
- 25 optical isomers thereof,
- with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.


30

These novel intermediates are described by the following formulas:

35

- 4 -

5 (II)




wherein

- A is a hydrogen or hydroxy; and
- 10 R₅ is H, -CH₂OD wherein D is hydrogen, acetate or benzoate, -CHO, Br, Cl, I, CN, -COOH, -COOalkyl or -CONR₆R₇ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched and R₆ and R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy
- 15 or R₆ and R₇ taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R₆ and R₇ cannot both be represented by C₁-C₆alkoxy.

20

(III)




25 wherein

- A is a hydrogen or hydroxy; and
- R₅ is H, Br, Cl, I, CN, -COOH, -COOalkyl
- 30 or -CONR₆R₇ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched and R₆ and R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy or R₆ and R₇ taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R₆ and R₇ cannot both be represented by C₁-C₆alkoxy.

35

- 5 -

5 (IV)




wherein

A is a hydrogen or hydroxy; and

- 10 R₅ is H, Br, Cl, I, CN, -COOH, -COOalkyl or
 -CONR₆R₇ wherein the alkyl moiety has from 1 to 6
 carbon atoms and is straight or branched and R₆ and
 R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy
 15 or R₆ and R₇ taken together with the nitrogen atom
 form a pyrrolidine, piperidine or morpholine, with
 the proviso that R₆ and R₇ cannot both be
 represented by C₁-C₆alkoxy.

20

(V)



wherein


- 25 Hal is Cl, Br or I;
 n is an integer of from 1 to 5;
 A is a hydrogen or hydroxy; and
 R₅ is H, CH₂OD wherein D is hydrogen, acetate or
 benzoate, CHO, Br, Cl, I, CN, -COOH or -CONR₆R₇
 30 wherein R₆ and R₇ are each independently H, C₁-
 C₆alkyl, C₁-C₆alkoxy or R₆ and R₇ taken together with
 the nitrogen atom form a pyrrolidine, piperidine or
 morpholine, with the proviso that R₆ and R₇ cannot
 both be represented by C₁-C₆alkoxy.

35

- 6 -

5

(VI)



wherein

10

Hal is Cl, Br or I;

n is an integer of from 1 to 5;

A is a hydrogen or hydroxy; and

R₅ is H, Br, Cl, I, CN, -COOH, -COOalkyl or


15

-CONR₆R₇ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched and R₆ and R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy or R₆ and R₇ taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R₆ and R₇ cannot both be

20

represented by C₁-C₆alkoxy.

(VII)



25

wherein

Hal is Cl, Br or I;

n is an integer of from 1 to 5;

30


A is a hydrogen or hydroxy;

R₅ is H, Br, Cl, I, CN, -COOH, -COOalkyl or

35


-CONR₆R₇ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched and R₆ and R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy or R₆ and R₇ taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R₆ and R₇ cannot both be represented by C₁-C₆alkoxy.

- 7 -




wherein

- 10 Hal is Cl, Br or I;
 n is an integer of from 1 to 5; and
 A is a hydrogen or hydroxy.




wherein A is a hydrogen or hydroxy.



wherein

- 25 Hal is Cl, Br or I;
 n is an integer of from 1 to 5;
 A is a hydrogen or hydroxy; and
 R₅ is H, CH₂OD wherein D is hydrogen, acetate or
 benzoate, CHO, Br, Cl, I, CN, -COOH, -COOalkyl or
 30 -CONR₆R₇ wherein the alkyl moiety has from 1 to 6
 carbon atoms and is straight or branched and R₆ and
 R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy
 or R₆ and R₇ taken together with the nitrogen atom
 form a pyrrolidine, piperidine or morpholine, with
 35 the proviso that R₆ and R₇ cannot both be
 represented by C₁-C₆alkoxy; and
 individual optical isomers thereof.

- 8 -




wherein

- W represents -C(=O)- or -CH(OH)-;
- 15 R₁ represents hydrogen or hydroxy;
- R₂ represents hydrogen; or
- R₁ and R₂ taken together form a second bond between the carbon atoms bearing R₁ and R₂;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R₅ is H, Br, Cl, I, CN or -CONR₆R₇ wherein R₆ and R₇ are each independently H, C₁-C₆alkyl, C₁-C₆alkoxy or R₆ and R₇ taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R₆ and R₇ cannot both be represented by C₁-C₆alkoxy;
- 25 A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R₁ and R₂ are taken together to form a second bond between the carbon atoms bearing R₁ and R₂ or where R₁ represented hydroxy, m is an integer 0.
- 30

35 In addition, the present invention provides novel processes for preparing the antihistaminic piperidine derivatives of formula


- 9 -



wherein

- W represents $-C(=O)-$ or $-CH(OH)-$;
- 15 R_1 represents hydrogen or hydroxy;
- R_2 represents hydrogen; or
- R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;
- each of A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual optical
- 25 isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0, comprising the steps of:


- 30 (a) reacting a cumene compound of the formula



35

wherein A is as defined above with a ω -halo compound of the formula

- 10 -



5 wherein B is halo or hydroxy, Hal represents Cl, Br or I and n is as defined above, in the presence of a suitable Lewis acid to produce a ω -halo cumylketone compound;


10 (b) reacting the ω -halo cumylketone compound with a suitable halogenating agent to give a ω -halo-halocumylketone compound;

15 (c) reacting the ω -halo-halocumylketone compound with a suitable cyanating agent to give a ω -halo-cyanocumylketone compound;

20 (d) reacting the ω -halo-cyanocumylketone compound with an appropriate straight or branched $\text{C}_1\text{-C}_6$ alcohol in the presence of a suitable anhydrous acid to give a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid imidate compound;

25 (e) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid imidate compound with water to give a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound;

30 (f) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound with a piperidine compound of the formula



35

- 11 -

wherein R_1 , R_2 and m are as defined above in the presence of a suitable non-nucleophilic base to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein
 5 R_3 is COOalkyl and W is $-C(=O)-$;

(g) optionally hydrolyzing the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOalkyl and W is $-C(=O)-$ to produce a ω' -piperidine- α' -
 10 hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$;

(h) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is
 15 COOalkyl and W is $-C(=O)-$ or the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$ with a suitable reducing agent to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of
 20 formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$; and

(i) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-$
 25 $COOH$ and W is $-CH(OH)-$ or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$ with an appropriate straight or branched C_1-C_6 alcohol in the presence of a suitable acid to
 30 produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$; and

(j) optionally reacting the ω' -piperidine- α' -keto- α,α -
 35 dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$, the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$, the ω' -piperidine- α' -hydroxy- α,α -

- 12 -


dimethylphenyl derivative of formula (I) wherein R_3 is $-\text{COOH}$ and W is $-\text{CH}(\text{OH})-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -
 5 COOalkyl and W is $-\text{CH}(\text{OH})-$ with an appropriate deprotecting reagent,

with the proviso that each of the hydroxy groups present in
 10 the compounds described in steps a-i are optionally protected or unprotected.

In addition, the present invention provides novel processes for preparing the antihistaminic piperidine derivatives of formula

15

20



25

30

35

- 13 -

wherein

- W represents $-C(=O)-$ or $-CH(OH)-$;
 R₁ represents hydrogen or hydroxy;
 5 R₂ represents hydrogen; or
 R₁ and R₂ taken together form a second bond between the
 carbon atoms bearing R₁ and R₂;
 n is an integer of from 1 to 5;
 m is an integer 0 or 1;
 10 R₃ is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has
 from 1 to 6 carbon atoms and is straight or branched;
 each of A is hydrogen or hydroxy; and
 pharmaceutically acceptable salts and individual optical
 isomers thereof, with the proviso that where R₁ and R₂
 15 are taken together to form a second bond between the
 carbon atoms bearing R₁ and R₂ or where R₁ represented
 hydroxy, m is an integer 0, comprising the steps of:

(a) reacting a ω -halo-halocumylketone compound with
 20 carbon dioxide under electrochemical reduction conditions to
 give a ω' -halo- α' -keto- α,α -dimethylphenylacetic compound;


(b) reacting the ω' -halo- α' -keto- α,α -
 dimethylphenylacetic compound compound with an appropriate
 25 straight or branched C₁-C₆ alcohol in the presence of a
 suitable anhydrous acid to give a ω' -halo- α' -keto- α,α -
 dimethylphenylacetic acid ester compound;

(c) reacting the ω' -halo- α' -keto- α,α -
 30 dimethylphenylacetic acid ester compound with a piperidine
 compound of the formula
 wherein R₁, R₂ and m are as defined above in the presence of
 a suitable non-nucleophilic base to produce a ω' -piperidine-
 α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein
 35 R₃ is COOalkyl and W = $-C(=O)-$;

(d) optionally hydrolyzing the ω' -piperidine- α' -keto-
 α,α -dimethylphenyl derivative of formula (I) wherein R₃ is

- 14 -

5



10

COOalkyl and W is $-C(=O)-$ to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$;

15

(e) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOalkyl and W is $-C(=O)-$ or the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$ with a suitable reducing agent to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$; and

20

25

(f) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$ with an appropriate straight or branched C_1 - C_6 alcohol in the presence of a suitable acid to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$; and

30

35


(g) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$, the ω' -piperidine- α' -keto- α,α -

- 15 -

dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -C(=O)-, the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is -CH(OH)- or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -CH(OH)- with an appropriate deprotecting reagent,

10 with the proviso that each of the hydroxy groups present in the compounds described in steps a-f are optionally protected or unprotected.

In addition, the present invention provides novel
15 processes for preparing the antihistaminic piperidine derivatives of formula



- 16 -

wherein

W represents $-C(=O)-$ or $-CH(OH)-$;

R_1 represents hydrogen or hydroxy;

5 R_2 represents hydrogen; or

R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;

n is an integer 3;


m is an integer 0 or 1;

10 R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched; each of A is hydrogen or hydroxy; and

pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0, comprising the steps of:


(a) reacting a cumyl compound of the formula

20



25 wherein A is as defined above with an appropriate cyclopropyl compound of the structure

30



wherein B is halo or hydroxy, in the presence of a suitable Lewis acid to produce a cyclopropyl cumylketone compound;

35 (b) reacting the cyclopropyl cumylketone compound with a suitable halogenating agent to give a cyclopropyl halocumylketone compound;


- 17 -

(c) reacting the cyclopropyl halocumylketone compound with carbon dioxide under electrochemical reduction conditions to give a cyclopropylketo- α,α -dimethylphenylacetic acid compound;

(d) reacting the cyclopropylketo- α,α -dimethylphenylacetic with an appropriate straight or branched C_1 - C_6 alcohol in the presence of a suitable anhydrous acid to give a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound;

(e) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound with a piperidine compound of the formula

20



wherein R_1 , R_2 and m are as defined above in the presence of a suitable non-nucleophilic base to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOalkyl and $W = -C(=O)-$;

(f) optionally hydrolyzing the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOalkyl and W is $-C(=O)-$ to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$;

35

(g) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOalkyl and W is $-C(=O)-$ or the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH

- 18 -

and W is -C(=O)- with a suitable reducing agent to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is -CH(OH)- or the ω' -
 5 piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -CH(OH)-; and

(h) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -
 10 COOH and W is -CH(OH)- or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is -C(=O)- with an appropriate straight or branched C_1 - C_6 alcohol in the presence of a suitable acid to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl
 15 derivative of formula (I) wherein R_3 is -COOalkyl and W is -CH(OH)- or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -C(=O)-; and

(i) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is -C(=O)-, the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -C(=O)-, the ω' -piperidine- α' -hydroxy- α,α -
 25 dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is -CH(OH)- or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOalkyl and W is -CH(OH)- with an appropriate deprotecting reagent,


30

with the proviso that each of the hydroxy groups present in the compounds described in steps a-h are optionally protected or unprotected.

35

Another embodiment of the present invention involves a process for preparing the piperidine derivatives of formula

- 19 -



wherein

- W represents $-C(=O)-$ or $-CH(OH)-$;
- 15 R_1 represents hydrogen or hydroxy;
- R_2 represents hydrogen; or
- R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;
- each of A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual optical
- 25 isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0, comprising the steps of:


- 30 (a) reacting a α,α -dimethylphenylacetic acid amide compound of the formula



wherein A is as defined above and R_6 and R_7 are each independently H, C_1-C_6alkyl , $C_1-C_6alkoxy$ or R_6 and R_7 taken


- 20 -

together with the nitrogen atom for a pyrrolidine, piperidine or morpholine, with the proviso that R_6 and R_7 cannot both be represented by C_1-C_6 alkoxy with a ω -halo compound of the formula



10 wherein B is halo or hydroxy, Hal represents Cl, Br or I and n is as defined above, in the presence of a suitable Lewis acid to produce a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound;

15 (b) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound with a piperidine compound of the formula



25 wherein R_1 and R_2 are as defined above in the presence of a suitable non-nucleophilic base to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (XI) wherein R_5 is $-\text{CONR}_6\text{R}_7$ wherein R_6 and R_7 are as defined
30 above;

(c) optionally hydrolyzing the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (XI) wherein R_5 is $-\text{CONR}_6\text{R}_7$ wherein R_6 and R_7 are as defined above to produce a
35 ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-\text{C}(=\text{O})-$;

(d) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$ with a suitable reducing agent to produce a
 5 ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$; and

(e) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -
 10 COOH and W is $-CH(OH)-$ or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$ with an appropriate straight or branched C_1-C_6 alcohol in the presence of a suitable acid to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl
 15 derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$; and


20 (f) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$, the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$, the ω' -piperidine- α' -hydroxy- α,α -
 25 dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ with an appropriate deprotecting reagent,

30

with the proviso that each of the hydroxy groups present in the compounds described in steps a-e are optionally protected or unprotected.

35 Another embodiment of the present invention involves a process for preparing the piperidine derivatives of formula


- 22 -



wherein

- W represents -C(=O)- or -CH(OH)-;
- 15 R₁ represents hydrogen or hydroxy;
- R₂ represents hydrogen; or
- R₁ and R₂ taken together form a second bond between the carbon atoms bearing R₁ and R₂;
- n is an integer of from 1 to 5;
- 20 m is an integer 0 or 1;
- R₃ is -COOH or -COOalkyl wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;
- each of A is hydrogen or hydroxy; and
- pharmaceutically acceptable salts and individual optical
- 25 isomers thereof, with the proviso that where R₁ and R₂ are taken together to form a second bond between the carbon atoms bearing R₁ and R₂ or where R₁ represented hydroxy, m is an integer 0, comprising the steps of:


- 30 (a) reacting a toluene compound of the formula



35

wherein A is as defined above with a ω-halo compound of the formula


- 23 -



- 5 wherein B is halo or hydroxy, Hal represents Cl, Br or I and n is as defined above, in the presence of a suitable Lewis acid to produce a ω -halo-tolylketone compound;
- 10 (b) reacting the ω -halo-tolylketone compound with a suitable base to give a cyclopropyl-tolylketone compound;
- 15 (c) reacting the cyclopropyl-tolylketone compound with a suitable halogenating agent to give a cyclopropyl-halotolylketone compound;
- (d) reacting the cyclopropyl-halotolylketone compound with a suitable cyanating agent to give a cyclopropyl cyanotolylketone compound;
- 20 (e) reacting the cyclopropyl cyanotolylketone compound with a suitable methylating agent to give a cyclopropyl cyanocumylketone compound;
- 25 (f) reacting the cyclopropyl cyanocumylketone compound with a suitable base to give a cyclopropylketo- α,α -dimethylphenylacetic acid amide;
- 30 (g) reacting the cyclopropylketo- α,α -dimethylphenylacetic acid amide with an appropriate straight or branched $\text{C}_1\text{-C}_6$ alcohol in the presence of a suitable anhydrous acid to give a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound;
- 35 (h) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound with a piperidine compound of the formula

- 24 -

5



10 wherein R_1 , R_2 and m are as defined above in the presence of a suitable non-nucleophilic base to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative;

15 (i) optionally hydrolyzing the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$;

20 (j) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-C(=O)-$ with a suitable reducing agent to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$; and

25 (k) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$ with an appropriate straight or
30 branched C_1-C_6 alcohol in the presence of a suitable acid to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (II) wherein R_3 is $-COOalkyl$ and W is
35 $-C(=O)-$; and


(l) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (II) wherein R_3 is -

- 25 -

COOH and W is $-C(=O)-$, the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (II) wherein R_3 is -COOalkyl and W is $-C(=O)-$, the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is -COOH and W is $-CH(OH)-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl of formula (I) wherein R_3 is -COOalkyl and W is $-CH(OH)-$ with an appropriate deprotecting reagent,

10 with the proviso that each of the hydroxy groups present in the compounds described in steps a-k are optionally protected or unprotected.

Another embodiment of the present invention involves a
15 process for preparing the piperidine derivatives of formula



- 26 -

wherein

W represents $-C(=O)-$ or $-\bar{C}H(OH)-$;

R_1 represents hydrogen or hydroxy;

5 R_2 represents hydrogen; or

R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;


n is an integer of from 1 to 5;

m is an integer 0 or 1;

10 R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched; each of A is hydrogen or hydroxy; and


pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0, comprising the steps of:

(a) reacting a phenylacetic acid ester compound of the
20 formula



25

wherein A is as defined above with a ω -halo compound of the formula



30

wherein B is halo or hydroxy, Hal represents Cl, Br or I and n is as defined above, in the presence of a suitable Lewis acid to produce a ω' -halo- α' -keto-phenylacetic acid ester compound;

35

(b) reacting the ω' -halo- α' -keto-phenylacetic acid ester compound with a suitable methylating agent in the

- 27 -

presence of a suitable base to give a cyclopropylketo- α,α -dimethylphenylacetic acid ester;

5 (c) purifying the cyclopropylketo- α,α -dimethylphenylacetic acid ester by distillation and/or recrystallization;

(d) reacting the cyclopropylketo- α,α -
10 dimethylphenylacetic acid ester with an appropriate straight or branched C_1 - C_6 alcohol in the presence of a suitable anhydrous acid to give a ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound;

15 (e) reacting the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound with a piperidine compound of the formula



wherein R_1 , R_2 and m are as defined above in the presence of a suitable non-nucleophilic base to produce a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-\text{COOalkyl}$ and W is $-\text{C}(=\text{O})-$;

30

(f) optionally hydrolyzing the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-\text{COOalkyl}$ and W is $-\text{C}(=\text{O})-$ to produce a ω' -piperidine- α' -
35 keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH and W is $-\text{C}(=\text{O})-$;

(g) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is COOH

- 28 -

and W is $-C(=O)-$ with a suitable reducing agent to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$; and

5

(h) optionally reacting the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$ with an appropriate straight or branched C_1-C_6 alcohol in the presence of a suitable acid to produce a ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ or a ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$; and

(i) optionally reacting the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-C(=O)-$, the ω' -piperidine- α' -keto- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOalkyl$ and W is $-C(=O)-$, the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl derivative of formula (I) wherein R_3 is $-COOH$ and W is $-CH(OH)-$ or the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl of formula (I) wherein R_3 is $-COOalkyl$ and W is $-CH(OH)-$ with an appropriate deprotecting reagent,

with the proviso that each of the hydroxy groups present in the compounds described in steps a-h are optionally protected or unprotected.

30

- 29 -

As used herein, the term "C₁-C₆alkyl" or "alkyl" refers to a straight or branched alkyl group having from 1 to 6 carbon atoms and as referred to herein are methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, tert-butyl, n-pentyl, neopentyl and n-hexyl. The term "C₁-C₆alkoxy" refers to a straight or branched alkoxy group having from 1 to 6 carbon atoms and as referred to herein are methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, sec-butoxy, tert-butoxy, n-pentoxy, neopentoxy and n-hexoxy. The term "Hal" or "halo" refers to a halogen group and includes Cl, Br or I.

15

- 30 -


The piperidine derivatives of the formula (IX) can form pharmaceutically acceptable salts. Pharmaceutically acceptable acid addition salts of the compounds of this invention are those of any suitable inorganic or organic acid. Suitable inorganic acids are, for example, hydrochloric, hydrobromic, sulfuric, and phosphoric acids. Suitable organic acids include carboxylic acids, such as, acetic, propionic, glycolic, lactic, pyruvic, malonic, succinic, fumaric, malic, tartaric, citric, cyclamic, ascorbic, maleic, hydroxymaleic, and dihydroxymaleic, benzoic, phenylacetic, 4-aminobenzoic, 4-hydroxybenzoic, anthranillic, cinnamic, salicylic, 4-aminosalicylic, 2-phenoxybenzoic, 2-acetoxybenzoic, and mandelic acid, sulfonic acids, such as, methanesulfonic, ethanesulfonic and β -hydroxyethanesulfonic acid. Non-toxic salts of the compounds of the above-identified formula formed with inorganic or organic bases are also included within the scope of this invention and include, for example, those of alkali metals, such as, sodium, potassium and lithium, alkaline earth metals, for example, calcium and magnesium, light metals of group IIIA, for example, aluminum, organic amines, such as, primary, secondary or tertiary amines, for example, cyclohexylamine, ethylamine, pyridine, methylaminoethanol and piperazine. The salts are prepared by conventional means as, for example, by treating a piperidine derivative of formula (I) with an appropriate acid or base.

30

35

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is hydrogen may be prepared as described in Scheme A. In Scheme A, all substituents are as previously defined unless otherwise indicated.

Scheme A



Scheme A provides various general synthetic procedures for preparing the novel intermediates of formula (II), formula (III) and formula (IV) wherein R₅ is hydrogen.

- 32 -

In step a, the appropriate toluene derivative of structure (1) is methylated to give the corresponding ethylbenzene derivative of structure (2).

For example, the appropriate toluene derivative of structure (1) is reacted with a slight molar excess of an appropriate methylating agent, such as iodomethane, chloromethane or bromomethane in the presence of a suitable non-nucleophilic base, such as potassium t-butoxide or sodium hydride. The reaction is typically conducted in a suitable organic solvent, such as diglyme, tert-butyl methyl ether or methylene chloride, for a period of time ranging from 30 minutes to 24 hours and at a temperature range of from -78°C to room temperature. The corresponding ethylbenzene derivative of structure (2) is recovered from the reaction zone by extractive methods as is known in the art and may be purified by distillation.

20

In step b, the appropriate ethylbenzene derivative of structure (2) is methylated to give the corresponding cumene derivative of structure (3) as described previously in step a, but using at least 2 molar equivalents of methylating agent.

In step c, the appropriate toluene derivative of structure (1) is dimethylated to give the corresponding cumene derivative of structure (3) as described previously in step a but using at least 2 molar equivalents of methylating agent.

In step d, the appropriate toluene derivative of structure (1) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo tolylketone compound of structure (4).

For example, the appropriate ω -halo tolylketone compound of structure (4) may be prepared by reacting an appropriate toluene derivative of structure (1) with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined, which are known in the art or are prepared by procedures well known in the art, under the general conditions of a Friedel-Crafts acylation using a suitable Lewis acid. The reaction is carried out in a solvent, such as carbon disulfide, 1,2-dichloroethane, n-hexane, acetonitrile, 1-nitropropane, nitromethane, diethyl ether and carbon tetrachloride, methylene chloride, tetrachloroethane or nitrobenzene with methylene chloride being the preferred solvent. The reaction time varies from about 1/2 hour to 25 hours, preferably 10 to 16 hours and the reaction temperature varies from about 0°C to 25°C. The corresponding ω -halo tolylketone compound of structure (4) is recovered from the reaction zone by an aqueous quench followed by extraction as is known in the art. The ω -halo tolylketone compound of structure (4) may be purified by procedures well known in the art, such as crystallization and/or distillation.

25

Alternatively, the appropriate toluene derivative of structure (1) may be acylated with the ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is hydroxy, Hal is Cl, Br or I and n is as previously defined in the presence of a Lewis acid to give the corresponding ω -halo tolylketone compound of structure (4) as described in *Arch. Pharm.* 306, 807 1973. In general, an appropriate toluene derivative of structure (1) and the ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is hydroxy, are melted together at about 50°C, then cooled to about 10°C after which a Lewis acid is added in an amount about 2.2 times the molar amount of the appropriate toluene derivative of structure (1) employed. The mixture is

35

- 34 -

heated at about 70°C for about 2 hours after which a 30% sodium acetate solution is added and extracted with ether. The organic layer is dried and the solvent evaporated to
5 give the corresponding ω -halo tolylketone compound of structure (4). The ω -halo tolylketone compound of structure (4) may be purified by procedures well known in the art, such as crystallization and/or distillation.

10 Suitable Lewis acids for the acylation reaction described in step d are well known and appreciated in the art. Examples of suitable Lewis acids are boron trichloride, aluminum chloride, titanium tetrachloride, boron trifluoride, tin tetrachloride, ferric chloride,
15 cobalt(II) chloride and zinc chloride, with aluminum chloride being preferred. The selection and utilization of suitable Lewis acids for the acylation reaction of step d is well known and appreciated by one of ordinary skill in the art.

20

The starting ω -halo compound of the structure Hal-(CH₂)_n-C(=O)-B, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined are commercially available or easily prepared by generally known methods.

25


While also not necessary for utilization in the acylation reaction of step d, the phenol functionality of those toluene derivatives of structure (1), wherein A is hydroxy may be protected with a suitable protecting group.
30 For example, suitable protecting groups for the phenolic hydroxy include methyl ether, 2-methoxyethoxymethyl ether (MEM), cyclohexyl ether, o-nitrobenzyl ether, 9-anthryl ether, t-butyldimethylsilyl ether, acetate, benzoate, methyl carbamate, benzyl carbamate, aryl pivaloate and aryl
35 methanesulfonate.

In step e, to appropriate toluene derivative of

- 35 -

structure (1) is acylated with an appropriate cyclopropyl compound of the structure

5




10 wherein B is as previously defined to give the corresponding cyclopropyl tolylketone derivative of structure (5) as described previously in step d.

15 In step f, the appropriate ethylbenzene derivative of structure (2) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo ethylphenylketone compound of structure (6) as described previously in step d.

20

In step g, the appropriate ethylbenzene derivative of structure (2) is acylated with an appropriate cyclopropyl compound of the structure

25



30

35

- 36 -


wherein B is as previously defined to give the corresponding cyclopropyl ethylphenylketone derivative of structure (7) as described previously in step e.

5

In step h, the appropriate cumene derivative of structure (3) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo cumylketone compound of structure (8) as described previously in step d.

In step i, to appropriate cumene derivative of structure (3) is acylated with an appropriate cyclopropyl compound of the structure

20



wherein B is as previously defined to give the corresponding cyclopropyl cumylketone derivative of structure (9) as described previously in step e.

25

In step j, the cyclopropyl functionality of the appropriate cyclopropyl tolylketone derivative of structure (5) is ring-opened to give the corresponding ω -halo tolylketone compound of structure (4) wherein $n = 3$.

30

For example, the appropriate cyclopropyl tolylketone derivative of structure (5) is reacted with an appropriate hydrogen halide in a suitable organic solvent, such as toluene, xylene and ethanol. The reaction is typically conducted at a temperature range of from room temperature to 70°C and for a period of time ranging from 20 minutes to 10 hours. The corresponding ω -halo tolylketone compound of structure (4) wherein $n = 3$ is isolated from the reaction

35

- 37 -

zone by evaporation of the solvent or may be stored in a solution of the hydrogen halide.

5 In step k, the appropriate ω -halo tolylketone compound of structure (4) wherein $n = 3$ is ring-closed to give the corresponding cyclopropyl tolylketone derivative of structure (5).

10 For example, the appropriate ω -halo tolylketone compound of structure (4) wherein $n = 3$ is reacted with an appropriate non-nucleophilic base, such as sodium hydroxide or potassium hydroxide in a suitable organic protic solvent, such as methanol or ethanol. The reaction is
15 typically conducted at a temperature range of from -10°C to room temperature and for a period of time ranging from 10 minutes to 5 hours. The corresponding cyclopropyl tolylketone derivative of structure (5) is isolated from the reaction zone by extractive methods as are known in the
20 art and may be purified by distillation.

 In step l, the cyclopropyl functionality of the appropriate cyclopropyl ethylphenylketone derivative of structure (7) is ring-opened to give the corresponding ω -
25 halo ethylphenylketone compound of structure (6) wherein $n = 3$ as described previously in step j.

 In step m, the appropriate ω -halo ethylphenylketone compound of structure (6) wherein $n = 3$ is ring-closed to
30 give the corresponding cyclopropyl ethylphenylketone derivative of structure (7) as described previously in step k.

 In step n, the cyclopropyl functionality of the appropriate cyclopropyl cumylketone derivative of structure
35 (9) is ring-opened to give the corresponding ω -halo cumylketone compound of structure (8) wherein $n = 3$ as described previously in step j.

- 38 -

In step o, the appropriate ω -halo cumylketone compound of structure (8) wherein $n = 3$ is ring-closed to give the
5 corresponding cyclopropyl cumylketone derivative of structure (9) as described previously in step k.

In step p, the appropriate ω -halo ethylphenylketone compound of structure (6) is methylated to give the
10 corresponding ω -halo cumylketone compound of structure (8) as described previously in step a.

In step q, the appropriate cyclopropyl tolylketone derivative of structure (5) is dimethylated to give the
15 corresponding cyclopropyl cumylketone derivative of structure (9) as described previously in step c.

In step r, the appropriate ω -halo tolylketone compound of structure (4) is methylated to give the corresponding ω -
20 halo ethylphenylketone compound of structure (6) as described previously in step a.

In step s, the appropriate ω -halo tolylketone compound of structure (4) is dimethylated to give the corresponding
25 ω -halo cumylketone compound of structure (8) as described previously in step c.

In step t, the appropriate cyclopropyl ethylphenylketone derivative of structure (7) is methylated
30 to give the corresponding cyclopropyl cumylketone derivative of structure (9) as described previously in step a.

In step u, the appropriate cyclopropyl tolylketone derivative of structure (5) is methylated to give the
35 corresponding cyclopropyl ethylphenylketone derivative of structure (7) as described previously in step a.

- 39 -

Starting materials for use in Scheme A are readily available to one of ordinary skill in the art.

5 The following examples present typical syntheses as described in Scheme A. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to
10 grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

15

Example 1

Step h: 4-Chloro-1-(4-isopropyl-phenyl)-butan-1-one
Slurry aluminum chloride (140.9g, 1.075mol) and 4-chlorobutyryl chloride (148g, 1.05mol) in methylene
20 chloride (1.0L) add, by dropwise addition, cumene (125g, 1.04mol) over a thirty minute period under a nitrogen atmosphere while maintaining the internal temperature between 5-8°C with an ice bath. Allow the stirred solution to come to room temperature and continue stirring under
25 nitrogen for 14 hours. Cautiously add the methylene chloride solution to 1L of crushed ice with stirring and add additional methylene chloride (400mL). Separate the organic phase and wash with 10% hydrochloric acid (3X300mL), water (3X300mL), 10% sodium bicarbonate
30 (3X300mL) and water (3X300mL). Dry (MgSO₄), filter and wash with methylene chloride (150mL). Evaporate the solvent to give the title compound (203g, 86%) as a clear oil which crystallizes on standing; mp 35-37°C.

35 ¹H NMR (300MHz, CDCl₃) δ 7.91 (d, J=8.2Hz, 2H), 7.31 (d, J=8.2Hz, 2H), 3.65 (t, J=6.3Hz, 2H), 3.13 (t, J=6.9Hz, 2H), 2.95 (p, J=6.9Hz, 1H), 2.20 (p, J=6.6Hz, 2H), 1.26 (d, J=6.9Hz, 6H); ¹³C NMR (75MHz, CDCl₃) δ 198.2, 154.4, 134.4,

- 40 -

128.1, 126.5, 44.5, 32.96, 34.0, 26.7, 23.5; IR (CDCl₃)
 2950, 2920, 1675, 1680, 1600, 1410, 1225 cm⁻¹; MS (GCCIMS
 (methane)) 255 (3), 251 (10), 227 (30 (M+H)), 225 (100
 5 (M+H)), 189 (70), 147 (95), 107 (13), 105 (40).

Anal. Calcd for C₁₃H₁₇OCl: C, 69.48; H, 7.62; Found: C,
 69.31; H, 7.39.

10

Example 2Step d: 4-Chloro-1-(4-methyl-phenyl)-butan-1-one

Suspend anhydrous AlCl₃ (156g, 1.15mol) in toluene (1500mL)
 and cool to 2-4°C. Add, by slow addition, a solution of 4-
 chlorobutyryl chloride (165.5g, 1.15mol) in toluene
 15 (300mL). Stir for 15 minutes and pour into stirring ice-
 water (2.5L). Stir for 30 hours, decant the toluene and
 extract the aqueous phase with toluene (700mL). Combine
 the organic layers and wash three times with water (1L, 1L,
 500mL). Evaporate the solvent in vacuo to give the title
 20 compound as a pale yellow oil (292.3g, 95%).

Example 3Step k: Cyclopropyl-p-tolyl-methanone

Dissolve potassium hydroxide (126g) in methanol (450mL),
 25 stir and cool in an ice-water bath. Add, by dropwise
 addition, a solution of 4-chloro-1-(4-methyl-phenyl)-butan-
 1-one (292g) in methanol (450mL). Stir for 20 minutes at
 8-10°C and partially evaporate the methanol in vacuo to
 give 400mL of a residue. Pour the residue, with stirring,
 30 into water (1500mL), filter the white solid and dry under
 vacuum to give the title compound as a white solid (190.8g,
 90%).

The following compounds can be prepared using the
 35 methodology depicted in Scheme A:

Cyclopropyl-(4-isopropyl-phenyl)-methanone;

- 41 -

Cyclopropyl-(4-ethyl-phenyl)-methanone; and

4-Chloro-1-(4-ethyl-phenyl)-butan-1-one.

5

10

15

20

25

30

35

- 42 -

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is OH, Cl, Br or I may be prepared as described
5 in Scheme B. In Scheme B, all substituents are as previously defined unless otherwise indicated.

10

15

20

25

30

35

:



Scheme B provides various general synthetic procedures for preparing the novel intermediates of formula (II),

- 44 -

formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is OH, Cl, Br or I.

- 5 In step a, the appropriate ω -halo cumylketone compound of structure (8) is halogenated to give the corresponding ω -halo-halocumylketone compound of structure (10).

For example, the appropriate ω -halo-halocumylketone
10 compound of structure (10) may be prepared by reacting an appropriate ω -halo cumylketone compound of structure (8) with a suitable halogenating agent optionally in the presence of a catalytic amount of a suitable initiator. Examples of suitable brominating agents are N-
15 bromosuccinimide, and 1,3-dibromo-5,5-dimethyl hydantoin, with N-bromosuccinimide being preferred. An example of suitable chlorinating agent is N-chlorosuccinimide and an example of a suitable iodinating agent is N-iodosuccinimide. Examples of suitable initiators are benzoyl peroxide, AIBN,
20 t-butyl peroxide and ultraviolet light. The reaction is carried out in a solvent, such as carbon tetrachloride, methylene chloride, 1,2-dichlorobenzene, 1,2-dichloroethane, ethyl formate or ethyl acetate, with carbon tetrachloride being the preferred solvent. The reaction
25 time varies from about 1/2 hour to 8 hours, preferably 1/2 to 2 hours and the reaction temperature varies from about 25°C to the reflux temperature of the solvent employed, preferably 70°C to 80°C. The corresponding ω -halo-halocumylketone compound of structure (10) is recovered
30 from the reaction zone by extractive methods as are known in the art followed by evaporation of the solvent.

In addition, the halogenation reaction of step a may be carried out in a 2-phase procedure. For example, the
35 appropriate ω -halo-halocumylketone compound of structure (10) may be prepared by reacting an appropriate ω -halo cumylketone compound of structure (8) with a suitable halogenating agent, such as sodium bromate/sodium bromide,

- 45 -

in a solvent mixture such as methylene chloride and water, catalyzing the reaction with, for example, ultraviolet light. The corresponding ω -halo-halocumylketone compound of structure (10) is recovered from the reaction zone by extractive methods as are known in the art followed by evaporation of the solvent.

The ω -halo-halocumylketone compound of structure (10) may dehydrohalogenate to the corresponding α -methylstyrene, giving various mixtures of ω -halo-halocumylketone compound of structure (10) and α -methylstyrene compounds. The α -methylstyrene compounds in such a mixture may be back-converted to ω -halo-halocumylketone compound of structure (10) by treatment with anhydrous hydrogen halide gas. Typically, a solution of the mixture of ω -halo-halocumylketone compound of structure (10) and α -methylstyrene compounds in a suitable organic solvent, such as methylene chloride or acetonitrile, is treated with a suitable anhydrous hydrogen halide gas, such as hydrogen chloride. The reaction is typically treated with the hydrogen halide gas for a period of time ranging from 30 minutes to 5 hours and at a temperature range of from 0°C to room temperature. The remediated ω -halo-halocumylketone compound of structure (10) may be isolated by evaporation of solvent, but may be stored as a solution in the organic solvent containing hydrogen halide gas.

In addition, halogen exchange of the benzylic halogen can be accomplished by thorough solvolysis in the presence of the appropriate hydrogen halide.

For example, the ω -chloro-halocumylketone compound of structure (10) can be prepared from the ω -bromo-halocumylketone compound of structure (10) by thorough aqueous solvolysis in the presence of hydrogen chloride.

- 46 -

In step b, the appropriate cyclopropyl cumylketone derivative of structure (9) is halogenated to give the corresponding cyclopropyl halocumylketone compound of structure (11) as described previously in step a.

In step c, the cyclopropyl functionality of the appropriate cyclopropyl halocumylketone compound of structure (11) is ring-opened to give the corresponding ω -halo-halocumylketone compound of structure (10) wherein $n = 3$ as described previously in Scheme A, step j.

In step d, the appropriate ω -halo ethylphenylketone compound of structure (6) is halogenated to give the corresponding ω -halo-haloethylphenylketone compound of structure (12) as described previously in step a.

In step e, the appropriate ω -halo tolylketone compound of structure (4) is halogenated to give the corresponding ω -halo halotolylketone compound of structure (13) as described previously in step a.

In step f, the appropriate cyclopropyl ethylphenylketone derivative of structure (7) is halogenated to give the corresponding cyclopropyl haloethylphenylketone compound of structure (14) as described previously in step a.

In step g, the appropriate cyclopropyl tolylketone derivative of structure (5) is halogenated to give the corresponding cyclopropyl halotolylketone of structure (15) as described previously in step a.

In step h, the appropriate cyclopropyl halotolylketone of structure (15) is ring-opened to give the corresponding ω -halo halotolylketone compound of structure (13) wherein $n = 3$ as described previously in Scheme A, step j.

- 47 -

In step i, the appropriate cyclopropyl haloethylphenylketone compound of structure (14) is ring-opened to give the corresponding ω -halo-haloethylphenylketone compound of structure (12) wherein n = 3 as described previously in Scheme A, step j.

In addition, the novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is OH may be prepared by solvolysis of the corresponding novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is Cl, Br or I, with, for example, tetrahydrofuran and water or any slightly acidic medium.

15

Starting materials for use in Scheme B are readily available to one of ordinary skill in the art.

The following examples present typical syntheses as described in Scheme B. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

30

Example 4

1-[4-(1-Bromo-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one

Step a, Method A:

Dissolve 4-chloro-1-(4-isopropyl-phenyl)-butan-1-one (2.10g, 9.35mmol) in carbontetrachloride (30mL), add N-bromosuccinimide (1.75g, 9.83mmol) and benzoylperoxide (3mg) and stir at reflux for 1 hour. Cool the reaction mixture, filter, wash with water and brine. Dry (MgSO₄), filter and

- 48 -

evaporate the solvent *in vacuo* to give the title compound as an amber oil.

5 Step a, Method B:

Dissolve 4-chloro-1-(4-isopropyl-phenyl)-butan-1-one (5.00g, 22.2mmol) and N-bromosuccinimide (4.1g, 23.0mmol) in carbon tetrachloride (25mL) and add AIBN radical initiator (300mg). Stir and maintain under a nitrogen atmosphere at 80-90°C or optionally irradiate with a sunlamp until a vigorous exotherm occurs at which point momentarily remove until reflux subsides and then reapply the heat. Reflux for 30 minutes and add another portion of N-bromosuccinimide (100mg) while maintaining reflux and reflux an additional 15 minutes. Cool to room temperature and precipitate the succinimide from the solution by allowing to stand overnight. Filter and wash the succinimide (2.25g) with carbon tetrachloride (20mL). Combine the filtrates and evaporate the solvent *in vacuo* to give the title compound as a yellow oil (6.80g, 100%).

¹H NMR (300MHz, CDCl₃) δ 7.935 (d, J=8.4Hz, 2H), 7.70 (d, J=8.4Hz, 2H), 3.66 (t, J=6.3Hz, 2H), 3.16 (t, J=6.8Hz, 2H), 2.21 (p, J=6.8Hz, 2H), 2.19 (s, 6H); ¹³C NMR (75MHz, CDCl₃) δ 198.1 (151.63), 135.8, 128.0, 126.0, 62.3, 44.5, 35.3, 35.1, 26.7; IR (neat) 2970, 2910, 1680, 1675, 1600, 1402, 1225, 1180 cm⁻¹.

Step a, Method C:

30 Dissolve 4-chloro-1-(4-isopropyl-phenyl)-butan-1-one (74.7g, 333mmol) in methylene chloride (250mL) and add sodium bromate (17.6g, 117mmol) in water (75mL) in a three-necked Morton flask equipped with an overhead stirrer. Cool the solution to 10°C and irradiate with two 150W incandescent flood lamps. Add, by dropwise addition, a solution of sodium bromide (24g, 233mmol) and stir for 2 hours. Illuminate for another 30 minutes, add sodium dithionate (2.0g), separate the organic phase, dry (MgSO₄)

- 49 -

and evaporate the solvent *in vacuo* to give the title compound (100g, 99%).

5 Step a, Method D:

Dissolve 1-[4-(1-bromo-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one (10.4g assayed at 67% by weight and containing 18wt% 1-[4-(2-propene)-phenyl]-4-chloro-butan-1-one) in methylene chloride (50mL) and sparge hydrogen chloride
10 through the solution for 70 minutes. Evaporate the solvent *in vacuo* to give a 3:1 mixture of 1-[4-(1-bromo-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one and 1-[4-(1-chloro-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one (11.6g).

15

Example 5

(4-Bromomethyl-phenyl)-cyclopropyl-methanone

Step g: Dissolve 4-chloro-1-(4-isopropyl-phenyl)-butan-1-one (20g, 124mmol) and 2,2'-Azolons (2-methylpropionitrile)
20 (0.5g) in methylene chloride (100mL) and cool to 5°C. Add a suspension of N-bromosuccinimide (12g) in methylene chloride (50mL) and irradiate with light (150 Watt lamp), maintaining the temperature at 5°C. After 2, 3 and 7 hour time periods, add additional N-bromosuccinimide (6g, 6g,
25 2.8g) and continue stirring. After 7.5 hours, wash with water (200mL) and with 0.4M sodium hydrogen carbonate (2X200mL). Dry (Na₂SO₄), evaporate the solvent *in vacuo* and recrystallize (hexane) to give the title compound as a crystalline solid (26.7g).

30

The following compounds can be prepared by procedures depicted in Scheme B:

[4-(1-bromoethyl)-phenyl]-cyclopropyl-methanone;

35


[4-(1-bromo-1-methyl-ethyl)-phenyl]-cyclopropyl-methanone;

1-[4-(1-bromomethyl)-phenyl]-4-chloro-butan-1-one; and

1-[4-(1-bromoethyl)-phenyl]-4-chloro-butan-1-one.

5 The novel intermediates of formula (VIII) and (IX) and
the novel intermediates of formula (II), formula (III),
formula (IV), formula (V), formula (VI) and formula (VII)
wherein R_5 is Cl, Br or I may also be prepared as described
in Scheme C. In Scheme C, all substituents are as
10 previously defined unless otherwise indicated.

Scheme C



- 51 -


Scheme C provides various general synthetic procedures for preparing the the novel intermediates of formula (VIII) and (IX) and novel intermediates of formula (II), formula
 5 (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is Cl, Br or I.

In step a, the appropriate α -methylstyrene compound of structure (16) is acylated with an appropriate ω -halo
 10 compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo- α -methylstyrene compound of structure (17) as described previously in Scheme A, step d.

15

In step b, the appropriate α -methylstyrene compound of structure (16) is acylated with an appropriate cyclopropyl compound of the structure

20



wherein B is as previously defined to give the
 25 corresponding cyclopropyl α -methylstyreneketone derivative of structure (18) as described previously in Scheme A, step e.

In step c, the appropriate ω -halo- α -methylstyrene
 30 compound of structure (17) wherein $n = 3$ is ring-closed to give the corresponding cyclopropyl α -methylstyreneketone derivative of structure (18) as described previously in Scheme A, step k.

35

In step d, the appropriate cyclopropyl α -methylstyreneketone derivative of structure (18) is ring-opened to give the corresponding ω -halo- α -methylstyrene

- 52 -

compound of structure (17) wherein $n = 3$ as described previously in Scheme A, step j.

5 In step e, the appropriate ω -halo- α -methylstyrene compound of structure (17) is hydrohalogenated to give the corresponding ω -halo-halocumylketone derivative of structure (10).

10 For example, the appropriate ω -halo- α -methylstyrene compound of structure (17) is treated with anhydrous hydrogen halide at a temperature range of from -50°C to room temperature, preferably 0°C -5°C and for a period of time ranging from 5 minutes to 2 hours. The ω -halo-
15 halocumylketone derivative of structure (10) is recovered from the reaction zone by purging with nitrogen.

 In step f, the appropriate ω -halo-halocumylketone derivative of structure (10) is dehydrohalogenated to give
20 the corresponding ω -halo- α -methylstyrene compound of structure (17) by treatment with base as is known in the art.

 In step g, the appropriate cyclopropyl α -
25 methylstyreneketone derivative of structure (18) is hydrohalogenated to give the corresponding cyclopropyl halocumylketone compound of structure (11) as described previously in step e.

30 In step h, the appropriate cyclopropyl halocumylketone compound of structure (11) is dehydrohalogenated to give the corresponding cyclopropyl α -methylstyreneketone derivative of structure (18) as described previously in step f.

35

- 53 -

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is CN may be prepared as described in Scheme D.

5 In Scheme D, all substituents are as previously defined unless otherwise indicated.

10

15


20

25

30


35

Scheme D



Scheme D provides various general synthetic procedures for preparing the novel intermediates of formula (II),

Scheme D Cont.



20 formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is CN.

25 In step a, the appropriate ω -halo-halocumylketone compound of structure (10) is cyanated to give the corresponding ω -halo-cyanocumylketone compound of structure (19).

30 For example, the appropriate ω -halo-cyanocumylketone compound of structure (19) may be prepared by reacting an appropriate ω -halo-halocumylketone compound of structure (10) with a suitable cyanating agent. Examples of suitable cyanating agents are trimethylsilyl cyanide, diethylaluminum cyanide and tetrabutylammonium cyanide, with trimethylsilyl cyanide being preferred. The reaction is carried out in a solvent, such as methylene chloride, tetrachloroethane and carbon tetrachloride, with methylene chloride being the preferred solvent. A catalytic amount of a suitable Lewis acid may also be employed in the reaction.

- 56 -

Examples of suitable Lewis acids are boron trichloride, aluminum chloride, titanium tetrachloride, boron trifluoride, tin tetrachloride and zinc chloride, with tin tetrachloride being preferred. The reaction time varies from about 1/2 hour to 8 hours, preferably 1/2 to 2 hours and the reaction temperature varies from about 0°C to room temperature, preferably room temperature. The ω -halo-cyanocumylketone compound of structure (16) is recovered from the reaction zone by an aqueous quench followed by extraction as is known in the art. The ω -halo-cyanocumylketone compound of structure (16) may be purified by procedures well known in the art, such as chromatography and crystallization.

15

In step b, the appropriate ω -halo cumylketone compound of structure (8) is cyanated to give the corresponding ω -halo-cyanocumylketone compound of structure (19).

20

For example, the ω -halo-cyanocumylketone compound of structure (19) may be prepared by reacting an appropriate the ω -halo cumylketone compound of structure (8) with a suitable cyanating agent. Examples of suitable cyanating agent are cyanogen chloride, cyanogen bromide and cyanogen iodide, with cyanogen chloride being preferred. The reaction is carried out according to the procedures outlined by Tanner and Bunce, J. Am. Chem. Soc., 91, 3028 (1969).

25

30

In step c, the appropriate cyclopropyl halocumylketone compound of structure (11) is cyanated to give the corresponding cyclopropyl cyanocumylketone compound of structure (20) as described previously in step a.

35

In step d, the appropriate cyclopropyl cumylketone derivative of structure (9) is cyanated to give the corresponding cyclopropyl cyanocumylketone compound of structure (20) as described previously in step b.

- 57 -

In step e, the appropriate ω -halo-haloethylphenylketone compound of structure (12) is cyanated to give the
5 corresponding ω -halo-cyanoethylphenylketone compound of structure (21) as described previously in step a.

In step f, the appropriate ω -halo-ethylphenylketone compound of structure (6) is cyanated to give the
10 corresponding ω -halo-cyanoethylphenylketone compound of structure (21) as described previously in step b.

In step g, the appropriate ω -halo halotolylketone compound of structure (13) is cyanated to give the
15 corresponding ω -halo cyanotolylketone compound of structure (22) as described previously in step a.

In step h, the appropriate ω -halo tolylketone compound of structure (4) is cyanated to give the corresponding ω -
20 halo cyanotolylketone compound of structure (22) as described previously in step b.

In step i, the appropriate cyclopropyl ethylphenylketone compound of structure (7) is cyanated to
25 give the corresponding cyclopropyl cyanoethylphenylketone compound of structure (23) as described previously in step b.

In step j, the appropriate cyclopropyl haloethylphenylketone compound of structure (14) is
30 cyanated to give the corresponding cyclopropyl cyanoethylphenylketone compound of structure (23) as described previously in step a.

35 In step k, the appropriate cyclopropyl tolylketone compound of structure (5) is cyanated to give the corresponding cyclopropyl cyanotolylketone compound of structure (24) as described previously in step b.

In step 1, the appropriate cyclopropyl halotolylketone of structure (15) is cyanated to give the corresponding
5 cyclopropyl cyanotolylketone compound of structure (24) as described previously in step a.

Starting materials for use in Scheme D are readily available to one of ordinary skill in the art.

10

The following examples present typical syntheses as described in Scheme D. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the
15 following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and
20 "μM" refers to micromolar.

Example 6

Step a: 2-[4-(4-chloro-butyl)-phenyl]-2-methyl-propionitrile

25 Dissolve 1-[4-(1-bromo-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one (2.00g, 6.59mmol) in anhydrous methylene chloride (20mL) and place under an argon atmosphere. Add trimethylsilyl cyanide (1.10mL, 8.25mmol) followed by tin (IV) chloride (0.20mL, 1.7mmol) via syringe. Stir at reflux
30 for 1 hour, add water (20mL) and stir for an additional 1/2 hour. Separate the layers and extract the aqueous layer with methylene chloride. Combine the organic layers, wash with brine, dry (MgSO₄), filter and evaporate the solvent in *vacuo*. Purify by silica gel chromatography (15% ethyl
35 acetate/hexane) to give the title compound as a white solid; mp 79-80°C.

Example 7

- 59 -

Step 1: (4-Cyclopropanecarbonyl-phenyl)-acetonitrile

Mix (4-bromomethyl-phenyl)-cyclopropyl-methanone (5.0g, 21mmol), potassium cyanide (2.0g, 30mmol), tetra-
5 butylammonium bromide (150mg), water (5mL) and acetonitrile (50mL). Mechanically stir at room temperature for 3 hours, pour into water (450mL) and stir overnight. Collect by filtration and recrystallize (hexane) to give the title compound as a white crystalline solid; mp 86-87°C.

10

The following compounds can be prepared by the synthetic procedures depicted in Scheme D:

15 2-(4-Cyclopropanecarbonyl-phenyl)-propionitrile;

2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionitrile;

[4-(4-Chloro-butyryl)-phenyl]-acetonitrile; and

20

2-[4-(4-Chloro-butyryl)-phenyl]-propionitrile.

25

30

35

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is CN may also be prepared as described in Scheme E. In Scheme E, all substituents are as previously defined unless otherwise indicated.

Scheme E



Scheme E provides alternative various general synthetic procedures for preparing the novel intermediates of formula

(II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is CN.

5 In step a, the appropriate phenylacetonitrile compound of structure (25) is methylated to give the corresponding 2-cyanoethylbenzene compound of structure (26) as described previously in Scheme A, step a.

10 Appropriate phenylacetonitrile compounds of structure (25) may be prepared from the corresponding benzyl halide by techniques and procedures well known by one of ordinary skill in the art and described previously in Scheme D, step a.

15 Appropriate benzyl halide compounds may be prepared from the corresponding toluene derivative of structure (1) as described previously in Scheme B, step a.

20 In step b, the appropriate 2-cyanoethylbenzene compound of structure (26) is methylated to give the corresponding 2-cyano-2-propylbenzene compound of structure (27) as described previously in Scheme A, step a.

25 Appropriate 2-cyanoethylbenzene compound of structure (26) may be prepared from the corresponding α -methylbenzyl halide by techniques and procedures well known by one of ordinary skill in the art and as described previously in step a.

30 Appropriate α -methylbenzyl halide compounds may be prepared from the corresponding ethylbenzene derivative of structure (2) as described previously in Scheme B, step a.


35 In step c, the appropriate phenylacetonitrile compound of structure (25) is dimethylated to give the corresponding 2-cyano-2-propylbenzene compound of structure (27) as described previously in Scheme A, step c.

In step d, the appropriate phenylacetonitrile compound of structure (25) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo cyanotolylketone compound of structure (22) as described previously in Scheme A, step d.

10

In step e, the appropriate phenylacetonitrile compound of structure (25) is acylated with an appropriate cyclopropyl compound of the structure

15



wherein B is as previously defined to give the corresponding cyclopropyl cyanotolylketone compound of structure (24) as described previously in Scheme A, step e.

20


In step f, the appropriate 2-cyanoethylbenzene compound of structure (26) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo-cyanoethylphenylketone compound of structure (21) as described previously in Scheme A, step d.

25

30

In step g, the appropriate 2-cyanoethylbenzene compound of structure (26) is acylated with an appropriate cyclopropyl compound of the structure

35



- 63 -

wherein B is as previously defined to give the corresponding cyclopropyl cyanoethylphenylketone compound of structure (23) as described previously in Scheme A, step
5 e.

In step h, the appropriate 2-cyano-2-propylbenzene compound of structure (27) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein
10 B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω -halo-cyanocumylketone compound of structure (19) as described previously in Scheme A, step d.


15 Appropriate 2-cyano-2-propylbenzene compound of structure (27) may be prepared from the corresponding α,α -dimethylbenzyl halide by techniques and procedures well known by one of ordinary skill in the art and as described previously in step a.

20

Appropriate α,α -dimethylbenzyl halide compounds may be prepared from the corresponding cumene derivative of structure (3) as described previously in Scheme B, step a.

25 In step i, the appropriate 2-cyano-2-propylbenzene compound of structure (27) is acylated with an appropriate cyclopropyl compound of the structure

30



wherein B is as previously defined to give the corresponding cyclopropyl cyanocumylketone compound of
35 structure (20) as described previously in Scheme A, step e.

In step j, the cyclopropyl functionality of the appropriate cyclopropyl cyanotolylketone compound of

- 64 -

structure (24) is ring-opened to give the corresponding ω -halo cyanotolylketone compound of structure (22) wherein $n = 3$ as described previously in Scheme A, step j.

5

In step k, the appropriate ω -halo cyanotolylketone compound of structure (22) wherein $n = 3$ is ring-closed to give the corresponding cyclopropyl cyanotolylketone compound of structure (24) as described previously in
10 Scheme A, step k.

In step l, the cyclopropyl functionality of the appropriate cyclopropyl cyanoethylphenylketone compound of structure (23) is ring-opened to give the corresponding ω -
15 halo-cyanoethylphenylketone compound of structure (21) wherein $n = 3$ as described previously in Scheme A, step j.

In step m, the appropriate ω -halo-cyanoethylphenylketone compound of structure (21) wherein n
20 $= 3$ is ring-closed to give the corresponding cyclopropyl cyanoethylphenylketone compound of structure (23) as described previously in Scheme A, step k.

In step n, the cyclopropyl functionality of the appropriate cyclopropyl cyanocumylketone compound of
25 structure (20) is ring-opened to give the corresponding ω -halo-cyanocumylketone compound of structure (19) wherein $n = 3$ as described previously in Scheme A, step j.

30 In step o, the appropriate ω -halo-cyanocumylketone compound of structure (19) is ring-closed to give the corresponding cyclopropyl cyanocumylketone compound of structure (20) as described previously in Scheme A, step k.

35 In step p, the appropriate ω -halo-cyanoethylphenylketone compound of structure (21) is methylated to give the corresponding ω -halo-

cyanocumylketone compound of structure (19) as described previously in Scheme A, step a.

- 5 In step q, the appropriate cyclopropyl cyanotolylketone compound of structure (24) is dimethylated to give the corresponding cyclopropyl cyanocumylketone compound of structure (20) as described previously in Scheme A, step c.
- 10 In step r, the appropriate ω -halo cyanotolylketone compound of structure (22) is methylated to give the corresponding ω -halo-cyanoethylphenylketone compound of structure (21) as described previously in Scheme A, step a.
- 15 In step s, the appropriate ω -halo cyanotolylketone compound of structure (22) is dimethylated to give the corresponding ω -halo-cyanocumylketone compound of structure (19) as described previously in Scheme A, step c.
- 20 In step t, the appropriate cyclopropyl cyanoethylphenylketone compound of structure (23) is methylated to give the corresponding cyclopropyl cyanocumylketone compound of structure (20) as described previously in Scheme A, step a.
- 25 In step u, the appropriate cyclopropyl cyanotolylketone compound of structure (24) is methylated to give the corresponding cyclopropyl cyanoethylphenylketone compound of structure (23) as described previously in Scheme A, step
- 30 a.

Starting materials for use in Scheme E are readily available to one of ordinary skill in the art.

- 35 The following examples present typical syntheses as described in Scheme E. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the

following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

Example 7

10 Step c: Cumyl cyanide

Place phenylacetonitrile (92.3mL, 0.800mol), tetra n-butylammonium chloride (4.45g of a 50% solution, 8.0mmol) and 50% aqueous sodium hydroxide solution (2.874 mole NaOH) into a 3-neck round-bottom flask, with a thermowell, 15 overheard stirrer, reflux condenser with a dry-ice/acetone trap and a sparge tube. Heat to 40-70C with stirring at 115 RPM (paddle stir blade), and bubble in methyl chloride gas (11.7g, 0.232 mole) over a 30 minute period. Turn off the methyl chloride addition and heat and stir overnight.

20

Sparge additional methyl chloride (35.4g, 0.700 mol) into the reaction mixture (heated to 35C) over a period of 2 hours. Stir the resulting mixture at ambient temperature for 22 hours and sparge additional methyl chloride (39.5g, 25 0.781mol) into the reaction mixture at a temperature of 40-70C (mostly at 55-60C). Sparge additional methyl chloride (8.7g, 0.172mol) into the reaction mixture and allow to cool to 30C. Remove the condenser and add deionized water (250mL) and heptane (250mL). Transfer to a separatory 30 funnel and draw off the aqueous (bottom) layer. Wash the remaining organic layer with fresh water (2X100mL), evaporate the solvent in vacuo to give a dark red oil.

Add the oil, 50% aqueous sodium hydroxide (79g, 0.988 mole) 35 and tetra n-butylammonium chloride (1.0g, 3.6mmol) to a 500mL 3-necked round bottom flask with a magnetic stir bar. Using the same experimental procedure described above, sparge in methyl chloride. Heat to 40-60C, stir and sparge in methyl chloride (20.5g, 0.40 mole) over 1 hour. Allow

- 67 -

the reaction mixture to cool, add deionized water (100g) and stir. Allow the layers to settle and remove the bottom layer by pipet. Repeat wash with additional water (100g) to give the title compound as a dark orange oil (111.0g, wet with water).

Example 8

Step g: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionitrile

Dissolve potassium t-butoxide (2.42g, 21.6mmol) in diglyme (8mL), cool to 10°C and slowly add with mechanical stirring, a solution of (4-cyclopropanecarbonyl-phenyl)-acetonitrile (2g, 10.8mmol) and methyl iodide (1.5mL, 24.0mmol) in diglyme (10mL). After 10 minutes, add additional potassium t-butoxide (0.3g, 2.6mmol) in diglyme (2mL) and stir for a total of 45 minutes. Pour into a mixture of water (100mL) and ethyl acetate (50mL) and adjust the pH to 1.5-2 with dilute hydrochloric acid.

Separate the organic phase and extract the aqueous phase with ethyl acetate (50mL). Combine the organic phases and wash with brine (2X100mL). Dry (Na₂SO₄), evaporate the solvent *in vacuo* and recrystallize (ethyl ether/hexane) to give the title compound as a yellow solid; mp 80-82°C.

The following compounds can be prepared by procedures depicted in Scheme E:

- (4-Cyclopropanecarbonyl-phenyl)-acetonitrile;
- 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionitrile;
- 2-(4-Cyclopropanecarbonyl-phenyl)-propionitrile;
- [4-(4-Chloro-butyryl)-phenyl]-acetonitrile; and
- 2-[4-(4-Chloro-butyryl)-phenyl]-propionitrile.

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is COOalkyl may also be prepared as described in Scheme F. In Scheme F, all substituents are as previously defined unless otherwise indicated.

Scheme F

10


15

20

25

30

35



Scheme F provides alternative various general synthetic procedures for preparing the novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R₅ is COOalkyl.

In step a, the appropriate phenylacetic acid ester compound of structure (28) is methylated to give the corresponding α -methylphenylacetic acid ester compound of structure (29) as described previously in Scheme A, step a.

Appropriate phenylacetic acid ester compounds of structure (28) are prepared from the corresponding phenylacetic acid compounds by standard esterification reactions which are well known by one of ordinary skill in the art.

Appropriate phenylacetic acid compounds may be prepared by hydrolysis of the corresponding phenylacetone nitrile compounds of structure (25) by techniques and procedures well known and appreciated by one of ordinary skill in the art, such as base hydrolysis. Alternatively, the phenylacetic acid compounds may be prepared by electrochemical carboxylation of the corresponding benzyl halide as described in Scheme H, step h.

In step b, the appropriate α -methylphenylacetic acid ester compound of structure (29) is methylated to give the corresponding α,α -dimethylphenylacetic acid ester compound of structure (30) as described previously in Scheme A, step a.

Alternatively α -methylphenylacetic acid ester compound of structure (29) are prepared for the corresponding α -methylphenylacetic acid compounds by standard esterification reactions which are well known by one of ordinary skill in the art as described in step a.

Appropriate α -methylphenylacetic acid compounds may be prepared by hydrolysis of the corresponding 2-cyanoethylbenzene compound of structure (26) as described previously in step a. Alternatively, the α -methylphenylacetic acid compounds may be prepared by electrochemical carboxylation of the corresponding α -methylbenzyl halide as described in Scheme H, step h.

10 In step c, the appropriate phenylacetic acid ester compound of structure (28) is dimethylated to give the corresponding α,α -dimethylphenylacetic acid ester compound of structure (30) as described previously in Scheme A, step c.

15 Alternatively α,α -dimethylphenylacetic acid ester compound of structure (30) are prepared for the corresponding α,α -dimethylphenylacetic acid compounds by standard esterification reactions which are well known by one of ordinary skill in the art as described in step a.

Appropriate α,α -dimethylphenylacetic acid compounds may be prepared by hydrolysis of the corresponding 2-cyano-2-propylbenzene compound of structure (27) as described previously in step a. Alternatively, the α,α -dimethylphenylacetic acid compounds may be prepared by electrochemical carboxylation of the corresponding α,α -dimethylbenzyl halide as described in Scheme H, step h. Appropriate α,α -dimethylbenzyl halide compounds may be prepared by hydrohalogenation of the corresponding α -methylstyrene as described previously in Scheme C, step e.


In step d, the appropriate phenylacetic acid ester compound of structure (28) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -

- 71 -

keto-phenylacetic acid ester compound of structure (34) as described previously in Scheme A, step d.

- 5 In step e, the appropriate phenylacetic acid ester compound of structure (28) is acylated with an appropriate cyclopropyl compound of the structure

10




- 15 wherein B is as previously defined to give the corresponding cyclopropylketo-phenylacetic acid ester compound of structure (33) as described previously in Scheme A, step e.

- 20 In step f, the appropriate α -methylphenylacetic acid ester compound of structure (26) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (30) as described previously in Scheme A, step d.

25

- In step g, the appropriate α -methylphenylacetic acid ester compound of structure (29) is acylated with an appropriate cyclopropyl compound of the structure

30



- 35 wherein B is as previously defined to give the corresponding cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) as described previously in Scheme A, step e.

- 72 -

In step h, the appropriate α,α -dimethylphenylacetic acid ester compound of structure (30) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -keto- α,α -di-methylphenylacetic acid ester compound of structure (31) as described previously in Scheme A, step d.

10


Appropriate α,α -dimethylphenylacetic acid ester compound of structure (30) are prepared for the corresponding α,α -dimethylphenylacetic acid compounds by standard esterification reactions which are well known by one of ordinary skill in the art as described in step a.

Appropriate α,α -dimethylphenylacetic acid compounds may be prepared by hydrolysis of the corresponding 2-cyano-2-propylbenzene compound of structure (27) as described previously in step a.

20

In step i, the appropriate α,α -dimethylphenylacetic acid ester compound of structure (30) is acylated with an appropriate cyclopropyl compound of the structure

25



wherein B is as previously defined to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) as described previously in Scheme A, step e.

30

In step j, the appropriate ω' -halo- α' -keto- α,α -methylphenylacetic acid ester compound of structure (33) is methylated to give the corresponding ω' -halo- α' -keto- α,α -

35

- 73 -

di-methylphenylacetic acid ester compound of structure (32) as described previously in Scheme A, step a.

5 In step k, the cyclopropyl functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) is ring-opened to give the corresponding ω' -halo- α' -keto- α,α -di-methylphenylacetic acid ester compound of structure (31) wherein $n = 3$ as
10 described previously in Scheme A, step j.

 In step l, the appropriate ω' -halo- α' -keto- α,α -di-methylphenylacetic acid ester compound of structure (31) wherein $n = 3$ is ring-closed to give the corresponding
15 cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) as described previously in Scheme A, step k.

 In step m, the appropriate ω' -halo- α' -keto-phenylacetic
20 acid ester compound of structure (34) is dimethylated to give the corresponding ω' -halo- α' -keto- α,α -di-methylphenylacetic acid ester compound of structure (31) as described previously in Scheme A, step c.

25 In step n, the appropriate ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) is methylated to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) as described previously in Scheme A, step a.

30

 In step o, the cyclopropyl functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) is ring-opened to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid
35 ester compound of structure (33) wherein $n = 3$ as described previously in Scheme A, step j.

- 74 -

In step p, the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) wherein $n = 3$ is ring-closed to give the corresponding
5 cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) as described previously in Scheme A, step k.

In step q, the appropriate cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) is
10 methylated to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) as described previously in Scheme A, step a.

In step r, the appropriate cyclopropylketo-phenylacetic
15 acid ester compound of structure (36) is dimethylated to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) as described previously in Scheme A, step c.

20 In step s, the cyclopropyl functionality of the appropriate cyclopropylketo-phenylacetic acid ester compound of structure (36) is ring-opened to give the corresponding ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) wherein $n = 3$ as described
25 previously in Scheme A, step j.

In step t, the appropriate ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) wherein $n = 3$ as is ring-closed to give the corresponding cyclopropylketo-
30 phenylacetic acid ester compound of structure (36) as described previously in Scheme A, step k.

In step u, the appropriate cyclopropylketo-phenylacetic acid ester compound of structure (36) is methylated to give
35 the corresponding cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) as described previously in Scheme A, step a.

- 75 -

Starting materials for use in Scheme F are readily available to one of ordinary skill in the art.

5 The following examples present typical syntheses as described in Scheme F. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to
10 grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

15

Example 9

Step c: 2-Methyl-2-phenylpropionate, methyl ester

Equip a two liter, 3-necked, round bottom flask with a thermowell with a thermometer, heating mantle, mechanical
20 agitator, gas inlet for MeCl, rubber septum for sampling by syringe and a cryoscopic condensing system. The condensing system is composed of an 18 inch inner helical coil/outer jacket condenser chilled to -50C with refrigerated acetone topped with a dry ice cold finger having approximately 100
25 square inches of chilled surface area. The cold finder is vented through a drying tube filled with drying agent and MeCl is supplied from a lecture bottle mounted on a digital balance. The feed rate can be accurately controlled using a needle valve and monitored by rotometer. The rotometer is
30 calibrated with MeCl to give an average response of 2.5mg/min/scale division. Phenylacetic acid, ethyl ester is supplied via 1/16 inch stainless steel tubing inserted through the rubber sampling septum by a HPLC pump from a 1 liter bottle mounted on a digital balance. The bottle is
35 sealed with a septum and vented through a drying tube filled with drying agent. The temperature is controlled using a thermowatch to regulate the heating mantle. If cooling is required, it is accomplished either by immersing

- 76 -

the reaction flask in a water bath or simply by removing the mantle.

- 5 The phenylacetic acid, ethyl ester pump is primed with phenylacetic acid containing 1 st% t-butanol and the phenylacetic acid, ethyl ester balance is zeroed. The MeCl balance is zoned and a 200g sample of 60% NaH is weighed into a wide mouth plastic jar in a nitrogen filled glove
10 bag and is transferred to the reaction vessel through a funnel (sampling septum is removed). Through the same funnel is added anhydrous glyme (800mL) and the septum (pierced by the 1/16 inch phenylacetic acid, ethyl ester feed tube) is replaced. The mixture is agitated and heated
15 to 50C while MeCl (40g) is introduced. When the reaction mixture reaches 50C, the continuous addition of phenylactic acid, ethyl ester/t-butanol at 1 mL/min and MeCl at approximately 0.62g/min. is initiated. Samples of about p.l mL are withdrawn at intervals using a disposable
20 syringe fitted with an 8 inch needle. A portion of the sample (5-15 drops depending on the accumulation of product) is dissovled in 25% aqueous acetonitrile (5mL) and analyzed immediately. The reaction is continued for an additional 2 hours at 50C and then at ambient temperature
25 overnight.

- In the apparatus described above, agitate NaH (180g of 60%) and anhydrous glyme (800mL) and heat to 50C. Add MeCl (52g) along with methyl phenylacetate (20g). Stir for 1
30 hour at 50C, then add, by continuous addition, methyl phenylacetate (0.8mL/min) and MeCl (approximately 0.53g/min). Stir for 1 hour, stop the additoina and continue heating for 1.5 hours. Resume the additions and run for 45 minutes. Allow to agitate at ambient
35 temperature overnight. Heat the reaction to 50C and resume the addition of methyl phenylacetate (0.4mL/min) and MeCl (approximately 0.27g/min). When a total of 246g of methyl phenylacitrate has been added, stop the addition and agitate

- 77 -

overnight. Distill the glyme at 1 atm. until the pot temperature reaches 125C. Cool the residue and pour into water (1L) containing acetic acid (100mL). Filter through
5 filter aid and separate the phases. Distill the organic phase through a 10-plate Oldershaw™ column fitted with a reflux splitting head at 4mm Hg. Collect 10mL at a 5:2 reflux ratio and discard. Collect the title compound at a 2:1 reflux ratio and head temperature of 93C (100g).

10

Example 10

Step d: [4-(4-Chloro-butyryl)-phenyl]-acetic acid, ethyl ester and [3-(4-Chloro-butyryl)-phenyl]-acetic acid, ethyl ester

- 15 Method A: Load a 3-neck flask with sublimed AlCl₃ (293g, 2.08mmol) and heptane (400mL). Cool to below 5°C and slowly add chlorobutyryl chloride (125mL), keeping the temperature below 5°C. Add phenylethyl acetate (160mL), keeping the temperature below 10°C and stir overnight.
- 20 Decant the heptane layer and dissolve the residue in methylene chloride (400mL). Slowly pour the methylene chloride solution into a mixture of concentrated hydrochloric acid (200mL) and cracked ice. Separate the organic phase, wash with water (1L), followed by 5% sodium
- 25 hydrogen carbonate (1L). Evaporate the solvent *in vacuo* to give a red oil (243g).

Dissolve the red oil (243g) in methylene chloride (250mL) and sparge with hydrogen chloride gas for 1.5 hours and
30 evaporate the solvent *in vacuo* to give the title compound as a 50:50 mixture of para and meta isomers (243g).

Method B: Place aluminum chloride (293g) and methylene chloride (300mL) in a 1L, 3-neck round bottom flask with a
35 thermowell and equipped with a thermometer, mechanical stirrer, reflux condenser, equilibrating dropping funnel and ice bath. Cool to 10C and add, by dropwise addition, 4-chlorobutyryl chloride (169g), keeping the temperature

- 78 -

below 10C. After addition is complete, add, by dropwise addition, phenylethyl acetate (164g), keeping the temperature below 10C. Heat the reaction to 40C for 16 hours, slowly pour into a mechanically agitated 4L beaker containing ice/water (2000g) and stir for 1 hour. Separate the layers, extract the water phase with methylene chloride (50mL), filter the combined organic phases through a 1/4 inch thick bed of filter aid and extract sequentially with water (100mL) and 10 wt% Na₂CO₃ (200mL). Re-extract the carbonate solution with fresh methylene chloride (50mL) and wash the combined methylene chloride solutions with water (100mL). Distill off solvent at atmospheric pressure until the pot temperature reaches 120C. Cool the residue and dilute with 2B absolute ethanol (200mL). Heat the solution to 70C and sparge in anhydrous HCl (20g) over 10 minutes. After 40 minutes, cool the reaction and hold overnight under nitrogen. Evaporate the solvent in vacuo to give the title compound (258g).

20

Example 11

Step k: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester

25 Method A: Dissolve 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (100g) in xylene (500mL) and ethanol (100mL) and heat to 70°C. Sparge the atmosphere of the reaction with hydrogen chloride gas (24.6g) over 220 hours. Evaporate the solvent *in vacuo* to
30 give the title compound.

Method B: Add a solution of 5M HCl in acetonitrile (50mL, 9g of HCl, 247mmol) to 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (25.5g, 98mmol) and seal
35 in a 100mL flask with a rubber septum. Heat to 50°C for 4 hours, dilute with toluene (50mL), wash with water (50mL), aqueous 10% Na₂CO₃ (50mL) and then water (50mL). Evaporate the solvent *in vacuo* to give the title compound as an oil

- 79 -

(27.2g).

Method C: Place 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (86g, 330mmol) and dry acetonitrile (70mL) in a 250mL 3-neck round-bottom flask equipped with a magnetic stirbar, thermomotoer, gas inlet and distillation head connected to a balloon by way of a T fitting for pressure control. Slowly warm the reaction mixture with stirring to 60°C while sparging excess HCl into the reaction mixture for 6 hours, dilute with toluene (50mL), wash with water (50mL), aqueous 10% Na₂CO₃ (50mL) and then water (50mL). Evaporate the solvent *in vacuo* to give the title compound.

15

Method D: Place 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (91g, 350mmol) in a 1 L 3-neck round-bottom flask equipped with a magnetic stirbar, thermometer, gas inlet, and distillation head connected to a balloon by way of a T fitting for pressure control. Slowly sparge in anhydrous HCl, keeping the balloon slightly inflated. After 10 minutes, add acetonitrile (590mL), heat to 65°C and add excess HCl over 7 hours. Heat the mixture and remove acetonitrile/HCl overhead. After 500mL of acetonitrile is removed, add mixed xylene (200mL) and continue the distillation. Add additional xylene (200m) and after a total of 640mL of solvent has been removed (pot = 130°C and overhead = 130°C), add ethanol 2B (100mL). Remove the ethanol by distillation to give the title compound as a oil (330g).

Method E: Place 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (98g, 410mmol) and xylenes (600mL) in a 1L 3-neck round-bottom flask equipped with a magnetic stirbar, thermomoter, gas inlet and distillation head connected to a balloon by way of a T fitting for pressure control. Heat the reaction mixture to 80°C and slowly sparge in anhydrous HCl. After 100

- 80 -

minutes, add ethanol 2B (100mL) and HCl (26g) and heat to 35°C for 2 hours. Remove the ethanol and HCl by distillation with aspirator vacuum (pot = 35°C, overhead = 30°C) to give the title compound as a solution in xylene.

Method F: Place 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (500g) in a 4L Hastelloy reactor equipped with a gas inlet, overhead stirrer, temperature control and dip pipe for sampling. Heat the oil to 60C and evacuate the head space. Add HCl raising the pressure to 10psig and react for 80-300 minutes. Vent the excess HCl and sparge the oil with nitrogen for 5 minutes to give the title compound.

Method G: Fit a 2L 3-neck round bottom flask with an overhead paddle stirrer, a gas sparge tube (with fritted end to disperse gas) and a reflux condenser (with drying tube on top, filled with drying agent). Fit the bottom of the flask with a heating mantle and put 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (78.10g, 0.300 mol), xylenes (400mL) and absolute 2B ethanol (90mL) into the flask. Stir to dissolve all the solids at ambient temperature. Sparge hydrogen chloride from a lecture bottle (38.36g, 1.052 mol) into the stirred solution without external heating over a 15 minute period. Replace the sparge tube with a glass stopper and heat the solution by mantle, with stirring, at 40-79C for 45 minutes and 79C for 15 minutes. Replace the reflux condenser with a simple still head fitted with a thermometer and condenser. Collect 200 mL of distillate (80-138C at atmospheric pressure) and allow the remaining light yellow solution to cool to give a mixture of the title compound and xylenes.

35

Example

- 81 -

Step t: (4-Cyclopropanecarbonyl-phenyl)-acetic acid, ethyl ester and (3-Cyclopropanecarbonyl-phenyl)-acetic acid, ethyl ester

- 5 Dissolve the mixture of [4-(4-chloro-butyryl)-phenyl]-acetic acid, ethyl ester and [3-(4-chloro-butyryl)-phenyl]-acetic acid, ethyl ester (650g) in 2B ethanol (1250mL). Add, by dropwise addition, a solution of 2B ethanolic KOH (168g in 1000mL), keeping the temperature below 10C. After
10 the addition, stir magnetically for 5 hours at -10C. Bring the mixture to pH 6 with acetic acid (5mL) and filter through a celite™ pre-coat. Evaporate the solvent in vacuo to give the title compound as an oil (538g).

15

Example 12

Step d: [4-(4-Chloro-butyryl)-phenyl]-acetic acid, 2-ethylhexyl ester

- Mix 2-ethyl-1-hexanol (6.5g, 5mol), triethylamine (50.5g, 0.5mol) and methylene chloride (50mL). Add, by dropwise
20 addition, 2-phenylacetyl chloride (5mol) and warm to 50°C. Stir at room temperature overnight, filter and wash the filtercake with methylene chloride (50mL). Combine the organic phases and wash with 5% aqueous hydrochloric acid (50mL) and water. Dry (MgSO₄), evaporate the solvent in
25 *vacuo* and purify by distillation to give 2-phenylacetic acid, 2-(2-ethylhexyl) ester.

- Mix chlorobutyryl chloride (16.9g) and AlCl₃ (29.3g) at room temperature. Add 2-phenylacetic acid, 2-ethylhexyl ester
30 (27.6g), keeping the temperature below 10°C. Heat at 35°C for 24 hours, quench in ice water (200g). Separate the organic phase, dry (MgSO₄) and evaporate the solvent in *vacuo*. Dilute the residue with ethanol (150mL), add hydrogen chloride (5g) and heat to 75°C. After 2.5 hours,
35 add another 5g of hydrogen chloride and stir at 75°C for 24 hours. Evaporate the solvent in *vacuo* to give the title compound.

- 82 -

Example 13

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester and 2-[3-(4-Chloro-butyryl)-phenyl]-2-

5 methyl-propionic acid, ethyl ester

Place aluminum chloride (58.4g, 438mmol) and methylene chloride (100mL) in a 250 mL 3 neck flask equipped with a condenser, thermometer, and overhead stirrer. Cool to 10C and add, by dropwise addition, 4-chlorobutyryl chloride
10 (32.4g, 230mmol), keeping the temperature below 10C. Add, by dropwise addition, ethyl dimethylphenylacetate (40g, 208mmol) at 10C. After the addition, slowly warm the mixture to room temperature and then heat at reflux for 17 hours. Quench the reaction into ice (400g) and stir for 1
15 hour. Extract with methylene chloride (2X25mL), wash with water (25mL), 10% aqueous sodium carbonate (25mL) and water (25mL). Evaporate the solvent in vacuo to give a red oil (58.7g).

20 Dissolve the red oil (58.7g) in 2B ethanol (40mL) and place in a 250mL round bottom flask equipped with an overhead stirrer, condenser, thermometer and gas inlet tube. Add anhydrous HCl (3g, 80mmol) with vigorous stirring and heat to 70C for 1 hour. Evaporate the solvent in vacuo to give
25 the title compound as a yellow oil (59g).

Example 14

Step 1: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester

30 Dissolve a mixture of 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester and 2-[3-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester (59g) in 2B ethanol (100mL) and add, by dropwise addition, a solution of KOH (49.4g of 85%) in 2B ethanol (250mL),
35 keeping the temperature below 15C. After the addition, warm the reaction mixture to room temperature and stir magnetically for 1 hour. Bring to pH 6 with acetic acid and filter through a celite™ pre-coat. Evaporate the

solvent in vacuo to give a mixture of 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester as an oil (57.1g) Purify by one of the following methods:

Method A: Pack a 31/32 in. I.D. vacuum jacked and silvered column with 53 inches of 1 in. diameter, 316 stainless steel packing. For high temperature distillation, the column is fitted with an adiabatic jacket composed of an inner layer of 1 in. fiber glass wrapped with heat tape in an upper and lower zone and finally covered with 2 in. fiber glass insulation. The upper zone is heated at 135C and the lower zone at 185C. The magnetic reflux splitting head is controlled by a reflux timer and fitted with a standard thermometer for monitoring overhead temperature. Vacuum is supplied by a system composed of a pump protected by a dry ice trap and fitted with a McLeod gage for monitoring the overhead pressure. The 1L distillation pot is heated with an electric mantel at 65 volts, agitated magnetically and fitted with a mercury manometer for monitoring bottoms pressure, and a thermocouple for monitoring bottoms temperature.

The still pot is charged with 265 g each of m- and p-xylene and fitted with a rubber septum for sampling by syringe. The xylene mixture is heated at total reflux and atmosphere pressure with the temperature 135C at the head and 139C in the bottoms. Samples are withdrawn for analysis by collecting a few drops of distillate and extracting about 1mL from the pot. The still is sampled after 3 hours and again after 18 hours for calibration by GC and theoretical plate calculations using the Fenske correlation and a relative volatility, $\alpha=1.0209$.

Charge the mixture of 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-

- 84 -

cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (901.2g) to the still pot and heat at total reflux until the column has equilibrated. Take a forecut at 2:1
5 reflux ratio and increase the reflux ratio to 5:1 and the 2-(3-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester stripped. Cool and release vacuum and allow to sit overnight. Add bis(2-ethylhexyl)phthalate (dioctyl phthalate) (100mL) to the still pot and restart the still
10 as before. Once the still has equilibrated, collect mixed fractions of 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester at 10:1 reflux ratio. Once the overheads are free of 2-(3-
15 cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester by GC analysis, reduce the reflux ratio to 2:1 and collect the title compound.

Method B: Place crude mixture of 2-(4-
20 cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester (4872g) on a rotary evaporator and strip of volatiles to an end point of 85C, 15mm to give a brown oil (4006g). Charge a 3L round bottom three
25 neck flask equipped with magnetic stirbar, thermometer and distillation head with stripped crude mixture of 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester. Distill the oil at 0.5mm Hg
30 and discard a light fraction boiling at 25-130C (pot temp - 105-165C, 9.5g). Continue distilling the oil at 0.5mm Hg and collect a second fraction boiling at 130-150C (pot temperature 165-190, 3217g).

35 Place the crude flash distilled product (1000g) in a 4L Hastelloy™ reactor equipped with Camille™ control along with water (500mL) and ethanol 2B (2L). Heat the mixture to 40C while agitating at 400 rpm. Set the reactor jacket to cool

- 85 -

the contents at approximately 12C/hour to a final temperature of 0C after a clear solution is observed. Then set the jacket to cool the reactor contents at approximately 12C/hour to a final temperature of -15C and hold at that temperature for more than one hour. Filter the slurry, wash with cold (-15C) ethanol, cold heptanes (-15C) and dry to give a solid (507g). Purify by recrystallization from mixed heptanes as above to give the title compound (503g) after drying.

Example 15

Step h and step l: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester and 2-(3-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester

Method A: Place aluminum chloride (586g, 4.4moles) and methylene chloride (300mL) in a 2L 3-neck round bottom flask equipped with an overhead stirrer, dry ice condenser, and nitrogen atmosphere. Cool to 10C and add, by dropwise addition, chlorobutyryl chloride (338g, 2.4moles), keeping the temperature below 15C. After addition is complete, add, by dropwise addition, ethyl 2-methyl-2-phenylpropionate (384g, 2mol), keeping the temperature below 15C. After addition was complete, warm the reaction mixture to 22C and stir for 1 hour. Raise the temperature to 90C, stir for 90 minutes, cool to room temperature and slowly pour into a 6L stirred flask containing ice/water (4kg). Filter through a celite™ precoat, separate the organic phase and wash the aqueous phase with methylene chloride (50mL). Evaporate the solvent in vacuo to give a mixture of 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester and 2-[3-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester.

Dissolve the mixture of 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester and 2-[3-(4-chloro-

-86-

butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester in 2B ethanol (400mL) and place in a 3L 3-neck round bottom flask equipped with an overhead stirrer, gas inlet and reflux condenser. Add anhydrous HCl (50g) and stir the mixture at 70C for 1 hour. Cool the solution to 15C and add, by dropwise addition, aqueous 50% NaOH (260g), keeping the temperature below 15C. After the addition, stir the mixture an addition 1 hour at 22C. Add toluene (700mL) followed by acetic acid (2g) and then water (500mL). Separate the layers and evaporate the solvent in vacuo to give the title compound as a yellow oil (551g).

Method B: Place aluminum chloride (458g, 3.4mole) and methylene chloride (234mL) in a 2L 3neck round bottom flask equipped with an overhead stirrer, dry ice condenser and nitrogen atmosphere. Cool to 10C and add, by dropwise addition, 4-chlorobutyryl chloride (264g, 1.9mol), keeping the temperature below 15C. After addition is complete, add, by dropwise addition, ethyl 2-methyl-2-phenylpropionate (300g, 1.56mol), keeping the temperature below 15C. After the addition is complete, warm the reaction mixture to 24C and stir for 1 hour. Raise the temperature to 57C for 2 hours, cool to room temperature and slowly pour into a 6L stirred flask containing ice/water (3.1kg). Filter through a celite™ precoat and separate the phases. Evaporate the solvent in vacuo to give an oil.

Dissolve the oil in 2B ethanol (312mL) and place in a 3L 3 neck round bottom flask equiped with an overhead stirrer, gas inlet and reflux condenser. Add anhydrous HCl (39g) and stir the mixture at 70C for 1 hour. Cool to 50C and add, by dropwise addition, aqueous 20% NaOH (641g), keeping the temperature below 50C. After the addition, stir the mixture for an additional 1 hour at 50C, cool to room temperature and neutralize with acetic acid (6.25g).

- 87 -

Separate the layers and evaporate the solvent in vacuo to give the title compound (391g).

5

Example 16

Step h and step 1: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, 2-ethylhexyl ester

Mix methylene chloride (50mL), 2-ethylhexyl alcohol (130g, 1mol) and triethylamine (50g, 0.5mol). Add, by dropwise
10 addition, ethyl dimethylphenylacetyl chloride (91g, 0.5mol). Heat the reaction mixture to 68C for 1 hour, add methylene chloride (100mL) and stir overnight. Remove the solids by filtration, wash with methylene chloride (50mL), combine with the liquid organics, wash with aqueous 5% HCl,
15 (50mL), water (50mL) and dry over MgSO₄. Evaporate the solvent in vacuo and purify by distillation (119 C at 1mmHg) (105g, 76%).

Place aluminum chloride (29.3g) and methylene chloride
20 (30mL) in a 250mL round bottom flask with an overhead stirrer, temperature control, condenser, additional funnel and nitrogen atmosphere. Add, by dropwise addition, chlorobutyryl chloride (16.9g), keeping the temperature below 10C. After addition is complete, warm the reaction
25 mixture to 36C and hold for 24 hours. Quench the reaction mixture into ice/water (200g) and extract with methylene chloride (50mL). Wash the organics with water (50mL) and dry (MgSO₄). Evaporate the solvent in vacuo to give an oil (30g). Place the oil in a 250mL flask equipped with an
30 overhead stirrer, gas inlet, condenser and thermometer. Add 2B ethanol (150mL) followed by anhydrous HCl (5g). Heat the reaction mixture to 76C for 2.5 hours then add additional HCl (5g). Heat the reaction mixture at 76C for 22 hours, evaporate the solvent in vacuo to give an oil.
35 Dissolve the oil in 2B ethanol (100mL), treat with solid KOH (10g) and heat at reflux for 2 hours.

- 88 -

Example 17

Step m and step 1: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, ethyl ester

- 5 Dissolve 2-[4-(4-chloro-butyryl)-phenyl]-acetic acid, ethyl ester (28.5g) in toluene (50mL) and evaporate the solvent *in vacuo* to remove traces of ethanol. Dissolve the residue in diglyme (50mL) and add, by dropwise addition, to a suspension of sodium hydride (12.2g of a 60% suspension in
10 mineral oil) slurried in diglyme (150mL) containing methyl chloride (10g). Slowly add methyl chloride (10g) and stir for 15 minutes. Filter through filter aid, wash filtercake with acetonitrile and evaporate the solvent. Remove meta-isomer by distillation (150°C @ 1mm) and crystallize
15 (ethanol) to give the title compound (93%).

Example 18

Step f and step : 2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid, ethyl ester and 2-(3-Cyclopropanecarbonyl-phenyl)-propionic acid, ethyl ester

- 20 Dissolve 2-phenylpropionic acid (30g) in 2B ethanol (100mL) and add anhydrous HCl (10g). Allow to sit for 48-72 hours, evaporate the solvent *in vacuo* and purify by distillation to give ethyl 2-phenylpropionate (31g); bp 100C at 6mm.

25

- Place aluminum chloride (49.4g, 0.371mole) and methylene chloride (50mL) in a 250mL 3-neck round bottom flask equipped with an overhead stirrer, addition funnel and thermometer. Cool to less than 10C and add, by dropwise
30 addition, chlorobutyrylchloride (23.8g, 0.202mol), keeping the temperature below 10C. After addition is complete, add, by dropwise addition, ethyl 2-phenylpropionate (30g, 0.17mol), keeping the temperature below 10C. Stir at room temperature for 1 hour then heat at reflux for 14 hours.
35 Quench into ice/water (350g) and filter through a celite™ pre-coat. Separate the layers and evaporate the solvent *in vacuo* to give a red oil.

- 89 -

Dissolve the red oil in 2B ethanol (35mL) and place in a round bottom flask with a condenser and gas inlet. Add anhydrous HCl (4.3g) and heat the solution to 70C for 1 hour. Cool the solution to 10C and add, by dropwise addition, 20% aqueous sodium hydroxide. Separate the layers and evaporate the solvent in vacuo to give an oil.

Re-treat the oil with HCl in 2B ethanol as above, cool to 10C and treat with a 20% solution of sodium ethoxide in ethanol. Neutralize with acetic acid, filter the solids and evaporate the solvent in vacuo. Purify by distillation to give the title compound; bp 161-167 at 1.2mm.

15

Example 19

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester and 2-[3-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester

Place AlCl₃ (146.5g, 1.1mol) and methylene chloride (75mL) in a 3-neck, 500mL round-bottomed flask equipped with an overhead stirrer, bottom drop valve, thermometer, condenser and temperature control and cool to 15°C. Add, by dropwise addition, 4-chlorobutyryl chloride (84.5g, 0.6mol), keeping the temperature below 15°C. Add, by dropwise addition, ethyl 2-methyl-2-phenylpropionate (96g, 0.5mol), keeping the temperature below 15° C. After addition is complete, stir the reaction mixture at 22°C for 1 hour, then heat at reflux (57°C) for 2 hours. Add the reaction mixture, by dropwise addition, by way of the bottom drop valve, to water (500mL) at 95°C contained in a 2L 3 neck flask equipped with a magnetic stirbar, thermometer and distillation head. During addition, hold the reaction mixture at 70°C by allowing the methylene chloride to distill overhead. After the quench is complete, separate the the organic layer, dry (MgSO₄) and evaporate the solvent in vacuo to give the title compound (150g).

The novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is CONR_6R_7 may also be prepared as described in Scheme G. In Scheme G, all substituents are as previously defined unless otherwise indicated.

Scheme G

10


15

20

25

30

35



- 91 -

Scheme G provides alternative various general synthetic procedures for preparing the novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is $CONR_6R_7$.

In step a, the appropriate phenylacetic acid amide compound of structure (37) is methylated to give the corresponding α -methylphenylacetic acid amide compound of structure (38) as described previously in Scheme A, step a.

Appropriate phenylacetic acid amide compound of structure (37) are prepared from the corresponding phenylacetic acid by standard amide-forming reactions as are known in the art. The appropriate phenylacetic acids may be prepared by hydrolysis of the corresponding 2-cyano-2-propylbenzene compound of structure (27) by techniques and procedures well known and appreciated by one of ordinary skill in the art.

20

In step b, the appropriate α -methylphenylacetic acid amide compound of structure (38) is methylated to give the corresponding α,α -dimethylphenylacetic acid amide compound of structure (39) as described previously in Scheme A, step a.

25

Appropriate α -methylphenylacetic acid amide compound of structure (38) are prepared from the corresponding α -methylphenylacetic acid by standard amide-forming reactions as are known in the art as as described in step a.

30

In step c, the appropriate phenylacetic acid amide compound of structure (37) is dimethylated to give the corresponding α,α -dimethylphenylacetic acid amide compound of structure (39) as described previously in Scheme A, step c.

35

In step d, the appropriate phenylacetic acid amide compound of structure (37) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) as described previously in Scheme A, step d.

10 In step e, the appropriate phenylacetic acid amide compound of structure (37) is acylated with an appropriate cyclopropyl compound of the structure



wherein B is as previously defined to give the corresponding cyclopropylketo-phenylacetic acid amide compound of structure (45) as described previously in Scheme A, step e.

In step f, the appropriate α -methylphenylacetic acid amide compound of structure (38) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) as described previously in Scheme A, step d.

In step g, the appropriate α -methylphenylacetic acid amide compound of structure (38) is acylated with an appropriate cyclopropyl compound of the structure



- 93 -

wherein B is as previously defined to give the corresponding cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) as described previously in
 5 Scheme A, step e.


In step h, the appropriate α,α -dimethylphenylacetic acid amide compound of structure (39) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I
 10 and n is as previously defined to give the corresponding ω' -halo- α' -keto- α,α -di-methylphenylacetic acid amide compound of structure (40) as described previously in Scheme A, step d.

15

Appropriate α,α -dimethylphenylacetic acid amide compound of structure (39) are prepared from the corresponding α,α -dimethylphenylacetic acid by standard amide-forming reactions as are known in the art as as
 20 described in step a.

In step i, the appropriate α,α -dimethylphenylacetic acid amide compound of structure (39) is acylated with an appropriate cyclopropyl compound of the structure

25



wherein B is as previously defined to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in
 30 Scheme A, step e.

35

In step j, the appropriate ω' -halo- α' -keto- α,α -methylphenylacetic acid amide compound of structure (42) is methylated to give the corresponding ω' -halo- α' -keto- α,α -

di-methylphenylacetic acid amide compound of structure (40) as described previously in Scheme a, step a.

5 In step k, the cyclopropyl functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) is ring-opened to give the corresponding ω' -halo- α' -keto- α,α -di-methylphenylacetic acid amide compound of structure (40) wherein $n = 3$ as
10 described previously in Scheme A, step j.

 In step l, the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound of structure (40) wherein $n = 3$ is ring-closed to give the corresponding
15 cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in Scheme A, step k.

 In step m, the appropriate ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) is dimethylated to
20 give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound of structure (40) as described previously in Scheme A, step c.

25 In step n, the appropriate ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) is methylated to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) as described previously in Scheme A, step a.

30

 In step o, the cyclopropyl functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) is ring-opened to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid
35 amide compound of structure (42) wherein $n = 3$ as described previously in Scheme A, step j.

- 95 -

In step p, the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) wherein $n = 3$ is ring-closed to give the corresponding
5 cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) as described previously in Scheme A, step k.

In step q, the appropriate cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) is
10 methylated to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in Scheme A, step a.

In step r, the appropriate cyclopropylketo-phenylacetic
15 acid amide compound of structure (45) is dimethylated to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in Scheme A, step c.

20 In step s, the cyclopropyl functionality of the appropriate cyclopropylketo-phenylacetic acid amide compound of structure (45) is ring-opened to give the corresponding ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) wherein $n = 3$ as described
25 previously in Scheme A, step j.

In step t, the appropriate ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) wherein $n = 3$ is ring-closed to give the corresponding cyclopropylketo-
30 phenylacetic acid amide compound of structure (45) as described previously in Scheme A, step k.

In step u, the appropriate cyclopropylketo-phenylacetic acid amide compound of structure (45) is methylated to give
35 the corresponding cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) as described previously in Scheme A, step a.

Starting materials for use in Scheme G are readily available to one of ordinary skill in the art.

5 The following examples present typical syntheses as described in Scheme G. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to
10 grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

15

Example 20

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, N-methoxy-N-methylamide

Dissolve 2-methyl-2-phenyl-propionic acid (15.0g, 91.2mmol)
20 in toluene (80mL) and add, by dropwise addition over 5 minutes, thionyl chloride (15mL, 206mmol). Stir at room temperature overnight, add additional thionyl chloride (3mL, 41.1mmol) and heat to reflux for 1 hour. Remove excess thionyl chloride by azeotropic distillation with
25 toluene (40mL). Add toluene (20mL) to the reaction mixture along with a solution of potassium carbonate (28.0g, 203mmol) in water (40mL). Add, by dropwise addition, a solution of N,O-dimethylhydroxylamine hydrochloride (8.9g, 91.2mmol) in water (20mL) without cooling and stir for 2
30 hours. Add tert-butylmethyl ether (75mL) followed by slow addition of aqueous HCl (2N, 75mL) with vigorous stirring. Separate the organic layer and wash with aqueous HCl (2N, 75mL), saturated sodium hydrogen carbonate (25mL) and brine (50mL). Dry the organic layer over (Na₂SO₄), filter,
35 evaporate the filtrate *in vacuo* and purify by vacuum distillation to give 2-methyl-2-phenyl-propionic acid, N-methoxy-N-methylamide (18.0g, 95%); bp 91-103°C/5mm Hg.

- 97 -

MS (CI, CH₄) m/e 208 (M⁺+1, 100), 119.

Slurry AlCl₃ (10.15g, 76.1mmol) and methylene chloride
5 (45mL) under a nitrogen atmosphere at room temperature.
Add 4-chlorobutyryl chloride (4.27mL, 38.1mmol), stir for
20 minutes and add, by dropwise addition over 10 minutes, a
solution of 2-methyl-2-phenyl-propionic acid, N-methoxy-N-
methylamide (6.58g, 31.7mmol) in methylene chloride (15mL).
10 Stir at room temperature for 45 minutes, then heat at 30-
35°C for 7 hours. Pour into ice water (150mL) and separate
the layers. Wash the aqueous layer with water (3X75mL),
combine the aqueous layers and extract with methylene
chloride (2X75mL). Combine the organic layers and dry
15 (Na₂SO₄). Filter, evaporate the filtrate *in vacuo* and purify
by silica gel chromatography (3:1 hexane/ethyl acetate) to
give the title compound (6.19g, 63%) as a light yellow oil.

MS (CI, CH₄) m/e 312 (M⁺+1), 276.

20

Example 21

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, dimethylamide

Dissolve 2-methyl-2-phenyl-propionic acid (15.0g, 91.2mmol)
25 in toluene (80mL) and add, by dropwise addition over 5
minutes, thionyl chloride (15mL, 206mmol). Stir at room
temperature overnight, add additional thionyl chloride
(3mL, 41.1mmol) and heat to reflux for 1 hour. Remove
excess thionyl chloride by azeotropic distillation with
30 toluene (40mL). Add toluene (20mL) to the reaction mixture
along with a solution of potassium carbonate (28.0g,
203mmol) in water (40mL). Add, by dropwise addition, a 40%
aqueous solution of dimethylamine hydrochloride (20mL,
0.18mol) without cooling and stir for 2 hours. Add tert-
35 butylmethyl ether (75mL) following by slow addition of
aqueous HCl (2N, 75mL) with vigorous stirring. Separate
the organic layer and wash with aqueous HCl (2N, 75mL),
saturated sodium hydrogen carbonate (25mL) and brine

- 98 -

(50mL). Dry the organic layer over (Na_2SO_4), filter, evaporate the filtrate *in vacuo* and purify by crystallization to give 2-methyl-2-phenyl-propionic acid, dimethylamide
5 (15.35g, 88%) as a white solid; mp 57-59°C.

Anal. Calcd for $\text{C}_{12}\text{H}_{17}\text{NO}$: C, 75.35; H, 8.96; N, 7.32;
Found: C, 75.12; H, 8.86; N, 7.26.

10 Add AlCl_3 (1.12g, 8.40mmol) to carbon tetrachloride (6mL) under a nitrogen atmosphere at room temperature. Add 4-chlorobutyryl chloride (0.49mL, 4.37mmol), stir for 15 minutes and add, by dropwise addition over 3 minutes, a solution of 2-methyl-2-phenyl-propionic acid, dimethylamide
15 (0.64g, 3.36mmol) in carbon tetrachloride (6mL). Stir at room temperature for 17 hours, dilute with methylene chloride (10mL), pour into ice water (50mL) and separate the layers. Wash the aqueous layer with methylene chloride (2X70mL), 5% aqueous sodium hydrogen carbonate, combine the
20 organic layers and dry (Na_2SO_4). Filter, evaporate the filtrate *in vacuo* and purify by silica gel chromatography (5:2 hexane/ethyl acetate) to give the title compound (0.72g, 72%) as a light yellow oil.

25

Example 22

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, pyrrolidineamide

Dissolve 2-methyl-2-phenyl-propionic acid (15.0g, 91.2mmol) in toluene (80mL) and add, by dropwise addition over 5
30 minutes, thionyl chloride (15mL, 206mmol). Stir at room temperature overnight, add additional thionyl chloride (3mL, 41.1mmol) and heat to reflux for 1 hour. Remove excess thionyl chloride by azeotropic distillation with toluene (40mL). Add toluene (20mL) to the reaction mixture
35 along with a solution of potassium carbonate (28.0g, 203mmol) in water (40mL). Add, by dropwise addition, pyrrolidine (7.61mL, 91mmol) without cooling and stir for 2 hours. Add tert-butylmethyl ether (75mL) following by slow

- 99 -

addition of aqueous HCl (2N, 75mL) with vigorous stirring. Separate the organic layer and wash with aqueous HCl (2N, 75mL), saturated sodium hydrogen carbonate (25mL) and brine
 5 (50mL). Dry the organic layer over (Na₂SO₄), filter, evaporate the filtrate *in vacuo* and purify by crystallization to give 2-methyl-2-phenyl-propionic acid, pyrrolidineamide (18.28g, 92%) as a solid; mp 96-97°C.

10 Anal. Calcd for C₁₄H₁₉NO: C, 77.38; H, 8.81; N, 6.45; Found: C, 77.21; H, 8.70; N, 6.41.

Add AlCl₃ (8.31g, 62.3mmol) to carbon tetrachloride (65mL) under a nitrogen atmosphere at room temperature. Add 4-
 15 chlorobutyryl chloride (03.5mL, 31.2mmol), stir for 15 minutes and add, by dropwise addition over 15 minutes, a solution of 2-methyl-2-phenyl-propionic acid, pyrrolidineamide (5.64g, 26.0mmol) in carbon tetrachloride (60mL). Stir at room temperature for 17 hours, pour into
 20 ice water (100mL) and separate the layers. Wash the aqueous layer with methylene chloride (2X70mL), 5% aqueous sodium hydrogen carbonate, combine the organic layers and dry (Na₂SO₄). Filter, evaporate the filtrate *in vacuo* and purify by silica gel chromatography (5:2 hexane/ethyl
 25 acetate) to give the title compound (6.55g, 78%) as a light yellow oil.

Example 23

Step 1: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-
 30 propionic acid, N-methoxy-N-methylamide

Add potassium hydroxide (13g) to 2-[4-(4-chloro-butyryl-phenyl)-2-methyl-propionamide, N-methoxy-N-methylamide (96.6mmol) and stir at room temperature for 40 minutes, filter and wash the filtercake with ethanol. Evaporate the
 35 ethanol in vacuo, dissolve in methylene chloride (100mL), wash with water (50mL), 5% sodium hydrogen carbonate (50mL) and water (50mL). Evaporate the solvent in vacuo, removing water with toluene azeotrope. Purify the product by

- 100 -

distillation followed by recrystallization (heptane) to give the title compound (7.4g).

5 The following compounds can be prepared by procedures depicted in Scheme G:

(4-cyclopropanecarbonyl-phenyl)-acetic acid, N-methoxy-N-methylamide;

10

(4-cyclopropanecarbonyl-phenyl)-acetic acid, dimethylamide;

(4-cyclopropanecarbonyl-phenyl)-acetic acid, pyrrolidineamide;

15

2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid, N-methoxy-N-methylamide;

2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid,

20 dimethylamide;

2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid, pyrrolidineamide;

25 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, N-methoxy-N-methylamide;

2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, dimethylamide;

30

2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid, pyrrolidineamide;

35 [4-(4-Chloro-butyryl)-phenyl]-acetic acid, N-methoxy-N-methylamide;

[4-(4-Chloro-butyryl)-phenyl]-acetic acid, dimethylamide;

- 101 -

[4-(4-Chloro-butyryl)-phenyl]-acetic acid, pyrroldineamide;

2-[4-(4-Chloro-butyryl)-phenyl]-propionic acid, N-methoxy-
5 N-methylamide;

2-[4-(4-Chloro-butyryl)-phenyl]-propionic acid,
dimethylamide;

10 2-[4-(4-Chloro-butyryl)-phenyl]-propionic acid,
pyrroldineamide;

2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, N-
methoxy-N-methylamide;
15

2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid,
dimethylamide;


2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid,
20 pyrroldineamide;

25


30

35

Scheme H




Scheme H Cont.



3c

35

Scheme H Cont.



31

Scheme H provides various general synthetic procedures for preparing the novel intermediates of formula (II), formula (III), formula (IV), formula (V), formula (VI) and formula (VII) wherein R_5 is COOH , COOalkyl or CONR_6R_7 .

In step a, the nitrile functionality of the appropriate ω -halo-cyanocumylketone compound of structure (19) is

converted to the corresponding ester by reaction with an appropriate C₁ to C₆ alcohol to give the corresponding ω'-halo-α'-keto-α,α-dimethylphenylacetic acid ester compound
5 of structure (31).

For example, the ω'-halo-α'-keto-α,α-dimethylphenylacetic acid ester compound of structure (31) may be prepared by reacting an appropriate ω-halo-
10 cyanocumylketone compound of structure (19) with an appropriate C₁-C₆ alcohol in the presence of a suitable anhydrous acid followed by treatment with water. Examples of appropriate alcohols are methanol, ethanol, propanol, and the like, with methanol being preferred. Examples of
15 appropriate acids are hydrogen chloride and hydrogen bromide, with hydrogen chloride being preferred. The reaction time varies from about 1/2 hour to 48 hours, preferably 3 to 5 hours and the reaction temperature varies from about -20°C to room temperature, preferably -10°C to
20 0°C. The ω'-halo-α'-keto-α,α-dimethylphenylacetic acid ester compound of structure (28) is recovered from the reaction zone by evaporation of the solvent followed by extraction as is known in the art. The ω'-halo-α'-keto-α,α-dimethylphenylacetic acid ester compound of structure
25 (31) may be purified by procedures well known in the art, such as chromatography.

In step b, the nitrile functionality of the appropriate ω-halo-cyanocumylketone compound of structure (19) is
30 converted to the corresponding amide to give the ω'-halo-α'-keto-α,α-dimethylphenylacetic acid amide compound of structure (40) wherein R₆ and R₇ are both hydrogen.

For example, hydrolysis may be achieved by using a
35 suitable acid, such as concentrated hydrochloric acid as is known in the art.

- 106 -

In step c, the carboxy ester functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31) is hydrolyzed to give the
5 corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

For example, hydrolysis may be achieved by using a suitable non-nucleophilic base, such as sodium methoxide in
10 methanol as is known in the art. Other methods known in the art for ester cleavage include potassium carbonate in methanol, methanolic ammonia, potassium carbonate, potassium hydroxide, calcium hydroxide, sodium hydroxide, magnesium hydroxide, sodium hydroxide/pyridine in methanol,
15 potassium cyanide in ethanol and sodium hydroxide in aqueous alcohols, with potassium hydroxide being preferred. The reaction is typically carried out in an aqueous lower alcohol solvent, such as methanol, ethanol, isopropyl alcohol, n-butanol, 2-ethoxyethanol or ethylene glycol or
20 pyridine, at temperatures ranging from room temperature to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 100 hours.

In step d, the carboxy functionality of the appropriate
25 ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure
30 (31).

For example, one such method involves reacting an appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) with an excess of an appropriate
35 C₁-C₆ alcohol which is straight or branched in the presence of a small amount of mineral acid, such as hydrochloric acid or sulfuric acid, hydrochloric acid being preferred, at reflux. Another suitable method involves reacting an

appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) with an excess of diazomethane in a suitable solvent such as ether at room temperature to
5 give the methyl ester. In addition, the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (28) may also be prepared by reacting an appropriate ω' -halo- α' -keto- α,α -di-methylphenylacetic acid compound of structure (46) with an excess of 2,2-dimethoxypropane in a
10 suitable solvent such as methanol at 0°C to room temperature to give the methyl ester. Another suitable method involves first reacting an appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) with thionyl chloride in a suitable solvent such as
15 methylene chloride to give an intermediate acid chloride, followed by addition of a suitable C₁ to C₆ alcohol which is straight or branched. Another suitable method involves the alkylation of the carboxylate anion with an appropriate electrophile, such as dimethyl sulfate or ethyl bromide, to
20 give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31). Such methods are well known in the art and are described in *J. Org. Chem.*, 29, 2490-2491 (1964).

25 Alternatively, step k and step d may be combined and the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (34) wherein n = 3 may be prepared from the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (50).

30 Alternatively, step p, step k and step d may be combined and the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31) wherein n = 3 may be prepared from the corresponding cyclopropyl cyanocumylketone compound of structure (20).

35

In step e, the nitrile functionality of the appropriate ω -halo-cyanocumylketone compound of structure (19) is converted to the corresponding carboxy to give the ω' -halo-

- 108 -

α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

5 For example, hydrolysis may be achieved by using a suitable acid, such as concentrated hydrochloric acid as is known in the art.

In step f, the amide functionality of the appropriate
10 ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound of structure (40) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

15 For example, hydrolysis may be achieved by using a suitable non-nucleophilic base, such as sodium methoxide in methanol as is known in the art. Other methods known in the art for ester cleavage include potassium carbonate in
20 methanol, methanolic ammonia, potassium carbonate, potassium hydroxide, calcium hydroxide, sodium hydroxide, magnesium hydroxide, sodium hydroxide/pyridine in methanol, potassium cyanide in ethanol and sodium hydroxide in aqueous alcohols, with potassium hydroxide being preferred. The
25 reaction is typically carried out in an aqueous lower alcohol solvent, such as methanol, ethanol, isopropyl alcohol, n-butanol, 2-ethoxyethanol or ethylene glycol or pyridine, at temperatures ranging from room temperature to the reflux temperature of the solvent, and the reaction
30 time varies from about 1/2 hour to 100 hours.

In step g, the carboxy functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) may be amidated by techniques and procedures
35 well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound of structure (40).

- 109 -

In step h, the α -halo functionality of the appropriate ω -halo-halocumylketone compound of structure (10) is carboxylated to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

For example, a solution of the appropriate ω -halo-halocumylketone compound of structure (10) and a suitable catalyst, such as tetraethylammonium bromide, in a suitable polar aprotic organic solvent, such as acetonitrile, N,N-dimethylacetamide, 1-methyl-2-pyrrolidinone or dimethylformamide, are placed in a jacketed glass cell and fitted with an expanded silver mesh cathode, magnesium anode and carbon dioxide delivery tube. Rotation of the electrodes provides stirring, while electrical contact with the electrodes is made via spring loaded sliding carbon brushes placed against the concentric metal shafts (insulated from each other with a length of plastic tubing) onto which the electrodes are mounted. Carbon dioxide is introduced into the cell at pressures of 1-10 atm, for a period of time ranging from 30 minutes to 50 hours and at a temperature range of from -30°C to 50°C . The corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) is isolated, after acidification with a suitable mineral acid, such as hydrochloric acid, by extractive methods as are known in the art.

It is preferred that the ω -halo functionality of the appropriate ω -halo-halocumylketone compound of structure (10) for use in step h be a ω -chloro.

Alternatively, the treatment of appropriate ω -halo-halocumylketone compound of structure (10) with a transition metal catalyst such as palladium, nickel or cobalt, optionally in the presence of a phosphine catalysis using low to modest pressures of carbon monoxide as described by Stahly et al. in U.S. Patent 4,990,658, 1991

- 110 -

also provides the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

5 In step i, the appropriate the amide functionality of the appropriate ω' -halo- α' -keto- α,α -di-methylphenylacetic acid amide compound of structure (40) is converted to the corresponding ester to give the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31).

10

For example, the appropriate ω' -halo- α' -keto- α,α -di-methylphenylacetic acid amide compound of structure (40) is reacted with an appropriate hydrogen halide in an appropriate organic solvent such as ethanol. The reaction
15 is typically conducted at a temperature range of from room temperature to reflux and for a period of time ranging from 5 minutes to 1 hour. The ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31) is recovered from the reaction zone by extractive methods
20 as is known in the art.

In step j, the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) wherein $n = 3$ is ring-closed to give the corresponding
25 cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in Scheme A, step k.

In step k, the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) is
30 ring-opened to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) wherein $n = 3$ as described previously in Scheme A, step j.

In step l, the nitrile functionality of the appropriate
35 cyclopropyl cyanocumylketone compound of structure (20) is converted to the corresponding ester by reaction with an appropriate C_1 to C_6 alcohol to give the cyclopropylketo-

- 111 -

α,α -dimethylphenylacetic acid ester compound of structure (32) as described previously in step a.

5 In step m, the nitrile functionality of the appropriate cyclopropyl cyanocumylketone compound of structure (20) is converted to the corresponding amide to give the ω' -halo- α' -keto- α,α -di-methylphenylacetic acid amide compound of structure (41) wherein R_6 and R_7 are both hydrogen as
10 described previously in step b.

 In step n, the carboxy ester functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure (32) is hydrolyzed to give the
15 corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step c.

 In step o, the carboxy functionality of the appropriate
20 cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid ester compound of structure
25 (32) as described previously in step d.

 In step p, the nitrile functionality of the appropriate cyclopropyl cyanocumylketone compound of structure (20) is converted to the corresponding carboxy to give the
30 cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step e.

 In step q, the amide functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid amide
35 compound of structure (41) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the corresponding cyclopropylketo- α,α -

- 112 -

dimethylphenylacetic acid compound of structure (47) as described previously in step f.

5 In addition, step q and step k may be combined and the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) wherein $n = 3$ may be prepared from the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in
10 Scheme A, step j.

 In step r, the carboxy functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) may be amidated by techniques and procedures
15 well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) as described previously in step g.

20 In step s, the α -halo functionality of the appropriate cyclopropyl halocumylketone compound of structure (11) is carboxylated to give the corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step h.

25 In step t, the appropriate the amide functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetic acid amide compound of structure (41) is converted to the corresponding ester to give the cyclopropylketo- α,α -
30 dimethylphenylacetic acid ester compound of structure (32) as described previously in step i.

 In step u, the nitrile functionality of the appropriate ω -halo-cyanoethylphenylketone compound of structure (21) is
35 converted to the corresponding ester by reaction with an appropriate C_1 to C_6 alcohol to give the ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) as described previously in step a.

- 113 -

In step v, the nitrile functionality of the appropriate ω -halo-cyanoethylphenylketone compound of structure (21) is converted to the corresponding amide to give the ω' -halo-
5 α' -keto- α -methylphenylacetic acid amide compound of structure (42) wherein R_6 and R_7 are both hydrogen as described previously in step b.

In step w, the carboxy ester functionality of the
10 appropriate ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) is hydrolyzed to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) as described previously in step c.

15

In step x, the carboxy functionality of the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) may be esterified by techniques and procedures well known and appreciated by one of ordinary
20 skill in the art to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) as described previously in step d.

Alternatively, step ee and step x may be combined and
25 the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (33) wherein $n = 3$ may be prepared from the corresponding cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in step d.

30

Alternatively, step jj, step ee and step x may be combined and the ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (33) wherein $n = 3$ may be prepared from the corresponding cyclopropyl
35 cyanoethylphenylketone compound of structure (23) as described previously in step d.

- 114 -

In step y, the nitrile functionality of the appropriate ω -halo-cyanoethylphenylketone compound of structure (21) is converted to the corresponding carboxy to give the ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) as described previously in step e.

In step z, the amide functionality of the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) as described previously in step f.

In step aa, the carboxy functionality of the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) may be amidated by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) as described previously in step g.

In step bb, the α -halo functionality of the appropriate ω -halo-haloethylphenylketone compound of structure (12) is carboxylated to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) as described previously in step h.

In step cc, the appropriate the amide functionality of the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid amide compound of structure (42) is converted to the corresponding ester to give the ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) as described previously in step i.

In step dd, the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) wherein $n = 3$ is ring-closed to give the corresponding

- 115 -

cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in Scheme A, step k.

5 In step ee, the appropriate cyclopropylketo- α -methylphenylacetic acid compound of structure (49) is ring-opened to give the corresponding ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) wherein $n = 3$ as described previously in Scheme A, step j.

10

 In step ff, the nitrile functionality of the appropriate cyclopropyl cyanoethylphenylketone compound of structure (23) is converted to the corresponding ester by reaction with an appropriate C_1 to C_6 alcohol to give the
15 cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) as described previously in step a.

 In step gg, the nitrile functionality of the appropriate cyclopropyl cyanoethylphenylketone compound of
20 structure (23) is converted to the corresponding amide to give the cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) wherein R_6 and R_7 are both hydrogen as described previously in step b.

25 In step hh, the carboxy ester functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) is hydrolyzed to give the corresponding cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in step
30 c.

 In step ii, the carboxy functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid compound of structure (49) may be esterified by techniques
35 and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo- α -methylphenylacetic acid ester compound of structure (35) as described previously in step d.

In step jj, the nitrile functionality of the appropriate cyclopropyl cyanoethylphenylketone compound of structure (23) is converted to the corresponding carboxy to give the cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in step e.

In step kk, the amide functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the corresponding cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in step f.

In addition, step kk and step ee may be combined and the ω' -halo- α' -keto- α -methylphenylacetic acid compound of structure (48) wherein $n = 3$ may be prepared from the corresponding cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) as described previously in Scheme A, step j.

In step ll, the carboxy functionality of the appropriate cyclopropylketo- α -methylphenylacetic acid compound of structure (49) may be amidated by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo- α -methylphenylacetic acid amide compound of structure (44) as described previously in step g.

In step mm, the α -halo functionality of the appropriate cyclopropyl haloethylphenylketone compound of structure (14) is carboxylated to give the corresponding cyclopropylketo- α -methylphenylacetic acid compound of structure (49) as described previously in step h.

In step nn, the appropriate the amide functionality of the appropriate ω' -halo- α' -keto- α -methylphenylacetic acid

- 117 -

amide compound of structure (42) is converted to the corresponding ester to give the ω' -halo- α' -keto- α -methylphenylacetic acid ester compound of structure (33) as described previously in step i.

In step oo, the nitrile functionality of the appropriate ω -halo cyanotolylketone compound of structure (22) is converted to the corresponding ester by reaction with an appropriate C_1 to C_6 alcohol to give the ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) as described previously in step a.

In step pp, the nitrile functionality of the appropriate ω -halo cyanotolylketone compound of structure (22) is converted to the corresponding amide to give the ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) wherein R_6 and R_7 are both hydrogen as described previously in step b.

20

In step qq, the carboxy ester functionality of the appropriate ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) is hydrolyzed to give the corresponding ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) as described previously in step c.

In step rr, the carboxy functionality of the appropriate ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) as described previously in step d.

35

Alternatively, step yy and step rr may be combined and the ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) wherein $n = 3$ may be prepared from the

- 118 -

corresponding ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) as described previously in step d.

5

Alternatively, step ddd, step yy and step rr may be combined the ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) wherein $n = 3$ may be prepared from the corresponding cyclopropyl cyanotolylketone compound of structure (24) as described previously in step d.

10 In step ss, the nitrile functionality of the appropriate ω -halo cyanotolylketone compound of structure (22) is converted to the corresponding carboxy to give the ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) as described previously in step e.

20 In step tt, the amide functionality of the appropriate ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) as described previously in step f.

25

In step uu, the carboxy functionality of the appropriate ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) may be amidated by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) as described previously in step g.

35 In step vv, the α -halo functionality of the appropriate ω -halo halotolylketone compound of structure (13) is carboxylated to give the corresponding ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) as described previously in step h.

In step ww, the appropriate the amide functionality of the appropriate ω' -halo- α' -keto-phenylacetic acid amide compound of structure (43) is converted to the corresponding ester to give the ω' -halo- α' -keto-phenylacetic acid ester compound of structure (34) as described previously in step i.

10 In step xx, the appropriate ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) wherein $n = 3$ is ring-closed to give the corresponding cyclopropylketo-phenylacetic acid compound of structure (51) as described previously in Scheme A, step k.

15 In step yy, the appropriate cyclopropylketo-phenylacetic acid compound of structure (51) is ring-opened to give the corresponding ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) wherein $n = 3$ as described previously in Scheme A, step j.

20 In step zz, the nitrile functionality of the appropriate cyclopropyl cyanotolylketone compound of structure (24) is converted to the corresponding ester by reaction with an appropriate C_1 to C_6 alcohol to give the cyclopropylketo-phenylacetic acid ester compound of structure (36) as described previously in step a.

30 In step aaa, the nitrile functionality of the appropriate cyclopropyl cyanotolylketone compound of structure (24) is converted to the corresponding amide to give the cyclopropylketo-phenylacetic acid amide compound of structure (45) wherein R_6 and R_7 are both hydrogen as described previously in step b.

35 In step bbb, the carboxy ester functionality of the appropriate cyclopropylketo-phenylacetic acid ester compound of structure (36) is hydrolyzed to give the

- 120 -

corresponding cyclopropylketo-phenylacetic acid compound of structure (51) as described previously in step c.

5 In step ccc, the carboxy functionality of the appropriate cyclopropylketo-phenylacetic acid compound of structure (51) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo-
10 phenylacetic acid ester compound of structure (36) as described previously in step d.

 In step ddd, the nitrile functionality of the appropriate cyclopropyl cyanotolylketone compound of
15 structure (24) is converted to the corresponding carboxy to give the cyclopropylketo-phenylacetic acid compound of structure (51) as described previously in step e.

 In step eee, the amide functionality of the appropriate
20 cyclopropylketo-phenylacetic acid amide compound of structure (45) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the corresponding cyclopropylketo-phenylacetic acid compound of structure (51) as described previously in step f.

25

 In addition, step yy and step eee may be combined and the ω' -halo- α' -keto-methylphenylacetic acid compound of structure (50) wherein $n = 3$ may be prepared from the corresponding cyclopropylketo-phenylacetic acid amide
30 compound of structure (45) as described previously in Scheme A, step j.

 In step fff, the carboxy functionality of the appropriate cyclopropylketo-phenylacetic acid compound of
35 structure (51) may be amidated by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding cyclopropylketo-phenylacetic

- 121 -

acid amide compound of structure (45) as described previously in step g.

5 In step ggg, the α -halo functionality of the appropriate cyclopropyl halotolylketone of structure (15) is carboxylated to give the corresponding cyclopropylketo-phenylacetic acid compound of structure (51) as described previously in step h.

10

In step hhh, the appropriate the amide functionality of the appropriate cyclopropylketo-phenylacetic acid amide compound of structure (45) is converted to the corresponding ester to give the cyclopropylketo-phenylacetic acid ester compound of structure (36) as described previously in step i.

15

Starting materials for use in Scheme H are readily available to one of ordinary skill in the art.

20

The following examples present typical syntheses as described in Scheme H. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

25

30

Example 24

Step a: 2-[4-(4-Chloro-buteryl)-phenyl]-2-methyl-propionic acid, methyl ester

35 Place anhydrous methanol (5mL) under argon, cool to 0°C and add hydrogen chloride until saturated. Add 2-[4-(4-chloro-buteryl)-phenyl]-2-methyl-propionitrile (103mg, 4.12mmol), remove the ice bath and stir for 5 hours at room temperature. Allow to stand at -10°C overnight, and stir

- 122 -

an additional 3 hours at room temperature. Pour into cracked ice (20g) and allow to stand for 5 minutes. Evaporate the solvent *in vacuo* to 1/2 volume, dilute with water and extract with methylene chloride (3X). Combine the organic layers, wash with saturated sodium hydrogen carbonate and brine. Dry (MgSO₄), filter and evaporate the solvent *in vacuo*. Extract the residue into hot hexane (12mL), filter hot and evaporate the solvent *in vacuo* to give the title compound as a colorless oil (97mg, 83%).

Example 25

Step d: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester

Add anhydrous hydrogen chloride gas (18.0g) to anhydrous ethanol DB (210g) by purging the solution. Add this hot solution (60°C) to a solution of 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid (31g, 115.6mmol) and reflux under a nitrogen atmosphere for 2.5 hours. Evaporate the solvent *in vacuo*, dissolve the residue in methylene chloride (150mL) and wash with water (2X100mL). Dry (MgSO₄), filter through silica gel, washing the gel with methylene chloride (250mL). Combine the organic washings and evaporate the solvent *in vacuo* to give the title compound as a colorless oil (33.3g, 97%).

¹H NMR (300MHz, CDCl₃) δ 7.96 (d, J=8.3Hz, 2H), 7.45 (d, J=8.3Hz, 2H), 4.15 (q, J=7.1Hz, 2H), 3.70 (t, J=6.6Hz, 2H), 3.19 (t, J=6.8Hz, 2H), 2.25 (p, J=6.6Hz, 2H), 1.61 (s, 6H), 1.20 (q, J=7.1Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 198.4, 176.0, 150.3, 135.1, 128.1, 126.0, 61.0, 46.8, 44.6, 35.2, 26.7, 26.3, 14.0; IR (neat) 2978, 1728, 1686, 1606, 1254, 1231, 1148, 1097 cm⁻¹.

Anal. Calcd for C₁₆H₂₁O₃Cl: C, 64.75; H, 7.13; Found: C, 64.24; H, 7.18.

Example 26

- 123 -

Step d: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, methyl ester

Dissolve 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid (6.2g, 23.1mmol) in hot methanolic solution of anhydrous hydrogen chloride (42mL of a methanol containing 3.2g of anhydrous hydrogen chloride). Reflux for 42 minutes, evaporate the solvent *in vacuo*, dissolve the residue in methylene chloride and wash with water. Dry (MgSO₄), filter through silica gel, washing the gel with methylene chloride. Combine the organic washings and evaporate the solvent *in vacuo* to give the title compound as a clear oil (6.21g, 94%).

¹H NMR (30MHz, CDCl₃) δ 7.95 (d, J=8.5Hz, 2H), 7.44 (d, J=8.5Hz, 2H), 3.66 (s, 3H), 3.67 (t, J=6.6Hz, 2H), 3.17 (t, J=6.6Hz, 2H), 2.30 (p, J=6.6Hz, 2H), 1.61 (s, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 198.0, 176.2, 149.8, 135.0, 128.0, 125.8, 52.4, 46.9, 44.7, 35.3, 26.8, 26.5.

20

Anal. Calcd for C₁₅H₁₉O₃Cl: C, 63.72; H, 6.77;
Found: C, 63.50; H, 6.67.

Example 27

25 Step d: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, methyl ester

Mix 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic acid (10.0g, 37.3mmol) and anhydrous potassium carbonate (3.5g, 25.3mmol). Heat to 40°C in acetonitrile (100mL) and stir under a nitrogen atmosphere. Add dimethyl sulfate (13.3g, 105mmol) and reflux for 45 minutes. Evaporate the solvent *in vacuo*, dissolve the residue in ethyl acetate (50mL) and wash with water (4X50mL). Dry (MgSO₄), filter through silica gel and evaporate the solvent *in vacuo* to give the title compound (6.4g, 89%).

Example 28

- 124 -

Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid

Fit a jacketed glass cell of about 6L capacity with a rotating expanded silver mesh cathode/magnesium anode assembly, a carbon dioxide delivery tube, and a stainless steel thermocouple. Load the cell with acetonitrile (5.8L) and tetraethylammonium bromide (26g). Sparge with carbon dioxide and cool in cooling bath. When the contents of the cell reach -10°C, add hydrogen chloride remediated 1-[4-(1-bromo-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one and 1-[4-(1-chloro-1-methyl-ethyl)-phenyl]-4-chloro-butan-1-one (424.9g, 53.5 mole % bromo and 20.4 mole % chloro by HPLC analysis, 1087 mmol total active tertiary benzylic halide) and perform electrolysis at a controlled current of 8 amps (20 mA cm⁻²) for 6 hours. Drain the contents, acidify with chilled aqueous 6M hydrochloric acid, extract, evaporate the solvent *in vacuo* and recrystallize to give the title compound (186g, 64%); 78.5-80.3°C.

20

¹H NMR (300MHz, CDCl₃) δ 10.5 (br s, 2H), 7.96 (d, J=8.2Hz, 2H), 7.50 (d, J=8.2Hz, 2H), 3.67 (t, J=6.8Hz, 2H), 3.17 (t, J=6.8Hz, 2H), 2.22 (m, J=6.7Hz, 2H), 1.63 (s, 6H); ¹³C NMR (75MHz, CDCl₃) δ 198.2, 181.9, 149.0, 135.2, 128.1, 126.1, 46.7, 44.7, 35.3, 26.9, 26.7; MS (CIMS (Methane)) 271 (3), 269 (11), 233 (100), 187 (75).

25

Anal. Calcd for C₁₄H₁₇O₃Cl: C, 62.57; H, 6.38; Found: C, 63.10; H, 6.59.

30

Example 29Step h: 2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid

Fit a jacketed glass cell of about 50mL capacity with an expanded silver mesh cathode (14 cm² geometric area), a roughly concentric magnesium sacrificial anode, a tube to deliver carbon dioxide gas and a magnetic stir bar. Add a solution of hydrogen chloride remediated 1-[4-(1-bromo-1-

35

- 125 -

methylethylphenyl-4-chlorobutan-1-one and 1-[4-(1-chloro-1-methylethyl)phenyl]-4-chlorobutan-1-one (2.79g, 89 mole %, 3:1 ratio of tertiary benzylic bromide to tertiary benzylic chloride by NMR, approximately 8.6mmol total active tertiary benzylic halide) in acetonitrile (45mL) and tetraethylammonium bromide (0.19g). Close the cell and cool to -10°C with a continuous carbon dioxide sparge for 169 minutes at an average current density of 13 mA cm⁻². Warm to contents of the cell to ambient temperature, drain the contents, acidify with chilled aqueous 6M hydrochloric acid, extract and evaporate the solvent *in vacuo* to give the title compound (1.53g, 66%).

15

Example 30Step h: 2-[4-(4-Chlorobutyryl)phenyl]-2-methylpropionic acid

Fit a jacketed glass cell of 50mL capacity with an expanded silver mesh cathode (14 cm² geometric area), a roughly concentric magnesium sacrificial anode, a tube to deliver carbon dioxide gas, and a magnetic stir bar. Cool the cell to -10°C under carbon dioxide. Add a solution of tetraethylammonium chloride (40mL of a 0.02M solution in dimethylformamide) and 1-[4-(1-chloro-1-methylethyl)phenyl]-4-chlorobutan-1-one (2.91g, 85% pure by NMR, 9.81mmol) and carry out electrolysis for 178 minutes at an average current density of 12.4 mA cm⁻²: the total charge passed is equal to 98% of the calculated theoretical two electron value. Warm the contents of the cell to ambient temperature, drain the contents, acidify with chilled aqueous 6M hydrochloric acid, extract and evaporate the solvent *in vacuo* to give the title compound (1.89g, 72%).

Example 31

35 Step m: 2-(4-Cyclopropanecarbonylphenyl)-2-methylpropionamide

Dissolve 2-(4-cyclopropanecarbonylphenyl)-2-methylpropionitrile (100mg) in aqueous ethanolic potassium

- 126 -

hydroxide (2mL) (prepared from ethanol (5mL), water (5mL) and solid potassium hydroxide (1.5g). Stir overnight at room temperature, then heat at reflux for 6 hours. Cool
 5 and evaporate the solvent *in vacuo* to give the title compound.

Example 32

Step t: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-
 10 propionic acid, ethyl ester

Dissolve 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-propionamide (100mg) in ethanol and bubble in hydrochloride gas for 5 minutes while stirring. Reflux for 10 hours, distill off the ethanol and extract into ethyl acetate.
 15 Evaporate the solvent *in vacuo* to give the title compound as an oil (50mg).

Example 33

Step k and step q: 2-[4-(4-Bromo-butyryl)-phenyl]-2-methyl-
 20 propionic acid

Treat 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-N-methyl-N-methoxy-propionamide (0.15g, 0.53mmol) with 48% HBr (1mL) for 2 hours at 80°C. Cool to room temperature, dilute with water (5mL) and neutralize with aqueous sodium hydrogen
 25 carbonate until pH 7. Extract with methylene chloride (3X15mL), dry (Na₂SO₄), filter and evaporate the solvent *in vacuo*. Purify by silica gel chromatography (3:1 hexane/ethyl acetate) to give the title compound (0.15g, 95%).

30

¹H NMR (CDCl₃) δ 7.97 (d, 2H), 7.51 (d, 2H), 3.53 (t, 2H), 3.16 (t, 2H), 2.30 (quin, 2H), 1.60 (s, 6H); ¹³C NMR (CDCl₃) δ 198.4, 181.8, 149.5, 131.0, 128.3, 126.3, 46.6, 36.5, 33.6, 26.9, 26.1; MS (CI) (M⁺+H) 303 (100), 315
 35 (98), 233 (80).

Example 34

- 127 -

Step p: 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid

Combine 2-(4-cyclopropanecarbonyl-phenyl)-2-methyl-
 5 propionitrile (0.5g) in 12.5% sodium hydroxide (20mL) and ethanol (12.5mL). Heat to reflux for 21 hours, cool and remove the ethanol by vacuum distillation. Extract the residual aqueous suspension with methylene chloride (40mL), acidify the aqueous phase with 20% HCl and extract with
 10 methylene chloride (2X40mL). Combine the organic phases, dry (Na₂SO₄) and evaporate the solvent *in vacuo* to give the title compound as a crystalline solid (350mg, 70%); mp 83-85°C.

15 ¹H NMR (CDCl₃) δ 7.50-8.00 (4H, d), 2.66 (1H, m), 1.62 (6H, s), 1.24 (2H, m), 1.04 (2H, m).

The following compounds can be prepared by using the procedures depicted in Scheme H:

20

(4-Cyclopropanecarbonyl-phenyl)-acetic acid;

2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid;

25 2-(4-Cyclopropanecarbonyl-phenyl)-2-methyl-propionic acid;

[4-(4-Chloro-butyryl)-phenyl]-acetic acid;

2-[4-(4-Chloro-butyryl)-phenyl]-propionic acid;

30

2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid;

(4-Cyclopropanecarbonyl-phenyl)-acetic acid, ethyl ester;

35 2-(4-Cyclopropanecarbonyl-phenyl)-propionic acid, ethyl ester;

[4-(4-Chloro-butyryl)-phenyl]-acetic acid, ethyl ester;

- 128 -

2-[4-(4-Chloro-butyryl)-phenyl]-propionic acid, ethyl ester;

5

2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionic acid, ethyl ester;

(4-Cyclopropanecarbonyl-phenyl)-acetamide;

10

2-(4-Cyclopropanecarbonyl-phenyl)-propionamide;

[4-(4-Chloro-butyryl)-phenyl]-acetamide;

15 2-[4-(4-Chloro-butyryl)-phenyl]-propionamide; and

2-[4-(4-Chloro-butyryl)-phenyl]-2-methyl-propionamide.


In addition, the novel intermediate of formula (II) wherein R₅ is COOH may be prepared as described in Scheme I. In Scheme I, all substituents are as previously defined unless otherwise indicated.

25

30

35

Scheme I



$D' = -C(=O)CH_3 \text{ or } -C(=O)C_6H_5$

- 130 -


Scheme I provides a general synthetic procedure for preparing the novel intermediate of formula (II) wherein R_5 is COOH.

5

In step a, the neophyl acetate of benzoate of structure (53) is acylated with an appropriate ω -halo compound of the structure $\text{Hal}-(\text{CH}_2)_n-\text{C}(=\text{O})-\text{B}$, wherein B is Hal or hydroxy, Hal is Cl, Br or I and n is as previously defined to give the corresponding ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) as described previously in Scheme A, step d.


The neophyl acetate of benzoate of structure (53) is prepared by reacting a methallyl halide of structure

20



wherein Hal is Cl, Br or I with sodium acetate or sodium benzoate in a suitable organic solvent such as 1-methyl-2-pyrrolidinone. The reactants are heated at a temperature of approximately 100 to 130°C and the corresponding to give the methallyl acetate or benzoate of structure

30




wherein D' is $-\text{C}(=\text{O})\text{CH}_3$ or $-\text{C}(=\text{O})\text{C}_6\text{H}_5$ which is collected by distillation.

35

A benzene compound of structure


- 131 -



5

wherein A is defined above is then alkylated with the methylallyl acetate or benzoate of structure

10




15

wherein D' is $-C(=O)CH_3$ or $-C(=O)C_6H_5$

to give the neophyl acetate or benzoate of structure (53) as described previously in Scheme A, step d.

20

In step a₂, the neophyl acetate or benzoate of structure (53) is acylated with an appropriate cyclopropyl compound of the structure



25

wherein B is as previously defined to give the corresponding cyclopropyl neophyl acetate or benzoate of structure (55) as described previously in Scheme A, step e.

30

In step b₁, the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) wherein n = 3 is ring-closed to give the corresponding cyclopropyl neophyl acetate or benzoate of structure (55) as described previously in Scheme A, step k.

35

In step b₂, the appropriate cyclopropyl neophyl acetate or benzoate of structure (55) is ring-opened to give the corresponding ω' -halo- α' -keto-(2-methylpropanol)benzene

- 132 -

acetate or benzoate compound of structure (54) wherein $n = 3$ as described previously in Scheme H, step j.

5 In step c_1 , the acetate or benzoate functionality of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) is hydrolyzed with concentrated hydrochloric acid in ethanol at reflux temperature for a period of time ranging from 1-
10 10 hours. The corresponding ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) is recovered from the reaction zone by extractive methods as is known in the art.

15 In step c_2 , the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) wherein $n = 3$ is ring closed and the acetate or benzoate functionality hydrolyzed with base to give the cyclopropyl neophyl alcohol compound of structure (57).

20 For example, the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) wherein $n = 3$ is reacted with 40% aqueous tetrabutylammonium hydroxide and 50% aqueous sodium
25 hydroxide at reflux temperature for a period of time ranging from 5-72 hours. The cyclopropyl neophyl alcohol compound of structure (57) may be recovered from the reaction zone by extractive methods as are known in the art.

30 In step c_3 , the acetate or benzoate functionality of the appropriate cyclopropyl neophyl acetate or benzoate of structure (55) is hydrolyzed to give the corresponding cyclopropyl neophyl alcohol of structure (57).

35 For example, the appropriate cyclopropyl neophyl acetate or benzoate of structure (55) is reacted with 50% aqueous sodium hydroxide at reflux temperature for a period

of time ranging from 5 minutes to 5 hours. The corresponding cyclopropyl neophyl alcohol of structure (57) is recovered from the reaction zone by extractive methods as are known in the art.

In step d₁, the ω' -halo- α' -keto-(2-methylpropanol)benzene acetate or benzoate compound of structure (54) is converted to the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

For example, the appropriate cyclopropyl neophyl alcohol of structure (54) may be reacted with ruthenium chloride/sodium periodate in a suitable organic solvent such as acetonitrile and/or carbon tetrachloride, ruthenium chloride/sodium hypochloride in a suitable solvent such as acetic acid/water, potassium permanganate in a suitable solvent such as acetic acid/water, fuming nitric acid in acetic acid or sodium nitrite/concentrated nitric acid in acetic acid. The reactants are typically mixed stirred together at a temperature range of 10°C to 50°C and for a period of time ranging from 30 minutes to 10 hours. The corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (46) is recovered from the reaction zone by extractive methods as is known in the art.

In step d₂, the ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) is converted to the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46).

For example, the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) may be oxidized with potassium permanganate in suitable acid solvent such as acetic acid. The reactants are typically reacted at a temperature range of from about 0°C to 5°C for a period of time ranging from 30 minutes to 10 hours.

- 134 -
- 134 -

The corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) is recovered from the reaction zone by extractive methods as are known in the art and may be purified by recrystallization. Other oxidizing reagents suitable for the oxidation of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) to the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) are nitric acid, chromium (IV) oxide, nitrogen dioxide, ruthenium (VIII) oxide, nickel peroxide, silver oxide, t-butyl chromate, xenic acid

In step d₃, the hydroxymethyl functionality of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) is oxidized with a variety of oxidizing agents and methods to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58).

20

One such method involves a procedure in which the hydroxymethyl functionality of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) is oxidized to the corresponding aldehyde functionality using, for example, Swern Oxidation conditions (dimethyl sulfoxide, oxalyl chloride and triethylamine), as is known in the art. The Swern Oxidation is carried out in a suitable aprotic organic solvent such as methylene chloride at temperatures ranging from about -78°C to room temperature, and the reaction time varies from about 1/2 hours to 8 hours. Other suitable reagents for the oxidation of the hydroxyethyl functionality of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) to the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) are Dess-Martin reagent, chromium (IV) oxide, nickel peroxide, sodium dichromate, potassium dichromate, t-butyl chromate, silver oxide, argentic picolinate, manganese

35

- 135 -

dioxide, lead tetraacetate, dicyclohexylcarbodiimide, 2,3-dichloro-5,6-dicyanoquinone, tetrachloro-1,2-benzoquinone, 2,2,6,6-tetramethylpiperidiny-1-oxyl (TEMPO) or quinolinium
5 chlorochromate.

In step d₄, the hydroxymethyl functionality of the appropriate cyclopropyl neophyl alcohol of structure (57) is oxidized to give the corresponding cyclopropylketo- α,α -
10 dimethylphenylacetaldehyde compound of structure (59) as described previously in step d₃.

In step d₅, the appropriate cyclopropyl neophyl alcohol of structure (57) is converted to the corresponding
15 cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step d₂.

In step d₆, the appropriate cyclopropyl neophyl acetate or benzoate of structure (55) is converted to the
20 corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step d₁.

In step e₁, the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) wherein n
25 = 3 is ring-closed to give the corresponding cyclopropyl neophyl alcohol of structure (57) as described previously in Scheme H, step j.

In step e₂, the appropriate cyclopropyl neophyl alcohol of structure (57) is ring-opened to give the corresponding
30 ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (56) wherein n = 3 as described previously in Scheme H, step k.

35

In step f₁, the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) wherein n = 3 is ring-closed to give the corresponding

cyclopropylketo- α,α -dimethylphenylacetaldehyde compound of structure (59) as described previously in Scheme H, step j.

5 In step f_2 , the appropriate cyclopropylketo- α,α -dimethylphenylacetaldehyde compound of structure (59) is ring-opened to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) wherein $n = 3$ as described previously in Scheme H, step k.

10

 In step g_1 , the aldehyde functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) is oxidized to give the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid
15 compound of structure (46).

 For example, the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) is reacted with, for example, potassium permanganate. The
20 potassium permanganate oxidation is carried out in a suitable acidic medium such as hydrochloric acid/acetone at a temperature ranging from about 0°C to room temperature and the reaction time varies from about 1/2 hour to 8 hours. Other suitable reagents for the oxidation of the
25 ω' -halo- α' -keto- α,α -dimethylphenylacetaldehyde compound of structure (58) to the corresponding ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) are chromium (IV) oxide, silver (I) oxide, silver oxide, argentic picolinate, peroxide, nitric acid, m-
30 chloroperbenzoic acid and peracetic acid.

 In step g_2 , the aldehyde functionality of the appropriate cyclopropylketo- α,α -dimethylphenylacetaldehyde compound of structure (59) is oxidized to give the
35 corresponding cyclopropylketo- α,α -dimethylphenylacetic acid compound of structure (47) as described previously in step g_1 .

- 137 -

Starting materials for use in Scheme I are readily available to one of ordinary skill in the art.

5 The following examples present typical syntheses as described in Scheme I. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to
10 grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

15

Example 35

Step a₁: 2-(4-(4-Chloro-1-oxo-butyl))-phenyl-2-methyl
propanyl acetate

20 Mix 1-methyl-2-pyrrolidinone (400mL), sodium acetate (205g, 2.5mol), stir at heat to 100°C in a reaction flask which is fitted with a distillation head. Add, by dropwise addition, methylallyl chloride (181g, 2.0mol) over 1 hour. Heat the pot to 120°C for 30 minutes collect methallyl
25 acetate by distillation (193g).

Mix methallyl acetate (228g, 2.0mol) and benzene (1L) and cool to 5°C. Add aluminum chloride (266g, 2.0mol) over approximately 30 minutes while maintaining the temperature
30 below 10°C. Add, in portions of 50mL to 80mL each, to a 5°C mixture of aluminum chloride (15g) in benzene (600mL). After addition is complete, stir at 0-3°C for 1/2 hour, pour onto ice (2kg) and separate the organic layer. Wash with water (2X300mL), dry (Na₂SO₄), and distill to give
35 neophyl acetate.

Dissolve neophyl acetate (150g, 0.78mol) in methylene chloride (390mL) and cool to 5°C. Add anhydrous aluminum

- 138 -

chloride (104g, 0.78mol) at such a rate that the temperature is maintained below 10°C. Cool the reaction mixture to 5°C. Dissolve anhydrous aluminum chloride (122g) in methylene chloride (390mL) and cool to 5°C. Add 4-chlorobutyryl chloride (132g, 0.94mol) at such a rate that the temperature is kept below 10°C. Cool the reaction to 5°C and add the neophyl acetate-aluminum chloride solution in one portion and stir between -5°C and 5°C for 19 hours. Pour slowly over crushed ice (1.5kg), separate the organic phase and wash with water (3X300mL), cold aqueous potassium carbonate (10%, 300mL) and water (300mL). Evaporate the solvent *in vacuo* and filter to give the title compound as a light-brown oil (221.1g, 95.6%).

15

¹H NMR (300MHz, CDCl₃) δ 1.34 (6H, s), 1.95 (3H, s), 2.18 (2H, quent.), 3.13 (2H, t), 3.65 (2H, t), 4.12 (2H, s), 7.43, 7.90 (2H each, d).

20

Example 36

Step b₁: 2-(4-(1-Oxo-1-cyclopropanyl)-phenyl-2-methylpropanyl acetate

Mix 2-(4-(4-chloro-1-oxo-butyl))-phenyl-2-methyl propanyl acetate (37.0g, 0.125mol), tetrabutylammonium hydroxide (8.1g of a 40% aqueous solution), methylene chloride (300mL) and 50% sodium hydroxide (40mL). Stir vigorously at room temperature for 4 hours, add water (100mL) and separate the organic layer. Wash with water (2X100mL), dry (MgSO₄) and evaporate the solvent *in vacuo* to give the title compound (29.9g).

¹H NMR (300MHz, CDCl₃) δ 1.00, 1.19 (2H each, m), 1.34 (6H, s), 1.95 (3H, s), 2.65 (1H, m), 4.13 (2H, s), 7.44, 7.95 (2H each, d).

35

Example 37

- 139 -

Step c₁: 2-(4-(4-Chloro-1-oxobutyl))-phenyl-2-methylpropanol
 Mix 2-(4-(4-chloro-1-oxo-butyl))-phenyl-2-methyl propanyl
 acetate, concentrated hydrochloric acid (555mL), and
 5 ethanol (2.5L) and reflux for 2.5 hours under a nitrogen
 atmosphere. Evaporate the solvent *in vacuo* and take the
 residue up in methylene chloride (1L). Wash sequentially
 with water (2X400mL), aqueous potassium carbonate (10%,
 200mL) and water (300mL). Evaporate the solvent *in vacuo* to
 10 give the title compound as a light-brown oil (200g, 90%).

¹H NMR (300MHz, CDCl₃) δ 1.35 (6H, s), 2.21 (2H, quent.)
 3.15, (2H, t), 3.64 (2H, s), 3.66 (2H, s), 7.48, 7.93 (2H
 each, d).
 15

Example 38

Step c₂: 2-(4-(1-Oxo-1-cyclopropanyl))-phenyl-2-
methylpropanol
 20 Mix 2-(4-(4-chloro-1-oxobutyl))-phenyl-2-methylpropanol
 (101g, 0.34mol), methylene chloride (800mL), 40% aqueous
 solution of tetrabutylammonium hydroxide (33g), and 50%
 aqueous solution of sodium hydroxide (162mL) and reflux for
 48 hours. Add water (300mL), separate the organic phase
 25 and wash with water (2X300mL). Dry (MgSO₄) and evaporate
 the solvent *in vacuo* to give the title compound as a light-
 brown oil (71.1g, 96%).

Example 39

30
Step c₃: 2-(4-(1-Oxo-1-cyclopropanyl))-phenyl-2-
methylpropanol
 Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanyl
 acetate (4.16g, 14mmol), ethanol (50mL) and water (5mL).
 35 Add 50% aqueous sodium hydroxide (4.48mL, 56mmol). Stir
 and heat at reflux for 30 minutes then remove the ethanol *in*
vacuo. Extract the aqueous residue with methylene chloride
 (2X25mL), wash with water (2X25mL), dry (MgSO₄) and

- 140 -

evaporate the solvent *in vacuo* to give the title compound as a brown oil (2.91g, 95.3%).

- 5 ^1H NMR (300MHz, CDCl_3) δ 1.03, 1.20 (2H each, m), 1.35 (6H, s), 1.70 (1H, t, br), 2.66 (1H, m), 3.64 (2H, d), 7.48, 7.98 (2H each, d).

Example 40

10

Step d₂: 2-(4-(4-Chloro-2-oxo-butyl))-phenyl-2-methylpropionic acid

- Mix powdered potassium permanganate (39.5g, 0.25mol), water (34mL) and acetic acid (200mL). Stir and cool at 0°C, then
15 add 85% phosphoric acid (4.2g). Stir vigorously and add 2-(4-(4-chloro-1-oxo-butyl))-phenyl-2-methylpropanol (24.5g, 0.1mol) in acetic acid (50mL) at such a rate as to keep the temperature below 5°C. Stir for 5.5 hours below 5°C, add
20 ice water (300mL), then sodium metabisulfite (45g) in small portions until the dark brown mixture becomes colorless. Extract the aqueous solution with methylene chloride (3X150mL), wash with water (100mL) then extract with 20% aqueous potassium carbonate (2X150mL). Wash the aqueous phase with methylene chloride (50mL), cool in an ice-bath
25 and acidify carefully with concentrated hydrochloric acid until pH 3. Extract with methylene chloride (2X150mL), wash with water (2X80mL) and dry (MgSO_4). Evaporate the solvent *in vacuo* to give the title compound as a crystalline solid (21.25g).

30

- ^1H NMR (300MHz, CDCl_3) δ 1.63 (6H, s), 2.22 (2H, quent.), 3.17 (2H, t), 3.67 (2H, t), 7.50, 7.92 (2H each, d), 12.3 (1H, s, br).

35

Example 41

Step d₅: 2-(4-(1-Oxo-1-cyclopropanyl))-phenyl-2-methylpropionic acid

Method A:

Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanol
5 (1.46g, 6.7mmol), ruthenium chloride (0.036g, 0.17mmol),
acetonitrile (14mL), carbon tetrachloride (14mL) and water
(20mL). Stir vigorously and add sodium periodate (5.85g)
in one portion. Stir at room temperature for one hour
longer, partition between methylene chloride (20mL) and
10 water (5mL), separate the organic layer, extract the
aqueous layer with methylene chloride (15mL) and wash the
combined methylene chloride layers with water (15mL) and
extract with 20% aqueous potassium carbonate (2X25mL).
Cool the base solution in an ice-bath, acidify carefully
15 with concentrated hydrochloride acid to pH 3 and extract
into methylene chloride (2X30mL). Wash with water (15mL),
dry (MgSO₄) and evaporate the solvent *in vacuo* to give the
title compound as a yellow oil (1.41g, 90%).

20 ¹H NMR (300MHz, CDCl₃) δ 1.04, 1.23 (2H each, d), 1.63 (6H,
s), 2.65 (1H, m), 7.50, 7.99 (2H each, d).

Method B:

Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanol
25 (10.9g, 50mmol), ruthenium chloride (0.032g, 0.16mmol),
acetic acid (100ml) and water (25mL). Cool to 10°C and
add, by dropwise addition, an aqueous solution of sodium
hypochloride (70ml), stirring vigorously over a 30-minute
period. Stir below 10°C for 30 minutes longer, evaporate
30 most of the solvent *in vacuo* and take the residue up in
methylene chloride (120mL). Wash the methylene chloride
solution with water (2X40mL) and extract with 20% aqueous
potassium carbonate (2X50mL). Cool the base solution in an
ice-bath, acidify carefully with concentrated hydrochloride
35 acid to pH 3 and extract into methylene chloride (2X50mL).
Wash the organic layer with water (40mL), dry (MgSO₄) and
evaporate the solvent *in vacuo* to give the title compound as
a light-yellow oil (5.46g, 47%).

Method C:

Mix potassium permanganate (3.61g, 22.8mmol), water (2mL)
5 and acetic acid (10mL). Stir and cool to 10°C and add 85%
phosphoric acid (500mg). Add, by dropwise addition, a
solution 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-
methylpropanol (1.66g, 7.6mmol) in acetic acid (5mL) over 5
minutes. Stir below 10°C for 1 hour and then at room
10 temperature for 5 hours. Add water (20mL) followed by
addition of Na₂S₂O₅ in small portions until the solution
becomes colorless. Extract with methylene chloride
(2X50mL), wash the methylene chloride solution with water
(30mL) and then extract with 10% aqueous potassium
15 carbonate (2X50mL). Cool the base solution in an ice-bath,
acidify carefully with concentrated hydrochloric acid to
pH 3 and extract with methylene chloride (2X50mL). Wash
the organic layer with water (20mL), dry (MgSO₄) and
evaporate the solvent *in vacuo* to give the title compound as
20 a colorless needles (1.20g, 68%).

¹H NMR (300MHz, CDCl₃) δ 1.00 (4H, d), 1.50 (6H, s), 7.49,
8.00 (2H each, d), 12.6 (1H, s, br).

25 Method D:

Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanol
(2.30g, 10.6mmol), acetic acid (5.5mL) and fuming nitric
acid (6.5mL). Stir and heat at 48-50°C for 2 hours, cool
and add ice water (20mL) followed by methylene chloride
30 (60mL). Separate the organic layer, wash with water
(2X20mL) and extract into 10% aqueous potassium carbonate
(2X40mL). Wash the alkaline solution with methylene
chloride (10mL) and cool in an ice-bath. Acidify carefully
with concentrated hydrochloric acid to pH 3, extract with
35 methylene chloride (2X40mL), wash the combined organic
layers with water (20mL), dry (MgSO₄) and evaporate the
solvent *in vacuo* to give the title compound as light-yellow
needles (1.89g, 77%).

- 143 -

Method E:

Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanol
5 (2.26g, 10.4mmol), sodium nitrite (60mg), acetic acid (5mL)
and concentrated nitric acid (6mL, d=1.42, 70%, 94mmol).
Stir and heat at 48-50°C for 2 hours, cool and dilute with
ice water (20mL). Extract into methylene chloride
(2X30mL), wash the combined organic layers with water
10 (2X20mL) and extract into 10% aqueous potassium carbonate
(2X40mL). Wash the alkaline solution with methylene
chloride (10mL) and cool in an ice-bath. Acidify carefully
with concentrated hydrochloric acid to pH 3 and extract
into methylene chloride (2X40mL). Wash the combined
15 organic layers with water (20mL), dry (MgSO₄) and evaporate
the solvent *in vacuo* to give the title compound as light
yellow needles (2.01g, 83%).

Example 42


20

Step d₆: 2-(4-(1-Oxo-1-cyclopropanyl))-phenyl-2-
methylpropionic acid

Mix 2-(4-(1-oxo-1-cyclopropanyl))-phenyl-2-methylpropanyl
acetate (5.0g, 0.0197mol), sodium nitrite (100mg), acetic
25 acid (10mL) and concentrated nitric acid (8.7mL, d=1.42,
70%, 0.137mol). Stir and heat at 48-50°C for 5.5 hours,
cool and dilute with ice water (40mL). Extract into
methylene chloride (2X70mL), wash the combined methylene
chloride extracts with water (2X50mL) and reduce the volume
30 to 50mL *in vacuo*. Extract with 10% aqueous potassium
carbonate (2X50mL), wash the base solution with methylene
chloride (20mL) and cool in an ice-bath. Acidify carefully
with concentrated hydrochloric acid to pH 3 and extract
into methylene chloride (2X60mL). Wash the combined
35 methylene chloride extracts with water (30mL), dry (MgSO₄)
and evaporate the solvent *in vacuo* to give the title compound
as a crystalline solid (4.12g, 90%).

The novel intermediates of formula (X) wherein R_5 is H, Br, Cl, I, CN, $-\text{COOH}$, $-\text{COOalkyl}$ or $-\text{CONR}_6\text{R}_7$ may be prepared as described in Scheme J. In Scheme J, all substituents 5 are as previously defined unless otherwise indicated.

Scheme J



35 Scheme J provides various general synthetic procedures for preparing the novel intermediates of formula (X) wherein R_5 is H, Br, Cl, I, CN, $-\text{COOH}$, $-\text{COOalkyl}$ or $-\text{CONR}_6\text{R}_7$.

In step a, the ketone functionality of the appropriate ω -halo-halocumylketone compound of structure (10) is
5 reduced to give the corresponding ω -halo-halocumylalcohol compound of structure (60).

For example, reduction of the appropriate ω -halo-halocumylketone compound of structure (10), using, for
10 example, a suitable reducing agent such as sodium borohydride, potassium borohydride, sodium cyanoborohydride, or tetramethylammonium borohydride is carried out in lower alcohol solvents, such as, methanol, ethanol, isopropyl alcohol or n-butanol at temperatures
15 ranging from about 0°C to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 8 hours. Other suitable reducing agents are, for example, lithium tri-tert-butylaluminumhydride and diisobutylaluminum hydride. These reduction reactions are
20 carried out in suitable solvents diethyl ether, tetrahydrofuran or dioxane at temperatures ranging from about 0°C to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 8 hours.

25 Catalytic reduction may also be employed in the preparation of appropriate ω -halo-halocumylalcohol compound of structure (60) from an appropriate ω -halo-halocumylketone compound of structure (10), using hydrogen gas in the presence of a suitable catalyst such as Raney
30 nickel, palladium, platinum or rhodium catalysts in lower alcohol solvents, such as, methanol, ethanol, isopropyl alcohol or n-butanol or acetic acid or their aqueous mixtures, or by the use of aluminum isopropoxide in isopropyl alcohol.

35

In addition, a chiral reduction of the appropriate ω -halo-halocumylketone compound of structure (10), using, for example, (+)-8-chlorodiisopinocampheylborane gives the

- 146 -

corresponding (R)- ω -halo-halocumylalcohol compound of structure (60) and (-)-B-chlorodiisopinocampheylborane gives the corresponding (S)- ω -halo-halocumylalcohol compound of structure (60). Other suitable chiral reducing agents are, (R) and (S)-oxazaborolidine/BH₃, potassium 9-O-(1,2:5,6-di-O-isopropylidene- α -D-glucofuranosyl)-9-boratabicyclo[3.3.1]nonane, (R) and (S)-B-3-pinanyl-9-borabicyclo[3.3.1]nonane, NB-Enantride, Lithium (R)-(+ and (S)-(-)-2,2'-dihydroxy-1,1'-binaphthyl alkoxyl aluminum hydride, (R)-(+ and (S)-(-)-2,2'-dihydroxy-6,6'-dimethylbiphenyl borane-amine complex, tris([(1S,2S,5R)-2-isopropyl-5-methyl-cyclohex-1-yl)methyl]aluminum, [(1R,3R)-2,2-dimethylbicyclo[2.2.1]hept-3-yl)methyl]beryllium chloride, (R)-BINAP-ruthenium complex/H₂ and 6,6'-bis(diphenylphosphino)-3,3'-dimethoxy-2,2',4,4'-tetramethyl-1,1'-biphenyl.

In step b, the ketone functionality of the appropriate ω -halo-cyanocumylketone compound of structure (19) is reduced to give the corresponding ω -halo-cyanocumylalcohol compound of structure (61) as described previously in step a.

In step c, the ketone functionality of the appropriate ω -halo-cyanocumylketone compound of structure (8) is reduced to give the corresponding ω -halo-cyanocumylalcohol compound of structure (62) as described previously in step a.

30

In step d, the α -halo functionality of the appropriate ω -halo-halocumylalcohol compound of structure (60) is cyanated to give the corresponding ω -halo-cyanocumylalcohol compound of structure (61) as described previously in Scheme D, step a.

35

In step e, the appropriate ω -halo-cyanocumylalcohol compound of structure (62) is cyanated to give the

corresponding ω -halo-cyanocumylalcohol compound of structure (61) as described previously in Scheme D, step b.

5 In step f, the appropriate appropriate ω -halo-cyanocumylalcohol compound of structure (62) is halogenated to give the corresponding ω -halo-halocumylalcohol compound of structure (60) as described previously in Scheme B, step a.

10

 In step g, the α -halo functionality of the appropriate ω -halo-halocumylalcohol compound of structure (60) is converted to the corresponding carboxy to give the ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (64) as described previously in Scheme H, step h.

15

 In step h, the nitrile functionality of the appropriate ω -halo-cyanocumylalcohol compound of structure (61) is converted to the corresponding ester to give the ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid ester compound of structure (63) as described previously in Scheme H, step a.

20

 In step i, the nitrile functionality of the appropriate ω -halo-cyanocumylalcohol compound of structure (61) is converted to the corresponding acid to give the ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (64) as described previously in Scheme H, step e.

25

 In step j, the nitrile functionality of the appropriate ω -halo-cyanocumylalcohol compound of structure (61) is converted to the corresponding amide to give the ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid amide compound of structure (65) wherein R_6 and R_7 are each hydrogen as described previously in Scheme H, step b.

30

35

 In step k, the ketone functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid ester compound of structure (31) is reduced to give the

corresponding ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid ester compound of structure (63) as described previously in step a.

5

In step l, the ketone functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid compound of structure (46) is reduced to give the corresponding ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (64) as described previously in step a.

10

In step m, the ketone functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenylacetic acid amide compound of structure (40) is reduced to give the corresponding ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid amide compound of structure (65) as described previously in step a.

15

In step n, the carboxy ester functionality of the appropriate ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid ester compound of structure (63) is hydrolyzed to give the corresponding ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (64) as described previously in Scheme H, step c.

20

In step o, the carboxy functionality of the appropriate ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (64) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid ester compound of structure (63) as described previously in Scheme H, step d.

25

In step p, the carboxy functionality of the appropriate ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid compound of structure (65) may be amidated by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -halo- α' -

30

- 149 -


hydroxy- α,α -dimethylphenylacetic acid amide compound of structure (57) as described previously in Scheme H, step g.

5 In step q, the amide functionality of the appropriate ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic acid amide compound of structure (65) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the ω' -halo- α' -hydroxy- α,α -dimethylphenylacetic
10 acid compound of structure (64) as described previously in Scheme H, step f.

In addition, the novel intermediates of formula (X) wherein R_5 is $-\text{CH}_2\text{OD}$ may be prepared as described in Scheme
15 K. In Scheme K, all substituents are as previously defined unless otherwise indicated.

Scheme K Cont.

20



25

D = H, $-\text{C}(=\text{O})\text{CH}_3$, $-\text{C}(=\text{O})\text{C}_6\text{H}_5$,

30

35

- 150 -

In Scheme K, the ketone functionality of the appropriate ω' -halo- α' -keto-(2-methylpropanol)benzene compound of structure (60) is reduced to give the
5 corresponding ω' -halo- α' -hydroxy-(2-methylpropanol)benzene compound of structure (66) as described previously in Scheme J, step a.

10

15

20

25

30

35

- 151 -

The novel intermediates of formula (XI) wherein R_5 is hydrogen, CN, COOalkyl or CONR_6R_7 may be prepared as described in Scheme L. In Scheme L, all substituents are
5 as previously defined unless otherwise indicated.

10

15


20

25

30

35

Scheme L



Scheme L provides various general synthetic procedures for preparing the novel intermediates of formula (XI) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇.

In step a, the ω' -halo functionality of the appropriate ω' -halo- α' -keto- α,α -dimethylphenyl compound of structure (67) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ is alkylated with the appropriate piperidine compound of structure (68) to give the corresponding ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇.

10

For example, the ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ may be prepared by reacting the appropriate ω' -halo- α' -keto- α,α -dimethylphenyl compound of structure (67) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ with the appropriate piperidine compound of structure (68) in a suitable solvent preferably in the presence of a suitable non-nucleophilic base and optionally in the presence of a catalytic amount of an iodide source, such as potassium or sodium iodide. The reaction time varies from about 4 to 120 hours and the reaction temperature varies from about 70°C to the reflux temperature of the solvent. Suitable solvent for the alkylation reaction include alcohol solvents such as, methanol, ethanol, isopropyl alcohol, or n-butanol; ketone solvents, such as, cyclohexanone, methyl isobutyl ketone; hydrocarbon solvents, such as, benzene, toluene or xylene; halogenated hydrocarbons, such as, chlorobenzene or methylene chloride or dimethylformamide. Suitable non-nucleophilic bases for the alkylation reaction include inorganic bases, for example, sodium bicarbonate, potassium carbonate, or potassium bicarbonate or organic bases, such as, a trialkylamine, for example, triethylamine or pyridine, or an excess of an appropriate piperidine compound of structure (68) may be used.

For those piperidine compounds of structure (68), wherein R_1 is hydroxy, it is preferred that R_1 be

unprotected for utilization in the alkylation reaction of step a, but those hydroxy functionalities present in the piperidine compounds of structure (68), wherein R₁ is hydroxy may be protected with a suitable protecting group. The selection and utilization of suitable protecting groups for the piperidine compounds of structure (68), wherein R₁ is hydroxy is well known by one of ordinary skill in the art and is described in "Protective Groups in Organic Syntheses", Theodora W. Greene, Wiley (1981). For example, suitable protecting groups for those hydroxy functionalities present include ethers such as tetrahydrothiopyranyl, tetrahydrothiofuranyl, 2-(phenylselenyl)ethyl ether, o-nitrobenzyl ether, trimethylsilyl ether, isopropyldimethylsilyl ether, t-butylldimethylsilyl ether, t-butylldiphenylsilyl ether, tribenzylsilyl ether, triisopropylsilyl ether; and esters, such as acetate ester, isobutyrate ester, pivaloate ester, adamantate ester, benzoate ester, 2,4,6-trimethylbenzoate (mesitoate) ester, methyl carbonate, p-nitrophenyl carbonate, S-benzyl thiocarbonate and N-phenylcarbamate.

The piperidine compounds of structure (68) are readily available to one of ordinary skill in the art and are described in United States Patent No. 4,254,129, March 3, 1981, United States Patent No. 4,254,130, March 3, 1981, United States Patent No. 4,285,958, April 25, 1981 and United States Patent No. 4,550,116, Oct. 29, 1985. The piperidine compounds of structure (68) wherein R₁ and R₂ form a second bond between the carbon atoms bearing R₁ and R₂ may be prepared by dehydration of the corresponding compound wherein R₁ is hydroxy by procedures generally known in the art, such as refluxing in strongly acidic solution.

35

The piperidine compounds of structure (68) include the limitations provided for previously for piperidine derivatives of formula (I) and (XI) in that when R₁ and R₂

are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.

5

In step b, the ω' -halo functionality of the appropriate ω -halo- α' -hydroxy- α,α -dimethylphenyl compound of structure (70) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ is alkylated with the appropriate piperidine compound of structure (68) to give the corresponding ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compound of structure (71) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ as described previously in step a.

15

In step c, the ketone functionality of the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ is reduced to give the corresponding ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compound of structure (71) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇.

20

For example, reduction of the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇, using, for example, a suitable reducing agent such as sodium borohydride, potassium borohydride, sodium cyanoborohydride, or tetramethylammonium borohydride is carried out in lower alcohol solvents, such as, methanol, ethanol, isopropyl alcohol or n-butanol at temperatures ranging from about 0°C to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 8 hours. Other suitable reducing agents are, for example, lithium tri-tert-butylaluminumhydride and diisobutylaluminum hydride. These reduction reactions are carried out in suitable solvents diethyl ether, tetrahydrofuran or dioxane at temperatures ranging from about 0°C to the reflux temperature of the solvent, and the reaction time varies from about 1/2 hour to 8 hours.

30

35

Catalytic reduction may also be employed in the preparation of appropriate ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compound of structure (71) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ from an appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇, using hydrogen gas in the presence of a suitable catalyst such as Raney nickel, palladium, platinum or rhodium catalysts in lower alcohol solvents, such as, methanol, ethanol, isopropyl alcohol or n-butanol or acetic acid or their aqueous mixtures, or by the use of aluminum isopropoxide in isopropyl alcohol.

15

Reduction using sodium borohydride or potassium borohydride is preferred over catalytic reduction for those ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ and wherein R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 .

In addition, a chiral reduction of the appropriate ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇, using, for example, (+)-B-chlorodiisopinocampheylborane gives the corresponding (R)- ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇ and (-)-B-chlorodiisopinocampheylborane gives the corresponding (S)- ω' -piperidine- α' -keto- α,α -dimethylphenyl compound of structure (69) wherein R_5 is hydrogen, CN, COOalkyl or CONR₆R₇. Other suitable chiral reducing agents are, (R) and (S)-oxazaborolidine/BH₃, potassium 9-O-(1,2:5,6-di-O-isopropylidene- α -D-glucofuransoyl)-9-boratabicyclo[3.3.1]nonane, (R) and (S)-B-3-pinanyl-9-borabicyclo[3.3.1]nonane, NB-Enantride, Lithium (R)-(+)- and (S)-(-)-2,2'-dihydroxy-1,1'-binaphthyl alkoxyl aluminum hydride, (R)-(+)- and (S)-(-)-2,2'-

- 157 -

5 dihydroxy-6,6'-dimethylbiphenyl borane-amine complex,
tris[[(1S,2S,5R)-2-isopropyl-5-methyl-cyclohex-1-
yl)methyl]aluminum, [[(1R,3R)-2,2-
dimethylbicyclo[2.2.1]hept-3-yl)methyl]beryllium chloride,
(R)-BINAP-ruthenium complex/H₂ and 6,6'-
bis(diphenylphosphino)-3,3'-dimethoxy-2,2',4,4'-
tetramethyl-1,1'-biphenyl.

10 Starting materials for use in Scheme L are readily
available to one of ordinary skill in the art.

The following examples present typical syntheses as
described in Scheme K. These examples are understood to be
15 illustrative only and are not intended to limit the scope
of the present invention in any way. As used herein, the
following terms have the indicated meanings: "g" refers to
grams; "mmol" refers to millimoles; "mL" refers to
milliliters; "bp" refers to boiling point; "°C" refers to
20 degrees Celsius; "mm Hg" refers to millimeters of mercury;
"μL" refers to microliters; "μg" refers to micrograms; and
"μM" refers to micromolar.

25

30

35

- 158 -

Example 43

Step a: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -dimethylbenzeneacetic acid methyl ester
5 Mix methyl 4'-(4-chloro-1-oxobutyl)- α,α -dimethylbenzene acetate (0.335mol), α,α -diphenyl-4-piperidinemethanol (101.8g, 0.335mol), potassium hydrogen carbonate (83.8g, 0.838mol), potassium iodide (1.00g, 0.006mol), toluene
10 (600mL) and water (220mL). Stir at reflux for 72 hours, add toluene (200mL) and deionized water (100mL). Filter through filter aid while at 80°C and separate the organic phase. Dry (MgSO₄), filter and purify by chromatography to give the title compound.

15

Example 44

Step a: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -dimethylbenzeneacetic acid ethyl ester

20 Method A: Remove the still head from the reaction flask containing a solution of ethyl 4'-(4-chloro-1-oxobutyl)- α,α -dimethylbenzene acetate and xylenes obtained from Example 11, Method G and reattach a reflux condenser. At ambient temperature, add azacyclonol free base which has
25 been recrystallized from toluene (178.28g, 0.660mol) and stir at 175 RPM while heating by heating mantle. After the temperature of the reaction slurry reaches 137 (approximately 30 minutes), stir the reaction for 5.5 hours, maintaining the temperature between 137-144C.

30 Remove the heating mantle, add mixed xylenes (100mL) and allow the reaction slurry to cool to 64C. Increase the stirring rate to 300 RPM and add glacial acetic acid (15.17g, 0.253mol). Maintain the temperature at 64-69C for
35 1.9 hours by heating with mantle, cool from 64-60C over a period of 15 minutes; and from 60-50C over a period of 32 minutes; from 50-42C over a period of 33 minutes. Filter at 42C by suction through a 350 mL coarse sintered glass filter funnel and wash the filtercake with mixed xylenes

- 159 -

(200mL) at ambient temperature. Allow the filtrate to stand at ambient temperature overnight then place in a 1L flask. Add isopropanol (40mL) and attached an overhead paddle stirrer. With stirring at 150 RPM, slowly add 37% aqueous concentrated HCl at ambient temperature, adding 2.00g during the first 17 minutes, adding a total of 33.13 g of HCl over 245 minutes. After the slurry has been digested, collect the solids by suction filtration through a 350mL coarse sintered glass funnel and wash the filtercake with fresh xylenes (200mL) and then with n-heptane (100mL). Dry the filtercake under vacuum at 47C for 2.5 days to give the title compound as an off-white solid (141.17g, 81%).

15

Concentrate the filtrate by rotary evaporator to give a thick residue of solids and syrup (23.78g) Add acetone (68g) and agitate by swirling until the syrup dissolves or releases as a solid. Collect the solids by suction filtration through a medium sintered glass funnel, wash with fresh acetone (17g) and dry under vacuum to give the title compound as a light tan solid (3.75g).

Method B: Place the solution of ethyl 4'-(4-chloro-1-oxobutyl)- α,α -dimethylbenzene acetate and xylenes obtained from Example 11, Method G in a 1L, 3-neck round bottom flask and add azacyclonol free base recrystallied from toluene (192.2g, 0.719mol). Stir the resulting slurry by overhead stirrer and heat to 140C for 5.5 hours. Allow to cool to ambient temperature and add a mixture of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -dimethylbenzeneacetic acid ethyl ester hydrochloride (33.8g, 0.0318 mol) and azacyclonol hydrochloride (0.0534mol), slurried in mixed xylenes (100mL). Reheat the resulting slurry to 135C with stirring and then allow to cool slowly to ambient temperature.

- 160 -

Vacuum filter and wash the filtercake with xylenes. Dry the filtercake under vacuum to give a solid (122.4g). Concentrate the filtrate by rotary evaporator to a weight
5 of 486g and add, by dropwise addition, 91g (2.75g, 0.0753mol) of a solution of HCl gas (5.6g) in absolute 2B ethanol (180mL) at 70-80C over a 1.5 hour period. Cool slowly to 30C and filter by vacuum. Wash the filtercake with mixed xylenes and dry under vacuum at 50C to give the
10 title compound as a solid (49.1g).

To the filtrate from the second filtercake, add absolute 2B ethanol (100mL), heat to 50C and sparge gaseous HCl (about 5g) into the solution. Add additional mixed xylenes
15 (170mL) and absolute 2B ethanol (100mL) and heat to 70C. Sparge in additional HCl gas until the total HCl added is 10g (0.274mol). Cool to 50C and stir for 2 hours then cool to ambient temperature and stir overnight.

20 Distill a total of 240mL of ethanol and xylenes from the slurry under reduced pressure (80mm, with pot temperature from 50 to 70C). Cool to 30C over a 1 hour period and filter by vacuum. Wash the filtercake with toluene and dry under vacuum at 50C to give the title compound as a solid
25 (119.2g).

Method C: Place ethyl 4'-(4-chloro-1-oxobutyl)- α,α -dimethylbenzene acetate (15.00g, 49.53mmol), azacyclonol free base (29.66g, 49.53mmol) and mixed xylenes (60mL) in a
30 250mL 1-neck round bottom flask fitted with a magnetic stir bar and reflux condenser. Heat the reaction mixture to reflux over a period of 15 minutes and then continue at reflux for 5.5 hours. Cool to ambient temperature and then to ice/water bath temperature. Separate the solids from
35 the orange xylenes solution by suction filtration through a coarse sintered glass funnel, wash the filtercake with cold xylenes (25mL) and dry in a vacuum oven at 60C to give the title compound as an off-white solid (16.21g).

- 161 -

- Method D: Place azacyclonol free base (35.00g, 125.68mmol), ethyl 4'-(4-chloro-1-oxobutyl)- α,α -
- 5 dimethylbenzene acetate (17.30g, 57.13mmol) and mixed xylenes (60mL) into a 250mL round bottom flask. Heat to reflux by mantel in 13 minutes and stir by magnetic bar and heat at reflux for 6.3 hours. Remove the heat from the reaction flask and cool by ice/water bath. Filter the cold
- 10 reaction slurry by suction through a coarse sintered glass funnel and wash the filtercake with fresh mixed xylenes (40mL). Vacuum dry the filtercake at 40C overnight to give the title compound as a solid (17.87g).
- 15 Add concentrated 37% HCl (2.18g, 22.1mmol) to the filtrate, stirred by magnetic bar. Stir overnight at ambient temperature, filter through suction through a coarse sintered glass funnel and wash the filtercake with fresh mixed xylenes (35mL). Vacuum dry the filtercake at 50C to
- 20 give the title compound as a solid (8.23g).
- Add concentrated 37% HCl (6.42g, 65.2mmol) to the filtrate stirred by magnetic bar. Add mixed xylenes (70mL) and filter through a coarse sintered glass funnel, at ambient
- 25 temperature. Wash the filtercake with fresh mixed xylenes (50mL) and vacuum dry the filtercake to give the title compound as a solid (27.25g).
- Purify by recrystallization as follows: Mix the title
- 30 compound (15g), absolute 2B ethanol (45mL) and n-heptane (90mL) in a 500 mL round bottom flask with a magnetic stir bar. Heat at reflux with stirring for 30 minutes, cool by ice/water bath and collect the solids by suction filtration through a coarse sintered glass funnel. Wash the
- 35 filtercake with 3:1 n-heptane/ethanol (24mL) and dry under vacuum at 55C to give the title compound as a white solid.

Example 45

- 162 -

Step c: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]- α,α -dimethylbenzeneacetic acid

Add sodium borohydride (0.105g, 2.77mmol) to a solution of
 5 sodium hydroxide (0.053g, 1.33mmol) in deionized water
 (2mL) and add, by dropwise addition, to a solution of 4-[4-
 [4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -
 dimethylbenzeneacetic acid hydrochloride (0.70g, 1.31mmol)
 in ethanol (30mL). Stir at room temperature for 3.5 hours
 10 at pH 7-8. Evaporate the solvent *in vacuo* and stir the
 residue with methylene chloride (15mL) and deionized water
 (15mL). Dry (MgSO₄), acidify to pH 3 with gaseous hydrogen
 chloride and evaporate the solvent. Add ether with
 stirring, filter the white solid and wash with additional
 15 ether. Dry to give the title compound.

Example 46

Step c: (R)-4-[4-[4-(Hydroxydiphenylmethyl)-1-
 20 piperidinyl]-1-hydroxybutyl]- α,α -dimethylbenzeneacetic,
 ethyl ester

Dissolve (+)-B-chlorodiisopinocampheylborane (2.5g,
 7.8mmol) in anhydrous tetrahydrofuran (5mL). Add a
 solution of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-
 1-oxobutyl]- α,α -dimethylbenzeneacetic, ethyl ester (2g,
 25 3.54mmol) in anhydrous tetrahydrofuran (5mL). Stir at room
 temperature for 3 days and cool to 0°C. Add water (1mL)
 and 30% hydrogen peroxide (2mL) and stir for 20 minutes.
 Add methylene chloride (30mL) and wash with brine (30mL),
 then aqueous sodium hydrogen carbonate (30mL), then brine
 30 (30mL). Dry (MgSO₄), evaporate the solvent *in vacuo* and
 purify by chromatography to give the title compound.

Example 47

Step c: (S)-4-[4-[4-(Hydroxydiphenylmethyl)-1-
 35 piperidinyl]-1-hydroxybutyl]- α,α -dimethylbenzeneacetic
 acid, ethyl ester

Dissolve (-)-B-chlorodiisopinocampheylborane (2.5g,
 7.8mmol) in anhydrous tetrahydrofuran (5mL). Add a

solution of 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -dimethylbenzeneacetic acid, methyl ester (3.54mmol) in anhydrous tetrahydrofuran (5mL). Stir at
 5 room temperature for 3 days and cool to 0°C. Add water (1mL) and 30% hydrogen peroxide (2mL) and stir for 20 minutes. Add methylene chloride (30mL) and wash with brine (30mL), then aqueous sodium hydrogen carbonate (30mL), then brine (30mL). Dry (MgSO₄), evaporate the solvent *in vacuo*
 10 and purify by chromatography to give the title compound.

Example 48

Step a: N,N-Dimethyl-2-(4-{4-[4-hydroxy-diphenylmethyl)-piperidin-1-yl]-butyryl}-phenyl)-isobutyramide

15 Dissolve N,N-dimethyl-2-[4-(4-chlorobutyryl)-phenyl]-isobutyramide (1.00g, 3.38mmol) in xylene (3mL) and add α,α -diphenyl-4-piperidinemethanol (1.09g, 4.07mmol) and potassium hydrogen carbonate (0.68g, 6.76mmol) in water (2.5mL). Heat at 100°C for 20 hours, remove hot water by
 20 pipette, dilute with ethyl acetate (20mL) and wash with water (20mL). Cool the organic layer to room temperature, dry (MgSO₄), evaporate the solvent *in vacuo* and purify by silica gel chromatography (9:1 ethyl acetate/methanol) and recrystallize (ethyl acetate/hexane) to give the title
 25 compound (1.13g, 63%) as a crystalline solid; mp 135-137°C.

Example 49

Step c: N,N-Dimethyl-2-(4-{1-hydroxy-4-[4-hydroxy-diphenylmethyl)-piperidin-1-yl]-butyryl}-phenyl)-

30 isobutyramide

Dissolve N,N-dimethyl-2-(4-{4-[4-hydroxy-diphenylmethyl)-piperidin-1-yl]-butyryl}-phenyl)-isobutyramide (3.00g, 5.69mmol) in ethanol (30mL), cool using an ice/water bath and add sodium borohydride (0.87g, 23.04mmol) in
 35 tetrahydrofuran (10mL). Remove the cold bath and stir at room temperature for 2.5 hours. Add water (25mL) and ethyl acetate (25mL) and separate the layers. Extract the aqueous layer with ethyl acetate (20mL), dry (MgSO₄) and

- 164 -

evaporate the solvent *in vacuo* to give the title compound (3.06g, 100%) as a white foam; mp 166-169°C.

5 MS (CI, CH₄) m/e 529 (M⁺+1), 280, 183.

Anal. Calcd for C₃₄H₄₄N₂O₃•0.3H₂O: C, 77.24; H, 8.39; N, 5.30; Found: C, 76.99; H, 8.36; N, 5.17.

10

Example 50

Step a: N-Methoxy-N-methyl-2-(4-{4-[4-hydroxy-diphenylmethyl)-piperidin-1-yl]-butyryl}-phenyl)-isobutyramide

Dissolve N-methoxy-N-methyl-2-[4-(4-chlorobutyryl)-phenyl]-
15 isobutyramide (1.44g, 4.62mmol) in 2:1 xylene/water (5mL) and add α,α-diphenyl-4-piperidinemethanol (1.36g, 5.07mmol) and potassium hydrogen carbonate (0.93g, 9.24mmol). Heat at 108°C for 22 hours, remove hot water by pipette, cool to room temperature and stir for 2 days. Evaporate the
20 solvent *in vacuo* and purify by silica gel chromatography (10:1 ethyl acetate/methanol) and recrystallize (ethyl acetate) to give the title compound (1.77g, 71%) as a white crystalline solid; mp 159-160.5°C.

25 MS (CI, CH₄) m/e 543 (M⁺+1), 293, 250, 183.

Anal. Calcd for C₃₄H₄₂N₂O₄•0.3H₂O: C, 74.50; H, 7.83; N, 5.11; Found: C, 74.75; H, 7.96; N, 5.15.

30

Example 51

Step c: N-Methoxy-N-methyl-2-(4-{1-hydroxy-4-[4-hydroxy-diphenylmethyl)-piperidine-1-yl]-butyryl}-phenyl)-isobutyramide

Dissolve N-methoxy-N-methyl-2-(4-{4-[4-hydroxy-
35 diphenylmethyl)-piperidin-1-yl]-butyryl}-phenyl)-isobutyramide (8.83g, 16.27mmol) in 3.5:1 methanol/tetrahydrofuran (85mL). Add sodium borohydride (0.62g, 16.27mmol) in 8 portions over 20 minutes at room

- 165 -

temperature. Stir at room temperature for 2 hours, evaporate the solvent *in vacuo*, dissolve the residue in ethyl acetate (60mL) and add water (25m). Stir at room
 5 temperature for 10 minutes, separate the layers and wash the organic layer with brine (2X25mL). Combine the organic layers, extract with ethyl acetate (35mL), dry (Na₂SO₄), evaporate the solvent *in vacuo* and dry to give the title compound (8.89g, 100%) as a foam; mp 80-83°C.

10

MS (CI, CH₄) m/e 545 (M⁺+1), 280, 236, 183.

Anal. Calcd for C₃₄H₄₄N₂O₄•0.2H₂O: C, 74.47; H, 8.16; N, 5.12; Found: C, 74.08; H, 8.16; N, 4.93.

15

Example 52

Step a: 1-[4-(1,1-Dimethyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-phenyl]-4-[4-hydroxy-diphenylmethyl)-piperidine-1-yl]-butan-1-one

- 20 Dissolve 4-chloro-1-[4-(1,1-dimethyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-phenyl]-butan-1-one (6.88g, 21.38mmol) in xylene (14mL) and add a suspension of α,α -diphenyl-4-piperidinemethanol hydrochloride (6.50g, 23.51mmol) and potassium carbonate (6.14g, 4.44mmol) in water (30mL).
- 25 Heat at 100°C for 24 hours, cool to room temperature, add methylene chloride (100mL) and separate the layers. Extract the aqueous layer with methylene chloride (100mL), wash with water (150mL), dry (Na₂SO₄), evaporate the solvent *in vacuo* and purify by silica gel chromatography (4:1 ethyl
- 30 acetate/methanol) to give the title compound (8.20g, 70%) as an off-white solid.

Anal. Calcd for C₃₆H₄₄N₂O₃•2H₂O: C, 77.72; H, 8.04; N, 5.08; Found: C, 77.38; H, 7.91; N, 4.93.

35

Example 53

Step c: 2-(4-{1-Hydroxy-4-[4-hydroxydiphenylmethyl)-

- 166 -

piperidin-1-yl]-butyl)-phenyl)-2-methyl-1-pyrrolidin-1-yl-
propan-1-one

Dissolve 1-[4-(1,1-dimethyl-2-oxo-2-pyrrolidin-1-yl-ethyl)-
 5 phenyl]-4-[4-hydroxy-diphenylmethyl)-piperidine-1-yl]-
 butan-1-one (0.55g, 1.00mmol) in methanol (10mL) and add
 sodium borohydride (38mg, 1.00mmol) at 10°C. Stir at room
 temperature for 2 hours, evaporate the solvent *in vacuo* and
 dissolve the residue in methylene chloride (60mL). Add
 10 water (10mL) and stir for 10 minutes. Separate the layers,
 wash with brine (5mL), dry (Na₂SO₄) and evaporate the
 solvent *in vacuo* to give the title compound (0.53g, 96%) as a
 white foam; mp 87-93°C.

15 Example 54

Step a: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-
oxobutyl]- α,α -dimethylbenzeneacetic acid, ethyl ester
hydrochloride

Dissolve 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic
 20 acid, ethyl ester (15.0g, 49.53mmol) and α,α -diphenyl-4-
 piperidinemethanol (29.66g, 106.4mmol) in xylene (60mL).
 Reflux for 5.5 hours, cool in an ice bath, filter and wash
 with cold xylenes (25mL). Filter the filtrate through
 silica gel (20g) and wash the gel with xylenes (40mL). Add
 25 xylene (60mL) and concentrated hydrochloric acid (6.45g,
 65.6mmol) with stirring. Add additional xylenes (40mL) and
 stir for 2 hour. Filter, wash with xylene (50mL), vacuum
 dry and slurry with a mixture of ethanol (60mL) and hexane
 (120mL) at 70-72°C for 30 minutes. Filter, wash with 3:1
 30 v/v solution of n-heptane/ethanol (30mL) and dry to give
 the title compound as a light white solid (19.7g, 70%); mp
 206-208°C.

¹H NMR (300MHz, CDCl₃) δ 7.90 (d, J=8.7Hz, 2H), 7.47 (m,
 35 4H), 7.41 (d, J=8.7Hz, 2H), 7.27 (m, 4H), 7.15 (m, 4H),
 4.10 (q, J=7.1Hz, 2H), 2.93 (m, 4H), 2.37 (m, 3H), 2.2
 (broad s, 1H), 1.92 (m, 4H), 1.59 (s, 6H), 1.39 (m, 4H),
 1.16 (t, J=7.1Hz, 3H); ¹³C NMR (75MHz, CDCl₃) δ 199.5,

- 167 -

176.1, 149.8, 146.0, 135.5, 128.2, 128.1, 126.4, 125.9,
 125.7, 79.4, 61.0, 57.8, 53.9, 46.7, 44.1, 36.3, 26.3,
 26.2, 21.9, 14.0; IR (CDCl₃) 3514, 2945, 1726, 1682, 1446,
 5 1254, 1147 1097 cm⁻¹;

Anal. Calcd for C₃₄H₄₁O₄N•HCl: C, 72.39; H, 7.50; N, 2.48;
 Found: C, 71.68; H, 7.52; N, 2.34.

10

Example 55

Step a: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetic acid, methyl ester hydrochloride

Dissolve 2-[4-(4-chloro-butyryl)-phenyl]-2-methyl-propionic
 15 acid, methyl ester (2.82g, 10.0mmol) and α,α-diphenyl-4-piperidinemethanol (5.58g, 21.0mmol) in toluene (20mL). Reflux for 29 hours, cool in an ice bath, filter, filter the filtrate through silica gel (5g) and wash the gel with toluene (10mL). Evaporate the solvent *in vacuo* and dissolve
 20 the residue in ethyl ether (100mL). Add anhydrous hydrogen chloride and filter to give the title compound as an off-white powder (4.2g, 76%); mp 165-175°C.

¹H NMR (300MHz, CDCl₃) δ 7.93 (d, J=8.3Hz, 2H), 7.47 (m, 4H), 7.42 (d, J=8.3Hz, 2H), 7.30 (m, 4H), 7.18 (m, 2H), 3.64 (s, 3H), 2.96 (m, 4H), 2.42 (m, 4H), 1.96 (m, 4H), 1.62 (s, 6H), 1.41 (m, 4H); ¹³C NMR (75MHz, CDCl₃) δ 199.1, 176.3, 149.4, 145.8, 135.5, 128.1, 128.0, 127.7, 126.3, 125.7, 1225.6, 79.4, 57.9, 54.0, 52.4, 46.9, 44.1, 36.4,
 30 26.4, 26.3, 22; MS (CI/NH₃) 514 (100 (M+H)), 293 (4), 268 (7).

Anal. Calcd for C₃₃H₃₉O₄N•HCl: C, 72.05; H, 7.33; N, 2.55;
 Found: C, 71.85; H, 7.23, N, 2.33.

35

Example 56

Step c: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-

- 168 -

hydroxybutyl]- α,α -dimethylbenzeneacetic acid, methyl ester hydrochloride

Dissolve 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]- α,α -dimethylbenzeneacetic acid, methyl ester hydrochloride (550mg, 1.00mmol) in methanol (5mL) and add sodium borohydride (62.8mg) in three batches. Stir for 1 hour, add 50% aqueous sodium hydroxide (800mg) and heat to reflux with stirring. After 3 hours, cool to -10°C , add approximately 1.5mL of 6N HCl over 10 minutes, filter the solid and wash with ice water (12mL) such that the final filtrate is pH=5. Dry the resulting solid *in vacuo* ($50-60^{\circ}\text{C}$, 10^{-1} mm) overnight to give the title compound (515mg, 94%); mp $165-180^{\circ}\text{C}$.

15

^1H NMR (300MHz, 5% MeOD_4 in CDCl_3) δ 7.50 (d, $J=7.3\text{Hz}$, 4H), 7.30 (m, 8H), 7.18 (t, $J=7.0\text{Hz}$, 2H), 4.66 (t, $J=5.3\text{Hz}$, 1H), 3.47 (m, 6H), 2.97 (m, 2H), 2.69 (m, 3H), 1.6-2.2 (m, 6H), 1.55 (s, 6H); ^{13}C NMR (75MHz, 5% MeOD_4 in CDCl_3) δ 179.1, 145.3, 143.8, 142.3, 128.2, 126.6, 125.7, 125.5, 125.4, 78.4 (bis benzylic), 72.5 (benzylic), 57.4, 53.2, 46.2, 24.2, 35.9, 26.6, 24.1, 20.8; MS (CI/ NH_3) 502 (100 (M+H)), 280 (5), 200 (10).

25

Example 57

Step c: 2-(4-(1-Hydroxy-4-(4-(hydroxydiphenylmethyl)-1-piperidinyl)-butyl)-phenyl)-2-methyl-propanol

Dissolve 2-(4-(1-oxo-4-(4-(hydroxydiphenylmethyl)-1-piperidinyl)-butyl)-phenyl)-2-methylpropanol in methanol (450mL) and stir for 15 minutes at room temperature. Add, by dropwise addition, a solution of sodium borohydride (2.25g, 0.06mol) in water (10mL) over 15 minutes. Stir for another 30 minutes and cool in an ice-bath. Slowly add concentrated hydrochloric acid (4mL) and water (8mL) and stir for an additional 20 minutes. Evaporate the solvent *in vacuo* and partition the residue between methylene chloride (150mL) and water (70mL). Separate the organic phase and

- 169 -

extract the aqueous phase with methylene chloride (25mL). Wash the combined organic layers with water (2X50mL), evaporate the solvent *in vacuo* and recrystallize (acetone) to
5 give the title compound as white needles (9.53g, 79%).

¹H NMR (300MHz, DMSO-d₆) δ 7.50 (4H, m), 7.23 (8H, m), 7.12 (2H, m), 5.34 (1H, s, br), 4.65 (1H, t), 4.45 (1H, s), 3.38 (2H, t), 2.60 (2H, m), 2.44 (2H, m), 2.20 (2H, t), 1.62
10 (2H, t), 1.50 (6H, m), 1.98 (6H, s); ¹³C NMR (DMSO-d₆) δ 147.2, 146.0, 143.4, 127.6, 125.6, 125.5, 125.2, 78.4, 72.0, 70.9, 58.0, 53.6, 53.5, 43.6, 38.0, 30.5, 25.9, 25.5, 23.1.

15

Alternatively, the novel intermediates of formula (XI) may be prepared as described in Scheme M. In Scheme M, all substituents are as previously defined unless otherwise indicated.


20

25

30

35

Scheme M



35

Scheme M provides various alternative general synthetic procedures for preparing the novel intermediates of formula (XI).

- 171 -

In step a, the appropriate ω' -piperidine-2-methylethylphenyl compound of structure (72) is cyanated to
5 give the corresponding ω' -piperidine- α,α -dimethylphenylacetonitrile compound of structure (73) as described previously in Scheme D, step b.

In step b, the appropriate ω' -piperidine-2-methylethylphenyl compound of structure (72) is halogenated
10 to give the corresponding ω' -piperidine- α,α -dimethylbenzyl halide compound of structure (74) as described previously in Scheme B, step a.

In step c, the nitrile functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetonitrile compound of structure (73) is converted to the
15 corresponding ester to give the ω' -piperidine- α,α -dimethylphenylacetic acid ester compound of structure (75)
20 as described previously in Scheme H, step a.

In step d, the halo functionality of the appropriate ω' -piperidine- α,α -dimethylbenzyl halide compound of structure (74) is converted to the corresponding carboxy to
25 give the ω' -piperidine- α,α -dimethylphenylacetic acid compound of structure (76) as described previously in Scheme H, step h.

In step e, the nitrile functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetonitrile compound of structure (73) is converted to the
30 corresponding carboxy to give the ω' -piperidine- α,α -dimethylphenylacetic acid compound of structure (76) as described previously in Scheme H, step e.

35

In step f, the nitrile functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetonitrile compound of structure (73) is converted to the

- 172 -

corresponding amide to give the ω' -piperidine- α,α -dimethylphenylacetic acid amide compound of structure (77) wherein R_6 and R_7 are each hydrogen as described previously
5 in Scheme H, step b.

In step g, the carboxy ester functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetic acid ester compound of structure (75) is hydrolyzed to give the
10 corresponding ω' -piperidine- α,α -dimethylphenylacetic acid compound of structure (76) as described previously in Scheme H, step c.

In step h, the carboxy functionality of the appropriate
15 ω' -piperidine- α,α -dimethylphenylacetic acid compound of structure (76) may be esterified by techniques and procedures well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -piperidine- α,α -dimethylphenylacetic acid ester compound of structure
20 (75) as described previously in Scheme H, step d.

In step i, the carboxy functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetic acid compound of structure (76) may be amidated by techniques and procedures
25 well known and appreciated by one of ordinary skill in the art to give the corresponding ω' -piperidine- α,α -dimethylphenylacetic acid amide compound of structure (77) as described previously in Scheme H, step g.

30 In step j, the amide functionality of the appropriate ω' -piperidine- α,α -dimethylphenylacetic acid amide compound of structure (77) is converted to the corresponding acid by acid hydrolysis as is known in the art to give the ω' -piperidine- α,α -dimethylphenylacetic acid compound of
35 structure (76) as described previously in Scheme H, step f.

Starting materials for use in Scheme M are readily available to one of ordinary skill in the art.

The following examples present typical syntheses as described in Scheme M. These examples are understood to be illustrative only and are not intended to limit the scope of the present invention in any way. As used herein, the following terms have the indicated meanings: "g" refers to grams; "mmol" refers to millimoles; "mL" refers to milliliters; "bp" refers to boiling point; "°C" refers to degrees Celsius; "mm Hg" refers to millimeters of mercury; "μL" refers to microliters; "μg" refers to micrograms; and "μM" refers to micromolar.

Example 58

15 Step g: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetic acid hydrochloride
Dissolve 4-[4-[4-(hydroxydiphenylmethyl)-1-piperidinyl]-1-oxobutyl]-α,α-dimethylbenzeneacetic acid methyl ester (0.131mol) in methanol (2.5L) and add 10% sodium hydroxide
20 (769mL, 1.92mol). Stir at reflux for 1.5 hours, cool to 68°C and evaporate the solvent *in vacuo* to a residue. Add chloroform (1L) and stir until the solids are dissolved. Separate the organic phase and extract the aqueous phase with chloroform (3X300mL). Combine the organic phases, dry
25 (MgSO₄) and evaporate the solvent *in vacuo* to give a residue. Treat the residue with ethereal HCl, filter and dry to give the title compound.

Example 59

30 Step j: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid
Dissolve N-methoxy-N-methyl-2-(4-{1-hydroxy-4-[4-hydroxy-diphenylmethyl)-piperidine-1-yl]-butyryl}-phenyl)-isobutyramide (8.35g, 15.33mmol) in isopropanol (50mL) and
35 add potassium hydroxide (8.63g, 153.7mmol). Heat to reflux for 2 hours, add additional potassium hydroxide (4.35g, 77.5mmol) and heat at reflux for an additional 16 hours. Cool to room temperature, treat with concentrated HCl by

- 174 -

dropwise addition until pH = 3. Dilute with water (100mL), stir vigorously for 2 hours, add ethyl acetate (30mL) and stir for 1 hour. Filter to give the title compound (7.15g, 87%) as an off-white solid.

MS (CI, CH₄) m/e 502 (M⁺+1), 107.

Anal. Calcd for C₃₂H₃₉NO₄•HCl•2.6H₂O: C, 65.70; H, 7.61; N, 2.39; Found: C, 65.25; H, 7.70; N, 2.36.

Example 60

Step j: 4-[4-[4-(Hydroxydiphenylmethyl)-1-piperidinyl]-1-hydroxybutyl]-α,α-dimethylbenzeneacetic acid

Dissolve N,N-dimethyl-2-(4-{1-hydroxy-4-[4-hydroxy-diphenylmethyl)-piperidin-1-yl]-butyry}-phenyl)-isobutyramide (15.33mmol) in isopropanol (50mL) and add potassium hydroxide (8.63g, 153.7mmol). Heat to reflux for 2 hours, add additional potassium hydroxide (4.35g, 77.5mmol) and heat at reflux for an additional 16 hours. Cool to room temperature, treat with concentrated HCl by dropwise addition until pH = 3. Dilute with water (100mL), stir vigorously for 2 hours, add ethyl acetate (30mL) and stir for 1 hour. Filter to give the title compound (41%).

As one skilled in the art would appreciate, the compounds depicted in Schemes A through M which bear hydroxy or phenolic functionalities may be protected prior to use in the synthesis depicted in Schemes A through M using suitable protecting groups. For example, suitable protecting groups for the phenolic hydroxy include methyl ether, 2-methoxyethoxymethyl ether (MEM), cyclohexyl ether, o-nitrobenzyl ether, 9-anthryl ether, t-butyldimethylsilyl ether, acetate, benzoate, methyl carbamate, benzyl carbamate, aryl pivaloate and aryl methanesulfonate.

As one skilled in the art would appreciate, the compounds depicted in Schemes A through M which bear α-

- 175 -

ketone functionalities may be protected prior to use in the synthesis depicted in Schemes A through M using suitable protecting groups. The selection and utilization of
5 suitable protecting groups for ketone groups is well known by one of ordinary skill in the art and is described in "Protective Groups in Organic Syntheses", Theodora W. Greene, Wiley (1981). For example, suitable protecting groups for ketone functionalities include acyclic acetals
10 and ketals such as dimethyl acetal, cyclic acetals and ketals such as 1,3-dioxanes and 1,3-dioxolanes, dithio acetals and ketals such as 1,3-dithiane and 1,3-dithiolane, hemithio acetals and ketals, O-substituted cyanohydrins, substituted hydrozones, imines, oxazolidines,
15 imidazolidines and thiazolidines.

As one skilled in the art would appreciate, the compounds depicted in Schemes A through M which bear protected hydroxy and/or ketone functionalities may be
20 reacting with appropriate deprotecting agents prior to use in any of the steps depicted in Schemes A through M. The selection and utilization of appropriate deprotecting reagents is well known by one of ordinary skill in the art and is described in "Protective Groups in Organic
25 Syntheses", Theodora W. Greene, Wiley (1981). Examples of appropriate deprotecting reagents are mineral acids, strong organic acids, Lewis acids, aqueous mineral bases, catalytic hydrogenation and the like.

30 For example, cleavage of β -methoxyethoxymethyl (MEM) protecting groups on any of the compounds depicted in Schemes A through M which bear protected hydroxy ketone functionalities, for example, can be achieved by using trifluoroacetic acid at room temperature or using 5 to 8
35 equivalents of powdered anhydrous zinc bromide in methylene chloride at about 25°C by the general procedure of E. J. Corey et al., *Tetrahedron Letters*, 11, 809-812 1976.

In addition, the individual (R) and (S) isomers of the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) can be prepared by techniques and procedures well known and appreciated by one of ordinary skill in the art.

For example, the mixture of (R) and (S) isomers of the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) may be subjected to chiral chromatography to give the corresponding individual (R)- ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) and (S)- ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71).

15

In addition, the individual (R) and (S) isomers of the ω -halo- α' -hydroxy- α,α -dimethylphenyl compound of structure (70) and the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) can be prepared by techniques and procedures well known and appreciated by one of ordinary skill in the art and described in "Enantiomers, Racemates, and Resolutions", Jacques, Collet and Wilen, Wiley (1981).

25

One such method involves reacting the mixture of (R) and (S) isomers of the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) with appropriate chiral acids to give the corresponding mixture of diastereomeric acid addition salts. The individual (R)- ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl chiral acid addition salt compounds of structure (71) and (S)- ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl chiral acid addition salt compounds of structure (71) are obtained by recrystallization and the individual ω' -piperidine-(R)- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) and ω' -piperidine-(S)- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) are obtained by subjecting the individual ω' -piperidine-(R)- α' -hydroxy- α,α -dimethylphenyl chiral acid

30

35


addition salt compounds of structure (71) and ω' -piperidine-(S)- α' -hydroxy- α,α -dimethylphenyl chiral acid addition salt compounds of structure (71) to base in order
 5 to free the piperidine nitrogen from the acid addition complex. Examples of suitable chiral acids are tartaric acid (+), (-), O,O'-dibenzoyltartaric acid (+), (-), O,O'-di-p-toluyltartaric acid (+), (-), 2-Nitrotartranillic acid (+), (-), mandelic acid (+), (-), malic acid (+), (-), 2-
 10 phenoxypropionic acid (+), hydratropic acid (+), (-), N-acetylleucine (-), (+), N-(α -methylbenzyl)succinamide (+), (-), N-(α -methylbenzyl)phthalamic acid (+), (-), camphor-10-sulfonic acid (+), 3-bromocamphor-9-sulfonic acid (+), (-), camphor-3-sulfonic acid (+), quinic acid (+), (-), Di-
 15 O-isopropylidene-2-oxo-L-gulononic acid (-), Lasalocid (-), 1,1'-binaphthyl-2,2'-phosphoric acid (+), (-), chloestenonesulfonic acid.

In addition, the individual (R) and (S) isomers of the
 20 ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) can be prepared by reacting the mixture of (R) and (S) isomers of the ω' -piperidine- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) with appropriate organic chiral acids to give the corresponding mixture of
 25 diastereomeric acid esters. The individual ω' -piperidine-(R)- α' -ester- α,α -dimethylphenyl compounds of structure (71) and ω' -piperidine-(S)- α' -ester- α,α -dimethylphenyl compounds of structure (71) are obtained by recrystallization or chromatography and the individual ω' -piperidine-(R)- α' -
 30 hydroxy- α,α -dimethylphenyl compounds of structure (71) and ω' -piperidine-(S)- α' -hydroxy- α,α -dimethylphenyl compounds of structure (71) are obtained by subjecting the individual ω' -piperidine-(R)- α' -ester- α,α -dimethylphenyl compounds of structure (71) and ω' -piperidine-(S)- α' -ester- α,α -
 35 dimethylphenyl compounds of structure (71) to hydrolysis conditions.

- 178 -

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A compound of the formula




wherein

- W represents $-C(=O)-$ or $-CH(OH)-$;
- R_1 represents hydrogen or hydroxy;
- R_2 represents hydrogen; or
- R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;
- n is an integer of from 1 to 5;
- m is an integer 0 or 1;
- R_5 is Br, Cl, I, CN or $-CONR_6R_7$ wherein R_6 and R_7 are each independently H, C_1 - C_6 alkyl, C_1 - C_6 alkoxy or R_6 and R_7 taken together with the nitrogen atom form a pyrrolidine, piperidine or morpholine, with the proviso that R_6 and R_7 cannot both be represented by C_1 - C_6 alkoxy;
- A is hydrogen or hydroxy; and

- 179 -

pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.

2. Use of the compound of claim 1 to prepare a compound of the formula



wherein

W represents $-C(=O)-$ or $-CH(OH)-$;

R_1 represents hydrogen or hydroxy;

R_2 represents hydrogen; or

R_1 and R_2 taken together form a second bond between the carbon atoms bearing R_1 and R_2 ;

n is an integer of from 1 to 5;

m is an integer 0 or 1;

R_3 is $-COOH$ or $-COOalkyl$ wherein the alkyl moiety has from 1 to 6 carbon atoms and is straight or branched;

- 180 -

each of A is hydrogen or hydroxy; and

pharmaceutically acceptable salts and individual optical isomers thereof, with the proviso that where R_1 and R_2 are taken together to form a second bond between the carbon atoms bearing R_1 and R_2 or where R_1 represented hydroxy, m is an integer 0.

