



HU000035884T2

(19) **HU**(11) Lajstromszám: **E 035 884**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 07 753102** (51) Int. Cl.: **G01N 33/564** (2006.01)
(22) A bejelentés napja: **2007. 03. 15.** **G01N 33/50** (2006.01)
C07K 2/00 (2006.01)
C07K 4/00 (2006.01)
(96) Az európai bejelentés bejelentési száma: **EP 20070753102**
(97) Az európai bejelentés közzétételi adatai: **EP 2001496 A2** **2007. 09. 27.** (86) A nemzetközi (PCT) bejelentési szám: **PCT/US 07/006451**
(97) Az európai szabadalom megadásának meghirdetési adatai: **EP 2001496 B1** **2017. 05. 31.** (87) A nemzetközi közzétételi szám: **WO 07109056**

(30) Elsőbbségi adatok: 782508 P 2006. 03. 15. US	(73) Jogosult(ak): THE BRIGHAM AND WOMEN'S HOSPITAL, INC., Boston, MA 02115 (US)
(72) Feltaláló(k): STOSSEL, Thomas, P., Belmont, MA 02478 (US) MAGNUSSON OSBORN, Anna, Charlotta, Teresia, Brookline, MA 02445 (US) TARKOWSKI, Andrej, S-413 46 Goteborg (SE)	(74) Képviselő: Danubia Szabadalmi és Jogi Iroda Kft., Budapest

(54) **Gelszolin alkalmazása gyulladós betegségek diagnosztizálására és kezelésére**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 001 496 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
31.05.2017 Bulletin 2017/22

(51) Int Cl.:
G01N 33/564 ^(2006.01) **C07K 2/00** ^(2006.01)
C07K 4/00 ^(2006.01) **G01N 33/50** ^(2006.01)

(21) Application number: **07753102.8**

(86) International application number:
PCT/US2007/006451

(22) Date of filing: **15.03.2007**

(87) International publication number:
WO 2007/109056 (27.09.2007 Gazette 2007/39)

(54) **USE OF GELSOLIN TO DIAGNOSE AND TREAT INFLAMMATORY DISEASES**

VERWENDUNG VON GELSOLIN ZUR DIAGNOSE UND BEHANDLUNG VON
ENTZÜNDUNGSERKRANKUNGEN

UTILISATION DE GELSOLINE AFIN DE DIAGNOSTIQUER ET DE TRAITER DES MALADIES
INFLAMMATOIRES

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE
SI SK TR**

(56) References cited:
**WO-A-02/059604 WO-A-03/088811
WO-A-2004/082617 WO-A-2005/046454
WO-A-2005/112970 US-A- 6 040 147
US-A1- 2006 009 386**

(30) Priority: **15.03.2006 US 782508 P**

(43) Date of publication of application:
17.12.2008 Bulletin 2008/51

(73) Proprietor: **THE BRIGHAM AND WOMEN'S
HOSPITAL, INC.
Boston, MA 02115 (US)**

(72) Inventors:
• **STOSSEL, Thomas, P.
Belmont, MA 02478 (US)**
• **MAGNUSSON OSBORN, Anna, Charlotta, Teresia
Brookline, MA 02445 (US)**
• **TARKOWSKI, Andrej
S-413 46 Goteborg (SE)**

(74) Representative: **Williams, Gareth Owen
Marks & Clerk LLP
62-68 Hills Road
Cambridge CB2 1LA (GB)**

- **CANDIANO GIOVANNI ET AL: "Gelsolin secretion in interleukin-4-treated bronchial epithelia and in asthmatic airways." AMERICAN JOURNAL OF RESPIRATORY AND CRITICAL CARE MEDICINE 1 NOV 2005, vol. 172, no. 9, 1 November 2005 (2005-11-01), pages 1090-1096, XP002527805 ISSN: 1073-449X**
- **ROTHENBACH PATRICIA A ET AL: "Recombinant plasma gelsolin infusion attenuates burn-induced pulmonary microvascular dysfunction." JOURNAL OF APPLIED PHYSIOLOGY (BETHESDA, MD. : 1985) JAN 2004, vol. 96, no. 1, January 2004 (2004-01), pages 25-31, XP002529488 ISSN: 8750-7587**
- **AIDINIS VASSILIS ET AL: "Cytoskeletal rearrangements in synovial fibroblasts as a novel pathophysiological determinant of modeled rheumatoid arthritis." PLOS GENETICS OCT 2005, vol. 1, no. 4, October 2005 (2005-10), page e48, XP002527807 ISSN: 1553-7404**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 001 496 B1

Description**Field of the Invention**

5 **[0001]** The invention is directed to diagnostic and therapeutic uses of gelsolin.

Background of the Invention

10 **[0002]** Inflammation is the body's response to injury, infection, or molecules perceived by the immune system as foreign. Inflammation is characterized by pain, swelling and altered function of the affected tissue. Although the ability to mount an inflammatory response is essential for survival, the ability to control inflammation is also necessary for health. Inflammatory diseases are characterized by activation of the immune system in a tissue or an organ to abnormal levels that may lead to abnormal function and/or disease in the tissue or organ.

15 **[0003]** Inflammatory diseases are a major cause of morbidity and mortality throughout the world. They affect various organs and tissues such as blood vessels, heart, brain, nerves, joints, skin, the lung, eye, gastrointestinal tract, kidneys, thyroid, adrenals, the pancreas, liver, and muscle. Treatments of inflammatory diseases have drawn a great deal of attention from the pharmaceutical industry. A recurrent theme in discussions of treatment options for inflammatory disorders is the inadequacy of the standard of care. Management and treatments are seeing improvements but there are no cures. The most common approach to treating inflammatory disorders in the last decade has addressed the pro-inflammatory role of cytokines with compounds that bind to these molecules or their receptors.

20 **[0004]** Despite recent advances, current therapies for inflammatory diseases still entail alleviating symptoms and reducing inflammation with non-specific drugs, slowing disease progression with disease-modifying agents, and improving the quality of life with lifestyle modifications, all while contending with side effects and resistance to medications. Better treatment options with less potential for side effects are needed. Because the outcome of treatment depends on a proper diagnosis, it is important to have proper tests to diagnose inflammatory diseases and to monitor the treatment of those diseases. A proper diagnosis permits a physician to institute proper and timely therapy. Proper monitoring of treatment allows the physician to decide on the course of treatment and to advise patients and their families about the expected disease course.

25 **[0005]** Thus, there is also a strong incentive to identify new improved tests and approaches to diagnose and to evaluate treatments of inflammatory diseases.

30 **[0006]** Gelsolin, first discovered as an intracellular actin-binding protein involved in cell motility (Yin, H. L. & Stossel, T. P. (1979) *Nature* 281, 583-6), has been recently implicated in a number of diseases. While the true function of plasma gelsolin is not known, clinical and animal studies have shown that depletion of plasma gelsolin by injury and inflammation is associated with adverse outcomes. The proposed mechanism of gelsolin depletion is that it binds abundant actin in cells exposed by tissue breakdown. More recently, gelsolin was found to bind bioactive inflammatory mediators, lyso-phosphatidic acid, diadenosine phosphate, AP peptide (a peptide implicated in the pathogenesis of Alzheimer's disease), platelet-activating factor and possibly others.

35 **[0007]** US 2006/0009386 and WO 2005/112970 relates to the use of gelsolin to treat infections and to monitor the treatment of infections.

40 **[0008]** WO 2005/046454 describes the administration of gelsolin to patients infected with gram-negative bacteria in order to control septic shock in a patient resulting from endotoxins triggered by release of LPS molecules from the bacterial infection.

45 **[0009]** WO 02/059604 discloses methods and compositions for screening, diagnosis and prognosis of Multiple Sclerosis (MS). The publication refers to a number of potential markers relating to Multiple Sclerosis which are differentially present in a sample from a subject having MS compared with a sample from a subject free from MS. One such marker is identified as gelsolin.

[0010] WO 2004/082617 describes biological markers for rheumatoid arthritis, and their use to diagnose and treat rheumatoid arthritis, or monitor progression of the disease. Table 2 gives a list of proteins which are decreased in subjects having rheumatoid arthritis, with one protein among many being referred to as gelsolin.

50 **[0011]** WO 03/088811 discloses the use of gelsolin as a prognostic marker of atherosclerotic diseases.

[0012] US 6040147 describes methods for characterizing an individual's risk profile of developing a future cardiovascular disorder by obtaining a level of the marker of systemic inflammation in the individual, in particular a C-reactive protein, a cytokine or a cellular adhesion molecule.

55 **[0013]** Candiano et al. (2005); *American Journal of Respiratory and Critical Care Medicine*; Vol. 172, No. 9 suggest that gelsolin might improve the fluidity of airway surface liquid in asthma by breaking down filamentous actin that may be released in large amounts by dying cells during inflammation.

[0014] Rothenbach et al. (2004); *Journal of Applied Physiology*; Vol. 96; No. 1 describes a study investigating the role of plasma gelsolin in the pathophysiology of inflammation-induced lung injury.

[0015] Aidinis et al. (2005) PLoS Genetics; Vol. 1; No. 4 identifies gelsolin-driven synovial fibroblast cytoskeletal rearrangements as a novel pathophysiological determinant of rheumatoid arthritis.

Summary of the Invention

5

[0016] The present invention relates to a method for characterizing a preferably apparently healthy subject's risk profile of developing a future inflammatory disease, comprising: comparing a level of gelsolin in a body fluid (for example blood, plasma, serum, urine, synovial fluid, or alveolar fluid), or tissue (for example a joint, gastrointestinal, thyroid, adrenal, vascular, pulmonary, renal, cardiac, skin, ocular, brain, pancreatic, liver, nerve, or muscle tissue) obtained from the subject to a predetermined value (preferably of about 250 mg/L of plasma or less), and characterizing the subject's risk profile of developing an inflammatory disease based upon the level of gelsolin in comparison to the predetermined value, wherein a level of gelsolin at or below the predetermined level is indicative that the subject is at an elevated risk of developing the inflammatory disease, and wherein a level of gelsolin at or above the predetermined level is indicative that the subject is not at an elevated risk of developing the inflammatory disease, wherein the subject does not have any signs or symptoms of an infection, sepsis or an actin-related disorder.

10

15

[0017] Another aspect of the present invention relates to A method for characterizing a preferably apparently healthy subject's risk profile of developing a future inflammatory disease, comprising comparing the level of the gelsolin in a sample obtained from a subject to a first predetermined value (preferably about 250 mg/L of plasma or lower) to establish a first risk value,

20

comparing the level of a second marker of inflammation in a sample obtained from a subject to a second predetermined value to establish a second risk value, and characterizing the subject's risk profile of developing the inflammatory disease based upon the combination of the first risk value and the second risk value, wherein the combination of the first risk value and second risk value establishes a third risk value different from said first and second risk values, wherein the subject does not have any signs or symptoms of an infection, sepsis or an actin-related disorder.

25

[0018] Another aspect of the present invention relates to gelsolin optionally in combination with a second agent, for use in a method for treating a subject at risk of an inflammatory disease who is known to have a below-normal level of gelsolin; in an effective amount to reduce the risk of the subject developing the inflammatory disease or in an effective amount to raise the level of gelsolin in the subject above a predetermined value.

30

[0019] Gelsolin (GSN), specifically cytoplasmic gelsolin (cGSN), in addition to being an intracellular actin-binding protein involved in cell motility, is also an abundant secretory protein (Yin, H. L., Kwiatkowski, D. J., Mole, J. E. & Cole, F. S. (1984) J Biol Chem 259, 5271-6). The exported isoform of gelsolin, designated plasma gelsolin (pGSN), has 25 additional amino acids and originates from alternative splicing of a single gene (Kwiatkowski, D. J., Stossel, T. P., Orkin, S. H., Mole, J. E., Cohen, H. R. & Yin, H. L. (1986) Nature 323, 455-8).

35

[0020] This invention is based on the surprising discovery that plasma gelsolin levels are reduced in blood samples from human subjects with an inflammatory disease, rheumatoid arthritis (RA). These findings support the hypothesis that reductions in plasma gelsolin levels reflect the primary injury inflicted on joint tissues by the causative agency of rheumatoid arthritis and precede joint pain and destruction by the resultant inflammatory process. These observations provide a basis for treatment with gelsolin to prevent and/or suppresses the manifestations of inflammatory diseases. One correlate of these observations is that monitoring of plasma gelsolin levels could become part of the management strategy of rheumatoid arthritis.

40

[0021] Without intending to be bound by any particular mechanism or theory, it is believed that gelsolin might be exerting its protective effect by inhibiting mediators of inflammation. Thus, the invention is directed to methods of using gelsolin to diagnose inflammatory diseases and to monitor the effect of therapy. The invention also involves the use of gelsolin to treat inflammation and inflammatory diseases.

45

[0022] According to one aspect of the invention, a method for characterizing a subject's risk profile of developing a future inflammatory disease (e.g., rheumatoid arthritis in some preferred embodiments) is provided. The method comprises obtaining a level of gelsolin in the subject and comparing the level of the gelsolin to a predetermined value. The subject's risk profile of developing an inflammatory disease (e.g., rheumatoid arthritis) is characterized based upon the level of gelsolin in comparison to the predetermined value. A level of gelsolin at or below the predetermined level is indicative that the subject is at an elevated risk of developing the inflammatory disease and a level of gelsolin at or above the predetermined level is indicative that the subject is not at an elevated risk of developing the inflammatory disease.

50

[0023] In some embodiments, the method further comprises performing one or more tests to evaluate the inflammatory disease. Evaluating an inflammatory disease may involve measuring a level of a marker of inflammation in the subject. Examples of markers of inflammation include but are not limited to CRP, soluble intercellular adhesion molecule (sICAM-1), ICAM 3, BL-CAM, LFA-2, VCAM-1, NCAM, PECAM, fibrinogen, serum amyloid A (SAA), lipoprotein associated phospholipase A2 (LpPIA2), sCD40 ligand (sCD4OL), myeloperoxidase, Interleukin-6 (IL-6), or Interleukin-8 (IL-8).

55

[0024] According to another aspect of the invention, a method for characterizing a subject's risk profile of developing a future inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method comprises

obtaining a level of gelsolin in the subject and comparing the level of the gelsolin to a first predetermined value to establish a first risk value. A level of a second marker of inflammation in the subject is obtained and the level of the second marker of inflammation is compared to a second predetermined value to establish a second risk value. The subject's risk profile of developing the inflammatory disease is characterized based upon the combination of the first risk value and the second risk value, wherein the combination of the first risk value and second risk value establishes a third risk value different from said first and second risk values.

[0025] In some embodiments, the subject is an apparently healthy subject.

[0026] In some embodiments, the first predetermined value may be a plurality of predetermined gelsolin level ranges, one of a plurality of ranges being below about 250 mg/L of plasma and another of said ranges being above about 250 mg/L of plasma, and the comparing step comprises determining in which of said plurality of predetermined gelsolin level ranges said subject's gelsolin level falls.

[0027] According to another aspect of the present disclosure, a method for treating a subject having or at risk of developing an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method comprises administering an effective amount of gelsolin to the subject in need of such a treatment to treat the subject.

[0028] According to another aspect of the disclosure, a method for treating a subject having or at risk of developing an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method comprises administering an effective amount of gelsolin to the subject in need of such a treatment to raise the level of gelsolin in the subject above a predetermined value.

[0029] In some embodiments, the subject is otherwise free of indications calling for treatment with gelsolin. The gelsolin preferably is administered orally, sublingually, buccally, intranasally, intravenously, intramuscularly, intraarticularly, intraperitoneally, subcutaneously, or topically. The gelsolin may be administered prophylactically.

[0030] In some embodiments, the treatment methods further comprise administering a second agent for treating the inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). Examples of agents for treating the inflammatory disease include but are not limited to Alclufenac, Alciometasone Dipropionate, Algestone Acetonide, Alpha Amylase, Amcinafal, Amcinafide, Amfenac Sodium, Amiprilose Hydrochloride, Anakinra, Aniolac, Anitrazafen, Apazone, Balsalazide Disodium, Bendazac, Benoxaprofen, Benzydamine Hydrochloride, Bromelains, Broperamol, Budesonide, Carprofen, Cicloprofen, Cintazone, Cliprofen, Clobetasol Propionate, Clobetasone Butyrate, Clopirac, Cloticasone Propionate, Cormethasone Acetate, Cortodoxone, Cyclooxygenase-2 (COX-2) inhibitor, Deflazacort, Desonide, Desoximetasone, Dexamethasone Dipropionate, Diclofenac Potassium, Diclofenac Sodium, Diflorasone Diacetate, Diflumidone Sodium, Diflunisal, Difluprednate, Diftalone, Dimethyl Sulfoxide, Drocinonide, Endryson, Enlimomab, Enolicam Sodium, Epirizole, Etodolac, Etofenamate, Felbinac, Fenamole, Fenbufen, Fenclofenac, Fenclorac, Fendosal, Fenpivalone, Fentiazac, Flazalone, Fluazacort, Flufenamic Acid, Flumizole, Flunisolide Acetate, Flunixin, Flunixin Meglurnine, Fluocortin Butyl, Fluorometholone Acetate, Fluquazone, Flurbiprofen, Fluretofen, Fluticasone Propionate, Furaprofen, Furobufen, Halcinonide, Halobetasol Propionate, Halopredone Acetate, Ibufenac, Ibuprofen, Ibuprofen Aluminum, Ibuprofen Piconol, Ilonidap, Indomethacin, Indomethacin Sodium, Indoprofen, Indoxole, Intrazole, Isoflupredone Acetate, Isoxepac, Isoxicam, Ketoprofen, Lofemizole Hydrochloride, Lornoxicam, Loteprednol Etabonate, Meclofenamate Sodium, Meclofenamic Acid, Meclorison Dibutyrate, Mefenamic Acid, Mesalarnine, Meseclazone, Methylprednisolone Suleptanate, Morniflumate, Nabumetone, Naproxen, Naproxen Sodium, Naproxol, Nimazone, Olsalazine Sodium, Orgotein, Orpanoxin, Oxaprozin, Oxyphenbutazone, Paranyline Hydrochloride, Pentosan Polysulfate Sodium, Phenbutazone Sodium Glycerate, Pirfenidone, Piroxicam, Piroxicam Cinnamate, Piroxicam Olamine, Pirprofen, Prednazate, Prifelone, Prodolic Acid, Proquazone, Proxazole, Proxazole Citrate, Rimexolone, Romazarit, Salcolex, Salnacedin, Salsalate, Sanguinarium Chloride, Seclazone, Sermetacin, Sudoxicam, Sulindac, Suprofen, Talmetacin, Talniflumate, Talosalate, Tebufelone, Tenidap, Tenidap Sodium, Tenoxicam, Tesicam, Tesimide, Tetrydamine, Tiopinac, Tixocortol Pivalate, Tolmetin, Tolmetin Sodium, Triclonide, Triflumidate, Zidometacin, or Zomepirac Sodium.

[0031] Anti-inflammatory agents also include Cyclooxygenase-2 (COX-2) inhibitors. Cyclooxygenase is an enzyme complex present in most tissues that produces various prostaglandins and thromboxanes from arachidonic acid. Non-steroidal, anti-inflammatory drugs exert most of their anti-inflammatory, analgesic and antipyretic activity and inhibit hormone-induced uterine contractions and certain types of cancer growth through inhibition of the cyclooxygenase (also known as prostaglandin G/H synthase and/or prostaglandin-endoperoxide synthase). Initially, only one form of cyclooxygenase was known, the "constitutive enzyme" or cyclooxygenase-1 (COX-1). It was originally identified in bovine seminal vesicles.

[0032] Cyclooxygenase-2 (COX-2) has been cloned, sequenced and characterized initially from chicken, murine and human sources (See, e.g., U.S. Patent 5,543,297, issued August 6, 1996 to Cromlish, et al., and assigned to Merck Frosst Canada, Inc., Kirkland, CA, entitled: "Human cyclooxygenase-2 cDNA and assays for evaluating cyclooxygenase-2 activity"). This enzyme is distinct from the COX-1. COX-2, is rapidly and readily inducible by a number of agents including mitogens, endotoxin, hormones, cytokines and growth factors. As prostaglandins have both physiological and pathological roles, it is believed that the constitutive enzyme, COX-1, is responsible, in large part, for endogenous basal release of prostaglandins and hence is important in their physiological functions such as the maintenance of gastroin-

testinal integrity and renal blood flow. By contrast, it is believed that the inducible form, COX-2, is mainly responsible for the pathological effects of prostaglandins where rapid induction of the enzyme would occur in response to such agents as inflammatory agents, hormones, growth factors, and cytokines. Therefore, it is believed that a selective inhibitor of COX-2 has similar antiinflammatory, antipyretic and analgesic properties to a conventional non-steroidal antiinflammatory drug, and in addition inhibits hormone-induced uterine contractions and also has potential anti-cancer effects, but with reduced side effects. In particular, such COX-2 inhibitors are believed to have a reduced potential for gastrointestinal toxicity, a reduced potential for renal side effects, a reduced effect on bleeding times and possibly a decreased potential to induce asthma attacks in aspirin-sensitive asthmatic subjects, and are therefore useful according to the present invention.

[0033] A number of selective COX-2 inhibitors are known in the art. Examples of selective COX-2 inhibitors include, for example, celecoxib (Celebrex®), valdecoxib (Bextra®) and rofecoxib (Vioxx®). Selective COX-2 inhibitors also include, but are not limited to, COX-2 inhibitors described in U.S. Patent 5,474,995 "Phenyl heterocycles as COX-2 inhibitors"; U.S. Patent 5,521,213 "Diaryl bicyclic heterocycles as inhibitors of cyclooxygenase-2"; U.S. Patent 5,536,752 "Phenyl heterocycles as COX-2 inhibitors"; U.S. Patent 5,550,142 "Phenyl heterocycles as COX-2 inhibitors"; U.S. Patent 5,552,422 "Aryl substituted 5,5 fused aromatic nitrogen compounds as anti-inflammatory agents"; U.S. Patent 5,604,253 "N-benzylindol-3-yl propanoic acid derivatives as cyclooxygenase inhibitors"; U.S. Patent 5,604,260 "5-methanesulfonamido-1-indanones as an inhibitor of cyclooxygenase-2"; U.S. Patent 5,639,780 N-benzyl indol-3-yl butanoic acid derivatives as cyclooxygenase inhibitors"; U.S. Patent 5,677,318 Diphenyl-1,2-3-thiadiazoles as anti-inflammatory agents"; U.S. Patent 5,691,374 "Diaryl-5-oxygenated-2-(5H)-furanones as COX-2 inhibitors"; U.S. Patent 5,698,584 "3,4-diaryl-2-hydroxy-2,5-dihydrofurans as prodrugs to COX-2 inhibitors"; U.S. Patent 5,710,140 "Phenyl heterocycles as COX-2 inhibitors"; U.S. Patent 5,733,909 "Diphenyl stilbenes as prodrugs to COX-2 inhibitors"; U.S. Patent 5,789,413 "Alkylated styrenes as prodrugs to COX-2 inhibitors"; U.S. Patent 5,817,700 "Bisaryl cyclobutenes derivatives as cyclooxygenase inhibitors"; U.S. Patent 5,849,943 "Stilbene derivatives useful as cyclooxygenase-2 inhibitors"; U.S. Patent 5,861,419 "Substituted pyridines as selective cyclooxygenase-2 inhibitors"; U.S. Patent 5,922,742 "Pyridiny1-2-cyclopenten-1-ones as selective cyclooxygenase-2 inhibitors"; U.S. Patent 5,925,631 "Alkylated styrenes as prodrugs to COX-2 inhibitors"; all of which are commonly assigned to Merck Frosst Canada, Inc. (Kirkland, CA). Additional COX-2 inhibitors are also described in U.S. Patent 5,643,933, assigned to G. D. Searle & Co. (Skokie, IL), entitled: "Substituted sulfonylphenyl-heterocycles as cyclooxygenase-2 and 5-lipoxygenase inhibitors."

[0034] A number of the above-identified COX-2 inhibitors are prodrugs of selective COX-2 inhibitors, and exert their action by conversion in vivo to the active and selective COX-2 inhibitors. The active and selective COX-2 inhibitors formed from the above-identified COX-2 inhibitor prodrugs are described in detail in WO 95/00501, published January 5, 1995, WO 95/18799, published July 13, 1995 and U.S. Patent 5,474,995, issued December 12, 1995. Given the teachings of U.S. Patent 5,543,297, entitled: "Human cyclooxygenase-2 cDNA and assays for evaluating cyclooxygenase-2 activity," a person of ordinary skill in the art would be able to determine whether an agent is a selective COX-2 inhibitor or a precursor of a COX-2 inhibitor, and therefore part of the present invention.

[0035] In some embodiments, the method further comprises administering a second agent for treating rheumatoid arthritis. Examples of agents for treating rheumatoid arthritis include but are not limited to hydroxychloroquine (Plaquenil), chloroquine (Aralen), methotrexate, sulfasalazine (Azulfidine), Lefhmomide (Arava), azathioprine (Imuran), penicillamine (Cuprimine or Depen), Gold salts (Ridaura or Aurolate), minocycline (Dynacin or Minocin), cyclosporine (Neoral or Sandimmune) cyclophosphamide (Cytoxan or Neosar), Etanercept (Enbrel), Infliximab (Remicade), Ahakinra (Kineret) or Adalimumab (Humira).

[0036] According to another aspect of the disclosure, a method for treating a subject to reduce the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method comprises selecting a subject on the basis that the subject is known to have a below-normal level of gelsolin and administering to the subject an effective amount of gelsolin and/or a second agent to reduce the risk of the subject developing the inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments).

[0037] According to another aspect of the disclosure, a method for treating a subject to reduce the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method comprises selecting a subject on the basis that the subject is known to have a below-normal level of gelsolin and administering an effective amount of gelsolin and/or a second agent to the subject to raise the level of gelsolin in the subject above a predetermined value.

[0038] In some embodiments, the method further comprises administering to the subject a second agent for treating the inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). Examples of agents for treating the inflammatory disease and rheumatoid arthritis are listed above.

[0039] According to yet another aspect of the disclosure, a method for treating a subject with a below-normal level of gelsolin is provided. The method comprises treating the subject with a first therapy for treating or reducing the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). A level of gelsolin in the subject is obtained. The level of gelsolin is compared to a predetermined value corresponding to a predetermined level of gelsolin (e.g., in an apparently healthy control population). If the predetermined level of gelsolin is not reached, the subject is treated

with a second agent for treating or reducing the risk of the inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) until the predetermined level of gelsolin is reached.

[0040] A "below-normal level of gelsolin" is a gelsolin level is at least 10% less than the measured mean level for a given population of subjects. The mean gelsolin level can depend upon the particular population of subjects. For example, an apparently healthy population will have a different "normal" range of gelsolin than will a population of subjects which have had a prior condition. In some embodiments, the gelsolin level is at least 10% less than the measured mean level for a given population of subjects. In other embodiments, the gelsolin level is at least 20% less than the measured mean level for a given population of subjects. In still other embodiments, the gelsolin level is at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95%, 99% or 100% less than the measured mean level for a given population of subjects. In one of the embodiments, the gelsolin level is below about 250 mg/L of plasma. In other important embodiments, the gelsolin level is below about 2.4 μ M/L (micromoles/Liter) of plasma.

[0041] In some embodiments the subject is otherwise free of indications calling for treatment with the agent. When the agent is gelsolin, a subject free of indications calling for treatment with gelsolin is a subject who has no signs or symptoms calling for treatment with gelsolin. Gelsolin is indicated for the treatment of sepsis and infections. Gelsolin is also indicated for the treatment of actin-related disorders such as Adult

[0042] Respiratory Distress Syndrome (ARDS), fulminant hepatic necrosis, acute renal failure, muscle injury, disorders characterized by elevated levels of BUN and/or creatinine. Actin-related disorders are known to those of ordinary skill in the art.

[0043] In other embodiments, the subject is apparently healthy. As used herein an "apparently healthy subject" is a subject who has no signs and/or symptoms of a disease.

[0044] According to another aspect of the disclosure, a method for evaluating the efficacy of a therapy for treating or reducing the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) in a subject is provided. The method comprises obtaining a level of gelsolin in a subject undergoing therapy with an agent to treat or reduce the risk of inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). The level of gelsolin obtained is compared to a predetermined value corresponding to a level of gelsolin (e.g., in an apparently healthy control population). A determination of whether the level of gelsolin is above the predetermined level is indicative of whether the therapy is efficacious.

[0045] In some embodiments, obtaining a level of the gelsolin is repeated so as to monitor the subject's level of the gelsolin over time.

[0046] The therapy may be with gelsolin, Alclofenac, Alclometasone Dipropionate, Algestone Acetonide, Alpha Amylase, Amcinafal, Arncinafide, Amfenac Sodium, Amiprilose Hydrochloride, Anakinra, Anirolac, Anitrazafen, Apazone, Balsalazide Disodium, Bendazac, Benoxaprofen, Benzydamine Hydrochloride, Bromelains, Broperamol, Budesonide, Carprofen, Cicloprofen, Cintazone, Cliprofen, Clobetasol Propionate, Clobetasone Butyrate, Clopirac, Cloticasone Propionate, Cormethasone Acetate, Cortodoxone, Cyclooxygenase-2 (COX-2) inhibitor, Deflazacort, Desonide, Desoximetasone, Dexamethasone Dipropionate, Diclofenac Potassium, Diclofenac Sodium, Diflorasone Diacetate, Diflumidone Sodium, Diflunisal, Difluprednate, Diftalone, Dimethyl Sulfoxide, Drocinnonide, Endryson, Enlimomab, Enolicam Sodium, Epirizole, Etodolac, Etofenamate, Felbinac, Fenamole, Fenbufen, Fenclofenac, Fenclorac, Fendosal, Fempipalone, Fentia7ac, Flazalone, Fluazacort, Flufenamic Acid, Flumizole, Flunisolid Acetate, Flunixin, Flunixin Meglumine, Fluocortin Butyl, Fluorometholone Acetate, Fluquazone, Flurbiprofen, Fluretofen, Fluticasone Propionate, Furaprofen, Furobufen, Halcinonide, Halobetasol Propionate, Halopredone Acetate, Ibufenac, Ibuprofen, Ibuprofen Aluminum, Ibuprofen Piconol, Ilonidap, Indomethacin, Indomethacin Sodium, Indoprofen, Indoxole, Intrazole, Isoflupredone Acetate, Isoxepac, Isoxicam, Ketoprofen, Lofemizole Hydrochloride, Lornoxicam, Loteprednol Etabonate, Meclofenamate Sodium, Meclofenamic Acid, Meclorison Dibutyrate, Mefenamic Acid, Mesalamine, Meseclazone, Methylprednisolone Suleptanate, Morniflumate, Nabumetone, Naproxen, Naproxen Sodium, Naproxol, Nimazone, Olsalazine Sodium, Orgotein, Orpanoxin, Oxaprozin, Oxyphenbutazone, Paranyline Hydrochloride, Pentosan Polysulfate Sodium, Phenbutazone Sodium Glycerate, Pirfenidone, Piroxicam, Piroxicam Cinnamate, Piroxicam Olamine, Pirprofen, Prednazate, Prifelone, Prodolic Acid, Proquazone, Proxazole, Proxazole Citrate, Rimexolone, Romazarit, Salcolex, Salnacedin, Salsalate, Sanguinarium Chloride, Seclazone, Sermetacin, Sudoxicam, Sulindac, Suprofen, Talmecacin, Talniflumate, Talosalate, Tebufelone, Tenidap, Tenidap Sodium, Tenoxicam, Tesicam, Tesimide, Tetrydamine, Tiopinac, Tixocortol Pivalate, Tolmetin, Tolmetin Sodium, Triclonide, Triflurnidate, Zidometacin, Zomepirac Sodium, hydroxychloroquine (Plaquenil), chloroquine (Aralen), methotrexate, sulfasalazine (Azulfidine), Leflunomide (Arava), azathioprine (Imuran), penicillamine (Cuprimine or Depen), Gold salts (Ridaura or Aurolate), minocycline (Dynacin or Minocin), cyclosporine (Neoral or Sandimmune) cyclophosphamide (Cytoxan or Neosar), Etanercept (Enbrel), Infliximab (Remicade), Anakinra (Kineret) or Adalimumab (Humira).

[0047] According to still another aspect of the disclosure, a method for deciding on the course of a therapy in a subject is provided. The method comprises obtaining a level of gelsolin in a subject undergoing a therapy to treat or reduce the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). The level of gelsolin is compared to a predetermined value corresponding to a level of gelsolin (e.g., in an apparently healthy control population). Whether

the level of gelsolin obtained is at or above or at or below the predetermined level is determined and the course of therapy is decided based on such determination. In some embodiments, obtaining a level of gelsolin is repeated so as to monitor the subject's level of gelsolin over time.

[0048] The following embodiments apply to various aspects of the invention set forth herein unless indicated otherwise.

[0049] The inflammatory disease may be arthritis, rheumatoid arthritis, asthma, inflammatory bowel disease (Crohn's disease or ulcerative colitis), chronic obstructive pulmonary disease (COPD), allergic rhinitis, vasculitis (polyarteritis nodosa, temporal arteritis, Wegener's granulomatosis, Takayasu's arteritis, or Behcet syndrome), inflammatory neuropathy, psoriasis, systemic lupus erythematosus (SLE), chronic thyroiditis, Hashimoto's thyroiditis, Addison's disease, polymyalgia rheumatica, Sjogren's syndrome, or Churg-Strauss syndrome. In some important embodiments, the inflammatory disease is rheumatoid arthritis.

[0050] The level of gelsolin may be in a body fluid of the subject. Examples of body fluids include but are not limited to blood, plasma, serum, urine, synovial fluid, or alveolar fluid.

[0051] The level of gelsolin may be in a body tissue of the subject. The body tissue may be joint, gastrointestinal, thyroid, adrenal, vascular, pulmonary, renal, cardiac, skin, ocular, brain pancreatic, liver, nerve, or muscle tissue. In some embodiments, the subject is an apparently healthy subject.

[0052] In some embodiments, the predetermined value is 250 mg/L of plasma or lower. In some embodiments, the predetermined value of gelsolin is about 240 mg/L, 230 mg/L, 220 mg/L, 210 mg/L, 200 mg/L, 190 mg/L, 180 mg/L, 170 mg/L, 160 mg/L, 150 mg/L, 140 mg/L, 130 mg/L, 120 mg/L, 110 mg/L, 100 mg/L, 90 mg/L, 80 mg/L, 70 mg/L, 60 mg/L, 50 mg/L, 40 mg/L, 30 mg/L, 20 mg/L, or 10 mg/L of plasma or lower.

[0053] In some other embodiments, the predetermined value is 2.4 μ M/L of plasma or lower. In some embodiments, the predetermined value of gelsolin is about 2.3 μ M/L, 2.2 μ M/L, 2.1 μ M/L, 2.0 μ M/L, 1.9 μ M/L, 1.8 μ M/L, 1.7 μ M/L, 1.6 μ M/L, 1.5 μ M/L, 1.4 μ M/L, 1.3 μ M/L, 1.2 μ M/L, 1.1 μ M/L, 1.0 μ M/L, 0.9 μ M/L, 0.8 μ M/L, 0.7 μ M/L, 0.6 μ M/L, 0.5 μ M/L, 0.4 μ M/L, 0.3 μ M/L, 0.2 μ M/L of plasma or lower.

[0054] Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention. The invention is capable of other embodiments and of being practiced or of being carried out in various ways. Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including", "comprising", or "having", "containing", "involving", and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

[0055] These and other aspects of the inventions, as well as various advantages and utilities will be apparent with reference to the Detailed Description of the Invention. Each aspect of the invention can encompass various embodiments as will be understood.

Brief Description of the Drawings

[0056]

FIG. 1 is a histogram showing that plasma gelsolin (pGSN) concentration is decreased in patients with rheumatoid arthritis (RA) compared to healthy controls and lower in synovial fluid than in blood in patients with RA.

FIG. 2 is an immunoblot with an antibody specific for the plasma isoform of gelsolin showing that the gelsolin present in the synovial fluid (SF) of RA patients is composed mainly of the plasma isoform.

FIG. 3 is a set of histograms showing that plasma gelsolin (pGSN) concentrations are decreased in mouse models of septic arthritis: A) Staphylococcus aureus (S. aureus) induced arthritis, B) sepsis induced by Staphylococcus aureus (S. aureus) and C) Streptococcus agalactiae (Str. agalactiae) induced septic arthritis. The decrease occurred at the earliest time-point tested (two days post-inoculation).

[0057] It is to be understood that the drawings are not required for enablement of the invention.

Detailed Description of the Invention

[0058] This invention is based on the surprising discovery that plasma gelsolin levels are reduced in blood samples from human subjects with an inflammatory disease, rheumatoid arthritis (RA). It is hypothesized that plasma gelsolin levels fall in response to the initial (unknown) injury inflicted by the agency causing RA. Therefore, peripheral gelsolin replacement can ameliorate the secondary injury mediated by various inflammatory cells. It is believed that the pattern of gelsolin depletion to predict disease and the use of gelsolin to treat disease is true for inflammatory diseases in general.

[0059] The present disclosure involves, in some aspects, administering gelsolin to a subject to treat an inflammatory disease in the subject. The term "treat" or "treatment" is intended to include prophylaxis, amelioration, prevention or cure from the disease.

[0060] As used herein the term "subject" means any mammal that may be in need of treatment. Subjects include but are not limited to: humans, non-human primates, cats, dogs, sheep, pigs, horses, cows, rodents such as mice, hamsters, and rats. Preferred subjects are human subjects.

[0061] As used herein the term "gelsolin" encompasses wild type gelsolin (GenBank accession No.: X04412), isoforms, analogs, variants, fragments or functional derivatives of gelsolin.

[0062] Gelsolin (GSN), unlike other mammalian proteins, has both cytoplasmic (cGSN) and secreted or exported isoforms, also called plasma gelsolin (pGSN), which are derived by alternative splicing of the message from a single gene (Sun et al. *J. Biol. Chem.* 274:33179-33182 (1999)). As used herein, gelsolin isoforms include versions of gelsolin with some small differences in their amino acid sequences, usually a splice variant or the result of some posttranslational modification.

[0063] Gelsolin encompasses native as well as synthetic and recombinant gelsolin and gelsolin analogs. Gelsolin is an abundant secretory protein (Yin, H. L., Kwiatkowski, D. J., Mole, J. E. & Cole, F. S. (1984) *J Biol Chem* 259, 5271-6). The exported isoform of gelsolin, pGSN, has 25 additional amino acids and originates from alternative splicing of a single gene (Kwiatkowski, D. J., Stossel, T. P., Orkin, S. H., Mole, J. E., Colten, H. R. & Yin, H. L. (1986) *Nature* 323, 455-8). Recombinant human gelsolin (rhGSN) (Biogen IDEC, Inc., Cambridge, MA) is produced in *E. coli*, and though it has the same primary structure as the native protein, under standard conditions of purification, it differs from natural human plasma gelsolin by a disulfide bond that is present in the natural protein. The recombinant protein is, therefore, properly oxidized after purification, and its structure and functions are indistinguishable from human plasma gelsolin (Wen et. al., *Biochemistry* 35:9700-9709 (1996)). In some of the important therapeutic aspects and embodiments of the disclosure, the use of rhGSN is preferred. In some of the important diagnostic aspects and embodiments of the disclosure, the use of pGSN is preferred.

[0064] A "gelsolin analog" refers to a compound substantially similar in function to either the native gelsolin or to a fragment thereof. Gelsolin analogs include biologically active amino acid sequences substantially similar to the gelsolin sequences and may have substituted, deleted, elongated, replaced, or otherwise modified sequences that possess bioactivity substantially similar to that of gelsolin. For example, an analog of gelsolin is one which does not have the same amino acid sequence as gelsolin but which is sufficiently homologous to gelsolin so as to retain the bioactivity of gelsolin. Bioactivity can be determined, for example, by determining the properties of the gelsolin analog and/or by determining the ability of the gelsolin analog to treat or prevent rheumatoid arthritis. One example of a gelsolin bioactivity assay is gelsolin's ability to stimulate actin nucleation. Gelsolin bioactivity assays are described in the Example and are known to those of ordinary skill in the art.

[0065] A "fragment" is meant to include any portion of a gelsolin molecule which provides a segment of gelsolin which maintains the bioactivity of gelsolin; the term is meant to include gelsolin fragments which are made from any source, such as, for example, from naturally-occurring peptide sequences, synthetic or chemically-synthesized peptide sequences, and genetically engineered peptide sequences.

[0066] A "variant" of gelsolin is meant to refer to a compound substantially similar in structure and bioactivity either to native gelsolin, or to a fragment thereof. The term variant encompasses the gelsolin family of proteins. The gelsolin family of proteins is a group of actin binding proteins sharing repeats of about 15kDa homologous domains that adopt a similar fold. Examples gelsolin family proteins include but are not limited to advillin, villin, capG, flightless proteins, fragmin, severin, adseverin, protovillin, and supervillin.

[0067] A "functional derivative" of gelsolin is a derivative which possesses a bioactivity that is substantially similar to the bioactivity of gelsolin. By "substantially similar" is meant activity which is quantitatively different but qualitatively the same. For example, a functional derivative of gelsolin could contain the same amino acid backbone as gelsolin but also contains other modifications such as post-translational modifications such as, for example, bound phospholipids, or covalently linked carbohydrate, depending on the necessity of such modifications for the performance of the diagnostic assay or therapeutic treatment. As used herein, the term is also meant to include a chemical derivative of gelsolin. Such derivatives may improve gelsolin's solubility, absorption, biological half life, etc. The derivatives may also decrease the toxicity of gelsolin, or eliminate or attenuate any undesirable side effect of gelsolin, etc. Chemical moieties capable of mediating such effects are disclosed in Remington's *Pharmaceutical Sciences* (1980). Procedures for coupling such moieties to a molecule such as gelsolin are well known in the art. The term "functional derivative" is intended to include the "fragments," "variants," "analogues," or "chemical derivatives" of gelsolin.

[0068] The disclosure involves in some aspects, methods for treating an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) in a subject. The subject is known to have, is suspected of having, or is at risk of having the inflammatory disease. The gelsolin is administered in an amount effective to treat the inflammatory disease in the subject.

[0069] A response to a treatment method of the disclosure can, for example, be measured by determining the physi-

ological effects of the treatment, such as the decrease or lack of symptoms following administration of the treatment.

[0070] In another aspect of the disclosure, a method for monitoring therapy in a subject is provided. The method involves obtaining a level of gelsolin in a subject undergoing therapy to treat an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). The level of gelsolin is compared to a predetermined value corresponding to a control level of gelsolin (e.g., in an apparently healthy population). A determination of whether the level of gelsolin is at or below a predetermined level is indicative of whether the subject would benefit from continued therapy with the same therapy or would benefit from a change in therapy. In some embodiments, obtaining a level of gelsolin is repeated so as to monitor the subject's levels of gelsolin over time. In some embodiments, the subject may have been undergoing the therapy for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 weeks or more. In some embodiments, the subject may have been undergoing the therapy for at least 3, 4, 5, 6 months, or more.

[0071] A change in therapy with gelsolin refers to an increase in the dose of the gelsolin, a switch from one gelsolin to another gelsolin, a switch from gelsolin to another agent, the addition of another agent to the gelsolin therapeutic regimen, or a combination thereof.

[0072] According to another aspect of the disclosure, a method for evaluating the efficacy of a therapy for treating or reducing the risk of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) is provided. The method involves obtaining a level of gelsolin in a subject undergoing therapy to treat the inflammatory disease. The level of gelsolin is compared to a predetermined value corresponding to a control level of gelsolin (e.g., in an apparently healthy population). A determination that the level of gelsolin is at or above a predetermined level is indicative that the therapy is efficacious. In some embodiments, the subject may have been undergoing the therapy for at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 weeks or more. In some embodiments, the subject may have been undergoing the therapy for at least 3, 4, 5, 6 months, or more.

[0073] One aspect of the disclosure is directed to the measurement of gelsolin to guide treatments in order to improve outcome in subjects. On-therapy levels of gelsolin have predictive value for response to treatments of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). The on-therapy levels of gelsolin are additive to prior art predictors of outcome of the disease.

[0074] Subjects who would benefit from this aspect of this disclosure are subjects who are undergoing therapy to treat or prevent the inflammatory disease such as, for example, rheumatoid arthritis (i.e., a subject "on-therapy"). A subject on-therapy is a subject who already has been diagnosed and is in the course of treatment with a therapy for treating an inflammatory disease such as rheumatoid arthritis. The therapy can be any of the therapeutic agents referred to herein. The therapy also can be non-drug treatments. In important embodiments, the therapy is one which increases levels of gelsolin. In a particularly important embodiment, the therapy is a therapy with gelsolin. Preferred subjects are human subjects. The subject most likely to benefit from this disclosure is a human subject on-therapy and who has a gelsolin level at or below about 250 mg/L (or 2.4 μ M/L) of plasma.

[0075] In some embodiments, the subject already has the disease. In some embodiments, the subject may be at an elevated risk of having the disease.

[0076] Risk factors for diseases are known to those of ordinary skill in the art. For example, risk factors for rheumatoid arthritis include: age (between 25 and 45 years), female gender, Caucasian or native American ethnicity, obesity, and a positive family history. The degree of risk of rheumatoid arthritis depends on the multitude and the severity or the magnitude of the risk factors that the subject has. Risk charts and prediction algorithms are available for assessing the risk of inflammatory diseases such as rheumatoid arthritis in a subject based on the presence and severity of risk factors. In some embodiments, the subject who is at an elevated risk of having the inflammatory disease may be an apparently healthy subject. An apparently healthy subject is a subject who has no signs or symptoms of disease.

[0077] Other methods of assessing the risk of an inflammatory disease in a subject are known by those of ordinary skill in the art.

[0078] The preferred treatment of the instant disclosure is gelsolin. Gelsolin may be administered alone, in a pharmaceutical composition or combined with other therapeutic regimens. Gelsolin and optionally other therapeutic agent(s) may be administered simultaneously or sequentially. When the other therapeutic agents are administered simultaneously they can be administered in the same or separate formulations, but are administered at the same time. The other therapeutic agents may be administered sequentially with one another and with gelsolin when the administration of the other therapeutic agents and the gelsolin is temporally separated. The separation in time between the administration of these compounds may be a matter of minutes or it may be longer.

[0079] In practicing certain methods of the present invention, it is required to obtain a level of gelsolin in a subject. This level then is compared to a predetermined value, wherein the level of gelsolin in comparison to the predetermined value is indicative of the likelihood that the subject will benefit from continued therapy. The subject then can be characterized in terms of the net benefit likely to be obtained from a change in therapy.

[0080] The level of the gelsolin for the subject can be obtained by any art recognized method. Typically, the level is determined by measuring the level of gelsolin in a body fluid, for example, blood, serum, plasma, lymph, saliva, urine, synovial fluid and the like. The level can be determined by ELISA, or other immunoassays or other conventional techniques

for determining the presence of gelsolin. Conventional methods may include sending a sample(s) of a subject's body fluid to a commercial laboratory for measurement. Methods for measuring gelsolin are described in the Example.

[0081] The invention also involves comparing the level of gelsolin for the subject with a predetermined value. The predetermined value can take a variety of forms. It can be single cut-off value, such as a median or mean. It can be established based upon comparative groups, such as, for example, where the risk in one defined group is double the risk in another defined group. It can be a range, for example, where the tested population is divided equally (or unequally) into groups, such as a low-risk group, a medium-risk group and a high-risk group, or into quartiles, the lowest quartile being subjects with the highest risk and the highest quartile being subjects with the lowest risk, or into tertiles the lowest tertile being subjects with the highest risk and the highest tertile being subjects with the lowest risk. The predetermined value may be a cut-off value which is predetermined by the fact that a group having a gelsolin level no less than the cut-off value demonstrates a statistically significant increase in the risk of developing an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) as compared to a comparative group. In some embodiments the comparative group is a group having a lower level of gelsolin.

[0082] The predetermined value can depend upon the particular population of subjects selected. For example, an apparently healthy population may have a different 'normal' range of gelsolin than will populations of subjects which have other conditions. Accordingly, the predetermined values selected may take into account the category in which a subject falls. Appropriate ranges and categories can be selected with no more than routine experimentation by those of ordinary skill in the art. The preferred body fluid is blood. In some embodiments, the predetermined value of gelsolin is about 250 mg/L of plasma or lower. In some embodiments, the predetermined value of gelsolin is about 240 mg/L, 230 mg/L, 220 mg/L, 210 mg/L, 200 mg/L, 190 mg/L, 180 mg/L, 170 mg/L, 160 mg/L, 150 mg/L, 140 mg/L, 130 mg/L, 120 mg/L, 110 mg/L, 100 mg/L, 90 mg/L, 80 mg/L, 70 mg/L, 60 mg/L, 50 mg/L, 40 mg/L, 30 mg/L, 20 mg/L, or 10 mg/L of plasma or lower.

[0083] In some embodiments, the predetermined value of gelsolin is about 2.4 μ M/L of plasma or lower. In some embodiments, the predetermined value of gelsolin is about 2.3 μ M/L, 2.2 μ M/L, 2.1 μ M/L, 2.0 μ M/L, 1.9 μ M/L, 1.8 μ M/L, 1.7 μ M/L, 1.6 μ M/L, 1.5 μ M/L, 1.4 μ M/L, 1.3 μ M/L, 1.2 μ M/L, 1.1 μ M/L, 1.0 μ M/L, 0.9 μ M/L, 0.8 μ M/L, 0.7 μ M/L, 0.6 μ M/L, 0.5 μ M/L, 0.4 μ M/L, 0.3 μ M/L, 0.2 μ M/L of plasma or lower.

[0084] An important predetermined value of gelsolin is a value that is the average for a healthy subject population (i.e., subjects who have no signs and symptoms of disease). The predetermined value will depend, of course, upon the characteristics of the subject population in which the subject lies. In characterizing risk, numerous predetermined values can be established.

[0085] Presently, there are commercial sources which produce reagents for assays for gelsolin. These include, for example, Cytoskeleton (Denver, CO), Sigma (St. Louis, MO) and Calbiochem (San Diego, CA).

[0086] In some embodiments, the invention further comprises measuring the level of gelsolin together with a level of a second marker of an inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments). Markers for inflammatory diseases are known to those of ordinary skill in the art and examples of which are described above. Examples of markers for rheumatoid arthritis include, for example, anti cyclic citrullinated peptide (anti-CCP) antibodies, HLA-DR4, and C-reactive protein (CRP). A level of gelsolin in the subject is obtained. The level of gelsolin is compared to a predetermined value to establish a first risk value. A level of the second marker inflammatory disease in the subject is also obtained. The level of the second marker inflammatory disease in the subject is compared to a second predetermined value to establish a second risk value. The subject's risk profile of developing the inflammatory disease (e.g., rheumatoid arthritis in preferred embodiments) then is characterized based upon the combination of the first risk value and the second risk value, wherein the combination of the first risk value and second risk value establishes a third risk value different from the first and second risk values. In some embodiments, the third risk value is greater than either of the first and second risk values. The preferred subjects for testing and predetermined values are as described above. The disease may be any of the inflammatory disease described above.

[0087] The disclosure provides methods for determining whether a subject will benefit from continued therapy or would benefit from a change in therapy. The benefit is typically a reduction in the signs and symptoms or a faster recovery from the manifestations of the disease. Signs, symptoms and manifestations of disease are known to those of ordinary skill in the art. For example, in rheumatoid arthritis, signs and symptoms of the disease include pain, swelling and tenderness in the affected joint(s).

[0088] These methods have important implications for patient treatment and also for the clinical development of new therapies. Determining whether a subject will benefit from continued therapy or would benefit from a change in therapy is clinically useful. One example of clinical usefulness of the methods of this invention includes identifying subjects who are less likely or more likely to respond to a therapy. The methods of the disclosure are also useful in predicting or determining that a subject would benefit from continued therapy or would benefit from a change in therapy. Health care practitioners select therapeutic regimens for treatment based upon the expected net benefit to the subject. The net benefit is derived from the risk to benefit ratio. The present invention permits the determination of whether a subject will benefit from continued therapy or would benefit from a change in therapy, thereby aiding the physician in selecting a

therapy.

[0089] Another example of clinical usefulness, in the case of human subjects for example, includes aiding clinical investigators in the selection for clinical trials of subjects with a high likelihood of obtaining a net benefit. It is expected that clinical investigators now will use the present invention for determining entry criteria for clinical trials.

[0090] A subject who would benefit from continued therapy is a subject whose on-therapy level of gelsolin reaches a certain predetermined value or whose level of gelsolin is increasing. Predetermined values of gelsolin are described above. A subject who would benefit from a change in therapy is a subject whose on-therapy level of the gelsolin did not reach a certain predetermined value or whose on-therapy level of gelsolin is not increasing.

[0091] As used herein, a "change in therapy" refers to an increase or decrease in the dose of the existing therapy, a switch from one therapy to another therapy, an addition of another therapy to the existing therapy, or a combination thereof. A switch from one therapy to another may involve a switch to a therapy with a high risk profile but where the likelihood of expected benefit is increased. In some embodiments, preferred therapies are therapies that increase the levels of gelsolin. A subject who would benefit from a change in therapy by increasing the dose of the existing therapy is a subject who, for example, was on the therapy but was not receiving the maximum tolerated dose or the maximum allowed dose of the therapy and whose level of gelsolin did not reach a certain predetermined value. In such instances the dose of the existing therapy is increased until the level of gelsolin reaches a certain predetermined value. In some instances, the dose of the existing therapy is increased from the existing dose to a higher dose that is not the maximum tolerated dose nor the maximum allowed dose of the therapy. In other instances, the dose is increased to the maximum tolerated or to the maximum allowed dose of the therapy. A subject who would benefit from a change in therapy by decreasing the dose of the existing therapy is, for example, a subject whose on-therapy level of gelsolin reaches or can reach a certain predetermined value with a lower dose of the therapy.

[0092] A subject who would benefit from a switch from one therapy to another therapy is, for example, a subject who was on the maximum tolerated dose or the maximum allowed dose of the therapy and whose level of gelsolin did not reach a certain predetermined value. Another example is a subject was not on the maximum tolerated or the maximum allowed dose of the therapy but was determined by a health care practitioner to more likely benefit from another therapy. Such determinations are based, for example, on the development in the subject of unwanted side effects on the initial therapy or a lack of response to the initial therapy.

[0093] A subject who would benefit from a change in therapy by the addition of another therapy to the existing therapy is, for example, a subject who was on a therapy but whose level of gelsolin did not reach a certain predetermined value. In such instances, another therapy is added to the existing therapy. The therapy that is added to the existing therapy can have a different mechanism of action in increasing the level of gelsolin than the existing therapy. In some instances, a combination of the aforementioned changes in therapy may be used.

[0094] The disclosure also provides methods for determining the efficacy of a therapy. The efficacy is typically the efficacy of the therapy in increasing the level of gelsolin. This is sometimes also referred to as a positive response or a favorable response. Efficacy can be determined by a gelsolin blood test(s) to determine whether gelsolin levels are increased as a result of therapy. In some embodiments efficacy determination is based on the efficacy of a therapy in increasing both gelsolin and normalizing levels of markers of inflammation and/or normalizing white blood cell (WBC) counts.

[0095] The gelsolin measurement is typically reported in $\mu\text{M/L}$ (micromoles/Liter), mg/dl (milligrams/deciliter), or mg/L (milligrams/Liter).

[0096] The disclosure also provides methods for deciding on the course of a therapy in a subject undergoing therapy for an inflammatory disease such as rheumatoid arthritis. Such a course of therapy is decided on the basis of the level of gelsolin. In some embodiments, the subject already has the disease or is at risk of having the inflammatory disease. In some embodiments, the subject is at an elevated risk of having the inflammatory disease the subject has one or more risk factors to have the disease.

[0097] The amount of a treatment may be varied for example by increasing or decreasing the amount of gelsolin or pharmacological agent or a therapeutic composition, by changing the therapeutic composition administered, by changing the route of administration, by changing the dosage timing and so on. The effective amount will vary with the particular condition being treated, the age and physical condition of the subject being treated, the severity of the condition, the duration of the treatment, the nature of the concurrent therapy (if any), the specific route of administration, and like factors are within the knowledge and expertise of the health practitioner. For example, an effective amount can depend upon the duration the individual has had the inflammatory disease.

[0098] An effective amount is a dosage of the therapeutic agent sufficient to provide a medically desirable result. An effective amount may also, for example, depend upon the degree to which an individual has abnormally decreased levels of gelsolin. It should be understood that the therapeutic agents of the invention are used to treat or prevent the inflammatory disease (such as rheumatoid arthritis), that is, they may be used prophylactically in subjects at risk of developing the inflammatory disease (such as rheumatoid arthritis). Thus, an effective amount is that amount which can lower the risk of, slow or perhaps prevent altogether the development of the inflammatory disease (such as rheumatoid

arthritis). It will be recognized when the therapeutic agent is used in acute circumstances, it is used to prevent one or more medically undesirable results that typically flow from such adverse events.

5 [0099] The factors involved in determining an effective amount are well known to those of ordinary skill in the art and can be addressed with no more than routine experimentation. It is generally preferred that a maximum dose of the pharmacological agents of the invention (alone or in combination with other therapeutic agents) be used, that is, the highest safe dose according to sound medical judgment. It will be understood by those of ordinary skill in the art however, that a patient may insist upon a lower dose or tolerable dose for medical reasons, psychological reasons or for virtually any other reasons.

10 [0100] The therapeutically effective amount of a pharmacological agent of the invention is that amount effective to treat the inflammatory disease. For example, in the case of rheumatoid arthritis, the desired response is inhibiting the progression of rheumatoid arthritis. This may involve only slowing the progression of rheumatoid arthritis temporarily, although more preferably, it involves halting the progression of the rheumatoid arthritis permanently. This can be monitored by routine diagnostic methods known to those of ordinary skill in the art. The desired response to treatment of rheumatoid arthritis also can be delaying the onset or even preventing the onset of rheumatoid arthritis.

15 [0101] The pharmacological agents of the invention are preferably sterile and contain an effective amount of gelsolin for producing the desired response in a unit of weight or volume suitable for administration to a subject. The doses of pharmacological agents administered to a subject can be chosen in accordance with different parameters, in particular in accordance with the mode of administration used and the state of the subject. Other factors include the desired period of treatment. In the event that a response in a subject is insufficient at the initial doses applied, higher doses (or effectively higher doses by a different, more localized delivery route) may be employed to the extent that patient tolerance permits. The dosage of a pharmacological agent may be adjusted by the individual physician or veterinarian, particularly in the event of any complication. A therapeutically effective amount typically varies from 0.01 mg/kg to about 1000 mg/kg, preferably from about 0.1 mg/kg to about 500 mg/kg, and most preferably from about 0.2 mg/kg to about 250 mg/kg, in one or more dose administrations daily, for one or more days.

20 [0102] Gelsolin and optionally other therapeutics may be administered per se or in the form of a pharmaceutically acceptable salt.

25 [0103] Various modes of administration are known to those of ordinary skill in the art which effectively deliver the pharmacological agents of the invention to a desired tissue, cell, or bodily fluid. The administration methods are discussed elsewhere in the application. The invention is not limited by the particular modes of administration disclosed herein. Standard references in the art (e.g., Remington's Pharmaceutical Sciences, 20th Edition, Lippincott, Williams and Wilkins, Baltimore MD, 2001) provide modes of administration and formulations for delivery of various pharmaceutical preparations and formulations in pharmaceutical carriers. Other protocols which are useful for the administration of pharmacological agents of the invention will be known to one of ordinary skill in the art, in which the dose amount, schedule of administration, sites of administration, mode of administration and the like vary from those presented herein.

30 [0104] Administration of pharmacological agents of the invention to mammals other than humans, e.g. for testing purposes or veterinary therapeutic purposes, is carried out under substantially the same conditions as described above. It will be understood by one of ordinary skill in the art that this invention is, applicable to both human and animal diseases. Thus, this invention is intended to be used in husbandry and veterinary medicine as well as in human therapeutics.

35 [0105] When administered, the pharmaceutical preparations of the invention are applied in pharmaceutically-acceptable amounts and in pharmaceutically-acceptable compositions. The term "pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients. Such preparations may routinely contain salts, buffering agents, preservatives, compatible carriers, and optionally other therapeutic agents. When used in medicine, the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts may conveniently be used to prepare pharmaceutically-acceptable salts thereof and are not excluded from the scope of the invention. Such pharmacologically and pharmaceutically-acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, maleic, acetic, salicylic, citric, formic, malonic, succinic, and the like. Also, pharmaceutically-acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts.

40 [0106] A pharmacological agent or composition may be combined, if desired, with a pharmaceutically-acceptable carrier. The term "pharmaceutically-acceptable carrier" as used herein means one or more compatible solid or liquid fillers, diluents or encapsulating substances which are suitable for administration into a human. The term "carrier" denotes an organic or inorganic ingredient, natural or synthetic, with which the active ingredient is combined to facilitate the application. The components of the pharmaceutical compositions also are capable of being co-mingled with the pharmacological agents of the invention, and with each other, in a manner such that there is no interaction which would substantially impair the desired pharmaceutical efficacy.

45 [0107] The pharmaceutical compositions may contain suitable buffering agents, as described above, including: acetate, phosphate, citrate, glycine, borate, carbonate, bicarbonate, hydroxide (and other bases) and pharmaceutically acceptable salts of the foregoing compounds. The pharmaceutical compositions also may contain, optionally, suitable preservatives,

such as: benzalkonium chloride, chlorobutanol, parabens and thimerosal.

[0108] The pharmaceutical compositions may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active agent into association with a carrier, which constitutes one or more accessory ingredients. In general, the compositions are prepared by uniformly and intimately bringing the active compound into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product.

[0109] The compounds, when it is desirable to deliver them systemically, may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0110] Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0111] Alternatively, the active compounds may be in powder form for constitution with a suitable vehicle (e.g., saline, buffer, or sterile pyrogen-free water) before use.

[0112] Compositions suitable for oral administration may be presented as discrete units, such as capsules, tablets, pills, lozenges, each containing a predetermined amount of the active compound (e.g., gelsolin). Other compositions include suspensions in aqueous liquids or non-aqueous liquids such as a syrup, elixir, an emulsion, or a gel.

[0113] Pharmaceutical preparations for oral use can be obtained as solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are, in particular, fillers such as sugars, including lactose, sucrose, mannitol, sorbitol or cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate. Optionally the oral formulations may also be formulated in saline or buffers, i.e. EDTA for neutralizing internal acid conditions or may be administered without any carriers.

[0114] Also specifically contemplated are oral dosage forms of the above component or components. The component or components may be chemically modified so that oral delivery of the derivative is efficacious. Generally, the chemical modification contemplated is the attachment of at least one moiety to the component molecule itself, where said moiety permits (a) inhibition of proteolysis; and (b) uptake into the blood stream from the stomach or intestine. Also desired is the increase in overall stability of the component or components and increase in circulation time in the body. Examples of such moieties include: polyethylene glycol, copolymers of ethylene glycol and propylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone and polyproline. Abuchowski and Davis, 1981, "Soluble Polymer-Enzyme Adducts" In: Enzymes as Drugs, Hochenberg and Roberts, eds., Wiley-Interscience, New York, NY, pp. 367-383; Newmark, et al., 1982, J. Appl. Biochem. 4:185-189. Other polymers that could be used are poly-1,3-dioxolane and poly-1,3,6-tioxocane. Preferred for pharmaceutical usage, as indicated above, are polyethylene glycol moieties.

[0115] For the component (or derivative) the location of release may be the stomach, the small intestine (the duodenum, the jejunum, or the ileum), or the large intestine. One skilled in the art has available formulations which will not dissolve in the stomach, yet will release the material in the duodenum or elsewhere in the intestine. Preferably, the release will avoid the deleterious effects of the stomach environment, either by protection of gelsolin or by release of the biologically active material beyond the stomach environment, such as in the intestine.

[0116] To ensure full gastric resistance a coating impermeable to at least pH 5.0 is essential. Examples of the more common inert ingredients that are used as enteric coatings are cellulose acetate trimellitate (CAT), hydroxypropylmethylcellulose phthalate (HPMCP), HPMCP 50, HPMCP 55, polyvinyl acetate phthalate (PVAP), Eudragit L30D, Aquateric, cellulose acetate phthalate (CAP), Eudragit L, Eudragit S, and Shellac. These coatings may be used as mixed films.

[0117] A coating or mixture of coatings can also be used on tablets, which are not intended for protection against the stomach. This can include sugar coatings, or coatings which make the tablet easier to swallow. Capsules may consist of a hard shell (such as gelatin) for delivery of dry therapeutic i.e. powder; for liquid forms, a soft gelatin shell may be used. The shell material of cachets could be thick starch or other edible paper. For pills, lozenges, molded tablets or tablet triturates, moist massing techniques can be used.

[0118] The therapeutic can be included in the formulation as fine multi-particulates in the form of granules or pellets of particle size about 1 mm. The formulation of the material for capsule administration could also be as a powder, lightly compressed plugs or even as tablets. The therapeutic could be prepared by compression.

[0119] Colorants and flavoring agents may all be included. For example, gelsolin may be formulated (such as by liposome or microsphere encapsulation) and then further contained within an edible product, such as a refrigerated beverage containing colorants and flavoring agents.

[0120] One may dilute or increase the volume of the therapeutic with an inert material. These diluents could include carbohydrates, especially mannitol, lactose, anhydrous lactose, cellulose, sucrose, modified dextrans and starch. Certain inorganic salts may also be used as fillers including calcium triphosphate, magnesium carbonate and sodium chloride. Some commercially available diluents are Fast-Flo, Emdex, STA-Rx 1500, Emcompress and Avicell.

[0121] Disintegrants may be included in the formulation of the therapeutic into a solid dosage form. Materials used as disintegrants include but are not limited to starch, including the commercial disintegrant based on starch, Explotab. Sodium starch glycolate, Amberlite, sodium carboxymethylcellulose, ultramylopectin, sodium alginate, gelatin, orange peel, acid carboxymethyl cellulose, natural sponge and bentonite may all be used. Another form of the disintegrants are the insoluble cationic exchange resins. Powdered gums may be used as disintegrants and as binders and these can include powdered gums such as agar, Karaya or tragacanth. Alginic acid and its sodium salt are also useful as disintegrants.

[0122] Binders may be used to hold the therapeutic agent together to form a hard tablet and include materials from natural products such as acacia, tragacanth, starch and gelatin. Others include methyl cellulose (MC), ethyl cellulose (EC) and carboxymethyl cellulose (CMC). Polyvinyl pyrrolidone (PVP) and hydroxypropylmethyl cellulose (HPMC) could both be used in alcoholic solutions to granulate the therapeutic.

[0123] An anti-frictional agent may be included in the formulation of the therapeutic to prevent sticking during the formulation process. Lubricants may be used as a layer between the therapeutic and the die wall, and these can include but are not limited to; stearic acid including its magnesium and calcium salts, polytetrafluoroethylene (PTFE), liquid paraffin, vegetable oils and waxes. Soluble lubricants may also be used such as sodium lauryl sulfate, magnesium lauryl sulfate, polyethylene glycol of various molecular weights, Carbowax 4000 and 6000.

[0124] Glidants that might improve the flow properties of the drug during formulation and to aid rearrangement during compression might be added. The glidants may include starch, talc, pyrogenic silica and hydrated silicoaluminate.

[0125] To aid dissolution of the therapeutic into the aqueous environment a surfactant might be added as a wetting agent. Surfactants may include anionic detergents such as sodium lauryl sulfate, dioctyl sodium sulfosuccinate and dioctyl sodium sulfonate. Cationic detergents might be used and could include benzalkonium chloride or benzethonium chloride. The list of potential non-ionic detergents that could be included in the formulation as surfactants are lauromacrogol 400, polyoxyl 40 stearate, polyoxyethylene hydrogenated castor oil 10, 50 and 60, glycerol monostearate, polysorbate 40, 60, 65 and 80, sucrose fatty acid ester, methyl cellulose and carboxymethyl cellulose. These surfactants could be present in the formulation of gelsolin either alone or as a mixture in different ratios.

[0126] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

[0127] Microspheres formulated for oral administration may also be used. Such microspheres have been well defined in the art. All formulations for oral administration should be in dosages suitable for such administration.

[0128] For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

[0129] For administration by inhalation, the compounds for use according to the present invention may be conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebulizer, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g. gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0130] Also contemplated herein is pulmonary delivery of gelsolin. Gelsolin is delivered to the lungs of a mammal while inhaling and traverses across the lung epithelial lining to the blood stream. Other reports of inhaled molecules include Adjei et al., 1990, *Pharmaceutical Research*, 7:565-569; Adjei et al., 1990, *International Journal of Pharmaceutics*, 63:135-144 (leuprolide acetate); Braquet et al., 1989, *Journal of Cardiovascular Pharmacology*, 13(suppl. 5):143-146 (endothelin-1); Hubbard et al., 1989, *Annals of Internal Medicine*, Vol. III, pp. 206-212 (al-antitrypsin); Smith et al., 1989, *J. Clin. Invest.* 84:1145-1146 (a-1 -proteinase); Oswein et al., 1990, "Aerosolization of Proteins", *Proceedings of Symposium on Respiratory Drug Delivery II*, Keystone, Colorado, March, (recombinant human growth hormone); Debs et al., 1988, *J. Immunol.* 140:3482-3488 (interferon- γ and tumor necrosis factor alpha) and Platz et al., U.S. Patent No. 5,284,656 (granulocyte colony stimulating factor). A method and composition for pulmonary delivery of drugs for systemic effect is described in U.S. Patent No. 5,451,569, issued September 19, 1995 to Wong et al.

[0131] Contemplated for use in the practice of this invention are a wide range of mechanical devices designed for

pulmonary delivery of therapeutic products; including but not limited to nebulizers, metered dose inhalers, and powder inhalers, all of which are familiar to those skilled in the art.

5 [0132] Some specific examples of commercially available devices suitable for the practice of this invention are the Ultravent nebulizer, manufactured by Mallinckrodt, Inc., St. Louis, Missouri; the Acorn II nebulizer, manufactured by Marquest Medical Products, Englewood, Colorado; the Ventolin metered dose inhaler, manufactured by Glaxo Inc., Research Triangle Park, North Carolina; and the Spinhaler powder inhaler, manufactured by Fisons Corp., Bedford, Massachusetts.

10 [0133] All such devices require the use of formulations suitable for the dispensing of gelsolin. Typically, each formulation is specific to the type of device employed and may involve the use of an appropriate propellant material, in addition to the usual diluents, adjuvants and/or carriers useful in therapy. Also, the use of liposomes, microcapsules or microspheres, inclusion complexes, or other types of carriers is contemplated. Chemically modified gelsolin may also be prepared in different formulations depending on the type of chemical modification or the type of device employed.

15 [0134] Formulations suitable for use with a nebulizer, either jet or ultrasonic, will typically comprise gelsolin dissolved in water at a concentration of about 0.1 to 25 mg of biologically active gelsolin per mL of solution. The formulation may also include a buffer and a simple sugar (e.g., for gelsolin stabilization and regulation of osmotic pressure). The nebulizer formulation may also contain a surfactant, to reduce or prevent surface induced aggregation of the gelsolin caused by atomization of the solution in forming the aerosol.

20 [0135] Formulations for use with a metered-dose inhaler device will generally comprise a finely divided powder containing the gelsolin suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as a chlorofluorocarbon, a hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol, and, 1,1,1,2-tetrafluoroethane, or combinations thereof. Suitable surfactants include sorbitan trioleate and soya lecithin. Oleic acid may also be useful as a surfactant.

25 [0136] Formulations for dispensing from a powder inhaler device will comprise a finely divided dry powder containing gelsolin and may also include a bulking agent, such as lactose, sorbitol, sucrose, or mannitol in amounts which facilitate dispersal of the powder from the device, e.g., 50 to 90% by weight of the formulation. The gelsolin should most advantageously be prepared in particulate form with an average particle size of less than 10 μ m (or microns), most preferably 0.5 to 5 μ m, for most effective delivery to the distal lung.

30 [0137] Nasal (or intranasal) delivery of a pharmaceutical composition of the present invention is also contemplated. Nasal delivery allows the passage of a pharmaceutical composition of the present invention to the blood stream directly after administering the therapeutic product to the nose, without the necessity for deposition of the product in the lung. Formulations for nasal delivery include those with dextran or cyclodextran.

35 [0138] For nasal administration, a useful device is a small, hard bottle to which a metered dose sprayer is attached. In one embodiment, the metered dose is delivered by drawing the pharmaceutical composition of the present invention solution into a chamber of defined volume, which chamber has an aperture dimensioned to aerosolize and aerosol formulation by forming a spray when a liquid in the chamber is compressed. The chamber is compressed to administer the pharmaceutical composition of the present invention. In a specific embodiment, the chamber is a piston arrangement. Such devices are commercially available.

40 [0139] Alternatively, a plastic squeeze bottle with an aperture or opening dimensioned to aerosolize an aerosol formulation by forming a spray when squeezed is used. The opening is usually found in the top of the bottle, and the top is generally tapered to partially fit in the nasal passages for efficient administration of the aerosol formulation. Preferably, the nasal inhaler will provide a metered amount of the aerosol formulation, for administration of a measured dose of the drug.

45 [0140] The compounds may also be formulated in rectal or vaginal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

[0141] In addition to the formulations described previously, the compounds may also be formulated as a depot preparation. Such long acting formulations may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

50 [0142] The pharmaceutical compositions also may comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers such as polyethylene glycols.

55 [0143] Suitable liquid or solid pharmaceutical preparation forms are, for example, aqueous or saline solutions for inhalation, microencapsulated, encochleated, coated onto microscopic gold particles, contained in liposomes, nebulized, aerosols, pellets for implantation into the skin, or dried onto a sharp object to be scratched into the skin. The pharmaceutical compositions also include granules, powders, tablets, coated tablets, (micro)capsules, suppositories, syrups, emulsions, suspensions, creams, drops or preparations with protracted release of active compounds, in whose preparation excipients and additives and/or auxiliaries such as disintegrants, binders, coating agents, swelling agents, lubricants, flavorings,

sweeteners or solubilizers are customarily used as described above. The pharmaceutical compositions are suitable for use in a variety of drug delivery systems. For a brief review of methods for drug delivery, see Langer, Science 249:1527-1533, 1990. The therapeutic agent(s), including specifically but not limited to gelsolin, may be provided in particles. Particles as used herein means nano or microparticles (or in some instances larger) which can consist in whole or in part of gelsolin or the other therapeutic agent(s) as described herein. The particles may contain the therapeutic agent(s) in a core surrounded by a coating, including, but not limited to, an enteric coating. The therapeutic agent(s) also may be dispersed throughout the particles. The therapeutic agent(s) also may be adsorbed into the particles. The particles may be of any order release kinetics, including zero order release, first order release, second order release, delayed release, sustained release, immediate release, and any combination thereof, etc. The particle may include, in addition to the therapeutic agent(s), any of those materials routinely used in the art of pharmacy and medicine, including, but not limited to, erodible, nonerodible, biodegradable, or nonbiodegradable material or combinations thereof. The particles may be microcapsules which contain the gelsolin in a solution or in a semi-solid state. The particles may be of virtually any shape.

[0144] Both non-biodegradable and biodegradable, polymeric materials can be used in the manufacture of particles for delivering the therapeutic agent(s). Such polymers may be natural or synthetic polymers. The polymer is selected based on the period of time over which release is desired. Bioadhesive polymers of particular interest include bioerodible hydrogels described by H.S. Sawhney, C.P. Pathak and J.A. Hubei' in *Macromolecules*, (1993) 26:581-587. These include polyhyaluronic acids, casein, gelatin, gluten, polyanhydrides, polyacrylic acid, alginate, chitosan, poly(methyl methacrylates), poly(ethyl methacrylates), poly(butylmethacrylate), poly(isobutyl methacrylate), poly(hexylmethacrylate), poly(isodecyl methacrylate), poly(lauryl methacrylate), poly(phenyl methacrylate), poly(methyl acrylate), poly(isopropyl acrylate), poly(isobutyl acrylate), and poly(octadecyl acrylate).

[0145] The therapeutic agent(s) may be contained in controlled release systems. The term "controlled release" is intended to refer to any drug-containing formulation in which the manner and profile of drug release from the formulation are controlled. This refers to immediate as well as non-immediate release formulations, with non-immediate release formulations including but not limited to sustained release and delayed release formulations. The term "sustained release" (also referred to as "extended release") is used in its conventional sense to refer to a drug formulation that provides for gradual release of a drug over an extended period of time, and that preferably, although not necessarily, results in substantially constant blood levels of a drug over an extended time period. The term "delayed release" is used in its conventional sense to refer to a drug formulation in which there is a time delay between administration of the formulation and the release of the drug therefrom. "Delayed release" may or may not involve gradual release of drug over an extended period of time, and thus may or may not be "sustained release."

[0146] Use of a long-term sustained release implant may be particularly suitable for treatment of chronic conditions. "Long-term" release, as used herein, means that the implant is constructed and arranged to deliver therapeutic levels of the active ingredient for at least 7 days, and preferably 30-60 days. Long-term sustained release implants are well-known to those of ordinary skill in the art and include some of the release systems described above.

[0147] For topical administration to the eye, nasal membranes, mucous membranes or to the skin, the gelsolin may be formulated as ointments, creams or lotions, or as a transdermal patch or intraocular insert or iontophoresis. For example, ointments and creams can be formulated with an aqueous or oily base alone or together with suitable thickening and/or gelling agents. Lotions can be formulated with an aqueous or oily base and, typically, further include one or more emulsifying agents, stabilizing agents, dispersing agents, suspending agents, thickening agents, or coloring agents. (See, e.g., U.S. 5,563,153, entitled "Sterile Topical Anesthetic Gel", issued to Mueller, D., et al., for a description of a pharmaceutically acceptable gel-based topical carrier.)

[0148] In general, the gelsolin is present in a topical formulation in an amount ranging from about 0.01% to about 30.0% by weight, based upon the total weight of the composition. Preferably, the gelsolin is present in an amount ranging from about 0.5 to about 30% by weight and, most preferably, the gelsolin is present in an amount ranging from about 0.5 to about 10% by weight. In one embodiment, the compositions of the invention comprise a gel mixture to maximize contact with the surface of the localized pain and minimize the volume and dosage necessary to alleviate the localized pain. GELFOAM® (a methylcellulose-based gel manufactured by Upjohn Corporation) is a preferred pharmaceutically acceptable topical carrier. Other pharmaceutically acceptable carriers include iontophoresis for transdermal drug delivery.

[0149] The disclosure also contemplates the use of kits. In some aspects of the disclosure, the kit can include a pharmaceutical preparation vial, a pharmaceutical preparation diluent vial, and gelsolin. The vial containing the diluent for the pharmaceutical preparation is optional. The diluent vial contains a diluent such as physiological saline for diluting what could be a concentrated solution or lyophilized powder of gelsolin. The instructions can include instructions for mixing a particular amount of the diluent with a particular amount of the concentrated pharmaceutical preparation, whereby a final formulation for injection or infusion is prepared. The instructions may include instructions for treating a subject with an effective amount of gelsolin. It also will be understood that the containers containing the preparations, whether the container is a bottle, a vial with a septum, an ampoule with a septum, an infusion bag, and the like, can contain indicia such as conventional markings which change color when the preparation has been autoclaved or otherwise

sterilized.

[0150] The present invention is further illustrated by the following Examples, which in no way should be construed as further limiting..

5 Examples

[0151] Plasma gelsolin (pGSN) is a secreted protein that circulates in the extracellular fluids of humans at concentrations averaging 250 mg/l. Diverse types of tissue injury lead to prolonged reductions in plasma gelsolin levels. Following severe tissue injury encountered in severe trauma, burns, sepsis, major surgery and hematopoietic stem cell transplant patients, declines in gelsolin (GSN) levels to approximately less than 25% of normal precede and, therefore, predict critical care complications measured by assisted ventilation requirements, length of intensive care unit residence and overall hospital stays, death and specific sequelae such as secondary lung injury (e.g. adult respiratory distress syndrome (ARDS), acute lung injury (ALI), multiple organ dysfunction syndromes (MODS)). Similar plasma gelsolin reductions in animal models precede lung permeability changes and inflammation, and infusion of recombinant plasma gelsolin ameliorates these effects.

Example 1: Plasma gelsolin (pGSN) concentration is decreased in patients with rheumatoid arthritis (RA) compared to healthy controls and lower in synovial fluid than in blood in patients with RA.

[0152] We measured plasma gelsolin (pGSN) levels in patients with rheumatoid arthritis (RA) to experimentally test the hypothesis that plasma gelsolin levels fall in response to the initial (unknown) injury inflicted by the agency causing rheumatoid arthritis. As shown in Figure 1, circulating (pGSN) levels were significantly lower in patients with RA compared to matched healthy controls (103 ± 23 versus 142 ± 29 , $P = 0.0002$, FIG. 1). pGSN levels were similar in both male and females, and were not dependent on the age of the patients or on the duration of arthritis. Circulating pGSN levels were inversely correlated to the levels of C-reactive protein ($r=-0.272$, $p=0.026$).

Materials and Methods:

[0153] Patients: Plasma and synovial fluid samples were collected from 82 RA patients attending the Rheumatology clinics, at Sahlgrenska University Hospital in Gothenburg, for acute joint effusion. RA was diagnosed according to the American College of Rheumatology criteria. At the time of synovial fluid and blood sampling all the patients received non-steroidal anti-inflammatory drugs. Recent radiographs of the hands and feet were obtained for all the patients. Presence of bone erosions defined as the loss of cortical definition of the joint, was recorded in proximal interphalangeal, metacarpophalangeal, carpus, interphalangeal and metatarsophalangeal joints of forefeet. Presence of one erosion was sufficient to fulfil requirement of an erosive disease. Presence of rheumatoid factor of any of immunoglobulin isotypes tested (IgM,-A, -G) was considered as positive. Control blood samples ($n=87$) were obtained from the blood donors attending Blood Transfusion Unit of Sahlgrenska University Hospital and matching the RA patients for age and gender.

[0154] Collection and preparation of samples: Synovial fluid was obtained from knee joints by arthrocentesis, aseptically aspirated and transferred into sodium citrate (0.129mol/l ; pH 7.4) containing tubes. Blood samples were simultaneously obtained from the cubital vein and collected into sodium citrate medium. Collected blood and synovial fluid samples were centrifuged at 800 g for 15 minutes, aliquoted, and stored frozen at -70°C until use.

[0155] Measurement of pGSN concentration in plasma and synovial fluid by pyrene actin-nucleation assay: pGSN is activated by calcium and binds two actin monomers to form a nucleus from which actin polymerizes in pointed (slowest growing) end direction. Pyrene actin was prepared by derivatizing actin with N-pyrenyliodoacetamide (Molecular Probes, Eugene Oregon). Before use pyrene actin was diluted in depolymerization buffer (Buffer A: 0.5 mM ATP, 5 mM β -mercaptoethanol, 0.2 mM Tris, 0.2 mM CaCl_2 , pH 7.4) to 20 μM , stored 1h at 37°C to reach monomer equilibrium and centrifuged at 250,000 g, 4°C for 30 minutes, in an Optima™ TL Ultracentrifuge (Beckman) to pellet any remnant filamentous actin. The supernatant was withdrawn and stored in an ice water bath until used. Platelet poor plasma to be analyzed was diluted 1:5 in buffer B (0.1 M KCl , 0.2 mM MgCl_2 , 0.2 mM CaCl_2 , 0.5 mM ATP, 10 mM Tris, 0.5 mM mercaptoethanol, pH 7.4). Pyrene actin fluorescence was recorded using a luminescence fluorometer (FluoroMax-2®, JobinYvon-Spex Instruments S.A., Inc). Excitation and emission wavelengths were 366 and 386 nm, respectively. Pyrene actin was added to a final concentration of 1 μM in 280 μl buffer B containing 0.4 μM phalloidin, 1.5 mM CaCl_2 and 5 μl of diluted sample in 6x50 mm borosilicate glass culture tubes (Kimble). Nucleation was monitored over 240 s in the fluorometer following a fast vortex. The linear slope of the fluorescence increase was calculated between 100-200s. All the samples were run in duplicates. Polymerization rate in each sample was converted to pGSN concentration by use of a standard curve of recombinant human pGSN.

[0156] Statistics: The level of pGSN in the blood and synovial fluid samples were expressed as mean \pm SD. Comparison between the matched blood and synovial fluid samples were analyzed by the paired t-test. Comparison of pGSN levels

was also performed between the patient blood samples and the healthy controls. For further comparison patient material was stratified according to radiological findings (erosive RA vs. non-erosive RA). Differences in pGSN levels in the blood and synovial fluid between the groups were calculated separately employing the Mann-Whitney U test. For the evaluation of possible influence of the ongoing treatment on the pGSN levels, patient material was stratified according to DMARD treatment (treated vs. untreated). Comparison between the groups was performed using the Mann-Whitney U test. For all the statistical evaluation of the results, P-values below 0.05 were considered statistically significant.

Example 2: The gelsolin isoform present in synovial fluid (SF) of RA patients is the plasma isoform of gelsolin (pGSN)

[0157] Since gelsolin is produced as both an intracellular and extracellular isoform, each of which have the capacity to induce actin polymerization, we analyzed the origin of the gelsolin activity by immunoblotting with an antibody specific for the plasma isoform (a-pGSN). Gelsolin present in SF is composed mainly of the plasma isoform (Fig. 2). Plasma origin of gelsolin was also supported by a correlation between gelsolin levels in plasma and synovial fluid in the matched pair of samples ($r=0.39$, $p=0.0006$). However, pGSN activity in the SF samples from RA patients were significantly lower than that of plasma (pg/ml , 69 ± 18 vs. 103 ± 23 , $p = 0.04$) indicating local consumption.

Materials and Methods

[0158] Immunoblotting: The platelet poor plasmas or synovial fluids were diluted 1:100 in 1X sample buffer (SB, 20% glycerol, 4.6% Tris, 0.25 M Tris-HCl, 0.01% Bromphenol blue, 10% v/v 2-Mercaptoethanol, pH 6.8) for test of gelsolin isoform, vortexed briefly and boiled at 100 °C for 10 minutes. Samples (10 μl) were run on 10% SDS-PAGE (sodium dodecylsulfate polyacrylamide gelelectrophoresis) gels in a modified Laemmli system. Platelet lysate ($2 \times 10^8/\text{ml}$, 5 μl) and human recombinant pGSN were run as negative and positive controls for pGSN respectively. Proteins were separated for 1.5 hour at 120V. Immobilon P membranes (PVDF, 0.45 μm ; Millipore Corp., Bedford, MA) were soaked in methanol for 1 minute and transfer buffer (192 mM glycine, 25 mM Tris, 0.1% Sodium Dodecyl Sulfate, 20% Methanol) for 5 minutes, before transfer. The transfer was carried out at variable voltage, IA for 90 minutes. The membranes were blocked overnight in PBS containing 0.05% Tween-20, 5% Carnation nonfat dry milk and 0.05% sodium azide pH 7.4 at 4°C. For pGSN a polyclonal antibody specific for the human plasma isoform was used (1:2000, 2h, produced in the lab). For all isoforms of GSN a primary monoclonal 2c4 anti-gelsolin antibody was used (1:2500, 2h, produced in the lab). Secondary antibodies used were rabbit-anti-mouse IgG (H+L)-HRP (1:5000, 80 min) and goat-anti-mouse IgG (H+L)-HRP conjugate respectively (1:3300, 80 min, Bio-Rad, Hercules, CA). Membranes were washed 3 times in PBS containing 0.05% Tween-20 between incubations.

[0159] Chemiluminescence detection was done using SuperSignal® West Pico Chemiluminescent Substrate for detection of HRP (PIERCE, Rockford, IL). The membranes were exposed to 2 ml each of SuperSignal® West Pico Stable Peroxide Solution and SuperSignal® West Pico Luminol/Enhancer Solution for 2 minutes. HyBlot CL autoradiography film (Denville Scientific, Inc., Metuchen, NJ) was exposed to the membrane for 1 minute in a FBXC 810 autoradiography cassette (FischerBiotech, Pittsburg, PA). The film was developed using a M35A X-OMAT Processor (Kodak). Example 3: pGSN is decreased at early stages of an experimental mouse model of septic arthritis/sepsis.

[0160] Levels of circulating pGSN were decreased both during streptococcal and staphylococcal infection. This decrease occurred early during the time course of disease, beginning 2 days post-bacterial inoculation both in arthritic and septic groups of animals. Prior to injection (day 0), the average pGSN level of the mice was 200 ± 20 mg/L. By day 2 pGSN levels decreased to 138 ± 16 mg/L for the arthritic *Staphylococcus aureus* (*S. aureus*) treated (7×10^6 cfu/mouse) animals, 157 ± 15 mg/L for the septic *S. aureus* injected animals (3.5×10^7 cfu/mouse), and 152 ± 15 mg/L for the animals injected with *Streptococcus agalactiae* (*Str. agalactiae*) (1×10^7 cfu/mouse). The pGSN levels at day 9 were 141 ± 21 , 151 ± 15 , and 124 ± 13 mg/L, respectively. Only the animals injected with streptococcal bacteria displayed further decrease of the circulating pGSN levels at day 9. pGSN decrease was not related to the intensity of the infection since pGSN concentrations following administration of septic doses (5 times higher than the arthritic ones), which result in higher mortality, were not lower than those observed in the arthritic-dosed mice.

Materials and Methods

[0161] Induction of *S. aureus* arthritis and sepsis: Female 5-6 weeks old NMRI mice were purchased from ALAB (Stockholm, Sweden) and maintained in the animal facility of the Department of Rheumatology, University of Goteborg. They were housed 10-11 animals per cage under standard conditions of temperature and light and fed standard laboratory chow and water ad libitum. *S. aureus*, strain LS-1, originally isolated from a swollen joint of a spontaneously arthritic NZBxW mouse, strain Newman, as well as *Str. agalactiae* strain 6313, a clinical isolate belonging to serotype III, were used for the induction of septic arthritis and sepsis. Bacteria were kept frozen at -20°C, in PBS (0.13 M sodium chloride, 10 mM sodium phosphate, pH 7.4), containing 5% BSA and 10% dimethyl sulfoxide, until used. Before the injection, the

EP 2 001 496 B1

bacterial solution was thawed, washed twice with PBS, and adjusted with PBS to the desired concentration. Mice were injected into the tail vein with a suspension of *S. aureus* or *Str. agalactiae* in 0.2 ml PBS. Viable counts in the leftover solution were determined to check the exact number of bacteria injected and presented as colony forming units (cfu/ml).

[0162] All the mice were monitored individually during the observation period of 8-9 days by assessment of joint appearance, weight, general appearance, alertness, and skin abnormalities. Blood samples were obtained from the tail vein in a sterile tube without anticoagulant and remained to clot for 6-8 hours. The samples were centrifuged at 3000xg for 15 min, serum was aliquoted and kept frozen at -70° C until used.

[0163] Experimental protocol: Twenty mice obtained intravenously (i.v.) a septic (LS-1, 4x10⁷ cfu/mouse, n=10) or arthritogenic (7x10⁶ cfu/mouse, n=10) dose of *S. aureus*. Additional 10 mice obtained intravenously a septic dose of *Str. agalactiae* (1x10⁷ cfu/mouse). On days -3, 2, 4, 6, and 9 blood samples were collected by the tail vein incision for the determination pGSN levels. On day 9 all mice were sacrificed by cervical dislocation. pGSN level was determined with the same method as for the human samples (pyrene actin nucleation assay).

Example 4: pGSN supplementation to mice delays development of arthritis.

[0164] Supplementation of mice with recombinant gelsolin was performed over 7 days with 24h interval starting immediately before bacterial inoculation (4x10⁷/m1). Five mice received gelsolin while 7 mice received PBS and were used as controls. During the first 9 days of staphylococcal infection no difference was observed in gelsolin-treated and the control groups with respect to weight loss, survival rate or development of arthritis. However, fewer of the gelsolin-treated mice developed arthritis on day 3 (1 treated vs 4 non-treated - See Table 1).

Table 1. Dynamics of weight, survival and development of arthritis in NMRI mice infected intravenously with *S. aureus*.

G = plasma gelsolin, C = Control.

	Day 1	Day 3	Day 5	Day 7	Day 9	Day 28	
Gelsolin	31,5g	30,0g	27,2g	25,3g	25,7g	30,8g	Mean weight
PBS controls	31,2	28,5	27,7	22,9	25,4	32,4	Mean weight
	5G/7C	5G/7C	5G/7C	5G/7C	3G/6C	2G/6C	Survival
	1G/0C	1 G/4C	2G/4C	3G/4C	2G/1C	2G/2C	Arthritis

Materials and Methods

[0165] Twelve mice were injected intravenously (i.v.) with a septic dose of *S. aureus* (Newman, 4x10⁷ cfu/mouse). Five mice received recombinant human pGSN infra-peritoneally (i.p.) during days 0-7 and the remaining 7 mice received PBS. The mice were monitored individually up to day 28 and signs of arthritis, weight, general appearance, and alertness were followed.

[0166] Recombinant pGSN was produced in *E. coli*, refolded with GS SG, formulated in 0.1 M NaCl and 1 mM CaCl₂ by Biogen-Idec, Inc. (Cambridge, MA) and kept at -70° C prior to use. Mice were injected with recombinant pGSN i.p. (6 mg/mouse) one hour prior to the bacterial challenge and continued with 24-hours interval (3 mg/mouse) during 7 days. The controls received a corresponding volume of PBS.

Example 5:

[0167] The hypothesis that administration of gelsolin could impact upon the inflammatory processes is tested in a rodent model of collagen induced arthritis (CIA), an autoimmune model that resembles rheumatoid arthritis. CIA is inducible in inbred DBA/1 male mice by priming intradermally with heterologous or homologous collagen II (about 50 microgram) in Freund's complete adjuvant and 2 weeks later boosting with the same amount of collagen II in Freund's incomplete adjuvant. The arthritis develops approximately 3 weeks after the priming dose and reaches its maximum within 8 weeks post priming. The mice have high levels of collagen II specific antibodies, collagen II specific T cells as well as signs of systemic inflammation (e.g. production of IL-6, TNF etc). Locally in the joints one observes both overwhelming inflammatory infiltrates (consisting of T cells, macrophages, neutrophils and fibroblasts) as well as severe destruction of cartilage and subchondral bone. These features mimic well the process seen in human rheumatoid arthritis (Myers et al., Life Sciences 61, p 1861-1878, 1997). In a therapeutic test one set of test animals receives, for example, subcutaneously 8 mg of bovine serum albumin or 8 mg human recombinant plasma gelsolin once on day ten from the start of therapy (1x) or three doses on days 2, 5 and 10 (3x). This route of administration and dosing has previously been

shown to raise gelsolin levels depleted 50% by sepsis to normal. Several parameters of the disease (e.g., clinical signs and symptoms, onset, progression, severity, and remission of symptoms) are measured. In summary, our findings support the two aspects of the hypothesis posed, namely, that reductions in plasma gelsolin levels precede manifestations of inflammatory diseases such as rheumatoid arthritis and that systemic treatment with plasma gelsolin prevents and/or suppresses these manifestations. One clinical correlate of these observations is that serial monitoring of plasma gelsolin levels could become part of the management strategy of inflammatory diseases such as rheumatoid arthritis, flagging when to intensify therapy. Another correlate is that prophylactic elevation of plasma gelsolin levels might protect patients from the sequelae of inflammation.

[0168] Although not intending to be bound by any particular mechanism or theory, it is presumed that plasma gelsolin is depleted from the blood during inflammatory diseases (such as rheumatoid arthritis (RA)) and is localized or sequestered at the site of injury/inflammation (the joint space in rheumatoid arthritis). It is believed that plasma gelsolin functions at the site of inflammation by binding to inflammatory mediators and prevents them from causing further damage by inhibiting their actions on cellular receptors. This is supported by the fact that plasma gelsolin in vitro binds to some inflammatory mediators such as lysophosphatidic acid (LPA), A β 3 (Alzheimer) peptide, diadenosine 5',5"-P $_1$,P $_3$ -triphosphate (Ap $_3$ A), fibronectin, fibrinogen and lipopolysaccharide (LPS) and decreases certain cellular responses to platelet activating factor (PAF). Additionally, damaged cells at the inflamed joint release actin and gelsolin binds to actin. Gelsolin may have a protective role by severing filamentous actin that might otherwise be toxic. We believe that gelsolin has an anti-inflammatory effect, but when the inflammation is severe and prolonged as in RA, gelsolin supplementation might be beneficial in reducing or treating the inflammation. Sequestration or recruitment of gelsolin to the site of inflammation (such as the joint space in rheumatoid arthritis) could explain why the plasma levels are reduced although other explanations are also possible.

Equivalents

[0169] The foregoing written specification is considered to be sufficient to enable one ordinarily skilled in the art to practice the invention. The present invention is not to be limited in scope by the example(s) provided, since the example(s) are intended as mere illustrations of one or more aspects of the invention. Other functionally equivalent embodiments are considered within the scope of the invention. Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description. Each of the limitations of the invention can encompass various embodiments of the invention. It is, therefore, anticipated that each of the limitations of the invention involving any one element or combinations of elements can be included in each aspect of the invention. This invention is not limited in its application to the details of construction and the arrangement of components set forth or illustrated in the drawings. The invention is capable of other embodiments and of being practiced or of being carried out in various ways.

[0170] Also, the phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of "including," "comprising," or "having," "containing", "involving", and variations thereof herein, is meant to encompass the items listed thereafter and equivalents thereof as well as additional items.

Claims

1. A method for characterizing a healthy subject's risk profile of developing a future inflammatory disease, comprising:

comparing a level of gelsolin in a body fluid (for example blood, plasma, serum, urine, synovial fluid, or alveolar fluid), or tissue (for example a joint, gastrointestinal, thyroid, adrenal, vascular, pulmonary, renal, cardiac, skin, ocular, brain, pancreatic, liver, nerve, or muscle tissue) obtained from the subject to a predetermined value (preferably of about 250 mg/L of plasma or less), and

characterizing the subject's risk profile of developing an inflammatory disease based upon the level of gelsolin in comparison to the predetermined value, wherein a level of gelsolin at or below the predetermined level is indicative that the subject is at an elevated risk of developing the inflammatory disease, and wherein a level of gelsolin at or above the predetermined level is indicative that the subject is not at an elevated risk of developing the inflammatory disease,

wherein the subject does not have any signs or symptoms of an infection, sepsis or an actin-related disorder.

2. The method of claim 1, wherein the inflammatory disease is arthritis, rheumatoid arthritis, asthma, inflammatory bowel disease (Crohn's disease or ulcerative colitis), chronic obstructive pulmonary disease (COPD), allergic rhinitis, vasculitis (polyarteritis nodosa, temporal arteritis, Wegener's granulomatosis, Takayasu's arteritis, or Behcet syndrome), inflammatory neuropathy, psoriasis, systemic lupus erythematosus (SLE), chronic thyroiditis, Hashimoto's

thyroiditis, Addison's disease, polymyalgia rheumatica, Sjogren's syndrome, or Churg-Strauss syndrome.

3. The method of claim 1 further comprising performing one or more tests to evaluate the inflammatory disease, preferably wherein the test is measuring a level of a marker of inflammation in the subject, for example wherein the marker of inflammation is CRP, soluble intercellular adhesion molecule (sICAM-1), ICAM 3, BL-CAM, LFA-2, VCAM-I, NCAM, PECAM, fibrinogen, serum amyloid A (SAA), lipoprotein associated phospholipase A2 (LpPIA2), sCD40 ligand (sCD40L), myeloperoxidase, Interleukin-6 (IL-6), or Interleukin-8 (IL-8).
4. A method for characterizing a healthy subject's risk profile of developing a future inflammatory disease, comprising:
 - comparing the level of the gelsolin in a sample obtained from a subject to a first predetermined value (preferably about 250 mg/L of plasma or lower) to establish a first risk value,
 - comparing the level of a second marker of inflammation in a sample obtained from a subject to a second predetermined value to establish a second risk value, and
 - characterizing the subject's risk profile of developing the inflammatory disease based upon the combination of the first risk value and the second risk value, wherein the combination of the first risk value and second risk value establishes a third risk value different from said first and second risk values, wherein the subject does not have any signs or symptoms of an infection, sepsis or an actin-related disorder.
5. The method of claim 4, wherein the first predetermined value is a plurality of predetermined gelsolin level ranges, one of said plurality being below about 250 mg/L of plasma and another of said ranges being above about 250 mg/L of plasma, and wherein said comparing step comprises determining in which of said plurality of predetermined gelsolin level ranges said subject's gelsolin level falls.
6. Gelsolin, optionally in combination with a second agent for treating an inflammatory disease, for use in a method for treating a subject at risk of developing an inflammatory disease, who is preferably otherwise free of indications calling for treatment with gelsolin, preferably wherein the gelsolin is plasma gelsolin (pGSN), cytoplasmic gelsolin (cGSN), advillin, villin, capG, flightless proteins, fragmin, severin, adseverin, protovillin, or supervillin.
7. The gelsolin of claim 6, wherein the inflammatory disease is arthritis, rheumatoid arthritis, asthma, inflammatory bowel disease (Crohn's disease or ulcerative colitis), chronic obstructive pulmonary disease (COPD), allergic rhinitis, vasculitis (polyarteritis nodosa, temporal arteritis, Wegener's granulomatosis, Takayasu's arteritis, or Behcet syndrome), inflammatory neuropathy, psoriasis, systemic lupus erythematosus (SLE), chronic thyroiditis, Hashimoto's thyroiditis, Addison's disease, polymyalgia rheumatica, Sjogren's syndrome, or Churg-Strauss syndrome.
8. The gelsolin of claim 6, wherein the second agent for treating the inflammatory disease is Alclufenac, Alclometasone Dipropionate, Algestone Acetonide, Alpha Amylase, Amcinafal, Amcinafide, Amfenac Sodium, Amiprilose Hydrochloride, Anakinra, Anirolac, Anitrazafen, Apazone, Balsalazide Disodium, Bendazac, Benoxaprofen, Benzylamine Hydrochloride, Bromelains, Broperamol, Budesonide, Carprofen, Cicloprofen, Cintazone, Cliprofen, Clobetasol Propionate, Clobetasone Butyrate, Clopirac, Cloticasone Propionate, Cormethasone Acetate, Cortodoxone, Cyclooxygenase-2 (COX-2) inhibitor, Deflazacort, Desonide, Desoximetasone, Dexamethasone Dipropionate, Diclofenac Potassium, Diclofenac Sodium, Diflorasone Diacetate, Diflumidone Sodium, Diflunisal, Difluprednate, Diftalone, Dimethyl Sulfoxide, Drocinnonide, Endrysone, Enlimomab, Enolicam Sodium, Epirizole, Etodolac, Etofenamate, Felbinac, Fenamole, Fenbufen, Fenclofenac, Fenclorac, Fendosal, Fempipalone, Fentiazac, Flazalone, Fluzacort, Flufenamic Acid, Flumizole, Flunisolide Acetate, Flunixin, Flunixin Meglumine, Fluocortin Butyl, Fluorometholone Acetate, Fluquazone, Flurbiprofen, Fluretofen, Fluticasone Propionate, Furaprofen, Furobufen, Halcinonide, Halobetasol Propionate, Halopredone Acetate, Ibufenac, Ibuprofen, Ibuprofen Aluminum, Ibuprofen Piconol, Ilonidap, Indomethacin, Indomethacin Sodium, Indoprofen, Indoxole, Intrazole, Isoflupredone Acetate, Isoxepac, Isoxicam, Ketoprofen, Lofemizole Hydrochloride, Lornoxicam, Loteprednol Etabonate, Meclofenamate Sodium, Meclofenamic Acid, Meclorison DIBUTYRATE, Mefenamic Acid, Mesalamine, Meseclazone, Methylprednisolone Suleptanate, Morniflumate, Nabumetone, Naproxen, Naproxen Sodium, Naproxol, Nimazone, Olsalazine Sodium, Orgotein, Orpanoxin, Oxaprozin, Oxyphenbutazone, Paranyline Hydrochloride, Pentosan Polysulfate Sodium, Phenbutazone Sodium Glycerate, Pirfenidone, Piroxicam, Piroxicam Cinnamate, Piroxicam Olamine, Pirprofen, Prednazate, Prifelone, Prodolic Acid, Proquazone, Proxazole, Proxazole Citrate, Rimexolone, Romazarit, Salcolex, Salnacedin, Salsalate, Sanguinarium Chloride, Seclazone, Sermetacin, Sudoxicam, Sulindac, Suprofen, Talmetacin, Talniflumate, Talosalate, Tebufelone, Tenidap, Tenidap Sodium, Tenoxicam, Tesicam, Tesimide, Tetrydamine, Tiopinac, Tixocortol Pivalate, Tolmetin, Tolmetin Sodium, Triclonide, Triflumidate, Zidometacin, or Zomepirac Sodium.

9. The gelsolin of claim 6 further comprising administering a second agent for treating rheumatoid arthritis.
10. The gelsolin of claim 9, wherein the second agent for treating rheumatoid arthritis is hydroxychloroquine (Plaquenil), chloroquine (Aralen), methotrexate, sulfasalazine (Azulfidine), Leflunomide (Arava), azathioprine (Imuran), penicillamine (Cuprimine or Depen), Gold salts (Ridaura or Aurolate), minocycline (Dynacin or Minocin), cyclosporine (Neoral or Sandimmune) cyclophosphamide (Cytoxan or Neosar), Etanercept (Enbrel), Infliximab (Remicade), Anakinra (Kineret) or Adalimumab (Humira).
11. Gelsolin optionally in combination with a second agent, for use in a method for treating a subject at risk of an inflammatory disease who is known to have a below-normal level of gelsolin; in an effective amount to reduce the risk of the subject developing the inflammatory disease.
12. Gelsolin optionally in combination with a second agent, for use in a method for treating a subject at risk of an inflammatory disease who is known to have a below-normal level of gelsolin; in an effective amount to raise the level of gelsolin in the subject above a predetermined value.

Patentansprüche

1. Verfahren zum Charakterisieren des Risikoprofils eines gesunden Patienten der Entwicklung einer zukünftigen entzündlichen Erkrankung, umfassend:

Vergleichen eines Pegels von Gelsolin in einer Körperflüssigkeit (beispielsweise Blut, Plasma, Serum, Urin, Synovialflüssigkeit oder Alveolarflüssigkeit) oder Gewebe (beispielsweise Gelenk-, gastrointestinales, Thyroid-, adrenales, vaskuläres, pulmonales, renales, Herz-, Haut-, Okular-, Hirn-, pankreatisches, Leber-, nervales, oder Muskelgewebe), das von einem Patienten erhalten wird, mit einem vorbestimmten Wert (bevorzugt etwa 250 mg/l Plasma oder weniger), und

Charakterisieren des Risikoprofils für den Patienten der Entwicklung einer entzündlichen Erkrankung auf der Grundlage des Pegels von Gelsolin im Vergleich zu dem vorbestimmten Wert, wobei der Pegel an Gelsolin bei oder unterhalb des vorbestimmten Pegels kennzeichnend dafür ist, dass der Patient ein erhöhtes Risiko der Entwicklung einer entzündlichen Erkrankung hat, und wobei der Pegel an Gelsolin bei oder oberhalb des vorbestimmten Pegels kennzeichnend dafür ist, dass der Patient kein erhöhtes Risiko der Entwicklung einer entzündlichen Erkrankung hat,

wobei der Patient keinerlei Anzeichen oder Symptome einer Infektion, Sepsis oder Aktin-bezogenen Erkrankung hat.

2. Verfahren nach Anspruch 1, wobei die entzündliche Erkrankung ist: Arthritis, rheumatoide Arthritis, Asthma, chronisch entzündliche Darmerkrankungen (Morbus Crohn oder Colitis ulcerosa), Chronisch Obstruktive Lungenerkrankung (COPD), allergische Rhinitis, Vaskulitis (Polyarteritis nodosa, temporäre Arterienentzündung, Wegener Granulomatosis, Takayasu Arterienentzündung oder Behcet Syndrom), entzündliche Neuropathie, Schuppenflechte, systemischer Lupus Erythematosus (SLE), chronische Thyreoiditis, Hashimoto Thyreoiditis, Morbus Addison, Poly-myalgia Rheumatica, Sjögren Syndrom oder Churg Strauss Syndrom.

3. Verfahren nach Anspruch 1, umfassend das Ausführen von einem oder mehreren Tests zur Einschätzung der entzündlichen Erkrankung, wobei der Test bevorzugt das Messen eines Pegels eines Markers der Entzündung in dem Patienten ist, beispielsweise wobei der Marker der Entzündung ist: CRP, lösliches interzelluläres Adhäsionsmolekül (sICAM-1), ICAM 3, BL-CAM, LFA-2, VCAM ist-I, NCAM, PECAM, Fibrinogen, Serum-Amyloid A (SAA), Lipoprotein in Verbindung mit Phospholipase A2 (LpPIA2), sCD40 Ligand (sCD40L), Myeloperoxidase, Interleukin-6 (IL-6) oder Interleukin-8 (IL-8).

4. Verfahren zum Charakterisieren des Risikoprofils eines gesunden Patienten der Entwicklung einer zukünftigen entzündlichen Erkrankung, umfassend:

Vergleichen des Pegels von Gelsolin in einer Probe, die von einem Patienten erhalten wird, mit einem ersten vorbestimmten Wert (vorzugsweise von ungefähr 250 mg/L Plasma oder weniger), um einen ersten Risikowert zu erstellen,

Vergleichen des Pegels von Gelsolin in einer Probe, die von einem Patienten erhalten wird, mit einem zweiten vorbestimmten Wert, um einen zweiten Risikowert zu erstellen, und

EP 2 001 496 B1

Charakterisieren des Risikoprofils der Entwicklung einer entzündlichen Erkrankung auf der Grundlage der Kombination des ersten Risikowertes und des zweiten Risikowertes, wobei die Kombination des ersten Risikowertes und des zweiten Risikowertes einen dritten Risikowert ergibt, der sich von den ersten und zweiten Risikowerten unterscheidet,

wobei der Patient keinerlei Anzeichen oder Symptome einer Infektion, Sepsis oder Aktin-bezogenen Erkrankung hat.

5. Verfahren nach Anspruch 4, wobei der erste vorbestimmte Wert eine Mehrzahl von vorbestimmten Bereichen des Pegels von Gelsolin ist, einer der Mehrzahl unterhalb von etwa 250 mg/l Plasma und ein anderer dieser Bereiche oberhalb von etwa 250 mg/l Plasma liegt und wobei der Schritt des Vergleichens ein Bestimmen umfasst, in denen von der Mehrzahl von vorbestimmten Bereichen des Pegels von Gelsolin der Pegel von Gelsolin des Patienten abfällt.
6. Gelsolin, wahlweise in Kombination mit einem zweiten Mittel zum Behandeln einer entzündlichen Erkrankung zur Verwendung in einem Verfahren zum Behandeln eines Patienten mit Risiko der Entwicklung einer entzündlichen Erkrankung, der ansonsten vorzugsweise frei ist von Anzeichen, die eine Behandlung mit Gelsolin erfordern, vorzugsweise wobei das Gelsolin Plasma-Gelsolin (pGSN), zytoplasmatisches Gelsolin (cGSN), Advillin, Villin, CapG, Flightless-Proteine, Fragmin, Severin, Adseverin, Protovillin oder Supervillin ist.
7. Gelsolin nach Anspruch 6, wobei die entzündliche Erkrankung ist: Arthritis, rheumatoide Arthritis, Asthma, chronisch entzündliche Darmerkrankungen (Morbus Crohn oder Colitis ulcerosa), Chronisch Obstruktive Lungenerkrankung (COPD), allergische Rhinitis, Vaskulitis (Polyarteritis nodosa, temporäre Arterienentzündung, Wegener Granulomatosis, Takayasu Arterienentzündung oder Behcet Syndrom), entzündliche Neuropathie, Schuppenflechte, systemischer Lupus Erythematosus (SLE), chronische Thyreoiditis, Hashimoto Thyreoiditis, Morbus Addison, Polymyalgia Rheumatica, Sjögren Syndrom oder Churg Strauss Syndrom.
8. Gelsolin nach Anspruch 6, wobei das zweite Mittel zum Behandeln der entzündlichen Erkrankung ist: Alclofenac, Alclometason Dipropionat, Algoston Acetonid, Alpha-Amylase, Amcinafal, Amcinafid, Amfenac Natrium, Amiprilos Hydrochlorid, Anakinra, Aniolac, Anitrazafen, Apazon, Balsalazid Dinatrium, Bendazac, Benoxaprofen, Benzylamin Hydrochlorid, Bromelains, Broperamol, Budesonid, Carprofen, Cicloprofen, Cintazone, Cliprofen, Clobetasol Propionat, Butyrat Clobetasone, Clopirac, Cloticasone Propionat, Cormethasone Acetat, Cortodoxone, Cyclooxygenase-2 (COX-2)-Hemmer, Deflazacort, Desonide, Desoximetasone, Dexamethason Dipropionat, Diclofenac Kalium, Diclofenac Natrium, Diflorasone Diacetat, Diflumidone Natrium, Diflunisal, Difluprednate, Diftalone, Dimethyl Sulfoxid, Drocinonide, Endryson, Enlimomab, Enolicam Natrium, Epirizole, Etodolac, Etofenamate, Felbinac, Fenamole, Fenbufen, Fenclofenac, Fenclorac, Fendosal, Fempipalone, Fentiazac, Flazalone, Fluazacort, Flufenaminsäure, Flumizole, Flunisolide Acetat, Flunixin, Flunixin Meglumine, Fluocortin Butyl, Fluorometholone Acetat, Fluquazone, Flurbiprofen, Fluretofen, Fluticasone Propionate, Furaprofen, Furobufen, Halcinonide, Halobetasol Propionat, Halopredone Acetat, Ibufenac, Ibuprofen, Ibuprofen Aluminium, Ibuprofen Piconol, Ilonidap, Indomethacin, Indomethacin Natrium, Indoprofen, Indoxole, Intrazole, Isoflupredone Acetat, Isoxepac, Isoxicam, Ketoprofen, Lofemizole Hydrochlorid, Lornoxicam, Loteprednol Etabonate, Meclofenamate Natrium, Meclofenaminsäure, Meclo-risone Dibutyrate, Mefenaminsäure, Mesalazin, Meseclazone, Methylprednisolon Suleptanate, Morniflumate, Nabumetone, Naproxen, Naproxen-Natrium, Naproxol, Nimazone, Olsalazine Natrium, Orgotein, Orpanoxin, Oxaprozin, Oxyphenbutazone, Paranyline-Hydrochlorid, Pentosan Polysulfat Natrium, Phenbutazone Natrium Glycerate, Pirofenidol, Piroxicam, Piroxicam Cinnamate, Piroxicam Olamine, Pirprofen, Prednazate, Prifelone, Prodolinsäure, Proquazone, Proxazole, Proxazolcitrat, Rimexolone, Romazarit, Salcolex, Salnacedin, Salsalate, Sanguinarium-Chlorid, Seclazone, Sermetacin, Sudoxicam, Sulindac, Suprofen, Talmetacin, Talniflumate, Talosalate, Tebufelone, Tenidap, Tenidap Natrium, Tenoxicam, Tesicam, Tesimide, Tetrydamine, Tiopinac, Tixocortol Pivalate, Tolmetin, Tolmetin Natrium, Triclonide, Triflumidate, Zidometacin, oder Zomepirac Natrium.
9. Gelsolin nach Anspruch 6, ferner umfassend das Verabreichen eines zweiten Mittels zum Behandeln von rheumtoider Arthritis.
10. Verfahren nach Anspruch 9, wobei das zweite Mittel zum Behandeln von rheumtoider Arthritis ist: Hydroxychlorchinon (Plaquenil), Chlorchinon (Aralen), Methotrexat, Sulfasalazin (Azulfidin), Leflunomid (Arava), Azathioprin (Imuran), Penicillamin (Cuprimin oder Depen), Gold-Salze (Ridaura oder Aurolat), Minocyclin (Dynacin oder Minocin), Cyclosporin (Neoral oder Sandimmun) Cyclophosphamid (Cytoxan oder Neosar), Etanercept (Enbrel), Infliximab (Remicade), Anakinra (Kineret) oder Adalimumab (Humira).
11. Gelsolin, wahlweise in Kombination mit einem zweiten Mittel zur Verwendung in einem Verfahren zum Behandeln

eines Patienten mit Risiko einer entzündlichen Erkrankung, von dem bekannt ist, dass er Gelsolin unterhalb des normalen Pegels aufweist;
in einer wirksamen Menge zum Vermindern des Risikos des Patienten, die entzündliche Erkrankung zu entwickeln.

- 5 12. Gelsolin, wahlweise in Kombination mit einem zweiten Mittel zur Verwendung in einem Verfahren zum Behandeln eines Patienten mit Risiko einer entzündlichen Erkrankung, von dem bekannt ist, dass er Gelsolin unterhalb des normalen Pegels aufweist;
in einer wirksamen Menge zum Erhöhen des Pegels des Gelsolins in dem Patienten oberhalb eines vorbestimmten Wertes.

10

Revendications

- 15 1. Méthode pour la caractérisation d'un profil de risque pour un sujet en bonne santé de développement d'une maladie inflammatoire future, comprenant:

la comparaison d'un niveau de gelsoline dans un fluide corporel (par exemple sang, plasma, sérum, urine, fluide synovial ou fluide alvéolaire) ou un tissu (par exemple un tissu d'articulation, gastrointestinal, thyroïdien, surrénal, vasculaire, pulmonaire, rénal, cardiaque, de peau, oculaire, de cerveau, pancréatique, de foie, de nerf ou de muscle) obtenu à partir du sujet à une valeur prédéterminée (de préférence d'environ 250 mg/l de plasma ou moins), et

20

la caractérisation du profil de risque pour le sujet de développement d'une maladie inflammatoire sur la base du niveau de gelsoline en comparaison avec la valeur prédéterminée, dans laquelle un niveau de gelsoline au niveau prédéterminé ou en dessous est une indication que le sujet est à un risque élevé de développement de la maladie inflammatoire et dans laquelle un niveau de gelsoline au niveau prédéterminé ou au-dessus est une indication que le sujet n'est pas à un risque élevé de développement de la maladie inflammatoire dans laquelle le sujet ne présente pas de signes ou de symptômes d'une infection, d'une septicémie ou d'un trouble lié à l'actine.

25

- 30 2. Méthode selon la revendication 1, dans laquelle la maladie inflammatoire est une arthrite, une polyarthrite rhumatoïde, un asthme, une maladie intestinale inflammatoire (maladie de Crohn ou colite ulcéreuse), une maladie pulmonaire obstructive chronique (COPD), une rhinite allergique, une vascularite (polyartérite noueuse, artérite temporale, granulomatose de Wegener, artérite de Takayasu ou syndrome de Behcet), une neuropathie inflammatoire, un psoriasis, un lupus érythémateux systémique (SLE), une thyroïdite chronique, une thyroïdite de Hashimoto, une maladie de Addison, une pseudo-polyarthrite rhizomélique, un syndrome de Sjogren ou un syndrome de Churg-Strauss.

35

3. Méthode selon la revendication 1 comprenant en outre la réalisation d'un ou de plusieurs tests pour évaluer la maladie inflammatoire, de préférence dans laquelle le test est la mesure d'un niveau d'un marqueur d'inflammation chez le sujet, par exemple dans laquelle le marqueur d'inflammation est CRP, une molécule d'adhésion intracellulaire soluble (sICAM-1), ICAM 3, BL-CAM, LFA-2, VCAM-I, NCAM, PECAM, le fibrinogène, l'amyloïde A sérique (SAA), une phospholipase associée à une lipoprotéine A2 (LpPIA2), le ligand sCD40 (sCD40L), une myéloperoxydase, l'interleukine-6 (IL-6) ou l'interleukine-8 (IL-8).

40

- 45 4. Méthode pour la caractérisation d'un profil de risque pour un sujet en bonne santé de développement d'une maladie inflammatoire future, comprenant:

la comparaison du niveau de la gelsoline dans un échantillon obtenu à partir d'un sujet à une première valeur prédéterminée (de préférence d'environ 250 mg/l de plasma ou moins) pour établir une première valeur de risque, la comparaison du niveau d'un deuxième marqueur d'inflammation dans un échantillon obtenu à partir d'un sujet à une deuxième valeur prédéterminée pour établir une deuxième valeur de risque, et

50

la caractérisation du profil de risque pour le sujet de développement de la maladie inflammatoire sur la base de la combinaison de la première valeur de risque et de la deuxième valeur de risque, dans laquelle la combinaison de la première valeur de risque et de la deuxième valeur de risque établit une troisième valeur de risque différente desdites première et deuxième valeurs de risque, dans laquelle le sujet ne présente pas de signes ou de symptômes d'une infection, d'une septicémie ou d'un trouble lié à l'actine.

55

5. Méthode selon la revendication 4, dans laquelle la première valeur prédéterminée est une pluralité d'intervalles de niveaux de gelsoline prédéterminés, un de ladite pluralité étant en dessous d'environ 250 mg/l de plasma et un autre desdits intervalles étant au-dessus d'environ 250 mg/l de plasma et dans laquelle ladite étape de comparaison comprend la détermination dans laquelle de ladite pluralité d'intervalles de niveaux de gelsoline prédéterminés ledit niveau de gelsoline du sujet se trouve.
6. Gelsoline, optionnellement en combinaison avec un deuxième agent pour le traitement d'une maladie inflammatoire, pour une utilisation dans une méthode pour le traitement d'un sujet à risque de développement d'une maladie inflammatoire, qui est de préférence autrement exempt d'indications exigées pour un traitement avec la gelsoline, de préférence où la gelsoline est une gelsoline plasmatique (pGSN), une gelsoline cytoplasmique (cGSN), advilline, villine, capG, des protéines flightless, fragmine, sévérine, adsévérine, protovilline ou supervilline.
7. Gelsoline selon la revendication 6, dans laquelle la maladie inflammatoire est une arthrite, une polyarthrite rhumatoïde, un asthme, une maladie intestinale inflammatoire (maladie de Crohn ou colite ulcéreuse), une maladie pulmonaire obstructive chronique (COPD), une rhinite allergique, une vascularite (polyarthrite noueuse, artérite temporale, granulomatose de Wegener, artérite de Takayasu ou syndrome de Behcet), une neuropathie inflammatoire, un psoriasis, un lupus érythémateux systémique (SLE), une thyroïdite chronique, une thyroïdite de Hashimoto, une maladie de Addison, une pseudo-polyarthrite rhizomélique, un syndrome de Sjogren ou un syndrome de Churg-Strauss.
8. Gelsoline selon la revendication 6, dans laquelle le deuxième agent pour le traitement de la maladie inflammatoire est alclofénac, aclométasone dipropionate, algestone acétonide, alpha amylase, amcinafal, amcinafide, amfénac sodium, amiprilose chlorhydrate, anakinra, anirolac, anitrazafène, apazone, balsalazide disodium, bendazac, benoxaprofène, benzydamine chlorhydrate, bromelaïnes, bropéramole, budésonide, carprofène, cycloprofène, cintazone, cliprofène, clobétasol propionate, clobétasone butyrate, clopirac, cloticasone propionate, corméthasone acétate, cortodoxone, inhibiteur de cyclooxygénase-2 (COX-2), déflazacort, désoude, désouimétasone, dexaméthasone dipropionate, diclofénac potassium, diclofénac sodium, diflorasone diacétate, diflumidone sodium, diflunisal, difluprednate, diftalone, diméthylsulfoxyde, drocinonide, endrysonne, enlimomab, énoïcicam sodium, épirizole, étodolac, étofénamate, felbinac, fénamole, fenbufène, fencolfénac, fenclorac, fendosal, fempipalone, fentiazac, flazalone, fluazacort, acide flufénamique, flumizole, flunisolid acétate, flunixin, flunixin méglumine, fluocortine butyle, fluorométholone acétate, fluquazone, flurbiprofène, flurétofène, fluticasone propionate, furaprofène, furobufène, halcinonide, halobétasol propionate, halopredone acétate, ibufénac, ibuprofène, ibuprofène aluminium, ibuprofène piconol, ilonidap, indométhacine, indométhacine sodium, indoprofène, indoxole, intrazole, isofluprednone acétate, isoxepac, isoxicam, kétoprofène, lofémizole chlorhydrate, lornoxicam, lotéprednol étabonate, méclofénamate sodium, acide méclofénamique, méclorisone dibutyrate, acide méfénamique, mésalamine, méseclazone, méthylprednisolone suleptanate, morniflumate, nabumétone, naproxène, naproxène sodium, naproxol, nimazone, olsalazine sodium, orgotéine, orpanoxine, oxaprozine, oxyphenbutazone, paranyline chlorhydrate, pentosane polysulfate sodium, phenbutazone sodium glycérate, pifénidone, piroxicam, piroxicam cinnamate, piroxicam olamine, pirprofène, prednazate, prifénone, acide prodolique, proquazone, proxazole, proxazole citrate, riméxolone, romazarit, salcolex, salnacédine, salsalate, sanguinarium chlorure, séclazone, sermétacine, sudoxicam, sulindac, suprofène, talmétacine, talniflumate, talosalate, tébufélone, ténidap, ténidap sodium, ténoxycam, tésicam, tésimide, tétrydamine, tiopinac, tixocortol pivalate, tolmétine, tolmétine sodium, triclone, triflumidate, zidométacine ou zomépirac sodium.
9. Gelsoline selon la revendication 6 comprenant en outre l'administration d'un deuxième agent pour le traitement d'une polyarthrite rhumatoïde.
10. Gelsoline selon la revendication 9, dans laquelle le deuxième agent pour le traitement d'une polyarthrite rhumatoïde est hydroxychloroquine (Plaquenil), chloroquine (Aralen), méthotrexate, sulfasalazine (Azulfidine), léflunomide (Ara-va), azathioprine (Imuran), pénicillamine (Cuprimine ou Depen), des sels d'or (Ridaura ou Aurolate), minocycline (Dynacin ou Minocin), cyclosporine (Neoral ou Sandimmune), cyclophosphamide (Cytoxan ou Neosar), étanercept (Enbrel), infliximab (Remicade), anakinra (Kineret) ou adalimumab (Humira).
11. Gelsoline optionnellement en combinaison avec un deuxième agent, pour une utilisation dans une méthode pour le traitement d'un sujet à risque d'une maladie inflammatoire qui est connu pour avoir un niveau en dessous de la normale de gelsoline; dans une quantité efficace pour réduire le risque que le sujet développe la maladie inflammatoire.
12. Gelsoline optionnellement en combinaison avec un deuxième agent, pour une utilisation dans une méthode pour

EP 2 001 496 B1

le traitement d'un sujet à risque d'une maladie inflammatoire qui est connu pour avoir un niveau en dessous de la normale de gelsoline;
dans une quantité efficace pour élever le niveau de gelsoline chez le sujet au-dessus d'une valeur prédéterminée.

5

10

15

20

25

30

35

40

45

50

55




Fig. 1




Fig. 2




Fig. 3

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 20060009386 A [0007]
- WO 2005112970 A [0007]
- WO 2005046454 A [0008]
- WO 02059604 A [0009]
- WO 2004082617 A [0010]
- WO 03088811 A [0011]
- US 6040147 A [0012]
- US 5543297 A, Cromlish [0032] [0034]
- US 5474995 A [0033] [0034]
- US 5521213 A [0033]
- US 5536752 A [0033]
- US 5550142 A [0033]
- US 5552422 A [0033]
- US 5604253 A [0033]
- US 5604260 A [0033]
- US 5677318 A [0033]
- US 5691374 A [0033]
- US 5698584 A [0033]
- US 5710140 A [0033]
- US 5733909 A [0033]
- US 5789413 A [0033]
- US 5817700 A [0033]
- US 5849943 A [0033]
- US 5861419 A [0033]
- US 5922742 A [0033]
- US 5925631 A [0033]
- US 5643933 A [0033]
- WO 9500501 A [0034]
- WO 9518799 A [0034]
- US 5284656 A, Platz [0130]
- US 5451569 A, Wong [0130]
- US 5563153 A [0147]

Non-patent literature cited in the description

- YIN, H. L. ; STOSSEL, T. P. *Nature*, 1979, vol. 281, 583-6 [0006]
- CANDIANO et al. *American Journal of Respiratory and Critical Care Medicine*, 2005, vol. 172 (9 [0013]
- ROTHENBACH et al. *Journal of Applied Physiology*, 2004, vol. 96 (1 [0014]
- AIDINIS et al. *PLoS Genetics*, 2005, vol. 1 (4 [0015]
- YIN, H. L. ; KWIATKOWSKI, D. J. ; MOLE, J. E. ; COLE, F. S. *J Biol Chem*, 1984, vol. 259, 5271-6 [0019] [0063]
- KWIATKOWSKI, D. J. ; STOSSEL, T. P. ; ORKIN, S. H. ; MOLE, J. E. ; COHEN, H. R. ; YIN, H. L. *Nature*, 1986, vol. 323, 455-8 [0019]
- SUN et al. *J. Biol. Chem.*, 1999, vol. 274, 33179-33182 [0062]
- KWIATKOWSKI, D. J. ; STOSSEL, T. P. ; ORKIN, S. H. ; MOLE, J. E. ; COLTEN, H. R. ; YIN, H. L. *Nature*, 1986, vol. 323, 455-8 [0063]
- WEN. *Biochemistry*, 1996, vol. 35, 9700-9709 [0063]
- *Remington's Pharmaceutical Sciences*, 1980 [0067]
- *Remington's Pharmaceutical Sciences*. Lippincott, Williams and Wilkins, 2001 [0103]
- Soluble Polymer-Enzyme Adducts. **ABUCHOWSKI ; DAVIS**. *Enzymes as Drugs*. Wiley-Interscience, 1981, 367-383 [0114]
- **NEWMARK et al.** *J. Appl. Biochem.*, 1982, vol. 4, 185-189 [0114]
- **ADJEI et al.** *Pharmaceutical Research*, 1990, vol. 7, 565-569 [0130]
- **ADJEI et al.** *International Journal of Pharmaceutics*, 1990, vol. 63, 135-144 [0130]
- **BRAQUET et al.** *Journal of Cardiovascular Pharmacology*, 1989, vol. 13 (5), 143-146 [0130]
- **HUBBARD et al.** *Annals of Internal Medicine*, 1989, vol. III, 206-212 [0130]
- **SMITH et al.** *J. Clin. Invest.*, 1989, vol. 84, 1145-1146 [0130]
- **OSWEIN et al.** *Aerosolization of Proteins. Proceedings of Symposium on Respiratory Drug Delivery*, 1990, vol. II [0130]
- **DEBS et al.** *J. Immunol.*, 1988, vol. 140, 3482-3488 [0130]
- **LANGER.** *Science*, 1990, vol. 249, 1527-1533 [0143]
- **H.S. SAWHNEY ; C.P. PATHAK ; J.A. HUBEI.** *Macromolecules*, 1993, vol. 26, 581-587 [0144]
- **MYERS et al.** *Life Sciences*, 1997, vol. 61, 1861-1878 [0167]

Szabadalmi igénypontok

1. Eljárás egészséges alanyok jövőbeli gyulladásos betegség kialakulására vonatkozó kockázati profiljának jellemzésére, amely eljárás tartalmazza a következőket:

az alanytól származó testfolyadékban (például vér, plazma, szérum, vizelet, szinoviális folyadék, vagy alveoláris folyadék) vagy azővenben (például izületi, emésztőrendszeri, pajzsmirigy-, mellékvese-, érendszeri, tüdő-, vese-, szív-, bőr-, szem-, agy-, hasnyálmirigy-, máj-, ideg- vagy izomszövet) a gelszolin szintjének összehasonlítása egy előre meghatározott értékkel (előnyösen kb. 250 mg/l plazma értékkel vagy kevesebbel), és

alanyok gyulladásos betegség kialakulására vonatkozó kockázati profiljának jellemzése a gelszolin szintjének az előre meghatározott értékkel történő összehasonlítás alapján, ahol az előre meghatározott érték szintjén vagy az alatt lévő gelszolin szint azt jelzi, hogy az alany gyulladásos betegség kialakulás magas kockázatának van kitéve, és ahol az előre meghatározott érték szintjén vagy felette lévő gelszolin szint azt jelzi, hogy az alany nincs gyulladásos betegség kialakulás magas kockázatának kitéve,

ahol az alany fertőzés, szepszis vagy aktívhoz kapcsolódó rendellenesség jeleit vagy tüneteit nem mutatja.

2. Az 1. igénypont szerinti eljárás, ahol a gyulladásos betegség izületi gyulladás, reumás izületi gyulladás, asztma, gyulladásos bélbetegség (Crohn-betegség vagy fekélyes vastagbélgyulladás), krónikus obstruktív tüdőbetegség (COPD), allergiás nátha, vaszkulitisz (poliarteritis nodosa, temporalis arteritis, Wegener-granulomatosis, Takayasu-arteritis vagy Behcet-szindróma), gyulladásos neuropátia, pikkelysömör, szisztémás lupus erythematosus (SLE), krónikus pajzsmirigy-gyulladás, Hashimoto-pajzsmirigy-gyulladás, Addison-kór, reumás polimialgia, Sjogren-szindróma vagy Churg-Strauss-szindróma.

3. Az 1. igénypont szerinti eljárás, amely tartalmazza továbbá egy vagy több vizsgálat elvégzését a gyulladásos betegség értékelésére, előnyösen ahol a vizsgálat a gyulladás egy markerének szintjét méri az alanyban, például ahol a gyulladás markere a CRP, oldható intercelluláris adhéziós molekula (sICAM-1), ICAM 3, BL-CAM, LFA-2, VCAM-1, NCAM, PECAM, fibrinogén, szérum amiloid A (SAA), lipoproteinhez kapcsolódó foszfolipáz A2 (LpPLA2), sCD40 ligandum (sCD40L), mieloperoxidáz, interleukin-6 (IL-6) vagy interleukin-8 (IL-8).

4. Eljárás egészséges alanyok jövőbeli gyulladásos betegség kialakulására vonatkozó kockázati profiljának jellemzésére, amely eljárás tartalmazza a következőket:

az alanytól származó mintában a gelszolin szintjének összehasonlítása egy első előre meghatározott értékkel (előnyösen kb. 250 mg/l plazma értékkel vagy kevesebbel), és így első kockázati értéket állapítunk meg,

az alanytól származó mintában a gyulladás egy második markerének szintjének összehasonlítása egy második előre meghatározott értékkel, és így második kockázati értéket állapítunk meg, és

alanyok gyulladásos betegség kialakulására vonatkozó kockázati profiljának jellemzése az első kockázati érték és a második kockázati érték kombinációjának alapján, ahol az első kockázati érték és a második



kockázati érték kombinációja egy harmadik, az első és második kockázati értéktől eltérő kockázati értéket állapít meg.

ahol az alany fertőzés, szepszis vagy aktínhoz kapcsolódó rendellenesség jeleit vagy tüneteit nem mutatja.

5. A 4. igénypont szerinti eljárás, ahol az első előre meghatározott érték előre meghatározott gelszolin szint tartományok sokasága, ezen sokaság egyike kb. 250 mg/l plazma érték alatti és a tartományok közül egy másik kb. 250 mg/l plazma érték feletti, és ahol az összehasonlító lépés tartalmazza annak meghatározását, hogy az előre meghatározott gelszolin szint tartomány sokaság közül melyikbe esik az alany gelszolin szintje.

6. Gelszolin, adott esetben kombinációban egy második, gyulladásos betegség kezeléséhez megfelelő szerrel, gyulladásos betegség kialakulásának kockázatát hordozó alany kezelésére szolgáló eljárásban történő alkalmazásra, amely alany előnyösen egyébként mentes olyan indikációktól, amely a gelszolinrel történő kezelést igényelné, előnyösen ahol a gelszolin plazma gelszolin (pGSN), citoplazmás gelszolin (cGSN), advillin, villin, capG, „flightless” fehérjék, fragmin, severin, adseverin, protovillin vagy supervillin.

7. A 6. igénypont szerinti gelszolin, ahol a gyulladásos betegség izületi gyulladás, reumás izületi gyulladás, asztma, gyulladásos bélbetegség (Crohn-betegség vagy fekélyes vastagbélgyulladás), krónikus obstruktív tüdőbetegség (COPD), allergiás nátha, vaszkulitisz (poliarteritis nodosa, temporalis arteritis, Wegener-granulomatosis, Takayasu-arteritis, vagy Behcet-szindróma), gyulladásos neuropátia, pikkelysömör, szisztémás lupus erythematosus (SLE), krónikus pajzsmirigy-gyulladás, Hashimoto-pajzsmirigy-gyulladás, Addison-kór, reumás polimialgia, Sjogren-szindróma vagy Churg-Strauss-szindróma.

8. A 6. igénypont szerinti gelszolin, ahol második, gyulladásos betegség kezeléséhez megfelelő szer az Aklufenak, Alklometazon-dipropionát, Algeszton-acetonid, Alfa-amiláz, Amcinafal, Amcinafid, Amfenak-nátrium, Amiprilóz-hidroklorid, Anakinra, Anilolak, Anitrazafen, Apazon, Balszalazid-dinátrium, Bendazak, Benoxaprofen, Benzidamin-hidroklorid, Bromelainok, Broperamol, Budeszonid, Karprofén, Cikloprofén, Clotazon, Kliprofén, Klobetazol-propionát, Klobetazon-butirát, Klopirak, Klotikazon-propionát, Kometazon-acetát, Kortodoxon, Ciklooxigenáz-2 (COX-2) inhibitor, Deflazacort, Dezonid, Dezoximetazon, Dexametazon-dipropionát, Diklofenak-kálium, Diklofenak-nátrium, Diflorazon-diacetát, Diflumidon-nátrium, Diflunizal, Difluprednát, Diftalon, Dimetil-szulfoxid, Drocinonid, Endrizon, Enlimomab, Enolkam-nátrium, Epirizol, Etodolak, Etofenamát, Felbinak, Fenamol, Fenbufén, Fenklofenak, Fenklorak, Fendozal, Fempipalon, Fentiazak, Flazalon, Fluazakort, Flufenaminsav, Flumizol, Flunizolid-acetát, Flunixin, Flumixin-meglumin, Fluokortin-butil, Fluorometolon-acetát, Fluquazon, Flurbiprofén, Fluretofén, Flufikazon-propionát, Furaprofén, Furobufén, Halcinonid, Halobetazol-propionát, Halopredon-acetát, Ibufenak, Ibuprofén, Ibuprofén-alumínium, Ibuprofén-pikonoil, Iionidap, Indometacin, Indometacin-nátrium, Indoprofén, Indoxol, Intrazol, Izoflupredon-acetát, Izoxepak, Izoxikam, Ketoprofén, Lofemizol-hidroklorid, Lomoxikam, Loteprednol-etabonát, Meklofenamát-nátrium, Meclofenaminsav, Meklorizon-dibutirát, Mefenaminsav, Mezalamin, Mezeklazon, Metilprednizolon-szuleptanát, Morniflumát, Nabumeton, Naproxén, Naproxén-nátrium, Naproxol, Nimazon, Olszalazin-nátrium, Örgöteín, Orpanoxin, Oxaprozin, Oxifenbutazon, Paramilin-hidroklorid, Pentozán-poliszulfát-nátrium, Fenbutazon-nátrium-glicerát, Pirtfenidon, Piroxikam, Piroxikam-cinnamát, Piroxikam-olamin, Pirprofén, Prednazát, Prifelon, Prodolinsav, Proquazon, Proxazol, Proxazol-citrát, Rimexolon, Romazarit, Szalkolex,

Szahnacedin, Szalszalát, Szanguinarium-klorid, Szeklazon, Szermetacín, Szudoxikam, Szulindak, Szuprofen, Talmetacin, Talniflumát, Talosalát, Tébufelon, Tenidap, Tenidap-nátrium, Tenoxicam, Tezikam, Tesimid, Tetrídamin, Tiopinak, Tixokortol-pivalát, Tolmetin, Tolmetin-nátrium, Triklonid, Triflumidát, Zidometacin vagy Zomepirak-nátrium.

9. A 6. igénypont szerinti gelszolin, amely tartalmazza továbbá reumás ízületi gyulladás kezeléséhez megfelelő második szer adagolását.

10. A 9. igénypont szerinti gelszolin, ahol a reumás ízületi gyulladás kezeléséhez megfelelő második szer a hidroxiklorokin (Plaquenil), klorokin (Aralen), metotrexát, szulfazalazin (Azulfidine), leflunomid (Arava), azatioprin (Imuran), penicillamin (Cuprimine vagy Depen), aranysók (Ridaura vagy Aurolate), minociklin (Dynacin vagy Minocin), ciklosporin (Neoral vagy Sandimmune) ciklofoszfamid (Cytosan vagy Neosar), Etanercept (Enbrel), Infliximab (Remicade), Anakinra (Kineret) vagy Adalimumab (Humira).

11. Gelszolin, adott esetben kombinációban egy második szerrel, gyulladásos betegség kockázatát hordozó alany kezelésére szolgáló eljárásban történő alkalmazásra, amely alanyról ismert, hogy a normálnál kisebb gelszolin szinttel rendelkezik, ahhoz hatékony mennyiségben, hogy csökkentse az alanyban a gyulladásos betegség kialakulásának kockázatát.

12. Gelszolin, adott esetben kombinációban egy második szerrel, gyulladásos betegség kockázatát hordozó alany kezelésére szolgáló eljárásban történő alkalmazásra, amely alanyról ismert, hogy a normálnál kisebb gelszolin szinttel rendelkezik, ahhoz hatékony mennyiségben, hogy az alanyban a gelszolin szintjét egy előre meghatározott értéknél magasabbra emelje.