

(51) International Patent Classification:
A61K 31/519 (2006.01)(21) International Application Number:
PCT/US2011/038544(22) International Filing Date:
31 May 2011 (31.05.2011)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/349,959 31 May 2010 (31.05.2010) US(71) Applicant (for all designated States except US): **INTRACELLULAR THERAPIES, INC.** [US/US]; 3690 Broadway, New York, New York 10032 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LI, Peng** [CN/US]; Intra-Cellular Therapies, Inc., 3690 BROADWAY, New York, New York 10032 (US). **ZHENG, Hailin** [CN/US]; Intra-Cellular Therapies, Inc., 3690 Broadway, New York, New York 10032 (US). **ZHAO, Jun** [CN/US]; Intra-Cellular Therapies, Inc., 3690 Broadway, New York, New York 10032 (US). **WENNOGLE, Lawrence, P.** [US/US]; Intra-Cellular Therapies, Inc., 3690 Broadway, New York, New York 10032 (US).(74) Agents: **LA, Brittany** et al.; Hoxie & Associates LLC, 75 Main Street, Suite 301, Millburn, NJ 07041 (US).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: ORGANIC COMPOUNDS



(57) Abstract: The present invention relates to (1- or 2- and/or 5 and/or 7-substituted)-(3-oxy)-(4H, 4-imino, 4-thioxo or 4-oxo)-pyrazolo[3,4-d]pyrimidin-6-ones, e.g., a compound of formula II as described below, processes for their production, their use as pharmaceuticals and pharmaceutical compositions comprising them. Of particular interest are novel compounds useful as inhibitors of phosphodiesterase 1 (PDE1), e.g., in the treatment of diseases involving disorders of the dopamine D1 receptor intracellular pathway, such as Parkinson's disease, depression, narcolepsy, damage to cognitive function, e.g., in schizophrenia, or disorders that may be ameliorated through enhanced progesterone-signaling pathway, e.g., female sexual dysfunction.

ORGANIC COMPOUNDS

This application claims priority from U.S. Provisional Application No. 61/349,959, filed May 31, 2010, the contents of which are hereby incorporated by reference in their entirety.

5

TECHNICAL FIELD

[0001] The present invention relates to (1- or 2- and/or 5 and/or 7-substituted)-(3-oxy)-(4H, 4-imino, 4-thioxo or 4-oxo)-pyrazolo[3,4-d]pyrimidin-6-ones, e.g., a compound of formula II as described below, processes for their production, their use as pharmaceuticals and pharmaceutical compositions comprising them. Of particular interest are novel compounds useful as inhibitors of phosphodiesterase 1 (PDE1), e.g., in the treatment of diseases involving disorders of the dopamine D1 receptor intracellular pathway, such as Parkinson's disease, depression, narcolepsy, damage to cognitive function, e.g., in schizophrenia, or disorders that may be ameliorated through enhanced progesterone-signaling pathway, e.g., female sexual dysfunction.

15

BACKGROUND OF THE INVENTION

[0002] Eleven families of phosphodiesterases (PDEs) have been identified but only PDEs in Family I, the Ca^{2+} -calmodulin-dependent phosphodiesterases (CaM-PDEs), have been shown to mediate both the calcium and cyclic nucleotide (e.g. cAMP and cGMP) signaling pathways. The three known CaM-PDE genes, PDE1A, PDE1B, and PDE1C, are all expressed in central nervous system tissue. PDE1A is expressed throughout the brain with higher levels of expression in the CA1 to CA3 layers of the hippocampus and cerebellum and at a low level in the striatum. PDE1A is also expressed in the lung and heart. PDE1B is predominately expressed in the striatum, dentate gyrus, olfactory tract and cerebellum, and its expression correlates with brain regions having high levels of dopaminergic innervation. Although PDE1B is primarily expressed in the central nervous system, it may be detected in the heart. PDE1C is primarily expressed in olfactory epithelium, cerebellar granule cells, and striatum. PDE1C is also expressed in the heart and vascular smooth muscle.

25
30

[0003] Cyclic nucleotide phosphodiesterases decrease intracellular cAMP and cGMP signaling by hydrolyzing these cyclic nucleotides to their respective inactive 5'-monophosphates (5' AMP and 5' GMP). CaM-PDEs play a critical role in

mediating signal transduction in brain cells, particularly within an area of the brain known as the basal ganglia or striatum. For example, NMDA-type glutamate receptor activation and/or dopamine D2 receptor activation result in increased intracellular calcium concentrations, leading to activation of effectors such as calmodulin-dependent kinase II (CaMKII) and calcineurin and to activation of CaM-PDEs, resulting in reduced cAMP and cGMP. Dopamine D1 receptor activation, on the other hand, leads to activation of nucleotide cyclases, resulting in increased cAMP and cGMP. These cyclic nucleotides in turn activate protein kinase A (PKA; cAMP-dependent protein kinase) and/or protein kinase G (PKG; cGMP-dependent protein kinase) that phosphorylate downstream signal transduction pathway elements such as DARPP-32 (dopamine and cAMP-regulated phosphoprotein) and cAMP responsive element binding protein (CREB). Phosphorylated DARPP-32 in turn inhibits the activity of protein phosphatase-1 (PP-1), thereby increasing the state of phosphorylation of substrate proteins such as progesterone receptor (PR), leading to induction of physiologic responses. Studies in rodents have suggested that inducing cAMP and cGMP synthesis through activation of dopamine D1 or progesterone receptor enhances progesterone signaling associated with various physiological responses, including the lordosis response associated with receptivity to mating in some rodents. See Mani, et al., Science (2000) 287: 1053, the contents of which are incorporated herein by reference.

[0004] CaM-PDEs can therefore affect dopamine-regulated and other intracellular signaling pathways in the basal ganglia (striatum), including but not limited to nitric oxide, noradrenergic, neurotensin, CCK, VIP, serotonin, glutamate (e.g., NMDA receptor, AMPA receptor), GABA, acetylcholine, adenosine (e.g., A2A receptor), cannabinoid receptor, natriuretic peptide (e.g., ANP, BNP, CNP), DARPP-32, and endorphin intracellular signaling pathways.

[0005] Phosphodiesterase (PDE) activity, in particular, phosphodiesterase 1 (PDE1) activity, functions in brain tissue as a regulator of locomotor activity and learning and memory. PDE1 is a therapeutic target for regulation of intracellular signaling pathways, preferably in the nervous system, including but not limited to a dopamine D1 receptor, dopamine D2 receptor, nitric oxide, noradrenergic, neurotensin, CCK, VIP, serotonin, glutamate (e.g., NMDA receptor, AMPA receptor), GABA, acetylcholine, adenosine (e.g., A2A receptor), cannabinoid

receptor, natriuretic peptide (e.g., ANP, BNP, CNP), endorphin intracellular signaling pathway and progesterone signaling pathway. For example, inhibition of PDE1B should act to potentiate the effect of a dopamine D1 agonist by protecting cGMP and cAMP from degradation, and should similarly inhibit dopamine D2

5 receptor signaling pathways, by inhibiting PDE1 activity. Chronic elevation in intracellular calcium levels is linked to cell death in numerous disorders, particularly in neurodegenerative diseases such as Alzheimer's, Parkinson's and Huntington's Diseases and in disorders of the circulatory system leading to stroke and myocardial infarction. PDE1 inhibitors are therefore potentially useful in diseases characterized

10 by reduced dopamine D1 receptor signaling activity, such as Parkinson's disease, restless leg syndrome, depression, narcolepsy and cognitive impairment. PDE1 inhibitors are also useful in diseases that may be alleviated by the enhancement of progesterone-signaling such as female sexual dysfunction.


[0006] There is thus a need for compounds that selectively inhibit PDE1 activity,

15 especially PDE1A and/or PDE1B activity.

SUMMARY OF THE INVENTION

[0007] The invention provides (1- or 2- and/or 5 and/or 7-substituted)-(3-oxy)-(4H, 4-imino, 4-thioxo or 4-oxo)-pyrazolo[3,4-d]pyrimidin-6-ones, e.g., a compound

20 of formula II:



Formula II

wherein

- (i) Q is $-C(=O)-$, $-C(=S)-$, $-C(=N(R_6))-$ or $-C(R_{14})(R_{15})-$;
- 25 (ii) R_1 is H or C_{1-6} alkyl (e.g., methyl or ethyl);
- (iii) R_2 is
- H,

C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
 haloC₁₋₆alkyl (e.g., trifluoromethyl or 2,2,2-trifluoroethyl),
 N(R₁₄)(R₁₅)-C₁₋₆alkyl (e.g., 2-(dimethylamino)ethyl or 2-aminopropyl),
 arylC₀₋₆alkyl (e.g., phenyl or benzyl), wherein said aryl is optionally substituted with one or more C₁₋₆alkoxy, for example, C₁₋₆alkoxyarylC₀₋₆alkyl (e.g., 4-methoxybenzyl), heteroarylC₀₋₆alkyl (e.g., pyridinylmethyl), wherein said heteroaryl is optionally substituted with one or more C₁₋₆alkoxy (e.g., C₁₋₆alkoxyheteroarylC₁₋₆alkyl);
 -G-J wherein G is a single bond or C₁₋₆alkylene (e.g., methylene) and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino, for example,
 -C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., -C₀₋₄alkyl-cyclopentyl, -C₀₋₄alkyl-cyclohexyl or -C₀₋₄alkyl-cyclopropyl), wherein said cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl),
 -C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., -C₀₋₄alkyl-pyrrolidinyl, for example, -C₀₋₄alkylpyrrolidin-3-yl) wherein said heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methylpyrrolidin-2-yl, 1-methylpyrrolidin-2-yl-methyl or 1-methylpyrrolidin-3-yl-methyl);

(iv) R₃ is

1) -D-E-F wherein:

D is a single bond, C₁₋₆alkylene (e.g., methylene), or arylC₁₋₆alkylene (e.g., benzylene or -CH₂C₆H₄-);

E is

5

a single bond,

C₁₋₄alkylene (e.g., methylene, ethynylene, prop-2-yn-1-ylene),

10

C₀₋₄alkylarylene (e.g., phenylene or -C₆H₄-, -benzylene- or -CH₂C₆H₄-), wherein the arylene group is optionally substituted with halo (e.g., Cl or F),

heteroarylene (e.g., pyridinylene or pyrimidinylene),

aminoC₁₋₆alkylene (e.g., -CH₂N(H)-),

amino (e.g., -N(H)-);

15

C₃₋₈cycloalkylene optionally containing one or more heteroatom selected from N or O (e.g., piperidinylene),

F is

H,

halo (e.g., F, Br, Cl),

C₁₋₆alkyl (e.g., isopropyl or isobutyl),

20

haloC₁₋₆alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl),

C₃₋₈cycloalkyl optionally containing one or more atom selected from a group consisting of N, S or O (e.g.,

25

cyclopentyl, cyclohexyl, piperidinyl, pyrrolidinyl,

tetrahydro-2H-pyran-4-yl, or morpholinyl), and

optionally substituted with one or more C₁₋₆alkyl (e.g., methyl or isopropyl), for example, 1-methylpyrrolidin-

2-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, piperidin-2-yl, 1-

methylpiperidin-2-yl, 1-ethylpiperidin-2-yl,

30

heteroaryl optionally substituted with one or more C₁₋₆alkyl,

halo (e.g., fluoro) or haloC₁₋₆alkyl (e.g., pyridyl (for

example, pyrid-2-yl), pyrimidinyl (for example,

pyrimidin-2-yl), thiadiazolyl (for example, 1,2,3-

thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example, pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl, 4-methylimidazolyl, 1-methylimidazol-2-yl)), triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-yl),
 5 C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazolyl);

C₁₋₆alkoxy,

-O-haloC₁₋₆alkyl (e.g., -O-CF₃),

C₁₋₆alkylsulfonyl (for example, methylsulfonyl or -

S(O)₂CH₃),

10 -C(O)-R₁₃, wherein R₁₃ is -N(R₁₄)(R₁₅), C₁₋₆alkyl (e.g., methyl), -OC₁₋₆alkyl (e.g., -OCH₃), haloC₁₋₆alkyl


(trifluoromethyl), aryl (e.g., phenyl), or heteroaryl;
 -N(R₁₄)(R₁₅);

or

15 2) a substituted heteroarylC₁₋₆alkyl, e.g., substituted with haloC₁₋₆alkyl;

or

3) attached to one of the nitrogens on the pyrazolo portion of
 Formula II and is a moiety of Formula A



Formula A

wherein:

25 X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen
 (e.g., Cl or F); and

R₁₀ is

halogen (e.g., fluoro or chloro),

C₁₋₆alkyl,
C₃₋₈cycloalkyl,
heteroC₃₋₈cycloalkyl (e.g., pyrrolidinyl or
piperidinyl),
5 haloC₁₋₆alkyl (e.g., trifluoromethyl),
aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for
example, pyrid-2-yl) or e.g., thiadiazolyl (for
example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g.,
pyrazolyl (for example, pyrazol-1-yl) or
10 imidazolyl (for example, imidazol-1-yl)),
triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g.,
tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-
methyl-1,2,4-oxadiazol), pyrazolyl (e.g.,
pyrazol-1-yl),
15 wherein said aryl, heteroaryl, cycloalkyl or
heterocycloalkyl is optionally substituted with
one or more C₁₋₆alkyl (e.g., methyl), halogen
(e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g.,
trifluoromethyl), hydroxy, carboxy, -SH, or an
20 additional aryl or heteroaryl (e.g., biphenyl or
pyridylphenyl),
C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl),
arylcarbonyl (e.g., benzoyl),
heteroarylcarbonyl,
25 C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl),
Aminocarbonyl,
-N(R₁₄)(R₁₅);
preferably R₁₀ is phenyl, pyridyl, piperidinyl or
pyrrolidinyl optionally substituted with the
30 substituents previously defined, e.g. optionally
substituted with halo or alkyl;
provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
respectively, is not present;

(v) R₄ is:

C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl),
 C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
 C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-
 3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-
 1-yl), morpholinyl),
 -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or
 -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-
 yl)

wherein said aryl or heteroaryl is optionally substituted with
 one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-
 hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group
 (e.g., biphenyl-4-ylmethyl);

(vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form.

[0007] In a particular embodiment, the invention further provides a compound of
 Formula II as follows:

2.1 Formula II or 2.1, wherein Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -
 C(R₁₄)(R₁₅)-;

2.2 Formula II or 2.1, wherein Q is -C(=O)-;

2.3 Formula II or 2.1, wherein Q is -C(=S)-;

2.4 Formula II or 2.1, wherein Q is -C(=N(R₆))-;

2.5 Formula II, wherein Q is -C(R₁₄)(R₁₅)-;

2.6 Formula II or any of 2.1-2.5, wherein R₁ is H or C₁₋₆alkyl (e.g.,
 methyl or ethyl);

2.7 Formula II or any of 2.1-2.6, wherein R₁ is H;

2.8 Formula II or any of 2.1-2.6, wherein R₁ is C₁₋₆alkyl (e.g., methyl
 or ethyl);

2.9 Formula II or any of 2.5-2.8, wherein R₂ is:

H,

C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-
 dimethylpropyl) wherein said alkyl group is optionally

substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),

haloC₁₋₆alkyl (e.g., trifluoromethyl or 2,2,2-trifluoroethyl),

5 N(R₁₄)(R₁₅)-C₁₋₆alkyl (e.g., 2-(dimethylamino)ethyl or 2-aminopropyl),

arylC₀₋₆alkyl (e.g., phenyl or benzyl), wherein said aryl is optionally substituted with one or more C₁₋₆alkoxy, for example, C₁₋₆alkoxyarylC₀₋₆alkyl (e.g., 4-methoxybenzyl),

10 heteroarylC₀₋₆alkyl (e.g., pyridinylmethyl), wherein said heteroaryl is optionally substituted with one or more C₁₋₆alkoxy (e.g., C₁₋₆alkoxyheteroarylC₁₋₆alkyl);

-G-J wherein G is a single bond or C₁₋₆alkylene (e.g., methylene) and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein

15 the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino, for example,

-C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., -C₀₋₄alkyl-cyclopentyl, -C₀₋₄alkyl-cyclohexyl or -C₀₋₄alkyl-cyclopropyl),

20 wherein said cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl),

-C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., -C₀₋₄alkyl-pyrrolidinyl, for example, -C₀₋₄alkylpyrrolidin-3-yl) wherein said heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methylpyrrolidin-2-yl, 1-methylpyrrolidin-2-yl-methyl or 1-methylpyrrolidin-3-yl-methyl);

25

30

2.10 Formula II or any of 2.5-2.9, wherein R₂ is:

C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally

- substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- haloC₁₋₆alkyl (e.g., trifluoromethyl or 2,2,2-trifluoroethyl),
- 5 N(R₁₄)(R₁₅)-C₁₋₆alkyl (e.g., 2-(dimethylamino)ethyl or 2-aminopropyl),
- arylC₀₋₆alkyl (e.g., phenyl or benzyl), wherein said aryl is optionally substituted with one or more C₁₋₆alkoxy, for example, C₁₋₆alkoxyarylC₀₋₆alkyl (e.g., 4-methoxybenzyl),
- 10 heteroarylC₀₋₆alkyl (e.g., pyridinylmethyl), wherein said heteroaryl is optionally substituted with one or more C₁₋₆alkoxy (e.g., C₁₋₆alkoxyheteroarylC₁₋₆alkyl);
- G-J wherein G is a single bond or C₁₋₆alkylene (e.g., methylene) and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein
- 15 the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino, for example,
- C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., -C₀₋₄alkyl-cyclopentyl, -C₀₋₄alkyl-cyclohexyl or -C₀₋₄alkyl-cyclopropyl), wherein said
- 20 cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl),
- C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., -C₀₋₄alkyl-pyrrolidinyl, for example, -C₀₋₄alkylpyrrolidin-3-yl) wherein said
- 25 heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methyl-pyrrolindin-2-yl, 1-methyl-pyrrolindin-2-yl-methyl or 1-methyl-pyrrolindin-3-yl-methyl);
- 30 2.11 Formula II or any of 2.5-2.9, wherein R₂ is H;
- 2.12 Formula II or any of 2.5-2.10, wherein R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo

- (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl);
- 2.13 Formula II or any of 2.5-2.10, wherein R₂ is haloC₁₋₆alkyl (e.g., trifluoromethyl or 2,2,2-trifluoroethyl);
- 5 2.14 Formula II or any of 2.5-2.10, wherein R₂ is N(R₁₄)(R₁₅)-C₁₋₆alkyl (e.g., 2-(dimethylamino)ethyl, 2-aminopropyl);
- 2.15 Formula II or any of 2.5-2.10, wherein R₂ is arylC₀₋₆alkyl (e.g., phenyl or benzyl), wherein said aryl is optionally substituted with one or more C₁₋₆alkoxy, for example, C₁₋₆alkoxyarylC₀₋₆alkyl (e.g., 4-methoxybenzyl);
- 10 2.16 Formula II or any of 2.5-2.10, wherein R₂ is heteroarylC₀₋₆alkyl (e.g., pyridinylmethyl), wherein said heteroaryl is optionally substituted with one or more C₁₋₆alkoxy (e.g., C₁₋₆alkoxyheteroarylC₁₋₆alkyl);
- 15 2.17 Formula II or any of 2.5-2.10, wherein R₂ is -G-J wherein G is a single bond or C₁₋₆alkylene (e.g., methylene) and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino, for example:
- 20 -C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., -C₀₋₄alkyl-cyclopentyl, -C₀₋₄alkyl-cyclohexyl or -C₀₋₄alkyl-cyclopropyl), wherein said cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl),
- 25 -C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., -C₀₋₄alkyl-pyrrolidinyl, for example, -C₀₋₄alkylpyrrolidin-3-yl) wherein said heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methyl-pyrrolindin-2-yl, 1-methyl-pyrrolindin-2-yl-methyl or 1-methyl-pyrrolindin-3-yl-methyl);
- 30 2.18 Formula II or any of 2.5-2.10, wherein R₂ is -G-J wherein G is a single bond and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g.,

- oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino;
- 2.19 Formula II or any of 2.5-2.10, wherein R₂ is -G-J wherein G is C₁₋₆alkylene (e.g., methylene) and J is C₃₋₈cycloalkyl or heteroC₃₋₈cycloalkyl (e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino;
- 2.20 Formula II or any of 2.5-2.10, wherein R₂ is -C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., -C₀₋₄alkyl-cyclopentyl, -C₀₋₄alkyl-cyclohexyl or -C₀₋₄alkyl-cyclopropyl), wherein said cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl), -C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., -C₀₋₄alkyl-pyrrolidinyl, for example, -C₀₋₄alkylpyrrolidin-3-yl) wherein said heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methylpyrrolidin-2-yl, 1-methylpyrrolidin-2-yl-methyl or 1-methylpyrrolidin-3-yl-methyl);
- 2.21 Formula II or any of 2.5-2.10, wherein R₂ is 2,2-dimethylpropyl;
- 2.22 Formula II or any of 2.5-2.10, wherein R₂ is isobutyl;
- 2.23 Formula II or any of 2.5-2.10, wherein R₂ is 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl;
- 2.24 Formula II or any of 2.5-2.10, wherein R₂ is cyclopentyl;
- 2.25 Formula II or any of 2.5-2.10, wherein R₂ is -C₀₋₄alkyl-pyrrolidinyl wherein the pyrrolidinyl is optionally substituted with one or more C₁₋₆alkyl, e.g., 1-methylpyrrolidin-3-yl, 1-methylpyrrolidin-2-yl, 1-methylpyrrolidin-2-yl-methyl or 1-methylpyrrolidin-3-yl-methyl);
- 2.26 Formula II or any of 2.5-2.25, wherein R₃ is -D-E-F wherein:
D is
a single bond,
C₁₋₆alkylene (e.g., methylene), or

arylC₁₋₆alkylene (e.g., benzylene or -CH₂C₆H₄-);

E is

a single bond,

C₁₋₄alkylene (e.g., methylene, ethynylene, prop-2-yn-1-ylene),

C₀₋₄alkylarylene (e.g., phenylene or -C₆H₄-, -benzylene- or -CH₂C₆H₄-), wherein the arylene group is optionally substituted with halo (e.g., Cl or F),

heteroarylene (e.g., pyridinylene or pyrimidinylene),

aminoC₁₋₆alkylene (e.g., -CH₂N(H)-),

amino (e.g., -N(H)-);

C₃₋₈cycloalkylene optionally containing one or more heteroatom selected from N, S or O (e.g., piperidinylene),

F is

H,

halo (e.g., F, Br, Cl),

C₁₋₆alkyl (e.g., isopropyl or isobutyl),


haloC₁₋₆alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl),

C₃₋₈cycloalkyl optionally containing one or more atom selected from a group consisting of N, S or O (e.g., cyclopentyl, cyclohexyl, piperidinyl, pyrrolidinyl, tetrahydro-2*H*-pyran-4-yl, or morpholinyl), and optionally substituted with one or more C₁₋₆alkyl (e.g., methyl or isopropyl), for example, 1-methylpyrrolidin-2-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, piperidin-2-yl, 1-methylpiperidin-2-yl, 1-ethylpiperidin-2-yl,

heteroaryl optionally substituted with one or more C₁₋₆alkyl, halo (e.g., fluoro) or haloC₁₋₆alkyl (e.g., pyridyl (for example, pyrid-2-yl), pyrimidinyl (for example, pyrimidin-2-yl), thiadiazolyl (for example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example,

- pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl, 4-methylimidazolyl, 1-methylimidazol-2-yl)), triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazol);
- 5 C₁₋₆alkoxy,
 -O-haloC₁₋₆alkyl (e.g., -O-CF₃),
 C₁₋₆alkylsulfonyl (for example, methylsulfonyl or -S(O)₂CH₃),
 -C(O)-R₁₃, wherein R₁₃ is -N(R₁₄)(R₁₅), C₁₋₆alkyl (e.g.,
 10 methyl), -OC₁₋₆alkyl (e.g., -OCH₃), haloC₁₋₆alkyl (trifluoromethyl), aryl (e.g., phenyl), or heteroaryl,
 -N(R₁₄)(R₁₅);
- 2.27 Formula 2.26, wherein R₁₃ is -N(R₁₄)(R₁₅), C₁₋₆alkyl (e.g., methyl), -OC₁₋₆alkyl (e.g., -OCH₃), haloC₁₋₆alkyl (trifluoromethyl), aryl (e.g., phenyl), or heteroaryl;
- 15 2.28 Formula 2.26, wherein D is C₁₋₆alkylene (e.g., methylene), E is C₃₋₈cycloalkylene optionally containing one or more heteroatom selected from N, S or O (e.g., piperidinylene) and F is C₁₋₆alkyl (e.g., isopropyl or isobutyl), for example, R₃ is isopropylpiperidin-1-ylmethyl;
- 20 2.29 Formula II or any of 2.5-2.25, wherein R₃ is a substituted heteroarylC₀₋₆alkyl, e.g., substituted with haloC₁₋₆alkyl, for example;
- 25 2.30 Formula II or any of 2.5-2.25, wherein R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen

(e.g., Cl or F); and

5

R₁₀ is

halogen (e.g., fluoro or chloro),

C₁₋₆alkyl,

C₃₋₈cycloalkyl,

10

heteroC₃₋₈cycloalkyl (e.g., pyrrolidinyl or

piperidinyl),

haloC₁₋₆alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for

example, pyrid-2-yl) or e.g., thiadiazolyl (for

example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g.,

15

pyrazolyl (for example, pyrazol-1-yl) or

imidazolyl (for example, imidazol-1-yl, 4-

methylimidazolyl, 1-methylimidazol-2-yl)),

triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g.,

tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-

20

methyl-1,2,4-oxadiazol), pyrazolyl (e.g.,

pyrazol-1-yl),

wherein said aryl, heteroaryl, cycloalkyl or

heterocycloalkyl is optionally substituted with

one or more C₁₋₆alkyl (e.g., methyl), halogen

25

(e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g.,

trifluoromethyl), hydroxy, carboxy, -SH or an

additional aryl or heteroaryl (e.g., biphenyl or

pyridylphenyl)

C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl),

30

arylcarbonyl (e.g., benzoyl),

heteroarylcarbonyl,

C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl),


aminocarbonyl

- N(R₁₄)(R₁₅);
- preferably R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl,
optionally substituted with one or more C₁₋₆alkyl
(e.g., methyl), halogen (e.g., chloro or fluoro),
5 haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy,
carboxy, -SH or an additional aryl or heteroaryl
(e.g., biphenyl or pyridylphenyl);
provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
respectively, is not present;
- 10 2.31 Formula 2.30, wherein X, Y and/or Z are independently nitrogen
and R₈, R₉, R₁₁ and R₁₂ are H provided that when X, Y or Z are
nitrogen, R₈, R₉ and R₁₀, respectively, are not present;
- 2.32 Formula 2.30, wherein R₃ is X, Y and Z are C and R₈, R₉, R₁₁ and
R₁₂ are H;
- 15 2.33 Formulae 2.30, 2.31 or 2.32, wherein R₁₀ is
halogen (e.g., fluoro or chloro),
C₁₋₆alkyl,
C₃₋₈cycloalkyl,
heteroC₃₋₈cycloalkyl (e.g., pyrrolidinyl or piperidinyl),
20 haloC₁₋₆alkyl (e.g., trifluoromethyl),
aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for example,
pyrid-2-yl) or e.g., thiadiazolyl (for example, 1,2,3-
thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example,
pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl),
25 triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-
yl), C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazol),
pyrazolyl (e.g., pyrazol-1-yl),
wherein said aryl, heteroaryl, cycloalkyl or heterocycloalkyl is
optionally substituted with one or more C₁₋₆alkyl (e.g.,
30 methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g.,
trifluoromethyl), hydroxy, carboxy, -SH or an additional
aryl or heteroaryl (e.g., biphenyl or pyridylphenyl),
C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl),

- arylcarbonyl (e.g., benzoyl),
 heteroarylcarbonyl,
 C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl),
 aminocarbonyl,
 5 -N(R₁₄)(R₁₅);
- 2.34 Any of formulae 2.30-2.33, wherein R₁₀ is halogen (e.g., fluoro or chloro);
 2.35 Any of formulae 2.30-2.33, wherein R₁₀ is haloC₁₋₆alkyl (e.g., trifluoromethyl);
- 10 2.36 Any of formulae 2.30-2.33, wherein R₁₀ is C₃₋₈cycloalkyl;
 2.37 Any of formulae 2.30-2.33, wherein R₁₀ is C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl);
 2.38 Any of formulae 2.30-2.33, wherein R₁₀ is arylcarbonyl (e.g., benzoyl);
- 15 2.39 Any of formulae 2.30-2.33, wherein R₁₀ is heteroarylcarbonyl;
 2.40 Any of formulae 2.30-2.33, wherein R₁₀ is C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl);
 2.41 Any of formulae 2.30-2.33, wherein R₁₀ is aminocarbonyl;
 2.42 Any of formulae 2.30-2.33, wherein R₁₀ is aryl (e.g., phenyl),
- 20 wherein said aryl is optionally substituted with one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl);
- 2.43 Any of formulae 2.30-2.33, wherein R₁₀ is heteroaryl (e.g., pyridyl,
- 25 (for example, pyrid-2-yl) or e.g., thiadiazolyl (for example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example, pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl), triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazol), pyrazolyl (e.g., pyrazol-1-yl),
- 30 wherein said heteroaryl is optionally substituted with one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl);


- 2.44 Any of formulae 2.30-2.34, wherein R₁₀ is triazolyl (e.g., 1,2,4-triazol-1-yl);
- 2.45 Any of formulae 2.30-2.34, wherein R₁₀ is pyridyl (e.g., pyrid-2-yl) optionally substituted with one or more C₁₋₆alkyl (e.g., methyl),
 5 halogen (e.g., chloro or fluoro);
- 2.46 Any of formulae 2.30-2.34, wherein R₁₀ is pyridyl (e.g., pyrid-2-yl);
- 2.47 Any of formulae 2.30-2.34, wherein R₁₀ is 6-fluoro-pyridyl (e.g., 6-fluoro-pyrid-2-yl);
- 10 2.48 Any of formulae 2.30-2.33, wherein R₁₀ is -N(R₁₄)(R₁₅);
- 2.49 Formula II or any of 2.1-2.48, wherein R₄ is:
 C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl),
 C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
 C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example
 15 pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl),
 -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally
 20 substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);
- 2.50 Formula II or any of 2.1-2.49, wherein R₄ is:
 C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
 25 C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl),
 -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally
 30 substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl).

- 2.51 Formula 2.50, wherein R₄ is C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl);
- 2.52 Formula 2.50, wherein R₄ is C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl);
- 5 2.53 Formula 2.50, wherein R₄ is C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl);
- 2.54 Formula 2.50, wherein R₄ is -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);
- 10 2.55 Formula 2.50, wherein R₄ is phenyl optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);
- 15 2.56 Formula 2.50, wherein R₄ is phenyl;
- 2.57 Formula 2.50, wherein R₄ is 4-fluorophenyl;
- 20 2.58 Formula 2.50, wherein R₄ is 4-hydroxyphenyl;
- 2.59 Formula 2.50, wherein R₄ is C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
- 2.60 Any of the preceding formulae wherein R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;
- 25 2.61 Any of the preceding formulae wherein R₆ is C₃₋₈cycloalkyl;
- 2.62 Any of the preceding formulae wherein R₆ is H;
- 2.63 Any of the preceding formulae wherein R₆ is C₁₋₆alkyl (e.g., methyl or ethyl);
- 2.64 Any of the preceding formulae wherein R₁₄ and R₁₅ are independently H or C₁₋₆alkyl;
- 30 2.65 Any of the preceding formulae wherein R₁₄ and R₁₅ are H;
- 2.66 Any of the preceding formulae wherein the Compound is selected from:



2.67 any of the preceding formulae wherein the compounds inhibit
 5 phosphodiesterase-mediated (e.g., PDE1-mediated, especially
 PDE1B-mediated) hydrolysis of cGMP, e.g., with an IC_{50} of less
 than $1\mu M$, preferably less than 25 nM, preferably less than 10 nM
 in an immobilized-metal affinity particle reagent PDE assay, for
 example, as described in Example 5;

10 **[0008]** In another aspect, the Compound of Formula II is a Compound of Formula
 II(a):




Formula II(a)

wherein

15 (i) Q is $-C(=O)-$, $-C(=S)-$, $-C(=N(R_6))-$ or $-C(R_{14})(R_{15})-$;

- (ii) R_1 is H or C_{1-6} alkyl (e.g., methyl or ethyl);
- (iii) R_2 is C_{1-6} alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxy C_{1-6} alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- (iv) R_3 is attached to one of the nitrogens on the pyrazolo portion of Formula II(a) and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R_8 , R_9 , R_{11} and R_{12} are independently H or halogen

(e.g., Cl or F); and

R_{10} is

halogen (e.g., fluoro or chloro),

C_{1-6} alkyl,

C_{3-8} cycloalkyl,

hetero C_{3-8} cycloalkyl (e.g., pyrrolidinyl or piperidinyl),

halo C_{1-6} alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for example, pyrid-2-yl) or e.g., thiadiazolyl (for

example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g.,

pyrazolyl (for example, pyrazol-1-yl) or

imidazolyl (for example, imidazol-1-yl)),

triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g.,

tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazol), pyrazolyl (e.g., pyrazol-1-yl),

wherein said aryl, heteroaryl, cycloalkyl or

heterocycloalkyl is optionally substituted with

one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl)

C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl), arylcarbonyl (e.g., benzoyl), heteroarylcarbonyl, C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl), Aminocarbonyl, -N(R₁₄)(R₁₅);

preferably R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl, optionally substituted with one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl);

provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀, respectively, is not present;

(v) R₄ is:

C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl), C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl), C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl), -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally

substituted with one or more halo (e.g., 4-fluorophenyl),
hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy
or another aryl group (e.g., biphenyl-4-ylmethyl);


(vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;

5

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form.

[0010] In another aspect, the Compound of Formula II is a compound of Formula II(b):



10

Formula II(b)

wherein:

(i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;


(ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);

(iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyc₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),

15

(iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(b) and is a moiety of Formula A

20



Formula A

wherein:

X, Y and Z are, independently, N or C,

5 R₈, R₉, R₁₁ and R₁₂ are independently H or halogen
(e.g., Cl or F); and

R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl, or pyrrolidinyl
optionally substituted with one or more C₁₋₆alkyl
(e.g., methyl), halogen (e.g., chloro or fluoro),
10 haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy,
carboxy, -SH or an additional aryl or heteroaryl
(e.g., biphenyl or pyridylphenyl);
provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
respectively, is not present;

15 (v) R₄ is:


C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl),
C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example
pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for
20 example, piperidin-1-yl), morpholinyl),
-C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆
alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-
3-yl) wherein said aryl or heteroaryl is optionally
substituted with one or more halo (e.g., 4-fluorophenyl),
25 hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy
or another aryl group (e.g., biphenyl-4-ylmethyl);

(vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form.


30 **[0011]** In another aspect, the Compound of Formula II is a compound of Formula
II(c):



Formula II(c)

wherein:

- 5 (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyc₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- 10 (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(c) and is a moiety of Formula A



Formula A

wherein:

- 15 X, Y and Z are, independently, N or C,
- R₈, R₉, R₁₁ and R₁₂ are independently H or halogen (e.g., Cl or F); and
- 20 R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl, pyrrolidinyl optionally substituted with one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro),

haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy,
 carboxy, -SH or an additional aryl or heteroaryl
 (e.g., biphenyl or pyridylphenyl);
 provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
 respectively, is not present;

(v) R₄ is:


-C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆
 alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-
 3-yl) wherein said aryl or heteroaryl is optionally
 substituted with one or more halo (e.g., 4-fluorophenyl),
 hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy
 or another aryl group (e.g., biphenyl-4-ylmethyl);

(vi) R₆ is H or C₁₋₆alkyl (e.g., methyl or ethyl);

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form.

[0012] In another aspect, the Compound of Formula II is a Compound of Formula
 II(d):




Formula II(d)

wherein:

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally
 substituted with one or more halo (e.g., fluoro) or hydroxy
 (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),

- (iv) R_3 is attached to one of the nitrogens on the pyrazolo portion of Formula II(d) and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R_8 , R_9 , R_{11} and R_{12} are independently H or halogen (e.g., Cl or F); and

R_{10} is phenyl or pyridyl, e.g., 2-pyridyl, pyrrolidinyl optionally substituted with one or more C_{1-6} alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), halo C_{1-6} alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl);

provided that when X, Y or Z is nitrogen, R_8 , R_9 or R_{10} , respectively, is not present;

- (v) R_4 is :

C_{3-8} cycloalkyl (e.g., cyclopentyl or cyclohexyl),

C_{3-8} heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl),

$-C_{0-6}$ alkylaryl (e.g., phenyl or benzyl) or $-C_{0-6}$


alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C_{1-6} alkyl, C_{1-6} alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);

- (vi) R_6 is H, C_{1-6} alkyl (e.g., methyl or ethyl);
 (vii) R_{14} and R_{15} are independently H or C_{1-6} alkyl,

in free or salt form.

[0013] In another aspect, the Compound of Formula II is a Compound of Formula


5 II(e):



Formula II(e)

wherein:

- (i) Q is $-C(=O)-$, $-C(=S)-$, $-C(=N(R_6))-$ or $-C(R_{14})(R_{15})-$;
 10 (ii) R_1 is H or C_{1-6} alkyl (e.g., methyl or ethyl);
 (iii) R_2 is C_{1-6} alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxy C_{1-6} alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
 15 (iv) R_3 is attached to one of the nitrogens on the pyrazolo portion of Formula II(e) and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,


R₈, R₉, R₁₁ and R₁₂ are independently H or halogen
(e.g., Cl or F); and

R₁₀ is phenyl, pyridyl (e.g., 2-pyridyl), pyrrolidinyl
(e.g., pyrrolidin-3-yl or pyrrolidin-1-yl) or
piperidinyl (e.g., piperidin-1-yl), optionally
substituted with one or more C₁₋₆alkyl (e.g., methyl)
or halogen (e.g., chloro or fluoro);

provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
respectively, is not present;

- (v) R₄ is phenyl optionally substituted with one or more halo (e.g.,
4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl,
C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);
- (vi) R₆ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,
- in free or salt form.


[0014] In still another aspect, the Compound of Formula II is a Compound of
Formula II(f):



Formula II(f)

wherein:

- (i) Q is -C(=O)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl);
- (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of
Formula II(f) and is a moiety of Formula A



Formula A

wherein:

5

X, Y and Z are C;

R₈, R₉, R₁₁ and R₁₂ are H; and

R₁₀ is pyridyl (e.g., 2-pyridyl), optionally substituted with one or more halogen (e.g., fluoro);

(v) R₄ is

10

phenyl optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl), preferably phenyl;


C₁₋₆alkyl (e.g., n-propyl),

15

C₃₋₈cycloalkyl (e.g., cyclopentyl),

in free or salt form.

[0015] In still another embodiment, the compound of the invention as hereinbefore described is selected from any of the following:



in free or salt form.

[0016] Preferably, the compound of Formula II or any of II(a)-II(e) is a compound


5 wherein Q is -C(=O)-.

[0017] In a particular embodiment, the Compound of any of Formulae II, II(a)-II(f), 2.1-2.67 contains the proviso that said compound is not:

- (a) 7-[(2,4-dichlorophenyl)methyl]-3-ethoxy-2,5-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- 10 (b) 7-[(4-chlorophenyl)methyl]-3-ethoxy-2,5-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (c) 2-[(2,5-dichlorophenyl)methyl]-3-ethoxy-5,7-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (d) 2-[(4-bromophenyl)methyl]-3-ethoxy-5,7-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- 15 (e) 3-[(4-methoxyphenyl)methoxy]-2,5,7-trimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (f) 1-[(3-bromophenyl)methyl]-5,7-dimethyl-3-(2-propen-1-yloxy)-1H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione.

20 **[0018]** If not otherwise specified or clear from context, the following terms herein have the following meanings:

- 5 (a) "Alkyl" as used herein is a saturated or unsaturated hydrocarbon moiety, preferably saturated, preferably having one to six carbon atoms, which may be linear or branched, and may be optionally mono-, di- or tri- substituted, e.g., with halogen (e.g., chloro or fluoro), hydroxy, or carboxy.
- 10 (b) "Cycloalkyl" as used herein is a saturated or unsaturated nonaromatic hydrocarbon moiety, preferably saturated, preferably comprising three to eight carbon atoms, at least some of which form a nonaromatic mono- or bicyclic, or bridged cyclic structure, and which may be optionally substituted, e.g., with halogen (e.g., chloro or fluoro), hydroxy, or carboxy. Wherein the cycloalkyl optionally contains one or more atoms selected from N and O and/or S, said cycloalkyl may optionally be a heterocycloalkyl.
- 15 (c) "Heterocycloalkyl" is, unless otherwise indicated, saturated or unsaturated nonaromatic hydrocarbon moiety, preferably saturated, preferably comprising three to nine carbon atoms, at least some of which form a nonaromatic mono- or bicyclic, or bridged cyclic structure, wherein at least one carbon atom is replaced with N, O or S, which heterocycloalkyl may be optionally substituted, e.g., with
- 20 halogen (e.g., chloro or fluoro), hydroxy, or carboxy.
- 25 (d) "Aryl" as used herein is a mono or bicyclic aromatic hydrocarbon, preferably phenyl, optionally substituted, e.g., with alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloalkyl (e.g., trifluoromethyl), hydroxy, carboxy, or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl).
- 30 (e) "Heteroaryl" as used herein is an aromatic moiety wherein one or more of the atoms making up the aromatic ring is sulfur or nitrogen rather than carbon, e.g., pyridyl or thiadiazolyl, which may be optionally substituted, e.g., with alkyl, halogen, haloalkyl, hydroxy or carboxy.
- (f) Wherein E is phenylene, the numbering is as follows:



(g) It is intended that wherein the substituents end in “ene”, for example, alkylene, phenylene or arylalkylene, said substituents are intended to bridge or be connected to two other substituents. Therefore, methylene is intended to be $-\text{CH}_2-$ and phenylene intended to be $-\text{C}_6\text{H}_4-$ and arylalkylene is intended to be $-\text{C}_6\text{H}_4-\text{CH}_2-$ or $-\text{CH}_2-\text{C}_6\text{H}_4-$.

[0019] The term “substituted,” as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. Similarly, the substituents defined for the Compounds of the Invention are intended to result in stable compounds.

[0020] Compounds of the Invention may exist in free or salt form, e.g., as acid addition salts. In this specification unless otherwise indicated, language such as “Compounds of the Invention” is to be understood as embracing the compounds described herein, e.g., (1- or 2- and/or 5 and/or 7-substituted)-(3-oxy)-(4H, 4-imino, 4-thioxo or 4-oxo)-pyrazolo[3,4-d]pyrimidin-6-ones, e.g., a compound of formula II, or any of II(a)-II(f), or 2.1-2.67, in any form, for example free or acid addition salt form, or where the compounds contain acidic substituents, in base addition salt form.

The Compounds of the Invention are intended for use as pharmaceuticals, therefore pharmaceutically acceptable salts are preferred. Salts which are unsuitable for pharmaceutical uses may be useful, for example, for the isolation or purification of free Compounds of the Invention or their pharmaceutically acceptable salts, are therefore also included. In a particular embodiment, the salt of the compounds of the invention is a formic acid addition salt.

[0021] Compounds of the Invention may in some cases also exist in prodrug form. A prodrug form is compound which converts in the body to a Compound of the Invention. For example, when the Compounds of the Invention contain hydroxy (or carboxy) substituents, these substituents may form physiologically hydrolysable and acceptable esters, e.g., C_{1-4} alkyl carboxylic acid ester. As used herein,

- “physiologically hydrolysable and acceptable ester” means esters of Compounds of the Invention which are hydrolysable under physiological conditions to yield acids (in the case of Compounds of the Invention which have hydroxy substituents) or alcohols (in the case of Compounds of the Invention which have carboxy substituents) which
- 5 are themselves physiologically tolerable at doses to be administered. Therefore, wherein the Compound of the Invention contains a hydroxy group, for example, Compound-OH, the acyl ester prodrug of such compound, i.e., Compound-O-C(O)-C₁₋₄alkyl, can hydrolyze in the body to form physiologically hydrolysable alcohol (Compound-OH) on the one hand and acid on the other (e.g., HOC(O)-C₁₋₄alkyl).
- 10 Alternatively, wherein the Compound of the Invention contains a carboxylic acid, for example, Compound-C(O)OH, the acid ester prodrug of such compound, Compound-C(O)O-C₁₋₄alkyl can hydrolyze to form Compound-C(O)OH and HO-C₁₋₄alkyl. As will be appreciated, the term thus embraces conventional pharmaceutical prodrug forms.
- 15 **[0022]** The invention also provides methods of making the Compounds of the Invention and methods of using the Compounds of the Invention for treatment of diseases and disorders as set forth below (especially treatment of diseases characterized by reduced dopamine D1 receptor signaling activity, such as Parkinson’s disease, Tourette’s Syndrome, Autism, fragile X syndrome, ADHD,
- 20 restless leg syndrome, depression, cognitive impairment of schizophrenia, narcolepsy and diseases that may be alleviated by the enhancement of progesterone-signaling such as female sexual dysfunction), or a disease or disorder such as psychosis or glaucoma). This list is not intended to be exhaustive and may include other diseases and disorders as set forth below.
- 25 **[0023]** In another embodiment, the invention further provides a pharmaceutical composition comprising a Compound of the Invention, e.g., (1- or 2- and/or 5 and/or 7-substituted)-(3-oxy)-(4H, 4-imino, 4-thioxo or 4-oxo)-pyrazolo[3,4-d]pyrimidin-6-ones, e.g., a compound of formula II, or any of II(a)-II(f), or 2.1-2.67, in free or pharmaceutically acceptable salt form, in admixture with a pharmaceutically
- 30 acceptable carrier.

DETAILED DESCRIPTION OF THE INVENTION

Methods of Making Compounds of the Invention

[0024] The Compounds of the Invention and their pharmaceutically acceptable salts may be made using the methods as described and exemplified herein and by methods similar thereto and by methods known in the chemical art. Such methods include, but not limited to, those described below. In the description of the synthetic methods described herein, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the conditions standard for that reaction, which should be readily recognized by one skilled in the art. Therefore, at times, the reaction may require to be run at elevated temperature or for a longer or shorter period of time. It is understood by one skilled in the art of organic synthesis that functionality present on various portions of the molecule must be compatible with the reagents and reactions proposed. If not commercially available, starting materials for these processes may be made by procedures, which are selected from the chemical art using techniques which are similar or analogous to the synthesis of known compounds. In particular, the intermediates and starting materials for the Compounds of the Invention may be prepared by methods and processes as described in WO 2007/143705; WO 2009/073210; PCT/US2009/06444; and PCT/US2009/06443. All references cited herein are hereby incorporated by reference in their entirety.

[0025] The Compounds of the Invention include their enantiomers, diastereoisomers and racemates, as well as their polymorphs, hydrates, solvates and complexes. Some individual compounds within the scope of this invention may contain double bonds. Representations of double bonds in this invention are meant to include both the E and the Z isomer of the double bond. In addition, some compounds within the scope of this invention may contain one or more asymmetric centers. This invention includes the use of any of the optically pure stereoisomers as well as any combination of stereoisomers.

[0026] As will be appreciated by those skilled in the art, the Compounds of the Invention may exhibit keto-enol tautomerization. Therefore, the invention as defined in the present invention is to be understood as embracing both the structures as set forth herewith and their tautomeric forms.

[0027] It is also intended that the Compounds of the Invention encompass their stable and unstable isotopes. Stable isotopes are nonradioactive isotopes which contain one additional neutron compared to the abundant nuclides of the same species (i.e.,

element). It is expected that the activity of compounds comprising such isotopes would be retained, and such compound would also have utility for measuring pharmacokinetics of the non-isotopic analogs. For example, the hydrogen atom at a certain position on the Compounds of the Invention may be replaced with deuterium (a stable isotope which is non-radioactive). Examples of known stable isotopes include, but not limited to, deuterium, ^{13}C , ^{15}N , ^{18}O . Alternatively, unstable isotopes, which are radioactive isotopes which contain additional neutrons compared to the abundant nuclides of the same species (i.e., element), e.g., ^{123}I , ^{131}I , ^{125}I , ^{11}C , ^{18}F , may replace the corresponding abundant species, e.g., I, C and F respectively.

Another example of useful isotope of the compound of the invention is the ^{11}C isotope. These radio isotopes are useful for radio-imaging and/or pharmacokinetic studies of the compounds of the invention. Methods of making isotopes of PDE1 inhibitors disclosed in WO 2011/043816, the contents of which are incorporated by reference in their entirety, may be used for making the isotopes of the compounds of the current invention.

[0028] Melting points are uncorrected and (dec) indicates decomposition.

Temperature are given in degrees Celsius ($^{\circ}\text{C}$); unless otherwise stated, operations are carried out at room or ambient temperature, that is, at a temperature in the range of 18-25 $^{\circ}\text{C}$. Chromatography means flash chromatography on silica gel; thin layer chromatography (TLC) is carried out on silica gel plates. NMR data is in the delta values of major diagnostic protons, given in parts per million (ppm) relative to tetramethylsilane (TMS) as an internal standard. Conventional abbreviations for signal shape are used. Coupling constants (J) are given in Hz. For mass spectra (MS), the lowest mass major ion is reported for molecules where isotope splitting results in multiple mass spectral peaks. Solvent mixture compositions are given as volume percentages or volume ratios. In cases where the NMR spectra are complex, only diagnostic signals are reported.


[0029] Terms and abbreviations:

BOC = di-*tert*-butyl dicarbonate,
BuLi = n-butyllithium
Bu^tOH = *tert*-butyl alcohol,
CAN = ammonium cerium (IV) nitrate,
DIPEA = diisopropylethylamine,

- DMF = N,N-dimethylformamide,
 DMSO = dimethyl sulfoxide,
 Et₂O = diethyl ether,
 EtOAc = ethyl acetate,
 5 equiv. = equivalent(s),
 h = hour(s),
 HPLC = high performance liquid chromatography,
 K₂CO₃ = potassium carbonate,
 LDA = lithium diisopropylamide
 10 MeOH = methanol,
 NaHCO₃ = sodium bicarbonate,
 NBS = N-bromosuccinimide
 NCS = N-chlorosuccinimide
 NH₄OH = ammonium hydroxide,
 15 Pd₂(dba)₃ = tris[dibenzylideneacetone]dipalladium(0)
 PMB = p-methoxybenzyl,
 POCl₃ = phosphorous oxychloride,
 SOCl₂ = thionyl chloride,
 TFA = trifluoroacetic acid,
 20 THF = tetrahydrofuran.


[0030] The synthetic methods in this invention are illustrated below. The significances for the R groups are as set forth in any of the formulae above, e.g., for formula II, II(a)-II(f) unless otherwise indicated.

- [0031] In an aspect of the invention, intermediate compounds of formula **IIb** can be
 25 synthesized by reacting a compound of formula **IIa** with a dicarboxylic acid, acetic anhydride and acetic acid mixing with heat, e.g., to about 90°C for about 3 hours and then cooled.




wherein R₁ is H or C₁₋₄alkyl [e.g., methyl].

[0032] Intermediate **IIc** can be prepared by for example reacting intermediate of **IIb** with for example a chlorinating compound such as POCl₃, sometimes with small amounts of water and heated, e.g., heating to about 80°C for about 4 hours and then cooled.




Intermediate **IIc** may be formed by reacting intermediate **IIc** with for example a P¹-L in a solvent such as DMF and a base such as K₂CO₃, sodium bicarbonate, cesium carbonate, sodium hydroxide, triethylamine, diisopropylethylamine or the like at room temperature or with heating.




wherein P¹ is a protective group [e.g., *p*-methoxybenzyl group (PMB) or BOC]; L is a leaving group such as a halogen, mesylate, or tosylate. Preferably, P¹ is PMB and the base is potassium carbonate.


[0033] Intermediate **Ile** may be prepared by reacting intermediate **IIc** with hydrazine or hydrazine hydrate in a solvent such as methanol and with heating, e.g. refluxed for about 4 hours and then cooled.



[0034] Intermediate **IIf** can be synthesized by for example reacting intermediate **Ile** with POCl₃ and DMF (**Reaction 1**):




- [0035] Intermediate **IIg** may be formed by reacting intermediate **IIf** with for example a $\text{R}_3\text{-L}$ in a solvent such as DMF and a base such as K_2CO_3 , sodium bicarbonate, cesium carbonate, triethylamine or the like, preferably K_2CO_3 , at room temperature or with heating (**Reaction 2**):




wherein R_3 is as defined previously [e.g. -D-E-F or moiety of Formula A]; L is a leaving group such as a halogen, mesylate, or tosylate.

- 10 [0036] Intermediate **IIh** may be formed by reacting intermediate **IIg** with for example C_2Cl_6 , NCS, NBS or I_2 in a solvent such as THF and a base such as LiHMDS, LDA or BuLi at low temperature (**Reaction 3**):




- 15 [0037] Intermediate **IIi** (Compound of the Invention with $\text{R}_2 = \text{P}^1$) may be formed by reacting intermediate **IIh** with for example a R_4OH in a solvent such as DMF and a base such as K_2CO_3 with heating. Sometimes, R_4OH itself can be used as a solvent (**Reaction 4**):




wherein R₄ is as defined previously in any of Formulae II or II(a)-II(f).

- [0038] Intermediate **IIj** (i.e., Compound of the Invention wherein R₂ = H) may be synthesized from intermediate **IIIi** by removing the protective group P¹ with an appropriate method. For example, if P¹ is a *p*-methoxybenzyl group, it can be removed by for example TFA/TFMSA in methylene chloride at room temperature (**Reaction 5**):



- Wherein P¹ is BOC, the compound may be deprotected by using acid such as hydrochloric acid or TFA.


[0039] The Compounds of the Invention may be formed by reacting intermediate **IIj** with for example a R₂-L in a solvent such as DMF and a base such as K₂CO₃ at room temperature or with heating (**Reaction 6**):




- wherein L is a leaving group such as a halogen, mesylate, or tosylate.

[0040] Alternatively, the Compounds of the Invention may be prepared in the order as described below.


- [0041] Intermediate **IIIa** may be formed by reacting intermediate **IIc** with for example a R₂-L in a solvent such as DMF and a base such as K₂CO₃ at room temperature or with heating (**Reaction 7**).




[0042] Intermediate **IIIb** may be prepared by reacting intermediate **IIIa** with anhydrous hydrazine or hydrazine hydrate in a solvent such as methanol and heated for several hours and then cooled (**Reaction 8**):




[0043] Intermediate **IIIc** may be formed by for example reacting intermediate **IIIb** with POCl₃ and DMF (**Reaction 10**):




[0044] Intermediate **IIId** may be formed by reacting intermediate **IIIc** with for example a R₃-L in a solvent such as DMF and a base such as K₂CO₃, sodium bicarbonate, cesium carbonate, triethylamine or the like, preferably K₂CO₃, at room temperature or with heating (**Reaction 11**):




[0045] Intermediate **IIIe** may be formed by reacting intermediate **IIId** with for example C₂Cl₆, NCS, NBS or I₂ in a solvent such as THF and a base such as LiHMDS, LDA or BuLi at low temperature (**Reaction 12**):




- [0046] Compound of Formula II may be formed by reacting intermediate IIIe with for example a R₄OH in a solvent such as DMF and a base such as K₂CO₃ with heating. Sometimes, R₄OH itself can be used as a solvent (**Reaction 13**):




- [0047] The 4-thioxo Compounds of the Invention, i.e., any of Formulae II or II(a)-II(f), wherein Q is C(=S) may then be prepared by reacting the dione Compounds of the Invention, i.e., any of Formulae II or II(a)-II(f), wherein Q is C(=O) with P₄S₁₀ in a microwave vial in the presence of a base, e.g., pyridine, and heating the mixture to an elevated temperature, e.g., in a microwave, e.g., to about 150°C.



- [0048] The 4-imino Compounds of the Invention, i.e., any of Formulae II or II(a)-II(f), wherein Q is C(=N(R₆)) may in turn be converted from the thioxo compound (i.e., any of Formulae II or II(a)-II(f), wherein with Q is X(=S)) by reacting the 4-thioxo compound with NH₂(R₆) in the presence of HgCl₂, e.g., in a solvent such as THF, and heating the reaction mixture to an elevated temperature, e.g., in a microwave, e.g., to about 110°C.



- [0049]** The Compounds of the Invention, any of Formulae II or II(a)-II(f), wherein Q is C(R₁₄)(R₁₅) may also be prepared by reacting the dione Compounds of the Invention, i.e., any of Formulae II or II(a)-II(f), wherein Q is C(=O) with a reducing agent, e.g., diisobutylaluminum hydride (DIBAL-H), lithium aluminum hydride, sodium borohydride, preferably, DIBAL-H.



- [0050]** The invention thus provides methods of making a 4-thioxo Compounds of the Invention, i.e., any of Formulae II or II(a)-II(f), wherein Q is C(=S) as hereinbefore described, for example, comprising reacting a Compound of Formula II wherein Q is C(=O), with P₄S₁₀ in the presence of a base, e.g., pyridine, and heating the reaction mixture to an elevated temperature, e.g., to >50°C, e.g., >100°C, e.g., >150°C, for example, in a microwave to about 150°C.


- [0051]** The invention also provides methods of making 4-imino Compounds of the Invention, e.g., any Compound of Formula II, wherein Q is C(=N(R₆)) as hereinbefore described, for example, comprising reacting the Compound of Formula II, wherein Q is C(=S), with NH₂(R₆) in the presence of HgCl₂, e.g., in a solvent such as THF, and heating the reaction mixture in a microwave, e.g., to >50°C, e.g., >75°C, e.g., >100°C, for example, in a microwave to about 110°C.

- [0052]** The invention also provides methods of making Compounds of Formula II, wherein Q is CH₂ comprising reacting a Compound of Formula II, wherein Q is C(=O), with a reducing agent, e.g., diisobutylaluminum hydride (DIBAL-H), lithium aluminum hydride, sodium borohydride, preferably, DIBAL-H.

[0053] Alternatively, the Compounds of the Invention, e.g., a Compound of Formula II wherein Q is C(=S), C(=N(R₆)) or CH₂, may be prepared first before attaching on R₁, R₂, R₃ and/or R₄.


[0054] The invention thus provides methods of making Compounds of the Invention, wherein Q is C(=O) and the remaining substituents are as described above, for example, comprising

(i) reacting Intermediate IIIe with a compound of formula R₄OH, wherein R₄ is as hereinbefore described under basic conditions with heating, for example wherein the 2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione is a compound of Formula IIIe:



wherein R₁, R₂ and R₃ are as defined above; and/or

(ii) reacting intermediate IIj with a compound of formula L-R₂ wherein L is a leaving group, e.g., halogen, mesylate, or tosylate, and R₂ is as hereinbefore described, for example wherein R₂ is isobutyl, N-methylpyrrolidin-3-yl, or neopentyl group, e.g., under basic conditions:



wherein R₁, R₃ and R₄ are as defined above, e.g., with reference to Formula II.

Methods of using Compounds of the Invention

[0055] The Compounds of the Invention are useful in the treatment of diseases characterized by disruption of or damage to cAMP and cGMP mediated pathways, e.g., as a result of increased expression of PDE1 or decreased expression of cAMP and cGMP due to inhibition or reduced levels of inducers of cyclic nucleotide synthesis, such as dopamine and nitric oxide (NO). By preventing the degradation of cAMP and cGMP by PDE1B, thereby increasing intracellular levels of cAMP and cGMP, the Compounds of the Invention potentiate the activity of cyclic nucleotide synthesis inducers.

[0056] The invention provides methods of treatment of any one or more of the following conditions:

- (i) Neurodegenerative diseases, including Parkinson's disease, restless leg tremors, dyskinesias, Huntington's disease, Alzheimer's disease, and drug-induced movement disorders;
- (ii) Mental disorders, including depression, attention deficit disorder, attention deficit hyperactivity disorder, bipolar illness, anxiety, sleep disorders, e.g., narcolepsy, cognitive impairment, dementia, Tourette's syndrome, autism, fragile X syndrome, psychostimulant withdrawal, and drug addiction;
- (iii) Circulatory and cardiovascular disorders, including cerebrovascular disease, stroke, congestive heart disease, hypertension or pulmonary hypertension, and sexual dysfunction;
- (iv) Respiratory and inflammatory disorders, including asthma, chronic obstructive pulmonary disease, and allergic rhinitis, as well as autoimmune and inflammatory diseases;
- (v) Any disease or condition characterized by low levels of cAMP and/or cGMP (or inhibition of cAMP and/or cGMP signaling pathways) in cells expressing PDE1; and/or
- (vi) Any disease or condition characterized by reduced dopamine D1 receptor signaling activity,

comprising administering an effective amount of a Compound of the Invention, e.g., compound according to any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, or a composition comprising such a

Compound of the Invention, to a human or animal patient in need thereof. This method also encompasses administering an effective amount of a Compound of the Invention in free or pharmaceutically acceptable salt form. In another aspect, the invention provides a method of treatment of the conditions disclosed above

5 comprising administering a therapeutically effective amount of compound according to any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, or a composition comprising the same, to a human or animal patient in need thereof.

[0057] In an especially preferred embodiment, the invention provides methods of
10 treatment or prophylaxis for narcolepsy. In this embodiment, PDE 1 Inhibitors may be used as a sole therapeutic agent, but may also be used in combination or for co-administration with other active agents. Thus, the invention further comprises a method of treating narcolepsy comprising administering simultaneously, sequentially, or contemporaneously administering therapeutically effective amounts of

15 (i) a PDE 1 Inhibitor of the Invention, e.g., a compound according to any of Formulae II, II(a)-II(f) or 2.1-2.67; and
(ii) a compound to promote wakefulness or regulate sleep, e.g., selected from
(a) central nervous system stimulants-amphetamines and amphetamine like compounds, e.g., methylphenidate, dextroamphetamine, methamphetamine,
20 and pemoline; (b) modafinil, (c) antidepressants, e.g., tricyclics (including imipramine, desipramine, clomipramine, and protriptyline) and selective serotonin reuptake inhibitors (including fluoxetine and sertraline); and/or (d) gamma hydroxybutyrate (GHB),

in free or pharmaceutically acceptable salt form, to a human or animal patient in need
25 thereof. In another embodiment, the invention provides methods of treatment or prophylaxis for narcolepsy as herein before described, wherein the PDE1 inhibitor is in a form of a pharmaceutical composition.

[0058] In another embodiment, the invention further provides methods of
treatment or prophylaxis of a condition which may be alleviated by the enhancement
30 of the progesterone signaling comprising administering an effective amount of a Compound of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form. Disease or condition that may be ameliorated by enhancement of progesterone signaling include, but are not limited to, female

sexual dysfunction, secondary amenorrhea (e.g., exercise amenorrhoea, anovulation, menopause, menopausal symptoms, hypothyroidism), pre-menstrual syndrome, premature labor, infertility, for example infertility due to repeated miscarriage, irregular menstrual cycles, abnormal uterine bleeding, osteoporosis, autoimmune disease, multiple sclerosis, prostate enlargement, prostate cancer, and hypothyroidism. For example, by enhancing progesterone signaling, the PDE 1 inhibitors may be used to encourage egg implantation through effects on the lining of uterus, and to help maintain pregnancy in women who are prone to miscarriage due to immune response to pregnancy or low progesterone function. The novel PDE 1 inhibitors, e.g., as described herein, may also be useful to enhance the effectiveness of hormone replacement therapy, e.g., administered in combination with estrogen/estradiol/estriol and/or progesterone/progestins in postmenopausal women, and estrogen-induced endometrial hyperplasia and carcinoma. The methods of the invention are also useful for animal breeding, for example to induce sexual receptivity and/or estrus in a nonhuman female mammal to be bred.

[0059] In this embodiment, PDE 1 Inhibitors may be used in the foregoing methods of treatment or prophylaxis as a sole therapeutic agent, but may also be used in combination or for co-administration with other active agents, for example in conjunction with hormone replacement therapy. Thus, the invention further comprises a method of treating disorders that may be ameliorated by enhancement of progesterone signaling comprising administering simultaneously, sequentially, or contemporaneously administering therapeutically effective amounts of

- (i) a PDE 1 Inhibitor of the Invention, e.g., A Compound of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, and
- (ii) a hormone, e.g., selected from estrogen and estrogen analogues (e.g., estradiol, estriol, estradiol esters) and progesterone and progesterone analogues (e.g., progestins)

in free or pharmaceutically acceptable salt form, to a human or animal patient in need thereof.

[0060] The invention also provides a method for enhancing or potentiating dopamine D1 intracellular signaling activity in a cell or tissue comprising contacting said cell or tissue with an amount of a Compound of the Invention sufficient to inhibit PDE1B activity.

[0061] The invention also provides a method for enhancing or potentiating progesterone signaling activity in a cell or tissue comprising contacting said cell or tissue with an amount of a Compound of the Invention sufficient to inhibit PDE1B activity.

- 5 [0062] The invention also provides a method for treating a PDE1-related, especially PDE1B-related disorder, a dopamine D1 receptor intracellular signaling pathway disorder, or disorders that may be alleviated by the enhancement of the progesterone signaling pathway in a patient in need thereof comprising administering to the patient an effective amount of a Compound of the Invention that inhibits
- 10 PDE1B, wherein PDE1B activity modulates phosphorylation of DARPP-32 and/or the GluR1 AMPA receptor.

[0063] “The Compound of the Invention” referred to above includes any of the compounds disclosed herein, e.g., a Compound of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form.

- 15 [0064] In another aspect, the invention also provides a method for the treatment for glaucoma or elevated intraocular pressure comprising topical administration of a therapeutically effective amount of a phosphodiesterase type I (PDE1) Inhibitor of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, in an ophthalmically compatible carrier to the eye of a patient in
- 20 need thereof. However, treatment may alternatively include a systemic therapy. Systemic therapy includes treatment that can directly reach the bloodstream, or oral methods of administration, for example.

- [0065] The invention further provides a pharmaceutical composition for topical ophthalmic use comprising a PDE1 inhibitor; for example an ophthalmic solution,
- 25 suspension, cream or ointment comprising a PDE1 Inhibitor of the Invention, e.g., a Compound of Formula II, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or ophthalmologically acceptable salt form, in combination or association with an ophthalmologically acceptable diluent or carrier.

- [0066] Optionally, the PDE1 inhibitor may be administered sequentially or
- 30 simultaneously with a second drug useful for treatment of glaucoma or elevated intraocular pressure. Where two active agents are administered, the therapeutically effective amount of each agent may be below the amount needed for activity as monotherapy. Accordingly, a subthreshold amount (i.e., an amount below the level

necessary for efficacy as monotherapy) may be considered therapeutically effective and also may also be referred alternatively as an effective amount. Indeed, an advantage of administering different agents with different mechanisms of action and different side effect profiles may be to reduce the dosage and side effects of either or both agents, as well as to enhance or potentiate their activity as monotherapy.

[0067] The invention thus provides the method of treatment of a condition selected from glaucoma and elevated intraocular pressure comprising administering to a patient in need thereof an effective amount, e.g., a subthreshold amount, of an agent known to lower intraocular pressure concomitantly, simultaneously or sequentially with an effective amount, e.g., a subthreshold amount, of a PDE1 Inhibitor of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, such that amount of the agent known to lower intraocular pressure and the amount of the PDE1 inhibitor in combination are effective to treat the condition. In one embodiment, one or both of the agents are administered topically to the eye. Thus the invention provides a method of reducing the side effects of treatment of glaucoma or elevated intraocular pressure by administering a reduced dose of an agent known to lower intraocular pressure concomitantly, simultaneously or sequentially with an effective amount of a PDE1 inhibitor. However, methods other than topical administration, such as systemic therapeutic administration, may also be utilized.

[0068] The optional additional agent or agents for use in combination with a PDE1 inhibitor may, for example, be selected from the existing drugs comprise typically of instillation of a prostaglandin, pilocarpine, epinephrine, or topical beta-blocker treatment, e.g. with timolol, as well as systemically administered inhibitors of carbonic anhydrase, e.g. acetazolamide. Cholinesterase inhibitors such as physostigmine and echothiopate may also be employed and have an effect similar to that of pilocarpine. Drugs currently used to treat glaucoma thus include, e.g.,

1. Prostaglandin analogs such as latanoprost (Xalatan), bimatoprost (Lumigan) and travoprost (Travatan), which increase uveoscleral outflow of aqueous humor. Bimatoprost also increases trabecular outflow.
2. Topical beta-adrenergic receptor antagonists such as timolol, levobunolol (Betagan), and betaxolol, which decrease aqueous humor production by the ciliary body.

3. Alpha₂-adrenergic agonists such as brimonidine (Alphagan), which work by a dual mechanism, decreasing aqueous production and increasing uveo-scleral outflow.
4. Less-selective sympathomimetics like epinephrine and dipivefrin (Propine) increase outflow of aqueous humor through trabecular meshwork and possibly through uveoscleral outflow pathway, probably by a beta₂-agonist action.
5. Miotic agents (parasympathomimetics) like pilocarpine work by contraction of the ciliary muscle, tightening the trabecular meshwork and allowing increased outflow of the aqueous humour.
6. Carbonic anhydrase inhibitors like dorzolamide (Trusopt), brinzolamide (Azopt), acetazolamide (Diamox) lower secretion of aqueous humor by inhibiting carbonic anhydrase in the ciliary body.
7. Physostigmine is also used to treat glaucoma and delayed gastric emptying.

[0069] For example, the invention provides pharmaceutical compositions comprising a PDE1 Inhibitor of the Invention and an agent selected from (i) the prostanoids, unoprostone, latanoprost, travoprost, or bimatoprost; (ii) an alpha adrenergic agonist such as brimonidine, apraclonidine, or dipivefrin and (iii) a muscarinic agonist, such as pilocarpine. For example, the invention provides ophthalmic formulations comprising a PDE-1 Inhibitor of the Invention together with bimatoprost, abrimonidine, brimonidine, timolol, or combinations thereof, in free or ophthalmologically acceptable salt form, in combination or association with an ophthalmologically acceptable diluent or carrier. In addition to selecting a combination, however, a person of ordinary skill in the art can select an appropriate selective receptor subtype agonist or antagonist. For example, for alpha adrenergic agonist, one can select an agonist selective for an alpha 1 adrenergic receptor, or an agonist selective for an alpha₂ adrenergic receptor such as brimonidine, for example. For a beta-adrenergic receptor antagonist, one can select an antagonist selective for either β_1 , or β_2 , or β_3 , depending on the appropriate therapeutic application. One can also select a muscarinic agonist selective for a particular receptor subtype such as M₁-M₅.

[0070] The PDE 1 inhibitor may be administered in the form of an ophthalmic composition, which includes an ophthalmic solution, cream or ointment. The

ophthalmic composition may additionally include an intraocular-pressure lowering agent.

[0071] In yet another example, the PDE-1 Inhibitors disclosed may be combined with a subthreshold amount of an intraocular pressure-lowering agent which may be a bimatoprost ophthalmic solution, a brimonidine tartrate ophthalmic solution, or brimonidine tartrate/timolol maleate ophthalmic solution.

[0072] In addition to the above-mentioned methods, it has also been surprisingly discovered that PDE1 inhibitors are useful to treat psychosis, for example, any conditions characterized by psychotic symptoms such as hallucinations, paranoid or bizarre delusions, or disorganized speech and thinking, e.g., schizophrenia, schizoaffective disorder, schizophreniform disorder, psychotic disorder, delusional disorder, and mania, such as in acute manic episodes and bipolar disorder. Without intending to be bound by any theory, it is believed that typical and atypical antipsychotic drugs such as clozapine primarily have their antagonistic activity at the dopamine D2 receptor. PDE1 inhibitors, however, primarily act to enhance signaling at the dopamine D1 receptor. By enhancing D1 receptor signaling, PDE1 inhibitors can increase NMDA receptor function in various brain regions, for example in nucleus accumbens neurons and in the prefrontal cortex. This enhancement of function may be seen for example in NMDA receptors containing the NR2B subunit, and may occur e.g., via activation of the Src and protein kinase A family of kinases.

[0073] Therefore, the invention provides a new method for the treatment of psychosis, e.g., schizophrenia, schizoaffective disorder, schizophreniform disorder, psychotic disorder, delusional disorder, and mania, such as in acute manic episodes and bipolar disorder, comprising administering an effective amount of a phosphodiesterase-1 (PDE1) Inhibitor of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, to a patient in need thereof.

[0074] PDE 1 Inhibitors may be used in the foregoing methods of treatment prophylaxis as a sole therapeutic agent, but may also be used in combination or for co-administration with other active agents. Thus, the invention further comprises a method of treating psychosis, e.g., schizophrenia, schizoaffective disorder, schizophreniform disorder, psychotic disorder, delusional disorder, or mania,

comprising administering simultaneously, sequentially, or contemporaneously administering therapeutically effective amounts of:

(i) a PDE 1 Inhibitor of the invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form; and

5 (ii) an antipsychotic, e.g.,

Typical antipsychotics, e.g.,

Butyrophenones, e.g. Haloperidol (Haldol, Serenace),

Droperidol (Droleptan);

Phenothiazines, e.g., Chlorpromazine (Thorazine, Largactil),

10 Fluphenazine (Prolixin), Perphenazine (Trilafon),

Prochlorperazine (Compazine), Thioridazine (Mellaril,

Melleril), Trifluoperazine (Stelazine), Mesoridazine,

Periciazine, Promazine, Triflupromazine (Vesprin),

Levomepromazine (Nozinan), Promethazine (Phenergan),

15 Pimozide (Orap);

Thioxanthenes, e.g., Chlorprothixene, Flupenthixol (Depixol,

Fluanxol), Thiothixene (Navane), Zuclopenthixol (Clopixol,

Acuphase);

Atypical antipsychotics, e.g.,

20 Clozapine (Clozaril), Olanzapine (Zyprexa), Risperidone

(Risperdal), Quetiapine (Seroquel), Ziprasidone (Geodon),

Amisulpride (Solian), Paliperidone (Invega), Aripiprazole

(Abilify), Bifeprunox; norclozapine,

in free or pharmaceutically acceptable salt form, to a patient in need thereof.

25 **[0075]** In a particular embodiment, the Compounds of the Invention are particularly useful for the treatment or prophylaxis of schizophrenia.

[0076] Compounds of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67,

in free or pharmaceutically acceptable salt form, are particularly useful for the

treatment of Parkinson's disease, schizophrenia, narcolepsy, glaucoma and female

30 sexual dysfunction.

[0077] In still another aspect, the invention provides a method of lengthening or enhancing growth of the eyelashes by administering an effective amount of a prostaglandin analogue, e.g., bimatoprost, concomitantly, simultaneously or

sequentially with an effective amount of a PDE1 inhibitor of the Invention, e.g., a e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, to the eye of a patient in need thereof.

[0078] In yet another aspect, the invention provides a method for the treatment of
5 traumatic brain injury comprising administering a therapeutically effective amount of a Compound of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, to a patient in need thereof. Traumatic brain injury (TBI) encompasses primary injury as well as secondary injury, including both focal and diffuse brain injuries. Secondary injuries are multiple, parallel,
10 interacting and interdependent cascades of biological reactions arising from discrete subcellular processes (e.g., toxicity due to reactive oxygen species, overstimulation of glutamate receptors, excessive influx of calcium and inflammatory upregulation) which are caused or exacerbated by the inflammatory response and progress after the initial (primary) injury. Abnormal calcium homeostasis is believed to be a critical
15 component of the progression of secondary injury in both grey and white matter. For a review of TBI, see Park et al., CMAJ (2008) 178(9):1163-1170, the contents of which are incorporated herein in their entirety. Studies have shown that the cAMP-PKA signaling cascade is downregulated after TBI and treatment of PDE IV inhibitors such as rolipram to raise or restore cAMP level improves histopathological
20 outcome and decreases inflammation after TBI. As Compounds of the present invention is a PDE1 inhibitor, it is believed that these compounds are also useful for the treatment of TBI, e.g., by restoring cAMP level and/or calcium homeostasis after traumatic brain injury.

[0079] The present invention also provides
25 (i) a Compound of the Invention for use as a pharmaceutical, for example for use in any method or in the treatment of any disease or condition as hereinbefore set forth,
(ii) the use of a Compound of the Invention in the manufacture of a medicament for treating any disease or condition as hereinbefore
30 set forth,
(iii) a pharmaceutical composition comprising a Compound of the Invention in combination or association with a pharmaceutically acceptable diluent or carrier, and

- (iv) a pharmaceutical composition comprising a Compound of the Invention in combination or association with a pharmaceutically acceptable diluent or carrier for use in the treatment of any disease or condition as hereinbefore set forth.

5 [0080] Therefore, the invention provides use of a Compound of the Invention, e.g., any of Formulae II, II(a)-II(f) or 2.1-2.67, in free or pharmaceutically acceptable salt form, for the manufacture of a medicament for the treatment or prophylactic treatment of the following diseases: Parkinson's disease, restless leg, tremors, dyskinesias, Huntington's disease, Alzheimer's disease, and drug-induced movement disorders;
10 depression, attention deficit disorder, attention deficit hyperactivity disorder, bipolar illness, anxiety, sleep disorder, narcolepsy, cognitive impairment, dementia, Tourette's syndrome, autism, fragile X syndrome, psychostimulant withdrawal, and/or drug addiction; cerebrovascular disease, stroke, congestive heart disease, hypertension, pulmonary hypertension, and/or sexual dysfunction; asthma, chronic
15 obstructive pulmonary disease, and/or allergic rhinitis, as well as autoimmune and inflammatory diseases; and/or female sexual dysfunction, exercise amenorrhoea, anovulation, menopause, menopausal symptoms, hypothyroidism, pre-menstrual syndrome, premature labor, infertility, irregular menstrual cycles, abnormal uterine bleeding, osteoporosis, multiple sclerosis, prostate enlargement, prostate cancer,
20 hypothyroidism, estrogen-induced endometrial hyperplasia or carcinoma; and/or any disease or condition characterized by low levels of cAMP and/or cGMP (or inhibition of cAMP and/or cGMP signaling pathways) in cells expressing PDE1, and/or by reduced dopamine D1 receptor signaling activity; and/or any disease or condition that may be ameliorated by the enhancement of progesterone signaling; comprising
25 administering an effective amount of a Compound of the Invention, or a pharmaceutical composition comprising a Compound of the Invention, to a patient in need of such treatment.

[0081] The invention also provides use of a Compound of the Invention for the manufacture of a medicament for the treatment or prophylactic treatment of:

- 30 a) glaucoma or elevated intraocular pressure,
b) psychosis, for example, any conditions characterized by psychotic symptoms such as hallucinations, paranoid or bizarre delusions, or disorganized speech and thinking, e.g., schizophrenia, schizoaffective

disorder, schizophreniform disorder, psychotic disorder, delusional disorder, and mania, such as in acute manic episodes and bipolar disorder, or

c) traumatic brain injury.

5 [0082] The invention further provides use of the Compound of the Invention for lengthening or enhancing growth of the eyelashes.

[0083] The words "treatment" and "treating" are to be understood accordingly as embracing prophylaxis and treatment or amelioration of symptoms of disease as well as treatment of the cause of the disease.

10 [0084] Compounds of the Invention are in particular useful for the treatment of Parkinson's disease, narcolepsy and female sexual dysfunction.

[0085] For methods of treatment, the word "effective amount" is intended to encompass a therapeutically effective amount to treat a specific disease or disorder.

15 [0086] The term "pulmonary hypertension" is intended to encompass pulmonary arterial hypertension.

[0087] The term "patient" include human or non-human (i.e., animal) patient. In particular embodiment, the invention encompasses both human and nonhuman. In another embodiment, the invention encompasses nonhuman. In other embodiment, the term encompasses human.

20 [0088] The term "comprising" as used in this disclosure is intended to be open-ended and does not exclude additional, unrecited elements or method steps.

[0089] Compounds of the Invention may be used as a sole therapeutic agent, but may also be used in combination or for co-administration with other active agents. For example, as Compounds of the Invention potentiate the activity of D1 agonists, such as dopamine, they may be simultaneously, sequentially, or contemporaneously administered with conventional dopaminergic medications, such as levodopa and levodopa adjuncts (carbidopa, COMT inhibitors, MAO-B inhibitors), dopamine agonists, and anticholinergics, e.g., in the treatment of a patient having Parkinson's disease. In addition, the novel PDE 1 inhibitors of the Invention, e.g., the Compounds of the Invention as described herein, may also be administered in combination with estrogen/estradiol/estriol and/or progesterone/progestins to enhance the effectiveness of hormone replacement therapy or treatment of estrogen-induced endometrial hyperplasia or carcinoma.

25
30

[0090] Dosages employed in practicing the present invention will of course vary depending, e.g. on the particular disease or condition to be treated, the particular Compound of the Invention used, the mode of administration, and the therapy desired. Compounds of the Invention may be administered by any suitable route, including orally, parenterally, transdermally, or by inhalation, but are preferably administered orally. In general, satisfactory results, e.g. for the treatment of diseases as hereinbefore set forth are indicated to be obtained on oral administration at dosages of the order from about 0.01 to 2.0 mg/kg. In larger mammals, for example humans, an indicated daily dosage for oral administration will accordingly be in the range of from about 0.75 to 150 mg, conveniently administered once, or in divided doses 2 to 4 times, daily or in sustained release form. Unit dosage forms for oral administration thus for example may comprise from about 0.2 to 75 or 150 mg, e.g. from about 0.2 or 2.0 to 50, 75 or 100 mg of a Compound of the Invention, together with a pharmaceutically acceptable diluent or carrier therefor.


[0091] Pharmaceutical compositions comprising Compounds of the Invention may be prepared using conventional diluents or excipients and techniques known in the galenic art. Thus oral dosage forms may include tablets, capsules, solutions, suspensions and the like.

EXAMPLES

[0092] The synthetic methods for various Compounds of the Present Invention are illustrated below. Other compounds of the Invention and their salts may be made using the methods as similarly described below and/or by methods similar to those generally described in the detailed description and by methods known in the chemical art.

Example 1

3-(cyclopentyloxy)-2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione



Step 1) 2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione

- 5 **[0093]** 2-(4-(6-fluoropyridin-2-yl)benzyl)-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (422 mg, 1.2 mmol), isobutyl iodide (276 μ L, 2.4 mmol) and potassium carbonate (497 mg, 3.6 mmol) are placed in a Biotage microwave vial, and then DMF (6 mL) is added. The mixture is heated in a Biotage microwave instrument at 140 °C for 30 min. The mixture is diluted with water (150 mL), and then extracted with methylene chloride three times. The combined organic phase is evaporated to dryness to give 460 mg of crude product as pale yellow solids, which is used in the next step without further purification. MS (ESI) m/z 408.2 $[M+H]^+$.

Step2) 3-chloro-2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione


- 15 **[0094]** 2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (450 mg, 1.1 mmol) and hexachloroethane (523 mg, 2.2 mmol) are dissolved in 5 mL of methylene chloride, and then 1.0 M LiHMDS (2.2 mL, 2.2 mmol) in THF is added dropwise. After the reaction mixture is stirred at room temperature for 10 min, water is added, and then extracted with methylene chloride three times. The combined organic phase is evaporated to dryness to give 520 mg of crude product as off-white solids, which is used in the next step without further purification. MS (ESI) m/z 442.2 $[M+H]^+$.

Step 3) 3-(cyclopentyloxy)-2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione

[0095] 3-chloro-2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (20 mg, 0.045 mmol), cesium carbonate (22 mg, 0.068 mmol) and cyclopentanol (0.3 mL) are placed in a Biotage microwave vial. The vial is sealed, and then heated in a Biotage microwave instrument at 150 °C for 4h. The reaction mixture is purified with a semi-preparative HPLC to give 10.4 mg of pure product as pale yellow solids (Purity: 97.7%). MS (ESI) m/z 492.2 [M+H]⁺.

Example 2


2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-3-phenoxy-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione



[0096] 3-chloro-2-(4-(6-fluoropyridin-2-yl)benzyl)-7-isobutyl-5-methyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (40 mg, 0.095 mmol), potassium carbonate (40 mg, 0.28 mmol) and phenol (26 mg, 0.28 mmol) are placed in a Biotage microwave vial, and then 0.4 mL of dioxane is added. The vial is sealed, and then heated in a Biotage microwave instrument at 150 °C for 3h. The reaction mixture is diluted with DMF, and then filtered. The filtrate is purified with a semi-preparative HPLC to give 32 mg of pure product as pale yellow crystals (Purity: 99.9%). MS (ESI) m/z 500.2 [M+H]⁺.

Example 3

5-methyl-7-neopentyl-3-phenoxy-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione



Step 1) 5-methyl-7-neopentyl-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione

- 5 **[0097]** 5-methyl-7-neopentyl-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (236 mg, 1.0 mmol), 4-(4-(chloromethyl)phenyl)pyridine (305 mg, 1.5 mmol) and potassium carbonate (414 mg, 3.0 mmol) are suspended in 3 mL of DMF. The reaction mixture is stirred at room temperature overnight, and then diluted with 100 mL of water. The mixture is extracted with methylene chloride four times (40 mL ×
- 10 4). The combined organic phase is dried over anhydrous sodium sulfate. After filtration, the filtrate is evaporated to dryness to give 510 mg of crude product, which is used in the next step without further purification. MS (ESI) m/z 404.2 $[M+H]^+$.

Step 2) 3-chloro-5-methyl-7-neopentyl-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione


- 15 **[0098]** 5-methyl-7-neopentyl-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (500 mg, 1.24 mmol) and hexachloroethane (587 mg, 2.48 mmol) are dissolved in 8 mL of methylene chloride, and then 1.0 M LiHMDS (2.48 mL, 2.48 mmol) in THF is added dropwise. After the reaction mixture is stirred
- 20 at room temperature for 10 min, 100 mL of deionized water is added, and then extracted with methylene chloride four times (30 mL × 4). The combined organic phase is dried over anhydrous sodium sulfate. After filtration, the filtrate is evaporated to dryness to give 500 mg of crude product as brown solids, which is used in the next step without further purification. MS (ESI) m/z 438.2 $[M+H]^+$.

Step 3) 5-methyl-7-neopentyl-3-phenoxy-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione

[0099] 3-chloro-5-methyl-7-neopentyl-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione (30 mg, 0.069 mmol), potassium carbonate (29 mg, 0.2 mmol) and phenol (20 mg, 0.20 mmol) are placed in a Biotage microwave vial, and then 0.3 mL of dioxane is added. The vial is sealed, and then heated in a Biotage microwave instrument at 150 °C for 3h. The reaction mixture is diluted with DMF, and then filtered. The filtrate is purified with a semi-preparative HPLC to give 23 mg of pure product as pale yellow solids. MS (ESI) m/z 496.2 [M+H]⁺.

Example 4

10 **5-methyl-7-neopentyl-3-propoxy-2-(4-(pyridin-4-yl)benzyl)-2H-pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione**



15 [00100] The synthesis method is analogous to **Example 3** wherein n-propanol is added in step 3 instead of phenol. MS (ESI) m/z 462.3 [M+H]⁺.

Example 5

Measurement of PDE1B inhibition *in vitro* using IMAP Phosphodiesterase Assay

20 **Kit**

[00101] Phosphodiesterase 1B (PDE1B) is a calcium/calmodulin dependent phosphodiesterase enzyme that converts cyclic guanosine monophosphate (cGMP) to 5'-guanosine monophosphate (5'-GMP). PDE1B can also convert a modified cGMP substrate, such as the fluorescent molecule cGMP-fluorescein, to the corresponding GMP-fluorescein. The generation of GMP-fluorescein from cGMP-fluorescein can be quantitated, using, for example, the IMAP (Molecular Devices, Sunnyvale, CA) immobilized-metal affinity particle reagent.

[00102] Briefly, the IMAP reagent binds with high affinity to the free 5'-phosphate that is found in GMP-fluorescein and not in cGMP-fluorescein. The resulting GMP-fluorescein – IMAP complex is large relative to cGMP-fluorescein. Small fluorophores that are bound up in a large, slowly tumbling, complex can be distinguished from unbound fluorophores, because the photons emitted as they fluoresce retain the same polarity as the photons used to excite the fluorescence.

[00103] In the phosphodiesterase assay, cGMP-fluorescein, which cannot be bound to IMAP, and therefore retains little fluorescence polarization, is converted to GMP-fluorescein, which, when bound to IMAP, yields a large increase in fluorescence polarization (Δmp). Inhibition of phosphodiesterase, therefore, is detected as a decrease in Δmp .

[00104] Enzyme assay

Materials: All chemicals are available from Sigma-Aldrich (St. Louis, MO) except for IMAP reagents (reaction buffer, binding buffer, FL-GMP and IMAP beads), which are available from Molecular Devices (Sunnyvale, CA).

Assay: 3',5'-cyclic-nucleotide-specific bovine brain phosphodiesterase (Sigma, St. Louis, MO) is reconstituted with 50% glycerol to 2.5 U/ml. One unit of enzyme will hydrolyze 1.0 μ mole of 3',5'-cAMP to 5'-AMP per min at pH 7.5 at 30°C. One part enzyme is added to 1999 parts reaction buffer (30 μ M CaCl₂, 10 U/ml of calmodulin (Sigma P2277), 10mM Tris-HCl pH 7.2, 10mM MgCl₂, 0.1% BSA, 0.05% NaN₃) to yield a final concentration of 1.25mU/ml. 99 μ l of diluted enzyme solution is added into each well in a flat bottom 96-well polystyrene plate to which 1 μ l of test compound dissolved in 100% DMSO is added. The compounds are mixed and pre-incubated with the enzyme for 10 min at room temperature.

[00105] The FL-GMP conversion reaction is initiated by combining 4 parts enzyme and inhibitor mix with 1 part substrate solution (0.225 μ M) in a 384-well microtiter plate. The reaction is incubated in dark at room temperature for 15 min. The reaction is halted by addition of 60 μ l of binding reagent (1:400 dilution of IMAP beads in binding buffer supplemented with 1:1800 dilution of antifoam) to each well of the 384-well plate. The plate is incubated at room temperature for 1 hour to allow IMAP binding to proceed to completion, and then placed in an Envision multimode microplate reader (PerkinElmer, Shelton, CT) to measure the fluorescence polarization (Δmp).

[00106] A decrease in GMP concentration, measured as decreased Δ mp, is indicative of inhibition of PDE activity. IC_{50} values are determined by measuring enzyme activity in the presence of 8 to 16 concentrations of compound ranging from 0.0037 nM to 80,000 nM and then plotting drug concentration versus Δ mp, which
5 allows IC_{50} values to be estimated using nonlinear regression software (XLFit; IDBS, Cambridge, MA).

[00107] The Compounds of the Invention may be selected and tested in this assay to show PDE1 inhibitory activity. Exemplified compounds are shown to have IC_{50} activities of less than 100 μ M, in some instances less than 100nM, e.g., in some
10 instances, less than 25nM, in other instances, less than 10nM.

Example 6

PDE1 inhibitor effect on sexual response in female rats


[00108] The effect of PDE1 inhibitors on Lordosis Response in female rats
15 may be measured as described in Mani, et al., Science (2000) 287: 1053.
Ovariectomized and cannulated wild-type rats are primed with 2 μ g estrogen followed 24 hours later by intracerebroventricular (icv) injection of progesterone (2 μ g), PDE1 inhibitors of the present invention (0.1mg, 1.0mg or 2.5mg) or sesame oil vehicle (control). The rats are tested for lordosis response in the presence of male rats.
20 Lordosis response is quantified by the lordosis quotient (LQ = number of lordosis/10 mounts x 100). The LQ for estrogen-primed female rats receiving Compounds of the Invention, at 0.1 mg, are expected to be similar to estrogen-primed rats receiving progesterone and higher than for estrogen-primed rats receiving vehicle.

25

CLAIMS

What is claimed is:

1. A compound of Formula II:



Formula II

wherein

(i) Q is $-C(=O)-$, $-C(=S)-$, $-C(=N(R_6))-$ or $-C(R_{14})(R_{15})-$;

(ii) R_1 is H or C_{1-6} alkyl (e.g., methyl or ethyl);

(iii) R_2 is

H,

C_{1-6} alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-

dimethylpropyl) wherein said alkyl group is optionally

substituted with one or more halo (e.g., fluoro) or hydroxy

(e.g., hydroxy C_{1-6} alkyl, for example 1-hydroxyprop-2-yl or

3-hydroxy-2-methylpropyl),

halo C_{1-6} alkyl (e.g., trifluoromethyl or 2,2,2-trifluoroethyl),

$N(R_{14})(R_{15})-C_{1-6}$ alkyl (e.g., 2-(dimethylamino)ethyl or 2-

aminopropyl),

aryl C_{0-6} alkyl (e.g., phenyl or benzyl), wherein said aryl is

optionally substituted with one or more C_{1-6} alkoxy, for

example, C_{1-6} alkoxyaryl C_{0-6} alkyl (e.g., 4-methoxybenzyl),

heteroaryl C_{0-6} alkyl (e.g., pyridinylmethyl), wherein said

heteroaryl is optionally substituted with one or more C_{1-6}

alkoxy (e.g., C_{1-6} alkoxyheteroaryl C_{1-6} alkyl);

$-G-J$ wherein G is a single bond or C_{1-6} alkylene (e.g.,

methylene) and J is C_{3-8} cycloalkyl or hetero C_{3-8} cycloalkyl

(e.g., oxetan-2-yl, pyrrolidin-3-yl, pyrrolidin-2-yl) wherein the cycloalkyl and heterocycloalkyl group are optionally substituted with one or more C₁₋₆alkyl or amino, for example,

- 5 –C₀₋₄alkyl-C₃₋₈cycloalkyl (e.g., –C₀₋₄alkyl-cyclopentyl, –C₀₋₄alkyl-cyclohexyl or –C₀₋₄alkyl-cyclopropyl), wherein said cycloalkyl is optionally substituted with one or more C₁₋₆alkyl or amino (for example, 2-aminocyclopentyl or 2-aminocyclohexyl),
- 10 –C₀₋₄alkyl-C₃₋₈heterocycloalkyl (e.g., –C₀₋₄alkyl-pyrrolidinyl, for example, –C₀₋₄alkylpyrrolidin-3-yl) wherein said heterocycloalkyl is optionally substituted with C₁₋₆alkyl (e.g., methyl), for example, 1-methylpyrrolidin-3-yl, 1-methyl-
- 15 pyrrolindin-2-yl, 1-methyl-pyrrolindin-2-yl-methyl or 1-methyl-pyrrolindin-3-yl-methyl);

(iv) R₃ is


4) –D-E-F wherein:

- 20 D is a single bond, C₁₋₆alkylene (e.g., methylene), or arylC₁₋₆alkylene (e.g., benzylene or –CH₂C₆H₄–);
- E is
- a single bond,
- C₁₋₄alkylene (e.g., methylene, ethynylene, prop-2-yn-1-ylene),
- 25 C₀₋₄alkylarylene (e.g., phenylene or –C₆H₄–, –benzylene– or –CH₂C₆H₄–), wherein the arylene group is optionally substituted with halo (e.g., Cl or F),
- heteroarylene (e.g., pyridinylene or pyrimidinylene),
- aminoC₁₋₆alkylene (e.g., –CH₂N(H)–),
- 30 amino (e.g., –N(H)–);
- C₃₋₈cycloalkylene optionally containing one or more heteroatom selected from N or O (e.g., piperidinylene),
- F is

- H,
 halo (e.g., F, Br, Cl),
 C₁₋₆alkyl (e.g., isopropyl or isobutyl),
 haloC₁₋₆alkyl (e.g., trifluoromethyl),
 5 aryl (e.g., phenyl),
 C₃₋₈cycloalkyl optionally containing one or more atom
 selected from a group consisting of N, S or O (e.g.,
 cyclopentyl, cyclohexyl, piperidiny, pyrrolidiny,
 tetrahydro-2*H*-pyran-4-yl, or morpholiny), and
 10 optionally substituted with one or more C₁₋₆alkyl (e.g.,
 methyl or isopropyl), for example, 1-methylpyrrolidin-
 2-yl, pyrrolidin-1-yl, pyrrolidin-2-yl, piperidin-2-yl, 1-
 methylpiperidin-2-yl, 1-ethylpiperidin-2-yl,
 heteroaryl optionally substituted with one or more C₁₋₆alkyl,
 15 halo (e.g., fluoro) or haloC₁₋₆alkyl (e.g., pyridyl (for
 example, pyrid-2-yl), pyrimidinyl (for example,
 pyrimidin-2-yl), thiadiazolyl (for example, 1,2,3-
 thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example,
 pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl,
 20 4-methylimidazolyl, 1-methylimidazol-2-yl)), triazolyl
 (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-yl),
 C₁₋₆alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazolyl);
 C₁₋₆alkoxy,
 -O-haloC₁₋₆alkyl (e.g., -O-CF₃),
 25 C₁₋₆alkylsulfonyl (for example, methylsulfonyl or -
 S(O)₂CH₃),
 -C(O)-R₁₃, wherein R₁₃ is -N(R₁₄)(R₁₅), C₁₋₆alkyl (e.g.,
 methyl), -OC₁₋₆alkyl (e.g., -OCH₃), haloC₁₋₆alkyl
 (trifluoromethyl), aryl (e.g., phenyl), or heteroaryl;
 30 -N(R₁₄)(R₁₅);
 or
 5) a substituted heteroarylC₁₋₆alkyl, e.g., substituted with haloC₁₋₆alkyl;

or

- 6) attached to one of the nitrogens on the pyrazolo portion of Formula II and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen (e.g., Cl or F); and

R₁₀ is

halogen (e.g., fluoro or chloro),

C₁₋₆alkyl,

C₃₋₈cycloalkyl,

heteroC₃₋₈cycloalkyl (e.g., pyrrolidinyl or piperidinyl),

haloC₁₋₆alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for

example, pyrid-2-yl) or e.g., thiadiazolyl (for

example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g.,

pyrazolyl (for example, pyrazol-1-yl) or

imidazolyl (for example, imidazol-1-yl)),

triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g.,

tetrazol-5-yl), C₁₋₆alkyloxadiazolyl (e.g., 5-

methyl-1,2,4-oxadiazol), pyrazolyl (e.g.,

pyrazol-1-yl),

wherein said aryl, heteroaryl, cycloalkyl or

heterocycloalkyl is optionally substituted with

one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH, or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl),
 C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl), arylcarbonyl (e.g., benzoyl), heteroarylcarbonyl, C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl), Aminocarbonyl,
 -N(R₁₄)(R₁₅); preferably R₁₀ is phenyl, pyridyl, piperidiny or pyrrolidiny optionally substituted with the substituents previously defined, e.g. optionally substituted with halo or alkyl;
 provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀, respectively, is not present;

(v) R₄ is:

C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl),
 C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
 C₃₋₈heterocycloalkyl (e.g., pyrrolidiny (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidiny (for example, piperidin-1-yl), morpholiny),
 -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or
 -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl)
 wherein said aryl or heteroaryl is optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);


(vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form, provided that said compound is not:

- (a) 7-[(2,4-dichlorophenyl)methyl]-3-ethoxy-2,5-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (b) 7-[(4-chlorophenyl)methyl]-3-ethoxy-2,5-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- 5 (c) 2-[(2,5-dichlorophenyl)methyl]-3-ethoxy-5,7-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (d) 2-[(4-bromophenyl)methyl]-3-ethoxy-5,7-dimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- (e) 3-[(4-methoxyphenyl)methoxy]-2,5,7-trimethyl-2H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione;
- 10 (f) 1-[(3-bromophenyl)methyl]-5,7-dimethyl-3-(2-propen-1-yloxy)-1H-Pyrazolo[3,4-d]pyrimidine-4,6(5H,7H)-dione.
2. The compound according to claim 1, which compound is selected from any one of the following:
- 15

A) a Compound of Formula II(a):




Formula II(a)

20 wherein

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally
- 25 substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),

- (iv) R_3 is attached to one of the nitrogens on the pyrazolo portion of Formula II(a) and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R_8 , R_9 , R_{11} and R_{12} are independently H or halogen (e.g., Cl or F); and

R_{10} is

halogen (e.g., fluoro or chloro),

C_{1-6} alkyl,

C_{3-8} cycloalkyl,

hetero C_{3-8} cycloalkyl (e.g., pyrrolidinyl or piperidinyl),


halo C_{1-6} alkyl (e.g., trifluoromethyl),

aryl (e.g., phenyl) or heteroaryl (e.g., pyridyl, (for example, pyrid-2-yl) or e.g., thiadiazolyl (for example, 1,2,3-thiadiazol-4-yl), diazolyl (e.g., pyrazolyl (for example, pyrazol-1-yl) or imidazolyl (for example, imidazol-1-yl)), triazolyl (e.g., 1,2,4-triazol-1-yl), tetrazolyl (e.g., tetrazol-5-yl), C_{1-6} alkyloxadiazolyl (e.g., 5-methyl-1,2,4-oxadiazol), pyrazolyl (e.g., pyrazol-1-yl),

wherein said aryl, heteroaryl, cycloalkyl or heterocycloalkyl is optionally substituted with one or more C_{1-6} alkyl (e.g., methyl), halogen

- (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl)
- 5 C₁₋₆alkyl sulfonyl (e.g., methyl sulfonyl),
 arylcarbonyl (e.g., benzoyl),
 heteroarylcarbonyl,
 C₁₋₆alkoxycarbonyl, (e.g., methoxycarbonyl),
 Aminocarbonyl,
- 10 -N(R₁₄)(R₁₅);
 preferably R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl,
 optionally substituted with one or more C₁₋₆alkyl
 (e.g., methyl), halogen (e.g., chloro or fluoro),
 haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy,
 15 carboxy, -SH or an additional aryl or heteroaryl
 (e.g., biphenyl or pyridylphenyl);
 provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
 respectively, is not present;
- (v) R₄ is:
- 20 C₁₋₆alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl),
 C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
 C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example
 pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for
 example, piperidin-1-yl), morpholinyl),
 25 -C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆
 alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-
 3-yl) wherein said aryl or heteroaryl is optionally
 substituted with one or more halo (e.g., 4-fluorophenyl),
 hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy
 30 or another aryl group (e.g., biphenyl-4-ylmethyl);
- (vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl) or C₃₋₈cycloalkyl;
- (vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,
- in free or salt form;


B) a Compound of formula II(b):



Formula II(b)

5 wherein:

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyc₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(b) and is a moiety of Formula A



Formula A

wherein:


X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen (e.g., Cl or F); and

5 R_{10} is phenyl or pyridyl, e.g., 2-pyridyl, or pyrrolidinyl optionally substituted with one or more C_{1-6} alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), halo C_{1-6} alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl); provided that when X, Y or Z is nitrogen, R_8 , R_9 or R_{10} , respectively, is not present;

- (v) R_4 is:
- 10 C_{1-6} alkyl (e.g., methyl, isopropyl, isobutyl, n-propyl), C_{3-8} cycloalkyl (e.g., cyclopentyl or cyclohexyl), C_{3-8} heterocycloalkyl (e.g., pyrrolidinyl (for example pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for example, piperidin-1-yl), morpholinyl),
- 15 $-C_{0-6}$ alkylaryl (e.g., phenyl or benzyl) or $-C_{0-6}$ alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-3-yl) wherein said aryl or heteroaryl is optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C_{1-6} alkyl, C_{1-6} alkoxy
- 20 or another aryl group (e.g., biphenyl-4-ylmethyl);
- (vi) R_6 is H, C_{1-6} alkyl (e.g., methyl or ethyl) or C_{3-8} cycloalkyl;
- (vii) R_{14} and R_{15} are independently H or C_{1-6} alkyl,
- in free or salt form;


25 C) a Compound of Formula II(c):



Formula II(c)

wherein:

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyc₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(c) and is a moiety of Formula A



Formula A

wherein:

X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen (e.g., Cl or F); and

R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl, pyrrolidinyl optionally substituted with one or more C₁₋₆alkyl (e.g., methyl), halogen (e.g., chloro or fluoro), haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy, carboxy, -SH or an additional aryl or heteroaryl (e.g., biphenyl or pyridylphenyl);

provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀, respectively, is not present;


- (v) R₄ is:
-C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-

3-yl) wherein said aryl or heteroaryl is optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);

- 5 (vi) R₆ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
 (vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form;

D) a Compound of Formula II(d):




10

Formula II(d)

wherein:

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
 (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
 15 (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyC₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
 20 (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(d) and is a moiety of Formula A



Formula A

wherein:

5

X, Y and Z are, independently, N or C,

R₈, R₉, R₁₁ and R₁₂ are independently H or halogen
(e.g., Cl or F); and

10

R₁₀ is phenyl or pyridyl, e.g., 2-pyridyl, pyrrolidinyl
optionally substituted with one or more C₁₋₆alkyl
(e.g., methyl), halogen (e.g., chloro or fluoro),
haloC₁₋₆alkyl (e.g., trifluoromethyl), hydroxy,
carboxy, -SH or an additional aryl or heteroaryl
(e.g., biphenyl or pyridylphenyl);

15

provided that when X, Y or Z is nitrogen, R₈, R₉ or R₁₀,
respectively, is not present;

(v) R₄ is :

20

C₃₋₈cycloalkyl (e.g., cyclopentyl or cyclohexyl),
C₃₋₈heterocycloalkyl (e.g., pyrrolidinyl (for example
pyrrolidin-3-yl or pyrrolidin-1-yl), piperidinyl (for
example, piperidin-1-yl), morpholinyl),
-C₀₋₆alkylaryl (e.g., phenyl or benzyl) or -C₀₋₆
alkylheteroaryl (e.g., pyrid-4-yl, pyrid-2-yl or pyrazol-
3-yl) wherein said aryl or heteroaryl is optionally
substituted with one or more halo (e.g., 4-fluorophenyl),
hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy
or another aryl group (e.g., biphenyl-4-ylmethyl);


25

(vi) R₆ is H, C₁₋₆alkyl (e.g., methyl or ethyl);

(vii) R₁₄ and R₁₅ are independently H or C₁₋₆alkyl,

in free or salt form.


E) a Compound of Formula II(e):



Formula II(e)

wherein:

- (i) Q is -C(=O)-, -C(=S)-, -C(=N(R₆))- or -C(R₁₄)(R₁₅)-;
- (ii) R₁ is H or C₁₋₆alkyl (e.g., methyl or ethyl);
- (iii) R₂ is C₁₋₆alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-dimethylpropyl) wherein said alkyl group is optionally substituted with one or more halo (e.g., fluoro) or hydroxy (e.g., hydroxyc₁₋₆alkyl, for example 1-hydroxyprop-2-yl or 3-hydroxy-2-methylpropyl),
- (iv) R₃ is attached to one of the nitrogens on the pyrazolo portion of Formula II(e) and is a moiety of Formula A




Formula A

wherein:

- X, Y and Z are, independently, N or C,
- R₈, R₉, R₁₁ and R₁₂ are independently H or halogen (e.g., Cl or F); and

- 5 R_{10} is phenyl, pyridyl (e.g., 2-pyridyl), pyrrolidinyl
 (e.g., pyrrolidin-3-yl or pyrrolidin-1-yl) or
 piperidinyl (e.g., piperidin-1-yl), optionally
 substituted with one or more C_{1-6} alkyl (e.g., methyl)
 or halogen (e.g., chloro or fluoro);
 provided that when X, Y or Z is nitrogen, R_8 , R_9 or R_{10} ,
 respectively, is not present;
 (v) R_4 is phenyl optionally substituted with one or more halo (e.g.,
 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C_{1-6} alkyl,
 10 C_{1-6} alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl);
 (vi) R_6 is H or C_{1-6} alkyl (e.g., methyl or ethyl);
 (vii) R_{14} and R_{15} are independently H or C_{1-6} alkyl,
 in free or salt form;


15 F) a Compound of Formula II(f):



Formula II(f)

wherein:

- 20 (i) Q is $-C(=O)-$;
 (ii) R_1 is H or C_{1-6} alkyl (e.g., methyl or ethyl);
 (iii) R_2 is C_{1-6} alkyl (e.g., isopropyl, isobutyl, 2-methylbutyl or 2,2-
 dimethylpropyl);
 (iv) R_3 is attached to one of the nitrogens on the pyrazolo portion of
 Formula II(f) and is a moiety of Formula A



Formula A

wherein:

5

X, Y and Z are C;

R₈, R₉, R₁₁ and R₁₂ are H; and

R₁₀ is pyridyl (e.g., 2-pyridyl), optionally substituted with one or more halogen (e.g., fluoro);

(v) R₄ is

10

phenyl optionally substituted with one or more halo (e.g., 4-fluorophenyl), hydroxy (e.g., 4-hydroxyphenyl), C₁₋₆alkyl, C₁₋₆alkoxy or another aryl group (e.g., biphenyl-4-ylmethyl), preferably phenyl;

C₁₋₆alkyl (e.g., n-propyl),

15

C₃₋₈cycloalkyl (e.g., cyclopentyl),

in free or salt form.

3. The compound according to any of claims 1-2, wherein Q is -C(=O) in free or salt form.


20

4. The compound according to any of claims 1-3, wherein R₂ is C₁₋₆alkyl in free or salt form.

5. The compound according to any of claims 1-4 wherein R₄ is phenyl or cyclopentyl in free or salt form.

25

6. The compound according to any of claims 1-5 wherein said compound is selected from any of the following:



in free or salt form.

- 5 7. A pharmaceutical composition comprising a compound according to any of claims 1-6, in free or a pharmaceutically acceptable salt form, in admixture with a pharmaceutically acceptable diluent or carrier.
8. A method for the treatment or prophylaxis of any of the following conditions:
- 10 Parkinson's disease, restless leg, tremors, dyskinesias, Huntington's disease, Alzheimer's disease, and drug-induced movement disorders; depression, attention deficit disorder, attention deficit hyperactivity disorder, bipolar illness, anxiety, sleep disorder, narcolepsy, cognitive impairment, dementia, Tourette's syndrome, autism, fragile X syndrome, psychostimulant withdrawal, and/or drug addiction;
- 15 cerebrovascular disease, stroke, congestive heart disease, hypertension, pulmonary hypertension, and/or sexual dysfunction; asthma, chronic obstructive pulmonary disease, and/or allergic rhinitis, as well as autoimmune and inflammatory diseases; and/or female sexual dysfunction, exercise amenorrhoea, anovulation, menopause, menopausal symptoms, hypothyroidism, pre-menstrual syndrome, premature
- 20 labor, infertility, irregular menstrual cycles, abnormal uterine bleeding, osteoporosis, multiple sclerosis, prostate enlargement, prostate cancer,

- hypothyroidism, estrogen-induced endometrial hyperplasia or carcinoma; and/or any disease or condition characterized by low levels of cAMP and/or cGMP (or inhibition of cAMP and/or cGMP signaling pathways) in cells expressing PDE1, and/or by reduced dopamine D1 receptor signaling activity; and/or any disease or condition that may be ameliorated by the enhancement of progesterone signaling; comprising administering a therapeutically effective amount of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, or a pharmaceutical composition according to claim 7, to a patient in need of such treatment.
9. The method of claim 8, wherein the condition is Parkinson's disease.
10. The method of claim 8, wherein the condition is cognitive impairment.
11. The method of claim 8, wherein the condition is narcolepsy.
12. The method of claim 11 further comprising administering a compound or compounds selected from central nervous system stimulants, modafinil, antidepressants, and gamma hydroxybutyrate, to a patient in need thereof.
13. The method of claim 8, wherein said condition is female sexual dysfunction.
14. The method of claim 13, further comprising administering a compound or compounds selected from a group consisting of estradiol, estriol, estradiol esters, progesterone and progestins to a patient in need thereof.
15. A method for the treatment for glaucoma or elevated intraocular pressure comprising topical administration of a therapeutically effective amount of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, to a patient in need of such treatment.

16. A pharmaceutical composition for topical ophthalmic use comprising a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form.
- 5 17. A method for the treatment or prophylaxis of psychosis including schizophrenia, schizoaffective disorder, schizophreniform disorder, psychotic disorder, delusional disorder, and mania, such as in acute manic episodes and bipolar disorder, comprising administering an effective amount of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free
10 or pharmaceutically acceptable salt form.
18. Use of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, for the manufacture of a medicament for the treatment or prophylactic treatment of a
15 condition set forth in any one of claims 8-14.
19. Use of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, for the manufacture of a medicament for the treatment or prophylactic treatment of
20 glaucoma or elevated intraocular pressure.
20. Use of a compound according to any of claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, for the manufacture of a medicament for the treatment or prophylactic treatment of
25 psychosis, e.g., schizophrenia, schizoaffective disorder, schizophreniform disorder, psychotic disorder, delusional disorder, and mania, such as in acute manic episodes and bipolar disorder.
21. A pharmaceutical composition comprising a Compound according to any of
30 claims 1-6, provided that proviso (a) through (f) does not apply, in free or pharmaceutically acceptable salt form, in combination or association with a pharmaceutically acceptable diluent or carrier for use in the treatment of any disease or condition according to any of claims 8-14.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 11/38544

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 31/519 (2011.01)

USPC - 514/258.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

USPC: 514/258.1

IPC(8): A61K 31/519 (2011.01)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

USPC: 514/49 514/50; 514/186; 514/259.5; 514/262.1; 514/264.11; 514/269; 514/274; 514/205; 544/256; 544/262

(see search terms below)

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST: PGPB, USPT, USOC, EPAB, JPAB

Google: Scholar/patents: pde1\$ inhibitors bicyclic pyrazolo pyrimidine diones pyrrolo pyrrole alkoxy phosphodiesterase crystal structure docking

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO2007/143705 A1 (WENNOGLE et al) 13 December 2007 (13.12.2007) para [0008]	1-3
Y	US 2005/0075795 A1 (PANDIT) 07 April 2005 (07.04.2005) para [0002];[0008];[0009];[0011]-[0013];[0106];[0107];[0116]	1-3

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

02 September 2011 (02.09.2011)

Date of mailing of the international search report

16 SEP 2011

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 11/38544

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4-21
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.