

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 December 2003 (18.12.2003)

PCT

(10) International Publication Number  
**WO 03/104397 A2**

- (51) International Patent Classification<sup>7</sup>: C12N
- (21) International Application Number: PCT/US03/17174
- (22) International Filing Date: 29 May 2003 (29.05.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
10/159,856 31 May 2002 (31.05.2002) US
- (71) Applicant (for all designated States except US): **ISIS PHARMACEUTICALS INC.** [US/US]; 2292 Faraday Avenue, Carlsbas, CA 92008 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **FREIER, Susan, M.** [US/US]; 2946 Renault Street, San Diego, CA 92122 (US).  
**DOBIE, Kenneth, W.** [US/US]; 703 Stratford Ct., #4, Del Mar, CA 92014 (US).
- (74) Agents: **SHUSTER, Michael, J.** et al.; Fenwick & West LLP, Silicon Valley Center, 801 California Street, Mountain View, CA 94041 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



**WO 03/104397 A2**

(54) Title: ANTISENSE MODULATION OF G PROTEIN-COUPLED RECEPTOR KINASE 6 EXPRESSION

(57) Abstract: Antisense compounds, compositions and methods are provided for modulating the expression of G protein-coupled receptor kinase 6. The compositions comprise antisense compounds, particularly antisense oligonucleotides, targeted to nucleic acids encoding G protein-coupled receptor kinase 6. Methods of using these compounds for modulation of G protein-coupled receptor kinase 6 expression and for treatment of diseases associated with expression of G protein-coupled receptor kinase 6 are provided.

**ANTISENSE MODULATION OF G PROTEIN-COUPLED RECEPTOR KINASE 6  
EXPRESSION**

**FIELD OF THE INVENTION**

The present invention provides compositions and methods for modulating the expression of G protein-coupled receptor kinase 6. In particular, this invention relates to compounds, particularly oligonucleotides, specifically hybridizable with nucleic acids encoding G protein-coupled receptor kinase 6. Such compounds have been shown to modulate the expression of G protein-coupled receptor kinase 6.

**BACKGROUND OF THE INVENTION**

G protein-coupled receptors (GPCRs) are 7-transmembrane domain-containing proteins activated by a wide variety of agonists (hormones, neurotransmitters, opioids, odorants, light) and these receptors serve to transduce the ligand-binding or activating event into an intracellular signal. Upon agonist stimulation, GPCRs catalyze nucleotide exchange (GDP to GTP) and thereby activate intracellular G-proteins, and the GPCR must be turned off to prevent continuing G-protein activation. G protein-coupled receptor kinases (GRKs) are primarily responsible for the rapid loss of GPCR responsiveness in spite of continuous presence of the agonist, in a process known as homologous desensitization. Phosphorylation of activated GPCRs by GRKs and by other second messenger-activated receptor-specific protein kinases, such as

protein kinase C (PKC), leads to a shortened physiologic response. Activated receptor phosphorylation promotes binding of regulatory arrestin proteins, which prevent further receptor-coupling to G-proteins, thereby impeding additional rounds of receptor-activated signal transduction (Palczewski, 5 *Eur. J. Biochem.*, 1997, 248, 261-269).

To date, there are six members of the functionally related serine/threonine G protein-coupled receptor kinase family, and this family is subdivided into two groups: the 10 first comprises the closely related GRK2 and GRK3, and the second includes GRK1, GRK4, GRK5 and GRK6. The amino terminal domains of all GRKs are highly similar in sequence and is believed to be involved in the interaction with activated GPCRs, and, in GRK4, GRK5 and GRK6, may also be involved in 15 binding phosphatidylinositol 4,5 bisphosphate (Palczewski, *Eur. J. Biochem.*, 1997, 248, 261-269).

In a screen of a human heart cDNA library using DNA sequences encoding the catalytic domains of the bovine  $\beta_2$ -adrenergic receptor kinase ( $\beta$ ARK) and  $\beta$ ARK2, a human G 20 protein-coupled receptor kinase 6 cDNA clone was isolated. Two major RNA species of approximately 2.4- and 3-kilobases were observed by Northern blot analysis. The larger transcript appeared to be expressed at comparable levels in brain, skeletal muscle and pancreas, with lower levels found in lung, 25 placenta, heart and kidney. The smaller transcript was found predominantly in skeletal muscle with lower levels in brain and pancreas (Benovic and Gomez, *J. Biol. Chem.*, 1993, 268, 19521-19527). Independently, a degenerate PCR approach based on amino acid sequences conserved in known receptor kinases 30 was used to amplify and a DNA fragment and subsequently isolate a full-length cDNA clone of human G protein-coupled receptor kinase 6 (also known as GPRK6 and GRK6) from a

neutrophil cDNA library. In Northern blot analyses, two transcripts of approximately 2.1- and 2.9-kilobases were detected in all tissues tested, with strongest expression in placenta and skeletal muscle (Haribabu and Snyderman, *Proc. Natl. Acad. Sci. U. S. A.*, 1993, 90, 9398-9402). By rat-human cell hybrid analysis, the G protein-coupled receptor kinase 6 gene was mapped to chromosomal region 5q35 near a cluster of growth factor and neurotransmitter receptors (Bullrich et al., *Cytogenet. Cell Genet.*, 1995, 70, 250-254). A pseudogene related to G protein-coupled receptor kinase 6 was also identified and mapped to a distinct location on chromosome 13 (Bullrich et al., *Cytogenet. Cell Genet.*, 1995, 70, 250-254; Gagnon and Benovic, *Gene*, 1997, 184, 13-19; Haribabu and Snyderman, *Proc. Natl. Acad. Sci. U. S. A.*, 1993, 90, 9398-9402).

The rat and mouse homologues of G protein-coupled receptor kinase 6 have also been cloned (Fehr et al., *Brain Res. Brain Res. Protoc.*, 1999, 3, 242-251; Neill et al., *Endocrinology*, 1996, 137, 3942-3947) and four distinct mouse isoforms, mGRK6-A through mGRK6-D, generated by alternative splicing, have been identified (Moepps et al., *Genomics*, 1999, 60, 199-209). Splice variant mGRK6-A but not mGRK6-B or mGRK6-C, was demonstrated to associate with and phosphorylate the Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor (NHERF) involved in regulation of cellular pH and sodium concentration (Hall et al., *J. Biol. Chem.*, 1999, 274, 24328-24334).

Plasma membrane association is an important feature of GRKs. Palmitoylation of G protein-coupled receptor kinase 6 increases its activity both by increasing its hydrophobicity and localizing it near its membrane-bound substrates, and by increasing its kinase catalytic activity (Stoffel et al., *Biochemistry*, 1998, 37, 16053-16059; Stoffel et al., *J. Biol. Chem.*, 1994, 269, 27791-27794).

G protein-coupled receptor kinase 6 phosphorylates and desensitizes a number of agonist-activated G protein-coupled receptors. Rhodopsin and the  $\beta_2$ -adrenergic and m2 muscarinic cholinergic receptors are stimulus-dependent substrates for G protein-coupled receptor kinase 6 (Loudon and Benovic, *J. Biol. Chem.*, **1994**, *269*, 22691-22697). Also demonstrated to be substrates for phosphorylation by G protein-coupled receptor kinase 6 are the  $\alpha_{1B}$ -adrenergic receptor (Diviani et al., *J. Biol. Chem.*, **1996**, *271*, 5049-5058) and the muscarinic M3 acetylcholine receptor (Willets et al., *Mol. Pharmacol.*, **2001**, *60*, 321-330).

In NG108-15 mouse neuroblastoma x rat glioma hybrid cells, overexpression of G protein-coupled receptor kinase 6 increases the extent of  $\delta$ -opioid receptor desensitization upon agonist stimulation, suggesting a potential role for G protein-coupled receptor kinase 6 in drug tolerance (Willets and Kelly, *Eur. J. Pharmacol.*, **2001**, *431*, 133-141). Levels of G protein-coupled receptor kinase 6 protein are upregulated in rat brains in response to chronic treatment with the  $\mu$ -opioid receptor agonist sufentanil, further implicating G protein-coupled receptor kinase 6 in the short- and long-term adaptive changes in  $\mu$ -opioid receptor activity that contribute to development of analgesic tolerance in living animals (Hurle, *J. Neurochem.*, **2001**, *77*, 486-492).

The activity of G protein-coupled receptor kinase 6 may be modulated by phosphorylation and is inhibited by calmodulin and calcium. G protein-coupled receptor kinase 6 possesses a putative autophosphorylation motif and although it is poorly autophosphorylated *in vitro*, the protein has been shown to undergo autophosphorylation *in vivo* in COS-7 cells (Milcent et al., *Biochem. Biophys. Res. Commun.*, **1999**, *259*, 224-229). Ca<sup>2+</sup>/calmodulin acts as a potent inhibitor of G protein-

coupled receptor kinase 6 activity (Pronin et al., *J. Biol. Chem.*, 1997, 272, 18273-18280).

Impairment in  $\beta$ -adrenergic receptor activation and alterations in vascular responsiveness to  $\beta$ -adrenergic agonists is believed to play an important role in development and/or maintenance of hypertension. GRK activity is increased in human lymphocytes from hypertensive subjects, and while this has not been attributed to altered activity of G protein-coupled receptor kinase 6 specifically, a possible role for G protein-coupled receptor kinase 6 in hypertension has also not been conclusively ruled out (Gros et al., *J. Clin. Invest.*, 1997, 99, 2087-2093).

In human myometrium, hormonal receptor desensitization occurs as a loss of response to agonists such as prostaglandin  $\gamma$ , oxytocin and  $\beta_2$ -adrenergic agonists. G protein-coupled receptor kinase 6 is expressed at a much higher level in pregnant term myometrium than in non-pregnant uterine tissue, and it has been suggested that G protein-coupled receptor kinase 6 may regulate uterine contractility at term (Brenninkmeijer et al., *J. Endocrinol.*, 1999, 162, 401-408).

During hematopoietic cell development *in vitro*, expression and activity of G protein-coupled receptor kinase 6 mRNA and protein were reported to be differentially regulated in a cell type-specific manner. Thus, G protein-coupled receptor kinase 6 appears to play an important role in modulating GPCR function in immune cells (Loudon et al., *Blood*, 1996, 88, 4547-4557). Proinflammatory chemokines and cytokines and their receptors are known to play an important role in chronic inflammatory diseases by initiating and maintaining the local inflammatory process through recruitment of monocytes and lymphocytes. Alterations in GRK activity have been demonstrated in T cell activation, hypertension, and

heart disease, and the use of antagonists of GPCRs has been proven effective in reducing the onset and severity of arthritis. Thus, it has been suggested that in chronic inflammatory diseases such as rheumatoid arthritis, 5 proinflammatory signaling pathways mediated through GPCRs could be less efficiently turned off by the GRK/arrestin desensitization process. Lending support for this idea is the observation that a significant decrease in GRK activity is mirrored by a decrease in G protein-coupled receptor kinase 6 10 protein levels in peripheral blood mononuclear cells from patients with rheumatoid arthritis, and reduced GRK activity is associated with increased sensitivity to  $\beta_2$ -adrenergic activation (Lombardi et al., *Faseb J.*, 1999, 13, 715-725).

To date, investigative strategies aimed at modulating G 15 protein-coupled receptor kinase 6 function have involved the use of function blocking monoclonal antibodies, dominant negative mutants, a protein kinase C (PKC) inhibitor, and antisense expression vectors and oligonucleotides.

The monoclonal antibody (mAb) A16/17 has been 20 demonstrated to recognize the GRK4, GRK5, and G protein-coupled receptor kinase 6 proteins. In HEK 293 cells co-transfected with and overexpressing the angiotensin II type 1A receptor and one of the GRKs, the mAb A16/17 functionally inhibits G protein-coupled receptor kinase activity, reducing 25 agonist-induced phosphorylation of the receptor (Oppermann et al., *Proc. Natl. Acad. Sci. U. S. A.*, 1996, 93, 7649-7654). The same mAb was used to block G protein-coupled receptor kinase activity (measured as the level of phosphorylation of rhodopsin) in membrane extracts of late pregnant rat 30 myometrium, demonstrating that  $\beta$ -adrenergic signaling at parturition may be triggered by GRKs (Simon et al., *Endocrinology*, 2001, 142, 1899-1905).

G protein-coupled receptor kinase 6 can phosphorylate the rat follicle stimulating hormone receptor (rFSHR), and a kinase-deficient dominant negative mutant GRK6-(K215M,K216M) effectively inhibits agonist-induced phosphorylation of the rFSHR (Lazari et al., *Mol. Endocrinol.*, 1999, 13, 866-878).

The thromboxane A<sub>2</sub> receptor mediates vasoconstriction, mitogenesis, and platelet aggregation, and has been shown to undergo rapid agonist-induced desensitization. The role of GRKs in the phosphorylation and desensitization of the alpha isoform of this receptor was investigated, and it was discovered that a known PKC inhibitor, GF 109203X, also inhibited GRKs, including G protein-coupled receptor kinase 6. Because GF 109203X acts as an inhibitor of multiple kinases, it has been suggested that this compound may interact with the nucleotide-binding site common to all kinases, but the mechanism by which it inhibits G protein coupled-receptor kinase 6 remains to be elucidated (Zhou et al., *J. Pharmacol. Exp. Ther.*, 2001, 298, 1243-1251).

Another compound known to inhibit PKC, Ro 32-0432, was found to also inhibit G protein coupled-receptor kinase 6 (Aiyar et al., *Eur. J. Pharmacol.*, 2000, 403, 1-7). Again, the mechanism of inhibition of this inhibitor of multiple kinases remains unclear, but in this same study, the involvement of G protein coupled-receptor kinase 6 in calcitonin gene-related peptide (CGRP)-mediated desensitization was confirmed and further tested using antisense oligonucleotides. An antisense oligonucleotide, 18 nucleotides in length, targeted to the initiation codon of the G protein coupled-receptor kinase 6 mRNA was found to significantly reverse CGRP-mediated desensitization (Aiyar et al., *Eur. J. Pharmacol.*, 2000, 403, 1-7).

A phosphorothioate antisense oligonucleotide, 17 nucleotides in length and targeted to the initiation codon of

the G protein coupled-receptor kinase 6 mRNA was used to investigate the role of G protein coupled-receptor kinase 6 in desensitization of histamine H<sub>2</sub> receptors in the human gastric carcinoma cell line MKN-45, and it was concluded that β-  
5 adrenergic receptor kinase 1 but not G protein coupled-receptor kinase 6, is involved in desensitization of H<sub>2</sub> receptors (Nakata et al., *Digestion*, 1996, 57, 406-410).

A construct expressing the G protein coupled-receptor kinase 6 cDNA in the antisense orientation was used as a  
10 specific inhibitor of the agonist-induced phosphorylation of rFSHR to demonstrate that the desensitization of the follicle stimulating hormone response is mediated by the GRK/arrestin system (Troispoux et al., *Mol. Endocrinol.*, 1999, 13, 1599-1614).

15 Disclosed and claimed in US Patents 5,532,151 and 5,591,618 is a purified and isolated polynucleotide that is a genomic DNA, a cDNA, a plasmid insert DNA, or a wholly or partially chemically synthesized DNA, encoding a mammalian G protein-coupled receptor kinase 6 (GRK6) enzyme, a host cell  
20 stably transformed or transfected with said DNA, a method for producing GRK6 enzyme, a purified and isolated mammalian GRK6 enzyme, a mammalian GRK6 enzyme encoded by a full length DNA sequence which hybridizes to the non-coding strand corresponding to the DNA representing the coding sequence of  
25 the G protein-coupled receptor kinase 6 gene. Antisense oligonucleotides are generally disclosed (Chantry et al., 1997; Chantry et al., 1996).

Disclosed and claimed in US Patent 6,255,069 is an essentially pure cDNA encoding wild-type GRK5 or GRK6,  
30 dominant negative mutations of wild-type GRK5 or GRK6 and fragments thereof for use as probes, a recombinant DNA expression vector containing said cDNA, a process for preparing an essentially pure G protein-coupled kinase, and a

cell line comprising Sf9 insect cells capable of expressing a G protein-coupled receptor kinase comprising wild-type GRK5 or GRK6, dominant negative mutations of wild-type GRK5 or GRK6 and fragments thereof for use as probes. Antisense  
5 oligonucleotides are generally disclosed (Benovic et al., 2001).

Disclosed and claimed in PCT Publication WO 00/18895 is a peptide comprising a peptide derivative of the  $\alpha$ D region of a protein kinase, wherein said protein has between about five  
10 and about thirty amino acids or amino acid analogs and said peptide modulates activity of the protein kinase, and wherein the protein kinase is a member of a protein kinase family selected from a group of which the G protein-coupled receptor kinases and specifically G protein-coupled receptor kinase 6  
15 are members. Further claimed is a method of identifying a peptide which modulates the activity of a protein kinase, a method of modulating the activity of a protein kinase in a subject, a method of detecting a ligand that binds to the  $\alpha$ D region of a protein kinase, and an antibody as well as a  
20 method of producing antibodies that immunologically bind to the  $\alpha$ D region of a protein kinase (Ben-Sasson, 2000).

Currently, there are no known therapeutic agents which effectively inhibit the synthesis of G protein-coupled receptor kinase 6.

25 Consequently, there remains a long felt need for agents capable of effectively inhibiting G protein-coupled receptor kinase 6 function.

Antisense technology is emerging as an effective means for reducing the expression of specific gene products and may  
30 therefore prove to be uniquely useful in a number of therapeutic, diagnostic, and research applications for the modulation of G protein-coupled receptor kinase 6 expression.

The present invention provides compositions and methods for modulating G protein-coupled receptor kinase 6 expression.

## 5 SUMMARY OF THE INVENTION

The present invention is directed to compounds, particularly antisense oligonucleotides, which are targeted to a nucleic acid encoding G protein-coupled receptor kinase 6, and which modulate the expression of G protein-coupled  
10 receptor kinase 6. Pharmaceutical and other compositions comprising the compounds of the invention are also provided. Further provided are methods of modulating the expression of G protein-coupled receptor kinase 6 in cells or tissues comprising contacting said cells or tissues with one or more  
15 of the antisense compounds or compositions of the invention. Further provided are methods of treating an animal, particularly a human, suspected of having or being prone to a disease or condition associated with expression of G protein-coupled receptor kinase 6 by administering a therapeutically  
20 or prophylactically effective amount of one or more of the antisense compounds or compositions of the invention.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention employs oligomeric compounds,  
25 particularly antisense oligonucleotides, for use in modulating the function of nucleic acid molecules encoding G protein-coupled receptor kinase 6, ultimately modulating the amount of G protein-coupled receptor kinase 6 produced. This is accomplished by providing antisense compounds which  
30 specifically hybridize with one or more nucleic acids encoding G protein-coupled receptor kinase 6. As used herein, the terms "target nucleic acid" and "nucleic acid encoding G protein-coupled receptor kinase 6" encompass DNA encoding G

protein-coupled receptor kinase 6, RNA (including pre-mRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds which specifically hybridize to it is generally referred to as "antisense". The functions of DNA to be interfered with include replication and transcription. The functions of RNA to be interfered with include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translocation of the RNA to sites within the cell which are distant from the site of RNA synthesis, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of G protein-coupled receptor kinase 6. In the context of the present invention, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene. In the context of the present invention, inhibition is the preferred form of modulation of gene expression and mRNA is a preferred target.

It is preferred to target specific nucleic acids for antisense. "Targeting" an antisense compound to a particular nucleic acid, in the context of this invention, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present invention, the target is a nucleic acid molecule encoding G protein-

coupled receptor kinase 6. The targeting process also includes determination of a site or sites within this gene for the antisense interaction to occur such that the desired effect, e.g., detection or modulation of expression of the protein, will result. Within the context of the present invention, a preferred intragenic site is the region encompassing the translation initiation or termination codon of the open reading frame (ORF) of the gene. Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes have a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG, and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function *in vivo*. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). It is also known in the art that eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular set of conditions. In the context of the invention, "start codon" and "translation initiation codon" refer to the codon or codons that are used *in vivo* to initiate translation of an mRNA molecule transcribed from a gene encoding G protein-coupled receptor kinase 6, regardless of the sequence(s) of such codons.

It is also known in the art that a translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA,

respectively). The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon.

The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Other target regions include the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA or corresponding nucleotides on the gene, and the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA or corresponding nucleotides on the gene. The 5' cap of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap. The 5' cap region may also be a preferred target region.

Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as

"introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. mRNA splice sites, i.e., intron-  
5 exon junctions, may also be preferred target regions, and are particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular mRNA splice product is implicated in disease. Aberrant fusion junctions due to rearrangements or deletions  
10 are also preferred targets. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts". It has also been found that introns can be effective, and therefore preferred, target regions for antisense compounds targeted, for example,  
15 to DNA or pre-mRNA.

It is also known in the art that alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are  
20 transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and extronic regions.

Upon excision of one or more exon or intron regions or  
25 portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative  
30 splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

It is also known in the art that variants can be produced through the use of alternative signals to start or stop transcription and that pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites.

Once one or more target sites have been identified, oligonucleotides are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

In the context of this invention, "hybridization" means hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases. For example, adenine and thymine are complementary nucleobases which pair through the formation of hydrogen bonds. "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides. For example, if a nucleotide at a certain position of an oligonucleotide is capable of hydrogen bonding with a nucleotide at the same position of a DNA or RNA molecule, then the oligonucleotide and the DNA or RNA are considered to be complementary to each other at that position. The oligonucleotide and the DNA or RNA are complementary to each other when a sufficient number of corresponding positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically

hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of complementarity or precise pairing such that stable and specific binding occurs between the oligonucleotide and the DNA or RNA target. It is  
5 understood in the art that the sequence of an antisense compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable.

An antisense compound is specifically hybridizable when binding of the compound to the target DNA or RNA molecule  
10 interferes with the normal function of the target DNA or RNA to cause a loss of activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological  
15 conditions in the case of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed. It is preferred that the antisense compounds of the present invention comprise at least 80% sequence complementarity to a target region  
20 within the target nucleic acid, moreover that they comprise 90% sequence complementarity and even more comprise 95% sequence complementarity to the target region within the target nucleic acid sequence to which they are targeted. For example, an antisense compound in which 18 of 20 nucleobases  
25 of the antisense compound are complementary, and would therefore specifically hybridize, to a target region would represent 90 percent complementarity. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using basic local alignment  
30 search tools (BLAST programs) (Altschul et al., *J. Mol. Biol.*, 1990, 215, 403-410; Zhang and Madden, *Genome Res.*, 1997, 7, 649-656).

Antisense and other compounds of the invention, which

hybridize to the target and inhibit expression of the target, are identified through experimentation, and representative sequences of these compounds are hereinbelow identified as preferred embodiments of the invention. The sites to which  
5 these preferred antisense compounds are specifically hybridizable are hereinbelow referred to as "preferred target regions" and are therefore preferred sites for targeting. As used herein the term "preferred target region" is defined as at least an 8-nucleobase portion of a target region to which  
10 an active antisense compound is targeted. While not wishing to be bound by theory, it is presently believed that these target regions represent regions of the target nucleic acid which are accessible for hybridization.

While the specific sequences of particular preferred  
15 target regions are set forth below, one of skill in the art will recognize that these serve to illustrate and describe particular embodiments within the scope of the present invention. Additional preferred target regions may be identified by one having ordinary skill.

20 Target regions 8-80 nucleobases in length comprising a stretch of at least eight (8) consecutive nucleobases selected from within the illustrative preferred target regions are considered to be suitable preferred target regions as well.

Exemplary good preferred target regions include DNA or  
25 RNA sequences that comprise at least the 8 consecutive nucleobases from the 5'-terminus of one of the illustrative preferred target regions (the remaining nucleobases being a consecutive stretch of the same DNA or RNA beginning immediately upstream of the 5'-terminus of the target region  
30 and continuing until the DNA or RNA contains about 8 to about 80 nucleobases). Similarly good preferred target regions are represented by DNA or RNA sequences that comprise at least the 8 consecutive nucleobases from the 3'-terminus of one of the

illustrative preferred target regions (the remaining nucleobases being a consecutive stretch of the same DNA or RNA beginning immediately downstream of the 3'-terminus of the target region and continuing until the DNA or RNA contains  
5 about 8 to about 80 nucleobases). One having skill in the art, once armed with the empirically-derived preferred target regions illustrated herein will be able, without undue experimentation, to identify further preferred target regions. In addition, one having ordinary skill in the art will also be  
10 able to identify additional compounds, including oligonucleotide probes and primers, that specifically hybridize to these preferred target regions using techniques available to the ordinary practitioner in the art.

Antisense compounds are commonly used as research  
15 reagents and diagnostics. For example, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes. Antisense compounds are also used, for example, to distinguish  
20 between functions of various members of a biological pathway. Antisense modulation has, therefore, been harnessed for research use.

For use in kits and diagnostics, the antisense compounds of the present invention, either alone or in combination with  
25 other antisense compounds or therapeutics, can be used as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

Expression patterns within cells or tissues treated with  
30 one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease

association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other  
5 compounds which affect expression patterns.

Examples of methods of gene expression analysis known in the art include DNA arrays or microarrays (Brazma and Vilo, *FEBS Lett.*, 2000, 480, 17-24; Celis, et al., *FEBS Lett.*, 2000, 480, 2-16), SAGE (serial analysis of gene expression) (Madden, 10 et al., *Drug Discov. Today*, 2000, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, *Methods Enzymol.*, 1999, 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., *Proc. Natl. Acad. Sci. U. S. A.*, 2000, 97, 1976-81), protein arrays  
15 and proteomics (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Jungblut, et al., *Electrophoresis*, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., *FEBS Lett.*, 2000, 480, 2-16; Larsson, et al., *J. Biotechnol.*, 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et  
20 al., *Anal. Biochem.*, 2000, 286, 91-98; Larson, et al., *Cytometry*, 2000, 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, *Curr. Opin. Microbiol.*, 2000, 3, 316-21), comparative genomic hybridization (Carulli, et al., *J. Cell Biochem. Suppl.*, 1998,  
25 31, 286-96), FISH (fluorescent *in situ* hybridization) techniques (Going and Gusterson, *Eur. J. Cancer*, 1999, 35, 1895-904) and mass spectrometry methods (reviewed in To, *Comb. Chem. High Throughput Screen*, 2000, 3, 235-41).

The specificity and sensitivity of antisense is also  
30 harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotide drugs, including ribozymes,

have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

In the context of this invention, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. This term includes oligonucleotides composed of naturally-occurring nucleobases, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally-occurring portions which function similarly. Such modified or substituted oligonucleotides are often preferred over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for nucleic acid target and increased stability in the presence of nucleases.

While antisense oligonucleotides are a preferred form of antisense compound, the present invention comprehends other oligomeric antisense compounds, including but not limited to oligonucleotide mimetics such as are described below. The antisense compounds in accordance with this invention preferably comprise from about 8 to about 80 nucleobases (i.e. from about 8 to about 80 linked nucleosides). Particularly preferred antisense compounds are antisense oligonucleotides from about 8 to about 50 nucleobases, even more preferably those comprising from about 12 to about 30 nucleobases. Antisense compounds include ribozymes, external guide sequence (EGS) oligonucleotides (oligozymes), and other short catalytic RNAs or catalytic oligonucleotides which hybridize to the target nucleic acid and modulate its expression.

Antisense compounds 8-80 nucleobases in length comprising a stretch of at least eight (8) consecutive nucleobases selected from within the illustrative antisense compounds are considered to be suitable antisense compounds as well.

5 Exemplary preferred antisense compounds include DNA or RNA sequences that comprise at least the 8 consecutive nucleobases from the 5'-terminus of one of the illustrative preferred antisense compounds (the remaining nucleobases being a consecutive stretch of the same DNA or RNA beginning  
10 immediately upstream of the 5'-terminus of the antisense compound which is specifically hybridizable to the target nucleic acid and continuing until the DNA or RNA contains about 8 to about 80 nucleobases). Similarly preferred antisense compounds are represented by DNA or RNA sequences  
15 that comprise at least the 8 consecutive nucleobases from the 3'-terminus of one of the illustrative preferred antisense compounds (the remaining nucleobases being a consecutive stretch of the same DNA or RNA beginning immediately  
20 downstream of the 3'-terminus of the antisense compound which is specifically hybridizable to the target nucleic acid and continuing until the DNA or RNA contains about 8 to about 80 nucleobases). One having skill in the art, once armed with the empirically-derived preferred antisense compounds  
25 illustrated herein will be able, without undue experimentation, to identify further preferred antisense compounds.

Antisense and other compounds of the invention, which hybridize to the target and inhibit expression of the target, are identified through experimentation, and representative  
30 sequences of these compounds are herein identified as preferred embodiments of the invention. While specific sequences of the antisense compounds are set forth herein, one of skill in the art will recognize that these serve to

illustrate and describe particular embodiments within the scope of the present invention. Additional preferred antisense compounds may be identified by one having ordinary skill.

5           As is known in the art, a nucleoside is a base-sugar combination. The base portion of the nucleoside is normally a heterocyclic base. The two most common classes of such heterocyclic bases are the purines and the pyrimidines. Nucleotides are nucleosides that further include a phosphate  
10   group covalently linked to the sugar portion of the nucleoside. For those nucleosides that include a pentofuranosyl sugar, the phosphate group can be linked to either the 2', 3' or 5' hydroxyl moiety of the sugar. In forming oligonucleotides, the phosphate groups covalently link  
15   adjacent nucleosides to one another to form a linear polymeric compound. In turn, the respective ends of this linear polymeric structure can be further joined to form a circular structure, however, open linear structures are generally preferred. In addition, linear structures may also have  
20   internal nucleobase complementarity and may therefore fold in a manner as to produce a double stranded structure. Within the oligonucleotide structure, the phosphate groups are commonly referred to as forming the internucleoside backbone of the oligonucleotide. The normal linkage or backbone of RNA  
25   and DNA is a 3' to 5' phosphodiester linkage.

          Specific examples of preferred antisense compounds useful in this invention include oligonucleotides containing modified  
backbones or non-natural internucleoside linkages. As defined in this specification, oligonucleotides having modified  
30   backbones include those that retain a phosphorus atom in the backbone and those that do not have a phosphorus atom in the backbone. For the purposes of this specification, and as sometimes referenced in the art, modified oligonucleotides

that do not have a phosphorus atom in their internucleoside backbone can also be considered to be oligonucleosides.

Preferred modified oligonucleotide backbones include, for example, phosphorothioates, chiral phosphorothioates, 5 phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates, 5'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, 10 thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, selenophosphates and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein one or more internucleotide linkages is a 3' to 3', 5' to 5' or 2' to 15 2' linkage. Preferred oligonucleotides having inverted polarity comprise a single 3' to 3' linkage at the 3'-most internucleotide linkage i.e. a single inverted nucleoside residue which may be abasic (the nucleobase is missing or has a hydroxyl group in place thereof). Various salts, mixed 20 salts and free acid forms are also included.

Representative United States patents that teach the preparation of the above phosphorus-containing linkages include, but are not limited to, U.S.: 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5,177,196; 5,188,897; 5,264,423; 25 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455,233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563,253; 5,571,799; 5,587,361; 5,194,599; 5,565,555; 5,527,899; 5,721,218; 5,672,697 and 5,625,050, certain of which are 30 commonly owned with this application, and each of which is herein incorporated by reference.

Preferred modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are

formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These include those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; riboacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH<sub>2</sub> component parts.

Representative United States patents that teach the preparation of the above oligonucleosides include, but are not limited to, U.S.: 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264,562; 5,264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596,086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623,070; 5,663,312; 5,633,360; 5,677,437; 5,792,608; 5,646,269 and 5,677,439, certain of which are commonly owned with this application, and each of which is herein incorporated by reference.

In other preferred oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine

backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S.: 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen *et al.*, *Science*, **1991**, 254, 1497-1500.

Most preferred embodiments of the invention are oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular -CH<sub>2</sub>-NH-O-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-O-CH<sub>2</sub>- [known as a methylene (methylimino) or MMI backbone], -CH<sub>2</sub>-O-N(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-N(CH<sub>3</sub>)-N(CH<sub>3</sub>)-CH<sub>2</sub>- and -O-N(CH<sub>3</sub>)-CH<sub>2</sub>-CH<sub>2</sub>- [wherein the native phosphodiester backbone is represented as -O-P-O-CH<sub>2</sub>-] of the above referenced U.S. patent 5,489,677, and the amide backbones of the above referenced U.S. patent 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced U.S. patent 5,034,506.

Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S- or N-alkynyl; or O-alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may be substituted or unsubstituted C<sub>1</sub> to C<sub>10</sub> alkyl or C<sub>2</sub> to C<sub>10</sub> alkenyl and alkynyl. Particularly preferred are O[(CH<sub>2</sub>)<sub>n</sub>O]<sub>m</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>OCH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>NH<sub>2</sub>, O(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>, O(CH<sub>2</sub>)<sub>n</sub>ONH<sub>2</sub>, and O(CH<sub>2</sub>)<sub>n</sub>ON[(CH<sub>2</sub>)<sub>n</sub>CH<sub>3</sub>]<sub>2</sub>, where n and m are from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C<sub>1</sub> to C<sub>10</sub> lower alkyl, substituted lower alkyl, alkenyl, alkynyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH<sub>3</sub>, OCN, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, SOCH<sub>3</sub>, SO<sub>2</sub>CH<sub>3</sub>, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, heterocycloalkyl, heterocycloalkaryl,

aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy (2'-O-CH<sub>2</sub>CH<sub>2</sub>OCH<sub>3</sub>, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin *et al.*, *Helv. Chim. Acta*, 1995, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification includes 2'-dimethylaminoxyethoxy, i.e., a O(CH<sub>2</sub>)<sub>2</sub>ON(CH<sub>3</sub>)<sub>2</sub> group, also known as 2'-DMAOE, as described in examples hereinbelow, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethyl-aminoethoxy-ethyl or 2'-DMAEOE), i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>, also described in examples hereinbelow.

Other preferred modifications include 2'-methoxy (2'-O-CH<sub>3</sub>), 2'-aminopropoxy (2'-OCH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>), 2'-allyl (2'-CH<sub>2</sub>-CH=CH<sub>2</sub>), 2'-O-allyl (2'-O-CH<sub>2</sub>-CH=CH<sub>2</sub>) and 2'-fluoro (2'-F). The 2'-modification may be in the arabino (up) position or ribo (down) position. A preferred 2'-arabino modification is 2'-F. Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures include, but are not limited to, U.S.: 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514,785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646,265; 5,658,873; 5,670,633; 5,792,747; and 5,700,920, certain of which are

commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

A further preferred modification includes Locked Nucleic Acids (LNAs) in which the 2'-hydroxyl group is linked to the 3' or 4' carbon atom of the sugar ring thereby forming a bicyclic sugar moiety. The linkage is preferably a methylene (-CH<sub>2</sub>-)<sub>n</sub> group bridging the 2' oxygen atom and the 4' carbon atom wherein n is 1 or 2. LNAs and preparation thereof are described in WO 98/39352 and WO 99/14226.

Oligonucleotides may also include nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleobases include the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleobases include other synthetic and natural nucleobases such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl (-C≡C-CH<sub>3</sub>) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine (1H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-

clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridoindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent  
5 No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch *et al.*, *Angewandte Chemie*, International Edition, 1991, 30, 613, and those disclosed by Sanghvi, Y.S., Chapter 15,  
15 *Antisense Research and Applications*, pages 289-302, Crooke, S.T. and Lebleu, B., ed., CRC Press, 1993. Certain of these nucleobases are particularly useful for increasing the binding affinity of the oligomeric compounds of the invention. These include 5-substituted pyrimidines, 6-azapyrimidines and N-2,  
20 N-6 and O-6 substituted purines, including 2-aminopropyl-adenine, 5-propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds., *Antisense Research and*  
25 *Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

Representative United States patents that teach the  
30 preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include, but are not limited to, the above noted U.S. 3,687,808, as well as U.S.: 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066;

5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177;  
5,525,711; 5,552,540; 5,587,469; 5,594,121, 5,596,091;  
5,614,617; 5,645,985; 5,830,653; 5,763,588; 6,005,096; and  
5,681,941, certain of which are commonly owned with the

5 instant application, and each of which is herein incorporated  
by reference, and United States patent 5,750,692, which is  
commonly owned with the instant application and also herein  
incorporated by reference.

Another modification of the oligonucleotides of the  
10 invention involves chemically linking to the oligonucleotide  
one or more moieties or conjugates which enhance the activity,  
cellular distribution or cellular uptake of the  
oligonucleotide. The compounds of the invention can include  
conjugate groups covalently bound to functional groups such as  
15 primary or secondary hydroxyl groups. Conjugate groups of the  
invention include intercalators, reporter molecules,  
polyamines, polyamides, polyethylene glycols, polyethers,  
groups that enhance the pharmacodynamic properties of  
oligomers, and groups that enhance the pharmacokinetic  
20 properties of oligomers. Typical conjugate groups include  
cholesterols, lipids, phospholipids, biotin, phenazine,  
folate, phenanthridine, anthraquinone, acridine, fluoresceins,  
rhodamines, coumarins, and dyes. Groups that enhance the  
pharmacodynamic properties, in the context of this invention,  
25 include groups that improve oligomer uptake, enhance oligomer  
resistance to degradation, and/or strengthen sequence-specific  
hybridization with RNA. Groups that enhance the pharmaco-  
kinetic properties, in the context of this invention, include  
groups that improve oligomer uptake, distribution, metabolism  
30 or excretion. Representative conjugate groups are disclosed  
in International Patent Application PCT/US92/09196, filed  
October 23, 1992 the entire disclosure of which is incor-  
porated herein by reference. Conjugate moieties include but

are not limited to lipid moieties such as a cholesterol moiety (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86, 6553-6556), cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660, 306-309; 5 Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20, 533-538), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10, 10 1111-1118; Kabanov et al., *FEBS Lett.*, 1990, 259, 327-330; Svinarchuk et al., *Biochimie*, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654; 15 Shea et al., *Nucl. Acids Res.*, 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., *Biochim. 20 Biophys. Acta*, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277, 923-937). Oligonucleotides of the invention may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, 25 ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepam, indomethacin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial 30 or an antibiotic. Oligonucleotide-drug conjugates and their preparation are described in United States Patent Application 09/334,130 (filed June 15, 1999) which is incorporated herein by reference in its entirety.

Representative United States patents that teach the preparation of such oligonucleotide conjugates include, but are not limited to, U.S.: 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717, 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 10 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241, 5,391,723; 5,416,203, 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 15 5,688,941, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference.

It is not necessary for all positions in a given compound to be uniformly modified, and in fact more than one of the 20 aforementioned modifications may be incorporated in a single compound or even at a single nucleoside within an oligonucleotide. The present invention also includes antisense compounds which are chimeric compounds. "Chimeric" antisense compounds or "chimeras," in the context of this 25 invention, are antisense compounds, particularly oligonucleotides, which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region 30 wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, increased stability and/or increased binding affinity for the target nucleic acid.

An additional region of the oligonucleotide may serve as a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA  
5 duplex. Activation of RNase H, therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of oligonucleotide inhibition of gene expression. The cleavage of RNA:RNA hybrids can, in like fashion, be accomplished through the actions of endoribonucleases, such as interferon-  
10 induced RNaseL which cleaves both cellular and viral RNA. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA  
15 target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art.

Chimeric antisense compounds of the invention may be formed as composite structures of two or more  
20 oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide mimetics as described above. Such compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not  
25 limited to, U.S.: 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

30 The antisense compounds used in accordance with this invention may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including, for

example, Applied Biosystems (Foster City, CA). Any other means for such synthesis known in the art may additionally or alternatively be employed. It is well known to use similar techniques to prepare oligonucleotides such as the  
5 phosphorothioates and alkylated derivatives.

The compounds of the invention may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor-targeted molecules, oral,  
10 rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption-assisting formulations include, but are not limited to, U.S.: 5,108,921; 5,354,844; 5,416,016;  
15 5,459,127; 5,521,291; 5,543,158; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and  
20 5,595,756, each of which is herein incorporated by reference.

The antisense compounds of the invention encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal, including a human, is capable of providing (directly  
25 or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to prodrugs and pharmaceutically acceptable salts of the compounds of the invention, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents.

30 The term "prodrug" indicates a therapeutic agent that is prepared in an inactive form that is converted to an active form (i.e., drug) within the body or cells thereof by the action of endogenous enzymes or other chemicals and/or

conditions. In particular, prodrug versions of the oligonucleotides of the invention are prepared as SATE [(S-acetyl-2-thioethyl) phosphate] derivatives according to the methods disclosed in WO 93/24510 to Gosselin *et al.*,  
5 published December 9, 1993 or in WO 94/26764 and U.S. 5,770,713 to Imbach *et al.*

The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the invention: i.e., salts that retain the  
10 desired biological activity of the parent compound and do not impart undesired toxicological effects thereto.

Pharmaceutically acceptable base addition salts are formed with metals or amines, such as alkali and alkaline earth metals or organic amines. Examples of metals used as  
15 cations are sodium, potassium, magnesium, calcium, and the like. Examples of suitable amines are N,N'-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, dicyclohexylamine, ethylenediamine, N-methylglucamine, and procaine (see, for example, Berge *et al.*, "Pharmaceutical Salts," *J. of Pharma Sci.*, 1977, 66, 1-  
20 19). The base addition salts of said acidic compounds are prepared by contacting the free acid form with a sufficient amount of the desired base to produce the salt in the conventional manner. The free acid form may be regenerated by  
25 contacting the salt form with an acid and isolating the free acid in the conventional manner. The free acid forms differ from their respective salt forms somewhat in certain physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free acid for  
30 purposes of the present invention. As used herein, a "pharmaceutical addition salt" includes a pharmaceutically acceptable salt of an acid form of one of the components of the compositions of the invention. These include organic or

inorganic acid salts of the amines. Preferred acid salts are the hydrochlorides, acetates, salicylates, nitrates and phosphates. Other suitable pharmaceutically acceptable salts are well known to those skilled in the art and include basic salts of a variety of inorganic and organic acids, such as, for example, with inorganic acids, such as for example hydrochloric acid, hydrobromic acid, sulfuric acid or phosphoric acid; with organic carboxylic, sulfonic, sulfo or phospho acids or N-substituted sulfamic acids, for example acetic acid, propionic acid, glycolic acid, succinic acid, maleic acid, hydroxymaleic acid, methylmaleic acid, fumaric acid, malic acid, tartaric acid, lactic acid, oxalic acid, gluconic acid, glucaric acid, glucuronic acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, salicylic acid, 4-aminosalicylic acid, 2-phenoxybenzoic acid, 2-acetoxybenzoic acid, embonic acid, nicotinic acid or isonicotinic acid; and with amino acids, such as the 20 alpha-amino acids involved in the synthesis of proteins in nature, for example glutamic acid or aspartic acid, and also with phenylacetic acid, methanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, ethane-1,2-disulfonic acid, benzenesulfonic acid, 4-methylbenzenesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 2- or 3-phosphoglycerate, glucose-6-phosphate, N-cyclohexylsulfamic acid (with the formation of cyclamates), or with other acid organic compounds, such as ascorbic acid. Pharmaceutically acceptable salts of compounds may also be prepared with a pharmaceutically acceptable cation. Suitable pharmaceutically acceptable cations are well known to those skilled in the art and include alkaline, alkaline earth, ammonium and quaternary ammonium cations. Carbonates or hydrogen carbonates are also possible.

For oligonucleotides, preferred examples of

pharmaceutically acceptable salts include but are not limited to (a) salts formed with cations such as sodium, potassium, ammonium, magnesium, calcium, polyamines such as spermine and spermidine, etc.; (b) acid addition salts formed with  
5 inorganic acids, for example hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, nitric acid and the like; (c) salts formed with organic acids such as, for example, acetic acid, oxalic acid, tartaric acid, succinic acid, maleic acid, fumaric acid, gluconic acid, citric acid,  
10 malic acid, ascorbic acid, benzoic acid, tannic acid, palmitic acid, alginic acid, polyglutamic acid, naphthalenesulfonic acid, methanesulfonic acid, p-toluenesulfonic acid, naphthalenedisulfonic acid, polygalacturonic acid, and the like; and (d) salts formed from  
15 elemental anions such as chlorine, bromine, and iodine.

The antisense compounds of the present invention can be utilized for diagnostics, therapeutics, prophylaxis and as research reagents and kits. For therapeutics, an animal, preferably a human, suspected of having a disease or disorder  
20 which can be treated by modulating the expression of G protein-coupled receptor kinase 6 is treated by administering antisense compounds in accordance with this invention. The compounds of the invention can be utilized in pharmaceutical compositions by adding an effective amount of an antisense  
25 compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the antisense compounds and methods of the invention may also be useful prophylactically, e.g., to prevent or delay infection, inflammation or tumor formation, for example.

30 The antisense compounds of the invention are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding G protein-coupled receptor kinase 6, enabling sandwich and other assays to easily be constructed to

exploit this fact. Hybridization of the antisense oligonucleotides of the invention with a nucleic acid encoding G protein-coupled receptor kinase 6 can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabelling of the oligonucleotide or any other suitable detection means. Kits using such detection means for detecting the level of G protein-coupled receptor kinase 6 in a sample may also be prepared.

10 The present invention also includes pharmaceutical compositions and formulations which include the antisense compounds of the invention. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration. Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration.

Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful. Preferred topical formulations include those in which

the oligonucleotides of the invention are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoylphosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoylphosphatidyl ethanolamine DOTMA). Oligonucleotides of the invention may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters include but are not limited arachidonic acid, oleic acid, eicosanoic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a C<sub>1-10</sub> alkyl ester (e.g. isopropylmyristate IPM), monoglyceride, diglyceride or pharmaceutically acceptable salt thereof. Topical formulations are described in detail in United States patent application 09/315,298 filed on May 20, 1999 which is incorporated herein by reference in its entirety.

Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitabets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the invention are administered in

conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts include chenodeoxycholic acid (CDCA) and ursodeoxychenodeoxycholic acid (UDCA), cholic acid, dehydrocholic acid, deoxycholic acid, glucolic acid, glycholic acid, glycodeoxycholic acid, taurocholic acid, taurodeoxycholic acid, sodium tauro-24,25-dihydro-fusidate and sodium glycodihydrofusidate. Preferred fatty acids include arachidonic acid, undecanoic acid, oleic acid, lauric acid, caprylic acid, capric acid, myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein, dilaurin, glyceryl 1-monocaprinate, 1-dodecylazacycloheptan-2-one, an acylcarnitine, an acylcholine, or a monoglyceride, a diglyceride or a pharmaceutically acceptable salt thereof (e.g. sodium). Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the invention may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents include poly-amino acids; polyimines; polyacrylates; polyalkylacrylates, polyoxethanes, polyalkylcyanoacrylates; cationized gelatins, albumins, starches, acrylates, polyethyleneglycols (PEG) and starches; polyalkylcyanoacrylates; DEAE-derivatized polyimines, pullulans, celluloses and starches. Particularly preferred complexing agents include chitosan, N-trimethylchitosan, poly-L-lysine, polyhistidine, polyornithine, polyspermines,

protamine, polyvinylpyridine, polythiodiethylamino-  
methylethylene P(TDAE), polyaminostyrene (e.g. p-amino),  
poly(methylcyanoacrylate), poly(ethylcyanoacrylate),  
poly(butylcyanoacrylate), poly(isobutylcyanoacrylate),  
5 poly(isohexylcynaoacrylate), DEAE-methacrylate, DEAE-  
hexylacrylate, DEAE-acrylamide, DEAE-albumin and DEAE-dextran,  
polymethylacrylate, polyhexylacrylate, poly(D,L-lactic acid),  
poly(DL-lactic-co-glycolic acid (PLGA), alginate, and  
polyethyleneglycol (PEG). Oral formulations for  
10 oligonucleotides and their preparation are described in detail  
in United States applications 08/886,829 (filed July 1, 1997),  
09/108,673 (filed July 1, 1998), 09/256,515 (filed February  
23, 1999), 09/082,624 (filed May 21, 1998) and 09/315,298  
(filed May 20, 1999), each of which is incorporated herein by  
15 reference in their entirety.

Compositions and formulations for parenteral, intrathecal  
or intraventricular administration may include sterile aqueous  
solutions which may also contain buffers, diluents and other  
suitable additives such as, but not limited to, penetration  
20 enhancers, carrier compounds and other pharmaceutically  
acceptable carriers or excipients.

Pharmaceutical compositions of the present invention  
include, but are not limited to, solutions, emulsions, and  
liposome-containing formulations. These compositions may be  
25 generated from a variety of components that include, but are  
not limited to, preformed liquids, self-emulsifying solids and  
self-emulsifying semisolids.

The pharmaceutical formulations of the present invention,  
which may conveniently be presented in unit dosage form, may  
30 be prepared according to conventional techniques well known in  
the pharmaceutical industry. Such techniques include the step  
of bringing into association the active ingredients with the  
pharmaceutical carrier(s) or excipient(s). In general, the

formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

5           The compositions of the present invention may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels, suppositories, and enemas. The compositions of the present invention may also be formulated  
10 as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

15           In one embodiment of the present invention the pharmaceutical compositions may be formulated and used as foams. Pharmaceutical foams include formulations such as, but not limited to, emulsions, microemulsions, creams, jellies and liposomes. While basically similar in nature these  
20 formulations vary in the components and the consistency of the final product. The preparation of such compositions and formulations is generally known to those skilled in the pharmaceutical and formulation arts and may be applied to the formulation of the compositions of the present invention.

25

#### Emulsions

The compositions of the present invention may be prepared and formulated as emulsions. Emulsions are typically heterogenous systems of one liquid dispersed in another in the  
30 form of droplets usually exceeding 0.1  $\mu\text{m}$  in diameter (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199; Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman,

Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., Volume 1, p. 245; Block in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 2, p. 335; Higuchi et al., in  
5 *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 301). Emulsions are often biphasic systems comprising two immiscible liquid phases intimately mixed and dispersed with each other. In general, emulsions may be of either the water-in-oil (w/o) or the oil-in-water  
10 (o/w) variety. When an aqueous phase is finely divided into and dispersed as minute droplets into a bulk oily phase, the resulting composition is called a water-in-oil (w/o) emulsion. Alternatively, when an oily phase is finely divided into and dispersed as minute droplets into a bulk aqueous phase, the  
15 resulting composition is called an oil-in-water (o/w) emulsion. Emulsions may contain additional components in addition to the dispersed phases, and the active drug which may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Pharmaceutical  
20 excipients such as emulsifiers, stabilizers, dyes, and anti-oxidants may also be present in emulsions as needed. Pharmaceutical emulsions may also be multiple emulsions that are comprised of more than two phases such as, for example, in the case of oil-in-water-in-oil (o/w/o) and water-in-oil-in-  
25 water (w/o/w) emulsions. Such complex formulations often provide certain advantages that simple binary emulsions do not. Multiple emulsions in which individual oil droplets of an o/w emulsion enclose small water droplets constitute a w/o/w emulsion. Likewise a system of oil droplets enclosed in  
30 globules of water stabilized in an oily continuous phase provides an o/w/o emulsion.

Emulsions are characterized by little or no thermodynamic stability. Often, the dispersed or discontinuous phase of the

emulsion is well dispersed into the external or continuous phase and maintained in this form through the means of emulsifiers or the viscosity of the formulation. Either of the phases of the emulsion may be a semisolid or a solid, as  
5 is the case of emulsion-style ointment bases and creams. Other means of stabilizing emulsions entail the use of emulsifiers that may be incorporated into either phase of the emulsion. Emulsifiers may broadly be classified into four categories: synthetic surfactants, naturally occurring  
10 emulsifiers, absorption bases, and finely dispersed solids (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Synthetic surfactants, also known as surface active  
15 agents, have found wide applicability in the formulation of emulsions and have been reviewed in the literature (Rieger, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285; Idson, in *Pharmaceutical Dosage Forms*, Lieberman,  
20 Rieger and Banker (Eds.), Marcel Dekker, Inc., New York, N.Y., 1988, volume 1, p. 199). Surfactants are typically amphiphilic and comprise a hydrophilic and a hydrophobic portion. The ratio of the hydrophilic to the hydrophobic nature of the surfactant has been termed the  
25 hydrophile/lipophile balance (HLB) and is a valuable tool in categorizing and selecting surfactants in the preparation of formulations. Surfactants may be classified into different classes based on the nature of the hydrophilic group: nonionic, anionic, cationic and amphoteric (Rieger, in  
30 *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 285).

Naturally occurring emulsifiers used in emulsion formulations include lanolin, beeswax, phosphatides, lecithin and acacia. Absorption bases possess hydrophilic properties such that they can soak up water to form w/o emulsions yet  
5 retain their semisolid consistencies, such as anhydrous lanolin and hydrophilic petrolatum. Finely divided solids have also been used as good emulsifiers especially in combination with surfactants and in viscous preparations. These include polar inorganic solids, such as heavy metal  
10 hydroxides, nonswelling clays such as bentonite, attapulgate, hectorite, kaolin, montmorillonite, colloidal aluminum silicate and colloidal magnesium aluminum silicate, pigments and nonpolar solids such as carbon or glyceryl tristearate.

A large variety of non-emulsifying materials are also  
15 included in emulsion formulations and contribute to the properties of emulsions. These include fats, oils, waxes, fatty acids, fatty alcohols, fatty esters, humectants, hydrophilic colloids, preservatives and antioxidants (Block, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker  
20 (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199).

Hydrophilic colloids or hydrocolloids include naturally  
25 occurring gums and synthetic polymers such as polysaccharides (for example, acacia, agar, alginic acid, carrageenan, guar gum, karaya gum, and tragacanth), cellulose derivatives (for example, carboxymethylcellulose and carboxypropylcellulose), and synthetic polymers (for example, carbomers, cellulose  
30 ethers, and carboxyvinyl polymers). These disperse or swell in water to form colloidal solutions that stabilize emulsions by forming strong interfacial films around the dispersed-phase

droplets and by increasing the viscosity of the external phase.

Since emulsions often contain a number of ingredients such as carbohydrates, proteins, sterols and phosphatides that may readily support the growth of microbes, these formulations often incorporate preservatives. Commonly used preservatives included in emulsion formulations include methyl paraben, propyl paraben, quaternary ammonium salts, benzalkonium chloride, esters of p-hydroxybenzoic acid, and boric acid. Antioxidants are also commonly added to emulsion formulations to prevent deterioration of the formulation. Antioxidants used may be free radical scavengers such as tocopherols, alkyl gallates, butylated hydroxyanisole, butylated hydroxytoluene, or reducing agents such as ascorbic acid and sodium metabisulfite, and antioxidant synergists such as citric acid, tartaric acid, and lecithin.

The application of emulsion formulations via dermatological, oral and parenteral routes and methods for their manufacture have been reviewed in the literature (Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Emulsion formulations for oral delivery have been very widely used because of ease of formulation, as well as efficacy from an absorption and bioavailability standpoint (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Idson, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 199). Mineral-oil base laxatives, oil-soluble vitamins and high fat nutritive preparations are among the materials that have commonly been administered orally as o/w emulsions.

In one embodiment of the present invention, the compositions of oligonucleotides and nucleic acids are formulated as microemulsions. A microemulsion may be defined as a system of water, oil and amphiphile which is a single optically isotropic and thermodynamically stable liquid solution (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Typically microemulsions are systems that are prepared by first dispersing an oil in an aqueous surfactant solution and then adding a sufficient amount of a fourth component, generally an intermediate chain-length alcohol to form a transparent system. Therefore, microemulsions have also been described as thermodynamically stable, isotropically clear dispersions of two immiscible liquids that are stabilized by interfacial films of surface-active molecules (Leung and Shah, in: *Controlled Release of Drugs: Polymers and Aggregate Systems*, Rosoff, M., Ed., 1989, VCH Publishers, New York, pages 185-215). Microemulsions commonly are prepared via a combination of three to five components that include oil, water, surfactant, cosurfactant and electrolyte. Whether the microemulsion is of the water-in-oil (w/o) or an oil-in-water (o/w) type is dependent on the properties of the oil and surfactant used and on the structure and geometric packing of the polar heads and hydrocarbon tails of the surfactant molecules (Schott, in *Remington's Pharmaceutical Sciences*, Mack Publishing Co., Easton, PA, 1985, p. 271).

The phenomenological approach utilizing phase diagrams has been extensively studied and has yielded a comprehensive knowledge, to one skilled in the art, of how to formulate microemulsions (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245; Block, in

*Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 335). Compared to conventional emulsions, microemulsions offer the advantage of solubilizing water-insoluble drugs in a formulation of thermodynamically stable droplets that are formed spontaneously.

Surfactants used in the preparation of microemulsions include, but are not limited to, ionic surfactants, non-ionic surfactants, Brij 96, polyoxyethylene oleyl ethers, polyglycerol fatty acid esters, tetraglycerol monolaurate (ML310), tetraglycerol monooleate (MO310), hexaglycerol monooleate (PO310), hexaglycerol pentaoleate (PO500), decaglycerol monocaprate (MCA750), decaglycerol monooleate (MO750), decaglycerol sequioleate (SO750), decaglycerol decaoleate (DAO750), alone or in combination with cosurfactants. The cosurfactant, usually a short-chain alcohol such as ethanol, 1-propanol, and 1-butanol, serves to increase the interfacial fluidity by penetrating into the surfactant film and consequently creating a disordered film because of the void space generated among surfactant molecules. Microemulsions may, however, be prepared without the use of cosurfactants and alcohol-free self-emulsifying microemulsion systems are known in the art. The aqueous phase may typically be, but is not limited to, water, an aqueous solution of the drug, glycerol, PEG300, PEG400, polyglycerols, propylene glycols, and derivatives of ethylene glycol. The oil phase may include, but is not limited to, materials such as Captex 300, Captex 355, Capmul MCM, fatty acid esters, medium chain (C8-C12) mono, di, and tri-glycerides, polyoxyethylated glyceryl fatty acid esters, fatty alcohols, polyglycolized glycerides, saturated polyglycolized C8-C10 glycerides, vegetable oils and silicone oil.

Microemulsions are particularly of interest from the standpoint of drug solubilization and the enhanced absorption of drugs. Lipid based microemulsions (both o/w and w/o) have been proposed to enhance the oral bioavailability of drugs, including peptides (Constantinides *et al.*, *Pharmaceutical Research*, 1994, 11, 1385-1390; Ritschel, *Meth. Find. Exp. Clin. Pharmacol.*, 1993, 13, 205). Microemulsions afford advantages of improved drug solubilization, protection of drug from enzymatic hydrolysis, possible enhancement of drug absorption due to surfactant-induced alterations in membrane fluidity and permeability, ease of preparation, ease of oral administration over solid dosage forms, improved clinical potency, and decreased toxicity (Constantinides *et al.*, *Pharmaceutical Research*, 1994, 11, 1385; Ho *et al.*, *J. Pharm. Sci.*, 1996, 85, 138-143). Often microemulsions may form spontaneously when their components are brought together at ambient temperature. This may be particularly advantageous when formulating thermolabile drugs, peptides or oligonucleotides. Microemulsions have also been effective in the transdermal delivery of active components in both cosmetic and pharmaceutical applications. It is expected that the microemulsion compositions and formulations of the present invention will facilitate the increased systemic absorption of oligonucleotides and nucleic acids from the gastrointestinal tract, as well as improve the local cellular uptake of oligonucleotides and nucleic acids within the gastrointestinal tract, vagina, buccal cavity and other areas of administration.

Microemulsions of the present invention may also contain additional components and additives such as sorbitan monostearate (Grill 3), Labrasol, and penetration enhancers to improve the properties of the formulation and to enhance the absorption of the oligonucleotides and nucleic acids of the

present invention. Penetration enhancers used in the microemulsions of the present invention may be classified as belonging to one of five broad categories - surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-  
5 surfactants (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p. 92). Each of these classes has been discussed above.

#### Liposomes

10 There are many organized surfactant structures besides microemulsions that have been studied and used for the formulation of drugs. These include monolayers, micelles, bilayers and vesicles. Vesicles, such as liposomes, have attracted great interest because of their specificity and the  
15 duration of action they offer from the standpoint of drug delivery. As used in the present invention, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers.

Liposomes are unilamellar or multilamellar vesicles which  
20 have a membrane formed from a lipophilic material and an aqueous interior. The aqueous portion contains the composition to be delivered. Cationic liposomes possess the advantage of being able to fuse to the cell wall. Non-cationic liposomes, although not able to fuse as efficiently  
25 with the cell wall, are taken up by macrophages *in vivo*.

In order to cross intact mammalian skin, lipid vesicles must pass through a series of fine pores, each with a diameter less than 50 nm, under the influence of a suitable transdermal gradient. Therefore, it is desirable to use a liposome which  
30 is highly deformable and able to pass through such fine pores.

Further advantages of liposomes include; liposomes obtained from natural phospholipids are biocompatible and biodegradable; liposomes can incorporate a wide range of water

and lipid soluble drugs; liposomes can protect encapsulated drugs in their internal compartments from metabolism and degradation (Rosoff, in *Pharmaceutical Dosage Forms*, Lieberman, Rieger and Banker (Eds.), 1988, Marcel Dekker, Inc., New York, N.Y., volume 1, p. 245). Important  
5 considerations in the preparation of liposome formulations are the lipid surface charge, vesicle size and the aqueous volume of the liposomes.

Liposomes are useful for the transfer and delivery of  
10 active ingredients to the site of action. Because the liposomal membrane is structurally similar to biological membranes, when liposomes are applied to a tissue, the liposomes start to merge with the cellular membranes and as the merging of the liposome and cell progresses, the liposomal  
15 contents are emptied into the cell where the active agent may act.

Liposomal formulations have been the focus of extensive investigation as the mode of delivery for many drugs. There is growing evidence that for topical administration, liposomes  
20 present several advantages over other formulations. Such advantages include reduced side-effects related to high systemic absorption of the administered drug, increased accumulation of the administered drug at the desired target, and the ability to administer a wide variety of drugs, both  
25 hydrophilic and hydrophobic, into the skin.

Several reports have detailed the ability of liposomes to deliver agents including high-molecular weight DNA into the skin. Compounds including analgesics, antibodies, hormones and high-molecular weight DNAs have been administered to the  
30 skin. The majority of applications resulted in the targeting of the upper epidermis.

Liposomes fall into two broad classes. Cationic liposomes are positively charged liposomes which interact with

the negatively charged DNA molecules to form a stable complex. The positively charged DNA/liposome complex binds to the negatively charged cell surface and is internalized in an endosome. Due to the acidic pH within the endosome, the liposomes are ruptured, releasing their contents into the cell cytoplasm (Wang *et al.*, *Biochem. Biophys. Res. Commun.*, **1987**, 147, 980-985).

Liposomes which are pH-sensitive or negatively-charged, entrap DNA rather than complex with it. Since both the DNA and the lipid are similarly charged, repulsion rather than complex formation occurs. Nevertheless, some DNA is entrapped within the aqueous interior of these liposomes. pH-sensitive liposomes have been used to deliver DNA encoding the thymidine kinase gene to cell monolayers in culture. Expression of the exogenous gene was detected in the target cells (Zhou *et al.*, *Journal of Controlled Release*, **1992**, 19, 269-274).

One major type of liposomal composition includes phospholipids other than naturally-derived phosphatidylcholine. Neutral liposome compositions, for example, can be formed from dimyristoyl phosphatidylcholine (DMPC) or dipalmitoyl phosphatidylcholine (DPPC). Anionic liposome compositions generally are formed from dimyristoyl phosphatidylglycerol, while anionic fusogenic liposomes are formed primarily from dioleoyl phosphatidylethanolamine (DOPE). Another type of liposomal composition is formed from phosphatidylcholine (PC) such as, for example, soybean PC, and egg PC. Another type is formed from mixtures of phospholipid and/or phosphatidylcholine and/or cholesterol.

Several studies have assessed the topical delivery of liposomal drug formulations to the skin. Application of liposomes containing interferon to guinea pig skin resulted in a reduction of skin herpes sores while delivery of interferon via other means (*e.g.* as a solution or as an emulsion) were

ineffective (Weiner *et al.*, *Journal of Drug Targeting*, 1992, 2, 405-410). Further, an additional study tested the efficacy of interferon administered as part of a liposomal formulation to the administration of interferon using an aqueous system, and concluded that the liposomal formulation was superior to aqueous administration (du Plessis *et al.*, *Antiviral Research*, 1992, 18, 259-265).

Non-ionic liposomal systems have also been examined to determine their utility in the delivery of drugs to the skin, in particular systems comprising non-ionic surfactant and cholesterol. Non-ionic liposomal formulations comprising Novasome™ I (glyceryl dilaurate/cholesterol/polyoxyethylene-10-stearyl ether) and Novasome™ II (glyceryl distearate/cholesterol/polyoxyethylene-10-stearyl ether) were used to deliver cyclosporin-A into the dermis of mouse skin. Results indicated that such non-ionic liposomal systems were effective in facilitating the deposition of cyclosporin-A into different layers of the skin (Hu *et al.* *S.T.P.Pharma. Sci.*, 1994, 4, 6, 466).

Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids that, when incorporated into liposomes, result in enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome (A) comprises one or more glycolipids, such as monosialoganglioside G<sub>M1</sub>, or (B) is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. While not wishing to be bound by any particular theory, it is thought in the art that, at least for sterically stabilized liposomes containing gangliosides, sphingomyelin, or PEG-derivatized lipids, the

enhanced circulation half-life of these sterically stabilized liposomes derives from a reduced uptake into cells of the reticuloendothelial system (RES) (Allen *et al.*, *FEBS Letters*, 1987, 223, 42; Wu *et al.*, *Cancer Research*, 1993, 53, 3765).

5 Various liposomes comprising one or more glycolipids are known in the art. Papahadjopoulos *et al.* (*Ann. N.Y. Acad. Sci.*, 1987, 507, 64) reported the ability of monosialoganglioside G<sub>M1</sub>, galactocerebroside sulfate and phosphatidylinositol to improve blood half-lives of liposomes. 10 These findings were expounded upon by Gabizon *et al.* (*Proc. Natl. Acad. Sci. U.S.A.*, 1988, 85, 6949). U.S. Patent No. 4,837,028 and WO 88/04924, both to Allen *et al.*, disclose liposomes comprising (1) sphingomyelin and (2) the ganglioside G<sub>M1</sub> or a galactocerebroside sulfate ester. U.S. Patent No. 15 5,543,152 (Webb *et al.*) discloses liposomes comprising sphingomyelin. Liposomes comprising 1,2-*sn*-dimyristoylphosphatidylcholine are disclosed in WO 97/13499 (Lim *et al.*).

Many liposomes comprising lipids derivatized with one or 20 more hydrophilic polymers, and methods of preparation thereof, are known in the art. Sunamoto *et al.* (*Bull. Chem. Soc. Jpn.*, 1980, 53, 2778) described liposomes comprising a nonionic detergent, 2C<sub>12</sub>15G, that contains a PEG moiety. Illum *et al.* (*FEBS Lett.*, 1984, 167, 79) noted that hydrophilic coating of 25 polystyrene particles with polymeric glycols results in significantly enhanced blood half-lives. Synthetic phospholipids modified by the attachment of carboxylic groups of polyalkylene glycols (e.g., PEG) are described by Sears (U.S. Patent Nos. 4,426,330 and 4,534,899). Klivanov *et al.* 30 (*FEBS Lett.*, 1990, 268, 235) described experiments demonstrating that liposomes comprising phosphatidylethanolamine (PE) derivatized with PEG or PEG stearate have significant increases in blood circulation half-

lives. Blume *et al.* (*Biochimica et Biophysica Acta*, 1990, 1029, 91) extended such observations to other PEG-derivatized phospholipids, e.g., DSPE-PEG, formed from the combination of distearoylphosphatidylethanolamine (DSPE) and PEG. Liposomes  
5 having covalently bound PEG moieties on their external surface are described in European Patent No. EP 0 445 131 B1 and WO 90/04384 to Fisher. Liposome compositions containing 1-20 mole percent of PE derivatized with PEG, and methods of use thereof, are described by Woodle *et al.* (U.S. Patent Nos.  
10 5,013,556 and 5,356,633) and Martin *et al.* (U.S. Patent No. 5,213,804 and European Patent No. EP 0 496 813 B1). Liposomes comprising a number of other lipid-polymer conjugates are disclosed in WO 91/05545 and U.S. Patent No. 5,225,212 (both to Martin *et al.*) and in WO 94/20073 (Zalipsky *et al.*)  
15 Liposomes comprising PEG-modified ceramide lipids are described in WO 96/10391 (Choi *et al.*). U.S. Patent Nos. 5,540,935 (Miyazaki *et al.*) and 5,556,948 (Tagawa *et al.*) describe PEG-containing liposomes that can be further derivatized with functional moieties on their surfaces.

20 A limited number of liposomes comprising nucleic acids are known in the art. WO 96/40062 to Thierry *et al.* discloses methods for encapsulating high molecular weight nucleic acids in liposomes. U.S. Patent No. 5,264,221 to Tagawa *et al.* discloses protein-bonded liposomes and asserts that the  
25 contents of such liposomes may include an antisense RNA. U.S. Patent No. 5,665,710 to Rahman *et al.* describes certain methods of encapsulating oligodeoxynucleotides in liposomes. WO 97/04787 to Love *et al.* discloses liposomes comprising antisense oligonucleotides targeted to the raf gene.

30 Transfersomes are yet another type of liposomes, and are highly deformable lipid aggregates which are attractive candidates for drug delivery vehicles. Transfersomes may be described as lipid droplets which are so highly deformable

that they are easily able to penetrate through pores which are smaller than the droplet. Transfersomes are adaptable to the environment in which they are used, e.g. they are self-optimizing (adaptive to the shape of pores in the skin), self-repairing, frequently reach their targets without fragmenting, and often self-loading. To make transfersomes it is possible to add surface edge-activators, usually surfactants, to a standard liposomal composition. Transfersomes have been used to deliver serum albumin to the skin. The transfersome-mediated delivery of serum albumin has been shown to be as effective as subcutaneous injection of a solution containing serum albumin.

Surfactants find wide application in formulations such as emulsions (including microemulsions) and liposomes. The most common way of classifying and ranking the properties of the many different types of surfactants, both natural and synthetic, is by the use of the hydrophile/lipophile balance (HLB). The nature of the hydrophilic group (also known as the "head") provides the most useful means for categorizing the different surfactants used in formulations (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

If the surfactant molecule is not ionized, it is classified as a nonionic surfactant. Nonionic surfactants find wide application in pharmaceutical and cosmetic products and are usable over a wide range of pH values. In general their HLB values range from 2 to about 18 depending on their structure. Nonionic surfactants include nonionic esters such as ethylene glycol esters, propylene glycol esters, glyceryl esters, polyglyceryl esters, sorbitan esters, sucrose esters, and ethoxylated esters. Nonionic alkanolamides and ethers such as fatty alcohol ethoxylates, propoxylated alcohols, and ethoxylated/propoxylated block polymers are also included in

this class. The polyoxyethylene surfactants are the most popular members of the nonionic surfactant class.

If the surfactant molecule carries a negative charge when it is dissolved or dispersed in water, the surfactant is classified as anionic. Anionic surfactants include  
5 carboxylates such as soaps, acyl lactylates, acyl amides of amino acids, esters of sulfuric acid such as alkyl sulfates and ethoxylated alkyl sulfates, sulfonates such as alkyl benzene sulfonates, acyl isethionates, acyl taurates and  
10 sulfosuccinates, and phosphates. The most important members of the anionic surfactant class are the alkyl sulfates and the soaps.

If the surfactant molecule carries a positive charge when it is dissolved or dispersed in water, the surfactant is  
15 classified as cationic. Cationic surfactants include quaternary ammonium salts and ethoxylated amines. The quaternary ammonium salts are the most used members of this class.

If the surfactant molecule has the ability to carry  
20 either a positive or negative charge, the surfactant is classified as amphoteric. Amphoteric surfactants include acrylic acid derivatives, substituted alkylamides, N-alkylbetaines and phosphatides.

The use of surfactants in drug products, formulations and  
25 in emulsions has been reviewed (Rieger, in *Pharmaceutical Dosage Forms*, Marcel Dekker, Inc., New York, NY, 1988, p. 285).

#### Penetration Enhancers

30 In one embodiment, the present invention employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides, to the skin of animals. Most drugs are present in solution in both ionized

and nonionized forms. However, usually only lipid soluble or lipophilic drugs readily cross cell membranes. It has been discovered that even non-lipophilic drugs may cross cell membranes if the membrane to be crossed is treated with a penetration enhancer. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs.

Penetration enhancers may be classified as belonging to one of five broad categories, *i.e.*, surfactants, fatty acids, bile salts, chelating agents, and non-chelating non-surfactants (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92). Each of the above mentioned classes of penetration enhancers are described below in greater detail.

Surfactants: In connection with the present invention, surfactants (or "surface-active agents") are chemical entities which, when dissolved in an aqueous solution, reduce the surface tension of the solution or the interfacial tension between the aqueous solution and another liquid, with the result that absorption of oligonucleotides through the mucosa is enhanced. In addition to bile salts and fatty acids, these penetration enhancers include, for example, sodium lauryl sulfate, polyoxyethylene-9-lauryl ether and polyoxyethylene-20-cetyl ether) (Lee *et al.*, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92); and perfluorochemical emulsions, such as FC-43. Takahashi *et al.*, *J. Pharm. Pharmacol.*, 1988, 40, 252).

Fatty acids: Various fatty acids and their derivatives which act as penetration enhancers include, for example, oleic acid, lauric acid, capric acid (n-decanoic acid), myristic acid, palmitic acid, stearic acid, linoleic acid, linolenic acid, dicaprinate, tricaprinate, monoolein (1-monooleoyl-*rac*-

glycerol), dilaurin, caprylic acid, arachidonic acid, glycerol 1-monocaprato, 1-dodecylazacycloheptan-2-one, acylcarnitines, acylcholines, C<sub>1-10</sub> alkyl esters thereof (e.g., methyl, isopropyl and t-butyl), and mono- and di-glycerides thereof  
5 (i.e., oleate, laurate, caprate, myristate, palmitate, stearate, linoleate, etc.) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, p.92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; El Hariri et al., *J. Pharm. Pharmacol.*, 1992, 44, 651-  
10 654).

Bile salts: The physiological role of bile includes the facilitation of dispersion and absorption of lipids and fat-soluble vitamins (Brunton, Chapter 38 in: Goodman & Gilman's  
15 *The Pharmacological Basis of Therapeutics*, 9th Ed., Hardman et al. Eds., McGraw-Hill, New York, 1996, pp. 934-935). Various natural bile salts, and their synthetic derivatives, act as penetration enhancers. Thus the term "bile salts" includes any of the naturally occurring components of bile as well as  
20 any of their synthetic derivatives. The bile salts of the invention include, for example, cholic acid (or its pharmaceutically acceptable sodium salt, sodium cholate), dehydrocholic acid (sodium dehydrocholate), deoxycholic acid (sodium deoxycholate), glucholic acid (sodium glucholate),  
25 glycholic acid (sodium glycocholate), glycodeoxycholic acid (sodium glycodeoxycholate), taurocholic acid (sodium taurocholate), taurodeoxycholic acid (sodium taurodeoxycholate), chenodeoxycholic acid (sodium chenodeoxycholate), ursodeoxycholic acid (UDCA), sodium tauro-  
30 24,25-dihydro-fusidate (STDHF), sodium glycodihydrofusidate and polyoxyethylene-9-lauryl ether (POE) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Swinyard, Chapter 39 In: *Remington's Pharmaceutical Sciences*,

18th Ed., Gennaro, ed., Mack Publishing Co., Easton, PA, 1990, pages 782-783; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Yamamoto et al., *J. Pharm. Exp. Ther.*, 1992, 263, 25; Yamashita et al., *J. Pharm. Sci.*, 1990, 79, 579-583).

Chelating Agents: Chelating agents, as used in connection with the present invention, can be defined as compounds that remove metallic ions from solution by forming complexes therewith, with the result that absorption of oligonucleotides through the mucosa is enhanced. With regards to their use as penetration enhancers in the present invention, chelating agents have the added advantage of also serving as DNase inhibitors, as most characterized DNA nucleases require a divalent metal ion for catalysis and are thus inhibited by chelating agents (Jarrett, *J. Chromatogr.*, 1993, 618, 315-339). Chelating agents of the invention include but are not limited to disodium ethylenediaminetetraacetate (EDTA), citric acid, salicylates (e.g., sodium salicylate, 5-methoxysalicylate and homovanilate), *N*-acyl derivatives of collagen, laureth-9 and *N*-amino acyl derivatives of beta-diketones (enamines) (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92; Muranishi, *Critical Reviews in Therapeutic Drug Carrier Systems*, 1990, 7, 1-33; Buur et al., *J. Control Rel.*, 1990, 14, 43-51).

Non-chelating non-surfactants: As used herein, non-chelating non-surfactant penetration enhancing compounds can be defined as compounds that demonstrate insignificant activity as chelating agents or as surfactants but that nonetheless enhance absorption of oligonucleotides through the alimentary mucosa (Muranishi, *Critical Reviews in Therapeutic*

*Drug Carrier Systems*, 1990, 7, 1-33). This class of penetration enhancers include, for example, unsaturated cyclic ureas, 1-alkyl- and 1-alkenylazacyclo-alkanone derivatives (Lee et al., *Critical Reviews in Therapeutic Drug Carrier Systems*, 1991, page 92); and non-steroidal anti-inflammatory agents such as diclofenac sodium, indomethacin and phenylbutazone (Yamashita et al., *J. Pharm. Pharmacol.*, 1987, 39, 621-626).

Agents that enhance uptake of oligonucleotides at the cellular level may also be added to the pharmaceutical and other compositions of the present invention. For example, cationic lipids, such as lipofectin (Junichi et al, U.S. Patent No. 5,705,188), cationic glycerol derivatives, and polycationic molecules, such as polylysine (Lollo et al., PCT Application WO 97/30731), are also known to enhance the cellular uptake of oligonucleotides.

Other agents may be utilized to enhance the penetration of the administered nucleic acids, including glycols such as ethylene glycol and propylene glycol, pyrrols such as 2-pyrrol, azones, and terpenes such as limonene and menthone.

#### Carriers

Certain compositions of the present invention also incorporate carrier compounds in the formulation. As used herein, "carrier compound" or "carrier" can refer to a nucleic acid, or analog thereof, which is inert (*i.e.*, does not possess biological activity *per se*) but is recognized as a nucleic acid by *in vivo* processes that reduce the bioavailability of a nucleic acid having biological activity by, for example, degrading the biologically active nucleic acid or promoting its removal from circulation. The coadministration of a nucleic acid and a carrier compound, typically with an excess of the latter substance, can result

in a substantial reduction of the amount of nucleic acid recovered in the liver, kidney or other extracirculatory reservoirs, presumably due to competition between the carrier compound and the nucleic acid for a common receptor. For  
5 example, the recovery of a partially phosphorothioate oligonucleotide in hepatic tissue can be reduced when it is coadministered with polyinosinic acid, dextran sulfate, polycytidic acid or 4-acetamido-4'isothiocyano-stilbene-2,2'-disulfonic acid (Miyao *et al.*, *Antisense Res. Dev.*, 1995, 5,  
10 115-121; Takakura *et al.*, *Antisense & Nucl. Acid Drug Dev.*, 1996, 6, 177-183).

#### Excipients

In contrast to a carrier compound, a "pharmaceutical  
15 carrier" or "excipient" is a pharmaceutically acceptable solvent, suspending agent or any other pharmacologically inert vehicle for delivering one or more nucleic acids to an animal. The excipient may be liquid or solid and is selected, with the planned manner of administration in mind, so as to provide for  
20 the desired bulk, consistency, *etc.*, when combined with a nucleic acid and the other components of a given pharmaceutical composition. Typical pharmaceutical carriers include, but are not limited to, binding agents (*e.g.*, pregelatinized maize starch, polyvinylpyrrolidone or  
25 hydroxypropyl methylcellulose, *etc.*); fillers (*e.g.*, lactose and other sugars, microcrystalline cellulose, pectin, gelatin, calcium sulfate, ethyl cellulose, polyacrylates or calcium hydrogen phosphate, *etc.*); lubricants (*e.g.*, magnesium stearate, talc, silica, colloidal silicon dioxide, stearic  
30 acid, metallic stearates, hydrogenated vegetable oils, corn starch, polyethylene glycols, sodium benzoate, sodium acetate, *etc.*); disintegrants (*e.g.*, starch, sodium starch glycolate,

etc.); and wetting agents (e.g., sodium lauryl sulphate, etc.).

Pharmaceutically acceptable organic or inorganic excipient suitable for non-parenteral administration which do not deleteriously react with nucleic acids can also be used to formulate the compositions of the present invention. Suitable pharmaceutically acceptable carriers include, but are not limited to, water, salt solutions, alcohols, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

Formulations for topical administration of nucleic acids may include sterile and non-sterile aqueous solutions, non-aqueous solutions in common solvents such as alcohols, or solutions of the nucleic acids in liquid or solid oil bases. The solutions may also contain buffers, diluents and other suitable additives. Pharmaceutically acceptable organic or inorganic excipients suitable for non-parenteral administration which do not deleteriously react with nucleic acids can be used.

Suitable pharmaceutically acceptable excipients include, but are not limited to, water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose, polyvinylpyrrolidone and the like.

#### Other Components

The compositions of the present invention may additionally contain other adjunct components conventionally found in pharmaceutical compositions, at their art-established usage levels. Thus, for example, the compositions may contain additional, compatible, pharmaceutically-active materials such

as, for example, antipruritics, astringents, local anesthetics or anti-inflammatory agents, or may contain additional materials useful in physically formulating various dosage forms of the compositions of the present invention, such as  
5 dyes, flavoring agents, preservatives, antioxidants, opacifiers, thickening agents and stabilizers. However, such materials, when added, should not unduly interfere with the biological activities of the components of the compositions of the present invention. The formulations can be sterilized  
10 and, if desired, mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously interact with the nucleic acid(s)  
15 of the formulation.

Aqueous suspensions may contain substances which increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

20 Certain embodiments of the invention provide pharmaceutical compositions containing (a) one or more antisense compounds and (b) one or more other chemotherapeutic agents which function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to  
25 daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bis-chloroethylnitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine,  
30 procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine,

5-azacytidine, hydroxyurea, deoxycoformycin, 4-hydroxyperoxycyclophosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16),  
5 trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and diethylstilbestrol (DES). See, generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed. 1987, pp. 1206-1228, Berkow et al., eds., Rahway, N.J. When used with the compounds of the invention, such chemotherapeutic agents  
10 may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and  
15 oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the invention. See,  
20 generally, *The Merck Manual of Diagnosis and Therapy*, 15th Ed., Berkow et al., eds., 1987, Rahway, N.J., pages 2499-2506 and 46-49, respectively). Other non-antisense chemotherapeutic agents are also within the scope of this invention. Two or more combined compounds may be used  
25 together or sequentially.

In another related embodiment, compositions of the invention may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted  
30 to a second nucleic acid target. Numerous examples of antisense compounds are known in the art. Two or more combined compounds may be used together or sequentially.

The formulation of therapeutic compositions and their

subsequent administration is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several  
5 months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition  
10 rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC<sub>50</sub>s found to be effective in *in vitro* and *in vivo* animal models. In general, dosage is from 0.01 ug to 100 g per kg of body weight, and may be given once or more  
15 daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be  
20 desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 ug to 100 g per kg of body weight, once or more daily, to once every 20 years.

25 While the present invention has been described with specificity in accordance with certain of its preferred embodiments, the following examples serve only to illustrate the invention and are not intended to limit the same.

**EXAMPLES****Example 1****5 Nucleoside Phosphoramidites for Oligonucleotide Synthesis  
Deoxy and 2'-alkoxy amidites**

2'-Deoxy and 2'-methoxy beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial sources (e.g. Chemgenes, Needham MA or Glen Research, Inc. Sterling VA).

10 Other 2'-O-alkoxy substituted nucleoside amidites are prepared as described in U.S. Patent 5,506,351, herein incorporated by reference. For oligonucleotides synthesized using 2'-alkoxy amidites, optimized synthesis cycles were developed that incorporate multiple steps coupling longer wait times relative  
15 to standard synthesis cycles.

The following abbreviations are used in the text: thin layer chromatography (TLC), melting point (MP), high pressure liquid chromatography (HPLC), Nuclear Magnetic Resonance (NMR), argon (Ar), methanol (MeOH), dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>),  
20 triethylamine (TEA), dimethyl formamide (DMF), ethyl acetate (EtOAc), dimethyl sulfoxide (DMSO), tetrahydrofuran (THF).

Oligonucleotides containing 5-methyl-2'-deoxycytidine (5-Me-dC) nucleotides were synthesized according to published methods (Sanghvi, et. al., *Nucleic Acids Research*, 1993, 21,  
25 3197-3203) using commercially available phosphoramidites (Glen Research, Sterling VA or ChemGenes, Needham MA) or prepared as follows:

**30 Preparation of 5'-O-Dimethoxytrityl-thymidine intermediate for  
5-methyl dC amidite**

To a 50 L glass reactor equipped with air stirrer and Ar gas line was added thymidine (1.00 kg, 4.13 mol) in anhydrous pyridine (6 L) at ambient temperature. Dimethoxytrityl (DMT)

chloride (1.47 kg, 4.34 mol, 1.05 eq) was added as a solid in four portions over 1 h. After 30 min, TLC indicated approx. 95% product, 2% thymidine, 5% DMT reagent and by-products and 2 % 3',5'-bis DMT product ( $R_f$  in EtOAc 0.45, 0.05, 0.98, 0.95 respectively). Saturated sodium bicarbonate (4 L) and  $\text{CH}_2\text{Cl}_2$  were added with stirring (pH of the aqueous layer 7.5). An additional 18 L of water was added, the mixture was stirred, the phases were separated, and the organic layer was transferred to a second 50 L vessel. The aqueous layer was extracted with additional  $\text{CH}_2\text{Cl}_2$  (2 x 2 L). The combined organic layer was washed with water (10 L) and then concentrated in a rotary evaporator to approx. 3.6 kg total weight. This was redissolved in  $\text{CH}_2\text{Cl}_2$  (3.5 L), added to the reactor followed by water (6 L) and hexanes (13 L). The mixture was vigorously stirred and seeded to give a fine white suspended solid starting at the interface. After stirring for 1 h, the suspension was removed by suction through a 1/2" diameter teflon tube into a 20 L suction flask, poured onto a 25 cm Coors Buchner funnel, washed with water (2 x 3 L) and a mixture of hexanes-  $\text{CH}_2\text{Cl}_2$  (4:1, 2x3 L) and allowed to air dry overnight in pans (1" deep). This was further dried in a vacuum oven (75°C, 0.1 mm Hg, 48 h) to a constant weight of 2072 g (93%) of a white solid, (mp 122-124°C). TLC indicated a trace contamination of the bis DMT product. NMR spectroscopy also indicated that 1-2 mole percent pyridine and about 5 mole percent of hexanes was still present.

**Preparation of 5'-O-Dimethoxytrityl-2'-deoxy-5-methylcytidine intermediate for 5-methyl-dC amidite**

To a 50 L Schott glass-lined steel reactor equipped with an electric stirrer, reagent addition pump (connected to an addition funnel), heating/cooling system, internal thermometer and an Ar gas line was added 5'-O-dimethoxytrityl-thymidine

(3.00 kg, 5.51 mol), anhydrous acetonitrile (25 L) and TEA (12.3 L, 88.4 mol, 16 eq). The mixture was chilled with stirring to  $-10^{\circ}\text{C}$  internal temperature (external  $-20^{\circ}\text{C}$ ). Trimethylsilylchloride (2.1 L, 16.5 mol, 3.0 eq) was added  
5 over 30 minutes while maintaining the internal temperature below  $-5^{\circ}\text{C}$ , followed by a wash of anhydrous acetonitrile (1 L). Note: the reaction is mildly exothermic and copious hydrochloric acid fumes form over the course of the addition. The reaction was allowed to warm to  $0^{\circ}\text{C}$  and the reaction  
10 progress was confirmed by TLC (EtOAc-hexanes 4:1;  $R_f$  0.43 to 0.84 of starting material and silyl product, respectively). Upon completion, triazole (3.05 kg, 44 mol, 8.0 eq) was added the reaction was cooled to  $-20^{\circ}\text{C}$  internal temperature (external  $-30^{\circ}\text{C}$ ). Phosphorous oxychloride (1035 mL, 11.1 mol,  
15 2.01 eq) was added over 60 min so as to maintain the temperature between  $-20^{\circ}\text{C}$  and  $-10^{\circ}\text{C}$  during the strongly exothermic process, followed by a wash of anhydrous acetonitrile (1 L). The reaction was warmed to  $0^{\circ}\text{C}$  and stirred for 1 h. TLC indicated a complete conversion to the  
20 triazole product ( $R_f$  0.83 to 0.34 with the product spot glowing in long wavelength UV light). The reaction mixture was a peach-colored thick suspension, which turned darker red upon warming without apparent decomposition. The reaction was cooled to  $-15^{\circ}\text{C}$  internal temperature and water (5 L) was  
25 slowly added at a rate to maintain the temperature below  $+10^{\circ}\text{C}$  in order to quench the reaction and to form a homogenous solution. (Caution: this reaction is initially very strongly exothermic). Approximately one-half of the reaction volume (22 L) was transferred by air pump to another vessel, diluted  
30 with EtOAc (12 L) and extracted with water (2 x 8 L). The combined water layers were back-extracted with EtOAc (6 L). The water layer was discarded and the organic layers were concentrated in a 20 L rotary evaporator to an oily foam. The

foam was coevaporated with anhydrous acetonitrile (4 L) to remove EtOAc. (note: dioxane may be used instead of anhydrous acetonitrile if dried to a hard foam). The second half of the reaction was treated in the same way. Each residue was  
5 dissolved in dioxane (3 L) and concentrated ammonium hydroxide (750 mL) was added. A homogenous solution formed in a few minutes and the reaction was allowed to stand overnight (although the reaction is complete within 1 h).

TLC indicated a complete reaction (product  $R_f$  0.35 in  
10 EtOAc-MeOH 4:1). The reaction solution was concentrated on a rotary evaporator to a dense foam. Each foam was slowly redissolved in warm EtOAc (4 L; 50°C), combined in a 50 L glass reactor vessel, and extracted with water (2 x 4L) to remove the triazole by-product. The water was back-extracted  
15 with EtOAc (2 L). The organic layers were combined and concentrated to about 8 kg total weight, cooled to 0°C and seeded with crystalline product. After 24 hours, the first crop was collected on a 25 cm Coors Buchner funnel and washed repeatedly with EtOAc (3 x 3L) until a white powder was left  
20 and then washed with ethyl ether (2 x 3L). The solid was put in pans (1" deep) and allowed to air dry overnight. The filtrate was concentrated to an oil, then redissolved in EtOAc (2 L), cooled and seeded as before. The second crop was collected and washed as before (with proportional solvents)  
25 and the filtrate was first extracted with water (2 x 1L) and then concentrated to an oil. The residue was dissolved in EtOAc (1 L) and yielded a third crop which was treated as above except that more washing was required to remove a yellow oily layer.

30 After air-drying, the three crops were dried in a vacuum oven (50°C, 0.1 mm Hg, 24 h) to a constant weight (1750, 600 and 200 g, respectively) and combined to afford 2550 g (85%) of a white crystalline product (MP 215-217°C) when TLC and NMR

spectroscopy indicated purity. The mother liquor still contained mostly product (as determined by TLC) and a small amount of triazole (as determined by NMR spectroscopy), bis DMT product and unidentified minor impurities. If desired, 5 the mother liquor can be purified by silica gel chromatography using a gradient of MeOH (0-25%) in EtOAc to further increase the yield.

**Preparation of 5'-O-Dimethoxytrityl-2'-deoxy-N<sup>4</sup>-benzoyl-5-methylcytidine penultimate intermediate for 5-methyl dC amidite**

Crystalline 5'-O-dimethoxytrityl-5-methyl-2'-deoxycytidine (2000 g, 3.68 mol) was dissolved in anhydrous DMF (6.0 kg) at ambient temperature in a 50 L glass reactor vessel equipped with an air stirrer and argon line. Benzoic anhydride (Chem Impex not Aldrich, 874 g, 3.86 mol, 1.05 eq) was added and the reaction was stirred at ambient temperature for 8 h. TLC (CH<sub>2</sub>Cl<sub>2</sub>-EtOAc; CH<sub>2</sub>Cl<sub>2</sub>-EtOAc 4:1; R<sub>f</sub> 0.25) indicated approx. 92% complete reaction. An additional amount 20 of benzoic anhydride (44 g, 0.19 mol) was added. After a total of 18 h, TLC indicated approx. 96% reaction completion. The solution was diluted with EtOAc (20 L), TEA (1020 mL, 7.36 mol, ca 2.0 eq) was added with stirring, and the mixture was extracted with water (15 L, then 2 x 10 L). The aqueous layer 25 was removed (no back-extraction was needed) and the organic layer was concentrated in 2 x 20 L rotary evaporator flasks until a foam began to form. The residues were coevaporated with acetonitrile (1.5 L each) and dried (0.1 mm Hg, 25°C, 24 h) to 2520 g of a dense foam. High pressure liquid 30 chromatography (HPLC) revealed a contamination of 6.3% of N<sup>4</sup>, 3'-O-dibenzoyl product, but very little other impurities.

The product was purified by Biotage column chromatography (5 kg Biotage) prepared with 65:35:1 hexanes-EtOAc-TEA (4L).

The crude product (800 g), dissolved in CH<sub>2</sub>Cl<sub>2</sub> (2 L), was applied to the column. The column was washed with the 65:35:1 solvent mixture (20 kg), then 20:80:1 solvent mixture (10 kg), then 99:1 EtOAc:TEA (17kg). The fractions containing the product were collected, and any fractions containing the product and impurities were retained to be resubjected to column chromatography. The column was re-equilibrated with the original 65:35:1 solvent mixture (17 kg). A second batch of crude product (840 g) was applied to the column as before. The column was washed with the following solvent gradients: 65:35:1 (9 kg), 55:45:1 (20 kg), 20:80:1 (10 kg), and 99:1 EtOAc:TEA (15 kg). The column was reequilibrated as above, and a third batch of the crude product (850 g) plus impure fractions recycled from the two previous columns (28 g) was purified following the procedure for the second batch. The fractions containing pure product combined and concentrated on a 20L rotary evaporator, co-evaporated with acetone and dried (0.1 mm Hg, 48 h, 25°C) to a constant weight of 2023 g (85%) of white foam and 20 g of slightly contaminated product from the third run. HPLC indicated a purity of 99.8% with the balance as the diBenzoyl product.

**[5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-deoxy-N<sup>4</sup>-benzoyl-5-methylcytidin-3'-O-yl]-2-cyanoethyl-N,N-diisopropylphosphoramidite (5-methyl dC amidite)**

5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-deoxy-N<sup>4</sup>-benzoyl-5-methylcytidine (998 g, 1.5 mol) was dissolved in anhydrous DMF (2 L). The solution was co-evaporated with toluene (300 ml) at 50°C under reduced pressure, then cooled to room temperature and 2-cyanoethyl tetraisopropylphosphorodiamidite (680 g, 2.26 mol) and tetrazole (52.5 g, 0.75 mol) were added. The mixture was shaken until all tetrazole was dissolved, N-methylimidazole (15 ml) was added and the mixture was left at

room temperature for 5 hours. TEA (300 ml) was added, the mixture was diluted with DMF (2.5 L) and water (600 ml), and extracted with hexane (3 x 3 L). The mixture was diluted with water (1.2 L) and extracted with a mixture of toluene (7.5 L) and hexane (6 L). The two layers were separated, the upper layer was washed with DMF-water (7:3 v/v, 3 x 2 L) and water (3 x 2 L), and the phases were separated. The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and rotary evaporated. The residue was co-evaporated with acetonitrile (2 x 2 L) under reduced pressure and dried to a constant weight (25 °C, 0.1mm Hg, 40 h) to afford 1250 g an off-white foam solid (96%).

## 2'-Fluoro amidites

### 2'-Fluorodeoxyadenosine amidites

2'-fluoro oligonucleotides were synthesized as described previously [Kawasaki, et. al., *J. Med. Chem.*, 1993, 36, 831-841] and United States patent 5,670,633, herein incorporated by reference. The preparation of 2'-fluoropyrimidines containing a 5-methyl substitution are described in US Patent 5,861,493. Briefly, the protected nucleoside N6-benzoyl-2'-deoxy-2'-fluoroadenosine was synthesized utilizing commercially available 9-beta-D-arabinofuranosyladenine as starting material and whereby the 2'-alpha-fluoro atom is introduced by a S<sub>N</sub>2-displacement of a 2'-beta-triflate group. Thus N6-benzoyl-9-beta-D-arabinofuranosyladenine was selectively protected in moderate yield as the 3',5'-ditetrahydropyranyl (THP) intermediate. Deprotection of the THP and N6-benzoyl groups was accomplished using standard methodologies to obtain the 5'-dimethoxytrityl-(DMT) and 5'-DMT-3'-phosphoramidite intermediates.

### 2'-Fluorodeoxyguanosine

The synthesis of 2'-deoxy-2'-fluoroguanosine was

accomplished using tetraisopropylidisiloxanyl (TPDS) protected 9-beta-D-arabinofuranosylguanine as starting material, and conversion to the intermediate isobutyryl-arabinofuranosylguanosine. Alternatively, isobutyryl-arabinofuranosylguanosine was prepared as described by Ross et al., (Nucleosides & Nucleosides, 16, 1645, 1997). Deprotection of the TPDS group was followed by protection of the hydroxyl group with THP to give isobutyryl di-THP protected arabinofuranosylguanine. Selective O-deacylation and triflation was followed by treatment of the crude product with fluoride, then deprotection of the THP groups. Standard methodologies were used to obtain the 5'-DMT- and 5'-DMT-3'-phosphoramidites.

#### 15           **2'-Fluorouridine**

Synthesis of 2'-deoxy-2'-fluorouridine was accomplished by the modification of a literature procedure in which 2,2'-anhydro-1-beta-D-arabinofuranosyluracil was treated with 70% hydrogen fluoride-pyridine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-Fluorodeoxycytidine**

2'-deoxy-2'-fluorocytidine was synthesized via amination of 2'-deoxy-2'-fluorouridine, followed by selective protection to give N4-benzoyl-2'-deoxy-2'-fluorocytidine. Standard procedures were used to obtain the 5'-DMT and 5'-DMT-3'phosphoramidites.

#### **2'-O-(2-Methoxyethyl) modified amidites**

30           2'-O-Methoxyethyl-substituted nucleoside amidites (otherwise known as MOE amidites) are prepared as follows, or alternatively, as per the methods of Martin, P., (Helvetica Chimica Acta, 1995, 78, 486-504).

**Preparation of 2'-O-(2-methoxyethyl)-5-methyluridine  
intermediate**

2,2'-Anhydro-5-methyl-uridine (2000 g, 8.32 mol), tris(2-  
5 methoxyethyl)borate (2504 g, 10.60 mol), sodium bicarbonate  
(60 g, 0.70 mol) and anhydrous 2-methoxyethanol (5 L) were  
combined in a 12 L three necked flask and heated to 130 °C  
(internal temp) at atmospheric pressure, under an argon  
atmosphere with stirring for 21 h. TLC indicated a complete  
10 reaction. The solvent was removed under reduced pressure  
until a sticky gum formed (50-85°C bath temp and 100-11 mm Hg)  
and the residue was redissolved in water (3 L) and heated to  
boiling for 30 min in order the hydrolyze the borate esters.  
The water was removed under reduced pressure until a foam  
15 began to form and then the process was repeated. HPLC  
indicated about 77% product, 15% dimer (5' of product attached  
to 2' of starting material) and unknown derivatives, and the  
balance was a single unresolved early eluting peak.

The gum was redissolved in brine (3 L), and the flask was  
20 rinsed with additional brine (3 L). The combined aqueous  
solutions were extracted with chloroform (20 L) in a heavier-  
than continuous extractor for 70 h. The chloroform layer was  
concentrated by rotary evaporation in a 20 L flask to a sticky  
foam (2400 g). This was coevaporated with MeOH (400 mL) and  
25 EtOAc (8 L) at 75°C and 0.65 atm until the foam dissolved at  
which point the vacuum was lowered to about 0.5 atm. After  
2.5 L of distillate was collected a precipitate began to form  
and the flask was removed from the rotary evaporator and  
stirred until the suspension reached ambient temperature.  
30 EtOAc (2 L) was added and the slurry was filtered on a 25 cm  
table top Buchner funnel and the product was washed with EtOAc  
(3 x 2 L). The bright white solid was air dried in pans for  
24 h then further dried in a vacuum oven (50°C, 0.1 mm Hg, 24

h) to afford 1649 g of a white crystalline solid (mp 115.5-116.5°C).

The brine layer in the 20 L continuous extractor was further extracted for 72 h with recycled chloroform. The chloroform was concentrated to 120 g of oil and this was combined with the mother liquor from the above filtration (225 g), dissolved in brine (250 mL) and extracted once with chloroform (250 mL). The brine solution was continuously extracted and the product was crystallized as described above to afford an additional 178 g of crystalline product containing about 2% of thymine. The combined yield was 1827 g (69.4%). HPLC indicated about 99.5% purity with the balance being the dimer.

#### 15 **Preparation of 5'-O-DMT-2'-O-(2-methoxyethyl)-5-methyluridine penultimate intermediate**

In a 50 L glass-lined steel reactor, 2'-O-(2-methoxyethyl)-5-methyl-uridine (MOE-T, 1500 g, 4.738 mol), lutidine (1015 g, 9.476 mol) were dissolved in anhydrous acetonitrile (15 L). The solution was stirred rapidly and chilled to -10°C (internal temperature). Dimethoxytriphenylmethyl chloride (1765.7 g, 5.21 mol) was added as a solid in one portion. The reaction was allowed to warm to -2°C over 1 h. (Note: The reaction was monitored closely by TLC (EtOAc) to determine when to stop the reaction so as to not generate the undesired bis-DMT substituted side product). The reaction was allowed to warm from -2 to 3°C over 25 min. then quenched by adding MeOH (300 mL) followed after 10 min by toluene (16 L) and water (16 L). The solution was transferred to a clear 50 L vessel with a bottom outlet, vigorously stirred for 1 minute, and the layers separated. The aqueous layer was removed and the organic layer was washed successively with 10% aqueous citric acid (8 L) and water (12

L). The product was then extracted into the aqueous phase by washing the toluene solution with aqueous sodium hydroxide (0.5N, 16 L and 8 L). The combined aqueous layer was overlaid with toluene (12 L) and solid citric acid (8 moles, 1270 g) was added with vigorous stirring to lower the pH of the aqueous layer to 5.5 and extract the product into the toluene. The organic layer was washed with water (10 L) and TLC of the organic layer indicated a trace of DMT-O-Me, bis DMT and dimer DMT.

The toluene solution was applied to a silica gel column (6 L sintered glass funnel containing approx. 2 kg of silica gel slurried with toluene (2 L) and TEA (25 mL)) and the fractions were eluted with toluene (12 L) and EtOAc (3 x 4 L) using vacuum applied to a filter flask placed below the column. The first EtOAc fraction containing both the desired product and impurities were resubjected to column chromatography as above. The clean fractions were combined, rotary evaporated to a foam, coevaporated with acetonitrile (6 L) and dried in a vacuum oven (0.1 mm Hg, 40 h, 40°C) to afford 2850 g of a white crisp foam. NMR spectroscopy indicated a 0.25 mole % remainder of acetonitrile (calculates to be approx. 47 g) to give a true dry weight of 2803 g (96%). HPLC indicated that the product was 99.41% pure, with the remainder being 0.06 DMT-O-Me, 0.10 unknown, 0.44 bis DMT, and no detectable dimer DMT or 3'-O-DMT.

**Preparation of [5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-5-methyluridin-3'-O-yl]-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE T amidite)**

5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-5-methyluridine (1237 g, 2.0 mol) was dissolved in anhydrous DMF (2.5 L). The solution was co-evaporated with toluene (200 ml) at 50°C under reduced pressure, then cooled

to room temperature and 2-cyanoethyl tetraisopropylphosphorodiamidite (900 g, 3.0 mol) and tetrazole (70 g, 1.0 mol) were added. The mixture was shaken until all tetrazole was dissolved, N-methylimidazole (20 ml) was added and the solution was left at room temperature for 5 hours. TEA (300 ml) was added, the mixture was diluted with DMF (3.5 L) and water (600 ml) and extracted with hexane (3 x 3L). The mixture was diluted with water (1.6 L) and extracted with the mixture of toluene (12 L) and hexanes (9 L). The upper layer was washed with DMF-water (7:3 v/v, 3x3 L) and water (3x3 L). The organic layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and evaporated. The residue was co-evaporated with acetonitrile (2 x 2 L) under reduced pressure and dried in a vacuum oven (25°C, 0.1mm Hg, 40 h) to afford 1526 g of an off-white foamy solid (95%).

**Preparation of 5'-O-Dimethoxytrityl-2'-O-(2-methoxyethyl)-5-methylcytidine intermediate**

To a 50 L Schott glass-lined steel reactor equipped with an electric stirrer, reagent addition pump (connected to an addition funnel), heating/cooling system, internal thermometer and argon gas line was added 5'-O-dimethoxytrityl-2'-O-(2-methoxyethyl)-5-methyl-uridine (2.616 kg, 4.23 mol, purified by base extraction only and no scrub column), anhydrous acetonitrile (20 L), and TEA (9.5 L, 67.7 mol, 16 eq). The mixture was chilled with stirring to -10°C internal temperature (external -20°C). Trimethylsilylchloride (1.60 L, 12.7 mol, 3.0 eq) was added over 30 min. while maintaining the internal temperature below -5°C, followed by a wash of anhydrous acetonitrile (1 L). (Note: the reaction is mildly exothermic and copious hydrochloric acid fumes form over the course of the addition). The reaction was allowed to warm to 0°C and the reaction progress was confirmed by TLC (EtOAc,  $R_f$

0.68 and 0.87 for starting material and silyl product, respectively). Upon completion, triazole (2.34 kg, 33.8 mol, 8.0 eq) was added the reaction was cooled to -20°C internal temperature (external -30°C). Phosphorous oxychloride (793 mL, 8.51 mol, 2.01 eq) was added slowly over 60 min so as to maintain the temperature between -20°C and -10°C (note: strongly exothermic), followed by a wash of anhydrous acetonitrile (1 L). The reaction was warmed to 0°C and stirred for 1 h, at which point it was an off-white thick suspension. TLC indicated a complete conversion to the triazole product (EtOAc,  $R_f$  0.87 to 0.75 with the product spot glowing in long wavelength UV light). The reaction was cooled to -15°C and water (5 L) was slowly added at a rate to maintain the temperature below +10°C in order to quench the reaction and to form a homogenous solution. (Caution: this reaction is initially very strongly exothermic). Approximately one-half of the reaction volume (22 L) was transferred by air pump to another vessel, diluted with EtOAc (12 L) and extracted with water (2 x 8 L). The second half of the reaction was treated in the same way. The combined aqueous layers were back-extracted with EtOAc (8 L) The organic layers were combined and concentrated in a 20 L rotary evaporator to an oily foam. The foam was coevaporated with anhydrous acetonitrile (4 L) to remove EtOAc. (note: dioxane may be used instead of anhydrous acetonitrile if dried to a hard foam). The residue was dissolved in dioxane (2 L) and concentrated ammonium hydroxide (750 mL) was added. A homogenous solution formed in a few minutes and the reaction was allowed to stand overnight

TLC indicated a complete reaction ( $\text{CH}_2\text{Cl}_2$ -acetone-MeOH, 20:5:3,  $R_f$  0.51). The reaction solution was concentrated on a rotary evaporator to a dense foam and slowly redissolved in warm  $\text{CH}_2\text{Cl}_2$  (4 L, 40°C) and transferred to a 20 L glass

extraction vessel equipped with a air-powered stirrer. The organic layer was extracted with water (2 x 6 L) to remove the triazole by-product. (Note: In the first extraction an emulsion formed which took about 2 h to resolve). The water  
5 layer was back-extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 x 2 L), which in turn was washed with water (3 L). The combined organic layer was concentrated in 2 x 20 L flasks to a gum and then recrystallized from EtOAc seeded with crystalline product. After sitting overnight, the first crop was collected on a 25  
10 cm Coors Buchner funnel and washed repeatedly with EtOAc until a white free-flowing powder was left (about 3 x 3 L). The filtrate was concentrated to an oil recrystallized from EtOAc, and collected as above. The solid was air-dried in pans for 48 h, then further dried in a vacuum oven (50°C, 0.1mm Hg, 17  
15 h) to afford 2248 g of a bright white, dense solid (86%). An HPLC analysis indicated both crops to be 99.4% pure and NMR spectroscopy indicated only a faint trace of EtOAc remained.

**Preparation of 5'-O-dimethoxytrityl-2'-O-(2-methoxyethyl)-N4-benzoyl-5-methyl-cytidine penultimate intermediate:**  
20

Crystalline 5'-O-dimethoxytrityl-2'-O-(2-methoxyethyl)-5-methyl-cytidine (1000 g, 1.62 mol) was suspended in anhydrous DMF (3 kg) at ambient temperature and stirred under an Ar atmosphere. Benzoic anhydride (439.3 g, 1.94 mol) was added  
25 in one portion. The solution clarified after 5 hours and was stirred for 16 h. HPLC indicated 0.45% starting material remained (as well as 0.32% N4, 3'-O-bis Benzoyl). An additional amount of benzoic anhydride (6.0 g, 0.0265 mol) was added and after 17 h, HPLC indicated no starting material was  
30 present. TEA (450 mL, 3.24 mol) and toluene (6 L) were added with stirring for 1 minute. The solution was washed with water (4 x 4 L), and brine (2 x 4 L). The organic layer was partially evaporated on a 20 L rotary evaporator to remove 4 L

of toluene and traces of water. HPLC indicated that the bis benzoyl side product was present as a 6% impurity. The residue was diluted with toluene (7 L) and anhydrous DMSO (200 mL, 2.82 mol) and sodium hydride (60% in oil, 70 g, 1.75 mol) was added in one portion with stirring at ambient temperature over 1 h. The reaction was quenched by slowly adding then washing with aqueous citric acid (10%, 100 mL over 10 min, then 2 x 4 L), followed by aqueous sodium bicarbonate (2%, 2 L), water (2 x 4 L) and brine (4 L). The organic layer was concentrated on a 20 L rotary evaporator to about 2 L total volume. The residue was purified by silica gel column chromatography (6 L Buchner funnel containing 1.5 kg of silica gel wetted with a solution of EtOAc-hexanes-TEA(70:29:1)). The product was eluted with the same solvent (30 L) followed by straight EtOAc (6 L). The fractions containing the product were combined, concentrated on a rotary evaporator to a foam and then dried in a vacuum oven (50°C, 0.2 mm Hg, 8 h) to afford 1155 g of a crisp, white foam (98%). HPLC indicated a purity of >99.7%.

20

**Preparation of [5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N<sup>4</sup>-benzoyl-5-methylcytidin-3'-O-yl]-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE 5-Me-C amidite)**

5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N<sup>4</sup>-benzoyl-5-methylcytidine (1082 g, 1.5 mol) was dissolved in anhydrous DMF (2 L) and co-evaporated with toluene (300 ml) at 50 °C under reduced pressure. The mixture was cooled to room temperature and 2-cyanoethyl tetraisopropylphosphorodiamidite (680 g, 2.26 mol) and tetrazole (52.5 g, 0.75 mol) were added. The mixture was shaken until all tetrazole was dissolved, N-methylimidazole (30 ml) was added, and the mixture was left at room temperature for 5 hours. TEA (300 ml) was added, the mixture

was diluted with DMF (1 L) and water (400 ml) and extracted with hexane (3 x 3 L). The mixture was diluted with water (1.2 L) and extracted with a mixture of toluene (9 L) and hexanes (6 L). The two layers were separated and the upper  
5 layer was washed with DMF-water (60:40 v/v, 3 x 3 L) and water (3 x 2 L). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated. The residue was co-evaporated with acetonitrile (2 x 2 L) under reduced pressure and dried in a vacuum oven (25 °C, 0.1mm Hg, 40 h) to afford 1336 g of an off-white foam  
10 (97%).

**Preparation of [5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N<sup>6</sup>-benzoyladenoin-3'-O-yl]-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE A amdite)**

5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-  
15 methoxyethyl)-N<sup>6</sup>-benzoyladenoin (purchased from Reliable Biopharmaceutical, St. Lois, MO), 1098 g, 1.5 mol) was dissolved in anhydrous DMF (3 L) and co-evaporated with toluene (300 ml) at 50 °C. The mixture was cooled to room temperature and 2-cyanoethyl tetraisopropylphosphorodiamidite  
20 (680 g, 2.26 mol) and tetrazole (78.8 g, 1.24 mol) were added. The mixture was shaken until all tetrazole was dissolved, N-methylimidazole (30 ml) was added, and mixture was left at room temperature for 5 hours. TEA (300 ml) was added, the mixture was diluted with DMF (1 L) and water (400 ml) and  
25 extracted with hexanes (3 x 3 L). The mixture was diluted with water (1.4 L) and extracted with the mixture of toluene (9 L) and hexanes (6 L). The two layers were separated and the upper layer was washed with DMF-water (60:40, v/v, 3 x 3 L) and water (3 x 2 L). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>),  
30 filtered and evaporated to a sticky foam. The residue was co-evaporated with acetonitrile (2.5 L) under reduced pressure and dried in a vacuum oven (25 °C, 0.1mm Hg, 40 h) to afford 1350 g of an off-white foam solid (96%).

**Preparation of [5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N<sup>4</sup>-isobutyrylguanosin-3'-O-yl]-2-cyanoethyl-N,N-diisopropylphosphoramidite (MOE G amidite)**

5           5'-O-(4,4'-Dimethoxytriphenylmethyl)-2'-O-(2-methoxyethyl)-N<sup>4</sup>-isobutyrylguanosine (purchased from Reliable Biopharmaceutical, St. Louis, MO, 1426 g, 2.0 mol) was dissolved in anhydrous DMF (2 L). The solution was co-evaporated with toluene (200 ml) at 50 °C, cooled to room  
10 temperature and 2-cyanoethyl tetraisopropylphosphorodiamidite (900 g, 3.0 mol) and tetrazole (68 g, 0.97 mol) were added. The mixture was shaken until all tetrazole was dissolved, N-methylimidazole (30 ml) was added, and the mixture was left at room temperature for 5 hours. TEA (300 ml) was added, the  
15 mixture was diluted with DMF (2 L) and water (600 ml) and extracted with hexanes (3 x 3 L). The mixture was diluted with water (2 L) and extracted with a mixture of toluene (10 L) and hexanes (5 L). The two layers were separated and the upper layer was washed with DMF-water (60:40, v/v, 3x3 L).  
20 EtOAc (4 L) was added and the solution was washed with water (3 x 4 L). The organic layer was dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and evaporated to approx. 4 kg. Hexane (4 L) was added, the mixture was shaken for 10 min, and the supernatant liquid was decanted. The residue was co-evaporated with acetonitrile (2  
25 x 2 L) under reduced pressure and dried in a vacuum oven (25 °C, 0.1mm Hg, 40 h) to afford 1660 g of an off-white foamy solid (91%).

**2'-O-(Aminoxyethyl) nucleoside amidites and 2'-O-(dimethylaminoxyethyl) nucleoside amidites**

2'-(Dimethylaminoxyethoxy) nucleoside amidites

2'-(Dimethylaminoxyethoxy) nucleoside amidites (also known in the art as 2'-O-(dimethylaminoxyethyl) nucleoside

amidites) are prepared as described in the following paragraphs. Adenosine, cytidine and guanosine nucleoside amidites are prepared similarly to the thymidine (5-methyluridine) except the exocyclic amines are protected with a benzoyl moiety in the case of adenosine and cytidine and with isobutyryl in the case of guanosine.

**5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine**  
O<sup>2</sup>-2'-anhydro-5-methyluridine (Pro. Bio. Sint., Varese, Italy, 100.0g, 0.416 mmol), dimethylaminopyridine (0.66g, 0.013eq, 0.0054mmol) were dissolved in dry pyridine (500 ml) at ambient temperature under an argon atmosphere and with mechanical stirring. tert-Butyldiphenylchlorosilane (125.8g, 119.0mL, 1.1eq, 0.458mmol) was added in one portion. The reaction was stirred for 16 h at ambient temperature. TLC (R<sub>f</sub> 0.22, EtOAc) indicated a complete reaction. The solution was concentrated under reduced pressure to a thick oil. This was partitioned between CH<sub>2</sub>Cl<sub>2</sub> (1 L) and saturated sodium bicarbonate (2 x 1 L) and brine (1 L). The organic layer was dried over sodium sulfate, filtered, and concentrated under reduced pressure to a thick oil. The oil was dissolved in a 1:1 mixture of EtOAc and ethyl ether (600mL) and cooling the solution to -10°C afforded a white crystalline solid which was collected by filtration, washed with ethyl ether (3 x 200 mL) and dried (40°C, 1mm Hg, 24 h) to afford 149g of white solid (74.8%). TLC and NMR spectroscopy were consistent with pure product.

**5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine**

In the fume hood, ethylene glycol (350 mL, excess) was added cautiously with manual stirring to a 2 L stainless steel pressure reactor containing borane in tetrahydrofuran (1.0 M,

2.0 eq, 622 mL). (Caution : evolves hydrogen gas). 5'-O-tert-Butyldiphenylsilyl-O<sup>2</sup>-2'-anhydro-5-methyluridine (149 g, 0.311 mol) and sodium bicarbonate (0.074 g, 0.003 eq) were added with manual stirring. The reactor was sealed and heated in an oil bath until an internal temperature of 160 °C was reached and then maintained for 16 h (pressure < 100 psig). The reaction vessel was cooled to ambient temperature and opened. TLC (EtOAc, R<sub>f</sub> 0.67 for desired product and R<sub>f</sub> 0.82 for ara-T side product) indicated about 70% conversion to the product. The solution was concentrated under reduced pressure (10 to 1mm Hg) in a warm water bath (40-100°C) with the more extreme conditions used to remove the ethylene glycol. (Alternatively, once the THF has evaporated the solution can be diluted with water and the product extracted into EtOAc). The residue was purified by column chromatography (2kg silica gel, EtOAc-hexanes gradient 1:1 to 4:1). The appropriate fractions were combined, evaporated and dried to afford 84 g of a white crisp foam (50%), contaminated starting material (17.4g, 12% recovery) and pure reusable starting material (20g, 13% recovery). TLC and NMR spectroscopy were consistent with 99% pure product.

**2'-O-([2-phthalimidoxy)ethyl]-5'-t-butyldiphenylsilyl-5-methyluridine**

5'-O-tert-Butyldiphenylsilyl-2'-O-(2-hydroxyethyl)-5-methyluridine (20g, 36.98mmol) was mixed with triphenylphosphine (11.63g, 44.36mmol) and N-hydroxyphthalimide (7.24g, 44.36mmol) and dried over P<sub>2</sub>O<sub>5</sub> under high vacuum for two days at 40°C. The reaction mixture was flushed with argon and dissolved in dry THF (369.8mL, Aldrich, sure seal bottle). Diethyl-azodicarboxylate (6.98mL, 44.36mmol) was added dropwise to the reaction mixture with the rate of addition maintained such that the resulting deep red

coloration is just discharged before adding the next drop. The reaction mixture was stirred for 4 hrs., after which time TLC (EtOAc:hexane, 60:40) indicated that the reaction was complete. The solvent was evaporated in vacuo and the residue purified by flash column chromatography (eluted with 60:40 EtOAc:hexane), to yield 2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine as white foam (21.819 g, 86%) upon rotary evaporation.

10           **5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine**

2'-O-([2-phthalimidoxy)ethyl]-5'-*t*-butyldiphenylsilyl-5-methyluridine (3.1g, 4.5mmol) was dissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (4.5mL) and methylhydrazine (300mL, 4.64mmol) was added dropwise at -10°C to 0°C. After 1 h the mixture was filtered, the filtrate washed with ice cold CH<sub>2</sub>Cl<sub>2</sub>, and the combined organic phase was washed with water and brine and dried (anhydrous Na<sub>2</sub>SO<sub>4</sub>). The solution was filtered and evaporated to afford 2'-O-(aminooxyethyl) thymidine, which was then dissolved in MeOH (67.5mL). Formaldehyde (20% aqueous solution, w/w, 1.1 eq.) was added and the resulting mixture was stirred for 1 h. The solvent was removed under vacuum and the residue was purified by column chromatography to yield 5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine as white foam (1.95 g, 78%) upon rotary evaporation.

**5'-O-*tert*-Butyldiphenylsilyl-2'-O-[N,N dimethylaminooxyethyl]-5-methyluridine**

30           5'-O-*tert*-butyldiphenylsilyl-2'-O-[(2-formadoximinoxy)ethyl]-5-methyluridine (1.77g, 3.12mmol) was dissolved in a solution of 1M pyridinium *p*-toluenesulfonate

(PPTS) in dry MeOH (30.6mL) and cooled to 10°C under inert atmosphere. Sodium cyanoborohydride (0.39g, 6.13mmol) was added and the reaction mixture was stirred. After 10 minutes the reaction was warmed to room temperature and stirred for 2 h. while the progress of the reaction was monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). Aqueous NaHCO<sub>3</sub> solution (5%, 10mL) was added and the product was extracted with EtOAc (2 x 20 mL). The organic phase was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to dryness. This entire procedure was repeated with the resulting residue, with the exception that formaldehyde (20% w/w, 30 mL, 3.37 mol) was added upon dissolution of the residue in the PPTS/MeOH solution. After the extraction and evaporation, the residue was purified by flash column chromatography and (eluted with 5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine as a white foam (14.6g, 80%) upon rotary evaporation.

#### 2'-O-(dimethylaminoxyethyl)-5-methyluridine

Triethylamine trihydrofluoride (3.91mL, 24.0mmol) was dissolved in dry THF and TEA (1.67mL, 12mmol, dry, stored over KOH) and added to 5'-O-tert-butyldiphenylsilyl-2'-O-[N,N-dimethylaminoxyethyl]-5-methyluridine (1.40g, 2.4mmol). The reaction was stirred at room temperature for 24 hrs and monitored by TLC (5% MeOH in CH<sub>2</sub>Cl<sub>2</sub>). The solvent was removed under vacuum and the residue purified by flash column chromatography (eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub>) to afford 2'-O-(dimethylaminoxyethyl)-5-methyluridine (766mg, 92.5%) upon rotary evaporation of the solvent.

#### 5'-O-DMT-2'-O-(dimethylaminoxyethyl)-5-methyluridine

2'-O-(dimethylaminoxyethyl)-5-methyluridine (750 mg, 2.17 mmol) was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at

40°C, co-evaporated with anhydrous pyridine (20 mL), and dissolved in pyridine (11 mL) under argon atmosphere. 4-dimethylaminopyridine (26.5 mg, 2.60 mmol) and 4,4'-dimethoxytrityl chloride (880 mg, 2.60 mmol) were added to the pyridine solution and the reaction mixture was stirred at room temperature until all of the starting material had reacted. Pyridine was removed under vacuum and the residue was purified by column chromatography (eluted with 10% MeOH in CH<sub>2</sub>Cl<sub>2</sub> containing a few drops of pyridine) to yield 5'-O-DMT-2'-O-(dimethylamino-oxyethyl)-5-methyluridine (1.13g, 80%) upon rotary evaporation.

**5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

5'-O-DMT-2'-O-(dimethylaminooxyethyl)-5-methyluridine (1.08 g, 1.67 mmol) was co-evaporated with toluene (20 mL), N,N-diisopropylamine tetrazonide (0.29 g, 1.67 mmol) was added and the mixture was dried over P<sub>2</sub>O<sub>5</sub> under high vacuum overnight at 40°C. This was dissolved in anhydrous acetonitrile (8.4 mL) and 2-cyanoethyl-N,N,N<sup>1</sup>,N<sup>1</sup>-tetraisopropylphosphoramidite (2.12 mL, 6.08 mmol) was added. The reaction mixture was stirred at ambient temperature for 4 h under inert atmosphere. The progress of the reaction was monitored by TLC (hexane:EtOAc 1:1). The solvent was evaporated, then the residue was dissolved in EtOAc (70mL) and washed with 5% aqueous NaHCO<sub>3</sub> (40mL). The EtOAc layer was dried over anhydrous Na<sub>2</sub>SO<sub>4</sub>, filtered, and concentrated. The residue obtained was purified by column chromatography (EtOAc as eluent) to afford 5'-O-DMT-2'-O-(2-N,N-dimethylaminooxyethyl)-5-methyluridine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite] as a foam (1.04g, 74.9%) upon rotary evaporation.

**2'-(Aminooxyethoxy) nucleoside amidites**

2'-(Aminooxyethoxy) nucleoside amidites (also known in the art as 2'-O-(aminooxyethyl) nucleoside amidites) are prepared as described in the following paragraphs. Adenosine, cytidine and thymidine nucleoside amidites are prepared similarly.

10 **N2-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite]**

The 2'-O-aminooxyethyl guanosine analog may be obtained by selective 2'-O-alkylation of diaminopurine riboside. Multigram quantities of diaminopurine riboside may be purchased from Schering AG (Berlin) to provide 2'-O-(2-ethylacetyl) diaminopurine riboside along with a minor amount of the 3'-O-isomer. 2'-O-(2-ethylacetyl) diaminopurine riboside may be resolved and converted to 2'-O-(2-ethylacetyl)guanosine by treatment with adenosine deaminase. (McGee, D. P. C., Cook, P. D., Guinosso, C. J., WO 94/02501 A1 940203.) Standard protection procedures should afford 2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine and 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-ethylacetyl)-5'-O-(4,4'-dimethoxytrityl)guanosine which may be reduced to provide 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-(2-hydroxyethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine. As before the hydroxyl group may be displaced by N-hydroxyphthalimide via a Mitsunobu reaction, and the protected nucleoside may be phosphitylated as usual to yield 2-N-isobutyryl-6-O-diphenylcarbamoyl-2'-O-([2-phthalamidoxy]ethyl)-5'-O-(4,4'-dimethoxytrityl)guanosine-3'-[(2-cyanoethyl)-N,N-diisopropylphosphoramidite].

**2'-dimethylaminoethoxyethoxy (2'-DMAEOE) nucleoside amidites**

2'-dimethylaminoethoxyethoxy nucleoside amidites (also known in the art as 2'-O-dimethylaminoethoxyethyl, i.e., 2'-O-CH<sub>2</sub>-O-CH<sub>2</sub>-N(CH<sub>2</sub>)<sub>2</sub>, or 2'-DMAEOE nucleoside amidites) are prepared as follows. Other nucleoside amidites are prepared similarly.

**2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

2[2-(Dimethylamino)ethoxy]ethanol (Aldrich, 6.66 g, 50 mmol) was slowly added to a solution of borane in tetrahydrofuran (1 M, 10 mL, 10 mmol) with stirring in a 100 mL bomb. (Caution: Hydrogen gas evolves as the solid dissolves). O<sup>2</sup>-, 2'-anhydro-5-methyluridine (1.2 g, 5 mmol), and sodium bicarbonate (2.5 mg) were added and the bomb was sealed, placed in an oil bath and heated to 155°C for 26 h. then cooled to room temperature. The crude solution was concentrated, the residue was diluted with water (200 mL) and extracted with hexanes (200 mL). The product was extracted from the aqueous layer with EtOAc (3 x 200 mL) and the combined organic layers were washed once with water, dried over anhydrous sodium sulfate, filtered and concentrated. The residue was purified by silica gel column chromatography (eluted with 5:100:2 MeOH/CH<sub>2</sub>Cl<sub>2</sub>/TEA) as the eluent. The appropriate fractions were combined and evaporated to afford the product as a white solid.

**5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine**

To 0.5 g (1.3 mmol) of 2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyl uridine in anhydrous pyridine (8 mL), was added TEA (0.36 mL) and dimethoxytrityl chloride (DMT-Cl, 0.87 g, 2 eq.) and the reaction was stirred for 1 h. The reaction mixture was poured into water (200 mL) and extracted

with CH<sub>2</sub>Cl<sub>2</sub> (2 x 200 mL). The combined CH<sub>2</sub>Cl<sub>2</sub> layers were washed with saturated NaHCO<sub>3</sub> solution, followed by saturated NaCl solution, dried over anhydrous sodium sulfate, filtered and evaporated. The residue was purified by silica gel column chromatography (eluted with 5:100:1 MeOH/CH<sub>2</sub>Cl<sub>2</sub>/TEA) to afford the product.

**5'-O-Dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)-ethyl]-5-methyl uridine-3'-O-(cyanoethyl-N,N-diisopropyl)phosphoramidite**

10

Diisopropylaminotetrazolide (0.6 g) and 2-cyanoethoxy-N,N-diisopropyl phosphoramidite (1.1 mL, 2 eq.) were added to a solution of 5'-O-dimethoxytrityl-2'-O-[2(2-N,N-dimethylaminoethoxy)ethyl]-5-methyluridine (2.17 g, 3 mmol) dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) under an atmosphere of argon. The reaction mixture was stirred overnight and the solvent evaporated. The resulting residue was purified by silica gel column chromatography with EtOAc as the eluent to afford the title compound.

15  
20

**Example 2**

**Oligonucleotide synthesis**

Unsubstituted and substituted phosphodiester (P=O) oligonucleotides are synthesized on an automated DNA synthesizer (Applied Biosystems model 394) using standard phosphoramidite chemistry with oxidation by iodine.

Phosphorothioates (P=S) are synthesized similar to phosphodiester oligonucleotides with the following exceptions: thiation was effected by utilizing a 10% w/v solution of 3H-1,2-benzodithiole-3-one 1,1-dioxide in acetonitrile for the oxidation of the phosphite linkages. The thiation reaction step time was increased to 180 sec and preceded by the normal capping step. After cleavage from the CPG column and

30

deblocking in concentrated ammonium hydroxide at 55°C (12-16 hr), the oligonucleotides were recovered by precipitating with >3 volumes of ethanol from a 1 M NH<sub>4</sub>OAc solution.

Phosphinate oligonucleotides are prepared as described in  
5 U.S. Patent 5,508,270, herein incorporated by reference.

Alkyl phosphonate oligonucleotides are prepared as described in U.S. Patent 4,469,863, herein incorporated by reference.

3'-Deoxy-3'-methylene phosphonate oligonucleotides are  
10 prepared as described in U.S. Patents 5,610,289 or 5,625,050, herein incorporated by reference.

Phosphoramidite oligonucleotides are prepared as described in U.S. Patent, 5,256,775 or U.S. Patent 5,366,878, herein incorporated by reference.

15 Alkylphosphonothioate oligonucleotides are prepared as described in published PCT applications PCT/US94/00902 and PCT/US93/06976 (published as WO 94/17093 and WO 94/02499, respectively), herein incorporated by reference.

3'-Deoxy-3'-amino phosphoramidate oligonucleotides are  
20 prepared as described in U.S. Patent 5,476,925, herein incorporated by reference.

Phosphotriester oligonucleotides are prepared as described in U.S. Patent 5,023,243, herein incorporated by reference.

25 Borano phosphate oligonucleotides are prepared as described in U.S. Patents 5,130,302 and 5,177,198, both herein incorporated by reference.

### Example 3

#### 30 Oligonucleoside Synthesis

Methylenemethylimino linked oligonucleosides, also identified as MMI linked oligonucleosides, methylenedimethylhydrazo linked oligonucleosides, also identified as MDH linked

oligonucleosides, and methylenecarbonylamino linked  
oligonucleosides, also identified as amide-3 linked  
oligonucleosides, and methyleneaminocarbonyl linked oligo-  
nucleosides, also identified as amide-4 linked oligonucleo-  
sides, as well as mixed backbone compounds having, for  
5 instance, alternating MMI and P=O or P=S linkages are prepared  
as described in U.S. Patents 5,378,825, 5,386,023, 5,489,677,  
5,602,240 and 5,610,289, all of which are herein incorporated  
by reference.

10 Formacetal and thioformacetal linked oligonucleosides are  
prepared as described in U.S. Patents 5,264,562 and 5,264,564,  
herein incorporated by reference.

Ethylene oxide linked oligonucleosides are prepared as  
described in U.S. Patent 5,223,618, herein incorporated by  
15 reference.

#### **Example 4**

##### **PNA Synthesis**

Peptide nucleic acids (PNAs) are prepared in accordance  
20 with any of the various procedures referred to in Peptide  
Nucleic Acids (PNA): Synthesis, Properties and Potential  
Applications, *Bioorganic & Medicinal Chemistry*, 1996, 4, 5-23.  
They may also be prepared in accordance with U.S. Patents  
5,539,082, 5,700,922, and 5,719,262, herein incorporated by  
25 reference.

#### **Example 5**

##### **Synthesis of Chimeric Oligonucleotides**

Chimeric oligonucleotides, oligonucleosides or mixed  
30 oligonucleotides/oligonucleosides of the invention can be of  
several different types. These include a first type wherein  
the "gap" segment of linked nucleosides is positioned between  
5' and 3' "wing" segments of linked nucleosides and a second

"open end" type wherein the "gap" segment is located at either the 3' or the 5' terminus of the oligomeric compound.

Oligonucleotides of the first type are also known in the art as "gapmers" or gapped oligonucleotides. Oligonucleotides of the second type are also known in the art as "hemimers" or "wingmers".

**[2'-O-Me]--[2'-deoxy]--[2'-O-Me] Chimeric  
Phosphorothioate Oligonucleotides**

Chimeric oligonucleotides having 2'-O-alkyl phosphorothioate and 2'-deoxy phosphorothioate oligonucleotide segments are synthesized using an Applied Biosystems automated DNA synthesizer Model 394, as above. Oligonucleotides are synthesized using the automated synthesizer and 2'-deoxy-5'-dimethoxytrityl-3'-O-phosphoramidite for the DNA portion and 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite for 5' and 3' wings. The standard synthesis cycle is modified by incorporating coupling steps with increased reaction times for the 5'-dimethoxytrityl-2'-O-methyl-3'-O-phosphoramidite. The fully protected oligonucleotide is cleaved from the support and deprotected in concentrated ammonia (NH<sub>4</sub>OH) for 12-16 hr at 55°C. The deprotected oligo is then recovered by an appropriate method (precipitation, column chromatography, volume reduced *in vacuo* and analyzed spectrophotometrically for yield and for purity by capillary electrophoresis and by mass spectrometry.

**[2'-O-(2-Methoxyethyl)]--[2'-deoxy]--[2'-O-(Methoxyethyl)] Chimeric Phosphorothioate  
Oligonucleotides**

[2'-O-(2-methoxyethyl)]--[2'-deoxy]--[2'-O-(methoxyethyl)] chimeric phosphorothioate oligonucleotides were prepared as per the procedure above for the 2'-O-methyl chimeric oligonucleotide, with the substitution of 2'-O-

(methoxyethyl) amidites for the 2'-O-methyl amidites.

[2'-O-(2-Methoxyethyl)Phosphodiester]--[2'-deoxy  
Phosphorothioate]--[2'-O-(2-Methoxyethyl) Phosphodiester]

5 **Chimeric Oligonucleotides**

[2'-O-(2-methoxyethyl phosphodiester)--[2'-deoxy  
phosphorothioate]--[2'-O-(methoxyethyl) phosphodiester]  
chimeric oligonucleotides are prepared as per the above  
procedure for the 2'-O-methyl chimeric oligonucleotide with  
10 the substitution of 2'-O-(methoxyethyl) amidites for the 2'-O-  
methyl amidites, oxidation with iodine to generate the  
phosphodiester internucleotide linkages within the wing  
portions of the chimeric structures and sulfurization  
utilizing 3,4-dihydro-2H-benzothiole-3-one 1,1-dioxide (Beaucage  
15 Reagent) to generate the phosphorothioate internucleotide  
linkages for the center gap.

Other chimeric oligonucleotides, chimeric  
oligonucleosides and mixed chimeric  
oligonucleotides/oligonucleosides are synthesized according to  
20 United States patent 5,623,065, herein incorporated by  
reference.

**Example 6**

**Oligonucleotide Isolation**

25 After cleavage from the controlled pore glass solid  
support and deblocking in concentrated ammonium hydroxide at  
55°C for 12-16 hours, the oligonucleotides or oligonucleosides  
are recovered by precipitation out of 1 M NH<sub>4</sub>OAc with >3  
volumes of ethanol. Synthesized oligonucleotides were  
30 analyzed by electrospray mass spectroscopy (molecular weight  
determination) and by capillary gel electrophoresis and judged  
to be at least 70% full length material. The relative amounts  
of phosphorothioate and phosphodiester linkages obtained in

the synthesis was determined by the ratio of correct molecular weight relative to the -16 amu product (+/-32 +/-48). For some studies oligonucleotides were purified by HPLC, as described by Chiang *et al.*, *J. Biol. Chem.* **1991**, *266*, 18162-5 18171. Results obtained with HPLC-purified material were similar to those obtained with non-HPLC purified material.

### Example 7

#### Oligonucleotide Synthesis - 96 Well Plate Format

10 Oligonucleotides were synthesized via solid phase P(III) phosphoramidite chemistry on an automated synthesizer capable of assembling 96 sequences simultaneously in a 96-well format. Phosphodiester internucleotide linkages were afforded by oxidation with aqueous iodine. Phosphorothioate  
15 internucleotide linkages were generated by sulfurization utilizing 3,4-dihydro-2H-benzothiazole-3-one 1,1-dioxide (Beaucage Reagent) in anhydrous acetonitrile. Standard base-protected beta-cyanoethyl-diisopropyl phosphoramidites were purchased from commercial vendors (e.g. PE-Applied Biosystems, Foster  
20 City, CA, or Pharmacia, Piscataway, NJ). Non-standard nucleosides are synthesized as per standard or patented methods. They are utilized as base protected beta-cyanoethyl-diisopropyl phosphoramidites.

Oligonucleotides were cleaved from support and  
25 deprotected with concentrated  $\text{NH}_4\text{OH}$  at elevated temperature (55-60°C) for 12-16 hours and the released product then dried *in vacuo*. The dried product was then re-suspended in sterile water to afford a master plate from which all analytical and test plate samples are then diluted utilizing robotic  
30 pipettors.

### Example 8

**Oligonucleotide Analysis - 96-Well Plate Format**

The concentration of oligonucleotide in each well was assessed by dilution of samples and UV absorption spectroscopy. The full-length integrity of the individual products was evaluated by capillary electrophoresis (CE) in either the 96-well format (Beckman P/ACE™ MDQ) or, for individually prepared samples, on a commercial CE apparatus (e.g., Beckman P/ACE™ 5000, ABI 270). Base and backbone composition was confirmed by mass analysis of the compounds utilizing electrospray-mass spectroscopy. All assay test plates were diluted from the master plate using single and multi-channel robotic pipettors. Plates were judged to be acceptable if at least 85% of the compounds on the plate were at least 85% full length.

15

**Example 9****Cell culture and oligonucleotide treatment**

The effect of antisense compounds on target nucleic acid expression can be tested in any of a variety of cell types provided that the target nucleic acid is present at measurable levels. This can be routinely determined using, for example, PCR or Northern blot analysis. The following cell types are provided for illustrative purposes, but other cell types can be routinely used, provided that the target is expressed in the cell type chosen. This can be readily determined by methods routine in the art, for example Northern blot analysis, ribonuclease protection assays, or RT-PCR.

25

T-24 cells:

30

The human transitional cell bladder carcinoma cell line T-24 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). T-24 cells were routinely cultured in

complete McCoy's 5A basal media (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal calf serum (Invitrogen Corporation, Carlsbad, CA), penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Invitrogen Corporation, Carlsbad, CA). Cells were routinely passaged by  
5 trypsinization and dilution when they reached 90% confluence. Cells were seeded into 96-well plates (Falcon-Primaria #3872) at a density of 7000 cells/well for use in RT-PCR analysis.

For Northern blotting or other analysis, cells may be  
10 seeded onto 100 mm or other standard tissue culture plates and treated similarly, using appropriate volumes of medium and oligonucleotide.

#### A549 cells:

15 The human lung carcinoma cell line A549 was obtained from the American Type Culture Collection (ATCC) (Manassas, VA). A549 cells were routinely cultured in DMEM basal media (Invitrogen Corporation, Carlsbad, CA) supplemented with 10% fetal calf serum (Invitrogen Corporation, Carlsbad, CA),  
20 penicillin 100 units per mL, and streptomycin 100 micrograms per mL (Invitrogen Corporation, Carlsbad, CA). Cells were routinely passaged by trypsinization and dilution when they reached 90% confluence.

#### 25 NHDF cells:

Human neonatal dermal fibroblast (NHDF) were obtained from the Clonetics Corporation (Walkersville, MD). NHDFs were routinely maintained in Fibroblast Growth Medium (Clonetics Corporation, Walkersville, MD) supplemented as recommended by  
30 the supplier. Cells were maintained for up to 10 passages as recommended by the supplier.

#### HEK cells:

Human embryonic keratinocytes (HEK) were obtained from the Clonetics Corporation (Walkersville, MD). HEKs were routinely maintained in Keratinocyte Growth Medium (Clonetics Corporation, Walkersville, MD) formulated as recommended by the supplier. Cells were routinely maintained for up to 10 passages as recommended by the supplier.

Treatment with antisense compounds:

When cells reached 70% confluency, they were treated with oligonucleotide. For cells grown in 96-well plates, wells were washed once with 100  $\mu$ L OPTI-MEM<sup>TM</sup>-1 reduced-serum medium (Invitrogen Corporation, Carlsbad, CA) and then treated with 130  $\mu$ L of OPTI-MEM<sup>TM</sup>-1 containing 3.75  $\mu$ g/mL LIPOFECTIN<sup>TM</sup> (Invitrogen Corporation, Carlsbad, CA) and the desired concentration of oligonucleotide. After 4-7 hours of treatment, the medium was replaced with fresh medium. Cells were harvested 16-24 hours after oligonucleotide treatment.

The concentration of oligonucleotide used varies from cell line to cell line. To determine the optimal oligonucleotide concentration for a particular cell line, the cells are treated with a positive control oligonucleotide at a range of concentrations. For human cells the positive control oligonucleotide is selected from either ISIS 13920 (**TCCGTCATCGCTCCTCAGGG**, SEQ ID NO: 1) which is targeted to human H-ras, or ISIS 18078, (**GTGCGCGGAGCCCGAAATC**, SEQ ID NO: 2) which is targeted to human Jun-N-terminal kinase-2 (JNK2). Both controls are 2'-O-methoxyethyl gapmers (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone. For mouse or rat cells the positive control oligonucleotide is ISIS 15770, **ATGCATTCTGCCCCAAGGA**, SEQ ID NO: 3, a 2'-O-methoxyethyl gapmer (2'-O-methoxyethyls shown in bold) with a phosphorothioate backbone which is targeted to both mouse and

rat c-raf. The concentration of positive control oligonucleotide that results in 80% inhibition of c-Ha-ras (for ISIS 13920) or c-raf (for ISIS 15770) mRNA is then utilized as the screening concentration for new oligonucleotides in subsequent experiments for that cell line. If 80% inhibition is not achieved, the lowest concentration of positive control oligonucleotide that results in 60% inhibition of H-ras or c-raf mRNA is then utilized as the oligonucleotide screening concentration in subsequent experiments for that cell line. If 60% inhibition is not achieved, that particular cell line is deemed as unsuitable for oligonucleotide transfection experiments. The concentrations of antisense oligonucleotides used herein are from 50 nM to 300 nM.

15

**Example 10****Analysis of oligonucleotide inhibition of G protein-coupled receptor kinase 6 expression**

Antisense modulation of G protein-coupled receptor kinase 6 expression can be assayed in a variety of ways known in the art. For example, G protein-coupled receptor kinase 6 mRNA levels can be quantitated by, e.g., Northern blot analysis, competitive polymerase chain reaction (PCR), or real-time PCR (RT-PCR). Real-time quantitative PCR is presently preferred. RNA analysis can be performed on total cellular RNA or poly(A)+ mRNA. The preferred method of RNA analysis of the present invention is the use of total cellular RNA as described in other examples herein. Methods of RNA isolation are taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*, Volume 1, pp. 4.1.1-4.2.9 and 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993. Northern blot analysis is routine in the art and is taught in, for example, Ausubel, F.M. et al., *Current Protocols in Molecular Biology*,

20  
25  
30

Volume 1, pp. 4.2.1-4.2.9, John Wiley & Sons, Inc., 1996.

Real-time quantitative (PCR) can be conveniently accomplished using the commercially available ABI PRISM™ 7700 Sequence Detection System, available from PE-Applied Biosystems, Foster  
5 City, CA and used according to manufacturer's instructions.

Protein levels of G protein-coupled receptor kinase 6 can be quantitated in a variety of ways well known in the art, such as immunoprecipitation, Western blot analysis

(immunoblotting), ELISA or fluorescence-activated cell sorting

10 (FACS). Antibodies directed to G protein-coupled receptor

kinase 6 can be identified and obtained from a variety of

sources, such as the MSRS catalog of antibodies (Aerie

Corporation, Birmingham, MI), or can be prepared via

conventional antibody generation methods. Methods for

15 preparation of polyclonal antisera are taught in, for example,

Ausubel, F.M. et al., (*Current Protocols in Molecular Biology*,

Volume 2, pp. 11.12.1-11.12.9, John Wiley & Sons, Inc., 1997).

Preparation of monoclonal antibodies is taught in, for

example, Ausubel, F.M. et al., (*Current Protocols in Molecular*

20 *Biology*, Volume 2, pp. 11.4.1-11.11.5, John Wiley & Sons,

Inc., 1997).

Immunoprecipitation methods are standard in the art and

can be found at, for example, Ausubel, F.M. et al., (*Current*

*Protocols in Molecular Biology*, Volume 2, pp. 10.16.1-

25 10.16.11, John Wiley & Sons, Inc., 1998). Western blot

(immunoblot) analysis is standard in the art and can be found

at, for example, Ausubel, F.M. et al., (*Current Protocols in*

*Molecular Biology*, Volume 2, pp. 10.8.1-10.8.21, John Wiley &

Sons, Inc., 1997). Enzyme-linked immunosorbent assays (ELISA)

30 are standard in the art and can be found at, for example,

Ausubel, F.M. et al., (*Current Protocols in Molecular Biology*,

Volume 2, pp. 11.2.1-11.2.22, John Wiley & Sons, Inc., 1991).

**Example 11****Poly(A)+ mRNA isolation**

Poly(A)+ mRNA was isolated according to Miura et al.,  
(*Clin. Chem.*, 1996, 42, 1758-1764). Other methods for  
5 poly(A)+ mRNA isolation are taught in, for example, Ausubel,  
F.M. et al., (*Current Protocols in Molecular Biology*, Volume  
1, pp. 4.5.1-4.5.3, John Wiley & Sons, Inc., 1993). Briefly,  
for cells grown on 96-well plates, growth medium was removed  
from the cells and each well was washed with 200  $\mu$ L cold PBS.  
10 60  $\mu$ L lysis buffer (10 mM Tris-HCl, pH 7.6, 1 mM EDTA, 0.5 M  
NaCl, 0.5% NP-40, 20 mM vanadyl-ribonucleoside complex) was  
added to each well, the plate was gently agitated and then  
incubated at room temperature for five minutes. 55  $\mu$ L of  
lysate was transferred to Oligo d(T) coated 96-well plates  
15 (AGCT Inc., Irvine CA). Plates were incubated for 60 minutes  
at room temperature, washed 3 times with 200  $\mu$ L of wash buffer  
(10 mM Tris-HCl pH 7.6, 1 mM EDTA, 0.3 M NaCl). After the  
final wash, the plate was blotted on paper towels to remove  
excess wash buffer and then air-dried for 5 minutes. 60  $\mu$ L of  
20 elution buffer (5 mM Tris-HCl pH 7.6), preheated to 70°C, was  
added to each well, the plate was incubated on a 90°C hot plate  
for 5 minutes, and the eluate was then transferred to a fresh  
96-well plate.

Cells grown on 100 mm or other standard plates may be  
25 treated similarly, using appropriate volumes of all solutions.

**Example 12****Total RNA Isolation**

Total RNA was isolated using an RNEASY 96™ kit and  
30 buffers purchased from Qiagen Inc. (Valencia, CA) following  
the manufacturer's recommended procedures. Briefly, for cells  
grown on 96-well plates, growth medium was removed from the

cells and each well was washed with 200  $\mu$ L cold PBS. 150  $\mu$ L Buffer RLT was added to each well and the plate vigorously agitated for 20 seconds. 150  $\mu$ L of 70% ethanol was then added to each well and the contents mixed by pipetting three times up and down. The samples were then transferred to the RNEASY 96<sup>TM</sup> well plate attached to a QIAVAC<sup>TM</sup> manifold fitted with a waste collection tray and attached to a vacuum source. Vacuum was applied for 1 minute. 500  $\mu$ L of Buffer RW1 was added to each well of the RNEASY 96<sup>TM</sup> plate and incubated for 15 minutes and the vacuum was again applied for 1 minute. An additional 500  $\mu$ L of Buffer RW1 was added to each well of the RNEASY 96<sup>TM</sup> plate and the vacuum was applied for 2 minutes. 1 mL of Buffer RPE was then added to each well of the RNEASY 96<sup>TM</sup> plate and the vacuum applied for a period of 90 seconds. The Buffer RPE wash was then repeated and the vacuum was applied for an additional 3 minutes. The plate was then removed from the QIAVAC<sup>TM</sup> manifold and blotted dry on paper towels. The plate was then re-attached to the QIAVAC<sup>TM</sup> manifold fitted with a collection tube rack containing 1.2 mL collection tubes. RNA was then eluted by pipetting 170  $\mu$ L water into each well, incubating 1 minute, and then applying the vacuum for 3 minutes.

The repetitive pipetting and elution steps may be automated using a QIAGEN Bio-Robot 9604 (Qiagen, Inc., Valencia CA). Essentially, after lysing of the cells on the culture plate, the plate is transferred to the robot deck where the pipetting, DNase treatment and elution steps are carried out.

30

**Example 13**

### Real-time Quantitative PCR Analysis of G protein-coupled receptor kinase 6 mRNA Levels

Quantitation of G protein-coupled receptor kinase 6 mRNA levels was determined by real-time quantitative PCR using the  
5 ABI PRISM™ 7700 Sequence Detection System (PE-Applied Biosystems, Foster City, CA) according to manufacturer's instructions. This is a closed-tube, non-gel-based, fluorescence detection system which allows high-throughput quantitation of polymerase chain reaction (PCR) products in  
10 real-time. As opposed to standard PCR in which amplification products are quantitated after the PCR is completed, products in real-time quantitative PCR are quantitated as they accumulate. This is accomplished by including in the PCR reaction an oligonucleotide probe that anneals specifically  
15 between the forward and reverse PCR primers, and contains two fluorescent dyes. A reporter dye (e.g., FAM or JOE, obtained from either PE-Applied Biosystems, Foster City, CA, Operon Technologies Inc., Alameda, CA or Integrated DNA Technologies Inc., Coralville, IA) is attached to the 5' end of the probe  
20 and a quencher dye (e.g., TAMRA, obtained from either PE-Applied Biosystems, Foster City, CA, Operon Technologies Inc., Alameda, CA or Integrated DNA Technologies Inc., Coralville, IA) is attached to the 3' end of the probe. When the probe and dyes are intact, reporter dye emission is quenched by the  
25 proximity of the 3' quencher dye. During amplification, annealing of the probe to the target sequence creates a substrate that can be cleaved by the 5'-exonuclease activity of Taq polymerase. During the extension phase of the PCR amplification cycle, cleavage of the probe by Taq polymerase  
30 releases the reporter dye from the remainder of the probe (and hence from the quencher moiety) and a sequence-specific fluorescent signal is generated. With each cycle, additional reporter dye molecules are cleaved from their respective

probes, and the fluorescence intensity is monitored at regular intervals by laser optics built into the ABI PRISM™ 7700 Sequence Detection System. In each assay, a series of parallel reactions containing serial dilutions of mRNA from untreated control samples generates a standard curve that is used to quantitate the percent inhibition after antisense oligonucleotide treatment of test samples.

Prior to quantitative PCR analysis, primer-probe sets specific to the target gene being measured are evaluated for their ability to be "multiplexed" with a GAPDH amplification reaction. In multiplexing, both the target gene and the internal standard gene GAPDH are amplified concurrently in a single sample. In this analysis, mRNA isolated from untreated cells is serially diluted. Each dilution is amplified in the presence of primer-probe sets specific for GAPDH only, target gene only ("single-plexing"), or both (multiplexing). Following PCR amplification, standard curves of GAPDH and target mRNA signal as a function of dilution are generated from both the single-plexed and multiplexed samples. If both the slope and correlation coefficient of the GAPDH and target signals generated from the multiplexed samples fall within 10% of their corresponding values generated from the single-plexed samples, the primer-probe set specific for that target is deemed multiplexable. Other methods of PCR are also known in the art.

PCR reagents were obtained from Invitrogen Corporation, (Carlsbad, CA). RT-PCR reactions were carried out by adding 20  $\mu$ L PCR cocktail (2.5x PCR buffer (-MgCl<sub>2</sub>), 6.6 mM MgCl<sub>2</sub>, 375  $\mu$ M each of dATP, dCTP, dGTP and dTTP, 375 nM each of forward primer and reverse primer, 125 nM of probe, 4 Units RNase inhibitor, 1.25 Units PLATINUM® Taq, 5 Units MuLV reverse transcriptase, and 2.5x ROX dye) to 96-well plates

containing 30  $\mu$ L total RNA solution. The RT reaction was carried out by incubation for 30 minutes at 48°C. Following a 10 minute incubation at 95°C to activate the PLATINUM® Taq, 40 cycles of a two-step PCR protocol were carried out: 95°C for 15  
5 seconds (denaturation) followed by 60°C for 1.5 minutes (annealing/extension).

Gene target quantities obtained by real time RT-PCR are normalized using either the expression level of GAPDH, a gene whose expression is constant, or by quantifying total RNA  
10 using RiboGreen™ (Molecular Probes, Inc. Eugene, OR). GAPDH expression is quantified by real time RT-PCR, by being run simultaneously with the target, multiplexing, or separately. Total RNA is quantified using RiboGreen™ RNA quantification reagent from Molecular Probes. Methods of RNA quantification  
15 by RiboGreen™ are taught in Jones, L.J., et al, (Analytical Biochemistry, 1998, 265, 368-374).

In this assay, 170  $\mu$ L of RiboGreen™ working reagent (RiboGreen™ reagent diluted 1:350 in 10mM Tris-HCl, 1 mM EDTA, pH 7.5) is pipetted into a 96-well plate containing 30  
20  $\mu$ L purified, cellular RNA. The plate is read in a CytoFluor 4000 (PE Applied Biosystems) with excitation at 480nm and emission at 520nm.

Probes and primers to human G protein-coupled receptor kinase 6 were designed to hybridize to a human G protein-coupled receptor kinase 6 sequence, using published sequence  
25 information (GenBank accession number NM\_002082.1, incorporated herein as SEQ ID NO:4). For human G protein-coupled receptor kinase 6 the PCR primers were:

forward primer: CCCTGGTCAACCCTCAAACA (SEQ ID NO: 5)  
30 reverse primer: CGGATTATTGCTGGGCACAT (SEQ ID NO: 6) and the PCR probe was: FAM-TCCGGACTCCCCTCATAACAATAGAC-TAMRA (SEQ ID NO: 7) where FAM is the fluorescent dye and TAMRA is

the quencher dye. For human GAPDH the PCR primers were:

forward primer: GAAGGTGAAGGTCGGAGTC (SEQ ID NO:8)

reverse primer: GAAGATGGTGATGGGATTTTC (SEQ ID NO:9) and the PCR

probe was: 5' JOE-CAAGCTTCCCGTTCTCAGCC- TAMRA 3' (SEQ ID NO:

5 10) where JOE is the fluorescent reporter dye and TAMRA is the quencher dye.

#### Example 14

#### 10 Northern blot analysis of G protein-coupled receptor kinase 6 mRNA levels

Eighteen hours after antisense treatment, cell monolayers were washed twice with cold PBS and lysed in 1 mL RNAZOL™

(TEL-TEST "B" Inc., Friendswood, TX). Total RNA was prepared

15 following manufacturer's recommended protocols. Twenty

micrograms of total RNA was fractionated by electrophoresis

through 1.2% agarose gels containing 1.1% formaldehyde using a

MOPS buffer system (AMRESCO, Inc. Solon, OH). RNA was

transferred from the gel to HYBOND™-N+ nylon membranes

20 (Amersham Pharmacia Biotech, Piscataway, NJ) by overnight

capillary transfer using a Northern/Southern Transfer buffer

system (TEL-TEST "B" Inc., Friendswood, TX). RNA transfer was

confirmed by UV visualization. Membranes were fixed by UV

cross-linking using a STRATALINKER™ UV Crosslinker 2400

25 (Stratagene, Inc, La Jolla, CA) and then probed using

QUICKHYB™ hybridization solution (Stratagene, La Jolla, CA)

using manufacturer's recommendations for stringent conditions.

To detect human G protein-coupled receptor kinase 6, a human G protein-coupled receptor kinase 6 specific probe was

30 prepared by PCR using the forward primer CCCTGGTCAACCCTCAAACA

(SEQ ID NO: 5) and the reverse primer CGGATTATTGCTGGGCACAT

(SEQ ID NO: 6). To normalize for variations in loading and

transfer efficiency membranes were stripped and probed for human glyceraldehyde-3-phosphate dehydrogenase (GAPDH) RNA (Clontech, Palo Alto, CA).

Hybridized membranes were visualized and quantitated using a PHOSPHORIMAGER™ and IMAGEQUANT™ Software V3.3 (Molecular Dynamics, Sunnyvale, CA). Data was normalized to GAPDH levels in untreated controls.

#### 10 Example 15

##### **Antisense inhibition of human G protein-coupled receptor kinase 6 expression by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap**

In accordance with the present invention, a series of oligonucleotides were designed to target different regions of the human G protein-coupled receptor kinase 6 RNA, using published sequences (GenBank accession number NM\_002082.1, incorporated herein as SEQ ID NO: 4). The oligonucleotides are shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number on the particular target sequence to which the oligonucleotide binds. All compounds in Table 1 are chimeric oligonucleotides ("gapmers") 20 nucleotides in length, composed of a central "gap" region consisting of ten 2'-deoxynucleotides, which is flanked on both sides (5' and 3' directions) by five-nucleotide "wings". The wings are composed of 2'-methoxyethyl (2'-MOE)nucleotides. The internucleoside (backbone) linkages are phosphorothioate (P=S) throughout the oligonucleotide. All cytidine residues are 5-methylcytidines. The compounds were analyzed for their effect on human G protein-coupled receptor kinase 6 mRNA levels by quantitative real-time PCR as described in other examples herein. Data are averages from two experiments in which A549 cells were treated with the oligonucleotides of the present

invention. The positive control for each datapoint is identified in the table by sequence ID number. If present, "N.D." indicates "no data".

5

Table 1

Inhibition of human G protein-coupled receptor kinase 6 mRNA levels by chimeric phosphorothioate oligonucleotides having 2'-MOE wings and a deoxy gap

ISIS #	REGION	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	% INHIB	SEQ ID NO	CONTROL SEQ ID NO
207121	Start Codon	4	55	ctcgagctccatgggctgtg	32	11	1
207122	Coding	4	82	gagtagcaccgtgttcgcta	52	12	1
207123	Coding	4	155	gagggaaactggagcatctgc	43	13	1
207124	Coding	4	174	tcttcgcactggctgatgtg	41	14	1
207125	Coding	4	193	ctcgaggctgagccgcagct	44	15	1
207126	Coding	4	210	aggctgtgatagtacagctc	42	16	1
207127	Coding	4	219	cgctcgcacaggctgtgata	47	17	1
207128	Coding	4	259	ggcacagaactctcggaaca	31	18	1
207129	Coding	4	272	gtccggcctcgtggcacag	34	19	1
207130	Coding	4	341	ccttcgcttgtcatccggg	39	20	1
207131	Coding	4	385	accgctgtggctcagaaaat	79	21	1
207132	Coding	4	395	tgaggtcaggaccgctgtgg	22	22	1
207133	Coding	4	461	ctttgcagggaccctgctcc	43	23	1
207134	Coding	4	479	tgagttcctggaaaaggtct	39	24	1
207135	Coding	4	500	ggtactcgtgggtcagccgg	33	25	1
207136	Coding	4	551	aacggttgaagtagatgctg	50	26	1
207137	Coding	4	589	ggtcactggctgcctttcca	33	27	1
207138	Coding	4	628	acctttgccaggactcgggt	67	28	1
207139	Coding	4	725	cccctttccgcttcttgatc	34	29	1
207140	Coding	4	755	ggatctgcttctcgttcagc	56	30	1
207141	Coding	4	793	ggccaagctcactacaaacc	43	31	1
207142	Coding	4	816	gcgctccttggctcataggc	70	32	1
207143	Coding	4	831	agcaccaggcacagcgcgctc	53	33	1
207144	Coding	4	843	ttcatcagtgtcagcaccag	0	34	1
207145	Coding	4	853	gtcgcccccgcttcatcagtg	15	35	1
207146	Coding	4	879	cccatgtggtatagtgaggaa	63	36	1
207147	Coding	4	945	tcctccaggccacagcagat	35	37	1
207148	Coding	4	962	tgcgctcccgggtgcaggtcc	61	38	1
207149	Coding	4	976	gtccctgtacacgatgcgct	69	39	1
207150	Coding	4	1003	atccagcaagatgttctcgg	29	40	1
207151	Coding	4	1012	gccgtggtcatccagcaaga	57	41	1
207152	Coding	4	1049	catgcacagctagtcccagg	45	42	1
207153	Coding	4	1060	gccctcgggcacatgcacag	56	43	1
207154	Coding	4	1074	cctttgatggtctggccctc	46	44	1
207155	Coding	4	1096	gtaaccacgggtgccacac	62	45	1
207156	Coding	4	1108	ctccggagccatgtaaccca	40	46	1
207157	Coding	4	1122	tcattcttcaccacctccgg	50	47	1

207158	Coding	4	1140	gggctgaacgtgtaccgttc	42	48	1
207159	Coding	4	1193	gcgactggcctgcatcatc	20	49	1
207160	Coding	4	1314	aggagctgtgagcaaagtga	48	50	1
207161	Coding	4	1399	gttcagcttcttaaagagg	17	51	1
207162	Coding	4	1437	aacggcggctccagcatgcc	43	52	1
207163	Coding	4	1469	ccttgagtaaatggcctgg	41	53	1
207164	Coding	4	1485	tcaatgtccagaacatcctt	11	54	1
207165	Coding	4	1496	tagagaactgttcaatgtcc	60	55	1
207166	Coding	4	1515	agctccacgcccttgaccgt	36	56	1
207167	Coding	4	1549	aaacttctggtagaagtcc	16	57	1
207168	Coding	4	1571	ggatgggcacactgcctgtg	19	58	1
207169	Coding	4	1596	gtctccaccatctcgttctg	38	59	1
207170	Coding	4	1628	gcccaaagacattcagctct	19	60	1
207171	Coding	4	1669	ctggcccttccagtcagggt	25	61	1
207172	Coding	4	1690	cttttttaggaggtgcagggtg	13	62	1
207173	Coding	4	1736	agtttccacagcaatcttgg	25	63	1
207174	3'UTR	4	1825	gctcggagttagctgctaccg	52	64	1
207175	3'UTR	4	1854	gaagatcactgtgcaaaact	62	65	1
207176	3'UTR	4	1878	gccacgactgagctggacaa	70	66	1
207177	3'UTR	4	1888	tgttccccaggccacgactt	52	67	1
207178	3'UTR	4	1924	ctgaggggacggaggacactg	42	68	1
207179	3'UTR	4	1947	gccaaaactcagccaggccag	64	69	1
207180	3'UTR	4	1963	ggatggcccaggccctgccca	69	70	1
207181	3'UTR	4	2073	cggcgggaattaagaatgct	53	71	1
207182	3'UTR	4	2112	tggctgccccaggagccaa	39	72	1
207183	3'UTR	4	2154	tggcagtggctcctctgccca	0	73	1
207184	3'UTR	4	2201	tgtgaccaccctcagccatc	9	74	1
207185	3'UTR	4	2212	gaaggctcaggtgtgaccac	0	75	1
207186	3'UTR	4	2220	acagtgtgaaggctcagggt	65	76	1
207187	3'UTR	4	2304	acagcttggggcccagagg	52	77	1
207188	3'UTR	4	2357	ggcctagggcatacagaggc	55	78	1
207189	3'UTR	4	2372	ggccacttggcacaggcct	66	79	1
207190	3'UTR	4	2443	aaggcagaggcagacctatc	58	80	1
207191	3'UTR	4	2560	aatgtttgaggttgaccag	44	81	1
207192	3'UTR	4	2603	ttattgctgggcacatgtgt	80	82	1
207193	3'UTR	4	2664	tcaccacacaggtacacac	19	83	1
207194	3'UTR	4	2749	aactctgggaccaggatggc	39	84	1
207195	3'UTR	4	2760	catgctaccctaactggg	41	85	1
207196	3'UTR	4	2802	aaaattggctttaaatatgg	10	86	1
207197	3'UTR	4	2812	acgagtaataaaaattggct	50	87	1
207198	3'UTR	4	2827	tttacgttggacaaaacgag	35	88	1

As shown in Table 1, SEQ ID NOs 12, 13, 14, 15, 16, 17, 21, 23, 26, 28, 30, 31, 32, 33, 36, 38, 39, 41, 42, 43, 44, 45, 46, 47, 48, 50, 52, 53, 55, 64, 65, 66, 67, 68, 69, 70, 71, 76, 77, 78, 79, 80, 81, 82, 85 and 87 demonstrated at least 40% inhibition of human G protein-coupled receptor kinase 6 expression in this assay and are therefore preferred. The target sites to which these preferred sequences are

complementary are herein referred to as "preferred target regions" and are therefore preferred sites for targeting by compounds of the present invention. These preferred target regions are shown in Table 2. The sequences represent the reverse complement of the preferred antisense compounds shown in Table 1. "Target site" indicates the first (5'-most) nucleotide number of the corresponding target nucleic acid. Also shown in Table 2 is the species in which each of the preferred target regions was found.

10

**Table 2**  
**Sequence and position of preferred target regions identified in G protein-coupled receptor kinase 6.**

15

SITEID	TARGET SEQ ID NO	TARGET SITE	SEQUENCE	REV COMP OF SEQ ID	ACTIVE IN	SEQ ID NO
124756	4	82	tagcgaacacggtgctactc	12	<i>H. sapiens</i>	89
124757	4	155	gcagatgctccagttccctc	13	<i>H. sapiens</i>	90
124758	4	174	cacatcagccagtgccaaga	14	<i>H. sapiens</i>	91
124759	4	193	agctgcggtcagcctcgag	15	<i>H. sapiens</i>	92
124760	4	210	gagcgtgactatcacagcct	16	<i>H. sapiens</i>	93
124761	4	219	tatcacagcctgtgagcg	17	<i>H. sapiens</i>	94
124765	4	385	atcttctgagccacacgggt	21	<i>H. sapiens</i>	95
124767	4	461	ggagcagggtccctgcaaag	23	<i>H. sapiens</i>	96
124770	4	551	cagcatctacttcaaccggt	26	<i>H. sapiens</i>	97
124772	4	628	accgagtcctgggcaaagg	28	<i>H. sapiens</i>	98
124774	4	755	gctgaacgagaagcagatcc	30	<i>H. sapiens</i>	99
124775	4	793	ggttttagtgagcttgcc	31	<i>H. sapiens</i>	100
124776	4	816	gcctatgagaccaaggacgc	32	<i>H. sapiens</i>	101
124777	4	831	gacgcgtgtgctggtgct	33	<i>H. sapiens</i>	102
124780	4	879	ttccacatctaccatggg	36	<i>H. sapiens</i>	103
124782	4	962	ggacctgcaccgggagcgca	38	<i>H. sapiens</i>	104
124783	4	976	agcgcatcgtgtacagggac	39	<i>H. sapiens</i>	105
124785	4	1012	tcttgctggatgaccacggc	41	<i>H. sapiens</i>	106
124786	4	1049	cctgggactagctgtgcatg	42	<i>H. sapiens</i>	107
124787	4	1060	ctgtgcatgtgcccagggc	43	<i>H. sapiens</i>	108
124788	4	1074	gagggccagaccatcaaagg	44	<i>H. sapiens</i>	109
124789	4	1096	gtgtgggcaccgtgggttac	45	<i>H. sapiens</i>	110
124790	4	1108	tgggttacatggctccggag	46	<i>H. sapiens</i>	111
124791	4	1122	ccggaggtggtgaagaatga	47	<i>H. sapiens</i>	112
124792	4	1140	gaacggtacacgttcagccc	48	<i>H. sapiens</i>	113
124794	4	1314	tcactttgctcacagctcct	50	<i>H. sapiens</i>	114
124796	4	1437	ggcatgctggagccgctgt	52	<i>H. sapiens</i>	115
124797	4	1469	ccaggccatttactgcaagg	53	<i>H. sapiens</i>	116

124799	4	1496	ggacattgaacagttctcta	55	<i>H. sapiens</i>	117
124808	4	1825	cggtagcagctactccgagc	64	<i>H. sapiens</i>	118
124809	4	1854	agttttgcacagtgatcttc	65	<i>H. sapiens</i>	119
124810	4	1878	ttgtccactcaagtcgtggc	66	<i>H. sapiens</i>	120
124811	4	1888	aagtcgtggcctggggaaca	67	<i>H. sapiens</i>	121
124812	4	1924	cagtgtcctccgtccctcag	68	<i>H. sapiens</i>	122
124813	4	1947	ctggcctggctgagtttggc	69	<i>H. sapiens</i>	123
124814	4	1963	tggcagggcctgggccatcc	70	<i>H. sapiens</i>	124
124815	4	2073	agcattcttaattcccgcgcg	71	<i>H. sapiens</i>	125
124820	4	2220	acctgagccttcagcactgt	76	<i>H. sapiens</i>	126
124821	4	2304	cctctggggcccaaagctgt	77	<i>H. sapiens</i>	127
124822	4	2357	gcctctgtatgccctaggcc	78	<i>H. sapiens</i>	128
124823	4	2372	aggcctgtgcccagtgccc	79	<i>H. sapiens</i>	129
124824	4	2443	gataggtctgcctctgcctt	80	<i>H. sapiens</i>	130
124825	4	2560	ctggtcaaccctcaacatt	81	<i>H. sapiens</i>	131
124826	4	2603	acacatgtgccagcaataa	82	<i>H. sapiens</i>	132
124829	4	2760	cccagtgtaggtagcatg	85	<i>H. sapiens</i>	133
124831	4	2812	agccaatttttattactcgt	87	<i>H. sapiens</i>	134

As these "preferred target regions" have been found by  
 5 experimentation to be open to, and accessible for,  
 hybridization with the antisense compounds of the present  
 invention, one of skill in the art will recognize or be able  
 to ascertain, using no more than routine experimentation,  
 further embodiments of the invention that encompass other  
 10 compounds that specifically hybridize to these sites and  
 consequently inhibit the expression of G protein-coupled  
 receptor kinase 6.

#### 15 Example 16

##### Western blot analysis of G protein-coupled receptor kinase 6 protein levels

Western blot analysis (immunoblot analysis) is carried  
 out using standard methods. Cells are harvested 16-20 h after  
 20 oligonucleotide treatment, washed once with PBS, suspended in  
 Laemmli buffer (100 ul/well), boiled for 5 minutes and loaded  
 on a 16% SDS-PAGE gel. Gels are run for 1.5 hours at 150 V,

and transferred to membrane for western blotting. Appropriate primary antibody directed to G protein-coupled receptor kinase 6 is used, with a radiolabeled or fluorescently labeled secondary antibody directed against the primary antibody species. Bands are visualized using a PHOSPHORIMAGER™ (Molecular Dynamics, Sunnyvale CA).

**What is claimed is:**

1. A compound 8 to 80 nucleobases in length targeted to a nucleic acid molecule encoding G protein-coupled receptor kinase 6, wherein said compound specifically hybridizes with said nucleic acid molecule encoding G protein-coupled receptor kinase 6 and inhibits the expression of G protein-coupled receptor kinase 6.

2. The compound of claim 1 which is an antisense oligonucleotide.

3. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified internucleoside linkage.

4. The compound of claim 3 wherein the modified internucleoside linkage is a phosphorothioate linkage.

5. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified sugar moiety.

6. The compound of claim 5 wherein the modified sugar moiety is a 2'-O-methoxyethyl sugar moiety.

7. The compound of claim 2 wherein the antisense oligonucleotide comprises at least one modified nucleobase.

8. The compound of claim 7 wherein the modified nucleobase is a 5-methylcytosine.

9. The compound of claim 2 wherein the antisense oligonucleotide is a chimeric oligonucleotide.

10. A compound 8 to 80 nucleobases in length which specifically hybridizes with at least an 8-nucleobase portion of a preferred target region on a nucleic acid molecule encoding G protein-coupled receptor kinase 6.

11. A composition comprising the compound of claim 1 and a pharmaceutically acceptable carrier or diluent.

12. The composition of claim 11 further comprising a colloidal dispersion system.

13. The composition of claim 11 wherein the compound is an antisense oligonucleotide.

14. A method of inhibiting the expression of G protein-coupled receptor kinase 6 in cells or tissues comprising contacting said cells or tissues with the compound of claim 1 so that expression of G protein-coupled receptor kinase 6 is inhibited.

15. A method of treating an animal having a disease or condition associated with G protein-coupled receptor kinase 6 comprising administering to said animal a therapeutically or prophylactically effective amount of the compound of claim 1 so that expression of G protein-coupled receptor kinase 6 is inhibited.

16. The method of claim 15 wherein the disease or condition is rheumatoid arthritis.

17. The method of claim 15 wherein the disease or condition is drug addiction.

18. The method of claim 15 wherein the disease or condition is a disorder of uterine contractility.

19. The method of claim 15 wherein the disease or condition is hypertension.

20. The method of claim 15 wherein the disease or condition arises from aberrant hematopoiesis.

## SEQUENCE LISTING

<110> Susan M. Freier  
Kenneth W. Dobie

<120> ANTISENSE MODULATION OF G PROTEIN-COUPLED RECEPTOR KINASE 6  
EXPRESSION

<130> RTS-0365

<160> 134

<210> 1

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 1

tccgtcatcg ctctcaggg

20

<210> 2

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 2

gtgcgcgcgga gcccgaaatc

20

<210> 3  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 3  
 atgcattctg cccccaagga 20

<210> 4  
 <211> 2848  
 <212> DNA  
 <213> H. sapiens

<220>

<220>  
 <221> CDS  
 <222> (63)...(1793)

<400> 4  
 cgcggcactc ggcgcgcatc ccggccggcg gcgcgccggc gggccaggcg gcgccacagc . 60

cc atg gag ctc gag aac atc gta gcg aac acg gtg cta ctc aag gcc 107  
 Met Glu Leu Glu Asn Ile Val Ala Asn Thr Val Leu Leu Lys Ala  
 1 5 10 15

cgg gaa ggt ggc ggt gga aat cgc aaa ggc aaa agc aag aaa tgg cgg 155  
 Arg Glu Gly Gly Gly Gly Asn Arg Lys Gly Lys Ser Lys Lys Trp Arg  
 20 25 30

cag atg ctc cag ttc cct cac atc agc cag tgc gaa gag ctg cgg ctc 203  
 Gln Met Leu Gln Phe Pro His Ile Ser Gln Cys Glu Glu Leu Arg Leu  
 35 40 45

agc ctc gag cgt gac tat cac agc ctg tgc gag cgg cac gcc att ggg	251
Ser Leu Glu Arg Asp Tyr His Ser Leu Cys Glu Arg His Ala Ile Gly	
50 55 60	
cgc ctg ctg ttc cga gag ttc tgt gcc acg agg ccg gag ctg agc cgc	299
Arg Leu Leu Phe Arg Glu Phe Cys Ala Thr Arg Pro Glu Leu Ser Arg	
65 70 75	
tgc gtc gcc ttc ctg gat ggg gtg gcc gag tat gaa gtg acc ccg gat	347
Cys Val Ala Phe Leu Asp Gly Val Ala Glu Tyr Glu Val Thr Pro Asp	
80 85 90 95	
gac aag cgg aag gca tgt ggg cgg cac gta acg cag aat ttt ctg agc	395
Asp Lys Arg Lys Ala Cys Gly Arg His Val Thr Gln Asn Phe Leu Ser	
100 105 110	
cac acg ggt cct gac ctc atc cct gag gtc ccc egg cag ctg gtg acg	443
His Thr Gly Pro Asp Leu Ile Pro Glu Val Pro Arg Gln Leu Val Thr	
115 120 125	
aac tgc acc cag cgg ctg gag cag ggt ccc tgc aaa gac ctt ttc cag	491
Asn Cys Thr Gln Arg Leu Glu Gln Gly Pro Cys Lys Asp Leu Phe Gln	
130 135 140	
gaa ctc acc cgg ctg acc cac gag tac ctg agc gtg gcc cct ttt gcc	539
Glu Leu Thr Arg Leu Thr His Glu Tyr Leu Ser Val Ala Pro Phe Ala	
145 150 155	
gac tac ctc gac agc atc tac ttc aac cgt ttc ctg cag tgg aag tgg	587
Asp Tyr Leu Asp Ser Ile Tyr Phe Asn Arg Phe Leu Gln Trp Lys Trp	
160 165 170 175	
ctg gaa agg cag cca gtg acc aaa aac acc ttc agg caa tac cga gtc	635
Leu Glu Arg Gln Pro Val Thr Lys Asn Thr Phe Arg Gln Tyr Arg Val	
180 185 190	
ctg ggc aaa ggt ggc ttt ggg gag gtg tgc gcc tgc cag gtg cgg gcc	683
Leu Gly Lys Gly Gly Phe Gly Glu Val Cys Ala Cys Gln Val Arg Ala	
195 200 205	

aca ggt aag atg tat gcc tgc aag aag cta gag aaa aag cgg atc aag 731  
 Thr Gly Lys Met Tyr Ala Cys Lys Lys Leu Glu Lys Lys Arg Ile Lys  
 210 215 220

aag cgg aaa ggg gag gcc atg gcg ctg aac gag aag cag atc ctg gag 779  
 Lys Arg Lys Gly Glu Ala Met Ala Leu Asn Glu Lys Gln Ile Leu Glu  
 225 230 235

aaa gtg aac agt agg ttt gta gtg agc ttg gcc tac gcc tat gag acc 827  
 Lys Val Asn Ser Arg Phe Val Val Ser Leu Ala Tyr Ala Tyr Glu Thr  
 240 245 250 255

aag gac gcg ctg tgc ctg gtg ctg aca ctg atg aac ggg ggc gac ctc 875  
 Lys Asp Ala Leu Cys Leu Val Leu Thr Leu Met Asn Gly Gly Asp Leu  
 260 265 270

aag ttc cac atc tac cac atg ggc cag gct ggc ttc ccc gaa gcg cgg 923  
 Lys Phe His Ile Tyr His Met Gly Gln Ala Gly Phe Pro Glu Ala Arg  
 275 280 285

gcc gtc ttc tac gcc gcc gag atc tgc tgt ggc ctg gag gac ctg cac 971  
 Ala Val Phe Tyr Ala Ala Glu Ile Cys Cys Gly Leu Glu Asp Leu His  
 290 295 300

cgg gag cgc atc gtg tac agg gac ctg aag ccc gag aac atc ttg ctg 1019  
 Arg Glu Arg Ile Val Tyr Arg Asp Leu Lys Pro Glu Asn Ile Leu Leu  
 305 310 315

gat gac cac ggc cac atc cgc atc tct gac ctg gga cta gct gtg cat 1067  
 Asp Asp His Gly His Ile Arg Ile Ser Asp Leu Gly Leu Ala Val His  
 320 325 330 335

gtg ccc gag ggc cag acc atc aaa ggg cgt gtg ggc acc gtg ggt tac 1115  
 Val Pro Glu Gly Gln Thr Ile Lys Gly Arg Val Gly Thr Val Gly Tyr  
 340 345 350

atg gct ccg gag gtg gtg aag aat gaa cgg tac acg ttc agc cct gac 1163  
 Met Ala Pro Glu Val Val Lys Asn Glu Arg Tyr Thr Phe Ser Pro Asp

355	360	365	
tgg tgg gcg ctc ggc tgc ctc ctg tac gag atg atc gca ggc cag tcg			1211
Trp Trp Ala Leu Gly Cys Leu Leu Tyr Glu Met Ile Ala Gly Gln Ser			
370	375	380	
ccc ttc cag cag agg aag aag aag atc aag cgg gag gag gtg gag cgg			1259
Pro Phe Gln Gln Arg Lys Lys Lys Ile Lys Arg Glu Glu Val Glu Arg			
385	390	395	
ctg gtg aag gag gtc ccc gag gag tat tcc gag cgc ttt tcc ccg cag			1307
Leu Val Lys Glu Val Pro Glu Glu Tyr Ser Glu Arg Phe Ser Pro Gln			
400	405	410	415
gcc cgc tca ctt tgc tca cag ctc ctc tgc aag gac cct gcc gaa cgc			1355
Ala Arg Ser Leu Cys Ser Gln Leu Leu Cys Lys Asp Pro Ala Glu Arg			
420	425	430	
ctg ggg tgt cgt ggg ggc agt gcc cgc gag gtg aag gag cac ccc ctc			1403
Leu Gly Cys Arg Gly Gly Ser Ala Arg Glu Val Lys Glu His Pro Leu			
435	440	445	
ttt aag aag ctg aac ttc aag cgg ctg gga gct ggc atg ctg gag ccg			1451
Phe Lys Lys Leu Asn Phe Lys Arg Leu Gly Ala Gly Met Leu Glu Pro			
450	455	460	
ccg ttc aag cct gac ccc cag gcc att tac tgc aag gat gtt ctg gac			1499
Pro Phe Lys Pro Asp Pro Gln Ala Ile Tyr Cys Lys Asp Val Leu Asp			
465	470	475	
att gaa cag ttc tct acg gtc aag ggc gtg gag ctg gag cct acc gac			1547
Ile Glu Gln Phe Ser Thr Val Lys Gly Val Glu Leu Glu Pro Thr Asp			
480	485	490	495
cag gac ttc tac cag aag ttt gcc aca ggc agt gtg ccc atc ccc tgg			1595
Gln Asp Phe Tyr Gln Lys Phe Ala Thr Gly Ser Val Pro Ile Pro Trp			
500	505	510	
cag aac gag atg gtg gag acc gag tgc ttc caa gag ctg aat gtc ttt			1643

Gln	Asn	Glu	Met	Val	Glu	Thr	Glu	Cys	Phe	Gln	Glu	Leu	Asn	Val	Phe		
			515					520					525				
ggg	ctg	gat	ggc	tca	ggt	ccc	cca	gac	ctg	gac	tgg	aag	ggc	cag	cca		1691
Gly	Leu	Asp	Gly	Ser	Val	Pro	Pro	Asp	Leu	Asp	Trp	Lys	Gly	Gln	Pro		
			530					535					540				
cct	gca	cct	cct	aaa	aag	gga	ctg	ctg	cag	aga	ctc	ttc	agt	cgc	caa		1739
Pro	Ala	Pro	Pro	Lys	Lys	Gly	Leu	Leu	Gln	Arg	Leu	Phe	Ser	Arg	Gln		
			545					550					555				
gat	tgc	tgt	gga	aac	tgc	agc	gac	agc	gag	gaa	gag	ctc	ccc	acc	cgc		1787
Asp	Cys	Cys	Gly	Asn	Cys	Ser	Asp	Ser	Glu	Glu	Glu	Leu	Pro	Thr	Arg		
560					565					570					575		
ctc	tag	ccccagccc	gaggccccca	ccagcagttg	gcggtagcag	ctactccgag											1843
Leu																	
cgccgtttac	agttttgcac	agtgatcttc	cccattgtcc	actcaagtcg	tggcctgggg												1903
aacacagacg	gagctgtccc	cagtgtcctc	cgtcctcag	ccctggcct	ggetgagttt												1963
ggcagggcct	gggccatccc	tgggacaaag	gtgcgtccct	tcagctcttc	tccgtggagc												2023
tgggggcttt	ctgtatttat	gtatattgtac	gaatgtatat	agcgaccaga	gcattcttaa												2083
ttcccgcgcg	agacctggcg	ccccgcctt	ggctcctggg	ggcagccagc	cctggctgga												2143
gagcgggacg	tggcagagga	gccactgcca	aactcaaggc	tcctctggcc	cagcttgat												2203
ggctgagggt	ggtcacacct	gagccttcag	cactgtgctg	ccaccccgcc	ctctgagtaa												2263
gactcgtgcc	tccccctgct	gccctgggct	caggctgcta	cctctggggc	ccaaagctgt												2323
cccttctcag	tgcttgtcag	cgctgggtct	ggggcctctg	tatgccctag	gcctgtgcca												2383
aagtggccag	agattgggct	gcctgtgata	cccatcagcc	cactgccccg	gccggcccag												2443

ataggctctgc ctctgccttc cagctcccac agcctgggtcc ctgatactgg gctctgtcct 2503  
 gcagacacct ctttcagaaa cgcccaagcc cagccctagg aggggggtggg gcatccctgg 2563  
 tcaaccctca aacattccgg actcccctca taacaataga cacatgtgcc cagcaataat 2623  
 ccgccccttc ctgtgtgctc ctgtgggggtg cgtgctctct gtgtgtacct gtgtgggtga 2683  
 aggggatagg gcgaggctgt gcctgtgccc cagggtcccag ccctggccct tcccagactg 2743  
 tgatggccat cctgggtcca gtgttagggt agcatgggat tacagggccc tgttttttcc 2803  
 atatttaaag ccaattttta ttactcgttt tgtccaacgt aaaag 2848

<210> 5  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 5  
 ccctgggtcaa ccctcaaaca 20

<210> 6  
 <211> 20  
 <212> DNA  
 <213> Artificial Sequence

<220>

<223> PCR Primer

<400> 6  
 cggattattg ctgggcacat 20

<210> 7

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 7

tccggactcc cctcataaca atagac

26

<210> 8

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 8

gaaggatgaag gtcggagtc

19

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Primer

<400> 9

gaagatggtg atgggatttc

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR Probe

<400> 10

caagcttccc gttctcagcc

20

<210> 11

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 11

ctcgagctcc atgggctgtg

20

<210> 12

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 12

gagtagcacc gtgttcgcta

20

<210> 13

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 13

gaggggaactg gagcatctgc

20

<210> 14

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 14

tcttcgcact ggctgatgtg

20

<210> 15

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 15

ctcgaggctg agccgcagct

20

<210> 16

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 16

aggctgtgat agtcacgctc

20

<210> 17

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 17

cgctcgaca ggctgtgata

20

<210> 18

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 18

ggcacagaac tctcggaaca

20

<210> 19

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 19

gctccggcct cgtggcacag

20

<210> 20

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 20

ccttccgctt gtcattccggg

20

<210> 21

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 21

accctgtgg ctcagaaaat

20

<210> 22

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 22

tgaggtcagg acccgtgtgg

20

<210> 23

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 23

ctttgcaggg accctgctcc

20

<210> 24

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 24

tgagttcctg gaaaaggtct

20

<210> 25

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 25

ggtactcgtg ggtcagccgg

20

<210> 26

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 26

aacggttgaa gtagatgctg

20

<210> 27

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 27

ggtcactggc tgcctttcca

20

<210> 28

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 28

acctttgcc aggactcgg

20

<210> 29

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 29

cccctttccg cttcttgatc

20

<210> 30

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 30

ggatctgctt ctcgttcagc

20

<210> 31

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 31

ggccaagctc actacaaacc

20

<210> 32

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 32

gcgtccttgg tctcataggc

20

<210> 33

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 33

agcaccaggc acagcgcgctc

20

<210> 34

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 34

ttcatcagtg tcagcaccag

20

<210> 35

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 35

gtcgcccccg ttcatcagtg

20

<210> 36

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 36

cccatgtggt agatgtggaa

20

<210> 37

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 37

tcctccaggc cacagcagat

20

<210> 38

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 38

tgcgctcccc gtgcaggtcc

20

<210> 39

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 39

gtccctgtac acgatgcgct

20

<210> 40

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 40

atccagcaag atggttctcgg

20

<210> 41

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 41

gccgtggtca tccagcaaga

20

<210> 42

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 42

catgcacagc tagtcccagg

20

<210> 43

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 43

gccctcgggc acatgcacag

20

<210> 44

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 44

cctttgatgg tctggccctc

20

<210> 45

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 45

gtaaccacacg gtgcccacac

20

<210> 46

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 46

ctccggagcc atgtaaccca

20

<210> 47

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 47

tcattcttca ccacctcggg

20

<210> 48

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 48

gggctgaacg tgtaccgttc

20

<210> 49

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 49

gcgactggcc tgcgatcatc

20

<210> 50

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 50

aggagctgtg agcaaagtga

20

<210> 51

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 51

gttcagcttc ttaaagaggg

20

<210> 52

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 52

aacggcggct ccagcatgcc

20

<210> 53

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 53

ccttgcagta aatggcctgg

20

<210> 54

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 54

tcaatgtcca gaacatcctt

20

<210> 55

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 55

tagagaactg ttcaatgtcc

20

<210> 56

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 56

agctccacgc ccttgaccgt

20

<210> 57

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 57

aaacttctgg tagaagtcct

20

<210> 58

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 58

ggatgggcac actgcctgtg

20

<210> 59

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 59

gtctccacca tctcgttctg

20

<210> 60

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 60

gccccaaagac attcagctct

20

<210> 61

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 61

ctggcccttc cagtccaggt

20

<210> 62

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 62

cttttttagga ggtgcaggtg

20

<210> 63

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 63

agtttccaca gcaatcttgg

20

<210> 64

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 64

gctcggagta gctgctaccg

20

<210> 65

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 65

gaagatcact gtgcaaaact

20

<210> 66

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 66

gccacgactt gagtggacaa

20

<210> 67

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 67

tgttccccag gccacgactt

20

<210> 68

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 68

ctgagggacg gaggacactg

20

<210> 69

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 69

gccaaactca gccaggccag

20

<210> 70

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 70

ggatggccca ggcctgcca

20

<210> 71

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 71

cggcgggaat taagaatgct

20

<210> 72

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 72

tggctgcccc caggagccaa

20

<210> 73

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 73

tggcagtggc tcctctgcc

20

<210> 74

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 74

tgtgaccacc ctcagccatc

20

<210> 75

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 75

gaaggctcag gtgtgaccac

20

<210> 76

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 76

acagtgctga aggctcaggt

20

<210> 77

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 77

acagctttgg gccccagagg

20

<210> 78

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 78

ggcctagggc atacagaggc

20

<210> 79

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 79

ggccactttg gcacaggcct

20

<210> 80

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 80

aaggcagagg cagacctatc

20

<210> 81

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 81

aatgtttgag gggtgaccag

20

<210> 82

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 82

ttattgctgg gcacatgtgt

20

<210> 83

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 83

tcaccacac aggtacacac

20

<210> 84

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 84

aacactggga ccaggatggc

20

<210> 85

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 85

catgctaccc taacactggg

20

<210> 86

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 86

aaaattggct ttaaatatgg

20

<210> 87

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 87

acgagtaata aaaattggct

20

<210> 88

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Antisense Oligonucleotide

<400> 88

tttacgttgg acaaaacgag

20

<210> 89

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 89

tagcgaacac ggtgctactc

20

<210> 90

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 90

gcagatgctc cagttccctc

20

<210> 91

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 91

cacatcagcc agtgcaaga

20

<210> 92

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 92

agctgaggct cagcctcgag

20

<210> 93

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 93

gagcgtgact atcacagcct

20

<210> 94

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 94

tatcacagcc tgtgcgagcg

20

<210> 95

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 95

atcttctgag ccacacgggt

20

<210> 96

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 96

ggagcagggt ccctgcaaag

20

<210> 97

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 97

cagcatctac ttcaaccggt

20

<210> 98

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 98

accgagtcct gggcaaaggt

20

<210> 99

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 99

gctgaacgag aagcagatcc

20

<210> 100

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 100

ggttttagt gagcttggcc

20

<210> 101

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 101

gcctatgaga ccaaggacgc

20

<210> 102

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 102

gacgcgctgt gcctgggtgct

20

<210> 103

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 103

ttccacatct accacatggg

20

<210> 104

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 104

ggacctgcac cgggagcgca

20

<210> 105

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 105

agcgcacatcgt gtacagggac

20

<210> 106

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 106

tcttgctgga tgaccacggc

20

<210> 107

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 107

cctgggacta gctgtgcatg

20

<210> 108

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 108

ctgtgcatgt gcccgagggc

20

<210> 109

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 109

gagggccaga ccatcaaagg

20

<210> 110

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 110

gtgtgggcac cgtgggttac

20

<210> 111

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 111

tgggttacat ggctccggag

20

<210> 112

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 112

ccggaggtgg tgaagaatga

20

<210> 113

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 113

gaacggtaca cgttcagccc

20

<210> 114

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 114

tcactttgct cacagctcct

20

<210> 115

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 115

ggcatgctgg agccgccggt

20

<210> 116

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 116

ccaggccatt tactgcaagg

20

<210> 117

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 117

ggacattgaa cagttctcta

20

<210> 118

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 118

cggtagcagc tactccgagc

20

<210> 119

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 119  
agttttgcac agtgatcttc 20

<210> 120  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 120  
ttgtccactc aagtcgtggc 20

<210> 121  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 121  
aagtcgtggc ctggggaaca 20

<210> 122  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 122  
cagtgtcctc cgtccctcag 20

<210> 123

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 123

ctggcctggc tgagtttggc

20

<210> 124

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 124

tggcagggcc tggccatcc

20

<210> 125

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 125

agcattctta attcccgccg

20

<210> 126

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 126  
acctgagcct tcagcactgt 20

<210> 127  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 127  
cctctggggc ccaaagctgt 20

<210> 128  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 128  
gcctctgtat gcctaggcc 20

<210> 129  
<211> 20  
<212> DNA  
<213> H. sapiens

<220>

<400> 129  
aggcctgtgc caaagtggcc 20

<210> 130  
<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 130

gataggtctg cctctgcctt

20

<210> 131

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 131

ctggtcaacc ctcaaacatt

20

<210> 132

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 132

acacatgtgc ccagcaataa

20

<210> 133

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 133

cccagtggtta gggtagcatg

20

<210> 134

<211> 20

<212> DNA

<213> H. sapiens

<220>

<400> 134

agccaatttt tattactcgt

20