

(19) **DANMARK**

(10) **DK/EP 1694846 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: *C 12 N 15/31 (2006.01)* *C 07 K 14/32 (2006.01)* *C 12 N 1/21 (2006.01)*
C 12 N 9/28 (2006.01) *C 12 N 15/56 (2006.01)* *C 12 N 15/75 (2006.01)*
C 12 P 21/02 (2006.01) *C 12 N 1/21 (2006.01)* *C 12 R 1/07 (2006.01)*
- (45) Oversættelsen bekendtgjort den: **2014-11-17**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2014-10-01**
- (86) Europæisk ansøgning nr.: **04803012.6**
- (86) Europæisk indleveringsdag: **2004-12-10**
- (87) Den europæiske ansøgnings publiceringsdag: **2006-08-30**
- (86) International ansøgning nr.: **DK2004000859**
- (87) Internationalt publikationsnr.: **WO2005056799**
- (30) Prioritet: **2003-12-10 DK 200301824**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU MC NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (72) Opfinder: **NIELSEN, Allan, Kent, Rolf Krakes Allé 19, 2860 Søborg, Danmark**
RASMUSSEN, Michael, Dolberg, Syv højvaenge 151, 2625 Vallensbæk, Danmark
- (74) Fuldmægtig i Danmark: **Novozymes A/S Patents, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (54) Benævnelse: **Celle med forbedret sekretion formidlet af MrgA protein eller en homolog**
- (56) Fremdragne publikationer:
WO-A-94/19471
WO-A-99/04007
CHEN L ET AL: "Bacillus subtilis MrgA is a Dps(PexB) homologue: evidence for metalloregulation of an oxidative-stress gene." MOLECULAR MICROBIOLOGY. OCT 1995, vol. 18, no. 2, October 1995 (1995-10), pages 295-300, XP002319117 ISSN: 0950-382X
VAN WELY K H ET AL: "The carboxyl terminus of the Bacillus subtilis SecA is dispensable for protein secretion and viability." MICROBIOLOGY (READING, ENGLAND) OCT 2000, vol. 146 (Pt 10), October 2000 (2000-10), pages 2573-2581, XP002319104 ISSN: 1350-0872 cited in the application
DATABASE EMBL 8 September 1993 (1993-09-08), CHEN L. ET AL.: "Bacillus subtilis MrgA (mrgA) gene, complete cds and orf, partial cds." XP002319105 Database accession no. L19547

Description**FIELD OF THE INVENTION**

[0001] In the industrial production of secreted polypeptides it is of interest to achieve a product yield as high as possible. Accordingly, it is highly desirable to remove any potential bottlenecks from the secretory machinery of production cells. To this end it is well-known that it can be advantageous to overexpress one or more gene(s) encoding protein(s) involved with secretion, e.g., PrsA protein or functional homologues thereof. The present invention relates to a *Bacillus* cell which overexpresses a heterologous MrgA protein or functional MrgA protein homologue, with an amino acid sequence being at least 70% identical to SEQ ID No. 2.

BACKGROUND OF THE INVENTION

[0002] WO 99/04007 discloses a method for enhancing secretion of heterologous proteins, such as enzymes, growth factors or hormones, in a Gram-positive microorganism, in particular in *Bacillus*, by overexpression of the secretion factors SecDF (cf. abstract; page 2, lines 12-31). WO 94/19471 discloses a method for enhancing secretion of a heterologous exoprotein in Gram-positive bacteria such as *Bacillus* sp. by overexpression of the PrsA protein in the bacterial host (cf. abstract; page 4, lines 8-25). Still, there remains a need to find alternative ways of enhancing secretion of heterologous proteins.

[0003] The MrgA protein of *Bacillus subtilis* was originally classified as a Dps(PexB) homologue, encoded by a metalloregulated oxidative-stress gene (metallo regulated gene) *mrgA*. One purported function of the MrgA protein in *B. subtilis* is to bind DNA under conditions of oxidative stress and to protect the DNA against damage (Chen L, Helmann JD. 1995. *Bacillus subtilis* MrgA is a Dps(PexB) homologue: evidence for metalloregulation of an oxidative stress gene. *Mol Microbiol* 18: 295-300).

[0004] A *B. subtilis mrgA* deletion mutant only had a somewhat reduced overall level of secreted proteins, and it was therefore broadly concluded that MrgA is not involved in protein secretion in *B. subtilis*, (van Wely KH, Swaving J, Klein M, Freudl R,

Driessen AJ. 2000. The carboxyl terminus of the *Bacillus subtilis* SecA is dispensable for protein secretion and viability. *Microbiology* 146: 2573-81).

[0005] However, the present inventors have found, as demonstrated herein, that MrgA is in fact involved in secretion in *Bacillus*, and that a higher expression of *mrgA* leads to a higher secretion of an exoenzyme, exemplified below by improved secretion of a heterologous alpha-amylase.

SUMMARY OF THE INVENTION

[0006] Severe secretion stress was imposed on a *Bacillus subtilis* cell by overexpressing an exoenzyme, the alpha-amylase AmyQ of *Bacillus amyloliquefaciens*, encoded by a plasmid-borne constitutively expressed gene in the cell. DNA microarray analyses revealed an increased expression of the general stress protein *mrgA* as a response to the imposed secretion stress.

[0007] The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis*. Three PCR reactions were carried out with three different upstream PCR primers, each comprising the sequence of a synthetic constitutive promoter of different strength. The three amplified PCR fragments were integrated into the chromosome of individual *B. subtilis* strains, resulting in three recombinant strains, each expressing *mrgA* from the native locus, and from the integrated *mrgA* copy, which was transcribed from the synthetic promoter.

[0008] The three strains overexpressing *mrgA*, and a corresponding control strain, were then transformed with the plasmid pKTH10, which carries and constitutively overexpresses the gene encoding the alpha-amylase AmyQ of *B. amyloliquefaciens*.

[0009] The yields of secreted AmyQ amylase from the transformed strains were determined after 1 weeks cultivation in 200 ml BPX culture flasks. For each of the three MrgA overexpressing strains, and the control strain, three independent isolates were analysed in triplicate to determine the amylase yields. The yields of secreted amylase from the MrgA overexpressing strains were 27% - 44% higher than the yield from the control strain.

[0010] Accordingly, in a first aspect the invention relates to a recombinant *Bacillus* cell for production of a heterologous exoprotein of interest, wherein the cell comprises a nucleic acid construct encoding the heterologous exoprotein of interest and:

1. a) a heterologous promoter operably linked with at least one gene encoding MrgA protein with an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2; or
2. b) at least one heterologous gene encoding MrgA protein with an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2.

[0011] In a second aspect, the invention relates to a method for producing a heterologous exoprotein of interest, comprising the steps of:

1. a) cultivating a cell as defined in the first aspect; and
2. b) recovering the heterologous exoprotein.

[0012] In the present context, a functional homologue of the MrgA protein is a protein, which when expressed at a higher level in a cell, leads to an increased secretion of an exoenzyme, such as an alpha-amylase, when compared with an otherwise identical cell with normal expression of the MrgA functional homologue cultivated under essentially identical conditions. In addition, the functional homologue of the MrgA protein shares an amino acid sequence identity with the MrgA protein of at least 50%, preferably 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, or most preferably 99% when aligned as described above.

DEFINITIONS

[0013] In accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (herein

"Sambrook et al., 1989") DNA Cloning: A Practical Approach, Volumes I and II/D.N. Glover ed. 1985); Oligonucleotide Synthesis (M.J. Gait ed. 1984); Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds (1985)); Transcription And Translation (B.D. Hames & S.J. Higgins, eds. (1984)); Animal Cell Culture (R.I. Freshney, ed. (1986)); Immobilized Cells And Enzymes (IRL Press, (1986)); B. Perbal, A Practical Guide To Molecular Cloning (1984).

[0014] A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized in vitro, or prepared from a combination of natural and synthetic molecules.

[0015] A "nucleic acid molecule" or "nucleotide sequence" refers to the phosphate ester polymeric form of ribonucleosides (adenosine, guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine, deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules") in either single stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and RNA-RNA helices are possible. The term nucleic acid molecule, and in particular DNA or RNA molecule, refers only to the primary and secondary structure of the molecule, and does not limit it to any particular tertiary or quaternary forms. Thus, this term includes double-stranded DNA found, inter alia, in linear or circular DNA molecules (e.g., restriction fragments), plasmids, and chromosomes. In discussing the structure of particular double-stranded DNA molecules, sequences may be described herein according to the normal convention of giving only the sequence in the 5' to 3' direction along the nontranscribed strand of DNA (i.e., the strand having a sequence homologous to the mRNA). A "recombinant DNA molecule" is a DNA molecule that has undergone a molecular biological manipulation.

[0016] A DNA "coding sequence" or an "open reading frame (ORF)" is a double-stranded DNA sequence which is transcribed and translated into a polypeptide in a cell in vitro or in vivo when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxyl) terminus.

A coding sequence can include, but is not limited to, prokaryotic sequences, cDNA from eukaryotic mRNA, genomic DNA sequences from eukaryotic (e.g., mammalian) DNA, and even synthetic DNA sequences. If the coding sequence is intended for expression in a eukaryotic cell, a polyadenylation signal and transcription termination sequence will usually be located 3' to the coding sequence.

[0017] An expression vector is a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments may include promoter and terminator sequences, and optionally one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, and the like. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

[0018] Transcriptional and translational control sequences are DNA regulatory sequences, such as promoters, enhancers, terminators, and the like, that provide for the expression of a coding sequence in a host cell. In eukaryotic cells, polyadenylation signals are control sequences.

[0019] A "secretory signal sequence" is a DNA sequence that encodes a polypeptide (a "secretory peptide" that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

[0020] The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

[0021] A chromosomal gene is rendered non-functional if the polypeptide that the gene encodes can no longer be expressed in a functional form. Such non-functionality of a gene can be induced by a wide variety of genetic manipulations as known in the art, some of which are described in Sambrook et al. *vide supra*. Partial

deletions within the ORF of a gene will often render the gene non-functional, as will mutations.

[0022] The term "an expressible copy of a chromosomal gene" is used herein as meaning a copy of the ORF of a chromosomal gene, wherein the ORF can be expressed to produce a fully functional gene product. The expressible copy may not be transcribed from the native promoter of the chromosomal gene, it may instead be transcribed from a foreign or heterologous promoter, or it may indeed be promoterless and expressed only by transcriptional read-through from a gene present upstream of the 5' end of the ORF. Transcriptional read-through is intended to have the same meaning here as the generally recognized meaning in the art.

[0023] "Operably linked", when referring to DNA segments, indicates that the segments are arranged so that they function in concert for their intended purposes, e.g. transcription initiates in the promoter and proceeds through the coding segment to the terminator.

[0024] A coding sequence is "under the control" of transcriptional and translational control sequences in a cell when RNA polymerase transcribes the coding sequence into mRNA, which is then trans-RNA spliced and translated into the protein encoded by the coding sequence.

[0025] "Heterologous" DNA refers to DNA not naturally located in the cell, or in a chromosomal site of the cell. Preferably, the heterologous DNA includes a gene foreign to the cell.

[0026] As used herein the term "nucleic acid construct" is intended to indicate any nucleic acid molecule of cDNA, genomic DNA, synthetic DNA or RNA origin. The term "construct" is intended to indicate a nucleic acid segment which may be single- or double-stranded, and which may be based on a complete or partial naturally occurring nucleotide sequence encoding a polypeptide of interest. The construct may optionally contain other nucleic acid segments.

[0027] The nucleic acid construct encoding the polypeptide may suitably be of

genomic or cDNA origin, for instance obtained by preparing a genomic or cDNA library and screening for DNA sequences coding for all or part of the polypeptide by hybridization using synthetic oligonucleotide probes in accordance with standard techniques (cf. Sambrook et al., *supra*).

[0028] The nucleic acid construct encoding the polypeptide may also be prepared synthetically by established standard methods, e.g. the phosphoramidite method described by Beaucage and Caruthers, *Tetrahedron Letters* 22 (1981), 1859 - 1869, or the method described by Matthes et al., *EMBO Journal* 3 (1984), 801 - 805. According to the phosphoramidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in suitable vectors.

[0029] Furthermore, the nucleic acid construct may be of mixed synthetic and genomic, mixed synthetic and cDNA or mixed genomic and cDNA origin prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate), the fragments corresponding to various parts of the entire nucleic acid construct, in accordance with standard techniques. The nucleic acid construct may also be prepared by polymerase chain reaction using specific primers, for instance as described in US 4,683,202 or Saiki et al., *Science* 239 (1988), 487 - 491.

[0030] The term nucleic acid construct may be synonymous with the term "expression cassette" when the nucleic acid construct contains the control sequences necessary for expression of a coding sequence of the present invention

[0031] The term "control sequences" is defined herein to include all components which are necessary or advantageous for expression of the coding sequence of the nucleic acid sequence. Each control sequence may be native or foreign to the nucleic acid sequence encoding the polypeptide. Such control sequences include, but are not limited to, a leader, a polyadenylation sequence, a propeptide sequence, a promoter, a signal sequence, and a transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of

introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the nucleic acid sequence encoding a polypeptide.

[0032] The control sequence may be an appropriate promoter sequence, a nucleic acid sequence which is recognized by a host cell for expression of the nucleic acid sequence. The promoter sequence contains transcription and translation control sequences which mediate the expression of the polypeptide. The promoter may be any nucleic acid sequence which shows transcriptional activity in the host cell of choice and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0033] The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleic acid sequence encoding the polypeptide. Any terminator which is functional in the host cell of choice may be used in the present invention.

[0034] The control sequence may also be a polyadenylation sequence, a sequence which is operably linked to the 3' terminus of the nucleic acid sequence and which, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence which is functional in the host cell of choice may be used in the present invention.

[0035] The control sequence may also be a signal peptide coding region, which codes for an amino acid sequence linked to the amino terminus of the polypeptide which can direct the expressed polypeptide into the cell's secretory pathway of the host cell. The 5' end of the coding sequence of the nucleic acid sequence may inherently contain a signal peptide coding region naturally linked in translation reading frame with the segment of the coding region which encodes the secreted polypeptide. Alternatively, the 5' end of the coding sequence may contain a signal peptide coding region which is foreign to that portion of the coding sequence which encodes the secreted polypeptide. A foreign signal peptide coding region may be required where the coding sequence does not normally contain a signal peptide coding region. Alternatively, the foreign signal peptide coding region may simply

replace the natural signal peptide coding region in order to obtain enhanced secretion of the exoprotein relative to the natural signal peptide coding region normally associated with the coding sequence. The signal peptide coding region may be obtained from a glucoamylase or an amylase gene from an *Aspergillus* species, a lipase or proteinase gene from a *Rhizomucor* species, the gene for the alpha-factor from *Saccharomyces cerevisiae*, an amylase or a protease gene from a *Bacillus* species, or the calf preprochymosin gene. However, any signal peptide coding region capable of directing the expressed polypeptide into the secretory pathway of a host cell of choice may be used in the present invention.

[0036] The control sequence may also be a propeptide coding region, which codes for an amino acid sequence positioned at the amino terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to mature active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding region may be obtained from the *Bacillus subtilis* alkaline protease gene (*aprE*), the *Bacillus subtilis* neutral protease gene (*nprT*), the *Saccharomyces cerevisiae* alpha-factor gene, or the *Myceliophthora thermophilum* laccase gene (WO 95/33836).

[0037] It may also be desirable to add regulatory sequences which allow the regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those which cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems would include the *lac*, *tac*, and *trp* operator systems.

[0038] Examples of suitable promoters for directing the transcription of the gene(s) of the present invention, especially in a bacterial host cell, are the promoters obtained from the *E. coli* *lac* operon, the *Streptomyces coelicolor* agarase gene (*dagA*), the *Bacillus subtilis* levansucrase gene (*sacB*), the *Bacillus subtilis* alkaline protease gene, the *Bacillus licheniformis* alpha-amylase gene (*amyL*), the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), the *Bacillus amyloliquefaciens* BAN amylase gene, the

Bacillus licheniformis penicillinase gene (penP), the *Bacillus subtilis* xylA and xylB genes, and the prokaryotic beta-lactamase gene (Villa-Kamaroff et al., 1978, Proceedings of the National Academy of Sciences USA 75:3727-3731), as well as the tac promoter (DeBoer et al., 1983, Proceedings of the National Academy of Sciences USA 80:21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242:74-94; and in Sambrook et al., 1989, supra.

[0039] An effective signal peptide coding region for bacterial host cells is the signal peptide coding region obtained from the maltogenic amylase gene from *Bacillus* NCIB 11837, the *Bacillus stearothermophilus* alpha-amylase gene, the *Bacillus licheniformis* subtilisin gene, the *Bacillus licheniformis* beta-lactamase gene, the *Bacillus stearothermophilus* neutral proteases genes (nprT, nprS, nprM), and the *Bacillus subtilis* PrsA gene. Further signal peptides are described by Simonen and Palva, 1993, Microbiological Reviews 57:109-137.

[0040] The present invention also describes to recombinant expression vectors comprising a nucleic acid sequence, a promoter, and transcriptional and translational stop signals. The various nucleic acid and control sequences described above may be joined together to produce a recombinant expression vector which may include one or more convenient restriction sites to allow for insertion or substitution of the nucleic acid sequence encoding the polypeptide at such sites. Alternatively, the nucleic acid sequence may be expressed by inserting the nucleic acid sequence or a nucleic acid construct comprising the sequence into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression, and possibly secretion.

[0041] The recombinant expression vector may be any vector (e.g., a plasmid or virus) which can be conveniently subjected to recombinant DNA procedures and can bring about the expression of the nucleic acid sequence. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vectors may be linear or closed circular plasmids. The vector may be an autonomously replicating vector, i.e., a vector which exists as an

extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g., a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one which, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. The vector system may be a single vector or plasmid or two or more vectors or plasmids which together contain the total DNA to be introduced into the genome of the host cell, or a transposon.

[0042] The vectors preferably contain one or more selectable markers which permit easy selection of transformed cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0043] Antibiotic selectable markers confer antibiotic resistance to such antibiotics as ampicillin, kanamycin, chloramphenicol, tetracycline, neomycin, hygromycin or methotrexate.

[0044] The vectors preferably contain an element(s) that permits stable integration of the vector, or of a smaller part of the vector, into the host cell genome or autonomous replication of the vector in the cell independent of the genome of the cell.

[0045] The vectors, or smaller parts of the vectors such as amplification units of the present invention, may be integrated into the host cell genome when introduced into a host cell. For chromosomal integration, the vector may rely on the nucleic acid sequence encoding the polypeptide or any other element of the vector for stable integration of the vector into the genome by homologous or nonhomologous recombination.

[0046] Alternatively, the vector may contain additional nucleic acid sequences for directing integration by homologous recombination into the genome of the host cell. The additional nucleic acid sequences enable the vector to be integrated into the host cell genome at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should

preferably contain a sufficient number of nucleic acids, such as 100 to 1,500 base pairs, preferably 400 to 1,500 base pairs, and most preferably 800 to 1,500 base pairs, which are highly homologous with the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding nucleic acid sequences; specific examples of encoding sequences suitable for site-specific integration by homologous recombination are given in WO 02/00907 (Novozymes, Denmark).

[0047] On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination. These nucleic acid sequences may be any sequence that is homologous with a target sequence in the genome of the host cell, and, furthermore, may be non-encoding or encoding sequences. The copy number of a vector, an expression cassette, an amplification unit, a gene or indeed any defined nucleotide sequence is the number of identical copies that are present in a host cell at any time. A gene or another defined chromosomal nucleotide sequence may be present in one, two, or more copies on the chromosome. An autonomously replicating vector may be present in one, or several hundred copies per host cell.

[0048] An amplification unit of the invention is a nucleotide sequence that can integrate into the chromosome of a host cell, whereupon it can increase in number of chromosomally integrated copies by duplication or multiplication. The unit comprises an expression cassette as defined herein comprising at least one copy of a gene of interest and an expressible copy of a chromosomal gene, as defined herein, of the host cell. When the amplification unit is integrated into the chromosome of a host cell, it is defined as that particular region of the chromosome which is prone to being duplicated by homologous recombination between two directly repeated regions of DNA. The precise border of the amplification unit with respect to the flanking DNA is thus defined functionally, since the duplication process may indeed duplicate parts of the DNA which was introduced into the chromosome as well as parts of the endogenous chromosome itself, depending on the exact site of recombination within the repeated regions. This principle is illustrated in Janni re et al. (1985, Stable gene amplification in the chromosome of *Bacillus subtilis*. *Gene*, 40: 47-55).

[0049] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, pACYC184, pUB110, pE194, pTA1060, and pAMBeta1. Examples of origin of replications for use in a yeast host cell are the 2 micron origin of replication, the combination of CEN6 and ARS4, and the combination of CEN3 and ARS1. The origin of replication may be one having a mutation which makes its functioning temperature-sensitive in the host cell (see, e.g., Ehrlich, 1978, Proceedings of the National Academy of Sciences USA 75:1433).

[0050] The present invention also describes recombinant host cells, comprising a nucleic acid sequence, which are advantageously used in the recombinant production of the polypeptides. The term "host cell" encompasses any progeny of a parent cell which is not identical to the parent cell due to mutations that occur during replication.

[0051] The cell is preferably transformed with a vector comprising a nucleic acid sequence of the invention followed by integration of the vector into the host chromosome. "Transformation" means introducing a vector comprising a nucleic acid sequence of the present invention into a host cell so that the vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector. Integration is generally considered to be an advantage as the nucleic acid sequence is more likely to be stably maintained in the cell. Integration of the vector into the host chromosome may occur by homologous or non-homologous recombination as described above.

[0052] The transformation of a bacterial host cell may, for instance, be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, Molecular General Genetics 168:111-115), by using competent cells (see, e.g., Young and Spizizin, 1961, Journal of Bacteriology 81:823-829, or Dubnar and Davidoff-Abelson, 1971, Journal of Molecular Biology 56:209-221), by electroporation (see, e.g., Shigekawa and Dower, 1988, Biotechniques 6:742-751), or by conjugation (see, e.g., Koehler and Thorne, 1987, Journal of Bacteriology 169:5771-5278).

[0053] The transformed or transfected host cells described above are cultured in a suitable nutrient medium under conditions permitting the expression of the desired polypeptide, after which the resulting polypeptide is recovered from the cells, or the culture broth.

[0054] The medium used to culture the cells may be any conventional medium suitable for growing the host cells, such as minimal or complex media containing appropriate supplements. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. in catalogues of the American Type Culture Collection). The media are prepared using procedures known in the art (see, e.g., references for bacteria and yeast; Bennett, J.W. and LaSure, L., editors, *More Gene Manipulations in Fungi*, Academic Press, CA, 1991).

[0055] If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it is recovered from cell lysates. The polypeptide are recovered from the culture medium by conventional procedures including separating the host cells from the medium by centrifugation or filtration, precipitating the proteinaceous components of the supernatant or filtrate by means of a salt, e.g. ammonium sulphate, purification by a variety of chromatographic procedures, e.g. ion exchange chromatography, gelfiltration chromatography, affinity chromatography, or the like, dependent on the type of polypeptide in question.

[0056] The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide.

[0057] The polypeptides may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing (IEF), differential solubility (e.g., ammonium

sulfate precipitation), or extraction (see, e.g., Protein Purification, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).

[0058] In the present context, the term "substantially pure polypeptide" means a polypeptide preparation which contains at the most 10% by weight of other polypeptide material with which it is natively associated (lower percentages of other polypeptide material are preferred, e.g. at the most 8% by weight, at the most 6% by weight, at the most 5% by weight, at the most 4% at the most 3% by weight, at the most 2% by weight, at the most 1% by weight, and at the most ½% by weight). Thus, it is preferred that the substantially pure polypeptide is at least 92% pure, i.e. that the polypeptide constitutes at least 92% by weight of the total polypeptide material present in the preparation, and higher percentages are preferred such as at least 94% pure, at least 95% pure, at least 96% pure, at least 96% pure, at least 97% pure, at least 98% pure, at least 99%, and at the most 99.5% pure. The polypeptides disclosed herein are preferably in a substantially pure form. In particular, it is preferred that the polypeptides disclosed herein are in "essentially pure form", i.e. that the polypeptide preparation is essentially free of other polypeptide material with which it is natively associated. This can be accomplished, for example, by preparing the polypeptide by means of well-known recombinant methods. Herein, the term "substantially pure polypeptide" is synonymous with the terms "isolated polypeptide" and "polypeptide in isolated form".

[0059] In the present context, the homology between two amino acid sequences or between two nucleotide sequences is described by the parameter "identity". For purposes of the present invention, alignments of sequences and calculation of homology scores may be done using a full Smith-Waterman alignment, useful for both protein and DNA alignments. The default scoring matrices BLOSUM50 and the identity matrix are used for protein and DNA alignments respectively. The penalty for the first residue in a gap is -12 for proteins and -16 for DNA, while the penalty for additional residues in a gap is -2 for proteins and -4 for DNA. Alignment may be made with the FASTA package version v20u6 (W. R. Pearson and D. J. Lipman (1988), "Improved Tools for Biological Sequence Analysis", PNAS 85:2444-2448, and W. R. Pearson (1990) "Rapid and Sensitive Sequence Comparison with FASTP and FASTA", Methods in Enzymology, 183:63-98).

[0060] Multiple alignments of protein sequences may be made using "ClustalW" (Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, positions-specific gap penalties and weight matrix choice. *Nucleic Acids Research*, 22:4673-4680). Multiple alignment of DNA sequences may be done using the protein alignment as a template, replacing the amino acids with the corresponding codon from the DNA sequence.

[0061] In the present context, a functional homologue of the MrgA protein is a protein, which when expressed at a higher level in a cell, leads to an increased secretion of an exoenzyme, such as an alpha-amylase, when compared with a cell with normal expression of the MrgA functional homologue cultivated under essentially identical conditions. In addition, the functional homologue of the MrgA protein shares an amino acid sequence identity with the MrgA protein of at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, or most preferably 99% when aligned as described above.

[0062] In the present context, the term "allelic variant" denotes any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene. Allelic variants are included in the present definition of functional homologues.

[0063] The MrgA protein or functional homologue thereof may be a wild-type protein identified and isolated from a natural source. Such wild-type proteins may be specifically screened for by standard techniques known in the art. Furthermore, the MrgA protein or functional homologue thereof may be prepared by the DNA shuffling technique, such as described in J.E. Ness et al. *Nature Biotechnology* 17, 893-896 (1999). Moreover, the MrgA protein or functional homologue thereof may be an artificial variant. Such artificial variants may be constructed by standard techniques known in the art, such as by site-directed/random mutagenesis. In one embodiment of the invention, amino acid changes (in the artificial variant as well as in wild-type

polypeptides) are of a minor nature, that is conservative amino acid substitutions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of one to about 30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to about 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0064] Examples of conservative substitutions are within the group of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine, valine and methionine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine and threonine). Amino acid substitutions which do not generally alter the specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, In, *The Proteins*, Academic Press, New York. The most commonly occurring exchanges are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly as well as these in reverse.

[0065] It will be apparent to those skilled in the art that such modifications can be made outside the regions critical to the function of the molecule and still result in an active polypeptide. Amino acid residues essential to the activity of the polypeptide encoded by the nucleotide sequence of the invention, and therefore preferably not subject to modification, such as substitution, may be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (see, e.g., Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, mutations are introduced at every positively charged residue in the molecule, and the resultant mutant molecules are tested for activity to identify amino acid residues that are critical to the activity of the molecule. Sites of substrate-enzyme interaction can also be determined by analysis of the three-dimensional structure as determined by such techniques as nuclear magnetic resonance analysis, crystallography or photoaffinity labelling (see, e.g., de Vos et al., 1992, *Science* 255:

306-312; Smith et al., 1992, Journal of Molecular Biology 224: 899-904; Wlodaver et al., 1992, FEBS Letters 309: 59-64).

[0066] Moreover, a nucleotide sequence encoding a polypeptide described herein may be modified by introduction of nucleotide substitutions which do not give rise to another amino acid sequence of the polypeptide encoded by the nucleotide sequence, but which correspond to the codon usage of the host organism intended for production of the enzyme.

[0067] The introduction of a mutation into the nucleotide sequence to exchange one nucleotide for another nucleotide may be accomplished by site-directed mutagenesis using any of the methods known in the art. Particularly useful is the procedure, which utilizes a supercoiled, double stranded DNA vector with an insert of interest and two synthetic primers containing the desired mutation. The oligonucleotide primers, each complementary to opposite strands of the vector, extend during temperature cycling by means of Pfu DNA polymerase. On incorporation of the primers, a mutated plasmid containing staggered nicks is generated. Following temperature cycling, the product is treated with DpnI which is specific for methylated and hemimethylated DNA to digest the parental DNA template and to select for mutation-containing synthesized DNA. Other procedures known in the art may also be used. For a general description of nucleotide substitution, see, e.g., Ford et al., 1991, Protein Expression and Purification 2: 95-107.

DETAILED DESCRIPTION OF THE INVENTION

[0068] The first aspect of the invention relates to a recombinant *Bacillus* cell for production of a heterologous exoprotein of interest, wherein the cell comprises a nucleic acid construct encoding the heterologous exoprotein of interest and:

1. a) a heterologous promoter operably linked with at least one gene encoding MrgA protein with an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2; or
2. b) at least one heterologous gene encoding MrgA protein with an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2.

[0069] The cell of the invention produces greater amounts of MrgA protein or a functional homologue thereof, than the parent cell. A comparison should be made by cultivating the cell of the invention as well as the parent cell under essentially identical conditions, and comparing the amount of MrgA protein by any standard method in the art. Preferably the cell of the invention produces at least 5% more MrgA than the parent, more preferably at least 10%, still more preferably at least 20%, and most preferably at least 50% more MrgA protein or a functional homologue thereof than the parent. Such overproduction may be accomplished by standard means known to the art, *e.g.*, use of multicopy plasmids, multiple copies of the genes encoding MrgA or a functional homologue thereof, and/or the protein of interest, in the chromosome of the host, combined with altering the regulatory elements to increase expression, *e.g.*, use of strong promoter(s), use of multiple promoters, use of enhancers, and so forth.

[0070] As the inventors show herein, a cell of the first aspect is capable of producing greater amounts of a protein of interest than the corresponding parent cell, when both are cultivated under essentially identical conditions.

[0071] Accordingly, a preferred embodiment of the invention relates to the cell of the first aspect, which produces greater amounts of a protein of interest than the parent cell. Preferably the protein of interest is an intracellular protein or an exoprotein. Preferably the cell of the invention secretes greater amounts of an exoprotein of interest than the parent cell. Preferably the cell of the invention secretes at least 5% more exoprotein than the parent, more preferably at least 10% more, still more preferably at least 20% more, and most preferably at least 50% more exoprotein than the parent. The amount of produced or secreted protein of interest from each cell may be determined by any suitable assay in the art; a non-limiting example is shown below with the determination of secreted amounts of the exoprotein alpha-amylase.

[0072] The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source. The host cell may be a unicellular microorganism, *e.g.*, a prokaryote, or a non-unicellular microorganism, *e.g.*, a eukaryote. Useful unicellular cells are bacterial cells such as gram positive bacteria including, but not

limited to, a *Bacillus* cell, e.g., *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*; or a *Streptomyces* cell, e.g., *Streptomyces lividans* or *Streptomyces murinus*, or gram negative bacteria such as *E. coli* and *Pseudomonas sp.* In a preferred embodiment, the bacterial host cell is a *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus stearothermophilus* or *Bacillus subtilis* cell. In one preferred embodiment, the bacterial host cell is a prokaryotic cell, preferably a Gram-positive prokaryotic cell, and more preferably the bacterial Gram positive cell is a species of the genus *Bacillus*, preferably selected from the group consisting of *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*.

[0073] A preferred embodiment relates to the cell of the invention, which is a bacterial cell, preferably a prokaryotic cell, more preferably a Gram-positive cell, and most preferably of the genus *Bacillus*; still more preferably it is of a species chosen from the group consisting of *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis*.

[0074] The protein of interest may be endogenous or exogenous to the host cell, it may be a homologous protein, or a heterologous protein.

[0075] A preferred embodiment relates to the cell, wherein said protein of interest is a protease, a lipase, a cutinase, an amylase, a galactosidase, a pullulanase, a cellulase, a glucose isomerase, a protein disulphide isomerase, a CGT'ase (cyclodextrin gluconotransferase), a phytase, a glucose oxidase, a glucosyl transferase, lactase, bilirubin oxidase, a xylanase, an antigenic microbial or protozoan protein, a bacterial protein toxin, a microbial surface protein, or a viral protein.

[0076] An evolutionary homologue of the MrgA protein, an allelic variant, an artificial

variant, a shuffled variant, a species variant, and so forth, are all referred to as a "functional homologue" or the MrgA protein in the present description, and the inventors envision that such functional homologue protein(s) will be equally effective in the cell of the invention.

[0077] Specifically, a preferred embodiment relates to the cell, wherein the MrgA protein or functional homologue thereof comprises an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2, preferably at least 75%, 80%, 85%, 90%, 95%, 97%, or even 99% identical to the amino acid sequence shown in SEQ ID NO:2.

[0078] Another preferred embodiment relates to the cell of the invention, wherein the MrgA protein or functional homologue thereof comprises or consists of the amino acid sequence shown in SEQ ID NO:2.

[0079] Still another preferred embodiment relates to the cell of the invention, which comprises at least one exogenous copy of a polynucleotide encoding MrgA protein or a functional homologue thereof comprising an amino acid sequence which is at least 70% identical to the amino acid sequence shown in SEQ ID NO:2, preferably at least 75%, 80%, 85%, 90%, 95%, 97%, or even 99% identical to the amino acid sequence shown in SEQ ID NO:2.

[0080] In a preferred embodiment the cell of the invention comprises at least one exogenous copy of a polynucleotide encoding MrgA protein or a functional homologue thereof comprising or consisting of the amino acid sequence shown in SEQ ID NO:2.

[0081] As described above, and exemplified herein, one preferred embodiment relates to a cell, wherein at least one exogenous copy of a gene encoding the MrgA protein or a functional homologue thereof is transcribed from one or more heterologous and/or artificial promoter.

[0082] In a preferred cell, at least one exogenous copy of a gene encoding the MrgA

protein or a functional homologue thereof is integrated into the genome of the cell; or is present on an extra-chromosomal construct, preferably a plasmid.

[0083] Another aspect of the invention relates to a method for producing a heterologous exoprotein of interest, comprising the steps of:

1. a) cultivating a cell as defined in the first aspect; and
2. b) recovering the heterologous exoprotein.

[0084] In the production methods of the present invention, the cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

[0085] The polypeptides may be detected using methods known in the art that are specific for the polypeptides. These detection methods may include use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide as described herein.

[0086] The polypeptides may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, centrifugation, filtration, extraction, spray-drying,

evaporation, precipitation, or extraction (see, e.g., Protein Purification, J.-C. Janson and Lars Ryden, editors, VCH Publishers, New York, 1989).

[0087] In a preferred embodiment of the method of the second aspect, the manipulated cell produces greater amounts of a protein of interest than the non-manipulated parent cell. Preferably the protein of interest is an intracellular protein or an exoprotein.

[0088] Another preferred embodiment relates to the method of the second aspect, wherein said method comprises the steps of:

1. a) identifying a gene from the parent cell that encodes MrgA protein or a functional homologue thereof; and
2. b) manipulating the cell to increase the expression of the gene identified in step (a), whereby the manipulated progeny cell expresses greater amounts of MrgA protein or functional homologue thereof, than the non-manipulated cell.

EXAMPLES

Materials and Methods

Strains

[0089]

B.subtilis 168: F. Kunst et. al. "The complete genome sequence of the Gram-positive bacterium *Bacillus subtilis*". Nature (1997) 390:249-256.

B.subtilis AN53: *B.subtilis* 168 with plasmid pKTH10 and P920mrgA integrated in the *amyE* locus (this study).

B.subtilis AN36: *B.subtilis* 168 with plasmid p920mrgA integrated into the *amyE* locus (this study).

B.subtilis AN42: *B.subtilis* 168 with plasmid p740mrgA integrated into the *amyE* locus (this study).

B.subtilis AN50: *B.subtilis* 168 with plasmid p726mrgA integrated into the *amyE* locus (this study).

B.subtilis AN55: *B.subtilis* 168 with plasmid pKTH10 and P740mrgA integrated in the *amyE* locus (this study).

B.subtilis AN57: *B.subtilis* 168 with plasmid pKTH10 and P726mrgA integrated in the *amyE* locus (this study).

B.subtilis AN83: *B.subtilis* 168 with plasmid pKTH10 (this study).

B.subtilis AN214: *B.subtilis* 168 (pel::PconsBAN)

B.subtilis AN217: *B.subtilis* 168 (pel::PconsBAN; amyE::P726mrgA)

B.subtilis AN218: *B.subtilis* 168 (pel::PconsBAN; amyE::P740mrgA)

B.subtilis AN219: *B.subtilis* 168 (pel::PconsBAN; amyE::P920mrgA)

[0090] Competent cells were prepared and transformed as described by Yasbin, R.E., Wilson, G.A. and Young, F.E. (1975) Transformation and transfection in lysogenic strains of *Bacillus subtilis* : evidence for selective induction of prophage in competent cells. J. Bacteriol, 121:296-304.

Plasmids

[0091]

pKTH10: Vehmaanpera J, Steinborn G, Hofemeister J.: " Genetic manipulation of *Bacillus amyloliquefaciens*." J Biotechnol. 1991. 19(2-3):221-40. This plasmid constitutively express the *Bacillus amyloliquefaciens* alpha-amylase (AmyQ).

pDG268neo: This plasmid is a pDG268 derivative, which is unable to replicate in *Bacillus subtilis* (Antoniewski, C., Savelli, B., and Stragier, P., 1990, J. Bact 172). The plasmid contains the chloramphenicol (cam) resistance marker next to *SfiI* and *BamHI* restriction enzyme recognition sequences, flanked by the "5'" and "3'" portions of the *amyE* locus of *Bacillus subtilis*. This plasmid is used for introduction of the MrgA expression cassette and the cam marker into the *amyE* locus of *B. subtilis* via double homologous recombination cross-over. The sequence of pDG268neo is shown in SEQ ID NO: 3.

pAN213: This plasmid is a pDG268 derivative (Antoniewski, C., Savelli, B., and Stragier, P., 1990, J. Bact 172) which is unable to replicate in *Bacillus subtilis*. The plasmid contains the erythromycin resistance marker next to *sacII* and *StyI* restriction enzyme recognition sequences. All of this sequence is flanked by the "5'" and "3'" portions of the pectate lyase (pel) locus of *Bacillus subtilis* 168.

pAN213ban: The amyQ gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers AN162 and AN163c. The upstream primer (AN162) encoded the synthetic promoter PconsBAN. The PCR product was cut with restriction enzymes *Sac*II and *Mlu*I and ligated to the large *Sac*II-*Mlu*I fragment of pAN213, resulting in plasmid pAN213ban. This plasmid is used for introduction of the AmyQ expression cassette and the erm marker into the *pel* locus of *B. subtilis* 168 via a double cross-over event. The sequence of pAN213ban is shown in SEQ ID NO: 16.

p920mrgA: The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers p920mrgaF2 (SEQ ID NO: 4) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p920mrgaF2; SEQ ID NO: 4) comprised a synthetic constitutive promoter, P920 (SEQ ID NO: 6). The PCR product shown in SEQ ID NO: 7 was cut with restriction enzymes *Sfi*I and *Bam*HI and ligated to the large *Sfi*I-*Bam*HI fragment of pDG268neo, resulting in plasmid p920mrgA.

p740mrgA: The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers p740mrgaF2 (SEQ ID NO: 8) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p740mrgaF2; SEQ ID NO: 8) comprised a synthetic constitutive promoter, P740 (SEQ ID NO: 9). The PCR product shown in SEQ ID NO: 10 was cut with restriction enzymes *Sfi*I and *Bam*HI and ligated to the large *Sfi*I-*Bam*HI fragment of pDG268neo, resulting in plasmid p740mrgA.

p726mrgA: The *mrgA* was amplified by PCR from the chromosome of *B. subtilis* by use of primers p726mrgaF2 (SEQ ID NO: 11) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p726mrgaF2; SEQ ID NO: 11) comprised the constitutive synthetic promoter P726 (SEQ ID NO: 12). The PCR product shown in SEQ ID NO: 13 was cut with restriction enzymes *Sfi*I and *Bam*HI and ligated to the large *Sfi*I-*Bam*HI fragment of pDG268neo, resulting in plasmid p726mrgA.

Primers:**[0092]**

P920mrgaF (SEQ ID NO: 4):

ctgaggccaattagccaagtttattcttgacattaggggaacatgcatgatat

aataggtaaagtaaacagatcacaaggaggacgttatc

P740mrgaF (SEQ ID NO: 8):

ctgaggccaattagggccggaagttgtgacacagctccaggatacaaat

ataatgggtcgactaaacagatcacaaggaggacgttatc

P726mrgaF (SEQ ID NO: 11):

: ctgaggccaattagggccgaggtgagatttgacactagtaggctacgggac

tataatcggggaagtaaacagatcacaaggaggacgttatc

MBmrgaR2 (SEQ ID NO: 5): tgaaggatccacgcgtccagcagacagaaagcag

AN162 (SEQ ID NO: 14):

agactgtccgcggtgtaaaaaataggaataaaggggggtgacatttttactgatatgtata

atataatttgataagaaaatgag

AN163c (SEQ ID NO: 15): gcatacacgcgtgtgtcacacctgatgccgacc

General molecular biology methods

[0093] Unless otherwise mentioned the DNA manipulations and transformations were performed using standard methods of molecular biology (Sambrook et al. (1989) Molecular cloning: A laboratory manual, Cold Spring Harbor lab., Cold Spring Harbor, NY; Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995; Harwood, C. R., and Cutting, S. M. (eds.) "Molecular Biological Methods for Bacillus". John Wiley and Sons, 1990).

[0094] Enzymes for DNA manipulations were used according to the specifications of the suppliers (e.g. restriction endonucleases, ligases etc. are obtainable from New England Biolabs, Inc.).

Media**[0095]**

LB agar (as described in Ausubel, F. M. et al. (eds.) "Current protocols in Molecular Biology". John Wiley and Sons, 1995).

LBP is LB agar supplemented with 0.05 M potassium phosphate, pH 7.0

LBPG is LB agar supplemented with 0.5% Glucose and 0.05 M potassium phosphate, pH 7.0.

LBPSK is LB agar supplemented with 0.05 M potassium phosphate, pH 7.0 and 1% of skimmed milk.

BPX media is described in EP 0 506 780 (WO 91/09129).

Fermentations

[0096] Fermentations to evaluate the amylase yields were performed in shakeflasks with 100 ml BPX at 37°C, 300 rpm for seven days. Culture volumes of 10 ml were harvested and centrifuged at 10.000 g to remove cells and debris. The clear supernants were used for assaying alpha-amylase activity.

Assay for alpha-amylase activity

[0097] Alpha-amylase activity was determined by a method employing an enzymatic colorimetric test with 4,6-ethylidene(G₇)-p-nitrophenyl(G₁)-alpha,D-maltoheptaoside (ethylidene-G₇PNP) as substrate (Boehringer Mannheim, Germany art. 1442309). Under a specified set of conditions (temp., pH, reaction time, buffer conditions) 1 mg of a given a-amylase will hydrolyse a certain amount of substrate and a yellow colour will be produced. The colour intensity is measured at 405 nm. The measured absorbance is directly proportional to the activity of the alpha-amylase in question under a given set of conditions.

EXAMPLE 1

[0098] The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers p920mrgaF2 (SEQ ID NO: 4) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p920mrgaF2) comprised the constitutive synthetic promoter P920 (SEQ ID NO:6). The PCR product (SEQ ID NO: 7) was cut with restriction enzymes *Sfi*I and *Bam*HI and ligated to the large *Sfi*I-*Bam*HI fragment of pDG268neo, resulting in plasmid p920mrgA. The ligation-mixture of p920mrgA, described in the Material and Methods section, was introduced by transformation into the *B. subtilis* 168 strain, and the transformants were plated on LBPSK media supplemented with 6 microg/ml chloramphenicol to select for integrants. Transformants growing on these plates have integrated the plasmid in the *amy* locus, either by a single (Cam⁺ kan⁺) or double cross-over event (cam⁺ kan⁻). Transformants were re-streaked on LBPSK/cam media

with and without 20 microg/ml kanamycine. Strains where double cross-over events had occurred were $\text{cam}^+ \text{kan}^-$. These strains no longer showed the tell-tale clearing zones; which signified that that integration in, and disruption of, the *amy* gene, had taken place. The site of integration was verified by PCR, the integrated copy of *mrgA* was verified by sequence analysis, and the strain was named AN36. AN36 was transformed with plasmid pKTH10 which constitutively expresses the alpha-amylase AmyQ of *Bacillus amyloliquefaciens*. The resulting strain was named AN53. Yields of amylase from AN53 were determined in triplicate from three independent isolates, and compared to the yield of amylase from the control strain AN83. Results are shown in table 1; the AN53 strain constitutively expressing *mrgA* from the synthetic promoter has an increased alpha-amylase yield, which on average is 13% higher than the control strain, AN83, which only comprises a wild-type copy of the *mrgA* gene.

Table 1. Yields of amylase from AN53 were determined in triplicate from three independent isolates, and compared to the yield of amylase from the control strain AN83. The average yields of each strain are also shown.

Strain	Amylase activity (relative)	average	average (in %)
AN53-1.1	15,7	13,5	113%
AN53-1.2	15,8		
AN53-1.3	14,9		
AN53-2.1	12,5		
AN53-2.2	8,65		
AN53-2.3	Nd		
AN53-3.1	Nd		
AN53-3.2	Nd		
AN53-3.3	Nd		
AN83-1.1	10,6	11,9	100%
AN83-1.2	15,7		
AN83-1.3	13,8		

Strain	Amylase activity (relative)	average	average (in %)
AN83-2.1	8,7		
AN83-2.2	7,72		
AN83-2.3	10,1		
AN83-3.1	14,1		
AN83-3.2	Nd		
AN83-3.3	14,6		
Nd: Not determined.			

EXAMPLE 2

[0099] The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers p740mrgaF2 (SEQ ID NO: 8) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p740mrgaF2) comprised the constitutive synthetic promoter P740 (SEQ ID NO: 9). The PCR product (SEQ ID NO: 10) was cut with restriction enzymes *SfiI* and *BamHI*, and ligated to the large *SfiI*-*BamHI* fragment of pDG268neo, resulting in plasmid p740mrgA. The ligation-mixture of p740mrgA, described in the Material and Methods section, was introduced by transformation into the *B. subtilis* 168 strain, and the transformants were plated on LBPSK media supplemented with 6 microg/ml chloramphenicol (cam) to select for integrants. Transformants growing on these plates have integrated the plasmid in the *amy* locus, either by a single (Cam^+ kan^+) or double cross-over event (cam^+ kan^-). The transformants were re-streaked on LBPSK/cam media with and without 20 microg/ml kanamycine. Strains where double cross-over events had occurred were cam^+ kan^- . These strains no longer showed the tell-tale clearing zones; which signified that that integration in, and disruption of, the *amy* gene, had taken place. The site of integration was verified by PCR, the integrated copy of *mrgA* was verified by sequence analysis, and the strain was named AN42. AN42 was transformed with plasmid pKTH10 which constitutively express the alpha-amylase AmyQ of *Bacillus amyloliquefaciens*. The resulting strain was named AN55. Yields of amylase from AN55 were determined in triplicate from three independent isolates, and compared to yield of amylase from the control strain AN83. Results are shown in table 2; the AN55 strain constitutively expressing *mrgA*

from the synthetic promoter has an increased alpha-amylase yield, which on average is 21% higher than the control strain, AN83, which only comprises a wild-type copy of the *mrgA* gene.

Table 2. Yields of amylase from AN55 were determined in triplicate from three independent isolates, and compared to the yield of amylase from the control strain AN83. The average yields of each strain are also shown.

Strain	Amylase activity (relative)	average	average (in %)
AN55-1.1	11,5	14,4	121%
AN55-1.2	10,8		
AN55-1.3	13,3		
AN55-2.1	17,4		
AN55-2.2	17,3		
AN55-2.3	15,6		
AN55-3.1	11,2		
AN55-3.2	14,8		
AN55-3.3	18,1		
AN83-1.1	10,6	11,9	100%
AN83-1.2	15,7		
AN83-1.3	13,8		
AN83-2.1	8,7		
AN83-2.2	7,72		
AN83-2.3	10,1		
AN83-3.1	14,1		
AN83-3.2	Nd		
AN83-3.3	14,6		
Nd: Not determined.			

EXAMPLE 3

[0100] The *mrgA* gene was amplified by PCR from the chromosome of *B. subtilis* by use of primers p726mrgaF2 (SEQ ID NO: 11) and MBmrgaR2 (SEQ ID NO: 5). The upstream primer (p726mrgaF2) comprised the constitutive synthetic promoter P726 (SEQ ID NO: 12). The PCR product (SEQ ID NO: 13) was cut with restriction enzymes *Sfi*I and *Bam*HI and ligated to the large *Sfi*I-*Bam*HI fragment of pDG268neo, resulting in plasmid p726mrgA. The ligation-mixture of p726mrgA, described in the Material and Methods section, was introduced by transformation into the *B. subtilis* 168 strain, and the transformants were plated on LBPSK media supplemented with 6 microg/ml chloramphenicol to select for integrants. Transformants growing on these plates have integrated the plasmid in the *amy* locus, either by a single ($\text{Cam}^+ \text{kan}^+$) or double cross-over event ($\text{cam}^+ \text{kan}^-$). The transformants were then re-streaked on LBPSK/cam media with and without 20 microg/ml kanamycine. Strains where double cross-over events had occurred were $\text{cam}^+ \text{kan}^-$. These strains no longer showed the tell-tale clearing zones; which signified that that integration in, and disruption of, the *amy* gene, had taken place. The site of integration was verified by PCR, the integrated copy of *mrgA* was verified by sequence analysis, and the strain was named AN50. AN50 was then transformed with plasmid pKTH10, which constitutively express the alpha-amylase AmyQ of *Bacillus amyloliquefaciens*. The resulting strain was named AN57. Yields of amylase from AN57 were determined in triplicate from three independent isolates, and compared to the yield of amylase from the control strain AN83. Results are shown in table 3; the AN55 strain constitutively expressing *mrgA* from the synthetic promoter has an increased alpha-amylase yield, which on average is 40% higher than the control strain, AN83, which only comprises a wild-type copy of the *mrgA* gene.

Table 3. Yields of amylase from AN57 were determined in triplicate from three independent isolates, and compared to the yield of amylase from the control strain AN83. The average yields of each strain are also shown.

Strain	Amylase activity (relative)	average	average (in %)
AN57-1.1	15,5	16,6	140%
AN57-1.2	10,6		
AN57-1.3	17,3		
AN57-2.1	17,4		

Strain	Amylase activity (relative)	average	average (in %)
AN57-2.2	20,7		
AN57-2.3	13,9		
AN57-3.1	15,1		
AN57-3.2	17		
AN57-3.3	22		
AN83-1.1	10,6	11,9	100%
AN83-1.2	15,7		
AN83-1.3	13,8		
AN83-2.1	8,7		
AN83-2.2	7,72		
AN83-2.3	10,1		
AN83-3.1	14,1		
AN83-3.2	Nd		
AN83-3.3	14,6		
Nd: Not determined.			

EXAMPLE 4

[0101] pAN213ban, described in the Material and Methods section, was introduced by transformation into the *B. subtilis* 168 strain and plated on LBPGS media supplemented with 5 microgram/ml erythromycin to select for integrants.

Transformants on these plates have integrated the plasmid in the pel locus, either by a single ($erm^+ kan^+$) or double cross-over event ($erm^+ kan^-$). Transformants were re-streaked on LBPGS/erm media with and without 20 microgram/ml kanamycine.

Strains where double cross-over events occurred were scored as $erm^+ kan^-$. These strains showed larger than wildtype clearing zones as an indication of $P_{consBAN}$ -amyQ integration and expression. The site of integration was verified by PCR. The resulting strain was named AN214. AN50 was transformed with chromosomal DNA from AN214 and transformants with the genotype ($pel::P_{consBAN}$, erm ; $amyE::P726mrgA$,

cam) were scored on plates. The resulting strain was verified by PCR and named AN217. Yield of amylase from AN217 was determined in duplicate from four independent isolates and compared to yield of amylase from the control strain AN214 (table 4).

Table 4. Yield of amylase from AN217 determined in duplicates from four independent isolates and compared to yield of amylase from the control strain AN214.

Strain	Amylase activity (relative)	average	
AN214-1.1	10,5	14,0	100%
AN214-1.2	10,9		
AN214-2.1	14,5		
AN214-2.2	13,6		
AN214-3.1	14,6		
AN214-3.2	16,1		
AN214-4.1	14,1		
AN214-4.2	17,9		
AN217-1.1	17,1	16,7	119%
AN217-1.2	16,5		
AN217-2.1	14,9		
AN217-2.2	14,6		
AN217-3.1	19,1		
AN217-3.2	18,1		
AN217-4.1	16,2		
AN217-4.2	17,2		

EXAMPLE 5

[0102] AN36 was transformed with chromosomal DNA from AN214 and transformants with the genotype (pel::P_{cons}BAN, erm; amyE::P920mrgA, cam) were scored on plates. The resulting strain was verified by PCR and named AN219. AN42

were transformed with chromosomal DNA from AN214 and transformants with the genotype (*pel::P_{cons}BAN*, *erm*; *amyE::P740mrgA*, *cam*) were scored on plates. The resulting strain was verified by PCR and named AN218. Yield of amylase from AN214, AN218 and AN219 were determined in duplicate from four independent isolates of each strain (table 5).

Table 5. Relative average yields of amylase from AN214, AN218 and AN219, determined in duplicate from four independent isolates of each strain.

Strain	Amylase activity
AN214	100
AN218	107
AN219	106

SEQUENCE LISTING

[0103]

<110> Novozymes A/S

<120> A cell with improved secretion mediated by MrgA protein or homologue

<130> 10527.204-WO

<160> 16

<170> PatentIn aversion 3.2

<210> 1

< 211> 815

< 212> DNA

< 213> *Bacillus subtilis* 168

<220>

< 221> misc_feature

< 222> (201)..(659)

< 223> MrgA encoding sequence

<400> 1

```

aagaatttgc gataccgat cggaagggc atcaagctca cctgctggt ccgatcgctt 60
tttccttggg ctgctggga gtctatctg aagaaaaagc tattcagctg atctaaatta 120
taattattat aatttagtat tgatttttat ttagtatatg atataattaa gtcaacagat 180
cacaaggagg acgttatctt atgaaaactg aaaacgcaa aacaaatcaa acattagttg 240
agaattcact gaacacacaa ttatcaaact ggtttctttt atactotaag ctccaccgtt 300
tccattggta tgtgaaaggg cctcatttct ttacattgca cgagaaattt gaagaacttt 360
atgaccatgc ggctgaaaca gtggatacca tcgctgagcg cctgctggcg attggcggac 420
agcctgttgc cacagtgaag gaatacactg agcatgcac tatcacagac gcgggaaacg 480
aaacatcagc atcagaaatg gtacaagcat tggtaaacga ctacaaacaa atcagcagcg 540
aatctaaatt cgtgatcggc ctggctgaag aaaatcaaga caatgcgaca gcggacttgt 600
ttgtcggatt aattgaagaa gttgaaaaac aagtgtggat gctttctctt tatttagggg 660
aacaaaaaag ctgaacctta atcgggttca gctttttgtt tttcttttagc ttgaactgct 720
ttctgtctgc ttggtcagtg ttgcgttcaa cgttttcggt tttcccttcc gcagcacttg 780
gattgttggt ttatctccga cttttaagtc tttgt 815

```

<210> 2

< 211> 153

< 212> PRT

< 213> Bacillus subtilis 168

<220>

< 221> PEPTIDE

< 222> (1)..(153)

< 223> MrgA protein

<400> 2

Met Lys Thr Glu Asn Ala Lys Thr Asn Gln Thr Leu Val Glu Asn Ser
1 5 10 15

Leu Asn Thr Gln Leu Ser Asn Trp Phe Leu Leu Tyr Ser Lys Leu His
20 25 30

Arg Phe His Trp Tyr Val Lys Gly Pro His Phe Phe Thr Leu His Glu
35 40 45

Lys Phe Glu Glu Leu Tyr Asp His Ala Ala Glu Thr Val Asp Thr Ile
50 55 60

Ala Glu Arg Leu Leu Ala Ile Gly Gly Gln Pro Val Ala Thr Val Lys
65 70 75 80

Glu Tyr Thr Glu His Ala Ser Ile Thr Asp Gly Gly Asn Glu Thr Ser
85 90 95

Ala Ser Glu Met Val Gln Ala Leu Val Asn Asp Tyr Lys Gln Ile Ser
100 105 110

Ser Glu Ser Lys Phe Val Ile Gly Leu Ala Glu Glu Asn Gln Asp Asn
115 120 125

Ala Thr Ala Asp Leu Phe Val Gly Leu Ile Glu Glu Val Glu Lys Gln
130 135 140

Val Trp Met Leu Ser Ser Tyr Leu Gly
145 150

<210> 3

< 211> 8644

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Plasmid pDG268neo

<400> 3

aacaaaattc tccagtcttc acatcggttt gaaaggagga agcgaagaa tgaagtaaga 60
gggatttttg actccgaagt aagtcttcaa aaaatcaaat aaggagtgtc aagaatgttt 120
gcaaaacgat tcaaaacctc ttactgccc ttattcgctg gatttttatt gctgtttcat 180

ttggttctgg caggaccggc ggctgcgagt gctgaaacgg cgaacaaac gaatgagctt	240
acagcacctg ccatcaaaag cggaaccatt ctccatgcac ggaattgggc gttcaatacg	300
ttaaaacaca atatgaagga tattcatgat gcaggatata cagccattca gacatctccg	360
attaaccaag taaaggaagg gaatcaagga gataaaagca tgcgaaactg gtactggctg	420
tatcagccga catcgatca aattggcaac cgttacttag gtactgaaca agaattfaaa	480
gaaatgtgtg cagccgctga agaatatggc ataaaggcca ttgttgacgc gcggccggcg	540
atccatacac aaaaaaacgc tgtgcccttt aaccgcacag cgttttttta ttgattaacg	600
cgttgccgct tctgcgttaa caagtccgct tccatacaag ttctgtcttc ctaaactagt	660
tgcggtattc ttagatgat ttogaatttg tacattagac caagatgggt tottttgttt	720
aacaagggcg gcgcacctg caacatgagg agtagccatc gatgtaccgt ttaagctggc	780
atatgttgaa cctgggtatg tgctctgcac gtttaccccg ggtgcgacaa tgtcaaggcc	840
tgcgccatac tgtgaaaagc tagcgcggtt gttgttttga tcagttagctc cgactgccat	900
tgcgttcgca tagcgcgcgc gatagctgat tgagcctgca cctgaattcc cagatgcgc	960
tacaacaaga acgcctctag aagtcgcgct attaacagct tgctcgagtg tggcacttgg	1020
cgaagggctt cctaaactca aattagcaac gtgcctgcca ttgttccctg cccattccaa	1080
tccttgggca atcgagctga ccgaacctga accgctcgcc cctaggactt taacagcgta	1140
tagctcagcg ctccggcgta cgcgaagaac gccaatcgaa ttgtttaaag cagcgatcgt	1200
cccgccaca tgcgtgccat gccattccc atcttgagtc gacggtccc ctggtacaaa	1260
gcttgcgcca ccacgaatat ttagatctgg atgagtggat atccctgtat cgaggacagc	1320
aacttttaca ccagaacctg tcaatccacg gttatgggca gctggggctt gcacacggct	1380
aattcccat ggtaccgatt gcgccattgt cgttacttct gcctcctctt caatataaga	1440
aatcgctgga tcgagttcaa gcgcgtccac atcttctggg cttaactcaa cggataaaac	1500
aggaatcgtt tcaaatcat gaagcaattc aatttcgact tctcttctc cagagagaat	1560
ggcgacctcg tcatttgctt ctacttgctt taaaaactca ctgacagctt cctgctcatt	1620
aaagccaatt aaatatTTTT cttttgcttc ttcagcagcc gatgcgatcg atgaactaaa	1680
agcaacagaa atgagtagtg cgggtgcttg gacaattttc cccaacgggt tottcattcg	1740
gtttccctcc tcatttttat agagctccat aatacataat tttcaaacctg ataaaatgat	1800
ttttcataaa tccattagac ggtgcaaata tatgttttta atgttcttcg tttttaggca	1860
tcctccttt caagataaat aatttataca ctattctatt ggaatcttaa tcattccaat	1920
agaaaaatat gtaatgatta taataagtc gcttcttata ataaatatat ttacatatcc	1980
atttaatact acatcatgtt aggtatagta aggctatcaa ggggtgctta atttctactt	2040

gtaacaatgt attggcatat tatatatgtga attgagaaaa ttaaatacag cgataattca	2100
catgaacaag ttcatttgga gttatatattt caaatattca aggttggtgct tgtatgtcat	2160
tctatagtta gataagcatt tgaggtagag tccgtccgaa tatatttgta atctgaagaa	2220
ggttcaaaca tattttctata taacgtattc tttttttgta gttcttactt ttgaggggog	2280
ttacaattca aagatattat ctttaattaa gottaacatt aataattctt caattgcaac	2340
aaaaaaagca cttttatcta aggtttcatc ttacgtttcg agggccccctc cattttctta	2400
tacaaattat attatacata tcagtaaaat aatgtcaacc cccctttatt cttttttttt	2460
acacagcgga cagtctggac agcaggccct taaggccaat tctcatgttt gacagcttat	2520
catcggaat agttaccctt attatcaaga taagaaagaa aaggattttt cgctacgctc	2580
aatccctta aaaaaacaca aaagaccaca ttttttaatg tggcttttat tcttcaacta	2640
aagcaccat tagttcaaca aacgaaaatt ggataaagtg ggatattttt aaaatatata	2700
tttatgttac agtaatatg acttttaaaa aaggattgat tctaataag aaagcagaca	2760
agtaagcctc ctaaattcac tttagataaa aatttaggag gcatatcaaa tgaactttaa	2820
taaaattgat ttagacaatt ggaagagaaa agagatattt aatcattatt tgaaccaaca	2880
aacgactttt agtataacca cagaaattga tattagtgtt ttataccgaa acataaaaca	2940
agaaggatat aaattttacc ctgcatttat tttcttagtg acaaggtga taaactcaaa	3000
tacagctttt agaactgggt acaatagcga cggagagtta ggttattggg ataagttaga	3060
gccactttat acaatttttg atggtgtatc taaaacattc tctggtattt ggactcctgt	3120
aaagaatgac ttcaaagagt tttatgattt atacctttct gatgtagaga aatataatgg	3180
ttcggggaaa ttgtttccca aaacacctat aacctgaaat gctttttctc tttctattat	3240
tccatggact tcatttactg ggtttaactt aaatatcaat aataatagta attaccttct	3300
acctattatt acagcaggaa aattcattaa taaaggtaat tcaatatatt taccgctatc	3360
tttacaggta catcattctg tttgtgatgg ttatcatgca ggattgttta tgaactctat	3420
tcaggaattg tcagataggc ctaatgactg gcttttataa tatgagataa tgccgactgt	3480
actttttaca gtccgttttc taatgtcact aacctgcccc gttagttgaa gaagggtttt	3540
atattacagc tccagatcct ctacgcgga cgcacgtgg ccggcatcac cggcgccaca	3600
ggtgcggttg ctggcgcta tatcgccgac atcaccgatg gggaagatcg ggctcgccac	3660
ttcgggctca tgagcgcttg tttcggcgtg ggtatggtgg caggccccgt ggccggggga	3720
ctgttgggog ccatctcctt gcatgcacca ttccttgagg cggcggtgct caacggcctc	3780
aacctactac tgggctgctt cctaatagcag gagtgcgata agggagagcg tcgacatgga	3840

tgagcgatga	tgatatccgt	ttaggctggg	cggtgatagc	ttctcgttca	ggcagtacgc	3900
ctcttttctt	ttccagacct	gaggaggcg	gaaatgggtg	gaggttcccg	gggaaaagcc	3960
aaataggcga	tcgcgggagt	gctttatttg	aagatcaggc	tatcactgcg	gtcaatagat	4020
ttcacaatgt	gatggctgga	cagcctgagg	aactctcgaa	cccgaatgga	aacaaccaga	4080
tatttatgaa	tcagcgcggc	tcacatggcg	ttgtgctggc	aatgcaggt	tcacctctcg	4140
ttcttatcaa	tacggcaaca	aaattgcttg	atggcaggta	tgacaataaa	gctggagcgg	4200
gttcatttca	agtgaacgat	ggtaaactga	caggcacgat	caatgccagg	tctgtagctg	4260
tgctttatcc	tgatgatatt	gcaaaaagcg	ctcatgtttt	ccttgagaat	tacaaaaag	4320
gtgtaacaca	ttctttcaat	gatcaactga	cgattacctt	gcgtgcagat	gcgaatacaa	4380
caaaaagcgt	ttatcaaadc	aataatggac	cagacgacag	gcgtttaagg	atggagatca	4440
attcacaatc	ggaaaaggag	atccaatttg	gcaaaacata	caccatcatg	ttaaaaggaa	4500
cgaacagtga	tggtgtaacg	aggaccgaga	aatacagttt	tggttaaaaga	gatccagcgt	4560
cggccaaaaac	catcggttat	caaaatccga	atcattggag	ccaggtaaatt	gcttatatct	4620
ataaacatga	tgggagccga	gtaattgaat	tgaccggatc	ttggcctgga	aaaccaatga	4680
ctaaaaatgc	agacggaatt	tacacgctga	cgctgcctgc	ggacacggat	acaaccaacg	4740
caaaagtgat	ttttaataat	ggcagcgccc	aagtgcocgg	tcagaatcag	cctggctttg	4800
attacgtgct	aatgggttta	tataatgact	cggtcttaag	cggttctctt	ccccattgag	4860
ggcaaggcta	gacgggactt	accgaaagaa	accatcaatg	atggtttctt	ttttgttcat	4920
aaatcagaca	aaacttttct	cttgcaaaag	tttgtgaagt	gttgcacaaat	ataaatgtga	4980
aatacttcac	aaacaaaaag	acatcaaaga	gaaacatacc	ctgcaaggat	gctgatattg	5040
tctgcatttg	cgccggagca	aaccaaaaaac	ctggtgagac	acgccttgaa	ttagtagaaa	5100
agaacttgaa	gattttcaaa	ggcatcgtta	gtgaagtcac	ggcgagcggg	tttgacggca	5160
ttttcttagt	cggtaacaat	cctcgttaaa	ggacaaggac	ctgagcggaa	gtgtatcgta	5220
cagtagacgg	agtatactag	tatagtctat	agtcctgga	attattatat	ttatctccga	5280
cgatattctc	atcagtgaag	tcagctgga	gttcttttagc	aaattttttt	attagctgaa	5340
cttagtatta	gtggggccgc	tgataattac	taatactagg	agaagttaat	aaatacgtaa	5400
ccaacatgat	taacaattat	tagaggtcac	cgttcaaaat	ggtatgcgtt	ttgacacatc	5460
cactatatat	ccgtgtcgtt	ctgtccactc	ctgaatccca	ttccagaaat	totctagcga	5520
ttccagaagt	ttctcagagt	cggaaagttg	accagacatt	acgaactggc	acagatggtc	5580
ataaactgaa	ggaagatctg	attgcttaac	tgcttcagtt	aagaccgaag	cgctcgtcgt	5640
ataacagatg	cgatgatgca	gaccaatcaa	catggcacct	gccattgcta	cctgtacagt	5700

caaggatggt agaaatgttg tcggtccttg cacacgaata ttacgccatt tgcctgcata	5760
ttcaaacagc tcttctacga taagggcaca aatcgcatcg tggaacgttt gggcttctac	5820
cgatttagca gtttgataca ctttctctaa gtatccacct gaatcataaa tcggcaaaat	5880
agagaaaaat tgaccatgtg taagcggccca atctgattcc acctgagatg cataatctag	5940
tagaatctct tcgctatcaa aattcacttc caccttcac tcaccgggtg tccattcatg	6000
gctgaactct gcttctctg ttgacatgac acacatcatc tcaatatccg aatagggcc	6060
atcagtctga cgaccaagag agccataaac accaatagcc ttaacatcat ccccatattt	6120
atccaatatt cgttctctaa tttcatgaac aatcttcatt ctttcttctc tagtcattat	6180
tattgggtcca ttcactattc tcattccctt ttcagataat tttagatttg cttttctaaa	6240
taagaatatt tggagagcac cgttcttatt cagctattaa taactcgtct tctaagcat	6300
catggtctca cttttccact ttttgtcttg tccactaaaa ccttgattt ttcattgaa	6360
taaatgctac tattaggaca cataatatta aaagaaacc ccatctattt agttatttgt	6420
ttagtcaact ataactttaa cagatggggt ttttctgtgc aacbaatttt aagggttttc	6480
aatactttaa aacacataca taccaacact tcaacgcacc ttcagcaac taaaataaaa	6540
atgacgttat ttctatatgt atcaagataa gaaagaacaa gttcaaaacc atcaaaaaaa	6600
gacacctttt caggtgcttt ttttatttta taaactcatt cctgatctc cccatactcc	6660
tccaatccaa agctatttag aaagattact atatcctcaa acaggcggtg accggcctct	6720
tcacgggaa tgcgcgcgac cttcagcatc gccggcatgt cccctggcg gacgggaagt	6780
atccagctcg aggtcgggcc gcgttgctgg cgtttttcca taggtccgc cccctgacg	6840
agcatcacia aaatcgacgc tcaagtcaga ggtggcgaaa ccgacagga ctataaagat	6900
accaggcgtt tccccctgga agctccctcg tgcgctctcc tgttccgacc ctgccgctta	6960
cggatacct gtccgccttt ctcccttcgg gaagcgtggc gctttctcat agctcacgct	7020
gtaggatatct cagttcggtg taggtcgttc gctccaagct gggctgtgtg caccgaaccc	7080
cgttcagcc cgaccgctgc gccttatccg gtaactatcg tcttgagtcc aacccggtaa	7140
gacacgaact atcgccactg gcagcagcca ctggtaacag gattagcaga gcgaggtatg	7200
taggcggtgc tacagagttc ttgaagtggg ggcctaacta cggtacact agaaggacag	7260
tatttggtat ctgcgctctg ctgaagccag ttaccttcgg aaaaagagtt ggtagctctt	7320
gatccggcaa acaaaccacc gctggtagcg gtgggttttt tgtttgcaag cagcagatta	7380
cgcgcagaaa aaaaggatct caagaagatc ctttgatctt ttctacggg tctgacgctc	7440
agtggaaaga aaactcacgt taagggattt tggcatgag attatcaaaa aggatcttca	7500

```

cctagatcct tttaaattaa aaatgaagtt ttaaataaat cttaaagtata tatgagtaaa 7560
cttgggtctga cagttaccpa tgcttaataca gtgaggcacc tatctcagcg atctgtctat 7620
ttcgttcate catagttgcc tgactccccg tcgtgtagat aactacgata cgggagggct 7680
taccatctgg cccagtgct gcaatgatac cgcgagaccc acgtcaccg gctccagatt 7740
tatcagcaat aaaccagcca gccggaagg cagagcgag aagtggtcct gcaactttat 7800
ccgcctccat ccagttctatt aattgttgcc gggagctag agtaagtagt tcgccagtta 7860
atagtttgcg caacgttggt gccattgctg caggcatcgt ggtgtcacgc tcgtcgtttg 7920
gtatggcttc attcagctcc ggttcccaac gatcaaggcg agttacatga tccccatgt 7980
tgtgcaaaaa agcgggttagc tccttcggtc ctccgatcgt tgtcagaagt aagttggccg 8040
cagtgttata actcatgggt atggcagpac tgcataattc tcttactgtc atgccatccg 8100
taagatgctt ttctgtgact ggtgagtact caaccaagtc attctgagaa tagtgtatgc 8160
ggcgaccgag ttgctcttgc ccggcgtaaa cccgggataa taccgcgcca catagcagaa 8220
ctttaaagt gctcatcatt ggaaaacgtt cttcggggcg aaaactctca aggatcttac 8280
cgctgttgag atccagttcg atgtaacca ctcggtgcacc caactgatct tcagcatctt 8340
ttactttcac cagcgtttct gggtgagcaa aaacaggaag gcaaaatgcc gcaaaaaagg 8400
gaataagggc gacacggaaa tgttgaatac tcatactctt cttttttcaa tattattgaa 8460
gcatttatca gggttattgt ctcatgagcg gatacatatt tgaatgtatt tagaaaaata 8520
aacaatatag ggttcgcgc acatttcccc gaaaagtgcc acctgacgtc taagaaacca 8580
ttattatcat gacattaacc tataaaaaata ggcgtatcac gaggeccctt cgtcttcaag 8640
aatt 8644

```

<210> 4

< 211> 91

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer p920mrgaF2

<400> 4

```

ctgaggccaa ttaggccaag ttattcttg acattaggga acatgcatga tataataggt 60
aaagtaaaca gatcacaagg aggacgttat c 91

```

<210> 5

< 211> 34

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer MBmrgaR2

<400> 5

tgaaggatcc acgcgtccag cagacagaaa gcag 34

<210> 6

< 211> 49

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Promoter P920

<400> 6

aagtttattc ttgacattag ggaacatgca tgatataata ggtaaagta 49

<210> 7

< 211> 642

< 212> DNA

< 213> Artificial sequence

<220>

< 223> PCR product of promoter 920 and mrgA

<400> 7

ctgaggcctt aaggccaag tttattcttg acattaggga acatgcatga tataataggt	60
aaagtaaaca gatcacaagg aggacgttat cttatgaaaa ctgaaaacgc aaaaacaaat	120
caaacattag ttgagaattc actgaacaca caattatcaa actggtttct tttatactct	180
aagctccacc gtttccattg gtatgtgaaa ggcctcatt tctttacatt gcacgagaaa	240
tttgaagaac tttatgacca tgcgctgaa acagtggata ccacgctga ggcctgctg	300
gcgattggcg gacagcctgt tgccacagtg aaagaatata ctgagcatgc atctatcaca	360
gacggcggaa acgaaacatc agcatcagaa atggtacaag cattggtaaa cgactacaaa	420
caaatcagca gcgaatctaa attcgtgacg ggccctggctg aagaaaatca agacaatgcg	480
acagcggact tgtttgtcgg attaattgaa gaagttgaaa aacaagtgtg gatgctttcc	540
tcttatttag ggtaacaaaa aagctgaacc ttaatcgggt tcagcttttt gttttttctt	600
agcttgaact gctttctgtc tgcttgacgc gtggatcctt ca	642

<210> 8

< 211> 91

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer p740mrgaF2

<400> 8

ctgaggccaa ttaggcccg aagtttgttg acacagctcc aggatacaaa tataatgggt 60

cgactaaaca gatcacaagg aggacgttat c 91

<210> 9 ,

< 211> 49

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Promoter P740

<400> 9

cggaagtttg ttgacacagc tccaggatac aaatataatg ggtcgagta 49

<210> 10

< 211> 642

< 212> DNA

< 213> Artificial sequence

<220>

< 223> PCR product of promoter 740 and mrgA

<400> 10

ctgaggcctt aagggcccg aagtttgttg acacagctcc aggatacaaa tataatgggt 60

cgagtaaaca gatcacaagg aggacgttat cttatgaaaa ctgaaaacgc aaaaacaaat 120

caaacattag ttgagaattc actgaacaca caattatcaa actggtttct tttatactct 180

aagctccacc gtttccattg gtatgtgaaa gggcctcatt tctttacatt gcacgagaaa 240

tttgaagaac tttatgacca tgcggctgaa acagtggata ccacgctga gcgctgctg 300

gcgattggcg gacagcctgt tgccacagtg aaagaatata ctgagcatgc atctatcaca 360

gacggcggaa acgaaacatc agcatcagaa atggtacaag cattggtaaa cgactacaaa 420

caaatacagca gcgaatctaa attcgtgatc ggcttggtg aagaaaatca agacaatgcg 480

acagcggact tgtttgtcgg attaattgaa gaagttgaaa aacaagtgtg gatgctttcc 540

tcttatttag ggtaacaaaa aagctgaacc ttaatcgggt tcagcttttt gttttttctt 600

agcttgaact gctttctgtc tgcttgacgc gtggatcctt ca 642

<210> 11

< 211> 91

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer p726mrgaF2

<400> 11

ctgaggccaa ttaggccgag gtgagatttg acactagtag gctacgggac tataatgcgg 60

gaagtaaaca gatacaagg aggacgttat c 91

<210> 12

< 211> 49

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Promoter P726

<400> 12

gaggtgagat ttgacactag taggctacgg gactataatg cggaagta 49

<210> 13

< 211> 642

< 212> DNA

< 213> Artificial sequence

<220>

< 223> PCR product of promoter P726 and mrgA

<400> 13

ctgaggcctt aaggccgag gtgagatttg acactagtag gctacgggac tataatgcgg 60

gaagtaaaca gatacaagg aggacgttat cttatgaaa ctgaaaacgc aaaaacaaat 120

caaacattag ttgagaattc actgaacaca caattatcaa acttggtttct ttataactct 180

aagctccacc gtttccattg gtatgtgaaa gggcctcatt tctttacatt gcacgagaaa 240

tttgaagaac tttatgacca tgcggtgaa acagtgata ccacgctga gcgctgctg 300

gcgattggcg gacagcctgt tgccacagt aaagaatata ctgagcatgc atctatcaca 360

gacggcgga acgaaacac agcatcagaa atggtacaag cattggtaaa cgactacaaa 420

caaatcagca gcaatctaa attcgtgatc ggcttgctg aagaaaatca agacaatgcg 480

acagcggact tgtttgtcgg attaattgaa gaagttgaaa aacaagtgtg gatgctttcc 540

tcttatttag ggtaacaaaa aagctgaacc ttaatcgggt tcagcttttt gttttttctt 600

agcttgaact gctttctgtc tgcttgacgc gtggatcctt ca 642

<210> 14

< 211> 88

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer AN162

<400> 14

```
agactgtccg cgggtgtaaaa aataggaata aagggggggtt gacattattt tactgatatg    60
tataatataa tttgtataag aaaatgag                                     88
```

<210> 15

< 211> 32

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Primer AN163c

<400> 15

```
gcatacacgc gttgtcacac ctgatgccga cc    32
```

<210> 16

< 211> 8152

< 212> DNA

< 213> Artificial sequence

<220>

< 223> Plasmid pAN213ban

<400> 16

ccgcgggtgta	aaaaatagga	ataaaggggg	gttgac'atta	ttttactgat	atgtataata	60
taattttgtat	aagaaaatga	gagggagagg	aaacatgatt	caaaaaacgaa	agcggacagt	120
ttcgtttcaga	cttgtgctta	tgtgcacgct	gttattttgtc	agttttgccga	ttacaaaaaac	180
atcacgcgta	aatggcacgc	tgatgcagta	ttttgaatgg	tatacgcgga	acgacggcca	240
gcattggaaa	cgattgcaga	atgatgcgga	acattttatcg	gatatcggaa	tcactgccgt	300
ctggattcct	ccgcataca	aaggattgag	ccaatccgat	aacggatacg	gaccttatga	360
tttgtatgat	ttaggagaat	tccagcaaaa	agggacggtc	agaacgaaat	acggcacaaa	420
atcacagcct	caagatgcga	tcggctcact	gcattcccgg	aacgtccaag	tatacggaga	480
tgtgtttttg	aatcataagg	ctggtgctga	tgcaacagaa	gatgtaactg	ccgtcgaaat	540
caatccggcc	aatagaaatc	aggaaacttc	ggaggaatat	caaatacaag	cgtggacgga	600
ttttcgtttt	ccgggccgtg	gaaacacgta	cagtgttttt	aaatggcatt	ggtatcattt	660
cgacgcgagcg	gactgggatg	aatcccggaa	gatcagccgc	atctttaagt	ttcgtgggga	720
aggaaaagcg	tgggattggg	aagtatcaag	tgaaaaacggc	aactatgact	attttaatgta	780
tgctgatgct	gactacgacc	accctgatgt	cgtggcagag	acaaaaaaat	ggggtatctg	840
gtatgcgaat	gaactgtcat	tagacggctt	ccgtattgat	gccgccaaac	atattaaatt	900
ttcattttctg	cgtgattggg	ttcaggcggg	cagacaggcg	acgggaaaag	aaatgtttac	960
ggttgccggag	tattggcaga	ataatgccgg	gaaactcgaa	aactacttga	ataaaaacaag	1020
ctttaatcaa	tccgtgtttg	atgttccgct	tcattttcaat	ttacaggcgg	cttcctcaca	1080
aggaggcgga	tatgatatag	ggcgtttgct	ggacgggtacc	gttgtgtcca	ggcatccgga	1140
aaaggcggtt	acattttgtt	aaaatcatga	cacacagccg	ggacagtcac	tggaaatcgac	1200
agtccaaact	tggtttaaac	cgcttgcata	cgccctttatt	ttgacaagag	aatccgggta	1260
tcctcagggtg	ttctatgggg	atatgtacgg	gacaaaaggg	acatcgccaa	aggaaattcc	1320
ctcactgaaa	gataatatag	agccgatttt	aaaagcgcgt	aaggagtacg	catacggggc	1380
ccagcacgat	tatatggacc	acccggatgt	gatcggatgg	acgagggaag	gtgacagctc	1440
cgccgccaaa	tcaggtttgg	ccgctttaat	cacggacgga	cccgccggat	caaagcggat	1500
gtatgccggc	ctgaaaaatg	ccggcgagac	atggtatgac	ataacgggca	accgttcaga	1560
tactgtaaaa	atcggatctg	acggctgggg	agagtttcat	gtaaacgatg	ggtccgtctc	1620
catttatgtt	cagaaataag	gtaataaaaa	aacacctcca	agctgagtgc	gggtatcagc	1680
ttggagggtgc	gtttattttt	tcagccgtat	gacaaggtcg	gcatcagggtg	tgacaacgcg	1740
tgatccagac	cagttccctg	agcttccgtc	agtcggatcc	cattgcggaa	aatagtcata	1800
ggcatccctgg	aattcaatgt	tgogaataat	gacgttatca	ctcttgattt	ggaagtttcc	1860
tcccacgact	ttagcgttag	tccttgaaac	gacgatcgtc	gtgtttgcag	ggatatccac	1920
catgaccgct	gctttttggg	ttttctgaga	gcgtgctctc	gcttcttctt	gtgttcccca	1980
cggctctttt	ttgcccatag	tgctaggatc	ataggctttc	aaatatttgt	ccaaatcata	2040
ctccggatct	ttatagtcac	ttaggccaag	cggcttcaga	ttgtcatcca	cgttcattgtc	2100
aatcgttccc	ttgatataaa	tgatttttgg	cgttgtgttc	gtttccttcc	ctaatagccga	2160
gacaagctgg	tttctgttgc	tgacgggtata	cacatttgag	gaggatgctt	ttgatccgcc	2220
tgtcgtgccg	gtcaggtacg	cgccccagcc	atcattggat	cccaacgtct	ggtggcctaa	2280
atcagctcg	ttcgcccgag	ctggagtcaa	tcctaaaaac	aaagccgtag	ctaacaatcaa	2340
aagggcctcg	tgatacgcc	attttttatag	gttaaatgtca	tgataataat	ggtttcttag	2400
acgtcagggtg	gcacttttgc	gggaaatgtg	cgcggaaccc	ctatttggtt	atttttctaa	2460
atacattcaa	atatgtatcc	gctcatgaga	caataaccct	gataaatgct	tcaataatat	2520
tgaaaaagga	agagtatgag	tattcaacat	ttccgtgtcg	cccttattcc	cttttttgcg	2580
gcatttttgc	ttcctgtttt	tgctcaccac	gaaacgcgtg	tgaaagtaaa	agatgctgaa	2640

gatcagtttg	gtgcacgagt	gggttacatc	gaactggatc	tcaacagcgg	taagatcctt	2700
gagagttttc	gccccgaaga	acgtttttcc	atgatgagca	cttttaaaagt	tctgctatgt	2760
ggcgcggtat	tatcccggtg	tgacgccggg	caagagcaac	tgggtcgccg	catacactat	2820
tctcagaatg	acttggttga	gtactcacca	gtcacagaaa	agcatcttac	ggatggcatg	2880
acagtaagag	aattatgcag	tgctgccata	accatgagtg	ataacactgc	ggccaactta	2940
cttcttgacaa	cgatcggagg	accgaaggag	ctaaccgctt	ttttgcacaa	catgggggat	3000
catgtaactc	gccttgatcg	ttgggaaccg	gagctgaatg	aagccatacc	aaacgacgag	3060
cgtgacacca	cgatgcctgc	agcaatggca	acaacgtttg	gcaaactatt	aactggcgaa	3120
ctacttactc	tagcttcccg	gcaacaatta	atagactgga	tggaggcgga	taaagttgca	3180
ggaccacttc	tgcgctcggc	ccttcggct	ggctggttta	ttgctgataa	atctggagcc	3240
ggtgagcgtg	ggtctcgcg	tatcattgca	gcactggggc	cagatggtaa	gccctcccg	3300
atcgtagtta	tctacacgac	ggggagtcag	gcaactatgg	atgaacgaaa	tagacagatc	3360
gctgagatag	gtgcctcact	gattaagcat	tggttaactgt	cagaccaagt	ttactcatat	3420
atacttttag	ttgattttaa	acttcatttt	taatttaaaa	ggatctaggt	gaagatcctt	3480
tttgataatc	tcatgaccaa	aatcccttaa	cgtgagtttt	cgttccactg	agcgtcagac	3540
cccgtagaaa	agatcaaaag	atcttcttga	gatccttttt	ttctgcgcgt	aatctgctgc	3600
ttgcaaacaa	aaaaaccacc	gctaccagcg	gtggttttgt	tgcgggatca	agagctacca	3660
actctttttc	tgcaggtaac	tggtctcagc	agagcgagca	taccaaatac	tgtccttcta	3720
gtgtagccgt	agttaggcca	ccacttcaag	aactctgtag	caccgcctac	atacctcgct	3780
ctgtaaatcc	tgttaccagt	ggctgctgoc	agtggcgata	agtcgtgtct	taccgggttg	3840
gactcaagac	gatagttacc	ggataaggcg	cagcggtcgg	gctgaacggg	gggttcgtgc	3900
acacagccca	cttgggagcg	aacgacctac	accgaactga	gatacctaca	gcgtgagcta	3960
tggagaaagc	ccacgcttcc	cgaagggaga	aaggcggaca	ggtatccggg	aagcggcagg	4020
gtcggaaacag	gagagcgcac	gagggagctt	ccagggggaa	acgcctggta	tctttatagt	4080
cctgtcgggt	ttcgccacct	ctgacttgag	cgtcgatttt	tgtgatgctc	gtcagggggg	4140
gggagcctat	ggaaaaacgc	cagcaacgcg	gcccgaacct	gagctggata	cttcccgtcc	4200
gccaggggga	catgccggcg	atgctgaagg	tcgcgcgcac	tcccgatgaa	gaggccggtt	4260
accgcctggt	tgaggatata	gtaatctttc	taaatagctt	tggattggag	gagtatgggg	4320
agatcaggga	atgagtttat	aaaaataaaa	aagcacctga	aaagggtgct	ttttttgatg	4380
gttttgaact	gttcttttct	tatcttgata	catatagaaa	taacgtcatt	tttattttag	4440
ttgctgaaag	gtgcgttgaa	gtgttggtat	gtatgtgttt	taaagtattg	aaaaccctta	4500
aaattgggtg	cacagaaaaa	ccccatctgt	taaagttata	agtgactaaa	caaataacta	4560
aatagatggg	ggtttctttt	aatattatgt	gtcctaatag	tagcatttat	tcagatgaaa	4620
aatcaagggt	tttagtgagc	aagacaaaaa	gtggaaaaagt	gagaccatga	tgcttaggaa	4680
gacgagttat	taatagctga	ataagaacgg	tgctctccaa	atattcttat	ttagaaaaagc	4740
aaatctaaaa	ttatctgaaa	agggaatgag	aatagtgaat	ggaccaataa	taatgactag	4800
agaagaaaaa	atgaagattg	ttcatgaaat	taaggaaacga	atattggata	aatattggga	4860
tgatgttaag	gctattgggt	tttatggctc	ctttggctgt	cagactgatg	ggccctattc	4920
ggatattag	atgatgtgtg	tcattgtcaac	agaggaagca	gagttcagcc	atgaatggac	4980
aaccggtgag	tggaaaggtg	aagtgaattt	tgatagcgaa	gagattctac	tagattatgc	5040
atctcagggt	gaatcagatt	ggccgcttac	acatggtcaa	tttttctcta	ttttgccgat	5100
ttatgattca	ggtggatact	tagagaaaagt	gtatcaaact	gctaaaatcg	tagaagccca	5160
aacgttccac	gatgcgattt	gtgcccttat	cgtagaagag	ctgtttgaat	atgcaggcaa	5220
atggcgtaat	attcgtgtgc	aaggaccgac	aacatttcta	ccatccttga	ctgtacaggt	5280
agcaatggca	ggtgccatgt	tgattgggtct	gcatactcgc	atctgttata	cgaacgagcg	5340
ttcggctctta	actgaagcag	ttaaagcaatc	agatcttctc	tcaggttatg	accatctgtg	5400
ccagttcgta	atgtctggtc	aactttccga	ctctgagaaa	cttctggaat	cgctagagaa	5460
tttctggaat	gggattcagg	agtggacaga	acgacacgga	tatatagtgg	atgtgtcaaa	5520
acgcatacca	ttttgaacga	tgacctctaa	taattgttaa	tcattgttgt	tacgtattta	5580
ttactttctc	ctagtattag	taattatcag	cggcccccact	aataactaagt	tcagctaata	5640
aaaaaatttg	ctaaagaact	cagctggat	ttcactgatg	agaatatcgt	cggagataaaa	5700
tataataatt	ccacggacta	tagactatac	tagtatactc	cgtctactgt	acgatacact	5760
tcogctcagg	tccttgctct	ttaacgagga	ttgttaccga	ctaagaaaat	gccgtcaaat	5820
ccgctcgcca	tgaactcag	tcgaaccgca	cccgttgat	ttataacatt	tgatttcaca	5880
ttagcagaag	catcaatcga	tcocatgcaga	gacggcgctcc	agccgacaga	agagctcagc	5940
ccgtttgcag	ccgatgcgtt	gatctgtgtg	ccgttcagca	acgtgccgga	gtcatataaa	6000
gccgttcccc	cgtgaatac	gctgatcggt	ttagcagctg	acagtcocgg	tacgtcaatg	6060
acattgtttt	gggcatagat	tttagatgac	tttccgattc	cccattgcata	gctaaaaagga	6120
taacttgaag	agcttgtgct	tccttcataa	tagttgttgt	atacgtgcac	ttgcccgaag	6180
cggactctcg	gcgcgcgctg	gacaatat	ttatagcggt	tatgatgcag	cgtaattttt	6240
aatttgcctg	catcggaggt	tttgcgtgca	cttgatccga	aaatggagct	tttatcatga	6300

tcgtgataat	agttgtagga	catcgtgata	tagttagcac	cgttggaagc	atccgtttgg	6360
ccgcatgggt	gctgataatt	tcttcataaa	tatttcggtg	atgtgctgtc	cggacgcgaa	6420
ccgtgtcggc	gatataggcg	ccagcaaccg	cacctgtggc	gccggtgatg	ccggccacga	6480
tgcgtccggc	gtagaggatc	tggagctgta	atataaaaac	cttcttcaac	taacggggca	6540
ggttagtga	attagaaaac	cgactgtaaa	aagtacagtc	ggcattatct	catattataa	6600
aagccagtca	ttaggcctat	ctgacaattc	ctgaatagag	ttcataaaca	atcctgcatg	6660
ataaccatca	caaacagaat	gatgtacctg	taaagatagc	ggtaaatata	ttgaattacc	6720
tttattaatg	aattttcctg	ctgtaataat	gggtagaagg	taattactat	tattattgat	6780
atttaagtta	aaccagtaa	atgaagtcca	tggttatgtc	tttgtatccc	gtttgtatta	6840
cttgatcctt	taactctggc	aacctcaaaa	attgaatgag	acatgctaca	cctccggata	6900
ataaatatat	ataaacgtat	atagatttca	taaagtctaa	cacactagac	ttatttactt	6960
cgtaattaag	tcgttaaac	gtgtgctcta	cgacccaaa	ctataaaac	ttagaacttt	7020
tcttttttta	caagaaaaaa	gaaattagat	aaatctctca	tatcttttat	tcaataatcg	7080
catccgattg	cagtataaat	ttaacgatca	ctcatcatgt	tcataattat	cagagctcgt	7140
gctataatta	tactaatttt	ataaggagga	aaaaatatgg	gcatttttag	tatttttgta	7200
atcagcacag	ttcattatca	accaaacaaa	aaataagtgg	ttataatgaa	tcgttaataa	7260
gcaaaattca	tataaccaaa	ttaaagaggg	ttataatgaa	cgagaaaaat	ataaaacaca	7320
gtcaaaactt	tattacttca	aaacataata	tagataaaat	aatgacaaat	ataagattaa	7380
atgaacatga	taatatcttt	gaaatcggct	caggaaaagg	ccattttacc	cttgaattag	7440
taaagagggtg	taatttcgta	actgccattg	aaatagacca	taaattatgc	aaaactacag	7500
aaaaataaact	tgttgatcac	gataatttcc	aagttttaaa	caaggatata	ttgcagttta	7560
aatttcctaa	aaaccaatcc	tataaaatat	atggtaatat	accttataac	ataagtcagg	7620
atataatacg	caaaattggt	tttgatagta	tagctaata	gattttattta	atcgtggaat	7680
acgggttttg	taaaagatta	ttaaatacaa	aacgctcatt	ggcattactt	ttaatggcag	7740
aagttgatata	ttctatatta	agtatgggtc	caagagaata	ttttcatcct	aaacctaaag	7800
tgaatagctc	acttatcaga	ttaagtagaa	aaaaatcaag	aatatcacac	aaagataaac	7860
aaaagtataa	ttatttcggt	atgaaatggg	ttaacaaaga	atacaagaaa	atatttacaa	7920
aaaatcaatt	taacaattcc	ttaaaacatg	caggaattga	cgattttaa	aatattagct	7980
ttgaacaatt	cttatctctt	ttcaatagct	ataaattatt	taataagtaa	gttaagggat	8040
gcataaaactg	catcccttaa	cttggttttc	gtgtgcctat	tttttgtaa	tcgacctgca	8100
ggcatgcaag	ctttttcaat	tcacccgtca	cagtcctcag	atgattgatc	ac	8152

KRAV

1 En rekombinant *Bacillus*-celle til fremstilling af et heterologt exoprotein af interesse, hvor cellen omfatter et nukleinsyrekonstrukt, der koder for det heterologe exoprotein af interesse og:

- a) en heterolog promoter, der er operabelt forbundet med i det mindste et gen, der koder for MrgA-protein med en aminosyresekvens, som er i det mindste 70% identisk med aminosyresekvensen vist i SEQ ID NO:2; eller
- b) i det mindste et heterologt gen, der koder for MrgA-protein med en aminosyresekvens, som er i det mindste 70% identisk med aminosyresekvensen vist i SEQ ID NO:2.

2. Celle ifølge krav 1, som er valgt fra gruppen bestående af *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus coagulans*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus stearothermophilus*, *Bacillus subtilis*, og *Bacillus thuringiensis*.

3. Celle ifølge ethvert af kravene 1 - 2, hvor det heterologe exoprotein er en protease, en lipase, en cutinase, en amylase, en galactosidase, en pullulanase, en cellulase, en glucoseisomerase, en proteindisulfidisomerase, en CGT'ase (cyclodextrinluconotransferase), en phytase, en glucoseoxidase, en glucosyltransferase, en laktase, en bilirubinoxidase, en xylanase, et antigen mikrobielt eller protozoanprotein, et bakterielt proteintoxin, et mikrobielt overfladeprotein eller et viralt protein.

4. Celle ifølge ethvert af kravene 1 - 3, hvor MrgA-proteinet omfatter aminosyresekvensen vist i SEQ ID NO:2.

5. Celle ifølge ethvert af kravene 1 - 4, som omfatter i det mindste et exogent gen, der koder for MrgA-proteinet omfattende en aminosyresekvens, som er i det mindste 70% identisk med aminosyresekvensen vist i SEQ ID NO:2.

6. Celle ifølge ethvert af kravene 1 - 5, som omfatter i det mindste et exogent gen, der koder for MrgA-protein omfattende aminosyresekvensen som vist i SEQ ID NO:2.

7. Celle ifølge ethvert af kravene 1 - 6, som omfatter i det mindste et exogent gen, der koder for MrgA omfattende en polynukleotidsekvens, som er i det mindste 70% identisk med sekvensen vist i SEQ ID NO:1.

8. Celle ifølge ethvert af kravene 1 - 7, hvor i det mindste et exogent gen, der koder for et MrgA-protein med en aminosyresekvens, som er i det mindste 70% identisk med aminosyresekvensen vist i SEQ ID NO:2, er integreret i cellens genom.

9. Celle ifølge ethvert af kravene 1 - 7, hvor i det mindste et exogent gen, der koder for et MrgA-protein med en aminosyresekvens, som er i det mindste 70% identisk med aminosyresekvensen vist i SEQ ID NO:2, er til stede på et ekstrakromosonalt konstrukt, fortrinsvis et plasmid.

10. Fremgangsmåde til fremstilling af et heterologt exoprotein af interesse, omfattende trinene:

- a) dyrkning af cellen ifølge ethvert af kravene 1 - 9; og
- b) indvinding af det heterologe exoprotein.