



(86) **Date de dépôt PCT/PCT Filing Date:** 2006/03/02
(87) **Date publication PCT/PCT Publication Date:** 2006/09/14
(45) **Date de délivrance/Issue Date:** 2016/07/12
(85) **Entrée phase nationale/National Entry:** 2007/08/31
(86) **N° demande PCT/PCT Application No.:** EP 2006/002095
(87) **N° publication PCT/PCT Publication No.:** 2006/094773
(30) **Priorités/Priorities:** 2005/03/03 (EP05075519.8);
2006/01/05 (EP06075024.7)

(51) **Cl.Int./Int.Cl.** **A01H 1/00** (2006.01),
A01H 1/02 (2006.01), **A01H 4/00** (2006.01),
A01H 5/00 (2006.01), **C12N 15/00** (2006.01),
C12N 5/04 (2006.01)
(72) **Inventeurs/Inventors:**
VAN DUN, CORNELIS MARIA PETRUS, NL;
DIRKS, ROBERT HELENE GHISLAIN, NL
(73) **Propriétaire/Owner:**
RIJK ZWAAN ZAADTEELT EN ZAADHANDEL B.V., NL
(74) **Agent:** GOWLING WLG (CANADA) LLP

(54) **Titre : PSEUDO-RETROCROISEMENT**
(54) **Title: NEAR REVERSE BREEDING**

(57) **Abrégé/Abstract:**

The present invention relates to a method for producing a homozygous non-human organism from a heterozygous non-human organism, which homozygous organism can be crossed to obtain a hybrid, comprising providing a heterozygous starting organism; allowing the organism to produce SDR-O cells through meiosis, which cells originate from second division restitution; regenerating SDR-O organisms from the SDR-O cells; and producing the homozygous organism from the SDR-O organisms thus obtained. The invention further relates to a method for producing a hybrid, comprising crossing a first homozygous organism that is produced according to the above method with a second homozygous organism. Furthermore, the invention provides homozygous non-human organism and hybrid non-human organisms obtainable by these methods. The invention relates in particular to plants.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 September 2006 (14.09.2006)

PCT

(10) International Publication Number
WO 2006/094773 A3

(51) International Patent Classification:
C12N 15/82 (2006.01)

(21) International Application Number:
PCT/EP2006/002095

(22) International Filing Date: 2 March 2006 (02.03.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
05075519.8 3 March 2005 (03.03.2005) EP
06075024.7 5 January 2006 (05.01.2006) EP

(71) Applicant (*for all designated States except US*): **RIJK ZWAAN ZAADTEELT EN ZAADHANDEL B.V.** [NL/NL]; Burg. Crezeelaan 40, NL-3678 KX De Lier (NL).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **VAN DUN, Cornelis, Maria, Petrus** [NL/NL]; Faunaberg 40, NL-4708 CC Roosendaal (NL). **DIRKS, Robert, Helene, Ghislain** [NL/NL]; Lollestraat 16, NL-4731 GL Oudenbosch (NL).

(74) Agent: **VAN KOOLJ, Adriaan**; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report*

(88) Date of publication of the international search report:
15 February 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: NEAR REVERSE BREEDING

(57) Abstract: The present invention relates to a method for producing a homozygous non-human organism from a heterozygous non-human organism, which homozygous organism can be crossed to obtain a hybrid, comprising providing a heterozygous starting organism; allowing the organism to produce SDR-O cells through meiosis, which cells originate from second division restitution; regenerating SDR-O organisms from the SDR-O cells; and producing the homozygous organism from the SDR-O organisms thus obtained. The invention further relates to a method for producing a hybrid, comprising crossing a first homozygous organism that is produced according to the above method with a second homozygous organism. Furthermore, the invention provides homozygous non-human organism and hybrid non-human organisms obtainable by these methods. The invention relates in particular to plants.



WO 2006/094773 A3

NEAR REVERSE BREEDING

The present invention relates to a method for producing a homozygous non-human organism from a heterozygous non-human organism, which homozygous organism can be crossed to obtain a hybrid. The invention relates in particular to plants.

Plant breeding is one of the fundamental occupations of mankind that is of pivotal importance for providing domesticated species that feed the world. Plant breeding is by consequence old and was originally based on selecting and propagating those plants that were outperforming in local selection fields.

Contemporary plant breeding is highly depending on knowledge of genetics and technologically supported by methods such as doubled haploids (DH) (see fi. Haploids in Crop Improvement II eds; Palmer C, Keller W, and Kasha K (2005) in: Biotechnology in Agriculture and Forestry 56 Eds; Nagata T, Lörz H, and Widholm J. Springer-Verlag Berlin Heidelberg New York, ISBN 3-500-22224-3) and molecular markers (see fi. De Vienne ed. (2003) Molecular Markers in Plant Genetics and Biotechnology. Science publishers Inc. Enfield, NH USA. ISBN 1-57808-239-0).

Genetic mechanisms during sexual reproduction have evolved to increase genetic variation which enhances the chances of survival of a species in a changing environment. Meiotic recombination, independent chromosome assortment and the mating system are main contributing factors in this respect. However, in plant breeding these mechanisms may act counterproductive especially in those cases when genetically heterozygous plants have been identified with high agronomic or horticultural value. Redistribution of genetic factors results in the generation of genetically dissimilar and

therefore heterogeneous plants and by consequence loss of commercially desirable traits.

In order to counter this effect, a number of technologies is available to the plant breeder. One possibility is to propagate plants vegetatively which leads to a complete preservation of their genetic composition as multiplication occurs exclusively through mitosis. For many plant species, *in vitro* tissue culture is being used to vegetatively propagate plants although other methods like producing cuttings *in vivo* may be applicable as well.

A disadvantage of vegetative propagation when compared to propagation through seeds is the fact that it is labour intensive and thereby costly. Furthermore, it is difficult to store plants for longer periods of time posing logistic problems and the risks of infections of the plant material with pathogens, like viruses, is considerably larger as compared to a situation in which plant material is propagated through seeds.

As an alternative, vegetative propagation may be achieved through the formation of asexual seeds, which is generally referred to as apomixis. Apomixis which occurs naturally in a number of species may be induced in sexually propagating plant species by genetic engineering. Currently, however, the genes responsible for the different steps of apomixis i.e. apomeiosis, parthenogenesis and autonomous endosperm development have not yet been identified and may interact in a complicated manner. Therefore, although the potential of apomixis technology for plant breeding is widely recognised for already a long period of time, proof of concept is still awaited.

As yet another alternative, use can be made of reverse breeding technology as described in WO-03017753. Reverse breeding is based on the suppression of meiotic

recombination through genetic engineering and the subsequent production of doubled haploid plants (DHs) derived from spores containing unrecombined parental chromosomes. These DHs differ with respect to their genetic composition solely
5 as a consequence of the independent parental chromosome assortment which occurred during meiosis. Therefore, it is sufficient to make use of one co-dominant, polymorphic marker per chromosome to determine which of the DHs or lines derived therefrom should be combined through crossing to reconstruct
10 the genetic composition of the original starting plant. As such, application of reverse breeding technology allows genetic preservation of any selected fertile plant through seeds even if its genetic composition is unknown.

However, a disadvantage of this technology is the
15 fact that complete suppression of meiotic recombination results in the absence of chiasmata and thereby inappropriate chromosome segregation during meiosis I which could lead to aneuploidy of the gametes and thereby reduced viability. When no chiasmata are formed during meiosis I, every chromosome
20 has an independent 50% chance to move to either one of the poles. This means that the theoretical chance to make a spore with a full chromosome complement is $(\frac{1}{2})^x$ wherein x represents the haploid chromosome number. The frequency of balanced gametes therefore decreases with increasing haploid
25 chromosome number.

Although many crop species have a relatively low chromosome number (e.g. cucumber has 7 chromosomes per haploid genome and spinach has only 6) there are also economically important species with relatively high
30 chromosome numbers like tomato, one of the largest vegetable crops, which has 12 chromosomes per haploid genome. This technical constraint significantly reduces the efficiency of reverse breeding technology. Therefore, a clear need in the

art exists for alternative methods which allow preservation of genetic composition in sexual offspring.

It is therefore the object of the present invention to provide a method for preserving the genetic composition of a parent organism, in particular a parent plant.

In the research that led to the present invention it was surprisingly found that by making use of plants regenerated from unreduced spores, for example as can be obtained as a result of a second division restitution (SDR) event, so-called "SDR-0 cells", it is possible to provide such method. This method can also be performed on other non-human organisms than plants, such as fungi or fish, and with other reproductive cells, such as gametes.

The invention thus relates to a method for producing a homozygous non-human organism from a heterozygous non-human organism, which homozygous organism can be crossed to obtain a hybrid, comprising:

- a) providing a heterozygous starting organism;
- b) allowing the organism to produce SDR-0 cells originating from second division restitution;
- c) regenerating SDR-0 organisms from the SDR-0 cells; and
- d) producing the homozygous organism from the SDR-0 organisms thus obtained.

This method will be called herein "Near Reverse Breeding".

Unreduced spores are formed preferentially as a consequence of the omission of the second meiotic division. This natural phenomenon is known as Second Division Restitution or SDR. SDR can occur during sexual reproduction in plants concomitantly with regular meiotic events. The Near Reverse Breeding technology of the invention exploits SDR events by specifically selecting unreduced spores that are

produced through natural or engineered SDR for regeneration. The resulting plants, called SDR-0 plants, which are largely homozygous, are, in a preferred embodiment used to produce DHs. However, the level of homozygosity in the SDR-0 plants
5 can also be increased through inbreeding steps or secondary SDR events or combinations thereof.

Molecular markers which are polymorphic between the paternal and maternal genomes of the starting plant can be used to identify those SDR-0 plants and DHs derived therefrom
10 which are essentially complementary with respect to their genetic composition and which upon crossing result in the near complete reconstruction of the genetic make-up of the original starting plant.

The reconstruction is "near complete" as a
15 consequence of meiotic recombination during the formation of the SDR-0 events and during the formation of the DHs derived therefrom. The reconstructed hybrids will genetically differ to some extent from each other as well as from the original starting hybrid plant. However, this variation is strongly
20 reduced in comparison to a situation in which the DHs are derived directly from a regular meiotic event. Moreover, the DHs are genetically fixed which means there is no room for further selection.

The advantage of integrating an SDR event in this
25 process is that the selection for genetic complementarity occurs in a two step process. The first step is concentrated on the proximal regions of the chromosomes i.e. including the centromeres. The second step is directed towards the distal ends of the chromosomes i.e. those regions which were
30 exchanged due to recombination. This delayed genetic fixation reduces the complexity and increases the chances of finding largely complementary genotypes especially when molecular markers are available for selection.

A further advantage of this approach is the fact that SDR is a natural process which occurs during sexual reproduction and which can be exploited as such without further need to interfere with sexual reproduction processes.

Various embodiments of the invention relate to a method for producing a homozygous plant from a heterozygous plant, which homozygous plant can be crossed to obtain a hybrid, comprising: a) subjecting a heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells, wherein the genetic modification is by stable incorporation into the genome of a genetic element increasing the number of second division restitution events in the plant, and wherein the environmental stress is selected from temperature stress, NO₂, nitrous oxide N₂O, or combinations thereof; b) regenerating SDR-0 plants from the SDR-0 cells; and c) producing the homozygous plant from the SDR-0 plants thus obtained. Some embodiments relate to a method for producing a hybrid comprising crossing a first homozygous plant as defined herein with a second homozygous plant.

Various embodiments of the invention relate to a method for introgressing a trait in a plant with a desired genetic background, comprising the steps of: a) crossing a first plant having the desired genetic background with a second plant having the desired trait to obtain a heterozygous starting plant; b) subjecting the heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells, wherein the genetic modification is by stable incorporation into the genome of a genetic element increasing the number of second division restitution events in the plant, and wherein the environmental stress is selected from temperature stress, NO₂, nitrous oxide N₂O, or combinations thereof; c) regenerating SDR-0 plants from the SDR-0 cells; d) selecting the SDR-0 plants that have the desired trait and the desired background; and e) producing homozygous plants from the SDR-0 plants selected.

Various embodiments of the invention relate to use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a homozygous plant that is partially complementary to a parental line of said heterozygous starting plant.

Various embodiments of the invention relate to use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a hybrid plant that genetically resembles the heterozygous starting plant such that the genetic resemblance with the heterozygous starting plant comprises the hybrid having at least 50% of the heterozygosity of the starting plant.

Figures 1 and 2 schematically illustrate the difference between a normal meiotic event (followed by the formation of DHs) and an SDR event. In both figures four chromosomal pairs have been depicted and the homologs are shown in light, respectively, dark rod like structures where the black circles on the rods depict the centromeres. The figures merely serve to illustrate the principle of this invention. It is obvious to those skilled in the art that in practice the number of chromosomal pairs involved depends on the species involved. Furthermore, the points of crossing-over are variable with respect to their number/chromosome and their position on the chromosome.

Figure 1 depicts a normal meiosis and the doubling (which can be spontaneous or induced) of the chromosomes after both meiotic divisions took place. In this case crossing-over has led to the occurrence of two parental and two recombinant chromosomes per pair. In addition, due to the independent assortment of the two homologous chromosomes of each pair, it is obvious that many genetically different spores/gametes can be formed.

In **Figure 1**, only three arbitrary outcomes of this process have been depicted. After regenerating plants from such spores, DHs can be obtained. The production of DHs is of utmost importance in contemporary plant breeding and is applied as an established technology for most crops. Plants regenerated from diploid spores will

6b

be further named DH-O i.e. doubled haploid as the primary regenerant from a spontaneous or induced diploidized spore originating from a normal meiotic event. The term "SDR-O" is used for the

primary regenerant from the cell or spore that lacked the second meiotic division. DH-0 plants when self-pollinated will give rise to progeny (DH-1) plants which are genetically 100% identical and completely fixed in all the alleles. So, 5 although the spores (gametes) that are formed on the DH-0 plants underwent again meiosis and recombination, no genetic rearrangements can take place. This therefore means that this so-called "pure line" is immortalised by the fact that no segregation can take place. Such a line can however 10 phenotypically show different appearance when grown in different conditions such as low or high temperatures, or for instance in different climatic zones. The differences that can be observed are however valid for all "members" of the line in other words there will be no "intra-line" variation. 15 There may, however, be differences between different pure lines (DH-1), which is called "inter-line" variation.

Figure 2 depicts an SDR event. In contrast to **Figure 1** where a spontaneous or induced chromosome doubling took place after completion of meiosis, the occurrence of diploid 20 spores is caused by the absence of the second meiotic division.

The fundamental difference between DHs and diploid SDR plants is illustrated by the fact that heterozygous segments are present on the different chromosomes in the SDR 25 plant, whereas DHs are fully homozygous. It should further be noted that in SDR plants all chromosomal pairs are homozygous with respect to their centromeric regions. Heterozygosity if present, resides at the distal chromosomal ends. This contrasts to another aberrant meiotic event called First 30 Division Restitution or FDR which is characterised by the absence of the first meiotic division and which results in heterozygosity at the centromeric regions for all chromosomal pairs.

In the theoretical case shown in **Figure 2**, the starting plant (donor-plant) that was used to generate DHs and SDRs, respectively, contains homologous chromosomes that are completely heterozygous. This means that all alleles of the genes carried by those chromosomes are polymorphic. In practice however this is highly unlikely and therefore this case exemplifies the most extreme heterozygous situation.

From **Figure 2** it is also clear that for an SDR event the crossing-over points of each chromosomal pair determines the ratio between homozygous loci and heterozygous loci. This ratio increases for those SDR-events which on average have their crossing-over positions located more towards the telomere whereas it decreases when on average the crossing over positions are located more towards the centromer. With the availability of sufficient molecular markers these crossing-over points can easily be determined for each SDR event.

The extent of crossing-over for each chromosomal arm is limited by the position of the centromere. It should further be noted that in case the residual heterozygosity is relatively low the SDR events resemble the occurrence of RIL's and BIL's but in heterozygous forms.

SDR is just one form of a broader class of phenomena leading to the formation of unreduced spores/gametes (Veilleux, Plant Breeding Reviews 3, 253-288 (1985)) describes the mechanisms by which unreduced gametes are formed and provides a list of the occurrence of unreduced gametes in crops plants. At that time mainly two different classes of unreduced gametes were recognised namely SDR and FDR. Recently a third class of unreduced gametes has been published named Indeterminate Meiotic Restitution or IMR (Lim et al. (2001) Theor. Appl. Genet. 103:219-230).

For the purpose of this invention only SDR is relevant. SDR occurs naturally in a wide variety of crops as is evidenced by the list presented by Veilleux, supra, and other independent research (Lim K et al. (2004) Breeding Science 54: 13-18). Interestingly, it was found that in pepper the frequency of SDR 2n gametes (pollen) increased from less than 1% to up to 10,5% (average) by 48 hours exposure of the plants to 11°C (Zhang X et al. (2002) Journal of Horticultural Science & Biotechnology 78: (1) 84-88). The maximum frequency of SDR occurrence was measured to be 81.3%. It is thus possible to increase the number of SDR events by external stimuli.

The occurrence of 2n spores or gametes is not restricted to the male gametophyte but there is also evidence that it occurs at the level of the female gametophyte. For example Zagorcheva L (Genetics and Plant Breeding 9(5), 386-399 (1976)) reported the occurrence of deviations of macrosporo-and macrogametogenesis in cucumber.

Near Reverse Breeding exploits the occurrence of unreduced spores which are the result of SDR in order to reconstruct the genetic composition of heterozygous starting plant material to a large extent. Basically, in order to reconstruct the original hybrid plant one starts to select those SDR-0 plants which are complementary for the centromeric region of each chromosome. This can be achieved by genotyping the centromeric region for each chromosome using polymorphic molecular markers. As SDR-0 regenerants are homozygous at the centromeric region for each chromosome, fully (or partly) complementary plants can readily be identified in this way. The probability to find a complementary SDR-0 plant (on the basis of centromere complementarity, and without taking any residual

heterozygosity into account) when 1 random SDR-0 plant has been selected is $(1/2)^x$ where x is the number of chromosomes.

In case the original parental lines are to be reconstructed, the probability of finding one parent is $(1/2)^{x-1}$ and for the other parent the probability is $(1/2)^x$. As stated earlier, such "near parental" lines can also be considered to be Back-cross Inbred Lines (BIL's) whereby the introgression segments are generated for both parents. These complementary plants can subsequently be crossed in order to reconstruct the genotype of the original hybrid plant.

However, since, due to meiotic recombination, the distal regions of the chromosomal arms of the SDR-0 plants will be heterozygous (in case of a single crossing-over per chromosome arm) segregation will occur in the reconstructed hybrid which may lead to a certain level of non-uniformity. When the genetic information located in the distal regions of the complementary SDR-0 events is not contributing significantly towards phenotypic variation because recombination has occurred at relatively distant chromosomal positions and therefore gene content is low or because the allelic variation does not contribute significantly to phenotypic variation, the reconstructed hybrid will be relatively uniform.

In a preferred embodiment of the invention SDR-0 events are generated in which the meiotic recombination occurred exclusively at the extreme telomeric regions of the chromosomes. In such events the chromosomes have been physically recombined but not genetically.

In order to obtain genetically fully homogeneous reconstructed hybrid plants, DHs are produced from each of the complementary SDR-0 events. This principle which is schematically illustrated in **Figure 3** depicts the formation

of spores/gametes that would occur for the plant that is regenerated from SDR-0 event 3 from **Figure 2**.

The spores which are formed can be regenerated and upon chromosome doubling DHs can be produced. By using
5 molecular markers one can select those DHs which are genetically very similar to either one of the parental lines. This is illustrated in **Figure 4** by the chromosomal set which has been doubled.

Due to segregation, the distal ends of the
10 chromosomes (from the recombination break point to the telomer) of these doubled haploid plants will on average contain 50% of the genetic information of each parent. When two randomly chosen doubled haploid plants derived from complementary SDR-0 events are combined through crossing, the
15 proximal arm regions will be 100% heterozygous whereas the distal arm regions will be 50% heterozygous in the reconstructed hybrid plant. As SDR-0 events are assumed to be 60% homozygous on average (Carputo, D. et al. (2003) Genetics 163, 287-294), reconstruction with complementary events will
20 result in 80% heterozygosity on average ($100\% \times 60\% + 50\% \times 40\%$) and 20% homozygosity. The homozygosity between the proximal and distal crossing-over point will be skewed to the genotype of the SDR-0 regenerant which was homozygous for this region, whereas distal to the distal crossing-over
25 point, the homozygosity will be equal for both parental genotypes.

It is clear to the person skilled in the art that the mentioned figures relating to the percentages of heterozygosity represent extreme values i.e. starting from a
30 plants which is 100% heterozygous which means all alleles of the genes carried on the chromosomes are polymorphic. In practice this is highly unlikely and therefore the percentages of heterozygosity will on average be lower.

11a

The invention further relates to a method for producing a homozygous plant from a heterozygous plant, which homozygous plant can be crossed to obtain a hybrid, comprising:

a) subjecting a heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells;

b) regenerating SDR-0 plants from the SDR-0 cells; and

c) producing the homozygous plant from the SDR-0 plants thus obtained.

The invention further relates to a method for introgressing a trait in a plant with a desired genetic background, comprising the steps of:

a) crossing a first plant having the desired genetic background with a second plant having the desired trait to obtain a heterozygous starting plant;

b) subjecting the heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells;

c) regenerating SDR-0 plants from the SDR-0 cells;

d) selecting the SDR-0 plants that have the desired trait and the desired background; and

e) producing homozygous plants from the SDR-0 plants selected.

The invention further relates to a use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a homozygous plant that is partially complementary to a parental line of said heterozygous starting plant.

The invention further relates to a use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a hybrid plant that genetically resembles the heterozygous starting plant such that the genetic resemblance with the heterozygous starting plant comprises the hybrid having at least 50% of the heterozygosity of the starting plant.

The invention further relates to a method for producing a hybrid, comprising crossing a first homozygous organism that is produced according to the invention with a second homozygous organism. According to the invention it is possible to reconstruct the genetic make-up of the original hybrid whereas, in addition, variants can be obtained which approach the genetic make-up of the original hybrid.

In a first embodiment, the second homozygous organism is at least partially complementary to the first homozygous organism such that the resulting hybrid resembles the heterozygous starting organism. Suitably, the resemblance with the heterozygous starting organism comprises the hybrid having at least 50% of the heterozygosity of the starting organism. Preferably, the resemblance in heterozygosity is any percentage between 50 and 100%, more specifically the resemblance in heterozygosity is any percentage between 50% and 60%, preferably between 60% and 70%, more preferably between 70% and 80%, even more preferably between 80% and 90% and most preferably between 90% and 100%. The phrase "any percentage" is intended to cover each and every percentage within the stated range even though the particular percentage is not explicitly mentioned.

Alternatively, the second homozygous organism is selected such that the resulting hybrid outperforms the original heterozygous starting organism. The "original heterozygous starting organism" is the organism that is used in step a) of claim 1.

The variation of the relative level of homozygosity as well as its parental origin in the reconstructed hybrid plant depends on the crossing-over point within each chromosomal arm of each parent. By using molecular markers both SDR-0 events as well as DHs derived therefrom can be

selected which upon hybridisation result in F1 hybrids with relatively low levels of homozygosity.

In this embodiment, for the SDR-0 events one should select for those having on average more distal crossing-overs
5 whereas for the DHs one should select those being most complementary.

On the other hand, however, it may be desirable to select those events which have a relatively large distance between the proximal and distal crossing-over point to
10 introduce homozygosity in the F1 hybrid which can be skewed to either parent.

From the above it is clear that Near Reverse Breeding enables a plant breeder to reconstruct a hybrid plant to a large extent but that significant variation due to different
15 levels and origins of the alleles which are homozygous can also be obtained between the experimental F1 hybrids which may result in improvement of the performance of the original (starting) hybrid plant.

In addition, it is possible to produce hybrids which
20 are complementary only for a given subset of chromosomes. This can be done by selecting those SDR-0 events which on the basis of their centromeric genotype are complementary for the desired chromosomal pairs and identical for the others. Such hybrids produced from so-called near-substitution lines will
25 be largely homozygous for non-complementary chromosomal pairs with similar levels of heterozygosity at the distal regions of the chromosomes as compared to the complementary chromosomal pairs. In its most extreme form, fully non-complementary double haploids can be crossed which leads
30 to hybrid plants with heterozygosity limited to the distal parts of the chromosomes.

In order to exploit SDR for Near Reverse Breeding technology, use can be made of the spontaneous occurrence of

SDR events or one can induce SDR events through e.g. genetic engineering.

Another embodiment of the inventions relates to backcrossing. Backcrossing is a plant breeding term which is well known to the person skilled in the art and which refers to a procedure in which a specific trait is introgressed into a plant line with a desired genetic make-up (the so-called genetic background). The desired outcome of this procedure is a plant line which is genetically nearly identical to the starting plant line but to which only the genetic factors underlying the desired trait are added through recombination. In order to achieve this objective the plants carrying the desired genetic background and the desired trait are crossed and the offspring is selected for plants carrying the trait as well as, as much as possible of the genetic background. As with each backcross cycle the increment of the desired genetic background is on average halved, in practice it takes 4 to 5 breeding cycles to complete the backcrossing procedure.

Near reverse breeding allows to speed up backcrossing as the procedure to reach homozygosity takes only two steps. The first step is identical to a traditional backcrossing procedure i.e. crossing of the plants carrying the desired genetic background and the desired trait. The resulting hybrid is used to generate SDR-0 events which are selected for the desired trait which can be based on the phenotype or marker assisted and the desired genetic background based on centromer markers specific for the genetic background.

The second step involves the production of DHs which are selected for the maximum amount of genetic background using molecular markers. Near reverse breeding fastens the procedure to come to homozygosity but allows two selection steps. When sufficient genetic markers are available the desired backcrossing product can be obtained much more

efficient as compared with the traditional procedure as the number of time consuming breeding cycles can significantly be reduced.

With the near reverse breeding method of the invention
5 CMS can be introgressed into a desired background in a very efficient manner. In order to apply near reverse breeding for CMS transfer, the CMS donor line is preferably genetically dissimilar to the line that has to be converted to male sterility for a large number of nuclear genetic markers so
10 that the difference between the chromosomes of the CMS and the fertile donor can be more easily determined.

In order to convert a pure line or a desired inbred line (homozygous or nearly homozygous) into a similar line but with a CMS background, a first cross is made by pollination of
15 the said CMS with pollen of the desired line. The resulting F1 progeny contains CMS and 50% of the chromosomes of the desired line. The F1 progeny plants are induced to perform SDR-meiosis during gynogenesis either by treatment with chemicals such as nitrous oxide or stress conditions such as low temperature.
20 SDR may also occur spontaneously during gynogenesis.

In case androgenesis is applied, use has to be made of a restorer gene which can suppress the effect of the male sterility inducing cytoplasm.

The resulting SDR-0 plants are genetically analysed
25 using DNA markers which are polymorphic for the centromeric regions of the F1 hybrid. SDR-0 plants which exclusively contain the centromeric regions of the line which has to be converted into a CMS are selected. In case large numbers of discriminating markers covering the genome are available, one
30 can select for those SDR-0 events which on average have the most proximal crossing over positions on the chromosomes. This largely converts a selected line into a CMS in a single step. If needed, this backcross cycle can be reiterated.

The spontaneous occurrence of SDR events may be enhanced by specific abiotic stress conditions such as heat or cold shocks. Such stress conditions are known to enhance the formation of unreduced, diploid spores (Zhang X et al., (2002) 5 Journal of Horticultural Science & Biotechnology 78: (1) 84-88). In addition, it is possible to induce the formation of 2n pollen by applying nitrous oxide (N₂O) gas (Okazaki, K. et al. (2005) Euphytica, 143, 101-114).

When diploid spores are produced through male meiosis 10 it is also possible to enrich for these cells through flow cytometry and fluorescence activated cell sorting. Such technologies are in itself well known to the person skilled in the art and have been applied to microspores in the past (Deslauriers C et al., (1991) Flow cytometric characterisation 15 and sorting of cultured Brassica napus microspores. Biochem. Biophys. Acta: 1091, 165-172).

In addition, it is known that diploid spores or pollen are larger in size than their haploid peers (Lernmi G and Negri V. (1994) Research on 2n pollen production in Lotus 20 tenuis an I.M.G.V. of Perugia University. Lotus newsletter Vol; 25, pp 24-27). Surprisingly, the mere fact that diploid spores are physically different from haploid spores makes it possible to enrich specifically for diploid spores through flow cytometry and fluorescence activated cell sorting.

25 Furthermore, the optimisation of the environmental conditions which lead to the formation of diploid spores can easily be done using the different sorting technologies.

Regeneration of the unreduced spores may occur through androgenesis, gynogenesis or parthenogenesis by prick 30 pollination in a manner similar to the regeneration of reduced spores. In case parthenogenesis by prick pollination is applied, it may be advantageous to use diploid pollen, especially for those species which have a low tolerance for

modifications of the ratio of maternal to paternal genomes in endosperm. For most, if not all crops these regeneration protocols are described in the literature and known to the person skilled in the art.

5 Once plants are regenerated from diploid spores, molecular markers can be used to determine whether or not the centromeric regions of the chromosomal pairs are homozygous or heterozygous which is diagnostic for SDR or FDR events, respectively. In this way SDR events can be readily selected.

10 Different genetic approaches are known to a person skilled in the art which allow interference with gene functions involved in the second cell division of meiosis. Such interference can either be through mutagenesis or transgenesis. Transgenic approaches aim at the stable or
15 transient introduction of a DNA fragment which modifies the second division of meiosis leading to diploid spores of the SDR type. This modification can occur through interference with genetic factors involved in meiotic processes especially those involved in the second cell division. The interference
20 can be established through specific down-regulation of gene expression based on post-transcriptional gene silencing (PTGS). PTGS can be achieved through RNA-interference (RNAi) or virus-induced gene silencing (VIGS).

 Yet in an other approach, the interference can be
25 established through the over-expression of proteins which exert a dominant negative effect on the second division of meiosis leading to SDR.

 Irrespective of the approach taken, the target gene needs to be known at the molecular level. A number of
30 recessive mutants have been described of potato (pcpc, osos, fcfc) which result in an SDR-type of meiosis (Carputo, D. et al (2003) Genetics 163, 287-294). Also for maize the elongatel

mutation leads to the absence of meiosis II (Barell, PJ and Grossniklaus, U. (2005) Plant J. 43, 309-320).

Although the genes which have been mutated in these specific examples have not yet been identified at the
5 molecular level the skilled person will be able to do so and therefore these and other yet unknown genes are excellent candidates to achieve SDR in target species using molecular suppression technologies. The present invention relates to the general principle of near reverse breeding and the fact that
10 not all possible embodiments of inducing SDR in a starting organism have been described is not relevant for the invention.

Alternatives to the genes described above can be found in genes like DUET (Venkata Reddy et al. (2003) Development
15 130, 5975-5987) and CYC1;2 (Wang et al. (2004) Plant Physiology 136, 4127-4135) which have been described for *Arabidopsis thaliana* and which upon mutation lead to an aberrant form of meiosis.

The diploid meiotic products in these mutants are
20 SDR-like and therefore DUET and CYC1;2 as well as their functional homologues in other plant species are candidate target genes to achieve an SDR-type of meiosis.

Yet another candidate target gene is TETRASPORE/STUD (Yang et al (2003) Plant J. 34, 229-240) which upon knock out
25 leads to absence of cell division after meiosis. Diploid regenerants of microspores of a tetraspore/stud mutant can be SDR-like.

Once SDR-0 plant have been obtained they can be further characterised molecularly. Initially, the haplotypes
30 of the centromeric regions can be determined which provides insight in the level of complementarity of the homologous chromosomes between the SDR-0 plants. Depending on the application one can select fully or partially complementary

SDR-0 plants. The SDR-0 plants may subsequently be used to generate more dense haplotype maps. This can be achieved using molecular typing technologies well known to the person skilled in the art. Examples are RFLP (Restriction Fragment Length Polymorphism (Beckmann, J.S. and Soller, M (1983) Theor. and Appl. Genet. 67, 35-43)), RAPD (Random Amplified Polymorphic DNA (Welsh, J. and McClelland, M. (1990) Nucleic Acids Res. 19, 961-866)), AFLP (Amplified Fragment Length Polymorphism (Vos, P. et al (1995) Nucleic Acids Res. 23, 4407-4414)) or SFP (Single Feature Polymorphism (Borovitz, J. et al (2003) Genome Research 13, 513-523). These technologies can be used without prior knowledge of the nature of the DNA polymorphisms.

In case the DNA polymorphisms are characterised e.g. as SNPs use can be made of a plethora of SNP detection technologies (Kwok, P.Y. and Chen, X (2003) Curr. Issues Mol. Biol. 5, 43-60). The haplotype maps can be used to select plants for DH production. The DHs obtained can be haplotyped for the regions which were heterozygous in the SDR-0 starting plant. Such analysis provides the genetic information which can be used to steer the ratio heterozygosity/homozygosity in the F1 hybrid and it provides the possibility to select for either of the parental origins of the homozygous regions of the genome of the F1 hybrid.

The probability of finding at least one complementary combination of two homozygous organisms (a combination that after crossing can 'resynthesize' the starting organism), is a function of the haploid chromosomal number x of a given species and the number k of homozygous organisms produced from a heterozygous starting organism in which SDR has taken place.

When the haploid chromosomal number of a given crop species is expressed as x , the maximum number of SDR-0 genotypes, considering only the chromosomal regions proximal

of the crossing over points and thus including the centromeric region, which are obtained from a plant of that crop species is 2^x . The probability that one randomly chosen pair of SDR-0 plants, or a DH plant derived therefrom, from this population will, upon crossing, result in an F1 hybrid which has a genotype near identical to the genotype which produced the SDR-0 events (original genotype) is $(2^x - 1)/2^x$. In case a total number of k SDR-0 plants is produced, there exists a number of $1/2k(k-1)$ combinations of 2 genetically distinct SDR-0 plants or DH plants derived therefrom which can be crossed. The probability for any randomly chosen combination of 2 SDR-0 plants or DH plants derived therefrom to be complementary is $1/2^x$. Thus, the probability for any randomly chosen combination of 2 SDR-0 plants or DHs therefrom that they are not complementary is $1 - 1/2^x = (2^x - 1)/2^x$. In case of k SDR-0s or DHs derived therefrom, $1/2k(k-1)$ combinations can be made and therefore the probability that within this SDR-0-population or DHs derived therefrom no complementary plants can be found is $((2^x - 1)/2^x)^{(1/2k(k-1))}$ and therefore the probability that at least one complementary combination of two SDR-0 plants or DHs derived therefrom can be found is $1 - ((2^x - 1)/2^x)^{(1/2k(k-1))}$. The result of this analysis is shown in table 1 below.

Table 1: *The probability of finding at least one combination of two complementary SDR-0 or DHs derived therefrom, using near reverse breeding technology as a function of the haploid chromosome number x and the number of available randomly produced SDR-0 plants k*

x/k	2	4	8	16	24	32	48	64	128	256
7	0.008	0.046	0.197	0.610	0.885	0.980	1.000	1.000	1.000	1.000
9	0.002	0.012	0.053	0.209	0.417	0.621	0.890	0.981	1.000	1.000
11	0.000	0.003	0.014	0.057	0.126	0.215	0.424	0.626	0.981	1.000
12	0.000	0.001	0.007	0.029	0.065	0.114	0.241	0.388	0.863	1.000

This analysis shows that the original genotype can be largely resynthesized as an F1 hybrid according to the present invention with high probability using 48 SDR-0 plants or DHs derived therefrom for plant species with a haploid chromosomal number of 7 like cucumber, 128 for a plant species with a haploid chromosomal number of 9 like cauliflower and 256 for a plant species with a haploid chromosomal number of 12 like tomato, melon and sweet pepper. These figures illustrate the fact that the number of SDR-0 plants required for applying near reverse breeding for these crop species is relatively low and therefore the generation of such numbers is highly feasible, especially in an industrial environment. It is obvious to the person skilled in the art that such calculations can be made for any plant species for which the haploid chromosomal number is known.

"SDR-0 plants or cells" as used in this application is intended to relate to the plants and cells resulting from unreduced gametes. Such unreduced gametes may in turn be the result of a Second Division Restitution (SDR) event, but may also be the product of an aberrant form of meiosis which leads to diploids.

The invention thus provides the means to use organisms, in particular plants, and their progenies, regenerated from SDR or SDR-like unreduced gametes, for example for the purpose of genotype reconstruction, for
5 producing near complementary parental lines, for making near chromosome substitution lines and for optimizing F1 hybrids by introduction of homo- and heterozygous genomic segments.

The present invention will be further illustrated in the non-limiting examples that follow and that refer to the
10 following figures:

Figure 5 shows the AFLP Patterns of typical F2 plants of cucumber. Every horizontal line represents one individual plant. Every vertical column represents a linkage group. Light grey segments represent heterozygous area's, black and dark
15 areas represent respective homozygous areas.

Figure 6 shows the AFLP analysis of typical DH lines of cucumber. Every horizontal line represents one individual plant. Every vertical column represents a linkage group. Only black and dark area's are present as expected in DHs Light
20 grey segments are absent.

Figure 7 shows the AFLP analysis of typical SDR-0 plants in cucumber. Every horizontal line represents 1 individual plant. Every vertical column represents a linkage group. Light grey segments represent heterozygous area's,
25 black and dark areas represent respective homozygous areas.

Figure 8 shows pollen collected from pepper plants treated with low temperature. Panel A represents the pollen of the cold-treated plants whereas panel B represents the pollen of the control plant as observed through the light microscope.
30

EXAMPLES

EXAMPLE 1

Demonstration of the occurrence of a relatively low level of heterozygosity in cucumber (*Cucumis sativus* L.) plants obtained through gynogenesis

By making haploids and doubled haploids in cucumber according to EP-374755 it was found by using AFLP analysis (carried out according to EP-534858) that among the expected doubled haploids, a certain percentage proved to be not originated from haploid megaspores but rather to be derived from an unreduced megaspore (2n). This is well demonstrated in **Figures 5, 6 and 7** which shown that the originally presumed doubled haploids (**Figure 6**) still contain heterozygous sectors, which by definition is not possible in true doubled haploids.

This example shows that through gynogenesis in cucumber DHs can be obtained but that in addition plants are obtained which show some regions which are heterozygous. This level of heterozygosity is strongly reduced as compared to the F2 generation and is therefore most likely caused by an SDR-like mechanism.

20

EXAMPLE 2

Enhancement of the formation of unreduced spores/gametes in sweet pepper (*Capsicum annuum* L.)

In order to increase the frequency of unreduced spore/gamete formation, cold stress was applied as an inducer. For this purpose, flowering plants of sweet pepper containing pre-meiotic floral buds and growing at 23°C were exposed for 5 days to 11°C according to Zhang et al. (2002) *Journal of Horticultural Science & Biotechnology* 78, 84-88. After this cold shock, the buds were harvested and pollen were extracted by opening the anthers using dissecting forceps and scalpel. The pollen were subsequently transferred on a microscopic glass slide and stained for viability using a drop of

aceto-carmin. Cover slides were put on top of the suspension which was investigated using light-microscopy. As a control, pollen was collected from sweet pepper plants which were grown at 23°C.

5 **Figure 8** shows representative examples of the morphologies of the pollen collected from the cold-treated plants (8A) versus the control plants (8B). As can be seen, the number of pollen with an increased size indicative for being derived from unreduced spores is strongly increased for
10 the cold-treated plant. In this particular example it was estimated that the % of unreduced spores mounted up to 25 due to the cold treatment. As such the enhancement of the formation of unreduced spores by temperature stress is shown to be highly feasible.

15

EXAMPLE 3

Enrichment of naturally occurring diploid spores in broccoli (*Brassica oleracea* L.) through flow cytometry

For broccoli (and other plant species) it is known
20 that diploid spores are larger than haploid spores. In order to determine whether it is feasible to enrich for diploid spores using flow cytometry different mixtures were made of spores obtained from diploid and tetraploid plants. Spores were isolated by grinding floral buds (3-4 mm in size) in a
25 buffer solution (8.2 g/L NaCl, 1.9 g/L Na₂HPO₄·2H₂O, 0.3 g/L NaH₂PO₄·2H₂O, pH=7.4) and subsequent filtering through a 110 µm filter. The cells were washed twice with the extraction buffer and counted.

The following mixtures were made of haploid versus
30 diploid cells: 1:1, 10:1 and 100:1. Sorting was carried out on the basis of the parameters granularity, vitality and size. Enrichment of diploid cells was obtained for all ratios and based on each of the three parameters. In a next experiment a

natural population of haploid microspores was sorted based on the above mentioned parameters. It was found that diploid microspores could be sorted which were estimated to be present in a frequency of 0.7% in the microspore population. **Figure 9** shows the purified diploid microspores (lower row) obtained in this experiment compared to normal haploid microspores as observed through the light microscope. Pictures were obtained by courtesy of Ing. M. Vennik; TNO Leiden (The Netherlands).

10 **EXAMPLE 4**

Construction of RNA interference vectors which can be used to downregulate candidate target gene Tetraspore

In order to downregulate the activity of a candidate target gene in a particular plant species, use can be made of RNA interference. For that purpose DNA fragments of the Tetraspore of *Arabidopsis thaliana* were inserted into pKANNIBAL (Wesley et al. (2001) *The Plant Journal* 27, 581-590) such that upon expression in plants an RNA molecule is formed which can fold back upon itself thus forming a hairpin structure that triggers the specific degradation of homologous RNA.

The vector pKANNIBAL contains an intron positioned downstream of the CaMV 35S promoter and upstream from a octopine synthase polyadenylation signal. At either side of the intron a multiple cloning site is positioned which allows convenient insertion of the left and right arm of DNA corresponding to the RNA interference target in a inverted orientation relative to each other. Upon transcription the intron is removed by splicing and the left and right arm fold back on each other forming the double stranded RNA.

Two cDNA fragments were isolated corresponding to the 5'-end of the gene (588 bp, forward primer: 5'-ACC TCC GAG AAC TCC GTT AAG-3', reverse primer: 5'-TGC CTG CTT TCT ACC ACT

TC-3') and the middle part of the gene (679 bp, forward primer: 5'-TTC TCA AGT GGC AAG GTG TC-3'; reverse primer: 5'-ATC CCT CTT TGG TGG AGT AG-3'). Restriction enzyme recognition sites were generated at both sides of each
5 fragment which allowed the insertion of the cDNA fragments as an inverted repeat in pKANNIBAL. The left arm for both fragments is engineered as an XhoI-KpnI fragment whereas the right arm for both fragments is engineered as a HindIII-XbaI fragment. As a final step the complete hairpin cassettes,
10 containing the both Tetraspore sequences as inverted repeat, are inserted separately into a T-DNA of a binary vector called pART27 which contains the neomycin phosphotransferase II gene as selectable marker for plant transformation. The integrity of the T-DNA was confirmed by sequence analysis.

15 The resulting binary vectors, denominated pRZ226 for the 5'-fragment and pRZ219 for the 3'-fragment are transferred into *Agrobacterium tumefaciens* using a triparental mating procedure. In case the sequence similarity between a fragment of a cloned candidate target gene and the specific gene that
20 needs to be downregulated by RNA interference is too low, the method described above can be used to make a similar construct containing DNA fragments which are sufficiently homologous to the gene that needs to be downregulated.

25 **EXAMPLE 5**

Transformation of *Arabidopsis thaliana* with pRZ226 and pRZ219

Agrobacterium tumefaciens strain C58 containing the plant transformation vectors pRZ226 and pRZ219 were grown overnight in LB medium containing streptomycin(100 mg/L) and
30 spectinomycin (300 mg/L) to select for the vectors and rifampicin (40 mg/L) and gentamycin (25 mg/L) to select for the *Agrobacterium tumefaciens* C58 background at 29°C.

In order to produce transgenic plants, the *Arabidopsis* floral dip method was used largely as described by Desfeux et al (2000) *Plant Physiology* 123, 895-904. The bacterial cells were resuspended in floral dip solution (50 g sucrose + 500
5 µl silwett L-77 surfactant per litre milliQ). Bolting plants, containing multiple floral buds, were submerged into the dipping solution containing the *Agrobacterium* cells at an OD between 1.0 and 1.5 during 5-10 seconds with gentle agitation. After inoculation, the plants were contained in a plastic
10 container to keep high humidity under low light conditions for a day and subsequently, seeds were grown on the plants.

Transformants were selected by germinating surface sterilised seeds in 0.1 % agarose layered upon half-strength MS plates containing 50 mg/L kanamycin. Kanamycin resistant
15 seedlings were transferred to soil in a greenhouse.

EXAMPLE 6

Naturally occurring SDR event in *Brassica oleracea*

Hybrids produced on red cabbage (*Brassica oleracea*)
20 carrying Ogura cytoplasmic male sterility (CMS) through a wide cross were screened for the occurrence of diploid plants which were phenotypically nearly identical to the original red cabbage CMS plant and which therefore are the result of *in vivo* gynogenesis. These plants which can easily be
25 distinguished from true hybrid plants on the basis of their phenotype were indeed identified with low frequency.

Such plants originate through parthenogenesis from reduced gametes which doubled spontaneously or from unreduced gametes. In order to discriminate between these possibilities,
30 the DNA derived from such plant was analysed using AFLP (carried out according to EP-534 858). The analysis demonstrated the presence of a relatively low frequency of heterozygous loci indicating that this plant originated from

an unreduced gamete. This example shows that through *in vivo* gynogenesis in *Brassica oleracea*, plants can be obtained with a relatively low level of heterozygosity which is likely caused by an SDR-like mechanism.

5 **Fig. 10** shows the AFLP fingerprint of a *Brassica oleracea* plant derived from an unreduced gamete. Grey lines indicates a homozygous marker call from one parent, dark grey from the other. Light grey indicates a hybrid marker locus.

10 **EXAMPLE 7**

Near reverse breeding in maize

Incorporation of nucleic acids in the genome of maize are routine procedures known in the art and methods how to achieve this have been described (EP-801134; US-5,489,520; EP-
15 97.114654.3). Using any one of the transformation methods described in these patent applications, nucleic acid sequences were introduced that confer a specific inhibitory effect on a gene or genes which are involved in the second meiotic division, in particular elongat1 (Barell, PJ and
20 Grossniklaus, U. (2005) *Plant J.* 43, 309-320) and which as a consequence lead to the occurrence of aberrant meiotic events of the SDR type. The frequency of SDR-events obtained sometimes differs between independent transformants as a
25 consequence of different genomic sites of integration of the transgenic nucleic acid sequences.

In another experiment the frequency of SDR spores was increased by treatment of maize plants with low temperatures or by applying nitrous oxide gas as described by Kato, A and Birchler, JA (2006) *J. Hered.* 1, 39-44.

30 As a consequence of the activity of the said inhibitory nucleic acids or application of low temperatures or nitrous oxide treatments, numerous microspores respectively macrospores, were produced which are of the SDR-type.

The cell population thus produced was enriched for the presence of SDR-0 microspores by using flow cytometry and fluorescence activated cell sorting based on the fact that SDR microspores are larger in size as compared to normal haploid
5 microspores.

The microspores or macrospores which were produced as a consequence of an SDR-event contain a diploid set of chromosomes. These diploid microspores or macrospores were the starting material for producing SDR-0 regenerants.

10 Haploid maize plants have also been obtained following natural and artificial pollination with a haploid inducer. In this case seeds are obtained that contain haploid embryos see fi. Rotarencu V (2002) Production of matroclinous maize haploids following natural and artificial pollination
15 with a haploid inducer. Maize Genetics Cooperation News Letter 76: 16. The mentioned protocols to produce DH maize plants have also be applied to produce SDR-0 maize embryo''s from SDR-0 cells, of which the formation is induced by the treatments specified in this example. In order to obtain
20 the proper balance between the maternal and paternal genomes in the endosperm of the SDR-0 kernels, use has also been made of a tetraploid inducer line.

In the specific experiment in which transgenesis is applied to induce SDR, transformants that contained a single
25 copy of the transgene were preferred. After obtaining a transgenic line containing a nucleic acid construct that largely inhibits the second meiotic division, this line was used in crosses in order to avoid repetitive transformation events. In that case the frequency of normal pollen containing
30 haploid gametes was low but still sufficient to be used for crossing.

In a further experiment, an inhibitory nucleic acid construct was used which is controlled by a promoter that is

chemically inducible by e.g. dexamethasone (Bohner, S. et al (1999) *Plant J.* 19, 87-95).

Haploids in maize were then routinely obtained from microspores (Pescitelli S and Petolino J (1988) *Plant Cell Reports* 7: 441-444; Coumans M et al., (1989) *Plant Cell Reports* 7: 618-621; Pescitelli S et al., (1989) *Plant Cell Reports* 7: 673-676; Buter B (1997) *In vitro* haploid production in maize. In: *In Vitro Haploid Production in Higher plants*, vol 4, 37-71. Kluwer Academic Publishers. Eds.; S Jain, S Sopory & R Veilleux).

Haploid maize plants were also be obtained following natural and artificial pollination with a haploid inducer (Rotarenco V (2002) *Maize Genetics Cooperation News Letter* 76: 16). In this case seeds were obtained that contain haploid embryos.

The above-mentioned protocols to produce DH maize plants were applied to produce SDR-0 maize plants from SDR-0 cells, of which the formation is induced transgenically or by the other treatments specified in this example.

The SDR-0 maize plants were genetically characterised with respect to the centromeric regions of each chromosome using a molecular marker which is polymorphic for these regions in the original starting plant. For the detection of the polymorphism different technologies, selected from CAPS, dCAPS, Invader, pyrosequencing, taqman were used.

Thus, couples of SDR-0 maize plants were identified which are complementary for all chromosomal pairs with respect to the marker scores of the centromeric regions. Such couples of SDR-0 maize plants were then used to produce doubled haploid plants for each SDR-0 complementary plant. The individual DHs obtained from each complementary SDR-0 plant were crossed pairwise i.e. the two DHs which are used to produce a maize F1 hybrid originate from either one of the

complementary SDR-0 events. These crosses were made reciprocally. The F1 hybrids thus produced were evaluated in field trials for agronomic performance in which the starting F1 hybrid was used as a control.

- 5 It was found that the agronomic performance of the hybrid obtained through near reverse breeding of the invention is similar to the performance of the original hybrid.

EXAMPLE 8Near reverse breeding in cucumber

Gynogenesis is a well established technology for cucumber and is carried out according to the method described
5 in EP 0 374 755. Spontaneous SDR events which occur during gynogenesis when applied according to EP 0 374 755 led to the formation of diploid regenerants which have some residual heterozygosity. These SDR-0 cucumber plants were identified using AFLP analysis carried out according to the method
10 provided in EP 534858.

The SDR-0 cucumber plants thus obtained were genetically characterised with respect to the centromeric regions of each chromosome using molecular markers which are polymorphic for these regions in the original starting plant.
15 For the detection of the polymorphisms, known technologies selected from CAPS, dCAPS, Invader, pyrosequencing, taqman were used. This resulted in the identification of couples of SDR-0 cucumber plants which were complementary for all chromosomal pairs with respect to the marker scores of the
20 centromeric regions. Such couples of SDR-0 cucumber plants were subsequently used to produce DH plants for each SDR-0 complementary plant according to the method described in EP 0 374 755.

Spontaneous occurring SDR events were discarded and
25 only true doubled haploid plants were used in further steps. In case haploid plants were obtained the chromosome number is doubled e.g. by applying colchicine. The chromosome doubling can also occur spontaneously. The individual DH plants obtained from each complementary SDR-0 plant were subsequently
30 crossed pairwise i.e. the two DHs which were used to produce a cucumber F1 hybrid originated from either one of the complementary SDR-0 events. These crosses were made reciprocally. The F1 hybrids thus produced were evaluated in

trials for agronomic performance in which the starting F1 hybrid is used as a control.

It was found that the agronomic performance of the hybrid obtained through near reverse breeding of the invention
5 is similar to the performance of the original F1 hybrid.

CLAIMS

1. Method for producing a homozygous plant from a heterozygous plant, which homozygous plant can be crossed to obtain a hybrid, comprising:
 - a) subjecting a heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells, wherein the genetic modification is by stable incorporation into the genome of a genetic element increasing the number of second division restitution events in the plant, and wherein the environmental stress is selected from temperature stress, NO₂, nitrous oxide N₂O, or combinations thereof;
 - b) regenerating SDR-0 plants from the SDR-0 cells; and
 - c) producing the homozygous plant from the SDR-0 plants thus obtained.
2. The method as claimed in claim 1, wherein the genetic modification is transient.
3. The method as claimed in claim 1 or 2, wherein the homozygous plants are produced by doubled haploid techniques, inbreeding, second division restitution, or combinations thereof.
4. Method for producing a hybrid, comprising crossing a first homozygous plant that is produced according to any one of claims 1-3 with a second homozygous plant.
5. The method as claimed in claim 4, wherein the second homozygous plant is at least partially complementary to the first homozygous plant such that the resulting hybrid genetically resembles the heterozygous starting plant such that the genetic resemblance with the heterozygous starting plant comprises the hybrid having at least 50% of the heterozygosity of the starting plant.
6. The method as claimed in claim 5, wherein the genetic resemblance in heterozygosity is any percentage between 50 and 100%.

7. The method as claimed in claim 5 or 6, wherein the genetic resemblance in heterozygosity is any percentage between 50% and 60%.
8. The method as claimed in claim 5 or 6, wherein the genetic resemblance in heterozygosity is any percentage between 60% and 70%.
9. The method as claimed in claim 5 or 6, wherein the genetic resemblance in heterozygosity is any percentage between 70% and 80%.
10. The method as claimed in claim 5 or 6, wherein the genetic resemblance in heterozygosity is any percentage between 80% and 90%.
11. The method as claimed in claim 5 or 6, wherein the genetic resemblance in heterozygosity is any percentage between 90% and 100%.
12. The method as claimed in claim 4, wherein the second homozygous plant is selected such that the resulting hybrid outperforms the heterozygous starting plant.
13. The method as claimed in claim 4, wherein the second homozygous plant is selected such that only a subset of its chromosomes is complementary to the corresponding chromosomes of the first homozygous plant.
14. The method as claimed in claim 13, wherein the remainder of the chromosomes is identical between the first and second homozygous plants.
15. Method for introgressing a trait in a plant with a desired genetic background, comprising the steps of:
 - a) crossing a first plant having the desired genetic background with a second plant having the desired trait to obtain a heterozygous starting plant;
 - b) subjecting the heterozygous starting plant to genetic modification or environmental stress to produce an above-average number of second division restitution-0 (SDR-0) cells, wherein the genetic modification is by stable incorporation into the genome of a genetic element increasing the number of second division restitution events

in the plant, and wherein the environmental stress is selected from temperature stress, NO₂, nitrous oxide N₂O, or combinations thereof;

- c) regenerating SDR-0 plants from the SDR-0 cells;
- d) selecting the SDR-0 plants that have the desired trait and the desired background; and
- e) producing homozygous plants from the SDR-0 plants selected.

16. The method as claimed in claim 15, wherein the genetic modification is transient.

17. The method as claimed in any one of claim 15 or 16, wherein the homozygous plants are produced by doubled haploid techniques, inbreeding, second division restitution, or combinations thereof.

18. The method as claimed in any one of claims 15-17, wherein the selection is performed on the basis of visual inspection or by means of molecular markers.

19. Use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a homozygous plant that is partially complementary to a parental line of said heterozygous starting plant.

20. Use of second division restitution-0 (SDR-0) cells originating from second division restitution in a heterozygous starting plant, for producing a hybrid plant that genetically resembles the heterozygous starting plant such that the genetic resemblance with the heterozygous starting plant comprises the hybrid having at least 50% of the heterozygosity of the starting plant.

Fig. 1A






Fig. 1B

Meiosis 2: formation of spores/gametes



3 possible examples of spores/gametes



Corresponding "Doubled Haploids"





Fig. 2

Second division does not take place = second division restitution




4 examples of spores/gametes (SDR-0):
note the partial heterozygosity


(1)

(2)

(3)








(4)







Note that in the 3th and 4th example the basic make up of the chromosome sets are derived from the respective parents; such sets resemble "BIL's" but in heterozygous state. Examples 1 and 2 resemble "RIL's" in a heterozygous state.

Fig. 3A

(3)



Meiosis 1: chromosome
doubling



Meiosis 1: recombination
took place

Fig. 3B-1

Possible gametes from 1 recombination event
from 1 SDR event




Fig. 3B-2




Fig. 3B-3




Fig. 3B-5




Fig. 4

Figure 4. Doubled haploid (left) form a specific spore/gamete from an SDR-0 event resembling one of the original parental lines (right).



Fig. 5




Fig. 6




Fig. 7




Fig. 8A




Fig. 8B




Fig. 9




Fig. 10

