



(86) Date de dépôt PCT/PCT Filing Date: 2010/12/15
(87) Date publication PCT/PCT Publication Date: 2011/06/23
(45) Date de délivrance/Issue Date: 2018/06/12
(85) Entrée phase nationale/National Entry: 2012/06/08
(86) N° demande PCT/PCT Application No.: DK 2010/050344
(87) N° publication PCT/PCT Publication No.: 2011/072697
(30) Priorités/Priorities: 2009/12/17 (US61/287,321);
2009/12/17 (DK PA 2009 01340)


(51) Cl.Int./Int.Cl. *C07D 471/04* (2006.01),
A61K 31/4196 (2006.01), *A61K 31/437* (2006.01),
A61P 25/18 (2006.01), *A61P 25/28* (2006.01)

(72) Inventeurs/Inventors:
PUSCHL, ASK, DK;
NIELSEN, JACOB, DK;
KEHLER, JAN, DK;
KILBURN, JOHN PAUL, DK;
MARIGO, MAURO, DK;
LANGGARD, MORTEN, DK

(73) Propriétaire/Owner:
H. LUNDBECK A/S, DK

(74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : DERIVES HETEROAROMATIQUES D'ARYLTRIAZOLE EN TANT QU'INHIBITEURS DE L'ENZYME PDE10A
(54) Title: HETEROAROMATIC ARYL TRIAZOLE DERIVATIVES AS PDE10A ENZYME INHIBITORS



(57) **Abrégé/Abstract:**

This invention is directed to compounds (Formula 1), which are PDE10A enzyme inhibitors. The invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. The present invention also provides processes for the preparation of the compounds of formula (I). The present invention further provides a method of treating a subject suffering from a neurodegenerative disorder comprising administering to the subject a therapeutically effective amount of a compound of formula (I). The present invention also provides a method of treating a subject suffering from a drug addiction comprising administering to the subject a therapeutically effective amount of a compound of formula (I). The present invention further provides a method of treating a subject suffering from a psychiatric disorder comprising administering to the subject a therapeutically effective amount of a compound of formula (I).

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
23 June 2011 (23.06.2011)

PCT

(10) International Publication Number
WO 2011/072697 A1

(51) International Patent Classification:

C07D 471/04 (2006.01) *A61P 25/18* (2006.01)
A61K 31/4196 (2006.01) *A61P 25/28* (2006.01)
A61K 31/437 (2006.01)

(21) International Application Number:

PCT/DK2010/050344

(22) International Filing Date:

15 December 2010 (15.12.2010)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

PA 2009 01340 17 December 2009 (17.12.2009) DK
 61/287,321 17 December 2009 (17.12.2009) US

(71) **Applicant** (for all designated States except US): **H. LUNDBECK A/S** [DK/DK]; Ottiliavej 9, DK-2500 Valby (DK).

(72) **Inventors; and**

(75) **Inventors/Applicants** (for US only): **PÜSCHL, Ask** [DK/DK]; Harsdorffsvej 7A, 4.tv., DK-1874 Frederiksberg C (DK). **NIELSEN, Jacob** [DK/DK]; Dannebrogsgade 17, t.th., DK-1660 København V (DK). **KEHLER, Jan** [DK/DK]; Nymøllevej 28, DK-2800 Lyngby (DK). **KILBURN, John, Paul** [GB/DK]; Kærsgårdvej 7, DK-4690 Haslev (DK). **MARIGO, Mauro** [IT/DK]; Jyttevej 23, DK-2740 Skovlunde (DK). **LANGGÅRD, Morten** [DK/DK]; Rågevej 20, DK-2600 Glostrup (DK).

(74) **Common Representative:** **H. LUNDBECK A/S**; Corporate Patents and Trademarks, Ottiliavej 9, DK-2500 Valby (DK).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) **Title:** HETEROAROMATIC ARYL TRIAZOLE DERIVATIVES AS PDE10A ENZYME INHIBITORS

(57) **Abstract:** This invention is directed to compounds (Formula 1), which are PDE10A enzyme inhibitors. The invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of the invention and a pharmaceutically acceptable carrier. The present invention also provides processes for the preparation of the compounds of formula (I). The present invention further provides a method of treating a subject suffering from a neurodegenerative disorder comprising administering to the subject a therapeutically effective amount of a compound of formula (I). The present invention also provides a method of treating a subject suffering from a drug addiction comprising administering to the subject a therapeutically effective amount of a compound of formula (I). The present invention further provides a method of treating a subject suffering from a psychiatric disorder comprising administering to the subject a therapeutically effective amount of a compound of formula (I).



WO 2011/072697 A1

Heteroaromatic aryl triazole derivatives as PDE10A enzyme inhibitors

Field of the Invention

- 5 The present invention provides heteroaromatic compounds that are PDE10A enzyme inhibitors, and as such are useful to treat neurodegenerative and psychiatric disorders. The present invention also provides pharmaceutical compositions comprising compounds of the invention and methods of treating disorders using the compounds of the invention.

10

Background of the Invention

15

- The cyclic nucleotides cyclic-adenosine monophosphate (cAMP) and cyclic-guanosine monophosphate (cGMP) function as intracellular second messengers regulating a vast array of processes in neurons. Intracellular cAMP and cGMP are generated by adenylyl and guanylyl cyclases, and are degraded by cyclic nucleotide phosphodiesterases (PDEs). Intracellular levels of cAMP and cGMP are controlled by intracellular signaling, and stimulation/repression of adenylyl and guanylyl cyclases in response to GPCR activation is a well characterized way of controlling cyclic nucleotide concentrations (Antoni, F.A. *Front. Neuroendocrinol.* **2000**, 21, 103-132). cAMP and cGMP levels in turn control activity of cAMP- and cGMP-dependent kinases as well as other proteins with cyclic nucleotide response elements, which through subsequent phosphorylation of proteins and other processes regulate key neuronal functions such as synaptic transmission, neuronal differentiation and survival.

30

There are 21 phosphodiesterase genes that can be divided into 11 gene families. There are ten families of adenylyl cyclases, two of guanylyl cyclases, and eleven of phosphodiesterases. PDEs are a class of intracellular enzymes

that regulate levels of cAMP and cGMP via hydrolysis of the cyclic nucleotides into their respective nucleotide monophosphates. Some PDEs degrade cAMP, some cGMP and some both. Most PDEs have a widespread expression and have roles in many tissues, while some are more tissue-specific.

5

Phosphodiesterase 10A (PDE10A) is a dual-specificity phosphodiesterase that can convert both cAMP to AMP and cGMP to GMP (Loughney, K. *et al. Gene* **1999**, 234, 109-117; Fujishige, K. *et al. Eur. J. Biochem.* **1999**, 266, 1118-1127 and Soderling, S. *et al. Proc. Natl. Acad. Sci.* **1999**, 96, 7071-10 7076). PDE10A is primarily expressed in the neurons in the striatum, n. accumbens and in the olfactory tubercle (Kotera, J. *et al. Biochem. Biophys. Res. Comm.* **1999**, 261, 551-557 and Seeger, T.F. *et al. Brain Research*, **2003**, 985, 113-126).

15 Mouse PDE10A is the first identified member of the PDE10 family of phosphodiesterases (Fujishige, K. *et al. J. Biol. Chem.* **1999**, 274, 18438-18445 and Loughney, K. *et al. Gene* **1999**, 234, 109-117) and N-terminal splice variants of both the rat and human genes have been identified (Kotera, J. *et al. Biochem. Biophys. Res. Comm.* **1999**, 261, 551-557 and Fujishige, K. 20 *et al. Eur. J. Biochem.* **1999**, 266, 1118-1127). There is a high degree of homology across species. PDE10A is uniquely localized in mammals relative to other PDE families. mRNA for PDE10 is highly expressed in testis and brain (Fujishige, K. *et al. Eur J Biochem.* **1999**, 266, 1118-1127; Soderling, S. *et al. Proc. Natl. Acad. Sci.* **1999**, 96, 7071-7076 and Loughney, K. *et al.* 25 *Gene* **1999**, 234,109-117). These studies indicate that within the brain, PDE10 expression is highest in the striatum (caudate and putamen), n. accumbens and olfactory tubercle. More recently, an analysis has been made of the expression pattern in rodent brain of PDE10A mRNA (Seeger, T.F. *et al. Abst. Soc. Neurosci.* **2000**, 26, 345.10) and PDE10A protein (Menniti, F.S. *et al.* 30 William Harvey Research Conference 'Phosphodiesterase in Health and Disease', Porto, Portugal, Dec. 5-7, 2001).

PDE10A is expressed at high levels by the medium spiny neurons (MSN) of the caudate nucleus, the accumbens nucleus and the corresponding neurons

of the olfactory tubercle. These constitute the core of the basal ganglia system. The MSN has a key role in the cortical-basal ganglia-thalamocortical loop, integrating convergent cortical/thalamic input, and sending this integrated information back to the cortex. MSN express two functional classes
5 of neurons: the D₁ class expressing D₁ dopamine receptors and the D₂ class expressing D₂ dopamine receptors. The D₁ class of neurons is part of the 'direct' striatal output pathway, which broadly functions to facilitate behavioral responses. The D₂ class of neurons is part of the 'indirect' striatal output pathway, which functions to suppress behavioral responses that compete with
10 those being facilitated by the 'direct' pathway. These competing pathways act like the brake and accelerator in a car. In the simplest view, the poverty of movement in Parkinson's disease results from over-activity of the 'indirect' pathway, whereas excess movement in disorders such as Huntington's disease represent over-activity of the direct pathway. PDE10A regulation of
15 cAMP and/or cGMP signaling in the dendritic compartment of these neurons may be involved in filtering the cortico/thalamic input into the MSN. Furthermore, PDE10A may be involved in the regulation of GABA release in the substantia nigra and globus pallidus (Seeger, T.F. *et al. Brain Research*, **2003**, 985, 113-126).

20 Dopamine D₂ receptor antagonism is well established in the treatment of schizophrenia. Since the 1950's, dopamine D₂ receptor antagonism has been the mainstay in psychosis treatment and all effective antipsychotic drugs antagonise D₂ receptors. The effects of D₂ are likely to be mediated primarily
25 through neurons in the striatum, n. accumbens and olfactory tubercle, since these areas receive the densest dopaminergic projections and have the strongest expression of D₂ receptors (Konradi, C. and Heckers, S. *Society of Biological Psychiatry*, **2001**, 50, 729-742). Dopamine D₂ receptor agonism leads to decrease in cAMP levels in the cells where it is expressed through
30 adenylate cyclase inhibition, and this is a component of D₂ signalling (Stoof, J. C.; Keabian J. W. *Nature* **1981**, 294, 366-368 and Neve, K. A. *et al. Journal of Receptors and Signal Transduction* **2004**, 24, 165-205). Conversely, D₂ receptor antagonism effectively increases cAMP levels, and this effect could be mimicked by inhibition of cAMP degrading phosphodiesterases.

Most of the 21 phosphodiesterase genes are widely expressed; therefore inhibition is likely to have side effects. Because PDE10A, in this context, has the desired expression profile with high and relatively specific expression in
5 neurons in striatum, n. accumbens and olfactory tubercle, PDE10A inhibition is likely to have effects similar to D₂ receptor antagonism and therefore have antipsychotic effects.

While PDE10A inhibition is expected to mimic D₂ receptor antagonism in part,
10 it might be expected to have a different profile. The D₂ receptor has signalling components besides cAMP (Neve, K. A. *et al. Journal of Receptors and Signal Transduction* **2004**, 24, 165-205), wherefore interference with cAMP through PDE10A inhibition may negatively modulate rather than directly antagonise dopamine signaling through D₂ receptors. This may reduce the
15 risk of the extrapyramidal side effects that are seen with strong D₂ antagonism. Conversely, PDE10A inhibition may have some effects not seen with D₂ receptor antagonism. PDE10A is also expressed in D₁ receptors expressing striatal neurons (Seeger, T. F. *et al. Brain Research*, **2003**, 985, 113-126). Since D₁ receptor agonism leads to stimulation of adenylate cyclase and
20 resulting increase in cAMP levels, PDE10A inhibition is likely to also have effects that mimic D₁ receptor agonism. Finally, PDE10A inhibition will not only increase cAMP in cells, but might also be expected to increase cGMP levels, since PDE10A is a dual specificity phosphodiesterase. cGMP activates a number of target protein in cells like cAMP and also interacts with the cAMP
25 signalling pathways. In conclusion, PDE10A inhibition is likely to mimic D₂ receptor antagonism in part and therefore has antipsychotic effect, but the profile might differ from that observed with classical D₂ receptor antagonists.

The PDE10A inhibitor papaverine is shown to be active in several
30 antipsychotic models. Papaverine potentiated the cataleptic effect of the D₂ receptor antagonist haloperidol in rats, but did not cause catalepsy on its own (WO 03/093499). Papaverine reduced hyperactivity in rats induced by PCP, while reduction of amphetamine induced hyperactivity was insignificant (WO 03/093499). These models suggest that PDE10A inhibition has the classic

antipsychotic potential that would be expected from theoretical considerations. WO 03/093499 further discloses the use of selective PDE10 inhibitors for the treatment of associated neurologic and psychiatric disorders. Furthermore, PDE10A inhibition reverses subchronic PCP-induced deficits in attentional
5 set-shifting in rats (Rodefer *et al. Eur. J. Neurosci.* **2005**, 4, 1070-1076). This model suggests that PDE10A inhibition might alleviate cognitive deficits associated with schizophrenia.

The tissue distribution of PDE10A indicates that PDE10A inhibitors can be
10 used to raise levels of cAMP and/or cGMP within cells that express the PDE10 enzyme, especially neurons that comprise the basal ganglia, and the PDE10A inhibitors of the present invention would therefore be useful in treating a variety of associated neuropsychiatric conditions involving the basal ganglia such as neurological and psychiatric disorders, schizophrenia, bipolar
15 disorder, obsessive compulsive disorder, and the like, and may have the benefit of not possessing unwanted side effects, which are associated with the current therapies on the market.

Furthermore, recent publications (WO 2005/120514, WO 2005012485, Cantin
20 *et al*, Bioorganic & Medicinal Chemistry Letters 17 (2007) 2869–2873) suggest that PDE10A inhibitors may be useful for treatment of obesity and non-insulin dependent diabetes.

With respect to inhibitors of PDE10A, EP 1250923 discloses the use of
25 selective PDE10 inhibitors in general, and papaverine in particular, for the treatment of certain neurologic and psychiatric disorders.

WO 05/113517 discloses benzodiazepine stereospecific compounds as inhibitors of phosphodiesterase, especially types 2 and 4, and the prevention
30 and treatment of pathologies involving a central and/or peripheral disorder. WO 02/88096 discloses benzodiazepine derivatives and their uses as inhibitors of phosphodiesterase, especially type 4 in the therapeutic field. WO 04/41258 discloses benzodiazepinone derivatives and their uses as inhibitors of phosphodiesterase, especially type 2 in the therapeutic field.

Pyrrolodihydroisoquinolines and variants thereof are disclosed as inhibitors of PDE10 in WO 05/03129 and WO 05/02579. Piperidinyl-substituted quinazolines and isoquinolines that serve as PDE10 inhibitors are disclosed in
5 WO 05/82883. WO 06/11040 discloses substituted quinazoline and isoquinoline compounds that serve as inhibitors of PDE10. US 20050182079 discloses substituted tetrahydroisoquinolinyl derivatives of quinazoline and isoquinoline that serve as effective phosphodiesterase (PDE) inhibitors. In particular, US 20050182079 relates to said compounds, which are selective
10 inhibitors of PDE10. Analogously, US 20060019975 discloses piperidine derivatives of quinazoline and isoquinoline that serve as effective phosphodiesterase (PDE) inhibitors. US 20060019975 also relates to compounds that are selective inhibitors of PDE10. WO 06/028957 discloses cinnoline derivatives as inhibitors of phosphodiesterase type 10 for the
15 treatment of psychiatric and neurological syndromes.

However, these disclosures do not pertain to the compounds of the invention, which are structurally unrelated to any of the known PDE10 inhibitors (Kehler, J. *et al. Expert Opin. Ther. Patents* **2007**, 17, 147-158 and Kehler, J. *et al.*
20 *Expert Opin. Ther. Patents* **2009**, 19, 1715-1725), and which have now been found by the inventors to be highly active and selective PDE10A enzyme inhibitors.

Compounds comprising a -CH₂-S- linker and where further HET-1 is either
25 imidazo[1,2-a]pyridine or imidazo[1,2-a]pyrimidine are disclosed in publicly available chemical libraries. These compounds are therefore disclaimed.

The compounds of the invention may offer alternatives to current marketed treatments for neurodegenerative and/or psychiatric disorders, which are not
30 efficacious in all patients. Hence, there remains a need for alternative methods of treatment.

Summary of the Invention


The objective of the present invention is to provide compounds that are selective PDE10A enzyme inhibitors.

A further objective of the present invention is to provide compounds which have such activity, and which have improved solubility, metabolic stability and/or bioavailability compared to prior art compounds.


Another objective of the invention is to provide an effective treatment, in particular long-term treatment, of a human patient, without causing the side effects typically associated with current therapies for neurological and psychiatric disorders.

Further objectives of the invention will become apparent upon reading the present specification.

Accordingly, in one aspect the present invention relates to compounds of formula **I**:



wherein **HET-1** is a heteroaromatic group of formula **II** containing from 2 to 4 nitrogen atoms:



wherein Y can be N or CH, Z can be N or C, and wherein **HET-1** may optionally be substituted with up to three substituents R7, R8 and R9

individually selected from H; C₁-C₆ alkyl such as Me; halogen such as chlorine and bromine; cyano; halo(C₁-C₆)alkyl such as trifluoromethyl; aryl such as phenyl; alkoxy, such as methoxy, dimethoxy, ethoxy, methoxy-ethoxy and ethoxy-methoxy, and C₁-C₆ hydroxyalkyl such as CH₂CH₂OH, and wherein *

5 denotes the attachment point,


-L- is a linker selected from -S-CH₂-, -CH₂-S-, -CH₂-CH₂-, -CH=CH-, and -C≡C-;

10 R₁ is selected from H; C₁-C₆ alkyl such as methyl, ethyl, 1-propyl, 2-propyl, isobutyl; C₁-C₆ alkyl(C₃-C₈)cycloalkyl such as cyclopropylmethyl; C₁-C₆ hydroxyalkyl such as hydroxyethyl; CH₂CN; CH₂C(O)NH₂; C₁-C₆ arylalkyl such as benzyl and 4-chlorobenzyl; and C₁-C₆ alkyl-heterocycloalkyl such as tetrahydropyran-4-yl-methyl and 2-morpholin-4-yl-ethyl;

15

wherein Q is a phenyl, optionally substituted with 1, 2 or three substituents or Q is a monocyclic 5-membered or 6-membered heteroaromatic group containing 1 or 2 heteroatoms, preferred Q is chosen among structures of the formula, wherein "*" denotes the attachment point:

20



Wherein R2-R6 are each selected independently from H; C₁-C₆ alkoxy such as methoxy; and halogen such as chlorine or fluorine;

5 with the proviso that the compound is not 1H-Benzimidazole, 2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(2-pyrazinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; 1H-Benzimidazole, 2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]methyl]thio]-; 1H-Benzimidazole, 1-ethyl-5-(1-piperidinylsulfonyl)-2-[[[3-(2-thienyl)-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 6-methyl-
10 2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(3-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 8-methyl-2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; Imidazo[1,2-a]pyridine, 6-chloro-2-[[[3-(2-thienyl)-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(4-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 6-methyl-2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 2-
15 [[[3-(2-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 6-chloro-2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; 3H-Imidazo[4,5-b]pyridine, 2-[[[3-phenyl-1H-1,2,4-triazol-5-yl]thio]methyl]-; or 1H-Benzimidazole, 2-[[[3-(2-furanyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-;
20 and tautomers and pharmaceutically acceptable salts thereof, and polymorphic forms thereof.

In separate embodiments of the invention, the compound of formula I is
25 selected among the specific compounds disclosed in the Experimental Section herein.

The invention further provides a compound of formula I, or a pharmaceutically acceptable salt thereof, for use as a medicament.

30

In another aspect, the present invention provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I and a pharmaceutically acceptable carrier, diluent or excipient.

The invention further provides the use of a compound of formula I, or a pharmaceutically acceptable salt thereof, for the preparation of a medicament for the treatment of a neurodegenerative or psychiatric disorder.

- 5 Furthermore, in yet another aspect, the present invention provides a method of treating a subject suffering from a neurodegenerative disorder, comprising administering to the subject a therapeutically effective amount of a compound of formula I. In a still further aspect, the present invention provides a method of treating a subject suffering from a psychiatric disorder, comprising
10 administering to the subject a therapeutically effective amount of a compound of formula I. In another embodiment, the present invention provides a method of treating a subject suffering from a drug addiction, such as an alcohol, amphetamine, cocaine, or opiate addiction.

15

Detailed Description of the Invention

Definition of Substitutents

- As used in the context of the present invention, the terms "halo" and "halogen"
20 are used interchangeably and refer to fluorine, chlorine, bromine or iodine.

- The term "C₁-C₆ alkyl" refers to a straight-chain or branched saturated hydrocarbon having from one to six carbon atoms, inclusive. Examples of such groups include, but are not limited to, methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, 2-methyl-1-butyl, and n-hexyl. The
25 expression "C₁-C₆ hydroxyalkyl" refers to a C₁-C₆ alkyl group as defined above which is substituted with one hydroxy group. The term "halo(C₁-C₆)alkyl" refers to a C₁-C₆ alkyl group as defined above which is substituted with up to three halogen atoms, such as trifluoromethyl.

30

The expression "C₁-C₆ alkoxy" refers to a straight-chain or branched saturated alkoxy group having from one to six carbon atoms, inclusive, with the open valency on the oxygen. Examples of such groups include, but are not limited to, methoxy, ethoxy, n-butoxy, 2-methyl-pentoxy and n-hexyloxy.

The term "C₃-C₈ cycloalkyl" typically refers to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl. The expression "C₁-C₆ alkyl(C₃-C₈)cycloalkyl" refers to a C₃-C₈ cycloalkyl as defined above which is substituted with a straight-chain or branched C₁-C₆ alkyl. Examples of such groups include, but are not limited to, cyclopropylmethyl.

The term "heterocycloalkyl" refers to a four to eight membered ring containing carbon atoms and up to three N, O or S atoms, provided that the four to eight membered ring does not contain adjacent O or adjacent S atoms. The open valency is on either the heteroatom or carbon atom. Examples of such groups include, but are not limited to, azetidiny, oxetanyl, piperaziny, morpholinyl, thiomorpholinyl and [1,4]diazepanyl. The term "hydroxyheterocycloalkyl" refers to a heterocycloalkyl as defined above which is substituted with one hydroxy group. The term "C₁-C₆ alkyl-heterocycloalkyl" refers to a heterocycloalkyl as defined above which is substituted with a C₁-C₆ alkyl group. Examples of such groups include, but are not limited to, tetrahydropyran-4-yl-methyl and 2-morpholin-4-yl-ethyl.

The term "aryl" refers to a phenyl ring, optionally substituted with halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy or halo(C₁-C₆)alkyl as defined above. Examples of such groups include, but are not limited to, phenyl and 4-chlorophenyl.

The term "C₁-C₆arylalkyl" refers to an aryl as defined above which is substituted with a straight-chain or branched C₁-C₆ alkyl. Examples of such groups include, but are not limited to, benzyl and 4-chlorobenzyl.


Additionally, the present invention further provides certain embodiments of the invention, that are described below.

30

In one embodiment of the invention, **HET-1** is a heteroaromatic group of formula **II** containing 2 nitrogen atoms. In another embodiment of the invention, **HET-1** is a heteroaromatic group of formula **II** containing 3 nitrogen

atoms. In yet another embodiment of the invention, **HET-1** is a heteroaromatic group of formula **II** containing 4 nitrogen atoms.

HET-1 is preferably chosen among the following heteroaromatic groups,
5 wherein “*” denotes the attachment point:



10

In a further embodiment one or more of the hydrogen atoms of the compound of formula I have been substituted by deuterium. In particular hydrogen has been replaced by deuterium when R₇- R₉ is methyl or methoxy.

15 In separate embodiments of the invention, the compound of formula I is selected among the following specific compounds, in the form of the free base, one or more tautomers thereof or a pharmaceutically acceptable salt thereof. Table 1 lists compounds of the invention and the corresponding IC₅₀ values determined as described in the section “PDE10A inhibition assay”.
20 Each of the compounds constitutes an individual embodiment, of the present invention.

It should be understood that the various aspects, embodiments, implementations and features of the invention mentioned herein may be
25 claimed separately, or in any combination, as illustrated by the following non-limiting examples.

Table 1: Compounds of the invention and IC₅₀ values

<u>Compound</u>	IC50 (nM)
8-Methoxy-5-methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine	52
5-Methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine	200
5-Methyl-2-(1-methyl-5-phenyl-1H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine	280
8-Methoxy-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	25
8-Methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	170
5,7-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrimidine	59
5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	15
8-Methoxy-5-methyl-2-[2-(5-phenyl-2-propyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	37
8-Methoxy-5-methyl-2-{2-[5-phenyl-2-(2,2,2-trifluoro-ethyl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyridine	180
5-Methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	220
8-Methoxy-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	560
{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-acetic acid ethyl ester	81
2-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol	23
5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	6.7
5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-c]pyrimidine	36

8-Ethyl-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	7,5
8-Ethyl-5-methyl-2-[2-(5-phenyl-2-propyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine	10
8-Ethyl-2-[2-(2-isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridine	40
3-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile	45
3-{5-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile	14
2-[2-(2-Isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridine	56
3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile	25
3-{2-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-4-phenyl-imidazol-1-yl}-propylamine	150
3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propylamine	100
2-{2-[2-(2-Methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazol-3-yl]-ethyl}-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine	14
8-Methoxy-2-{2-[2-(2-methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazol-3-yl]-ethyl}-5-methyl-[1,2,4]triazolo[1,5-a]pyridine	62
5,8-Dimethyl-2-[2-(2-methyl-5-thiophen-3-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	5.3
2-[2-(5-Furan-2-yl-1-methyl-1H-[1,2,4]triazol-3-yl)-ethyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine	38
2-[(E)-2-(5-Furan-2-yl-1-methyl-1H-[1,2,4]triazol-3-yl)-vinyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine	60
5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-4-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	25
5,8-Dimethyl-2-{2-[2-methyl-5-(5-methyl-thiazol-2-yl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyrazine	120

5,8-Dimethyl-2-{2-[2-methyl-5-(4-methyl-thiazol-2-yl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyrazine	76
5,8-Dimethyl-2-[2-(2-methyl-5-oxazol-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	120
5,8-Dimethyl-2-[2-(2-methyl-5-thiophen-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	5.6
5,8-Dimethyl-2-[2-(2-methyl-5-pyrimidin-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	53
5,8-Dimethyl-2-[2-(2-methyl-5-pyridin-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	39
5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-5-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	26
5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	36
5,8-Dimethyl-2-[2-(2-methyl-5-pyridin-3-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine	110
2-[2-(5-Furan-2-yl-2-methyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine	4.6
5,8-Dimethyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyrazine	53
2-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol	69

In a particular embodiment of the present invention the compounds of the present invention have an IC₅₀ value of less than 50 nM, such as in the range of 0.2 – 20 nM, particularly in the range of 0.2 – 10 nM, such as in the range of 0.2 – 5 nM or in the range of 0.2 – 1 nM.

Pharmaceutically Acceptable Salts

The present invention also comprises salts of the compounds, typically, pharmaceutically acceptable salts. Such salts include pharmaceutically acceptable acid addition salts. Acid addition salts include salts of inorganic acids as well as organic acids.

Representative examples of suitable inorganic acids include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, sulfamic, nitric acids and the like. Representative examples of suitable organic acids include formic, acetic, 5 trichloroacetic, trifluoroacetic, propionic, benzoic, cinnamic, citric, fumaric, glycolic, itaconic, lactic, methanesulfonic, maleic, malic, malonic, mandelic, oxalic, picric, pyruvic, salicylic, succinic, methane sulfonic, ethanesulfonic, tartaric, ascorbic, pamoic, bismethylene salicylic, ethanedisulfonic, gluconic, citraconic, aspartic, stearic, palmitic, EDTA, glycolic, p-aminobenzoic, 10 glutamic, benzenesulfonic, p-toluenesulfonic acids, theophylline acetic acids, as well as the 8-halothephyllines, for example 8-bromothephylline and the like. Further examples of pharmaceutically acceptable inorganic or organic acid addition salts include the pharmaceutically acceptable salts listed in Berge, S.M. et al., *J. Pharm. Sci.* 1977, 66, 2.

15

Furthermore, the compounds of this invention may exist in unsolvated as well as in solvated forms with pharmaceutically acceptable solvents such as water, ethanol and the like. In general, the solvated forms are considered equivalent to the unsolvated forms for the purposes of this invention.

20

Therapeutically effective amount

In the present context, the term "therapeutically effective amount" of a compound means an amount sufficient to cure, alleviate or partially arrest the clinical manifestations of a given disease and its complications in a therapeutic intervention comprising the administration of said compound. An 25 amount adequate to accomplish this is defined as "therapeutically effective amount". Effective amounts for each purpose will depend on the severity of the disease or injury as well as the weight and general state of the subject. It will be understood that determining an appropriate dosage may be achieved using routine experimentation, by constructing a matrix of values and testing 30 different points in the matrix, which is all within the ordinary skills of a trained physician.

In the present context, the term "treatment" and "treating" means the management and care of a patient for the purpose of combating a condition, such as a disease or a disorder. The term is intended to include the full spectrum of treatments for a given condition from which the patient is suffering, such as administration of the active compound to alleviate the symptoms or complications, to delay the progression of the disease, disorder or condition, to alleviate or relief the symptoms and complications, and/or to cure or eliminate the disease, disorder or condition as well as to prevent the condition, wherein prevention is to be understood as the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of the active compounds to prevent the onset of the symptoms or complications. Nonetheless, prophylactic (preventive) and therapeutic (curative) treatments are two separate aspects of the invention. The patient to be treated is preferably a mammal, in particular a human being.

Pharmaceutical compositions

The present invention further provides a pharmaceutical composition comprising a therapeutically effective amount of a compound of formula I and a pharmaceutically acceptable carrier or diluent. The present invention also provides a pharmaceutical composition comprising a therapeutically effective amount of one of the specific compounds disclosed in the Experimental Section herein and a pharmaceutically acceptable carrier or diluent.

The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers, diluents or excipients, in either single or multiple doses. The pharmaceutical compositions according to the invention may be formulated with pharmaceutically acceptable carriers or diluents as well as any other known adjuvants and excipients in accordance with conventional techniques such as those disclosed in Remington: The Science and Practice of Pharmacy, 19th Edition, Gennaro, Ed., Mack Publishing Co., Easton, PA, 1995.

The pharmaceutical compositions may be specifically formulated for administration by any suitable route such as oral, rectal, nasal, pulmonary, topical (including buccal and sublingual), transdermal, intracisternal, intraperitoneal, vaginal and parenteral (including subcutaneous, intramuscular, intrathecal, intravenous and intradermal) routes. It will be appreciated that the route will depend on the general condition and age of the subject to be treated, the nature of the condition to be treated and the active ingredient.

Pharmaceutical compositions for oral administration include solid dosage forms such as capsules, tablets, dragees, pills, lozenges, powders and granules. Where appropriate, the compositions may be prepared with coatings such as enteric coatings or they may be formulated so as to provide controlled release of the active ingredient such as sustained or prolonged release according to methods well known in the art. Liquid dosage forms for oral administration include solutions, emulsions, suspensions, syrups and elixirs. Pharmaceutical compositions for parenteral administration include sterile aqueous and nonaqueous injectable solutions, dispersions, suspensions or emulsions as well as sterile powders to be reconstituted in sterile injectable solutions or dispersions prior to use. Other suitable administration forms include, but are not limited to, suppositories, sprays, ointments, creams, gels, inhalants, dermal patches and implants.

Typical oral dosages range from about 0.001 to about 100 mg/kg body weight per day. Typical oral dosages also range from about 0.01 to about 50 mg/kg body weight per day. Typical oral dosages further range from about 0.05 to about 10 mg/kg body weight per day. Oral dosages are usually administered in one or more dosages, typically, one to three dosages per day. The exact dosage will depend upon the frequency and mode of administration, the sex, age, weight and general condition of the subject treated, the nature and severity of the condition treated and any concomitant diseases to be treated and other factors evident to those skilled in the art.

The formulations may also be presented in a unit dosage form by methods known to those skilled in the art. For illustrative purposes, a typical unit dosage form for oral administration may contain from about 0.01 to about 1000 mg, from about 0.05 to about 500 mg, or from about 0.5 mg to about 200 mg.

For parenteral routes such as intravenous, intrathecal, intramuscular and similar administration, typical doses are in the order of half the dose employed for oral administration.

The present invention also provides a process for making a pharmaceutical composition comprising mixing a therapeutically effective amount of a compound of formula I and at least one pharmaceutically acceptable carrier or diluent. In an embodiment, of the present invention, the compound utilized in the aforementioned process is one of the specific compounds disclosed in the Experimental Section herein.

The compounds of this invention are generally utilized as the free substance or as a pharmaceutically acceptable salt thereof. One example is an acid addition salt of a compound having the utility of a free base. When a compound of formula I contains a free base such salts are prepared in a conventional manner by treating a solution or suspension of a free base of formula I with a molar equivalent of a pharmaceutically acceptable acid. Representative examples of suitable organic and inorganic acids are described above.

For parenteral administration, solutions of the compounds of formula I in sterile aqueous solution, aqueous propylene glycol, aqueous vitamin E or sesame or peanut oil may be employed. Such aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. The aqueous solutions are particularly suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. The compounds of formula I may be readily incorporated into known sterile aqueous media using standard techniques known to those skilled in the art.

Suitable pharmaceutical carriers include inert solid diluents or fillers, sterile aqueous solutions and various organic solvents. Examples of solid carriers include lactose, terra alba, sucrose, cyclodextrin, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and lower alkyl ethers of cellulose. Examples of liquid carriers include, but are not limited to, syrup, peanut oil, olive oil, phospholipids, fatty acids, fatty acid amines, polyoxyethylene and water. Similarly, the carrier or diluent may include any sustained release material known in the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax. The pharmaceutical compositions formed by combining the compounds of formula I and a pharmaceutically acceptable carrier are then readily administered in a variety of dosage forms suitable for the disclosed routes of administration. The formulations may conveniently be presented in unit dosage form by methods known in the art of pharmacy.

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules or tablets, each containing a predetermined amount of the active ingredient, and optionally a suitable excipient. Furthermore, the orally available formulations may be in the form of a powder or granules, a solution or suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsion.

If a solid carrier is used for oral administration, the preparation may be tableted, placed in a hard gelatin capsule in powder or pellet form or it may be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will range from about 25 mg to about 1 g per dosage unit. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatin capsule or sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

The pharmaceutical compositions of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by

mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine prepare tablets. Examples of adjuvants or diluents comprise: corn starch, potato starch, talcum, magnesium stearate, gelatin, lactose, gums, and the
5 like. Any other adjuvants or additives usually used for such purposes such as colorings, flavorings, preservatives etc. may be used provided that they are compatible with the active ingredients.

Treatment of Disorders

10 As mentioned above, the compounds of formula I are PDE10A enzyme inhibitors and as such are useful to treat associated neurological and psychiatric disorders.

The invention thus provides a compound of formula I or a pharmaceutically
15 acceptable acid addition salt thereof, as well as a pharmaceutical composition containing such a compound, for use in the treatment of a neurodegenerative disorder, psychiatric disorder or drug addiction in mammals including humans; wherein the neurodegenerative disorder is selected from the group consisting of Alzheimer's disease, multi-infarct
20 dementia, alcoholic dementia or other drug-related dementia, dementia associated with intracranial tumors or cerebral trauma, dementia associated with Huntington's disease or Parkinson's disease, or AIDS-related dementia; delirium; amnesic disorder; post-traumatic stress disorder; mental retardation; a learning disorder, for example reading disorder, mathematics disorder, or a
25 disorder of written expression; attention-deficit/hyperactivity disorder; and age-related cognitive decline; and wherein the psychiatric disorder is selected from the group consisting of schizophrenia, for example of the paranoid, disorganized, catatonic, undifferentiated, or residual type; schizophreniform disorder; schizoaffective disorder, for example of the delusional type or the
30 depressive type; delusional disorder; substance-induced psychotic disorder, for example psychosis induced by alcohol, amphetamine, cannabis, cocaine, hallucinogens, inhalants, opioids, or phencyclidine; personality disorder of the

paranoid type; and personality disorder of the schizoid type; and wherein the drug addiction is an alcohol, amphetamine, cocaine, or opiate addiction.

The compounds of formula I or pharmaceutically acceptable salts thereof may
5 be used in combination with one or more other drugs in the treatment of
diseases or conditions for which the compounds of the present invention have
utility, where the combination of the drugs together are safer or more effective
than either drug alone. Additionally, the compounds of the present invention
may be used in combination with one or more other drugs that treat, prevent,
10 control, ameliorate, or reduce the risk of side effects or toxicity of the
compounds of the present invention. Such other drugs may be administered,
by a route and in an amount commonly used therefore, contemporaneously or
sequentially with the compounds of the present invention. Accordingly, the
pharmaceutical compositions of the present invention include those that
15 contain one or more other active ingredients, in addition to the compounds of
the present invention. The combinations may be administered as part of a unit
dosage form combination product, or as a kit or treatment protocol wherein
one or more additional drugs are administered in separate dosage forms as
part of a treatment regimen.

20

The present invention provides a method of treating a mammal, including a
human, suffering from a neurodegenerative disorder selected from a cognition
disorder or movement disorder, which method comprises administering to the
subject a therapeutically effective amount of a compound of formula I.

25

This invention further provides a method of treating a neurodegenerative
disorder or condition in a mammal, including a human, which method
comprises administering to said mammal an amount of a compound of
formula I effective in inhibiting PDE10.

30

This invention also provides a method of treating a subject suffering from a
psychiatric disorder, which method comprises administering to the subject a
therapeutically effective amount of a compound of formula I. Examples of
psychiatric disorders that can be treated according to the present invention

include, but are not limited to, schizophrenia, for example of the paranoid, disorganized, catatonic, undifferentiated, or residual type; schizophreniform disorder; schizoaffective disorder, for example of the delusional type or the depressive type; delusional disorder; substance-induced psychotic disorder, 5 for example psychosis induced by alcohol, amphetamine, cannabis, cocaine, hallucinogens, inhalants, opioids, or phencyclidine; personality disorder of the paranoid type; and personality disorder of the schizoid type; and the anxiety disorder is selected from panic disorder; agoraphobia; a specific phobia; social phobia; obsessive-compulsive disorder; post-traumatic stress disorder; 10 acute stress disorder; and generalized anxiety disorder.

It has been found that the compounds of formula I or pharmaceutically acceptable salts thereof may advantageously be administered in combination with at least one neuroleptic agent (which may be a typical or an atypical 15 antipsychotic agent) to provide improved treatment of psychiatric disorders such as schizophrenia. The combinations, uses and methods of treatment of the invention may also provide advantages in treatment of patients who fail to respond adequately or who are resistant to other known treatments.

20 The present invention thus provides a method of treating a mammal suffering from a psychiatric disorder, such as schizophrenia, which method comprises administering to the mammal a therapeutically effective amount of a compound of formula I, either alone or as combination therapy together with at least one neuroleptic agent.

25

The term "neuroleptic agent" as used herein refers to drugs, which have the effect on cognition and behaviour of antipsychotic agent drugs that reduce confusion, delusions, hallucinations, and psychomotor agitation in patients with psychoses. Also known as major tranquilizers and antipsychotic drugs, 30 neuroleptic agents include, but are not limited to: typical antipsychotic drugs, including phenothiazines, further divided into the aliphatics, piperidines, and piperazines, thioxanthenes (e.g., cisordinol), butyrophenones (e.g., haloperidol), dibenzoxazepines (e.g., loxapine), dihydroindolones (e.g., molindone), diphenylbutylpiperidines (e.g., pimozide), and atypical

antipsychotic drugs, including benzisoxazoles (e.g., risperidone), sertindole, olanzapine, quetiapine, osanetant and ziprasidone.

Particularly preferred neuroleptic agents for use in the invention are
5 sertindole, olanzapine, risperidone, quetiapine, aripiprazole, haloperidol, clozapine, ziprasidone and osanetant.

The present invention further provides a method of treating a subject suffering from a cognition disorder, which method comprises administering to the
10 subject a therapeutically effective amount of a compound of formula I. Examples of cognition disorders that can be treated according to the present invention include, but are not limited to, Alzheimer's disease, multi-infarct dementia, alcoholic dementia or other drug-related dementia, dementia associated with intracranial tumors or cerebral trauma, dementia associated
15 with Huntington's disease or Parkinson's disease, or AIDS-related dementia; delirium; amnesic disorder; post-traumatic stress disorder; mental retardation; a learning disorder, for example reading disorder, mathematics disorder, or a disorder of written expression; attention-deficit/hyperactivity disorder; and age-related cognitive decline.

20

This invention also provides a method of treating a movement disorder, which method comprises administering to the subject a therapeutically effective amount of a compound of formula I. Examples of movement disorders that can be treated according to the present invention include, but are not limited
25 to, Huntington's disease and dyskinesia associated with dopamine agonist therapy. This invention further provides a method of treating a movement disorder selected from Parkinson's disease and restless leg syndrome, which comprises administering to the subject a therapeutically effective amount of a compound of formula I.

30

This invention also provides a method of treating a mood disorder, which method comprises administering to the subject a therapeutically effective amount of a compound of formula I. Examples of mood disorders and mood episodes that can be treated according to the present invention include, but

are not limited to, major depressive episode of the mild, moderate or severe type, a manic or mixed mood episode, a hypomanic mood episode; a depressive episode with a typical features; a depressive episode with melancholic features; a depressive episode with catatonic features; a mood
5 episode with postpartum onset; post-stroke depression; major depressive disorder; dysthymic disorder; minor depressive disorder; premenstrual dysphoric disorder; post-psychotic depressive disorder of schizophrenia; a major depressive disorder superimposed on a psychotic disorder such as delusional disorder or schizophrenia; a bipolar disorder, for example bipolar I
10 disorder, bipolar II disorder, and cyclothymic disorder. It is understood that a mood disorder is a psychiatric disorder.

This invention further provides a method of treating a drug addiction, for example an alcohol, amphetamine, cocaine, or opiate addiction, in a mammal,
15 including a human, which method comprises administering to said mammal an amount of a compound of formula I effective in treating drug addiction.

This invention also provides a method of treating a drug addiction, for example an alcohol, amphetamine, cocaine, or opiate addiction, in a mammal,
20 including a human, which method comprises administering to said mammal an amount of a compound of formula I effective in inhibiting PDE10.

The term "drug addiction", as used herein, means an abnormal desire for a drug and is generally characterized by motivational disturbances such a
25 compulsion to take the desired drug and episodes of intense drug craving.

Drug addiction is widely considered a pathological state. The disorder of addiction involves the progression of acute drug use to the development of drug-seeking behavior, the vulnerability to relapse, and the decreased, slowed
30 ability to respond to naturally rewarding stimuli. For example, The Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV) has categorized three stages of addiction: preoccupation/anticipation, binge/intoxication, and withdrawal/negative affect. These stages are characterized, respectively, everywhere by constant cravings and

preoccupation with obtaining the substance; using more of the substance than necessary to experience the intoxicating effects; and experiencing tolerance, withdrawal symptoms, and decreased motivation for normal life activities.

- 5 This invention further provides a method of treating a disorder comprising as a symptom a deficiency in attention and/or cognition in a mammal, including a human, which method comprises administering to said mammal an amount of a compound of formula I effective in treating said disorder.
- 10 Other disorders that can be treated according to the present invention are obsessive/compulsive disorders, Tourette's syndrome and other tic disorders.

As used herein, and unless otherwise indicated, a "neurodegenerative disorder or condition" refers to a disorder or condition that is caused by the
15 dysfunction and/or death of neurons in the central nervous system. The treatment of these disorders and conditions can be facilitated by administration of an agent which prevents the dysfunction or death of neurons at risk in these disorders or conditions and/or enhances the function of damaged or healthy neurons in such a way as to compensate for the loss of
20 function caused by the dysfunction or death of at-risk neurons. The term "neurotrophic agent" as used herein refers to a substance or agent that has some or all of these properties.

Examples of neurodegenerative disorders and conditions that can be treated
25 according to the present invention include, but are not limited to, Parkinson's disease; Huntington's disease; dementia, for example Alzheimer's disease, multi-infarct dementia, AIDS-related dementia, and Fronto temporal Dementia; neurodegeneration associated with cerebral trauma; neurodegeneration associated with stroke, neurodegeneration associated with cerebral infarct;
30 hypoglycemia-induced neurodegeneration; neurodegeneration associated with epileptic seizure; neurodegeneration associated with neurotoxin poisoning; and multi-system atrophy.

In one embodiment of the present invention, the neurodegenerative disorder or condition involves neurodegeneration of striatal medium spiny neurons in a mammal, including a human.

- 5 In a further embodiment of the present invention, the neurodegenerative disorder or condition is Huntington's disease.

In another embodiment, the invention provides a method of treating a subject to reduce body fat or body weight, or to treat non-insuline demanding diabetes mellitus (NIDDM), metabolic syndrome, or glucose intolerance, comprising
10 administering to a subject in need thereof a therapeutically effective amount of a compound of formula I. In preferred embodiments, the subject is human, the subject is overweight or obese and the antagonist is administered orally. In another preferred embodiment, the method further comprising administering a
15 second therapeutic agent to the subject, preferably an anti-obesity agent, e.g., rimonabant, orlistat, sibutramine, bromocriptine, ephedrine, leptin, pseudoephedrine, or peptide YY3-36, or analogs thereof.

The term "metabolic syndrome" as used herein refers to a constellation of
20 conditions that place people at high risk for coronary artery disease. These conditions include type 2 diabetes, obesity, high blood pressure, and a poor lipid profile with elevated LDL ("bad") cholesterol, low HDL ("good") cholesterol, and elevated triglycerides. All of these conditions are associated with high blood insulin levels. The fundamental defect in the metabolic
25 syndrome is insulin resistance in both adipose tissue and muscle.

30

Headings and sub-headings are used herein for convenience only, and should not be construed as limiting the invention in any way.

The use of any and all examples, or exemplary language (including “for instance”, “for example”, “e.g.”, and “as such”) in the present specification is intended merely to better illuminate the invention, and does not pose a
5 limitation on the scope of invention unless otherwise indicated.

The citation and incorporation of patent documents herein is done for convenience only, and does not reflect any view of the validity, patentability and/or enforceability of such patent documents.


10

The present invention includes all modifications and equivalents of the subject-matter recited in the claims appended hereto, as permitted by applicable law.

15 Experimental Section

Preparation of the compounds of the invention

20




25

Compounds of the general formula I of the invention may be prepared as described in the following reaction schemes. Unless otherwise indicated, in the reaction schemes and discussion that follow, **HET-1**, R₁-R₉, -L-, Z and Y are as defined above.

30

Compounds of formula I, wherein -L- is -S-CH₂-, can be prepared by the coupling of a nucleophile of formula **V** or **Va** with an electrophile of formula **VI**, where X is a leaving group, e.g. Cl, Br, I, methanesulfonyl, 4-toluenesulfonyl, as shown in scheme 1. In the reaction between **Va** and **VI**, alkylation of the

sulfur atom of **Va** with **VI** and ring closure to form the fused bicyclic triazole ring both take place under the same reaction conditions in a one-pot procedure.



5

Scheme 1.

This reaction is typically carried out in a solvent such as 1-propanol, toluene, DMF, or acetonitrile, optionally in the presence of a carbonate base such as potassium carbonate or a tertiary amine base such as triethylamine or diisopropylethylamine (DIPEA), at a temperature ranging from about 0° C to about 200° C, optionally under pressure in a closed vessel. Other suitable solvents include benzene, chloroform, dioxane, ethyl acetate, 2-propanol and xylene. Alternatively, solvent mixtures such as toluene/2-propanol can be used.


Compounds of formula **V** are either commercially available or can be prepared as described in the literature, see for example Brown *et al. Aust. J. Chem.* **1978**, 31, 397-404; Yutilov *et al. Khim. Geter. Soedin.* **1988**, 799-804; Wilde *et al. Bioorg. Med. Chem. Lett.* **1995**, 5, 167-172; Kidwai *et al. J. Korean Chem. Soc.* **2005**, 49, 288-291. Compounds of formula **Va** can be prepared as described in WO 96/01826 from the corresponding 1,2-diaminopyridines by

20

reaction with thiocarbonyldiimidazole in a suitable solvent, such as chloroform, at a suitable temperature, such as room temperature or +40 °C. The requisite 1,2-diaminopyridines are readily available from the corresponding commercially available 2-aminopyridines by reaction with a suitable *N*-amination reagent, such as *O*-(mesitylsulfonyl)hydroxylamine, in a suitable solvent, such as chloroform, at a suitable temperature, such as 0 °C or room temperature, see WO 96/01826.

2-Halomethyl-4-(aryl)-1H-triazoles of formula **VI** can be prepared by halogenation of the corresponding 2-hydroxymethyl-4-(aryl)-1H-triazoles using a suitable reagent, e.g. thionyl chloride, phosphorous trichloride, or phosphorous tribromide, optionally using a suitable solvent such as dichloromethane, using methods well known to chemists skilled in the art. The requisite 2-hydroxymethyl-4-(aryl)-1H-imidazoles can be prepared by methods known in the art (see for example Browne, E. J.; *Australian Journal of Chemistry* **1971**, 24(2), 393-403.; Browne, E. J.; Nunn, E. E.; Polya, John B. *Journal of the Chemical Society [Section] C: Organic* **1970**, (11), 1515-18. Becker, Heinz G. O.; Goermar, G.; Timpe, Hans J. *Journal für Praktische Chemie (Leipzig)* **1970**, 312(4), 610-21. Behringer, Hans; Ramert, Reiner. *Justus Liebigs Annalen der Chemie* **1975**, (7-8), 1264-71. Moderhack, Dietrich, *Liebigs Annalen der Chemie* **1984**, (1), 48-65.).

Compounds of formula **I**, wherein -L- is -CH₂-S -, can be prepared by the coupling of a nucleophile of formula **XII** with an electrophile of formula **VIII** as shown in scheme 2.




Scheme 2.

5 This reaction is typically carried out in a solvent such as 1-propanol, toluene, DMF, or acetonitrile, optionally in the presence of a carbonate base such as potassium carbonate or a tertiary amine base such as triethylamine or diisopropylethylamine (DIPEA), at a temperature ranging from about 0° C to about 200° C, optionally under pressure in a closed vessel. Other suitable
10 solvents include benzene, chloroform, dioxane, ethyl acetate, 2-propanol and xylene. Alternatively, solvent mixtures such as toluene/2-propanol can be used.

Some electrophiles of formula **VIII** are commercially available, and many others are known in the art, see for example JP 59176277. The electrophile **VIII**, where X is a leaving group, e.g. Cl, Br, I, methanesulfonyl, 4-toluenesulfonyl, can also be prepared by conversion of the primary alcohol of compounds of formula **VII** to said leaving group by methods known to chemists skilled in the art. Said methods can for example be selected from reacting compounds of formula **VII** with thionyl chloride, phosphorous trichloride, phosphorous tribromide, methanesulfonyl chloride, or 4-toluenesulfonyl chloride optionally in the presence of a suitable solvent, such as dichloromethane or 1,2-dichloroethane, and optionally in the presence of a base, such as triethylamine, diisopropylethylamine, or pyridine. Alternatively,

- electrophiles of formula **VIII** can be prepared by reacting commercially available aromatic amines of formula **IX** with 1,3-dihaloacetones of formula **XI**, e.g. 1,3-dichloroacetone, in a suitable solvent, such as 1,2-dimethoxyethane or ethanol, at a suitable temperature, such as room temperature or reflux.
- 5 Some electrophiles of formula **VII** are commercially available, and many others are known in the art, see for example Tsuchiya, T.; Sashida, H. *J. Chem. Soc., Chem. Commun.* **1980**, 1109-1110; Tsuchiya, T.; Sashida, H.; Konoshita, A. *Chem. Pharm. Bull.* **1983**, 31, 4568-4572. Alternatively, alcohols of formula **VII** can be prepared by reacting commercially available aromatic
- 10 amines of formula **IX** with a suitable *N*-amination reagent, such as *O*-(mesitylsulfonyl)hydroxylamine, in a suitable solvent, such as chloroform, at a suitable temperature, such as 0 °C or room temperature, see WO 96/01826, to yield compounds of formula **X**. Said compounds of formula **X** can be converted into compounds of formula **VII** by reaction with methyl glycolate
- 15 followed by reduction of the methyl ester to the requisite alcohol using a suitable reducing agent such as lithium aluminium hydride in a suitable solvent such as diethyl ether or tetrahydrofuran using methods known to chemists skilled in the art.
- 20 Compounds of formula **XII** are either commercially available or can be prepared as described in the literature, see e.g. Hoggarth, Eric. *Journal of the Chemical Society* **1949**, 1160-3. Losse, Gunter; Hessler, Willi; Barth, Alfred. *Chemische Berichte* **1958**, 91, 150-7. Potts, K. T.; Burton, H. R.; Roy, S. K. *Journal of Organic Chemistry* **1966**, 31(1), 265-73. ale, Harry L.; Piala, Joseph
- 25 J.; *Journal of Medicinal Chemistry* **1966**, 9(1), 42-6. Lalezari, I.; Sharghi, N. *Journal of Heterocyclic Chemistry* **1966**, 3(3), 336-7. Durant, Graham J. *Journal of the Chemical Society [Section] C: Organic* **1967**, (1), 92-4. US-3962237. Barnikow, Guenter; Ebeling, Horst. *Zeitschrift für Chemie* **1980**, 20(2), 55-6. WO-2009087218. US-2003162812. WO-2000012489. Baxter,
- 30 Andrew; et. al. *Bioorganic & Medicinal Chemistry Letters* **2003**, 13(16), 2625-2628. Dhiman, A. M.; Wadodkar, K. N.; Patil, S. D. *Indian Journal of Chemistry, Section B: Organic Chemistry Including Medicinal Chemistry* **2001**, 40B(7), 636-639.

Compounds of formula I, wherein R1 is not hydrogen, can be prepared by the alkylation of a compounds of formula I, wherein R1 is hydrogen, with an alkyl halide of formula XIII as shown in scheme 3.



5

Scheme 3.


This reaction is typically carried out in a suitable solvent, such as dimethylformamide, dimethylacetamide, or acetonitrile, in the presence of a suitable base such as a carbonate base, e.g. potassium carbonate, or a tertiary amine base, e.g. triethylamine or diisopropylethylamine (DIPEA), at a temperature ranging from about 0° C to about 100° C.

Compounds of formula I, wherein -L- is -CH=CH- or -CH₂-CH₂- can be prepared by the reaction sequence shown in scheme 4.

15

Compounds where -L- is $-\text{C}\equiv\text{C}-$ can be prepared by the reaction sequence shown in scheme 5


35



Scheme 4.

- 5 Specifically, compounds of formula **I**, wherein -L- is -CH₂-CH₂- can be prepared by reduction of an alkene of formula **I**, wherein -L- is -CH=CH-, by hydrogenation using a transition metal catalyst, such as palladium metal, together with a hydrogen source, such as hydrogen gas, ammonium hydrogen carbonate, or cyclohexadiene. Said alkenes of formula **I**, wherein -L- is -CH=CH- can be prepared by the Wittig reaction between a phosphonium salt of formula **XIV** and an aldehyde of formula **XV** in a suitable solvent, such as tetrahydrofuran, in the presence of a suitable base, such as 1,8-diazabicyclo[5.4.0]undec-7-ene. Phosphonium salt of formula **XIV** are readily available by reaction of compounds of formula **VIII** (see scheme 2 above) with triphenylphosphine by methods known to chemists skilled in the art. Aldehydes of formula **XV** are readily available by oxidation of alcohols of formula **VII** (see scheme 2 above) by methods known to chemists skilled in the art, e.g. by reacting alcohols of formula **VII** with a suitable oxidizing agent, such as Dess-Martin periodinane, in a suitable solvent, such as dichloromethane or 1,2-dichloroethane.

Compounds of formula **I**, wherein L is a triple bond (etynylene), can be prepared by a coupling reaction between an triazolyl alkyne of formula **XVII** with an heteroaryl halide of formula **XVI** or by the reverse coupling between an heteroaryl alkyne of formula **XVIII** with an triazolyl halide of formula **XIX** as shown in scheme 5.



Scheme 5.

This reaction is typically carried out in a suitable solvent, such as tetrahydrofuran and performed by mixing the heteroaryl halide with the heteroaryl alkyne together with a suitable catalyst e.g. Copper(I) iodide with a phosphine ligand e.g. 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride dichloromethane complex and an organic base like triethylamine and then heating the reaction in a sealed vial at 120 °C for 15 minutes (MicroWave).

The invention disclosed herein is further illustrated by the following non-limiting examples.

General Methods

Analytical LC-MS data were obtained using one of the following methods.

Method A:

A PE Sciex® API 150EX instrument equipped with atmospheric pressure photo ionisation and a Shimadzu™ LC-8A/SLC-10A LC system was used. Column: 4.6 x 30 mm Waters Symmetry C18 column with 3.5 µm particle size; Column temperature: 60 °C; Solvent system: A = water/trifluoroacetic acid (100:0.05) and B = water/ acetonitrile/trifluoroacetic acid (5:95:0.035); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 2.4 minutes and with a flow rate of 3.3 mL/min.

Method B:

A PE Sciex® API 300 instrument equipped with atmospheric pressure photo ionisation and a Waters UPLC system was used. Column: Acquity® UPLC BEH C18 1.7 µm, 2.1 x 50 mm (Waters); Column temperature: 60 °C; Solvent system: A = water/trifluoroacetic acid (100:0.05) and B = water/acetonitrile/trifluoroacetic acid (5:95:0.035); Method: Linear gradient elution with A:B = 90:10 to 0:100 in 1.0 minutes and with a flow rate of 1.2 mL/min.

Method C:

A PE Sciex® API 150EX instrument equipped with atmospheric pressure photo ionisation and a Shimadzu™ LC-8A/SLC-10A LC system was used. Column: 4.6 x 30 mm Waters Symmetry C₁₈ column with 3.5 µm particle size; Column temperature: 60 °C; Solvent system: A= water/trifluoroacetic acid (99.95:0.05) and B = methanol/trifluoroacetic acid (99.965:0.035); Method: Linear gradient elution with A:B = 83:17 to 0:100 in 2.4 minutes and with a flow rate of 3.0 mL/min.

Preparative LC-MS-purification was performed on a PE Sciex® API 150EX instrument with atmospheric pressure chemical ionization. Column: 50 X 20 mm YMC ODS-A with 5 µm particle size; Method: Linear gradient elution with A:B = 80:20 to 0:100 in 7 minutes and with a flow rate of 22.7 mL/minute. Fraction collection was performed by split-flow MS detection.

¹H NMR spectra were recorded at 500.13 MHz on a Bruker Avance™ AV500 instrument or at 250.13 MHz on a Bruker Avance™ DPX250 instrument. TMS


was used as internal reference standard. Chemical shift values are expressed in ppm. The following abbreviations are used for multiplicity of NMR signals: s = singlet, d = doublet, t = triplet, q = quartet, qui = quintet, h = heptet, dd = double doublet, dt = double triplet, dq = double quartet, tt = triplet of triplets, m = multiplet, br s = broad singlet and br = broad signal.

Abbreviations are in accordance with to the ACS Style Guide: "The ACS Styleguide – A manual for authors and editors" Janet S. Dodd, Ed. 1997, ISBN: 0841234620

General: p-Toluene-sulfonyl hydrazide (98%) was from Avocado.

Preparation of intermediates

2-Methyl-5-phenyl-2H-1,2,4-triazole-3-carbaldehyde



N-Methylhydrazine (1.46 mL, 27.5 mmol) was added to a stirred solution of 2-Phenyl-oxazol-4-one (4.03 g, 25.0 mmol) (prepared as described in Yushiyuki *et al. Synthesis*. 2004, 1359-1363) in abs Ethanol (12 mL) at rt under Ar. Exothermic reaction. The reaction mixture was stirred at rt 1h. The solvent was evaporated off to produce 4.64 g of a yellow/white solid of crude 2-Methyl-5-phenyl-2H-1,2,4-triazol-3-yl)-methanol.

Dess-Martin periodinane (11.3 g, 26.7 mmol) was added in one portion to (2-Methyl-5-phenyl-2H-1,2,4-triazol-3-yl)-methanol (4.60 g, 24.3 mmol) dissolved in Methylene chloride (72 mL) at 0 °C under Ar. After 2h, the reaction was diluted with DCE (100 ml) and extracted with sat NaHCO₃ (100 ml). The aq phase was discarded and the organic phase was washed with brine and was

rotovaped. The crude product was purified by silica gel chromatography (Eluent: 20-50% EtOAc in heptane). Yield: 3.20 g of the title compound as a solid. H-NMR: (CDCl₃) δ 10.08 (s, 1H), 8.13 (m, 2H), 7.50-7.42 (m, 3H), 4.26 (s, 3H).

5 The Following compounds were prepared analogously:

2-Ethyl-5-phenyl-2H-1,2,4-triazole-3-carbaldehyde from Ethyl-hydrazine and 2-Phenyl-oxazol-4-one. H-NMR: (DMSO-d₆) δ 10.01 (s, 1H), 8.07 (d, 2H), 7.50 (m, 3H), 4.59 (q, 2H), 1.42 (t, 3H).

10 **5-Phenyl-2-propyl-2H-1,2,4-triazole-3-carbaldehyde**, from Propyl-hydrazine and 2-Phenyl-oxazol-4-one. H-NMR: (CDCl₃) δ 10.07 (s, 1H), 8.17 (d, 2H), 7.48 (m, 3H), 4.59 (m, 2H), 1.95 (m, 2H) 1.00 (t, 3H).


2-Isopropyl-5-phenyl-2H-1,2,4-triazole-3-carbaldehyde, from Isopropyl-hydrazine and 2-Phenyl-oxazol-4-one. H-NMR: (DMSO-d₆) δ 10.02 (s, 1H), 8.08 (d, 2H), 7.50 (m, 3H), 5.37 (m, 1H), 1.51 (d, 6H).

15 **5-Phenyl-2-(2,2,2-trifluoro-ethyl)-2H-1,2,4-triazole-3-carbaldehyde**, from (2,2,2-Trifluoro-ethyl)-hydrazine and 2-Phenyl-oxazol-4-one. H-NMR: (CDCl₃) δ 10.07 (s, 1H), 8.18 (d, 2H), 7.50 (m, 3H), 5.28 (m, 2H).

20 **3-(5-Formyl-3-phenyl-1,2,4-triazol-1-yl)-propionitrile**, from 3-Hydrazino-propionitrile and 2-Phenyl-oxazol-4-one. H-NMR: (DMSO-d₆) δ 10.03 (s, 1H), 8.08 (d, 2H), 7.52 (m, 3H), 4.86 (t, 2H), 3.20 (t, 2H).

(5-Formyl-3-phenyl-1,2,4-triazol-1-yl)-acetic acid ethyl ester, from Hydrazino-acetic acid ethyl ester and 2-Phenyl-oxazol-4-one. H-NMR: (CDCl₃) δ 10.05 (s, 1H), 8.15 (d, 2H), 7.50 (m, 3H), 5.38 (s, 2H), 4.28 (m, 2H), 1.32 (t, 3H).

25 **Example 2 2-(2-Methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazole-3-carbaldehyde**



2-HYDROXYETHYLHYDRAZINE (1.60 g, 21.0 mmol) was added to a stirred
 5 solution of 2-Phenyl-oxazol-4-one (3.08 g, 19.1 mmol) in Ethanol (9.6 mL) at rt
 under Ar. Exothermic reaction. The reaction mixture was stirred at rt ON. The
 solvent was evaporated off. Yield: 4.24 g of **2-(5-Hydroxymethyl-3-phenyl-
 1,2,4-triazol-1-yl)-ethanol**.

Sodium hydride (60% in mineral oil) (60:40, Sodium hydride:Mineral Oil, 1.36
 10 g) was added in portions, to a stirred solution of 2-(5-Hydroxymethyl-3-phenyl-
 1,2,4-triazol-1-yl)-ethanol (2.92 g, 13.3 mmol) in N,N-Dimethylformamide (100
 mL) at rt. The reaction was stirred at rt 5 min under Ar. Benzyl bromide (1.35
 mL, 11.3 mmol) was added and the solution was stirred at rt under Ar 1 h. The
 reaction mixture was quenched by adding conc HCl (1 ml) and then EtOAc
 15 (200 ml) and brine (100 ml). The phases were separated and the org phase
 was rotovaped and the crude product was purified by silica gel chromatography
 (Eluent: 20-50% EtOAc in heptane). Yield: 1.39 g of 2-(5-Benzyloxymethyl-3-
 phenyl-[1,2,4]triazol-1-yl)-ethanol as an oil.

Sodium hydride (60% in mineral oil) (60:40, Sodium hydride:Mineral Oil, 197.7
 20 mg) was added in portions, to a stirred solution of 2-(5-Benzyloxymethyl-3-
 phenyl-1,2,4-triazol-1-yl)-ethanol (1.39 g, 4.49 mmol) in N,N-
 Dimethylformamide (45 mL) at rt. The reaction was stirred at rt 5 min under
 Ar. Methyl iodide (0.3077 mL, 4.942 mmol) was added and the solution was
 stirred at rt under Ar 1 h. More NaH (200 mg) and MeI (0.1 ml) was added.
 25 This addition was repeated after aprox 1 h.

EtOAc (100 ml) and brine (100 ml) was added and the phases were
 separated. The org phase was washed with more brine, dried (Na₂SO₄) and


the solvent evaporated off. The crude product was purified by silica gel chromatography (20-100% EtOAc in heptane). Yield: 1.23 g of **5-Benzyloxymethyl-1-(2-methoxy-ethyl)-3-phenyl-1H-1,2,4-triazole** as an oil.

- 5 Pd/C (10%) (9:1, carbon black:Palladium, 250 mg) was added, to a solution of 5-Benzyloxymethyl-1-(2-methoxy-ethyl)-3-phenyl-1H-1,2,4-triazole (1.23 g, 3.80 mmol) in Methanol (72 mL) and TFA (2 ml) at rt. The reaction was hydrogenated at 3 bar using a Parr shaker ON. More catalyst (200 mg) was added and the hydrogenation was cont ON. The catalyst was filtered off and
- 10 the solvent was evaporated off. The crude product was purified by silica gel chromatography (25-100% EtOAc in heptane). Yield: 0.84 g of **[2-(2-Methoxy-ethyl)-5-phenyl-2H-1,2,4-triazol-3-yl]-methanol** as an oil.

- Dess-Martin periodinane (1530 mg, 3.60 mmol) was added to a stirred suspension of [2-(2-Methoxy-ethyl)-5-phenyl-2H-1,2,4-triazol-3-yl]-methanol
- 15 (840 mg, 3.6 mmol) dissolved in Methylene chloride (36.0 mL) at rt under Ar. The solution was stirred ON. Some solid was filtered off and discarded. Sat NaHCO₃ (35 ml) and more DCM (35 ml) were added to the DCM solution. The solvent was evaporated off and the crude product was purified by silica gel chromatography (Eluent: 50% EtOAc in heptane). Yield: 0.76 g of the title
- 20 compound as an oil. H-NMR: (CDCl₃) δ 10.04 (s, 1H), 8.13 (m, 2H), 7.45 (m, 3H), 4.82 (t, 2H), 3.83 (t, 2H), 3.32 (s, 3H).

Preparation of other intermediates

2-Chloromethyl-5,7-dimethyl-[1,2,4]triazolo[1,5-a]pyrimidine



To a solution of 4,6-Dimethyl-pyrimidin-2-ylamine (25 g, 200 mmol) in 400 mL
 5 of CH_2Cl_2 was added dropwise a solution of hydroxylamine-2,4,6-Trimethyl-
 benzenesulfonate (105 g, 488 mmol) in 300 mL of CH_2Cl_2 at 0°C , and the
 mixture was stirred at 0°C for 1 hour and filtered. The solid collected was
 washed with CH_2Cl_2 (100 mL) to give 1-Amino-4,6-dimethyl-1H-pyrimidin-2-
 ylidene-ammonium 2,4,6-Trimethyl-benzenesulfonate (40 g, yield:62%).

10

A mixture of 1-Amino-4,6-dimethyl-1H-pyrimidin-2-ylidene-ammonium 2,4,6-
 Trimethyl-benzenesulfonate (40 g, 0.1 mol) and NaOH (10 g, 0.2 mol) in 500
 mL of EtOH was stirred at $50\sim 60^\circ\text{C}$ for 1 hour. After chloroacetic acid methyl
 ester (16.6 g, 0.15 mol) was added, the resultant mixture was stirred at reflux
 15 for 4 hours. After being concentrated under reduce pressure, the residue was
 diluted with water (1000 mL) and extracted with CH_2Cl_2 (300 mL $\times$ 3). The
 combined organic layers were washed with brine (200 mL), dried over
 Na_2SO_4 , filtered, and concentrated under vacuum. The residue was purified
 by column chromatography on silica gel (petroleum ether/EtOAc = 2/1) to give
 20 2 g of 2-Chloromethyl-5,7-dimethyl-[1,2,4]triazolo[1,5-a]pyrimidine in 9% yield.

¹H NMR (300 MHz, DMSO-*d*₆): δ 8.55 (s, 1H), 6.25 (s, 2H), 4.05 (s, 3H), 3.95 (s, 3H); LC-MS (MH⁺): *m/z* = 196.9, *t*_R (min, method A) = 0.52

The following intermediates were prepared analogously:

5

7-Chloro-2-chloromethyl-5,8-dimethyl-[1,2,4]triazolo[1,5-*c*]pyrimidine from 6-Chloro-2,5-dimethyl-pyrimidine-4-ylamine prepared as described by Henze *et al.* J. Org. Chem **1952**, 17, 1320-1327. 3.2% yield, LC-MS: *m/z* = 231.5 (MH⁺), *t*_R = 1.13 min, method C

10

2-Chloromethyl-5,8-dimethyl-[1,2,4]-triazolo[1,5-*a*]pyrazine from 2-amino-3,6-dimethylpyrazine. 60% yield, ¹H NMR (500 MHz, CDCl₃): δ 7.91 (s, 1H), 4.87 (s, 2H), 2.91 (s, 3H), 2.74 (s, 3H), LC-MS: *m/z* = 196.9 (MH⁺), *t*_R = 0.64 min, method A

15

2-Chloromethyl-5,8-dimethyl-[1,2,4]triazolo[1,5-*a*]pyridine from 6-Chloro-5-ethyl-2-methyl-pyrimidin-4-ylamine. 21 % yield, LC-MS: *m/z* = 245.0 (MH⁺), *t*_R = 0.72 min, method A

20

2-Chloromethyl-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-*a*]pyridine from 3-Methoxy-6-methyl-pyridin-2-ylamine. 64%, ¹H NMR (500 MHz, DMSO-*d*₆): δ 7.11-7.08 (d, 1H), 7.01-6.98 (d, 1H), 4.93 (s, 2H), 3.98 (s, 3H), 2.61 (s, 3H)

25

2-Chloromethyl-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-*a*]pyridine from 2-AMINO-6-METHYLPYRIDINE, LC-MS: *m/z* = 181.8 (MH⁺), *t*_R = 0.64 min, method A.


30

2-Chloromethyl-8-methyl-[1,2,4]triazolo[1,5-*a*]pyridine from 2-AMINO-3-METHYLPYRIDINE, LC-MS: *m/z* = xx (MH⁺), *t*_R = xx min, method xx. CHINA!

2-Chloromethyl-8-methoxy-[1,2,4]triazolo[1,5-*a*]pyridine from 2-AMINO-3-METHOXYPYRIDINE, LC-MS: *m/z* = 197.8 (MH⁺), *t*_R = 0.40 min, method B.

2-Chloromethyl-8-ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridine from 2-AMINO-3-ETHYL-6-METHYLPYRIDINE, LC-MS: $m/z = 209.8$ (MH^+), $t_R = 0.60$ min, method B.


5 **(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-ylmethyl)-triphenyl-phosphonium; chloride**




A solution of 2-chloromethyl-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine (1.351 g, 6.87 mmol) and triphenylphosphine (1.80 g, 6.87 mmol) in acetonitrile 150 mL was heated at reflux for 12 h. The solvents were removed *in vacuo* and the residue slurried in ether, filtered and dried to yield (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-ylmethyl)-triphenyl-phosphonium; chloride as an off white solid (2.412 g, 74.9%). LC-MS: $m/z = 423.2$ ($[M-Cl]^+$), $t_R = 0.86$ min, method A.

15

The following intermediates were prepared analogously:




20 (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: $m/z = 422.2$ (MH^+), $t_R = 1.02$ min, method A



(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: $m/z = 438.4$ (MH^+), $t_R = 0.96$ min, method A


5



(5-Methyl-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride


from 2-Chloromethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: $m/z =$

10 408.4 (MH^+), $t_R = 0.88$ min, method A




(8-Methyl-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-8-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS:

15 $m/z = 408.2$ (MH^+), $t_R = 0.59$ min, method B.




20 (5,7-Dimethyl-[1,2,4]triazolo[1,5-a]pyrimidin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-5,7-dimethyl-[1,2,4]triazolo[1,5-a]pyrimidine, LC-MS: $m/z = 423.3$ (MH^+), $t_R = 0.85$ min, method A



(8-Methoxy-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-8-methoxy-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: $m/z = 423.9$ (MH^+), $t_R = 0.55$ min, method B

5



(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-ylmethyl)-triphenyl-phosphonium chloride from 2-Chloromethyl-8-ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: $m/z = 423.9$ (MH^+), $t_R = 0.55$ min, method B


10

Preparation of the compounds of the invention

15

Example 1

5-Methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine



20

25

Triethylamine (0.7602 mL, 5.454 mmol) was added to a solution of [2-Chloromethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridine (495 mg, 2.73 mmol) in N,N-Dimethylformamide (12 mL, 150 mmol) and the solution was stirred approx 2 mins. 5-Phenyl-2H-1,2,4-triazole-3-thiol (532 mg, 3.00 mmol) was added to the mixture as a solid. The reaction mixture was subsequently heated at 60 °C


for 0.5 hours. The mixture was rotovaped. EtOAc (60 mL) and sat. bicarbonate (20 mL) were added to the residue and the phases were separated. The organic phase was washed with brine (20 mL). After drying (MgSO₄), the solvent was removed under reduced pressure. The crude product was purified by silica gel chromatography (Eluent: EtOAc:N-heptane 50-100%). This afforded 542 mg (61%) of the title compound as a white solid. LC-MS: m/z = 323.1 (MH⁺), t_R = 0.52 min, Method B. ¹H NMR (600 MHz, DMSO-d₆): δ 7.96 (d, 2H), 7.63 (m, 1H), 7.60 (t, 1H) 7.50 (m, 3H), 7.06 (d, 1H), 4.68 (s, 2H), 2.67 (s, 3H).

The following compound was prepared analogously:

8-Methoxy-5-methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine (**2**), LC-MS: m/z = 353.4 (MH⁺), t_R = 0.57 min, Method B.

Example 2

5-Methyl-2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine



Methyl iodide (142 mg, 1.00 mmol) dissolved in DMF (0.5 ml) was added to a stirred suspension of 5-Methyl-2-(5-phenyl-2H-1,2,4-triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine (322 mg, 1.00 mmol) and Potassium carbonate (276 mg, 2.0 mmol) in N,N-Dimethylformamide (3.5 mL) at rt, under Ar. The solution was stirred at 3 h and then quenched by adding Sat bicarbonate sol and extracting with EtOAc. The Organic phase was dried (MgSO₄) and was rotovaped. The crude product was purified by HPLC. This afforded 10 mg of the title compound. LC-MS: m/z = 337.1 (MH⁺), t_R = 0.63


min, Method B. ^1H NMR (600 MHz, DMSO-d_6): δ 7.96 (d, 2H), 7.63 (d, 1H), 7.60 (t, 1H) 7.50-7.40 (m, 3H), 7.07 (d, 1H), 4.68 (s, 2H), 2.66 (s, 3H).

The following compound was synthesized analogously:

5,8-Dimethyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-
 5 [1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 338.1 (MH^+), t_R = 1.38 minutes,
 Method C

Example 3

5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-
 10 [1,2,4]triazolo[1,5-a]pyrazine



15

1,8-Diazabicyclo[5.4.0]undec-7-ene (0.16 mL, 1.07 mmol) was added to a stirred suspension of (5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-ylmethyl)-triphenyl-phosphonium; chloride (0.490 g, 1.07 mmol) and 2-Methyl-5-phenyl-
 20 2H-1,2,4-triazole-3-carbaldehyde (0.200 g, 1.07 mmol) in Tetrahydrofuran (6.9 mL) in dry THF (8 mL) and the mixture was stirred at room temperature under an atmosphere of Argon overnight. DCM (50 mL) was added and the organic phase was extracted with water (2 x 50 mL), dried (MgSO_4) and the solvent was evaporated off.

25 The remanens was dissolved in DCM, and purified by silica gel chromatography (Eluent: EtOAc). Yield: 270 mg (76%) mg of the intermediate as a cis/trans mixture.

This material (270 mg, 0.81 mmol) was dissolved in DCM (10 mL) and MeOH
 30 (80 mL). The solution was filtered and was hydrogenated at rt on an H-cube through a Pd/C CatCart column at 1 bar hydrogen pressure. The solvent was

evaporated off, and the crude product was purified by silica gel chromatography (Eluent: EtOAc). Yield: 138 mg (51%) of the title compound as a solid. LC-MS: m/z = 334.5 (MH^+), t_R = 1.16 min, Method C. ^1H NMR (600 MHz, DMSO- d_6): δ 8.0-7.9 (m, 3H), 7.45-7.35 (m, 3H), 3.90 (s, 3H), 3.44 (t, 2H), 3.35 (t, 2H), 2.74 (s, 3H), 2.64 (s, 3H).

The following compounds were prepared analogously:

8-Methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine dihydrochloride, LC-MS: m/z = 318.9 (MH^+), t_R = 0.52 min, Method B.

5,7-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrimidine, LC-MS: m/z = 334.5 (MH^+), t_R = 1.05 min, Method C.

8-Methoxy-5-methyl-2-[2-(5-phenyl-2-propyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine dihydrochloride, LC-MS: m/z = 377.5 (MH^+), t_R = 0.66 min, Method B.

8-Methoxy-5-methyl-2-{2-[5-phenyl-2-(2,2,2-trifluoro-ethyl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 416.4 (MH^+), t_R = 0.71 min, Method B.

8-Methoxy-5-methyl-2-{2-[5-phenyl-2-(2,2,2-trifluoro-ethyl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 319.2 (MH^+), t_R = 1.14 min, Method C.

8-Methoxy-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 335.3 (MH^+), t_R = 1.09 min, Method C.

5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-acetic acid ethyl ester, LC-MS: m/z = 420.6 (MH^+), t_R = 0.65 min, Method B.

5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 333.2 (MH^+), t_R = 1.24 min, Method C.

5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-c]pyrimidine, LC-MS: m/z = 334.5 (MH^+), t_R = 1.18 min, Method C.

- 5,8-Dimethyl-2-[2-(2-methyl-5-thiophen-3-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 340,1 (MH⁺), t_R = 1,29 min, Method C.
- 2-[2-(5-Furan-2-yl-1-methyl-1H-[1,2,4]triazol-3-yl)-ethyl]-5,8-dimethyl-
5 [1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 324,1 (MH⁺), t_R = 1,17 min, Method C.
- 2-[(E)-2-(5-Furan-2-yl-1-methyl-1H-[1,2,4]triazol-3-yl)-vinyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 322,1 (MH⁺), t_R = 1,46 min, Method C.
- 10 5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-4-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 341,1 (MH⁺), t_R = 0,97 min, Method C.
- 5,8-Dimethyl-2-{2-[2-methyl-5-(5-methyl-thiazol-2-yl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 355,1 (MH⁺), t_R = 1,21 min,
15 Method C.
- 5,8-Dimethyl-2-{2-[2-methyl-5-(4-methyl-thiazol-2-yl)-2H-[1,2,4]triazol-3-yl]-ethyl}-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 355,1 (MH⁺), t_R = 1,18 min, Method C.
- 5,8-Dimethyl-2-[2-(2-methyl-5-oxazol-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-
20 [1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 325,1 (MH⁺), t_R = 0,87 min, Method C.
- 5,8-Dimethyl-2-[2-(2-methyl-5-thiophen-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 340,1 (MH⁺), t_R = 1,24 min, Method C.
- 25 5,8-Dimethyl-2-[2-(2-methyl-5-thiophen-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 340,1 (MH⁺), t_R = 1,26 min, Method C.
- 5,8-Dimethyl-2-[2-(2-methyl-5-pyrimidin-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 336,2 (MH⁺), t_R = 0,79 min,
30 Method C.
- 5,8-Dimethyl-2-[2-(2-methyl-5-pyridin-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 335,2 (MH⁺), t_R = 0,78 min, Method C.

5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-5-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 341,1 (MH⁺), t_R = 1.0 min, Method C.

5,8-Dimethyl-2-[2-(2-methyl-5-thiazol-2-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 341,1 (MH⁺), t_R = 1,01 min, Method C.

5,8-Dimethyl-2-[2-(2-methyl-5-pyridin-3-yl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 335,2 (MH⁺), t_R = 0,7 min, Method C.

10 2-[2-(5-Furan-2-yl-2-methyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 324,1 (MH⁺), t_R = 1,09 min, Method C.


2-[2-(5-Furan-2-yl-2-methyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 324,1 (MH⁺), t_R = 1,08 min, Method C.

5,8-Dimethyl-2-(5-phenyl-2H-[1,2,4]triazol-3-yl)sulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyrazine LC-MS: m/z = 338,1 (MH⁺), t_R = 1,38 min, Method C.

20 2-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol LC-MS: m/z = 364,2 (MH⁺), t_R = 1,3 min, Method C.

Example 4

8-Ethyl-2-(2-{5-[(Z)-1-eth-(E)-ylidene-penta-2,4-dienyl]-2-propyl-2H-[1,2,4]triazol-3-yl}-ethyl)-5-methyl-[1,2,4]triazolo[1,5-a]pyridine



1,8-Diazabicyclo[5.4.0]undec-7-ene (0.150 mL, 1.00 mmol) was added to a stirred suspension of 8-Ethyl-5-methyl-[1,2,4]triazolo [1,5-a]pyridin-2-

ylmethyl)-triphenyl phosphonium; chloride (472 mg, 1.00 mmol) and 5-Phenyl-2-propyl-2H-1,2,4-triazole-3-carbaldehyde (215 mg, 1.00 mmol) in dry THF (20 mL) and the mixture was stirred at room temperature under an atmosphere of Argon 2 days. The solvent was removed *in vac*. The remanens
5 was dissolved in DCM, and purified by silica gel chromatography (Eluent: 0-100% EtOAc in n-heptane). Yield: 290 mg of the intermediate (as a single isomer) as a white solid (78%). LC-MS: m/z = 373.5 (MH^+), t_R = 0.93 min, Method B.

10 This material (290 mg, 0.78 mmol) was dissolved in DMF (7.8 mL) and p-Toluenesulfonylhydrazide (430 mg, 2.3 mmol) was added and the reaction was stirred at 120 °C 8h under Ar. The solution was allowed to reach room temperature and stirred overnight. More p-Toluenesulfonyl hydrazide (0.08 g)
15 was added, and the reaction mixture was at 120 °C ON under Ar. DMF was evaporated and the residue was dissolved in EtOAc (50 ml) and extracted with sat NaHCO₃ (2 x 25 ml). The organic phase was washed with brine, dried (MgSO₄) and was rotovaped. The crude product was purified by silica gel chromatography (Eluent: 50-100% EtOAc in n-heptane). Yield: 140 mg (48%). of the title compound as a solid. LC-MS: m/z = 375.3 (MH^+), t_R = 1.68
20 min, Method C. ¹H NMR (600 MHz, CDCl₃): δ 8.08 (d, 2H), 7.40 (d, 2H), 7.34 (m, 1H), 7.17 (d, 1H), 6.70 (d, 1H), 4.12 (m, 2H), 3.55 (m, 2H), 3.38 (m, 2H), 3.00 (m, 2H), 2.68 (s, 3H), 1.88 (m, 2H), 1.34 (t, 3H), 0.93 (t, 3H).

The following compounds were prepared analogously:

25

8-Methoxy-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 349.5 (MH^+), t_R = 0.59 min, Method B.

8-Ethyl-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 347.1 (MH^+), t_R = 0.59 min, Method B.

8-Ethyl-2-[2-(2-isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 375.2 (MH^+), t_R = 1.67 min, Method C.

3-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile, LC-MS: m/z = 388.5 (MH^+), t_R = 1.18 min, Method C.

5 3-{5-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile, LC-MS: m/z = 386.6 (MH^+), t_R = 1.32 min, Method C.


2-[2-(2-Isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 377.4 (MH^+), t_R = 1.45 min, Method C.

3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile, LC-MS: m/z = 373.4 (MH^+), t_R = 0.56 min, Method B.

2-{2-[2-(2-Methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazol-3-yl]-ethyl}-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine, LC-MS: m/z = 378.6 (MH^+), t_R = 1.42 min, Method C. Mp = 141-143 C.

Example 5

20 **2-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol**



1M of Lithium tetrahydroaluminate in Tetrahydrofuran (5.0 mL) was added to a stirred solution of {5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-1,2,4-triazol-1-yl}-acetic acid ethyl ester (950 mg, 2.2 mmol) in Tetrahydrofuran (3 mL) at rt. The reaction is exothermic. The solution was stirred at rt under Ar 3 h. The solution was diluted with THF (10 ml) and


quenched by adding wet Na₂SO₄. The solution was filtered through dry Na₂SO₄, and was rotovaped. The crude product was purified by silica gel chromatography (Eluent: 0-30% MeOH in EtOAc). Yield: 710 mg (75%) of the title compound as a clear oil. LC-MS: m/z = 379.4 (MH⁺), t_R = 0.51 min, Method B.

The following compound was prepared analogously:

8-Methoxy-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine, LC-MS: m/z = 349.5 (MH⁺), t_R = 0.59 min, Method B.

Example 6

3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propylamin




3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-1,2,4-triazol-1-yl}-propionitrile (28 mg, 0.075 mmol) was dissolved in 2 M of Ammonia in Methanol (15 mL) and hydrogenated on an H-cube through a column of Raney-Nickel (50 bar hydrogen, rt). The solvent was evaporated off. The crude product was purified by silica gel chromatography (Eluent: 10% MeOH, 10% triethylamine, 80% EtOAc). Yield: 10 mg (35%) of the title compound as a clear oil. LC-MS: m/z = 377.3 (MH⁺), t_R = 0.81 min, Method C.

The following compound was prepared analogously:

3-{5-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propylamine, LC-MS: m/z = 390.2 (MH^+), t_R = 0.48 min, Method B.

5 Example 7

2-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol



10

Iodotrimethylsilane (92.0 μ L, 0.646 mmol) was added to a stirred solution of 2-{2-[2-(2-Methoxy-ethyl)-5-phenyl-2H-1,2,4-triazol-3-yl]-ethyl}-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine (61 mg, 0.16 mmol) in Chloroform (5 mL) under an atmosphere of Argon at rt. The solution was stirred at rt 4h. More Iodotrimethylsilane (184 μ L, 1.29 mmol) was added and the mixture was stirred at rt ON. MeOH (10 ml) and solid sodium sulfite (0.5 g) was added and the mixture was stirred 30 min, filtered and evaporated. The crude product was purified by silica gel chromatography (Eluent: 0-5% MeOH in EtOAc). Yield: 27 mg (46%) of the title compound as a solid. LC-MS: m/z = 364.5 (MH^+), t_R = 1.30 min, Method C.

20

Pharmacological Testing

PDE10A enzyme

Active PDE10A enzyme is prepared in a number of ways for use in PDE assays (Loughney, K. *et al. Gene* **1999**, 234, 109-117; Fujishige, K. *et al. Eur J Biochem.* **1999**, 266, 1118-1127 and Soderling, S. *et al. Proc. Natl. Acad. Sci.* **1999**, 96, 7071-7076). PDE10A can be expressed as full-length proteins or as truncated proteins, as long as they express the catalytic domain. PDE10A can be prepared in different cell types, for example insect cells or E.

25

coli. An example of a method to obtain catalytically active PDE10A is as follows: The catalytic domain of human PDE10A (amino acids 440-779 from the sequence with accession number NP 006652) is amplified from total human brain total RNA by standard RT-PCR and is cloned into the BamH1 and Xho1 sites of the pET28a vector (Novagen). Expression in coli is performed according to standard protocols. Briefly, the expression plasmids are transformed into the BL21 (DE3) E. coli strain, and 50 mL cultures inoculated with the cells allowed to grow to an OD600 of 0.4-0.6 before protein expression is induced with 0.5mM IPTG. Following induction, the cells are incubated overnight at room temperature, after which the cells are collected by centrifugation. Cells expressing PDE10A are resuspended in 12 mL (50 mM TRIS-HCl-pH8.0, 1 mM MgCl₂ and protease inhibitors). The cells are lysed by sonication, and after all cells are lysed, TritonX100 is added according to Novagen protocols. PDE10A is partially purified on Q sepharose and the most active fractions were pooled.

PDE10A inhibition assay

A PDE10A assay may for example, be performed as follows: The assay is performed in 60 uL samples containing a fixed amount of the relevant PDE enzyme (sufficient to convert 20-25% of the cyclic nucleotide substrate), a buffer (50 mM HEPES7.6; 10mM MgCl₂; 0.02% Tween™20), 0.1 mg/ml BSA, 225 pCi of ³H-labelled cyclic nucleotide substrate, tritium labeled cAMP to a final concentration of 5 nM and varying amounts of inhibitors. Reactions are initiated by addition of the cyclic nucleotide substrate, and reactions are allowed to proceed for one hr at room temperature before being terminated through mixing with 15 uL 8 mg/mL yttrium silicate SPA beads (Amersham™). The beads are allowed to settle for one hr in the dark before the plates are counted in a Wallac 1450 Microbeta counter. The measured signal can be converted to activity relative to an uninhibited control (100 %) and IC₅₀ values can be calculated using the Xlfit extension to EXCEL.

Phencyclidine (PCP) induced hyperactivity

Male mice (NMRI, Charles River) weighing 20-25g are used. Eight mice are used in each group receiving the test compound (5 mg/kg) plus PCP (2.3

mg/kg) including the parallel control groups receiving the vehicle of the test compound plus PCP or vehicle injections only. The injection volumen is 10 ml/kg. The experiment is made in normal light conditions in an undisturbed room. The test substance is injected per oss 60 min before injection of PCP, 5 which is administered subcutaneous.


Immediately after injection of PCP the mice are placed individually in special designed test cage (20 cm x 32 cm). The activity is measured by 5X8 infrared light sources and photocells spaced by 4 cm. The light beams cross the cage 10 1.8 cm above the bottom of the cage. Recording of a motility count requires interruption of adjacent light beams, thus avoiding counts induced by stationary movements of the mice.

Motility is recorded in 5 min intervals for a period of 1 hour. The drug effect is 15 calculated on the total counts during the 1 hour behavioral test period in the following manner:

The mean motility induced by vehicle treatment in the absence of PCP is used as baseline. The 100 per cent effect of PCP is accordingly calculated to be total motility counts minus baseline. The response in groups receiving test 20 compound is thus determined by the total motility counts minus baseline, expressed in per cent of the similar result recorded in the parallel PCP control group. The per cent responses are converted to per cent inhibition.


CLAIMS

1. A compound having the structure I



5

wherein **HET-1** is a heteroaromatic group of formula **II** containing from 2 to 4 nitrogen atoms:



- 10 wherein Y can be N or CH, Z can be N or C, and wherein **HET-1** may optionally be substituted with up to three substituents R2-R4 individually selected from hydrogen; C₁-C₆ alkyl; halogen; cyano; halo(C₁-C₆)alkyl; aryl; alkoxy; and hydroxy(C₁-C₆)alkyl, and wherein * denotes the attachment point,

- 15 **Q** is a is phenyl, optionally substituted with one to three substituents individually selected from H, C₁-C₆ alkoxy and halogen, or a monocyclic 5-membered or 6-membered heteroaromatic group containing 1 or 2 heteroatoms

- L- is a linker selected from -S-CH₂-, -CH₂-S-, -CH₂-CH₂-, -CH=CH-, and
20 -C≡C- ,

R₁ is selected from H; C₁-C₆ alkyl; C₁-C₆ alkyl(C₃-C₈)cycloalkyl*; hydroxy(C₁-C₆)alkyl; CH₂CN; CH₂C(O)NH₂; aryl(C₁-C₆)alkyl; and C₁-C₆ alkyl-heterocycloalkyl*, wherein * denotes the attachment point,

with the proviso that the compound is not 1H-Benzimidazole, 2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(2-pyrazinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; 1H-Benzimidazole, 2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)methyl]thio]-; 1H-Benzimidazole, 1-ethyl-5-(1-piperidinylsulfonyl)-2-[[[3-(2-thienyl)-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 6-methyl-2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(3-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 8-methyl-2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; Imidazo[1,2-a]pyridine, 6-chloro-2-[[[3-(2-thienyl)-1H-1,2,4-triazol-5-yl]thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(4-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 6-methyl-2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; 1H-Benzimidazole, 2-[[[3-(2-pyridinyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-; Imidazo[1,2-a]pyridine, 6-chloro-2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; 3H-Imidazo[4,5-b]pyridine, 2-[[[(3-phenyl-1H-1,2,4-triazol-5-yl)thio]methyl]-; or 1H-Benzimidazole, 2-[[[3-(2-furanyl)-1H-1,2,4-triazol-5-yl]methyl]thio]-;

and tautomers and pharmaceutically acceptable salts thereof, with the proviso that when the linker (L) is -CH₂-S-, then HET-1 is neither imidazo[1,2-a]pyridine nor imidazo[1,2-a]pyrimidine.


2. The compound of Claim 1 in which **HET-1** is an imidazo[1,2-a]pyrimidine moiety.
3. The compound of Claim 1 in which **HET-1** is a [1,2,4]triazolo[1,5-a]pyridine moiety.
4. The compound of Claim 1 in which **HET-1** is an imidazo[1,2-a]pyridine moiety.
5. The compound of Claim 1 in which **HET-1** is an imidazo[4,5-b]pyrimidine moiety.
6. The compound of Claim 1 in which **HET-1** is a pyrazolo[1,5-a]pyridine moiety.

7. The compound of Claim 1 in which **HET-1** is a [1,2,4]triazolo[1,5-a]pyrimidine moiety.
8. The compound of Claim 1 in which **HET-1** is a [1,2,4]triazolo[1,5-a]pyrazine moiety.
- 5 9. The compound of Claim 1 in which **HET-1** is a [1,2,4]triazolo[1,5-c]pyrimidine moiety.
- 10 10. The compound of any one of Claims 1 to 9 wherein Q is phenyl, thiophene, furan, thiazole, pyrazole, pyridine, pyrimidine or pyrazine.
11. The compound of any one of Claims 1 to 10 wherein R2, R3 and R4 are all hydrogen.
12. The compound of any one of Claims 1 to 10 wherein at least one of R2, R3 and R4 is C₁-C₆ alkyl.
13. The compound of Claim 12 wherein C₁-C₆ alkyl is methyl.
14. The compound of any one of Claims 1 to 10 wherein at least one of R2, R3 and R4 is halogen.
- 15 15. The compound of Claim 14 wherein halogen is chlorine or bromine.
16. The compound of Claim 1, wherein the compound is 8-Methoxy-5-methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
- 20 17. The compound of Claim 1, wherein the compound is 5-Methyl-2-(5-phenyl-2H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
- 25 18. The compound of Claim 1, wherein the compound is 5-Methyl-2-(1-methyl-5-phenyl-1H-[1,2,4]triazol-3-ylsulfanylmethyl)-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.

19. The compound of Claim 1, wherein the compound is 8-Methoxy-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
20. The compound of Claim 1, wherein the compound is 8-Methoxy-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
21. The compound of Claim 1, wherein the compound is 8-Methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
22. The compound of Claim 1, wherein the compound is 5,7-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrimidine and pharmaceutically acceptable salts thereof.
23. The compound of Claim 1, wherein the compound is 5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyrazine and pharmaceutically acceptable salts thereof.
24. The compound of Claim 1, wherein the compound is 8-Methoxy-5-methyl-2-[2-(5-phenyl-2-propyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
25. The compound of Claim 1, wherein the compound is 5-Methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
26. The compound of Claim 1, wherein the compound is 8-Methoxy-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
27. The compound of Claim 1, wherein the compound is {5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-acetic acid ethyl ester and pharmaceutically acceptable salts thereof.

28. The compound of Claim 1, wherein the compound is 2-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-ethanol and pharmaceutically acceptable salts thereof.
29. The compound of Claim 1, wherein the compound is 5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
30. The compound of Claim 1, wherein the compound is 5,8-Dimethyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-c]pyrimidine and pharmaceutically acceptable salts thereof.
31. The compound of Claim 1, wherein the compound is 8-Ethyl-5-methyl-2-[2-(2-methyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
32. The compound of Claim 1, wherein the compound is 8-Ethyl-5-methyl-2-[2-(5-phenyl-2-propyl-2H-[1,2,4]triazol-3-yl)-ethyl]-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
33. The compound of Claim 1, wherein the compound is 8-Ethyl-2-[2-(2-isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-5-methyl-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
34. The compound of Claim 1, wherein the compound is 3-{5-[2-(8-Methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile and pharmaceutically acceptable salts thereof.
35. The compound of Claim 1, wherein the compound is 3-{5-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile and pharmaceutically acceptable salts thereof.
36. The compound of Claim 1, wherein the compound is 2-[2-(2-Isopropyl-5-phenyl-2H-[1,2,4]triazol-3-yl)-ethyl]-8-methoxy-5-methyl-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.

37. The compound of Claim 1, wherein the compound is 3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propionitrile and pharmaceutically acceptable salts thereof.
38. The compound of Claim 1, wherein the compound is 3-{2-[2-(8-Ethyl-5-methyl-[1,2,4]triazolo[1,5-a]pyridin-2-yl)-ethyl]-4-phenyl-imidazol-1-yl}-propylamine and pharmaceutically acceptable salts thereof.
39. The compound of Claim 1, wherein the compound is 3-{5-[2-(5,8-Dimethyl-[1,2,4]triazolo[1,5-a]pyrazin-2-yl)-ethyl]-3-phenyl-[1,2,4]triazol-1-yl}-propylamine and pharmaceutically acceptable salts thereof.
40. The compound of Claim 1, wherein the compound is 2-{2-[2-(2-Methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazol-3-yl]-ethyl}-5,8-dimethyl-[1,2,4]triazolo[1,5-a]pyrazine and pharmaceutically acceptable salts thereof.
41. The compound of Claim 1, wherein the compound is 8-Methoxy-2-{2-[2-(2-methoxy-ethyl)-5-phenyl-2H-[1,2,4]triazol-3-yl]-ethyl}-5-methyl-[1,2,4]triazolo[1,5-a]pyridine and pharmaceutically acceptable salts thereof.
42. A compound of any one of Claims 1 to 41, but without proviso, or a pharmaceutically acceptable salt thereof for use as a medicament.



I