



(86) Date de dépôt PCT/PCT Filing Date: 2006/12/05
(87) Date publication PCT/PCT Publication Date: 2007/06/21
(85) Entrée phase nationale/National Entry: 2008/06/13
(86) N° demande PCT/PCT Application No.: EP 2006/011653
(87) N° publication PCT/PCT Publication No.: 2007/068377
(30) Priorité/Priority: 2005/12/17 (DE10 2005 060 466.8)

(51) Cl.Int./Int.Cl. *C07D 231/12* (2006.01),
A01N 43/10 (2006.01), *A01N 43/48* (2006.01),
C07D 213/61 (2006.01), *C07D 333/28* (2006.01)


(71) Demandeur/Applicant:
BAYER CROPSCIENCE AG, DE

(72) Inventeurs/Inventors:
DUNKEL, RALF, FR;
ELBE, HANS-LUDWIG, DE;
GREUL, JOERG NICO, DE;
GAYER, HERBERT, DE;
KRUEGER, BERND-WIELAND, DE;
VOERSTE, ARND, DE;
...

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : CARBOXAMIDES POUR LUTTER CONTRE LES MICRO-ORGANISMES DANS LA PROTECTION DES
PLANTES ET DES MATERIAUX

(54) Title: CARBOXAMIDES FOR CONTROLLING MICRO-ORGANISMS IN PLANT AND MATERIAL PROTECTION



(57) Abrégé/Abstract:

The invention relates to novel carboxamides of formula (I), wherein X, M, Q, Z and A have the meanings cited in the description. The invention also relates to several methods for producing said substances, to the use thereof controlling undesired micro-organisms, to novel intermediate products and to the production thereof.




(72) Inventeurs(suite)/Inventors(continued): WACHENDORFF-NEUMANN, ULRIKE, DE; DAHMEN, PETER, DE

Carboxamides

A b s t r a c t

Novel carboxamides of the formula (I)



in which X, M, Q, Z and A are as defined in the description,


a plurality of processes for preparing these compounds and their use for controlling unwanted microorganisms, and also novel intermediates and their preparation.

Carboxamides

The present invention relates to novel carboxamides, to a plurality of processes for their preparation and to the use for controlling harmful microorganisms in crop protection and in the protection of materials.

It is already known that numerous carboxamides have fungicidal properties (cf., for example, WO 03/070705, EP-A 0 545 099 and JP-A 9-132567). The activity of the compounds described in these publications is good; however, at low application rates it is sometimes unsatisfactory.

This invention now provides novel carboxamides of the formula (I)



in which

X represents OR^1 , SR^2 , NHR^3 , NR^4R^5 , SOR^6 , SO_2R^6 ,

15 R^1 represents C_1 - C_{12} -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_8 -haloalkyl, hydroxy- C_1 - C_8 -alkyl, (C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, bis(C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, (C_1 - C_4 -alkoxy)carbonyl- $(C_1$ - C_4 -alkyl); in each case optionally halogen-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted phenyl- $(C_1$ - C_2 -alkyl) or hetaryl- $(C_1$ - C_2 -alkyl); in each case optionally C_1 - C_4 -alkoxy-, halogen-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted (C_1 - C_6 -alkyl)carbonyl, (C_3 - C_8 -cycloalkyl)carbonyl, phenylcarbonyl or hetarylcarbonyl,

20 R^2 represents C_1 - C_{12} -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_8 -haloalkyl, hydroxy- C_1 - C_8 -alkyl, (C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, bis(C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, cyano- $(C_1$ - C_4 -alkyl), $-CH_2S-CN$, (C_1 - C_4 -alkoxy)carbonyl- $(C_1$ - C_4 -alkyl); in each case optionally halogen-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted phenyl- $(C_1$ - C_2 -alkyl) or hetaryl- $(C_1$ - C_2 -alkyl); in each case optionally C_1 - C_4 -alkoxy-, halogen-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted (C_1 - C_6 -alkyl)carbonyl, (C_3 - C_8 -cycloalkyl)carbonyl, phenylcarbonyl or hetarylcarbonyl,

25 R^3 , R^4 and R^5 independently of one another represent amino, hydroxyl, cyano, C_1 - C_{12} -alkyl, C_1 - C_8 -alkoxy, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_6 -alkylamino, di(C_1 - C_6 -alkyl)amino, hydroxy- C_1 - C_8 -alkyl, (C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, bis(C_1 - C_4 -alkoxy)- C_1 - C_8 -alkyl, cyano- $(C_1$ - C_4 -alkyl), phenoxy, phenylamino, benzyloxy or benzylamino,

30 R^6 represents C_1 - C_{12} -alkyl or C_1 - C_8 -haloalkyl,

M represents a phenyl, thiophene, pyridine, pyrimidine, pyridazine or pyrazine ring, each of which is monosubstituted by R^7 , or represents a thiazole ring which is substituted by R^8 ,


R^7 represents hydrogen, fluorine, chlorine, methyl, isopropyl, methylthio or trifluoromethyl,

- R^8 represents hydrogen, methyl, methylthio or trifluoromethyl,
- Q represents a direct bond, C_1 - C_4 -alkylene, C_2 - C_4 -alkenylene, O, S, SO, SO_2 or NR^9 ,
- R^9 represents hydrogen, C_1 - C_8 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl or C_3 - C_6 -cycloalkyl,
- Z represents Z^1 , Z^2 , Z^3 , Z^4 , Z^5 or Z^6 ,
- Z^1 represents phenyl which is optionally mono- to pentasubstituted by identical or different substituents,
- Z^2 represents pyridinyl which is optionally mono- to trisubstituted by identical or different substituents,
- Z^3 represents cycloalkyl or bicycloalkyl, each of which is optionally mono- or polysubstituted by identical or different halogen, alkyl and/or $-(CR^{10}R^{11})_mSiR^{12}R^{13}R^{14}$ substituents,
- Z^4 represents unsubstituted C_2 - C_{20} -alkyl or represents C_1 - C_{20} -alkyl which is mono- or polysubstituted by identical or different halogen, alkylthio, alkylsulphinyl, alkylsulphonyl, alkoxy, alkylamino, dialkylamino, haloalkylthio, haloalkylsulphinyl, haloalkylsulphonyl, haloalkoxy, haloalkylamino, halodialkylamino, $-SiR^{12}R^{13}R^{14}$ and/or C_3 - C_6 -cycloalkyl substituents, where the cycloalkyl moiety for its part may optionally be mono- or polysubstituted by identical or different halogen and/or C_1 - C_4 -alkyl substituents,
- Z^5 represents C_2 - C_{20} -alkenyl or C_2 - C_{20} -alkynyl, each of which is optionally mono- or polysubstituted by identical or different halogen, alkylthio, alkylsulphinyl, alkylsulphonyl, alkoxy, alkylamino, dialkylamino, haloalkylthio, haloalkylsulphinyl, haloalkylsulphonyl, haloalkoxy, haloalkylamino, halodialkylamino, $-SiR^{12}R^{13}R^{14}$ and/or C_3 - C_6 -cycloalkyl substituents, where the cycloalkyl moiety for its part may optionally be mono- or polysubstituted by identical or different halogen and/or C_1 - C_4 -alkyl substituents,
- Z^6 represents an optionally mono- or polysubstituted saturated or unsaturated 3- to 7-membered ring which contains a silicon atom as ring member, in which case Q represents a direct bond or C_1 - C_4 -alkylene,
- R^{10} represents hydrogen or C_1 - C_4 -alkyl,
- R^{11} represents hydrogen or C_1 - C_4 -alkyl,
- m represents 0, 1, 2 or 3,
- R^{12} and R^{13} independently of one another represent hydrogen, C_1 - C_8 -alkyl, C_1 - C_8 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl or C_1 - C_6 -haloalkyl,
- R^{14} represents hydrogen, C_1 - C_8 -alkyl, C_1 - C_8 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl, C_3 - C_6 -cycloalkyl, or represents in each case optionally substituted phenyl or phenylalkyl,

or

M-Q-Z together represent 1H-2,3-dihydroinden-4-yl, 1,3-dihydro-2-benzofuran-4-yl or 1,3-dihydro-2-benzothien-4-yl, each of which is optionally mono- to trisubstituted by methyl,

5 A represents one of the radicals A1 to A19 below



- 10 R^{15} represents hydrogen, cyano, halogen, nitro, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_3 - C_6 -cycloalkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy or C_1 - C_4 -haloalkylthio having in each case 1 to 5 halogen atoms, aminocarbonyl or aminocarbonyl- C_1 - C_4 -alkyl,
- R^{16} represents hydrogen, halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy or C_1 - C_4 -alkylthio,
- R^{17} represents hydrogen, C_1 - C_4 -alkyl, hydroxy- C_1 - C_4 -alkyl, C_2 - C_6 -alkenyl, C_3 - C_6 -cycloalkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkylthio- C_1 - C_4 -alkyl, C_1 - C_4 -haloalkoxy- C_1 - C_4 -alkyl having in each case 1 to 5 halogen atoms, or phenyl,
- 15 R^{18} and R^{19} independently of one another, represent hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{20} represents halogen, cyano or C_1 - C_4 -alkyl, or C_1 - C_4 -haloalkyl or C_1 - C_4 -haloalkoxy having in
- 20 each case 1 to 5 halogen atoms,
- R^{21} and R^{22} independently of one another represent hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{23} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,

- R^{24} represents hydrogen, halogen, hydroxy, cyano, C_1 - C_6 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy or C_1 - C_4 -haloalkylthio having in each case 1 to 5 halogen atoms,
- R^{25} represents halogen, hydroxyl, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkylthio or C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms,
- R^{26} represents hydrogen, halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms, C_1 - C_4 -alkylsulphinyl or C_1 - C_4 -alkylsulphonyl,
- R^{27} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{28} represents C_1 - C_4 -alkyl,
- Q^1 represents S (sulphur), SO, SO₂ or CH₂,
- p represents 0, 1 or 2, where R^{22} represents identical or different radicals if p represents 2,
- R^{29} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{30} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{31} and R^{32} independently of one another represent hydrogen, halogen, amino, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{33} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{34} and R^{35} independently of one another represent hydrogen, halogen, amino, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{36} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{37} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di- $(C_1$ - C_4 -alkyl)amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{38} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{39} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di- $(C_1$ - C_4 -alkyl)amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{40} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{41} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{42} represents hydrogen or C_1 - C_4 -alkyl,
- R^{43} represents halogen or C_1 - C_4 -alkyl,
- R^{44} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{45} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{46} represents halogen, hydroxyl, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkylthio or C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms,
- R^{47} represents hydrogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, hydroxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylsulphonyl, di- $(C_1$ - C_4 -alkyl)aminosulphonyl, C_1 - C_6 -alkylcarbonyl or in each case optionally substituted phenylsulphonyl or benzoyl,

- R^{48} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{49} represents hydrogen, halogen, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{50} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 5 R^{51} represents C_1 - C_4 -alkyl.

Furthermore, it has been found that carboxamides of the formula (I) are obtained by one of the processes described below.

- 10 (a) If X represents OR^1 , carboxamides of the formula (II-a)



in which M, Q, Z and A are as defined above

are, if appropriate, in the presence of a base, reacted with an alkylating agent of the formula (III-a)

- 15 LG^1-R^1 (III-a)

in which

R^1 is as defined above and

LG^1 represents a leaving group

or with a Meerwein salt of the formula (III-d)


- 20 $(R^1)_3O^+ E^-$ (III-d)

in which

R^1 is as defined above and


E represents BF_4 , SbF_6 or $SbCl_6$,

- 25 giving carboxamides of the formula (I-a)



in which R^1 , M, Q, Z and A are as defined above.


- 30 (b) If X represents SR^2 , SOR^6 or SO_2R^6 , in a first step carboxamides of the formula (II-a)



in which M, Q, Z and A are as defined above

are, if appropriate in the presence of a diluent, reacted with a thionating agent,

5 thus giving carboxamides of the formula (II-b)



in which M, Q, Z and A are as defined above

10 which, in a second step, are, if appropriate in the presence of a base, reacted with an alkylating agent of the formula (III-b) or the formula (III-c)





in which

R^2 and R^6 are as defined above,

15 LG^2 represents a leaving group,

LG^6 represents a leaving group,

giving carboxamides of the formula (I-b) or the formula (I-c)




20

in which R^2 , R^6 , M, Q, Z and A are as defined above,

and, in a third step, the carboxamides of the formula (I-c) are then reacted with an oxidizing agent, if appropriate in the presence of a diluent, if appropriate in the presence of


an acid binder and if appropriate in the presence of a catalyst, thus giving the carboxamides of the formula (I-d)



in which



5 R^6 , M, Q, Z and A are as defined above and
n represents 1 or 2.

(c) If X represents NHR^3 or NR^4R^5 , carboxamides of the formula (II-b)



10 in which M, Q, Z and A are as defined above

are, if appropriate in the presence of a base, reacted with an amidating agent of the formula (IV-a) or the formula (IV-b)





15

in which

R^3 , R^4 and R^5 are as defined above and

LG^3 represents a leaving group,

20 giving carboxamides of the formula (I-e) or the formula (I-f)



in which R^3 , R^4 , R^5 , M, Q, Z and A are as defined above.

Finally, it has been found that the novel carboxamides of the formula (I) have very good microbicidal properties and can be used for controlling unwanted microorganisms both in crop protection and in the protection of materials.

If appropriate, the compounds according to the invention can be present as mixtures of different possible isomeric forms, in particular of stereoisomers, such as, for example, E and Z, threo and erythro, and also optical isomers, and, if appropriate, also of tautomers. What is claimed are both the E and the Z isomers, and also the threo and erythro and the optical isomers, and the mixtures of these isomers and the possible tautomeric forms.

The formula (I) provides a general definition of the biphenylthiazolecarboxamides according to the invention. Preferred radical definitions of the formulae mentioned above and below are stated below. These definitions apply to the end products of the formula (I) and likewise to all intermediates.

- X preferably represents OR^1 .
X also preferably represents SR^2 .
20 X also preferably represents NHR^3 or NR^4R^5 .
X also preferably represents SOR^6 or SO_2R^6 .

R^1 preferably represents C_1 - C_8 -alkyl, C_2 - C_6 -alkenyl, C_2 - C_6 -alkynyl, C_1 - C_4 -haloalkyl, hydroxy- C_1 - C_6 -alkyl, $(C_1$ - C_2 -alkoxy)- C_1 - C_6 -alkyl, bis(C_1 - C_2 -alkoxy)- C_1 - C_6 -alkyl, $(C_1$ - C_2 -alkoxy)carbonyl- $(C_1$ - C_2 -alkyl); in each case optionally fluorine-, chlorine-, bromine-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl- $(C_1$ - C_2 -alkyl) or thienyl- $(C_1$ - C_2 -alkyl); in each case optionally C_1 - C_4 -alkoxy, fluorine-, chlorine-, bromine-, C_1 - C_4 -alkyl- or C_1 - C_4 -haloalkyl-substituted $(C_1$ - C_4 -alkyl)-carbonyl, $(C_3$ - C_6 -cycloalkyl)carbonyl, phenylcarbonyl, pyridinylcarbonyl or thienyl-carbonyl.

R^1 particularly preferably represents methyl, ethyl, n-, isopropyl, n-, i-, s-, t-butyl, allyl, propargyl, trifluoromethyl, difluoromethyl, trichloromethyl, dichloromethyl, $-CH_2CF_3$, pentafluoroethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, dimethoxymethyl, dimethoxyethyl, diethoxymethyl, diethoxyethyl, methoxycarbonylmethyl, methoxycarbonyl-ethyl, ethoxycarbonylmethyl, ethoxycarbonyl-ethyl; in each case optionally fluorine-, chlorine-, bromine-, methyl- or trifluoromethyl-substituted

benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl-(C₁-C₂-alkyl) or thienyl-(C₁-C₂-alkyl); in each case optionally methoxy-, ethoxy-, fluorine-, chlorine-, bromine-, methyl-, trifluoromethyl-substituted methylcarbonyl, ethylcarbonyl, cyclopropylcarbonyl, cyclohexylcarbonyl, phenylcarbonyl, pyridinylcarbonyl or thienylcarbonyl.

5 R¹ very particularly preferably represents methyl, ethyl, n-propyl, n-butyl, n-hexyl, n-octyl, n-decyl, n-dodecyl, allyl, propargyl, trifluoromethyl, trichloromethyl, -CH₂CF₃, pentafluoroethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, dimethoxymethyl, dimethoxyethyl, diethoxymethyl, diethoxyethyl, methoxy-carbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl; in each
10 case optionally fluorine-, chlorine-, bromine-, methyl- or trifluoromethyl-substituted benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl-(C₁-C₂-alkyl) or thienyl-(C₁-C₂-alkyl) (with emphasis 6-chloro-3-pyridinylmethyl, 2,6-dichloro-4-pyridinylmethyl, 4,5-dibromo-2-thienyl); in each case optionally methoxy-, ethoxy-, fluorine-, chlorine-, bromine-, methyl-, trifluoromethyl-substituted methylcarbonyl, ethylcarbonyl, cyclopropylcarbonyl, cyclohexylcarbonyl,
15 phenylcarbonyl, pyridinylcarbonyl or thienylcarbonyl (with emphasis 6-chloro-3-pyridinylcarbonyl, 2,6-dichloro-4-pyridinylcarbonyl, 4,5-dibromo-2-thienylcarbonyl).

R² preferably represents C₁-C₈-alkyl, dodecyl, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₄-haloalkyl, hydroxy-C₁-C₆-alkyl, (C₁-C₂-alkoxy)-C₁-C₆-alkyl, bis(C₁-C₂-alkoxy)-C₁-C₆-alkyl, cyano-
20 (C₁-C₄-alkyl), -CH₂S-CN, (C₁-C₂-alkoxy)carbonyl-(C₁-C₂-alkyl); in each case optionally fluorine-, chlorine-, bromine-, C₁-C₄-alkyl- or C₁-C₄-haloalkyl-substituted benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl-(C₁-C₂-alkyl) or thienyl-(C₁-C₂-alkyl); in each case optionally C₁-C₄-alkoxy-, fluorine-, chlorine-, bromine-, C₁-C₄-alkyl- or C₁-C₄-haloalkyl-substituted (C₁-C₄-alkyl)carbonyl, (C₃-C₆-cycloalkyl)carbonyl, phenylcarbonyl,
25 pyridinylcarbonyl or thienylcarbonyl.

R² particularly preferably represents methyl, ethyl, n-, isopropyl, n-, i-, s-, t-butyl, n-hexyl, n-octyl, n-decyl, dodecyl, allyl, propargyl, trifluoromethyl, difluoromethyl, trichloromethyl, dichloromethyl, -CH₂CF₃, pentafluoroethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, dimethoxymethyl,
30 dimethoxyethyl, diethoxymethyl, diethoxyethyl, cyanomethyl, cyanoethyl, -CH₂S-CN, methoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl; in each case optionally fluorine-, chlorine-, bromine-, methyl- or trifluoromethyl-substituted benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl-(C₁-C₂-alkyl) or thienyl-(C₁-C₂-alkyl); in each case optionally methoxy-, ethoxy-, fluorine-, chlorine-,
35 bromine-, methyl-, trifluoromethyl-substituted methylcarbonyl, ethylcarbonyl,

cyclopropylcarbonyl, cyclohexylcarbonyl, phenylcarbonyl, pyridinylcarbonyl or thienylcarbonyl.

5 R² very particularly preferably represents methyl, ethyl, n-propyl, n-butyl, n-hexyl, n-octyl, n-decyl, n-dodecyl, allyl, propargyl, trifluoromethyl, trichloromethyl, -CH₂CF₃, pentafluoroethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, dimethoxymethyl, dimethoxyethyl, diethoxymethyl, diethoxyethyl, cyanomethyl, cyanoethyl, -CH₂S-CN, methoxycarbonylmethyl, methoxycarbonylethyl, ethoxycarbonylmethyl, ethoxycarbonylethyl; in each case optionally
10 fluorine-, chlorine-, bromine-, methyl- or trifluoromethyl-substituted benzyl, phenyl-1-ethyl, phenyl-2-ethyl, pyridinyl-(C₁-C₂-alkyl) or thienyl-(C₁-C₂-alkyl) (with emphasis 6-chloro-3-pyridinylmethyl, 2,6-dichloro-4-pyridinylmethyl, 4,5-dibromo-2-thienyl); in each case optionally methoxy-, ethoxy-, fluorine-, chlorine-, bromine-, methyl-, trifluoromethyl-substituted methylcarbonyl, ethylcarbonyl, cyclopropylcarbonyl, cyclohexylcarbonyl, phenylcarbonyl, pyridinylcarbonyl or thienylcarbonyl (with emphasis 6-chloro-3-pyridinylcarbonyl, 2,6-dichloro-4-pyridinylcarbonyl, 4,5-dibromo-2-thienylcarbonyl).
15

R³, R⁴ and R⁵ independently of one another preferably represent hydrogen, amino, hydroxyl, cyano, C₁-C₈-alkyl, C₁-C₆-alkoxy, C₂-C₆-alkenyl, C₂-C₆-alkynyl, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, hydroxy-C₁-C₆-alkyl, (C₁-C₂-alkoxy)-C₁-C₆-alkyl, bis(C₁-C₂-alkoxy)-C₁-C₆-alkyl, cyano-(C₁-C₄-alkyl), phenoxy, phenylamino, benzyloxy or benzylamino.
20


R³, R⁴ and R⁵ independently of one another particularly preferably represent hydrogen, amino, hydroxyl, cyano, methyl, ethyl, n-, isopropyl, n-, i-, s-, t-butyl, methoxy, ethoxy, n-, isopropoxy, n-, i-, s-, t-butoxy, allyl, propargyl, methylamino, ethylamino, n-, isopropylamino, n-, i-, s-, t-butylamino, dimethylamino, diethylamino, hydroxymethyl, hydroxyethyl, hydroxypropyl, methoxymethyl, methoxyethyl, ethoxymethyl, ethoxyethyl, dimethoxymethyl, dimethoxyethyl, diethoxymethyl, diethoxyethyl, cyanomethyl, phenoxy, phenylamino, benzyloxy or benzylamino.
25

R⁶ preferably represents C₁-C₄-alkyl or C₁-C₄-haloalkyl.

30 R⁶ particularly preferably represents methyl, ethyl, n-, isopropyl, n-, i-, s-, t-butyl, trifluoromethyl, difluoromethyl, trichloromethyl, dichloromethyl, -CH₂CF₃, pentafluoroethyl.

R⁶ very particularly preferably represents methyl, ethyl, trifluoromethyl, trichloromethyl, -CH₂CF₃, pentafluoroethyl.

35 M preferably represents one of the cycles below



where the bond marked “*” is linked to the amide and the bond marked “#” is linked to the radical R.

- 5 M particularly preferably represents a cycle selected from the group consisting of M-1, M-2, M-3, M-4, M-5, M-6, M-9, M-10 and M-11.
- M very particularly preferably represents a cycle selected from the group consisting of M-1, M-2, M-5, M-6, M-9, M-10 and M-11.
- 10 M especially preferably represents the cycle M-1.
- M furthermore especially preferably represents the heterocycle M-2.
- M furthermore especially preferably represents the heterocycle M-5.
- M furthermore especially preferably represents the heterocycle M-6.
- M furthermore especially preferably represents the heterocycle M-9.
- 15 M furthermore especially preferably represents the heterocycle M-10.
- M furthermore especially preferably represents the heterocycle M-11.
- R⁷ preferably represents hydrogen.
- R⁷ furthermore, if M represents M-1, preferably represents fluorine, where fluorine is particularly preferably located in the 4-, 5- or 6-position, very particularly preferably in the 4- or 6-position, especially in the 4-position, of the anilide radical.
- 20 R⁷ furthermore, if M represents M-1, preferably represents chlorine, where chlorine is particularly preferably located in the 5-position of the anilide radical.
- R⁷ furthermore, if M represents M-1, preferably represents methyl, where methyl is particularly preferably located in the 3-position of the anilide radical.
- 25

- R⁷ furthermore, if M represents M-1, preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 4- or 5-position of the anilide radical.
- R⁷ furthermore, if M represents M-2, M-3 or M-4, preferably represents chlorine, where chlorine is particularly preferably located in the 5-position (M-2, M-3) or in the 3-position (M-4).
- R⁷ furthermore, if M represents M-2, M-3 or M-4, preferably represents methyl, where methyl is particularly preferably located in the 5-position (M-2, M-3) or in the 3-position (M-4).
- R⁷ furthermore, if M represents M-5, M-6, M-7 or M-8, preferably represents fluorine, where fluorine is particularly preferably located in the 6-position (M-5, M-6) or in the 3-position (M-7, M-8).
- R⁷ furthermore, if M represents M-5, M-6, M-7 or M-8, preferably represents chlorine, where chlorine is particularly preferably located in the 6-position (M-5, M-6) or in the 3-position (M-7, M-8).
- R⁷ furthermore, if M represents M-5, M-6, M-7 or M-8, preferably represents methyl, where methyl is particularly preferably located in the 4-position (M-5) or in the 3-position (M-6, M-7, M-8).
- R⁷ furthermore, if M represents M-9, preferably represents methyl, where methyl is particularly preferably located in the 3-position.
- R⁷ furthermore, if M represents M-9, preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 3-position.
- R⁷ furthermore, if M represents M-12, preferably represents methyl, where methyl is particularly preferably located in the 4-position.
- R⁷ furthermore, if M represents M-12, preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 4-position.
- R⁷ furthermore, if M represents M-13, preferably represents methyl, where methyl is particularly preferably located in the 3-position.
- R⁷ furthermore, if M represents M-13, preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 3-position.
- R⁷ furthermore, if M represents M-14, preferably represents methyl, where methyl is particularly preferably located in the 3-position.
- R⁷ furthermore, if M represents M-14, preferably represents trifluoromethyl, where trifluoromethyl is particularly preferably located in the 3-position.
- R⁸ preferably represents hydrogen.
- R⁸ furthermore preferably represents methyl.
- R⁸ furthermore preferably represents trifluoromethyl.

- Q preferably represents a direct bond.
- Q furthermore preferably represents $-\text{CH}_2-$, $-(\text{CH}_2)_2-$, $-(\text{CH}_2)_3-$, $-\text{CH}(\text{CH}_3)-$, $-\text{C}(\text{CH}_3)_2-$,
particularly preferably represents $-\text{CH}_2-$, $-(\text{CH}_2)_2-$, $-\text{CH}(\text{CH}_3)-$.
- 5 Q furthermore preferably represents $-\text{CH}=\text{CH}-$, $-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{CH}(\text{CH}_3)-\text{CH}=\text{CH}-$,
particularly preferably represents $-\text{CH}=\text{CH}-$, $-\text{CH}_2-\text{CH}=\text{CH}-$.
- Q furthermore preferably represents O.
- Q furthermore preferably represents S.
- Q furthermore preferably represents SO.
- 10 Q furthermore preferably represents SO_2 .
- Q furthermore preferably represents NR^9 , particularly preferably NH.
- R^9 preferably represents hydrogen, C_1 - C_6 -alkyl, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl, C_1 - C_3 -alkylthio- C_1 - C_3 -alkyl or C_3 - C_6 -cycloalkyl.
- 15 R^9 particularly preferably represents hydrogen, methyl, ethyl, n- or isopropyl, n-, sec-, iso- or
tert-butyl, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl,
ethylthiomethyl, methylthioethyl, ethylthioethyl or cyclopropyl.
- R^9 very particularly preferably represents hydrogen, methyl, ethyl, n- or isopropyl, iso- or tert-
butyl, methoxymethyl or methylthiomethyl.
- 20
- Z preferably represents Z^1 .
- Z^1 preferably represents phenyl which is optionally mono- to pentasubstituted by identical or
different substituents, where the substituents are in each case selected from the list W^1 .
- Z^1 particularly preferably represents monosubstituted phenyl, where the substituents are
25 selected from the list W^1 .
- Z^1 also particularly preferably represents phenyl which is disubstituted by identical or
different substituents, where the substituents are selected from the list W^1 .
- Z^1 also particularly preferably represents phenyl which is trisubstituted by identical or
different substituents, where the substituents are selected from the list W^1 .
- 30 Z^1 very particularly preferably represents phenyl which is monosubstituted in the 4-position,
where the substituents are selected from the list W^1 .
- Z^1 very particularly preferably represents phenyl which is disubstituted by identical or
different substituents in the 3,4-position, where the substituents are selected from the list
 W^1 .

- Z^1 very particularly preferably represents phenyl which is disubstituted by identical or different substituents in the 2,3-position, where the substituents are selected from the list W^1 .
- 5 Z^1 very particularly preferably represents phenyl which is disubstituted by identical or different substituents in the 2,4-position, where the substituents are selected from the list W^1 .
- Z^1 very particularly preferably represents phenyl which is disubstituted by identical or different substituents in the 3,5-position, where the substituents are selected from the list W^1 .
- 10 Z^1 very particularly preferably represents phenyl which is trisubstituted by identical or different substituents in the 2,4,6-position, where the substituents are selected from the list W^1 .
- 15 W^1 represents halogen, cyano, nitro, amino, hydroxyl, formyl, carboxy, carbamoyl, thio-carbamoyl;
in each case straight-chain or branched alkyl, hydroxyalkyl, oxoalkyl, alkoxy, alkoxyalkyl, alkylthioalkyl, dialkoxyalkyl, alkylthio, alkylsulphanyl or alkylsulphonyl having in each case 1 to 8 carbon atoms;
in each case straight-chain or branched alkenyl or alkenyloxy having in each case 2 to 6 carbon atoms;
20 in each case straight-chain or branched haloalkyl, haloalkoxy, haloalkylthio, haloalkylsulphanyl or haloalkylsulphonyl having in each case 1 to 6 carbon atoms and 1 to 13 identical or different halogen atoms;
in each case straight-chain or branched haloalkenyl or haloalkenyloxy having in each case 2 to 6 carbon atoms and 1 to 11 identical or different halogen atoms;
25 in each case straight-chain or branched alkylamino, dialkylamino, alkylcarbonyl, alkylcarbonyloxy, alkoxycarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, arylalkylaminocarbonyl, dialkylaminocarbonyloxy having 1 to 6 carbon atoms in the respective hydrocarbon chains, alkenylcarbonyl or alkynylcarbonyl having 2 to 6 carbon atoms in the respective hydrocarbon chains;
30 cycloalkyl or cycloalkyloxy having in each case 3 to 6 carbon atoms;
in each case doubly attached alkylene having 3 or 4 carbon atoms, oxyalkylene having 2 or 3 carbon atoms or dioxyalkylene having 1 or 2 carbon atoms, each of which radicals is optionally mono- to tetrasubstituted by identical or different substituents from the group
35 consisting of fluorine, chlorine, oxo, methyl, trifluoromethyl and ethyl;
or the groupings $-(CR^{10}R^{11})_mSiR^{12}R^{13}R^{14}$ or $-C(Q^2)=N-Q^3$, in which


- Q² represents hydrogen, hydroxyl or alkyl having 1 to 4 carbon atoms, haloalkyl having 1 to 4 carbon atoms and 1 to 9 fluorine, chlorine and/or bromine atoms or cycloalkyl having 1 to 6 carbon atoms and
- Q³ represents hydroxyl, amino, methylamino, phenyl, benzyl or represents in each case optionally cyano-, hydroxyl-, alkoxy-, alkylthio-, alkylamino-, dialkylamino- or phenyl-substituted alkyl or alkoxy having 1 to 4 carbon atoms, or represents alkenyloxy or alkynyloxy having in each case 2 to 4 carbon atoms, and also phenyl, phenoxy, phenylthio, benzoyl, benzoylethenyl, cinnamoyl, heterocyclyl or phenylalkyl, phenylalkyloxy, phenylalkylthio or heterocyclylalkyl having in each case 1 to 3 carbon atoms in the respective alkyl moieties, each of which radicals is optionally mono- to trisubstituted in the ring moiety by halogen and/or straight-chain or branched alkyl or alkoxy having 1 to 4 carbon atoms.
- W¹ preferably represents fluorine, chlorine, bromine, methyl, ethyl, n- or i-propyl, n-, i-, s- or t-butyl, methoxy, ethoxy, n- or i-propoxy, trifluoromethyl, trifluoroethyl, difluoromethoxy, trifluoromethoxy, difluorochloromethoxy, trifluoroethoxy, in each case doubly attached difluoromethylenedioxy or tetrafluoroethylenedioxy, or the groupings -CH₂Si(CH₃)₃, -Si(CH₃)₃ or -C(Q²)=N-Q³, in which
- Q² represents hydrogen, methyl, ethyl or trifluoromethyl and
- Q³ represents hydroxyl, methoxy, ethoxy, propoxy or isopropoxy.
- Z preferably represents Z².
- Z² preferably represents 2-pyridinyl, 3-pyridinyl or 4-pyridinyl, each of which is optionally mono- to trisubstituted by identical or different substituents, where the substituents are in each case selected from the list W².
- Z² particularly preferably represents in each case monosubstituted 2-pyridinyl, 3-pyridinyl or 4-pyridinyl, where the substituents are in each case selected from the list W².
- Z² also particularly preferably represents 2-pyridinyl, 3-pyridinyl or 4-pyridinyl, each of which is disubstituted by identical or different substituents, where the substituents are in each case selected from the list W².
- Z² also particularly preferably represents 2-pyridinyl, 3-pyridinyl or 4-pyridinyl, each of which is trisubstituted by identical or different substituents, where the substituents are in each case selected from the list W².
- Z² very particularly preferably represents 2-pyridinyl which is monosubstituted in the 5-position or 3-pyridinyl which is monosubstituted in the 6-position, where the substituents are in each case selected from the list W².

- Z^2 very particularly preferably represents 2-pyridinyl which is disubstituted by identical or different substituents in the 3,5-position, where the substituents are selected from the list W^2 .
- Z^2 very particularly preferably represents 3-pyridinyl which is disubstituted by identical or different substituents in the 4,6-position, where the substituents are selected from the list W^2 .
- Z^2 very particularly preferably represents 4-pyridinyl which is disubstituted by identical or different substituents in the 3,5-position, where the substituents are selected from the list W^2 .
- W^2 represents hydrogen, halogen, cyano, nitro, C_1 - C_6 -alkyl, C_2 - C_4 -alkenyl, C_2 - C_4 -alkynyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -alkylsulphinyl, C_1 - C_4 -alkylsulphonyl, C_3 - C_6 -cycloalkyl; represents C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylthio, C_1 - C_4 -haloalkylsulphinyl, C_1 - C_4 -haloalkylsulphonyl having in each case 1 to 5 halogen atoms; represents $-SO_2NR^{52}R^{53}$, $-C(=X)R^{54}$, $-Si(R^{55})_3$, C_2 - C_4 -alkenylene- $-Si(R^{55})_3$, C_2 - C_4 -alkynylene- $-Si(R^{55})_3$, $-NR^{52}R^{53}$, $-CH_2-NR^{52}R^{53}$, in which
- R^{52} represents hydrogen, C_1 - C_4 -alkyl or $-C(=X)R^{54}$,
- R^{53} represents hydrogen, C_1 - C_4 -alkyl or $-C(=X)R^{54}$,
- R^{52} and R^{53} furthermore together with the nitrogen atom to which they are attached form a saturated heterocycle which has 5 to 8 ring atoms and is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and C_1 - C_4 -alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR^{56} ,
- R^{54} represents hydrogen, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy or $-NR^{57}R^{58}$,
- R^{55} represents hydrogen, C_1 - C_8 -alkyl, C_1 - C_8 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl or C_1 - C_6 -haloalkyl, where the three radicals R^{55} may in each case be identical or different,
- R^{56} represents hydrogen or C_1 - C_6 -alkyl,
- R^{57} represents hydrogen or C_1 - C_4 -alkyl,
- R^{58} represents hydrogen or C_1 - C_4 -alkyl,
- R^{57} and R^{58} furthermore together with the nitrogen atom to which they are attached represent a saturated heterocycle having 5 to 8 ring atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of halogen and C_1 - C_4 -alkyl, where the heterocycle may contain 1 or 2 further non-adjacent heteroatoms from the group consisting of oxygen, sulphur and NR^{56} ,

- W² preferably represents hydrogen, fluorine, chlorine, bromine, cyano, nitro, methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl, allyl, propargyl, methoxy, ethoxy, n- or isopropoxy, n-, iso-, sec- or tert-butoxy, methylthio, ethylthio, n- or isopropylthio, n-, iso-, sec- or tert-butylthio, methylsulphinyl, ethylsulphinyl, n- or isopropylsulphinyl, n-, iso-, sec- or tert-butylsulphinyl, methylsulphonyl, ethylsulphonyl, n- or iso-propylsulphonyl, n-, iso-, sec- or tert-butylsulphonyl, cyclopropyl, cyclopentyl, cyclohexyl, trifluoromethyl, difluoromethyl, trichloromethyl, trifluoroethyl, trifluoromethoxy, difluoromethoxy, trichloromethoxy, difluoromethylthio, difluorochloromethylthio, trifluoromethylthio, trifluoromethylsulphinyl, trifluoromethylsulphonyl, -SO₂NMe₂, -C(=X)R⁵⁴, -Si(R⁵⁵)₃, -CH=CH-Si(R⁵⁵)₃, -CH₂-CH=CH-Si(R⁵⁵)₃, -CH=CH-CH₂-Si(R⁵⁵)₃, -C≡C-Si(R⁵⁵)₃, -CH₂-C≡C-Si(R⁵⁵)₃, -C≡C-CH₂-Si(R⁵⁵)₃, -CH₂-C≡C-CH₂-Si(R⁵⁵)₃, -NR⁵²R⁵³, -CH₂-NR⁵²R⁵³.
- R⁵² preferably represents hydrogen, methyl, ethyl, n- or isopropyl or -C(=X)R⁵⁴.
- R⁵² particularly preferably represents hydrogen or methyl.
- R⁵³ preferably represents hydrogen, methyl, ethyl, n- or isopropyl or -C(=X)R⁵⁴.
- R⁵³ particularly preferably represents hydrogen or methyl.
- R⁵² and R⁵³ furthermore together with the nitrogen atom to which they are attached preferably form a saturated heterocycle from the group consisting of morpholine, thiomorpholine and piperazine which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine and methyl, where the piperazine may be substituted on the second nitrogen atom by R⁵⁶.
- R⁵⁴ preferably represents hydrogen, methyl, ethyl, n- or isopropyl, methoxy, ethoxy, n- or isopropoxy or -NR⁵⁷R⁵⁸.
- R⁵⁴ particularly preferably represents hydrogen, methyl, ethyl, methoxy, ethoxy or -NR⁵⁷R⁵⁸.
- R⁵⁵ preferably represents methyl, ethyl, methoxy, ethoxy, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl, ethylthiomethyl, methylthioethyl or ethylthioethyl, where the three radicals R⁵⁵ may in each case be identical or different.
- R⁵⁵ particularly preferably represents methyl, methoxy, methoxymethyl or methylthiomethyl, where the three radicals R⁵⁵ may in each case be identical or different.
- R⁵⁵ very particularly preferably represents methyl.
- R⁵⁶ preferably represents hydrogen or C₁-C₄-alkyl.

- R⁵⁶ particularly preferably represents hydrogen, methyl, ethyl, n- or isopropyl, n-, iso-, sec- or tert-butyl.
- R⁵⁷ preferably represents hydrogen, methyl, ethyl, n- or isopropyl.
- 5 R⁵⁷ particularly preferably represents hydrogen or methyl.
- R⁵⁸ preferably represents hydrogen, methyl, ethyl, n- or isopropyl.
- R⁵⁸ particularly preferably represents hydrogen or methyl.
- 10 R⁵⁷ and R⁵⁸ furthermore together with the nitrogen atom to which they are attached preferably form a saturated a heterocycle from the group consisting of morpholine, thiomorpholine and piperazine which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine and methyl, where the piperazine may be substituted on the second nitrogen atom by R⁷.
- 15 Z also preferably represents Z³.
- Z³ preferably represents cycloalkyl or bicycloalkyl having in each case 3 to 10 carbon atoms, each of which radicals is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of halogen, C₁-C₄-alkyl, -CH₂Si(CH₃)₃ and/or -Si(CH₃)₃.
- 20 Z³ particularly preferably represents cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, bicyclo[2.2.1]heptyl or bicyclo[2.2.2]octyl, each of which is optionally mono- to tetrasubstituted by identical or different substituents from the group consisting of chlorine, methyl, -CH₂Si(CH₃)₃ and/or -Si(CH₃)₃.
- 25 Z³ very particularly preferably represents chlorine- and methyl-substituted cyclopropyl.
- Z also preferably represents Z⁴.
- Z⁴ preferably represents unsubstituted C₂-C₂₀-alkyl or represents C₁-C₂₀-alkyl which is mono- or polysubstituted by identical or different substituents from the group consisting of
- 30 fluorine, chlorine, bromine, iodine, C₁-C₆-alkylthio, C₁-C₆-alkylsulphinyl, C₁-C₆-alkylsulphonyl, C₁-C₆-alkoxy, C₁-C₆-alkylamino, di(C₁-C₆-alkyl)amino, C₁-C₆-haloalkylthio, C₁-C₆-haloalkylsulphinyl, C₁-C₆-haloalkylsulphonyl, C₁-C₆-haloalkoxy, C₁-C₆-haloalkylamino, halo-di(C₁-C₆-alkyl)amino, -SiR¹²R¹³R¹⁴ and/or C₃-C₆-cycloalkyl, where the cycloalkyl moiety for its part may optionally be mono- to tetrasubstituted by identical or different
- 35 substituents from the group consisting of fluorine, chlorine, bromine, iodine, C₁-C₄-alkyl and/or C₁-C₄-haloalkyl.

- Z^4 particularly preferably represents unsubstituted C_2 - C_{20} -alkyl.
- Z^4 also particularly preferably represents C_1 - C_{20} -alkyl which is substituted by fluorine, chlorine, bromine, iodine, C_1 - C_6 -alkylthio, C_1 - C_4 -alkylsulphinyl, C_1 - C_4 -alkylsulphonyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino, C_1 - C_4 -haloalkylthio, C_1 - C_4 -haloalkylsulphinyl, C_1 - C_4 -haloalkylsulphonyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylamino, halo-di(C_1 - C_4 -alkyl)amino having in each case 1 to 9 fluorine, chlorine and/or bromine atoms, $-\text{SiR}^6\text{R}^7\text{R}^8$, cyclopropyl, dichlorocyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl; very particularly preferably C_1 - C_{20} -alkyl which is substituted by fluorine, chlorine, methylthio, ethylthio, n- or isopropylthio, n-, iso-, sec-, tert-butylthio, pentylthio, hexylthio, methylsulphonyl, ethylsulphonyl, n- or isopropylsulphonyl, n-, iso-, sec-, tert-butylsulphonyl, methoxy, ethoxy, n- or isopropoxy, n-, iso-, sec-, tert-butoxy, methylamino, ethylamino, n- or isopropylamino, n-, iso-, sec-, tert-butylamino, dimethylamino, diisopropylamino, trifluoromethylthio, trifluoromethoxy, $-\text{SiR}^{12}\text{R}^{13}\text{R}^{14}$, cyclopropyl, dichlorocyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.
- 15 Z also preferably represents Z^5 .
- Z^5 preferably represents C_2 - C_{20} -alkenyl or C_2 - C_{20} -alkynyl, each of which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine, iodine, C_1 - C_6 -alkylthio, C_1 - C_6 -alkylsulphinyl, C_1 - C_6 -alkylsulphonyl, C_1 - C_6 -alkoxy, C_1 - C_6 -alkylamino, di(C_1 - C_6 -alkyl)amino, C_1 - C_6 -haloalkylthio, C_1 - C_6 -haloalkylsulphinyl, C_1 - C_6 -haloalkylsulphonyl, C_1 - C_6 -haloalkoxy, C_1 - C_6 -haloalkylamino, halo-di(C_1 - C_6 -alkyl)amino, $-\text{SiR}^{12}\text{R}^{13}\text{R}^{14}$ and/or C_3 - C_6 -cycloalkyl, where the cycloalkyl moiety for its part may optionally be mono- to tetrasubstituted by identical or different substituents from the group consisting of fluorine, chlorine, bromine, iodine, C_1 - C_4 -alkyl and/or C_1 - C_4 -haloalkyl.
- 20 Z^5 particularly preferably represents C_2 - C_{20} -alkenyl or C_2 - C_{20} -alkynyl, each of which is optionally substituted by fluorine, chlorine, bromine, iodine, C_1 - C_6 -alkylthio, C_1 - C_4 -alkylsulphinyl, C_1 - C_4 -alkylsulphonyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylamino, di(C_1 - C_4 -alkyl)amino, C_1 - C_4 -haloalkylthio, C_1 - C_4 -haloalkylsulphinyl, C_1 - C_4 -haloalkylsulphonyl, C_1 - C_4 -haloalkoxy, C_1 - C_4 -haloalkylamino, halo-di(C_1 - C_4 -alkyl)amino having in each case 1 to 9 fluorine, chlorine and/or bromine atoms, $-\text{SiR}^{12}\text{R}^{13}\text{R}^{14}$, cyclopropyl, dichlorocyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl.
- 30 Z^5 very particularly preferably represents C_2 - C_{20} -alkenyl or C_2 - C_{20} -alkynyl.
- Z also preferably represents Z^6 .
- Z^6 preferably represents one of the rings below



in which R⁵⁹ represents hydrogen or methyl.

5 R¹⁰ preferably represents hydrogen or methyl.

R¹⁰ particularly preferably represents hydrogen.

R¹¹ preferably represents hydrogen or methyl.

R¹¹ particularly preferably represents hydrogen.

10

m preferably represents 0, 1 or 2.

R¹² and R¹³ independently of one another preferably represent C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₃-alkoxy-C₁-C₃-alkyl or C₁-C₃-alkylthio-C₁-C₃-alkyl.

15 R¹² and R¹³ independently of one another particularly preferably represent methyl, ethyl, methoxy, ethoxy, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl, ethylthiomethyl, methylthioethyl or ethylthioethyl.

R¹² and R¹³ independently of one another very particularly preferably represent methyl, methoxy, methoxymethyl or methylthiomethyl.

20 R¹² and R¹³ especially preferably each represent methyl.

R¹⁴ preferably represents C₁-C₆-alkyl, C₁-C₆-alkoxy, C₁-C₃-alkoxy-C₁-C₃-alkyl, C₁-C₃-alkylthio-C₁-C₃-alkyl, C₃-C₆-cycloalkyl, phenyl or benzyl.

25 R¹⁴ particularly preferably represents methyl, ethyl, n- or isopropyl, n-, sec-, iso- or tert-butyl, methoxy, ethoxy, n- or isopropoxy, n-, sec-, iso- or tert-butoxy, methoxymethyl, ethoxymethyl, methoxyethyl, ethoxyethyl, methylthiomethyl, ethylthiomethyl, methylthioethyl, ethylthioethyl, cyclopropyl, phenyl or benzyl.

R¹⁴ very particularly preferably represents methyl, ethyl, n- or isopropyl, iso- or tert-butyl, methoxy, isopropoxy, iso- or tert-butoxy, methoxymethyl, methylthiomethyl or phenyl.

30 R¹⁴ especially preferably represents methyl, ethyl, n- or isopropyl, iso- or tert-butyl, methoxy, isopropoxy, iso- or tert-butoxy.

R¹⁴ most preferably represents methyl.

M-Q-Z also preferably together represents 1,1,3-trimethyl-1H-2,3-dihydroinden-4-yl, 1,3-dimethyl-1H-2,3-dihydroinden-4-yl, 1,1,3-trimethyl-1,3-dihydro-2-benzofuran-4-yl, 1,3-dimethyl-1,3-dihydro-2-benzofuran-4-yl, 1,1,3-trimethyl-1,3-dihydro-2-benzothien-4-yl or 1,3-dimethyl-1,3-dihydro-2-benzothien-4-yl.

M-Q-Z also particularly preferably together represents 1,1,3-trimethyl-1H-2,3-dihydroinden-4-yl.

A preferably represents one of the radicals A1, A2, A3, A4, A5, A6, A9, A10, A11, A12 or A17.

A particularly preferably represents one of the radicals A1, A2, A4, A5, A6, A9, A11, A16, A17.

A very particularly preferably represents the radical A1.

A furthermore very particularly preferably represents the radical A2.

15 A furthermore very particularly preferably represents the radical A4.

A furthermore very particularly preferably represents the radical A5.

A furthermore very particularly preferably represents the radical A6.

A furthermore very particularly preferably represents the radical A9.

A furthermore very particularly preferably represents the radical A11.

20 A furthermore very particularly preferably represents the radical A16.

A furthermore very particularly preferably represents the radical A17.

R¹⁵ preferably represents hydrogen, cyano, fluorine, chlorine, bromine, iodine, methyl, ethyl, isopropyl, methoxy, ethoxy, methylthio, ethylthio, cyclopropyl, C₁-C₂-haloalkyl, C₁-C₂-haloalkoxy having in each case 1 to 5 fluorine, chlorine and/or bromine atoms, trifluoromethylthio, difluoromethylthio, aminocarbonyl, aminocarbonylmethyl or aminocarbonylethyl.

R¹⁵ particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, isopropyl, monofluoromethyl, monofluoroethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, dichloromethyl, cyclopropyl, methoxy, ethoxy, trifluoromethoxy, trichloromethoxy, methylthio, ethylthio, trifluoromethylthio or difluoromethylthio.

R¹⁵ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, isopropyl, monofluoromethyl, monofluoroethyl, difluoromethyl, Trifluoromethyl, difluorochloromethyl or trichloromethyl.

R¹⁵ especially preferably represents methyl, difluoromethyl, trifluoromethyl or 1-fluoroethyl.

- R¹⁶ preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, methoxy, ethoxy, methylthio or ethylthio.
- R¹⁶ particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine or methyl.
- 5 R¹⁶ very particularly preferably represents hydrogen, fluorine, chlorine or methyl.
- R¹⁷ preferably represents hydrogen, methyl, ethyl, n-propyl, isopropyl, C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms, hydroxymethyl, hydroxyethyl, cyclopropyl, cyclopentyl, cyclohexyl or phenyl.
- 10 R¹⁷ particularly preferably represents hydrogen, methyl, ethyl, isopropyl, trifluoromethyl, difluoromethyl, hydroxymethyl, hydroxyethyl or phenyl.
- R¹⁷ very particularly preferably represents hydrogen, methyl, trifluoromethyl or phenyl.
- R¹⁷ especially preferably represents methyl.
- 15 R¹⁸ and R¹⁹ independently of one another preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R¹⁸ and R¹⁹ independently of one another particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl.
- 20 R¹⁸ and R¹⁹ independently of one another very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl or trichloromethyl.
- R¹⁸ and R¹⁹ especially preferably each represent hydrogen.
- 25 R²⁰ preferably represents fluorine, chlorine, bromine, cyano, methyl, ethyl, C₁-C₂-haloalkyl or C₁-C₂-haloalkoxy having in each case 1 to 5 fluorine, chlorine and/or bromine atoms.
- R²⁰ particularly preferably represents fluorine, chlorine, bromine, cyano, methyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy.
- 30 R²⁰ very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl or trifluoromethoxy.
- R²⁰ especially preferably represents methyl or trifluoromethyl.
- R²¹ and R²² independently of one another preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 35

R^{21} and R^{22} independently of one another particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl, difluorochloromethyl or trichloromethyl.

R^{21} and R^{22} independently of one another very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, difluoromethyl, trifluoromethyl or trichloromethyl.

R^{21} and R^{22} especially preferably each represent hydrogen.

R^{23} preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl or C_1 - C_2 -haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

R^{23} particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl or trifluoromethyl.

R^{23} very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl or trifluoromethyl.

R^{24} preferably represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C_1 - C_4 -alkyl, C_1 - C_2 -haloalkyl, C_1 - C_2 -haloalkoxy or C_1 - C_2 -haloalkylthio having in each case 1 to 5 fluorine, chlorine and/or bromine atoms.

R^{24} particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, difluoromethyl, trifluoromethyl, difluorochloromethyl, trichloromethyl, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, trifluoromethylthio, difluoromethylthio, difluorochloromethylthio or trichloromethylthio.

R^{24} very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, difluoromethyl, trifluoromethyl or trichloromethyl.

R^{24} especially preferably represents iodine, methyl, difluoromethyl or trifluoromethyl.

R^{25} preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, C_1 - C_4 -alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C_1 - C_2 -haloalkyl or C_1 - C_2 -haloalkoxy having in each case 1 to 5 fluorine, chlorine and/or bromine atoms.

R^{25} particularly preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, cyano, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy or trichloromethoxy.

- R²⁵ very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 5 R²⁶ preferably represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, C₁-C₄-alkyl, methoxy, ethoxy, methylthio, ethylthio, C₁-C₂-haloalkyl or C₁-C₂-haloalkoxy having in each case 1 to 5 fluorine, chlorine and/or bromine atoms, C₁-C₂-alkylsulphinyl or C₁-C₂-alkylsulphonyl.
- 10 R²⁶ particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl, trichloromethyl, methoxy, ethoxy, methylthio, ethylthio, trifluoromethoxy, difluoromethoxy, difluorochloromethoxy, trichloromethoxy, methylsulphinyl or methylsulphonyl.
- 15 R²⁶ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, trichloromethyl, methylsulphinyl or methylsulphonyl.
- R²⁶ especially preferably represents hydrogen.
- R²⁷ preferably represents methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 20 R²⁷ particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R²⁸ preferably represents methyl or ethyl.
- R²⁸ particularly preferably represents methyl.
- 25 Q¹ preferably represents S (sulphur), SO₂ or CH₂.
- Q¹ particularly preferably represents S (sulphur) or CH₂.
- Q¹ very particularly preferably represents S (sulphur).
- 30 p preferably represents 0 or 1.
- p particularly preferably represents 0.
- R²⁹ preferably represents methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 35 R²⁹ particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

- R²⁹ very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 5 R³⁰ preferably represents methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R³⁰ particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 10 R³⁰ very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R³¹ and R³² independently of one another preferably represent hydrogen, fluorine, chlorine, bromine, amino, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 15 R³¹ and R³² independently of one another particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R³¹ and R³² independently of one another very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R³¹ and R³² especially preferably each represent hydrogen.
- 20 R³³ preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R³³ particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 25 R³³ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R³³ especially preferably represents methyl.
- R³⁴ and R³⁵ independently of one another preferably represent hydrogen, fluorine, chlorine, bromine, amino, nitro, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 30 R³⁴ and R³⁵ independently of one another particularly preferably represent hydrogen, fluorine, chlorine, bromine, nitro, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 35 R³⁴ and R³⁵ independently of one another very particularly preferably represent hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.

R³⁴ and R³⁵ especially preferably each represent hydrogen.

- 5 R³⁶ preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R³⁶ particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R³⁶ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 10 R³⁶ especially preferably represents methyl.
- R³⁷ preferably represents hydrogen, fluorine, chlorine, bromine, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, cyano, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 15 R³⁷ particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R³⁷ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 20 R³⁷ especially preferably represents amino, methylamino, dimethylamino, methyl or trifluoromethyl.
- R³⁸ preferably represents fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 25 R³⁸ particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R³⁸ very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R³⁸ especially preferably represents methyl, trifluoromethyl or difluoromethyl.
- 30 R³⁹ preferably represents hydrogen, fluorine, chlorine, bromine, amino, C₁-C₄-alkylamino, di(C₁-C₄-alkyl)amino, cyano, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 35 R³⁹ particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, cyano, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.

- R³⁹ very particularly preferably represents hydrogen, fluorine, chlorine, bromine, amino, methylamino, dimethylamino, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R³⁹ especially preferably represents amino, methylamino, dimethylamino, methyl or trifluoromethyl.
- 5 R⁴⁰ preferably represents fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R⁴⁰ particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 10 R⁴⁰ very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R⁴⁰ especially preferably represents methyl, trifluoromethyl or difluoromethyl.
- R⁴¹ preferably represents fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 15 R⁴¹ particularly preferably represents fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- R⁴¹ very particularly preferably represents fluorine, chlorine, bromine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- 20 R⁴² preferably represents hydrogen, methyl or ethyl.
- R⁴² particularly preferably represents methyl.
- R⁴³ preferably represents fluorine, chlorine, bromine, methyl or ethyl.
- 25 R⁴³ particularly preferably represents fluorine, chlorine or methyl.
- R⁴⁴ preferably represents methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R⁴⁴ particularly preferably represents methyl, ethyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 30 R⁴⁴ very particularly preferably represents methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R⁴⁴ especially preferably represents methyl or trifluoromethyl.
- 35 R⁴⁵ preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.

- R⁴⁵ particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl or trifluoromethyl.
- 5 R⁴⁶ preferably represents fluorine, chlorine, bromine, iodine, hydroxyl, C₁-C₄-alkyl, methoxy, ethoxy, methylthio, ethylthio, difluoromethylthio, trifluoromethylthio, C₁-C₂-haloalkyl or C₁-C₂-haloalkoxy having in each case 1 to 5 fluorine, chlorine and/or bromine atoms.
- R⁴⁶ particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 10 R⁴⁶ very particularly preferably represents fluorine, chlorine, bromine, iodine, methyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R⁴⁷ preferably represents hydrogen, methyl, ethyl, C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms, C₁-C₂-alkoxy-C₁-C₂-alkyl, hydroxymethyl, hydroxyethyl, methylsulphonyl or dimethylaminosulphonyl.
- 15 R⁴⁷ particularly preferably represents hydrogen, methyl, ethyl, trifluoromethyl, methoxymethyl, ethoxymethyl, hydroxymethyl or hydroxyethyl.
- R⁴⁷ very particularly preferably represents methyl or methoxymethyl.
- 20 R⁴⁸ preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R⁴⁸ particularly preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl, trifluoromethyl, difluoromethyl or trichloromethyl.
- R⁴⁸ very particularly preferably represents hydrogen or methyl.
- 25 R⁴⁹ preferably represents hydrogen, fluorine, chlorine, bromine, iodine, cyano, methyl, ethyl, isopropyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- R⁴⁹ particularly preferably represents hydrogen, fluorine, chlorine, bromine, cyano, methyl, ethyl, isopropyl, trifluoromethyl, difluoromethyl, difluorochloromethyl or trichloromethyl.
- 30 R⁴⁹ very particularly preferably represents hydrogen, methyl, difluoromethyl or trifluoromethyl.
- R⁵⁰ preferably represents hydrogen, fluorine, chlorine, bromine, methyl, ethyl or C₁-C₂-haloalkyl having 1 to 5 fluorine, chlorine and/or bromine atoms.
- 35 R⁵⁰ particularly preferably represents hydrogen, fluorine, chlorine, bromine, iodine, methyl or trifluoromethyl.

R^{50} very particularly preferably represents hydrogen.

R^{51} preferably represents methyl, ethyl, n-propyl or isopropyl.

R^{51} particularly preferably represents methyl or ethyl.

5


Preference is given to those compounds of the formula (I) in which all radicals each have the meanings mentioned above as being preferred.

Particular preference is given to those compounds of the formula (I) in which all radicals each have the meanings mentioned above as being particularly preferred.

10

Preferred, and in each case to be understood as a sub-group of the compounds of the formula (I) mentioned above, are the following groups of novel carboxamides:


Group 1: Carboxamides of the formula (I-a)



15


in which R^1 , M, Q, Z and A are as defined above.

Group 2: Carboxamides of the formula (I-b)



20 in which R^2 , M, Q, Z and A are as defined above.


Group 3: Carboxamides of the formula (I-e)



in which R^3 , M, Q, Z and A are as defined above.


25

Group 4: Carboxamides of the formula (I-f)



in which R^4 , R^5 , M, Q, Z and A are as defined above.

Group 5: Carboxamides of the formula (I-d)



5

in which n, R^6 , M, Q, Z and A are as defined above. n preferably represents 2.

Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-1.

10 Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-2.

Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-5.

15 Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-6.

Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-9.


Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-10.

20 Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which M represents M-11.

Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which Q represents a direct bond and Z represents Z^1 .

25 Preference is given to carboxamides of the formula (I) and of groups 1 to 5 in which Q represents a direct bond and Z represents Z^4 .

Z^4 especially preferably represents the grouping




G¹ represents hydrogen or methyl,

G² represents hydrogen or methyl,

G³ represents methyl or ethyl.

Z⁴ also preferably represents one of the groupings G1 to G8



particularly preferably G3, G5 or G7.

The definition C₁-C₂₀-alkyl comprises the largest range defined here for an alkyl radical.

10 Specifically, this definition comprises the meanings methyl, ethyl, n-, isopropyl, n-, iso-, sec-, tert-butyl, and also in each case all isomeric pentyls, hexyls, heptyls, octyls, nonyls, decyls, undecyls, dodecyls, tridecyls, tetradecyls, pentadecyls, hexadecyls, heptadecyls, octadecyls, nonadecyls and eicosyls. A preferred range is C₂-C₁₂-alkyl, such as ethyl and straight-chain or branched propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl and dodecyl, particularly preferably

15 straight-chain or branched C₃-C₁₀-alkyl, such as propyl, 1-methylethyl, butyl, 1-methylpropyl, 2-methylpropyl, 1,1-dimethylethyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,2-dimethylpropyl, 1,1-dimethylpropyl, 2,2-dimethylpropyl, 1-ethylpropyl, n-hexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 1,1-dimethylbutyl, 2,2-dimethylbutyl, 3,3-dimethylbutyl, 1,1,2-trimethylpropyl,

20 1,2,2-trimethylpropyl, 1-ethylbutyl, 2-ethylbutyl, 1-ethyl-3-methylpropyl, n-heptyl, 1-methylhexyl, 1-ethylpentyl, 2-ethylpentyl, 1-propylbutyl, octyl, 1-methylheptyl, 2-methylheptyl, 1-ethylhexyl, 2-ethylhexyl, 1-propylpentyl, 2-propylpentyl, nonyl, 1-methyloctyl, 2-methyloctyl, 1-ethylheptyl, 2-ethylheptyl, 1-propylhexyl, 2-propylhexyl, decyl, 1-methylnonyl, 2-methylnonyl, 1-ethyloctyl, 2-ethyloctyl, 1-propylheptyl and 2-propylheptyl, in particular propyl, 1-methylethyl, butyl, 1-

25 methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,1-dimethylethyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, pentyl, 1-methylbutyl, 1-ethylpropyl, hexyl, 3-methylpentyl, heptyl, 1-methylhexyl, 1-ethyl-3-methylbutyl, 1-methylheptyl, 1,2-dimethylhexyl, 1,3-dimethyloctyl, 4-methyloctyl, 1,2,2,3-tetramethylbutyl, 1,3,3-trimethylbutyl, 1,2,3-trimethylbutyl, 1,3-dimethylpentyl, 1,3-dimethylhexyl, 5-methyl-3-hexyl, 2-methyl-4-heptyl, 2,6-dimethyl-4-heptyl and 1-methyl-2-

30 cyclopropylethyl.

Halogen-substituted alkyl is, for example, chloromethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl, dichlorofluoromethyl, chlorodifluoromethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl, 5 pentafluoroethyl, 3-chloro-1-methylbutyl, 2-chloro-1-methylbutyl, 1-chlorobutyl, 3,3-dichloro-1-methylbutyl, 3-chloro-1-methylbutyl, 1-methyl-3-trifluoromethylbutyl, 3-methyl-1-trifluoromethylbutyl.

The substituent $-\text{SiR}^{12}\text{R}^{13}\text{R}^{14}$ preferably represents the following radicals: SiMe_3 , SiMe_2Et , 10 $\text{SiMe}_2\text{CHMe}_2$, $\text{SiMe}_2\text{CH}_2\text{CHMe}_2$, $\text{SiMe}_2\text{CH}_2\text{CMe}_3$, $\text{SiMe}_2\text{OCHMe}_2$, $\text{SiMe}_2\text{OCH}_2\text{CHMe}_2$, SiMe_2OMe , $\text{SiMe}_2\text{CMe}_3$, $\text{SiMe}_2\text{CH}_2\text{CH}_2\text{Me}$.


The definition $\text{C}_2\text{-C}_{20}$ -alkenyl comprises the largest range defined here for an alkenyl radical. Specifically, this definition comprises the meanings ethenyl, n-, isopropenyl, n-, iso-, sec-, tert- 15 butenyl, and also in each case all isomeric pentenyls, hexenyls, heptenyls, octenyls, nonenyls, decenyls, undecenyls, dodecenyls, tridecenyls, tetradecenyls, pentadecenyls, hexadecenyls, heptadecenyls, octadecenyls, nonadecenyls and eicosenyls, 1-methyl-1-propenyl, 1-ethyl-1-butenyl, 2,4-dimethyl-1-pentenyl, 2,4-dimethyl-2-pentenyl.

20 The definition $\text{C}_2\text{-C}_{20}$ -alkynyl comprises the largest range defined here for an alkynyl radical. Specifically, this definition comprises the meanings ethynyl, n-, isopropynyl, n-, iso-, sec-, tert-butynyl, and also in each case all isomeric pentynyls, hexynyls, heptynyls, octynyls, nonynyls, decynyls, undecynyls, dodecynyls, tridecynyls, tetradecynyls, pentadecynyls, hexadecynyls, heptadecynyls, octadecynyls, nonadecynyls and eicosynyls.

25 Optionally substituted radicals may be mono- or polysubstituted, where in the case of polysubstitution, the substituents may be identical or different. Thus, the definition dialkylamino also embraces an amino group which is substituted asymmetrically by alkyl, such as, for example, methylethylamino.

30 Halogen-substituted radicals, such as, for example, haloalkyl, are mono- or polyhalogenated. In the case of polyhalogenation, the halogen atoms may be identical or different. Here, halogen represents fluorine, chlorine, bromine and iodine, in particular fluorine, chlorine and bromine.

35 The processes according to the invention can be illustrated by the general reaction scheme below:



The formula (II-a) provides a general definition of the carboxamides required as starting materials for carrying out the process (a) according to the invention and also the first step of the process (b) according to the invention. In this formula (II-a), M, Q, Z and A preferably, particularly preferably, very particularly preferably and especially preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for these radicals.

10 The carboxamides of the formula (II-a) are known and/or can be prepared by known processes (cf., for example, EP-A 0 545 099, WO 03/070705, WO 01/42223, JP-A 01-290662 and US 5,093,347).

The formula (III-a) provides a general definition of the alkylating agents furthermore required as starting materials for carrying out the process (a) according to the invention. In this formula (III-a),
 15 R^1 preferably, particularly preferably, very particularly preferably and especially preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for this radical. LG^1 represents a leaving group, preferably halogen, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate), particularly preferably chlorine,
 20 bromine, iodine, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate).

Alkylating agents of the formula (III-a) are known.

The formula (III-d) provides a general definition of the Meerwein salts furthermore required as starting materials for carrying out the process (a) according to the invention. In this formula (III-d)

5 R^1 preferably, particularly preferably, very particularly preferably and especially preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for this radical.


10 Meerwein salts of the formula (III-d) are known.

The formula (II-b) provides a general definition of the carboxamides which occur as intermediates when carrying out the process (b) according to the invention or which are required as starting materials for carrying out the process (c) according to the invention. In this formula (II-b), M, Q, Z

15 and A preferably, particularly preferably, very particularly preferably and especially preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for these radicals.

20 Some of the carboxamides of the formula (II-b) are known and/or they can be prepared by known processes (cf., for example, WO 01/42223).

Novel, and thus also part of the subject-matter of the present application, are carboxamides of the formula (II-c)



25

in which

A^1 represents A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17 and A19,

M, Q and Z are as defined above,

30 except for compounds in which

M represents M-1,

R^7 represents hydrogen,

Q represents a direct bond,

Z represents phenyl which is substituted in the 4-position by halogen,

A^1 represents A^1 ,

R^{15} represents C_1 - C_3 -haloalkyl,

R^{16} represents hydrogen,

R^{17} represents C_1 - C_3 -alkyl, C_1 - C_3 -alkoxy- C_1 - C_3 -alkyl, C_1 - C_3 -haloalkyl, C_1 - C_3 -haloalkoxy-

5 C_1 - C_3 -alkyl.

In this formula (II-c), M, Q and Z preferably, particularly preferably, very particularly preferably and especially preferably have those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being
10 preferred, particularly preferred, etc., for these radicals. A^1 preferably, particularly preferably, very particularly preferably and especially preferably has those meanings which have already been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for the radical A. The preferred ranges apply correspondingly to the excluded compounds.

15

The formula (III-b) and the formula (III-c) provide general definitions of the alkylating agents furthermore required as starting materials for carrying out the process (b) according to the invention. In these formulae (III-b) and (III-c), R^2 and R^6 preferably, particularly preferably, very particularly preferably and especially preferably have those meanings which have already been
20 mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for these radicals. LG^2 represents a leaving group, preferably halogen, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate), particularly preferably chlorine, bromine, iodine, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate). LG^6 represents a leaving group,
25 preferably halogen, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate), particularly preferably chlorine, bromine, iodine, triflate (trifluoromethylsulphonate), tosylate (p-toluenesulphonate).

Alkylating agents of the formula (III-b) and the formula (III-c) are known.

30

The formula (IV-a) and the formula (IV-b) provide general definitions of the amidating agents furthermore required as starting materials for carrying out the process (c) according to the invention. In these formulae (IV-a) and (IV-b) R^3 , R^4 and R^5 preferably, particularly preferably, very particularly preferably and especially preferably have those meanings which have already
35 been mentioned in connection with the description of the compounds of the formula (I) according to the invention as being preferred, particularly preferred, etc., for these radicals. LG^3 represents a

leaving group, preferably hydrogen, amino, hydroxyl, trialkylsilyl, particularly preferably hydrogen, trimethylsilyl.

Amidating agents of the formula (IV-a) and the formula (IV-b) are known.

5

Suitable diluents for carrying out the processes (a) and (c) according to the invention and the second step of the process (b) according to the invention are all inert organic solvents. These preferably include aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or decalin; halogenated
10 hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, tetrachloromethane, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; ketones, such as acetone, butanone, methyl isobutyl ketone or cyclohexanone; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitrile;
15 amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; mixtures thereof with water or pure water.

The processes (a) and (c) according to the invention and the second step of the process (b) according to the invention are, if appropriate, carried out in the presence of a suitable acid
20 acceptor. Suitable acid acceptors are all customary inorganic or organic bases. These preferably include alkaline earth metal or alkali metal hydrides, hydroxides, amides, alkoxides, acetates, carbonates or bicarbonates, such as, for example, sodium hydride, sodium amide, lithium diisopropylamide, sodium methoxide, sodium ethoxide, potassium tert-butoxide, sodium hydroxide, potassium hydroxide, sodium acetate, sodium carbonate, potassium carbonate,
25 potassium bicarbonate, sodium bicarbonate or ammonium carbonate, and also tertiary amines, such as trimethylamine, triethylamine, tributylamine, N,N-dimethylaniline, N,N-dimethylbenzylamine, pyridine, N-methylpiperidine, N-methylmorpholine, N,N-dimethylaminopyridine, diaza-bicyclooctane (DABCO), diazabicyclononene (DBN) or diazabicycloundecene (DBU).

30 The first step of the process (b) according to the invention is carried out in the presence of a thionating agent. Suitable thionating agents are all reagents customary for such reactions. Preference is given to using sulphur, phosphorus pentasulphide (P_2S_5 , P_4S_{10}) or Lawessons reagent.

Suitable diluents for carrying out the process (b) according to the invention are all inert organic
35 solvents. These preferably include aliphatic, alicyclic or aromatic hydrocarbons, such as, for example, petroleum ether, hexane, heptane, cyclohexane, methylcyclohexane, benzene, toluene, xylene or

decalin; halogenated hydrocarbons, such as, for example, chlorobenzene, dichlorobenzene, dichloromethane, chloroform, carbon tetrachloride, dichloroethane or trichloroethane; ethers, such as diethyl ether, diisopropyl ether, methyl t-butyl ether, methyl t-amyl ether, dioxane, tetrahydrofuran, 1,2-dimethoxyethane, 1,2-diethoxyethane or anisole; ketones, such as acetone, butanone, methyl isobutyl ketone or cyclohexanone; nitriles, such as acetonitrile, propionitrile, n- or i-butyronitrile or benzonitriles; amides, such as N,N-dimethylformamide, N,N-dimethylacetamide, N-methylformanilide, N-methylpyrrolidone or hexamethylphosphoric triamide; mixtures thereof with water or pure water.

10 Suitable oxidizing agents for carrying out the third step of the process (b) according to the invention are all oxidizing agents customarily used for oxidizing sulphur. Particularly suitable are hydrogen peroxide, organic peracids, such as, for example, peracetic acid, m-chloroperbenzoic acid, p-nitroperbenzoic acid or atmospheric oxygen.

15 Suitable diluents for carrying out the third step of the process (b) according to the invention are inert organic solvents. Preference is given to using hydrocarbons, such as, benzene, toluene, hexane or petroleum ether; chlorinated hydrocarbons, such as dichloromethane, 1,2-dichloroethane, chloroform, carbon tetrachloride or chlorobenzene; ethers, such as diethyl ether, dioxane or tetrahydrofuran; carboxylic acids, such as acetic acid or propionic acid; or dipolar aprotic solvents, such as acetonitrile, acetone, ethyl acetate or dimethylformamide.

If appropriate, the third step of the process (b) according to the invention may be carried out in the presence of an acid binder. Suitable acid binders are all organic and inorganic acid binders customarily used. Preference is given to using alkaline earth metal or alkali metal hydroxides, acetates or carbonates, such as, for example, calcium hydroxide, sodium hydroxide, sodium acetate or sodium carbonate.

If appropriate, the third step of the process (b) according to the invention can be carried out in the presence of a suitable catalyst. Suitable catalysts are all metal salt catalysts customarily used for such sulphur oxidations. In this context, ammonium molybdate and sodium tungstate may be mentioned by way of example.

When carrying out the processes (a) and (c) according to the invention and the second step of the process (b) according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, the reaction is carried out at temperatures of from 0°C to 150°C, preferably at temperatures of from 20°C to 110°C.

When carrying out the third step of the process (b) according to the invention, the reaction temperatures can be varied within a relatively wide range. In general, the reaction is carried out at temperatures between -20°C and $+70^{\circ}\text{C}$, preferably at temperatures between 0°C and $+50^{\circ}\text{C}$.

5

For carrying out the process (a) according to the invention for preparing compounds of the formula (I-a), in general from 0.8 to 5 mol, preferably from 1 to 3 mol, of alkylating agent of the formula (III-a) or from 0.1 to 1.5 mol, preferably from 0.25 to 1 mol, of Meerwein salt of the formula (III-d) are employed per mole of carboxamide of the formula (II-a).

10

For carrying out the first step of the process (b) according to the invention for preparing the compounds of the formula (II-b), in general from 0.8 to 1.2 mol, preferably equimolar amounts, of thionating agent are employed per mole of carboxamide of the formula (II-a).

15

For carrying out the second step of the process (b) according to the invention for preparing the compounds of the formula (I-b) or the formula (I-c), in general from 0.8 to 5 mol, preferably from 1 to 3 mol, of alkylating agent of the formula (III-b) or the formula (III-c) are employed per mole of carboxamide of the formula (II-b).

20

For carrying out the third step of the process (b) according to the invention for preparing the compounds of the formula (I-d), in general from 0.8 to 1.2 mol, preferably equimolar amounts, of oxidizing agent are employed per mole of carboxamide of the formula (I-c) if the oxidation of the sulphur is to be interrupted at the sulfoxide stage. For oxidation to the sulphone, in general from 1.8 to 3.0 mol, preferably double molar amounts, of oxidizing agent are employed per mole of carboxamide of the formula (I-c).

25

For carrying out the process (c) according to the invention for preparing the compounds of the formula (I-e) or the formula (I-f), in general from 0.8 to 5 mol, preferably from 1 to 3 mol, of amidating agent of the formula (IV-a) or the formula (IV-b) are employed per mole of carboxamide of the formula (II-b).

30

The processes according to the invention are generally carried out under atmospheric pressure. However, it is also possible to operate under elevated or reduced pressure – in general between 0.1 bar and 10 bar.

35

The compounds according to the invention exhibit a potent microbicidal activity and can be employed in crop protection and in the protection of materials for controlling undesirable microorganisms such as fungi and bacteria.

- 5 Fungicides can be employed in crop protection for controlling Plasmodiophoromycetes, Oomycetes, Chytridiomycetes, Zygomycetes, Ascomycetes, Basidiomycetes and Deuteromycetes.

Bactericides can be employed in crop protection for combating Pseudomonadaceae, Rhizobiaceae, Enterobacteriaceae, Corynebacteriaceae and Streptomycetaceae.

10

Examples which may be mentioned, but not by limitation, of some pathogens of fungal and bacterial diseases which come under the abovementioned general terms are:

diseases caused by powdery mildew pathogens, such as, for example

Blumeria species such as, for example, Blumeria graminis;

- 15 Podosphaera species such as, for example, Podosphaera leucotricha;

Sphaerotheca species such as, for example, Sphaerotheca fuliginea;

Uncinula species such as, for example, Uncinula necator;

diseases caused by rust pathogens such as, for example,

- 20 Gymnosporangium species such as, for example, Gymnosporangium sabinae

Hemileia species such as, for example, Hemileia vastatrix;

Phakopsora species such as, for example, Phakopsora pachyrhizi and Phakopsora meibomia;

Puccinia species such as, for example, Puccinia recondita or Puccinia graminis;

Uromyces species such as, for example, Uromyces appendiculatus;

25

diseases caused by pathogens from the Oomycetes group such as, for example,

Bremia species such as, for example, Bremia lactucae;

Peronospora species such as, for example, Peronospora pisi or P. brassicae;

Phytophthora species such as, for example, Phytophthora infestans;

- 30 Plasmopara species such as, for example, Plasmopara viticola;

Pseudoperonospora species such as, for example, Pseudoperonospora humuli or

Pseudoperonospora cubensis;

Pythium species such as, for example, Pythium ultimum;

- 35 leaf spot diseases and leaf wilts caused by, for example,

Alternaria species such as, for example, Alternaria solani;

- Cercospora species such as, for example, *Cercospora beticola*;
Cladosporium species such as, for example, *Cladosporium cucumerinum*;
Cochliobolus species such as, for example, *Cochliobolus sativus*
(conidial form: *Drechslera*, syn: *Helminthosporium*);
- 5 *Colletotrichum* species such as, for example, *Colletotrichum lindemuthianum*;
Cycloconium species such as, for example, *Cycloconium oleaginum*;
Diaporthe species such as, for example, *Diaporthe citri*;
Elsinoe species such as, for example, *Elsinoe fawcettii*;
Gloeosporium species such as, for example, *Gloeosporium laeticolor*;
- 10 *Glomerella* species such as, for example, *Glomerella cingulata*;
Guignardia species such as, for example, *Guignardia bidwelli*;
Leptosphaeria species such as, for example, *Leptosphaeria maculans*;
Magnaporthe species such as, for example, *Magnaporthe grisea*;
Mycosphaerella species such as, for example, *Mycosphaerella graminicola* and *Mycosphaerella*
15 *fijiensis*;
Phaeosphaeria species such as, for example, *Phaeosphaeria nodorum*;
Pyrenophora species such as, for example, *Pyrenophora teres*;
Ramularia species such as, for example, *Ramularia collo-cygni*;
Rhynchosporium species such as, for example, *Rhynchosporium secalis*;
- 20 *Septoria* species such as, for example, *Septoria apii*;
Typhula species such as, for example, *Typhula incarnata*;
Venturia species such as, for example, *Venturia inaequalis*;
- root and stem diseases caused by, for example,
- 25 *Corticium* species such as, for example, *Corticium graminearum*;
Fusarium species such as, for example, *Fusarium oxysporum*;
Gaeumannomyces species such as, for example, *Gaeumannomyces graminis*;
Rhizoctonia species such as, for example, *Rhizoctonia solani*;
Tapesia species such as, for example, *Tapesia acuformis* or *Tapesia yallundae*;
- 30 *Thielaviopsis* species such as, for example, *Thielaviopsis basicola*;
- ear and panicle diseases (including maize cobs), caused by, for example,
- Alternaria* species such as, for example, *Alternaria* spp.;
- Aspergillus* species such as, for example, *Aspergillus flavus*;
- 35 *Cladosporium* species such as, for example, *Cladosporium cladosporioides*.;
- Claviceps* species such as, for example, *Claviceps purpurea*;

Fusarium species such as, for example, *Fusarium culmorum*;
Gibberella species such as, for example, *Gibberella zeae*;
Monographella species such as, for example, *Monographella nivalis*;

- 5 diseases caused by smuts such as, for example,
Sphacelotheca species such as, for example, *Sphacelotheca reiliana*;
Tilletia species such as, for example, *Tilletia caries*;
Urocystis species such as, for example, *Urocystis occulta*;
Ustilago species such as, for example, *Ustilago nuda*;
10 fruit rots caused by, for example,
Aspergillus species such as, for example, *Aspergillus flavus*;
Botrytis species such as, for example, *Botrytis cinerea*;
Penicillium species such as, for example, *Penicillium expansum* and *Penicillium purpurogenum*;
15 Sclerotinia species such as, for example, *Sclerotinia sclerotiorum*;
Verticillium species such as, for example, *Verticillium alboatrum*;

- seed- and soil-borne rot and wilts, and seedling diseases, caused by, for example,
Fusarium species such as, for example, *Fusarium culmorum*;
20 Phytophthora species such as, for example, *Phytophthora cactorum*;
Pythium species such as, for example, *Pythium ultimum*;
Rhizoctonia species such as, for example, *Rhizoctonia solani*;
Sclerotium species such as, for example, *Sclerotium rolfsii*;

- 25 cancers, galls and witches' broom disease, caused by, for example,
Nectria species such as, for example, *Nectria galligena*;

- wilts caused by, for example,
Monilinia species such as, for example, *Monilinia laxa*;

- 30 deformations of leaves, flowers and fruits, caused by, for example,
Taphrina species such as, for example, *Taphrina deformans*;

- degenerative diseases of woody species, caused by, for example,
35 Esca species such as, for example, *Phaeoconiella clamydospora* and *Phaeoacremonium aleophilum* and *Fomitiporia mediterranea*;

diseases of flowers and seeds, caused by, for example,

Botrytis species such as, for example, Botrytis cinerea;

diseases of the plant tubers, caused by, for example,

5 Rhizoctonia species such as, for example, Rhizoctonia solani;

Helminthosporium species such as, for example, Helminthosporium solani;

diseases caused by bacterial pathogens such as, for example,

Xanthomonas species such as, for example, Xanthomonas campestris pv. oryzae;

10 Pseudomonas species such as, for example, Pseudomonas syringae pv. lachrymans;

Erwinia species such as, for example, Erwinia amylovora.

The following diseases of soybeans can preferably be controlled:

Fungal diseases on leaves, stems, pods and seeds caused by, for example,

15 alternaria leaf spot (Alternaria spec. atrans tenuissima), anthracnose (Colletotrichum gloeosporoides dematium var. truncatum), brown spot (Septoria glycines), cercospora leaf spot and blight (Cercospora kikuchii), choanephora leaf blight (Choanephora infundibulifera trispora (syn.)), dactuliophora leaf spot (Dactuliophora glycines), downy mildew (Peronospora manshurica), drechslera blight (Drechslera glycini), frog-eye leaf spot (Cercospora sojae),
20 leptosphaerulina leaf spot (Leptosphaerulina trifolii), phyllosticta leaf spot (Phyllosticta sojaecola), pod and stem blight (Phomopsis sojae) powdery mildew (Microsphaera diffusa), pyrenochaeta leaf spot (Pyrenochaeta glycines), rhizoctonia aerial, foliage, and web blight (Rhizoctonia solani), rust (Phakopsora pachyrhizi), scab (Sphaceloma glycines), stemphylium leaf blight (Stemphylium botryosum), target spot (Corynespora cassiicola);

25

fungal diseases on roots and the stem base caused by, for example,

black root rot (Calonectria crotalariae), charcoal rot (Macrophomina phaseolina), fusarium blight or wilt, root rot, and pod and collar rot (Fusarium oxysporum, Fusarium orthoceras, Fusarium semitectum, Fusarium equiseti), mycoleptodiscus root rot (Mycoleptodiscus terrestris), neocosmospora (Neocosmospora vasinfecta), pod and stem blight (Diaporthe phaseolorum), stem canker (Diaporthe phaseolorum var. caulivora), phytophthora rot (Phytophthora megasperma), brown stem rot (Phialophora gregata), pythium rot (Pythium aphanidermatum, Pythium irregulare, Pythium debaryanum, Pythium myriotylum, Pythium ultimum), rhizoctonia root rot, stem decay, and damping-off (Rhizoctonia solani), sclerotinia stem decay (Sclerotinia sclerotiorum), sclerotinia
35 southern blight (Sclerotinia rolfii), thielaviopsis root rot (Thielaviopsis basicola).

The active compounds according to the invention also have a potent strengthening effect in plants. They are therefore suitable for mobilizing the plants' defences against attack by undesired microorganisms.

5 Plant-strengthening (resistance-inducing) compounds are understood as meaning, in the present context, those substances which are capable of stimulating the defence system of plants in such a way that, when subsequently inoculated with undesired microorganisms, the treated plants display a substantial degree of resistance to these microorganisms.

10 In the present case, undesired microorganisms are understood as meaning phytopathogenic fungi, bacteria and viruses. Thus, the compounds according to the invention can be employed for protecting plants against attack by the abovementioned pathogens within a certain period of time after the treatment. The period of time within which their protection is effected is generally extended from 1 to 28 days, preferably from 1 to 14 days, particularly preferably from 1 to 7 days,
15 after the plants have been treated with the active compounds.

The fact that the active compounds, at the concentrations required for the controlling of plant diseases, are well tolerated by plants permits the treatment of aerial plant parts, of vegetative propagation material and seed, and of the soil.

20

In this context, the active compounds according to the invention can be employed particularly successfully for controlling cereal diseases such as, for example, against Puccinia species and of diseases in viticulture, fruit production and vegetable production such as, for example against Botrytis, Venturia or Alternaria species.

25

The active compounds according to the invention are also suitable for increasing the yield. Moreover, they display a low degree of toxicity and are well tolerated by plants.

If appropriate, the active compounds according to the invention can also be used in certain
30 concentrations and application rates as herbicides, for influencing plant growth and for controlling animal pests. If appropriate, they can also be employed as intermediates and precursors for the synthesis of further active compounds.

All plants and plant parts can be treated in accordance with the invention. Plants are understood as
35 meaning, in the present context, all plants and plant populations, such as desired and undesired wild plants or crop plants (including naturally occurring crop plants). Crop plants may be plants

which can be obtained by conventional breeding and optimization methods or else by biotechnological and genetic engineering methods or by combinations of these methods, including the transgenic plants and including the plant varieties capable or not capable of being protected by Plant Breeders' rights. Plant parts are understood as meaning all aerial and subterranean parts and
5 organs of the plants, such as shoot, leaf, flower and root, examples which may be mentioned being leaves, needles, stalks, stems, flowers, fruiting bodies, fruits and seeds, and also roots, tubers and rhizomes. The plant parts also include harvested material and vegetative and generative propagation material, for example cuttings, tubers, rhizomes, slips and seeds.

10 The treatment according to the invention with the active compounds, of the plants and plant parts, is carried out directly or by acting on their environment, habitat, or store by the customary treatment methods, for example by immersion, spraying, vaporizing, fogging, broadcasting, painting on and, in the case of propagation material, in particular in the case of seeds, furthermore by coating with one or more coats.

15

In the protection of materials, the substances according to the invention can be employed for protecting industrial materials against attack and destruction by undesired microorganisms.

In the present context, industrial materials are understood as meaning non live materials which
20 have been made for use in technology. For example, industrial materials which are to be protected by active compounds according to the invention from microbial modification or destruction can be glues, sizes, paper and board, textiles, leather, timber, paints and plastic articles, cooling lubricants and other materials which are capable of being attacked or destroyed by microorganisms. Parts of production plants, for example cooling-water circuits, which can be adversely affected by the
25 multiplication of microorganisms may also be mentioned within the materials to be protected. Industrial materials which may be mentioned with preference for the purposes of the present invention are glues, sizes, paper and board, leather, timber, paints, cooling lubricants and heat-transfer fluids, especially preferably wood.

30 Microorganisms which are capable of bringing about a degradation or modification of the industrial materials and which may be mentioned are, for example, bacteria, fungi, yeasts, algae and slime organisms. The active compounds according to the invention are preferably active against fungi, in particular moulds, wood-discolouring and wood-destroying fungi (Basidiomycetes) and against slime organisms and algae.

35

Examples which may be mentioned are microorganisms of the following genera:

- Alternaria such as *Alternaria tenuis*,
Aspergillus such as *Aspergillus niger*,
Chaetomium such as *Chaetomium globosum*,
Coniophora such as *Coniophora puetana*,
5 *Lentinus* such as *Lentinus tigrinus*,
Penicillium such as *Penicillium glaucum*,
Polyporus such as *Polyporus versicolor*,
Aureobasidium such as *Aureobasidium pullulans*,
Sclerophoma such as *Sclerophoma pityophila*,
10 *Trichoderma* such as *Trichoderma viride*,
Escherichia such as *Escherichia coli*,
Pseudomonas such as *Pseudomonas aeruginosa*,
Staphylococcus such as *Staphylococcus aureus*.
- 15 Depending on their respective physical and/or chemical properties, the active compounds can be converted to the customary formulations, such as solutions, emulsions, suspensions, powders, foams, pastes, granules, aerosols, very fine capsules in polymeric substances and in coating compositions for seed, and also ULV cold- and warm-fogging formulations.

These formulations are produced in a known manner, for example by mixing the active compounds
20 with extenders, that is, liquid solvents, pressurized liquefied gases and/or solid carriers, optionally with the use of surface-active agents, that is emulsifiers and/or dispersants, and/or foam formers. If the extender used is water, it is also possible to employ for example organic solvents as cosolvents. Suitable liquid solvents are essentially: aromatics, such as xylene, toluene or alkyl-naphthalenes, chlorinated aromatics or chlorinated aliphatic hydrocarbons, such as chlorobenzenes, chloroethylenes
25 or methylene chloride, aliphatic hydrocarbons, such as cyclohexane or paraffins, for example mineral oil fractions, alcohols, such as butanol or glycol as well as their ethers and esters, ketones, such as acetone, methyl ethyl ketone, methyl isobutyl ketone or cyclohexanone, strongly polar solvents, such as dimethylformamide and dimethyl sulphoxide, and also water. Liquefied gaseous extenders or carriers are those liquids which are gaseous at ambient temperature and at atmospheric pressure, for
30 example aerosol propellants such as halogenated hydrocarbons and also butane, propane, nitrogen and carbon dioxide. As solid carriers there are suitable: for example ground natural minerals, such as kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth, and ground synthetic minerals, such as finely divided silica, alumina and silicates. As solid carriers for granules there are suitable: for example crushed and fractionated natural rocks such as calcite, pumice, marble,
35 sepiolite and dolomite, and also synthetic granules of inorganic and organic meals, and granules of

organic material such as sawdust, coconut shells, maize cobs and tobacco stalks. As emulsifiers and/or foam formers there are suitable: for example non-ionic and anionic emulsifiers, such as polyoxyethylene fatty acid esters, polyoxyethylene fatty alcohol ethers, for example alkylaryl polyglycol ethers, alkylsulphonates, alkyl sulphates, arylsulphonates and protein hydrolysates. As
5 dispersants there are suitable: for example lignosulphite waste liquors and methylcellulose.

Tackifiers such as carboxymethylcellulose and natural and synthetic polymers in the form of powders, granules or latices, such as gum arabic, polyvinyl alcohol and polyvinyl acetate, as well as natural phospholipids, such as cephalins and lecithins, and synthetic phospholipids, can be used in the
10 formulations. Other possible additives are mineral and vegetable oils.

It is possible to use colorants such as inorganic pigments, for example iron oxide, titanium oxide and Prussian Blue, and organic dyestuffs, such as alizarin dyestuffs, azo dyestuffs and metal phthalocyanine dyestuffs, and trace nutrients such as salts of iron, manganese, boron, copper, cobalt,
15 molybdenum and zinc.

The formulations in general contain between 0.1 and 95% by weight of active compound, preferably between 0.5 and 90%.

20 The active compounds according to the invention, as such or in their formulations, can also be used as a mixture with known fungicides, bactericides, acaricides, nematocides, or insecticides, for example, to improve the activity spectrum or prevent the development of resistance. In many instances, synergistic effects are obtained, i.e. the activity of the mixture exceeds the activity of the individual components.

25

Examples of co-components in mixtures are the following compounds:

Fungicides:

1) Nucleic acid synthesis inhibitors: for example benalaxyl, benalaxyl-M, bupirimate, clozylacon,
30 dimethirimol, ethirimol, furalaxyl, hymexazol, mefenoxam, metalaxyl, metalaxyl-M, ofurace, oxadixyl, oxolinic acid;

2) mitosis and cell division inhibitors: for example benomyl, carbendazim, diethofencarb, ethaboxam, fuberidazole, pencycuron, thiabendazole, thiophanate-methyl, zoxamide;

3) respiration inhibitors (inhibitors of the respiratory chain):

35 3.1) inhibitors which act on complex I of the respiratory chain: for example diflumetorim;

- 3.2) inhibitors which act on complex II of the respiratory chain: for example boscalid/nicobifen, carboxin, fenfuram, flutolanil, furametpyr, furmecyclox, mepronil, oxycarboxin, penthiopyrad, thifluzamide;
- 3.3) inhibitors which act on complex III of the respiratory chain: for example amisulbrom, azoxystrobin, cyazofamid, dimoxystrobin, enestrobin, famoxadone, fenamidone, fluoxastrobin, kresoxim-methyl, metominostrobin, orysastrobin, picoxystrobin, pyraclostrobin, trifloxystrobin;
- 4) decouplers: for example dinocap, fluazinam, meptyldinocap;
- 5) ATP production inhibitors: for example fentin acetate, fentin chloride, fentin hydroxide, silthiofam;
- 6) amino acid and protein biosynthesis inhibitors: for example andoprím, blasticidin-S, cyprodinil, kasugamycin, kasugamycin hydrochloride hydrate, mepanipirim, pyrimethanil;
- 7) signal transduction inhibitors: for example fenpiclonil, fludioxonil, quinoxifen;
- 8) lipid and membrane synthesis inhibitors: for example biphenyl, chlozolate, edifenphos, iodocarb, iprobenfos, iprodione, isoprothiolane, procymidone, propamocarb, propamocarb hydrochloride, pyrazophos, tolclofos-methyl, vinclozolin;
- 9) inhibitors of ergosterol biosynthesis: for example aldimorph, azaconazole, bitertanol, bromuconazole, cyproconazole, diclobutrazole, difenoconazole, diniconazole, diniconazole-M, dodemorph, dodemorph acetate, epoxiconazole, etaconazole, fenarimol, fenbuconazole, fenhexamid, fenpropidin, fenpropimorph, fluquinconazole, flurprimidol, flusilazole, flutriafol, furconazole, furconazole-cis, hexaconazole, imazalil, imazalil sulfate, imibenconazole, ipconazole, metconazole, myclobutanil, naftifine, nuarimol, oxpoconazole, paclobutrazol, pefurazoate, penconazole, prochloraz, propiconazole, prothioconazole, pyributicarb, pyrifenox, simeconazole, spiroxamine, tebuconazole, terbinafine, tetraconazole, triadimefon, triadimenol, tridemorph, triflumizole, triforine, triticonazole, uniconazole, viniconazole, voriconazole;
- 10) cell wall synthesis inhibitors: for example benthiavalicarb, dimethomorph, flumorph, iprovalicarb, polyoxins, polyoxorim, validamycin A;
- 11) melanin biosynthesis inhibitors: for example carpropamid, diclocymet, fenoxanil, phthalide, pyroquilon, tricyclazole;
- 12) resistance inductors: for example acibenzolar-S-methyl, probenazole, tiadinil;
- 13) compounds with multi-site activity: for example Bordeaux mixture, captafol, captan, chlorothalonil, copper naphthenate, copper oxide, copper oxychloride, copper preparations such as, for example, copper hydroxide, copper sulphate, dichlofluanid, dithianon, dodine, dodine free base, ferbam, fluorofolpet, folpet, guazatine, guazatine acetate, iminoctadine, iminoctadine albesilate, iminoctadine triacetate, mancozeb, maneb, metiram, metiram zinc, oxine-copper, propineb, sulphur and sulphur preparations such as, for example, calcium polysulphide, thiram, tolylfluanid, zineb, ziram;

14) a compound selected from the following enumeration: N-methyl-(2E)-2-(2-{[6-(3-chloro-2-methylphenoxy)-5-fluoropyrimidin-4-yl]oxy}phenyl)-2-(methoxyimino)acetamide, N-methyl-(2E)-2-{2-[(1E)-1-(3-{[(E)-1-fluoro-2-phenylvinyl]oxy}phenyl)ethylidene]amino}oxy)methyl]phenyl}-2-(methoxyimino)acetamide, 1-(4-chlorophenyl)-2-(1H-1,2,4-triazol-1-yl)cycloheptanol, 1-[(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl-1H-imidazole-1-carboxylate, 2-(4-chlorophenyl)-N-{2-[3-methoxy-4-(prop-2-yn-1-yloxy)phenyl]ethyl}-2-(prop-2-yn-1-yloxy)acetamide, 2,3,5,6-tetrachloro-4-(methylsulphonyl)pyridine, 2-butoxy-6-iodo-3-propyl-4H-chromen-4-one, 2-chloro-N-(1,1,3-trimethyl-2,3-dihydro-1H-inden-4-yl)nicotinamide, 2-phenylphenol and salts thereof, 3,4,5-trichloropyridine-2,6-dicarbonitrile, 3,4-dichloro-N-(2-cyanophenyl)isothiazole-5-carboxamide, 3-[5-(4-chlorophenyl)-2,3-dimethylisoxazolidin-3-yl]pyridine, 5-chloro-6-(2,4,6-trifluorophenyl)-N-[(1R)-1,2,2-trimethylpropyl][1,2,4]triazolo[1,5-a]pyrimidine-7-amine, 5-chloro-7-(4-methylpiperidin-1-yl)-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine, 5-chloro-N-[(1R)-1,2-dimethylpropyl]-6-(2,4,6-trifluorophenyl)[1,2,4]triazolo[1,5-a]pyrimidine-7-amine, 8-hydroxyquinoline sulphate, benthiazole, bethoxazin, capsimycin, carvone, quinomethionate, cufraneb, cyflufenamid, cymoxanil, dazomet, debacarb, dichlorophen, diclomezine, dicloran, difenzoquat, difenzoquat methylsulphate, diphenylamine, ferimzone, flumetover, fluopicolide, fluoroimide, flusulfamide, fosetyl-aluminium, fosetyl-calcium, fosetyl-sodium, hexachlorobenzene, irumamycin, methasulfocarb, methyl (2-chloro-5-{(1E)-N-[(6-methylpyridin-2-yl)methoxy]ethanimidoyl}benzyl)carbamate, methyl (2E)-2-{2-[(cyclopropyl[(4-methoxyphenyl)imino]methyl}thio)methyl]phenyl}-3-methoxyacrylate, methyl 1-(2,2-dimethyl-2,3-dihydro-1H-inden-1-yl)-1H-imidazole-5-carboxylate, methyl 3-(4-chlorophenyl)-3-{[N-(isopropoxycarbonyl)valyl]amino}propanoate, methyl isothiocyanate, metrafenone, mildiomycin, N-(3',4'-dichloro-5-fluorobiphenyl-2-yl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide, N-(3-ethyl-3,5,5-trimethylcyclohexyl)-3-(formylamino)-2-hydroxybenzamide, N-(4-chloro-2-nitrophenyl)-N-ethyl-4-methylbenzenesulphonamide, N-[(5-bromo-3-chloropyridin-2-yl)methyl]-2,4-dichloronicotinamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2,4-dichloronicotinamide, N-[1-(5-bromo-3-chloropyridin-2-yl)ethyl]-2-fluoro-4-iodonicotinamide, N-[2-(4-{3-(4-chlorophenyl)prop-2-yn-1-yl]oxy}-3-methoxyphenyl)ethyl]-N²-(methylsulphonyl)valinamide, N-{(Z)-[(cyclopropylmethoxy)imino][6-(difluoromethoxy)-2,3-difluorophenyl]methyl}-2-phenylacetamide, N-{2-[3-chloro-5-(trifluoromethyl)pyridin-2-yl]ethyl}-2-(trifluoromethyl)benzamide, natamycin, nickel dimethyl dithiocarbamate, nitrothal-isopropyl, O-{1-[(4-methoxyphenoxy)methyl]-2,2-dimethylpropyl} 1H-imidazole-1-carbothioate, octhilinone, oxamocarb, oxyfentiin, pentachlorophenol and salts, phosphoric acid and its salts, piperalin, propamocarb fosetilate, propanosine-sodium, proquinazid, pyrrolnitrine, quintozone, tecloftalam, tecnazene, triazoxide, trichlamide, zarilamid.

Bactericides:

Bronopol, dichlorophen, nitrapyrin, nickel dimethyl dithiocarbamate, kasugamycin, othilinone, furancarboxylic acid, oxytetracyclin, probenazole, streptomycin, tecloftalam, copper sulphate and other copper preparations.

5

Insecticides/Acaricides/Nematicides:

1. Acetylcholine esterase (AChE) inhibitors

1.1 Carbamates (for example alanycarb, aldicarb, aldoxycarb, allyxycarb, aminocarb, azamethi-
phos, bendiocarb, benfuracarb, bufencarb, butacarb, butocarboxim, butoxycarboxim, carbaryl,
10 carbofuran, carbosulfan, chloethocarb, coumaphos, cyanofenphos, cyanophos, dimetilan,
ethiofencarb, fenobucarb, fenothiocarb, formetanate, furathiocarb, isoprocarb, metam-sodium,
methiocarb, methomyl, metolcarb, oxamyl, pirimicarb, promecarb, propoxur, thiodicarb, thiofanox,
triazamate, trimethacarb, XMC, xylylcarb)

1.2 Organophosphates (for example acephate, azamethiphos, azinphos (-methyl, -ethyl),
15 bromophos-ethyl, bromfenvinfos (-methyl), butathiofos, cadusafos, carbophenothion,
chlorethoxyfos, chlorfenvinphos, chlormephos, chlorpyrifos (-methyl/-ethyl), coumaphos,
cyanofenphos, cyanophos, chlorfenvinphos, demeton-S-methyl, demeton-S-methylsulphon,
dialifos, diazinon, dichlofenthion, dichlorvos/DDVP, dicrotophos, dimethoate, dimethylvinphos,
dioxabenzofos, disulfoton, EPN, ethion, ethoprophos, etrimfos, famphur, fenamiphos, fenitrothion,
20 fensulfothion, fenthion, flupyrzofos, fonofos, formothion, fosmethilan, fosthiazate, heptenophos,
iodofenphos, iprobenfos, isazofos, isofenphos, isopropyl O-salicylate, isoxathion, malathion,
mecarbam, methacrifos, methamidophos, methidathion, mevinphos, monocrotophos, naled,
omethoate, oxydemeton-methyl, parathion (-methyl/-ethyl), phenthoate, phorate, phosalone,
phosmet, phosphamidon, phosphocarb, phoxim, pirimiphos (-methyl/-ethyl), profenofos,
25 propaphos, propetamphos, prothiofos, prothoate, pyraclofos, pyridaphenthion, pyridathion,
quinalphos, sebufos, sulfotep, sulprofos, tebupirimfos, temephos, terbufos, tetrachlorvinphos,
thiometon, triazophos, trichlorfon, vamidothion)

2. Sodium channel modulators/voltage-dependent sodium channel blockers

2.1 Pyrethroids (for example acrinathrin, allethrin (d-cis-trans, d-trans), beta-cyfluthrin, bifenthrin,
30 bioallethrin, bioallethrin-S-cyclopentyl isomer, bioethanomethrin, biopermethrin, bioresmethrin,
chlovaporthrin, cis-cypermethrin, cis-resmethrin, cis-permethrin, clocythrin, cycloprothrin, cyflu-
thrin, cyhalothrin, cypermethrin (alpha-, beta-, theta-, zeta-), cyphenothrin, DDT, deltamethrin,
empenthrin (1R isomer), esfenvalerate, etofenprox, fenfluthrin, fenpropathrin, fenpyrithrin, fen-
valerate, flubrocycythrinate, flucythrinate, flufenprox, flumethrin, fluvalinate, fubfenprox, gamma-
35 cyhalothrin, imiprothrin, kadethrin, lambda-cyhalothrin, metofluthrin, permethrin (cis-, trans-),
phenothrin (1R trans-isomer), prallethrin, profluthrin, protrifenbute, pyresmethrin, resmethrin, RU

- 15525, silafluofen, tau-fluvalinate, tefluthrin, terallethrin, tetramethrin (1R isomer), tralomethrin, transfluthrin, ZXI 8901, pyrethrins (pyrethrum))
- 2.2 Oxadiazines (for example indoxacarb)
3. *Acetylcholine receptor agonists/antagonists*
- 5 3.1 Chloronicotinylns/neonicotinoids (for example acetamiprid, clothianidin, dinotefuran, imidacloprid, nitenpyram, nithiazine, thiacloprid, thiamethoxam)
- 3.2 Nicotine, bensultap, cartap
4. *Acetylcholine receptor modulators*
- 4.1 Spinosyns (for example spinosad)
- 10 5. *GABA-controlled chloride channel antagonists*
- 5.1 Cyclodiene organochlorines (for example camphechlor, chlordane, endosulfan, gamma-HCH, HCH, heptachlor, lindane, methoxychlor)
- 5.2 Fiproles (for example acetoprole, ethiprole, fipronil, vaniliprole)
6. *Chloride channel activators*
- 15 6.1 Mectins (for example abamectin, avermectin, emamectin, emamectin benzoate, ivermectin, milbemectin, milbemycin)
7. *Juvenile hormone mimetics*
- (for example diofenolan, epofenonane, fenoxycarb, hydroprene, kinoprene, methoprene, pyriproxifen, triprene)
- 20 8. *Ecdysone agonists/disruptors*
- 8.1 Diacylhydrazines (for example chromafenozide, halofenozide, methoxyfenozide, tebufenozide)
9. *Chitin biosynthesis inhibitors*
- 9.1 Benzoylureas (for example bistrifluron, chlofluaazuron, diflubenzuron, fluazuron, flucycloxuron, flufenoxuron, hexaflumuron, lufenuron, novaluron, noviflumuron, penfluron, teflubenzuron, triflumuron)
- 25 9.2 Buprofezin
- 9.3 Cyromazine
10. *Inhibitors of oxidative phosphorylation, ATP disruptors*
- 10.1 Diafenthiuron
- 30 10.2 Organotins (for example azocyclotin, cyhexatin, fenbutatin oxide)
11. *Uncouplers of oxidative phosphorylation by interrupting the H-proton gradient*
- 11.1 Pyrroles (for example chlorfenapyr)
- 11.2 Dinitrophenols (for example binapacyrl, dinobuton, dinocap, DNOC)
12. *Site-I electron transport inhibitors*
- 35 12.1 METIs (for example fenazaquin, fenpyroximate, pyrimidifen, pyridaben, tebufenpyrad, tolfenpyrad)

- 12.2 Hydramethylnone
- 12.3 Dicofol
- 13. *Site-II electron transport inhibitors*
- 13.1 Rotenone
- 5 14. *Site-III electron transport inhibitors*
- 14.1 Acequinocyl, fluacrypyrim
- 15. *Microbial disruptors of the insect gut membrane*
- Bacillus thuringiensis strains
- 16. *Fat synthesis inhibitors*
- 10 16.1 Tetronic acids (for example spiroadiclofen, spiromesifen)
- 16.2 Tetramic acids [for example 3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl carbonate (also known as: carbonic acid, 3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl ester, CAS Reg. No.: 382608-10-8) and carbonic acid, cis-3-(2,5-dimethylphenyl)-8-methoxy-2-oxo-1-azaspiro[4.5]dec-3-en-4-yl ethyl ester (CAS Reg. No.: 203313-25-1)]
- 15 17. *Carboxamides*
- (for example flonicamid)
- 18. *Octopaminergic agonists*
- (for example amitraz)
- 20 19. *Inhibitors of magnesium-stimulated ATPase*
- (for example propargite)
- 20. *Ryanodin receptor agonists,*
- 20.1 Benzoic acid dicarboxamides
- [for example N²-[1,1-dimethyl-2-(methylsulfonyl)ethyl]-3-iodo-N¹-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide (CAS Reg. No.: 272451-65-7), flubendiamide]
- 25 20.2 Anthranilamides (for example DPX E2Y45 = 3-bromo-N-{4-chloro-2-methyl-6-[(methylamino)-carbonyl]phenyl}-1-(3-chloropyridin-2-yl)-1H-pyrazole-5-carboxamide)
- 21. *Nereistoxin analogues*
- 30 (for example thiocyclam hydrogen oxalate, thiosultap sodium)
- 22. *Biologicals, hormones or pheromones*
- (for example azadirachtin, Bacillus spec., Beauveria spec., codlemone, Metarrhizium spec., Paecilomyces spec., thuringiensin, Verticillium spec.)
- 23. *Active compounds with unknown or unspecific mechanisms of action*
- 35 23.1 Fumigants (for example aluminium phosphide, methyl bromide, sulphuryl fluoride)
- 23.2 Selective antifeedants (for example cryolite, flonicamid, pymetrozine)

23.3 Mite growth inhibitors (for example clofentezine, etoxazole, hexythiazox)

23.4 Amidoflumet, benclothiaz, benzoximate, bifenazate, bromopropylate, buprofezin, quino-
methionate, chlordimeform, chlorobenzilate, chloropicrin, clothiazoben, cycloprene, cyflumetofen,
5 dicyclanil, fenoxacrim, fentrifanil, flubenzimine, flufenerim, flutenzin, gossyplure,
hydramethylnone, japonilure, metoxadiazone, petroleum, piperonyl butoxide, potassium oleate,
pyrafluprole, pyridalyl, pyriprole, sulfluramid, tetradifon, tetrasul, triarathene, verbutin,
furthermore the compound 3-methyl-phenyl propylcarbamate (tsumacide Z), the compound 3-(5-chloro-
3-pyridinyl)-8-(2,2,2-trifluoroethyl)-8-azabicyclo[3.2.1]octane-3-carbonitrile (CAS Reg. No. 185982-80-
3) and the corresponding 3-endo isomer (CAS Reg. No. 185984-60-5) (cf. WO 96/37494, WO
10 98/25923), and preparations which contain insecticidally active plant extracts, nematodes, fungi or
viruses.

A mixture with other known active compounds such as herbicides, or with fertilizers and growth
regulators, safeners or semiochemicals is also possible.

15

In addition, the compounds of the formula (I) according to the invention also have very good
antimycotic activity. They have a very broad antimycotic spectrum of action, in particular against
dermatophytes and budding fungi, moulds and diphasic fungi (for example against *Candida* species
such as *Candida albicans*, *Candida glabrata*) and *Epidermophyton floccosum*, *Aspergillus* species
20 such as *Aspergillus niger* and *Aspergillus fumigatus*, *Trichophyton* species such as *Trichophyton*
mentagrophytes, *Microsporon* species such as *Microsporon canis* and *audouinii*. The enumeration
of these fungi is no restriction whatsoever of the mycotic spectrum which can be controlled and is
provided by illustration only.

25 The active compounds can be employed as such, in the form of their formulations or the use forms
prepared therefrom, such as ready-to-use solutions, suspensions, wettable powders, pastes, soluble
powders, dusts and granules. They are applied in the customary manner, for example by pouring,
spraying, atomizing, broadcasting, dusting, foaming, painting on and the like. It is furthermore
possible to apply the active compounds by the ultra-low-volume method, or to inject the active
30 compound preparation or the active compound itself into the soil. The seed of the plant can also be
treated.

When employing the active compounds according to the invention as fungicides, the application
rates can be varied within a substantial range, depending on the type of application. In the
35 treatment of plant parts, the application rates of active compound are generally between 0.1 and
10 000 g/ha, preferably between 10 and 1000 g/ha. For the treatment of seed, the application rates

of active compound are generally between 0.001 and 50 g per kilogram of seed, preferably between 0.01 and 10 g per kilogram of seed. For treating the soil, the application rates of active compound are generally between 0.1 and 10 000 g/ha, preferably between 1 and 5000 g/ha.

- 5 As already mentioned above, all plants and their parts can be treated in accordance with the invention. In a preferred embodiment, plant species and plant varieties which are found in the wild or are obtained by traditional biological breeding methods, such as hybridization or protoplast fusion, and parts of the former are treated. In a further preferred embodiment, transgenic plants and plant varieties which have been obtained by recombinant methods, if appropriate in combination
10 with traditional methods (genetically modified organisms) and their parts are treated. The term "parts" or "parts of plants" or "plant parts" has been illustrated above.

Particularly preferably, plants of the plant cultivars which are in each case commercially available or in use are treated according to the invention. Plant cultivars are understood as meaning plants
15 with new properties ("traits") which have been obtained by conventional cultivation, by mutagenesis or else by recombinant DNA techniques. These may be cultivars, breeds, biotypes or genotypes.

Depending on the plant species or plant cultivars, their location and growth conditions (soils,
20 climate, vegetation period, nutrition), the treatment according to the invention may also result in superadditive ("synergistic") effects. Thus, for example, reduced application rates and/or extensions of the activity spectrum and/or an increase in the activity of the substances and compositions that can be used according to the invention, better plant growth, more developed root system, higher resistance of the plant variety or plant cultivar, increased growth of shoots, higher
25 plant vitality, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil salinity, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, larger fruit, increased plant size, greener leaf colour, earlier blossoming, better quality and/or a higher nutritional value of the harvested products, higher sugar concentration in the fruits, better storage stability and/or processability of the harvested products
30 which exceed the effects which were actually to be expected are possible.


The preferred transgenic plants or plant cultivars (i.e. those obtained by genetic engineering) which are to be treated according to the invention include all plants which, as a result of the recombinant modification, received genetic material which imparted particularly advantageous
35 useful properties ("traits") to these plants. Examples of such properties are better plant growth, increased tolerance to high or low temperatures, increased tolerance to drought or to water or soil

salinity, increased flowering performance, easier harvesting, accelerated maturation, higher harvest yields, better quality and/or a higher nutritional value of the harvested products, better storage stability and/or processability of the harvested products. Further and particularly emphasized examples of such properties are a better defence of the plants against animal and microbial pests, such as against insects, mites, phytopathogenic fungi, bacteria and/or viruses, and also increased tolerance of the plants to certain herbicidally active compounds. Examples of transgenic plants which may be mentioned are the important crop plants, such as cereals (wheat, rice), maize, soya beans, potatoes, cotton, tobacco, oilseed rape and also fruit plants (with the fruits apples, pears, citrus fruits and grapes), and particular emphasis is given to maize, soya beans, potatoes, cotton, tobacco and oilseed rape. Traits that are emphasized in particular are increased defence of the plants against insects, arachnids, nematodes, slugs and snails as the result of toxins formed in the plants, in particular those formed in the plants by the genetic material from *Bacillus thuringiensis* (for example by the genes CryIA(a), CryIA(b), CryIA(c), CryIIA, CryIIIA, CryIIIB2, Cry9c, Cry2Ab, Cry3Bb and CryIF and also combinations thereof) (hereinbelow referred to as “Bt plants”). Traits which are also particularly emphasized are the increased defence of plants against fungi, bacteria and viruses by systemic acquired resistance (SAR), systemin, phytoalexins, elicitors and resistance genes and the correspondingly expressed proteins and toxins. Traits that are furthermore particularly emphasized are the increased tolerance of the plants to certain herbicidally active compounds, for example imidazolinones, sulphonylureas, glyphosate or phosphinothricin (for example the “PAT” gene). The genes which impart the desired traits in question can also be present in combination with one another in the transgenic plants. Examples of “Bt plants” which may be mentioned are maize varieties, cotton varieties, soya bean varieties and potato varieties which are sold under the trade names YIELD GARD® (for example maize, cotton, soya beans), KnockOut® (for example maize), StarLink® (for example maize), Bollgard® (cotton), Nucoton® (cotton) and NewLeaf® (potato). Examples of herbicide-tolerant plants which may be mentioned are maize varieties, cotton varieties and soya bean varieties which are sold under the trade names Roundup Ready® (tolerance to glyphosates, for example maize, cotton, soya bean), Liberty Link® (tolerance to phosphinothricin, for example oilseed rape), IMI® (tolerance to imidazolinones) and STS® (tolerance to sulphonylureas, for example maize). Herbicide-resistant plants (plants bred in a conventional manner for herbicide tolerance) which may be mentioned also include the varieties sold under the name Clearfield® (for example maize). Of course, these statements also apply to plant cultivars having these genetic traits or genetic traits still to be developed, which cultivars will be developed and/or marketed in the future.

The plants listed can be treated according to the invention in a particularly advantageous manner with the compounds of the general formula (I) or the active compound mixtures according to the

invention. The preferred ranges stated above for the active compounds or mixtures also apply to the treatment of these plants. Particular emphasis is given to the treatment of plants with the compounds or mixtures specifically mentioned in the present text.


- 5 The preparation and the use of the active compounds according to the invention is illustrated by the examples below.

Preparation examplesPreparation of compound 9














- 5 At 0°C, 0.1 g (3.6 mmol) of sodium hydride (80% pure) and 1.2 g (8.3 mmol) of methyl iodide in 5 ml of dimethylformamide are added to a solution consisting of 1.0 g (2.8 mmol) of N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-thiocarboxamide in 10 ml of dimethylformamide. The reaction mixture is slowly warmed to room temperature and stirred at this temperature for 1 hour. For work-up, the reaction mixture is poured into water and extracted three times
- 10 with in each case 50 ml of ethyl acetate. The combined organic phases are washed with saturated sodium chloride solution, dried over sodium sulphate and concentrated under reduced pressure. Column chromatography (cyclohexane/ethyl acetate 4:1) gives 0.7 g (1.9 mmol, 67% of theory) of methyl N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carbimidothioate [logP (pH 2.3) = 4.75].





















15









Analogous to this example and in accordance with the general descriptions of the processes according to the invention, it is possible to obtain the compounds of the formula (I) listed in Table 1 below.

20 **Table 1**

No.	X	M	Q	Z	A	logP (pH 2.3)
1	SO ₂ CH ₃	M-1, R ⁷ = H	direct bond			


No.	X	M	Q	Z	A	logP (pH 2.3)
2	$\text{SCH}_2\text{CH}(\text{OCH}_3)_2$	M-1, $\text{R}^7 = \text{H}$	direct bond			
3	$\text{S}(\text{CH}_2)_2\text{OCH}_3$	M-1, $\text{R}^7 = \text{H}$	direct bond			4.62
4	$\text{S}(\text{CH}_2)_3\text{OH}$	M-1, $\text{R}^7 = \text{H}$	direct bond			
5	$\text{SCH}_2\text{C}(\text{O})\text{NHCH}_3$	M-1, $\text{R}^7 = \text{H}$	direct bond			
6	SCH_2CN	M-1, $\text{R}^7 = \text{H}$	direct bond			4.15
7	SCH_2SCN	M-1, $\text{R}^7 = \text{H}$	direct bond			4.55
8	$\text{SCH}_2\text{C}\equiv\text{CH}$	M-1, $\text{R}^7 = \text{H}$	direct bond			4.65
9	SCH_3	M-1, $\text{R}^7 = \text{H}$	direct bond			4.75

No.	X	M	Q	Z	A	logP (pH 2.3)
10	$S(CH_2)_2OH$	M-1, $R^7 = H$	direct bond			3.23
11	$SCH_2CO_2C_2H_5$	M-1, $R^7 = H$	direct bond			4.66
12		M-1, $R^7 = H$	direct bond			4.95
13	$S(CH_2)_{11}CH_3$	M-1, $R^7 = H$	direct bond			7.36
14		M-1, $R^7 = H$	direct bond			
15		M-1, $R^7 = H$	direct bond			
16	$SC(O)CH_2OCH_3$	M-1, $R^7 = H$	direct bond			
17		M-1, $R^7 = H$	direct bond			

No.	X	M	Q	Z	A	logP (pH 2.3)
18	$\text{SC(O)C(CH}_3)_3$	M-1, $\text{R}^7 = \text{H}$	direct bond			5.81
19		M-1, $\text{R}^7 = \text{H}$	direct bond			6.52
20		M-1, $\text{R}^7 = \text{H}$	direct bond			6.66


Preparation of starting materials of the formula (II-a)

5 Preparation of compound (II-7)



0.3 g (1.4 mmol) of phosphorus pentasulphide is added to a solution consisting of 0.48 g (1.4 mmol) of N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-carboxamide in 20 ml of toluene, and the reaction mixture is stirred at 70°C for 2 hours. For work-up, the reaction mixture is poured into water and extracted once with 20 ml of toluene and twice with in each case 20 ml of ethyl acetate. The combined organic phases are dried over sodium sulphate and concentrated under reduced pressure. Column chromatography (gradient cyclohexane/ethyl acetate) gives 0.32 g (0.89 mmol, 63% of theory) of N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-thiocarboxamide [logP (pH 2.3) = 3.66].


Preparation of compound (II-8)


















0.9 g (8.4 mmol) of phenylhydrazine in 10 ml of dimethylformamide is added to a solution consisting of 1.0 g (2.7 mmol) of N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-1H-pyrazole-4-thiocarboxamide in 10 ml of methanol. The reaction mixture is stirred at 70°C for 16 hours. For work-up, the reaction mixture is poured into water and extracted three times with in each case 20 ml of dichloromethane. The combined organic phases are dried over sodium sulphate and concentrated under reduced pressure. Column chromatography (gradient cyclohexane/ethyl acetate) gives 0.42 g (0.96 mmol, 35% of theory) of N-(2-bicyclo[2.2.1]hept-2-ylphenyl)-3-(difluoromethyl)-1-methyl-N'-phenyl-1H-pyrazole-4-carbohydrazonamide [logP (pH 2.3) = 3.66].

Analogously to this example and in accordance with the general descriptions of the processes according to the invention, it is possible to obtain the compounds of the formula (III) listed in Table 2 below.

Table 2



No.	X	M	Q	Z	A	logP (pH 2.3)
II-1	S	M-1, R ⁷ = H	direct bond	G5		3.70 (at pH 7.5)
II-2	S	M-1, R ⁷ = 4-F	direct bond			

No.	X	M	Q	Z	A	logP (pH 2.3)
II-3	N-OCH ₃	M-1, R ⁷ = H	direct bond			
II-4	N-CH ₃	M-1, R ⁷ = H	direct bond			
II-5	N-NH ₂	M-1, R ⁷ = H	direct bond			
II-6	N-N(CH ₃) ₂	M-1, R ⁷ = H	direct bond			
II-7	S	M-1, R ⁷ = H	direct bond			3.66
II-8		M-1, R ⁷ = H	direct bond			3.66
II-9	N-OH	M-1, R ⁷ = H	direct bond			2.88

The logP values given in the Tables and Preparation Examples above are determined in accordance with EEC Directive 79/831 Annex V.A8 by HPLC (High Performance Liquid Chromatography) on a reversed-phase column (C 18). Temperature: 43°C.

The determination is carried out in the acidic range at pH 2.3 using the mobile phases 0.1% aqueous phosphoric acid and acetonitrile; linear gradient from 10% acetonitrile to 90% acetonitrile.

5 The determination is carried out in the neutral range at pH 7.5 using the mobile phases 0.01-molar aqueous phosphate buffer solution and acetonitrile; linear gradient from 10% acetonitrile to 90% acetonitrile.

Calibration is carried out using unbranched alkan-2-ones (having 3 to 16 carbon atoms), with known logP values (determination of the logP values by the retention times using linear interpolation between two successive alkanones).

10 The lambda max values were determined in the maxima of the chromatographic signals using the UV spectra from 200 nm to 400 nm.

Use examples

Example A

15 **Sphaerotheca Test (cucumber)/protective**

Solvents: 24.5 parts by weight of acetone

24.5 parts by weight of dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether




20 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier, and the concentrate is diluted with water to the desired concentration.

To test for protective activity, young plants are sprayed with the active compound preparation at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of *Sphaerotheca fuliginea*. The plants are then placed in a greenhouse at
25 about 23°C and a relative atmospheric humidity of about 70%.

Evaluation is carried out 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table A

Sphaerotheca Test (cucumber)/protective

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(10) 	100	100
(11) 	100	100
(20) 	100	100

Example B





5 **Venturia Test (apple)/protective**

Solvents: 24.5 parts by weight of acetone
 24.5 parts by weight of dimethylacetamide
Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is
10 mixed with the stated amounts of solvents and emulsifier and the concentrate is diluted with water
to the desired concentration. To test for protective activity, young plants are sprayed with the
active compound preparation at the stated application rates. After the spray coating has dried on,
the plants are inoculated with an aqueous conidia suspension of the apple scab pathogen *Venturia*
inaequalis and then remain in an incubation cabin at about 20°C and 100% relative atmospheric
15 humidity for 1 day. The plants are then placed in a greenhouse at about 21°C and a relative
atmospheric humidity of about 90%. Evaluation is carried out 10 days after the inoculation. 0%

means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table B
Venturia Test (apple)/protective

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(10) 	100	100
(11) 	100	100
(19) 	100	99
(20) 	100	100

5

Example C
Botrytis Test (bean)/protective



Solvents: 24.5 parts by weight of acetone
 24.5 parts by weight of dimethylacetamide

10 Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier and the concentrate is diluted with water

to the desired concentration. To test for protective activity, young plants are sprayed with the active compound preparation at the stated application rate. After the spray coating has dried on, 2 small pieces of agar colonized by *Botrytis cinerea* are placed onto each leaf. The inoculated plants are placed in a dark chamber at about 20°C and 100% relative atmospheric humidity. The size of the infected areas on the leaves is evaluated 2 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table C
Botrytis Test (bean)/protective

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(10) 	250	100
(11) 	250	100

10

Example D
Uromyces Test (bean)/protective





- Solvents:
- 24.5 parts by weight of acetone
- 24.5 parts by weight of dimethylacetamide
- Emulsifier:
- 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvents and emulsifier and the concentrate is diluted with water to the desired concentration. To test for protective activity, young plants are sprayed with the active compound preparation at the stated application rate. After the spray coating has dried on, the plants are inoculated with an aqueous spore suspension of the bean rust pathogen *Uromyces appendiculatus* and then remain in an incubation cabinet at about 20°C and 100% relative atmospheric humidity for 1 day. The plants are then placed in a greenhouse at about 21°C and a relative atmospheric humidity of about 90%. Evaluation is carried out 10 days after the

20

inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table D
Uromyces Test (bean)/protective

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(10) 	100	100
(11) 	100	100
(19) 	100	96
(20) 	100	99

5

Example E





Alternaria Test (tomato)/protective

Solvent: 49 parts by weight of N,N-dimethylformamide
Emulsifier: 1 part by weight of alkylaryl polyglycol ether

10 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier and the concentrate is diluted with water to the desired concentration. To test for protective activity, young tomato plants are sprayed with

the active compound preparation at the stated application rate. 1 day after the treatment, the plants are inoculated with a spore suspension of *Alternaria solani* and then remain at 100% relative atmospheric humidity and 20°C for 24 h. The plants then remain at 96% relative atmospheric humidity and a temperature of 20°C. Evaluation is carried out 7 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table E
Alternaria Test (tomato)/protective

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(10) 	500	100
(11) 	500	95
(19) 	500	90
(20) 	500	80

10 Example F




Pyrenophora teres Test (barley)/protective

Solvent: 50 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier and the concentrate is diluted with water to the desired concentration. To test for protective activity, young plants are sprayed with the active compound preparation at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Pyrenophora teres*. The plants remain in an incubation cabin at 20°C and 100% relative atmospheric humidity for 48 hours. The plants are then placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of about 80%. Evaluation is carried out 8 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.

Table F
***Pyrenophora teres* Test (barley)/protective**

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(11) 	1000	100
(19) 	1000	92
(20) 	1000	100

Example G**Puccinia Test (wheat)/protective**




Solvent: 50 parts by weight of N,N-dimethylacetamide

Emulsifier: 1 part by weight of alkylaryl polyglycol ether

- 5 To produce a suitable preparation of active compound, 1 part by weight of active compound is mixed with the stated amounts of solvent and emulsifier and the concentrate is diluted with water to the desired concentration. To test for protective activity, young plants are sprayed with the active compound preparation at the stated application rate. After the spray coating has dried on, the plants are sprayed with a conidia suspension of *Puccinia recondita*. The plants remain in an
- 10 incubation cabin at 20°C and 100% relative atmospheric humidity for 48 hours. The plants are then placed in a greenhouse at a temperature of about 20°C and a relative atmospheric humidity of about 80% to promote the development of rust pustules. Evaluation is carried out 10 days after the inoculation. 0% means an efficacy which corresponds to that of the control, whereas an efficacy of 100% means that no infection is observed.


15

Table G**Puccinia Test (wheat)/protective**

Active compound according to the invention	Application rate of active compound in ppm	Efficacy in %
(11) 	1000	100
(19) 	1000	93
(20) 	1000	100

Patent claims

1. Carboxamides of the formula (I)



5 in which

X represents OR^1 , SR^2 , NHR^3 , NR^4R^5 , SOR^6 , SO_2R^6 ,

10 R^1 represents $\text{C}_1\text{-C}_{12}\text{-alkyl}$, $\text{C}_2\text{-C}_8\text{-alkenyl}$, $\text{C}_2\text{-C}_8\text{-alkynyl}$, $\text{C}_1\text{-C}_8\text{-haloalkyl}$, hydroxy-
 $\text{C}_1\text{-C}_8\text{-alkyl}$, $(\text{C}_1\text{-C}_4\text{-alkoxy})\text{-C}_1\text{-C}_8\text{-alkyl}$, $\text{bis}(\text{C}_1\text{-C}_4\text{-alkoxy})\text{-C}_1\text{-C}_8\text{-alkyl}$, $(\text{C}_1\text{-C}_4\text{-alkoxy})\text{carbonyl-(C}_1\text{-C}_4\text{-alkyl)}$; in each case optionally halogen-, $\text{C}_1\text{-C}_4\text{-alkyl-}$ or $\text{C}_1\text{-C}_4\text{-haloalkyl-substituted phenyl-(C}_1\text{-C}_2\text{-alkyl)}$ or $\text{hetaryl-(C}_1\text{-C}_2\text{-alkyl)}$; in each case optionally $\text{C}_1\text{-C}_4\text{-alkoxy-}$, halogen-, $\text{C}_1\text{-C}_4\text{-alkyl-}$ or $\text{C}_1\text{-C}_4\text{-haloalkyl-substituted (C}_1\text{-C}_6\text{-alkyl)carbonyl}$, $(\text{C}_3\text{-C}_8\text{-cycloalkyl)carbonyl}$, phenylcarbonyl or hetarylcarbonyl ,

15 R^2 represents $\text{C}_1\text{-C}_{12}\text{-alkyl}$, $\text{C}_2\text{-C}_8\text{-alkenyl}$, $\text{C}_2\text{-C}_8\text{-alkynyl}$, $\text{C}_1\text{-C}_8\text{-haloalkyl}$, hydroxy- $\text{C}_1\text{-C}_8\text{-alkyl}$, $(\text{C}_1\text{-C}_4\text{-alkoxy})\text{-C}_1\text{-C}_8\text{-alkyl}$, $\text{bis}(\text{C}_1\text{-C}_4\text{-alkoxy})\text{-C}_1\text{-C}_8\text{-alkyl}$, cyano- $(\text{C}_1\text{-C}_4\text{-alkyl})$, $-\text{CH}_2\text{S-CN}$, $(\text{C}_1\text{-C}_4\text{-alkoxy})\text{carbonyl-(C}_1\text{-C}_4\text{-alkyl)}$; in each case optionally halogen-, $\text{C}_1\text{-C}_4\text{-alkyl-}$ or $\text{C}_1\text{-C}_4\text{-haloalkyl-substituted phenyl-(C}_1\text{-C}_2\text{-alkyl)}$ or $\text{hetaryl-(C}_1\text{-C}_2\text{-alkyl)}$; in each case optionally $\text{C}_1\text{-C}_4\text{-alkoxy-}$, halogen-, $\text{C}_1\text{-C}_4\text{-alkyl-}$ or $\text{C}_1\text{-C}_4\text{-haloalkyl-substituted (C}_1\text{-C}_6\text{-alkyl)carbonyl}$, $(\text{C}_3\text{-C}_8\text{-cycloalkyl)carbonyl}$, phenylcarbonyl or hetarylcarbonyl ,

20 R^3 , R^4 and R^5 independently of one another represent amino, hydroxyl, cyano, $\text{C}_1\text{-C}_{12}\text{-alkyl}$, $\text{C}_1\text{-C}_8\text{-alkoxy}$, $\text{C}_2\text{-C}_8\text{-alkenyl}$, $\text{C}_2\text{-C}_8\text{-alkynyl}$, $\text{C}_1\text{-C}_6\text{-alkylamino}$, $\text{di(C}_1\text{-C}_6\text{-alkyl)amino}$, hydroxy- $\text{C}_1\text{-C}_8\text{-alkyl}$, $(\text{C}_1\text{-C}_4\text{-alkoxy})\text{-C}_1\text{-C}_8\text{-alkyl}$, $\text{bis(C}_1\text{-C}_4\text{-alkoxy)-C}_1\text{-C}_8\text{-alkyl}$, cyano- $(\text{C}_1\text{-C}_4\text{-alkyl})$, phenoxy, phenylamino, benzyloxy or benzylamino,

25

 R^6 represents $\text{C}_1\text{-C}_{12}\text{-alkyl}$ or $\text{C}_1\text{-C}_8\text{-haloalkyl}$,M represents a phenyl, thiophene, pyridine, pyrimidine, pyridazine or pyrazine ring, each of which is monosubstituted by R^7 , or represents a thiazole ring which is substituted by R^8 ,

30

 R^7 represents hydrogen, fluorine, chlorine, methyl, isopropyl, methylthio or trifluoromethyl, R^8 represents hydrogen, methyl, methylthio or trifluoromethyl,Q represents a direct bond, $\text{C}_1\text{-C}_4\text{-alkylene}$, $\text{C}_2\text{-C}_4\text{-alkenylene}$, O, S, SO, SO_2 or NR^9 ,


- R^9 represents hydrogen, C_1 - C_8 -alkyl, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -haloalkenyl, C_2 - C_6 -haloalkynyl or C_3 - C_6 -cycloalkyl,
- Z represents Z^1 , Z^2 , Z^3 , Z^4 , Z^5 or Z^6 ,
- 5 Z^1 represents phenyl which is optionally mono- to pentasubstituted by identical or different substituents,
- Z^2 represents pyridinyl which is optionally mono- to trisubstituted by identical or different substituents,
- 10 Z^3 represents cycloalkyl or bicycloalkyl, each of which is optionally mono- or polysubstituted by identical or different halogen, alkyl and/or $-(CR^{10}R^{11})_mSiR^{12}R^{13}R^{14}$ substituents,
- 15 Z^4 represents unsubstituted C_2 - C_{20} -alkyl or represents C_1 - C_{20} -alkyl which is mono- or polysubstituted by identical or different halogen, alkylthio, alkylsulphinyl, alkylsulphonyl, alkoxy, alkylamino, dialkylamino, haloalkylthio, haloalkylsulphinyl, haloalkylsulphonyl, haloalkoxy, haloalkylamino, halodialkylamino, $-SiR^{12}R^{13}R^{14}$ and/or C_3 - C_6 -cycloalkyl substituents, where the cycloalkyl moiety for its part may optionally be mono- or polysubstituted by identical or different halogen and/or C_1 - C_4 -alkyl substituents,
- 20 Z^5 represents C_2 - C_{20} -alkenyl or C_2 - C_{20} -alkynyl, each of which is optionally mono- or polysubstituted by identical or different halogen, alkylthio, alkylsulphinyl, alkylsulphonyl, alkoxy, alkylamino, dialkylamino, haloalkylthio, haloalkylsulphinyl, haloalkylsulphonyl, haloalkoxy, haloalkylamino, halodialkylamino, $-SiR^{12}R^{13}R^{14}$ and/or C_3 - C_6 -cycloalkyl substituents, where the cycloalkyl moiety for its part may optionally be mono- or polysubstituted by identical or different halogen and/or C_1 - C_4 -alkyl substituents,
- 25 Z^6 represents an optionally mono- or polysubstituted saturated or unsaturated 3- to 7-membered ring which contains a silicon atom as ring member, in which case Q represents a direct bond or C_1 - C_4 -alkylene,
- R^{10} represents hydrogen or C_1 - C_4 -alkyl,
- 30 R^{11} represents hydrogen or C_1 - C_4 -alkyl,
- m represents 0, 1, 2 or 3,
- R^{12} and R^{13} independently of one another represent hydrogen, C_1 - C_8 -alkyl, C_1 - C_8 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl or C_1 - C_6 -haloalkyl,
- 35 R^{14} represents hydrogen, C_1 - C_8 -alkyl, C_1 - C_8 -alkoxy, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylthio- C_1 - C_4 -alkyl, C_2 - C_8 -alkenyl, C_2 - C_8 -alkynyl, C_1 - C_6 -haloalkyl, C_2 - C_6 -

haloalkenyl, C₂-C₆-haloalkynyl, C₃-C₆-cycloalkyl, or represents in each case optionally substituted phenyl or phenylalkyl,

or

M-Q-Z together represent 1H-2,3-dihydroinden-4-yl, 1,3-dihydro-2-benzofuran-4-yl or 1,3-dihydro-2-benzothien-4-yl, each of which is optionally mono- to trisubstituted by methyl,

A represents one of the radicals A1 to A19 below



R¹⁵ represents hydrogen, cyano, halogen, nitro, C₁-C₄-alkyl, C₁-C₄-alkoxy, C₁-C₄-alkylthio, C₃-C₆-cycloalkyl, C₁-C₄-haloalkyl, C₁-C₄-haloalkoxy or C₁-C₄-haloalkylthio having in each case 1 to 5 halogen atoms, aminocarbonyl or aminocarbonyl-C₁-C₄-alkyl,

R¹⁶ represents hydrogen, halogen, cyano, C₁-C₄-alkyl, C₁-C₄-alkoxy or C₁-C₄-alkylthio,


R¹⁷ represents hydrogen, C₁-C₄-alkyl, hydroxy-C₁-C₄-alkyl, C₂-C₆-alkenyl, C₃-C₆-cycloalkyl, C₁-C₄-alkylthio-C₁-C₄-alkyl, C₁-C₄-alkoxy-C₁-C₄-alkyl, C₁-C₄-haloalkyl, C₁-C₄-haloalkylthio-C₁-C₄-alkyl, C₁-C₄-haloalkoxy-C₁-C₄-alkyl having in each case 1 to 5 halogen atoms, or phenyl,

R¹⁸ and R¹⁹ independently of one another, represent hydrogen, halogen, C₁-C₄-alkyl or C₁-C₄-haloalkyl having 1 to 5 halogen atoms,

- R^{20} represents halogen, cyano or C_1 - C_4 -alkyl, or C_1 - C_4 -haloalkyl or C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms,
- R^{21} and R^{22} independently of one another represent hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- 5 R^{23} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{24} represents hydrogen, halogen, hydroxy, cyano, C_1 - C_6 -alkyl, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy or C_1 - C_4 -haloalkylthio having in each case 1 to 5 halogen atoms,
- R^{25} represents halogen, hydroxyl, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, 10 C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkylthio or C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms,
- R^{26} represents hydrogen, halogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms, C_1 - C_4 -alkylsulphinyl or C_1 - C_4 -alkylsulphonyl,
- 15 R^{27} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{28} represents C_1 - C_4 -alkyl,
- Q^1 represents S (sulphur), SO, SO₂ or CH₂,
- p represents 0, 1 or 2, where R^{22} represents identical or different radicals if p represents 2,
- 20 R^{29} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{30} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{31} and R^{32} independently of one another represent hydrogen, halogen, amino, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{33} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms, 25
- R^{34} and R^{35} independently of one another represent hydrogen, halogen, amino, nitro, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{36} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- 30 R^{37} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di-(C_1 - C_4 -alkyl)amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{38} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{39} represents hydrogen, halogen, amino, C_1 - C_4 -alkylamino, di-(C_1 - C_4 -alkyl)amino, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- 35 R^{40} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
- R^{41} represents halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,

- 5
10
15
20
- R^{42} represents hydrogen or C_1 - C_4 -alkyl,
 R^{43} represents halogen or C_1 - C_4 -alkyl,
 R^{44} represents C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{45} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{46} represents halogen, hydroxyl, C_1 - C_4 -alkyl, C_1 - C_4 -alkoxy, C_1 - C_4 -alkylthio, C_1 - C_4 -haloalkyl, C_1 - C_4 -haloalkylthio or C_1 - C_4 -haloalkoxy having in each case 1 to 5 halogen atoms,
 R^{47} represents hydrogen, cyano, C_1 - C_4 -alkyl, C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms, C_1 - C_4 -alkoxy- C_1 - C_4 -alkyl, hydroxy- C_1 - C_4 -alkyl, C_1 - C_4 -alkylsulphonyl, di(C_1 - C_4 -alkyl)-aminosulphonyl, C_1 - C_6 -alkylcarbonyl or in each case optionally substituted phenylsulphonyl or benzoyl,
 R^{48} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{49} represents hydrogen, halogen, cyano, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{50} represents hydrogen, halogen, C_1 - C_4 -alkyl or C_1 - C_4 -haloalkyl having 1 to 5 halogen atoms,
 R^{51} represents C_1 - C_4 -alkyl.

2. Carboxamides according to Claim 1 of the formulae (I-a), (I-b), (I-e), (I-f) and (I-d)




25 in which

R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , M, Q, Z and A are as defined in Claim 1,

n represents 1 or 2.

3. Process for preparing carboxamides of the formula (I) according to Claim 1, characterized in that

- (a) if X represents OR^1 , carboxamides of the formula (II-a)



in which M, Q, Z and A are as defined in Claim 1
are, if appropriate, in the presence of a base, reacted with an alkylating agent of the formula (III-a)



in which

R^1 is as defined in Claim 1 and

LG^1 represents a leaving group

or with a Meerwein salt of the formula (III-d)




in which

R^1 is as defined in Claim 1 and


E represents BF_4 , SbF_6 or $SbCl_6$,

giving carboxamides of the formula (I-a)



in which R^1 , M, Q, Z and A are as defined in Claim 1,


- (b) if X represents SR^2 , SOR^6 or SO_2R^6 , in a first step carboxamides of the formula (II-a)



in which M, Q, Z and A are as defined in Claim 1

are, if appropriate in the presence of a diluent, reacted with a thionating agent,

thus giving carboxamides of the formula (II-b)



in which M, Q, Z and A are as defined in Claim 1

5

which, in a second step, are, if appropriate in the presence of a base, reacted with an alkylating agent of the formula (III-b) or the formula (III-c)



10

in which



R^2 and R^6 are as defined in Claim 1,

LG^2 represents a leaving group,

LG^6 represents a leaving group,

15


giving carboxamides of the formula (I-b) or the formula (I-c)



in which R^2 , R^6 , M, Q, Z and A are as defined in Claim 1,

20

and, in a third step, the carboxamides of the formula (I-c) are then reacted with an oxidizing agent, if appropriate in the presence of a diluent, if appropriate in the presence of an acid binder and if appropriate in the presence of a catalyst, thus giving the carboxamides of the formula (I-d)




in which

R^6 , M, Q, Z and A are as defined in Claim 1 and


n represents 1 or 2,

- 5 (c) if X represents NHR^3 or NR^4R^5 , carboxamides of the formula (II-b)



in which M, Q, Z and A are as defined in Claim 1

are, if appropriate in the presence of a base, reacted with an amidating agent of the
10 formula (IV-a) or the formula (IV-b)





in which

R^3 , R^4 and R^5 are as defined in Claim 1 and

15 LG^3 represents a leaving group,


giving carboxamides of the formula (I-e) or the formula (I-f)



20 in which R^3 , R^4 , R^5 , M, Q, Z and A are as defined in Claim 1.

4. Compositions for controlling unwanted microorganisms, characterized in that they comprise at least one carboxamide of the formula (I) according to Claim 1, in addition to extenders and/or surfactants.

5. Use of carboxamides of the formula (I) according to Claim 1 for controlling unwanted microorganisms.
6. Method for controlling unwanted microorganisms, characterized in that carboxamides of the formula (I) according to Claim 1 are applied to the microorganisms and/or their habitat.
7. Process for preparing compositions for controlling unwanted microorganisms, characterized in that carboxamides of the formula (I) according to Claim 1 are mixed with extenders and/or surfactants.
8. Carboxamides of the formula (II-c)



in which

- A¹ represents A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, A11, A12, A13, A14, A15, A16, A17 and A19, as defined in Claim 1,
- M, Q and Z are as defined in Claim 1,
- except for compounds in which
- M represents M-1,
- R⁷ represents hydrogen,
- Q represents a direct bond,
- Z represents phenyl which is substituted in the 4-position by halogen,
- A¹ represents A1,
- R¹⁵ represents C₁-C₃-haloalkyl,
- R¹⁶ represents hydrogen,
- R¹⁷ represents C₁-C₃-alkyl, C₁-C₃-alkoxy-C₁-C₃-alkyl, C₁-C₃-haloalkyl, C₁-C₃-haloalkoxy-C₁-C₃-alkyl.

Fetherstonhaugh
Ottawa, Canada
Patent Agents

