



(86) Date de dépôt PCT/PCT Filing Date: 2009/07/07  
(87) Date publication PCT/PCT Publication Date: 2010/01/14  
(85) Entrée phase nationale/National Entry: 2010/12/30  
(86) N° demande PCT/PCT Application No.: US 2009/049744  
(87) N° publication PCT/PCT Publication No.: 2010/005922  
(30) Priorité/Priority: 2008/07/07 (US61/129,578)

(51) Cl.Int./Int.Cl. *C07C 311/01* (2006.01),  
*A61K 31/195* (2006.01), *A61K 31/662* (2006.01),  
*A61P 3/04* (2006.01), *C07C 311/15* (2006.01),  
*C07F 9/38* (2006.01)  
(71) Demandeurs/Applicants:  
FASGEN, INC., US;  
THE JOHNS HOPKINS UNIVERSITY, US  
(72) Inventeurs/Inventors:  
TOWNSEND, CRAIG A., US;  
KUHAJDA, FRANCIS, US;  
MEDGHALCHI, SUSAN, US  
(74) Agent: ROBIC

(54) Titre : NOUVEAUX COMPOSES, COMPOSITIONS PHARMACEUTIQUES LES CONTENANT, ET LEURS  
PROCEDES D'UTILISATION ET DE PREPARATION

(54) Title: NOVEL COMPOUNDS, PHARMACEUTICAL COMPOSITIONS CONTAINING SAME, METHODS OF USE  
FOR SAME, AND METHODS FOR PREPARING SAME





**FIG. 1**



(57) **Abrégé/Abstract:**

The present invention relates to a novel class of compounds comprising formula I, wherein n is 0 or 1. A is NR<sup>1</sup>, O, or S, wherein R<sup>1</sup> is H, hydroxyl, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, alkenyl, aryl, alkylaryl or arylalkyl. X is a carboxylate, a phosphonate, or a phosphate



**(57) Abrégé(suite)/Abstract(continued):**

residue, or a C<sub>1</sub>-C<sub>10</sub> alkyl residue optionally substituted with a carboxylate, phosphonate or phosphate residue. Y is a C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, halide, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl, or a heterocyclic ring and is optionally substituted with one or more halides. Z is a H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring and is optionally substituted with one or more C<sub>1</sub>-C<sub>10</sub> alkyl groups, C<sub>1</sub>-C<sub>10</sub> alkoxy groups, hydroxyl groups, cyano groups, carboxylate groups, halides, aryl groups, alkylaryl groups, arylalkyl groups, cycloalkyl groups, cycloalkenyl groups or heterocyclic rings.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
14 January 2010 (14.01.2010)(10) International Publication Number  
**WO 2010/005922 A1**(51) International Patent Classification:  
*A01N 41/06* (2006.01) *A61K 31/18* (2006.01)(21) International Application Number:  
PCT/US2009/049744(22) International Filing Date:  
7 July 2009 (07.07.2009)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/129,578 7 July 2008 (07.07.2008) US(71) Applicants (for all designated States except US): **FAS-GEN, INC.** [US/US]; 800 West Baltimore Street, Suite 150, Baltimore, Maryland 21201 (US). **THE JOHNS HOPKINS UNIVERSITY** [US/US]; 3400 N. Charles Street, Baltimore, Maryland 21218 (US).



(72) Inventors; and

(75) Inventors/Applicants (for US only): **TOWNSEND, Craig A.** [US/US]; 3400 N. Charles Street, Baltimore,Maryland 21218 (US). **KUHAJDA, Francis** [US/US]; 800 W. Baltimore Street, Suite 150, Baltimore, Maryland 21224 (US). **MEDGHALCHI, Susan** [US/US]; 800 W. Baltimore Street, Suite 150, Baltimore, Maryland 21201 (US).(74) Agents: **SUMNER, J. Eric** et al.; Fox Rothschild LLP, 2000 Market Street, 10th Floor, Philadelphia, Pennsylvania 19103 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: NOVEL COMPOUNDS, PHARMACEUTICAL COMPOSITIONS CONTAINING SAME, METHODS OF USE FOR SAME, AND METHODS FOR PREPARING SAME

**FIG. 1**

(57) Abstract: The present invention relates to a novel class of compounds comprising formula I, wherein n is 0 or 1. A is NR<sup>1</sup>, O, or S, wherein R<sup>1</sup> is H, hydroxyl, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, alkenyl, aryl, alkylaryl or arylalkyl. X is a carboxylate, a phosphonate, or a phosphate residue, or a C<sub>1</sub>-C<sub>10</sub> alkyl residue optionally substituted with a carboxylate, phosphonate or phosphate residue. Y is a C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, halide, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl, or a heterocyclic ring and is optionally substituted with one or more halides. Z is a H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring and is optionally substituted with one or more C<sub>1</sub>-C<sub>10</sub> alkyl groups, C<sub>1</sub>-C<sub>10</sub>alkoxy groups, hydroxyl groups, cyano groups, carboxylate groups, halides, aryl groups, alkylaryl groups, arylalkyl groups, cycloalkyl groups, cycloalkenyl groups or heterocyclic rings.

**WO 2010/005922 A1**

**(84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

# **NOVEL COMPOUNDS, PHARMACEUTICAL COMPOSITIONS CONTAINING SAME, METHODS OF USE FOR SAME, AND METHODS FOR PREPARING SAME**

## **Priority Filing**

**[0001]** This application claims priority from U.S. Provisional Application No. 61/129,578, which was filed on July 7, 2008 and is incorporated herein by reference.

## **Field of the Invention**

**[0002]** The present invention relates to novel compounds, pharmaceutical compositions containing the same, and methods of use for a variety of therapeutically valuable uses including, but not limited to, treating obesity by inhibiting the activity of Glycerol 3-phosphate acyltransferase (GPAT).

## **Background of the Invention**

**[0003]** The incidence of obesity and other diseases associated with an increased triacylglycerol mass is increasingly recognized as a significant public health issue. Obesity is currently estimated by the World Health Organization to affect at least 400 million adults worldwide. In the U.S. alone, there are estimates that approximately two-thirds of adults are overweight or obese. Various diseases are associated with obesity, including type-2 diabetes, hypertension, cardiovascular diseases, nonalcoholic fatty liver disease, and certain types of cancer.

**[0004]** Even though there is a clear need for effective and widely available anti-obesity therapeutics, only two such drugs approved for long-term use in the U.S.: Orlistat functions by blocking the absorption of dietary fat, and sibutramine affects the central nervous system, reducing energy intake and increasing energy use. Although not completely ineffective, each of these drugs displays limited efficacy and produces undesirable side effects.

[0005] Anti-obesity drugs currently in development utilize a wide variety of mechanisms, involving both central and peripheral targets. Alteration of lipid metabolism, by decreasing the de novo synthesis of triglycerides while increasing oxidation of stored fats, is a peripheral mechanism. This approach, based on weight loss effects observed with the compounds C75, cerulenin, and hGH<sub>(177-191)</sub>, may be highly valuable in developing anti-obesity drugs.

[0006] Glycerol 3-phosphate acyltransferase (GPAT) catalyzes the rate-limiting step of glycerolipid biosynthesis, the acylation of glycerol 3-phosphate with saturated long chain acyl-CoAs. At present, there are four identified GPAT family members: GPAT1, a mitochondrial isoform catalyzing the bulk of hepatic triglyceride synthesis; GPAT2, a second mitochondrial isoform that synthesizes triglycerides but is less responsive to dietary control; GPAT3, localized to the endoplasmic reticulum, is responsible for the bulk of triglyceride synthesis in adipocytes, small intestine, kidney, and heart; and GPAT4, a microsomal isoform whose function is not completely elucidated. The mitochondrial isoform of glycerol-3-phosphate acyltransferase-1 (mtGPAT) catalyzes the esterification of long chain acyl-CoAs with *sn*-glycerol-3-phosphate to produce lysophosphatidic acid (LPA). This reaction is thought to constitute the first committed and rate-limiting step of glycerolipid biosynthesis. The purported mechanism of this reaction is similar to that of a serine protease, with the primary hydroxyl group of glycerol-3-phosphate taking the place of serine in the catalytic triad. Next, LPA is esterified further to produce phosphatidic acid, a precursor of various phospholipids including triacylglycerol (TAG), the main component of animal fat. In addition to obesity, high TAG levels in the bloodstream have been linked to several diseases, notably atherosclerosis and pancreatitis.

[0007] It has been shown that mtGPAT1 displays a strong preference for incorporating palmitoyl-CoA (16:0), thereby primarily producing saturated phospholipids, whereas the other

two enzymes are not selective. Of the three isoforms of GPAT, only mtGPAT1 is affected by changes in diet or exercise. When excess calories are available from a high-carbohydrate diet, mtGPAT1 mRNA expression increases, resulting in greater mtGPAT1 activity. It has been shown that mice that remain stationary for ten hours following a prolonged exercise regimen experience an increase in mtGPAT1 activity compared to mice that did not exercise at all, resulting in a significant overshoot of triacylglycerol (TAG) synthesis. MtGPAT1-deficient mice exhibit lower hepatic TAG levels and secrete less very low density lipoprotein (VLDL) than control mice. In contrast, rat hepatocytes with 2.7-fold increased mtGPAT1 activity demonstrated a significant increase in *de novo* synthesis of diacylglycerol. Overexpression of mtGPAT1 *in vivo*, as expected from the previous result, causes the levels of accumulated TAG and diacylglycerol (DAG) in mouse liver to rise dramatically to 12-fold and 7-fold that of normal levels. In addition to producing a certain amount of TAG dependent on the amount of active enzyme present, mtGPAT1 activity is essential for controlling the partitioning of fatty-acyl CoAs to  $\beta$ -oxidation or glycerolipid synthesis.


[0008] Both mtGPAT1 and carnitine palmitoyltransferase-1 (CPT-1), the enzyme that catalyzes the rate-limiting step of  $\beta$ -oxidation, are located on the outer mitochondrial membrane. This suggests that there is a competition between these enzymes for fatty acyl-CoA substrates. AMP-activated protein kinase (AMPK), which inactivates acetyl-CoA carboxylase (ACC) by phosphorylation, appears to acutely regulate both of these enzymes. Inactivation of ACC by AMPK prevents the buildup of malonyl-CoA, an allosteric suppressor of CPT-1, resulting in an increase in  $\beta$ -oxidation. AMPK inhibits mtGPAT1 as well, thereby decreasing the amount of TAG produced. The relationship between these two processes has been demonstrated *in vivo*. Feeding mtGPAT1-knockout mice a high-fat, high-sugar diet to induce obesity resulted in an

increase in oxidation as the long-chain acyl-CoA substrates were partitioned away from the TAG synthetic pathway toward CPT-1 and  $\beta$ -oxidation. MtGPAT1 overexpression in rat hepatocytes produced an 80% reduction in fatty acid oxidation coupled to an increase in phospholipid biosynthesis. Overexpression *in vivo* resulted in a decrease in  $\beta$ -oxidation as well.

[0009] The evidence suggesting that a drop in mtGPAT activity leads to a decrease in TAG levels as well as an increase in the amount of  $\beta$ -oxidation suggests that inhibition of this enzyme with a small molecule could be an effective treatment for obesity, diabetes, and other health problems associated with increased TAG synthesis. There is a need, therefore, for small molecules which can inhibit mtGPAT and other GPAT isoforms. Such compounds might be used for treating obesity or inducing weight loss.

### Summary of the Invention

[0010] The present invention relates to a novel class of compounds comprising formula I:



wherein n is either 0 or 1. A is selected from the group consisting of  $\text{NR}^1$ , O, and S, wherein  $\text{R}^1$  is either a H, hydroxyl,  $\text{C}_1\text{-C}_{10}$  alkyl,  $\text{C}_1\text{-C}_{10}$  alkoxy, alkenyl, aryl, alkylaryl or arylalkyl. X is selected from the group consisting of a carboxylate residue, a phosphonate residue, a phosphate residue, or a  $\text{C}_1\text{-C}_{10}$  alkyl residue which is optionally substituted with one or more carboxylate, phosphonate or phosphate residues. Y is selected from the group consisting of  $\text{C}_1\text{-C}_{20}$  alkyl, alkenyl, halide, hydroxyl,  $\text{C}_1\text{-C}_{20}$  alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl, or a heterocyclic ring and may optionally be substituted at one or more positions with a halide. Z is

selected from the group consisting of a H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring. In embodiments, where Z is an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring, the ring moiety may be substituted with one or more substituent groups selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, C<sub>1</sub>-C<sub>10</sub> alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring.

[0011] Based on the foregoing, one or more compounds of the present invention, either alone or in combination with another active ingredient, may be synthesized and administered as a therapeutic composition using dosage forms and routes of administration contemplated herein or otherwise known in the art. Dosaging and duration will further depend upon the factors provided herein and those ordinarily considered by one of skill in the art. To this end, determination of a therapeutically effective amounts are well within the capabilities of those skilled in the art, especially in light of the detailed disclosure and examples provided herein.

### Description of the Figures

[0012] Figure 1 illustrates a first reaction scheme for manufacturing compounds of the instant invention, particularly compounds 5a-5d disclosed herein.

[0013] Figure 2 illustrates a second reaction scheme for manufacturing compounds of the instant invention, particularly compounds 5e-5f disclosed herein.

[0014] Figure 3 illustrates a third reaction scheme for manufacturing compounds of the instant invention, particularly compounds 13a-13f disclosed herein.

[0015] Figure 4 illustrates a fourth reaction scheme for manufacturing compounds of the instant invention, particularly compounds 15a-15i disclosed herein.

[0016] Figure 5 illustrates a fifth reaction scheme for manufacturing compounds of the instant invention, particularly compounds 17a-17f disclosed herein.

[0017] Figure 6 illustrates a sixth reaction scheme for manufacturing compounds of the instant invention, particularly compounds 21a-21c disclosed herein.

[0018] Figure 7 illustrates a first reaction scheme for manufacturing compounds of the instant invention, particularly compounds 24a-24f disclosed herein.

[0019] Figure 8 illustrates a reaction scheme for manufacturing compounds 4a-t, disclosed herein.

[0020] Figure 9 illustrates a reaction scheme for manufacturing compounds 7a-t.

[0021] Figure 10 illustrates FSG67 inhibition of acylglyceride synthesis in 3T3-L1 adipocytes. The concentration dependent reduction of triglyceride synthesis is reflected in phase-contrast photomicrographs of cultured cells showing a corresponding reduction in lipid droplet accumulation (x 400).

[0022] Figure 11 illustrates acute FSG67 treatment of lean and DIO mice reduced body weight and decreased food consumption without conditioned taste aversion. Body weight and food intake were measured following a single 20 mg/kg ip dose of FSG67 in lean or DIO mice, 8 per group. (a) FSG67 treated lean mice (grey bar) lost  $3.7 \pm 0.9\%$  ( $1.0 \pm 0.2$  g); fasted mice lost  $15.5 \pm 0.7\%$  ( $3.9 \pm 0.2$  g) (black bar). The reduction in body mass of both treated and fasted mice was significant compared to the vehicle control mice (white bar) that gained  $2.5 \pm 0.5\%$  ( $0.6 \pm$

0.1 g) ( $p < 0.0001$  2-tailed t-test). **(b)** FSG67 treatment reduced food consumption to 33% of vehicle control ( $1.4 \pm 0.2$  g, grey bar, versus  $4.2 \pm 0.2$  g white bar,  $p < 0.0001$ , 2-tailed t-test). **(c)** FSG67 treated DIO mice (grey bar) lost  $4.3 \pm 0.5\%$  ( $1.7 \pm 0.2$  g) of body mass, fasted mice (black bar) lost  $5.3 \pm 0.4\%$  ( $2.1 \pm 0.2$  g) and vehicle controls (white bar) lost a  $2.5 \pm 0.6\%$  ( $1.0 \pm 0.2$  g). Compared to the vehicle controls, the weight loss was significant in both the FSG67 treated ( $p = 0.026$ , 2-tailed t-test) and fasted ( $p = 0.002$ , 2-tailed t-test) mice. **(d)** FSG67 reduced food consumption to 41.6% of vehicle control ( $0.5 \pm 0.1$  g, grey bar versus  $1.2 \pm 0.3$  g, white bar,  $p = 0.043$ , 2-tailed t-test). **(e)** FSG67 did not induce conditioned taste aversion in mice. CTA testing using a two bottle choice paradigm in groups of 8 lean mice did not produce a significant reduction in saccharine intake at 5 mg/kg ( $p = 0.12$ ) or 20 mg/kg ( $p = 0.10$ ) (2-tailed t-tests). Thus, the FSG67 effect on food intake was likely a specific effect on appetite rather than an induction of sickness behavior. All data are expressed as means  $\pm$  SEM. (\*,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ ).

**[0023]** Figure 12 illustrates chronic FSG67 treatment of DIO mice reversibly reduces body weight and food intake while enhancing fatty acid oxidation. **(a)** DIO mice, 4 per group, were treated with daily FSG67 5 mg/kg ip (red) or vehicle control (black) for 20 d (black arrow indicates termination of treatment) and were then allowed to regain their weight. The FSG67 treated mice lost  $10.3 \pm 0.6\%$  of body mass during treatment (days 0-19) compared to an increase of  $4.0 \pm 0.5\%$  for vehicle controls ( $p > 0.0001$ , 2-way ANOVA analysis). The FSG67 weight loss was reversible with treated animals returning to original weight at day 32. **(b)** Food consumption was significantly reduced during FSG67 treatment ( $2.6 \pm 0.1$  g/d) compared to vehicle controls ( $3.1 \pm 0.1$  g/d) ( $p = 0.0008$ , 2-way ANOVA). Following cessation of treatment at day 20, food consumption increased in the FSG67 treatment group to  $3.5 \pm 0.1$  g/d representing a significant

increase in food intake compared to vehicle controls  $3.2 \pm 0.1$  g/d ( $p=0.006$ , 2-way ANOVA). **(c)** Following 3 days of acclimatization in the calorimeter, 8 DIO mice per group were treated with FSG67 5 mg/kg ip (red) or vehicle control (black) daily for 16 days, along with a group pair-fed to the FSG67 treated animals (blue). The FSG67-treated animals lost  $9.5 \pm 0.6\%$  and pair-fed lost  $5.5 \pm 0.9\%$  of body mass while the vehicle controls increased by  $3.5 \pm 1.3\%$ . The weight loss in the FSG67 treated animals was significant compared to both vehicle controls and pair-fed animals ( $p<0.0001$ , 2-way ANOVA). **(d)** FSG67 treatment (red) reduced average daily food consumption by 33% ( $2.0 \pm 0.1$  g/d) compared to vehicle controls (black)  $3.1 \pm 0.1$  g/d ( $p<0.0001$ , 2-way ANOVA). **(e)** FSG67 treatment increased the average  $VO_2$  to  $106.5 \pm 1.1\%$  of the pre-treatment value (red line) compared to a reduction of  $89.9 \pm 1.1\%$  for the pair-fed group (blue line) ( $p<0.0001$  2-way ANOVA) consistent with increased energy utilization. **(f)** In contrast, the average RER was lower for the FSG67 treated DIO mice ( $0.732 \pm 0.002$ ) (red line) compared to ( $0.782 \pm 0.006$ ) (blue line) for the pair-fed group ( $p<0.0001$ , 2-way ANOVA) indicating increased reliance on fatty acids for fuel.

**[0024]** Figures 13 illustrates pharmacological GPAT inhibition reduced adiposity and down-regulated lipogenic gene expression in DIO mice. **(a)** Q-NMR analysis of FSG67 treated or vehicle control animals 10 per group. FSG67 treated animals (checkered bars) exhibited a significant reduction in fat mass (4.0 g) compared to vehicle controls (white bars) while lean and water mass were unaffected ( $p<0.0001$ , 2-tailed t-test). At the conclusion of the experiment, the vehicle control mice weighed 4.4 g more than the FSG67 controls ( $p=0.0014$ , 2-tailed t-test). **(b)** Real-time RT-PCR analysis of lipogenic gene expression in FSG67 treated (checkered bars), vehicle control (white bars), and pair-fed (black bars) DIO mice (from experiment shown in Fig. 4c). FSG67 reduced the expression of ACC1 ( $p=0.0005$  vs. control,  $p=0.0004$  vs. pair-fed), FAS

( $p=0.0001$  vs. control,  $p=0.0007$  vs. pair-fed),  $\text{PPAR}\gamma$  ( $p=0.032$  vs. control,  $p=0.0019$  vs. pair-fed), and GPAT ( $p=0.0034$  vs. control,  $p=0.0002$  vs. pair-fed) were all down regulated in white adipose tissue. Data are analyzed with 2-tailed t-tests. (\*,  $p<0.05$ ; \*\*,  $p<0.01$ ; \*\*\*,  $p<0.001$ ).

**[0025]** Figure 14 illustrates FSG67 treatment reduced hepatic steatosis and serum triglyceride and glucose levels. Oil red-stained histological sections of liver from **(a)** vehicle control, **(b)** pair-fed, and **(c)** FSG67-treated DIO mice from the 16-day treatment experiment in Fig. 4. Note intracytoplasmic large and small droplet fat accumulation most prominent in the vehicle control (a). Pair feeding reduced steatosis, whereas FSG67-treated animals showed almost complete amelioration of fat accumulation. **(d)** Average serum triglyceride, cholesterol, and glucose measurements from vehicle control, FSG67-treated, and pair-fed mice from the same animals. FSG67-treated animals had significantly reduced serum glucose levels ( $153.3 \pm 10.5$  mg/dL) compared to pair-fed mice ( $189.0 \pm 20.3$  mg/dL,  $p=0.047$ ) and vehicle controls ( $200.6 \pm 22.2$  mg/dL,  $p=0.031$  2-way ANOVA). The reduction in triglyceride levels were not statistically significant; cholesterol levels were unaffected. Data are expressed as means  $\pm$  SEM. (\*,  $p<0.05$ ; \*\*,  $p<0.01$ ; \*\*\*,  $p<0.001$ ).

**[0026]** Figure 15 illustrates Intracerebroventricular (icv) FSG67 treatment reduced food consumption and body weight. **(a)** FSG67 or vehicle was administered icv to groups of 6 lean mice. One day following treatment, mouse weight was significantly reduced by both 100 (vertical bars) and 320 nmole (checkered bar) doses ( $p=0.016$ ,  $p=0.0003$ , 2-tailed t-tests). **(b)** A significant reduction in food intake occurred only in the 320 nmole group (checkered bar) ( $p=0.005$ , 2-tailed t-test). (\*,  $p<0.05$ ; \*\*,  $p<0.01$ ; \*\*\*,  $p<0.001$ ).

[0027] Figure 16 illustrates acute and chronic FSG67 treatment altered hypothalamic neuropeptide expression. **(a)** Real-time RT-PCR analysis of hypothalamic neuropeptides were conducted in lean mice treated with a single 20 mg/kg dose of FSG67 (from Fig 3a). NPY was significantly reduced in the FSG67 treated group (grey bar) compared to fasted mice (black bar) ( $p=0.016$ ), while AGRP expression was diminished compared to both vehicle control (white bar) ( $p=0.02$ ) and fasted mice ( $p=0.0009$ ). Expression of POMC and CART were unaffected. **(b)** Similar analysis from 16 d treated DIO mice (from Fig. 4c) showed a reduction of NPY expression in both FSG67 treated ( $p=0.0074$ ) and pair-fed controls ( $p=0.0057$ ). The expression of AGRP, POMC, and CART were unaffected. Data are analyzed with 2-tailed t-tests. (\*,  $p<0.05$ ; \*\*,  $p<0.01$ ; \*\*\*,  $p<0.001$ ).

[0028] Figure 17 illustrates dose response of FSG67 in DIO mice. Groups of DIO mice were treated daily with FSG67 ip at doses indicated or vehicle. Over the 5 day course, 5 mg/kg was the minimum dose that led to a significant weight loss of 3.9% compared to vehicle controls ( $p=0.008$ , 2-way ANOVA). (\*  $p<0.05$ ; \*\*,  $p<0.01$ ; \*\*\*,  $p<0.001$ ).

[0029] Figure 18 illustrates FSG67 treatment of DIO mice for Q-NMR analysis. DIO mice (10 per group) treated daily for 10 d with FSG67 (5 mg/kg) lost significant body mass (6.1 g, 13.1%) compared to vehicle controls (1.1 g, 2.4%) ( $p<0.0001$ , 2-way ANOVA).

[0030] Figure 19 illustrates FSG67 treatment increases UCP2 expression in liver and WAT. Real-time RT-PCR expression analysis of LCPT-1 and UCP2 expression in liver and white adipose tissue. UCP2 expression was increased in the **(a)** liver ( $p=0.043$  vs. control) and **(b)** WAT ( $p=0.013$ , vs. pair-fed) of DIO mice treated with FSG67 for 16 d (see Fig. 4C). L-CPT-

1 expression was not affected by FSG67 treatment or pair-feeding. Data were analyzed with two-tailed t-tests,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ .

[0031] Figure 20 illustrates FSG67 treatment down-regulated hepatic lipogenic genes. Real-time RT-PCR expression analysis of lipogenic gene expression in the liver of DIO mice treated with FSG67 for 16 d (see Fig. 12C). FAS expression was reduced compared to both vehicle and pair-fed animals ( $p = 0.0016$  vs. control,  $p = 0.018$  vs. pair-fed) while ACC1 was reduced compared to pair-fed animals ( $p = 0.037$ ). GPAT expression was unaffected. Data were analyzed with two-tailed t-tests,  $p < 0.05$ ; \*\*,  $p < 0.01$ ; \*\*\*,  $p < 0.001$ .

### Detailed Description of the Invention

[0032] Definitions

[0033] As used herein, “an alkyl group” denotes both straight and branched carbon chains with one or more carbon atoms, but reference to an individual radical such as “propyl” embraces only the straight chain radical, a branched chain isomer such as “isopropyl” specifically referring to only the branched chain radical. A “substituted alkyl” is an alkyl group wherein one or more hydrogens of the alkyl group are substituted with one or more substituent groups as otherwise defined herein.

[0034] As used herein, “an alkoxy group” refers to a group of the formula alkyl-O-, where alkyl is as defined herein. A “substituted alkoxy” is an alkoxy group wherein one or more hydrogens are substituted with one or more of the substituent groups otherwise defined herein.

[0035] As used herein, “alkenyl” refers to a partially unsaturated alkyl radical derived by the removal of one or more hydrogen atoms from a alkyl chain such that it contains at least one carbon-carbon double bond.

[0036] As used herein, “an aryl group” denotes a structure derived from an aromatic ring containing six carbon atoms. Examples include, but are not limited to a phenyl or benzyl radical and derivatives thereof.

[0037] As used herein, “arylalkyl” denotes an aryl group having one or more alkyl groups not at the point of attachment of the aryl group.

[0038] As used herein, “alkylaryl” denotes an aryl group having an alkyl group at the point of attachment.

[0039] As used herein, “carboxylate” denotes salt or ester of an organic acid, containing the radical  $-\text{COOR}$ , wherein R may be, but is not limited to, a H, an alkyl group, an alkenyl group, or any other residue otherwise known in the art.

[0040] As used herein, “carboxylic acid” denotes an organic functional group comprising the following structure:  $-\text{COOH}$  or  $-\text{CO}_2\text{H}$ .

[0041] As used herein, “cyano” denotes an organic functional group comprising the following structure:  $-\text{C}\equiv\text{N}$ .

[0042] As used herein, “cycloalkyl” refers to a monovalent or polycyclic saturated or partially unsaturated cyclic non-aromatic group containing all carbon atoms in the ring structure, which may be substituted with one or more substituent groups defined herein. In certain non-limiting embodiments the number of carbons comprising the cycloalkyl group may be between 3 and 7.

[0043] As used herein, “cycloalkenyl” refers to a partially unsaturated cycloalkyl radical derived by the removal of one or more hydrogen atoms from a cycloalkyl ring system such that it contains at least one carbon-carbon double bond.

[0044] As used herein, “halogen” or “halide” denotes any one or more of a fluorine, chlorine, bromine, or iodine atoms.


[0045] As used herein, “heterocyclic” refers to a monovalent saturated or partially unsaturated cyclic aromatic or non-aromatic carbon ring group which contains at least one heteroatom, in certain embodiments between 1 to 4 heteroatoms, which may be but is not limited to one or more of the following: nitrogen, oxygen, sulfur, phosphorus, boron, chlorine, bromine, or iodine. In further non-limiting embodiments, the heterocyclic ring may be comprised of between 1 and 10 carbon atoms.

[0046] As used herein, “hydroxyl” denotes an organic functional group comprising the following structure:  $\text{—OH}$ .

[0047] As used herein, “phosphonate” denotes an organic functional group comprising the following structure:  $\text{—PO}_3\text{H}_2$  or  $\text{—PO(OH)}_2$ .

[0048] As used herein, “phosphate” denotes an organic functional group comprising the following structure:  $\text{—OPO}_3\text{H}_2$  or  $\text{—OPO(OH)}_2$ .

[0049] The present invention relates to novel compounds, pharmaceutical compositions containing the same, and methods of use by inhibiting the enzymatic activity of Glycerol 3-phosphate acyltransferase (GPAT). Such compounds, compositions, and methods have a variety of therapeutically valuable uses including, but not limited to, treating obesity. The class of compounds of the present invention are comprised of formula I:




wherein n is either 0 or 1. A is selected from the group consisting of NR<sup>1</sup>, O, and S, wherein R<sup>1</sup> is either a H, hydroxyl, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, alkenyl, aryl, alkylaryl or arylalkyl. X is selected from the group consisting of a carboxylate residue, a phosphonate residue, a phosphate residue, or a C<sub>1</sub>-C<sub>10</sub> alkyl residue which is optionally substituted with one or more carboxylate, phosphonate or phosphate residues. Y is selected from the group consisting of C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, halide, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl, or a heterocyclic ring. In embodiments where Y is a C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl, or a heterocyclic ring, it is optionally substituted at one or more positions with a halide. Z is selected from the group consisting of a H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring. In embodiments, where Z is an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring, the ring moiety may be substituted with one or more substituent groups selected from a C<sub>1</sub>-C<sub>10</sub> alkyl group, C<sub>1</sub>-C<sub>10</sub> alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring.

**[0050]** In certain embodiments, X is comprised of either a carboxylic acid residue or a phosphonate residue. In alternative embodiments, X may include a C<sub>1</sub>-C<sub>10</sub> alkyl group, which is substituted at one or more positions with either a phosphonate residue or carboxylate. In further embodiments, the alkyl group may comprise between 1 and 3 carbons. In any of the foregoing, X may be positioned on the phenyl ring in either the ortho, meta, or para position with respect to the sulfonyl linker. As shown below, in certain non-limiting embodiments X occupies either the ortho or meta position.


[0051] In further non-limiting embodiments, Y is comprised of a C<sub>1</sub>-C<sub>20</sub> alkyl group, which may be either a CH<sub>3</sub>, C<sub>5</sub>H<sub>11</sub>, C<sub>8</sub>H<sub>17</sub>, C<sub>9</sub>H<sub>19</sub>, C<sub>14</sub>H<sub>29</sub>, and C<sub>16</sub>H<sub>33</sub>. Alternatively, Y may be comprised of an aryl ring system, which is optionally substituted with one or more halogen atoms. In even further alternative embodiments, Y is comprised of an alkylaryl residue, wherein the alkyl moiety connects the aryl ring to the Y position. The alkyl chain may have between 1 to 3 carbon atoms, with certain embodiments having 1 or 2 carbon atoms. The aryl residue in this latter embodiment may be substituted with one or more halogen atoms.

[0052] In even further non-limiting embodiments, Z is either a hydrogen atom, a hydroxyl group, a halogen atom, an optionally substituted aryl group or an optionally substituted heterocyclic ring. In any of the foregoing, Z may be position on the phenyl ring in either the ortho, meta, or para position with respect to the sulfonyl linker. As shown below, in certain non-limiting embodiments Z occupies either the meta or para position with respect to the sulfonyl linker of the phenyl ring. In even further embodiments, Z occupies either the meta or para position with respect to both the sulfonyl linker and X positions.


[0053] Based on the foregoing, one compound of the instant invention is C-67 or FSG67 and is comprised of the following structure:





[0054] In another embodiment, the compounds of the instant invention may be comprised of the following structures:




[0055] In an even further embodiment, the compounds of the instant invention may be comprised of one or more of the following:











wherein each of n, X, Y, and Z are any of the embodiments defined above.

[0057] In alternative embodiments of formula I, n is comprised of 0. To this end, certain compounds of the instant invention may be represented by formula III:




wherein each of A, X, Y, and Z are any of the embodiments defined above.

[0058] In even further embodiments of formula I, X is comprised of a carboxylic acid residue at either the ortho, meta or para positions with respect to the sulfonyl linker of the phenyl ring. Accordingly, certain compounds of the instant invention may be represented by formula IVa:





wherein each of n, A, Y, and Z are any of the embodiments defined above.

[0059] While the carboxylic acid residue may occupy either the ortho, meta, or para positions, in certain embodiments it occupies the ortho position with respect to the sulfonyl linker. To this end, certain compounds of the instant invention may be represented by formula IVb:




wherein each of n, A, Y, and Z are any of the embodiments defined above.


**[0060]** Similarly, although it may occupy either the ortho, meta, or para positions, in certain compounds of the instant invention Z occupies either the meta or the para positions with respect to both the sulfonyl linker and X, as set forth below in formulas IVc and IVd:





wherein each of n, A, Y, and Z are any of the embodiments defined above.


**[0061]** Based on the foregoing structures of formulas IVc-d compounds of the instant invention may be comprised of one or more of the following:







**[0063]** Accordingly, compounds of the instant invention may be comprised of one or more of the following:



[0064] Without seeking to limit the possible scope of use of the foregoing compounds, the clinical therapeutic indications envisioned include, but are not limited to, treatment of obesity or the induction of weight loss. One or more small molecules, or pharmaceutical salts thereof, of the present invention may be synthesized and administered as a composition used to treat and/or prevent obesity by targeted GPAT activity, in particular mtGPAT activity, and/or by stimulating fatty acid oxidation. Compounds of the present invention may be synthesized using methods known in the art or as otherwise specified herein.

[0065] Unless otherwise specified, a reference to a particular compound of the present invention includes all isomeric forms of the compound, to include all diastereomers, tautomers, enantiomers, racemic and/or other mixtures thereof. Unless otherwise specified, a reference to a particular compound also includes ionic, salt, solvate (e.g., hydrate), protected forms, and prodrugs thereof. To this end, it may be convenient or desirable to prepare, purify, and/or handle a corresponding salt of the active compound, for example, a pharmaceutically-acceptable salt. Examples of pharmaceutically acceptable salts are discussed in Berge et al., 1977, "Pharmaceutically Acceptable Salts," J. Pharm. Sci., Vol. 66, pp. 1-19, the contents of which are incorporated herein by reference.

[0066] Based on the foregoing, one or more compounds of the present invention, either alone or in combination with another active ingredient, may be synthesized and administered as a therapeutic composition. The compositions of the present invention can be presented for administration to humans and other animals in unit dosage forms, such as tablets, capsules, pills, powders, granules, sterile parenteral solutions or suspensions, oral solutions or suspensions, oil in water and water in oil emulsions containing suitable quantities of the compound, suppositories and in fluid suspensions or solutions. To this end, the pharmaceutical compositions may be

formulated to suit a selected route of administration, and may contain ingredients specific to the route of administration. Routes of administration of such pharmaceutical compositions are usually split into five general groups: inhaled, oral, transdermal, parenteral and suppository. In one embodiment, the pharmaceutical compositions of the present invention may be suited for parenteral administration by way of injection such as intravenous, intradermal, intramuscular, intrathecal, or subcutaneous injection. Alternatively, the composition of the present invention may be formulated for oral administration as provided herein or otherwise known in the art.

[0067] As used in this specification, the terms "pharmaceutical diluent" and "pharmaceutical carrier," have the same meaning. For oral administration, either solid or fluid unit dosage forms can be prepared. For preparing solid compositions such as tablets, the compound can be mixed with conventional ingredients such as talc, magnesium stearate, dicalcium phosphate, magnesium aluminum silicate, calcium sulfate, starch, lactose, acacia, methylcellulose and functionally similar materials as pharmaceutical diluents or carriers. Capsules are prepared by mixing the compound with an inert pharmaceutical diluent and filling the mixture into a hard gelatin capsule of appropriate size. Soft gelatin capsules are prepared by machine encapsulation of a slurry of the compound with an acceptable vegetable oil, light liquid petrolatum or other inert oil.

[0068] Fluid unit dosage forms or oral administration such as syrups, elixirs, and suspensions can be prepared. The forms can be dissolved in an aqueous vehicle together with sugar or another sweetener, aromatic flavoring agents and preservatives to form a syrup. Suspensions can be prepared with an aqueous vehicle with the aid of a suspending agent such as acacia, tragacanth, methylcellulose and the like.

[0069] For parenteral administration fluid unit dosage forms can be prepared utilizing the compound and a sterile vehicle. In preparing solutions the compound can be dissolved in water for injection and filter sterilized before filling into a suitable vial or ampoule and sealing.

Adjuvants such as a local anesthetic, preservative and buffering agents can be dissolved in the vehicle. The composition can be frozen after filling into a vial and the water removed under vacuum. The lyophilized powder can then be sealed in the vial and reconstituted prior to use.

[0070] Dose and duration of therapy will depend on a variety of factors, including (1) the patient's age, body weight, and organ function (e.g., liver and kidney function); (2) the nature and extent of the disease process to be treated, as well as any existing significant co-morbidity and concomitant medications being taken, and (3) drug-related parameters such as the route of administration, the frequency and duration of dosing necessary to effect a cure, and the therapeutic index of the drug. In general, the dose will be chosen to achieve serum levels of 1 ng/ml to 100 ng/ml with the goal of attaining effective concentrations at the target site of approximately 1 gg/ml to 10 µg/ml. Using factors such as this, a therapeutically effective amount may be administered so as to ameliorate the targeted symptoms of and/or treat or prevent obesity or diseases related thereto. Determination of a therapeutically effective amount is well within the capabilities of those skilled in the art, especially in light of the detailed disclosure and examples provided herein.

### **Examples**

[0071] *Example 1 - Chemical Syntheses of compounds 5a-5d*

[0072] Synthesis of compounds 5a-5d was performed using Scheme 1, as illustrated in figure 1 herein.

[0073] Reaction conditions: (a) NBS, hv, CH<sub>3</sub>CN; (b) NaN<sub>3</sub>, EtOH, reflux; (d) C<sub>9</sub>H<sub>19</sub>SO<sub>2</sub>Cl or C<sub>5</sub>H<sub>11</sub>SO<sub>2</sub>Cl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to room temperature; (e) K<sup>+</sup>O<sup>-</sup>*t*-Bu, Et<sub>2</sub>O, H<sub>2</sub>O, 0 °C to room temperature.

[0074] The first series of compounds was derived from the variously substituted methyl methylbenzoates. The *meta*- and *para*-amines were made by following a literature protocol. (Okada, Y. et al., Bromination by means of sodium monobromoisocyanurate (SMBI). *Org. Biomolec. Chem.* **2003**, *1*, 2506-2511.) Following radical bromination of the methyl group with NBS in CH<sub>3</sub>CN, the bromide was displaced by refluxing with NaN<sub>3</sub> in EtOH. Under Staudinger conditions, the azide was reduced to the free amine **3**, which could then be coupled to 1-pentane- or 1-nonanesulfonyl chloride, prepared as described. (Blotny, G., A new, mild preparation of sulfonyl chlorides, *Tet. Lett.* **2003**, *44*, 1499-1501.) Finally, the methyl ester **4** was converted to the carboxylate product **5** by reaction with potassium *t*-butoxide in Et<sub>2</sub>O with water present.

[0075] **General Procedure for 4a-d.** To a stirring solution of the appropriate amine **3a-c** (1.2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) at 0 °C, the sulfonyl chloride (1.3 mmol) was added dropwise, followed by Et<sub>3</sub>N (1.3 mmol). The reaction mixture was allowed to warm to room temperature, where it was stirred for 2-3 h. Saturated NH<sub>4</sub>Cl solution was added to quench the reaction, and the mixture was extracted with 3 × 10 mL CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layers were dried over MgSO<sub>4</sub>, concentrated *in vacuo*, and the products were purified by flash chromatography (20% EtOAc in hexanes).

[0076] **General Procedure for 5a-d.** To a stirring suspension of potassium *t*-butoxide (5.88 mmol) in Et<sub>2</sub>O (15mL) cooled to 0 °C, was added water (1.4 mmol) via syringe. The slurry was stirred for 5 min, and **4a-d** (0.67 mmol) was added. The mixture was stirred at room

temperature until starting material disappeared by TLC analysis (20% EtOAc in hexanes). Ice water was added until 2 clear layers formed. The aqueous layer was separated and acidified with 1 M HCl. The product was then extracted with Et<sub>2</sub>O (3 × 20 mL) and evaporated *in vacuo* to afford **5a-d**.

[0077]        **4-(Pentylsulfonamidomethyl)benzoic acid 5a.** mp = 188-189 °C; <sup>1</sup>H NMR (MeOD) δ 8.02 (d, *J* = 8.4 Hz, 2H), 7.50 (d, *J* = 8.1 Hz, 2H), 4.31 (s, 2H), 2.95 (t, *J* = 8.1 Hz, 2H), 1.73 (m, 2H), 1.33 (m, 4H), 0.91 (t, *J* = 6.9 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 169.5, 145.0, 131.1, 131.0, 128.8, 53.6, 47.2, 31.4, 24.3, 23.2, 14.0; HRMS (FAB) calcd for C<sub>13</sub>H<sub>20</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 286.11131; found, 286.1111.

[0078]        **4-(Nonylsulfonamidomethyl)benzoic acid 5b.** mp = 178-180 °C; <sup>1</sup>H NMR (MeOD) δ 8.03 (d, *J* = 8.4 Hz, 2H), 7.51 (d, *J* = 8.1 Hz, 2H), 4.32 (s, 2H), 2.94 (t, *J* = 7.8 Hz, 2H), 1.71 (m, 2H), 1.30 (m, 12H), 0.92 (t, *J* = 6.9 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 167.0, 143.6, 129.5, 129.3, 127.5, 51.5, 45.4, 31.2, 28.6, 28.5, 28.4, 27.4, 23.0, 22.0, 13.9; HRMS (FAB) calcd for C<sub>17</sub>H<sub>28</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 342.17391; found, 342.17447.

[0079]        **3-(Pentylsulfonamidomethyl)benzoic acid 5c.** mp = 160-161 °C; <sup>1</sup>H NMR (MeOD) δ 8.08 (s, 1H), 7.97 (d, *J* = 7.8 Hz, 1H), 7.63 (d, *J* = 7.8 Hz, 1H), 7.48 (t, *J* = 7.8 Hz, 1H), 4.31 (s, 2H), 2.92 (t, *J* = 8.1 Hz, 2H), 1.72 (m, 2H), 1.33 (m, 4H), 0.91 (t, *J* = 6.9 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 169.5, 140.2, 133.5, 132.3, 130.1, 129.8, 129.7, 53.6, 47.1, 31.4, 24.3, 23.1, 14.0; HRMS (FAB) calcd for C<sub>13</sub>H<sub>18</sub>NO<sub>3</sub>S [M – OH]<sup>+</sup>, 268.10074; found, 268.09988.

[0080]        **3-(Nonylsulfonamidomethyl)benzoic acid 5d.** mp = 150-151 °C; <sup>1</sup>H NMR (MeOD) δ 8.08 (s, 1H), 7.97 (d, *J* = 7.6 Hz, 1H), 7.63 (d, *J* = 7.6 Hz, 1H), 7.48 (t, *J* = 7.6 Hz, 1H), 4.31 (s, 2H), 2.91 (t, *J* = 8.0 Hz, 2H), 1.70 (m, 2H), 1.28 (m, 12H), 0.92 (t, *J* = 7.2 Hz); <sup>13</sup>C

NMR (MeOD)  $\delta$  169.5, 140.2, 133.5, 132.3, 130.1, 129.9, 129.7, 53.7, 47.2, 32.9, 30.4, 30.3, 30.1, 29.2, 24.6, 23.6, 14.4; HRMS (FAB) calcd for C<sub>17</sub>H<sub>28</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 342.17391; found, 342.17333.

**[0081]**        *Example 2 - Synthesis of Compounds 5e and 5f*

**[0082]**        Synthesis of compounds 5e-5f was performed using Scheme 2, as illustrated in figure 2 herein.

**[0083]**        Reaction conditions: (a) NH<sub>3</sub>, MeOH, reflux; (b) NaH, RSO<sub>2</sub>Cl, DMF, 0 °C to room temperature; (c) NaOH, THF/H<sub>2</sub>O, 0 °C to room temperature.

**[0084]**        The *ortho*-substituted carboxylates required a different approach than the *meta*- and *para*- compounds. Indolinone **6**, formed in a reaction between the *ortho*-bromide and ammonia gas in MeOH, (Kovtunenkov, V. A., et al.; Condensation of o-(bromomethyl)benzoic acid with amines, *Ukrainskii Khimicheskii Zhurnal* **1983**, 49, 1099-1103) was coupled to the alkane sulfonyl chlorides with NaH in DMF, and the resulting  $\gamma$ -lactam bond was readily cleaved with NaOH in THF/H<sub>2</sub>O to produce carboxylic acids **5e** and **5f**.

**[0085]**        **General Procedure for 7a-b.** 1.5 mmol **6** was added to DMF (8 mL), and the solution was cooled to 0 °C. NaH (1.65 mmol) was added, followed by the sulfonyl chloride (1.8 mmol), and the mixture was stirred and allowed to warm to room temperature. Reaction progress was monitored by TLC (25% MeOH in CHCl<sub>3</sub>). When complete, saturated ammonium chloride solution was added (80 mL), the product was extracted with EtOAc (3 × 20 mL), dried over MgSO<sub>4</sub>, and evaporated *in vacuo*. The product was purified by flash chromatography (2% MeOH in CHCl<sub>3</sub>).

[0086] **General Procedure for 5e-f.** **7a-b** (0.66 mmol) was dissolved in THF (3 mL), and the solution was cooled to 0 °C. 1 M NaOH (1 mL, 10 equiv) was then added, and the solution was stirred and warmed to room temperature. Reaction progress was monitored by TLC (1:1 EtOAc:hexanes). When starting material had completely reacted, saturated NaHCO<sub>3</sub> (30 mL) was added, and the solution was washed with EtOAc. The aqueous phase was acidified to pH 3 with 1 M HCl, and product was extracted with EtOAc, dried over MgSO<sub>4</sub>, and evaporated *in vacuo*.

[0087] **2-(Pentylsulfonamidomethyl)benzoic acid 5e.** mp = 100 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.0 (s, 1H), 7.87 (d, *J* = 8.1 Hz, 1H), 7.60 (m, 2H), 7.39 (m, 2H), 4.51 (d, *J* = 6.3 Hz, 2H), 2.92 (t, *J* = 7.8 Hz, 2H), 1.61 (m, 2H), 1.25 (m, 4H), 0.84 (t, *J* = 6.9 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 170.3, 140.8, 133.6, 132.3, 131.3, 130.7, 128.8, 53.6, 46.5, 31.3, 24.3, 23.1, 14.0; HRMS (FAB) calcd for C<sub>13</sub>H<sub>20</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 286.11131; found, 286.11103.

[0088] **2-(Nonylsulfonamidomethyl)benzoic acid 5f.** mp = 79-82 °C; <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 8.04 (d, *J* = 7.2 Hz, 1H), 7.60 (m, 2H), 7.43 (t, *J* = 6.8 Hz, 1H), 4.60 (s, 2H), 2.89 (t, *J* = 8.0 Hz, 2H), 1.66 (m, 2H), 1.28 (m, 12H), 0.92 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 168.3, 139.7, 132.1, 130.4, 129.8, 129.0, 127.2, 51.7, 44.1, 31.3, 28.7, 28.6, 28.5, 27.6, 23.1, 22.1, 14.0; HRMS (FAB) calcd for C<sub>17</sub>H<sub>28</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 342.17391; found, 342.17478.

[0089] **Example 3 - Synthesis of Compounds 13a-13f**

[0090] Synthesis of compounds 13a-13f was performed using Scheme 3, as illustrated in figure 3 herein.

[0091] Reaction conditions: (a) NBS, hv, CH<sub>3</sub>CN; (b) P(OEt)<sub>3</sub>, reflux; (c) H<sub>2</sub>SO<sub>4</sub>, EtOH, reflux; (d) C<sub>9</sub>H<sub>19</sub>SO<sub>2</sub>Cl or C<sub>5</sub>H<sub>11</sub>SO<sub>2</sub>Cl, pyridine, CH<sub>3</sub>CN, 0 °C to room temperature; (e) TMSBr, CH<sub>2</sub>Cl<sub>2</sub>, room temperature.

[0092] The synthesis of the alkyl phosphonates **13a-f** commenced with the protection of the starting toluidines as the bis-acylated aniline **8** (Brown, J. J.; Brown, R. K. Preparation of o- and p-acetamidobenzaldehydes, *Can. J. Chem.* **1955**, 33, 1819-1823). Free-radical bromination with NBS in CH<sub>3</sub>CN afforded benzyl bromide **9**, which was converted to phosphonate **10** through Arbuzov reaction with triethyl phosphite. The aniline was unmasked by exposure to a refluxing acidic solution of EtOH. Following coupling of the amine with the alkane sulfonyl chloride to produce sulfonamide **12**, the phosphonic acid moiety was revealed by treatment with TMSBr in CH<sub>2</sub>Cl<sub>2</sub> followed by methanolysis.

[0093] **General Procedure for 9a-c.** **8a-c** (31.3 mmol) was dissolved in CH<sub>3</sub>CN (150mL) and NBS (31.3 mmol) was added. The solution was then heated to reflux with a 275 W Sunlamp. Reaction progress was monitored by TLC (30% EtOAc in hexanes). The solution was then cooled, evaporated *in vacuo*, and the mixture was purified by flash chromatography (30% EtOAc in hexanes).

[0094] **General Procedure for 10a-c.** **9a-c** (22.2 mmol) was dissolved in P(OEt)<sub>3</sub> (25 mL, 6.6 equiv), and the solution was heated to reflux for 18 h with a reflux condenser heated to 50 °C. Reaction progress was monitored by TLC (30% EtOAc in hexanes). The reaction mixture was then cooled, and P(OEt)<sub>3</sub> was removed *in vacuo*. The product was then purified by flash chromatography (2% MeOH in CHCl<sub>3</sub>).

[0095]        **General Procedure for 11a-c.** Concentrated H<sub>2</sub>SO<sub>4</sub> (3 mL) was added to a stirring solution of **10a-c** (9.7 mmol) in EtOH (60 mL). The solution was heated to reflux for 18 h. Reaction progress was monitored by TLC (5% MeOH in CHCl<sub>3</sub>). The solution was diluted with water (100 mL), washed with EtOAc (30 mL), and the aqueous phase was brought to pH 9 with saturated NaHCO<sub>3</sub> solution. The product was extracted with EtOAc (3 × 30 mL), the combined organic layers were dried over MgSO<sub>4</sub>, and solvent was removed *in vacuo*.

[0096]        **General Procedure for 12a-b.** **11a** (1.36 mmol) was dissolved in CH<sub>3</sub>CN (3.3 mL), then pyridine (10.8 mmol) was added. The solution was cooled to 0 °C, and sulfonyl chloride (1.63 mmol) was added slowly by syringe. The solution was allowed to warm to room temperature. Reaction progress was monitored by TLC (5% MeOH in CHCl<sub>3</sub>). When complete, the reaction was quenched by adding saturated NaHCO<sub>3</sub> solution. The product was extracted with EtOAc (3 × 5 mL), washed with 1 N HCl, and the combined organic extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo*. The product was purified by flash chromatography (2% MeOH in CHCl<sub>3</sub>).

[0097]        **General Procedure for 12c-f.** Sulfonyl chloride (4.9 mmol) was added dropwise to a solution of **11b-c** (3.3 mmol) in CH<sub>3</sub>CN (13 mL) at 0 °C. Et<sub>3</sub>N (3.63 mmol) was added dropwise, and the solution was stirred and allowed to warm to room temperature. Reaction progress was monitored by TLC (10% MeOH in CHCl<sub>3</sub>). When complete (about 2 h), the reaction was quenched by adding saturated sodium bicarbonate solution. The product was extracted with EtOAc (3 × 10 mL), and the combined organic extracts were dried over MgSO<sub>4</sub> and concentrated *in vacuo*. Flash chromatography (2% MeOH in CHCl<sub>3</sub>) afforded the product.

[0098] **General Procedure for 13a-f.** TMSBr (8.6 mmol) was added to a solution of **12a-f** (0.277 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2 mL), and the solution was stirred at room temperature. After 24 h, the reaction was quenched by adding MeOH (3 × 1.6 mL). The solution was concentrated *in vacuo*, and dissolved in saturated NaHCO<sub>3</sub> solution (10 mL). This solution was washed with Et<sub>2</sub>O (5 mL), then acidified with 1 N HCl. The product was extracted with Et<sub>2</sub>O (3 × 5 mL), and the combined organic extracts were dried over MgSO<sub>4</sub> and dried *in vacuo*.

[0099] **2-(Pentylsulfonamido)benzylphosphonic acid 13a.** <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.73 (s, 1H), 7.38 (d, *J* = 8.0 Hz, 1H), 7.25 (m, 2H), 7.13 (t, *J* = 7.6 Hz, 1H), 3.14 (t, *J* = 8.0 Hz, 2H), 3.10 (d, *J* = 20.8 Hz, 2H), 1.71 (m, 2H), 1.32 (m, 4H), 0.85 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 136.2 (d, *J* = 5.8 Hz), 131.6 (d, *J* = 6.4 Hz), 127.6 (d, *J* = 10.0 Hz), 127.1 (d, *J* = 3.4 Hz), 125.0 (d, *J* = 2.8 Hz), 123.7 (d, *J* = 3.0 Hz), 52.3, 32.5 (d, *J* = 130.3 Hz), 29.4, 22.7, 21.3, 13.3.

[00100] **2-(Nonylsulfonamido)benzylphosphonic acid 13b.** mp = 104-106 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 9.80 (s, 1H), 7.37 (d, *J* = 8.0 Hz, 1H), 7.25 (m, 2H), 7.15 (t, *J* = 7.6 Hz, 1H), 3.14 (t, *J* = 7.6 Hz, 2H), 3.09 (d, *J* = 21.2 Hz, 2H), 1.68 (m, 2H), 1.35 (m, 2H), 1.22 (m, 8H), 0.85 (t, *J* = 6.8 Hz); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 136.3 (d, *J* = 5.7 Hz), 131.8 (d, *J* = 6.5 Hz), 127.7 (d, *J* = 8.7 Hz), 127.3 (d, *J* = 3.3 Hz), 125.2 (d, *J* = 2.6 Hz), 123.8 (d, *J* = 3.0 Hz), 52.3, 32.6 (d, *J* = 130.5 Hz), 31.2, 28.6, 28.5, 28.4, 27.4, 23.2, 22.0, 13.9.

[00101] **3-(Pentylsulfonamido)benzylphosphonic acid 13c.** mp = 127-128 °C; <sup>1</sup>H NMR (MeOD) δ 7.27 (t, *J* = 8.0 Hz, 1H), 7.22 (s, 1H), 7.11 (m, 2H), 3.11 (d, *J* = 21.6 Hz, 2H), 3.09 (t, *J* = 7.8 Hz, 2H), 1.78 (m, 2H), 1.38 (m, 4H), 0.91 (t, *J* = 6.4 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 139.5 (d, *J* = 3.3 Hz), 136.0 (d, *J* = 9.3 Hz), 130.3 (d, *J* = 3.4 Hz), 126.8 (d, *J* = 5.9 Hz), 122.5 (d, *J* =

6.5 Hz), 119.3 (d,  $J = 3.4$  Hz), 51.9, 35.8 (d,  $J = 134.2$  Hz), 31.2, 24.2, 23.1, 14.0; HRMS (FAB) calcd for  $C_{12}H_{21}NO_5PS$   $[M + H]^+$ , 322.08781; found, 322.08830.

**[00102] 3-(Nonylsulfonamido)benzylphosphonic acid 13d.** mp = 149-150 °C;  $^1H$  NMR (MeOD)  $\delta$  7.27 (t,  $J = 8.0$  Hz, 1H), 7.22 (s, 1H), 7.11 (m, 2H), 3.11 (d,  $J = 21.6$  Hz, 2H), 3.08 (t,  $J = 7.6$  Hz, 2H), 1.77 (m, 2H), 1.33 (m, 12H), 0.91 (t,  $J = 6.4$  Hz, 3H);  $^{13}C$  NMR (MeOD)  $\delta$  139.5 (d,  $J = 3.0$  Hz), 130.3 (d,  $J = 3.0$  Hz), 126.8 (d,  $J = 6.1$  Hz), 122.5, (d,  $J = 6.3$  Hz), 119.3 (d,  $J = 3.5$  Hz), 51.9, 35.8 (d,  $J = 134.0$  Hz), 32.9, 30.3, 30.2, 30.1, 29.1, 24.5, 23.6, 14.3; HRMS (FAB) calcd for  $C_{16}H_{29}NO_5PS$   $[M + H]^+$ , 378.15041; found, 378.14975.

**[00103] 4-(Pentylsulfonamido)benzylphosphonic acid 13e.** mp = 198-200 °C;  $^1H$  NMR (MeOD)  $\delta$  7.29 (dd,  $J = 8.4, 2.4$  Hz, 2H), 7.20 (d,  $J = 8.4$  Hz, 2H), 3.10 (d,  $J = 21.6$  Hz, 2H), 3.05 (t,  $J = 8.0$  Hz, 2H), 1.78 (m, 2H), 1.34 (m, 4H), 0.90 (t,  $J = 7.2$  Hz, 3H);  $^{13}C$  NMR (MeOD)  $\delta$  137.9 (d,  $J = 3.7$  Hz), 131.8 (d,  $J = 6.3$  Hz), 130.6 (d,  $J = 9.6$  Hz), 121.4 (d,  $J = 2.9$  Hz), 51.8, 35.1 (d,  $J = 134.6$  Hz), 31.2, 24.2, 23.1, 14.0; HRMS (FAB) calcd for  $C_{12}H_{20}NO_5PS$   $[M]^+$ , 321.07998; found, 321.07934.

**[00104] 4-(Nonylsulfonamido)benzylphosphonic acid 13f.** mp = 201-203 °C;  $^1H$  NMR (MeOD)  $\delta$  7.29 (dd,  $J = 8.8, 2.4$  Hz, 2H), 7.20 (d,  $J = 8.4$  Hz, 2H), 3.09 (d,  $J = 21.2$ , 2H), 3.05 (t,  $J = 8.0$  Hz, 2H), 1.77 (m, 2H), 1.29 (m, 12H), 0.91 (t,  $J = 7.2$  Hz, 3H);  $^{13}C$  NMR (MeOD)  $\delta$  137.9 (d,  $J = 3.3$  Hz), 131.8 (d,  $J = 6.5$  Hz), 130.6 (d,  $J = 9.3$  Hz), 121.4 (d,  $J = 3.0$  Hz), 51.8, 35.1 (d,  $J = 134.5$  Hz), 32.9, 30.3, 30.2, 30.1, 29.1, 24.5, 14.4; HRMS (FAB) calcd for  $C_{16}H_{29}NO_5PS$   $[M + H]^+$ , 378.15041; found, 378.14945.

**[00105] Example 4 - Synthesis of Compounds 15a-15i**

[00106] Synthesis of compounds 15a-15i was performed using Scheme 4, as illustrated in figure 4 herein.

[00107] Reaction conditions: (a) RSO<sub>2</sub>Cl, pyridine, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to rt; (b) K<sup>+</sup>O<sup>-</sup>*t*-Bu, Et<sub>2</sub>O, H<sub>2</sub>O, 0 °C to room temperature.

[00108] Compounds **15a-i** were synthesized by coupling the commercially available starting aniline with a variety of sulfonyl chlorides. The resulting sulfonamides **14a-i** were then converted to the final products by hydrolysis with potassium *t*-butoxide and water in ether. Aromatic sulfonyl chlorides were used as well as the saturated C<sub>9</sub> chain in an attempt to mimic the CoA portion of the acyl-CoA substrate, as opposed to the alkyl chain.

[00109] **General Procedure for 14a-i.** To a stirring solution of the aniline starting material (3.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (12 mL) at 0° C was added pyridine (7.5 equiv) was added. The sulfonyl chloride (1.2 equiv) was then added slowly via syringe. The solution was stirred and allowed to warm to room temperature. Reaction progress was monitored by TLC (20% EtOAc in hexanes). When complete, the reaction was poured into saturated NaHCO<sub>3</sub> solution (45 mL), extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL), and washed with 1 M HCl (50 mL). The combined organic phases were concentrated *in vacuo*, and recrystallization from EtOAc / hexanes afforded **14a-i**.

[00110] **General Procedure for 15a-i.** To a stirring suspension of potassium *t*-butoxide (5.88 mmol) in Et<sub>2</sub>O (15mL) cooled to 0 °C, was added water (1.4 mmol) via syringe. The slurry was stirred for 5 min, and **14a-i** (0.67 mmol) was added. The mixture was stirred at room temperature until starting material disappeared by TLC analysis (20% EtOAc in hexanes). Ice water was added until 2 clear layers formed. The aqueous layer was separated and acidified with

1 M HCl. The product was then extracted with Et<sub>2</sub>O (3 × 20 mL) and evaporated *in vacuo* to afford **15a-i**.

[00111] **4-(Nonylsulfonamido)benzoic acid 15a.** mp = 193-194 °C; <sup>1</sup>H NMR (MeOD) δ 7.99 (d, *J* = 8.0 Hz, 2H), 7.31 (d, *J* = 8.4 Hz, 2H), 3.17 (t, *J* = 8.0 Hz, 2H), 1.78 (m, 2H), 1.40 (m, 2H), 1.28 (m, 10H), 0.89 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 169.3, 144.3, 132.3, 126.7, 118.8, 52.4, 32.9, 30.2, 30.2, 30.0, 28.9, 24.4, 23.6, 14.3; HRMS (FAB) calcd for C<sub>16</sub>H<sub>25</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 327.15043; found, 327.14957.

[00112] **4-(Phenylsulfonamido)benzoic acid 15b.** mp = 186-188 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 12.72 (br s, 1H), 10.82 (br s, 1H), 7.80 (m, 4H), 7.61 (t, *J* = 6.8 Hz, 1H), 7.56 (t, *J* = 8.0 Hz, 2H), 7.20 (t, *J* = 7.2 Hz, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.6, 141.9, 142.0, 133.2, 130.7, 129.4, 126.6, 125.6, 118.2.; HRMS (FAB) calcd for C<sub>13</sub>H<sub>11</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 277.04088; found, 277.04077.

[00113] **4-(4-Chlorophenylsulfonamido)benzoic acid 15c.** mp = 254-256 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 12.76 (br s, 1H), 10.86 (br s, 1H), 7.81 (d, *J* = 6.4 Hz, 4H), 7.65 (d, *J* = 7.2 Hz, 2H), 7.18 (d, *J* = 6.8 Hz, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.6, 141.5, 138.1, 138.0, 130.7, 129.5, 128.5, 125.9, 118.4; HRMS (FAB) calcd for C<sub>13</sub>H<sub>11</sub>ClNO<sub>4</sub>S [M + H]<sup>+</sup>, 312.00973; found, 312.00859.

[00114] **3-(Nonylsulfonamido)benzoic acid 15d.** mp = 183-184 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.03 (br s, 1H), 9.98 (s, 1H), 7.81 (s, 1H), 7.64 (m, 1H), 7.44 (m, 2H), 3.07 (t, *J* = 7.6 Hz, 2H), 1.65 (m, 2H), 1.21 (m, 12H), 0.83 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.8, 138.7, 131.8, 129.5, 124.3, 123.2, 119.7, 50.5, 31.1, 28.5, 28.5, 28.3, 27.1, 22.9, 22.0, 13.8; HRMS (FAB) calcd for C<sub>16</sub>H<sub>26</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 328.15826; found, 328.15640.

[00115]        **3-(Phenylsulfonamido)benzoic acid 15e.** mp = 203-204 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.02 (br s, 1H), 10.51 (br s, 1H), 7.75 (d, *J* = 7.2 Hz, 2H), 7.68 (s, 1H), 7.56 (m, 4H), 7.34 (m, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.6, 139.2, 137.9, 133.0, 131.7, 129.4, 129.3, 126.5, 124.8, 124.0, 120.5; HRMS (FAB) calcd for C<sub>13</sub>H<sub>11</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 277.04088; found, 277.04054.

[00116]        **3-(4-Chlorophenylsulfonamido)benzoic acid 15f.** mp = 242-243 °C; <sup>1</sup>H NMR (MeOD) δ 7.76 (d, *J* = 8.8 Hz, 4H), 7.52 (d, *J* = 8.4 Hz, 2H), 7.34 (m, 2H); <sup>13</sup>C NMR (MeOD) δ 168.9, 140.2, 139.5, 139.0, 133.0, 130.3, 130.3, 129.8, 127.0, 126.4, 123.1; HRMS (FAB) calcd for C<sub>13</sub>H<sub>10</sub>ClNO<sub>4</sub>S [M]<sup>+</sup>, 311.00191; found, 311.00152.

[00117]        **C67 - 2-(Nonylsulfonamido)benzoic acid 15g.** mp = 122-124 °C; <sup>1</sup>H NMR (MeOD) δ 8.11 (d, *J* = 8.0 Hz, 1H), 7.73 (d, *J* = 8.4 Hz, 1H), 7.59 (t, *J* = 7.6 Hz, 1H), 7.16 (t, *J* = 7.6 Hz, 1H), 3.18 (t, *J* = 8.0 Hz, 2H), 1.71 (m, 2H), 1.24 (m, 12H), 0.88 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 171.3, 142.4, 135.7, 133.1, 123.8, 118.8, 117.0, 52.3, 32.9, 30.2, 30.1, 29.9, 28.8, 24.4, 23.6, 14.4; HRMS (FAB) calcd for C<sub>16</sub>H<sub>25</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 327.15043; found, 327.15044.

[00118]        **2-(Phenylsulfonamido)benzoic acid 15h.** mp = 213-215 °C; <sup>1</sup>H NMR (MeOD) δ 7.95 (d, *J* = 8.0 Hz, 1H), 7.81 (d, *J* = 8.0 Hz, 2H), 7.69 (d, *J* = 8.4 Hz, 1H), 7.56 (t, *J* = 7.6 Hz, 1H), 7.49 (m, 3H), 7.09 (t, *J* = 7.6 Hz, 1H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 169.7, 139.7, 138.5, 134.4, 133.5, 131.5, 129.4, 126.8, 123.3, 118.4, 116.7; HRMS (FAB) calcd for C<sub>13</sub>H<sub>11</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 277.04088; found, 277.04124.

[00119]        **2-(4-Chlorophenylsulfonamido)benzoic acid 15i.** mp = 202-203 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.98 (br s, 1H), 11.12 (br s, 1H), 7.90 (d, *J* = 8.0 Hz, 1H), 7.81 (d, *J* = 8.8 Hz, 2H), 7.63 (d, *J* = 8.8 Hz, 2H), 7.54 (t, *J* = 7.6 Hz, 1H), 7.48 (d, *J* = 8.4 Hz, 1H), 7.14 (t, *J* = 7.2

Hz, 1H);  $^{13}\text{C}$  NMR (MeOD)  $\delta$  171.1, 141.3, 140.6, 139.0, 135.4, 132.8, 130.3, 129.9, 124.7, 120.6, 118.3; HRMS (FAB) calcd for  $\text{C}_{13}\text{H}_{10}\text{ClINO}_4\text{S}$   $[\text{M}]^+$ , 311.00191; found, 311.00136.

**[00120]**      *Example 5 - Synthesis of Compounds 17a-17f*

**[00121]**      Synthesis of compounds 17a-17f was performed using Scheme 5, as illustrated in figure 5 herein.

**[00122]**      Reaction conditions: (a)  $\text{RSO}_2\text{Cl}$ , pyridine,  $\text{CH}_2\text{Cl}_2$ , 0 °C to rt; (b)  $\text{K}^+\text{O}^-t\text{-Bu}$ ,  $\text{Et}_2\text{O}$ ,  $\text{H}_2\text{O}$ , 0 °C to room temperature.

**[00123]**      Compounds **17a-f** were designed to probe the effect of linkers of different length in the aryl sulfonamide portion of the molecule. These were produced in the same manner as compounds **15a-i**, starting with the commercially available aniline and coupling to either benzylsulfonyl chloride or phenylethylsulfonyl chloride with pyridine in methylene chloride to yield sulfonamides **16a-f**. The methyl esters were then converted to the carboxylic acids **17a-f** with potassium *t*-butoxide and water in ether.

**[00124]**      **General Procedure for 16a-f.** To a stirring solution of the aniline starting material (3.3 mmol) in  $\text{CH}_2\text{Cl}_2$  (12 mL) at 0 °C was added pyridine (7.5 equiv). The sulfonyl chloride (1.2 equiv) was then added slowly via syringe. The solution was stirred and allowed to warm to room temperature. Reaction progress was monitored by TLC (20% EtOAc in hexanes). When complete, the reaction was poured into saturated  $\text{NaHCO}_3$  (45 mL), extracted with  $\text{CH}_2\text{Cl}_2$  ( $3 \times 15$  mL), and washed with 1 M HCl (50 mL). The combined organic phases were concentrated *in vacuo*, and the resulting solid was recrystallized from EtOAc / hexanes to afford **16a-f**.

[00125]        **General Procedure for 17a-f.** To a stirring suspension of potassium *t*-butoxide (5.88 mmol) in Et<sub>2</sub>O (15 mL) cooled to 0 °C, was added water (1.4 mmol) via syringe. The slurry was stirred for 5 min, and **16a-f** (0.67 mmol) was added. The mixture was stirred at room temperature until starting material disappeared by TLC analysis (20% EtOAc in hexanes). Ice water was added until 2 clear layers formed. The aqueous layer was separated and acidified with 1 M HCl. The product was then extracted with Et<sub>2</sub>O (3 × 20 mL) and evaporated *in vacuo* to afford **17a-f**.

[00126]        **4-(Phenylmethylsulfonamido)benzoic acid 17a.** mp = 221-223 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 12.72 (br s, 1H), 10.29 (s, 1H), 7.88 (d, *J* = 8.4 Hz, 2H), 7.33 (m, 3H), 7.24 (m, 4H), 4.56 (s, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.8, 142.7, 130.9, 130.8, 129.2, 128.3, 128.3, 124.8, 117.2, 57.1; HRMS (FAB) calcd for C<sub>14</sub>H<sub>14</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 292.06435; found, 292.06397.

[00127]        **4-(2-Phenylethylsulfonamido)benzoic acid 17b.** mp = 222-223 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 12.74 (br s, 1H), 10.38 (s, 1H), 7.90 (d, *J* = 8.0 Hz, 2H), 7.26 (m, 2H), 7.23 (m, 2H), 7.18 (m, 3H), 3.48 (t, *J* = 6.4, 2H), 2.98 (t, *J* = 6.4 Hz, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.8, 142.4, 137.8, 130.8, 128.4, 128.3, 126.5, 125.2, 117.8, 51.9, 29.0; HRMS (FAB) calcd for C<sub>15</sub>H<sub>16</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 306.08000; found, 306.07892.

[00128]        **3-(Phenylmethylsulfonamido)benzoic acid 17c.** mp = 205-206 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.02 (br s, 1H), 10.06 (s, 1H), 7.79 (s, 1H), 7.64 (d, *J* = 7.2 Hz, 1H), 7.42 (m, 2H), 7.33 (m, 3H), 7.25 (m, 2H), 4.48 (s, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.9, 138.7, 131.8, 130.9, 129.4, 129.3, 128.3, 128.2, 124.1, 122.9, 119.4, 57.0; HRMS (FAB) calcd for C<sub>14</sub>H<sub>14</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 292.06435; found, 292.06448.

[00129] **3-(2-Phenylethylsulfonamido)benzoic acid 17d.** mp = 199-200 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.06 (s, 1H), 10.11 (s, 1H), 7.85 (s, 1H), 7.67 (d, *J* = 6.8 Hz, 1H), 7.48 (m, 2H), 7.24 (m, 2H), 7.17 (m, 3H), 3.38 (t, *J* = 8.0 Hz, 2H), 2.99 (t, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 166.8, 138.5, 137.9, 131.9, 129.6, 128.4, 128.3, 126.5, 124.6, 123.8, 120.2, 51.7, 29.0; HRMS (FAB) calcd for C<sub>15</sub>H<sub>16</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 306.08000; found, 306.08051.

[00130] **2-(Phenylmethylsulfonamido)benzoic acid 17e.** mp = 216-219 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.86 (br s, 1H), 10.68 (s, 1H), 7.99 (d, *J* = 7.6 Hz, 1H), 7.58 (m, 2H), 7.32 (m, 3H), 7.19 (m, 3H), 4.69 (s, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 169.6, 140.7, 134.6, 131.5, 130.7, 128.8, 128.4, 128.3, 122.4, 117.2, 115.4, 57.2; HRMS (FAB) calcd for C<sub>14</sub>H<sub>13</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 291.05653; found, 291.05655.

[00131] **2-(2-Phenylethylsulfonamido)benzoic acid 17f.** mp = 157-159 °C; <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 13.90 (br s, 1H), 10.74 (br s, 1H), 7.98 (d, *J* = 8.0 Hz, 1H), 7.61 (d, *J* = 4.4 Hz, 2H), 7.20 (m, 2H), 7.16 (m, 4H), 3.61 (t, *J* = 8.0 Hz, 2H), 2.98 (t, *J* = 8.0 Hz, 2H); <sup>13</sup>C NMR (DMSO-d<sub>6</sub>) δ 169.7, 140.3, 137.5, 134.6, 131.6, 128.3, 128.2, 126.5, 122.6, 117.7, 115.9, 52.0, 28.9; HRMS (FAB) calcd for C<sub>15</sub>H<sub>16</sub>NO<sub>4</sub>S [M + H]<sup>+</sup>, 306.08000; found, 306.07886.

[00132] *Example 6 - Synthesis of Compounds 21a-21c*

[00133] Synthesis of compounds 21a-21c was performed using Scheme 6, as illustrated in figure 6 herein.

[00134] Reaction conditions: (a) diethyl phosphite, Et<sub>3</sub>N, Pd(PPh<sub>3</sub>)<sub>4</sub>, EtOH, reflux; (b) H<sub>2</sub>SO<sub>4</sub>, EtOH, reflux; (c) C<sub>8</sub>H<sub>17</sub>SO<sub>2</sub>Cl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C to room temperature; (d) TMSBr, CH<sub>2</sub>Cl<sub>2</sub>, room temperature.

[00135] The synthesis of aryl phosphonic acids **21a-c** is shown in **Scheme 6**. Aryl bromide **18** underwent palladium-catalyzed aryl halide coupling with diethyl phosphite to install the phosphonate functionality. (Gooßen, L. J., et. al.; Dezfuli, M. K. Practical Protocol for the Palladium-Catalyzed Synthesis of Arylphosphonates from Bromoarenes and Diethyl Phosphite, *Synlett* **2005**, 3, 445). The aniline was then deprotected by refluxing in acidic ethanol, and the free amine was coupled with commercially-available octanesulfonyl chloride to produce **20**. The final compound was then obtained by deprotecting the diethyl phosphonate with TMSBr.

[00136] **General Procedure for 19a-c.** The starting bromide **18** (1.96 mmol) was added to a round-bottomed flask containing diethyl phosphite (2.35 mmol), tetrakis(triphenylphosphine)palladium (0) (0.04 mmol), Et<sub>3</sub>N (2.94 mmol), and EtOH (8 mL), and the solution was heated to reflux overnight (16 h). The solution was then diluted with 30 mL EtOAc, washed with 50 mL saturated NaHCO<sub>3</sub> solution, 50 mL H<sub>2</sub>O, dried over MgSO<sub>4</sub>, and concentrated *in vacuo*. The product was then purified by flash chromatography (EtOAc).

[00137] **4-(Octylsulfonamido)phenylphosphonic acid 21a.** mp = 185-187 °C; <sup>1</sup>H NMR (MeOD) δ 7.75 (dd, *J* = 12.8, 8.0 Hz, 2H), 7.33 (dd, *J* = 8.0, 3.2 Hz, 2H), 3.15 (t, *J* = 8.0 Hz, 2H), 1.78 (m, 2H), 1.39 (m, 2H), 1.28 (m, 8H), 0.90 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 143.0 (d, *J* = 3.6 Hz), 133.4 (d, *J* = 11.0 Hz), 127.7 (d, *J* = 190 Hz), 119.1 (d, *J* = 15.2 Hz), 52.4, 32.8, 30.0, 29.9, 29.0, 24.5, 23.6, 14.3; HRMS (FAB) calcd for C<sub>14</sub>H<sub>25</sub>NO<sub>5</sub>PS [M + H]<sup>+</sup>, 350.11911; found, 350.11869.

[00138] **3-(Octylsulfonamido)phenylphosphonic acid 21b.** mp = 112-114 °C; <sup>1</sup>H NMR (MeOD) δ 7.72 (d, *J* = 14.8 Hz, 1H), 7.55 (m, 1H), 7.44 (m, 2H), 3.12 (t, *J* = 8.0 Hz, 2H), 1.78 (m, 2H), 1.39 (m, 2H), 1.27 (m, 8H), 0.90 (t, *J* = 7.2 Hz, 3H); <sup>13</sup>C NMR (MeOD) δ 139.6 (d, *J* =

18.2 Hz), 134.5 (d,  $J = 184$  Hz), 130.5 (d,  $J = 16.1$  Hz), 127.3 (d,  $J = 9.5$  Hz), 123.8 (d,  $J = 3.0$  Hz), 122.8 (d,  $J = 11.6$  Hz), 52.2, 32.7, 30.0, 29.9, 29.0, 24.4, 23.5, 14.3; HRMS (FAB) calcd for  $C_{14}H_{25}NO_5PS$   $[M + H]^+$ , 350.11911; found, 350.11879.

[00139]        **2-(Octylsulfonamido)phenylphosphonic acid 21c.** mp = 92-94 °C;  $^1H$  NMR (MeOD)  $\delta$  7.70 (m, 2H), 7.54 (t,  $J = 8.4$  Hz, 1H), 7.21 (t,  $J = 7.5$  Hz, 1H), 3.18 (t,  $J = 7.8$  Hz, 2H), 1.77 (m, 2H), 1.23 (m, 10H), 0.89 (t,  $J = 7.5$  Hz, 3H);  $^{13}C$  NMR (MeOD)  $\delta$  141.8 (d,  $J = 7.0$  Hz), 134.3 (d,  $J = 2.7$  Hz), 134.1 (d,  $J = 6.8$  Hz), 120.4 (d,  $J = 178$  Hz), 119.6 (d,  $J = 10.8$  Hz), 52.6, 32.8, 29.9, 29.9, 29.0, 24.3, 23.5, 14.3; HRMS (FAB) calcd for  $C_{14}H_{25}NO_5PS$   $[M + H]^+$ , 350.11911; found, 350.11826.

[00140]        *Example 7 - Synthesis of Compounds 24a-24f*

[00141]        Synthesis of compounds 24a-24f was performed using Scheme 7, as illustrated in figure 7 herein.

[00142]        Reaction conditions: (a)  $RSO_2Cl$ , pyridine,  $CH_2Cl_2$ , 0 °C to room temperature; (b)  $K^+O^-t-Bu$ ,  $Et_2O$ ,  $H_2O$ , 0 °C to room temperature.

[00143]        Compounds **24a-c**, based on **15g**, were designed as probes to examine the effect of installing different length alkylsulfonamides on the *ortho*-substituted analogs. It was believed that the compound with the saturated  $C_{16}$ -chain (**24c**) would exhibit significantly greater inhibitory activity than **15g**, as the enzyme demonstrates a marked preference for palmitoyl-CoA over other long-chain acyl-CoAs.<sup>13</sup> Compounds **24d-f** were designed to examine the role of an electronegative group at the 4-position of the benzene ring, which could possibly mimic the electron density of the secondary alcohol on glycerol-3-phosphate. All of these compounds (**24a-f**) were produced with the same reaction sequence used to produce **15a-f** and **17a-f**.

[00144]        **General Procedure for 23a-f.** To a stirring solution of the aniline starting material (3.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (12 mL) at 0 °C was added pyridine (7.5 equiv). The sulfonyl chloride (1.2 equiv) was then added slowly via syringe. The solution was stirred and allowed to warm to room temperature. Reaction progress was monitored by TLC (20% EtOAc in hexanes). When complete, the reaction was poured into saturated NaHCO<sub>3</sub> solution (45 mL), extracted with CH<sub>2</sub>Cl<sub>2</sub> (3 × 15 mL), and washed with 1 M HCl (50 mL). The combined organic phases were concentrated *in vacuo*, and separated by flash chromatography (20 % EtOAc in hexanes) to afford **23a-f**.

[00145]        **General Procedure for 24a-f.** To a stirring suspension of potassium *t*-butoxide (5.88 mmol) in Et<sub>2</sub>O (15 mL) cooled to 0 °C was added water (1.4 mmol) via syringe. The slurry was stirred for 5 min, and **23a-f** (0.67 mmol) was added. The mixture was stirred at room temperature until starting material disappeared by TLC analysis (20% EtOAc in hexanes). Ice water was added until two clear layers formed. The aqueous layer was separated and acidified with 1 M HCl, and the product was extracted with Et<sub>2</sub>O (3 × 20 mL) and evaporated *in vacuo*. If necessary, the product was then recrystallized (EtOAc / hexanes) to afford pure **24a-f**.

[00146]        **2-(Methylsulfonylamido)benzoic acid 24a.** mp = 187-189 °C; <sup>1</sup>H NMR (MeOD) δ 8.11 (d, *J* = 8.0 Hz, 1H), 7.70 (d, *J* = 8.0 Hz, 1H), 7.60 (t, *J* = 7.2 Hz, 1H), 7.17 (t, *J* = 7.6 Hz, 1H), 3.08 (s, 3H); <sup>13</sup>C NMR (MeOD) δ 171.2, 142.2, 135.7, 133.0, 123.9, 119.2, 117.3, 39.9; HRMS (FAB) calcd for C<sub>8</sub>H<sub>9</sub>NO<sub>4</sub>S [M]<sup>+</sup>, 215.02523; found, 215.02576.

[00147]        **2-(Tetradecylsulfonylamido)benzoic acid 24b.** mp = 120-122 °C; <sup>1</sup>H NMR (MeOD) δ 8.12 (d, *J* = 8.0 Hz, 1H), 7.74 (d, *J* = 8.4 Hz, 1H), 7.59 (t, *J* = 7.6 Hz, 1H), 7.17 (t, *J* = 7.6 Hz, 1H), 3.19 (t, *J* = 8.0 Hz, 2H), 1.70 (m, 2H), 1.29 (m, 22H), 0.91 (t, *J* = 6.8 Hz, 3H); <sup>13</sup>C

NMR (MeOD)  $\delta$  171.4, 142.5, 135.7, 133.1, 123.8, 118.8, 117.0, 52.2, 33.0, 30.7, 30.7, 30.7, 30.6, 30.5, 30.4, 30.2, 29.9, 28.8, 24.4, 23.7, 14.4.

[00148]      **2-(Hexadecylsulfonamido)benzoic acid 24c.** mp = 126-128 °C;  $^1\text{H}$  NMR (MeOD)  $\delta$  8.12 (d,  $J$  = 7.6 Hz, 1H), 7.74 (d,  $J$  = 8.4 Hz, 1H), 7.59 (t,  $J$  = 8.0 Hz, 1H), 7.17 (t,  $J$  = 7.6 Hz, 1H), 3.19 (t,  $J$  = 7.6 Hz, 2H), 1.73 (m, 2H), 1.23 (m, 26H), 0.91 (t,  $J$  = 7.2 Hz, 3H);  $^{13}\text{C}$  NMR (MeOD)  $\delta$  169.8, 140.7, 134.6, 131.6, 122.4, 117.3, 115.6, 50.9, 31.2, 29.0, 29.0, 29.0, 29.0, 29.0, 28.9, 28.8, 28.7, 28.5, 28.2, 27.0, 22.8, 22.0, 13.8; HRMS (FAB) calcd for  $\text{C}_{23}\text{H}_{40}\text{NO}_4\text{S}$   $[\text{M} + \text{H}]^+$ , 426.26781; found, 426.26825.

[00149]      **5-Chloro-2-(nonylsulfonamido)benzoic acid 24d.** mp = 101-103 °C;  $^1\text{H}$  NMR (MeOD)  $\delta$  8.05 (d,  $J$  = 2.8 Hz, 1H), 7.74 (d,  $J$  = 9.2 Hz, 1H), 7.59 (dd,  $J$  = 9.2, 2.8 Hz, 1H), 3.21 (t,  $J$  = 8.0 Hz, 2H), 1.72 (m, 2H), 1.23 (m, 12H), 0.90 (t,  $J$  = 7.2 Hz, 3H);  $^{13}\text{C}$  NMR (MeOD)  $\delta$  170.1, 141.2, 135.5, 132.4, 128.9, 120.6, 118.0, 52.5, 32.9, 30.2, 30.1, 29.9, 28.8, 24.4, 23.6, 14.4; HRMS (FAB) calcd for  $\text{C}_{16}\text{H}_{24}\text{ClNO}_4\text{S}$   $[\text{M}]^+$ , 361.11146; found, 361.11063.

[00150]      **5-Hydroxy-2-(octylsulfonamido)benzoic acid 24e.** mp = 142-144 °C;  $^1\text{H}$  NMR (MeOD)  $\delta$  7.56 (d,  $J$  = 8.8 Hz, 1H), 7.51 (d,  $J$  = 2.8 Hz, 1H), 7.03 (dd,  $J$  = 8.8, 2.8 Hz, 1H), 3.07 (t,  $J$  = 8.0 Hz, 2H), 1.68 (m, 2H), 1.23 (m, 10H), 0.89 (t,  $J$  = 7.2 Hz, 3H);  $^{13}\text{C}$  NMR (MeOD)  $\delta$  171.0, 154.6, 134.1, 122.8, 121.9, 119.2, 118.5, 53.1, 32.8, 29.9, 29.8, 28.8, 24.3, 23.6, 14.4.

[00151]      **5-Fluoro-2-(octylsulfonamido)benzoic acid 24f.** mp = 141-143 °C;  $^1\text{H}$  NMR (MeOD)  $\delta$  7.77 (m, 2H), 7.38 (m, 1H), 3.17 (t,  $J$  = 8.0 Hz, 2H), 1.71 (m, 2H), 1.23 (m, 10H), 0.88 (t,  $J$  = 7.2 Hz, 3H);  $^{13}\text{C}$  NMR (MeOD)  $\delta$  170.2 (d,  $J$  = 1.8 Hz), 159.2 (d,  $J$  = 241 Hz), 138.6 (d,  $J$  = 2.7 Hz), 122.7 (d,  $J$  = 22.7 Hz), 121.5 (d,  $J$  = 7.6 Hz), 119.0 (d,  $J$  = 6.9 Hz), 118.8 (d,  $J$  =

24.1 Hz), 52.4, 32.8, 29.9, 29.9, 28.8, 24.4, 23.6, 14.3; HRMS (FAB) calcd for C<sub>15</sub>H<sub>22</sub>FNO<sub>4</sub>S [M]<sup>+</sup>, 331.12536; found, 331.12445.

**[00152]**      *Example 8 – Synthesis of Compounds 4a-t and 7a-t*

**[00153]**      Synthesis of compounds 4a-t and 7a-t was performed using Schemes illustrated in figures 8 and 9, respectively, herein.

**[00154]**      General Suzuki Reaction Experimental - 0.247 mmol aryl bromide was placed into a vial flushed with argon, and a solution of 10 mg Pd(PPh<sub>3</sub>)<sub>4</sub> in 0.40 mL toluene was added, followed by 0.25 mL 2M Na<sub>2</sub>CO<sub>3</sub> solution. The solution was stirred at room temperature for 5 min, and then a solution of the boronic acid (1.25 equiv) in 0.40 mL MeOH was added. The vial was capped and heated to 90 °C for 24 h. The reaction was then cooled to room temperature and diluted with CH<sub>2</sub>Cl<sub>2</sub>, the organic phase was separated from the aqueous phase, and the organic phase was concentrated *in vacuo*. The crude product was purified by column chromatography (EtOAc/hexanes) to yield the desired bis-aryl product.

**[00155]**      **General Procedure for 4a-t and 7a-t.** To a stirring suspension of potassium *t*-butoxide (2.00 mmol) in Et<sub>2</sub>O (8 mL) cooled to 0 °C, was added water (0.4 mmol) via syringe. The slurry was stirred for 5 min, and **3a-t** or **6a-t** (0.2 mmol) was added. The mixture was stirred at room temperature until starting material disappeared by TLC analysis (20% EtOAc in hexanes). Ice water was added until 2 clear layers formed. The aqueous layer was separated and acidified with 1 M HCl. The product was then extracted with Et<sub>2</sub>O (3 × 20 mL) and evaporated *in vacuo* to afford **4a-t** and **7a-t**. If further purification was necessary, the product was purified by flash chromatography (1: 1: 8 AcOH: EtOAc: hexanes).

[00156] (4a)  $^1\text{H}$  NMR (DMSO- $\text{d}_6$ )  $\delta$  8.00 (d,  $J = 8.0$  Hz, 1H), 7.57 (m, 1H), 7.48 (s, 1H), 7.41 (m, 3H), 6.96 (d,  $J = 8.0$  Hz, 1H), 3.08 (t,  $J = 8.0$  Hz, 2H), 1.60 (m, 2H), 1.18 (m, 8H), 0.80 (t,  $J = 7.2$  Hz, 3H).

[00157] (4b)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.21 (d,  $J = 8.4$  Hz, 1H), 7.97 (s, 1H), 7.68 (s, 1H), 7.62 (d,  $J = 8.4$  Hz, 1H), 7.49 (m, 2H), 3.25 (t,  $J = 8.4$  Hz, 2H), 1.76 (m, 2H), 1.38 (m, 2H), 1.23 (m, 8H), 0.87 (t,  $J = 7.2$  Hz, 3H).

[00158] (4c)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.15 (d,  $J = 8.0$  Hz, 1H), 7.89 (d,  $J = 1.6$  Hz, 1H), 7.65 (d,  $J = 6.8$  Hz, 2H), 7.49 (d,  $J = 6.8$  Hz, 2H), 7.35 (dd,  $J = 8.0, 2.0$  Hz, 1H), 3.14 (t,  $J = 8.0$  Hz, 2H), 1.74 (m, 2H), 1.35 (m, 2H), 1.22 (m, 8H), 0.86 (t,  $J = 6.8$  Hz, 3H).

[00159] (4d)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.26 (m, 1H), 8.23 (d,  $J = 8.4$  Hz, 1H), 8.07 (m, 1H), 8.01 (d,  $J = 1.5$  Hz, 1H), 7.92 (m, 1H), 7.66 (t,  $J = 7.8$  Hz, 1H), 7.49 (dd,  $J = 8.4, 1.8$  Hz, 1H), 3.23 (t,  $J = 7.8$  Hz, 2H), 2.69 (s, 3H), 1.77 (m, 2H), 1.40 (m, 2H), 1.22 (m, 8H), 0.89 (t,  $J = 7.2$  Hz, 3H).

[00160] (4e)  $^1\text{H}$  NMR (DMSO- $\text{d}_6$ )  $\delta$  8.08 (m, 3H), 7.82 (m, 3H), 7.46 (dd,  $J = 7.6, 1.6$  Hz, 1H), 3.27 (t,  $J = 7.6$  Hz, 2H), 2.60 (s, 3H), 1.60 (m, 2H), 1.30 (m, 2H), 1.16 (m, 8H), 0.79 (t,  $J = 6.8$  Hz, 3H).

[00161] (4f)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.23 (d,  $J = 8.4$  Hz, 1H), 8.03 (d,  $J = 1.6$  Hz, 1H), 7.87 (m, 4H), 7.50 (dd,  $J = 8.4, 1.6$  Hz, 1H), 3.24 (t,  $J = 8.0$  Hz, 2H), 1.76 (m, 2H), 1.38 (m, 2H), 1.23 (m, 8H), 0.87 (t,  $J = 7.2$  Hz, 3H).

[00162] (4g)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.06 (d,  $J = 8.4$  Hz, 1H), 7.45 (m, 5H), 7.23 (m, 5H), 7.12 (dd,  $J = 8.0, 1.2$  Hz, 1H), 2.61 (t,  $J = 8.0$  Hz, 2H), 1.56 (m, 2H), 1.26 (m, 10H), 0.91 (t,  $J = 7.2$  Hz, 3H).

[00163] (4h)  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  8.08 (d,  $J = 8.4$  Hz, 1H), 7.81 (m, 7H), 7.49 (m, 3H), 7.39 (t,  $J = 7.2$  Hz, 1H), 3.31 (t,  $J = 8.0$  Hz, 2H), 1.64 (m, 2H), 1.31 (m, 2H), 1.18 (m, 8H), 0.79 (t,  $J = 6.8$  Hz, 3H).

[00164] (4i)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.12 (d,  $J = 8.4$  Hz, 1H), 7.93 (s, 1H), 7.40 (m, 2H), 7.28 (d,  $J = 8.4$  Hz, 1H), 7.11 (m, 2H), 3.85 (s, 3H), 3.25 (t,  $J = 7.8$  Hz, 2H), 1.71 (m, 2H), 1.22 (m, 10H), 0.87 (t,  $J = 6.9$  Hz, 3H).

[00165] (4j)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.12 (d,  $J = 8.4$  Hz, 1H), 7.94 (d,  $J = 1.6$  Hz, 1H), 7.60 (d,  $J = 6.8$  Hz, 2H), 7.37 (dd,  $J = 8.4, 1.6$  Hz, 1H), 7.02 (d,  $J = 6.8$  Hz, 2H), 3.84 (s, 3H), 3.20 (t,  $J = 7.6$  Hz, 2H), 1.73 (m, 2H), 1.36 (m, 2H), 1.24 (m, 8H), 0.86 (t,  $J = 7.2$  Hz, 3H).

[00166] (4k)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.17 (m, 1H), 7.93 (m, 1H), 7.52 (m, 1H), 7.44 (m, 1H), 7.32 (m, 2H), 7.22 (m, 1H), 3.22 (t,  $J = 8.0$  Hz, 2H), 1.75 (m, 2H), 1.34 (2H), 1.22 (m, 8H), 0.83 (t,  $J = 6.8$  Hz, 3H).

[00167] (4l)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.19 (d,  $J = 8.4$  Hz, 1H), 7.97 (s, 1H), 7.50 (m, 2H), 7.43 (t,  $J = 8.0$  Hz, 2H), 7.16 (m, 1H), 3.22 (t,  $J = 7.6$  Hz, 2H), 1.76 (m, 2H), 1.37 (m, 2H), 1.23 (m, 8H), 0.87 (t,  $J = 7.2$  Hz, 3H).

[00168] (4m)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.17 (d,  $J = 8.0$  Hz, 1H), 7.94 (d,  $J = 1.6$  Hz, 1H), 7.71 (m, 2H), 7.40 (dd,  $J = 8.4, 2.0$  Hz, 1H), 7.22 (t,  $J = 8.8$  Hz, 2H), 3.21 (t,  $J = 7.6$  Hz, 2H), 1.75 (m, 2H), 1.37 (m, 2H), 1.24 (m, 8H), 0.87 (t,  $J = 7.2$  Hz, 3H).

[00169] (4n)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.12 (m, 1H), 8.01 (m, 1H), 7.34 (m, 2H), 7.22 (m, 1H), 6.93 (m, 2H), 3.26 (t,  $J = 8.0$  Hz, 2H), 1.73 (m, 2H), 1.35 (m, 2H), 1.18 (m, 8H), 0.85 (t,  $J = 7.2$  Hz, 3H).

[00170] (4o)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.15 (d,  $J = 8.4$  Hz, 1H), 7.95 (s, 1H), 7.38 (d,  $J = 8.4$  Hz, 1H), 7.29 (t,  $J = 7.6$  Hz, 1H), 7.11 (m, 2H), 6.85 (m, 1H), 3.20 (t,  $J = 8.0$  Hz, 2H), 1.74 (m, 2H), 1.35 (m, 2H), 1.24 (m, 8H), 0.85 (t,  $J = 7.2$  Hz, 3H).

[00171] (4p) 8.10 (d,  $J = 8.0$  Hz, 1H), 7.92 (s, 1H), 7.53 (m, 2H), 7.36 (m, 1H), 6.89 (m, 2H), 3.19 (t,  $J = 7.6$  Hz, 2H), 1.72 (m, 2H), 1.34 (m, 2H), 1.19 (m, 8H), 0.84 (t,  $J = 6.8$  Hz, 3H).

[00172] (4q)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.10 (t,  $J = 8.4$  Hz, 1H), 8.01 (d,  $J = 8.4$  Hz, 1H), 7.53 (m, 2H), 7.43 (m, 1H), 7.18 (m, 1H), 3.22 (t,  $J = 8.0$  Hz, 2H), 1.74 (m, 2H), 1.36 (m, 2H), 1.19 (m, 8H), 0.82 (t,  $J = 7.2$  Hz, 3H).

[00173] (4r)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.84 (s, 1H), 8.57 (s, 1H), 8.21 (d,  $J = 8.4$  Hz, 1H), 8.13 (d,  $J = 7.6$  Hz, 1H), 7.92 (d,  $J = 2.0$  Hz, 1H), 7.56 (m, 1H), 7.40 (dd,  $J = 8.0, 2.0$  Hz, 1H), 3.15 (t,  $J = 8.0$  Hz, 2H), 1.77 (m, 2H), 1.36 (m, 2H), 1.22 (m, 8H), 0.86 (t,  $J = 6.8$  Hz, 3H).

[00174] (4s)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.15 (d,  $J = 8.4$  Hz, 1H), 7.88 (s, 1H), 7.52 (s, 2H), 7.42 (s, 1H), 7.33 (d,  $J = 8.0$  Hz, 1H), 3.22 (t,  $J = 7.6$  Hz, 2H), 1.72 (m, 2H), 1.35 (m, 2H), 1.23 (m, 8H), 0.84 (t,  $J = 7.2$  Hz, 3H).

[00175] (4t)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.18 (d,  $J = 8.0$  Hz, 1H), 7.79 (s, 1H), 7.60 (d,  $J = 2.0$  Hz, 1H), 7.43 (m, 2H), 7.19 (dd,  $J = 8.0, 1.6$  Hz, 1H), 3.23 (t,  $J = 7.6$  Hz, 2H), 1.74 (m, 2H), 1.36 (m, 2H), 1.21 (m, 8H), 0.87 (t,  $J = 7.2$  Hz, 3H).

[00176] (7a)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.17 (d,  $J = 2.4$  Hz, 1H), 7.76 (d,  $J = 8.4$  Hz, 1H), 7.60 (dd,  $J = 8.4, 2.4$  Hz, 1H), 7.49 (d,  $J = 8.0$  Hz, 1H), 7.37 (m, 3H), 3.21 (t,  $J = 8.0$  Hz, 2H), 1.76 (m, 2H), 1.20 (m, 10H), 0.86 (t,  $J = 7.2$  Hz, 3H).

[00177] (7b)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.29 (s, 1H), 7.81 (m, 2H), 7.57 (m, 1H), 7.52 (d,  $J = 7.6$  Hz, 1H), 7.42 (t,  $J = 8.0$  Hz, 1H), 7.34 (d,  $J = 8.0$  Hz, 1H), 3.22 (t,  $J = 8.0$  Hz, 2H), 1.72 (m, 2H), 1.23 (m, 10H), 0.83 (t,  $J = 7.2$  Hz, 3H).

[00178] (7c)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.32 (d,  $J = 2.0$  Hz, 1H), 7.81 (m, 2H), 7.58 (d,  $J = 8.8$  Hz, 2H), 7.43 (d,  $J = 8.4$  Hz, 2H), 3.22 (t,  $J = 8.0$  Hz, 2H), 1.73 (m, 2H), 1.35 (m, 2H), 1.20 (m, 8H), 0.84 (t,  $J = 7.2$  Hz, 3H).

[00179] (7d)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.37 (s, 1H), 8.19 (s, 1H), 7.97 (d,  $J = 7.8$  Hz, 1H), 7.81 (m, 3H), 7.57 (t,  $J = 7.5$  Hz, 1H), 3.21 (t,  $J = 7.8$  Hz, 2H), 2.66 (s, 3H), 1.74 (m, 2H), 1.20 (m, 10H), 0.83 (t,  $J = 7.2$  Hz, 3H).

[00180] (7e)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.42 (d,  $J = 2.4$  Hz, 1H), 8.12 (d,  $J = 8.4$  Hz, 2H), 7.99 (d,  $J = 8.4$  Hz, 1H), 7.89 (d,  $J = 8.7$  Hz, 1H), 7.81 (d,  $J = 8.4$  Hz, 2H), 3.29 (t,  $J = 8.0$  Hz, 2H), 2.66 (s, 3H), 1.77 (m, 2H), 1.37 (m, 2H), 1.24 (m, 8H), 0.89 (t,  $J = 7.2$  Hz, 3H).

[00181] (7f)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.40 (s, 1H), 7.89 (m, 1H), 7.80 (m, 5H), 3.23 (t,  $J = 8.0$  Hz, 2H), 1.73 (m, 2H), 1.36 (m, 2H), 1.21 (m, 8H), 0.84 (t,  $J = 6.8$  Hz, 3H).

[00182] (7g)  $^1\text{H}$  NMR (MeOD)  $\delta$  7.89 (d,  $J = 2.0$  Hz, 1H), 7.51 (d,  $J = 8.8$  Hz, 1H), 7.38 (m, 4H), 7.19 (m, 4H), 7.09 (m, 2H), 3.10 (t,  $J = 8.0$  Hz, 2H), 1.65 (m, 2H), 1.21 (m, 10H), 0.86 (t,  $J = 6.8$  Hz, 3H).

[00183] (7h)  $^1\text{H}$  NMR ( $\text{DMSO-d}_6$ )  $\delta$  8.31 (d,  $J = 2.4$  Hz, 1H), 8.01 (dd,  $J = 8.8, 2.4$  Hz, 1H), 7.74 (m, 7H), 7.48 (t,  $J = 8.0$  Hz, 2H), 7.38 (t,  $J = 8.0$  Hz, 1H), 3.33 (t,  $J = 7.6$  Hz, 2H), 1.63 (m, 2H), 1.31 (m, 2H), 1.20 (m, 8H), 0.81 (t,  $J = 6.8$  Hz, 3H).

[00184] (7i)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.24 (s, 1H), 7.71 (m, 2H), 7.28 (m, 2H), 6.98 (m, 2H), 3.77 (s, 3H), 3.17 (t,  $J = 8.0$  Hz, 2H), 1.70 (m, 2H), 1.19 (m, 10H), 0.83 (t,  $J = 7.2$  Hz, 3H).

[00185] (7j)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.26 (s, 1H), 7.71 (m, 2H), 7.46 (d,  $J = 8.8$  Hz, 2H), 6.93 (d,  $J = 8.8$  Hz, 2H), 3.79 (s, 3H), 3.15 (t,  $J = 8.0$  Hz, 2H), 1.70 (m, 2H), 1.20 (m, 10H), 0.81 (t,  $J = 7.2$  Hz, 3H).

[00186] (7k)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.27 (s, 1H), 7.78 (d,  $J = 8.4$  Hz, 1H), 7.70 (d,  $J = 8.4$  Hz, 1H), 7.42 (t,  $J = 8.0$  Hz, 1H), 7.32 (m, 1H), 7.19 (m, 2H), 3.19 (t,  $J = 7.6$  Hz, 2H), 1.71 (m, 2H), 1.32 (m, 2H), 1.16 (m, 8H), 0.81 (t,  $J = 7.2$  Hz, 3H).

[00187] (7l)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.31 (t,  $J = 1.2$  Hz, 1H), 7.79 (s, 2H), 7.41 (m, 2H), 7.30 (dd,  $J = 8.8, 1.6$  Hz, 1H), 7.05 (dt,  $J = 8.8, 1.6$  Hz, 1H), 3.20 (t,  $J = 8.0$  Hz, 2H), 1.71 (m, 2H), 1.22 (m, 2H), 1.16 (m, 8H), 0.81 (t,  $J = 6.8$  Hz, 3H).

[00188] (7m)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.28 (d,  $J = 2.0$  Hz, 1H), 7.74 (m, 2H), 7.57 (dd,  $J = 8.8, 4.8$  Hz, 2H), 7.12 (t,  $J = 8.4$  Hz, 2H), 3.17 (t,  $J = 8.0$  Hz, 2H), 1.70 (m, 2H), 1.31 (m, 2H), 1.18 (m, 8H), 0.81 (t,  $J = 6.8$  Hz, 3H).

[00189] (7n)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.33 (s, 1H), 7.75 (dd,  $J = 8.4, 2.4$  Hz, 1H), 7.69 (d,  $J = 8.4$  Hz, 1H), 7.28 (dd,  $J = 8.0, 1.6$  Hz, 1H), 7.15 (dt,  $J = 7.6, 1.6$  Hz, 1H), 6.90 (m, 2H), 3.17 (t,  $J = 8.0$  Hz, 2H), 1.75 (m, 2H), 1.23 (m, 10H), 0.86 (t,  $J = 7.2$  Hz, 3H).

[00190] (7o)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.31 (s, 1H), 7.77 (s, 2H), 7.24 (t,  $J = 8.0$  Hz, 1H), 7.04 (m, 2H), 6.79 (d,  $J = 8.0$  Hz, 1H), 3.19 (t,  $J = 8.0$  Hz, 2H), 1.72 (m, 2H), 1.33 (m, 2H), 1.21 (m, 8H), 0.82 (t,  $J = 6.8$  Hz, 3H).

[00191] (7p)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.32 (s, 1H), 7.64 (m, 2H), 7.46 (m, 2H), 6.86 (m, 2H), 3.13 (t,  $J = 8.0$  Hz, 2H), 1.73 (m, 2H), 1.34 (m, 2H), 1.20 (m, 8H), 0.83 (t,  $J = 6.8$  Hz, 3H).

[00192] (7q)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.36 (d,  $J = 2.4$  Hz, 1H), 7.80 (dd,  $J = 8.4, 2.0$  Hz, 1H), 7.73 (d,  $J = 8.4$  Hz, 1H), 7.39 (m, 2H), 7.10 (dd,  $J = 5.2, 3.6$  Hz, 1H), 3.19 (t,  $J = 8.0$  Hz, 2H), 1.74 (m, 2H), 1.38 (m, 2H), 1.22 (m, 8H), 0.85 (t,  $J = 7.2$  Hz, 3H).

[00193] (7r) 8.86 (s, 1H), 8.55 (d,  $J = 4.4$  Hz, 1H), 8.42 (s, 1H), 8.17 (d,  $J = 8.0$  Hz, 1H), 7.90 (m, 2H), 7.57 (m, 1H), 3.25 (t,  $J = 8.0$  Hz, 2H), 1.77 (m, 2H), 1.37 (m, 2H), 1.24 (m, 8H), 0.86 (t,  $J = 7.2$  Hz, 3H).

[00194] (7s)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.21 (s, 1H), 7.75 (m, 2H), 7.45 (d,  $J = 1.6$  Hz, 2H), 7.32 (t,  $J = 1.6$  Hz, 1H), 3.22 (t,  $J = 8.0$  Hz, 2H), 1.74 (m, 2H), 1.33 (m, 2H), 1.19 (m, 8H), 0.83 (t,  $J = 7.2$  Hz, 3H).

[00195] (7t)  $^1\text{H}$  NMR (MeOD)  $\delta$  8.16 (d,  $J = 2.4$  Hz, 1H), 7.74 (d,  $J = 8.8$  Hz, 1H), 7.54 (dd,  $J = 8.4, 2.4$  Hz, 1H), 7.50 (s, 1H), 7.34 (s, 2H), 3.20 (t,  $J = 8.0$  Hz, 2H), 1.75 (m, 2H), 1.34 (m, 2H), 1.20 (m, 8H), 0.84 (t,  $J = 6.8$  Hz, 3H).

[00196] **Example 8 - In Vitro Testing**


[00197] The compounds produced as described above were evaluated for their ability to inhibit the acylation of glycerol-3-phosphate *in vitro*. The acylation reaction between  $^{14}\text{C}$ -labelled glycerol-3-phosphate and palmitoyl-CoA, initiated by adding mtGPAT, was measured in

the presence of varying concentrations of the inhibitor by scintillation counting as described in more detail below.

**[00198]** A mitochondrial preparation of glycerol 3-phosphate acyltransferase was added to the incubation mixture containing  $^{14}\text{C}$ -labeled glycerol 3-phosphate, palmitoyl-CoA, and varying inhibitor concentrations to initiate the reaction. After ten min, the reaction was terminated by adding chloroform, methanol, and 1% perchloric acid. Five minutes later, more chloroform and perchloric acid were added, and the upper aqueous layer was removed. After washing three times with 1% perchloric acid, the organic layer was evaporated under nitrogen, and the amount of  $^{14}\text{C}$  present was counted to determine the extent of reaction inhibition. Data points were recorded in triplicate, and  $\text{IC}_{50}$  values were calculated based on the amount of test inhibitor necessary to produce 50% of mtGPAT activity observed in the absence of inhibitor but in the presence of DMSO vehicle control.

**[00199]** Results for compounds 5a-f, 13a-f, 15a-i, 17a-f, 21a-c, and 24a-f are summarized in Tables 1-3 below. The results for each of the compounds 4a-t and 7a-t are summarized individually below.

**Table 1.** *In Vitro* Anti-mtGPAT1 Activity of Sulfonamides **5a-f** and **13a-f**




| Compound  | X                           | Y                              | n | $\text{IC}_{50}$ ( $\mu\text{M}$ ) $\pm$ SD |
|-----------|-----------------------------|--------------------------------|---|---------------------------------------------|
| <b>5a</b> | <i>p</i> -CO <sub>2</sub> H | C <sub>5</sub> H <sub>11</sub> | 1 | 72.0 $\pm$ 1.7                              |
| <b>5b</b> | <i>p</i> -CO <sub>2</sub> H | C <sub>9</sub> H <sub>19</sub> | 1 | 43.9 $\pm$ 6.3                              |

|            |                                                          |                                |   |             |
|------------|----------------------------------------------------------|--------------------------------|---|-------------|
| <b>5c</b>  | <i>m</i> -CO <sub>2</sub> H                              | C <sub>5</sub> H <sub>11</sub> | 1 | 88.5 ± 1.7  |
| <b>5d</b>  | <i>m</i> -CO <sub>2</sub> H                              | C <sub>9</sub> H <sub>19</sub> | 1 | 28.5 ± 1.6  |
| <b>5e</b>  | <i>o</i> -CO <sub>2</sub> H                              | C <sub>5</sub> H <sub>11</sub> | 1 | 61.9 ± 13.5 |
| <b>5f</b>  | <i>o</i> -CO <sub>2</sub> H                              | C <sub>9</sub> H <sub>19</sub> | 1 | 22.7 ± 1.1  |
| <b>13a</b> | <i>o</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>5</sub> H <sub>11</sub> | 0 | 41.4 ± 8.4  |
| <b>13b</b> | <i>o</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>9</sub> H <sub>19</sub> | 0 | 30.6 ± 6.2  |
| <b>13c</b> | <i>m</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>5</sub> H <sub>11</sub> | 0 | 45.3 ± 9.0  |
| <b>13d</b> | <i>m</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>9</sub> H <sub>19</sub> | 0 | 23.7 ± 0.7  |
| <b>13e</b> | <i>p</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>5</sub> H <sub>11</sub> | 0 | 47.7 ± 9.6  |
| <b>13f</b> | <i>p</i> -CH <sub>2</sub> PO <sub>3</sub> H <sub>2</sub> | C <sub>9</sub> H <sub>19</sub> | 0 | 30.7 ± 5.4  |

**[00200]** Data obtained from benzoic acids **5a-f** indicate that in all cases, regardless of the position of the carboxylate with respect to the sulfonamide, the longer C<sub>9</sub> alkyl chain resulted in greater inhibition than the C<sub>5</sub> saturated chain. The most effective orientation between the acid and sulfonamide appeared to be *ortho*-substitution, as **5f** (IC<sub>50</sub> = 22.7 μM) is a better inhibitor than either **5b** (IC<sub>50</sub> = 43.9 μM) or **5d** (IC<sub>50</sub> = 28.5 μM). The assay data from phosphonic acids **13a-f** also indicated that the longer C<sub>9</sub> alkyl chain is more effective. In this series of compounds, however, there is no significant difference in activity between the different orientations of the phosphonic acid and the alkyl sulfonamide moiety. The most active compound of this class was **13d** (IC<sub>50</sub> = 23.7 μM), the *meta*-substituted phosphonic acid, though not by much over **13b** (IC<sub>50</sub> = 30.6 μM) and **13f** (IC<sub>50</sub> = 30.7 μM).

**Table 2.** *In Vitro* Anti-mtGPAT1 Activity of Sulfonamides **15a-i** and **17a-f**




| Compound         | X                           | Y                                | IC <sub>50</sub> (μM) ± SD |
|------------------|-----------------------------|----------------------------------|----------------------------|
| <b>15a</b>       | <i>p</i> -CO <sub>2</sub> H | C <sub>9</sub> H <sub>19</sub>   | 29.1 ± 4.3                 |
| <b>15b</b>       | <i>p</i> -CO <sub>2</sub> H | Ph                               | 41.9 ± 5.3                 |
| <b>15c</b>       | <i>p</i> -CO <sub>2</sub> H | 4-ClPh                           | 33.7 ± 1.3                 |
| <b>15d</b>       | <i>m</i> -CO <sub>2</sub> H | C <sub>9</sub> H <sub>19</sub>   | 24.2 ± 2.9                 |
| <b>15e</b>       | <i>m</i> -CO <sub>2</sub> H | Ph                               | 38.3 ± 7.6                 |
| <b>15f</b>       | <i>m</i> -CO <sub>2</sub> H | 4-ClPh                           | 23.6 ± 1.2                 |
| <b>15g (C67)</b> | <i>o</i> -CO <sub>2</sub> H | C <sub>9</sub> H <sub>19</sub>   | 8.1 ± 0.7                  |
| <b>15h</b>       | <i>o</i> -CO <sub>2</sub> H | Ph                               | 40.5 ± 2.6                 |
| <b>15i</b>       | <i>o</i> -CO <sub>2</sub> H | 4-ClPh                           | 33.5 ± 2.5                 |
| <b>17a</b>       | <i>p</i> -CO <sub>2</sub> H | CH <sub>2</sub> Ph               | 64.5 ± 11.6                |
| <b>17b</b>       | <i>p</i> -CO <sub>2</sub> H | C <sub>2</sub> H <sub>4</sub> Ph | 63.0 ± 12.9                |
| <b>17c</b>       | <i>m</i> -CO <sub>2</sub> H | CH <sub>2</sub> Ph               | 52.1 ± 9.0                 |
| <b>17d</b>       | <i>m</i> -CO <sub>2</sub> H | C <sub>2</sub> H <sub>4</sub> Ph | 50.3 ± 4.4                 |
| <b>17e</b>       | <i>o</i> -CO <sub>2</sub> H | CH <sub>2</sub> Ph               | 40.7 ± 1.2                 |
| <b>17f</b>       | <i>o</i> -CO <sub>2</sub> H | C <sub>2</sub> H <sub>4</sub> Ph | 46.4 ± 4.5                 |

**[00201]** The distance between the benzene ring and the sulfonamide sulfur does not appear to have a significant effect on the inhibitory activity of these compounds, as there is effectively no difference between one methylene and two methylene linkers. It is apparent, however, that the *ortho*-substituted compounds containing these linker methylenes (**17e-f**) are more effective than the other substituted benzoic acids (**17a-d**). For the *meta*- and *para*-compounds, inhibitory activity is greater when the benzene ring is directly attached to the sulfur, although the *ortho*-

compounds are all similar. The addition of a *para*-chloride on the benzene ring leads to slight increases in activity for the *para*- (**15c**), *meta*- (**15f**), and *ortho*-compounds (**15i**). Compounds **15a**, **15d**, and **15g** were easily obtainable targets, which allowed for examination of the effect of the methylene linker between the benzene ring and the sulfonamide in **5a-f**. For every substitution, these compounds were the most effective GPAT inhibitors, with the *ortho*-compound (**15g**, **C67**) demonstrating the greatest activity ( $IC_{50} = 8.1 \mu M$ ). Based on these results, a long alkyl chain is preferable to a simple benzene ring.

**Table 3.** *In Vitro* Anti-mtGPAT1 Activity of Sulfonamides **21a-c** and **24a-f**










| Compound   | X                                        | Y                               | Z  | $IC_{50} (\mu M) \pm SD$ |
|------------|------------------------------------------|---------------------------------|----|--------------------------|
| <b>21a</b> | <i>p</i> -PO <sub>3</sub> H <sub>2</sub> | C <sub>8</sub> H <sub>17</sub>  | H  | 33.3 ± 3.8               |
| <b>21b</b> | <i>m</i> -PO <sub>3</sub> H <sub>2</sub> | C <sub>8</sub> H <sub>17</sub>  | H  | 25.3 ± 5.4               |
| <b>21c</b> | <i>o</i> -PO <sub>3</sub> H <sub>2</sub> | C <sub>8</sub> H <sub>17</sub>  | H  | 25.7 ± 2.5               |
| <b>24a</b> | CO <sub>2</sub> H                        | CH <sub>3</sub>                 | H  | 28.6 ± 4.6               |
| <b>24b</b> | CO <sub>2</sub> H                        | C <sub>14</sub> H <sub>29</sub> | H  | 6.9 ± 0.5                |
| <b>24c</b> | CO <sub>2</sub> H                        | C <sub>16</sub> H <sub>33</sub> | H  | 7.8 ± 0.8                |
| <b>24d</b> | CO <sub>2</sub> H                        | C <sub>9</sub> H <sub>19</sub>  | Cl | 11.5 ± 0.7               |
| <b>24e</b> | CO <sub>2</sub> H                        | C <sub>8</sub> H <sub>17</sub>  | OH | 38.2 ± 4.1               |
| <b>24f</b> | CO <sub>2</sub> H                        | C <sub>8</sub> H <sub>17</sub>  | F  | 29.5 ± 2.6               |






[00202] In view of the increased inhibitory activity of **15g**, two other compound series were prepared. The first, **21a-c**, probes the effectiveness of an aryl phosphonic acid in place of






the benzoic acid moiety. *In vitro*, the *ortho*-substituted acid (**21c**) is less active than **15g**, and substitution of the phosphonic acid moiety does not appear to significantly affect activity (**Table 3**). The other compounds produced (**24a-f**) indicate the importance of chain length of the alkyl sulfonamide, as well as the effect of adding heteroatoms *para*- to the sulfonamide. It appears that the longer chain is very important to the activity of these compounds, as a C<sub>1</sub>-chain (**24a**) results in significantly less *in vitro* activity than the C<sub>9</sub> chain. Compounds **24b** and **24c** were produced to determine if the naturally-favored C<sub>16</sub> chain is preferred in these compounds over other chain lengths, including the C<sub>14</sub> chain. In this case, there is no observed preference for the C<sub>16</sub> compound over other long chains, in contrast to that observed with the natural acyl-CoA substrates.






**[00203]** Results for compounds 4a-t and 7a-t, which were developed using the methods described above, include the following:






|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 107% at 6.25 ug/ml                  |             | 7.8 ± 1.1 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 115% at 6.25 ug/ml                  |             | 6.8 ± 0.5 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |






|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 89% at 1.56 ug/ml                   |             | 9.8 ± 0.9 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 83% at 0.098 ug/ml                  |             | 8.3 ± 0.4 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 98% at 6.25 ug/ml                   |             | 12.6 ± 2.1 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 83% at 0.098 ug/ml                  |             | 8.9 ± 1.1 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 92% at 0.395 ug/ml                  |             | 25.7 ± 0.4 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |






|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 82% at 0.098 ug/ml                  |             | 8.1±1.0 ug/ml         |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 113% at 6.25 ug/ml                  |             | 8.4 ± 0.2 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 112% at 6.25 ug/ml                  |             | 7.4 ± 0.2 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 101% at 1.56 ug/ml                  |             | 6.7 ± 0.2 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 86% at 1.56 ug/ml                   |             | 5.7 ± 0.2 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |





|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 126% at 0.098 ug/ml                 |             | 5.5 ± 0.3 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 124% at 0.395 ug/ml                 |             | 6.1 ± 0.3 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 104% at 0.098 ug/ml                 |             | 12.1 ± 1.3 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 89% at 100ug/ml                     |             | 303 ± 47 ug/ml        |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 95% at 0.395 ug/ml                  |             | 6.3 ± 0.3 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 119% at 0.098 ug/ml                 |             | 30.6 ± 0.8 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 91% at 0.098 ug/ml                  |             | 9.8 ± 0.7 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 89% at 0.395 ug/ml                  |             | 8.0 ± 1.0 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 104% at 1.56 ug/ml                  |             | 29.8 ± 2.6 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 87% at 0.098 ug/ml                  |             | 8.8 ± 0.8 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

|                                                                                          |                         |                                     |             |                       |
|------------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|         | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                          | Neg                     | 100% at 0.395 ug/ml                 |             | 10.2 ± 0.9 ug/ml      |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                          | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|        | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                          | Neg                     | 90% at 0.098 ug/ml                  |             | 7.9 ± 0.8 ug/ml       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                          | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|       | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                          | Neg                     | 109% at 0.098 ug/ml                 |             | 25.7 ± 3.2 ug/ml      |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                          | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|       | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                          |                         | % at ug/ml                          |             | 23.4 ± 1.0 ug/ml      |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                          | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
| <br>1 | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                          | Not Tested              | Not Tested                          |             |                       |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                          | Neg                     | 103% at 1.56 ug/ml                  |             | 12.7 ± 0.7 ug/ml      |
|                                                                                          |                         |                                     |             |                       |
|                                                                                          | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                          | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     | Neg                     | 102% at 6.25 ug/ml                  |             | 21.2 ± 3.1 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 8.4 ± 1.7 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 8.7 ± 1.4 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 22.7 ± 1.0 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 11.7 ± 0.8 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 10.3 ± 0.9 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 8.8 ± 2.4 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 8.4 ± 1.9 ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 25.4 ± 1.6 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 22.5 ± 0.5 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

|                                                                                     |                         |                                     |             |                       |
|-------------------------------------------------------------------------------------|-------------------------|-------------------------------------|-------------|-----------------------|
|    | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 24.7 ± 1.7ug/ml       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|   | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | 26.8 ± 1.4 ug/ml      |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | ug/ml                 |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |
|  | FAS (IC <sub>50</sub> ) | <sup>14</sup> C (IC <sub>50</sub> ) |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | CPT I Stim              | Weight Loss                         |             |                       |
|                                                                                     | Not Tested              | Not Tested                          |             |                       |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | FAO SC 150              | FAO Max                             |             | GPAT IC <sub>50</sub> |
|                                                                                     |                         | % at ug/ml                          |             | ug/ml                 |
|                                                                                     |                         |                                     |             |                       |
|                                                                                     | SA/MH (MIC)             | SA/Tsoy(MIC)                        | EF/MH (MIC) | EF/Tsoy(MIC)          |
|                                                                                     | ug/ml                   | ug/ml                               | ug/ml       | ug/ml                 |

[00204]      *Example 9 - In Vivo Testing*

[00205]      *Experimental Procedures*

[00206] *DIO and lean mouse models.* All animal experimentation was done in accordance with guidelines on animal care and use as established by the Johns Hopkins University School of Medicine IACUC. DIO C57BL6J male mice were obtained from Jackson Laboratory (Bar Harbor, ME) and fed a synthetic diet comprised of 60% calories from fat, 20% from carbohydrate, and 20% from protein (5.2 kcal/g) post-weaning through the experimental procedures (D12492i, Research Diets, Inc., New Brunswick, NJ). For lean animal studies, twelve-week old C57BL6J male mice (Jackson Laboratory, Bar Harbor, ME) were fed rodent chow comprised of 13% calories from fat, 58% from carbohydrate, and 29% from protein (4.1 kcal/g) (Prolab RMH 2500, PMI Nutrition International Inc., Brentwood, MO). Mice were maintained in 12 hr light-dark cycle at 25°C for 1 week for acclimatization prior to treatment. In all studies, FSG67 (FASgen, Inc., Baltimore, MD) was dissolved in RPMI 1640 (Invitrogen, Carlsbad, CA).

[00207] For acute studies, 6 DIO or lean mice were treated with a single dose of FSG67 (20 mg/kg, i.p.) approximately 3 hrs past lights-on. Animal weights and food consumption were measured 18 h after treatment. Following euthanization, the hypothalamuses were harvested to measure orexigenic and anorexigenic neuropeptide gene expression. In the chronic studies, DIO mice, 4-10 animals per group, were treated daily with FSG67 (5 mg/kg, i.p.) or with RMPI vehicle for the days indicated. Body weight and food intake were measured daily. In one study, a cohort of mice was pair-fed with amounts consumed by the FSG67-treated animals and mice were monitored with indirect calorimetry (Oxymax Equal Flow System®, Columbus Instruments, Columbus, OH). Measurements of VO<sub>2</sub> (ml/kg/hr) and VCO<sub>2</sub> (ml/kg/hr) were performed and recorded every 15 min. The respiratory exchange ratio (RER) was calculated by Oxymax software, version 5.9, and is defined as ratio of VCO<sub>2</sub> to VO<sub>2</sub> 33. After completion of

the treatment course, animals were euthanized by CO<sub>2</sub> inhalation 4 hrs following the final dose of FSG67. Tissues were harvested immediately for RNA extraction; serum was collected and analyzed for glucose, cholesterol, and triglyceride measurements (Bioanalytics, Gaithersburg, MD). Fresh liver tissue was snap frozen in liquid N<sub>2</sub>, sectioned, and stained with hematoxylin and Oil Red O to visualize triglyceride droplets.

**[00208]**      *Chronic lateral cerebroventricle cannulas.* For experiments requiring intracerebroventricular (i.c.v.) administration of compounds, mice were outfitted unilaterally with chronic indwelling cannulas aimed at the lateral cerebroventricle. After mice recovered from surgeries for one week, cannula placements were assessed by measuring food intake in response to i.c.v. neuropeptide Y (NPY, American Peptide Co., CA). Mice were given NPY (0.25 nmol/2 µl injection) or sterile 0.9% saline vehicle via the i.c.v. cannula, and allowed 1-h access to grain-based pellets during the light phase. Mice that ate at least 0.5 g of food after NPY were used in the experiments. Eleven mice were given a 2 µL injection of RPMI-1640 without glucose (Cambrex, MD) for vehicle control. Three days later, six mice received a 100 nmole dose of FSG67 in the vehicle while 5 mice received 320 nmoles of compound.

**[00209]**      *Q-NMR assessment of adiposity.* Following 10 days of FSG67 treatment or vehicle by ip administration, the DIO mice were euthanized and carcasses were stored at -80°C. Carcasses were thawed for Q-NMR analysis. Measurement of fat, lean, and water mass was performed using an EchoMRI-100™ (Echo Medical Systems, Houston, TX) in the Molecular and Comparative Pathobiology Phenotyping Core.

**[00210]**      *Conditioned taste aversion.* Ten days before testing, eighteen male C57/BL6 mice were placed on a schedule of 2 h daytime access to water. On the test day, mice were divided

into three groups and were given access to 0.15% sodium saccharin rather than water for 30 min. Immediately after saccharin access, mice were injected ip with RPMI vehicle or FSG67 (5 and 20 mg/kg body wt) and were allowed water access for the remaining 90 min. Twenty-four hours later, mice were given 2h access to a two-bottle choice test of 0.15% saccharin vs. water. Intakes of both solutions were recorded, and data were expressed as saccharin preference (100 X saccharin intake/saccharin intake + water intake).

[00211] *Real-time RT-PCR.* Hypothalamus, liver, and WAT of DIO and lean mice were harvested and immediately frozen in liquid nitrogen. Total RNA was isolated and real-time quantitative RT-PCR was performed as previously described (13). Gene-specific primer pairs were designed using Primer3 software ([http://www-genome.wi.mit.edu/cgi-bin/primer/primer3\\_www.cgi/](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi/)). The sequences of the primer pairs are listed in Supplemental Data Table 1.

[00212] *3T3-L1 Adipocytes* 3T3-L1 cells were differentiated into adipocytes as described 34. Seven days post-differentiation, cells were treated with FSG67 at indicated concentrations for 18 h, then labeled with [<sup>14</sup>C]palmitate for 2 h. Following Folch extraction, lipids were subjected to polar and non-polar thin-layer chromatography 35. Triglyceride and phosphatidylcholine fractions were quantified with phosphorimaging (Storm 840, Molecular Dynamics, Piscataway, NJ).

[00213] *Statistical analysis.* All data are presented as means  $\pm$  standard error of the mean. IC50 determinations were performed with linear regression. Two-tailed unpaired t-tests or two-way ANOVA tests were performed as indicated using Prism 4.0 (Graph Pad Software, San Diego, CA).

[00214] *FSG67 reduces acylglyceride synthesis in mouse 3T3-L1 adipocytes.*

[00215] Mouse 3T3-L1 adipocytes were used to test the effect FSG67 on acylglyceride synthesis *in vitro*. 3T3-L1 adipocytes, at 7 days post-differentiation, were treated with FSG67 at concentrations of 7.6  $\mu\text{M}$  to 61  $\mu\text{M}$  (2.5 - 20  $\mu\text{g/ml}$ ) and the IC<sub>50</sub> values for inhibition of triglyceride and phosphatidylcholine synthesis were determined using linear regression. The IC<sub>50</sub> values were 33.9  $\mu\text{M}$  for cellular triglyceride synthesis ( $p=0.023$ ,  $r^2 = 0.86$ ,  $n=3$ ) and 36.3  $\mu\text{M}$  for phosphatidylcholine synthesis ( $p=0.015$ ,  $r^2 = 0.89$ ,  $n=3$ ). As phosphatidylcholine was the predominant phospholipid synthesized in the 3T3-L1 adipocytes, it is representative of overall cellular phospholipid synthesis. These IC<sub>50</sub> values are similar to the reported IC<sub>50</sub> value of 24.7  $\mu\text{M}$  for mouse mitochondrial GPAT activity 12. Consistent with its inhibition of acylglyceride synthesis, Figure 10 shows the dose-dependent reduction of triglyceride accumulation in 3T3-L1 adipocytes 48 h following FSG67 treatment. Note the decrease in lipid droplets in the FSG67 treated cells compared to vehicle treated controls. Thus, FSG67 inhibits cellular acylglyceride synthesis with an IC<sub>50</sub> similar to its inhibition of GPAT activity in mitochondrial preparations. In keeping with these biochemical observations, FSG67 substantially reduced triglyceride accumulation in cultured adipocytes. Taken together, these results demonstrate that FSG67 inhibits cellular GPAT activity.

[00216] Acute FSG67 treatment of lean and DIO mice reduced body weight, and decreased food consumption without conditioned taste aversion. Since FSG67 reduced acylglyceride synthesis *in vitro*, we tested both lean and DIO mice with a single dose of FSG67 (20 mg/kg i.p.) to examine the acute effect on animal weight and feeding behavior. In addition, we performed conditioned taste aversion (CTA) testing to determine if FSG67 triggers a CTA response that might suggest malaise as the cause of reduced food intake. Eight DIO and lean

mice were treated with FSG67 at the beginning of dark cycle. Within 24 h, the lean mice injected with FSG67 lost  $3.7 \pm 0.9\%$  ( $1.0 \pm 0.2$  g) of body mass while fasted mice lost  $15.5 \pm 0.7\%$  ( $3.9 \pm 0.2$  g) (Fig. 11a). The reduction in body mass of both groups was significant compared to vehicle controls which gained  $2.5 \pm 0.5\%$  ( $0.6 \pm 0.1$ g) ( $p < 0.0001$ , 2-tailed t-test). FSG67 treatment also reduced food intake to 33% of vehicle control ( $p < 0.0001$  2-tailed t-test) (Fig. 11b).

**[00217]** GPAT inhibition with FSG67 decreased body weight in DIO mice consuming a high fat diet. FSG67 treated DIO mice lost  $4.3 \pm 0.5\%$  ( $1.7 \pm 0.2$  g) of body mass versus a  $5.3 \pm 0.4\%$  ( $2.1 \pm 0.2$  g) loss for fasted mice (Fig. 11c). Compared to the vehicle control mice which lost  $2.5 \pm 0.6\%$  ( $1.0 \pm 0.2$  g) the weight loss was significant in both the FSG67 treated ( $p = 0.026$ , 2-tailed t-test) and fasted mice ( $p = 0.002$ , 2-tailed t-test). FSG67 significantly reduced food consumption in the DIO mice to 41.6% of vehicle control (Fig. 11d). While the average food intake between the DIO and lean vehicle control groups is substantially different (1.2 and 4.2 g, respectively) ( $p < 0.0001$ , 2-tailed t-test), the relative reduction of food intake following FSG67 treatment is not different between the DIO (41.6% of vehicle control) and lean mice (33% of vehicle control) ( $p = 0.19$ , Fisher's exact test). CTA testing in groups of 8 lean mice using a two bottle choice paradigm showed that FSG67 failed to produce a significant reduction in saccharin intake at 5 mg/kg ( $p = 0.12$ ) or 20 mg/kg ( $p = 0.10$ , 2-tailed t-test). Thus, the reduction in food intake from FSG67 was not due to sickness behavior (Fig. 11e). No overt toxicity was noted from the FSG67 treatment of the lean or DIO mice. These data demonstrate a clear anorexigenic effect of pharmacological GPAT inhibition in both lean and DIO mice with accompanying reduction in animal weight.

**[00218]** *Chronic FSG67 treatment of DIO mice reversibly reduced body weight and food consumption, and increased fatty acid oxidation.* To determine the dose of FSG67 suitable for

chronic treatment, we performed a 5-day dose ranging study in DIO mice, four per group, with daily intraperitoneal doses of 1, 2, and 5 mg/kg (Fig. 17). The 5 mg/kg dose led to significant weight loss of 3.9% compared to vehicle controls ( $p=0.008$ , 2-way ANOVA). This dose was chosen for the subsequent chronic treatment experiments.

**[00219]** The first chronic treatment experiment was designed to test if weight loss induced by FSG67 was reversible. Four DIO mice per group were treated with FSG67 or vehicle for 20 days. For the entire 32 d trial, weight and food consumption were recorded daily until the FSG67 treated animals regained their original weight. During FSG67 treatment (days 0-20), the mice lost  $10.3 \pm 0.6\%$  of their body mass while controls gained  $4.0 \pm 0.5\%$  ( $p<0.0001$ , 2-way ANOVA) (Fig 12a). Average food consumption was reduced during FSG67 treatment ( $2.6 \pm 0.1$  g/d, days 1-20) compared to vehicle controls ( $3.1 \pm 0.1$  g) ( $p=0.0008$ , 2-way ANOVA) (Fig 12b). Following cessation of treatment, food consumption increased in the FSG67 treatment group to an average of  $3.5 \pm 0.1$  g/d (days 21-32) representing a significant increase in food intake compared to vehicle controls  $3.2 \pm 0.1$  g/d ( $p=0.006$ , 2-way ANOVA). The FSG67 treated animals achieved their average pre-treatment weight 11 days following termination of treatment (Fig 12a).

**[00220]** In the second chronic treatment study, indirect calorimetry was utilized to study changes in metabolism during GPAT inhibition. DIO mice (8 per group) were treated with FSG67 (5 mg/kg, ip), or pair-fed to FSG67 treated animals. Indirect calorimetry was utilized to measure changes in oxygen consumption ( $VO_2$ ) and respiratory exchange ratio (RER) between pair-fed and treated animals. After 16 days of treatment, the FSG67 treated mice lost  $9.5 \pm 0.6\%$  of body mass, pair-fed lost  $5.5 \pm 0.9\%$ , while vehicle controls gained  $3.5 \pm 1.3\%$  (Fig. 12c). The weight loss in the FSG67 treated animals was significant compared to both vehicle controls and

pair-fed animals ( $p < 0.0001$ , 2-way ANOVA). FSG67 treatment again significantly reduced food consumption by 33%,  $2.0 \pm 0.1$  g/d in the FSG67 treated group compared to  $3.1 \pm 0.1$  g/d for vehicle controls ( $p < 0.0001$ , 2-way ANOVA) (Fig. 12d). FSG67 treatment increased the average  $\text{VO}_2$  to  $106.5 \pm 1.1\%$  of pre-treatment value. This value was significantly increased compared to pair-fed mice, which displayed a reduction in  $\text{VO}_2$  to  $89.9 \pm 1.1\%$  of the pre-treatment value ( $p < 0.0001$ , 2-way ANOVA) (Fig. 12e). RER was reduced in FSG67 treated mice ( $0.732 \pm 0.002$ ) compared to pair-fed ( $0.782 \pm 0.006$ ) ( $p < 0.0001$ , 2-way ANOVA) (Fig. 12f) indicating increased use of fatty acids for fuel in the FSG67 treated animals. The combination of increased  $\text{VO}_2$  and reduced RER in the FSG67 treated animals are consistent with increased fatty acid oxidation and energy utilization which likely contribute to their reduced body mass compared to the pair-fed controls.

**[00221]      *Pharmacological GPAT inhibition reduced adiposity and down-regulated lipogenic gene expression in DIO mice.*** Since FSG67 increased fatty acid oxidation and reduced food intake in DIO mice, we next used Q-NMR analysis to measure lean, fat and water mass in FSG67 treated and control mice to determine the composition of the tissue loss with FSG67 treatment. In an additional chronic treatment experiment, 10 DIO mice were treated with FSG67 (5 mg/kg/d, ip) and 10 received vehicle for 10 days. The FSG67 treated mice lost  $6.1 \pm 0.9$  g ( $13.1 \pm 1.9\%$ ) while vehicle controls lost  $1.1 \pm 0.4$  g ( $2.3 \pm 0.8\%$ ) ( $p < 0.0001$ , 2-way ANOVA). (Fig. 18) Q-NMR analysis demonstrated a 4.0 g reduction in fat mass in the FSG67 treated animals compared to vehicle control ( $p < 0.0001$ , 2-tailed t-test) but no significant change in lean or water mass (Fig. 13a). At the conclusion of the experiment, the FSG67 treated mice weighed 4.4 g less than the vehicle controls, which could be accounted for by the 4.0 g difference in fat mass. Thus, GPAT inhibition selectively reduces adiposity in DIO mice.

[00222] To further explore the mechanism responsible for the reduction in adipose tissue mass, we used real-time RT-PCR to measure the expression of the following key lipogenic genes in white adipose tissue from vehicle control, pair-fed, and FSG67 treated DIO mice from the second indirect calorimetry trial (see Fig. 12c): fatty acid synthase (FAS), responsible for the *de novo* reductive synthesis of fatty acid 13, acetyl-CoA carboxylase 1 (ACC1), the cytoplasmic isoform of ACC expressed in lipogenic organs that synthesizes malonyl-CoA used as a substrate of FAS for fatty acid synthesis 14, peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) a key transcription factor for adipogenesis 15, lipid partitioning 16, and postprandial lipid storage 17, and GPAT. After 16 days of treatment, real-time RT-PCR analysis of white adipose tissue from FSG67 treated animals showed substantial down-regulation of ACC1 (p=0.0005 vs. control, p=0.0004 vs. pair-fed), FAS (p=0.0001 vs. control, p=0.0007 vs. pair-fed), PPAR $\gamma$  (p=0.032 vs. control, p=0.0019 vs. pair-fed), and GPAT (p=0.0034 vs. control, p=0.0002 vs. pair-fed) (Fig. 13b). Interestingly, uncoupling protein-2 (UCP2) expression was increased in both liver (p=0.043 vs. control) and white adipose tissue (p=0.013, vs. pair-fed) of the FSG67 treated animals which could also contribute to increased fatty acid oxidation 18; L-CPT-1 expression was unaffected. (Fig. 19). Thus, pharmacological GPAT inhibition not only increases fatty acid oxidation and reduces food intake, but up-regulates UCP2 in liver and white adipose tissue while down-regulating lipogenic gene expression in white adipose tissue, all of which should favor a selective decrease in adiposity.

[00223] *FSG67 substantially reduced serum glucose and triglyceride levels while resolving hepatic steatosis in DIO mice.* Consistent with the systemic reduction in adiposity, GPAT inhibition reversed hepatic steatosis in DIO mice. Oil red-O staining of frozen sections of liver showed marked steatosis characterized by large and small droplet triglyceride accumulation

in the vehicle treated animals (Fig 14a). Steatosis was reduced in the pair-fed animals (Fig 14b) with nearly complete resolution with FSG67 treatment (Fig 14c). No inflammation, necrosis, or hepatocellular injury was identified. Real-time RT-PCR expression analysis of the hepatic lipogenic genes, ACC1, FAS, and GPAT showed a significant reduction in FAS ( $p=0.0016$  vs. control,  $p=0.018$  vs. pair-fed) and ACC1 ( $p=0.037$  vs. pair-fed) expression but not GPAT, indicating a down-regulation of *de novo* fatty acid synthesis with FSG67 treatment. (Fig. 20) In addition to the reduction of tissue triglycerides, serum glucose levels were reduced ( $153.3 \pm 10.5$  mg/dL) compared with both vehicle control mice ( $200.6 \pm 22.2$  mg/dL,  $p=0.03$  2-way ANOVA) and pair-fed ( $189.0 \pm 20.3$  mg/dL,  $p=0.04$ , 2-way ANOVA). The reduction in serum triglyceride levels seen in the FSG67 treated DIO mice ( $111.3 \pm 10.9$  mg/dL) compared to pair-fed ( $138.5 \pm 9.8$  mg/dL) or vehicle controls ( $138.8 \pm 13.5$  mg/dL) were not statistically significant. Cholesterol levels remained unchanged (Fig. 14d). The resolution of the hepatic steatosis in FSG67 treated mice may have contributed to the normalization of blood glucose levels.

**[00224]      *Intracerebroventricular (icv) FSG67 treatment reduced food consumption and body weight.*** We administered FSG67 icv to determine whether GPAT inhibition acts centrally to reduce food intake. Lean mice were treated with FSG67 icv at doses 100 and 320 nmoles (approximately 300- and 100-fold less than the 5 mg/kg single day systemic dose). Within 24 h, mice treated with 100 nmoles lost  $0.75 \pm 0.4$  g ( $p=0.016$ ) while the 320 nmole group lost  $1.8 \pm 0.3$  g ( $p=0.0003$ ); vehicle controls gained  $0.43 \pm 0.1$  g and  $0.33 \pm 0.1$  g respectively (Fig 15a). The animal weight was regained within 48 h without a significant rebound (data not shown). Significant reduction in chow intake only occurred in the 320 nmole treatment group ( $3.8 \pm 0.1$  g control,  $2.5 \pm 0.3$  g FSG67,  $p=0.0051$ ) (Fig 15b). Within 48 h, the animals began eating normally with slight hyperphagia in the 320 nmole group on days 3 and 4 (data not shown). These data

indicate that the reduction in food consumption accompanying GPAT inhibition may have a significant contribution from the CNS. Moreover, the occurrence of weight loss without a reduction of food intake in the 100 nmole group suggests a central effect on metabolism independent of changes in food intake behaviors.

**[00225]      *Acute and chronic FSG67 treatment altered hypothalamic neuropeptide***


***expression.*** Hypothalamic peptide expression was measured in the lean and DIO mice treated with a single dose of FSG67 (see Fig 11) and in the chronically treated DIO mice (see Fig 12) to further assess the mechanism responsible for reduced food intake. In the lean mice treated with a single dose of FSG67, the expression of the orexigenic hypothalamic neuropeptide neuropeptide-Y (NPY) was significantly reduced compared to the fasted animals ( $p=0.012$ , 2-tailed t-test) while agouti-related protein (AgRP) expression was substantially diminished compared to both fasted ( $p=0.020$ , 2-tailed t-test) and vehicle controls ( $p=0.0009$ , 2-tailed t-test) consistent with the acute reduction in food intake (Fig. 16a). Conversely, the anorexigenic neuropeptides, pro-opiomelanocortin (POMC) and cocaine-amphetamine-related transcript (CART) mRNA levels were not affected by food deprivation or acute FSG67 treatment. In contrast to the findings in lean mice, single dose FSG67 treatment of DIO mice significantly increased AgRP expression over that in the vehicle controls and food-deprived animals (data not shown). Notably, food deprivation did not result in increased levels of hypothalamic NPY or AgRP message in DIO mice as was seen with the lean animals. This pattern of increased orexigenic neuropeptide expression with treatment is consistent with a hunger response and may indicate a rebound of orexigenic peptide expression in the DIO mice or could represent a further example of dysregulated neuropeptide signaling in DIO mice 19. In the chronically treated DIO mice, however, hypothalamic neuropeptide analysis showed a significant reduction in NPY expression

in both FSG67 ( $p=0.0052$ , 2-tailed t-test) and pair-fed animals ( $p=0.0074$ , 2-tailed t-test) compared to vehicle controls (Fig. 16b). This profile was more similar to the acutely treated lean mice, and may reflect normalization of the appetite response in the chronically treated DIO mice.

## CLAIMS

We Claim:

- 1) A compound comprising a formula I:



wherein

n is either 0 or 1;

A is selected from the group consisting of  $\text{NR}^1$ , O, and S, wherein  $\text{R}^1$  is selected from the group consisting of H, hydroxyl,  $\text{C}_1\text{-C}_{10}$  alkyl,  $\text{C}_1\text{-C}_{10}$  alkoxy, alkenyl, aryl, alkylaryl and arylalkyl;

X is selected from the group consisting of a carboxylate residue, a phosphonate residue, a phosphate residue, and a  $\text{C}_1\text{-C}_{10}$  alkyl residue which is optionally substituted with one or more residues selected from the group consisting of a carboxylate residue, a phosphonate residue and a phosphate residue;

Y is selected from the group consisting of  $\text{C}_1\text{-C}_{20}$  alkyl, alkenyl, halide, hydroxyl,  $\text{C}_1\text{-C}_{20}$  alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl and a heterocyclic ring, any of which are optionally substituted at one or more positions with a halogen; and


Z is selected from the group consisting of H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring, any of which are is optionally substituted at one or more positions with one or a combination of substitution groups selected from the group consisting of a  $\text{C}_1\text{-C}_{10}$  alkyl group,  $\text{C}_1\text{-C}_{10}$  alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group,



an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring.

- 2) The compound of claim 1 wherein A is comprised of  $\text{NR}^1$  wherein  $\text{R}^1$  is a hydrogen.
- 3) The compound of claim 1 wherein X is comprised of a carboxylic acid residue.
- 4) The compound of claim 1 wherein X is comprised of a phosphonate residue.
- 5) The compound of claim 1 wherein X is comprised of a methyl or ethyl residue substituted with a phosphonate residue or carboxylate residue.
- 6) The compound of claim 1 wherein X is in the ortho or meta position with respect to the sulfonyl linker of the phenyl ring.
- 7) The compound of claim 1 wherein Y is comprised of a  $\text{C}_1\text{-C}_{20}$  alkyl group selected from the group consisting of  $\text{CH}_3$ ,  $\text{C}_5\text{H}_{11}$ ,  $\text{C}_8\text{H}_{17}$ ,  $\text{C}_9\text{H}_{19}$ ,  $\text{C}_{14}\text{H}_{29}$ , and  $\text{C}_{16}\text{H}_{33}$ .
- 8) The compound of claim 1 wherein Y is selected from the group consisting of an aryl, alkylaryl, and arylalkyl residue, any of which is optionally substituted with one or more halogen atoms.
- 9) The compound of claim 8 wherein Y is comprised of 4-ClPh.
- 10) The compound of claim 1 wherein Y is comprised of an alkylaryl residue having  $\text{C}_1\text{-C}_3$  carbon atoms.
- 11) The compound of claim 10 wherein the aryl moiety of the alkylaryl residue is substituted with one or more halogen atoms.
- 12) The compound of claim 1 wherein Z is selected from the group consisting of H, F, Cl, or OH.

- 14) The compound of claim 1 selected from the group consisting of:







wherein

$n$  is either 0 or 1;

$A$  is selected from the group consisting of  $NR^1$ , O, and S, wherein  $R^1$  is selected from the group consisting of H, hydroxyl,  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy, alkenyl, aryl, alkylaryl and arylalkyl;

Y is selected from the group consisting of C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, halide, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl and a heterocyclic ring, any of which are optionally substituted at one or more positions with a halogen; and

Z is selected from the group consisting of H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring, any of which are optionally substituted at one or more positions with one or a combination of substitution groups selected from the group consisting of a C<sub>1-10</sub> alkyl group, C<sub>1-10</sub> alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring.

16) The compound of claim 15 wherein A is comprised of NR<sup>1</sup> wherein R<sup>1</sup> is a hydrogen.

17) The compound of claim 15 wherein Y is comprised of a C<sub>1</sub>-C<sub>20</sub> alkyl group selected from the group consisting of CH<sub>3</sub>, C<sub>5</sub>H<sub>11</sub>, C<sub>8</sub>H<sub>17</sub>, C<sub>9</sub>H<sub>19</sub>, C<sub>14</sub>H<sub>29</sub>, and C<sub>16</sub>H<sub>33</sub>.


18) The compound of claim 15 wherein Y is selected from the group consisting of an aryl, alkylaryl, and arylalkyl residue, any of which is optionally substituted with one or more halogen atoms.


19) The compound of claim 18 wherein Y is comprised of 4-ClPh.


20) The compound of claim 15 wherein Y is comprised of a alkylaryl residue having C<sub>1</sub>-C<sub>3</sub> carbon atoms.

21) The compound of claim 20 wherein the aryl moiety of the alkylaryl residue is substituted with one or more halogen atoms.


- 22) The compound of claim 15 wherein Z is selected from the group consisting of H, F, Cl, OH, an optionally substituted aryl group and an optionally substituted heterocyclic ring.
- 23) The compound of claim 15 wherein COOH is in an ortho or meta position with respect to the sulfonyl linker of the phenyl ring.
- 24) The compound of claim 15 wherein Z is in either a meta or para position with respect to the sulfonyl linker of the phenyl ring.
- 25) The compound of claim 15 selected from the group consisting of:







26) A compound comprising a formula IVb:



wherein

n is either 0 or 1;


A is selected from the group consisting of NR<sup>1</sup>, O, and S, wherein R<sup>1</sup> is selected from the group consisting of H, hydroxyl, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, alkenyl, aryl, alkylaryl and arylalkyl;


Y is selected from the group consisting of C<sub>1</sub>-C<sub>20</sub> alkyl, alkenyl, halide, hydroxyl, C<sub>1</sub>-C<sub>20</sub> alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl and a heterocyclic ring, any of which are optionally substituted at one or more positions with a halogen; and

Z is selected from the group consisting of H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring, any of which are optionally substituted at one or more positions with one or a combination of substitution groups selected from the group consisting of a C<sub>1-10</sub> alkyl group, C<sub>1-10</sub> alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group and a heterocyclic ring.


- 27) The compound of claim 26 wherein A is comprised of NR<sup>1</sup> wherein R<sup>1</sup> is a hydrogen.
- 28) The compound of claim 26 wherein Y is comprised of a C<sub>1</sub>-C<sub>20</sub> alkyl group selected from the group consisting of CH<sub>3</sub>, C<sub>5</sub>H<sub>11</sub>, C<sub>8</sub>H<sub>17</sub>, C<sub>9</sub>H<sub>19</sub>, C<sub>14</sub>H<sub>29</sub>, and C<sub>16</sub>H<sub>33</sub>.
- 29) The compound of claim 26 wherein Y is selected from the group consisting of an aryl, alkylaryl, and arylalkyl residue, any of which is optionally substituted with one or more halogen atoms.
- 30) The compound of claim 29 wherein Y is comprised of 4-ClPh.
- 31) The compound of claim 26 wherein Y is comprised of a alkylaryl residue having C<sub>1</sub>-C<sub>3</sub> carbon atoms.
- 32) The compound of claim 31 wherein the aryl moiety of the alkylaryl residue is substituted with one or more halogen atoms.
- 33) The compound of claim 26 wherein Z is selected from the group consisting of H, F, Cl, OH, an optionally substituted aryl group and an optionally substituted heterocyclic ring.
- 34) The compound of claim 26 wherein Z is in an ortho position with respect to the sulfonyl linker of the phenyl ring.

35) The compound of claim 26 selected from the group consisting of:





36) A compound comprising a formula V:



wherein

$n$  is either 0 or 1;

$m$  is either 0, 1, 2, or 3;

A is selected from the group consisting of  $\text{NR}^1$ , O, and S, wherein  $\text{R}^1$  is selected from the group consisting of H, hydroxyl,  $\text{C}_1\text{-C}_{10}$  alkyl,  $\text{C}_1\text{-C}_{10}$  alkoxy, alkenyl, aryl, alkylaryl and arylalkyl;

Y is selected from the group consisting of  $\text{C}_1\text{-C}_{20}$  alkyl, alkenyl, halide, hydroxyl,  $\text{C}_1\text{-C}_{20}$  alkoxy, aryl, alkylaryl, arylalkyl, cycloalkyl, cycloalkenyl and a heterocyclic ring, any of which are optionally substituted at one or more positions with a halogen; and

Z is selected from the group consisting of H, a hydroxyl group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring, any of which are optionally substituted at one or more positions with one or a combination of substitution groups selected from the group consisting of a  $\text{C}_{1-10}$  alkyl group,  $\text{C}_{1-10}$  alkoxy group, a hydroxyl group, a cyano group, a carboxylate group, a halide, an aryl group, an alkylaryl group, an arylalkyl group, a cycloalkyl group, a cycloalkenyl group or a heterocyclic ring.

37) The compound of claim 36 wherein A is comprised of  $\text{NR}^1$  wherein  $\text{R}^1$  is a hydrogen.


38) The compound of claim 36 wherein Y is comprised of a  $\text{C}_1\text{-C}_{20}$  alkyl group selected from the group consisting of  $\text{CH}_3$ ,  $\text{C}_5\text{H}_{11}$ ,  $\text{C}_8\text{H}_{17}$ ,  $\text{C}_9\text{H}_{19}$ ,  $\text{C}_{14}\text{H}_{29}$ , and  $\text{C}_{16}\text{H}_{33}$ .

39) The compound of claim 36 wherein Y is comprised of an aryl residue.


40) The compound of claim 39 wherein the aryl residue is substituted with one or more halogen atoms.


41) The compound of claim 36 wherein Y is comprised of a alkylaryl residue having  $\text{C}_1\text{-C}_3$  carbon atoms.


- 42) The compound of claim 41 wherein the aryl moiety of the alkylaryl residue is substituted with one or more halogen atoms.
- 43) The compound of claim 36 wherein Z is selected from the group consisting of H, F, Cl, OH, an optionally substituted aryl group and an optionally substituted heterocyclic ring.
- 44) The compound of claim 36 wherein  $(\text{CH}_2)_m\text{PO}_3\text{H}_2$  is in an ortho position with respect to the sulfonyl linker of the phenyl ring.
- 45) The compound of claim 36 wherein Z is in a para position with respect to the sulfonyl linker of the phenyl ring.
- 46) The compound of claim 36 selected from the group consisting of:




- 47) A pharmaceutical composition comprising a pharmaceutical diluent and a compound according to any of claims 1, 15, 26, and 36
- 48) The pharmaceutical composition of claim 47 wherein the compound is selected from the group consisting of:









49) The pharmaceutical composition of claim 47 wherein the compound is selected from the group consisting of:




50) A method of inducing weight loss in a subject comprising administering an effective amount of a pharmaceutical composition according to claim 47 to the subject.


51) The method of claim 50 wherein the pharmaceutical composition includes one or more compound selected from the group consisting of:






54) The method of claim 53 wherein the pharmaceutical composition includes one or more compound selected from the group consisting of:






55) The method of claim 53 wherein the pharmaceutical composition includes one or more compounds selected from the group consisting of:




56) A method of increasing fatty acid oxidation in a subject comprising administering an effective amount of a pharmaceutical composition according to claim 47 to the subject.

57) The method of claim 56 wherein the pharmaceutical composition includes one or more compound selected from the group consisting of:





1 / 17

**FIG. 1**

2 / 17




FIG. 2

3 / 17






FIG. 3

4 / 17



- 14a** *p*-, R = C<sub>9</sub>H<sub>19</sub>  
**14b** *p*-, R = Ph  
**14c** *p*-, R = 4-ClPh  
**14d** *m*-, R = C<sub>9</sub>H<sub>19</sub>  
**14e** *m*-, R = Ph  
**14f** *m*-, R = 4-ClPh  
**14g** *o*-, R = C<sub>9</sub>H<sub>19</sub>  
**14h** *o*-, R = Ph  
**14i** *o*-, R = 4-ClPh



- 15a** *p*-, R = C<sub>9</sub>H<sub>19</sub>  
**15b** *p*-, R = Ph  
**15c** *p*-, R = 4-ClPh  
**15d** *m*-, R = C<sub>9</sub>H<sub>19</sub>  
**15e** *m*-, R = Ph  
**15f** *m*-, R = 4-ClPh  
**15g** *o*-, R = C<sub>9</sub>H<sub>19</sub>  
**15h** *o*-, R = Ph  
**15i** *o*-, R = 4-ClPh

FIG. 4

5 / 17






FIG. 5

6 / 17

**FIG. 6**

7 / 17

**FIG. 7**

8 / 17




FIG. 8

9 / 17





FIG. 9

10 / 17

**FIG. 10**

11 / 17




FIG. 11

12 / 17






FIG. 12

13 / 17

**FIG. 13****FIG. 14**

14 / 17




FIG. 15






FIG. 16



15 / 17

**FIG. 17**

16 / 17

**FIG. 18**

17 / 17

**FIG. 19****FIG. 20**




**FIG. 1**

