

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3762499 T5**

(12) **Rettet oversættelse af
europæisk patentskrift**

-
- (51) Int.Cl.: **C 12 N 15/81 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2024-08-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2023-12-20**
- (86) Europæisk ansøgning nr.: **19711486.1**
- (86) Europæisk indleveringsdag: **2019-03-04**
- (87) Den europæiske ansøgnings publiceringsdag: **2021-01-13**
- (86) International ansøgning nr.: **US2019020526**
- (87) Internationalt publikationsnr.: **WO2019173204**
- (30) Prioritet: **2018-03-06 US 201862639183 P**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **DANISCO US INC, 925 Page Mill Road, Palo Alto, CA 94304, USA**
- (72) Opfinder: **Qi, Min, 11 Stuyvesant Drive, Hockessin, Delaware 19707, USA**
TEUNISSEN, Paula Johanna Maria, 925 Page Mill Road, Palo Alto, California 94304, USA
ZHU, Quinn Qun, 544 Revere Road West, Chester, Pennsylvania 19382, USA
- (74) Fuldmægtig i Danmark: **RWS Group, Europa House, Chiltern Park, Chiltern Hill, Chalfont St Peter, Bucks SL9 9FG, Storbritannien**
- (54) Benævnelse: **REDUKTION I ACETATPRODUKTION VED HJÆLP AF GÆR, DER OVERUDTRYKKER PAB1**
- (56) Fremdragne publikationer:
WO-A1-2015/148272
US-A- 5 985 575
US-A1- 2012 276 587
US-A1- 2014 273 144
FRANCESCA MARTANI ET AL: "The Saccharomyces cerevisiae poly(A) binding protein Pab1 as a target for eliciting stress tolerant phenotypes", SCIENTIFIC REPORTS, vol. 5, no. 1, 14 December 2015 (2015-12-14), XP055584725, DOI: 10.1038/srep18318
KYLIE D. SWISHER ET AL: "Localization to, and Effects of Pbp1, Pbp4, Lsm12, Dhh1, and Pab1 on Stress Granules in Saccharomyces cerevisiae", PLOS ONE, vol. 5, no. 4, 2 April 2010 (2010-04-02), page e10006, XP055584715, DOI: 10.1371/journal.pone.0010006
M. SONDEREGGER ET AL: "Metabolic Engineering of a Phosphoketolase Pathway for Pentose Catabolism in Saccharomyces cerevisiae", APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 70, no. 5, 1 May 2004 (2004-05-01), pages 2892-2897, XP055552748, US ISSN: 0099-2240, DOI: 10.1128/AEM.70.5.2892-2897.2004
KYLIE D. SWISHER ET AL: "Localization to, and Effects of Pbp1, Pbp4, Lsm12, Dhh1, and Pab1 on Stress Granules in Saccharomyces cerevisiae", PLOS ONE, vol. 5, no. 4, 2 April 2010 (2010-04-02), pages e10006, XP055584715, DOI: 10.1371/journal.pone.0010006

Fortsættes ...

M. SONDEREGGER ET AL: "Metabolic Engineering of a Phosphoketolase Pathway for Pentose Catabolism in *Saccharomyces cerevisiae*"; APPLIED AND ENVIRONMENTAL MICROBIOLOGY, vol. 70, no. 5, 1 May 2004 (2004-05-01), US, pages 2892 - 2897, XP055552748, ISSN: 0099-2240, DOI: 10.1128/AEM.70.5.2892-2897.2004

DESCRIPTION

TECHNICAL FIELD

[0001] The present compositions and methods relate to modified yeast that over-expresses polyadenylate-binding protein (PAB1). The yeast produces a decreased amount of acetate compared to their parental cells. Such yeast is particularly useful for large-scale ethanol production from starch substrates, where acetate is an undesirable by-product.

BACKGROUND

[0002] First-generation yeast-based ethanol production converts sugars into fuel ethanol. The annual fuel ethanol production by yeast is about 90 billion liters worldwide (Gombert, A.K. and van Maris, A.J. (2015) *Curr. Opin. Biotechnol.* 33:81-86). It is estimated that about 70% of the cost of ethanol production is the feedstock. Since the production volume is so large, even small yield improvements have massive economic impact across the industry.

[0003] The phosphoketolase (PKL) pathway has been genetically engineered into yeast to increase ethanol production, as described in WO2015148272 (Miasnikov et al.). Unfortunately, the engineered strains also produce more acetate than the parental yeast. Acetate is not a desirable by-product as it has negative effects on yeast growth and fermentation. In addition, acetate reduces the pH of left-over water from fermentation and distillation, referred to as backset, which is typically reused for liquefaction of a subsequent batch of substrate. As a result, ethanol producers must adjust the pH of the backset (or liquefact) or increase the amount of fresh water used for liquefaction.

[0004] US2014/273144A1 relates to the use of phosphoketolase and phosphotransacetylase for production of acetyl-coenzyme A derived compounds.

[0005] The need exists to control the amount of acetate produced by yeast, particularly engineered yeast that tend to produce an increased amount of acetate.

SUMMARY

[0006] The present invention is defined by the appended claims. The present compositions and methods relate to modified yeast that over-express the PAB1 polypeptide as defined in the appended claims.

DETAILED DESCRIPTION

I. Definitions

[0007] Prior to describing the present yeast and methods in detail, the following terms are defined for clarity. Terms not defined should be accorded their ordinary meanings as used in the relevant art.

[0008] As used herein, the term "alcohol" refers to an organic compound in which a hydroxyl functional group (-OH) is bound to a saturated carbon atom.

[0009] As used herein, the terms "yeast cells," "yeast strains," or simply "yeast" refer to organisms from the phyla Ascomycota and Basidiomycota. Exemplary yeast is budding yeast from the order Saccharomycetales. Particular examples of yeast are *Saccharomyces* spp., including but not limited to *S. cerevisiae*. Yeast include organisms used for the production of fuel alcohol as well as organisms used for the production of potable alcohol, including specialty and proprietary yeast strains used to make distinctive-tasting beers, wines, and other fermented beverages.

[0010] As used herein, the phrase "engineered yeast cells," "variant yeast cells," "modified yeast cells," or similar phrases, refer to yeast that include genetic modifications and characteristics described herein. Variant/modified yeast do not include naturally occurring yeast.

[0011] As used herein, the terms "polypeptide" and "protein" (and their respective plural forms) are used interchangeably to refer to polymers of any length comprising amino acid residues linked by peptide bonds. The conventional one-letter or three-letter codes for amino acid residues are used herein and all sequence are presented from an N-terminal to C-terminal direction. The polymer can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art.

[0012] As used herein, functionally and/or structurally similar proteins are considered to be "related proteins," or "homologs." Such proteins can be derived from organisms of different genera and/or species, or different classes of organisms (e.g., bacteria and fungi), or artificially designed. Related proteins also encompass homologs determined by primary sequence analysis, determined by secondary or tertiary structure analysis, or determined by immunological cross-reactivity, or determined by their functions.

[0013] As used herein, the term "homologous protein" refers to a protein that has similar activity and/or structure to a reference protein. It is not intended that homologs necessarily be evolutionarily related. Thus, it is intended that the term encompass the same, similar, or corresponding enzyme(s) (*i.e.*, in terms of structure and function) obtained from different organisms. In some embodiments, it is desirable to identify a homolog that has a quaternary, tertiary and/or primary structure similar to the reference protein. In some embodiments, homologous proteins induce similar immunological response(s) as a reference protein. In some embodiments, homologous proteins are engineered to produce enzymes with desired activity(ies).

[0014] The degree of homology between sequences can be determined using any suitable method known in the art (see, *e.g.*, Smith and Waterman (1981) *Adv. Appl. Math.* 2:482; Needleman and Wunsch (1970) *J. Mol. Biol.*, 48:443; Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85:2444; programs such as GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package (Genetics Computer Group, Madison, WI); and Devereux et al. (1984) *Nucleic Acids Res.* 12:387-95).

[0015] For example, PILEUP is a useful program to determine sequence homology levels. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pair-wise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng and Doolittle, (Feng and Doolittle (1987) *J. Mol. Evol.* 35:351-60). The method is similar to that described by Higgins and Sharp ((1989) *CABIOS* 5:151-53). Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps. Another example of a useful algorithm is the BLAST algorithm, described by Altschul et al. ((1990) *J. Mol. Biol.* 215:403-10) and Karlin et al. ((1993) *Proc. Natl. Acad. Sci. USA* 90:5873-87). One particularly useful BLAST program is the WU-BLAST-2 program (see, *e.g.*, Altschul et al. (1996) *Meth. Enzymol.* 266:460-80). Parameters "W," "T," and "X" determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word-length (W) of 11, the BLOSUM62 scoring matrix (see, *e.g.*, Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M⁵, N⁻⁴, and a comparison of both strands.

[0016] As used herein, the phrases "substantially similar" and "substantially identical," in the context of at least two nucleic acids or polypeptides, typically means that a polynucleotide or polypeptide comprises a sequence that has at least about 70% identity, at least about 75% identity, at least about 80% identity, at least about 85% identity, at least about 90% identity, at least about 91% identity, at least about 92% identity, at least about 93% identity, at least about 94% identity, at least about 95% identity, at least about 96% identity, at least about 97% identity, at least about 98% identity, or even at least about 99% identity, or more, compared to the reference (*i.e.*, wild-type) sequence. Percent sequence identity is calculated using CLUSTAL W algorithm with default parameters. See Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680. Default parameters for the CLUSTAL W algorithm are:

Gap opening penalty:	10.0
----------------------	------

Gap extension penalty:	0.05
Protein weight matrix:	BLOSUM series
DNA weight matrix:	ILTB
Delay divergent sequences %:	40
Gap separation distance:	8
DNA transitions weight:	0.50
List hydrophilic residues:	GPSNDQEKR
Use negative matrix:	OFF
Toggle Residue specific penalties:	ON
Toggle hydrophilic penalties:	ON
Toggle end gap separation penalty	OFF

[0017] Another indication that two polypeptides are substantially identical is that the first polypeptide is immunologically cross-reactive with the second polypeptide. Typically, polypeptides that differ by conservative amino acid substitutions are immunologically cross-reactive. Thus, a polypeptide is substantially identical to a second polypeptide, for example, where the two peptides differ only by a conservative substitution. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions (e.g., within a range of medium to high stringency).

[0018] As used herein, the term "gene" is synonymous with the term "allele" in referring to a nucleic acid that encodes and directs the expression of a protein or RNA. Vegetative forms of filamentous fungi are generally haploid, therefore a single copy of a specified gene (*i.e.*, a single allele) is sufficient to confer a specified phenotype. The term "allele" is generally preferred when an organism contains more than one similar genes, in which case each different similar gene is referred to as a distinct "allele."

[0019] As used herein, "constitutive" expression refers to the production of a polypeptide encoded by a particular gene under essentially all typical growth conditions, as opposed to "conditional" expression, which requires the presence of a particular substrate, temperature, or the like to induce or activate expression.

[0020] As used herein, the term "expressing a polypeptide" and similar terms refers to the cellular process of producing a polypeptide using the translation machinery (e.g., ribosomes) of the cell.

[0021] As used herein, "over-expressing a polypeptide," "increasing the expression of a polypeptide," and similar terms, refer to expressing a polypeptide at higher-than-normal levels compared to those observed with parental or "wild-type cells that do not include a specified genetic modification.

[0022] As used herein, an "expression cassette" refers to a DNA fragment that includes a promoter, and amino acid coding region and a terminator (*i.e.*, promoter: :amino acid coding region: :terminator) and other nucleic acid sequence needed to allow the encoded polypeptide to be produced in a cell. Expression cassettes can be exogenous (*i.e.*, introduced into a cell) or endogenous (*i.e.*, extant in a cell).

[0023] As used herein, the terms "fused" and "fusion" with respect to two DNA fragments, such as a promoter and the coding region of a polypeptide refer to a physical linkage causing the two DNA fragments to become a single molecule.

[0024] As used herein, the terms "wild-type" and "native" are used interchangeably and refer to genes, proteins or strains found in nature, or that are not intentionally modified for the advantage of the presently described yeast.

[0025] As used herein, the term "protein of interest" refers to a polypeptide that is desired to be expressed in modified yeast. Such a protein can be an enzyme, a substrate-binding protein, a surface-active protein, a structural protein, a selectable marker, a signal transducer, a receptor, a transporter, a transcription factor, a translation factor, a co-factor, or the like, and can be expressed. The protein of interest is encoded by an endogenous gene or a heterologous gene (*i.e.*, gene of interest") relative to the parental strain. The protein of interest can be expressed intracellularly or as a secreted protein.

[0026] As used herein, "disruption of a gene" refers broadly to any genetic or chemical manipulation, *i.e.*, mutation, that substantially prevents a cell from producing a function gene product, *e.g.*, a protein, in a host cell. Exemplary methods of disruption include complete or partial deletion of any portion of a gene, including a polypeptide-coding sequence, a promoter, an enhancer, or another regulatory element, or mutagenesis of the same, where mutagenesis encompasses substitutions, insertions, deletions, inversions, and combinations and variations, thereof, any of which mutations substantially prevent the production of a function gene product. A gene can also be disrupted using CRISPR, RNAi, antisense, or any other method that abolishes gene expression. A gene can be disrupted by deletion or genetic manipulation of non-adjacent control elements. As used herein, "deletion of a gene," refers to its removal from the genome of a host cell. Where a gene includes control elements (*e.g.*, enhancer elements) that are not located immediately adjacent to the coding sequence of a gene, deletion of a gene refers to the deletion of the coding sequence, and optionally adjacent enhancer elements, including but not limited to, for example, promoter and/or terminator sequences, but does not require the deletion of non-adjacent control elements. Deletion of a gene also refers to the deletion a part of the coding sequence, or a part of promoter immediately or not immediately adjacent to the coding sequence, where there is no functional activity of the interested gene existed in the engineered cell.

[0027] As used herein, the terms "genetic manipulation" and "genetic alteration" are used interchangeably and refer to the alteration/change of a nucleic acid sequence. The alteration can include but is not limited to a substitution, deletion, insertion or chemical modification of at

least one nucleic acid in the nucleic acid sequence.

[0028] As used herein, a "functional polypeptide/protein" is a protein that possesses an activity, such as an enzymatic activity, a binding activity, a surface-active property, a signal transducer, a receptor, a transporter, a transcription factor, a translation factor, a co-factor, or the like, and which has not been mutagenized, truncated, or otherwise modified to abolish or reduce that activity. Functional polypeptides can be thermostable or thermolabile, as specified.

[0029] As used herein, "a functional gene" is a gene capable of being used by cellular components to produce an active gene product, typically a protein. Functional genes are the antithesis of disrupted genes, which are modified such that they cannot be used by cellular components to produce an active gene product, or have a reduced ability to be used by cellular components to produce an active gene product.

[0030] As used herein, yeast cells have been "modified to prevent the production of a specified protein" if they have been genetically or chemically altered to prevent the production of a functional protein/polypeptide that exhibits an activity characteristic of the wild-type protein. Such modifications include, but are not limited to, deletion or disruption of the gene encoding the protein (as described, herein), modification of the gene such that the encoded polypeptide lacks the aforementioned activity, modification of the gene to affect post-translational processing or stability, and combinations, thereof.

[0031] As used herein, "attenuation of a pathway" or "attenuation of the flux through a pathway," *i.e.*, a biochemical pathway, refers broadly to any genetic or chemical manipulation that reduces or completely stops the flux of biochemical substrates or intermediates through a metabolic pathway. Attenuation of a pathway may be achieved by a variety of well-known methods. Such methods include but are not limited to: complete or partial deletion of one or more genes, replacing wild-type alleles of these genes with mutant forms encoding enzymes with reduced catalytic activity or increased K_m values, modifying the promoters or other regulatory elements that control the expression of one or more genes, engineering the enzymes or the mRNA encoding these enzymes for a decreased stability, misdirecting enzymes to cellular compartments where they are less likely to interact with substrate and intermediates, the use of interfering RNA, and the like.

[0032] As used herein, "aerobic fermentation" refers to growth in the presence of oxygen.

[0033] As used herein, "anaerobic fermentation" refers to growth in the absence of oxygen.

[0034] As used herein, the expression "end of fermentation" refers to the stage of fermentation when the economic advantage of continuing fermentation to produce a small amount of additional alcohol is exceeded by the cost of continuing fermentation in terms of fixed and variable costs. In a more general sense, "end of fermentation" refers to the point where a fermentation will no longer produce a significant amount of additional alcohol, *i.e.*, no more than about 1% additional alcohol, or no more substrate left for further alcohol production.

[0035] As used herein, the expression "carbon flux" refers to the rate of turnover of carbon molecules through a metabolic pathway. Carbon flux is regulated by enzymes involved in metabolic pathways, such as the pathway for glucose metabolism and the pathway for maltose metabolism.

[0036] As used herein, the singular articles "a," "an" and "the" encompass the plural referents unless the context clearly dictates otherwise. The following abbreviations/acronyms have the following meanings unless otherwise specified:

EC

enzyme commission

PKL

phosphoketolase

PTA

phosphotransacetylase

AADH

acetaldehyde dehydrogenases

ADH

alcohol dehydrogenase

EtOH

ethanol

AA

α -amylase

GA

glucoamylase

$^{\circ}$ C

degrees Centigrade

bp

base pairs

DNA

deoxyribonucleic acid

ds or DS

dry solids

g or gm

gram

g/L

grams per liter

H₂O

water

HPLC

high performance liquid chromatography

hr or h

hour

kg	kilogram
M	molar
mg	milligram
mL or ml	milliliter
min	minute
mM	millimolar
N	normal
nm	nanometer
PAB1	polyadenylate-binding protein
PCR	polymerase chain reaction
ppm	parts per million
Δ	relating to a deletion
μ g	microgram
μ L and μ l	microliter
μ M	micromolar

II. Modified yeast cells having increased PAB1 expression

[0037] Described are modified yeast and methods having a genetic alteration that results in the production of increased amounts of PAB1 polypeptides compared to corresponding (*i.e.*, otherwise-identical) parental cells. PAB1 is the major poly(A)-binding protein in yeast. It is an approximately 577- amino acid residue, multifunctional-protein that mediates many cellular functions associated with the 3'-poly(A)-tail of messenger RNAs (see, e.g., Brune, C. et al. (2005) RNA. 11:517-531). Over-expression of PAB1 increased stress tolerance of engineered yeast (Martani, F. et al., (2015), Sci. Rep. 5: 18318; doi: 10.1038/srep18318). No association has heretofore been made between PAB1 expression and acetate reduction in engineered

yeast.

[0038] Applicants have discovered that yeast cells over-expressing PAB1 polypeptides produce a decreased amount of acetate compared to otherwise-identical parental cells. Decreased acetate is desirable as acetate adversely affects yeast growth and fermentation and additionally results in backset that has a lower than desirable pH, requiring pH adjustment or the use of more fresh water to dilute the backset.

[0039] In some embodiments, the increase in the amount of PAB1 polypeptides produced by the modified cells is an increase of at least 20%, at least 30%, at least 40%, at least 50%, at least 70%, at least 100%, at least 150%, at least 200%, at least 500%, at least 1000%, or more, especially at later stage of fermentation, compared to the amount of PAB1 polypeptides produced by parental cells grown under the same conditions.

[0040] In some embodiments, the increase in the amount of PAB1 polypeptides produced by the modified cells is at least 2-fold, at least 5-fold, at least 10-fold, at least 20-fold, at least 30-fold or more, compared to the amount of PAB1 polypeptides produced by parental cells grown under the same conditions.

[0041] In some embodiments, the increase in the strength of the promoter used to control expression of the PAB1 polypeptides produced by the modified cells is at least 2-fold, at least 5-fold, at least 10-fold, at least 13-fold, at least 20-fold, at least 30-fold, or more, compared to strength of the native promoter controlling PAB1 expression, based on the amount of mRNA produced. RNAseq data (see, e.g., Wang, Z. et al. (2009) Nature Rev. Gen. 10:57-63) indicates that the EFB1 promoter used for the exemplified PAB1 expression cassette is approximately 13-times stronger than the normal PAB1 promoter.

[0042] In some embodiments, the decrease in acetate production by the modified cells is a decrease of at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, or at least 50%, more, compared to the amount of acetate produced by parental cells grown under the same conditions.

[0043] Preferably, increased PAB1 expression is achieved by genetic manipulation using sequence-specific molecular biology techniques, as opposed to chemical mutagenesis, which is generally not targeted to specific nucleic acid sequences. However, chemical mutagenesis is not excluded as a method for making modified yeast cells.

[0044] In some embodiments, the present compositions and methods involve introducing into yeast cells a nucleic acid capable of directing the over-expression, or increased expression, of a PAB1 polypeptide. Particular methods include but are not limited to (i) introducing an exogenous expression cassette for producing the polypeptide into a host cell, optionally in addition to an endogenous expression cassette, (ii) substituting an exogenous expression cassette with an endogenous cassette that allows the production of an increased amount of the polypeptide, (iii) modifying the promoter of an endogenous expression cassette to increase

expression, (iv) increase copy number of the same or different cassettes for over-expression of PAB1, and/or (v) modifying any aspect of the host cell to increase the half-life of the polypeptide in the host cell.

[0045] In some embodiments, the parental cell that is modified already includes a gene of interest, such as a gene encoding a selectable marker, carbohydrate-processing enzyme, or other polypeptide. In some embodiments, a gene of introduced is subsequently introduced into the modified cells.

[0046] In some embodiments, the parental cell that is modified already includes an engineered pathway of interest, such as a PKL pathway to increases ethanol production, or any other pathway to increase alcohol production.

[0047] The amino acid sequence of the exemplified *S. cerevisiae* PAB1 polypeptide is shown, below, as SEQ ID NO: 1:

```
MADITDKTAEQLENLNIQDDQKQAATGSESQSVENSSASLYVGDLEPSVSEAHLYDIFSPIG
SVSSIRVCRDAITKTSLGYAYVNFNDHEAGRKAIEQLNYTPIKGRLCRIMWSQRDPSLRKKG

SGNIFIKNLHPDIDNKALYDTFSVFGDILSSKIATDENGKSKGFGFVHFEEEGAAKEAIDAL
NGMLLNGQEIIYVAPHLRSRKERDSQLEETKAHYTNLYVKNINSETTDEQFQELFAKFGPIVSA
SLEKDADGKLGKGFVNYEKHEDAVKAVEALNDSELNGEKLYVGRAQKKNERMHVLKQYEA
YRLEKMAKYQGVNLFVKNLDDSDVDEKLEEEFAPYGTITSAKVMRTENGKSKGFGFVCFSTP
EEATKAITEKNQQIVAGKPLYVAIAQRKDVRRSOLAQQIQARNQMRYYQATAAAAAAAGMP
GQFMPPMFYGVMPPRGVPFNGPNPQQMNPMGGMPKNGMPPQFRNGPVYGVPPQGGFPRNAND
NNQFYQQQRQALGEQLYKKVSAKTSNEEAAGKITGMILDLPQEVFPLESEDELFEQHYKE
ASAAYESFKKEQEQQTEQA
```

[0048] The NCBI database includes over 100 entries for *S. cerevisiae* PAB 1 polypeptides. Natural variations in the amino acid sequence are not expected to affect its function. In addition, based on such BLAST and Clustal W data, it is apparent that the exemplified *S. cerevisiae* PAB1 polypeptide shares a high degree of sequence identity to polypeptides from other organisms, and over-expression of functionally and/or structurally similar proteins, homologous proteins and/or substantially similar or identical proteins, is expected to produce similar beneficial results.

[0049] In particular embodiments of the present compositions and methods, the amino acid sequence of the PAB 1 polypeptide that is over-expressed in modified yeast cells has at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or even at least about 99% identity, to SEQ ID NO: 1.

III. Modified yeast cells having increased PAB1 expression in combination with genes

of an exogenous PKL pathway

[0050] Increased expression of PAB 1 can be combined with expression of genes in the PKL pathway to reduce the production of elevated amounts of acetate that is associated with introducing an exogenous PKL pathway into yeast.

[0051] Engineered yeast cells having a heterologous PKL pathway have been previously described in WO2015148272 (Miasnikov et al.). These cells express heterologous phosphoketolase (PKL), phosphotransacetylase (PTA) and acetylating acetyl dehydrogenase (AADH), optionally with other enzymes, to channel carbon flux away from the glycerol pathway and toward the synthesis of acetyl-CoA, which is then converted to ethanol. Such modified cells are capable of increased ethanol production in a fermentation process when compared to otherwise-identical parent yeast cells.

IV. Combination of increased active PAB1 production with other mutations that affect alcohol production

[0052] In some embodiments, in addition to expressing increased amounts of active PAB1 polypeptides, optionally in combination with introducing an exogenous PKL pathway, the present modified yeast cells include additional modifications that affect ethanol production.

[0053] The modified cells may further include mutations that result in attenuation of the native glycerol biosynthesis pathway and/or reuse glycerol pathway, which are known to increase alcohol production. Methods for attenuation of the glycerol biosynthesis pathway in yeast are known and include reduction or elimination of endogenous NAD-dependent glycerol 3-phosphate dehydrogenase (GPD) or glycerol phosphate phosphatase activity (GPP), for example by disruption of one or more of the genes *GPD1*, *GPD2*, *GPP1* and/or *GPP2*. See, e.g., U.S. Patent Nos. 9,175,270 (Elke et al.), 8,795,998 (Pronk et al.) and 8,956,851 (Argyros et al.). Methods to enhance the reuse glycerol pathway by over expression of glycerol dehydrogenase (*GCY1*) and dihydroxyacetone kinase (*DAK1*) to convert glycerol to dihydroxyacetone phosphate (Zhang et al; J Ind Microbiol Biotechnol (2013) 40:1153-1160).

[0054] The modified yeast may further feature increased acetyl-CoA synthase (also referred to acetyl-CoA ligase) activity (EC 6.2.1.1) to scavenge (*i.e.*, capture) acetate produced by chemical or enzymatic hydrolysis of acetyl-phosphate (or present in the culture medium of the yeast for any other reason) and converts it to Ac-CoA. This partially reduces the undesirable effect of acetate on the growth of yeast cells and may further contribute to an improvement in alcohol yield. Increasing acetyl-CoA synthase activity may be accomplished by introducing a heterologous acetyl-CoA synthase gene into cells, increasing the expression of an endogenous acetyl-CoA synthase gene and the like.

[0055] In some embodiments the modified cells may further include a heterologous gene encoding a protein with NAD⁺-dependent acetylating acetaldehyde dehydrogenase activity and/or a heterologous gene encoding a pyruvate-formate lyase. The introduction of such genes in combination with attenuation of the glycerol pathway is described, e.g., in U.S. Patent No. 8,795,998 (Pronk et al.). In some embodiments of the present compositions and methods the yeast expressly lacks a heterologous gene(s) encoding an acetylating acetaldehyde dehydrogenase, a pyruvate-formate lyase or both.

[0056] In some embodiments, the present modified yeast cells may further over-express a sugar transporter-like (STL1) polypeptide to increase the uptake of glycerol (see, e.g., Ferreira *et al.*, 2005; Dušková *et al.*, 2015 and WO 2015023989 A1).

[0057] In some embodiments, the present modified yeast cells further include a butanol biosynthetic pathway. In some embodiments, the butanol biosynthetic pathway is an isobutanol biosynthetic pathway. In some embodiments, the isobutanol biosynthetic pathway comprises a polynucleotide encoding a polypeptide that catalyzes a substrate to product conversion selected from the group consisting of: (a) pyruvate to acetolactate; (b) acetolactate to 2,3-dihydroxyisovalerate; (c) 2,3-dihydroxyisovalerate to 2-ketoisovalerate; (d) 2-ketoisovalerate to isobutyraldehyde; and (e) isobutyraldehyde to isobutanol. In some embodiments, the isobutanol biosynthetic pathway comprises polynucleotides encoding polypeptides having acetolactate synthase, keto acid reductoisomerase, dihydroxy acid dehydratase, ketoisovalerate decarboxylase, and alcohol dehydrogenase activity.

[0058] In some embodiments, the modified yeast cells comprising a butanol biosynthetic pathway further comprise a modification in a polynucleotide encoding a polypeptide having pyruvate decarboxylase activity. In some embodiments, the yeast cells comprise a deletion, mutation, and/or substitution in an endogenous polynucleotide encoding a polypeptide having pyruvate decarboxylase activity. In some embodiments, the polypeptide having pyruvate decarboxylase activity is selected from the group consisting of: PDC1, PDC5, PDC6, and combinations thereof. In some embodiments, the yeast cells further comprise a deletion, mutation, and/or substitution in one or more endogenous polynucleotides encoding FRA2, ALD6, ADH1, GPD2, BDH1, and YMR226C.

V. Combination of increased expression PAB1 with other beneficial mutations

[0059] In some embodiments, in addition to increased expression of PAB 1 polypeptides, optionally in combination with other genetic modifications that benefit alcohol production, the present modified yeast cells further include any number of additional genes of interest encoding proteins of interest. Additional genes of interest may be introduced before, during, or after genetic manipulations that result in the increased production of active HAC1 polypeptides. Proteins of interest, include selectable markers, carbohydrate-processing enzymes, and other commercially-relevant polypeptides, including but not limited to an enzyme selected from the

group consisting of a dehydrogenase, a transketolase, a phosphoketolase, a transaldolase, an epimerase, a phytase, a xylanase, a β -glucanase, a phosphatase, a protease, an α -amylase, a β -amylase, a glucoamylase, a pullulanase, an isoamylase, a cellulase, a trehalase, a lipase, a pectinase, a polyesterase, a cutinase, an oxidase, a transferase, a reductase, a hemicellulase, a mannanase, an esterase, an isomerase, a pectinases, a lactase, a peroxidase and a laccase. Proteins of interest may be secreted, glycosylated, and otherwise-modified.

VI. Use of the modified yeast for increased alcohol production

[0060] The present compositions and methods include methods for increasing alcohol production and/or reducing glycerol production, in fermentation reactions. Such methods are not limited to a particular fermentation process. The present engineered yeast is expected to be a "drop-in" replacement for convention yeast in any alcohol fermentation facility. While primarily intended for fuel alcohol production, the present yeast can also be used for the production of potable alcohol, including wine and beer.

VII. Yeast cells suitable for modification

[0061] Yeasts are unicellular eukaryotic microorganisms classified as members of the fungus kingdom and include organisms from the phyla Ascomycota and Basidiomycota. Yeast that can be used for alcohol production include, but are not limited to, *Saccharomyces* spp., including *S. cerevisiae*, as well as *Kluyveromyces*, *Lachancea* and *Schizosaccharomyces* spp. Numerous yeast strains are commercially available, many of which have been selected or genetically engineered for desired characteristics, such as high alcohol production, rapid growth rate, and the like. Some yeasts have been genetically engineered to produce heterologous enzymes, such as glucoamylase or α -amylase.

VII. Substrates and products

[0062] Alcohol production from a number of carbohydrate substrates, including but not limited to corn starch, sugar cane, cassava, and molasses, is well known, as are innumerable variations and improvements to enzymatic and chemical conditions and mechanical processes. The present compositions and methods are believed to be fully compatible with such substrates and conditions.

[0063] Alcohol fermentation products include organic compound having a hydroxyl functional group (-OH) is bound to a carbon atom. Exemplary alcohols include but are not limited to methanol, ethanol, n-propanol, isopropanol, n-butanol, isobutanol, n-pentanol, 2-pentanol, isopentanol, and higher alcohols. The most commonly made fuel alcohols are ethanol, and butanol.

[0064] These and other aspects and embodiments of the present yeast strains and methods will be apparent to the skilled person in view of the present description. The following examples are intended to further illustrate, but not limit, the compositions and methods.

EXAMPLES

Example 1

Materials and methods

Liquefact preparation:

[0065] Liquefact (corn mash slurry) was prepared by adding 600 ppm of urea, 0.124 SAPU/g ds FERMGEN™ 2.5x (acid fungal protease), 0.33 GAU/g ds CS4 (a variant *Trichoderma* glucoamylase) and 1.46 SSCU/g ds AKAA (*Aspergillus kawachii* α-amylase), adjusted to a pH of 4.8.

Serum vial assays:

[0066] 2 mL of YPD in 24-well plates were inoculated with yeast cells and the cultures allowed to grow overnight to an OD between 25-30. 5 mL liquefact (5.6g) was transferred to serum vials (Chemglass, Catalog No. CG-4904-01) and yeast was added to each vial to a final OD of about 0.3. The lids of the vials were installed and punctured with needle (BD, Catalog No. 305111) for ventilation (to release CO₂), then incubated at 32°C with shaking at 200 RPM for 55 hours.

AnKom assays:

[0067] 300 µL of concentrated yeast overnight culture was added to each of a number ANKOM bottles filled with 50 g prepared liquefact (see above) to a final OD of 0.3. The bottles were then incubated at 32°C with shaking at 150 RPM for 55 hours.

HPLC analysis:

[0068] Samples of the cultures from serum vials and AnKom assays were collected in Eppendorf tubes by centrifugation for 12 minutes at 14,000 RPM. The supernatants were filtered using 0.2 µM PTFE filters and then used for HPLC (Agilent Technologies 1200 series) analysis with the following conditions: Bio-Rad Aminex HPX-87H columns, running temperature of 55C. 0.6 ml/min isocratic flow 0.01 N H₂SO₄, 2.5 µl injection volume. Calibration standards were used for quantification of the of acetate, ethanol, glycerol, glucose and other molecules. All values are reported in g/L.

Example 2

Preparation of a PAB1 expression cassette

[0069] The PAB1 gene of *Saccharomyces cerevisiae* was codon optimized and synthesized to generate the artificial gene, "PAB1s." The EFB1 promoter (YAL003W locus; SEQ ID NO: 3) and TPI terminator (YDR050C locus; SEQ ID NO: 4) were operably linked to the coding sequence to generate EFB1Pro::PAB1s::Tpi1Ter expression cassette.. This expression cassette was introduced at the upstream region of the AAP1 locus (YHR047C) of either (i) FERMAX™ Gold (Martrex Inc., Minnesota, USA; herein abbreviated, "FG"), a well-known fermentation yeast used in the grain ethanol industry, or (ii) FG-PKR, engineered FG yeast having a heterologous phosphoketolase (PKL) pathway involving the expression of phosphoketolase (PKL), phosphotransacetylase (PTA) and acetylating acetyl dehydrogenase (AADH) as described in WO2015148272 (Miasnikov et al.) using CRISPR Cas9 technology. The expected insertion of the PAB1 expression cassette in the two parental strains was confirmed by PCR.

[0070] The amino acid sequence of the PAB1 polypeptide is shown, below, as SEQ ID NO: 1:

```
MADI TDKTAEQLENLNIQDDQKQAATGSESQSVENS SASLYVGDLEPSVSEAHLYDIFSPIG
SVSSIRVCRDAITKTS LGYAYVNFNDHEAGRKAIEQLNYTPIKGR LCRIMWSQRDPSLRKKG
SGNIFIKNLHPDIDNKALYDTFSVFGDILSSKIATDENGKSKGFGFVHFEEEGAAKEAIDAL
NGMLLNGQEIIYVAPHL SRKERDSQLEETKAHYTNLYVKINSETTDEQFQELFAKFGPIVSA
SLEKDADGKLGKGFVNYEKHEDAVKAVEALNDSELNGEKLYVGRAQKKNERMHVLKKQYEA
YRLEKMAKYQG VNLVFNLDSDVDEKLEEEFAPYGTITSAKVMRTENGKSKGFGFVCFSTP
EEATKAITEKNQQIVAGKPLYVAIAQRKDVRRS QLAQQIQARNQMRYQQATAAAAAAAGMP
GQFMPPMFYGVMP RGVPEFNGPNPQQMNPMGGMPKNGMPPQFRNGPVYGVPPQGGFPRNAND
NNQFYQQQRQALGEQLYKKVSAKTSNEEAAGKITGMILDLPPQEVEFPLLESDDELFEQHYKE
ASAAYESFKKEQEQQTEQA
```

[0071] The PAB1-coding region of the PAB1s gene is shown, below, as SEQ ID NO: 2:

```
ATGGCTGACATCACCGACAAGACTGCCGAACAATTGGAAAACCTGAACATTCAAGACGATCA
GAAGCAAGCTGCCACTGGTTC CGAATCTCAATCCGTCGAAAACCTCTTCGGCTTCTTTGTACC
```

TTGGTGACTTAGAACCATCTGTCTCCGAAGCTCACTTGTACGACATCTTCTCTCCAATCGGT
 TCCGTCTCTTCCATCAGAGTTTGTCTGTGATGCTATCACCAAGACCAGCTTGGGTTACGCTTA
 CGTCAACTTTAACGACCATGAAGCTGGCAGAAAGGCTATCGAACAAATTGAACTACACTCCAA
 TCAAGGGTAGACTATGTAGAATCATGTGGTCTCAACGTGATCCATCCTTGAGAAAGAAAGGT
 TCTGGCAACATCTTCATCAAGAACTTGCATCCAGACATTGACAACAAGGCTTTGTACGATAC
 TTTCAAGTGTCTTTGGTGACATCTTATCTTCCAAGATTGCTACCGACGAAAACGGAAAGTCCA
 AGGGTTTTGGCTTTGTTCACTTCGAAGAGGAAGGTGCTGCCAAGGAAGCTATCGATGCTTTG
 AATGGTATGCTATTGAACGGTCAAGAAATCTACGTTGCTCCTCACTTGTCCAGAAAGGAACG
 TGACTCTCAATTGGAAGAGACCAAGGCTCACTACACCAACTTGTACGTCAAGAATATCAACT
 CCGAAACTACCGACGAACAATTTCAAGAATTGTTTGCCAAGTTTGGTCCAATTGTTTCTGCT
 TCCTTGAAAAGGATGCCGACGGCAAGTTGAAGGGTTTCGGTTTCGTCAACTACGAAAAGCA
 CGAAGACGCTGTCAAGGCTGTGGAAGCCTTGAACGATTCCGAACTAAATGGCGAAAAGTTGT
 ACGTTGGTAGAGCACAGAAAAAGAACGAACGTATGCACGCTCTGAAGAAACAATACGAAGCT
 TACAGATTGGAAGAGATGGCCAAGTACCAAGGTGTTAATTTGTTTCGTCAAGAACTTGGACGA
 TTCTGTGATGACGAAAAGTTGGAAGAGGAATTTGCTCCATACGGTACTATCACCTCTGCCA
 AGGTCATGAGAACCGAAAACGGTAAGTCCAAGGGTTTTGGCTTCGTCTGTTTCTCTACTCCA
 GAGGAAGCTACCAAGGCTATCACCGAAAAGAATCAACAGATTGTTGCTGGCAAGCCATTGTA
 CGTTGCCATTGCTCAAAGAAAGGACGTCAGACGTTCTCAACTAGCTCAACAGATCCAAGCCA
 GAAACCAAATGAGATACCAACAGGCTACTGCTGCCGCTGCAGCTGCCGCTGCAGGTATGCCA
 GGTCAATTCATGCCTCCAATGTTTTACGGTGTCTATGCCTCCAAGAGGTGTTCCATTCAACGG
 TCCAAACCCTCAACAGATGAATCCAATGGGTGGAATGCCAAAGAACGGTATGCCTCCACAAT
 TCAGAAACGGTCCAGTCTACGGTGTCCCTCCACAAGGTGGCTTTCCAAGAAACGCTAACGCAC
 AACAAATCAATTCTACCAACAGAAGCAAAGACAAGCTTTGGGTGAACAATTGTACAAGAAAGT
 TTCTGCCAAGACTTCCAACGAAGAAGCTGCAGGCAAGATCACTGGTATGATCTTGGATTTAC
 CACCTCAAGAAGTCTTCCCCTATTGGAATCCGACGAATTGTTGGAACAACACTACAAGGAG
 GCTTCTGCTGCCTACGAATCTTTCAAAAAGGAACAAGAGCAACAGACTGAACAAGCCTAA

[0072] The EFB1 promoter region used for PAB1s over-expression shown, below, as SEQ ID NO: 3:

TATGTTGTACCTAAATCAATACCGACAGCTTTTGACATATTATCTGTTATTTACTTGAATTT
 TTGTTTCTTGTAATACTTGATTACTTTTCTTTTGATGTGCTTATCTTACAAATAGAGAAAAT
 AAAACAACCTTAAGTAAGAAATTGGGAAACGAACTACAACCTCAATCCCTTCTCGAAGATACAT
 CAATCCACCCCTTATATAACCTTGAAGTCTCGAAACGATCAGCTAATCTAAATGGCCCCC
 TTCTTTTTGGGTTCTTCTCTCCCTCTTGCCGCGGATGGAACGTTCTGGAAAAGAAGAATA
 ATTTAATTACTTTCTCAACTAAAATCTGGAGAAAAACGCAAATGACAGCTTCTAAACGTTT
 CGTGTGCTTTCTTCTAGAAATGTTCTGGAAAGTTTACAACAATCCACAAGAACGAAAATGCC
 GTTGACAATGATGAAACCATCATCCACACACCGCGCACACGTTTATTTCTTTTTCTGAA
 TTTTTTTTTTCCGCCATTTTCAACCAAGGAAATTTTTTTTTCTTAGGGCTCAGAACCTGCAGG

TGAAGAAGCGCTTTAGAAATCAAAGCACACGCAACAATTTGTCGACAACCGAGCCTTTGAA
 GAAAAAATTTTTTACATTGTCGCCTCTAAATAAATAGTTTTAAGGTTATCTACCCACTATATT
 TAGTTGGTTCTTTTTTTTTTTCCTTCTACTCTTTATCTTTTTTACCTCATGCTTTCTACCTTTC
 AGCACTGAAGAGTCCAACCGAATATATACACACATA

[0073] The Tpi1 terminator region used for PAB1s over-expression shown, below, as SEQ ID NO: 4:

ATTAATATAATTATATAAAAAATATTATCTTCTTTTTCTTTATATCTAGTGTTATGTAA
 AATAAATTGATGACTACGGAAAGCTTTTTTATATTGTTTCTTTTTTCATTCTGAGCCACTTAA
 ATTTTCGTGAATGTTCTTGTAAGGGACGGTAGATTTACAAGTGATACAACAAAAAGCAAGGCG
 CTTTTTCTAATAAAAAAGAAGAAAAGCATTAAACAATTGAACACCTCTATATCAACGAAGAAT
 ATTACTTTGTCTCTAAATCCTTGTAATAATGTGTACGATCTCTATATGGGTTACTCATAAGTG
 TACCGAAGACTGCATTGAAAGTTTATGTTTTTTCCTGAGGCGTCATTTTCGCGTTGAGAA
 GATGTTCTTATCCAAATTTCAA

Example 3

Alcohol production using yeast that over-express PAB1

[0074] Strains over-expressing PAB1 were tested in a small vial assay, containing 5.6 g liquefact, and then in an Ankom assay, containing 50 g liquefact, as described in Example 1. Fermentations were performed at 32°C for 55 hours. Samples from the end of fermentation were analyzed by HPLC. The results are summarized in Tables 1 and 2.

Table 1. HPLC results from a small vial assays

Strain features	Glycerol (g/L)	Acetate (g/L)	Glucose (g/L)	Ethanol (g/L)	Acetate reduction (%)
FG	13.767	1.275	0.291	143.839	-0-
FG-PAB1	14.707	1.063	0.173	143.039	16.6%
FG-PKL	12.811	1.744	0.583	145.101	-0-
FG-PKL-PAB1	13.554	1.308	0.661	144.524	25.0%

Table 2. HPLC results from an Ankom assay

Strain features	Glucose (g/L)	Glycerol (g/L)	Acetate (g/L)	Ethanol (g/L)	Acetate reduction (%)
FG	0.32	13.15	0.90	140.20	-0-
FG-PAB1	0.31	13.91	0.79	140.11	12.2%
FG-PKL	0.61	11.94	1.42	141.45	-0-

Strain features	Glucose (g/L)	Glycerol (g/L)	Acetate (g/L)	Ethanol g/L	Acetate reduction (%)
FG-PKL-PAB1	0.71	12.69	1.06	140.33	25.4%

[0075] Over-expression of PAB 1 resulted in a decrease in acetate production of about 12-17% in FG yeast, which is recognized as a robust, high-ethanol-producing yeast for the fuel ethanol industry, while not being a genetically engineered organism. Over-expression of PAB 1 resulted in a decrease of up to almost twice as much in FG yeast engineered to have an exogenous PKL pathway. These results demonstrate that PAB 1 over-expression is beneficial for reducing acetate, in general, but is particularly beneficial for reducing the increased amount of acetate produced by yeast having a PKL pathway.

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2015148272A \[0003\] \[0051\] \[0069\]](#)
- [US2014273144A1 \[0004\]](#)
- [US9175270B \[0053\]](#)
- [US8795998B \[0053\] \[0055\]](#)
- [US8956851B \[0053\]](#)
- [WO2015023989A1 \[0056\]](#)

Non-patent literature cited in the description

- GOMBERT, A.K.VAN MARIS. A.J.Curr. Opin. Biotechnol., 2015, vol. 33, 81-86 [0002]
- SMITHWATERMAN Adv. Appl. Math., 1981, vol. 2, 482- [0014]
- NEEDLEMAN WUNSCH J. Mol. Biol., 1970, vol. 48, 443- [0014]
- PEARSON LIPMAN Proc. Natl. Acad. Sci. USA, 1988, vol. 85, 2444- [0014]
- DEVEREUX et al. Nucleic Acids Res., 1984, vol. 12, 387-95 [0014]
- FENG DOOLITTLE J. Mol. Evol., 1987, vol. 35, 351-60 [0015]
- HIGGINS SHARP CABIOS, 1989, vol. 5, 151-53 [0015]
- ALTSCHUL et al. J. Mol. Biol., 1990, vol. 215, 403-10 [0015]
- KARLIN et al. Proc. Natl. Acad. Sci. USA, 1993, vol. 90, 5873-87 [0015]
- ALTSCHUL et al. Meth. Enzymol., 1996, vol. 266, 460-80 [0015]
- HENIKOFF HENIKOFF Proc. Natl. Acad. Sci. USA, 1989, vol. 89, 10915- [0015]
- THOMPSON et al. Nucleic Acids Res., 1994, vol. 22, 4673-4680 [0016]
- BRUNE, C. et al. RNA., 2005, vol. 11, 517-531 [0037]
- MARTANI, F. et al. Sci. Rep., 2015, vol. 5, 18318- [0037]
- WANG, Z. et al. Nature Rev. Gen., 2009, vol. 10, 57-63 [0041]
- ZHANG et al. J Ind Microbiol Biotechnol, 2013, vol. 40, 1153-1160 [0053]

Patentkrav

1. Modificerede gærceller afledt fra modergærceller, hvilke modificerede celler omfatter en genetisk ændring, der forårsager, at de modificerede celler producerer en forøget mængde af PAB1-polypeptider sammenlignet med modercellerne, hvor de modificerede celler under fermentering producerer en reduceret mængde acetat sammenlignet med mængden af acetat, der produceres af modercellerne under identiske fermenteringsbetingelser, og hvor de modificerede celler yderligere omfatter et eller flere heterologe gener i phosphoketolase-reaktionsvejen, hvor den genetiske ændring omfatter indførelse i modercellerne af en nukleinsyre, der er i stand til at styre ekspressionen af et PAB1-polypeptid til et niveau over niveauet for modercellen, der dyrkes under tilsvarende betingelser.

2. Modificerede celler ifølge krav 1, hvor den genetiske ændring omfatter:

(a) indførelse af en exogen ekspressionkassette til produktion af PAB1-polypeptidet i værtscellen;

(b) modificering af promotoren for en endogen ekspressionskassette til forøgelse af ekspression af PAB1-polypeptidet; eller

(c) forøgelse af kopiantallet i den samme eller forskellige kassetter til overekspression af PAB1.

3. Modificerede celler ifølge krav 1, hvor den genetiske ændring omfatter indførelsen af en ekspressionskassette til ekspression af et PAB1-polypeptid.

4. Modificerede celler ifølge krav 1, hvor generne i phosphoketolase-reaktionsvejen er valgt fra gruppen, der består af phosphoketolase, phosphotransacetylase og acetylerende acetyldehydrogenase.

5. Modificerede celler ifølge et hvilket som helst af kravene 1-4, hvor størrelsen af stigningen i ekspression af PAB1-

polypeptidet er mindst ca. 200 % sammenlignet med niveauet af ekspresion i modercellerne, der dyrkes under tilsvarende betingelser.

5 6. Modificerede celler ifølge et hvilket som helst af kravene 1-4, hvor størrelsen af stigningen i produktionen af mRNA, der koder for PAB1-polypeptidet, er mindst ca. 400 % sammenlignet med niveauet af ekspresion i modercellerne, der dyrkes under tilsvarende betingelser.

10

7. Modificerede celler ifølge et hvilket som helst af kravene 1-6, hvor cellerne yderligere omfatter et exogent gen, der koder for et carbohydratprocesserende enzym.

15 8. Modificerede celler ifølge et hvilket som helst af kravene 1-7, som yderligere omfatter en ændring i glycerolreaktionsvejen og/eller acetyl-CoA-reaktionsvejen.

20 9. Modificerede celler ifølge et hvilket som helst af kravene 1-8, som yderligere omfatter en alternativ reaktionsvej til fremstilling af ethanol.

10. Modificerede celler ifølge et hvilket som helst af kravene 1-9, hvor cellerne er fra en *Saccharomyces* spp.

25

11. Fremgangsmåde til reducere af produktionen af acetat fra gærceller, der dyrkes på et carbohydratsubstrat, som omfatter: indførelse i modergærceller af en genetisk ændring, der øger produktionen af PAB1-polypeptider sammenlignet med mængden, der produceres i modercellerne, hvor den genetiske ændring omfatter indførelse i modercellerne af en nukleinsyre, der er i stand til at styre ekspresionen af et PAB1-polypeptid til et niveau over niveauet for modercellen, der dyrkes under tilsvarende betingelser.

35

12. Fremgangsmåde ifølge krav 11, hvor cellerne, der har den indførte genetiske ændring, er modificerede celler ifølge et hvilket som helst af kravene 1-10.

13. Fremgangsmåde ifølge krav 11 eller 12, hvor reduktionen i acetatproduktion er mindst 10 %, mindst 15 %, mindst 20 % eller mindst 25 %.

5

14. Fremgangsmåde ifølge et hvilket som helst af kravene 11-13, hvor PAB1-polyeptider overudtrykkes med mindst 200 %.

15. Fremgangsmåde ifølge et hvilket som helst af kravene 11-10 13, hvor PAB1-polyeptider overudtrykkes mindst 15 gange.

SEKVENSLISTE

Sekvenslisten er udeladt af skriftet og kan hentes fra det Europæiske Patent Register.

The Sequence Listing was omitted from the document and can be downloaded from the European Patent Register.

