




US 20170165362A1

(19) **United States**(12) **Patent Application Publication**
Greene et al.(10) **Pub. No.: US 2017/0165362 A1**(43) **Pub. Date: Jun. 15, 2017**(54) **INNOVATIVE DISCOVERY OF
THERAPEUTIC, DIAGNOSTIC, AND
ANTIBODY COMPOSITIONS RELATED TO
PROTEIN FRAGMENTS OF ARGINYL-TRNA
SYNTHETASES**(71) Applicants: **aTyr Pharma, Inc.**, San Diego, CA
(US); **Pangu BioPharma Limited**,
Hong Kong (CN)(72) Inventors: **Leslie Ann Greene**, San Diego, CA
(US); **Kyle P. Chiang**, Cardiff, CA
(US); **Fei Hong**, San Diego, CA (US);
Alain P. Vasserot, Carlsbad, CA (US);
Wing-Sze Lo, Hong Kong (HK); **Jeffrey
D. Watkins**, Encinitas, CA (US);
Cheryl L. Quinn, Minneapolis, MN
(US); **John D. Mendlein**, Encinitas, CA
(US)(21) Appl. No.: **15/332,495**(22) Filed: **Oct. 24, 2016****Related U.S. Application Data**(63) Continuation of application No. 14/593,620, filed on
Jan. 9, 2015, now Pat. No. 9,593,322, which is acontinuation of application No. 13/696,042, filed on
Feb. 8, 2013, now Pat. No. 8,961,961, filed as appli-
cation No. PCT/US2011/034838 on May 2, 2011.(60) Provisional application No. 61/330,785, filed on May
3, 2010, provisional application No. 61/330,790, filed
on May 3, 2010, provisional application No. 61/330,
778, filed on May 3, 2010.**Publication Classification**(51) **Int. Cl.**
A61K 39/395 (2006.01)
A61K 9/19 (2006.01)
(52) **U.S. Cl.**
CPC **A61K 39/395** (2013.01); **A61K 9/19**
(2013.01); **A61K 2121/00** (2013.01)(57) **ABSTRACT**Provided are compositions comprising newly identified pro-
tein fragments of aminoacyl-tRNA synthetases, polynucle-
otides that encode them and complements thereof, related
agents, and methods of use thereof in diagnostic, drug
discovery, research, and therapeutic applications.




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B



FIG. 3C

**INNOVATIVE DISCOVERY OF
THERAPEUTIC, DIAGNOSTIC, AND
ANTIBODY COMPOSITIONS RELATED TO
PROTEIN FRAGMENTS OF ARGINYL-TRNA
SYNTHETASES**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a Continuation of U.S. application Ser. No. 14/593,620, filed Jan. 9, 2015; which is a Continuation of U.S. application Ser. No. 13/696,042, filed Feb. 8, 2013, now granted U.S. Pat. No. 8,961,961, issued Feb. 24, 2015; which is a U.S. National Phase Application of International Patent Application No. PCT/US2011/034838, filed May 2, 2011, which claims the benefit under 35 U.S.C. §119(e) of U.S. provisional patent application No. 61/330,785 filed on May 3, 2010, U.S. provisional patent application No. 61/330,790 filed on May 3, 2010, and U.S. provisional patent application No. 61/330,778 filed on May 3, 2010, the entire contents of each of which, are incorporated herein by reference.

**STATEMENT REGARDING SEQUENCE
LISTING**

[0002] The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is ATYR_035_03US_ST25.txt. The text file is about 145 KB, was created on Oct. 24, 2016, and is being submitted electronically via EFS-Web.

TECHNICAL FIELD

[0003] The present invention relates generally to compositions comprising newly identified protein fragments of aminoacyl-tRNA synthetases and other proteins, polynucleotides that encode them and complements thereof, related agents, and methods of use thereof in diagnostic, drug discovery, research, and therapeutic applications.

BACKGROUND

[0004] For over four decades, aminoacyl-tRNA synthetases (AARSs) were thought of as essential housekeeping proteins that catalyze the aminoacylation of tRNA molecules as part of the decoding of genetic information during the process of protein translation. AARSs have been extensively studied in this respect, and many of their full-length sequences were cloned for sequence analysis and to provide a rich source of biochemical experimentation. Some fragments of AARSs, and other proteins, however, possess unexpected activities not associated with aminoacylation, including extracellular signaling activities that modulate pathways beyond protein translation. Generally, these unexpected activities are not observed in the context of the full-length or parental protein sequences; instead, they are observed following removal or resection of AARS protein fragments from their parental sequences, or by expressing and sufficiently purifying fragment AARS sequences and then testing for novel, non-synthetase related activities.

[0005] While the full-length sequences of AARS have been known for some time, no systematic experimental analysis has been conducted to elucidate such AARS protein fragments, or protein fragments from related or associated

proteins, or to evaluate the potential role of the full length AARS proteins for novel biological activities outside of the context of amino acid synthesis. In portions of this specification, such AARS protein fragments, AARS domains, or AARS alternative splice variants are referred to herein as “resectins”. In its broadest context, the term “resectin” refers to a portion of a protein which has been excised or restricted (either by means of proteolysis, alternative splicing, mutagenesis, or recombinant genetic engineering) from the context of its native full-length or parental protein sequence, which often otherwise masks its novel biological activities. Likewise, no systematic experimental analysis has been conducted to explore the use of such resectins as biotherapeutic agents, diagnostic agents, or drug targets in the treatment of various medical conditions, or their potential association with human diseases. As essential housekeeping genes with a known function in mammals that is critical to life, AARSs were neither considered as drug targets in mammals, nor were they parsed out by standard genomic sequencing, bioinformatic, or similar efforts to identify resectins having non-synthetase activities. Standard biochemical research efforts have similarly been directed away from characterizing the biological properties of AARS resectins and their potential therapeutic and diagnostic relevance, mainly due to the previously understood role of their corresponding full-length parental AARSs.

BRIEF DESCRIPTION OF THE DRAWINGS

[0006] FIG. 1 shows the domain structure of the Arginyl aminoacyl tRNA synthetase overlaid with the relative positions and sizes of the N-terminal AARS polypeptides identified from mass spectrometry analysis shown schematically.

[0007] FIGS. 2A-2C show the domain structure of the Arginyl aminoacyl tRNA synthetase overlaid with the relative positions and sizes of the C-terminal AARS polypeptides shown schematically. FIG. 2A representing fragments identified from mass spectrometry analysis, FIG. 2B representing the fragments identified from deep sequencing of transcriptomes, and FIG. 2C representing fragments identified from bioinformatics analysis.

[0008] FIGS. 3A-3C show the domain structure of the Arginyl aminoacyl tRNA synthetase overlaid schematically with the relative positions and sizes of the Internal AARS polypeptides shown schematically. FIG. 3A representing fragments identified from mass spectrometry analysis, FIG. 3B representing the fragments identified from deep sequencing of transcriptomes, and FIG. 3C representing fragments identified from bioinformatics analysis.

BRIEF SUMMARY OF THE INVENTION

[0009] Embodiments of the present invention relate generally to the discovery of protein fragments of aminoacyl-tRNA synthetases (AARSs), which possess non-canonical biological activities, such as extracellular signaling activities, and/or other characteristics of therapeutic and diagnostic relevance. The AARSs are universal and essential elements of the protein synthesis machinery found in all organisms, but human AARSs and their associated proteins have naturally-occurring resected variants, with potent cell signaling activities that contribute to normal functioning of humans. The activities of these protein fragments are distinct from the protein synthesis activities commonly known for AARSs, and the present invention includes the discovery

and development of these resected proteins as new biotherapeutic agents, new discovery research reagents, and as new antigens/targets for directed biologics and diagnostic agents that can be used to potentially treat or diagnose a wide variety of human diseases, such as inflammatory, hematological, neurodegenerative, autoimmune, hematopoietic, cardiovascular, and metabolic diseases or disorders.

[0010] The AARS protein fragment(s) of the present invention may therefore be referred to as “resectins,” or alternatively as “appendacrine.” As noted above, the term “resectin” derives from the process of excising or resecting a given AARS protein fragment from the context of its full-length parent AARS sequence, which typically masks its non-canonical activities. In certain instances, the AARS protein fragments and polynucleotides of the present invention were identified through the occurrence of this resection process, whether naturally-occurring (e.g., proteolytic, splice variant), artificially-induced, or predicted. The term “appendacrine” derives from a combination of “append” (from Latin—appender) and to “separate” or “discern” (from Greek—crines),” and also reflects the separation of one or more appended domains of the AARS protein fragments from their corresponding full-length or parent AARS sequences.

[0011] Although a few AARS fragments have been previously shown to have non-synthetase activities, the expression, isolation, purification, and characterization of such fragments for biotherapeutic, discovery, or diagnostic utility is limited, and persons skilled in the art would not have readily appreciated such activities to associate with each member of the entire family of AARSs, or with alternative fragments. Here, a methodical approach was utilized to discover and verify AARS protein fragments for the 20 mitochondrial and 20 cytosolic AARSs (and associated proteins) for biotherapeutic discovery and diagnostic utility. For instance, certain of the present AARS protein fragment (s) and polynucleotides that encode them are identified from biological samples using mass spectrometry (MS), mainly to identify proteolytic fragments, and others were identified by deep sequencing techniques, mainly to identify splice variants. Other AARS protein fragment(s) are identified using in silico predictions of amino acid sequences, such as by computationally comparing synthetases from humans and lower organisms along with key demarcations (e.g., protease sites); this approach utilized sequence analysis of the full-length AARS based on specific criteria to discern proteolytic fragments and functional domains possessing non-canonical biological activities.

[0012] Novel resectins of the AARSs are unexpected, and their differential expression is also unexpected. Specific resections are typically seen under different treatments (e.g., from cells grown in media with or without serum), at different stages of growth (e.g., adult brain vs. fetal brain) and for different tissue types (e.g., pancreas vs. liver). The pattern of expression is not the same for all aminoacyl tRNA synthetases despite the fact that the canonical functions for all aminoacyl tRNA synthetases are needed in the same cell locations and in relatively proportional amounts. One would not expect the levels of an aminoacyl tRNA synthetase activity to increase without an increase in the amounts of other aminoacyl tRNA synthetase activities at the same time. The mass spectrometry and deep sequencing data indicates that aminoacyl tRNA synthetase resectins do have varying levels and do occur in different sites and at different stages.

[0013] In addition, AARS protein fragments can be expressed and purified to sufficiently high purity to discern their biological properties. Previously, fragments were often not of sufficient purity, folding, and stability to enable proper biological characterization of non-synthetase activities. Cell based assays, for instance, are used in conjunction with sufficiently pure, stable, soluble and folded resectins to reveal their important biotherapeutic, discovery or diagnostic activities.

[0014] In particular, embodiments of the present invention relate to protein fragments of Arginyl tRNA synthetases, related agents and compositions of biotherapeutic, discovery, or diagnostic utility, and methods of use thereof. The compositions of the present invention are useful in a variety of diagnostic, drug discovery, and therapeutic applications, as described herein. Preferably, the AARS proteins and fragments are purified and stored in suitable condition to the extent required for such biotherapeutic, discovery, or diagnostic uses.

[0015] Certain embodiments include therapeutic compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least about 100, 90, 80, 70, 60, 50 or 40 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein the protein fragment has an EC_{50} of about 10 nM or less for a cell-based non-canonical biological activity and has a solubility of at least about 10 mg/ml, and wherein the composition has a purity of at least about 95% on a protein basis and substantially undetectable levels of endotoxin. In specific embodiments, the composition is substantially serum free. Various assays for measuring such features of resectins are described herein and may be used to define aspects of the invention. In certain aspects, these features will be preferable for biotherapeutic utility of the AARS protein fragments described herein.

[0016] Certain embodiments include compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least about 100, 90, 80, 70, 60, 50 or 40 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, and has a solubility of at least about 5 mg/ml, and wherein the composition has a purity of at least about 95% on a protein basis, and less than about 10 EU/mg protein endotoxin. In one aspect, the composition is a therapeutic composition. In specific embodiments, the composition is substantially serum free. In some embodiments the AARS protein fragment comprises a non-canonical activity. In some embodiments, the non-canonical biological activity is selected from modulation of extracellular signaling, modulation of cell proliferation, modulation of cell differentiation, modulation of gene transcription, modulation of cytokine production or activity, modulation of cytokine receptor activity, and modulation of inflammation. In some embodiments, the AARS protein fragment has an EC_{50} of less than about 1 nM, about 5 nM, about 10 nM, about 50 nM, about 100 nM or about 200 nM for a cell-based non-canonical biological activity.

[0017] In certain embodiments the AARS protein fragment is fused to a heterologous polypeptide. In some embodiments, the AARS fusion protein substantially retains a non-canonical activity of the AARS protein fragment. In some embodiments, the AARS fusion protein suppresses a non-canonical activity of the AARS protein fragment. In some embodiments, the heterologous polypeptide is attached to the N-terminus of the AARS protein fragment. In

some embodiments, the heterologous polypeptide is attached to the C-terminus of the AARS protein fragment. In one aspect of any of these embodiments the heterologous polypeptide is selected from the group consisting of purification tags, epitope tags, targeting sequences, signal peptides, membrane translocating sequences, and PK modifiers.

[0018] In certain embodiments, the composition comprises an AARS protein fragment at a concentration of least about 10 mg/mL. In certain embodiments the composition comprises an AARS protein fragment which is at least 90% monodisperse. In certain embodiments the composition comprises less than about 3% high molecular weight aggregated proteins. In certain embodiments the composition exhibits less than 3% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C. In certain embodiments the composition exhibits less than 3% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at room temperature.

[0019] Various assays for measuring such features of resectins are described herein and may be used to define aspects of the invention. In certain aspects, these features will be preferable for biotherapeutic utility of the AARS protein fragments described herein.

[0020] Certain embodiments include compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that differs from an amino acid sequence set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9 by substitution, deletion, and/or addition of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 amino acids, wherein the altered protein fragment substantially retains a non-canonical activity of the unaltered protein, or has a dominant negative phenotype in relation to the non-canonical activity, wherein the protein fragment has a solubility of at least about 5 mg/mL, and wherein the composition has a purity of at least about 95% on a protein basis and less than about 10 EU/mg protein endotoxin. In specific embodiments, the composition is substantially serum free.

[0021] Other embodiments include compositions, comprising an isolated antibody that specifically binds to an isolated aminoacyl-tRNA synthetase (AARS) protein fragment as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein affinity of the antibody for the AARS protein fragment is about 10× stronger than its affinity for a corresponding full-length AARS polypeptide. One of the surprising aspects of the present invention includes certain resectins possessing “new” surfaces accessible to antibody or other directed biologics, whereas the full length AARS “hides” or covers these surfaces with other sequences or adjacent domains. The process of resecting can also create greater aqueous accessibility for revealing previously unidentified biological activities. Some embodiments include compositions, comprising an isolated antibody that specifically binds to an isolated aminoacyl-tRNA synthetase (AARS) protein fragment as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein the antibody has an affinity of at least about 10 nM for the AARS protein fragment, and an affinity of at least about 100 nM for a corresponding full-length AARS polypeptide. In some embodiments, the antibody binds to an epitope located within an AARS polypeptide unique splice junction as set forth in any of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or to an amino acid sequence C-terminal of this splice site. In certain embodiments, the antibody antagonizes the non-

canonical activity of the AARS protein fragment. Such antagonists may optionally bind the corresponding parental or full-length AARS.

[0022] Other aspects relate to bioassay systems, comprising a substantially pure aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, and a binding partner that binds to the AARS protein fragment. In one aspect, the binding partner is selected from the group consisting of a cellular surface receptor protein, nucleic acid, lipid membrane, cell regulatory protein, enzyme, and transcription factor. Optionally, such a receptor may be part of a cell, preferably a cell relevant to the revealed biology of the resectin.

[0023] Certain embodiments include cellular compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, and an engineered population of cells in which at least one cell comprises a polynucleotide encoding said AARS protein fragment. In one aspect, the cells are capable of growing in a serum free medium.

[0024] Also included are detection systems, comprising a substantially pure aminoacyl-tRNA synthetase (AARS) protein fragment of at least 50 or 100 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, a cell that comprises a cell-surface receptor or an extracellular portion thereof that binds to the protein fragment, and a molecule of less than about 2000 daltons, or a second polypeptide, which modulates binding or interaction between the AARS protein fragment and the extracellular receptor.

[0025] Particular embodiments include diagnostic systems, comprising a substantially pure aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, and a cell that comprises a cell-surface receptor or an extracellular portion thereof that binds to the AARS protein fragment, wherein the system or cell comprises an indicator molecule that allows detection of a change in the levels or activity of the cell-surface receptor or extracellular portion thereof.

[0026] Certain embodiments include cellular growth devices, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, an engineered population of cells in which at least one cell comprises a polynucleotide encoding said AARS protein fragment, at least about 10 liters of serum-free cell media, and a sterile container. In specific embodiments, the cells utilized for any of the methods or compositions described herein are capable of growing in serum-free media, optionally with an antibiotic and an inducer.

[0027] Some embodiments relate to antisense or RNA interference (RNAi) agents, comprising a sequence that is targeted against a unique splice junction of an AARS splice variant as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9.

[0028] Also included are therapeutic compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an

amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein the protein fragment specifically binds to a binding partner and has a solubility of at least about 5 mg/ml, and wherein the composition has a purity of at least about 95% on a protein basis. In some aspects, the composition may have less than 10 EU endotoxin/mg protein.

[0029] Also included are compositions, comprising an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that is at least 80%, 85%, 90%, 95%, 98%, or 100% identical to an amino acid sequence set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein the protein fragment has a solubility of at least about 5 mg/ml, and wherein the composition has a purity of at least about 95% on a protein basis and less than 10 EU endotoxin/mg protein. In any of these embodiments, the compositions may comprise an AARS protein fragment that is at least about 50%, about 60%, about 70%, about 80%, about 90% or about 95% monodisperse with respect to its apparent molecular mass. In another aspect of any of these embodiments, the compositions comprise less than about 10% (on a protein basis) high molecular weight aggregated proteins, or less than about 5% high molecular weight aggregated proteins, or less than about 4% high molecular weight aggregated proteins, or less than about 3% high molecular weight aggregated proteins, or less than 2% high molecular weight aggregated proteins, or less than about 1% high molecular weight aggregated proteins.

[0030] In another aspect of any of these embodiments, the compositions exhibits less than about 10% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C., or less than about 5% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C., or less than about 3% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C., or less than about 2% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C., or less than about 1% aggregation when stored at a concentration of at least 10 mg/mL in PBS for one week at 4° C.

[0031] Certain embodiments include compositions, comprising a substantially pure aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, and at least one covalently or non-covalently moiety attached thereto. In some embodiments, the moiety is a detectable label. In some embodiments, the moiety is a water soluble polymer. In some embodiments, the moiety is PEG. In one aspect of any of these embodiments, the moiety is attached to the N-terminus of the protein fragment. In one aspect of any of these embodiments, the moiety is attached to the C-terminus of the protein fragment.

[0032] Particular embodiments include compositions, comprising a solid substrate attached to an isolated aminoacyl-tRNA synthetase (AARS) protein fragment of at least 70 amino acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or a biologically active fragment or variant thereof, wherein the protein fragment has a solubility of at least about 5 mg/ml, and the composition has a purity of at least about 95% on a protein basis.

[0033] Also included are compositions, comprising a binding agent that specifically binds to an isolated amino-

acyl-tRNA synthetase (AARS) protein fragment as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, wherein the binding agent has an affinity of at least about 1 nM for the protein fragment. In one aspect, the binding agent binds to an epitope located within an AARS polypeptide unique splice junction as set forth in any of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or to an amino acid sequence C-terminal of this splice site. In some embodiments, the binding agent antagonizes a non-canonical activity of the AARS polypeptide.

[0034] Certain embodiments include isolated aminoacyl-tRNA synthetase (AARS) polypeptides, comprising an amino acid sequence of an AARS protein fragment as described herein, an amino acid sequence encoded by an AARS polynucleotide as described herein, or a variant or fragment thereof. Certain AARS polypeptides comprise an amino acid sequence that is at least 80%, 85%, 90%, 95%, 98%, or 100% identical to an AARS reference sequence as disclosed in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. Certain AARS polypeptides consist essentially of an amino acid sequence that is at least 80%, 85%, 90%, 95%, 98%, or 100% identical to an AARS reference sequence as disclosed in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. In certain embodiments, the polypeptide comprises a non-canonical biological activity. In specific embodiments, the non-canonical biological activity is selected from modulation of cell signaling (e.g., extracellular signaling), modulation of cell proliferation, modulation of cell migration, modulation of cell differentiation, modulation of apoptosis or cell death, modulation of angiogenesis, modulation of cell binding, modulation of cellular metabolism, modulation of cellular uptake, modulation of gene transcription, or secretion, modulation of cytokine production or activity, modulation of cytokine receptor activity, and modulation of inflammation.

[0035] Other aspects include antibodies and other binding agents that exhibit binding specificity for an isolated AARS polypeptide as described herein, a binding partner of the AARS polypeptide, or the complex of both. In some embodiments, the affinity of the antibody or binding agent for the AARS polypeptide is about 10× stronger than its affinity for a corresponding full-length AARS polypeptide. In specific embodiments, the binding agent is selected from a peptide, peptide mimetic, an adnectin, an aptamer, and a small molecule. In certain embodiments, the antibody or binding agent antagonizes a non-canonical activity of the AARS polypeptide. In other embodiments, the antibody or binding agent agonizes a non-canonical activity of the AARS polypeptide.

[0036] Certain embodiments include isolated aminoacyl-tRNA synthetase (AARS) polynucleotides, comprising a nucleotide sequence of an AARS polynucleotide as described herein, a nucleotide sequence that encodes an AARS protein fragment as described herein, or a variant, a fragment, or a complement thereof. Certain AARS polynucleotides comprise a nucleotide sequence that is at least 80%, 85%, 90%, 95%, 98%, or 100% identical to an AARS reference polynucleotide, or a complement thereof, as disclosed in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. In some embodiments, the nucleotide sequence is codon optimized for bacterial expression. In one aspect, the nucleotide sequence is at least 80% identical a polynucleotide sequence disclosed in Table E2.

[0037] Specific AARS polynucleotides consist essentially of a nucleotide sequence that is at least 80%, 85%, 90%, 95%, 98%, or 100% identical to an AARS reference polynucleotide, or a complement thereof, as disclosed in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. Other AARS polynucleotides comprise or consist essentially of a nucleotide sequence that specifically hybridizes to an AARS reference polynucleotide, as disclosed in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. In certain embodiments, the polynucleotide is selected from a primer, a probe, and an antisense oligonucleotide. In specific embodiments, the primer, probe, or antisense oligonucleotide is targeted to a specific or unique splice junction, and/or sequence 3' of this splice site within an AARS polynucleotide.

[0038] Certain embodiments include methods of determining presence or levels of an AARS protein fragment in a sample, comprising contacting the sample with one or more binding agents that specifically bind to an AARS protein fragment as described herein, detecting the presence or absence of the binding agent, and thereby determining the presence or levels of the AARS protein fragment. Other embodiments include methods of determining presence or levels of an AARS protein fragment in a sample, comprising analyzing the sample with a detector that is capable of specifically identifying a protein fragment as described herein, and thereby determining the presence or levels of the AARS protein fragment. In specific embodiments, the detector is a mass spectrometer (MS), a flow cytometer, a protein imaging device, an enzyme-linked immunosorbent assays (ELISA), or a protein microarray. Certain embodiments comprise comparing the presence or levels of the AARS protein fragment to a control sample or a predetermined value. Certain embodiments comprise characterizing the state of the sample to distinguish it from the control. In specific embodiments, the sample and control comprise a cell or tissue, and the method comprises distinguishing between cells or tissues of different species, cells of different tissues or organs, cells at different cellular developmental states, cells at different cellular differentiation states, cells at different physiological states, or healthy and diseased cells. For instance, selected resectins may be more abundant under conditions such as stress or insult.

[0039] Certain embodiments include discovery methods of, and related compositions for, identifying a compound that specifically binds to an aminoacyl-tRNA synthetase (AARS) polypeptide as described herein, or one or more of its cellular binding partners, comprising a) combining the AARS polypeptide or its cellular binding partner or both with at least one test compound under suitable conditions, and b) detecting binding of the AARS polypeptide or its cellular binding partner or both to the test compound, thereby identifying a compound that specifically binds to the AARS polypeptide or its cellular binding partner or both. In certain embodiments, the test compound is a polypeptide or peptide, an antibody or antigen-binding fragment thereof, a peptide mimetic, or a small molecule. In certain embodiments, the test compound agonizes a non-canonical biological activity of the AARS polypeptide or its cellular binding partner. In other embodiments, the test compound antagonizes a non-canonical biological activity of the AARS polypeptide or its cellular binding partner. Certain embodiments include a compound identified by the above-method, such as an agonist (e.g., small molecule, peptide).

[0040] Certain embodiments include methods of determining presence or levels of a polynucleotide sequence of an AARS splice variant in a sample, comprising contacting the sample with one or more oligonucleotides that specifically hybridize to an AARS polynucleotide as described herein, detecting the presence or absence of the oligonucleotides in the sample, and thereby determining the presence or levels of the polynucleotide sequence of the AARS splice variant. Other embodiments include methods of determining presence or levels of a polynucleotide sequence of an AARS splice variant in a sample, comprising contacting the sample with at least two oligonucleotides that specifically amplify an AARS polynucleotide as described herein, performing an amplification reaction, detecting the presence or absence of an amplified product, and thereby determining presence or levels of the polynucleotide sequence of the AARS splice variant. In specific embodiments, the oligonucleotide(s) specifically hybridize to or specifically amplify a splice junction that is unique to the AARS splice variant. Certain embodiments include comparing the presence or levels of the AARS protein fragment or splice variant to a control sample or a predetermined value. Certain embodiments include characterizing the state of the sample to distinguish it from the control. In specific embodiments, the sample and control comprise a cell or tissue, and the method comprises distinguishing between cells or tissues of different species, cells of different tissues or organs, cells at different cellular developmental states, cells at different cellular differentiation states, or healthy and diseased cells.

[0041] Some embodiments include pharmaceutical compositions, comprising an AARS polynucleotide described herein, an AARS polypeptide described herein, a binding agent as described herein, or a compound identified by the above-method or described herein, and a pharmaceutically acceptable excipient or carrier.

[0042] Certain embodiments include methods of modulating a cellular activity of a cell, comprising contacting the cell with an AARS polynucleotide described herein, an AARS polypeptide described herein, a binding agent described herein, a compound of the above-method or described herein, or a pharmaceutical composition described herein. In specific embodiments, the cellular activity is selected from cell proliferation, cell migration, cell differentiation, apoptosis or cell death, cell signaling, angiogenesis, cell binding, cellular uptake, cell secretion, metabolism, cytokine production or activity, cytokine receptor activity, gene transcription, and inflammation. In one aspect, the cell is selected from the group consisting of pre-adipocytes, bone marrow, neutrophils, blood cells, hepatocytes, astrocytes, mesenchymal stem cells, and skeletal muscle cells.

[0043] In certain embodiments, the cell is in a subject. Certain embodiments comprise treating the subject, wherein the subject has a condition associated with a neoplastic disease, an immune system disease or condition, an infectious disease, a metabolic disease, an inflammatory disorder, neuronal/neurological disease, a muscular/cardiovascular disease, a disease associated with aberrant hematopoiesis, a disease associated with aberrant angiogenesis, or a disease associated with aberrant cell survival.

[0044] Also included are processes for manufacturing a pharmaceutical compound, comprising: a) performing an in vitro screen of one or more candidate compounds in the presence an AARS protein fragment of at least 70 amino

acids that comprises an amino acid sequence as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, to identify a compound that specifically binds to the AARS protein fragment; b) performing a cell-based or biochemical or receptor assay with the compound identified in step a), to identify a compound that modulates one or more non-canonical activities of the AARS protein fragment; c) optionally assessing the structure-activity relationship (SAR) of the compound identified in step b), to correlate its structure with modulation of the non-canonical activity, and optionally derivatizing the compound to alter its ability to modulate the non-canonical activity; and d) producing sufficient amounts of the compound identified in step b), or the derivatized compound in step c), for use in humans, thereby manufacturing the pharmaceutical compound.

[0045] Other embodiments include processes for manufacturing a pharmaceutical compound, comprising: a) performing an in vitro screen of one or more candidate compounds in the presence a cell-surface receptor or an extracellular portion thereof that specifically binds to an AARS protein fragment of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, to identify a compound that specifically binds to the cell-surface receptor or extracellular portion thereof; b) performing a cell-based or biochemical or receptor assay with the compound identified in step a), to identify a compound that modulates one or more non-canonical activities of the AARS protein fragment; c) optionally assessing the structure-activity relationship (SAR) of the compound identified in step b), to correlate its structure with modulation of the non-canonical activity, and optionally derivatizing the compound to alter its ability to modulate the non-canonical activity; and d) producing sufficient amounts of the compound identified in step b), or the derivatized compound in step c), for use in humans, thereby manufacturing the pharmaceutical compound.

[0046] Some embodiments include a cellular composition, comprising an engineered population of cells in which at least one cell comprises a polynucleotide encoding a heterologous full length aminoacyl-tRNA synthetase (AARS) protein, wherein the cells are capable of growing in a serum-free medium. In one aspect, the full length aminoacyl-tRNA synthetase (AARS) protein comprises a heterologous purification or epitope tag to facilitate purification of an AARS protein fragment. In another aspect, the full length aminoacyl-tRNA synthetase (AARS) protein comprises a heterologous proteolysis site to enable production of the AARS protein fragment upon cleavage.

[0047] Some embodiments include a method for producing an AARS polypeptide as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2 in situ within a cell, comprising: i) expressing a heterologous full length aminoacyl-tRNA synthetase (AARS) protein within the cell, wherein the cell comprises a protease capable of cleaving the heterologous full length aminoacyl-tRNA synthetase (AARS) protein to produce the AARS polypeptide.

[0048] Some embodiments include a method for producing an AARS polypeptide as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2 comprising contacting an isolated full length aminoacyl-tRNA synthetase (AARS) protein with a protease that is capable of cleaving the full length aminoacyl-tRNA synthetase (AARS) protein and producing an AARS polypeptide.

[0049] Some embodiments include an engineered full length aminoacyl-tRNA synthetase (AARS) protein com-

prising a heterologous proteolysis site to enable the proteolytic generation of an AARS protein fragment as set forth in any of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9 or Table E2.

[0050] Some embodiments include a composition, comprising an isolated full length aminoacyl-tRNA synthetase protein, wherein the composition has a purity of at least about 95% on a protein basis, less than about 10 EU endotoxin/mg protein, and is substantially serum free. In one aspect, the full length aminoacyl-tRNA synthetase protein is present at a concentration of at least 10 mg/mL, and is at least 90% monodisperse.

[0051] A further embodiment includes a method of treating a disease or disorder mediated by the dysregulation of the expression, activity or spatiotemporal location of a tRNA synthetase via the administration of an AARS protein fragment, or nucleic acid encoding the AARS protein fragment, as set forth in any of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, or Table E2. In one aspect of this embodiment, the disease is selected cancer, neuropathy, diabetes, and inflammatory disorders.

DETAILED DESCRIPTION OF THE INVENTION

[0052]

TABLE OF CONTENTS

I.	OVERVIEW	15
II.	DEFINITIONS	16
III.	PURIFIED AARS PROTEIN FRAGMENTS AND VARIANTS	28
IV.	AARS POLYNUCLEOTIDES	67
V.	ANTIBODIES	79
VI.	ANTIBODY ALTERNATIVES AND OTHER BINDING AGENTS	84
VII.	BIOASSAYS AND ANALYTICAL ASSAYS	89
VIII.	EXPRESSION AND PURIFICATION SYSTEMS	91
IX.	DIAGNOSTIC METHODS AND COMPOSITIONS	104
X.	ANTISENSE AND RNAI AGENTS	119
	A. ANTISENSE AGENTS	120
	B. RNA INTERFERENCE AGENTS	128
XI.	DRUG DISCOVERY	136
XII.	METHODS OF USE	144
XIII.	PHARMACEUTICAL FORMULATIONS, ADMINISTRATION AND KITS	148
XIV.	EXAMPLES	157

I. OVERVIEW

[0053] The current invention is directed, at least in part, to the discovery of novel AARS polypeptides, and methods for their preparation and use, which represent the transformation of native wild type proteins into new forms that exhibit markedly different characteristics compared to the naturally occurring full length Arginyl tRNA synthetase genes. Such AARS polypeptides were identified based on extensive sequence, and mass spectrometry analysis of expressed Arginyl tRNA synthetase in different tissues, followed by the systematic production and testing of each potential AARS polypeptide to identify protein sequences that represent stable and soluble protein domains which exhibit novel biological activities, and favorable therapeutic drug characteristics.

[0054] Based on this analysis at least three new novel families of AARS polypeptides derived from Arginyl tRNA synthetase have been identified.

[0055] In one aspect, such Arginyl RNA synthetase derived AARS polypeptides comprise polypeptide sequences comprising approximately comprising amino acids 1-71 of Arginyl tRNA synthetase.

[0056] In a second aspect, such Arginyl tRNA synthetase derived AARS polypeptides comprise polypeptide sequences comprising approximately comprising amino acids 168-530 of Arginyl tRNA synthetase.

[0057] In a third aspect, such Arginyl tRNA synthetase derived AARS polypeptides comprise polypeptide sequences comprising approximately comprising amino acids 551-660 of Arginyl tRNA synthetase.

[0058] These new AARS polypeptide families represent novel, previously unknown protein products which exhibit inter alia i) novel biological activity, ii) favorable protein stability and aggregation characteristics, and iii) the ability to be expressed and produced at high level in prokaryotic expression systems, which are materially different characteristics not found in the intact wild type protein.

II. DEFINITIONS

[0059] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, preferred methods and materials are described. For the purposes of the present invention, the following terms are defined below.

[0060] The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0061] By “about” is meant a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

[0062] An “agonist” refers to a molecule that intensifies or mimics an activity. For example, a non-canonical biological activity of an AARS, or another protein. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition that modulates the activity of an AARS either by directly interacting with the AARS or its binding partner, or by acting on components of the biological pathway in which the AARS participates. Included are partial and full agonists.

[0063] As used herein, the term “amino acid” is intended to mean both naturally occurring and non-naturally occurring amino acids as well as amino acid analogs and mimetics. Naturally occurring amino acids include the 20 (L)-amino acids utilized during protein biosynthesis as well as others such as 4-hydroxyproline, hydroxylysine, desmosine, isodesmosine, homocysteine, citrulline and ornithine, for example. Non-naturally occurring amino acids include, for example, (D)-amino acids, norleucine, norvaline, p-fluorophenylalanine, ethionine and the like, which are known to a person skilled in the art. Amino acid analogs include modified forms of naturally and non-naturally occurring amino acids. Such modifications can include, for example, substi-

tution or replacement of chemical groups and moieties on the amino acid or by derivitization of the amino acid. Amino acid mimetics include, for example, organic structures which exhibit functionally similar properties such as charge and charge spacing characteristic of the reference amino acid. For example, an organic structure which mimics Arginine (Arg or R) would have a positive charge moiety located in similar molecular space and having the same degree of mobility as the e-amino group of the side chain of the naturally occurring Arg amino acid. Mimetics also include constrained structures so as to maintain optimal spacing and charge interactions of the amino acid or of the amino acid functional groups. Those skilled in the art know or can determine what structures constitute functionally equivalent amino acid analogs and amino acid mimetics.

[0064] In certain aspects, the use of non-natural amino acids can be utilized to modify (e.g. increase) a selected non-canonical activity of an AARS protein fragment, or to alter the in vivo or in vitro half-life of the protein. Non-natural amino acids can also be used to facilitate (selective) chemical modifications (e.g., pegylation) of an AARS protein. For instance, certain non-natural amino acids allow selective attachment of polymers such as PEG to a given protein, and thereby improve their pharmacokinetic properties.

[0065] Specific examples of amino acid analogs and mimetics can be found described in, for example, Roberts and Vellaccio, *The Peptides: Analysis, Synthesis, Biology*, Eds. Gross and Meinhofer, Vol. 5, p. 341, Academic Press, Inc., New York, N.Y. (1983), the entire volume of which is incorporated herein by reference. Other examples include peralkylated amino acids, particularly permethylated amino acids. See, for example, *Combinatorial Chemistry*, Eds. Wilson and Czarnik, Ch. 11, p. 235, John Wiley & Sons Inc., New York, N.Y. (1997), the entire book of which is incorporated herein by reference. Yet other examples include amino acids whose amide portion (and, therefore, the amide backbone of the resulting peptide) has been replaced, for example, by a sugar ring, steroid, benzodiazepine or carbo cycle. See, for instance, *Burger's Medicinal Chemistry and Drug Discovery*, Ed. Manfred E. Wolff, Ch. 15, pp. 619-620, John Wiley & Sons Inc., New York, N.Y. (1995), the entire book of which is incorporated herein by reference. Methods for synthesizing peptides, polypeptides, peptidomimetics and proteins are well known in the art (see, for example, U.S. Pat. No. 5,420,109; M. Bodanzsky, *Principles of Peptide Synthesis* (1st ed. & 2d rev. ed.), Springer-Verlag, New York, N.Y. (1984 & 1993), see Chapter 7; Stewart and Young, *Solid Phase Peptide Synthesis*, (2d ed.), Pierce Chemical Co., Rockford, Ill. (1984), each of which is incorporated herein by reference). Accordingly, the AARS polypeptides of the present invention may be composed of naturally occurring and non-naturally occurring amino acids as well as amino acid analogs and mimetics.

[0066] The term “antagonist” refers to a molecule that reduces or attenuates an activity. For example, a non-canonical biological activity of an AARS, or another protein. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition that modulates the activity of an AARS or its binding partner, either by directly interacting with the AARS or its binding partner or by acting on components of the biological pathway in which the AARS participates. Included are partial and full antagonists.

[0067] The term “aminoacyl-tRNA synthetase” (AARS) refers generally to enzymes that in their natural or wild-type form are capable of catalyzing the esterification of a specific amino acid or its precursor to one of all its compatible cognate tRNAs to form an aminoacyl-tRNA. In this “canonical” activity, aminoacyl-tRNA synthetases catalyze a two-step reaction: first, they activate their respective amino acid by forming an aminoacyl-adenylate, in which the carboxyl of the amino acid is linked in to the alpha-phosphate of ATP by displacing pyrophosphate, and then, when the correct tRNA is bound, the aminoacyl group of the aminoacyl-adenylate is transferred to the 2' or 3' terminal OH of the tRNA.

[0068] Class I aminoacyl-tRNA synthetases typically have two highly conserved sequence motifs. These enzymes aminoacylate at the 2'-OH of an adenosine nucleotide, and are usually monomeric or dimeric. Class II aminoacyl-tRNA synthetases typically have three highly conserved sequence motifs. These enzymes aminoacylate at the 3'-OH of the same adenosine, and are usually dimeric or tetrameric. The active sites of class II enzymes are mainly made up of a seven-stranded anti-parallel β -sheet flanked by α -helices. Although phenylalanine-tRNA synthetase is class II, it aminoacylates at the 2'-OH.

[0069] AARS polypeptides include sources of mitochondrial and cytoplasmic forms of tyrosyl-tRNA synthetase (TyrRS), a tryptophanyl-tRNA synthetase (TrpRS), a glutaminyl-tRNA synthetase (GlnRS), a glycyl-tRNA synthetase (GlyRS), a histidyl-tRNA synthetase (HisRS), a seryl-tRNA synthetase (SerRS), a phenylalanyl-tRNA synthetase (PheRS), an alanyl-tRNA synthetase (AlaRS), an asparaginyl-tRNA synthetase (AsnRS), an aspartyl-tRNA synthetase (AspRS), a cysteinyl-tRNA synthetase (CysRS), a glutamyl-tRNA synthetase (GluRS), a prolyl-tRNA synthetase (ProRS), an arginyl-tRNA synthetase (ArgRS), an isoleucyl-tRNA synthetase (IleRS), a leucyl-tRNA synthetase (LeuRS), a lysyl-tRNA synthetase (LysRS), a threonyl-tRNA synthetase (ThrRS), a methionyl-tRNA synthetases (MetRS), or a valyl-tRNA synthetase (ValRS). The wild-type or parental sequences of these AARS polypeptides are known in the art.

[0070] By “coding sequence” is meant any nucleic acid sequence that contributes to the code for the polypeptide product of a gene. By contrast, the term “non-coding sequence” refers to any nucleic acid sequence that does not contribute to the code for the polypeptide product of a gene.

[0071] Throughout this specification, unless the context requires otherwise, the words “comprise,” “comprises,” and “comprising” will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

[0072] By “consisting of” is meant including, and limited to, whatever follows the phrase “consisting of” Thus, the phrase “consisting of” indicates that the listed elements are required or mandatory, and that no other elements may be present. By “consisting essentially of” is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the phrase “consisting essentially of” indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be

present depending upon whether or not they materially affect the activity or action of the listed elements.

[0073] The recitation “endotoxin free” or “substantially endotoxin free” relates generally to compositions, solvents, and/or vessels that contain at most trace amounts (e.g., amounts having no clinically adverse physiological effects to a subject) of endotoxin, and preferably undetectable amounts of endotoxin. Endotoxins are toxins associated with certain bacteria, typically gram-negative bacteria, although endotoxins may be found in gram-positive bacteria, such as *Listeria monocytogenes*. The most prevalent endotoxins are lipopolysaccharides (LPS) or lipo-oligo-saccharides (LOS) found in the outer membrane of various Gram-negative bacteria, and which represent a central pathogenic feature in the ability of these bacteria to cause disease. Small amounts of endotoxin in humans may produce fever, a lowering of the blood pressure, and activation of inflammation and coagulation, among other adverse physiological effects.

[0074] Therefore, in pharmaceutical production of AARS polypeptides, it is often desirable to remove most or all traces of endotoxin from drug products and/or drug containers, because even small amounts may cause adverse effects in humans. A depyrogenation oven may be used for this purpose, as temperatures in excess of 300° C. are typically required to break down most endotoxins. For instance, based on primary packaging material such as syringes or vials, the combination of a glass temperature of 250° C. and a holding time of 30 minutes is often sufficient to achieve a 3 log reduction in endotoxin levels. Other methods of removing endotoxins are contemplated, including, for example, chromatography and filtration methods, as described herein and known in the art. Also included are methods of producing AARS polypeptides in and isolating them from eukaryotic cells such as mammalian cells to reduce, if not eliminate the risk of endotoxins being present in a composition of the invention. Preferred are methods of producing AARS polypeptides in and isolating them from serum free cells. Such compositions comprising AARS polypeptides, represent new formulations which exhibit novel and new biological and therapeutic characteristics not found in AARS polypeptide compositions contaminated with serum or endotoxin which have the potential to bind to and alter the novel biological properties of the AARS polypeptides.

[0075] Endotoxins can be detected using routine techniques known in the art. For example, the Limulus Ameobocyte Lysate assay, which utilizes blood from the horseshoe crab, is a very sensitive assay for detecting presence of endotoxin, and reagents, kits and instrumentation for the detection of endotoxin based on this assay are commercially available, for example from the Lonza Group. In this test, very low levels of LPS can cause detectable coagulation of the limulus lysate due a powerful enzymatic cascade that amplifies this reaction. Endotoxins can also be quantitated by enzyme-linked immunosorbent assay (ELISA). To be substantially endotoxin free, endotoxin levels may be less than about 0.001, 0.005, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.08, 0.09, 0.1, 0.5, 1.0, 1.5, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, or 10 EU/mg of protein. Typically, 1 ng lipopolysaccharide (LPS) corresponds to about 1-10 EU.

[0076] In certain embodiments, the “purity” of any given agent (e.g., AARS protein fragment) in a composition may be specifically defined. For instance, certain compositions may comprise an agent that is at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% pure, including all decimals

in between, as measured, for example and by no means limiting, by high pressure liquid chromatography (HPLC), a well-known form of column chromatography used frequently in biochemistry and analytical chemistry to separate, identify, and quantify compounds.

[0077] As used herein, the terms “function” and “functional” and the like refer to a biological, enzymatic, or therapeutic function.

[0078] By “gene” is meant a unit of inheritance that may occupy a specific locus on a chromosome and consists of transcriptional and/or translational regulatory sequences and/or a coding region and/or non-translated sequences (i.e., introns, 5' and 3' untranslated sequences).

[0079] “Homology” refers to the percentage number of amino acids that are identical or constitute conservative substitutions. Homology may be determined using sequence comparison programs such as GAP (Deveraux et al., 1984, *Nucleic Acids Research* 12, 387-395), which is incorporated herein by reference. In this way sequences of a similar or substantially different length to those cited herein could be compared by insertion of gaps into the alignment, such gaps being determined, for example, by the comparison algorithm used by GAP.

[0080] The term “host cell” includes an individual cell or cell culture that can be or has been a recipient of any recombinant vector(s), isolated polynucleotide, or polypeptide of the invention. Host cells include progeny of a single host cell, and the progeny may not necessarily be completely identical (in morphology or in total DNA complement) to the original parent cell due to natural, accidental, or deliberate mutation and/or change. A host cell includes cells transfected or infected in vivo or in vitro with a recombinant vector or a polynucleotide of the invention. A host cell which comprises a recombinant vector of the invention is a recombinant host cell.

[0081] By “isolated” is meant material that is substantially or essentially free from components that normally accompany it in its native state. For example, an “isolated polynucleotide,” as used herein, includes a polynucleotide that has been purified from the sequences that flank it in its naturally-occurring state, e.g., a DNA fragment which has been removed from the sequences that are normally adjacent to the fragment. Alternatively, an “isolated peptide” or an “isolated polypeptide” and the like, as used herein, includes the in vitro isolation and/or purification of a peptide or polypeptide molecule from its natural cellular environment, and from association with other components of the cell; i.e., it is not significantly associated with in vivo substances.

[0082] The term “mRNA” or sometimes refer by “mRNA transcripts” as used herein, include, but not limited to pre-mRNA transcript(s), transcript processing intermediates, mature mRNA(s) ready for translation and transcripts of the gene or genes, or nucleic acids derived from the mRNA transcript(s). Transcript processing may include splicing, editing and degradation. As used herein, a nucleic acid derived from an mRNA transcript refers to a nucleic acid for whose synthesis the mRNA transcript or a subsequence thereof has ultimately served as a template. A cDNA reverse transcribed from an mRNA, an RNA transcribed from that cDNA, a DNA amplified from the cDNA, an RNA transcribed from the amplified DNA, etc., are all derived from the mRNA transcript and detection of such derived products is indicative of the presence and/or abundance of the original transcript in a sample. Thus, mRNA derived

samples include, but are not limited to, mRNA transcripts of the gene or genes, cDNA reverse transcribed from the mRNA, cRNA transcribed from the cDNA, DNA amplified from the genes, RNA transcribed from amplified DNA, and the like.

[0083] “Non-canonical” activity as used herein, refers generally to either i) a new activity possessed by an AARS polypeptide of the invention that is not possessed to any significant degree by the intact native full length parental protein, or ii) an activity that was possessed by the intact native full length parental protein, where the AARS polypeptide either exhibits a significantly higher (i.e. at least 20% greater) specific activity compared to the intact native full length parental protein, or exhibits the activity in a new context; for example by isolating the activity from other activities possessed by the intact native full length parental protein. In the case of AARS polypeptides, non-limiting examples of non-canonical activities include extracellular signaling, RNA-binding, amino acid-binding, modulation of cell proliferation, modulation of cell migration, modulation of cell differentiation (e.g., hematopoiesis, neurogenesis, myogenesis, osteogenesis, and adipogenesis), modulation of gene transcription, modulation of apoptosis or other forms of cell death, modulation of cell signaling, modulation of cellular uptake, or secretion, modulation of angiogenesis, modulation of cell binding, modulation of cellular metabolism, modulation of cytokine production or activity, modulation of cytokine receptor activity, modulation of inflammation, and the like.

[0084] The term “half maximal effective concentration” or “EC₅₀” refers to the concentration of an AARS protein fragment, antibody or other agent described herein at which it induces a response halfway between the baseline and maximum after some specified exposure time; the EC₅₀ of a graded dose response curve therefore represents the concentration of a compound at which 50% of its maximal effect is observed. In certain embodiments, the EC₅₀ of an agent provided herein is indicated in relation to a “non-canonical” activity, as noted above. EC₅₀ also represents the plasma concentration required for obtaining 50% of a maximum effect in vivo. Similarly, the “EC₉₀” refers to the concentration of an agent or composition at which 90% of its maximal effect is observed. The “EC₉₀” can be calculated from the “EC₅₀” and the Hill slope, or it can be determined from the data directly, using routine knowledge in the art. In some embodiments, the EC₅₀ of an AARS protein fragment, antibody, or other agent is less than about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, or 100 nM. Preferably, biotherapeutic composition will have an EC₅₀ value of about 1 nM or less.

[0085] The term “modulating” includes “increasing” or “stimulating,” as well as “decreasing” or “reducing,” typically in a statistically significant or a physiologically significant amount as compared to a control. Accordingly a “modulator” may be an agonist, an antagonist, or any mixture thereof depending upon the conditions used. An “increased” or “enhanced” amount is typically a “statistically significant” amount, and may include an increase that is 1.1, 1.2, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30 or more times (e.g., 500, 1000 times) (including all integers and decimal points in between and above 1, e.g., 1.5, 1.6, 1.7, 1.8, etc.) the amount produced by no composition (the absence of an agent or compound) or a control composition. A “decreased”

or reduced amount is typically a “statistically significant” amount, and may include a 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% decrease in the amount produced by no composition (the absence of an agent or compound) or a control composition, including all integers in between. As one non-limiting example, a control in comparing canonical and non-canonical activities could include the AARS protein fragment of interest compared to its corresponding full-length AARS, or a fragment AARS having comparable canonical activity to its corresponding full-length AARS. Other examples of “statistically significant” amounts are described herein.

[0086] By “obtained from” is meant that a sample such as, for example, a polynucleotide extract or polypeptide extract is isolated from, or derived from, a particular source of the subject. For example, the extract can be obtained from a tissue or a biological fluid isolated directly from the subject. “Derived” or “obtained from” can also refer to the source of a polypeptide or polynucleotide sequence. For instance, an AARS sequence of the present invention may be “derived” from the sequence information of an AARS proteolytic fragment or AARS splice variant, or a portion thereof, whether naturally-occurring or artificially generated, and may thus comprise, consist essentially of, or consist of that sequence

[0087] The terms “polypeptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues and to variants and synthetic and naturally occurring analogues of the same. Thus, these terms apply to amino acid polymers in which one or more amino acid residues are synthetic non-naturally occurring amino acids, such as a chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally-occurring amino acid polymers and naturally occurring chemical derivatives thereof. Such derivatives include, for example, post-translational modifications and degradation products including pyroglutamyl, iso-aspartyl, proteolytic, phosphorylated, glycosylated, oxidized, isomerized, and deaminated variants of the AARS reference fragment.

[0088] The recitations “sequence identity” or, for example, comprising a “sequence 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity.

[0089] Terms used to describe sequence relationships between two or more polynucleotides or polypeptides include “reference sequence,” “comparison window,” “sequence identity,” “percentage of sequence identity” and

“substantial identity.” A “reference sequence” is at least 12 but frequently 15 to 18 and often at least 25 monomer units, inclusive of nucleotides and amino acid residues, in length. Because two polynucleotides may each comprise (1) a sequence (i.e., only a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) a sequence that is divergent between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a “comparison window” to identify and compare local regions of sequence similarity. A “comparison window” refers to a conceptual segment of at least 6 contiguous positions, usually about 50 to about 100, more usually about 100 to about 150 in which a sequence is compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. The comparison window may comprise additions or deletions (i.e., gaps) of about 20% or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, Wis., USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., 1997, *Nucl. Acids Res.* 25:3389. A detailed discussion of sequence analysis can be found in Unit 19.3 of Ausubel et al., “Current Protocols in Molecular Biology,” John Wiley & Sons Inc, 1994-1998, Chapter 15.

[0090] Calculations of sequence similarity or sequence identity between sequences (the terms are used interchangeably herein) are performed as follows. To determine the percent identity of two amino acid sequences, or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second amino acid or nucleic acid sequence for optimal alignment and non-homologous sequences can be disregarded for comparison purposes). In certain embodiments, the length of a reference sequence aligned for comparison purposes is at least 30%, preferably at least 40%, more preferably at least 50%, 60%, and even more preferably at least 70%, 80%, 90%, 100% of the length of the reference sequence. The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in the first sequence is occupied by the same amino acid residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position.

[0091] The percent identity between the two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences.

[0092] The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch, (1970, *J. Mol. Biol.* 48: 444-453) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package, using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. A particularly preferred set of parameters (and the one that should be used unless otherwise specified) are a Blossum 62 scoring matrix with a gap penalty of 12, a gap extend penalty of 4, and a frame shift gap penalty of 5.

[0093] The percent identity between two amino acid or nucleotide sequences can be determined using the algorithm of E. Meyers and W. Miller (1989, *Cabios*, 4: 11-17) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

[0094] The nucleic acid and protein sequences described herein can be used as a "query sequence" to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al., (1990, *J. Mol. Biol.* 215: 403-10). BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997, *Nucleic Acids Res.* 25: 3389-3402). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

[0095] The term "solubility" refers to the property of an agent provided herein to dissolve in a liquid solvent and form a homogeneous solution. Solubility is typically expressed as a concentration, either by mass of solute per unit volume of solvent (g of solute per kg of solvent, g per dL (100 mL), mg/ml, etc.), molarity, molality, mole fraction or other similar descriptions of concentration. The maximum equilibrium amount of solute that can dissolve per amount of solvent is the solubility of that solute in that solvent under the specified conditions, including temperature, pressure, pH, and the nature of the solvent. In certain embodiments, solubility is measured at physiological pH. In certain embodiments, solubility is measured in water or a physiological buffer such as PBS. In certain embodiments, solubility is measured in a biological fluid (solvent) such as blood or serum. In certain embodiments, the temperature can be about room temperature (e.g., about 20, 21, 22, 23, 24, 25° C.) or about body temperature (37° C.). In certain embodiments, an agent such as an AARS protein fragment has a solubility of at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6,

0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, or 30 mg/ml at room temperature or at 37° C.

[0096] A "splice junction" as used herein includes the region in a mature mRNA transcript or the encoded polypeptide where the 3' end of a first exon joins with the 5' end of a second exon. The size of the region may vary, and may include 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more (including all integers in between) nucleotide or amino acid residues on either side of the exact residues where the 3' end of one exon joins with the 5' end of another exon. An "exon" refers to a nucleic acid sequence that is represented in the mature form of an RNA molecule after either portions of a precursor RNA (introns) have been removed by cis-splicing or two or more precursor RNA molecules have been ligated by trans-splicing. The mature RNA molecule can be a messenger RNA or a functional form of a non-coding RNA such as rRNA or tRNA. Depending on the context, an exon can refer to the sequence in the DNA or its RNA transcript. An "intron" refers to a non-coding nucleic acid region within a gene, which is not translated into a protein. Non-coding intronic sections are transcribed to precursor mRNA (pre-mRNA) and some other RNAs (such as long noncoding RNAs), and subsequently removed by splicing during the processing to mature RNA.

[0097] A "splice variant" refers to a mature mRNA and its encoded protein that are produced by alternative splicing, a process by which the exons of the RNA (a primary gene transcript or pre-mRNA) are reconnected in multiple ways during RNA splicing. The resulting different mRNAs may be translated into different protein isoforms, allowing a single gene to code for multiple proteins.

[0098] A "subject," as used herein, includes any animal that exhibits a symptom, or is at risk for exhibiting a symptom, which can be treated or diagnosed with an AARS polynucleotide or polypeptide of the invention. Also included are subjects for which it is desirable to profile levels of AARS polypeptides and/or polynucleotides of the invention, for diagnostic or other purposes. Suitable subjects (patients) include laboratory animals (such as mouse, rat, rabbit, or guinea pig), farm animals, and domestic animals or pets (such as a cat or dog). Non-human primates and, preferably, human patients, are included.

[0099] "Treatment" or "treating," as used herein, includes any desirable effect on the symptoms or pathology of a disease or condition that can be effected by the non-canonical activities of an AARS polynucleotide or polypeptide, as described herein, and may include even minimal changes or improvements in one or more measurable markers of the disease or condition being treated. Also included are treatments that relate to non-AARS therapies, in which an AARS sequence described herein provides a clinical marker of treatment. "Treatment" or "treating" does not necessarily indicate complete eradication or cure of the disease or condition, or associated symptoms thereof. The subject receiving this treatment is any subject in need thereof. Exemplary markers of clinical improvement will be apparent to persons skilled in the art.

[0100] The practice of the present invention will employ, unless indicated specifically to the contrary, conventional methods of molecular biology and recombinant DNA techniques within the skill of the art, many of which are described below for the purpose of illustration. Such tech-

niques are explained fully in the literature. See, e.g., Sambrook, et al., *Molecular Cloning: A Laboratory Manual* (3rd Edition, 2000); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *Oligonucleotide Synthesis: Methods and Applications* (P. Herdewijn, ed., 2004); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds., 1985); *Nucleic Acid Hybridization: Modern Applications* (Buzdin and Lukyanov, eds., 2009); *Transcription and Translation* (B. Hames & S. Higgins, eds., 1984); *Animal Cell Culture* (R. Freshney, ed., 1986); Freshney, R. I. (2005) *Culture of Animal Cells, a Manual of Basic Technique*, 5th Ed. Hoboken N.J., John Wiley & Sons; B. Perbal, *A Practical Guide to Molecular Cloning* (3rd Edition 2010); Farrell, R., *RNA Methodologies: A Laboratory Guide for Isolation and Characterization* (3rd Edition 2005), *Methods of Enzymology: DNA Structure Part A: Synthesis and Physical Analysis of DNA* Methods in Enzymology, Academic Press; *Using Antibodies: A Laboratory Manual: Portable Protocol NO. 1* by Edward Harlow, David Lane, Ed Harlow (1999, Cold Spring Harbor Laboratory Press, ISBN 0-87969-544-7); *Antibodies: A Laboratory Manual* by Ed Harlow (Editor), David Lane (Editor) (1988, Cold Spring Harbor Laboratory Press, ISBN 0-87969-3, 4-2), 1855. *Handbook of Drug Screening*, edited by Ramakrishna Seethala, Prabhavathi B. Fernandes (2001, New York, N.Y., Marcel Dekker, ISBN 0-8247-0562-9); and *Lab Ref: A Handbook of Recipes, Reagents, and Other Reference Tools for Use at the Bench*, Edited Jane Roskams and Linda Rodgers, (2002, Cold Spring Harbor Laboratory, ISBN 0-87969-630-3).

[0101] All publications, patents and patent applications cited herein are hereby incorporated by reference in their entirety.

III. PURIFIED AARS PROTEIN FRAGMENTS AND VARIANTS FOR THERAPEUTICS AND OTHER APPLICATIONS

[0102] Surprisingly, and unlike their full-length parental sequences that are known only for their aminoacylation-activities, it has been found that AARS fragments possess biological activities important for biotherapeutic, discovery and diagnostic applications. Embodiments of the present invention therefore include full length proteins, mature protein isoforms and protein fragments of aminoacyl-tRNA synthetases (AARS), in addition to biologically active variants and fragments thereof. In certain embodiments, the proteins and fragments may arise through endogenous proteolysis, in vitro proteolysis, splice variation, or in silico prediction, among other mechanisms.

[0103] The AARS protein fragments described herein, and variants thereof, may possess at least one “non-canonical” biological activity. The AARS protein fragment(s) of the present invention are also referred to herein as “AARS polypeptides” or “AARS reference polypeptides.” In certain embodiments, the AARS polypeptides provided herein comprise or consist essentially of all or a portion of the AARS polypeptide “reference sequence(s)” as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9 below, which represent the amino acid sequence(s) of various fragments of Arginyl tRNA synthetases. Mouse and human AARS protein sequences are highly related, typically differing by no more than a few amino acids within an entire sequence, a particular domain, or a particular protein fragment.

N-Terminal AARS Polypeptides: (Tables 1, 2 & 3)

[0104]

TABLE 1

AARS polypeptides identified by MS			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
ArgRS ^{N1}	Protein/ Human/ 1-398	MDVLVSECSARLLQEEIEIKSL TAEIDRLKNCGLGASPNLEQL QEENLKLKYRLNLRKSLQAER NKPTKNMINIISRLQEVFGHAI KAAYPDLENPPLLVTPSQAKF GDYQCNSAMGISQMLKTEQKV NPRIEAINITKHLPDNECIEKV EIAGPGFINVHLRKDFVSEQLT SLLVNGVQLPALGENKKVIVDF SSPNIKEMHVGHLRSTIIGES ISRLFEFAGYDVLRLNHVGDWG TQFGMLIAHLQDKFPDYLTVSP PIGDLQVFPYKESKKRFDTEEF KKRAYQCIVLLQGGKNPDITKAW KLICDVSRQELNKKIYDALDVSL IERGESFYQDRMNDIVKEFEDR GFVQVDDGRKIVFVPGCSIPLT IVKSDGGTYTDSDLAAIKQRL FE	SEQ.ID. NO. 12
		ATGGACGCTACTGGTGTCTGAGT GCTCCGCGCGCTGCTGCAGCA GGAGAAGAGATTAATCTCTGT ACTGCTGAAATTGACCGGTGA AAACTGTGGCTGTTTAGGAGC TTCTCCAAATTTGGAGCAGTTA CAAGAAGAAAATTTAAATTA AGTATCGACTGAATATTCTTCG AAAGAGTCTTCAGGCAGAAAGG AACAAACCAACTAAAAATATGA TTAATATTATTAGCCGCCTACA AGAGGTCTTTGGTTCATGCAATT AAGGCTGCATATCCAGATTGG AAAATCCTCCTCTGCTAGTGAC ACCAAGTCAGCAGGCCAAGTTT GGGGACTATCAGTGTAAATAGTG CTATGGGTATTTCTCAGATGCT CAAAACCAAGGAACAGAAAGTT AATCCAAGAGAAATGTCTGA ACATTACCAACACCTCCAGAG CAATGAATGTATGAAAAAGTT GAAATTGCTGGTCTGGTTT TTAATGTCCACTTAAGAAAGGA TTTTGTATCAGAACAAATGACC AGTCTTCTAGTGAATGGAGTTC AACTACCTGCTCTGGGAGAGAA TAAAAAGGTTATAGTTGACTTT TCCTCCCTAATATAGCTAAAG AGATGCATGTAGGCACCTGAG GTCAACTATCATAGGAGAGAGT ATAAGCCGCCCTCTTGAATTTG CAGGGTATGACGTGCTCAGGTT AAATCATGTAGGAGACTGGGG ACCCAGTTTGGCATGCTCATGC CTCACCTGCAAGACAAATTTCC AGATTATCTAACAGTTTTCACCT CCTATTGGGGATCTTCAGGTCT TTTATAAGGAATCTAAGAGAG GTTTGATACTGAGGAGGAATTT AAGAAGCGAGCATATCAGTGTG TAGTTCTGCTCCAGGGTAAAA CCCAGATATTACAAAGCTTTGG AAGCTTATCTGTGATGCTCTCC GCCAAGAGTTAAATAAATCTTA TGATGCATTGGACGCTCTCTTA ATAGAGAGAGGGGAATCCTCT ATCAAGATAGGATGAATGATAT	SEQ.ID. NO. 13

TABLE 1-continued

TGTAAGGAATTGAAGATAGA GGATTGTGCAGGTGGATGATG GCAGAAAGATTGTATTTGTCCC AGGGTGTTCATACCATTAAACC ATAGTAAATCAGATGGAGGTT ATACCTATGATACATCTGACCT GGCTGCTATTAAACAAAGACTA TTTGAG		
B ArgRS ^{N1} Mass spec peptides detected and inferred linking peptides		
Type/ species	Sequence	SEQ. ID. NO.
Protein/ mouse	AAYPDLENPPLIVTPSQQPK	SEQ. ID. NO. 14
Protein/ mouse	FGDYQCNSAMGISQMLKAKEQKVS PRE IAENITKHLPNNKYIDKVEIAGPFIN VHLRK	SEQ. ID. NO. 15
Protein/ mouse	DFVSEQLTSLLVNGVQLPVLGDKEK	SEQ. ID. NO. 16
Protein/ mouse	VIVDFSSPNIK	SEQ. ID. NO. 17
Protein/ mouse	EMHVGHLRSTIIIGESMSRLFEFAGYDV LRLNHVGDWGTQFGMLIAHLQDKFPDY LTVSPPIGDLQAFYKESKKRFDADDEF KKRAYQCVVLLQSKNPDIMKAWNLICD VSREEFKKIYDALDITLIERGESFYQD RMKDIVKEFEDKGFVQVDDGRKIVFVP GC	SEQ. ID. NO. 18
Protein/ mouse	SVPLTIVKSDGGYTYDTSDLAAIK	SEQ. ID. NO. 19

C ArgRS ^{N1} Concatenated sequences based on mass spec peptides detected		
Type/ species	Sequence	SEQ. ID. NO.
Protein/ mouse	<u>AAYPDLENPPLIVTPSQQPKFGDYQCN</u> SAMGISQMLKAKEQKVS PREIAENITK HLPNNKYIDKVEIAGPFINVHLRKDF <u>VSEQLTSLLVNGVOLPVLGDKEKVI</u> <u>DFSSPNIKEMHVGHLRSTIIIGESMSRL</u> FEFAGYDVLRNLHVGDWGTQFGMLIAH LQDKFPDYLTVPPIGDLQAFYKESKK RFDADDEFKKRAYQCVVLLQSKNPDIM KAWNLICDVSREEFKKIYDALDITLIE RGESEFYQDRMKDIVKEFEDKGFVQVDD GRKIVFVPGCSVPLTIVKSDGGYTYDT <u>SDLAAIK</u>	SEQ. ID. NO. 20

TABLE 2

AARS polypeptides and alternative transcripts identified by Deep Sequencing			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ. ID. NO.
ArgRS ^{N3}	Protein/ Human/ 1-353 + 20 aa	MDVLVSECSARLLQEEIEKSL TAEIDRLKNCGLGASPNLEQL QEENLKLKYRLNLRKSLQAE NKPTKNMINIISRLQEVFGHAI KAAYPDLENPPLLVTPSQQAKF	SEQ. ID. NO. 21

TABLE 2-continued

GDYQCNSAMGISQMLKTKEQKV NPRIAENITKHLPDNECIEKV EIAGPGFINVHLRKDFVSEQLT SLLVNGVQLPALGENKKVIVDF SSPNIAKEMHVGHLRSTIIIGES ISRLFEFAGYDVLRNLHVGDWGT TQFGMLIAHLQDKFPDYLTVP PIGDLQVYKESKKRFDTEEEF KKRAYQCVVLLQGNPDITKAW KLICDVSRLQELNKIYDALDVSL IERGESFYQDRMNDIVKEFEDR GKSLKHVRVKQCASWIFWEKD			
ArgRS ^{N3}	DNA/ Human	ATGGACGTACTGGTGTCTGAGT GCTCCGCGCGGCTGCTGCAGCA GGAAGAAGAGATTAAATCTCTG ACTGCTGAAATTGACCGGTTGA AAAACTGTGGCTGTTTAGGAGC TTCTCCAAATTGGAGCAGTTA CAAGAAGAAAATTTAAATTA AGTATCGACTGAATATTTCTCG AAAGAGTCTTCAGGCAGAAAGG AACAAACCAACTAAAATATGA TTAACATTATTAGCCGCCTACA AGAGGTCTTTGGTCATGCAATT AAGGCTGCATATCCAGATTGG AAAATCCTCCTCTGCTAGTGAC ACCAAGTCAGCAGGCCAAGTTT GGGGACTATCAGTGTAAATAGTG CTATGGGTATTTCTCAGATGCT CAAAACCAAGGAACAGAAAGTT AATCCAAGAGAAATGTGCGAAA ACATTACCAACACCTCCCAGA CAATGAATGTATTGAAAAAGTT GAAATTGCTGCTCTGTTTGA TTAATGTCCACTTAAGAAAGGA TTTTGTATCAGAACAAATTGACC AGTCTTCTAGTGAATGGAGTTC AACTACCTGCTCTGGGAGAGAA TAAAAAGGTTATAGTTGACTTT TCCTCCCTTAATATAGCTAAAG AGATGCATGTAGGCCACCTGAG GTCACCTATCATAGGAGAGAT ATAAGCCGCCTCTTTGAATTTG CAGGTATGACGTGCTCAGGTT AAATCATGTAGGAGACTGGGG ACCCAGTTTGGCATGCTCATCG CTCACCTGCAAGACAAATTTCC AGATTATCTAACAGTTTCACT CCTATTGGGATCTTCAGGTCT TTTATAAGGAATCTAAGAAAG GTTTGATCTGAGGAGGAAATTT AAGAAGCGAGCATATCAGTGTG TAGTTCTGCTCCAGGGTAAAAA CCCAGATATTACAAAGCTTGG AAGCTTATCTGTGATGCTCCC GCCAAGAGTTAAATAAATCTA TGATGCATTGGACGCTCTCTTA ATAGAGAGAGGGGAATCCTTCT ATCAAGATAGGATGAATGATAT TGTAAGGAATTTGAAGATAGA GGAAAAAGTTTAAACACGTTT GGGTGAACAGTGCGCCCTCATG GATCTTCTGGGAGAGGACTAA	SEQ. ID. NO. 22
ArgRS ^{N4}	Protein/ Human/ 1-160	MDVLVSECSARLLQEEIEKSL TAEIDRLKNCGLGASPNLEQL QEENLKLKYRLNLRKSLQAE NKPTKNMINIISRLQEVFGHAI KAAYPDLENPPLLVTPSQQAKF GDYQCNSAMGISQMLKTKEQKV NPRIAENITKHLPDNECIEKV EIAGPG	SEQ. ID. NO. 23
ArgRS ^{N4}	DNA/ Human	ATGGACGTACTGGTGTCTGAGT GCTCCGCGCGGCTGCTGCAGCA GGAAGAAGAGATTAAATCTCTG	SEQ. ID. NO. 24

TABLE 2-continued

ACTGCTGAAATTGACCGTTGA AAAACTGTGGCTGTTAGGAGC TTCTCCAAATTTGGAGCAGTTA CAAGAAGAAAATTTAAATTA AGTATCGACTGAATATTCTTCG AAAGAGTCTTCAGGCAGAAAGG AACAAACCACTAAAAATATGA TTAACATTATTAGCCGCCTACA AGAGGTCTTTGGTCATGCAATT AAGGCTGCATATCCAGATTTGG AAAATCCTCCTCTGCTAGTGAC ACCAAGTCAGCAGGCCAAGTTT GGGGACTATCAGTGAATAGTG CTATGGGTATTCTCAGATGCT CAAAACCAAGGAACAGAAAGTT AATCCAAGAGAAATTGCTGAAA ACATTACCAACACCTCCAGAA CAATGAATGTATTGAAAAGTT GAAATGTCTGGTCTCGTTAA			
B			
AARS polypeptides unique splice junctions			
Name	Type/ species	Amino acid and Nucleic Acid Sequences in the vicinity of the unique splice junction	SEQ.ID. NO.
R1-AS02	DNA/ Human/	ATTGTAAGGAATTTGAAGATA GAG GAAAAAGTTTAAACACG TTCGGGT	SEQ.ID. NO. 25
	Protein/ Human/	IVKEFEDRGKSLKHVR	SEQ.ID. NO. 26
R1-AS03	DNA/ Human/	GAAAAAGTTGAAATTGCTGGTC CTG GTTAAATCATGTAGGAGA CTGGGGG	SEQ.ID. NO. 27
	Protein/ Human/	EKVEIAGPG	SEQ.ID. NO. 28

TABLE 3

AARS polypeptides and nucleic acids identified by Bioinformatics			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
ArgRS ^{N2}	Protein/ Human/ 1-71	MDVLVSECSARLLQEEIEIKSL TAEIDRLKNCGLGASPNLEQL QEENLKLKYRLNLRKSLQAER NKPTK	SEQ.ID. NO. 29
ArgRS ^{N2}	DNA/ Human/	ATGGACGTACTGGTGTCTGAGT GCTCCGCGCGCTGCTGCAGCA GGAAGAAGAGATTAATCTCTG ACTGCTGAAATTGACCGTTGA AAAAGTGTGGCTGTTAGGAGC TTCTCCAAATTTGGAGCAGTTA CAAGAAGAAAATTTAAATTA AGTATCGACTGAATATTCTTCG AAAGAGTCTTCAGGCAGAAAGG AACAAACCACTAAA	SEQ.ID. NO. 30

C-Terminal AARS Polypeptides: (Tables 4, 5 & 6)

[0105]

TABLE 4

A			
AARS polypeptides identified by MS			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
ArgRS ^{C1}	Protein/ Human/ 73-660	MINIISRLQEVFGHAIKAAAYPD LENPPLLVTSPQQAKFGDYQCN SAMGISQMLKTKEQKVNPREIA ENITKHLDPNECIEKVEIAGPG FINVHLRKDFVSEQLTSLLVNG VQLPALGENKKVIVDFSSPNIA KEMHVGHRLSTIIGESISRLFE FAGYDVLRLNHNVDWGTQFGML IAHLQDKFPDYLTVSPPIGDLQ VFYKESKKRFPDEEFKKRAYQ CVVLLQGKNPDI TKAWKLICDV SRQELNKIYDALDVS LIERGES FYQDRMNDIVKEFEDRGFVQVD DGRKIVFVPGCSIPLTTIVKSDG GYTYDTSDLAAIKQRLFEEKAD MIIYVVDNGQSVHFQTI FAAAA MIGWYDPKVTRV F HAGFVVLG EDKKKFKTRSGETVRLMDLLGE GLKRSMDKLKEKERDKVLTABE LNAAQTSVAYGCIKYADLSHNR LNDYIFSPDKMLDDRGNTAAYL LYAFTRIRSIARLANIDEEMLQ KAARETKILLDHEKEWKLGRCI LRPFEILQKILDDLFLHTLCDY IYELATAFTEFYDSCYCVKEKDR QTGKILKVNMMRWMLLCEAVALV MAKGFIDILGIKPVRM	SEQ.ID. NO. 43
ArgRSC1	DNA/ Human/	ATGATTAACATTATTAGCCGCC TACAAGAGGTCTTTGGTCATGC AATTAAGGCTGCATATCCAGAT TTGGAAATCCTCCTCTGCTAG TGACACCAAGTCAGCAGGCCAA GTTTGGGGACTATCAGTGTAA AGTGTATGGGTATTCTCAGA TGCTCAAAACCAAGGAACAGAA AGTTAATCCAGAGAAATTGCT GAAAACATTACCAACACCTCC CAGACAATGAATGTATTGAAA AGTTGAAATTGCTGGTCTCTGGT TTTATTAATGTCCACTTAAGAA AGGATTTTGTATCAGAACAAAT GACCAGTCTTCTAGTGAATGGA GTTCAACTACCTGCTCTGGGAG AGAATAAAAAGGTTATAGTTGA CTTTTCCTCCCTAATATAGCT AAAGAGATGCATGTAGGCCACC TGAGGTCACTATCATAGGAGA GAGTATAAGCCGCCTCTTTGAA TTTGCAGGGTATGACGTGCTCA GGTTAAATCATGTAGGAGACTG GGGGACCCAGTTTGGCATGCTC ATCGCTCACCTGCAAGACAAAT TTCCAGATTATCTAACAGTTTC ACCTCCTATTGGGGATCTTCAG GTCTTTTATAAGGAATCTAAGA AGAGGTTTGATCTGAGGAGGA ATTTAAGAAGCGAGCATATCAG TGTGTAGTTCTGCTCCAGGGTA AAAACCCAGATATTACAAAAGC TTGGAAGCTTATCTGTGATGTC TCCCGCCAAGAGTTAAATAAAA TCTATGATGCATTGGACGTCTC TTTAATAGAGAGAGGGGAATCC TTCTATCAAGATAGGATGAATG ATATTGTAAGGAATTTGAAGA	SEQ.ID. NO. 44

TABLE 4-continued

TAGAGGATTTGTGCAGGTGGAT		
GATGGCAGAAAGATTGTATTTG		
TCCCAGGGTGTTCCATACCATT		
AACCATAGTAAAAATCAGATGGA		
GGTTATACCTATGATACATCTG		
ACCTGGCTGCTATTAAACAAAG		
ACTATTTGAGGAAAAAGCAGAT		
ATGATTATCTATGTTGTGGACA		
ATGGACAATCTGTGCACTTCCA		
GACAATATTTGCTGCTGCTCAA		
ATGATTGGTTGGTATGACCTTA		
AAGTAACTCGAGTCTTCCATGC		
TGGATTTGGTGTGGTCTAGGG		
GAAGACAAGAAAAAGTTAAAA		
CACGTTCCGGGTGAAACAGTGCG		
CCTCATGGATCTTCTGGGAGAA		
GGACTAAACGATCCATGGACA		
AGTTGAAGGAAAAAGAAAGAGA		
CAAGGTCTTAACGCAGAGGAA		
TTGAATGCTGCTCAGACATCCG		
TTGCGTATGGCTGCATCAATA		
TGCTGACCTTTCCATAACCGG		
TTGAATGACTACATCTTCTCCT		
TTGACAAAATGCTAGATGACAG		
AGGAAATACAGCTGCTTACTTG		
TTGTATGCCTTCACTAGATCA		
GGTCTATTGCACGCTCTGGCCAA		
TATTGATGAAGAAATGCTCCAA		
AAAGCTGCTCGAGAAACCAAGA		
TTCTTTTGATCATGAGAAGGA		
ATGGAACTAGGCCGGTGCAAT		
TTACGGTTCCTGAGATTCTGC		
AAAAGATTTTAGATGACTTATT		
TCTCCACACTCTCTGTGATTAT		
ATATATGAGCTGGCAACTGCTT		
TCACAGAGTTCTATGATAGCTG		
CTACTGTGTGGAGAAAGATAGA		
CAGACTGGAAAAATATTGAAGG		
TGAACATGTGGCGTATGCTGCT		
ATGTGAAGCAGTAGCTGCTGTC		
ATGGCCAAGGGTTTGATATCC		
TGGGAATAAAACCTGTCCAAG		
GATGTAA		

B
ArgRS1^{C1}
Mass spec peptides detected
and inferred linking peptides

Type/ species	Sequence	SEQ. ID. NO.
Protein/ mouse	LQEVFGCAIR	SEQ. ID. NO. 45
Protein/ mouse	AAYPDLENPPLIVTPSQQPK	SEQ. ID. NO. 46
Protein/ mouse	FGDYQCN	SEQ. ID. NO. 47
Protein/ mouse	SAMGISQMLKAKEQKVS PREIAENITKHL PNNKYIDKVEIAGPGFINVHLRKDFVSEQ LTSLLVNGVQLPVLGDKKEK	SEQ. ID. NO. 48
Protein/ mouse	VIVDFSSPNIAK	SEQ. ID. NO. 49
Protein/ mouse	EMHVGHLRSTIIGESMSR	SEQ. ID. NO. 50
Protein/ mouse	LFEFAGYDVLK	SEQ. ID. NO. 51
Protein/ mouse	LNHVGDWGTQFGMLIAHLQDK	SEQ. ID. NO. 52

TABLE 4-continued

Protein/ mouse	FPDYLTVSPPIGDLQAFYK	SEQ. ID. NO. 53
Protein/ mouse	ESKKR	SEQ. ID. NO. 54
Protein/ mouse	FDADEEFKKRAYQCVVLLQSK	SEQ. ID. NO. 55
Protein/ mouse	NPDIMK	SEQ. ID. NO. 56
Protein/ mouse	AWNLICDVSR	SEQ. ID. NO. 57
Protein/ mouse	EEFK	SEQ. ID. NO. 58
Protein/ mouse	KIYDALDITLIER	SEQ. ID. NO. 59
Protein/ mouse	GESFYQDRMKDIVKEFEDKGFVQVDDGRK	SEQ. ID. NO. 60
Protein/ mouse	IVFVPGCSVPLTIVKSDGGYTYDTSDLAA IK	SEQ. ID. NO. 61
Protein/ mouse	QRLFEEKANKIIYVVDNGQAIHFQTIFAA AQMIGWYDPKVTLVTHVGFVVLGEDKKK FKTRSGETVRLMDLLEELKRSMDKLKEK ER	SEQ. ID. NO. 62
Protein/ mouse	DKVLTEEEKL	SEQ. ID. NO. 63
Protein/ mouse	AAQTSVAYGCIK	SEQ. ID. NO. 64
Protein/ mouse	YADLSHNRLNDYIFSPDKMLDDR	SEQ. ID. NO. 65
Protein/ mouse	GNTAAYLLYAFTR	SEQ. ID. NO. 66
Protein/ mouse	IRSIAR	SEQ. ID. NO. 67
Protein/ mouse	LANIDEAMLQR	SEQ. ID. NO. 68
Protein/ mouse	AARETKIILDHEKEWKLGRCILRFPEILQ KILDDLFLHTLCDYIYELATTTFEYDSC YCVEKDRQTGKVLKVNMR	SEQ. ID. NO. 69
Protein/ mouse	MLLCEAAVAVMAK	SEQ. ID. NO. 70
Protein/ mouse	GFDILGIKPVQR	SEQ. ID. NO. 71
Protein/ mouse	AAYPDLENPPLIVTPSQQPK	SEQ. ID. NO. 72
Protein/ mouse	FGDYQCNSAMGISQMLKAKEQKVS PREIA ENITKHL PNNKYIDKVEIAGPGFINVHLR K	SEQ. ID. NO. 73
Protein/ mouse	DFVSEQLTSLVNGVQLPVLGDKKEK	SEQ. ID. NO. 74
Protein/ mouse	VIVDFSSPNIAK	SEQ. ID. NO. 75
Protein/ mouse	EMHVGHLRSTIIGESMSR	SEQ. ID. NO. 76

TABLE 4-continued

Protein/ mouse	LFEFAGYDVLR	SEQ.ID. NO. 77
Protein/ mouse	LNHVGDWGTQFGMLIAHLQDK	SEQ.ID. NO. 78
Protein/ mouse	FPDYLTVPPIGDLQAFYK	SEQ.ID. NO. 79
Protein/ mouse	ESKKRFDADEEFKKRAYQCVVLLQSKNPD IMK	SEQ.ID. NO. 80
Protein/ mouse	AWNLICDVSR	SEQ.ID. NO. 81
Protein/ mouse	EEFKKIYDALDITLIERGESFYQDRMKDI VKEFEDKGFVQVDDGRK	SEQ.ID. NO. 82
Protein/ mouse	IVFVPGCSVPLTIVKSDGGYTYDTSDLAA IK	SEQ.ID. NO. 83
Protein/ mouse	QRLFEEKANKIIYVVDNGQAIHFQTIFAA AQMIGWYDPKVTLVTHVGFVVLGEDKKK FKTRSGETVRLMDLLEEGLRKSRMDKLKEK ER	SEQ.ID. NO. 84
Protein/ mouse	DKVLTEEEKL	SEQ.ID. NO. 85
Protein/ mouse	AAQTSVAYGCIKYADLSHNRLNDYIFSFD KMLDDR	SEQ.ID. NO. 86
Protein/ mouse	GNTAAYLLYAFTR	SEQ.ID. NO. 87
Protein/ mouse	IRSIAR	SEQ.ID. NO. 88
Protein/ mouse	LANIDEAMLQR	SEQ.ID. NO. 89
Protein/ mouse	AARETK	SEQ.ID. NO. 90
Protein/ mouse	IILDHEKEWK	SEQ.ID. NO. 91
Protein/ mouse	LGRCILRFPEILQKILDDLFLHTLCDYIY ELATTTFEYDSCYCYVEKDRQTGKVLKVN MWR	SEQ.ID. NO. 92
Protein/ mouse	MLLCEAVALAVMAK	SEQ.ID. NO. 93
Protein/ mouse	LQEVFGCAIRAAYPDLENPPLIVTPSQQP KFGDYQCNSAMGISQMLK	SEQ.ID. NO. 94
Protein/ mouse	AKEQKVS PREIAENITKHLPPNNK	SEQ.ID. NO. 95
Protein/ mouse	YIDKVEIAGPGFINVHLR	SEQ.ID. NO. 96
Protein/ mouse	DFVSEQLTSLLVNGVQLPVLGDKEKVIVD FSSPNIK	SEQ.ID. NO. 97
Protein/ mouse	EMHVGHLRSTIIGESMSR	SEQ.ID. NO. 98
Protein/ mouse	LFEFAGYDVLR	SEQ.ID. NO. 99
Protein/ mouse	LNHVGDWGTQFGMLIAHLQDK	SEQ.ID. NO. 100

TABLE 4-continued

Protein/ mouse	FPDYLTVPPIGDLQAFYK	SEQ.ID. NO. 101
Protein/ mouse	ESKKRFDADEEFKKR	SEQ.ID. NO. 102
Protein/ mouse	AYQCVVLLQSK	SEQ.ID. NO. 103
Protein/ mouse	NPDIMK	SEQ.ID. NO. 104
Protein/ mouse	AWNLICDVSR	SEQ.ID. NO. 105
Protein/ mouse	EEFK	SEQ.ID. NO. 106
Protein/ mouse	IYDALDITLIER	SEQ.ID. NO. 107
Protein/ mouse	GESFYQDRMKDIVK	SEQ.ID. NO. 108
Protein/ mouse	EFEDKGFVQVDDGR	SEQ.ID. NO. 109
Protein/ mouse	KIVFVPGCSVPLTIVKSDGGYTYDTSDLA AIK	SEQ.ID. NO. 110
Protein/ mouse	QRLFEEKANKIIYVVDNGQAIHFQTIFAA AQMIGWYDPKVTLVTHVGFVVLGEDKKK FKTRSGETVR	SEQ.ID. NO. 111
Protein/ mouse	LMDLLEEGLRK	SEQ.ID. NO. 112
Protein/ mouse	SMDKLKEKER	SEQ.ID. NO. 113
Protein/ mouse	DKVLTEEEKL	SEQ.ID. NO. 114
Protein/ mouse	AAQTSVAYGCIK	SEQ.ID. NO. 115
Protein/ mouse	YADLSHNR	SEQ.ID. NO. 116
Protein/ mouse	LN DYIFSFDK	SEQ.ID. NO. 117
Protein/ mouse	MLDDR	SEQ.ID. NO. 118
Protein/ mouse	GNTAAYLLYAFTR	SEQ.ID. NO. 119
Protein/ mouse	IRSIAR	SEQ.ID. NO. 120
Protein/ mouse	LANIDEAMLQR	SEQ.ID. NO. 121
Protein/ mouse	AARETK	SEQ.ID. NO. 122
Protein/ mouse	IILDHEKEWK	SEQ.ID. NO. 123
Protein/ mouse	LGRCILRFPEILQK	SEQ.ID. NO. 124
Protein/ mouse	ILDDLFLHTLC	SEQ.ID. NO. 125

TABLE 4-continued

Protein/ mouse	DYIYELATTFTEFYDSCYVEKDRQTGKV LKVNMMWR	SEQ. ID. NO. 126
Protein/ mouse	MLLCEAVAAMAKGFDILGIKPVQR	SEQ. ID. NO. 127
C ArgRS1^{C1} Concatenated sequences based on mass spec peptides detected		
Type/ species	Sequence	SEQ. ID. NO.
Protein/ mouse	<u>LQEVFGCAIRAAYPDLENPPLIVTPSQOP</u> <u>KFGDYQCNSAMGISQMLKAKEQKVS PREI</u> <u>AENITKHLNNKYIDKVEIAGPGFINVHL</u> <u>RKDFVSEQLTSLLVNGVQLPVLGDKEKVI</u> <u>VDFSSPNIAKEMHVGHIRSTIIGESMSRL</u> <u>FEFAGYDVLRLNHVGDWGTQFGMLIAHLQ</u> <u>DKFPDYLTVSPPIGDLQAFYKESKKRFDA</u> <u>DEEFKKRAYQCVVLLQSKNPDIMKAWNLI</u> <u>CDVSREEFKKIYDALDITLIERGESFYQD</u> <u>RMKDIVKEFEDKGFVQVDDGRKIVFVPGC</u> <u>SVPLTIVKSDGGYTYDTSDLAAIKQRLFE</u> <u>EKANKI IYVVDNGQAIHFQTI FAAAQMI</u> <u>GYDPKVTLVTHVGFVVLGDKKKFKTRS</u> <u>GETVRLMDLLEGLKRSMDKLKEKERDKV</u> <u>LTEEELKAAQTSVAYGCIKYADLSHNRLN</u> <u>DYIFSFDKMLDDRGNATAAYLLYAFTRIRS</u> <u>IARLANIDEAMLQRAARETKI LDHEKEW</u> <u>KLGRCILRFPEILQKILDDLFLHTLCDYI</u> <u>YELATTFTEFYDSCYVEKDRQTGKVLK</u> <u>NMWRMLLCEAVAAMAKGFDILGIKPVQR</u>	SEQ. ID. NO. 128
Protein/ mouse	<u>AAYPDLENPPLIVTPSQOPKFGDYQCNSA</u> <u>MGISQMLKAKEQKVS PREIAENITKHLPN</u> <u>NKYIDKVEIAGPGFINVHLRKDFVSEQLT</u> <u>SLLVNGVQLPVLGDKEKVI VDFSSPNIAK</u> <u>EMHVGHIRSTIIGESMSRLFEFAGYDVLRL</u> <u>LNHVGDWGTQFGMLIAHLQDKFPDYLTVS</u> <u>PPIGDLQAFYKESKKRFDADEEFKKRAYQ</u> <u>CVVLLQSKNPDIMKAWNLI CDVSREEFKK</u> <u>IYDALDITLIERGESFYQDRMKD IVKEFE</u> <u>DKGFVQVDDGRKIVFVPGCSVPLTIVKSD</u> <u>GGYTYDTSDLAAIKQRLFEEKANKI IYV</u> <u>DNGQAIHFQTI FAAAQMI GWDPKVTLV</u> <u>HVGFVVLGDKKKFKTRSGETVRLMDLL</u> <u>EEGLKRSMDKLKEKERDKVLTTEEELKAAQ</u> <u>TSVAYGCIKYADLSHNRLNDYIFSFDKML</u> <u>DDRGNATAAYLLYAFTRIRS IARLANIDEA</u> <u>MLQRAARETKI LDHEKEWKLGRCILRFP</u> <u>EILQKILDDLFLHTLCDYIYELATTTTEF</u> <u>YDSCYVEKDRQTGKVLKVNMMWRMLLCEA</u> <u>VAAVMAK</u>	SEQ. ID. NO. 129
Protein/ mouse	<u>LQEVFGCAIRAAYPDLENPPLIVTPSQOP</u> <u>KFGDYQCNSAMGISQMLKAKEQKVS PREI</u> <u>AENITKHLNNKYIDKVEIAGPGFINVHL</u> <u>RKDFVSEQLTSLLVNGVQLPVLGDKEKVI</u> <u>VDFSSPNIAKEMHVGHIRSTIIGESMSRL</u> <u>FEFAGYDVLRLNHVGDWGTQFGMLIAHLQ</u> <u>DKFPDYLTVSPPIGDLQAFYKESKKRFDA</u> <u>DEEFKKRAYQCVVLLQSKNPDIMKAWNLI</u> <u>CDVSREEFKKIYDALDITLIERGESFYQD</u> <u>RMKDIVKEFEDKGFVQVDDGRKIVFVPGC</u> <u>SVPLTIVKSDGGYTYDTSDLAAIKQRLFE</u> <u>EKANKI IYVVDNGQAIHFQTI FAAAQMI</u> <u>GYDPKVTLVTHVGFVVLGDKKKFKTRS</u> <u>GETVRLMDLLEGLKRSMDKLKEKERDKV</u> <u>LTEEELKAAQTSVAYGCIKYADLSHNRLN</u> <u>DYIFSFDKMLDDRGNATAAYLLYAFTRIRS</u> <u>IARLANIDEAMLQRAARETKI LDHEKEW</u> <u>KLGRCILRFPEILQKILDDLFLHTLCDYI</u> <u>YELATTFTEFYDSCYVEKDRQTGKVLK</u> <u>NMWRMLLCEAVAAMAKGFDILGIKPVQR</u>	SEQ. ID. NO. 130

TABLE 5

AARS polypeptides and alternative transcripts identified by Deep Sequencing			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ. ID. NO.
ArgRS ^{C3}	Protein/ Human/ 463-660	MDLLGEGLKRSMDKLKEKERDK VLTAEEELNAAQTSVAYGCIKYA DLSHNRLNDYIFSFDKMLDDRGN NTAAYLLYAFTRIRS IARLANI DEEMLQKAARETKILLDHEKEW KLGRCILRFPEILQKILDDLFL HTLCDYIYELATAFTEFYDSCY VEKDRQTGKILKVNMMWRMLLC EAVAAMAKGFDILGIKPVQRM	SEQ. ID. NO. 131
ArgRS ^{C3}	DNA/ Human	ATGGATCTTCTGGGAGAAGGAC TAAACGATCCATGGACAAGTT GAAGGAAAAAGAGACAAAG GTCTTAAGTCAGAGGAATTGA ATGCTGCTCAGACATCCGTTGC GTATGGCTGCATCAATATGCT GACCTTTCCCATACCGGTTGA ATGACTACATCTTCTCCTTGA CAAAATGCTAGATGACAGAGGA AATACAGCTGCTTACTTGTGT ATGCCTTCACTAGATCAGGTC TATTGCAGCTGCGCAATAT GATGAAGAAATGCTCCAAAAAG CTGCTCGAGAAACCAAGATTCT TTTGATCATGAGAGGAATGG AACTAGGCCGTCATTTTAC GGTTCCTGAGATTCTGCAAAA GATTTTAGATGACTTATTCTC CACACTCTCTGTGATTATATAT ATGAGCTGGCACTGCTTTTAC AGAGTTCTATGATAGCTGCTAC TGTGTGGAGAAAGATAGACAGA CTGGAAAAATATGAAGGTGAA CATGTGGCGTATGCTGCTATGT GAAGCAGTAGCTGCTGCTATGG CCAAGGGTTTGATATCCTGGG AATAAACTGTCCAAAGGATG TAA	SEQ. ID. NO. 132
ArgRS ^{C4}	Protein/ Human/ 247-660	MLIAHLQDKFPDYLTVSPPIGD LQVYFYESKKRFDTEEFKKRA YQCVVLLQGKNPDIKAWKLCI DVSRLQELNKIYDALDVSLIERG ESFYQDRMNDIVKEFEDRGFVQ VDDGRKIVFVPGCSIPLTIKVS DGGYTYDTSDLAAIKQRLFEEK ADMIIYVVDNGQSVHFQTI FAA AQMIQWYDPKVTRVFHAGFGV LGEDKKKFKTRSGETVRLMDLL GEGLKRSMDKLKEKERDKVLT EELNAAQTSVAYGCIKYADLSH NRLNDYIFSFDKMLDDRGNATA YLLYAFTRIRS IARLANIDEEM LQKAARETKI LDHEKEWKLGR CILRFPEILQKILDDLFLHTLC DYIYELATAFTEFYDSCYVEK DRQTGKILKVNMMWRMLLCEA VAMAKGFDILGIKPVQRM	SEQ. ID. NO. 133
ArgRS ^{C4}	DNA/ Human	ATGCTCATCGCTCACCTGCAAG ACAAATTTCCAGATTATCTAAC AGTTTACCTCCTATTGGGGAT CTTCAGGTCTTTTATAAGGAAT CTAAGAAAGAGGTTTGATCTGA GGAGGAATTTAAGACGAGCAG TATCAGTGTGTAGTCTGTCTCC AGGGTAAAAACCCAGATATTAC AAAAGCTTGAAGCTTATCTGT GATGTCTCCCGCCAGAGTTAA ATAAATCTATGATGCATTGGA	SEQ. ID. NO. 134

TABLE 5-continued

CGTCTCTTTAATAGAGAGAGGG GAATCCTTCTATCAAGATAGGA TGAATGATATTGTAAAGGAATT TGAAGATAGAGGATTGTGCAG GTGGATGATGGCAGAAAGATTG TATTTGTCCCAGGGTGTTCAT ACCATTAACCATAGTAAAAATCA GATGGAGGTTATACCTATGATA CATCTGACCTGGCTGCTATTAA ACAAAGACTATTGAGGAAAAA GCAGATATGATTATCTATGTTG TGGACAATGGACAATCTGTGCA CTTCCAGACAATATTGCTGCT GCTCAATGATTGGTTGGTATG ACCCATAAGTAACTCGAGTCTT CCATGCTGGATTGGTGTGGTG CTAGGGGAAGACAAGAAAAAGT TTAAACACGTTCCGGTGAAAC AGTGCGCCTCATGGATCTTCTG GGAGAAGGACTAAAACGATCCA TGGACAAGTTGAAGGAAAAAGA AAGAGACAAGGTCTTAAGTGCA GAGGAATTGAATGCTGCTCAGA CATCCGTTGCGTATGGCTGCAT CAAAATATGCTGACCTTTCCCAT AACCAGTTGAATGACTACATCT TCTCCTTTGACAAAATGCTAGA TGACAGAGGAAATACAGCTGCT TACTTGTGTATGCCTTCACTA GAATCAGGTCTATTGCACGCT GGCCAATATTGATGAAGAAATG CTCCAAAAGCTGCTCGAGAAA CCAAGATTCTTTGGATCATGA GAAGGAATGGAACCTAGGCCGG TGCATTTTACGGTTCCTGAGA TTCTGCAAAAGATTTAGATGA CTTATTTCTCCACTCTCTGT GATTATATATGAGCTGGCAA CTGCTTTCACAGAGTCTATGA TAGCTGCTACTGTGTGGAGAAA GATAGACAGACTGGAATAATAT TGAAGGTGAACATGTGGCGTAT GCTGCTATGTGAAGCAGTAGCT GCTGTCATGGCCAGGGGTTTG ATATCCTGGGAATAAACCTGT CCAAAGGATGTAA			
B AARS polypeptides unique splice junctions			
Name	Type/ species	Amino acid and Nucleic Acid Sequences in the vicinity of the unique splice junction	SEQ.ID. NO.
R1- AS02	DNA/ Human/ Protein/ Human/	ATTGTAAAGGAATTGAAGATA GAG GAAAAAGTTTAAACACG TTCGGGT	SEQ.ID. NO. 135
R1- AS03	DNA/ Human/ Protein/ Human/	GAAAAAGTTGAAATGCTGGTC CTG GTTAAATCATGTAGGAGA CTGGGGG	SEQ.ID. NO. 136

TABLE 6

AARS polypeptides and nucleic acids identified by Bioinformatics			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
ArgRS ^{C2}	Protein/ Human/ 551-660	DEEMLQKAARETKILLDHEKEW KLGRCILRFPEILQKILLDDLFL HTLCDYIYELATAFTEFYDSCY CVEKDRQTGKILKVNMMWRMLLC EAVAAMAKGFDILGIKPVQRM	SEQ.ID. NO. 137
ArgRS ^{C2}	DNA/ Human/	GATGAAGAAATGCTCCAAAAAG CTGCTCGAGAAACCAAGATTCT TTTGGATCATGAGAAGGAATGG AAACTAGGCCGGTGCATTTTAC GGTTCCCTGAGATTCTGCAAAA GATTTTAGATGACTTATTCTC CACACTCTCTGTGATTATATAT ATGAGCTGGCAACTGCTTTTAC AGAGTTCTATGATAGCTGCTAC TGTGTGGAGAAAGATAGACAGA CTGGAAAAATATTGAAGGTGAA CATGTGGCGTATGCTGCTATGT GAAGCAGTAGCTGCTGCTATGG CCAAGGGGTTGATATCCTGGG AATAAAACCTGTCCAAGGATG TAA	SEQ.ID. NO. 138

Internal AARS Polypeptides: (Tables 7, 8 & 9)

[0106]

TABLE 7

A AARS polypeptides identified by MS			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
ArgRS ^{I1}	Protein/ Human/ 168-530	KDFVSEQLTSLVNGVQLPALG ENKKVIVDFSSPNIKEMHVGH LRSTIIGESISRLFEFAGYDVL RLNHVGDWGTQFGLIAHLQDK FPDYLTVSPPIGDLQVFYKESK KRFDTEEEFKKRAYQCVLLQG KNPDITKAWKLCVSRQELNK IYDALDVSLIERGESFYQDRMN DIVKEFEDRGFVQVDDGRKIVF VPGCSIPLTIVKSDGGYTYDTS DLAAIKQRLFEKADMIYVVD NGQSVHFQTI FAAAQMIGWYDP KVTRVVFHAGFGVVLGEDKKKPK TRSGETVRLMDLLGEGLKRSMD KLKEKERDKVLTAEELNAAQTS VAYGCIKYADLSHNLNDYIFS FDMKLLDRGNT	SEQ.ID. NO. 151
ArgRS ^{I1}	DNA/ Human/	AAGGATTTTGTATCAGAACAAT TGACCAGTCTTCTAGTGAATGG AGTTCAACTACCTGCTCTGGGA GAGAATAAAAAGGTTATAGTTG ACTTTTCTCCCTAATATAGC TAAAGAGATGCATGTAGGCCAC CTGAGGTCAACTATCATAGGAG AGAGTATAAGCCGCTCTTTGA ATTTGCAGGGTATGACGTGCTC AGGTTAAATCATGTAGGAGACT GGGGGACCCAGTTTGGCATGCT CATCGCTCACCTGCAAGACAAA TTTCCAGATTATCTAACAGTTT	SEQ.ID. NO. 152

TABLE 7-continued

A			
AARS polypeptides identified by MS			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
		CACCTCCTATTGGGGATCTTCA GGTCTTTTATAAGGAATCTAAG AAGAGGTTTGATACTGAGGAGG AATTTAAGAAGCGAGCATATCA GTGTGTAGTTCTGCTCCAGGGT AAAAACCCAGATATTACAAAAG CTTGGAAGCTTATCTGTGATGT CTCCCGCCAAGAGTTAAATAAA ATCTATGATGCATTGGACGTCT CTTTAATAGAGAGAGGGGAATC CTTCTATCAAGATAGGATGAAT GATATTGTAAAGGAATTTGAAG ATAGAGGATTTGTGCAGGTGGA TGATGGCAGAAAGATTGTATTT GTCCCAGGGTGTTCATACCAT TAACCATAGTAAATCAGATGG AGGTTATACCTATGATACATCT GACCTGGCTGCTATTAAACAAA GACTATTTGAGGAAAAAGCAGA TATGATTATCTATGTTGTGGAC AATGGACAATCTGTGCACTTCC AGACAATATTTGCTGCTGCTCA AATGATTGGTTGGTATGACCCCT AAAGTAACTCGAGTCTTCCATG CTGGATTTGGTGTGGTGTAGG GGAAGACAAGAAAAAGTTTAA ACACGTTCCGGTGAAACAGTGC GCCTCATGGATCTTCTGGGAGA AGGACTAAACGATCCATGGAC AAGTTGAAGGAAAAAGAAAGAG ACAAGGTCTTAACTGCAGAGGA ATTGAATGCTGCTCAGACATCC GTTGCGTATGGCTGCATCAAAT ATGCTGACCTTCCCATAACCG GTTGAATGACTACATCTTCTCC TTTGACAAAATGCTAGATGACA GAGGAAATACA	

B		
ArgRS ^{fl}		
Mass spec peptides detected and inferred linking peptides		
Type/ species	Sequence	SEQ.ID. NO.
Protein/ mouse	VIVDFSSPNIAK	SEQ.ID. NO. 153
Protein/ mouse	EMHVGHLRSTIIGESMSRLFEFAGYDVLR LNHVGDWGTQFGMLIAHLQDKFPDYLTVS PPIGDLQAFYKESKKRFDADDEEFKKRAYQ CVVLLQSKNPDIMKAWNLICDVSREEFKK IYDALDITLIERGESFYQDRMKDIVKEFE DKGFVQVDDGRKIVFVPGCSVPLTIVK	SEQ.ID. NO. 154
Protein/ mouse	SDGGYTYDSDLAAIK	SEQ.ID. NO. 155

C		
ArgRS ^{fl}		
Concatenated sequences based on mass spec peptides detected		
Type/ species	Sequence	SEQ.ID. NO.
Protein/ mouse	VIVDFSSPNIAKEMHVGHLRSTIIGESMS RLFEFAGYDVLRNLNVGDWGTQFGMLIAH LQDKFPDYLTVSPPIGDLQAFYKESKKRF	SEQ.ID. NO. 156

TABLE 7-continued

A			
AARS polypeptides identified by MS			
Name	Type/ species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ.ID. NO.
		DADEEFKKRAYQCVVLLQSKNPDIMKAWN LICDVSREEFKKIYDALDITLIERGESFY QDRMKDIVKEFEDKGFVQVDDGRKIVFVP GCSVPLTIVKSDGGYTYDSDLAAIK	

TABLE 8

AARS polypeptides and alternative transcripts identified by Deep Sequencing			
Name	Type / species / Residues	Amino acid and Nucleic Acid Sequences	SEQ. ID. NO.
②			
② indicates text missing or illegible when filed			

TABLE 8B

AARS polypeptides unique splice junctions			
Name	Type / species / Residues	Amino acid and Nucleic Acid Sequences in the vicinity of the unique splice junction	SEQ. ID. NO.
②			
② indicates text missing or illegible when filed			

TABLE 9

AARS polypeptides and nucleic acids identified by Bioinformatics			
Name	Type / species/ Residues	Amino acid and Nucleic Acid Sequences	SEQ. ID. NO.
②			
② indicates text missing or illegible when filed			

[0107] “Protein fragments,” or the amino acid sequence of protein fragments, such as proteolytic fragments or splice variant fragments, can be characterized, identified, or derived according to a variety of techniques. For instance, splice variants can be identified by techniques such as deep sequencing (see, e.g., Xing et al., *RNA*. 14:1470-1479, 2008; and Zhang et al., *Genome Research*. 17:503-509, 2007). As a further example, protein fragments such as proteolytic fragments can be identified in vitro, such as by incubating full-length or other AARS polypeptides with selected proteases, or they can be identified endogenously (e.g., in vivo). In certain embodiments, protein fragments such as endogenous proteolytic fragments can be generated or identified, for instance, by recombinantly expressing full-length or other AARS polypeptides in a selected microorganism or eukaryotic cell that has been either modified to contain one or more selected proteases, or that naturally contains one or more proteases that are capable of acting on a selected

AARS polypeptide, and isolating and characterizing the endogenously produced protein fragments therefrom.

[0108] In certain embodiments, protein fragments such as endogenous (e.g., naturally-occurring) proteolytic fragments can be generated or identified, for instance, from various cellular fractions (e.g., cytosolic, membrane, nuclear) and/or growth medium of various cell-types, including, for example, immune cells such as monocytes, dendritic cells, macrophages (e.g., RAW 264.7 macrophages), neutrophils, eosinophils, basophils, and lymphocytes, such as B-cells and T-cells (e.g., CD4+ helper and CD8+ killer cells), including primary T-cells and T-cell lines such as Jurkat T-cells, as well as natural killer (NK) cells.

[0109] In certain embodiments, protein fragments such as endogenous proteolytic fragments, however generated, can be identified by techniques such as mass-spectrometry, or equivalent techniques. Once an in vitro or endogenously identified protein fragment has been generated or identified, it can be mapped or sequenced, and, for example, cloned into an expression vector for recombinant production, or produced synthetically.

[0110] A wide variety of proteases can be used to produce, identify, derive, or characterize the sequence of AARS protein fragments such as proteolytic fragments. Generally, proteases are usually classified according to three major criteria: (i) the reaction catalyzed, (ii) the chemical nature of the catalytic site, and (iii) the evolutionary relationship, as revealed by the structure. General examples of proteases or proteinases, as classified by mechanism of catalysis, include aspartic proteases, serine proteases, cysteine proteases, and metalloproteases.

[0111] Most aspartic proteases belong to the pepsin family. This family includes digestive enzymes, such as pepsin and chymosin, as well as lysosomal cathepsins D and processing enzymes such as renin, and certain fungal proteases (e.g., penicillopepsin, rhizopuspepsin, endothiapepsin). A second family of aspartic proteases includes viral proteinases such as the protease from the AIDS virus (HIV), also called retropepsin.

[0112] Serine proteases include two distinct families. First, the chymotrypsin family, which includes the mammalian enzymes such as chymotrypsin, trypsin, elastase, and kallikrein, and second, the subtilisin family, which includes the bacterial enzymes such as subtilisin. The general 3D structure between these two families is different, but they have the same active site geometry, and catalysis proceeds via the same mechanism. The serine proteases exhibit different substrate specificities, differences which relate mainly to amino acid substitutions in the various enzyme subsites (substrate residue interacting sites). Some serine proteases have an extended interaction site with the substrate whereas others have a specificity that is restricted to the P1 substrate residue.

[0113] The cysteine protease family includes the plant proteases such as papain, actinidin, and bromelain, several mammalian lysosomal cathepsins, the cytosolic calpains (calcium-activated), as well as several parasitic proteases (e.g., *Trypanosoma*, *Schistosoma*). Papain is the archetype and the best studied member of the family. Recent elucidation of the X-ray structure of the Interleukin-1-beta Converting Enzyme has revealed a novel type of fold for cysteine proteinases.

[0114] The metalloproteases are one of the older classes of proteases, found in bacteria, fungi, and higher organisms.

They differ widely in their sequences and their 3D structures, but the great majority of enzymes contain a zinc atom that is catalytically active. In some cases, zinc may be replaced by another metal such as cobalt or nickel without loss of proteolytic activity. Bacterial thermolysin has been well characterized and its crystallographic structure indicates that zinc is bound by two histidines and one glutamic acid. Many metalloproteases contain the sequence motif HEXXH, which provides two histidine ligands for the zinc. The third ligand is either a glutamic acid (thermolysin, neprilysin, alanyl aminopeptidase) or a histidine (astacin, serralsin).

[0115] Illustrative proteases include, for example, achromopeptidase, aminopeptidase, ancrod, angiotensin converting enzyme, bromelain, calpain, calpain I, calpain II, carboxypeptidase A, carboxypeptidase B, carboxypeptidase G, carboxypeptidase P, carboxypeptidase W, carboxypeptidase Y, caspase 1, caspase 2, caspase 3, caspase 4, caspase 5, caspase 6, caspase 7, caspase 8, caspase 9, caspase 10, caspase 11, caspase 12, caspase 13, cathepsin B, cathepsin C, cathepsin D, cathepsin E, cathepsin G, cathepsin H, cathepsin L, chymopapain, chymase, chymotrypsin, clostripain, collagenase, complement C1r, complement C1s, complement Factor D, complement factor I, cucumisin, dipeptidyl peptidase IV, elastase (leukocyte), elastase (pancreatic), endoproteinase Arg-C, endoproteinase Asp-N, endoproteinase Glu-C, endoproteinase Lys-C, enterokinase, factor Xa, ficin, furin, granzyme A, granzyme B, HIV Protease, IGase, kallikrein tissue, leucine aminopeptidase (general), leucine aminopeptidase (cytosol), leucine aminopeptidase (microsomal), matrix metalloprotease, methionine aminopeptidase, neutrase, papain, pepsin, plasmin, prolidase, pronase E, prostate specific antigen, protease alkalophilic from *Streptomyces griseus*, protease from *Aspergillus*, protease from *Aspergillus saitoi*, protease from *Aspergillus sojae*, protease (*B. licheniformis*) (alkaline or alcalase), protease from *Bacillus polymyxa*, protease from *Bacillus* sp., protease from *Rhizopus* sp., protease S, proteasomes, proteinase from *Aspergillus oryzae*, proteinase 3, proteinase A, proteinase K, protein C, pyroglutamate aminopeptidase, rennin, rennin, streptokinase, subtilisin, thermolysin, thrombin, tissue plasminogen activator, trypsin, tryptase and urokinase.

[0116] Certain embodiments relate to isolated AARS polypeptides, comprising, consisting essentially of, or consisting of amino acid sequences that have been derived from endogenous, naturally-occurring AARS polypeptide fragments, and pharmaceutical compositions comprising said fragments, and methods of use thereof. These and related embodiments can be generated or identified in vivo, ex vivo, and/or in vitro. In certain preferred in vitro embodiments, AARS proteolytic fragments are generated or identified by incubating an AARS polypeptide, such as a full-length AARS polypeptide, with one or more isolated human proteases, mainly those proteases that are endogenous or natural to humans, such as elastase and others described herein and known in the art. Other embodiments relate to isolated AARS polypeptides, comprising, consisting essentially of, or consisting of amino acid sequences that have been derived from endogenous, naturally-occurring AARS splice variants, and pharmaceutical compositions comprising said fragments, and methods of use thereof. Essentially, AARS protein fragment can be isolated from samples that have been exposed to proteases, whether in vivo or in vitro.

[0117] In certain embodiments, AARS protein fragments can be identified by techniques such as mass-spectrometry, or equivalent techniques. Merely by way of illustration and not limitation, in certain embodiments the proteomes from various cell types, tissues, or body fluids from a variety of physiological states (e.g., hypoxia, diet, age, disease) or fractions thereof may be separated by 1D SDS-PAGE and the gel lanes cut into bands at fixed intervals; after which the bands may be optionally digested with an appropriate protease, such as trypsin, to release the peptides, which may then be analyzed by 1D reverse phase LC-MS/MS. The resulting proteomic data may be integrated into so-called peptographs, which plot, in the left panel, sequence coverage for a given protein in the horizontal dimension (N to C terminus, left to right) versus SDS-PAGE migration in the vertical dimension (high to low molecular weight, top to bottom). The specific peptide fragments can then be sequenced or mapped. In certain embodiments, the AARS reference fragment may be characterized by its unique molecular weight, as compared, for example, to the molecular weight of the corresponding full-length AARS.

[0118] As noted above, embodiments of the present invention include the AARS polypeptides set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9. Also included are “variants” of the AARS reference polypeptides. The recitation polypeptide “variant” refers to polypeptides that are distinguished from a reference AARS polypeptide by the addition, deletion, and/or substitution of at least one amino acid residue, and which typically retain (e.g., mimic) or modulate (e.g., antagonize) one or more non-canonical activities of a reference AARS polypeptide.

[0119] Moreover human Arginyl tRNA synthetases include several hundred highly related polymorphic forms, and these are known in the art to be at least partially functionally interchangeable. It would thus be a routine matter to select a naturally occurring variant of Arginyl tRNA synthetase, including, for example the single nucleotide polymorphic forms listed in Table A to create an AARS polypeptide containing one or more amino acid changes based on the sequence of any of the homologues, orthologs, and naturally-occurring isoforms of human as well as other species of Arginyl tRNA synthetase.

TABLE A

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs118184696	A/T	rs2290632	A/T
rs118180032	A/C	rs2290631	A/G
rs117979436	C/T	rs2290630	A/C
rs117938085	C/T	rs2290629	A/G
rs117910938	A/G	rs2290628	A/G
rs117904947	C/T	rs2290627	A/G
rs117805893	C/T	rs2290626	C/T
rs117586708	G/T	rs2290625	A/G
rs116881881	A/G	rs2161173	C/G

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs116754739	A/G	rs2112326	A/G
rs116692940	G/T	rs1870052	G/T
rs116690232	C/T	rs1136673	G/T
rs116688540	C/T	rs1136672	G/T
rs116664558	A/G	rs1136671	C/T
rs116606943	C/T	rs1136669	C/T
rs116604336	A/G	rs1136668	A/C
rs116570152	G/T	rs1136667	C/T
rs116531890	A/G	rs1059443	A/G
rs116464791	A/T	rs918406	C/T
rs116380137	A/C	rs680677	C/G
rs116361226	C/T	rs653568	A/G
rs116257593	A/G	rs632097	A/G
rs116183920	A/G	rs590784	A/C
rs116071286	A/C	rs582906	G/T
rs115948568	A/G	rs244903	A/G
rs115905321	A/G	rs244902	C/T
rs115901063	A/G	rs244901	C/G
rs115887332	A/T	rs244900	A/G
rs115864473	C/T	rs244898	C/T
rs115825016	A/C	rs244897	C/T
rs115760164	A/T	rs244896	G/T
rs115683699	C/T	rs244889	A/C
rs115667735	A/G	rs244888	A/T
rs115473823	A/G	rs193466	C/T
rs115461823	G/T	rs184957	A/T
rs115365149	C/G	rs168647	A/G
rs115263344	C/T	rs165948	C/T
rs115238954	C/T	rs11269831	- /TTTCTGAT TTACGTGAAT CACTTAGTAC ATAT
rs115197001	A/G	rs35585158	- /TTTCTGAT TTACGTGAAT CACTTAGTAC ATAT
rs115167734	C/T	rs6863186	G/T
rs114935064	A/G	rs6555816	C/T

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs114905260	C/G	rs5873114	-/GTTA
rs114795201	C/T	rs4976620	A/G
rs114671846	A/G	rs4976619	A/C
rs114625871	C/T	rs4976618	-/A/T
rs114619091	C/T	rs4976617	C/T
rs114504027	A/G	rs4976616	A/T
rs114270274	A/C	rs4976615	A/T
rs114038151	A/G	rs4976614	C/G
rs114037780	A/G	rs4976560	A/G
rs114006453	C/G	rs4976559	C/T
rs113862186	A/G	rs3797711	A/C
rs113782038	A/G	rs3756602	C/T
rs113720769	A/G	rs3756601	C/T
rs113717899	-/CA	rs3084550	-/ACTA
rs113695809	A/G	rs2445786	A/G
rs113646872	A/G	rs2445785	C/T
rs113605026	C/T	rs2432645	A/G
rs113580372	C/G	rs2432644	A/G
rs113554809	A/G	rs2432643	G/T
rs113486302	C/G	rs2432642	C/G
rs113470525	A/G	rs2307771	-/C
rs113440432	C/T	rs2305737	A/C
rs113378496	A/G	rs2305736	A/G
rs113362439	A/G	rs2305735	A/G
rs113304455	A/T	rs2305734	A/T
rs113175122	A/G	rs2305733	A/G
rs113173551	-/G	rs2305732	C/T
rs113076954	A/G	rs2305731	C/T
rs113072198	A/C	rs2305730	A/G
rs112942704	C/T	rs2305729	C/T
rs112929633	C/G	rs2305728	A/G
rs112915607	A/C	rs2305727	A/G
rs112866258	A/G	rs2305726	A/G
rs112853562	-/TT	rs2305725	A/C
rs112609654	C/T	rs2290633	A/G
rs112554320	A/G	rs11287832	-/T

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs112467769	A/G	rs11134516	A/G
rs112386681	A/G	rs11134515	C/T
rs112374318	C/T	rs10675915	-/TAT
rs112316281	A/C	rs10557485	-/TTT
rs112301722	-/TT	rs10533468	-/CCCC
rs112297730	A/C	rs10516047	A/G
rs112290637	A/G	rs10223174	C/T
rs112132879	C/T	rs10075737	C/T
rs112031493	C/G	rs10075191	A/C
rs112030251	C/T	rs10069598	A/G
rs112026739	C/T	rs10063775	A/G
rs111802241	C/T	rs10042894	A/G
rs111744682	A/T	rs10042188	A/T
rs111718700	C/T	rs9764041	A/T
rs111573970	C/T	rs9717186	A/C
rs111545111	A/G	rs9647575	A/G
rs111462618	C/G	rs9647574	A/G
rs111413807	-/T	rs9647573	G/T
rs111400978	G/T	rs9647572	A/G
rs111236454	A/G	rs9647571	C/G
rs80346848	C/T	rs9331569	-/TTT
rs80313082	G/T	rs7737859	A/G
rs80241471	A/G	rs7729863	A/T
rs80206048	C/T	rs7721460	A/G
rs80066923	C/T	rs7720641	C/T
rs80054466	A/C	rs7714332	A/C
rs79948711	C/T	rs7709842	C/T
rs79719603	A/G	rs7704557	A/G
rs79557735	G/T	rs7704206	G/T
rs79523408	A/G	rs7700690	C/T
rs79321362	C/T	rs6898877	C/G
rs79283789	A/C	rs6898527	A/G
rs79201400	C/T	rs6890233	C/T
rs79154165	-/G	rs6887036	A/G
rs79116025	C/T	rs6882371	A/G

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs79113634	G/T	rs6874879	A/G
rs79103980	C/T	rs6873664	A/C
rs79062764	A/C	rs6868421	A/C
rs78985547	A/G	rs12521951	C/T
rs78849762	G/T	rs12521858	G/T
rs78656507	A/G	rs12521577	C/T
rs78655764	C/T	rs12520913	A/T
rs78604201	A/C	rs12520653	G/T
rs78602242	A/G	rs12520552	A/G
rs78511125	A/T	rs12519228	C/T
rs78390101	-/TTTT	rs12519090	A/C
rs78363759	A/G	rs12516801	A/C
rs78325695	A/C	rs12516460	A/G
rs78274997	G/T	rs12516424	C/T
rs78268250	-/T	rs12515318	A/G
rs78238764	A/G	rs12514748	A/G
rs78217880	A/T	rs12189091	A/T
rs78203348	G/T	rs12188304	C/T
rs78069156	C/T	rs12187995	A/C
rs78039114	C/T	rs12153772	C/G
rs77959391	A/G	rs12153564	A/G
rs77936688	A/C	rs12153563	C/T
rs77827529	A/C	rs11958698	C/T
rs77824483	A/G	rs11958018	C/T
rs77804607	C/T	rs11749654	A/C
rs77771045	A/G	rs11748918	G/T
rs77661711	A/C	rs11746080	A/G
rs77628308	G/T	rs11746078	A/G
rs77607421	G/T	rs11743840	A/G
rs77576032	A/C	rs11743757	C/G
rs77570331	G/T	rs11738672	C/G
rs77483531	-/TC	rs11738671	C/T
rs77319475	C/T	rs11557637	A/C
rs77292057	C/G	rs11557636	A/G
rs77204454	A/C	rs11557634	C/T
rs77178198	C/T	rs11557633	C/T

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs77101561	C/T	rs11453962	-/T
rs77020089	-/T	rs11430234	-/T
rs77005211	G/T	rs11423188	-/T
rs76943812	A/T	rs11345301	-/A
rs76848592	C/T	rs11311315	-/T
rs76727748	A/G	rs11291393	-/A
rs76700391	A/T	rs34446466	A/G
rs76691939	G/T	rs34442207	-/T
rs76685579	C/T	rs34318878	-/T
rs76633933	A/C	rs34274489	-/T
rs76584439	A/C	rs34271397	-/T
rs76532941	C/T	rs34225874	-/T
rs76530903	C/T	rs34121907	-/TAT
rs76484666	G/T	rs34104604	-/G
rs76424455	G/T	rs33990018	-/A
rs76401404	A/T	rs33980252	-/T
rs76376959	A/C	rs33920269	-/TTT
rs76217402	C/G	rs28688070	A/G
rs76174192	A/C	rs28571618	C/T
rs76163401	C/T	rs28537325	A/T
rs76121814	C/T	rs28483077	A/G
rs76112678	A/G	rs17732105	A/G
rs76011402	A/T	rs17732057	C/T
rs75972441	A/G	rs17732045	A/G
rs75643003	A/T	rs17732015	A/G
rs75441465	C/T	rs17731981	A/G
rs75409719	C/T	rs17664614	A/G
rs75399022	A/C	rs17664559	A/G
rs75384112	G/T	rs17664493	C/T
rs75307167	G/T	rs17633878	C/T
rs75249584	G/T	rs17633806	A/G
rs75218784	C/T	rs17633733	A/T
rs75168852	A/C	rs17552440	C/T
rs75140107	G/T	rs17552419	C/T
rs75106564	C/T	rs17552018	C/T

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs75095020	A/T	rs13182927	A/G
rs75031790	C/G	rs13172925	A/G
rs74936396	A/G	rs13172906	C/G
rs74936344	A/T	rs13172658	C/G
rs74897611	C/T	rs13172502	A/C
rs74885621	A/G	rs13158178	A/G
rs74765456	A/G	rs13155873	C/T
rs74744069	C/T	rs13154956	A/T
rs74660183	A/C	rs12654784	A/C
rs74537477	A/G	rs12522512	A/G
rs74487766	A/G	rs57791972	-/CC
rs74468174	A/G	rs57477209	C/T
rs74406691	A/C	rs56891451	-/A
rs74389345	A/T	rs56654743	-/T
rs73802341	C/T	rs56361177	G/T
rs73802340	C/T	rs56341346	A/G
rs73326643	A/G	rs56309278	C/T
rs73316547	A/G	rs56064173	A/G
rs73316530	A/G	rs55966278	G/T
rs73316525	A/G	rs55694682	A/T
rs73316519	C/T	rs55645650	G/T
rs73316518	C/T	rs41283169	C/T
rs73314851	C/G	rs41283167	A/C
rs73314846	A/G	rs36049925	-/AG
rs73314836	A/G	rs36015007	-/A
rs73314833	C/G	rs35898318	-/TTT
rs73314824	A/T	rs35886946	-/T
rs72836363	C/T	rs35738082	-/C
rs72836362	C/T	rs35722978	-/T
rs72836358	C/T	rs35717210	-/T
rs72830975	C/T	rs35643354	-/A
rs72830973	C/G	rs35641348	-/G
rs72830969	C/G	rs35560247	-/T
rs72830965	G/T	rs35468758	-/T
rs72830962	A/G	rs35466646	-/CCCC
rs72830959	C/G	rs35439881	-/AG

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs72532703	-/T	rs35365870	-/A
rs72496362	-/T	rs35319547	-/A
rs72431656	-/A	rs35254979	-/T
rs72411315	-/CC	rs35237798	-/T
rs72334054	-/TAT	rs35170210	-/T
rs72276928	-/T	rs35093178	-/T
rs72227613	-/CA	rs35001933	-/AA
rs72153989	-/T	rs34864760	-/AA
rs72104437	-/AA	rs34791845	-/T
rrs71992136	-/A	rs34723836	A/T
rs71990776	-/ATT	rs34579916	-/A
rs71977003	-/TAGT	rs34501328	-/T
rs71972722	-/T	rs34484619	-/T
rs71905728	-/T	rs34578180	-/A/CA/G
rs71896554	-/AC	rs34669829	-/G/GTTA
rs71796598	-/AAAAA	rs67225121	-/CACA/CAC ACA
rs71779923	-/A	rs61156257	-/CACACA
rs71738365	-/T	rs72412281	-/CACACA
rs71603865	A/T	rs72083953	-/ACACAC
rs71603864	A/G	rs70976491	-/AAAAAAAA AAAAAAAA AAAAAAAA AAAA
rs71589837	A/G	rs61532333	-/TTTT
rs71588659	AAA/TAT	rs61525394	-/TAT
rs71303324	-/AAT	rs61456455	A/G
rs70976490	-/A	rs60743864	-/AA
rs70976489	-/AAA	rs60614962	-/T
rs67907149	-/T	rs60594519	A/G
rs67885474	-/C	rs60514038	C/T
rs67562347	-/T	rs60492623	C/T
rs67012869	C/T	rs60401511	-/AA
rs66757470	-/T	rs60295313	A/C
rs66745652	-/T	rs59359069	A/G
rs62385664	A/G	rs58186106	-/A
rs62385663	G/T	rs58133689	-/C

TABLE A-continued

Human Arginyl tRNA synthetase SNPs			
Gene Bank Accession Number	Nucleotide Change	Gene Bank Accession Number	Nucleotide Change
rs62385662	A/G	rs57885645	-/G
rs62385661	A/C	rs57821179	-/T
rs62385660	C/T	rs61532333	-/TTTT
rs61744940	A/G	rs61525394	-/TAT
		rs61456455	A/G
		rs60743864	-/AA

[0120] In certain embodiments, a polypeptide variant is distinguished from a reference polypeptide by one or more substitutions, which may be conservative or non-conservative, as described herein and known in the art. In certain embodiments, the polypeptide variant comprises conservative substitutions and, in this regard, it is well understood in the art that some amino acids may be changed to others with broadly similar properties without changing the nature of the activity of the polypeptide.

[0121] In certain embodiments, a variant polypeptide includes an amino acid sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or more sequence identity or similarity to a corresponding sequence of an AARS reference polypeptide, as described herein, and substantially retains the non-canonical activity of that reference polypeptide. Also included are sequences differing from the reference AARS sequences by the addition, deletion, or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150 or more amino acids but which retain the properties of the reference AARS polypeptide. In certain embodiments, the amino acid additions or deletions occur at the C-terminal end and/or the N-terminal end of the AARS reference polypeptide. In certain embodiments, the amino acid additions include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50 or more wild-type residues (i.e., from the corresponding full-length AARS polypeptide) that are proximal to the C-terminal end and/or the N-terminal end of the AARS reference polypeptide.

[0122] In certain embodiments, variant polypeptides differ from the corresponding AARS reference sequences by at least 1% but less than 20%, 15%, 10% or 5% of the residues. (If this comparison requires alignment, the sequences should be aligned for maximum similarity. “Looped” out sequences from deletions or insertions, or mismatches, are considered differences.) The differences are, suitably, differences or changes at a non-essential residue or a conservative substitution. In certain embodiments, the molecular weight of a variant AARS polypeptide differs from that of the AARS reference polypeptide by about 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, or more.

[0123] Also included are biologically active “fragments” of the AARS reference polypeptides, i.e., biologically active fragments of the AARS protein fragments. Representative biologically active fragments generally participate in an

interaction, e.g., an intramolecular or an inter-molecular interaction. An inter-molecular interaction can be a specific binding interaction or an enzymatic interaction. An inter-molecular interaction can be between an AARS polypeptide and a cellular binding partner, such as a cellular receptor or other host molecule that participates in the non-canonical activity of the AARS polypeptide. In some embodiments, AARS proteins, variants, and biologically active fragments thereof, bind to one or more cellular binding partners with an affinity of at least about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM. The binding affinity of an AARS protein fragment for a selected cellular binding partner, particularly a binding partner that participates in a non-canonical activity, is typically stronger than that of the AARS protein fragment’s corresponding full-length AARS polypeptide, by at least about 1.5x, 2x, 2.5x, 3x, 3.5x, 4x, 4.5x, 5x, 6x, 7x, 8x, 9x, 10x, 15x, 20x, 25x, 30x, 40x, 50x, 60x, 70x, 80x, 90x, 100x, 200x, 300x, 400x, 500x, 600x, 700x, 800x, 900x, 1000x or more (including all integers in between). The binding affinity of an AARS protein fragment for a binding partner that participates in at least one canonical activity of an AARS is typically weaker than that of the AARS protein fragment’s corresponding full-length AARS polypeptide, by at least about 1.5x, 2x, 2.5x, 3x, 3.5x, 4x, 4.5x, 5x, 6x, 7x, 8x, 9x, 10x, 15x, 20x, 25x, 30x, 40x, 50x, 60x, 70x, 80x, 90x, 100x, 200x, 300x, 400x, 500x, 600x, 700x, 800x, 900x, 1000x or more.

[0124] Typically, biologically active fragments comprise a domain or motif with at least one activity of an AARS reference polypeptide and may include one or more (and in some cases all) of the various active domains, and include fragments having a non-canonical activity. In some cases, biologically active fragments of an AARS polypeptide have a biological activity that is unique to the particular, truncated fragment, such that the full-length AARS polypeptide may not have that activity. In certain cases, the biological activity may be revealed by separating the biologically active AARS polypeptide fragment from the other full-length AARS polypeptide sequences, or by altering certain residues of the full-length AARS wild-type polypeptide sequence to unmask the biologically active domains.

[0125] A biologically active fragment of an AARS reference polypeptide can be a polypeptide fragment which is, for example, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 220, 240, 260, 280, 300, 320, 340, 360, 380, 400, 450, 500, 550, 600, 650, 700, 750 or more contiguous or non-contiguous amino acids, including all integers (e.g., 101, 102, 103) and ranges (e.g., 50-100, 50-150, 50-200) in between, of the amino acid sequences set forth in any one of the AARS reference polypeptides described herein, but typically exclude the full-length AARS. In certain embodiments, a biologically active fragment comprises a non-canonical activity-related sequence, domain, or motif. In certain embodiments, the C-terminal or N-terminal region of any AARS reference polypeptide may be truncated by about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, or 700 or more amino acids, or by about 10-50, 20-50, 50-100, 100-150, 150-200, 200-250, 250-300, 300-350, 350-400, 400-

450, 450-500, 500-550, 550-600, 600-650, 650-700 or more amino acids, including all integers and ranges in between (e.g., 101, 102, 103, 104, 105), so long as the truncated AARS polypeptide retains the non-canonical activity of the reference polypeptide. Typically, the biologically-active fragment has no less than about 1%, about 5%, about 10%, about 25%, or about 50% of an activity of the biologically-active (i.e., non-canonical activity) AARS reference polypeptide from which it is derived. Exemplary methods for measuring such non-canonical activities are described in the Examples.

[0126] As noted above, an AARS polypeptide may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of an AARS reference polypeptide can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985, *Proc. Natl. Acad. Sci. USA*, 82: 488-492), Kunkel et al., (1987, *Methods in Enzymol.*, 154: 367-382), U.S. Pat. No. 4,873,192, Watson, J. D. et al., (“Molecular Biology of the Gene”, Fourth Edition, Benjamin/Cummings, Menlo Park, Calif., 1987) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff et al., (1978) Atlas of Protein Sequence and Structure (Natl. Biomed. Res. Found., Washington, D.C.).

[0127] Similarly it is within the skill in the art to address and/or mitigate immunogenicity concerns if they arise using an AARS polypeptide, e.g., by the use of automated computer recognition programs to identify potential T cell epitopes, and directed evolution approaches to identify less immunogenic forms.

[0128] Methods for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property are known in the art. Such methods are adaptable for rapid screening of the gene libraries generated by combinatorial mutagenesis of AARS polypeptides. Recursive ensemble mutagenesis (REM), a technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify AARS polypeptide variants (Arkin and Yourvan (1992) *Proc. Natl. Acad. Sci. USA* 89: 7811-7815; Delgrave et al., (1993) *Protein Engineering*, 6: 327-331). Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be desirable as discussed in more detail below.

[0129] Biologically active truncated and/or variant AARS polypeptides may contain conservative amino acid substitutions at various locations along their sequence, as compared to a reference AARS amino acid residue. Additionally, naturally occurring variants of AARS proteins have been sequenced, and are known in the art to be at least partially functionally interchangeable. It would thus be a routine matter to select an amino acid position to introduce a conservative, or non-conservative mutation into an AARS polypeptide based on naturally occurring sequence variation among the known AARS protein homologues, orthologs, and naturally-occurring isoforms of human as well as other species of an AARS protein.

[0130] A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art, which can be generally sub-classified as follows:

[0131] Acidic: The residue has a negative charge due to loss of H ion at physiological pH and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH. Amino acids having an acidic side chain include glutamic acid and aspartic acid.

[0132] Basic: The residue has a positive charge due to association with H ion at physiological pH or within one or two pH units thereof (e.g., histidine) and the residue is attracted by aqueous solution so as to seek the surface positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium at physiological pH. Amino acids having a basic side chain include arginine, lysine and histidine.

[0133] Charged: The residues are charged at physiological pH and, therefore, include amino acids having acidic or basic side chains (i.e., glutamic acid, aspartic acid, arginine, lysine and histidine).

[0134] Hydrophobic: The residues are not charged at physiological pH and the residue is repelled by aqueous solution so as to seek the inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. Amino acids having a hydrophobic side chain include tyrosine, valine, isoleucine, leucine, methionine, phenylalanine and tryptophan.

[0135] Neutral/polar: The residues are not charged at physiological pH, but the residue is not sufficiently repelled by aqueous solutions so that it would seek inner positions in the conformation of a peptide in which it is contained when the peptide is in aqueous medium. Amino acids having a neutral/polar side chain include asparagine, glutamine, cysteine, histidine, serine and threonine.

[0136] This description also characterizes certain amino acids as “small” since their side chains are not sufficiently large, even if polar groups are lacking, to confer hydrophobicity. With the exception of proline, “small” amino acids are those with four carbons or less when at least one polar group is on the side chain and three carbons or less when not. Amino acids having a small side chain include glycine, serine, alanine and threonine. The gene-encoded secondary amino acid proline is a special case due to its known effects on the secondary conformation of peptide chains. The structure of proline differs from all the other naturally-occurring amino acids in that its side chain is bonded to the nitrogen of the α -amino group, as well as the α -carbon. Several amino acid similarity matrices are known in the art (see e.g., PAM120 matrix and PAM250 matrix as disclosed for example by Dayhoff et al., 1978, A model of evolutionary change in proteins). Matrices for determining distance relationships In M. O. Dayhoff, (ed.), Atlas of protein sequence and structure, Vol. 5, pp. 345-358, National Biomedical Research Foundation, Washington D.C.; and by Gonnet et al., (*Science*, 256: 14430-1445, 1992), however, include proline in the same group as glycine, serine, alanine and threonine. Accordingly, for the purposes of the present invention, proline is classified as a “small” amino acid.

[0137] The degree of attraction or repulsion required for classification as polar or nonpolar is arbitrary and, therefore,

amino acids specifically contemplated by the invention have been classified as one or the other. Most amino acids not specifically named can be classified on the basis of known behavior.

[0138] Amino acid residues can be further sub-classified as cyclic or non-cyclic, and aromatic or non-aromatic, self-explanatory classifications with respect to the side-chain substituent groups of the residues, and as small or large. The residue is considered small if it contains a total of four carbon atoms or less, inclusive of the carboxyl carbon, provided an additional polar substituent is present; three or less if not. Small residues are, of course, always non-aromatic. Dependent on their structural properties, amino acid residues may fall in two or more classes. For the naturally-occurring protein amino acids, sub-classification according to this scheme is presented in Table B.

TABLE B

Amino acid sub-classification	
Sub-classes	Amino acids
Acidic	Aspartic acid, Glutamic acid
Basic	Noncyclic: Arginine, Lysine; Cyclic: Histidine
Charged	Aspartic acid, Glutamic acid, Arginine, Lysine, Histidine
Small	Glycine, Serine, Alanine, Threonine, Proline
Polar/neutral	Asparagine, Histidine, Glutamine, Cysteine, Serine, Threonine
Polar/large	Asparagine, Glutamine
Hydrophobic	Tyrosine, Valine, Isoleucine, Leucine, Methionine, Phenylalanine, Tryptophan
Aromatic	Tryptophan, Tyrosine, Phenylalanine
Residues that influence chain orientation	Glycine and Proline

[0139] Conservative amino acid substitution also includes groupings based on side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. For example, it is reasonable to expect that replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the properties of the resulting variant polypeptide. Whether an amino acid change results in a functional truncated and/or variant AARS polypeptide can readily be determined by assaying its non-canonical activity, as described herein. Conservative substitutions are shown in Table C under the heading of exemplary substitutions. Amino acid substitutions falling within the scope of the invention, are, in general, accomplished by selecting substitutions that do not differ significantly in their effect on maintaining (a) the structure of the peptide backbone in the area of the substitution, (b) the charge or hydrophobicity of the molecule at the target site, (c) the bulk of the side chain, or (d) the biological function. After the substitutions are introduced, the variants are screened for biological activity.

TABLE C

Exemplary Amino Acid Substitutions		
Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala	Val, Leu, Ile	Val
Arg	Lys, Gln, Asn	Lys
Asn	Gln, His, Lys, Arg	Gln
Asp	Glu	Glu
Cys	Ser	Ser
Gln	Asn, His, Lys,	Asn
Glu	Asp, Lys	Asp
Gly	Pro	Pro
His	Asn, Gln, Lys, Arg	Arg
Ile	Leu, Val, Met, Ala, Phe, Norleu	Leu
Leu	Norleu, Ile, Val, Met, Ala, Phe	Ile
Lys	Arg, Gln, Asn	Arg
Met	Leu, Ile, Phe	Leu
Phe	Leu, Val, Ile, Ala	Leu
Pro	Gly	Gly
Ser	Thr	Thr
Thr	Ser	Ser
Trp	Tyr	Tyr
Tyr	Trp, Phe, Thr, Ser	Phe
Val	Ile, Leu, Met, Phe, Ala, Norleu	Leu

[0140] Alternatively, similar amino acids for making conservative substitutions can be grouped into three categories based on the identity of the side chains. The first group includes glutamic acid, aspartic acid, arginine, lysine, histidine, which all have charged side chains; the second group includes glycine, serine, threonine, cysteine, tyrosine, glutamine, asparagine; and the third group includes leucine, isoleucine, valine, alanine, proline, phenylalanine, tryptophan, methionine, as described in Zubay, G., *Biochemistry*, third edition, Wm.C. Brown Publishers (1993).

[0141] Thus, a predicted non-essential amino acid residue in a truncated and/or variant AARS polypeptide is typically replaced with another amino acid residue from the same side chain family. Alternatively, mutations can be introduced randomly along all or part of an AARS coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for an activity of the parent polypeptide to identify mutants which retain that activity. Following mutagenesis of the coding sequences, the encoded peptide can be expressed recombinantly and the activity of the peptide can be determined. A “non-essential” amino acid residue is a residue that can be altered from the reference sequence of an embodiment polypeptide without abolishing or substantially altering one or more of its activities. Suitably, the alteration does not substantially abolish one of these activities, for example, the activity is at least 20%, 40%, 60%, 70% or 80% 100%, 500%, 1000% or more of the reference AARS sequence. An “essential” amino acid residue is a residue that, when altered from the reference sequence of an AARS polypeptide, results in abolition of an activity of the parent molecule such that less than 20% of the reference activity is present. For example, such essential amino acid residues include those that are conserved in AARS polypeptides across different species, including those sequences that are conserved in the active binding site(s) or motif(s) of AARS polypeptides from various sources.

[0142] In general, polypeptides and fusion polypeptides (as well as their encoding polynucleotides) are isolated. An “isolated” polypeptide or polynucleotide is one that is removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from

some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

[0143] Certain embodiments also encompass dimers of AARS polypeptides. Dimers may include, for example, homodimers between two identical AARS polypeptides, heterodimers between two different AARS polypeptides (e.g., a full-length YRS polypeptide and a truncated YRS polypeptide; a truncated YRS polypeptide and a truncated WRS polypeptide), and/or heterodimers between an AARS polypeptide and a heterologous polypeptide. Certain heterodimers, such as those between an AARS polypeptide and a heterologous polypeptide, may be bi-functional, as described herein.

[0144] Also included are monomers of AARS polypeptides, including isolated AARS polypeptides monomers that do not substantially dimerize with a second AARS polypeptide, whether due to one or more substitutions, truncations, deletions, additions, chemical modifications, or a combination of these alterations. In certain embodiments, monomeric AARS polypeptides possess biological activities, including non-canonical activities, which are not possessed by dimeric or multimeric AARS polypeptide complexes.

[0145] Certain embodiments of the present invention also contemplate the use of modified AARS polypeptides, including modifications that improved the desired characteristics of an AARS polypeptide, as described herein. Modifications of AARS polypeptides of the invention include chemical and/or enzymatic derivatizations at one or more constituent amino acid, including side chain modifications, backbone modifications, and N- and C-terminal modifications including acetylation, hydroxylation, methylation, amidation, and the attachment of carbohydrate or lipid moieties, cofactors, and the like. Exemplary modifications also include pegylation of an AARS polypeptide (see, e.g., Veronese and Harris, *Advanced Drug Delivery Reviews* 54: 453-456, 2002; and Pasut et al., *Expert Opinion. Ther. Patents* 14(6) 859-894 2004, both herein incorporated by reference).

[0146] PEG is a well-known polymer having the properties of solubility in water and in many organic solvents, lack of toxicity, and lack of immunogenicity. It is also clear, colorless, odorless, and chemically stable. For these reasons and others, PEG has been selected as the preferred polymer for attachment, but it has been employed solely for purposes of illustration and not limitation. Similar products may be obtained with other water-soluble polymers, including without limitation; polyvinyl alcohol, other poly(alkylene oxides) such as poly(propylene glycol) and the like, poly(oxyethylated polyols) such as poly(oxyethylated glycerol) and the like, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1,3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride, and polyaminoacids. One skilled in the art will be able to select the desired polymer based on the desired dosage, circulation time, resistance to proteolysis, and other considerations.

[0147] In particular a wide variety of PEG derivatives are both available and suitable for use in the preparation of PEG-conjugates. For example, NOF Corp.'s PEG reagents sold under the trademark SUNBRIGHT® Series provides numerous PEG derivatives, including methoxypolyethylene

glycols and activated PEG derivatives such as methoxy-PEG amines, maleimides, N-hydroxysuccinimide esters, and carboxylic acids, for coupling by various methods to the N-terminal, C-terminal or any internal amino acid of the AARS polypeptide. Nektar Therapeutics' Advanced PEGylation technology also offers diverse PEG-coupling technologies to potentially improve the safety and efficacy of an AARS polypeptide based therapeutic.

[0148] A search of patents, published patent applications, and related publications will also provide those skilled in the art reading this disclosure with significant possible PEG-coupling technologies and PEG-derivatives. For example, U.S. Pat. Nos. 6,436,386; 5,932,462; 5,900,461; 5,824,784; and 4,904,584; the contents of which are incorporated by reference in their entirety, describe such technologies and derivatives, and methods for their manufacture.

[0149] In certain aspects, chemoselective ligation technology may be utilized to modify AARS polypeptides of the invention, such as by attaching polymers in a site-specific and controlled manner. Such technology typically relies on the incorporation of chemoselective anchors into the protein backbone by either chemical or recombinant means, and subsequent modification with a polymer carrying a complementary linker. As a result, the assembly process and the covalent structure of the resulting protein-polymer conjugate may be controlled, enabling the rational optimization of drug properties, such as efficacy and pharmacokinetic properties (see, e.g., Kochendoerfer, *Current Opinion in Chemical Biology* 9:555-560, 2005).

[0150] In other embodiments, fusion proteins of AARS polypeptide to other proteins are also included, and these fusion proteins may increase the AARS polypeptide's biological activity, secretion, targeting, biological life, the ability to penetrate cellular membranes or the blood brain barrier, or pharmacokinetic properties. Examples of fusion proteins that improve pharmacokinetic properties ("PK modifiers") include without limitation, fusions to human albumin (Osborn et al.: *Eur. J. Pharmacol.* 456(1-3): 149-158, (2002)), antibody Fc domains, poly Glu or poly Asp sequences, and transferrin. Additionally, fusion with conformationally disordered polypeptide sequences composed of the amino acids Pro, Ala, and Ser ("PASylation") or hydroxyethyl starch (sold under the trademark HESYLATION®) provides a simple way to increase the hydrodynamic volume of the AARS polypeptide. This additional extension adopts a bulky random structure, which significantly increases the size of the resulting fusion protein. By this means the typically rapid clearance of smaller AARS polypeptides via kidney filtration is retarded by several orders of magnitude. Additionally use of Ig G fusion proteins has also been shown to enable some fusion protein proteins to penetrate the blood brain barrier (Fu et al., (2010) *Brain Res.* 1352:208-13).

[0151] Examples of fusion proteins that improve penetration across cellular membranes include fusions to membrane translocating sequences. In this context, the term "membrane translocating sequences" refers to naturally occurring and synthetic amino acid sequences that are capable of membrane translocation across a cellular membrane. Representative membrane translocating sequences include those based on the naturally occurring membrane translocating sequences derived from the Tat protein, and homeotic transcription protein Antennapedia, as well as synthetic membrane translocating sequences based in whole or part on poly Arginine and Lysine residues. Representative membrane

translocating sequences include for example those disclosed in the following patents, U.S. Pat. No. 5,652,122; U.S. Pat. No. 5,670,617; U.S. Pat. No. 5,674,980; U.S. Pat. No. 5,747,641; U.S. Pat. No. 5,804,604; U.S. Pat. No. 6,316,003; U.S. Pat. No. 7,585,834; U.S. Pat. No. 7,312,244; U.S. Pat. No. 7,279,502; U.S. Pat. No. 7,229,961; U.S. Pat. No. 7,169,814; U.S. Pat. No. 7,453,011; U.S. Pat. No. 7,235,695; U.S. Pat. No. 6,982,351; U.S. Pat. No. 6,605,115; U.S. Pat. No. 7,306,784; U.S. Pat. No. 7,306,783; U.S. Pat. No. 6,589,503; U.S. Pat. No. 6,348,185; U.S. Pat. No. 6,881,825; U.S. Pat. No. 7,431,915; WO0074701A2; WO2007111993A2; WO2007106554A2; WO02069930A1; WO03049772A2; WO03106491A2; and WO2008063113A1.

[0152] It will be appreciated that a flexible molecular linker (or spacer) optionally may be interposed between, and covalently join, the AARS polypeptide and any of the fusion proteins disclosed herein.

[0153] Additionally in some embodiments, the AARS polypeptide can include synthetic, or naturally occurring secretion signal sequences, derived from other well characterized secreted proteins. In some embodiments such proteins, may be processed by proteolytic cleavage to form the AARS polypeptide in situ. Such fusions proteins include for example fusions of AARS polypeptide to ubiquitin to provide a new N-terminal amino acid, or the use of a secretion signal to mediate high level secretion of the AARS polypeptide into the extracellular medium, or N, or C-terminal epitope tags to improve purification or detection.

[0154] The AARS polypeptides described herein may be prepared by any suitable procedure known to those of skill in the art, such as by recombinant techniques. In addition to recombinant production methods, polypeptides of the invention may be produced by direct peptide synthesis using solid-phase techniques (Merrifield, *J. Am. Chem. Soc.* 85:2149-2154 (1963)). Protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Alternatively, various fragments may be chemically synthesized separately and combined using chemical methods to produce the desired molecule.

IV. AARS POLYNUCLEOTIDES

[0155] Embodiments of the present invention include polynucleotides that encode one or more newly identified protein fragments of an aminoacyl-tRNA synthetase (AARS), in addition to complements, variants, and fragments thereof. In certain embodiments, an AARS polynucleotide encodes all or a portion of the AARS polypeptide reference sequence(s) as set forth in Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9, which represent splice variants, proteolytic fragments, or other type of fragments of Arginyl tRNA synthetase. Certain embodiments include polynucleotides, encoding polypeptides or proteins that comprise the sequence of one or more splice junctions of those splice variants, in addition to complements, variants, and fragments thereof. In certain embodiments, typically due to the singular nature of a selected AARS splice variant, which combines exons in a new or exceptional way, the AARS polynucleotide references sequences comprise a unique or exceptional splice junction. Certain embodiments exclude a corresponding full-length AARS polynucleotide.

[0156] Also included within the AARS polynucleotides of the present invention are primers, probes, antisense oligonucleotides, and RNA interference agents that comprise all or a portion of these reference polynucleotides, which are complementary to all or a portion of these reference polynucleotides, or which specifically hybridize to these reference polynucleotides, as described herein.

[0157] The term “polynucleotide” or “nucleic acid” as used herein designates mRNA, RNA, cRNA, cDNA or DNA. The term typically refers to polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The terms “DNA” and “polynucleotide” and “nucleic acid” refer to a DNA molecule that has been isolated free of total genomic DNA of a particular species. Therefore, an isolated DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Also included are non-coding polynucleotides (e.g., primers, probes, oligonucleotides), which do not encode an AARS polypeptide. Included within the terms “DNA segment” and “polynucleotide” are DNA segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phagemids, phage, viruses, and the like.

[0158] Additional coding or non-coding sequences may, but need not, be present within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials. Hence, the polynucleotides of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably.

[0159] It is therefore contemplated that a polynucleotide fragment of almost any length may be employed; with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. Included are polynucleotides of about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 41, 43, 44, 45, 46, 47, 48, 49, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 220, 240, 260, 270, 280, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 2100, 2200, 2300, 2400, 2500, 2600, 2700, 2800, 2900, 3000 or more (including all integers in between) bases in length, including any portion or fragment (e.g., greater than about 6, 7, 8, 9, or 10 nucleotides in length) of an AARS reference polynucleotide (e.g., base number X-Y, in which X is about 1-3000 or more and Y is about 10-3000 or more), or its complement.

[0160] Embodiments of the present invention also include “variants” of the AARS reference polynucleotide sequences. Polynucleotide “variants” may contain one or more substitutions, additions, deletions and/or insertions in relation to a reference polynucleotide. Generally, variants of an AARS reference polynucleotide sequence may have at least about 30%, 40% 50%, 55%, 60%, 65%, 70%, generally at least about 75%, 80%, 85%, desirably about 90% to 95% or more, and more suitably about 98% or more sequence identity to

that particular nucleotide sequence as determined by sequence alignment programs described elsewhere herein using default parameters. In certain embodiments, variants may differ from a reference sequence by about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 60, 70, 80, 90, 100 (including all integers in between) or more bases. In certain embodiments, such as when the polynucleotide variant encodes an AARS polypeptide having a non-canonical activity, the desired activity of the encoded AARS polypeptide is not substantially diminished relative to the unmodified polypeptide. The effect on the activity of the encoded polypeptide may generally be assessed as described herein.

[0161] Certain embodiments include polynucleotides that hybridize to a reference AARS polynucleotide sequence, or to their complements, under stringency conditions described below. As used herein, the term “hybridizes under low stringency, medium stringency, high stringency, or very high stringency conditions” describes conditions for hybridization and washing. Guidance for performing hybridization reactions can be found in Ausubel et al., (1998, *supra*), Sections 6.3.1-6.3.6. Aqueous and non-aqueous methods are described in that reference and either can be used.

[0162] Reference herein to low stringency conditions include and encompass from at least about 1% v/v to at least about 15% v/v formamide and from at least about 1 M to at least about 2 M salt for hybridization at 42° C., and at least about 1 M to at least about 2 M salt for washing at 42° C. Low stringency conditions also may include 1% Bovine Serum Albumin (BSA), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridization at 65° C., and (i) 2×SSC, 0.1% SDS; or (ii) 0.5% BSA, 1 mM EDTA, 40 mM NaHPO₄ (pH 7.2), 5% SDS for washing at room temperature. One embodiment of low stringency conditions includes hybridization in 6× sodium chloride/sodium citrate (SSC) at about 45° C., followed by two washes in 0.2×SSC, 0.1% SDS at least at 50° C. (the temperature of the washes can be increased to 55° C. for low stringency conditions).

[0163] Medium stringency conditions include and encompass from at least about 16% v/v to at least about 30% v/v formamide and from at least about 0.5 M to at least about 0.9 M salt for hybridization at 42° C., and at least about 0.1 M to at least about 0.2 M salt for washing at 55° C. Medium stringency conditions also may include 1% Bovine Serum Albumin (BSA), 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridization at 65° C., and (i) 2×SSC, 0.1% SDS; or (ii) 0.5% BSA, 1 mM EDTA, 40 mM NaHPO₄ (pH 7.2), 5% SDS for washing at 60-65° C. One embodiment of medium stringency conditions includes hybridizing in 6×SSC at about 45° C., followed by one or more washes in 0.2×SSC, 0.1% SDS at 60° C. High stringency conditions include and encompass from at least about 31% v/v to at least about 50% v/v formamide and from about 0.01 M to about 0.15 M salt for hybridization at 42° C., and about 0.01 M to about 0.02 M salt for washing at 55° C.

[0164] High stringency conditions also may include 1% BSA, 1 mM EDTA, 0.5 M NaHPO₄ (pH 7.2), 7% SDS for hybridization at 65° C., and (i) 0.2×SSC, 0.1% SDS; or (ii) 0.5% BSA, 1 mM EDTA, 40 mM NaHPO₄ (pH 7.2), 1% SDS for washing at a temperature in excess of 65° C. One embodiment of high stringency conditions includes hybridizing in 6×SSC at about 45° C., followed by one or more washes in 0.2×SSC, 0.1% SDS at 65° C. One embodiment

of very high stringency conditions includes hybridizing in 0.5 M sodium phosphate, 7% SDS at 65° C., followed by one or more washes in 0.2×SSC, 1% SDS at 65° C.

[0165] Other stringency conditions are well known in the art and a skilled artisan will recognize that various factors can be manipulated to optimize the specificity of the hybridization. Optimization of the stringency of the final washes can serve to ensure a high degree of hybridization. For detailed examples, see Ausubel et al., *supra* at pages 2.10.1 to 2.10.16 and Sambrook et al. (1989, *supra*) at sections 1.101 to 1.104.

[0166] While stringent washes are typically carried out at temperatures from about 42° C. to 68° C., one skilled in the art will appreciate that other temperatures may be suitable for stringent conditions. Maximum hybridization rate typically occurs at about 20° C. to 25° C. below the T_m for formation of a DNA-DNA hybrid. It is well known in the art that the T_m is the melting temperature, or temperature at which two complementary polynucleotide sequences dissociate. Methods for estimating T_m are well known in the art (see Ausubel et al., *supra* at page 2.10.8).

[0167] In general, the T_m of a perfectly matched duplex of DNA may be predicted as an approximation by the formula: $T_m = 81.5 + 16.6 (\log_{10} M) + 0.41 (\% \text{ G+C}) - 0.63 (\% \text{ formamide}) - (600/\text{length})$ wherein: M is the concentration of Na⁺, preferably in the range of 0.01 molar to 0.4 molar; % G+C is the sum of guanosine and cytosine bases as a percentage of the total number of bases, within the range between 30% and 75% G+C; % formamide is the percent formamide concentration by volume; length is the number of base pairs in the DNA duplex. The T_m of a duplex DNA decreases by approximately 1° C. with every increase of 1% in the number of randomly mismatched base pairs. Washing is generally carried out at $T_m - 15^\circ \text{C.}$ for high stringency, or $T_m - 30^\circ \text{C.}$ for moderate stringency.

[0168] In one example of a hybridization procedure, a membrane (e.g., a nitrocellulose membrane or a nylon membrane) containing immobilized DNA is hybridized overnight at 42° C. in a hybridization buffer (50% deionized formamide, 5×SSC, 5×Denhardt's solution (0.1% ficoll, 0.1% polyvinylpyrrolidone and 0.1% bovine serum albumin), 0.1% SDS and 200 mg/mL denatured salmon sperm DNA) containing a labeled probe. The membrane is then subjected to two sequential medium stringency washes (i.e., 2×SSC, 0.1% SDS for 15 min at 45° C., followed by 2×SSC, 0.1% SDS for 15 min at 50° C.), followed by two sequential higher stringency washes (i.e., 0.2×SSC, 0.1% SDS for 12 min at 55° C. followed by 0.2×SSC and 0.1% SDS solution for 12 min at 65-68° C.

[0169] As noted above, certain embodiments relate to AARS polynucleotides that encode an AARS polypeptide. Among other uses, these embodiments may be utilized to recombinantly produce a desired AARS polypeptide or variant thereof, or to express the AARS polypeptide in a selected cell or subject. It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides may bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to differences in codon usage are specifically contemplated by the present invention, for example polynucleotides that are optimized for human and/or primate codon selection.

[0170] Therefore, multiple polynucleotides can encode the AARS polypeptides of the invention. Moreover, the polynucleotide sequence can be manipulated for various reasons. Examples include but are not limited to the incorporation of preferred codons to enhance the expression of the polynucleotide in various organisms (see generally Nakamura et al., *Nuc. Acid. Res.* (2000) 28 (1): 292). In addition, silent mutations can be incorporated in order to introduce, or eliminate restriction sites, decrease the density of CpG dinucleotide motifs (see for example, Kameda et al., *Biochem. Biophys. Res. Commun.* (2006) 349(4): 1269-1277) or reduce the ability of single stranded sequences to form stem-loop structures: (see, e.g., Zuker M., *Nucl. Acid Res.* (2003); 31(13): 3406-3415). In addition, mammalian expression can be further optimized by including a Kozak consensus sequence [i.e., (a/g)cc(a/g)ccATGg] at the start codon. Kozak consensus sequences useful for this purpose are known in the art (Mantyh et al. *PNAS* 92: 2662-2666 (1995); Mantyh et al. *Prot. Exp. & Purif.* 6,124 (1995)).

[0171] The polynucleotides of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a polynucleotide fragment of almost any length may be employed; with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

[0172] Polynucleotides and fusions thereof may be prepared, manipulated and/or expressed using any of a variety of well-established techniques known and available in the art. For example, polynucleotide sequences which encode polypeptides of the invention, or fusion proteins or functional equivalents thereof, may be used in recombinant DNA molecules to direct expression of an AARS polypeptide in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences that encode substantially the same or a functionally equivalent amino acid sequence may be produced and these sequences may be used to clone and express a given polypeptide.

[0173] As will be understood by those of skill in the art, it may be advantageous in some instances to produce polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce a recombinant RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence. Such polynucleotides are commonly referred to as "codon-optimized." Any of the polynucleotides described herein may be utilized in a codon-optimized form. In certain embodiments, a polynucleotide can be codon optimized for use in specific bacteria such as *E. coli* or yeast such as *S. cerevisiae* (see, e.g., Burgess-Brown et al., *Protein Expr Purif.* 59:94-102, 2008; Emolaeva M D (2001) *Curr. Iss. Mol. Biol.* 3 (4) 91-7; Welch et al., *PLoS ONE* 4(9): e7007 doi:10.1371/journal.pone.0007002).

[0174] Moreover, the polynucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter polypeptide encoding sequences for a variety of reasons, including but not limited

to, alterations which modify the cloning, processing, expression and/or activity of the gene product.

[0175] According to another aspect of the invention, polynucleotides encoding polypeptides of the invention may be delivered to a subject *in vivo*, e.g., using gene therapy techniques. Gene therapy refers generally to the transfer of heterologous nucleic acids to the certain cells, target cells, of a mammal, particularly a human, with a disorder or conditions for which such therapy is sought. The nucleic acid is introduced into the selected target cells in a manner such that the heterologous DNA is expressed and a therapeutic product encoded thereby is produced.

[0176] Various viral vectors that can be utilized for gene therapy as taught herein include adenovirus, herpes virus, vaccinia, adeno-associated virus (AAV), or, preferably, an RNA virus such as a retrovirus. Preferably, the retroviral vector is a derivative of a murine or avian retrovirus, or is a lentiviral vector. The preferred retroviral vector is a lentiviral vector. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), SIV, BIV, HIV and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified and generated. By inserting a zinc finger derived-DNA binding polypeptide sequence of interest into the viral vector, along with another gene that encodes the ligand for a receptor on a specific target cell, for example, the vector may be made target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a protein (dimer). Illustrative targeting may be accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector containing the zinc finger-nucleotide binding protein polynucleotide.

[0177] Since recombinant retroviruses are defective, they require assistance in order to produce infectious vector particles. This assistance can be provided, for example, by using helper cell lines that contain plasmids encoding all of the structural genes of the retrovirus under the control of regulatory sequences within the LTR. These plasmids are missing a nucleotide sequence which enables the packaging mechanism to recognize an RNA transcript for encapsulation. Helper cell lines which have deletions of the packaging signal include but are not limited to PSI.2, PA317 and PA12, for example. These cell lines produce empty virions, since no genome is packaged. If a retroviral vector is introduced into such cells in which the packaging signal is intact, but the structural genes are replaced by other genes of interest, the vector can be packaged and vector virion produced. The vector virions produced by this method can then be used to infect a tissue cell line, such as NIH 3T3 cells, to produce large quantities of chimeric retroviral virions.

[0178] "Non-viral" delivery techniques for gene therapy can also be used including, for example, DNA-ligand complexes, adenovirus-ligand-DNA complexes, direct injection of DNA, CaPO_4 precipitation, gene gun techniques, electroporation, liposomes, lipofection, and the like. Any of these methods are widely available to one skilled in the art

and would be suitable for use in the present invention. Other suitable methods are available to one skilled in the art, and it is to be understood that the present invention can be accomplished using any of the available methods of transfection. Lipofection can be accomplished by encapsulating an isolated DNA molecule within a liposomal particle and contacting the liposomal particle with the cell membrane of the target cell. Liposomes are self-assembling, colloidal particles in which a lipid bilayer, composed of amphiphilic molecules such as phosphatidyl serine or phosphatidyl choline, encapsulates a portion of the surrounding media such that the lipid bilayer surrounds a hydrophilic interior. Unilamellar or multilamellar liposomes can be constructed such that the interior contains a desired chemical, drug, or, as in the instant invention, an isolated DNA molecule.

[0179] In another aspect, polynucleotides encoding polypeptides of the invention may be used to express and deliver an AARS polypeptide via cell therapy. Accordingly in another aspect, the current invention includes a cell therapy for treating a disease or disorder, comprising administering a host cell expressing, or capable of expressing, an AARS polypeptide.

[0180] Cell therapy involves the administration of cells which have been selected, multiplied and pharmacologically treated or altered (i.e. genetically modified) outside of the body (Bordignon, C. et al, *Cell Therapy: Achievements and Perspectives* (1999), Haematologica, 84, pp. 1110-1149). Such host cells include for example, primary cells, including macrophages, and stem cells which have been genetically modified to express an AARS polypeptide. The aim of cell therapy is to replace, repair or enhance the biological function of damaged tissues or organs.

[0181] The use of transplanted cells has been investigated for the treatment of numerous endocrine disorders such as anemia and dwarfism, hematological disorders, kidney and liver failure, pituitary and CNS deficiencies and diabetes mellitus (Uludag et al., *Technology of Mammalian Cell Encapsulation* (2000), *Advanced Drug Delivery Reviews*, 42, pp. 29-64). Transplanted cells may function by releasing bioactive compounds such as an AARS polypeptide of the invention, to replace endogenous AARS polypeptides which are absent or produced in insufficient quantities in an effected system.

[0182] Embodiments of the present invention also include oligonucleotides, whether for detection, amplification, antisense therapies, or other purpose. For these and related purposes, the term "oligonucleotide" or "oligo" or "oligomer" is intended to encompass a singular "oligonucleotide" as well as plural "oligonucleotides," and refers to any polymer of two or more of nucleotides, nucleosides, nucleobases or related compounds used as a reagent in the amplification methods of the present invention, as well as subsequent detection methods. The oligonucleotide may be DNA and/or RNA and/or analogs thereof.

[0183] The term oligonucleotide does not necessarily denote any particular function to the reagent, rather, it is used generically to cover all such reagents described herein. An oligonucleotide may serve various different functions, e.g., it may function as a primer if it is capable of hybridizing to a complementary strand and can further be extended in the presence of a nucleic acid polymerase, it may provide a promoter if it contains a sequence recognized by an RNA polymerase and allows for transcription, and it may function to prevent hybridization or impede primer extension if

appropriately situated and/or modified. An oligonucleotide may also function as a probe, or an antisense agent. An oligonucleotide can be virtually any length, limited only by its specific function, e.g., in an amplification reaction, in detecting an amplification product of the amplification reaction, or in an antisense or RNA interference application. Any of the oligonucleotides described herein can be used as a primer, a probe, an antisense oligomer, or an RNA interference agent.

[0184] The term "primer" as used herein refers to a single-stranded oligonucleotide capable of acting as a point of initiation for template-directed DNA synthesis under suitable conditions defined, for example, by buffer and temperature, in the presence of four different nucleoside triphosphates and an agent for polymerization, such as a DNA or RNA polymerase or reverse transcriptase. The length of the primer, in any given case, depends on, for example, the intended use of the primer, and generally ranges from about 15 to 30 nucleotides, although shorter and longer primers may be used. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to hybridize with such template. The primer site is the area of the template to which a primer hybridizes. The primer pair is a set of primers including a 5' upstream primer that hybridizes with the 5' end of the sequence to be amplified and a 3' downstream primer that hybridizes with the complement of the 3' end of the sequence to be amplified.

[0185] The term "probe" as used herein includes a surface-immobilized or soluble but capable of being immobilized molecule that can be recognized by a particular target. See, e.g., U.S. Pat. No. 6,582,908 for an example of arrays having all possible combinations of probes with 10, 12, and more bases. Probes and primers as used herein typically comprise at least 10-15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 nucleotides of an AARS reference sequence or its complement. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the knowledge in the art and the specification, including the tables, figures, and Sequence Listing, may be used.

[0186] Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview N.Y.; Ausubel, F. M. et al. (1987) *Current Protocols in Molecular Biology*, Greene Publ. Assoc. & Wiley-Intersciences, New York N.Y.; Innis, M. et al. (1990) *PCR Protocols. A Guide to Methods and Applications*, Academic Press, San Diego Calif. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge Mass.).

[0187] Oligonucleotides for use as primers or probes may be selected using software known in the art. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up

to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas Tex.) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope.

[0188] The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge Mass.) allows the user to input a “mispriming library,” in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user’s specific needs.) The Prime-Gen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described herein.

[0189] In certain embodiments, oligonucleotides can be prepared by stepwise solid-phase synthesis, employing methods detailed in the references cited above, and below with respect to the synthesis of oligonucleotides having a mixture or uncharged and cationic backbone linkages. In some cases, it may be desirable to add additional chemical moieties to the oligonucleotide, e.g., to enhance pharmacokinetics or to facilitate capture or detection of the compound. Such a moiety may be covalently attached, typically to a terminus of the oligomer, according to standard synthetic methods. For example, addition of a polyethyleneglycol moiety or other hydrophilic polymer, e.g., one having 10-100 monomeric subunits, may be useful in enhancing solubility. One or more charged groups, e.g., anionic charged groups such as an organic acid, may enhance cell uptake.

[0190] A variety of detectable molecules may be used to render an oligonucleotide, or protein detectable, such as a radioisotopes, fluorochromes, dyes, enzymes, nanoparticles, chemiluminescent markers, biotin, or other monomer known in the art that can be detected directly (e.g., by light emission) or indirectly (e.g., by binding of a fluorescently-labeled antibody).

[0191] Radioisotopes provide examples of detectable molecules that can be utilized in certain aspects of the present invention. Several radioisotopes can be used as detectable molecules for labeling nucleotides or proteins, including, for example, ^{32}P , ^{33}P , ^{35}S , ^3H , and ^{125}I . These radioisotopes have different half-lives, types of decay, and levels of energy which can be tailored to match the needs of a particular protocol. For example, ^3H is a low energy emitter which

results in low background levels, however this low energy also results in long time periods for autoradiography. Radioactively labeled ribonucleotides, deoxyribonucleotides and amino acids are commercially available. Nucleotides are available that are radioactively labeled at the first, or α , phosphate group, or the third, or γ , phosphate group. For example, both [α - ^{32}P] dATP and [γ - ^{32}P] dATP are commercially available. In addition, different specific activities for radioactively labeled nucleotides are also available commercially and can be tailored for different protocols.

[0192] Other examples of detectable molecules that can be utilized to detect an oligonucleotide include fluorophores. Several fluorophores can be used for labeling nucleotides including, for example, fluorescein, tetramethylrhodamine, Texas Red, and a number of others (e.g., Haugland, *Handbook of Fluorescent Probes*-9th Ed., 2002, Molec. Probes, Inc., Eugene Oreg.; Haugland, *The Handbook: A Guide to Fluorescent Probes and Labeling Technologies*-10th Ed., 2005, Invitrogen, Carlsbad, Calif.).

[0193] As one example, oligonucleotides may be fluorescently labeled during chemical synthesis, since incorporation of amines or thiols during nucleotide synthesis permit addition of fluorophores. Fluorescently labeled nucleotides are commercially available. For example, uridine and deoxyuridine triphosphates are available that are conjugated to ten different fluorophores that cover the spectrum. Fluorescent dyes that can be bound directly to nucleotides can also be utilized as detectable molecules. For example, FAM, JOE, TAMRA, and ROX are amine reactive fluorescent dyes that have been attached to nucleotides and are used in automated DNA sequencing. These fluorescently labeled nucleotides, for example, ROX-ddATP, ROX-ddCTP, ROX-ddGTP and ROX-ddUTP, are commercially available.

[0194] Non-radioactive and non-fluorescent detectable molecules are also available. As noted above, biotin can be attached directly to nucleotides and detected by specific and high affinity binding to avidin or streptavidin which has been chemically coupled to an enzyme catalyzing a colorimetric reaction (such as phosphatase, luciferase, or peroxidase). Digoxigenin labeled nucleotides can also similarly be used for non-isotopic detection of nucleic acids. Biotinylated and digoxigenin-labeled nucleotides are commercially available.

[0195] Very small particles, termed nanoparticles, also can be used to label oligonucleotide probes. These particles range from 1-1000 nm in size and include diverse chemical structures such as gold and silver particles and quantum dots. When irradiated with angled incident white light, silver or gold nanoparticles ranging from 40-120 nm will scatter monochromatic light with high intensity. The wavelength of the scattered light is dependent on the size of the particle. Four to five different particles in close proximity will each scatter monochromatic light, which when superimposed will give a specific, unique color. The particles are being manufactured by companies such as Genicon Sciences (Carlsbad, Calif.). Derivatized silver or gold particles can be attached to a broad array of molecules including, proteins, antibodies, small molecules, receptor ligands, and nucleic acids. For example, the surface of the particle can be chemically derivatized to allow attachment to a nucleotide.

[0196] Other types of nanoparticles that can be used for detection of a detectable molecule include quantum dots. Quantum dots are fluorescing crystals 1-5 nm in diameter that are excitable by light over a large range of wavelengths. Upon excitation by light having an appropriate wavelength,

these crystals emit light, such as monochromatic light, with a wavelength dependent on their chemical composition and size. Quantum dots such as CdSe, ZnSe, InP, or InAs possess unique optical properties; these and similar quantum dots are available from a number of commercial sources (e.g., NN-Labs, Fayetteville, Ark.; Ocean Nanotech, Fayetteville, Ark.; Nanoco Technologies, Manchester, UK; Sigma-Aldrich, St. Louis, Mo.).

[0197] Many dozens of classes of particles can be created according to the number of size classes of the quantum dot crystals. The size classes of the crystals are created either 1) by tight control of crystal formation parameters to create each desired size class of particle, or 2) by creation of batches of crystals under loosely controlled crystal formation parameters, followed by sorting according to desired size and/or emission wavelengths. Two examples of references in which quantum dots are embedded within intrinsic silicon epitaxial layers of semiconductor light emitting/detecting devices are U.S. Pat. Nos. 5,293,050 and 5,354,707 to Chapple Sokol, et al.

[0198] In certain embodiments, oligonucleotide primers or probes may be labeled with one or more light-emitting or otherwise detectable dyes. The light emitted by the dyes can be visible light or invisible light, such as ultraviolet or infrared light. In exemplary embodiments, the dye may be a fluorescence resonance energy transfer (FRET) dye; a xanthene dye, such as fluorescein and rhodamine; a dye that has an amino group in the alpha or beta position (such as a naphthylamine dye, 1-dimethylaminonaphthyl-5-sulfonate, 1-anilino-8-naphthalene sulfonate and 2-p-touidiny-6-naphthalene sulfonate); a dye that has 3-phenyl-7-isocyanatocoumarin; an acridine, such as 9-isothiocyanatoacridine and acridine orange; a pyrene, a benzoxadiazole and a stilbene; a dye that has 3-(ϵ -carboxypentyl)-3'-ethyl-5,5'-dimethyloxycarbocyanine (CYA); 6-carboxy fluorescein (FAM); 5&6-carboxyrhodamine-110 (R110); 6-carboxyrhodamine-6G (R6G); N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA); 6-carboxy-X-rhodamine (ROX); 6-carboxy-4',5'-dichloro-2',7'-dimethoxyfluorescein (JOE); ALEXA FLUOR™; Cy2; Texas Red and Rhodamine Red; 6-carboxy-2',4,7,7'-tetrachlorofluorescein (TET); 6-carboxy-2',4,4',5',7,7'-hexachlorofluorescein (HEX); 5-carboxy-2',4',5',7'-tetrachlorofluorescein (ZOE); NAN; NED; Cy3; Cy3.5; Cy5; Cy5.5; Cy7; and Cy7.5; IR800CW, ICG, Alexa Fluor 350; Alexa Fluor 488; Alexa Fluor 532; Alexa Fluor 546; Alexa Fluor 568; Alexa Fluor 594; Alexa Fluor 647; Alexa Fluor 680, or Alexa Fluor 750.

[0199] The AARS polynucleotides and oligonucleotides of the present invention can be used in any of the therapeutic, diagnostic, research, or drug discovery compositions and methods described herein.

V. ANTIBODIES

[0200] According to another aspect, the present invention further provides antibodies that exhibit binding specificity for an AARS polypeptide, or its native cellular binding partner (i.e. cellular receptor, lipid, carbohydrate, protein, or nucleic acid binding partner), or complex thereof, and methods of using the same. The term antibody includes the various variations of the same, such as FABs, human antibodies, modified human antibodies, single chains, nonhuman antibodies, and other derivatives of the immunoglobulin fold that underlie immune system ligands for antigens, as described herein and known in the art. Antibodies can be

used in any of the therapeutic, diagnostic, drug discovery, or protein expression/purification methods and compositions provided herein.

[0201] Certain antibodies of the present invention differ from certain previously made antibodies because they can distinguish between the AARS protein fragments of Table(s) 1-3, or Table(s) 4-6, or Table(s) 7-9 and their corresponding full-length AARS, typically by binding with greater affinity to the AARS protein fragments than to the corresponding full-length AARS. Generally, such antibodies may bind to unique sequences or structures generated or revealed by splice variations, proteolysis, or other cellular processing that generates an AARS protein fragment of the invention (e.g., post translational processing, including but not limited to phosphorylation and other modifications that change protein structure). In some aspects the antibodies may bind to sequences around a unique splice junction (for example to one or more regions of at least 5 contiguous amino acids selected from the splice junction sequences listed in Tables 2B, 5B, or 8B, or alternatively to any amino acid sequence C-terminal of this splice site, for example as listed in Tables 2B, 5B, or 8B. For example, such antibodies may have binding specificity to one or more non-solvent exposed faces that are exposed in the AARS protein fragment but not in the full-length AARS, or sequences that are not found or are otherwise inaccessible in the full-length AARS. Antibodies may also bind to unique three-dimensional structures that result from differences in folding between the AARS protein fragment and the full-length AARS. Such differences in folding may be localized (e.g., to a specific domain or region) or globalized. As one example, folding of AARS protein fragments may generate unique continuous or discontinuous epitopes that are not found in the corresponding or parent AARS. Examples also include antibodies that specifically bind to N- or C-termini generated by splice variations, proteolysis, or other cellular processing; such termini may be unique compared to the full-length AARS or may not be exposed for antibody binding in the full-length versions due to their termini being completely or partially buried in the overall structure of the larger AARS parent molecule.

[0202] In some embodiments, antibodies provided herein do not form aggregates, have a desired solubility, and/or have an immunogenicity profile that is suitable for use in humans, as described herein and known in the art. Also included are antibodies that are suitable for production work, such as to purify the AARS protein fragments described herein. Preferably, active antibodies can be concentrated to at least about 10 mg/ml and optionally formulated for biotherapeutic uses.

[0203] In certain embodiments, antibodies are effective for modulating one or more of the non-canonical activities mediated by an AARS polypeptide of the invention. In certain embodiments, for example, the antibody is one that binds to an AARS polypeptide and/or its binding partner, inhibits their ability to interact with each other, and/or antagonizes the non-canonical activity of the AARS polypeptide. In certain embodiments, for example, the antibody binds to the cellular binding partner of an AARS polypeptide, and mimics the AARS polypeptide activity, such as by increasing or agonizing the non-canonical activity mediated by the AARS polypeptide. Accordingly, antibodies may be used to diagnose, treat, or prevent diseases, disorders or

other conditions that are mediated by an AARS polypeptide of the invention, such as by antagonizing or agonizing its activity partially or fully.

[0204] An antibody, or antigen-binding fragment thereof, is said to “specifically bind,” “immunologically bind,” and/or is “immunologically reactive” to a polypeptide of the invention if it reacts at a detectable level (within, for example, an ELISA assay) with the polypeptide, and does not react detectably in a statistically significant manner with unrelated polypeptides under similar conditions. In certain instances, a binding agent does not significantly interact with a full-length version of the AARS polypeptide.

[0205] Immunological binding, as used in this context, generally refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of binding such as immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. See, e.g., Davies et al. (1990) *Annual Rev. Biochem.* 59:439-473. In certain illustrative embodiments, an antibody has an affinity for an AARS protein fragment of at least about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM. In certain embodiments, the affinity of the antibody for an AARS protein fragment is stronger than its affinity for a corresponding full-length AARS polypeptide, typically by about 1.5×, 2×, 2.5×, 3×, 3.5×, 4×, 4.5×, 5×, 6×, 7×, 8×, 9×, 10×, 15×, 20×, 25×, 30×, 40×, 50×, 60×, 70×, 80×, 90×, 100×, 200×, 300×, 400×, 500×, 600×, 700×, 800×, 900×, 1000× or more (including all integers in between). In certain embodiments, an antibody as an affinity for a corresponding full-length AARS protein of at least about 0.05, 0.1, 0.25, 0.5, 0.75, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 μ M. In certain embodiments, an antibody binds weakly or substantially undetectably to a full-length AARS protein.

[0206] An “antigen-binding site,” or “binding portion” of an antibody, refers to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable (“V”) regions of the heavy (“H”) and light (“L”) chains. Three highly divergent stretches within the V regions of the heavy and light chains are referred to as “hypervariable regions” which are interposed between more conserved flanking stretches known as “framework regions,” or “FRs”. Thus the term “FR” refers to amino acid sequences which are naturally found between and adjacent to hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as “complementarity-determining regions,” or “CDRs.”

[0207] Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. Monoclonal antibodies

specific for a polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Also included are methods that utilize transgenic animals such as mice to express human antibodies. See, e.g., Neuberger et al., *Nature Biotechnology* 14:826, 1996; Lonberg et al., *Handbook of Experimental Pharmacology* 113:49-101, 1994; and Lonberg et al., *Internal Review of Immunology* 13:65-93, 1995. Particular examples include the VELOCIMMUNE® platform by REGERNEREX® (see, e.g., U.S. Pat. No. 6,596,541). Antibodies can also be generated or identified by the use of phage display or yeast display libraries (see, e.g., U.S. Pat. No. 7,244,592; Chao et al., *Nature Protocols* 1:755-768, 2006). Non-limiting examples of available libraries include cloned or synthetic libraries, such as the Human Combinatorial Antibody Library (HuCAL), in which the structural diversity of the human antibody repertoire is represented by seven heavy chain and seven light chain variable region genes. The combination of these genes gives rise to 49 frameworks in the master library. By superimposing highly variable genetic cassettes (CDRs=complementarity determining regions) on these frameworks, the vast human antibody repertoire can be reproduced. Also included are human libraries designed with human-donor-sourced fragments encoding a light-chain variable region, a heavy-chain CDR-3, synthetic DNA encoding diversity in heavy-chain CDR-1, and synthetic DNA encoding diversity in heavy-chain CDR-2. Other libraries suitable for use will be apparent to persons skilled in the art. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

[0208] An “Fv” fragment can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent $V_H::V_L$ heterodimer including an antigen-binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule. See, e.g., Inbar et al. (1972) *Proc. Nat. Acad. Sci. USA* 69:2659-2662; Hochman et al. (1976) *Biochem* 15:2706-2710; and Ehrlich et al. (1980) *Biochem* 19:4091-4096.

[0209] A single chain Fv (“sFv”) polypeptide is a covalently linked $V_H::V_L$ heterodimer which is expressed from a gene fusion including V_H - and V_L -encoding genes linked by a peptide-encoding linker. Huston et al. (1988) *PNAS USA* 85(16):5879-5883. A number of methods have been described to discern chemical structures for converting the naturally aggregated—but chemically separated—light and heavy polypeptide chains from an antibody V region into an sFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

[0210] Each of the above-described molecules includes a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain FR set which provide support to the CDRS and define the spatial relationship of the CDRs relative to each other. As used herein, the term “CDR set” refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are

denoted as “CDR1,” “CDR2,” and “CDR3” respectively. An antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (e.g., a CDR1, CDR2 or CDR3) is referred to herein as a “molecular recognition unit.” Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen-binding site.

[0211] As used herein, the term “FR set” refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRS. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, all V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain “canonical” structures—regardless of the precise CDR amino acid sequence. Further, certain FR residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

[0212] Certain embodiments include single domain antibody (sdAbs or “nanobodies”), which refer to an antibody fragment consisting of a single monomeric variable antibody domain (see, e.g., U.S. Pat. Nos. 5,840,526; 5,874,541; 6,005,079, 6,765,087, 5,800,988; 5,874,541; and 6,015,695). Such sdABs typically have a molecular weight of about 12-15 kDa. In certain aspects, sdABs and other antibody molecules can be derived or isolated from the unique heavy-chain antibodies of immunized camels and llamas, often referred to as camelids. See, e.g., Conrath et al., *JBC*. 276:7346-7350, 2001.

[0213] A number of “humanized” antibody molecules comprising an antigen-binding site derived from a non-human immunoglobulin have been described, including chimeric antibodies having rodent V regions and their associated CDRs fused to human constant domains (Winter et al. (1991) *Nature* 349:293-299; Lobuglio et al. (1989) *Proc. Nat. Acad. Sci. USA* 86:4220-4224; Shaw et al. (1987) *J Immunol.* 138:4534-4538; and Brown et al. (1987) *Cancer Res.* 47:3577-3583), rodent CDRs grafted into a human supporting FR prior to fusion with an appropriate human antibody constant domain (Riechmann et al. (1988) *Nature* 332:323-327; Verhoeven et al. (1988) *Science* 239:1534-1536; and Jones et al. (1986) *Nature* 321:522-525), and rodent CDRs supported by recombinantly veneered rodent FRs (European Patent Publication No. 519,596, published Dec. 23, 1992). These “humanized” molecules are designed to minimize unwanted immunological response toward rodent antihuman antibody molecules which limits the duration and effectiveness of therapeutic applications of those moieties in human recipients. See, e.g., U.S. Pat. Nos. 5,530,101; 5,585,089; 5,693,762; 6,180,370; and 7,022,500.

[0214] The antibodies of the present invention can be used in any of the therapeutic, diagnostic, drug discovery, protein purification, and analytical methods and compositions described herein.

VI. ANTIBODY ALTERNATIVES AND OTHER BINDING AGENTS

[0215] According to another aspect, the present invention further provides antibody alternatives or other binding agents, such as soluble receptors, adnectins, peptides, peptide mimetics, small molecules, aptamers, etc., that exhibit binding specificity for an AARS polypeptide or its cellular binding partner as disclosed herein, or to a portion, variant or derivative thereof, and compositions and methods of using same. Binding agents can be used in any of the therapeutic, diagnostic, drug discovery, or protein expression/purification, and analytical methods and compositions described herein. Biologic-based binding agents such as adnectins, soluble receptors, avimers, and trinectins are particularly useful.

[0216] In certain embodiments, such binding agents are effective for modulating one or more of the non-canonical activities mediated by an AARS polypeptide of the invention. In some embodiments, for example, the binding agent is one that binds to an AARS polypeptide and/or its binding partner, inhibits their ability to interact with each other, and/or antagonizes the non-canonical activity of the AARS polypeptide. In certain embodiments, for example, the binding agent binds to the cellular binding partner of an AARS polypeptide, and mimics the AARS polypeptide activity, such as by increasing or agonizing the non-canonical activity mediated by the AARS polypeptide. Accordingly, such binding agents may be used to diagnose, treat, or prevent diseases, disorders or other conditions that are mediated by an AARS polypeptide of the invention, such as by antagonizing or agonizing its activity partially or fully.

[0217] A binding agent is said to “specifically bind” to an AARS polypeptide of the invention, or its cellular binding partner, if it reacts at a detectable level (within, for example, an ELISA assay) with the polypeptide or its cellular binding partner, and does not react detectably in a statistically significant manner with unrelated polypeptides under similar conditions. In certain instances, a binding agent does not significantly interact with a full-length version of the AARS polypeptide. In certain illustrative embodiments, a binding agent has an affinity for an AARS protein fragment or its cellular binding partner of at least about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM. In certain embodiments, the affinity of the binding agent for an AARS protein fragment is stronger than its affinity for a corresponding full-length AARS polypeptide, typically by about 1.5x, 2x, 2.5x, 3x, 3.5x, 4x, 4.5x, 5x, 6x, 7x, 8x, 9x, 10x, 15x, 20x, 25x, 30x, 40x, 50x, 60x, 70x, 80x, 90x, 100x, 200x, 300x, 400x, 500x, 600x, 700x, 800x, 900x, 1000x or more (including all integers in between). In certain embodiments, a binding agent has an affinity for a corresponding full-length AARS protein of at least about 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 μ M.

[0218] As noted above, “peptides” are included as binding agents. The term peptide typically refers to a polymer of amino acid residues and to variants and synthetic analogues of the same. In certain embodiments, the term “peptide”

refers to relatively short polypeptides, including peptides that consist of about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, or 50 amino acids, including all integers and ranges (e.g., 5-10, 8-12, 10-15) in between, and interact with an AARS polypeptide, its cellular binding partner, or both. Peptides can be composed of naturally-occurring amino acids and/or non-naturally occurring amino acids, as described herein.

[0219] In addition to peptides consisting only of naturally-occurring amino acids, peptidomimetics or peptide analogs are also provided. Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed “peptide mimetics” or “peptidomimetics” (Luthman, et al., *A Textbook of Drug Design and Development*, 14:386-406, 2nd Ed., Harwood Academic Publishers (1996); Joachim Grante, *Angew. Chem. Int. Ed. Engl.*, 33:1699-1720 (1994); Fauchere, J., *Adv. Drug Res.*, 15:29 (1986); Veber and Freidinger *TINS*, p. 392 (1985); and Evans, et al., *J. Med. Chem.* 30:229 (1987)). A peptidomimetic is a molecule that mimics the biological activity of a peptide but is no longer peptidic in chemical nature. Peptidomimetic compounds are known in the art and are described, for example, in U.S. Pat. No. 6,245,886.

[0220] The present invention also includes peptoids. Peptoid derivatives of peptides represent another form of modified peptides that retain the important structural determinants for biological activity, yet eliminate the peptide bonds, thereby conferring resistance to proteolysis (Simon, et al., *PNAS USA*, 89:9367-9371, 1992). Peptoids are oligomers of N-substituted glycines. A number of N-alkyl groups have been described, each corresponding to the side chain of a natural amino acid. The peptidomimetics of the present invention include compounds in which at least one amino acid, a few amino acids or all amino acid residues are replaced by the corresponding N-substituted glycines. Peptoid libraries are described, for example, in U.S. Pat. No. 5,811,387.

[0221] A binding agent may also include one or more small molecules. A “small molecule” refers to an organic compound that is of synthetic or biological origin (biomolecule), but is typically not a polymer. Organic compounds refer to a large class of chemical compounds whose molecules contain carbon, typically excluding those that contain only carbonates, simple oxides of carbon, or cyanides. A “biomolecule” refers generally to an organic molecule that is produced by a living organism, including large polymeric molecules (biopolymers) such as peptides, polysaccharides, and nucleic acids as well, and small molecules such as primary secondary metabolites, lipids, phospholipids, glycolipids, sterols, glycerolipids, vitamins, and hormones. A “polymer” refers generally to a large molecule or macromolecule composed of repeating structural units, which are typically connected by covalent chemical bond.

[0222] In certain embodiments, a small molecule has a molecular weight of less than 1000-2000 Daltons, typically between about 300 and 700 Daltons, and including about 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 or 2000 Daltons. Small molecule libraries are described elsewhere herein.

[0223] Aptamers are also included as binding agents (see, e.g., Ellington et al., *Nature*, 346, 818-22, 1990; and Tuerk et al., *Science*, 249, 505-10, 1990). Examples of aptamers

included nucleic acid aptamers (e.g., DNA aptamers, RNA aptamers) and peptide aptamers. Nucleic acid aptamers refer generally to nucleic acid species that have been engineered through repeated rounds of in vitro selection or equivalent method, such as SELEX (systematic evolution of ligands by exponential enrichment), to bind to various molecular targets such as small molecules, proteins, nucleic acids, and even cells, tissues and organisms. See, e.g., U.S. Pat. Nos. 6,376,190; and 6,387,620 Hence, included are nucleic acid aptamers that bind to the AARS polypeptides described herein and/or their cellular binding partners.

[0224] Peptide aptamers typically include a variable peptide loop attached at both ends to a protein scaffold, a double structural constraint that typically increases the binding affinity of the peptide aptamer to levels comparable to that of an antibody's (e.g., in the nanomolar range). In certain embodiments, the variable loop length may be composed of about 10-20 amino acids (including all integers in between), and the scaffold may include any protein that has good solubility and compacity properties. Certain exemplary embodiments may utilize the bacterial protein Thioredoxin-A as a scaffold protein, the variable loop being inserted within the reducing active site (-Cys-Gly-Pro-Cys-loop in the wild protein), with the two cysteines lateral chains being able to form a disulfide bridge. Methods for identifying peptide aptamers are described, for example, in U.S. Application No. 2003/0108532. Hence, included are peptide aptamers that bind to the AARS polypeptides described herein and/or their cellular binding partners. Peptide aptamer selection can be performed using different systems known in the art, including the yeast two-hybrid system.

[0225] Also included are ADNECTINS™, AVIMERS™, anaphones and anticalins that specifically bind to an AARS protein fragment of the invention. ADNECTINS™ refer to a class of targeted biologics derived from human fibronectin, an abundant extracellular protein that naturally binds to other proteins. See, e.g., U.S. Application Nos. 2007/0082365; 2008/0139791; and 2008/0220049. ADNECTINS™ typically consists of a natural fibronectin backbone, as well as the multiple targeting domains of a specific portion of human fibronectin. The targeting domains can be engineered to enable an ADNECTIN™ to specifically recognize a therapeutic target of interest, such as an AARS protein fragment of the invention.

[0226] AVIMERS™ refer to multimeric binding proteins or peptides engineered using in vitro exon shuffling and phage display. Multiple binding domains are linked, resulting in greater affinity and specificity compared to single epitope immunoglobulin domains. See, e.g., Silverman et al., *Nature Biotechnology*, 23:1556-1561, 2005; U.S. Pat. No. 7,166,697; and U.S. Application Nos. 2004/0175756, 2005/0048512, 2005/0053973, 2005/0089932 and 2005/0221384.

[0227] Also included are designed ankyrin repeat proteins (DARPs), which include a class of non-immunoglobulin proteins that can offer advantages over antibodies for target binding in drug discovery and drug development. Among other uses, DARPs are ideally suited for in vivo imaging or delivery of toxins or other therapeutic payloads because of their favorable molecular properties, including small size and high stability. The low-cost production in bacteria and the rapid generation of many target-specific DARPs make the DARPin approach useful for drug discovery. Addition-

ally, DARPin can be easily generated in multispecific formats, offering the potential to target an effector DARPin to a specific organ or to target multiple receptors with one molecule composed of several DARPins. See, e.g., Stumpp et al., *Curr Opin Drug Discov Devel.* 10:153-159, 2007; U.S. Application No. 2009/0082274; and PCT/EP2001/10454.

[0228] Certain embodiments include “monobodies,” which typically utilize the 10th fibronectin type III domain of human fibronectin (FNfn10) as a scaffold to display multiple surface loops for target binding. FNfn10 is a small (94 residues) protein with a β -sandwich structure similar to the immunoglobulin fold. It is highly stable without disulfide bonds or metal ions, and it can be expressed in the correctly folded form at a high level in bacteria. The FNfn10 scaffold is compatible with virtually any display technologies. See, e.g., Baton et al., *Protein Eng.* 15:1015-20, 2002; and Wojcik et al., *Nat Struct Mol Biol.*, 2010; and U.S. Pat. No. 6,673,901.

[0229] Anticalins refer to a class of antibody mimetics, which are typically synthesized from human lipocalins, a family of binding proteins with a hypervariable loop region supported by a structurally rigid framework. See, e.g., U.S. Application No. 2006/0058510. Anticalins typically have a size of about 20 kDa. Anticalins can be characterized by a barrel structure formed by eight antiparallel β -strands (a stable β -barrel scaffold) that are pairwise connected by four peptide loops and an attached α -helix. In certain aspects, conformational deviations to achieve specific binding are made in the hypervariable loop region(s). See, e.g., Skerra, *FEBS J.* 275:2677-83, 2008, herein incorporated by reference.

VII. BIOASSAYS AND ANALYTICAL ASSAYS FOR DRUG RELEASE ASSAYS AND PRODUCT SPECIFICATIONS, DIAGNOSTICS, AND REAGENTS

[0230] Also included are bioassays that relate to the AARS protein fragments and related agents as therapeutic and diagnostic reagents. Examples include bioassays and analytical assays that measure purity, biological activity, affinity, solubility, pH, endotoxin levels, among others, many of which are described herein. Also included are assays that establish dose response curves and/or provide one or more bases for comparison between different batches of agents. Batch comparisons can be based on any one or more of chemical characterization, biological characterization, and clinical characterization. For protein agents, also included are methods of evaluating the potency, stability, pharmacokinetics, and immunogenicity of a selected agent. Among other uses, these and other methods can be used for lot releasing testing of biologic or chemical agents, including the AARS protein fragments, antibodies, binding agents, polynucleotides such as antisense agents and vectors, and others described herein.

[0231] Certain embodiments include the use of bioaffinity assays. Such assays can be used to assess the binding affinity, for example, between an AARS protein fragment and a cellular binding partner, or between an AARS protein fragment and an antibody. Binding affinity can also be measured between an AARS protein fragment and an alternate binding agent such as a candidate or lead test compound (e.g., small molecule modulator of an AARS), or between an AARS cellular binding partner and a candidate or lead test

compound. Certain exemplary binding affinity assays may utilize ELISA assays, as described herein and known in the art. Certain assays utilize high-performance receptor binding chromatography (see, e.g., Roswall et al., *Biologicals.* 24:25-39, 1996). Other exemplary binding affinity assays may utilize surface plasmon resonance (SPR)-based technologies. Examples include BIACore technologies, certain of which integrate SPR technology with a microfluidics system to monitor molecular interactions in real time at concentrations ranging from pM to mM. Also included are KINEXA™ assays, which provide accurate measurements of binding specificity, binding affinity, and binding kinetics/rate constants.

[0232] Certain embodiments relate to immunoassays for evaluating or optimizing the immunogenicity of protein agents. Examples include ex vivo human cellular assays and in vitro immuno-enzymatic assays to provide useful information on the immunogenic potential of a therapeutic protein. Ex vivo cell-response assays can be used, for example, to reproduce the cellular co-operation between antigen-presenting cells (APCs) and T-cells, and thereby measure T-cells activation after contact with a protein of interest. Certain in vitro enzymatic assays may utilize a collection of recombinant HLA-DR molecules that cover a significant portion of a relevant human population, and may include automated immuno-enzymatic assays for testing the binding of peptides (stemming from the fragmentation of the therapeutic protein) with the HLA-DR molecules. Also included are methods of reducing the immunogenicity of a selected protein, such as by using these and related methods to identify and then remove or alter one or more T-cell epitopes from a protein agent.

[0233] Also included are biological release assays (e.g., cell-based assays) for measuring parameters such as specific biological activities, including non-canonical biological activities, and cytotoxicity. Certain specific biological assays include, for example, cell-based assays that utilize a cellular binding partner (e.g., cell-surface receptor) of a selected AARS protein fragment, which is functionally coupled to a readout, such as a fluorescent or luminescent indicator of a non-canonical biological activity, as described herein. For instance, specific embodiments include a cell that comprises a cell-surface receptor or an extracellular portion thereof that binds to an AARS protein fragment, wherein the cell comprises a detector or readout. Also included are in vivo biological assays to characterize the pharmacokinetics of an agent, such as an AARS polypeptide or antibody, typically utilizing engineered mice or other mammal (see, e.g., Lee et al., *The Journal of Pharmacology.* 281:1431-1439, 1997). Examples of cytotoxicity-based biological assays include release assays (e.g., chromium or europium release assays to measure apoptosis; see, e.g., von Zons et al., *Clin Diagn Lab Immunol.* 4:202-207, 1997), among others, which can assess the cytotoxicity of AARS protein fragments, whether for establishing dose response curves, batch testing, or other properties related to approval by various regulatory agencies, such as the Food and Drug Administration (FDA).

[0234] Such assays can be used, for example, to develop a dose response curve for a selected AARS protein fragment or other agent, and/or to compare the dose response curve of different batches of proteins or other agents. A dose-response curve is an X-Y graph that relates the magnitude of a stressor to the response of a receptor; the response may be a

physiological or biochemical response, such as a non-canonical biological activity in a cell in vitro or in a cell or tissue in vivo, a therapeutically effective amount as measured in vivo (e.g., as measured by EC_{50}), or death, whether measured in vitro or in vivo (e.g., cell death, organismal death). Death is usually indicated as an LD_{50} , a statistically-derived dose that is lethal to 50% of a modeled population, though it can be indicated by LC_{01} (lethal dose for 1% of the animal test population), LC_{100} (lethal dose for 100% of the animal test population), or LC_{LO} (lowest dose causing lethality). Almost any desired effect or endpoint can be characterized in this manner.

[0235] The measured dose of a response curve is typically plotted on the X axis and the response is plotted on the Y axis. More typically, the logarithm of the dose is plotted on the X axis, most often generating a sigmoidal curve with the steepest portion in the middle. The No Observable Effect Level (NOEL) refers to the lowest experimental dose for which no measurable effect is observed, and the threshold dose refers to the first point along the graph that indicates a response above zero. As a general rule, stronger drugs generate steeper dose response curves. For many drugs, the desired effects are found at doses slightly greater than the threshold dose, often because lower doses are relatively ineffective and higher doses lead to undesired side effects. For in vivo generated dose response curves, a curve can be characterized by values such as $\mu\text{g/kg}$, mg/kg , or g/kg of body-weight, if desired.

[0236] For batch comparisons, it can be useful to calculate the coefficient of variation (CV) between different dose response curves of different batches (e.g., between different batches of AARS protein fragments, antibodies, or other agents), in part because the CV allows comparison between data sets with different units or different means. For instance, in certain exemplary embodiments, two or three or more different batches of AARS protein fragments or other agents have a CV between them of less than about 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, or 1% for a 4, 5, 6, 7, or 8 point dose curve. In certain embodiments, the dose response curve is measured in a cell-based assay, and its readout relates to an increase or a decrease in a selected non-canonical activity of the AARS protein fragment. In certain embodiments, the dose response curve is measured in a cell release assay or animal model (e.g., mouse model), and its readout relates to cell death or animal death. Other variations will be apparent to persons skilled in the art.

VIII. EXPRESSION AND PURIFICATION SYSTEMS

[0237] Embodiments of the present invention include methods and related compositions for expressing and purifying the AARS protein fragments or other polypeptide-based agents of the invention. Such recombinant AARS polypeptides can be conveniently prepared using standard protocols as described for example in Sambrook, et al., (1989, *supra*), in particular Sections 16 and 17; Ausubel et al., (1994, *supra*), in particular Chapters 10 and 16; and Coligan et al., *Current Protocols in Protein Science* (John Wiley & Sons, Inc. 1995-1997), in particular Chapters 1, 5 and 6. As one general example, AARS polypeptides may be prepared by a procedure including one or more of the steps of: (a) preparing a construct comprising a polynucleotide sequence that encodes a AARS polypeptide and that is

operably linked to a regulatory element; (b) introducing the construct into a host cell; (c) culturing the host cell to express the AARS polypeptide; and (d) isolating the AARS polypeptide from the host cell.

[0238] AARS polynucleotides are described elsewhere herein. In order to express a desired polypeptide, a nucleotide sequence encoding the polypeptide, or a functional equivalent, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook et al., *Molecular Cloning, A Laboratory Manual* (1989), and Ausubel et al., *Current Protocols in Molecular Biology* (1989).

[0239] A variety of expression vector/host systems are known and may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems, including mammalian cell and more specifically human cell systems.

[0240] The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector—enhancers, promoters, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUE-SCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSPORT1 plasmid (Gibco BRL, Gaithersburg, Md.) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

[0241] In bacterial systems, a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster, *J. Biol.*

Chem. 264:5503-5509 (1989)); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[0242] Certain embodiments may employ *E. coli*-based expression systems (see, e.g., Structural Genomics Consortium et al., *Nature Methods*. 5:135-146, 2008). These and related embodiments may rely partially or totally on ligation-independent cloning (LIC) to produce a suitable expression vector. In specific embodiments, protein expression may be controlled by a T7 RNA polymerase (e.g., pET vector series). These and related embodiments may utilize the expression host strain BL21(DE3), a λ DE3 lysogen of BL21 that supports T7-mediated expression and is deficient in lon and ompT proteases for improved target protein stability. Also included are expression host strains carrying plasmids encoding tRNAs rarely used in *E. coli*, such as ROSETTA™ (DE3) and Rosetta 2 (DE3) strains. Cell lysis and sample handling may also be improved using reagents sold under the trademarks BENZONASE® nuclease and BUGBUSTER® Protein Extraction Reagent. For cell culture, auto-inducing media can improve the efficiency of many expression systems, including high-throughput expression systems. Media of this type (e.g., OVERNIGHT EXPRESS™ Autoinduction System) gradually elicit protein expression through metabolic shift without the addition of artificial inducing agents such as IPTG. Particular embodiments employ hexahistidine tags (such as those sold under the trademark HIS•TAG® fusions), followed by immobilized metal affinity chromatography (IMAC) purification, or related techniques. In certain aspects, however, clinical grade proteins can be isolated from *E. coli* inclusion bodies, without or without the use of affinity tags (see, e.g., Shimp et al., *Protein Expr Purif.* 50:58-67, 2006). As a further example, certain embodiments may employ a cold-shock induced *E. coli* high-yield production system, because over-expression of proteins in *Escherichia coli* at low temperature improves their solubility and stability (see, e.g., Qing et al., *Nature Biotechnology*. 22:877-882, 2004).

[0243] Also included are high-density bacterial fermentation systems. For example, high cell density cultivation of *Ralstonia eutropha* allows protein production at cell densities of over 150 g/L, and the expression of recombinant proteins at titers exceeding 10 g/L.

[0244] In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al., *Methods Enzymol.* 153:516-544 (1987). Also included are *Pichia pandoris* expression systems (see, e.g., Li et al., *Nature Biotechnology*. 24, 210-215, 2006; and Hamilton et al., *Science*, 301:1244, 2003). Certain embodiments include yeast systems that are engineered to selectively glycosylate proteins, including yeast that have humanized N-glycosylation pathways, among others (see, e.g., Hamilton et al., *Science*. 313:1441-1443, 2006; Wildt et al., *Nature Reviews Microbiol.* 3:119-28, 2005; and Gerngross et al., *Nature-*

Biotechnology. 22:1409-1414, 2004; U.S. Pat. Nos. 7,629,163; 7,326,681; and 7,029,872). Merely by way of example, recombinant yeast cultures can be grown in Fernbach Flasks or 15 L, 50 L, 100 L, and 200 L fermentors, among others.

[0245] In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6:307-311 (1987)). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi et al., *EMBO J.* 3:1671-1680 (1984); Broglie et al., *Science* 224:838-843 (1984); and Winter et al., *Results Probl. Cell Differ.* 17:85-105 (1991)). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, e.g., Hobbs in McGraw Hill, *Yearbook of Science and Technology*, pp. 191-196 (1992)).

[0246] An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia ni* cells. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia ni* cells in which the polypeptide of interest may be expressed (Engelhard et al., *Proc. Natl. Acad. Sci. U.S.A.* 91:3224-3227 (1994)). Also included are baculovirus expression systems, including those that utilize SF9, SF21, and *T. ni* cells (see, e.g., Murphy and Piwnicka-Worms, *Curr Protoc Protein Sci.* Chapter 5:Unit5.4, 2001). Insect systems can provide post-translation modifications that are similar to mammalian systems.

[0247] In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci. U.S.A.* 81:3655-3659 (1984)). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

[0248] Examples of useful mammalian host cell lines include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells sub-cloned for growth in suspension culture, Graham et al., *J. Gen Virol.* 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); mouse sertoli cells (TM4, Mather, *Biol. Reprod.* 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells

(BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TR1 cells (Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2). Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR-CHO cells (Urlaub et al., *PNAS USA* 77:4216 (1980)); and myeloma cell lines such as NSO and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki and Wu, *Methods in Molecular Biology*, Vol. 248 (B. K.C. Lo, ed., Humana Press, Totowa, N.J., 2003), pp. 255-268. Certain preferred mammalian cell expression systems include CHO and HEK293-cell based expression systems. Mammalian expression systems can utilize attached cell lines, for example, in T-flasks, roller bottles, or cell factories, or suspension cultures, for example, in 1 L and 5 L spinners, 5 L, 14 L, 40 L, 100 L and 200 L stir tank bioreactors, or 20/50 L and 100/200 L WAVE bioreactors, among others known in the art.

[0249] Also included is cell-free expression of proteins. These and related embodiments typically utilize purified RNA polymerase, ribosomes, tRNA and ribonucleotides; these reagents may be produced by extraction from cells or from a cell-based expression system.

[0250] Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf et al., *Results Probl. Cell Differ.* 20:125-162 (1994)).

[0251] In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, post-translational modifications such as acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as yeast, CHO, HeLa, MDCK, HEK293, and W138, in addition to bacterial cells, which have or even lack specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

[0252] For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous

expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type. Transient production, such as by transient transfection or infection, can also be employed. Exemplary mammalian expression systems that are suitable for transient production include HEK293 and CHO-based systems.

[0253] Any number of selection systems may be used to recover transformed or transduced cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223-232 (1977)) and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817-823 (1990)) genes which can be employed in tk- or apt-cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler et al., *Proc. Natl. Acad. Sci. U.S.A.* 77:3567-70 (1980)); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin et al., *J. Mol. Biol.* 150:1-14 (1981)); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, supra). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. U.S.A.* 85:8047-51 (1988)). The use of visible markers has gained popularity with such markers as green fluorescent protein (GFP) and other fluorescent proteins (e.g., RFP, YFP), anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (see, e.g., Rhodes et al., *Methods Mol. Biol.* 55:121-131 (1995)).

[0254] Embodiments of the present invention also include high-throughput protein production systems, or micro-production systems. Certain aspects may utilize, for example, hexa-histidine fusion tags for protein expression and purification on metal chelate-modified slide surfaces or Magne-His Ni-Particles (see, e.g., Kwon et al., *BMC Biotechnol.* 9:72, 2009; and Lin et al., *Methods Mol Biol.* 498:129-41, 2009)). Also included are high-throughput cell-free protein expression systems (see, e.g., Sitaraman et al., *Methods Mol Biol.* 498:229-44, 2009). These and related embodiments can be used, for example, to generate microarrays of AARS protein fragment(s), which can then be used for screening libraries to identify agents that interact with the AARS protein fragment(s).

[0255] A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using binding agents or antibodies such as polyclonal or monoclonal antibodies specific for the product, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), western immunoblots, radioimmunoassays (RIA), and fluorescence activated cell sorting (FACS). These and other assays are described, among other places, in Hampton et al., *Serological Methods, a Laboratory Manual* (1990) and Maddox et al., *J. Exp. Med.* 158:1211-1216 (1983).

[0256] A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

[0257] Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. Certain specific embodiments utilize serum free cell expression systems. Examples include HEK293 cells and CHO cells that can be grown on serum free medium (see, e.g., Rosser et al., *Protein Expr. Purif* 40:237-43, 2005; and U.S. Pat. No. 6,210,922).

[0258] The protein produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides of the invention may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification and/or detection of soluble proteins. Examples of such domains include cleavable and non-cleavable affinity purification and epitope tags such as avidin, FLAG tags, poly-histidine tags (e.g., 6xHis), cMyc tags, V5-tags, glutathione S-transferase (GST) tags, and others.

[0259] The protein produced by a recombinant cell can be purified and characterized according to a variety of techniques known in the art. Exemplary systems for performing protein purification and analyzing protein purity include fast protein liquid chromatography (FPLC) (e.g., AKTA and Bio-Rad FPLC systems), high-pressure liquid chromatography (HPLC) (e.g., Beckman and Waters HPLC). Exemplary chemistries for purification include ion exchange chromatography (e.g., Q, S), size exclusion chromatography, salt gradients, affinity purification (e.g., Ni, Co, FLAG, maltose, glutathione, protein A/G), gel filtration, reverse-phase, ceramic HYPERD® ion exchange chromatography, and hydrophobic interaction columns (HIC), among others known in the art. Also included are analytical methods such as SDS-PAGE (e.g., coomassie, silver stain), immunoblot, Bradford, and ELISA, which may be utilized during any step of the production or purification process, typically to measure the purity of the protein composition.

[0260] Also included are methods of concentrating AARS protein fragments, and composition comprising concentrated soluble proteins. In different aspects such concentrated solutions of AARS polypeptides may comprise pro-

teins at a concentration of about 5 mg/mL; or about 8 mg/mL; or about 10 mg/mL; about 15 mg/mL; or about 20 mg/mL.

[0261] In one aspect such compositions may be substantially monodisperse, meaning that the AARS polypeptide compositions exist primarily (i.e. at least about 90%, or greater) in one apparent molecular weight form when assessed for example, by size exclusion chromatography, dynamic light scattering, or analytical ultracentrifugation.

[0262] In another aspect, such compositions have a purity (on a protein basis) of at least about 90%, or in some aspects at least about 95% purity, or in some embodiments, at least 98% purity. Purity may be determined via any routine analytical method as known in the art.

[0263] In another aspect, such compositions have a high molecular weight aggregate content of less than about 10%, compared to the total amount of protein present, or in some embodiments such compositions have a high molecular weight aggregate content of less than about 5%, or in some aspects such compositions have a high molecular weight aggregate content of less than about 3%, or in some embodiments a high molecular weight aggregate content of less than about 1%. High molecular weight aggregate content may be determined via a variety of analytical techniques including for example, by size exclusion chromatography, dynamic light scattering, or analytical ultracentrifugation.

[0264] In certain embodiments, as noted herein, the AARS polypeptide compositions have an endotoxin content of less than about 10 EU/mg of AARS polypeptide, or less than about 5 EU/mg of AARS polypeptide, less than about 3 EU/mg of AARS polypeptide, or less than about 1 EU/mg of AARS polypeptide.

[0265] Examples of concentration approaches contemplated herein include lyophilization, which is typically employed when the solution contains few soluble components other than the protein of interest. Lyophilization is often performed after HPLC run, and can remove most or all volatile components from the mixture. Also included are ultrafiltration techniques, which typically employ one or more selective permeable membranes to concentrate a protein solution. The membrane allows water and small molecules to pass through and retains the protein; the solution can be forced against the membrane by mechanical pump, gas pressure, or centrifugation, among other techniques.

[0266] In certain embodiments, the reagents, AARS protein fragments, or related agents (e.g., antibodies) have a purity of at least about 90%, as measured according to routine techniques in the art. In certain embodiments, such as diagnostic compositions or certain therapeutic compositions, the AARS compositions of the present invention have a purity of at least about 95%. In specific embodiments, such as therapeutic or pharmaceutical compositions, the AARS compositions of the present invention have a purity of at least about 97% or 98% or 99%. In other embodiments, such as when being used as reference or research reagents, AARS protein fragments can be of lesser purity, and may have a purity of at least about 50%, 60%, 70%, or 80%. Purity can be measured overall or in relation to selected components, such as other proteins, e.g., purity on a protein basis.

[0267] Purified AARS protein fragments can also be characterized according to their biological characteristics. Examples include binding affinity or binding kinetics to a selected ligand (e.g., a cellular binding partner of the AARS protein fragment such as a cell-surface receptor or an

extracellular domain thereof), and the presence or levels of one or more canonical or non-canonical biological activity, as described herein. Binding affinity and binding kinetics can be measured according to a variety of techniques known in the art, such as Biacore® and related technologies that utilize surface plasmon resonance (SPR), an optical phenomenon that enables detection of unlabeled interactants in real time. SPR-based biosensors can be used in determination of active concentration, screening and characterization in terms of both affinity and kinetics. The presence or levels of one or more canonical or non-canonical biological activities can be measured according to cell-based assays, including those that utilize a cellular binding partner (e.g., cell-surface receptor) of a selected AARS protein fragment, which is functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator of a non-canonical biological activity, as described herein.

[0268] In certain embodiments, as noted above, the AARS polypeptide compositions are about substantially endotoxin free, including, for example, about 95% endotoxin free, preferably about 99% endotoxin free, and more preferably about 99.99% endotoxin free. The presence of endotoxins can be detected according to routine techniques in the art, as described herein. In specific embodiments, the AARS compositions are made from a eukaryotic cell such as a mammalian or human cell in substantially serum free media.

[0269] In certain embodiments, the AARS polypeptide compositions comprise less than about 10% wt/wt high molecular weight aggregates, or less than about 5% wt/wt high molecular weight aggregates, or less than about 2% wt/wt high molecular weight aggregates, or less than about or less than about 1% wt/wt high molecular weight aggregates.

[0270] Also included are protein-based analytical assays and methods, which can be used to assess, for example, protein purity, size, solubility, and degree of aggregation, among other characteristics. Protein purity can be assessed a number of ways. For instance, purity can be assessed based on primary structure, higher order structure, size, charge, hydrophobicity, and glycosylation. Examples of methods for assessing primary structure include N- and C-terminal sequencing and peptide-mapping (see, e.g., Allen et al., *Biologicals*. 24:255-275, 1996)). Examples of methods for assessing higher order structure include circular dichroism (see, e.g., Kelly et al., *Biochim Biophys Acta*. 1751:119-139, 2005), fluorescent spectroscopy (see, e.g., Meagher et al., *J. Biol. Chem.* 273:23283-89, 1998), FT-IR, amide hydrogen-deuterium exchange kinetics, differential scanning calorimetry, NMR spectroscopy, immunoreactivity with conformationally sensitive antibodies. Higher order structure can also be assessed as a function of a variety of parameters such as pH, temperature, or added salts. Examples of methods for assessing protein characteristics such as size include analytical ultracentrifugation and size exclusion HPLC (SEC-HPLC), and exemplary methods for measuring charge include ion-exchange chromatography and isoelectric focusing. Hydrophobicity can be assessed, for example, by reverse-phase HPLC and hydrophobic interaction chromatography HPLC. Glycosylation can affect pharmacokinetics (e.g., clearance), conformation or stability, receptor binding, and protein function, and can be assessed, for example, by mass spectrometry and nuclear magnetic resonance (NMR) spectroscopy.

[0271] As noted above, certain embodiments include the use of SEC-HPLC to assess protein characteristics such as purity, size (e.g., size homogeneity) or degree of aggregation, and/or to purify proteins, among other uses. SEC, also including gel-filtration chromatography (GFC) and gel-permeation chromatography (GPC), refers to a chromatographic method in which molecules in solution are separated in a porous material based on their size, or more specifically their hydrodynamic volume, diffusion coefficient, and/or surface properties. The process is generally used to separate biological molecules, and to determine molecular weights and molecular weight distributions of polymers. Typically, a biological or protein sample (such as a protein extract produced according to the protein expression methods provided herein and known in the art) is loaded into a selected size-exclusion column with a defined stationary phase (the porous material), preferably a phase that does not interact with the proteins in the sample. In certain aspects, the stationary phase is composed of inert particles packed into a dense three-dimensional matrix within a glass or steel column. The mobile phase can be pure water, an aqueous buffer, an organic solvent, or a mixture thereof. The stationary-phase particles typically have small pores and/or channels which only allow molecules below a certain size to enter. Large particles are therefore excluded from these pores and channels, and their limited interaction with the stationary phase leads them to elute as a "totally-excluded" peak at the beginning of the experiment. Smaller molecules, which can fit into the pores, are removed from the flowing mobile phase, and the time they spend immobilized in the stationary-phase pores depends, in part, on how far into the pores they penetrate. Their removal from the mobile phase flow causes them to take longer to elute from the column and results in a separation between the particles based on differences in their size. A given size exclusion column has a range of molecular weights that can be separated. Overall, molecules larger than the upper limit will not be trapped by the stationary phase, molecules smaller than the lower limit will completely enter the solid phase and elute as a single band, and molecules within the range will elute at different rates, defined by their properties such as hydrodynamic volume. For examples of these methods in practice with pharmaceutical proteins, see Bruner et al., *Journal of Pharmaceutical and Biomedical Analysis*. 15: 1929-1935, 1997.

[0272] Protein purity for clinical applications is also discussed, for example, by Anicetti et al. (*Trends in Biotechnology*. 7:342-349, 1989). More recent techniques for analyzing protein purity include, without limitation, the LabChip GXII, an automated platform for rapid analysis of proteins and nucleic acids, which provides high throughput analysis of titer, sizing, and purity analysis of proteins. In certain non-limiting embodiments, clinical grade proteins such as protein fragments and antibodies can be obtained by utilizing a combination of chromatographic materials in at least two orthogonal steps, among other methods (see, e.g., *Therapeutic Proteins: Methods and Protocols*. Vol. 308, Eds., Smales and James, Humana Press Inc., 2005). Typically, protein agents (e.g., AARS protein fragments, antibodies, binding agents) and other agents (e.g., antisense, RNAi, small molecules) are substantially endotoxin-free, as measured according to techniques known in the art and described herein.

[0273] Protein solubility assays are also included. Such assays can be utilized, for example, to determine optimal

growth and purification conditions for recombinant production, to optimize the choice of buffer(s), and to optimize the choice of AARS protein fragments or variants thereof. Solubility or aggregation can be evaluated according to a variety of parameters, including temperature, pH, salts, and the presence or absence of other additives. Examples of solubility screening assays include, without limitation, microplate-based methods of measuring protein solubility using turbidity or other measure as an end point, high-throughput assays for analysis of the solubility of purified recombinant proteins (see, e.g., Stenvall et al., *Biochim Biophys Acta*. 1752:6-10, 2005), assays that use structural complementation of a genetic marker protein to monitor and measure protein folding and solubility in vivo (see, e.g., Wigley et al., *Nature Biotechnology*. 19:131-136, 2001), and electrochemical screening of recombinant protein solubility in *Escherichia coli* using scanning electrochemical microscopy (SECM) (see, e.g., Nagamine et al., *Biotechnology and Bioengineering*. 96:1008-1013, 2006), among others. AARS protein fragments with increased solubility (or reduced aggregation) can be identified or selected for according to routine techniques in the art, including simple in vivo assays for protein solubility (see, e.g., Maxwell et al., *Protein Sci*. 8:1908-11, 1999).

[0274] Protein solubility and aggregation can also be measured by dynamic light scattering techniques. Aggregation is a general term that encompasses several types of interactions or characteristics, including soluble/insoluble, covalent/noncovalent, reversible/irreversible, and native/denatured interactions and characteristics. For protein therapeutics, the presence of aggregates is typically considered undesirable because of the concern that aggregates may cause an immunogenic reaction (e.g., small aggregates), or may cause adverse events on administration (e.g., particulates). Dynamic light scattering refers to a technique that can be used to determine the size distribution profile of small particles in suspension or polymers such as proteins in solution. This technique, also referred to as photon correlation spectroscopy (PCS) or quasi-elastic light scattering (QELS), uses scattered light to measure the rate of diffusion of the protein particles. Fluctuations of the scattering intensity can be observed due to the Brownian motion of the molecules and particles in solution. This motion data can be conventionally processed to derive a size distribution for the sample, wherein the size is given by the Stokes radius or hydrodynamic radius of the protein particle. The hydrodynamic size depends on both mass and shape (conformation). Dynamic scattering can detect the presence of very small amounts of aggregated protein (<0.01% by weight), even in samples that contain a large range of masses. It can also be used to compare the stability of different formulations, including, for example, applications that rely on real-time monitoring of changes at elevated temperatures. Accordingly, certain embodiments include the use of dynamic light scattering to analyze the solubility and/or presence of aggregates in a sample that contains an AARS protein fragment, antibody, or other agent of the invention.

IX. DIAGNOSTIC METHODS AND COMPOSITIONS

[0275] AARS agents such as AARS protein fragments, AARS polynucleotides, and antibodies and other binding agents described herein can be used in diagnostic assays and

diagnostic compositions. Included are biochemical, histological, and cell-based methods and compositions, among others.

[0276] These and related embodiments include the detection of the AARS polynucleotide sequence(s) or corresponding AARS polypeptide sequence(s) or portions thereof of one or more newly identified AARS protein fragments, also referred to as AARS polypeptides. For instance, certain aspects include detection of the AARS polynucleotide sequence(s) or corresponding polypeptide sequence(s) or portions thereof of one or more newly identified AARS splice variants, and/or one or more splice junctions of those splice variants. In certain embodiments, the polynucleotide or corresponding polypeptide sequence(s) of at least one of the splice junctions is unique to that particular AARS splice variant.

[0277] Also included is the direct detection of AARS protein fragments, including splice variants, proteolytic fragments, and others. In certain embodiments, the presence or levels of one or more newly identified AARS protein fragments associate or correlate with one or more cellular types or cellular states. Hence, the presence or levels of an AARS polypeptide or polynucleotide can be used to distinguish between different cellular types or different cellular states. The presence or levels of AARS protein fragments or their related polynucleotides can be detected according to polynucleotide and/or polypeptide-based diagnostic techniques, as described herein and known in the art.

[0278] Certain aspects can employ the AARS protein fragments, antibody, or AARS polynucleotides as part of a companion diagnostic method, typically to assess whether a subject or population subjects will respond favorably to a specific medical treatment. For instance, a given AARS therapeutic agent (e.g., protein fragment, antisense, RNAi, antibody, binding agent) could be identified as suitable for a subject or certain populations of subjects based on whether the subject(s) have one or more selected biomarkers for a given disease or condition. Examples of biomarkers include serum/tissue markers as well as markers that can be identified by medical imaging techniques. In certain embodiments, a naturally-occurring AARS protein fragment (or its corresponding polynucleotide) may itself provide a serum and/or tissue biomarker that can be utilized to measure drug outcome or assess the desirability of drug use in a specific subject or a specific population of subjects. In certain aspects, the identification of an AARS polypeptide or polynucleotide reference sequence may include characterizing the differential expression of that sequence, whether in a selected subject, selected tissue, or otherwise, as described herein and known in the art.

[0279] Certain of the methods provided herein rely on the differential expression of an AARS polypeptide or polynucleotide to characterize the condition or state of a cell, tissue, or subject, and to distinguish it from another cell, tissue, or subject. Non-limiting examples include methods of detecting the presence or levels of an AARS polypeptide or polynucleotide in a biological sample to distinguish between cells or tissues of different species, cells of different tissues or organs, cellular developmental states such as neonatal and adult, cellular differentiation states, conditions such as healthy, diseased and treated, intracellular and extracellular fractions, in addition to primary cell cultures and other cell cultures, such as immortalized cell cultures.

[0280] Differential expression includes a statistically significant difference in one or more gene expression levels of an AARS polynucleotide or polypeptide reference sequence compared to the expression levels of the same sequence in an appropriate control. The statistically significant difference may relate to either an increase or a decrease in expression levels, as measured by RNA levels, protein levels, protein function, or any other relevant measure of gene expression such as those described herein. Also included is a comparison between an AARS polynucleotide or polypeptide of the invention and a full-length or wild-type cytosolic or mitochondrial AARS sequence, typically of the same or corresponding type. Differential expression can be detected by a variety of techniques in the art and described herein, including polynucleotide and polypeptide based techniques, such as real-time PCR, subtractive hybridization, polynucleotide and polypeptide arrays, and others.

[0281] A result is typically referred to as statistically significant if it is unlikely to have occurred by chance. The significance level of a test or result relates traditionally to a frequentist statistical hypothesis testing concept. In simple cases, statistical significance may be defined as the probability of making a decision to reject the null hypothesis when the null hypothesis is actually true (a decision known as a Type I error, or “false positive determination”). This decision is often made using the p-value: if the p-value is less than the significance level, then the null hypothesis is rejected. The smaller the p-value, the more significant the result. Bayes factors may also be utilized to determine statistical significance (see, e.g., Goodman S., *Ann Intern Med* 130:1005-13, 1999).

[0282] In more complicated, but practically important cases, the significance level of a test or result may reflect an analysis in which the probability of making a decision to reject the null hypothesis when the null hypothesis is actually true is no more than the stated probability. This type of analysis allows for those applications in which the probability of deciding to reject may be much smaller than the significance level for some sets of assumptions encompassed within the null hypothesis.

[0283] In certain exemplary embodiments, statistically significant differential expression may include situations wherein the expression level of a given AARS sequence provides at least about a 1.2×, 1.3×, 1.4×, 1.5×, 1.6×, 1.7×, 1.8×, 1.9×, 2.0×, 2.2×, 2.4×, 2.6×, 2.8×, 3.0×, 4.0×, 5.0×, 6.0×, 7.0×, 8.0×, 9.0×, 10.0×, 15.0×, 20.0×, 50.0×, 100.0×, or greater difference in expression (i.e., differential expression that may be higher or lower expression) in a suspected biological sample as compared to an appropriate control, including all integers and decimal points in between (e.g., 1.24×, 1.25×, 2.1×, 2.5×, 60.0×, 75.0×, etc.). In certain embodiments, statistically significant differential expression may include situations wherein the expression level of a given AARS sequence provides at least about 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000 percent (%) or greater difference in expression (i.e., differential expression that may be higher or lower) in a suspected biological sample as compared to an appropriate control, including all integers and decimal points in between.

[0284] As an additional example, differential expression may also be determined by performing Z-testing, i.e., calculating an absolute Z score, as described herein and known

in the art (see Example 1). Z-testing is typically utilized to identify significant differences between a sample mean and a population mean. For example, as compared to a standard normal table (e.g., a control tissue), at a 95% confidence interval (i.e., at the 5% significance level), a Z-score with an absolute value greater than 1.96 indicates non-randomness. For a 99% confidence interval, if the absolute Z is greater than 2.58, it means that $p < 0.01$, and the difference is even more significant—the null hypothesis can be rejected with greater confidence. In these and related embodiments, an absolute Z-score of 1.96, 2, 2.58, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more, including all decimal points in between (e.g., 10.1, 10.6, 11.2, etc.), may provide a strong measure of statistical significance. In certain embodiments, an absolute Z-score of greater than 6 may provide exceptionally high statistical significance.

[0285] Substantial similarly relates generally to the lack of a statistically significant difference in the expression levels between the biological sample and the reference control. Examples of substantially similar expression levels may include situations wherein the expression level of a given SSCIGS provides less than about a 0.05×, 0.1×, 0.2×, 0.3×, 0.4×, 0.5×, 0.6×, 0.7×, 0.8×, 0.9×, 1.0×, 1.1×, 1.2×, 1.3×, or 1.4× difference in expression (i.e., differential expression that may be higher or lower expression) in a suspected biological sample as compared to a reference sample, including all decimal points in between (e.g., 0.15×, 0.25×, 0.35×, etc.). In certain embodiments, differential expression may include situations wherein the expression level of a given AARS sequence provides less than about 0.25, 0.5, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50 percent (%) difference in expression (i.e., differential expression that may be higher or lower) in a suspected biological sample as compared to a reference sample, including all decimal points in between.

[0286] In certain embodiments, such as when using an Affymetrix Microarray to measure the expression levels of an AARS polynucleotide or polypeptide reference sequence, differential expression may also be determined by the mean expression value summarized by Affymetrix Microarray Suite 5 software (Affymetrix, Santa Clara, Calif.), or other similar software, typically with a scaled mean expression value of 1000.

[0287] Embodiments of the present invention include methods of detecting the presence or levels of an AARS polynucleotide or polypeptide reference sequence or a portion thereof to distinguish between cells or tissues or other biological sample of a different organism or species, wherein the presence or levels of that sequence associates with a selected organism or species. General examples include methods of distinguishing between humans and any combination of bacteria, fungi, plants, and other non-human animals. Included within animals are methods of distinguishing between humans and any combination of vertebrates and invertebrates, including vertebrates such as fish, amphibians, reptiles, birds, and non-human mammals, and invertebrates such as insects, mollusks, crustaceans, and corals. Included within non-human mammals are methods of distinguishing between humans and any combination of non-human mammals from the Order Afrosoricida, Macroscelidea, Tubulidentata, Hyracoidea, Proboscidea, Sirenia, Cingulata, Pilosa, Scandentia, Dermoptera, Primates, Rodentia, Lagomorpha, Erinaceomorpha, Soricomorpha, Chiroptera, Pholidota, Cetacea, Carnivora, Perissodactyla,

or Artiodactyla. Included within the Primate Order are monkeys, apes, gorillas, and chimpanzees, among others known in the art. Accordingly, the presence or levels of an AARS polynucleotide or polypeptide reference sequence or variant, as described herein, may be used to identify the source of a given biological sample, such as a cell, tissue, or organ, by distinguishing between any combination of these organisms, or by distinguishing between humans and any one or more of these organisms, such as a panel of organisms. In certain embodiments, the source of a given biological sample may also be determined by comparing the presence or levels of an AARS sequence or a portion thereof to a pre-determined value.

[0288] Embodiments of the present invention include methods of detecting the presence or levels of an AARS polynucleotide or polypeptide reference sequence or a portion thereof to distinguish between cells or other biological samples that originate from different tissues or organs. Non-limiting examples include methods of distinguishing between a cell or other biological sample that originates from any combination of skin (e.g., dermis, epidermis, subcutaneous layer), hair follicles, nervous system (e.g., brain, spinal cord, peripheral nerves), auditory system or balance organs (e.g., inner ear, middle ear, outer ear), respiratory system (e.g., nose, trachea, lungs), gastroesophageal tissues, the gastrointestinal system (e.g., mouth, esophagus, stomach, small intestines, large intestines, rectum), vascular system (e.g., heart, blood vessels and arteries), liver, gallbladder, lymphatic/immune system (e.g., lymph nodes, lymphoid follicles, spleen, thymus, bone marrow), uro-genital system (e.g., kidneys, ureter, bladder, urethra, cervix, Fallopian tubes, ovaries, uterus, vulva, prostate, bulbourethral glands, epididymis, prostate, seminal vesicles, testicles), musculoskeletal system (e.g., skeletal muscles, smooth muscles, bone, cartilage, tendons, ligaments), adipose tissue, mammary tissue, and the endocrine system (e.g., hypothalamus, pituitary, thyroid, pancreas, adrenal glands). Hence, based on the association of an AARS polynucleotide or polypeptide sequence as described herein, these methods may be used to identify or characterize the tissue or organ from which a cell or other biological sample is derived.

[0289] Embodiments of the present invention include methods of detecting the presence or levels of an AARS polynucleotide or polypeptide reference sequence or a portion thereof to distinguish between or characterize the developmental or differentiation state of the cell. Also included are methods of differentiating between germ cells, stem cells, and somatic cells. Examples of developmental states include neonatal and adult. Examples of cellular differentiation states include all of the discreet and identifiable stages between a totipotent cell, a pluripotent cell, a multipotent progenitor stem cell and a mature, fully differentiated cell.

[0290] A totipotent cell has total potential, typically arises during sexual and asexual reproduction, and includes and spores and zygotes, though in certain instances cells can dedifferentiate and regain totipotency. A pluripotent cell includes a stem cell that has the potential to differentiate into any of the three germ layers, including the endoderm (interior stomach lining, gastrointestinal tract, the lungs), the mesoderm (muscle, bone, blood, urogenital), and the ectoderm (epidermal tissues and nervous system). Multipotent progenitor cells are typically capable of differentiating into

a limited number of tissue types. Examples of multipotent cells include, without limitation, hematopoietic stem cells (adult stem cells) from the bone marrow that give rise to immune cells such as red blood cells, white blood cells, and platelets, mesenchymal stem cells (adult stem cells) from the bone marrow that give rise to stromal cells, fat cells, and various types of bone cells, epithelial stem cells (progenitor cells) that give rise to the various types of skin cells, and muscle satellite cells (progenitor cells) that contribute to differentiated muscle tissue. Accordingly, the presence or levels of particular AARS polynucleotide or polypeptide sequence (e.g., splice junction of an AARS splice variant, AARS proteolytic fragment), can be used to distinguish between or characterize the above-noted cellular differentiation states, as compared to a control or a predetermined level.

[0291] Embodiments of the present invention include methods of detecting the presence or levels of an AARS polynucleotide or polypeptide reference sequence to characterize or diagnose the condition of a cell, tissue, organ, or subject, in which that condition may be characterized as healthy, diseased, at risk for being diseased, or treated. For such diagnostic purposes, the term “diagnostic” or “diagnosed” includes identifying the presence or nature of a pathologic condition, characterizing the risk of developing such a condition, and/or measuring the change (or no change) of a pathologic condition in response to therapy. Diagnostic methods may differ in their sensitivity and specificity. In certain embodiments, the “sensitivity” of a diagnostic assay refers to the percentage of diseased cells, tissues or subjects which test positive (percent of “true positives”). Diseased cells, tissues or subjects not detected by the assay are typically referred to as “false negatives.” Cells, tissues or subjects that are not diseased and which test negative in the assay may be termed “true negatives.” In certain embodiments, the “specificity” of a diagnostic assay may be defined as one (1) minus the false positive rate, where the “false positive” rate is defined as the proportion of those samples or subjects without the disease and which test positive. While a particular diagnostic method may not provide a definitive diagnosis of a condition, it suffices if the method provides a positive indication that aids in diagnosis.

[0292] In certain instances, the presence or risk of developing a pathologic condition can be diagnosed by comparing the presence or levels of one or more selected AARS polynucleotide or polypeptide reference sequences or portions thereof that correlate with the condition, whether by increased or decreased levels, as compared to a suitable control. A “suitable control” or “appropriate control” includes a value, level, feature, characteristic, or property determined in a cell or other biological sample of a tissue or organism, e.g., a control or normal cell, tissue or organism, exhibiting, for example, normal traits, such as the absence of the condition. In certain embodiments, a “suitable control” or “appropriate control” is a predefined value, level, feature, characteristic, or property. Other suitable controls will be apparent to persons skilled in the art. Examples of diseases and conditions are described elsewhere herein.

[0293] Embodiments of the present invention include AARS polynucleotide or nucleic acid-based detection techniques, which offer certain advantages due to sensitivity of detection. Hence, certain embodiments relate to the use or detection of AARS polynucleotides as part of a diagnostic method or assay. The presence and/or levels of AARS

polynucleotides may be measured by any method known in the art, including hybridization assays such as Northern blot, quantitative or qualitative polymerase chain reaction (PCR), quantitative or qualitative reverse transcriptase PCR (RT-PCR), microarray, dot or slot blots, or in situ hybridization such as fluorescent in situ hybridization (FISH), among others. Certain of these methods are described in greater detail below.

[0294] AARS polynucleotides such as DNA and RNA can be collected and/or generated from blood, biological fluids, tissues, organs, cell lines, or other relevant sample using techniques known in the art, such as those described in Kingston. (2002 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y. (see, e.g., as described by Nelson et al. *Proc Natl Acad Sci USA*, 99: 11890-11895, 2002) and elsewhere. Further, a variety of commercially available kits for constructing RNA are useful for making the RNA to be used in the present invention. RNA may be constructed from organs/tissues/cells procured from normal healthy subjects; however, this invention also contemplates construction of RNA from diseased subjects. Certain embodiments contemplate using any type of organ from any type of subject or animal. For test samples RNA may be procured from an individual (e.g., any animal, including mammals) with or without visible disease and from tissue samples, biological fluids (e.g., whole blood) or the like.

[0295] In certain embodiments, amplification or construction of cDNA sequences may be helpful to increase detection capabilities. The instant disclosure, as well as the art, provides the requisite level of detail to perform such tasks. In one exemplary embodiment, whole blood is used as the source of RNA and accordingly, RNA stabilizing reagents are optionally used, such as PAX tubes, as described, for example, in Thach et al., *J. Immunol. Methods*. December 283(1-2):269-279, 2003 and Chai et al., *J. Clin. Lab Anal.* 19(5):182-188, 2005 (both of which are incorporated by reference). Complementary DNA (cDNA) libraries can be generated using techniques known in the art, such as those described in Ausubel et al. (2001 *Current Protocols in Molecular Biology*, Greene Publ. Assoc. Inc. & John Wiley & Sons, Inc., NY, N.Y.); Sambrook et al. (1989 *Molecular Cloning*, Second Ed., Cold Spring Harbor Laboratory, Plainview, N.Y.); Maniatis et al. (1982 *Molecular Cloning*, Cold Spring Harbor Laboratory, Plainview, N.Y.) and elsewhere. Further, a variety of commercially available kits for constructing cDNA libraries are useful for making the cDNA libraries of the present invention. Libraries can be constructed from organs/tissues/cells procured from normal, healthy subjects.

[0296] Certain embodiments may employ hybridization methods for detecting AARS polynucleotide sequences. Methods for conducting polynucleotide hybridization assays have been well developed in the art. Hybridization assay procedures and conditions will vary depending on the application and are selected in accordance with the general binding methods known including those referred to in: Maniatis et al. *Molecular Cloning: A Laboratory Manual* (2nd Ed. Cold Spring Harbor, N.Y., 1989); Berger and Kimmel *Methods in Enzymology*, Vol. 152, Guide to Molecular Cloning Techniques (Academic Press, Inc., San Diego, Calif., 1987); Young and Davis, *PNAS*. 80: 1194 (1983). Methods and apparatus for carrying out repeated and controlled hybridization reactions have been described in

U.S. Pat. Nos. 5,871,928, 5,874,219, 6,045,996 and 6,386,749, 6,391,623 each of which are incorporated herein by reference

[0297] Certain embodiments may employ nucleic acid amplification methods for detecting AARS polynucleotide sequences. The term “amplification” or “nucleic acid amplification” refers to the production of multiple copies of a target nucleic acid that contains at least a portion of the intended specific target nucleic acid sequence. The multiple copies may be referred to as amplicons or amplification products. In certain embodiments, the amplified target contains less than the complete target gene sequence (introns and exons) or an expressed target gene sequence (spliced transcript of exons and flanking untranslated sequences). For example, specific amplicons may be produced by amplifying a portion of the target polynucleotide by using amplification primers that hybridize to, and initiate polymerization from, internal positions of the target polynucleotide. Preferably, the amplified portion contains a detectable target sequence that may be detected using any of a variety of well-known methods.

[0298] “Selective amplification” or “specific amplification,” as used herein, refers to the amplification of a target nucleic acid sequence according to the present invention wherein detectable amplification of the target sequence is substantially limited to amplification of target sequence contributed by a nucleic acid sample of interest that is being tested and is not contributed by target nucleic acid sequence contributed by some other sample source, e.g., contamination present in reagents used during amplification reactions or in the environment in which amplification reactions are performed.

[0299] The term “amplification conditions” refers to conditions permitting nucleic acid amplification according to the present invention. Amplification conditions may, in some embodiments, be less stringent than “stringent hybridization conditions” as described herein. Oligonucleotides used in the amplification reactions of the present invention hybridize to their intended targets under amplification conditions, but may or may not hybridize under stringent hybridization conditions. On the other hand, detection probes of the present invention typically hybridize under stringent hybridization conditions. Acceptable conditions to carry out nucleic acid amplifications according to the present invention can be easily ascertained by someone having ordinary skill in the art depending on the particular method of amplification employed.

[0300] Many well-known methods of nucleic acid amplification require thermocycling to alternately denature double-stranded nucleic acids and hybridize primers; however, other well-known methods of nucleic acid amplification are isothermal. The polymerase chain reaction (U.S. Pat. Nos. 4,683,195; 4,683,202; 4,800,159; 4,965,188), commonly referred to as PCR, uses multiple cycles of denaturation, annealing of primer pairs to opposite strands, and primer extension to exponentially increase copy numbers of the target sequence. In a variation called RT-PCR, reverse transcriptase (RT) is used to make a complementary DNA (cDNA) from mRNA, and the cDNA is then amplified by PCR to produce multiple copies of DNA.

[0301] As noted above, the term “PCR” refers to multiple amplification cycles that selectively amplify a target nucleic acid species. Included are quantitative PCR (qPCR), real-time PCR, reverse transcription PCR (RT-PCR) and quan-

titative reverse transcription PCR (qRT-PCR) is well described in the art. The term “pPCR” refers to quantitative polymerase chain reaction, and the term “qRT-PCR” refers to quantitative reverse transcription polymerase chain reaction. qPCR and qRT-PCR may be used to amplify and simultaneously quantify a targeted cDNA molecule. It enables both detection and quantification of a specific sequence in a cDNA pool, such as a selected AARS gene or transcript.

[0302] The term “real-time PCR” may use DNA-binding dye to bind to all double-stranded (ds) DNA in PCR, causing fluorescence of the dye. An increase in DNA product during PCR therefore leads to an increase in fluorescence intensity and is measured at each cycle, thus allowing DNA concentrations to be quantified. However, dsDNA dyes such as SYBR Green will bind to all dsDNA PCR products. Fluorescence is detected and measured in the real-time PCR thermocycler, and its geometric increase corresponding to exponential increase of the product is used to determine the threshold cycle (“Ct”) in each reaction.

[0303] The term “Ct Score” refers to the threshold cycle number, which is the cycle at which PCR amplification has surpassed a threshold level. If there is a higher quantity of mRNA for a particular gene in a sample, it will cross the threshold earlier than a lowly expressed gene since there is more starting RNA to amplify. Therefore, a low Ct score indicates high gene expression in a sample and a high Ct score is indicative of low gene expression.

[0304] Certain embodiments may employ the ligase chain reaction (Weiss, *Science*, 254: 1292, 1991), commonly referred to as LCR, which uses two sets of complementary DNA oligonucleotides that hybridize to adjacent regions of the target nucleic acid. The DNA oligonucleotides are covalently linked by a DNA ligase in repeated cycles of thermal denaturation, hybridization and ligation to produce a detectable double-stranded ligated oligonucleotide product.

[0305] Another method is strand displacement amplification (Walker, G. et al., 1992, *Proc. Natl. Acad. Sci. USA* 89:392-396; U.S. Pat. Nos. 5,270,184 and 5,455,166), commonly referred to as SDA, which uses cycles of annealing pairs of primer sequences to opposite strands of a target sequence, primer extension in the presence of a dNTPoS to produce a duplex hemiphosphorothioated primer extension product, endonuclease-mediated nicking of a hemimodified restriction endonuclease recognition site, and polymerase-mediated primer extension from the 3' end of the nick to displace an existing strand and produce a strand for the next round of primer annealing, nicking and strand displacement, resulting in geometric amplification of product. Thermophilic SDA (tSDA) uses thermophilic endonucleases and polymerases at higher temperatures in essentially the same method (European Pat. No. 0 684 315).

[0306] Other amplification methods include, for example: nucleic acid sequence based amplification (U.S. Pat. No. 5,130,238), commonly referred to as NASBA; one that uses an RNA replicase to amplify the probe molecule itself (Lizardi, P. et al., 1988, *BioTechnol.* 6: 1197-1202), commonly referred to as Q β replicase; a transcription based amplification method (Kwoh, D. et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177); self-sustained sequence replication (Guatelli, J. et al., 1990, *Proc. Natl. Acad. Sci. USA* 87: 1874-1878); and, transcription mediated amplification (U.S. Pat. Nos. 5,480,784 and 5,399,491), commonly referred to as TMA. For further discussion of known ampli-

fication methods see Persing, David H., 1993, “In Vitro Nucleic Acid Amplification Techniques” in *Diagnostic Medical Microbiology: Principles and Applications* (Persing et al., Eds.), pp. 51-87 (American Society for Microbiology, Washington, D.C.).

[0307] Illustrative transcription-based amplification systems of the present invention include TMA, which employs an RNA polymerase to produce multiple RNA transcripts of a target region (U.S. Pat. Nos. 5,480,784 and 5,399,491). TMA uses a “promoter-primer” that hybridizes to a target nucleic acid in the presence of a reverse transcriptase and an RNA polymerase to form a double-stranded promoter from which the RNA polymerase produces RNA transcripts. These transcripts can become templates for further rounds of TMA in the presence of a second primer capable of hybridizing to the RNA transcripts. Unlike PCR, LCR or other methods that require heat denaturation, TMA is an isothermal method that uses an RNase H activity to digest the RNA strand of an RNA:DNA hybrid, thereby making the DNA strand available for hybridization with a primer or promoter-primer. Generally, the RNase H activity associated with the reverse transcriptase provided for amplification is used.

[0308] In an illustrative TMA method, one amplification primer is an oligonucleotide promoter-primer that comprises a promoter sequence which becomes functional when double-stranded, located 5' of a target-binding sequence, which is capable of hybridizing to a binding site of a target RNA at a location 3' to the sequence to be amplified. A promoter-primer may be referred to as a “T7-primer” when it is specific for T7 RNA polymerase recognition. Under certain circumstances, the 3' end of a promoter-primer, or a subpopulation of such promoter-primers, may be modified to block or reduce primer extension. From an unmodified promoter-primer, reverse transcriptase creates a cDNA copy of the target RNA, while RNase H activity degrades the target RNA. A second amplification primer then binds to the cDNA. This primer may be referred to as a “non-T7 primer” to distinguish it from a “T7-primer.” From this second amplification primer, reverse transcriptase creates another DNA strand, resulting in a double-stranded DNA with a functional promoter at one end. When double-stranded, the promoter sequence is capable of binding an RNA polymerase to begin transcription of the target sequence to which the promoter-primer is hybridized. An RNA polymerase uses this promoter sequence to produce multiple RNA transcripts (i.e., amplicons), generally about 100 to 1,000 copies. Each newly-synthesized amplicon can anneal with the second amplification primer. Reverse transcriptase can then create a DNA copy, while the RNase H activity degrades the RNA of this RNA:DNA duplex. The promoter-primer can then bind to the newly synthesized DNA, allowing the reverse transcriptase to create a double-stranded DNA, from which the RNA polymerase produces multiple amplicons. Thus, a billion-fold isothermal amplification can be achieved using two amplification primers.

[0309] In certain embodiments, other techniques may be used to evaluate RNA transcripts of the transcripts from a particular cDNA library, including microarray analysis (Han, M., et al., *Nat Biotechnol.* 19: 631-635, 2001; Bao, P., et al., *Anal Chem.* 74: 1792-1797, 2002; Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-19, 1996; and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-55, 1997) and SAGE (serial analysis of gene expression). Like MPSS, SAGE is digital and can generate a large number of signature

sequences. (see e.g., Velculescu, V. E., et al., *Trends Genet.* 16: 423-425., 2000; Tuteja R. and Tuteja N. *Bioessays.* 2004 August; 26(8):916-22), although orders of magnitude fewer than that are available from techniques such as MPSS.

[0310] In certain embodiments, the term “microarray” includes a “nucleic acid microarray” having a substrate-bound plurality of nucleic acids, hybridization to each of the plurality of bound nucleic acids being separately detectable. The substrate can be solid or porous, planar or non-planar, unitary or distributed. Nucleic acid microarrays include all the devices so called in Schena (ed.), *DNA Microarrays: A Practical Approach* (Practical Approach Series), Oxford University Press (1999); Nature Genet. 21(1) (suppl.): 1-60 (1999); Schena (ed.), *Microarray Biochip: Tools and Technology*, Eaton Publishing Company/BioTechniques Books Division (2000). Nucleic acid microarrays may include a substrate-bound plurality of nucleic acids in which the plurality of nucleic acids are disposed on a plurality of beads, rather than on a unitary planar substrate, as described, for example, in Brenner et al., *Proc. Natl. Acad. Sci. USA* 97(4): 1665-1670 (2000). Examples of nucleic acid microarrays may be found in U.S. Pat. Nos. 6,391,623, 6,383,754, 6,383,749, 6,380,377, 6,379,897, 6,376,191, 6,372,431, 6,351,712, 6,344,316, 6,316,193, 6,312,906, 6,309,828, 6,309,824, 6,306,643, 6,300,063, 6,287,850, 6,284,497, 6,284,465, 6,280,954, 6,262,216, 6,251,601, 6,245,518, 6,263,287, 6,251,601, 6,238,866, 6,228,575, 6,214,587, 6,203,989, 6,171,797, 6,103,474, 6,083,726, 6,054,274, 6,040,138, 6,083,726, 6,004,755, 6,001,309, 5,958,342, 5,952,180, 5,936,731, 5,843,655, 5,814,454, 5,837,196, 5,436,327, 5,412,087, and 5,405,783, the disclosures of which are incorporated by reference.

[0311] Additional examples include nucleic acid arrays that are commercially available from Affymetrix (Santa Clara, Calif.) under the brand name GENECHIP™. Further exemplary methods of manufacturing and using arrays are provided in, for example, U.S. Pat. Nos. 7,028,629; 7,011,949; 7,011,945; 6,936,419; 6,927,032; 6,924,103; 6,921,642; and 6,818,394.

[0312] The present invention as related to arrays and microarrays also contemplates many uses for polymers attached to solid substrates. These uses include gene expression monitoring, profiling, library screening, genotyping and diagnostics. Gene expression monitoring and profiling methods and methods useful for gene expression monitoring and profiling are shown in U.S. Pat. Nos. 5,800,992, 6,013,449, 6,020,135, 6,033,860, 6,040,138, 6,177,248 and 6,309,822. Genotyping and uses therefore are shown in U.S. Ser. Nos. 10/442,021, 10/013,598 (U.S. Application No. 2003/0036069), and U.S. Pat. Nos. 5,925,525, 6,268,141, 5,856,092, 6,267,152, 6,300,063, 6,525,185, 6,632,611, 5,858,659, 6,284,460, 6,361,947, 6,368,799, 6,673,579 and 6,333,179. Other methods of nucleic acid amplification, labeling and analysis that may be used in combination with the methods disclosed herein are embodied in U.S. Pat. Nos. 5,871,928, 5,902,723, 6,045,996, 5,541,061, and 6,197,506.

[0313] As will be apparent to persons skilled in the art, certain embodiments may employ oligonucleotides, such as primers or probes, for amplification or detection, as described herein. Oligonucleotides of a defined sequence and chemical structure may be produced by techniques known to those of ordinary skill in the art, such as by chemical or biochemical synthesis, and by in vitro or in vivo expression from recombinant nucleic acid molecules, e.g.,

bacterial or viral vectors. In certain embodiments, an oligonucleotide does not consist solely of wild-type chromosomal DNA or the in vivo transcription products thereof.

[0314] Oligonucleotides or primers may be modified in any way, as long as a given modification is compatible with the desired function of a given oligonucleotide. One of ordinary skill in the art can easily determine whether a given modification is suitable or desired for any given oligonucleotide of the present invention. Relevant AARS oligonucleotides are described in greater detail elsewhere herein.

[0315] While the design and sequence of oligonucleotides depends on their function as described herein, several variables are generally taken into account. Among the most relevant are: length, melting temperature (T_m), specificity, complementarity with other oligonucleotides in the system, G/C content, polypyrimidine (T, C) or polypurine (A, G) stretches, and the 3'-end sequence. Controlling for these and other variables is a standard and well known aspect of oligonucleotide design, and various computer programs are readily available to screen large numbers of potential oligonucleotides for optimal ones.

[0316] Certain embodiments therefore include methods for detecting a target AARS polynucleotide in a sample, the polynucleotide comprising the sequence of a reference AARS polynucleotide, as described herein, comprising a) hybridizing the sample with a probe comprising a sequence complementary to the target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. Also included are methods for detecting a target AARS polynucleotide in a sample, the polynucleotide comprising the sequence of a reference AARS polynucleotide, as described herein, comprising a) amplifying the target polynucleotide or fragment thereof, and b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof. Specific embodiments relate to the detection of AARS splice variants, such as by detecting a unique splice junction of the splice variant, whether by hybridization, amplification, or other detection method.

[0317] Embodiments of the present invention include a variety of AARS polypeptide-based detection techniques, including antibody-based detection techniques. Included in these embodiments are the use of AARS polypeptides to generate antibodies or other binders, which may then be used in diagnostic methods and compositions to detect or quantitate selected AARS polypeptides in a cell or other biological sample, typically from a subject.

[0318] Certain embodiments may employ standard methodologies and detectors such as western blotting and immunoprecipitation, enzyme-linked immunosorbent assays (ELISA), flow cytometry, and immunofluorescence assays (IFA), which utilize an imaging device. These well-known methods typically utilize one or more monoclonal or polyclonal antibodies as described herein that specifically bind to a selected AARS polypeptide of the invention, or a unique region of that AARS polypeptide, and generally do not bind significantly to other AARS polypeptides, such as a full-length AARS polypeptide. In certain embodiments, the unique region of the AARS polypeptide may represent a

unique three-dimensional structure that is possessed by a newly identified protein fragment of an AARS.

[0319] Certain embodiments may employ “arrays,” such as “microarrays.” In certain embodiments, a “microarray” may also refer to a “peptide microarray” or “protein microarray” having a substrate-bound collection or plurality of polypeptides, the binding to each of the plurality of bound polypeptides being separately detectable. Alternatively, the peptide microarray may have a plurality of binders, including but not limited to monoclonal antibodies, polyclonal antibodies, phage display binders, yeast 2 hybrid binders, and aptamers, which can specifically detect the binding of the AARS polypeptides described herein. The array may be based on autoantibody detection of these AARS polypeptides, as described, for example, in Robinson et al., *Nature Medicine* 8(3):295-301 (2002). Examples of peptide arrays may be found in WO 02/31463, WO 02/25288, WO 01/94946, WO 01/88162, WO 01/68671, WO 01/57259, WO 00/61806, WO 00/54046, WO 00/47774, WO 99/40434, WO 99/39210, and WO 97/42507 and U.S. Pat. Nos. 6,268,210, 5,766,960, and 5,143,854, each of which are incorporated by reference.

[0320] Certain embodiments may employ MS or other molecular weight-based methods for diagnostically detecting AARS polypeptide sequences. Mass spectrometry (MS) refers generally to an analytical technique for determining the elemental composition of a sample or molecule. MS may also be used for determining the chemical structures of molecules, such as peptides and other chemical compounds.

[0321] Generally, the MS principle consists of ionizing chemical compounds to generate charged molecules or molecule fragments, and then measuring their mass-to-charge ratios. In an illustrative MS procedure: a sample is loaded onto the MS instrument, and undergoes vaporization, the components of the sample are ionized by one of a variety of methods (e.g., by impacting them with an electron beam), which results in the formation of positively charged particles, the positive ions are then accelerated by a magnetic field, computations are performed on the mass-to-charge ratio (m/z) of the particles based on the details of motion of the ions as they transit through electromagnetic fields, and, detection of the ions, which in step prior were sorted according to m/z .

[0322] An illustrative MS instruments has three modules: an ion source, which converts gas phase sample molecules into ions (or, in the case of electrospray ionization, move ions that exist in solution into the gas phase); a mass analyzer, which sorts the ions by their masses by applying electromagnetic fields; and a detector, which measures the value of an indicator quantity and thus provides data for calculating the abundances of each ion present.

[0323] The MS technique has both qualitative and quantitative uses, including identifying unknown compounds, determining the isotopic composition of elements in a molecule, and determining the structure of a compound by observing its fragmentation. Other uses include quantifying the amount of a compound in a sample or studying the fundamentals of gas phase ion chemistry (the chemistry of ions and neutrals in a vacuum). Included are gas chromatography-mass spectrometry (GC/MS or GC-MS), liquid chromatography mass spectrometry (LC/MS or LC-MS), and ion mobility spectrometry/mass spectrometry (IMS/MS or IMMS). Accordingly, MS techniques may be used according to any of the methods provided herein to measure

the presence or levels of an AARS polypeptide of the invention in a biological sample, and to compare those levels to a control sample or a pre-determined value.

[0324] Certain embodiments may employ cell-sorting or cell visualization or imaging devices/techniques to detect or quantitate the presence or levels of AARS polynucleotides or polypeptides. Examples include flow cytometry or FACS, immunofluorescence analysis (IFA), and in situ hybridization techniques, such as fluorescent in situ hybridization (FISH).

[0325] Certain embodiments may employ conventional biology methods, software and systems for diagnostic purposes. Computer software products of the invention typically include computer readable medium having computer-executable instructions for performing the logic steps of the method of the invention. Suitable computer readable medium include floppy disk, CD-ROM/DVD/DVD-ROM, hard-disk drive, flash memory, ROM/RAM, magnetic tapes and etc. The computer executable instructions may be written in a suitable computer language or combination of several languages. Basic computational biology methods are described in, for example Setubal and Meidanis et al., *Introduction to Computational Biology Methods* (PWS Publishing Company, Boston, 1997); Salzberg, Searles, Kasif, (Ed.), *Computational Methods in Molecular Biology*, (Elsevier, Amsterdam, 1998); Rashidi and Buehler, *Bioinformatics Basics: Application in Biological Science and Medicine* (CRC Press, London, 2000) and Ouelette and Bzevanis *Bioinformatics: A Practical Guide for Analysis of Gene and Proteins* (Wiley & Sons, Inc., 2nd ed., 2001). See U.S. Pat. No. 6,420,108.

[0326] Certain embodiments may employ various computer program products and software for a variety of purposes, such as probe design, management of data, analysis, and instrument operation. See, U.S. Pat. Nos. 5,593,839, 5,795,716, 5,733,729, 5,974,164, 6,066,454, 6,090,555, 6,185,561, 6,188,783, 6,223,127, 6,229,911 and 6,308,170.

[0327] The whole genome sampling assay (WGSA) is described, for example in Kennedy et al., *Nat. Biotech.* 21, 1233-1237 (2003), Matsuzaki et al., *Gen. Res.* 14: 414-425, (2004), and Matsuzaki, et al., *Nature Methods* 1:109-111 (2004). Algorithms for use with mapping assays are described, for example, in Liu et al., *Bioinformatics.* 19: 2397-2403 (2003) and Di et al. *Bioinformatics.* 21:1958 (2005). Additional methods related to WGSA and arrays useful for WGSA and applications of WGSA are disclosed, for example, in U.S. Patent Application Nos. 60/676,058 filed Apr. 29, 2005, 60/616,273 filed Oct. 5, 2004, Ser. No. 10/912,445, 11/044,831, 10/442,021, 10/650,332 and 10/463,991. Genome wide association studies using mapping assays are described in, for example, Hu et al., *Cancer Res.*; 65(7):2542-6 (2005), Mitra et al., *Cancer Res.*, 64(21): 8116-25 (2004), Butcher et al., *Hum Mol Genet.*, 14(10): 1315-25 (2005), and Klein et al., *Science.* 308(5720):385-9 (2005).

[0328] Additionally, certain embodiments may include methods for providing genetic information over networks such as the Internet as shown, for example, in U.S. application Ser. Nos. 10/197,621, 10/063,559 (United States Publication Number 2002/0183936), Ser. Nos. 10/065,856, 10/065,868, 10/328,818, 10/328,872, 10/423,403, and 60/482,389.

X ANTISENSE AND RNAI AGENTS

[0329] Embodiments of the present invention also include antisense oligonucleotides and RNAi agents that target the AARS polynucleotide sequences, and methods of use thereof to reduce expression of a selected AARS transcript and/or protein fragment. Certain embodiments relate to targeting one or more splice junctions (often unique) that generate a splice variant, AARS protein fragment of instant invention. Also included are methods of antisense or RNAi inhibition that target certain splice forms, either to encourage or discourage splicing of a selected protein fragment. In certain preferred embodiments, the splice junctions that generate the AARS protein fragments are over-expressed with respect to particular tissues, and are unique to that splice variant. In these and related embodiments, such splice variants are not the only source of cytosolic AARS activity in the targeted cell type. For instance, certain splice variants to be targeted may represent about 10% to 50% of the total copy number of the AARS RNA splice variants in a given cell or tissue, and preferably about 1-10% of the total copy number of the AARS RNA splice variants in a given cell or tissue. Splice variants that are about <1% of the total copy number of the AARS RNA splice variants in a given cell or tissue may also be targeted.

[0330] In certain embodiments, the antisense or RNAi agent does not target the full-length protein, because such full-length proteins are responsible for a key step in protein synthesis, and thereby avoids lethality that often results from wild-type AARS knockouts. Certain of the methods described herein can therefore be used to avoid undesired effects such as toxicities in both chronic and acute treatments, and to selectively modulate the non-canonical activities of the AARS protein fragment. However, certain embodiments may generically target AARS sequences, including full-length AARS sequences, such as to kill or substantially derange the cell physiology of a target cell or tissue.

[0331] In certain embodiments, the AARS splice variant to be targeted possesses a non-canonical biological activity. In some embodiments, the AARS splice variant has reduced or undetectable canonical AARS activity, and the antisense or RNAi-related method more specifically modulates its non-canonical activity. In certain embodiments, the antisense or RNAi-related agents can be combined with a targeted or local delivery approach to lessen systemic undesired effects to non-targeted cells or tissues. Among others described herein, exemplary cells or tissues that could be targeted this way include cancer cells, and cells to tissues that lend themselves to localized targeting, such as tumors or epithelia via topical application.

[0332] A. Antisense Agents

[0333] The terms “antisense oligomer” or “antisense compound” or “antisense oligonucleotide” are used interchangeably and refer to a sequence of cyclic subunits, each bearing a base-pairing moiety, linked by intersubunit linkages that allow the base-pairing moieties to hybridize to a target sequence in a nucleic acid (typically an RNA) by Watson-Crick base pairing, to form a nucleic acid:oligomer heteroduplex within the target sequence, and typically thereby prevent translation of that RNA. Also included are methods of use thereof to modulate expression of a selected AARS transcript, such as a splice variant or proteolytic fragment, and/or its corresponding polypeptide.

[0334] Antisense oligonucleotides may contain between about 8 and 40 subunits, typically about 8-25 subunits, and preferably about 12 to 25 subunits. In certain embodiments, oligonucleotides may have exact sequence complementarity to the target sequence or near complementarity, as defined below. In certain embodiments, the degree of complementarity between the target and antisense targeting sequence is sufficient to form a stable duplex. The region of complementarity of the antisense oligomers with the target RNA sequence may be as short as 8-11 bases, but is preferably 12-15 bases or more, e.g., 12-20 bases, or 12-25 bases, including all integers in between these ranges. An antisense oligomer of about 14-15 bases is generally long enough to have a unique complementary sequence in targeting the selected AARS gene. In certain embodiments, a minimum length of complementary bases may be required to achieve the requisite binding T_m , as discussed herein.

[0335] In certain embodiments, antisense oligomers as long as 40 bases may be suitable, where at least a minimum number of bases, e.g., 10-12 bases, are complementary to the target sequence. In general, however, facilitated or active uptake in cells is optimized at oligomer lengths less than about 30. For certain oligomers, described further below, an optimum balance of binding stability and uptake generally occurs at lengths of 18-25 bases. Included are antisense oligomers (e.g., PNAs, LNAs, 2'-OMe, MOE) that consist of about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 bases, in which at least about 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 contiguous or non-contiguous bases are complementary to their AARS target sequence, or variants thereof.

[0336] In certain embodiments, antisense oligomers may be 100% complementary to an AARS nucleic acid target sequence, or it may include mismatches, e.g., to accommodate variants, as long as a heteroduplex formed between the oligomer and AARS nucleic acid target sequence is sufficiently stable to withstand the action of cellular nucleases and other modes of degradation which may occur in vivo. The term “target sequence” refers to a portion of the target RNA against which the oligonucleotide is directed, that is, the sequence to which the oligonucleotide will hybridize by Watson-Crick base pairing of a complementary sequence. In certain embodiments, the target sequence may be a contiguous region of an AARS mRNA (e.g., a unique splice junction of an AARS mRNA), or may be composed of non-contiguous regions of the mRNA.

[0337] Oligomer backbones which are less susceptible to cleavage by nucleases are discussed below. Mismatches, if present, are less destabilizing toward the end regions of the hybrid duplex than in the middle. The number of mismatches allowed will depend on the length of the oligomer, the percentage of G:C base pairs in the duplex, and the position of the mismatch(es) in the duplex, according to well understood principles of duplex stability. Although such an antisense oligomer is not necessarily 100% complementary to the AARS nucleic acid target sequence, it is effective to stably and specifically bind to the target sequence, such that a biological activity of the nucleic acid target, e.g., expression of AARS protein(s), is modulated.

[0338] The stability of the duplex formed between an oligomer and a target sequence is a function of the binding T_m and the susceptibility of the duplex to cellular enzymatic

cleavage. The T_m of an antisense oligonucleotide with respect to complementary-sequence RNA may be measured by conventional methods, such as those described by Hames et al., *Nucleic Acid Hybridization*, IRL Press, 1985, pp. 107-108 or as described in Miyada C. G. and Wallace R. B., 1987, Oligonucleotide hybridization techniques, *Methods Enzymol.* Vol. 154 pp. 94-107. In certain embodiments, antisense oligomer may have a binding T_m , with respect to a complementary-sequence RNA, of greater than body temperature and preferably greater than 50° C. T_m 's in the range 60-80° C. or greater are preferred. According to well known principles, the T_m of an oligomer compound, with respect to a complementary-based RNA hybrid, can be increased by increasing the ratio of C:G paired bases in the duplex, and/or by increasing the length (in base pairs) of the heteroduplex. At the same time, for purposes of optimizing cellular uptake, it may be advantageous to limit the size of the antisense oligomer. For this reason, compounds that show high T_m (50° C. or greater) at a length of 25 bases or less are generally preferred over those requiring greater than 25 bases for high T_m values.

[0339] Antisense oligomers can be designed to block or inhibit translation of mRNA or to inhibit natural pre-mRNA splice processing, or induce degradation of targeted mRNAs, and may be said to be “directed to” or “targeted against” a target sequence with which it hybridizes. In certain embodiments, the target sequence may include any coding or non-coding sequence of an AARS mRNA transcript, and may thus be within an exon or within an intron. In certain embodiments, the target sequence is relatively unique or exceptional among AARSs (e.g., a full-length AARS) and is selective for reducing expression of a selected AARS protein fragment, such as a proteolytic fragment or splice variant. In certain embodiments, the target site includes a 3' or 5' splice site of a pre-processed mRNA, or a branch point. The target sequence for a splice site may include an mRNA sequence having its 5' end 1 to about 25 to about 50 base pairs downstream of a splice acceptor junction or upstream of a splice donor junction in a pre-processed mRNA. In certain embodiments, a target sequence may include a splice junction of an alternatively splice AARS mRNA, such as a splice junction that does not occur in the full-length AARS, or is unique or exceptional to that transcript, in that it either does not occur or only seldom occurs in other AARS splice variants. An oligomer is more generally said to be “targeted against” a biologically relevant target, such as reference AARS polynucleotide, when it is targeted against the nucleic acid of the target in the manner described herein.

[0340] An oligonucleotide is typically complementary to a target sequence, such as a target DNA or RNA. The terms “complementary” and “complementarity” refer to polynucleotides (i.e., a sequence of nucleotides) related by the base-pairing rules. For example, the sequence “A-G-T,” is complementary to the sequence “T-C-A.” Complementarity may be “partial,” in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or, there may be “complete” or “total” complementarity (100%) between the nucleic acids. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. While perfect complementarity is often desired, some embodiments can include one or more but preferably 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3,

2, or 1 mismatches with respect to the target sequence. Variations at any location within the oligomer are included. In certain embodiments, variations in sequence near the termini of an oligomer are generally preferable to variations in the interior, and if present are typically within about 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1 nucleotides of the 5' and/or 3' terminus.

[0341] The term “targeting sequence” or in certain embodiments “antisense targeting sequence” refers to the sequence in an oligonucleotide that is complementary (meaning, in addition, substantially complementary) to the target sequence in the DNA or RNA target molecule. The entire sequence, or only a portion, of the antisense compound may be complementary to the target sequence. For example, in an oligonucleotide having 20-30 bases, about 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, or 29 may be targeting sequences that are complementary to the target region. Typically, the targeting sequence is formed of contiguous bases, but may alternatively be formed of non-contiguous sequences that when placed together, e.g., from opposite ends of the oligonucleotide, constitute sequence that spans the target sequence.

[0342] Target and targeting sequences are described as “complementary” to one another when hybridization occurs in an antiparallel configuration. A targeting sequence may have “near” or “substantial” complementarity to the target sequence and still function for the purpose of the present invention, that is, it may still be functionally “complementary.” In certain embodiments, an oligonucleotide may have at most one mismatch with the target sequence out of 10 nucleotides, and preferably at most one mismatch out of 20. Alternatively, an oligonucleotide may have at least about 80%, 85%, 90% sequence homology, and preferably at least 95% sequence homology, with an AARS reference polynucleotide sequence described herein, or its complement.

[0343] An oligonucleotide “specifically hybridizes” to a target polynucleotide if the oligomer hybridizes to a target (e.g., an AARS reference polynucleotide or its complement) under physiological conditions, with a T_m substantially greater than 45° C., preferably at least 50° C., and typically 60° C.-80° C. or higher. Such hybridization preferably corresponds to stringent hybridization conditions. At a given ionic strength and pH, the T_m is the temperature at which 50% of a target sequence hybridizes to a complementary polynucleotide. Again, such hybridization may occur with “near” or “substantial” complementarity of the antisense oligomer to the target sequence, as well as with exact complementarity.

[0344] The terms specifically binds or specifically hybridizes refer generally to an oligonucleotide probe or polynucleotide sequence that not only binds to its intended target gene sequence in a sample under selected hybridization conditions, but does not bind significantly to other target sequences in the sample, and thereby discriminates between its intended target and all other targets in the target pool. A probe that specifically hybridizes to its intended target sequence may also detect concentration differences under the selected hybridization conditions, as described herein.

[0345] A “nuclease-resistant” oligomeric molecule (oligomer) refers to one whose backbone is substantially resistant to nuclease cleavage, in non-hybridized or hybridized form; by common extracellular and intracellular nucleases in the body; that is, the oligomer shows little or no nuclease

cleavage under normal nuclease conditions in the body to which the oligomer is exposed.

[0346] A “heteroduplex” refers to a duplex between an oligonucleotide and the complementary portion of a target polynucleotide, such as a target DNA or RNA. A “nuclease-resistant heteroduplex” refers to a heteroduplex formed by the binding of an oligomer to its complementary target, such that the heteroduplex is substantially resistant to *in vivo* degradation by intracellular and extracellular nucleases, such as RNaseH, which are capable of cutting double-stranded RNA/RNA or RNA/DNA complexes.

[0347] A “subunit” of an oligonucleotide refers to one nucleotide (or nucleotide analog) unit. The term may refer to the nucleotide unit with or without the attached intersubunit linkage, although, when referring to a “charged subunit”, the charge typically resides within the intersubunit linkage (e.g., a phosphate or phosphorothioate linkage or a cationic linkage).

[0348] The cyclic subunits of an oligonucleotide may be based on ribose or another pentose sugar or, in certain embodiments, alternate or modified groups. Examples of modified oligonucleotide backbones include, without limitation, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates including 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates including 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Also contemplated are peptide nucleic acids (PNAs), locked nucleic acids (LNAs), 2'-O-Methyl oligonucleotides (2'-OMe), 2'-methoxyethoxy oligonucleotides (MOE), among other oligonucleotides known in the art.

[0349] The purine or pyrimidine base pairing moiety is typically adenine, cytosine, guanine, uracil, thymine or inosine. Also included are bases such as pyridin-4-one, pyridin-2-one, phenyl, pseudouracil, 2,4,6-trimethylthioxy benzene, 3-methyl uracil, dihydrouridine, naphthyl, amino-phenyl, 5-alkylcytidines (e.g., 5-methylcytidine), 5-alkyluridines (e.g., ribothymidine), 5-halouridine (e.g., 5-bromouridine) or 6-azapyrimidines or 6-alkylpyrimidines (e.g. 6-methyluridine), propyne, quinosine, 2-thiouridine, 4-thiouridine, wybutosine, wybutosine, 4-acetyluridine, 5-(carboxyhydroxymethyl)uridine, 5'-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluridine, β -D-galactosylquenosine, 1-methyladenosine, 1-methylinosine, 2,2-dimethylguanosine, 3-methylcytidine, 2-methyladenosine, 2-methylguanosine, N6-methyladenosine, 7-methylguanosine, 5-methoxyaminomethyl-2-thiouridine, 5-methylaminomethyluridine, 5-methylcarbonylmethyluridine, 5-methylxyluridine, 5-methyl-2-thiouridine, 2-methylthio-N6-isopentenyladenosine, β -D-mannosylquenosine, uridine-5-oxyacetic acid, 2-thiocytidine, threonine derivatives and others (Burgin et al., 1996, *Biochemistry*, 35, 14090; Uhlman & Peyman, *supra*). By “modified bases” in this aspect is meant nucleotide bases other than adenine (A), guanine (G), cytosine (C), thymine (T), and uracil (U), as illustrated above; such bases can be used at any position in the antisense molecule. Persons skilled in the art will appreciate that depending on the uses of the oligomers, Ts and Us

are interchangeable. For instance, with other antisense chemistries such as 2'-O-methyl antisense oligonucleotides that are more RNA-like, the T bases may be shown as U.

[0350] As noted above, certain oligonucleotides provided herein include peptide nucleic acids (PNAs). Peptide nucleic acids (PNAs) are analogs of DNA in which the backbone is structurally homomorphous with a deoxyribose backbone, consisting of N-(2-aminoethyl) glycine units to which pyrimidine or purine bases are attached. PNAs containing natural pyrimidine and purine bases hybridize to complementary oligonucleotides obeying Watson-Crick base-pairing rules, and mimic DNA in terms of base pair recognition (Egholm, Buchardt et al. 1993). The backbone of PNAs is formed by peptide bonds rather than phosphodiester bonds, making them well-suited for antisense applications (see structure below). The backbone is uncharged, resulting in PNA/DNA or PNA/RNA duplexes that exhibit greater than normal thermal stability. PNAs are not recognized by nucleases or proteases.

[0351] PNAs may be produced synthetically using any technique known in the art. PNA is a DNA analog in which a polyamide backbone replaces the traditional phosphate ribose ring of DNA. Despite a radical structural change to the natural structure, PNA is capable of sequence-specific binding in a helix form to DNA or RNA. Characteristics of PNA include a high binding affinity to complementary DNA or RNA, a destabilizing effect caused by single-base mismatch, resistance to nucleases and proteases, hybridization with DNA or RNA independent of salt concentration and triplex formation with homopurine DNA. Panagene™ has developed its proprietary Bts PNA monomers (Bts; benzothiazole-2-sulfonyl group) and proprietary oligomerisation process. The PNA oligomerisation using Bts PNA monomers is composed of repetitive cycles of deprotection, coupling and capping. Panagene's patents to this technology include U.S. Pat. No. 6,969,766, U.S. Pat. No. 7,211,668, U.S. Pat. No. 7,022,851, U.S. Pat. No. 7,125,994, U.S. Pat. No. 7,145,006 and U.S. Pat. No. 7,179,896. Representative United States patents that teach the preparation of PNA compounds include, but are not limited to, U.S. Pat. Nos. 5,539,082; 5,714,331; and 5,719,262, each of which is herein incorporated by reference. Further teaching of PNA compounds can be found in Nielsen et al., *Science*, 1991, 254, 1497.

[0352] Also included are “locked nucleic acid” subunits (LNAs). The structures of LNAs are known in the art: for example, Wengel, et al., *Chemical Communications* (1998) 455; *Tetrahedron* (1998) 54, 3607, and *Accounts of Chem. Research* (1999) 32, 301; Obika, et al., *Tetrahedron Letters* (1997) 38, 8735; (1998) 39, 5401, and *Bioorganic Medicinal Chemistry* (2008) 16, 9230.

[0353] Oligonucleotides may incorporate one or more LNAs; in some cases, the compounds may be entirely composed of LNAs. Methods for the synthesis of individual LNA nucleoside subunits and their incorporation into oligonucleotides are known in the art: U.S. Pat. Nos. 7,572,582; 7,569,575; 7,084,125; 7,060,809; 7,053,207; 7,034,133; 6,794,499; and 6,670,461. Typical intersubunit linkers include phosphodiester and phosphorothioate moieties; alternatively, non-phosphorous containing linkers may be employed. A preferred embodiment is an LNA containing compound where each LNA subunit is separated by a DNA subunit (i.e., a deoxyribose nucleotide). Further preferred

compounds are composed of alternating LNA and DNA subunits where the intersubunit linker is phosphorothioate.

[0354] Certain oligonucleotides may comprise morpholino-based subunits bearing base-pairing moieties, joined by uncharged or substantially uncharged linkages. The terms “morpholino oligomer” or “PMO” (phosphoramidate- or phosphorodiamidate morpholino oligomer) refer to an oligonucleotide analog composed of morpholino subunit structures, where (i) the structures are linked together by phosphorus-containing linkages, one to three atoms long, preferably two atoms long, and preferably uncharged or cationic, joining the morpholino nitrogen of one subunit to a 5' exocyclic carbon of an adjacent subunit, and (ii) each morpholino ring bears a purine or pyrimidine or an equivalent base-pairing moiety effective to bind, by base specific hydrogen bonding, to a base in a polynucleotide.

[0355] Variations can be made to this linkage as long as they do not interfere with binding or activity. For example, the oxygen attached to phosphorus may be substituted with sulfur (thiophosphorodiamidate). The 5' oxygen may be substituted with amino or lower alkyl substituted amino. The pendant nitrogen attached to phosphorus may be unsubstituted, monosubstituted, or disubstituted with (optionally substituted) lower alkyl. The purine or pyrimidine base pairing moiety is typically adenine, cytosine, guanine, uracil, thymine or inosine. The synthesis, structures, and binding characteristics of morpholino oligomers are detailed in U.S. Pat. Nos. 5,698,685, 5,217,866, 5,142,047, 5,034,506, 5,166,315, 5,521,063, and 5,506,337, and PCT Appn. Nos. PCT/US07/11435 (cationic linkages) and U.S. Ser. No. 08/012,804 (improved synthesis), all of which are incorporated herein by reference.

[0356] The morpholino subunits may also be linked by non-phosphorus-based intersubunit linkages, as described further below, where at least one linkage is modified with a pendant cationic group as described above. Other oligonucleotide analog linkages which are uncharged in their unmodified state but which could also bear a pendant amine substituent could be used. For example, a 5' nitrogen atom on a morpholino ring could be employed in a sulfamide linkage or a urea linkage (where phosphorus is replaced with carbon or sulfur, respectively) and modified in a manner analogous to the 5'-nitrogen atom in structure (b3) above.

[0357] Certain embodiments include substantially uncharged morpholino oligomers, such as a substantially uncharged phosphorodiamidate-linked morpholino oligomer. A substantially uncharged, phosphorus containing backbone in an oligonucleotide analog is one in which a majority of the subunit linkages, e.g., between 50-100%, typically at least 60% to 100% or 75% or 80% of its linkages, are uncharged at physiological pH, and contain a single phosphorous atom. Examples of morpholino oligonucleotides having phosphorus-containing backbone linkages include phosphoroamidate and phosphorodiamidate-linked morpholino oligonucleotides. Certain embodiments may contain positively charged groups at preferably about 10%-50% of their backbone linkages.

[0358] Properties of the morpholino-based subunits include, for example, the ability to be linked in an oligomeric form by stable, uncharged or positively charged backbone linkages, the ability to support a nucleotide base (e.g., adenine, cytosine, guanine, thymidine, uracil and hypoxanthine) such that the polymer formed can hybridize with a complementary-base target nucleic acid, including target

RNA, T_m values above about 45° C. in relatively short oligonucleotides (e.g., 10-15 bases), the ability of the oligonucleotide to be actively or passively transported into mammalian cells, and the ability of the antisense oligonucleotide:RNA heteroduplex to resist RNase and RNaseH degradation, respectively.

[0359] In certain embodiments, a substantially uncharged oligonucleotide may be modified to include charged linkages, e.g., up to about 1 per every 2-5 uncharged linkages, such as about 4-5 per every 10 uncharged linkages. In certain embodiments, optimal improvement in antisense activity may be seen when about 25% of the backbone linkages are cationic. In certain embodiments, enhancement may be seen with a small number e.g., 10-20% cationic linkages, or where the number of cationic linkages are in the range 50-80%, such as about 60%. In certain embodiments the cationic backbone charges may be further enhanced by distributing the bulk of the charges close of the “center-region” backbone linkages of the antisense oligonucleotide, e.g., in a 20-mer oligonucleotide with 8 cationic backbone linkages, having at least 70% of these charged linkages localized in the 10 centermost linkages.

[0360] Oligonucleotides that target one or more portions of an AARS polynucleotide reference sequence or its complement may be used in any of the therapeutic, diagnostic, or drug screening methods described herein and apparent to persons skilled in the art.

[0361] B. RNA Interference Agents

[0362] Certain embodiments relate to RNA interference (RNAi) agents that target one or more mRNA transcripts of an aminoacyl-tRNA synthetase (AARS) reference polynucleotide, including fragments and splice variants thereof. Also included are methods of use thereof to modulate the levels of a selected AARS transcript, such as an AARS splice variant or endogenous proteolytic fragment.

[0363] The term “double-stranded” means two separate nucleic acid strands comprising a region in which at least a portion of the strands are sufficiently complementary to hydrogen bond and form a duplex structure. The term “duplex” or “duplex structure” refers to the region of a double stranded molecule wherein the two separate strands are substantially complementary, and thus hybridize to each other. “dsRNA” refers to a ribonucleic acid molecule having a duplex structure comprising two complementary and anti-parallel nucleic acid strands (i.e., the sense and antisense strands). Not all nucleotides of a dsRNA must exhibit Watson-Crick base pairs; the two RNA strands may be substantially complementary. The RNA strands may have the same or a different number of nucleotides.

[0364] In certain embodiments, a dsRNA is or includes a region which is at least partially complementary to the target RNA. In certain embodiments, the dsRNA is fully complementary to the target RNA. It is not necessary that there be perfect complementarity between the dsRNA and the target, but the correspondence must be sufficient to enable the dsRNA, or a cleavage product thereof, to direct sequence specific silencing, such as by RNAi cleavage of the target RNA. Complementarity, or degree of homology with the target strand, is typically most critical in the antisense strand. While perfect complementarity, particularly in the antisense strand, is often desired some embodiments can include one or more but preferably 6, 5, 4, 3, 2, or fewer mismatches with respect to the target RNA. The mismatches are most tolerated in the terminal regions, and if present are prefer-

ably in a terminal region or regions, e.g., within 6, 5, 4, or 3 nucleotides of the 5' and/or 3' terminus. The sense strand need only be substantially complementary with the antisense strand to maintain the overall double-strand character of the molecule.

[0365] As used herein, “modified dsRNA” refers to a dsRNA molecule that comprises at least one alteration that renders it more resistant to nucleases (e.g., protein kinase) than an identical dsRNA molecule that recognizes the same target RNA. Modified dsRNAs may include a single-stranded nucleotide overhang and/or at least one substituted nucleotide.

[0366] As used herein, a “nucleotide overhang” refers to the unpaired nucleotide or nucleotides that protrude from the duplex structure when a 3'-end of one RNA strand extends beyond the 5'-end of the other complementary strand, or vice versa. “Blunt” or “blunt end” means that there are no unpaired nucleotides at that end of the dsRNA, i.e., no nucleotide overhang. A “blunt ended” dsRNA is a dsRNA that is double stranded over its entire length, i.e., no nucleotide overhang at either end of the molecule.

[0367] The term “terminal base pair,” as used herein, refers to the last nucleotide base pair on one end of the duplex region of a double-stranded molecule. For example, if a dsRNA or other molecule is blunt ended (i.e., has no nucleotide overhangs), the last nucleotide base pairs at both ends of the molecule are terminal base pairs. Where a dsRNA or other molecule has a nucleotide overhang at one or both ends of the duplex structure, the last nucleotide base pair(s) immediately adjacent the nucleotide overhang(s) is the terminal base pair at that end(s) of the molecule.

[0368] In certain embodiments, the methods provided herein may utilize double-stranded ribonucleic acid (dsRNA) molecules as modulating agents, for reducing expression of an AARS transcript such as a selected fragment or splice variant. dsRNAs generally comprise two single strands. One strand of the dsRNA comprises a nucleotide sequence that is substantially identical to a portion of the target gene or target region (the “sense” strand), and the other strand (the “complementary” or “antisense” strand) comprises a sequence that is substantially complementary to a portion of the target region. The strands are sufficiently complementary to hybridize to form a duplex structure. In certain embodiments, the complementary RNA strand may be less than 30 nucleotides, less than 25 nucleotides in length, or even 19 to 24 nucleotides in length. In certain aspects, the complementary nucleotide sequence may be 20-23 nucleotides in length, or 22 nucleotides in length.

[0369] In certain embodiments, at least one of the RNA strands comprises a nucleotide overhang of 1 to 4 nucleotides in length. In other embodiments, the dsRNA may further comprise at least one chemically modified nucleotide. In certain aspects, a dsRNA comprising a single-stranded overhang of 1 to 4 nucleotides may comprise a molecule wherein the unpaired nucleotide of the single-stranded overhang that is directly adjacent to the terminal nucleotide pair contains a purine base. In other aspects, the last complementary nucleotide pairs on both ends of a dsRNA are a G-C pair, or, at least two of the last four terminal nucleotide pairs are G-C pairs.

[0370] Certain embodiments of the present invention may comprise microRNAs. Micro-RNAs represent a large group of small RNAs produced naturally in organisms, some of which regulate the expression of target genes. Micro-RNAs

are formed from an approximately 70 nucleotide single-stranded hairpin precursor transcript by Dicer. (V. Ambros et al. *Current Biology* 13:807, 2003). Certain micro-RNAs may be transcribed as hairpin RNA precursors, which are then processed to their mature forms by Dicer enzyme.

[0371] Certain embodiments may also employ short-interfering RNAs (siRNA). In certain embodiments, the first strand of the double-stranded oligonucleotide contains two more nucleoside residues than the second strand. In other embodiments, the first strand and the second strand have the same number of nucleosides; however, the first and second strands may be offset such that the two terminal nucleosides on the first and second strands are not paired with a residue on the complementary strand. In certain instances, the two nucleosides that are not paired are thymidine residues.

[0372] Also included are short hairpin RNAs (shRNAs) and micro RNAs (miRNAs). A double-stranded structure of an shRNA is formed by a single self-complementary RNA strand, and RNA duplex formation may be initiated either inside or outside the cell. MicroRNAs (miRNAs) are small non-coding RNAs of 20-22 nucleotides, typically excised from ~70 nucleotide foldback RNA precursor structures known as pre-miRNAs.

[0373] In instances when the modulating agent comprises siRNA, the agent should include a region of sufficient homology to the target region, and be of sufficient length in terms of nucleotides, such that the siRNA agent, or a fragment thereof, can mediate down regulation of the target RNA. It will be understood that the term “ribonucleotide” or “nucleotide” can, in the case of a modified RNA or nucleotide surrogate, also refer to a modified nucleotide, or surrogate replacement moiety at one or more positions. Thus, an siRNA agent is or includes a region which is at least partially complementary to the target RNA, as described herein.

[0374] In addition, an siRNA modulating agent may be modified or include nucleoside surrogates. Single stranded regions of an siRNA agent may be modified or include nucleoside surrogates, e.g., the unpaired region or regions of a hairpin structure, e.g., a region which links two complementary regions, can have modifications or nucleoside surrogates. Modification to stabilize one or more 3'- or 5'-terminus of an siRNA agent, e.g., against exonucleases, or to favor the antisense siRNA agent to enter into RISC are also useful. Modifications can include C3 (or C6, C7, C12) amino linkers, thiol linkers, carboxyl linkers, non-nucleotidic spacers (C3, C6, C9, C12, abasic, triethylene glycol, hexaethylene glycol), special biotin or fluorescein reagents that come as phosphoramidites and that have another DMT-protected hydroxyl group, allowing multiple couplings during RNA synthesis.

[0375] siRNA agents may include, for example, molecules that are long enough to trigger the interferon response (which can be cleaved by Dicer (Bernstein et al. 2001. *Nature*, 409:363-366) and enter a RISC (RNAi-induced silencing complex)), in addition to molecules which are sufficiently short that they do not trigger the interferon response (which molecules can also be cleaved by Dicer and/or enter a RISC), e.g., molecules which are of a size which allows entry into a RISC, e.g., molecules which resemble Dicer-cleavage products. An siRNA modulating agent, or a cleavage product thereof, can down regulate a target gene, e.g., by inducing RNAi with respect to a target RNA, preferably an AARS target such as a selected splice variant.

[0376] Each strand of an siRNA agent can be equal to or less than 35, 30, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, or 15 nucleotides in length. The strand is preferably at least 19 nucleotides in length. For example, each strand can be between 21 and 25 nucleotides in length. Preferred siRNA agents have a duplex region of 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotide pairs, and one or more overhangs, preferably one or two 3' overhangs, of 2-3 nucleotides.

[0377] In addition to homology to target RNA and the ability to down regulate a target gene, an siRNA agent may have one or more of the following properties: it may, despite modifications, even to a very large number, or all of the nucleosides, have an antisense strand that can present bases (or modified bases) in the proper three dimensional framework so as to be able to form correct base pairing and form a duplex structure with a homologous target RNA which is sufficient to allow down regulation of the target, e.g., by cleavage of the target RNA; it may, despite modifications, even to a very large number, or all of the nucleosides, still have "RNA-like" properties, i.e., it may possess the overall structural, chemical and physical properties of an RNA molecule, even though not exclusively, or even partly, of ribonucleotide-based content. For example, an siRNA agent can contain, e.g., a sense and/or an antisense strand in which all of the nucleotide sugars contain e.g., 2' fluoro in place of 2' hydroxyl. This deoxyribonucleotide-containing agent can still be expected to exhibit RNA-like properties. While not wishing to be bound by theory, the electronegative fluorine prefers an axial orientation when attached to the C2' position of ribose. This spatial preference of fluorine can, in turn, force the sugars to adopt a C_{3'}-endo pucker. This is the same puckering mode as observed in RNA molecules and gives rise to the RNA-characteristic A-family-type helix. Further, since fluorine is a good hydrogen bond acceptor, it can participate in the same hydrogen bonding interactions with water molecules that are known to stabilize RNA structures. Generally, it is preferred that a modified moiety at the 2' sugar position will be able to enter into H-bonding which is more characteristic of the OH moiety of a ribonucleotide than the H moiety of a deoxyribonucleotide.

[0378] A "single strand RNAi agent" as used herein, is an RNAi agent which is made up of a single molecule. It may include a duplexed region, formed by intra-strand pairing, e.g., it may be, or include, a hairpin or pan-handle structure. Single strand RNAi modulating agents are preferably antisense with regard to the target molecule. A single strand RNAi agent should be sufficiently long that it can enter the RISC and participate in RISC mediated cleavage of a target mRNA. A single strand RNAi agent is at least 14, and more preferably at least 15, 20, 25, 29, 35, 40, or 50 nucleotides in length. It is preferably less than 200, 100, or 60 nucleotides in length.

[0379] Hairpin RNAi modulating agents may have a duplex region equal to or at least 17, 18, 19, 20, 21, 22, 23, 24, or 25 nucleotide pairs. The duplex region may preferably be equal to or less than 200, 100, or 50, in length. Certain ranges for the duplex region are 15-30, 17 to 23, 19 to 23, and 19 to 21 nucleotides pairs in length. The hairpin may have a single strand overhang or terminal unpaired region, preferably the 3', and preferably of the antisense side of the hairpin. In certain embodiments, overhangs are 2-3 nucleotides in length.

[0380] Certain modulating agents utilized according to the methods provided herein may comprise RNAi oligonucle-

otides such as chimeric oligonucleotides, or "chimeras," which contain two or more chemically distinct regions, each made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotide compound. These oligonucleotides typically contain at least one region wherein the oligonucleotide is modified so as to confer upon the oligonucleotide increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate oligodeoxynucleotides. Chimeric oligonucleotides may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleotides and/or oligonucleotide mimetics as described above. Such oligonucleotides have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures include, but are not limited to, U.S. Pat. Nos. 5,013,830; 5,149,797; 5,220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; 5,700,922; and 5,955,589, each of which is herein incorporated by reference. In certain embodiments, the chimeric oligonucleotide is RNA-DNA, DNA-RNA, RNA-DNA-RNA, DNA-RNA-DNA, or RNA-DNA-RNA-DNA, wherein the oligonucleotide is between 5 and 60 nucleotides in length.

[0381] In one aspect of the invention RNAi agents relate to an oligonucleotide comprising at least one ligand tethered to an altered or non-natural nucleobase. A large number of compounds can function as the altered base. The structure of the altered base is important to the extent that the altered base should not substantially prevent binding of the oligonucleotide to its target, e.g., mRNA. In certain embodiments, the altered base is difluorotolyl, nitropyrrolyl, nitroimidazolyl, nitroindolyl, naphthalenyl, anthracenyl, pyridinyl, quinolinyl, pyrenyl, or the divalent radical of any one of the non-natural nucleobases described herein. In certain embodiments, the non-natural nucleobase is difluorotolyl, nitropyrrolyl, or nitroimidazolyl. In certain embodiments, the non-natural nucleobase is difluorotolyl. A wide variety of ligands are known in the art and are amenable to the present invention. For example, the ligand can be a steroid, bile acid, lipid, folic acid, pyridoxal, B12, riboflavin, biotin, aromatic compound, polycyclic compound, crown ether, intercalator, cleaver molecule, protein-binding agent, or carbohydrate. In certain embodiments, the ligand is a steroid or aromatic compound. In certain instances, the ligand is cholesteryl.

[0382] In other embodiments, the RNAi agent is an oligonucleotide tethered to a ligand for the purposes of improving cellular targeting and uptake. For example, an RNAi agent may be tethered to an antibody, or antigen binding fragment thereof. As an additional example, an RNAi agent may be tethered to a specific ligand binding molecule, such as a polypeptide or polypeptide fragment that specifically binds a particular cell-surface receptor.

[0383] In other embodiments, the modulating agent comprises a non-natural nucleobase, as described herein. In certain instances, the ribose sugar moiety that naturally occurs in nucleosides is replaced with a hexose sugar. In certain aspects, the hexose sugar is an allose, altrose, glucose, mannose, gulose, idose, galactose, talose, or a derivative thereof. In a preferred embodiment, the hexose is a D-hexose. In certain instances, the ribose sugar moiety that

naturally occurs in nucleosides is replaced with a polycyclic heteroalkyl ring or cyclohexenyl group. In certain instances, the polycyclic heteroalkyl group is a bicyclic ring containing one oxygen atom in the ring. In certain instances, the polycyclic heteroalkyl group is a bicyclo[2.2.1]heptane, a bicyclo[3.2.1]octane, or a bicyclo[3.3.1]nonane. Examples of modified RNAi agents also include oligonucleotides containing modified backbones or non-natural internucleoside linkages, as described herein.

[0384] The present invention further encompasses oligonucleotides employing ribozymes. Synthetic RNA molecules and derivatives thereof that catalyze highly specific endoribonuclease activities are known as ribozymes. (see, e.g., U.S. Pat. No. 5,543,508 to Haseloff et al., and U.S. Pat. No. 5,545,729 to Goodchild et al.) The cleavage reactions are catalyzed by the RNA molecules themselves. In naturally occurring RNA molecules, the sites of self-catalyzed cleavage are located within highly conserved regions of RNA secondary structure (Buzayan et al., *Proc. Natl. Acad. Sci. U.S.A.*, 1986, 83, 8859; Forster et al., *Cell*, 1987, 50, 9). Naturally occurring autocatalytic RNA molecules have been modified to generate ribozymes which can be targeted to a particular cellular or pathogenic RNA molecule with a high degree of specificity. Thus, ribozymes serve the same general purpose as antisense oligonucleotides (i.e., modulation of expression of a specific gene) and, like oligonucleotides, are nucleic acids possessing significant portions of single-strandedness.

[0385] In certain instances, the RNAi agents or antisense oligonucleotides for use with the methods provided herein may be modified by non-ligand group. A number of non-ligand molecules have been conjugated to oligonucleotides in order to enhance the activity, cellular distribution, cellular targeting, or cellular uptake of the oligonucleotide, and procedures for performing such conjugations are available in the scientific literature. Such non-ligand moieties have included lipid moieties, such as cholesterol (Letsinger et al., *Proc. Natl. Acad. Sci. USA*, 1989, 86:6553), arginine-rich peptides, cholic acid (Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1994, 4:1053), a thioether, e.g., hexyl-5-tritylthiol (Manoharan et al., *Ann. N.Y. Acad. Sci.*, 1992, 660:306; Manoharan et al., *Bioorg. Med. Chem. Lett.*, 1993, 3:2765), a thiocholesterol (Oberhauser et al., *Nucl. Acids Res.*, 1992, 20:533), an aliphatic chain, e.g., dodecandiol or undecyl residues (Saison-Behmoaras et al., *EMBO J.*, 1991, 10:111; Kabanov et al., *FEBS Lett.*, 1990, 259:327; Svinarchuk et al., *Biochimie*, 1993, 75:49), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651; Shea et al., *Nucl. Acids Res.*, 1990, 18:3777), a polyamine or a polyethylene glycol chain (Manoharan et al., *Nucleosides & Nucleotides*, 1995, 14:969), or adamantane acetic acid (Manoharan et al., *Tetrahedron Lett.*, 1995, 36:3651), a palmitoyl moiety (Mishra et al., *Biochim. Biophys. Acta*, 1995, 1264:229), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Croke et al., *J. Pharmacol. Exp. Ther.*, 1996, 277:923). Representative United States patents that teach the preparation of such oligonucleotide conjugates have been listed above. Typical conjugation protocols involve the synthesis of oligonucleotides bearing an aminolinker at one or more positions of the sequence. The amino group is then reacted with the molecule being conjugated using appropriate coupling or activating reagents. The conjugation reaction may

be performed either with the oligonucleotide still bound to the solid support or following cleavage of the oligonucleotide in solution phase. Purification of the oligonucleotide conjugate by HPLC typically affords the pure conjugate.

[0386] Additional examples of RNAi agents may be found in U.S. Application Publication Nos. 2007/0275465, 2007/0054279, 2006/0287260, 2006/0035254, 2006/0008822, which are incorporated by reference. Also included are vector delivery systems that are capable of expressing the AARS-targeting sequences described herein. Included are vectors that express siRNA or other duplex-forming RNA interference molecules.

[0387] A vector or nucleic acid construct system can comprise a single vector or plasmid, two or more vectors or plasmids, which together contain the total DNA to be introduced into the genome of the host cell, or a transposon. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. In the present case, the vector or nucleic acid construct is preferably one which is operably functional in a mammalian cell, such as a muscle cell. The vector can also include a selection marker such as an antibiotic or drug resistance gene, or a reporter gene (i.e., green fluorescent protein, luciferase), that can be used for selection or identification of suitable transformants or transfectants. Exemplary delivery systems may include viral vector systems (i.e., viral-mediated transduction) including, but not limited to, retroviral (e.g., lentiviral) vectors, adenoviral vectors, adeno-associated viral vectors, and herpes viral vectors, among others known in the art.

XI. DRUG DISCOVERY

[0388] Certain embodiments relate to the use of AARS polypeptides, antibodies, or polynucleotides in drug discovery, typically to identify agents that modulate one or more of the non-canonical activities of the reference AARS polypeptide, e.g., the AARS protein fragment. For example, certain embodiments include methods of identifying one or more "cellular binding partners" of an AARS reference polypeptide, such as a cellular protein, lipid, nucleic acid or other host molecule that directly or physically interacts with the AARS polypeptide. Particular examples include for example cell-surface receptors, such as GPCRs, protein-protein interaction domains, and extracellular or intracellular domains thereof.

[0389] Also included are methods of identifying host molecules that participate in one or more non-canonical activities of the AARS polypeptide, including molecules that directly or indirectly interact with the cellular binding partner, and either regulate its role in a non-canonical activity, or are regulated by the binding partner. Such host molecules include both upstream and downstream components of the non-canonical pathway, typically related by about 1, 2, 3, 4, 5 or more identifiable steps in the pathway, relative to the cellular binding partner/AARS protein interaction.

[0390] Certain aspects include methods of identifying a compound (e.g., polypeptide) or other agent that agonizes or antagonizes the non-canonical activity of an AARS reference polypeptide or active variant thereof, such as by interacting with the AARS polypeptide and/or one or more of its cellular binding partners. Also included are methods of identifying agents that modulate the expression (e.g., splicing) of AARS splice variants, or modulate the activity of

proteases that otherwise regulate the production of endogenous AARS protein fragments (resectins) at the protein level.

[0391] Certain embodiments therefore include methods of identifying a binding partner of an AARS reference polypeptide, comprising a) combining the AARS polypeptide with a biological sample under suitable conditions, and b) detecting specific binding of the AARS polypeptide to a binding partner, thereby identifying a binding partner that specifically binds to the AARS reference polypeptide. Also included are methods of screening for a compound that specifically binds to an AARS reference polypeptide or a binding partner of the AARS polypeptide, comprising a) combining the polypeptide or the binding partner with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide or the binding partner to the test compound, thereby identifying a compound that specifically binds to the polypeptide or its binding partner. In certain embodiments, the compound is a polypeptide or peptide. In certain embodiments, the compound is a small molecule or other (e.g., non-biological) chemical compound. In certain embodiments, the compound is a peptide mimetic.

[0392] Any method suitable for detecting protein-protein interactions may be employed for identifying cellular proteins that interact with an AARS reference polypeptide, interact with one or more of its cellular binding partners, or both. Examples of traditional methods that may be employed include co-immunoprecipitation, cross-linking, and co-purification through gradients or chromatographic columns of cell lysates or proteins obtained from cell lysates, mainly to identify proteins in the lysate that interact with the AARS polypeptide.

[0393] In these and related embodiments, at least a portion of the amino acid sequence of a protein that interacts with an AARS polypeptide or its binding partner can be ascertained using techniques well known to those of skill in the art, such as via the Edman degradation technique. See, e.g., Creighton *Proteins: Structures and Molecular Principles*, W. H. Freeman & Co., N.Y., pp. 34-49, 1983. The amino acid sequence obtained may be used as a guide for the generation of oligonucleotide mixtures that can be used to screen for gene sequences encoding such proteins. Screening may be accomplished, for example, by standard hybridization or PCR techniques, as described herein and known in the art. Techniques for the generation of oligonucleotide mixtures and the screening are well known. See, e.g., Ausubel et al. *Current Protocols in Molecular Biology* Green Publishing Associates and Wiley Interscience, N.Y., 1989; and Innis et al., eds. *PCR Protocols: A Guide to Methods and Applications* Academic Press, Inc., New York, 1990.

[0394] Additionally, methods may be employed in the simultaneous identification of genes that encode the binding partner or other polypeptide. These methods include, for example, probing expression libraries, in a manner similar to the well known technique of antibody probing of lambda-gt11 libraries, using labeled AARS protein, or another polypeptide, peptide or fusion protein, e.g., a variant AARS polypeptide or AARS domain fused to a marker (e.g., an enzyme, fluor, luminescent protein, or dye), or an Ig-Fc domain.

[0395] One method that detects protein interactions in vivo, the two-hybrid system, is described in detail for illustration only and not by way of limitation. One example

of this system has been described (Chien et al., *PNAS USA* 88:9578-9582, 1991) and is commercially available from Clontech (Palo Alto, Calif.).

[0396] Briefly, utilizing such a system, plasmids may be constructed that encode two hybrid proteins: one plasmid consists of nucleotides encoding the DNA-binding domain of a transcription activator protein fused to an AARS reference nucleotide sequence (or, in certain embodiments, its binding partner), or a variant thereof, and the other plasmid consists of nucleotides encoding the transcription activator protein's activation domain fused to a cDNA (or collection of cDNAs) encoding an unknown protein(s) that has been recombined into the plasmid as part of a cDNA library. The DNA-binding domain fusion plasmid and the activator cDNA library may be transformed into a strain of the yeast *Saccharomyces cerevisiae* that contains a reporter gene (e.g., HBS or lacZ) whose regulatory region contains the transcription activator's binding site. Either hybrid protein alone cannot activate transcription of the reporter gene: the DNA-binding domain hybrid cannot because it does not provide activation function and the activation domain hybrid cannot because it cannot localize to the activator's binding sites. Interaction of the two hybrid proteins reconstitutes the functional activator protein and results in expression of the reporter gene, which is detected by an assay for the reporter gene product.

[0397] The two-hybrid system or other such methodology may be used to screen activation domain libraries for proteins that interact with the "bait" gene product. By way of example, and not by way of limitation, an AARS reference polypeptide or variant may be used as the bait gene product. An AARS binding partner may also be used as a "bait" gene product. Total genomic or cDNA sequences are fused to the DNA encoding an activation domain. This library and a plasmid encoding a hybrid of a bait AARS gene product fused to the DNA-binding domain are co-transformed into a yeast reporter strain, and the resulting transformants are screened for those that express the reporter gene.

[0398] A cDNA library of the cell line from which proteins that interact with bait AARS gene products are to be detected can be made using methods routinely practiced in the art. For example, the cDNA fragments can be inserted into a vector such that they are translationally fused to the transcriptional activation domain of GAL4. This library can be co-transformed along with the bait gene-GAL4 fusion plasmid into a yeast strain, which contains a lacZ gene driven by a promoter that contains GAL4 activation sequence. A cDNA encoded protein, fused to GAL4 transcriptional activation domain, that interacts with bait gene product will reconstitute an active GAL4 protein and thereby drive expression of the HIS3 gene. Colonies, which express HIS3, can be detected by their growth on Petri dishes containing semi-solid agar based media lacking histidine. The cDNA can then be purified from these strains, and used to produce and isolate the bait AARS gene-interacting protein using techniques routinely practiced in the art.

[0399] Also included are three-hybrid systems, which allow the detection of RNA-protein interactions in yeast. See, e.g., Hook et al., *RNA* 11:227-233, 2005. Accordingly, these and related methods can be used to identify a cellular binding partner of an AARS polypeptide, and to identify other proteins or nucleic acids that interact with the AARS polypeptide, the cellular binding partner, or both.

[0400] Certain embodiments relate to the use of interactome screening approaches. Particular examples include protein domain-based screening (see, e.g., Boxem et al., *Cell*. 134:534-545, 2008; and Yu et al., *Science*. 322:10-110, 2008).

[0401] As noted above, once isolated, binding partners can be identified and can, in turn, be used in conjunction with standard techniques to identify proteins or other compounds with which it interacts. Certain embodiments thus relate to methods of screening for a compound that specifically binds to the binding partner of an AARS reference polypeptide, comprising a) combining the binding partner with at least one test compound under suitable conditions, and b) detecting binding of the binding partner to the test compound, thereby identifying a compound that specifically binds to the binding partner. In certain embodiments, the test compound is a polypeptide. In certain embodiments, the test compound is a chemical compound, such as a small molecule compound or peptide mimetic.

[0402] Certain embodiments include methods of screening for a compound that modulates the activity of an AARS reference polypeptide, comprising a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide. Certain embodiments include methods of screening for a compound that modulates the activity of a binding partner of an AARS reference polypeptide, comprising a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the binding partner, b) assessing the activity of the binding partner in the presence of the test compound, and c) comparing the activity of the binding partner in the presence of the test compound with the activity of the binding partner in the absence of the test compound, wherein a change in the activity of the binding partner in the presence of the test compound is indicative of a compound that modulates the activity of the binding partner. Typically, these and related embodiments include assessing a selected non-canonical activity that is associated with the AARS polypeptide or its binding partner. Included are in vitro and in vivo conditions, such as cell culture conditions.

[0403] Certain embodiments include methods of screening a compound for effectiveness as a full or partial agonist of an AARS reference polypeptide or an active fragment or variant thereof, comprising a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample, typically by measuring an increase in the non-canonical activity of the AARS polypeptide. Certain methods include a) exposing a sample comprising a binding partner of the AARS polypeptide to a compound, and b) detecting agonist activity in the sample, typically by measuring an increase in the selected non-canonical activity of the AARS polypeptide. Certain embodiments include compositions that comprise an agonist compound identified by the method and a pharmaceutically acceptable carrier or excipient.

[0404] Also included are methods of screening a compound for effectiveness as a full or partial antagonist of an

AARS reference polypeptide, comprising a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample, typically by measuring a decrease in the non-canonical activity of the AARS polypeptide. Certain methods include a) exposing a sample comprising a binding partner of the AARS polypeptide to a compound, and b) detecting antagonist activity in the sample, typically by measuring a decrease in the selected non-canonical activity of the AARS polypeptide. Certain embodiments include compositions that comprise an antagonist compound identified by the method and a pharmaceutically acceptable carrier or excipient.

[0405] In certain embodiments, in vitro systems may be designed to identify compounds capable of interacting with or modulating an AARS reference sequence or its binding partner. Certain of the compounds identified by such systems may be useful, for example, in modulating the activity of the pathway, and in elaborating components of the pathway itself. They may also be used in screens for identifying compounds that disrupt interactions between components of the pathway; or may disrupt such interactions directly. One exemplary approach involves preparing a reaction mixture of the AARS polypeptide and a test compound under conditions and for a time sufficient to allow the two to interact and bind, thus forming a complex that can be removed from and/or detected in the reaction mixture.

[0406] In vitro screening assays can be conducted in a variety of ways. For example, an AARS polypeptide, a cellular binding partner, or test compound(s) can be anchored onto a solid phase. In these and related embodiments, the resulting complexes may be captured and detected on the solid phase at the end of the reaction. In one example of such a method, the AARS polypeptide and/or its binding partner are anchored onto a solid surface, and the test compound(s), which are not anchored, may be labeled, either directly or indirectly, so that their capture by the component on the solid surface can be detected. In other examples, the test compound(s) are anchored to the solid surface, and the AARS polypeptide and/or its binding partner, which are not anchored, are labeled or in some way detectable. In certain embodiments, microtiter plates may conveniently be utilized as the solid phase. The anchored component (or test compound) may be immobilized by non-covalent or covalent attachments. Non-covalent attachment may be accomplished by simply coating the solid surface with a solution of the protein and drying. Alternatively, an immobilized antibody, preferably a monoclonal antibody, specific for the protein to be immobilized may be used to anchor the protein to the solid surface. The surfaces may be prepared in advance and stored.

[0407] To conduct an exemplary assay, the non-immobilized component is typically added to the coated surface containing the anchored component. After the reaction is complete, un-reacted components are removed (e.g., by washing) under conditions such that any specific complexes formed will remain immobilized on the solid surface. The detection of complexes anchored on the solid surface can be accomplished in a number of ways. For instance, where the previously non-immobilized component is pre-labeled, the detection of label immobilized on the surface indicates that complexes were formed. Where the previously non-immobilized component is not pre-labeled, an indirect label can be used to detect complexes anchored on the surface; e.g., using a labeled antibody specific for the previously non-immobi-

lized component (the antibody, in turn, may be directly labeled or indirectly labeled with a labeled anti-Ig antibody).

[0408] Alternatively, the presence or absence of binding of a test compound can be determined, for example, using surface plasmon resonance (SPR) and the change in the resonance angle as an index, wherein an AARS polypeptide or a cellular binding partner is immobilized onto the surface of a commercially available sensorchip (e.g., manufactured by BIACORE™) according to a conventional method, the test compound is contacted therewith, and the sensorchip is illuminated with a light of a particular wavelength from a particular angle. The binding of a test compound can also be measured by detecting the appearance of a peak corresponding to the test compound by a method wherein an AARS polypeptide or a cellular binding partner is immobilized onto the surface of a protein chip adaptable to a mass spectrometer, a test compound is contacted therewith, and an ionization method such as MALDI-MS, ESI-MS, FAB-MS and the like is combined with a mass spectrometer (e.g., double-focusing mass spectrometer, quadrupole mass spectrometer, time-of-flight mass spectrometer, Fourier transformation mass spectrometer, ion cyclotron mass spectrometer and the like).

[0409] In certain embodiments, cell-based assays, membrane vesicle-based assays, or membrane fraction-based assays can be used to identify compounds that modulate interactions in the non-canonical pathway of the selected AARS polypeptide. To this end, cell lines that express an AARS polypeptide and/or a binding partner, or a fusion protein containing a domain or fragment of such proteins (or a combination thereof), or cell lines (e.g., COS cells, CHO cells, HEK293 cells, HeLa cells etc.) that have been genetically engineered to express such protein(s) or fusion protein(s) can be used. Test compound(s) that influence the non-canonical activity can be identified by monitoring a change (e.g., a statistically significant change) in that activity as compared to a control or a predetermined amount.

[0410] For embodiments that relate to antisense and RNAi agents, for example, also included are methods of screening a compound for effectiveness in altering expression of an AARS reference polynucleotide, comprising a) exposing a sample comprising the AARS reference polynucleotide to a compound such as a potential antisense oligonucleotide, and b) detecting altered expression of the AARS polynucleotide. In certain non-limiting examples, these and related embodiments can be employed in cell-based assays or in cell-free translation assays, according to routine techniques in the art. Also included are the antisense and RNAi agents identified by such methods.

[0411] Antibodies to AARS protein fragments can also be used in screening assays, such as to identify an agent that specifically binds to an AARS, confirm the specificity or affinity of an agent that binds to an AARS protein fragment, or identify the site of interaction between the agent and the AARS protein fragment. Included are assays in which the antibody is used as a competitive inhibitor of the agent. For instance, an antibody that specifically binds to an AARS protein fragment with a known affinity can act as a competitive inhibitor of a selected agent, and be used to calculate the affinity of the agent for the AARS protein fragment. Also, one or more antibodies that specifically bind to known epitopes or sites of an AARS protein fragment can be used as a competitive inhibitor to confirm whether or not the

agent binds at that same site. Other variations will be apparent to persons skilled in the art.

[0412] Also included are any of the above methods, or other screening methods known in the art, which are adapted for high-throughput screening (HTS). HTS typically uses automation to run a screen of an assay against a library of candidate compounds, for instance, an assay that measures an increase or a decrease in a non-canonical activity, as described herein.

[0413] Any of the screening methods provided herein may utilize small molecule libraries or libraries generated by combinatorial chemistry. Libraries of chemical and/or biological mixtures, such as fungal, bacterial, or algal extracts, are known in the art and can be screened with any of the assays of the invention. Examples of methods for the synthesis of molecular libraries can be found in: (Carell et al., 1994a; Carell et al., 1994b; Cho et al., 1993; DeWitt et al., 1993; Gallop et al., 1994; Zuckermann et al., 1994).

[0414] Libraries of compounds may be presented in solution (Houghten et al., 1992) or on beads (Lam et al., 1991), on chips (Fodor et al., 1993), bacteria, spores (Ladner et al., U.S. Pat. No. 5,223,409, 1993), plasmids (Cull et al., 1992) or on phage (Cwirla et al., 1990; Devlin et al., 1990; Felici et al., 1991; Ladner et al., U.S. Pat. No. 5,223,409, 1993; Scott and Smith, 1990). Embodiments of the present invention encompass the use of different libraries for the identification of small molecule modulators of one or more AARS protein fragments, their cellular binding partners, and/or their related non-canonical activities. Libraries useful for the purposes of the invention include, but are not limited to, (1) chemical libraries, (2) natural product libraries, and (3) combinatorial libraries comprised of random peptides, oligonucleotides and/or organic molecules.

[0415] Chemical libraries consist of structural analogs of known compounds or compounds that are identified as “hits” or “leads” via natural product screening. Natural product libraries are derived from collections of microorganisms, animals, plants, or marine organisms which are used to create mixtures for screening by: (1) fermentation and extraction of broths from soil, plant or marine microorganisms or (2) extraction of plants or marine organisms. Natural product libraries include polyketides, non-ribosomal peptides, and variants (non-naturally occurring) thereof. See, e.g., Cane et al., *Science* 282:63-68, 1998. Combinatorial libraries may be composed of large numbers of peptides, oligonucleotides or organic compounds as a mixture. They are relatively easy to prepare by traditional automated synthesis methods, PCR, cloning or proprietary synthetic methods.

[0416] More specifically, a combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical “building blocks” such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

[0417] For a review of combinatorial chemistry and libraries created therefrom, see, e.g., Huc, I. and Nguyen, R. (2001) *Comb. Chem. High Throughput Screen* 4:53-74; Lepre, C.A. (2001) *Drug Discov. Today* 6:133-140; Peng, S.

X. (2000) *Biomed. Chromatogr.* 14:430-441; Bohm, H. J. and Stahl, M. (2000) *Curr. Opin. Chem. Biol.* 4:283-286; Barnes, C. and Balasubramanian, S. (2000) *Curr. Opin. Chem. Biol.* 4:346-350; Lepre, Enjalbal, C., et al., (2000) *Mass Spectrom. Rev.* 19:139-161; Hall, D. G., (2000) *Nat. Biotechnol.* 18:262-262; Lazo, J. S., and Wipf, P. (2000) *J. Pharmacol. Exp. Ther.* 293:705-709; Houghten, R. A., (2000) *Ann. Rev. Pharmacol. Toxicol.* 40:273-282; Kobayashi, S. (2000) *Curr. Opin. Chem. Biol.* (2000) 4:338-345; Kopylov, A. M. and Spiridonova, V. A. (2000) *Mol. Biol. (Mosk)* 34:1097-1113; Weber, L. (2000) *Curr. Opin. Chem. Biol.* 4:295-302; Dolle, R. E. (2000) *J. Comb. Chem.* 2:383-433; Floyd, C D., et al., (1999) *Prog. Med. Chem.* 36:91-168; Kundu, B., et al., (1999) *Prog. Drug Res.* 53:89-156; Cabilly, S. (1999) *Mol. Biotechnol.* 12:143-148; Lowe, G. (1999) *Nat. Prod. Rep.* 16:641-651; Dolle, R. E. and Nelson, K. H. (1999) *J. Comb. Chem.* 1:235-282; Czarnick, A. W. and Keene, J. D. (1998) *Curr. Biol.* 8:R705-R707; Dolle, R. E. (1998) *Mol. Divers.* 4:233-256; Myers, P. L., (1997) *Curr. Opin. Biotechnol.* 8:701-707; and Pluckthun, A. and Cortese, R. (1997) *Biol. Chem.* 378:443.

[0418] Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 MPS, 390 MPS, Advanced Chem Tech, Louisville Ky., Symphony, Rainin, Woburn, Mass., 433A Applied Biosystems, Foster City, Calif., 9050 Plus, Millipore, Bedford, Mass.). In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, N.J., Asinex, Moscow, Ru, Tripos, Inc., St. Louis, Mo., Chem-Star, Ltd., Moscow, RU, 3D Pharmaceuticals, Exton, Pa., Martek Biosciences, Columbia, Md., etc.).

XII. METHODS OF USE

[0419] Embodiments of the present invention include therapeutic methods of treatment. Accordingly, the AARS agents described herein, including AARS polypeptides, AARS polynucleotides, AARS polynucleotide-based vectors, AARS expressing host cells, antisense oligonucleotides, RNAi agents, as well as binding agents such as peptides, antibodies and antigen-binding fragments, peptide mimetics and other small molecules, can be used to treat a variety of non-limiting diseases or conditions associated with the non-canonical activities of a reference AARS. Examples of such non-canonical activities include modulation of extracellular signaling, modulation of cell proliferation, modulation of cell migration, modulation of cell differentiation (e.g., hematopoiesis, neurogenesis, myogenesis, osteogenesis, and adipogenesis), modulation of apoptosis or other forms of cell death, modulation of angiogenesis, modulation of cell binding, modulation of cellular metabolism, modulation of cytokine production or activity, modulation of cytokine receptor activity, modulation of cellular uptake, or secretion, immunomodulation, modulation of inflammation, modulation of metabolic processes such as glucose control, and the like.

[0420] Included are polynucleotide-based therapies, such as antisense therapies and RNAi interference therapies, which typically relate to reducing the expression of a target molecule, such as an endogenous fragment of an AARS, or a cellular binding partner of an AARS polypeptide, which otherwise contributes to its non-canonical activity. Antisense or RNAi therapies typically antagonize the non-canonical activity, such as by reducing expression of the AARS reference polypeptide. Also included are polypeptides or

peptides, antibodies or antigen-binding fragment, peptide mimetics, or other small molecule-based therapies, which either agonize or antagonize the non-canonical activity of an AARS reference polypeptide, such as by interacting directly with the AARS polypeptide, its cellular binding partner(s), or both.

[0421] These and related embodiments include methods of using the AARS agents or compositions of the present invention for treating a cell, tissue or subject. The cells or tissues that may be treated or modulated by the present invention are preferably mammalian cells or tissues, or more preferably human cells or tissues. Such cells or tissues can be of a healthy state or of a diseased state.

[0422] In certain embodiments, for example, methods are provided for modulating therapeutically relevant cellular activities including, but not limited to, cellular metabolism, cell differentiation, cell proliferation, cellular uptake, cell secretion, cell death, cell mobilization, cell migration, gene transcription, mRNA translation, cell impedance, immune responses, inflammatory responses, and the like, comprising contacting a cell with an AARS agent or composition as described herein. In certain embodiments, the cell is in a subject. Accordingly, the AARS compositions may be employed in treating essentially any cell or tissue or subject that would benefit from modulation of one or more such activities.

[0423] The AARS agents and compositions may also be used in any of a number of therapeutic contexts including, for example, those relating to the treatment or prevention of neoplastic diseases, immune system diseases or conditions (e.g., autoimmune diseases and inflammation), infectious diseases, metabolic diseases, neuronal/neurological diseases, muscular/cardiovascular diseases, diseases associated with aberrant hematopoiesis, diseases associated with aberrant myogenesis, diseases associated with aberrant neurogenesis, diseases associated with aberrant adipogenesis, diseases associated with aberrant osteogenesis, diseases associated with aberrant angiogenesis, diseases associated with aberrant cell survival, diseases associated with aberrant lipid uptake, diseases associated with aging (e.g. hearing loss, peripheral or autonomic neuropathies, senile dementia, retinopathy) and others.

[0424] For example, in certain illustrative embodiments, the AARS compositions of the invention may be used to modulate angiogenesis, e.g., via modulation of endothelial cell proliferation and/or signaling. Endothelial cell proliferation and/or signaling may be monitored using an appropriate cell line (e.g., human microvascular endothelial lung cells (HMVEC-L) and human umbilical vein endothelial cells (HUVEC)), and using an appropriate assay (e.g., endothelial cell migration assays, endothelial cell proliferation assays, tube-forming assays, matrigel plug assays, etc.), many of which are known and available in the art.

[0425] Therefore, in related embodiments, the compositions of the invention may be employed in the treatment of essentially any cell or tissue or subject that would benefit from modulation of angiogenesis. For example, in some embodiments, a cell or tissue or subject experiencing or susceptible to angiogenesis (e.g., an angiogenic condition) may be contacted with a suitable composition of the invention to inhibit an angiogenic condition. In other embodiments, a cell or tissue experiencing or susceptible to insufficient angiogenesis (e.g., an angiostatic condition) may be

contacted with an appropriate composition of the invention in order to interfere with angiostatic activity and/or promote angiogenesis.

[0426] Also included are methods of modulating hematopoiesis and related conditions. Examples of hematopoietic processes that may be modulated by the AARS polypeptides of the invention include, without limitation, the formation of myeloid cells (e.g., erythroid cells, mast cells monocytes/macrophages, myeloid dendritic cells, granulocytes such as basophils, neutrophils, and eosinophils, megakaryocytes, platelets) and lymphoid cells (e.g., natural killer cells, lymphoid dendritic cells, B-cells, and T-cells). Certain specific hematopoietic processes include erythropoiesis, granulopoiesis, lymphopoiesis, megakaryopoiesis, thrombopoiesis, and others. Also included are methods of modulating the trafficking or mobilization of hematopoietic cells, including hematopoietic stem cells, progenitor cells, erythrocytes, granulocytes, lymphocytes, megakaryocytes, and thrombocytes.

[0427] The methods of modulating hematopoiesis may be practiced in vivo, in vitro, ex vivo, or in any combination thereof. These methods can be practiced on any biological sample, cell culture, or tissue that contains hematopoietic stem cells, hematopoietic progenitor cells, or other stem or progenitor cells that are capable of differentiating along the hematopoietic lineage (e.g., adipose tissue derived stem cells). For in vitro and ex vivo methods, stem cells and progenitor cells, whether of hematopoietic origin or otherwise, can be isolated and/or identified according to the techniques and characteristics described herein and known in the art.

[0428] The compositions of the invention may also be useful as immunomodulators for treating anti- or pro-inflammatory indications by modulating the cells that mediate, either directly or indirectly, autoimmune and/or inflammatory diseases, conditions and disorders. The utility of the compositions of the invention as immunomodulators or modulators of inflammation can be monitored using any of a number of known and available techniques in the art including, for example, migration assays (e.g., using leukocytes or lymphocytes) or cell viability assays (e.g., using B-cells, T-cells, monocytes or NK cells).

[0429] "Inflammation" refers generally to the biological response of tissues to harmful stimuli, such as pathogens, damaged cells (e.g., wounds), and irritants. The term "inflammatory response" refers to the specific mechanisms by which inflammation is achieved and regulated, including, merely by way of illustration, immune cell activation or migration, cytokine production, vasodilation, including kinin release, fibrinolysis, and coagulation, among others described herein and known in the art.

[0430] Clinical signs of chronic inflammation are dependent upon duration of the illness, inflammatory lesions, cause and anatomical area affected. (see, e.g., Kumar et al., Robbins Basic Pathology-8th Ed., 2009 Elsevier, London; Miller, L M, Pathology Lecture Notes, Atlantic Veterinary College, Charlottetown, PEI, Canada). Chronic inflammation is associated with a variety of pathological conditions or diseases, including, for example, allergies, Alzheimer's disease, anemia, aortic valve stenosis, arthritis such as rheumatoid arthritis and osteoarthritis, cancer, congestive heart failure, fibromyalgia, fibrosis, heart attack, kidney failure, lupus, pancreatitis, stroke, surgical complications, inflammatory lung disease, inflammatory bowel disease, athero-

sclerosis, neurological disorders, diabetes, metabolic disorders, obesity, and psoriasis, among others described herein and known in the art. Hence, AARS compositions may be used to treat or manage chronic inflammation, modulate any of one or more of the individual chronic inflammatory responses, or treat any one or more diseases or conditions associated with chronic inflammation.

[0431] Criteria for assessing the signs and symptoms of inflammatory and other conditions, including for purposes of making differential diagnosis and also for monitoring treatments such as determining whether a therapeutically effective dose has been administered in the course of treatment, e.g., by determining improvement according to accepted clinical criteria, will be apparent to those skilled in the art and are exemplified by the teachings of e.g., Berkow et al., eds., The Merck Manual, 16th edition, Merck and Co., Rahway, N.J., 1992; Goodman et al., eds., Goodman and Gilman's The Pharmacological Basis of Therapeutics, 10th edition, Pergamon Press, Inc., Elmsford, N.Y., (2001); Avery's Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics, 3rd edition, ADIS Press, Ltd., Williams and Wilkins, Baltimore, Md. (1987); Ebadi, Pharmacology, Little, Brown and Co., Boston, (1985); Osolci al., eds., Remington's Pharmaceutical Sciences, 18th edition, Mack Publishing Co., Easton, Pa. (1990); Katzung, Basic and Clinical Pharmacology, Appleton and Lange, Norwalk, Conn. (1992).

[0432] In other embodiments, the AARS compositions of the invention may be used to modulate cellular proliferation and/or survival and, accordingly, for treating or preventing diseases, disorders or conditions characterized by abnormalities in cellular proliferation and/or survival. For example, in certain embodiments, the AARS compositions may be used to modulate apoptosis and/or to treat diseases or conditions associated with abnormal apoptosis. Apoptosis can be monitored by any of a number of available techniques known and available in the art including, for example, assays that measure fragmentation of DNA, alterations in membrane asymmetry, activation of apoptotic caspases and/or release of cytochrome C and AIF.

[0433] The progress of these and other therapies (e.g., ex vivo therapies) can be readily monitored by conventional methods and assays and based on criteria known to the physician or other persons of skill in the art.

XIII. PHARMACEUTICAL FORMULATIONS, ADMINISTRATION AND KITS

[0434] Embodiments of the present invention include AARS polynucleotides, AARS polypeptides, host cells expressing AARS polypeptides, binding agents, modulatory agents, or other compounds described herein, formulated in pharmaceutically-acceptable or physiologically-acceptable solutions for administration to a cell or an animal, either alone, or in combination with one or more other modalities of therapy. It will also be understood that, if desired, the compositions of the invention may be administered in combination with other agents as well, such as, e.g., other proteins or polypeptides or various pharmaceutically-active agents. There is virtually no limit to other components that may also be included in the compositions, provided that the additional agents do not adversely affect the modulatory or other effects desired to be achieved.

[0435] In the pharmaceutical compositions of the invention, formulation of pharmaceutically-acceptable excipients

and carrier solutions is well-known to those of skill in the art, as is the development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including e.g., oral, parenteral, intravenous, intranasal, subcutaneous, and intramuscular administration and formulation.

[0436] In certain applications, the pharmaceutical or therapeutic compositions of the invention do not stimulate an immune reaction. In other embodiments, the pharmaceutical or therapeutic compositions of the invention, typically comprising one or more AARS polypeptides or polynucleotides, stimulate an immune reaction, such as by serving as an adjuvant in a vaccine or related composition, or being present in a composition together with a separate adjuvant or agent stimulates an immune response.

[0437] In certain embodiments, the AARS agents such as AARS polypeptides, AARS polynucleotides, and antibodies have a solubility that is desirable for the particular mode of administration, such intravenous administration. Examples of desirable solubilities include at least about 1 mg/ml, at least about 10 mg/ml, at least about 25 mg/ml, and at least about 50 mg/ml.

[0438] In certain applications, the pharmaceutical compositions disclosed herein may be delivered via oral administration to a subject. As such, these compositions may be formulated with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard- or soft-shell gelatin capsule, or they may be compressed into tablets, or they may be incorporated directly with the food of the diet.

[0439] In certain circumstances it will be desirable to deliver the pharmaceutical compositions disclosed herein parenterally, subcutaneously, intravenously, intramuscularly, intra-arterially, intrathecally, intraparenchymally, intraventricularly, intraurethrally, intrasternally, intracranially, intrasynovially, or even intraperitoneally as described, for example, in U.S. Pat. No. 5,543,158; U.S. Pat. No. 5,641,515 and U.S. Pat. No. 5,399,363 (each specifically incorporated herein by reference in its entirety). Suitable devices for parenteral administration include needle (including microneedle) injectors, needle-free injectors, and infusion techniques.

[0440] Solutions of the active compounds as free base or pharmacologically acceptable salts may be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions may also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof and in oils. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0441] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions (U.S. Pat. No. 5,466,468, specifically incorporated herein by reference in its entirety). In all cases the form should be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage and should be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and/or vegetable oils. Proper fluidity may be maintained, for example, by the use of a coating, such as lecithin,

by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. The prevention of the action of microorganisms can be facilitated by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

[0442] For parenteral administration in an aqueous solution, for example, the solution should be suitably buffered if necessary and the liquid diluent first rendered isotonic with sufficient saline or glucose. These particular aqueous solutions are especially suitable for intravenous, intramuscular, subcutaneous and intraperitoneal administration. In this connection, a sterile aqueous medium that can be employed will be known to those of skill in the art in light of the present disclosure. For example, one dosage may be dissolved in 1 ml of isotonic NaCl solution and either added to 1000 ml of hypodermoclysis fluid or injected at the proposed site of infusion (see, e.g., *Remington's Pharmaceutical Sciences*, 15th Edition, pp. 1035-1038 and 1570-1580). Some variation in dosage will necessarily occur depending on the condition of the subject being treated. The person responsible for administration will, in any event, determine the appropriate dose for the individual subject. Moreover, for human administration, preparations should meet sterility, pyrogenicity, and the general safety and purity standards as required by FDA Office of Biologics standards.

[0443] Sterile injectable solutions can be prepared by incorporating the active compounds in the required amount in the appropriate solvent with the various other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum-drying and freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0444] The compositions disclosed herein may be formulated in a neutral or salt form. Pharmaceutically-acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like. Upon formulation, solutions will be administered in a manner compatible with the dosage formulation and in such amount as is therapeutically effective. The formulations are easily administered in a variety of dosage forms such as injectable solutions, drug-release capsules, and the like.

[0445] As used herein, "carrier" includes any and all solvents, dispersion media, vehicles, coatings, diluents, antibacterial and antifungal agents, isotonic and absorption

delaying agents, buffers, carrier solutions, suspensions, colloids, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

[0446] The phrase “pharmaceutically-acceptable” refers to molecular entities and compositions that do not produce an allergic or similar untoward reaction when administered to a human. The preparation of an aqueous composition that contains a protein as an active ingredient is well understood in the art. Typically, such compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified.

[0447] In certain embodiments, the pharmaceutical compositions may be delivered by intranasal sprays, inhalation, and/or other aerosol delivery vehicles. Methods for delivering genes, polynucleotides, and peptide compositions directly to the lungs via nasal aerosol sprays have been described e.g., in U.S. Pat. No. 5,756,353 and U.S. Pat. No. 5,804,212 (each specifically incorporated herein by reference in its entirety). Likewise, the delivery of drugs using intranasal microparticle resins (Takenaga et al., 1998) and lysophosphatidyl-glycerol compounds (U.S. Pat. No. 5,725,871, specifically incorporated herein by reference in its entirety) are also well-known in the pharmaceutical arts. Likewise, transmucosal drug delivery in the form of a polytetrafluoroethylene support matrix is described in U.S. Pat. No. 5,780,045 (specifically incorporated herein by reference in its entirety).

[0448] The pharmaceutical compositions may be formulated to be immediate and/or sustained release. Sustained release compositions include delayed, modified, pulsed, controlled, targeted and programmed release. Thus the compositions may be formulated as a suspension or as a solid, semi-solid, or thixotropic liquid for administration as an implanted depot providing sustained release of the AARS polynucleotides, AARS polypeptides, binding agents, modulatory agents and other active agents. Examples of such formulations include without limitation, drug-coated stents and semi-solids and suspensions comprising drug-loaded poly(DL-lactic-co-glycolic)acid (PGLA), poly(DL-lactide-co-glycolide) (PLG) or poly(lactide) (PLA) lamellar vesicles or microparticles, hydrogels (Hoffman A S: *Ann. N.Y. Acad. Sci.* 944: 62-73 (2001)), poly-amino acid nanoparticles systems, sold under the trademark MEDUSA® developed by Flamel Technologies Inc., non aqueous gel systems sold under the trademark ATRIGEL® developed by Atrix, Inc., and Sucrose Acetate Isobutyrate Extended Release formulations sold under the trademark SABER® developed by Durect Corporation, and lipid-based systems developed by SkyePharma and sold under the trademark DEPOFOAM®.

[0449] Sustained release devices capable of delivering desired doses of the pharmaceutical compositions over extended periods of time are known in the art. For example, U.S. Pat. Nos. 5,034,229; 5,557,318; 5,110,596; 5,728,396; 5,985,305; 6,113,938; 6,156,331; 6,375,978; and 6,395,292; teach osmotically-driven devices capable of delivering an active agent formulation, such as a solution or a suspension, at a desired rate over an extended period of time (i.e., a

period ranging from more than one week up to one year or more). Other exemplary sustained release devices include regulator-type pumps that provide constant flow, adjustable flow, or programmable flow of beneficial agent formulations, which are available from Medtronic including the Intrathecal pumps sold under the trademark SYNCHROMED INFUSION SYSTEM®, the Johnson and Johnson systems sold under the trademark CODMAN® division pumps, and INSET® technologies pumps. Further examples of devices are described in U.S. Pat. Nos. 6,283,949; 5,976,109; 5,836,935; and 5,511,355.

[0450] In certain embodiments, the delivery may occur by use of liposomes, nanocapsules, microparticles, microspheres, lipid particles, vesicles, and the like, for the introduction of the compositions of the present invention into suitable host cells. In particular, the compositions of the present invention may be formulated for delivery either encapsulated in a lipid particle, a liposome, a vesicle, a nanosphere, a nanoparticle or the like. The formulation and use of such delivery vehicles can be carried out using known and conventional techniques.

[0451] In certain embodiments, the agents provided herein may be attached to a pharmaceutically acceptable solid substrate, including biocompatible and biodegradable substrates such as polymers and matrices. Examples of such solid substrates include, without limitation, polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ -ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as poly(lactic-co-glycolic acid) (PLGA) and the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), poly-D-(-)-3-hydroxybutyric acid, collagen, metal, hydroxyapatite, bioglass, aluminate, bioceramic materials, and purified proteins.

[0452] In one particular embodiment, the solid substrate comprises biodegradable polymers sold under the trademark ATRIGEL™ (QLT, Inc., Vancouver, B.C.). The ATRIGEL® drug delivery system consists of biodegradable polymers dissolved in biocompatible carriers. Pharmaceuticals may be blended into this liquid delivery system at the time of manufacturing or, depending upon the product, may be added later by the physician at the time of use. When the liquid product is injected into the subcutaneous space through a small gauge needle or placed into accessible tissue sites through a cannula, water in the tissue fluids causes the polymer to precipitate and trap the drug in a solid implant. The drug encapsulated within the implant is then released in a controlled manner as the polymer matrix biodegrades with time.

[0453] Pharmaceutical compositions for use in the present invention may also be administered topically, (intra)dermally, or transdermally to the skin or mucosa. Typical formulations for this purpose include gels, hydrogels, lotions, solutions, creams, ointments, dusting powders, dressings, foams, films, skin patches, wafers, implants, sponges, fibers, bandages, and microemulsions. Liposomes may also be used. Typical carriers include alcohol, water, mineral oil, liquid petrolatum, white petrolatum, glycerin, polyethylene glycol, and propylene glycol. Penetration enhancers may be incorporated—see, e.g., Finnin and Morgan: *J. Pharm. Sci.* 88(10): 955-958, (1999). Other means of

topical administration include delivery by electroporation, iontophoresis, phonophoresis, sonophoresis, and microneedle or needle-free injection for example using the systems sold under the trademarks POWDERJECT™, and BIOJECT™.

[0454] Methods of formulation are well known in the art and are disclosed, for example, in Remington: The Science and Practice of Pharmacy, Mack Publishing Company, Easton, Pa., 20th edition, ISBN: 0683306472 (2000). The compositions and agents provided herein may be administered according to the methods of the present invention in any therapeutically effective dosing regimen. The dosage amount and frequency are selected to create an effective level of the agent without harmful effects. The effective amount of a compound of the present invention will depend on the route of administration, the type of warm-blooded animal being treated, and the physical characteristics of the specific warm-blooded animal under consideration. These factors and their relationship to determining this amount are well known to skilled practitioners in the medical arts. This amount and the method of administration can be tailored to achieve optimal efficacy but will depend on such factors as weight, diet, concurrent medication and other factors which those skilled in the medical arts will recognize.

[0455] In particular embodiments, the amount of a composition or agent administered will generally range from a dosage of from about 0.1 to about 100 mg/kg/day, and typically from about 0.1 to 10 mg/kg where administered orally or intravenously. In particular embodiments, a dosage is 5 mg/kg or 7.5 mg/kg. In various embodiments, the dosage is about 50-2500 mg per day, 100-2500 mg/day, 300-1800 mg/day, or 500-1800 mg/day. In one embodiment, the dosage is between about 100 to 600 mg/day. In another embodiment, the dosage is between about 300 and 1200 mg/day. In particular embodiments, the composition or agent is administered at a dosage of 100 mg/day, 240 mg/day 300 mg/day, 600 mg/day, 1000 mg/day, 1200 mg/day, or 1800 mg/day, in one or more doses per day (i.e., where the combined doses achieve the desired daily dosage). In related embodiments, a dosage is 100 mg bid, 150 mg bid, 240 mg bid, 300 mg bid, 500 mg bid, or 600 mg bid. In various embodiments, the composition or agent is administered in single or repeat dosing. The initial dosage and subsequent dosages may be the same or different.

[0456] In certain embodiments, a composition or agent is administered in a single dosage of 0.1 to 10 mg/kg or 0.5 to 5 mg/kg. In other embodiments, a composition or agent is administered in a dosage of 0.1 to 50 mg/kg/day, 0.5 to 20 mg/kg/day, or 5 to 20 mg/kg/day.

[0457] In certain embodiments, a composition or agent is administered orally or intravenously, e.g., by infusion over a period of time of about, e.g., 10 minutes to 90 minutes. In other related embodiments, a composition or agent is administered by continuous infusion, e.g., at a dosage of between about 0.1 to about 10 mg/kg/hr over a time period. While the time period can vary, in certain embodiments the time period may be between about 10 minutes to about 24 hours or between about 10 minutes to about three days.

[0458] In particular embodiments, an effective amount or therapeutically effective amount is an amount sufficient to achieve a total concentration of the composition or agent in the blood plasma of a subject with a C_{max} of between about 0.1 µg/ml and about 20 µg/ml or between about 0.3 µg/ml and about 20 µg/ml. In certain embodiments, an oral dosage

is an amount sufficient to achieve a blood plasma concentration (C_{max}) of between about 0.1 µg/ml to about 5 µg/ml or between about 0.3 µg/ml to about 3 µg/ml. In certain embodiments, an intravenous dosage is an amount sufficient to achieve a blood plasma concentration (C_{max}) of between about 1 µg/ml to about 10 µg/ml or between about 2 µg/ml and about 6 µg/ml. In a related embodiment, the total concentration of an agent in the blood plasma of the subject has a mean trough concentration of less than about 20 µg/ml and/or a steady state concentration of less than about 20 µg/ml. In a further embodiment, the total concentration of an agent in the blood plasma of the subject has a mean trough concentration of less than about 10 µg/ml and/or a steady state concentration of less than about 10 µg/ml.

[0459] In yet another embodiment, the total concentration of an agent in the blood plasma of the subject has a mean trough concentration of between about 1 ng/ml and about 10 µg/ml and/or a steady state concentration of between about 1 ng/ml and about 10 µg/ml. In one embodiment, the total concentration of an agent in the blood plasma of the subject has a mean trough concentration of between about 0.3 µg/ml and about 3 µg/ml and/or a steady state concentration of between about 0.3 µg/ml and about 3 µg/ml.

[0460] In particular embodiments, a composition or agent is administered in an amount sufficient to achieve in the mammal a blood plasma concentration having a mean trough concentration of between about 1 ng/ml and about 10 µg/ml and/or a steady state concentration of between about 1 ng/ml and about 10 µg/ml. In related embodiments, the total concentration of the agent in the blood plasma of the mammal has a mean trough concentration of between about 0.3 µg/ml and about 3 µg/ml and/or a steady state concentration of between about 0.3 µg/ml and about 3 µg/ml.

[0461] In particular embodiments of the present invention, the effective amount of a composition or agent, or the blood plasma concentration of composition or agent is achieved or maintained, e.g., for at least 15 minutes, at least 30 minutes, at least 45 minutes, at least 60 minutes, at least 90 minutes, at least 2 hours, at least 3 hours, at least 4 hours, at least 8 hours, at least 12 hours, at least 24 hours, at least 48 hours, at least 3 days, at least 4 days, at least 5 days, at least 6 days, at least one week, at least 2 weeks, at least one month, at least 2 months, at least 4 months, at least 6 months, at least one year, at least 2 years, or greater than 2 years.

[0462] In certain polypeptide-based embodiments, the amount of polypeptide administered will typically be in the range of about 0.1 µg/kg to about 0.1 mg/kg to about 50 mg/kg of patient body weight. Depending on the type and severity of the disease, about 0.1 µg/kg to about 0.1 mg/kg to about 50 mg/kg body weight (e.g., about 0.1-15 mg/kg/dose) of polypeptide can be an initial candidate dosage for administration to the patient, whether, for example, by one or more separate administrations, or by continuous infusion. For example, a dosing regimen may comprise administering an initial loading dose of about 4 mg/kg, followed by a weekly maintenance dose of about 2 mg/kg of the polypeptide, or about half of the loading dose. However, other dosage regimens may be useful. A typical daily dosage might range from about 0.1 µg/kg to about 1 µg/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs.

[0463] In particular embodiments, the effective dosage achieves the blood plasma levels or mean trough concentration of a composition or agent described herein. These may be readily determined using routine procedures.

[0464] Embodiments of the present invention, in other aspects, provide kits comprising one or more containers filled with one or more of the polypeptides, polynucleotides, antibodies, multiunit complexes, compositions thereof, etc., of the invention, as described herein. The kits can include written instructions on how to use such compositions (e.g., to modulate cellular signaling, angiogenesis, cancer, inflammatory conditions, diagnosis etc.).

[0465] The kits herein may also include a one or more additional therapeutic agents or other components suitable or desired for the indication being treated, or for the desired diagnostic application. An additional therapeutic agent may be contained in a second container, if desired. Examples of additional therapeutic agents include, but are not limited to anti-neoplastic agents, anti-inflammatory agents, antibacterial agents, antiviral agents, angiogenic agents, etc.

[0466] The kits herein can also include one or more syringes or other components necessary or desired to facilitate an intended mode of delivery (e.g., stents, implantable depots, etc.).

[0467] All publications, patent applications, and issued patents cited in this specification are herein incorporated by reference as if each individual publication, patent application, or issued patent were specifically and individually indicated to be incorporated by reference.

[0468] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims. The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

XIV. EXAMPLES

[0469] GENERAL METHODS Unless indicated otherwise in the examples below, the following general methods for gene optimization, small and large scale protein expression, protein purification, transcriptional profiling and screening were used to make and characterize the AARS polypeptides described in the Examples below.

Gene Synthesis and Cloning into Expression Vectors

[0470] Polynucleotide sequences encoding epitope tagged versions of the AARS polypeptides were codon optimized and cloned into bacterial expression vectors using the methods listed below.

[0471] In method (1), *E. coli* codon-optimized DNA (Welch et al., PLoS ONE 4(9): e7007 doi:10.1371/journal.pone.0007002) encoding each AARS polypeptide is synthesized by DNA 2.0 (Menlo Park, Calif.), and two versions of each AARS polypeptide are synthesized, containing either an N-terminal, or C-terminal combined epitope tag comprising both a six histidine tag and V5 epitope tag.

[0472] DNA encoding the N-terminally tagged AARS polypeptides is synthesized with a 5' extension encoding in

5' to 3' orientation, a ribosome binding site (rbs (underlined below)), NdeI restriction site, six histidine tag, and a V5 epitope tag,

(SEQ. ID. No. 1)
(AGGAGGTA~~AAACAT~~ATGCATCATCATCATCATCACGGTAAGCCTATC
CCTAACCC~~TTTGCTCGGTCGATTCTACG~~),

which is fused in frame to the predicted AARS polypeptide open reading frame. In cases where the AARS polypeptide comprises a predicted native initiation methionine (ATG) residue, or the first amino acid residue of the predicted AARS polypeptide is Met, this was deleted. At the end of the predicted AARS polypeptide open reading frame, two stop codons and a XhoI site (TAATGACTCGAG) (SEQ. ID. No. 2) are added.

[0473] DNA encoding the C-terminally tagged AARS polypeptides is synthesized with a 5' extension encoding a rbs (underlined below) and NdeI restriction site that either recapitulates the predicted native start codon for the AARS polypeptide, or inserts an ATG in frame with the predicted AARS polypeptide open reading frame,

(SEQ. ID. No. 3)
(AGGAGATA~~AAACAT~~ATG)

In different embodiments, the ribosome binding site can comprise the sequences "AGGAGGTA~~AAACAT~~" (SEQ. ID. No. 4), "AGGAGATA~~AAACAT~~" (SEQ. ID. No. 5), or GAAGGAGATATACAT (SEQ. ID. No. 6). At the 3' end of the predicted AARS polypeptide open reading frame, a 3' extension is synthesized which encodes in 5' to 3' order, a V5 epitope tag, six histidine tag, two stop codons and a XhoI site, (GGTAAGCCTATCCCTAACCCCTCTCCTCGGTCTCGATTCTACGCACCACCATCATCACCATTAAATGACTCGAG) (SEQ. ID. No. 7), which is fused in frame to the predicted AARS polypeptide open reading frame. If the AARS polypeptide included a predicted native stop codon, this was deleted.

[0474] Synthesized DNA sequences encoding the AARS polypeptides are subcloned into pJExpress411 vector (DNA 2.0). After sequencing to confirm synthesis of the correct product, expression vectors are transformed into bacteria for protein expression as described more fully below.

[0475] In method (2), *E. coli* codon-optimized DNA (Ermolaeva M D (2001) Curr. Iss. Mol. Biol. 3 (4) 91-7) encoding each AARS polypeptide is synthesized by GENEWIZ (South Plainfield, N.J.). Each polynucleotide sequence encoding the AARS polypeptide was synthesized with short 5' and 3' extensions comprising unique restriction sites for subsequent cloning.

[0476] Specifically a BamHI restriction site was inserted at the 5' end of the predicted open reading frame. In cases where the AARS polypeptide comprises a predicted native initiation methionine residue (ATG), or the first amino acid residue of the predicted AARS polypeptide is Met, this was deleted. Additionally a XhoI restriction site was inserted at the 3' end of the predicted open reading frame. In cases where the AARS polypeptide comprises a predicted native stop codon, this was deleted.

[0477] After restriction digestion, the resulting DNA sequences are subcloned into modified pET-24b vectors (EMD, Gibbstown, N.J.) containing either an N-terminal (pET24b_N-6xHisN5), or C-terminal (pET24b_C-V5/6x

His) combined epitope tag comprising both a six histidine and V5 epitope tag (vector modification by GENEWIZ, (South Plainfield, N.J.).

[0478] After restriction digestion, and cloning, the DNA encoding the N-tagged AARS polypeptide is cloned into the N-tagged vector (pET24b_N-6xHisN5), which comprises a 5' DNA sequence encoding six histidines and a V5 epitope tag, (CATATGCATCATCATCATCACGGTAAGCCTATCCCTAACCCCTCTCTCGGTCTCGATTC TACGGGATCC) (SEQ. ID. No. 8), in frame with an initiation codon (ATG) embedded within the NdeI restriction site. This 5' extension is fused to the predicted AARS polypeptide open reading frame through a short 2 amino acid linker (GS).

[0479] At the 3' end of the predicted open reading frame, the DNA encoding the N-tagged AARS polypeptide comprises a DNA sequence encoding a 2 amino acid extension (LE) followed by two termination codons (CTCGAGTAATGA) (SEQ. ID. No. 9).

[0480] After restriction digestion, and cloning, the DNA encoding the C-tagged AARS polypeptide cloned into the C-tagged vector (pET24b_C-V5/6)(His), comprises a 5' sequence encoding an initiation codon (ATG) embedded within the NdeI restriction site which is fused to the predicted AARS polypeptide open reading frame through a short 2 amino acid linker (GS), (CATATGGGATCC) (SEQ. ID. No. 10).

[0481] At the 3' end of the predicted open reading frame, the DNA encoding the C-tagged AARS polypeptide comprises a 3' DNA sequence encoding a short linker 2 amino acid linker (LE) followed by a V5 epitope tag followed by six histidines, and two stop codons,

(SEQ. ID. No. 11)
CTCGAGGGTAAGCCTATCCCTAACCCCTCTCTCGGTCTCGATTCTACG
CACCACCACCACCACCACTAATGA

AARS Polypeptide Expression, Purification and Biophysical Characterization

[0482] 6xHis-tagged AARS polypeptides are expressed in bacteria in a medium-throughput format and/or in larger scale flask cultures depending upon the amount of protein required. AARS polypeptides are purified using affinity and ion exchange chromatography as described below, and as specified for specific experiments.

[0483] Bacterial Cultures:

[0484] 100 ng of expression vector comprising codon optimized DNA encoding each AARS polypeptide (as described above) is transformed into BL21(DE3) (EMD chemicals, cat. no. 69450) competent *E. coli* bacteria at 42° C. for 30 seconds in PCR plates. C41(DE3) (Lucigen, cat. no. 60442), HMS174(DE3) (EMD chemicals, cat. no. 69453) and Origami2(DE3) (EMD chemicals, cat. no. 71345) strains are also evaluated. The plates are placed on ice for 2 minutes and 100 μ L of SOC medium is added, followed by a 1-hour incubation at 37° C. 5 mL of auto-induction medium (EMD chemicals, cat. no. 71491) supplemented with kanamycin (100 μ g/mL) is added into each well of a 24-well block (Qiagen, cat. no. 19583). The transformation reactions are added to the individual wells, the block is sealed with adhesive film (VWR, cat. no. 60941-078) and incubated overnight at 250 rpm in a 37° C. shaker. When low temperature (25° C.) conditions are used, incubation is carried out for 48 hours instead.

[0485] For larger scale expression, 200 mL of auto-induction medium supplemented with kanamycin (100 μ g/mL) is added into 500-mL Erlenmeyer flasks with vent caps (Corning, cat. no. 431401). The transformation reactions are added to the individual flasks and incubated for 30 hours at 250 rpm in a 37° C. shaker.

[0486] Protein Isolation:

[0487] After the culture reached stationary phase (typical OD₆₀₀ of 3-6), the blocks are centrifuged at 3600xg for 10 minutes. The medium is carefully aspirated and the blocks are frozen at -80° C. or -20° C. for 10 minutes. The blocks are then allowed to thaw at room temperature and 1 mL lysis buffer (100 mL Bugbuster supplemented with 200 μ L lyso-nase (EMD chemicals, cat. no. 71370) and protease inhibitors "complete mini EDTA-free" (Roche, cat. no. 11 836 170 001)) is added into each well. The pellets are resuspended by repeat pipetting until no clump is visible and transferred to eppendorf tubes, followed by a 10-20 minute incubation on a shaker at room temperature. After centrifugation at 16,000 g for 10 minutes at 4° C., the lysates are loaded onto a TurboFilter 96 Plate included in the Ni-NTA Superflow 96 BioRobot Kit (Qiagen, cat. no. 969261) and centrifuged at 500 g for 5-10 minutes.

[0488] For larger scale expression, the stationary phase culture is transferred into 500-mL bottles and centrifuged at 6,000 g for 10 minutes. The medium is decanted and the pellet is stored at -80° C. or -20° C. before further processing. The pellet is then allowed to thaw at room temperature and 20 mL lysis buffer is added into each bottle. The pellets are resuspended by repeat pipetting until no clump is visible, followed by 20 minute incubation on a shaker at room temperature. After centrifugation at 10,000 g for 30 minutes at 4° C., the lysates are transferred to clean tubes or bottles. If trace amounts of debris are carried over during the transfer, the sample is centrifuged again or passed through a 0.45 μ m cellulose acetate membrane (Corning, cat. no. 430314) for further clarification.

[0489] Affinity Purification:

[0490] A QIAFilter 96 Plate is loaded with 200 μ L Ni-NTA Superflow slurry included in the Ni-NTA Superflow 96 BioRobot Kit and the resin is equilibrated by adding 600 μ L binding buffer (20 mM sodium phosphate, 500 mM sodium chloride and 10 mM imidazole, pH 7.5). A vacuum of -15 in. Hg is applied until all the buffer has passed through the resin. The clarified cell lysates from the previous step are then loaded onto the QIAFilter® 96 Plate and allowed to bind for 5 minutes. A vacuum of -3 in. Hg is applied for approximately 5 minutes until all the samples have passed through the resin. The resin is then washed with 1 mL binding buffer, followed by two washes with 1 mL binding buffer containing 0.1% Triton X-100. The resin is then washed 10 times with 1 mL binding buffer without Triton X-100. The bound 6xHis-tagged AARS polypeptides are eluted with 450 μ L elution buffer (20 mM sodium phosphate, 500 mM sodium chloride and 500 mM imidazole, pH 7.5) and stored at 4° C.

[0491] For larger scale expression, an empty disposable column "Poly-Prep" (Bio-Rad, cat. no. 731-1550) is loaded with 1 mL Ni-NTA Superflow slurry (Qiagen, cat. no. 30450) and the 0.5 mL resin is equilibrated by adding 5 mL binding buffer. The clarified cell lysate from the previous step is then loaded onto the column and allowed to pass through by gravity. The resin is first washed with 50 mL binding buffer plus 0.1% Triton X-100, then washed with 50

mL binding buffer without Triton X-100. The bound 6xHis-tagged AARS polypeptides are eluted with 2 mL elution buffer and stored at 4° C.

[0492] Desalting and Polishing Steps:

[0493] For AARS polypeptides with a molecular mass of >10 kDa, the Omega 10K membrane of an AcroPrep 96 filter plate (Pall, cat. no. 5034) is rinsed with 20 µL 1xPBS and the plate is placed onto a vacuum manifold (>10 in Hg) until all the liquid passes through. The eluates from the previous step (Ni-NTA) are dispensed into each well and the vacuum applied until all the liquid passes through. These steps are repeated until the total eluate volume (450 µL) has been processed. AARS polypeptides are recovered by adding 180 µL of 1xPBS pH 7.4 to each well, pipetting up and down 10 times carefully and then transferred to a clean block. This step is repeated to yield a total volume of 360 µL per well and the block is stored at 4° C. For AARS polypeptides with a molecular mass of <10 kDa, the eluates from Ni-NTA are loaded onto an Amicon Ultra-15 Centrifugal Filter Unit with Ultracel-3 membrane (Millipore, cat. no. UFC900308), followed by the addition of 10 mL 1xPBS and a centrifugation at 3,600 g for 10-30 minutes until the volume is less than 360 µL. The samples are recovered and 1xPBS is added to a final volume of 360 µL.

[0494] In order to remove endotoxins, an AcroPrep Advance filter plate with Mustang Q membrane (Pall, cat. no. 8171) is rinsed with 300 µL of 1xPBS and centrifuged at 1,000 g for 5 minutes to remove the buffer. The desalted AARS polypeptides (360 µL/well) are added to the filter plate and incubated on a shaker for 5-10 minutes. The plate is then centrifuged at 1,000 g for 5-10 minutes and the flow through fractions containing the AARS polypeptides are collected and stored at 4° C.

[0495] For larger scale expression, the eluates from Ni-NTA are loaded onto an Amicon Ultra-15 Centrifugal Filter Unit with Ultracel-3 or Ultracel-10 membrane (Millipore, cat. no. UFC900308 or UFC901008) depending on the molecular weight of the AARS polypeptide and then centrifuged at 3,600 g for 10-30 minutes until the volume is reduced to 250 µL. The samples are mixed in 10 mL 1xPBS, pH 7.4 and centrifuged again at 3,600 g for 10-30 minutes until the volume is about 250 µL. This step is repeated one more time, the supernatants are recovered and 1xPBS is added to a final volume of 1.5 mL.

[0496] In order to remove endotoxins, a Sartobind Q 5 strong anion exchanger membrane (Sartorius, cat. no. Q5F) is flushed with 1 mL 1xPBS and the AARS polypeptides are slowly passed through the membrane using a plastic syringe. The flow through fraction containing the AARS polypeptides is collected in a 96-deep well block that is sealed and stored at 4° C.

[0497] 6xHis-tagged AARS polypeptides expressed in bacteria and found in inclusion bodies are purified using affinity chromatography and a series of refolding steps, as described below.

[0498] Bacterial Cultures:

[0499] 100 ng of plasmid encoding each AARS polypeptide is transformed into BL21(DE3) (EMD chemicals, cat. no. 69450) or C41(DE3) (Lucigen, cat. no. 60442) competent *E. coli* bacteria at 42° C. for 30 seconds in PCR plates. The plates are placed on ice for 2 minutes and 100 µL of SOC medium is added, followed by a 1-hour incubation at 37° C. 5 mL of auto-induction medium (EMD chemicals, cat. no. 71491) supplemented with kanamycin (100 µg/mL)

is added into each well of a 24-well block (Qiagen, cat. no. 19583). The transformation reactions are added to the individual wells, the block is sealed with adhesive film (VWR, cat. no. 60941-078) and incubated overnight at 250 rpm in a 37° C. shaker.

[0500] For larger scale expression, 200 mL of auto-induction medium supplemented with kanamycin (100 µg/mL) is added into 500-mL Erlenmeyer flasks with vent caps (Corning, cat. no. 431401). The transformation reactions are added to the individual flasks and incubated for 30 hours at 250 rpm in a 37° C. shaker.

[0501] Isolation:

[0502] After the cultures reach stationary phase (typical OD₆₀₀ of 3-6), the blocks are centrifuged at 3,600×g for 10 minutes. The medium is carefully aspirated and the blocks are frozen at -80° C. or -20° C. for 10 minutes. The blocks are then allowed to thaw at room temperature and 1 mL lysis buffer (100 mL Bugbuster supplemented with 200 µL lysozyme (EMD chemicals, cat. no. 71370) and protease inhibitor "complete mini EDTA-free" (Roche, cat. no. 11 836 170 001)) is added into each well. The pellets are resuspended by repeat pipetting until no clump is visible and transferred to eppendorf tubes, followed by a 10-20 minute incubation on a shaker at room temperature. After centrifugation at 16,000×g for 10 minutes at 4° C., the soluble lysates are discarded and the inclusion bodies are thoroughly resuspended in denaturing binding buffer (20 mM sodium phosphate, 500 mM sodium chloride, 6 M guanidine hydrochloride, 10 mM imidazole, pH 7.5). The samples are centrifuged at 16,000 g for 10 minutes and the supernatants loaded onto a TurboFilter 96 Plate included in the Ni-NTA Superflow 96 BioRobot Kit (Qiagen, cat. no. 969261) followed by centrifugation at 500 g for 5-10 minutes. The filtrates are collected in a clean 96-well block (Greiner, cat. no. 780286).

[0503] For larger scale expression, the stationary phase culture is transferred into 500-mL bottles and centrifuged at 6,000 g for 10 minutes. The medium is decanted and the pellet is stored at -80° C. or -20° C. before further processing. The pellet is then allowed to thaw at room temperature and 20 mL lysis buffer is added into each bottle. The pellets are resuspended by repeat pipetting until no clump is visible, followed by 20 minute incubation on a shaker at room temperature. After centrifugation at 10,000 g for 30 minutes at 4° C., the soluble lysates are discarded and the insoluble inclusion bodies thoroughly resuspended in denaturing binding buffer.

[0504] Affinity Purification:

[0505] A QIAFilter 96 Plate is loaded with 200 µL Ni-NTA Superflow slurry included in the Ni-NTA Superflow 96 BioRobot Kit and the resin is equilibrated by adding 600 µL denaturing binding buffer (see above). A vacuum of -15 in. Hg is applied until all of the buffer passes through the resin. The clarified denatured samples from the previous step are then loaded onto the QIAFilter® 96 Plate and allowed to bind for 5 minutes. A vacuum of approximately 3 inches of mercury is applied for approximately 5 minutes until all the samples pass through the resin. The resin is then washed with 1 mL denaturing binding buffer, followed by five washes with 1 mL denaturing binding buffer containing 0.1% Triton X-100. The resin is then washed 15 times with 1 mL denaturing binding buffer without Triton X-100. The bound 6xHis-tagged AARS polypeptides are then eluted with 450 µL denaturing elution buffer (20 mM sodium

phosphate, 500 mM sodium chloride, 6 M guanidine hydrochloride and 500 mM imidazole, pH 7.5) and stored at 4° C.

[0506] For larger scale expression, an empty disposable column "Poly-Prep" (Bio-Rad, cat. no. 731-1550) is loaded with 1 mL Ni-NTA Superflow slurry (Qiagen, cat. no. 30450) and the 0.5 mL resin is equilibrated by adding 5 mL denaturing binding buffer (see above). The denatured inclusion bodies from the previous step are then loaded onto the column and allowed to pass through by gravity. The resin is first washed with 50 mL denaturing binding buffer plus 0.1% Triton X-100, then washed with 50 mL denaturing binding buffer without Triton X-100. The bound 6xHis-tagged AARS polypeptides are eluted with 2 mL denaturing elution buffer and stored at 4° C.

[0507] Refolding:

[0508] For AARS polypeptides >10 kDa, the Omega 10K membrane of an AcroPrep 96 filter plate (Pall, cat. no. 5034) is rinsed with 20 μ L, 1xPBS and the plate is placed onto a vacuum manifold (>10 in. Hg) until all the liquid passes through. The eluates from the previous step (Ni-NTA) are dispensed into each well and the vacuum applied until all the liquid passes through. These steps are repeated until the total eluate volume (450 μ L) has been processed. AARS polypeptides are recovered by adding 200 μ L, of refolding buffer containing 50 mM Tris, 250 mM sodium chloride, 10 mM potassium chloride, 2 mM magnesium chloride, 2 mM calcium chloride, 400 mM sucrose, 500 mM arginine, 1 mM DTT and 0.01% polysorbate 80, pH 7.4) to each well, pipetting up and down 10 times carefully, and then transferred to a clean block. This step is repeated to yield a total volume of 400 μ L per well and the block is placed on the shaker overnight at 4° C. For AARS polypeptides <10 kDa, the eluates from Ni-NTA are loaded onto an Amicon Ultra-15 Centrifugal Filter Unit with Ultracel-3 membrane (Millipore, cat. no. UFC900308), followed by the addition of 10 mL refolding buffer and a centrifugation at 3,600 g for 10-30 minutes until the volume is less than 400 μ L. The samples are recovered and extra refolding buffer is added to a final volume of 400 μ L. The samples are transferred to a 96-well block, sealed with film and placed on a shaker overnight at 4° C.

[0509] For larger scale cultures, the eluates from Ni-NTA are loaded onto an Amicon Ultra-15 centrifugal filter unit with Ultracel-3 or Ultracel-10 membrane (Millipore, cat. no. UFC900308 or UFC901008 depending on the molecular weight of the AARS polypeptide) and then centrifuged at 3,600 g for 10-30 minutes until the volume is reduced to about 500 μ L. For AARS polypeptides with pI >7, the samples are diluted 20-fold in the following buffer: 50 mM sodium acetate, 10 mM sodium chloride, 0.4 mM potassium chloride, 1 mM EDTA, 400 mM sucrose, 500 mM arginine, 1 mM DTT and 0.01% polysorbate 80, pH 6.0. For AARS polypeptides with pI<7, the samples are diluted 20-fold in the following buffer: 50 mM Tris, 250 mM sodium chloride, 10 mM potassium chloride, 2 mM magnesium chloride, 2 mM calcium chloride, 400 mM sucrose, 500 mM arginine, 1 mM DTT and 0.01% polysorbate 80, pH 8.0. The samples are incubated on a shaker at 4° C. overnight.

[0510] Desalting and Polishing Steps:

[0511] After overnight incubation, the 96-well block is centrifuged at 3,600 g to remove any potential aggregates. The supernatants are then subjected to buffer exchange with 1xPBS (Invitrogen, cat. no. 10010). For AARS polypeptides >10 kDa, the Omega 10K membrane of an AcroPrep 96 filter

plate is rinsed with 20 μ L 1xPBS and the plate is placed onto a vacuum manifold (>10 in. Hg) until all the liquid passes through. The samples in the refolding buffer are dispensed into each well and the vacuum applied until all the liquid passes through. These steps are repeated until the total sample volume (400 μ L) has been processed. AARS polypeptides are recovered by adding 180 μ L of 1xPBS pH 7.4 to each well, pipetting up and down 10 times carefully, and then transferred to a clean block. This step is repeated to yield a total volume of 360 μ L per well and the block is stored at 4° C. For AARS polypeptides <10 kDa, the refolded samples are loaded onto an Amicon Ultra-15 Centrifugal Filter Unit with Ultracel-3 membrane (Millipore, cat. no. UFC900308) followed by the addition of 10 mL 1xPBS and centrifugation at 3,600 g for 10-30 minutes until the volume is less than 360 μ L. The samples are recovered and 1xPBS is added to a final volume of 360 μ L.

[0512] In order to remove endotoxins, an AcroPrep Advance filter plate with Mustang Q membrane (Pall, cat. no. 8171) is rinsed with 300 μ L of 1xPBS and centrifuged at 1,000 g for 5 minutes to remove the buffer. The AARS polypeptides (360 μ L/well) are added to the filter plate and incubated on a shaker for 5-10 minutes. The plate is then centrifuged at 1,000 g for 5-10 minutes and the flow through fractions containing the AARS polypeptides are collected and stored at 4° C.

[0513] For larger scale cultures, after overnight incubation, the refolded samples are centrifuged at 10,000 g for 10 minutes to remove any insoluble aggregates. The supernatant is loaded onto an Amicon Ultra-15 Centrifugal Filter Unit and centrifuged at 3,600 g until the volume is reduced to 250 μ L. The samples are mixed in 10 mL 1x PBS and centrifuged again at 3,600 g for 10-30 minutes until the volume is about 250 μ L. Note that the pH of 1xPBS is adjusted to match the pH of the refolding buffer, either pH 6.0 or pH 8.0. This step is repeated one more time, the supernatants are recovered and 1xPBS is added to a final volume of 1.5 mL.

[0514] In order to remove endotoxins, a Sartobind Q 5 strong anion exchanger membrane (Sartorius, cat. no. Q5F) is flushed with 1 mL 1xPBS and the AARS polypeptides are slowly passed through the membrane using a plastic syringe. The flow through fraction containing the AARS polypeptides is collected in a 96-deep well block that is sealed and stored at 4° C.

[0515] Biophysical Characterization:

[0516] All purified AARS polypeptides are analyzed by SDS-PAGE, their concentration determined based on A_{280} and calculated extinction coefficient (ProtParam on ExPASy server). Endotoxin levels are measured by the QCL-1000 Endpoint Chromogenic LAL assay (Lonza, cat. no. 50-648U) according to the manufacturer's instructions.

[0517] Dynamic Light Scattering:

[0518] A Wyatt Technology DynaPro 99 instrument and the temperature controller (20° C.) are warmed up for 15 minutes before the experiment followed by connection of the Dynamics software to the instrument. The acquisition time is set to 10 seconds for multiple acquisitions and the laser power is set to 100%. The quartz cuvette is washed thoroughly with deionized water and methanol before the addition of the protein sample (15 μ L at a concentration of approximately 1 mg/mL in PBS). Air bubbles are removed by tapping the cuvette before it is inserted into the holder with the frosted side to the left. If the intensity is too high

(warning message shown on the screen), the sample is further diluted with PBS until the intensity is decreased to a normal range. The data collected include hydrodynamic radius, polydispersity, predicted average molecular weight, percentage of intensity and percentage of mass.

[0519] Size Exclusion Chromatography:

[0520] The protein sample is diluted to a concentration of about 5-10 mg/mL in PBS before being loaded into a 100 μ L sample loop on the General Electric AKTA FPLC. The Superdex 200 10/300 GL size exclusion column (General Electric, cat. no. 17-5175-01) is used for separation. The column is first equilibrated with 1.5 column volume (CV) of 1 $\times$ PBS buffer, followed by sample injection. The column is run in 1 CV of 1 $\times$ PBS buffer (isocratic flow) with absorbance at 280 nm monitoring. The peak area is integrated and the percentage calculated with the Unicorn software. The elution volume is used to estimate the molecular weight based on comparison with gel filtration calibration kits (General Electric, cat. no. 28-4038-41 and 28-4038-42).

[0521] Protein Recovery upon Storage at High Concentration:

[0522] 10 μ L of the AARS polypeptides concentrated to ≥ 10 mg/mL using an Amicon Ultra-15 filter unit (Millipore, cat. no. UFC901024 or UFC900324 depending on molecular weight) are transferred to a clean microcentrifuge tube. The sample is stored at room temperature for one week followed by centrifugation at 16,000 g for 10 minutes to pellet any precipitates. The concentration of the supernatant is determined by a Bradford protein assay and compared to the concentration measured prior to the week-long exposure to room temperature. The recovery is expressed as percentage of the starting concentration.

[0523] Characterization of AARS Polypeptides by LC-MS:

[0524] Purified AARS polypeptides (1 mg/mL) are diluted 1:10 into 0.1% formic acid and 0.6 μ g protein is loaded with a Dionex autosampler onto a C4 capillary column. The capillary column is prepared by cutting 150 mm of fused silica tubing (0.36 mm OD by 0.1 mm ID, Polymicro Technologies, cat. no. 2000023). The capillary is pulled at one end with a Suter Instrument Laser Fiber Puller and cut with a fused silica cutter to generate a 5 μ m tip. The capillary is packed to the length of 75 mm with C4 resin (5 μ m, 300 Å, Michrom, cat. no. PM5/64300/00) using pressure bomb. The LC-MS analysis is performed on an ThermoFisher LTQ ion trap mass spectrometer coupled to a Dionex Ultimate3000 HPLC system. The analyte is eluted from the column using a 35-minute gradient of 5-70% acetonitrile in 0.1% formic acid at a flow rate of 0.9 μ L/min. The LTQ is operated on a full MS scan mode (300-2,000 m/z) with a spray voltage of 2.5 kV.

[0525] Data collection and analysis: raw mass spectrometry data are stored in RAW files generated by XCalibur running on the LTQ XL mass spectrometer. The MS spectra of the major peaks on the chromatograph are further analyzed with ThermoFisher deconvoluting algorithm ProMass to obtain the AARS polypeptide molecular weights.

Functional Analysis of AARS Polypeptides

[0526] Transcriptional Profiling

[0527] Background and Therapeutic Relevance:

[0528] In addition to traditional target identification techniques, genomic tools have recently emerged as important approaches to aid in elucidating the mechanism of action of

AARS polypeptides and can provide direct insight into therapeutic relevance early in the drug discovery process. To facilitate an understanding of potential therapeutic utility, primary human cell types are cultured with AARS polypeptides and transcriptional profiling is assessed at two separate time points following incubation with AARS polypeptides.

[0529] The cell types chosen for transcriptional profiling are based on the pluripotent capabilities of the cells in question and potential to identify AARS polypeptides of direct therapeutic value. For example, Mesenchymal stem cells (MSCs) can differentiate into osteogenic, adipogenic, chondrogenic, myocardial, or neural lineages when exposed to specific stimuli, making them attractive for understanding the potential relevance of the AARS polypeptides to a broad range of cell types, and diseases.

[0530] In addition to supporting hematopoietic cells, marrow stromal cells can also be induced to differentiate into cells of different connective tissue lineage, such as bone, cartilage, and fat. The potential of Human Mesenchymal stem cells (hMSCs) to maintain multipotency and proliferate extensively in vitro provides new avenues for cell-based therapy in the restoration of damaged or diseased tissue. Recent reports also indicate that hMSCs are capable of cell fate crossing germ layer boundaries. In addition to differentiating into multi-lineages of the mesoderm, these cells can also differentiate into neurons of ectodermal origin and hepatocyte-like cells of endodermal origin. During the process of differentiation, these cells may modify expression patterns of certain lineage specific transcripts.

[0531] Accordingly the ability of specific AARS polypeptides to modulate specific patterns of genes in hMSCs in a time dependent manner demonstrates that these proteins play potentially significant roles in a broad array of differentiation pathways, as well as diseases and disorders resulting from the dysfunction, or deterioration of these processes, or the corresponding cell types. Moreover AARS polypeptides with the ability to modulate gene transcription in MSCs have significant therapeutic utility to enable the in vitro or in vivo modulation of hematopoiesis, neurogenesis, myogenesis, osteogenesis, and adipogenesis, as well as in a broad range of disorders and diseases, including for example inflammatory responses, autoimmunity, cancer, neuronal degeneration, muscular dystrophy, osteoporosis, and lipodystrophy.

[0532] Human Skeletal Muscle Cells (HSkMC) can undergo differentiation to exhibit actin and myosin myofilaments, and have been used in the study of genetic muscular diseases such as Malignant Hyperthermia. HSkMC also have the potential to act as a cardiac graft, mending damage to the heart. Recently, cultured Human Skeletal Muscle cells have been used in micro gravity experiments to study the effects of low gravity environments on Human Skeletal Muscle.

[0533] Accordingly the ability of specific AARS polypeptides to modulate specific patterns of genes in HSkMC in a time dependent manner demonstrates that these proteins play potentially significant roles in the processes of myogenesis, as well as diseases and disorders resulting from the dysfunction, or deterioration of these processes as well as muscle cell development or metabolism. Accordingly AARS polypeptides with the ability to modulate gene transcription in muscle cells have therapeutic utility in a broad range of diseases including for example, the treatment of metabolic

disease, cachexia, various muscle wasting conditions, as well as musculoskeletal diseases.

[0534] Methods:

[0535] The ability of AARS polypeptides to modulate gene expression is assessed using a high-throughput microfluidic real-time quantitative PCR (RT-qPCR) approach (Fluidigm Corporation). (See Petriv et al., (2010) PNAS (doi/10.1073/pnas.1009320107) in Human Marrow Stromal Cells (HMSC) and Human Skeletal Muscle Cells (HskMC). In the experiments reported here, Human HskMC (Cat #150-050 and HMSC (Cat #492-050 were purchased from Cell Applications. HMSC cells are cryopreserved at second passage and can be cultured and propagated to 10 population doublings. Here HMSC in the 6th Passage are used. Human Skeletal Muscle Cells (HskMC) are cryopreserved at second passage and can be cultured and propagated for at least 15 population doublings. In the experiments reported here HskMC at passage 6 post harvest from normal human donor are used.

[0536] In both cases, cells are plated at 50000 cells/mL in 100 μ L, volume of growth media and exposed to AARS polypeptides at a concentration of 250 nM, or as otherwise indicated below, for 24 hours and 72 hours. Controls include Differentiation media with a standard cocktail to promote (1)

Adipogenesis, (2) Osteogenesis, (3) Chondrogenesis and (4) Skeletal muscle myotube formation. Additional controls include untreated wells containing only growth media. Two wells were run for each Differentiation control. Controls: all media was made utilizing DMEM as the basal media. Standard literature was followed and Differentiation media was purchased from Cell Applications. Per the vendor, differentiation media contained the following additives: Skeletal muscle differentiation cocktail: FBS, insulin, glutamine, FGF, EGF; Adipogenesis cocktail: insulin, dexamethasone and IBMX; Osteogenesis cocktail: FBS, dexamethasone, ascorbate 2 phosphate, beta-glycerophosphate; Chondrogenesis cocktail: insulin, ascorbate-2-phosphate, and TGF- β 1.

[0537] Standard protocols for using an ABI (Applied Biosystems, Item # AM1728) TAQMAN® Gene Expression Cells-to-CT™ Kit are utilized to lyse cells and harvest genomic material. An ABI Pre-Amp Mix (Applied Biosystems, Item #4391128) is used to initiate pre-amplification. Gene specific primers are created using a Primer 3 program and purchased from IDT technologies. Fluidigm profiling arrays (Item # BMK-M-96.96) were used for actual quantitative PCR with standard Fluidigm loading reagents and pipetting devices. Table E1 below lists the genes profiled.

TABLE E1

List of genes assessed in transcriptional profiling			
Compiled Unique List	refseq_nt	Full_name__	Synonyms
ABCA1	NM_005502	ATP-binding cassette, sub-family A (ABC1), member 1	ABC-1 ABC1 CERP FLJ14958 HDLDT1 MGC164864 MGC165011 TGD
ACTB	NM_001101	actin, beta	PS1TP5BP1
ACTG1	NM_001614	actin, gamma 1	ACT ACTG DFNA20 DFNA26
ACVR2B	NM_001106	activin A receptor, type IIB	ACTRIIB ActR-IIB MGC116908
APOA1	NM_000039	apolipoprotein A-I	MGC117399
ARNT	NM_178427	aryl hydrocarbon receptor nuclear translocator	HIF-1beta HIF1B HIF1BETA TANGO1bHLHe2
BAD	NM_032989	BCL2-associated agonist of cell death	BBC2 BCL2L8
BCL2	NM_000657	B-cell CLL/lymphoma 2	Bcl-2
BMP2	NM_001200	bone morphogenetic protein 2	BMP2A
BMP4	NM_130851	bone morphogenetic protein 4	BMP2B BMP2B1 MCOPS6 OFC11 ZYME
C3AR1	NM_004054	complement component 3a receptor 1	AZ3B C3AR HNFA09
CASP3	NM_032991	caspase 3, apoptosis-related cysteine peptidase	CPP32 CPP32B SCA-1
CAV1	NM_001753	caveolin 1, caveolae protein, 22 kDa	BSCL3 CGL3 MSTP085 VIP21
CDH5	NM_001795	cadherin 5, type 2 (vascular endothelium)	7B4 CD144 FLJ17376
CFLAR	NM_003879	CASP8 and FADD-like apoptosis regulator	CASH CASP8AP1 CLARP Casper FLAME FLAME-1 FLAME1 FLIP I-FLICE MRIT c-FLIP c-FLIPL c-FLIPR c-FLIPS
COMP	NM_000095	cartilage oligomeric matrix protein	EDM1 EPD1 MED MGC131819 MGC149768 PSACH THBS5
CSF1	NM_172212	colony stimulating factor 1 (macrophage)	MCSF MGC31930
CTGF	NM_001901	connective tissue growth factor	CCN2 HCS24 IGFBP8 MGC102839 NOV2
CTNNB1	NM_001904	catenin (cadherin-associated protein), beta 1, 88 kDa	CTNNB DKFZp686D02253 FLJ25606 FLJ37923

TABLE E1-continued

List of genes assessed in transcriptional profiling			
Compiled Unique List	refseq_nt	Full_name__	Synonyms
DAAM1	NM_014992	dishevelled associated activator of morphogenesis 1	FLJ41657 KIAA0666
ELN	NM_001081755	elastin	FLJ38671 FLJ43523 SVAS WBS WS
ENO1	NM_001428	enolase 1, (alpha)	ENO1L1 MPB1 NNE PPH
FABP3	NM_004102	fatty acid binding protein 3, muscle and heart (mammary-derived growth inhibitor)	FABP11 H-FABP MDGI O-FABP
FAK	NM_001199649	focal adhesion kinase	fak1
FGF4	NM_002007	fibroblast growth factor 4	HBGF-4 HST HST-1 HSTF1 K-FGF KFGF
FIGF	NM_004469	c-fos induced growth factor (vascular endothelial growth factor D)	VEGF-D VEGFD
FLT1	NM_002019	fms-related tyrosine kinase 1 (vascular endothelial growth factor/vascular permeability factor receptor)	FLT VEGFR1
FOXA1	NM_004496	forkhead box A1	HNF3A MGC33105 TCF3A
GAPDH	NM_002046	glyceraldehyde-3-phosphate dehydrogenase	G3PD GAPD MGC88685
GFAP	NM_002055	glial fibrillary acidic protein	FLJ45472
SLC2A4	NM_001042	solute carrier family 2 (facilitated glucose transporter), member 4	GLUT4
HAND1	NM_004821	heart and neural crest derivatives expressed 1	Hxt Thing1 bHLHa27 eHand
HIF1A	NM_181054	hypoxia inducible factor 1, alpha subunit (basic helix-loop-helix transcription factor)	HIF-1alpha HIF1 HIF1-ALPHA MOP1 PASD8 bHLHe78
HK2	NM_000189	hexokinase 2	DKFZp686M1669 HKII HXK2
HMGB1	NM_002128	high-mobility group box 1	DKFZp686A04236 HMG1 HMG3 SBP-1
HNF4A	NM_178850	hepatocyte nuclear factor 4, alpha	FLJ39654 HNF4 HNF4a7 HNF4a8 HNF4a9 HNF4alpha MODY1 MODY1 NR2A1 NR2A21 TCF1 TCF14
HPRT1	NM_000194	hypoxanthine phosphoribosyltransferase 1	HGPRT HPRT
HSPB1	NM_001540	heat shock 27 kDa protein 1	CMT2F DKFZp586P1322 HMN2B HS.76067 HSP27 HSP28 Hsp25 SRP27
ICAM1	NM_000201	intercellular adhesion molecule 1	BB2 CD54 P3.58
IFNG	NM_000619	interferon, gamma	IFG IFI
IGF1	NM_001111285	insulin-like growth factor 1 (somatomedin C)	IGF-I IGF1A IGF1
IGF2	NM_001127598	insulin-like growth factor 2 (somatomedin A)	C11orf43 FLJ22066 FLJ44734 INSIGF pp9974
IGFBP3	NM_001013398	insulin-like growth factor binding protein 3	BP-53 IBP3
IGFBP5	NM_000599	insulin-like growth factor binding protein 5	IBP5
IKKB	NM_001556	inhibitor of kappa light polypeptide gene enhancer in B-cells, kinase beta	FLJ33771 FLJ36218 FLJ38368 FLJ40509 IKK-beta IKK2 IKKB MGC131801 NFKB1KB
IL10	NM_000572	interleukin 10	CSIF IL-10 IL10A MGC126450 MGC126451 TGIF
IL1B	NM_000576	interleukin 1, beta	IL-1 IL1-BETA IL1F2
IL3	NM_000588	interleukin 3 (colony-stimulating factor, multiple)	IL-3 MCGF MGC79398 MGC79399 MULTI-CSF

TABLE E1-continued

List of genes assessed in transcriptional profiling			
Compiled Unique List	refseq_nt	Full_name__	Synonyms
IL4	NM_172348	interleukin 4	BCGF-1 BCGF1 BSF1 IL-4 MGC79402
IL5	NM_000879	interleukin 5 (colony-stimulating factor, eosinophil)	EDF IL-5 TRF
IL6R	NM_181359	interleukin 6 receptor	CD126 IL-6R-1 IL-6R-alpha IL6RA MGC104991
IL8	NM_000584	interleukin 8	CXCL8 GCP-1 GCP1 LECT LUCT LYNAP MDNCF MONAP NAF NAP-1 NAP1
ITGA5	NM_002205	integrin, alpha 5 (fibronectin receptor, alpha polypeptide)	CD49e FNRA VLA5A
KDR	NM_002253	kinase insert domain receptor (a type III receptor tyrosine kinase)	CD309 FLK1 VEGFR VEGFR2
LEP	NM_000230	leptin	FLJ94114 OB OBS
LPL	NM_000237	lipoprotein lipase	HDLCQ11 LIPD
MAPK11	NM_002751	mitogen-activated protein kinase 11	P38B P38BETA2 PRKM11 SAPK2 SAPK2B p38-2 p38Beta
MMP1	NM_002421	matrix metalloproteinase 1 (interstitial collagenase)	CLG CLGN
MMP3	NM_002422	matrix metalloproteinase 3 (stromelysin 1, progelatinase)	CHDS6 MGC126102 MGC126103 MGC126104 MMP-3 SL-1 STMY STMY1 STR1
MYH1	NM_005963	myosin, heavy chain 1, skeletal muscle, adult	MGC133384 MYHSA1 MYHa MyHC-2X/D MyHC-2x
MYH11	NM_022844	myosin, heavy chain 11, smooth muscle	AAT4 DKFZp686D10126 DKFZp686D19237 FAA4 FLJ35232 MGC126726 MGC32963 SMHC SMMHC
MYH7	NM_000257	myosin, heavy chain 7, cardiac muscle, beta	CMD1S CMH1 DKFZp451F047 MGC138376 MGC138378 MPD1 MYHCB SPMD SPMM
MYOD1	NM_002478	myogenic differentiation 1	MYF3 MYOD PUM bHLHc1
NFATC1	NM_172390	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 1	MGC138448 NF-ATC NFAT2 NFATc
NFATC2	NM_173091	nuclear factor of activated T-cells, cytoplasmic, calcineurin-dependent 2	NFAT1 NFATP
NFKB1	NM_003998	nuclear factor of kappa light polypeptide gene enhancer in B-cells 1	DKFZp686C01211 EBP-1 KBF1 MGC54151 NF-kappa-B NF-kappaB NFKB-p105 NFKB-p50 p105 p50
NOS2	NM_000625	nitric oxide synthase 2, inducible	HEP-NOS INOS NOS NOS2A
NOTCH1	NM_017617	notch 1	TAN1 hN1
NR3C1	NM_001024094	nuclear receptor subfamily 3, group C, member 1 (glucocorticoid receptor)	GCCR GCR GR GRL
NRP2	NM_201279	neuropilin 2	MGC126574 NP2 NPN2 PRO2714 VEGF165R2
PAX7	NM_013945	paired box 7	FLJ37460 HUP1 PAX7B RMS2
PDGFB	NM_033016	platelet-derived growth factor beta polypeptide (simian sarcoma viral (v-sis) oncogene homolog)	FLJ12858 PDGF2 SIS SSV c-sis
PDK4	NM_002612	pyruvate dehydrogenase kinase, isozyme 4	FLJ40832
PLA2G1B	NM_000928	phospholipase A2, group IB (pancreas)	MGC119834 MGC119835 PLA2 PLA2A PPLA2
PLIN1	NM_002666	lipid droplet associated protein	Perilipin
PPARG	NM_138712	peroxisome proliferator-activated receptor gamma	CIMT1 GLM1 NR1C3 PPARG1 PPARG2 PPARgamma
QARS	NM_005051	glutaminyl-tRNA synthetase	GLNRS PRO2195
RHOA	NM_001664	ras homolog gene family, member A	ARH12 ARHA RHO12 RHOH12
RUNX1	NM_001754	runt-related transcription factor 1	AML1 AML1-EVI-1 AMLCR1 CBFA2 EVI-1 PEBP2aB
RXRA	NM_002957	retinoid X receptor, alpha	FLJ00280 FLJ00318 FLJ16020 FLJ16733 MGC102720 NR2B1

TABLE E1-continued

List of genes assessed in transcriptional profiling			
Compiled Unique List	refseq_nt	Full_name__	Synonyms
SERPINE1	NM_001165413	serpin peptidase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 1	PAI PAI-1 PAI1 PLANH1
SMAD2	NM_005901	SMAD family member 2	JV18 JV18-1 MADH2 MADR2 MGC22139 MGC34440 hMAD-2 hSMAD2
SMAD4	NM_005359	SMAD family member 4	DPC4 JIP MADH4
TERT	NM_198255	telomerase reverse transcriptase	EST2 TCS1 TP2 TRT hEST2
TGFB1	NM_000660	transforming growth factor, beta 1	CED DPD1 LAP TGFB TGFBeta
TGFB3	NM_003239	transforming growth factor, beta 3	ARVD FLJ16571 TGF-beta3
THBS4	NM_003248	thrombospondin 4	TSP4
TNF	NM_000594	tumor necrosis factor	DIF TNF-alpha TNFA TNFSF2
TUBB	NM_178014	tubulin, beta	M40 MGC117247 MGC16435 OK/SW-cl.56 TUBB1 TUBB5
TUBB1	NM_030773	tubulin, beta 1	tubulin isoform beta (1)
TUBG1	NM_001070	tubulin, gamma 1	GCP-1 TUBG TUBGCP1
VCAM1	NM_080682	vascular cell adhesion molecule 1	CD106 DKFZp779G2333 INCAM-100 MGC99561
VEGFA	NM_003376	vascular endothelial growth factor A	MGC70609 MVCD1 VEGF VPF
VIM	NM_003380	vimentin	FLJ36605
WISP1	NM_080838	WNT1 inducible signaling pathway protein 1	CCN4 WISP1c WISP1i WISP1tc
WNT1	NM_005430	wingless-type MMTV integration site family, member 1	INT1

[0538] Bioinformatics Analysis:

[0539] Data retrieved in .csv format from the Biomark machine by Fluidigm is converted to a tabular format including sample, mRNA, and replicate information along with the raw fluorescence value. PCR reactions that failed are marked as missing. Multiple experiments were combined after normalizing to total expression of mRNA species. All measured mRNA expression is filtered based on the requirement of detection in at least 2 of all of the biological replicates tested. We assessed technical, biological and set deviation mean in entire dataset.

[0540] For data analysis Ct values for all genes of interest are first normalized to the averaged Ct values for house-keeping genes from the corresponding sample to obtain ΔCt values ($\Delta Ct = Ct_{\text{gene}} - Ct_{\text{average housekeeping genes}}$). Genes from each sample are then normalized to the same gene in untreated control to obtain $\Delta\Delta Ct$ values ($\Delta\Delta Ct = \Delta Ct_{\text{control sample}} - \Delta Ct_{\text{treated sample}}$).

[0541] To obtain fold change values up-regulated genes (i.e. $\Delta\Delta Ct$ s greater than 0) are subject to the following calculation: Fold Change = $2^{-\Delta\Delta Ct}$. For down-regulated genes (i.e. $\Delta\Delta Ct$ s less than 0): Fold Change = $-(2^{-|\Delta\Delta Ct|})$.

Cellular Proliferation Assays (Assays A1-A11 in the Data Tables Below)

[0542] Background and Therapeutic Relevance:

[0543] The ability to modulate the rate of cellular proliferation and apoptosis of different cell types represents a fundamental property of many therapeutic compounds, and is of direct relevance to the treatment and prevention of a broad range of diseases and disorders.

[0544] Accordingly AARS polypeptides with the ability to modulate the rate of cellular proliferation and or apoptosis have significant therapeutic utility in a broad range of diseases including, as growth factors, and differentiation factors for stem cells, and in treatment regimens to enhance or suppress the proliferation of specific cell types of interest in vivo or in vitro, including for example, haemopoietic cells, immunomodulatory cells, cancer, and for the treatment and prevention of diseases associated with aging, including for example neurodegeneration, peripheral neuropathy, and loss of muscular and soft tissue tone.

[0545] Methods:

[0546] Effects of the AARS polypeptides on cellular proliferation is assessed using one or more of the methods listed below, and as more specifically elaborated in the methods below.

[0547] Hoechst 33432.

[0548] Standard cell counts to assess proliferation are performed using Hoechst 33432, which is a cell-permeant nuclear counterstain that emits blue fluorescence when bound to dsDNA. It is available as a solution (Invitrogen Cat # H-3570) that is used at a final concentration of 1 ug/mL in either media or PBS. Cells are grown in 96 well plates in the presence of AARS polypeptides for a standard growth time of 48 hours, or longer depending on cell type and as described in the examples below.

[0549] ATP-Lite.

[0550] Cellular ATP levels correlate with cellular health and can be readily determined using a variety of commercially available kits. ATP-lite (Perkin-Elmer, Cat #6016947 Boston, Mass. 02481) which is a homogenous mixture of

lysis solution and ATP-detection reagent. is pre-mixed before use and is used 1:1 v:v ratio with cultured cells. Plates are incubated for 5 minutes to promote lysis and plates are measured using a luminescent plate reader. Cells are grown in 96 well plates in the presence of AARS polypeptides for a standard growth time of 48 hours, or longer depending on cell type and as described in the examples below.

[0551] ALAMARBLUE® (Resazurin) is a cell viability indicator which is based on the redox state of the cells. Resazurin, the active ingredient, is a nontoxic, cell permeable compound that is blue in color and virtually nonfluorescent when present in its oxidized form. However upon entering normal viable cells, resazurin is rapidly reduced to resorufin, which produces a red fluorescence signal. Viable cells continuously convert resazurin to resorufin, thereby generating a quantitative measure of viability—and cytotoxicity. The lack of toxicity allows long-term exposure of cells to resazurin without negative impact; cells grown in the presence of resazurin were found to produce similar numbers of viable cells as control cells, as determined by flow cytometric analysis.

[0552] Measurements are made by adding a solution of Resazurin/ALAMARBLUE® to cells, incubating them for 1-4 hours, and reading the fluorescence or absorbance. The amount of fluorescence or absorbance is proportional to the number of living cells and corresponds to the cells metabolic activity. Damaged and nonviable cells have lower innate metabolic activity and thus generate a proportionally lower signal than healthy cells. After incubation with ALAMARBLUE®, samples can readily be measured on fluorescence and absorbance instrumentation. For fluorescence readings: 530 nm excitation and 590 nm emission filter settings are used.

[0553] Cells are grown in 96 well plates in the presence of AARS polypeptides for a standard growth time of 48 hours, or longer depending on cell type and as described in the examples below.

Acetylated LDL Uptake in HEPG2C3A Human Hepatocyte Cells. (Assay B1 in the Data Tables Below)

[0554] Background and Therapeutic Relevance:

[0555] LDL is the major carrier of cholesterol in the blood, accounting for more than 60% of the cholesterol in plasma. In humans, the hepatic LDL receptor is responsible for clearing around 70% of plasma LDL from circulation. Internalized LDL is degraded to free cholesterol and amino acids in the lysosome. The liver is the most important organ for LDL catabolism and LDL receptor activity in humans. LDL that is not internalized and remains in circulation can be transported by endothelial cells into the vessel wall, resulting in the formation of atherosclerotic plaques. Circulating LDL can also be taken up by macrophages and this can also contribute to the formation of plaques. Increasing LDL uptake into hepatic tissue is thought to be beneficial to human health and finding safe and efficacious therapeutics that may the positively regulate this process may provide new therapies for cardiovascular and metabolic diseases. To investigate whether the unique properties of AARS polypeptides can regulate uptake of acetylated LDL, a standard assay for measuring acetylated LDL uptake is employed in HepG2C3a cells.

[0556] Accordingly AARS polypeptides with the ability to modulate LDL uptake have significant therapeutic utility in a broad range of diseases including for example, the treat-

ment of hypercholesteremia, hyperlipidemia, type 1 and 2 diabetes, metabolic syndrome, and vascular diseases including atherosclerosis

[0557] Methods:

[0558] HEPG2C3a cells (ATCC# CRL-10741) are maintained in Eagles Minimal Essential (EMEM) medium supplemented with 10% FBS (HyClone Cat#SH30910.03), 50 u/mL penicillin/50 µg/mL streptomycin, (Invitrogen) in 15 mL medium in 75 mL flasks. Cells are grown at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. HEPG2C3a express the LDL-receptor and are competent for acetylated LDL uptake when grown on clear bottom collagen coated plates. A 100 µL volume of cells is plated on collagen coated plates (Invitrogen Cat#A11428) overnight in complete medium (above) at a cell density of 50,000 cells/mL. Cells are washed once with PBS (Invitrogen Cat#10010) and 80 µL of serum free EMEM is added to each well. AARS polypeptides at a final concentration of 250 nM per well are added in a consistent volume in sterile PBS to each well. A unique AARS polypeptide is placed in each well. Cells are serum starved and exposed to the AARS polypeptides for 16 hours. Following the 16 hour incubation, the supernatant is collected and soluble ICAM is measured using a standard ELISA kit from RND Systems (Cat # DY643), and serum free media supplemented with 5 µg/mL ac-LDL (Alexa Fluor 488 labeled Cat # L23380, Invitrogen) is added to each well. Following a 2 hour incubation at 37° C. 5% CO₂, cells are washed twice with sterile PBS before 100 µL PBS is added to each well for quantification. Plates were analyzed for total fluorescent intensity using a bottom read on a Victor X5 fluorescent plate reader (Perkin Elmer) at an excitation wavelength centered around 485 nm, and an emission wavelength centered around 535 nm. Cells are stained with Hoechst dye and fluorescent intensity 405 nm Excitation/ 450 nm Emission is read to confirm total cell number is consistent across the plate.

Regulation of Human Neutrophil Oxidative Burst and Elastase Production (Assays C1-C3 in the Data Tables Below)

[0559] Neutrophil Oxidative Burst

[0560] Background and Therapeutic Relevance:

[0561] Phagocytosis by polymorphonuclear neutrophils and monocytes constitutes an essential arm of host defense against infections by microorganisms including bacteria and fungi. The phagocytic process can be separated into several major stages: chemotaxis (migration of phagocytes to inflammatory sites), attachment of particles to the cell surface of phagocytes, ingestion (phagocytosis) and intracellular killing by oxygen-dependent (oxidative burst) and oxygen-independent mechanisms. Reduced or missing burst activity is observed in inborne defects like the chronic granulomatous disease (CGD). CGD is a heterogeneous group of inherited disorders that usually manifests itself during the first two years of life. The disease is characterized by repeated and life-threatening infections caused by bacterial and fungal organisms. These infections typically consist of pneumonia, lymphadenitis, or abscesses that involve lymph nodes, lungs, and liver. The NADPH oxidase is the enzyme system responsible for producing superoxide anion, which is quickly converted to hydrogen peroxide and

hydroxyl radicals. Abnormalities in the constituent peptides of the NADPH oxidase enzyme system lead to the dysfunctions characteristic of CGD. Neutrophils from CGD patients fail to produce a significant oxidative burst following stimulation. Different forms of CGD are described (classical X-linked CGD and autosomal recessive patterns). The oxidative burst of granulocytes is impaired in transplantation, later stages of HIV infection, and in the elderly, making these populations more susceptible to secondary infection and exacerbations of inflammatory disease. Various immunomodulators (e.g., cytokines (GM-CSF, G-CSF, TNF) or drugs) also seem to have effects on the oxidative burst. There is the potential for proteins with the ability to up-regulate or down-regulate oxidative burst in a therapeutic fashion to be useful for a variety of different disease states.

[0562] Methods:

[0563] The protein kinase C ligand phorbol 12-myristate 13-acetate (PMA) can be utilized in this assay as an agonist of the oxidative burst process. Heparinized whole blood is mixed with sterile dextran (0.6% final concentration) for 1 hour and allowed to separate into layers. The lower layer contains neutrophil, monocytes and red blood cells. An ammonium chloride lysis step is utilized to remove all RBCs and a 97% pure population of neutrophils with approximately 3% monocyte contamination remains following lysis step. Upon stimulation, granulocytes and monocytes produce reactive oxygen metabolites (superoxide anion, hydrogen peroxide, hypochlorous acid) which destroy bacteria inside the phagosome. Formation of the reactive oxidants during the oxidative burst can be monitored by the addition and oxidation of Amplex Red. The percentage of cells having produced reactive oxygen radicals are then analyzed as well as their mean fluorescence intensity using a fluorescent plate reader. The typical time course for this reaction is 10 minutes, with obvious burst being seen by 2 minutes and a drop off of signal being seen by 20 minutes. This assay can be run in agonist mode in the absence of PMA or in antagonist mode, with concomitant administration of AARS polypeptides and PMA at a concentration that is below the EC50 for this compound.

Regulation of Human Neutrophil Elastase Production

[0564] Background and Therapeutic Relevance:

[0565] Neutrophil elastase is a serine protease that has been implicated as having a specific role in the development of a wide range of human diseases, including inflammatory disorders of the lung and cardiovascular system. Although its key physiologic role is in innate host defense, it can also participate in tissue remodeling and possesses secretagogue actions that are now recognized as important to local inflammatory signals. Neutrophil elastase activity has been implicated in the development of emphysema for several decades, however only relatively recently has a pathogenetic function been ascribed to this serine proteinase in situations where excessive extracellular matrix deposition occurs. The use of genetically manipulated animal models is starting to uncover the potential ways in which its actions might influence fibrotic lung repair. Emerging evidence suggests that the engagement of cellular pathways with more direct effects on fibrogenic mediator generation and collagen synthesis appears to underpin the actions of neutrophil elastase in promoting lung matrix accumulation. Human neutrophil elastase is also present within atherosclerotic plaques where it contributes to matrix degradation and weakening of the

vessel wall associated with the complications of aneurysm formation and plaque rupture. It is joined by other extracellular proteases in these actions but the broad range of substrates and potency of this enzyme coupled with activity associated with neutrophil degranulation single this disruptive protease out as therapeutic target in atherosclerotic disease.

[0566] Methods:

[0567] This assay uses the ENZCHEK® Elastase Assay Kit (Invitrogen Catalog # E-12056). Neutrophils are prepared from fresh human blood using a 6% dextran solution and red blood cells are lysed before plating cells in RPMI media (media should be un-supplemented with no serum, no antibiotics). A 1.0 mg/mL stock solution of the DQ elastin substrate is prepared by adding 1.0 mL of deionized water (dH₂O) directly to one of the three vials containing the lyophilized substrate and mixing to dissolve. 1× Reaction Buffer is prepared by diluting 6 mL of the 10× Reaction Buffer in 54 mL dH₂O. A 100 µg/mL working solution of the DQ elastin substrate is prepared by diluting the DQ elastin stock solution tenfold in 1× Reaction Buffer. Porcine pancreatic elastase stock solution is prepared by making a 100 U/mL stock solution in dH₂O. To assay for elastase activity, 50 µL of 1× Reaction Buffer is pipette into each assay well containing 500,000 neutrophils/mL in a 30 µL volume. 8 µL of each AARS polypeptide is added per well, and the sample incubated for 20 minutes at 37° C. 50 µL of 100 µg/mL DQ elastin working solution is added to each well and mixed. Samples are incubated at room temperature, protected from light, for 30 minutes. Fluorescence intensity in a fluorescence microplate reader equipped with standard fluorescein filters (ex 485/Em 535) fluorescence may be measured over multiple time points.

Binding to Toll-Like Receptors and Activation of NFκB (Assays D1-D4 in the Data Tables Below)

[0568] Background and Therapeutic Relevance:

[0569] Macrophages are major players in the innate immune system and express a large repertoire of different classes of pattern recognition receptors (PRRs), including the family of Toll-like receptors (TLRs) which are powerful regulators and controllers of the immune response.

[0570] Stimulation of TLRs by microbial pathogens and endogenous ligands initiates signaling cascades that induce the secretion of pro-inflammatory cytokines and effector cytokines that direct downstream adaptive immune responses. Endogenous ligands, as well as microbial components, are recognized by and can activate TLRs, raising the possibility that these receptors may be critical targets for the development of new therapies for multiple diseases.

[0571] Accordingly AARS polypeptides that modulate TLR receptor activity, have therapeutic utility in a broad range of diseases and disorders including for example, inflammatory diseases and disorders, autoimmune diseases, tissue transplantation/organ rejection, cancer prevention or treatment, the modulation of haematopoiesis and infection.

[0572] Measurement of TLR Activation in RAW-BLUE Cells

[0573] Mouse macrophages sold under the trademark RAW-BLUE™ cells (Invivogen, Catalog code: raw-sp) express all TLRs except TLR5 and include a secreted embryonic alkaline phosphatase (SEAP) gene which is inducible by NF-κB and AP-1 transcription factors. Upon TLR stimulation, RAW-BLUE™ cells activate NF-κB and/

or AP-1 leading to the secretion of SEAP which is measurable when using SEAP detection medium.

[0574] Methods:

[0575] RAW-BLUE™ cells are washed twice with PBS, trypsinized and resuspended in fresh Growth Medium (Growth Medium: DMEM, 4.5 g/l glucose, 10% heat-inactivated fetal bovine serum (30 minutes at 56° C.), 100 mg/mL ZEOCIN™, 2 mM L-glutamine). Cells are plated at a concentration of 50,000 cells/well in a 96 well plate in a total volume of 100 μ L, and AARS polypeptides, controls, or AARS polypeptides (+LPS) are added to each well at the concentrations shown in the experiments outlined below. Cells are incubated at 37° C. in a 5% CO₂ incubator for 18 hours. On experimental day 2, SEAP detection medium (QUANTI-BLUE™) (Invivogen Catalog code: rep-qbl) is prepared following the instructions and 120 μ L is added per well to a clear flat-bottom 96-well plate, and cell supernatant is added (20 μ L). Samples are incubated at 37° C. for about 30 minutes to up to 2 hours. SEAP levels are determined using a spectrophotometer and reading absorbance at 650 nM.

[0576] To detect AARS polypeptides that specifically block TLR activation this assay can be modified to identify potential TLR antagonists. In this case AARS polypeptides are added to the cells at a final concentration of about 250 nM per well, (or as otherwise specified in the Examples below) 1 hour prior to adding 50 ng/mL LPS. Cells are incubated and SEAP detected as described above. PBS control wells with no LPS or AARS polypeptide alone added are used to find the basal level of TLR stimulation at the time of the measurement. Control wells are pretreated with PBS and known TLR agonists and antagonists. The ratio of the background subtracted [PBS plus LPS signal] to [AARS polypeptide plus LPS signal] is used to determine percent antagonism.

[0577] Human TLR Screening in Hek293 Cells

[0578] Human HEK293 cells are genetically modified and sold under the trademark HEK-Blue™ TLR cells (Invivogen). The TLR2 and TLR4 versions of this cell type selectively express all TLR2 or TLR4 and include a secreted embryonic alkaline phosphatase (SEAP) reporter gene under the control of an IFN-beta minimal promoter which is fused to five NF-kB and AP-1 transcription factors binding sites. With the use of specific TLR 2 or 4 agonists (respectively), HEK-BLUE™ TLR2 and HEK-BLUE™ TLR4 cells activate NF-kB and/or AP-1 leading to the secretion of SEAP which is measurable when using SEAP detection reagent. The HEK-BLUE™ TLR2 cells are co-transfected with the LPS co-receptor protein CD14 to enhance TLR2 responsiveness and improve signal quality. The parent cell expresses endogenous levels of TLR1, 3, 5, 6 and also NOD1.

[0579] Methods:

[0580] HEK-BLUE™-TLR2 or HEK-BLUE™-TLR4 cells are washed twice with PBS, trypsinized and resuspended in fresh Growth Medium (Growth Medium: DMEM, 4.5 g/L glucose, 10% heat-inactivated fetal bovine serum (30 minutes at 56° C.), 100 mg/mL ZEOCIN™, 2 mM L-glutamine). Cells are plated at a concentration of 50,000 cells/well in a 96 well plate in a total volume of 100 μ L, and AARS polypeptides, controls, or AARS polypeptides (+LPS) are added to each well at the concentrations shown in the experiments outlined below. Cells are incubated at 37° C. in a 5% CO₂ incubator for 18 hours. On experimental day

2, SEAP detection medium (QUANTI-BLUE™) (Invivogen Catalog code: rep-qbl) is prepared following the instructions and 120 μ L is added per well to a clear flat-bottom 96-well plate, and cell supernatant is added (20 μ L). Samples are incubated at 37° C. for about 30 minutes to up to 2 hours. SEAP levels are determined using a spectrophotometer and reading absorbance at 650 nM. Control wells are pretreated with PBS and known TLR agonists such as UltraPure LPS (TLR-4) or PAM3CSK4 (TLR-2). The ratio of the background subtracted [PBS plus LPS signal] to [AARS polypeptide plus LPS signal] is used to determine percent agonism.

Cytokine Release (Assays E1-E16 in the Data Tables Below)

[0581] Background and Therapeutic Relevance:

[0582] Cytokines are a diverse set of small cell signaling protein molecules that are used extensively for intercellular communication, and play significant roles in normal body homeostasis, including immunomodulation and regulation. Accordingly AARS polypeptides that modulate the release, or biological activities of cytokines, have therapeutic utility in a broad range of diseases and disorders including for example, inflammatory diseases and disorders, autoimmune diseases, tissue transplantation/organ rejection, cancer prevention or treatment, the modulation of haematopoiesis and infection.

[0583] Cytokine Release from Cells in Culture

[0584] Methods:

[0585] Test cells are seeded into a 24-well plate at density of about 1 million cells/well in 1 mL of growth media. Cells are treated with either AARS polypeptide (at the concentrations shown in the examples below) or an equal volume of PBS and incubated overnight at 37° with 5% CO₂. Following cell treatment, samples are centrifuged at 4° C. in a swinging bucket centrifuge at 2,000 $\times$ g for 5 minutes. Media is carefully removed so as to not disturb the cell pellet and transferred to a new tube. Samples are assayed immediately or snap frozen in liquid nitrogen for subsequent analysis. Cytokine release (including the cytokines MIF, IL-8, IL-10, Serpin E1, GM-CSF, GRO, IL-1 alpha, IL-1 beta, IL-1ra, IL-6, MCP-1, MIP-1, RANTES and TNF-alpha) is determined using commercially available kits (R&D Systems, Inc, MN, USA) or via a contract research organization (MD Biosciences (St. Paul, Minn.).

[0586] Cytokine Release from Human Whole Blood

[0587] Methods:

[0588] Human whole blood is obtained from normal human donors and collected with heparin in standard collection tubes. Blood is used on the same day as it is collected to ensure adequate cell health. Blood is mixed gently and plated in an 100 μ L volume into 96 well polycarbonate V bottom plates. AARS polypeptides are added and slowly mixed into blood 2 $\times$ using a multichannel pipet set on 50 μ L. Filter tips are used for all experimentation and full PPE is worn. All experimentation occurs in a dedicated biosafety hood that is suitable for experimentation with human blood. Blood is incubated overnight at 37° C. with 5% CO₂. Following cell treatment, samples are centrifuged in a swinging bucket centrifuge at 2,000 $\times$ g for 5 minutes. Supernatant is collected for cytokine ELISAs ELISA are performed as described previously.

[0589] Cytokine Release from PBMCs

[0590] Methods:

[0591] To isolate peripheral blood mononuclear cells freshly isolated human whole blood is gently layered over Sigma HISTOPAQUE®-1077 at a ratio of 1:1 in 50 mL conical tubes at room temperature. Layered samples are centrifuged at 400×g in a swinging bucket clinical centrifuge for 30 minutes at room temperature with no brake. The white cellular layer at the interface between the plasma and density gradient is then removed by pipet. These peripheral blood mononuclear cells are washed twice with RPMI-1640 (Invitrogen #22400-105) by dilution and centrifugation for 10 minutes at 250×g. The washed PBMC were resuspended in RPMI-1640+10% FBS and plated at 1×10^6 cells/mL.

[0592] Cytokine Release from Human Synoviocytes

[0593] Background and Therapeutic Relevance:

[0594] A large number of studies have demonstrated that IL-6 and IL-8 are overproduced in several diseases, and thus may play a fundamental role in the pathogenesis of inflammatory disease. IL-6 activates endothelial cell production, leading to the release of IL-8 and monocyte chemoattractant protein, expression of adhesion molecules, and recruitment of leukocytes to inflammatory sites. These cytokines are expressed in cell types associated with inflammatory disease, including cells involved in the pathogenesis of systemic juvenile arthritis, systemic lupus erythematosus, Crohn's disease, and rheumatoid arthritis. One of the most important systemic actions of cytokine production is the induction of the acute phase response. Acute phase proteins are produced primarily by the liver and include proteins that promote the immune response through activation of complement, induction of proinflammatory cytokines, and stimulation of neutrophil chemotaxis. Alternatively, the acute phase response can be helpful, and acute-phase proteins, such as proteinase antagonists, opsonins, and procoagulants, help limit tissue destruction by resolving inflammation. In particular, IL-6 can stimulate synoviocyte proliferation and osteoclast activation, leading to synovial pannus formation and repair. IL-6 acts with IL-1 to increase production of matrix metalloproteinases, which may contribute to joint and cartilage destruction. However, IL-6 may also have protective effects in the joint, as suggested by the finding that this cytokine induces the expression of the tissue inhibitor of metalloproteinase and stimulates proteoglycan synthesis when injected into the joints of mice with antigen-induced arthritis. Human Fibroblast-Like Synoviocytes-Rheumatoid Arthritis (HFLS-RA) are isolated from synovial tissues obtained from patients with Rheumatoid Arthritis (RA). They are cryopreserved at second passage and can be cultured and propagated at least 5 population doublings. HFLS are long known for their role in joint destruction by producing cytokines and metalloproteinases that contribute to cartilage degradation.

[0595] Accordingly AARS polypeptides with the ability to modulate the growth, differentiation, or cytokine release profile of fibroblast-like synoviocytes-rheumatoid arthritis (HFLS-RA) have therapeutic utility in a broad range of diseases including for example, the treatment of inflammatory diseases and disorders including systemic juvenile arthritis, systemic lupus erythematosus, Crohn's disease, and rheumatoid arthritis.

[0596] Methods:

[0597] HFLS-RA, adult cells (Cell Applications Cat #408RA-05a) are maintained in Synoviocyte Growth

Medium (Cell Applications Cat #415-50) in 15 mL medium in 125 mL flasks for 1 passage before use. Cells are maintained at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. An 80 µL volume of cells is plated overnight in growth medium at a cell density of about 50,000 cells/mL. AARS polypeptides at a final concentration of 250 nM per well (or as otherwise indicated in the examples below) are added in sterile PBS to each well following overnight adherence. Control wells contain untreated cells and are incubated with an equivalent volume of PBS. Cells are exposed to proteins or PBS in basal media (Cell Applications Cat #310-470) for 24 hours. Supernatant is removed and IL-8, IL-6 and TNFα ELISA assays are run according to manufacturer's instructions (RND Systems, Cat # DY206 and DY-208, DY-210 Duo-set kits). Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Plates are read on a fluorescent plate reader and viability/proliferation is expressed as a function of resorufin associated fluorescence of AARS polypeptide treated wells divided by resorufin associated fluorescence of PBS only treated wells.

[0598] Human Astrocyte Proliferation and Inflammatory Cytokine Production

[0599] Background and Therapeutic Relevance:

[0600] Human astrocytes (HA) are derived from human cerebral cortex. They are cryopreserved at second passage and can be cultured and propagated 10 population doublings. HA are the most abundant cells in the central nervous system and they perform many functions such as provision of mechanical support and nutrients to neurons, and removal of wastes from neurons. In addition to playing a critical support role for optimal neuronal functioning, they also provide biochemical support of endothelial cells which form the blood-brain barrier. Recent studies have shown that astrocytes are capable of regulating neurogenesis by instructing the stem cells to adopt a neuronal fate and controlling the function of single synapses, participate actively in the transfer and storage of information in the brain. Recognition of the importance of astrocytes in nervous system functioning is increasing, HA can serve as useful in vitro model for exploring the diversity of astrocytes functions. Astrocytes have been shown to proliferate in response to IL6 and TNFα. In addition, these cells are capable of making their own IL6 and TNFα. Thus AARS polypeptides which modulate the proliferation and cytokine production in HA have therapeutic utility in a variety of neurological diseases including neuro-inflammation, neurodegeneration, tumorigenesis of the brain, and brain ischemia and repair.

[0601] Methods:

[0602] Human Astrocytes (HA) from Cell Applications (Cat #882K-05f) are maintained in Cell Applications HA Cell Growth Medium (Cat #821-500) according to manufacturer's instructions. Cells are maintained at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. An 80 µL volume of cells is plated on collagen coated plates overnight in complete medium (above) at a cell density of 50,000 cells/mL. Cells

are washed once with PBS and 80 μ L of serum free growth media is added to each well. AARS polypeptides at a final concentration of 250 nM per well (or as otherwise described in the examples below) are added in a consistent volume in sterile PBS to each well. Cells are exposed to AARS polypeptides for 48 hours and spent media is removed for cytokine assessment (as described previously). Cells are exposed to proteins or PBS in basal media (Cell Applications Cat #310-470) for 48 hours. Supernatant is removed and IL-8 and IL-6 ELISA assays are run according to manufacturer's instructions (RND Systems, Cat # DY206 and DY-208, DY-210 Duo-set kits). Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Plates are read on a fluorescent plate reader and viability/proliferation is expressed as a function of resorufin associated fluorescence of AARS polypeptide treated wells divided by resorufin associated fluorescence of PBS only treated wells.

[0603] Human Lung Microvascular Endothelial Cell (HLMVEC) Proliferation and Inflammatory Cytokine Production

[0604] Background and Therapeutic Relevance:

[0605] The pulmonary vasculature is of great physiological/pathological significance. It is now recognized to be a tissue composed of metabolically active, functionally responsive cells, that interact with circulating substrates and formed elements in ways that regulate the composition of systemic arterial blood, affect target organ functions, and contribute to thrombosis, hemostasis and immune reactions, as well as tumor metastasis. Human lung microvascular endothelial cells (HLMVEC) exhibit elevated expression of chemoattractant cytokines and cell adhesion molecules that provide critical cues for directed migration of leucocytes into the lung during acute lung injury. This primary cell type can be useful tool for studying various aspects of pathology and biology of the pulmonary microvasculature in vitro. Alteration in the structure and function of the microvasculature in response to inflammatory stimuli is believed to be a key factor in organ damage and under appropriate conditions, may provide a stimulus for repair. A significant cause of these vascular alterations is the induction of an inflammatory reaction involving leukocyte infiltration. A variety of studies focused on granulocyte adhesion to the endothelium have revealed that leukocyte recruitment and emigration involves a well-orchestrated adhesion cascade. The adhesion cascade begins when the granulocyte attaches to the endothelium and begins to roll in the direction of fluid flow at a low velocity. As the granulocyte rolls, it becomes activated, subsequently firmly adheres to the endothelium, and migrates across the endothelium into the extravascular space. These adhesion events are mediated, in part, by molecular interactions that occur between CAMs on the surface of the granulocytes and cognate glycoproteins present on the endothelium. A variety of studies have revealed that the endothelial cell adhesion molecule E-selectin can interact with SLe^x-type glycan presenting granulocyte ligands to mediate the attachment and rolling steps of the adhesion cascade. The downstream steps of the cascade involve the interaction of endothelial-expressed intercellular adhesion molecule with granulocyte-expressed CD18 integrins.

[0606] Thus AARS polypeptides which modulate proliferation and/or cytokine production of human lung micro-

vascular endothelial cells have therapeutic utility in a variety of vascular and pulmonary diseases including inflammatory and obstructive lung diseases including for example, pulmonary hypertension, chronic obstructive pulmonary disease, idiopathic pulmonary fibrosis, and asthma.

[0607] Methods:

[0608] HLMVEC (Cell Applications, Catalog #540-05) are maintained in Cell Applications Microvascular Endothelial Cell Growth Medium (Cat #111-500). For appropriate growth, an Attachment Factor Solution containing collagen (Cell Applications, Catalog #123-100), is used to coat plates and flasks before plating cells. Cells are maintained at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. A 80 μ L volume of cells is plated on collagen coated plates overnight in complete medium (above) at a cell density of 50,000 cells/mL. Cells are washed once with PBS and 80 μ L of serum free growth media is added to each well. AARS polypeptides at a final concentration of 250 nM per well (or as otherwise described in the examples below) are added in a consistent volume in sterile PBS to each well. Cells are exposed to AARS polypeptides for 48 hours and spent media is removed for ELISA for cell adhesion molecules and cytokine assessment (as described previously). Cell adhesion molecules including soluble VCAM and/or ICAM are measured using a standard ELISA kit from RND Systems (Cat # DY643 and DY720 respectively). Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Plates are read on a fluorescent plate reader and viability/proliferation is expressed as a function of resorufin associated fluorescence of AARS polypeptide treated wells divided by resorufin associated fluorescence of PBS only treated wells.

Cell Adhesion ((Assays F1-F7 in the Data Tables Below)

[0609] Background and Therapeutic Relevance:

[0610] Cell Adhesion Molecules (CAMs) are proteins located on the cell surface which are involved with the binding with other cells or with the extracellular matrix (ECM) in the process called cell adhesion. These proteins are typically transmembrane receptors and are composed of three domains: an intracellular domain that interacts with the cytoskeleton, a transmembrane domain, and an extracellular domain that interacts either with other CAMs of the same kind (homophilic binding) or with other CAMs or the extracellular matrix (heterophilic binding). Most of the CAMs belong to four protein families: Ig (immunoglobulin) superfamily (IgSF CAMs), the integrins, the cadherins, and the selectins. The immunoglobulin superfamily (IgSF) cell adhesion molecules are calcium-independent transmembrane glycoproteins, including: neural cell adhesion molecules (NCAMs), intercellular cell adhesion molecules (ICAMs), vascular cell adhesion molecule (VCAM), platelet-endothelial cell adhesion molecule (PECAM-1), endothelial cell-selective adhesion molecule (ESAM), junctional adhesion molecule (JAMs), nectins, and other cell adhesion molecules.

[0611] Cell adhesion molecules are cell surface glycoproteins that are critical for leukocyte adhesion to the sinusoidal endothelium and transmigration and cytotoxicity in a variety of inflammatory liver diseases. ICAM-1 plays an important

role in inflammation, and the increased expression of ICAM-1 on endothelial cells is reflected in the activation of endothelial cells. ICAM-1 is of particular importance since it mediates firm endothelial adhesion and facilitates leukocyte transmigration. Studies have shown that there is an upregulation of ICAM-1 on both sinusoidal cells and hepatocytes in inflammatory liver conditions such as hepatitis B viral infection, autoimmune liver disorders, alcoholic hepatitis, and liver allograft rejection.

[0612] Thus AARS polypeptides which modulate cell adhesion molecule production and cell adhesion to endothelial cells have therapeutic utility in a variety of inflammatory diseases including for example, cardiovascular diseases, atherosclerosis, autoimmunity and pulmonary hypertension.

[0613] Methods:

[0614] Human umbilical vein cells (ATCC, Cat # CRL-2873) (HUVEC) are seeded at a concentration of about 1.2×10^5 cells/well in 12 well plates coated with human fibronectin attachment solution in the suggested ATCC media and supplements and grown according to manufacturer's instructions. Cells are stimulated with AARS polypeptides at the indicated concentrations, or PBS alone, and incubated overnight in growth media. Human acute monocytic leukemia (THP-1 (TIB-202)), cells are resuspended into 0.1% BSA/RPMI serum free medium with calcein AM (6 μ L/mL; Invitrogen Cat # C1430) and incubated for 30 minutes. Labeled cells are collected and resuspended in RPMI medium containing 10% FBS, and the density adjusted to 2×10^6 cells/mL.

[0615] 100 μ L (2×10^5) labeled THP-1 cells are placed into each well of the HUVEC monolayer in 1 mL of growth media and incubated for 15 minutes. The wells are washed twice with PBS to remove unbound cells, and then the cells are read by fluorescent plate reader with an Excitation wavelength of 488 nm and an Emission wavelength of 530 nm.

Cellular Differentiation (Assays G1-G4 in the Data Tables Below)

[0616] Adipocyte Differentiation and Proliferation in Primary Human Pre-Adipocyte Cells.

[0617] Background and therapeutic relevance: Both obesity and lipodystrophy are commonly associated with pathologies including diabetes and cardiovascular diseases. It is now recognized that adipose tissue is an endocrine organ that secretes a wide variety of factors, and dysregulated secretion affects adipogenesis as well as whole-body glucose/insulin homeostasis. Excess adipose tissue leading to obesity has become a severe public health threat. Adipose tissue development can be affected by genetic background, hormonal balance, diet, and physical activity. Adipose tissue mass can increase when fat cells are increased in size due to higher triacylglycerol accumulation. In addition, an increase in fat cell number, arising from differentiation of precursor cells into adipocytes, can also occur even in adults as observed in severe human obesity and in rodents fed a high-carbohydrate or high-fat diet. Adipocytes specifically are thought to arise from mesenchymal cells that undergo the commitment and differentiation process, adipogenesis. Pre-adipocyte cell lines can undergo adipocyte differentiation upon treatment with adipogenic agents comprised of synthetic glucocorticoid, dexamethasone (DEX), isobutylmethylxanthine (IBMX), and insulin, have been valuable in these studies. Peroxisome proliferator-activated receptor γ

(PPAR γ) and CCAAT enhancer-binding protein (C/EBP) family of transcription factors have been firmly established to play critical roles in adipocyte differentiation. Early during adipocyte differentiation, C/EBP β and C/EBP δ are induced by DEX and IBMX, respectively, which together then induce PPAR γ and C/EBP α to activate various adipocyte markers that are required for adipocyte function. Other transcription factors have also been reported to either positively or negatively regulate adipogenesis and various growth factors and hormones can affect adipocyte differentiation by regulating expression of adipogenic transcription factors. In fact, in addition to being the main site for energy storage in mammals by storing triacylglycerol and releasing fatty acids in times of need, adipose tissue secretes a wide array of molecules that are involved in diverse physiological processes including immune response, vascular function, and energy homeostasis. Cytokines such as TNF- α and IL-6 are secreted from adipocytes. Some of these factors may also affect growth and development of adipose tissue by auto-crine/paracrine action.

[0618] Thus AARS polypeptides which have the ability to modulate the differentiation and/or proliferation of normal human pre-adipocytes have therapeutic utility in a broad range of diseases including for example, the treatment and prevention of metabolic disease, cardiovascular diseases, obesity and lipodystrophies, as well as the long term complications of diabetes.

[0619] Methods:

[0620] HPAd (human pre-adipocytes) (Cell Application Cat #803sD) are maintained according to vendor instructions. For culturing, cells are thawed quickly, and transferred immediately into 15 mL of Adipocyte Growth Medium (Cell Application Cat #811M-250) and plated into a standard sterile tissue culture treated flask. Media is replaced with fresh Adipocyte Growth Medium every other day until cell is >60% confluent. Cells are grown at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. Cells are plated in clear bottom black walled 96 well tissue culture treated assay plates for differentiation at a concentration of about 50,000 cells/mL. AARS polypeptides at a final concentration of 250 nM per well (or as otherwise indicated in the Examples below) are added to each assay well. All cells are maintained in growth media for 2 days with the exception of the positive controls which are stimulated with adipogenic differentiation media (Cell Applications Cat #811D-250). Cells are exposed to AARS polypeptides for 48 hours. Cell adhesion molecules including soluble VCAM and/or ICAM are measured using a standard ELISA kit from RND Systems (Cat # DY643 and DY720 respectively). Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Plates are read on a fluorescent plate reader and viability/proliferation is expressed as a function of resorufin associated fluorescence of AARS polypeptide treated wells divided by resorufin associated fluorescence of PBS only treated wells. Fresh media is added and differentiation is maintained for 16 days post initial media exchange, with fresh media exchanged every other day to maintain cell health. On day 15, cells are placed in serum free media. On day 16, differentiation to mature adipocytes is assessed with Nile Red (Invitrogen,

concentration of 3 μ M final) staining and quantified with a fluorescent plate reader with the appropriate wavelengths. To perform this assay cells are fixed with 10% paraformaldehyde, washed in PBS and permeabilized in PBS containing 0.5% BSA and 0.1% Triton X-100. Cell proliferation is assessed with an intensity measurement on a fluorescent reader with Hoechst dye 33432 at a concentration of 1 μ g/mL final, as described previously. Adipogenesis is expressed as intensity of Nile Red signal. Hoechst dye signal is used to assess cellular number.

[0621] Human Skeletal Muscle Cell Differentiation and Proliferation.

[0622] Background and Therapeutic Relevance:

[0623] The development of skeletal muscle is a multistep process that involves the determination of pluripotential mesodermal cells to give rise to myoblasts, withdrawal of the myoblasts from the cell cycle and differentiation into muscle cells, and finally growth and maturation of skeletal muscle fibers. Skeletal muscle differentiation involves myoblast alignment, elongation, and fusion into multinucleate myotubes, together with the induction of regulatory and structural muscle-specific genes. At the molecular level, myogenic commitment and muscle-specific gene expression involve the skeletal muscle-specific helix-loop-helix (bHLH) MyoD family of proteins, which includes MyoD, myogenin, myf-5, and MRF4, and the myocyte enhancer-binding factor 2 (MEF2). The DNA binding activity of MyoD family proteins is attenuated by Id, which forms complexes with E2a gene products in proliferating cells and is down-regulated when they are induced to differentiate. The decision to differentiate into myotubes is influenced negatively by several factors. Treatment of myoblasts with fetal bovine serum, basic fibroblast growth factor 2, or transforming growth factor (31 is known to inhibit differentiation of myoblasts. Myogenesis is also regulated negatively by oncogenes such as c-myc, c-jun, c-fos, H-ras, and E1a. There is very little information regarding the signaling that is triggered in the myoblast upon serum withdrawal which leads to the induction of the MyoD family gene expression and to muscle differentiation. Myogenic differentiation appears to depend on the activation of integrins present on the plasma membrane of myoblasts suggesting the operation of an "outside-in" biochemical pathway in which integrin is the upstream molecular species. Interactions of insulin-like growth factor (IGF)-I and -II with their receptors are also positive regulators of skeletal muscle differentiation.

[0624] Accordingly AARS polypeptides with the ability to modulate muscle development have therapeutic utility in a broad range of diseases including for example, the treatment of metabolic disease, cachexia, various muscle wasting conditions, as well as musculoskeletal disease where muscle atrophy plays a key role in the pathogenesis and symptomatology. Human Skeletal Muscle Cells (HskMC) can undergo differentiation to exhibit actin and myosin myofilaments. HskMC have been used in the study of genetic muscular diseases such as Malignant Hyperthermia. HskMC also have the potential to act as a cardiac graft, mending damage to the heart, and thus AARS polypeptides with the ability to modulate muscle development also have utility as in vitro and in vivo regulators of myogenesis.

[0625] Methods:

[0626] To assess the potential role of AARS polypeptides in this process, a standard assay of skeletal muscle cell

differentiation was employed. For this assay, Human Adult Skeletal Muscle Cells (HskMC, Cell Application Cat #150-050) are isolated from healthy human donors from limbal skeletal muscle. Cells are maintained in HskMC Growth Medium (Cell Applications, Cat #151-500). These cells can be cultured and propagated for at least 15 population doublings. For differentiation, cells are maintained in growth media for one passage and then plated at 50,000 cells per mL media in to 96 well clear bottom black walled TC treated plates treated with collagen at 100 μ L per well. Cells are allowed to adhere overnight. AARS polypeptides in PBS, or PBS alone, is added to each well at a final concentration of 250 nM protein (or as otherwise indicated in the examples below). Control wells received the same volume of Differentiation Media (Cell Applications Cat #151D-250) at this time. Cells are incubated with protein or differentiation media for 48 hours. At 48 hours, cell culture supernatant is collected from all wells and differentiation media is added at a volume of 150 μ L to the entire plate with the exception of control wells which are maintained in growth media only. Supernatant is utilized to assess cytokine production including IL6 and IL8 as described previously. Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Cells are monitored under the microscope and media is exchanged for fresh Differentiation media every 2 days. On Day 10, media is removed and cells are fixed with 10% paraformaldehyde for 30 minutes. Cells are permeabilized with 0.1% Triton X-100 in PBS for 15 minutes and cells are stained with TR-Labeled phalloidin and Hoechst 33432 (as described previously) to define actin and nuclei respectively. Nuclear intensity is used to determine cell proliferation in each well and phalloidin intensity is used to determine total actin content. Cells are also stained with alpha actin skeletal muscle antibody (GenTex Cat # GTX101362). Digital photos using a fluorescent microscope as well as visual inspections and scoring are made of all wells.

[0627] Human Bone Marrow Mesenchymal Stem Differentiation and Proliferation.

[0628] Background and Therapeutic Relevance:

[0629] Mesenchymal stem cells (MSCs) are multipotent stem cells that can differentiate into a variety of cell types, including osteoblasts, chondrocytes, myocytes, adipocytes, beta-pancreatic islets cells, and potentially, neuronal cells. Many different events contribute to the commitment of the MSC to other lineages including the coordination of a complex network of transcription factors, cofactors and signaling intermediates from numerous pathways. MSCs are of intense therapeutic interest because they represent a population of cells with the potential treat a wide range of acute and degenerative diseases.

[0630] Moreover AARS polypeptides with the ability to modulate the differentiation of MSCs into different developmental pathways have significant therapeutic utility to enable the in vitro or in vivo modulation of hematopoiesis, neurogenesis, myogenesis, osteogenesis, and adipogenesis, as well as in a broad range of disorders and diseases, including for example inflammatory responses, autoimmunity, cancer, neuronal degeneration, muscular dystrophy, osteoporosis, and lipodystrophy. Human MSCs are immunoprivileged, and represent an advantageous cell type for allogeneic transplantation, reducing the risks of rejection and complications of transplantation. Recently, there have also

been significant advances in the use of autologous mesenchymal stem cells to regenerate human tissues, including cartilage and meniscus, tendons, and bone fractures. Many studies have also investigated the use of MSCs for gene therapy, including transplantation of MSCs transfected with vascular endothelial growth factor for the improvement of heart function after MI in rats, MSCs as vehicles for interferon- β delivery into tumors in mice and gene therapy with MSCs expressing BMPs to promote bone formation. Accordingly due to the intense interest as MSCs as direct and modified therapeutics, as well as the potential of AARS polypeptides to act as therapeutic agents to regulate the differentiation of MSCs in vivo, AARS polypeptides were tested as potential inducers of MSC proliferation and differentiation.

[0631] Methods:

[0632] hMSC (human marrow stromal cells) (Cell Application Cat #492-050) are maintained according to vendor instructions. For culturing, cells are thawed quickly, and transferred immediately into 15 mL of Marrow Stromal cell Growth Medium (Cell Application Cat #419-500) and plated into a standard sterile tissue culture treated flask. Media is replaced with fresh Marrow Stromal cell Growth Medium every other day until cells are >60% confluent. Cells are grown at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. Cells are plated in clear bottom black walled 96 well tissue culture treated assay plates for differentiation at a concentration of 50,000 cells/mL. tRNA synthetase derived proteins at a final concentration of 250 nM per well (or as otherwise specified in the Examples below) are added to each assay well. All cells are maintained in growth media for 2 days with the exception of the positive controls, which was stimulated with osteogenic or chondrogenic differentiation media (StemPro, Invitrogen, Cat # A10072-01 and A10071-01 respectively). Cells are exposed to AARS polypeptides for 48 hours. Soluble VCAM is measured using a standard ELISA kit from RND Systems (Cat # DY643). Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Plates are read on a fluorescent plate reader and viability/proliferation is expressed as a function of resorufin associated fluorescence of AARS polypeptide treated wells divided by resorufin associated fluorescence of PBS only treated wells. Following an assessment of cell viability, resazurin is removed with two media exchanges and 0.5× differentiation media is added to all wells. Differentiation is monitored by visual inspections of all wells for 10 days post media exchange, with fresh media exchanged every other day to maintain cell health. Differentiation was assessed with alkaline phosphatase staining using ELF-97 stain (Invitrogen Cat# E6601) at day 10 post first differentiation exchange. (Yang et al, Nature Protocols (6) 187-213 (2011) doi:10.1038/nprot.2010.189).

[0633] Human Pulmonary Artery Smooth Muscle Cell (hPASC) Proliferation and Differentiation.

[0634] Background and Therapeutic Relevance:

[0635] Pulmonary artery smooth muscle cells (PASMCs) in normal human adult lung blood vessels are mostly quiescent, non-migratory and are largely committed to executing their contractile function in the lung. However, PASMCs

are not terminally differentiated and possess the ability to modulate their phenotype and exit their quiescent state in response to changing local environmental cues. This differentiation state may occur in development, tissue injury, and vessel remodeling in response to changes in tissue demand. Pulmonary hypertension (PH) is associated with a variety of underlying conditions including an increase in peripheral pulmonary vascular resistance as a result of increased vascular tone and PASMC contractility and vascular remodeling. Vascular remodeling involves PASMC growth, synthesis of matrix material, and alterations in cell-cell and cell-matrix interactions in the walls of small pulmonary arteries (PAs), which lead to increased thickness of the smooth muscle component of the vessel wall and abnormal muscularization of the normally nonmuscularized, distal PAs. This process contributes to reduced lumen diameter and increased peripheral resistance. Although the precise role of the PASMCs in the initial cause of the disease is controversial, the changes that occur play a key role in the clinical consequences of the disease. A crucial step in studying cellular differentiation is identifying a set of cell-specific or cell-selective genes that contribute to the differentiated function(s) of the cell. A variety of smooth muscle cell (SMC) genes have been identified that serve as useful markers of the relative state of differentiation or maturation of the vascular SMCs, such as SM alpha-actin, SM MHC, h1-calponin, SM22-alpha, desmin, metavinculin, smothelin and others. The most widely used marker is SM alpha-actin, partially because of the commercial availability of a number of very high-affinity and highly selective antibodies for this protein. Whether changes in PASMCs result from their inherent characteristics or from dysregulation of molecular events that govern PASMC growth remains an open question. However determining the regulatory cues and managing potential dis-regulation provides significant therapeutic insight to managing a variety of vascular and pulmonary diseases including pulmonary hypertension, vascular diseases.

[0636] Thus AARS polypeptides which have the ability to modulate the differentiation and/or proliferation of normal human PASMCs derived from adult humans have therapeutic utility in a variety of vascular and pulmonary diseases including inflammatory and obstructive lung diseases including for example, pulmonary hypertension, chronic obstructive pulmonary disease, idiopathic pulmonary fibrosis, and asthma.

[0637] Methods:

[0638] HPASC (Cell Applications Cat #352-05a) are maintained in HPASC growth media (Cell Applications Cat #352-05a) in 15 mL medium in 125 mL flasks for 1 passage before use. Cells are maintained at 37° C., 5% CO₂, in a humidified environment and utilized in BSL2 certified tissue culture hoods using sterile technique and appropriate personal protective equipment including goggles, gloves and lab coats. An 80 μ L volume of cells is plated on collagen coated overnight in growth medium at a cell density of 50,000 cells/mL. AARS polypeptides were added in sterile PBS to each well at a final concentration of 250 nM (or as otherwise specified in the Examples below). Control wells held only an equivalent volume of PBS. Positive control samples were incubated with vendor supplied HPASC differentiation media (Cell Applications Cat #311D-250). Cells are exposed to AARS polypeptides or PBS in basal media (Cell Applications Cat #310-470) for 48 hours fol-

lowed by a media exchange to differentiation media for the entire plate. Supernatant is collected and utilized to assess cytokine production including IL6 and IL8 as described previously. Proliferation is assessed with Resazurin as described previously by adding fresh media containing Resazurin to plates following supernatant removal and incubating for three hours at 37° C. Cells are monitored for 10 days with a media exchange every other day. Differentiation is assessed after fixation as described above, and permeabilization with 0.1% Triton X-100, by quantifying staining to smooth muscle actin- α staining using an anti-SMA- α antibody (GeneTex Cat #GTX101362) and an Alexa 405 conjugated secondary antibody. Proliferation is assessed with Hoechst staining after cell fixation in 10% formaldehyde for 30 minutes. Hoechst dye is read using a bottom reading fluorescent plate reader with an excitation wavelength (Ex) of 405 nm, and an emission wavelength (Em) of 450 nm. Total actin staining is assessed via the use of an Alexa-488 labeled phalloidin stain (Invitrogen Cat# A12379).

Analysis of the Binding of AARS Polypeptides to Cells (Assays H1-H10 in the Data Tables Below)

[0639] Background and Therapeutic Relevance:

[0640] The binding of AARS polypeptides to specific cell types demonstrates that the cell type in question expresses specific receptors for the AARS polypeptide in question. Depending upon the cell type in question, cell binding implies a potential role for the AARS polypeptide in regulating the activity or behavior of the cell, or similar types of cell, *in vivo*. Specific examples of such regulatory roles include for example, the binding and modulation of B-cells and T-cells (immunomodulation/chemotaxis/autoimmunity/inflammation); HepG2 cells (control of metabolism, cholesterol uptake or metabolism); THP-1, Jurkat, Raji cells (immunomodulation/chemotaxis/autoimmunity/inflammation), platelets (thrombopoiesis), 3T3L1 adipocytes (lipogenesis/metabolism), and C2C12 mouse myoblasts (myogenesis, osteogenesis).

[0641] Binding to Blood Cells

[0642] Methods:

[0643] Blood is collected in EDTA tubes from healthy donors. 2 mL whole blood is placed into 5 mL Falcon FACS tube. 2 mL of staining buffer (PBS+2% FBS) is added, vortexed 3-5 seconds, centrifuged for 5 minutes at 300 $\times$ g. The supernatant aspirated, the wash repeated, and the pellet resuspended in 2 mL of staining buffer.

[0644] 100 μ L of washed blood is transferred to clean 5 mL FACS sample tubes. His6- or V5-His6-tagged AARS polypeptides are added to tubes at the concentrations indicated in the specific experiments outlined below and incubated on ice for 45 minutes. After incubation, antibodies for the different cell type surface markers (BD Pharmingen Cat Nos. 560910, 555398, 555415, 340953, 560361), and FITC labeled anti-V5 tag antibody (V5-FITC, Invitrogen Cat # R96325) or FITC labeled anti-His6 antibody (AbCam Cat #ab1206) are added to tubes, incubated in the dark on ice 30 minutes. After incubation 2 mL of BD FACS Lysing Solution (cat #349202) was added to tubes. Samples are vortexed, and placed on ice for 15 minutes. Samples are washed with 1 $\times$ 2 mL PBS and resuspended in 2 mL of 2% formaldehyde in PBS prior to FACS analysis. AARS polypeptides that bind greater than 25% of a cellular population, where antibody alone has no significant signal, is deemed a hit.

[0645] Platelet Binding Assays:

[0646] 50 μ L of washed blood is transferred to clean 5 mL FACS sample tubes, His6- or V5-His6-tagged AARS polypeptides are added to tubes at the concentrations indicated in the specific experiments outlined below and tubes are placed on ice for 45 minutes. 20 μ L CD61 pan platelet antibody (BD Pharmingen, Cat #555754) and 0.5 μ L anti-V5-FITC labeled antibody (Invitrogen, R96325) or FITC labeled anti-His6 antibody (AbCam Cat #ab1206) are added to each tube. Tubes are placed on ice and protected from light for 30 minutes. Samples are brought up to a total volume in 2 mL of 1% formaldehyde in PBS and analyzed by flow cytometry within 24 hours. AARS polypeptides that bind greater than 25% of a cellular population, where antibody alone has no significant signal, is deemed a hit.

[0647] Binding to Cells in Culture:

[0648] Approximately 1 $\times$ 10⁶ cells in 100 μ L complete RPMI medium are placed into 5 mL FACS tubes. His6- or V5-His6-tagged AARS polypeptides are added to tubes at the concentrations indicated in the specific experiments outlined below and tubes are placed on ice for 45 minutes. Cell samples are washed twice with 1 mL staining buffer (PBS+2% FBS), and then 0.54 of anti-V5-FITC antibody (Invitrogen R96325) or FITC labeled anti-His6 antibody (AbCam Cat #ab1206) in staining buffer with 200 μ g/mL human IgG, is added and the samples incubated on ice, protected from light, for 30 minutes. Samples are washed twice with 1mL staining buffer, and then brought up to a total volume in 2 mL of 1% formaldehyde in PBS and analyzed by flow cytometry within 24 hours. AARS polypeptides that bind greater than 25% of a cellular population, where antibody alone has no significant signal, is deemed a hit.

Animal Studies: Modulation of Haematopoiesis and Circulating Cytokines

[0649] Background and Therapeutic Relevance:

[0650] Hematopoiesis (alternatively haemopoiesis or hemopoiesis) is the formation of blood cellular components. All cellular blood components are derived from haematopoietic stem cells (HSCs) which reside in the medulla of the bone (bone marrow) and have the unique ability to give rise to all of the different mature blood cell types. HSCs are self-renewing: when they proliferate, at least some of their daughter cells remain as HSCs, so the pool of stem cells does not become depleted. The other daughters of HSCs (myeloid and lymphoid progenitor cells), however can each commit to any of the alternative differentiation pathways that lead to the production of one or more specific types of blood cells, but cannot themselves self-renew. A change in the blood components in response to exposure to an AARS polypeptide therefore suggests that the AARS polypeptide is capable of modulating hematopoiesis, and regulating the development of haematopoietic stem cells.

[0651] All blood cells can be divided into three lineages; Erythroid cells, lymphocytes and myelocytes.

[0652] Erythroid cells are the oxygen carrying red blood cells. Both reticulocytes and erythrocytes are functional and are released into the blood. Accordingly a reticulocyte count estimates the rate of erythropoiesis, and a change in red blood cell count suggests that an AARS polypeptide modulates erythropoiesis.

[0653] Lymphocytes are the cornerstone of the adaptive immune system. They are derived from common lymphoid progenitors. The lymphoid lineage is primarily composed of

T-cells and B-cells (types of white blood cells). Accordingly a change in white blood cell count or composition in response to exposure to an AARS polypeptide suggests that that the AARS polypeptide modulates lymphopoiesis.

[0654] Myelocytes, which include granulocytes, megakaryocytes and macrophages, and are derived from common myeloid progenitors, are involved in a variety of roles, including innate immunity, adaptive immunity, and blood clotting. Accordingly a change in myeloid cell count or composition in response to exposure to an AARS polypeptide suggests that the AARS polypeptide modulates myelopoiesis. The same rationale can be used to establish whether the AARS polypeptides modulate granulopoiesis, by measuring changes in granulocyte number in response to exposure to the AARS polypeptides. A role for the AARS polypeptide in modulating megakaryocytopoiesis may be inferred by a change in megakaryocyte or platelet composition or number in the blood.

[0655] Cytokine release in either wild type mice, or in various animal model systems of inflammation, provides an initial assessment of the potential ability of the AARS polypeptides to modulate inflammatory responses. The role of AARS polypeptides in modulating acute chronic inflammatory processes for example, can be readily assessed using a mouse model of diet induced obesity (DIO). The DIO model centers upon placing rodents on a high fat diet for several months leading to increased obesity, insulin resistance and immune system dysfunction. A particular consequence of this immune system dysregulation results in increased production of proinflammatory cytokines in DIO animals leading to a condition of chronic systemic inflammation. There is a growing body of evidence suggesting that low grade inflammation contributes to the development and maintenance of obesity and a diabetic phenotype that is similarly observed in the human condition termed metabolic syndrome. As such, the ability of AARS polypeptides to modulate the immune system and restore homeostatic balance towards a resolution of this chronic inflammatory state would be particularly beneficial in numerous diseases and disorders including but not limited to the treatment and prevention of the symptoms and side effects of metabolic disease, diabetes, cardiovascular diseases, atherosclerosis, obesity, as well as various autoimmune diseases and disorders, including for example, multiple sclerosis, vascular and allergic disorders.

[0656] Methods:

[0657] Male wild type control (C57BL/6) or diet induced obesity mice (C57BL/6NHsd) are purchased from Harlan (Indianapolis, Ind.) and housed individually. DIO mice are fed a high fat diet (Cat. #TD.06414-60% kcal from fat) and control mice are fed a normal diet (Cat. #2018S-18% kcal from fat). DIO mice are placed on the high fat diet starting at 6 weeks of age for a total of 10 weeks. Both DIO and control mice are allowed to feed and drink ad libitum. At 16 weeks of age, mice are sorted and randomized into groups of 5 animals based on weight. On day 2, mice are weighed and tail vein bled (100 μ L) for pre-treatment complete blood count (CBC) analysis. On day 1, mice are weighed and intravenously injected via the tail vein with vehicle (PBS) or individual AARS polypeptides at 10 mg/kg. Four hours post-injection, mice are facial vein bled (150-200 μ L) for subsequent cytokine analysis. On days 2, 3, & 4, mice are intravenously dosed as on day 1. On day 5, mice are weighed, terminated and blood are collected by heart punc-

ture for Complete Blood Count (CBC analysis) (plasma-EDTA) and cytokine examination (serum).

[0658] CBC and Cytokine Analysis:

[0659] Complete blood counts are analyzed from blood draws preceding injections (day -2) and 24 hours after the final injection (day 5). CBC values are assessed for total white blood cell counts and overall red blood cell morphology. White blood cells are further characterized by total and fractional percentage of neutrophils, lymphocytes, monocytes, eosinophils, & basophils. Red blood cell breakdown included measurements of hemoglobin (dL), hematocrit (%), mean corpuscular volume (fL), mean corpuscular hemoglobin, mean corpuscular hemoglobin concentration (%), and total platelet count ($10^3/\mu$ L). CBC analysis is performed by Antech Diagnostics (Fishers, Ind.).

[0660] Circulating cytokine levels are examined at 4 hours post-injection (day 1) and 24 hours after the final injection (day 5). Serum is isolated, snap frozen and sent to Rules Based Medicine (Austin, Tex.) for multi-analyte profiling. Serum samples are analyzed using the RodentMap panel encompassing 59 unique biomarkers including Apo A-1, CD40, CD40-L, CRP, ET-1, eotaxin, EGF, Factor VII, fibrinogen, FGF-9, FGF-basic, GST- α , GCP-2, GM-CSF, KC/GRO α , haptoglobin, IgA, IFN γ , IP-10, IL-1 α , IL-1 β , IL-10, IL-11, IL-12p70, IL-17A, IL-18, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, LIF, lymphotactin, M-CSF-1, MIP-1 α , MIP-1 β , MIP-1 γ , MIP-2, MIP-3 β , MDC, MMP-9, MCP-1, MCP-3, MCP-5, MPO, myoglobin, SAP, SGOT, SCF, RANTES, TPO, tissue factor, TIMP-1, TNF- α , VCAM-1, VEGF-A, and vWF. A change in cytokine levels was counted as a hit if the cytokine increased by at least 2-fold or decreased by at least 50% compared to vehicle controls.

Example 1

Identification of Proteolytic Fragments and Products of Alternative Splicing from AARSs Using Protein Topography and Migration Analysis Platform

[0661] To identify AARS fragments from cell lines, conditioned media and tissues, samples are prepared in the following ways:

[0662] Mouse Macrophage (RAW 264.7), Cytosol and Conditioned Media:

[0663] Cells are treated with serum free DMEM media at a density of 15×10^6 cells/flasks. After 48 hours conditioned media and cell pellets are collected and processed. 200 μ g of protein from secreted and cytosolic proteomic fractions are separated by SDS-PAGE and gel slices are prepared for analysis by mass spectrometry.

[0664] Mouse Pancreas Tissue:

[0665] The pancreas from three mice are chopped, dounce homogenized, and sonicated in PBS with protease inhibitors. Cytosolic proteome is isolated by centrifugation and 200 μ g of protein is separated by SDS-PAGE and gel slices are prepared for analysis by mass spectrometry.

[0666] Mouse Liver Tissue:

[0667] Three mouse livers are chopped, dounced homogenized, and sonicated in PBS with protease inhibitors. Cytosolic proteome is isolated by centrifugation and 200 μ g of protein is separated by SDS-PAGE and gel slices are prepared for analysis by mass spectrometry.

[0668] In-gel digests are analyzed by LTQ XL ion trap mass spectrometer (ThermoFisher) equipped with ultimate

3000 μ LC system (Dionex). The samples are first loaded on PepTrap (michrom) for 10 min with 5% Acetonitrile in 0.1% formic acid using Dionex autosampler. Then the samples are analyzed with a 100 μ m (inner diameter) fused silica capillary column containing 10 cm of C18 resin (michrom). Peptides are eluted from the column into mass spectrometer with a flow rate of 0.45 μ l/min using a linear gradient of 5-33.5% acetonitrile in 0.1% formic acid within 110 min.

[0669] LTQ is operated in data-dependent scanning mode such that one full MS scan is followed by seven MS/MS scans of the seven most abundant ions. Dynamic exclusion is enabled with repeat count equals to 1, repeat duration equals to 20 seconds, exclusion list size is 300 and exclusion duration is 60 seconds.

[0670] After LC-MS/MS analysis, the raw data is searched with BioWorks3.3.1(SEQUEST) using a concatenated target/decoy variant of the mouse IPI database. The SEQUEST data are filtered and sorted with DTASelect. Tables 1, 4 and 7 show sequences identified in this way.

Example 2

Identification of Splice Variants Using Deep Sequencing

[0671] Splice variants of the aminoacyl tRNA synthetase are identified using high throughput sequencing of cDNA libraries enriched for aminoacyl tRNA synthetase transcripts. The cDNA templates are prepared from total RNA extracts of tissues such as human adult and fetal brains and enriched for aminoacyl tRNA synthetase transcripts by using primer sequences specific for all annotated exons of all annotated human aminoacyl tRNA synthetases and their associated proteins.

[0672] Human Total RNAs are obtained from Clontech. For cell line and mouse tissue samples, total RNAs are extracted using RNA Extract II Kit (MN). Genomic DNA is digested in the total RNA samples by DNAase I. To obtain mature messenger RNAs (mRNAs), the RNA samples are enriched twice by binding polyA+ RNA and digestion of RNA without 5'-cap by 5'-phosphate dependent exonuclease. Complementary DNA (cDNA) is synthesized from mature RNAs using primers that anneal to exon sequences of aminoacyl tRNA synthetase genes. A transcriptome enriched for aminoacyl tRNA synthetase genes is amplified by multiplex PCR using the aminoacyl tRNA synthetase-exon specific cDNA and different combinations of aminoacyl tRNA synthetase-exon primers. The double-stranded aminoacyl tRNA synthetase-enriched transcriptome PCR products are enzymatically repaired at both ends before adding A-overhangs to the 3' ends of the repaired fragments. Sequencing adaptors and index sequences are then added to the aminoacyl tRNA synthetase-enriched transcriptome PCR products to generate cDNA libraries for deep sequencing with Illumina's Multiplex Sequencing Kit. In brief, the aminoacyl tRNA synthetase-enriched transcriptome PCR products with 3'-A overhangs are ligated to the InPE adaptor oligonucleotides provided in the kits. Index sequences are added to the PCR products with InPE adaptors. To obtain enough DNA fragments for deep sequencing, the PCR products with index sequences are further amplified by PCR. Aminoacyl tRNA synthetase-enriched cDNA libraries with different indexes are pooled and sequenced using an Illumina DNA sequencing machine to get 50 base pair end reads. Sequencing reads are mapped to human or mouse

genome for identification of alternative splicing events. "Splicemap" software is used to identify splice junctions.

[0673] Deep sequencing of these cDNAs are performed to generate about 1 million sequencing reads of about 50 nucleotides in length. The sequences specific for exons of the aminoacyl tRNA synthetases are queried against annotated exon junctions and new exon junctions are identified as alternative splicing events.

[0674] The columns in Tables 2, 5, and 8 labeled "5' exon" and "3'exon" indicate, when present, which exons are fused together in the cDNA sequence. Tables 2, 5, and 8 show sequences that were identified for alternative splice events, transcripts containing such splice events, and the polypeptides expressed by those transcripts. Alternative splice variants identified by deep sequencing are identified in Tables 2, 5, and 8 as those ones in which there are numbers greater than zero in the columns labeled as "Sequencing reads" in the human adult or fetal brain.

Example 3

Identification of AARS Polypeptides Using Bioinformatics

[0675] AARS protein fragments (resectin or appendacrine peptides) are identified using bioinformatics. Amino acid sequences of the full length human aminoacyl tRNA synthetase are aligned with the full length amino acid sequence of its ortholog from the bacterium *Escherichia coli* using a program such as FASTA or the BLASTP program from the NCBI. Resectin sequences from the human proteins are identified as sequences covering regions where there are gaps in the bacterial sequence in the alignment, or regions with low homology between the two species. The peptide, and corresponding DNA sequences in Tables 3, 6, and 9 include examples identified in this way.

Example 4

Differential Expression of AARS Polypeptides Identified by Mass Spectrometry

[0676] The PROTOmap technique is used as described in Example 1 to compare the differential expression of Arginyl tRNA synthetases in different tissues/cell types (refer to Tables 1, 4, and 7 for sequences and comparisons): Aminoacyl-tRNA synthetase resectin expression is compared between mouse liver tissue and mouse pancreas tissue. Aminoacyl-tRNA synthetase resectin expression is compared between cytosol of RAW264.7 and conditioned media from RAW264.7 cells harvested after 48 hours of serum starvation.

Example 5

Differential Expression of AARS Polypeptides Identified by Deep Sequencing

[0677] To test for differential expression of splice events, the deep sequencing is done for cDNAs prepared from different tissues.

[0678] Expression of specific alternative splice events for aminoacyl tRNA synthetases is unexpected and indicates biological importance. The variation in relative number of reads seen in the deep sequencing of different transcriptome samples indicates that alternative splice events of aminoacyl

tRNA synthetases are differentially regulated and not just artifacts due to sample handling.

Example 6

Antibody Screening

[0679] To facilitate the discovery of antibodies displaying preferential binding to specific aminoacyl tRNA synthetase fragments (e.g., ≥ 10 -fold higher affinity when compared to the parental full length enzyme), a human antibody phage display library is screened by AbD Serotec (a division of MORPHOSYSTM, Martinsried/Planegg, Germany) using affinity enrichment techniques (panning). Antibodies enriched after multiple rounds of screening with the aminoacyl tRNA synthetase fragments are subsequently characterized by ELISA for reactivity to the fragments, and to the parental, full length enzyme. Clones demonstrating preferential binding (e.g., ≥ 10 -fold higher affinity) to the aminoacyl tRNA synthetase fragments are further characterized.

[0680] If the necessary specificity is not achieved at the end of this process, subtraction strategies, such as pre-adsorption steps with the full length enzyme and/or counter-screening, are used to eliminate cross reacting antibodies and drive the selection process towards the unique epitope(s) on the aminoacyl tRNA synthetase fragments.

form the canonical transcripts. These different PCR products are excised and purified from the gel and ligated into a standard cloning vector for DNA sequence analysis. Alternative splicing variants are identified as different sequences from the canonical transcripts. Splice variants identified by this systematic PCR approach are shown in Tables 2, 5 and 8.

Example 8

Codon Optimization of Selected AARS Polynucleotides

[0682] Representative AARS polypeptides (summarized in Table E2) are selected for further biochemical, biophysical and functional characterization based on one or more of the following criteria, i) the identification of AARS polypeptide proteolytic fragments, ii) the identification of AARS polypeptide splice variants, iii) the identification of AARS polypeptides by bioinformatic analysis, iv) evidence of differential expression of specific AARS polypeptides, v) the domain structure of the AARS protein, vi) the size of the AARS polypeptide, and vii) the minimization of similar duplicative sequences.

TABLE E2

Summary of AARS Polypeptides Selected for Codon Optimization and Bacterial Expression					
AARS Polypeptide Name	SEQ. ID Nos. for Epitope Tagged AARS polypeptides	SEQ. ID. Nos. for AARS Polynucleotides	Residues of AARS protein	Location of epitope tag	Cloning/synthesis method used
ArgRS ^{N1}	SEQ. ID. NO. 31	SEQ. ID. NO. 39	1-398	N-terminal	1
ArgRS ^{N1}	SEQ. ID. NO. 32	SEQ. ID. NO. 39	1-398	C-terminal	1
ArgRS ^{N2}	SEQ. ID. NO. 33	SEQ. ID. NO. 40	1-71	N-terminal	1
ArgRS ^{N2}	SEQ. ID. NO. 34	SEQ. ID. NO. 40	1-71	C-terminal	1
ArgRS ^{N3}	SEQ. ID. NO. 35	SEQ. ID. NO. 41	1-353 + 20 aa	N-terminal	1
ArgRS ^{N3}	SEQ. ID. NO. 36	SEQ. ID. NO. 41	1-353 + 20 aa	C-terminal	1
ArgRS ^{N4}	SEQ. ID. NO. 37	SEQ. ID. NO. 42	1-160	N-terminal	1
ArgRS ^{N4}	SEQ. ID. NO. 38	SEQ. ID. NO. 42	1-160	C-terminal	1
ArgRS ^{C1}	SEQ. ID. NO. 139	SEQ. ID. NO. 147	73-660	N-terminal	1
ArgRS ^{C1}	SEQ. ID. NO. 140	SEQ. ID. NO. 147	73-660	C-terminal	1
ArgRS ^{C2}	SEQ. ID. NO. 141	SEQ. ID. NO. 148	551-660	N-terminal	1
ArgRS ^{C2}	SEQ. ID. NO. 142	SEQ. ID. NO. 148	551-660	C-terminal	1
ArgRS ^{C3}	SEQ. ID. NO. 143	SEQ. ID. NO. 149	463-660	N-terminal	1
ArgRS ^{C3}	SEQ. ID. NO. 144	SEQ. ID. NO. 149	463-660	C-terminal	1
ArgRS ^{C4}	SEQ. ID. NO. 145	SEQ. ID. NO. 150	247-660	N-terminal	1
ArgRS ^{C4}	SEQ. ID. NO. 146	SEQ. ID. NO. 150	247-660	C-terminal	1
ArgRS ^{I1}	SEQ. ID. NO. 157	SEQ. ID. NO. 159	168-530	N-terminal	1
ArgRS ^{I1}	SEQ. ID. NO. 158	SEQ. ID. NO. 159	168-530	C-terminal	1

Example 7

Identification of Splice Variants Using Systematic PCR

[0681] cDNA templates for PCR reactions are reverse transcribed from total RNA extracts of tissues or cells (e.g., human brain, IMR-32 and HEK293T). PCR reactions are performed using aminoacyl tRNA synthetase specific primers, pairing a forward primer (FP1) designed to anneal to the 5' untranslated region or exons in the 5' half of the gene with a reverse primer (RP1) designed to anneal to exons in the 3' half of the gene or the 3'UTR. Amplified DNA products are analyzed by agarose gel electrophoresis to identify PCR products that are a different size than the fragment amplified

[0683] Polynucleotides encoding the selected AARS polypeptides listed in Table E2, along with the appropriate N or C-terminal epitope tag, are synthesized and cloned as described in the General Materials and Methods section using the gene synthesis methodology listed in Table E2.

Example 9

Small Scale Bacterial Expression and Purification

[0684] The AARS polypeptides listed in Table E2 are expressed in *E. coli* as described in the General Materials and Methods section. The relative expression of soluble and inclusion body localized AARS polypeptides is summarized in Table E3 below.

TABLE E3

Summary of AARS Polypeptide Bacterial Expression Characteristics			
AARS Polypeptide	Location of Epitope Tag	Amount of Protein Recovered from Soluble Fraction	Amount of Protein Recovered from Inclusion Bodies
ArgRS ^{N1}	N-terminal	+	ND
ArgRS ^{N1}	C-terminal	+	ND
ArgRS ^{N2}	N-terminal	+++++	ND
ArgRS ^{N2}	C-terminal	+++++	ND
ArgRS ^{N3}	N-terminal	+	ND
ArgRS ^{N3}	C-terminal	+	ND
ArgRS ^{N4}	N-terminal	+	ND
ArgRS ^{N4}	C-terminal	+	+
ArgRS ^{C1}	N-terminal	+	ND
ArgRS ^{C1}	C-terminal	+	ND
ArgRS ^{C2}	N-terminal	+	++
ArgRS ^{C2}	C-terminal	+	++
ArgRS ^{C3}	N-terminal	+	ND
ArgRS ^{C3}	C-terminal	+	ND
ArgRS ^{C4}	N-terminal	+	ND
ArgRS ^{C4}	C-terminal	+	ND
ArgRS ^{I1}	N-terminal	+	+++
ArgRS ^{I1}	C-terminal	+	++

“+” represents 0-1 mg/L AARS polypeptide expression

“++” represents 1-5 mg/L AARS polypeptide expression;

“+++” represents 5-10 mg/L AARS polypeptide expression;

“++++” represents 10-15 mg/L AARS polypeptide expression;

“+++++” represents ≥15 mg/L AARS polypeptide expression;

ND: not determined

[0685] Surprisingly, the protein expression data demonstrates the existence of at least one protein domain that exhibits high level expression of soluble protein when expressed in *E. coli*. Specifically the data demonstrates that the AARS polypeptide ArgRS1^{N2}, (amino acids 1-71), defines the boundary of a novel protein domain that is highly expressed in *E. coli*.

Example 10

Large Scale Production of AARS Polypeptides

[0686] Representative AARS polypeptides are prepared in larger amounts to enable further functional and biophysical characterization. The AARS polypeptides listed in Table E4 are expressed in *E. coli* in large scale culture as described in the General Materials and Methods section. The yields, and specified biophysical characteristics, for each expressed soluble protein are summarized below in Table E4.

TABLE E4

Summary of representative AARS Polypeptide yield and biophysical characterization								
AARS Polypeptide	Location of Epitope Tag	Yield [mg/L] ⁽¹⁾	Purity [%]	Endotoxin [EU/mg]	Molecular Weight	Working stock conc [mg/ml]	Stability [percent recovery] ⁽²⁾	Aggregation [DLS]
ArgRS ^{N2}	C-terminal	4.5	95	0.8	ND	4.24	ND	ND
ArgRS ^{I1}	C-terminal	0.7	99	<0.1	ND	0.16	ND	ND
ArgRS ^{C2}	C-terminal	1.1	99	<0.1	ND	8.4	ND	ND

Notes

⁽¹⁾ Yield determined by measuring protein recovery after last purification step

⁽²⁾ Determined as percent recovery of non aggregated material after 1 week at 25° C.

Key:

ND: Not Determined

[0687] The results from these studies establish that representative AARS proteins from the ArgRS1^{N2}, ArgRS1^{I1} and

ArgRS1^{C2} families of AARS proteins can be successfully expressed in *E. coli*, and purified to high purity, with low endotoxin contamination levels.

Example 11

Transcriptional Profiling of Representative AARS Polypeptides

[0688] To test for the ability of the AARS polypeptides to modulate gene expression, selected AARS polypeptides were incubated with mesenchymal stem cells or human skeletal muscle cells for the times and at the concentrations shown in Table E5.

TABLE E5

Transcriptional profiling of representative AARS Polypeptides in Mesenchymal Stem Cells (MSC) or Human Skeletal Muscle Cells (HSkMC)						
Cell type and Exposure Time						
Test Sample Description	Concen- nM	MSC 24 hours	MSC 72 hours	HSkMC 24 hours	HSkMC 72 hours	
AARS Polypeptides	Location of Epitope Tag					
ArgRS ^{N2}	N-terminal	250	4	3	3	2
ArgRS ^{N2}	C-terminal	250	1	1	5	7
ArgRS ^{C2}	N-terminal	250	2	5	5	10
ArgRS ^{C2}	C-terminal	250	0	4	7	3
ArgRS ^{I1}	N-terminal	250	2	5	7	3
ArgRS ^{I1}	C-terminal	250	5	3	10	5
Controls						
Average across all AARS polypeptides screened		3	5	6	7	
Osteogenesis cocktail		17	20	11	16	
Chondrogenesis cocktail		17	19	14	19	
Adipogenesis cocktail		19	15	16	18	
SKMC Pos Ctrl		11	8	5	4	
Untreated		0	0	1	1	

[0689] In Table E5, the numbers in each column represent the number of genes which were modulated, either positively or negatively by at least 4 fold compared to the control genes, as described in the general methods section. The data shows that specific forms of the AARS polypeptides tested have the surprising ability to regulate the transcription, and hence potentially modulate the developmental fate or differentiation status when added to either Mesenchymal Stem

Cells (MSC) and/or Human Skeletal Muscle Cells (HSkMC). Shaded cells with bolded numbers in the table

represent examples where the AARS polypeptide exhibits a significant impact on the regulation of gene transcription in the cell lines and times indicated in the table.

[0690] It is concluded that ArgRS1^{N2}, ArgRS1^{C2}, and ArgRS1^{I1} appear to be major regulators of Mesenchymal Stem Cell and/or human skeletal muscle cell gene expression.

Example 12

Functional Profiling of AARS Polypeptides

[0691] To test for the ability of the AARS polypeptides to modulate a range of phenotypic processes, selected AARS polypeptides were incubated with the cell types, and the conditions provided in the general methods section, and Tables E5 and E6.

TABLE E6

Key to Assays and criteria for indicating a hit	
Proliferation assays	
Source and cell type	Assay Number
Human megakaryocytic leukemia cells/Mo7e	A1
Human acute promyelocytic leukemia cells/HL60	A2
Human lymphoblast (cancer cell line)/RPMI8226	A3
Human mesenchymal stem cells/hMSC	A4
Human astrocytes	A5
Human bone marrow aspirate cells/Bone Marrow Cells	A6
Human bone marrow aspirate cells/Bone Marrow Cells (Long Term Culture)	A7
Human Synoviocyte/HFLS-SynRA	A8
Human pre-adipocyte cells/hPAD	A9
Human pulmonary artery smooth muscle cell/hPASMC	A10
Human skeletal muscle cell/hSKMC	A11
Data analysis for proliferation assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be proliferative if greater than 3 SD away from the PBS value in the positive direction. A tRNA synthetase derived AARS polypeptide was considered to be cytotoxic if greater than 3 SD away from the PBS value in the negative direction. A cytotoxic compound was utilized as a negative control and the average value for this was always greater than 3 SD away from PBS average value.	
Cellular differentiation and phenotype assays	
Assay Description	Assay Number
Human hepatocyte (HepG2C3a cells) acetylated LDL uptake	B1
Data analysis for ac-LDL uptake assay was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be a modulator of ac-LDL uptake if greater than 2 SD away from the PBS value in the positive or negative direction. A visual check to confirm plate reader results was made using a fluorescent microscope.	
Human Neutrophil assays	
Assay Description	Assay Number
Neutrophil Elastase	C1
Neutrophil oxidative burst (agonist)	C2
Neutrophil oxidative burst (antagonist)	C3
Data analysis for neutrophil assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be a modulator of neutrophil elastase production or oxidative burst biology if greater than 2 SD away from the PBS value in the positive or negative direction.	
Modulation of Toll-like receptors (TLR)	
Assay Description	Assay Number
TLR activation in RAW BLUE cells	D1
TLR antagonism in RAW BLUE cells	D2
Activation of hTLR2	D3
Activation of hTLR4	D4
Data analysis for TLR modulation assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be a modulator of TLR specific biology if greater than 3 SD away from the PBS value in the positive or negative direction. Positive controls,	

TABLE E6-continued

Key to Assays and criteria for indicating a hit	
including LPS and detection reagent were always significantly distinct and >3 SD from PBS average value.	
Cytokine Release	
Assay Description	Assay Number
Human Synovocyte cytokine production (IL6 release)	E1
Human pulmonary artery smooth muscle cell (hPASMC) cytokine production (IL6 release)	E2
Human skeletal muscle cell (hSKMC) cytokine production (IL6 release)	E3
Human Astrocyte cytokine production (IL6 release)	E4
Whole blood IL6 release	E5
Human pulmonary artery smooth muscle cell (hPASMC) cytokine production (IL8release) 72 h Incubation	E6
Z Assay Description	Assay Number
IL8 production	
Human Synovocyte cytokine production (IL8 release)	E7
Human pulmonary artery smooth muscle cell (hPASMC) cytokine production (IL8release)	E8
Human skeletal muscle cell (hSKMC) cytokine production (IL8 release)	E9
Human Astrocyte cytokine production (IL8 release)	E10
Human hepatocyte (HepG2C3a cells) IL8 release	E11
Human acute promyelocytic leukemia cells/HL60 (IL8 release)	E12
Human lymphoblast (cancer cell line)/RPMI8226 (IL8 Release)	E13
TNF alpha production	
Human Synovocyte cytokine production (TNF alpha release)	E14
Whole blood TNF alpha release	E15
IL10 Release	
Human acute promyelocytic leukemia cells/HL60 IL10 release	E16
Human Primary Blood Mononuclear cells (IL10 Release)	E17
Data analysis for cytokine release assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be a modulator of cytokine production or cytokine related biology if the measured value was greater than 2 SD away from the PBS value in the positive or negative direction. A protein standard (specific to each assay kit) was run on every plate to insure good assay quality. Only assays with protein standard curves that had an R ² value of > than 0.9 were chosen for data analysis.	
Cell Adhesion and Chemotaxis	
Assay Description	Assay Number
Monocyte THP 1/Human umbilical vein endothelial cell (HUVEC) cell adhesion	F1
Human hepatocyte (HepG2C3a cells) (ICAM release)	F2
Human lung microvascular endothelial cell (HLMVEC) cell adhesion regulation (ICAM release)	F3
Human umbilical vein endothelial cell (HUVEC) cell adhesion regulation (VCAM release)	F4
Human mesenchymal stem cell (hMSC) cell adhesion regulation (VCAM release)	F5
Human skeletal muscle cell (hSKMC) cell adhesion regulation (VCAM release)	F6
Human pulmonary artery smooth muscle cell (hPASMC) cell adhesion regulation (VCAM release)	F7
Data analysis for cell adhesion regulation assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. AARS polypeptides were considered to be a modulator of cell adhesion or a regulator of biology related to cell adhesion if the measured value was greater than 2 SD away from the PBS value in the positive or negative direction. In the case of the ELISA assays, a protein standard (specific to each assay kit) was run on every plate to insure good assay quality. Only assays with protein standard curves that had an R ² value of > than 0.9 were chosen for data analysis.	

TABLE E6-continued

Key to Assays and criteria for indicating a hit	
Cellular Differentiation	
Assay Description	Assay Number
Human pre-adipocyte (hPAD) cell differentiation	G1
Human skeletal muscle (hSKMC) cell differentiation	G2
Human mesenchymal stem (hMSC) cell differentiation	G3
Human pulmonary artery smooth muscle cell (hPASMC) differentiation	G4
Data analysis for cellular differentiation assays was performed by dividing the numerical value in the assay well by the average PBS value for the assay plate. Differentiation assays were scored based upon fluorescent intensity of particular antibodies as described in the methods section. AARS polypeptides were considered to be a modulator of cellular differentiation if an intensity value for a specific marker of differentiation was greater than 2 SD away from the PBS value in the positive or negative direction in a given treated well. For the hSKMC analysis, digital photos were taken of all wells and photos were scored in a blinded fashion by three people using a 4 point scoring system where a score of "4" indicated intense skeletal muscle actin staining and obvious myotube formation and a score of "1" indicated a lack of any differentiation or a suppression of differentiation. The average value from visual scoring was used and only wells with an average value of >3 were considered hits. Differentiation control treated wells in this assay typically scored >2, while PBS treated wells scored <2.	
Cell Binding	
Assay Description	Assay Number
PBMC	H1
Primary T cell	H2
Primary B cell	H3
Primary Monocyte	H4
HepG2	H5
3T3L1	H6
C2C12	H7
THP1	H8
Jurkat	H9
Raji	H10
AARS polypeptides were considered to be binding to a particular cell type if the mean cell bound fluorescence intensity was greater than 2 SD away from the reagent control values for that cell type.	

TABLE E7

Results of Functional Profiling studies of AARS Polypeptides			
AARS Polypeptides	Location of Epitope Tag	Concentration [nM]	Assay Hits
ArgRS ^{N2}	N-terminal	250	A5 (Proliferation)
ArgRS ^{N2}	C-terminal	250	
ArgRS ^{I1}	N-terminal	250	A6 (Proliferation) D2 (Modulation of Toll-like receptors) E4 (Cytokine Release) F5 (Cell Adhesion and Chemotaxis)
ArgRS ^{I1}	C-terminal	250	E4 (Cytokine Release) F5 (Cell Adhesion and Chemotaxis)
ArgRS ^{C2}	N-terminal	250	A6 (Proliferation)
ArgRS ^{C2}	C-terminal	250	

[0692] It is concluded that ArgRS1^{N2}, ArgRS1^{I1}, and ArgRS1^{C2} appear to be major regulators of cytokine release, proliferation, and cell adhesion and chemotaxis. Of note is that in many cases there is a significant difference in the relative activity of the N-terminally tagged and C-terminally tagged versions of the AARS polypeptides both in the

transcriptional profiling experiments, as well as in the phenotypic screening experiments. This data is consistent with the hypothesis that with these AARS polypeptides, the novel biological activity is suppressed when the AARS polypeptide is part of the intact tRNA synthetase, or translationally fused to another protein, but that this biological activity is revealed when the AARS polypeptides have a free amino or free carboxy terminus.

[0693] When viewed in the context of the transcriptional profiling studies, the phenotypic screening data demonstrates that the AARS polypeptides ArgRS1^{N2} (amino acids 1-71), ArgRS1^{I1} (amino acids 168-530) and ArgRS1^{C2} (amino acids 551-660) define the boundaries of a three novel protein domains that are highly active in a broad array of phenotypic screening assays.

[0694] Accordingly it is concluded that AARS polypeptides comprising amino acids 1-71, or 168-530, or 551-660 of Arginyl tRNA synthetase define the approximate boundaries (i.e. within about +/-5 amino acids) of a three distinct novel, highly active AARS polypeptide domains, that are i) highly functionally active, ii) can be readily made and produced in *E. coli*, and iii) exhibit favorable protein stability and aggregation characteristics.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 159

<210> SEQ ID NO 1

<211> LENGTH: 77

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 1

aggaggtaaa acatatgcat catcatcatc atcacggtaa gcctatccct aaccctttgc 60

tcggtctcga ttctacg 77

<210> SEQ ID NO 2

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 2

taatgactcg ag 12

<210> SEQ ID NO 3

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 3

aggagataaa acatatg 17

<210> SEQ ID NO 4

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 4

aggaggtaaa acat 14

<210> SEQ ID NO 5

<211> LENGTH: 14

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 5

aggagataaa acat 14

<210> SEQ ID NO 6

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 6

gaaggagata tacat 15

-continued

```

<210> SEQ ID NO 7
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 7

ggtaagccta tccctaacc tctcctcggg ctcgattcta cgcaccacca tcatcaccat   60
taatgactcg ag                                                    72

<210> SEQ ID NO 8
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 8

catatgcatc atcatcatca tcacggtaag cctatcccta accctctcct cggtctcgat   60
tctacgggat cc                                                    72

<210> SEQ ID NO 9
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 9

ctcgagtaat ga                                                    12

<210> SEQ ID NO 10
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 10

catatgggat cc                                                    12

<210> SEQ ID NO 11
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide

<400> SEQUENCE: 11

ctcgagggta agcctatccc taacctcttc ctcgggtctcg attctacgca ccaccaccac   60
caccactaat ga                                                    72

<210> SEQ ID NO 12
<211> LENGTH: 398
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1          5          10          15

```

-continued

Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
 20 25 30
 Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
 35 40 45
 Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
 50 55 60
 Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile Ile Ser Arg Leu
 65 70 75 80
 Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr Pro Asp Leu Glu
 85 90 95
 Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala Lys Phe Gly Asp
 100 105 110
 Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Thr Lys
 115 120 125
 Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys His
 130 135 140
 Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile Ala Gly Pro Gly
 145 150 155 160
 Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu Thr
 165 170 175
 Ser Leu Leu Val Asn Gly Val Gln Leu Pro Ala Leu Gly Glu Asn Lys
 180 185 190
 Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met His
 195 200 205
 Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Ile Ser Arg Leu
 210 215 220
 Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly Asp
 225 230 235 240
 Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys Phe
 245 250 255
 Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Val Phe
 260 265 270
 Tyr Lys Glu Ser Lys Lys Arg Phe Asp Thr Glu Glu Glu Phe Lys Lys
 275 280 285
 Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Gly Lys Asn Pro Asp Ile
 290 295 300
 Thr Lys Ala Trp Lys Leu Ile Cys Asp Val Ser Arg Gln Glu Leu Asn
 305 310 315 320
 Lys Ile Tyr Asp Ala Leu Asp Val Ser Leu Ile Glu Arg Gly Glu Ser
 325 330 335
 Phe Tyr Gln Asp Arg Met Asn Asp Ile Val Lys Glu Phe Glu Asp Arg
 340 345 350
 Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe Val Pro Gly
 355 360 365
 Cys Ser Ile Pro Leu Thr Ile Val Lys Ser Asp Gly Gly Tyr Thr Tyr
 370 375 380
 Asp Thr Ser Asp Leu Ala Ala Ile Lys Gln Arg Leu Phe Glu
 385 390 395

<210> SEQ ID NO 13

<211> LENGTH: 1194

-continued

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

```

atggacgtac tgggtgtctga gtgctccgcg cggctgctgc agcaggaaga agagattaaa    60
tctctgactg ctgaaattga ccggttgaaa aactgtggct gtttaggagc ttctccaaat    120
ttggagcagt tacaagaaga aaatttaaaa ttaaagtatc gactgaatat tcttcgaaag    180
agtcttcagg cagaaggaa caaaccaact aaaaatatga ttaacattat tagccgccta    240
caagaggctt ttggtcatgc aattaaggct gcatatccag atttgaaaa tctcctctg    300
ctagtgcac caagtcagca ggccaagttt ggggactatc agtgtaatag tgctatgggt    360
atttctcaga tgctcaaac caaggaacag aaagttaatc caagagaaat tgctgaaac    420
attaccaaac acctcccaga caatgaatgt attgaaaag ttgaaattgc tggctcctgt    480
tttattaatg tccacttaag aaaggatttt gtatcagaac aattgaccag tcttctagtg    540
aatggagttc aactacctgc tctgggagag aataaaaagg ttatagtga ctttctctcc    600
cctaatatag ctaagagat gcatgtaggc cacctgaggt caactatcat aggagagagt    660
ataagccgcc tctttgaatt tgcagggtat gacgtgctca gggttaatca tgtaggagac    720
tgggggagccc agtttgcat gctcatcgct cacctgcaag acaaatttcc agattatcta    780
acagtttca ctcctattgg ggatcttcag gtcttttata aggaatctaa gaagaggttt    840
gatactgagg aggaatttaa gaagcgagca tatcagtgtg tagttctgct ccagggtaaa    900
aaccagata ttacaaaagc ttggaagctt atctgtgatg tctcccgcca agagttaaat    960
aaaatctatg atgcattgga cgtctcttta atagagagag gggaatcctt ctatcaagat   1020
aggatgaatg atattgtaaa ggaatttgaa gatagaggat ttgtgcaggt ggatgatggc   1080
agaaagattg tatttgtccc aggggtgttc ataccattaa ccatagtaaa atcagatgga   1140
ggttatacct atgatacatc tgacctggct gctattaaac aaagactatt tgag           1194

```

<210> SEQ ID NO 14

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 14

```

Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Ile Val Thr Pro Ser
1          5              10              15
Gln Gln Pro Lys
          20

```

<210> SEQ ID NO 15

<211> LENGTH: 59

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 15

```

Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu
1          5              10              15
Lys Ala Lys Glu Gln Lys Val Ser Pro Arg Glu Ile Ala Glu Asn Ile
          20              25              30
Thr Lys His Leu Pro Asn Asn Lys Tyr Ile Asp Lys Val Glu Ile Ala
          35              40              45

```

-continued

Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys
50 55

<210> SEQ ID NO 16
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 16

Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln
 1 5 10 15
 Leu Pro Val Leu Gly Asp Lys Glu Lys
 20 25

<210> SEQ ID NO 17
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 17

Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys
 1 5 10

<210> SEQ ID NO 18
 <211> LENGTH: 164
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 18

Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met
 1 5 10 15
 Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His
 20 25 30
 Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln
 35 40 45
 Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu
 50 55 60
 Gln Ala Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu
 65 70 75 80
 Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn
 85 90 95
 Pro Asp Ile Met Lys Ala Trp Asn Leu Ile Cys Asp Val Ser Arg Glu
 100 105 110
 Glu Phe Lys Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg
 115 120 125
 Gly Glu Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe
 130 135 140
 Glu Asp Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe
 145 150 155 160
 Val Pro Gly Cys

<210> SEQ ID NO 19
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 19

-continued

Ser	Val	Pro	Leu	Thr	Ile	Val	Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp
1				5					10					15	

Thr	Ser	Asp	Leu	Ala	Ala	Ile	Lys
			20				

<210> SEQ ID NO 20
 <211> LENGTH: 304
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 20

Ala	Ala	Tyr	Pro	Asp	Leu	Glu	Asn	Pro	Pro	Leu	Ile	Val	Thr	Pro	Ser
1				5					10					15	

Gln	Gln	Pro	Lys	Phe	Gly	Asp	Tyr	Gln	Cys	Asn	Ser	Ala	Met	Gly	Ile
			20					25					30		

Ser	Gln	Met	Leu	Lys	Ala	Lys	Glu	Gln	Lys	Val	Ser	Pro	Arg	Glu	Ile
		35					40					45			

Ala	Glu	Asn	Ile	Thr	Lys	His	Leu	Pro	Asn	Asn	Lys	Tyr	Ile	Asp	Lys
	50					55					60				

Val	Glu	Ile	Ala	Gly	Pro	Gly	Phe	Ile	Asn	Val	His	Leu	Arg	Lys	Asp
65					70					75				80	

Phe	Val	Ser	Glu	Gln	Leu	Thr	Ser	Leu	Leu	Val	Asn	Gly	Val	Gln	Leu
			85						90					95	

Pro	Val	Leu	Gly	Asp	Lys	Glu	Lys	Val	Ile	Val	Asp	Phe	Ser	Ser	Pro
		100						105					110		

Asn	Ile	Ala	Lys	Glu	Met	His	Val	Gly	His	Leu	Arg	Ser	Thr	Ile	Ile
		115					120					125			

Gly	Glu	Ser	Met	Ser	Arg	Leu	Phe	Glu	Phe	Ala	Gly	Tyr	Asp	Val	Leu
	130					135					140				

Arg	Leu	Asn	His	Val	Gly	Asp	Trp	Gly	Thr	Gln	Phe	Gly	Met	Leu	Ile
145				150						155				160	

Ala	His	Leu	Gln	Asp	Lys	Phe	Pro	Asp	Tyr	Leu	Thr	Val	Ser	Pro	Pro
			165						170					175	

Ile	Gly	Asp	Leu	Gln	Ala	Phe	Tyr	Lys	Glu	Ser	Lys	Lys	Arg	Phe	Asp
		180						185					190		

Ala	Asp	Glu	Glu	Phe	Lys	Lys	Arg	Ala	Tyr	Gln	Cys	Val	Val	Leu	Leu
		195					200					205			

Gln	Ser	Lys	Asn	Pro	Asp	Ile	Met	Lys	Ala	Trp	Asn	Leu	Ile	Cys	Asp
	210					215					220				

Val	Ser	Arg	Glu	Glu	Phe	Lys	Lys	Ile	Tyr	Asp	Ala	Leu	Asp	Ile	Thr
225					230					235				240	

Leu	Ile	Glu	Arg	Gly	Glu	Ser	Phe	Tyr	Gln	Asp	Arg	Met	Lys	Asp	Ile
			245					250					255		

Val	Lys	Glu	Phe	Glu	Asp	Lys	Gly	Phe	Val	Gln	Val	Asp	Asp	Gly	Arg
		260						265					270		

Lys	Ile	Val	Phe	Val	Pro	Gly	Cys	Ser	Val	Pro	Leu	Thr	Ile	Val	Lys
		275					280					285			

Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr	Ser	Asp	Leu	Ala	Ala	Ile	Lys
	290					295					300				

<210> SEQ ID NO 21
 <211> LENGTH: 373
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 21

```

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1      5      10      15
Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
20      25      30
Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
35      40      45
Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
50      55      60
Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile Ile Ser Arg Leu
65      70      75      80
Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr Pro Asp Leu Glu
85      90      95
Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala Lys Phe Gly Asp
100     105     110
Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Thr Lys
115     120     125
Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys His
130     135     140
Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile Ala Gly Pro Gly
145     150     155     160
Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu Thr
165     170     175
Ser Leu Leu Val Asn Gly Val Gln Leu Pro Ala Leu Gly Glu Asn Lys
180     185     190
Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met His
195     200     205
Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Ile Ser Arg Leu
210     215     220
Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly Asp
225     230     235     240
Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys Phe
245     250     255
Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Val Phe
260     265     270
Tyr Lys Glu Ser Lys Lys Arg Phe Asp Thr Glu Glu Glu Phe Lys Lys
275     280     285
Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Gly Lys Asn Pro Asp Ile
290     295     300
Thr Lys Ala Trp Lys Leu Ile Cys Asp Val Ser Arg Gln Glu Leu Asn
305     310     315     320
Lys Ile Tyr Asp Ala Leu Asp Val Ser Leu Ile Glu Arg Gly Glu Ser
325     330     335
Phe Tyr Gln Asp Arg Met Asn Asp Ile Val Lys Glu Phe Glu Asp Arg
340     345     350
Gly Lys Ser Leu Lys His Val Arg Val Lys Gln Cys Ala Ser Trp Ile
355     360     365
Phe Trp Glu Lys Asp
370

```

-continued

```

<210> SEQ ID NO 22
<211> LENGTH: 1122
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

atggacgtac tgggtgtctga gtgctccgcg cggctgctgc agcaggaaga agagattaaa    60
tctctgactg ctgaaattga ccggttgaaa aactgtggct gtttaggagc ttctccaaat    120
ttggagcagt tacaagaaga aaatttataa ttaaagtatc gactgaatat tcttcgaaag    180
agtcttcagg cagaaggaa caaaccaact aaaaatatga ttaacattat tagccgccta    240
caagaggtct ttggtcatgc aattaaggct gcatatccag atttgaaaa tcctcctctg    300
ctagtgcac caagtcagca ggccaagttt ggggactatc agtgtaatag tgctatgggt    360
atctctcaga tgctcaaac caaggaacag aaagttaatc caagagaaat tgctgaaac    420
attacaaac acctcccaga caatgaatgt attgaaaaag ttgaaattgc tggctcctggt    480
tttattaatg tccacttaag aaaggatttt gtatcagaac aattgaccag tcttctagt    540
aatggagttc aactacctgc tctgggagag aataaaaagg ttatagttga cttttcctcc    600
cctaatatag ctaaagagat gcatgtaggc cacctgaggt caactatcat aggagagagt    660
ataagccgcc tctttgaatt tgcagggtat gacgtgctca gggtaaatca tgtaggagac    720
tggggggacc agtttgcat gctcatcgct cacctgcaag acaaatttcc agattatcta    780
acagtttcac ctctatttgg ggatcttcag gtcttttata aggaatctaa gaagaggttt    840
gatactgagg aggaatttaa gaagcgagca tatcagtgtg tagttctgct ccagggtaaa    900
aaccagata ttacaaaagc ttggaagctt atctgtgatg tctcccgcca agagttaaat    960
aaaatctatg atgcattgga cgtctcttta atagagagag gggaatcctt ctatcaagat   1020
aggatgaatg atattgtaaa ggaatttgaa gatagaggaa aaagtttaaa acacgttcgg   1080
gtgaaacagt gcgcctcatg gatcttctgg gagaaggact aa                        1122

```

```

<210> SEQ ID NO 23
<211> LENGTH: 160
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1          5          10          15
Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
20         25         30
Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Asn
35         40         45
Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
50         55         60
Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile Ile Ser Arg Leu
65         70         75         80
Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr Pro Asp Leu Glu
85         90         95
Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala Lys Phe Gly Asp
100        105        110
Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Thr Lys
115        120        125

```

-continued

Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys His
130 135 140

Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile Ala Gly Pro Gly
145 150 155 160

<210> SEQ ID NO 24

<211> LENGTH: 483

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

```
atggacgtac tgggtgtctga gtgctccgcg cggctgctgc agcaggaaga agagattaaa    60
tctctgactg ctgaaattga ccggttgaaa aactgtggct gtttaggagc ttctccaaat    120
ttggagcagt tacaagaaga aaatttataa ttaaagtatc gactgaatat tcttcgaaag    180
agtcttcagg cagaaaggaa caaaccaact aaaaatatga ttaacattat tagccgecta    240
caagaggtct ttggtcatgc aattaaggct gcatatccag atttgaaaaa tcctcctctg    300
ctagtgcacac caagtcagca ggccaagttt ggggactatc agtgtaatag tgctatgggt    360
atttctcaga tgctcaaaac caaggaacag aaagttaatc caagagaaat tgctgaaaac    420
attaccaaac acctcccaga caatgaatgt attgaaaaag ttgaaattgc tggtcctggt    480
taa                                                    483
```

<210> SEQ ID NO 25

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

```
attgtaaagg aatttgaaga tagaggaaaa agttttaaac acgttcgggt    50
```

<210> SEQ ID NO 26

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Ile Val Lys Glu Phe Glu Asp Arg Gly Lys Ser Leu Lys His Val Arg
1 5 10 15

<210> SEQ ID NO 27

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

```
gaaaaagttg aaattgctgg tcctgggttaa atcatgtagg agactggggg    50
```

<210> SEQ ID NO 28

<211> LENGTH: 9

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28

Glu Lys Val Glu Ile Ala Gly Pro Gly
1 5

-continued

<210> SEQ ID NO 29
 <211> LENGTH: 71
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29

```

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1           5           10           15
Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
                20           25           30
Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
                35           40           45
Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
                50           55           60
Glu Arg Asn Lys Pro Thr Lys
65           70

```

<210> SEQ ID NO 30
 <211> LENGTH: 213
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30

```

atggacgtac tgggtgtctga gtgctccgcg cggctgctgc agcaggaaga agagattaaa      60
tctctgactg ctgaaattga ccggttgaaa aactgtggct gtttaggagc ttctccaaat      120
ttggagcagt tacaagaaga aaatttaaaa ttaaagtatc gactgaatat tcttcgaaag      180
agtcttcagg cagaaaggaa caaaccaact aaa                               213

```

<210> SEQ ID NO 31
 <211> LENGTH: 418
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
 N-terminal 6xHis affinity tag

<400> SEQUENCE: 31

```

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
1           5           10           15
Gly Leu Asp Ser Thr Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu
                20           25           30
Leu Gln Gln Glu Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg
                35           40           45
Leu Lys Asn Cys Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu
                50           55           60
Gln Glu Glu Asn Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys
65           70           75           80
Ser Leu Gln Ala Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile
                85           90           95
Ile Ser Arg Leu Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr
                100          105          110
Pro Asp Leu Glu Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala
                115          120          125
Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met
130          135          140

```

-continued

```

Leu Lys Thr Lys Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn
145                      150                      155                      160

Ile Thr Lys His Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile
                      165                      170                      175

Ala Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser
                      180                      185                      190

Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln Leu Pro Ala Leu
                      195                      200                      205

Gly Glu Asn Lys Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala
210                      215                      220

Lys Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser
225                      230                      235                      240

Ile Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn
                      245                      250                      255

His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu
                      260                      265                      270

Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp
                      275                      280                      285

Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Thr Glu Glu
290                      295                      300

Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Gly Lys
305                      310                      315                      320

Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu Ile Cys Asp Val Ser Arg
                      325                      330                      335

Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu Asp Val Ser Leu Ile Glu
                      340                      345                      350

Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Asn Asp Ile Val Lys Glu
                      355                      360                      365

Phe Glu Asp Arg Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val
370                      375                      380

Phe Val Pro Gly Cys Ser Ile Pro Leu Thr Ile Val Lys Ser Asp Gly
385                      390                      395                      400

Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys Gln Arg Leu
                      405                      410                      415

Phe Glu

```

```

<210> SEQ ID NO 32
<211> LENGTH: 418
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
      C-terminal 6xHis affinity tag

```

```

<400> SEQUENCE: 32

```

```

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1                      5                      10                      15

Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
20                      25                      30

Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
35                      40                      45

Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
50                      55                      60

```

-continued

Glu	Arg	Asn	Lys	Pro	Thr	Lys	Asn	Met	Ile	Asn	Ile	Ile	Ser	Arg	Leu
65					70					75					80
Gln	Glu	Val	Phe	Gly	His	Ala	Ile	Lys	Ala	Ala	Tyr	Pro	Asp	Leu	Glu
			85						90					95	
Asn	Pro	Pro	Leu	Leu	Val	Thr	Pro	Ser	Gln	Gln	Ala	Lys	Phe	Gly	Asp
			100					105					110		
Tyr	Gln	Cys	Asn	Ser	Ala	Met	Gly	Ile	Ser	Gln	Met	Leu	Lys	Thr	Lys
		115					120					125			
Glu	Gln	Lys	Val	Asn	Pro	Arg	Glu	Ile	Ala	Glu	Asn	Ile	Thr	Lys	His
	130					135					140				
Leu	Pro	Asp	Asn	Glu	Cys	Ile	Glu	Lys	Val	Glu	Ile	Ala	Gly	Pro	Gly
145					150					155					160
Phe	Ile	Asn	Val	His	Leu	Arg	Lys	Asp	Phe	Val	Ser	Glu	Gln	Leu	Thr
			165						170					175	
Ser	Leu	Leu	Val	Asn	Gly	Val	Gln	Leu	Pro	Ala	Leu	Gly	Glu	Asn	Lys
			180					185					190		
Lys	Val	Ile	Val	Asp	Phe	Ser	Ser	Pro	Asn	Ile	Ala	Lys	Glu	Met	His
		195						200				205			
Val	Gly	His	Leu	Arg	Ser	Thr	Ile	Ile	Gly	Glu	Ser	Ile	Ser	Arg	Leu
	210					215					220				
Phe	Glu	Phe	Ala	Gly	Tyr	Asp	Val	Leu	Arg	Leu	Asn	His	Val	Gly	Asp
225					230					235					240
Trp	Gly	Thr	Gln	Phe	Gly	Met	Leu	Ile	Ala	His	Leu	Gln	Asp	Lys	Phe
			245						250					255	
Pro	Asp	Tyr	Leu	Thr	Val	Ser	Pro	Pro	Ile	Gly	Asp	Leu	Gln	Val	Phe
			260					265					270		
Tyr	Lys	Glu	Ser	Lys	Lys	Arg	Phe	Asp	Thr	Glu	Glu	Glu	Phe	Lys	Lys
		275					280					285			
Arg	Ala	Tyr	Gln	Cys	Val	Val	Leu	Leu	Gln	Gly	Lys	Asn	Pro	Asp	Ile
	290					295					300				
Thr	Lys	Ala	Trp	Lys	Leu	Ile	Cys	Asp	Val	Ser	Arg	Gln	Glu	Leu	Asn
305					310					315					320
Lys	Ile	Tyr	Asp	Ala	Leu	Asp	Val	Ser	Leu	Ile	Glu	Arg	Gly	Glu	Ser
			325						330					335	
Phe	Tyr	Gln	Asp	Arg	Met	Asn	Asp	Ile	Val	Lys	Glu	Phe	Glu	Asp	Arg
			340					345					350		
Gly	Phe	Val	Gln	Val	Asp	Asp	Gly	Arg	Lys	Ile	Val	Phe	Val	Pro	Gly
		355					360					365			
Cys	Ser	Ile	Pro	Leu	Thr	Ile	Val	Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr
	370					375					380				
Asp	Thr	Ser	Asp	Leu	Ala	Ala	Ile	Lys	Gln	Arg	Leu	Phe	Glu	Gly	Lys
385					390					395					400
Pro	Ile	Pro	Asn	Pro	Leu	Leu	Gly	Leu	Asp	Ser	Thr	His	His	His	His
				405					410					415	

His His

<210> SEQ ID NO 33

<211> LENGTH: 91

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with N-terminal 6xHis affinity tag

-continued

<400> SEQUENCE: 33

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
1 5 10 15
Gly Leu Asp Ser Thr Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu
20 25 30
Leu Gln Gln Glu Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg
35 40 45
Leu Lys Asn Cys Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu
50 55 60
Gln Glu Glu Asn Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys
65 70 75 80
Ser Leu Gln Ala Glu Arg Asn Lys Pro Thr Lys
85 90

<210> SEQ ID NO 34

<211> LENGTH: 91

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
C-terminal 6xHis affinity tag

<400> SEQUENCE: 34

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1 5 10 15
Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
20 25 30
Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
35 40 45
Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
50 55 60
Glu Arg Asn Lys Pro Thr Lys Gly Lys Pro Ile Pro Asn Pro Leu Leu
65 70 75 80
Gly Leu Asp Ser Thr His His His His His
85 90

<210> SEQ ID NO 35

<211> LENGTH: 393

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
N-terminal 6xHis affinity tag

<400> SEQUENCE: 35

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
1 5 10 15
Gly Leu Asp Ser Thr Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu
20 25 30
Leu Gln Gln Glu Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg
35 40 45
Leu Lys Asn Cys Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu
50 55 60
Gln Glu Glu Asn Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys
65 70 75 80

-continued

```

Ser Leu Gln Ala Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile
      85                      90
Ile Ser Arg Leu Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr
      100                    105          110
Pro Asp Leu Glu Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala
      115                    120          125
Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met
      130                    135          140
Leu Lys Thr Lys Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn
      145                    150          155          160
Ile Thr Lys His Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile
      165                    170          175
Ala Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser
      180                    185          190
Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln Leu Pro Ala Leu
      195                    200          205
Gly Glu Asn Lys Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala
      210                    215          220
Lys Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser
      225                    230          235          240
Ile Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn
      245                    250          255
His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu
      260                    265          270
Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp
      275                    280          285
Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Thr Glu Glu
      290                    295          300
Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Gly Lys
      305                    310          315          320
Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu Ile Cys Asp Val Ser Arg
      325                    330          335
Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu Asp Val Ser Leu Ile Glu
      340                    345          350
Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Asn Asp Ile Val Lys Glu
      355                    360          365
Phe Glu Asp Arg Gly Lys Ser Leu Lys His Val Arg Val Lys Gln Cys
      370                    375          380
Ala Ser Trp Ile Phe Trp Glu Lys Asp
      385                    390

<210> SEQ ID NO 36
<211> LENGTH: 393
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
      C-terminal 6xHis affinity tag

<400> SEQUENCE: 36

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
1          5          10          15
Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
20          25          30

```

```
<210> SEQ ID NO 37
<211> LENGTH: 180
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
```

-continued

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
N-terminal 6xHis affinity tag

<400> SEQUENCE: 37

```

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
 1           5           10           15

Gly Leu Asp Ser Thr Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu
 20           25           30

Leu Gln Gln Glu Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg
 35           40           45

Leu Lys Asn Cys Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu
 50           55           60

Gln Glu Glu Asn Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys
 65           70           75           80

Ser Leu Gln Ala Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile
 85           90           95

Ile Ser Arg Leu Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr
100          105          110

Pro Asp Leu Glu Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala
115          120          125

Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met
130          135          140

Leu Lys Thr Lys Glu Gln Lys Val Asn Pro Arg Glu Ile Ala Glu Asn
145          150          155          160

Ile Thr Lys His Leu Pro Asp Asn Glu Cys Ile Glu Lys Val Glu Ile
165          170          175

Ala Gly Pro Gly
180

```

<210> SEQ ID NO 38

<211> LENGTH: 180

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
C-terminal 6xHis affinity tag

<400> SEQUENCE: 38

```

Met Asp Val Leu Val Ser Glu Cys Ser Ala Arg Leu Leu Gln Gln Glu
 1           5           10           15

Glu Glu Ile Lys Ser Leu Thr Ala Glu Ile Asp Arg Leu Lys Asn Cys
 20           25           30

Gly Cys Leu Gly Ala Ser Pro Asn Leu Glu Gln Leu Gln Glu Glu Asn
 35           40           45

Leu Lys Leu Lys Tyr Arg Leu Asn Ile Leu Arg Lys Ser Leu Gln Ala
 50           55           60

Glu Arg Asn Lys Pro Thr Lys Asn Met Ile Asn Ile Ile Ser Arg Leu
 65           70           75           80

Gln Glu Val Phe Gly His Ala Ile Lys Ala Ala Tyr Pro Asp Leu Glu
 85           90           95

Asn Pro Pro Leu Leu Val Thr Pro Ser Gln Gln Ala Lys Phe Gly Asp
100          105          110

Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Thr Lys
115          120          125

```

-continued

Glu	Gln	Lys	Val	Asn	Pro	Arg	Glu	Ile	Ala	Glu	Asn	Ile	Thr	Lys	His
130						135					140				
Leu	Pro	Asp	Asn	Glu	Cys	Ile	Glu	Lys	Val	Glu	Ile	Ala	Gly	Pro	Gly
145				150						155				160	
Gly	Lys	Pro	Ile	Pro	Asn	Pro	Leu	Leu	Gly	Leu	Asp	Ser	Thr	His	His
				165					170					175	
His	His	His	His												
			180												

<210> SEQ ID NO 39
 <211> LENGTH: 1203
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase polynucleotide

<400> SEQUENCE: 39

gacgtactgg tcacgcaatg cagcgcacgc ttactccaac aggaagaaga aatcaaatct	60
ctgacggccg agatcgaccg tctgaagaat tgtggctgtt tgggtgagag cccgaacctg	120
gagcagctgc aggaagaaaa tctgaaactg aagtaccgcc tgaatattct gcgtaaatcc	180
ttgcaggcag agcgtaacaa accgaccaag aacatgatta atattattag ccgtttgcag	240
gaagtgtttg gccacgccat caaggcggcg tatccggatc tggagaaccc tccgtgtctg	300
gtgaccccg aaccaacaagc taaatttggg tactaccagt gtaactctgc aatgggtatt	360
agccagatgc tgaaaaccaa agaacagaaa gtgaaccgcg gcgagatcgc ggagaacatc	420
accaagcacc tgccggacaa cgagtgcatt gagaaggtgg agatcgagg tccgggtttc	480
atcaatgtcc atctgcgtaa ggatttcgtt tccgaacaac tgaccagcct gttggtgaat	540
ggcgttcagc tgcttgccct gggtgagaac aagaaagtca ttgtggactt tagcagccct	600
aatattgcga aagagatgca cgtgggccac ctgcgtagca ccattattgg cgagtcgac	660
agccgcctgt ttgagttcgc gggttatgat gtcttgctgc tgaatcatgt tgggtgactgg	720
ggtacgcagt tcggcatgct gatcgcgcat ctgcaggata agttccaga ctatttgacg	780
gtcagcccg cgattggcga tctgcaagtg ttttacaag agagcaagaa gcgctttgac	840
accgaagaag aattcaagaa gcgtgcctat caatgcgttg ttctgctgca aggaagaat	900
ccggacatta ctaaggcgtg gaaattgatc tgcgacgtca gccgtcaaga actgaacaaa	960
atctacgacg cgctggatgt tagcctgatc gagcgtgggt agagcttcta tcaggatcgt	1020
atgaacgata ttgttaaaga gttcgaggac cgcggttttg ttcaagtga cgatggtcgc	1080
aaaattgtct ttgtccgggg ttgcagcatt ccgctgacga tcgtgaagtc tgatggcgg	1140
tacacctacg atacgtccga cctggctgca atcaacaac gtctgttcga ataatgactc	1200
gag	1203

<210> SEQ ID NO 40
 <211> LENGTH: 222
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase polynucleotide

<400> SEQUENCE: 40

-continued

gacgtactgg ttagcgaatg ctccgcacgc ctgctgcaac aagaagaaga gatcaaaagc	60
ctgaccgccc agatcgaccg tctgaagaat tgcggctgtc tgggtgcatc tccgaacctg	120
gaacagctgc aggaagagaa tctgaagctg aagtaccgtt tgaacatttt gcgtaaaagc	180
ctgcaagcgg agcgcaacaa accgcagaaa taatgactcg ag	222

<210> SEQ ID NO 41
 <211> LENGTH: 1128
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase polynucleotide

<400> SEQUENCE: 41

gacgtactgg ttagcgaatg ctccgcacgc ctgctgcaac aagaagaaga aattaaatcc	60
ctgacggctg agattgaccg tctgaagaac tgtggttgcc tgggtgagag cccgaatctg	120
gagcagttgc aagaagagaa cctgaaactg aaatatcgtc tgaatatctt gcgcaagagc	180
ctgcaagcag agcgcaacaa gccgaccaag aacatgatta atatcatcag ccgcctgcaa	240
gaagtctttg gtcacgcgat caaggcagca taccgggacc tggaaaatcc gccgttgctg	300
gtgaccccca gccagcaggc gaagtccggc gactaccaat gcaattccgc tatgggcatt	360
agccagatgc tgaaaacgaa agaacaaaag gtcaatccgc gcgagatcgc ggaaaacatt	420
acgaagcacc tgccggacaa cgagtgtatt gagaaagtgg agatcgccgg tccaggtttc	480
atcaatgttc acctgcgtaa agatttcgtg tctgagcagt tgaccagcct gctggtgaac	540
ggcgttcaac tgccggcgct gggtgagaac aagaaggatg ttgtcgactt ttctagcccg	600
aacatcgcca aagagatgca tgcgggcat ctgcgttcca ccattattgg tgaaagcatc	660
agccgtctgt tcgagtttgc gggctacgat gttctgcgct tgaaccacgt cggtgactgg	720
gggtactcaat tcggtatgct gatcgacat ttgcaggata aatttcggga ctacctgacc	780
gtgagccccc caattggtga ttgacaggtc ttttaccaaag aaagcaagaa acgctttgat	840
acggaagaag aattcaagaa acgtgcctat cagtgtgttg ttctgctgca gggcaagaat	900
cctgatatca ccaaaagcgt gaaactgac tgcgatgta gccgtcaaga gctgaataag	960
atztatgacg ccctggacgt tagcctgac gagcgtggcg agagcttcta tcaggaccgt	1020
atgaatgata tcgttaaaga gtttgaggac cgcggcaaaa gcttgaagca cgtgcgtgtg	1080
aaacagtgcg cgtcgtggat tttctgggaa aaggattaat gactcgag	1128

<210> SEQ ID NO 42
 <211> LENGTH: 489
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase polynucleotide

<400> SEQUENCE: 42

gacgtactgg ttagcgaatg ctccgcacgc ctgctgcaac aagaagaaga gattaagagc	60
ctgacggcag agatcgaccg cctgaagaac tgcggttgtc tgggtgagag cccgaacctg	120
gaacagctgc aagaagaaaa tctgaaactg aagtatcgtc tgaatatctt gcgtaagagc	180
ttgcaggctg aacgtaacaa gccaaccaag aacatgatta acatcatttc ccgtctgcaa	240

-continued

```

gaggtctttg gtcacgcgat taaggccgcg tacccggatc tggagaatcc gccgctgctg   300
gttaccctt cccaacaagc gaaattcggc gactaccagt gcaatagcgc aatgggcatc   360
agccagatgc tgaaaaccaa agagcagaaa gtgaatccgc gcgaaattgc agagaacatt   420
acgaaacatc tgccggataa tgagtgtatc gagaaagttg agatcgccgg cccgggttaa   480
tgactcgag                                     489

```

```

<210> SEQ ID NO 43
<211> LENGTH: 588
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 43

```

```

Met Ile Asn Ile Ile Ser Arg Leu Gln Glu Val Phe Gly His Ala Ile
 1             5             10             15

Lys Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Leu Val Thr Pro
 20             25             30

Ser Gln Gln Ala Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly
 35             40             45

Ile Ser Gln Met Leu Lys Thr Lys Glu Gln Lys Val Asn Pro Arg Glu
 50             55             60

Ile Ala Glu Asn Ile Thr Lys His Leu Pro Asp Asn Glu Cys Ile Glu
 65             70             75             80

Lys Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys
 85             90             95

Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln
100             105             110

Leu Pro Ala Leu Gly Glu Asn Lys Lys Val Ile Val Asp Phe Ser Ser
115             120             125

Pro Asn Ile Ala Lys Glu Met His Val Gly His Leu Arg Ser Thr Ile
130             135             140

Ile Gly Glu Ser Ile Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val
145             150             155             160

Leu Arg Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu
165             170             175

Ile Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro
180             185             190

Pro Ile Gly Asp Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys Arg Phe
195             200             205

Asp Thr Glu Glu Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu
210             215             220

Leu Gln Gly Lys Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu Ile Cys
225             230             235             240

Asp Val Ser Arg Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu Asp Val
245             250             255

Ser Leu Ile Glu Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Asn Asp
260             265             270

Ile Val Lys Glu Phe Glu Asp Arg Gly Phe Val Gln Val Asp Asp Gly
275             280             285

Arg Lys Ile Val Phe Val Pro Gly Cys Ser Ile Pro Leu Thr Ile Val
290             295             300

Lys Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile

```

-continued

305	310	315	320
Lys Gln Arg Leu Phe Glu Glu Lys Ala Asp Met Ile Ile Tyr Val Val	325	330	335
Asp Asn Gly Gln Ser Val His Phe Gln Thr Ile Phe Ala Ala Ala Gln	340	345	350
Met Ile Gly Trp Tyr Asp Pro Lys Val Thr Arg Val Phe His Ala Gly	355	360	365
Phe Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr Arg Ser	370	375	380
Gly Glu Thr Val Arg Leu Met Asp Leu Leu Gly Glu Gly Leu Lys Arg	385	390	395
Ser Met Asp Lys Leu Lys Glu Lys Glu Arg Asp Lys Val Leu Thr Ala	405	410	415
Glu Glu Leu Asn Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys	420	425	430
Tyr Ala Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe	435	440	445
Asp Lys Met Leu Asp Asp Arg Gly Asn Thr Ala Ala Tyr Leu Leu Tyr	450	455	460
Ala Phe Thr Arg Ile Arg Ser Ile Ala Arg Leu Ala Asn Ile Asp Glu	465	470	475
Glu Met Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu Leu Asp His	485	490	495
Glu Lys Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile	500	505	510
Leu Gln Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys Asp Tyr	515	520	525
Ile Tyr Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp Ser Cys Tyr	530	535	540
Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys Val Asn Met	545	550	555
Trp Arg Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys Gly	565	570	575
Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg Met	580	585	

<210> SEQ ID NO 44

<211> LENGTH: 1767

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

```

atgattaaca ttattagccg cctacaagag gtctttggtc atgcaattaa ggctgcatat    60
ccagatttgg aaaatcctcc tctgctagtg acaccaagtc agcaggccaa gtttggggac    120
tatcagtgta atagtgtat ggggtatttct cagatgctca aaaccaagga acagaaaagt    180
aatccaagag aaattgctga aaacattacc aaacacctcc cagacaatga atgtattgaa    240
aaagttgaaa ttgctggtcc tgggttttatt aatgtccact taagaaagga ttttgtatca    300
gaacaattga ccagtcttct agtgaatgga gttcaactac ctgctctggg agagaataaa    360
aaggttatag ttgacttttc ctccccta atagctaaag agatgcatgt aggccacctg    420
aggtcaacta tcataggaga gagtataagc cgcctctttg aatttcagg gtatgacgtg    480

```

-continued

```

ctcaggttaa atcatgtagg agactggggg acccagtttg gcatgctcat cgctcacctg 540
caagacaaat ttccagatta tctaacagtt tcacctccta ttggggatct tcagggtcttt 600
tataaggaat ctaagaagag gtttgatact gaggaggaat ttaagaagcg agcatatcag 660
tgtgtagttc tgctccaggg taaaaacca gatattacaa aagcttgga gcttatctgt 720
gatgtctccc gccaagagtt aaataaaatc tatgatgcat tggacgtctc tttaatagag 780
agaggggaat ccttctatca agataggatg aatgatattg taaaggaatt tgaagataga 840
ggatttgtgc aggtggatga tggcagaaag attgtatttg tcccagggtg ttccatacca 900
ttaaccatag taaaatcaga tggaggttat acctatgata catctgacct ggctgctatt 960
aaacaagac tatttgagga aaaagcagat atgattatct atgttgtgga caatggacaa 1020
tctgtgcact tccagacaat atttgtgct gctcaaatga ttggttgga tgacctaaa 1080
gtaactcgag tcttccatgc tggatttggg gtggtgctag gggaagacaa gaaaaagttt 1140
aaaacacgtt cgggtgaac agtgcgcctc atggatcttc tgggagaagg actaaaacga 1200
tccatggaca agttgaagga aaaagaaaga gacaaggtct taactgcaga ggaattgaat 1260
gctgctcaga catccgttgc gtatggctgc atcaaatatg ctgaccttcc ccataaccgg 1320
tgtaatgact acatcttctc ctttgacaaa atgctagatg acagaggaaa tacagctgct 1380
tacttgttgt atgccttcac tagaatcagg tctattgcac gtctggccaa tattgatgaa 1440
gaaatgctcc aaaaagctgc tcgagaaacc aagattcttt tggatcatga gaaggaatgg 1500
aaactaggcc ggtgcatttt acggttcctt gagattctgc aaaagatttt agatgactta 1560
tttctccaca ctctctgtga ttatatatat gagctggcaa ctgcttccac agagttctat 1620
gatagctgct actgtgtgga gaaagataga cagactggaa aaatattgaa ggtgaacatg 1680
tggcgtatgc tgctatgtga agcagtagct gctgtcatgg ccaaggggtt tgatatcctg 1740
ggaataaaac ctgtccaaag gatgtaa 1767

```

```

<210> SEQ ID NO 45
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 45

```

```

Leu Gln Glu Val Phe Gly Cys Ala Ile Arg
1           5           10

```

```

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 46

```

```

Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Ile Val Thr Pro Ser
1           5           10           15

```

```

Gln Gln Pro Lys
           20

```

```

<210> SEQ ID NO 47
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

-continued

<400> SEQUENCE: 47

Phe Gly Asp Tyr Gln Cys Asn
1 5

<210> SEQ ID NO 48

<211> LENGTH: 77

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 48

Ser Ala Met Gly Ile Ser Gln Met Leu Lys Ala Lys Glu Gln Lys Val
1 5 10 15

Ser Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys His Leu Pro Asn Asn
20 25 30

Lys Tyr Ile Asp Lys Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val
35 40 45

His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val
50 55 60

Asn Gly Val Gln Leu Pro Val Leu Gly Asp Lys Glu Lys
65 70 75

<210> SEQ ID NO 49

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 49

Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys
1 5 10

<210> SEQ ID NO 50

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 50

Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met
1 5 10 15

Ser Arg

<210> SEQ ID NO 51

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 51

Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg
1 5 10

<210> SEQ ID NO 52

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 52

Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala
1 5 10 15

His Leu Gln Asp Lys
20

-continued

<210> SEQ ID NO 53
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 53

Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala
1 5 10 15

Phe Tyr Lys

<210> SEQ ID NO 54
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 54

Glu Ser Lys Lys Arg
1 5

<210> SEQ ID NO 55
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 55

Phe Asp Ala Asp Glu Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val Val
1 5 10 15

Leu Leu Gln Ser Lys
20

<210> SEQ ID NO 56
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 56

Asn Pro Asp Ile Met Lys
1 5

<210> SEQ ID NO 57
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 57

Ala Trp Asn Leu Ile Cys Asp Val Ser Arg
1 5 10

<210> SEQ ID NO 58
<211> LENGTH: 4
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 58

Glu Glu Phe Lys
1

<210> SEQ ID NO 59
<211> LENGTH: 13
<212> TYPE: PRT

-continued

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 59

Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg
1 5 10

<210> SEQ ID NO 60

<211> LENGTH: 29

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 60

Gly Glu Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe
1 5 10 15

Glu Asp Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys
20 25

<210> SEQ ID NO 61

<211> LENGTH: 31

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 61

Ile Val Phe Val Pro Gly Cys Ser Val Pro Leu Thr Ile Val Lys Ser
1 5 10 15

Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys
20 25 30

<210> SEQ ID NO 62

<211> LENGTH: 89

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 62

Gln Arg Leu Phe Glu Glu Lys Ala Asn Lys Ile Ile Tyr Val Val Asp
1 5 10 15

Asn Gly Gln Ala Ile His Phe Gln Thr Ile Phe Ala Ala Ala Gln Met
20 25 30

Ile Gly Trp Tyr Asp Pro Lys Val Thr Leu Val Thr His Val Gly Phe
35 40 45

Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr Arg Ser Gly
50 55 60

Glu Thr Val Arg Leu Met Asp Leu Leu Glu Glu Gly Leu Lys Arg Ser
65 70 75 80

Met Asp Lys Leu Lys Glu Lys Glu Arg
85

<210> SEQ ID NO 63

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 63

Asp Lys Val Leu Thr Glu Glu Glu Leu Lys
1 5 10

<210> SEQ ID NO 64

<211> LENGTH: 12

<212> TYPE: PRT

-continued

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 64

Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys
1 5 10

<210> SEQ ID NO 65

<211> LENGTH: 23

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 65

Tyr Ala Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe
1 5 10 15

Asp Lys Met Leu Asp Asp Arg
20

<210> SEQ ID NO 66

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 66

Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg
1 5 10

<210> SEQ ID NO 67

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 67

Ile Arg Ser Ile Ala Arg
1 5

<210> SEQ ID NO 68

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 68

Leu Ala Asn Ile Asp Glu Ala Met Leu Gln Arg
1 5 10

<210> SEQ ID NO 69

<211> LENGTH: 77

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 69

Ala Ala Arg Glu Thr Lys Ile Ile Leu Asp His Glu Lys Glu Trp Lys
1 5 10 15

Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu
20 25 30

Asp Asp Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala
35 40 45

Thr Thr Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp
50 55 60

Arg Gln Thr Gly Lys Val Leu Lys Val Asn Met Trp Arg
65 70 75

-continued

<210> SEQ ID NO 70
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 70

Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys
1 5 10

<210> SEQ ID NO 71
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 71

Gly Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg
1 5 10

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 72

Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Ile Val Thr Pro Ser
1 5 10 15

Gln Gln Pro Lys
20

<210> SEQ ID NO 73
<211> LENGTH: 59
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 73

Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu
1 5 10 15

Lys Ala Lys Glu Gln Lys Val Ser Pro Arg Glu Ile Ala Glu Asn Ile
20 25 30

Thr Lys His Leu Pro Asn Asn Lys Tyr Ile Asp Lys Val Glu Ile Ala
35 40 45

Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys
50 55

<210> SEQ ID NO 74
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 74

Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln
1 5 10 15

Leu Pro Val Leu Gly Asp Lys Glu Lys
20 25

<210> SEQ ID NO 75
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 75

Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys
1 5 10

<210> SEQ ID NO 76

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 76

Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met
1 5 10 15

Ser Arg

<210> SEQ ID NO 77

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 77

Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg
1 5 10

<210> SEQ ID NO 78

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 78

Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala
1 5 10 15

His Leu Gln Asp Lys
20

<210> SEQ ID NO 79

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 79

Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala
1 5 10 15

Phe Tyr Lys

<210> SEQ ID NO 80

<211> LENGTH: 32

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 80

Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu Phe Lys Lys Arg Ala
1 5 10 15

Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn Pro Asp Ile Met Lys
20 25 30

<210> SEQ ID NO 81

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 81

Ala Trp Asn Leu Ile Cys Asp Val Ser Arg
1 5 10

<210> SEQ ID NO 82

<211> LENGTH: 46

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 82

Glu Glu Phe Lys Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu
1 5 10 15

Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu
20 25 30

Phe Glu Asp Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys
35 40 45

<210> SEQ ID NO 83

<211> LENGTH: 31

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 83

Ile Val Phe Val Pro Gly Cys Ser Val Pro Leu Thr Ile Val Lys Ser
1 5 10 15

Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys
20 25 30

<210> SEQ ID NO 84

<211> LENGTH: 89

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 84

Gln Arg Leu Phe Glu Glu Lys Ala Asn Lys Ile Ile Tyr Val Val Asp
1 5 10 15

Asn Gly Gln Ala Ile His Phe Gln Thr Ile Phe Ala Ala Ala Gln Met
20 25 30

Ile Gly Trp Tyr Asp Pro Lys Val Thr Leu Val Thr His Val Gly Phe
35 40 45

Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr Arg Ser Gly
50 55 60

Glu Thr Val Arg Leu Met Asp Leu Leu Glu Glu Gly Leu Lys Arg Ser
65 70 75 80

Met Asp Lys Leu Lys Glu Lys Glu Arg
85

<210> SEQ ID NO 85

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 85

Asp Lys Val Leu Thr Glu Glu Glu Leu Lys
1 5 10

<210> SEQ ID NO 86

-continued

<211> LENGTH: 35
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 86

Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala Asp Leu
1 5 10 15

Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys Met Leu
20 25 30

Asp Asp Arg
35

<210> SEQ ID NO 87
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 87

Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg
1 5 10

<210> SEQ ID NO 88
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 88

Ile Arg Ser Ile Ala Arg
1 5

<210> SEQ ID NO 89
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 89

Leu Ala Asn Ile Asp Glu Ala Met Leu Gln Arg
1 5 10

<210> SEQ ID NO 90
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 90

Ala Ala Arg Glu Thr Lys
1 5

<210> SEQ ID NO 91
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 91

Ile Ile Leu Asp His Glu Lys Glu Trp Lys
1 5 10

<210> SEQ ID NO 92
<211> LENGTH: 61
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 92

Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu
1 5 10 15
Asp Asp Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala
20 25 30
Thr Thr Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp
35 40 45
Arg Gln Thr Gly Lys Val Leu Lys Val Asn Met Trp Arg
50 55 60

<210> SEQ ID NO 93

<211> LENGTH: 13

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 93

Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys
1 5 10

<210> SEQ ID NO 94

<211> LENGTH: 47

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 94

Leu Gln Glu Val Phe Gly Cys Ala Ile Arg Ala Ala Tyr Pro Asp Leu
1 5 10 15
Glu Asn Pro Pro Leu Ile Val Thr Pro Ser Gln Gln Pro Lys Phe Gly
20 25 30
Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys
35 40 45

<210> SEQ ID NO 95

<211> LENGTH: 23

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 95

Ala Lys Glu Gln Lys Val Ser Pro Arg Glu Ile Ala Glu Asn Ile Thr
1 5 10 15
Lys His Leu Pro Asn Asn Lys
20

<210> SEQ ID NO 96

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 96

Tyr Ile Asp Lys Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val His
1 5 10 15
Leu Arg

<210> SEQ ID NO 97

<211> LENGTH: 37

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 97

-continued

Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln
1 5 10 15

Leu Pro Val Leu Gly Asp Lys Glu Lys Val Ile Val Asp Phe Ser Ser
20 25 30

Pro Asn Ile Ala Lys
35

<210> SEQ ID NO 98
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 98

Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met
1 5 10 15

Ser Arg

<210> SEQ ID NO 99
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 99

Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg
1 5 10

<210> SEQ ID NO 100
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 100

Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala
1 5 10 15

His Leu Gln Asp Lys
20

<210> SEQ ID NO 101
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 101

Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala
1 5 10 15

Phe Tyr Lys

<210> SEQ ID NO 102
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 102

Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu Phe Lys Lys Arg
1 5 10 15

<210> SEQ ID NO 103
<211> LENGTH: 11
<212> TYPE: PRT

-continued

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 103

Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys
1 5 10

<210> SEQ ID NO 104

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 104

Asn Pro Asp Ile Met Lys
1 5

<210> SEQ ID NO 105

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 105

Ala Trp Asn Leu Ile Cys Asp Val Ser Arg
1 5 10

<210> SEQ ID NO 106

<211> LENGTH: 4

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 106

Glu Glu Phe Lys
1

<210> SEQ ID NO 107

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 107

Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg
1 5 10

<210> SEQ ID NO 108

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 108

Gly Glu Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys
1 5 10

<210> SEQ ID NO 109

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 109

Glu Phe Glu Asp Lys Gly Phe Val Gln Val Asp Asp Gly Arg
1 5 10

<210> SEQ ID NO 110

<211> LENGTH: 32

-continued

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 110

Lys Ile Val Phe Val Pro Gly Cys Ser Val Pro Leu Thr Ile Val Lys
1 5 10 15Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys
20 25 30

<210> SEQ ID NO 111

<211> LENGTH: 68

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 111

Gln Arg Leu Phe Glu Glu Lys Ala Asn Lys Ile Ile Tyr Val Val Asp
1 5 10 15Asn Gly Gln Ala Ile His Phe Gln Thr Ile Phe Ala Ala Ala Gln Met
20 25 30Ile Gly Trp Tyr Asp Pro Lys Val Thr Leu Val Thr His Val Gly Phe
35 40 45Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr Arg Ser Gly
50 55 60Glu Thr Val Arg
65

<210> SEQ ID NO 112

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 112

Leu Met Asp Leu Leu Glu Glu Gly Leu Lys Arg
1 5 10

<210> SEQ ID NO 113

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 113

Ser Met Asp Lys Leu Lys Glu Lys Glu Arg
1 5 10

<210> SEQ ID NO 114

<211> LENGTH: 10

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 114

Asp Lys Val Leu Thr Glu Glu Glu Leu Lys
1 5 10

<210> SEQ ID NO 115

<211> LENGTH: 12

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 115

Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys

-continued

1	5	10
---	---	----

<210> SEQ ID NO 116
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 116

Tyr Ala Asp Leu Ser His Asn Arg
1 5

<210> SEQ ID NO 117
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 117

Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys
1 5 10

<210> SEQ ID NO 118
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 118

Met Leu Asp Asp Arg
1 5

<210> SEQ ID NO 119
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 119

Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg
1 5 10

<210> SEQ ID NO 120
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 120

Ile Arg Ser Ile Ala Arg
1 5

<210> SEQ ID NO 121
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 121

Leu Ala Asn Ile Asp Glu Ala Met Leu Gln Arg
1 5 10

<210> SEQ ID NO 122
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 122

-continued

Ala Ala Arg Glu Thr Lys
1 5

<210> SEQ ID NO 123
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 123

Ile Ile Leu Asp His Glu Lys Glu Trp Lys
1 5 10

<210> SEQ ID NO 124
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 124

Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys
1 5 10

<210> SEQ ID NO 125
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 125

Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys
1 5 10

<210> SEQ ID NO 126
<211> LENGTH: 36
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 126

Asp Tyr Ile Tyr Glu Leu Ala Thr Thr Phe Thr Glu Phe Tyr Asp Ser
1 5 10 15

Cys Tyr Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Val Leu Lys Val
20 25 30

Asn Met Trp Arg
35

<210> SEQ ID NO 127
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 127

Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp
1 5 10 15

Ile Leu Gly Ile Lys Pro Val Gln Arg
20 25

<210> SEQ ID NO 128
<211> LENGTH: 580
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

<400> SEQUENCE: 128

Leu Gln Glu Val Phe Gly Cys Ala Ile Arg Ala Ala Tyr Pro Asp Leu

-continued

1	5	10	15
Glu Asn Pro Pro Leu Ile Val Thr Pro Ser Gln Gln Pro Lys Phe Gly	20	25	30
Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Ala	35	40	45
Lys Glu Gln Lys Val Ser Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys	50	55	60
His Leu Pro Asn Asn Lys Tyr Ile Asp Lys Val Glu Ile Ala Gly Pro	65	70	75
Gly Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu	85	90	95
Thr Ser Leu Leu Val Asn Gly Val Gln Leu Pro Val Leu Gly Asp Lys	100	105	110
Glu Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met	115	120	125
His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met Ser Arg	130	135	140
Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly	145	150	155
Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys	165	170	175
Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala	180	185	190
Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu Phe Lys	195	200	205
Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn Pro Asp	210	215	220
Ile Met Lys Ala Trp Asn Leu Ile Cys Asp Val Ser Arg Glu Glu Phe	225	230	235
Lys Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg Gly Glu	245	250	255
Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe Glu Asp	260	265	270
Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe Val Pro	275	280	285
Gly Cys Ser Val Pro Leu Thr Ile Val Lys Ser Asp Gly Gly Tyr Thr	290	295	300
Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys Gln Arg Leu Phe Glu Glu	305	310	315
Lys Ala Asn Lys Ile Ile Tyr Val Val Asp Asn Gly Gln Ala Ile His	325	330	335
Phe Gln Thr Ile Phe Ala Ala Ala Gln Met Ile Gly Trp Tyr Asp Pro	340	345	350
Lys Val Thr Leu Val Thr His Val Gly Phe Gly Val Val Leu Gly Glu	355	360	365
Asp Lys Lys Lys Phe Lys Thr Arg Ser Gly Glu Thr Val Arg Leu Met	370	375	380
Asp Leu Leu Glu Glu Gly Leu Lys Arg Ser Met Asp Lys Leu Lys Glu	385	390	395
Lys Glu Arg Asp Lys Val Leu Thr Glu Glu Glu Leu Lys Ala Ala Gln	405	410	415

-continued

Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala Asp Leu Ser His Asn
 420 425 430
 Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys Met Leu Asp Asp Arg
 435 440 445
 Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg Ile Arg Ser
 450 455 460
 Ile Ala Arg Leu Ala Asn Ile Asp Glu Ala Met Leu Gln Arg Ala Ala
 465 470 475 480
 Arg Glu Thr Lys Ile Ile Leu Asp His Glu Lys Glu Trp Lys Leu Gly
 485 490 495
 Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu Asp Asp
 500 505 510
 Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala Thr Thr
 515 520 525
 Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp Arg Gln
 530 535 540
 Thr Gly Lys Val Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys Glu
 545 550 555 560
 Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile Lys
 565 570 575
 Pro Val Gln Arg
 580

<210> SEQ ID NO 129

<211> LENGTH: 558

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 129

Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu Ile Val Thr Pro Ser
 1 5 10 15
 Gln Gln Pro Lys Phe Gly Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile
 20 25 30
 Ser Gln Met Leu Lys Ala Lys Glu Gln Lys Val Ser Pro Arg Glu Ile
 35 40 45
 Ala Glu Asn Ile Thr Lys His Leu Pro Asn Asn Lys Tyr Ile Asp Lys
 50 55 60
 Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val His Leu Arg Lys Asp
 65 70 75 80
 Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val Gln Leu
 85 90 95
 Pro Val Leu Gly Asp Lys Glu Lys Val Ile Val Asp Phe Ser Ser Pro
 100 105 110
 Asn Ile Ala Lys Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile
 115 120 125
 Gly Glu Ser Met Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu
 130 135 140
 Arg Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile
 145 150 155 160
 Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro
 165 170 175
 Ile Gly Asp Leu Gln Ala Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp

-continued

180					185					190				
Ala	Asp	Glu	Glu	Phe	Lys	Lys	Arg	Ala	Tyr	Gln	Cys	Val	Val	Leu
	195						200					205		Leu
Gln	Ser	Lys	Asn	Pro	Asp	Ile	Met	Lys	Ala	Trp	Asn	Leu	Ile	Cys
	210					215					220			Asp
Val	Ser	Arg	Glu	Glu	Phe	Lys	Lys	Ile	Tyr	Asp	Ala	Leu	Asp	Ile
	225				230					235				Thr
Leu	Ile	Glu	Arg	Gly	Glu	Ser	Phe	Tyr	Gln	Asp	Arg	Met	Lys	Asp
				245					250					255
Val	Lys	Glu	Phe	Glu	Asp	Lys	Gly	Phe	Val	Gln	Val	Asp	Asp	Gly
			260					265				270		Arg
Lys	Ile	Val	Phe	Val	Pro	Gly	Cys	Ser	Val	Pro	Leu	Thr	Ile	Val
		275					280					285		Lys
Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr	Ser	Asp	Leu	Ala	Ala	Ile
	290					295					300			Lys
Gln	Arg	Leu	Phe	Glu	Glu	Lys	Ala	Asn	Lys	Ile	Ile	Tyr	Val	Val
	305				310					315				Asp
Asn	Gly	Gln	Ala	Ile	His	Phe	Gln	Thr	Ile	Phe	Ala	Ala	Ala	Gln
				325					330					Met
Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val	Thr	Leu	Val	Thr	His	Val	Gly
			340					345					350	Phe
Gly	Val	Val	Leu	Gly	Glu	Asp	Lys	Lys	Lys	Phe	Lys	Thr	Arg	Ser
		355					360					365		Gly
Glu	Thr	Val	Arg	Leu	Met	Asp	Leu	Leu	Glu	Glu	Gly	Leu	Lys	Arg
	370					375					380			Ser
Met	Asp	Lys	Leu	Lys	Glu	Lys	Glu	Arg	Asp	Lys	Val	Leu	Thr	Glu
	385				390					395				Glu
Glu	Leu	Lys	Ala	Ala	Gln	Thr	Ser	Val	Ala	Tyr	Gly	Cys	Ile	Lys
			405						410					415
Ala	Asp	Leu	Ser	His	Asn	Arg	Leu	Asn	Asp	Tyr	Ile	Phe	Ser	Phe
		420						425					430	Asp
Lys	Met	Leu	Asp	Asp	Arg	Gly	Asn	Thr	Ala	Ala	Tyr	Leu	Leu	Tyr
		435					440					445		Ala
Phe	Thr	Arg	Ile	Arg	Ser	Ile	Ala	Arg	Leu	Ala	Asn	Ile	Asp	Glu
	450					455					460			Ala
Met	Leu	Gln	Arg	Ala	Ala	Arg	Glu	Thr	Lys	Ile	Ile	Leu	Asp	His
	465				470					475				Glu
Lys	Glu	Trp	Lys	Leu	Gly	Arg	Cys	Ile	Leu	Arg	Phe	Pro	Glu	Ile
			485					490						Leu
Gln	Lys	Ile	Leu	Asp	Asp	Leu	Phe	Leu	His	Thr	Leu	Cys	Asp	Tyr
		500						505					510	Ile
Tyr	Glu	Leu	Ala	Thr	Thr	Phe	Thr	Glu	Phe	Tyr	Asp	Ser	Cys	Tyr
		515					520					525		Cys
Val	Glu	Lys	Asp	Arg	Gln	Thr	Gly	Lys	Val	Leu	Lys	Val	Asn	Met
	530					535					540			Trp
Arg	Met	Leu	Leu	Cys	Glu	Ala	Val	Ala	Ala	Val	Met	Ala	Lys	
	545				550					555				

<210> SEQ ID NO 130

<211> LENGTH: 580

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

-continued

<400> SEQUENCE: 130

```

Leu Gln Glu Val Phe Gly Cys Ala Ile Arg Ala Ala Tyr Pro Asp Leu
1          5          10          15
Glu Asn Pro Pro Leu Ile Val Thr Pro Ser Gln Gln Pro Lys Phe Gly
20          25          30
Asp Tyr Gln Cys Asn Ser Ala Met Gly Ile Ser Gln Met Leu Lys Ala
35          40          45
Lys Glu Gln Lys Val Ser Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys
50          55          60
His Leu Pro Asn Asn Lys Tyr Ile Asp Lys Val Glu Ile Ala Gly Pro
65          70          75          80
Gly Phe Ile Asn Val His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu
85          90          95
Thr Ser Leu Leu Val Asn Gly Val Gln Leu Pro Val Leu Gly Asp Lys
100          105          110
Glu Lys Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met
115          120          125
His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met Ser Arg
130          135          140
Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly
145          150          155          160
Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys
165          170          175
Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala
180          185          190
Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu Phe Lys
195          200          205
Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn Pro Asp
210          215          220
Ile Met Lys Ala Trp Asn Leu Ile Cys Asp Val Ser Arg Glu Glu Phe
225          230          235          240
Lys Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg Gly Glu
245          250          255
Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe Glu Asp
260          265          270
Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe Val Pro
275          280          285
Gly Cys Ser Val Pro Leu Thr Ile Val Lys Ser Asp Gly Gly Tyr Thr
290          295          300
Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys Gln Arg Leu Phe Glu Glu
305          310          315          320
Lys Ala Asn Lys Ile Ile Tyr Val Val Asp Asn Gly Gln Ala Ile His
325          330          335
Phe Gln Thr Ile Phe Ala Ala Ala Gln Met Ile Gly Trp Tyr Asp Pro
340          345          350
Lys Val Thr Leu Val Thr His Val Gly Phe Gly Val Val Leu Gly Glu
355          360          365
Asp Lys Lys Lys Phe Lys Thr Arg Ser Gly Glu Thr Val Arg Leu Met
370          375          380
Asp Leu Leu Glu Glu Gly Leu Lys Arg Ser Met Asp Lys Leu Lys Glu

```

-continued

385	390	395	400
Lys Glu Arg Asp	Lys Val Leu Thr Glu Glu Glu Leu Lys Ala Ala Gln		
	405	410	415
Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala Asp Leu Ser His Asn			
	420	425	430
Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys Met Leu Asp Asp Arg			
	435	440	445
Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg Ile Arg Ser			
	450	455	460
Ile Ala Arg Leu Ala Asn Ile Asp Glu Ala Met Leu Gln Arg Ala Ala			
	465	470	475
Arg Glu Thr Lys Ile Ile Leu Asp His Glu Lys Glu Trp Lys Leu Gly			
	485	490	495
Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu Asp Asp			
	500	505	510
Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala Thr Thr			
	515	520	525
Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp Arg Gln			
	530	535	540
Thr Gly Lys Val Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys Glu			
	545	550	555
Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile Lys			
	565	570	575
Pro Val Gln Arg			
	580		

<210> SEQ ID NO 131

<211> LENGTH: 198

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 131

Met Asp Leu Leu Gly Glu Gly Leu Lys Arg Ser Met Asp Lys Leu Lys			
1	5	10	15
Glu Lys Glu Arg Asp Lys Val Leu Thr Ala Glu Glu Leu Asn Ala Ala			
	20	25	30
Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala Asp Leu Ser His			
	35	40	45
Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys Met Leu Asp Asp			
	50	55	60
Arg Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg Ile Arg			
	65	70	75
Ser Ile Ala Arg Leu Ala Asn Ile Asp Glu Glu Met Leu Gln Lys Ala			
	85	90	95
Ala Arg Glu Thr Lys Ile Leu Leu Asp His Glu Lys Glu Trp Lys Leu			
	100	105	110
Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu Asp			
	115	120	125
Asp Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala Thr			
	130	135	140
Ala Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp Arg			
	145	150	155
			160

-continued

Gln Thr Gly Lys Ile Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys
 165 170 175

Glu Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile
 180 185 190

Lys Pro Val Gln Arg Met
 195

<210> SEQ ID NO 132
 <211> LENGTH: 597
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 132

```

atggatcttc tgggagaagg actaaaacga tccatggaca agttgaagga aaaagaaaga      60
gacaagggtct taactgcaga ggaattgaat gctgctcaga catccgttgc gtatggctgc      120
atcaaatatg ctgacctttc ccataaccgg ttgaatgact acatcttctc ctttgacaaa      180
atgcttagatg acagagggaaa tacagctgct tacttggtgt atgccttcac tagaatcagg      240
tctattgcac gtctggccaa tattgatgaa gaaatgctcc aaaaagctgc tcgagaaacc      300
aagattcttt tggatcatga gaaggaatgg aaactaggcc ggtgcatttt acggttcct      360
gagattctgc aaaagatttt agatgactta tttctccaca ctctctgtga ttatatatat      420
gagctggcaa ctgctttcac agagttctat gatagctgct actgtgtgga gaaagataga      480
cagactggaa aaatattgaa ggtgaacatg tggcgtatgc tgctatgtga agcagtagct      540
gctgtcatgg ccaaggggtt tgatatcctg ggaataaaac ctgtccaaag gatgtaa      597

```

<210> SEQ ID NO 133
 <211> LENGTH: 414
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 133

```

Met Leu Ile Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val
1           5           10           15

Ser Pro Pro Ile Gly Asp Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys
20          25          30

Arg Phe Asp Thr Glu Glu Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val
35          40          45

Val Leu Leu Gln Gly Lys Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu
50          55          60

Ile Cys Asp Val Ser Arg Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu
65          70          75          80

Asp Val Ser Leu Ile Glu Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met
85          90          95

Asn Asp Ile Val Lys Glu Phe Glu Asp Arg Gly Phe Val Gln Val Asp
100         105         110

Asp Gly Arg Lys Ile Val Phe Val Pro Gly Cys Ser Ile Pro Leu Thr
115         120         125

Ile Val Lys Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala
130         135         140

Ala Ile Lys Gln Arg Leu Phe Glu Glu Lys Ala Asp Met Ile Ile Tyr
145         150         155         160

Val Val Asp Asn Gly Gln Ser Val His Phe Gln Thr Ile Phe Ala Ala

```

-continued

165								170				175			
Ala	Gln	Met	Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val	Thr	Arg	Val	Phe	His
			180					185					190		
Ala	Gly	Phe	Gly	Val	Val	Leu	Gly	Glu	Asp	Lys	Lys	Lys	Phe	Lys	Thr
			195				200					205			
Arg	Ser	Gly	Glu	Thr	Val	Arg	Leu	Met	Asp	Leu	Leu	Gly	Glu	Gly	Leu
	210					215					220				
Lys	Arg	Ser	Met	Asp	Lys	Leu	Lys	Glu	Lys	Glu	Arg	Asp	Lys	Val	Leu
225					230					235					240
Thr	Ala	Glu	Glu	Leu	Asn	Ala	Ala	Gln	Thr	Ser	Val	Ala	Tyr	Gly	Cys
				245					250					255	
Ile	Lys	Tyr	Ala	Asp	Leu	Ser	His	Asn	Arg	Leu	Asn	Asp	Tyr	Ile	Phe
			260					265					270		
Ser	Phe	Asp	Lys	Met	Leu	Asp	Asp	Arg	Gly	Asn	Thr	Ala	Ala	Tyr	Leu
		275					280					285			
Leu	Tyr	Ala	Phe	Thr	Arg	Ile	Arg	Ser	Ile	Ala	Arg	Leu	Ala	Asn	Ile
	290					295					300				
Asp	Glu	Glu	Met	Leu	Gln	Lys	Ala	Ala	Arg	Glu	Thr	Lys	Ile	Leu	Leu
305					310					315					320
Asp	His	Glu	Lys	Glu	Trp	Lys	Leu	Gly	Arg	Cys	Ile	Leu	Arg	Phe	Pro
			325						330					335	
Glu	Ile	Leu	Gln	Lys	Ile	Leu	Asp	Asp	Leu	Phe	Leu	His	Thr	Leu	Cys
		340						345					350		
Asp	Tyr	Ile	Tyr	Glu	Leu	Ala	Thr	Ala	Phe	Thr	Glu	Phe	Tyr	Asp	Ser
		355					360					365			
Cys	Tyr	Cys	Val	Glu	Lys	Asp	Arg	Gln	Thr	Gly	Lys	Ile	Leu	Lys	Val
	370					375					380				
Asn	Met	Trp	Arg	Met	Leu	Leu	Cys	Glu	Ala	Val	Ala	Ala	Val	Met	Ala
385					390					395					400
Lys	Gly	Phe	Asp	Ile	Leu	Gly	Ile	Lys	Pro	Val	Gln	Arg	Met		
			405						410						

<210> SEQ ID NO 134

<211> LENGTH: 1245

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 134

```

atgctcatcg ctcacctgca agacaaattt ccagattatc taacagtttc acctcctatt    60
ggggatcttc aggtctttta taaggaatct aagaagaggt ttgatactga ggaggaattt    120
aagaagcgag catatcagtg tgtagttctg ctccagggta aaaaccgaga tattacaaaa    180
gcttggaagc ttatctgtga tgtctccgc caagagttaa ataaaatcta tgatgcattg    240
gacgtctctt taatagagag aggggaatcc ttctatcaag ataggatgaa tgatattgta    300
aaggaaattg aagatagagg atttgtgcag gtggatgatg gcagaaagat tgtatttgtc    360
ccagggtgtt ccataccatt aaccatagta aaatcagatg gaggttatac ctatgataca    420
tctgacctgg ctgctattaa acaagacta ttgaggaaa aagcagatat gattatctat    480
gttgtggaca atggacaatc tgtgcacttc cagacaatat ttgctgctgc tcaaagtatt    540
ggttggtatg accctaaagt aactcgagtc ttccatgctg gatttggtgt ggtgctaggg    600
gaagacaaga aaaagtttaa aacacgttcg ggtgaaacag tgcgcctcat ggatcttctg    660

```

-continued

```

ggagaaggac taaaacgac catggacaag ttgaaggaaa aagaaagaga caaggtotta 720
actgcagagg aattgaatgc tgctcagaca tccgttgctg atggetgcat caaatatgct 780
gacctttccc ataaccggtt gaatgactac atcttctcct ttgacaaaat gctagatgac 840
agaggaaata cagctgctta cttgtgtgat gccttcaact gaatcaggtc tattgcacgt 900
ctggccaata ttgatgaaga aatgctccaa aaagctgctc gagaaaccaa gattcttttg 960
gatcatgaga aggaatggaa actaggcccg tgcattttac ggttccctga gattctgcaa 1020
aagatttttag atgacttatt tctccacact ctctgtgatt atatatatga gctggcaact 1080
gctttcacag agttctatga tagctgctac tgtgtggaga aagatagaca gactggaaaa 1140
atattgaagg tgaacatgtg gcgtatgctg ctatgtgaag cagtagctgc tgcatggcc 1200
aaggggtttg atatcctggg aataaaacct gtccaaagga tgtaa 1245

```

```

<210> SEQ ID NO 135
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 135

```

```

attgtaaagg aatttgaaga tagaggaaaa agtttaaac acgttcgggt 50

```

```

<210> SEQ ID NO 136
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 136

```

```

gaaaaagttg aaattgctgg tcctgggttaa atcatgtagg agactggggg 50

```

```

<210> SEQ ID NO 137
<211> LENGTH: 110
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 137

```

```

Asp Glu Glu Met Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu Leu
1           5           10          15
Asp His Glu Lys Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe Pro
20          25          30
Glu Ile Leu Gln Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys
35          40          45
Asp Tyr Ile Tyr Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp Ser
50          55          60
Cys Tyr Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys Val
65          70          75          80
Asn Met Trp Arg Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala
85          90          95
Lys Gly Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg Met
100         105         110

```

```

<210> SEQ ID NO 138
<211> LENGTH: 333
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

-continued

<400> SEQUENCE: 138

```

gatgaagaaa tgctccaaaa agctgctcga gaaaccaaga ttcttttggga tcatgagaag    60
gaatggaaac taggccggtg cattttacgg ttccctgaga ttctgcaaaa gatttttagat    120
gacttatttc tccacactct ctgtgattat atatatgagc tggcaactgc ttccacagag    180
ttctatgata gctgctactg tgtggagaaa gatagacaga ctggaaaaat attgaaggtg    240
aacatgtggc gtatgtctgt atgtgaagca gtagctgctg tcatggccaa ggggtttgat    300
atcctgggaa taaaacctgt ccaaaggatg taa                                333

```

<210> SEQ ID NO 139

<211> LENGTH: 608

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
N-terminal 6xHis affinity tag

<400> SEQUENCE: 139

```

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
 1             5             10             15
Gly Leu Asp Ser Thr Ile Asn Ile Ile Ser Arg Leu Gln Glu Val Phe
 20             25             30
Gly His Ala Ile Lys Ala Ala Tyr Pro Asp Leu Glu Asn Pro Pro Leu
 35             40             45
Leu Val Thr Pro Ser Gln Gln Ala Lys Phe Gly Asp Tyr Gln Cys Asn
 50             55             60
Ser Ala Met Gly Ile Ser Gln Met Leu Lys Thr Lys Glu Gln Lys Val
 65             70             75             80
Asn Pro Arg Glu Ile Ala Glu Asn Ile Thr Lys His Leu Pro Asp Asn
 85             90             95
Glu Cys Ile Glu Lys Val Glu Ile Ala Gly Pro Gly Phe Ile Asn Val
100             105             110
His Leu Arg Lys Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val
115             120             125
Asn Gly Val Gln Leu Pro Ala Leu Gly Glu Asn Lys Lys Val Ile Val
130             135             140
Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met His Val Gly His Leu
145             150             155             160
Arg Ser Thr Ile Ile Gly Glu Ser Ile Ser Arg Leu Phe Glu Phe Ala
165             170             175
Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly Asp Trp Gly Thr Gln
180             185             190
Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu
195             200             205
Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Val Phe Tyr Lys Glu Ser
210             215             220
Lys Lys Arg Phe Asp Thr Glu Glu Glu Phe Lys Lys Arg Ala Tyr Gln
225             230             235             240
Cys Val Val Leu Leu Gln Gly Lys Asn Pro Asp Ile Thr Lys Ala Trp
245             250             255
Lys Leu Ile Cys Asp Val Ser Arg Gln Glu Leu Asn Lys Ile Tyr Asp
260             265             270

```

-continued

Ala	Leu	Asp	Val	Ser	Leu	Ile	Glu	Arg	Gly	Glu	Ser	Phe	Tyr	Gln	Asp	275	280	285
Arg	Met	Asn	Asp	Ile	Val	Lys	Glu	Phe	Glu	Asp	Arg	Gly	Phe	Val	Gln	290	295	300
Val	Asp	Asp	Gly	Arg	Lys	Ile	Val	Phe	Val	Pro	Gly	Cys	Ser	Ile	Pro	305	310	315
Leu	Thr	Ile	Val	Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr	Ser	Asp	325	330	335
Leu	Ala	Ala	Ile	Lys	Gln	Arg	Leu	Phe	Glu	Glu	Lys	Ala	Asp	Met	Ile	340	345	350
Ile	Tyr	Val	Val	Asp	Asn	Gly	Gln	Ser	Val	His	Phe	Gln	Thr	Ile	Phe	355	360	365
Ala	Ala	Ala	Gln	Met	Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val	Thr	Arg	Val	370	375	380
Phe	His	Ala	Gly	Phe	Gly	Val	Val	Leu	Gly	Glu	Asp	Lys	Lys	Lys	Phe	385	390	395
Lys	Thr	Arg	Ser	Gly	Glu	Thr	Val	Arg	Leu	Met	Asp	Leu	Leu	Gly	Glu	405	410	415
Gly	Leu	Lys	Arg	Ser	Met	Asp	Lys	Leu	Lys	Glu	Lys	Glu	Arg	Asp	Lys	420	425	430
Val	Leu	Thr	Ala	Glu	Glu	Leu	Asn	Ala	Ala	Gln	Thr	Ser	Val	Ala	Tyr	435	440	445
Gly	Cys	Ile	Lys	Tyr	Ala	Asp	Leu	Ser	His	Asn	Arg	Leu	Asn	Asp	Tyr	450	455	460
Ile	Phe	Ser	Phe	Asp	Lys	Met	Leu	Asp	Asp	Arg	Gly	Asn	Thr	Ala	Ala	465	470	475
Tyr	Leu	Leu	Tyr	Ala	Phe	Thr	Arg	Ile	Arg	Ser	Ile	Ala	Arg	Leu	Ala	485	490	495
Asn	Ile	Asp	Glu	Glu	Met	Leu	Gln	Lys	Ala	Ala	Arg	Glu	Thr	Lys	Ile	500	505	510
Leu	Leu	Asp	His	Glu	Lys	Glu	Trp	Lys	Leu	Gly	Arg	Cys	Ile	Leu	Arg	515	520	525
Phe	Pro	Glu	Ile	Leu	Gln	Lys	Ile	Leu	Asp	Asp	Leu	Phe	Leu	His	Thr	530	535	540
Leu	Cys	Asp	Tyr	Ile	Tyr	Glu	Leu	Ala	Thr	Ala	Phe	Thr	Glu	Phe	Tyr	545	550	555
Asp	Ser	Cys	Tyr	Cys	Val	Glu	Lys	Asp	Arg	Gln	Thr	Gly	Lys	Ile	Leu	565	570	575
Lys	Val	Asn	Met	Trp	Arg	Met	Leu	Leu	Cys	Glu	Ala	Val	Ala	Ala	Val	580	585	590
Met	Ala	Lys	Gly	Phe	Asp	Ile	Leu	Gly	Ile	Lys	Pro	Val	Gln	Arg	Met	595	600	605

<210> SEQ ID NO 140
 <211> LENGTH: 608
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
 C-terminal 6xHis affinity tag

<400> SEQUENCE: 140

Met	Ile	Asn	Ile	Ile	Ser	Arg	Leu	Gln	Glu	Val	Phe	Gly	His	Ala	Ile
1			5						10					15	

Lys	Ala	Ala	Tyr	Pro	Asp	Leu	Glu	Asn	Pro	Pro	Leu	Leu	Val	Thr	Pro	
			20					25					30			
Ser	Gln	Gln	Ala	Lys	Phe	Gly	Asp	Tyr	Gln	Cys	Asn	Ser	Ala	Met	Gly	
		35					40					45				
Ile	Ser	Gln	Met	Leu	Lys	Thr	Lys	Glu	Gln	Lys	Val	Asn	Pro	Arg	Glu	
		50				55					60					
Ile	Ala	Glu	Asn	Ile	Thr	Lys	His	Leu	Pro	Asp	Asn	Glu	Cys	Ile	Glu	
65					70					75					80	
Lys	Val	Glu	Ile	Ala	Gly	Pro	Gly	Phe	Ile	Asn	Val	His	Leu	Arg	Lys	
				85					90					95		
Asp	Phe	Val	Ser	Glu	Gln	Leu	Thr	Ser	Leu	Leu	Val	Asn	Gly	Val	Gln	
			100					105					110			
Leu	Pro	Ala	Leu	Gly	Glu	Asn	Lys	Lys	Val	Ile	Val	Asp	Phe	Ser	Ser	
		115					120					125				
Pro	Asn	Ile	Ala	Lys	Glu	Met	His	Val	Gly	His	Leu	Arg	Ser	Thr	Ile	
		130				135					140					
Ile	Gly	Glu	Ser	Ile	Ser	Arg	Leu	Phe	Glu	Phe	Ala	Gly	Tyr	Asp	Val	
145					150					155					160	
Leu	Arg	Leu	Asn	His	Val	Gly	Asp	Trp	Gly	Thr	Gln	Phe	Gly	Met	Leu	
			165					170						175		
Ile	Ala	His	Leu	Gln	Asp	Lys	Phe	Pro	Asp	Tyr	Leu	Thr	Val	Ser	Pro	
			180					185					190			
Pro	Ile	Gly	Asp	Leu	Gln	Val	Phe	Tyr	Lys	Glu	Ser	Lys	Lys	Arg	Phe	
		195				200						205				
Asp	Thr	Glu	Glu	Glu	Phe	Lys	Lys	Arg	Ala	Tyr	Gln	Cys	Val	Val	Leu	
		210				215					220					
Leu	Gln	Gly	Lys	Asn	Pro	Asp	Ile	Thr	Lys	Ala	Trp	Lys	Leu	Ile	Cys	
225				230						235					240	
Asp	Val	Ser	Arg	Gln	Glu	Leu	Asn	Lys	Ile	Tyr	Asp	Ala	Leu	Asp	Val	
			245						250					255		
Ser	Leu	Ile	Glu	Arg	Gly	Glu	Ser	Phe	Tyr	Gln	Asp	Arg	Met	Asn	Asp	
			260					265					270			
Ile	Val	Lys	Glu	Phe	Glu	Asp	Arg	Gly	Phe	Val	Gln	Val	Asp	Asp	Gly	
		275				280						285				
Arg	Lys	Ile	Val	Phe	Val	Pro	Gly	Cys	Ser	Ile	Pro	Leu	Thr	Ile	Val	
		290				295					300					
Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr	Ser	Asp	Leu	Ala	Ala	Ile	
305					310					315					320	
Lys	Gln	Arg	Leu	Phe	Glu	Glu	Lys	Ala	Asp	Met	Ile	Ile	Tyr	Val	Val	
			325						330					335		
Asp	Asn	Gly	Gln	Ser	Val	His	Phe	Gln	Thr	Ile	Phe	Ala	Ala	Ala	Gln	
			340					345					350			
Met	Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val								

-continued

Glu Glu Leu Asn Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys
 420 425 430
 Tyr Ala Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe
 435 440 445
 Asp Lys Met Leu Asp Asp Arg Gly Asn Thr Ala Ala Tyr Leu Leu Tyr
 450 455 460
 Ala Phe Thr Arg Ile Arg Ser Ile Ala Arg Leu Ala Asn Ile Asp Glu
 465 470 475 480
 Glu Met Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu Leu Asp His
 485 490 495
 Glu Lys Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile
 500 505 510
 Leu Gln Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys Asp Tyr
 515 520 525
 Ile Tyr Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp Ser Cys Tyr
 530 535 540
 Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys Val Asn Met
 545 550 555 560
 Trp Arg Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys Gly
 565 570 575
 Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg Met Gly Lys Pro Ile
 580 585 590
 Pro Asn Pro Leu Leu Gly Leu Asp Ser Thr His His His His His His
 595 600 605

<210> SEQ ID NO 141
 <211> LENGTH: 131
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
 N-terminal 6xHis affinity tag

<400> SEQUENCE: 141

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
 1 5 10 15
 Gly Leu Asp Ser Thr Asp Glu Glu Met Leu Gln Lys Ala Ala Arg Glu
 20 25 30
 Thr Lys Ile Leu Leu Asp His Glu Lys Glu Trp Lys Leu Gly Arg Cys
 35 40 45
 Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu Asp Asp Leu Phe
 50 55 60
 Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala Thr Ala Phe Thr
 65 70 75 80
 Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp Arg Gln Thr Gly
 85 90 95
 Lys Ile Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys Glu Ala Val
 100 105 110
 Ala Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile Lys Pro Val
 115 120 125
 Gln Arg Met
 130

<210> SEQ ID NO 142

-continued

<211> LENGTH: 131
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
C-terminal 6xHis affinity tag

<400> SEQUENCE: 142

Met Asp Glu Glu Met Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu
1 5 10 15
Leu Asp His Glu Lys Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe
20 25 30
Pro Glu Ile Leu Gln Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu
35 40 45
Cys Asp Tyr Ile Tyr Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp
50 55 60
Ser Cys Tyr Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys
65 70 75 80
Val Asn Met Trp Arg Met Leu Leu Cys Glu Ala Val Ala Ala Val Met
85 90 95
Ala Lys Gly Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg Met Gly
100 105 110
Lys Pro Ile Pro Asn Pro Leu Leu Gly Leu Asp Ser Thr His His His
115 120 125
His His His
130

<210> SEQ ID NO 143
<211> LENGTH: 218
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
N-terminal 6xHis affinity tag

<400> SEQUENCE: 143

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
1 5 10 15
Gly Leu Asp Ser Thr Asp Leu Leu Gly Glu Gly Leu Lys Arg Ser Met
20 25 30
Asp Lys Leu Lys Glu Lys Glu Arg Asp Lys Val Leu Thr Ala Glu Glu
35 40 45
Leu Asn Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala
50 55 60
Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys
65 70 75 80
Met Leu Asp Asp Arg Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe
85 90 95
Thr Arg Ile Arg Ser Ile Ala Arg Leu Ala Asn Ile Asp Glu Glu Met
100 105 110
Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu Leu Asp His Glu Lys
115 120 125
Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln
130 135 140
Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr
145 150 155 160

-continued

Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val
 165 170 175

Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys Val Asn Met Trp Arg
 180 185 190

Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp
 195 200 205

Ile Leu Gly Ile Lys Pro Val Gln Arg Met
 210 215

<210> SEQ ID NO 144
 <211> LENGTH: 218
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
 C-terminal 6xHis affinity tag

<400> SEQUENCE: 144

Met Asp Leu Leu Gly Glu Gly Leu Lys Arg Ser Met Asp Lys Leu Lys
 1 5 10 15

Glu Lys Glu Arg Asp Lys Val Leu Thr Ala Glu Glu Leu Asn Ala Ala
 20 25 30

Gln Thr Ser Val Ala Tyr Gly Cys Ile Lys Tyr Ala Asp Leu Ser His
 35 40 45

Asn Arg Leu Asn Asp Tyr Ile Phe Ser Phe Asp Lys Met Leu Asp Asp
 50 55 60

Arg Gly Asn Thr Ala Ala Tyr Leu Leu Tyr Ala Phe Thr Arg Ile Arg
 65 70 75 80

Ser Ile Ala Arg Leu Ala Asn Ile Asp Glu Glu Met Leu Gln Lys Ala
 85 90 95

Ala Arg Glu Thr Lys Ile Leu Leu Asp His Glu Lys Glu Trp Lys Leu
 100 105 110

Gly Arg Cys Ile Leu Arg Phe Pro Glu Ile Leu Gln Lys Ile Leu Asp
 115 120 125

Asp Leu Phe Leu His Thr Leu Cys Asp Tyr Ile Tyr Glu Leu Ala Thr
 130 135 140

Ala Phe Thr Glu Phe Tyr Asp Ser Cys Tyr Cys Val Glu Lys Asp Arg
 145 150 155 160

Gln Thr Gly Lys Ile Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys
 165 170 175

Glu Ala Val Ala Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile
 180 185 190

Lys Pro Val Gln Arg Met Gly Lys Pro Ile Pro Asn Pro Leu Leu Gly
 195 200 205

Leu Asp Ser Thr His His His His His His
 210 215

<210> SEQ ID NO 145
 <211> LENGTH: 434
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
 N-terminal 6xHis affinity tag

<400> SEQUENCE: 145

Met 1	His	His	His	His 5	His	His	Gly	Lys	Pro 10	Ile	Pro	Asn	Pro	Leu 15	Leu
Gly	Leu	Asp	Ser 20	Thr	Leu	Ile	Ala	His 25	Leu	Gln	Asp	Lys 30	Phe	Pro	Asp
Tyr	Leu	Thr	Val 35	Ser	Pro	Pro	Ile 40	Gly	Asp	Leu	Gln	Val 45	Phe	Tyr	Lys
Glu	Ser	Lys	Lys	Arg	Phe	Asp 55	Thr	Glu	Glu	Glu	Phe 60	Lys	Lys	Arg	Ala
Tyr 65	Gln	Cys	Val	Val	Leu 70	Leu	Gln	Gly	Lys	Asn 75	Pro	Asp	Ile	Thr	Lys 80
Ala	Trp	Lys	Leu	Ile 85	Cys	Asp	Val	Ser	Arg 90	Gln	Glu	Leu	Asn	Lys 95	Ile
Tyr	Asp	Ala	Leu 100	Asp	Val	Ser	Leu	Ile 105	Glu	Arg	Gly	Glu	Ser 110	Phe	Tyr
Gln	Asp	Arg	Met 115	Asn	Asp	Ile	Val 120	Lys	Glu	Phe	Glu	Asp 125	Arg	Gly	Phe
Val	Gln	Val	Asp 130	Asp	Gly 135	Arg	Lys	Ile	Val	Phe 140	Val	Pro	Gly	Cys	Ser
Ile 145	Pro	Leu	Thr	Ile	Val 150	Lys	Ser	Asp	Gly	Gly 155	Tyr	Thr	Tyr	Asp	Thr 160
Ser	Asp	Leu	Ala 165	Ala	Ile	Lys	Gln	Arg	Leu 170	Phe	Glu	Glu	Lys	Ala 175	Asp
Met	Ile	Ile	Tyr 180	Val	Val	Asp	Asn	Gly 185	Gln	Ser	Val	His 190	Phe	Gln	Thr
Ile	Phe	Ala 195	Ala	Ala	Gln	Met	Ile 200	Gly	Trp	Tyr	Asp	Pro 205	Lys	Val	Thr
Arg	Val	Phe 210	His	Ala	Gly 215	Phe	Gly	Val	Val	Leu	Gly 220	Glu	Asp	Lys	Lys
Lys 225	Phe	Lys	Thr	Arg	Ser 230	Gly	Glu	Thr	Val	Arg 235	Leu	Met	Asp	Leu	Leu 240
Gly	Glu	Gly	Leu 245	Lys	Arg	Ser	Met	Asp	Lys 250	Leu	Lys	Glu	Lys	Glu	Arg 255
Asp	Lys	Val	Leu 260	Thr	Ala	Glu	Glu	Leu 265	Asn	Ala	Ala	Gln	Thr 270	Ser	Val
Ala	Tyr	Gly 275	Cys	Ile	Lys	Tyr	Ala 280	Asp	Leu	Ser	His 285	Asn	Arg	Leu	Asn
Asp	Tyr 290	Ile	Phe	Ser	Phe	Asp 295	Lys	Met	Leu	Asp	Asp 300	Arg	Gly	Asn	Thr
Ala 305	Ala	Tyr	Leu	Leu	Tyr 310	Ala	Phe	Thr	Arg	Ile 315	Arg	Ser	Ile	Ala	Arg 320
Leu	Ala	Asn	Ile 325	Asp	Glu	Glu	Met	Leu 330	Gln	Lys	Ala	Ala	Arg	Glu	Thr 335
Lys	Ile	Leu	Leu 340	Asp	His	Glu	Lys	Glu 345	Trp	Lys	Leu	Gly	Arg 350	Cys	Ile
Leu	Arg	Phe 355	Pro	Glu	Ile	Leu	Gln 360	Lys	Ile	Leu	Asp	Asp 365	Leu	Phe	Leu
His 370	Thr	Leu	Cys	Asp	Tyr 375	Ile	Tyr	Glu	Leu	Ala 380	Thr	Ala	Phe	Thr	Glu
Phe 385	Tyr	Asp	Ser	Cys	Tyr 390	Cys	Val	Glu	Lys	Asp 395	Arg	Gln	Thr	Gly	Lys 400

-continued

Ile Leu Lys Val Asn Met Trp Arg Met Leu Leu Cys Glu Ala Val Ala
405 410 415

Ala Val Met Ala Lys Gly Phe Asp Ile Leu Gly Ile Lys Pro Val Gln
420 425 430

Arg Met

<210> SEQ ID NO 146

<211> LENGTH: 434

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
C-terminal 6xHis affinity tag

<400> SEQUENCE: 146

Met Leu Ile Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val
1 5 10 15

Ser Pro Pro Ile Gly Asp Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys
20 25 30

Arg Phe Asp Thr Glu Glu Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val
35 40 45

Val Leu Leu Gln Gly Lys Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu
50 55 60

Ile Cys Asp Val Ser Arg Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu
65 70 75 80

Asp Val Ser Leu Ile Glu Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met
85 90 95

Asn Asp Ile Val Lys Glu Phe Glu Asp Arg Gly Phe Val Gln Val Asp
100 105 110

Asp Gly Arg Lys Ile Val Phe Val Pro Gly Cys Ser Ile Pro Leu Thr
115 120 125

Ile Val Lys Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala
130 135 140

Ala Ile Lys Gln Arg Leu Phe Glu Glu Lys Ala Asp Met Ile Ile Tyr
145 150 155 160

Val Val Asp Asn Gly Gln Ser Val His Phe Gln Thr Ile Phe Ala Ala
165 170 175

Ala Gln Met Ile Gly Trp Tyr Asp Pro Lys Val Thr Arg Val Phe His
180 185 190

Ala Gly Phe Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr
195 200 205

Arg Ser Gly Glu Thr Val Arg Leu Met Asp Leu Leu Gly Glu Gly Leu
210 215 220

Lys Arg Ser Met Asp Lys Leu Lys Glu Lys Glu Arg Asp Lys Val Leu
225 230 235 240

Thr Ala Glu Glu Leu Asn Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys
245 250 255

Ile Lys Tyr Ala Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe
260 265 270

Ser Phe Asp Lys Met Leu Asp Asp Arg Gly Asn Thr Ala Ala Tyr Leu
275 280 285

Leu Tyr Ala Phe Thr Arg Ile Arg Ser Ile Ala Arg Leu Ala Asn Ile
290 295 300

-continued

Asp Glu Glu Met Leu Gln Lys Ala Ala Arg Glu Thr Lys Ile Leu Leu
 305 310 315 320
 Asp His Glu Lys Glu Trp Lys Leu Gly Arg Cys Ile Leu Arg Phe Pro
 325 330 335
 Glu Ile Leu Gln Lys Ile Leu Asp Asp Leu Phe Leu His Thr Leu Cys
 340 345 350
 Asp Tyr Ile Tyr Glu Leu Ala Thr Ala Phe Thr Glu Phe Tyr Asp Ser
 355 360 365
 Cys Tyr Cys Val Glu Lys Asp Arg Gln Thr Gly Lys Ile Leu Lys Val
 370 375 380
 Asn Met Trp Arg Met Leu Leu Cys Glu Ala Val Ala Ala Val Met Ala
 385 390 395 400
 Lys Gly Phe Asp Ile Leu Gly Ile Lys Pro Val Gln Arg Met Gly Lys
 405 410 415
 Pro Ile Pro Asn Pro Leu Leu Gly Leu Asp Ser Thr His His His His
 420 425 430
 His His

<210> SEQ ID NO 147

<211> LENGTH: 1773

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase polynucleotide

<400> SEQUENCE: 147

```

attaacatca tctcgcgct gcaagaagta ttcggccacg cgatcaaggc agcctaccg      60
gatctggaaa acccgccgct gttggtgacc ccgtcccagc aagcgaagtt cggcgactac      120
cagtgcaaca gcgcgatggg tattagccaa atgctgaaaa ctaaagaaca aaaggttaac      180
ccgcgcgaga ttgccgagaa tatcactaag cacttgccgg acaacgagtg catcgagaaa      240
gttgaaatcg cgggtccagg ttctattaat gtgcacctgc gcaaggattt tgtcagcgaa      300
caattgacga gcctgctggt gaattgtgtc cagctgccgg cgctgggtga aaacaagaaa      360
gtgatcgctg actttagcag ccctaataat gcgaaagaaa tgcacgtggg ccacctgcgt      420
agcacgatca tcggcgagtc gatttcccgc ctgtttgaat ttgccggtta tgatgttctg      480
cgtctgaatc acgtgggtga ttggggcacc cagtttggtg tgctgatcgc ccattctgcag      540
gataagtttc cggattacct gaccgtttct ccgccaatg gtgacttgca ggtcttttac      600
aaagaaagca agaaacgctt cgacacggaa gaagagttaa agaagcgcgc atatcaatgc      660
gtcgttctgc tgcagggcaa gaaccgggac atcacgaaag cctggaaaact gatttgtgac      720
gttagccgtc aggagttgaa taaaatttat gatgcgctgg atgtgtcgtc gatcgagcgt      780
ggcgagtcct tctatcagga ccgtatgaat gatattgtga aagaattcga ggaccgtggt      840
ttcgttcaag ttgatgacgg tcgtaagatc gtgttcgttc cgggctgtag catcccgtg      900
acgattgtta agagcgacgg tggttatacc tacgacacga gcgatctggc ggcgattaag      960
caacgtctgt ttgaagagaa agcagacatg attatctacg tcgtggacaa cggccaatct     1020
gtccacttcc aaaccatctt tgcgggtgca caaatgattg gctggtagca cccgaaagtt     1080
acccgtgttt tccatgcggg ttccgtgtgc gtgttgggag aagataagaa gaaattcaag     1140
acccgtagcg gtgaaacggt gcgtctgatg gacctgctgg gcgagggtct gaaacgtagc     1200

```

-continued

```

atggataagc tgaaagagaa agagcgcgac aaagtctga ccgctgaaga gctgaacgcg 1260
gctcagactt ccgtcgctta cggttgtatc aagtaacgag acctgagcca taatcgtctg 1320
aacgactaca ttttcagctt cgataagatg ctggatgatc gcggtaacac cgcagcgtat 1380
ctgctgtatg cattcacgcg tatccgcagc atcgcgcggt tggcgaacat tgatgaagag 1440
atgctgcaga aagcagcccg tgaaacaaa attctgctgg atcatgagaa agagtggaag 1500
ctgggcccgtt gcattctgcg ctttccggag atcttgca gaattttgga cgatctgttc 1560
ttgcacacc tgtgtgatta catttacgag ctggcgacgg cattcacga gttttatgac 1620
agctgctatt gtgtggagaa agaccgcaa accggcaaga tcctgaaagt caatatgtgg 1680
cgtatgctgc tgtgcgaagc cgttgcggct gtgatggcga agggctttga catcctgggt 1740
atcaaacggg ttcagcgtat gtaatgactc gag 1773

```

```

<210> SEQ ID NO 148
<211> LENGTH: 342
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase
polynucleotide

```

```

<400> SEQUENCE: 148

```

```

gacgaagaga tgctgcagaa ggccgctcgc gaaacaaaa tcctgctgga tcatgagaaa 60
gaatggaagc tgggtcgtg taccctgcgc ttcccgaaa tcctgcagaa gattctggac 120
gacctgttct tgcacacct gtgtgattat atctatgagc tggcgacggc ctttacggag 180
ttttacgata gctgctactg cgtggagaaa gaccgtcaa ccggcaagat tctgaaagt 240
aacatgtggc gtatgctgct gtgcgaagca gtcgcggctg tgatggcgaa gggcttcgac 300
atcttgggta ttaaacgggt tcagcgtatg taatgactcg ag 342

```

```

<210> SEQ ID NO 149
<211> LENGTH: 603
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase
polynucleotide

```

```

<400> SEQUENCE: 149

```

```

gacctgctgg gcgagggcct gaaacgcagc atggataagt tgaaagagaa agagcgcgac 60
aaagtgctga ccgcggaaga gctgaatgct gcacagacca gcgtcgcgta tggttgtatc 120
aagtatgcag atctgtctca caatcgctg aacgattaca ttttcagctt cgacaagatg 180
ctggatgacc gtggcaatac ggccgcctat ctgctgtatg cgtttaccgg catccgttcc 240
atcgcacgtc tggcgaacat tgacgaagag atgctgcaga aggcggcacg cgaaaccaag 300
attctgttgg accatgagaa agaatgaaa ctgggtcgtt gtattctgcg tttccggag 360
atcttgcaaa agatcttgga cgacctgtt ctgcacactc tgtgcgatta catctacgaa 420
ctggcgacgg cttttaccga gttttacgat agctgctact gtgtggagaa ggaccgtcaa 480
acgggtaaaa tcctgaaagt taacatgtgg cgtatgctgc tgtgcgaagc tgcgcgcgc 540
gttatggcga agggtttcga tattctgggc attaaacggg ttcagcgtat gtaatgactc 600
gag 603

```

-continued

```

<210> SEQ ID NO 150
<211> LENGTH: 1251
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase
      polynucleotide

<400> SEQUENCE: 150

ctgatcgccc acctgcaaga caaatccct gactacctga ccgtagccc gccgatcggt    60
gacctgcaag tgttttacia agagagcaag aagcgctttg ataccgaaga agagttcaag    120
aaacgcgcgt accagtgtgt tgtgctgctg caaggaaga atccggatat caccaaagca    180
tggaagctga tttgcgacgt tagccgtcag gagctgaaca aaatctatga cgcactggac    240
gtttccctga tcgaacgcgg tgagagcttt taccaagacc gtatgaatga tattgtgaaa    300
gaatttgagg accgcggttt tgtccaagtt gatgatggtc gtaagattgt gttcgtgccg    360
ggctgctcga tcccgtgac gattgtgaaa agcgatggtg gctatactta cgacaccagc    420
gatctggctg cgattaagca gcgtctgttc gaagagaaag ctgatatgat tatctatgtt    480
gtggacaacg gccagagcgt ccatttccag acgatcttcg cggcagcaca aatgattggt    540
tggtacgacc caaaggtgac ccgtgtgttt cacgcgggtt tcggcgttgt cctgggtgag    600
gataagaaga agttcaagac ccgtagcggc gaaacggtec gtttgatgga tctgctgggt    660
gaaggcctga aacgctcgat ggacaagttg aaagagaaag agcgcgacaa ggttctgacc    720
gccgaagagc tgaatgcggc gaaaacctcc gtcgcgtacg gttgtatcaa gtatgcggac    780
ctgtctcata atcgtctgaa tgattatata ttttctttcg acaaaatgct ggatgatcgt    840
ggcaacacgg cggcctacct gctgtacgct ttcacccgca tccgtagcat tgcacgctg    900
gcgaacattg atgaagagat gctgcaaaag gctgcacgtg aaacaaaaat tctgctggac    960
cacgagaaag agtggaaact gggccgttgt attctgcgtt ttctgagat cttgcagaaa   1020
atcctggaag atttgttect gcacacgctg tgtgactaca tttatgagct ggcaactgcg   1080
ttcacggaat tctacgacag ctgctattgc gtcgagaagg accgtcagac cgtaagatc   1140
ctgaaagtta acatgtggcg tatgttgctg tgcgaagccg ttgccgcggt tatggccaaa   1200
ggttttgaca tcttgggtat taaaccggtc cagcgcgatg aatgactcga g           1251

```

```

<210> SEQ ID NO 151
<211> LENGTH: 363
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 151

Lys Asp Phe Val Ser Glu Gln Leu Thr Ser Leu Leu Val Asn Gly Val
1          5          10          15

Gln Leu Pro Ala Leu Gly Glu Asn Lys Lys Val Ile Val Asp Phe Ser
20          25          30

Ser Pro Asn Ile Ala Lys Glu Met His Val Gly His Leu Arg Ser Thr
35          40          45

Ile Ile Gly Glu Ser Ile Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp
50          55          60

Val Leu Arg Leu Asn His Val Gly Asp Trp Gly Thr Gln Phe Gly Met
65          70          75          80

```

-continued

Leu Ile Ala His Leu Gln Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser
 85 90 95
 Pro Pro Ile Gly Asp Leu Gln Val Phe Tyr Lys Glu Ser Lys Lys Arg
 100 105 110
 Phe Asp Thr Glu Glu Glu Phe Lys Lys Arg Ala Tyr Gln Cys Val Val
 115 120 125
 Leu Leu Gln Gly Lys Asn Pro Asp Ile Thr Lys Ala Trp Lys Leu Ile
 130 135 140
 Cys Asp Val Ser Arg Gln Glu Leu Asn Lys Ile Tyr Asp Ala Leu Asp
 145 150 155 160
 Val Ser Leu Ile Glu Arg Gly Glu Ser Phe Tyr Gln Asp Arg Met Asn
 165 170 175
 Asp Ile Val Lys Glu Phe Glu Asp Arg Gly Phe Val Gln Val Asp Asp
 180 185 190
 Gly Arg Lys Ile Val Phe Val Pro Gly Cys Ser Ile Pro Leu Thr Ile
 195 200 205
 Val Lys Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala
 210 215 220
 Ile Lys Gln Arg Leu Phe Glu Glu Lys Ala Asp Met Ile Ile Tyr Val
 225 230 235 240
 Val Asp Asn Gly Gln Ser Val His Phe Gln Thr Ile Phe Ala Ala Ala
 245 250 255
 Gln Met Ile Gly Trp Tyr Asp Pro Lys Val Thr Arg Val Phe His Ala
 260 265 270
 Gly Phe Gly Val Val Leu Gly Glu Asp Lys Lys Lys Phe Lys Thr Arg
 275 280 285
 Ser Gly Glu Thr Val Arg Leu Met Asp Leu Leu Gly Glu Gly Leu Lys
 290 295 300
 Arg Ser Met Asp Lys Leu Lys Glu Lys Glu Arg Asp Lys Val Leu Thr
 305 310 315 320
 Ala Glu Glu Leu Asn Ala Ala Gln Thr Ser Val Ala Tyr Gly Cys Ile
 325 330 335
 Lys Tyr Ala Asp Leu Ser His Asn Arg Leu Asn Asp Tyr Ile Phe Ser
 340 345 350
 Phe Asp Lys Met Leu Asp Asp Arg Gly Asn Thr
 355 360

<210> SEQ ID NO 152

<211> LENGTH: 1089

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 152

```

aaggattttg tatcagaaca attgaccagt cttctagtga atggagttca actacctgct      60
ctgggagaga ataaaaaggt tatagttgac ttttcctccc ctaatatagc taaagagatg      120
catgtaggcc acctgaggtc aactatcata ggagagagta taagccgcct ctttgaattt      180
gcagggtatg acgtgctcag gttaaatacat gtaggagact gggggaccca gtttgcatg      240
ctcatcgctc acctgcaaga caaatttcca gattatctaa cagtttcacc tcctattggg      300
gatcttcagg tcttttataa ggaatctaag aagagggttg atactgagga ggaatttaag      360
aagcgagcat atcagtgtgt agttctgctc cagggtaaaa acccagatat tacaaaagct      420

```

-continued

```

tggaagctta tctgtgatgt ctcccgccaa gagttaaata aaatctatga tgcattggac 480
gtctctttta tagagagagg ggaatccttc tatcaagata ggatgaatga tattgtaaag 540
gaatttgaag atagaggatt tgtgcagggt gatgatggca gaaagattgt atttgtccca 600
gggtgttcca taccattaac catagtaaaa tcagatggag gttataccta tgatacatct 660
gacctggctg ctattaaaca aagactatct gaggaanaag cagatatgat tatctatgtt 720
gtggacaatg gacaatctgt gcacttcag acaatatttg ctgctgctca aatgattggt 780
tggtatgacc ctaaagtaac tcgagtcttc catgctggat ttggtgtggt gctaggggaa 840
gacaagaaaa agtttaaaac acgttcgggt gaaacagtgc gcctcatgga tcttctggga 900
gaaggactaa aacgatccat ggacaagttg aaggaaaaag aaagagacaa ggtcttaact 960
gcagaggaaat tgaatgctgc tcagacatcc gttgcgtatg gctgcatcaa atatgctgac 1020
ctttcccata accggttgaa tgactacatc ttctcctttg acaaaatgct agatgacaga 1080
ggaaataca 1089

```

```

<210> SEQ ID NO 153
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 153

```

```

Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys
1           5           10

```

```

<210> SEQ ID NO 154
<211> LENGTH: 172
<212> TYPE: PRT
<213> ORGANISM: Mus musculus

```

```

<400> SEQUENCE: 154

```

```

Glu Met His Val Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met
1           5           10           15
Ser Arg Leu Phe Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His
20          25          30
Val Gly Asp Trp Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln
35          40          45
Asp Lys Phe Pro Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu
50          55          60
Gln Ala Phe Tyr Lys Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu
65          70          75          80
Phe Lys Lys Arg Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn
85          90          95
Pro Asp Ile Met Lys Ala Trp Asn Leu Ile Cys Asp Val Ser Arg Glu
100         105         110
Glu Phe Lys Lys Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg
115         120         125
Gly Glu Ser Phe Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe
130         135         140
Glu Asp Lys Gly Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe
145         150         155         160
Val Pro Gly Cys Ser Val Pro Leu Thr Ile Val Lys
165         170

```

-continued

<210> SEQ ID NO 155

<211> LENGTH: 16

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 155

Ser Asp Gly Gly Tyr Thr Tyr Asp Thr Ser Asp Leu Ala Ala Ile Lys
1 5 10 15

<210> SEQ ID NO 156

<211> LENGTH: 200

<212> TYPE: PRT

<213> ORGANISM: Mus musculus

<400> SEQUENCE: 156

Val Ile Val Asp Phe Ser Ser Pro Asn Ile Ala Lys Glu Met His Val
1 5 10 15

Gly His Leu Arg Ser Thr Ile Ile Gly Glu Ser Met Ser Arg Leu Phe
20 25 30

Glu Phe Ala Gly Tyr Asp Val Leu Arg Leu Asn His Val Gly Asp Trp
35 40 45

Gly Thr Gln Phe Gly Met Leu Ile Ala His Leu Gln Asp Lys Phe Pro
50 55 60

Asp Tyr Leu Thr Val Ser Pro Pro Ile Gly Asp Leu Gln Ala Phe Tyr
65 70 75 80

Lys Glu Ser Lys Lys Arg Phe Asp Ala Asp Glu Glu Phe Lys Lys Arg
85 90 95

Ala Tyr Gln Cys Val Val Leu Leu Gln Ser Lys Asn Pro Asp Ile Met
100 105 110

Lys Ala Trp Asn Leu Ile Cys Asp Val Ser Arg Glu Glu Phe Lys Lys
115 120 125

Ile Tyr Asp Ala Leu Asp Ile Thr Leu Ile Glu Arg Gly Glu Ser Phe
130 135 140

Tyr Gln Asp Arg Met Lys Asp Ile Val Lys Glu Phe Glu Asp Lys Gly
145 150 155 160

Phe Val Gln Val Asp Asp Gly Arg Lys Ile Val Phe Val Pro Gly Cys
165 170 175

Ser Val Pro Leu Thr Ile Val Lys Ser Asp Gly Gly Tyr Thr Tyr Asp
180 185 190

Thr Ser Asp Leu Ala Ala Ile Lys
195 200

<210> SEQ ID NO 157

<211> LENGTH: 384

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with
N-terminal 6xHis affinity tag

<400> SEQUENCE: 157

Met His His His His His His Gly Lys Pro Ile Pro Asn Pro Leu Leu
1 5 10 15

Gly Leu Asp Ser Thr Lys Asp Phe Val Ser Glu Gln Leu Thr Ser Leu
20 25 30

-continued

Leu	Val	Asn	Gly	Val	Gln	Leu	Pro	Ala	Leu	Gly	Glu	Asn	Lys	Lys	Val
		35					40					45			
Ile	Val	Asp	Phe	Ser	Ser	Pro	Asn	Ile	Ala	Lys	Glu	Met	His	Val	Gly
	50					55					60				
His	Leu	Arg	Ser	Thr	Ile	Ile	Gly	Glu	Ser	Ile	Ser	Arg	Leu	Phe	Glu
65					70					75					80
Phe	Ala	Gly	Tyr	Asp	Val	Leu	Arg	Leu	Asn	His	Val	Gly	Asp	Trp	Gly
				85					90					95	
Thr	Gln	Phe	Gly	Met	Leu	Ile	Ala	His	Leu	Gln	Asp	Lys	Phe	Pro	Asp
			100					105					110		
Tyr	Leu	Thr	Val	Ser	Pro	Pro	Ile	Gly	Asp	Leu	Gln	Val	Phe	Tyr	Lys
		115					120					125			
Glu	Ser	Lys	Lys	Arg	Phe	Asp	Thr	Glu	Glu	Glu	Phe	Lys	Lys	Arg	Ala
	130					135					140				
Tyr	Gln	Cys	Val	Val	Leu	Leu	Gln	Gly	Lys	Asn	Pro	Asp	Ile	Thr	Lys
145					150					155					160
Ala	Trp	Lys	Leu	Ile	Cys	Asp	Val	Ser	Arg	Gln	Glu	Leu	Asn	Lys	Ile
				165					170					175	
Tyr	Asp	Ala	Leu	Asp	Val	Ser	Leu	Ile	Glu	Arg	Gly	Glu	Ser	Phe	Tyr
			180					185					190		
Gln	Asp	Arg	Met	Asn	Asp	Ile	Val	Lys	Glu	Phe	Glu	Asp	Arg	Gly	Phe
		195					200					205			
Val	Gln	Val	Asp	Asp	Gly	Arg	Lys	Ile	Val	Phe	Val	Pro	Gly	Cys	Ser
	210					215					220				
Ile	Pro	Leu	Thr	Ile	Val	Lys	Ser	Asp	Gly	Gly	Tyr	Thr	Tyr	Asp	Thr
225					230					235					240
Ser	Asp	Leu	Ala	Ala	Ile	Lys	Gln	Arg	Leu	Phe	Glu	Glu	Lys	Ala	Asp
				245					250					255	
Met	Ile	Ile	Tyr	Val	Val	Asp	Asn	Gly	Gln	Ser	Val	His	Phe	Gln	Thr
			260					265					270		
Ile	Phe	Ala	Ala	Ala	Gln	Met	Ile	Gly	Trp	Tyr	Asp	Pro	Lys	Val	Thr
		275					280					285			
Arg	Val	Phe	His	Ala	Gly	Phe	Gly	Val	Val	Leu	Gly	Glu	Asp	Lys	Lys
	290					295					300				
Lys	Phe	Lys	Thr	Arg	Ser	Gly	Glu	Thr	Val	Arg	Leu	Met	Asp	Leu	Leu
305					310					315					320
Gly	Glu	Gly	Leu	Lys	Arg	Ser	Met	Asp	Lys	Leu	Lys	Glu	Lys	Glu	Arg
				325					330					335	
Asp	Lys	Val	Leu	Thr	Ala	Glu	Glu	Leu	Asn	Ala	Ala	Gln	Thr	Ser	Val
			340					345					350		
Ala	Tyr	Gly	Cys	Ile	Lys	Tyr	Ala	Asp	Leu	Ser	His	Asn	Arg	Leu	Asn
		355					360					365			
Asp	Tyr	Ile	Phe	Ser	Phe	Asp	Lys	Met	Leu	Asp	Asp	Arg	Gly	Asn	Thr
	370					375					380				

<210> SEQ ID NO 158

<211> LENGTH: 384

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Arginyl tRNA Synthetase polypeptide with C-terminal 6xHis affinity tag

<400> SEQUENCE: 158

-continued

Met 1	Lys	Asp	Phe	Val 5	Ser	Glu	Gln	Leu	Thr 10	Ser	Leu	Leu	Val	Asn 15	Gly
Val	Gln	Leu	Pro 20	Ala	Leu	Gly	Glu	Asn 25	Lys	Lys	Val	Ile	Val 30	Asp	Phe
Ser	Ser	Pro 35	Asn	Ile	Ala	Lys	Glu 40	Met	His	Val	Gly	His 45	Leu	Arg	Ser
Thr	Ile 50	Ile	Gly	Glu	Ser	Ile 55	Ser	Arg	Leu	Phe	Glu 60	Phe	Ala	Gly	Tyr
Asp 65	Val	Leu	Arg	Leu	Asn 70	His	Val	Gly	Asp	Trp 75	Gly	Thr	Gln	Phe	Gly 80
Met	Leu	Ile	Ala	His 85	Leu	Gln	Asp	Lys 90	Phe	Pro	Asp	Tyr	Leu	Thr 95	Val
Ser	Pro	Pro	Ile 100	Gly	Asp	Leu	Gln	Val 105	Phe	Tyr	Lys	Glu	Ser 110	Lys	Lys
Arg	Phe	Asp 115	Thr	Glu	Glu	Glu	Phe 120	Lys	Lys	Arg	Ala	Tyr 125	Gln	Cys	Val
Val	Leu 130	Leu	Gln	Gly	Lys	Asn 135	Pro	Asp	Ile	Thr	Lys 140	Ala	Trp	Lys	Leu
Ile 145	Cys	Asp	Val	Ser	Arg 150	Gln	Glu	Leu	Asn	Lys 155	Ile	Tyr	Asp	Ala	Leu 160
Asp	Val	Ser	Leu 165	Ile	Glu	Arg	Gly	Glu 170	Ser	Phe	Tyr	Gln	Asp	Arg 175	Met
Asn	Asp	Ile	Val 180	Lys	Glu	Phe	Glu	Asp 185	Arg	Gly	Phe	Val	Gln 190	Val	Asp
Asp	Gly	Arg 195	Lys	Ile	Val	Phe	Gly 200	Pro	Gly	Cys	Ser	Ile 205	Pro	Leu	Thr
Ile	Val 210	Lys	Ser	Asp	Gly 215	Tyr	Thr	Tyr	Asp 220	Thr	Ser	Asp	Leu	Ala	
Ala 225	Ile	Lys	Gln	Arg	Leu 230	Phe	Glu	Glu	Lys	Ala 235	Asp	Met	Ile	Ile	Tyr 240
Val	Val	Asp	Asn 245	Gly	Gln	Ser	Val	His	Phe 250	Gln	Thr	Ile	Phe	Ala 255	Ala
Ala	Gln	Met	Ile 260	Gly	Trp	Tyr	Asp	Pro 265	Lys	Val	Thr	Arg 270	Val	Phe	His
Ala	Gly	Phe 275	Gly	Val	Val	Leu	Gly 280	Glu	Asp	Lys	Lys	Lys 285	Phe	Lys	Thr
Arg	Ser 290	Gly	Glu	Thr	Val	Arg 295	Leu	Met	Asp	Leu	Leu 300	Gly	Glu	Gly	Leu
Lys 305	Arg	Ser	Met	Asp 310	Lys	Leu	Lys	Glu	Lys	Glu 315	Arg	Asp	Lys	Val	Leu 320
Thr	Ala	Glu	Glu	Leu 325	Asn	Ala	Ala	Gln	Thr	Ser	Val	Ala	Tyr	Gly 335	Cys
Ile	Lys	Tyr	Ala 340	Asp	Leu	Ser	His	Asn 345	Arg	Leu	Asn	Asp	Tyr 350	Ile	Phe
Ser	Phe	Asp 355	Lys	Met	Leu	Asp	Asp 360	Arg	Gly	Asn	Thr	Gly 365	Lys	Pro	Ile
Pro	Asn 370	Pro	Leu	Leu	Gly	Leu	Asp 375	Ser	Thr	His	His	His	His	His	His

```
<211> LENGTH: 1101
```

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Codon optimized Arginyl tRNA Synthetase
        polynucleotide

<400> SEQUENCE: 159

aaagacttcg tgagcgaaca actgacgtcc ctgctggtaa acggcgtgca actgccggca      60
ctgggcgaga acaagaaagt tatcgtcgac tttagcagcc cgaacattgc gaaagaaatg      120
catgtgggcc acctgcgtc caccattatt ggtgagagca tctcccgctc gttegagttt      180
gcaggttacg atgttctgcg cctgaatcac gtcggtgact ggggcacgca atttggcatg      240
ctgacgcctc atctgcagga caagtttccg gattacttga ccgtttccgc accgattggc      300
gacttgcagg tgttttacia agagagcaag aagcgtttcg ataccgaaga agagttcaag      360
aaacgcgcac accagtgcgt cgtcctgctg caaggaaga atcctgatat taccaaagcc      420
tgaaaactga tttgtgatgt tagccgtcaa gaactgaata aaatttacga tgcgctggac      480
gtgtccctga tcgaacgtgg tgaaagcttc tatcaggacc gcatgaatga catcggtaaa      540
gaatttgagg accgtggttt cgttcaagtg gatgacggtc gcaagattgt tttcgtgccg      600
ggttgctcta ttccgtgac gatcgtgaag agcgatggcg gttataccta cgacatttct      660
gatctggcag cgattaaaca gcgtctgttt gaagagaagg ccgacatgat catctatgtc      720
gttgataacg gccagagcgt tcatttccag acgattttcg cggcggctca gatgatcggc      780
tggtatgacg cgaagtcac gcgtgtgttc caccggggtt ttggtgtggt tctgggtgag      840
gataagaaga agttcaagac ccgtagcggg gagactgtgc gtctgatgga cctgctgggt      900
gaaggcctga agcgtagcat ggacaagttg aaagagaaaag agcgtgataa agtcctgacg      960
gcggaagagt tgaacgcagc gcaaacaccg gttgcctacg gttgtatcaa atatgcggac      1020
ctgagccaca atcgcttgaa cgattatata ttcagctttg acaaaatgct ggaacgaccg      1080
ggtaacacct aatgactcga g                                     1101

```

1-125. (canceled)

126. A therapeutic composition, comprising a pharmaceutically-acceptable carrier and an antibody or antigen-binding fragment thereof that exhibits binding specificity for an isolated aminoacyl-tRNA synthetase (AARS) polypeptide or an epitope selected from Table(s) 1-3, or Table(s) 4-6, and Table(s) 7-9, wherein the composition has a purity of at least about 90% on a protein basis and less than about 10 EU endotoxin/mg protein.

127. The therapeutic composition of claim **126**, wherein the AARS polypeptide is selected from Table E7.

128. The therapeutic composition of claim **126**, wherein the antibody or antigen binding fragment thereof is a monoclonal antibody.

129. The therapeutic composition of claim **126**, wherein the antibody is a humanized antibody.

130. The therapeutic composition of claim **126**, wherein the antibody or antigen-binding fragment thereof is an Fv fragment or a single chain Fv (sFv) polypeptide.

131. The therapeutic composition of claim **126**, wherein the composition is a sterile, freeze-dried powder.

132. The therapeutic composition of claim **126**, wherein the composition is a sterile injectable solution.

133. The therapeutic composition of claim **126**, wherein the composition is buffered and comprises an isotonic agent.

134. The therapeutic composition of claim **126**, wherein the composition is suitable for intravenous, intramuscular, subcutaneous, or intraperitoneal administration.

135. The therapeutic composition of claim **126**, wherein the composition comprises a surfactant.

136. The therapeutic composition of claim **126**, wherein the composition has a purity of at least about 95% on a protein basis.

137. A method of modulating a cellular activity of a cell, or protein, comprising contacting the cell or protein with a therapeutic composition of claim **126**.

138. The method of claim **137**, wherein the cellular activity is selected from cell proliferation, cell differentiation, cell signaling, gene transcription, angiogenesis, cell binding, metabolism, cytokine production or activity, cytokine receptor activity, and inflammation.

139. The method of claim **137**, wherein the cell type is selected from the group consisting of pre-adipocytes, bone marrow, neutrophils, blood cells, hepatocytes, astrocytes, mesenchymal stem cells, and skeletal muscle cells.

140. The method of claim **137**, wherein the cell is in a subject.

141. The method of claim **140**, comprising treating the subject, wherein the subject has a condition associated with a neoplastic disease, an immune system disease or condition, an inflammatory disorder, an infectious disease, a metabolic disease, neuronal/neurological disease, a muscular/cardiovascular disease, a disease associated with aberrant hematopoiesis, a disease associated with aberrant angiogenesis, or a disease associated with aberrant cell survival.

* * * * *