



(86) Date de dépôt PCT/PCT Filing Date: 1998/05/14
 (87) Date publication PCT/PCT Publication Date: 1998/11/19
 (45) Date de délivrance/Issue Date: 2009/05/05
 (85) Entrée phase nationale/National Entry: 1999/11/12
 (86) N° demande PCT/PCT Application No.: US 1998/009788
 (87) N° publication PCT/PCT Publication No.: 1998/051821
 (30) Priorité/Priority: 1997/05/15 (US60/046,529)

(51) Cl.Int./Int.Cl. *G01N 33/68* (2006.01),
G01N 33/574 (2006.01), *C12Q 1/68* (2006.01)
 (72) Inventeur/Inventor:
GIORDANO, ANTONIO, US
 (73) Propriétaire/Owner:
THOMAS JEFFERSON UNIVERSITY, US
 (74) Agent: BERESKIN & PARR LLP

(54) Titre : DETERMINATION DE NIVEAUX DE P27 INHIBITEUR DE KINASE DEPENDANT DE LA CYCLINE COMME
FACTEUR DE PRONOSTIC CHEZ DES PATIENTS ATTEINTS DE CANCER
 (54) Title: DETERMINATION OF CYCLIN-DEPENDENT KINASE INHIBITOR P27 LEVELS AS A PROGNOSTIC
FACTOR IN CANCER PATIENTS

(57) **Abrégé/Abstract:**

Methods of grading tumors and prognosticating survival rates of cancer patients by determination of p27 expression levels in tissues samples from tumors are provided.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, G01N 33/574	A1	(11) International Publication Number: WO 98/51821 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/US98/09788 (22) International Filing Date: 14 May 1998 (14.05.98) (30) Priority Data: 60/046,529 15 May 1997 (15.05.97) US (71) Applicant: THOMAS JEFFERSON UNIVERSITY [US/US]; 11th and Walnut Streets, Philadelphia, PA 19107 (US). (72) Inventor: GIORDANO, Antonio; 303 Ginger Lane, Philadel- phia, PA (US). (74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: DETERMINATION OF CYCLIN-DEPENDENT KINASE INHIBITOR P27 LEVELS AS A PROGNOSTIC FACTOR IN CANCER PATIENTS		
(57) Abstract Methods of grading tumors and prognosticating survival rates of cancer patients by determination of p27 expression levels in tissues samples from tumors are provided.		

**DETERMINATION OF CYCLIN-DEPENDENT KINASE INHIBITOR
P27 LEVELS AS A PROGNOSTIC FACTOR IN CANCER PATIENTS**

Introduction

This invention was made in the course of research
5 sponsored by the National Institutes of Health. The U.S.
Government may have certain rights in this invention.

Background of the Invention

The eukaryotic cell cycle is controlled by protein
complexes composed of cyclins and cyclin-dependent kinases
10 (cdks). The regulatory function of cdks is achieved by
phosphorylation of key substrates, such as the members of the
retinoblastoma gene family. Activity of cdks is regulated
by post-translational modification and by the association or
dissociation with inhibitory subunits designated cyclin-
15 dependent-kinase inhibitors (CKIs). Two families of these
inhibitors have been identified in mammalian cells. Members
of each of the two families share a high percentage of
sequence homology, in addition to their specificity of
interaction with cdks. The first family, which includes p21
20 (also known as Cip1, Pic1, Sdi1, mda6 and Waf1), p27 (also
known as Ick, Kip1 and Pic2) and p57 (also known as Kip2)
preferentially inhibit cdk2. The second family, which
includes p16 (also known as Ink4A, Mts1, Cdkn2 and Cdkn4i),
p15 (also known as Ink4B and Mts2), p18 (also known as Ink4C
25 and Ink6A) and p19/p20 (also known as Ink4D and Ink6B)
preferentially bind to and inhibit cdk4 and cdk6. Results

- 2 -

from several studies indicate CKIs to be the products of potential tumor-suppressor genes (MacLachlan et al. (1995) *Crit. Rev. Eukaryotic Gene Expression* 5:127-156). p27 was first identified in complexes with cdk2/cyclin E in Transforming Growth Factor- β (TGF- β) arrested cells (Koff et al. (1991) *Cell* 66:1217-1228). p27 protein associates with cyclin E/cdk2 and Cyclin A/cdk2 complexes and inhibits their activities. It is a negative cell cycle regulator implicated in G1 phase arrest by TGF- β cell contact, inhibition agents that increase the level of cyclic AMP, staurosporin, lovastatin, tamoxifen and rapamycin. Overexpression of p27 protein in mammalian cells induces a G1 block of the cell cycle (Polyak et al. (1994) *Cell* 79: 59-66; Toyoshima, T. and Hunter, T. (1994) *Cell* 78: 67-74). In addition, high levels of p27 have been found in quiescent cells thus suggesting a role for p27 in maintaining cells in G0 (Nourse et al. (1994) *Nature* 372:570-573). Levels of p27 decrease as cells reenter the cell cycle, mostly due to ubiquitin-proteasome dependent degradation (Pagano et al. (1995) *Science* 269:682-685). No structural alteration of p27 gene has been reported in human neoplasms to date (Bullrich et al. (1995) *Cancer Research* 55: 1199-1205; Cordon-Cardo, C. (1995) *Am. J. Pathol.* 147:1-16; Kawamat et al. (1995) *Cancer Research* 55:2266-2269; Pietenpol et al. (1995) *Cancer Research* 55:1206-1210; Ponce-Castañeda et al. (1995) *Cancer Research* 55:1211-1214; Stegmaier et al. (1996) *Cancer Research* 56:1413-1417). Thus, despite its potential role as a tumor suppressor, p27 gene does not appear to be mutated in human tumors.

However, recent evidence suggests an involvement of this CKI in neoplastic transformation. For example, p27 deficient mice have been shown to develop tumors of the pituitary gland with 100% of penetrance (Fero et al. (1996) *Cell* 85:733-774; Kiyokawa et al. (1996) *Cell* 85:721-732). In addition, p27 has been shown to be a target of the Adenovirus E1A (Mal et al.

(1996) *Nature* 380:262-265). Further, HPV E7 oncoproteins have been shown to dissociate p27 from the cyclin/cdk complexes (Zerfass-Thome et al. (1996) *Oncogene* 2323-2330). p27 has also been implicated as a regulator of drug resistance in 5 solid tumors and has been suggested as a target for development of antagonists which may be useful as chemosensitizers in conjunction with conventional anticancer therapy. St. Croix et al. (1996) *Nature Medicine* 2(11):1204-1210.

10 It has now been found that levels of p27 expression in tumor cells correlate with the degree of malignancy of the tumor and can be used in prognosticating overall survival time in cancer patients.

Summary of the Invention

15 An object of the present invention is to provide a method of determining the degree of malignancy or grading of a tumor in a patient by determining levels of p27 expression in tumor cells isolated from the patient.

20 Another object of the present invention is to provide a method of prognosticating overall survival time in cancer patients by determining levels of p27 expression in tumor cells isolated from the patient.

Detailed Description of the Invention

25 Lung cancer is the greatest single cause of cancer-related deaths in Western countries and despite continuing research efforts into new therapeutic strategies, survival in patients suffering from lung cancer (approximately 13%) has not improved over the past two decades. Only approximately 30% of patients suffering from lung cancer are eligible for 30 radical lung resection, and only about one third of these patients are free from disease 5 years after surgery (Shottenfeld, D., "Epidemiology of lung cancer", in *Lung Cancer: Principles and Practice*, ed. by Pass H.I., Mitchell J.B., Johnson, Turrisi AT. (1996). Lippincot-Raven

- 4 -

Publishers, Philadelphia). Thus, accurate determination of the cancer's malignant potential is important in order to select an appropriate course of therapy. This determination, known as "grading", is typically carried out
5 by examination of the character and appearance of tumor sections by skilled pathologists and is based upon a number of histological criteria. However, a significant problem in the use of such histological criteria is that they are subjective, and variability exists not only between different
10 pathologists but, oftentimes, within a single pathologist's reading of the same tissue specimen. In addition, there is heterogeneity within the tumor itself in both primary and metastatic sites. Accordingly, there is a need for more accurate determination of the malignant potential of a tumor.

15 It has now been found that expression levels of cell cycle regulators such as p27 in tumor cells correlate with the degree of malignancy (grading) ($p=0.01$). Further, expression levels of p27 in tumor cells were shown to correlate with survival time in cancer patients. Thus, a simple
20 immunohistochemical assay measuring p27 expression on formalin fixed-paraffin embedded specimens can now be used to quickly and reproducibly to grade tumor and to evaluate the prognosis of cancer patients.

Accordingly, the present invention provides a method of
25 grading tumors by determination of the levels of p27 expression in a tissue sample from a tumor. Tissue samples of a tumor can be obtained by various means including, but not limited, surgical resectioning or removal and biopsies. Once the tissue is obtained, levels of p27 expression in the sample
30 are determined preferably via an immunohistochemical assay. As will be obvious to those of skill in the art upon this disclosure, however, other means of measuring p27 expression levels in tissues can be used. The levels are then compared to established control levels of p27 expression in normal
35 noncancerous tissue, nonmetastatic cancerous tissue and metastatic cancerous tissue so that tissue sample can be graded.

The present invention also provides a method for prognosticating the survival rates of cancer patients by determination of p27 levels in tumor tissue obtained from these patients. For example, the determination of p27 levels in tissue samples from patients affected by non small cell lung cancer was demonstrated to have prognostic value in identifying those patients most likely to benefit from radical lung resection. By "non small cell lung cancer or NSCLC", it is meant all forms of lung cancer except small cell lung cancer (SCLC) and includes, but is not limited to squamous cell carcinomas, adenocarcinomas, bronchioalveolar carcinomas and large cell carcinomas. p27 protein expression levels were measured in a series of non small cell lung cancer specimens. Immunohistochemistry and western blot analysis were performed on each specimen. Tumors expressing low to undetectable levels of p27 were found to contain high p27 degradation activity. Expression levels were then correlated with the outcomes of these patients. It was found that p27 was a prognostic factor correlating with the overall survival times (p=.0012).

More specifically, routine histological assays were performed on 108 samples of non small cell lung cancers and evaluated independently by p27 immunostaining. Sixty-two of these tumors were identified as adenocarcinomas while 46 were squamous carcinomas. Sixty-two of the tumors (57.4%) were classified as stage I, 18 of the tumors (16.7%) were classified as stage II and 28 tumors (25.9%) were classified as stage III. Thirty-two of the tumors were considered low differentiated (29.6%), 42 of the tumors were medium differentiated (38.9%) and 34 of the tumors were well differentiated (31.5%).

Immunoreactivity for p27 was found in both normal and neoplastic tissues. p27 immunostaining in both tissues was always nuclear, with a low to absent background. p27 expression in normal lung tissue was detected in bronchial epithelia (mostly in ciliated cells) and in the adjacent glands. Pneumocytes displayed moderate nuclear

immunoreactivity. p27 immunostaining was also observed in 96 (88.9%) of the neoplastic specimens. The percentage of positive cells in the remaining 12 specimens (11.1%) ranged from 0 to 5%. No statistically significant difference in the p27 expression was found between adenocarcinomas and squamous carcinomas. Specificity of the staining was confirmed by blocking with a p27 specific antibody.

p27 expression was also analyzed in these specimens by western blot. Protein extracts were prepared from all of the 108 frozen samples. p27 protein was found at varying levels in all of the specimens examined. Twelve of the 108 specimens, however, had undetectable levels of p27 protein expression. To confirm that the same amount of protein was present in each lysate and that the samples were not degraded, the same blot was normalized with the monoclonal antibody heat shock protein 72/73 (HSP 72/73; Oncogene Science, Uniondale, NY). Results from western blotting correlated statistically with results from the immunohistochemical assays ($p < 0.001$).

Northern blot analysis was performed to determine if the different levels of p27 protein were related to difference in transcriptional levels. However, similar amounts of p27 transcript were found to be present in all the examined specimens.

Accordingly, degradation assays were performed to determine whether the p27 degradation pathway was enhanced in tumors with low to undetectable p27 expression. In these experiments, 1 gram of each frozen human tissue sample (6 representative samples) was sectioned and quickly homogenized at 15000 rpm in 1 ml of ice cold doubly distilled water. The sample was frozen and thawed 3 times. Purified recombinant p27 was incubated either alone or in the presence of extracts from tumor expressing different p27 levels for 1 hour, 3 hours, 6 hours, 9 hours and 12 hours. It was found that tumors with high p27 expression level presented lower degradation activity, as compared with tumors with low or undetectable p27 expression level, which presented high or

- 7 -

very high degradation activity. Thus, low levels of p27 protein in tumors correlate with high p27 degradation activity.

The role of proteasome in this degradation was then
5 determined. In these experiments, samples with low to undetectable p27 expression were ultracentrifuged to eliminate the proteasome particles. Purified recombinant p27 was then incubated in the presence of extracts from the proteasome depleted supernatant and in the same supernatant after
10 readdition of the proteasome-rich pellet. No degradation was detected in the sample in which the proteasome was left out. p27 degradation was found to be restored in the sample in which the proteasome was added back to the supernatant.

Specimens were then divided into three groups based upon
15 p27 expression levels. The first group (referred to as non expressors) contained 12 specimens (11.1%) with up to 5% of cells positive for p27. The second group (referred to as low expressors) contained 56 (51.8%) specimens with up to 50% of cells positive for p27. The third group contained 40
20 specimens (37%) with p27 immunostaining detected in more than 50% of positive cells (referred to as high expressors). Analysis of the data utilizing such necessarily arbitrary cuts-off was highly statistically significant and, therefore, functionally operative. A significant negative relationship
25 was found between p27 expression and the histological grading ($p=0.01$) assigned to the specimen. In addition, p27 expression was associated significantly with overall survival. The median survival time in low expressor patients was 20 months as compared to 30 months in high expressor patients and
30 only 14 months in non expressor patients. The five year survival rate also differed statistically among the three groups ($p=.0012$). The low expressor group had an overall survival of 14% as compared to 25% for the high expressor group. All 12 non expressor patients were dead within 30
35 months following surgery. Accordingly, p27 levels served as a prognostic factor correlating with the overall survival time of the patients ($p=0.0012$). This correlation was strengthened

by the relationship found between p27 expression levels and the degree of malignancy (grading) ($p=0.01$).

Similar results have been obtained for p27 in a cohort of one hundred and forty-nine patients with carcinoma of the colon or rectum. Accordingly, in addition to non small cell lung cancer, the determination of p27 expression levels can be used in grading of other types of tumors and in prognosticating survival rates in cancer patients suffering from these tumors. The following nonlimiting examples are being provided to further illustrate the present invention.

EXAMPLES

Example 1: Population study

One hundred and eight non small cell lung cancer specimens were obtained from patients who underwent a surgical resection (lobectomy or pneumonectomy) in the departments of Thoracic Surgery of the V. Monaldi Hospital and of the II University of Naples (Italy) between 1988 and 1992. All specimens were from patients who had not received chemo- or radio-therapy prior to surgical resection. Outcome data were collected from hospital charts and from periodic interviews with patients and their relatives. The follow up period was 48 months from the date of surgery (for survivors). Patients who died of causes other than lung cancer were not included in the study. The mean age was 60.1. Gender was unevenly distributed with females accounting for only 7.4% of the population (8 patients). The histological diagnoses and classifications of the tumors were based on the WHO criteria (World Health Organization, 1981). The postsurgical pathologic TNM stage was determined according to the guidelines of the American Joint Committee on Cancer (Beahrs et al. (1992). 4th ed. Chicago, American Joint Committee on Cancer, 115-122). After surgical resection each tumor specimen was divided into two parts. The first part was frozen instantly for the extraction of RNAs and proteins. The second part was formalin fixed immediately and then paraffin

embedded for routine and immunohistochemical investigation.

Example 2: Immunohistochemistry

Immunohistochemistry sections from each specimen were
5 cut at 3-5 μm , mounted on glass and dried overnight at 37°C.
All sections were deparaffinized in xylene, rehydrated through
a graded alcohol series and washed in PBS. This buffer was
used for all subsequent washes and for dilution of the
antibodies. Sections were heated twice in a microwave oven
10 for 5 minutes, each at 700 W, in 10 mmole/L citrate buffer (pH
6); sequentially quenched in 0.5% hydrogen peroxide; and
blocked with diluted 10% normal horse anti-mouse serum (Vector
Laboratories, Burlingame, CA). A monoclonal antibody raised
against p27 (Transduction Laboratories) was used (dilution
15 1:100). The incubation time was 60 minutes at room
temperature. After washing in PBS, slides were incubated with
diluted horse anti-mouse biotinylated antibody (Vector
Laboratories) for 30 minutes at room temperature. All slides
were processed by the ABC method (Vector Laboratories) for 30
20 minutes at room temperature. Diaminobenzidine was used as the
final chromogen, and hematoxylin was used as the nuclear
counterstain. To evaluate the specificity of the antibody,
the protein used to generate it and an irrelevant protein were
both pre-adsorbed to the antibody. Negative controls for each
25 tissue section were prepared by leaving out the primary
antibody. Immunohistochemistry was performed on normal
bronchial specimens to establish the normal p27 expression
pattern of this tissue for use as the positive control. All
samples were processed under the same conditions. The
30 staining pattern of the protein was evaluated and scored for
the percentage of positive nuclei by two separate
pathologists: a score of 1 refers to less than 10% positive
cells; a score of 2 refers to a range of from 10% to 50%
positive cells; and a score of 3 refers to more than 50%
35 positive cells. At least 20 high power fields were chosen
randomly and 2000 cells were counted.

WO 98/51821

PCT/US98/09788

- 10 -

Example 3: Western Blot

One gram of each frozen lung cancer tissue sample was sectioned and quickly homogenized at 4°C in 250 mM NaCl, 50 mM Tris (pH 7.4), 5 mM EDTA, 0.1% (v/v) Triton X-100,TM 1 mM phenylmethylsulfonyl fluoride, 50 mM NaF, 0.5 mM Na₂VO₄, 10 mg/ml leupeptin and 50 mg/ml aprotinin. The homogenates were then spun at 13,000 x g at 4°C and total protein in the extracts was determined. Fifty milligrams of protein was denatured by boiling in 2X Laemmli sample buffer and separated
10 by electrophoresis in a 15% sodium dodecyl sulfate-polyacrylamide gel, followed by electrophoretic transfer of the proteins to a PVDF membrane (Millipore, Bedford, MA) in CAPS buffer (10 mM CAPS, 20% methanol, pH 11). The membrane was then blocked with 5% milk in TBS-T buffer (2
15 mM Tris, 13.7 mM NaCl, 0.1% Tween-20,TM pH 7.6) and washed in TBS-T. Primary antibody was incubated with the membrane in 3% milk and then washed in TBS-T. Sheep anti-mouse antibody coupled to horseradish peroxidase was incubated with the membrane and then washed in TBS-T. The presence of secondary
20 antibody bound to the membrane was detected using the ECL system (Dupont NEN, Boston, MA).

Example 4: Northern Blot Analysis

Cytoplasmic RNA was extracted using the RNAzol method (CINNA/BIOTECH, Friendswood, TX) after homogenization from
25 13 non small cell lung cancer samples (including samples representative of all the coded p27 expressing groups). Northern blots were performed in accordance with procedures described by Claudio et al. (1994). *Cancer Res* 54: 5556-5560.

Example 5: Degradation assay

30 One gram of each frozen human tissue sample (6 representative samples) was sectioned and quickly homogenized at 15,000 rpm in 1 ml of ice cold doubly distilled water. The sample was frozen and thawed 3 times. The lysate was spun down at 15,000 rpm for 45 minutes at 4°C. The supernatant was

SUBSTITUTE SHEET (RULE 26)

- 11 -

retrieved and frozen at -80°C. This method of preparation of total extract preserves ubiquitinating enzymes. Purified histidine-tagged p27 (0.06 ml) was incubated at 37°C for various time periods in 30 ml of degradation mix containing 5 100 mg of protein human tissue extracts, 50 mM Tris-HCL (pH 8.0), 5 mM MgCl₂, and 1 mM DTT, 2 mM ATP, 10 mM creatine phosphate and 5 mM ubiquitin. Degradation of p27 was analyzed by immunoblotting with p27 mAb. In order to remove the proteasome, tumor samples were ultracentrifuged at 100,000 g 10 for 6 hours in accordance with procedures described by Pagano M et al. (1995) *Science* 269: 682-685.

Example 6: Statistical analysis

Patient survival data were used to determine a correlation between p27 expression levels and overall survival 15 time. Survival curves were constructed using Kaplan-Meier analysis. Statistical significance of these data was measured by the Mantel-Cox test. Possible associations between the different variables of the analyzed tumor samples were tested by the chi square test.

What is claimed is:

1. A method of grading a non-small cell lung tumor comprising:

- 5 (a) obtaining a tissue sample from the tumor;
(b) measuring a level of p27 protein in the tissue sample;
(c) comparing the measured p27 protein level to establish control levels of p27 protein; and
10 (d) assigning a pathological grade to the tumor based upon the level of p27 protein in the tissue sample.

2. A method of prognosticating a survival time of a cancer patient afflicted with non-small cell lung carcinoma comprising:

- 15 (a) obtaining a tissue sample from a non-small cell lung tumor of the cancer patient;
(b) measuring a level of p27 protein in the tissue sample; and
20 (c) predicting a survival time of the cancer patient based upon the measured level of p27 protein, said protein level being indicative of patient survival time, wherein the measured level of p27 protein and the patient survival time is a positive relationship.

25

3. A method of grading a tumor comprising:

- (a) obtaining a tissue sample from a tumor;
(b) measuring a level of p27 protein in the tissue sample;
30 (c) comparing the measured p27 protein level to established control levels of p27 protein; and
(d) assigning a pathological grade to the tumor based upon the level of p27 protein in the tissue sample.

4. A method of prognosticating a survival time of a cancer patient comprising:

(a) obtaining a tissue sample from a tumor of a cancer patient;

5 (b) measuring a level of p27 protein in the tissue sample; and

(c) predicting a survival time of the cancer patient based upon the measured level of p27 protein, said protein level being indicative of patient survival time,
10 wherein the measured level of p27 protein and the patient survival time is a positive relationship.