




(86) **Date de dépôt PCT/PCT Filing Date:** 2011/03/02  
(87) **Date publication PCT/PCT Publication Date:** 2011/09/09  
(45) **Date de délivrance/Issue Date:** 2017/12/05  
(85) **Entrée phase nationale/National Entry:** 2012/08/16  
(86) **N° demande PCT/PCT Application No.:** JP 2011/055430  
(87) **N° publication PCT/PCT Publication No.:** 2011/108751  
(30) **Priorité/Priority:** 2010/03/03 (JP2010-046370)

(51) **Cl.Int./Int.Cl. A01N 37/38** (2006.01),  
**A01N 37/24** (2006.01), **A01N 37/30** (2006.01),  
**A01N 37/34** (2006.01), **A01N 37/36** (2006.01),  
**A01N 37/46** (2006.01), **A01N 43/40** (2006.01),  
**A01N 43/56** (2006.01), **A01N 43/76** (2006.01),  
**A01N 43/78** (2006.01), **A01N 43/80** (2006.01),  
**A01N 43/84** (2006.01), **A01N 45/02** (2006.01),  
**A01N 47/32** (2006.01), **A01N 53/00** (2006.01)

(72) **Inventeurs/Inventors:**  
KIGUCHI, SO, JP;  
TANAKA, SOICHI, JP;  
OZAWA, MAYUKO, JP;  
IWATA, ATSUSHI, JP

(73) **Propriétaire/Owner:**

(54) **Titre : COMPOSITIONS DESTINEES A CONTROLER UNE MALADIE FONGIQUE DE VEGETAUX RENFERMANT DE LA  
MANDESTROBINE ET DU BOSCALIDE ET METHODE DE CONTROLER DES MALADIES FONGIQUES DE VEGETAUX**  
(54) **Title: PLANT FUNGAL DISEASE CONTROLLING COMPOSITION CONTAINING MANDESTROBIN AND BOSCALID AND  
METHOD FOR CONTROLLING PLANT FUNGAL DISEASES**



(1)

(57) **Abrégé/Abstract:**


The present invention provides a composition having an excellent controlling activity on plant disease. The composition comprising the compound represented by the formula 1 (see formula 1) and one or more carboxamide fungicidal compound selected from the group (A) shows an excellent controlling activity on a plant disease; group (A): a group consisting of thifluzamide, oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, and bixafen. Of particular suitability is the carboxamide fungicidal compound boscalid.

(73) **Propriétaires(suite)/Owners(continued):**SUMITOMO CHEMICAL COMPANY, LIMITED, JP

(74) **Agent:** KIRBY EADES GALE BAKER

## ABSTRACT

The present invention provides a composition having an excellent controlling activity on plant disease. The composition comprising the compound represented by the formula (1)



and one or more carboxamide fungicidal compound selected from the group (A) shows an excellent controlling activity on a plant disease;

group (A): a group consisting of thifluzamide, oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, and bixafen. Of particular suitability is the carboxamide fungicidal compound boscalid.

PLANT FUNGAL DISEASE CONTROLLING COMPOSITION  
CONTAINING MANDESTROBIN AND BOSCALID AND  
METHOD FOR CONTROLLING PLANT FUNGAL DISEASES

5

Technical Field

The present invention relates to a plant disease controlling composition and a method for controlling a plant disease.

10


Background Art

Hitherto, there has been provided compounds as an active ingredient for a composition for controlling plant disease (see e.g., WO 99/24413 pamphlet; WO 03/070705 pamphlet; and The Pesticide Manual - 15th edition (BCPC published) ISBN 1901396188).

15

Also there has been provided a compound of the formula

(1):



20 (see e.g., WO 95/27693 pamphlet and WO 02/10101 pamphlet).


## Disclosure of Invention

An object of the present invention is to provide a composition having an excellent control effect on a plant disease.

5 The present inventors have intensively studied to find a composition having an excellent control effect on a plant disease. As a result, they have found that a composition comprising a compound represented by the formula (1) and one or more carboxamide fungicidal compound selected from  
10 the following group (A) shows a synergistic activity, and thus has an excellent control effect on a plant disease, and therefore the present invention has been completed.

The present invention provides:

[1] A plant fungal diseases controlling composition  
15 comprising a compound represented by the formula (1):




and one or more carboxamide fungicidal compound selected from the following group (A):

group (A): a group consisting of boscalid.

[2] The plant fungal diseases controlling composition according to the above [1], wherein a weight ratio of the compound represented by the formula (1) to the carboxamide fungicidal compound is that of the compound represented by the formula (1)/the carboxamide fungicidal compound = 0.0125/1 to 500/1.

[3] The plant fungal diseases controlling composition according to the above [1] or [2], wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

[4] A method for controlling plant fungal diseases which comprises applying an effective amount of each of the compound of the formula (1):



and one or more carboxamide fungicidal compound selected from the following group (A) to a plant or a soil for cultivating the plant,

group (A): a group consisting of boscalid.


[5] The method for controlling plant fungal diseases comprising applying a compound represented by formula (1)

and one or more carboxamide fungicidal compound selected from group (A) to a seed.

[6] The method for controlling plant fungal diseases according to the above [4] or [5], wherein a weight ratio of the compound represented by the formula (1) to the carboxamide fungicidal compound is that of the compound represented by the formula (1)/the carboxamide fungicidal compound = 0.0125/1 to 500/1.

[7] The method for controlling plant fungal diseases according to any one of the above [4] to [6], wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

[8] A use of a combination of the compound represented by the formula (1):



15

and one or more carboxamide fungicidal compound selected from the following group (A) for controlling plant fungal diseases,


group (A): a group consisting of boscalid.

20

The present invention enables to control a plant disease.

#### Best Mode for Carrying Out the Invention

5 A plant disease controlling composition of the present invention (hereinafter, referred to as a composition of the present invention) comprises a compound represented by the formula (1):



10 (hereinafter, referred to as an amide compound of the present invention) and one or more carboxamide compound selected from the following group (A) (hereinafter, referred to as a carboxamide compound of the present invention),

15 group (A): a group consisting of thifluzamide, oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, and bixafen.

20 The present amide compounds are those described in for example, WO 95/27693 pamphlet and WO 02/10101 pamphlet, and thus can be prepared according to the method described

therein.

The present amide compound has one asymmetric carbon. Herein, a compound represented by the formula (1) wherein an enantiomer having R- absolute configuration is enriched  
5 is referred to as the amide compound having R- absolute configuration.

The present amide compound encompasses the following compounds:

a compound represented by the formula (1) which  
10 contains an enantiomer having R- absolute configuration in .70% and more;

a compound represented by the formula (1) which contains an enantiomer having R- absolute configuration in  
90% and more;

15 a compound represented by the formula (1) which contains an enantiomer having R- absolute configuration in 95% and more.

Thi fluzamide, oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil,  
20 tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, and bixafen used in the present invention are all known compounds. Thi fluzamide, oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, mepronil,  
25 tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid,

diclocymet, mandipropamid, dimethomorph, flumorph, and penthiopyrad and are described in for example, "The PESTICIDE MANUAL - 15th EDITION (BCPC published) ISBN 1901396188", pages 1119, 847, 871, 76, 473, 580, 676, 729, 5 1080, 121, 535, 533, 167, 340, 705, 377, 531 and 877 respectively. Isotianil and bixafen are described in for example, WO 99/24413 pamphlet and WO 03/070705 pamphlet respectively. These compounds are either commercially available, or can be prepared by a known method.

10 The weight ratio of the present amide compound to the present carboxamide compound in the composition of the present invention is usually that of the present amide compound/the present carboxamide compound = 0.0125/1 to 500/1, preferably 0.025/1 to 100/1, and more preferably 15 0.1/1 to 10/1.

Although the composition of the present invention may be a mixture as itself of the present amide compound and the present carboxamide compound, the composition of the present invention is usually prepared by mixing the present 20 amide compound, the present carboxamide compound and an inert carrier, and if necessary, adding a surfactant or other pharmaceutical additive, and then formulating into the form of oil solution, emulsifiable concentrate, flowable formulation, wettable powder, granulated wettable 25 powder, dust formulation, granules and so on. Such

formulations can be used by itself or with an addition of another inert component as an agent for controlling a plant disease.

Usually, the composition of the present invention can contain 0.1 to 99 % by weight, preferably 0.2 to 90 % by weight, and more preferably 1 to 80 % by weight of the present amide compound and the present carboxamide compound in total.

Examples of a solid carrier used on the formulation include finely-divided powder or particles of clay consisting of minerals (e.g., kaolin clay, attapulgite clay, bentonite, montmorillonite, acid clay, pyrophyllite, talc, diatomaceous earth, or calcite), natural organic substances (e.g., corncob powder, or walnut shell powder), synthetic organic substances (e.g., urea), salts (e.g., calcium carbonate, or ammonium sulfate), synthetic inorganic substances (e.g., synthetic hydrous silicon oxide) and so on. Examples of a liquid carrier include aromatic hydrocarbons (e.g., xylene, alkyl benzene, or methylnaphthalene), alcohols (e.g., 2-propanol, ethylene glycol, propylene glycol, or ethylene glycol monoethyl ether), ketones (e.g., acetone, cyclohexanone, or isophorone), vegetable oils (e.g., soybean oil, or cotton oils), petroleum-derived aliphatic hydrocarbons, esters, dimethylsulfoxide, acetonitrile and water.

Examples of the surfactant include anionic surfactant (e.g., alkyl sulfate salts, alkylaryl sulfate salts, dialkyl sulfosuccinate salts, polyoxyethylene alkylaryl ether phosphates, lignin sulfonate, or naphthalenesulfonate formaldehyde polycondensation), nonionic surfactant (e.g., polyoxyethylene alkylaryl ether, polyoxyethylene alkyl polyoxypropylene block copolymer, or sorbitan fatty acid ester) and cationic surfactant (e.g., alkyltrimethyl ammonium salts).

Examples of the other pharmaceutical additives include water-soluble polymer (e.g., polyvinyl alcohol, or polyvinyl pyrrolidone), polysaccharides (e.g. arabic gum, alginic acid and salts thereof, CMC (carboxymethyl-cellulose), or xanthan gum), inorganic substances (e.g., aluminum magnesium silicate, or alumina-sol), antiseptic agent, coloring agent, and PAP (isopropyl acid phosphate), and stabilizing agent (e.g., BHT).

The composition of the present invention can also be prepared by separately formulating the present amide compound and the present carboxamide compound into different formulations by the above procedures, if necessary, further diluting each of them with water, thereafter, mixing the separately prepared different formulations or the dilute solutions.

The composition of the present invention may further contain one or more other fungicides and/or insecticides.

The composition of the present invention is used to control a plant disease by applying it to a plant or a soil  
5 for cultivating the plant.

The plant diseases which can be controlled by the present invention are exemplified below:

Rice diseases: blast (*Magnaporthe oryzae*),  
helminthosporium leaf spot (*Cochliobolus miyabeanus*),  
10 sheath blight (*Rhizoctonia solani*) and bakanae disease  
(*Gibberella fujikuroi*);

Diseases of barley, wheat, oats and rye: powdery  
mildew (*Erysiphe graminis*), Fusarium head blight (*Fusarium  
graminearum*, *F. avenaceum*, *F. culmorum*, *F. asiaticum*,  
15 *Microdochium nivale*), rust (*Puccinia striiformis*,  
*P. graminis*, *P. recondite*, *P. hordei*), snow blight  
(*Typhula* sp., *Micronectriella nivalis*), loose smut  
(*Ustilago tritici*, *U. nuda*), bunt (*Tilletia caries*),  
eyespot (*Pseudocercospora herpotrichoides*), scald  
20 (*Rhynchosporium secalis*), leaf blotch (*Septoria tritici*),  
glume blotch (*Leptosphaeria nodorum*) and net blotch  
(*Pyrenophora teres Drechsler*);

Citrus diseases: melanose (*Diaporthe citri*), scab  
(*Elsinoe fawcetti*) and Penicillium rot (*Penicillium  
25 digitatum*, *P. italicum*);

Apple diseases: blossom blight (*Monilinia mali*),  
canker (*Valsa ceratosperma*), powdery mildew (*Podosphaera*  
*leucotricha*), Alternaria leaf spot (*Alternaria alternata*  
apple pathotype), scab (*Venturia inaequalis*), bitter rot  
5 (*Colletotrichum acutatum*) and late blight (*Phytophthora*  
*cactorum*);

Pear diseases: scab (*Venturia nashicola*, *V. pirina*),  
black spot (*Alternaria alternata* Japanese pear pathotype),  
rust (*Gymnosporangium haraeum*) and late blight  
10 (*Phytophthora cactorum*);

Peach diseases: brown rot (*Monilinia fructicola*),  
scab (*Cladosporium carpophilum*) and Phomopsis rot  
(*Phomopsis* sp.);

Grapes diseases: anthracnose (*Elsinoe ampelina*), ripe  
15 rot (*Glomerella cingulata*), powdery mildew (*Uncinula*  
*necator*), rust (*Phakopsora ampelopsidis*), black rot  
(*Guignardia bidwellii*), downy mildew (*Plasmopara viticola*)  
and Gray mold (*Botrytis cinerea*);

Diseases of Japanese persimmon: anthracnose  
20 (*Gloeosporium kaki*) and leaf spot (*Cercospora kaki*,  
*Mycosphaerella nawae*);

Diseases of gourd family: anthracnose (*Colletotrichum*  
*lagenarium*), powdery mildew (*Sphaerotheca fuliginea*), gummy  
stem blight (*Mycosphaerella melonis*), Fusarium wilt  
25 (*Fusarium oxysporum*), downy mildew (*Pseudoperonospora*

*cubensis*), Phytophthora rot (*Phytophthora* sp.) and damping-off (*Pythium* sp.);

Tomato diseases: early blight (*Alternaria solani*), leaf mold (*Cladosporium fulvum*) and late blight  
5 (*Phytophthora infestans*);

Egg plant disease: brown spot (*Phomopsis vexans*) and powdery mildew (*Erysiphe cichoracearum*);

Diseases of Cruciferous Vegetables: *Alternaria* leaf spot (*Alternaria japonica*), white spot (*Cercospora brassicae*), clubroot (*Plasmodiophora brassicae*), and downy  
10 mildew (*Peronospora parasitica*);

Rapeseed diseases: *Sclerotinia* rot (*Sclerotinia sclerotiorum*), black spot (*Alternaria brassicae*), powdery mildew (*Erysiphe cichoracearum*), blackleg (*Leptosphaeria maculans*);  
15

Welsh onion diseases: rust (*Puccinia allii*);

Soybean diseases: purple seed stain (*Cercospora kikuchii*), Sphaceloma scab (*Elsinoe glycines*), pod and stem blight (*Diaporthe phaseolorum* var. *sojae*), rust (*Phakopsora pachyrhizi*) and phytophthora stem rot (*Phytophthora sojae*);  
20

Adzuki-bean diseases: Gray mold (*Botrytis cinerea*), *Sclerotinia* rot (*Sclerotinia sclerotiorum*);

Kidney bean diseases: Gray mold (*Botrytis cinerea*), *Sclerotinia* rot (*Sclerotinia sclerotiorum*), anthracnose  
25 (*Colletotrichum lindemthianum*);

Peanut diseases: leaf spot (*Cercospora personata*),  
brown leaf spot (*Cercospora arachidicola*) and southern  
blight (*Sclerotium rolfsii*);

Garden pea diseases: powdery mildew (*Erysiphe pisi*);

5 Potato diseases: early blight (*Alternaria solani*) and  
late blight (*Phytophthora infestans*);

Strawberry diseases: powdery mildew (*Sphaerotheca  
humuli*);

10 Tea diseases: net blister blight (*Exobasidium  
reticulatum*), white scab (*Elsinoe leucospila*), gray blight  
(*Pestalotiopsis* sp.) and anthracnose (*Colletotrichum theae-  
sinensis*);

Cotton diseases: fusarium wilt (*Fusarium oxysporum*),  
damping-off (*Rhizoctonia solani*);

15 Tobacco diseases: brown spot (*Alternaria longipes*),  
powdery mildew (*Erysiphe cichoracearum*), anthracnose  
(*Colletotrichum tabacum*), downy mildew (*Peronospora  
tabacina*) and late blight (*Phytophthora nicotianae*);

20 Sugar beet diseases: Cercospora leaf spot (*Cercospora  
beticola*), leaf blight (*Thanatephorus cucumeris*), Root rot  
(*Aphanidermatum cochlioides*);

Rose diseases: black spot (*Diplocarpon rosae*) and  
powdery mildew (*Sphaerotheca pannosa*);

25 Chrysanthemum diseases: leaf blight (*Septoria  
chrysanthemi-indici*) and white rust (*Puccinia horiana*);

Various plant diseases: diseases caused by *Pythium* spp. (*Pythium aphanidermatum*, *Pythium debarianum*, *Pythium graminicola*, *Pythium irregulare*, *Pythium ultimum*), Gray mold (*Botrytis cinerea*), Sclerotinia rot (*Sclerotinia sclerotiorum*),

Japanese radish diseases: *Alternaria* leaf spot (*Alternaria brassicicola*);

Turfgrass diseases: dollar spot (*Sclerotinia homeocarpa*), brown patch and large patch (*Rhizoctonia solani*); and

Banana diseases: Sigatoka disease (*Mycosphaerella fijiensis*, *Mycosphaerella musicola*, *Pseudocercospora musae*).

Examples of the plants to which the composition of the present invention can be applied are as follows:

Crops: corn, rice, wheat, barley, rye, oat, sorghum, cotton, soybean, adzuki-bean, kidney bean, peanut, buckwheat, beet, rapeseed, sunflower, sugar cane, and tobacco, etc.;

Vegetables: solanaceous vegetables (eggplant, tomato, pimento, pepper, and potato, etc.), cucurbitaceous vegetables (cucumber, pumpkin, zucchini, watermelon, melon, and squash, etc.), cruciferous vegetables (Japanese radish, white turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, leaf mustard, broccoli, and cauliflower, etc.), asteraceous vegetables (burdock, crown daisy, artichoke,

and lettuce, etc.), liliaceous vegetables (green onion, onion, garlic, and asparagus), ammiaceous vegetables (carrot, parsley, celery, and parsnip, etc.), chenopodiaceous vegetables (spinach, and Swiss chard, etc.),  
 5 lamiaceous vegetables (*Perilla frutescens*, mint, and basil, etc.), strawberry, sweet potato, *Dioscorea japonica*, and colocasia, etc.;

Flowers;

Foliage plants;

10 Turfgrass;

Fruits: pomaceous fruits (apple, pear, Japanese pear, Chinese quince, and quince, etc.), stone fleshy fruits (peach, plum, nectarine, *Prunus mume*, cherry fruit, apricot, and prune, etc.), citrus fruits (*Citrus unshiu*, orange, lemon, lime, and grapefruit, etc.), nuts (chestnut, walnuts, hazelnuts, almond, pistachio, cashew nuts, and macadamia nuts, etc.), berries (blueberry, cranberry, blackberry, and raspberry, etc.), grape, kaki fruit, olive, Japanese plum, banana, coffee, date palm, and coconuts, etc.; and

20 Trees other than fruit trees: tea, mulberry, flowering plant, roadside trees (ash, birch, dogwood, Eucalyptus, *Ginkgo biloba*, lilac, maple, *Quercus*, poplar, Judas tree, *Liquidambar formosana*, plane tree, zelkova, Japanese arborvitae, fir wood, hemlock, juniper, *Pinus*, *Picea*, and  
 25 *Taxus cuspidate*), etc.

The aforementioned "plants" include plants to which resistance has been imparted by genetic recombination.

Exemplary embodiments of the composition of the present invention are as follows:

5 a composition comprising the present amide compound and thifluzamide wherein a weight ratio thereof is that of the present amide compound/thifluzamide = 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and thifluzamide wherein a weight ratio thereof is that of the present amide compound/thifluzamide = 0.025/1 to 100/1;

a composition comprising the present amide compound and thifluzamide wherein a weight ratio thereof is that of the present amide compound/thifluzamide = 0.1/1 to 10/1;

15 a composition comprising the present amide compound and oxadixyl wherein a weight ratio thereof is that of the present amide compound/oxadixyl = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and oxadixyl wherein a weight ratio thereof is that of the present amide compound/oxadixyl = 0.025/1 to 100/1;

a composition comprising the present amide compound and oxadixyl wherein a weight ratio thereof is that of the present amide compound/oxadixyl = 0.1/1 to 10/1;

25 a composition comprising the present amide compound and pencycuron wherein a weight ratio thereof is that of

the present amide compound/pencycuron = 0.0125/1 to 500/1;

a composition comprising the present amide compound and pencycuron wherein a weight ratio thereof is that of the present amide compound/pencycuron = 0.025/1 to 100/1;

5 a composition comprising the present amide compound and pencycuron wherein a weight ratio thereof is that of the present amide compound/pencycuron = 0.1/1 to 10/1;

a composition comprising the present amide compound and benalaxyl-M wherein a weight ratio thereof is that of the present amide compound/benalaxyl-M = 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and benalaxyl-M wherein a weight ratio thereof is that of the present amide compound/benalaxyl-M = 0.025/1 to 100/1;

a composition comprising the present amide compound and benalaxyl-M wherein a weight ratio thereof is that of the present amide compound/benalaxyl-M = 0.1/1 to 10/1;

15 a composition comprising the present amide compound and fenhexamid wherein a weight ratio thereof is that of the present amide compound/fenhexamid = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and fenhexamid wherein a weight ratio thereof is that of the present amide compound/fenhexamid = 0.05/1 to 20/1;

a composition comprising the present amide compound and fenhexamid wherein a weight ratio thereof is that of the present amide compound/fenhexamid M = 0.2/1 to 5/1;

25

a composition comprising the present amide compound and furametpyr wherein a weight ratio thereof is that of the present amide compound/furametpyr = 0.0125/1 to 500/1;

5 a composition comprising the present amide compound and furametpyr wherein a weight ratio thereof is that of the present amide compound/furametpyr = 0.025/1 to 100/1;

a composition comprising the present amide compound and furametpyr wherein a weight ratio thereof is that of the present amide compound/furametpyr = 0.1/1 to 10/1;

10 a composition comprising the present amide compound and isopyrazam wherein a weight ratio thereof is that of the present amide compound/isopyrazam = 0.0125/1 to 500/1;

a composition comprising the present amide compound and isopyrazam wherein a weight ratio thereof is that of the present amide compound/isopyrazam = 0.05/1 to 20/1;

15 a composition comprising the present amide compound and isopyrazam wherein a weight ratio thereof is that of the present amide compound/isopyrazam = 0.5/1 to 5/1;

a composition comprising the present amide compound and isotianil wherein a weight ratio thereof is that of the present amide compound/isotianil = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and isotianil wherein a weight ratio thereof is that of the present amide compound/isotianil = 0.025/1 to 100/1;

a composition comprising the present amide compound and isotianil wherein a weight ratio thereof is that of the present amide compound/isotianil = 0.1/1 to 10/1;

5 a composition comprising the present amide compound and mepronil wherein a weight ratio thereof is that of the present amide compound/mepronil = 0.0125/1 to 500/1;

a composition comprising the present amide compound and mepronil wherein a weight ratio thereof is that of the present amide compound/mepronil = 0.025/1 to 100/1;

10 a composition comprising the present amide compound and mepronil wherein a weight ratio thereof is that of the present amide compound/mepronil = 0.1/1 to 10/1;

a composition comprising the present amide compound and tecloftalam wherein a weight ratio thereof is that of  
15 the present amide compound/tecloftalam = 0.0125/1 to 500/1;

a composition comprising the present amide compound and tecloftalam wherein a weight ratio thereof is that of the present amide compound/tecloftalam = 0.025/1 to 100/1;

20 a composition comprising the present amide compound and tecloftalam wherein a weight ratio thereof is that of the present amide compound/tecloftalam = 0.1/1 to 10/1;

a composition comprising the present amide compound and boscalid wherein a weight ratio thereof is that of the present amide compound/boscalid = 0.0125/1 to 500/1;

a composition comprising the present amide compound and boscalid wherein a weight ratio thereof is that of the present amide compound/boscalid = 0.025/1 to 100/1;

5 a composition comprising the present amide compound and boscalid wherein a weight ratio thereof is that of the present amide compound/boscalid = 0.1/1 to 10/1;

a composition comprising the present amide compound and fluopyra wherein a weight ratio thereof is that of the present amide compound/fluopyra = 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and fluopyra wherein a weight ratio thereof is that of the present amide compound/fluopyra = 0.05/1 to 20/1;

a composition comprising the present amide compound and fluopyra wherein a weight ratio thereof is that of the present amide compound/fluopyra = 0.2/1 to 5/1;

15 a composition comprising the present amide compound and fluopicolide wherein a weight ratio thereof is that of the present amide compound/fluopicolide = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and fluopicolide wherein a weight ratio thereof is that of the present amide compound/fluopicolide = 0.025/1 to 100/1;

a composition comprising the present amide compound and fluopicolide wherein a weight ratio thereof is that of the present amide compound/fluopicolide = 0.1/1 to 10/1;

25

a composition comprising the present amide compound and carpropamid wherein a weight ratio thereof is that of the present amide compound/carpropamid = 0.0125/1 to 500/1;

5 a composition comprising the present amide compound and carpropamid wherein a weight ratio thereof is that of the present amide compound/carpropamid = 0.025/1 to 100/1;

a composition comprising the present amide compound and carpropamid wherein a weight ratio thereof is that of the present amide compound/carpropamid = 0.1/1 to 10/1;

10 a composition comprising the present amide compound and diclocymet wherein a weight ratio thereof is that of the present amide compound/diclocymet = 0.0125/1 to 500/1;

a composition comprising the present amide compound and diclocymet wherein a weight ratio thereof is that of  
15 the present amide compound/diclocymet = 0.025/1 to 100/1;

a composition comprising the present amide compound and diclocymet wherein a weight ratio thereof is that of the present amide compound/diclocymet = 0.1/1 to 10/1;

20 a composition comprising the present amide compound and mandipropamid wherein a weight ratio thereof is that of the present amide compound/mandipropamid = 0.0125/1 to 500/1;

a composition comprising the present amide compound and mandipropamid wherein a weight ratio thereof is that of  
25 the present amide compound/mandipropamid = 0.025/1 to

100/1;

a composition comprising the present amide compound and mandipropamid wherein a weight ratio thereof is that of the present amide compound/mandipropamid = 0.1/1 to 10/1;

5 a composition comprising the present amide compound and dimethomorph wherein a weight ratio thereof is that of the present amide compound/dimethomorph = 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and dimethomorph wherein a weight ratio thereof is that of the present amide compound/dimethomorph = 0.025/1 to 100/1;

a composition comprising the present amide compound and dimethomorph wherein a weight ratio thereof is that of the present amide compound/dimethomorph = 0.1/1 to 10/1;

15 a composition comprising the present amide compound and flumorph wherein a weight ratio thereof is that of the present amide compound/flumorph = 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and flumorph wherein a weight ratio thereof is that of the present amide compound/flumorph = 0.025/1 to 100/1;

a composition comprising the present amide compound and flumorph wherein a weight ratio thereof is that of the present amide compound/flumorph = 0.1/1 to 10/1;

25 a composition comprising the present amide compound and penthiopyrad wherein a weight ratio thereof is that of

the present amide compound/penthiopyrad = 0.0125/1 to 500/1;

a composition comprising the present amide compound and penthiopyrad wherein a weight ratio thereof is that of the present amide compound/penthiopyrad = 0.05/1 to 20/1;

a composition comprising the present amide compound and penthiopyrad wherein a weight ratio thereof is that of the present amide compound/penthiopyrad = 0.2/1 to 5/1;

a composition comprising the present amide compound and bixafen wherein a weight ratio thereof is that of the present amide compound/bixafen = 0.0125/1 to 500/1;

a composition comprising the present amide compound and bixafen wherein a weight ratio thereof is that of the present amide compound/bixafen = 0.05/1 to 20/1; and

a composition comprising the present amide compound and bixafen wherein a weight ratio thereof is that of the present amide compound/bixafen = 0.2/1 to 5/1.

The method for controlling a plant disease of the present invention (hereinafter, referred to as the method for controlling of the present invention) is carried out by applying an effective amount of each of the present amide compound and the present carboxamide compound to the plants or the soil for cultivating the plant.

Such plants may be, for example, foliage of plant, seeds of plant, or bulbs of plant. The bulbs herein are

intended to mean bulb, corm, rootstock, tubera, tuberous root and rhizophore.

In the method for controlling of the present invention, the present amide compound and the present carboxamide  
5 compound may be applied separately around the same time to the plant or the soil for cultivating the plant, but is usually applied as the composition of the present invention in terms of application convenience.

In the method for controlling of the present invention,  
10 examples of the method of applying the present amide compound and the carboxamide compound include foliage treatment, soil treatment, root treatment and seed treatment.

Such foliage treatment includes, for example, a method  
15 of applying the composition of the present invention to a surface of the plant to be cultivated by a foliage application or a stem application.

Such root treatment includes, for example, a method of soaking a whole or a root of the plant into a medicinal  
20 solution comprising the present amide compound and the present carboxamide compound, and a method of attaching a solid formulation comprising the present amide compound, the present carboxamide compound and the solid carrier to a root of the plant.

Such soil treatment includes, for example, soil broadcast, soil incorporation, and irrigation of the medicinal solution to a soil.

Such seed treatment includes, for example, an application of the composition of the present invention to a seed or a bulb of the plant to be prevented from the plant disease, specifically, for example, a spray treatment by spraying a suspension of the composition of the present invention in a mist form to a surface of a seed or a surface of a bulb, a smear treatment by smearing the wettable powder, the emulsifiable concentrate or the flowable formulation of the composition of the present invention with additional small amounts of water or as itself to a seed or a bulb, an immersion treatment of a seed into a solution of the composition of the present invention for a given time, a film-coating treatment, and a pellet-coating treatment.

Each dose of the present amide compound and the present carboxamide compound in the method for controlling of the present invention may be varied depending on a kind of plant to be treated, a kind or a frequency of an occurrence of a plant disease as a control subject, a dosage form, a treatment period, a treatment method, a treatment site, a climate condition, etc. In case of an application to a foliage of the plant or a soil for

cultivating the plant, a total amount of the present amide compound and the carboxamide compound is usually 1 to 500 g, preferably 2 to 200 g, and more preferably 10 to 100 g, per 1000 m<sup>2</sup>. Each dose of the present amide compound and the present carboxamide compound in the treatment for seed is usually 0.001 to 10 g, and preferably 0.01 to 1 g, per 1kg of seeds.

The emulsifiable concentrate, the wettable powder or the flowable formulation, etc., is usually applied by dilution with water, and then spreading them. In this case, usually, each concentration of the present amide compound and the present carboxamide compound contain 0.0005 to 2% by weight, and preferably 0.005 to 1% by weight of the present amide compound and the present carboxamide compound in total. The dust formulation or the granular formulation, etc, is usually applied as itself without dilution.

#### EXAMPLES

Next, the present invention is described in more detail below by the following examples including formulation examples and a test example, but the present invention should not be construed to be limited thereto.

The formulation examples are given below. It is to be noted that in the formulation examples, the term ''part'' indicates ''part by weight''.

## Formulation 1

5 Parts of the present amide compound, 5 parts of thifluzamide, 35 parts of a mixture of white carbon and polyoxyethylene alkylether sulfate ammonium salts (weight ratio 1:1), and 55 parts of water were mixed and the resulting solution was then subjected to fine grinding according to a wet grinding method, so as to obtain a flowable formulation. The same above operations were carried out with oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, or bixafen instead of thifluzamide, so as to obtain various types of flowable formulations.

15

## Formulation 2

10 Parts of the present amide compound, 5 parts of thifluzamide and 1.5 parts of sorbitan trioleate were mixed into 28 parts of an aqueous solution that contained 2 parts of polyvinyl alcohol, and the mixed solution was then subjected to fine grinding according to a wet grinding method. Thereafter, 45.50 parts of an aqueous solution that contained 0.05 parts of xanthan gum and 0.1 part of aluminum magnesium silicate was added to the resultant product, and 10 parts of propylene glycol was further added

25

thereto. The obtained mixture was blended by stirring, so as to obtain the flowable formulation. The same above operations were carried out with oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, or bixafen instead of thifluzamide, so as to obtain various types of flowable formulations.

10           Formulation 3

10           10 Parts of the present amide compound, 40 parts of thifluzamide, 3 parts of calcium lignosulfonate, 2 parts of sodium lauryl sulfate, and 45 parts of synthetic hydrous silicon oxide were fully crushed and mixed, so as to obtain wettable powders. The same above operations were carried out with oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isopyrazam, isotianil, mepronil, tecloftalam, boscalid, fluopyram, fluopicolide, carpropamid, diclocymet, mandipropamid, dimethomorph, flumorph, penthiopyrad, or bixafen instead of thifluzamide, so as to obtain various types of wettable powders.

The test examples are given below.

Test Example 1

25           True leaf of cucumber is punched out with cork borer to 13mm in diameter to prepare a leaf disk. In 24 well

microwell plate that is dispensed with 1ml 0.8% water agar, the leaf disk is placed such that the upper side of the leaf is in an upward direction. Thereto is added 20 micro liter a testing solution prepared by mixing a dimethyl sulfoxide solution of the present compound (racemate) and a dimethyl sulfoxide solution of isopyrazam, boscalid or penthiopyrad to a predetermined concentration to treat the leaf disk.

After confirming that the testing medical solution is dried, conidium of gray mold fungus (*Botrytis cinerea*) is suspended into potato dextrose broth (DIFCO) in a density of about  $10^5$  conidium/mL and is then subjected to a spray inoculation. After leaving the leaf disk to stand in a growth chamber set up at 15°C for four days, an onset area on the leaf is measured and then calculated an onset area rate (hereinafter, referred to as an onset area rate of treated group).

The same operation is carried out with 20 micro liter water instead of 20 micro liter a testing medicine solution to calculate an onset area rate (hereinafter, referred to an onset area rate of non-treated group).

A preventive value is calculated from the above onset area rate of treated group and the onset area rate of non-treated group by the following equation:

$$\text{Preventive value (\%)} = 100 \times (A-B)/A$$

wherein

A: an onset area rate of treated group

B: an onset area rate of non-treated group

5 The results are shown in Tables 1, 2 and 3.

Table 1

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	isopyrazam	
1	2.5	0.5	95
2	1.0	5.0	97.5

Table 2

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	boscalid	
1	2.5	0.5	100
2	1.0	5.0	100

10 Table 3

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	penthioopyrad	
1	2.5	0.5	100
2	1.0	5.0	92.5

#### Test Example 2

The same operations as described in Test Example 1 are carried out with oxadixyl, pencycuron, benalaxyl-M, 15 fenhexamid, furametpyr, isotianil, fluopyram, diclocymet or dimethomorph instead of isopyrazam, boscalid or penthiopyrad, so as to calculate respective preventive

values.

Also for comparison, the same operations as described in Test Example 1 are carried out with the exception that the testing medicine solution is substituted with a predetermined concentration of each dimethyl sulfoxide solution of the present compound (racemate), oxadixyl, pencycuron, benalaxyl-M, fenhexamid, furametpyr, isotianil, fluopyram, diclocymet or dimethomorph, so as to calculate respective preventive values.

The results are shown in Tables 4 to 12.

Table 4

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	oxadixyl	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	10
	-	5.0	15

Table 5

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	pencycuron	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	10
	-	5.0	15

Table 6

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	benalaxyl-M	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	10
	-	5.0	15

Table 7

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	fenhexamid	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	20
	-	5.0	57

5 Table 8

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	furametpyr	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	15
	-	5.0	20

Table 9

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	isotianil	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	10
	-	5.0	25

Table 10

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	fluopyram	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	35
	-	5.0	45

5 Table 11

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	diclocymet	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56
	1.0	-	46
	-	0.5	10
	-	5.0	15

Table 12

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	dimethomorph	
1	2.5	0.5	100
2	1.0	5.0	100
	2.5	-	56

	1.0	-	46
	-	0.5	10
	-	5.0	15

Next, the Reference Examples are given below.

#### Reference Examples

For comparison, the same operations as described in  
 5 Test Example 1 are carried out with the exception that the  
 testing medicine solution is substituted with a  
 predetermined concentration of each dimethyl sulfoxide  
 solution of isopyrazam, boscalid or penthiopyrad, so as to  
 calculate respective preventive values.

10 The results are shown in Tables 13 to 15.

Table 13

	treatment concentration (ppm)	preventive value (%)
	isopyrazam	
	0.5	20
	5.0	40

Table 14


	treatment concentration (ppm)	preventive value (%)
	boscalid	
	0.5	49
	5.0	56

15 Table 15

	treatment concentration (ppm)	preventive value (%)
	penthiopyrad	
	0.5	39
	5.0	45

## CLAIMS

1. A plant fungal diseases controlling composition comprising a compound represented by the formula (1):




5

and a carboxamide fungicidal compound, boscalid.

2. The plant fungal diseases controlling composition according to claim 1, wherein a weight ratio of the compound represented by the formula (1) to the carboxamide fungicidal compound is that of the compound represented by the formula (1)/the carboxamide fungicidal compound = 0.0125/1 to 500/1.


3. The plant fungal diseases controlling composition according to claim 1 or 2, wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

4. A method for controlling plant fungal diseases which comprises applying an effective amount of each of the compound of the formula (1):



and a carboxamide fungicidal compound, boscalid, to a plant or a soil for cultivating the plant.

5. A method for controlling plant fungal diseases  
 5 comprising applying a compound represented by formula (1):




and a carboxamide fungicidal compound, boscalid, to a seed.


6. The method for controlling plant fungal diseases according to claim 4 or 5, wherein a weight ratio of the  
 10 compound represented by the formula (1) to the carboxamide fungicidal compound is that of the compound represented by the formula (1)/the carboxamide fungicidal compound = 0.0125/1 to 500/1.

7. The method for controlling plant fungal diseases  
 15 according to any one of claims 4 to 6, wherein the compound represented by the formula (1) is that represented by the formula (1) having R- absolute configuration.

8. A use of a combination of the compound represented by the formula (1):



and a carboxamide fungicidal compound, boscalid, for  
5 controlling plant fungal diseases.



(1)