




(86) Date de dépôt PCT/PCT Filing Date: 2016/11/11
(87) Date publication PCT/PCT Publication Date: 2017/05/18
(45) Date de délivrance/Issue Date: 2019/10/22
(85) Entrée phase nationale/National Entry: 2018/04/24
(86) N° demande PCT/PCT Application No.: KR 2016/013029
(87) N° publication PCT/PCT Publication No.: 2017/082688
(30) Priorité/Priority: 2015/11/13 (KR10-2015-0159637)

(51) Cl.Int./Int.Cl. *C07D 417/12* (2006.01),
A61K 31/427 (2006.01), *A61K 31/497* (2006.01),
C07D 417/14 (2006.01)

(72) Inventeurs/Inventors:
LEE, HYUNG-GEUN, KR;
KIM, IL-HWAN, KR;
JUNG, MYUNGGI, KR;
KIM, HYO SHIN, KR;
LEE, CHUN HO, KR;
JUN, SUN AH, KR;
YOON, JI SUNG, KR;
KIM, SUNG-YOUNG, KR

(73) Propriétaire/Owner:

(54) Titre : BLOQUEUR DES CANAUX SODIQUES
(54) Title: SODIUM CHANNEL BLOCKER



(57) Abrégé/Abstract:

The present invention relates to a compound represented by Chemical Formula, or a pharmaceutically acceptable salt thereof. The compound according to the present invention can be usefully used for the prevention or treatment of sodium channel blocker-

(73) Propriétaires(suite)/Owners(continued):DAEWOONG PHARMACEUTICAL CO., LTD., KR


(74) Agent: RICHES, MCKENZIE & HERBERT LLP

(57) Abrégé(suite)/Abstract(continued):
related diseases.

(see above formula)

[ABSTRACT]

The present invention relates to a compound represented by Chemical Formula, or a pharmaceutically acceptable salt thereof. The compound according to the present invention can be usefully used for the prevention or treatment of sodium channel blocker-related diseases.



[DESCRIPTION]**[TITLE OF THE INVENTION]****SODIUM CHANNEL BLOCKER****[Technical Field]**

5 The present invention relates to a compound having sodium ion channel blocking activity, a process for preparing the same and use thereof.

[Background Art]

 Voltage-gated sodium (Nav) channels are found in all excitable cells, including muscle and nerve cells of the central and peripheral nervous systems.

10 The sodium channels are essential for the initiation and propagation of electrical signals in the nervous system. Therefore, proper function of the sodium channels is essential to the normal function of the nerves. Ultimately, abnormal Nav channels play an important role in various diseases such as epilepsy, arrhythmia, myotonia, ataxia, multiple sclerosis, irritable bowel syndrome,

15 urinary incontinence, visceral pain, depression, and pains. Currently, ten types of Nav channels are found in Human (Nav 1.1-1.9, Nax). Among them, four channels of Nav1.3, Nav1.7, Nav1.8 and Nav 1.9 are known to have a close connection with pain signaling, and thus are recognized as an important analgesic drug target.

20

 A total of 10 types of Nav channels are found so far as summarized in Table 1 below. Among the 10 types, 9 types of Nav 1.1 to Nav1.9 form channels, among which NG 1.3, Nav 1.6, Nav 1.7, Nav 1.8 and Nav 1.9 are expressed in DRG.

25 [Table 1]

Type	Gene	Distribution tissue	TTX IC-50 nM	Indications
Nav1.1	SCN1A	CNS / PNS	10	Pain, epilepsy, neurodegeneration
Nav1.2	SCN2A	CNS	10	neurodegeneration, epilepsy
Nav1.3	SCN3A	CNS	15	Pain, epilepsy
Nav1.4	SCN4A	Sk. muscle	25	myotonia
Nav1.5	SCN5A	Heart	2000	arrhythmia
Nav1.6	SCN6A	CNS / PNS	6	pain, motor disorder
Nav1.7	SCN7A	PNS	25	pain, neuroendocrine disorder
Nav1.8	SCN8A	PNS	50000	pain

Nav1.9	SCN9A	PNS	1000	pain
--------	-------	-----	------	------

In particular, Nav1.7 is known to be preferentially expressed in dorsal root ganglia (DRG) and sympathetic ganglia. In the sensory ganglia DRG, Nav1.7 channel is expressed in A- or C-fiber neurons, but frequently distributed
5 in small neurons having a deep connection with pain. In particular, 85% of DRG are present in cells defined as nociceptors. This fact indicates that Nav1.7 has a close connection with pain.

The fact that Nav1.7 channel has a close connection with pain is well
10 demonstrated in the results of not only animal studies but also human disease studies. The results of animal studies indicated that, when inflammation occurs, the gene transcript of Nav1.7 significantly increases and the expression of proteins also increases. This increase in transcript is believed to be attributable to an increase in NGF. The increased expression of Nav1.7 is believed to be
15 the direct cause of an increase in excitability of sensory cells. In particular, when the gene of the Nav1.7 channel is removed or reduced, inflammatory pain is greatly reduced. However, animal studies do not indicate that the removal or reduction of the Nav1.7 channel gene reduces neuropathic pain. However, there are many evidences that Nav1.7 is involved in neuropathic pain in
20 humans.

Survey results for lineages that feel severe pain or no pain give many answers to pain studies. Particularly, these results directly indicate the importance of Nav1.7 in causing pain. There are two types of inherited diseases
25 that cause severe pain. In the case of erythromelalgia or erythermalgia among these diseases, severe pain is sometimes felt for a few hours when the body is slightly warm or takes exercises. In some cases, the skin becomes red, and the hand, the foot or the face swell. The results of genetic research indicated that SCN9A (the human gene name of Nav1.7) is present at chromosomal sites
30 associated with diseases. Nine mutations of Nav1.7 were found until now. These mutations lower activation threshold or result in slow deactivation of the channel. Thus, these mutations can easily generate action potential even upon

depolarization of some neurons (see Dib-Hajj, S D. et al., Trends in Neurosci., 30, 555-563:(2007)).

In the case of paroxysmal extreme pain disorder (PEPD) that is
 5 another inherited disease, pain is felt through life and caused when the bowels
 are evacuated or the anal region is stimulated. In addition to pain, the leg
 becomes red. As is known in the art, in PEPD, eight mutations occur in Nav1.7.
 These mutations occur mainly in sites that cause inactivation. The Nav channel
 has an inactivation ball in the linker between domains III and IV, and a peptide
 10 receiving region in the linker between the S5 and S6 segments of domains III
 and IV. Interestingly, mutations that cause PEPD all occur in these two regions.
 It appears that these cause a problem in the inactivation of Nav1.7. As expected,
 these mutations cause a problem in the inactivation of Nav1.7, resulting in slow
 deactivation of the channel (see Fertleman, C. R. et al., Neuron, 52, 767-774
 15 (2006)). Thus, the amount of electric current that enters through the channel
 increases.

Still another inherited disease is congenital indifference to pain (CIP).
 This disease results from mutation of the Nav1.7 channel and exist in Pakistani
 20 and Chinese lineages. Persons suffering from this disease feel no pain (see
 Cox, J. J. et al., Nature, 444, 894-898 (2006)). CIP causes the loss of function
 of the Nav1.7 channel. Particularly, a mutation in this channel inhibits the
 expression of this channel. Thus, this channel is not expressed (see Cox, J. J.
 et al., Nature, 444, 894-898 (2006)). Interestingly, the knock-out of Nav1.7 does
 25 not influence other sensations. However, it influences the olfactory sensation.
 This fact directly indicates that Nav1.7 does not overlap with other channels in
 pain transmission and the function thereof is not compensated for by other Nav
 channels.

30 As described above for the above diseases, when a mutation in the
 Nav1.7 channel causes a gain of function, severe pain is felt, and when it
 causes a loss of function, pain is relieved. This is a good clinical example that

directly shows that the Nav1.7 channel is the major cause of pain. Thus, it is considered that an antagonist that inhibits this channel will naturally result in a pain-relieving effect.

5 However, if the Nav1.7 channel antagonist inhibits a plurality of Nav channels including the Nav1.7 channel, it can show adverse effects of various CNS disturbances, such as blurring of vision, dizziness, vomiting and depression. Particularly, if it inhibits the Nav1.5 channel, it can cause cardiac arrhythmia and heart failure, which threaten life. For these reasons, selective
10 inhibition of the Nav1.7 channels is very important.

Pains can be largely classified into three: acute pain, inflammatory pain, and neuropathic pain. Acute pain plays an important protective function of maintaining the safety of organisms from stimuli that can cause tissue injury.
15 Thus, it is generally temporary and intense. On the other hand, inflammatory pain can be longer lasting, and the intensity thereof further increases. Inflammatory pain is mediated by various substances that are released during inflammation, including substance P, histamine, acids, prostaglandin, bradykinin, CGRP, cytokines, ATP and other substances. The third pain is neuropathic and
20 involves nerve injury or a nerve injury caused by viral infection. It causes reconstitution of circuits with neuron proteins to cause pathological "sensitization", which can result in chronic pain that is lasting for several years. This type of pain does not provide an advantage of adaptability and is difficult to treat by current therapy.

25 Particularly, neuropathic pain and intractable pain are great medical problems that have not been solved. Several hundred million patients are suffering from severe pain that is not well inhibited by current therapeutic methods. Drugs that are currently used for the treatment of pain include
30 NSAIDS, COX-2 inhibitors, opioids, tricyclic antidepressants and anticonvulsants. Neuropathic pain is particularly difficult to treat, because it does not well respond to opioids until a high dose is reached. Currently,

gabapentin is most widely used as a therapeutic agent against neuropathic pain, but it is effective for 60% of the patients and is not greatly effective. This drug is generally safe, but is problematic in terms of sedative action at high doses.

5 Accordingly, studies on the discovery of new regulators of the Nav1.7 channel and the use thereof for the treatment of acute pain, chronic acute, inflammatory pain and neuropathic pain have been actively conducted by global pharmaceutical companies, including Merck, AstraZeneca and the like (see US2010-0197655; US2012-0010183; WO2013-086229; WO2013-177224; 10 US2012-0238579; WO2007-145922).

 In view of the above, as a result of studying novel compounds, the present inventors has found that a compound having a chemical structure different from sodium channel blockers reported so far not only has excellent 15 sodium channel blocking effects, thereby completing the present invention. The compounds belonging to the present invention mainly have sodium channel inhibitory activity on their own, but do not exclude a possibility of exhibiting a pharmacological action as an efficacious agent by a special body environment or by products of metabolic process, after absorption into the body.

20 **[DETAILED DESCRIPTION OF THE INVENTION]**


[Technical Problem]

 It is one object of the present invention to provide a compound having a blocking effect against sodium ion channels, particularly Nav 1.7, a process for its preparation and its use.

25 **[Technical Solution]**

 In order to achieve the above objects, the present invention provides a compound represented by Chemical Formula 1 below, or a pharmaceutically acceptable salt thereof:

[Chemical Formula 1]



in Chemical Formula 1,

R_1 is hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, halogen, or cyano,

5 R_2 and R_3 are each independently hydrogen, or halogen,

R_4 is C_{5-10} heteroaryl containing one or two elements each independently selected from the group consisting of N, S and O,

wherein the C_{5-10} heteroaryl is unsubstituted or substituted with C_{1-4} alkyl or halogen,

10 R_5 is $-CH_2CH_2-N(R_7)(R_8)$, or $-CH_2CH_2CH_2-N(R_7)(R_8)$, R_6 is hydrogen, or C_{1-4} alkyl; or R_5 and R_6 together form C_{3-5} alkylene, $(C_{2-4}$ alkylene)- $N(R_9)$ -(C_{2-4} alkylene), or $(C_{2-4}$ alkylene)-O-(C_{2-4} alkylene),

wherein the C_{3-5} alkylene, or C_{2-4} alkylene is each independently unsubstituted or substituted with one or two R_{10} ,

15 R_7 , R_8 , and R_9 are each independently hydrogen, or C_{1-4} alkyl,

R_{10} is C_{1-4} alkyl, C_{1-4} alkoxy, halogen, amino, $NH(C_{1-4}$ alkyl), $N(C_{1-4}$ alkyl)₂, $NHCO(C_{1-4}$ alkyl), or pyrrolidinyl,


X_1 is C- R' , or N, wherein R' is hydrogen, or halogen,

X_2 is CH, or N, and

20 X_3 is N- R'' , wherein R'' is hydrogen, or C_{1-4} alkyl.

Preferably, the compound represented by Chemical Formula 1 is represented by Chemical Formula 1' below:

[Chemical Formula 1']



wherein,

R_1 is hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, C_{1-4} haloalkyl, C_{1-4} haloalkoxy, halogen, or cyano,

5 R_2 and R_3 are each independently hydrogen, or halogen,

R_4 is C_{5-10} heteroaryl containing one or two elements each independently selected from the group consisting of N, S and O,

R_{11} is hydrogen, or C_{1-4} alkyl,

R_{12} is hydrogen, C_{1-4} alkyl, C_{1-4} alkoxy, halogen, amino, $NH(C_{1-4}$ alkyl),

10 $N(C_{1-4}$ alkyl) $_2$, $NHCO(C_{1-4}$ alkyl), or pyrrolidinyl,

X_1 is C-R', or N, wherein R' is hydrogen, or halogen,

X_2 is CH, or N, and

X_3 is N-R'', wherein R'' is hydrogen, or C_{1-4} alkyl,

X_4 is a bond, NH, $N(C_{1-4}$ alkyl), or o (oxygen), and

15 n is an integer of 1 to 4.


Preferably, R_1 is hydrogen, methyl, methoxy, trifluoromethyl, difluoromethoxy, trifluoromethoxy, fluoro, chloro, or cyano.

20 Preferably, R_2 and R_3 are each independently hydrogen, fluoro, or chloro.

Preferably, R_4 is thiazolyl, isoxazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridinyl, pyrimidinyl, or thiadiazolyl, wherein the R_4 is unsubstituted or
25 substituted with C_{1-4} alkyl, or halogen.

Preferably, R_5 and R_6 together form C_{3-5} alkylene, $(C_{2-4}$ alkylene)- $N(R_9)-(C_{2-4}$ alkylene), or $(C_{2-4}$ alkylene)- $O-(C_{2-4}$ alkylene), wherein the C_{3-5} alkylene or C_{2-4} alkylene is each independently unsubstituted or substituted with methyl, methoxy, fluoro, amino, $NHCH_3$, $N(CH_3)_2$, $N(CH_2CH_3)_2$, $NHCOCH_3$, or pyrrolidinyl.

Preferably, R_5 , R_6 , and R_5 and R_6 together with the nitrogen to which they are attached form any one selected from the group consisting of:



wherein, R_{10} is as defined above.

Preferably, R_5 is $-CH_2CH_2-NH-CH_3$, $-CH_2CH_2CH_2-NH-CH_3$, or $-CH_2CH_2-N(CH_3)_2$, and R_6 is hydrogen, or methyl.

Preferably, X_1 is CH, CF, or N, X_2 is CH, or N, provided that both X_1 and X_2 are not N.

Preferably, X_3 is NH.

Representative examples of the compound represented by Chemical Formula 1 or a pharmaceutically acceptable salt thereof are as follows:

1) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

2) 5-chloro-4-((4-chloro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

- 3) (R)-5-chloro-4-((4-chloro-2-(2-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 4) (R)-5-chloro-4-((4-chloro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 5) (R)-N-(1-(5-chloro-2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)phenyl)pyrrolidin-3-yl)acetamide,
- 6) 5-chloro-4-((4-chloro-2-(3-(diethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 7) 4-((2-([1,3'-bipyrrolidin]-1'-yl)-4-chlorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 8) 5-chloro-2-fluoro-4-((4-methyl-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 9) (S)-5-chloro-4-((4-chloro-2-(3-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 10) 4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 11) 3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 12) 3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 13) 2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 14) 4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15) 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 16) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 17) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 18) 5-chloro-4-((5-chloro-3-(3-(methylamino)pyrrolidin-1-yl)pyridin-2-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

- 19) 5-chloro-4-((6-chloro-2-(3-(methylamino)pyrrolidin-1-yl)pyridin-3-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 21) 5-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 22) (R)-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 23) (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 24) (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25) (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15 26) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 27) (R)-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 28) (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 20 29) (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30) (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 31) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 32) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 33) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 30 34) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-3-chloro-N-(thiazol-4-yl)benzenesulfonamide,

35) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

36) 4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

5 37) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

38) 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

10 39) 4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

40) 3-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

41) (R)-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

15 42) (R)-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

43) (R)-3-chloro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

20 44) (R)-3,5-difluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

45) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

46) 2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

25 47) 3-chloro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

48) 5-chloro-2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

30 49) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

50) 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

- 51) 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 52) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 53) 2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 54) 3-chloro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 55) 5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 56) (R)-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 57) (R)-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15 58) (R)-3-chloro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 59) (R)-3,5-difluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 60) (R)-5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 20 61) 3-chloro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 62) 5-chloro-2-fluoro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 63) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 64) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 65) 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30 66) 5-chloro-4-((4-(difluoromethoxy)-2-(4-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

- 67) 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 68) (S)-3-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 69) (S)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 70) (S)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 71) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-methoxyphenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 10 72) (S)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 73) (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 15 74) (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 75) (R)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20 76) 4-((2-(1,4-diazepan-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 77) 5-chloro-4-((4-cyano-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 25 78) (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide,
- 79) (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide,
- 80) (S)-3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30 81) (S)-5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

- 82) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)azetidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 83) 4-((2-(3-aminoazetidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 5 84) 5-chloro-4-((2-(3-(dimethylamino)azetidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 85) N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)azetidin-3-yl)acetamide,
- 86) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 87) 5-chloro-2-fluoro-4-((2-(3-methoxypyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 88) (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide,
- 15 89) 3-chloro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 90) 3-chloro-4-((2-(3-methoxypyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 91) (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide,
- 20 92) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 93) 3-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 94) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 95) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 96) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,
- 30 97) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-

(methylamino)ethyl)amino)-4-

(trifluoromethyl)phenyl)amino)benzenesulfonamide,

98) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

99) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

100) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,

101) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,

102) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

103) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyridin-2-yl)benzenesulfonamide,

104) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,

105) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,

106) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,

107) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide,

108) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,

- 109) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)benzenesulfonamide,
- 110) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,
- 111) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,
- 112) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 113) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,
- 114) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,
- 115) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,
- 116) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,
- 117) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide,
- 118) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 119) 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,
- 120) 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 121) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 122) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

- 123) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 124) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,
- 125) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,
- 126) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,
- 127) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 128) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluorothiazol-2-yl)benzenesulfonamide,
- 129) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 130) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 131) 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 132) 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 133) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,
- 134) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,
- 135) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-

(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide,

136) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide,

137) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,

138) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,

139) 5-chloro-N-(5-chlorothiazol-2-yl)-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluorobenzenesulfonamide,

140) 5-chloro-N-(5-chlorothiazol-2-yl)-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)phenyl)amino)-2-fluorobenzenesulfonamide,

141) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylthiazol-2-yl)benzenesulfonamide,

142) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(oxazol-2-yl)benzenesulfonamide,

143) N-(5-(tert-butyl)isoxazol-3-yl)-5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

144) N-(5-(tert-butyl)isoxazol-3-yl)-5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

145) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1,2,4-thiadiazol-5-yl)benzenesulfonamide,

146) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1-methyl-1H-pyrazol-3-yl)benzenesulfonamide,

147) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-4-yl)benzenesulfonamide,

148) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,

149) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrazin-2-yl)benzenesulfonamide,

150) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,

5 151) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-5-yl)benzenesulfonamide,

152) 5-chloro-2-fluoro-4-((4-fluoro-2-((2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

153) 5-chloro-4-((2-((2-(dimethylamino)ethyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide, and

154) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(3-(methylamino)propyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide.

In addition, the compounds of the present invention may exist in the form of salts, especially pharmaceutically acceptable salts. As salts, salts commonly used in the art, such as acid addition salts formed by pharmaceutically acceptable free acids can be used without limitation. The term “pharmaceutically acceptable salt” as used herein refers to any organic or inorganic addition salt of the compound represented by Chemical Formula 1, whose concentration is relatively non-toxic and harmless to a patient and activates effectively and whose side effects do not degrade the beneficial efficacy of the above compound.

As the free acid, an organic acid and an inorganic acid can be used. Examples of the inorganic acids include hydrochloric acid, phosphoric acid, sulfuric acid, nitric acid, tartaric acid and the like. Examples of the organic acids include methanesulfonic acid, p-toluenesulfonic acid, acetic acid, trifluoroacetic acid, maleic acid, succinic acid, oxalic acid, benzoic acid, tartaric acid, fumaric acid, mandelic acid, propionic acid, citric acid, lactic acid, glycolic acid, gluconic acid, galacturonic acid, glutamic acid, glutaric acid, glucuronic acid, aspartic acid, ascorbic acid, carbonic acid, vanillic acid, hydroiodic acid and the like, but are not limited thereto.

In addition, a pharmaceutically acceptable metal salt can be obtained by a conventional method using a base. For example, a compound represented by Chemical Formula 1 is dissolved in an excessive amount of an alkali metal hydroxide or an alkaline earth metal hydroxide solution, the non-soluble salt is
5 filtered, and the filtrate is evaporated and dried to obtain a pharmaceutically acceptable metal salt. At this time, it is particularly preferable to prepare a sodium salt, a potassium salt or a calcium salt as the metal salt.


10 A pharmaceutically unacceptable salt or solvate of the compound of Chemical Formula 1 may be used as an intermediate when preparing the compound of Chemical Formula 1, or the pharmaceutically acceptable salt or the solvate thereof.

15 Further, the compound of Chemical Formula 1 according to the present invention includes not only pharmaceutically acceptable salts thereof, but also solvates such as hydrates that can be prepared therefrom, and includes all possible stereoisomers, but are not limited thereto. The solvate and the stereoisomer of the compound of Chemical Formula 1 may be prepared
20 from the compound of Chemical Formula 1 using common methods known in the art.

In addition, the compound of Chemical Formula 1 according to the present invention may be prepared either in a crystalline form or in a non-crystalline form, and when the compound of Chemical Formula 1 is prepared in
25 a crystalline form, it may be optionally hydrated or solvated. In the present invention, the compound of Chemical Formula 1 may not only include a stoichiometric hydrate, but also include a compound containing various amounts of water. The solvate of the compound of Chemical Formula 1
30 according to the present invention includes both stoichiometric solvates and non-stoichiometric solvates.

Furthermore, as an example, the present invention can produce the compound represented by Chemical Formula 1 through Reaction Scheme 1 below.

[Reaction Scheme 1]



5

in Reaction Scheme 1, X_1 to X_4 , R_1 to R_6 , and n are as previously defined in Chemical Formula 1.

According to the steps i, ii and iii, the respective compounds represented by Chemical Formulas 1-3 and 1-6 can be prepared as intermediates. The step i is preferably reacted at 50 to 60°C in the presence of potassium carbonate, and the solvent is preferably dimethylformamide. The step ii is preferably carried out at 50 to 60°C in the presence of sodium hyposulfite, and as the solvent, a mixed solvent of ethanol and water (1:1) is preferably used. The step iii is preferably carried out at -78°C in the presence of lithium hexamethyldisilylamide, and the solvent is preferably tetrahydrofuran.

The intermediate represented by Chemical Formula 1-7 can be prepared according to the step iv. The above step is a step of reacting the compound represented by Chemical Formula 1-3 with the compound represented by Chemical Formula 1-6. The reaction is preferably carried out at 120°C in the presence of palladium diacetate, bis(diphenylphosphino)binaphthyl and cesium carbonate, and the solvent is preferably dioxane.

The compound represented by Chemical Formula 1 can be obtained according to the step v. The reaction is preferably carried out at 60 to 80°C in the presence of hydrochloric acid, and the solvent is preferably methanol. Also, depending on the type of substituent of R₁ to R₆, if it is necessary to protect in
5 the reaction of the steps i to iv, the steps i to iv may be carried out in a state of being protected with a protective group, which can be then removed in the step v.

Further, the present invention provides a pharmaceutical composition
10 for preventing or treating diseases, which is effective for sodium channel blocking activity, comprising the compound represented by Chemical Formula 1, or a pharmaceutically acceptable salt, hydrate, solvate or isomer thereof as an active ingredient.

15 In this case, the disease includes acute pain, chronic pain, neuropathic pain, postoperative pain, migraine, arthralgia, neuropathy, nerve damage, diabetic neuropathy, neuropathic disease, epilepsy, arrhythmia, myotonia, ataxia, multiple sclerosis, irritable bowel syndrome, urinary incontinence, visceral pain, depression, erythralgia, PEPD (paroxysmal extreme
20 pain disorder) or the like.

As used herein, the term "prevention" refers to any act to delay or inhibit occurrence, spread or recurrence of the above-mentioned diseases by administration of the composition of the present invention, and "treatment"
25 refers to any act to improve or change the symptoms of the above diseases for the better by administration of the composition of the present invention.

The pharmaceutical composition according to the present invention can be formulated in types for oral or parenteral administrations according to a
30 standard pharmaceutical practice. These formulations may contain additives such as pharmaceutically acceptable carrier, adjuvant or diluent in addition to the active ingredient.

Suitable carriers include, for example, physiological saline, polyethylene glycol, ethanol, vegetable oil, and isopropyl myristate and the like. Diluents include, for example, lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine and the like, but are not limited thereto. Further, the compounds of the present invention can be dissolved in oils, propylene glycol or other solvents commonly used in the preparation of injection solutions. Furthermore, the compounds of the present invention can be formulated in ointments or creams for topical application.

10

A preferred dose of the compound of the present invention may be varied according to the condition and weight of a patient, the severity of a disease, the type of a drug, and the route and duration of administration, but it may be suitably selected by those skilled in the art. In order to achieve the desirable effects, however, the compound of the present invention may be administrated daily at a dose of 0.0001 to 100 mg/kg (body weight), and preferably 0.001 to 100 mg/kg (body weight). The administration may be performed once a day or in divided doses each day through an oral or parenteral route.

20

Depending on the method of administration, the Pharmaceutical composition may contain the compound of the present invention in an amount of 0.001 to 99% by weight, preferably 0.01 to 60% by weight.

25

The pharmaceutical composition according to the present invention may be administered to mammals such as a rat, a mouse, a domestic animal, a human, through various routes. The administration may be carried out through all possible methods, for example, oral, rectal, intravenous, intramuscular, subcutaneous, intra-endometrial, intracerebroventricular injection.

30

[ADVANTAGEOUS EFFECTS]


The compound represented by Chemical Formula 1 according to the present invention or a pharmaceutically acceptable salt, hydrate, solvate or

isomer thereof can be usefully used for the prevention or treatment of sodium channel blocker-related diseases.

[DETAILED DESCRIPTION OF THE EMBODIMENTS]

Below, the present invention will be described in more detail by way of examples. However, these examples are provided for illustrative purposes only, and should not be construed as limiting the scope of the present invention to these examples.

Example 1: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



Step 1) Preparation of tert-butyl (1-(2-amino-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate

2,4-difluoro-1-nitrobenzene (2.0 g, 12.6 mmol) and tert-butylmethyl(pyrrolidin-3-yl)carbamate (2.5 g, 1.0 eq.) were dissolved in DMF (20 mL), and then K_2CO_3 (2.6 g, 1.5 eq.) was added thereto. While maintaining the internal temperature at 60 to 70°C, the reaction mixture was stirred for 2 hours. When the reaction solution became dark yellow, the completion of the reaction was confirmed by TLC. After cooling to room temperature, ethyl acetate (EA)/H₂O was added and stirred, and then the layers were separated. $MgSO_4$ was added to the separated organic layer, which was stirred, dried and then filtered. The filtrate was concentrated under reduced pressure, and the residue was dissolved in EtOH (10 mL) and distilled water (10 mL), to which $Na_2S_2O_4$ (13.0 g, 6 eq.) was added. While maintaining the internal temperature at 60 to 70°C, the reaction mixture was stirred for 2 hours. When the yellow color of the reaction solution faded and became almost colorless, the completion of the reaction was confirmed by TLC. After cooling to room temperature, distilled

water (50 mL) was added and extracted twice with EA (100 mL). MgSO₄ was added to the organic layer, which was stirred, dried and then filtered. The filtrate was concentrated under reduced pressure, and the resulting residue was separated by column chromatography (n-Hexane/EA = 3/1) to obtain the title compound (2.0 g, 51.1%).

¹H NMR (MeOD): 6.73(m, 1H), 6.57(t, 1H), 3.23(m, 1H), 3.10(m, 2H), 2.94(m, 1H), 2.91(s, 3H), 2.25(m, 1H), 1.99(m, 1H)

Step 2) Preparation of tert-butyl thiazol-4-yl carbamate

Thiazol-4-carboxylic acid (5.0 g, 38.8 mmol) was dissolved in t-BuOH (100 mL), and then TEA (8.1 mL, 1.5 eq.) and DPPA (7.1 mL, 1.5 eq.) were added thereto. While maintaining the internal temperature at 90 to 100°C, the reaction mixture was stirred for 3 days, and then the completion of the reaction was confirmed by TLC. The product was concentrated under reduced pressure, distilled water (50 mL) was added, and extracted twice with EA (100 mL). MgSO₄ was added to the organic layer, which was stirred, dried and then filtered. The filtrate was concentrated under reduced pressure, and the residue was added to a small amount of EA and slurried. The resulting solid was filtered to obtain a white title compound (4.0 g, 51.5%).

¹H NMR (MeOD): 8.73(s, 1H), 7.24(s, 1H), 1.52(s, 9H)

Step 3) Preparation of tert-butyl ((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(thiazol-4-yl)carbamate

Tert-butyl thiazol-4-yl carbamate (4.0 g, 20.0 mmol) prepared in the step 2) was added to a reaction vessel and the inside of the vessel was replaced with nitrogen gas. After dissolving in THF (32 mL), the solution was cooled to -78°C using dry ice-acetone. After cooling, LiHMDS (22.4 mL, 1.5 eq.) was slowly added and the reaction mixture was stirred for 30 minutes. 4-Bromo-5-chloro-2-fluorobenzenesulfonyl chloride (6.0 g, 1.0 eq.) was dissolved in THF (10 mL) and then slowly added to the reaction solution. The reaction mixture was stirred overnight and the completion of the reaction was confirmed by TLC. Distilled water (50 mL) was added and extracted twice with EA (100 mL).

MgSO₄ was added to the organic layer, which was stirred, dried and then filtered. The filtrate was concentrated under reduced pressure, and the residue was crystallized with THF/n-Hexane to obtain the title compound (4.4 g, 59.0%).

1H NMR (MeOD): 9.00(s, 1H), 8.22(d, 1H), 7.90(d, 1H), 7.78(s, 1H),
5 1.35(s, 9H)

Step 4) Preparation of tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate

10 Tert-butyl (1-(2-amino-5-fluorophenyl)pyrrolidin-3-
yl)(methyl)carbamate (0.5 g, 1.1 mmol) prepared in the step 1) and tert-butyl
((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(thiazol-4-yl)carbamate (0.9 g, 1.2
eq.) prepared in the step 3) were dissolved in 1,4-dioxane (10 mL).
Pd(OAc)₂(0.03 g, 0.1 eq), rac-BINAP(0.19 g, 0.2 eq.) and Cs₂CO₃(1.5 g, 3.0 eq.)
15 were added to the reaction solution. After reacting at 120°C for 30 minutes
using a microwave initiator, the completion of the reaction was confirmed by
TLC. Distilled water (50 mL) was added and extracted twice with EA (100 mL).
MgSO₄ was added to the organic layer, which was stirred, filtered and then
dried. The filtrate was concentrated under reduced pressure, and the residue
20 was separated by column chromatography (EA/n-Hexane= 1/1). This procedure
was repeated twice to obtain the title compound (2.0 g, 88.2%).

¹H NMR (MeOD): 8.95(s, 1H), 7.94(d, 1H), 7.65(s, 1H), 7.14(t, 1H), 6.70(d, 1H), 6.64(t, 1H), 6.07(d, 1H), 3.40(m, 1H), 3.28(m, 2H), 3.16(m, 1H), 2.64(s, 3H), 2.06(m, 1H), 1.89(m, 1H), 1.41(s, 9H), 1.36(s, 9H)

Step 5) Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride


To tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (2.0 g, 2.9 mmol) prepared in the step 4) was added 1.25 M HCl in MeOH(15 mL). After stirring the mixture overnight while heating to 40

to 50°C, the completion of the reaction was confirmed by TLC. The product was concentrated, methylene chloride (15 mL) was added to the obtained residue, stirred for 1 hour, and the produced solid was filtered to give the title compound (0.9 g, 58.8%).

5 ¹H NMR (MeOD): 8.73(s, 1H), 7.75(d, 1H), 7.12(t, 1H), 7.00(s, 1H), 6.69(d, 1H), 6.67(t, 1H), 6.05(d, 1H), 3.73(m, 1H), 3.54(m, 1H), 3.45(m, 1H), 3.38(m, 1H), 3.26(m, 1H), 2.63(s, 3H), 2.31(m, 1H), 1.96(m, 1H)

Hereinafter, the compounds of Examples 2 to 9 were prepared in the
10 same manner as described in Example 1, except that the reactants corresponding to the structures of the compounds to be produced were used.


Example 2: Preparation of 5-chloro-4-((4-chloro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride
15



¹H NMR (500 MHz, MeOD): 8.79(s, 1H), 7.74(d, 1H), 7.09(d, 1H), 7.02(s, 1H), 6.98(s, 1H), 6.92(d, 1H), 6.13(d, 1H), 3.76(m, 1H), 3.53(m, 3H), 2.64(s, 3H), 2.31(m, 1H), 2.01(m, 1H), 1.27(m, 1H)

20


Example 3: Preparation of (R)-5-chloro-4-((4-chloro-2-(2-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



¹H NMR (500 MHz, MeOD): 8.73(d, 1H), 7.84(d, 1H), 7.38(d, 1H), 7.30(d, 1H), 7.28(d, 1H), 7.04(d, 1H), 6.99(d, 1H), 3.40(m, 3H), 3.15(m, 2H), 3.07(m, 1H), 2.86(m, 1H), 0.93(m, 3H)

5


Example 4: Preparation of (R)-5-chloro-4-((4-chloro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



¹H NMR (500 MHz, MeOD): 8.75(d, 1H), 7.80(d, 1H), 7.28(d, 1H), 7.20(d, 1H), 7.17(m, 1H), 7.03(d, 1H), 6.58(d, 1H), 3.44(m, 1H), 3.02(m, 1H), 2.94(m, 1H), 2.78(m, 1H), 2.67(s, 3H), 2.08(m, 1H), 1.79(m, 1H), 1.50(m, 1H), 1.41(m, 1H), 0.87(m, 1H)


15

Example 5: Preparation of (R)-N-(1-(5-chloro-2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)phenyl)pyrrolidin-3-yl)acetamide hydrochloride




¹H NMR (500 MHz, MeOD): 8.78(m, 1H), 7.76(d, 1H), 7.17(d, 1H), 7.08(m, 1H), 7.02(s, 1H), 6.92(m, 1H), 6.15(t, 1H), 4.26(m, 1H), 3.55(m, 1H), 3.35(m, 1H), 3.25(m, 1H), 2.27(m, 1H), 1.93(m, 1H), 1.90(s, 3H)

5 **Example 6: Preparation of 5-chloro-4-((4-chloro-2-(3-(diethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride**




10 ¹H NMR (500 MHz, MeOD): 8.78(d, 1H), 7.77(d, 1H), 7.12(d, 1H), 7.03(m, 1H), 6.99(dd, 1H), 6.03(d, 1H), 4.08(m, 1H), 3.52(m, 3H), 3.17(m, 4H), 3.05(m, 1H), 2.39(m, 1H), 2.01(m, 1H), 1.26(m, 6H)

15 **Example 7: Preparation of 4-((2-([1,3'-bipyrrolidin]-1'-yl)-4-chlorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride**




¹H NMR (500 MHz, MeOD): 8.76(d, 1H), 7.77(d, 1H), 7.11(d, 1H), 7.02(m, 2H), 6.97(m, 1H), 6.07(d, 1H), 3.91(m, 1H), 3.50(m, 5H), 3.22(m, 1H), 3.05(m, 2H), 2.36(m, 1H), 1.97(m, 5H)

20 **Example 8: Preparation of 5-chloro-2-fluoro-4-((4-methyl-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-**

yl)benzenesulfonamide hydrochloride

1H NMR (500 MHz, MeOD): 8.76(d, 1H), 7.74(d, 1H), 7.02(m, 2H),
 6.91(s, 1H), 6.85(d, 1H), 6.16(d, 1H), 3.78(m, 1H), 3.45(m, 3H), 3.23(m, 1H),
 2.65(s, 3H), 2.35(s, 3H), 2.34(m, 1H), 2.01(m, 1H)

Example 9: Preparation of (S)-5-chloro-4-((4-chloro-2-(3-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride




10

1H NMR (500 MHz, MeOD): 8.73(d, 1H), 7.79(d, 1H), 7.29(d, 1H),
 7.20(m, 2H), 7.03(d, 1H), 6.54(d, 1H), 3.88(m, 1H), 3.44(s, 3H), 3.04(m, 1H),
 2.97(m, 1H), 2.80(m, 1H), 1.27(m, 3H)

15

Example 10: Preparation of 4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



Intermediate tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(thiazole-

4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate was prepared in the same manner as described in the steps 1 to 4 of Example 1, except that 4-bromobenzenesulfonyl chloride was used instead of 4-bromo-5-chloro-2-fluorobenzenesulfonyl chloride in the step 3 of Example 1.


5

To the obtained intermediate was added 1.25 M HCl in MeOH (15 mL). After stirring the mixture overnight while heating to 40 to 50°C, the completion of the reaction was confirmed by TLC. The product was concentrated, and the obtained residue was separated and purified by PLC to
10 obtain 0.05 g of the target compound (yield: 48%).

¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.57(d, 2H), 7.07(t, 1H), 6.92(s, 1H), 6.66-6.54(m, 4H), 3.42-3.30(m, 4H), 3.14-3.13(m, 1H), 2.42(s, 3H), 2.18-2.16(m, 1H), 1.84-1.40(m, 1H)


15 Hereinafter, the compounds of Examples 11 to 91 were prepared in the same manner as described in Example 10, except that the reactants corresponding to the structures of the compounds to be produced were used.

20 **Example 11: Preparation of 3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide**




¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.74(s, 1H), 7.47(d, 1H), 7.05(t, 1H), 6.93(s, 1H), 6.63(d, 1H), 6.54(t, 1H), 6.33(d, 1H), 3.40-3.37(m, 2H),
25 3.17-3.11(m, 3H), 2.31(s, 3H), 2.08-2.00(m, 1H), 1.72-1.68(m, 1H)

Example 12: Preparation of 3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-

yl)benzenesulfonamide

1H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.43(d, 2H), 7.04(s, 1H),
 6.63(t, 1H), 6.75(d, 1H), 6.60(t, 1H), 3.66-3.63(m, 1H), 3.49-3.35(m, 3H), 3.08-
 5 3.04(m, 1H), 2.63(s, 3H), 2.31-2.28(m, 1H), 1.96-1.93(m, 1H)

Example 13: Preparation of 2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




10

1H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.56(t, 1H), 7.03(t, 1H),
 6.84(s, 1H), 6.63(d, 1H), 6.55(t, 1H), 6.40(d, 1H), 6.23(d, 1H), 3.35-3.30(m, 3H),
 3.18-3.16(m, 2H), 2.36(s, 3H), 2.11-2.05(m, 1H), 1.78-1.75(m, 1H)

15


Example 14: Preparation of 4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.60(d, 2H), 7.16(d, 1H),
 6.92(s, 1H), 6.77-6.68(m, 4H), 3.38-3.61(m, 1H), 3.35-3.33(m, 2H), 3.28-3.27(m

1H), 3.09-3.06(m, 1H), 2.36(s, 3H), 2.36-2.35(m, 1H), 1.85-1.83(m, 1H)


Example 15: Preparation of 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.62(s, 1H), 7.41(d, 2H), 6.87(s, 1H), 6.76-6.75(m, 1H), 6.69-6.67(m, 2H), 3.59(s, 1H), 3.49-3.47(m, 2H), 3.25-3.23(m, 1H), 2.99-2.98(m, 1H), 2.59(s, 3H), 2.29-2.27(m, 1H), 1.99-1.97(m, 1H)

10

Example 16: Preparation of 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.58(t, 1H), 7.16(d, 1H), 6.84(s, 1H), 6.84-6.83(m, 2H), 6.49(d, 1H), 6.35(d, 1H), 3.38-3.34(m, 3H), 3.23-3.21(m, 1H), 3.12-3.10(m, 1H), 2.41(s, 3H), 2.18-2.16(m, 1H), 1.85-1.83(m, 1H)

15

Example 17: Preparation of 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


20



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.74(d, 1H), 7.15(d, 1H), 6.85(s, 1H), 6.75-6.74(m, 2H), 6.04(d, 1H), 3.45-3.38(m, 3H), 3.19-3.17(m, 2H), 2.40(s, 3H), 2.13-2.12(m, 1H), 1.80-1.79(m, 1H)


5

Example 18: Preparation of 5-chloro-4-((5-chloro-3-(3-(methylamino)pyrrolidin-1-yl)pyridin-2-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(d, 1H), 8.30(d, 1H), 7.96(d, 1H), 7.84(d, 1H), 7.53(d, 1H), 6.87(d, 1H), 3.44(m, 3H), 3.10(m, 2H), 2.50(s, 3H), 2.36(m, 1H), 1.95(m, 1H)


Example 19: Preparation of 5-chloro-4-((6-chloro-2-(3-(methylamino)pyrrolidin-1-yl)pyridin-3-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.76(s, 1H), 7.34(s, 1H), 6.90(m, 2H), 6.70(s, 1H), 6.01(d, 1H), 3.45(m, 3H), 2.99(m, 2H), 2.49(s, 3H),

2.11(m, 2H)


Example 20: Preparation of 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.67(d, 1H), 7.78(d, 1H), 7.26(d, 1H), 7.16(m, 2H), 6.81(d, 1H), 6.21(d, 1H), 3.39(m, 3H), 3.17(m, 2H), 2.43(s, 3H), 2.13(m, 1H), 1.94(m, 1H)

10

Example 21: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.63(d, 1H), 7.73(d, 1H), 7.07(d, 1H), 6.70(t, 2H), 6.62(t, 2H), 6.03(d, 1H), 3.40(m, 3H), 3.23(m, 2H), 2.42(s, 3H), 2.11(m, 1H), 1.83(m, 1H)

15

Example 22: Preparation of (R)-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


20



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.61(d, 2H), 7.17(t, 1H), 6.89(s, 1H), 6.77-6.75(m, 3H), 6.74-6.73(m, 1H), 3.34-3.30(m, 1H), 2.88-2.59(m, 4H), 2.32(s, 3H), 1.82-1.70(m, 2H), 1.45-1.44(m, 2H)

5

Example 23: Preparation of (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




10

¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.61(t, 1H), 7.20(t, 1H), 6.87(d, 1H), 6.85-6.81(m, 2H), 6.78(d, 1H), 6.67(d, 1H), 3.34(s, 1H), 2.92-2.79(m, 4H), 2.48(s, 3H), 1.89-1.87(m, 2H), 1.79-1.78(m, 1H), 1.47-1.46(m, 1H)

Example 24: Preparation of (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


15



¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.80(s, 1H), 7.54(s, 1H), 7.24-7.21(m, 1H), 6.89-6.77(m, 4H), 3.01-2.97(m, 1H), 2.68-2.41(m, 3H), 2.30(s,

3H), 1.86-1.84(m, 1H), 1.68-1.66(m, 1H), 1.40-1.35(m, 2H)


Example 25: Preparation of (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.64(s, 1H), 7.41(d, 2H), 6.85(d, 1H), 6.75-6.66(m, 3H), 3.26-3.24(m, 1H), 2.88-2.78(m, 4H), 2.55(s, 3H), 1.87-1.86(m, 1H) 1.85-1.83(m, 1H), 1.74-1.72(m, 1H), 1.37-1.35(m, 1H)

10


Example 26: Preparation of (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.66(s, 1H), 7.77(d, 1H), 7.22(t, 1H), 6.89-6.84(m, 2H), 6.81(s, 1H), 6.35(d, 1H), 3.38-3.36(m, 1H), 3.03-3.01(m, 1H), 2.63-2.61(m, 1H), 2.48-2.44(m, 2H), 2.37(s, 3H), 1.89-1.87(m, 1H), 1.68-1.66(m, 1H), 1.32-1.32(m, 2H)

20


Example 27: Preparation of (R)-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.56(d, 2H), 7.04(t, 1H), 6.92(s, 1H), 6.63-6.53(m, 4H), 3.24-3.21(m, 3H), 3.14-3.10(m, 2H), 2.31(s, 3H), 2.11-2.08(m, 1H), 1.73-1.70(m, 1H)

5


Example 28: Preparation of (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.57-7.53(m, 1H), 7.05-7.03(m, 1H), 6.83(s, 1H), 6.62(d, 1H), 6.55-6.53(m, 1H), 6.40(d, 1H), 6.23(d, 1H), 3.34-3.30(m, 2H), 3.28-3.27(m, 1H), 3.18-3.15(m, 2H), 2.36(s, 3H), 2.13-2.10(m, 1H), 1.76-1.74(m, 1H)

15


Example 29: Preparation of (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.75(s, 1H), 7.47-7.45(m,


1H), 7.06-7.04(m, 1H), 6.92(s, 1H), 6.62(d, 1H), 6.54(d, 1H), 6.32(d, 1H), 3.40-3.37(m, 2H), 3.17-3.11(m, 3H), 2.33(s, 3H), 2.08-2.05(m, 1H), 1.72-1.69(m, 1H)

Example 30: Preparation of (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




1H NMR (500 MHz, MeOD): 8.65(s, 1H), 7.37(d, 2H), 6.77-6.74(m, 2H), 6.69(d, 1H), 6.56-6.52(m, 1H), 3.41-3.26(m, 3H), 3.26-3.24(m, 1H), 3.05-3.02(m, 1H), 2.48(s, 3H), 2.20-2.18(m, 1H), 1.83-1.81(m, 1H)

Example 31: Preparation of (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.67(s, 1H), 7.72(d, 1H), 7.04(t, 1H), 6.81(d, 1H), 6.62(t, 1H), 5.98(d, 1H), 5.48(s, 1H), 3.43-3.36(m, 2H), 3.39-3.36(m, 1H), 3.30-3.26(m, 2H), 2.36(s, 3H), 2.36-2.10(m, 1H), 1.77-1.73(m, 1H)


Example 32: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.73(d, 1H), 7.06(t, 1H), 6.83(s, 1H), 6.64(d, 1H), 6.57(t, 1H), 6.01(d, 1H), 3.50-3.40(m, 3H), 3.28-3.23(m, 2H), 2.48(s, 3H), 2.20-2.19(m, 1H), 1.95-1.93(m, 1H)

5


Example 33: Preparation of 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.72(s, 1H), 7.58(t, 1H), 7.10(t, 1H), 6.93(m, 1H), 6.68(d, 1H), 6.62(d, 1H), 6.45(d, 1H), 6.33(d, 1H), 3.79(m, 1H), 3.50(m, 2H), 3.30(m, 1H), 3.20(m, 1H), 2.29(m, 1H), 1.95(m, 1H)

10


Example 34: Preparation of 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-3-chloro-N-(thiazol-4-yl)benzenesulfonamide



15


¹H NMR (500 MHz, MeOD): 8.70(d, 1H), 7.75(s, 1H), 7.47(d, 1H), 7.07(t, 1H), 7.07(t, 1H), 6.64(d, 1H), 6.58(t, 1H), 6.37(d, 1H), 3.71(m, 1H), 3.59(m, 2H), 3.25(m, 1H), 3.13(m, 1H), 2.13(m, 1H), 1.77(m, 1H)

Example 35: Preparation of 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.72(s, 1H), 7.74(d, 1H), 7.10(t, 1H),
 5 6.98(d, 1H), 6.68(d, 1H), 6.63(t, 1H), 6.06(d, 1H), 3.73(m, 1H), 3.53(m, 2H),
 3.24(m, 2H), 2.23(m, 1H), 1.87(m, 1H)


Example 36: Preparation of 4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



10

1H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.58(d, 2H), 7.14(d, 1H),
 7.05(t, 1H), 6.96(d, 1H), 6.91-6.87(m, 2H), 6.75(d, 2H), 3.34-3.30(m, 2H), 3.18-
 3.15(m, 2H), 3.00-2.97(m, 1H), 2.35(s, 3H), 2.15-2.12(m, 1H), 1.75-1.71(m, 1H)


15 **Example 37: Preparation of 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide**



1H NMR (500 MHz, MeOD): 8.69(d, 1H), 7.65(t, 1H), 7.35(d, 1H),


7.20(m, 2H), 6.87(d, 1H), 6.74(d, 1H), 6.66(d, 1H), 3.65(m, 1H), 3.47(m, 1H),
3.37(m, 2H), 3.03(m, 1H), 2.60(s, 3H), 2.30(m, 1H), 2.02(m, 1H)

Example 38: Preparation of 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




1H NMR (500 MHz, MeOD): 8.67(d, 1H), 7.80(d, 1H), 7.53(d, 1H),
7.25(d, 1H), 7.17(s, 1H), 7.13(d, 1H), 6.89(d, 1H), 6.59(d, 1H), 3.35(m, 2H),
3.28(m, 1H), 3.12(m, 2H), 2.37(s, 3H), 2.11(m, 1H), 1.75(m, 1H)

Example 39: Preparation of 4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.58(t, 1H), 7.12(d, 1H),
6.70(t, 1H), 6.66(m, 2H), 6.50(d, 1H), 6.38(d, 1H), 3.59(m, 1H), 3.41(m, 2H),
3.32(m, 1H), 3.12(m, 1H), 2.54(s, 3H), 2.24(m, 1H), 1.93(m, 1H)


Example 40: Preparation of 3-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.76(s, 1H), 7.47(d, 1H), 7.09(d, 1H), 6.81(t, 2H), 6.65(m, 2H), 6.39(d, 1H), 3.40(m, 3H), 3.18(m, 2H), 2.44(s, 3H), 2.15(m, 1H), 1.81(m, 1H)

5


Example 41: Preparation of (R)-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.54(d, 2H), 7.02(t, 1H), 6.92(s, 1H), 6.60(d, 1H), 6.52-6.49(m, 3H), 3.57-3.53(m, 1H), 3.49-3.46(m, 1H), 3.26-3.21(m, 3H), 2.10-2.06(m, 1H), 2.00-1.94(m, 1H)

10


Example 42: Preparation of (R)-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



15


¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.54(t, 1H), 7.01(t, 1H), 6.90(d, 1H), 6.59(t, 1H), 6.50(d, 1H), 6.32(d, 1H), 6.17(d, 1H), 3.43-3.41(m, 2H), 3.26-3.21(m, 2H), 2.21-1.94(m, 3H)

Example 43: Preparation of (R)-3-chloro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.73(s, 1H), 7.44(d, 1H),
 5 7.04(t, 1H), 6.92(s, 1H), 6.61(d, 1H), 6.53(t, 1H), 6.28(d, 1H), 3.47-3.41(m, 3H),
 3.26-3.24(m, 2H), 2.12-1.97(m, 2H)

Example 44: Preparation of (R)-3,5-difluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




10

1H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.34(d, 2H), 7.04(s, 1H),
 6.92(t, 1H), 6.57(d, 1H), 6.47(t, 1H), 3.56-3.25(m, 4H), 3.13-3.11(m, 1H), 2.04-
 1.99(m, 2H)

15


Example 45: Preparation of (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



1H NMR (500 MHz, MeOD): 8.70(s, 1H), 7.70(d, 1H), 7.04(t, 1H),
 6.96(d, 1H), 6.65(d, 1H), 6.55(t, 1H), 5.92(d, 1H), 3.46-3.40(m, 3H), 3.37-


3.33(m, 2H), 2.15-1.96(m, 2H)

Example 46: Preparation of 2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.69(d, 1H), 7.65(t, 1H), 7.08(lm, 2H), 6.88(m, 3H), 6.74(m, 1H), 3.56(m, 1H), 3.37(m, 2H), 3.34(m, 1H), 3.17(m, 1H), 3.16(s, 3H), 2.35(m, 1H), 2.05(m, 1H)

Example 47: Preparation of 3-chloro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.67(d, 1H), 7.84(d, 1H), 7.63(d, 1H), 7.19(d, 1H), 7.05(m, 2H), 6.87(s, 1H), 6.79(m, 1H), 3.60(m, 1H), 3.52(m, 1H), 3.33(m, 1H), 3.26(m, 1H), 3.20(m, 1H), 2.56(s, 3H), 2.27(m, 1H), 1.90(m, 1H)


Example 48: Preparation of 5-chloro-2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,



¹H NMR (500 MHz, MeOD): 8.65(s, 1H), 7.81(d, 1H), 7.10(m, 1H), 7.05(m, 1H), 6.84(m, 3H), 3.53(m, 3H), 3.25(m, 2H), 2.56(s, 3H), 2.26(m, 1H), 1.91(m, 1H)


5

Example 49: Preparation of 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.57(t, 1H), 7.14-7.09(m, 2H), 6.97(d, 1H), 6.92-6.90(m, 1H), 6.84(s, 1H), 6.55(d, 1H), 6.40(d, 1H), 3.42-3.41(m, 1H), 3.34-3.30(m, 1H), 3.26-3.21(m, 2H), 3.04-3.02(m, 1H), 2.42(s, 3H), 2.20-2.17(m, 1H), 1.85-1.81(m, 1H)


Example 50: Preparation of 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.66(s, 1H), 7.76(s, 1H), 7.48(d, 1H),


7.15-7.10(m, 2H), 6.96(d, 1H), 6.89-6.86(m, 1H), 6.83(s, 1H), 6.52(d 1H), 3.30-3.25(m, 3H), 3.11-3.07(m, 2H), 2.34(s, 3H), 2.10-2.07(m, 1H), 1.73-1.69(m, 1H)

Example 51: Preparation of 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.65(s, 1H), 7.43(d, 2H), 7.04(d, 1H), 6.93-6.87(m, 2H), 6.75(s, 1H), 6.64(d, 1H), 3.63-3.60(m, 1H), 3.45-3.43(m, 2H), 3.19-3.17(m, 1H), 2.96-2.92(m, 1H), 2.61(s, 3H), 2.32-2.28(m, 1H), 1.98-1.95(m, 1H)


Example 52: Preparation of 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.66(s, 1H), 7.74(d, 1H), 7.20(t, 1H), 7.12(d, 1H), 6.96(d, 1H), 6.92-6.89(m, 1H), 6.79(s, 1H), 6.12(d, 1H), 3.30(s, 3H), 3.16-3.11(m, 2H), 2.40(s, 3H), 2.15-2.12(m, 1H), 1.79-1.78(m, 1H)

20


Example 53: Preparation of 2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.71(s, 1H), 7.55(t, 1H), 7.02(d, 1H), 6.89(s, 1H), 6.49(m, 3H), 6.33(d, 1H), 3.78(s, 3H), 3.67(m, 1H), 3.42(m, 1H), 3.38(m, 2H), 3.10(m, 1H), 2.58(s, 3H), 2.27(m, 1H), 1.99(m, 1H)

5


Example 54: Preparation of 3-chloro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



10

¹H NMR (500 MHz, MeOD): 8.70(d, 1H), 7.74(d, 1H), 7.45(d, 1H), 7.00(d, 1H), 6.94(d, 1H), 6.49(m, 2H), 6.37(d, 1H), 3.78(s, 3H), 3.47(m, 1H), 3.38(m, 2H), 3.27(m, 1H), 3.16(m, 1H), 2.46(s, 3H), 2.17(m, 1H), 1.83(m, 1H)


Example 55: Preparation of 5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.71(d, 1H), 7.72(d, 1H), 7.01(d, 1H), 6.89(d, 1H), 6.51(m, 2H), 6.48(d, 1H), 3.81(s, 3H), 3.67(m, 1H), 3.45(m, 3H),

3.20(m, 1H), 2.59(s, 3H), 2.27(m, 1H), 1.96(m, 1H)


Example 56: Preparation of (R)-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.59(d, 2H), 7.13(d, 1H), 6.93(s, 1H), 6.78(d, 2H), 6.62(d, 2H), 3.77(s, 3H), 3.25(m, 1H), 2.87-2.65(m, 4H), 2.39(s, 3H), 1.85(m, 1H), 1.70(m, 1H), 1.54(m, 2H)

10

Example 57: Preparation of (R)-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.70(s, 1H), 7.58(t, 1H), 7.12(d, 1H), 6.88(s, 1H), 6.65-6.58(m, 3H), 6.49(d, 1H), 3.78(s, 3H), 3.25(m, 1H), 2.89-2.77(m, 4H), 2.47(s, 3H), 1.88(m, 2H), 1.46(m, 2H)

15

Example 58: Preparation of (R)-3-chloro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


20



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.77(s, 1H), 7.51(d, 1H), 7.17(d, 1H), 6.90(s, 1H), 6.72(d, 1H), 6.67(d, 2H), 3.78(s, 3H), 2.96-2.94(m, 1H), 2.70-2.45(m, 3H), 2.33(s, 3H), 1.88(m, 1H), 1.68(m, 1H), 1.44-1.37(m, 3H)


5

Example 59: Preparation of (R)-3,5-difluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.40(d, 2H), 6.80(t, 2H), 6.63(s, 1H), 6.56(d, 1H), 3.74(s, 3H), 2.90-2.70(m, 4H), 2.53(s, 3H), 1.86-1.28(m, 5H)


Example 60: Preparation of (R)-5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.68(s, 1H), 7.74(d, 1H), 7.15(d, 1H), 6.81(s, 1H), 6.69(d, 2H), 6.31(d, 1H), 3.79(s, 3H), 2.99(m, 1H), 2.67-2.49(m,

3H), 2.39(s, 3H), 1.88(m, 1H), 1.68(m, 1H), 1.35-1.28(m, 3H)

Example 61: Preparation of 3-chloro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




5

¹H NMR (500 MHz, MeOD): 8.70(s, 1H), 7.81(s, 1H), 7.54(d, 1H), 7.24(d, 1H), 7.00(s, 1H), 6.89-6.79(m, 3H), 2.98-2.92(m, 4H), 2.37(m, 4H), 2.20(s, 3H)

10


Example 62: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



15

¹H NMR (500 MHz, MeOD): 8.71(s, 1H), 7.77(d, 1H), 7.24(t, 1H), 6.98(s, 1H), 6.91-6.82(m, 2H), 6.40(d, 1H), 2.96(m, 4H), 2.36(m, 4H), 2.15(s, 3H)


Example 63: Preparation of (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 8.62(d, 1H), 7.79(d, 1H), 6.98(m, 2H), 6.52(m, 2H), 6.23(m, 2H), 3.22(m, 3H), 3.14(m, 1H), 2.70(m, 1H), 2.23(s, 6H), 1.78(m, 2H)


5

Example 64: Preparation of (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, CDCl₃): 8.63(s, 1H), 7.83(d, 1H), 7.17(m, 3H), 6.99(s, 1H), 6.64(s, 1H), 6.56(d, 1H), 3.25(m, 4H), 2.81(m, 1H), 2.24(s, 6H), 1.85(m, 2H)

Example 65: Preparation of 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.71(s, 1H), 7.72(d, 1H), 7.28(m, 3H), 6.06(s, 1H), 6.06(d, 1H), 3.37(m, 4H), 2.66(m, 4H), 2.37(s, 3H)

Example 66: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(4-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




5

¹H NMR (500 MHz, MeOD): 8.72(s, 1H), 7.78(s, 1H), 7.28(d, 1H), 6.98-6.82(m, 4H), 6.49(d, 1H), 3.39-3.30(m, 4H), 2.97(m, 4H), 2.25(s, 3H)

Example 67: Preparation of 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


10



¹H NMR (500 MHz, MeOD): 8.71(s, 1H), 7.81(d, 1H), 7.35(d, 1H), 7.03-6.98(m, 3H), 6.59(d, 1H), 4.09(d, 1H), 2.97(m, 4H), 2.42(m, 4H), 2.22(s, 3H)

15


Example 68: Preparation of (S)-3-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 8.66(d, 1H), 7.80(d, 1H), 7.52(dd, 1H), 7.28(s, 1H), 7.14(m, 2H), 7.05(d, 1H), 6.89(d, 1H), 6.64(s, 1H), 3.22(m, 2H), 3.16(m, 2H), 2.79(m, 1H), 2.23(s, 6H), 2.10(m, 1H), 2.06(m, 1H)


5

Example 69: Preparation of (S)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.70(d, 1H), 7.56(t, 1H), 7.06(t, 1H), 6.90(d, 1H), 6.65(d, 1H), 6.57(t, 1H), 6.43(d, 1H), 6.30(d, 1H), 3.37(m, 3H), 3.24(m, 1H), 3.13(m, 1H), 2.44(s, 3H), 2.18(m, 1H), 2.00(m, 1H)


Example 70: Preparation of (S)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.71(s, 1H), 7.73(d, 1H), 7.07(t, 1H),


6.92(s, 1H), 6.67(d, 1H), 6.59(t, 1H), 6.01(d, 1H), 3.45(m, 3H), 3.23(m, 2H),
2.46(s, 3H), 2.17(m, 1H), 1.83(m, 1H)

Example 71: Preparation of (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-methoxyphenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, CDCl₃): 8.58(d, 1H), 7.76(d, 1H), 6.99(m, 2H),
6.37(m, 2H), 6.26(m, 2H), 3.80(s, 3H), 3.25-3.21(m, 5H), 2.76(m, 1H), 2.21(s,
6H), 2.05(m, 1H)

Example 72: Preparation of (S)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 8.63(d, 1H), 7.79(d, 1H), 7.04(m, 1H),
6.96(d, 1H), 6.57(m, 2H), 6.50(d, 1H), 6.29(m, 2H), 3.22(m, 4H), 2.75(m, 1H),
2.22(s, 6H), 2.05(m, 1H), 1.78(m, 1H)


Example 73: Preparation of (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 8.64(s, 1H), 7.79(d, 1H), 7.01(t, 1H), 6.98(s, 1H), 6.51(m, 2H), 6.23(m, 2H), 3.24(m, 5H), 2.71(m, 1H), 2.20(s, 6H), 1.77(m, 1H)

5

Example 74: Preparation of (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, CDCl₃): 8.64(d, 1H), 7.84(d, 1H), 7.24(d, 1H), 7.12(m, 2H), 6.98(d, 1H), 6.66(s, 1H), 6.54(d, 1H), 3.23(m, 4H), 2.83(m, 1H), 2.23(s, 6H), 1.90(m, 2H)

10

Example 75: Preparation of (R)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide


15



¹H NMR (500 MHz, CDCl₃): 8.62(d, 1H), 7.79(d, 1H), 7.05(d, 1H),

6.98(d, 1H), 6.57(s, 1H), 6.50(t, 1H), 6.31(m, 2H), 3.24(m, 3H), 3.15(m, 1H),
2.74(m, 1H), 2.21(s, 6H), 1.77(m, 2H)


Example 76: Preparation of 4-((2-(1,4-diazepan-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.76(d, 1H), 7.20(m, 1H),
6.97(d, 1H), 6.92(s, 1H), 6.82(t, 1H), 6.22(d, 1H), 3.30(m, 2H), 3.20(t, 2H),
3.10(m, 4H), 1.95(m, 2H)

10

Example 77: Preparation of 5-chloro-4-((4-cyano-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.69(s, 1H), 7.80(d, 1H), 7.29(s, 1H),
7.24(m, 2H), 6.90(s, 1H), 6.33(d, 1H), 3.42(m, 3H), 3.16(m, 2H), 2.47(s, 3H),
2.18(m, 1H), 1.85(m, 1H)

15

Example 78: Preparation of (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide


20



¹H NMR (500 MHz, CDCl₃): 9.25(broad, 1H), 8.64(d, 1H), 7.84(d, 1H), 7.18(m, 2H), 7.02(d, 1H), 6.69(s, 1H), 6.52(d, 1H), 5.60(d, 1H), 4.50(m, 1H), 3.37(m, 1H), 3.27(m, 1H), 3.09(m, 2H), 2.27(m, 1H), 1.95(s, 3H), 1.77(m, 1H)

5


Example 79: Preparation of (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide



¹H NMR (500 MHz, CDCl₃): 9.34(s, 1H), 8.69(d, 1H), 7.79(d, 1H), 7.52(d, 1H), 7.28(d, 1H), 7.18(m, 2H), 7.07(s, 1H), 6.86(d, 1H), 6.69(s, 1H), 5.59(d, 1H), 4.51(m, 1H), 3.49(s, 3H), 3.33(m, 1H), 3.26(m, 1H), 3.06(m, 2H), 2.27(m, 1H), 1.94(s, 3H), 1.75(m, 1H)

15


Example 80: Preparation of (S)-3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.70(s, 1H), 7.82(s, 1H), 7.55(d, 1H), 7.23(m, 3H), 6.95(s, 1H), 6.69(d, 1H), 3.60(m, 1H), 3.44(m, 2H), 3.30(m, 1H), 3.15(m, 1H), 2.57(s, 3H), 2.24(m, 1H), 1.94(m, 1H)

5

Example 81: Preparation of (S)-5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, MeOD): 8.65(s, 1H), 7.77(d, 1H), 7.24(d, 1H), 7.19(s, 1H), 7.16(d, 1H), 6.74s, 1H), 6.24(d, 1H), 3.57(m, 1H), 3.45(m, 2H), 3.30(m, 1H), 3.17(m, 1H), 2.54(s, 3H), 2.22(m, 1H), 1.93(m, 1H)

10

Example 82: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)azetidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


15



¹H NMR (500 MHz, MeOD): 8.86(s, 1H), 8.17(s, 1H), 7.07(d, 1H),

6.84(d, 1H), 6.79(d, 1H), 6.40(m, 1H), 6.33(s, 1H), 3.93-3.70 (m, 4H), 3.35(m, 1H), 3.23(s, 3H)


Example 83: Preparation of 4-((2-(3-aminoazetidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, MeOD): 8.84(s, 1H), 8.07(s, 1H), 7.11(d, 1H), 6.85(d, 1H), 6.81(d, 1H), 6.40(m, 1H), 6.33(s, 1H), 4.03-3.82 (m, 4H), 3.42(m, 1H)

10

Example 84: Preparation of 5-chloro-4-((2-(3-(dimethylamino)azetidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




15

¹H NMR (500 MHz, MeOD): 8.76(s, 1H), 8.10(s, 1H), 7.07(d, 1H), 6.84(d, 1H), 6.79(d, 1H), 6.40(m, 1H), 6.33(s, 1H), 3.99-3.65 (m, 4H), 3.33(m, 1H), 3.12(s, 6H)

Example 85: Preparation of N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)azetidin-3-yl)acetamide


20



¹H NMR (500 MHz, MeOD): 8.56(s, 1H), 8.00(s, 1H), 7.00(d, 1H), 6.77(d, 1H), 6.71(d, 1H), 6.30(m, 1H), 6.29(s, 1H), 3.91-3.55 (m, 4H), 3.35(m, 1H), 2.13(s, 3H)

5

Example 86: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide




¹H NMR (500 MHz, CDCl₃): 10.93(s, 1H), 8.74(d, 1H), 7.81(d, 1H), 6.99(t, 1H), 6.50(m, 2H), 6.24(m, 2H), 3.92(t, 1H), 3.32(m, 2H), 3.26(s, 3H), 3.11(m, 2H), 2.01(m, 1H), 1.92(m, 1H), 1.69(m, 1H)

10

Example 87: Preparation of 5-chloro-2-fluoro-4-((2-(3-methoxypyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide


15



¹H NMR (500 MHz, CDCl₃): 10.79(s, 1H), 8.74(d, 1H), 7.85(d, 1H), 7.25(m, 2H), 7.15(s, 1H), 7.11(d, 1H), 6.94(d, 1H), 6.66(s, 1H), 6.56(d, 1H),


3.97(t, 1H), 3.31(m, 2H), 3.30(s, 3H), 3.12(m, 2H), 2.04(m, 2H)

Example 88: Preparation of (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide




¹H NMR (500 MHz, CDCl₃): 10.63(s, 1H), 8.70(d, 1H), 7.78(d, 1H), 7.01(m, 1H), 6.94(d, 1H), 6.51(m, 2H), 6.30(s, 1H), 6.17(d, 1H), 5.78(d, 1H), 4.44(m, 1H), 3.38(m, 1H), 3.27(m, 1H), 3.10(m, 1H), 3.03(m, 1H), 2.15(m, 1H), 1.90(s, 3H), 1.75(m, 1H)

Example 89: Preparation of 3-chloro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 10.74(s, 1H), 8.77(s, 1H), 7.72(s, 1H), 7.42(d, 1H), 7.01(m, 2H), 6.53(m, 3H), 6.18(s, 1H), 3.90(s, 1H), 3.34(m, 2H), 3.25(s, 3H), 3.12(m, 2H), 1.99(m, 1H), 1.89(m, 1H)


Example 90: Preparation of 3-chloro-4-((2-(3-methoxypyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



¹H NMR (500 MHz, CDCl₃): 10.38(s, 1H), 8.76(s, 1H), 7.78(s, 1H), 7.50(d, 1H), 7.26(d, 1H), 7.14(m, 2H), 7.02(s, 1H), 6.91(d, 1H), 6.65(s, 1H), 3.97(s, 1H), 3.33(m, 5H), 3.12(m, 2H), 2.03(m, 2H)


5

Example 91: Preparation of (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide



¹H NMR (500 MHz, CDCl₃): 10.43(broad, 1H), 8.76(s, 1H), 7.71(s, 1H), 7.41(s, 1H), 7.03(s, 2H), 6.54(m, 3H), 6.25(s, 1H), 5.90(s, 1H), 4.40(s, 1H), 3.28(m, 2H), 2.98(m, 3H), 2.15(m, 1H), 2.03-1.74(m, 3H)

Example 92: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



Step 1) Preparation of tert-butyl (2-((2-amino-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate(iii)

2,4-difluoro-1-nitrobenzene (i, 2.0 g, 12.6 mmol) and tert-butylmethyl(2-(methylamino)ethyl)carbamate (ii, 2.4 g, 1.0 eq.) were dissolved in DMF (20 mL), and then K_2CO_3 (2.6 g, 1.5 eq.) was added thereto. While maintaining the internal temperature at 60 to 70°C, the reaction mixture was stirred for 2 hours. When the reaction solution became dark yellow, the completion of the reaction was confirmed by TLC. After cooling to room temperature, EA/H₂O was added and stirred, and then the layers were separated. MgSO₄ was added to the separated organic layer, which was stirred, filtered and then dried. The filtrate was concentrated under reduced pressure, and the residue was dissolved in MeOH (0.13g, 0.1 eq.) and then Pd/C (0.13 g, 0.1 eq.) was added. The inside was replaced with hydrogen gas, and the reaction mixture was stirred at room temperature for 6 hours. When the yellow color of the reaction solution faded and became almost colorless, the completion of the reaction was confirmed by TLC. The metal catalyst was filtered through Celite[™]. The filtrate was concentrated under reduced pressure, and the obtained residue was separated by column chromatography (Hx/EA = 3/1) to obtain the target compound (iii, 2.5 g, 66.9%).

¹H NMR (500 MHz, CDCl₃): 6.76(d, 1H), 6.64(m, 2H), 3.36(m, 2H), 2.94(m, 2H), 2.86(m, 3H), 2.68(s, 3H), 1.45(s, 9H)

Step 2) Preparation of tert-butyl thiazol-4-yl carbamate (v)

Thiazol-4-carboxylic acid (iv, 5.0 g, 38.8 mmol) was dissolved in t-

BuOH (100 mL), and then TEA (8.1 mL, 1.5 eq.) and DPPA (7.1 mL, 1.5 eq.) were added thereto. While maintaining the internal temperature at 90 to 100°C, the reaction mixture was stirred for 3 days, and then the completion of the reaction was confirmed by TLC. The reaction was concentrated under reduced pressure, to which H₂O (50 mL) was added and extracted twice with ethyl acetate (EA, 100 mL). MgSO₄ was added to the organic layer, which was stirred, filtered and dried. The filtrate was concentrated under reduced pressure, and the residue was added to a small amount of EA and slurried. The resulting solid was filtered to obtain a white title compound (v, 4.0 g, 51.5%).

10 ¹H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.24(s, 1H), 1.52(s, 9H)

Step 3) Preparation of tert-butyl ((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(thiazol-4-yl)carbamate (vii)

Tert-butyl thiazol-4-yl carbamate (v, 4.0 g, 20.0 mmol) was added to a reaction vessel and the inside of the vessel was replaced with nitrogen gas. After dissolving in THF (32 mL), the solution was cooled to -78°C using dry ice-acetone. After cooling, LiHMDS (22.4 mL, 1.5 eq.) was slowly added and the reaction mixture was stirred for 30 minutes. Then, 4-bromo-5-chloro-2-fluorobenzenesulfonyl chloride (vi, 6.0 g, 1.0 eq.) was dissolved in THF (10 mL) and then slowly added to the reaction solution. The reaction mixture was stirred overnight and the completion of the reaction was confirmed by TLC. H₂O (50 mL) was added and extracted twice with ethyl acetate (EA, 100 mL). MgSO₄ was added to the organic layer, which was stirred, filtered and dried. The filtrate was concentrated under reduced pressure, and the residue was crystallized with THF/n-hexane to obtain the title compound (vii, 4.4 g, 59.0%).

25 ¹H NMR (500 MHz, MeOD): 9.00(s, 1H), 8.22(d, 1H), 7.90(d, 1H), 7.78(s, 1H), 1.35(s, 9H)

Step 4) Preparation of tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl(methyl)carbamate(viii)

Tert-butyl (2-((2-amino-5-fluorophenyl)(methyl)amino)ethyl(methyl)carbamate (iii, 10.0 g, 33.7 mmol) and tert-butyl ((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(thiazol-4-yl)carbamate (vii, 13.0 g, 1.0 eq.) were dissolved in 1,4-dioxane (200 mL). Pd(OAc)₂ (0.7 g, 0.1 eq), rac-BINAP (4.11 g, 0.2 eq.), and Cs₂CO₃ (21.2 g, 2.0 eq.) were added to the reaction solution. While maintaining the internal temperature at 90 to 100°C, the reaction mixture was stirred for 5 hours, and then the completion of the reaction was confirmed by TLC. H₂O (100 mL) was added and extracted twice with ethyl acetate (EA, 1000 mL). MgSO₄ was added to the organic layer, which was stirred, filtered and then dried. The filtrate was concentrated under reduced pressure, and the residue was separated by column chromatography using a mobile phase of EA/Hex=1/4 to obtain the title compound (16.0 g, 69.1%).

¹H NMR (500 MHz, MeOD): 8.95(d, 1H), 7.96(d, 1H), 7.68(s, 1H), 7.26(s, 1H), 6.95(t, 1H), 6.8(s, 1H), 6.39(s, 1H), 3.27(s, 2H), 3.14(s, 2H), 2.79(s, 3H), 2.70(d, 3H), 1.40(s, 9H), 1.37(s, 9H)


Step 5) Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride

To tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl(methyl)carbamate (vii, 14.0 g, 20.3 mmol) was added 1 M HCl in ethyl acetate (200 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered, and MC (200 mL) was added to the obtained residue and stirred for 1 hour. The resulting solid was filtered to obtain the target compound (9.1 g, 85.3%).

¹H NMR (500 MHz, MeOD): 8.73(d, 1H), 7.79(d, 1H), 7.26(dd, 1H), 7.03(m, 2H), 6.90(td, 1H), 6.43(d, 1H), 3.26(t, 2H), 3.09(t, 2H), 2.70(s, 3H), 2.65(s, 3H)

Example 93: Preparation of 3-chloro-4-((2-((2-


(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide



An intermediate was prepared in the same manner as described in Example 92, except that N,N,N'-trimethylethane-1,2-diamine was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate (ii) and 4-bromo-5-chlorobenzene sulfonyl chloride was used instead of 4-bromo-5-chloro-2-fluorobenzenesulfonyl chloride (vi). To the obtained intermediate tert-butyl ((3-chloro-4-((2-((dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)phenyl)sulfonyl)(thiazol-4-yl)carbamate (0.05 g, 0.09 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 48.0%).

¹H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.81(s, 1H), 7.54(d, 1H), 7.21(dd, 1H), 7.00(m, 2H), 6.82(dd, 1H), 6.75(d, 1H), 3.20(t, 2H), 2.93(t, 2H), 2.70(s, 3H), 2.55(s, 3H)

Example 94: Preparation of 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide




An intermediate was prepared in the same manner as described in

Example 92, except that N,N,N'-trimethylethane-1,2-diamine was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate (ii). To the obtained intermediate tert-butyl ((5-chloro-4-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluorophenyl)sulfonyl)(thiazol-4-yl)carbamate (0.05 g, 0.08 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 49.8%).

¹H NMR (500 MHz, MeOD): 8.72(s, 1H), 7.76(d, 1H), 7.2(t, 1H), 6.95(m, 2H), 6.80(t, 1H), 6.38(d, 1H), 3.01(t, 2H), 2.68(s, 3H), 2.40(t, 2H), 2.15(s, 3H)


Example 95: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 92, except that 2-fluoro-1-nitro-4-(trifluoromethyl)benzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-(trifluoromethyl)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 55.5%).

¹H NMR (500 MHz, MeOD): 8.74(s, 1H), 7.88(d, 1H), 7.54(s, 1H), 7.42(m, 2H), 7.06(s, 1H), 7.02(d, 1H), 3.35(t, 2H), 3.16(t, 2H), 2.73(s, 3H), 2.70(s, 3H)

Example 96: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide hydrochloride



5


N-(2,4-dimethoxybenzyl)-5-fluoropyrimidin-2-amine (v) was prepared instead of the step 2 of Example 92. Specifically, 2-chloro-5-fluoropyrimidine (iv, 0.48 g, 3.62 mmol), (2,4-dimethoxyphenyl)methaneamine (0.60 g, 1.0 eq.) and trimethylamine (0.76 mL, 1.5 eq.) were dissolved in EtOH (10 mL). After stirring the mixture stirred overnight while heating to 70 to 80°C, the completion of the reaction was confirmed by TLC. The solution was concentrated under reduced pressure, and the obtained residue was separated by column chromatography using a mobile phase of EA/Hex=1/2 to obtain 0.32 g (yield 34%) of the target compound (v).

15 ¹H NMR (500 MHz, MeOD): 8.20(s, 2H), 7.13(d, 1H), 6.51(s, 1H), 6.41(d, 1H), 4.44(s, 2H), 3.82(s, 3H), 3.76(s, 3H)

The target compound was prepared in the same manner as described in steps 1, 3, 4 and 5 of Example 92, except that N-(2,4-dimethoxybenzyl)-5-fluoropyrimidin-2-amine prepared above was used instead of tert-butylthiazol-4-ylcarbamate (v) in Example 92.

20 ¹H NMR (500 MHz, MeOD): 8.43(m, 2H), 7.99(t, 1H), 7.27(dd, 1H), 7.03(d, 1H), 6.88(t, 1H), 6.45(d, 1H), 3.20(t, 2H), 3.10(t, 2H), 2.71(s, 3H), 2.67(s, 3H)

Example 97: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)benzenesulfonamide hydrochloride



5

An intermediate was prepared in the same manner as described in Example 96, except that 2-fluoro-1-nitro-4-(trifluoromethyl)benzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-


10 (trifluoromethyl)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target

15 compound (0.02 g, 58.3%).

¹H NMR (500 MHz, MeOD): 8.47(s, 1H), 8.29(s, 1H), 8.05(m, 1H), 7.53(d, 1H), 7.41(m, 2H), 7.01(t, 1h), 3.45(t, 2H), 3.16(t, 2H), 2.74(s, 3H), 2.69(s, 3H)

20

Example 98: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 92, except that 4-(difluoromethoxy)-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-

5 fluorophenyl)amino)-5-(difluoromethoxy)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 55.7%).

10 ¹H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.80(d, 1H), 7.29(d, 1H), 7.02(s, 2H), 6.94(d, 1H), 6.85(t, 1H), 6.55(d, 1H), 3.25(t, 2H), 3.10(t, 2H), 2.70(s, 3H), 2.66(s, 3H)

Example 99: Preparation of 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride

15



An intermediate was prepared in the same manner as described in Example 92, except that 3-fluoro-4-nitrobenzonitrile was used instead of 2,4-


20 difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-cyanophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was

25 confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.1%).

¹H NMR (500 MHz, MeOD): 8.75(s, 1H), 7.91(d, 1H), 7.62(d, 1H), 7.47(t, 1H), 7.32(d, 1H), 7.17(d, 1H), 7.07(d, 1H), 3.28(t, 2H), 3.17(t, 2H), 2.71(s,

3H), 2.70(s, 3H)


Example 100: Preparation of 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 3-fluoro-4-nitrobenzonitrile was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-cyanophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.7%).

¹H NMR (500 MHz, MeOD): 8.43(s, 2H), 8.10(d, 1H), 7.63(s, 1H), 7.48(d, 1H), 7.36(d, 1H), 7.17(d, 1H), 3.34-3.32(m, 2H), 3.19-3.16(m, 2H), 2.72(s, 3H), 2.71(s, 3H)


Example 101: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 4-(difluoromethoxy)-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(difluoromethoxy)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.2%).

¹H NMR (500 MHz, MeOD): 8.42(s, 2H), 8.01(d, 1H), 7.33-7.17(m, 3H), 6.71(d, 1H), 3.35-3.32(m, 2H), 3.13-3.10(m, 2H), 2.68(s, 3H), 2.67(s, 3H)


Example 102: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.7%).

¹H NMR (500 MHz, MeOD): 8.42(s, 2H), 8.00(d, 1H), 7.30(d, 1H), 7.03(d, 1H), 6.94(d, 1H), 6.69(s, 1H), 6.52(d, 1H), 3.34-3.32(m, 2H), 3.12-3.10(m, 2H), 2.70(s, 3H), 2.67(s, 3H)

5 **Example 103: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyridin-2-yl)benzenesulfonamide hydrochloride**



N-(2,4-dimethoxybenzyl)-5-fluoropyridin-2-amine (v) was prepared
 10 instead of the step 2 of Example 92. Specifically, 5-fluoropyridin-2-amine (iv, 0.55 g, 0.01 mmol) and 2,4-dimethoxybenzaldehyde (0.45 g, 0.9 eq.) were dissolved in DCM (10 mL). After stirring at room temperature for 1 hour, sodium triacetoxyborohydride (1.0 g, 1 eq.) was added three times at 15 minute intervals. After stirring the mixture stirred overnight, the completion of the
 15 reaction was confirmed by TLC. H₂O (10 mL) was added and extracted twice with dichloromethane (10 mL). MgSO₄ was added to the organic layer, which was stirred, filtered and dried. The filtrate was concentrated under reduced pressure, and then the resulting residue was separated by column chromatography using a mobile phase of EA/Hex=1/4 to obtain 0.65 g (yield
 20 51%) of N-(2,4-dimethoxybenzyl)-5-fluoropyridin-2-amine(v).


¹H NMR (500 MHz, CDCl₃): 7.90(d, 1H), 7.16(m, 2H), 6.46(d, 1H), 6.41(d, 1H), 6.34(m, 1H), 4.36(s, 2H), 3.81(s, 3H), 3.77(s, 3H)

The target compound was prepared in the same manner as
 25 described in steps 1, 3, 4 and 5 of Example 92, except that N-(2,4-

dimethoxybenzyl)-5-fluoropyridin-2-amine prepared above was used instead of tert-butylthiazol-4-ylcarbamate (v) in Example 92.

¹H NMR (500 MHz, MeOD): 8.05(d, 1H), 7.89(d, 1H), 7.51(d, 1H), 7.26-6.87(m, 4H), 6.42(d, 1H), 3.38-3.25(m, 2H), 3.10-3.09(m, 2H), 2.69(s, 3H),
5 2.65(s, 3H)

Example 104: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide hydrochloride



10

An intermediate was prepared in the same manner as described in Example 103, except that pyridine-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((5-chloro-2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(pyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.0%).


15

¹H NMR (500 MHz, MeOD): 8.01-7.92(m, 2H), 7.29-7.24(m, 3H), 7.03(d, 2H), 6.87(d, 1H), 6.46(d, 1H), 3.42-3.40(m, 2H), 3.11-3.08(m, 2H), 2.70(s, 3H), 2.66(s, 3H)

20

Example 105: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide hydrochloride


25



An intermediate was prepared in the same manner as described in Example 103, except that thiazole-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((5-chloro-2-((2-chloro-4-
 5 (N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target
 10 compound (0.02 g, 59.8%).

¹H NMR (500 MHz, MeOD): 7.84(d, 2H), 7.28-7.04(m, 3H), 6.89(d, 1H), 6.74(d, 1H), 6.46(d, 1H), 3.26-3.24(m, 2H), 3.11-3.08(m, 2H), 2.71(s, 3H), 2.66(s, 3H)

15 **Example 106: Preparation of 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide hydrochloride**




An intermediate was prepared in the same manner as described in Example 103, except that 3-fluoro-4-nitrobenzonitrile was used instead of 2,4-
 20 difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was

added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.0%).

5 ¹H NMR (500 MHz, MeOD): 8.05-8.01(m, 2H), 7.62(s, 1H), 7.55-7.46(m, 2H), 7.33(d, 1H), 7.14(d, 1H), 3.34-3.32(m, 2H), 3.18-3.17(m, 2H), 2.72(s, 3H), 2.71(s, 3H)


10 **Example 107: Preparation of 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide hydrochloride**



An intermediate was prepared in the same manner as described in Example 103, except that 3-fluoro-4-nitrobenzonitrile was used instead of 2,4-difluoro-1-nitrobenzene (i), and pyridine-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(pyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-cyanophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 57.2%).

20 ¹H NMR (500 MHz, MeOD): 8.05(d, 1H), 7.97-7.86(s, 2H), 7.62(s, 1H), 7.46(d, 1H), 7.33-7.19(m, 3H), 7.01-6.97(m, 1H), 3.32-3.31(m, 2H), 3.19-3.16(m, 2H), 2.72(s, 3H), 2.71(s, 3H)


Example 108: Preparation of 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-

yl)benzenesulfonamide hydrochloride

An intermediate was prepared in the same manner as described in Example 103, except that 3-fluoro-4-nitrobenzonitrile was used instead of 2,4-difluoro-1-nitrobenzene (i), and thiazole-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-cyanophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.2%).

¹H NMR (500 MHz, MeOD): 7.96(d, 1H), 7.62(s, 1H), 7.46(d, 1H), 7.28-7.14(m, 3H), 6.77(d, 1H), 3.32-3.30(m, 2H), 3.20-3.17(m, 2H), 2.72(s, 3H), 2.71(s, 3H)

Example 109: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)benzenesulfonamide hydrochloride



20

An intermediate was prepared in the same manner as described in Example 103, except that 2-fluoro-1-nitro-4-(trifluoromethyl)benzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-


5-fluorophenyl)amino)-5-

(trifluoromethyl)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.2%).

¹H NMR (500 MHz, MeOD): 8.05(d, 1H), 7.99(d, 1H), 7.53(m, 2H), 7.43(s, 2H), 7.17(m, 1H), 7.02(d, 1H), 3.47(t, 2H), 3.17(t, 2H), 2.74(s, 3H), 2.70(s, 3H)

10

Example 110: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide hydrochloride




15 An intermediate was prepared in the same manner as described in Example 103, except that 2-fluoro-1-nitro-4-(trifluoromethyl)benzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and pyridin-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(pyridin-2-yl)sulfamoyl-5-fluorophenyl)amino)-5-

20 (trifluoromethyl)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.8%).

¹H NMR (500 MHz, MeOD): 8.02(d, 1H), 7.98(t, 1H), 7.86(t, 1H), 7.53(s, 1H), 7.40(m, 2H), 7.31(d, 1H), 7.05(d, 1H), 6.99(m, 1H), 3.34(t, 2H), 3.17(t, 2H), 2.74(s, 3H), 2.71(s, 3H)


Example 111: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide hydrochloride



5 An intermediate was prepared in the same manner as described in Example 103, except that 2-fluoro-1-nitro-4-(trifluoromethyl)benzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and thiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-
10 chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(trifluoromethyl)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction
15 solution was filtered to obtain the target compound (0.02 g, 58.6%).

¹H NMR (500 MHz, MeOD): 7.93(d, 1H), 7.53(s, 1H), 7.41(m, 2H), 7.13(d, 1H), 7.07(d, 1H), 6.77(d, 1H), 3.34(t, 2H), 3.18(t, 2H), 2.75(s, 3H), 2.71(s, 3H)


20 **Example 112: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride**



An intermediate was prepared in the same manner as described in Example 103, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 61.3%).

¹H NMR (500 MHz, MeOD): 8.06(d, 1H), 7.91(d, 1H), 7.51(m, 1H), 7.30(m, 2H), 7.22(t, 1H), 7.15(m, 2H), 6.70(d, 1H), 3.26(t, 2H), 3.11(t, 2H), 2.67(s, 6H)

Example 113: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 103, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and pyridine-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(pyridin-2-yl)sulfamoyl)-5-

fluorophenyl)amino)phenyl)(methylamino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 61.6%).

¹H NMR (500 MHz, MeOD): 8.03(d, 1H), 7.96(d, 1H), 7.90(d, 1H), 7.30(t, 3H), 7.23(t, 1H), 7.17(t, 1H), 7.04(m, 1H), 6.73(d, 1H), 3.26(t, 2H), 3.12(t, 2H), 2.68(s, 3H), 2.67(s, 3H)


Example 114: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and thiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methylamino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 61.3%).


¹H NMR (500 MHz, MeOD): 7.86(d, 1H), 7.30(m, 2H), 7.20(m, 2H), 7.12(d, 1H), 6.75(m, 2H), 3.26(t, 2H), 3.12(t, 2H), 2.70(s, 3H), 2.68(s, 3H)

Example 115: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-

fluoropyridin-2-yl)benzenesulfonamide hydrochloride

An intermediate was prepared in the same manner as described in Example 103, except that 4-(difluoromethoxy)-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(difluoromethyl)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.2%).

¹H NMR (500 MHz, MeOD): 8.06(d, 1H), 7.90(d, 1H), 7.52(m, 1H), 7.29(d, 1H), 7.16(m, 1H), 7.02(d, 1H), 6.94(m, 1H), 6.84(t, 1H), 6.53(d, 1H), 3.26(t, 2H), 3.10(t, 2H), 2.70(s, 3H), 2.66(s, 3H)

Example 116: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide hydrochloride


20

An intermediate was prepared in the same manner as described in Example 103, except that 4-(difluoromethoxy)-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and thiazol-2-amine was used instead

of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(difluoromethoxy)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.7%).

¹H NMR (500 MHz, MeOD): 7.85(d, 1H), 7.29(d, 1H), 7.11(d, 1H), 7.03(d, 1H), 6.94(d, 1H), 6.85(t, 1H), 6.75(d, 1H), 6.58(d, 1H), 3.23(t, 2H), 3.12(t, 2H), 2.72(s, 3H), 2.67(s, 3H)

Example 117: Preparation of 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 103, except that 4-(difluoromethoxy)-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and pyridine-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(pyridine-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(difluoromethoxy)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.3%).

¹H NMR (500 MHz, MeOD): 8.04(d, 1H), 7.95(d, 1H), 7.90(t, 1H),

7.30(m, 2H), 7.06(m, 2H), 6.93(d, 1H), 6.84(t, 1H), 6.56(d, 1H), 3.26(t, 2H), 3.11(t, 2H), 2.70(s, 3H), 2.67(s, 3H)


Example 118: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-(methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.5%).

¹H NMR (500 MHz, MeOD): 8.05(d, 1H), 7.86(d, 1H), 7.53-7.50(m, 1H), 7.18-7.14(m, 2H), 6.80-6.73(m, 2H), 6.35(d, 1H), 3.82(s, 3H), 3.25-3.23(m, 2H), 3.09-3.07(m, 2H), 2.67(s, 3H), 2.64(s, 3H)

Example 119: Preparation of 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 103, except that 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and thiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.0%).

¹H NMR (500 MHz, MeOD): 7.81(d, 1H), 7.20(d, 1H), 7.11(d, 1H), 6.81-6.73(m, 3H), 6.40(d, 1H), 3.82(s, 3H), 3.30-3.28(m, 2H), 3.25-3.24(m, 2H), 2.69(s, 3H), 2.66(s, 3H)

15

Example 120: Preparation of 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride




20

An intermediate was prepared in the same manner as described in Example 92, except that 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-

fluorophenyl)amino)-5-methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 55.5%).

¹H NMR (500 MHz, MeOD): 8.73(d, 1H), 7.76(d, 1H), 7.18(d, 1H), 7.02(d, 1H), 6.82(d, 1H), 6.75(d, 1H), 6.36(d, 1H), 3.85(s, 3H), 3.26-3.24(m, 2H), 3.09-3.07(m, 2H), 2.69(s, 3H), 2.64(s, 3H)

Example 121: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 92, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)phenyl(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 57.5%).

¹H NMR (500 MHz, MeOD): 8.74(d, 1H), 7.81(d, 1H), 7.33-7.18 (m, 4H), 7.04(d, 1H), 6.70(d, 1H), 3.26-3.24(m, 2H), 3.12-3.09(m, 2H), 2.69(s, 3H), 2.66(s, 3H)

Example 122: Preparation of 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((4-methoxy-2-(methyl(2-


(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.5%).

¹H NMR (500 MHz, MeOD): 8.42(s, 2H), 7.96(d, 1H), 7.18 (d, 1H), 6.82(d, 1H), 6.76(d, 1H), 6.36(d, 1H), 3.81(s, 3H), 3.27-3.25(m, 2H), 3.17-3.08(m, 2H), 2.70(s, 3H), 2.65(s, 3H)

Example 123: Preparation of 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride




20

An intermediate was prepared in the same manner as described in Example 92, except that 4-chloro-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-

(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-chlorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 55.8%).

¹H NMR (500 MHz, MeOD): 8.74(s, 1H), 7.81(d, 1H), 7.31-7.14 (m, 3H), 7.04(d, 1H), 6.64(d, 1H), 3.26-3.24(m, 2H), 3.11-3.10(m, 2H), 2.69(s, 3H), 2.66(s, 3H)


Example 124: Preparation of 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 4-chloro-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((5-chloro-2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.3%).

¹H NMR (500 MHz, MeOD): 8.42(s, 2H), 8.01(d, 1H), 7.27(d, 2H), 7.15-7.13(m, 1H), 6.64(d, 1H), 3.27-3.25(m, 2H), 3.13-3.11(m, 2H), 2.70(s, 3H), 2.67(s, 3H)

Example 125: Preparation of 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide hydrochloride



5

An intermediate was prepared in the same manner as described in Example 103, except that 4-chloro-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((5-chloro-2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyridin-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.3%).


10

15

¹H NMR (500 MHz, MeOD): 8.06(d, 1H), 7.91(d, 1H), 7.54-7.50(m, 2H), 7.27-7.13(m, 4H), 6.63(d, 1H), 3.27-3.25(m, 2H), 3.11-3.09(m, 2H), 2.69(s, 3H), 2.66(s, 3H)

20


Example 126: Preparation of 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 4-chloro-2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i) and thiazole-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((5-chloro-2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(thiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.8%).

¹H NMR (500 MHz, MeOD): 7.87(d, 1H), 7.27(d, 2H), 7.15-7.12(m, 2H), 6.75-6.68(m, 2H), 3.26-3.25(m, 2H), 3.13-3.10(m, 2H), 2.71(s, 3H), 2.68(s, 3H)


Example 127: Preparation of 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 5-chlorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(5-chlorothiazol-2-yl)-N-(2,4-dimethoxybenzyl)sulfamoyl)-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.2%).

¹H NMR (500 MHz, MeOD): 7.82(d, 1H), 7.29(d, 1H), 7.27(s, 1H), 7.05(d, 1H), 6.90(t, 1H), 6.48(d, 1H), 3.27-3.25(m, 2H), 3.12-3.09(m, 2H), 2.71(s, 3H), 2.67(s, 3H)

5 **Example 128: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluorothiazol-2-yl)benzenesulfonamide hydrochloride**




An intermediate was prepared in the same manner as described in
 10 Example 103, except that 5-fluorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluorothiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring
 15 the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.8%).

¹H NMR (500 MHz, MeOD): 7.82(t, 1H), 7.30(t, 1H), 7.05-6.91(m, 3H), 6.48(d, 1H), 3.27-3.25(m, 2H), 3.12-3.10(m, 2H), 2.72(s, 3H), 2.67(s, 3H)

20


Example 129: Preparation of 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 5-chlorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv) and 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(5-chlorothiazol-2-yl)-N-(2,4-dimethoxybenzyl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.8%).

¹H NMR (500 MHz, MeOD): 7.84(d, 1H), 7.33(d, 1H), 7.40(d, 1H), 7.19(m, 3H), 6.75(d, 1H), 3.26(t, 2H), 3.13(t, 2H), 2.70(s, 3H), 2.69(s, 3H)

Example 130: Preparation of 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 103, except that 5-chlorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv) and 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl

(2-((2-((2-chloro-4-(N-(5-chlorothiazol-2-yl)-N-(2,4-dimethoxybenzyl)sulfamoyl)-5-fluorophenyl)amino)-5-methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 58.7%).

¹H NMR (500 MHz, MeOD): 7.79(d, 1H), 7.20(d, 1H), 7.19(s, 1H), 6.80(d, 1H), 6.75(dd, 1H), 6.41(d, 1H), 3.82(s, 3H), 3.26(t, 2H), 3.10(t, 2H), 2.70(s, 3H), 2.68(s, 3H)


Example 131: Preparation of 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 5-fluorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv) and 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluorothiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl) carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.5%).

¹H NMR (500 MHz, MeOD): 7.84(d, 1H), 7.31(m, 2H), 7.22(m, 2H), 7.00(s, 1H), 6.76(d, 1H), 3.27(t, 2H), 3.13(t, 2H), 2.70(s, 3H), 2.69(s, 3H)


Example 132: Preparation of 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 103, except that 5-fluorothiazol-2-amine was used instead of 5-fluoropyridin-2-amine (iv) and 2-fluoro-4-methoxy-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluorothiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-methoxyphenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.3%).

¹H NMR (500 MHz, MeOD): 7.79(t, 1H), 7.22(t, 1H), 6.99(s, 1H), 6.86(d, 1H), 6.78(dd, 1H), 6.40(d, 1H), 3.83(s, 3H), 3.34(t, 2H), 3.11(t, 2H), 2.75(s, 3H), 2.67(s, 3H)

Example 133: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide hydrochloride



Tert-butyl

((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(5-

methyloxazol-3-yl)carbamate (vii) was prepared instead of steps 2 and 3 of Example 92. Specifically, 5-methylisoxazol-3-amine (iv, 1.00 g, 10.19 mmol), 4-bromo-5-chloro-2-fluorobenzenesulfonyl chloride (vi, 3.14 g, 1.0 eq.) and pyridine (2.4 mL, 3.0 eq.) were dissolved in DCM (25 mL). After stirring the mixture stirred overnight, the completion of the reaction was confirmed by TLC. H₂O (30 mL) was added and extracted twice with dichloromethane (10 mL). MgSO₄ was added to the reaction and extracted twice with ethyl acetate. MgSO₄ was added to the organic layer, which was stirred, filtered and then dried. The filtrate was concentrated under reduced pressure, and then the resulting residue was separated by column chromatography using a mobile phase of EA/Hex=1/1 to obtain 1.0 g (yield 27%) of the target compound (v).

¹H NMR (500 MHz, CDCl₃): 8.66(d, 1H), 7.96(d, 1H), 7.44(d, 1H),

5.88(broad, 1H), 2.33(s, 3H)

The above-prepared 4-bromo-5-chloro-2-fluoro-N-(5-methylisoxazol-3-yl)benzenesulfonamide (v, 1.00 g, 2.71 mmol), N,N-dimethylaminopyridine (0.06 g, 0.2 eq.) and di-tert-butyl dicarbonate (1.1 mL, 2.0 eq.) were dissolved in tetrahydrofuran (20 mL). After stirring the reaction solution overnight at room temperature, and the completion of the reaction was confirmed by TLC. H₂O (30 mL) was added to the reaction product, and the mixture was extracted twice with ethyl acetate. MgSO₄ was added to the organic layer, which was stirred, filtrated and then dried. The filtrate was concentrated under reduced pressure, and the obtained residue was separated by column chromatography using a

mobile phase of EA/Hex=1/2 to obtain 0.40 g (yield 31%) of the target compound (vii).

¹H NMR (500 MHz, MeOD): 8.19(d, 1H), 7.93(d, 1H), 6.34(s, 1H), 2.49(s, 3H), 1.36(s, 9H)

5


The target compounds were prepared in the same manner as described in steps 1, 4 and 5 of Example 1, except that the above-prepared tert-butyl ((4-bromo-5-chloro-2-fluorophenyl)sulfonyl)(5-methylisoxazol-3-yl)carbamate (vii) was used.

10

¹H NMR (500 MHz, MeOD): 7.8(d, 1H), 7.34-7.31(m, 2H), 7.25(t, 1H), 7.21-7.19(m, 1H), 6.71(d, 1H), 6.07(s, 1H), 2.70(s, 3H), 2.67(s, 3H), 2.32(s, 3H)

Example 134: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide hydrochloride

15




An intermediate was prepared in the same manner as described in Example 133, except that 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(5-methylisoxazol-3-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 65.6%).

25

¹H NMR (500 MHz, MeOD): 7.79(d, 1H), 7.21(t, 1H), 7.00(d, 1H), 6.98(t, 1H), 6.41(d, 1H), 5.85(s, 1H), 3.24-3.23(m, 2H), 3.05-3.03(m, 2H), 2.69(s,

3H), 2.58(s, 3H), 2.22(s, 3H)

Example 135: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 133, except that tert-butyl 3-amino-5-methyl-1H-pyrazole-1-carboxylate was used instead of 5-methylisoxazol-3-amine (iv). To the obtained intermediate tert-butyl 3-((N-(tert-butoxycarbonyl)-4-(2-(2-((tert-butoxycarbonyl)(methyl)amino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-5-chloro-2-fluorophenyl)sulfonamido)-5-methyl-1H-pyrazole-1-carboxylate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 64.7%).

¹H NMR (500 MHz, MeOD): 7.86(1H), 7.27(t, 1H), 7.06(d, 1H), 6.89(t, 1H), 6.47(d, 1H), 6.08(d, 1H), 3.15-3.12(m, 2H), 2.71(s, 3H), 2.67(s, 3H), 2.34(s, 3H)

20


Example 136: Preparation of 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 133, except that tert-butyl 3-amino-5-methyl-1H-pyrazole-1-carboxylate was used instead of 5-methylisoxazol-3-amine (iv), and 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl 3-((N-(tert-butoxycarbonyl)-4-(2-(2-((tert-butoxycarbonyl)(methyl)amino)ethyl)(methyl)amino)phenyl)amino)-5-chloro-2-fluorophenyl)sulfonamido)-5-methyl-1H-pyrazole-1-carboxylate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 65.7%).

¹H NMR (500 MHz, MeOD): 7.88(d, 1H), 7.35-7.21(m, 4H), 6.72(d, 1H), 6.08(s, 1H), 3.35-3.33(m, 2H), 3.14-3.11(m, 2H), 2.72(s, 3H), 2.68(s, 3H), 2.34(s, 3H)

Example 137: Preparation of 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide hydrochloride




An intermediate (viii) was prepared in the same manner as described in Example 103, except that N,N,N'-trimethylethane-1,2-diamine was used instead of tert-butylmethyl(2-(methylamino)ethyl)carbamate (ii). To the obtained intermediate

5 intermediate 5-chloro-N-(2,4-dimethoxybenzyl)-4-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(5-fluoropyridine-2-yl)benzene sulfonamide (viii, 0.05 g, 0.08 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC.

10 The reaction solution was filtered to obtain the target compound (0.02 g, 48.3%).

¹H NMR (500 MHz, MeOD): 8.06(d, 1H), 7.88(d, 1H), 7.53(m, 1H), 7.25(dd, 1H), 7.16(dd, 1H), 7.04(m, 1H), 6.88(m, 1H), 6.35(d, 1H), 3.33(t, 2H), 3.20(t, 2H), 2.92(s, 6H), 2.71(s, 3H)

15 **Example 138: Preparation of 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide hydrochloride**



An intermediate was prepared in the same manner as described in


20 Example 137, except that thiazol-2-amine was used instead of 5-fluoropyridine-

2-amine (iv). To the obtained intermediate 5-chloro-N-(2,4-dimethoxybenzyl)-4-
 ((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-
 (thiazol-2-yl)benzene sulfonamide (0.05 g, 0.08 mmol) was added 1 M HCl in
 ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to
 5 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction
 solution was filtered to obtain the target compound (0.02 g, 48.4%).

¹H NMR (500 MHz, MeOD): 7.83(d, 1H), 7.27(m, 1H), 7.12(t, 1H),
 7.04(m, 1H), 6.90(t, 1H), 6.75(d, 1H), 6.40(d, 1H), 3.34(t, 2H), 3.22(t, 2H),
 2.83(s, 6H), 2.73(s, 3H)

10

**Example 139: Preparation of 5-chloro-N-(5-chlorothiazol-2-yl)-4-
 ((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-
 fluorobenzenesulfonamide hydrochloride**




15 An intermediate was prepared in the same manner as described in
 Example 137, except that 5-chlorothiazol-2-amine was used instead of 5-
 fluoropyridine-2-amine (iv). To the obtained intermediate 5-chloro-N-(5-
 chlorothiazol-2-yl)-N-(2,4-dimethoxybenzyl)-4-((2-((2-
 (dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluorobenzene
 20 sulfonamide (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL).
 After stirring the mixture stirred overnight while heating to 50 to 60°C, the
 completion of the reaction was confirmed by TLC. The reaction solution was
 filtered to obtain the target compound (0.02 g, 47.9%).

¹H NMR (500 MHz, MeOD): 7.82(d, 1H), 7.28(t, 1H), 7.19(d, 1H),
 25 7.05(d, 1H), 6.89(t, 1H), 6.41(d, 1H), 3.30(t, 2H), 3.23(t, 2H), 2.84(s, 6H), 2.74(s,
 3H)

Example 140: Preparation of 5-chloro-N-(5-chlorothiazol-2-yl)-4-


((2-((2-(dimethylamino)ethyl)(methyl)amino)phenyl)amino)-2-fluorobenzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in
 5 Example 137, except that 5-chlorothiazol-2-amine was used instead of 5-fluoropyridine-2-amine (iv) and 2-fluoro-1-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate 5-chloro-N-(5-chlorothiazol-2-yl)-N-(2,4-dimethoxybenzyl)-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)phenyl)amino)-2-fluorobenzene
 10 sulfonamide (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 48.2%).

1H NMR (500 MHz, MeOD): 7.84(d, 1H), 7.31(t, 2H), 7.25(t, 1H),
 15 7.20(t, 2H), 6.40(d, 1H), 3.36(t, 2H), 3.24(t, 2H), 2.86(s, 6H), 2.72(s, 3H)

Example 141: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylthiazol-2-yl)benzenesulfonamide hydrochloride



20


An intermediate was prepared in the same manner as described in
 Example 103, except that 5-methylthiazol-2-amine was used instead of 5-fluoropyridine-2-amine (iv). To the obtained intermediate tert-butyl 2-((2-((2-

chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-methylthiazol-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 53.0%).

¹H NMR (500 MHz, MeOD): 7.83(d, 1H), 7.29(dd, 1H), 7.05(dd, 1H), 6.89(t, 1H), 6.82(s, 1H), 6.47(d, 1H), 3.27(t, 2H), 2.11(t, 2H), 2.72(s, 3H), 2.67(s, 3H), 2.24(s, 3H)

10

Example 142: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(oxazol-2-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 133, except that oxazole-2-amine was used instead of 5-methylisoxazole-3-amine (iv). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(oxazol-2-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate (0.03 g, 0.05 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.01 g, 47.5%).

¹H NMR (500 MHz, MeOD): 7.87(d, 1H), 7.25-7.22(m, 2H), 7.01 (d, 1H), 6.85-6.84(m, 1H), 6.42(d, 1H), 5.48(s, 1H), 3.25-3.24(m, 2H), 3.09-3.07(m, 2H), 2.71(s, 3H), 2.61(s, 3H)

Example 143: Preparation of N-(5-(tert-butyl)isoxazol-3-yl)-5-


chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride



5 An intermediate was prepared in the same manner as described in Example 133, except that 5-(tert-butyl) isoxazole-3-amine was used instead of 5-methylisoxazole-3-amine (iv). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(5-tert-butyl)isoxazol-3-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)ethyl)(methyl)carbamate
 10 (0.03 g, 0.04 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.01 g, 46.4%).

15 ¹H NMR (500 MHz, MeOD): 7.78(d, 1H), 7.19-7.16(m, 1H), 6.97(d, 1H), 6.82-6.79(m, 1H), 6.42(d, 1H), 5.87(s, 1H), 3.30-3.29(m, 2H), 2.67(s, 3H), 2.64(s, 3H), 1.23(s, 9H)


Example 144: Preparation of N-(5-(tert-butyl)isoxazol-3-yl)-5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide hydrochloride
 20



An intermediate was prepared in the same manner as described in Example 133, except that 5-(tert-butyl)isoxazole-3-amine was used instead of 5-methylisoxazole-3-amine (iv) and 1-fluoro-2-nitrobenzene was used instead of 2,4-difluoro-1-nitrobenzene (i). To the obtained intermediate tert-butyl (2-((2-((4-
 5 (N-(tert-butoxycarbonyl)-N-(5-tert-butyl)isoxazol-3-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)phenyl)(methyl)amino)ethyl)(methyl)carbamate (0.03 g, 0.04 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.01 g, 46.0%).

¹H NMR (500 MHz, MeOD): 7.83(d, 1H), 7.23(d, 2H), 7.14-7.10(m, 2H), 6.70(d, 1H), 5.89(s, 1H), 3.30-3.27(m, 2H), 3.13-3.11(m, 2H), 2.66(s, 6H), 1.23(s, 9H)

Example 145: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1,2,4-thiadiazol-5-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 133, except that 1,2,4-thiadiazol-5-amine was used instead of 5-methylisoxazole-3-amine (iv) and tert-butyl methyl(pyrrolidin-3-yl)carbamate

was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(1,2,4-thiadiazol-5-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.0%).

¹H NMR (500 MHz, MeOD): 8.39(s, 1H), 7.80(d, 1H), 7.14(t, 1H), 6.73(d, 1H), 6.68(t, 1H), 6.08(d, 1H), 3.78(t, 1H), 3.58(dd, 1H), 3.48(m, 1H), 3.40(dd, 1H), 2.67(s, 3H), 2.34(m, 1H), 1.99(m, 1H)


Example 146: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1-methyl-1H-pyrazol-3-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 133, except that 1-methyl-1H-pyrazol-3-amine was used instead of 5-methylisoxazole-3-amine (iv) and tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(1-methyl-1H-pyrazol-3-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.1%).

¹H NMR (500 MHz, MeOD): 7.68(d, 1H), 7.38(s, 1H), 7.13(t, 1H), 6.75(dd, 1H), 6.86(t, 1H), 6.06(d, 1H), 6.00(s, 1H), 3.78(t, 1H), 3.70(s, 3H), 3.56(dd, 1H), 3.48(m, 1H), 3.41(dd, 1H), 3.39(s, 3H), 2.36(m, 1H), 1.98(m, 1H)


Example 147: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-4-yl)benzenesulfonamide hydrochloride



5 An intermediate was prepared in the same manner as described in Example 133, except that pyrimidin-4-amine was used instead of 5-methylisoxazole-3-amine (iv) and tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To
10 the obtained intermediate tert-butyl ((4-((2-(3-((tert-butoxycarbonyl)(methyl)amino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluorophenyl)sulfonyl)(pyrimidin-4-yl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC.
15 The reaction solution was filtered to obtain the target compound (0.02 g, 56.2%).

¹H NMR (500 MHz, MeOD): 8.45(s, 1H), 8.44(s, 1H), 7.95(d, 1H), 7.12(d, 1H), 7.00(t, 1H), 6.72(dd, 1H), 6.66(td, 1H), 6.04(d, 1H), 3.78(t, 1H), 3.56(dd, 1H), 3.47(m, 1H), 3.40(dd, 1H), 2.65(s, 3H), 2.31(m, 1H), 1.98(m, 1H)


20 **Example 148: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide hydrochloride**



An intermediate was prepared in the same manner as described in Example 96, except that tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((2-chloro-4-(N-(2,4-dimethoxybenzyl)-N-(5-fluoropyrimidin-2-yl)sulfamoyl)-5-fluorophenyl)amino) -5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 59.8%).

¹H NMR (500 MHz, MeOD): 8.46(s, 2H), 7.94(d, 1H), 7.11(td, 1H), 6.72(d, 1H), 6.66(td, 1H), 6.04(d, 1H), 3.77(t, 1H), 3.57(dd, 1H), 3.48(m, 1H), 3.41(dd, 1H), 2.66(s, 3H), 2.33(m, 1H), 1.99(m, 1H)


Example 149: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrazin-2-yl)benzenesulfonamide hydrochloride



An intermediate was prepared in the same manner as described in Example 96, except that 2-chloropyrazine was used instead of 2-chloro-5-fluoropyrimidine (iv) and tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((2-chloro-4-(N-(2,4-dimethoxyphenyl)-N-(pyrazin-2-yl)sulfamoyl)-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 60.9%).

¹H NMR (500 MHz, MeOD): 8.29(s, 1H), 8.15(s, 1H), 8.09(s, 1H), 7.88(d, 1H), 7.09(dd, 1H), 6.70(dd, 1H), 6.64(t, 1H), 6.05(d, 1H), 5.45(d, 1H), 3.76(m, 1H), 3.54(dd, 1H), 3.44(m, 2H), 2.67(s, 3H), 2.29(m, 1H), 1.90(m, 1H)


5 **Example 150: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide hydrochloride**



An intermediate was prepared in the same manner as described in
 10 Example 133, except that tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)-N-(5-methylisoxazol-3-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.06 mmol) was added 1 M HCl in ethyl acetate
 15 (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.1%).

¹H NMR (500 MHz, MeOD): 7.80(d, 1H), 7.14(dd, 1H), 6.73(d, 1H), 6.69(dd, 1H), 6.09(d, 1H), 6.07(s, 1H), 3.76(t, 1H), 3.58(m, 2H), 3.43(m, 2H),
 20 2.66(s, 3H), 2.32(s, 3H), 2.05(m, 2H)

Example 151: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-5-yl)benzenesulfonamide hydrochloride




An intermediate was prepared in the same manner as described in Example 133, except that pyrimidin-5-amine was used instead of 5-methylisoxazol-3-amine (iv) and tert-butyl methyl(pyrrolidin-3-yl)carbamate was used instead of tert-butyl methyl(pyrrolidin-3-yl)carbamate(ii). To the obtained intermediate tert-butyl (1-(2-((4-(N-(tert-butoxycarbonyl)(pyrimidin-5-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)pyrrolidin-3-yl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 56.2%).

¹H NMR (500 MHz, MeOD): 8.87(s, 1H), 8.59(d, 2H), 7.79(d, 1h), 7.12(td, 1H), 6.73(d, 1H), 6.67(td, 1H), 6.07(d, 1H), 3.76(t, 1H), 3.50(m, 2H), 3.54(m, 2H), 2.67(s, 3H), 2.32(m, 1H), 1.98(m, 1H)

15

Example 152: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-((2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



20


An intermediate was prepared in the same manner as described in Example 92, except that tert-butyl (2-aminoethyl)(methyl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate(ii). To the obtained intermediate tert-butyl (2-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-

yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)amino)ethyl)(methyl)carbamate (0.05 g, 0.07 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC.

5 The reaction solution was filtered to obtain the target compound (0.02 g, 56.9%).

¹H NMR (500 MHz, MeOD): 8.72(s, 1H), 7.75(d, 1H), 7.07(t, 1H), 7.00(s, 1H), 6.63(d, 1H), 6.49(t, 1H), 6.05(d, 1H), 3.46(t, 2H), 3.17(t, 2H), 2.70(s, 3H)

10 **Example 153: Preparation of 5-chloro-4-((2-((2-(dimethylamino)ethyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide hydrochloride**




An intermediate was prepared in the same manner as described in

15 Example 92, except that N,N-dimethylethane-1,2-diamine(3-(methylamino)propyl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate (ii). To the obtained intermediate tert-butyl (2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(2-(dimethylamino)ethyl)carbamate (0.02 g, 0.0003 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.01 g, 70.5%).

¹H NMR (500 MHz, MeOD): 8.73(s, 1H), 7.75(d, 1H), 7.08(t, 1H), 7.01(s, 1H), 6.66(d, 1H), 6.50(t, 1H), 6.04(d, 1H), 3.53(t, 2H), 3.30(t, 2H), 2.90(s, 6H)

25

Example 154: Preparation of 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(3-(methylamino)propyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide hydrochloride



5 An intermediate was prepared in the same manner as described in Example 92, except that tert-butyl methyl(3-(methylamino)propyl)carbamate was used instead of tert-butyl methyl(2-(methylamino)ethyl)carbamate (ii). To the obtained intermediate tert-butyl (3-((2-((4-(N-(tert-butoxycarbonyl)-N-(thiazol-4-yl)sulfamoyl)-2-chloro-5-fluorophenyl)amino)-5-fluorophenyl)(methyl)amino)propyl)(methyl)carbamate (0.05 g, 0.7 mmol) was added 1 M HCl in ethyl acetate (5 mL). After stirring the mixture stirred overnight while heating to 50 to 60°C, the completion of the reaction was confirmed by TLC. The reaction solution was filtered to obtain the target compound (0.02 g, 55.9%).

15 ¹H NMR (500 MHz, MeOD): 8.74(s, 1H), 7.78(d, 1h), 7.25(m, 1H), 7.03(m, 2H), 6.88(m, 1H), 6.29(d, 1H), 3.06(d, 2H), 2.80(d, 2H), 2.70(s, 3H), 2.60(s, 3H), 1.81(d, 2H)

Experimental Example

20 In order to measure the activities of the inventive compounds as antagonists, an experiment of blocking effect against sodium ion channel 1.7 (Nav1.7) and sodium ion channel 1.5 (Nav1.5) was carried out as follows.

1) Cell culture

25 The hNav 1.7 HEK 293 cell line was a cell line in which a human sodium ion channel 1.7 gene (type IX voltage-gated sodium channel alpha subunit) was stably expressed in human embryonic kidney (HEK) 293 cells, and was purchased from Millipore. The culture medium used was prepared by adding 1% 100X NEAA and 10% heat-inactivated FBS to DMEM F-12, and then

adding 1% P/S as an antibiotic thereto. G-418 as a restriction enzyme was added during subculture. The hNav1.7 HEK293 cells were cultured at a confluence of about 80% in a 5% CO₂ incubator at 37°C in a T75 flask for 2 or 3 days, and detached from the flask by treatment with 0.05% trypsin solution.
5 Then, the cells were collected by centrifugation and used in the experiment.

The hNav 1.5 HEK 293 cell line was a cell line in which a human sodium ion channel 1.5 gene (Homo sapiens sodium channel, voltage-gated, type V, alpha subunit (SCN 5 A)) was stably expressed in human embryonic
10 kidney (HEK) 293 cells, and was purchased from Creacell. The culture medium used was prepared by mixing 2% 100X L-glutamine and 10% heat-inactivated FBS to DMEM, and then adding 1% P/S as an antibiotic thereto. G-418 as a restriction enzyme was added during subculture, and the hNav1.5 HEK293 cells were cultured at a confluence of about 80% in a 5% CO₂ incubator at 37°C in a
15 T75 flask for 2 or 3 days, and detached from the flask by treatment with 0.05% trypsin solution. Then, the cells were collected by centrifugation and used in the experiment.

2) Preparation of compound samples

20 The compounds prepared in the Examples of the present invention were dissolved in dimethyl sulfoxide (DMSO) and used in the experiment. 90 mM and 10 mM DMSO stock solutions were prepared from each of the compounds and diluted in an extracellular solution (4 mM KCl, 138 mM NaCl, 1 mM MgCl₂, 1.8 mM CaCl₂, 5.6 mM Glucose, 10 mM HEPES, pH 7.45) at
25 various concentrations, so that the final concentration of DMSO was 0.3% or less.

3) Measurement of sodium ion channel blocking effects

In order to measure the sodium ion channel blocking effect, an
30 IonFlux16 Auto patch clamp system (Fluxion, Inc.) and a plate for exclusive use were used. The cells were distributed in an extracellular solution (4 mM KCl, 138 mM NaCl, 1 mM MgCl₂, 1.8 mM CaCl₂, 5.6 mM glucose, 10 mM HEPES,

pH 7.45), and then dispensed in the specified region of the plate, and each of the prepared compound samples was diluted at various concentrations, and then dispensed in the specified region of the plate. After the dispensation of the cells, the compound samples and an intracellular solution (100 mM CsF, 45 mM CsCl, 5 mM NaCl, 5 mM EGTA, 10 mM HEPES, pH 7.2) in the plate has been completed, the plate was mounted in the patch clamp system, and whether the compounds inhibited the ion channel was measured according to a set program.

Specifically, eight concentrations per compound were set, and percent inhibition was determined by calculating the percentage of inhibition of the peak current, generated after treating the cells with each concentration of the compound for 50 seconds, relative to the peak current generated before treatment with the compound, and the IC₅₀ value was calculated using the Sigma plot program. The results of the calculation are shown in Tables 2 to 5 below.

[Table 2]

Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)
1	0.035	21	0.066	41	> 1	61	> 1
2	0.026	22	> 1	42	> 1	62	> 1
3	0.103	23	0.357	43	> 1	63	> 10
4	0.044	24	0.049	44	> 1	64	1.174
5	0.120	25	n/a	45	0.236	65	0.541
6	0.182	26	0.066	46	0.686	66	0.160
7	0.125	27	> 1	47	0.375	67	0.028
8	0.489	28	> 0.3	48	0.086	68	0.103
9	1.505	29	0.362	49	0.609	69	0.241
10	0.396	30	> 1	50	0.328	70	0.050
11	1.007	31	0.044	51	n/a	71	0.435
12	0.412	32	0.017	52	0.061	72	0.046
13	0.372	33	> 1	53	> 1	73	0.100
14	0.447	34	0.241	54	> 1	74	0.278
15	0.369	35	0.205	55	0.012	75	0.126
16	0.108	36	0.051	56	> 1	76	0.149
17	0.053	37	0.058	57	> 1	77	0.032
18	> 1	38	0.027	58	> 1	78	0.212
19	> 1	39	0.053	59	> 1	79	0.216
20	0.021	40	0.013	60	0.073	80	0.028

[Table 3]

Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)	Example	Nav1.7 (IC ₅₀)
---------	----------------------------	---------	----------------------------	---------	----------------------------	---------	----------------------------

81	0.023	101	0.10	121	0.06	141	0.05
82		102	0.11	122	0.47	142	0.36
83		103	0.04	123	0.05	143	0.19
84		104	0.11	124	0.12	144	>10
85		105	0.02	125	0.08	145	1.40
86	0.277	106	0.23	126	0.02	146	0.39
87	0.140	107	0.46	127	0.008	147	0.25
88	1.929	108	0.02	128	0.009	148	0.11
89	1.216	109	0.05	129	0.004	149	0.30
90	0.352	110	0.07	130	0.006	150	0.36
91	6.581	111	0.01	131	0.027	151	>10
92	0.04	112	0.22	132	0.184	152	
93	0.35	113	0.59	133	0.67	153	
94	0.10	114	0.02	134	0.02	154	
95	0.03	115	0.07	135	0.24		
96	0.08	116	0.05	136	>10		
97	0.06	117	0.12	137	0.30		
98	0.04	118	0.20	138	0.24		
99	0.12	119	0.02	139	0.06		
100	1.09	120	0.14	140	3.67		

[Table 4]

Example	Nav1.5(IC ₅₀)	Example	Nav1.5(IC ₅₀)	Example	Nav1.5(IC ₅₀)
1	> 10	24	> 10	40	> 10
2	> 10	26	> 10	48	> 10
3	6.723	32	> 10	52	> 10
9	> 10	36	> 10	55	> 10
10	> 10	37	> 10	67	> 10
21	> 3	39	> 10		


[Table 5]

Example	Nav1.5(IC ₅₀)	Example	Nav1.5(IC ₅₀)	Example	Nav1.5(IC ₅₀)
92	25.03	101	> 10	128	1.83
95	8.2	102	> 10	148	> 10
96	> 10	103	> 10	149	> 10
97	> 10	104	> 10	150	> 10
98	> 10	105	> 10	151	> 10
99	> 10	106	> 10		
100	> 10	127	5.54		

We Claim:

1. A compound represented by Chemical Formula 1 below, or a pharmaceutically acceptable salt thereof:

5 [Chemical Formula 1]



in Chemical Formula 1,

R₁ is hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, halogen, or cyano,

10 R₂ and R₃ are each independently hydrogen, or halogen,

R₄ is C₅₋₁₀ heteroaryl containing one or two elements each independently selected from the group consisting of N, S and O,

wherein the C₅₋₁₀ heteroaryl is unsubstituted or substituted with C₁₋₄ alkyl, or halogen,

15 R₅ is -CH₂CH₂-N(R₇)(R₈), or -CH₂CH₂CH₂-N(R₇)(R₈), R₆ is hydrogen, or C₁₋₄ alkyl; or R₅ and R₆ together form C₃₋₅ alkylene, (C₂₋₄ alkylene)-N(R₉)-(C₂₋₄ alkylene), or (C₂₋₄ alkylene)-O-(C₂₋₄ alkylene),

wherein the C₃₋₅ alkylene, or C₂₋₄ alkylene is each independently unsubstituted or substituted with one or two R₁₀,

20 R₇, R₈, and R₉ are each independently hydrogen, or C₁₋₄ alkyl,

R₁₀ is C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, amino, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHCO(C₁₋₄ alkyl), or pyrrolidinyl,

X₁ is C-R', or N, wherein R' is hydrogen, or halogen,


X₂ is CH, or N, and

25 X₃ is N-R'', wherein R'' is hydrogen, or C₁₋₄ alkyl.

2. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

the compound represented by Chemical Formula 1 is represented by Chemical Formula 1' below:

[Chemical Formula 1']



5 wherein,

R₁ is hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ haloalkyl, C₁₋₄ haloalkoxy, halogen, or cyano,

R₂ and R₃ are each independently hydrogen, or halogen,

10 R₄ is C₅₋₁₀ heteroaryl containing one or two elements each independently selected from the group consisting of N, S and O,

R₁₁ is hydrogen, or C₁₋₄ alkyl,

R₁₂ is hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, halogen, amino, NH(C₁₋₄ alkyl), N(C₁₋₄ alkyl)₂, NHCO(C₁₋₄ alkyl), or pyrrolidinyl,

X₁ is C-R', or N, wherein R' is hydrogen, or halogen,

15 X₂ is CH, or N, and

X₃ is N-R'', wherein R'' is hydrogen, or C₁₋₄ alkyl,

X₄ is a bond, NH, N(C₁₋₄ alkyl), or O(oxygen), and

n is an integer of 1 to 4.

20 3. The compound or pharmaceutically acceptable salt thereof according to claim 1 or claim 2, wherein

R₁ is hydrogen, methyl, methoxy, trifluoromethyl, difluoromethoxy, trifluoromethoxy, fluoro, chloro, or cyano.

25 4. The compound or pharmaceutically acceptable salt thereof according to claim 1 or claim 2, wherein

R_2 and R_3 are each independently hydrogen, fluoro, or chloro.

5. The compound or pharmaceutically acceptable salt thereof according to claim 1 or claim 2, wherein


5 R_4 is thiazolyl, isoxazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridinyl, pyrimidinyl, or thiadiazolyl, and
the R_4 is unsubstituted or substituted with C_{1-4} alkyl, or halogen.

6. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

10 R_5 and R_6 together form C_{3-5} alkylene, $(C_{2-4}$ alkylene)- $N(R_9)$ -(C_{2-4} alkylene), or $(C_{2-4}$ alkylene)-O-(C_{2-4} alkylene), and
the C_{3-5} alkylene or C_{2-4} alkylene is each independently unsubstituted or substituted with methyl, methoxy, fluoro, amino, $NHCH_3$, $N(CH_3)_2$,
15 $N(CH_2CH_3)_2$, $NHCOCH_3$, or pyrrolidinyl.

7. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

20 R_5 and R_6 together with the nitrogen to which they are attached form any one selected from the group consisting of the following:



8. The compound or pharmaceutically acceptable salt thereof according to claim 1, wherein

25 R_5 is $-CH_2CH_2-NH-CH_3$, $-CH_2CH_2CH_2-NH-CH_3$, or $-CH_2CH_2-N(CH_3)_2$,

and

R₆ is hydrogen, or methyl.

9. The compound or pharmaceutically acceptable salt thereof according to
5 claim 1 or claim 2, wherein

X₁ is CH, CF, or N, X₂ is CH, or N, provided that both X₁ and X₂ are
not N.

10. The compound or pharmaceutically acceptable salt thereof according to
10 claim 1 or claim 2, wherein

X₃ is NH.

11. The compound or pharmaceutically acceptable salt thereof according to
claim 1, wherein

15 the compound is any one selected from the group consisting of:

- 1) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-
yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 2) 5-chloro-4-((4-chloro-2-(3-(methylamino)pyrrolidin-1-
yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20 3) (R)-5-chloro-4-((4-chloro-2-(2-methylpiperazin-1-
yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 4) (R)-5-chloro-4-((4-chloro-2-(3-(methylamino)piperidin-1-
yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 5) (R)-N-(1-(5-chloro-2-((2-chloro-5-fluoro-4-(N-(thiazol-4-
25 yl)sulfamoyl)phenyl)amino)phenyl)pyrrolidin-3-yl)acetamide,
- 6) 5-chloro-4-((4-chloro-2-(3-(diethylamino)pyrrolidin-1-
yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 7) 4-((2-([1,3'-bipyrrolidin]-1'-yl)-4-chlorophenyl)amino)-5-chloro-2-
fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 30 8) 5-chloro-2-fluoro-4-((4-methyl-2-(3-(methylamino)pyrrolidin-1-
yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 9) (S)-5-chloro-4-((4-chloro-2-(3-methylpiperazin-1-

- yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 10) 4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 11) 3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 12) 3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 13) 2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 14) 4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15 15) 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 16) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 17) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 18) 5-chloro-4-((5-chloro-3-(3-(methylamino)pyrrolidin-1-yl)pyridin-2-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20 19) 5-chloro-4-((6-chloro-2-(3-(methylamino)pyrrolidin-1-yl)pyridin-3-yl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 21) 5-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 25 22) (R)-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 23) (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30 24) (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25) (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-

- yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 26) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 27) (R)-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 28) (R)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 29) (R)-3-chloro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 30) (R)-3,5-difluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 31) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 32) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15 33) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 34) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-3-chloro-N-(thiazol-4-yl)benzenesulfonamide,
- 20 35) 4-((2-(3-aminopyrrolidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 36) 4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 37) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 38) 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 39) 4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 30 40) 3-chloro-4-((4-(difluoromethoxy)-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 41) (R)-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-

- (thiazol-4-yl)benzenesulfonamide,
- 42) (R)-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 43) (R)-3-chloro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 44) (R)-3,5-difluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 45) (R)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-fluoropyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 46) 2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 47) 3-chloro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 48) 5-chloro-2-fluoro-4-((3-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 15 49) 2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 50) 3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 20 51) 3,5-difluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 52) 5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 53) 2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 54) 3-chloro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 55) 5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30 56) (R)-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 57) (R)-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-

- yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 58) (R)-3-chloro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 59) (R)-3,5-difluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 5 60) (R)-5-chloro-2-fluoro-4-((4-methoxy-2-(3-(methylamino)piperidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 61) 3-chloro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 10 62) 5-chloro-2-fluoro-4-((4-fluoro-2-(4-methylpiperazin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 63) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 64) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 15 65) 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 66) 5-chloro-4-((4-(difluoromethoxy)-2-(4-methylpiperazin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 20 67) 5-chloro-2-fluoro-4-((2-(4-methylpiperazin-1-yl)-4-(trifluoromethoxy)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 68) (S)-3-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 69) (S)-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 25 70) (S)-5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 71) (S)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-methoxyphenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 30 72) (S)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,

- 73) (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 74) (R)-5-chloro-4-((2-(3-(dimethylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 5 75) (R)-5-chloro-4-((4-(difluoromethoxy)-2-(3-(dimethylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 76) 4-((2-(1,4-diazepan-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 10 77) 5-chloro-4-((4-cyano-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 78) (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide,
- 79) (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-(trifluoromethyl)phenyl)pyrrolidin-3-yl)acetamide,
- 15 80) (S)-3-chloro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 81) (S)-5-chloro-2-fluoro-4-((2-(3-(methylamino)pyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 20 82) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)azetidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 83) 4-((2-(3-aminoazetidin-1-yl)-4-fluorophenyl)amino)-5-chloro-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 84) 5-chloro-4-((2-(3-(dimethylamino)azetidin-1-yl)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 25 85) N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)azetidin-3-yl)acetamide,
- 86) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 30 87) 5-chloro-2-fluoro-4-((2-(3-methoxypyrrolidin-1-yl)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 88) (R)-N-(1-(2-((2-chloro-5-fluoro-4-(N-(thiazol-4-

yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide,
89) 3-chloro-4-((4-fluoro-2-(3-methoxypyrrolidin-1-yl)phenyl)amino)-
N-(thiazol-4-yl)benzenesulfonamide,
90) 3-chloro-4-((2-(3-methoxypyrrolidin-1-yl)-4-
5 (trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
91) (R)-N-(1-(2-((2-chloro-4-(N-(thiazol-4-
yl)sulfamoyl)phenyl)amino)-5-fluorophenyl)pyrrolidin-3-yl)acetamide,
92) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
10 93) 3-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-
fluorophenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
94) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-
fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
95) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-
15 (trifluoromethyl)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
96) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyrimidin-2-
yl)benzenesulfonamide,
97) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-
20 (methylamino)ethyl)amino)-4-
(trifluoromethyl)phenyl)amino)benzenesulfonamide,
98) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-
yl)benzenesulfonamide,
25 99) 5-chloro-4-((4-cyano-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-
yl)benzenesulfonamide,
100) 5-chloro-4-((4-cyano-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-
30 yl)benzenesulfonamide,
101) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-
(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-

yl)benzenesulfonamide,

102) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

103) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluoropyridin-2-yl)benzenesulfonamide,

104) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,

105) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,

106) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,

107) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide,

108) 5-chloro-4-((4-cyano-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,

109) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)benzenesulfonamide,

110) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,

111) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)-4-(trifluoromethyl)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,

112) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

113) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(pyridin-2-yl)benzenesulfonamide,

114) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,

- 115) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,
- 116) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,
- 117) 5-chloro-4-((4-(difluoromethoxy)-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(pyridin-2-yl)benzenesulfonamide,
- 118) 5-chloro-2-fluoro-N-(5-fluoropyridin-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 119) 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-2-yl)benzenesulfonamide,
- 120) 5-chloro-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 121) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,
- 122) 5-chloro-2-fluoro-N-(5-fluoropyrimidin-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
- 123) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide,
- 124) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,
- 125) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,
- 126) 5-chloro-4-((4-chloro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,
- 127) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-fluoro-2-

(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
 128) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-fluorothiazol-2-yl)benzenesulfonamide,
 5 129) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
 130) 5-chloro-N-(5-chlorothiazol-2-yl)-2-fluoro-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
 131) 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
 10 132) 5-chloro-2-fluoro-N-(5-fluorothiazol-2-yl)-4-((4-methoxy-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,
 133) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,
 15 134) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,
 135) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide,
 20 136) 5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methyl-1H-pyrazol-3-yl)benzenesulfonamide,
 137) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(5-fluoropyridin-2-yl)benzenesulfonamide,
 138) 5-chloro-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-2-yl)benzenesulfonamide,
 139) 5-chloro-N-(5-chlorothiazol-2-yl)-4-((2-((2-(dimethylamino)ethyl)(methyl)amino)-4-fluorophenyl)amino)-2-fluorobenzenesulfonamide,
 30 140) 5-chloro-N-(5-chlorothiazol-2-yl)-4-((2-((2-

(dimethylamino)ethyl)(methyl)amino)phenyl)amino)-2-fluorobenzenesulfonamide,

141) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(5-methylthiazol-2-yl)benzenesulfonamide,

142) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)-N-(oxazol-2-yl)benzenesulfonamide,

143) N-(5-(tert-butyl)isoxazol-3-yl)-5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

144) N-(5-(tert-butyl)isoxazol-3-yl)-5-chloro-2-fluoro-4-((2-(methyl(2-(methylamino)ethyl)amino)phenyl)amino)benzenesulfonamide,

145) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1,2,4-thiadiazol-5-yl)benzenesulfonamide,

146) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(1-methyl-1H-pyrazol-3-yl)benzenesulfonamide,

147) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-4-yl)benzenesulfonamide,

148) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-fluoropyrimidin-2-yl)benzenesulfonamide,

149) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrazin-2-yl)benzenesulfonamide,

150) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(5-methylisoxazol-3-yl)benzenesulfonamide,

151) 5-chloro-2-fluoro-4-((4-fluoro-2-(3-(methylamino)pyrrolidin-1-yl)phenyl)amino)-N-(pyrimidin-5-yl)benzenesulfonamide,

152) 5-chloro-2-fluoro-4-((4-fluoro-2-((2-(methylamino)ethyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide,

153) 5-chloro-4-((2-((2-(dimethylamino)ethyl)amino)-4-fluorophenyl)amino)-2-fluoro-N-(thiazol-4-yl)benzenesulfonamide, and

154) 5-chloro-2-fluoro-4-((4-fluoro-2-(methyl(3-(methylamino)propyl)amino)phenyl)amino)-N-(thiazol-4-yl)benzenesulfonamide.

12. A pharmaceutical composition for preventing or treating sodium channel blocker-related diseases comprising the compound or pharmaceutically acceptable salt thereof according to any one of claims 1 to 11 as an active ingredient.

5

13. The pharmaceutical composition according to claim 12, wherein the sodium channel blocker-related diseases are acute pain, chronic pain, neuropathic pain, postoperative pain, migraine, arthralgia, neuropathy, nerve damage, diabetic neuropathy, neuropathic disease, epilepsy, arrhythmia, myotonia, ataxia, multiple sclerosis, irritable bowel syndrome, urinary incontinence, visceral pain, depression, erythralgia, or PEPD (paroxysmal extreme pain disorder).

10

