

A&A Ref: 149659

PUBLICATION PARTICULARS AND ABSTRACT
(Section 32(3)(a) - Regulations 22(1)(g) and 31)

21	01	PATENT APPLICATION NO	22	LODGING DATE	43	ACCEPTANCE DATE
----	----	-----------------------	----	--------------	----	-----------------

2004/1221

16 February 2004

27.10.04

51	INTERNATIONAL CLASSIFICATION	NOT FOR PUBLICATION
----	------------------------------	---------------------

A61K, C07D

CLASSIFIED BY: ISA

71	FULL NAME(S) OF APPLICANT(S)
----	------------------------------

Aventis Pharma Deutschland GmbH

72	FULL NAME(S) OF INVENTOR(S)
----	-----------------------------

SCHWINK, Lothar
GOSSEL, Matthias
ROSSE, Gerard

STENGELIN, Siegfried
WALSER, Armin

EARLIEST PRIORITY CLAIMED	COUNTRY	NUMBER	DATE
	33 DE	31 101 39 416.0	32 17 August 2001

NOTE: The country must be indicated by its International Abbreviation - see schedule 4 of the Regulations

54	TITLE OF INVENTION
----	--------------------

Aminoalkyl-substituted aromatic bicyclic compounds, method for the production thereof and their use
as medicaments

57	ABSTRACT (NOT MORE THAN 150 WORDS)
----	------------------------------------

NUMBER OF SHEETS	79
------------------	----

The sheet(s) containing the abstract is/are attached.

If no classification is furnished, Form P.9 should accompany this form.
The figure of the drawing to which the abstract refers is attached.



(67) Abstract: The invention relates to aminoalkyl-substituted aromatic bicyclic compounds and to their physiologically acceptable salts and physiologically functional derivatives. Disclosed are compounds of formula (I), wherein the radicals have the meaning cited in the description. Further disclosed are the physiologically acceptable salts of said compounds and to a method for the production thereof. Said compounds can be suitably used as anorectic drugs.

AS ORIGINALLY FILED

Aminoalkyl-substituted aromatic bicyclic compounds, methods for their preparation and
their use as pharmaceuticals

5

The invention relates to aminoalkyl-substituted aromatic bicyclic compounds and to the physiologically acceptable salts and physiologically functional derivatives thereof.


10

Structurally similar nonaromatic bicyclic compounds with pharmacological action have already been described in the prior art (for example in WO 01/21577).

It was the object of the present invention to provide compounds which cause a reduction in weight in mammals and which are suitable for preventing and treating obesity.

15

The invention therefore relates to compounds of the formula I,



20

in which

A

is (C₁-C₈)alkyl, (C₀-C₈)alkylenearyl;

a 3- to 12-membered mono- or bicyclic ring which may contain one or more heteroatoms from the group consisting of N, O and S and the 3- to 12-

25

membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃, OCF₃, CN, (C₁-C₆)alkyl, aryl, CON(R₃₇)(R₃₈), N(R₃₉)(R₄₀), OH, O-(C₁-C₆)alkyl, S-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

X is a bond, C(R8)(R9), C(OR10)(R11), O, N(R12), S, SO, SO₂, CO;

R8, R9, R10, R11, R12 are independently of one another H, (C₁-C₆)alkyl;

5 D is N, C(R41);

E is N, C(R42);

G is N, C(R43);

10

L is N, C(R44);

R1, R2, R3, R41, R42, R43, R44 are independently of one another

15

H, F, Cl, Br, J, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)alkyl, (C₁-C₄)-alkoxyalkyl, S-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₃-C₈)-cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkenyl, O-(C₃-C₈)-cycloalkenyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₈)alkylenearyl, S-aryl, N(R13)(R14), SO₂-CH₃, COOH, COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21),
20 a 5- to 7-membered heterocycle having 1-4 heteroatoms;

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

25

or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R15, R16 are independently of one another H, (C₁-C₆)alkyl,

30

or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R17, R19 are independently of one another H, (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24), O;

5

R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

10 W is N, C(R25);

R25 is H, (C₁-C₆)alkyl, aryl, a bond to Y;

T is N, C(R26);

15

R26 is H, (C₁-C₆)alkyl, aryl, (C₀-C₈)alkylenearyl, a bond to Y;

U is O, S, N(R27), -C(R30)=N-, -N=C(R31)-;

20 R27, R30, R31 are independently of one another H, (C₁-C₆)alkyl, a bond to Y;

Y is (C₁-C₈)alkylene, in which one or more carbons may be replaced by O, S, SO, SO₂, C(R32)(R33), CO, C(R34)(OR35) or N(R36);

25 R32, R33, R34, R35, R36 are independently of one another H, (C₁-C₆)alkyl, aryl;

R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form

30

a 3- to 8-membered ring in which one or more carbons may be replaced by O, N or S and the 3- to 8-membered ring may carry further substituents such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH, O-(C₁-C₆)alkyl or NHCO(C₁-C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

5

Preference is given to compounds of the formula I, in which one or more radicals have the following meaning:

A is (C₂-C₇)alkyl, (C₀-C₃)alkylenearyl;

10

a 4- to 10-membered mono- or bicyclic ring which may contain one or more heteroatoms from the group consisting of N, O and S, and the 4- to 10-membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), O-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

15

X is a bond, C(R8)(R9), O, N(R12), S, SO₂;

R8, R9, R12 are independently of one another H, (C₁-C₆)alkyl;

20 D

is N, C(R41);

E

is N, C(R42);

G

is N, C(R43);

25

L

is N, C(R44);

where the total number of the nitrogen atoms defined by D, E, G and L is 0, 1 or 2;

30

R1, R2, R3, R41, R42, R43, R44 are independently of one another

H, F, Cl, Br, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₃)-

alkylenearyl, S-aryl, N(R13)(R14), SO₂-CH₃, COO-(C₁-C₆)alkyl,
CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21);

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

5

or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

10 R15, R16 are independently of one another H, (C₁-C₆)alkyl,

or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

15 R17, R19 are independently of one another H, (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24), O;

20

R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

25 W is N, C(R25);

R25 is H, (C₁-C₆)alkyl, aryl;

T is C(R26);

30 R26 is H, (C₁-C₆)alkyl, aryl, a bond to Y;

U is O, S, N(R27), -N=C(R31)-;

R27, R31 are independently of one another H, (C₁-C₆)alkyl, a bond to Y;

Y is (C₁-C₄)alkylene, in which a carbon may be replaced by SO₂, C(R32)(R33), CO or N(R36);

5

R32, R33, R36 are independently of one another H, (C₁-C₆)alkyl, aryl;

R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

10

or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form a 4- to 7-membered ring in which one or more carbons may be replaced by O, N or S and the 4- to 7-membered ring may carry further substituents such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH or NHCO(C₁-C₆)alkyl;

15

R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

20

Particular preference is given to compounds of the formula I, in which one or more radicals have the following meaning:

A is (C₃-C₇)alkyl, (C₀-C₂)alkylenearyl;

25

a 5- to 10-membered mono- or bicyclic ring which may contain 0, 1 or 2 heteroatoms from the group consisting of N, O and S, and the 5- to 10-membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, O-(C₁-C₆)alkyl or NHCO(C₁-C₆)alkyl;

30

X is a bond, C(R8)(R9), O, N(R12);

R8, R9, R12 are independently of one another H, (C₁-C₆)alkyl;

D is N, C(R41);

E is N, C(R42);

5 G is N, C(R43);

L is N, C(R44);

where the total number of the nitrogen atoms defined by D, E, G and L is 0 or 1;

10

R1, R2, R3, R41, R42, R43, R44 are independently of one another

H, F, Cl, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, O-(C₃-C₈)cycloalkyl, (C₀-C₂)alkylenearyl, -O-(C₀-C₃)alkylenearyl, N(R13)(R14), COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21);

15

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

R15, R16 are independently of one another H, (C₁-C₆)alkyl,

20 R17, R19 are independently of one another H, (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24);

25

R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

30 W is N, C(R25);

R25 is H, (C₁-C₆)alkyl;

T is C(R26);

R26 is H, (C₁-C₆)alkyl, a bond to Y;

5 U is O, S, N(R27);

R27 is H, (C₁-C₆)alkyl, a bond to Y;

10 Y is (C₁-C₃)alkylene, in which a carbon may be replaced by SO₂, C(R32)(R33) or CO;

R32, R33 are independently of one another H, (C₁-C₆)alkyl, aryl;

15 R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

or R6 and Y or R6 and R7 together with the nitrogen to which they are bonded form a 5- to
or 6-membered ring in which one or more carbons may be replaced by O or
N and the 5- or 6-membered ring may carry further substituents such as (C₁-
20 C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH or NHCO(C₁-C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

25 The invention relates to compounds of the formula I in the form of their racemates, enantiomer-enriched mixtures and pure enantiomers and to their diastereomers and mixtures thereof.

The substituents R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, R12, R13, R14, R15,
30 R16, R17, R18, R19, R20, R21, R22, R25, R26, R27, R30, R31, R32, R33, R34, R35, R36, R37, R38, R39, R40, R41, R42, R43 and R44 may have straight-chain, branched or optionally halogenated alkyl, alkylene, alkenyl and alkynyl radicals.

The term "aryl" means a phenyl or naphthyl group. The term "ring" means a cyclic structure which may be aromatic, partly saturated or completely saturated.

The optional ring formation of R⁶, Y and the nitrogen to which they are bonded can be illustrated by examples 6 and 16 without limiting the general description mentioned above.

5

Pharmaceutically acceptable salts are particularly suitable for medical applications, due to their greater solubility in water compared with the starting or base compounds. Said salts must have a pharmaceutically acceptable anion or cation. Suitable pharmaceutically acceptable acid addition salts of the compounds of the invention are salts of inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, metaphosphoric acid, nitric acid, sulfonic acid and sulfuric acid and also of organic acids such as, for example, acetic acid, benzenesulfonic acid, benzoic acid, citric acid, ethanesulfonic acid, fumaric acid, gluconic acid, glycolic acid, isethionic acid, lactic acid, lactobionic acid, maleic acid, malic acid, methanesulfonic acid, succinic acid, p-toluenesulfonic acid, tartaric acid and trifluoroacetic acid. For medicinal purposes, particular preference is given to using the chlorine salt. Suitable pharmaceutically acceptable basic salts are ammonium salts, alkali metal salts (such as sodium salts and potassium salts) and alkaline earth metal salts (such as magnesium salts and calcium salts).

10

15

20

Salts having a pharmaceutically unacceptable anion are likewise included within the scope of the present invention as useful intermediates for preparing or purifying pharmaceutically acceptable salts and/or for use in nontherapeutic applications, for example in-vitro applications.

25

The term "physiologically functional derivative" used herein relates to any physiologically acceptable derivative of an inventive compound of the formula I, for example an ester which on administration to a mammal, for example humans, is capable of forming (directly or indirectly) a compound of the formula I or an active metabolite thereof.

30

The physiologically functional derivatives also include prodrugs of the compounds of the invention. Such prodrugs may be metabolized in vivo to a compound of the invention. These prodrugs may or may not be active themselves.

The compounds of the invention may also be present in various polymorphous forms, for example as amorphous and crystalline polymorphous forms. All polymorphous forms of the compounds of the invention are included within the scope of the invention and are another aspect of the invention.

5

All references to "compound(s) according to formula (I)" refer hereinbelow to a compound/compounds of the formula (I) as described above and also to their salts, solvates and physiologically functional derivatives as described herein.

- 10 The amount of a compound according to formula (I) which is required in order to attain the desired biological effect depends on a number of factors, for example the specific compound selected, the intended use, the type of administration and the clinical state of the patient. In general, the daily dose is in the range from 0.3 mg to 100 mg (typically from 3 mg to 50 mg) per day per kilogram of body weight, for example 3-10 mg/kg/day. An
- 15 intravenous dose can be, for example, in the range from 0.3 mg to 1.0 mg/kg and can be administered in a suitable manner as an infusion of 10 ng to 100 ng per kilogram per minute. Suitable infusion solutions for these purposes may contain, for example, from 0.1 ng to 10 mg, typically from 1 ng to 10 mg per milliliter. Individual doses may contain, for example, from 1 mg to 10 g of the active compound. Thus, ampoules for injections can
- 20 contain, for example, from 1 mg to 100 mg, and orally administerable individual dose formulations such as, for example, tablets or capsules can contain, for example, from 1.0 to 1000 mg, typically from 10 to 600 mg. In the case of pharmaceutically acceptable salts, the abovementioned masses relate to the mass of the free compound on which the salt is based. The compound used for the prophylaxis or therapy of the abovementioned conditions may
- 25 be the compounds according to formula (I) themselves, but they are preferably present in the form of a pharmaceutical composition together with an acceptable carrier. The carrier must be naturally acceptable, in the sense that it is compatible with the other ingredients of said composition and is not harmful to the patient's health. The carrier may be a solid or a liquid or both and is preferably formulated with the compound as an individual dose, for
- 30 example as a tablet which may contain from 0.05% to 95% by weight of the active compound. Further pharmaceutically active substances may also be present, including further compounds according to formula (I). The pharmaceutical compositions of the invention may be prepared according to any of the known pharmaceutical methods which

essentially comprise mixing the ingredients with pharmacologically acceptable carriers and/or excipients.

Pharmaceutical compositions of the invention are those which are suitable for oral, rectal, topical, peroral (e.g. sublingual) and parenteral (e.g. subcutaneous, intramuscular, intradermal or intravenous) administration, although the most suitable manner of administration depends in each individual case on the nature and severity of the condition to be treated and on the nature of the compound according to formula (I) used in each case. Sugar-coated formulations and sugar-coated delayed-release formulations, too, are included within the scope of the invention. Preference is given to acid-resistant and enteric formulations. Suitable enteric coatings include cellulose acetate phthalate, polyvinyl acetate phthalate, hydroxypropylmethylcellulose phthalate and anionic polymers of methacrylic acid and methyl methacrylate.

Suitable pharmaceutical compounds for oral administration may be present in separate units as, for example, capsules, cachets, lozenges or tablets, which in each case contain a particular amount of the compound according to formula (I); as powders or granules; as solution or suspension in an aqueous or nonaqueous liquid; or as an oil-in-water or water-in-oil emulsion. As already mentioned, said compositions can be prepared according to any suitable pharmaceutical method which includes a step in which the active compound and the carrier (which may comprise one or more additional components) are contacted. In general, the compositions are prepared by uniform and homogeneous mixing of the active compound with a liquid and/or finely dispersed solid carrier, after which the product is shaped, if necessary. Thus a tablet, for example, may be prepared by pressing or shaping a powder or granules of the compound, where appropriate with one or more additional components. Pressed tablets can be prepared by tableting the compound in free-flowing form, for example a powder or granules, mixed, where appropriate, with a binder, lubricant, inert diluent and/or one or more surface active/dispersing agents in a suitable machine. Shaped tablets can be prepared by shaping the pulverulent compound, moistened with an inert liquid diluent, in a suitable machine.

Pharmaceutical compositions which are suitable for peroral (sublingual) administration include lozenges which contain a compound according to formula (I) with a flavoring,

usually sucrose and gum arabic or tragacanth, and pastilles which comprise the compound in an inert base such as gelatin and glycerol or sucrose and gum arabic.

Suitable pharmaceutical compositions for parenteral administration preferably comprise sterile aqueous preparations of a compound according to formula (I) which are preferably isotonic with the blood of the intended recipient. These preparations are preferably administered intravenously, although they may also be administered subcutaneously, intramuscularly or intradermally as an injection. Said preparations may preferably be prepared by mixing the compound with water and rendering the obtained solution sterile and isotonic with the blood. Injectable compositions of the invention generally contain from 0.1 to 5% by weight of the active compound.

Suitable pharmaceutical compositions for rectal administration are preferably present as individual dose suppositories. These may be prepared by mixing a compound according to formula (I) with one or more conventional solid carriers, for example cocoa butter, and shaping the resulting mixture.

Suitable pharmaceutical compositions for topical application to the skin are preferably present as ointment, cream, lotion, paste, spray, aerosol or oil. Carriers which may be used are petroleum jelly, lanolin, polyethylene glycols, alcohols and combinations of two or more of these substances. In general, the active compound is present at a concentration of from 0.1 to 15%, for example from 0.5 to 2%, by weight of the composition.

Transdermal administration is also possible. Suitable pharmaceutical compositions for transdermal administration may be present as individual patches which are suitable for long-term close contact with the epidermis of the patient. Such patches suitably contain the active compound in an optionally buffered aqueous solution, dissolved and/or dispersed in an adhesive or dispersed in a polymer. A suitable active compound concentration is from approx. 1% to 35%, preferably approx. 3% to 15%. A particular possibility is the release of the active compound by electrotransport or iontophoresis, as described, for example, in Pharmaceutical Research, 2(6): 318 (1986).

The compounds of the formula I are distinguished by beneficial actions on the metabolism of lipids, and they are particularly suitable for weight reduction and, after weight reduction, for maintaining a reduced weight in mammals and as anorectic agents. The compounds are distinguished by their low toxicity and their few side effects. The compounds may be employed alone or in combination with other weight-reducing or anorectic active compounds. Further anorectic active compounds of this kind are mentioned, for example, in the Rote Liste, Chapter 01 under weight-reducing agents/appetite suppressants, and may also include those active compounds which increase the energy turnover of the organism and thus lead to weight reduction or else those which influence the general metabolism of said organism such that increased calorie intake does not cause an enlargement of the fat depots and a normal calorie intake causes a reduction in the fat depots of said organism. The compounds are suitable for the prophylaxis and, in particular, for the treatment of problems of excess weight or obesity. The compounds are furthermore suitable for the prophylaxis and, in particular, for the treatment of type II diabetes, of arteriosclerosis and for the normalization of lipid metabolism and for the treatment of high blood pressure. The compounds act as MCH antagonists and are also suitable for the treatment of paresthesia and other psychiatric indications such as, for example, depressions, anxieties, anxiety neuroses, schizophrenia and also for the treatment of disorders associated with the circadian rhythm and for the treatment of drug abuse.

In a further aspect of the invention, the compounds of the formula I may be administered in combination with one or more further pharmacologically active substances which may be selected, for example, from the group consisting of antidiabetics, antiadipose agents, blood-pressure-lowering active compounds, lipid reducers and active compounds for the treatment and/or prevention of complications caused by diabetes or associated with diabetes.

Suitable antidiabetics include insulins, amylin, GLP-1 and GLP-2 derivatives such as, for example, those disclosed by Novo Nordisk A/S in WO 98/08871 and also oral hypoglycemic active compounds.

Said oral hypoglycemic active compounds preferably include sulfonyl ureas, biguanidines, meglitinides, oxadiazolidinediones, thiazolidinediones, glucosidase inhibitors, glucagon

receptor antagonists, GLP-1 agonists, potassium channel openers such as, for example, those disclosed by Novo Nordisk A/S in WO 97/26265 and WO 99/03861, insulin sensitizers, activators of insulin receptor kinase, inhibitors of liver enzymes involved in the stimulation of gluconeogenesis and/or glycogenolysis, for example glycogen phosphorase
 5 inhibitors, modulators of glucose uptake and glucose elimination, lipid metabolism-modifying compounds such as antihyperlipidemic active compounds and antilipidemic active compounds, for example HMGCoA-reductase inhibitors, inhibitors of cholesterol transport/cholesterol uptake, inhibitors of the reabsorption of bile acid or inhibitors of microsomal triglyceride transfer protein (MTP), compounds which reduce food intake,
 10 PPAR and RXR agonists and active compounds which act on the ATP-dependent potassium channel of beta cells.

In one embodiment of the present invention, the present compounds are administered in combination with insulin.

In another embodiment, the compounds of the invention are administered in combination
 15 with a sulfonylurea such as, for example, tolbutamide, glibenclamide, glimepiride, glipizide, gliquidone, glisoxepide, glibornuride or gliclazide.

In another embodiment, the compounds of the present invention are administered in combination with a biguanidine such as, for example, metformin.

In another embodiment, the compounds of the present invention are administered in
 20 combination with a meglitinide such as, for example, repaglinide.

In yet another embodiment, the compounds of the present invention are administered in combination with a thiazolidinedione such as, for example, troglitazone, ciglitazone, pioglitazone, rosiglitazone or the compounds disclosed by Dr. Reddy's Research Foundation in WO 97/41097, in particular 5-[[4-[(3,4-dihydro-3-methyl-4-oxo-2-quinazolinylmethoxy)]-
 25 phenyl]methyl]-2,4-thiazolidinedione.

In another embodiment, the compounds of the present invention are administered in combination with an α -glucosidase inhibitor such as, for example, miglitol or acarbose.

In another embodiment, the compounds of the present invention are administered in combination with an active compound which acts on the ATP-dependent potassium channel of the beta cells, such as, for example, tolbutamide, glibenclamide, glimepiride, glipizide, gliclazide or repaglinide.

- 5 In yet another embodiment, the compounds of the present invention are administered in combination with an antihyperlipidemic active compound or an antilipidemic active compound such as, for example, cholestyramine, colestipol, clofibrate, fenofibrate, gemfibrozil, lovastatin, pravastatin, simvastatin, atorvastatin, cerivastatin, fluvastatin, probucol, ezetimibe or dextrothyroxine.
- 10 In another embodiment, the compounds of the present invention are administered in combination with more than one of the aforementioned compounds, for example in combination with a sulfonylurea and metformin, a sulfonylurea and acarbose, repaglinide and metformin, insulin and a sulfonylurea, insulin and metformin, insulin and troglitazone, insulin and lovastatin, etc.
- 15 Furthermore, the compounds of the invention may be administered in combination with one or more antiadipose agents or appetite-controlling active compounds.

Such active compounds may be selected from the group consisting of CART agonists, NPY antagonists, MC4 agonists, orexin antagonists, H3 agonists, TNF agonists, CRF agonists, CRF BP antagonists, urocortin agonists, $\beta 3$ agonists, MSH (melanocyte-stimulating hormone) agonists, CCK agonists, serotonin re-uptake inhibitors, mixed serotonin and noradrenalin reuptake inhibitors, 5HT modulators, MAO inhibitors, bombesin agonists, galanin antagonists, growth hormone, growth-hormone-releasing compounds, TRH agonists, uncoupling protein 2 or 3 modulators, leptin agonists, dopamine agonists (bromocriptine, doprexin), lipase/amylase inhibitors, cannabinoid
20 receptor 1 antagonists, modulators of acylation-stimulating protein (ASP), PPAR modulators, RXR modulators, hCNTF mimetics or TR- β agonists.

In one embodiment of the invention, the antiadipose agent is leptin or modified leptin.

In another embodiment, the antiadipose agent is dexamphetamine or amphetamine.

In another embodiment, the antiadipose agent is fenfluramine or dexfenfluramine.

In yet another embodiment, the antiadipose agent is sibutramine or the mono- and bis-demethylated active metabolite of sibutramine.

In another embodiment, the antiadipose agent is orlistate.

- 5 In another embodiment, the antiadipose agent is mazindol, diethylpropione or phentermine.

Furthermore, the compounds of the present invention may be administered in combination with one or more antihypertensive active compounds. Examples of antihypertensive active compounds are betablockers such as alprenolol, atenol, timolol, pindolol, propanolol and metoprolol, ACE (angiotensin-converting enzyme) inhibitors such as, for example,
 10 benazepril, captopril, enalapril, fosinopril, lisinopril, quinapril and rampril, calcium channel blockers such as nifedipine, felodipine, nicardipine, isradipine, nimodipine, diltiazem and verapamil, and also alphablockers such as doxazosin, urapidil, prazosin and terazosin. Furthermore, reference may be made to Remington: The Science and Practice of Pharmacy, 19th edition, Gennaro, editor, Mack Publishing Co., Easton, PA, 1995.

- 15 It is self-evident that every suitable combination of the compounds of the invention with one or more of the aforementioned compounds and optionally one or more other pharmacologically active substances is to be regarded as covered by the scope of protection of the present invention.

The activity of the compounds was assayed as follows:

- 20 Biological test model:

The anorectic action was tested on female NMRI mice. After removal of feed for 17 hours, the preparation to be tested was administered by gavage. The animals were housed singly and, with free access to drinking water, they were offered evaporated milk 30 minutes after administration of the preparation. The consumption of evaporated milk was determined
 25 and the general behavior of the animals were monitored every half an hour for 7 hours. The measured milk consumption was compared to that of vehicle-treated control animals.

Table 1: Anorectic action, measured as a reduction in the cumulative milk consumption by treated animals compared with control animals

Example	Oral dose [mg/kg]	Number of animals/ cumulative milk consumption by treated animals N / [ml]	Number of animals/ cumulative milk consumption by control animals N / [ml]	Reduction in cumulative milk consumption as % of the control
Example 1	30	5/2.28	5/3.26	30
Example 4	10	5/2.74	5/4.44	38

The table indicates that the compounds of the formula I exhibit very good anorectic action.

In two simultaneously published articles in Nature (Nature 400, 261-264, 1999; Nature
 5 400, 265-269, 1999, see enclosure), two groups separately described a highly specific
 receptor for melanin-concentrating hormone (MCH). MCH takes over important functions
 in the control of food intake. Compounds acting on the MCH receptor therefore have
 anorectic action and are suitable for the treatment of obesity. The test for anorectic action
 of the inventive compounds of the formula I was therefore carried out as follows.

Functional measurements for determination of IC₅₀

Cloning of the cDNA for human MCH receptor, preparation of a recombinant HEK293
 cell line expressing human MCH receptor and functional measurements with said
 15 recombinant cell line were carried out according to the description by Audinot et al. (J.
 Biol. Chem. 276, 13554-13562, 2001). In contrast to the reference, however, plasmid
 pEAK8 from EDGE Biosystems (USA) was used for constructing the expression vector. A
 transformed HEK cell line named "PEAK Stable Cells" (likewise from EDGE Biosystems)
 served as host for transfection. The functional measurements of cellular calcium flow, after
 20 addition of agonists (MCH), in the presence of the ligand of the invention was carried out

with the aid of the FLIPR instrument from Molecular Devices (USA), using the manufacturer's protocols.

Table 2 shows the results from the cellular assay.

Table 2:

Example	IC50 / μM
1	0.15
2	0.15
3	0.29
4	0.13
5	0.50
6	2.34
7	0.45
8	1.90
9	0.10
10	0.11
11	0.14
13	2.50
14	0.30
15	0.18
16	0.33
17	2.14


18	1.04
19	0.70
22	4.42
24	0.86
26	0.92
29	2.91
33	1.24
63	0.57
65	0.50
71	2.65
72	0.32
73	0.14
76	4.25
77	0.70
78	2.75
79	2.13
80	3.36
81	2.69
84	0.40
86	2.78

105	1.0
106	0.20
107	1.0
108	0.43
109	1.29

The examples and preparation methods listed below serve to illustrate the invention but without limiting it.

Example 1


5 1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



Carbonyldiimidazole (5.12 g) was added to a solution cooled to 0°C of 1-dimethyl-aminoethyl-5-aminoindole (6.30 g) in dimethylformamide (50 ml). After 10 minutes, 4-aminodiphenyl ether (5.84 g) was added and the reaction mixture was heated to 80°C for 2
 10 hours. After cooling, the reaction was diluted with ethyl acetate and washed with water. The organic phase was dried over magnesium sulfate, filtered and concentrated. The residue was purified by chromatography on silica gel (eluent: dichloromethane/methanol 9:1). Thus the product having a molecular weight of 414.15 (C₂₅H₂₆N₄O₂); MS (ESI): 415
 15 (M+H⁺) was obtained.

Example 2


1-(4-Butoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea



The compound was prepared from 4-butoxyaniline and 1-dimethylaminoethyl-5-aminoindole, as described in Example 1. Thus the product having a molecular weight of
 5 394.52 ($C_{23}H_{30}N_4O_2$); MS (ESI): 395 ($M+H^+$) was obtained.

Example 3

1-(1-Methyl-2-pyrrolidin-1-ylmethyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



10

The compound was prepared from 4-aminodiphenyl ether and 1-methyl-2-pyrrolidin-1-ylmethyl-1H-indol-5-ylamine, as described in Example 1. Thus the product having a molecular weight of 440.55 ($C_{27}H_{28}N_4O_2$); MS (ESI): 441 ($M+H^+$) was obtained.

15 1-Methyl-2-pyrrolidin-1-ylmethyl-1H-indol-5-ylamine

Formic acid (0.11 ml) was added to a suspension of 1-methyl-5-nitro-2-pyrrolidin-1-ylmethyl-1H-indole (150 mg), ethanol (2 ml) and palladium(II) hydroxide on carbon (20%, 30 mg) and the suspension was heated to 60°C for 5 minutes. After gas production had ceased, the suspension was stirred for another 20 minutes and the catalyst was filtered off.

20 The filtrate was concentrated and distributed between saturated sodium carbonate solution and methyl tert-butyl ether. The organic phase was removed, dried over magnesium sulfate

and concentrated. Thus the product having a molecular weight of 229.33 ($C_{14}H_{19}N_3$); MS (ESI): 230 ($M+H^+$) was obtained.

1-Methyl-5-nitro-2-pyrrolidin-1-ylmethyl-1H-indole

- 5 Mesyl chloride (92 mg) was added dropwise to a solution cooled to 0°C of (1-methyl-5-nitro-1H-indol-2-yl)methanol (121 mg) in dichloromethane (10 ml) and triethylamine (0.17 ml). After 15 minutes, pyrrolidine (142 mg) was added and the solution was then stirred at room temperature for 1 hour. The reaction solution was washed with saturated sodium carbonate solution, dried over magnesium sulfate and concentrated. The residue was
- 10 purified via chromatography on silica gel (eluent: ethyl acetate/triethylamine 99:1). Thus the product having a molecular weight of 259.31 ($C_{14}H_{17}N_3O_2$); MS (ESI): 260 ($M+H^+$) was obtained.

(1-Methyl-5-nitro-1H-indol-2-yl)methanol

- 15 Sulfuric acid (96% strength, 0.64 ml) was added dropwise to a suspension cooled to 0°C of lithium aluminum hydride in tetrahydrofuran (50 ml) within 20 minutes. After 20 minutes, a solution of ethyl 1-methyl-5-nitro-1H-indole 2-carboxylate (1.85 g) in tetrahydrofuran (40 ml) was added dropwise. After 30 minutes, water (2 ml) was added. After 30 minutes, the resulting precipitate was filtered off and the filtrate was concentrated. The crude
- 20 product was purified via chromatography on silica gel (eluent: n-heptane/ethyl acetate 3:2). Thus the product having a molecular weight of 206.20 ($C_{10}H_{10}N_2O_3$); MS (ESI): 207 ($M+H^+$) was obtained.


Ethyl 1-methyl-5-nitro-1H-indole 2-carboxylate

- 25 A suspension of ethyl 5-nitro-1H-indole 2-carboxylate (2.34 g), potassium carbonate (3.45 g), methyl iodide (2.13 g) and acetonitrile (30 ml) was kept at 60°C for 6 hours. After cooling to room temperature, water was added and the precipitated product was isolated by

filtration. Thus the product having a molecular weight of 248.24 ($C_{12}H_{12}N_2O_4$); MS (ESI): 249 ($M+H^+$) was obtained.

Example 4

5 1-[1-(2-Dimethylaminoethyl)-1H-benzimidazol-5-yl]-3-(4-phenoxyphenyl)urea



Zinc dust (250 mg) was added to a solution of 1-[4-(2-dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (50 mg) in dichloromethane (10 ml) and glacial acetic acid (1 ml). After 10 minutes, the inorganic material was filtered off via kieselguhr. The filtrate was washed with a sodium carbonate solution (10% strength), dried over magnesium sulfate and concentrated. The residue was taken up in dichloromethane (5 ml) and ethanol (5 ml) and admixed with dimethylformamide dimethyl acetal (0.3 ml) and formic acid (0.3 ml). Dichloromethane was evaporated by heating the mixture by means of a hot-air gun. The remaining mixture was concentrated and distributed between dichloromethane and a sodium carbonate solution (10% strength). The organic phase was removed, dried and concentrated. The residue was purified by preparative HPLC. Thus the product having a molecular weight of 415.50 ($C_{24}H_{25}N_5O_2$); MS (ESI): 416 ($M+H^+$) was obtained. Melting point of the hydrochloride: 213-215 °C.

1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea

A solution of 2-dimethylaminoethylamine in dimethylformamide (1M, 2 ml) and 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea (200 mg) was stirred for 48 hours. The mixture was distributed between dichloromethane and a sodium carbonate solution (10%

strength). The organic phase was dried and concentrated. The residue was recrystallized from toluene. Melting point: 178-180°C.


1-(4-Fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea

- 5 4-Fluoro-3-nitrophenyl isocyanate (2.2 mmol) was added to a solution of 4-phenoxyaniline (2 mmol) in dimethylformamide (20 ml). After 2 days, the reaction mixture was distributed between dichloromethane and a saturated sodium carbonate solution. The organic phase was dried and concentrated. The residue was purified via chromatography on silica gel (eluent: ethyl acetate/dichloromethane 95:5) and subsequent recrystallization from ethyl
10 acetate/hexane. Melting point: 174-176°C.

Example 5

1-[1-(2-Dimethylaminoethyl)-2-methyl-1H-benzimidazol-5-yl]-3-(4-isopropoxyphenyl)-
urea

15



- 1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-isopropoxyphenyl)urea (75 mg) was reduced using zinc dust, as described in Example 4. The reaction product was dissolved in methanol and admixed with triethyl orthoacetate (0.5 ml) and glacial acetic acid (0.2 ml). The mixture was heated under reflux for 5 minutes. Volatile components
20 were removed. The residue was distributed between dichloromethane and a sodium carbonate solution. The organic phase was dried and concentrated. The residue was purified by preparative HPLC. Thus the product having a molecular weight of 395.51 ($C_{22}H_{29}N_5O_2$); MS (ESI): 396 ($M+H^+$) was obtained.

1-[4-(2-Dimethylaminoethoxy)-3-nitrophenyl]-3-(4-isopropoxyphenyl)urea


The compound was obtained from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 2-dimethylaminoethylamine as in Example 4. The compound was reacted further without purification.

1-(4-Fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea

The compound was obtained from 4-fluoro-3-nitrophenyl isocyanate and 4-isopropoxyaniline as in Example 4. Melting point: 170-172°C.

Example 6

1-[1-(1-Ethylpyrrolidin-2-ylmethyl)-2-methyl-1H-benzoimidazol-5-yl]-3-(4-isopropoxyphenyl)urea




The compound was prepared from 1-{4-[(1-ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-isopropoxyphenyl)urea, as described in Example 5. Thus the product having a molecular weight of 435.57 ($C_{25}H_{33}N_5O_2$); MS (ESI): 436 ($M+H^+$) was obtained. Melting point: (ethyl acetate/hexane): 185-187°C.

1-{4-[(1-Ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-isopropoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-ethylpyrrolidin-2-ylmethylamine, as described in Example 4, and reacted further without any further purification.

5 Example 7

1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-urea



- 10 The compound was prepared from 1-(4-isopropoxyphenyl)-3-[3-nitro-4-(2-piperidin-1-ylethylamino)phenyl]urea, as described in Example 5. Thus the product having a molecular weight of 435.57 ($C_{25}H_{33}N_5O_2$); MS (ESI): 436 ($M+H^+$) was obtained.


1-(4-Isopropoxyphenyl)-3-[3-nitro-4-(2-piperidin-1-ylethylamino)phenyl]urea

- 15 The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-(2-aminoethyl)piperidine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate): 157-159°C.

Example 8

1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-morpholin-4-yl-ethyl)-1H-benzoimidazol-5-yl]urea

20



The compound was prepared from 1-(4-isopropoxyphenyl)-3-[4-(2-morpholin-4-ylethylamino)-3-nitrophenyl]urea, as described in Example 5. Thus the product having a molecular weight of 437.55 ($C_{24}H_{31}N_5O_3$); MS (ESI): 438 ($M+H^+$) was obtained.

5


1-(4-Isopropoxyphenyl)-3-[4-(2-morpholin-4-yl-ethylamino)-3-nitrophenyl]urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-isopropoxyphenyl)urea and 1-(2-aminoethyl)morpholine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate): 191-193°C.

10

Example 9

1-(4-Isopropoxyphenyl)-3-[2-methyl-1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]urea



15

The compound was prepared from 1-[3-nitro-4-(2-pyrrolidin-1-ylethylamino)phenyl]-3-(4-phenoxyphenyl)urea, as described in Example 5. Thus the product having a molecular weight of 455.56 ($C_{27}H_{29}N_5O_2$); MS (ESI): 456 ($M+H^+$) was obtained.


1-[3-Nitro-4-(2-pyrrolidin-1-ylethylamino)phenyl]-3-(4-phenoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 1-(2-aminoethyl)pyrrolidine (60°C, 5 h), as described in Example 4. Melting point (ethyl acetate/hexane): 179-181°C.

5

Example 10

1-[2-Methyl-1-(2-dimethylaminoethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




The compound was prepared from 1-[4-(2-dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea, as described in Example 5. Thus the product having a molecular weight of 429.53 (C₂₅H₂₇N₅O₂); MS (ESI): 430 (M+H⁺) was obtained.

10

Example 11

1-(4-Phenoxyphenyl)-3-[1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]urea




15

The compound was prepared from 1-[3-nitro-4-(2-pyrrolidin-1-ylethylamino)phenyl]-3-(4-phenoxyphenyl)urea, as described in Example 4. Thus the product having a molecular weight of 441.54 (C₂₆H₂₇N₅O₂); MS (ESI): 442 (M+H⁺) was obtained.

Example 12

1-[2-Benzyl-1-(2-dimethylaminoethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




5

1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (75 mg) was reduced as described in Example 4. The crude product was treated with phenylacetic acid (0.33 mmol), activated with HATU (0.33 mmol), and diisopropylamine (0.7 mmol) in dimethylformamide (1.5 ml) for 3 hours. The reaction mixture was distributed between dichloromethane and a sodium carbonate solution (10% strength). The organic phase was dried and concentrated. The residue was heated under reflux in trifluoroacetic acid (1 ml), water (1 ml) and acetonitrile (0.5 ml) for 5 minutes. Volatile components were evaporated and the residue was purified by preparative HPLC. Thus the product having a molecular weight of 505.63 ($C_{31}H_{31}N_5O_2$); MS (ESI): 506 ($M+H^+$) was obtained.

15

Example 13


1-[1-(2-Dimethylaminoethyl)-2-phenyl-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



1-[4-(2-Dimethylaminoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (50 mg) was reduced as described in Example 4. After filtration via kieselguhr, benzaldehyde (0.2 ml) was added to the filtrate. The reaction mixture was washed with a sodium carbonate solution (10% strength), dried and admixed with manganese dioxide (0.5 g). After 15 minutes, the inorganic material was filtered off and the filtrate was concentrated. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 491.60 ($C_{30}H_{29}N_5O_2$); MS (ESI): 492 ($M+H^+$) was obtained.

10 Example 14


1-[2-Ethyl-1-(2-pyrrolidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



1-[4-(2-Pyrrolidinoethylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethyl orthopropionate according to Example 5. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.

20 Example 15

1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea




1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethyl
 5 orthoacetate, as described in Example 5. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.

1-[2-Methyl-1-(2-piperidin-1-ylethyl)-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea

10 The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 1-(2-aminoethyl)piperidine (60°C, 4 h), as described in Example 4. Melting point (ethyl acetate/hexane): 163-165°C.

Example 16

15 1-[1-(1-Ethylpyrrolidin-2-ylmethyl)-2-methyl-1H-benzoimidazol-5-yl]-3-(4-phenoxyphenyl)urea



1-{4-[(1-Ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-phenoxyphenyl)urea was reduced as described in Example 4. The crude product was reacted with triethylorthoacetate, as described in Example 5. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 469.59 ($C_{28}H_{31}N_5O_2$); MS (ESI): 470 ($M+H^+$) was obtained.


1-{4-[(1-Ethylpyrrolidin-2-ylmethyl)amino]-3-nitrophenyl}-3-(4-phenoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and C-(1-ethylpyrrolidin-2-yl)methylamine (60°C, 4 h), as described in Example 4.

10 Melting point (ethyl acetate/hexane): 129-132°C.

Example 17

1-(2-Dimethylaminomethyl-1H-benzoimidazol-5-yl)-3-(4-phenoxyphenyl)urea



15

1-[4-(2,4-Dimethoxybenzylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea (75 mg) was reduced as described in Example 4. The reduced product was reacted with dimethylaminoacetic acid (1 mmol), HATU (1 mmol) and diisopropylamine (2 mmol) in dimethylformamide (3 ml). After 3 hours, the mixture was distributed between ethyl acetate and a sodium carbonate solution. The organic phase was dried and concentrated. The crude product was purified by preparative HPLC. Thus the intermediate (N-{2-amino-5-[3-(4-phenoxyphenyl)ureido]phenyl}-2-dimethylaminoacetamide) having a molecular weight of 419.49 ($C_{23}H_{25}N_5O_3$); MS (ESI): 420 ($M+H^+$) was obtained.

20

This material was heated under reflux with pivalic acid and volatile components were then removed under a high vacuum. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 401.47 ($C_{23}H_{23}N_5O_2$); MS (ESI): 402 ($M+H^+$) was obtained.

5


1-[4-(2,4-Dimethoxybenzylamino)-3-nitrophenyl]-3-(4-phenoxyphenyl)urea

The compound was prepared from 1-(4-fluoro-3-nitrophenyl)-3-(4-phenoxyphenyl)urea and 2,4-dimethoxybenzylamine (60°C, 12 h), as described in Example 4. Melting point (ethyl acetate): 214-216°C.

10

Example 18

1-[1-(2-Dimethylaminoethyl)-2,3-dimethyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



- 15 The compound was prepared from 1-(2-dimethylaminoethyl)-2,3-dimethyl-1H-indol-5-ylamine and 4-phenoxyaniline, as described in Example 1. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 442.57 ($C_{27}H_{30}N_4O_3$); MS (ESI): 443 ($M+H^+$) was obtained.

20 1-(2-Dimethylaminoethyl)-2,3-dimethyl-1H-indol-5-ylamine

The compound was obtained by hydrogenation of [2-(2,3-dimethyl-5-nitroindol-1-yl)ethyl]dimethylamine, as described in Example 3. Thus the product having a molecular weight of 231.34 ($C_{14}H_{21}N_3$); MS (ESI): 232 ($M+H^+$) was obtained.


[2-(2,3-Dimethyl-5-nitroindol-1-yl)ethyl]dimethylamine

Sodium hydride (50% strength in oil; 0.8 g) was added to 2,3-dimethyl-5-nitro-1H-indole (1 g) in tetrahydrofuran (10 ml) at 0°C. After 30 minutes at room temperature,

- 5 dimethylaminoethyl chloride (hydrochloride; 1.1 g) was added and the mixture was then heated at 65°C for two hours. The cooled reaction solution was extracted with dichloromethane. The organic phase was dried and concentrated. The crude product was purified via chromatography on silica gel (eluent: dichloromethane/methanol 9:1). Thus the product having a molecular weight of 261.33 ($C_{14}H_{19}N_3O_2$); MS (ESI): 262 ($M+H^+$)
- 10 was obtained.

Example 19

1-[1-(2-Dimethylaminoethyl)-2-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



- 15 The compound was prepared from 1-(2-dimethylaminoethyl)-2-methyl-1H-indol-5-ylamine and 4-phenoxyaniline, as described in Example 1. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 428.54 ($C_{26}H_{28}N_4O_3$); MS (ESI): 428 ($M+H^+$) was obtained.

20 1-(2-Dimethylaminoethyl)-2-methyl-1H-indol-5-ylamine


The compound was obtained by hydrogenation of [2-(2-methyl-5-nitroindol-1-yl)ethyl]dimethylamine, as described in Example 3. Thus the product having a molecular weight of 217.32 ($C_{13}H_{19}N_3$); MS (ESI): 218 ($M+H^+$) was obtained.

[2-(2-Methyl-5-nitroindol-1-yl)ethyl]dimethylamine


The compound was prepared from 2-methyl-5-nitro-1H-indole and dimethylaminoethyl chloride (hydrochloride) as in Example 18. Thus the product having a molecular weight of
5 247.30 ($C_{13}H_{17}N_3O_2$); MS (ESI): 248 ($M+H^+$) was obtained.

Table 3:


Examples of the formula I



5 where the moiety x_1 is














and x_2 is









10







and x_2 is listed in the column denoted "aniline" of the table below.






Ex-ample	Name	Aniline	Molecular formula	Mol-ecular weight	[M+H] ⁺
20	1-[4-(Cyclohexylmethyl-amino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₃₅ N ₅ O	433.60	434
21	1-[4-(Cyclohexylmethyl-amino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₃₅ N ₅ O	433.60	434
22	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-pyrrolidin-1-ylphenyl)urea		C ₂₃ H ₂₉ N ₅ O	391.52	392
23	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2,5-dimethylpyrrolidin-1-yl)phenyl]urea		C ₂₅ H ₃₃ N ₅ O	419.57	420
24	1-[4-(3,6-Dihydro-2H-pyridin-1-yl)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₄ H ₂₉ N ₅ O	403.53	404
25	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2,6-dimethylmorpholin-4-yl)phenyl]urea		C ₂₅ H ₃₃ N ₅ O ₂	435.57	436

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
26	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-thiomorpholin-4-yl-phenyl)urea		C23H29N5OS	423.58	424
27	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-methylpiperidin-1-yl)phenyl]urea		C25H33N5O	419.57	420
28	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-ethylpiperidin-1-yl)phenyl]urea		C26H35N5O	433.60	434
29	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-methylpiperidin-1-yl)phenyl]urea		C25H33N5O	419.57	420
30	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3,3-dimethylpiperidin-1-yl)phenyl]urea		C26H35N5O	433.60	434






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
31	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3,5-dimethylpiperidin-1-yl)phenyl]urea		C ₂₆ H ₃₅ N ₅ O	433.60	434
32	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(4-phenylpiperidin-1-yl)phenyl]urea		C ₃₀ H ₃₅ N ₅ O	481.65	482
33	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(4-methylpiperidin-1-yl)phenyl]urea		C ₂₅ H ₃₃ N ₅ O	419.57	420
34	1-(4-Azepan-1-ylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₃₃ N ₅ O	419.57	420
35	1-[4-(Benzylmethylamino)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₇ H ₃₁ N ₅ O	441.58	442
36	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3[4-(methylphenethylamino)-phenyl]urea		C ₂₈ H ₃₃ N ₅ O	455.61	456

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
37	1-[4-(Butylmethyl- amino)phenyl]-3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₂₄ H ₃₃ N ₅ O	407.56	408
38	1-[4-(Benzylbutyl- amino)phenyl]-3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₃₀ H ₃₇ N ₅ O	483.66	484
39	1-(4- Dibutylaminophenyl)- 3-[1-(2- dimethylaminoethyl)- 1H-indol-5-yl]urea		C ₂₇ H ₃₉ N ₅ O	449.64	450
40	1-[1-(2-Dimethyl- aminoethyl)-1H-indol- 5-yl]-3-[(4aR,8aS)-4- (octahydroisoquinolin- 2-yl)phenyl]urea		C ₂₈ H ₃₇ N ₅ O	459.64	460
41	1-[1-(2- Dimethylaminoethyl)- 1H-indol-5-yl]-3-[4-(2- methylpyrrolidin-1- yl)phenyl]urea		C ₂₄ H ₃₁ N ₅ O	405.55	406






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
42	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(5-ethyl-2-methylpiperidin-1-yl)phenyl]urea		C ₂₇ H ₃₇ N ₅ O	447.63	448
43	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(methylpyridin-3-ylmethylamino)phenyl]urea		C ₂₆ H ₃₀ N ₆ O	442.57	443
44	1-[4-(3-Azabicyclo[3.2.2]non-3-yl)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₇ H ₃₅ N ₅ O	445.61	446
45	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-isopropylpyrrolidin-1-yl)phenyl]urea		C ₂₆ H ₃₅ N ₅ O	433.60	434
46	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-isobutylpyrrolidin-1-yl)phenyl]urea		C ₂₇ H ₃₇ N ₅ O	447.63	448
47	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-phenylpyrrolidin-1-yl)phenyl]urea		C ₂₉ H ₃₃ N ₅ O	467.62	468







Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
48	1-[1-(2-Dimethyl-aminoethyl)-1H-indol-5-yl]-3-[4-(3-trifluoromethylpiperidin-1-yl)-phenyl]urea		C ₂₅ H ₃₀ F ₃ N ₅ O	473.55	474
49	1-[1-(2-Dimethyl-aminoethyl)-1H-indol-5-yl]-3-[(4aR,8aR)-4-(octahydroisoquinolin-2-yl)phenyl]urea		C ₂₈ H ₃₇ N ₅ O	459.64	460
50	1-[4-(3,4-Dihydro-1H-isoquinolin-2-yl)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₈ H ₃₁ N ₅ O	453.59	454
51	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(((1S,5R)-1,3,3-trimethyl-6-azabicyclo[3.2.1]oct-6-yl)phenyl]urea		C ₂₉ H ₃₉ N ₅ O	473.67	474
52	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,6-dimethylphenyl)urea		C ₂₅ H ₃₄ N ₄ O ₂	422.58	423






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
53	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-methoxyphenyl)urea		C24H32N4O3	424.55	425
54	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-methylphenyl)urea		C24H32N4O2	408.55	409
55	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,5-dimethylphenyl)urea		C25H34N4O2	422.58	423
56	1-(3,5-Dichloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C23H28Cl2N4O	463.41	463
57	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-nitrophenyl)urea		C23H29N5O4	439.52	440
58	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3-methylphenyl)urea		C24H32N4O2	408.55	409

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
59	1-Benzyl-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]-1-(4-isobutoxyphenyl)urea		C ₃₀ H ₃₆ N ₄ O ₂	484.65	485
60	1-(3-Chloro-4-isobutoxy-5-methylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₄ H ₃₁ ClN ₄ O ₂	442.99	443
61	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-nitrophenyl)urea		C ₂₃ H ₂₉ N ₅ O ₄	439.52	440
62	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2,3-dimethylphenyl)urea		C ₂₅ H ₃₄ N ₄ O ₂	422.58	423
63	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(2-fluoro-4-isobutoxyphenyl)urea		C ₂₃ H ₂₉ FN ₄ O ₂	412.51	413






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
64	1-(3-Chloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₃ H ₂₉ ClN ₄ O ₂	428.97	429
65	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(3-fluoro-4-isobutoxyphenyl)urea		C ₂₃ H ₂₉ FN ₄ O ₂	412.51	413
66	1-(2-Chloro-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₃ H ₂₉ ClN ₄ O ₂	428.97	429
67	Methyl 5-{3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]ureido}-2-isobutoxy benzoate		C ₂₅ H ₃₂ N ₄ O ₄	452.56	453
68	1-(3-Cyano-4-isobutoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₄ H ₂₉ N ₅ O ₂	419.53	420
69	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-3,5-dimethylphenyl)urea		C ₂₅ H ₃₄ N ₄ O ₂	422.58	423







Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
70	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxy-2-trifluoromethylphenyl) urea		C ₂₄ H ₂₉ F ₃ N ₄ O	462.52	463
71	1-(4-Butylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₃ H ₃₀ N ₄ O	378.52	379
72	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-isobutoxyphenyl)urea		C ₂₃ H ₃₀ N ₄ O ₂	394.52	395
73	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(pyridin-3-yloxy)phenyl]urea		C ₂₄ H ₂₅ N ₅ O ₂	415.50	416
74	1-(3-Cyclopentyloxy-4-methoxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₃₂ N ₄ O ₃	436.56	437






Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
75	1-(4-Benzenesulfonyl-2-nitrophenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₂₅ N ₅ O ₅ S	507.57	508
76	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(2-methoxyphenoxy)-phenyl]urea		C ₂₆ H ₂₈ N ₄ O ₃	444.54	445
77	1-[4-(3-Chlorophenoxy)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₂₅ ClN ₄ O ₂	448.96	449
78	1-Biphenyl-4-yl-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₂₆ N ₄ O	398.51	399
79	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(2-methoxy-4-phenylaminophenyl)-urea		C ₂₆ H ₂₉ N ₅ O ₂	443.55	444
80	1-(4-Benzyloxyphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429

Ex- am- ple	Name	Aniline	Molecular formula	Mol.- weight	[M+H] ⁺
81	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4'-fluorobiphenyl-4-yl)urea		C ₂₅ H ₂₅ FN ₄ O	416.50	417
82	1-(4-Benzylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₈ N ₄ O	412.54	413
83	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-pyridin-4-ylmethylphenyl)urea		C ₂₅ H ₂₇ N ₅ O	413.53	414
84	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-p-tolyloxyphenyl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
85	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-phenylsulfanyphenyl)urea		C ₂₅ H ₂₆ N ₄ OS	430.58	431

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
86	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-trifluoromethylphenoxy)phenyl]urea		C ₂₆ H ₂₅ F ₃ N ₄ O	482.51	483
87	1-(4-Butyl-2-methylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₄ H ₃₂ N ₄ O	392.55	393
88	1-(4'-Cyanobiphenyl-4-yl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₅ N ₅ O	423.52	424
89	1-[4-(4-Chlorophenoxy)-2-trifluoromethylphenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₄ ClF ₃ N ₄	516.95	517
90	1-[3-Chloro-4-(pyrimidin-2-yloxy)phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₃ H ₂₃ ClN ₆ O ₂	450.93	451
91	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(5-methoxy-2-methylbiphenyl-4-yl)urea		C ₂₇ H ₃₀ N ₄ O ₂	442.57	443

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
92	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(piperidine-1-sulfonyl)phenyl]urea		C ₂₄ H ₃₁ N ₅ O ₃ S	469.61	470
93	Ethyl 5-(4-{3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]ureido}phenyl)-2-methylfuran-3-carboxylate		C ₂₇ H ₃₀ N ₄ O ₄	474.56	475
94	1-(4-Benzooxazol-2-ylphenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₆ H ₂₅ N ₅ O ₂	439.52	440
95	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(piperidine-1-carbonyl)phenyl]urea		C ₂₅ H ₃₁ N ₅ O ₂	433.56	434
96	1-[3-Cyano-4-(3-trifluoromethylphenyl)sulfanyl]phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₇ H ₂₄ F ₃ N ₅ O	523.58	524

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
97	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-heptafluoropropyl-sulfanyphenyl)urea		C ₂₂ H ₂₁ F ₇ N ₄ O	522.49	523
98	1-(4-Benzenesulfonyl-3-chlorophenyl)-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C ₂₅ H ₂₅ ClN ₄ O ₃	497.02	497
99	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(pyrimidin-2-yloxy)phenyl]urea		C ₂₃ H ₂₄ N ₆ O ₂	416.49	417
100	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(2-methoxybiphenyl-4-yl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
101	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(6-methoxybiphenyl-3-yl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
102	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-[1,3]dithiolan-2-ylphenyl)urea		C ₂₂ H ₂₆ N ₄ OS ₂	426.61	427

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
103	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(thiophen-2-ylsulfanyl)phenyl]urea		C23H24N4OS2	436.60	437
104	3-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-1-(4-methoxyphenyl)-1-methylurea		C21H26N4O2	366.47	367
105	1-[4-(2-Chlorophenoxy)-phenyl]-3-[1-(2-dimethylaminoethyl)-1H-indol-5-yl]urea		C25H25ClN4O2	448.96	449
106	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(6-phenoxy-pyridin-3-yl)urea		C24H25N5O2	415.50	416
107	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-m-tolyloxyphenyl)urea		C26H28N4O2	428.54	429

Ex- am- ple	Name	Aniline	Molecular formula	Mol- ecular weight	[M+H] ⁺
108	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-(4-o-tolyloxyphenyl)urea		C ₂₆ H ₂₈ N ₄ O ₂	428.54	429
109	1-[1-(2-Dimethylaminoethyl)-1H-indol-5-yl]-3-[4-(3-methoxyphenoxy)-phenyl]urea		C ₂₆ H ₂₈ N ₄ O ₃	444.54	445

The molecule ion peak ([M+H]⁺) was taken from ESI mass spectra.

- 5 The examples 20 – 51 and 71 -109 were prepared according to Example 1.

Synthesis of Examples 52 – 70

- Carbonyldiimidazole (0.25 mmol) was added to 1-(2-dimethylaminoethyl)-1H-indol-5-ylamine (0.25 mmol) in dimethylformamide (1 ml) at 0°C. After 1 hour at room temperature, the reaction solution was cooled again to 0°C and the appropriate aminophenol (0.25 mmol) was added. After 15 hours at room temperature, cesium carbonate (0.5 mmol) and isobutyl iodide (0.5 mmol) were added and the solution was heated at 80°C for 2 hours. The reaction solutions were filtered and the filtrate was washed with sodium bicarbonate (5% strength) and sodium chloride solution (5% strength). The organic phase was dried and concentrated. The crude product was purified by preparative HPLC. Thus the product having the molecular weight indicated in Table 3 and the molecule ion peak of the mass spectrum, likewise indicated in Table 3, was obtained.

Precursors of Examples 20-51

A mixture of 4-fluoronitrobenzene (0.35 mmol), potassium carbonate (0.7 mmol), the appropriate amine and dimethylformamide (1 ml) was heated to 100°C for three hours. The reaction solution was filtered and washed with sodium chloride solution (5% strength). The organic phase was dried and concentrated. The 4-nitroaniline obtained as crude product was dissolved in glacial acetic acid (1 ml) and zinc dust (0.25 g) was added. After a reaction time of 3 hours, the reaction solution was diluted with ethyl acetate (10 ml), filtered and the filtrate was washed with sodium chloride solution (5% strength). The filtrate was dried and concentrated. The obtained crude product, 4-substituted aniline, was reacted further without any further purification.

The following 4-nitroanilines were prepared:

1-(4-nitrophenyl)azocan

cyclohexylmethyl-(4-nitrophenyl)amine

1-(4-nitrophenyl)pyrrolidine

2,5-dimethyl-1-(4-nitrophenyl)pyrrolidine

1-(4-nitrophenyl)-1,2,3,6-tetrahydropyridine

2,6-dimethyl-4-(4-nitrophenyl)morpholine

4-(4-nitrophenyl)thiomorpholine

2-methyl-1-(4-nitrophenyl)piperidine

2-ethyl-1-(4-nitrophenyl)piperidine

3-methyl-1-(4-nitrophenyl)piperidine

3,3-dimethyl-1-(4-nitrophenyl)piperidine

3,5-dimethyl-1-(4-nitrophenyl)piperidine

1-(4-nitrophenyl)-4-phenylpiperidine

4-methyl-1-(4-nitrophenyl)piperidine

2-(4-nitrophenyl)-1,2,3,4-tetrahydroisoquinoline

1-(4-nitrophenyl)azepan

benzylmethyl-(4-nitrophenyl)amine

5 methyl-(4-nitrophenyl)phenethylamine

butylmethyl-(4-nitrophenyl)amine

benzylbutyl-(4-nitrophenyl)amine

dibutyl-(4-nitrophenyl)amine

(4aR,8aS)-2-(4-nitrophenyl)decahydroisoquinoline

10 2-methyl-1-(4-nitrophenyl)pyrrolidine

5-ethyl-2-methyl-1-(4-nitrophenyl)piperidine

methyl-(4-nitrophenyl)pyridine-3-ylmethylamine

3-(4-nitrophenyl)-3-azabicyclo[3.2.2]nonane

2-isopropyl-1-(4-nitrophenyl)pyrrolidine

15 2-isobutyl-1-(4-nitrophenyl)pyrrolidine

1-(4-nitrophenyl)-3-phenylpyrrolidine

1-(4-nitrophenyl)-3-trifluoromethylpiperidine

(4aR,8aR)-2-(4-nitrophenyl)decahydroisoquinoline

(1S,5R)-1,3,3-trimethyl-6-(4-nitrophenyl)-6-azabicyclo[3.2.1]octane

20 All of the 4-nitroanilines listed above showed the expected molecule ion peak in the ESI mass spectrum.

The following 4-substituted anilines were prepared:

- 4-azocan-1-ylphenylamine
- N-cyclohexyl-N-methylbenzene-1,4-diamine
- 4-pyrrolidin-1-ylphenylamine
- 4-(2,5-dimethylpyrrolidin-1-yl)phenylamine
- 5 4-(3,6-dihydro-2H-pyridin-1-yl)phenylamine
- 4-(2,6-dimethylmorpholin-4-yl)phenylamine
- 4-thiomorpholin-4-ylphenylamine
- 4-(2-methylpiperidin-1-yl)phenylamine
- 4-(2-ethylpiperidin-1-yl)phenylamine
- 10 4-(3-methylpiperidin-1-yl)phenylamine
- 4-(3,3-dimethylpiperidin-1-yl)phenylamine
- 4-(3,5-dimethylpiperidin-1-yl)phenylamine
- 4-(4-phenylpiperidin-1-yl)phenylamine
- 4-(4-methylpiperidin-1-yl)phenylamine
- 15 4-(3,4-dihydro-1H-isoquinolin-2-yl)phenylamine
- 4-azepan-1-ylphenylamine
- N-benzyl-N-methylbenzene-1,4-diamine
- N-methyl-N-phenethylbenzene-1,4-diamine
- N-butyl-N-methylbenzene-1,4-diamine
- 20 N-benzyl-N-butylbenzene-1,4-diamine
- N,N-dibutylbenzene-1,4-diamine

(4aR,8aS)-4-(octahydroisoquinolin-2-yl)phenylamine

4-(2-methylpyrrolidin-1-yl)phenylamine

4-(5-ethyl-2-methylpiperidin-1-yl)phenylamine

N-methyl-N-pyridin-3-ylmethylbenzene-1,4-diamine

5 4-((1S,5R)-1,3,3-trimethyl-6-azabicyclo[3.2.1]oct-6-yl)phenylamine

4-(3-azabicyclo[3.2.2]non-3-yl)phenylamine

4-(2-isopropylpyrrolidin-1-yl)phenylamine

4-(2-isobutylpyrrolidin-1-yl)phenylamine

4-(3-phenylpyrrolidin-1-yl)phenylamine


10 4-(3-trifluoromethylpiperidin-1-yl)phenylamine

(4aR,8aR)-4-(octahydroisoquinolin-2-yl)phenylamine.

All of the 4-substituted anilines listed above showed the expected molecule ion peak in the ESI mass spectrum.

15 Example 110


4-Phenoxyphenyl [1-(2-dimethylaminoethyl)-1H-indol-5-yl]carbamate



The compound was prepared according to Example 1 by reacting the carbonyldiimidazole-activated indolamine with deprotonated 4-phenoxyphenol. Thus the product having a
20 molecular weight of 415.50 (C₂₅H₂₅N₃O₃); MS (ESI): 416 (M+H⁺) was obtained.

Example 111

1-(2-Imidazol-1-ylmethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



Mesyl chloride (47 μ l) was added to 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea (0.2 g) and triethylamine (0.16 ml) in dichloromethane (4 ml) at 0°C. After 10 minutes, imidazole (185 mg) was added. After 12 hours, the reaction solution was washed with sodium chloride solution, dried and concentrated. The crude product was purified by preparative HPLC. Thus the product having a molecular weight of 437.51 ($C_{26}H_{23}N_5O_2$); MS (ESI): 438 ($M+H^+$) was obtained.

1-(2-Hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea


(5-Amino-1-methyl-1H-indol-2-yl)methanol was reacted with 4-phenoxyaniline and carbonyldiimidazole, as described in Example 1. Thus the product having a molecular weight of 387.44 ($C_{23}H_{21}N_3O_3$); MS (ESI): 388 ($M+H^+$) was obtained.

(5-Amino-1-methyl-1H-indol-2-yl)methanol

(1-Methyl-5-nitro-1H-indol-2-yl)methanol was hydrogenated as described in Example 3. Thus the product having a molecular weight of 176.22 ($C_{10}H_{12}N_2O$); MS (ESI): 177 ($M+H^+$) was obtained.

Example 112

1-[1-Methyl-2-(2-methyl-4,5-dihydroimidazol-1-ylmethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea




The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 2-methyl-4,5-dihydroimidazole, as described in Example 111.

Thus the product having a molecular weight of 453,55 ($C_{27}H_{27}N_5O_2$); MS (ESI): 454

5 ($M+H^+$) was obtained.

Example 113


1-(2-Cyclohexylaminomethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea



10 The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and cyclohexylamine, as described in Example 111. Thus the product having a molecular weight of 468.60 ($C_{29}H_{32}N_4O_2$); MS (ESI): 469 ($M+H^+$) was obtained.

Example 114

15 1-[2-(3-Dimethylaminopyrrolidin-1-ylmethyl)-1-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea




The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 3-dimethylaminopyrrolidine, as described in Example 111. Thus the product having a molecular weight of 483.62 ($C_{29}H_{33}N_5O_2$); MS (ESI): 484 ($M+H^+$) was obtained.

5

Example 115


1-[2-(4-Hydroxypiperidin-1-ylmethyl)-1-methyl-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea



The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 4-hydroxypiperidin, as described in Example 111. Thus the product having a molecular weight of 470.58 ($C_{28}H_{30}N_4O_3$); MS (ESI): 471 ($M+H^+$) was obtained.

Example 116

1-[1-Methyl-2-(4-phenylpiperidin-1-ylmethyl)-1H-indol-5-yl]-3-(4-phenoxyphenyl)urea




The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and 4-phenylpiperidine, as described in Example 111. Thus the product having a molecular weight of 530.68 ($C_{34}H_{34}N_4O_2$); MS (ESI): 531 ($M+H^+$) was obtained.

5

Example 117


N-(1-{1-Methyl-5-[3-(4-phenoxyphenyl)ureido]-1H-indol-2-ylmethyl}pyrrolidin-3-yl)acetamide



10 The compound was prepared from 1-(2-hydroxymethyl-1-methyl-1H-indol-5-yl)-3-(4-phenoxyphenyl)urea and pyrrolidin-3-ylacetamide, as described in Example 111. Thus the product having a molecular weight of 497.60 ($C_{29}H_{31}N_5O_3$); MS (ESI): 498 ($M+H^+$) was obtained.

15 Example 118

1-(4-Phenoxyphenyl)-3-(2-pyrrolidin-1-ylmethylbenzofuran-5-yl)urea



The compound was prepared from 2-pyrrolidin-1-ylmethylbenzofuran-5-ylamine and 4-phenoxyaniline, as described in Example 1. Thus the product having a molecular weight of 427.51 ($C_{26}H_{25}N_3O_3$); MS (ESI): 428 ($M+H^+$) was obtained.

20

2-Pyrrolidin-1-ylmethylbenzofuran-5-ylamine

The compound was prepared by hydrogenation of 1-(5-nitrobenzofuran-2-ylmethyl)pyrrolidine, as described in Example 3. Thus the product having a molecular weight of 216.29 ($C_{13}H_{16}N_2O$); MS (ESI): 217 ($M+H^+$) was obtained.

1-(5-Nitrobenzofuran-2-ylmethyl)pyrrolidine


The compound was prepared from (5-nitrobenzofuran-2-yl)methanol, as described in Example 3. Thus the product having a molecular weight of 246.27 ($C_{13}H_{14}N_2O_3$); MS (ESI): 247 ($M+H^+$) was obtained.

(5-Nitrobenzofuran-2-yl)methanol

The compound was prepared by reduction of methyl 5-nitrobenzofuran 2-carboxylate, as described in Example 3. Thus the product having a molecular weight of 193.16 ($C_9H_7NO_4$); MS (ESI): 194 ($M+H^+$) was obtained.

Example 119

1-(4-Phenoxyphenyl)-3-(2-pyrrolidin-1-ylmethylbenzo[b]thiophen-5-yl)urea



The compound was prepared from 2-pyrrolidin-1-ylmethylbenzo[b]thiophen-5-ylamine and 4-phenoxyaniline, as described in Example 1. Thus the product having a molecular weight of 443.57 ($C_{26}H_{25}N_3O_2S$); MS (ESI): 444 ($M+H^+$) was obtained.

2-Pyrrolidin-1-ylmethylbenzo[b]thiophen-5-ylamine

The compound was prepared by hydrogenation of 1-(5-nitrobenzo[b]thiophen-2-ylmethyl)pyrrolidine, as described in Example 3. Thus the product having a molecular weight of 232.35 ($C_{13}H_{16}N_2S$); MS (ESI): 233 ($M+H^+$) was obtained.

1-(5-Nitrobenzo[b]thiophen-2-ylmethyl)pyrrolidine

The compound was prepared from (5-nitrobenzo[b]thiophen-2-yl)methanol, as described in Example 3. Thus the product having a molecular weight of 262.33 ($C_{13}H_{14}N_2O_2S$); MS (ESI): 263 ($M+H^+$) was obtained.


(5-Nitrobenzo[b]thiophen-2-yl)methanol

The compound was prepared by reduction of methyl 5-nitrobenzo[b]thiophene 2-carboxylate, as described in Example 3. Thus the product having a molecular weight of 209.23 ($C_9H_7NO_3S$); MS (ESI): 210 ($M+H^+$) was obtained.

In general, all of the basic compounds described were obtained either as free bases or in the form of a salt of one of the following acids: formic acid, trifluoroacetic acid or hydrochloric acid.

We claim:

1. A compound of the formula I,



in which

10 A is (C₁-C₈)alkyl, (C₀-C₈)alkylenearyl;
a 3- to 12-membered mono- or bicyclic ring which may contain one or more
heteroatoms from the group consisting of N, O and S and the 3- to 12-
membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃,
15 OCF₃, CN, (C₁-C₆)alkyl, aryl, CON(R₃₇)(R₃₈), N(R₃₉)(R₄₀), OH, O-(C₁-
C₆)alkyl, S-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

X is a bond, C(R₈)(R₉), C(OR₁₀)(R₁₁), O, N(R₁₂), S, SO, SO₂, CO;

20 R₈, R₉, R₁₀, R₁₁, R₁₂ are independently of one another H, (C₁-C₆)alkyl;

D is N, C(R₄₁);

E is N, C(R₄₂);

25 G is N, C(R₄₃);

L is N, C(R44);

R1, R2, R3, R41, R42, R43, R44 are independently of one another

5 H, F, Cl, Br, J, OH, CF₃, NO₂, CN, OCF₃, O-(C₁-C₆)alkyl, (C₁-C₄)-alkoxyalkyl, S-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₂-C₆)alkenyl, (C₃-C₈)cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₃-C₈)cycloalkenyl, O-(C₃-C₈)cycloalkenyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₈)alkylenearyl, S-aryl, N(R13)(R14), SO₂-CH₃, COOH, COO-(C₁-C₆)alkyl,
 10 CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21),
 a 5- to 7-membered heterocycle having 1-4 heteroatoms;

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

15 or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R15, R16 are independently of one another H, (C₁-C₆)alkyl,

20 or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R17, R19 are independently of one another H, (C₁-C₆)alkyl;

25

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24), O;

30 R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

W is N, C(R25);

R25 is H, (C₁-C₆)alkyl, aryl, a bond to Y;

T is N, C(R26);

5

R26 is H, (C₁-C₆)alkyl, aryl, (C₀-C₈)alkylenearyl, a bond to Y;

U is O, S, N(R27), -C(R30)=N-, -N=C(R31)-;

10 R27, R30, R31 are independently of one another H, (C₁-C₆)alkyl, a bond to Y;

Y is (C₁-C₈)alkylene, in which one or more carbons may be replaced by O, S, SO, SO₂, C(R32)(R33), CO, C(R34)(OR35) or N(R36);

15 R32, R33, R34, R35, R36 are independently of one another H, (C₁-C₆)alkyl, aryl;

R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form

20

a 3- to 8-membered ring in which one or more carbons may be replaced by O, N or S and the 3- to 8-membered ring may carry further substituents such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH, O-(C₁-C₆)alkyl or NHCO(C₁-C₆)alkyl;

25 R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

30 2. A compound of the formula I as claimed in claim 1, wherein

A is (C₂-C₇)alkyl, (C₀-C₃)alkylenearyl;

a 4- to 10-membered mono- or bicyclic ring which may contain one or more heteroatoms from the group consisting of N, O and S, and the 4- to 10-membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), O-(C₁-C₆)alkyl, or NHCO(C₁-C₆)alkyl;

X is a bond, C(R8)(R9), O, N(R12), S, SO₂;

R8, R9, R12 are independently of one another H, (C₁-C₆)alkyl;

D is N, C(R41);

E is N, C(R42);

G is N, C(R43);

L is N, C(R44);

where the total number of the nitrogen atoms defined by D, E, G and L is 0, 1 or 2;

R1, R2, R3, R41, R42, R43, R44 are independently of one another

H, F, Cl, Br, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, O-(C₃-C₈)cycloalkyl, (C₂-C₆)alkynyl, (C₀-C₈)alkylenearyl, -O-(C₀-C₃)-alkylenearyl, S-aryl, N(R13)(R14), SO₂-CH₃, COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21);

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

or R13 and R14 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

R15, R16 are independently of one another H, (C₁-C₆)alkyl,

or R15 and R16 together with the nitrogen atom to which they are bonded form a 5- to 6-membered ring, where, in the case of the 6-membered ring, a CH₂ group may be replaced by O or S;

5 R17, R19 are independently of one another H, (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24), O;

10

R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

15 W is N, C(R25);

R25 is H, (C₁-C₆)alkyl, aryl;

T is C(R26);

20

R26 is H, (C₁-C₆)alkyl, aryl, a bond to Y;

U is O, S, N(R27), -N=C(R31)-;

25 R27, R31 are independently of one another H, (C₁-C₆)alkyl, a bond to Y;

Y is (C₁-C₄)alkylene, in which a carbon may be replaced by SO₂, C(R32)(R33), CO or N(R36);

30 R32, R33, R36 are independently of one another H, (C₁-C₆)alkyl, aryl;

R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form a 4- to 7-membered ring in which one or more carbons may be replaced by O, N or S and the 4- to 7-membered ring may carry further substituents such as (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH or NHCO(C₁-C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

and the physiologically acceptable salts thereof.

3. A compound of the formula I as claimed in either of claims 1 and 2, wherein

A is (C₃-C₇)alkyl, (C₀-C₂)alkylenearyl;
 a 5- to 10-membered mono- or bicyclic ring which may contain 0, 1 or 2 heteroatoms from the group consisting of N, O and S, and the 5- to 10-membered ring may carry further substituents such as F, Cl, Br, NO₂, CF₃, (C₁-C₆)alkyl, aryl, O-(C₁-C₆)alkyl or NHCO(C₁-C₆)alkyl;

X is a bond, C(R8)(R9), O, N(R12);

R8, R9, R12 are independently of one another H, (C₁-C₆)alkyl;

D is N, C(R41);

E is N, C(R42);

G is N, C(R43);

L is N, C(R44);

where the total number of the nitrogen atoms defined by D, E, G and L is 0 or 1;

R1, R2, R3, R41, R42, R43, R44 are independently of one another

H, F, Cl, CF₃, NO₂, O-(C₁-C₆)alkyl, (C₁-C₆)alkyl, O-(C₃-C₈)cycloalkyl, (C₀-C₂)alkylenearyl, -O-(C₀-C₃)alkylenearyl, N(R13)(R14), COO-(C₁-C₆)alkyl, CON(R15)(R16), N(R17)CO(R18), N(R19)SO₂(R20), CO(R21);

5

R13, R14 are independently of one another H, (C₁-C₆)alkyl,

R15, R16 are independently of one another H, (C₁-C₆)alkyl,

10 R17, R19 are independently of one another H, (C₁-C₆)alkyl;

R18, R20, R21 are independently of one another (C₁-C₆)alkyl, aryl;

B is N(R24);

15

R24 is H, (C₁-C₆)alkyl;

R5 is H, (C₁-C₆)alkyl;

20 W is N, C(R25);

R25 is H, (C₁-C₆)alkyl;

T is C(R26);

25

R26 is H, (C₁-C₆)alkyl, a bond to Y;

U is O, S, N(R27);

30 R27 is H, (C₁-C₆)alkyl, a bond to Y;

Y is (C₁-C₃)alkylene, in which a carbon may be replaced by SO₂, C(R32)(R33) or CO;

R32, R33 are independently of one another H, (C₁-C₆)alkyl, aryl;

5 R6, R7 are independently of one another H, (C₁-C₆)alkyl, (C₃-C₇)cycloalkyl,

or R6 and Y or R6 and R7 together with the nitrogen atom to which they are bonded form
a 5- or 6-membered ring in which one or more carbons may be replaced by
O or N and the 5- or 6-membered ring may carry further substituents such as
10 (C₁-C₆)alkyl, aryl, CON(R37)(R38), N(R39)(R40), OH or NHCO(C₁-
C₆)alkyl;

R37, R38, R39, R40 are independently of one another H, (C₁-C₆)alkyl;

15 and the physiologically acceptable salts thereof.

4. A pharmaceutical comprising one or more of the compounds as claimed in one or
more of claims 1 to 3.

20

5. A pharmaceutical comprising one or more of the compounds as claimed in one or
more of claims 1 to 3 and one or more anorectic active substances.

25 6. The use of the compounds as claimed in one or more of claims 1 to 3 for preparing
a medicament for the prophylaxis or treatment of obesity.

7. The use of the compounds as claimed in one or more of claims 1 to 3 for preparing
30 a medicament for the prophylaxis or treatment of type II diabetes.

8. A compound as claimed in one or more of claims 1 to 3 in combination with at least one further anorectic active substance for application as medicament for the prophylaxis or treatment of obesity.

5

9. A compound as claimed in one or more of claims 1 to 3 in combination with at least one further anorectic active substance for application as medicament for the prophylaxis or treatment of type II diabetes.

10

10. A method for preparing a pharmaceutical comprising one or more of the compounds as claimed in one or more of claims 1 to 3, which comprises mixing the active substance with a pharmaceutically suitable carrier and bringing said mixture into a form suitable for administration.

15

11. A substance or composition for use in a method for the prophylaxis or treatment of obesity, said substance or composition comprising the compounds as claimed in one or more of claims 1 to 3, and said method comprising administering said substance or composition.

20

12. A substance or composition for use in a method for the prophylaxis or treatment of type II diabetes, said substance or composition comprising the compounds as claimed in one or more of claims 1 to 3, and said method comprising administering said substance or composition.

25

13. A compound as claimed in claim 1, substantially as herein described and illustrated.

5 14. A pharmaceutical as claimed in claim 4 or claim 5, substantially as herein described and illustrated.

10 15. Use as claimed in claim 6 or claim 7, substantially as herein described and illustrated.

15 16. A substance or composition for use in a method of prophylaxis or treatment as claimed in claim 8 or claim 9 or claim 11 or claim 12, substantially as herein described and illustrated.

20 17. A method as claimed in claim 10, substantially as herein described and illustrated.

25 18. A new compound, a new pharmaceutical, a new use of a compound as claimed in any one of claims 1 to 3, a substance or composition for a new use in a method of prophylaxis or treatment, or a new method for preparing a pharmaceutical, substantially as herein described.