

(51) International Patent Classification:
C12N 5/0775 (2010.01)(21) International Application Number:
PCT/IL2013/050136(22) International Filing Date:
13 February 2013 (13.02.2013)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data:
61/597,899 13 February 2012 (13.02.2012) US
61/597,909 13 February 2012 (13.02.2012) US(71) Applicant: **GAMIDA-CELL LTD.** [IL/IL]; 5 Nahum
Hafzadi Street, Ofer Building Givat Shaul, 95484 Jerus-
alem (IL).(72) Inventors: **PELED, Tony**; 19 Bareket Street, 90805
Mevaseret Zion (IL). **STEINHARDT, Yair**; 11 Ben Mai-
mon Boulevard, 92262 Jerusalem (IL).(74) Agents: **G. E. EHRlich (1995) LTD.** et al.; 11 Mena-
chem Begin Road, 5268104 Ramat Gan (IL).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).**Declarations under Rule 4.17:**

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

(54) Title: CULTURING OF MESENCHYMAL STEM CELLS

FIG. 27(57) **Abstract:** Methods of culturing mesenchymal stem cells are provided. The methods comprise culturing MSCs in a medium comprising nicotinamide and fibroblast growth factor 4 (FGF4). Populations of mesenchymal stem cells generated using the methods described herein and uses thereof are also provided.

CULTURING OF MESENCHYMAL STEM CELLS

FIELD AND BACKGROUND OF THE INVENTION

The present invention, in some embodiments thereof, relates to methods of
5 expanding mesenchymal stem cells and cell populations generated thereby.

Mesenchymal stem cells (MSCs) are non-hematopoietic cells that are capable of
differentiating into specific types of mesenchymal or connective tissues including
adipose, osseous, cartilaginous, elastic, neuronal, hepatic, pancreatic, muscular, and
fibrous connective tissues. The specific differentiation pathway which these cells enter
10 depends upon various influences from mechanical influences and/or endogenous
bioactive factors, such as growth factors, cytokines, and/or local microenvironmental
conditions established by host tissues.

MSCs reside in a diverse host of tissues throughout the adult organism and
possess the ability to 'regenerate' cell types specific for these tissues. Examples of these
15 tissues include adipose tissue, umbilical cord blood, periosteum, synovial membrane,
muscle, dermis, pericytes, blood, bone marrow and trabecular bone.

The multipotent character of mesenchymal stem cells make these cells an
attractive therapeutic tool and candidate for transplantation, capable of playing a role in
a wide range of clinical applications in the context of both cell and gene therapy
20 strategies. Mesenchymal cells may be used to enhance hematopoietic engraftment post-
transplantation, to correct inherited and acquired disorders of bone and cartilage, for
implantation of prosthetic devices in connective and skeletal tissue, and as vehicles for
gene therapy.

In culture, expanded MSC express a panel of key markers including CD105
25 (endoglin, SH2), CD73 (ecto-5' nucleotidase, SH3, SH4), CD166 (ALCAM), CD29
(β 1-integrin), CD44 (H-CAM), and CD90 (Thy-1). In contrast to hematopoietic stem
cells they lack CD45, CD34 and CD133 expression.

MSC can be identified by their ability to form colony forming units-fibroblast
(CFU-F) *in vitro*. However, these cells are heterogeneous with respect to their
30 proliferation and differentiation capacity. At least two morphologically distinct MSC
populations have been identified that differ not only in size but also in their cell division
rate and differentiation capacity. In addition, analysis of single cell-derived MSC

colonies from adult bone marrow revealed differential capacity of colonies to undergo osteogenic, adipogenic, and chondrogenic differentiation.

In most cases, unfractionated bone marrow-derived cells are used as the starting population for the culture of MSC. This isolation method relies on the adherence of fibroblast-like cells to a plastic surface and the removal of non-adherent hematopoietic cells. The resulting cells are poorly defined and give rise not only to heterogeneous MSC populations but also to osteoblasts and/or osteoprogenitor cells, fat cells, reticular cells, macrophages, and endothelial cells. To define the starting population more precisely, surface markers such as SH2 (CD105), SH3/SH4 (CD73), SSEA-4 and the low affinity nerve growth factor receptor (CD271), which enrich for MSC, have been employed [Simmons P.J et al. (1991) *Blood* 78:55–62; Conconi MT et al., (2006) *Int J Mol Med* 18:1089–96; Gang EJ et al., (2007) *Blood* 109:1743–51; Liu PG, (2005); *Zhongguo Shi Yan Xue Ye Xue Za Zhi* 13:656–9; Quirici N, et al., (2002) *Exp. Hematol* 30:783–91].

Another example of a cell surface antigen which has been targeting for isolating homogeneous populations of mesenchymal stem cells is stromal precursor antigen-1 (STRO-1). The STRO-1 antigen is expressed on the surface of approximately 10–20 % of adult human BM that includes all CFU-F, Glycophorin-A nucleated red cells, and a small subset of CD19 B-cells, but is not expressed on hematopoietic stem and progenitor cells (HSC) (Simmons and Torok-Storb, 1991). STRO-1 is widely regarded as a marker of early mesenchymal/stromal precursor cells, because it has been strongly linked to mesenchymal cell clonogenicity, plasticity, and other progenitor cell characteristics [Psaltis et al., (2010), *Journal of Cellular Physiology*, 530-540]. High co-expression of STRO-1 (STRO-1^{Bright}) with other surface markers, such as CD106, CD49a, CD146 or STRO-3 has been shown to greatly increase the cloning efficiency of BM MNC (Gronthos et al., 2008, *Methods Molecular Biology*, 449:45-57]. Freshly isolated STRO-1^{Bright} BM MNC also possess other hallmark features characteristic of multipotent stem cells, including in vivo quiescence, high telomerase activity, and an undifferentiated phenotype. Moreover, this population of cells lacks hematopoietic stem cell (CD34), leukocyte (CD45), and erythroid (Glycophorin-A) associated markers.

More recently, platelet derived growth factor receptor- β (PDGF-R β ; CD140b) was identified as a selective marker for the isolation of clonogenic MSC [Buhning HJ,

(2007) Ann N Y Acad Sci 1106:262–71]. Other reports demonstrated a 9.5-fold enrichment of MSC in bone marrow cells with prominent aldehyde dehydrogenase activity [Gentry T et al., (2007) Cytotherapy 9:259–74].

Even though MSCs multiply relatively easily in vitro, their proliferative potential and their stem cell characteristics are continuously decreased during prolonged culture. For example, it has been shown that expansion in culture leads to premature senescence (the process of aging characterized by continuous morphological and functional changes). Cells became much larger with irregular and flat shape and the cytoplasm became more granular. These senescence-associated effects are continuously acquired from the onset of in vitro culture (PLoS ONE, May 2008 | Volume 3 | Issue 5 | e2213). As a result, the successful manufacturing for commercialization of large batches from one donor of homogenous MSCs that preserve their characteristics following expansion in culture remains a challenge.

Due to the low or absent expression of MHC molecules, especially class II on mesenchymal stem cells, these cells may be considered immune privileged, thus paving the way for allogeneic transplantation of such cells for the treatment of a wide range of disorders. Accordingly, improved methods of expanding banks of mesenchymal stem cells have become an important factor for commercializing their use.

The role of growth factors in increasing proliferation and survival in MSCs has been widely studied over the past few years and many factors have been proposed for increasing the expansion efficiency of these cells.

For example, many protocols relating to the expansion of MSCs include culturing in the presence of basic fibroblast growth factor (b-FGF) (Vet Res Commun. 2009 Dec;33(8):811-21). It has been shown that b-FGF not only maintains MSC proliferation potential, it also retains osteogenic, adipogenic and chondrogenic differentiation potentials through the early mitogenic cycles.

Vascular endothelial growth factor (VEGF) has also been shown to increase MSC proliferation [Pons et al., Biochem Biophys Res Commun 2008, 376:419-422].

Exogenous addition of Hepatocyte growth factor (HGF) to MSC populations has been shown to affect proliferation, migration and differentiation (Furge et al., Oncogene 2000, 19:5582-5589).

Another proposed growth factor for increasing the expansion of MSCs is Platelet derived growth factor (PDGF) shown to be a potent mitogen of MSCs [Kang et al., J Cell Biochem 2005, 95:1135-1145].

Epidermal growth factor (EGF) and heparin-binding EGF have both been shown to promote ex vivo expansion of MSCs without triggering differentiation into any specific lineage [Tamama et al., Stem Cells 2006, 24:686-695; Krampera et al., Blood 2005, 106:59-66]. In addition to its mitogenic effect on MSCs, EGF also increases the number of colony-forming units by 25% [Tamama et al., J Biomed Biotechnol 2010, 79:5385].

Other have suggested the use of Wnt signalling agonists for expanding MSCs based on experiments which study Wnt signaling proliferation in MSCs. Canonical Wnt signalling was shown to maintain stem cells in an undifferentiated but self-renewing state. Addition of Wnt3a by activating the canonical Wnt pathway increased both proliferation and survival while preventing differentiation into the osteoblastic lineage in MSCs [Boland et al., J Cell Biochem 2004, 93:1210-1230].

The choice of growth factors to be used on MSCs was initially determined based on previously existing knowledge about the effect of a particular growth factor on cell morphogenesis. This was done with the dual pursuit of expanding MSCs and causing them to differentiate into the lineage that it was known to favor. Transforming growth factor beta (TGF β), for example, is known to influence cells from the chondrogenic lineage in vivo, promoting initial stages of mesenchymal condensation, prechondrocyte proliferation, production of extracellular matrix and cartilage-specific molecule deposition, while inhibiting terminal differentiation. When applied to MSCs, cells show increased proliferation and a bias towards the chondrogenic lineage [Bonewald et al., J Cell Biochem 1994, 55:350-357; Longobardi L, J Bone Miner Res 2006, 21:626-636].

BMP-3, another member of the transforming growth factor beta family, known to enhance bone differentiation was shown to increase MSC proliferation threefold [Stewart A et al., Cell Physiol 2010, 223:658-666].

Nicotinamide (NA), the amide form of niacin (vitamin B3), is a base-exchange substrate and a potent inhibitor of NAD(+)-dependent enzymes endowed with mono- and poly-ADP-ribosyltransferase activities. ADP-ribosylation is implicated in the

modification of a diverse array of biological processes (Corda D, Di Girolamo M. 2003;22(9):1953-1958; Rankin PW, *et al.*, J Biol Chem. 1989;264:4312-4317; Banasik M. *et al.*, J Biol Chem. 1992;267:1569-1575; Ueda K, Hayaishi O, Annu Rev Biochem. 1985;54:73-100; Smith S. Trends Biochem Sci. 2001;26:174-179; Virág L, Szabó C. Pharm. Reviews. 2002;54:375-429).

WO 07/063545 discloses the use of nicotinamide for the expansion of hematopoietic stem and/or progenitor cell populations.

WO 03/062369 discloses the use of nicotinamide, and other inhibitors of CD38, for the inhibition of differentiation in *ex-vivo* expanding stem and progenitor cells. However, WO 03/062369 does not teach administration of nicotinamide for particular time intervals.

U.S. Patent Application No. 20050260748 teaches isolation and expansion of mesenchymal stem cells with nicotinamide in the presence of a low calcium concentration.

Additional background art includes Farre *et al.*, Growth Factors, 2007 Apr;25(2):71-6.

SUMMARY OF THE INVENTION

According to an aspect of some embodiments of the present invention there is provided a method of culturing mesenchymal stem cells (MSCs) comprising culturing a population of the MSCs in a medium comprising nicotinamide and fibroblast growth factor 4 (FGF4), thereby culturing MSCs.

According to an aspect of some embodiments of the present invention there is provided a method of expanding a population of mesenchymal stem cells, the method comprising culturing a seeded population of mesenchymal stem cells for a period of time sufficient for cell expansion, wherein for at least a portion of the period of time the culturing is effected in a medium devoid of nicotinamide; and for at least a second portion of the period of time, the culturing is effected in a medium comprising nicotinamide and FGF4, thereby generating an expanded population of mesenchymal stem cells.

According to an aspect of some embodiments of the present invention there is provided a method of generating cells useful for transplantation into a subject, the method comprising:

- (a) culturing mesenchymal stem cells according to the methods described
5 herein to generate a population of cultured mesenchymal stem cells;
- (b) contacting the population of cultured mesenchymal stem cells with a differentiating agent, thereby generating cells useful for transplantation into a subject.

According to an aspect of some embodiments of the present invention there is provided a method of generating cells useful for transplantation, the method
10 comprising:

- (a) expanding mesenchymal stem cells according to the methods described herein; and
- (b) contacting the mesenchymal stem cells with a differentiating agent, thereby
generating cells useful for transplantation.

15 According to an aspect of some embodiments of the present invention there is provided an isolated population of mesenchymal stem cells generated according to the methods described herein.

An isolated population of differentiated cells generated according to the methods described herein.

20 According to an aspect of some embodiments of the present invention there is provided a method of treating a disease or disorder, the method comprising transplanting to a subject in need thereof a therapeutically effective amount of the isolated population of cells described herein, thereby treating the disease or disorder.

According to an aspect of some embodiments of the present invention there is
25 provided a cell culture comprising mesenchymal stem cells and a medium which comprises nicotinamide and FGF4.

According to some embodiments of the invention, the medium comprises DMEM.

According to some embodiments of the invention, the medium comprises serum
30 or platelet lysate.

According to some embodiments of the invention, the mesenchymal stem cells are derived from a tissue selected from the group consisting of bone marrow, adipose tissue, placenta and umbilical cord blood.

According to some embodiments of the invention, the nicotinamide is selected
5 from the group consisting of nicotinamide, a nicotinamide analog, a nicotinamide metabolite, a nicotinamide analog metabolite and derivatives thereof.

According to some embodiments of the invention, the culturing is effected on a plastic surface.

According to some embodiments of the invention, the population of MSCs is
10 comprised in a heterogeneous population of cells.

According to some embodiments of the invention, at least 70 % of the heterogeneous population of cells are MSCs.

According to some embodiments of the invention, the calcium concentration of the medium is greater than 1.8 mM.

According to some embodiments of the invention, the culturing is effected for at
15 least 1 week.

According to some embodiments of the invention, the culturing is effected for at least 3 passages.

According to some embodiments of the invention, the concentration of the
20 nicotinamide is 1-20 mM.

According to some embodiments of the invention, the medium is devoid of platelet derived growth factor (PDGF).

According to some embodiments of the invention, the expanding is effected under conditions that do not induce differentiation of the mesenchymal stem cells.

According to some embodiments of the invention, the seeded population of
25 mesenchymal stem cells were seeded in an absence of nicotinamide.

According to some embodiments of the invention, the seeded population of mesenchymal stem cells were seeded in a presence of nicotinamide.

According to some embodiments of the invention, the medium is devoid of
30 platelet derived growth factor (PDGF).

According to some embodiments of the invention, the medium devoid of nicotinamide is devoid of FGF4.

According to some embodiments of the invention, the medium devoid of nicotinamide comprises FGF4

According to some embodiments of the invention, the culturing in the medium comprising nicotinamide is effected prior to the culturing in the medium devoid of
5 nicotinamide.

According to some embodiments of the invention, the culturing in the medium devoid of the nicotinamide is effected prior to the culturing in the medium comprising nicotinamide.

According to some embodiments of the invention, the culturing in the medium
10 comprising nicotinamide is effected for at least one day.

According to some embodiments of the invention, the culturing in the medium comprising nicotinamide is effected for at least one week.

According to some embodiments of the invention, the culturing in the medium devoid of nicotinamide is effected for at least one day.

According to some embodiments of the invention, the culturing in the medium
15 devoid of nicotinamide is effected for at least one week.

According to some embodiments of the invention, the culturing in the medium comprising nicotinamide is effected in a medium comprising calcium, wherein a concentration of the calcium is greater than 1.8 mM.

According to some embodiments of the invention, the culturing in the medium
20 devoid of nicotinamide is effected in a medium comprising calcium, wherein a concentration of the calcium is greater than 1.8 mM.

According to some embodiments of the invention, the differentiation agent comprises a growth factor.

According to some embodiments of the invention, the differentiation agent
25 comprises a polynucleotide which encodes the differentiation agent.

According to some embodiments of the invention, the polynucleotide encodes bone morphogenic protein 2 (BMP2).

According to some embodiments of the invention, the isolated population of
30 mesenchymal stem cells is substantially homogeneous.

According to some embodiments of the invention, at least 40 % of the cells express VCAM1/CD106.

According to some embodiments of the invention, at least 90 % of the cells have a diameter less than 20 μm .

According to some embodiments of the invention, the isolated population of mesenchymal stem cells are less granular than mesenchymal stem cells generated under
5 identical conditions but in an absence of nicotinamide.

According to some embodiments of the invention, less than 30 % of the cells express CD45, more than 95 % of the cells express CD90 and more than 90 % of the cells express CD105 and CD44.

According to some embodiments of the invention, the disease or disorder is
10 selected from the group consisting of a bone or cartilage disease, a neurodegenerative disease, a cardiac disease, a hepatic disease, cancer, nerve damage, autoimmune disease, GvHD, wound healing and tissue regeneration.

According to some embodiments of the invention the mesenchymal stem cells cultured with nicotinamide and/or nicotinamide and FGF4 secrete increased levels of
15 growth factors, and reduced levels of pro-inflammatory factors into the medium.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable
20 methods and materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this
25 invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. In case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

BRIEF DESCRIPTION OF THE DRAWINGS

The patent or application file contains at least one drawing executed in color. Copies of this patent or patent application publication with color drawing(s) will be provided by the Office upon request and payment of the necessary fee.

5 Some embodiments of the invention are herein described, by way of example only, with reference to the accompanying drawings and images. With specific reference now to the drawings in detail, it is stressed that the particulars shown are by way of example and for purposes of illustrative discussion of embodiments of the invention. In this regard, the description taken with the drawings makes apparent to those skilled in
10 the art how embodiments of the invention may be practiced.

In the drawings:

FIG. 1 is a bar graph illustrating that basic fibroblast growth factor (bFGF) has a negative effect on the ability of nicotinamide to increase proliferation of mesenchymal stem cells.

15 FIGs. 2A-B illustrate that heparin-binding EGF-like growth factor (HB-EGF) has a negative effect on the ability of nicotinamide to increase proliferation of two different batches of mesenchymal stem cells.

FIGs. 3A-D are bar graphs illustrating the synergistic activity of nicotinamide (NAM) and FGF4 on expansion of mesenchymal stem cells. Four different batches of
20 MSC cultures were treated with FGF4 (50ng/ml), NAM (5mM) or a combination of FGF4+NAM. Cumulative cell counts at the indicated passages are shown.

FIGs. 4A-B are graphs illustrating that nicotinamide (NAM) preserves the undifferentiated state of MSCs cultured with FGF4. Two different batches of MSC cultures were treated with FGF4 (50 ng/ml), NAM (5 mM) or a combination of
25 FGF4+NAM. Cell size was analyzed by Cedex cell counter.

FIGs. 5A-D are graphs illustrating that cells expanded with a combination of NAM+FGF4 are undifferentiated MSCs (CD105+CD45-). Four different batches of MSC cultures were treated with FGF4 (50ng/ml), NAM (5 mM) or a combination of FGF4+NAM. Percent of MSC (CD105+CD45-) was analyzed by FACS.

30 FIGs. 6A-D are bar graphs illustrating inconsistent results obtained following expansion of MSC with NAM+PDGF-BB. Four different batches of MSC cultures

were treated with PDGF-BB (50 ng/ml), NAM (5 mM) or a combination of PDGF-BB+NAM. Cumulative cell counts at the indicated passages are shown.

FIGs. 7A-D are graphs illustrating that MSC cultures treated with PDGF-BB or a combination of PDGF-BB + NAM comprise a higher fraction of cells other than MSCs that contaminates the cultures as compared to MSC cultured in the absence of PDGF-BB. Four different batches of MSC cultures were treated with PDGF-BB (50 ng/ml), NAM (5 mM) or a combination of PDGF-BB+NAM. Percent of MSC (CD105+CD45-) was analyzed by FACS.

FIGs. 8A-B are bar graphs illustrating a consistent synergistic effect between NAM and FGF4 in contrast to the absence of a synergistic or additive effect between FGF4 and PDGF-BB. Further, a combination of NAM, FGF4 and PDGF-BB had an adverse effect on MSC expansion. MSC cultures were treated with PDGF-BB (50ng/ml), FGF4 (50ng/ml) and NAM (5mM) or a combination of two or three factors, as indicated. Cumulative cell counts at the indicated passages are shown.

FIGs. 9A-B are graphs illustrating that PDGF-BB supports expansion of cells other than MSCs in MSC cultures. This effect is not alleviated by NAM and/or FGF4. MSC cultures were treated with PDGF-BB (50 ng/ml), FGF4 (50 ng/ml) and NAM (5 mM) or a combination of two or three factors, as indicated. Percent of MSC (CD105+CD45-) was analyzed by FACS.

FIGs. 10A-H are photographs of day 34 MSC cultures illustrating that PDGF-BB supports expansion of cells other than MSC in MSC cultures. This effect is not alleviated by NAM and/or FGF4. MSC cultures were treated with PDGF-BB (50 ng/ml), FGF4 (50 ng/ml) and NAM (5 mM) or a combination of two or three factors, as indicated.

FIG. 11 is a bar graph illustrating % of BM derived adherent cells expressing mesenchymal stem cells markers in culture seeded +/- NAM, prior to the first passage. Mononuclear cells were isolated from bone marrow using Ficoll and the "plastic adherence" method in the presence or absence of Nicotinamide. Non-adherent cells were washed away 3-4 days later and the media was replaced every 3-4 days. FACS analysis was performed in order to obtain expression levels of surface molecules prior to the first passage (8 days post-seeding).

FIGs. 12A-C are graphs illustrating phenotypic characterization of adipose tissue derived mesenchymal stem cells after six passages in different concentrations of nicotinamide.

FIG. 13 is a bar graph illustrating phenotypic characterization of bone marrow
5 derived mesenchymal stem cells following the first passage of cultures treated +/- different concentration of nicotinamide. Mononuclear cells were isolated from bone marrow using Ficoll and the "plastic adherence" method. Non-adherent cells were washed away 3-4 days later and the media was replaced every 3-4 days. FACS analysis was performed in order to obtain expression levels of surface molecules following the
10 first passage (8 days post-seeding).

FIG. 14 is a bar graph illustrating the effect of different concentrations of nicotinamide (added at passage 3, and at each subsequent passage) on the number of MSC at passage 6. Nicotinamide substantially improved adipose derived mesenchymal stem cell expansion in culture.

15 FIG. 15 is a graph illustrating the effect of nicotinamide on bone marrow derived mesenchymal stem cell expansion. Nicotinamide was added from the initiation of the culture and at each subsequent passage.

FIG. 16 is a graph illustrating the effect of different concentrations of nicotinamide on adipose derived mesenchymal stem cell expansion. Nicotinamide was
20 added from passage 3 and at each subsequent passage.

FIGs. 17A-B illustrate that mesenchymal stem cells cultured in nicotinamide, expand more rapidly than mesenchymal stem cells cultured under identical conditions, but in the absence of nicotinamide.

FIG. 18 is a graph illustrating the effect of nicotinamide on the cumulative cell
25 count of mesenchymal stem cells cultured in a large batch.

FIG. 19 is a bar graph illustrating the results of one of two experiments performed illustrating that the effect of nicotinamide on mesenchymal stem cell proliferation is not dependent on the particular batch of serum used.

FIGs. 20A-C are graphs and plots illustrating the beneficial effect of culturing in
30 the presence of nicotinamide on cell size and granularity. Figure 20C shows that cells grown in the presence of nicotinamide are smaller and less granular (most of the cells are in the red circle), as oppose to cells grown without nicotinamide which are larger

and more granular (black circle). For Figure 20A, the concentration of nicotinamide used was 5 mM.

FIGs. 21A-B are graphs and plots illustrating that mesenchymal stem cells grown in the presence of nicotinamide are less granular than mesenchymal stem cells grown in the absence of nicotinamide under identical conditions.

FIG. 22 is a graph illustrating that culturing MSCs in the presence of nicotinamide (5 mM) increases the cumulative CFU-F count.

FIGs. 23A-D are photographs illustrating that nicotinamide reduces the amount of senescence of mesenchymal stem cells. Bone marrow-derived mesenchymal stem cells were cultured for 5 passages +/- 5mM NAM. The cells were fixated and X-Gal staining was performed to detect senescent cells (blue stain).

FIGs. 24A-B are bar graphs illustrating that nicotinamide modulates expression of surface markers on mesenchymal stem cells – VCAM1/CD106 (Figure 24A) and CD54 (Figure 24B). Note the enhanced expression of VCAM1/CD106 and reduced expression of CD54 in cells grown in the presence of nicotinamide;

FIGs. 25A-B are photographs illustrating results of an *in vitro* wound healing assay which was performed on MSCs cultured with (Figure 25B) or without (Figure 25A) nicotinamide at passage 3. Wound healing was observed 4 days post wound formation.

FIG. 26 is a graph illustrating the effect of nicotinamide on bone marrow derived mesenchymal stem cell Doubling Time. Nicotinamide was added from the initiation of the culture and at each subsequent passage.

FIG. 27 is a graph illustrating the effect of nicotinamide, with and without FGF4 on bone marrow-derived MSC proliferation, through 5 passages of culture. Bone marrow derived mesenchymal stem cells were isolated using Ficoll and plastic adherence method, and cultured for several passages with fetal bovine serum. -NAM - FGF4= controls (light blue circles), -NAM +FGF4= culture with 50ng/ml FGF4 (dark blue circles), +NAM -FGF4= culture with 5mM NAM (pink circles), +NAM +FGF4= culture with 50ng/ml FGF4 and 5mM NAM (red circles). Note the synergic effect of Nicotinamide and FGF4 added together throughout all passages of MSC proliferation;

FIG. 28 is a bar graph illustrating enhancement of Hepatocyte Growth Factor (HGF) content of conditioned medium from nicotinamide and FGF4-treated MSC

cultures. Bone marrow mesenchymal stem cells were isolated using Ficoll and plastic adherence method, and cultured for several passages with fetal bovine serum, with added nicotinamide and FGF4 (+NAM+FGF4), with added nicotinamide (+NAM-FGF4) and without added nicotinamide or FGF (-NAM-FGF4). Twenty four hours before passage 4, the media was changed and fresh media without fetal bovine serum or FGF4 was added. The cultured media from passage 4 cultures was collected and assayed for HGF content by ELISA. -NAM, -FGF4= control; +NAM -FGF4 = 5mM NAM, +NAM +FGF4 = 5mM NAM + 50ng/ml FGF4. Note the significant effect of combined FGF4 and nicotinamide on HGF secretion;

FIG. 29 is a bar graph illustrating enhancement of Transforming Growth Factor- β (TGF β) content of conditioned medium from nicotinamide and FGF4-treated MSC cultures. Bone marrow mesenchymal stem cells were isolated and cultured as in FIG. 28 above. Cultured media was changed to medium without fetal bovine serum or FGF4 24 hours prior to passage 4, collected from the passage 4 cultures and assayed for TGF β content by ELISA. -NAM, -FGF4= control; +NAM -FGF4 = 5mM NAM, +NAM +FGF4 = 5mM NAM + 50ng/ml FGF4. Note the significant effect of combined FGF4 and nicotinamide on TGF β secretion;

FIG. 30 is a bar graph illustrating enhancement of Keratinocyte Growth Factor (KGF) content of conditioned medium from nicotinamide and FGF4-treated MSC cultures. Bone marrow mesenchymal stem cells were isolated and cultured as in FIG. 28 above. Cultured media was changed to medium without fetal bovine serum or FGF4 24 hours prior to passage 4, collected from the passage 4 cultures and assayed for KGF content by ELISA. -NAM, -FGF4=control; +NAM -FGF4 = 5mM NAM, +NAM +FGF4 = 5mM NAM + 50ng/ml FGF4. Note the significant effect of combined FGF4 and nicotinamide on KGF secretion;

FIG. 31 is a bar graph illustrating reduction of cytokine IL-6 (IL-6) content of conditioned medium from nicotinamide and FGF4-treated MSC cultures. Bone marrow mesenchymal stem cells were isolated and cultured as in FIG. 28 above. Cultured media was changed to medium without fetal bovine serum or FGF4 24 hours prior to passage 4, collected from the passage 4 cultures and assayed for IL-6 content by ELISA. -NAM, -FGF4= control; +NAM -FGF4 = 5mM NAM, +NAM +FGF4 = 5mM

NAM + 50ng/ml FGF4. Note the significant reduction by combined FGF4 and nicotinamide on IL-6 secretion;

FIG. 32 is a graph illustrating the effect of nicotinamide, with and without FGF4 on adipose-derived MSC proliferation, through 4 passages of culture. Adipose derived mesenchymal stem cells were isolated using collagenase digestion and plastic adherence method, and cultured for several passages with fetal bovine serum. -NAM -FGF4= controls (blue diamonds), +NAM -FGF4= culture with 5mM NAM (red squares), +NAM +FGF4= culture with 50ng/ml FGF4 and 5mM NAM (green triangles). Note the synergic effect of Nicotinamide and FGF4 added together on adipose-derived MSC proliferation;

FIG. 33 is a graph detailing the effect of nicotinamide, with and without FGF4 on nucleated cell proliferation in adipose-derived MSC proliferation at passage 4. Adipose-derived mesenchymal stem cells were isolated and cultured as described in FIG. 32. -NAM -FGF4= controls, +NAM -FGF4= culture with 5mM/ml NAM, +NAM +FGF4= culture with 50ng/ml FGF4 and 5mM/ml NAM. Note the synergic effect of nicotinamide and FGF4 together on proliferation of total nucleated cells in the culture;

FIG. 34 is a bar graph illustrating the beneficial effect of culturing adipose derived MSCs in the presence of nicotinamide and FGF4 on the size of the cultured mesenchymal stem cells. Adipose-derived mesenchymal stem cells were isolated and cultured as described in FIG. 32. Cell size was analyzed by Cedex cell counter. -NAM -FGF4= controls, +NAM -FGF4= culture with 5mM/ml NAM, +NAM +FGF4= culture with 50ng/ml FGF4 and 5mM/ml NAM. Note that smaller size of MSC cells grown in the presence of nicotinamide, and even smaller MSCs grown with nicotinamide and FGF4;

FIG. 35 is a bar graph showing the effect of nicotinamide with and without FGF4 on differentiation of ex-vivo expanded hematopoietic cells. Umbilical cord-derived early progenitor (CD 133+) hematopoietic cells were isolated using CD133 microbeads and CliniMACS (Milentyi, Inc), and cultured for 3 weeks in MEM α supplemented with 50ng/ml early acting cytokines and fetal bovine serum, $\pm$ 2.5 or 5mM nicotinamide (NAM), $\pm$ 10, 50 or 200ng/ml FGF4. After three weeks culture CD38-CD133+ cells were stained and counted by FACS. Column 1= Control: -NAM, -FGF4, Column 2= + 2.5mM NAM, Column 3= + 5 mM NAM, Column 4= + 10 ng/ml

FGF4, Column 5= +50 ng/ml FGF4, Column 6= +200 ng/ml FGF4, Column 7= +2.5 mM NAM, +10 ng/ml FGF4, Column 8= +2.5 mM NAM, +50 ng/ml FGF4, Column 9= +2.5 mM NAM, +200 ng/ml FGF4, Column 10= +5 mM NAM, +10 ng/ml FGF4, Column 11= +5 mM NAM, +50 ng/ml FGF4, Column 12= +5 mM NAM, +200 ng/ml FGF4.

Note the significantly greater fraction of undifferentiated early progenitors (CD38- CD133+) in the nicotinamide-treated cultures (columns 2 and 3), absence of any significant effect of FGF4 alone (columns 4-6) and absence of any significant effect of FGF4 on the nicotinamide-mediated inhibition of hematopoietic progenitor cell differentiation (columns 7-12);

FIG. 36 is a bar graph showing the effect of nicotinamide with and without FGF4 on differentiation of ex-vivo expanded hematopoietic cells. Umbilical cord-derived progenitor (CD 133+) hematopoietic cells were isolated and cultured, with and without nicotinamide and FGF4, as in FIG. 35. After three weeks culture CD38+ cells were stained and counted by FACS. Columns 1-12 as in FIG. 35. Note the significant inhibition of differentiation (CD38+ cells) in the nicotinamide-treated cultures (columns 2 and 3), absence of any significant effect of FGF4 alone on differentiation (columns 4-6) and absence of any significant effect of FGF4 on the nicotinamide-mediated inhibition of hematopoietic progenitor cell differentiation (columns 7-12);

FIG. 37 is a bar graph showing the effect of nicotinamide with and without FGF4 on myeloid lineage differentiation of ex-vivo expanded hematopoietic cells. Umbilical cord-derived progenitor (CD 133+) hematopoietic cells were isolated and cultured, with and without nicotinamide and FGF4, as in FIG. 35. After three weeks culture myeloid lineage differentiated (CD33+) cells were stained and counted by FACS. Columns 1-12 as in FIG. 35. Note the significant inhibition of myeloid lineage differentiation (CD33+ cells) in the nicotinamide-treated cultures (columns 2 and 3), moderate enhancement of myeloid lineage differentiation by FGF4 alone (columns 4-6) and absence of any significant effect of FGF4 on the nicotinamide-mediated inhibition of myeloid lineage hematopoietic cell differentiation (columns 7-12);

FIG. 38 is a bar graph showing the effect of nicotinamide with and without FGF4 on lymphoid lineage differentiation of ex-vivo expanded hematopoietic cells. Umbilical cord-derived progenitor (CD 133+) hematopoietic cells were isolated and

cultured, with and without nicotinamide and FGF4, as in FIG. 35. After three weeks culture lymphoid lineage differentiated (CD19+) cells were stained and counted by FACS. Columns 1-12 as in FIG. 35. Note the striking inhibition of lymphoid lineage differentiation (CD19+ cells) in the nicotinamide-treated cultures (columns 2 and 3),
5 absence of any significant effect on lymphoid lineage differentiation by FGF4 alone (columns 4-6) and absence of any significant effect of FGF4 on the nicotinamide-mediated inhibition of lymphoid lineage hematopoietic cell differentiation (columns 7-12);

10 DESCRIPTION OF SPECIFIC EMBODIMENTS OF THE INVENTION

The present invention, in some embodiments thereof, relates to methods of expanding mesenchymal stem cells and cell populations generated thereby.

Before explaining at least one embodiment of the invention in detail, it is to be understood that the invention is not necessarily limited in its application to the details set
15 forth in the following description or exemplified by the Examples. The invention is capable of other embodiments or of being practiced or carried out in various ways.

The multipotent character of mesenchymal stem cells (MSCs) make these cells an attractive therapeutic tool and candidate for transplantation, capable of playing a role in a wide range of clinical applications in the context of both cell and gene therapy
20 strategies. For example, mesenchymal stem cells may be used to enhance hematopoietic engraftment post-transplantation, to aid in tissue re-generation, to promote wound healing and to correct for a myriad of other inherited and acquired disorders. Efficient mesenchymal stem cell expansion protocols that do not have deleterious effects on the differentiation potential and target tissue engraftment potential of the cells are crucial to
25 the success of any of these strategies.

In addition, MSCs are attractive for clinical therapy in regenerative medicine and inflammatory conditions due to their ability to differentiate, provide trophic support, and modulate the innate immune response. The therapeutic potential of MSC is being tested in multiple clinical trials for indications such as bone and cartilage repair,
30 cardiac regeneration, critical limb ischemia, acute ischemic conditions, diabetes, Crohn's disease and graft vs host disease.

MSCs are immune-privileged and can be transplanted without the need for HLA matching between the donor and the recipient and therefore can be manufactured at large scale and marketed as an off-the-shelf cell product. The success of large scale batch production from one donor is highly dependent on donor and serum selection, the potential of seeded cells for prolonged expansion in culture and the duration of the manufacturing. Even though MSC multiply relatively easily in vitro, their proliferative potential is continuously decreased and their doubling time increases during culture. As a result, the successful manufacturing for commercialization of large batches of homogenous MSCs from one donor remains a challenge.

Whilst studying the effect of growth factors on MSC expansion, the present inventors found that growth factors such as basic FGF (bFGF), HB-EGF or platelet derived growth factor (PDGF) have a non-reproducible or even negative effect when cultured in the presence of nicotinamide on mesenchymal stem cell proliferation (Figures 1, 2, 6).

In sharp contrast, FGF4 surprisingly demonstrated a reproducible, synergistic activity together with nicotinamide on mesenchymal stem cell expansion/proliferation

As illustrated in Figures 3A-D, the present inventors demonstrated that nicotinamide potentiates the effect of FGF4 on the proliferation of mesenchymal stem cells.

In addition, the present inventors demonstrated an unexpected effect of nicotinamide on cell size of mesenchymal stem cells cultured with FGF4.

As illustrated in Figures 4A-B, MSCs generated by culturing in nicotinamide and FGF4 are smaller than mesenchymal stem cells cultured according to identical methods, but in the presence of FGF4 alone and similar to MSC cultured with nicotinamide alone. For example, between days 10-32, the mesenchymal stem cells which are cultured in nicotinamide and FGF4 are less than about 20 μm in diameter, whereas cells grown in the presence of FGF4, but the absence of nicotinamide are greater than 20 μm in diameter. Thus, nicotinamide imposed an undifferentiated state on MSC cultured with FGF4.

Whilst further reducing the present invention to practice, the present inventors demonstrated that percent of cells expressing the MSC marker, CD105+CD45- is preserved in cultures treated with nicotinamide and FGF4 (Figures 5A-D).

Further, the present inventors have found that the use of nicotinamide during particular stages of the selection and expansion protocol was advantageous to mesenchymal stem cell populations. Thus, for example, seeding mesenchymal stem cells in the presence of nicotinamide and high calcium concentrations increased their seeding efficacy, as noted by analyzing marker phenotype of the cells (Figures 11-13). Mesenchymal stem cells could be successfully expanded for at least six passages in the presence of nicotinamide without induction of differentiation, as illustrated by the surface marker composition of the cells (Figures 12A-C). Further, it was shown that nicotinamide promoted expansion of a more homogeneous, less granular population of MSCs (Figures 20A-C and 21A-B).

The present inventors experimentally showed that MSCs cultured with nicotinamide proliferate more rapidly and as a result, their doubling time (see Figure 26) is decreased and the cultures reach confluence in a substantially shorter period of time (Figures 14-17, 26 and 27). The proliferative effect of nicotinamide was also demonstrated in large cultures of MSCs (Figure 18). Further, the effect was not restricted to selected batches of serum (Figure 19), a substantial advantage for the manufacturing of larger batches of MSCs. Yet further, the present inventors have shown that the proliferative effects of nicotinamide in combination with fibroblast growth factor 4 (FGF4) are not observed for stem cells of non-mesenchymal origin, for example, hematopoietic stem or progenitor cells (e.g, CD133+) (see Example 10, and Figures 35-38 herein). Indeed, FGF4 alone or in combination with nicotinamide was without any effect on proliferation or differentiation of ex-vivo cultured hematopoietic stem or progenitor cells (see Figures 35-38, lanes 1 and 4-12).

Thus, according to one aspect of the present invention there is provided a method of culturing mesenchymal stem cells (MSCs) comprising culturing a population of the MSCs in a medium comprising nicotinamide and fibroblast growth factor 4 (FGF4).

Yet further, the present inventors have now found that culturing a mixed population of mesenchymal stem cells in the presence of nicotinamide enhances the mesenchymal stem cell phenotype, such that subsequent selection or pre-selection with a mesenchymal stem cell marker provides for a more homogeneous population of mesenchymal stem cells, thereby providing a method for obtaining enriched populations

of mesenchymal stem cell subsets. This was substantiated by the present inventors when they showed that culturing MSCs in the presence of nicotinamide increases expression of a particular adhesion molecule - Vascular cell adhesion protein 1 (VCAM1/CD106; see Figure 24A). Conversely, the present inventors have shown that culturing MSCs in the presence of nicotinamide decreased expression of a marker for cell senescence (CD54; see Figure 24B) thereby providing a method for obtaining enriched populations of mesenchymal stem cell by depleting the cell population for non-relevant cells.

The use of a selection or sorting step further enhances the stringency of sorting and selection specificity for MSCs and furthermore potentially reduces possible contamination from the starting material.

Thus, according to one aspect of the present invention there is provided a method of isolating mesenchymal stem cells (MSCs) from a mixed population of cells, comprising:

(a) culturing the mixed population of cells in a medium comprising nicotinamide; and

(b) selecting cells based on the expression of a cell surface molecule from the mixed population of cells, thereby selecting MSCs from a mixed population of cells.

The term "mesenchymal stem cell" or "MSC" is used interchangeably for adult cells which are not terminally differentiated, which can divide to yield cells that are either stem cells, or which, irreversibly differentiate to give rise to cells of a mesenchymal cell lineage, e.g., adipose, osseous, cartilaginous, elastic and fibrous connective tissues, myoblasts) as well as to tissues other than those originating in the embryonic mesoderm (e.g., neural cells) depending upon various influences from bioactive factors such as cytokines.

MSC cultures utilized by some embodiments of the invention preferably include three groups of cells which are defined by their morphological features: small and agranular cells (referred to as RS-1, hereinbelow), small and granular cells (referred to as RS-2, hereinbelow) and large and moderately granular cells (referred to as mature MSCs, hereinbelow). The presence and concentration of such cells in culture can be assayed by identifying a presence or absence of various cell surface markers, by using, for example, immunofluorescence, *in situ* hybridization, and activity assays.

When MSCs are cultured under the culturing conditions of some embodiments of the invention they exhibit negative staining for the hematopoietic stem cell markers CD34, CD11B, CD43 and CD45. A small fraction of cells (less than 10 %) may be dimly positive for CD31 and/or CD38 markers. In addition, mature MSCs may be dimly positive for the hematopoietic stem cell marker, CD117 (c-Kit), moderately positive for the osteogenic MSCs marker, Stro-1 [Simmons, P. J. & Torok-Storb, B. (1991). Blood 78, 5562] and positive for the thymocytes and peripheral T lymphocytes marker, CD90 (Thy-1). On the other hand, the RS-1 cells are negative for the CD117 and Stro1 markers and are dimly positive for the CD90 marker, and the RS-2 cells are negative for all of these markers.

Mesenchymal cells cultured under the culturing conditions of some embodiments of the invention can secrete biologically active factors into the medium. The present inventors have observed that medium collected from mesenchymal cells cultured with nicotinamide comprises elevated levels of growth factors and cytokines (e.g. hepatocyte growth factor, keratinocyte growth factor, transforming growth factor beta) and reduced levels of pro-inflammatory factors (e.g. IL6) (see Example 8 and Figures 28-31). Addition of FGF4 to the medium further increased the levels of growth factors in the medium, whilst further reducing the levels of IL6 in the culture medium. Isolated culture medium was observed to have a strong anti-inflammatory effect, as well as enhancing proliferation of cells in culture. Thus, the mesenchymal cells cultured under the culturing conditions secrete biologically active factors having anti-inflammatory and cell proliferative-enhancing activity into the medium.

According to a preferred embodiment of this aspect of the present invention, the mesenchymal stem cells are human.

According to another embodiment of this aspect of the present invention, the mesenchymal stem cells are isolated from newborn humans.

Mesenchymal stem cells may be isolated from various tissues including but not limited to bone marrow, peripheral blood, blood, placenta (e.g. fetal side of the placenta), cord blood, umbilical cord, amniotic fluid, placenta and from adipose tissue.

A method of isolating mesenchymal stem cells from peripheral blood is described by Kassis et al [Bone Marrow Transplant. 2006 May; 37(10):967-76]. A method of isolating mesenchymal stem cells from placental tissue is described by Zhang

et al [Chinese Medical Journal, 2004, 117 (6):882-887]. Methods of isolating and culturing adipose tissue, placental and cord blood mesenchymal stem cells are described by Kern et al [Stem Cells, 2006; 24:1294-1301].

Bone marrow can be isolated from the iliac crest of an individual by aspiration.

5 Low-density BM mononuclear cells (BMMNC) may be separated by a FICOL-PAQUE density gradient or by elimination of red blood cells using Hetastarch (hydroxyethyl starch). Preferably, mesenchymal stem cell cultures are generated by diluting BM aspirates (usually 20 ml) with equal volumes of Hank's balanced salt solution (HBSS; GIBCO Laboratories, Grand Island, NY, USA) and layering the diluted cells over about
10 10 ml of a Ficoll column (Ficoll-Paque; Pharmacia, Piscataway, NJ, USA). Following 30 minutes of centrifugation at 2,500 x g, the mononuclear cell layer is removed from the interface and suspended in HBSS. Cells are then centrifuged at 1,500 x g for 15 minutes and resuspended in a complete medium (MEM, α medium without deoxyribonucleotides or ribonucleotides; GIBCO); 20 % fetal calf serum (FCS) derived
15 from a lot selected for rapid growth of MSCs (Atlanta Biologicals, Norcross, GA); 100 units/ml penicillin (GIBCO), 100 μ g/ml streptomycin (GIBCO); and 2 mM L-glutamine (GIBCO).

Adipose tissue-derived MSCs can be obtained from any fat-containing tissue, for example, from epididymal fat or by liposuction and mononuclear cells can be isolated
20 manually by removal of the fat and fat cells, or using the Celution System (Cytori Therapeutics) following the same procedure as described above for preparation of MSCs.

As mentioned, the method is effected by culturing (i.e. *ex vivo* or *in vitro*) the mesenchymal stem cells in a medium comprising nicotinamide and FGF4.

25 According to this aspect of the present invention, the cells are cultured under conditions that do not induce differentiation (e.g. in the absence of differentiation factors or in the presence of a non-differentiating amount of differentiating factors).

The present invention contemplates directly culturing mesenchymal stem cells following isolation from their source or culturing populations of cells that have been
30 pre-selected for mesenchymal stem cells. Thus, the present invention contemplates culturing both heterogeneous populations of cells which comprise the MSCs and more homogeneous populations of cells, which have been enriched for MSCs, wherein more

than 70 %, more than 80 %, more than 90 % or more than 95 %, more than 98 % thereof are MSCs. Also, contemplated is the enriching for MSCs concomitant with the culturing as further described herein below.

It will be appreciated that the composition of the heterogeneous population of cells will be dependent on the source of the cells. Thus, for example, if the placenta is selected as the cell source, the heterogeneous population of cells will comprise placental cells as well as mesenchymal stem cells. If the bone marrow is selected as the cell source, the heterogeneous population of cells will comprise blood cells. However, as shown in Example 10, according to some embodiments of the present invention, culturing the mesenchymal stem cells under the culturing conditions of some embodiments of the invention (e.g, nicotinamide and FGF4 in combination) results in selective expansion of mesenchymal stem cell populations, while not having a concomitant proliferative effect on non-mesenchymal stem cell populations.

According to one method, the population of cells are cultured (*in vitro* or *ex vivo*) on polystyrene plastic surfaces (e.g. in a flask) so as to enrich for mesenchymal stem cells by removing non-adherent cells (i.e. non-mesenchymal stem cells). This method of enriching for MSCs may be effected prior to the culturing in nicotinamide and FGF4, concomitant with the culturing in nicotinamide and FGF4 and/or following the culturing in nicotinamide and FGF4.

Other methods of selecting for MSCs are known in the art including for example positive selection against mesenchymal stem cell markers and/or negative selection against hematopoietic stem and progenitor markers such as CD34, CD133, CD8, etc. Methods of determining protein cell-surface expression are well known in the art. Examples include immunological methods, such as, FACS analysis as well as biochemical methods (cell-surface labeling, e.g., radioactive, fluorescence, avidin-biotin).

It will be appreciated that a selecting stage may also be performed following the culturing in nicotinamide and FGF4. This may be effected as well as a preselection stage or instead of a preselection stage.

As used herein "nicotinamide" refers to nicotinamide as well as to products that are derived from nicotinamide, analogs thereof and metabolites of nicotinamide or nicotinamide analogs, such as, for example, NAD, NADH and NADPH.

As used herein, the phrase “nicotinamide analog” refers to any molecule that is known to act similarly to nicotinamide. Representative examples of nicotinamide analogs include, without limitation, benzamide, nicotinethioamide (the thiol analog of nicotinamide), nicotinic acid, α -amino-3-indolepropionic acid, and inhibitors of sirtuin
5 family of histone/protein deacetylases.

Examples of nitotinamide analog derivatives include, but are not limited to substituted benzamides, substituted nicotinamides and nicotinethioamides and N-substituted nicotinamides and nicotinthioamides.

In a particular embodiment, the nicotinamide is supplied at a concentration of at least about 1 mM to 20 mM. In other embodiment, the nicotinamide concentration is
10 supplied at a concentration of at least about 1 mM to 10 mM, e.g. about 2.5 mM, about 5 mM, about 7.5 mM.

Fibroblast growth factor 4, the FGF4 (map locus 11q13.3) gene product, FGF-4/HBGF-4/KFGF, is a 176 AA long protein derived by cleavage of the N-terminal 30
15 AAs of the precursor protein. FGF-4 contains a single N-linked glycosylation site. Unglycosylated FGF-4 is cleaved into two NH₂-terminally truncated peptides (13 and 15 kDa) that are more active with higher heparin affinity than wild-type protein.

According to a particular embodiment, the FGF4 is human FGF4.

Recombinant FGF4 protein is commercially available (e.g. from Sigma Aldrich,
20 where it is produced in baculovirus and cleaved at the N-terminal to yield a 148 AA protein; or from Invitrogen where it is produced in E.coli).

In a particular embodiment, the FGF4 is supplied to the culture at a concentration of at least about 1-1000 ng/ml. In other embodiment, the FGF4 concentration is supplied at a concentration of at least about 10-200 ng/ml, 10-100
25 ng/ml, e.g. about 50 ng/ml.

According to a particular embodiment, the culturing medium comprising both nicotinamide and FGF4 is devoid of additional growth factors such as PDGF, HB-EGF or bFGF (FGF2).

It will be appreciated that when referring to a medium being devoid of a
30 particular component, the present invention contemplates that the medium comprises this component, but at a concentration which is below its minimal activity. Thus, for example, certain mediums may comprise trace amounts of the above described growth

factors, however, the methods of the present invention relate to a medium being devoid of exogenously added growth factor beyond what is included in a commercial medium's formula, or that resulting from overall adjustment of medium component concentrations. Thus, according to a particular embodiment, the medium which comprises nicotinamide and FGF4 may comprise any one of the above mentioned additional growth factors but at a concentration less than 1 ng/ml.

A typical cell medium to which the nicotinamide and FGF4 may be added is Dulbecco's modified MEM (DMEM). Alternatively, the cell medium may be Ham's F12. Other contemplated mediums include HEM RPMI, F-12, and the like.

It will be noted that many of the culture media contain nicotinamide as a vitamin supplement for example, MEM α (8.19 μ M nicotinamide), RPMI (8.19 μ M nicotinamide), DMEM (32.78 μ M nicotinamide) and Glasgow's medium (16.39 μ M nicotinamide), however, the methods of the present invention relate to exogenously added nicotinamide supplementing any nicotinamide and/or nicotinamide moiety included the medium's formula, or that resulting from overall adjustment of medium component concentrations.

In an embodiment of the invention, the cell culture medium has a high calcium concentration of more than about 1.8 mM, more than about 2 mM, or more than about 5 mM. It will be appreciated that the calcium concentration is calculated as the total calcium concentration including that already present in the culture medium.

Thus, for example, if the medium is Dulbecco's modified MEM (DMEM) (which already has a calcium ion concentration of about 1.8 mM), no additional calcium needs to be added. If the cell medium is Ham's F12 which has a calcium ion concentration of about 0.9 mM, additional calcium should be added so the total calcium concentration is above 1.8 mM. In one embodiment, the source of the additional calcium may be serum.

During the culturing, the medium can contain supplements required for cellular metabolism such as glutamine and other amino acids, vitamins, minerals and useful proteins such as transferrin, and the like. The medium may also contain antibiotics to prevent contamination with yeast, bacteria, and fungi, such as penicillin, streptomycin, gentamicin, and the like. If cells are to be cultured, conditions should be close to

physiological conditions (preferably, a pH of about 6 to about 8, and a temperature of about 30 ° C to about 40 ° C).

Normoxia or hypoxia conditions are also contemplated.

According to one embodiment, the culture medium is devoid of serum (i.e. serum free medium) and comprises serum replacements including, but not limited to platelet lysate (during seeding and/or expansion).

According to still another embodiment the medium comprises about 10 % fetal bovine serum. Human serum is also contemplated.

The culturing according to this aspect of the present invention may be effected for a limited amount of time, such that no expansion takes place (e.g. during the seeding stage only) or may be effected for longer periods of time so as to allow for mesenchymal stem cell expansion (i.e. cell propagation), thereby obtaining increased quantities thereof

For each round of propagation, adherent cells may be harvested using trypsin/EDTA or by cell scraping, and dissociated by passage through a narrow Pasteur plastic pipette, and preferably replated at a density of about 100 to about 10,000 cells/cm².

According to this aspect of the present invention, a period of time sufficient for cell expansion may be taken to mean the length of time required for at least one cell to divide.

According to one embodiment, the culturing is effected for at least one day, at least two days, at least three days, at least four days, at least five days, at least six days, at least one week, at least two weeks, at least three weeks, at least four weeks or at least five weeks.

According to another embodiment, the culturing is not effected for more than ten weeks.

According to still another embodiment, the cells are allowed to expand for at least two population doublings, at least four population doublings, at least six population doublings, at least eight population doublings, at least ten population doublings, at least 15 population doublings, at least 20 population doublings, at least 25 population doublings, at least 30 population doublings, at least 35 population doublings, at least 40 population doublings, or at least 45 population doublings.

According to another embodiment, the cells are not allowed to expand for more than 50 population doublings.

The present invention contemplates additional methods of mesenchymal stem cell expansion as well as (or instead of) culturing in nicotinamide and FGF4.

5 Since the present inventors have found that when at least a portion of the time of the expansion process is effected in the presence of nicotinamide, increased numbers of mesenchymal stem cells are obtained, preferably additional methods of expansion include culturing in the presence of nicotinamide.

Thus, according to another aspect of the present invention there is provided a
10 method of expanding a population of mesenchymal stem cells, the method comprising culturing a seeded population of mesenchymal stem cells for a period of time sufficient for cell expansion, wherein for at least a portion of the period of time the culturing is effected in a medium devoid of nicotinamide; and for at least a second portion of the period of time, the culturing is effected in a medium comprising nicotinamide and
15 FGF4, thereby generating an expanded population of mesenchymal stem cells.

The term "expanding" as used herein refers to increasing the number of cells in the cell population due to cell replication.

According to this aspect of the present invention, the cells are expanded under conditions that do not induce differentiation (e.g. in the absence of differentiation
20 factors).

The seeded population of undifferentiated mesenchymal stem cells may be a heterogeneous population of cells or a purified population of mesenchymal stem cells, as further described herein above.

As mentioned, a medium being devoid of nicotinamide refers to a medium
25 comprising less than the minimal effective amount of nicotinamide (e.g. less than 0.5 mM, or more preferably less than 0.05 mM). Thus mediums comprising trace amounts of nicotinamide (as described herein above) may be used for this aspect of the present invention. Thus, according to a particular embodiment, the medium without exogenously added nicotinamide may comprise, before the addition of exogenous
30 nicotinamide as a supplement, nicotinamide at a concentration less than 0.5 mM or more preferably less than 0.05 mM.

According to one embodiment, the MSCs are at least 50 % purified, at least 75 % purified or at least 90 % purified.

The population of mesenchymal stem cells may be seeded (and also cultured) in any medium including those described herein above or those disclosed in U.S. Patent Application No. 20050260748, incorporated herein by reference.

The time ratio of culturing in the presence of nicotinamide and FGF4: culturing in the absence of nicotinamide may vary and may include all ratios from 1:99; 2:98; 3:97; 4:96; 5:95; 6:94; 7:93; 8:92; 9:91; 10:90; 11:89; 12:88; 13:87; 14:86; 15:85; 16:84; 17:83; 18:82; 19:81; 20:80; 21:79; 22:78; 23:77; 24:76; 25:75; 26:74; 27:73; 28:72; 29:71; 30:70; 31:69; 32:68; 33:67; 34:66; 35:65; 36:64; 37:63; 38:62; 39:61; 40:60; 41:59; 42:58; 43:57; 44:56; 45:55; 46:54; 47:53; 48:52; 49:51; 50:50; 51:49; 52:48; 53:47; 54:46; 55:45; 56:44; 57:43; 58:42; 59:41; 60:40; 61:39; 62:38; 63:37; 64:36; 65:35; 66:34; 67:33; 68:32; 69:31; 70:30; 71:29; 72:28; 73:27; 74:26; 75:25; 76:24; 77:23; 78:22; 79:21; 80:20; 81:19; 82:18; 83:17; 84:16; 85:15; 86:14; 87:13; 88:12; 89:11; 90:10; 91:9; 92:8; 93:7; 94:6; 95:5; 96:4; 97:3; 98:2; 99:1.

According to one embodiment, at least one full round of propagation is effected in the presence of nicotinamide.

It will be appreciated that the culturing in the medium comprising nicotinamide may be effected prior or following the culturing in the medium devoid of nicotinamide.

According to embodiments of the present invention, the medium which is devoid of nicotinamide comprises FGF4 (either at the same or a different concentration as the medium which comprises nicotinamide).

According to other embodiments of the present invention, the medium which is devoid of nicotinamide is further devoid of FGF4.

Further, the present inventors contemplate more than one culturing stage in the presence of nicotinamide and FGF4 interspersed with culturing stages in the absence of the nicotinamide and vice versa.

According to one embodiment, the culturing in the presence of nicotinamide and FGF4 is effected for at least one day, at least two days, at least three days, at least four days, at least five days, at least six days, at least one week, at least two weeks, at least three weeks, at least four weeks or at least five weeks.

According to another embodiment, the culturing in the absence of nicotinamide is effected for at least one day, at least two days, at least three days, at least four days, at least five days, at least six days, at least one week, at least two weeks, at least three weeks, at least four weeks or at least five weeks.

5 As mentioned, the second step of the purification process is selecting for MSCs based on the expression of a mesenchymal stem cell surface marker. The selection or sorting step may comprise selecting mesenchymal stem cells (MSC) from the mixed population of cells by means of one or more of such surface markers. The use of a selection or sorting step further enhances the stringency of sorting and selection
10 specificity for MSCs and furthermore potentially reduces possible contamination from the starting material.

Prior to sorting, the mixed cell populations are typically dispersed using cell dispersing agents. Preferably single cell populations are obtained. Examples of agents that may be used to disperse the cells include, but are not limited to collagenase, dispase,
15 accutase, trypsin (e.g. trypsin-EDTA), papain. Alternatively, or additionally trituration may also be performed to increase the dispersal of the cells.

According to a specific embodiment, the selecting is effected by selecting cells which express VCAM-1/CD106 (NP_001069.1) above a predetermined level.

According to another embodiment, the selecting is effected by selecting cells
20 which express at least one of CD105 (SH2), CD73 (SH3/4), CD44, CD90 (Thy-1), CD71, STRO-1, CD29, CD166, CD146, CD106 and CD271 above a predetermined level.

According to a particular embodiment, the surface marker is stromal precursor antigen-1 (STRO-1), CD105 or VCAM (CD106).

25 According to still another embodiment, the selecting is effected by selecting cells which express at least one of CD34, CD11B, CD43 and CD45 below a predetermined level.

A number of methods are known for selection or sorting based on antigen expression, and any of these may be used in the selection or sorting step described here.
30 In particularly preferred embodiments, the analysis is achieved using a flow cytometer and the cells are subsequently sorted based upon the specific light scattering and

fluorescent characteristics of each cell. Thus, the selection or sorting may be achieved by means of fluorescence activated cell sorting (FACS).

As is known in the art, FACS involves exposing cells to a reporter, such as a fluorescently labelled antibody, which binds to and labels antigens expressed by the cell. Methods of production of antibodies and labelling thereof to form reporters are known in the art, and described for example in Harlow and Lane. Antibodies that may be used for FACS analysis are taught in Schlossman S, Boumell L, et al, [Leucocyte Typing V. New York: Oxford University Press; 1995] and are widely commercially available. The cells are then passed through a FACS machine, which sorts the cells from each other based on the labeling.

Alternatively or in addition, magnetic cell sorting (MACS) or immunopanning may be employed to sort the cells.

As mentioned hereinabove, the mixed cell populations are analyzed by a Flow Cytometer, such as a laser scanning Cytometer. A Flow Cytometer typically consists of a laser light source, flow measurement chamber, and an optical system consisting of lenses, filters, and light detectors. Two photo-multiplier tubes (light detectors), one at 180 degrees and one at 90 degrees to the laser, are used to measure forward (FSC) and right-angle scatter (SSC), respectively. Three fluorescence detectors, each consisting of a filter and photomultiplier tube, are used to detect fluorescence. The three detectors sense green (FL1--530 nm), orange (FL2--585 nm), and red fluorescence (FL3--650 nm). Cells are identified by sort logic applied to all five of the detector signals (FSC, SSC, FL1, FL2, FL3) using a computer.

Exemplary Flow Cytometers that may be used in this aspect of the present invention are manufactured by companies such as Becton Dickinson (USA), Beckman Coulter (USA), Partec (Germany).

The FACS machine may be set such that cells of a particular forward scatter and/or side scatter are selected. Forward-scattered light (FSC) is proportional to cell-surface area or size. FSC is a measurement of mostly diffracted light and is detected just off the axis of the incident laser beam in the forward direction by a photodiode. FSC provides a suitable method of detecting particles greater than a given size independent of their fluorescence.

Side-scattered light (SSC) is proportional to cell granularity or internal complexity. SSC is a measurement of mostly refracted and reflected light that occurs at any interface within the cell where there is a change in refractive index. SSC is collected at approximately 90 degrees to the laser beam by a collection lens and then
5 redirected by a beam splitter to the appropriate detector.

Thus, for example, the present invention contemplates selecting cells which have a diameter below about 20 μm , by gating at a particular forward scatter and a particular granularity by gating at a particular side scatter.

The present invention contemplates selecting particular cell populations based
10 on the level of cell surface expression. Thus, in the case of FACS, the machine may be set such that cell populations gated for events stained with a fluorescent intensity between about 20-100 (dim), between about 100-500 (moderate) or between about 500-2000, or greater (bright). The following cell populations are contemplated by the present invention:

15 VCAM1 bright cells;
VCAM1 moderate cells;
VCAM1 dim cells;
STRO-1 bright cells;
STRO-1 moderate cells;
20 STRO-1 dim cells;
CD105 bright cells;
CD105 moderate cells;
CD105 dim cells;

It will be appreciated that cell populations may be selected based on expression
25 of more than one of the above mentioned markers – e.g. at least two of the above mentioned markers or at least three of the above mentioned markers.

The above described cell populations are typically enriched for cells that do not express CD45. Thus, according to another embodiment, less than 10 % of the cells in the above described cell populations express CD45 as measured by FACS.

30 According to still another embodiment, more than 90 % of the cells in the above described cell populations express CD90, as measured by FACS.

According to still another embodiment, more than 95 % of the cells in the above described cell populations express CD90, as measured by FACS.

According to still another embodiment, more than 90 % of the cells in the above described cell populations express CD44, as measured by FACS.

5 According to still another embodiment, more than 95 % of the cells in the above described cell populations express CD44, as measured by FACS.

As mentioned, additional steps are contemplated by the present inventors prior to, during or following the two step protocol described herein. Such additional steps may involve culturing on a plastic surface, as described herein above and/or additional expansion steps, for example, as described herein above re culturing in nicotinamide.

10 In some embodiments, the cells are selected according to cell size, for example, by a cell counter based on Trypan Blue exclusion and graphical analysis. Suitable cell counters include, but are not limited to Cedex counters (Roche Innovatis). The number of cells that may be cultured according to any of the methods of the present invention may be any number including small batches - e.g. 100×10^4 cells to larger batches – e.g. 100×10^{12} or 100×10^{13} cells.

When large batches are required, the cells are typically cultured in a bioreactor (or in multi-level industrial flasks), the size of which is selected according to the number of cells being cultured.

20 Examples of flasks and plates that may be used for growing MSCs in commercial quantities include for example Corning HYPERFlask™ Cell Culture Vessel, Corning CellSTACK™ Chambers, Corning HYPERStack™ Cell Culture Vessel, 40 stack chambers and NUNC Automatic Cell Factory Manipulator.

As used herein, the term “bioreactor” refers to any device in which biological and/or biochemical processes develop under monitored and controlled environmental and operating conditions, for example, pH, temperature, pressure, nutrient supply and waste removal. According to one embodiment of the invention, the basic classes of bioreactors suitable for use with the present invention include static bioreactors, stirred flask bioreactors, rotating wall bioreactors, hollow fiber bioreactors and direct perfusion bioreactors, as further described in WO 2005/007799, the contents of which are incorporated by reference.

The cultured population of cells generated using the methods described herein may be further treated following the culturing or stored (e.g. cryopreserved) in the presence of a cryopreservant. Such cryopreservants include dimethyl sulfoxide (DMSO), glycerol, and the like.

5 The cell populations generated following the culturing and/or the expansion method of the present invention may be used for a variety of purposes including research, for screening agents which affect the differentiation thereof and for therapeutic uses. Additionally, or alternatively, the cell populations may be stored (e.g. frozen) until required.

10 According to one embodiment, the mesenchymal stem cell populations generated using the methods disclosed herein may be used for further differentiation protocols.

Methods of differentiating mesenchymal stem cells towards various cell lineages are known in the art.

15 Differentiating cells may be obtained by culturing or differentiating MSC in a growth environment that enriches for cells with the desired phenotype, e.g. osteoblasts, adipocytes, etc. The culture may comprise agents that enhance differentiation to a specific lineage.

Osteogenic differentiation may be performed by plating cells and culturing to
20 confluency, then culturing in medium comprising .beta.-glycerol phosphate, ascorbic acid and retinoic acid (see Cowan et al. (2005) Tissue engineering 11, 645-658).

To induce adipogenic differentiation detached cells may be reseeded in 24 well plates (7×10^4 cells/ml) and treated with adipogenic medium for three weeks. Two exemplary adipogenic mediums are provided: DMEM supplemented with 0.05 mg/ml
25 Gentamicin, 2 mM L-glutamine, 10 % FBS, 0.5 μ M 3-isobutyl-1-methylxanthine (IBMX, Sigma), 0.5 μ M hydrocortisone (Sigma) and 60 μ M indomethacin (Sigma), or MSC adipogenic stimulatory supplements purchased from StemCell Technologies, as per manufacturer's instructions. Adipogenic differentiation may be assessed by oil-red staining: cells are fixed with methanol at -20 °C for 10 minutes and treated with 60 %
30 isopropanol for 3 minutes. Plates may be stained in oil-red-O (Sigma) for 10 minutes and rinsed in tap water. After rinsing plates may be counterstained with Mayer hematoxylin (Sigma) for 1 minute and rinsed in tap water.

Myocyte differentiation may be performed by plating cells and culturing to confluency, then culturing in medium comprising horse serum, dexamethasone, and hydrocortisone (see Eun et al. (2004) Stem Cells 22:617-624); or 5-azacytidine (see Fukuda et al. (2001) Artificial Organs 25:187).

5 Chondrocyte differentiation may be performed by plating cells and culturing to confluency, then culturing in medium comprising dexamethasone, ascorbic acid 2-phosphate, insulin, transferrin, selenous acid, with or without TGF- β_1 (see Williams et al. (2003) Tissue Engineering 9(4):679).

Neuronal differentiation is known in the art. For example, generation of neurons
10 and or oligodendrocytes from mesenchymal stem cells may be effected as described in U.S. Patent No. 20100021434 and 20090257987.

Alternatively, or additionally, the mesenchymal stem cells may be genetically modified so as to express an agent (e.g. a polypeptide, siRNA or miRNA) that is useful for treating a disease or alternatively that drives its differentiation towards a certain
15 lineage.

Thus, for example, the mesenchymal stem cells may be genetically modified to express bone morphogenic factor 2 (BMP2) in order to promote differentiation into bone.

Alternatively, the mesenchymal stem cells may be genetically modified to
20 express Pd-x in order to promote differentiation into pancreatic cells.

Since mesenchymal stem cells are known to home and migrate towards wounds, the cells may be used as carriers, transporting useful molecules to the site of injury. The useful molecules may be molecules that are inherently found inside the mesenchymal stem cells (e.g. growth factors) or may be artificially placed inside the cells (i.e. proteins
25 or polynucleotides transfected into the cells).

Both the differentiated and non-differentiated mesenchymal stem cell populations described herein may be used to treat a myriad of disorders, the particular disorders being selected according to the differentiation status of the cells.

Thus, according to another aspect of the present invention there is provided a
30 method of treating a disease or disorder, the method comprising transplanting to a subject in need thereof a therapeutically effective amount of the isolated population of cells the present invention.

According to one embodiment, the disease or disorder is selected from the group consisting of a bone or cartilage disease, a neurodegenerative disease, a cardiac disease, a hepatic disease, cancer, nerve damage, wound healing, autoimmune disease, graft versus host disease, spinal cord injury and tissue regeneration.

5 Bone defects suitable for treatment using the cells of the present invention include, but are not limited to osteogenesis imperfecta, fracture, congenital bone defects, and the like.

Further, the mesenchymal stem cells of the present invention can be implanted in a subject to provide osseous and connective tissue support of orthopedic and other
10 (e.g. dental) prosthetic devices, such as joint replacements and/or tooth implants.

The mesenchymal stem cells of the present invention can be used to treat CNS diseases.

Representative examples of CNS diseases or disorders that can be beneficially treated with the cells described herein include, but are not limited to, a pain disorder, a
15 motion disorder, a dissociative disorder, a mood disorder, an affective disorder, a neurodegenerative disease or disorder and a convulsive disorder.

More specific examples of such conditions include, but are not limited to, Parkinson's, ALS, Multiple Sclerosis, Huntingdon's disease, autoimmune encephalomyelitis, diabetic neuropathy, glaucomatous neuropathy, macular
20 degeneration, action tremors and tardive dyskinesia, panic, anxiety, depression, alcoholism, insomnia, manic behavior, Alzheimer's and epilepsy.

As mentioned, since MSCs can differentiate into cartilage, the mesenchymal stem cells of the present invention may be suitable for the treatment of joint conditions including, but not limited to osteoarthritis, rheumatoid arthritis, inflammatory arthritis,
25 chondromalacia, avascular necrosis, traumatic arthritis and the like.

Bone marrow-derived mesenchymal stem cells (MSCs) are known to interact with hematopoietic stem cells (HSCs) and immune cells, and represent potential cellular therapy to enhance allogeneic hematopoietic engraftment and prevent graft-versus-host disease (GVHD). When hematopoietic stem cell numbers were limited, human
30 engraftment of NOD-SCID mice was observed only after co-infusion of unrelated human MSCs, but not with co-infusion of mouse mesenchymal cell line. Unrelated human MSCs did not elicit T-cell activation in vitro and suppressed T-cell activation by

Tuberculin and unrelated allogeneic lymphocytes in a dose-dependent manner. Cell-free MSC culture supernatant, mouse stromal cells and human dermal fibroblasts did not elicit this effect. These preclinical data suggest that unrelated, human bone marrow-derived, culture-expanded MSCs may improve the outcome of allogeneic transplantation by promoting hematopoietic engraftment and limiting GVHD (Maitra B, et al Bone Marrow Transplant. 2004 33(6):597-604).

It is known that when MSCs are introduced into the infarcted heart, they can prevent deleterious remodeling and improve recovery. MSCs have been injected directly into the infarct, or they have been administered intravenously and seen to home to the site of injury. Examination of the interaction of allogeneic MSCs with cells of the immune system indicates little rejection by T cells. Persistence of allogeneic MSCs in vivo suggests their potential "off the shelf" therapeutic use for multiple recipients (Pittenger MF, et al Circ Res. 2004 Jul 9;95(1):9-20).

The use of ex-vivo expanded mesenchymal cells for transplantation has the following advantages:

It reduces the volume of blood or other tissue required for reconstitution of a recipient adult tissue system.

It enables storage of small number of unexpanded mesenchymal cells, for example, from cord blood or peripheral blood, for potential future use.

In the case of autologous transplantation of recipients with malignancies, contaminating tumor cells in autologous infusion often contribute to the recurrence of the disease. Selecting and expanding mesenchymal cells will reduce the load of tumor cells in the final transplant.

Tissue regeneration: Mesenchymal cell populations of the present invention can be used for the promotion of tissue regeneration. Transplantation of mesenchymal stem cells has great promise for benefits in regenerative medicine, autoimmune diseases, inflammatory conditions, acute and chronic ischemic conditions reconstructive surgery, tissue engineering, regenerating new tissues and naturally healing diseased or injured organs.

Gene therapy: For successful long-term gene therapy, a high frequency of genetically modified cells with transgenes stably integrated within their genome is an obligatory requirement. Presently, gene transfer into fresh stem and/or progenitor cells

is highly inefficient. The ability to store and process a selected population of mesenchymal cells *ex-vivo*, and enhance their homing and engraftment potential would provide for an increased probability of the successful use of genetically modified cell transplantation [Palmiter Proc Natl Acad Sci USA 91(4): 1219-1223, (1994)].

5 In any of the methods described herein the cells may be obtained from an autologous, semi-autologous or non-autologous (i.e., allogeneic or xenogeneic) human donor or embryo or cord/placenta. For example, cells may be isolated from a human cadaver or a donor subject.

The term semi-autologous refers to donor cells which are partially-mismatched to recipient cells at a major histocompatibility complex (MHC) class I or class II locus.

The cells of the present invention can be administered to the treated individual using a variety of transplantation approaches, the nature of which depends on the site of implantation.

15 According to one embodiment, the cells are not transplanted into the body in a medium comprising nicotinamide.

The cells may be transplanted to a damaged or healthy region of the tissue. In some cases the exact location of the damaged tissue area may be unknown and the cells may be inadvertently transplanted to a healthy region. In other cases, it may be preferable to administer the cells to a healthy region, thereby avoiding any further damage to that region. Whatever the case, following transplantation, the cells preferably migrate to the damaged area.

The term or phrase "transplantation", "cell replacement" or "grafting" are used interchangeably herein and refer to the introduction of the cells of the present invention to target tissue. As mentioned, the cells can be derived from the recipient or from an allogeneic, semi-allogeneic or xenogeneic donor. Other xeno-origins are also contemplated.

25 Cells of the present invention may be transplanted by means of direct injection into an organ, injection into the bloodstream, intraperitoneal injection, injection directly to lymphoid organs etc. Suitable methods of transplantation can be determined by monitoring the homing and engraftment of the implanted cells to the desired organ, the expression of desired organ-specific genes or markers, and the function of the derived organ of the subject. In the pancreas, for example, maintenance of euglycemia,

secretion of insulin and/or C peptide can be a measure of the restoration of function to a diabetic host animal following cell replacement therapy as disclosed hereinbelow. In the liver, for example, albumin synthesis can be monitored.

MSCs typically down regulate MHC class 2 and are therefore less immunogenic. Embryonal or newborn cells obtained from the cord blood, cord's Warton's jelly or placenta are further less likely to be strongly immunogenic and therefore less likely to be rejected, especially since such cells are immunosuppressive and immunoregulatory to start with.

Notwithstanding, since non-autologous cells may induce an immune reaction when administered to the body several approaches may be taken to reduce the likelihood of rejection of non-autologous cells. These include either administration of cells to privileged sites, or alternatively, suppressing the recipient's immune system, providing anti-inflammatory treatment which may be indicated to control autoimmune disorders to start with and/or encapsulating the non-autologous/semi-autologous cells in immunoisolating, semipermeable membranes before transplantation. Encapsulation techniques are generally classified as microencapsulation, involving small spherical vehicles and macroencapsulation, involving larger flat-sheet and hollow-fiber membranes (Uludag, H. et al. Technology of mammalian cell encapsulation. Adv Drug Deliv Rev. 2000; 42: 29-64).

Methods of preparing microcapsules are known in the arts and include for example those disclosed by Lu M Z, et al., Cell encapsulation with alginate and alpha-phenoxybenzylidene-acetylated poly(allylamine). Biotechnol Bioeng. 2000, 70: 479-83, Chang T M and Prakash S. Procedures for microencapsulation of enzymes, cells and genetically engineered microorganisms. Mol. Biotechnol. 2001, 17: 249-60, and Lu M Z, et al., A novel cell encapsulation method using photosensitive poly (allylamine alpha-cyanobenzylideneacetate). J. Microencapsul. 2000, 17: 245-51.

For example, microcapsules are prepared by complexing modified collagen with a ter-polymer shell of 2-hydroxyethyl methylacrylate (HEMA), methacrylic acid (MAA) and methyl methacrylate (MMA), resulting in a capsule thickness of 2-5 .mu.m. Such microcapsules can be further encapsulated with additional 2-5 .mu.m ter-polymer shells in order to impart a negatively charged smooth surface and to minimize plasma

protein absorption (Chia, S. M. et al. Multi-layered microcapsules for cell encapsulation Biomaterials. 2002 23: 849-56).

Other microcapsules are based on alginate, a marine polysaccharide (Sambanis, A. Encapsulated islets in diabetes treatment. Diabetes Technol. Ther. 2003, 5: 665-8) or its derivatives. For example, microcapsules can be prepared by the polyelectrolyte complexation between the polyanions sodium alginate and sodium cellulose sulphate with the polycation poly (methylene-co-guanidine) hydrochloride in the presence of calcium chloride.

It will be appreciated that cell encapsulation is improved when smaller capsules are used. Thus, the quality control, mechanical stability, diffusion properties, and in vitro activities of encapsulated cells improved when the capsule size was reduced from 1 mm to 400 μm (Canaple L. et al., Improving cell encapsulation through size control. J Biomater Sci Polym Ed. 2002; 13:783-96). Moreover, nanoporous biocapsules with well-controlled pore size as small as 7 nm, tailored surface chemistries and precise microarchitectures were found to successfully immunoisolate microenvironments for cells (Williams D. Small is beautiful: microparticle and nanoparticle technology in medical devices. Med Device Technol. 1999, 10: 6-9; Desai, T. A. Microfabrication technology for pancreatic cell encapsulation. Expert Opin Biol Ther. 2002, 2: 633-46).

Examples of immunosuppressive agents include, but are not limited to, methotrexate, cyclophosphamide, cyclosporine, cyclosporin A, chloroquine, hydroxychloroquine, sulfasalazine (sulphasalazopyrine), gold salts, D-penicillamine, leflunomide, azathioprine, anakinra, infliximab (REMICADETM), etanercept, TNF alpha blockers, a biological agent that targets an inflammatory cytokine, and Non-Steroidal Anti-Inflammatory Drug (NSAIDs). Examples of NSAIDs include, but are not limited to acetyl salicylic acid, choline magnesium salicylate, diflunisal, magnesium salicylate, salsalate, sodium salicylate, diclofenac, etodolac, fenoprofen, flurbiprofen, indomethacin, ketoprofen, ketorolac, meclofenamate, naproxen, nabumetone, phenylbutazone, piroxicam, sulindac, tolmetin, acetaminophen, ibuprofen, Cox-2 inhibitors and tramadol.

Cell populations of the present invention can be provided per se, along with the culture medium containing same, isolated from the culture medium, and combined with

a pharmaceutically acceptable carrier as well as with additional agents which may promote cell engraftment and/or organ function (e.g., immunosuppressing agents, antibiotics, growth factor). Hence, cell populations of the invention can be administered in a pharmaceutically acceptable carrier or diluent, such as sterile saline and aqueous buffer solutions. The use of such carriers and diluents is well known in the art.

Compositions of the present invention may, if desired, be presented in a pack or dispenser device, such as an FDA-approved kit, which may contain one or more unit dosage forms containing the active ingredient (e.g., cells). The pack may, for example, comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. The pack or dispenser device may also be accompanied by a notice in a form prescribed by a governmental agency regulating the manufacture, use, or sale of pharmaceuticals, which notice is reflective of approval by the agency of the form of the compositions for human or veterinary administration. Such notice, for example, may include labeling approved by the U.S. Food and Drug Administration for prescription drugs or of an approved product insert. Compositions comprising a preparation of the invention formulated in a pharmaceutically acceptable carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition, as further detailed above.

The cells prepared according to the methods of the present invention can be administered to the subject per se, seeded on a scaffold and/or in a pharmaceutical composition where it is mixed with suitable carriers or excipients.

As used herein, a "pharmaceutical composition" refers to a preparation of one or more of the active ingredients described herein with other chemical components such as physiologically suitable carriers and excipients. The purpose of a pharmaceutical composition is to facilitate administration of a compound to an organism.

Hereinafter, the phrases "physiologically acceptable carrier" and "pharmaceutically acceptable carrier," which may be used interchangeably, refer to a carrier or a diluent that does not cause significant irritation to an organism and does not abrogate the biological activity and properties of the administered compound. An adjuvant is included under these phrases.

Herein, the term "excipient" refers to an inert substance added to a pharmaceutical composition to further facilitate administration of an active ingredient.

Examples, without limitation, of excipients include calcium carbonate, calcium phosphate, various sugars and types of starch, cellulose derivatives, gelatin, vegetable oils, and polyethylene glycols.

Techniques for formulation and administration of drugs may be found in the latest edition of "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, PA, which is herein fully incorporated by reference.

Suitable routes of administration may, for example, include oral, rectal, transmucosal, especially transnasal, intestinal, or parenteral delivery, including intramuscular, subcutaneous, and intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections.

Alternately, one may administer the pharmaceutical composition in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into a tissue region of a patient.

Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries, which facilitate processing of the active ingredients into preparations that can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

For injection, the active ingredients of the pharmaceutical composition may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hank's solution, Ringer's solution, or physiological salt buffer. For transmucosal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

Pharmaceutical compositions suitable for use in the context of the present invention include compositions wherein the active ingredients are contained in an amount effective to achieve the intended purpose. More specifically, a "therapeutically effective amount" means an amount of active ingredients (e.g., a nucleic acid construct) effective to prevent, alleviate, or ameliorate symptoms of a disorder (e.g., ischemia) or prolong the survival of the subject being treated.

Determination of a therapeutically effective amount is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein.

For any preparation used in the methods of the invention, the dosage or the therapeutically effective amount can be estimated initially from in vitro and cell culture assays. For example, a dose can be formulated in animal models to achieve a desired concentration or titer. Such information can be used to more accurately determine useful
5 doses in humans.

Toxicity and therapeutic efficacy of the active ingredients described herein can be determined by standard pharmaceutical procedures in vitro, in cell cultures or experimental animals. The data obtained from these in vitro and cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The
10 dosage may vary depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration, and dosage can be chosen by the individual physician in view of the patient's condition. (See, e.g., Fingl, E. et al. (1975), "The Pharmacological Basis of Therapeutics," Ch. 1, p.1.)

Depending on the severity and responsiveness of the condition to be treated,
15 dosing can be of a single or a plurality of administrations, with course of treatment lasting from several days to several weeks, or until cure is effected or diminution of the disease state is achieved.

The amount of a composition to be administered will, of course, be dependent on the subject being treated, the severity of the affliction, the manner of administration,
20 the judgment of the prescribing physician, etc.

As used herein the term "method" refers to manners, means, techniques and procedures for accomplishing a given task including, but not limited to, those manners, means, techniques and procedures either known to, or readily developed from known manners, means, techniques and procedures by practitioners of the chemical,
25 pharmacological, biological, biochemical and medical arts.

As used herein, the term "treating" includes abrogating, substantially inhibiting, slowing or reversing the progression of a condition, substantially ameliorating clinical or aesthetical symptoms of a condition or substantially preventing the appearance of clinical or aesthetical symptoms of a condition.

30 Various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below find experimental support in the following examples.

Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

EXAMPLES

Reference is now made to the following examples, which together with the above descriptions, illustrate the invention in a non limiting fashion.

Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, "Molecular Cloning: A laboratory Manual" Sambrook et al., (1989); "Current Protocols in Molecular Biology" Volumes I-III Ausubel, R. M., ed. (1994); Ausubel et al., "Current Protocols in Molecular Biology", John Wiley and Sons, Baltimore, Maryland (1989); Perbal, "A Practical Guide to Molecular Cloning", John Wiley & Sons, New York (1988); Watson et al., "Recombinant DNA", Scientific American Books, New York; Birren et al. (eds) "Genome Analysis: A Laboratory Manual Series", Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; "Cell Biology: A Laboratory Handbook", Volumes I-III Cellis, J. E., ed. (1994); "Current Protocols in Immunology" Volumes I-III Coligan J. E., ed. (1994); Stites et al. (eds), "Basic and Clinical Immunology" (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (eds), "Selected Methods in Cellular Immunology", W. H. Freeman and Co., New York (1980); available immunoassays are extensively described in the patent and scientific literature, see, for example, U.S. Pat. Nos. 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521; "Oligonucleotide Synthesis" Gait, M. J., ed. (1984); "Nucleic Acid Hybridization" Hames, B. D., and Higgins S. J., eds. (1985); "Transcription and Translation" Hames, B. D., and Higgins S. J., Eds. (1984); "Animal Cell Culture"

Freshney, R. I., ed. (1986); "Immobilized Cells and Enzymes" IRL Press, (1986); "A Practical Guide to Molecular Cloning" Perbal, B., (1984) and "Methods in Enzymology" Vol. 1-317, Academic Press; "PCR Protocols: A Guide To Methods And Applications", Academic Press, San Diego, CA (1990); Marshak et al., "Strategies for Protein Purification and Characterization - A Laboratory Course Manual" CSHL Press (1996); all of which are incorporated by reference as if fully set forth herein. Other general references are provided throughout this document. The procedures therein are believed to be well known in the art and are provided for the convenience of the reader. All the information contained therein is incorporated herein by reference.

EXAMPLES

METHODS AND EXPERIMENTAL PROCEDURES

Mesenchymal stem cells Isolation: Bone marrow derived and adipose tissue derived mesenchymal cells were isolated based on their plastic adherence potential in expansion medium containing: High glucose DMEM and 10 % Fetal Bovine Serum (FBS, Hyclone, Logan, UT, USA) supplemented with 0.05 mg/ml Gentamicin (Sigma) and 2 mM L-glutamine (Biological Industries, Israel). Cells were allowed to adhere for 3-4 days and non-adherent cells were washed out with medium changes. The medium was further exchanged with fresh medium every 3-4 days.

Hematopoietic stem and progenitor cells: Umbilical cord-derived hematopoietic stem cells were isolated using CD133 microbeads and CliniMACS® separator (Miltenyi, Inc. Auburn, CA), and cultured for 3 weeks in MEMα supplemented with 50ng/ml TPO, IL6, SCF, Flt3, fetal bovine serum, ± 2.5 or 5mM nicotinamide, and/or 10, 50 or 200ng/ml FGF4. After 3 weeks in culture the cells were stained for surface markers (CD38, CD133, CD33, CD19) and the cell populations determined by FACS analysis (see below). Results are expressed as percentage of total population assayed.

Maintenance and expansion: Once adherent cells reached approximately 80-90 % confluency, they were detached with 0.25 % trypsin-EDTA (Sigma), washed twice in DMEM and 10 % Fetal Bovine Serum, with centrifugation, 400 g, 5 minutes, and replated at a 1:2 to 1:1000 dilution under the same culture conditions.

Measurement of cell size: Cell size was measured using Cedex Automated Cell Counter (Innovatis). The cells were diluted 1:2 in Trypan Blue (Sigma) and cell size was measured automatically under microscope.

Measurement of granularity: Following trypsin treatment, the cells were
5 analyzed for granularity by side scatter FACS.

Measurement of number of cells in culture: Cell number was measured using Cedex Automated Cell Counter (Innovatis). The cells were diluted 1:2 in Trypan Blue (Sigma) and cell number was measured automatically under microscope.

Surface antigen analysis: At different time points the cells were detached with
10 0.25 % trypsin-EDTA. The cells were washed with a PBS solution containing 1 % BSA, and stained (at 4 °C for 30 min) with either fluorescein isothiocyanate (FITC) or phycoerythrin (PE)-conjugated antibodies: 105 PE, 105 FITC (Serotec, Raleigh, NC), 45 FITC, 14 FITC, HLA-DR FITC, 106 PE, 31 PE (BD, Franklin Lakes NJ), 34 PE (Dako, Glostrup, Denmark), 73 PE, HLA class1 PE, 49b PE (Pharmingen, San Diego,
15 CA), 29 PE, 44 PE, 54 FITC, 59 PE, 90 PE (BioLegend, San Diego, CA). CD133 - (AC141) PE (Miltenyi, Auburn, CA), CD38 FITC (Dako, Glostrup, Denmark), CD19 FITC (BD Biosciences, Franklin Lakes NJ), CD33 FITC (BD Biosciences, Franklin Lakes NJ).

The cells were then washed in the above buffer and analyzed using a
20 FACScalibur^R flow cytometer (Becton Dickinson, Franklin Lakes NJ). The cells were passed at a rate of up to 1000 cells/second, using a 488 nm argon laser beam as the light source for excitation. Emission of 10000 cells was measured using logarithmic amplification, and analyzed using the CellQuest software (Becton Dickinson). Cells stained with FITC- and PE-conjugated isotype control antibodies were used to
25 determine background fluorescence.

CFU-F assay: Cultured MSCs were seeded in 6-well plates at density of 50-100 cells/cm² and maintained with DMEM and 10 % FBS. After 14 days the cells were fixed using 10 % cold Formalin (Sigma) and stained with Harris Hematoxylin (Sigma). Clones (cluster of more than 50 cells with evident epicenter) are stained blue-purple and
30 counted using microscope.

Senescence evaluation assay: Cultured MSCs were stained using the Senescence beta-Galactosidase Staining Kit (Cell Signaling). The cells are fixed and

stained for detection of beta-Galactosidase activity at pH 6 using X-Gal and incubation overnight in 37 °C in dry incubator.

In-vitro wound healing assay: Wound was performed in MSCs cultures at ~70 % confluence using 200 µl or 1000 µl tip. Four days later the cells were fixated using
5 10% cold Formalin (Sigma) and stained with Harris Hematoxylin (Sigma). The in-vitro wound healing process was evaluated using microscope.

Treatment of mesenchymal cultures with nicotinamide: Mesenchymal cultures were initiated as described above, and supplemented with nicotinamide 1-15 mM alone, or in combination with growth factors or growth factors alone, incubated at 37 °C in a
10 humidified atmosphere of 5 % CO₂ in air. At each passage and at each medium exchange, the cultures were supplemented with mesenchymal medium, nicotinamide and growth factors.

In some experiments, the adherent cells were cultured with or without nicotinamide and indicated factors, and 24 hours before passage 4, and the medium
15 replaced with medium without fetal bovine serum or FGF4.

In Vitro Migration Assay: RPMI plus 10 % FCS (0.6 ml) containing 100 ng/ml CXCL12 (R&D Systems) was placed into the lower chamber of a Costar 24-well “transwell” culture plate (Corning, Inc, Corning, NY). Cells (2×10^5) in 100-µl medium were introduced into the upper chamber, over a porous membrane (pore size, 5 µm).
20 After 4 hours, cells were collected from both chambers and counted by flow cytometry (FACSort, Becton Dickinson and Co, San Jose, CA, USA). Spontaneous migration was performed as a control without CXCL12 in the lower chamber.

In vivo analysis of homing: NOD/SCID mice (8-10 week old) (Harlan Ltd., Israel) were sub-lethally irradiated (at 375cGy at 67cGy/min) and 24 hours later
25 inoculated via the tail vein with either CFSE-labeled mesenchymal stem cells cultured in the presence of nicotinamide or CFSE-labeled mesenchymal stem cells cultured in the absence of nicotinamide. Mice were sacrificed at 24 hours post injection and bone marrow or other tissue samples were obtained. Homing of human cells was detected by flow cytometry via visualization of CFSE-stained cells over a background of unlabeled
30 murine cells. The bright fluorescence of CFSE is sufficient to separate labeled human cells from unlabeled murine cells by at least 1 log. To quantify homing of human progenitor cells, bone marrow cells were stained with APC-conjugated antihuman cell

marker monoclonal antibodies and CFSE⁺/cell marker cells were enumerated. For each sample 100,000 events are recorded and analyzed.

Transplantation of mesenchymal cells into NOD/SCID mice: NOD/SCID mice were bred and maintained in sterile intra-ventilated cages (Techniplast, Bugugiatte, Italy). Eight-week-old mice were sub-lethally irradiated as described above. Mice were inoculated via the tail vein with mesenchymal cells cultured in the presence or absence of nicotinamide. To avoid donor variability, mesenchymal cells from several units were pooled and used for expansion cultures as well as group injection. Mice were sacrificed at week 4, and marrow samples were obtained by flushing their femurs and tibias with IMDM at 4 °C. Flow cytometric analysis of NOD/SCID marrow cells was performed as described hereinabove, using monoclonal antibodies against human cell surface differentiation antigens to identify human cell engraftment.

Delayed Type Hypersensitivity Assay: BALB/C mice were sensitized with Oxazolone (4-ethoxymethylene-2-phenyloxazol-5-one), and 6 days later challenged with Oxazolone, injected into the ear. Immune modulation by candidate compositions, as indicated, was determined 24 hours following their topical administration, by measurement of ear thickness with a caliper.

Growth Factor Secretion: Medium from MSC cultured as indicated, and depleted of fetal bovine serum and FGF4 24 hours before passage 4 was collected and assayed by ELISA for growth factors and immune-related factors secreted into the medium (human growth factor HGF, transforming growth factor beta TGF- β , keratinocyte growth factor KGF and interleukin 6 IL-6). ELISA was carried out using solid phase sandwich ELISA kits specific for human KGF (R&D systems, cat#DKG00), IL-6 (R&D systems, cat#D6050), TGF- β 1 (R&D systems, cat#DB100B), HGF (R&D systems, cat#DHG00).

Keratinocyte Proliferation Assay: Normal human epidermal keratinocytes (Promocell, GmbH, Heidelberg, Germany) were cultured for one passage in keratinocyte growth medium, detached and reseeded in keratinocyte growth medium (containing 50% Supplement Mix) diluted, as indicated, with MSC conditioned medium. Medium was changed twice a week, and keratinocytes detached and counted after reaching 90% confluence.

Mixed Lymphocyte Reaction-Like Assay: Peripheral blood derived mononuclear cells (MNCs), containing greater than 50% T-cells, were isolated by buoyant density centrifugation and activated with 3 µg/ml phytohemagglutinin (PHA). Following activation, the cells were cultured with or without conditioned medium from MSC culture and other factors. Response of the PBMN cells to the activation by PHA was measured by the extent of TNF-alpha secretion into the medium cell (pg/ml), 72 hours after initial activation, measured by ELISA.

Statistics- The non-parametric Wilcoxon Rank Test was applied for testing differences between the study groups. All the tests applied were two-tailed, and a p value of $\leq 5\%$ was considered statistically significant. The data were analyzed using SAS software (SAS Institute, Cary, NC).

EXAMPLE 1

Analyzing of nicotinamide on mesenchymal stem cells cultured in the presence of growth factors

MATERIALS AND METHODS

Mesenchymal stem cells were selected and cultured in the presence of particular growth factors (basic fibroblast growth factor - bFGF, heparin binding epidermal growth factor - HB-EGF, fibroblast growth factor 4 - FGF-4 and platelet derived growth factor, homodimer, subunit B, PDGF-BB) in the presence and absence of nicotinamide for three or four passages and the number and size of the cells was calculated.

Two concentrations (10 and 50 ng/ml) of each one of the following factors were examined.

The experimental groups were as follows:

Group 1: Ctrl

Group 2: 10 ng/ml growth factor

Group 3: 50 ng/ml growth factor

Group 4: 5 mM NAM

Group 5: 5 mM NAM + 10 ng/ml growth factor

Group 6: 5 mM NAM + 50 ng/ml growth factor

In addition, the influence of the combination: 5 mM NAM + 50 ng/ml FGF4 + 50 ng/ml PDGF-BB was examined in comparison to an individual supplement.

RESULTS

Figure 1 illustrates that basic FGF has a negative effect on the ability of nicotinamide to increase proliferation of mesenchymal stem cells.

Figures 2A-B illustrate that heparin-binding EGF-like growth factor (HB-EGF) has a negative effect on the ability of nicotinamide to increase proliferation of mesenchymal stem cells.

Figures 3-5 illustrate that nicotinamide has a potentiating effect on the ability of FGF4 to increase proliferation of mesenchymal stem cells.

Figures 6A-D illustrate that PDGF-BB has an inconsistent effect on the ability of nicotinamide to increase proliferation of mesenchymal stem cells.

Figures 7A-D illustrate that MSC cultures treated with PDGF-BB or a combination of PDGF-BB + NAM are contaminated with a higher fraction of cells other than MSCs as compared with cultures treated without PDGF-BB.

Figures 8A-B, 9A-B and 10A-H illustrate that the combination of three factors – FGF4, nicotinamide and PDGF-BB has a detrimental effect on proliferation of mesenchymal stem cells as compared to the effect of FGF4 and nicotinamide in the absence of PDGF-BB.

Figure 27 illustrates the consistent synergic effect of combined nicotinamide and FGF4 on the expansion (cumulative cell count) throughout 5 passages.

EXAMPLE 2

The effect of Nicotinamide on bone marrow derived mesenchymal stem cell culture

The present inventors showed that nicotinamide increased the seeding efficacy (selection) of bone marrow derived MSCs. Phenotypic characterization of these cells after one passage in nicotinamide is shown in Figures 11 and 13. Figures 15, 17A-B, 22 and 26 illustrate the effect of nicotinamide on the expansion rate of bone marrow derived MSCs. Low concentrations of nicotinamide (e.g. 0.1 mM) had insignificant effect on the expansion rate of bone marrow derived MSCs (Figure not shown). Figures 20A-C and 21A-B illustrate that mesenchymal stem cells grown in the presence of nicotinamide are smaller and less granular than mesenchymal stem cells grown in the absence of nicotinamide under identical conditions.

EXAMPLE 3***Nicotinamide increases expansion of cultured adipose tissue-derived mesenchymal cells***

Phenotypic characterization of adipose tissue derived MSCs is shown in Figure 12. As illustrated in Figures 14 and 16, nicotinamide substantially improved adipose derived mesenchymal stem cell expansion in culture.

EXAMPLE 4***Nicotinamide increases tissue homing of cultured mesenchymal cells***

To evaluate the effect of nicotinamide on the homing of cultured mesenchymal cells, NOD/SCID mice are transplanted with either non-cultured mesenchymal cells, or with their total progeny following 3-weeks in culture with cytokines, with or without nicotinamide. Prior to transplantation, the cells are labeled with CFSE. Twenty-four hours post transplantation total CFSE-labeled cells and CFSE labeled mesenchymal cells that homed to the mouse bone marrow of the recipient mice are quantified by FACS.

Results indicate an effect of nicotinamide on tissue homing of mesenchymal cells, if the homing of nicotinamide-treated mesenchymal cells is significantly higher than the homing of non-cultured mesenchymal cells not subjected to nicotinamide

EXAMPLE 5***Nicotinamide increases functionality of chemokine receptors and adhesion molecules***

In order to determine the role of adhesion and related molecules in nicotinamide-mediated enhancement of homing and engraftment of cells, the effect of nicotinamide on *in-vitro* migration and the functionality of the adhesion molecules can be tested.

Using a trans-well migration assay, CXCL12-induced migration of non-cultured and cultured mesenchymal cells is tested, assessing the effects of nicotinamide on integrin and adhesion molecule function. Enhanced stimulation of migration in the nicotinamide treated cells, compared to the cells cultured without nicotinamide or non-cultured cells suggests that treatment of mesenchymal cells with nicotinamide can

potentially increase the responsiveness of adhesion molecules to their ligands, resulting in enhanced engraftment and homing potential of the nicotinamide-treated cells.

The functional quality of cell binding to adhesion molecules can also be investigated using shear flow analysis. The strong effect of nicotinamide on adhesion molecule-mediated binding and retention on substrate adhesion molecules can be evidenced by significantly enhanced percentage of initially settled cells resistant to removal by shear stress evident in the mesenchymal cells treated with nicotinamide.

EXAMPLE 6

Nicotinamide increases the SCID-repopulating capacity of cultured cells

Nicotinamide treatment is tested for ability to enhance homing and engraftment of transplanted cells by repopulation of NOD/SCID mice. To evaluate repopulating capacity, NOD/SCID mice are transplanted with non-cultured mesenchymal cells (n = 12) over a range of doses intended to achieve a sub-optimal transplantation, and subsequent non-engraftment in a fraction of mice or their progeny following 3-weeks expansion with cytokines. Human cell engraftment is evaluated 4-weeks post transplantation. Mice are scored as positively engrafted if 0.5 % of the recipient bone marrow cells expressed human CD45 antigen (CD45+). In the event that the presence of nicotinamide in culture results in superior and clear engraftment of mesenchymal cells in the mice at a predetermined dose range, while the untreated cells fail to engraft or engraft poorly, it can be concluded that nicotinamide is effective in enhancing engraftment and homing of transplanted mesenchymal cells.

EXAMPLE 7

Further analysis on the effect of nicotinamide on mesenchymal stem cells

MATERIALS AND METHODS

Mesenchymal stem cells were isolated using plastic adherence method, as described above and cultured for several passages with fetal bovine serum, $\pm$ NAM. The cells were selected in the presence of NAM.

At about 80 % confluence, adherent cells were collected following trypsin treatment, counted, characterized and re-seeded at a concentration of 1×10^3 cells/cm².

Measurement of VCAM1/CD106: Following Trypsin treatment the cells were analyzed for CD106 expression in FACS using anti-human CD106 PE antibodies.

Measurement of CD54: Following Trypsin treatment the cells were analyzed for CD54 expression in FACS using anti-human CD54 antibodies.

5 **RESULTS**

Figure 18 illustrates that the effect of nicotinamide on cell count was evident on large batches of mesenchymal stem cells indicating that large commercial batches of MSCs can be manufactured with less passages. This ensures better quality of the therapeutic cells due to shorter cultivation time and preservation of stem cells characteristics by nicotinamide.

Figure 19 illustrates that the effect of nicotinamide is not dependent on a particular batch of serum, and presents the results of one of two experiments performed. The cultures on these experiments were treated individually (each group was passaged upon reaching confluence).

15 The amount of senescent cells was reduced following culture in nicotinamide (Figures 23A-D).

Figure 24A illustrates that mesenchymal stem cells grown in the presence of nicotinamide express more VCAM1/CD106 adhesion molecule than mesenchymal stem cells grown in the absence of nicotinamide under identical conditions.

20 Figure 24B illustrates that mesenchymal stem cells grown in the presence of nicotinamide express less CD54 than mesenchymal stem cells grown in the absence of nicotinamide under identical conditions.

Figure 25 illustrates that mesenchymal stem cells grown in the presence of nicotinamide have higher ability to perform wound closure than mesenchymal stem cells grown in the absence of nicotinamide under identical conditions.

EXAMPLE 8

Effect of Combined Nicotinamide and FGF4 on Expression of Growth Factors in Cultured Mesenchymal Stem Cells

30 Culture of mesenchymal stem cells in the presence of nicotinamide and FGF4 provides a synergic increase in expansion potential of the mesenchymal stem cells, while maintaining the cells in an undifferentiated state (see Figures 3A-3D, 4A-4B, 5A-

5D and 27). In order to further characterize the MSCs expanded in these cultures, secretion of cytokines into the culture medium was measured by ELISA.

Figures 28 to 31 illustrate the significant increase in hepatocyte growth factor (HGF, Figure 28), transforming growth factor beta (TGF- β , Figure 29) and keratinocyte growth factor (KGF, Figure 30) with combined FGF4 and nicotinamide, compared to
5 nicotinamide alone. Figure 31 shows nicotinamide's reduction in pro-inflammatory interleukin 6 (IL-6) secreted, and the further reduction in IL6 with addition of FGF4.

When medium from mesenchymal stem cell cultured with nicotinamide or nicotinamide and FGF4 was assayed for effect on inflammation in the *in-vivo* delayed
10 hypersensitivity test, reduction in inflammatory response to challenge with the sensitizing allergen (Oxazolone) was clearly observed (data not shown). Further analysis in the *ex-vivo* mixed lymphocyte reaction-type assay clearly demonstrated the anti-inflammatory potential of the MSC culture medium from nicotinamide and nicotinamide with FGF4 in reducing secretion of TNF α by peripheral blood
15 mononuclear cells in response to activation with PHA (data not shown).

When medium from mesenchymal stem cell cultured with nicotinamide or nicotinamide and FGF4 was assayed for effect on keratinocyte proliferation *in-vitro*, significant induction of keratinocyte proliferation was clearly observed (data not shown).

20 Thus, adherent mesenchymal stem cells cultured with nicotinamide and nicotinamide in combination with FGF4 release biologically active factors into the medium, including factors having anti-inflammatory and proliferation-inducing activity.

EXAMPLE 9

25 *Effect of Combined Nicotinamide and FGF4 on Adipose-derived Mesenchymal Stem Cell in Culture*

Proliferation and cell size distribution in adipose-derived mesenchymal stem cells cultured with nicotinamide with or without additional FGF4 was assessed in up to 4 passages of the cultures.

30 Figures 32 to 33 illustrate the striking effect of combined nicotinamide and FGF4 on adipose derived adherent cell proliferation, expressed as the number of total

nucleated cells in the cultures, compared to controls as well as nicotinamide-treated cultures.

The size of mesenchymal stem cells in culture is often used as an indicator of the degree of differentiation of the MSCs, with the undifferentiated state more prevalent in the smaller size cells. Figure 34 illustrates the increased prevalence of smaller size cells in cultures of adipose derived MSCs exposed to nicotinamide, and the yet greater prevalence of smaller size cells among adipose derived MSCs exposed to nicotinamide and FGF4.

Thus, these results indicate that a combination of nicotinamide and FGF4 synergistically increases the rate of proliferation of adipose derived mesenchymal cells, while effectively maintaining the cells in an undifferentiated state.

EXAMPLE 10

Effect of Nicotinamide and FGF4 on Hematopoietic Stem Cell Differentiation

In order to determine whether or not the effects of combined nicotinamide and FGF4 on mesenchymal stem cells are a specific or generalized phenomenon, hematopoietic stem cells were cultured with and without FGF4 and nicotinamide, and the degree of differentiation of component sub-populations assessed according to cell surface markers.

MATERIALS AND METHODS

Umbilical cord-derived CD133+ hematopoietic stem cells were isolated and cultured for 3 weeks in medium supplemented with early-acting growth factors and fetal bovine serum, nicotinamide and/or FGF4. After 3 weeks in culture the cells were stained for surface markers (CD38, CD133, CD19) and the cell populations determined by FACS analysis. Cell proliferation was assessed by counting total cells at three weeks.

Results:

FGF4 does not affect differentiation of hematopoietic stem cells in culture:

When hematopoietic early progenitor cells (CD133+) are cultured for 3 weeks in the presence of 2.5 or 5 mM nicotinamide, the proportion of differentiated cells decreases significantly, and increase in the stem and progenitor cell fraction is clearly observed (see Figure 35, columns 1, 2 and 3). Exposure of the cells to 10 to 200 ng/ml

FGF4, on the other hand, is without any effect on the degree of differentiation at three weeks culture, as evidenced by the fraction of CD38 and CD133-expressing cells (see Figures 35 and 36, columns 1 and 4-6). Addition of FGF4, with or without nicotinamide was also without any discernible effect on total cell proliferation in cultures (results not shown).

Addition of FGF4 to cells cultured with nicotinamide neither improved nor reduced nicotinamides inhibition of hematopoietic stem cell differentiation, at any concentration of FGF4 (see, Figures 35-36, lanes 1 and 7-12).

FGF4 does not affect differentiation of HSC into committed myeloid or lymphoid lineage:

In order to determine whether committed hematopoietic stem cells were affected by exposure to FGF4 during culturing, the abundance of cells expressing CD33 and CD19, representing differentiated committed lineage myeloid and lymphoid cells, respectively, was measured in the three week cultures. While nicotinamide consistently reduced the committed cell fraction (see Figures 37 and 38, lanes 1-3 and 7-12), addition of FGF4, alone or in combination with nicotinamide, was without any effect on the abundance of committed myeloid or lymphoid cells (see Figures 37 and 38, lanes 1, 4-6 and 7-12).

Thus, these results clearly indicate that the proliferation-enhancing effects of FGF4 observed in mesenchymal stem cell culture, and the synergic effects of exposure of mesenchymal stem cells to combined nicotinamide and FGF4 are not a general phenomenon, and are not observed in ex-vivo hematopoietic stem cell cultures.

EXAMPLE 11

Combined use of nicotinamide +/- FGF4 followed by selection using VCAM1 /CD106. CD105 or STRO-1 for selection and expansion of mesenchymal stem cells

MATERIALS AND METHODS

Isolation: Bone marrow derived and adipose tissue derived mesenchymal cells are isolated based on their plastic adherence potential in expansion medium containing: High glucose DMEM and 10 % Fetal Bovine Serum (FBS, Hyclone, Logan, UT, USA), 0.05 mg/ml Gentamicin (Sigma) and 2 mM L-glutamine (Biological Industries, Israel).

Cells are allowed to adhere in the presence of nicotinamide +/- FGF4 for 3-4 days and non-adherent cells were washed out with medium changes.

Mesenchymal stem cells are cultured for several passages (1-3) with fetal bovine serum, + NAM, $\pm$ FGF4. The cells may be cultured on plastic adherent plates.

At about 80 % confluence, adherent cells are collected following trypsin treatment, counted, characterized and selected by CD105, CD106 or STRO-1 expression using directly or indirectly conjugated mouse anti-human antibodies (Miltenyi Biotec) and magnetic cell sorting (MACS) or Fluorescence-activated cell sorting (FACS) and re-seeded for further expansion.

Measurement of VCAM1/CD106, CD105 or STRO-1: following trypsin treatment the cells are analyzed for CD106 expression, STRO-1 expression or CD105 in FACS using anti-human CD106 PE antibodies, anti-human CD105 antibodies or anti-STRO-1 antibodies.

EXAMPLE 12

Combined use of nicotinamide +/-FGF4 preceded by selection using VCAM1 /CD106. CD105 or STRO-1 for selection of mesenchymal stem cells

MATERIALS AND METHODS

Selection of bone marrow derived and adipose tissue derived MSC: Bone marrow derived and adipose tissue derived mesenchymal cells are selected by CD105, CD106 or STRO-1 expression using directly or indirectly conjugated mouse anti-human antibodies (Miltenyi Biotec) and magnetic cell sorting (MACS). The selected cells are seeded at concentration of 6×10^3 cells/cm² in expansion medium containing: High glucose DMEM and 10 % Fetal Bovine Serum (FBS, Hyclone, Logan, UT, USA), 0.05 mg/ml Gentamicin (Sigma) and 2 mM L-glutamine (Biological Industries, Israel). Cells are allowed to adhere for 3-4 days and non-adherent cells were washed out with medium changes.

Culture of selected MSCs populations in NAM $\pm$ FGF4: Mesenchymal stem cells are cultured for several passages with fetal bovine serum, + NAM, $\pm$ FGF4.

It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for

brevity, described in the context of a single embodiment, may also be provided separately or in any suitable subcombination.

Although the invention has been described in conjunction with specific embodiments thereof, it is evident that many alternatives, modifications and variations will be apparent to those skilled in the art. Accordingly, it is intended to embrace all such alternatives, modifications and variations that fall within the spirit and broad scope of the appended claims. All publications, patents and patent applications and GenBank Accession numbers mentioned in this specification are herein incorporated in their entirety by reference into the specification, to the same extent as if each individual publication, patent or patent application or GenBank Accession number was specifically and individually indicated to be incorporated herein by reference. In addition, citation or identification of any reference in this application shall not be construed as an admission that such reference is available as prior art to the present invention.

WHAT IS CLAIMED IS:

1. A method of culturing mesenchymal stem cells (MSCs) comprising culturing a population of the MSCs in a medium comprising nicotinamide and fibroblast growth factor 4 (FGF4), thereby culturing MSCs.
2. The method of claim 1, wherein said medium comprises DMEM.
3. The method of claim 1, wherein said medium comprises serum or platelet lysate.
4. The method of claim 1, wherein the mesenchymal stem cells are derived from a tissue selected from the group consisting of bone marrow, adipose tissue, placenta and umbilical cord blood.
5. The method of claim 1, wherein said nicotinamide is selected from the group consisting of nicotinamide, a nicotinamide analog, a nicotinamide metabolite, a nicotinamide analog metabolite and derivatives thereof.
6. The method of claim 1, wherein said culturing is effected on a plastic surface.
7. The method of claim 1, wherein said population of MSCs is comprised in a heterogeneous population of cells.
8. The method of claim 7, wherein at least 70 % of said heterogeneous population of cells are MSCs.
9. The method of claim 1, wherein a calcium concentration of said medium is greater than 1.8 mM.

10. The method of claim 1, wherein said culturing is effected for at least 1 week.
11. The method of claim 1, wherein said culturing is effected for at least 3 passages.
12. The method of claim 1, wherein a concentration of said nicotinamide is 1-20 mM.
13. The method of claim 1, wherein said medium is devoid of platelet derived growth factor (PDGF).
14. A method of expanding a population of mesenchymal stem cells, the method comprising culturing a seeded population of mesenchymal stem cells for a period of time sufficient for cell expansion, wherein for at least a portion of said period of time said culturing is effected in a medium devoid of nicotinamide; and for at least a second portion of said period of time, said culturing is effected in a medium comprising nicotinamide and FGF4, thereby generating an expanded population of mesenchymal stem cells.
15. The method of claim 14, wherein said expanding is effected under conditions that do not induce differentiation of said mesenchymal stem cells.
16. The method of claim 14, wherein said seeded population of mesenchymal stem cells were seeded in an absence of nicotinamide.
17. The method of claim 14, wherein said seeded population of mesenchymal stem cells were seeded in a presence of nicotinamide.
18. The method of claim 14, wherein said medium is devoid of platelet derived growth factor (PDGF).

19. The method of claim 14, wherein said medium devoid of nicotinamide is devoid of FGF4.

20. The method of claim 14, wherein said medium devoid of nicotinamide comprises FGF4

21. The method of claim 14, wherein said culturing in said medium comprising nicotinamide is effected prior to said culturing in said medium devoid of nicotinamide.

22. The method of claim 14, wherein said culturing in said medium devoid of said nicotinamide is effected prior to said culturing in said medium comprising nicotinamide.

23. The method of claim 14, wherein said culturing in said medium comprising nicotinamide is effected for at least one day.

24. The method of claim 14, wherein said culturing in said medium comprising nicotinamide is effected for at least one week.

25. The method of claim 14, wherein said culturing in said medium devoid of nicotinamide is effected for at least one day.

26. The method of claim 14, wherein said culturing in said medium devoid of nicotinamide is effected for at least one week.

27. The method of claim 14, wherein said culturing in said medium comprising nicotinamide is effected in a medium comprising calcium, wherein a concentration of said calcium is greater than 1.8 mM.

28. The method of claim 14, wherein said culturing in said medium devoid of nicotinamide is effected in a medium comprising calcium, wherein a concentration of said calcium is greater than 1.8 mM.

29. A method of generating cells useful for transplantation into a subject, the method comprising:

(a) culturing mesenchymal stem cells according to the method of claim 1 to generate a population of cultured mesenchymal stem cells;

(b) contacting said population of cultured mesenchymal stem cells with a differentiating agent, thereby generating cells useful for transplantation into a subject.

30. A method of generating cells useful for transplantation, the method comprising:

(a) expanding mesenchymal stem cells according to the method of claim 14; and

(b) contacting the mesenchymal stem cells with a differentiating agent, thereby generating cells useful for transplantation.

31. The method of claims 29 or 30, wherein said differentiation agent comprises a growth factor.

32. The method of claims 29 or 30, wherein said differentiation agent comprises a polynucleotide which encodes said differentiation agent.

33. The method of claim 32, wherein said polynucleotide encodes bone morphogenic protein 2 (BMP2).

34. An isolated population of mesenchymal stem cells generated according to the method of claims 1 or 14.

35. An isolated population of differentiated cells generated according to the method of claims 29 or 30.

36. The isolated population of mesenchymal stem cells of claim 34, being substantially homogeneous.

37. The isolated population of mesenchymal stem cells of claim 34, wherein at least 40 % of the cells express VCAM1/CD106.

38. The isolated population of mesenchymal stem cells of claims 34, wherein at least 90 % of the cells have a diameter less than 20 μm .

39. The isolated population of mesenchymal stem cells of claim 34, being less granular than mesenchymal stem cells generated under identical conditions but in an absence of nicotinamide.

40. The isolated population of mesenchymal stem cells of claims 34, wherein less than 30 % of the cells express CD45, more than 95 % of the cells express CD90 and more than 90 % of the cells express CD105 and CD44.

41. A method of treating a disease or disorder, the method comprising transplanting to a subject in need thereof a therapeutically effective amount of the isolated population of cells of any of the claim 34 or 35, thereby treating the disease or disorder.

42. The method of claim 41, wherein said disease or disorder is selected from the group consisting of a bone or cartilage disease, a neurodegenerative disease, a cardiac disease, a hepatic disease, cancer, nerve damage, autoimmune disease, GvHD, wound healing and tissue regeneration.

43. A cell culture, comprising mesenchymal stem cells and a medium which comprises nicotinamide and FGF4.

FIG. 1

DA#156




FIG. 2A





FIG. 2B



2/38

FIG. 3B




FIG. 3D




FIG. 3A





FIG. 3C




CTRL FGF4 NAM 5 NAM 5+FGF4

Cell count

FIG. 4A

FIG. 4B

Cell size



5/38

CD105+45 Expression (%)




FIG. 5A

FIG. 5B




FIG. 5C

FIG. 5D

ctrl 50 ng/ml FGF4 NAM+50 ng/ml FGF4

FIG. 6A




FIG. 6B




FIG. 6C





FIG. 6D



7/38

FIG. 7A




FIG. 7B




FIG. 7C





FIG. 7D



% expression

FIG. 8B





FIG. 8A



DA#161



FIG. 9B

DA#158




FIG. 9A




FIG. 10A

ctrl




FIG. 10B

FGF-4




FIG. 10C

PDGF-BB






FIG. 10D


FGF4+PDGF-BB




NAM



NAM+FGF-4



NAM+PDGF-BB



NAM+FGF4+PDGF-BB




FIG. 10E

FIG. 10F

FIG. 10G

FIG. 10H

FIG. 11




FIG. 12A





FIG. 12B




FIG. 12C




FIG. 13

FIG. 14




FIG. 15

Bone marrow-MSCs - Cumulative cell count




FIG. 16

Adipose-MSCs - Cumulative cell count




FIG. 17A

Rapid expansion
(7 days after passage 2)



FIG. 17B

Number of cells in culture






FIG. 18





20/38




FIG. 21B





FIG. 21A



22/38




FIG. 22

FIG. 23A





FIG. 23B







FIG. 23C

FIG. 23D



24/38




FIG. 24A




FIG. 24B



FIG. 25A

NAM 0




FIG. 25B

NAM 5




FIG. 26

FIG. 27




FIG. 28

ELISA HGF




FIG. 29

ELISA TGFβ




FIG. 30

ELISA KGF




FIG. 31

ELISA IL6




FIG. 32




FIG. 33




FIG. 34




FIG. 35

The influence of FGF4 on ex-vivo expanded CD133+ cells differentiation




FIG. 36

The influence of FGF4 on ex-vivo expanded CD133+ cells differentiation

CD38+ (%)




FIG. 37

The influence of FGF4 on ex-vivo expanded CD133+ cells differentiation





FIG. 38

The influence of FGF4 on ex-vivo expanded CD133+ cells differentiation

CD19+ (%)



INTERNATIONAL SEARCH REPORT

International application No
PCT/IL2013/050136

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12N5/0775
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, BIOSIS, CHEM ABS Data, EMBASE, INSPEC, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KR 2009 0065814 A (UNIV DONGGUK IND ACAD COOP [KR]) 23 June 2009 (2009-06-23) paragraph [0104] - paragraph [0105] claims 1-10	1-43
X	----- "13th Annual Meeting on Surgical Research/13. Chirurgische Forschungstage; 10.-12. September 2009, HÄRsaaltrakt Klinikum Großhadern, Munich, Germany", LANGENBECK'S ARCHIVES OF SURGERY, SPRINGER, BERLIN, DE, vol. 394, no. 5, 30 July 2009 (2009-07-30), pages 915-970, XP019740458, ISSN: 1435-2451 Abstract 132 ----- -/--	1-43



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

18 April 2013

Date of mailing of the international search report

02/05/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bayer, Martin

INTERNATIONAL SEARCH REPORT

International application No
PCT/IL2013/050136

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BONORA-CENTELLES A ET AL: "Sequential Hepatogenic Transdifferentiation of Adipose Tissue-Derived Stem Cells: Relevance of Different Extracellular Signaling Molecules, Transcription Factors Involved, and Expression of New Key Marker Genes", CELL TRANSPLANTATION, COGNIZANT COMMUNICATION CORP, US, vol. 18, no. 12, 1 January 2009 (2009-01-01), pages 1319-1340, XP009136899, ISSN: 0963-6897, DOI: 10.3727/096368909X12483162197321 the whole document -----	1-43

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/IL2013/050136

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
KR 20090065814 A	23-06-2009	NONE	
